


**Ichthyofauna of Portugal: Taxonomic diversity, common
and scientific names of marine fishes**

**Ictiofauna de Portugal: Diversidade taxonómica, nomes
comuns e nomes científicos dos peixes marinhos**



**Miguel Carneiro, Rogélia Martins,
Francisco Reiner, Irineu Batista**

LISBOA
2019

FICHA TÉCNICA

Título

Ichthyofauna of Portugal: Taxonomic diversity, common and scientific names of marine fishes.
(Ictiofauna de Portugal: Diversidade taxonómica, nomes comuns e nomes científicos dos peixes marinhos).

Edição

Instituto Português do Mar e da Atmosfera, I.P.
Rua C do Aeroporto de Lisboa
1749-077 Lisboa – Portugal



Revisão técnica

Instituto Português do Mar e da Atmosfera, I.P.

ISBN eletrónico

978-972-9083-21-1

Referência Bibliográfica

CARNEIRO¹, M.; MARTINS¹, R.; REINER², F.; BATISTA¹, I., 2019. Ichthyofauna of Portugal: Taxonomic diversity, common and scientific names of marine fishes. IPMA, I.P., Vol. I, 376 p.

1 - IPMA - Instituto Português do Mar e da Atmosfera, I.P.

2 - Centro Português de Estudo dos Mamíferos Marinhos - Projecto Delfim e Centro Português de Actividades Subaquáticas.

Imagen da capa © EMEPC - Estrutura de Missão para a Extensão da Plataforma Continental.

Atribuição – Não Comercial – Compartilha Igual 4.0 Internacional (CC BY-NC-SA 4.0)

O IPMA, I.P. detém os direitos de autor de parte do conteúdo desta publicação, podendo o mesmo ser utilizado nos termos do contrato CC BY-NC-SA 4.0. Contudo, nos casos em que os autores são nominalmente identificados, os direitos de autor pertencem aos mesmos e não ao IPMA, I.P. e foram publicados sob um contrato de licença que não autoriza a copia desse material.

Contents / Índice

Volume I

Abstract / Resumo.....	6
Acronyms and abbreviations /	
Acrónimos e abreviaturas.....	7
Introduction / Introdução.....	10
Methods / Métodos.....	11
Results / Resultados.....	15
Myxinidae	15
Petromyzontidae.....	15
Hexanchidae.....	16
Chlamydoselachidae.....	16
Rhincodontidae.....	17
Odontaspidae.....	17
Mitsukurinidae.....	17
Lamnidae.....	18
Cetorhinidae.....	19
Alopiidae.....	19
Pentanchidae.....	20
Scyliorhinidae.....	21
Pseudotriakidae.....	22
Triakidae.....	23
Carcharhinidae.....	24
Sphyrnidae.....	28
Dalatiidae.....	29
Etmopteridae.....	30
Somniosidae.....	32
Oxynotidae.....	34
Centrophoridae.....	35
Squalidae.....	38
Echinorhinidae.....	39
Squatinidae.....	39
Torpedinidae.....	40
Rhinobatidae.....	41
Glaucostegidae.....	42
Pristidae.....	42
Rajidae.....	42
Arhynchobatidae.....	49
Dasyatidae.....	49
Gymnuridae.....	50
Myliobatidae.....	51
Rhinopteridae.....	52
Mobulidae.....	52
Chimaeridae.....	53
Rhinochimaeridae.....	54
Acipenseridae.....	55
Megalopidae.....	55
Albulidae.....	55
Halosauridae.....	56
Notacanthidae.....	57

Synaphobranchidae.....	58
Muraenidae.....	60
Chlopsidae.....	63
Derichthyidae.....	63
Ophichthidae.....	64
Muraenesocidae.....	65
Nettastomatidae.....	65
Congridae.....	66
Nemichthysidae.....	68
Serrivomeridae.....	68
Anguillidae.....	69
Cyematidae.....	69
Monognathidae.....	70
Eurypharyngidae.....	70
Saccopharyngidae.....	70
Clupeidae.....	72
Engraulidae.....	74
Alepocephalidae.....	74
Platypteroctidae.....	80
Ariidae.....	82
Argentinidae.....	82
Microstomatidae.....	83
Bathylagidae.....	84
Opisthoproctidae.....	84
Salmonidae.....	85
Gonostomatidae.....	86
Sternoptychidae.....	89
Phosichthyidae.....	92
Stomiidae.....	93
Ateleopodidae.....	104
Aulopidae.....	105
Chlorophthalmidae.....	105
Ipnopidae.....	105
Scopelarchidae.....	106
Notosudidae.....	107
Synodontidae.....	108
Bathysauridae.....	109
Paralepididae.....	109
Anotopteridae.....	112
Evermannellidae.....	112
Omosudidae.....	112
Alepisauridae.....	113
Neoscopelidae.....	113
Myctophidae.....	113
Lampridae.....	127
Lophotidae.....	127
Radiicephalidae.....	128
Trachipteridae.....	128
Regalecidae.....	128
Oreosomatidae.....	129
Parazenidae.....	129
Zeniontidae.....	130

Grammicolepididae.....	130
Zeidae.....	130
Stylephoridae.....	131
Bregmacerotidae.....	131
Macrouridae.....	131
Moridae.....	139
Melanonidae.....	142
Gadidae.....	142
Lotidae.....	144
Phycidae.....	146
Merlucciidae.....	148
Polymixiidae.....	149
Berycidae.....	149
Melamphaidae.....	150
Stephanoberycidae.....	153
Rondeletiidae.....	153
Barbourisiidae.....	153
Cetomimidae.....	153
Trachichthyidae.....	154
Diretmidae.....	155
Anoplogasteridae.....	156
Holocentridae.....	156
Ophidiidae.....	156
Carapidae.....	159
Bythitidae.....	160
Batrachoididae.....	162
Gempylidae.....	162
Trichiuridae.....	164
Scombridae.....	166
Scombrolabracidae.....	171
Centrolophidae.....	171
Nomeidae.....	172
Tetragonuridae.....	173
Stromateidae.....	173
Aulostomidae.....	174
Fistulariidae.....	174
Centriscidae.....	174
Syngnathidae.....	175
Dactylopteridae.....	178
Gobiidae.....	178
Xiphiidae.....	185
Istiophoridae.....	185
Citharidae.....	187
Scophthalmidae.....	187
Paralichthyidae.....	188
Bothidae.....	189
Pleuronectidae.....	191
Soleidae.....	191
Cynoglossidae.....	196
Atherinidae.....	197
Scomberesocidae.....	198
Belonidae.....	199

Hemiramphidae.....	200
Exocoetidae.....	200
Mugilidae.....	202
Gobiesocidae.....	205
Tripterygiidae.....	206
Labrisomidae.....	207
Clinidae.....	207
Blenniidae.....	207
Lophiidae.....	212
Antennariidae.....	213
Chaunacidae.....	214
Caulophrynididae.....	215
Neoceratiidae.....	215
Melanocetidae.....	215
Himantolophidae.....	216
Diceratiidae.....	216
Oneirodidae.....	216
Thaumatichthyidae.....	219
Centrophrynididae.....	219
Ceratiidae.....	219
Gigantactinidae.....	220
Linophrynididae.....	220
Molidae.....	222
Diodontidae.....	223
Tetraodontidae.....	224
Ostraciidae.....	226
Monacanthidae.....	226
Balistidae.....	227
Sebastidae.....	228
Setarchidae.....	230
Scorpaenidae.....	230
Triglidae.....	234
Peristediidae.....	237
Cottidae.....	237
Psychrolutidae.....	237
Cyclopteridae.....	238
Liparidae.....	238
Kyphosidae.....	238
Polyprionidae.....	239
Epigonidae.....	240
Howellidae.....	240
Moronidae.....	241
Serranidae.....	242
Callanthiidae.....	247
Priacanthidae.....	247
Apogonidae.....	248
Pomatomidae.....	248
Rachycentridae.....	249
Echeneidae.....	249
Carangidae.....	250
Coryphaenidae.....	258
Bramidae.....	259

Caristiidae.....	260
Lutjanidae.....	261
Lobotidae.....	262
Haemulidae.....	262
Sparidae.....	263
Sciaenidae.....	275
Mullidae.....	277
Chaetodontidae.....	278
Pomacanthidae.....	278
Cepolidae.....	278
Pomacentridae.....	278
Labridae.....	280
Scaridae.....	287
Zoarcidae.....	288
Anarhichadidae.....	289
Pholidae.....	289
Chiasmodontidae.....	289
Ammodytidae.....	291
Trachinidae.....	292
Uranoscopidae.....	293
Callionymidae.....	293
Draconettidae.....	295
Luvaridae.....	295
Acanthuridae.....	295
Sphyraenidae.....	295
Caproidae.....	296
Gasterosteidae.....	297
Global appraisal of taxonomic diversity /	
Avaliação global da diversidade taxonómica.....	299
Final Comments / Considerações finais.....	300
Acknowledgements / Agradecimentos.....	301
Alphabetical indexes of scientific and common names /	
Índices alfabéticos dos nomes científicos e comuns	
Index of scientific names /	
Índice dos nomes científicos.....	302
Index of the English names /	
Índice dos nomes Ingleses.....	327
Index of the Portuguese names /	
Índice dos nomes Portugueses.....	352
Volume II	
Bibliography and electronic references /	
Bibliografia e referências eletrónicas.....	377

Ichthyofauna of Portugal: Taxonomic diversity, common and scientific names of marine fishes

Ictiofauna de Portugal: Diversidade taxonómica, nomes comuns e nomes científicos dos peixes marinhos

Abstract

This paper presents an updated checklist of the common and scientific names of all taxonomically valid fish species that were reported in the Portuguese marine waters. It includes the most important bibliographic references of the works dedicated to the Portuguese ichthyofauna published since 1765, as well as the registration codes in databases (specimens, tissues or human observation).

This checklist includes 1349 species of which 1191 occur in territorial waters and in the three sub-areas of the Portuguese Exclusive Economic Zone, 71 solely in the Portuguese Continental Shelf Extension area and 87 whose presence in the studied area of this work were considered doubtful.

A total of 874 new Portuguese common names are proposed and the criteria considered in their assignment are presented.

Keywords: Ichthyofauna, Taxonomic diversity, Common and scientific names, Portuguese Exclusive Economic Zone, Extension of the Portuguese Continental Shelf.

Resumo

Este trabalho apresenta uma lista atualizada dos nomes comuns e científicos de todas as espécies de peixes, taxonomicamente válidas, que foram referenciadas nas águas marinhas de Portugal. Inclui as referências bibliográficas mais importantes dos trabalhos dedicados à ictiofauna portuguesa publicados desde 1765, bem como os códigos de registo em bases de dados (espécimes, tecidos ou de observação humana).

Nesta lista incluem-se 1349 espécies em que 1191 ocorrem em águas territoriais e nas três sub-áreas da Zona Económica Exclusiva Portuguesa, 71 exclusivamente na área da Extensão da Plataforma Continental Portuguesa e 87 cuja presença na área de estudo deste trabalho foi considerada duvidosa.

São propostos 874 novos nomes vulgares em português e apresentam-se os critérios considerados na atribuição destes nomes.

Palavras-chave: Ictiofauna, Diversidade taxonómica, Nomes comuns e científicos, Zona Económica Exclusiva Portuguesa, Extensão da Plataforma Continental Portuguesa.

Acronyms and abbreviations / Acrónimos e abreviaturas

- ASFIS** - List of Species for Fishery Statistics Purposes, FAO.
- AMS** - Australian Museum, Sydney (Australia).
- AMNH** - American Museum of Natural History, New York (USA).
- ANSP** - The Academy of Natural Sciences, Philadelphia, Pennsylvania (USA).
- ARC** - Atlantic Reference Centre, Saint Andrews, New Brunswick (Canada).
- AVG** - Aquário Vasco da Gama, Lisbon (Portugal).
- AZ** - Azores EEZ sub-area.
- BMNH** - Natural History Museum, London (UK) (see NHMUK; NHMUK is preferred abbreviation for specimen citation).
- CAS** - California Academy of Sciences, San Francisco (USA).
- CITES** - The Convention on International Trade in Endangered Species of Wild Fauna and Flora.
- CLCS** - Commission on the Limits of the Continental Shelf (United Nations).
- CMS** - The Conservation of Migratory Sharks.
- CSIC** - Consejo Superior de Investigaciones Científicas, Madrid (Spain).
- DATRAS** - Database of Trawl Surveys of ICES.
- DFO-NFLD** - Fisheries and Oceans Canada - Newfoundland.
- Diveboard** - Diveboard - Scuba diving citizen science observations.
- EEZ** - Economic Exclusive Zone
- EMBL** - European Molecular Biology Laboratory, Heidelberg (Germany).
- EMEPC** - Estrutura de Missão para a Extensão da Plataforma Continental [alternatively Portuguese Task Group for the Extension of the Continental Shelf], Paço de Arcos (Portugal).
- ENA** - European Nucleotide Archive (EMBL-EBI).
- FAO** - Food and Agriculture Organization, Rome (Italy).
- FMNH** - Field Museum of Natural History, Zoology Department, Chicago (USA).
- GNM** - Göteborgs Naturhistoriska Museum, Göteborg (Sweden).
- GNUB** - Global Names Usage Bank.
- IBM** - Instituto de Biologia Marítima, Lisbon (Portugal).
- ICES** - International Council for the Exploration of the Sea [alternatively CIEM - Conselho Internacional para a Exploração do Mar], Copenhagen (Denmark).
- ICM** - Instituto de Ciencias del Mar, Barcelona (Spain).
- IFREMER** - Institut Français de Recherche pour l'Exploitation de la Mer, Paris (France).
- IGFA** - International Game Fish Association, Dania Beach (USA).
- IMAR/DOP** - Instituto do Mar / Departamento de Oceanografia e Pescas, Açores (Portugal).
- INIP** - Instituto Nacional de Investigações das Pescas, Lisboa (Portugal) (Current as: IPMA) or Instituto Nacional de Investigação Pesqueira, Luanda (Angola).
- IPMA** - Instituto Português do Mar e da Atmosfera, Lisboa (Portugal).
- IORAS** - P.P. Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow (Russia).
- ISH** - Johann Heinrich von Thünen-Institut (vTI), Bundesforschungsinstitut für Ländliche Räume, Wald und Fischerei, Institut für Seefischerei, Hamburg (Germany).
- KU** - University of Kansas Biodiversity Institute, Kansas (USA).
- LACM** - Natural History Museum of Los Angeles County, Los Angeles (USA).

MAR - Mid-Atlantic Region.

MAR-ECO - The MAR-ECO international research project and an element of the Census of Marine Life, co-ordinated by the University of Bergen and Institute of Marine Research, Flødevigen (Norway).

MB - Museu Bocage [alternatively Museu Nacional de História Natural], Lisboa (Portugal).

MCM - Museu Carlos Machado, Ponta Delgada (Açores).

MCZ - Museum of Comparative Zoology, Cambridge, Massachusetts (USA).

MD - Madeira EEZ sub-area.

MHNUSC - Museo de Historia Natural da Universidade de Santiago de Compostela, Santiago de Compostela (España).

MNHN - Muséum national d'Histoire naturelle, Paris (France).

MOM - Musée Océanographique de Monaco (Monaco).

MZLU - Zoologiska museet, Lunds universitet [Museum of Zoology, Lund University], Lund (Sweden).

NCBI - National Center for Biotechnology Information, U.S. National Library of Medicine.

n.d. - no date.

NHMUK - See BMNH.

NM - nautic miles.

NMSZ - National Museums of Scotland, Edinburgh (Scotland).

NMW - Naturhistorisches Museum, Wien (Austria).

ONU - Organização das Nações Unidas.

PECS - Proposed Area for the Extension of the Portuguese Continental Shelf.

POGID - Plano de Ordenamento e Gestão das Ilhas Desertas.

POGIS - Plano de Ordenamento e Gestão das Ilhas Selvagens.

PT - mainland Portugal EEZ sub-area.

ROM - Royal Ontario Museum, Toronto (Canada).

SAIAB - South African Institute for Aquatic Biodiversity, Grahamstown (South Africa).

SAM - South African Museum, Cape Town (South Africa).

SDSC - SeamountsOnline (seamount biota) (CoML).

SMF - Senckenberg Forschungsinstitut und Naturmuseum [alternatively Senckenberg Research Institute and Natural History Museum], Frankfurt am Main (Germany).

SIO - Scripps Institution of Oceanography, Marine Vertebrate Collection, San Diego (USA).

SOC - Southampton Oceanography Center, Southampton (UK).

TCWC - Biodiversity Research and Teaching Collections, Texas (USA).

UN - United Nations.

USNM - National Museum of Natural History, Smithsonian Institution, Washington D.C.(USA).

vd. - *vide*.

VLIZ - Vlaams Instituut voor de Zee [alternatively Flanders Marine Institute], Ostend (Belgium).

VMM - Suffolk County Vanderbilt Museum, New York (USA).

WoRMS - World Register of Marine Species.

WWF - World Wildlife Fund, Gland (Switzerland).

YPM - Yale University, Peabody Museum of Natural History, New Haven (USA).

ZMB - Museum für Naturkunde, Berlin (Germany).

ZMH - Zoological Museum Hamburg, Hamburg (Germany).

ZMUB - Universitetsmuseet i Bergen [alternatively The University Museum of Bergen], Bergen (Norway).

ZMUC - Københavns Universitet, Zoologisk Museum [alternatively Zoological Museum, University of Copenhagen], Copenhagen (Denmark).

ZMUO - Zoologisk Museum Universitetet i Oslo, Oslo (Norway).

ZSM - Zoologische Staatssammlung München, München (Germany).

Introduction

The fishing activity in Portugal has a long tradition and involves multiple fishing communities along the Portuguese continental coast and the archipelagos of Madeira and Azores. The social and economic importance of this activity has led to a great diversity of common denominations of fish. These denominations result from the input of all stakeholders involved in the marketing chain which includes fishermen, retailers, fishmongers and consumers. They also represent a valuable linguistic heritage.

Much of this activity takes place in the Portuguese Exclusive Economic Zone (EEZ), which includes the sub-areas of mainland Portugal and the archipelagos of Azores and Madeira, and globally representing 1,727,408 km².

In 2009 it was submitted a proposal for the extension of the Portuguese continental shelf beyond 200 nautical miles to the UN which is under consideration in the Commission on the Limits of the Continental Shelf (CLCS). This extension would substantially increase this area to approximately 4,100,000 km².

Beyond 200 nautical miles the rights of sovereignty over the continental shelf comprises only the soil and subsoil (not the water column), throughout the natural extension of its land territory, to the outer edge of the continental margin.

The inclusion of the PECS (Portuguese Extension of the Continental Shelf) area in this work intents to evaluate the fish biodiversity in areas beyond the national jurisdiction.

In 2014 the first global checklist of fish species cited for Portugal was published (Carneiro *et al.*, 2014). Meanwhile, significant changes occurred both in the structure of the taxonomic classification and in the number of species due to the inclusion of new occurrence records, redefinition of the area beyond national jurisdiction, local or regional extinction of species or even corrections of former doubtful or erroneous records.

Introdução

A atividade pesqueira em Portugal tem uma longa tradição e envolve múltiplas comunidades piscatórias distribuídas ao longo da costa continental portuguesa e nos arquipélagos da Madeira e dos Açores. A importância social e económica dessa atividade levou a uma grande diversidade de denominações comuns dos peixes. Essas denominações resultam do contributo dos intervenientes económicos envolvidos na cadeia de comercialização e incluem pescadores, distribuidores, retalhistas, vendedores e consumidores e constituem um valioso património linguístico.

Grande parte desta atividade ocorre na Zona Económica Exclusiva Portuguesa (ZEE) que inclui as sub-áreas do continente e dos arquipélagos dos Açores e da Madeira, totalizando 1 727 408 km².

Em 2009, foi apresentada à ONU uma proposta para a extensão da plataforma continental portuguesa para além das 200 milhas marítimas, que está a ser avaliada na Comissão de Limites da Plataforma Continental (CLCS). Esta extensão aumentaria substancialmente esta área para cerca de 4 100 000 km².

Para além das 200 milhas marítimas, os direitos de soberania sobre a Plataforma Continental compreendem apenas o solo e o subsolo (e não a coluna de água), ao longo da extensão natural do território terrestre, até ao bordo exterior da margem continental.

Com a inclusão da área do PECS (Extensão da Plataforma Continental Portuguesa) neste trabalho procurou-se contribuir para a avaliação da biodiversidade dos peixes em áreas fora da atual jurisdição nacional.

Em 2014 foi publicada a primeira lista global de espécies de peixes cuja ocorrência foi citada para Portugal (Carneiro *et al.*, 2014). Entretanto, registaram-se alterações significativas quer na estrutura da classificação taxonómica, quer no número de espécies devido à inclusão de novos registos de ocorrência, redefinição da área além da jurisdição nacional, extinção local ou regional de espécies ou mesmo correções de anteriores registos duvidosos ou erróneos.

Thus, the main objective of this work was to provide an updated checklist of the Portuguese ichthyofauna based on the *California Academy of Sciences* (Van der Laan *et al.*, 2019) taxonomic classification.

It was also intended: (i) to give information on marketed fish species for the period of 2015-2018 based on the commercial catch records; (ii) to present the diversity of the Portuguese common names traditionally used in the different fishing communities and regions since the last quarter of the 18th century; and (iii) to contribute for the harmonization of the Portuguese common names.

Methods

In this work the criterion for the inclusion of a species in the checklist was based on its occurrence record in the following geographic areas: territorial waters, the Exclusive Economic Zone (EEZ) and the proposed PECS. In figure 1 the PECS area is highlighted.

In the current work, the classification system for taxonomic categories of Van der Laan *et al.* (2019) was followed. The taxonomic categories considered were: superclass, class, order, family and species. In each family, the species were alphabetically sorted and for each species was indicated: name, authority and year, English common name, Portuguese common names, citations and occurrence codes in the different area(s) included in this study.

The English common name of each fish species was based on the FAO-FIES (2018) of the Aquatic Science and Fisheries Information System (ASFIS) and the type FAO name (in bold) or optionally another name when available.

The traditional Portuguese name (the most usual Portuguese vernacular name or proposed Portuguese vernacular name for a future Portuguese FAO designation from Portugal in bold), followed by other designations when available.

Assim, o principal objetivo deste trabalho foi elaborar uma lista atualizada da ictiofauna portuguesa, com a inclusão de novas espécies, tendo por base a classificação taxonómica da *California Academy of Sciences* (Van der Laan *et al.*, 2019).

Pretendeu-se também: (i) disponibilizar informação sobre as espécies de peixes comercializadas durante o quadriénio 2015-2018, com base nos registos das capturas comerciais; (ii) apresentar a diversidade dos nomes vulgares portugueses tradicionalmente utilizados nas diferentes comunidades e regiões piscatórias desde o último quartel do século XVIII; e (iii) contribuir para a harmonização dos nomes portugueses vulgares dos peixes.

Métodos

Neste trabalho o critério de inclusão de uma espécie na lista baseou-se no registo da sua ocorrência nas seguintes áreas geográficas: as águas territoriais, a Zona Económica Exclusiva (ZEE) e a área proposta para a Extensão da Plataforma Continental Portuguesa (PECS). Na figura 1 encontra-se evidenciada a área PECS.

Foi seguido o sistema de classificação para categorias taxonómicas de Van der Laan *et al.* (2019). As categorias taxonómicas consideradas foram as seguintes: superclasse, classe, ordem, família e espécie. Em cada família, as espécies foram ordenadas alfabeticamente e para cada espécie foi indicado: nome, autoridade e ano, nome vulgar inglês, nomes vulgares portugueses, citações e códigos de ocorrência nas diferente(s) área(s) incluídas neste estudo.

O nome vulgar em inglês de cada espécie foi baseado na FAO-FIES (2018) do Sistema de Informação sobre Ciências Aquáticas e Pesca (ASFIS), indicando-se a denominação FAO (em negrito) ou opcionalmente outro nome, quando disponível.

O nome tradicional português (o mais comum (em negrito) ou o proposto para uma futura denominação FAO portuguesa (de Portugal) (a azul) e, quando disponíveis, outras denominações.




Fig. 1. Map of the study area - the Portuguese EEZ, that includes the territorial waters (grey lines) and the area proposed for the extension of the Portuguese continental shelf (yellow line) (source: EMEPC – Mission Structure for the Extension of the Continental Shelf).

[Mapa da área do estudo - ZEE portuguesa onde se incluem as águas territoriais (linhas a cinzento) e a área proposta para a Extensão da Plataforma Continental Portuguesa (PECS) (linhas a amarelo)(fonte: EMEPC – Estrutura de Missão para a Extensão da Plataforma Continental)]

The most important reference works followed for the Portuguese common names were: Albuquerque (1954–1956), Osório de Castro (1953, 1967), Nunes (1974), Sanches (1986, 1989) and the list of common names of the Portuguese legislation (Annex I and II of the “Portaria nº 587/2006” – Lista de Nomes Comerciais Autorizados de Produtos da Pesca e Aquicultura, and “Declaração de Rectificação no. 52/2006”).

Annex I of the “Portaria (Governamental Order) nº 21/2019 de 19 de março de 2019 (JO RA Azores)”.

The proposed Portuguese common name of each species was based on the following criteria:

- i. The common Portuguese name traditionally used in fishing communities mentioned in reference books or used in the official trade names. For example, *Sardina pilchardus* (Walbaum 1792) - Sardinha;

As principais obras de referência consultadas para os nomes vulgares portugueses foram: Albuquerque (1954-1956), Osório de Castro (1953, 1967), Nunes (1974), Sanches (1986, 1989) e a lista de nomes comuns da legislação portuguesa (Anexo I e II da Portaria nº 587/2006 - Lista de Nomes Comerciais Autorizados de Produtos da Pesca e Aquicultura e Declaração de Rectificação nº 52/2006).

Anexo I da Portaria n.º 21/2019 de 19 de março de 2019 (JO RA Açores).

O nome vulgar em português proposto para cada espécie baseou-se nos seguintes critérios:

- i. O nome vulgar em português tradicionalmente usado nas comunidades pesqueiras, mencionado em obras de referência ou utilizado nas denominações comerciais oficiais. Por exemplo, *Sardina pilchardus* (Walbaum 1792) - Sardinha;

- ii. The name of the person to whom the fish species is dedicated, the name of a scientific research vessel or the name of a scientific expedition. For example, *Bathytroctes michaelarsi* Koefoed 1927 - Celindra-de-Michael-Sars;
- iii. The Portuguese translation of the specific epithet of the scientific name. For example, *Deltentosteus quadrimaculatus* (Valenciennes 1837) - Caboz-de-quatro-manchas;
- iv. The Portuguese translation of the common name already existing in another language. For example, Copper shark (*Carcharhinus brachyurus* (Günther 1870)) - Tubarão-cobre;
- v. The name of the place where the holotype was caught for the first time. For example, *Sphyraena viridensis* Cuvier, 1829 - Bicuda-de-Cabo-Verde;
- vi. The name of the scientist(s) who first published the scientific species name. For example, *Diaphus adenomus* Gilbert 1905 - Peixe-lanterna-de-Gilbert;
- vii. The general external appearance, specific morphological features (position of certain specialized structures, filaments, fins, spines, photophores, illicium, etc.) or the *in vivo* coloration(s). For example, all members of family Linophrynidae are named Peixe-sestro due to the peculiar displacement of the vent or anal opening to the left of center (sinistral);
- viii. The type of habitat, the marine habitat division occupied, the geographic distribution area or the characteristic behaviour of the species. For example, *Cyclothona obscura* Brauer 1902 - Boqueirão-sombrio;
- ix. The combination of two or more criteria mentioned above. For example, *Nerophis lumbriciformis* (Jenyns 1835) - Marinha-lombriga.

Finally, the references (bibliographic or database) and codes (specimen, tissue or human observation) on the occurrence in the different area(s) are included. The codes were used in order to consolidate the bibliographic quotations of the most important works on the occurrence of the species in each area and sometimes they are the only reference of its presence in a given geographic area.

- ii. O nome da pessoa a quem a espécie é dedicada, o nome de uma embarcação de investigação científica ou o nome de uma expedição científica. Por exemplo, *Bathytroctes michaelarsi* Koefoed 1927 - Celindra-de-Michael-Sars;
- iii. A tradução portuguesa do restritivo específico do nome científico. Por exemplo, *Deltentosteus quadrimaculatus* (Valenciennes 1837) - Caboz-de-quatro-manchas;
- iv. A tradução portuguesa do nome comum já existente noutro idioma. Por exemplo, Copper shark (*Carcharhinus brachyurus* (Günther 1870)) - Tubarão-cobre;
- v. O nome do local onde o holótipo foi capturado pela primeira vez. Por exemplo, *Sphyraena viridensis* Cuvier, 1829 - Bicuda-de-Cabo-Verde;
- vi. O nome do(s) cientista(s) que primeiro publicou(aram) o nome científico da espécie. Por exemplo, *Diaphus adenomus* Gilbert 1905 - Peixe-lanterna-de-Gilbert;
- vii. O aspecto externo geral, características morfológicas específicas (posição de certas estruturas especializadas, filamentos, barbatanas, espinhos, fotóforos, *illicium*, etc.) ou a(s) coloração(ões) *in vivo*. Por exemplo, todos os membros da família Linophrynidae são denominados Peixe-sestro devido à posição peculiar da abertura anal à esquerda do eixo central;
- viii. O tipo e o andar do habitat marinho ocupado, a área de distribuição geográfica ou o comportamento característico da espécie. Por exemplo, *Cyclothona obscura* Brauer 1902 - Boqueirão-sombrio;
- ix. A combinação de dois ou mais critérios supra mencionados. Por exemplo, *Nerophis lumbriciformis* (Jenyns 1835) - Marinha-lombriga.

Seguem-se as referências (bibliográficas ou base de dados) e os códigos (espécime, tecido ou observação humana) sobre a ocorrência nas diferentes áreas. Os códigos foram incluídos a fim de consolidar as citações bibliográficas dos trabalhos mais importantes sobre a ocorrência das espécies em cada área e de modo a permitir um contínuo temporal de registos da sua ocorrência. Por vezes, são a única referência da sua presença numa determinada área geográfica.

All occurrence records in the PECS area are indicated. Furthermore, all species considered native due to its geographic distribution in one of the three EEZ sub-areas, but without specific occurrence records, are included and commented.

The sign (†) stands for an extinct, apparently extinct or regionally extinct species; ▲ stands for a species with commercial catches for the period of 2015-2018; the question mark (?) located after the name of a specific area indicates that the species was erroneously included in this area, misidentified or its current geographic distribution is different; if the question mark (?) is after a reference it indicates that the author had pointed out as a doubtful presence and/or have not a positive identification of the species and the record was questionable or more information was needed.

When the PECS acronym stands after the name of a particular area (e.g. Azores) it means that all references or codes refer to occurrence records in their area of extension; if this acronym is placed after a quotation indicates that it only refers to an occurrence in its area of extension.

Finally, where appropriate, comments are included on:

- The existence of type specimens: holotypes, syntypes, neotypes, paratypes and lectotypes caught in Portuguese waters or under Portuguese jurisdiction;
- Synonymy of invalid *taxa* designations;
- Indication of species that may be confused with the concerned species;
- The current geographic distributions of the species;
- The need of specimens to confirm the occurrence of the species in a certain area;
- Reasons for not including the species whose occurrence is considered doubtful;
- Existence of contradictory quotations on the presence of the species in a given area.

Todos os registos de ocorrência na área do PECS são indicados. Além disso, são também incluídas as espécies consideradas nativas, devido à sua distribuição geográfica numa das três sub-áreas da ZEE, embora não existam registos específicos da sua ocorrência.

O sinal (†) significa uma espécie extinta, aparentemente extinta ou regionalmente extinta; o sinal ▲ representa uma espécie com capturas comerciais durante o quadriénio de 2015-2018; o ponto de interrogação (?) localizado após o nome de determinada área indica que a espécie foi erroneamente incluída nessa área, mal identificada ou a sua atual distribuição geográfica é diferente; se o ponto de interrogação (?) estiver depois de uma citação indica que o autor apontou a sua presença e/ou identificação como duvidosas, sendo o registo da sua ocorrência questionável ou necessitando de mais informação.

Quando a sigla PECS se encontra após o nome de uma determinada área (por exemplo, Açores) isso significa que todas as referências ou códigos dizem respeito a ocorrências na respetiva área de extensão; se esta sigla estiver colocada depois de uma citação indica que diz respeito apenas a uma ocorrência na respetiva área de extensão.

Por último, sempre que oportuno, são incluídos comentários sobre:

- Existência de espécimes-tipo: holótipos, síntipos, neótipos, parátipos e lectótipos capturados em águas portuguesas ou sob jurisdição portuguesa;
- Sinonímia de denominações de *taxa* não válidas;
- Indicação de espécie(s) que se possa(m) ser confundida(s) com a espécie em causa;
- Atuais distribuições geográficas de ocorrência da espécie;
- A necessidade de espécimes para confirmação da ocorrência da espécie numa determinada área;
- Razões para a não inclusão da espécie cuja presença seja considerada duvidosa;
- Existência de citações contraditórias sobre a presença da espécie numa determinada área.

Results / Resultados

Superclass Myxinomorphi
Class Myxini
Order Myxiniformes
Family Myxinidae

Myxine glutinosa Linnaeus 1758

Common name: **Hagfish; Enguia-de-casulo.**

References: mainland Portugal ([MNHN-IC-1883-0150](#), 1882; [AVG CODC-Peixes PE/03](#), 1899; Bragança, 1903; Brauer, 1906; Seabra, 1911; Osório, 1917; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Vladýkov, 1979; Fernández *et al.*, 1981; Fernholm and Vladýkov, 1984; Sanches, 1986; Fischer *et al.*, 1987; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Garcia, 2015a; Fernholm and Mincarone, 2016; Mecklenburg *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22498, 1967).

Superclass Petromyzontomorphi
Class Petromyzonti
Order Petromyzontiformes
Family Petromyzontidae

Lampetra fluviatilis (Linnaeus 1758)

Common name: **European river lamprey; Lampreia-do-rio**, Lampreia, Lampreia-de-água-doce, Lampreia-fluvial.

References: mainland Portugal ([Baldaque da Silva, 1891](#); [Vieira, 1894a, b, 1897, 1901](#); Osório, 1896; Nobre, 1903, 1931, 1932, 1935; Seabra, 1911; Springer, 1911; Albuquerque, 1954-1956; Osório de Castro, 1967; Vladýkov, 1984; Sanches, 1986; Fischer *et al.*, 1987; Almaça, 1995; Carneiro *et al.*, 2014; Mateus *et al.*, 2016; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Froese and Pauly, 2019).

Petromyzon marinus Linnaeus 1758

Common name: **Sea lamprey; Lampreia-do-mar**, Lamprede, Lampreia, Lampreia-marinha.

References: mainland Portugal[▲] ([Carvalho, 1765](#); [Vandelli, 1771, 1797](#); [de Brito Capello, 1869a, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1894a, b, 1897, 1901](#); Nobre, 1903, 1931, 1932, 1935; Seabra, 1911; Springer, 1911; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Vladýkov, 1979, 1984; Sanches, 1986; Fischer *et al.*, 1987; Collares-Pereira *et al.*, 2000; Garnerot *et al.*, 2004; Carneiro *et al.*, 2014; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Froese and Pauly, 2019) and Madeira ([Ribeiro *et al.*, 2005a](#); Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Superclass Gnathostomata
Class Chondrichthyes
Sub-class Elasmobranchii
Order Hexanchiformes
Family Hexanchidae

Heptranchias perlo (Bonnaterre 1788)

Common name: **Sharpnose sevengill shark; Tubarão-de-sete-guelras**, Archote, Albafar-bravo, Bico-doce, Bico-doce-do-alto, Boca-doce, Boca-doce-do-alto, Cação, Cação-severino, Olho-branco, Rinquim, Severino, Tubarão, Tubarão-severino.

References: mainland Portugal ([de Brito Capello, 1869a, 1880](#); [Vieira, 1893, 1897](#); Nobre, 1894, 1935; Bragança, 1904a; Seabra, 1911; Springer, 1911; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Boeseman, 1984; Sanches, 1986; Fischer *et al.*, 1987; Ward *et al.*, 2008; Rodrigues 2009; Carneiro *et al.*, 2014; Moura *et al.*, 2015a), Azores[▲] ([Ferreira, 1939](#); [Collins, 1954](#); Osório de Castro, 1967; Dias *et al.*, 1976; Ehrich, 1977; Martins, 1982a; Boeseman, 1984; INIP, 1985a; Sanches, 1986; Dias, 1991; Arruda, 1997; Santos *et al.*, 1997; Fock *et al.*, 2002 (PECS); Kukuev, 2002 (PECS); Menezes *et al.*, 2006; Correia, 2009; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and

Stehmann, 2013; Carneiro *et al.*, 2014; Torres *et al.*, 2016; Das and Afonso, 2017; ICES, 2017a; Biscoito *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira (Lowe, 1837; Günther, 1870; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Ehrich, 1977; Boeseman, 1984; Sanches, 1986; Carneiro *et al.*, 2014; Compagno, 2016; POGID, 2016; Biscoito *et al.*, 2018). One specimen is in the MMF collections (MMF 36031, 1961).

Hexanchus griseus (Bonnaterre 1788)

Common name: **Bluntnose sixgill shark; Tubarão-albafar**, Albafar, Albaftara, Albafor, Albaftora, Albufara, Albufasa, Cação, Fernesura, Olho-grande, Olho-verde, Rinquim, Rosa, Tubarão, Tubarão-albatara, Tubarão-albafar-bravo, Tubarão-de-seis-guelras.

References: mainland Portugal▲ (Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Almeida and Roquette, 1892(?); Nobre, 1894, 1935; Vieira, 1897; Bragança, 1903, 1904a; Seabra, 1911; Springer, 1911; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981, 1987; Boeseman, 1984; Compagno, 1984a; Sanches, 1986; Borges *et al.*, 2002; ICES, 2005; Melendez and Macias, 2007; King *et al.*, 2008; Bordalo-Machado *et al.*, 2009; Correia, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Walls *et al.*, 2015a; Froese and Pauly, 2019), Azores (Ferreira, 1939; Collins, 1954; Osório de Castro, 1967; Wheeler, 1978; Martins, 1982a; Boeseman, 1984; Sanches, 1986; Dias *et al.*, 1990; Dias, 1991; Arruda, 1997; Santos *et al.*, 1997; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Menezes *et al.*, 2009; Porteiro *et al.*, 2010, 2013; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Besugo, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Walls *et al.*, 2015a; Compagno, 2016; Torres *et al.*, 2016; Das and Afonso, 2017; ICES, 2017a; Biscoito *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira (Lowe, 1837; Belloc, 1934; Noronha and Sarmento 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Boeseman, 1984; INIP, 1984a; Sanches, 1986; Carneiro *et al.*, 2014; Walls *et al.*, 2015a; Compagno, 2016; Biscoito *et al.*, 2018). According to Ebert and Stehmann (2013) small specimens of *Hexanchus griseus*, usually less than 120 cm in length, may be confused with *H. nakamurai*.

Hexanchus nakamurai Teng 1962

Common name: **Bluntnose sixgill shark; Tubarão-albafar-de-pontas-brancas**.

References: mainland Portugal (Boeseman, 1984(?); Compagno and Niem, 1998; Melendez and Macias, 2007; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019). This species is sometimes confused with *Hexanchus griseus* (Bonnaterre 1788) (Soldo, 2015). According to Ebert *et al.* (2013) the current geographic distribution of this species in the Eastern North and Central Atlantic includes the Bay of Biscay, off France, Spain, Gibraltar, Morocco, and possibly the Ivory Coast and Nigeria, and also the western Mediterranean. Specimens are needed for confirmation the presence of this species in mainland Portugal.

Family Chlamydoselachidae

Chlamydoselachus anguineus Garman 1884

Common name: **Frilled shark; Tubarão-cobra**, Cobra-da-fundura.

References: mainland Portugal (Bragança, 1903, 1904a; Springer, 1911; Osório, 1917; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Fischer *et al.*, 1981; Boeseman, 1984; Compagno, 1984a; Sanches, 1986; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2015a, b; Compagno, 2016; Froese and Pauly, 2019; ICES 2017; Froese and Pauly, 2019), Azores (Melo and Menezes, 2002; Kukuev, 2002 (PECS); Kukuev and Pavlov, 2008 (PECS); Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Das and Afonso, 2017; Biscoito *et al.*, 2018) and Madeira (Collet, 1890; Goode and Bean, 1895, 1896; Brauer, 1906; Belloc, 1934; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d; Albuquerque, 1954-1956; Osório de Castro, 1967; Boeseman, 1979, 1984; Cadenat and Blache 1981; Fischer *et al.*, 1981; Compagno, 1984a, 2016; Sanches, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Delgado *et al.*, 2017; Biscoito *et al.*, 2018; Froese and Pauly, 2019).

Order Orectolobiformes
Family Rhincodontidae

Rhincodon typus Smith 1828

Common name: **Whale shark; Tubarão-baleia**, Caldeirão, Malhado, Pintado, Tubarão-malhado, Tubarão-pintado.

References: mainland Portugal ([Rodrigues, 2009](#); [Carneiro et al., 2014](#)), Azores ([Santos et al., 1994a, 1995a, 1997](#); [Pinho and Pereira, 1995b](#); [Arruda, 1997](#); [Compagno 2001, 2016](#); [CITES, 2003](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Pierce et al., 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Compagno, 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)).

Order Lamniformes
Family Odontaspidae

Carcharias taurus Rafinesque 1810

Common name: **Smalltooth sand tiger shark; Tubarão-toiro**, Albafar, Dentudo, Tubarão, Tubarão-amarelo, Tubarão-de-areia, Tubarão-dentudo.

References: mainland Portugal ([Osório de Castro, 1967](#)), Azores ([ZMH 25521, 1938](#); [IMAR/DOP 1PAL-04-P05TAU, 2005](#); [Schroeter, 2016 \(PECS\)](#)) and Madeira ([Osório de Castro, 1967](#); [Fischer et al., 1981](#); [Walls et al., 2015b](#)). Since Osório de Castro (1967) no other author described the presence of this species in mainland Portugal and its occurrence in this EEZ was not yet confirmed by more recent data. The maps of distribution of this species reported by Fischer et al. (1981) and Walls et al. (2015b) included the area of Selvagens Islands (Madeira EEZ). According to Golani and Fricke (2018) and Eschmeyer et al. (2019) *C. taurus* is a circumglobal species in tropical through warm temperate seas. Specimens are needed to confirm its presence in mainland Portugal and Madeira.

Odontaspis ferox (Risso 1810)

Common name: **Sand tiger shark; Tubarão-areia**, Albafar, Tubarão, Tubarão-de-areia.

References: mainland Portugal ([Osório de Castro, 1967](#); [Graham et al., 2016](#)), Azores ([Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Naylor et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Pollard et al., 2015](#); [Compagno, 2016](#); [Graham et al., 2016](#); [Das and Afonso, 2017](#); [Barcelos et al., 2018](#); [Biscoito et al., 2018](#)) and Madeira ([Maul, 1955a](#); [Springer, 1979a](#); [Cadenat and Blache 1981](#); [Fischer et al., 1981](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Pollard et al., 2015](#); [Graham et al., 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). Two specimens are in the MMF collections (MMF 2478, 1941; MMF 2678, 1941).

Odontaspis noronhai (Maul 1955)

Common name: **Bigeye sand tiger shark; Tubarão-de-Noronha**

References: Madeira ([Maul, 1955a](#); [Springer, 1979a](#); [Cadenat and Blache 1981](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Carcharias noronhai* Maul 1955, synonym of *O. noronhai*, is in the MMF collections (MMF 2691, 1941) caught off Câmara de Lobos (Madeira).

Family Mitsukurinidae

Mitsukurina owstoni Jordan 1898

Common name: **Goblin shark; Tubarão-duede**, Demónio, Narigudo, Tubarão, Tubarão-demónio, Tubarão-narigudo.

References: mainland Portugal ([Bragança, 1903, 1904a, b](#); [Osório, 1909](#); [Seabra, 1911](#); [Springer, 1911](#); [Nobre, 1935](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Osório de Castro, 1954, 1967](#); [Springer, 1979a](#); [Cadenat and Blache 1981](#); [Fischer et al., 1981](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Prokofiev and Kukuev, 2009a](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Moura et al., 2015a, b](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)) and Madeira ([Cadenat and Blache 1981](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Ebert and Stehmann, 2013](#)).

2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018; Froese and Pauly, 2019). One specimen in the MMF collections (MMF 8196, 1956).

Family Lamnidae

Carcharodon carcharias (Linnaeus 1758)

Common name: **Great white shark; Tubarão-branco**, Cascarra, Comedor-de-homens, Gandú, Rinquim, Tubarão, Tubarão-azul, Tubarão-come-homens, Tubarão-de-São-Tomé, Tubarão-grande.
 References: mainland Portugal (Vandelli, 1771; Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Baldaque da Silva, 1891; Vieira, 1897; Bragança, 1904a; Seabra, 1911; Springer, 1911; Albuquerque, 1954-1956; Springer, 1979a; Fischer *et al.*, 1981, 1987; Compagno, 1984a, 2001, 2016; Quéro, 1984a; Sanches, 1986(?); Melendez and Macias, 2007; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; ICES, 2017a; Froese and Pauly, 2019), Azores (Ferreira, 1939; SAMS-SHARKS-002206, 1949; Collins, 1954; Osório de Castro, 1967; Saldanha, 1980, 1995; Ellis and McCosker, 1991; Bruun, 1992; Santos *et al.*, 1994a, 1995a, 1997; Arruda, 1997; Fowler *et al.*, 2005; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Compagno, 2016; Das and Afonso, 2017; Bischoff *et al.*, 2018) and Madeira (Lichtenstein, 1844; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Fischer *et al.*, 1981; Compagno 1984a, 2001, 2016; Quéro, 1984a; Sanches, 1986; Lloris *et al.*, 1991; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018). Wirtz *et al.* (2008) do not indicate the presence of this species as a coastal fish of Madeira, but confirm its existence recorded from the open sea. According to Froese and Pauly (2019) this species is absent of Madeira because there was a misunderstanding of the text by Wirtz *et al.* (2008).

Isurus oxyrinchus Rafinesque 1810

Common name: **Longfin mako; Tubarão-anequim**, Anequim, Anequim-marrachão, Aniquim, Arrequim, Cão, Enequim, Marrachão, Marracho, Marracho-azul, Melracho, Merracho, Panhaço, Peixe-agunha, Peixe-cão, Peixe-ruim, Rinquim, Sardo, Tubarão, Tubarão-azul, Tubarão-da-praia, Tubarão-palhaço.
 References: mainland Portugal▲ (Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Almeida and Roquette, 1892; Bragança, 1903, 1904a; Seabra, 1911; Springer, 1911; Nobre, 1935; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Springer, 1979a; Fischer *et al.*, 1981, 1987; Compagno, 1984a, 2001, 2016; Quéro, 1984a; Dinis, 1985; Sanches, 1986; Erzini *et al.*, 2001a; Santos *et al.*, 2002a; Coelho *et al.*, 2005; Fowler *et al.*, 2005; Melendez and Macias, 2007; Henriques *et al.*, 2008; Bordalo-Machado *et al.*, 2009; Correia, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2015b; ICES, 2017a; Froese and Pauly, 2019), Azores▲ (Hilgendorf, 1888a; Sampaio, 1904; Fowler, 1936a; Ferreira, 1937a, 1939; Bigelow and Schroeder, 1948; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1978; Springer, 1979a; Martins, 1982a; Compagno, 1984a, 2001, 2016; Quéro, 1984a; Sanches, 1986; Moreno and Morón, 1992; Pinho and Pereira, 1995a; Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Correia, 2009; Porteiro *et al.*, 2010, 2017(?); Barreiros and Gadig, 2011; Besugo, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Torres *et al.*, 2016; Das and Afonso, 2017; ICES 2017; Bischoff *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira▲ (Günther, 1870; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cadenat and Blache 1981; Compagno, 1984a, 2001, 2016; Quéro, 1984a; Springer, 1979a; Sanches, 1986; Wirtz *et al.*, 2008; Correia, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; POGID, 2016; ICES, 2017a; Bischoff *et al.*, 2018; Froese and Pauly, 2019).

Isurus paucus Guitart Manday 1966

Common name: **Longfin mako; Tubarão-anequim-de-gadanha**, Anequim, Marracho, Marajo-negro, Peixe-ruim, Tubarão-anequim.
 References: mainland Portugal (Muñoz-Chápuli, 1984; Quéro, 1984a; Compagno, 2001(?), 2016; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Reardon *et al.*, 2015; ICES, 2017a; Froese and Pauly, 2019(?)), Azores (Queiroz *et al.*, 2008; Porteiro *et al.*, 2010; Ebert and Stehmann, 2013(?); Martins, 2013; Carneiro *et al.*, 2014; Reardon *et al.*, 2015; Compagno, 2016; Das and Afonso, 2017) and Madeira (Ebert and Stehmann, 2013(?); Reardon *et al.*, 2015; Compagno, 2016). This species is sometimes confused with the more common *Isurus oxyrinchus* Rafinesque 1810 (Compagno 2001).

***Lamna nasus* (Bonnaterre 1788)**

Common name: **Porbeagle; Tubarão-sardo**, Agulha, Anequim, Aniquim, Arrequim, Barrilote, Cão, Enequim, Lobo-do-mar; Marcacho, Marracho, Melracho, Merracho, Nequim, Peixe-agulha, Peixe-cão, Peixe-ruim, Requim, Rinquim, Sardão, Sardo, Tubarão, Tubarão-anequim, Tubarão-azul, Tubarão-agulha, Tubarão-sarda, Tubarão-toupeira.

References: mainland Portugal ([Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, 1880](#); [Gervais and Boulart, 1877a](#); [Almeida and Roquette, 1892](#); [Vieira, 1897](#); [Bragança, 1904a](#); [Seabra, 1911](#); [Springer, 1911](#); [Osório, 1917](#); [Nobre, 1935, 1937](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Springer, 1979a](#); [Fischer et al., 1981, 1987](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Melendez and Macias, 2007](#); [Correia, 2009](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967](#); [Wheeler, 1978](#); [Martins, 1982a](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Dias, 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Fowler et al., 2005](#); [Correia, 2009](#); [Porteiro et al., 2010, 2017\(?\)](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Torres et al., 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Lowe, 1837](#); [Belloc, 1934](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Fischer et al., 1981](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [INIP, 1984a](#); [Sanches, 1986](#); [Severino, 2004](#); [Wirtz et al., 2008\(?\)](#); [Correia, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019\(?\)](#)).

Family Cetorhinidae

***Cetorhinus maximus* (Gunnerus 1765)**

Common name: **Basking shark; Tubarão-frade**, Albafar, Albafar-manso, Baleia, Carago, Frade, Padre, Peixe-carago, Peixe-frade, Peixe-padre, Peregrino, Rinquim, Soalheiro, Tubarão, Tubarão-baleia, Tubarão-peregrino, Tubarão-soalheiro.

References: mainland Portugal[▲] ([Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, 1880](#); [Vieira, 1897](#); [Bragança, 1903, 1904a](#); [Seabra, 1911](#); [Springer, 1911](#); [Nobre, 1935](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Springer, 1979a](#); [Fischer et al., 1981, 1987](#); [INIP, 1982a](#); [Compagno, 1984a, 2001](#); [Quéro, 1984a](#); [Sanches, 1986](#); [ICES, 2005](#); [Melendez and Macias, 2007](#); [Correia, 2009](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Sims et al., 2015](#); [Witt et al., 2016](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)), Azores ([Ferreira, 1939](#); [Collins, 1954](#); [Clarke, 1956](#); [Osório de Castro, 1967](#); [Wheeler, 1978](#); [Quéro, 1984a](#); [Lloris et al., 1991](#); [Santos et al., 1994a, 1995a, 1997](#); [Arruda, 1997](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Sims et al., 2015](#); [Compagno, 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#)) and Madeira ([Johnson, 1885](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Maul 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Sims et al., 2015](#); [Witt et al., 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). One specimen is in the MMF collections (MMF 606, 1937).

Family Alopiidae

***Alopias superciliosus* Lowe 1841**

Common name: **Pelagic thresher or Thresher; Tubarão-raposo-olhudo**, Peixe-rato, Peixe-rato-de-natura, Peixe-zorro, Raposo, Raposo-de-olhos-grandes, Raposo-olhudo, Romano, Romão, Tubarão-raposa, Tubarão-raposo, Tubarão-rato, Zorro-de-olhos-grandes.

References: mainland Portugal ([Antunes, 1970](#); [Fischer et al., 1981](#); [Muñoz-Chápuli, 1984](#); [Compagno, 1984a, 2001, 2016](#); [Quéro, 1984a](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [MNHN-IC-1989-0782, 1989](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Compagno, 1984a \(PECS\)\(?\)](#), [2001, 2016](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Simões, 1999](#); [Cardigos et al., 2005](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Martins, 2013](#); [Carneiro et al., 2014](#); [Fernandez-Carvalho et al., 2015](#); [Torres et al., 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#); [Fauconnet et al., 2019](#)) and Madeira ([Lowe, 1840a, 1841](#); [Springer, 1979a](#); [Cadenat](#)

and Blache 1981; Fischer *et al.*, 1981; Muñoz-Chápuli, 1984(?); Compagno, 1984a, 2001, 2016; Quéro, 1984a; Sanches, 1986; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Bischoff *et al.*, 2018; Froese and Pauly, 2019). The holotype (unique) of this species was caught off Madeira and its whereabouts are unknown. A second specimen is in the MMF collections (MMF 42302, 2012). It is a highly migratory species (Melendez and Macias, 2007).

***Alopias vulpinus* (Bonnaterre 1788)**

Common name: **Pelagic thresher or Bigeye thresher; Tubarão-raposo**, Albafar, Albafar-bravo, Alecrim, Anequim, Arrequim, Cavallo, Jogador-de-pau, Marracho, Peixe-alecrim, Peixe-cavalo, Peixe-raposo, Peixe-rato, Peixe-zorra, Peixe-zorro, Raposa, Raposo, Ratão, Rato, Rinquim, Roaz-bandeira, Roaz-de-bandeira, Romano, Romão, Tubarão, Tubarão-raposa, Tubarão-rato, Tubarão-zorra, Tubarão-zorro, Zorra, Zorro.

References: mainland Portugal▲ (Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, b, 1880; Almeida and Roquette, 1892; Vieira, 1893, 1897; Nobre, 1894, 1935; Bragança, 1904a; Seabra, 1911; Osório, 1917; Helling, 1940; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Springer, 1979a; Fischer *et al.*, 1981, 1987; Compagno, 1984a, 2001, 2016; Quéro, 1984a; Sanches, 1986; Santos *et al.*, 2002a, b, c; Melendez and Macias, 2007; Correia, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Ferreira, 1937a, 1939; Collins, 1954; Osório de Castro, 1967; Wheeler, 1978; Compagno, 1984a(?), 2001, 2016; Arruda, 1997; Santos *et al.*, 1997; Fowler *et al.*, 2005; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Torres *et al.*, 2016; Das and Afonso, 2017(?); Bischoff *et al.*, 2018; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1837, 1840a, 1841, 1849, 1852a, b; de Velloso, 1892; Belloc, 1934; Noronha and Sarmento, 1934; Nobre, 1935; Maul 1949d; Fowler, 1936a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cadenat and Blache 1981; Quéro 1984a; Sanches, 1986; Compagno, 2001, 2016; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018; Froese and Pauly, 2019). According to Das and Afonso (2017) the records of *A. vulpinus* in the Azores were quite probably a misidentification of *Alopias superciliosus* Lowe 1841.

Order Carcharhiniformes

Family Pentanchidae

***Apristurus laurussonii* (Saemundsson 1922)**

Common name: **Madeira catshark; Tubarão-gato-da-Madeira**.

References: mainland Portugal (?) (Ebert and Stehmann, 2013), Azores (Saldanha and Bischoff, 1997; Santos *et al.*, 1997; Desbruyères *et al.*, 2001; Melo and Menezes, 2002; Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Das and Afonso, 2017; Bischoff *et al.*, 2018) and Madeira (Cadenat & Maul, 1966; Springer, 1979b; Cadenat and Blache 1981; Compagno, 1984b, 2016; Quéro, 1984a; Sanches, 1986; Lloris *et al.*, 1991; Quéro *et al.*, 1993; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018; Froese and Pauly, 2019). The holotype (unique) of *Apristurus maderensis* Cadenat & Maul 1966 caught off Câmara de Lobos (Madeira), synonym of *A. laurussonii*, is in the MMF collections (MMF 18750, 1961). The distribution of this species in the Eastern Atlantic is discontinuous, from Greenland to Gabon. The distribution map reported by Ebert and Stehmann (2013) apparently includes the Gorringe Ridge, which belongs to the mainland Portugal EEZ. Thus, the occurrence of this species in mainland Portugal is questionable and it is not considered in the total number of valid species to this area.

***Apristurus manis* (Springer 1979)**

Common name: **Ghost catshark; Tubarão-gato-fantasma**.

References: Azores (Santos *et al.*, 1997; Melo and Menezes, 2002(?); IMAR/DOP COLETA 1687, 2004; IMAR/DOP SEDLOT-03-V04APR, 2004; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Das and Afonso, 2017(?)). Melo and Menezes (2002) and Das and Afonso (2017) mentioned that *Apristurus manis* is suspected to occur in the Azores.

***Apristurus microps* (Gilchrist 1922)**

Common name: **Smalleye catshark; Tubarão-gato-olhinho**.

References: Azores (Porteiro *et al.*, 2017).

Apristurus profundorum (Goode & Bean 1896)

Common name: **Deep-water catshark; Tubarão-gato-abissal**.

References: Azores (MAR-ECO 6251, 2004; MAR-ECO 8529, 2004; Bergstad *et al.*, 2008a (PECS); Carneiro *et al.*, 2014; Porteiro *et al.*, 2017).

Galeus atlanticus (Vaillant 1888)

Common name: **Atlantic sawtail catshark; Leitão-do-Atlântico**.

References: mainland Portugal (Rodrigues, 2009; Costa *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Moura *et al.*, 2015a, b; ICES, 2017a).

Galeus melastomus Rafinesque 1810

Common name: **Blackmouth catshark; Leitão**, Abarmão, Alfarica, Alitão, Boca-negra, Cação, Caçao-papoula, Cão-do-monte, Feiticeira, Gala, Leitão-do-mar, Litão, Papoila, Papoula, Peixe-gata, Peixelim, Pexelsim.

References: mainland Portugal▲ (Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Almeida and Roquette, 1892; Vieira, 1893, 1897; Nobre, 1894, 1903, 1935; Goode and Bean, 1895, 1896; Osório, 1896; Bragança, 1904a; Seabra, 1911; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Springer, 1979b; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; Martins, 1981; INIP, 1981a, 1982a, b; Compagno, 1984b, 2016; Dinis and Marecos, 1984; Quéro, 1984a; Dinis, 1985; Sanches, 1986; Martins *et al.*, 1992; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Erzini *et al.*, 2001b; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002a, b; Coelho *et al.*, 2003, 2005; Mendes *et al.*, 2004; Moura, 2004; Costa *et al.*, 2005, 2012; Fowler *et al.*, 2005; ICES, 2005; Sousa *et al.*, 2005, 2006; Costa, 2007; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Silva, 2008, 2010, 2011; Bordalo-Machado *et al.*, 2009; Correia, 2009; Rodrigues, 2009; Serena *et al.*, 2009a; Silva *et al.*, 2009; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Abella *et al.*, 2015; Moura *et al.*, 2015a, c, 2017b; Bañón *et al.*, 2016a; Gordon *et al.*, 2016; ICES, 2017a; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Dias, 1991; Arruda, 1997; Santos *et al.*, 1997; Hareide and Garnes, 2001 (PECS); Ebert and Stehmann, 2013(?); Carneiro *et al.*, 2014; Abella *et al.*, 2015; Compagno, 2016; Biscoito *et al.*, 2018) and Madeira (Lowe, 1837, 1843-1860; Günther, 1870; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d, 1976; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Cadenat and Blache 1981; Fischer *et al.*, 1981; INIP, 1982c; Sanches, 1986; Severino, 2004; Ferreira *et al.*, 2008; Pakhorukov, 2008 (PECS); Menezes *et al.*, 2009; Naylor *et al.*, 2012; Carneiro *et al.*, 2014; Abella *et al.*, 2015; Compagno, 2016; POGID, 2016; Biscoito *et al.*, 2018). Several specimens are in the MMF collections caught off at coast of mainland Portugal (MMF 22497, 1967, set of fifteen specimens).

Galeus murinus (Collett 1904)

Common name: **Mouse catshark; Leitão-da-Ilha**, Leitão-liso.

References: mainland Portugal (Ebert and Stehmann, 2013; Carneiro *et al.*, 2014) and Azores (Santos *et al.*, 1997; Menezes *et al.*, 2006, 2012; Fossen *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Barreiros and Gadig, 2011; Menezes and Giacomello, 2013; Carneiro *et al.*, 2014; Das and Afonso, 2017; ICES, 2017a).

Family Scyliorhinidae

Scyliorhinus canicula (Linnaeus 1758)

Common name: **Small-spotted catshark; Pata-roxa**, Bruxa, Cação, Caneja, Caneja-pata-roxa, Caneja-patarroxa, Cascarra, Feiticeira, Gato, Pata-rocha, Pata-roxa-pequena, Pataroxa, Patarraxa, Pata-roixa, Patarroixa, Patarroxa, Pinta-roixa, Pinta-roxa, Pintarroixa, Pintarroxa.

References: mainland Portugal▲ (Carvalho, 1765; Vandelli, 1771, 1797; Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Almeida and Roquette, 1892; Nobre, 1895a, 1903, 1904, 1935; Osório, 1896; Bragança, 1902, 1903, 1904a; Seabra, 1911; Murray and Hjort, 1912; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Springer, 1979b; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; Martins, 1981; INIP, 1981a, 1982a, b; Borges, 1984; Compagno, 1984b, 2016; Dinis and Marecos, 1984; Quéro, 1984a; Dinis, 1985; Sanches, 1986; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Erzini *et al.*, 2001a, b; Gomes *et al.*,

2001; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Santos *et al.*, 2002a, b; Coelho *et al.*, 2003, 2005; Mendes *et al.*, 2004; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Stergiou *et al.*, 2006; Costa, 2007; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Silva, 2008, 2010, 2011; Batista *et al.*, 2009; Rodrigues, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Moura *et al.*, 2015a, d, 2017a; Gordon *et al.*, 2016; Ryland and Lancaster, 2017; ICES, 2017a; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Arruda, 1997; Santos *et al.*, 1997; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013(?); Carneiro *et al.*, 2014; Compagno, 2016; Bischoff *et al.*, 2018) and Madeira (Lichtenstein, 1844; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Ribeiro, 2008; Carneiro *et al.*, 2014; Compagno, 2016; Bischoff *et al.*, 2018(?)). According to Maul (1976) it does not seem to live in Madeiran waters and Bischoff *et al.* (2018) refer that the presence of this species is considered as subject to confirmation.

***Scyliorhinus stellaris* (Linnaeus 1758)**

Common name: **Polkadot catshark; Pata-roxa-gata**, Bruxa, Cação, Carraça, Carrica, Cascarra, Feiticeira, Gata, Pata-roixa, Pata-roxa, Pata-roxa-denise; Pata-roxa-grande, Pataroxa, Patarôxa, Patarroixa, Patarroxa, Peixelim, Pexelim, Pinta-roixa, Pinta-roxa, Pintarroixa, Pintarroxa.

References: mainland Portugal▲ (Vandelli, 1771, 1797; Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Baldaque da Silva, 1891; Vieira, 1893, 1897; Nobre, 1894, 1935; Bragança, 1904a; Seabra, 1911; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Compagno, 1984b, 2016; Dinis and Marecos, 1984; Quéro, 1984a; Fischer *et al.*, 1981, 1987; INIP, 1982b; Sanches, 1986; Saldanha *et al.*, 1995; Melendez and Macias, 2007; Henriques *et al.*, 2008; Rodrigues, 2009; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Ellis *et al.*, 2015a; Gordon *et al.*, 2016; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Sampaio, 1904; Ribeiro, 1936; Arruda, 1997; Carneiro *et al.*, 2014) and Madeira (Osório de Castro, 1954, 1967; Ellis *et al.*, 2015a). The occurrence of this species in the Azores was firstly reported by Sampaio (1904), but Santos *et al.* (1997) questioned the identification of this specimen as belonging to the species *Scyllium catulus* Müller & Henle 1838 and considered that it could be *Dalatias licha* (Bonnaterre 1788). The presence in the Azores region is not yet confirmed and thus it is not considered in the total number of valid species in this area. The map of distribution of this species reported by Ellis *et al.* (2015a) included the area of Selvagens Islands (Madeira EEZ).

Family Pseudotriakidae

***Pseudotriakis microdon* de Brito Capello 1868**

Common name: **False catshark; Tubarão-mona**, Carago, Mamona, Mamôna, Mona, Peixe-carago, Rinquim, Tubarão.

References: mainland Portugal (de Brito Capello, 1867a, b, c, 1869a, 1880; Goode and Bean, 1895, 1896; Bragança, 1904a; Seabra, 1911; Nobre, 1935; Fowler, 1936a; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Compagno, 1984b, 2016; Quéro, 1984a; Sanches, 1986; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Kyne *et al.*, 2015; Roberts *et al.*, 2017; Froese and Pauly, 2019), Azores (Albert I^{er} de Mónaco, 1904; Fowler, 1936a; Ferreira, 1939; Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1967; Springer, 1979b; Fischer *et al.*, 1981; Compagno, 1984b, 2016; Quéro, 1984a; Sanches, 1986; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Menezes *et al.*, 2009; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Kyne *et al.*, 2015; Froese and Pauly, 2019; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018) and Madeira (NHMUK 1921.6.8.1, 1912-1936; Noronha 1926a; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Maul 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cadenat and Blache 1981; Fischer *et al.*, 1981; Compagno, 1984b, 2016; Quéro, 1984a; Sanches, 1986; Lloris *et al.*, 1991; Menezes *et al.*, 2009; Carneiro *et al.*, 2014; Kyne *et al.*, 2015; Bischoff *et al.*, 2018; Froese and Pauly, 2019). The specimen of *P. microdon* caught off Setúbal described by de Brito Capello in 1868 was preserved in the Museu Bocage and destroyed in the 1978 fire. A second specimen of this species is in the MMF collections (MMF 3270, 1951). Maul considers it as a rare species in Madeira.

Family Triakidae

Galeorhinus galeus (Linnaeus 1758)

Common name: **Tope shark; Perna-de-moça**, Bico-doce, Bico-doce-do-baixo, Boca-doce, Boca-doce-do-baixo, Cação, Caneja, Cascaria, Cascarra, Chanão, Charrão, Chião, Chona, Chonão, Choua, Chouão, Dentudo, Faguetão, Faquetão, Faticeira, Feiticeira, Foguetão, Gata, Lixa, Marracho, Peixe-gata, Peixelim, Peixe-ruim, Pexelim, Rinquim, Ruim, Tolho, Tubarão, Tubarão-da-sopa, Tubarão-perna-de-moça.

References: mainland Portugal▲ ([Vandelli, 1771, 1797](#); [Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a](#); [Gervais and Boulart, 1877a](#); [Nobre, 1894, 1904, 1935](#); [Vieira, 1897](#); [Bragança, 1904a](#); [Richard, 1904, 1905a, 1908](#); [Seabra, 1911](#); [Roule, 1912](#); [Fowler, 1936a](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Compagno, 1979, 1984b, 2016](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1982a](#); [Martins, 1981](#); [Branstetter, 1984a](#); [Muñoz-Chápuli, 1984](#); [Dinis, 1985](#); [Sanches, 1986](#); [Erzini et al., 2001a](#); [Coelho et al., 2005](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Melendez and Macias, 2007](#); [Rodrigues, 2009](#); [Chaves and Silva, 2013](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Moura et al., 2015b](#); [Ryland and Lancaster, 2017](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores▲ ([Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Richard, 1913](#); [Roule, 1919a](#); [Guiart, 1935](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Dias et al., 1976, 1990](#); [Martins, 1982a](#); [INIP, 1985a, b, c](#); [Dias and Monteiro, 1986](#); [Sanches, 1986](#); [Patzner et al., 1990, 1992](#); [Dias, 1991](#); [Arruda, 1992, 1997](#); [Krug, 1995](#); [Silva et al., 1994, 1995](#); [Silva and Menezes, 1995a, b](#); [Isidro, 1996](#); [Menezes, 1996](#); [Santos et al., 1997](#); [Morato-Gomes et al., 1998](#); [Cardigos et al., 2005](#); [Fowler et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Pham et al., 2013](#); [Carneiro et al., 2014](#); [Torres et al., 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#); [Almeida and Biscoito, 2019](#); [Fauconnet et al., 2019](#)) and Madeira▲ ([Lowe, 1837](#); [Belloc, 1934](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Compagno, 1979, 2016](#); [INIP, 1980, 1982c](#); [Cadenat and Blache 1981](#); [Branstetter, 1984a](#); [Muñoz-Chápuli, 1984](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Severino, 2004](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [POGID, 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)).

Mustelus asterias Cloquet 1819

Common name: **Starry smooth-hound; Cação-pintado**, Alitão, Alvarinho, Álvaro, Cação, Cação-pique, Caneja, Caneja-pintada, Corre, Corre-costa, Corre-costas, Esgalhudo, Feiticeira, Galhudo, Lixa.

References: mainland Portugal▲ ([Vandelli, 1797](#); [Barboza du Bocage and de Brito Capello, 1866](#); [Gervais and Boulart, 1877a](#); [Nobre, 1903](#); [Murray and Hjort, 1912](#); [Osório de Castro, 1967](#); [Compagno, 1979, 1984b](#); [Fischer et al., 1981, 1987](#); [Branstetter, 1984a](#); [Sanches, 1986](#); [Melendez and Macias, 2007](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Osório de Castro, 1967\(?\)](#); [Compagno, 2016\(?\)](#)) and Madeira ([Fowler, 1936a\(?\)](#); [Osório de Castro, 1967](#); [Compagno, 1979](#); [Branstetter, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Biscoito et al., 2018\(?\)](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). Osório de Castro (1967) and Compagno (2016) doubt the occurrence of this species in the Azores. The presence of this species in the Azores is not yet confirmed and thus it is not considered in the total number of valid species in this area.

Mustelus canis (Mitchill 1815)

Common name: **Dusky smooth-hound; Cação-cachorro**, Alitão, Cação, Cação-morraceiro, Caneja, Corre-costa, Corre-costas, Feiticeira, Picua, Pique, Tubarão-de-barbatanas-pretas.

References: mainland Portugal (?) ([de Brito Capello, 1880](#); [Rochebrune 1882](#); [Almeida and Roquette, 1892](#); [Bragança, 1904a](#); [Seabra, 1911](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#), [Osório de Castro, 1954](#)) and Madeira (?) ([Lowe, 1837](#); [Günther, 1870](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#)). Several authors indicate the occurrence of this species in Portuguese mainland and Madeira waters with different scientific names (*Mustelus laevis* (Rondel) or *Mustelus laevis* Risso). However, *Mustelus laevis* is currently synonyms of *Mustelus mustelus* (Linnaeus 1758). *M. canis* is allopatric with *M. mustelus* and sympatric with *M. norrisi* (Compagno, 1984b). The current geographic distribution of *M. canis* is restricted to the western Atlantic (Compagno 1984b; Eschmeyer et al., 2019; Froese and Pauly, 2019). According to Ebert and Stehmann (2013) this species has been sometimes

confused with *Mustelus mustelus* (Linnaeus 1758). Such as reported by Biscoito *et al.* (2018) this species should be excluded from the list of chondrichthyans of Madeira. Thus, this species is not considered in the total number of valid species.

***Mustelus mustelus* (Linnaeus 1758)**

Common name: **Dusky smooth-hound; Cação-liso**, Alitão, Cação, Cação-branco, Cação-morraceiro, Cação-morraqueiro, Cação-pique, Caneja, Caneja-lisa, Corre-costa, Corre-costas, Feiticeira, Galhudo, Picua, Pique, Tubarão-de-barbatanas-pretas.

References: mainland Portugal[▲] ([Carvalho, 1765\(?\)](#); [Vandelli, 1771](#); [Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, b, 1880](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1893, 1897, 1899](#); [Nobre, 1894, 1903, 1935](#); [Osório, 1896](#); [Bragança, 1904a](#); [Seabra, 1911](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Compagno, 1979, 1984b, 2016](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [Branstetter, 1984a](#); [Sanches, 1986](#); [Martins et al., 1992](#); [Erzini et al., 2001b](#); [Santos et al., 2002b](#); [Coelho et al., 2005](#); [ICES, 2005](#); [Melendez and Macias, 2007](#); [Henriques et al., 2008](#); [Batista et al., 2009](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Belloc, 1934](#); [Osório de Castro, 1967\(?\)](#); [Branstetter 1984a\(?\)](#); [Santos et al., 1997 \(?\)](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014\(?\)](#); [Compagno, 2016\(?\)](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019\(?\)](#)) and Madeira[▲] ([Lowe, 1837](#); [Vinciguerra, 1883](#); [de Velloso, 1892](#); [Roule, 1919a](#); [Belloc, 1934](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967\(?\)](#); [Compagno, 1979, 1984b, 2016](#); [Cadenat and Blache 1981](#); [Fischer et al., 1981](#); [Branstetter, 1984a](#); [INIP, 1980, 1982c, 1984b](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Compagno, et al., 2005 \(?\)](#); [Melendez and Macias, 2007](#); [Ferreira et al., 2008](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Compagno, 2016](#); [POGID, 2016](#); [Biscoito et al., 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019\(?\)](#)). As mentioned by Santos *et al.* (1997) there is only one known record of this species for the Azores based on Belloc (1934). Thus, the occurrence of this species in the Azores region needs further documentation.

***Mustelus punctulatus* Risso 1827**

Common name: **Blackspotted smooth-hound; Cação-ponteado**, Cação, Caneja.

References: mainland Portugal ([Branstetter, 1984a](#); [Compagno, 1984b](#); [Rodrigues, 2009](#); [Serena et al., 2009](#); [Ramos, 2010](#); [Carneiro et al., 2014](#); [Dulvy et al., 2015](#)) and Madeira ([Cadenat and Blache 1981](#); [Branstetter, 1984a](#); [Wirtz et al., 2008\(?\)](#); [Serena et al., 2009b](#); [Dulvy et al., 2015](#); [Compagno, 2016](#); [Froese and Pauly, 2019\(?\)](#)). According to Eschmeyer *et al.* (2019) *Mustelus mediterraneus* Quignard & Capapé 1827 is synonym of *M. punctulatus*. Cadenat and Blache (1981) reported a specimen caught off from the coast of Madeira and other authors include the area of Selvagens Islands in the distribution maps of this species ([Branstetter, 1984a](#); [Serena et al., 2009b](#); [Dulvy et al., 2015](#); [Compagno, 2016](#)). On the other hand, Wirtz *et al.* (2008), Biscoito *et al.* (2018) and Froese and Pauly (2019) considered doubtful the presence of this species in the Madeira Archipelago. Thus, the distribution area of this species should be re-evaluated because of its confusion with *Mustelus mustelus* (Linnaeus 1758).

Family Carcharhinidae

***Carcharhinus altimus* (Springer 1950)**

Common name: **Bignose shark; Tubarão-baboso**, Marracho, Marracho-baboso.

References: Madeira (?) ([Compagno, 2016\(?\)](#)). *C. altimus* is a circumglobal and highly migratory species in tropical and warm temperate seas (including Mediterranean Sea, Red Sea) ([Mancini et al., 2015](#); [Eschmeyer et al., 2019](#)). Compagno (2016) mentions as doubtful the occurrence of this species in Madeira. Specimens are needed to confirm its presence in Madeira. Thus, this species is not considered in the total number of valid species of Madeira.

***Carcharhinus brachyurus* (Günther 1870)**

Common name: **Copper shark; Tubarão-cobre**.

References: mainland Portugal ([Muñoz-Chápuli, 1984\(?\)](#); [IGFA 864-1988, 1987](#); [Carneiro et al., 2014](#); [Morey et al., 2015](#)) and Madeira (?) ([Muñoz-Chápuli, 1984\(?\)](#); [Naylor et al., 2012\(?\)](#); [Morey et al.,](#)

2015(?); Compagno, 2016(?)). According to Compagno (1984b) *C. brachyurus* is a tropical and subtropical species with a world-wide distribution, which occurs from inshore to offshore waters and from the surface to a depth of, at least, 100 m. This species is sometimes confused with other large *Carcharhinus* species, particularly with *C. obscurus* (Lesueur 1818) (Morey et al., 2015 and Compagno, 2016) and the scarce knowledge of this species in the Madeira area has not permitted a clear decision of its presence. However, according to Compagno (2016), the occurrence of *C. brachyurus* in the Madeira Archipelago is questionable and its presence in the Madeira EEZ needs further documentation. Thus, this species is not considered in the total number of valid species of Madeira.

Carcharhinus brevipinna (Valenciennes 1839)

Common name: **Graceful shark; Tubarão-tecelão**, Faqueita, Faqueta, Faquete, Jaqueta, Marracho, Tubarão.

References: mainland Portugal (Branstetter, 1984b; Quéro et al., 2003; Ebert and Stehmann, 2013(?); Carneiro et al., 2014; ICES, 2017a), Azores (?) (Branstetter, 1984b(?); Arruda, 1997(?); Carneiro et al., 2014(?)) and Madeira (pers. comm. Reiner: a specimen landed at Machico harbor, Aug. 16, 1982). This species is sometimes confused with other *Carcharhinus* species, particularly with *C. limbatus* (Valenciennes 1839) and possibly with *C. isodon* (Valenciennes 1839) (Compagno, 2016). Quéro et al. (2003) indicated the presence of the species off Portugal, without referring the geographical area of occurrence and Ebert and Stehmann (2013) reported possibly southern Portugal. Arruda (1997), Bristetter (1984b) and Carneiro et al. (2014) consider doubtful the occurrence of this species in the Azores, which needs further documentation and it is not considered in the total number of valid species of Azores.

Carcharhinus falciformis (Bibron 1839)

Common name: **Silky shark; Tubarão-luzidio**, Faqueira, Faqueta, Jaqueta, Marracho, Marracho-luzidio, Tubarão.

References: mainland Portugal (Fischer et al., 1987; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro et al., 2014; Compagno, 2016; ICES, 2017a), Azores (Arruda, 1997(?); Bristetter, 1984b(?); Ebert and Stehmann, 2013(?); Carneiro et al., 2014; Compagno, 2016; ICES, 2017a; Biscoito et al., 2018) and Madeira (Lowe, 1839, 1840b, 1843, 1844a, 1849; NHMUK 1851.1.9.14, 1850-1853; Noronha and Sarmento, 1934; Nobre, 1935; Fowler, 1936a; Compagno, 1979, 1984b, 2016; Fischer et al., 1981; Sanches, 1986; Lloris et al., 1991; Melendez and Macias, 2007; Wirtz et al., 2008(?); Ebert and Stehmann, 2013; Carneiro et al., 2014; Biscoito et al., 2018; Froese and Pauly, 2019(?)). This species is sometimes confused with *C. obscurus* (Lesueur 1818) (Compagno, 2016). The current geographic distribution of this species is circumglobal in tropical seas (including Caribbean Sea, Red Sea, Persian Gulf) (Eschmeyer et al., 2019). Ebert and Stehmann (2013) indicated its presence off Madeira, based on its known geographic distribution. Arruda (1997) and Bristetter (1984b) considered the presence of this species off the Azores as doubtful and Ebert and Stehmann (2013) considered it possible.

Carcharhinus galapagensis (Snodgrass & Heller 1905)

Common name: **Galapagos shark; Tubarão-dos-Galápagos**.

References: mainland Portugal (Compagno, 1984b(?), 2016(?); Bennett et al., 2003), Azores (Bruun and Azevedo, 1995; Santos et al., 1997; Bennett et al., 2003; Porteiro et al., 2010; Barreiros and Gadig, 2011; Afonso et al., 2013; Carneiro et al., 2014; Compagno, 2016(?); Das and Afonso, 2017; Biscoito et al., 2018) and Madeira (Compagno, 1984b, 2016; Wirtz et al., 2008; Carneiro et al., 2014; Biscoito et al., 2018; Froese and Pauly, 2019). This species is sometimes confused with *C. obscurus* (Lesueur 1818) (Compagno, 2016). The current geographic distribution of this species is worldwide in tropical waters, especially around offshore islands (Eschmeyer et al., 2019). According to Compagno (2016) the occurrence of this species in the Eastern Atlantic is off the Azores and off the coasts of Spain or Portugal, is possible. For mainland Portugal the presence of this species needs further documentation. One specimen is in the MMF collections (MMF 023409, 1940).

Carcharhinus isodon (Valenciennes 1839)

Common name: **Finetooth shark; Tubarão-dente-liso**.

References: Madeira (Branstetter, 1984b(?); Lloris et al., 1991). The current geographic distribution of this species in the Eastern Atlantic is Senegal and Guinea-Bissau (Eschmeyer et al., 2019) and Angola

(INIP, 1985d). This species is sometimes confused with *C. brevipinna* (Valenciennes 1839) but possibly also with *C. limbatus* (Valenciennes 1839)(Compagno, 2016).

***Carcharhinus leucas* (Valenciennes 1839)**

Common name: **Pigeye shark; Tubarão-buldogue**, Bico-doce, Boca-doce, Faqueita, Faqueta, Jaqueta, Marracho, Melracho, Merracho, Olho-branco, Olhudo-branco, Perna-de-moça, Rinquim, Sarda, Tubarão, Tubarão-grande, Tubarão-sarda.

References: mainland Portugal (?) ([Osório de Castro, 1954, 1967\(?\)](#)), Azores ([Osório de Castro, 1967](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Simpfendorfer and Burgess, 2009](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Compagno, 2016\(?\)](#)) and Madeira (?) ([Osório de Castro, 1967](#)). This species is sometimes confused with two other species of the genus *Carcharhinus*, namely *C. amboinensis* (Müller & Henle 1839) and *C. carcharias* (Linnaeus 1758) (Compagno, 2016). According to Eschmeyer et al. (2019) *C. leucas* is a circumglobal species in tropical seas. The only possible presence of this species in the mainland Portugal is reported by Osório de Castro (1967). Since then no other author described its occurrence in mainland Portugal. Compagno (2016) reports that this species may occur off the Azores (although the records may be based on *C. obscurus* or *C. galapagensis*). The only presence of *C. leucas* in the fishing grounds of Madeira was reported by Osório de Castro (1967) without indicating the geographic location. However, in 2000 it was indicated the presence of this species in the Canary Islands (human observation). Since no specimens or other reference for Madeira were found, the presence of this species is considered as questionable and subject to confirmation. Thus, this species is not considered in the total number of valid species to mainland Portugal and Madeira.

***Carcharhinus limbatus* (Valenciennes 1839)**

Common name: **Blacktip reef shark; Tubarão-de-pontas-negras**, Anequim, Faqueita, Faqueta, Faquete, Jaqueta, Marracho, Marracho-de-pontas-negras, Tubarão, Tubarão-pardo.

References: mainland Portugal ([Branstetter, 1984b](#); [Fischer et al., 1987](#); [Ebert and Stehmann, 2013](#); [Burgess and Branstetter, 2015](#); [Carneiro et al., 2014](#); [ICES, 2017a](#)), Azores (?) ([Branstetter, 1984b\(?\)](#); [Arruda, 1997\(?\)](#); [Carneiro et al., 2014\(?\)](#); [Biscoito et al., 2018](#)) and Madeira ([Lowe, 1839, 1840a, 1841, 1843, 1849](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Compagno, 1979, 1984b; 2016](#); [Fischer et al., 1981](#); [Branstetter, 1984b](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Melendez and Macias, 2007](#); [Wirtz et al., 2008\(?\)](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Burgess and Branstetter, 2015](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019\(?\)](#)). This species is sometimes confused with *C. brevipinna* (Müller & Henle 1839) (Compagno, 2016). Branstetter (1984b), Arruda (1997) and Carneiro et al. (2014) consider doubtful the occurrence of this species in the Azores and further documentation is needed. Thus, this species is not considered in the total number of valid species of the Azores. The holotype (unique) of *Carcharias microps* Lowe 1841 was caught off Madeira, synonym of *C. limbatus*, and its whereabouts are unknown. A second specimen is in the MMF collections (MMF 39543, 2008).

***Carcharhinus longimanus* (Poey 1861)**

Common name: **Oceanic whitetip shark; Tubarão-de-pontas-brancas**, Faqueta, Faquete, Jaqueta, Marracho, Marracho-de-pontas-brancas, Tubarão.

References: mainland Portugal ([Compagno, 1979\(?\)](#), [1984b, 2016](#); [Fischer et al., 1981](#); [Branstetter, 1984b](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Walls et al., 2015c](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores ([Ferreira, 1939](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Fischer et al., 1981](#); [Branstetter, 1984b](#); [Compagno, 1984b, 2016](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017\(?\)](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#)) and Madeira ([Maul, 1955a](#); [Osório de Castro, 1967](#); [Compagno, 1979, 1984b, 2016](#); [Cadenat and Blache 1981](#); [Fischer et al., 1981](#); [Branstetter, 1984b](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Walls et al., 2015c](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). Two specimens are in the MMF collections (collected in 1938 and 1953).

Carcharhinus obscurus (Lesueur 1818)

Common name: **Dusky shark; Tubarão-faqueta**, Faqueita, Faqueta, Faquete, Faquetão, Jaqueta, Marracho, Marracho-faqueta, Meiracho, Merracho, Peixe-romã, Rinquim, Romã, Tá-rá-tá-tá, Toiro, Touro, Tubarão, Tubarão-branco, Tubarão-toiro, Tubarão-touro.

References: mainland Portugal ([Osório de Castro, 1967](#); [Compagno, 1979, 1984b\(?\)](#), 2016; [Cadenat and Blache 1981](#); [Fischer et al., 1981, 1987](#); [Branstetter, 1984b](#); [Sanches, 1986](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013 \(?\)](#); [Carneiro et al., 2014](#); [ICES, 2017a](#)), Azores (?) ([Arruda, 1997](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014\(?\)](#); [Biscoito et al., 2018](#)) and Madeira ([Günther, 1870](#); [Belloc, 1934](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Maul 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Compagno, 1979, 1984b\(?\)](#), 2016; [Cadenat and Blache 1981](#); [Fischer et al., 1981](#); [Branstetter, 1984b](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Melendez and Macias, 2007\(?\)](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)).

The current geographic distribution of this species is nearly worldwide, tropical and subtropical seas (including Caribbean Sea, Red Sea) ([Eschmeyer et al., 2019](#)). According to Compagno (2016) and Melendez and Macias (2007), in Madeira some records of *C. obscurus* may be based on misidentification of *Carcharhinus galapagensis* (Snodgrass & Heller 1905). Furthermore, records of this species in the Azores may also be of the Galapagos shark (*C. galapagensis*), assuming that this species is distinct from *C. obscurus*. However, Ebert and Stehmann (2013), based on molecular studies, suggested that these two species may in fact be the same. The occurrence of this species in the Azores needs further documentation. Thus, this species is not considered in the total number of valid species of the Azores.

Carcharhinus plumbeus (Nardo 1827)

Common name: **Sandbar shark; Tubarão-corre-costa**, Faqueita, Faqueta, Faquete, Jaqueta, Lobo, Marracho, Marracho-branco, Melgacho, Tubarão, Tubarão-azul, Tubarão-lobo.

References: mainland Portugal▲ ([Osório de Castro, 1954, 1967](#); [Compagno, 1979, 1984b](#), 2016; [Cadenat and Blache 1981](#); [Fischer et al., 1981, 1987](#); [Branstetter, 1984b](#); [Sanches, 1986](#); [Melendez and Macias, 2007](#); [Henriques et al., 2008](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Ferretti et al., 2015a](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores ([Branstetter, 1984b\(?\)](#); [Sanches, 1986\(?\)](#); [Arruda, 1997\(?\)](#); [Carneiro et al., 2014](#); [ICES, 2017a](#); [Biscoito et al., 2018](#)) and Madeira ([Compagno, 1979, 2016](#); [Fischer et al., 1981](#); [Branstetter, 1984b\(?\)](#); [Sanches, 1986\(?\)](#); [Lloris et al., 1991](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014\(?\)](#); [Ferretti et al., 2015a](#); [Biscoito et al., 2018\(?\)](#)). According to Compagno (2016) this species is sometimes confused with four other species of the genus *Carcharhinus*, namely *C. altimus* (Springer 1950), *C. galapagensis* (Snodgrass & Heller 1905), *C. obscurus* (Lesueur 1818) and *C. leucas* (Valenciennes 1839). Specimens are needed for confirmation of the presence of this species in Madeira.

Carcharhinus signatus (Poey 1868)

Common name: **Night shark; Tubarão-da-noite**.

References: mainland Portugal ([ICES, 2017a](#)) and Madeira (?) ([Branstetter, 1984b\(?\)](#)). This species is sometimes confused with *C. falciformis* (Bibron 1839) (Compagno, 2016). The current geographic distribution of this species in the Eastern Atlantic is from Senegal to northern Namibia (Ebert and Stehmann, 2013, Compagno, 2016, Eschmeyer et al., 2019). The species *C. signatus* may occur in mainland Portugal as occasional vagrants (ICES, 2017a), but this requires confirmation. According to Branstetter (1984b) the occurrence of *C. signatus* in the Madeira Archipelago is questionable. Thus, this species is not considered in the total number of valid species of Madeira.

Galeocerdo cuvier (Péron & Lesueur 1822)

Common name: **Tiger shark; Tubarão-tigre**, Tigre, Tubarão.

References: mainland Portugal ([Compagno, 1979](#); [Branstetter, 1984b\(?\)](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores ([Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Simpfendorfer and Soldo, 2015](#); [Compagno, 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#)) and Madeira ([Fischer et al., 1981](#); [Lloris et al., 1991](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Simpfendorfer and Soldo, 2015](#); [Compagno, 2016\(?\)](#); [Biscoito et al., 2018\(?\)](#)). The distribution maps of this species reported by Fischer et al. (1981), Simpfendorfer and Soldo (2015) and Compagno (2016) included the distribution range of this species in the Selvagens

Islands (Madeira EEZ). Specimens are needed for confirmation of the presence of this species in Madeira (Biscoito *et al.*, 2018).

***Negaprion brevirostris* (Poey 1868)**

Common name: **Lemon shark; Tubarão-limão**, Cação-limão, Limão, Tubarão.

References: Madeira ([Branstetter, 1984b\(?\)](#); [Lloris *et al.*, 1991](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is the Atlantic and Eastern Pacific oceans. The only reported presence of *N. brevirostris* in Madeira was indicated by Lloris *et al.* (1991). Since then no other author described the presence of this species and its occurrence in this EEZ was not yet confirmed by more recent data. Specimens are needed to confirm its presence in mainland Portugal and Madeira.

***Prionace glauca* (Linnaeus 1758)**

Common name: **Blue shark; Tintureira**, Agulha, Bico-doce, Boca-doce, Caelha, Guelha, Luzidia, Peixe-agulha, Quelha, Rinquim, Tigre-azul, Tintureiro, Tubarão, Tubarão-azul, Valentina, Valetina, Veletina.

References: mainland Portugal[▲] ([Carvalho, 1765](#); [Vandelli, 1771, 1797](#); [Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, 1880](#); [Gervais and Boulart, 1877a](#); [Posser, 1880](#); [Almeida and Roquette, 1892](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Osório, 1896](#); [Vieira, 1897](#); [Bragança, 1902, 1903, 1904a](#); [Richard, 1904](#); [Seabra, 1911](#); [Roule, 1919a](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Compagno, 1979, 1984b, 2016](#); [Cadenat and Blache 1981](#); [Fischer *et al.*, 1981](#); [Branstetter, 1984b](#); [Muñoz-Chápuli, 1984](#); [Sanches, 1986](#); [Erzini *et al.*, 2001a, b](#); [Santos *et al.*, 2002b](#); [Coelho *et al.*, 2005](#); [Fowler *et al.*, 2005](#); [Costa, 2007](#); [Melendez and Macias, 2007](#); [Bordalo-Machado *et al.*, 2009](#); [Bordalo-Machado and Figueiredo, 2009](#); [Correia, 2009](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro *et al.*, 2014](#); [Vandeperre *et al.*, 2014](#); [Moura *et al.*, 2015b](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Roule, 1919a](#); [Belloc, 1934](#); [Fowler, 1936a](#); [Ferreira, 1937a, 1939](#); [Bigelow and Schroeder, 1948](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Parin, 1970](#); [Wheeler, 1978](#); [Fischer *et al.*, 1981](#); [Martins, 1982a](#); [Compagno, 1984b, 2016](#); [Branstetter, 1984b](#); [Martin 1986](#); [Sanches, 1986](#); [Lloris *et al.*, 1991](#); [Arruda, 1992, 1997](#); [Pinho and Pereira, 1995a](#); [Silva and Menezes, 1995b](#); [Santos *et al.*, 1995b, 1997](#); [Silva *et al.*, 1995](#); [Menezes, 1996](#); [Cardigos *et al.*, 2005](#); [Menezes *et al.*, 2006](#); [Correia, 2009](#); [Porteiro *et al.*, 2010](#); [Barreiros and Gadig, 2011](#); [Besugo, 2013](#); [Ebert and Stehmann, 2013](#); [Martins, 2013](#); [Carneiro *et al.*, 2014](#); [Vandeperre *et al.*, 2014](#); [Torres *et al.*, 2016](#); [Das and Afonso, 2017](#); [ICES, 2017a](#); [Biscoito *et al.*, 2018](#); [Fauconnet *et al.*, 2019](#)) and Madeira[▲] ([Lowe, 1837](#); [Belloc, 1934](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Compagno, 1979, 1984b, 2016](#); [Cadenat and Blache 1981](#); [Fischer *et al.*, 1981](#); [Branstetter, 1984b](#); [Muñoz-Chápuli, 1984](#); [Sanches, 1986](#); [Lloris *et al.*, 1991](#); [Severino, 2004](#); [Wirtz *et al.*, 2008](#); [Correia, 2009](#); [EMEPC, 2010](#); [Ebert and Stehmann, 2013](#); [Carneiro *et al.*, 2014](#); [Vandeperre *et al.*, 2014](#); [Almada *et al.*, 2015](#); [POGID, 2016](#); [POGIS, 2016](#); [Biscoito *et al.*, 2017, 2018](#); [Froese and Pauly, 2019](#)). One specimen caught off at the Portuguese continental slope in September 2009 is in the MB collections.

***Rhizoprionodon acutus* (Rüppell 1837)**

Common name: **Milk shark; Tubarão-bicudo**, Cação, Leopard, Sarda, Tubarão, Tubarão-leopardo.

References: Madeira ([Compagno, 1979, 1984b, 2016](#); [Cadenat and Blache 1981](#); [Fischer *et al.*, 1981](#); [Branstetter, 1984b\(?\)](#); [Sanches, 1986](#); [Lloris *et al.*, 1991](#); [Melendez and Macias, 2007](#); [Wirtz *et al.*, 2008\(?\)](#); [Carneiro *et al.*, 2014\(?\)](#); [Biscoito *et al.*, 2018\(?\)](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Rhizoprionodon terraenovae* (Richardson 1837) (Compagno, 2016). According to Eschmeyer *et al.* (2019) *R. acutus* is a circumglobal species in tropical through warm temperate seas. Biscoito *et al.* (2018) considered that the presence of this species is subject to confirmation.

Family Sphyrnidae

***Sphyraena lewini* (Griffith & Smith 1834)**

Common name: **Scalloped hammerhead; Tubarão-martelo-recortado**, Corno, Cornuda, Cornudo, Martelo, Peixe-corno, Peixe-cornudo, Peixe-martelo, Tubarão, Tubarão-martelo, Tubarão-mona.

References: mainland Portugal ([Quéro, 1984b](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro *et al.*, 2014](#); [Ferretti *et al.*, 2015b](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores ([Quéro, 1984b](#); [Arruda, 2007](#)).

1997(?); Baum *et al.*, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Ferretti *et al.*, 2015b; Porteiro *et al.*, 2017(?); Bischoff *et al.*, 2018; Froese and Pauly, 2019) and Madeira (Quéro, 1984b; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Ferretti *et al.*, 2015b; Bischoff *et al.*, 2018(?); Froese and Pauly, 2019). According to Compagno (2016) this species is sometimes confused with two other species of the genus *Sphyrna*, namely *S. mokarran* (Rüppell 1837) and *S. zygaena* (Linnaeus 1758). Santos *et al.* (1997) don't consider valid the record of this species in the Azores. The distribution map of this species reported by Ferretti *et al.* (2015b) included the area of Selvagens Islands (Madeira EEZ).

Sphyrna mokarran (Rüppell 1837)

Common name: **Great hammerhead; Tubarão-martelo-gigante**, Cândar, Corno, Cornuda, Cornuda-gigante, Cornudo, Peixe-corno, Peixe-cornudo, Peixe-martelo, Tubarão, Tubarão-martelo.

References: mainland Portugal (Denham *et al.*, 2007; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Asber *et al.*, 2015) and Madeira (Fischer *et al.*, 1981; Denham *et al.*, 2007; Ebert and Stehmann, 2013). According to Compagno (2016) this species is sometimes confused with three other species of the genus *Sphyrna*, namely *S. lewini* (Griffith & Smith 1834), *S. zygaena* (Linnaeus 1758) and *S. tudes* (Valenciennes 1822). The possible distribution of this species shown in the maps reported by Fischer *et al.* (1981), Denham *et al.* (2007) and Ebert and Stehmann (2013) partially included the area of Madeira and Selvagens islands, respectively.

Sphyrna zygaena (Linnaeus 1758)

Common name: **Smooth hammerhead; Tubarão-martelo**, Cândar, Corno, Cornuda, Cornudo, Esparto, Martelo, Peixe-corno, Peixe-cornuda, Peixe-cornudo, Peixe-martelo, Peixelim, Pexelim, Rinquim, Tubarão.

References: mainland Portugal (Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Almeida and Roquette, 1892; Vieira, 1897; Bragança, 1904a; Seabra, 1911; Nobre, 1935; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Gilbert 1979; Fischer *et al.*, 1981, 1987; Compagno, 1984b; Muñoz-Chápuli, 1984; Quéro, 1984b; Sanches, 1986; Santos *et al.*, 2002a; Melendez and Macias, 2007; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; ICES, 2017a; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Sampaio, 1904; Roule, 1919a; Fowler, 1936b; Ferreira, 1939; Bigelow and Schroeder, 1948; Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1967; Wheeler, 1978; Sanches, 1986; Arruda, 1992, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Carneiro *et al.*, 2014; Compagno, 2016 (PECS); Torres *et al.*, 2016; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira[▲] (Bowdich, 1825; Lowe, 1837, 1843-1860; Günther, 1870; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muñoz-Chápuli, 1984; Sanches, 1986; Severino, 2004; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Compagno, 2016; POGID, 2016; POGIS, 2016; Bischoff *et al.*, 2018; Froese and Pauly, 2019). According to Compagno (2016) this species is sometimes confused with two other species of the genus *Sphyrna*, namely *S. lewini* (Griffith & Smith 1834) and *S. mokarran* (Rüppell 1837).

Order Squaliformes

Family Dalatiidae

Dalatias licha (Bonnaterre 1788)

Common name: **Kitefin shark; Gata**, Carocho, Gata-lixa, Gato, Lixa, Lixa-de-lé, Lixa-de-pau, Lixa-de-ré, Negra, Pailona, Peixelim, Pexelim, Pichelim, Rinquim, Trabolha, Tubarão.

References: mainland Portugal[▲] (Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, b, 1880; Gervais and Boulart, 1877a; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1894, 1935; Vieira, 1897; Bragança, 1903, 1904a; Brauer, 1906; Seabra, 1911; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Tortonese, 1979; Fernández *et al.*, 1981; Compagno, 1984a; McEachran and Branstetter, 1984; Sanches, 1986; Fischer *et al.*, 1987; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002; Santos *et al.*, 2002b; Moura, 2004; ICES, 2005; Costa, 2007; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Silva, 2008; Bordalo-Machado *et al.*, 2009; Correia, 2009; Gordo, 2009; Rodrigues, 2009; Silva *et al.*, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Moura *et al.*, 2015a, b; Walls

and Guallart, 2015; Bañón *et al.*, 2016a; ICES, 2017a; Priede, 2017; Froese and Pauly, 2019), Azores (Cadenat, 1937; Ferreira, 1939; Bigelow and Schroeder, 1948; Collins, 1954; Clarke, 1956; Osório de Castro, 1967; Dias *et al.*, 1976, 1990; Saldanha, 1977a; Wheeler, 1978; Krefft and Tortonese, 1979; Fischer *et al.*, 1981; Muzavor and Westhaus, 1981; Martins, 1982a; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Shcherbachov *et al.*, 1985a; Dias and Monteiro, 1986; Sanches, 1986; Silva, 1986a,b, 1987, 1992; Dias, 1991; Lloris *et al.*, 1991; Clarke and Martins, 1995; Santos *et al.*, 1994a, 1995a, 1997; Silva *et al.*, 1995; Isidro, 1996; Menezes, 1996; Arruda, 1997; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Cardigos *et al.*, 2005; Fowler *et al.*, 2005; Menezes *et al.*, 2006; Correia, 2009; Porteiro *et al.*, 2010, 2013; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Naylor *et al.*, 2012; Besugo, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Walls and Guallart, 2015; Bañón *et al.*, 2016a; Torres *et al.*, 2016; Das and Afonso, 2017; ICES, 2017a; Biscoito *et al.*, 2018; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Lowe, 1837; Günther, 1870; Collett, 1890b; Goode and Bean, 1895, 1896; Roule, 1912; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Tortonese, 1979; Cadenat and Blache 1981; Fischer *et al.*, 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Melendez and Macias, 2007; Correia, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Walls and Guallart, 2015; Bañón *et al.*, 2016a; Delgado *et al.*, 2017; Biscoito *et al.*, 2018; Froese and Pauly, 2019).

Isistius brasiliensis (Quoy & Gaimard 1824)

Common name: **Cookie cutter shark**; **Tubarão-luminoso**, Cação-luminoso.

References: Madeira (Cadenat and Blache, 1981; Stevens, 2015). According to de Figueiredo Petean and de Carvalho (2018) and Eschmeyer *et al.* (2019) *I. brasiliensis* is a circumglobal species in tropical and warm temperate seas, although the records are patchy as mentioned by Stevens (2015). Only one specimen has been caught off Madeira (Cadenat and Blache, 1981) and its occurrence in Madeiran EEZ was not yet confirmed by more recent data.

Isistius plutodus Garrick & Springer 1964

Common name: **Largetooth cookiecutter shark**; **Tubarão-dentão**.

References: Azores (PECS) (Zidowitz *et al.*, 2004; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019). According to Eschmeyer *et al.* (2019) *I. plutodus* is a antiequatorial species in the Atlantic and western Pacific and it is probably a circumglobal species as reported by de Figueiredo Petean and de Carvalho (2018).

Squaliolus laticaudus Smith & Radcliffe 1912

Common name: **Spined pygmy shark**; **Tubarão-anão**, Cação-corcunda, Corcunda, Tolho-anão.

References: mainland Portugal (McEachran and Branstetter, 1984; Ebert and Stehmann, 2013(?); Carneiro *et al.*, 2014), Azores (Lloris *et al.*, 1991; Santos *et al.*, 1997; Aires da Silva *et al.*, 1998; Menezes *et al.*, 2006; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Naylor *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Compagno, 2016; Das and Afonso, 2017; ICES, 2017a; Biscoito *et al.*, 2018) and Madeira (FMNH 58862, 1924; Noronha, 1926a; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d; Albuquerque, 1954-1956; Osório de Castro, 1967; Krefft and Tortonese, 1979; Cadenat and Blache 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Aires da Silva *et al.*, 1998; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Biscoito *et al.*, 2018; Froese and Pauly, 2019). The holotype (unique) of *Squaliolus sarmenti* Noronha 1926 caught off Câmara de Lobos (Madeira), synonym of *S. laticaudus*, is in the FMNH collections (FMNH 58862, 1924). Ebert and Stehmann (2013) considered that the distribution of *S. laticaudus* in mainland Portugal is possible.

Family Etomopteridae

Centroscyllium fabricii (Reinhardt 1825)

Common name: **Black dogfish**; **Lixinha-negra**, Cação-torto.

References: mainland Portugal (Compagno, 1984a, 2016; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Azores (Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Naylor *et al.*, 2012; Menezes and Giacomello, 2013; Carneiro *et al.*, 2014; Das and Afonso, 2017)

and Madeira (?) ([Compagno, 2016\(?\)](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is Greenland, both sides of the North Atlantic and south-western Atlantic. According to Compagno (2016) the occurrence of *C. fabricii* in the Madeira Archipelago is questionable. The occurrence of this species in Madeira needs further documentation. Thus, this species is not considered in the total number of valid species (Madeira).

***Etomopterus princeps* Collet 1904**

Common name: **Great lanternshark; Lixinha-da-fundura-grada**, Xarinha-preta, Gata-negra, Raimudo-pequeno, Xarinha.

References: mainland Portugal ([Krefft and Tortonese, 1979](#); [Compagno, 1984a, 2016](#); [McEachran and Branstetter, 1984](#); [Saldanha et al., 1995](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [Jung et al., 2015](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Saldanha and Biscoito, 1997](#); [Santos et al., 1997](#); [Desbruyères et al., 2001](#); [Hareide and Garnes, 2001](#) (PECS); [Fowler et al., 2005](#); [Martins et al., 2006](#); [Menezes et al., 2006, 2009](#); [Bergstad et al., 2008a](#) (PECS); [Porteiro et al., 2010, 2017](#); [Barreiros and Gadig, 2011](#); [Menezes et al., 2012](#); [Besugo, 2013](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Jung et al., 2015](#); [Compagno, 2016](#); [Das and Afonso, 2017](#); [ICES, 2017a](#); [Biscoito et al., 2018](#); [Fauconnet et al., 2019](#)) and Madeira ([Freitas and Biscoito, 2007](#); [Naylor et al., 2012](#); [Carneiro et al., 2014](#); [Compagno, 2016\(?\)](#); [Biscoito et al., 2017, 2018](#)). The occurrence of this species in Madeira is known at least since 1984.

***Etomopterus pusillus* (Lowe 1839)**

Common name: **Smooth lanternshark; Xarinha-preta**, Cação, Gata, Gata-negra, Gata-nigra, Gata-preta, Lixinha, Lixinha-da-fundura, Lixinha-de-fundura, Peixelim, Pexelim, Quelmazinha, Raimudo-pequeno, Xara, Xara-preta, Xarinha.

References: mainland Portugal ([Bragança, 1904a](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Tortonese, 1979](#); [Martins, 1981](#); [INIP, 1982a, b](#); [Borges, 1984](#); [Compagno, 1984a](#); [McEachran and Branstetter, 1984](#); [Dinis, 1985](#); [Sanches, 1986](#); [Figueiredo et al., 1994](#); [Saldanha et al., 1995](#); [Erzini et al., 2001b](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002b](#); [Coelho et al., 2003, 2005](#); [Moura, 2004](#); [Coelho and Erzini, 2005, 2007](#); [ICES, 2005, 2006](#); [Costa, 2007](#); [Fernandes et al., 2007](#); [Silva, 2008, 2011](#); [Ward et al., 2008](#); [Bordalo-Machado et al., 2009](#); [Bordalo-Machado and Figueiredo, 2009](#); [Rodrigues, 2009](#); [Silva et al., 2009](#); [Silva, 2010](#); [Naylor et al., 2012](#); [Chaves and Silva, 2013](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Moura et al., 2015a, b](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores ([Roule, 1912, 1919](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Dias et al., 1976](#); [Krefft and Tortonese, 1979](#); [Compagno, 1984a](#); [McEachran and Branstetter, 1984](#); [INIP, 1985a, b](#); [Dias and Monteiro, 1986](#); [Sanches, 1986](#); [Dias, 1991](#); [Lloris et al., 1991](#); [Menezes, 1996](#); [Arruda, 1997](#); [Saldanha and Biscoito, 1997](#); [Santos et al., 1997](#); [Morato-Gomes et al., 1998](#); [Fowler et al., 2005](#); [Menezes et al., 2006, 2009](#); [Rosa et al., 2006](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Costa et al., 2012](#); [Menezes et al., 2012](#); [Naylor et al., 2012](#); [Besugo, 2013](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#); [ICES, 2017a](#); [Biscoito et al., 2018](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1833a, 1837, 1839, 1840b, 1844a, 1843-1860, 1849](#); [NHMUK 1855.11.29.27, 1854-1861](#); [Günther, 1870](#); [Collet 1890b](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Garman, 1913](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Tortonese, 1979](#); [INIP, 1980, 1982c](#); [Compagno, 1984a, 2016](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Ferreira et al., 2008](#); [Naylor et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [POGID, 2016](#); [Biscoito et al., 2017, 2018](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)). Two syntypes of *Acanthidium pusillum* Lowe 1839, synonym of *E. pusillus*, are in the NHMUK collections (NHMUK 1855.11.29.27) caught off Madeira.

***Etomopterus spinax* (Linnaeus 1758)**

Common name: **Velvet belly; Lixinha-da-fundura**, Cação, Lixinha, Lixinha-de-fundura, Quelmazinha; Peixe-prego.

References: mainland Portugal ([Vandelli, 1771](#); [de Brito Capello, 1880](#); [Bragança, 1904a](#); [Brauer, 1906](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Tortonese, 1979](#); [Fischer et al., 1981, 1987](#); [Compagno, 1984a, 2016](#); [McEachran and](#)

Branstetter, 1984; Sanches, 1986; Figueiredo *et al.* 1994; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002b; Moura, 2004; Coelho and Erzini, 2005; ICES, 2005; Sousa *et al.*, 2005, 2006; Menezes *et al.*, 2006; Neiva *et al.*, 2006; Rosa *et al.*, 2006; Costa, 2007; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Chaves, 2008; Silva, 2008, 2010, 2011; Bordalo-Machado *et al.*, 2009; Rodrigues, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Moura *et al.*, 2015a, b; Bañón *et al.*, 2016a; ICES, 2017a; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Richard, 1905b; Brian, 1912; Roule, 1912, 1919; Fowler, 1936a; Ferreira, 1939; Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1967; Compagno, 1984a; McEachran and Branstetter, 1984; Sanches, 1986; Silva *et al.*, 1995; Menezes, 1996; Arruda, 1997; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Fowler *et al.*, 2005; Melendez and Macias, 2007; Aranha *et al.*, 2009; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Das and Afonso, 2017; ICES, 2017a; Biscoito *et al.*, 2018; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1833a, 1849; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Grey, 1956; Krefft and Tortonese, 1979; Fischer *et al.*, 1981, 1987; Martins, 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Santos *et al.*, 2002a; Figueiredo *et al.*, 2005; Fowler *et al.*, 2005; Melendez and Macias, 2007; Moura *et al.*, 2008a, 2014; Bordalo-Machado *et al.*, 2009; Bordalo-Machado and Figueiredo, 2009; Correia, 2009; Gordo, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Catarino *et al.*, 2015; ICES, 2017a; Roberts *et al.*, 2017; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores (Regan 1908; Roule, 1912, 1919; Guiart, 1935; Fowler, 1936a; Ferreira, 1939; Bigelow and Schroeder, 1948; Albuquerque, 1954-1956; Collins, 1954; Grey, 1956; Osório de Castro, 1967; Golovan, 1976; Wheeler, 1978; Krefft and Tortonese, 1979; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Arruda, 1997; Saldanha and Biscoito, 1997; Santos *et al.*, 1997; Desbruyères *et al.*, 2001; Hareide and Garnes, 2001 (PECS); Fowler *et al.*, 2005; Menezes *et al.*, 2006, 2009; Bergstad *et al.*, 2008a (PECS); Correia, 2009; Porteiro *et al.*, 2010, 2017; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Besugo, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2014; Catarino *et al.*, 2015; Das and Afonso, 2017; Biscoito *et al.*, 2018; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira[▲] (de Brito Capello, 1872; Goode and Bean, 1895, 1896; Brauer, 1906; Belloc, 1934; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Grey, 1956; Krefft and Tortonese, 1979; Cadenat and Blache 1981; Fischer *et al.*, 1981; INIP, 1980, 1982c; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Severino, 2004; Correia, 2009; Naylor *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Catarino *et al.*, 2015; Biscoito *et al.*, 2017, 2018; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019). This species is sometimes confused with *C. owstonii* Garman 1906 (Compagno, 2016). The holotype of this species caught off mainland Portugal was preserved in the Museu Bocage (MB T.113) and destroyed in the 1978 fire.

***Centroscymnus owstonii* Garman 1906**

Common name: **Roughskin dogfish; Xara-preta-de-natura**, Pailona, Tubarão, Xara, Xara-preta.

References: mainland Portugal (Moura *et al.*, 2008a, b; Bordalo-Machado *et al.*, 2009; Gordo, 2009; Rodrigues, 2009; Naylor *et al.*, 2012; Carneiro *et al.*, 2014), Azores[▲] (Compagno, 1984a, 2016; Santos *et al.*, 1997; Menezes *et al.*, 2006, 2009; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Besugo, 2013;

Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira (NHMUK 1865.5.20.14, 1864-1877; Günther, 1870; Regan 1906; Garman, 1913; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1949d; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cadenat and Blache 1981; McEachran and Branstetter, 1984; Compagno, 1984a, 2016; Sanches, 1986; Lloris *et al.*, 1991; Naylor *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2017, 2018; Porteiro *et al.*, 2017; Froese and Pauly, 2019). This species is sometimes confused with *C. coelolepis* Barbosa du Bocage & de Brito Capello 1864 (Compagno, 2016). The holotype (unique) of *Centroscymnus cryptacanthus* Regan 1906 is in the NHMUK collections (NHMUK 1865.5.20.14) caught off Madeira, synonym of *C. owstonii*.

Centroselachus crepidater (Barbosa du Bocage & de Brito Capello 1864)

Common name: **Roughskin dogfish; Sapata-preta**, Sapata-de-natura, Tubarão.

References: mainland Portugal[▲] (Barboza du Bocage and de Brito Capello, 1864a, b, 1865, 1866; de Brito Capello, 1869a, 1880; Osório, 1888; Goode and Bean, 1895, 1896; Bragança, 1904a; Seabra, 1911; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Kreff and Tortonese, 1979; Fischer *et al.*, 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Bordalo-Machado *et al.*, 2009; Bordalo-Machado and Figueiredo, 2009; Correia, 2009; Gordo, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2015a, b; Bañón *et al.*, 2016a; ICES, 2017a; Roberts *et al.*, 2017; Froese and Pauly, 2019), Azores (Santos *et al.*, 1997; Menezes *et al.*, 2006, 2009; Rosa *et al.*, 2006; Porteiro *et al.*, 2010, 2017; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Besugo, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Compagno, 2016; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018) and Madeira (Günther, 1870; Goode and Bean, 1895, 1896; Brauer, 1906; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cadenat and Blache 1981; Fischer *et al.*, 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Porteiro *et al.*, 2010; Naylor *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; POGID, 2016; Bischoff *et al.*, 2018; Froese and Pauly, 2019). The holotype (unique) of *Centrophorus crepidater* Barbosa du Bocage & de Brito Capello 1864 caught off mainland Portugal was preserved in the Museu Bocage (MB T.112), synonym of *Centroselachus crepidater*, but was destroyed in the 1978 fire. One specimen caught off in the Portuguese continental slope in September 2009 is in the MB collections.

Scymnodalatias garricki Kukuev & Konovalenko 1988

Common name: **Azores dogfish; Tubarão-cão-dos-Açores**, Tubarão-dos-Açores, Peixe-cão-dos-Açores.

References: Azores (SAMS-SHARKS-012649, 1977; Kukuev and Konovalenko, 1988; Arruda, 1997; Santos *et al.*, 1997; Kukuev, 2006 (PECS); Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Séret *et al.*, 2015a; Compagno, 2016; Das and Afonso, 2017; Froese and Pauly, 2019).

Scymnodon ringens Barbosa du Bocage & de Brito Capello 1864

Common name: **Knifetooth dogfish; Arreganhada**, Boca, Bruxa, Negra, Peixelim, Pexelim, Risota, Tubarão, Xara, Xara-preta-de-focinho-comprido.

References: mainland Portugal (Barboza du Bocage and de Brito Capello, 1864a, b, 1865, 1866; de Brito Capello, 1869a, 1880; NHMUK 1867.7.23.3, 1864 - 1877; Almeida and Roquette, 1892; Goode and Bean, 1895, 1896; Vieira, 1897; Bragança, 1904a; Brauer, 1906; Seabra, 1911; Roule, 1912; Nobre, 1935; Fowler, 1936a; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Kreff and Tortonese, 1979; Cadenat and Blache 1981; Fernández *et al.*, 1981; Fischer *et al.*, 1981; Compagno, 1984a, 2016; Dinis and Marecos, 1984; McEachran and Branstetter, 1984; Sanches, 1986; Borges *et al.*, 2002; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Silva, 2008, 2011; Bordalo-Machado *et al.*, 2009; Bordalo-Machado and Figueiredo, 2009; Rodrigues, 2009; Silva *et al.*, 2009; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Moura *et al.*, 2015a, b; Bañón *et al.*, 2016a; ICES, 2017a; Roberts *et al.*, 2017; Froese and Pauly, 2019) and Madeira (de Brito Capello, 1872; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1949d; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Kreff and Tortonese, 1979; Cadenat and Blache 1981; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Compagno, 2016(?);

Delgado *et al.*, 2017; Bischoff *et al.*, 2018). One syntype of this species caught off mainland Portugal is in the NHMUK collections (NHMUK 1867.7.23.3).

***Somniosus microcephalus* (Bloch & Schneider 1801)**

Common name: **Greenland shark; Tubarão-da-Gronelândia**, Pailona, Tubarão, Tubarão-da-noite.

References: mainland Portugal (de Brito Capello, 1880; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Quéro *et al.*, 1976; Krefft and Tortonese, 1979(?); Compagno, 1984a(?), 2016(?); McEachran and Branstetter, 1984(?); Sanches, 1986; Francis *et al.*, 1988(?); Bordalo-Machado *et al.*, 2009; Correia, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013(?); Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (MAR-ECO 16601, 2004; MAR-ECO 16349, 2004; Fossen *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Das and Afonso, 2017; Bischoff *et al.*, 2018; Mecklenburg *et al.*, 2018) and Madeira (Quéro *et al.*, 1976; Sanches, 1986; Freitas and Bischoff, 2007; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018). One specimen is in the MMF collections (MMF 23429, 1983).

***Somniosus rostratus* (Risso 1827)**

Common name: **Little sleeper shark; Pailona**, Borracha, Tubarão, Trabolha-de-natura.

References: mainland Portugal (de Brito Capello, 1869a, 1880; Bragança, 1904a; Seabra, 1911; Nobre, 1935; Osório de Castro, 1967; Krefft and Tortonese, 1979; Cadenat and Blache 1981; Fischer *et al.*, 1981, 1987; Martins, 1981; McEachran and Branstetter, 1984; Compagno, 1984a(?), 2016; Sanches, 1986; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2015a; Guallart *et al.*, 2015; Bañón *et al.*, 2016a), Azores (Fowler *et al.*, 2005; Menezes *et al.*, 2006; Wenneck *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Barreiros and Gadig, 2011; Carneiro *et al.*, 2014; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018) and Madeira (Noronha, 1926a; Maul, 1955a; Krefft and Tortonese, 1979; Cadenat and Blache 1981; Fischer *et al.*, 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Melendez and Macias, 2007; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Guallart *et al.*, 2015; Bañón *et al.*, 2016a; Bischoff *et al.*, 2018; Froese and Pauly, 2019).

***Zameus squamulosus* (Günther 1877)**

Common name: **Smallmouth knifetooth dogfish; Arreganhada-de-focinho-comprido**, Arreganhada, Lixa-de-pau, Pailona, Sapata-preta, Tubarão, Xara, Xara-preta, Xara-preta-de-focinho-comprido.

References: mainland Portugal (Günther, 1880; Compagno, 1984a, 2016; Last and Stevens, 1994; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (IMAR/DOP 1AÇO-03-P96XAR, 1996; Menezes *et al.*, 2012; Besugo, 2013; Carneiro *et al.*, 2014; Compagno, 2016; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira (Günther, 1880; Noronha, 1926a; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Maul, 1949d; Albuquerque, 1954-1956; Osório de Castro, 1967; Krefft and Tortonese, 1979; Cadenat and Blache 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018; Froese and Pauly, 2019).

Family Oxynotidae

***Oxynotus centrina* (Linnaeus 1758)**

Common name: **Angular roughshark; Peixe-porco**, Cação, Gato, Peixe-gato, Peixe-rato, Peixe-gato, Porco, Porco-marinho, Rato, Rinquim, Tambor, Tubarão.

References: mainland Portugal▲ (Vandelli, 1771, 1797; Barboza du Bocage and de Brito Capello, 1866; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Almeida and Roquette, 1892; Nobre, 1894, 1895b, c, 1903, 1935; Goode and Bean, 1895, 1896; Osório, 1895, 1917; Vieira, 1897; Bragança, 1903, 1904a; Seabra, 1911; Murray and Hjort, 1912; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Tortonese, 1979; Fernández *et al.*, 1981, 1987; Fischer *et al.*, 1981; Martins, 1981; Compagno, 1984a, 2016; Quéro, 1984b; Sanches, 1986; ICES, 2005; Bradai *et al.*, 2007; Melendez and Macias, 2007; Henriques *et al.*, 2008; Rodrigues, 2009; Silva, 2010; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Soldo and Guallart, 2015; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967) and Madeira (Lowe, 1837; Noronha and Sarmento, 1934; Nobre, 1935; Osório de Castro, 1967; Fischer *et al.*, 1981; Bradai *et al.*, 2007; Ebert and Stehmann,

2013; Soldo and Guallart, 2015; Compagno, 2016; Biscoito *et al.*, 2018(?)). The only report on the presence of *O. centrina* in the fishing grounds of the Azores is indicated by Osório de Castro (1967). Current geographic distribution of this species is on the continental shelves of the Mediterranean Sea and, in the Eastern Atlantic, from North-western North Sea and English Channel to South Africa. Since then no other author describes the presence of this species in this region. Thus, the occurrence of this species in the Azores Archipelago is questionable and is not considered in the total number of valid species for the Azorean ichthyofauna.

***Oxynotus paradoxus* Trade 1929**

Common name: **Sailfin roughshark; Peixe-porco-de-vela**, Cação, Gato, Peixe-gato, Peixe-porco, Peixe-rato, Rato.

References: mainland Portugal (Kreff and Tortonese, 1979; Compagno, 1984a, 2016; Quéro, 1984b; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2015a; Soldo *et al.*, 2015; Bañón *et al.*, 2016a; Compagno, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Santos *et al.*, 1997; Azevedo *et al.*, 2003; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Soldo *et al.*, 2015; Compagno, 2016; Das and Afonso, 2017; Biscoito *et al.*, 2018) and Madeira (Quéro, 1984b; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Soldo *et al.*, 2015; Compagno, 2016; Biscoito *et al.*, 2018(?)).

Family Centrophoridae

***Centrophorus granulosus* (Bloch & Schneider 1801)**

Common name: **Gulper shark; Barroso**, Cação, Chapenta, Chapeuta, Chaputa, Guelmo, Lixa, Lixa-de-lei, Queime, Quelme, Quelme, Quelvacho, Ramudo, Rinquim, Sarraco, Sarroco, Tubarão, Xaputa.

References: mainland Portugal[▲] (Barboza du Bocage and de Brito Capello, 1864a, b, 1865, 1866; de Brito Capello, 1869a, b, 1880; Vinciguerra, 1883; Almeida and Roquette, 1892; Girard, 1892; Goode and Bean, 1895(?), 1896(?); Nobre, 1895a, 1935; Vieira, 1897; Bragança, 1903, 1904a; Brauer, 1906; Seabra, 1911; Helling, 1940; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Kreff and Tortonese, 1979; Fischer *et al.*, 1981, 1987; Martins, 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Figueiredo *et al.*, 1994, 2005; Saldanha *et al.*, 1995; Compagno and Niem, 1998; Borges *et al.*, 2002; Moura, 2004; Fowler *et al.*, 2005; ICES, 2005; Costa, 2007; Melendez and Macias, 2007; Moura *et al.*, 2008a, 2014, 2015b; Bordalo-Machado *et al.*, 2009; Correia, 2009; Gordo, 2009; Rodrigues, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; White *et al.*, 2013; Costa, 2014; Carneiro *et al.*, 2014; ICES, 2017a; Froese and Pauly, 2019), Azores[▲] (Neuville, 1897; Ferreira, 1939; Collins, 1954; Osório de Castro, 1967; INIP, 1985b; Dias, 1991; Arruda, 1997; Santos *et al.*, 1997; Fowler *et al.*, 2005; Menezes *et al.*, 2006; Correia, 2009; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Menezes *et al.*, 2012; Besugo, 2013; White *et al.*, 2013(?); Carneiro *et al.*, 2014; Moura *et al.*, 2014, 2015; Torres *et al.*, 2016; Das and Afonso, 2017; Biscoito *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira[▲] (Günther, 1870; Goode and Bean, 1895, 1896; Belloc, 1934; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul 1949d, 1955a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cadenat and Blache 1981; Fischer *et al.*, 1981; INIP, 1982c, 1984a, b, c; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Freitas and Biscoito, 2007; Ferreira *et al.*, 2008; Correia, 2009; Naylor *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2014; POGID, 2016; Biscoito *et al.*, 2017, 2018; Froese and Pauly, 2019). This species is sometimes confused with *Centrophorus uyato* (Rafinesque 1810) (Compagno, 2016). *Centrophorus niaukang* Teng 1959 is synonym of *C. granulosus*.

***Centrophorus lusitanicus* Barbosa du Bocage & de Brito Capello 1864**

Common name: **Lowfin gulper shark; Tubarão-lusitano**, Barroso, Chapeuta, Chaputa, Lixa-de-lei, Quelmo, Ramudo, Sarroco.

References: mainland Portugal (Barbosa du Bocage and de Brito Capello, 1864a, b(?), 1866(?); de Brito Capello, 1880; Goode and Bean, 1895, 1896; Vieira, 1897; Bragança, 1904a; Brauer, 1906; Seabra, 1911; Nobre, 1935; Osório de Castro, 1954; Kreff and Tortonese, 1979; Compagno, 1984a, 2016; Fischer *et al.*, 1981; McEachran and Branstetter, 1984; Bordalo-Machado *et al.*, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Moura *et al.*, 2015a; ICES, 2017a; Froese and Pauly, 2019),

Azores ([Das and Afonso, 2017](#); [Fauconnet et al., 2019](#)) and Madeira ([McEachran and Branstetter, 1984](#); [Carneiro et al., 2014](#); [Compagno, 2016\(?\)](#); [Das and Afonso, 2017](#)). Barbosa du Bocage and de Brito Capello (1864) described for the first time the species *Centrophorus lusitanicus* from Portugal and the specimen was deposited in the Natural History Museum in London (former syntype BMNH 1867.7.23.2). White *et al.* (2017a) considered that the original description and illustration of the specimen examined by Barbosa du Bocage and de Brito Capello corresponded to a junior synonym of *C. granulosus*. The former authors also concluded that the specimen in the Natural History Museum in London should not be considered a syntype of *C. lusitanicus* but a long-finned species. The former syntype of *C. lusitanicus* was re-evaluated as a paratype of *Centrophorus lesliei* White, Ebert & Naylor 2017 (NHMUK 1867.7.23.2, 1864 - 1877). According to Das and Afonso (2017) the presence of this species has only recently been confirmed from the Azores but is presumably rare. This species is sometimes confused with *C. granulosus* (Bloch & Schneider 1801) (Ebert and Stehmann, 2013).

Centrophorus machiquensis Maul 1955

Common name: No English common name; **Quelmo**.

References: Madeira ([Maul, 1955a](#); [Krefft and Tortonese, 1979](#); [Cadenat and Blache, 1981](#); [McEachran and Branstetter, 1984](#); [Lloris et al., 1991](#); [Biscoito et al., 2018](#)). The holotype (unique) of this species caught off Machico (Madeira) is in the MMF collections (MMF 3767, 1941). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2019) *C. machiquensis* is synonym of *Centrophorus granulosus* (Bloch & Schneider 1801). Cadenat and Blache (1981) considered as *Centrophorus cf. uyato – machiquensis*. Nevertheless, Biscoito *et al.* (2018) treated *C. machiquensis* as a valid species, based on its morphological and ecological characters.

Centrophorus squamosus (Bonnaterre 1788)

Common name: **Leafscale gulper shark**; **Lixa**, Arreganhada, Lixa-de-escama, Lixa-de-pau, Peixelim, Pexelim, Raimudo, Ramudo, Rinquim, Sapata, Sapata-preta, Tubarão, Xara, Xara-branca.

References: mainland Portugal▲ ([Carvalho, 1765\(?\)](#); [Barboza du Bocage and de Brito Capello, 1864a, 1866](#); [de Brito Capello, 1869a, 1880](#); [Günther, 1880](#); [Goode and Bean, 1895, 1896](#); [Vieira, 1897](#); [Bragança, 1902, 1903, 1904a](#); [Brauer, 1906](#); [Richard, 1908](#); [Seabra, 1911](#); [Roule, 1912](#); [Nobre, 1935](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Grey, 1956](#); [Kreff and Tortonese, 1979](#); [Fischer et al., 1981](#); [Compagno, 1984a](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986](#); [Figueiredo et al., 2005](#); [Fowler et al., 2005](#); [Moura et al. 2008a](#); [Bordalo-Machado et al., 2009](#); [Correia, 2009](#); [Gordo, 2009](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Veiga et al., 2013](#); [Carneiro et al., 2014](#); [Priede, 2017](#); [Froese and Pauly, 2019](#)), Azores▲ ([Hilgendorf, 1888a](#); [Richard, 1912](#); [Roule, 1912, 1919](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [Collignon and Aloncle, 1972](#); [Kreff and Tortonese, 1979](#); [Fischer et al., 1981](#); [Compagno, 1984a, 2016](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Hareide and Garnes, 2001 \(PECS\)](#); [Fowler et al., 2005](#); [Menezes et al., 2006, 2009, 2012](#); [Correia, 2009](#); [Barreiros and Gadig, 2011](#); [Porteiro et al., 2010, 2017](#); [Besugo, 2013](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Moura et al., 2015b](#); [Das and Afonso, 2017](#); [ICES, 2017a](#); [Biscoito et al., 2018](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira▲ ([Lowe, 1850, 1852](#); [NHMUK 1865.5.20.15, 1864-1877](#); [Johnson, 1868](#); [Günther, 1870, 1880](#); [Goode and Bean, 1895, 1896](#); [Belloc, 1934](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Grey, 1956](#); [Kreff and Tortonese, 1979](#); [Cadenat and Blache, 1981](#); [Fischer et al., 1981](#); [Compagno, 1984a, 2016](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Martins and Ferreira, 1995](#); [Severino, 2004](#); [Correia, 2009](#); [Severino et al., 2009](#); [Naylor et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Delgado et al., 2017](#); [ICES, 2017a](#); [Porteiro et al., 2017](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Machecephalus dumerili* Johnson 1868 caught off Madeira, synonym of *C. squamosus*, is in the NHMUK collections (NHMUK 1865.5.20.15).

Centrophorus uyato (Rafinesque 1810)

Common name: **Little gulper shark**; **Quelme**, Tubarão.

References: mainland Portugal ([NHMUK 1904.11.30.11, 1904-1925\(?\)](#); [NHMUK 1904.11.30.12, 1904-1925\(?\)](#); [Regan, 1906](#); [INIP, 1982a, b](#); [McEachran and Branstetter, 1984](#); [Compagno 1984a](#); [Fischer et al., 1987](#); [Last and Stevens, 1994](#); [Saldanha et al., 1995](#); [Guallart and Walls, 2015](#); [Moura et al., 2015a](#);

Froese and Pauly, 2019) and Madeira (NHMUK 1862.4.22.29, 1862; McEachran and Branstetter, 1984; Guallart and Walls, 2015; Biscoito et al., 2018). Two syntypes of *Centrophorus bragancae* Regan 1906, both caught off Sesimbra, are in the NHMUK collections (NHMUK 1904.11.30.11, and NHMUK 1904.11.30.12) but their current status is uncertain as *C. uyato*. The DNA barcoding analysis of the mitochondrial cytochrome oxidase subunit I (col) gene for species of *Centrophorus* suggests conspecificity of *C. uyato* and *Centrophorus zeehaani* White, Ebert & Compagno 2008 (Wienerroither et al., 2015).

Centrophorus cf. zeehaani White, Ebert & Compagno 2008

Common name: **Little gulper shark; Quelme-do-Zeehaan.**

References: Madeira (?) (Naylor et al., 2012). White et al. (2008), Naylor et al. (2012) and Wienerroither et al. (2015) considered *C. zeehaani* an endemic species of southern Australia. However, Eschmeyer et al. (2019) widen the geographic distribution of this species in the Eastern Indian Ocean and western Pacific: Andaman Islands and around southern Australia from New South Wales and Tasmania to western Australia. Naylor et al. (2012) evaluate a specimen from Madeira and provisionally designated as *C. cf. zeehaani*, which was confirmed by Wienerroither et al. (2015). These latter authors, based on the morphology, morphometry and DNA barcoding, concluded that *C. uyato* and *C. zeehaani* are conspecific. Thus, the occurrence of this species in Madeira Archipelago is questionable and is not considered in the total number of valid species for the Madeiran ichthyofauna.

Deania calcea (Lowe 1839)

Common name: **Birdbeak dogfish; Sapata**, Ferreta, Ferrête, Pífaro, Pife, Rinquim, Sapata, Sapata-branca, Tubarão, Tutia, Zapata-branca.

References: mainland Portugal▲ (Barboza du Bocage and Brito Capello, 1864a, b, 1866; de Brito Capello, 1869a, 1880; NHMUK 1867.7.23.1, 1864-1877; Nobre, 1894, 1935; Goode and Bean, 1895, 1896; Vieira, 1897; Bragança, 1902, 1903, 1904a; Brauer, 1906; Seabra, 1911; Roule, 1912; Helling, 1940; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández et al., 1981; Fischer et al., 1981; INIP, 1981a, 1982a; Martins, 1981; McEachran and Branstetter, 1984; Compagno, 1984a; Sanches, 1986; Compagno, et al., 1989; Figueiredo et al., 1994; Saldanha et al., 1995; Monteiro et al., 2001; Borges et al., 2002; Moura, 2004; ICES, 2005; Sousa et al., 2005; Costa, 2007; Fernandes et al., 2007; Henriques et al., 2008; Pakhorukov, 2008; Bordalo-Machado et al., 2009; Correia, 2009; Rodrigues, 2009; Silva et al., 2009; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; Christiansen et al., 2014; Moura et al., 2014, 2015a, b; OCEANA, 2014; Silva and Borges, 2014; Dureuil, 2015; Bañón et al., 2016a; ICES, 2017a; Froese and Pauly, 2019), Azores (Richard, 1905b; Roule, 1912, 1919a; Guiart, 1935; Fowler, 1936a; Ferreira, 1939; Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1954, 1967; Dias et al., 1976; Saldanha, 1977a, 1991, 1995; Kreff and Tortonese, 1979; INIP, 1985a, b, c; Sanches, 1986; Dias, 1991; Gonçalves 1994; Van Dover et al., 1996(?); Arruda, 1997; Santos et al., 1997; Morato-Gomes et al., 1998; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Fowler et al., 2005; Menezes et al., 2006, 2009; Martins et al., 2006; Rosa et al., 2006; Pakhorukov, 2008 (PECS); Barreiros and Gadig, 2011; Menezes et al., 2012; Besugo, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; ICES, 2017a; Porteiro et al., 2017; Das and Afonso, 2017; ICES, 2017a; Biscoito et al., 2018; Fauconnet et al., 2019) and Madeira▲ (Lowe, 1839, 1840b, 1843, 1844a, 1849; NHMUK 1861.5.19.33, 1854-1861; Günther, 1870; de Brito Capello, 1872; Goode and Bean, 1895, 1896; Garman, 1913; Belloc, 1934; Fowler, 1936a; Noronha and Sarmento, 1948; Maul 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Cadenat and Blache 1981; Fischer et al., 1981; Compagno, 1984a, 2016; McEachran and Branstetter, 1984; Sanches, 1986; Pakhorukov, 2008 (PECS); Ebert and Stehmann, 2013; Carneiro et al., 2014; Dureuil, 2015; POGID, 2016; Roberts et al., 2017; Biscoito et al., 2018; Froese and Pauly, 2019). Three syntypes of *Centrophorus crepidalbus* Barbosa du Bocage & de Brito Capello 1864 are in NHMUK (NHMUK 1867.7.23.1), MNHN (MNHN-IC-0000-4801), and ZMB collections (ZMB 6471), all caught off mainland Portugal. The holotype (unique) of *Acanthidium calceus* Lowe 1839, synonym of *D. calcea*, caught off Madeira is in NHMUK collections (NHMUK 1861.5.19.33).

Deania hystricosa (Garman 1906)

Common name: **Rough longnose dogfish; Sapata-áspera.**

References: mainland Portugal ([MNHN-IC-A-3929](#), 1881; Rodrigues, 2009; Carneiro *et al.*, 2014), Azores[▲] (Besugo, 2013; Ebert and Stehmann, 2013(?); Carneiro *et al.*, 2014; Ebert *et al.*, 2015; Compagno, 2016; ICES, 2017a; Bischoff *et al.*, 2018) and Madeira (Compagno, 1984a, 2016; Cadenat and Blache 1981; Cappetta *et al.*, 1985; Lloris *et al.*, 1991; Ferreira *et al.*, 2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Ebert *et al.*, 2015; Delgado *et al.*, 2017; Bischoff *et al.*, 2018; Froese and Pauly, 2019). The holotype of *Deania mauli* Cadenat & Blache 1981, synonym of *Deania hystricosa*, was caught off Câmara de Lobos (Madeira).

Deania profundorum (Smith & Ratcliffe 1912)

Common name: **Longsnout dogfish; Sapata-flecha**, Sapata, Sapata-branca, Sapata-guilha.

References: mainland Portugal ([Moura](#), 2004; Silva, 2008, 2010, 2011; [Bordalo-Machado](#) *et al.*, 2009; Gordo, 2009; Rodrigues, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Naylor *et al.*, 2012(?); Chaves and Silva, 2013; Carneiro *et al.*, 2014; OCEANA, 2014; Silva and Borges, 2014; Moura *et al.*, 2015a; ICES, 2017a), Azores[▲] (Silva *et al.*, 1995; Menezes, 1996; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Desbruyères *et al.*, 2001; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Naylor *et al.*, 2012(?); Besugo, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Compagno, 2016; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018; Fauconnet *et al.*, 2019) and Madeira (Cadenat and Blache 1981; Freitas and Bischoff, 2007; Sousa *et al.*, 2009; Carneiro *et al.*, 2014; Compagno, 2016(?); POGID, 2016; Bischoff *et al.*, 2017, 2018; Delgado *et al.*, 2017).

Family Squalidae

Squalus acanthias Linnaeus 1758

Common name: **Spotted spiny dogfish; Galhudo-malhado**, Alitão, Cação, Cação-ferrão, Cação-galhudo, Caúdo, Esgalhudo, Ferranho, Galhudo, Melca, Melga, Melga-de-ferrão, Tubarão.

References: mainland Portugal[▲] ([Vandelli](#), 1771, 1797; [Barboza du Bocage](#) and [de Brito Capello](#), 1866; [de Brito Capello](#), 1869a, 1880; [Gervais](#) and [Boulart](#), 1877a; [Nobre](#), 1894, 1903, 1935; [Osório](#), 1895; [Vieira](#), 1897; [Bragança](#), 1904a; [Seabra](#), 1911; [Helling](#), 1940; [Gonçalves](#), 1941; [Albuquerque](#), 1954-1956; [Osório de Castro](#), 1954, 1967; [Kreff](#) and [Tortonese](#), 1979; Compagno 1984a; McEachran and Branstetter, 1984; Sanches, 1986; Fischer *et al.*, 1987; Melendez and Macias, 2007; Correia, 2009; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Ryland and Lancaster, 2017; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores ([McEachran](#) and [Branstetter](#), 1984; Sanches, 1986; Arruda, 1997; Santos *et al.*, 1997(?); Ebert and Sthmann 2013(?); Carneiro *et al.*, 2014(?); Compagno, 2016(?); Bischoff *et al.*, 2018; Mecklenburg *et al.*, 2018(?)) and Madeira ([Bowdich](#), 1825; [Belloc](#), 1934; [Noronha](#) and [Sarmento](#), 1934; [Nobre](#), 1935; [Fowler](#), 1936a; [Albuquerque](#), 1954-1956; [Osório de Castro](#), 1967; [Kreff](#) and [Tortonese](#), 1979; Cadenat and Blache 1981; Fischer *et al.*, 1981; McEachran and Branstetter, 1984; Sanches, 1986; Lloris *et al.*, 1991; Ebert and Stehmann, 2013(?); Carneiro *et al.*, 2014; Compagno, 2016; Mecklenburg *et al.*, 2018). According to Ebert and Stehmann (2013), Compagno (2016) and Mecklenburg *et al.* (2018) the occurrence of *S. acanthias* in the Azores Archipelago is probable. Nevertheless, Bischoff *et al.* (2018) proposed that this species shouldn't be considered in the list of Squalidae from Madeira. Specimens are needed for confirmation the presence of this species in Madeira.

Squalus blainvillei (Risso 1827)

Common name: **Longnose spurdog; Galhudo**, Alitão, Cação, Cação-de-bico, Cação-ferrão, Caúdo, Esgalhudo, Ferranho, Galhudo, Galhudo-cubano, Melca, Melga, Melga-de-ferrão, Tubarão.

References: mainland Portugal ([Vandelli](#), 1797; [Barboza du Bocage](#) and [de Brito Capello](#), 1866; [de Brito Capello](#), 1869a; [Almeida](#) and [Roquette](#), 1892; [Vieira](#), 1897; [Bragança](#), 1902, 1904a; [Nobre](#), 1903, 1935; [Seabra](#), 1911; [Osório de Castro](#), 1967; [Maul](#), 1976; [Kreff](#) and [Tortonese](#), 1979; Fischer *et al.*, 1981, 1987; [Martins](#), 1981; INIP, 1981a, 1982b; Compagno, 1984a; [Dinis](#) and [Marecos](#), 1984; McEachran and Branstetter, 1984; Sanches, 1986; ICES, 2005; Melendez and Macias, 2007; [Henriques](#) *et al.*, 2008; Rodrigues, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (?) ([McEachran](#) and [Branstetter](#), 1984(?); [Arruda](#), 1997(?); Carneiro *et al.*, 2014(?)) and Madeira ([MNHN-IC-2005-2445](#), n.d.; [McEachran](#) and [Branstetter](#), 1984; Sanches, 1986(?)). According to McEachran and Branstetter (1984) and Arruda (1997) the occurrence of *S. blainvillei* in the Azores Archipelago is doubtful. The occurrence of this species in the Azores needs further documentation. Thus, this species

is not considered in the total number of valid species to the Azores. One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22417, 1967).

Squalus megalops (Macleay 1881)

Common name: **Picked spurdog; Galhudo-de-focinho-curto.**

References: mainland Portugal ([Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Cavanagh et al., 2015](#); [Compagno, 2016](#)). This species is sometimes confused with *Squalus acutipinnis* Regan 1908 (Compagno, 2016).

Squalus mitsukurii Jordan & Snyder 1903

Common name: **Shortspine spurdog; Galhudo-de-espinho-curto.**

References: mainland Portugal ([Cavanagh et al., 2009](#); [Compagno, 2016](#)). Cavanagh et al. (2009) reported that several species were previously considered to be con-specific with *S. mitsukurii*. These authors also consider that further investigation of this species from around the world will likely result in more taxa being recognized. Thus, the occurrence records to mainland Portugal should be revised.

Family Echinorhinidae

Echinorhinus brucus (Bonnaterre 1788)

Common name: **Bramble shark; Tubarão-prego**, Peixe-prego, Prego, Rinquim, Tubarão.

References: mainland Portugal ([Vandelli, 1797](#); [Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, 1880](#); [Almeida and Roquette, 1892](#); [Nobre, 1894, 1895a, 1935](#); [Vieira, 1897](#); [Bragança, 1903, 1904a](#); [Seabra, 1911](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Tortonese, 1979](#); [Fischer et al., 1981, 1987](#); [Compagno, 1984a, 2016](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986](#); [Melendez and Macias, 2007](#); [Correia, 2009](#); [Rodrigues, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967](#); [Wheeler, 1978](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986\(?\)](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Compagno, 2016](#)) and Madeira ([Osório de Castro, 1967](#); [Fischer et al., 1981](#); [McEachran and Branstetter, 1984](#); [Sanches, 1986\(?\)](#)). According to Eschmeyer et al. (2019) *E. brucus* is a cosmopolitan species (including Mediterranean Sea, southern North Sea). The distribution map of this species reported by Fischer et al. (1981) included the area of Selvagens Islands (Madeira EEZ), and McEachran and Branstetter (1984) reported occasional occurrence in the areas of mainland Portugal, Azores and Madeira. However, no recent occurrence records were reported for Madeira Island.

Order Squatiniformes

Family Squatinidae

Squatina aculeata Cuvier 1829

Common name: **Sawback angelshark; Anjo-espinhoso**, Anjo, Anjo-do-mar, Peixe-anjo, Raia, Raia-bastarda, Viola, Violão.

References: mainland Portugal (?) ([Compagno, 1984a](#); [Soldo and Bariche, 2015](#)) and Madeira (?) ([Soldo and Bariche, 2015](#)). According to Eschmeyer et al. (2019) the current geographic distribution of this species is the Mediterranean Sea and, in the Eastern Atlantic, from Morocco to Angola and Namibia. The distribution maps of this species reported by Compagno (1984a) and Soldo and Bariche (2015) included southern mainland Portugal region and Soldo and Bariche (2015) included the area of Selvagens Islands in the Madeira EEZ. The historical records of this species result from Portuguese landings data from the fleet operating off Morocco and Mauritania and occurrence records in mainland Portugal or Madeira, based on specimens observation, have not been reported. The geographic distribution of *S. aculeata* in mainland Portugal should be considered as a former range of occurrence. Thus, this species is not considered in the total number of valid species to mainland Portugal and Madeira.

Squatina oculata Bonaparte 1840

Common name: **Smoothback angelshark; Anjo-de-malhas**, Anjo, Anjo-do-mar, Peixe-anjo, Raia, Raia-bastarda, Viola, Violão.

References: mainland Portugal ([MNHN-IC-1905-0572](#), 1902; [SAMS-SHARKS-014378](#), no date; Compagno, 1984a, 2016; Carneiro et al., 2014; Ferretti et al., 2015c) and Madeira (?) ([Ferretti et al., 2015c](#)). According to Eschmeyer et al. (2019) the current geographic distribution of this species is the Mediterranean Sea and, in the Eastern Atlantic, from Morocco to Angola and Namibia. The distribution map of this species reported by Ferretti et al. (2015c) included mainland Portugal and the area of Selvagens Islands (Madeira EEZ). There are no recent records of captures of smoothback angelshark in mainland Portugal and can be considered a rare species with uncertain presence. Because *S. oculata* is difficult to identify to the species level, many of the records from fishermen reports are often assigned only to genus level (Ferretti et al., 2015c), which contributes to the absence of occurrence records and putative reduction of areas of former habitat. Occurrence records in Madeira EEZ based on specimens observation have not been reported. Thus, this species is not considered in the total number of valid species in Madeira.

***Squatina squatina* (Linnaeus 1758)**

Common name: **Angelshark; Anjo**, Anjo-do-mar, Lixa, Peixe-anjo, Raia, Raia-bastarda, Raia-tubarão, Tubarão-raia, Viola, Violão.

References: mainland Portugal ([Carvalho, 1765](#); [Vandelli, 1771\(?\)](#); [Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, b, 1880](#); [Gervais and Boulart, 1877a](#); [Posser, 1880](#); [Girard, 1892](#); [Nobre, 1985a, 1903, 1935](#); [Almeida and Roquette, 1892](#); [Osório, 1896](#); [Vieira, 1897](#); [Bragança, 1904a](#); [Seabra, 1911](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Tortonese, 1979](#); [Compagno, 1984a](#); [Roux, 1984](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [Melendez and Macias, 2007](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#); [Morey et al., 2019\(?\)](#)) and Madeira (?) ([Fischer et al., 1981\(?\)](#); [Morey et al., 2019\(?\)](#)). According to Eschmeyer et al. (2019) the current geographic distribution of this species in the Eastern Atlantic is from Norway to Canary Islands and Western Sahara. The distribution map of this species reported by Fischer et al. (1981) included the area of Selvagens Islands (Madeira EEZ). However, Morey et al. (2019) considered it possibly extinct in Norway and North Sea and considered uncertain the presence of this species in Madeira and in mainland Portugal. Moreover, the catches of angelshark in the northeast Atlantic are now rare, both in surveys and commercially (ICES, 2015a). Occurrence records in Madeira based on specimens observation have not been yet reported. Thus, specimens are needed to confirm the presence of this species in the Selvagens Islands and this species is not considered in the total number of valid species in Madeira.

Order Torpediniformes

Family Torpedinidae

***Tetronarce nobiliana* (Bonaparte 1835)**

Common name: **Electric ray; Tremelga-negra**, Arraia, Dormideira, Orega, Raia, Raião, Tormetim, Tormentim, Torpedo, Tramelga, Treme-mão, Tremão, Tremedeira, Tremelga.

References: mainland Portugal ([Carvalho, 1765](#); [Osório de Castro, 1954, 1967](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Dinis, 1985](#); [Fischer et al., 1987](#); [Erzini et al., 2001a](#); [Borges et al., 2002, 2003](#); [Coelho et al., 2005](#); [Costa, 2007](#); [Fernandes et al., 2007](#); [Melendez and Macias, 2007](#); [Henriques et al., 2008](#); [Silva et al., 2009](#); [Silva, 2010, 2011](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Séret, 2016a](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bigelow and Schroeder, 1953](#); [Osório de Castro, 1967](#); [Krefft and Stehmann 1979a](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Trunov and Kukuev, 2005](#); [Menezes et al., 2006](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#); [ICES, 2017a](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1837, 1843](#); [NHMUK 1852.8.30.22, 1850-1853](#); [Günther, 1870](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [INIP, 1984c](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Christiansen et al., 2009](#); [Carneiro et al., 2014](#); [Séret, 2016a](#); [Biscoito et al., 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The syntype of

Torpedo hebetans Lowe 1838, synonym of *T. nobiliana*, is in the NHMUK collections (NHMUK 1852.8.30.22) caught off Madeira.

***Torpedo marmorata* Risso 1810**

Common name: **Marbled electric ray; Tremelga-marmoreada**, Dormideira, Orega, Raia, Raião, Tormentim, Tormetim, Tormeutin, Torpedo, Tramelga, Tremão, Tremedeira, Tremelga, Treme-mão. References: mainland Portugal (de Brito Capello, 1869a, b, 1880; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1894, 1895a, 1904, 1935; Osório, 1895; Vieira, 1897; Bragança, 1903; Seabra, 1911; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muus and Dahlström, 1978; Krefft and Stehmann 1979a; Fernández et al., 1981; Stehmann and Bürkel, 1984a; Dinis, 1985; Sanches, 1986; Fischer et al., 1987; Erzini et al., 2001a; WWF, 2001; Borges et al., 2002; Palma et al., 2003; Gonçalves et al., 2004; Moura, 2004; Coelho et al., 2005; ICES, 2005; Ribeiro et al., 2006; Costa, 2007; Fernandes et al., 2007; Melendez and Macias, 2007; Chaves, 2008; Henriques et al., 2008; Abecasis et al., 2009; Batista et al., 2009; Silva et al., 2009; Silva, 2010; Costa et al., 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; Costa, 2014; OCEANA, 2014; Séret, 2016a; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1843, 1844b; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Krefft and Stehmann 1979a; Stehmann and Bürkel, 1984a; Sanches, 1986; Lloris et al., 1991; Araújo et al., 2005; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; POGIS, 2016; Séret, 2016a; Bischoff et al., 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019). The whereabouts of the syntype of *Torpedo picta* Lowe 1843, synonym of *T. marmorata*, caught off Madeira are unknown. One specimen is in the MMF collections (MMF 020632, 1964).

***Torpedo torpedo* (Linnaeus 1758)**

Common name: **Common torpedo; Tremelga-de-olhos**, Dormideira, Orega, Raia, Raião, Tormentim, Tormetim, Torpedo, Tramelga, Tremão, Tremedeira, Tremedor, Tremelga, Treme-mão, Trememão, Trometim.

References: mainland Portugal (Carvalho, 1765; Vandelli, 1771; de Brito Capello, 1869a, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1935; Vieira, 1897; Seabra, 1911; Helling, 1940; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Stehmann 1979a; Stehmann and Bürkel, 1984a; Sanches, 1986; Fischer et al., 1987; Capapé and Desoutter, 1990; Cabral, 1999; Erzini et al., 2001a; WWF, 2001; Borges et al., 2002, 2003; Palma et al., 2003; Coelho et al., 2005; Costa, 2007; Melendez and Macias, 2007; Henriques et al., 2008; Neves et al., 2008; Ribeiro et al., 2008; Abecassis et al., 2009; Batista et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; Séret, 2016a; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Santos et al., 1997; Carneiro et al., 2014; Séret, 2016a; Bischoff et al., 2018) and Madeira (Lowe, 1837; Günther, 1870; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1949d; Nunes, 1953, 1974; Osório de Castro, 1967; Sanches, 1986; Carneiro et al., 2014; Séret, 2016a; Bischoff et al., 2018). One specimen caught off at the Portuguese continental slope in May 2011 is in the MB collections.

Order Rhinopristiformes

Family Rhinobatidae

***Rhinobatos rhinobatos* (Linnaeus 1758)**

Common name: **Common guitarfish; Viola**, Cassapai, Facinho-de-raia, Focinho-da-raia, Focinho-de-raia, Guitarra, Peixe-viola, Rabeca, Raia, Raia-tubarão, Rebeca, Tubarão-raia, Violão.

References: mainland Portugal (Vandelli, 1771; de Brito Capello, 1869a, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1897, 1901; Nobre, 1903, 1904, 1935; Seabra, 1911; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Stehmann, 1979a; Fischer et al., 1981, 1987; McEachran and Capapé, 1984a; Sanches, 1986; Melendez and Macias, 2007; Ebert and Stehmann, 2013; Carneiro et al., 2014; Bradai and Soldo, 2015; Séret, 2016a; Newell, 2017(?); Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (†) (Osório de Castro, 1967; Notarbartolo di Sciara et al., 2007) and Madeira (†) (Osório de Castro, 1967; Notarbartolo di Sciara et al., 2007).

According to Eschmeyer et al. (2019) the current geographic distribution of this species is the

Mediterranean Sea and, in the Eastern Atlantic, from Bay of Biscay to Angola. Information regarding this species in the Eastern Atlantic is very limited and the only historical records of *R. rhinobatos* in the fishing grounds of the Azores and Madeira were indicated by Osório de Castro (1967). Notarbartolo di Sciara *et al.* (2007) reported the occurrence of this species offshore around the archipelagos of the Azores and Madeira, although its absence during research surveys and from commercial landings suggest that it is now extremely rare or even locally extinct. Thus, the occurrence of this species in the Azores and Madeira region needs further documentation and is not considered in the total number of valid species. It is a very rare species or even extinct in mainland Portugal. According to Newell (2017) *R. rhinobatos* and *Glaucostegus cemiculus* (Geoffroy St. Hilaire 1817) are not reported in the ICES DATRAS, which is a database of 45 years' worth of survey data including the Atlantic coasts of France, Spain, and Portugal, indicating that they have likely been historically rare in the North of the Strait of Gibraltar.

Family Glaucostegidae

***Glaucostegus cemiculus* (Geoffroy St. Hilaire 1817)**

Common name: **Blackchin guitarfish; Viola-barba-negra**, Guitarra, Peixe-viola, Rabeca, Raia, Rebeca.
References: mainland Portugal ([Krefft and Stehmann 1979a](#); [Fischer et al., 1981, 1987](#); [McEachran and Capapé, 1984a](#); [Robins et al., 1991](#); [Melendez and Macias, 2007](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Soldo and Bradai, 2015](#); [Séret, 2016a](#); [Newell, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#); [Kyne and Jabado, 2019](#)). It is a very rare species or even extinct in mainland Portugal [see remarks on *Rhinobatos rhinobatos* (Linnaeus 1758)].

Family Pristidae

***Pristis pristis* (Linnaeus 1758)**

Common name: **Common sawfish; Espadarte-serra**, Espadarte, Peixe-serra, Sarra, Serra, Tubarão-serra.
References: mainland Portugal (†) ([Carvalho, 1765](#); [Barboza du Bocage and de Brito Capello, 1866](#); [de Brito Capello, 1869a, 1880](#); [Posser, 1880](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Stehmann, 1979a](#); [Fischer et al., 1981, 1987](#); [Stehmann and Bürkel, 1984b](#); [Sanches, 1986](#); [Stehmann, 1990a](#); [Serena, 2005](#); [Melendez and Macias, 2007](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#) [formerly]) and Madeira (†) ([Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Fischer et al., 1981](#); [Stehmann and Bürkel, 1984b](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Séret, 2016a](#); [Biscoito et al., 2018\(?\)](#); [Eschmeyer et al., 2019](#) [formerly]; [Froese and Pauly, 2019](#) [formerly]). Apparently it is an extinct species. The last known fishing record in mainland Portugal dates between 1986 and 2001 (Correia and Smith, 2003). Thus, this species is not considered in the total number of valid species.

Order Rajiformes

Family Rajidae

***Amblyraja jensenii* (Bigelow & Schroeder 1950)**

Common name: **Shorttail skate; Raia-de-cauda-curta**.
References: Azores (PECS) ([Ebert and Stehmann, 2013](#); [Porteiro et al., 2017](#)).

***Amblyraja radiata* (Donovan 1808)**

Common name: **Starry ray; Raia-repregada**, Arraia, Rabo-azedo, Raia, Raia-radiada, Repregada, Repregado.
References: mainland Portugal ([Nobre, 1895c, 1935](#); [Osório, 1895](#); [Vieira, 1897](#); [Seabra, 1911](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [INIP, 1981b](#); [Sanches, 1986](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#)) and Azores (PECS) (?) ([TCWC 3450.01, 1960\(?\)](#); [Carneiro et al., 2014](#)). According to Kulka *et al.* (2015) some reports of this species in the southern North Sea are likely to be inaccurate, owing to confusion with *Raja clavata* Linnaeus 1758. The specimen preserved in the Biodiversity Research and Teaching Collections of the Texas A&M University (TCWC 3450.01, 1960) has the following geographic coordinates: 38°28.0'N, 39°28.0'W. However, the geographic coordinates of the trawler (37°37.0'N, 74°14.0'W) are inconsistent with those displayed for the occurrence record and it is

recognized in the Integrated Digitized Biocollections (iDigBio) that this record requires verification. Thus, this species is not considered in the total number of valid species for the Azores.

Dipturus batis (Linnaeus 1758)

Common name: **Blue skate; Raia-oirega**, Airoga, Arrai, Arraia, Bico-doce, Cirega, Eiriogo, Eiroga, Eiroga-preta, Oirega, Orega, Ourega, Pigareva, Preta, Rabo-azedo, Raia, Raia-airoga, Raia-aurega, Raia-manteiga; Raia-ouregaa, Raia-preta, Tintureira, Urega.

References: mainland Portugal ([Vandelli, 1771](#); [de Brito Capello, 1869a, 1880](#); [Vieira, 1897](#); [Seabra, 1911](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Muus and Dahlström, 1978](#); [Stehmann, 1979](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [Melendez and Macias, 2007](#); [Silva, 2010](#); [Costa et al., 2012](#); [Ebert and Stehmann, 2013\(?\)](#); [Christiansen et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Dias et al., 1976](#); [Silva et al., 1995](#); [Menezes, 1996, 2014b](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Naylor et al., 2012](#); [Ebert and Stehmann, 2013\(?\)](#); [Menezes and Giacomello, 2013](#); [Menezes, 2014](#); [Biscoito et al., 2018](#)) and Madeira ([Günther, 1870\(?\)](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Norman, 1935a](#); [Fowler, 1936a\(?\)](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Stehmann, 1979](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Ebert and Stehmann, 2013\(?\)](#); [McEachran, 2016](#); [Biscoito et al., 2018\(?\)](#); [Froese and Pauly, 2019](#)). Froese and Pauly (2019) questioned the taxonomic validity of the species *Raja macrorhynchus* Rafinesque 1810 and indicated it as a synonym of *D. batis*. Biscoito et al. (2018) re-evaluated the specimen (NHMUK 1860.8.30.7, 1854-1861) previously identified as *Raja macrorhynchus* Rafinesque 1810 by Günther and identified it as *Dipturus intermedius* (Parnell 1837). According to Weigmann (2016) and Das and Afonso (2017) *D. batis* is treated as a composite species comprising a small (*D. cf. flossada*) and a large (*D. cf. intermedia*) species as referred by Iglesias et al. (2010). The geographic distributions of *D. batis* and *D. intermedius* present some overlapping (Eschmeyer et al., 2019) in the area considered in this study. The occurrence records of this species in mainland Portugal, Azores and Madeira don't allow unequivocally say if it is *D. batis* or *D. intermedius*. Further investigation is needed to confirm its presence in these three areas.

Dipturus intermedius (Parnell 1837)

Common name: Flapper Skate; **Raia-delgada**.

References: mainland Portugal ([Ebert and Stehmann, 2013\(?\)](#)) and Azores ([Ebert and Stehmann, 2013\(?\)](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018\(?\)](#)) and Madeira ([NHMUK 1860.8.30.7, 1854-1861](#); [Günther, 1870](#); [Fowler, 1936a](#); [Ebert and Stehmann, 2013](#); [Biscoito et al., 2018](#)). This species was recently resurrected from the synonymy of *Dipturus batis* (Linnaeus, 1758) (Last et al., 2016a). According to Eschmeyer et al. (2019) the current geographic distribution of this species is the northeast Atlantic Ocean. The specimen NHMUK 1860.8.30.7 (1854-1861) currently in The Natural History Museum (London) is registered as *Raja macrorhynchus* Rafinesque, 1810. The current status of *Raja macrorhynchus* Rafinesque 1810 is uncertain as *Dipturus batis* (Linnaeus 1758) as referred by Eschmeyer et al. (2019). However, the specimen NHMUK 1860.8.30.7 was identified as *D. intermedius* as mentioned by Biscoito et al. (2018). According to Ebert and Stehmann (2013) the occurrence of this species in mainland Portugal and in the Azores is possible and Biscoito et al. (2018) considered it most probable in the Azores. However, the occurrence records of this species in mainland Portugal and Azores don't allow unequivocally say if it is *D. batis* or *D. intermedius*. Thus, specimens are needed to confirm its presence in the mainland Portugal and Azores and it is not considered in the total number of valid species in these areas. One specimen is in the MMF collections (MMF 2869, 1943).

Dipturus nidarosiensis (Storm 1881)

Common name: **Norwegian skate; Raia-norueguesa**.

References: mainland Portugal ([Eschmeyer et al., 2019](#); [Geraci et al., 2019](#)) and Madeira ([Carbonara et al., 2019](#)). Its a very rare species and its known as the only endemic species of *Dipturus* to the Northeastern Atlantic area (Cannas et al., 2010).

Dipturus oxyrinchus (Linnaeus 1758)

Common name: **Longnosed skate; Raia-bicuda**, Arado, Arrai, Arraia, Bico, Corva, Curva, Ferreiro, Rabo-azedo, Raia, Raia-corva, Raia-curva, Raia-longa.

References: mainland Portugal▲ (Vandelli, 1771; de Brito Capello, 1869a, c, 1880; Nobre, 1894, 1903, 1904, 1935; Vieira, 1897; Seabra, 1911; Helling, 1940; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann, 1979; Fernández et al., 1981; Martins, 1981; INIP, 1982a, b; Stehmann and Bürkel, 1984a; Sanches, 1986; Fischer et al., 1987; Saldanha et al., 1995; Erzini et al., 2001a; Monteiro et al., 2001; Borges et al., 2002, 2003; Moura, 2004; Coelho et al., 2005; ICES, 2005; Costa, 2007; Melendez and Macias, 2007; Silva, 2008, 2010, 2011; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; OCEANA, 2014; Silva and Borges, 2014; McEachran, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (INIP, 1985c; Sanches, 1986; Dias, 1991; Arruda, 1997; Santos et al., 1997; Fock et al., 2002 (PECS); Menezes, 2003; Menezes et al., 2006; Pakhorukov, 2008 (PECS); Porteiro et al., 2010; Menezes and Giacomello, 2013; Carneiro et al., 2014; Das and Afonso, 2017; Bischoff et al., 2018) and Madeira (Lowe, 1843, 1849; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann and Bürkel, 1984a; Sanches, 1986; Lloris et al., 1991; Melendez and Macias, 2007; Wirtz et al., 2008; Ebert and Stehmann, 2013; Carneiro et al., 2014; McEachran, 2016; Bischoff et al., 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019).

Dipturus sp. Stehmann: probably *Dipturus nidarosiensis* (Storm 1881)

Common name: **Norwegian skate; Raia-norueguesa**.

References: mainland Portugal (?) (Ebert and Stehmann, 2013(?)). *D. nidarosiensis* is a benthic species occurring along continental and insular slopes from 200 to 1000 m (McEachran, 2016) from southern Norway to northern Mauritania (McEachran, 2016; Froese and Pauly, 2019), and for Eschmeyer et al. (2019) it distributes from Iceland to South Africa. Additionally, Ramírez-Amaro et al. (2017), Geraci et al. (2019) and Eschmeyer et al. (2019) indicated its presence in the western and central Mediterranean Sea. Ebert and Stehmann (2013) referred four specimens taken by the Portuguese RV Capricórnio in 1117 m depth at 36°29.8'N, 8°08.0'W (off the Algarve) and provisionally identified as *Dipturus* sp. For Last et al. (2016a) *D. nidarosiensis* is probably a composite species in the eastern-north Atlantic as also mentioned by Ebert and Stehmann (2013) and Weigmann (2016). Thus, since there is no confirmation of the identification of the specimens caught off in the Algarve, the species *D. nidarosiensis* is not considered in the total number of valid species in mainland Portugal.

Leucoraja circularis (Couch 1838)

Common name: **Sandy ray; Raia-de-São-Pedro**, Arraia, Rabo-azedo, Raia, Raia-de-quatro-olhos, Raia-santiaga, São-Pedro.

References: mainland Portugal▲ (Vieira, 1897; Bragança, 1903; Seabra, 1911; Murray and Hjort 1912; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Stehmann, 1979; Stehmann and Bürkel, 1984a; Sanches, 1986; Fischer et al., 1987; Saldanha et al., 1995; Moura, 2004; ICES, 2005; Fernandes et al., 2007; Melendez and Macias, 2007; Silva, 2008, 2010, 2011; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; Silva and Borges, 2014; McEachran, 2016; Froese and Pauly, 2019) and Madeira (NHMUK 1860.8.30.3, 1854-1861; Gunther, 1870; Belloc, 1934; Noronha and Sarmento, 1934; Nobre, 1935; Osório de Castro, 1967; Bischoff et al., 2018).

Leucoraja fullonica (Linnaeus 1758)

Common name: **Shagreen ray; Raia-pregada**, Arraia, Cardadora, Pregada, Rabo-azedo, Raia, Raia-cardadora, Raia-de-picôs, Raia-espinhosa, Raia-legítima, Raia-rajada, Raia-verdadeira, Rajada.

References: mainland Portugal (Vandelli, 1771; de Brito Capello, 1869a, 1880; Seabra, 1911; Nobre, 1935; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann, 1979, 1990b; Stehmann and Bürkel, 1984a; Sanches, 1986; Fischer et al., 1987; Melendez and Macias, 2007; Henriques et al., 2008; Ebert and Stehmann, 2013; Carneiro et al., 2014; ICES, 2017a; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Dias et al., 1976, 1990; INIP, 1985b; Sanches, 1986; Dias, 1991; Silva et al., 1995; Menezes, 1996; Arruda, 1997; Santos et al., 1997; Menezes et al., 2006; Rosa et al., 2006; Porteiro et al., 2010; Barreiros and Gadig, 2011; Naylor et al., 2012; Carneiro et al., 2014; Das and Afonso, 2017; ICES, 2017a; Bischoff et al., 2018) and Madeira (Günther, 1870; Garman, 1913; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann, 1979; Stehmann and Bürkel, 1984a; Sanches,

1986; Lloris *et al.*, 1991; Melendez and Macias, 2007; Wirtz *et al.*, 2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Biscoito *et al.*, 2018; Froese and Pauly, 2019; Saad and Alkusairy, 2019).

***Leucoraja naevus* (Müller & Henle 1841)**

Common name: **Cuckoo ray; Raia-de-dois-olhos**, Arraia, Rabo-azedo, Raia-de-quatro-olhos.

References: mainland Portugal▲ (de Brito Capello, 1869a, 1880; Nobre, 1935; Helling, 1940; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Marques and Ré, 1978; Stehmann, 1979; Fernández *et al.*, 1981; Martins, 1981; INIP, 1981a, 1982b; Dinis and Marecos, 1984; Stehmann, and Bürkel, 1984a; Dinis, 1985; Sanches, 1986; Fischer *et al.*, 1987; Saldanha *et al.*, 1995; Erzini *et al.*, 2001a; Borges *et al.*, 2002; Machado *et al.*, 2004; Moura, 2004; Coelho *et al.*, 2005; Farias, 2005; ICES, 2005; Costa, 2007; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Henriques *et al.*, 2008; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Costa *et al.*, 2012; Maia *et al.*, 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Gordon *et al.*, 2016; McEachran, 2016; Ryland and Lancaster, 2017; ICES, 2017a; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Malacoraja kreffti* (Stehmann 1978)**

Common name: **Krefft's ray; Raia-de-Krefft**.

References: Azores (?) (Ebert and Stehmann, 2013(?)). According to Ebert and Stehmann (2013) and Eschmeyer *et al.* (2019) the current geographic distribution of this species is the eastern-north Atlantic. From off south-western Iceland, the Porcupine Seabight and the Northern Mid-Atlantic Ridge, and may also extend South to the Mid-Atlantic Ridge (Ebert and Stehmann, 2013; Stehmann and Orlov, 2015). Thus, the occurrence of this species in the Azores Archipelago is questionable and shall not be considered in the total number of valid species for the Azorean ichthyofauna.

***Neoraja iberica* Stehmann, Séret, Costa & Baro 2008**

Common name: Iberian pygmy skate; **Raia-pigmeia-ibérica**, Raia-ibérica-anã.

References: mainland Portugal (Stehmann *et al.*, 2008; Silva, 2011; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype of this species is in the MB collections (MB06-004869, 2006) and was collected at the Algarve coast (Portugal).

***Raja asterias* Delaroche 1809**

Common name: **Mediterranean starry ray; Raia-pintada**, Arraia, Mona, Pintada, Rabo-azedo, Raia, Raia-estrelada.

References: mainland Portugal (de Brito Capello, 1869a, b, 1880; Girard, 1892; Osório, 1896; Seabra, 1911; Helling, 1940; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann and Bürkel, 1984a; Sanches, 1986; Fischer *et al.*, 1987; Martins *et al.*, 1992; Erzini *et al.*, 2001a; Coelho *et al.*, 2005; Henriques *et al.*, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Günther, 1870).

***Raja brachyura* Lafont 1873**

Common name: **Blonde ray; Raia-ponteada**, Arraia, Mona, Pintada, Pontuada, Rabo-azedo, Raia, Raia-pintada, Raia-pontuada.

References: mainland Portugal▲ (Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Marques and Ré, 1978; Stehmann, 1979; Fernández *et al.*, 1981; Martins, 1981; INIP, 1981a, 1982b; Borges, 1984; Dinis and Marecos, 1984; Stehmann and Bürkel, 1984a; Dinis, 1985; Sanches, 1986; Fischer *et al.*, 1987; Erzini *et al.*, 2001a; Machado *et al.*, 2004; Moura, 2004; Coelho *et al.*, 2005; Farias, 2005; ICES, 2005; Sousa *et al.*, 2005; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Chaves, 2008, 2009; Batista *et al.*, 2009; Gonçalves *et al.*, 2010; Costa *et al.*, 2012; Pina-Rodrigues, 2012; McEachran, 2016; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Gordon *et al.*, 2016; Froese and Pauly, 2019), Azores (Fowler, 1936a; Collins, 1954; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Carneiro *et al.*, 2014; Das and Afonso, 2017; ICES, 2017a; Biscoito *et al.*, 2018) and Madeira (Günther, 1870; Stehmann, 1979; Stehmann & Bürkel 1984a; Sanches, 1986; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; McEachran 2016; POGID, 2016; Biscoito *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Raja clavata Linnaeus 1758

Common name: **Thornback ray; Raia-lenga**, Arraia, Brocheada, Langa, Lenga, Pinta, Pintada, Pregada, Rabo-azedo, Raia, Raia-brocheada, Raia-langa, Raia-pinta, Raia-pintada, Raia-pregada.

References: mainland Portugal[▲] ([Vandelli, 1771](#); [de Brito Capello, 1869a, 1880](#); [Baldaque da Silva, 1891](#); [Nobre, 1895a, 1935](#); [Vieira, 1897](#); [Seabra, 1911](#); [Murray and Hjort, 1912](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Marques and Ré, 1978](#); [Muus and Dahlström, 1978](#); [Stehmann, 1979](#); [INIP, 1981a, b, 1982a, b](#); [Fernández et al., 1981](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Stehmann and Bürkel, 1984a](#); [Dinis, 1985](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [Martins et al., 1992](#); [Saldanha et al., 1995](#); [Erzini et al., 2001a, b](#); [Gomes et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002b](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Machado et al., 2004](#); [Mendes et al., 2004](#); [Moura, 2004](#); [Coelho et al., 2005](#); [ICES, 2005](#); [Serra-Pereira, 2005](#); [Sousa et al., 2005](#); [Costa, 2007](#); [Fernandes et al., 2007](#); [Melendez and Macias, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [França et al., 2011](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Ebert and Stehmann, 2013](#); [Figueiredo and Serra-Pereira, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Gordon et al., 2016](#); [McEachran, 2016](#); [Figueiredo et al., 2017](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Belloc, 1934](#); [Fowler, 1936a](#); [Ribeiro, 1936](#); [Ferreira, 1939](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [INIP, 1985a, b, c](#); [Dias and Monteiro, 1986](#); [Sanches, 1986](#); [Dias et al., 1990](#); [Dias, 1991](#); [Dias and Cascalho, 1991](#); [Arruda, 1992, 1997](#); [Wirtz, 1994](#); [Silva and Menezes, 1995a, b](#); [Silva, 1995](#); [Menezes, 1996](#); [Santos et al., 1997](#); [Morato-Gomes et al., 1998](#); [Kukuev, 2002 \(PECS\)](#); [Fowler et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Naylor et al., 2012](#); [Pham et al., 2013](#); [Carneiro et al., 2014](#); [McEachran, 2016](#); [Torres et al., 2016](#); [Das and Afonso, 2017](#); [ICES, 2017a](#); [Biscoito et al., 2018](#); [Fauconnet et al., 2019](#)) and Madeira ([Günther, 1870](#); [Garman, 1913](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949d, 1976](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008\(?\)](#); [Carneiro et al., 2014](#); [McEachran, 2016](#); [POGID, 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019\(?\)](#)). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22313, 1967).

Raja maderensis Lowe 1839

Common name: **Madeiran ray; Raia-da-Madeira**, Arraia, Raia.

References: mainland Portugal ([Bragança, 1902](#); [Osório de Castro, 1954](#); [Maul, 1976](#); [SDSC 22955, 1967](#); [WWF, 2001](#); [Abecasis et al., 2009](#); [Carneiro et al., 2014](#)), Azores ([Hilgendorf, 1888a](#); [Regan, 1903a](#); [Sampaio, 1904](#); [Fowler, 1936](#); [Ribeiro, 1936](#); [Ferreira, 1939](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Ehrich, 1977 \(PECS\)\(?\)](#); [Stehmann, 1979](#); [Sanches, 1986](#); [Lloris et al., 1991\(?\)](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Fock et al., 2002 \(PECS\)](#); [Kukuev, 2002 \(PECS\)](#); [Pakhorukov, 2008 \(PECS\)](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013 \(PECS\)](#); [Carneiro et al., 2014](#); [Ball et al., 2016](#); [Weigmann, 2016](#); [Das and Afonso, 2017\(?\)](#); [Biscoito et al., 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019\(?\)](#)) and Madeira ([Lowe, 1837, 1839](#); [Duméril, 1865](#); [Günther, 1870](#); [de Brito Capello, 1872](#); [Garman, 1913](#); [Belloc, 1934](#); [Noronha and Sarmento 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949d, 1976](#); [Nunes, 1953, 1974](#); [Osório de Castro, 1954, 1967](#); [Stehmann, 1979](#); [INIP, 1984b, c, 1985c](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Pakhorukov, 2008 \(PECS\)](#); [Wirtz et al., 2008](#); [Christiansen et al., 2009\(?\)](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Ball et al., 2016](#); [Last et al., 2016b](#); [McEachran, 2016](#); [POGID, 2016](#); [Biscoito et al., 2017, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The whereabouts of the syntype of this species caught off Madeira are unknown. The record of a female of *R. maderensis* in the dataset of the Geographically tagged INSDC sequences, published by European Molecular Biology Laboratory (EMBL) and registered in the Global Biodiversity Information Facility indicates its occurrence in mainland Portugal (Occurrence ID - HM043185) (EMBL, 2014). In fact, it was a specimen landed in Peniche fishing harbour and caught off in the Ampère Seamount in 2007 (36°28.0'N, 11°35'W, Madeira EEZ) (Serra-Pereira et al., 2011). A second specimen of this species was caught off in the Gorringe Ridge (36°29.9'N, 11°33.0'W, mainland Portugal EEZ). According to Weigmann (2016) the isolated population at the Azores Archipelago is morphologically very similar to *R. maderensis* from off Madeira, but shows constant differences in colouration, which are either a variant of *R. clavata* or possibly a sub-species of *R. maderensis* (Stehmann and Bürkel, 1984a). Ball et al. (2016)

mention the presence of a morphotype of *R. clavata* in the Azores. Das and Afonso (2017) considered the records of *R. maderensis* in the Azores as *R. clavata*. One specimen caught off at the Gattysburg Seamount is in the MMF collections (MMF 22318, 1967).

***Raja microocellata* Montagu 1818**

Common name: **Small-eyed ray; Raia-zimbreira**, Arraia, Arrai, Rabo-azedo, Raia, Raia-branca, Raia-branca-legítima, Zimbreira, Zimbreiro.

References: mainland Portugal[▲] (de Brito Capello, 1869c, 1880; Vieira, 1897; Seabra, 1911; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann, 1979; Fernández et al., 1981; Dinis and Marecos, 1984; Stehmann and Bürkel, 1984a; Sanches, 1986; Machado et al., 2004; Moura, 2004; Henriques et al., 2008; Ebert and Stehmann, 2013; Carneiro et al., 2014; Ellis and Walls, 2015; Gordon et al., 2016; McEachran, 2016; Ryland and Lancaster, 2017; Baptista et al., 2019; Froese and Pauly, 2019) and Azores (?) (Dias et al., 1976; Arruda, 1997; Santos et al., 1997(?); Carneiro et al., 2014(?)). According to Stehmann and Bürkel (1984a) and Eschmeyer et al. (2019) the current geographic distribution of this species is restricted primarily to the coasts of eastern-north Atlantic, from the British Isles southwards to Gibraltar and north-western Africa (Western Sahara and Mauritania). There are no records of specimens caught off in the Azores and according to Santos et al. (1997) further documentation is needed. Thus, this species is not considered present in the Azores.

***Raja miraletus* Linnaeus 1758**

Common name: **Brown ray; Raia-de-quatro-olhos**, Arraia, Patelo, Rabo-azedo, Raia, Raia-azul, Raia-branca, Raia-de-espelhos, Raia-pintada, Raza.

References: mainland Portugal[▲] (Vandelli, 1771; de Brito Capello, 1869a, 1880; Nobre, 1903, 1935; Seabra, 1911; Helling, 1940; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Stehmann, 1979; INIP, 1981a, b, 1982a, b; Fernández et al., 1981; Martins, 1981; Dinis and Marecos, 1984; Stehmann and Bürkel, 1984a; Dinis, 1985; Sanches, 1986; Fischer et al., 1987; Saldanha et al., 1995; Erzini et al., 2001a; Borges et al., 2002, 2003; Machado et al., 2004; Mendes et al., 2004; Coelho et al., 2005; ICES, 2005; Costa, 2007; Fernandes et al., 2007; Melendez and Macias, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Batista et al., 2009; Silva et al., 2009; Silva, 2010, 2011; Costa et al., 2012; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Last and Séret, 2016; McEachran, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Günther, 1870; Garman, 1913; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Stehmann, 1979; Stehmann and Bürkel, 1984a; Sanches, 1986; Lloris et al., 1991; Melendez and Macias, 2007; Wirtz et al., 2008(?); Ebert and Stehmann, 2013; Carneiro et al., 2014; Last and Séret, 2016; McEachran, 2016; POGID, 2016; POGIS, 2016; Biscoito et al., 2018(?); Eschmeyer et al., 2019; Froese and Pauly, 2019). According to Maul (1976) the presence of this species in Madeiran waters seems unlikely. Wirtz et al. (2008) and Biscoito et al. (2018) consider doubtful the occurrence of *R. miraletus* in Madeira.

***Raja montagui* Fowler 1910**

Common name: **Mediterranean starry ray; Raia-manchada**, Arraia, Mona, Patelo, Pintada, Rabo-azedo, Raia, Raião, Raia-pintada, Zimbreiro.

References: mainland Portugal[▲] (de Brito Capello, 1880; Bragança, 1902; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Marques and Ré, 1978; Stehmann, 1979; Martins, 1981; INIP, 1982b; Borges, 1984; Dinis and Marecos, 1984; Stehmann and Bürkel, 1984a; Dinis, 1985; Sanches, 1986; Fischer et al., 1987; Saldanha et al., 1995; Borges et al., 2002; Prista et al., 2003; Machado et al., 2004; Moura, 2004; ICES, 2005; Serra-Pereira, 2005; Fernandes et al., 2007; Melendez and Macias, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Batista et al., 2009; Silva et al., 2009; Gonçalves et al., 2010; Silva, 2010, 2011; Costa et al., 2012; Pina-Rodrigues, 2012; Ebert and Stehmann, 2013; Carneiro et al., 2014; Costa, 2014; Gordon et al., 2016; McEachran, 2016; Ryland and Lancaster, 2017; ICES, 2017a; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967) and Madeira (Günther, 1870; Noronha and Sarmento, 1934; Nobre, 1935; Osório de Castro, 1954, 1967; Maul, 1976; Ellis et al., 2015b; Biscoito et al., 2018). The only record of *R. montagui* in the fishing grounds of the Azores was indicated by Osório de Castro (1967). Since then no other author described the presence of this species in the Azores. Thus, the occurrence of this species in the Azores Archipelago is questionable and is not considered in the total number of valid species for

the Azorean ichthyofauna. The distribution map of this species reported by Ellis *et al.* (2015b) included the area of Selvagens Islands (Madeira EEZ). Two specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22308, 1967 and 22314, 1967).

***Raja undulata* Lacepède 1802**

Common name: **Undulate ray; Raia-curva**, Arraia, Arraia-corva, Arraia-curva, Corva, Curva, Galega, Mosaica, Rabo-azedo, Raia-da-areia, Raia-corva, Raia-do-rio, Raia-galega, Raia-moira, Raia-mosaica, Raia-mosaico, Raia-preta, Raia-riscada.

References: mainland Portugal[▲] ([de Brito Capello, 1869a, b, c, 1880](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Bragança, 1902](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Stehmann, 1979](#); [Martins, 1981](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Coelho and Erzini, 2002, 2006](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Machado et al., 2004](#); [Coelho et al., 2005](#); [ICES, 2005](#); [Ribeiro et al., 2006, 2008](#); [Costa, 2007](#); [Henriques et al., 2007, 2008](#); [Melendez and Macias, 2007](#); [Moura et al., 2007, 2008c](#); [Neves et al., 2008](#); [Batista et al., 2009](#); [Chaves, 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Ebert and Stehmann, 2013](#); [Serra-Pereira et al., 2013, 2015](#); [Carneiro et al., 2014](#); [Figueiredo et al., 2014, 2015](#); [McEachran, 2016](#); [Ryland and Lancaster, 2017](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira (?) ([Günther, 1870](#); [Fowler, 1936a](#); [Osório de Castro, 1967](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *R. undulata* in the Eastern Atlantic is confined to British Isles and south-western North Sea to Senegal, mainly along coastline (Ebert and Stehmann, 2013). The specimen identified by Günther (1870) currently deposited in The Natural History Museum (London) was later identified as *Raja brachyura* Lafont 1873. Thus, this species is not considered in the total number of valid species for Madeira region.

***Rajella bathyphila* (Holt & Byrne 1908)**

Common name: **Deep-water ray; Raia-da-fundura**.

References: mainland Portugal ([Dulvy and Walls, 2015](#)) and Azores ([Stehmann, 1979](#); [Arruda, 1997](#); [Carneiro et al., 2014](#); [ICES, 2017a](#); [Almeida and Biscoito, 2019](#)).

***Rajella bigelowi* (Stehmann 1978)**

Common name: **Bigelow's ray; Raia-de-Bigelow**, Arraia, Raia.

References: mainland Portugal ([Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)) and Azores ([MOM-IC-0091-3717, 1911](#); [Roule, 1912, 1919](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Stehmann, 1971, 1978, 1990b, 1995](#); [Hureau and Monod, 1979](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Moore et al. 2003](#); [Orlov et al., 2006](#); [Porteiro et al., 2010, 2017](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#); [Quéro et al., 2017](#); [Froese and Pauly, 2019](#)). There are bibliographical references to the occurrence of *Raja ackleyi* Garman 1881 [currently *Rostroraja ackleyi* (Garman 1881)] in the Azores (Roule, 1912; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967). However, Stehmann (1978) re-evaluated the Roule specimen and renamed it as paratype of *R. bigelowi*.

***Rajella fyllae* (Lütken 1887)**

Common name: **Round ray; Raia-redonda**.

References: mainland Portugal ([Fernandes et al., 2007](#); [Landi et al., 2010](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *R. fyllae* is the North Atlantic and Arctic Seas to Nova Scotia and to France. This is the most southern occurrence record out of the current geographic distribution area and it is an uncommon catch of Portuguese bottom trawl fleets, during 2004, but the specimen was discarded on-board. One specimen is in the MB collections (MB06-005121).

***Rajella linteia* (Fries 1838)**

Common name: **Sailray; Raia-nevoeira**, Arraia, Nevoeira, Oirega, Orega, Ourega, Peixe-areia, Rabo-azedo, Raia, Teiroga.

References: mainland Portugal ([de Brito Capello, 1869a, 1880](#); [Baldaque da Silva, 1891](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Sanches, 1986](#)).

Rostroraja alba (Lacepède 1803)

Common name: **White skate; Raia-tairoga**, Arraia, Neveira, Nevoeira, Oirega, Oregá, Ourega, Raia, Raia-branca, Raia-teiroga, Tairoga, Teiroga.

References: mainland Portugal ([de Brito Capello, 1880](#); [Vieira, 1897](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Stehmann, 1979](#); [Borges, 1984](#); [Stehmann and Bürkel, 1984a](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [Moura, 2004](#); [ICES, 2005](#); [Melendez and Macias, 2007](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Ellis et al., 2015c](#); [Gordon et al., 2016](#); [McEachran, 2016](#); [ICES, 2017a](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967\(?\)](#); [Ellis et al., 2015c](#); [ICES, 2017a](#)) and Madeira ([Osório de Castro, 1967\(?\)](#); [Stehmann, 1979](#); [Hulley, 1986](#); [Lloris et al., 1991](#); [Ferreira et al., 2008](#); [Pakhorukov, 2008 \(PECS\)](#); [Carneiro et al., 2014](#); [Ellis et al., 2015c](#); [POGID, 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). Several specimens are in the MMF collections: MMF 22315, 1967 (set of two specimens caught at coast of mainland Portugal) and MMF 23650, 1984 and MMF 36215, 2004 (both caught off at Desertas Islands). The distribution map of this species reported by [Ellis et al. \(2015c\)](#) included the area of Selvagens Islands (Madeira EEZ).

Family Arhynchobatidae

Bathyraja pallida (Forster 1967)

Common name: **Pale ray; Raia-pálida**.

References: Azores ([MAR-ECO 16391, 2004 \(PECS\)](#); [Fossen et al., 2008](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#)).

Bathyraja richardsoni (Garrick 1961)

Common name: **Richardson's ray; Raia-de-Richardson**.

References: Azores ([Orlov et al., 2006](#); [Fossen et al., 2008](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 2010, 2017](#); [Menezes et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Stehmann and Farrell 2015](#); [Das and Afonso, 2017](#)).

Order Myliobatiformes

Family Dasyatidae

Bathytoshia lata (Garman 1880)

Common name: Brown stingray; **Uge-de-cardas**, Arraião, Raia, Raia-de-agulhão, Ratão, Uga, Uge, Uja, Usga.

References: mainland Portugal▲ ([Krefft and Stehmann, 1979b](#); [Fischer et al., 1981, 1987](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [Melendez and Macias, 2007](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Séret, 2016b](#); [Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967](#); [Serrão et al., 1997](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017\(?\)](#); [Biscoito et al., 2018](#)) and Madeira ([Günther, 1870](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft and Stehmann, 1979b](#); [Fischer et al., 1981](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [Lloris et al., 1991](#); [Melendez and Macias, 2007](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Last et al. 2016c](#); [Séret, 2016b](#); [Biscoito et al., 2018](#)). According to [Afonso et al. \(2013\)](#) the species *Dasyatis centroura* ([Mitchill 1815](#)), synonym of *Bathytoshia centroura* ([Mitchill 1815](#)), has been misidentified in the Azores as *Dasyatis pastinaca* ([Linnaeus 1758](#)). On the other hand, Eastern Atlantic populations of *B. centroura* refer to *B. lata* as mentioned by [Last et al. \(2016d\)](#).

Dasyatis pastinaca (Linnaeus 1758)

Common name: **Common stingray; Uge**, Arraia, Arraião, Chucho, Manteiga, Raia, Raia-de-agulhão, Raia-de-espinhos, Raia-manteiga, Ratão, Rato, Rato-do-mar, Ratona, Retomba, Uge-manteiga, Uge-vulgar, Uja, Ujo, Urce, Urcha, Urga, Urge, Urja, Urze, Usa, Usga.

References: mainland Portugal ([Vandelli, 1771](#); [de Brito Capello, 1869a, b, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1897](#); [Seabra, 1911](#); [Nobre, 1935, 1937](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Muus and Dahlström, 1978](#); [Krefft and Stehmann, 1979b](#); [Fischer et al., 1981, 1987](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [Melendez and Macias, 2007](#); [Ribeiro, 2008](#); [Batista et al., 2008](#); [Henriques et al.,](#)

2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Séret, 2016b; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Fowler, 1936a; Ferreira, 1939; Collins, 1954; Osório de Castro, 1967; Dias *et al.*, 1976; Harmelin and Harmelin-Vivien, 1979; McEachran and Capapé, 1984b; INIP, 1985a, c; Sanches, 1986; Barreiros 1990; Dias *et al.*, 1990; Nash *et al.*, 1990, 1991, 1994a; Patzner, 1990; Patzner *et al.*, 1990, 1992; Ré, 1990; Wirtz, 1990a, 1994; Dias, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Azevedo *et al.*, 1992; Santos *et al.*, 1994b, 1997; Azevedo, 1995; Gonçalves *et al.*, 1995; Santos and Nash, 1995; Menezes, 1996; Harmelin-Vivien *et al.*, 2001; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Afonso, 2016; Das and Afonso, 2017; ICES, 2017a; Bischoff *et al.*, 2018; Almeida and Bischoff, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837; Günther, 1870; de Velloso, 1892; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; INIP, 1982c, 1984b; McEachran and Capapé, 1984b; Augier, 1985; Sanches, 1986; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Quéro *et al.*, 2017; Bischoff *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Pteroplatytrygon violacea* (Bonaparte 1832)**

Common name: **Pelagic stingray; Uge-violeta**, Raia-pelágica, Ratão.

References: mainland Portugal (Santos *et al.*, 2002b; ICES, 2005; Fernandes *et al.*, 2007; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; ICES, 2017a), Azores (Pinho and Pereira, 1995a; Silva and Menezes, 1995b; Arruda, 1997; Santos *et al.*, 1997; Menezes *et al.*, 2006; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Das and Afonso, 2017; Bischoff *et al.*, 2018) and Madeira (Bischoff and Wirtz, 1994; Ellis 2007; Wirtz *et al.*, 2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bischoff *et al.*, 2018; Froese and Pauly, 2019).

***Taeniurus grabatus* (Geoffroy St.-Hilaire 1817)**

Common name: **Round stingray; Uge-redonda**, Ratão, Uge.

References: Azores (Wirtz, 1994; Saldanha, 1995; Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Carneiro *et al.*, 2014; Das and Afonso, 2017; Bischoff *et al.*, 2018) and Madeira (Bischoff and Wirtz, 1994; Mello, 2000; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Séret, 2016b; POGIS, 2016; Bischoff *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen is in the MMF collections (MMF 28784, 1997).

Family Gymnuridae

***Gymnura altavela* (Linnaeus 1758)**

Common name: **Spiny butterfly ray; Uge-manta**, Andorinha, Andorinha-do-mar, Avejão, Borboleta, Breamanta, Briamanta, Jaimanta, Jamanta, Raia, Raia-borboleta, Ratão, Uje-mata, Uge-manta, Urjamanta, Vela-latina.

References: mainland Portugal[▲] (Vandelli, 1771; de Brito Capello, 1880; Vieira, 1897; Seabra, 1911; Osório, 1917; Nobre, 1935; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Stehmann, 1979b; Fischer *et al.*, 1981, 1987; McEachran and Capapé, 1984b; Sanches, 1986; McEachran and Séret, 1990a; Melendez and Macias, 2007; Correia, 2009; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Séret, 2016b; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Vooren *et al.*, 2007; Walls *et al.*, 2015d) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1870; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Krefft and Stehmann, 1979b; Fischer *et al.*, 1981; McEachran and Capapé, 1984b; Sanches, 1986; McEachran and Séret, 1990a; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Séret, 2016b; Bischoff *et al.*, 2018; Froese and Pauly, 2019). One specimen is in the MMF collections (MMF 14040, 1958).

***Gymnura hirundo* (Lowe 1843)**

Common name: Madeira butterfly ray; **Uge-da-Madeira**, Andorinha, Andorinha-do-mar, Avejão, Jamanta, Jamanta-de-asas-redonda, Raia, Raia-borboleta, Raia-borboleta-da-Madeira, Ratão.

References: Azores (?) ([Osório de Castro, 1967](#)) and Madeira ([Lowe, 1843, 1844b, 1849](#); [Günther, 1870](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [McEachran and Séret, 1990a](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Biscoito et al., 2018\(?\)](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The only record of *G. hirundo* in the fishing grounds of the Azores was indicated by Osório de Castro (1967). The occurrence of this species in the Azores region needs further documentation. Thus, this species is not considered in the total number of valid species to the Azores. According to Eschmeyer et al. (2019) the current status of this species is valid but several authors questioned its validity.

Family Myliobatidae

***Aetomylaeus bovinus* (Geoffroy St. Hilaire 1817)**

Common name: **Bull ray**; **Ratão-bispo**, Bispo, Dromadeiro, Papa, Peixe-bispo, Peixe-raia, Peixe-ratão, Raia, Raia-ratão, Ratão.

References: mainland Portugal ([Vieira, 1897](#); [Seabra, 1911](#); [Nunes, 1953, 1974](#); [Osório de Castro, 1954, 1967](#); [Krefft and Stehmann, 1979b](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [McEachran and Séret, 1990a](#); [Erzini et al., 2001a](#); [Coelho et al., 2005](#); [Melendez and Macias, 2007](#); [Naylor et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Séret, 2016b](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1843-1860](#); [Günther, 1870](#); [Noronha and Sarmento, 1934](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1949](#); [Nunes, 1953, 1974](#); [Osório de Castro, 1954, 1967](#); [Krefft and Stehmann, 1979b](#); [Fischer et al., 1981](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [McEachran and Séret, 1990a](#); [Lloris et al., 1991](#); [Melendez and Macias, 2007](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGIS, 2016](#); [Séret, 2016b](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)).

***Myliobatis aquila* (Linnaeus 1758)**

Common name: **Common eagle ray**; **Ratão-água**, Águia-do-mar, Águia-marinha, Arrião, Arreião, Batão, Bispo, Chucho, Dormideiro, Peixe-bispo, Peixe-raia, Peixe-ratão, Raia, Raião, Raia-ratão, Rato, Ratão, Viúva.

References: mainland Portugal▲ ([Vandelli, 1771](#); [de Brito Capello, 1869a, b, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1894, 1903, 1935](#); [Vieira, 1897](#); [Seabra, 1911](#); [Helling, 1940](#); [Gonçalves, 1941](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Stehmann, 1979b](#); [INIP, 1981b](#); [Fischer et al., 1981, 1987](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [McEachran and Séret, 1990a](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Jorge et al., 2002](#); [Coelho et al., 2005](#); [ICES, 2005](#); [Henriques et al., 2007, 2008](#); [Neves et al., 2008](#); [Batista et al., 2009](#); [Correia, 2009](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Séret, 2016b](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a](#); [Roule, 1912, 1919](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Wheeler, 1978](#); [Harmelin and Harmelin-Vivien, 1979](#); [Krefft and Stehmann, 1979b](#); [Ré, 1979a, 1990](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [McEachran and Séret, 1990a](#); [Patzner et al., 1990, 1992](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Gonçalves et al., 1995](#); [Santos et al., 1997](#); [Harmelin-Vivien et al., 2001](#); [Cardigos et al., 2005](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1837, 1843-1860](#); [Günther, 1870](#); [Belloc, 1934](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft and Stehmann, 1979b](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [McEachran and Séret, 1990a](#); [Lloris et al., 1991](#); [Falcón et al., 2000](#); [Mello, 2000](#); [Araújo et al., 2005](#); [Melendez and Macias, 2007](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Séret, 2016b](#); [POGID, 2016](#); [POGIS, 2016](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)).

Family Rhinopteridae

Rhinoptera marginata (Geoffroy St. Hilaire 1817)

Common name: **Lusitanian cownose ray; Gavião-do-mar**, Bispo, Raia, Uja.

References: mainland Portugal ([Fischer et al., 1981, 1987](#); [McEachran and Capapé, 1984b](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Séret, 2016b](#); [Eschmeyer et al., 2019](#)), Azores ([Collins, 1954](#); [Osório de Castro, 1967](#); [Arruda, 1997](#); [Santos et al., 1997\(?\)](#); [Carneiro et al., 2014\(?\)](#)) and Madeira ([Osório de Castro, 1967](#)). According to [Santos et al. \(1997\)](#) the two specimens cited by [Collins \(1954\)](#) could be a misidentification of *Myliobatis aquila* (Linnaeus 1758). However, [Osório de Castro \(1967\)](#) mentions the occurrence of *R. marginata* in the fishing grounds of the Azores. Thus, the occurrence of this species in the Azores Archipelago is questionable and needs further documentation. The only records of *R. marginata* in the fishing grounds of Madeira were indicated by [Osório de Castro \(1967\)](#). The Lusitanian cownose ray is a demersal and semipelagic ray found in coastal waters over the continental shelf and around offshore islands, from shore line to about 30 m depth, on soft bottoms ([Séret, 2016b](#)) and the current geographic distribution in the Eastern Atlantic is mainly along coastline, but includes part of the Canary Islands ([Séret, 2016b](#)). It is reasonable to believe that the occurrence of this species in the Madeira region (namely on soft bottoms of Selvagens Islands) is probable. However, the occurrence of this species in Madeira needs further documentation.

Family Mobulidae

Mobula birostris (Walbaum 1792)

Common name: **Giant manta; Manta-gigante**, Diabo, Diabo-do-mar, Jamanta, Manta, Maroma, Marona, Morcego, Morcego-do-mar, Peixe-diabo, Peixe-morcego, Raia, Raia-cornuda, Raia-diabo, Raia-vampiro, Ugemanta, Ujamanta, Urjamanta, Vampiro.

References: mainland Portugal ([Gonçalves et al., 2004](#); [Abecasis et al., 2009](#); [Couturier et al., 2012](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Lawson et al., 2017](#)), Azores ([Collins, 1954](#); [Osório de Castro, 1967](#); [Ré, 1979a](#); [Wirtz, 1994, 2012](#); [Saldanha, 1995](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Garcia, 2008](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Barreiros and Gadig, 2011](#); [Couturier et al., 2012](#); [Ebert and Stehmann, 2013](#); [Gallagher et al., 2013](#); [Sobral, 2013](#); [Carneiro et al., 2014](#); [Sobral and Afonso, 2014](#); [Das and Afonso, 2017](#); [Lawson et al., 2017](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Maul, 1949d](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Nunes, 1953](#); [Osório de Castro, 1954, 1967](#); [Sanches, 1986](#); [McEachran and Séret, 1990b](#); [Wirtz et al., 2008](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Nortarbartolo di Sciara and Séret, 2016](#); [Lawson et al., 2017](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)).

Mobula mobular (Bonnaterre 1788)

Common name: **Devil fish; Manta-do-Mediterrâneo**, Boi, Cornuda, Diabo, Diabo-das-caraíbas, Diabo-do-mar, Jamanta, Jimanta, Peixe-diabo, Raia, Raia-cornuda, Ugemanta, Uja, Ujamanta, Uje, Urjamanta.

References: mainland Portugal ([Nobre, 1895a, 1935](#); [Seabra, 1911](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Stehmann, 1979c](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [Fischer et al., 1987](#); [McEachran and Séret, 1990b](#); [Santos et al., 2002a](#); [Fowler et al., 2005](#); [Melendez and Macias, 2007](#); [Ebert and Stehmann, 2013\(?\)](#); [Carneiro et al., 2014](#); [Lawson et al., 2017](#); [ICES, 2017a](#); [Froese and Pauly, 2019](#)), Azores ([Duhamel du Monceau, 1777](#); [Walbaum, 1792](#); [Lacepède, 1798](#); [van Beneden, 1892](#); [Silva 1893](#); [Sampaio, 1904](#); [Fowler, 1936a](#); [Ferreira, 1939](#); [Bigelow and Schroeder, 1953](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [McEachran and Capapé, 1984b](#); [Sanches, 1986](#); [Harmelin and Harmelin-Vivien, 1979](#); [Krefft and Stehmann, 1979c](#); [Wirtz, 1990a, 1991, 1994](#); [Lloris et al., 1991](#); [Saldanha, 1995](#); [Santos et al., 1994a, 1995a, 1997](#); [Arruda, 1997](#); [Harmelin-Vivien et al., 2001](#); [Fowler et al., 2005](#); [Melendez and Macias, 2007](#); [Porteiro et al., 2010](#); [Couturier et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Sobral and Afonso, 2014](#); [CMS/Sharks/MOS2/Doc.8.2.5, 2015](#); [Das and Afonso, 2017](#); [Lawson et al., 2017](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Johnson, 1885](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Maul, 1949](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Krefft and Stehmann, 1979c](#); [Sanches, 1986](#); [Wirtz, 1994](#); [Mello, 2000](#); [Fowler et al., 2005](#); [Wirtz et al., 2008](#); [Couturier et al., 2012\(?\)](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [CMS/Sharks/MOS2/Doc.8.2.5, 2015](#); [POGID, 2016](#); [POGIS, 2016](#); [Lawson et al., 2017](#); [Biscoito et al., 2018](#)).

2018; Froese and Pauly, 2019(?)). Last *et al.* (2016d), White *et al.* (2017b), Eschmeyer *et al.* (2019) and Froese and Pauly (2019) considered the species *Mobula japanica* (Müller & Henle 1841) as synonym of *M. mobular*. Lawson *et al.* (2017) reported the presence of this species in mainland Portugal as an extension of former range of occurrence.

***Mobula tarapacana* (Philippi 1892)**

Common name: **Chilean devil ray; Manta-cornuda**, Diabo-do-mar, Jamanta, Jimanta, Uge, Uja.

References: mainland Portugal (Lawson *et al.*, 2017), Azores (Wirtz, 1990a, 1991, 1994, 2012; Santos *et al.*, 1994a, 1995a; Afonso, 2001; Cardigos *et al.*, 2005; Garcia, 2008; Porteiro *et al.*, 2010; Barreiros and Gadig, 2011; Gallagher *et al.*, 2013; Sobral, 2013; Carneiro *et al.*, 2014; Sobral and Afonso, 2014; Thorrold *et al.*, 2014; CMS/Sharks/MOS2/Doc.8.2.5, 2015; Pardo *et al.*, 2016; Das and Afonso, 2017; Lawson *et al.*, 2017; Biscoito *et al.*, 2018) and Madeira (Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; CMS/Sharks/MOS2/Doc.8.2.5, 2015; Pardo *et al.*, 2016(?); Lawson *et al.*, 2017). Lawson *et al.* (2017) reported the presence of this species in mainland Portugal and Madeira as an extension of former range of occurrence. However, Biscoito *et al.* (2018) considered the presence of this species in Madeiran waters as in need of confirmation.

***Mobula thurstoni* (Lloyd 1908)**

Common name: **Smoothtail mobula; Manta-de-rabo-liso**.

References: Azores (Lawson *et al.*, 2017). According to Eschmeyer *et al.* (2019) *M. thurstoni* is a circumglobal species in warm seas. Lawson *et al.* (2017) mention new sightings of this species in the Azores.

Subclasse Holocephali

Order Chimaeriformes

Family Chimaeridae

***Chimaera monstrosa* Linnaeus 1758**

Common name: **Rabbit fish; Quimera-vulgar**, Bugio, Coelho, Papagaio, Papagaio-do-mar, Peixe-coelho, Peixe-rato, Ratazana, Rato, Rato-do-mar.

References: mainland Portugal (Carvalho, 1765; de Brito Capello, 1869a, 1880; Gervais and Boulart, 1877a; Posser, 1880; Osório, 1888, 1909, 1917; Nobre, 1894, 1903, 1935; Almeida and Roquette, 1892; Goode & Bean, 1895, 1896; Vieira, 1897; Bragança, 1903; Seabra, 1911; Roule, 1912; Helling, 1940; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muus and Dahlström, 1978; Krefft, 1979a, 1990a; Fernández *et al.*, 1981; INIP, 1982a; Stehmann and Bürkel, 1984c; Dinis, 1985; Sanches, 1986; Fischer *et al.*, 1987; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Erzini *et al.*, 2001b; Monteiro *et al.*, 2001; Santos *et al.*, 2002b; Borges *et al.*, 2002, 2003; Moura, 2004; Coelho *et al.*, 2005; ICES, 2005; Moura *et al.*, 2005, 2015a; Sousa *et al.*, 2005; Costa, 2007; Fernandes *et al.*, 2007; Melendez and Macias, 2007; Chaves, 2008, 2009; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Chaves and Silva, 2013; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Bañón *et al.*, 2016a; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (MNHN-IC-1884-0395, 1883; Vaillant, 1888; Goode and Bean, 1895, 1896; Neuville, 1900; Richard, 1900, 1910; Brauer, 1906; Roule, 1912, 1919; D'Ancona, 1931; Fowler, 1936a; Ferreira, 1939; Poll, 1947; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Collignon and Aloncle, 1972; Wheeler, 1978; Krefft, 1979a, 1990a; Stehmann and Bürkel, 1984c; Sanches, 1986; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Porteiro *et al.*, 2010, 2017; Barreiros and Gadig, 2011; Costa *et al.*, 2012; Menezes *et al.*, 2012; Ebert and Stehmann, 2013; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Das and Afonso, 2017(?); Biscoito *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Stehmann and Burkel, 1984; Sanches, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Didier, 2016; Biscoito *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of *Chimaera dubia* Osório 1909 caught off mainland Portugal, synonym of *C. monstrosa*, was preserved in the Museu Bocage (MB.T.111) and destroyed in the 1978 fire.

***Chimaera opalescens* Luchetti, Iglesias and Sellos 2011**

Common name: **Opal chimaera; Quimera-opala**.

References: Azores (?) ([Vieira and Cunha, 2014\(?\)](#); [Das and Afonso, 2017\(?\)](#)) and Madeira ([Freitas et al., 2017](#); [Biscoito et al., 2018](#)). Two specimens are in the MMF collections (MMF 18538, 1961 and MMF 20455, 1964). According to Das and Afonso (2017) it is suspected that records of *Chimaera monstrosa* Linnaeus 1758 from the Azores might be a misidentification of *C. opalescens*. This species has been observed near seamounts in North-East Atlantic (Vieira and Cunha, 2014) but the presence in the Azores region is not yet confirmed. Thus, the occurrence of this species in the Azores region is questionable, needs further confirmation and is not considered in the total number of valid species for this region.

***Hydrolagus affinis* (de Brito Capello 1868)**

Common name: **Smalleyed rabbitfish; Ratazana-da-fundura**, Bugio, Peixe-rato, Quimera, Quimera-da-fundura, Rato, Ratazana.

References: mainland Portugal ([de Brito Capello, 1867b, c, 1880](#); [Almeida and Roquette, 1892](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [Krefft, 1979a](#); [Stehmann and Burkel, 1984](#); [Sanches, 1986](#); [Hardy and Stehmann, 1990](#); [Moore et al., 2003](#); [Moura et al., 2005, 2015a](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)), Azores ([PENN-C 2859, 1970](#); [AMNH 78379, 1986](#) (PECS); [Desbruyères et al., 2001](#); [IMAR/DOP SEDLOP-04-O03HAF, 2003](#); [Martins et al., 2006](#); [Menezes et al., 2009, 2012](#); [Porteiro et al., 2010](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#); [Biscoito et al., 2018](#)) and Madeira ([Freitas et al., 2011](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Biscoito et al., 2018](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Chimaera affinis* de Brito Capello 1868 caught off Setúbal (mainland Portugal), synonym of *H. affinis*, was preserved in the Museu Bocage and destroyed in the 1978 fire.

***Hydrolagus lusitanicus* Moura, Figueiredo, Bordalo-Machado, Almeida & Gordo 2005**

Common name: No English common name; **Ratazana-lusitânica**, Peixe-coelho.

References: mainland Portugal ([Moura et al., 2005, 2015a](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype and the paratypes of this species, caught off mainland Portugal (36°35.0'N, 8°35.0'W) is in MB collections (MB06-004666, 2003).

***Hydrolagus mirabilis* (Collett 1904)**

Common name: **Large-eyed rabbitfish; Ratazana-olhuda**.

References: mainland Portugal ([Ebert and Stehmann, 2013](#); [Walls, 2015](#)) and Azores ([PENN-C 1459, n.d.](#); [PENN-C 1460, n.d.](#); [Saldanha, 1994](#); [Arruda, 1997](#); [IMAR/DOP SEDLOP-04-O03HMI, 2003](#); [Carneiro et al., 2014](#)). According to Saldanha and Biscoito (1997) the specimen cited as *H. mirabilis* by Saldanha (1994) was found to be a misidentification of *Hydrolagus pallidus* Hardy and Stehmann 1990. However, there are three occurrence records of this species in the Azores. Specimens are needed to confirm the presence of this species in the Azores.

***Hydrolagus pallidus* Hardy & Stehmann 1990**

Common name: Pale chimaera; **Ratazana-pálida**.

References: Azores ([Saldanha, 1994](#); [Van Dover et al., 1996](#); [Saldanha and Biscoito, 1997](#); [Santos et al., 1997](#); [Desbruyères et al., 2001](#); [Martins et al., 2006](#); [Menezes et al., 2009](#); [Porteiro et al., 2010, 2017](#); [Barreiros and Gadig, 2011](#); [Menezes et al., 2012](#); [Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#)).

Family Rhinochimaeridae

***Harriotta raleighana* Goode & Bean 1895**

Common name: **Pacific longnose chimaera; Peixe-rato-de-Raleigh**, Peixe-rato, Rato.

References: mainland Portugal (?) ([Osório de Castro, 1967\(?\)](#)) and Madeira ([Buscher and Walls, 2015](#); [Dagit et al., 2016](#)). Osório de Castro (1967) mentions the occurrence of *H. raleighana* in the mainland Portugal as doubtful and no specimens have yet been recorded. Thus, the occurrence of this species is questionable and is not considered in the total number of valid species in mainland Portugal. The distribution maps of this species reported by Buscher and Walls (2015) and Dagit et al. (2016) included the area of Selvagens Islands (Madeira EEZ).

Rhinochimaera atlantica Holt & Byrne 1909

Common name: **Straightnose rabbitfish; Peixe-rato-narigudo.**

References: mainland Portugal ([Ebert and Stehmann, 2013](#); [Carneiro et al., 2014](#)) and Azores ([Melo and Menezes, 2002](#); [Porteiro et al., 2010](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Das and Afonso, 2017](#)).

Class Actinopteri
Order Acipenseriformes
Family Acipenseridae

Acipenser sturio Linnaeus 1758

Common name: **Sturgeon; Esturjão, Esturgião, Esturião, Esturjão-comum, Esturjão-real, Peixe-da-cola, Peixe-do-caviar, Peixe-do-rei, Peixe-rei, Soilho, Solho, Solho-grande, Solho-rei, Sôlho.**

References: mainland Portugal (†) ([Carvalho, 1765](#); [Vandelli, 1771, 1797](#); [de Brito Capello, 1869a, 1880](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Nobre, 1894, 1931, 1932, 1935](#); [Vieira, 1894a, 1897](#); [Seabra, 1911](#); [Helling, 1940](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Daget, 1968](#); [Muus and Dahlström, 1978](#); [Svetovidov, 1979a](#), [1984a](#); [Fischer et al., 1987](#); [Rogado et al., 2005](#)). Regionally extinct ([Rogado et al., 2005](#)), the last record occurred during the early 1980s, in the Lower Guadiana (Almaça, 1988). Thus, this species is not considered in the total number of valid species.

Acipenser naccarii Bonaparte 1836

Common name: **Adriatic sturgeon; Esturjão-do-Adriático, Esturgião, Esturião, Esturjão, Peixe-da-cola, Peixe-do-caviar, Peixe-rei, Sôlho, Sôlho-rei.**

References: mainland Portugal (†) ([de Brito Capello, 1869a, 1880](#); [Osório, 1894](#); [Seabra, 1911](#); [Nobre, 1931, 1932, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967\(?\)](#)). Regionally extinct, the last bibliographic reference dates from Albuquerque (1954-1956) and Osório de Castro (1954). Thus, this species is not considered in the total number of valid species.

Order Elopiformes
Family Megalopidae

Megalops atlanticus Valenciennes 1847

Common name: **Tarpon; Tarpão-do-Atlântico, Peixe-prata, Peixe-prata-do-Atlântico, Tainha-congo, Tarpão.**

References: mainland Portugal ([Pereira and Saldanha, 1977](#); [Hureau and Monod, 1979](#); [Quéro et al., 1982, 1997, 1998](#); [Hureau, 1984](#); [Saldanha and Whitehead, 1990](#); [Carneiro et al., 2014](#); [Bañón et al., 2019a](#); [Froese and Pauly, 2019](#)), Azores ([Pereira and Saldanha, 1977](#); [Hureau and Monod, 1979](#); [Quéro et al., 1982](#); [Quéro, and Delmas 1982](#); [Hureau, 1984](#); [Saldanha and Whitehead, 1990](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Bañón et al., 2019a](#); [Froese and Pauly, 2019](#)) and Madeira ([Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Bañón et al., 2019a](#); [Froese and Pauly, 2019](#)).

Order Albuliformes
Family Albulidae

Pterothrissus gissu Hilgendorf 1877

Common name: Japanese gissu; **Gissu-japonês.**

References: mainland Portugal (?) ([Osório, 1917](#)). According to Hidaka et al. (2016) and Eschmeyer et al. (2019) the current geographic distribution of *P. gissu* is the Western North Pacific. However, two specimens caught off Portugal or Morocco in 1915, at the depth of 600 meters, were identified as *Bathythrissa dorsalis* Günther 1877 by Osório (1917) which is synonym of *Pterothrissus gissu* Hilgendorf 1877. Nevertheless, this author recognized small morphological differences between these specimens and the unique specimen identified by Günther (1877). These specimens were preserved in the Museu Bocage and destroyed in the 1978 fire. The current geographic distribution of *Pterothrissus gissu* Hilgendorf 1877 and *Nemoisis belloci* (Cadenat 1937) (Eastern Atlantic), together with the significant differences in morphological characters of the Osório specimens caught off in Portugal or Morocco in 1915, lead to admitting that these specimens belonged to the species *Nemoisis belloci* (Cadenat 1937)

[Longfin bonefish / Falso-badejo, Badejo, Falso-abadejo]. Thus, this species is not considered in the total number of valid species.

Order Notacanthiformes
Family Halosauridae

Aldrovandia affinis (Günther 1877)

Common name: Pale halosaur; **Peixe-lagarto-pálido**.

References: mainland Portugal ([Maul, 1976](#); [Sulak, 1990a](#); [Carneiro et al., 2014](#)), Azores ([Melo and Menezes, 2002](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Sulak, 1986](#), [1990a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Bañón et al., 2016b](#); [Smith, 2015b, 2016a](#); [Froese and Pauly, 2019](#)). Two specimens caught off at the coast of the mainland Portugal are in the MMF collections (MMF 22470, 1967, set of two specimens).

Aldrovandia gracilis Goode & Bean 1896

Common name: Slender halosaur; **Peixe-lagarto-delgado**.

References: Madeira (PECS) ([NHMUK 1995.8.4.67-69, 1972](#); [Carneiro et al., 2014](#); [Tighe, 2015a](#)).

Aldrovandia oleosa Sulak 1977

Common name: Oily halosaur; **Peixe-lagarto-oleoso**.

References: Azores ([Shcherbachov et al., 1985a](#) (PECS); [Kukuev, 2002](#) (PECS); [Carneiro et al., 2014](#) (PECS); [Tighe, 2015b](#) (PECS), [Bañón et al., 2016b](#)) and Madeira (PECS) ([Pakhorkov, 2008](#)).

Aldrovandia phalacra (Vaillant 1888)

Common name: **Hawaiian halosaurid fish; Peixe-lagarto-dos-Açores**.

References: mainland Portugal ([Maul, 1976](#); [Sulak, 1986, 1990a](#); [MNHN-IC-1988-0544, 1984](#); [Carneiro et al., 2014](#); [Bañón et al., 2016b](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1885-0382, 1883](#); [Vaillant, 1888, 1919](#); [Goode and Bean, 1895, 1896](#); [Perrier, 1903](#); [Brauer, 1906](#); [Roule, 1919a](#); [Fowler, 1936a](#); [Ferreira, 1942](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Harrison, 1966](#); [McDowell, 1973a](#); [Maul, 1976](#); [Harrison, 1979](#); [Sulak, 1986, 1990a](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002](#) (PECS); [Melo and Menezes, 2002](#); [Bergstad et al., 2008a](#) (PECS); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Pakhorkov, 2008](#) (PECS); [Smith, 2016a](#)). The lectotype of *Halosaurus phalacrus* Vaillant 1888, synonym of *A. phalacra*, caught off in the Azores is in the MNHN collections ([MNHN-IC-1885-0382, 1883](#)). One specimen caught off at the coast of the mainland Portugal is in the MMF collections (MMF 22506, 1967).

Aldrovandia rostrata (Günther 1878)

Common name: Sharpnose halosaur; **Peixe-lagarto-de-bico**.

References: Azores (PECS) ([Shcherbachov et al., 1985b](#); [Evseenko et al., 2004](#); [Carneiro et al., 2014](#); [Smith, 2016a](#)). One specimen of this species caught off at the Great Meteor Tablemount is in Shirshov Institute of Oceanology (SDSC 15910, 1982).

Halosauropsis macrochir (Günther 1878)

Common name: **Abyssal halosaur; Peixe-lagarto-abissal**.

References: mainland Portugal ([NHMUK 1887.12.7.237, 1873](#); [Sulak, 1986](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [Bañón et al., 2016b](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1885-0390, 1883](#); [Vaillant, 1888, 1919](#); [Collett, 1896a](#); [Richard, 1900, 1910](#); [Joubin, 1905](#); [Brauer, 1906](#); [Roule, 1919a](#); [Murray and Hort, 1912](#); [Koefoed, 1927](#); [Roule and Angel, 1933](#); [Zugmayer, 1933](#); [Fowler, 1936a](#); [Ferreira, 1942](#); [Collins, 1954](#); [Grey, 1956](#); [Albuquerque, 1954-1956](#); [Harrison, 1966](#); [Geistdoerfer et al., 1971](#); [McDowell, 1973a](#); [Sulak, 1986, 1990a](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Bergstad et al., 2008a](#) (PECS); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Sulak, 1986](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Halosaurus guentheri Goode & Bean 1896

Common name: Günther's halosaur; **Peixe-lagarto-de-Guenther**.

References: Madeira ([Smith, 2016a](#)). The distribution map of this species reported by Smith (2016a) included the area of Selvagens Islands (Madeira EEZ). The occurrence of this species in the Madeira Archipelago is questionable and needs further documentation.

Halosaurus johnsonianus Vaillant 1888

Common name: Johnson's halosaur; **Peixe-lagarto-de-Johnson**.

References: mainland Portugal ([Maul, 1976](#); [Harrison, 1979](#); [Sulak, 1986, 1990a](#); [Gordo et al., 1990](#); [Carneiro et al., 2014](#); [Bañón et al., 2016b](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1896a](#); [Richard, 1900, 1910](#); [Brauer, 1906](#); [Vaillant, 1919](#); [Koefoed, 1927](#); [Fowler, 1936a](#); [Collins, 1954](#); [Grey, 1956](#); [Albuquerque, 1954-1956](#); [Harrison, 1966](#); [Saldanha, 1977a](#); [Sulak, 1986](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Bañón et al., 2016b](#); [Froese and Pauly, 2019](#)) and Madeira ([Smith, 2016a](#)). The distribution map of this species reported by Smith (2016a) included the area of Selvagens Islands (Madeira EEZ). The occurrence of this species in the Madeira Archipelago is questionable and needs further documentation. One specimen caught at the coast of the mainland Portugal is in the MMF collections (MMF 22997, 1967).

Halosaurus ovenii Johnson 1864

Common name: Oven's halosaur; **Peixe-lagarto-de-Owen**.

References: mainland Portugal ([Costa and Reiner, 1978](#); [Fernandes et al., 2007](#); [Pais et al., 2009](#); [Carneiro et al., 2014](#); [Bañón et al., 2016b](#)), Azores ([Vaillant, 1888](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Harrison, 1966](#); [Maul, 1976](#); [McDowell, 1973a](#); [Maul, 1976](#); [Harrison, 1979](#); [Sulak, 1986, 1990a](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Pais et al., 2009](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a, b](#); [Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1863.12.12.1.](#), 1861-1890; [Johnson, 1864a, b](#); [Günther, 1868](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Maul, 1976](#); [Harrison, 1979](#); [Sulak, 1986, 1990a](#); [Lloris et al., 1991](#); [Pais et al., 2009](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a, b](#); [Smith, 2016a](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species is in NHMUK collections (NHMUK 1863.12.12.1.) caught off in Madeira.

Family Notacanthidae

Leptocephalus giganteus Castle 1959

Common name: Giant leptocephalus; sem nome comum.

References: mainland Portugal (?) ([NHMUK 1994.11.1.3931, 1966](#); [Carneiro et al., 2014](#)) and Madeira (?) ([NHMUK 1994.11.1.3922, 1966](#); [Carneiro et al., 2014](#)). According to Eschmeyer et al. (2019), *L. giganteus* is a larval Notocanthiform eel. This record should be considered provisional and is not considered in the total number of valid species.

Notacanthus bonaparte Risso 1840

Common name: Bonaparte's spiny eel; **Enguia-espinhosa-de-Bonaparte**.

References: mainland Portugal ([Saldanha, 1986](#); [Sulak, 1990a](#); [Saldanha et al., 1995](#); [Monteiro et al., 2001](#); [Silva et al., 2009](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Smith, 2016a](#); [Eschmeyer et al., 2019](#)), Azores (PECS) ([Bergstad et al., 2008a](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([Maul, 1955b](#); [Sulak, 1986](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Notacanthus chemnitzii Bloch 1788

Common name: **Chemnitz's spiny eel**; **Enguia-espinhosa-de-Chemnitz**.

References: mainland Portugal ([Osório, 1909](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Wheeler, 1979a](#); [Fernández et al., 1981](#); [Sulak, 1986](#); [Saldanha et al., 1995](#); [Monteiro et al., 2001](#); [Borges et al., 2002](#); [Moura, 2004](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)), Azores (PECS) ([Hareide and Garnes, 2001](#); [Porteiro et al., 2017](#); [Mecklenburg et al., 2018](#)) and Madeira ([Sulak, 1986](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Notacanthus melanoventralis* Osório 1909, synonym of *N. chemnitzii*, caught off in Sesimbra (mainland Portugal) was preserved in the Museu Bocage and destroyed in the 1978 fire.

Polyacanthonotus challengerii (Vaillant 1888)

Common name: **Challenger's spiny eel**; **Enguia-espinhosa-do-Challenger**.

References: mainland Portugal ([McCosker, 2015a](#)), Azores ([MNHN-IC-1982-0891, 1971](#); [Sulak, 1986, 1990a](#); [Lloris et al., 1991](#); [Bergstad et al., 2008a](#) (PECS); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#);

Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (MNHN-IC-1887-0130, 1883; Carneiro et al., 2014; Smith, 2016a; Porteiro et al., 2017).

Polyacanthonotus merretti Sulak, Crabtree & Hureau 1984

Common name: **Merrett's spiny eel**; **Enguia-espinhosa-de-Merrett**.

References: Madeira (McCosker, 2015b; Smith, 2016a). The distribution map of this species reported by Smith (2016a) included the area of Selvagens Islands (Madeira EEZ). Specimens are needed to confirm the presence of this species in Madeira.

Polyacanthonotus rissoanus (De Filippi & Verany 1857)

Common name: Smallmouth spiny eel; **Enguia-espinhosa-de-boca-pequena**.

References: mainland Portugal (MNHN-IC-1988-0555, 1984; Carneiro et al., 2014; Bañón et al., 2016a; Froese and Pauly, 2019) and Azores (Roule, 1919a; Fowler, 1936a; Collins, 1954; Albuquerque, 1954-1956; McDowell, 1973b; Hureau and Monod, 1979; Wheeler, 1979a; Sulak, 1986; Lloris et al., 1991; Santos et al., 1997; Desbruyères et al., 2001; Bergstad et al., 2008a (PECS); Porteiro et al., 2010, 2017; Carneiro et al., 2014; Almeida and Biscoito, 2019).

Order Anguilliformes

Family Synaphobranchidae

Dysomma brevirostre (Facciolà 1887)

Common name: **Batnose eel**; **Enguia-de-focinho-bolboso**.

References: mainland Portugal (Marques and Saldanha, 1998; Carneiro et al., 2014) and Madeira (Blache et al., 1970; Saldanha and Bauchot, 1986a; Smith and Castle, 1990; Lloris et al., 1991; Marques and Saldanha, 1998; Carneiro et al., 2014; Smith, 2016b; Froese and Pauly, 2019).

Dysommina proboscideus (Lea 1913)

Common name: No English common name; **Enguia-de-tromba-de-elefante**.

References: Azores (Lea, 1913 (PECS); MCZ 73374, 1978; Carneiro et al., 2014; Froese and Pauly, 2019 (PECS)). The holotype (unique) of *Leptocephalus proboscideus* Lea 1913, synonym of *D. proboscideus*, is in the ZMUB collections (ZMUB 5105) caught off in the Azores (PECS area).

Histiobranchus australis (Regan 1913)

Common name: Southern cut-throat eel; **Enguia-da-fundura-austral**.

References: Azores (MNHN-IC-2009-0504, 1971; MNHN-IC-2009-0503, 1971; Almeida and Biscoito, 2012, 2019; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (?) (Carneiro et al., 2014). This species was considered present in Madeira by Carneiro et al. (2014) based on a former occurrence record (34°6.0'N, 17°6.0'W) in GBIF which was lately eliminated. Thus, this species is not considered in the total number of valid species to Madeira.

Histiobranchus bathybius (Günther 1877)

Common name: **Deep-water arrowtooth eel**; **Enguia-da-fundura**, Moreão.

References: mainland Portugal (King et al., 2008; Carneiro et al., 2014; Christiansen et al., 2014; Karmovskaya and Rijnsdorp, 2015; Porteiro et al., 2017), Azores (Zugmayer, 1933; Fowler, 1936b; Bruun, 1937; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Saldanha and Bauchot, 1986a; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Bergstad et al., 2008a (PECS); Porteiro et al., 2010, 2017; Almeida and Biscoito, 2012; Carneiro et al., 2014; Karmovskaya and Rijnsdorp, 2015; Froese and Pauly, 2019) and Madeira (MNHN-IC-1992-0348, 1981; Armstrong et al., 1992; Davies and Merrett, 1998; Freitas et al., 2013; Carneiro et al., 2014; Karmovskaya and Rijnsdorp, 2015; Smith, 2016b; Porteiro et al., 2017; Priede, 2017).

Ilyophis blachei Saldanha & Merrett 1982

Common name: Blache's ooze eel; **Enguia-da-fundura-de-Blache**.

References: Azores (Saldanha and Biscoito, 1997; Porteiro et al., 2010; Carneiro et al., 2014).

Ilyophis brunneus Gilbert 1891

Common name: **Muddy arrowtooth eel**; **Enguia-da-fundura-castanha**.

References: Azores ([Saldanha and Bauchot, 1986a; Lloris et al., 1991; Arruda, 1997; Bergstad et al., 2008a \(PECS\); Carneiro et al., 2014; Porteiro et al., 2017; Froese and Pauly, 2019](#)) and Madeira ([Saldanha and Bauchot, 1986a; Lloris et al., 1991; Carneiro et al., 2014; Porteiro et al., 2017; Froese and Pauly, 2019](#)).

***Simenchelys parasitica* Gill 1879**

Common name: **Snubnosed eel; Fluta.**

References: mainland Portugal ([NHMUK 1934.12.19.15, 1912-1936; Saldanha and Bauchot, 1986a; Carneiro et al., 2014; Christiansen et al., 2014; Froese and Pauly, 2019](#)), Azores ([Collett, 1889a; Good and Bean, 1895; Richard, 1900, 1910; Perrier, 1903; Albert I^{er} de Mónaco, 1904; Brauer, 1906; Richard, 1910; Zugmayer, 1911b; Vaillant, 1919; Jaquet, 1920; Barnard, 1925; Fowler, 1936a; Ferreira, 1938; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Saldanha, 1977a; Saldanha and Bauchot, 1986a; Robins and Robins, 1989; Smith and Castle, 1990; Lloris et al., 1991; Arruda, 1992, 1997; Saldanha and Biscoito, 1997; Santos et al., 1997; Desbruyères et al., 2001; Porteiro et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Saldanha and Bauchot, 1986a; Lloris et al., 1991; Carneiro et al., 2014; Smith, 2016b; Biscoito et al., 2017; Froese and Pauly, 2019](#)). Several syntypes of *Conchognathus grimaldii* Collett 1889, synonym of *S. parasitica*, were caught off in Graciosa Islands, Faial, San Jorge, Pico, Flores and Corvo (Azores) and preserved in the MOM collections.

***Synaphobranchus affinis* Günther 1877**

Common name: **Grey cutthroat eel; Moreão-do-golfo.**

References: mainland Portugal ([Almeida et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019](#)), Azores ([Almeida and Biscoito, 2007; Bergstad et al., 2008a \(PECS\); Porteiro et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019](#)) and Madeira ([Almeida et al., 2010; Carneiro et al., 2014; POGID, 2016; Biscoito et al., 2017; Froese and Pauly, 2019](#)).

***Synaphobranchus brevidorsalis* Günther 1887**

Common name: Shortdorsal cutthroat eel; **Moreão-de-dorsal-curta.**

References: Azores ([Almeida and Biscoito, 2007; Porteiro et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019](#)).

***Synaphobranchus dolichorhynchus* (Lea 1913)**

Common name: No English common name; **Moreão-bicudo.**

References: Azores (PECS) ([Lea, 1913; Carneiro et al., 2014](#)). Adults not known.

***Synaphobranchus kaupii* Johnson 1862**

Common name: **Kaup's arrowtooth eel; Moreão-de-natura**, Congrinho, Moreão, Moreia, Moreia-de-natura.

References: mainland Portugal ([MOM-IC-0091-511, 1901; Bragança, 1903\(?\); Roule, 1916; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Maul, 1976; Fernández et al., 1981; INIP, 1981a; Martins, 1981; Merrett and Domanski, 1985; Saldanha and Bauchot, 1986a; Gordo et al., 1990; Saldanha et al., 1995; Monteiro et al., 2001; Borges et al., 2002; Moura, 2004; Fernandes et al., 2007; Gomes and Olim, 2007; King et al., 2008; Bordalo-Machado et al., 2009; Silva et al., 2009; Costa et al., 2012; Carneiro et al., 2014; Christiansen et al., 2014; Silva and Borges, 2014; Bañón et al., 2016a; Froese and Pauly, 2019](#)), Azores ([Vaillant, 1888, 1919; Goode and Bean, 1895, 1896; Collett, 1896a; Zugmayer, 1911b; Murray and Hjort, 1912; Lea, 1913; Roule, 1919a; Koefoed, 1927; Zugmayer, 1933; Guiart, 1935; Nobre, 1935; Bruun, 1936, 1937; Fowler, 1936a, b; Ferreira, 1938; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Wood and Williams, 1974; Saldanha, 1980; INIP, 1985a, b; Saldanha and Bauchot, 1986; Smith and Castle, 1990; Dias, 1991; Lloris et al., 1991; Arruda, 1992, 1997; Gonçalves and Santos, 1994; Silva and Menezes, 1995a; Menezes, 1996; Saldanha and Biscoito, 1997; Santos et al., 1997; Morato-Gomes et al., 1998; Desbruyères et al., 2001; Menezes et al., 2006, 2009, 2012; Martins et al., 2006; Rosa et al., 2006; Almeida and Biscoito, 2007, 2019; Bergstad et al., 2008a \(PECS\); Porteiro et al., 2010, 2017; Carneiro et al., 2014; Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1862.6.14.9, 1861-1890; Johnson, 1862a, 1863; Günther, 1870; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Grey, 1956; Maul, 1976; INIP,](#)

1984c, 1985c; Saldanha and Bauchot, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; POGID, 2016; Bischoff *et al.*, 2017; Froese and Pauly, 2019). Bragança (1903) reported the occurrence of *Synaphobranchus pinnatus* Gronow in mainland Portugal, caught off SW of the Espichel cape in 1903. This species is composed of two subspecies (var. *brevidorsalis* and var. *parvipinnis*). In 1903, the two subspecies were not yet identified, but taking into account the current geographic distribution of the respective species, we suppose that the specimen captured in 1903 should be the subspecies *Synaphobranchus pinnatus parvipinnis* Roule, 1916, synonym of *S. kaupii*. The holotype (unique) of *Synaphobranchus pinnatus parvipinnis* Roule 1916 is in the MOM collections (MOM-IC-0091-0511) caught off south-east of Portugal, synonym of *S. kaupii*, and a syntypes (set of two specimens) of this species is in NHMUK collections (NHMUK 1862.6.14.9) caught off in Madeira. One specimen caught off at the coast of the mainland Portugal is in the MMF collections (MMF 22469, 1967).

Family Muraenidae

Anarchias longicauda (Peter 1877)

Common name: Pygmy moray; **Moreia-de-cauda-longa**, Peixe-lobo.

References: Azores (Vaillant, 1919; Fowler, 1936a; Blache *et al.*, 1979a; Bauchot, 1986a; Smith and Böhlke, 1990; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Smith, 2016c; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Saldanha, 1968a; Blache *et al.*, 1979a; Bauchot, 1986a; Smith and Böhlke, 1990; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Smith, 2016c; Froese and Pauly, 2019).

Anarchias similis (Lea 1913)

Common name: Pygmy moray; **Moreia-anã**, Moreia-pigmeia.

References: Azores (MCZ 71062, 1978; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *A. similis* is the western Atlantic and Annabon islands.

Enchelycore anatina (Lowe 1838)

Common name: **Fangtooth moray**; **Moreia-serpente**, Moreia, Moreia-bico-de-pato, Moreia-víbora, Serpente, Serpente-do-mar, Víbora.

References: Azores[▲] (Hilgendorf, 1888a; Ferreira, 1938; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Blache *et al.*, 1979a; Böhlke, 1981; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Bauchot, 1986a; Böhlke *et al.*, 1989; Patzner *et al.*, 1990, 1992; Wirtz, 1990a, 1994; Lloris *et al.*, 1991; Arruda, 1992, 1997; Azevedo, 1995; Silva *et al.*, 1995; Menezes, 1996; Santos *et al.*, 1997; Böhlke, 2002; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Afonso, 2016; Smith, 2016c; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837; NHMUK 1869.3.7.1., 1864-1877; Günther, 1870; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Blache *et al.*, 1979a; Böhlke, 1981, 2002; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Bauchot, 1986a; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Falcón *et al.*, 2000; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Smith, 2016c; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The neotype of *Muraena anatina* Lowe 1838, synonym of *E. anatina*, is in NHMUK collections (NHMUK 1869.3.7.1) caught off in Madeira.

Gymnothorax afer Bloch 1795

Common name: **Dark moray**; **Moreão-escuro**, Moreão, Moreia, Moreia-verde.

References: Azores (?) (Fowler, 1936b; Osório de Castro, 1967(?); Arruda, 1997; Santos *et al.*, 1997(?); Carneiro *et al.*, 2014(?)) and Madeira (?) (Fowler, 1936b; Osório de Castro, 1967(?)). This species is cited for the Azores and Madeira by Fowler (1936b) based on three specimens: two collected in Faial and in Terceira Islands, by F.M. McGuire in 1879 and by W. Trelease in 1894, respectively, and the third in Funchal, Madeira, obtained by A.C. di Noronha. According to Wirtz *et al.* (2008) the specimen deposited at the USNM, catalogue number 94531, originally identified as *G. afer* was re-identified as *Gymnothorax unicolor* (Delaroche 1809). The two Azorean specimens, also deposited at the USNM, catalogue numbers 23320 and 94497, were also re-identified as *G. unicolor*. Thus, there is no records of *G. afer* for the Azores or Madeira.

Gymnothorax bacalladoi Böhlke & Brito 1987

Common name: **Canary moray; Moreia-das-Canárias.**

References: Madeira ([Brito et al., 2002](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Almada et al., 2009a](#); [Carneiro et al., 2014](#); [Biscoito 2015a](#); [Smith, 2016c](#); [Priede, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Gymnothorax maderensis (Johnson 1862)

Common name: **Sharktooth moray; Moreia-da-Madeira**, Moreão, Moreia.

References: Azores ([Vaillant, 1919](#); [Harmelin and Harmelin-Vivien, 1979](#); [Arruda, 1997](#); [Kukuev, 2002 \(PECS\)](#); [Carneiro et al., 2014](#); [Smith, 2016c](#)) and Madeira ▲ ([Lowe, 1837](#); [Kaup, 1856a](#); [Johnson, 1862a, 1863](#); [Günther, 1870](#); [de Brito Capello, 1872](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Blache et al., 1979a](#); [Fischer et al., 1981](#); [Böhlke, 1981](#); [Dooley et al., 1985](#); [Bauchot, 1986a](#); [Smith and Böhlke, 1990](#); [Lloris et al., 1991](#); [Delgado, 2008, 2015a](#); [Ferreira et al., 2008](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Smith, 2016c](#); [Carvalho-Filho and Paiva, 2017](#); [Froese and Pauly, 2019](#)).

Gymnothorax miliaris (Kaup 1856)

Common name: **Goldentail moray; Moreia-dourada**, Moreia.

References: Azores (?) ([MCZ 34270, 1989](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#)) and Madeira ([Smith, 2016c](#)). The specimen deposited at the Museum of Comparative Zoology, catalogue number 34270, originally identified as *G. miliaris*, was re-identified as *Gymnothorax unicolor* (Delaroche 1809). There are no records of *G. miliaris* for the Azores. Thus, this species is not considered in the total number of valid species to the Azores. According to Smith (2016c) the distribution map of *G. miliaris* includes the Selvagens Islands. The occurrence of this species in the Madeira Archipelago is questionable and needs further documentation.

Gymnothorax polygonius Poey 1875

Common name: **Polygon moray; Moreia-poligonal**, Moreão.

References: Madeira ([Bauchot, 1986a](#); [Lloris et al., 1991](#); [Ferreira et al., 2008](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Biscoito, 2015b](#); [Smith, 2016c](#); [Froese and Pauly, 2019](#)).

Gymnothorax unicolor (Delaroche 1809)

Common name: **Brown moray; Moreia-castanha**, Moreão, Moreão-castanho, Morena, Moreno, Morreão, Morrião.

References: mainland Portugal ([Almeida and Roquette, 1892](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bauchot, 1986a](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ▲ ([USNM 23320, 1879](#); [USNM 94497, 1894](#); [Sampaio, 1904](#); [Vaillant, 1919](#); [Fowler, 1936](#); [Ferreira, 1938](#); [Nobre, 1924, 1930, 1935](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Blache et al., 1979a](#); [Harmelin and Harmelin-Vivien, 1979](#); [Saldanha, 1980, 1982b, 1995](#); [Böhlke, 1981](#); [Martins, 1982a](#); [Dooley et al., 1985](#); [INIP, 1985c](#); [Bauchot, 1986a](#); [Fischer et al., 1987](#); [Patzner et al., 1990, 1992](#); [Ré, 1990](#); [Santos et al., 1990b, 1994a, 1995a, 1997](#); [Wirtz, 1990a, 1994](#); [Smith and Böhlke, 1990](#); [Azevedo et al., 1991, 1992](#); [Dias, 1991](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Gonçalves, 1993](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Silva et al., 1995](#); [Isidro, 1996](#); [Menezes, 1996](#); [Harmelin-Vivien et al., 2001](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Smith, 2016c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ▲ ([USNM 94531, n.d.](#); [Lowe, 1837](#); [Kaup, 1856a](#); [Johnson, 1862a](#); [Günther, 1870](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Blache et al., 1979a](#); [Böhlke, 1981](#); [Fischer et al., 1981](#); [Saldanha, 1982b](#); [INIP, 1984b](#); [Dooley et al., 1985](#); [Bauchot, 1986a](#); [Smith and Böhlke, 1990](#); [Lloris et al., 1991](#); [Andrade and Albuquerque, 1995](#); [Falcón et al., 2000](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGID, 2016](#); [POGIS, 2016](#); [Smith, 2016c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Gymnothorax vicinus (Castelnau 1855)

Common name: **Purplemouth moray; Moreão-amarelo**, Moreão, Moreia, Moreia-felia.

References: Azores (?) (Smith and Böhlke, 1990; IMAR/DOP 6FCO-01-P07MOR2, 2007; IMAR/DOP 3SMA-02-P08MOR2, 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (?) (Blache et al., 1979a; Böhlke, 1981; Fischer et al., 1981; Dooley et al., 1985; Bauchot, 1986a; Smith and Böhlke, 1990; Lloris et al., 1991; Wirtz et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019). According to Eschmeyer et al. (2019) the current geographic distribution of *G. vicinus* is the Western Atlantic and in the central Atlantic: Ascension and Saint Helena islands; Smith (2016c) also mentions in the Western Atlantic, from Bermuda, the Bahamas, and Florida south to Brazil and in the Eastern Atlantic, in the Cape Verde, Canary, Bay of Biafra, and Ascension Islands. According to Brito et al. (2014) the previous existing records for the Azores and Madeira islands were caused by misidentification, but it occurs in Canary Islands. Smith (2016c) also considered erroneous the records from the Azores and Madeira. However, the distribution map of this species reported by Smith (2016c) partially includes the Selvagens Islands area. There are no records of *G. vicinus* for Madeira and the occurrence of this species in the Madeira Archipelago is questionable and needs further documentation. Thus, this species is not considered in the total number of valid species to Madeira and the Azores.

***Muraena augusti* (Kaup 1856)**

Common name: **White-eyed moray; Moreia-da-Macaronésia**, Moreão, Moreia, Moreia-preta.

References: mainland Portugal (Abecasis et al., 2009; OCEANA, 2014), Azores[▲] (Regan, 1903a; Ferreira, 1938; Collins, 1954; Osório de Castro, 1967; Saldanha, 1980, 1995; Böhlke, 1981; Fischer et al., 1981; Dooley et al., 1985; Patzner et al., 1990, 1992; Wirtz, 1990a, 1994; Lloris et al., 1991; Azevedo et al., 1992; Azevedo, 1995; Barreiros, 1995; Santos et al., 1994a, 1995a, 1996, 1997; Arruda, 1997; Harmelin-Vivien et al., 2001; Brito et al., 2002; Cardigos et al., 2005; Menezes et al., 2006; Floeter et al., 2008; Bertoncini et al., 2010; Botelho et al., 2010; Porteiro et al., 2010; Diogo and Pereira, 2013, 2014; Carneiro et al., 2014; Delgado, 2015b; Smith, 2016c; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837; Richardson 1844-1848; Kaup, 1856b; Johnson, 1862a; NHMUK 1862.2.5.2, 1861-1890; Günther, 1870; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Böhlke, 1981; Fischer et al., 1981; Dooley et al. 1985; Lloris et al., 1991; Wirtz, 1994; Delgado, 1998, 2015b; Falcón et al., 2000; Mello, 2000; Brito et al., 2002; Araújo et al., 2005, 2008; Ribeiro et al., 2005b; Floeter et al., 2008; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Afonso, 2016; POGID, 2016; POGIS, 2016; Smith, 2016c; Froese and Pauly, 2019). This species is sometimes confused with *Muraena helena* Linnaeus 1758 (Smith, 2016c). The holotype (unique) of *Thyrsoidea atlantica* Johnson 1862, synonym of *M. augusti*, is in the NHMUK collections (NHMUK 1862.2.5.2) caught off at sea near Madeira. A second specimen recorded as *Thyrsoidea augusti* Kaup 1856 (no types known) was caught off in Madeira and is also synonym of *M. augusti*.

***Muraena helena* Linnaeus 1758**

Common name: **Mediterranean moray; Moreia, Cobra-do-mato, Morea, Moreão, Moreia-amarela, Moreia-legitima, Moreia-pintada, Moreia-preta, Morena, Murêa, Pilosa, Pintada.**

References: mainland Portugal[▲] (Carvalho, 1765; Vandelli, 1771; de Brito Capello, 1869a; Osório, 1888; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1894, 1895a, c, 1903, 1935; Vieira, 1901; Richard, 1904, 1908; Seabra, 1911; Osório de Castro, 1954, 1967; Blache et al., 1979a; Bauchot, 1986a; Fischer et al., 1987; Almeida, 1996; Erzini et al., 1996, 2001a; Borges et al., 2002; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Abecasis et al., 2009; Silva et al., 2009; Carneiro et al., 2014; Christiansen et al., 2014; OCEANA, 2014; Golani and Fricke, 2018; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Regan, 1903a; Sampaio, 1904; Fowler, 1936; Ferreira, 1938; Nobre, 1924, 1930; Ribeiro, 1936; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Wheeler, 1978; Blache et al., 1979a; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Böhlke, 1981; Fischer et al., 1981; Martins, 1982a; Dooley et al., 1985; INIP, 1985b, c; Bauchot, 1986a; Dias and Monteiro, 1986; Barreiros, 1990, 1995; Dias et al., 1990; Patzner et al., 1990, 1992; Santos et al., 1990b, 1994a, 1995a, 1997; Smith and Böhlke, 1990; Wirtz, 1990a, 1994; Azevedo et al., 1991, 1992; Dias, 1991; Lloris et al., 1991; Arruda et al., 1992; Santos, 1992a; Gonçalves, 1993; Gonçalves et al., 1995; Krug, 1995; Silva et al., 1994, 1995; Azevedo, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Arruda, 1997; Morato-Gomes et al., 1998; Sousa et al., 1999; Harmelin-Vivien et al.,

2001; Quigley and Flannery, 2004; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Afonso, 2016; Smith, 2016c; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Lowe, 1837; Johnson, 1863; Günther, 1870; Noronha and Sarmento, 1934; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Blache *et al.*, 1979a; INIP, 1980, 1982c, 1984b, c, 1985c; Böhlke, 1981; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Bauchot, 1986a; Smith and Böhlke, 1990; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Delgado, 1998; Mello, 2000; Araújo *et al.*, 2005; Ferreira *et al.*, 2008; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; Smith, 2016c; Biscoito *et al.*, 2017; Froese and Pauly, 2019).

***Muraena melanotis* (Kaup 1859)**

Common name: **Honeycomb moray; Moreia-pintada**, Moreia, Cobra.

References: Madeira (Fischer *et al.*, 1981; Smith, 2015a). According to Eschmeyer *et al.* (2019) the current geographic distribution of *M. melanotis* is the Western Atlantic (Saint Paul's Rocks) and, in the Eastern Atlantic, from Mauritania south to Angola, including Cape Verde Islands and islands in Gulf of Guinea, and also in the southern central Atlantic (Saint Helena Island). The distribution maps of this species is reported by Fischer *et al.* (1981), and Smith (2015a) partially includes the Selvagens Islands area (Madeira EEZ). Thus, the validity of the Madeira records are questionable and specimen are needed for confirmation.

Family Chlopsidae

***Chlopsis bicolor* Rafinesque 1810**

Common name: **Bicoloured false moray; Congrinho-bicolor**.

References: Azores (IMAR/DOP COLETA 70, 1983; SDSC 22973, 1967 (PECS), SDSC 22974, 1967 (PECS); Maul, 1976 (PECS); Ehrich, 1977 (PECS); Santos *et al.*, 1997; Fock *et al.*, 2002 (PECS); Kukuev, 2002 (PECS); Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Karmovskaya and Papaconstantinou, 2015a) and Madeira (Karmovskaya and Papaconstantinou, 2015a). Two specimens caught off at the Meteor Bank are in the MMF collections (MMF 22381, 1967 and MMF 22383, 1967).

***Chlopsis dentatus* (Seale 1917)**

Common name: **Mottled false moray; Congrinho-malhado**.

References: Azores PECS (NHMUK 1994.11.1.482, 1981) and Madeira (?) (Carvalho-Filho and Paiva, 2017). This species is reported to occur in Madeira (Carvalho-Filho and Paiva, 2017) but the occurrence record of the Museum History (London) mentions off SW Madeira Islands, in the Madeira abyssal Plain (NHMUK 1994.11.1.482, 32°22.998'N, 29°49.398'W). However, these geographic coordinates correspond to a position at 53 NM SSW of the Plato Seamount in the PECS of Azores. Thus, this species is not considered in the total number of valid species to Madeira EEZ.

Family Derichthyidae

***Derichthys serpentinus* Gill 1884**

Common name: **Narrownecked oceanic eel; Enguia-de-pescoço-fino**.

References: mainland Portugal (NHMUK 1998.8.9.12751, 1966; IFREMER Biocean 34589, 1969; Carneiro *et al.*, 2014; Karmovskaya and Papaconstantinou, 2015b), Azores (Post and Tesch, 1982; Karmovskaya, 1985; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Karmovskaya and Papaconstantinou, 2015b) and Madeira (Karmovskaya and Papaconstantinou, 2015b; Smith, 2016d; Porteiro *et al.*, 2017).

***Nessorhamphus ingolfianus* (Schmidt 1912)**

Common name: **Duckbill oceanic eel; Enguia-bico-de-pato**.

References: mainland Portugal (MCZ 153058, 1986; Bauchot and Saldanha, 1986a; ZSM-Pis-029407 / 534598 / 386186, 1990; Carneiro *et al.*, 2014; Papaconstantinou and Karmovskaya, 2015; Bañón *et al.*, 2016a; Froese and Pauly, 2019), Azores (Nielsen, 1974 (PECS); Lloris *et al.*, 1991; Arruda, 1997; Bauchot and Saldanha, 1986a; MAR-ECO 13211, 2004; Sutton *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Papaconstantinou and Karmovskaya, 2015; Roberts *et al.*, 2017; Froese and Pauly, 2019) and

Madeira (MCZ 64395, 1969; MCZ 64397, 1969; MCZ 153057, 1986; Bauchot and Saldanha, 1986a; Carneiro et al., 2014; Papaconstantinou and Karmovskaya, 2015; Smith, 2016d; Froese and Pauly, 2019). The syntypes of *Leptocephalus ingolfianus* Schmidt 1912, synonym of *N. ingolfianus*, is in the ZMUC collections (P32520-23, 1911) caught off Azorean PECS.

Family Ophichthidae

***Apterichtus anguiformis* (Peters 1877)**

Common name: Slender finless eel; **Cobra-de-boca-negra**.

References: Madeira (Floeter et al., 2008; Wirtz et al., 2008(?); MNHN-IC-B-2676, 1883; Carneiro et al., 2014; McCosker and Hibino, 2015; Froese and Pauly, 2019).

***Apterichtus caecus* (Linnaeus 1758)**

Common name: European finless eel; **Congrinho-da-areaia**.

References: Azores (Ferreira, 1938; SMF 7538, 1964; Pereira and Aguiar, 1979; Bauchot, 1986b, 1987; IMAR/DOP COLETA 103, 1989; IMAR/DOP COLETA 102, 1989; IMAR/DOP COLETA 101, 1990; Patzner et al., 1990, 1992; Wirtz, 1990a, 1991, 1994; Azevedo et al., 1991, 1992; Lloris et al., 1991; Arruda et al., 1992; Santos et al., 1994b, 1997; Azevedo, 1995; Saldanha, 1995; Arruda, 1997; Porteiro et al., 2010; Carneiro et al., 2014; McCosker and Hibino, 2015; McCosker, 2016; Froese and Pauly, 2019) and Madeira (MNHN-IC-1884-0426, 1883; Floeter et al., 2008; Wirtz et al., 2008; Carneiro et al., 2014; McCosker and Hibino, 2015; Froese and Pauly, 2019).

***Dalophis imberbis* (Delaroche 1809)**

Common name: Armless snake eel; **Cobra-imberbe**, Cobra.

References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Bauchot, 1986b; Fischer et al., 1987; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967). The only records of *D. imberbis* in the fishing grounds of Madeira were indicated by Osório de Castro (1967). The occurrence of this species in Madeira region needs further documentation.

***Echelus myrus* (Linnaeus 1758)**

Common name: **Painted eel**; **Cobra-de-orelhas**, Cobra-do-mar, Peixe-cobra

References: mainland Portugal (Dooley et al., 1985; Bauchot, 1986b; Fischer et al., 1987; Leiby 1990; ZSM-Pis-029321 / 534521 / 386109, 1990(?); Borges et al., 2002; ICES, 2005; Carneiro et al., 2014; McCosker, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Carneiro et al., 2014; Karmovskaya and Papaconstantinou, 2015c(?)). According to Wirtz et al. (2008), Froese and Pauly (2007) erroneously recorded this species from Madeira, allegedly on the basis of Leiby (1990), which does not, however, include such a record. The map of distribution of *E. myrus* reported by Karmovskaya and Papaconstantinou (2015c) includes the Selvagens Islands. The occurrence of this species in the Madeira Archipelago is questionable and needs further documentation.

***Myrichthys pardalis* (Valenciennes 1839)**

Common name: **Leopard eel**; **Cobra-leopardo**, Cobra, Cobra-de-água, Cobra-do-mar, Moreia, Moreia-pintada.

References: Madeira (Fischer et al., 1981; González Pérez, 2015a). According to Eschmeyer et al. (2019) the current geographic distribution of *M. pardalis* is the Eastern Atlantic. The distribution maps of this species reported by Fischer et al. (1981) and González Pérez (2015a) partially include the Selvagens Islands area (Madeira EEZ).

***Myrophis plumbeus* (Cope 1871)**

Common name: **Leaden worm eel**; **Cobra-chumbo**.

References: Azores (PECS) (NHMUK 1994.11.1.1413, 1970; Carneiro et al., 2014).

***Ophisurus serpens* (Linnaeus 1758)**

Common name: **Serpent eel**; **Cobra-do-mar**, Cobra, Enguia-do-mar, Órfia, Peixe-cobra, Serpente, Serpente-do-mar.

References: mainland Portugal (de Brito Capello, 1869a, 1880; Osório, 1888, 1917; Vieira, 1897, 1901; Bragança, 1903(?); Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández et al., 1981; Fischer et al., 1981, 1987; Cappetta et al., 1985;

Bauchot, 1986b; Erzini *et al.*, 2001b; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Palma *et al.*, 2003; Gomes and Olim, 2007; Henriques *et al.*, 2008; Bañón *et al.*, 2011; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Osório de Castro, 1967) and Madeira (Lowe, 1849; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1949; Albuquerque, 1954-1956; Osório de Castro, 1967; Fischer *et al.*, 1981; Bauchot, 1986b; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Bañón *et al.*, 2011; Carneiro *et al.*, 2014; McCosker, 2016; Froese and Pauly, 2019). The only records of *O. serpens* in the fishing grounds of the Azores were indicated by Osório de Castro (1967). The occurrence of this species in the Azores region is questionable and needs further documentation.

***Pisodonophis semicinctus* (Richardson 1848)**

Common name: Saddled snake eel; **Cobra-malhada**, Cobra, Cobra-do-mar, Moreia, Peixe-cobra, Serpente, Serpente-do-mar.

References: mainland Portugal. One specimen caught off near shore Peniche in 2008 was identified by the authors as *P. semicinctus* (unpublished data). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. semicinctus* is the Mediterranean Sea and Eastern Atlantic.

Family Muraenesocidae

***Cynoponticus ferox* Costa 1846**

Common name: **Guinean pike conger**; **Congro-branco-africano**, Congro-branco, Congro, Safio, Safio-branco.

References: Madeira (Fischer *et al.*, 1981; Karmovskaya and Papaconstantinou, 2015d). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. ferox* is the Mediterranean Sea and Gibraltar south to Angola. The distribution maps of this species reported by Fischer *et al.* (1981) and Karmovskaya and Papaconstantinou (2015d) partially include the Selvagens Islands area (Madeira EEZ). Thus, the occurrence of this species in the Madeira region is questionable and needs further documentation.

Family Nettastomatidae

***Facciolella oxyrhynchus* (Bellotti 1883)**

Common name: **Facciola's sorcerer**; **Cobra-cabeça-de-pato**.

References: mainland Portugal (ICM-CSIC 7764, n.d.; Saldanha, 1986; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002; ICES, 2005; Fernandes *et al.*, 2007; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Smith and Castle, 1982; IMAR/DOP COLETA 215, 1996; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Carneiro *et al.*, 2014) and Madeira (Saldanha, 1986; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Nettastoma melanurum* Rafinesque 1810**

Common name: Blackfin sorcerer; **Cobra-de-cauda-preta**, Cobra-bico-de-pato.

References: mainland Portugal (Osório, 1909; Saldanha, 1966a; Blache *et al.*, 1979b; Merrett and Saldanha, 1985; Saldanha, 1986; Saldanha *et al.*, 1995; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Chaves, 2008; Silva, 2008, 2010, 2011; Melo *et al.*, 2009; Silva *et al.*, 2009; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Priede, 2017; Froese and Pauly, 2019), Azores (Lea, 1913; Roule, 1919a; Fowler, 1936a; Ferreira, 1938; Collins, 1954; Smith and Castle, 1982; Smith, 1990a; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (NHMUK 1994.11.1.3474, 1966; MCZ 65443, 1970; Carneiro *et al.*, 2014; Smith, 2016e). The holotype of *Muraenosaurus guentheri* Osório 1909, synonym of *N. melanurum*, caught off in Cape Espichel (mainland Portugal) was preserved in the Museu Bocage (MB 2066) and destroyed in the 1978 fire.

***Venefica proboscidea* (Vaillant 1888)**

Common name: **Whipsnout sorcerer**; **Cobra-de-bico-comprido**.

References: mainland Portugal (Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Gomes and Olim, 2007; Carneiro *et al.*, 2014; Costa, 2014).

Family Congridae

Ariosoma balearicum (Delaroche 1809)

Common name: **Bandtooth conger; Congro-das-Baleares**, Bocavante, Coreano, Safio.

References: mainland Portugal ([Bauchot and Saldanha, 1986b](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Roule, 1919a](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Bauchot and Saldanha, 1986b](#); [Wirtz, 1994](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Menezes et al., 2006](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Fowler, 1923, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Bauchot and Saldanha, 1986b](#); [Araújo et al., 2005](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Bathyuroconger vicinus (Vaillant 1888)

Common name: Long-tooth conger; **Congro-castanho-dentão**, Congro, Congro-castanho, Safio, Safio-castanho.

References: mainland Portugal ([Sánchez, 1999](#); [ICES, 2005](#); [Carneiro et al., 2014](#)).

Conger conger (Linnaeus 1758)

Common name: **European conger; Congro**, Chamirro, Congra, Congra-de-natura, Congro-de-natura, Congro-europeu, Congro-grande, Congro-legítimo, Congro-vulgar, Cronga, Porca, Safio, Safio-preto, Samilo, Samiro, Samirro, Zafio.

References: mainland Portugal▲ ([Carvalho, 1765](#); [de Brito Capello, 1869a, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1893, 1897, 1901](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Osório, 1895](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Muus and Dahlström, 1978](#); [Dinis, 1981](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, b, 1982a, b](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Bauchot and Saldanha, 1986b](#); [Arruda et al., 1988](#); [Costa and Quéro, 1988](#); [Smith, 1990b](#); [Martins et al., 1992, 2015](#); [Figueiredo et al., 1994](#); [Saldanha et al., 1995](#); [Almeida, 1996](#); [Gonçalves et al., 1996, 1997](#); [Erzini et al., 1997, 2001a, b](#); [Cabral, 1999](#); [Gomes et al., 2001](#); [Monteiro et al., 2001](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Santos et al., 2002b, c](#); [Garnerot et al., 2004](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Ribeiro et al., 2006, 2008](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Neves et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Abecasis et al., 2009](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [OCEANA, 2014](#); [Bañón et al., 2016a](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores▲ ([Drouët, 1861b](#); [Ramos, 1869](#); [Hilgendorf, 1888a](#); [Regan, 1903a](#); [Sampaio, 1904](#); [Fowler, 1936](#); [Ribeiro, 1936](#); [Cadenat, 1937](#); [Ferreira, 1938](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Dias et al., 1976, 1990](#); [Muzanor and Westhaus, 1981](#); [Martins, 1982a](#); [INIP, 1985a, b, c](#); [Dias and Monteiro, 1986](#); [Silva, 1986a,b, 1992, 1995](#); [Barreiros, 1990, 1995](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1994a, 1995a, 1997](#); [Smith, 1990b](#); [Wirtz, 1990a, 1994](#); [Dias, 1991](#); [Dias and Cascalho, 1991](#); [Arruda, 1992, 1997](#); [Santos, 1992a](#); [Silva and Krug, 1992](#); [Gonçalves, 1993](#); [Krug, 1995](#); [Nash et al., 1994a](#); [Silva et al., 1994, 1995](#); [Azevedo, 1995](#); [Gonçalves et al., 1995](#); [Santos and Nash, 1995](#); [Silva and Menezes, 1995a, b](#); [Isidro, 1996](#); [Menezes, 1996](#); [Morato-Gomes et al., 1998](#); [Sousa et al., 1999](#); [Morato et al., 2001a](#); [Kukuev, 2002 \(PECS\)](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Porteiro et al., 2010, 2013](#); [Gomes-Pereira et al., 2012](#); [Besugo, 2013](#); [Pham et al., 2013](#); [Carneiro et al., 2014](#); [Diogo and Pereira, 2014](#); [Afonso, 2016](#); [Torres et al., 2016](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira▲ ([Lowe, 1837](#); [de Velloso, 1892](#); [Fowler, 1923, 1936a](#); [Noronha and Sarmento, 1934](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Osório de Castro, 1954, 1967](#); [Maul, 1967](#); [INIP, 1980, 1982c](#), [1984a, b, c](#); [Bauchot and Saldanha, 1986b](#); [Smith, 1990b](#); [Abreu et al., 1995](#); [Andrade and Albuquerque, 1995](#); [Ferreira et al., 2008](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Christiansen et al., 2009](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGID, 2016](#); [POGIS, 2016](#); [Biscoito et al., 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Conger triporiceps Kanazawa 1958

Common name: **Manytooth conger; Congro-dentudo**.

References: Madeira ([MCZ 71775, 1970](#); Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2019) the current geographic distribution of *C. triporiceps* is the Western Atlantic. However, there is a single historical occurrence record of this species in the Eastern Atlantic. The preserved specimen (larval) from Madeira deposited in the Museum of Comparative Zoology requires verification.

***Gnathophis codoniphorus* Maul 1972**

Common name: No English common name; **Congro-dos-Açores**.

References: Azores (PECS) ([Maul, 1972b, 1976](#); Ehrich, 1977 (PECS); Hureau and Monod, 1979; Bauchot and Saldanha, 1986b; Lloris *et al.*, 1991; Fock *et al.*, 2002; Pakhorukov, 2008; Carneiro *et al.*, 2014; Smith, 2016e; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira ([Lloris *et al.*, 1991](#)). The holotype and the paratypes of this species are in the MMF collections [MMF 22876, 1967 and MMF 22545, 1967 (set of two specimens), respectively] all caught off at Meteor Bank.

***Gnathophis mystax* (Delaroche 1809)**

Common name: Thinlip conger; **Coreano**, Safio.

References: mainland Portugal ([Saldanha, 1967](#); Palmer, 1971; Maul, 1976; Fischer *et al.*, 1981, 1987; Bauchot and Saldanha, 1986b; Moura, 2004; ICES, 2005; Silva, 2008, 2010; Silva *et al.*, 2009; Carneiro *et al.*, 2014; Smith, 2016e; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores ([Ehrich, 1977 \(PECS\)](#); IMAR/DOP 3SMI-01-P97CGR2, 1997; Fock *et al.*, 2002 (PECS); Kukuev, 2002 (PECS); Menezes *et al.*, 2006; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira ([Noronha and Sarmento, 1948](#); Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Bauchot and Saldanha, 1986b; Lloris *et al.*, 1991; Christiansen *et al.*, 2009; Carneiro *et al.*, 2014; Smith, 2016e; Froese and Pauly, 2019). Two specimens caught off at the coast of mainland Portugal and Josephine Bank are in the MMF collections (MMF 22800, 1967 and MMF 22801, 1967, respectively).

***Heteroconger longissimus* Günther 1870**

Common name: Brown garden eel; **Enguia-de-jardim**, Congro.

References: Madeira ([Maul, 1949d](#); Bauchot and Bassot, 1958; Hureau and Monod, 1979; Tortonese and Hureau, 1979; INIP, 1982c, 1984a, b, c; Saldanha, 1982a; Augier, 1985; Dooley *et al.*, 1985; Bauchot and Saldanha, 1986; Saldanha *et al.*, 1986; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Mello, 2000; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Bischoff, 2015c; POGID, 2016; POGIS, 2016; Smith, 2016e; Froese and Pauly, 2019).

***Paraconger macrops* (Günther 1870)**

Common name: Manypore conger; **Congro-de-natura**, Congra, Congro, Congro-da-areia, Congro-de-rolo, Congro-do-rolo, Cronga, Ortiga, Safio.

References: Azores ([Hilgendorf, 1888a](#); Ferreira, 1938; Collins, 1954; Saldanha, 1981, 1982a; Bauchot and Saldanha, 1986b; Lloris *et al.*, 1991; Arruda, 1992, 1997; Wirtz, 1994; Santos *et al.*, 1997; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019; Moreno-Borges *et al.*, 2019) and Madeira ([Günther, 1870](#); Kanazawa, 1961; Osório de Castro, 1954, 1967; Blache *et al.*, 1979c; Saldanha, 1982a; Bauchot and Saldanha, 1986b; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Smith, 2016e; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019; Moreno-Borges *et al.*, 2019).

***Paraconger notialis* Kanazawa 1961**

Common name: Guinean conger; **Congro-da-Guiné**, Congro, Congro-do-oeste-africano.

City.: Azores ([IMAR/DOP 6FCO-02-P07CGI2, 2007](#); IMAR/DOP 2TER-04-P12CGI2, 2012; IMAR/DOP 2FPI-01-P13CGI2, 2013).

***Pseudophichthys splendens* (Lea 1913)**

Common name: **Purplemouthed conger**; **Congro-de-boca-púrpura**, Coreano, Safio.

References: Azores ([MOM-IC-0091-0544, 1902](#); Roule, 1916, 1919a; Fowler, 1936a; Ferreira, 1938; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Blache *et al.*, 1979c; Bauchot and Saldanha, 1986b; Smith, 1990b; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Melo *et al.*, 2009; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Smith, 2016e; Froese and Pauly, 2019) and Madeira ([Bauchot and Saldanha, 1986b\(?\)](#); Smith, 2016e).

Family Nemichthidae

Avocettina infans (Günther 1878)

Common name: **Avocet snipe eel; Cobra-centopeia.**

References: mainland Portugal ([Osório de Castro, 1967](#); [NHMUK 2015.2.26.14, 1984 \(PECS\)](#)), Azores ([Vaillant, 1888](#); [Fowler, 1936a](#); [Ferreira, 1938](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Nielsen, 1986a](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Smith, 2016e](#)) and Madeira ([Osório de Castro, 1967](#); [Maul, 1971a](#); [Nielsen, 1986a](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Smith, 2016e](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)).

Labichthys carinatus Gill & Ryder 1883

Common name: Shortgut fintail snipe eel; **Cobra-de-crista.**

References: Azores (PECS) ([ZMH 118052, 1979](#); [Carneiro et al., 2014](#); [Smith, 2016e](#)) and Madeira ([ZMH 109394, 1971](#); [Smith, 2016e](#)).

Nemichthys curvirostris (Strömmman 1896)

Common name: Pale threadtail snipe eel; **Cobra-pálida-ponteada.**

References: mainland Portugal ([Ralph, 2015a](#)), Azores ([Nielsen and Smith, 1978](#); [Hureau and Monod, 1979](#); [Nielsen, 1986a](#); [Smith and Nielsen, 1989](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Ralph, 2015a](#); [Smith, 2016e](#); [Froese and Pauly, 2019](#)) and Madeira ([Hureau and Monod, 1979](#); [Nielsen, 1986a](#); [Carneiro et al., 2014](#); [Ralph, 2015a](#); [Smith, 2016e](#); [Froese and Pauly, 2019](#)). According to Ralph (2015a) this species is found off mainland Portugal.

Nemichthys scolopaceus Richardson 1848

Common name: **Slender snipe eel; Cobra-de-bico, Cobra.**

References: mainland Portugal ([Bragança, 1903](#); [Osório, 1917](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Martins, 1981](#); [Nielsen, 1986a](#); [Monteiro et al., 2001](#); [Borges et al., 2002](#); [Moura, 2004](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Silva, 2008, 2011](#); [Silva et al., 2009](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Karmovskaya and Papaconstantinou, 2015e](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912](#); [Roule, 1919a](#); [Roule and Bertin, 1929, 1931](#); [Fowler, 1936a](#); [Roule and Angel, 1933](#); [Belloc, 1938](#); [Ferreira, 1938](#); [Albuquerque, 1954-1956](#); [Koefoed, 1955](#); [Osório de Castro, 1967](#); [Nielsen and Smith, 1978](#); [Wheeler, 1978](#); [Post and Tesch, 1982](#); [Nielsen, 1986a](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Karmovskaya and Papaconstantinou, 2015e](#); [Smith, 2016e](#); [Almeida and Bischoff, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Richardson, 1848](#); [Lowe, 1850, 1851](#); [Günther, 1870](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Roule, 1919a](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Kanazawa and Maul, 1967](#); [Nielsen, 1986a](#); [Larsen 1990](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Karmovskaya and Papaconstantinou, 2015e](#); [Smith, 2016e](#); [Froese and Pauly, 2019](#)). The holotype of *Paravocettinops trilinearis* Kanazawa & Maul 1967 is in the MMF collections (MMF 21163) caught off in the coast of Madeira (stomach content) which is synonym of *N. scolopaceus* a second holotype (unique and lost) of *Leptorhynchus leuchtenbergi* Lowe 1851 caught off in Madeira is also synonym of *N. scolopaceus*.

Family Serrivomeridae

Serrivomer beanii Gill & Ryder 1883

Common name: **Bean's sawtooth eel; Cobra-couraça.**

References: mainland Portugal ([MOM-IC-0091-1216, 1910](#); [Zugmayer, 1911a, b](#); [YPM ICH 005169, 1959](#); [MCZ 65835, 1969](#); [USNM 315426, 1970](#); [Bauchot, 1986c](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1884-1070, 1883](#); [Vaillant, 1888](#); [Goode and Bean, 1895, 1896](#); [Perrier, 1903](#); [Brauer, 1906](#); [Roule and Bertin, 1929](#); [Roule and Angel, 1931, 1933](#); [Ferreira, 1938](#); [Collins, 1954](#); [Bauchot-Boutin, 1955](#); [Post and Tesch, 1982](#); [Bauchot, 1986c](#); [Tighe, 1989](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Bauchot et al., 1993](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Smith, 2016e](#); [Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Bauchot, 1986c](#); [Lloris et al., 1991](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#);

Smith, 2016e; Delgado *et al.*, 2017; Froese and Pauly, 2019). Two unique holotypes of *Nemichthys richardi* Vaillant 1888 and *Gavialiceps hasta* Zugmayer 1911 were caught off in the Azores (MNHN-IC-1884-1070) and west of the Azores (MOM-IC-0091-1216), respectively. *N. richardi* is synonym of *S. beanii* and *G. hasta* is uncertain as *S. beanii*.

***Serrivomer lanceolatoides* (Schmidt 1916)**

Common name: Bronze sawtooth eel; **Cobra-de-dentes-curtos**.

References: mainland Portugal ([Bauchot, 1986c](#); MCZ 159543, 1986; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores ([Post and Tesch, 1982](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Sutton *et al.*, 2008](#); [Porteiro *et al.*, 2010, 2017](#); Carneiro *et al.*, 2014; Smith, 2016e; Froese and Pauly, 2019) and Madeira ([Bauchot, 1986c](#); Carneiro *et al.*, 2014; Smith, 2016e; Porteiro *et al.*, 2017; Froese and Pauly, 2019).

Family Anguillidae

***Anguilla anguilla* (Linnaeus 1758)**

Common name: **European eel**; **Enguia**, Anguia, Anguila, Angula, Angulha, Brasino, Brazino, Civela, Cobra, Eirão, Eiro, Eiró, Eiró-do-mar, Eirol, Eirós, Enguia-europeia, Enguia-de-vidro, Enguia-do-rio, Enguia-grande, Enguia-macha, Enguia-macho, Enguia-prateada, Eró, Iró, Irós, Loura, Machão, Machinho, Meixão, Mexão, Pibala, Torneira, Traça, Tração.

References: mainland Portugal▲ ([Carvalho, 1765](#); [de Brito Capello, 1869a](#), 1880; [Osório, 1888](#), 1895; [Baldaque da Silva, 1891](#); [Vieira, 1894a](#), 1897, 1901; [Nobre, 1895a](#), c, 1903, 1931, 1932, 1935; [Bragança, 1903](#); [Seabra, 1911](#); [Gandolfi-Hornyold, 1914-16a](#), b; [Gonçalves, 1941](#); [Hellinng, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#), 1967; [Almaça, 1965b](#); [Bauchot, 1986d](#); [Fischer *et al.*, 1987](#); [Antunes *et al.*, 1988](#); [Arruda *et al.*, 1988](#); [Arruda, 1990](#); [Gordo *et al.*, 1990](#); [Smith, 1990c](#); [Cabral, 1999](#); [Garnerot *et al.*, 2004](#); ICES, 2005; [Ribeiro *et al.*, 2006, 2008, 2012](#); [Henriques *et al.*, 2008](#); [Neves *et al.*, 2008](#); [Veiga *et al.*, 2009](#); [Gonçalves *et al.*, 2010](#); [França *et al.*, 2011](#); [Carneiro *et al.*, 2014](#); [Smith, 2016a](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)), Azores ([Drouët, 1858](#), 1861a, b; [Ramos, 1869](#); [Günther, 1870](#); [Hilgendorf, 1888a](#); [Vaillant, 1888](#), 1898; [Collett, 1896a](#); [Sampaio, 1904](#); [Lea, 1913](#); [Nobre, 1924](#); [D'Ancona, 1929](#); [Fowler, 1936a](#); [Ferreira, 1934](#), 1938; [Ege, 1939](#); [Poll, 1947](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Wheeler, 1978](#); [Martins, 1982a](#); [Bauchot, 1986d](#); [Smith, 1990c](#); [Azevedo *et al.*, 1991](#), 1992; [Lloris *et al.*, 1991](#); [Arruda, 1992](#), 1997; [Azevedo, 1995](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1837](#); [Vaillant, 1919](#); [Noronha and Sarmento, 1934](#), 1948; [Nobre, 1935](#); [Fowler, 1936a](#); [Nunes, 1953](#), 1974; [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#), 1967; [Fischer *et al.*, 1981](#); [Bauchot, 1986d](#); [Smith, 1990c](#); [Lloris *et al.*, 1991](#); [Wirtz *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Smith, 2016a](#); [Froese and Pauly, 2019](#)).

***Anguilla rostrata* (Lesueur 1817)**

Common name: **American eel**; **Enguia-americana**.

References: mainland Portugal (PECS) ([ICES EEL 8302450, 1984](#)) and Azores (PECS) ([MCZ 160463, 1986](#); [Carneiro *et al.*, 2014](#)). *Anguilla rostrata* is widely distributed from the Northwest to western Central Atlantic, including Greenland (Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). However, the occurrence records of this species in the Eastern Atlantic are very rare.

Order Saccopharyngiformes

Family Cyematidae

***Cyema atrum* Günther 1878**

Common name: **Bobtail eel**; **Cobra-de-leme**.

References: mainland Portugal ([Bertin, 1937](#); [Albuquerque, 1954-1956](#); [YPM ICH 005163, 1959](#); [YPM ICH 005165, 1959](#); [YPM ICH 005166, 1959](#); [YPM ICH 005167, 1959](#); [Osório de Castro, 1967](#); [Saldanha and Bauchot, 1986b](#); [Carneiro *et al.*, 2014](#)), Azores ([Zugmayer, 1911b](#); [Murray and Hjort, 1912](#); [Lea, 1913](#); [Roule and Bertin, 1929](#); [Fowler, 1936a](#), b; [Bertin, 1937](#); [Grey, 1956](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Saldanha and Bauchot, 1986b](#); [Lloris *et al.*, 1991](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Vaillant, 1919](#); [Roule and Angel, 1930](#); [Zugmayer, 1933](#); [Fowler, 1936a,b](#); [Bertin, 1937](#); [Albuquerque, 1954-1956](#);

NHMUK 1994.11.1.3732, 1962; NHMUK 1994.11.1.3733, 1962; Osório de Castro, 1967; Maul, 1971a; Saldanha and Bauchot, 1986b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Family Monognathidae

Monognathus bertini Bertelsen & Nielsen 1987

Common name: Bertin's one-jawed eel; **Enguia-da-fundura-de-Bertin**.

References: mainland Portugal ([MCZ 64662](#), 1986; Carneiro *et al.*, 2014).

Monognathus boehlkei Bertelsen & Nielsen 1987

Common name: Böhlke's one-jawed eel; **Enguia-da-fundura-de-Böhlke**.

References: Azores (PECS) ([NHMUK 1984.12.3.2](#), 1973; Smith, 2016f(?)) and Madeira ([Bertelsen and Nielsen, 1987](#); Carneiro *et al.*, 2014).

Monognathus jesperseni Bertin 1936

Common name: No English common name; **Enguia-da-fundura-de-Jespersen**.

References: mainland Portugal ([NHMUK 1987.2.3.2](#), 1978). It is a very rare species.

Monognathus herringi Bertelsen & Nielsen 1987

Common name: Herring's one-jawed eel; **Enguia-da-fundura-de-Herring**.

References: Azores (PECS) ([NHMUK 1987.2.3.1](#), 1981; Bertelsen and Nielsen, 1987; Ralph, 2015b; Smith, 2016f(?)) and Madeira (?) ([Carneiro et al., 2014](#)). Bertelsen and Nielsen (1987) mentioned the occurrence of this species in west of Madeira (34°58.1'N, 32°56.5'W; holotype NHMUK 1987.2.3.1, 1981) but these geographic coordinates correspond to a position in the Mid-Atlantic Ridge, SW of Azores Islands (PECS area). Thereby, this species was erroneously included as presented in Madeira by Carneiro *et al.* (2014). Thus, this species is not considered in the total number of valid species to Madeira.

Monognathus nigeli Bertelsen & Nielsen 1987

Common name: Nigel's one-jawed eel; **Enguia-da-fundura-de-Nigel**.

References: Azores (PECS) ([ZMUC P2340540](#), 1981; Quéro *et al.*, 2003; Carneiro *et al.*, 2014; Smith, 2016f(?)).

Family Eurypharyngidae

Eurypharynx pelecanoides Vaillant 1882

Common name: **Pelican eel**; **Peixe-pelicano**, Pelicano, Enguia-pelicano.

References: mainland Portugal ([MNHN-IC-1912-0027](#), 1910; Nielsen and Bertelsen, 1986; Carneiro *et al.*, 2014; Smith, 2016f; Porteiro *et al.*, 2017; Froese and Pauly, 2019), Azores (Murray and Hjort, 1912; Roule, 1916; Roule and Angel, 1933; Bertin, 1934, 1938; Fowler, 1936b; Ferreira, 1938; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Post and Tesch, 1982; Nielsen and Bertelsen, 1986; Arruda, 1997; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Smith, 2016f; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira ([MOM-IC-0091-0592](#), 1901; Zugmayer, 1911b; Roule, 1914a, b, 1919a; Fowler, 1936a, b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1971a; Bauchot, 1979; Nielsen and Bertelsen, 1986; Lloris *et al.*, 1991; Davies and Merrett, 1998; Carneiro *et al.*, 2014; Smith, 2016f; Froese and Pauly, 2019). The holotype of *Eurypharynx richardi* Roule 1914, synonym of *E. pelecanoides*, was caught off south west of Madeira and preserved in the MOM collections but not found in 1995.

Family Saccopharyngidae

Saccopharynx ampullaceus (Harwood 1827)

Common name: Northern swallower eel; **Peixe-pelicano-do-Norte**, Enguia-pelicano.

References: mainland Portugal ([Girard, 1895](#); Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Bauchot, 1979; Bertelsen and Nielsen, 1986; Carneiro *et al.*, 2014; Smith, 2016f; Froese and Pauly, 2019), Azores ([Harwood, 1827 \(PECS\)](#); Roule and Angel, 1933; Fowler, 1936b; Albuquerque, 1954-1956; Böhlke, 1966; Osório de Castro, 1967; Nielsen and Bertelsen, 1985; Bertelsen and Nielsen, 1986; Krefft, 1987; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017; Carneiro

et al., 2014; Smith, 2016f; Froese and Pauly, 2019) and Madeira (NHMUK 1861.6.4.3., 1854 - 1861; Johnson, 1862c; Günther, 1870, 1887; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Bauchot, 1979; Bertelsen and Nielsen, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Smith, 2016f; Froese and Pauly, 2019). The lectotype of *Saccopharynx johnsoni* Bertin 1934, synonym of *S. ampullaceus*, was caught off in Madeira and is currently in the NHMUK collections (NHMUK 1861.6.4.3).

***Saccopharynx harrisoni* Beebe 1932**

Common name: No English common name; **Peixe-pelícano-de-Harrison**.

References: Madeira (Bertelsen and Nielsen, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2019) the current geographic distribution of *S. harrisoni* is the Western Atlantic. The distribution map of this species, reported by Bertelsen and Nielsen (1986), indicates the occurrence of this species in Madeira.

***Saccopharynx hjorti* Bertin 1938**

Common name: No English common name; **Peixe-pelícano-de-Hjort**.

References: Azores (PECS) (Bertin, 1938; Arruda, 1997; Carneiro *et al.*, 2014). The holotype (unique) of this species is in the ZMB collections (ZMB 6041) caught off SW of the Azores (34°59.0'N, 33°01.0'W).

***Saccopharynx paucovertebratis* Nielsen & Bertelsen 1985**

Common name: No English common name; **Peixe-pelícano-dos-Açores**.

References: Madeira (?) (Nielsen and Bertelsen, 1985, 1986; Carneiro *et al.*, 2014).

The holotype (unique) of this species is in the ZMUC collections (ZMUC P239898) caught off near the Azores. Bertelsen and Nielsen (1985) describe for the first time this species and indicate the occurrence near the Azores, north-eastern Atlantic, 32°56.0'N, 23°47.0'W. The same authors mention in 1986 that the only known specimen was caught off west of Madeira (Bertelsen and Nielsen, 1986). Both citations referred to the same specimen, and the geographic coordinates correspond to a position off the Azores, out of the PECS area and was erroneously included as presented in Madeira (Carneiro *et al.*, 2014). Thus, this species is not considered in the total number of valid species.

***Saccopharynx ramosus* Nielsen & Bertelsen 1985**

Common name: Shortline swallower eel; **Peixe-pelícano-de-cauda-ramosa**.

References: Azores (PECS) (NHMUK 1983.11.19.2, 1982; Carneiro *et al.*, 2014) and Madeira (?) (Bertelsen and Nielsen, 1986; Carneiro *et al.*, 2014; Smith, 2016f; Froese and Pauly, 2019). According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is the Atlantic and Indian Ocean. Bertelsen and Nielsen (1985) describe for the first time this species and indicate its occurrence in the North-eastern Atlantic, 30°18.4'N, 26°39.9'W and in 1986 they mention two records of *S. ramosus* in West of Madeira (Bertelsen and Nielsen, 1986). Smith (2016f) reports the occurrence of this species in the eastern-central Atlantic, known from SW of Madeira. Probably, the three citations referred to the same specimens (holotype: NHMUK 1983.11.19.2 and paratypes: SAM 23423) and the geographic coordinates correspond to a position south of the Azores, out of the PECS area (90 NM east of the Great Meteor Tablemount). Therefore, the validity of the Madeira records are questionable and specimen are needed for confirmation. Thus, this species is not considered in the total number of valid species to Madeira.

***Saccopharynx thalassa* Nielsen & Bertelsen 1985**

Common name: Hairytail swallower eel; **Peixe-pelícano-de-cauda-peluda**.

References: Madeira (Bertelsen and Nielsen, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Smith, 2016f; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Saccopharynx trilobatus* Nielsen & Bertelsen 1985**

Common name: No English common name; **Peixe-pelícano-de-cauda-trilobada**.

References: Azores (PECS) (NHMUK 1983.11.19.1, 1981; Nielsen and Bertelsen, 1985; Arruda, 1997; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of this species is in the NHMUK collections (NHMUK 1983.11.19.1.) and caught off SW of the Azores (32°58.0'N, 33°14.6'W).

Order Clupeiformes

Family Clupeidae

***Alosa alosa* (Linnaeus 1758)**

Common name: **Allis shad; Sável**, Saboga, Savaleta, Savalete, Save, Savelha, Savelinha.

References: mainland Portugal▲ ([Carvalho, 1756](#); [Vandelli, 1771, 1797](#); [de Brito Capello, 1869a, 1880](#); [Steindachner, 1868b](#); [Osório, 1888, 1896](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1893, 1894a, 1897, 1901](#); [Nobre, 1894, 1931, 1932, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979b](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Whitehead, 1984](#); [Dinis, 1985](#); [Dooley et al., 1985](#); [Martins et al., 1992, 2015](#); [Jorge et al., 2002](#); [Santos et al., 2002a, c](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Kottelat and Freyhof, 2007](#); [Henriques et al., 2008](#); [Ribeiro et al., 2008](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Munroe, 2016a](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)).

***Alosa fallax* (Lacepède 1803)**

Common name: **Twaite shad; Savelha**, Junqueira, Sabaia, Sabega, Sabelha, Saboga, Sabóia, Saramugo, Savaleta, Savatela, Sável, Savelinha, Savilhão, Sebatela, Sevelha.

References: mainland Portugal▲ ([de Brito Capello, 1869a, 1880](#); [Gervais and Boulart, 1876](#); [Osório, 1888, 1895, 1896](#); [Almeida and Roquette, 1892](#); [Nobre, 1895c, 1903, 1931, 1932, 1935](#); [Vieira, 1897, 1901](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Almaça, 1965b](#); [Svetovidov, 1979b](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982a, b](#); [Martins, 1981](#); [Whitehead, 1984](#); [Dinis, 1985](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Collares-Pereira et al., 2000](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Ribeiro et al., 2008, 2009](#); [Gonçalves et al., 2010](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Munroe, 2016a](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Barreiros et al., 2014](#)) and Madeira ([MNHN-IC-2005-2408, 1925](#); [Carneiro et al., 2014](#)).

***Clupea harengus* Linnaeus 1758**

Common name: **Atlantic herring; Arenque-do-Atlântico**, Arenque.

References: mainland Portugal▲ ([Almeida and Roquette, 1892](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *C. harengus* is the North Atlantic and adjacent Arctic areas (including Baltic Sea, North Sea). Albuquerque (1954-1956) and Osório de Castro (1967) reported the rare occurrence of this species in mainland Portugal.

***Harengula clupeola* Valenciennes, 1847**

Common name: **False herring; Sardinha-escamuda**, Espadilha.

References: mainland Portugal (?) ([Nobre, 1895c, 1935](#); [Osório, 1895](#); [Seabra, 1911](#)). Nobre (1895c) reported for the first time the occurrence of *Harengula latulus* Valenciennes, 1847, synonym of *H. clupeola*, caught off in Matosinhos in June 1893. Nobre (1935) also mentioned that this species is uncommon and appears along with sardine (*S. pilchardus*). However, according to Eschmeyer (2019) the current geographic distribution of *H. clupeola* is the Western Atlantic. Nevertheless, in the fish collections of the Museum national d'Histoire naturelle, there are nine preserved specimens of this species caught off NW Galicia and Normandy coast. Five of these specimens were designated as paralectotypes of *H. clupeola* [MNHN-IC-0000-3236, 1827 (Whitehead, 1967)]. Since Nobre (1935), no other author described the presence of this species in mainland Portugal and its occurrence in this EEZ was not yet confirmed by more recent data. Thus, this species is not considered in the total number of valid species.

***Sardina pilchardus* (Walbaum 1792)**

Common name: **European pilchard; Sardinha**, Arenque-da-costa-larga, Costeleta-na-grelha, Esquilha, Majuga, Manjua, Manjunga, Petinga, Salpicadinho-da-costa, Sardinha-de-lei, Sardinha-do-puto, Sardinha-do-reino, Sardinha-europeia, Sardinha-legítima, Sardinha-portuguesa, Sardinha-real, Sardinha-verdadeira, Sardinheta, Viva-da-costa, Vivinha, Vivinha-da-costa.

References: mainland Portugal[▲] ([Steindachner, 1868b](#); [de Brito Capello, 1869a, 1880](#); [Gervais and Boulart, 1877a](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1893, 1897, 1901](#); [Nobre, 1895a, c, 1903, 1935](#); [Osório, 1895](#); [Bragança, 1902](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979b](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [INIP, 1981a, 1982a, b](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Whitehead, 1984, 1985](#); [Dinis, 1985](#); [Dooley et al., 1985](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Erzini et al., 1996, 2001a](#); [Cabral, 1999](#); [Gomes et al., 2001](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2006](#); [Stergiou et al., 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Neves et al., 2008](#); [Ribeiro et al., 2008, 2012](#); [Batista et al., 2009](#); [Garrido et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Munroe, 2016a](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Vaillant, 1919](#); [Fage, 1920](#); [Ramalho, 1929a](#); [Fowler, 1936a](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Martins, 1982a](#); [Isidro, 1988, 1990 a, b, 1996](#); [Nash et al., 1990, 1994a](#); [Patzner et al., 1990, 1992](#); [Ré, 1990](#); [Arruda, 1992, 1997](#); [Santos et al., 1994b, 1997](#); [Azevedo, 1995](#); [Pinho et al., 1995](#); [Santos and Nash, 1995](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Diogo and Pereira, 2014](#); [Afonso, 2016](#); [Fauconnet et al., 2019](#) and Madeira[▲] ([Günther, 1868](#); [Lowe, 1837, 1843, 1844b](#); [de Velloso, 1892](#); [Regan, 1916](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1948a](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979b](#); [Fischer et al., 1981](#); [Whitehead, 1984, 1985](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Munroe, 2016a](#); [Froese and Pauly, 2019](#)). The whereabouts of the syntype of *Clupea laticosta* Lowe 1843, synonym of *S. pilchardus*, caught off Madeira are unknown.

***Sardinella aurita* Valenciennes 1847**

Common name: **Round sardinella; Sardinela-lombuda**, Arenque, Lombuda, Maromba, Sardinela, Sardinha, Sardinha-arinca, Sardinha-lombuda.

References: mainland Portugal ([SAIAB FC 11253, n.d.](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a](#); [Martins, 1981](#); [Whitehead, 1984](#); [Gourène and Teugels, 1990](#); [Borges et al., 2002](#); [ICES, 2005](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)), Azores (?) ([Osório de Castro, 1967](#)) and Madeira ([ANSP 95591, n.d.](#); [MNHN-IC-A-6228, 1882](#); [Fowler, 1923, 1936a](#); [Osório de Castro, 1967](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *S. aurita* is the Western and Eastern Atlantic, including Mediterranean Sea and Black Sea. The only record of this species in the fishing grounds of the Azores was indicated by Osório de Castro (1967). The presence of this species in Pico (Azores), in 2009, was reported (pers. comm. Magnus Lundgren), but its occurrence in the Azores region is questionable and needs further documentation.

***Sardinella maderensis* (Lowe 1838)**

Common name: **Madeiran sardinella; Sardinela-da-Madeira**, Arenque, Palheta, Sardinela, Sardinha, Sardinha-arinca, Sardinela-palheta.

References: mainland Portugal ([Whitehead, 1984](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#)), Azores[▲] ([Günther, 1868](#); [Regan, 1903a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Arruda, 1992, 1997](#); [Santos et al., 1997\(?\)](#); [Carneiro et al., 2014](#)) and Madeira[▲] ([Lowe, 1837](#); [NHMUK 1855.12.26.496-500, 1854-1861](#); [Von Martens, 1876](#); [Günther, 1868](#); [de Velloso, 1892](#); [Regan, 1917](#); [Fowler, 1936a](#); [Maul, 1948a](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [MNHN-IC-1956-0059, 1956](#); [Fischer et al., 1981](#); [Whitehead, 1984](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Munroe, 2016a](#); [POGIS, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The syntypes of *Clupea maderensis* Lowe 1838, synonym of *S. maderensis*, is in the NHMUK collections (NHMUK 1855.12.26.496-500, set of five specimens) caught off Madeira. According to Eschmeyer et al. (2019) the current geographic distribution of *S. maderensis* includes the Mediterranean Sea and, in the Eastern Atlantic, from Gibraltar south to Angola, including Madeira and Canary islands. Santos et al. (1997) mention that there is no

clear demonstration of the occurrence of the species in the Azores and the presence of this species in the region needs further documentation.

***Sprattus sprattus* (Linnaeus 1758)**

Common name: **European sprat; Espadilha**, Espadinha, Lampadinha, Lavadilha, Lavadinha, Lavalinha, Navalhinha, Navalhinho, Navalinha.

References: mainland Portugal▲ ([Vandelli, 1771](#); [Osório, 1895](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#); [1967](#); [Svetovidov, 1979b](#); [Fernández et al., 1981](#); [Martins, 1981](#); [INIP, 1982b](#); [Whitehead, 1984](#), [1985](#); [Dinis and Marecos 1984](#); [Dinis, 1985](#); [Fischer et al., 1987](#); [Arruda et al., 1988](#); [Jorge et al., 2002](#); [ICES, 2005](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Martins, 1982a](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#)) and Madeira ([Noronha and Sarmento, 1934](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#)). Several earlier literature citations indicate the presence of this species in the archipelagos of Madeira and the Azores. However, there is no clear demonstration of the occurrence records of this species in these regions. The occurrence of this species in the Azores and Madeira archipelagos is questionable and specimens are needed for confirmation.

Family Engraulidae

***Engraulis encrasicolus* (Linnaeus 1758)**

Common name: **European anchovy; Biqueirão**, Anchova, Biqueira, Biqueirão-branco, Biqueirão-negro, Boca-torta, Boqueirão, Chacareu, Chacaréu, Enchova, Mancatarina.

References: mainland Portugal▲ ([Carvalho, 1765\(?\)](#); [de Brito Capello, 1869a](#), [1880](#); [Gervais and Boulart, 1877a](#); [Vieira, 1893](#), [1897](#), [1901](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Nobre, 1895c](#), [1903](#), [1935](#); [Osório, 1895](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#), [1967](#); [Svetovidov, 1979c](#); [Fernández et al., 1981](#); [Fischer et al., 1981](#), [1987](#); [INIP, 1981a](#), [1982a](#), [b](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Whitehead, 1984](#), [1990](#); [Dinis, 1985](#); [Dooley et al., 1985](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Whitehead et al., 1988](#); [Martins et al., 1992](#), [1999](#), [2015](#); [Cabral, 1999](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2006](#); [Gomes and Olim, 2007](#); [Chaves, 2008](#), [2009](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Ribeiro et al., 2008](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Nizinski and Munroe, 2016](#); [Ryland and Lancaster, 2017](#); [Golani and Fricke, 2018](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Ramos, 1869](#); [Martins, 1982a](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#); [Nizinski and Munroe, 2016](#)) and Madeira ([Maul, 1948a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Whitehead, 1984\(?\)](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Nizinski and Munroe, 2016](#); [Biscoito et al., 2017](#); [Golani and Fricke, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Order Alepocephaliformes

Family Alepocephalidae

***Alepocephalus agassizii* Goode & Bean 1883**

Common name: Agassiz' slickhead; **Celindra-negra-de-Agassiz**.

References: mainland Portugal ([Mecklenburg et al., 2018](#)) and Azores ([Arruda, 1997](#); [MAR-ECO 2725](#), [2004](#); [MAR-ECO 3417](#), [2004](#); [MAR-ECO 9734](#), [2004](#); [MAR-ECO 10924](#), [2004](#); [MAR-ECO 2539](#), [2004](#); [MAR-ECO 13389](#), [2004](#); [Bergstad et al., 2008a](#) (PECS); [Carneiro et al., 2014](#); [Nielsen et al., 2015a](#); [Porteiro et al., 2017](#); [Mecklenburg et al., 2018](#)). The distribution map reported by Mecklenburg et al. (2018) indicated the occurrence of this species in mainland Portugal.

***Alepocephalus australis* Barnard 1923**

Common name: **Small scaled brown slickhead**; **Celindra-austral**.

References: Azores ([Arruda, 1997](#); [Melo and Menezes, 2002](#); [MAR-ECO 8097](#), [2004](#); [Bergstad et al., 2008a](#) (PECS); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#); [Almeida and Biscoito, 2019](#)).

Alepocephalus bairdii Goode & Bean 1879

Common name: **Baird's slickhead**; **Celindra-de-Baird**, Celindra, Triste-linda.

References: mainland Portugal[▲] ([Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976](#); [Krefft, 1979b](#); [Markle and Quéro, 1984](#); [Pakhorukov, 2008](#); [Bordalo-Machado et al., 2009](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Melo and Menezes, 2002](#); [Pakhorukov, 2008](#) (PECS); [Porteiro et al., 2010](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#)) and Madeira[▲] ([Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). Two specimens caught off at coast of mainland Portugal are in the MMF collections (MMF 22500, 1967, set of two specimens).

Alepocephalus productus Gill 1883

Common name: **Smalleye smooth-head**; **Celindra-de-olho-pequeno**.

References: mainland Portugal ([Markle and Quéro, 1984](#); [Carneiro et al., 2014](#); [Uiblein et al., 2015a](#); [Quéro et al., 2019](#)), Azores ([IMAR/DOP COLETA 1688, 2001](#); [Melo and Menezes, 2002](#); [Bergstad et al., 2008a](#) (PECS); [Porteiro et al., 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Uiblein et al., 2015a](#)) and Madeira ([Freitas et al., 2013](#)). Markle and Quéro (1984) mention the caught of three specimens of this species off Morocco (35°46.0'N, 8°16.0'W) but this geographic coordinates correspond to a position at mainland Portugal EEZ.

Alepocephalus rostratus Risso 1820

Common name: **Risso's smooth-head**; **Celindra-de-Risso**, Celindra, Triste-linda.

References: mainland Portugal ([Maul, 1976](#); [Krefft, 1979b](#); [Markle and Quéro, 1984](#); [MNHN-IC-1988-0500, 1984](#); [MNHN-IC-1988-0493, 1984](#); [Gordo et al., 1990](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Fabri et al., 2016](#); [Hartel and Orrell, 2016c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([MNHN-IC-1885-0157, 1833](#); [Vaillant, 1888, 1919](#); [Goode and Bean, 1895, 1896](#); [Perrier, 1903](#); [Brauer, 1906](#); [Murray and Hjort, 1912](#); [Koefoed, 1927](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [Maul, 1976](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Menezes et al., 2006, 2012](#); [Porteiro et al., 2010](#); [Besugo, 2013](#); [Carneiro et al., 2014](#)) and Madeira[▲] ([Johnson, 1862c](#); [Fowler, 1936a](#); [Maul, 1948a, 1976](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [INIP, 1980](#); [Severino, 2004](#); [Carneiro et al., 2014](#)). Five specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22465, 1967, set of five specimens).

Asquamiceps caeruleus Markle 1980

Common name: No English common name; **Celindra-de-cabeça-azul**.

References: mainland Portugal ([Hulley, 2015a](#)) and Madeira ([Hulley, 2015a](#)). In the Eastern Atlantic this species is distributed from 30°N to 21°S ([Hartel and Orrell, 2016b](#)) and according to Eschmeyer et al. (2019) *A. caeruleus* is a cosmopolitan species. The distribution map of this species reported by Hulley (2015a) included parts of mainland Portugal and the Madeira EEZ. However, no recent occurrence records have been reported for mainland Portugal or Madeira Island. Specimens are needed for confirmation.

Asquamiceps hjorti (Koefoed 1927)

Common name: Barethroat slickhead; **Celindra-de-Hjort**.

References: Azores (PECS) ([Koefoed, 1927](#); [Frøiland, 1979](#); [Markle, 1980](#); [Carneiro et al., 2014](#)). The holotype of this species is in the USNM collections (USNM 3335, 1910) caught off SW of the Azores (34°59'N, 33°1.0'W).

Asquamiceps velaris Zugmayer 1911

Common name: Fanfin smooth-head; **Celindra-de-leque**.

References: mainland Portugal ([MOM-IC-0091-1198, 1910](#); [Zugmayer, 1911a](#); [Belloc, 1949](#); [Markle, 1980](#); [Markle and Quéro, 1984](#); [Markle and Sazonov, 1990](#); [Carneiro et al., 2014](#)). The holotype (unique) of this species is in the MOM collections (MOM-IC-0091-1198) caught off mainland Portugal (36° 06' N, 9° W).

Bajacalifornia megalops (Lütken 1898)

Common name: Bigeye smooth-head; **Celindra-de-olho-grande**.

References: mainland Portugal ([Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [Nielsen and Hulley, 2015a](#)) and Azores ([ZMH 119101, 1979](#); [Markle and Quéro, 1984](#); [Krefft, 1985](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Bergstad et al., 2008a \(PECS\)](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2010, 2017](#); [Nielsen and Hulley, 2015a](#); [Froese and Pauly, 2019](#)) and Madeira ([Nielsen and Hulley, 2015a](#)). One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005304).

Bathylaco nigricans Goode & Bean 1896

Common name: Black warrior; **Celindra-negra**.

References: mainland Portugal ([Knudsen et al., 2015a](#)), Azores ([Hureau and Monod, 1979](#); [Markle and Quéro, 1984](#); [Krefft, 1985](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Knudsen et al., 2015a](#); [Froese and Pauly, 2019](#)) and Madeira ([Hureau and Monod, 1979](#); [Nielsen, 1979a](#); [Lloris et al., 1991](#); [NHMUK 2016.8.25.757-771, 1995](#); [Carneiro et al., 2014](#); [Knudsen et al., 2015a](#)).

Bathyprion danae Marshall 1966

Common name: **Fangtooth smooth-head**; **Celindra-de-Dana**.

References: mainland Portugal ([Nielsen and Hulley, 2015b](#)), Azores (PECS) ([MAR-ECO 6957, 2004](#); [MAR-ECO 5390, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Nielsen and Hulley, 2015b](#); [Porteiro et al., 2017](#)) and Madeira ([Markle, 1976](#); [Hureau and Monod, 1979](#); [Krefft, 1979b](#); [Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Nielsen and Hulley, 2015b](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)).

Bathytroctes macrolepis Günther 1887

Common name: **Koefoed's smooth-head**; **Celindra-escamuda**.

References: mainland Portugal ([NHMUK 1991.7.9.863, 1989](#); [Uiblein et al., 2015b](#)), Azores ([Parr 1951 \(PECS\)](#); [MCZ 36636, 1947](#); [Markle and Quéro, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#); [Uiblein et al., 2015b](#); [Froese and Pauly, 2019](#)) and Madeira ([Uiblein et al., 2015b](#)).

Bathytroctes michaelsarsi Koefoed 1927

Common name: **Michael Sars smooth-head**; **Celindra-de-Michael-Sars**.

References: mainland Portugal ([Markle and Quéro, 1984](#); [Carneiro et al., 2014](#); [Nielsen et al., 2015b](#); [Froese and Pauly, 2019](#)), Azores ([MAR-ECO 2697, 2004](#); [MAR-ECO 3403, 2004](#); [Bergstad et al., 2008a \(PECS\)](#); [Carneiro et al., 2014](#); [Nielsen et al., 2015b](#); [Porteiro et al., 2017 \(PECS\)](#)) and Madeira ([Nielsen et al., 2015b](#)).

Bathytroctes microlepis Günther 1878

Common name: **Smallscale smooth-head**; **Celindra-de-escama-pequena**.

References: mainland Portugal ([NHMUK 1887.12.7.233, 1873](#); [Günther, 1878](#); [Brauer, 1906](#); [MOM-IC-0091-1145, 1910](#); [Zugmayer, 1911a](#); [Nobre, 1935](#); [Grey, 1956](#); [Krefft, 1979b](#); [Markle and Quéro, 1984](#); [NHMUK 1991.7.9.862, 1989](#); [Carneiro et al., 2014](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)), Azores ([Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Krefft, 1979b](#); [Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species was caught off in south-eastern of Cape Vincent, 35°47.0'N, 8°23.0'W (mainland Portugal) and is in the NHMUK collections (NHMUK 1887.12.7.233). A second holotype (unique) of *Bathytroctes grimaldii* Zugmayer 1911, synonym of *B. microlepis*, was caught off in mainland Portugal (37°38.0'N, 10°53.0'W) and is in the MOM collections (MOM-IC-0091-1145).

Conocara fiolenti Sazonov & Ivanov 1979

Common name: **Fiolenti's smooth-head**; **Celindra-de-Fiolenti**.

References: Azores ([Markle and Quéro, 1984](#); [Crabtree and Sulak, 1986](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Conocara macropterum (Vaillant 1888)

Common name: **Longfin smooth-head**; **Celindra-de-barbatana-comprida**, Celindra, Triste-linda.

References: mainland Portugal ([Markle and Quéro, 1984; MNHN-IC-1988-0497, 1984; Carneiro et al., 2014; Hulley, 2015b; Porteiro et al., 2017; Froese and Pauly, 2019](#)), Azores ([Roule, 1916, 1919; Roule and Angel, 1933; Fowler, 1936a; Collins, 1954; Albuquerque, 1954-1956; Krefft, 1979b; Markle and Quéro, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Melo and Menezes, 2002; Bergstad et al., 2008a \(PECS\); Porteiro et al., 2010, 2017 \(PECS\); Carneiro et al., 2014; Hulley, 2015b; Froese and Pauly, 2019](#)) and Madeira ([Hulley, 2015b](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *C. macropterus* is the Atlantic ocean. The distribution map reported by Hulley (2015b) indicated the occurrence of this species in Madeira and Selvagens Islands. Its presence in the Madeira fauna has not been documented. The occurrence of this species in the Madeira Archipelago is questionable and needs further documentation.

***Conocara murrayi* (Koefoed 1927)**

Common name: Murray's smooth-head; **Celindra-de-Murray**.

References: mainland Portugal ([Koefoed, 1927; Markle and Quéro, 1984; Crabtree and Sulak, 1986; Carneiro et al., 2014; Knudsen et al., 2015b; Froese and Pauly, 2019](#)), Azores ([NHMUK 2009.7.20.8, 1970; ZMH 121217, 1982 \(PECS\); Markle and Quéro, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; MAR-ECO 3962, 2004 \(PECS\); Bergstad et al., 2008a \(PECS\); Porteiro et al., 2010, 2017 \(PECS\); Carneiro et al., 2014; Knudsen et al., 2015b; Froese and Pauly, 2019](#)) and Madeira ([Lloris et al., 1991; Knudsen et al., 2015b](#)).

***Conocara salmonicum* (Gill & Townsend 1897)**

Common name: Salmon smooth-head; **Celindra-salmonada**.

References: mainland Portugal ([Crabtree and Sulak, 1986\(?\)](#); Christiansen et al., 2014) and Madeira ([NHMUK 2016.8.25.752, 1995](#)).

***Conocara werneri* Nybelin 1946**

Common name: Werner's smooth-head; **Celindra-de-Werner**.

References: mainland Portugal ([GNM 1946-6587, 1946; Carneiro et al., 2014](#)). The holotype (unique) of this species is in the NHMG collections (NHMG Pi. 1300) caught off south of Cape Santa Maria, 35°43.0'N, 8°16.0'W (mainland Portugal).

***Einara edentula* (Alcock 1892)**

Common name: **Toothless smooth-head; Celindra-desdentada**, Celindra.

References: Azores ([MNHN-IC-1987-0241, 1969; Geistdoerfer et al., 1971; Markle, 1976; Hureau and Monod, 1979; Markle and Quéro, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019](#)) and Madeira ([Roule and Angel, 1931; Osório de Castro, 1967\(?\)](#); Markle and Quéro, 1984; Lloris et al., 1991; [NHMUK 1996.2.14.26-27, 1995; Carneiro et al., 2014; Froese and Pauly, 2019](#)). The holotype (unique) of *Alepocephalus edentulus atlanticus* Roule & Angel 1931, synonym of *E. edentula*, was caught off between Madeira and southern coast of Moroccois and is in the MOM collections (MOM-IC-0091-1352).

***Einara macrolepis* (Koefoed 1927)**

Common name: **Loosescale smooth-head; Celindra-de-escama-caduca**.

References: Azores ([Arruda, 1997; MAR-ECO 5627, 2004; Sutton et al., 2008; Carneiro et al., 2014; Porteiro et al., 2017 \(PECS\)](#)) and Madeira ([ZMH 109373, 1971; Krefft, 1979b; Markle and Quéro, 1984; Lloris et al., 1991; Carneiro et al., 2014; Hartel and Orrell, 2016c; Froese and Pauly, 2019](#)).

***Herwigia krefftii* (Nielsen & Larsen 1970)**

Common name: **Krefft's smooth-head; Celindra-de-Krefft**.

References: Azores ([Tortonese and Hureau, 1979; Arruda, 1997; Santos et al., 1997; Markle and Quéro, 1984; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1996.2.14.13, 1995; NHMUK 2004.2.3.61, 1995](#)).

***Leptochilichthys agassizii* Garman 1899**

Common name: Agassiz's tongueless smooth-head; **Celindra-de-Agassiz**.

References: mainland Portugal ([Markle and Quéro, 1984; Machida and Shiagoki, 1988\(?\)](#); Carneiro et al., 2014; Uiblein et al., 2015c), Azores (PECS) ([Post, 1987a](#)) and Madeira ([Uiblein et al., 2015c; Porteiro et al., 2017](#)).

Leptochilichthys pinguis (Vaillant 1886)

Common name: Vaillant's tongueless smooth-head; **Celindra-de-Vaillant**.

References: mainland Portugal ([Machida and Shiogaki, 1988](#)).

Leptoderma macrophthalmum Byrkjedal, Poulsen & Galbraith 2011

Common name: No English common name; **Celindra-dos-Açores**.

References: Azores (PECS) ([Byrkjedal et al., 2011](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *L. macrophthalmum*, caught off in the Mid Atlantic Ridge (42°47.0'N, 29°38.0'W), north of the Azores in 2004 and is in the ZMUB collections (ZMUB 19686).

Leptoderma macrops Vaillant 1886

Common name: **Grenadier smooth-head**; **Celindra-olhuda**.

References: mainland Portugal ([Hulley, 2015c](#); [Froese and Pauly, 2019](#)) and Madeira ([Hulley, 2015c](#)). In the Eastern Atlantic this species is distributed from southern Bay of Biscay to Namibia ([Froese and Pauly, 2019](#)). The distribution map reported by Hulley (2015c) indicated the occurrence of this species in parts of mainland Portugal and Selvagens Islands.

Mirognathus normani Parr 1951

Common name: **Norman's smooth-head**; **Celindra-de-Norman**.

References: mainland Portugal ([Markle and Quéro, 1984](#); [Moore et al., 2003](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). It is a very rare species but several earlier literature citations indicate the presence of this species in northern Portugal. However there is no clear demonstration of the occurrence records of this species in this region.

Narcetes erimelas Alcock 1890

Common name: No English common name; **Celindra-de-Goa**.

References: Azores ([MNHN-IC-2009-0463, 1971](#); [Bergstad et al., 2008a](#) (PECS); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#) (PECS); [Almeida and Biscoito, 2019](#)). The geographic coordinates presented (38° 0' 3.6" N ; 26° 21' 3.6" E) in the occurrence record of the Museum national d'Histoire naturelle are not compatible with the location in the Azores because the longitude must be 26° 21' 3.6" W.

Narcetes stomias (Gilbert 1890)

Common name: **Blackhead salmon**; **Celindra-de-cabeça-negra**.

References: Azores ([MNHN-IC-2009-0494, 1971](#); [Bergstad et al., 2008a](#) (PECS); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Almeida and Biscoito, 2019](#)).

Photostylus pycnopterus Beebe 1933

Common name: **Starry smooth-head**; **Celindra-luminosa**.

References: mainland Portugal ([MNHN-IC-1987-0256, 1969](#); [Markle, 1976](#); [Hureau and Monod, 1979](#); [Markle and Quéro, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Krefft, 1976, 1985](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Bergstad et al., 2008a](#) (PECS); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Rinoctes nasutus (Koefoed 1927)

Common name: **Abyssal smooth-head**; **Celindra-abissal**.

References: mainland Portugal ([Christiansen et al., 2014](#)), Azores ([Koefoed, 1927](#) (PECS); [MNHN-IC-2009-0492, 1971](#); [Hureau and Monod, 1979](#) (PECS); [Markle and Quéro, 1984](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bergstad et al., 2008a](#) (PECS); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#); [Almeida and Biscoito, 2019\(?\)](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Bathytroctes nasutus* Koefoed 1927, synonym of *R. nasutu*, is in the ZMUB collections (ZMUB 3208, 1910) caught off 92 NM north-eastern off Marsala Seamount (34°59.0'N, 33°01.0'W). Almeida and Biscoito (2019) mentioned *Rinoctes cf nasutus* (Koefoed, 1927).

Rouleina attrita (Vaillant 1888)

Common name: **Softskin smooth-head**; **Celindra-macia**.

References: mainland Portugal ([Krefft, 1979b](#); [Carneiro et al., 2014](#)), Azores ([MNHN-IC-1885-0166, 1883](#); [Vaillant, 1888, 1919](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Roule, 1916, 1919](#); [Fowler, 1936a](#);

Bertin, 1940; Collins, 1954; Grey, 1956; MNHN-IC-2009-0505, 1971; Markle, 1978; Hureau and Monod, 1979; Krefft, 1979b; Markle and Quéro, 1984; Lloris et al., 1991; Arruda, 1992, 1997; Santos et al., 1997; Melo and Menezes, 2002; Bergstad et al., 2008a (PECS); Porteiro et al., 2010, 2017; Carneiro et al., 2014; Roberts et al., 2017; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Grey, 1956; Markle, 1978; Krefft, 1979b; Lloris et al., 1991). The lectotype of *Bathytroctes attritus* Vaillant 1888, synonym of *R. attrita*, is in the MNHN collections (MNHN-IC-1885-0166) caught off SW the Azores.

***Rouleina maderensis* Maul 1948**

Common name: **Madeiran smooth-head; Celindra-da-Madeira**.

References: Azores (Maul, 1948a; Albuquerque, 1954-1956; Markle and Quéro, 1984; Lloris et al., 1991; Arruda, 1997; Melo and Menezes, 2002; Kukuev, 2002 (PECS); Carneiro et al., 2014; Porteiro et al., 2010, 2017; Froese and Pauly, 2019; Quéro et al., 2019) and Madeira (Maul, 1948a; Albuquerque, 1954-1956; Krefft, 1979b; Markle and Quéro, 1984; Markle and Sazonov, 1990; Lloris et al., 1991; Carneiro et al., 2014; Delgado et al., 2017; Froese and Pauly, 2019; Quéro et al., 2019). The holotype and six paratypes of this species are in the MMF collections (MMF 2698, 1948 (holotype) and MMF 50, MMF 2395, MMF 2396, MMF 2397, MMF 2699, MMF 2745) caught off Madeira.

***Talismania antillarum* (Goode & Bean 1896)**

Common name: **Antillean smooth-head; Celindra-das-Antilhas**.

References: mainland Portugal (?) (Hulley, 2015d(?)), Azores (?) (Hulley, 2015d(?)) and Madeira (?) (Hulley, 2015d(?)). According to Hartel and Orrell (2016c) the current geographic distribution of *T. antillarum* in the Eastern Atlantic is tropical, circumglobal, from about 35°N to 9°S and Froese and Pauly, 2019 indicated from 49°N to 35°S. The distribution map of this species presented by Hulley (2015d) includes the mainland Portugal, Azores and Madeira Islands. The presence of this species among the Portuguese marine ichthyofauna has not been firmly documented and it was not found any evidence on its occurrence in Portugal. Therefore, its occurrence in the mainland Portugal, Azores and Madeira archipelagos is questionable and needs further documentation. Thus, this species is not considered in the total number of valid species.

***Talismania homoptera* (Vaillant 1888)**

Common name: **Hairfin smooth-head; Celindra-da-Mauritânia**.

References: mainland Portugal (?) (Hulley, 2015e(?)) and Madeira (?) (Hulley, 2015e(?)). According to Hartel and Orrell (2016c) the current geographic distribution of *T. homoptera* in the Eastern Atlantic is from 31°N to 4°S. The distribution map of this species presented by Hulley (2015e) includes in part the mainland Portugal and Madeira Islands. The presence of this species in these areas has not been firmly documented and it was not found any evidence of its occurrence in Portugal. The occurrence of this species in the mainland Portugal and Madeira Archipelago is questionable and needs further documentation. Thus, this species is not considered in the total number of valid species.

***Talismania mekistonema* Sulak 1975**

Common name: **Threadfin smooth-head; Celindra-de-barbatana-filamentosa**.

References: Azores (Hureau and Monod, 1979; Tortonese and Hureau, 1979; Markle and Quéro, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Bergstad et al., 2008a (PECS); Porteiro et al., 2010, 2017; Carneiro et al., 2014; Froese and Pauly, 2019).

***Xenodermichthys copei* (Gill 1884)**

Common name: **Bluntsnout smooth-head; Celindra-de-focinho-curto**.

References: mainland Portugal (AVG CODC-Peixes PE/177, 1899; Bragança, 1903; Osório, 1909, 1917; Seabra, 1911; Zugmayer, 1914; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Markle and Quéro, 1984; Moura, 2004; ICES, 2005; Fernandes et al., 2007; Silva, 2008, 2010, 2011; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Silva and Borges, 2014; Nielsen et al., 2015c; Froese and Pauly, 2019), Azores (Collett, 1896a; Richard, 1900, 1910; Joubin, 1905; Roule, 1919a; Fowler, 1936a; Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Markle and Quéro, 1984; Krefft, 1985; Shcherbachov et al., 1985b (PECS); Lloris et al., 1991; Arruda, 1992, 1997; Santos et al., 1997; Pakhorukov, 2008 (PECS); Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Nielsen et al., 2015c; Froese and Pauly, 2019) and Madeira (NHMUK 2004.3.10.56, 1965; Maul, 1976; Markle and Quéro, 1984; Lloris et al., 1991; NHMUK 2016.8.25.751, 1995; Carneiro et al., 2014; Nielsen et al.,

2015c; Froese and Pauly, 2019). The holotype (unique) of *Aleposomus cyaneus* Zugmayer 1914, synonym of *X. copei*, is in the MOM collections (MOM-IC-0091-1324) caught off northern Portugal.

Family Platytroctidae

***Barbantus curvifrons* (Roule & Angel 1931)**

Common name: **Palebelly searsid; Celindra-de-tubo-de-ventre-claro.**

References: mainland Portugal (Quéro et al., 1984; Carneiro et al., 2014; Nielsen et al., 2015d; Orrell and Hartel, 2016; Froese and Pauly, 2019), Azores (PECS) (Carneiro et al., 2014; Nielsen et al., 2015d; Orrell and Hartel, 2016) and Madeira (Maul, 1957; Krefft, 1979b; Quéro et al., 1984, 1990a; Matsui and Rosenblatt, 1987; Lloris et al., 1991; Carneiro et al., 2014; Nielsen et al., 2015d; Orrell and Hartel, 2016; Froese and Pauly, 2019).

***Holtbyrnia anomala* Krefft 1980**

Common name: **Bighead searsid; Celindra-de-tubo-cabeçuda.**

References: Azores (Melo and Menezes, 2002; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Knudsen et al., 2015c; Orrell and Hartel, 2016) and Madeira (Swinney, 1996; Knudsen et al., 2015c).

***Holtbyrnia innesi* (Fowler 1934)**

Common name: **Teardrop tubeshoulder; Celindra-de-tubo-de-Innes.**

References: Azores (PECS) (MAR-ECO - HamPelFish - TI/SF MAR-ECO 4220, 1982) and Madeira (Maul, 1954a; Grey, 1956).

***Holtbyrnia macrops* Maul 1957**

Common name: **Bigeye searsid; Celindra-de-tubo-de-olhuda.**

References: mainland Portugal (Iwamoto, 2015a; Knudsen et al., 2015d; Orrell and Hartel, 2016; Porteiro et al., 2017), Azores (Lloris et al., 1991; Arruda, 1997; Quéro et al., 1984; Sutton et al., 2008; Carneiro et al., 2014; Iwamoto, 2015a; Porteiro et al., 2017; Froese and Pauly, 2019) and Madeira (Maul, 1957; Krefft, 1979b; Quéro et al., 1984, 1990a; Matsui and Rosenblatt, 1987; Lloris et al., 1991; Carneiro et al., 2014; Iwamoto, 2015a; Orrell and Hartel, 2016; Porteiro et al., 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Madeira (stomach content) is in the MMF collections (MMF 6314).

***Holtbyrnia melanocephala* (Vaillant 1888)**

Common name: No English common name; **Celindra-de-tubo-de-cabeça-negra.**

References: mainland Portugal (Nobre, 1935; Grey, 1956; ZSM-Pis-029128 / 534338 / 385926, 1992; ZSM-Pis-029144 / 534354 / 385942, 1992; ZSM-Pis-029246 / 534449 / 386037, 1992). According to Eschmeyer et al. (2019) the current geographic distribution of *H. melanocephala* is in Banc d'Arguin, Eastern Atlantic.

***Maulisia argipalla* Matsui & Rosenblatt 1979**

Common name: Palegold searsid; **Celindra-de-tubo-dourada.**

References: Azores (Santos et al., 1997; Melo and Menezes, 2002; MAR-ECO 5799, 2004; MAR-ECO 13445, 2004 (PECS); Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Nielsen, 2015a) and Madeira (Quéro et al., 1984; Matsui and Rosenblatt, 1987; Lloris et al., 1991; Carneiro et al., 2014; Nielsen, 2015a; Orrell and Hartel, 2016; Froese and Pauly, 2019).

***Maulisia mauli* Parr 1960**

Common name: **Maul's searsid; Celindra-de-tubo-de-Maul.**

References: Azores (MAR-ECO 3495, 2004; Sutton et al., 2008; Carneiro et al., 2014; Orrell and Hartel, 2016; Porteiro et al., 2017) and Madeira (Parr, 1960; Krefft, 1979b; Quéro et al., 1984; Matsui and Rosenblatt, 1987; Lloris et al., 1991; Carneiro et al., 2014; Orrell and Hartel, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Madeira (stomach content) is in the MMF collections (MMF 9594).

***Maulisia microlepis* Sazonov & Golovan 1976**

Common name: **Smallscale searsid; Celindra-de-tubo-de-escama-curta.**

References: mainland Portugal ([Mecklenburg et al., 2018](#)) and Azores ([Melo and Menezes, 2002](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#); [Mecklenburg et al., 2018](#)). The distribution map reported by Mecklenburg et al. (2018) indicated the occurrence of this species in mainland Portugal.

***Mentodus facilis* (Parr 1951)**

Common name: Dark searsid; **Celindra-de-tubo-escura**.

References: mainland Portugal ([Knudsen, 2015a](#)), Azores ([Knudsen, 2015a](#); [Orrell and Hartel, 2016](#)) and Madeira ([NHMUK 1996.8.7.2, 1995](#); [Swinney, 1996](#); [Carneiro et al., 2014](#); [Knudsen, 2015a](#); [Orrell and Hartel, 2016](#)). According to Eschmeyer et al. (2019) *M. facilis* is a cosmopolitan species. The distribution map of this species reported by Knudsen (2015) included parts of mainland Portugal. However, no recent occurrence records were reported for mainland Portugal. Specimens are needed for confirmation.

***Mentodus mesalirus* (Matsui & Rosenblatt 1987)**

Common name: Midridged searsid; **Celindra-de-tubo-de-nervura**.

References: Azores ([MAR-ECO - HamPelFish - TI/SF MAR-ECO 2687, 1979 \(PECS\)](#); [MAR-ECO - HamPelFish - TI/SF MAR-ECO 2686, 1979 \(PECS\)](#); [Carneiro et al., 2014](#); [Knudsen, 2015b](#); [Orrell and Hartel, 2016](#)). Specimens are needed for confirmation in the Azores EEZ.

***Mentodus rostratus* (Günther 1878)**

Common name: Beaked searsid; **Celindra-de-tubo-bicuda**.

References: Azores (?) ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Carneiro et al., 2014](#)) and Madeira (?) ([Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Carneiro et al., 2014](#)). According to Eschmeyer et al. (2019) *M. rostratus* is a cosmopolitan species. However, Matsui and Rosenblatt (1987) refer that the descriptions made by Murray and Hjort (1912) are not sufficient to allocate in this species the specimens described. Thus, this species is not considered in the total number of valid species.

***Normichthys operosus* Parr 1951**

Common name: **Multipore searsid; Celindra-de-tubo-multiporo**.

References: mainland Portugal ([Quéro et al., 1984](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#); [Froese and Pauly, 2019](#)), Azores ([MAR-ECO - HamPelFish - TI/SF MAR-ECO_2122, 1979 \(PECS\)](#); [Matsui and Rosenblatt, 1984](#); [Quéro et al., 1984](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Orrell and Hartel, 2016](#)).

***Perspasia kopua* (Phillipps 1942)**

Common name: Spangled tubeshoulder; **Celindra-de-tubo-estrelada**.

References: mainland Portugal (?) ([YPM ICH 004879, 1959](#); [YPM ICH 004880, 1959](#); [Carneiro et al., 2014](#)). There are two specimens caught off Portugal in the collections of the Peabody Museum of Natural History of Yale University (YPM ICH 004879 and YPM ICH 004880). However, the distribution of this species is circumglobal in the southern hemisphere (Matsui and Rosenblatt, 1987), excluding South America (Froese and Pauly, 2019) or South Africa (Eschmeyer et al., 2019). Thus, this species is not considered in the total number of valid species.

***Platyptroctes apus* Günther 1878**

Common name: **Legless searsid; Celindra-de-tubo-curva**.

References: mainland Portugal ([Zugmayer 1911b](#); [Fowler, 1936a](#); [Krefft, 1979b](#); [Quéro et al., 1984](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#); [Froese and Pauly, 2019](#)), Azores ([ZMH 117473, 1979](#); [SOC Discovery 740633_8_FIS_656001, 1970 \(PECS\)](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#)) and Madeira ([USNM 206893, 1971](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#)).

***Sagamichthys schnakenbecki* (Krefft 1953)**

Common name: **Schnakenbeck's searsid; Celindra-de-tubo-de-Schnakenbeck**.

References: mainland Portugal ([AMNH 36472, 1950\(?\)](#); [Quéro et al., 1984](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#); [Froese and Pauly, 2019](#)), Azores ([MCZ 128329, 1972](#); [Quéro et al., 1984](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Kukuev, 2002 \(PECS\)](#); [Porteiro et al., 1999, 2017](#); [Carneiro et al., 2014](#); [Orrell and Hartel, 2016](#)).

Hartel, 2016; Froese and Pauly, 2019) and Madeira (SOC Discovery 826200_8_FIS_372001, 1973; Carneiro et al., 2014; Orrell and Hartel, 2016).

***Searsia koefoedi* Parr 1937**

Common name: **Koefoed's searsid; Celindra-de-tubo-de-Koefoed.**

References: mainland Portugal (Quéro et al., 1984; MNHN-IC-1988-0502, 1984; MNHN-IC-1987-0253, 1969; Carneiro et al., 2014; Orrell and Hartel, 2016; Froese and Pauly, 2019), Azores (Koefoed, 1927; Parr, 1960; MNHN-IC-1987-0252, 1969; Sazonov, 1976; Matsui and Rosenblatt, 1987; Quéro et al., 1984; Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Orrell and Hartel, 2016; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Maul, 1948a, 1952b; NHMUK 1987.9.4.6, 1958; NHMUK 2002.11.17.4, 1962; MCZ 67160, 1972; MCZ 100979, 1972; Quéro et al., 1984; Matsui and Rosenblatt, 1987; Davies and Merrett, 1998; Carneiro et al., 2014; Orrell and Hartel, 2016; Froese and Pauly, 2019).

Order Siluriformes

Family Ariidae

***Galeichthys feliceps* Valenciennes 1840**

Common name: **White barbel; Bagre-barba-branca.**

References: Azores (?) (Lampe, 1914; Fowler, 1936; Collins, 1954; Arruda, 1997; Santos et al., 1997).

According to Eschmeyer et al. (2019) the distribution of *G. feliceps* includes south-eastern Atlantic and south-western Indian Ocean off southern Africa: Namibia to about East London (South Africa). Santos et al. (1997) mention that the record of this species for the Azores may have been due to a mislabelling of the specimens or jars. The occurrence of this species in the Azores is questionable and specimens are needed for confirmation. Thus, this species is not considered in the total number of valid species.

Order Argentiniformes

Family Argentinidae

***Argentina sphyraena* Linnaeus 1758**

Common name: **Argentine; Argentina-branca**, Argentina, Biqueirão, Biqueirão-branco.

References: mainland Portugal (de Brito Capello, 1869a, 1880; Goode and Bean, 1895, 1896; Seabra, 1911; Osório, 1917; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Muus and Dahlström, 1978; Cohen, 1979a, 1984, 1990; Fernández et al., 1981; INIP, 1981a, 1982a, b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Gomes et al., 2001; Borges et al., 2002; Jorge et al., 2002; ICES, 2005; Fernandes et al., 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Silva, 2008, 2010, 2011; Batista et al., 2009; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Hartel and Orrell, 2016a; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (?) (Nobre, 1924, 1930; Albuquerque, 1954-1956; Osório de Castro, 1967; Arruda, 1997; Santos et al., 1997(?); Carneiro et al., 2014) and Madeira (PECS) (Shcherbachev et al., 1985a; Carneiro et al., 2014). According to Eschmeyer et al. (2019) the distribution of *A. sphyraena* includes the Mediterranean Sea and Sea of Marmara and, in the Eastern Atlantic, from Iceland and Norway to Western Sahara. Santos et al. (1997) mention that the record of this species for the Azores may probably be a misidentification by Nobre (1924). The occurrence of this species in the Azores is questionable and needs further documentation. Thus, this species is not considered in the total number of valid species. Two specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22833, 1967, set of two specimens).

***Glossanodon leioglossus* (Valenciennes 1848)**

Common name: **Smalltoothed argentine; Argentina-de-dentes-pequenos, Argentina-dourada, Biqueirão, Biqueirão-branco.**

References: mainland Portugal (Guichet, 1966; Osório de Castro, 1967; Maul, 1976; Hureau and Monod, 1979; Fischer et al., 1987; Carneiro et al., 2014), Azores (PECS) (Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977 (PECS); Cohen, 1984, 1990; Fock et al., 2002; Kukuev, 2002 (PECS); Carneiro et al., 2014) and Madeira (Guichet, 1966; Osório de Castro, 1967(?); Maul, 1976 (PECS); Ehrich, 1977 (PECS); Hureau and Monod, 1979; Cohen, 1984; Carneiro et al., 2014; Froese and Pauly, 2019). Several

specimens caught off in different areas are in the MMF collections: two specimens at the Josephine Bank (MMF 22857, set of two specimens) and at the Meteor Bank (MMF 22540, 1967, set of twenty five specimens and MMF 22982, 1967, one specimen).

Family Microstomatidae

***Microstoma microstoma* (Risso 1810)**

Common name: **Slender argentine; Argentina-fina.**

References: Azores ([Cohen, 1960; MCZ 66009, 1964; Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014](#)) and Madeira ([Maul, 1948a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Cohen, 1984; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019](#)).

***Nansenia atlantica* Blache & Rossignol 1962**

Common name: No English common name; **Argentina-atlântica.**

References: Azores ([ZMH 118044, 1979 \(PECS\); ZMH 119388, 1979 \(PECS\); MAR-ECO 99999991, 2004 \(PECS\); MAR-ECO 99999995, 2004](#); [Sutton et al., 2008; Carneiro et al., 2014; Uiblein 2015a; Porteiro et al., 2017](#)).

***Nansenia boreacrassicauda* Poulsen 2015**

Common name: No English common name; **Argentina-boreal.**

References: Azores (PECS) ([ZMH 120833, 1982; Carneiro et al., 2014; Poulsen, 2015](#)). Poulsen (2015) re-evaluates several specimens of *Nansenia* and describes a new species - *Nansenia boreacrassicauda*. This re-evaluation leads to the reclassification of specimen ZMH 120833 (former ISH 967-1982), 268.0 mm SL (43°42.0'N, 28°27.0'W), previously identified as *Nansenia iberica* Matallanas 1985 for one of the paratypes of *N. boreacrassicauda*.

***Nansenia groenlandica* (Reinhardt 1840)**

Common name: Greenland argentine; **Argentina-da-Groenlândia.**

References: mainland Portugal ([Cohen, 1984; Carneiro et al., 2014; Froese and Pauly, 2019](#)), Azores ([ZMH 113994, 1975; ZMH 113999, 1975; Carneiro et al., 2014; Poulsen, 2015; Mecklenburg et al., 2018](#)) and Madeira ([NHMUK 1998.8.9.12752, 1958](#)).

***Nansenia iberica* Matallanas 1985**

Common name: No English common name; **Argentina-ibérica.**

References: Azores (PECS) (?) ([ZMH 120833, 1982\(?\)](#); [ZMH 120958, 1982\(?\)](#); [Carneiro et al., 2014](#)). According to Poulsen (2015) the occurrence records ZMH 120833 and ZMH 120958 corresponded to *N. boreacrassicauda* and *N. cf pelagica*, respectively. Thus, this species is not considered in the total number of valid species.

***Nansenia longicauda* Kawaguchi & Butler 1984**

Common name: No English common name; **Argentina-de-cauda-longa.**

References: Madeira ([ZMH 25146, 1966; ZMH 138021, 1982; Carneiro et al., 2014; Poulsen, 2015](#)). The distribution map reported by Poulsen (2015) shows the occurrence of this species off Madeira without indicating the geographic coordinates, which is probably within the Madeira EEZ.

***Nansenia oblita* (Facciolà 1887)**

Common name: No English common name; **Argentina-do-Mediterrâneo.**

References: mainland Portugal ([Eschmeyer et al., 2019](#)) and Madeira ([Poulsen, 2015](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *N. oblita* is the Greenland and North Atlantic, Mediterranean Sea and, in the Eastern Atlantic, from southern Ireland to Western Sahara. The distribution map reported by Poulsen (2015) shows the occurrence of this species off Madeira without indicating the geographic coordinates, which is probably within the Madeira EEZ.

Nansenia cf. pelagica

Common name: No English common name; **Argentina-pelágica.**

References: Azores (PECS) ([ZMH 120958, 1982; Poulsen, 2015](#)). According to Poulsen (2015) the occurrence record ZMH 120958 (former ISH 465-1982) corresponded to *N. cf. pelagica*.

Nansenia tenera Kawaguchi & Butler 1984

Common name: No English common name; **Argentina-dourada**.

References: Azores (?) ([MAR-ECO 4225, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017 \(PECS\)](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *N. tenera* is the south-eastern Atlantic and Porteiro *et al.* (2017) indicate scattered records in North Atlantic at about 55°N and northern Middle-Atlantic Ridge (MAR) in the Northeast Atlantic and around Cape Verde Islands. Gon (pers. comm., 2014, *in* Uiblein, 2015b) mentions the need for a revision of the taxonomy of the genus *Nansenia*, as there may be one or two hidden synonymies among the currently recognized species. Thus, the occurrence records of this species for the Azores should be re-evaluate and it is not considered in the total number of validated species.

Family Bathylagidae

Bathylagichthys greyae (Cohen 1958)

Common name: Grey's deepsea smelt; **Argentina-da-fundura-de-Grey**.

References: mainland Portugal ([Cohen, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 1999, 2010, 2017](#); [Carneiro et al., 2014](#)) and Madeira ([ZMH 6825, 1970](#); [ZMH 6826, 1970](#); [Hureau and Monod, 1979](#); [Cohen, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Bathylagus euryops Goode & Bean 1896

Common name: **Goiter blacksmelt; Argentina-da-fundura-negra**.

References: mainland Portugal ([IFREMER Biocean 34599, 1969](#); [Cohen, 1984](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Porteiro et al., 2017](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)), Azores ([ZMH 114097, 1975](#); [MAR-ECO 7385, 2004](#); [MAR-ECO 5566, 2004](#); [MAR-ECO 7674, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#); [Mecklenburg et al., 2018](#)) and Madeira ([MCZ 66229, 1969](#); [MCZ 66230, 1969](#); [MCZ 60353, 1969](#); [Cohen, 1984\(?\)](#); [Carneiro et al., 2014](#)).

Dolicholagus longirostris (Maul 1948)

Common name: Longsnout blacksmelt; **Argentina-da-fundura-de-focinho-comprido**.

References: mainland Portugal ([MCZ 159382, 1969](#); [Cohen, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([MCZ 159367, 1969](#); [MCZ 159379, 1969](#); [Porteiro et al., 1999, 2017](#); [Arkipov and Mamedov, 2008 \(PECS\)](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#)) and Madeira ([Maul, 1948a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [SOC Discovery 826300_8_FIS_770002, 1973](#); [Cohen, 1979b](#), [1984b](#), [1990b](#); [Lloris et al., 1991](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Bathylagus longirostris* Maul 1948, synonym of *D. longirostris*, caught off in Madeira is in the MMF collections.

Melanolagus bericoides (Borodin 1929)

Common name: Bigscale deepsea smelt; **Argentina-da-fundura-escamuda**.

References: mainland Portugal ([YPM ICH 004503, 1959](#); [Cohen, 1984](#); [Carneiro et al., 2014](#)), Azores ([IMAR/DOP COLETA 225, 2001](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017 \(PECS\)](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([SOC Discovery 826200_8_FIS_070001, 1973](#); [MCZ 61922, 1969](#); [Cohen, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Family Opisthoproctidae

Bathylychnops brachyrhynchus (Parr 1937)

Common name: No English common name; **Peixe-fantasma-dardo-listado**.

References: Azores (PECS) ([NHMUK 2016.6.15.5, 1961](#); [Carneiro et al., 2014](#)) and Madeira ([Hartel and Orrell, 2016b](#)). According to Eschmeyer *et al.* (2019) this species is apparently found in all oceans. Hartel and Orrell (2016b) indicate the occurrence of this rare species off Madeira without indicating the geographic coordinates. The occurrence of this species in Madeira is questionable and needs further documentation.

Bathylychnops exilis Cohen 1958

Common name: **Javelin spookfish; Peixe-fantasma-dardo**.

References: Azores (Cohen, 1984; MAR-ECO 5677, 2004; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019).

Dolichopteryx longipes (Vaillant 1888)

Common name: Brown-snout spookfish; **Peixe-fantasma-de-barbatanas-longas**.

References: Azores (IMAR/DOP COLETA 1693, 2001; Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Carneiro *et al.*, 2014) and Madeira (MCZ 66321, 1969; MCZ 66331, 1972; Cohen, 1984; Quéro, 1990a; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Monacoa grimaldii (Zugmayer 1911)

Common name: **Mirrorbelly**; **Peixe-sola-de-Grimaldi**, Peixe-sola.

References: mainland Portugal (MOM-IC-0091-1163, 1910; Zugmayer, 1911a, b; Fowler, 1936a; Albuquerque, 1954-1956; Cohen, 1984; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Murray and Hjort, 1912; Roule and Angel, 1933; Fowler, 1936a, b; Albuquerque, 1954-1956; Cohen, 1964a, 1984c, 1990; Geistdoerfer *et al.*, 1971; Hamer *et al.*, 1994; Arruda, 1997; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 1999, 2010, 2017; Carneiro *et al.*, 2014; Poulsen *et al.*, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Roule and Angel, 1933; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Cohen, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The syntype of *Opisthoproctus grimaldii* Zugmayer 1911, synonym of *M. grimaldii*, caught off in Portugal is in the MOM collections (MOM-IC-0091-1163).

Opisthoproctus soleatus Vaillant 1888

Common name: **Barrel-eye**; **Peixe-sola**.

References: mainland Portugal (Cohen, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Albert I^{er} de Mónaco, 1904; Murray and Hjort, 1912; Roule, 1916, 1919a; Schmidt, 1918; Roule and Angel, 1933; Fowler, 1936a, b; Albuquerque, 1954-1956; Cohen, 1964, 1984c; Arruda, 1997; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Maul, 1949a; Albuquerque, 1954-1956; Cohen, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Rhynchohyalus natalensis (Gilchrist & von Bonde 1924)

Common name: Glasshead barreleye; **Peixe-sola-cabeça-de-vidro**.

References: Azores (MCZ 66429, 1969; Post 1987a; Porteiro *et al.*, 1999; Carneiro *et al.*, 2014) and Madeira (Maul, 1946b; MCZ 96900, 1978; Cohen, 1979c, 1984c; Quéro, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Ophthalmostelon macropus* Maul 1946, synonym of *R. natalensis*, caught off in Madeira (stomach content) is in the MMF collections.

Order Salmoniformes

Family Salmonidae

Oncorhynchus mykiss (Walbaum 1792)

Common name: **Rainbow trout**; **Truta-arco-íris**, Arco-íris, Francesa, Truíta, Truta, Truta americana, Truta-francesa, Truta-pequena.

References: mainland Portugal[▲] (Nobre, 1931, 1932, 1935; Welcomme, 1988; Carneiro *et al.*, 2014; Froese and Pauly, 2019) Azores (AquaNIS, 1990; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (AquaNIS, 2002; Ribeiro *et al.*, 2005a; Carneiro *et al.*, 2014; Froese and Pauly, 2019). *O. mykiss* is an alien and invasive species introduced in Portugal. This species is primarily a freshwater fish, although sea-run populations, often known as steelhead, exist in some areas. However, populations of *O. mykiss* are capable of migrating to and surviving in the sea (Jonsson 2011). In Portugal all the occurrence records “take place” in the watersheds (mainland Portugal, Azores and Madeira Islands), but occasional and scattered catches are reported in estuaries and coastal areas (mainland Portugal). The introduction of this species in the Azores occurred in 1990 and in Madeira in 2002.

Salmo salar Linnaeus 1758

Common name: **Atlantic salmon**; **Salmão-do-Atlântico**, Nelma, Salmão, Salmão-europeu, Salmonete, Santantoninho, Santo-antoninho, Tranca.

References: mainland Portugal▲ ([Carvalho, 1756](#); [Vandelli, 1771, 1797](#); [de Brito Capello, 1869a, 1880](#); [Gervais and Boulart, 1876](#); [Baldaque da Silva, 1891](#); [Vieira, 1894a, 1897, 1901](#); [Seabra, 1911](#); [Nobre, 1931, 1932, 1935](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979d, 1984b](#); [Kottelat and Freyhof, 2007](#); [Carneiro et al., 2014](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)).

***Salmo trutta* Linnaeus 1758**

Common name: **Sea trout; Truta-marisca**, Cambo, Marinha, Margarida, Marisca, Palmeira, Pinta, Pintada, Relha, Relho, Salmonada, Salmoneira, Salmoneja, Salmonete, Sapeira, Truíta, Truta, Truta-assalmonada, Truta-de-água-doce, Truta-do-mar, Truta-do-rio, Truta-dos-rios, Truta-maior, Truta-marinha, Truta-prateada, Truta-salmonada, Truta-salmoneira, Truta-salmoneja, Truta-sapeira, Truta-vulgar, Vermelhinha.

References: mainland Portugal▲ ([Carvalho, 1765](#); [Vandelli, 1771](#); [de Brito Capello, 1869a, 1880](#); [Baldaque da Silva, 1891](#); [Osório, 1894](#); [Vieira, 1894a, 1897, 1901](#); [Nobre, 1895c, 1903, 1931, 1932, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Osório de Castro, 1954, 1967](#); [Blanc et al., 1971](#); [Svetovidov, 1979d, 1984b](#); [Fischer et al., 1987](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#)), Azores (PECS) ([MNHN-IC-1982-0480, 1979](#)) and Madeira (Ribeiro et al., 2005a; Froese and Pauly, 2019).

Order Stomiiformes

Family Gonostomatidae

***Bonapartia pedaliota* Goode & Bean 1896**

Common name: Longray fangjaw; **Boqueirão-de-raio-longo**.

References: mainland Portugal ([ZMH 103332, 1966](#); [MCZ 79875, 1969](#); [Badcock, 1984](#); [Carneiro et al., 2014](#)), Azores ([Roule and Angel, 1933](#); [Jespersen, 1933](#); [Fowler, 1936b](#); [Ferreira, 1942](#); [Belloc, 1949](#); [Albuquerque, 1954-1956](#); [Grey, 1964](#); [Mukhacheva, 1976](#); [Post, 1985](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#)) and Madeira ([MCZ 144673, 1969 \(PECS\)](#); [SOC Discovery 826200_8_FIS_075001, 1973](#); [Badcock, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Zaphotias photocephalus* Roule & Angel 1933, synonym of *B. pedaliota*, caught off in the Azores is in the MOM collections (MOM-IC-0091-1280).

***Cyclothona acclinidens* Garman 1899**

Common name: **Benttooth bristlemouth; Boqueirão-de-dentes-curvos**.

References: mainland Portugal ([YPM ICH 004535, 1959](#); [YPM ICH 004537, 1959](#); [YPM ICH 004539, 1959](#); [MCZ 144290, 1986](#); [Badcock, 1984](#); [Miya and Nishida, 1996](#); [Carneiro et al., 2014](#); [Kobyliansky et al., 2015a](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Witzell, 1973](#); [Quéro, 1975](#); [Witzell, 1979](#); [Badcock, 1984](#); [Miya and Nishida, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#)) and Madeira ([ZMH 105464, 1968](#); [SOC Discovery 826200_8_FIS_135004, 1973](#); [Badcock, 1984](#); [Miya and Nishida, 1996](#); [Carneiro et al., 2014](#); [Kobyliansky et al., 2015a](#); [Froese and Pauly, 2019](#)).

***Cyclothona alba* Brauer 1906**

Common name: Pale bristlemouth; **Boqueirão-branco**.

References: mainland Portugal ([ZMH 117223, 1979](#); [Badcock, 1984](#); [Miya and Nishida, 1996](#)), Azores ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Koefoed, 1960](#); [MCZ 144143, 1964](#); [Badcock, 1984](#); [Miya and Nishida, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Bekker et al., 1982](#); [Miya and Nishida, 1996](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Cyclothona braueri* Jespersen & Tåning 1926**

Common name: **Gerrick; Boqueirão-de-Brauer**.

References: mainland Portugal ([Jespersen and Taaning, 1926](#); [Fowler, 1936b](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Badcock, 1984](#); [Miya and Nishida, 1996](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Zugmayer, 1911b](#); [Roule, 1919a](#); [Roule and Angel, 1930](#); [Fowler, 1936a](#); [Ferreira, 1942](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Koefoed, 1960](#); [Geistdoerfer et al., 1971](#); [Saldanha,](#)

1977a; Bekker *et al.*, 1982; Badcock, 1984; Miya and Nishida, 1996; Arruda, 1997; Santos *et al.*, 1997; Arkhipov and Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Roule and Angel, 1930; Zugmayer, 1933; Fowler, 1936b; Albuquerque, 1954-1956; Maul, 1971a; Bekker *et al.*, 1982; Badcock, 1984; Miya and Nishida, 1996; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Cyclothona livida* Brauer 1902**

Common name: Bristlemouth; **Boqueirão-lívido**.

References: mainland Portugal (YPM ICH 004603, 1959; MCZ 143319, 1969; Badcock, 1984; MCZ 143318, 1986; Gordo *et al.*, 1990; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (MCZ 143321, 1969; Witzell, 1979; Porteiro *et al.*, 1999, 2010; Carneiro *et al.*, 2014) and Madeira (ZMH 105422, 1968; MCZ 143320, 1969; MCZ 143314, 1978; Badcock, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Cyclothona microdon* (Günther 1878)**

Common name: Veiled anglemouth; **Boqueirão-de-dentes-pequenos**.

References: mainland Portugal (MNHN-IC-1883-0063, 1882; Vaillant, 1888; Seabra, 1911; MNHN-IC-1912-0017, 1908; Albuquerque, 1954-1956; Badcock, 1984; YPM ICH 004619, 1959; ZMH 123503, 1981 (PECS); ZSM-Pis-029160 / 534370 / 385958, 1992; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores (NHMUK 1890.6.16.53, 1878-1892; MNHN-IC-1884-1083, 1883; MNHN-IC-1884-1084, 1883; Vaillant, 1888, 1919; Collett, 1896a; Richard, 1905a; Zugmayer, 1911b, 1933; Murray and Hjort, 1912; Roule and Angel, 1924, 1930, 1933; Fowler, 1936a; Bertin, 1940; Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Koefoed, 1960; Geistdoerfer *et al.*, 1971; Badcock, 1984; Arruda, 1992, 1997; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira (Zugmayer, 1911; ZMH 105427, 1968; MCZ 143077, 1969; Maul, 1971a; Badcock, 1984; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019). The holotype, paratype and two syntypes of *Neostoma quadrioculatum* Vaillant 1888, synonym of *C. microdon*, are in the MNHN and NHMUK collections [MNHN-IC-1884-1083 (holotype), MNHN-IC-1884-1084 and NHMUK 1890.6.16.53 (syntypes)] both caught off in the Azores and the MNHN-IC-1883-0063 (paratype) caught off in mainland Portugal.

***Cyclothona obscura* Brauer 1902**

Common name: Hidden bristlemouth; **Boqueirão-sombrio**.

References: Madeira (Badcock, 1982(?), 1984; Carneiro *et al.*, 2014; Ralph, 2015c; Froese and Pauly, 2019).

***Cyclothona pallida* Brauer 1902**

Common name: Tan bristlemouth; **Boqueirão-bicolor**.

References: mainland Portugal (YPM ICH 004684, 1959; Badcock, 1982, 1984; ZSM-Pis-029138 / 534348 / 385936, 1991; Miya and Nishida, 1996; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Murray and Hjort, 1912; Fowler, 1936a; Koefoed, 1960; Geistdoerfer *et al.*, 1971; Badcock, 1984; Miya and Nishida, 1996; Arruda, 1997; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (MCZ 142720, 1969; MCZ 142683, 1978; Maul, 1971a; Badcock, 1982, 1984; Bekker *et al.*, 1982; ZSM-Pis-029351 / 534549 / 386137, 1990; Miya and Nishida, 1996; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Cyclothona parapallida* Badcock 1982**

Common name: Shadow bristlemouth; **Boqueirão-sombreado**.

References: mainland Portugal (Miya and Nishida, 1996) and Madeira (Badcock, 1982; Miya, 1994; Miya and Nishida, 1996; Carneiro *et al.*, 2014).

***Cyclothona pseudopallica* Mukhacheva 1964**

Common name: Slender bristlemouth; **Boqueirão-delgado**.

References: mainland Portugal (Badcock, 1984; MCZ 142514, 1986; ZSM-Pis-028636 / 533859 / 385447, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Witzell, 1979; Badcock, 1984; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (MCZ 142500, 1969; MCZ 142225, 1978; Bekker *et al.*, 1982; Badcock, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Cyclothona signata Garman 1899

Common name: Showy bristlemouth; **Boqueirão-vistoso**.

References: mainland Portugal (?) ([VLIZ 318569](#), 1922; [VLIZ 318570](#), 1922; [Damas, 1934](#); [Carneiro et al., 2014](#)), Azores (?) ([Zugmayer, 1911b](#); [Murray and Hjort, 1912](#); [Roule, 1919a](#); [VLIZ 318563](#), 1922; [VLIZ 318564](#), 1922; [Damas, 1934](#); [Fowler, 1936a](#); [Carneiro et al., 2014](#)) and Madeira (?) ([Damas, 1934](#)). According to Miya and Nishida (1996) and Eschmeyer *et al.* (2019) the current geographic distribution of *C. signata* is the central and Eastern Pacific. Several authors mention the occurrence of the species *Cyclothona signata* Garman 1899 and the subspecies *Cyclothona signata alba* Brauer 1906 in Portuguese waters. The occurrence records of *C. signata* var. *alba* corresponded to *Cyclothona alba* Brauer 1906 and the occurrence records of *C. signata* are likely to belong to *C. alba* subspecies. Thus, this species is not considered in the total number of valid species.

Diplophos taenia Günther 1873

Common name: Pacific portholefish; **Boqueirão-comprido**.

References: mainland Portugal ([Witzell, 1979](#); [Badcock, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Santos et al., 1997](#); [Carneiro et al., 2014 \(PECS\)](#)) and Madeira ([MCZ 88494](#), 1970; [MCZ 97043](#), 1986; [Quéro et al., 1990b](#); [Lloris et al., 1991](#); [NHMUK 2016.8.25.755-756](#), 1995; [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Gonostoma atlanticum Norman 1930

Common name: **Atlantic fangjaw**; **Boqueirão-do-Atlântico**.

References: mainland Portugal ([Fernández et al., 1981](#); [Gordo et al., 1990](#); [ZSM-Pis-028850 / 534069](#) / [385657](#), 1991), Azores ([Witzell, 1979](#); [Arkhipov and Mamedov, 2008 \(PECS\)](#)) and Madeira ([Badcock, 1984](#); [ROM 68126](#), 1984 (PECS); [NHMUK 2004.2.3.77](#), 1990 (PECS); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Gonostoma denudatum Rafinesque 1810

Common name: Bristlemouth; **Boqueirão-do-Mediterrâneo**.

References: mainland Portugal ([Witzell, 1979](#); [Badcock, 1984](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Silva, 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912](#); [Roule and Angel, 1930](#); [Fowler, 1936a](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Koefoed, 1958](#); [Dias et al., 1976](#); [Badcock, 1984](#); [Post, 1985](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Kukuev, 2002 \(PECS\)](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Günther, 1864a](#); [Johnson, 1862c](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Norman, 1930](#); [Fowler, 1936a](#); [Badcock, 1984](#); [Quéro et al., 1990b](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Manducus maderensis (Johnson 1890)

Common name: No English common name; **Boqueirão-da-Madeira**.

References: Madeira ([NHMUK 1890.5.31.6.](#), 1878-1892; [Johnson, 1890](#); [Norman, 1930](#); [Fowler, 1936a, b](#); [Maul, 1948a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Witzell, 1979](#); [Badcock, 1984](#); [Quéro et al., 1990b](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Gonostoma maderense* Johnson 1890, synonym of *M. maderensis*, collected from the fish market, Funchal (Madeira) is in the NHMUK collections ([NHMUK 1890.5.31.6](#)).

Margrethia obtusirostra Jespersen & Tåning 1919

Common name: Bighead portholefish; **Boqueirão-cabeçudo**.

References: mainland Portugal ([NHMUK 2004.3.8.490](#), 1966; [MCZ 89211](#), 1969; [Badcock, 1984](#); [ZSM-Pis-029337 / 534536](#) / [386124](#), 1990; [ZSM-Pis-029395 / 534589](#) / [386177](#), 1990; [Moura, 2004](#); [Carneiro et al., 2014](#); [Kobylansky et al., 2015b](#); [Froese and Pauly, 2019](#)), Azores ([Koefoed, 1958](#); [Grey, 1964](#); [Mukhacheva, 1976](#); [Badcock, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 1999, 2010, 2017](#); [Carneiro et al., 2014](#); [Kobylansky et al., 2015b](#); [Froese and Pauly, 2019](#)) and Madeira ([Maurin et al., 1977](#); [Witzell, 1979](#); [Badcock, 1984](#); [Quéro et al., 1990b](#); [Carneiro et al., 2014](#); [Kobylansky et al., 2015b](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)).

Sigmops bathyphilus (Vaillant 1884)

Common name: **Spark anglemouth**; **Boqueirão-brilhante**.

References: mainland Portugal ([MNHN-IC-1912-0013](#), 1910; [NHMUK 1987.9.4.43](#), 1956; [YPM ICH 004730](#), 1959; [Witzell, 1979](#); [Badcock, 1984](#); [ZSM-Pis-029133 / 534343 / 385931](#), 1992; [ICES, 2005](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1885-0059](#), 1883; [Vaillant, 1888](#), 1919; [Goode and Bean, 1895](#), 1896; [Collett, 1896b](#); [Brauer, 1906](#); [Murray and Hjort, 1912](#); [Roule, 1919a](#); [Roule and Angel, 1923](#); [Barnard, 1925](#); [Fowler, 1936a](#); [Bertin, 1940](#); [Ferreira, 1942](#); [Belloc, 1949](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Koefoed, 1958](#); [Grey, 1964](#); [Badcock, 1984](#); [Post, 1985](#); [Arruda, 1992](#), 1997; [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 2010](#), 2017; [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Zugmayer, 1911b](#); [Fowler, 1936a, b](#); [Albuquerque, 1954-1956](#); [MCZ 141716](#), 1969; [Maul, 1971a](#); [Badcock, 1984](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Cyclothona grandis* Collett 1896 is in the MOM collections (not found in 1995) caught off west of São Miguel (Azores) is synonym of *S. bathyphilus*.

***Sigmops elongatus* (Günther 1878)**

Common name: **Elongated bristlemouth fish; Boqueirão-dentudo.**

References: mainland Portugal ([MOM-IC-0091-1180](#), 1910; [Zugmayer, 1911a](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Badcock, 1984](#); [Moura, 2004](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912](#) (PECS); [Roule and Angel, 1933](#); [Zugmayer, 1933](#); [Fowler, 1936a](#); [Ferreira, 1942](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Koefoed, 1958](#); [Grey, 1964](#); [Geistdoerfer et al., 1971](#); [Badcock, 1984](#); [Post, 1985](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 2010](#), 2017; [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Maul, 1971a](#); [Badcock, 1984](#); [Quéro et al., 1990b](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The syntypes of *Gonostoma polyphos* Zugmayer 1911, synonym of *S. elongatus*, are in the MOM collections ([MOM-IC-0091-1180](#) and [MOM-IC-0091-1195](#)) both caught off mainland Portugal.

Family Sternopychidae

***Argyripnus atlanticus* Maul 1952**

Common name: No English common name; **Pai-velho-do-Atlântico.**

References: Azores (PECS) ([Ehrich, 1977](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#)) and Madeira ([Maul, 1952b](#); [Ehrich, 1977](#); [Witzell, 1979](#); [Badcock, 1984](#); [Quéro et al., 1990c](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off at the Funchal Harbor (Madeira) is in the MMF collections (MMF 3226).

***Argyropelecus aculeatus* Valenciennes 1850**

Common name: Atlantic silver hatchetfish; **Pai-velho-prateado-do-Atlântico**, Pai-velho.

References: mainland Portugal ([ICM-CSIC 7794](#), n.d.; [NHMUK 2002.3.1.94-96](#), 1967; [MNHN-IC-1987-0305](#), 1969; [INIP, 1981a](#), 1982a(?); [Martins, 1981](#); [Badcock, 1984](#); [Saldanha et al., 1995](#); [Borges et al., 2002](#); [Moura, 2004](#); [ICES, 2005](#); [Pakhorukov, 2008](#); [Silva, 2008](#), 2011; [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-0000-1817](#), 1829; [Cuvier and Valenciennes, 1849](#); [Günther, 1864a](#); [Goode and Bean, 1895](#), 1896; [Collett, 1897](#), 1903, 1905; [Brauer, 1906](#); [Murray and Hjort, 1912](#); [Jespersen, 1915](#), 1934; [Borodin, 1931](#); [Zugmayer, 1933](#); [Fowler, 1936a](#); [Bertin, 1940](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Koefoed, 1962](#); [Schultz, 1964](#); [Osório de Castro, 1967](#); [Dias et al., 1976](#); [Pafort-Van Iersel, 1981](#); [Badcock, 1984](#); [Lloris et al., 1991](#); [Arruda, 1992](#), 1997; [Santos et al., 1997](#); [Fock et al., 2002](#) (PECS); [Pakhorukov, 2008](#) (PECS); [Porteiro et al., 1999](#), 2010, 2017; [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1850](#), 1852; [Noronha and Sarmento, 1948](#); [Maul, 1949a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Badcock, 1984](#); [Quéro et al., 1990c](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in the Azores is in the MNHN collections ([MNHN-IC-0000-1817](#)).

***Argyropelecus affinis* Garman 1899**

Common name: **Pacific hatchet fish; Pai-velho-do-Pacífico**, Pai-velho.

References: mainland Portugal ([GNM 1946-6123](#), 1946 (PECS); [GNM 1946-6124](#), 1946), Azores ([Roule and Angel, 1933](#) (PECS); [USNM 219955](#), 1979 (PECS); [Osório de Castro, 1967](#); [Baird, 1979](#) (PECS); [Badcock, 1984](#) (PECS); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira

(SMF 2070, 1898; Baird, 1979; SOC Discovery 826200 8 FIS 011003, 1973; Osório de Castro, 1967; Badcock, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). Folwer (1936a), Albuquerque (1954-1956) and Badcock (1984) indicate the occasional occurrence of *A. affinis* between the Azores and Madeira.

***Argyropelecus bocagei* Osório 1909**

Common name: No English common name; **Pai-velho-de-Bocage**.

References: mainland Portugal (?) (Osório, 1909; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956). Osório (1909) identified this species from one specimen caught off Setúbal (Portugal), at the depth of 600 meters. This unique holotype was preserved in the Museu Bocage and destroyed in the 1978 fire. This species is not considered in the total number of valid species, because its not possible to re-evaluate the taxonomic validity of this record.

***Argyropelecus gigas* Norman 1930**

Common name: **Hatchetfish; Pai-velho-gigante**, Pai-velho.

References: mainland Portugal (NHMUK 2003.4.25.219-265, 1966; ZSM-Pis-028797 / 534017 / 385605, 1991; Badcock, 1984; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019), Azores (NHMUK 2015.2.12.12-14, 1966; Badcock, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Maul, 1949a, 1952b; Albuquerque, 1954-1956; Osório de Castro, 1967; Badcock, 1984; Quéro *et al.*, 1990c; Carneiro *et al.*, 2014; Delgado *et al.*, 2017; Froese and Pauly, 2019).

***Argyropelecus hemigymnus* Cocco 1829**

Common name: **Half-naked hatchetfish; Pai-velho-manchado**, Pai-velho.

References: mainland Portugal (MNHN-IC-1883-0053, 1882; Vaillant, 1888; Goode and Bean, 1895, 1896; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Badcock, 1984; Saldanha *et al.*, 1995; Moura, 2004; ICES, 2005; Chaves, 2008; Silva, 2008, 2011; Carneiro *et al.*, 2014; Silva and Borges, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores (Brauer, 1906; Murray and Hjort, 1912; Jespersen, 1915; Roule, 1919a; Vaillant, 1919; Roule and Angel, 1924; Borodin, 1931; Zugmayer, 1933; Fowler, 1936a; Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Koefoed, 1962; Schultz, 1964; Osório de Castro, 1967; Baird, 1971; Pafort-Van Iersel, 1981; Bekker *et al.*, 1982; Badcock, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira (Lowe, 1850, 1852; Zugmayer, 1911b; Vaillant, 1919; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1971a; Bekker *et al.*, 1982; Badcock, 1984; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019).

***Argyropelecus olfersii* (Cuvier 1829)**

Common name: Silver hatchetfish; **Pai-velho-prateado**, Pai-velho.

References: mainland Portugal (MNHN-IC-1883-0052, 1882; Vaillant, 1888; Goode and Bean, 1895, 1896; Brauer, 1906; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Muus and Dahlström, 1978; INIP, 1982a; Badcock, 1984; IPMA 9121, 1998; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores (Lütken, 1892; Collett, 1896a; Regan, 1903a; Brauer, 1906; Murray and Hjort, 1912; Jespersen, 1915; Roule, 1919a; Vaillant, 1919; Nobre, 1924, 1930, 1935; Zugmayer, 1933; Roule and Angel, 1933; Nobre, 1935; Fowler, 1936a; Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Koefoed, 1962; Schultz, 1964; Osório de Castro, 1967; Baird, 1971; Pafort-Van Iersel, 1981; Badcock, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira (Lowe, 1850, 1852; Vaillant, 1888, 1919; Brauer, 1906; Norman, 1930; Nobre, 1935; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Quéro *et al.*, 1990c; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019).

***Argyropelecus sladeni* Regan 1908**

Common name: **Sladen's hatchet fish; Pai-velho-de-Sladen**.

References: Azores (MCZ 34993, 1931; Carneiro *et al.*, 2014) and Madeira (Bekker *et al.*, 1982; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Maurolicus amethystinopunctatus Cocco 1838

Common name: Pearlsides; **Peixe-luz**.

References: mainland Portugal ([MCZ 64712, 1969](#); [MCZ 92518, 1972](#); Pakhorukov, 2008; Carneiro et al., 2014), Azores ([Vaillant, 1919](#); [Collins, 1954](#); Parin and Kobyliansky, 1996; Santos et al., 1997; Pakhorukov, 2008 (PECS); Porteiro et al., 2010; Carneiro et al., 2014; Rees et al., 2016; Froese and Pauly, 2019) and Madeira ([MCZ 81214, 1969](#); [MCZ 92514, 1972](#); Carneiro et al., 2014). Rees et al. (2016) based on molecular data studies suggest that the type species for the genus, *M. muelleri* (Gmelin 1798), and the species *M. amethystinopunctatus* Cocco 1838 are conspecifics. However, Eschmeyer et al. (2019) considered these two species valid.

Maurolicus muelleri (Gmelin 1789)

Common name: **Silvery lightfish; Peixe-luz-de-Muller**.

References: mainland Portugal ([Muus and Dahlström, 1978](#); [INIP, 1981a, b, 1982a](#); Martins, 1981; Badcock, 1984; [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Gordo et al., 1990](#); ICES, 2005; Chaves, 2008; Henriques et al., 2008; [Garrido et al., 2009](#); [Silva et al., 2009](#); Carneiro et al., 2014; OCEANA, 2014; [Silva and Borges, 2014](#); [Bañón et al., 2016a](#); Froese and Pauly, 2019), Azores ([Fowler, 1936a](#); Albuquerque, 1954-1956; Koefoed, 1958; Grey, 1964; [Dias et al., 1976](#); Mukhacheva, 1981; Bekker et al., 1982; Badcock, 1984; [INIP, 1985b](#); [Lloris et al., 1991](#); Parin and Kobyliansky, 1996; Monteiro, 1996; Monteiro et al., 1996b; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Kukuev, 2002 (PECS); Diekmann et al., 2006 (PECS); Arkhipov and Mamedov, 2008 (PECS); Sutton et al., 2008; Carneiro et al., 2014; Rees et al., 2016; Froese and Pauly, 2019) and Madeira ([Quéro et al., 1990c](#); Carneiro et al., 2014; Froese and Pauly, 2019).

Polyipnus polli Schultz 1961

Common name: No English common name; **Pai-velho-redondo**.

References: Azores (PECS) ([MCZ 149500, 1978](#); Porteiro et al., 1999, 2017; Carneiro et al., 2014) and Madeira ([MCZ 135280, 1978](#); Quéro et al., 1990c; Carneiro et al., 2014; Froese and Pauly, 2019).

Sternopyx diaphana Hermann 1781

Common name: **Diaphanous hatchet fish; Pai-velho-diáfano**, Pai-velho.

References: mainland Portugal ([Brauer, 1906](#); [MNHN-IC-1912-0021, 1910](#); [Nobre, 1935](#); Albuquerque, 1954-1956; Badcock, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([Cuvier and Valenciennes, 1849](#); [MNHN-IC-1884-1072, 1883](#); [Vaillant, 1888, 1919](#); [Collett, 1896a](#); [Regan, 1903a](#); Brauer, 1906; Richard, 1910; [Zugmayer, 1911b](#), 1933; Murray and Hjort, 1912; Roule, 1911, 1919a; Jespersen, 1915; Roule and Angel, 1924, 1930; [Fowler, 1936a, b](#); [Belloc, 1938](#); Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Koefoed, 1962; Haedrich, 1964; [Schultz, 1964](#); Geistdoerfer et al., 1971; Pafort-Van Iersel, 1981; Badcock, 1984; Arruda, 1992, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([Goode and Bean, 1895\(?\)](#), 1896(?); [Vaillant, 1919](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936b](#); [Maul, 1949a, 1971a](#); Nunes, 1953, 1974; Albuquerque, 1954-1956; Badcock, 1984; Carneiro et al., 2014; Froese and Pauly, 2019).

Sternopyx pseudoboscura Baird 1971

Common name: **Highlight hatchetfish; Pai-velho-oblíquo**, Pai-velho.

References: mainland Portugal ([NHMUK 2007.2.26.40, 1966](#); Badcock, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([AMS I.12669, 1912](#); Pafort-Van Iersel, 1981; Badcock, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([ZMH 110847, 1971](#); Badcock, 1984; Carneiro et al., 2014; Froese and Pauly, 2019).

Valencienellus tripunctulatus (Esmark 1871)

Common name: Constellationfish; **Pai-velho-de-três-pontos**.

References: mainland Portugal ([YPM ICH 004835, 1959](#); Badcock, 1984; Carneiro et al., 2014; Kobyliansky and Herrera, 2015a; Bañón et al., 2016a; Froese and Pauly, 2019), Azores ([Murray and Hjort, 1912](#); Roule and Angel, 1923; [Fowler, 1936a](#); Albuquerque, 1954-1956; Koefoed, 1958; Grey, 1964; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Kobyliansky and Herrera, 2015a; Froese and Pauly, 2019) and Madeira (NHMUK 2003.3.18.21-23,

1962; Maul, 1971a; Bekker *et al.*, 1982; Badcock, 1984; Carneiro *et al.*, 2014; Kobyliansky and Herrera, 2015a; Froese and Pauly, 2019).

Family Phosichthyidae

***Ichthyococcus ovatus* (Cocco 1838)**

Common name: Ovate lightfish; **Peixe-luminoso-oval**.

References: mainland Portugal ([MNHN-IC-1883-0064](#), 1882; [Vaillant](#), 1888; [Goode and Bean](#), 1895, 1896; [Brauer](#), 1906; [Seabra](#), 1911; [Albuquerque](#), 1954-1956; [Badcock](#), 1984; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019), Azores ([Murray and Hjort](#), 1912; [Roule and Angel](#), 1924, 1930; [Fowler](#), 1936a; [Collins](#), 1954; [Albuquerque](#), 1954-1956; [Koefoed](#), 1958; [Grey](#), 1964; [Krefft](#), 1983; [Badcock](#), 1984; [Arruda](#), 1997; [Santos et al.](#), 1997; [Porteiro et al.](#), 1999, 2010, 2017; [Sutton et al.](#), 2008; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019) and Madeira ([MNHN-IC-1883-0061](#), 1882; [Badcock](#), 1984; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019).

***Pollimchthys mauli* (Poll 1953)**

Common name: Stareye lightfish; **Peixe-luminoso-de-Maul**.

References: mainland Portugal ([NHMUK 2004.3.8.496](#), 1966; [NHMUK 2003.7.11.18](#), 1984; [Iwamoto](#), 2015b), Azores ([MCZ 140261](#), 1972(PECS); [Sutton et al.](#), 2008(PECS); [Carneiro et al.](#), 2014(PECS); [Iwamoto](#), 2015b) and Madeira ([Iwamoto](#), 2015b). Specimens are needed for confirmation of its presence in Madeira EEZ.

***Polymetme corythaeola* (Alcock 1898)**

Common name: Rendezvous fish; **Peixe-luminoso-do-Índico**.

References: mainland Portugal ([Saldanha](#), 1966b; [Maul](#), 1976; [INIP](#), 1981a, 1982a, b; [Badcock](#), 1984; [Saldanha et al.](#), 1995; [Marques](#), 2001; [Monteiro et al.](#), 2001; [Borges et al.](#), 2002, 2003; [Moura](#), 2004; [EMBL JQ774548](#), 2006; [EMBL JQ774549](#), 2006; [ICES](#), 2005; [Fernandes et al.](#), 2007; [Gomes and Olim](#), 2007; [Silva](#), 2008, 2010, 2011; [Silva et al.](#), 2009; [Costa et al.](#), 2012; [Chaves and Silva](#), 2013; [Carneiro et al.](#), 2014; [Silva and Borges](#), 2014; [Froese and Pauly](#), 2019), Azores (PECS) ([SDSC 22957](#), 1967; [Maul](#), 1976; [Ehrich](#), 1977; [Badcock](#), 1984; [Fock et al.](#), 2002; [Kukuev](#), 2002; [Carneiro et al.](#), 2014) and Madeira ([SDSC 22956](#), 1967 (PECS); [Maul](#), 1976 (PECS); [Ehrich](#), 1977 (PECS); [Badcock](#), 1984; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019). Several specimens caught off in different areas are in the MMF collections. One specimen caught off coast of mainland Portugal (MMF 22866), one set of eleven specimens caught off at the Josephine Bank (MMF 22857, 1967) and one set of eleven specimens in the Meteor Bank (MMF 22541, 1967).

***Polymetme thaeocoryla* Parin & Borodulina 1990**

Common name: No English common name; **Peixe-luminoso-do-Atlântico**.

References: mainland Portugal ([Parin and Borodulina](#), 1990; [Harold](#), 2002; [Kobyliansky and Herrera](#), 2015b), Azores (PECS) ([Parin and Borodulina](#), 1990; [Carneiro et al.](#), 2014) and Madeira (PECS) ([Parin and Borodulina](#), 1990; [Carneiro et al.](#), 2014; [Kobyliansky and Herrera](#), 2015b).

***Vinciguerria attenuata* (Cocco 1838)**

Common name: Slender lightfish; **Peixe-luminoso-esguio**.

References: mainland Portugal ([Jespersen and Taaning](#), 1919, 1926; [Badcock](#), 1984; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019), Azores ([Zugmayer](#), 1911b, 1933; [Brauer](#), 1906; [Murray and Hjort](#), 1912; [Roule and Angel](#), 1924, 1930, 1933; [Fowler](#), 1936b; [Ferreira](#), 1942; [Collins](#), 1954; [Albuquerque](#), 1954-1956; [Koefoed](#), 1958; [Bekker et al.](#), 1982; [Badcock](#), 1984; [Post](#), 1985; [Arruda](#), 1997; [Santos et al.](#), 1997; [Porteiro et al.](#), 1999, 2010, 2017; [Sutton et al.](#), 2008; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019) and Madeira ([Fowler](#), 1936b; [Noronha and Sarmento](#), 1948; [Maul](#), 1949a, 1971a; [Albuquerque](#), 1954-1956; [Bekker et al.](#), 1982; [Badcock](#), 1984; [Carneiro et al.](#), 2014; [Froese and Pauly](#), 2019).

***Vinciguerria lucetia* (Garman 1899)**

Common name: Panama lightfish; **Peixe-luminoso-do-Panamá**.

References: Azores (?) ([Zugmayer](#), 1911b (PECS); [Murray and Hjort](#), 1912; [NHMUK 1934.12.19.6](#), 1912-1936; [Roule and Angel](#), 1924, 1930, 1933; [Fowler](#), 1936a; [Collins](#), 1954) and Madeira (?) ([Roule and Angel](#), 1930). According to Eschmeyer *et al.* (2019) the current geographic distribution of *V. lucetia* is

the Pacific Ocean. Several authors mention the occurrence of this species in the Azores and Madeira (larva), but according to Santos *et al.* (1997) the occurrence records of *V. lucetia* in the Azores corresponded to misidentification of *Vinciguerria attenuata* (Cocco 1838). Thus, this species is not considered in the total number of Madeira and Azores species.

***Vinciguerria nimbaria* (Jordan & Williams 1895)**

Common name: **Oceanic lightfish; Peixe-luminoso-oceânico.**

References: mainland Portugal ([MCZ 138991, 1986](#); [Quéro *et al.*, 1990d](#); [ZSM-Pis-028656 / 533879 / 385467, 1991](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)), Azores ([Zugmayer, 1911b, 1933](#); [Murray and Hjort, 1912](#); [Roule and Angel, 1924, 1930, 1933](#); [Ferreira, 1942](#); [Collins, 1954](#); [Koefoed, 1958](#); [Bekker *et al.*, 1982](#); [Badcock, 1984](#); [Lloris *et al.*, 1991](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Maul, 1949a](#); [MCZ 139200, 1969](#); [Badcock, 1984](#); [Lloris *et al.*, 1991](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)).

***Vinciguerria poweriae* (Cocco 1838)**

Common name: **Power's deep-water bristle-mouth fish; Peixe-luminoso-de-Power.**

References: mainland Portugal ([Jespersen and Taaning, 1926](#); [Badcock, 1984](#); [ICES, 2005](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)), Azores ([Koefoed, 1958](#); [Badcock, 1984](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010, 2017](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Maul, 1949a, 1971a](#); [Albuquerque, 1954-1956](#); [Bekker *et al.*, 1982](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)).

***Yarrella blackfordi* Goode & Bean 1896**

Common name: No English common name; **Peixe-luminoso-de-Blackford.**

References: Azores (PECS) ([Kukuev, 2002](#)).

Family Stomiidae

***Aristostomias grimaldii* Zugmayer 1913**

Common name: No English common name; **Peixe-dragão-de-Grimaldi.**

References: Azores ([MCZ 150372, 1978](#); [Porteiro *et al.*, 1999, 2010](#); [Carneiro *et al.*, 2014](#)) and Madeira ([Gibbs, 1984](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)).

***Aristostomias lunifer* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-lunar.**

References: Azores (PECS) ([ZMH 118630, 1979](#); [NHMUK 1998.8.9.18817, 1981](#); [Carneiro *et al.*, 2014](#)) and Madeira ([USNM 296721, 1971](#); [Gibbs, 1984](#); [Lloris *et al.*, 1991](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)).

***Aristostomias tittmanni* Welsh 1923**

Common name: Loosejaw; **Peixe-dragão-de-Tittmann.**

References: Azores ([NHMUK 1929.7.6.253, 1922](#); [Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Morrow, 1964b](#); [MCZ 149494, 1978](#); [Gibbs, 1984](#); [Lloris *et al.*, 1991](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 1999, 2010, 2017](#); [MAECO385-09, 2004](#); [Sutton *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Harold, 2015a](#); [Froese and Pauly, 2019](#)) and Madeira ([MCZ 131974, 1986](#); [Carneiro *et al.*, 2014](#); [Harold, 2015a](#); [Porteiro *et al.*, 2017](#)).

***Aristostomias xenostoma* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-das-Caraíbas.**

References: Azores (PECS) ([MNHN-IC-1938-0124, 1938](#); [SOC Discovery 1022203_86_FIS_012001, 1980](#); [Carneiro *et al.*, 2014](#)). One paratype of *A. xenostoma* caught off in the Azores is in the MNHN collections ([MNHN-IC-1938-0124](#)).

***Astronesthes atlanticus* Parin & Borodulina 1996**

Common name: No English common name; **Peixe-dragão-do-Atlântico.**

References: Madeira ([ZMH 24891, 1966](#); [Carneiro *et al.*, 2014](#)). One paratype of *A. atlanticus* caught off 39 NM South of Madeira Islands is in the ZMH collections ([ZMH 24891, 1966](#)).

***Astronesthes cyaneus* (Brauer 1902)**

Common name: No English common name; **Peixe-dragão-azulado**.

References: Azores (?) (PECS) ([SOC Discovery 1023214_85_FIS_013010, 1980](#); Carneiro et al., 2014). According to Eschmeyer et al. (2019) the current geographic distribution of *A. cyaneus* is the Indo-West Pacific: from Madagascar east to Hawaiian Islands, north to Ogasawara Islands. Several authors mention the occurrence of this species in the Atlantic, but there is uncertainty about the identification of the Azores specimen. Thus, this species is not considered in the total number of valid species.

***Astronesthes gemmifer* Goode & Bean 1896**

Common name: Snaggletooth; **Peixe-dragão-da-Terra Nova**.

References: mainland Portugal ([ZMH 107009, 1968](#); [USNM 301139, 1970](#); [ZMH 118203, 1979](#); Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([MCZ 133383, 1969](#); Gibbs, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Kukuev, 2002 (PECS); Sutton et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([Roule and Angel, 1933](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Maul, 1956b](#); [NHMUK 1962.12.12.54, 1958](#); Gibbs, 1984; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

***Astronesthes indicus* Brauer 1902**

Common name: Black snaggletooth; **Peixe-dragão-do-Índico**.

References: Madeira ([Gibbs, 1984](#); Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019). According to Eschmeyer et al. (2019) the current geographic distribution of *A. indicus* is the Indo-Pacific: East Africa and Persian Gulf east to Hawaiian Islands and Galapagos Archipelago, South to New Caledonia. However, several authors mention the occurrence of this species in the Atlantic, namely in Madeira and Canary islands ([Badcock, 1970](#); [Bordes et al., 1999](#), [Bordes Caballero et al., 2009](#)) and Cabo Verde ([Reiner, 1996](#)), and assuming that it is a circumglobal species ([Sutton and Hopkins, 1996](#); [Carpenter and De Angelis, 2016b](#)). Specimens are needed for confirmation in Madeira EEZ.

***Astronesthes leucopogon* Regan & Trewavas 1929**

Common name: No English common name; **Peixe-dragão-de-barbilho-branco**.

References: mainland Portugal ([YPM ICH 004518, 1959](#); Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([USNM 359070, 1979 \(PECS\)](#); [ZMH 121510, 1982 \(PECS\)](#); Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([MCZ 96569, 1969](#); Gibbs, 1984; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

***Astronesthes macropogon* Goodyear & Gibbs 1970**

Common name: No English common name; **Peixe-dragão-de-barbilho-graúdo**.

References: Madeira ([Gibbs, 1984](#); Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

***Astronesthes micropogon* Goodyear & Gibbs 1970**

Common name: Snaggletooth; **Peixe-dragão-de-barba-curta**.

References: Azores ([MCZ 91742, 1969](#); Santos et al., 1997; Porteiro et al., 1999, 2010; Carneiro et al., 2014) and Madeira ([USNM 359059, 1978](#); Gibbs, 1984, 1990; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

***Astronesthes neopogon* Regan & Trewavas 1929**

Common name: Snaggletooth; **Peixe-dragão-de-barbilho-plano**.

References: mainland Portugal ([INIP, 1982a](#); Gibbs, 1984; Carneiro et al., 2014), Azores ([Regan and Trewavas, 1929](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [MCZ 149502, 1978](#); Porteiro, 2005; Santos et al., 1997; Porteiro et al., 1999, 2010; Carneiro et al., 2014) and Madeira ([Regan and Trewavas, 1929](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Maul, 1956b](#); [ZMH 104221, 1982](#); Gibbs, 1984, 1990; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

***Astronesthes niger* Richardson 1845**

Common name: Snaggletooth; **Peixe-dragão-manchado**.

References: mainland Portugal ([USNM 301306, 1969](#); [USNM 301289, 1969](#); [INIP, 1982a](#); Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([Murray and Hjort, 1912 \(PECS\)](#); [Regan and Trewavas, 1929](#); [Fowler, 1936a](#); [Belloc, 1938](#); [Albuquerque, 1954-1956](#); Gibbs, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al.,

2014; Froese and Pauly, 2019) and Madeira (Lowe, 1850, 1852; MOM-IC-0091-1366, 1911; Fowler, 1936a; Albuquerque, 1954-1956; Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019). The holotype (unique) of *Astronesthes myriaster* Zugmayer 1913, synonym of *A. niger*, caught off in Madeira is in the MOM collections (MOM-IC-0091-1366).

***Astronesthes richardsoni* (Poey 1852)**

Common name: **Richardson's snaggletooth; Peixe-dragão-de-Richardson.**

References: mainland Portugal (Porteiro, 2005).

***Astronesthes similis* Parr 1927**

Common name: No English common name; **Peixe-dragão-de-bolbo-filamentoso.**

References: Azores (PECS) (MAR-ECO - HamPelFish - TI/SF MAR-ECO 6570, 1979; Carneiro et al., 2014).

According to Eschmeyer et al. (2019) the current geographic distribution of *A. similis* is the Western Atlantic. However, there are three specimens caught off in the tropical and temperate Eastern Atlantic in the MCZ (MCZ 132863 and MCZ 132862) and NHMUK (NHMUK 1929.1.4.83) collections. The location indicated in the NHMUK record is the North-east Atlantic Ocean, East of Portugal, but the geographic coordinates mentioned are outside of the mainland Portugal PECS area (370 NM west of the Portuguese continental coast).

***Astronesthes zharovi* Parin & Borodulina 1998**

Common name: No English common name; **Peixe-dragão-de-Zarov.**

References: Azores (PECS) (SOC Discovery 1023215_86_FIS_013006, 1980; SOC Discovery 1023217_86_FIS_013006, 1980; Carneiro et al., 2014).

***Bathophilus brevis* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-oval.**

References: Azores (Regan and Trewavas 1930; Fowler, 1936b; Albuquerque, 1954-1956; NHMUK 1998.8.9.18346, 1981; NHMUK 1998.8.9.18347, 1981) and Madeira (NHMUK 1929.7.6.19, 1921; MCZ 132781, 1986; MCZ 155709, 1986; MCZ 132780, 1986; MCZ 153111, 1986; Carneiro et al., 2014).

***Bathophilus digitatus* (Welsh 1923)**

Common name: Scaleless black dragonfish; **Peixe-dragão-iridescente.**

References: mainland Portugal (ZSM-Pis-029239 / 534442 / 386030, 1992; Kobylansky and Herrera, 2015c), Azores (NHMUK 1998.8.9.18748, 1981 (PECS); Lloris et al., 1991; MAR-ECO 4822, 2004 (PECS); Carneiro et al., 2014; Kobylansky and Herrera, 2015c; Porteiro et al., 2017) and Madeira (Regan and Trewavas, 1930; Fowler, 1936a; Gibbs, 1984; Gibbs and Barnett, 1990; Lloris et al., 1991; Carneiro et al., 2014; Kobylansky and Herrera, 2015c; Froese and Pauly, 2019).

***Bathophilus longipinnis* (Pappenheim 1914)**

Common name: Longfin dragonfish; **Peixe-dragão-de-barbatana-longa.**

References: mainland Portugal (ZMH 118388, 1979), Azores (Regan and Trewavas, 1930; Fowler, 1936b; Albuquerque, 1954-1956; SOC Discovery 1023215_86_FIS_067009, 1980 (PECS); MAR-ECO 7343, 2004; Sutton et al., 2008; Carneiro et al., 2014; Porteiro et al., 2017) and Madeira (SOC Discovery 826200 8 FIS 067009, 1973; Carneiro et al., 2014).

***Bathophilus metallicus* (Welsh 1923)**

Common name: No English common name; **Peixe-dragão-de-Welsh.**

References: mainland Portugal (NHMUK 1929.7.6.39, 1921; Harold, 2015b), Azores (NHMUK 1929.7.6.28, 1922; NHMUK 1929.7.6.45, 1922; MCZ 84809, 1964; MCZ 84816, 1969; MCZ 84827, 1969; MCZ 84829, 1969; Harold, 2015b) and Madeira (NHMUK 1929.7.6.34, 1921; NHMUK 1929.7.6.29-30, 1921; NHMUK 1929.7.6.35, 1921; NHMUK 1929.7.6.36-38, 1921; NHMUK 1929.7.6.40-41, 1921; NHMUK 1929.7.6.47, 1921; MCZ 132760, 1986; Harold, 2015b). The species *Trichostomias metallicus* Welsh 1923 is synonym of *B. metallicus*. In the original description Welsh (1923) mentioned that *T. metallicus* was close to *Trichostomias vaillanti* Zugmayer 1911, from which it differed chiefly in its much smaller eye, higher number of dorsal and anal rays, and the presence of four small teeth on the palatines. Froese and Pauly (2019) considered *Bathophilus metallicus* (Welsh 1923) as synonym of *Bathophilus vaillanti* (Zugmayer 1911). However, Eschmeyer et al. (2019) considered *Bathophilus metallicus* (Welsh 1923) as a valid species. Thus, some former occurrence records of *B. vaillanti* may

relate to *B. metallicus* or vice versa and an overlapping geographic distribution of these two species occurs. This subject deserves further detailed attention.

Bathophilus nigerrimus Giglioli 1882

Common name: Scaleless dragonfish; **Peixe-dragão-chicote**.

References: Azores (Regan and Trewavas, 1930; Fowler, 1936b; Albuquerque, 1954-1956; MCZ 149509, 1978; Arruda, 1887; Santos et al., 1997; Porteiro et al., 2010, 2017; Carneiro et al., 2014) and Madeira (Regan and Trewavas, 1930; Fowler, 1936b; MCZ 84843, 1978; Carneiro et al., 2014).

Bathophilus pawnee Parr 1927

Common name: **Pawnee dragonfish; Peixe-dragão-de-Pawnee**.

References: Azores (PECS) ([SOC Discovery 1023216 84 FIS 067004, 1980](#); [SOC Discovery 1022207 84 FIS 067006, 1980](#); Carneiro et al., 2014) and Madeira (Porteiro, 2005).

Bathophilus proximus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-das-Bermudas**.

References: Azores (PECS) ([SOC Discovery 1024104_84_FIS_067008, 1980](#); Carneiro et al., 2014).

Bathophilus schizochirus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-cinzento**.

References: Madeira ([Harold, 2015c](#)). According to Harold (2015c) *B. schizochirus* is a circumglobal species occurring in tropical to subtropical waters and Eschmeyer (2019) also mentions a circumglobal distribution in warm seas. The distribution map of this species reported by Harold (2015c) partially included the Madeira EEZ. However, Froese and Pauly (2019) indicate the current distribution of this species between 28°N and 17°S. Thus, the presence of this species in Madeira was not yet confirmed and specimens are needed for confirmation.

Bathophilus vaillanti (Zugmayer 1911)

Common name: Scaleless dragonfish; **Peixe-dragão-de-Vaillant**.

References: mainland Portugal ([MOM-IC-0091-1143, 1910](#); Zugmayer, 1911a, b; Fowler, 1936a; Albuquerque, 1954-1956; Gibbs, 1984; Psomadakis et al., 2012; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([MOM-IC-0091-1553, 1912 \(PECS\)](#); Regan and Trewavas, 1930; Roule and Angel, 1931, 1933; Fowler, 1936b; Albuquerque, 1954-1956; Koefoed, 1956; Dias et al., 1976; Gibbs, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([Regan and Trewavas, 1930](#); Fowler, 1936b; Albuquerque, 1954-1956; Gibbs, 1984; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019). Three unique holotypes - *Trichostomias vaillanti* Zugmayer 1911, *Trichostomias alberti* Roule & Angel 1931 and *Nasistomias curvatus* Koefoed 1956 (synonyms of *B. vaillanti*) - caught off in mainland Portugal are in the MOM (MOM-IC-0091-1143) and the specimens caught off in the Azores are in the MOM and ZMUB collections (MOM-IC-0091-1553 and ZMUB 5293, respectively).

Borostomias antarcticus (Lönnberg 1905)

Common name: Snaggletooth; **Peixe-dragão-do-Antártico**.

References: mainland Portugal ([IFREMER Biocean 34583, 1969](#); Carneiro et al., 2014), Azores ([ZMH 114114, 1975](#); [MAR-ECO 4151, 2004](#); Sutton et al., 2008; Carneiro et al., 2014; Porteiro et al., 2017).

Borostomias elucens (Brauer 1906)

Common name: No English common name; **Peixe-dragão-brilhante**.

References: mainland Portugal ([Harold, 2015d](#); [Ralph, 2015d](#)), Azores (PECS) ([SOC Discovery 1022203_86_FIS_076001, 1980](#); Carneiro et al., 2014; Harold, 2015d) and Madeira ([Gibbs, 1984](#); [Lloris et al., 1991](#); Carneiro et al., 2014; Harold, 2015d; Froese and Pauly, 2019). According to Eschmeyer et al. (2019) *B. elucens* is a cosmopolitan species with a widely tropical geographic distribution (Harold, 2015d). Ralph (2015d) indicates the occurrence of this species in mainland Portugal (on maps). Its presence among the Portuguese ichthyofauna is not yet documented. Specimens are needed for confirmation in mainland Portugal.

Borostomias mononema (Regan & Trewavas 1929)

Common name: Sickle snaggletooth; **Peixe-dragão-abutre**.

References: Madeira ([ZMH 104281, 1966](#); Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Chauliodus danae Regan & Trewavas 1929

Common name: **Dana viperfish**; **Peixe-dragão-do-Dana**, Peixe-víbora.

References: mainland Portugal ([GNM 1946-6476, 1946](#); [MNHN-IC-1987-0354, 1969](#); [MCZ 129206, 1972 \(PECS\)](#); [ZMH 118322, 1979](#); Gibbs, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Regan and Trewavas, 1929; Fowler, 1936b; Ege, 1948; Albuquerque, 1954-1956; Koefoed, 1956; Geistdoerfer *et al.*, 1971; Gibbs, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Regan and Trewavas, 1929; Fowler, 1936b; MCZ 131146, 1969; MCZ 101877, 1978; Gibbs, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Chauliodus sloani Bloch & Schneider 1801

Common name: **Sloane's viperfish**; **Peixe-dragão-de-Sloane**, Demónio, Peixe-demónio.

References: mainland Portugal ([Seabra, 1911](#); Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Fernández *et al.*, 1981; INIP, 1981b, 1982a; Martins, 1981; Gibbs, 1984; Gordo *et al.*, 1990; Borges *et al.*, 2002; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Silva, 2008, 2011; Silva *et al.*, 2009; Carneiro *et al.*, 2014; Silva and Borges, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores (Richard, 1905a; Brauer, 1906; Zugmayer, 1911b, 1933; Murray and Hjort, 1912 (PECS); Roule, 1919a; Roule and Angel, 1924; Regan and Trewavas, 1929, 1930; Fowler, 1936a; Ferreira, 1942; Ege, 1948; Collins, 1954; Albuquerque, 1954-1956; Koefoed, 1956; Morrow, 1964a; Geistdoerfer *et al.*, 1971; Dias *et al.*, 1976; Bekker *et al.*, 1982; Gibbs, 1984; Arruda, 1997; Santos *et al.*, 1997; Melo and Menezes, 2002; Porteiro *et al.*, 1999, 2010, 2017; Arkhipov and Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira ([Collett, 1890b](#); Goode and Bean, 1895, 1896; Brauer, 1906; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Maul, 1948a, 1971a; Albuquerque, 1954-1956; Gibbs, 1984; Davies and Merrett, 1998; Carneiro *et al.*, 2014; Delgado *et al.*, 2017; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005167).

Chirostomias pliopterus Regan & Trewavas 1930

Common name: Scaleless dragonfish; **Peixe-dragão-de-bolbo-papilar**.

References: mainland Portugal ([Gibbs, 1984](#); Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Regan and Trewavas, 1930; Gibbs, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Regan and Trewavas, 1930; Fowler, 1936b; Albuquerque, 1954-1956; Gibbs, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Echiostoma barbatum Lowe 1843

Common name: Threadfin dragonfish; **Peixe-dragão-barbatana-filamentosa**.

References: mainland Portugal ([ZMH 105984, 1968](#); Costa and Reiner, 1978; Gibbs, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores ([YPM ICH 003201, 1931](#); Parr, 1934; Gibbs, 1984; Melo and Menezes, 2002; Arruda, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Lowe, 1843, 1844b; Günther, 1864a; Goode and Bean, 1895, 1896; Brauer, 1906; NHMUK 1917.7.14.82, 1912-1936; Fowler, 1936a; Maul, 1948a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Madeira is in the NHMUK collections (NHMUK 1917.7.14.82).

Eustomias acinosus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-barbilho-acinoso**.

References: Azores (PECS) ([NHMUK 1998.8.9.18746, 1981](#)).

Eustomias braueri Zugmayer 1911

Common name: No English common name; **Peixe-dragão-lusitano**.

References: mainland Portugal ([MOM-IC-0091-1212, 1910](#); Zugmayer, 1911b; Fowler, 1936a; Albuquerque, 1954-1956; Bogutskaya, 2007a; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015a), Azores ([MCZ 149504, 1978](#); Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015a) and Madeira ([MCZ 132611, 1986](#); MCZ 66975, 1986; Swinney, 1990; Carneiro *et al.*,

2014; Papaconstantinou and Kobyliansky, 2015a). The holotype (unique) of this species caught off in mainland Portugal is in the MOM collections (MOM-IC-0091-1212).

Eustomias contiguus Gomon & Gibbs 1985

Common name: No English common name; **Peixe-dragão-bibolboso**.

References: Azores (PECS) ([USNM 322759, 1979](#); [ZMH 25001, 1979](#); [ZMH 25002, 1979](#); Gomon and Gibbs, 1985; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([ZMH 24999, 1968](#); Carneiro et al., 2014).

Eustomias drechseli Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-Drechsel**.

References: Azores (PECS) ([Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); Carneiro et al., 2014). There is a unique holotype of this species (ZMUC P201897) in the fish collection of the Zoological Museum of Copenhagen (Calabuig, 2014). This species is accepted according to Global Names Usage Bank (GNUB, 2017) and Eschmeyer et al. (2019).

Eustomias dubius Parr 1927

Common name: No English common name; **Peixe-dragão-ambíguo**.

References: Azores (PECS) ([SOC Discovery 1024307 86 FIS 261017, 1980](#); Carneiro et al., 2014). According to Eschmeyer et al. (2019) the current geographic distribution of *E. dubius* is the Western Atlantic, but it also occurs in the central Atlantic Ocean off the Azores (30 NM SSW of the Konstantinov Ridge).

Eustomias enbarbatus Welsh 1923

Common name: **Barbate dragonfish; Peixe-dragão-barbado**.

References: Azores (PECS) ([USNM 372074, 1979](#); [USNM 379984, 1979](#); Carneiro et al., 2014) and Madeira ([Swinney, 1990](#); [Porteiro, 2005](#)).

Eustomias filifer (Gilchrist 1906)

Common name: No English common name; **Peixe-dragão-de-barbilho-filífero**.

References: mainland Portugal ([Gibbs, 1984](#)), Azores (PECS) ([Regan and Trewavas, 1930](#); [USNM 322960, 1979](#); [ZMH 120730, 1982](#); [NHMUK 1998.8.9.17848, 1981](#); [Gibbs, 1984](#); Carneiro et al., 2014) and Madeira ([ZMH 104021, 1966](#); [NHMUK 2007.10.31.64, 1966](#); [Gibbs, 1984](#); [Swinney, 1990](#); [Lloris et al., 1991](#); Carneiro et al., 2014; Froese and Pauly, 2019).

Eustomias fissibarbis (Pappenheim 1914)

Common name: No English common name; **Peixe-dragão-de-barbilho-bifurcado**.

References: Azores (PECS) ([ZMH 118383, 1979](#); Carneiro et al., 2014; [Ralph, 2015e](#)) and Madeira ([Gibbs and Barnett, 1990](#); Carneiro et al., 2014; [Ralph, 2015e](#); Froese and Pauly, 2019).

Eustomias furcifer Regan & Trewavas 1930

Common name: Scaleless dragonfish; **Peixe-dragão-de-furfur**.

References: Azores ([Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bogutskaya, 2007b](#); Carneiro et al., 2014; [Harold, 2015e](#)) and Madeira ([Harold, 2015e](#)). Specimens are needed for confirmation in Madeira EEZ.

Eustomias lipochirus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-Santa Cruz**.

References: Azores (PECS) ([USNM 358647, 1979](#); Carneiro et al., 2014) and Madeira ([Gibbs and Barnett, 1990](#); [Swinney, 1990](#); Carneiro et al., 2014; Froese and Pauly, 2019).

Eustomias longibarba Parr 1927

Common name: No English common name; **Peixe-dragão-de-barbilho-crescido**.

References: Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF MAR-ECO 4630, 1979](#); Carneiro et al., 2014) and Madeira ([Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [ZMH 104064, 1966](#); [Gibbs, 1984](#); [Swinney, 1990](#); [Lloris et al., 1991](#); [NHMUK 1996.2.14.32, 1995](#) (PECS); Carneiro et al., 2014; Froese and Pauly, 2019).

Eustomias macronema Regan & Trewavas 1930

Common name: Bigbarb dragonfish; **Peixe-dragão-de-barbijo-grande**.

References: Madeira ([Gibbs, 1984](#); [Swinney, 1990](#); [Clarke, 2000](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Eustomias macrurus Regan & Trewavas 1930

Common name: Yellowstem dragonfish; **Peixe-dragão-de-barbijo-amarelo**.

References: Azores ([Santos et al., 1997](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#)).

Eustomias melanostigma Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-mancha-negra**.

References: Madeira ([ZMH 104068, 1966](#)).

Eustomias monodactylus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-bolbo-monodáctilo**.

References: Azores (PECS) ([USNM 322963, 1979](#); [Carneiro et al., 2014](#)) and Madeira ([Swinney, 1990](#); [Porteiro, 2005](#)).

Eustomias obscurus Vaillant 1884

Common name: Scaleless dragonfish; **Peixe-dragão-obscuro**.

References: mainland Portugal ([Parr, 1927](#); [ZMH 118533, 1979](#); [USNM 258813, 1970](#); [YPM ICH 004717, 1959](#); [Gibbs, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1885-0064, 1883](#); [Filhol, 1884](#); [Vaillant, 1888](#); [Goode and Bean, 1895, 1896](#); [Perrier, 1903](#); [Brauer, 1906](#); [Bertin, 1940](#); [Ferreira, 1942](#); [Regan and Trewavas, 1930](#); [Roule and Angel, 1933](#); [Zugmayer, 1933](#); [Fowler, 1936a](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Morrow and Gibbs, 1964](#); [Gibbs, 1984](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1929.7.6.95-97, 1921](#); [NHMUK 1929.7.6.91, 1921](#); [Maul, 1971a](#); [MCZ 132737, 1986](#); [Gibbs, 1984](#); [Swinney, 1990](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species and a holotype of *Eustomias zugmayeri* Parr 1927 are in the NHMUK and MOM collections (MNHN-IC-1885-0064 caught off in the Azores and MOM-IC-91 1015 caught off in mainland Portugal, respectively). *E. zugmayeri* is synonym of *E. obscurus*.

Eustomias parri Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-Parr**.

References: Azores ([YPM ICH 004038, 1931](#); [YPM ICH 008748, 1931](#); [Carneiro et al., 2014](#)).

Eustomias radicifilis Borodin 1930

Common name: No English common name; **Peixe-dragão-de-barbijo-ramificado**.

References: Azores (PECS) ([SOC Discovery 827000 8 FIS 261016, 1973](#); [Carneiro et al., 2014](#)).

Eustomias satterleei Beebe 1933

Common name: Twinray dragonfish; **Peixe-dragão-de-Satterlee**.

References: Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF MAR-ECO 4664, 1979](#); [Clarke, 2000](#); [Carneiro et al., 2014](#)).

Eustomias schiffi Beebe 1932

Common name: No English common name; **Peixe-dragão-de-Schiff**.

References: Azores (PECS) ([NHMUK 1998.8.9.17845, 1981](#); [NHMUK 1998.8.9.17905, 1981](#); [Porteiro, 2005](#); [Carneiro et al., 2014](#)).

Eustomias schmidti Regan & Trewavas 1930

Common name: Schmidt's dragonfish; **Peixe-dragão-de-Schmidt**.

References: Azores (PECS) ([USNM 292889, 1979](#); [SOC Discovery 1022825 84 FIS 261013, 1980](#); [SOC Discovery 1022825 85 FIS 261013, 1980](#); [SOC Discovery 1022825 86 FIS 261013, 1980](#); [Carneiro et al., 2014](#)) and Madeira ([ZMH 106069, 1968](#); [Gibbs, 1984](#); [Swinney, 1990](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Eustomias simplex Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-bolbo-simples**.

References: Azores (PECS) ([USNM 372019](#), 1979; [USNM 379335](#), 19779; SOC Discovery 1022804 85 FIS 261010, 1980; SOC Discovery 1022804 86 FIS 261010, 1980; Carneiro *et al.*, 2014) and Madeira (Gibbs, 1984; Swinney, 1990; Froese and Pauly, 2019).

Eustomias tenisoni Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-Tenison**.

References: Madeira (PECS) ([NHMUK 2007.2.26.13](#), 1990; Carneiro *et al.*, 2014).

Eustomias tetranema Zugmayer 1913

Common name: No English common name; **Peixe-dragão-da-Madeira**.

References: Azores ([MCZ 101714](#), 1984 (PECS); Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira ([MOM-IC-0091-1450](#), 1912; Zugmayer, 1913; Regan and Trewavas, 1930; Roule and Angel, 1933(?); Fowler, 1936a; Albuquerque, 1954-1956; ZMH 106055, 1968; NHMUK 1995.3.7.2, 1979; Gibbs, 1984; MCZ 97621, 1986; Swinney, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of this species caught off SW of Madeira (31°45.0'N, 20°17.0'W) is in the MOM collections.

Flagellostomias boureei (Zugmayer 1913)

Common name: Longbarb dragonfish; **Peixe-dragão-dos-Açores**.

References: mainland Portugal ([YPM ICH 004723](#), 1959; [MCZ 132486](#), 1972; Gibbs, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores ([MOM-IC-0091-1524](#), 1912; Zugmayer, 1913; Regan and Trewavas, 1930; Roule and Angel, 1933; Fowler, 1936b; Ferreira, 1942; Belloc, 1949; Morrow and Gibbs, 1964; Geistdoerfer *et al.*, 1971; Gibbs, 1984; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019) and Madeira (Regan and Trewavas, 1930; Fowler, 1936b; Gibbs, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Eustomias boureei* Zugmayer 1913, synonym of *F. boureei*, caught off south-west of the Azores (38°55.0'N, 34°7.50'W) is in the MOM collections ([MOM-IC-0091-1524](#)).

Grammatostomias circularis Morrow 1959

Common name: No English common name; **Peixe-dragão-de-mancha-redonda**.

References: Madeira ([ZMH 105793](#), 1968; Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Grammatostomias dentatus Goode & Bean 1896

Common name: No English common name; **Peixe-dragão-colmilhoso**.

References: mainland Portugal ([YPM ICH 004745](#), 1959; [YPM ICH 004830](#), 1959; Carneiro *et al.*, 2014), Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF MAR-ECO 4330](#), 1979; Carneiro *et al.*, 2014) and Madeira (Swinney, 1988; Carneiro *et al.*, 2014).

Grammatostomias flagellibarba Holt & Byrne 1910

Common name: No English common name; **Peixe-dragão-de-barbijo-filamentoso**.

References: mainland Portugal (PECS) ([NHMUK 2013.8.12.12-13](#), 1984), Azores (PECS) ([ZMH 119421](#), 1979; Gibbs, 1984(?)) and Madeira ([MCZ 70504](#), 1978; Swinney, 1988; Carneiro *et al.*, 2014).

Heterophotus ophistoma Regan & Trewavas 1929

Common name: Wingfin snaggletooth; **Peixe-dragão-de-boca-de-serpente**.

References: mainland Portugal ([NHMUK 2004.2.27.134](#), 1966).

Idiacanthus fasciola Peters 1877

Common name: Ribbon sawtail fish; **Peixe-dragão-de-cauda-serreada**.

References: mainland Portugal ([Nobre](#), 1935; GNM 1946-7576, 1946; NHMUK 1988.2.25.52, 1955; IFREMER Biocean 34545, 1969; NHMUK 2014.8.17.104, 1984 (PECS); Gibbs, 1984; ZSM-Pis-028703 / 533924 / 385512, 1991; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015b; Froese and Pauly, 2019), Azores ([Murray and Hjort](#), 1912 (PECS); Roule and Angel, 1924, 1933; Regan and Trewavas, 1930; Fowler, 1936a, b; Ferreira, 1942; Gibbs, 1964, 1984; Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015b; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira ([Maul](#), 1971a; MCZ 89605, 1978; MCZ 89603, 1978; Gibbs, 1984; Davies and Merrett, 1998; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015b; Froese and Pauly, 2019).

***Leptostomias gladiator* (Zugmayer 1911)**

Common name: Scaleless dragonfish; **Peixe-dragão-gladiador**.

References: mainland Portugal ([Gibbs, 1984](#); [Carneiro et al., 2014](#); [Papaconstantinou and Kobylansky, 2015c](#); [Froese and Pauly, 2019](#)), Azores ([Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Gibbs, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#); [Papaconstantinou and Kobylansky, 2015c](#); [Froese and Pauly, 2019](#)) and Madeira ([ZMH 105809, 1968](#); [ZMH 105828, 1968](#); [Gibbs, 1984](#); [Carneiro et al., 2014](#); [Papaconstantinou and Kobylansky, 2015c](#); [Froese and Pauly, 2019](#)).

***Leptostomias haplocaulus* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-de-barbijo-singelo**.

References: mainland Portugal ([YPM ICH 004751, 1959](#); [YPM ICH 004752, 1959](#); [ZMH 105783, 1968](#); [Carneiro et al., 2014](#)), Azores ([Gibbs, 1984](#); [Arruda, 1997](#); [IMAR/DOP COLETA 1694, 2001](#); [Melo and Menezes, 2002](#); [Porteiro et al., 2010](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Gibbs, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Leptostomias leptobolus* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-de-bolbo-alongado**.

References: Madeira ([ZMH 104372, 1966](#); [Porteiro, 2005](#)).

***Leptostomias longibarba* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-de-barbijo-longo**.

References: mainland Portugal ([Ralph, 2015f](#)), Azores ([Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Melo and Menezes, 2002](#); [Porteiro et al., 2010](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Ralph, 2015f](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *L. longibarba* is the North Atlantic. The distribution map of this species reported by Ralph (2015f) indicates the occurrence of this species in Madeira and mainland Portugal. Specimens are needed for confirmation in Madeira and mainland Portugal.

***Malacosteus niger* Ayres 1848**

Common name: **Stoplight loosejaw**; **Peixe-dragão-de-draga**.

References: mainland Portugal ([MNHN-IC-1987-0339, 1969](#); [Gibbs, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1885-0063, 1883](#); [Vaillant, 1888](#); [Goode and Bean, 1895, 1896](#); [Murray and Hjort, 1912](#); [Roule, 1919a](#); [Fowler, 1936a](#); [Ferreira, 1942](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [Morrow, 1964b](#); [Gibbs, 1984](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Roule and Angel, 1933](#); [Fowler, 1936b](#); [Maul, 1948a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Gibbs, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Melanostomias bartonbeani* Parr 1927**

Common name: **Scaleless black dragonfish**; **Peixe-dragão-de-Barton Bean**.

References: mainland Portugal ([Gibbs, 1984](#); [ZSM-Pis-029421 / 534608 / 386196, 1990](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Regan and Trewavas, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Gibbs, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Gibbs, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Melanostomias biseriatus* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-de-barbijo-bisseriado**.

References: Azores ([PECS](#)) ([SOC Discovery 1024117_84_FIS_518003, 1980](#); [SOC Discovery 1024405_84_FIS_518003, 1980](#); [Carneiro et al., 2014](#)) and Madeira ([ZMH 110463, 1971](#); [MCZ 70571, 1978](#); [Gibbs, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Melanostomias macrophotus* Regan & Trewavas 1930**

Common name: No English common name; **Peixe-dragão-luminoso**.

References: mainland Portugal ([ZMH 105820, 1968](#); [USNM 300329, 1970](#); [Gibbs, 1984](#); [Gibbs and Barnett, 1990](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([USNM 358661, 1979](#); [Gibbs, 1984](#);

Arruda, 1997; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019) and Madeira (MCZ 70575, 1978; Gibbs and Barnett, 1990; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Melanostomias melanopogon Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-barbilho-negro**.

References: Azores (Arruda, 1997; MAR-ECO 3449, 2004; Carneiro *et al.*, 2014).

Melanostomias melanops Brauer 1902

Common name: No English common name; **Peixe-dragão-de-Sumatra**.

References: Azores (PECS) (USNM 358821, 1979; USNM 358822, 1979; Carneiro *et al.*, 2014; Harold, 2015f) and Madeira (NHMUK BMNH 2007.10.31.11, 1966; MCZ 132284, 1969; Gibbs, 1984; ZSM-Pis-029424 / 534609 / 386197, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Harold, 2015f; Froese and Pauly, 2019).

Melanostomias spilorhynchus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-focinho-manchado**.

References: mainland Portugal (ZMH 105785, 1968; Harold, 2015g), Azores (Santos *et al.*, 1997; Harold, 2015g), and Madeira (ZMH 102886, 1964; ZMH 110418, 1971; MNHN-IC-1938-0120, 1938; Harold, 2015g). This species was previously considered a synonym of *Melanostomias bartonbeani* Parr 1927.

Melanostomias tentaculatus (Regan & Trewavas 1930)

Common name: Tentacle dragonfish; **Peixe-dragão-tentaculado**.

References: Azores (PECS) (USNM 358831, 1979; USNM 358662, 1979; MAR-ECO - HamPelFish - TI/SF_MAR-ECO_2675, 1982; Carneiro *et al.*, 2014) and Madeira (ZMH 104226, 1966; ZMH 104227, 1966; ZMH 105803, 1968; MCZ 132200, 1970; Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Melanostomias valdiviae Brauer 1902

Common name: **Valdivia black dragon fish; Peixe-dragão-de-Valdívia**.

References: mainland Portugal (ZMH 105911, 1968; Gibbs, 1984; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015d; Froese and Pauly, 2019), Azores (MCZ 132155, 1969; Gibbs, 1984; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015d) and Madeira (ZMH 104379, 1966; Gibbs, 1984; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015d; Froese and Pauly, 2019).

Neonesthes capensis (Gilchrist & von Bonde 1924)

Common name: Cape snaggletooth; **Peixe-dragão-do-Cabo**.

References: mainland Portugal (Gibbs, 1984; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015e; Froese and Pauly, 2019), Azores (YPM ICH 007712, 1931; Gibbs, 1984; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015e; Porteiro *et al.*, 2017; Froese and Pauly, 2019) and Madeira (MCZ 85924, 1969; Gibbs, 1984; MCZ 153108, 1986; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015e; Froese and Pauly, 2019).

Pachystomias microdon (Günther 1878)

Common name: Smalltooth dragonfish; **Peixe-dragão-dente-curto**.

References: mainland Portugal (Gibbs, 1984; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015f; Froese and Pauly, 2019), Azores (MAR-ECO 7399, 2004; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Papaconstantinou and Kobyliansky, 2015f; Porteiro *et al.*, 2017) and Madeira (Papaconstantinou and Kobyliansky, 2015f). The holotype (unique) of *Aristostomias brattstroemii* Koefoed 1956, synonym of *P. microdon*, caught off South of the Azores (out of the PECS) is in the ZMUB collections (ZMUB 5291).

Photonectes braueri (Zugmayer 1913)

Common name: Brauer's dragonfish; **Peixe-dragão-de-Brauer**.

References: Azores (MOM-IC-0091-1555, 1912; Zugmayer, 1913; Roule and Angel, 1933; Fowler, 1936b; Albuquerque, 1954-1956; Koefoed, 1956; Morrow and Gibbs, 1964; Gibbs, 1984; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019) and Madeira (Regan and Trewavas, 1930; Fowler, 1936b; Albuquerque, 1954-1956; Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019).

The holotype (unique) of *Melanostomias braueri* Zugmayer 1913, synonym of *P. braueri*, caught off West of the Azores is in the MOM collections (MOM-IC-0091-1555).

Photonectes dinema Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-bifilar**.

References: Azores (Gibbs, 1984; Lloris *et al.*, 1991; Arruda, 1997; Melo and Menezes, 2002; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira ([MCZ 55082](#), 1970; Gibbs, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Photonectes leucospilus Regan & Trewavas 1930

Common name: No English common name; **Peixe-dragão-de-pinta-branca**.

References: Azores (PECS) ([NHMUK 1998.8.9.17904](#), 1981; [NHMUK 1998.8.9.18230](#), 1981) and Madeira ([ZMH 110405](#), 1971).

Photonectes margarita (Goode & Bean 1896)

Common name: No English common name; **Peixe-dragão-boca-negra**.

References: Azores (MOM-IC-0091-1554, 1912; Zugmayer, 1913; Roule and Angel, 1931, 1933; Fowler, 1936b; Albuquerque, 1954-1956; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014) and Madeira (Gibbs and Barnett, 1990; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Echiostoma richardi* Zugmayer 1913, synonym of *P. margarita*, caught off West of the Azores (39°19.0'N, 35° 24.50'W) is in the MOM collections (MOM-IC-0091-1554).

Photonectes mirabilis Parr 1927

Common name: Blueband dragonfish; **Peixe-dragão-maravilhoso**.

References: Azores (PECS) ([ZMH 118251](#), 1979; Carneiro *et al.*, 2014) and Madeira ([MCZ 132040](#), 1972; Carneiro *et al.*, 2014).

Photonectes parvimanus Regan & Trewavas 1930

Common name: Fleshfin dragonfish; **Peixe-dragão-barbatana-curta**.

References: mainland Portugal ([YPM ICH 004758](#), 1959; [YPM ICH 004759](#), 1959; Carneiro *et al.*, 2014), Azores (PECS) ([Regan and Trewavas, 1930](#); Fowler, 1936b; Albuquerque, 1954-1956; Carneiro *et al.*, 2014) and Madeira ([Regan and Trewavas, 1930 \(PECS\)](#); ZMH 103970, 1966; ZMH 106090, 1968; MCZ 55096, 1969; Gibbs, 1984; Lloris *et al.*, 1991; Davies and Merrett, 1998; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Photonectes fimbria* Regan & Trewavas 1930, synonym of *P. parvimanus*, caught off West of Madeira is in the ZMUC collections (ZMUC P201920).

Photostomias atrox (Alcock 1890)

Common name: No English common name; **Peixe-dragão-atroz**.

References: Azores (PECS) ([Harold, 2015h](#)) and Madeira (Kenaley and Hartel, 2005; Carneiro *et al.*, 2014; Harold, 2015h).

Photostomias goodyeari Kenaley & Hartel 2005

Common name: No English common name; **Peixe-dragão-de-Goodyear**.

References: Azores ([MCZ 150373](#), 1978; Kenaley and Hartel, 2005; Carneiro *et al.*, 2014) and Madeira ([USNM 376101](#), 1971; Kenaley and Hartel, 2005; Carneiro *et al.*, 2014).

Photostomias guernei Collett 1889

Common name: Loosejaw; **Peixe-dragão-de-Guerne**.

References: mainland Portugal ([Nobre, 1935](#); Albuquerque, 1954-1956; Gibbs, 1984; Quéro *et al.*, 2003; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Froese and Pauly, 2019), Azores (MOM-IC-0091-50, 1887; Collett, 1889b, 1896a; Lütken, 1892; Goode and Bean, 1895, 1896; Richard, 1900; Perrier, 1903; Joubin, 1905; Brauer, 1906; Zugmayer, 1911b, 1933; Murray and Hjort, 1912 (PECS); Regan and Trewavas, 1930; Nobre, 1935; Fowler, 1936a, b; [MNHN-IC-1938-0123](#), 1938; Ferreira, 1942; Belloc, 1949; Collins, 1954; Koefoed, 1956; Morrow, 1964b; Gibbs, 1984; Arruda, 1992, 1997; Albuquerque, 1954-1956; Santos *et al.*, 1997; Kenaley and Hartel, 2005; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Regan and Trewavas, 1930; Maul, 1952b; USNM 225026, 1971; USNM 225027, 1971; Gibbs, 1984; Kenaley and Hartel, 2005; Carneiro *et al.*, 2014).

al., 2014; Froese and Pauly, 2019). The holotype (unique) of this species caught off in the Azores (38°34.50'N, 30°43.30'W) is in the MOM collections (MOM-IC-0091-0050). *Photostomias mirabilis* (Beebe 1933) is synonym of *P. guernei*.

***Rhadinesthes decimus* (Zugmayer 1911)**

Common name: Slender snaggletooth; **Peixe-dragão-delgado**.

References: mainland Portugal (Zugmayer, 1911a; Nybelin, 1946; Carneiro et al., 2014; Roberts et al., 2017; Froese and Pauly, 2019), Azores (USNM 272910, n.d (PECS); USNM 358792, 1979 (PECS); Arruda, 1997; Porteiro et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (MCZ 86132, 1969; Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019). The holotypes (uniques) of *Rhadinesthes jacobsoni* Nybelin 1946 and *Astronesthes decimus* Zugmayer 1911, which are synonyms of *R. decimus*, both caught off in mainland Portugal are in the NHMG collections (NHMG pi. ex. 1354) and MOM collections (MOM-IC-0091-0959).

***Stomias boa* (Risso 1810)**

Common name: **Boa dragonfish; Peixe-dragão-jibóia**.

References: mainland Portugal (Vaillant, 1888; Goode and Bean, 1895, 1896; Seabra, 1911; Osório, 1917; Albuquerque, 1954-1956; INIP, 1982a(?); Gibbs, 1984; Borges et al., 2002; ICES, 2005; Silva et al., 2009; Silva, 2010; Chaves and Silva, 2013; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Mecklenburg et al., 2018; Eschmeyer et al., 2019), Azores (Reinhardt, 1842; Zugmayer, 1911b; Murray and Hjort, 1912 (PECS); Ege, 1918, 1934; Roule and Angel, 1933; Bruun, 1936; Fowler, 1936a; Belloc, 1938; Collins, 1954; Albuquerque, 1954-1956; Koefoed, 1956; Gibbs, 1969; Backus et al., 1970; Shcherbachov and Novikova, 1976; Gibbs, 1984; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Mecklenburg et al., 2018) and Madeira (Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Maul, 1956b; Gibbs, 1984; Carneiro et al., 2014; Mecklenburg et al., 2018; Froese and Pauly, 2019).

***Stomias brevibarbus* Ege 1918**

Common name: No English common name; **Peixe-dragão-barbilho-curto**.

References: mainland Portugal (Ege, 1934; ZSM-Pis-028731 / 533952 / 385540, 1991; ZSM-Pis-028732 / 533953 / 385541, 1991; Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores (Ege, 1934; MCZ 128677, 1969; Santos et al., 1997; Porteiro et al., 1999, 2010; Carneiro et al., 2014; Priede, 2017) and Madeira (Ege, 1934; Maul, 1948a, 1956b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Gibbs, 1984; Lloris et al., 1991; Davies and Merrett, 1998; Carneiro et al., 2014; Froese and Pauly, 2019).

***Stomias longibarbus* (Brauer 1902)**

Common name: **Longbarb scaly dragonfish; Peixe-dragão-escamudo**.

References: mainland Portugal (ZMH 106044, 1968; Gibbs, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores (Murray and Hjort, 1912(?); NHMUK 2003.1.13.100, 1970; IMAR/DOP COLETA 8889, 2010; Carneiro et al., 2014) and Madeira (ZMH 110608, 1971; SOC Discovery 826200_8_FIS_511001, 1973; NHMUK 1998.8.9.8091, 1973; Gibbs, 1984; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

***Trigonolampa miriceps* Regan & Trewavas 1930**

Common name: Threelights dragonfish; **Peixe-dragão-listado**.

References: Azores (MCZ 114690, 1964; MCZ 62186, 1972; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Carneiro et al., 2014).

Order Ateleopodiformes

Family Ateleopodidae

***Guentherus altivelis* Osório 1917**

Common name: **Jellynose; Peixe-de-vela-alta**.

References: mainland Portugal (Trunov and Isarev, 1971; Costa and Reiner, 1977; Hureau and Monod, 1979; Paxton, 1986; Carneiro et al., 2014; Froese and Pauly, 2019).

Order Aulopiformes

Family Aulopidae

***Aulopus filamentosus* (Bloch 1792)**

Common name: **Royal flagfin; Lagarto-do-mar**, Lagarto-do-alto, Peixe-lagarto, Lagarto.

References: mainland Portugal ([Bragança, 1903](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Thompson and Gormon, 2016](#)), Azores ([Hilgendorf, 1888a](#); [Nobre, 1924, 1930](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [Shcherbachov et al., 1985b \(PECS\)](#); [Dias and Monteiro, 1986](#); [Silva and Menezes, 1995b](#); [Arruda, 1992, 1997](#); [Silva et al., 1995](#); [Menezes, 1996](#); [Santos et al., 1997](#); [Sousa et al., 1999](#); [Kukuev, 2002 \(PECS\)](#); [Diekmann et al., 2006 \(PECS\)](#); [Menezes et al., 2006](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Gomes-Pereira et al., 2014](#); [Thompson and Gormon, 2016](#)) and Madeira ([Lowe, 1837, 1843](#); [Günther, 1864a](#); [Fowler, 1923, 1936a](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Maul, 1946a, 1976 \(PECS\)](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Ehrich, 1977](#); [Fischer et al., 1981](#); [Sulak, 1984](#); [Lloris et al., 1991](#); [Ferreira et al., 2008](#); [Christiansen et al., 2009](#); [Carneiro et al., 2014](#); [POGID, 2016](#); [Thompson and Gormon, 2016](#); [Froese and Pauly, 2019](#)). Several specimens caught off in different areas are in the MMF collections: one specimen at the Josephine Bank (MMF 22857, 1967) and at the Meteor Bank (MMF 22347, 1967, one specimen; MMF 22378, 1967, set of three specimens; MMF 22404, 1967, set of four specimens; MMF 22348, 1967, one specimen; MMF 22349, 1967, one specimen; MMF 22874, 1967, one specimen and MMF 22370, 1967, one specimen).

Family Chlorophthalmidae

***Chlorophthalmus agassizi* Bonaparte 1840**

Common name: **Shortnose greeneye; Olho-verde, Olho-vivo**.

References: mainland Portugal ([Osório, 1917](#); [Fernández et al., 1981](#); [Sulak, 1984](#); [MNHN-IC-1988-0512, 1984](#); [Fischer et al., 1987](#); [Merrett, 1990](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [OCEANA, 2014](#); [Garcia, 2015b](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1885-0089, 1883](#); [MNHN-IC-1885-0090, 1883](#); [Vaillant, 1888](#); [Goode and Bean, 1895, 1896](#); [Perrier, 1903](#); [Fowler, 1936a](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Maul, 1976](#); [Ehrich, 1977](#); [Carvalho, 1982](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Fock et al., 2002 \(PECS\)](#); [Kukuev, 2002 \(PECS\)](#); [Moura, 2004](#); [Diekmann et al., 2006 \(PECS\)](#); [Menezes et al., 2006](#); [Arkhipov & Mamedov, 2008 \(PECS\)](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010, 2013](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Garcia, 2015b](#); [Russell et al., 2015](#)) and Madeira ([Maul, 1976](#); [Ehrich, 1977](#); [Sulak, 1984](#); [Carneiro et al., 2014](#); [Garcia, 2015](#); [Russell et al., 2015](#)). Nine specimens caught off at the Meteor Bank are in the MMF collections (MMF 22407, 1967, set of four specimens; MMF 22544, 1967, one specimen and MMF 22837, 1967, set of four specimens).

Family Ipnopidae

***Bathymicrops multispinis* Nielsen & Merrett 1992**

Common name: No English common name; **Peixe-tripé-espinhoso**.

References: Madeira (PECS) ([NHMUK 1997.1.2.2, 1990](#); [USNM 344624, 1990](#); [Carneiro et al., 2014](#)).

***Bathymicrops regis* Hjort & Koefoed 1912**

Common name: No English common name; **Peixe-tripé-real**.

References: mainland Portugal ([Christiansen et al., 2014](#)), Azores (PECS) ([SDSC 16042, 1982](#); [Shcherbachov et al., 1985a](#); [Carneiro et al., 2014](#)) and Madeira ([Nielsen, 1979b](#); [Sulak, 1984](#); [Merrett, 1990](#); [Lloris et al., 1991](#); [NHMUK 1996.2.14.30, 1995](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Bathypterois dubius* Vaillant 1888**

Common name: **Spiderfish; Peixe-tripé-filamentoso**, Peixe-tripé.

References: mainland Portugal ([Zugmayer, 1911](#); [Maul, 1976](#); [Sulak, 1977, 1984](#); [MNHN-IC-1988-0511, 1984](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)) and Azores ([MNHN-IC-1885-0121, 1883](#); [Vaillant, 1888, 1919](#); [Good and Bean, 1895](#); [Collett, 1896a](#); [Richard, 1900, 1910](#); [Perrier,](#)

1903; Brauer, 1906; Brian, 1908, 1912; Zugmayer, 1911; Roule, 1919a; Koefoed, 1927; Fowler, 1936a; Collins, 1954; Albuquerque, 1954-1956; Grey, 1956; Golovan, 1976; Maul, 1976; Saldanha, 1977a; Sulak, 1977, 1984; Shcherbachov *et al.*, 1985a; Arruda, 1992, 1997; Santos *et al.*, 1997; Kukuev, 2002 (PECS); Melo and Menezes, 2002; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The lectotype of this species caught off in the Azores is in MNHN collections (MNHN-IC-1885-121). Several specimens caught off in the mainland Portugal coast are in the MMF collections (MMF 22491, 1967, set of sixteen specimens).

***Bathypterois grallator* (Goode & Bean 1886)**

Common name: **Tripodfish; Peixe-tripé.**

References: mainland Portugal ([GNM 1946-6606, 1946](#); Nybelin, 1946; Sulak, 1977; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014; Bannermann *et al.*, 2015c), Azores ([MNHN-IC-1978-0500, 1971](#); Rannou and Gaborit-Rezzouk, 1976; Sulak, 1977, 1984; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Franco *et al.*, 2009; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Bannermann *et al.*, 2015c; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira ([Bannermann *et al.*, 2015c](#); Froese and Pauly, 2019). The holotype (unique) of *Skagerakia nilssoni* Nybelin 1946, synonym of *B. grallator*, caught off at 77 NM SSW of Cape Santa Maria (Algarve), 35°43.0'N, 8°16.0'W is in the NHMG collections.

***Bathypterois longipes* Günther 1878**

Common name: **Abyssal spiderfish; Peixe-tripé-abissal.**

References: mainland Portugal ([Sulak, 1977](#); NHMUK 1994.9.20.2-3, 1983; NHMUK 1991.7.9.864, 1989; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014), Azores ([Murray and Hjort, 1912 \(PECS\)](#); Fowler, 1936a (PECS); MCZ 36634, 1947; Nybelin, 1951; Geistdoerfer *et al.*, 1971; Sulak, 1977, 1984; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Kukuev, 2002 (PECS); Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira ([Sulak, 1977, 1984](#); NHMUK 2016.3.10.66-84, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Bathypterois phenax* Parr 1928**

Common name: Blackfin spiderfish; **Peixe-tripé-de-barbatana-negra.**

References: Azores ([Rannou and Gaborit-Rezzouk, 1976](#); Hureau and Monod, 1979; Sulak, 1984; Arruda, 1997; Santos *et al.*, 1997; Kukuev, 2002 (PECS); Bergstad *et al.*, 2008a (PECS); Franco *et al.*, 2009; Porteiro *et al.*, 2010, 2013, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Bathytyphlops marionae* Mead 1958**

Common name: Marion's spiderfish; **Peixe-tripé-de-Marion.**

References: Azores (PECS) ([Kukuev, 2002](#); Bannermann *et al.*, 2015d).

***Bathytyphlops sewelli* (Norman 1939)**

Common name: No English common name; **Peixe-tripé-de-Sewell.**

References: Azores ([MNHN-IC-1970-0030, 1969](#); Geistdoerfer *et al.*, 1970; Nielsen, 1979b; Sulak, 1984; Lloris *et al.*, 1991; Arruda, 1997; Franco *et al.*, 2009; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Bathytyphlops azorensis* Geistdoerfer, Hureau & Rannou 1970, synonym of *B. sewelli*, caught off at 88 NM SW of São Miguel (Azores) is in the MNHN collections (MNHN-IC-1970-0030).

***Discoverichthys praecox* Merrett & Nielsen 1987**

Common name: No English common name; **Peixe-tripé-temporão.**

References: Azores (PECS) ([NHMUK 1986.7.23.1., 1985](#); Merrett and Nielsen, 1987). The holotype (unique) of this species caught off at 150 NM East of Hyères Seamount, 31°7.30'N, 25°5.40'W (Monaco Basin) is in the NHMUK collections (NHMUK 1986.7.23.1).

Family Scopelarchidae

***Benthalbella infans* Zugmayer 1911**

Common name: **Zugmayer's pearleye; Olho-pérola-de-Zugmayer.**

References: mainland Portugal ([MOM-IC-0091-1188, 1910](#); Zugmayer, 1911a; Fowler, 1936a; YPM ICH 004882, 1959; Johnson, 1984; ZSM-Pis-029192 / 534400 / 385988, 1992; Carneiro *et al.*, 2014), Azores

(Roule and Angel, 1924; Merrett *et al.*, 1973; Johnson, 1974, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014) and Madeira (MCZ 70344, 1969; MCZ 70352, 1972; Johnson, 1984; Bekker *et al.*, 1982; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off at North of Madeira is in the MOM collections (MOM-IC-0091-1188).

***Rosenblattichthys hubbsi* Johnson 1974**

Common name: Hubb's pearleye; **Olho-pérola-de-Hubb.**

References: Azores (MCZ 52223, 1973 (PECS); MCZ 52225, 1973 (PECS); Carneiro *et al.*, 2014 (PECS); Bannermann *et al.*, 2015a). Bannermann *et al.* (2015a) include the Azorean EEZ in the distribution area of *R. hubbsi*.

***Scopelarchus analis* (Brauer 1902)**

Common name: Short fin pearleye; **Olho-pérola-de-barbatana-curta.**

References: mainland Portugal (GNM 1946-16269, 1946; GNM 1946-16270, 1946; MCZ 69303, 1969; MCZ 69304, 1969; Johnson, 1984; Carneiro *et al.*, 2014), Azores (MOM-IC-0091-765, 1914; MOM-IC-0091-757, 1914; Roule, 1916, 1919; Fowler, 1936a; Belloc, 1949; Albuquerque, 1954-1956; Koefoed, 1955; Johnson, 1974; Maul, 1979b; Bekker *et al.*, 1982; Johnson, 1984; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Maul, 1946a, 1979b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Bekker *et al.*, 1982; Johnson, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The syntypes of *Odontostomus perarmatus* Roule 1916, synonym of *S. analis*, both caught off at North-east of the Azores are in the MOM collections (MOM-IC-0091-757 and MOM-IC-0091-765 (set of three specimens)).

***Scopelarchus guentheri* Alcock 1896**

Common name: Staring pearleye; **Olho-pérola-de-Guenther.**

References: mainland Portugal (ZSM-Pis-029180 / 534388 / 385976, 1992) and Azores (Kukuev, 2002 (PECS); MAR-ECO 5644, 2004 (PECS); Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014).

***Scopelarchus michaelsarsi* Koefoed 1955**

Common name: Bigfin pearleye; **Olho-pérola-de-Michael Sars.**

References: Azores (MCZ 71113, 1973 (PECS); Carneiro *et al.*, 2014 (PECS); Ralph and Carpenter, 2015) and Madeira (Ralph and Carpenter, 2015). Ralph and Carpenter (2015) include the Azorean and Madeiran EEZs in the distribution area of *S. michaelsarsi*. Specimens are needed for confirmation its presence in the Azores and Madeira EEZ's.

Family Notosudidae

***Ahliasaurus berryi* Bertelsen, Krefft & Marshall 1976**

Common name: No English common name; **Peixe-vigia-de-Berry.**

References: mainland Portugal (Krefft, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Bertelsen *et al.*, 1976; Krefft, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (ZMH 24865, 1966; ZMH 24852, 1966; Krefft, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Scopelosaurus argenteus* (Maul 1954)**

Common name: **Waryfish; Peixe-vigia-prateado.**

References: mainland Portugal (Krefft, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Bertelsen *et al.*, 1976; Krefft, 1984; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Maul, 1954b; Hureau and Monod, 1979; Krefft, 1979c, 1984, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype and paratype of *Notosudis argenteus* Maul 1954, synonym of *S. argenteus*, both found on beach at Porto Santo Island (Madeira) are in the MMF collections (MMF 3998 and MMF 4341, respectively).

***Scopelosaurus lepidus* (Krefft & Maul 1955)**

Common name: **Blackfin wavyfish; Peixe-vigia-de-barbatana-preta.**

References: mainland Portugal (Krefft, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (MCZ 126962, 1972; Bertelsen *et al.*, 1976; Krefft, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Krefft, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Scopelosaurus smithii* Bean 1925**

Common name: No English common name; **Peixe-vigia-de-Smith.**

References: Azores (Arruda, 1997; MAR-ECO 3535, 2004; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017) and Madeira (Maul, 1954; Hureau and Monod, 1979; Krefft, 1979c, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019).

Family Synodontidae

***Synodus foetens* (Linnaeus 1766)**

Common name: Inshore lizardfish; **Peixe-lagarto-da-praia.**

References: Azores (PECS) (?) (ROM 23913, 1966; Carneiro *et al.*, 2014). According to Froese and Pauly (2019) and Eschmeyer *et al.* (2019) the current geographic distribution of *S. foetens* is the Western Atlantic. The occurrence record of this species off the Azores (ROM 23913) is in the PECS area (80 NM West of the Great Meteor Tablemount) and there is an inconsistency between the geographic coordinates registered (off the Azores) and the indicated locality (Atlantic Ocean off Florida). Thereby, this species was erroneously considered present in the Azores by Carneiro *et al.* (2014). Thus, this species is not considered in the total number of valid species.

***Synodus saurus* (Linnaeus 1758)**

Common name: **Atlantic lizardfish; Peixe-lagarto-do-Atlântico**, Bombô, Lagartijo-do-mar, Lagartixa-do-mar, Lagarto, Lagarto-da-costa, Lagarto-do-mar, Lagarto-de-rolo, Lagarto-do-rolo, Peixe-de-senhora, Peixe-lagarto, Pescada.

References: mainland Portugal (Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Antunes *et al.*, 1988; Carneiro *et al.*, 2014), Azores[▲] (Lowe, 1837; Collett, 1896a; Sampaio, 1904; Roule, 1919a; Vaillant, 1919; Fowler, 1936a; Collins, 1954; Albuquerque, 1954-1956; Anderson *et al.*, 1966; Osório de Castro, 1967; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a; Saldanha, 1980, 1995; Fischer *et al.*, 1981; Sulak, 1984; Patzner and Santos, 1990a, e; Barreiros, 1990, 1995; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Nash *et al.*, 1994a; Azevedo, 1995; Gonçalves *et al.*, 1995; Santos and Nash, 1995; Silva *et al.*, 1995; Menezes, 1996; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001b; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010, 2013; Diogo and Pereira, 2013a; Carneiro *et al.*, 2014; Afonso, 2016; Russell, 2016a; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837; Günther, 1864a; Vinciguerra, 1883; Goode and Bean, 1895, 1896; Fowler, 1936a, b; Norman, 1935b; Maul, 1946a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Augier, 1985; Sulak, 1984; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Russell, 2016a; Froese and Pauly, 2019).

***Synodus synodus* (Linnaeus 1758)**

Common name: **Redbarred lizardfish; Peixe-lagarto-de-rolo**, Lagarto, Lagarto-da-costa; Lagarto-de-rolo, Lagarto-do-rolo, Lagarto-do-mar.

References: Azores (Osório de Castro, 1967; Arruda, 1997; Ré, 1990; IMAR/DOP 2TER-02-P99LAG3, 1999; IMAR/DOP 6FCO-02-P13LAG3, 2013) and Madeira[▲] (NHMUK 1862.2.5.1., 1861-1890; Johnson, 1863; Günther, 1864a; Norman, 1935b; Fowler, 1936a; Maul, 1946a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nielsen, 1979b; Fischer *et al.*, 1981; INIP, 1984c; Sulak, 1984; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Russell, 2016a; Froese and Pauly, 2019). The holotype of *Saurus*

atlanticus Johnson 1863, synonym of *S. synodus*, caught off in Madeira is in the NHMUK collections (NHMUK 1862.2.5.1).

***Trachinocephalus myops* (Forster 1801)**

Common name: **Snakefish; Peixe-lagarto-de-focinho-rombo**, Aranha, Lagarto-do-mar, Lagarto-focinho-rombo, Peixe-aranha, Peixe-lagarto.

References: Azores ([MCZ 66756, 1984](#); [Carneiro et al., 2014](#)). According to Polanco *et al.* (2016) and Eschmeyer *et al.* (2019) the current geographic distribution of *T. myops* in the Eastern Atlantic is from Mauritania south to Gabon, including Saint Helena and Ascension islands. In the Museum of Comparative Zoology there is a preserved specimen of *T. myops* (catalogue number 66756) which was not included in the material examined in the study of Polanco *et al.* (2016).

Family Bathysauridae

***Bathysaurus ferox* Günther 1878**

Common name: **Deep-sea lizardfish; Peixe-lagarto-da-fundura**.

References: mainland Portugal ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Russell, 2015](#); [Froese and Pauly, 2019](#)), Azores ([Regan, 1903a, 1911](#); [Murray and Hjort, 1912 \(PECS\)](#); [Fowler, 1936a \(PECS\)](#); [Berg, 1940](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Saldanha, 1977a](#); [Sulak, 1984](#); [Lloris et al., 1991](#); [Santos et al., 1997](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Russell, 2015](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Sulak, 1984](#); [Mormede and Davies, 2003](#); [Carneiro et al., 2014](#); [Russell, 2015](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Macristium chavesi* Regan 1903, synonym of *B. ferox*, caught off in the Azores is in the NHMUK collections (apparently lost).

***Bathysaurus mollis* Günther 1878**

Common name: **Highfin lizardfish; Peixe-lagarto-da-costa**, Lagarto-da-costa, Peixe-lagarto.

References: mainland Portugal ([NHMUK 1994.9.20.4, 1983](#); [Carneiro et al., 2014](#)), Azores ([Roule, 1919a \(PECS\)](#); [Fowler, 1936a \(PECS\)](#); [Saldanha, 1977a, 1991](#); [MNHN-IC-1978-0180, 1971](#); [Hureau and Monod, 1979](#); [Lloris et al., 1991](#); [Pérès, 1992](#); [Arruda, 1997](#); [Santos et al., 1997](#); [MAR-ECO 6207, 2004](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Almeida and Biscoito, 2019](#)) and Madeira ([Hureau and Monod, 1979](#); [NHMUK 2003.6.11.17, 1989](#); [Sulak, 1990b](#); [Lloris et al., 1991](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Family Paralepididae

***Arctozenus risso* (Bonaparte 1840)**

Common name: **Spotted barracudina; Barracudina-pintada**.

References: mainland Portugal ([YPM ICH 005128, 1959](#); [ZMH 106009, 1968](#); [Post, 1984](#); [MCZ 159552, 1986](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#)), Azores ([Ege, 1930, 1953](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Rofen, 1966a](#); [Post, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Maul, 1945](#); [Wolff, 1971](#); [MCZ 67586, 1972](#); [MCZ 67732, 1972](#); [Post, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Lestidiops affinis* (Ege 1930)**

Common name: Barracudina; **Barracudina-ruça**, Barracudina.

References: Azores ([Ege, 1930](#); [Albuquerque, 1954-1956](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Ege, 1930](#); [Albuquerque, 1954-1956](#); [Maul, 1971a](#); [Post, 1984](#); [Carneiro et al., 2014](#)).

***Lestidiops jayakari* (Boulenger 1889)**

Common name: Pacific barracudina; **Barracudina-do-Pacífico**, Barracudina.

References: mainland Portugal ([Post, 1984](#); [Fischer et al., 1987](#)), Azores ([Ege, 1930, 1953](#); [Albuquerque, 1954-1956](#); [Rofen, 1966a](#); [Post, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Diekmann et al., 2006 \(PECS\)](#); [Arkhipov and Mamedov, 2008 \(PECS\)](#); [Sutton et al., 2008](#)) and Madeira ([Ege, 1930](#); [Maul, 1945, 1952b](#); [ZMH 104260, 1966](#); [MCZ 68103, 1970](#); [Post, 1984](#); [Lloris et al.,](#)

1991). The holotype (unique) of *Macroparalepis egei* Maul 1945, synonym of *L. jayakari*, caught off in Madeira (stomach content) is in the MMF collections (MMF 3003).

***Lestidiops pseudosphyraenoides* (Ege 1918)**

Common name: No English common name; **Barracudina-do-Atlântico-nordeste**, Barracudina. References: mainland Portugal (Carneiro et al., 2014; Froese and Pauly, 2019), Azores (Vaillant, 1919; MCZ 68073, 1964; MCZ 68074, 1964; MCZ 68075, 1964; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Johnson, 1862c; Günther, 1864a; Roule, 1919a; Vaillant, 1919; MCZ 68353, 1969; Carneiro et al., 2014).

***Lestidiops similis* (Ege 1933)**

Common name: No English common name; **Barracudina-análoga**, Barracudina. References: Azores (ZMH 114053, 1975; Carneiro et al., 2014; Carpenter, 2015).

***Lestidiops sphyrenoides* (Risso 1820)**

Common name: Barracudina; **Barracudina-do-Mediterrâneo**, Barracudina. References: mainland Portugal (Post, 1984; Fischer et al., 1987; Carneiro et al., 2014; Froese and Pauly, 2019), Azores (Ege, 1930, 1953; Fowler, 1936a; Albuquerque, 1954-1956; Post, 1984; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Günther, 1892; Goode and Bean, 1895, 1896; Brauer, 1906; Ege, 1930; Fowler, 1936a; Albuquerque, 1954-1956; Post, 1984; Carneiro et al., 2014; Porteiro et al., 2017; Froese and Pauly, 2019).

***Lestidium atlanticum* Borodin 1928**

Common name: **Atlantic barracudina**; **Barracudina-do-Atlântico**. References: mainland Portugal (Bannermann et al., 2015b), Azores (Arkipov and Mamedov, 2008 (PECS); Bannermann et al., 2015b) and Madeira (Bannermann et al., 2015b). Bannermann et al. (2015b) include the mainland Portugal EEZ in the distribution area of *L. atlanticum*.

***Macroparalepis affinis* Ege 1933**

Common name: Barracudina; **Barracudina-das-Canárias**, Barracudina. References: mainland Portugal (Post, 1984; Whitehead et al., 1984), Azores (Ege, 1957; ZMH 119544, 1979 (PECS); Bekker et al., 1982; Post, 1984; Arruda, 1997; Santos et al., 1997; Porteiro et al., 2010; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Maul, 1945; ZMH 104329, 1966; Post, 1984; Carneiro et al., 2014).

***Macroparalepis brevis* Ege 1933**

Common name: No English common name; **Barracudina-pequena**. References: Azores (Bekker et al., 1982; SOC Discovery 1126129_84_FIS_510002, 1985 (PECS); SOC Discovery 1126130_85_FIS_510002, 1985 (PECS); Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Post, 1984).

***Macroparalepis nigra* (Maul 1965)**

Common name: No English common name; **Barracudina-negra**. References: Madeira (Maul, 1965; Post, 1984; Lloris et al., 1991; Carneiro et al., 2014; Delgado et al., 2017; Froese and Pauly, 2019). The holotype (unique) of *Longisudis nigra* Maul 1965, synonym of *M. nigra*, caught off in Madeira (stomach content) is in the MMF collections (MMF 21171).

***Magnisudis atlantica* (Krøyer 1868)**

Common name: **Duckbill barracudina**; **Barracudina-bico-de-pato**, Barracudina. References: mainland Portugal (MOM-IC-0091-1162, 1910; Zugmayer, 1911a, b; Fowler, 1936a; MCZ 68500, 1972; Post, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores (Ege, 1930; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Diekmann et al., 2006 (PECS); Sutton et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Maul, 1945; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Post, 1984; Carneiro et al., 2014). The holotype (unique) of *Paralepis brevis* Zugmayer 1911, synonym of *M. atlantica*, caught off in mainland Portugal is in the MOM collections (MOM-IC-0091-1162).

***Paralepis brevirostris* (Parr 1928)**

Common name: No English common name; **Barracudina-de-focinho-curto**, Barracudina.

References: Azores ([MAR-ECO 3479, 2004](#); [MAR-ECO 14768, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([Maul, 1945, 1954b, 1955c](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [ZMH 105846, 1968](#); [Post, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)). The holotype (unique) of *Paralepis harryi* Maul 1954, synonym of *P. brevirostris*, is in the MMF collections (MMF 3863). It was obtained in the Funchal fish market, from the stomach of an black scabbardfish caught off in Madeira.

***Paralepis coregonoides* Risso 1820**

Common name: Lancet fish; **Barracudina-lanceta**, Barracudina, Fura-pão; Fura-pás, Peixe-grato.

References: mainland Portugal ([Nobre, 1903](#); [Seabra, 1911](#); [IFREMER Biocean 34521, 1969](#); [Post, 1984](#); [ZSM-Pis-029132 / 534342 / 385930, 1992](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)), Azores ([Ege, 1930, 1932, 1953](#); [Roule and Angel, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Haedrich, 1964](#); [Rofen, 1966a](#); [Post, 1984, 1987b](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Johnson, 1862b, c](#); [Ege, 1930](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936b](#); [Maul, 1945](#); [Albuquerque, 1954-1956](#); [MCZ 68745, 1969](#); [Post, 1984](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)).

***Paralepis elongata* (Brauer 1906)**

Common name: Barracudina; **Barracudina-translúcida**, Barracudina.

References: Azores ([PECS](#)) ([SOC Discovery 1023306_86_FIS_647004, 1980](#); [Carneiro et al., 2014](#); [Moore et al., 2015](#)) and Madeira ([Moore et al., 2015a](#)). According to Eschmeyer et al. (2019) *P. elongata* is a cosmopolitan species. The distribution map of this species reported by Moore et al. (2015a) includes the Madeira Islands area. Specimens are needed for its confirmation in Madeira.

***Paralepis speciosa* Bellotti 1878**

Common name: No English common name; **Barracudina-manchada**, Barracudina.

References: mainland Portugal ([ZSM-Pis-029372 / 534568 / 386156, 1990](#); [ZSM-Pis-029151 / 534361 / 385949, 1992](#)), Azores ([MCZ 128199, 1969](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([MCZ 128198, 1969](#); [ZSM-Pis-029349 / 534547 / 386135, 1990](#); [Carneiro et al., 2014](#)).

***Sudis atrox* Rofen 1963**

Common name: Fierce pike smelt; **Barracudina-feroz**, Barracudina.

References: Azores ([MCZ 67361, 1973 \(PECS\)](#); [MCZ 67382, 1973 \(PECS\)](#); [Carneiro et al., 2014 \(PECS\)](#); [Carpenter et al., 2015a](#)) and Madeira ([Carpenter et al., 2015a](#)). Carpenter et al. (2015a) include the Azorean and Madeiran EEZs in the distribution area of *S. atrox*.

***Sudis hyalina* Rafinesque 1810**

Common name: Barracudina; **Barracudina-imperial**, Barracudina.

References: mainland Portugal ([YPM ICH 005154, 1959](#); [YPM ICH 005155, 1959](#); [MCZ 68302, 1972](#); [Carneiro et al., 2014](#)), Azores ([MCZ 68570, 1964](#); [Shores 1969](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Arkhipov and Mamedov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Maul, 1945](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [MCZ 130971, 1978](#); [Post, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Uncisudis longirostra* Maul 1956**

Common name: No English common name; **Barracudina-da-Madeira**, Barracudina.

References: Madeira ([Maul, 1956b](#); [Post, 1984, 1990](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype of this species caught off in Madeira (stomach content) is in the MMF collections (MMF 5722).

***Uncisudis quadrimaculata* (Post 1969)**

Common name: Barracudina; **Barracudina-de-quatro-manchas**, Barracudina.

References: Azores ([MCZ 68562, 1964](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#)).

Family Anotopteridae

Anotopterus pharae Zugmayer 1911

Common name: **Daggetooth; Barracudina-dente-de-adaga.**

References: mainland Portugal ([MOM-IC-0091-1154](#), 1910; Zugmayer, 1911a; [YPM ICH 004511](#), 1959; [Maul](#) 1979c; Post, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([MCZ 73018](#), 1969; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014) and Madeira (Maul 1946a, 1952b, 1971a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; ZMH 110492, 1971; Post, 1984; Kukuev, 1998; Carneiro et al., 2014; Froese and Pauly, 2019). The holotype (unique) of this species caught off at 36 NM SW of the Gorringe Ridge, 36°30.0'N, 11°49.0'W (off mainland Portugal) is in the MOM collections ([MOM-IC-0091-1154](#)).

Family Evermannellidae

Coccarella atlantica (Parr 1928)

Common name: Atlantic sabretooth; **Dente-de-sabre-do-Atlântico.**

References: Azores ([Roule and Angel](#), 1930; [Fowler](#), 1936b; Albuquerque, 1954-1956; [MNHN-IC-2009-0405](#), 1971; Maul, 1979b; Johnson, 1984; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 2010; Carneiro et al., 2014; Almeida and Biscoito, 2019) and Madeira ([MOM-IC-0091-1733](#), 1926; [Roule](#), 1929; [Fowler](#), 1936b; Albuquerque, 1954-1956; ZMH 104355, 1966; Maul, 1979b; Johnson, 1984; Lloris et al., 1991; Carneiro et al., 2014). The records of the subspecies *Coccarella atrata atlantica* ([Roule](#) 1929) was raised to species *C. atlantica*. The holotype of *Coccarella atrata atlantica* [Roule](#) 1929 caught off Câmara de Lobos (Madeira) is in the MOM collections ([MOM-IC-0091-1733](#)).

Evermannella balbo (Risso 1820)

Common name: **Balbo sabretooth; Dente-de-sabre-de-Balbo.**

References: mainland Portugal ([SOC Discovery 1104202_85_FIS_263001](#), 1984 (PECS); [Johnson](#), 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([Schmidt](#), 1918; [Roule and Angel](#), 1930; [Fowler](#), 1936a; [Rofen](#), 1966b; [Arruda](#), 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([Schmidt](#) 1918; [Fowler](#), 1936a; [Maul](#), 1946a, 1979b; [Noronha and Sarmento](#), 1948; Albuquerque, 1954-1956; MCZ 128165, 1978; [Johnson](#), 1984; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019).

Evermannella melanoderma Parr 1928

Common name: Indian sabertooth; **Dente-de-sabre-ponteado.**

References: mainland Portugal ([NHMUK 2009.6.24.9](#), 1958 (PECS); [Johnson](#), 1982; Carneiro et al., 2014) and Madeira ([ZMH 104333](#), 1966; [MCZ 128175](#), 1978; [Johnson](#), 1982; [Swinney](#), 1994; Carneiro et al., 2014; Froese and Pauly, 2019). The records of the subspecies *Evermannella indica melanoderma*, [Parr](#) 1928 was raised to species *Evermannella indica* ([Johnson](#) 1982). According to [Swinney](#) (1994) and [Russell](#) (2016b) records of *E. indica* from the Atlantic are now known to be *Evermannella melanoderma*.

Odontostomops normalops (Parr 1928)

Common name: Undistinguished sabretooth; **Dente-de-sabre-olho-redondo.**

References: Madeira ([NHMUK 2003.9.16.94](#), 1958; [Nunoo](#) et al., 2015).

Family Omosudidae

Omosudis lowii Günther 1887

Common name: **Omosudid; Peixe-mandíbula-de-martelo.**

References: mainland Portugal ([YPM ICH 005138](#), 1959; [Maul](#), 1976; Post, 1984; Carneiro et al., 2014), Azores ([Roule and Angel](#), 1930, 1933; [Fowler](#), 1936b; Albuquerque, 1954-1956; [Ege](#), 1958; [Rofen](#), 1966b; Post, 1984; [Arruda](#), 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([NHMUK 1868.5.13.6](#), 1864-1877; [Günther](#), 1887; [Brauer](#), 1906; [Murray and Hjort](#), 1912; [MOM-IC-0091-1727](#), 1926; [Roule](#), 1929; [Fowler](#), 1936a, b; [Maul](#), 1946a, 1979b; [Noronha and Sarmento](#), 1948; Albuquerque, 1954-1956; Post, 1984; Lloris et al., 1991; Carneiro et al., 2014; Porteiro et al., 2017; Roberts et al., 2017; Froese and Pauly, 2019). The holotype of the subspecies *Omosudis lowei funchali* [Roule](#) 1929 caught off Funchal (Madeira) is in the

MOM collections (MOM-IC-0091-1727). It was raised to synonym of species *O. lowii*. One syntype of this species caught off Magdalena (Madeira) is in NHMUK collections (NHMUK 1868.5.13.6). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22496, 1967).

Family Alepisauridae

Alepisaurus brevirostris Gibbs 1960

Common name: **Short snouted lancetfish; Lírio-ferro-de-focinho-curto**, Peixe-cavalo.

References: Azores (USNM 200817, 1963; Haedrich, 1964; Santos *et al.*, 1997; Melo and Menezes, 2002; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Menezes *et al.*, 2012; Carneiro *et al.*, 2014).

Alepisaurus ferox Lowe 1833

Common name: **Long snouted lancetfish; Lírio-ferro-de-focinho-comprido**, Água, Bicuda-da-índia, Cavalo, Golpim, Lirão, Lírio, Lírio-ferro, Liro, Peixe-água, Peixe-cavalo.

References: mainland Portugal (Steindachner, 1868b; de Brito Capello, 1867d, 1880; Osório, 1888; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Post, 1984; Fischer *et al.*, 1987; Bordalo-Machado *et al.*, 2009; de Casamajor and Morandeau, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Haedrich, 1964; Osório de Castro, 1967; Parin, 1970; Post, 1984; Lloris *et al.*, 1991; Pinho and Pereira, 1995a; Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Arkhipov & Mamedov, 2008 (PECS); Porteiro *et al.*, 2010, 2017(?); Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Lowe, 1833b, 1835a, b, 1837; Cuvier and Valenciennes, 1849; NHMUK 1852.9.13.98, 1850-1853; Johnson, 1862c; Günther, 1864a; Collett, 1890b; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1946a, 1979b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Post, 1984; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; de Casamajor and Morandeau, 2013; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019). One syntype (?) (former holotype BMNH 1852.9.13.98) of this species caught off in Madeira is in the NHMUK collections (NHMUK 1852.9.13.98).

Order Myctophiformes

Family Neoscopelidae

Neoscopelus macrolepidotus Johnson 1863

Common name: **Large-scaled lantern fish; Peixe-lanterna-escamudo**, Lanterna-vermelha.

References: Azores (Kukuev, 2002 (PECS); Melo and Menezes, 2002; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira (NHMUK 1862.10.3.3, 1861-1890; Johnson, 1863; Goode and Bean, 1895, 1896; Brauer, 1906; Fowler, 1936a; Maul, 1946a, 1951b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Nielsen, 1979c; Hulley, 1984, 1990a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (?) of this species caught off in Madeira is in the NHMUK collections (NHMUK 1862.10.3.3).

Neoscopelus microchir Matsubara 1943

Common name: Shortfin neoscopelid; **Peixe-lanterna-de-barbatanas-curtas**.

References: mainland Portugal (Moura, 2004; Carneiro *et al.*, 2014) and Madeira (Maul, 1951b; Nielsen, 1979c; Hulley, 1984, 1990; Lloris *et al.*, 1991; Bañón *et al.*, 2002; Carneiro *et al.*, 2014; Delgado *et al.*, 2017; Froese and Pauly, 2019).

Scopelengys tristis Alcock 1890

Common name: **Pacific blackchin; Peixe-lanterna-do-Pacífico**.

References: Madeira (Maurin *et al.*, 1977; Nielsen, 1979c; Hulley, 1984, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Family Myctophidae

Benthosema glaciale (Reinhardt 1837)

Common name: **Glacier lantern fish; Peixe-lanterna-glacial**, Divertido, Escolarinho, Ferrreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (GNM 1946-7959, 1946; Osório de Castro, 1967; Hulley, 1984; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Bolin,

1959; Osório de Castro, 1967(?); Halliday, 1970; Geistdoerfer *et al.*, 1971; Nafpaktitis *et al.*, 1977; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (MCZ 126034, 1969; MCZ 126035, 1969; MCZ 126179, 1978; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Mecklenburg *et al.*, 2018).

***Benthosema suborbitale* (Gilbert 1913)**

Common name: **Smallfin lanternfish**; **Peixe-lanterna-corcunda**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (MCZ 125367, 1969; Krefft and Bekker, 1979; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016), Azores (Tåning, 1932b; Albuquerque, 1954-1956; Bolin, 1959; Osório de Castro, 1967; Nafpaktitis *et al.*, 1977; Bekker *et al.*, 1982; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Arkhipov & Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Tåning, 1932b; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984, 1990b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Bolinichthys indicus* (Nafpaktitis & Nafpaktitis 1969)**

Common name: Smoothcheek lanternfish; **Peixe-lanterna-do-Índico**, Romeirinho, Divertido, Escolarinho, Lobisomem.

References: mainland Portugal (NHMUK 2003.4.11.197-200, 1959; ZMH 117979, 1979; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019), Azores (Bolin, 1959; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (MCZ 124433, 1969; MCZ 124308, 1970; Maul, 1971a(?); Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Bolinichthys longipes* (Brauer 1906)**

Common name: Popeye lampfish; **Peixe-lanterna-barbatana-comprida**.

References: Madeira (?) (Brauer, 1906; Fowler, 1936a; Carneiro *et al.*, 2014; Roberts *et al.*, 2017). According to Eschmeyer *et al.* (2019) the current geographic distribution of *B. longipes* is the warm water areas of the Indo-Pacific: South Africa east to Hawaiian Islands, north to southern Japan, south to northern Australia and New Zealand EEZ; Eastern Pacific from off California to off Argentina/Chile. According to Halley and Duhamel (2009), the type specimen caught near Madeira (*Myctophum longipes* Brauer 1906, accepted as *B. longipes*) is apparently referable to *B. indicus*. The occurrence of this species in Madeira is very questionable and is not considered in the total number of valid species.

***Bolinichthys photothorax* (Parr 1928)**

Common name: Spurcheek lanternfish; **Peixe-lanterna-esporão**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: Madeira (Hulley and Duhamel, 2009; Carneiro *et al.*, 2014; Hulley and Paxton, 2016).

***Bolinichthys pyrsobolus* (Alcock 1890)**

Common name: Fiery lanternfish; **Peixe-lanterna-arco-íris**.

References: mainland Portugal (?) (Carneiro *et al.*, 2014). There are twelve specimens caught off Portugal in the Vertebrate Zoology Division - Ichthyology of the Yale Peabody Museum of Natural History cataloged as *B. pyrsobolus* (Alcock 1890). According to Hulley and Duhamel (2009) there has been considerable confusion in the identity of specimens in museum collections due to the inadequacy of the type description by Alcock (1890), coupled with the lumping of all Atlantic species of the genus under the name *Lepidophanes* (= *Bolinichthys*) *pyrsobolus* by Bolin (1959). On the other hand, these authors and Eschmeyer *et al.* (2019) reported the current geographic distribution of this species in the Indo-West Pacific: Indian Ocean to central and southern Pacific, north to Japan, south to Australia. Thus, the occurrence of this species in mainland Portugal reported by Carneiro *et al.* (2014) is probably erroneous and it is not considered in the total number of valid species.

***Bolinichthys supralateralis* (Parr 1928)**

Common name: Stubby lanternfish; **Peixe-lanterna-atarracado**.

References: mainland Portugal (PECS) ([NHMUK 2006.6.8.40, 1984](#)), Azores ([MAR-ECO 3444, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([MCZ 123791, 1969](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)).

Centrobranchus andreae (Lütken 1892)

Common name: Andre's lanternfish; [Peixe-lanterna-de-Andréas](#).

References: mainland Portugal (?) ([Brauer, 1906](#); [Nobre, 1935](#); [Krefft and Bekker, 1979](#)) and Azores (?) ([Lütken, 1892](#); [Brauer, 1906](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is nearly circumglobal in the tropical areas of the Indo-Pacific. Lütken (1892) indicates the capture of some specimens of *C. andreae* in the Atlantic, some of which in the Azores region. Krefft and Bekker (1979) mention that Lütken type locality appear most doubtful, referring to specimens caught off Portugal coast, without indicating any geographic coordinates. According to Yang *et al.* (1996) the current geographic distribution of *C. andreae* is Indo-Pacific and the South China Sea. Thus, these specimens could have been misidentified (?) and according to the current geographic distribution this species is not considered in the total number of valid species.

Centrobranchus choerocephalus Fowler 1904

Common name: No English common name; [Peixe-lanterna-de-focinho-comprido](#).

References: Azores (PECS) ([Murray and Hjort, 1912](#)). According to Eschmeyer *et al.* (2019) *C. choerocephalus* is a cosmopolitan species.

Centrobranchus nigrocellatus (Günther 1873)

Common name: Roundnose lanternfish; [Peixe-lanterna-de-focinho-redondo](#), Ferreiro, Peixinho-preto, Romeirinho.

References: Azores ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [MCZ 123282, 1973 \(PECS\)](#); [Arkipov & Mamedov, 2008 \(PECS\)](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)) and Madeira ([MCZ 123206, 1969](#); [MCZ 123208, 1969](#); [Hulley, 1984, 1990b](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

Ceratoscopelus maderensis (Lowe 1839)

Common name: Madeira lantern fish; [Peixe-lanterna-da-Madeira](#), Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([Brauer, 1906](#); [Nobre, 1935](#); [ZMH 103490, 1966](#); [Osório de Castro, 1967](#); [Martins, 1981](#); [ZMH 105679, 1968](#); [Hulley, 1984](#); [ZSM-Pis-029231 / 534434 / 386022, 1992](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Collet 1896](#); [Richard, 1905a](#); [Brauer, 1906](#); [Zugmayer, 1911b](#); [Roule, 1919a](#); [Vaillant, 1919](#); [Roule and Angel, 1930](#); [Nobre, 1935](#); [Fowler, 1936a,b](#); [Belloc, 1938](#); [Ferreira, 1942](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Backus et al., 1970](#); [Geistdoerfer et al., 1971](#); [Dias et al., 1976](#); [Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Hulley, 1984](#); [Chicharo and Teodósio, 1990](#); [Monteiro, 1996](#); [Monteiro et al., 1996b](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1839, 1840b, 1844a, 1849, 1850, 1852](#); [Jonhson, 1862c](#); [Günther, 1864a](#); [NHMUK 1865.1.19.2, 1864-1877](#); [Good and Bean, 1895](#); [Brauer, 1906](#); [Nobre, 1935](#); [Fowler, 1936a, b](#); [Maul, 1946a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft and Bekker, 1979](#); [Hulley, 1984](#); [Lloris et al., 1991](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Scopelus maderensis* Lowe 1839, synonym of *C. maderensis*, caught off in Madeira is in the NHMUK collections (NHMUK 1865.1.19.2).

Ceratoscopelus townsendi (Eigenmann & Eigenmann 1889)

Common name: Dogtooth lampfish; [Peixe-lanterna-de-Townsend](#), Ferreiro, Peixinho-preto, Romeirinho.

References: Azores ([Osório de Castro, 1967](#)) and Madeira ([NHMUK 2004.3.10.49-50, 1965](#); [Osório de Castro, 1967](#)). According to Eschmeyer *et al.* (2019) *C. townsendi* is a circumglobal species in tropical through cold temperate seas.

Ceratoscopelus warmingii (Lütken 1892)

Common name: Warming's lanternfish; [Peixe-lanterna-de-Warming](#), Divertido, Escolarinho, Lobisomem. Romeirinho.

References: mainland Portugal ([NHMUK 2004.3.8.504-505, 1966; MCZ 122224, 1969; Carneiro et al., 2014](#)), Azores ([Lütken, 1892\(?\)](#); [Brauer, 1906](#); [MCZ 121755, 1964](#); [Nafpaktitis et al., 1977](#); [Krefft and Bekker, 1979](#); [Bekker, 1983](#); [Lloris et al., 1991](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Arkhipov & Mamedov, 2008 \(PECS\)](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Roberts et al., 2017](#)) and Madeira ([Maul, 1946a, 1971a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Krefft and Bekker, 1979](#); [Hulley, 1984, 1990b](#); [Lloris et al., 1991](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Scopelus warmingii* (*Nyctophus*) Lütken 1892, synonym of *C. warmingii*, caught off SW of Azores, north-eastern Atlantic at 32°6.0'N, 39°28.0'W is in the ZMUC collections (ZMUC 39). Several authors ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#)) mention the occurrence of this species in the Azores based on Lütken (1892). These citations referred to the same specimen (ZMUC 39), and the geographic coordinates correspond to a position off the Azores, out of the PECS area. The first records of this species in the Azores were made by Richard H. Backus in 1969 (RV Atlantis II, research survey 049).

***Ctenoscopelus phengodes* (Lütken 1892)**

Common name: Bright lanternfish; **Peixe-lanterna-brilhante**.

References: Azores (?) ([Zugmayer, 1911b](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft and Bekker, 1979 \(PECS\)](#); [Lloris et al., 1991](#)). Based on morphological characters and phylogenetic analyses the species *Myctophum phengodes* (Lütken 1892) was relocated in the genus *Ctenoscopelus* ([Martin et al., 2018](#)). A single specimen of this species was found from north-east of the Azores ([Krefft and Bekker, 1979](#)). However, according to [Eschmeyer et al. \(2019\)](#) and [Froese and Pauly \(2019\)](#) *C. phengodes* is a southern circumglobal species. Thus, this species is not considered in the total number of valid species of the Azores.

***Dasyscopelus selenops* (Tåning 1928)**

Common name: **Wisner's lantern fish**; **Peixe-lanterna-de-Wisner**, Romeirinho.

References: Azores (PECS) ([ZMH 118315, 1979](#); [Arkhipov & Mamedov, 2008 \(PECS\)](#); [Carneiro et al., 2014](#)) and Madeira ([Maul, 1952b](#); [ZMH 104128, 1966](#); [MCZ 105328, 1969](#); [Hulley, 1984](#); [MCZ 105334, 1970](#); [Lloris et al., 1991](#); [Hulley and Paxton, 2016](#)). Based on morphological characters and phylogenetic analyses, the species *Myctophum selenops* Tåning 1928 was relocated in the genus *Dasyscopelus* ([Martin et al., 2018](#)).

***Diaphus adenomus* Gilbert 1905**

Common name: Firebrow lanternfish; **Peixe-lanterna-de-Gilbert**.

References: Madeira (?) ([Hulley, 1984](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The current geographic distribution of this species is the Atlantic and the Pacific, including Hawaiian Islands ([Eschmeyer et al., 2019](#)). [Carneiro et al. \(2014\)](#) and [Froese and Pauly \(2019\)](#) consider this species as native of Madeira based on [Hulley \(1984\)](#), but this author does not mention the occurrence of this species in this area. Thus, this species is not considered in the total number of valid species.

***Diaphus bertelseni* Nafpaktitis 1966**

Common name: Bertelsen's lanternfish; **Peixe-lanterna-de-Bertelsen**, Romeirinho.

References: mainland Portugal ([Krefft and Bekker, 1979](#); [Hulley, 1984](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)), Azores ([MAR-ECO 5313, 2004 \(PECS\)](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Porteiro et al., 2017](#)) and Madeira ([ZMH 105494, 19686](#); [Krefft and Bekker, 1979](#); [Hulley, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)).

***Diaphus brachycephalus* Tåning 1928**

Common name: **Short-headed lantern fish**; **Peixe-lanterna-de-cabeça-curta**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: mainland Portugal ([YPM ICH 004901, 1959](#); [YPM ICH 004902, 1959](#); [YPM ICH 004903, 1959](#); [Carneiro et al., 2014](#)), Azores ([ZMH 117673, 1979 \(PECS\)](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Hulley, 2015f](#)) and Madeira ([ZMH 103409, 1966](#); [ZMH 105342, 1968](#); [Krefft and Bekker, 1979](#); [Hulley, 1984, 2015d](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Diaphus dumerilii* (Bleeker 1856)**

Common name: Dumeril's lanternfish; **Peixe-lanterna-de-Duméril**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([Costa, 2004](#); [Carneiro et al., 2014](#)), Azores ([MCZ 121135, 1964](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Johnson, 1890](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft and Bekker, 1979](#); [Hulley, 1990b](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Diaphus effulgens* (Goode & Bean 1896)**

Common name: Headlight lanternfish; **Peixe-lanterna-de-focinho-luminoso**, Divertido, Escolarinho, Lobisomem.

References: mainland Portugal ([ZSM-Pis-028806 / 534025 / 385613, 1991](#)), Azores ([Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)) and Madeira ([Maul, 1952b](#); [MCZ 110113, 1969](#); [Krefft and Bekker, 1979](#); [Hulley, 1984, 1990b](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Diaphus holti* Tåning 1918**

Common name: Small lantern fish; **Peixe-lanterna-de-Holt**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: mainland Portugal ([ZMH 103388, 1966](#); [Krefft and Bekker, 1979](#); [Hulley, 1984](#); [Fischer et al., 1987](#); [ZSM-Pis-029189 / 534397 / 385985, 1992](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Bolin, 1959](#); [Nafpaktitis, 1968](#); [Backus et al., 1970](#); [Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Hulley, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([MCZ 120501, 1969](#); [MCZ 120506, 1970](#); [Hulley, 1984, 1990b](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Diaphus lucidus* (Goode & Bean 1896)**

Common name: Spotlight lanternfish; **Peixe-lanterna-cintilante**, Romeirinho.

References: Azores ([Carneiro et al., 2014](#) (PECS); [Hulley and Paxton, 2016](#)) and Madeira ([ZMH 103638, 1966](#); [ZMH 103592, 1966](#); [Krefft and Bekker, 1979](#); [Hulley, 1984, 1990b](#); [Lloris et al., 1991](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Diaphus luetkeni* (Brauer 1904)**

Common name: Luetken's lanternfish; **Peixe-lanterna-de-Lüetken**.

References: Azores ([PECS](#)) ([SOC Discovery 740614_8_FIS_214005, 1970](#); [Carneiro et al., 2014](#)) and Madeira ([MCZ 120148, 1969](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)).

***Diaphus metopoclampus* (Cocco 1829)**

Common name: Spothead lantern fish; **Peixe-lanterna-de-cabeça-manchada**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([ZMH 105630, 1968](#) (PECS); [Hulley, 1984](#); [Fischer et al., 1987](#); [Saldanha et al., 1995](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Papaconstantinou et al., 2015](#); [Froese and Pauly, 2019](#)), Azores ([Zugmayer, 1933](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Nafpaktitis, 1968](#); [Dias et al., 1976](#); [Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Hulley, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Papaconstantinou et al., 2015](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Osório de Castro, 1967\(?\)](#); [MCZ 120047, 1969](#); [MCZ 120048, 1969](#); [Hulley, 1984](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Papaconstantinou et al., 2015](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Diaphus mollis* Tåning 1928**

Common name: Soft lanternfish; **Peixe-lanterna-mole**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: mainland Portugal ([ZSM-Pis-028814 / 534033 / 385621, 1991](#); [ZSM-Pis-028815 / 534034 / 385622, 1991](#); [ZSM-Pis-028816 / 534035 / 385623, 1991](#)), Azores ([Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Hulley, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)) and Madeira ([ZMH 103630, 1966](#); [MCZ 119267, 1969](#)).

Krefft and Bekker, 1979; Hulley, 1984, 1990b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Diaphus perspicillatus* (Ogilby 1898)**

Common name: **Flatface lanternfish; Peixe-lanterna-luneta.**

References: Azores (MCZ 126609, 1964 (PECS); Carneiro *et al.*, 2014 (PECS); Ralph, 2015g). According to Eschmeyer *et al.* (2019) *D. perspicillatus* is a circumglobal species in tropical through temperate seas, but not in the Eastern Pacific. The distribution map of this species reported by Ralph (2015g) partially includes the Azores. Specimens are needed for confirmation in the Azores EEZ.

***Diaphus rafinesquii* (Cocco 1838)**

Common name: White-spotted lantern fish; **Peixe-lanterna-de-Rafinesque**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; NHMUK 1989.1.25.81, 1955; Osório de Castro, 1967; Hulley, 1984; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Murray and Hjort, 1912; Zugmayer, 1933; Fowler, 1936a, b; Albuquerque, 1954-1956; Bolin, 1959; Osório de Castro, 1967; Nafpaktitis, 1968; Geistdoerfer *et al.*, 1971; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (ZMH 104370, 1966; Osório de Castro, 1967; MCZ 102028, 1978; Hulley, 1984; Davies and Merrett, 1998; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Diaphus splendidus* (Brauer 1904)**

Common name: **Horned lanternfish; Peixe-lanterna-cornudo.**

References: Azores (PECS) (ZMH 119041, 1979; Carneiro *et al.*, 2014) and Madeira (Hulley and Paxton, 2016). According to Eschmeyer *et al.* (2019) *D. splendidus* is a circumglobal species in tropical seas. Hulley and Paxton (2016) indicate the distribution of this species in the area between 30°N and 23°S and Froese and Pauly (2019) referred the area between 30° and 19° N. The northernmost distribution limit reported by these authors includes the area of the Selvagens Islands. However, specimens are needed for confirmation in the Madeira Archipelago.

***Diaphus subtilis* Nafpaktitis 1968**

Common name: Flabby lanternfish; **Peixe-lanterna-flácido**, Romeirinho.

References: Azores (PECS) (SOC Discovery 827000_8_FIS_214010, 1973; SOC Discovery 1023204_84_FIS_214010, 1980; SOC Discovery 1023202_85_FIS_214010, 1980; Carneiro *et al.*, 2014) and Madeira (Krefft and Bekker, 1979; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Diaphus termophilus* Tåning 1928**

Common name: **Taaning's lantern fish; Peixe-lanterna-termófilo.**

References: mainland Portugal (PECS) (ZMH 105789, 1968) and Madeira (Krefft and Bekker, 1979; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016(?); Froese and Pauly, 2019).

***Diaphus theta* Eigenmann & Eigenmann 1890**

Common name: California headlightfish; **Peixe-lanterna-da-California.**

References: Madeira (?) (Vaillant, 1919; Fowler, 1936a). Vaillant (1919) indicates the occurrence of this species off Madeira, 30°27.0'N, 25°12.0'W, at surface. This geographic coordinates correspond to a position in the Madeira Abyssal Plain, but out of the PECS area. According to Eschmeyer *et al.* (2019) the current geographic distribution of *D. theta* is the North Pacific subarctic transition zone and California Current (north-eastern Pacific), waifs reaching Hawaiian Islands. Thus, this specimen could have been misidentified and according to the current geographic distribution of this species it is not considered in the total number of valid species.

***Diogenichthys atlanticus* (Tåning 1928)**

Common name: Atlantic lanternfish; **Peixe-lanterna-do-Atlântico**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (YPM ICH 004907, 1959; MCZ 117811, 1969; Hulley, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Schmidt, 1918; Bolin, 1959; Nafpaktitis *et al.*, 1977; Bekker *et al.*,

1982; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Diekmann *et al.*, 2006 (PECS); Arkhipov & Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967; MCZ 117842, 1970; Bekker *et al.*, 1982; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Diogenichthys laternatus* (Garman 1899)**

Common name: Diogenes lanternfish; **Peixe-lanterna-de-Diogenes**.

References: Azores (?) (Schmidt, 1918(?); Fowler, 1936a(?); Osório de Castro, 1967) and Madeira (?) (Osório de Castro, 1967(?)). Schmidt (1918) indicates the occurrence of this species in the Atlantic, in the Bay of Cadiz, off Azores and off Grand Banks, in 220-1600 m. Osório de Castro (1967) mentioned the occurrence of *D. laternatus* in Madeira as doubtful. Since Osório de Castro (1967) no other author described the presence of this species in Madeira and its occurrence in this EEZ was not yet confirmed by more recent data. According to Eschmeyer *et al.* (2019) the current geographic distribution of *D. laternatus* is the central and Eastern Pacific: California to Chile, west to Hawaiian Islands and equator at 170°W. Thus, this species is not considered in the total number of valid species.

***Electrona risso* (Cocco 1829)**

Common name: **Electric lantern fish**; **Peixe-lanterna-faísca**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (Krefft and Bekker, 1979; Hulley, 1984; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Brauer, 1906; Murray and Hjort, 1912; Zugmayer, 1933; Fowler, 1936b; Ferreira, 1942; Collins, 1954; Bolin, 1959; Osório de Castro, 1967; Backus *et al.*, 1970; Dias *et al.*, 1976; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Topal, 1987; Monteiro, 1996; Monteiro *et al.*, 1996a, b; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Arkhipov & Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Günther, 1864a; Brauer, 1906; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Gonichthysocco* (Cocco 1829)**

Common name: **Cocco's lanternfish**; **Peixe-lanterna-de-Cocco**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (NHMUK 2003.4.11.176, 1959; NHMUK 2004.3.8.501, 1966; Hulley, 1984; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Lütken, 1892; Collett, 1896a; Brauer, 1906; Murat and Hjort, 1912; Roule and Angel, 1933; Fowler, 1936a; Belloc, 1938; Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Bolin, 1959; Osório de Castro, 1967; Nafpaktitis *et al.*, 1977; Hulley, 1984; Arruda, 1992, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1864b; Good and Bean, 1895; Brauer, 1906; Fowler, 1936a; NHMUK 1948.8.9.1., 1937-1960; Maul, 1946a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Bekker *et al.*, 1982; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019). The holotype of *Alysia loricata* Lowe 1839, synonym of *G.occo*, caught off in Madeira is in the NHMUK collections (NHMUK 1948.8.9.1).

***Hygophum benoiti* (Cocco 1838)**

Common name: **Benoit's lanternfish**; **Peixe-lanterna-de-Benoit**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (MNHN-IC-1987-0409, 1969; ZMH 105353, 1968; Hulley, 1984; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Lütken, 1892; Goode and Bean, 1895, 1896; Brauer, 1906; Zugmayer, 1911b; Fowler, 1936a; Belloc, 1938; Albuquerque, 1954-1956; Bolin, 1959; Osório de Castro, 1967; Geistdoerfer *et al.*, 1971; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Brauer, 1906; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Hygophum hygomii* (Lütken 1892)**

Common name: **Bermuda lantern fish; Peixe-lanterna-das-Bermudas**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([ZMH 117527](#), 1979; Hulley, 1984; Fischer *et al.*, 1987; ZSM-Pis-029412 / 534601 / 386189, 1990; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Lütken, 1892; Goode and Bean, 1895, 1896; Brauer, 1906; Tåning, 1932a; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Arkhipov & Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira ([Maul](#), 1946a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Hygophum macrochir* (Günther 1864)**

Common name: Largefin lanternfish; **Peixe-lanterna-de-barbatanas-grandes**.

References: Azores ([SOC Discovery 1023216_84_FIS_375003](#), 1980 (PECS); Carneiro *et al.*, 2014; Hulley, 2015g) and Madeira ([Maul](#), 1971a; ZMH 6973, 1979 (PECS); Carneiro *et al.*, 2014).

***Hygophum reinhardtii* (Lütken 1892)**

Common name: **Reinhardt's lanternfish; Peixe-lanterna-de-Reinhardt**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([YPM ICH 004954](#), 1959; ZSM-Pis-028818 / 534037 / 385625, 1991; Carneiro *et al.*, 2014), Azores ([MCZ 114804](#), 1969; MCZ 114811, 1972; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Arkhipov & Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016) and Madeira (Lütken, 1892; Brauer, 1906; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1971a; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Roberts *et al.*, 2017; Froese and Pauly, 2019).

***Hygophum taaningi* Becker 1965**

Common name: **Tåning's lanternfish; Peixe-lanterna-de-Tåning**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: Azores ([Bolin](#), 1959; NHMUK 2017.7.13.62-67, 1973; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira ([ZMH 103571](#), 1966; MCZ 114590, 1969; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016).

***Lampadена anomala* Parr 1928**

Common name: **Anomalous lanternfish; Peixe-lanterna-anómalo**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: Azores ([MCZ 149506](#), 1984; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016) and Madeira ([MCZ 96916](#), 1978; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Porteiro *et al.*, 2017).

***Lampadena atlantica* Maul 1969**

Common name: Atlantic tail-light lanternfish; **Peixe-lanterna-romeirinho**, Romeirinho.

References: Azores ([MAR-ECO 3434](#), n.d.; MAR-ECO 7371, n.d.; NHMUK 2015.7.9.161, 1981 (PECS); ISH 909-1982 (PECS); ISH 911-1982 (PECS); IMAR/DOP 2FPI-04-P02LUA, 2002; Pakhorukov, 2008 (PECS); Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014) and Madeira ([ZMH 103567](#), 1966; ZMH 103343, 1966; ZMH 105496, 1968; Maul, 1969a; Krefft and Bekker, 1979; Hulley, 1984, 1990b; ZSM-Pis-029350 / 534548 / 386136, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Porteiro *et al.*, 2017; Froese and Pauly, 2019). The holotype of subspecies *Lampadena urophaos atlantica* Maul 1969, currently in the MMF collections (MMF 2464), was caught off at Câmara de Lobos (Madeira) and was raised to synonym of species *L. atlantica*.

***Lampadena chavesi* Collett 1905**

Common name: Chaves' lanternfish; **Peixe-lanterna-de-Chaves**, Divertido, Escolarinho, Lobisomem, Pirilampo, Romeirinho.

References: mainland Portugal ([YPM ICH 004958](#), 1959; ZMH 105700, 1968; Hulley, 1984; Moura, 2004; Carneiro *et al.*, 2014), Azores ([Collett](#), 1905; Brauer, 1906; Fowler, 1936a; Belloc, 1938; Barnard, 1948;

Collins, 1954; Osório de Castro, 1967; Krefft, 1970; Dias *et al.*, 1976; Nafpaktitis *et al.*, 1977; Bekker, 1983; Albuquerque, 1954-1956; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016) and Madeira (NHMUK 2004.3.10.39, 1965; Osório de Castro, 1967(?); Krefft and Bekker, 1979; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016). The holotype (unique) of *Lampadena chavesi* Collett 1905, synonym of *L. chavesi*, caught off in the Azores is in the ZMUC collections (ZMUC J693).

***Lampadena luminosa* (Garman 1899)**

Common name: **Luminous lanternfish**; **Peixe-lanterna-luminoso**, Pirilampo.

References: mainland Portugal (Osório de Castro, 1967) and Madeira (Maul, 1946a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro *et al.*, 2014). Since Osório de Castro (1967) no other author described the presence of this species in mainland Portugal and its occurrence in this EEZ was not yet confirmed by more recent data. Specimens are needed for confirmation in mainland Portugal.

***Lampadena speculigera* Goode & Bean 1896**

Common name: Mirror lanternfish; **Peixe-lanterna-espelhado**, Divertido, Escolarinho, Lobisomem, Pirilampo, Romeirinho.

References: mainland Portugal (Costa and Geistdoerfer, 1979; Hulley, 1984; Carneiro *et al.*, 2014), Azores (ZMH 108469, 1971; MCZ 114345, 1972; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014) and Madeira (MCZ 114327, 1969; MCZ 114350, 1972; Krefft and Bekker, 1979; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016).

***Lampanyctus alatus* Goode & Bean 1896**

Common name: Winged lanternfish; **Peixe-lanterna-alado**, Peixinho-preto, Romeirinho.

References: mainland Portugal (MCZ 113950, 1972; Hulley, 1984; ZSM-Pis-028734 / 533955 / 385543, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (PECS) (MCZ 113874, 1969; SOC Discovery 826500_8_FIS_461003, 1973; Arkhipov & Mamedov, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016) and Madeira (MCZ 113872, 1969; MCZ 113949, 1972; Hulley, 1984, 1990b; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Lampanyctus ater* Tåning 1928**

Common name: Dusky lanternfish; **Peixe-lanterna-escuro**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: mainland Portugal (MNHN-IC-1912-0035, 1910; MNHN-IC-1987-0415, 1969; Hulley, 1984; Zahuranec, 2000; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; IFREMER Biocean 34554, 1969; NHMUK 2003.1.13.66-84, 1970; Geistdoerfer *et al.*, 1971; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Zahuranec, 2000; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Maul, 1971a; Osório de Castro, 1967; LACM 34893-1, 1969; LACM 34894-1, 1969; Hulley, 1984; Davies and Merrett, 1998; Zahuranec, 2000; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Lampanyctus crocodilus* (Risso 1810)**

Common name: **Jewel lanternfish**; **Peixe-lanterna-crocodilo**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (Bragança, 1903; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; ZMH 103734, 1966; Osório de Castro, 1967; Hulley, 1984; Fischer *et al.*, 1987; ZSM-Pis-028742 / 533963 / 385551, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Lütken, 1892 (PECS); Brauer, 1906 (PECS); Fowler, 1936a (PECS); Osório de Castro, 1967; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Johnson, 1862b, c; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Lampanyctus cuprarius* Tåning 1928**

Common name: Atlantic blackcap lanternfish; **Peixe-lanterna-cobreado**, Romeirinho.

References: mainland Portugal ([MCZ 112934, 1972 \(PECS\)](#); [Hulley, 1984](#); [Carneiro et al., 2014](#)), Azores ([MCZ 151346, 1978](#); [Hulley, 1984](#); [Arruda, 1997](#); [Porteiro et al., 1999, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)) and Madeira ([ZMH 103945, 1966](#); [LACM 34907-1, 1969](#); [MCZ 112815, 1978](#); [Hulley, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)).

Lampanyctus festivus Tåning 1928

Common name: Festive lanternfish; [Peixe-lanterna-divertido](#), Divertido, Escolarinho, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([YPM ICH 004975, 1959](#); [LACM 34910-1, 1969](#); [Hulley, 1984](#); [Carneiro et al., 2014](#)), Azores ([Tåning, 1928](#); [Nafpaktitis et al., 1977](#); [Krefft and Bekker, 1979](#); [Bekker, 1983](#); [Hulley, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)) and Madeira ([MCZ 112491, 1969](#); [MCZ 112497, 1970](#); [Maul, 1971a](#); [Krefft and Bekker, 1979](#); [Hulley, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Porteiro et al., 2017](#)).

Lampanyctus gemmifer Goode & Bean 1896

Common name: No English common name; [Peixe-lanterna-nacarado](#).

References: Azores ([Murray and Hjort, 1912](#); [Zugmayer, 1933](#); [Fowler, 1936a, b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bolin, 1959](#); [Geistdoerfer et al., 1971](#); [Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Stefanescu et al., 1994](#); [Santos et al., 1997](#)) and Madeira ([Zugmayer, 1911b](#), [1933](#); [Fowler, 1936a, b](#)).

Lampanyctus intricarius Tåning 1928

Common name: Intricate lanternfish; [Peixe-lanterna-complexo](#), Divertido, Escolarinho, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([Tåning, 1928](#); [ZMH 105332, 1968](#); [LACM 34913-1, 1969](#); [Krefft and Bekker, 1979](#); [Hulley, 1984](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)), Azores ([Bolin, 1959](#); [Nafpaktitis et al., 1977](#); [Bekker, 1983](#); [Hulley, 1984](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([MCZ 112323, 1969](#); [Hulley, 1984](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

Lampanyctus lineatus Tåning 1928

Common name: Lanternfish; [Peixe-lanterna-riscado](#), Romeirinho.

References: mainland Portugal ([MCZ 112243, 1969](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#)), Azores ([Krefft and Bekker, 1979 \(PECS\)](#); [MAR-ECO 14894, 2004 \(PECS\)](#); [MAR-ECO 14866, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Porteiro et al., 2017](#)) and Madeira ([NHMUK 2004.3.10.1-2, 1965](#); [ZMH 103666, 1966](#); [Hulley, 1984](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Hulley and Paxton, 2016](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)). Froese and Pauly (2019) consider *Lampanyctodes lineatus* Tåning 1928 synonym of *Nannobrachium lineatum* Tåning 1928 and according to Eschmeyer et al. (2019) *Nannobrachium lineatum* Tåning 1928 is synonym of *L. lineatus*.

Lampanyctus macdonaldi (Goode & Bean 1896)

Common name: MacDonald's lanternfish; [Peixe-lanterna-de-MacDonald](#).

References: mainland Portugal ([NHMUK 2004.3.20.564-568, 1978 \(PECS\)](#); [Carneiro et al., 2014](#); [Munroe et al., 2015a](#)), Azores ([NHMUK 2004.8.18.78-86, 1980 \(PECS\)](#); [MAR-ECO 5533, 2004](#); [MAR-ECO 7364, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Munroe et al., 2015a](#); [Porteiro et al., 2017](#)) and Madeira ([Munroe et al., 2015a](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *L. macdonaldi* is the Atlantic and southern hemisphere. The distribution map of this species reported by Munroe et al. (2015a) indicates the occurrence of this species in the area of Madeira Islands. Specimens are needed for confirmation in Madeira Archipelago.

Lampanyctus niger (Günther 1887)

Common name: Black lantern fish; [Peixe-lanterna-preto](#).

References: mainland Portugal (?) ([YPM ICH 004979, 1959](#)) and Azores (?) ([Pappenheim, 1914](#); [Fowler, 1936a](#)). Pappenheim (1914) indicates the occurrence of this species in the Atlantic, south of Azores, 28°25.2'N, 34°19.8'W. This geographic coordinates correspond to a position in the Atlantis Fracture Zone, out of the PECS area. This species is also regarded as being present off mainland Portugal,

37°32.4'N, 12°57.0'W, and there is a specimen deposited at Yale University Peabody Museum. According to Eschmeyer *et al.* (2019) the current geographic distribution of *L. niger* is the Indo-West Pacific: Indonesia and Vietnam east to Hawaiian Islands and Tuamotu Archipelago, north to southern Japan, south to New Caledonia and northern New Zealand. Thus, this specimen from Yale University Peabody Museum could have been misidentified (probably confused with *Lampanyctus ater* Tåning 1928) and according to the current geographic distribution this species it is not considered in the total number of valid species.

***Lampanyctus photonotus* Parr 1928**

Common name: Dotback lanternfish; **Peixe-lanterna-lobisomem**, Divertido, Escolarinho, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (YPM ICH 004981, 1959; MCZ 111942, 1969; Krefft and Bekker, 1979; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016), Azores (Bolin, 1959; Nafpaktitis *et al.*, 1977; Krefft and Bekker, 1979; Bekker, 1983; Hulley, 1984; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Hulley and Paxton, 2016; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014) and Madeira (Maul, 1971a; MCZ 111870, 1978; Krefft and Bekker, 1979; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016).

***Lampanyctus pusillus* (Johnson 1890)**

Common name: Pygmy lanternfish; **Peixe-lanterna-anão**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (YPM ICH 004990, 1959; Hulley, 1984; ZMH 117083, 1979; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Zugmayer, 1911b; Roule and Angel, 1933; Fowler, 1936b; Belloc, 1938; Ferreira, 1942; Collins, 1954; Albuquerque, 1954-1956; Bolin, 1959; Osório de Castro, 1967; Geistdoerfer *et al.*, 1971; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (NHMUK 1890.5.31.8, 1878-1892; Johnson, 1890; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of *Scopelus pusillus* Johnson 1890, synonym of *L. pusillus*, caught off in Madeira is in the NHMUK collections (NHMUK 1890.5.31.8).

***Lepidophanes gaussi* (Brauer 1906)**

Common name: Gauss' lanternfish; **Peixe-lanterna-de-Gauss**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: mainland Portugal (YPM ICH 004946, 1959; ZMH 116740, 1979; Hulley, 1984; Carneiro *et al.*, 2014), Azores (Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Diekmann *et al.*, 2006 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016) and Madeira (Maul, 1971a; MCZ 109319, 1978; MCZ 109321, 1978; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016).

***Lepidophanes guentheri* (Goode & Bean 1896)**

Common name: Günther's lanternfish; **Peixe-lanterna-de-Günther**, Divertido, Escolarinho, Lobisomem, Peixinho-preto, Romeirinho.

References: Azores (MCZ 108641, 1964; Nafpaktitis *et al.*, 1977; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014) and Madeira (ZMH 103891, 1966; Osório de Castro, 1967; Hulley, 1984; Lloris *et al.*, 1991; Porteiro *et al.*, 2017; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Lobianchia dofleini* (Zugmayer 1911)**

Common name: Doflein's lantern fish; **Peixe-lanterna-de-Doflein**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (YPM ICH 005036, 1959; ZMH 116668, 1979; Hulley, 1984; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Bolin, 1959; Nafpaktitis, 1968; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Maul, 1946a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967;

Krefft and Bekker, 1979; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Lobianchia gemellarii* (Cocco 1838)**

Common name: **Gemellar's lanternfish**; **Peixe-lanterna-de-Gemellar**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([ZMH 103835](#), 1966; Hulley, 1984; Fischer *et al.*, 1987; ZSM-Pis-028658 / [533881](#) / [385469](#), 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores ([Murray and Hjort, 1912](#); [Roule and Angel, 1930](#); [Zugmayer, 1933](#); [Fowler, 1936a,b](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Nafpaktitis, 1968](#); [Nafpaktitis *et al.*, 1977](#); [Bekker, 1983](#); [Hulley, 1984](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 1999, 2010, 2017](#); [Sutton *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Hulley, 1984](#); [Carneiro *et al.*, 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Loweina interrupta* (Tåning 1928)**

Common name: Barebelly lanternfish; **Peixe-lanterna-ferreiro**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([YPM ICH 005049](#), 1959; Hulley, 1984; Carneiro *et al.*, 2014; [Ralph, 2015h](#)), Azores ([Nafpaktitis *et al.*, 1977](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 1999, 2010, 2017](#); [Sutton *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Ralph, 2015h](#)) and Madeira ([Maul, 1946a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#)).

***Loweina rara* (Lütken 1892)**

Common name: Rare lanternfish; **Peixe-lanterna-raro**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([Brauer, 1906\(?\)](#); [Nobre, 1935\(?\)](#); [Osório de Castro, 1967](#)), Azores ([MAR-EKO - HamPelFish - TI/SF_MAR-ECO_3303](#), 1979 (PECS); [Hulley, 1984](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 1999, 2010](#); [Arkhipov & Mamedov, 2008](#) (PECS); [Carneiro *et al.*, 2014](#); [Hulley and Paxton, 2016](#)) and Madeira ([Brauer, 1906](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Hulley, 1984](#); [Carneiro *et al.*, 2014](#); [Hulley and Paxton, 2016](#)). According to Froese and Pauly (2019), *L. rara* occurs in the Atlantic Ocean between 46°N and 26°S [in tropical through temperate seas (Eschmeyer *et al.* (2019))].

***Myctophum affine* (Lütken 1892)**

Common name: **Metallic lanternfish**; **Peixe-lanterna-metálico**, Ferreiro, Peixinho-preto, Romeirinho.

References: Azores (PECS) ([Murray and Hjort, 1912](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Carneiro *et al.*, 2014](#)) and Madeira (?) ([Osório de Castro, 1967\(?\)](#)). The only report of the presence of *M. affine* in the fishing grounds of Madeira is indicated by Osório de Castro (1967). Since then no other author describes the presence of this species in Madeira. Thus, the occurrence of this species in the Madeira Archipelago is questionable and is not considered in the total number of valid species for the Madeiran ichthyofauna.

***Myctophum nitidulum* Garman 1899**

Common name: Pearlyspotted lanternfish; **Peixe-lanterna-fosforescente**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: Azores ([MCZ 106197](#), 1969; [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 1999, 2010](#); [Arkhipov & Mamedov, 2008](#) (PECS); [Carneiro *et al.*, 2014](#)) and Madeira ([MCZ 106061](#), 1969; [ZMH 107695](#), 1970; [Hulley, 1984](#); [Carneiro *et al.*, 2014](#); [Hulley and Paxton, 2016](#); [Froese and Pauly, 2019](#)).

***Myctophum punctatum* Rafinesque 1810**

Common name: **Spotted lanternfish**; **Peixe-lanterna-manchado**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal ([Brauer, 1906](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft and Bekker, 1979](#); [Martins, 1981](#); [Hulley, 1984](#); [Fischer *et al.*, 1987](#); [Carneiro *et al.*, 2014](#); [Mecklenburg *et al.*, 2018](#); [Froese and Pauly, 2019](#)), Azores ([Lütken, 1892](#); [Collett, 1896a](#); [Brauer, 1906](#); [Murray and Hjort, 1912](#) (PECS); [Vaillant, 1919](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Belloc,](#)

1938; Collins, 1954; Albuquerque, 1954-1956; Bolin, 1959; Osório de Castro, 1967; Parin, 1970; Saldanha, 1977a; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Monteiro, 1996; Monteiro *et al.*, 1996a, b; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Arkhipov & Mamedov, 2008 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira (Lütken, 1892; Brauer, 1906; Nobre, 1935; Maul, 1946a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Mecklenburg *et al.*, 2018).

***Notolychnus valdiviae* (Brauer 1904)**

Common name: Topside lanternfish; **Peixe-lanterna-do-Valdívia**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho.

References: mainland Portugal (MCZ 104925, 1969; MCZ 105086, 1972; Hulley, 1984; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Bolin, 1959; Nafpaktitis *et al.*, 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Diekmann *et al.*, 2006 (PECS); Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (MCZ 104947, 1970; Maul, 1971a; MCZ 105076, 1972; Hulley, 1984; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Notoscopelus bolini* Nafpaktitis 1975**

Common name: Fanfare lanternfish; **Peixe-lanterna-de-Bolin**, Divertido, Escolarinho, Lobisomem.

References: mainland Portugal (ICM-CSIC 7258, n.d.; ZMH 106027, 1968; Maul, 1976; Hulley, 1984; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Kobylansky *et al.*, 2015c; Froese and Pauly, 2019), Azores (MCZ 104023, 1969; Maul, 1976; Nafpaktitis *et al.*, 1977; Hulley, 1984; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Kobylansky *et al.*, 2015c; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Kobylansky *et al.*, 2015c; Hulley and Paxton, 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *N. bolini* is the Eastern Atlantic and Mediterranean Sea. Kobylansky *et al.* (2015c) indicate the occurrence of this species in Madeira, and Hulley and Paxton (2016) off Canary Islands. However, the distribution maps of this species reported by the latter authors partially included the area of Selvagens Islands (Madeira EEZ). Thus, a specimen is needed for confirmation. One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22512, 1967).

***Notoscopelus caudispinosus* (Johnson 1863)**

Common name: Spinetail lanternfish; **Peixe-lanterna-de-cauda-espinhosa**, Divertido, Escolarinho, Lobisomem, Romeirinho.

References: Azores (Zugmayer, 1933; Collins, 1954; MCZ 104073, 1972; Bekker *et al.*, 1982; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (NHMUK 1862.2.5.3, 1861-1890; Johnson, 1863; Goode and Bean, 1895, 1896; MCZ 104054, 1969; MCZ 104063, 1970; Krefft and Bekker, 1979; Bekker *et al.*, 1982; Hulley, 1984; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Hulley and Paxton, 2016; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of *Scopelus caudispinosus* Johnson 1863, synonym of *N. caudispinosus*, caught off in Madeira is in the NHMUK collections (NHMUK 1862.2.5.3).

***Notoscopelus elongatus* (Costa 1844)**

Common name: No English common name; **Peixe-lanterna-do-Mediterrâneo**.

References: mainland Portugal (?) (MNHN-IC-1987-0444, 1969; IFREMER Biocean 34520, 1969; Carneiro *et al.*, 2014), Azores (?) (Zugmayer, 1933; Fowler, 1936b; Collins, 1954) and Madeira (?) (Brauer, 1906; Maul, 1946a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956). According to Eschmeyer *et al.* (2019) this species is endemic in Mediterranean Sea and Sea of Marmara. The occurrence records and bibliographic references of *Notoscopelus elongatus* for mainland Portugal and Madeira may refer eventually to *Notoscopelus elongatus* subsp. *kroyeri* (Malm 1861), synonym of *Notoscopelus kroyeri* (Malm 1861). Santos *et al.* (1997) mention that the specimens identified as *Lampanyctus elongatus* by Collins (1954) and *Myctophum (Lampanyctus) elongatus* by Zugmayer (1933) are misidentifications of *Notoscopelus kroeyerii* (Malm 1861). Thus, these specimens need to be revaluated (rank reclassified) and, according to the current geographic distribution, this species is not considered in the total number of valid species.

***Notoscopelus kroyeri* (Malm 1861)**

Common name: Krøyer's lanternfish; **Peixe-lanterna-de-Krøyer**, Divertido, Escolarinho, Lobisomem. References: mainland Portugal ([ZMH 104287](#), 1966; Hulley, 1984; [ZSM-Pis-029209 / 534414 / 386002](#), 1992; Carneiro et al., 2014; Mecklenburg et al., 2018; Froese and Pauly, 2019), Azores ([Zugmayer](#), 1933; Collins, 1954; MCZ 104268, 1972; Hulley, 1984; Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Mecklenburg et al., 2018) and Madeira ([ZSM-Pis-029348 / 534546 / 386134](#), 1990; Carneiro et al., 2014; Hulley and Paxton, 2016).

***Notoscopelus resplendens* (Richardson 1845)**

Common name: **Patchwork lanternfish**; **Peixe-lanterna-oliva**, Divertido, Escolarinho, Ferreiro, Lobisomem, Peixinho-preto, Romeirinho. References: mainland Portugal ([YPM ICH 005135](#), 1959; MCZ 103774, 1969; MCZ 103858, 1969; Hulley, 1984; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([Belloc](#), 1938; Dias et al., 1976; Nafpaktitis et al., 1977; Bekker et al., 1982; Becker, 1983; Hulley, 1984; Clarke et al., 1995; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira (Johnson, 1863; Günther, 1864a; Fowler, 1936a; Maul, 1946a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Nafpaktitis 1975; Krefft and Bekker, 1979; Hulley, 1984; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Protomyctophum arcticum* (Lütken 1892)**

Common name: Arctic lanternfish; **Peixe-lanterna-do-Ártico**, Divertido, Escolarinho, Lobisomem. References: mainland Portugal ([IFREMER Biocean 34590](#), 1969; Carneiro et al., 2014) and Azores ([MCZ 103011](#), 1964; Nafpaktitis et al., 1977; Bekker, 1983; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Carneiro et al., 2014).

***Scopelopsis multipunctatus* Brauer 1906**

Common name: Multispotted lanternfish; **Peixe-lanterna-sarapintado**. References: mainland Portugal (?) ([GMN 1946-7628](#), 1946) and Madeira (?) ([Brauer](#), 1906; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Carneiro et al., 2014). There is one specimen caught off at 36°39.60'N, 13°4.80'W, June 1, 1946 and preserved in the Gottenburg Natural History Museum identified by O. Nybelin (GMN Pisces 1946-7628). However, the current geographic distribution is circumglobal in the southern hemisphere (Hulley and Paxton, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). Thus, this species is not considered in the total number of valid species.

***Symbolophorus rufinus* (Tåning 1928)**

Common name: Rufous lanternfish; **Peixe-lanterna-de-Rufous**. References: Azores ([PECS](#)) ([MCZ 103498](#), 1973; SOC Discovery 1022209_86_FIS_785001, 1980; Diekmann et al., 2006; Carneiro et al., 2014).

***Symbolophorus veranyi* (Moreau 1888)**

Common name: **Large-scale lantern fish**; **Peixe-lanterna-de-Vérany**, Divertido, Escolarinho, Lobisomem, Romeirinho. References: mainland Portugal ([NHMUK 2003.4.11.113-115](#), 1959; Hulley, 1984; Fischer et al., 1987; [ZSM-Pis-029226 / 534429 / 386017](#), 1992; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([Lütken](#), 1892; Brauer, 1906; Tåning, 1932c; Nobre, 1935; Fowler, 1936a; Belloc, 1938; Collins, 1954; Albuquerque, 1954-1956; Bolin, 1959; Dias et al., 1976; Nafpaktitis et al., 1977; Bekker, 1983; Hulley, 1984; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019) and Madeira ([Lütken](#), 1892; Fowler, 1936a; Hulley, 1984; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

***Taaningichthys bathyphilus* (Tåning 1928)**

Common name: Deepwater lanternfish; **Peixe-lanterna-da-fundura**, Romeirinho. References: Azores ([MNHN-IC-1980-1379](#), 1979; Sutton et al., 2008; Carneiro et al., 2014; Hulley and Paxton, 2016; Porteiro et al., 2017) and Madeira ([NHMUK 2004.3.10.33](#), 1965; Hulley, 1984; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019).

Taaningichthys minimus (Tåning 1928)

Common name: **Waistcoat lanternfish; Peixe-lanterna-de-Cabo-Verde**, Escolarinho, Romeirinho.

References: mainland Portugal ([ZSM-Pis-029148 / 534358 / 385946, 1992](#)), Azores ([YPM ICH 007724, 1931; ZMH 114111, 1975; ZMH 118457, 1979; Arruda, 1997; Hulley, 1984; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019](#)) and Madeira ([ZMH 104184, 1966; ZMH 105938, 1968; Hulley, 1984; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019](#)).

Taaningichthys paurolychnus Davy 1972

Common name: Naked lanternfish; **Peixe-lanterna-nu**, Romeirinho.

References: Azores ([SOC Discovery 748201_8_FIS_854003, 1970 \(PECS\); SOC Discovery 1319802_85_FIS_854003, 1997; NHMUK 2010.10.27.2, 1981 \(PECS\); NHMUK 2003.1.16.125, 1981 \(PECS\); Carneiro et al., 2014](#)) and Madeira ([Krefft and Bekker, 1979; Hulley, 1984; Lloris et al., 1991; Carneiro et al., 2014; Hulley and Paxton, 2016; Froese and Pauly, 2019](#)).

Triphoturus nigrescens (Brauer 1904)

Common name: Highseas lampfish; **Peixe-lanterna-do-alto**.

References: Azores (?) ([Murray and Hjort, 1912](#)). There is one specimen of *Myctophum (Lampanyctus) micropterum*, caught at 32°12.0'N, 34°42.0'W, Stations 51, June 5-6, 1910 (Murray and Hjort, 1912). However, the current geographic distribution is the Indo-Pacific: East Africa east to Hawaiian Islands and tropical Eastern Pacific, north to southern Japan, south to Western Australia, New Caledonia and New Zealand (Eschmeyer et al., 2019). Thus, this species is not considered in the total number of valid species.

Order Lampriformes

Family Lampridae

Lampris guttatus (Brünnich 1788)

Common name: **Opah; Peixe-cravo**, Cravo, Joaninha, Lua, Mariposa, Opá, Rei, São-pedro, Sô, Sol.

References: mainland Portugal[▲] ([Steindachner, 1868a; Osório, 1894; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer et al., 1981, 1987; Palmer, 1986; Carneiro et al., 2014; Mecklenburg et al., 2018; Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967; Parin, 1970; Wheeler, 1978; Fischer et al., 1981; Palmer, 1986; Santos et al., 1994a, 1995a, 1997; Pinho and Pereira, 1995a; Arruda, 1997; Porteiro et al., 2010, 2017\(?\); Carneiro et al., 2014; Mecklenburg et al., 2018; Froese and Pauly, 2019](#)) and Madeira[▲] ([de Velloso, 1892\(?\); Nobre, 1935; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Palmer, 1979, 1986; Fischer et al., 1981; Quéro and Gayet, 1990; Lloris et al., 1991; Severino, 2004; Carneiro et al., 2014; Mecklenburg et al., 2018; Froese and Pauly, 2019](#)). *Zeus luna* Gmelin 1789 and *Lampris luna* (Gmelin 1789) are synonyms of *Lampris guttatus* (Brünnich 1788). Several authors mentioned the occurrence of *Lampris lauta* Lowe 1838 in Madeira but erroneously included it in the synonymy of *L. guttatus*.

Lampris lauta Lowe 1838

Common name: East Atlantic Opah; **Peixe-cravo-elegante**, Peixe-cravo.

References: Azores ([Underkoffler et al., 2018; Eschmeyer et al., 2019](#)) and Madeira ([Lowe, 1837, 1839, 1840b, 1843-1860, 1844a, 1849; Osório, 1894\(?\); Goode and Bean, 1895\(?\), 1896\(?\); Underkoffler et al., 2018\(?\)](#)). The holotype and neotype of this species was caught off in Madeira. The neotype is in the MMF collections [MMF 42252, 2012; caught off "North of Madeira Island" (Underkoffler et al., 2018)] and the whereabouts of this holotype are unknown.

Family Lophotidae

Lophotus lacepede Giorna 1809

Common name: **Crested oarfish; Peixe-remo-de-crista**, Peixe-remo, Remo.

References: mainland Portugal ([Osório de Castro, 1967\(?\); Saldanha, 1977b; Hureau and Monod, 1979; Palmer, 1986; Quiniou and Quéro, 1990; Carneiro et al., 2014; Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1863.8.27.1, 1861-1890; Johnson, 1863; Goode and Bean, 1895, 1896; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Palmer, 1979, 1986; Robins and Ray, 1986](#)).

Quiniou and Quéro, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Lophotes cristatus* Johnson 1863, synonym of *L. lacepede*, caught off in Madeira is in the NHMUK collections (NHMUK 1863.8.27.1).

Family Radiicephalidae

Radiicephalus elongatus Osório 1917

Common name: **Tapetail; Peixe-remo-de-cauda-fina**.

References: mainland Portugal (Palmer, 1986; Koeda and Ho, 2018), Azores (NHMUK 1967.10.2.2, 1966; Harrisson and Palmer, 1968; Palmer, 1979, 1986; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Koeda and Ho, 2018; Froese and Pauly, 2019) and Madeira (ZMH 103552, 1966; Palmer, 1986; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Family Trachipteridae

Trachipterus arcticus (Brünnich 1788)

Common name: **Dealfish; Peixe-tábua**, Rei-dos-arenques.

References: mainland Portugal (Nobre, 1935; Albuquerque, 1954-1956; Almaça, 1965a; Osório de Castro, 1967; Palmer, 1986; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019; Moore, 2019), Azores (Regan, 1903a; Albuquerque, 1954-1956, ZMH 114011, 1975; Wheeler, 1978; Palmer, 1986; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017(?); Carneiro *et al.*, 2014; Moritz *et al.*, 2015; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019; Moore, 2019) and Madeira (Lowe, 1850, 1852; Goode and Bean, 1895, 1896; NHMUK 1917.7.14.83., 1912-1936; Palmer, 1979, 1986; Aguiar and Quéro, 1990; Lloris *et al.*, 1991; de Casamajor and Morandeau, 2013; Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019; Moore, 2019). The holotype (unique) of *Trachypterus gryphurus* Lowe 1852, synonym of *T. arcticus*, caught off in Madeira is in the NHMUK collections (NHMUK 1917.7.14.83). Mecklenburg *et al.* (2018) mention that, in the Eastern Atlantic, differences between *T. arcticus* and *T. trachypterus* are not always clear, as in specimens from the Azores recently reported (Moritz *et al.*, 2015). Thus, the former authors consider the need of reviewing this family.

Trachipterus trachypterus (Gmelin 1789)

Common name: **Mediterranean dealfish; Peixe-tábua-do-Mediterrâneo**, Peixe-tábua.

References: mainland Portugal (Osório, 1917; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Farias *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Regan, 1903a; Albuquerque, 1954-1956; Osório de Castro, 1967; Arruda, 1997; ZMH 117932, 1979; ZMH 119183, 1979 (PECS); ZMH 119184, 1979 (PECS); Carneiro *et al.*, 2014) and Madeira (Lowe, 1850, 1852; Günther, 1861a; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro *et al.*, 2014).

Zu cristatus (Bonelli 1819)

Common name: **Scalloped ribbonfish; Peixe-tábua-recortado**.

References: mainland Portugal (Smith-Vaniz and Herrera, 2015a), Azores (Palmer, 1961, 1979, 1986; Parin, 1970; MCZ 84699, 1984; Aguiar and Quéro, 1990; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010; Carneiro *et al.*, 2014; Quigley and Henderson, 2014; Smith-Vaniz and Herrera, 2015a; Froese and Pauly, 2019) and Madeira (Palmer, 1979, 1986; Aguiar and Quéro, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Quigley and Henderson, 2014; Smith-Vaniz and Herrera, 2015a; Froese and Pauly, 2019).

Family Regalecidae

Regalecus glesne Ascanius 1772

Common name: **King of herrings; Rei-dos-Arenques**, Pai-da-sarda, Peixe-real, Regaleco, Regalengo, Relangueiro.

References: mainland Portugal (Frade, 1934; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Palmer, 1986; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Wheeler, 1978; Palmer, 1986; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017(?);

Carneiro *et al.*, 2014) and Madeira (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Palmer, 1979(?), 1986; Carneiro *et al.*, 2014).

Regalecus russellii (Cuvier 1816)

Common name: Oarfish; **Peixe-remo-do-Índico**.

References: Madeira (?) ([MOM-IC-0091-1199, 1912](#); Zugmayer, 1914; Roberts, 2012(?); Froese and Pauly, 2019(?)). The holotype (unique) of *Regalecus caudatus* Zugmayer 1914, synonym of *R. russellii* (Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), caught off SW of Madeira is in the MOM collections ([MOM-IC-0091-1199](#)). According to Eschmeyer *et al.* (2019) *R. russellii* is a circumglobal species in warm waters. Roberts (2012) re-evaluates the holotype (a small juvenile), whose characters don't allow the correct diagnostic because the counts of dorsal fin rays and vertebrae to the vent should provide identification as either *R. russellii* or *R. glesne*. Thus, this species is not considered in the total number of valid species.

Order Zeiformes

Family Oreosomatidae

Allocyttus verrucosus (Gilchrist 1906)

Common name: **Warty dory; Galo-verrugoso**.

References: Azores (PECS) ([Hareide and Garnes, 2001](#); [Pakhorukov, 2008](#); [Carneiro *et al.*, 2014](#)).

Neocyttus helgae (Holt & Byrne 1908)

Common name: **Irish oreo; Galo-irlandês**.

References: Azores ([MNHN-IC-2009-0472, 1971](#); Saldanha and Biscoito, 1997; Santos *et al.*, 1997; Melo and Menezes, 2002; Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019) and Madeira ([Maul, 1948b, 1949b, c](#); Osório de Castro, 1967; Wheeler, 1979c; Karrer 1986a; Lloris *et al.*, 1991; Quéro *et al.*, 1993, 2000; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Heemstra, 2016a; Porteiro *et al.*, 2017; Froese and Pauly, 2019). This species can be confused with *Neocyttus rhomboidalis* Gilchrist 1906. The holotype (unique) of *Crassispinus granulosus* Maul 1948, synonym of *N. helgae*, caught off near Porto Moniz, north side of Madeira is in the MMF collections.

Neocyttus rhomboidalis Gilchrist 1906

Common name: **Spiky oreo; Galo-espinhoso**.

References: Madeira (?) ([Maul, 1949b](#); [Albuquerque, 1954-1956](#); [Quéro, 1982d](#); [Carneiro *et al.*, 2014](#)?); Heemstra, 2016a). This species can be confused with *Neocyttus helgae* (Holt and Byrne, 1908). Maul (1949b) re-evaluated the specimen of *Crassispinus granulosus* Maul 1948 and considered it a synonym of *N. rhomboidalis* Gilchrist 1906, but that specimen is distinct from the other specimens described by other authors and is now considered a synonym of *N. helgae*. According to Eschmeyer *et al.* (2019) *N. rhomboidalis* is a circumglobal species in Southern seas. The occurrence of this species in the Madeira region is not yet confirmed and specimens are needed for validation in this archipelago. Thus, this species is not considered in the total number of valid species.

Family Parazenidae

Cytopsis rosea (Lowe 1843)

Common name: **Red dory; Galo-de-natura**, Galo, Peixe-galo, Peixe-galo-de-natura, Peixe-prata.

References: mainland Portugal ([Osório, 1917](#); NHMUK 1972.1.10.60-62, 1971; Fernández *et al.*, 1981; Martins, 1981; Quéro, 1986a; Quéro *et al.*, 1997, 1998; Borges *et al.*, 2002; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Chaves, 2008, 2009; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; OCEANA, 2014; Garcia, 2015c; Froese and Pauly, 2019), Azores ([Maul, 1976](#) (PECS); Ehrich, 1977 (PECS); IMAR/DOP COLETA 130, 2003; Fock *et al.*, 2002 (PECS); Kukuev, 2002 (PECS); Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Garcia, 2015c) and Madeira ([Lowe, 1843, 1844b](#); NHMUK 1852.9.13.104, 1850-1853; Günther, 1860; Johnson, 1863b; Goode and Bean, 1895, 1896; Brauer, 1906; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Ehrich, 1977; Wheeler, 1979d; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Garcia, 2015; Heemstra, 2016a; Roberts *et al.*, 2017). The syntype of *Zeus roseus* Lowe 1843, synonym of *C. rosea*, caught off in

Madeira is in the NHMUK collections (NHMUK 1852.9.13.104). One specimen caught off at Meteor Bank is in the MMF collections (MMF 22354, 1967).

Family Zeniontidae

Zenion hololepis (Goode & Bean 1896)

Common name: **Dwarf dory; Galito-rosado**, Peixe-prata.

References: mainland Portugal ([Martins et al., 2012](#); [Carneiro et al., 2014](#); [Ralph, 2015i](#); [Froese and Pauly, 2019](#)). The first specimen of this species in Portuguese continental waters was collected in 2008 and has associated the INSDC sequences JF718832 and JF718833 [European Nucleotide Archive (EMBL-EBI)].

Family Grammicolepididae

Grammicolepis brachiusculus Poey 1873

Common name: **Thorny tinselfish; Peixe-prata-espinhoso**.

References: mainland Portugal ([Vasconcelos et al., 2003](#); [KJ768237, 2009](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Garcia, 2015d](#)) and Azores ([Fock et al., 2002 \(PECS\)](#); [Kukuev, 2002 \(PECS\)](#); [Porteiro et al., 2010](#); [Barreiros et al., 2011](#); [Carneiro et al., 2014](#); [Garcia, 2015d](#); [Froese and Pauly, 2019](#)).

Family Zeidae

Zenopsis conchifer (Lowe 1852)

Common name: **Silvery John dory; Galo-branco**, Galinha, Galinha-do-mar, Galito, Galo, Galo-da-fundura, Galo-de-fundura, Galo-de-natura, Peixe-galinha, Peixe-galo, Peixe-galo-branco, Peixe-galo-da-fundura, Peixe-galo-de-fundura, Peixe-galo-de-natura, Peixe-prata.

References: mainland Portugal[▲] ([Saldanha, 1968b](#); [Maul, 1976](#); [Quéro et al., 1976, 1997, 1998](#); [Fischer et al., 1981](#); [Dinis, 1985](#); [Quéro, 1986a](#); [Borges et al., 2002, 2003](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Heemstra, 2016b](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Maul, 1976 \(PECS\)](#); [Ehrich, 1977 \(PECS\)](#); [Santos et al., 1994a](#), [1995a](#), [1997](#); [Silva et al., 1994](#); [Azevedo and Heemstra, 1995](#); [Silva and Menezes, 1995a](#); [Krug, 1995](#); [Arruda, 1997](#); [Kukuev, 2002 \(PECS\)](#); [Menezes et al., 2006](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#)) and Madeira ([Lowe, 1850, 1852](#); [Günther, 1860](#); [Johnson, 1863b](#); [de Brito Capello, 1871](#); [Nunes, 1953, 1974](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Quéro et al., 1976](#); [Ehrich, 1977](#); [Fischer et al., 1981](#); [Karrer and Post, 1990a](#); [Lloris et al., 1991](#); [Christiansen et al., 2009](#); [Carneiro et al., 2014](#); [Biscoito et al., 2017](#); [Froese and Pauly, 2019](#)). Several specimens caught off at the Meteor Bank are in the MMF collections (MMF 22374, 1967, set of four specimens, MMF 22375, 1967, set of six specimens and MMF 22402, 1967, set of five specimens).

Zeus faber Linnaeus 1758

Common name: **John dory; Galo-negro**, Afaquete, Alfaquete, Alfaqui, Alfaquim, Alfaquique, Carta, Enxarroco, Enxarrôco, Galo, Galito, Galo-cristo, Galo-de-São-Pedro, Peixe-galo, Peixe-galo-negro, Peixe-São-Pedro, São-Pedro, Xaroco.

References: mainland Portugal[▲] ([Carvalho, 1765](#); [Vandelli, 1771](#); [de Brito Capello, 1867a](#), [1869b](#), [1880](#); [Steindachner, 1868a](#); [Gervais and Boulart, 1877b](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1893, 1897, 1898](#); [Nobre, 1894, 1895a, c](#), [1903, 1935](#); [Osório, 1895](#); [Bragança, 1902](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Muus and Dahlström, 1978](#); [Wheeler, 1979d](#); [INIP, 1981a, b](#), [1982a, b](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Quéro, 1986a](#); [Martins et al., 1992, 2015](#); [Almeida, 1996](#); [Erzini et al., 2001a](#); [Gomes et al., 2001](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Stergiou et al., 2006](#); [Fernandes et al., 2007](#); [Henriques et al., 2007, 2008](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Silva, 2008, 2010, 2011](#); [Ward et al., 2008](#); [Abecasis et al., 2009](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Drouët, 1861b](#);

Ramos, 1869; Hilgendorf, 1888a; Regan, 1903a; Sampaio, 1904; Fowler, 1936b; Nunes, 1953, 1974; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Maul, 1976; Wheeler, 1979d; Martins, 1982a; INIP, 1985c; Dias and Monteiro, 1986; Quéro, 1986a; Dias, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1994a, 1995a, 1997; Silva *et al.*, 1994, 1995; Wirtz, 1994; Azevedo, 1995; Krug, 1995; Silva and Menezes, 1995b; Menezes, 1996; Kukuev, 2002 (PECS); Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Gomes-Pereira *et al.*, 2014; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1949c, 1976; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Wheeler, 1979d; Fischer *et al.*, 1981; Quéro, 1986a; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Heemstra, 2016b; Ryland and Lancaster, 2017; Froese and Pauly, 2019).

Order Stylephoriformes

Family Stylephoridae

***Stylephorus chordatus* Shaw 1791**

Common name: **Tube-eye; Peixe-cauda-de-fio.**

References: Azores (IMAR/DOP COLETA 1697, 2001; Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Carneiro *et al.*, 2014).

Order Gadiformes

Family Bregmacerotidae

***Bregmaceros atlanticus* Goode & Bean 1886**

Common name: Antenna codlet; **Peixe-antena-do-Atlântico.**

References: Madeira (?) (Maul 1952a(?); Goren and Galil, 2008; Iwamoto, 2010; Nedreaas *et al.*, 2015a).

According to Eschmeyer *et al.* (2019) *B. atlanticus* is a circumglobal species in tropical and warm temperate seas. The only two specimens of the genus *Bregmaceros* for Madeira were described by Maul (1952a). According to this author, the Madeira specimens show characters which strongly disagree with the three possible species *atlanticus*, *japonicus* and *macclellandi*, taking into account the characters and geographic distribution of these species, which were then included in the genus *Bregmaceros*. This genus contains fourteen valid nominal species (Eschmeyer *et al.*, 2019) and at least four undescribed species (Harold and Golani, 2016). According to Eschmeyer *et al.* (2019) only four species (*atlanticus*, *macclellandi*, *cantori* and *nectabanus*) are likely to be found in Madeira taking into account their current geographic distributions. However, the meristic and morphometric characters of *B. nectabanus* and *B. atlanticus* may lead to misidentifications (Harold and Golani, 2016). The occurrence of this species in the Madeira region is not yet confirmed and specimens are needed. Thus, this species is not considered in the total number of valid species.

Family Macrouridae

***Asthenomacrurus victoris* Sazonov & Shcherbachov 1982**

Common name: Victory whiptail; **Lagartixa-do-mar-de-Vitória.**

References: Azores (PECS) (MAR-ECO 6267, 2004; Iwamoto and Orlov, 2008; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017).

***Bathygadus favosus* Goode & Bean 1886**

Common name: No English common name; **Lagartixa-do-mar-favosa.**

References: mainland Portugal (MNHN-IC-1988-0562, 1984; MNHN-IC-1988-0556, 1984; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017), Azores (PECS) (Kukuev, 2002; MAR-ECO 9832, 2004; Bergstad *et al.*, 2008a; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017) and Madeira (NHMUK 1995.8.4.61, 1972 (PECS); Marshall, 1979; Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Iwamoto, 2016a; Froese and Pauly, 2019). This species is often confused with *Bathygadus melanobranchus* Vaillant 1888.

***Bathygadus melanobranchus* Vaillant 1888**

Common name: **Vaillant's grenadier; Lagartixa-do-mar-de-Vaillant**, Peixe-rato, Rato.

References: mainland Portugal (Marshall, 1979; Cohen *et al.*, 1990; Geistdoerfer, 1990; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Froese and Pauly, 2019), Azores (Collett, 1896a; Brauer, 1906; Richard, 1900, 1910; MOM-IC-0091-1513, 1912; Roule, 1919a, b; Roule and Angel, 1933; Fowler, 1936a; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Grey, 1956; Marshall and Iwamoto, 1973; Geistdoerfer, 1986; Cohen *et al.*, 1990; Arruda, 1992, 1997; Santos *et al.*, 1997; Melo and Menezes, 2002; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Roule and Angel, 1933; Fowler, 1936b; Maul, 1951a; Albuquerque, 1954-1956; Carneiro *et al.*, 2014; Iwamoto, 2016b). This species is often confused with *Bathygadus favosus* Goode & Bean 1886. The holotype (unique) of *Bathygadus vaillanti* Roule & Angel 1933, synonym of *B. melanobranchus*, caught off North of Hyères Seamount (32°23.0'N, 28°23.50'W), South Azores is in the MOM collections (MOM-IC-0091-1513).

***Cetonurus globiceps* (Vaillant 1884)**

Common name: Globehead grenadier; **Lagartixa-do-mar-de-cabeça-redonda**, Lagartixa, Peixe-lagartixa, Peixe-rato, Rata, Rato.

References: mainland Portugal (Geistdoerfer, 1986; Carneiro *et al.*, 2014; Iwamoto, 2015c), Azores (MNHN-IC-1886-0091, 1883; Vaillant, 1888, 1919; Goode and Bean, 1895, 1896; Murray and Hjort, 1912; Fowler, 1936a; Albuquerque, 1954-1956; Grey, 1956(?); Osório de Castro, 1967; Bauchot *et al.*, 1971; Marshall and Iwamoto, 1973; Marshall, 1979; Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Iwamoto, 2015c; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967; Geistdoerfer, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Iwamoto, 2015c).

***Coelorinchus caelorrhincus* (Risso 1810)**

Common name: Hollowsnout grenadier; **Lagartixa-do-mar-de-focinho-côncavo**, Lagartixa, Lagartixa-do-mar, Pagra, Peixe-furão; Praia, Rato-bicudo.

References: mainland Portugal (Osório, 1909, 1917; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Marshall, 1979; Fernández *et al.*, 1981; Fischer *et al.*, 1981; INIP, 1982b; Geistdoerfer, 1986, 1990; Cohen *et al.*, 1990; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Chaves, 2008; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; OCEANA, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a; Vaillant, 1888, 1919; Collett, 1890a; Goode and Bean, 1895, 1896; Regan, 1903a; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Bauchot *et al.*, 1971; Marshall and Iwamoto, 1973; Dias *et al.*, 1976; Wheeler, 1978; Shcherbachov *et al.*, 1985a; Cohen *et al.*, 1990; Geistdoerfer, 1990; Dias, 1991; Arruda, 1992, 1997; Silva *et al.*, 1995; Menezes, 1996; INIP, 1985a, b; Morato-Gomes *et al.*, 1998; Kukuev, 2002 (PECS); Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1839, 1840b, 1844a, 1849; Johnson, 1862a, b; Günther, 1862, 1887; Vinciguerra, 1883; Collett, 1890b; Goode and Bean, 1895, 1896; NHMUK 1917.7.14.85., 1912 - 1936; Osório, 1917; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Maul, 1951a, 1976 (PECS); Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Fischer *et al.*, 1981; INIP, 1982c, 1984a, b, c; Carneiro *et al.*, 2014; Iwamoto, 2016b). The holotype (unique) of *Macrourus atlanticus* Lowe 1839, synonym of *C. caelorrhincus*, caught off in Madeira is in the NHMUK collections (NHMUK 1917.7.14.85). One specimen caught off at the Josephine Bank is in the MMF collections (MMF 22411, 1967).

***Coelorinchus labiatus* (Köhler 1896)**

Common name: Spearsnouted grenadier; **Lagartixa-do-mar-de-focinho-comprido**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato.

References: mainland Portugal (Cohen *et al.*, 1990; Carneiro *et al.*, 2014; Fernandes *et al.*, 2015a; Bañón *et al.*, 2016a; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019), Azores (Vaillant, 1887, 1888, 1919; Brauer, 1906; Roule, 1916, 1919b; Koefoed, 1927; Fowler, 1936a; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Bauchot *et al.*, 1971; Marshall and Iwamoto, 1973; Cohen *et al.*, 1990; Arruda, 1992, 1997; Saldanha and Biscoito, 1997; Santos *et al.*, 1997; Desbruyères *et al.*, 2001; Melo and Menezes, 2002; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2013, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Fernandes *et al.*, 2015a; Mecklenburg *et al.*, 2018; Almeida and Biscoito, 2019;

Froese and Pauly, 2019) and Madeira (Cohen *et al.*, 1990; Fernandes *et al.*, 2015a; Iwamoto, 2016b). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. labiatus* is the Eastern Atlantic. The distribution maps of this species reported by Fernandes *et al.* (2015) and Iwamoto (2016b) partially included the area of Selvagens Islands (Madeira EEZ). However, specimens are needed for confirmation.

***Coelorinchus occa* (Goode & Bean 1885)**

Common name: **Swordsnout grenadier; Lagartixa-do-mar-de-focinho-curto**, Peixe-rato, Rato.

References: mainland Portugal (Marshall, 1979; Geistdoerfer, 1986, 1990; Borges *et al.*, 2002; Carneiro *et al.*, 2014), Azores (Vaillant, 1888; MOM-IC-0091-0607, 1902; MOM-IC-0091-0637, 1902; Roule, 1916; Marshall, 1979; Shcherbachov *et al.*, 1985b (PECS); Arruda, 1997; Menezes *et al.*, 2006; Pakhorukov, 2008 (PECS); Carneiro *et al.*, 2014). The syntype of *Coelorhynchus vaillanti* Roule 1916, synonym of *C. occa*, caught off in the Azores is in the MOM collections (MOM-IC-0091-0637).

***Coelorinchus parallelus* (Günther 1877)**

Common name: **Spiny grenadier; Lagartixa-do-mar-espinhosa**, Lagartixa, Lagartixa-do-mar.

References: Azores (?) (Zugmayer, 1933; Collins, 1954; Arruda, 1997). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. parallelus* is the Indo-West Pacific: Japan to Taiwan, East China Sea, South China Sea and South-West Pacific. The references of this species for the Azores may refer eventually to another species of the genus *Coelorinchus* and as a consequence this species is not considered in the total number of valid species of the Azores.

***Coryphaenoides armatus* (Hector 1875)**

Common name: **Abyssal grenadier; Lagartixa-do-mar-abissal**, Lagartixa, Lagartixa-de-natura, Lagartixa-do-mar, Peixe-rato, Rato.

References: mainland Portugal (Geistdoerfer, 1986; Cohen *et al.*, 1990; Davies and Merrett, 1998; King *et al.*, 2008; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014; Fernandes *et al.*, 2015b), Azores (Brauer, 1906(?); Brian, 1912; Murray and Hjort, 1912; Roule, 1916, 1919; Vaillant, 1919; Fowler, 1936a; Nybelin, 1951, 1957; Collins, 1954; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Geistdoerfer *et al.*, 1971; Marshall and Iwamoto, 1973; Geistdoerfer, 1986; Cohen *et al.*, 1990; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Fernandes *et al.*, 2015b; Almeida and Biscoito, 2019) and Madeira (Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Wolff, 1971; MNHN-IC-2008-2387, 1981; NHMUK 1991.7.9.868, 1989; Cohen *et al.*, 1990; Armstrong *et al.*, 1992; Freitas *et al.*, 2013; Carneiro *et al.*, 2014; Fernandes *et al.*, 2015b).

***Coryphaenoides brevibarbis* (Goode & Bean 1896)**

Common name: Shortbeard grenadier; **Lagartixa-do-mar-de-barbilho-curto**.

References: Azores (Marshall, 1979; Bauchot and Pras, 1980; Geistdoerfer, 1986; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019) and Madeira (Porteiro *et al.*, 2017).

***Coryphaenoides carapinus* Goode & Bean 1883**

Common name: Carapine grenadier; **Lagartixa-do-mar-carapina**, Lagartixa, Peixe-rato, Rato.

References: mainland Portugal (GNM Pisces 1946-7667, 1946; Marshall, 1979; Geistdoerfer, 1986, 1990; Carneiro *et al.*, 2014), Azores (Roule and Angel, 1933; Fowler, 1936a; Schroeder, 1940; Collins, 1954; Albuquerque, 1954-1956; Geistdoerfer *et al.*, 1971; Grey, 1956; Osório de Castro, 1967; Marshall and Iwamoto, 1973; Geistdoerfer, 1986; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019) and Madeira (Wolff, 1971; Geistdoerfer, 1986; Carneiro *et al.*, 2014).

***Coryphaenoides guentheri* (Vaillant 1888)**

Common name: **Günther's grenadier; Lagartixa-do-mar-de-Günther**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato.

References: mainland Portugal (NHMUK 1887.12.7.84, 1873; Grey, 1956; Osório de Castro, 1967; Cohen *et al.*, 1990; Geistdoerfer, 1986, 1990; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Iwamoto, 2016b; Froese and Pauly, 2019), Azores (Collett, 1896a; Brauer, 1906; Richard, 1900, 1910; Roule, 1919a;

Koefoed, 1927; Fowler, 1936; Collins, 1954; Grey, 1956; Osório de Castro, 1967; Geistdoerfer and Rannou, 1971; Marshall and Iwamoto, 1973; Cohen *et al.*, 1990; Arruda, 1992, 1997; Santos *et al.*, 1997; Melo and Menezes, 2002; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967; Cohen *et al.*, 1990; Nedreaas *et al.*, 2015b). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. guentheri* is the Greenland, eastern-north Atlantic and Mediterranean Sea. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. The distribution maps of this species reported by Nedreaas *et al.* (2015b) partially included the area of Selvagens Islands (Madeira EEZ). However, specimens are needed for confirmation.

***Coryphaenoides leptolepis* Günther 1877**

Common name: Ghostly grenadier; **Lagartixa-do-mar-pálida**, Lagartixa, Peixe-rato, Rato.

References: mainland Portugal (IFREMER Biocean 34969, 1969; Geistdoerfer, 1990; Carneiro *et al.*, 2014; Fernandes *et al.*, 2015c), Azores (Murray and Hjort, 1912; Roule 1919a, b(?); Fowler, 1936a; Nybelin, 1951, 1957; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Marshall and Iwamoto, 1973; Geistdoerfer, 1986; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a, b; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Fernandes *et al.*, 2015c) and Madeira (Davies and Merrett, 1998; Freitas *et al.*, 2013).

***Coryphaenoides mediterraneus* (Giglioli 1893)**

Common name: Mediterranean grenadier; **Lagartixa-do-mar-do-Mediterrâneo**, Peixe-rato, Rato.

References: mainland Portugal (GNM 1946-6456, 1946; GNM 1946-6458, 1946; Nybelin, 1948; Geistdoerfer, 1986; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Iwamoto, 2016b) and Azores (Roule, 1919a(?); Geistdoerfer and Rannou, 1971, 1972; Marshall and Iwamoto, 1973(?); Marshall, 1979; Bauchot and Pras, 1980; Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019).

***Coryphaenoides profundiculus* (Nybelin 1957)**

Common name: Deepwater grenadier; **Lagartixa-do-mar-da-fundura**.

References: mainland Portugal (NHMUK 1994.9.20.5, 1983; NHMUK 1991.7.9.865-867, 1989; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014), Azores (Murray and Hjort, 1912) and Madeira (Geistdoerfer, 1986; NHMUK 1991.7.9.871-872, 1989; NHMUK 1991.7.9.796, 1990 (PECS); Lloris *et al.*, 1991; Davies and Merrett, 1998; Carneiro *et al.*, 2014; Iwamoto, 2016b).

***Coryphaenoides rutilus* Günther 1878**

Common name: Madeiran grenadier; **Lagartixa-do-mar-cabeçuda**.

References: References: mainland Portugal (Iwamoto, 2015d), Azores (Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira (Maul, 1951a; Marshall, 1979; Cohen *et al.*, 1990; Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Iwamoto, 2015d; Froese and Pauly, 2019). The holotype of *Nematonurus macrocephalus* Maul 1951, synonym of *C. rutilus*, caught off in Madeira is in the MMF collections (MMF 898).

***Coryphaenoides rupestris* Gunnerus 1765**

Common name: Roundnose grenadier; **Lagartixa-da-rocha**.

References: mainland Portugal (Cohen *et al.*, 1990; Fernandes *et al.*, 2007; Bordalo-Machado *et al.*, 2009; OCEANA, 2014; Cook *et al.*, 2015; Bañón *et al.*, 2016a; Iwamoto, 2016b; Porteiro *et al.*, 2017; Froese and Pauly, 2019), Azores[▲] (IMAR/DOP COLETA 1690, 2001; Hareide and Garnes, 2001 (PECS); Melo and Menezes, 2002; Bergstad *et al.*, 2010 (PECS); Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014) and Madeira (NHMUK 1995.7.25.17, 1972 (PECS); Carneiro *et al.*, 2014; Porteiro *et al.*, 2017).

***Coryphaenoides thelestomus* Maul 1951**

Common name: Roughlip grenadier; **Lagartixa-do-mar-da-Madeira**, Lagartixa.

References: Madeira (Maul, 1951a, 1952b; Albuquerque, 1954-1956; Osório de Castro, 1967; Marshall, 1979; Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Iwamoto, 2016b; Froese and

[Pauly, 2019](#)). The holotype (unique) of this species caught off in Madeira is in the MMF collections obtained in the Funchal fish market. A second specimen is in the MMF collections (MMF 41365, 2010).

Coryphaenoides zaniophorus (Vaillant 1888)

Common name: **Thickbeard grenadier; Lagartixa-do-mar-de-barbilho-grosso**, Peixe-lagartixa.

References: mainland Portugal ([Geistdoerfer, 1986](#); [Carneiro et al., 2014](#)) and Madeira ([Geistdoerfer, 1986](#); [Cohen et al., 1990\(?\)](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Biscoito, 2015d](#)).

Echinomacrurus mollis Roule 1916

Common name: No English common name; **Lagartixa-do-mar-macia**.

References: mainland Portugal ([NHMUK 1993.9.13.10-19, 1989](#); [Christiansen et al., 2014](#)), Azores (PECS) ([GNM 1948-7017, 1948](#); [GNM 1948-7018, 1948](#)) and Madeira ([Geistdoerfer, 1986](#); [CAS ICH 83371, 1995](#) (PECS); [Carneiro et al., 2014](#)).

Gadomus arcuatus (Goode & Bean 1886)

Common name: Doublethread grenadier; **Lagartixa-do-mar-filamentosa**.

References: mainland Portugal ([Geistdoerfer, 1986, 1990](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Azores ([Geistdoerfer, 1990](#); [IMAR/DOP COLETA 1691, 2001](#); [Melo and Menezes, 2002](#); [Porteiro et al., 2010](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Delgado et al., 2017](#)).

Gadomus dispar (Vaillant 1888)

Common name: No English common name; **Lagartixa-do-mar-ímpar**.

References: mainland Portugal ([Saldanha et al., 1995](#); [Marques and Saldanha, 1998](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#)) and Azores ([Melo and Menezes, 2002](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)).

Gadomus longifilis (Goode & Bean 1885)

Common name: Threadfin grenadier; **Lagartixa-do-mar-de-bandeira**, Peixe-rato, Rato.

References: mainland Portugal ([Roule, 1919a](#); [Grey, 1956](#); [Maul, 1976](#); [Marshall, 1979](#); [Geistdoerfer, 1986, 1990](#); [Fernandes et al., 2007](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Iwamoto, 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1896a](#); [Richard, 1900, 1910](#); [Brauer, 1906](#); [Richard, 1910](#); [Roule, 1919a](#); [Koefoed, 1927](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Marshall and Iwamoto, 1973](#); [Marshall, 1979](#); [Geistdoerfer, 1990](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Iwamoto, 2016a](#)); Several specimens caught off in 1967 at the coast of the mainland Portugal are in the MMF collections (MMF 22476, 1967, set of sixteen specimens). One specimen caught off in October 2010 at the Portuguese continental slope is in the MB collections.

Hymenocephalus italicus Giglioli 1884

Common name: **Glasshead grenadier; Lagartixa-do-mar-prateada**, Lagartixa, Lagartixa-do-mar, Lagartixa-prateada, Peixe-rato, Rato.

References: mainland Portugal ([Osório, 1888](#); [Nobre, 1903, 1935](#); [Osório de Castro, 1967](#); [Grey, 1956](#); [Marshall, 1979](#); [Fernández et al., 1981](#); [INIP, 1982a, b](#); [Geistdoerfer, 1986, 1990](#); [Fischer et al., 1987](#); [Cohen et al., 1990](#); [Figueiredo et al., 1994](#); [Borges et al., 2002](#); [Santos et al., 2002b](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Silva and Borges, 2014](#); [Nedreaas et al., 2015c](#); [Iwamoto, 2016b](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1886-0056, 1883](#); [Vaillant, 1888](#); [Collett, 1896a](#); [Brauer, 1906](#); [Fowler, 1936a](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [Bauchot et al., 1971](#); [Marshall and Iwamoto, 1973](#); [Maul, 1976](#); [Saldanha, 1977a](#); [Cohen et al., 1990](#); [Geistdoerfer, 1990](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Nedreaas et al., 2015c](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-1883-0120, 1882](#); [Osório de Castro, 1967](#); [Cohen et al., 1990](#); [Carneiro et al., 2014](#); [Nedreaas et al., 2015c](#); [Froese and Pauly, 2019](#)).

Hymenogadus gracilis (Gilbert & Hubbs 1920)

Common name: Graceful grenadier; **Lagartixa-do-mar-elegante**.

References: mainland Portugal ([ENA MG717680.1, 2018](#); [ENA MG717681.1, 2018](#); [ENA MG717682.1, 2018](#)), Azores (PECS) ([Maul, 1976](#); [Ehrich, 1977](#); [Shcherbachov et al., 1985b](#); [Fock et al., 2002](#); [Kukuev, 2002](#); [Pakhorukov, 2008](#); [Carneiro et al., 2014](#)). The northernmost occurrence record of *H. gracilis* (38°0.0'N, 9°1.0'W) was reported in 2017 [European Nucleotide Archive (EMBL-EBI)]. Several specimens caught off at the Meteor Bank are in the MMF collections (MMF 22844, 1967, set of four specimens; MMF 22872, 1967, set of five specimens and MMF 22836, 1967, set of nine specimens).

***Macrourus berglax* Lacepède 1801**

Common name: **Roughhead grenadier**; **Lagartixa-do-mar-olho-de-cebola**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato

References: mainland Portugal ([Porteiro et al., 2017](#)), Azores (PECS) ([Porteiro et al., 2017](#)) and Madeira ([Porteiro et al., 2017](#)).

***Malacocephalus laevis* (Lowe 1843)**

Common name: **Softhead grenadier**; **Lagartixa-do-mar-redonda**, Batage, Lagartixa, Lagartixa-do-mar, Peixe-rato, Ratage, Rato, Rato-redondo.

References: mainland Portugal ([Osório, 1888, 1909](#); [Vieira, 1893, 1897, 1900](#); [Nobre, 1894, 1935](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Marshall, 1979](#); [INIP, 1981a, b, 1982a, b](#); [Fernández et al., 1981](#); [Martins, 1981](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Geistdoerfer, 1986, 1990](#); [Cohen et al., 1990](#); [Figueiredo et al., 1994](#); [Saldanha et al., 1995](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002b](#); [Moura, 2004](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Pakhorukov, 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [Costa, 2014](#); [OCEANA, 2014](#); [Silva and Borges, 2014](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Marshall and Iwamoto, 1973](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [INIP, 1985a, b](#); [Shcherbachov et al., 1985b \(PECS\)](#); [Cohen et al., 1990](#); [Dias et al., 1990](#); [Dias, 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Melo and Menezes, 2002](#); [Menezes et al., 2006, 2012](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1843, 1844b](#); [NHMUK 1855.11.29.4, 1854-1861](#); [Johnson, 1862b](#); [Günther, 1862](#); [de Brito Capello, 1872](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Maul, 1951a, 1976](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Ehrich, 1977](#); [Fischer et al., 1981\(?\)](#); [Cohen et al., 1990](#); [Carneiro et al., 2014](#); [POGID, 2016](#); [Roberts et al., 2017](#)). The holotype (unique) of *Macrourus laevis* Lowe 1843, synonym of *Malacocephalus laevis*, caught off in Madeira is in the NHMUK collections (NHMUK 1855.11.29.4). One specimen caught at the Meteor Bank is in the MMF collections (MMF 22397, 1967).

***Malacocephalus occidentalis* Goode & Bean 1885**

Common name: **Western softhead grenadier**; **Lagartixa-do-mar-occidental**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato.

References: Madeira (?) ([Iwamoto, 2016b](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *M. occidentalis* is the Western and Eastern Atlantic. The distribution maps of this species reported by Iwamoto (2016b) partially included the area of Selvagens Islands (Madeira EEZ). The occurrence of this species in this region is not yet confirmed and specimens are needed. Thus, this species is not considered in the total number of valid species.

***Nezumia aequalis* (Günther 1878)**

Common name: **Common Atlantic grenadier**; **Largatixa-do-mar-lisa**, Lagartixa, Lagartixa-do-mar, Lagartixa-lisa, Imperador, Imperador-de-natura, Peixe-rato, Rato, Rato-redondo.

References: mainland Portugal ([NHMUK 1887.12.7.85, 1873](#); [NHMUK 1887.12.7.86, 1873](#); [Günther, 1878](#); [Vinciguerra, 1883](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [Geistdoerfer, 1986, 1990](#); [Fischer et al., 1987](#); [Cohen et al., 1990](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1896a](#); [Brauer, 1906](#); [Roule, 1919a](#); [Vaillant, 1919](#); [Koefoed, 1927](#); [Nobre, 1935](#); [Fowler, 1936a](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Osório de Castro, 1967](#); [Marshall and Iwamoto, 1973](#); [Maul, 1976](#);

Marshall, 1979; Fischer *et al.*, 1981; Geistdoerfer, 1986, 1990; Cohen *et al.*, 1990; Lloris *et al.*, 1991; Arruda, 1992, 1997; Hamer *et al.*, 1994; Silva *et al.*, 1995; Menezes, 1996; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Melo and Menezes, 2002; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Almeida and Bischoff, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1843, 1844b; Günther, 1862; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Maul, 1951a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Maul, 1976; Fischer *et al.*, 1981; Geistdoerfer, 1986; Cohen *et al.*, 1990; Lloris *et al.*, 1991; Ferreira *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Iwamoto, 2016b; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Two syntypes of *Coryphaenoides aequalis* Günther 1878 both caught off south of mainland Portugal are in the NHMUK collections (NHMUK 1887.12.7.85 and NHMUK 1887.12.7.86). The whereabouts of the holotype (unique) of the *Macrourus serratus* Lowe 1843 caught off Madeira are unknown. *Coryphaenoides aequalis* is synonym of *N. aequalis* and *M. serratus* is uncertain as *N. aequalis*.

***Nezumia bairdii* (Goode & Bean 1877)**

Common name: **Marlin-spike grenadier**; **Lagartixa-do-mar-de-Baird**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato.

References: mainland Portugal (INIP, 1982a, b; Fernandes *et al.*, 2007; Chaves, 2008; Silva *et al.*, 2009; Silva and Borges, 2014), Azores (Roule and Angel, 1933; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; IFREMER Biocean 34775, 1969; Osório de Castro, 1967; Geistdoerfer *et al.*, 1970(?), 1971; Marshall and Iwamoto, 1973(?); Merrett, 1986; Cohen *et al.*, 1990(?); Arruda, 1997; Santos *et al.*, 1997(?); Carneiro *et al.*, 2014; Froese and Pauly, 2019(?)) and Madeira (?) (Roule and Angel, 1933; Carneiro *et al.*, 2014). According to Froese and Pauly (2019) the report of the specimen from Madeira probably refers to other species; *N. bairdii* is often confused with *Nezumia aequalis* (Günther 1878). Eschmeyer *et al.* (2019) reported the current geographic distribution of this species in the Atlantic, and Möller *et al.* (2010) indicated its occurrence in Greenland waters and elsewhere in the western North Atlantic. Thus, this species is not considered in the total number of valid species of Madeira EEZ.

***Nezumia longebarbata* (Roule & Angel 1933)**

Common name: Bluntnose grenadier; **Lagartixa-do-mar-de-focinho-curto**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato.

References: Madeira (MOM-IC-0091-1875, 1911; Roule and Angel, 1933; Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1967; Marshall, 1979; Geistdoerfer, 1986; Lloris *et al.*, 1991; Moore *et al.*, 2003; Carneiro *et al.*, 2014; Iwamoto, 2016b; Froese and Pauly, 2019).

***Nezumia sclerorhynchus* (Valenciennes 1838)**

Common name: **Roughtip grenadier**; **Lagartixa-do-mar-áspera**, Lagartixa, Lagartixa-áspera, Lagartixa-do-mar, Peixe-rato, Rato.

References: mainland Portugal (Brauer, 1906; MNHN-IC-1912-0044, 1908; Zugmayer, 1911a; NHMUK 1934.12.19.29, 1912-1936; Albuquerque, 1954-1956; Fernández *et al.*, 1981; Fischer *et al.*, 1987; Cohen *et al.*, 1990; Gordo *et al.*, 1990; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Moura, 2004; ICES, 2005; Sousa *et al.*, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Froese and Pauly, 2019), Azores (Vaillant, 1888; Collett, 1896a; Brauer, 1906; Roule, 1919a; Koefoed, 1927; Fowler, 1936a; Collins, 1954; Albuquerque, 1954-1956; Grey, 1956; Bauchot *et al.*, 1971; Marshall and Iwamoto, 1973; Marshall, 1979; Geistdoerfer, 1986, 1990; Cohen *et al.*, 1990; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Desbruyères *et al.*, 2001; Porteiro *et al.*, 2010; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Iwamoto, 2016b; Froese and Pauly, 2019) and Madeira (Geistdoerfer, 1986; Cohen *et al.*, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Iwamoto, 2016b; Froese and Pauly, 2019). Five syntypes of *Macrurus violaceus* Zugmayer 1911, synonym of *N. sclerorhynchus*, caught off south of mainland Portugal are in the NHMUK and MOM collections (NHMUK 1934.12.19.29 and MOM-IC-0091-1026 (set of four specimens)).

***Odontomacrurus murrayi* Norman 1939**

Common name: **Roundhead grenadier**; **Lagartixa-do-mar-de-Murray**, Cobaia, Peixe-rato, Rato.

References: Azores (Maul and Koefoed, 1950 (PECS); Lewis and Elder, 1953; Collins, 1954; Marshall, 1979; Geistdoerfer, 1986; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999,

2010, 2017; Melo and Menezes, 2002; Sutton *et al.*, 2008; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Maul and Koefoed, 1950; Maul, 1951a; Lewis and Elder, 1953; Albuquerque, 1954-1956; Osório de Castro, 1967; Marshall, 1979; Geistdoerfer, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype and two syntypes of *Phalacromacrurus pantherinus* Maul & Koefoed 1950, synonym of *O. murrayi*, caught off 90 NM north-east of Marsala Seamount, 34°59.0'N, 33°1.0'W are in the ZMUB collections (ZMUB 4473) and in the MMF collections, respectively.

***Pseudonezumia flagellicauda* (Koefoed 1927)**

Common name: No English common name; **Lagartixa-do-mar-de-chicote**.

References: mainland Portugal ([GNM 1946-7387](#), 1946; Porteiro *et al.*, 2017) and Azores ([Koefoed, 1927](#) (PECS); [MAR-ECO 6239](#), 2004; [MAR-ECO 6253](#), 2004; Bergstad *et al.*, 2008a (PECS); Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Priede, 2017).

***Sphagmacrurus grenadæ* (Parr 1946)**

Common name: Pugnose grenadier; **Lagartixa-do-mar-de-nariz-arrebitado**.

References: mainland Portugal (?) ([Maul, 1976](#); [Hureau and Monod, 1979](#); [MNHN-IC-1988-0594](#), 1984; [Geistdoerfer, 1986](#); [Carneiro *et al.*, 2014](#)) and Azores (?) (PECS) ([Shcherbachov *et al.*, 1985a](#); [Kukuev, 2002](#); [Carneiro *et al.*, 2014](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. grenadæ* is the Western Atlantic, from Gulf of Mexico to Brazil. This species is closely related to *Sphagmacrurus hirundo* (Collett 1896) (in the Eastern Atlantic) and *Sphagmacrurus pumiliceps* (Alcock 1894) (in Indian Ocean). One specimen caught off at the coast of the mainland Portugal is in the MMF collections (MMF 22477, 1967). However, as recommended by Froese and Pauly (2019), it is advisable a thorough study to distinguish these three species and re-evaluate the specimens identified as *S. grenadæ* caught off Eastern Atlantic. Consequently, this species is not considered in the total number of valid species for mainland Portugal and Azores.

***Sphagmacrurus hirundo* (Collett 1896)**

Common name: Swallow grenadier; **Lagartixa-do-mar-dos-Açores**, Lagartixa, Lagartixa-do-mar, Peixe-rato, Rato.

References: Azores ([MOM-IC-0091-133](#), 1888; Collett, 1896a; Richard, 1900, 1910; Brauer, 1906; Roule, 1919a, 1919b; Fowler, 1936a; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Marshall, 1979; Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; González *et al.*, 2014; Iwamoto, 2016b; Almeida and Biscoito, 2019; Frose and Pauly 2019) and Madeira (Geistdoerfer, 1986, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; González *et al.*, 2014; Iwamoto, 2016b; Frose and Pauly 2017; Porteiro *et al.*, 2017; Frose and Pauly 2019). One syntype of *Macrurus hirundo* Collett 1896, synonym of *S. hirundo*, caught off between Pico and São Jorge (Azores) is in the MOM collections (MOM-IC-0091-0133).

***Squalogadus modificatus* Gilbert & Hubbs 1916**

Common name: Tadpole whiptail; **Lagartixa-do-mar-girino**.

References: Azores (PECS) ([MAR-ECO 2873](#), 2004; Bergstad *et al.*, 2008a; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017).

***Trachonurus sulcatus* (Goode & Bean 1885)**

Common name: Bristly grenadier; **Largartixa-do-mar-arrepiada**, Lagartixa, Lagartixa-negra.

References: mainland Portugal ([Cohen *et al.*, 1990](#); [Carneiro *et al.*, 2014](#)) and Azores ([MNHN-IC-1886-0118](#), 1883; [MNHN-IC-1886-0187](#), 1883; [Vaillant, 1888](#); [Brauer, 1906](#); [Roule, 1916, 1919b](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [ZMH 108614, 1971](#) (PECS); [MNHN-IC-2009-0403](#), 1971; [Carneiro *et al.*, 2014](#); [Almeida and Biscoito, 2019](#); [Frose and Pauly 2019](#)). The lectotype of *Coryphaenoides asperimus* Vaillant 1888, synonym of *T. sulcatus*, caught off in the Azores is in the MNHN collections (MNHN-IC-1886-0118).

***Trachonurus villosus* (Günther 1877)**

Common name: Furry whiptail; **Lagartixa-do-mar-peluda**, Lagartixa-negra, Peixe-rato, Rato.

References: mainland Portugal (?) ([Carneiro et al., 2014](#)), Azores ([Vaillant, 1888](#); [Brauer, 1906](#); [Roule, 1916, 1919](#); [Fowler, 1936](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bauchot et al., 1971](#); [Marshall and Iwamoto, 1973](#); [Marshall, 1979](#); [Geistdoerfer, 1986, 1990](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1995.7.11.481, 1972 \(PECS\)](#); [Geistdoerfer, 1986, 1990](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The occurrence record of the specimen deposited in the History Museum (London), catalogue number 1995.7.11.481, mentions as local of catching 266 NM south-west of Cape São Vicente, (mainland Portugal). The geographic coordinates of this records corresponds to a position off Madeira, in the PECS of Madeira and was erroneously considered present in mainland Portugal by Carneiro et al., (2014). Thus, this species is not considered in the total number of valid species for mainland Portugal.

***Trachyrincus scabrus* (Rafinesque 1810)**

Common name: **Roughsnout grenadier; Peixe-lima**, Besouro, Besouro, Bezouro, Furão, Lima, Peixe-furão. References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Steindachner, 1868b](#); [Vinciguerra, 18883](#); [Goode and Bean, 1895, 1896](#); [Bragança, 1903](#); [MNHN-IC-1905-0582, 1903](#); [MNHN-IC-1905-0583, 1903](#); [Nobre, 1903, 1904, 1935](#); [Richard, 1904](#); [Osório, 1909, 1917](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Marshall, 1979](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a](#); [Martins, 1981](#); [Geistdoerfer, 1986, 1990](#); [Cohen et al., 1990](#); [Gordo et al., 1990](#); [Figueiredo et al., 1994](#); [Borges et al., 2002, 2003](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2006](#); [Fernandes et al., 2007](#); [King et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Bordalo-Machado et al., 2009](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967](#); [Kukuev, 2002 \(PECS\)](#)) and Madeira ([Osório de Castro, 1967](#); [Nedreaas et al., 2015d](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *T. scabrus* is the Mediterranean Sea and, in the Eastern Atlantic, from off Scotland to Cape Verde Islands. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores and Madeira without indicating the geographic location. The distribution map of this species reported by Nedreaas et al. (2015d) partially included the area of Selvagens Islands (Madeira EEZ). Specimens are needed for confirmation in Madeira and Azores archipelagos.

Family Moridae

***Antimora rostrata* (Günther 1878)**

Common name: **Blue antimora; Mora-azul**.

References: mainland Portugal ([NHMUK 1996.9.11.9, 1974](#), [Carneiro et al., 2014](#); [Nielsen et al., 2015e](#); [Bañón et al., 2016a](#)), Azores ([Trunov, 1992 \(PECS\)](#); [Martins et al., 2006](#); [Porteiro et al., 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Nielsen et al., 2015e](#); [Almeida and Biscoito, 2019](#)) and Madeira ([Freitas et al., 2013](#); [Nielsen et al., 2015e](#)).

***Eretmophorus kleinenbergi* Giglioli 1889**

Common name: No English common name; **Peixe-remo-de-Kleinenberg**.

References: Azores (PECS) ([ZMH 121033, 1982](#); [Carneiro et al., 2014](#)).

***Gadella imberbis* (Vaillant 1888)**

Common name: Beardless codling; **Abrótea-imberbe**, Laibeque.

References: mainland Portugal ([Biscoito and González, 2017](#)).

***Gadella maraldi* (Risso 1810)**

Common name: **Gadella; Abrótea-de-natura**, Abrótia, Abrótia-de-natura, Gadela.

References: mainland Portugal ([Bragança, 1902](#); [Osório, 1917](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [INIP, 1982a, b](#); [Cohen, 1986, 1990c](#); [Fischer et al., 1987](#); [Cohen et al., 1990](#); [Saldanha et al., 1995](#); [Borges et al., 2002, 2003](#); [Moura, 2004](#); [ICES, 2005](#); [González et al., 2008](#); [Silva, 2008, 2011](#); [Silva et al., 2009](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Silva and Borges, 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Regan, 1903a](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977 \(PECS\)](#); [Aguiar and Pereira, 1982](#); [Cohen, 1986 \(PECS\)](#); [Cohen et al., 1990](#); [Silva and Menezes, 1995a](#); [Menezes, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sousa et al., 1999](#); [Fock et al., 2002 \(PECS\)](#); [Kukuev, 2002 \(PECS\)](#); [Menezes et al., 2006](#); [González et al., 2008](#); [Porteiro et al., 2010](#); [Carneiro](#)

et al., 2014; Iwamoto and Cohen, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1840a, 1841, 1843, 1844b, 1849; Günther, 1862; Johnson, 1862b; Goode and Bean, 1895, 1896; NHMUK 1917.7.14.84, 1912-1936; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Maul, 1952a, 1976 (PECS); Albuquerque, 1954-1956; Ehrich, 1977; Cohen, 1979d, 1986, 1990c; Cohen *et al.*, 1990; Lloris *et al.*, 1991; González *et al.*, 2008; Christiansen *et al.*, 2009; Carneiro *et al.*, 2014; Iwamoto and Cohen, 2016; POGID, 2016; Biscoito *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Two unique holotypes of *Gadus gracilis* Lowe 1843 and *Merluccius ambiguus* Lowe 1841 were caught off Madeira. The holotype *G. gracilis* is in the NHMUK collections (NHMUK 1917.7.14.84) and the other one whereabouts are unknown. One specimen caught off in June 2009 at the Portuguese continental slope is in the MB collections (MB06-005171). Several specimens caught off at Josephine Bank (MMF 22802, 1967, set of two specimens and MMF 22849, 1967, set of seven specimens) and Meteor Bank (MMF 22870, 1967, one specimen) are in the MMF collections.

Guttigadus latifrons (Holt & Byrne 1908)

Common name: No English common name; **Abrótea-cabeçuda**, Abrótea-de-natura.

References: mainland Portugal (Moura, 2004; ICES, 2005) and Azores (Saldanha and Biscoito, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017 (PECS); Carneiro *et al.*, 2014; Bañón *et al.*, 2016a).

Halargyreus johnsonii Günther 1862

Common name: **Slender codling**; **Abrótea-delgada**, Abrótea-de-natura.

References: mainland Portugal (Girard, 1895; AVG CODC-Peixes PE/322, 1894; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Fernández *et al.*, 1981; Carneiro *et al.*, 2014), Azores (Trunov, 1992 (PECS); IMAR/DOP COLETA 1698, 2001, 2001; Melo and Menezes, 2002; Bergstad *et al.*, 2008a (PECS); Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Porteiro *et al.*, 2010, 2017; Almeida and Biscoito, 2019; Froese and Pauly, 2019 (PECS)) and Madeira (NHMUK 1861.6.4.2., 1854-1861; Günther, 1862; Goode and Bean, 1895, 1896; Brauer, 1906; Fowler, 1936a; Maul, 1952a; Osório de Castro, 1967; Cohen, 1979d, 1986; Cohen *et al.*, 1990; Lloris *et al.*, 1991; González *et al.*, 2008; Carneiro *et al.*, 2014; Iwamoto and Cohen, 2016; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Madeira (stomach content) is in the NHMUK collections (NHMUK 1861.6.4.2).

Laemonema robustum Johnson 1862

Common name: Robust cod; **Abrótea-robusta**, Abrótea, Abrótea-de-natura, Abrótia-de-natura.

References: Azores (IMAR/DOP 3SMI-02-P11MM06, 2011; Ralph, 2015j) and Madeira (NHMUK 1862.6.14.15, 1861-1890; Günther, 1862; Johnson, 1862a; de Brito Capello, 1872; Goode and Bean, 1895, 1896; Brauer, 1906; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1952a; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Cohen, 1979d, 1986; Smith and McKay, 1986; Lloris *et al.*, 1991; Meléndez and Markle, 1997; Ferreira *et al.*, 2008; Carneiro *et al.*, 2014; Ralph, 2015j; Iwamoto and Cohen, 2016; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Madeira is in the NHMUK collections (NHMUK 1862.6.14.15).

Laemonema yarrellii (Lowe 1838)

Common name: **Codling**; **Abrótea-de-Yarrell**, Abrótea, Abrótea-de-natura, Abrótea-do-alto, Abrótia, Abrótia-de-natura, Abrótia-do-alto, Bordião.

References: mainland Portugal (OCEANA, 2014); Azores (PECS) (Maul, 1976; Ehrich, 1977; Cohen, 1986, 1990c; Meléndez and Markle, 1997; Fock *et al.*, 2002; Kukuev, 2002; Carneiro *et al.*, 2014; Iwamoto and Cohen, 2016; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1843-1860; NHMUK 1855.12.26.482, 1854-1861; Günther, 1862; Goode and Bean, 1895, 1896; Brauer, 1906; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Ehrich, 1977; Cohen, 1979d, 1986; Biscoito and Maul, 1989; Lloris *et al.*, 1991; Meléndez and Markle, 1997; Carneiro *et al.*, 2014; Iwamoto and Cohen, 2016; Froese and Pauly, 2019). Two unique holotypes of *Phycis yarrellii* Lowe 1838 and *Laemonema curtipes* Biscoito & Maul 1989 are in the NHMUK and MMF collections (NHMUK 1855.12.26.482 and MMF 23858), respectively. *P. yarrellii* was caught off in Madeira and *L. curtipes* at Seine Seamount (33°45.8'N, 14°20.6'W) both are synonyms of *L. yarrellii*. Two specimens caught off at the Meteor Bank are in the MMF collections (MMF 22386, 1967 and MMF 22543, 1967).

***Lepidion guentheri* (Giglioli 1880)**

Common name: Guenther's codling; **Abrótea-de-Günther**, Abrótea, Abrótea-de-natura, Abrótia, Abrótia-de-natura, Juliana-pequena.

References: mainland Portugal (Goode and Bean, 1895, 1896; Brauer, 1906; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Cohen, 1979d; Gordo et al., 1990; González et al., 2008; Bordalo-Machado et al., 2009; Carneiro et al., 2014; Bañón et al., 2016a; Froese and Pauly, 2019), Azores (Vaillant, 1919; Collins, 1954; Templeman, 1970a, b; Cohen, 1979d, 1986; Gordo et al., 1990; Lloris et al., 1991; Stefanescu et al., 1991; Arruda, 1997; Santos et al., 1997; Melo and Menezes, 2002; Menezes et al., 2006, 2009, 2012; González et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Bañón et al., 2016a; Iwamoto and Cohen, 2016; Froese and Pauly, 2019) and Madeira (NHMUK 1862.4.22.9, 1861-1890; Günther, 1862, 1887; Johnson, 1862b; Giglioli, 1880; Goode and Bean, 1895, 1896; Brauer, 1906; Nobre, 1935; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1952a; Albuquerque, 1954-1956; Osório de Castro, 1967; Cohen, 1979d, 1986; Gordo et al., 1990; Lloris et al., 1991; González et al., 2008; Carneiro et al., 2014; Bañón et al., 2016a; Iwamoto and Cohen, 2016; Porteiro et al., 2017; Froese and Pauly, 2019). The holotype (unique) of *Haloporphyrus guentheri* Giglioli 1880, synonym of *L. guentheri*, caught off in Madeira is in the NHMUK collections (NHMUK 1862.4.22.9).

***Lepidion lepidion* (Risso 1810)**

Common name: **Mediterranean codling; Abrótea-do-Mediterrâneo**.

References: Azores (Richard, 1934; Guiart, 1935; Silva et al., 1995; Menezes, 1996; Santos et al., 1997; Menezes et al., 2006, 2009, 2012; González et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014) and Madeira (Günther, 1862; Johnson, 1862b; Vinciguerra, 1883(?); NHMUK 2015.11.5.10, 1986). According to Barros-García et al. (2016) *Lepidion eques* (Günther 1887) is a junior synonym of this species.

***Lepidion schmidti* Svetovidov 1936**

Common name: Giant lepidion; **Abrótea-de-Schmidt**.

References: Azores (PENN-C 1343, n.d.; IFREMER Biocean 49909, 1997; Saldanha and Biscoito, 1997; ZMH 121025, 1982; Desbruyères et al., 2001; Porteiro et al., 2010, 2017).

***Mora moro* (Risso 1810)**

Common name: **Common mora; Mora**, Abrótea-do-alto, Abrótia-do-alto, Busio-do-alto; Buzia, Búzia, Escamuda-branca, Juliana, Mariquita, Melga, Olhudo, Pimentel, Robaldo, Robaldo-branco, Robalo, Robalo-branco, Salmonete-preto.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; MNHN-IC-A-4798, 1881; Vaillant, 1888; Goode and Bean, 1895, 1896; Almeida and Roquette, 1892; Bragança, 1903; Richard, 1905b; Brauer, 1906; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Cohen, 1981, 1986; Fischer et al., 1981, 1987; Martins, 1981; Cohen et al., 1990; Saldanha et al., 1995; Monteiro et al., 2001; Borges et al., 2002; Moura, 2004; ICES, 2005; Gomes and Olim, 2007; King et al., 2008; Silva, 2008, 2010, 2011; Silva et al., 2009; Carneiro et al., 2014; Christiansen et al., 2014; Froese and Pauly, 2019), Azores▲ (Roule, 1919a; Saldanha, 1977a; Cohen, 1981, 1986; Fischer et al., 1981; INIP, 1985a, b, c; Cohen et al., 1990; Dias, 1991; Lloris et al., 1991; Gonçalves and Santos, 1994; Krug, 1995; Silva et al., 1994, 1995; Silva and Menezes, 1995a, b; Menezes, 1996; Arruda, 1997; Saldanha and Biscoito, 1997; Santos et al., 1997; Morato-Gomes et al., 1998; Desbruyères et al., 2001; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Melo and Menezes, 2002; Cardigos et al., 2005; Menezes et al., 2006, 2009, 2012; Martins et al., 2006; Rosa et al., 2006; González et al., 2008; Pakhorukov, 2008 (PECS); Porteiro et al., 2010, 2013, 2017; Besugo, 2013; Pham et al., 2013; Carneiro et al., 2014; Bañón et al., 2016a; Iwamoto and Cohen, 2016; Almeida and Biscoito, 2019; Eschmeyer et al., 2019; Fauconnet et al., 2019; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1843; Günther, 1862; Vinciguerra, 1883; de Velloso, 1892; Brauer, 1906; Schnakenbeck, 1933; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1952a, 1976; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Cohen, 1979d, 1981, 1986, 1990c; Fischer et al., 1981; INIP, 1982c, 1984a, b, c; Cohen et al., 1990; Lloris et al., 1991; Severino, 2004; Ferreira et al., 2008; González et al., 2008; Pakhorukov, 2008 (PECS); Carneiro et al., 2014; Bañón et al., 2016a; Iwamoto and Cohen, 2016; POGID, 2016; Biscoito et al., 2017; Delgado et al., 2017; Porteiro et al., 2017; Eschmeyer et al., 2019; Froese and Pauly, 2019). Four

specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22464, 1967, set of two specimens and MMF 22468, 1967, set of two specimens).

***Physiculus dalwigki* Kaup 1858**

Common name: **Black codling; Abrótea-de-Dalwigk**, Abrótea-de-natura, Abrótia, Abrótia-de-natura.
 References: mainland Portugal ([Carneiro et al., 2014](#); [OCEANA, 2014](#); [Lorance et al., 2015a](#)), Azores ([Maul, 1976 \(PECS\)](#); [Ehrich, 1977 \(PECS\)](#); [Aguiar and Pereira, 1982](#); [Shcherbachov et al., 1985b \(PECS\)](#); [Cohen, 1986 \(PECS\)](#), [1990c \(PECS\)](#); [Cohen et al., 1990 \(PECS\)](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sousa et al., 1999](#); [Fock et al., 2002 \(PECS\)](#); [Kukuev, 2002 \(PECS\)](#); [Menezes et al., 2006](#); [González et al., 2008](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Lorance et al., 2015a](#); [Bañón et al., 2016a \(PECS\)](#); [Iwamoto and Cohen, 2016 \(PECS\)](#); [Eschmeyer et al., 2019 \(PECS\)](#); [Froese and Pauly, 2019 \(PECS\)](#)) and Madeira ([Kaup, 1858](#); [Günther, 1862, 1880, 1887](#); [Vinciguerra, 1883\(?\)](#); [Vaillant, 1888](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Maul, 1952a](#); [Albuquerque, 1954-1956](#); [Ehrich, 1977](#); [Cohen, 1986, 1990c](#); [Cohen et al., 1990](#); [Lloris et al., 1991](#); [Bañón et al., 2002](#); [Ferreira et al., 2008](#); [González et al., 2008](#); [Carneiro et al., 2014](#); [Lorance et al., 2015a](#); [Bañón et al., 2016a](#); [Iwamoto and Cohen, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

***Rhynchogadus hepaticus* (Facciolà 1884)**

Common name: No English common name; **Abrótea-amuada**.
 References: Azores ([MAR-ECO - HamPelFish - TI/SF_MAR-ECO_1135, 1979 \(PECS\)](#); [Carneiro et al., 2014](#)).

Family Melanonidae

***Melanonus zugmayeri* Norman 1930**

Common name: **Arrowtail; Peixe-cauda-de-seta**.
 References: mainland Portugal ([ZMH 105749, 1968](#); [Cohen, 1986](#); [ZSM-Pis-029158 / 534368 / 385956, 1992](#); [Henriques et al., 2001](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Fernandes et al., 2015d](#); [Froese and Pauly, 2019](#)), Azores ([Koefoed, 1953](#); [Cohen, 1986](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Melo and Menezes, 2002](#); [Porteiro et al., 1999, 2010, 2017](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Fernandes et al., 2015d](#); [Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Maul, 1952a](#); [Albuquerque, 1954-1956](#); [Cohen, 1979d, 1986](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Fernandes et al., 2015d](#); [Delgado et al., 2017](#); [Froese and Pauly, 2019](#)). Two syntypes of *Melanonus macrostoma* Maul 1952, synonym of *M. zugmayeri*, caught off in Madeira are in the MMF collections (MMF 158 and MMF 159).

Family Gadidae

***Gadiculus argenteus* Guichenot 1850**

Common name: **Silvery pout; Badejinho**, Badejo.
 References: mainland Portugal ([Bragança, 1902](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976](#); [Fernández et al., 1981](#); [INIP, 1981a, b, 1982a, b](#); [Martins, 1981](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Svetovidov, 1986a](#); [Fischer et al., 1987](#); [Cohen et al., 1990](#); [Gordo et al., 1990](#); [Saldanha et al., 1995](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Silva and Borges, 2014](#); [Lloris and Iwamoto, 2016](#); [Froese and Pauly, 2019](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)) and Azores ([MNHN-IC-2009-0705, 1971](#); [MNHN-IC-2009-0706, 1971](#); [MNHN-IC-2009-0707, 1971](#); [Cohen et al., 1990](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sousa et al., 1999](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Almeida and Biscoito, 2019](#)).

***Gadus morhua* Linnaeus 1758**

Common name: **Atlantic cod; Bacalhau-do-Atlântico**, Bacalhau, Bacalhau-corrente, Bacalhau-crescido, Bacalhau-curado, Bacalhau-frescal, Bacalhau-da-Terranova, Bacalhau-da-Terra-Nova, Bacalhau-de-Viana, Bacalhau-miúdo, Bacalhau-verde, Bacalhuço, Balona, Cadalhau, Faneca, Fiel-amigo, Flausino, Folha, Folha-de-bacalhau, Gabilami, Juliana, Juliano, Miudeiro, Peixota, Peixota-de-bacalhau, Rabotinho, Raboto, Sanapaio, Sunapai, Tomecode, Trochuela.

References: mainland Portugal▲ and Azores ([Arruda, 1997](#)). One specimen (estimate length and weight: 850 mm TL and 6200 g) was caught off by xávega (beach seine) in Costa de Caparica (38°36.981'N, 9°14.376'W), on September 6, 2016. Other sporadic catches of this species off Portugal (Figueira da Foz and Peniche) has been reported. This is the southernmost record in Europe.

***Merlangius merlangus* (Linnaeus 1758)**

Common name: **Whiting; Badejo**, Bacalhau, Badejo-do-mar-negro, Corvelo.

References: mainland Portugal▲ (subspecies *merlangus*) ([Vandelli, 1771](#); [de Brito Capello, 1868, 1880](#); [Osório, 1888](#); [Nobre, 1894, 1903, 1935](#); [Seabra, 1911](#); [Murray and Hjort, 1912](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979e](#); [Fischer et al., 1987\(?\)](#); [Cohen et al., 1990](#); [Erzini et al., 2001a](#); [ICES, 2005](#); [Henriques et al., 2008](#); [Batista et al., 2009](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#); [Martins and Carneiro, 2018](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)) and Azores (?) ([Osório de Castro, 1967](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *Merlangius merlangus* (subspecies *merlangus*) is Mediterranean Sea, Sea of Marmara, Black Sea, north-eastern Atlantic and adjacent fringes of Arctic regions. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location. Specimens are needed for confirmation in the Azores Archipelago. Thus, this species is not considered in the total number of valid species to the Azores.

***Micromesistius poutassou* (Risso 1827)**

Common name: **Blue whiting; Verdinho**, Lacrau-do-mar, Maria-mole, Pechelim, Pichelim.

References: mainland Portugal▲ ([de Brito Capello, 1868, 1880](#); [Vieira, 1893, 1897, 1900](#); [Nobre, 1935](#); [Ehrenbaum, 1936](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979e, 1986a](#); [Fernández et al., 1981](#); [INIP, 1981a, b, 1982a, b](#); [Martins, 1981](#); [Fischer et al., 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Fischer et al., 1987](#); [Cohen, 1990d](#); [Cohen et al., 1990](#); [Martins et al., 1992](#); [Figueiredo et al., 1994](#); [Saldanha et al., 1995](#); [Erzini et al., 2001a, b](#); [Gomes et al., 2001](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002a, b](#); [Mendes et al., 2004](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Martins and Carneiro, 2018](#); [Mecklenburg et al., 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Osório de Castro, 1967](#); [Silva et al., 1995](#); [Menezes, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sousa et al., 1999](#); [Menezes et al., 2006](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#)) and Madeira ([Fischer et al., 1981](#)).

***Pollachius pollachius* (Linnaeus 1758)**

Common name: **Pollack; Juliania**, Abadejo, Abadiva, Bacalhau, Bacalhau-verdelho, Badejo, Badiva, Palouco-amarelo, Verdelho.

References: mainland Portugal▲ ([Carvalho, 1765](#); [de Brito Capello, 1868, 1880](#); [Steindachner, 1868b](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Vieira, 1893, 1897, 1900](#); [Osório, 1895](#); [Bragança, 1903](#); [Nobre, 1903, 1904, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Borges, 1984](#); [Costa and Quéro, 1988](#); [Martins et al., 1992](#); [Almeida, 1996](#); [Prista et al., 2003](#); [Thiel et al., 2003](#); [ICES, 2005](#); [Henriques et al., 2007, 2008](#); [Carneiro et al., 2014](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)) and Azores ([Drouët 1861b\(?\)](#); [Osório de Castro, 1967](#); [Martins, 1982a](#); [Arruda, 1997](#); [Santos et al., 1997](#)). This species can be confused with *Pollachius virens* (Linnaeus 1758).

***Raniceps raninus* (Linnaeus 1758)**

Common name: **Tadpole fish; Rainúnculo-negro**, Breta, Rainúnculo, Rainúnculo-preto.

References: mainland Portugal ([Osório, 1895, 1896](#); [Vieira, 1897, 1900](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Costa and Quéro, 1988](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#)).

***Trisopterus capelanus* (Lacepède 1800)**

Common name: Poor cod Mediterranean; **Fanecão-do-Mediterrâneo**.

References: mainland Portugal ([Fischer et al., 1987](#); [Gaemers, 2017](#)). Fischer et al., 1987 reported this species as *Trisopterus minutus capelanus* (Lacepède 1800) and Delling et al. (2011) resurrected this

subspecies from the synonymy of *T. minutus* to the species status *T. capelanus* based on molecular and morphological data. The previous geographic distribution of *T. capelanus* was in the Adriatic Sea, the West of Mediterranean Sea and the Atlantic coast of Morocco (Tortonese, 1970). More recently, Gaemers (2017) extended the distribution south range to Western Sahara and along the Atlantic coast of Portugal and at least up to and including the Ría de Arosa (Galicia, Spain). This species can easily be confused with *Trisopterus minutus* (Linnaeus 1758).

***Trisopterus luscus* (Linnaeus 1758)**

Common name: **Pouting; Faneca.**

References: mainland Portugal▲ ([Carvalho, 1765](#); [Vandelli, 1771](#); [de Brito Capello, 1868, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Osório, 1895, 1896](#); [Vieira, 1897, 1900](#); [Bragança, 1902](#); [Seabra, 1911](#); [Murray and Hjort, 1912](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Svetovidov, 1979e](#); [INIP, 1981a, b, 1982a, b](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Svetovidov, 1986a](#); [Costa and Quéro, 1988](#); [Cohen, 1990d](#); [Cohen et al., 1990](#); [Martins et al., 1992, 2015](#); [Almeida, 1996](#); [Gonçalves et al., 1996, 1997](#); [Erzini et al., 1997, 2001a](#); [Gomes et al., 2001](#); [Jorge et al., 2002](#); [Santos et al., 2002a](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Stergiou et al., 2006](#); [Fernandes et al., 2007](#); [Henriques et al., 2007, 2008](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Batista et al., 2009](#); [Gonçalves et al., 2010](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Gaemers, 2017](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Fischer et al., 1981](#); [INIP, 1982c](#); [Gaemers, 2017](#)). One specimen caught off at the Portuguese continental slope in March 2011 is in the MB collections (MB06-005325).

***Trisopterus minutus* (Linnaeus 1758)**

Common name: **Poor cod; Fanecão.**

References: mainland Portugal▲ ([Steindachner, 1868b](#); [Vieira, 1893, 1897, 1900](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Fernández et al., 1981](#); [INIP, 1981a, b, 1982b](#); [Martins, 1981](#); [Borge, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Svetovidov, 1986a](#); [Cohen et al., 1990](#); [Martins et al., 1992, 2015](#); [Almeida, 1996](#); [Gomes et al., 2001](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2006](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Gaemers, 2017](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Family Lotidae

***Brosme brosme* (Ascanius 1772)**

Common name: **Tusk; Bolota, Abrótea, Abrótia, Alecrim, Bacalhau, Brota, Lobelino, Lobelina, Lobina, Zarbo.**

References: Azores (PECS) ([Porteiro et al., 2017](#)).

***Ciliata mustela* (Linnaeus 1758)**

Common name: **Fivebeard rockling; Laibeque-de-cinco-barbillhos, Badião, Bedião, Bodião, Laibeque, Látego, Látego, Pico.**

References: mainland Portugal▲ ([de Brito Capello, 1868, 1880](#); [Osório, 1888, 1896](#); [Nobre, 1895c, 1935](#); [Vieira, 1900](#); [Bragança, 1902](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979e, 1986a](#); [Dinis, 1981](#); [Arruda et al., 1988](#); [Cohen et al., 1990](#); [Moreira et al., 1992](#); [Almeida, 1996](#); [Cabral, 1999](#); [Costa et al., 2002](#); [Jorge et al., 2002](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Henriques et al., 2007, 2008](#); [Neves et al., 2008](#); [Carneiro et al., 2014](#); [Fernandes et al., 2014](#); [Martins et al., 2015](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)).

***Gaidropsarus granti* (Regan 1903)**

Common name: Azores rockling; **Laibeque-dos-Açores**, Aranha-do-alto, Laibeque, Látego, Viúva-do-alto.

References: Azores▲ ([Regan, 1903a](#); [Collett, 1905](#); [Vaillant, 1919](#); [Fowler, 1936a\(?\)](#); [Svetovidov, 1979e, 1986a](#); [Dooley et al., 1985](#); [Cohen, 1990d](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bañón et al., 2002](#); [Porteiro et al., 2010](#); [Orsi Relini and Relini, 2013](#); [Carneiro et al., 2014](#); [Francisco et al., 2014](#);

Bañón *et al.*, 2016a; Lloris and Iwamoto, 2016; Eschmeyer *et al.*, 2019; Spinelli and Castríota, 2019) and Madeira (Menezes *et al.*, 2009; Carneiro *et al.*, 2014; Froese and Pauly, 2019; Spinelli and Castríota, 2019). The lectotype of *Onus granti* Regan 1903, synonym of *G. granti*, caught off in the Azores is in the NHMUK collections (NHMUK 1903.10.22.1).

***Gaidropsarus macrophthalmus* (Günther 1867)**

Common name: **Bigeye rockling**; **Laibeque-do-Atlântico**, common name as *Gaidropsarus biscayensis* (Collett, 1890); **Laibeque-olho-grande**.

References: mainland Portugal (Maul, 1976; Svetovidov, 1986a, b; Fischer *et al.*, 1987; Cohen *et al.*, 1990; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002; ICES, 2005; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Chaves, 2008; Silva, 2008, 2011; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Francisco *et al.*, 2014), Azores (Maul, 1976 (PECS); Hureau and Monod, 1979 (PECS); Svetovidov, 1986a, b; Kukuev, 2002 (PECS); Lloris and Iwamoto, 2016) and Madeira (Svetovidov, 1986a, b; Fischer *et al.*, 1987; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Francisco *et al.*, 2014; Lloris and Iwamoto, 2016). According to Barros-García *et al.* (2017) and Eschmeyer *et al.* (2019) the species *Gaidropsarus biscayensis* (Collett 1890) is a junior synonym of *G. macrophthalmus*. Several specimens caught off at the coast of the mainland Portugal are in the MMF collections (MMF 22495, 1967, set of thirteen specimens and MMF 22859, 1967, one specimen).

***Gaidropsarus mauli* Biscoito & Saldanha 2018**

Common name: No English common name; **Laibeque-de-Maul**.

References: Azores (Biscoito and Saldanha, 2018).

***Gaidropsarus mediterraneus* (Linnaeus 1758)**

Common name: **Shore rockling**; **Laibeque-do-Mediterrâneo**, Abrosto, Abrótea, Abrótea-da-poça, Abrótia, Barbaíscos, Barbisco, Bedião, Bodião, Guingau, Larote, Larote-de-cabeça-encangrenada, Latego, Latego-de-cabeça-encangrenada, Látego, Lulão, Paz-da-casa, Pico-de-el-rei, Raposa, common name as *Gaidropsarus guttatus* (Collet 1890): Abrótea-da-poça, Abrótia-da-poça, Aranha-da-pedra, Guingau, Laibeque, Viúva.

References: mainland Portugal▲ (de Brito Capello, 1880; Nobre, 1895c, 1903, 1935; Osório, 1895; Vieira, 1900; Seabra, 1911; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Svetovidov, 1979e, 1986a, b; Fernández *et al.*, 1981; Fischer *et al.*, 1987; Cohen *et al.*, 1990; Saldanha *et al.*, 1995(?); Almeida, 1996; Garnerot *et al.*, 2004; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Henriques *et al.*, 2007, 2008; Silva, 2008, 2010; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Francisco *et al.*, 2014; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores▲ (Hilgendorf, 1888a; Collett, 1890a, 1892, 1896, 1905; Vaillant, 1919; Nobre, 1924, 1930, 1935; Zugmayer, 1933; Fowler, 1936a; Belloc, 1949; Maul, 1952; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Arruda, 1979a, 1980b, 1992; Cohen and Russo, 1979; Svetovidov, 1979e; Dooley *et al.*, 1985; Cohen, 1990d; Cohen *et al.*, 1990; Patzner *et al.*, 1990, 1992; Azevedo *et al.*, 1991, 1992; Lloris *et al.*, 1991; Arruda *et al.*, 1992; Santos 1992a; Santos *et al.*, 1994b, 1997; Arruda, 1997; Wirtz, 1994; Azevedo, 1995; Gonçalves *et al.*, 1995; Morato *et al.*, 2001b; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Francisco *et al.*, 2014; Afonso, 2016; Lloris and Iwamoto, 2016; Barros-García *et al.*, 2017) and Madeira (Lowe, 1837; Günther, 1862; Fowler, 1936a; Maul, 1952a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Svetovidov, 1979e; Cohen, 1990d; Lloris *et al.*, 1991; EMEPC, 2010; Carneiro *et al.*, 2014; Francisco *et al.*, 2014; Almada *et al.*, 2015; Afonso, 2016; Lloris and Iwamoto, 2016; POGIS, 2016). One syntype of *Onus guttatus* Collett 1890, synonym of *G. mediterraneus*, caught off at Fayal (Azores) is in the MOM collections (MOM-IC-0091-3719). According to Barros-García *et al.* (2017) and Eschmeyer *et al.* (2019) the species *Gaidropsarus guttatus* (Collet 1890) is a junior synonym of *G. mediterraneus*.

***Gaidropsarus vulgaris* (Cloquet 1824)**

Common name: **Three-bearded rockling**; **Laibeque-de-três-barbilhos**, Abroito, Abrótea, Abrótea-da-poça, Abrótia, Abrótia-de-poça, Badião, Barbacho, Barbaíscos, Barbaixo, Barbisco, Bardo-do-mar, Bedião, Bodião, Caboz, Caralhete d'Elrei; Julianas, Laibeque, Larote, Larote-de-cabeça-encangrenada, Larote-de-cabeça-encarnada, Látego, Lulão, Paz-da-casa, Pica, Pica-de-el-rei, Pico, Pico-de-el-rei, Raposa.

References: mainland Portugal (de Brito Capello, 1868, 1880; Steindachner, 1868b; Osório, 1888, 1895; Vieira, 1893, 1897, 1900; Nobre, 1894, 1903, 1935; Bragança, 1902, 1903(?); Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Svetovidov, 1979e, 1986a; INIP, 1982a, b; Fischer et al., 1987; Antunes et al., 1988; Cohen et al., 1990; Saldanha et al., 1995; Borges et al., 2002, 2003; Moura, 2004; ICES, 2005; Fernandes et al., 2007; Henriques et al., 2007, 2008; Carneiro et al., 2014; Francisco et al., 2014; Lloris and Iwamoto, 2016; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (?) (Drouët 1861b; Hilgendorf, 1888a; Vaillant, 1919; Zugmayer, 1933; Fowler, 1936a, b; Osório de Castro, 1967; Arruda, 1992, 1997) and Madeira (MNHN-IC-1883-0097, 1882; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Carneiro et al., 2014; Lloris and Iwamoto, 2016). The current geographic distribution of *G. vulgaris* is in the North-eastern Atlantic from Norway and Iceland south to Gibraltar and also in the Western Baltic Sea, North Sea, Mediterranean Sea, and Sea of Marmara as reported by Eschmeyer et al. (2019). Santos et al. (1997) reviewed the former quotes for the Azores Archipelago and concluded that this species does not occur in the Azores. Thus, this species is not considered in the total number of valid species to the Azores. This species can be confused with *Gaidropsarus granti* (Regan 1903).

***Molva dypterygia* (Pennant 1784)**

Common name: **Blue ling; Maruca-azul**, Donzela, Donzela-azul, Lingue-azul.

References: mainland Portugal (Svetovidov, 1986a; Cohen et al., 1990; Carneiro et al., 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019) and Azores (Sousa et al., 1999; Hareide and Garnes, 2001 (PECS); Morato et al., 2001a; Carneiro et al., 2014; Lloris and Iwamoto, 2016).

***Molva macrophthalma* (Rafinesque 1810)**

Common name: **Spanish ling; Maruca-do-Mediterrâneo**, Abrótea, Abrótia, Donzela, Juliana, Juliana-dapiedra, Pescada-dos-Açores, Peixe-pau.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1900; Nobre, 1894, 1935; Osório, 1895; Bragança, 1903; Cligny, 1903; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Svetovidov, 1979e; Fernández et al., 1981; Martins, 1981; Fischer et al., 1987; Cohen, 1990d; Carneiro et al., 2014; Martins and Carneiro, 2018) and Azores▲ (Osório de Castro, 1967; Dias et al., 1976, 1990; INIP, 1985a, b; Dias, 1991; Silva et al., 1994, 1995; Clarke et al., 1995; Krug, 1995; Silva and Menezes, 1995a, b; Menezes, 1996; Arruda, 1997; Santos et al., 1997; Cardigos et al., 2005; Menezes et al., 2006; Rosa et al., 2006; Porteiro et al., 2010, 2013; Besugo, 2013; Pham et al., 2013; Carneiro et al., 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019).

***Molva molva* (Linnaeus 1758)**

Common name: **Ling; Maruca**, Abrótea, Abrótia, Bacalhau-de-água-doce, Donzela, Juliana, Lingue, Malva, Maruga.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; Vinciguerra, 1883; Almeida and Roquette, 1892; MNHN-IC-1905-0579, 1898; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Svetovidov, 1979e; Martins, 1981; Cohen et al., 1990; Martis et al., 1992; Borges et al., 2002; ICES, 2005; Henriques et al., 2008; Silva et al., 2009; Silva, 2010, 2011; Costa et al., 2012; Carneiro et al., 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Mecklenburg et al., 2018; Froese and Pauly, 2019) and Azores (Osório de Castro, 1967; INIP, 1985a; Dias, 1991; Arruda, 1997).

Family Phycidae

***Phycis blennoides* (Brünnich 1768)**

Common name: **Greater forkbeard; Abrótea-do-alto**, Abrota, Abrótea, Abrótea-da-fundura, Abrótea-de-fundura, Abrótea-de-vapor, Abrótia, Abrótia-do-alto, Abrótia-da-fundura, Abrótia-de-fundura, Abrótia-de-vapor, Brota, Juliana, Melga, Ricardo.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; Steindachner, 1868b; Vaillant, 1888; Baldaque da Silva, 1891; Vieira, 1893, 1897, 1900; Goode and Bean, 1895, 1896; Bragança, 1902, 1903; Brauer, 1906; Richard, 1908; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Svetovidov, 1979e; Fernández et al., 1981; Fischer et al., 1981,

1987; INIP, 1981a, b, 1982a, b; Martins, 1981; Dinis and Marecos, 1984; Dinis, 1985; Cohen, 1990d; Cohen *et al.*, 1990; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Erzini *et al.*, 2001b; Monteiro *et al.*, 2001; Borges *et al.*, 2002; Santos *et al.*, 2002b; Mendes *et al.*, 2004; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Silva, 2008, 2010, 2011; Bordalo-Machado *et al.*, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Bañón *et al.*, 2016a; Martins and Carneiro, 2018; Mecklenburg *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Drouët 1861b; Osório de Castro, 1967; Dias *et al.*, 1976; Martins, 1982a; INIP, 1985b; Silva, 1986a; Dias, 1991; Santos *et al.*, 1994a, 1995a, 1997; Silva *et al.*, 1994, 1995; Krug, 1995; Isidro, 1996; Menezes, 1996; Arruda, 1997; Morato-Gomes *et al.*, 1998; Hareide and Garnes, 2001 (PECS); Morato *et al.*, 2001a; Menezes *et al.*, 2006, 2009, 2012; Rosa *et al.*, 2006; Porteiro *et al.*, 2010, 2017; Besugo, 2013; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Lloris and Iwamoto, 2016; Froese and Pauly, 2019) and Madeira (Johnson, 1862c; Günther, 1862; de Brito Capello, 1872; Nobre, 1935; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1952a; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Cohen *et al.*, 1990; Carneiro *et al.*, 2014; Lloris and Iwamoto, 2016; Froese and Pauly, 2019).

Phycis phycis (Linnaeus 1766)

Common name: **Forkbeard; Abrótea-da-costa**, Abrota, Abrótea, Abrótea-da-costa, Abrótea-da-pedra, Abrótea-da-rocha, Abrótea-de-três-barbas, Abrótea-do-baixo, Abrótea-legítima, Abrótia, Abrótia-da-costa, Abrótia-da-pedra, Abrótia-de-três-barbas, Abrótia-do-baixo, Abrótia-legítima, Albrótea, Albrótia, Brotá.

References: mainland Portugal[▲] (de Brito Capello, 1868, 1880; Steindachner, 1868b; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1900; Nobre, 1894, 1895a, 1903, 1935; Osório, 1896; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Fischer *et al.*, 1981, 1987; Martins, 1981; INIP, 1982a, b; Dinis and Marecos, 1984; Dinis, 1985; Svetovidov, 1986a; Costa and Quéro, 1988; Cohen, 1990d; Cohen *et al.*, 1990; Martins *et al.*, 1992; Saldanha *et al.*, 1995; Almeida, 1996; Gonçalves *et al.*, 1996, 1997; Erzini *et al.*, 1997, 2001a; Borges *et al.*, 2002; Mendes *et al.*, 2004; Moura, 2004; ICES, 2005; Stergiou *et al.*, 2006; Fernandes *et al.*, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Abecasis *et al.*, 2009; Batista *et al.*, 2009; Silva *et al.*, 2009; Silva, 2010; Chaves and Silva, 2013; Carneiro *et al.*, 2014; OCEANA, 2014; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; McGuire, 1879; Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Sampaio, 1904; Vaillant, 1919; Nobre, 1924, 1930; Fowler, 1936b; Ribeiro, 1936; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Wood, 1974; Dias *et al.*, 1976, 1990; Maul, 1976 (PECS); Ehrich, 1977; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a; Fischer *et al.*, 1981; Muzanor and Westhaus, 1981; Martins, 1982b; INIP, 1985a, b, c; Dias and Monteiro, 1986; Silva, 1986a, b, 1992, 1995; Isidro, 1987, 1989, 1996; Cohen, 1990d (PECS); Cohen *et al.*, 1990; Barreiros, 1990; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, 1995a, 1997; Wirtz, 1990a, 1994; Dias, 1991; Dias and Cascalho, 1991; Monteiro, 1991; Azevedo *et al.*, 1991, 1992; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Silva and Krug, 1992; Azevedo, 1995; Krug, 1995; Silva *et al.*, 1994, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a, b; Kukuev, 2002 (PECS); Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010, 2013; Diogo and Pereira, 2013; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Afonso, 2016; Lloris and Iwamoto, 2016; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Lowe, 1837, 1843-1860, 1850, 1852; Günther, 1862; de Velloso, 1892; Goode and Bean, 1895, 1896; D'Ancona, 1933; Fowler, 1936a; Noronha and Sarmento, 1948; Maul, 1952a, 1976; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nunes, 1974; Ehrich, 1977; Svetovidov, 1979e; INIP, 1980, 1982c, 1984b, c, 1985c; Fischer *et al.*, 1981; Cohen *et al.*, 1990; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Ferreira *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Lloris and Iwamoto, 2016; POGID, 2016; POGIS, 2016; Biscoito *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Fowler (1936b) mentioned that two specimens of this species were caught off the Azores, Fayal in 1879 and Terceira in 1894. One specimen caught off at the Portuguese continental slope in October 2010 is in the MB collections

(MB06-005358). Three specimens caught off at the Meteor Bank in 1967 are in the MMF collections (MMF 22345, 1967, MMF 22346, 1967 and MMF 22372, 1967).

Family Merlucciidae

***Lyconus brachycolus* Holt & Byrne 1906**

Common name: No English common name; **Peixe-de-focinho-arreganhado**.

References: mainland Portugal ([Moura, 2004](#); [Carneiro et al., 2014](#)), Azores ([Post, 1987a \(PECS\)](#); [MAR-EKO 3336, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([Maul, 1951a](#); [Albuquerque, 1954-1956](#); [Marshall, 1979](#); [Svetovidov, 1979e, 1986a](#); [Maurin 1990](#); [Lloris et al., 1991, 2005](#); [Carneiro et al., 2014](#); [Fernandes et al., 2015d](#); [Lloris and Iwamoto, 2016](#); [Froese and Pauly, 2019](#)). The holotype and two paratypes of *Macruronus caninus* Maul 1951, synonyms of *L. brachycolus*, caught off in Madeira are in the MMF collections (MMF 3155, MMF 3156 and MMF 3157, respectively).

***Macruronus maderensis* Maul 1951**

Common name: No English common name; **Peixe-orelhudo-da-Madeira**.

References: Madeira ([Maul, 1951a](#); [Albuquerque, 1954-1956](#); [Marshall, 1979](#); [Svetovidov, 1979e](#); [Carneiro et al., 2014](#); [Lloris and Iwamoto, 2016](#); [Priede, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype of this species caught off Madeira (stomach content) is in the MMF collections (MMF 3155).

***Merluccius merluccius* (Linnaeus 1758)**

Common name: **European hake; Pescada-branca**, Escamanta, Marmota, Marmota-branca, Marmota-grada, Marmota-média, Morcego-do-mar, Peixota, Pescada, Pescada-branca-do-Atlântico, Pescada-comum, Pescada-comum-do-Atlântico, Pescada-da-costa, Pescada-europeia, Pescada-de-viana, Pescada-de-vigo, Pescada-marmota, Pescada-vulgar, Pescadinha, Pescadinha-de-rabo-na-boca, Pescadinha-marmota, Pissota.

References: mainland Portugal▲ ([Carvalho, 1765](#); [de Brito Capello, 1868, 1869b, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1893, 1897, 1900](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Osório, 1895](#); [Bragança, 1902](#); [Seabra, 1911](#); [Murray and Hjort, 1912](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Svetovidov, 1979e, 1986a](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Inada, 1981](#); [INIP, 1981a, b, 1982a, b](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Cohen et al., 1990](#); [Martins et al., 1992, 2015](#); [Figueiredo et al., 1994](#); [Saldanha et al., 1995](#); [Erzini et al., 1997, 2001a, b](#); [Gomes et al., 2001](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002a, b, c](#); [Coelho et al., 2003](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Mendes et al., 2004](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Stergiou et al., 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Fernandes et al., 2015e](#); [Bañón et al., 2016a](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1890a](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [MNHN-IC-1956-0019, 1955](#); [Isidro, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#); [Fernandes et al., 2015e](#)) and Madeira▲ ([Lowe, 1837, 1839, 1840a, b, 1841, 1844a, 1849](#); [de Velloso, 1892](#); [Collett, 1896](#); [Noronha and Sarmento, 1948](#); [Maul, 1951a](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Svetovidov, 1979e](#); [Fischer et al., 1981](#); [Inada, 1981](#); [Carneiro et al., 2014](#); [Fernandes et al., 2015e](#)). According to [Santos et al. \(1997\)](#) the occurrence records of *Merluccius merluccius* by [Fowler \(1936a\)](#), based on *Onus guttatus* by [Collett \(1890a\)](#), should be considered a synonym of *Gaidropsarusr guttatus* ([Collett 1890](#)). [Albuquerque \(1954-1956\)](#) also refers the presence of this species in the Azores based on [Fowler \(1936a\)](#). The distribution map of this species reported by [Fernandes et al. \(2015e\)](#) partially included the area of Selvagens Islands (Madeira EEZ). One specimen caught off at the Portuguese continental slope in March 2011 is in the MB collections (MB06-005337).

Order Polymixiiformes

Family Polymixiidae

***Polymixia nobilis* Lowe 1838**

Common name: **Stout beardfish; Salmonete-do-alto**, Salmonéta-do-alto, Barbudo, Salmonete.

References: Azores[▲] (Maul, 1976 (PECS); Ehrich, 1977 (PECS); Saldanha, 1977a; Hureau, 1986a; Quéro, 1990b; Lloris et al., 1991; Santos et al., 1995a, 1997; Silva and Menezes, 1995b; Silva et al., 1995; Menezes, 1996; Arruda, 1997; Fock et al., 2002 (PECS); Kukuev, 2002 (PECS); Porteiro et al., 2010; Carneiro et al., 2014; Moore, 2016a; Borden et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837, 1838; NHMUK 1855.11.29.10, 1854-1861; Günther, 1859, 1880(?), 1887; de Velloso, 1892; Vinciguerra, 1893; Goode and Bean, 1895, 1896; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Maul, 1954a, 1976; Osório de Castro, 1954, 1967; Ehrich, 1977; Nielsen, 1979d; Fischer et al., 1981; Hureau, 1981, 1986a; INIP, 1984c; Dooley et al., 1985; Quéro, 1990b; Lloris et al., 1991; Carneiro et al., 2014; Moore, 2016a; POGID, 2016; Borden et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019). The holotype of this species caught off in Madeira is in the NHMUK collections (NHMUK 1855.11.29.10). One specimen caught off at the Meteor Bank is in the MMF collections (MMF 22359, 1967).

Order Beryciformes

Family Berycidae

***Beryx decadactylus* Cuvier 1829**

Common name: **Alfonsino; Imperador**, Alfacim, Alfonsim, Alfonsim-de-costa-largal, Alfonsino, Alfoucim, Cardeal, Cardial, Folião, Melo, Realista, Regulado.

References: mainland Portugal[▲] (de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Günther, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1903, 1935; Goode and Bean, 1895, 1896; Cligny, 1903; Brauer, 1906; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nielsen, 1979i; Fernández et al., 1981; INIP, 1981a, 1982a; Martins, 1981; Dinis, 1985; Maul, 1986c; Fischer et al., 1987; Borges et al., 2002; Santos et al., 2002b; Moura, 2004; ICES, 2005; Fernandes et al., 2007; Gomes and Olim, 2007; Henriques et al., 2008; Silva, 2010; Chaves and Silva, 2013; Carneiro et al., 2014; Bañón et al., 2016a; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores[▲] (Steindachner, 1867; Hilgendorf, 1888a; Regan, 1903a; Sampaio, 1904; Vaillant, 1918, 1919; Mohr, 1927; Ramalho, 1929b; Fowler, 1936a, b; Bougis, 1945; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Woods and Sonoda, 1973; Maul, 1976 (PECS); 1986c; Wheeler, 1978; Nielsen, 1979i; Busakhin, 1982; Martins, 1982a; INIP, 1985a; Shcherbachov et al., 1985b (PECS); Dias et al., 1990; Dias, 1991; Lloris et al., 1991; Monteiro, 1991; Arruda, 1992, 1997; Silva and Krug, 1992; Krug, 1995; Silva et al., 1994, 1995; Clarke et al., 1995; Silva, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Santos et al., 1997; Morato-Gomes et al., 1998; Sousa et al., 1999; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Cardigos et al., 2005; Menezes et al., 2006; Rosa et al., 2006; Pakhorukov, 2008 (PECS); Porteiro et al., 2010, 2013; Besugo, 2013; Pham et al., 2013; Carneiro et al., 2014; Moore, 2016b; Froese and Pauly, 2019) and Madeira[▲] (Cuvier, 1829; Lowe, 1838, 1849; Günther, 1859, 1880; Steindachner, 1867; Von Martens, 1876; Goode and Bean, 1895, 1896; Brauer, 1906; Ramalho, 1929b; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Maul, 1954a, 1986c; Osório de Castro, 1954, 1967; Nielsen, 1979i; Fischer et al., 1981; INIP, 1984c; Dooley et al., 1985; Lloris et al., 1991; Ferreira et al., 2008; Carneiro et al., 2014; Moore, 2016b; POGID, 2016; Froese and Pauly, 2019).

***Beryx splendens* Lowe 1834**

Common name: **Splendid alfonsino; Imperador-de-costa-estreita**, Alfacim, Alfonsim, Alfonsim-de-costa-estreita, Alfoucim, Cardeal, Melo, Salão, Selongo.

References: mainland Portugal[▲] (Bragança, 1903; Osório, 1917; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Ehrich, 1977; Maul, 1986c; Borges et al., 2002; ICES, 2005; Fernandes et al., 2007; Chaves, 2008; Silva, 2008, 2010; Silva et al., 2009; Chaves and Silva, 2013; Carneiro et al., 2014; Bañón et al., 2016a; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Fowler, 1936a, b; Collins, 1954; Albuquerque, 1954-1956; Woods, 1957; Osório de Castro, 1967; Woods and Sonoda, 1973; Wheeler, 1978; Busakhin, 1982; INIP, 1985a, b;

Shcherbachev *et al.*, 1985b (PECS); Dias and Monteiro, 1986; Maul, 1986c; Dias *et al.*, 1990; Dias, 1991; Monteiro, 1991; Arruda, 1992, 1997; Silva, 1992, 1995; Silva *et al.*, 1994, 1995; Clarke *et al.*, 1995; Krug, 1995; Silva and Menezes, 1995a,b; Isidro, 1996; Menezes, 1996; Ehrich, 1977 (PECS); Saldanha and Bischoff, 1977b; Lloris *et al.*, 1991; Isidro, 1996; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Melo and Menezes, 2002; Cardigos *et al.*, 2005; Menezes *et al.*, 2006, 2009, 2012; Rosa *et al.*, 2006; Pakhorukov, 2008 (PECS); Porteiro *et al.*, 2010, 2013; Besugo, 2013; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Moore, 2016b; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1833a, 1837, 1838, 1843; NHMUK 1857.6.13.113, 1854-1861; Günther, 1859, 1880; Von Martens, 1876; Goode and Bean, 1895, 1896; Brauer, 1906; Borodin, 1934; Nobre, 1935; Fowler, 1936a, b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Maul, 1954a, 1986c; Osório de Castro, 154, 1967; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Shcherbachev *et al.*, 1985b; Ferreira *et al.*, 2008; Pakhorukov, 2008; Carneiro *et al.*, 2014; Moore, 2016b; POGID, 2016; Delgado *et al.*, 2017; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype of this species caught off in Madeira is in the NHMUK collections (NHMUK 1857.6.13.113). One specimen caught off at the Portuguese continental slope is in the MB collections (MB06-005161).

Family Melamphaidae

Melamphaes ebelingi Keene 1973

Common name: No English common name; **Peixe-rã-de-Ebeling**.

References: Madeira ([Kotlyar, 2012](#); [Harold, 2015i](#)).

Melamphaes indicus Ebeling 1962

Common name: No English common name; **Peixe-rã-do-Índico**.

References: Madeira (?) ([ZMH 105339, 1968](#); [MCZ 58093, 1970](#); [Carneiro et al., 2014](#)). There are two specimens of *M. indicus* in the ZMH and MCZ collections caught off in the Eastern Atlantic Ocean (Madeira EEZ). However, the current geographic distribution is Indo-Pacific from Madagascar east to Hawaiian Islands and Central America ([Eschmeyer et al., 2019](#)). Thus, the re-evaluation of those specimens is needed and this species is not considered in the total number of valid species.

Melamphaes falsidicus Kotlyar 2011

Common name: No English common name; **Peixe-rã-falso**, Peixe-rã, Rã.

References: Azores ([Kotlyar, 2011](#); [Carneiro et al., 2014](#)) and Madeira (?) ([Carneiro et al., 2014](#)). Kotlyar (2011) described the species *M. falsidicus* from the northern Atlantic Ocean, where it was sampled between 34° N and 58° N. According to Kotlyar (2011), this species was previously defined as *Melamphaes microps* (Günther 1878), which is probably a variant of *M. falsidicus*. Thus, a few records of the species *M. microps* off the Azores should probably be considered as *M. falsidicus*. The occurrence of this species in Madeira is doubtful and it is not considered in total number of valid species for this archipelago.

Melamphaes longivelis Parr 1933

Common name: Eyebrow bigscale; **Peixe-rã-sobrolho**, Peixe-rã, Rã.

References: mainland Portugal ([Maul, 1986a](#); [ZSM-Pis-029157 / 534367 / 385955, 1992](#); [ZSM-Pis-029197 / 534405 / 385993, 1992](#)), Azores ([Ebeling and Weed, 1973](#); [Maul, 1986a](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Maul, 1979d, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Melamphaes microps (Günther 1878)

Common name: Ridgehead; **Peixe-rã-olhinho**, Peixe-rã, Rã.

References: mainland Portugal ([Maul, 1986a](#)), Azores ([Ebeling and Weed, 1973](#); [Maul, 1986a](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Sutton et al., 2008](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019\(?\)](#)) and Madeira ([Maul, 1986a](#); [Froese and Pauly, 2019\(?\)](#)). For Eschmeyer *et al.* (2019) the current geographic distribution of *M. microps* is the North Atlantic between 30°N and 64°N. However, Froese and Pauly (2019) referred that in the North Atlantic (between 30°N and 60°N) specimens are probable variants of *M. falsidicus*.

Melamphaes polylepis Ebeling 1962

Common name: No English common name; **Peixe-rã-cabecinha**, Peixe-rã, Rã.

References: Azores (PECS) ([ZMH 121152](#), 1982; [ZMH 121153](#), 1982; [ZMH 121154](#), 1982; Carneiro et al., 2014). According to Kotlyar (2011) this species is characterized by circumtropical geographic range (Atlantic, Indian, western and central Pacific Ocean) and Eschmeyer et al. (2019) mention it as a circumglobal species in tropical and subtropical seas, in the Atlantic between the equator and 20°N. However, there are in the Zoologisches Institut und Museum Hamburg of the Hamburg University three records of specimens of *M. polylepis* caught off in the Eastern Atlantic, at the Azores PECS, between 44° and 47°N.

Melamphaes pumilus Ebeling 1962

Common name: No English common name; **Peixe-rã-anão**.

References: Azores (PECS) ([SOC Discovery 1126123_85_FIS_516001](#), 1985; Carneiro et al., 2014; Kotlyar, 2016a).

Melamphaes simus Ebeling 1962

Common name: Ridgehead; **Peixe-rã-nariz-achatado**, Peixe-rã, Rã.

References: mainland Portugal (YPM ICH 005174, 1959; Osório de Castro, 1967; Maul, 1979d, 1986a; Carneiro et al., 2014; Kotlyar, 2016b), Azores ([SIO 60-153](#), 1922; Ebeling 1962; Osório de Castro, 1967; Ebeling and Weed, 1973; Maul, 1986a; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017(?); Carneiro et al., 2014; Kotlyar, 2016b) and Madeira (Osório de Castro, 1967 ; ZMH 105695, 1968; Maul, 1979d, 1986a; Quéro, 1982a (PECS)(?); Carneiro et al., 2014; Kotlyar, 2016b).

Melamphaes suborbitalis (Gill 1883)

Common name: Shoulderspine bigscale; **Peixe-rã-de-ombro-espinhoso**, Peixe-rã, Rã.

References: mainland Portugal (Maul, 1986a; ZSM-Pis-028592 / 533815 / 385403, 1991; Carneiro et al., 2014), Azores ([SIO 60-152](#), 1922; Roule and Angel, 1933; Ebeling 1962; Osório de Castro, 1967; Ebeling and Weed, 1973; Maul, 1986a; Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017 (PECS); Carneiro et al., 2014) and Madeira ([SIO 58-104](#), 1957; [SIO 58-103](#), 1958; Osório de Castro, 1967; Maul, 1979d, 1986a; Lloris et al., 1991; Kotlyar, 2012; Carneiro et al., 2014).

Melamphaes typhlops (Lowe 1843)

Common name: Ridgehead; **Peixe-rã-da-Madeira**, Peixe-rã, Rã.

References: mainland Portugal (Roule, 1914a, 1919a; Nybelin, 1948; Koefoed, 1953; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1979d, 1986a; Carneiro et al., 2014; Kotlyar, 2016c), Azores (Zugmayer, 1911b; Fowler, 1936a; Koefoed, 1953 (PECS); Albuquerque, 1954-1956; Ebeling 1962; Osório de Castro, 1967; Ebeling and Weed, 1973; Maul, 1986; IORAS 03292, 1988 (PECS); Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Kotlyar, 2016c) and Madeira (Lowe, 1843, 1844b, 1850, 1852; Günther, 1864a, 1887; NHMUK 1864.11.8.1, 1864-1877; Goode and Bean, 1895, 1896; Norman, 1929; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1986a; Lloris et al., 1991; Kotlyar, 2004, 2016c; Carneiro et al., 2014). The neotype of *Metopias typhlops* Lowe 1843, synonym of *Melamphaes typhlops*, caught off in Madeira is in the NHMUK collections (NHMUK 1864.11.8.1).

Poromitra capito Goode & Bean 1883

Common name: Ridgehead; **Peixe-rã-de-nariz-espinhoso**, Peixe-rã, Rã.

References: mainland Portugal (Osório, 1906, 1909; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; ZMH 103400, 1966; Osório de Castro, 1967; Maul, 1979d, 1986a, 1990; Quéro, 1982a; Kotlyar, 2009a; Carneiro et al., 2014), Azores (Roule and Angel, 1933; Belloc, 1949; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Geistdoerfer et al., 1971; Ebeling and Weed, 1973; Maul, 1979d, 1986a; Arruda, 1997; Santos et al., 1997; Melo and Menezes, 2002; Kotlyar, 2004; Sutton et al., 2008; Kotlyar, 2009a; Porteiro et al., 2010, 2017; Menezes et al., 2012; Carneiro et al., 2014; Almeida and Biscoito, 2019) and Madeira (MOM-IC-0091-1394, 1911; MOM-IC-0091-1530, 1911; Roule and Angel, 1933; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Maul, 1954a, 1979d, 1986a, 1990; SIO 57-132, 1956; Osório de Castro, 1967; Quéro, 1982a; Lloris et al., 1991; Kotlyar, 2004, 2009a; de Casamajor and Morandieu, 2013; Carneiro et al., 2014; Froese and Pauly, 2019). The holotype (unique) of *Lophocephalus anthrax* Osório 1906 caught off western coast of

Portugal (stomach content) was preserved in the Museu Bocage [MB T-126 (145)] was destroyed in the 1978 fire. Five syntypes of *Melamphaes triceratops* Roule & Angel 1933 caught off west of Flores (Azores) are in the MOM collections (MOM-IC-0091-1394 (set of three specimens), MOM-IC-0091-1530 and MOM-IC-0091-1686). Both names are synonyms of *P. capito*.

***Poromitra crassiceps* (Günther 1878)**

Common name: **Crested bigscale; Peixe-rã-de-crista**, Peixe-rã, Rã.

References: mainland Portugal (?) (Bragança, 1903; Maul, 1986a), Azores (?) (MOM-IC-0091-728, 1905; Zugmayer, 1911a, b; Fowler, 1936; Belloc, 1949; Koefoed, 1953; Osório de Castro, 1967; Ebeling and Weed III, 1973; Maul, 1979d, 1986a; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Kotlyar, 2004; Sutton et al., 2008; Porteiro et al., 2010; Carneiro et al., 2014) and Madeira (?) (ZMH 109494, 1971(?); Maul, 1986a; Carneiro et al., 2014; Froese and Pauly, 2019). According to Kotlyar (2009b) the occurrence records and quotes of *P. crassiceps* from the north-eastern Atlantic may be related to *Poromitra nigriceps* (Zugmayer 1911) and to determine the correct identification of Atlantic specimens their re-examination is required. Thus, this species is not considered in the total number of valid species.

***Poromitra megalops* (Lütken 1878)**

Common name: Bigscale; **Peixe-rã-olhudo**, Peixe-rã, Rã.

References: mainland Portugal (SOC Discovery 1103601_84_FIS_658002, 1984; Maul, 1986a; Carneiro et al., 2014 (PECS)) and Azores (Lütken, 1878; Günther 1887; Goode and Bean, 1895, 1896; Brauer, 1906; Roule and Angel, 1930; Fowler, 1936a,b; Belloc, 1938; Koefoed, 1953; Osório de Castro, 1967; Ebeling and Weed, 1973; Nielsen, 1974; Maul, 1979d, 1986a; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Kotlyar, 2004, 2010; Sutton et al., 2008; Carneiro et al., 2014) and Madeira (Maul, 1986a). The holotype (unique) of *Melamphaes megalops* Lütken 1878, synonym of *P. megalops*, caught off south of Azores (stomach content) is in the ZMUC collections (ZMUC 84).

***Poromitra nigriceps* (Zugmayer 1911)**

Common name: No English common name; **Peixe-rã-de-cabeça-preta**, Peixe-rã, Rã.

References: mainland Portugal (Maul, 1986a; Carneiro et al., 2014; Froese and Pauly, 2019), Azores (Zugmayer, 1911a, b; Kotlyar, 2009b; Carneiro et al., 2014; Porteiro et al., 2017; Froese and Pauly, 2019) and Madeira (ZMH 109494; Maul, 1986a; Carneiro et al., 2014). One syntype of this species caught off in the Azores is in the MOM collections (MOM-IC-0091-0728).

***Scopeloberyx opisthotterus* (Parr 1933)**

Common name: Bigscale; **Peixe-rã-delgado**, Peixe-rã, Rã.

References: mainland Portugal (IFREMER Biocean 34576, 1969; Carneiro et al., 2014), Azores (MCZ 31621, 1928; Geistdoerfer et al., 1971; Maul, 1986a (PECS); Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Sutton et al., 2008; Carneiro et al., 2014) and Madeira (SOC Discovery 826300_8_FIS_777001, 1973; Quéro, 1982a; Carneiro et al., 2014).

***Scopeloberyx robustus* (Günther 1887)**

Common name: Longjaw bigscale; **Peixe-rã-robusto**, Peixe-rã, Rã.

References: mainland Portugal (MOM-IC-0091-1179, 1910; Zugmayer, 1911a, b; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1979d, 1986a, 1990; Carneiro et al., 2014), Azores (Roule, 1919a; Fowler, 1936a; Koefoed, 1953; Albuquerque, 1954-1956; Grey, 1956; Osório de Castro, 1967; Maul, 1986a; Arruda, 1997; Santos et al., 1997; Sutton et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Almeida and Biscoito, 2019 (PECS)) and Madeira (ZMH 110659, 1971; SOC Discovery 826300_8_FIS_777002, 1973; Maul, 1986a; Carneiro et al., 2014). The holotype (unique) of *Scopeloberyx opercularis* Zugmayer 1911, synonym of *S. robustus*, caught off in mainland Portugal is in the MOM collections (MOM-IC-0091-1179).

***Scopeloberyx rubriventer* (Koefoed 1953)**

Common name: Ridgehead; **Peixe-rã-dos-Açores**, Peixe-rã, Rã.

References: Azores (Koefoed, 1953; Ebeling and Weed, 1973; Maul, 1979d, 1986a; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Moore et al., 2003; Kotlyar, 2004; Porteiro et al., 2010; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019).

***Scopelogadus beanii* (Günther 1887)**

Common name: Bean's bigscale; **Peixe-rã-de-Bean**, Peixe-rã, Rã.

References: mainland Portugal ([Nobre, 1935](#); [Quéro, 1982b](#); [Maul, 1986a](#); [Smith-Vaniz et al., 2015a](#)), Azores ([MCZ 31655, 1928](#); [Borodin, 1931](#); [Ebeling and Weed, 1963](#); [Maul, 1986a](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Smith-Vaniz et al., 2015a](#); [Froese and Pauly, 2019](#)) and Madeira ([Maul, 1986a](#); [Carneiro et al., 2014](#); [Smith-Vaniz et al., 2015a](#); [Froese and Pauly, 2019](#)).

***Scopelogadus mizolepis* (Günther 1878)**

Common name: Ragged bigscale; **Peixe-rã-couraça**, Peixe-rã, Rã.

References: mainland Portugal ([Harold, 2015j](#)), Azores ([MCZ 42891, 1928](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017 \(PECS\)](#); [Carneiro et al., 2014](#); [Harold, 2015j](#)) and Madeira ([Harold, 2015j](#)). According to Kotlyar (2004) this species can be found in the Atlantic Ocean between 40°N and 20°S and Eschmeyer *et al.* (2019) mention it as a circumglobal species in tropical seas, except in the Eastern Pacific. Specimens are needed for confirmation in mainland Portugal and Madeira.

Family Stephanoberycidae

***Acanthochaenus luetkenii* Gill 1884**

Common name: Pricklefish; **Peixe-de-coroa-espinhosa**.

References: Azores ([IFREMER Biocean 34988, 1969](#); [MNHN-IC-1987-0462, 1969](#); [Geistdoerfer et al., 1971](#); [Nielsen, 1979g](#); [Maul, 1986a](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Porteiro et al., 2017](#)). Porteiro *et al.* (2017) indicate the occurrence of this species off Madeira without indication of geographic coordinates. Specimens are needed for confirmation in Madeira Archipelago.

Family Rondeletiidae

***Rondeletia bicolor* Goode & Bean 1895**

Common name: Hooked redmouth whalefish; **Peixe-baleia-bicolor**.

References: Azores (PECS) ([MCZ 62168, 1984](#); [Carneiro et al., 2014](#)).

***Rondeletia loricata* Abe & Hotta 1963**

Common name: Redmouth whalefish; **Peixe-baleia-de-boca-vermelha**.

References: Azores ([Paxton, 1973](#); [ZMH 118502, 1979 \(PECS\)](#); [ZMH 121514, 1982 \(PECS\)](#); [Santos et al., 1997](#); [MAR-ECO 3567, 2004](#); [Sutton et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Paxton and Trnski, 2016a](#)) and Madeira ([ZMH 110544, 1971](#); [NHMUK 1995.7.25.69, 1972 \(PECS\)](#); [Herring 1976 \(PECS\)](#); [Paxton, 1986 \(PECS\)](#); [Carneiro et al., 2014](#); [Paxton and Trnski, 2016a](#)).

Family Barbourisiidae

***Barbourisia rufa* Parr 1945**

Common name: Velvet whalefish; **Peixe-baleia-veludo**.

References: mainland Portugal ([Harold, 2015k](#)) and Azores ([pers. comm. Reiner: occurrence of one specimen off São Miguel island, April 2006](#); [Harold, 2015k](#)).

Family Cetomimidae

***Ataxolepis apus* Myers & Freihofer 1966**

Common name: No English common name; **Peixe-baleia-mosaico**.

References: Azores ([Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Roa-Varón and Iwamoto, 2019a](#)).

***Cetichthys indagator* (Rofen 1959)**

Common name: No English common name; **Peixe-baleia-do-Galathea**, Peixe-baleia.

References: Madeira ([Vieira, et al., 2012](#); [Carneiro et al., 2014](#)).

***Cetomimus gilli* Goode & Bean 1895**

Common name: Gill's flabby whalefish; **Peixe-baleia-de-Gill**.

References: Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF_MAR-ECO_5914, 1979; Carneiro et al., 2014](#)) and Madeira ([Paxton and Trnski, 2016b](#)). According to Eschmeyer *et al.* (2019) *C. gillii* is a cosmopolitan species. The distribution map of this species reported by Paxton and Trnski (2016b) partially included the area of Selvagens Islands (Madeira EEZ). However, specimens are needed for confirmation in Madeira Archipelago.

***Cetomimus hempeli* Maul 1969**

Common name: Whalefish; [Peixe-baleia-de-Hempel](#).

References: Madeira ([Maul, 1969b\(?\)](#); Paxton, 1979, 1986; Paxton and Blake, 1990; Lloris *et al.*, 1991; NHMUK 1996.2.14.38, 1995; Carneiro *et al.*, 2014; Paxton and Trnski, 2016b; Froese and Pauly, 2019). The holotype and paratype of this species are in the MMF collections (MMF 22568 and MMF 10474, 1956, respectively). The paratype was obtained in the Funchal fish market, from the stomach of a black scabbardfish caught off in Madeira.

***Cetostoma regani* Zugmayer 1914**

Common name: **Pink flabby whalefish**; [Peixe-baleia-rosa](#), Peixe-baleia.

References: mainland Portugal ([YPM ICH 004899, 1959](#); [YPM ICH 004900, 1959](#); Bertelsen, 1986b; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores ([NHMUK 2010.10.27.7, 1981](#) (PECS); Bertelsen, 1986b; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Paxton and Trnski, 2016b; Porteiro *et al.*, 2017; Froese and Pauly, 2019) and Madeira ([ZMH 105982, 1968](#); Bertelsen, 1986b; Paxton, 1979, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Paxton and Trnski, 2016b; Froese and Pauly, 2019).

***Ditropichthys storeri* (Goode & Bean 1895)**

Common name: No English common name; [Peixe-baleia-de-Storer](#), Peixe-baleia.

References: Azores (PECS) ([NHMUK 2004.2.19.48, 1958](#); [NHMUK 2004.9.3.2, 1985](#); Carneiro *et al.*, 2014; Harold, 2015I; Paxton and Trnski, 2016b) and Madeira ([Harold, 2015I](#); Paxton and Trnski, 2016b).

***Eutaeniophorus festivus* (Bertelsen & Marshall 1956)**

Common name: **Festive ribbonfish**; [Peixe-serpentina](#).

References: Azores (PECS) ([Paxton, 1979](#); MCZ 63033, 1984; MCZ 100933, 1984; Carneiro *et al.*, 2014) and Madeira ([Bertelsen, 1990c](#); Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Gyrinomimus myersi* Parr 1934**

Common name: Myers' flabby whalefish; [Peixe-baleia-de-Myers](#).

References: Azores ([MAR-ECO 7346, 2004](#); Carneiro *et al.*, 2014; Paxton and Trnski, 2016b; Porteiro *et al.*, 2017 (PECS)) and Madeira ([Paxton and Trnski, 2016b](#)).

***Mirapinna esau* Bertelsen & Marshall 1956**

Common name: **Hairyfish**; [Peixe-baleia-Esaú](#).

References: Azores (PECS) ([Bertelsen and Marshall, 1956](#); Paxton, 1979; Bertelsen, 1986b; MAR-ECO 7023, 2004; Carneiro *et al.*, 2014; Priede, 2017; Froese and Pauly, 2019). According to Johnson *et al.* (2009) the species *Proctichthys kreffti* Paxton 1989 is synonym of *M. esau*.

Order Trachichthyiformes

Family Trachichthyidae

***Gephyroberyx darwinii* (Johnson 1866)**

Common name: **Darwin's slimehead**; [Olho-de-vidro-imperatriz](#), Imperatriz, Olho-de-vidro, Peixe-vidro.

References: mainland Portugal ([INIP, 1981a](#); OCEANA, 2014), Azores[▲] ([Kukuev, 2002 \(PECS\)](#); IMAR/DOP COLETA 26, 2003; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira[▲] ([NHMUK 1866.6.1.2., 1864 - 1877](#); Johnson, 1866; Günther, 1887; Goode and Bean, 1895, 1896; Osório, 1917; Le Gall, 1932; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Maul, 1954a, 1986b; Nielsen, 1979h; Fischer *et al.*, 1981; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Moore, 2016b; Roberts *et al.*, 2017; Brown *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of *Trachichthys darwinii* Johnson 1866, synonym of *G. darwinii*, caught off in Madeira is in the NHMUK collections (NHMUK 1866.6.1.2).

***Hoplostethus atlanticus* Collett 1889**

Common name: **Orange roughy**; [Olho-de-vidro-laranja](#), Olho-de-vidro, Relógio.

References: mainland Portugal (ICES, 2005; Silva, 2008; Silva *et al.*, 2009; Chaves and Silva, 2013; Carneiro *et al.*, 2014; OCEANA, 2014; Silva and Borges, 2014; Bañón *et al.*, 2016a; Porteiro *et al.*, 2017; Martins and Carneiro, 2018), Azores[▲] (MOM-IC-0091-0101, 1888; Collett, 1889c, 1896a; MOM-IC-0091-0362, 1896; Goode and Bean, 1895, 1896; Richard, 1900, 1910; Brauer, 1906; Roule, 1916, 1919; Fowler, 1936a; Belloc, 1949; Collins, 1954; Albuquerque, 1954-1956; Woods and Sonoda, 1973; Hureau and Monod, 1979; Maul, 1986b; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Melo and Menezes, 2002; Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Moore, 2016b; Roberts *et al.*, 2017; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Moore, 2016b; Delgado *et al.*, 2017). The holotype (unique) of subspecies *Leiogaster atlanticus spinulosus* Roule 1916 caught off at Santa Cruz (Azores) is in the MOM collections (MOM-IC-0091-0362) and was raised to synonym of species *H. atlanticus*.

Hoplostethus cadenati Quéro 1974

Common name: Black slimehead; **Olho-de-vidro-preto**.

References: mainland Portugal (Moura, 2004; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019) and Madeira (Fischer *et al.*, 1981).

Hoplostethus mediterraneus Cuvier 1829

Common name: **Mediterranean slimehead**; **Olho-de-vidro-do-Mediterrâneo**, Alfonsim, Alfonsim-de-natura, Alfonsim-do-alto, Alfoucim, Olho-de-vidro, Pargo-do-alto, Relógio.

References: mainland Portugal[▲] (Bragança, 1902, 1903; Cligny, 1903; Osório, 1909, 1917; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976, 1986b; Fernández *et al.*, 1981; INIP, 1982a; Martins, 1981; Quéro, 1982c; Fischer *et al.*, 1987; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002b; Moura, 2004; ICES, 2005; Sousa *et al.*, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Pakhorukov, 2008; Silva, 2008, 2010, 2011; Ward *et al.*, 2008; Silva *et al.*, 2009; Carneiro *et al.*, 2014; Silva and Borges, 2014; Álvarez, 2015; Bañón *et al.*, 2016a; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (Cligny, n.d.; MNHN-IC-1887-0271, 1883; Hilgendorf, 1888a; Vaillant, 1888; Regan, 1903a; Brauer, 1906; Koefoed, 1927; Le Gall, 1932; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Woods and Sonoda, 1973; Wheeler, 1978; INIP, 1985b; Dias and Monteiro, 1986; Dias, 1991; Arruda, 1992, 1997; Santos *et al.*, 1994a, 1997; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Melo and Menezes, 2002; Pakhorukov, 2008 (PECS); Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Álvarez, 2015; Moore, 2016b; Almeida and Biscoito, 2019) and Madeira (Lowe, 1839, 1843-1860; Günther, 1859, 1887; Goode and Bean, 1895, 1896; Brauer, 1906; Le Gall, 1932; Nobre, 1935; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Maul, 1954a, 1976, 1986b; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Lloris *et al.*, 1991; Pakhorukov, 2008 (PECS); Carneiro *et al.*, 2014; Álvarez, 2015; Moore, 2016b; Delgado *et al.*, 2017; Froese and Pauly, 2019). One specimen caught off at the coast of mainland Portugal in 1967 is in the MMF collections (MMF 22509), a second specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005175).

Family Diretmidae

Diretmichthys parini (Post & Quéro 1981)

Common name: **Parin's spinyfish**; **Peixe-enfadado-de-Parin**.

References: mainland Portugal (Monteiro *et al.*, 2001; ICES, 2005), Azores (Kukuev, 2002 (PECS); Melo and Menezes, 2002; Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Ralph, 2015k) and Madeira (Post, 1986a; Lloris *et al.*, 1991; Quéro *et al.*, 1997; Arronte and Heredia, 2006; Carneiro *et al.*, 2014; Ralph, 2015k; Froese and Pauly, 2019).

Diretmoides pauciradiatus (Woods 1973)

Common name: **Silver spinyfin**; **Peixe-enfadado-barbatana-comprida**.

References: mainland Portugal (Moura, 2004) and Madeira (Post, 1986a(?); Lloris *et al.*, 1991; Delgado *et al.*, 2017). According to Eschmeyer *et al.* (2019) the current geographic distribution of *D. pauciradiatus* is tropical and temperate waters in all oceans except Eastern Pacific. However, the

occurrence in the southern coast of Portugal (Algarve) reported by Moura (2004) may indicate its northernmost distribution limit in the Eastern Atlantic.

Diretmus argenteus Johnson 1864

Common name: **Silver spinyfin; Peixe-enfadado-prateado**.

References: mainland Portugal ([Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Post, 1986a](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Roule, 1919a](#); [Zugmayer, 1933](#); [Fowler, 1936b](#); [Koefoed, 1953](#); [Albuquerque, 1954-1956](#); [Woods and Sonoda, 1973](#); [Post and Quéro, 1981](#); [Post, 1986a](#); [Hamer et al., 1994](#); [Clarke et al., 1995](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Borges et al., 2002](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1863.12.12.3, 1861-1890](#); [Johnson, 1864a, b](#); [Goode and Bean, 1895, 1896](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Maul, 1949c, d, 1954a](#); [Albuquerque, 1954-1956](#); [Hureau and Monod, 1979](#); [Post, 1986a](#); [Carneiro et al., 2014](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in Madeira is in the NHMUK collections ([NHMUK 1863.12.12.3](#)).

Family Anoplogasteridae

Anoplogaster cornuta (Valenciennes 1833)

Common name: **Common fangtooth; Peixe-ogre**.

References: mainland Portugal ([Osório, 1917](#); [ISH 57-1966, 1966](#); [Post, 1986b](#); [Carneiro et al., 2014](#)), Azores ([Borodin, 1931](#); [Zugmayer, 1933](#); [Woods and Sonoda, 1973](#); [Post, 1986b](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017](#); [Melo and Menezes, 2002](#); [Sutton et al., 2008](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Maul, 1954a](#); [Nielsen, 1979g](#); [Quéro, 1982b \(PECS\)\(?\)](#); [Post, 1986b](#); [Lloris et al., 1991](#); [Davies and Merrett, 1998](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Order Holocentriformes

Family Holocentridae

Corniger spinosus Agassiz 1831

Common name: **Spinycheek soldierfish; Peixe-soldado-espinhoso**.

References: Madeira ([Moore et al., 2015b](#); [Greenfield, 2016](#)). The distribution map of this species reported by Moore et al. (2015b) includes the Selvagens Islands area.

Sargocentron hastatum (Cuvier 1829)

Common name: **Red squirrelfish; Esquilo-real**, Esquilo, Peixe-esquilo, Peixe-rainha, Peixe-rei, Peixe-soldado, Rainha, Rei, Soldado.

References: mainland Portugal ([Jordano and Muruve, 1959](#); [Nielson, 1979f](#); [Ben-Tuvia, 1986a](#); [Carneiro et al., 2014](#); [Knudsen et al., 2015e](#); [Froese and Pauly, 2019](#)).

Order Ophidiiformes

Family Ophidiidae

Abyssobrotula galatheae Nielsen 1977

Common name: No English common name; **Abadejo-abissal**.

References: mainland Portugal ([NHMUK 1991.7.9.869, 1989](#)), Azores (PECS) ([Knudsen, 2015c](#)) and Madeira ([Knudsen, 2015c](#)). Specimens are needed for confirmation in the Azores and Madeira regions.

Acanthonus armatus Günther 1878

Common name: Bony-eared assfish; **Abadejo-espinhoso**.

References: Azores ([Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Knudsen, 2015d](#)).

Barathrites iris Zugmayer 1911

Common name: No English common name; **Abadejo-arco-íris**.

References: mainland Portugal ([Nielsen, 2015b](#)), Azores ([Zugmayer, 1911a](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Arruda, 1997](#); [Eschmeyer 1997](#); [Carneiro et al., 2014](#); [Nielsen, 2015b](#)) and Madeira ([Nielsen, 1979e](#), [1986b](#), [2015b](#); [Armstrong et al., 1992](#); [Froese and Pauly, 2019](#)). This species occurs SW of Azores,

out of the PECS area (723 NM South-west of Flores Island, 31°44.50'N, 42°39.00'W, Zugmayer, 1911a; MOM-IC-0091-905, 1905, as holotype of *B. iris*) and 175 NM ENE of Great Meteor Tablemount, 30°42.0' N, 25°12.0'W (MOM-IC-0091-436, 1897, as holotype of *Barathrites abyssorum* Roule 1916) and was erroneously considered present in the Azores (Carneiro *et al.*, 2014). However, Nielsen (2015b) includes the Azores in the distribution map.

***Bassozetus compressus* (Günther 1878)**

Common name: Abyssal assfish; [Abadejo-da-fundura](#).

References: mainland Portugal (?) ([NHMUK 1990.8.21.141](#), 1983; Carneiro *et al.*, 2014) and Madeira (?) ([NHMUK 1994.5.6.3](#), 1984; [NHMUK 1994.5.9.13](#), 1990 (PECS); Carneiro *et al.*, 2014). According to Tomiyama *et al.* (2016) and Eschmeyer *et al.* (2019) the current geographic distribution of *B. compressus* is from New Caledonia to Okinawa Islands. The previously occurring records that reported *B. compressus* to mainland Portugal and Madeira (Carneiro *et al.* 2014) were re-evaluated by Tomiyama *et al.* (2018) who consider this species as *B. nielseni*. Thus, this species is not considered in the total number of valid species.

***Bassozetus nielseni* Tomiyama, Takami & Fukui 2018**

Common name: Masked assfish; [Abadejo-de-Nielsen](#).

References: mainland Portugal ([NHMUK 1990.8.21.141](#), 1983; [NHMUK 1991.7.9.734](#), 1983; Tomiyama *et al.*, 2018) and Madeira ([NHMUK 1994.5.6.3](#), 1984; [NHMUK 1994.5.9.13](#), 1990 (PECS); Tomiyama *et al.*, 2018). According to Tomiyama *et al.* (2018) the current geographic distribution of *B. nielseni* is the North Atlantic and West Indian oceans.

***Bassozetus levistomatus* Machida 1989**

Common name: No English common name; [Abadejo-boca-doce](#).

References: Madeira (?) (PECS) ([NHMUK 1991.7.9.859](#), 1990; Carneiro *et al.*, 2014). The original occurrence record of this specimen indicated cf. *bassozetus* sp. (Zoology Accessions Register: Fishes: 1986 - 1994 of the Natural History Museum, London, page 109). Subsequently this occurrence record was modified to *Bassozetus levistomatus* Machida 1989 (Carneiro *et al.*, 2014) and on 30 November 2017 to *Bassozetus compressus* (Günther 1878). According to Tomiyama *et al.* (2018) specimens previously reported as *B. compressus* from the North Atlantic are now considered *B. nielseni*. The specimen 1991.7.9.859 was not included in the review made by Tomiyama *et al.* (2018). Thus, this species is not considered in the total number of valid species.

***Bassozetus taenia* (Günther 1887)**

Common name: No English common name; [Abadejo-ténia](#).

References: Madeira (PECS) ([Roule 1935](#); Grey, 1956; [NHMUK 1991.7.9.860](#), 1990; Carneiro *et al.*, 2014). Roule (1935) reported this species in Madeira, but without description or depth data (*in* Grey, 1956).

***Bathyonus laticeps* (Günther 1878)**

Common name: No English common name; [Abadejo-cabeçudo](#).

References: Azores ([Koefoed, 1932](#) (PECS); Grey, 1956 (PECS); Nielsen, 1979e(?), 2015c; Carneiro *et al.*, 2014 (PECS);) and Madeira ([Nielsen, 1986b](#); [NHMUK 2003.6.11.18](#), 1989; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Nielsen, 2015c; Froese and Pauly, 2019).

***Benthocometes robustus* (Goode & Bean 1886)**

Common name: No English common name; [Abadejo-cometa](#).

References: mainland Portugal ([ICM-CSIC 7765](#), n.d.; Moura, 2004; ICES, 2005; Silva, 2008; Carneiro *et al.*, 2014; OCEANA, 2014; Uiblein *et al.*, 2015d), Azores ([Santos *et al.*, 1997](#)(?); Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Uiblein *et al.*, 2015c; Gomes-Pereira *et al.*, 2017) and Madeira ([Uiblein *et al.*, 2015d](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *B. robustus* is the Mediterranean Sea and the Atlantic. The distribution map of this species reported by Uiblein *et al.* (2015d) partially included the area of Selvagens Islands (Madeira EEZ). However, specimens are needed for confirmation its occurrence in Madeira archipelago.

***Brotulotaenia brevicauda* Cohen 1974**

Common name: No English common name; [Abadejo-de-cauda-curta](#).

References: Azores (IMAR/DOP COLETA 1692, 2001; Melo and Menezes, 2002; Porteiro *et al.*, 2010; Menezes *et al.*, 2012; Carneiro *et al.*, 2014).

***Brotulotaenia crassa* Parr 1934**

Common name: Blue cusk eel; **Abadejo-grosso**.

References: mainland Portugal (Nielsen, 1986b; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019), Azores (Kukuev, 2002 (PECS); Melo and Menezes, 2002; MAR-ECO 7666, 2004; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014) and Madeira (Nielsen, 1979e, 1986b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017; Froese and Pauly, 2019).

***Brotulotaenia nigra* Parr 1933**

Common name: Dark cusk; **Abadejo-negro**.

References: Azores (PECS) (SOC Discovery 1023328_86_FIS_078001, 1980; Carneiro *et al.*, 2014) and Madeira (Nielsen, 1979e, 1986b; Lloris *et al.*, 1991; Nielsen *et al.*, 1999; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Holomycteronus squamosus* (Roule 1916)**

Common name: No English common name; **Abadejo-escamudo**.

References: mainland Portugal (Christiansen *et al.*, 2014), Azores (MOM-IC-0091-325, 1896; Roule, 1915, 1916, 1919; Fowler, 1936b; Belloc, 1949; Albuquerque, 1954-1956; Grey, 1956; Nielsen, 1986b, 2015d; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Bergstad *et al.*, 2008a (PECS); Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Nielsen, 2015d). The holotype (unique) *Grimaldichthys squamosus* Roule 1916, synonym of *H. squamosus*, caught off near Santa Maria (Azores) is in the MOM collections (MOM-IC-0091-0235).

***Lamprogrammus niger* Alcock 1891**

Common name: No English common name; **Abadejo-cristado**.

References: Azores (PECS) (MAR-ECO 2703, 2004; Bergstad *et al.*, 2008a; Carneiro *et al.*, 2014; Kobylansky and Orlov, 2015; Porteiro *et al.*, 2017).

***Monomitopus metriostoma* (Vaillant 1888)**

Common name: No English common name; **Abadejo-da-Mauritânia**.

References: mainland Portugal (Nielsen, 1986b, 2016a; Carneiro *et al.*, 2014) and Azores (MNHN-IC-1886-0521, 1883; Vaillant, 1888; Goode and Bean, 1895, 1896; Roule and Angel, 1921; Fowler, 1936b; Albuquerque, 1954-1956; Nielsen, 1979e, 1986b, 1990; Hureau and Nielsen, 1981; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The paratype of *Sirembo metriostoma* Vaillant 1888, synonym of *M. metriostoma*, caught off between Pico and São Jorge Islands (Azores) is in the MNHN collections (MNHN-IC-1886-0521).

***Ophidion barbatum* Linnaeus 1758**

Common name: **Snake blenny; Abadejo-cobrelo**, Barbudo, Cobrelo, Cobrelo-barbudo, Peixe-cobrelo, Peixe-cobrelo-barbudo.

References: mainland Portugal (Bragança, 1902, 1903; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nielsen, 1986b, 1990, 2016a; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro *et al.*, 2014).

***Parophidion vassali* (Risso 1810)**

Common name: No English common name; **Abadejo-vermelho**.

References: Azores (Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997; Matallanas and Casadevall, 1999; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira (Matallanas and Casadevall, 1990, 1999; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019(?)).

***Penopus microphthalmus* (Vaillant 1888)**

Common name: No English common name; **Abadejo-de-olho-pequeno**.

References: Azores (PECS) (MAR-ECO 12655, 2004; Bergstad *et al.*, 2008a; Carneiro *et al.*, 2014; Porteiro *et al.*, 2017).

Porogadus miles Goode & Bean 1885

Common name: Slender cuskeel; **Abadejo-fino**.

References: mainland Portugal ([Knudsen, 2015e](#)), Azores ([EMBL KX150868, 2004 \(PECS\)](#); [EMBL KX150923, 2004 \(PECS\)](#); [MAR-ECO 2661, 2004 \(PECS\)](#); [Bergstad et al., 2008a \(PECS\)](#); [Carneiro et al., 2014 \(PECS\)](#); [Knudsen, 2015e](#); [Porteiro et al., 2017 \(PECS\)](#)) and Madeira ([Knudsen, 2015e](#)). The previous quote by Carneiro et al. (2014) is updated to the Azorean PECS area. Specimens are needed for confirmation in mainland Portugal and Madeira regions.

Spectrunculus crassus (Vaillant 1888)

Common name: No English common name; **Abadejo-dos-Açores**.

References: Azores (PECS)([Grey, 1956](#) ; [MAR-ECO 8067, 2004](#); [MAR-ECO 8081, 2004](#); [MAR-ECO 2679, 2004](#); [MAR-ECO 13053, 2004](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)). Vaillant (1888) indicates the occurrence of *Bythites crassus* in the North-eastern Atlantic between Azores and France, 44°20.0'N, 17°11.0'W, depth 4255 meters [holotype (unique) in the MNHN collections (MNHN-IC-1886-0552)]. This geographic coordinates correspond to a position near the Azores - Biscay Rise, but out of the PECS area. Uiblein et al. (2008) demonstrated the overlapping occurrence of two closely related sister species, *S. crassus* and *S. grandis*, in the lower bathyal and abyssal of the Atlantic. According to Eschmeyer et al. (2019) the actual geographic distribution of this species is the Eastern-north Atlantic. The previous quote by Carneiro et al. (2014) is updated to the Azorean PECS area.

Spectrunculus grandis (Günther 1877)

Common name: Pudgy cuskeel; **Abadejo-grande**.

References: mainland Portugal ([GNM 1946-6250, 1946](#); [Nielsen, 1986b](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1976-0046, 1971](#); [ZMH 119213, 1979](#); [Nielsen and Hureau, 1980](#); [Hureau and Nielsen, 1981](#); [Nielsen, 1986b](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bergstad et al., 2008a \(PECS\)](#); [Menezes et al., 2009, 2012](#); [Porteiro et al., 2010, 2017 \(PECS\)](#); [Carneiro et al., 2014](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Nielsen, 1986b](#); [Armstrong et al., 1992](#); [Carneiro et al., 2014](#); [NHMUK 2017.3.9.38, 2017](#); [Froese and Pauly, 2019](#)). According to Eschmeyer et al. (2019) *S. grandis* is a cosmopolitan species. The co-existence of closely related species, *S. crassus* and *S. grandis* (Uiblein et al., 2008), may have skewed the respective geographic distributions in the Mid-Atlantic Ridge. Thus, the actual geographic distribution of this species may be masked by information relating to *S. crassus* (Porteiro et al., 2017).

Family Carapidae

Carapus acus (Brünnich 1768)

Common name: Pearl fish; **Peixe-pérola-cobrelo**, Cobrelo, Peixe-cobrelo, Peixe-pérola, Pérola.

References: mainland Portugal ([Bragança, 1903](#); [Osório, 1917](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Carneiro et al., 2014](#); [Uiblein et al 2015e](#); [Eschmeyer et al., 2019](#)), Azores ([Collins, 1954](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Uiblein et al 2015e](#)) and Madeira ([Lowe, 1843](#); [MOM-IC-0091-1723, 1897](#); [Vaillant, 1919](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Maul, 1949](#); [Albuquerque, 1954-1956](#); [Arnold, 1956](#); [Osório de Castro, 1967](#); [Wheeler, 1979b](#); [Trott and Olney, 1986](#); [Lloris et al., 1991](#); [Wirtz, 1994](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Uiblein et al 2015e](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Carapus birpex* Arnold 1956, synonym of *C. acus*, caught off near Madeira is in the MMF collections (MMF 2739).

Echiodon dentatus (Cuvier 1829)

Common name: No English common name; **Peixe-pérola-dentado**.

References: mainland Portugal ([Uiblein et al., 2015f](#)), Azores (PECS) ([SDSC 9309, 1967](#); [SDSC 23049, 1967](#); [SDSC 23050, 1967](#); [SDSC 23051, 1967](#); [Maul, 1976](#); [Ehrich, 1977](#); [SDSC 9310, 1998](#); [Fock et al., 2002](#); [Kukuev, 2002](#); [Carneiro et al., 2014](#); [Uiblein et al., 2015f](#)) and Madeira (PECS) ([SDSC 23048, 1967](#); [Maul, 1976](#); [Carneiro et al., 2014](#); [Uiblein et al., 2015f](#)). Several specimens caught off in different areas are in the MMF collections: two specimens at the Josephine Bank (MMF 22806, 1967) and at the Meteor Bank (MMF 22377, 1967, set of four specimens, MMF 22875, 1967, one specimen and MMF 22382, 1967, set of two specimens).

Echiodon drummondii Thompson 1837

Common name: No English common name; **Peixe-pérola-de-Drummond**.

References: Azores ([MCZ 149546, 1978](#); [Markle and Olney, 1990](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#)).

Family Bythitidae

Aphyonus gelatinosus Günther 1878

Common name: Gelatinous blindfish; **Abadejo-cego-gelatinoso**, Peixe-cego-gelatinoso.

References: Azores ([MNHN-IC-1973-0032, 1971](#); [Rannou et al., 1974](#); [Nielsen, 1986b, 2015e](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)).

Barathronus multidens Nielsen 1984

Common name: No English common name; **Abadejo-cego-dentudo**.

References: mainland Portugal (?) ([Carneiro et al., 2014](#)). According to Evseenko et al. (2018) and Eschmeyer et al. (2019) the actual geographic distribution of this species is the Western and eastern-north Atlantic and Mid-Atlantic Ridge. *B. multidens* is known from only a few specimens in the Atlantic Ocean and was erroneously considered present in mainland Portugal ([Carneiro et al., 2014](#)). Thus, this species is not considered in the total number of valid species for mainland Portugal.

Barathronus parfaiti (Vaillant 1888)

Common name: No English common name; **Abadejo-cego-de-Vaillant**.

References: Azores ([MNHN-IC-1886-0554, 1883 \(PECS\)](#); [Vaillant, 1888 \(PECS\)](#); [Brauer, 1906 \(PECS\)](#); [Roule, 1915, 1916, 1919a](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Nielsen, 1969, 1986b](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Nielsen et al., 2015f \(PECS\)](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Trachichthys darwinii* Johnson 1866, synonym of *B. parfaiti*, caught off in north-eastern Atlantic between Azores and France is in the MNHN collections ([MNHN-IC-1886-0554](#)).

Barathronus roulei n. sp.

Common name: No English common name; **Abadejo-cego-de-Roule**.

References: Azores ([Nielsen, 2019](#)). The holotype of this species was caught 23 NM SE off the Flores islands (Azores) and is preserved in the MOM collections (MOM POI-0003655, 1896).

Bellottia apoda Giglioli 1883

Common name: No English common name; **Brótula-coxa**.

References: mainland Portugal ([Maul, 1976](#); [Nielsen, 1986b, 1990b](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([PECS](#)) ([SDSC 23047, 1967](#); [Maul, 1976](#); [Nielsen, 1986b](#); [Nielsen et al., 1999](#); [Kukuev, 2002](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Nielsen, 1986b](#); [Gramitto and Coen, 1997](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). Five specimens caught at the coast of mainland Portugal and Meteor Bank are in the MMF collections ([MMF 22516, 1967](#), set of two specimens, [MMF 22860, 1967](#), set of two specimens and [MMF 22845, 1967](#), one specimen, respectively).

Cataetyx allenii (Byrne 1906)

Common name: No English common name; **Brótula-de-Allen**.

References: mainland Portugal ([Nielsen, 1986b](#); [Carneiro et al., 2014](#); [Uiblein et al., 2015g](#); [Bañón et al., 2016a](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype of *Grammonus leucos* Osório 1917 caught off in coast of Morocco and preserved in the Museu Bocage was destroyed in the 1978 fire. The current status of *Grammonus leucos* is uncertain as *C. allenii*.

Cataetyx laticeps Koefoed 1927

Common name: No English common name; **Brótula-de-cabeça-chata**.

References: mainland Portugal ([Christiansen et al., 2014](#); [Knudsen et al., 2015f](#)) and Azores ([MNHN-IC-1976-0066, 1971](#); [MNHN-IC-1976-0068, 1971](#); [Hureau and Nielsen, 1981](#); [Nielsen, 1986b](#); [Saldanha, 1994](#); [Van Dover et al., 1996](#); [Saldanha and Biscoito, 1997](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Bañón, 2001](#); [Desbruyères et al., 2001](#); [Melo and Menezes, 2002](#); [Bergstad et al., 2008a \(PECS\)](#); [Porteiro et al., 2008](#)).

2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Knudsen *et al.*, 2015f; Bañón *et al.*, 2016a; Almeida and Biscoito, 2019; Froese and Pauly, 2019).

***Grammonus ater* (Risso 1810)**

Common name: No English common name; **Brótula-negra**.

References: mainland Portugal (?) ([Nielsen *et al.*, 2014](#)) and Azores (?) ([Roule and Angel, 1930](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Arruda, 1997](#); [Carneiro *et al.*, 2014](#)). According to [Psomadakis *et al.* \(2012\)](#) and [Eschmeyer *et al.* \(2019\)](#) the current geographic distribution of *G. ater* is the Mediterranean Sea (endemic). However, [Nielsen *et al.* \(2014\)](#) referred the presence of this species in the Portuguese mainland coast, but this remains unverified. Several authors ([Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Arruda, 1997](#); [Carneiro *et al.*, 2014](#)) mentioned the occurrence of this species at the Azores, based on the description of post-larvae of *Pteridium atrum* by [Roule and Angel \(1930\)](#). [Cohen \(1964b\)](#) stated that the register for the Azores is due to misidentification of *Oligopus ater*. Thus, this species is not considered in the total number of valid species to the Azores and mainland Portugal.

***Leucobrotula adipata* Koefoed 1952**

Common name: No English common name; **Brótula-falsa-branca**.

References: mainland Portugal ([GNM 1946-6619, 1946](#); [YPM ICH 005187, 1959](#); [Nielsen, 1986b](#); [Carneiro *et al.*, 2014](#)), Azores ([Nielsen, 1986b](#); [Nielsen *et al.*, 1990](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)).

***Meteoria erythrops* Nielsen 1969**

Common name: No English common name; **Abadejo-cego-dos-Açores**.

References: Azores ([Nielsen, 1969, 1979f](#); [Lloris *et al.*, 1991](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#)). The holotype (unique) of this species is in the ZMH collections (ZMH 25142) and was caught off in the north-eastern Atlantic off Galiza (Spain). However, Nielsen (1969) refers two specimens caught off near the Azores.

***Meteoria longidorsalis* Nielsen 2016**

Common name: No English common name; **Abadejo-cego-de-dorsal-longa**.

References: Madeira ([NHMUK 2015.5.27.1, 1984](#); [Nielsen, 2016b](#)). One paratype of this species caught off 130 NM NW of Madeira is in the NHMUK collections (NHMUK 2015.5.27.1).

***Nybelinella brevianalis* Nielsen 2017**

Common name: No English common name; **Abadejo-cego-da-Madeira**.

References: Madeira (PECS) ([NHMUK 2016.7.29.11, 1990](#); [Nielsen, 2017](#)). The holotype (unique) of this species caught off in Madeira, in the PECS area (217 NM WSW) is in the NHMUK collections (NHMUK 2016.7.29.11).

***Nybelinella eriksoni* (Nybelin 1957)**

Common name: No English common name; **Abadejo-cego-de-Eriksson**.

References: mainland Portugal ([Van der Land *et al.*, 2001](#); [Carneiro *et al.*, 2014](#)), Azores (PECS) ([NHMUK 2016.7.29.1, 1981](#); [NHMUK 2016.7.29.8, 1985](#); [Carneiro *et al.*, 2014](#)) and Madeira ([NHMUK 2016.7.29.2-3, 1984](#); [NHMUK 2016.7.29.10, 1995](#) (PECS); [Nielsen, 2017](#)).

***Parabrotula plagiophthalma* Zugmayer 1911**

Common name: False cusk; **Brótula-falsa-dos-Açores**.

References: Azores ([Zugmayer, 1911a \(PECS\)](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Nielsen, 1968, 1986b](#); [Nielsen *et al.*, 1990](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([SOC 826300_8_FIS_646001, 1973](#); [Carneiro *et al.*, 2014](#)). The holotype (unique) of this species caught off north-east of Azores (43°2.4'N, 19°25.2'W) in the PECS area is in the MOM collections (MOM-IC-0091-0878).

***Sciadonus pedicellaris* Garman 1899**

Common name: No English common name; **Abadejo-cego-do-Pacifico**.

References: mainland Portugal (PECS) ([ZMH 120293, 1979](#)) and Azores ([GNM 1948-6610, 1948 \(PECS\)](#); [Nybelin, 1951, 1957](#); [Nielsen, 1969](#); [Rannou *et al.*, 1974](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Nielsen and Møller, 2008](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#)). The holotype (unique) of *Sciadonus*

kullenbergi Nybelin 1957 caught off in the Azores is in the GNM collections. According to Nielsen (2018) the species *Sciadonus galatheaee* (Nielsen 1969) is synonym of *S. pedicellaris*.

Order Batrachoidiformes

Family Batrachoididae

Halobatrachus didactylus (Bloch & Schneider 1801)

Common name: **Lusitanian toadfish; Charroco**, Chicharro, Encharrôco, Enxarroco, Manuel-carvalho, Xarrocó.

References: mainland Portugal▲ (Carvalho, 1765; de Brito Capello, 1868, 1869b, 1880; Steindachner, 1868a; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Nobre, 1895a, 1903, 1904, 1935; Bragança, 1902; Osório 1905; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Monod, 1979a; Fischer et al., 1981, 1987; Dooley et al., 1985; Roux, 1986a; Antunes et al., 1988; Halstead et al., 1990; Martins et al., 1992, 2015; Erzini et al., 1996, 2001a; Cabral, 1999; Borges et al., 2002, 2003; Santos et al., 2002a; Palma et al., 2003; Ribeiro et al., 2006, 2008; Stergiou et al., 2006; Henriques et al., 2007, 2008; Gomes and Olim, 2007; Neves et al., 2008; Batista et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Costa et al., 2012; Carneiro et al., 2014; Collette, 2015a; Collette and Greenfield, 2016; Martins and Carneiro, 2018; Bañón et al., 2019b; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Roux, 1986a; Lloris et al., 1991; Wirtz et al., 2008(?); Carneiro et al., 2014; Collette, 2015a; Eschmeyer et al., 2019; Froese and Pauly, 2019(?)).

Order Scombriformes

Family Gempylidae

Diplospinus multistriatus Maul 1948

Common name: **Striped escolar; Ecolar-raiado**.

References: Azores (ZMH 117612, 1979; Bekker et al., 1982; Diekmann et al., 2006 (PECS); Sutton et al., 2008; Carneiro et al., 2014; Ralph, 2015I; Parin and Nakamura, 2016; Froese and Pauly, 2019) and Madeira (NHMUK 1953.10.28.1, 1937-1960; Maul, 1948b; Parin, 1986b, 1990; Lloris et al., 1991; Parin and Nakamura, 1993, 2016; Carneiro et al., 2014; Ralph, 2015I; Froese and Pauly, 2019). The holotype and seven paratypes of this species caught off in Madeira are in the MMF and NHMUK collections [MMF 3063 (holotype), NHMUK 1953.10.28.1 and MMF 3064-69 (set of six specimens), respectively].

Gempylus serpens Cuvier 1829

Common name: **Snake mackerel; Ecolar-serpente**, Coelho-de-natura, Ecolar, Peixe-coelho-de-natura.

References: Azores (Lütken, 1880; Parin, 1986b; Lloris et al., 1991; Nakamura and Parin, 1993; Arruda, 1997; Santos et al., 1997; Arkhipov & Mamedov, 2008 (PECS); Porteiro et al., 2010; Carneiro et al., 2014; Parin and Nakamura, 2016; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Carneiro et al., 2014; Parin and Nakamura, 2016).

Lepidocybium flavobrunneum (Smith 1843)

Common name: **Ecolar; Ecolar-preto**, Bonito; Bonito-do-alto; Ecolar-de-natura, Ecolar-da-fundura.

References: mainland Portugal▲ (Noronha 1926b; Fischer et al., 1981; Nakamura and Parin, 1993; Carneiro et al., 2014; Collette et al., 2015a; Froese and Pauly, 2019), Azores▲ (Nakamura and Parin, 1993; Azevedo and Heemstra, 1995; Santos et al., 1997; Cardigos et al., 2005; Porteiro et al., 2010; Carneiro et al., 2014; Collette et al., 2015a; Parin and Nakamura, 2016; Froese and Pauly, 2019) and Madeira▲ (Noronha 1926; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer et al., 1981; Parin, 1986b; Lloris et al., 1991; Nakamura and Parin, 1993; Carneiro et al., 2014; Collette et al., 2015a; Parin and Nakamura, 2016; Froese and Pauly, 2019). The unique holotype of *Diplogonurus maderensis* Noronha 1926 was caught off near Câmara de Lobos (Madeira). The type, apparently the head (dried?) illustrated in Noronha (1926), was never received by the Carnegie Museum (Henn, 1928).

Nealotus tripes Johnson 1865

Common name: **Black snake mackerel; Escolar-coelho-de-natura**, Coelho-de-natura, Escolar, Peixe-coelho-de-natura.

References: Azores ([MCZ 83334, 1973 \(PECS\)](#); Nakamura and Parin, 1993; Carneiro et al., 2014; Parin and Nakamura, 2016; Froese and Pauly, 2019) and Madeira ([NHMUK 1862.10.3.2, 1861-1890](#); Johnson, 1865a, b; Goode and Bean, 1895, 1896; Brauer, 1906; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; MCZ 83258, 1970; MCZ 83259, 1970; Parin, 1986b; Lloris et al., 1991; Nakamura and Parin, 1993; Carneiro et al., 2014; Parin and Nakamura, 2016; Roberts et al., 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Madeira is in the NHMUK collections ([NHMUK 1862.10.3.2](#)).

Nesiarchus nasutus Johnson 1862

Common name: **Black gemfish; Escolar-geribé**, Coelho-de-natura, Escolar, Espada, Espada-preto, Geribé, Peixe-coelho, Peixe-coelho-de-natura, Peixe-espada, Peixe-espada-preto.

References: mainland Portugal ([de Brito Capello, 1867a, d, 1880](#); Steindachner, 1867; Osório, 1888; Almeida and Roquette, 1892; Girard, 1892; Goode and Bean, 1895, 1896; MNHN-IC-1905-0575, 1904; Brauer, 1906; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Parin, 1986b; Nakamura and Parin, 1993; ICES, 2005; Bordalo-Machado et al., 2009; Carneiro et al., 2014; Smith-Vaniz and Herrera, 2015b; Bañón et al., 2016a; Froese and Pauly, 2019), Azores ([Koefoed, 1927 \(PECS\)](#); MCZ 35343, 1931; Hureau and Monod, 1979 (PECS); Parin, 1986b; Nakamura and Parin, 1993; Arruda, 1997; Santos et al., 1997; IMAR/DOP COLETA 1160, 2007; Porteiro et al., 2010; IMAR/DOP CONDOR-05-O11GES1, 2011; Carneiro et al., 2014; Smith-Vaniz and Herrera, 2015b; Froese and Pauly, 2019) and Madeira ([Johnson, 1862a](#); Steindachner, 1867; Günther, 1887; Goode and Bean, 1895, 1896; Brauer, 1906; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; INIP, 1982c; Parin, 1986b; Lloris et al., 1991; de Casamajor and Morandeau, 2013; Carneiro et al., 2014; Smith-Vaniz and Herrera, 2015b; Roberts et al., 2017). The holotype (unique) of *Prometheus paradoxus* de Brito Capello 1867 was caught off in mainland Portugal but it is lost. *P. paradoxus* is synonym of *N. nasutus*.

Promethichthys prometheus (Cuvier 1832)

Common name: **Roudi escolar; Escolar-branco**, Coelho, Coelho-do-alto, Cuê, Escolar, Peixe-coelho, Peixe-coelho-do-alto.

References: mainland Portugal ([Fischer et al., 1981](#); Carneiro et al., 2014; Herrera and Smith-Vaniz, 2015a), Azores ([Collins, 1954](#); Albuquerque, 1954-1956; Osório de Castro, 1967; Parin, 1986b; Nakamura and Parin, 1993; Arruda, 1997; Santos et al., 1994a, 1995a, 1997; Kukuev, 2002 (PECS); Porteiro et al., 2010; Besugo, 2013; Carneiro et al., 2014; Herrera and Smith-Vaniz, 2015a; Froese and Pauly, 2019) and Madeira[▲] ([Lowe, 1837, 1839](#); Günther, 1860, 1887; Goode and Bean, 1895, 1896; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; INIP, 1980; Fischer et al., 1981; Parin, 1986b; Nakamura and Parin, 1993; Severino, 2004; Carneiro et al., 2014; Herrera and Smith-Vaniz, 2015a; Parin and Nakamura, 2016; POGID, 2016; Froese and Pauly, 2019).

Ruvettus pretiosus Cocco 1833

Common name: **Oilfish; Escolar**, Chocolate, Corvina-da-fundura, Cravo, Escolarinho, Peixe-chocolate, Peixe-cravo, Peixe-escolar, Peixe-óleo, Peixe-sabão, Sabão.

References: mainland Portugal[▲] ([de Brito Capello, 1867a, d, 1880](#); Steindachner, 1867; Osório, 1888; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892(?); Goode and Bean, 1895, 1896; Nobre, 1903, 1935, 1937; Brauer, 1906; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer et al., 1981; INIP, 1982a; Parin, 1986b; Fischer et al., 1987; Nakamura and Parin, 1993; Erzini et al., 2001b; Borges et al., 2002; Santos et al., 2002a, b; ICES, 2005; Fernandes et al., 2007; Gomes and Olim, 2007; Pakhorukov, 2008; Carneiro et al., 2014; Costa, 2014; Smith-Vaniz and Herrera, 2015c; Froese and Pauly, 2019), Azores[▲] ([Sampaio, 1904](#); Collins, 1954; Osório de Castro, 1967; Wheeler, 1978; Martins, 1982a; Santos et al., 1994a, 1995a, 1997; Arruda, 1997; Morato et al., 2001a; Kukuev, 2002 (PECS); Cardigos et al., 2005; Menezes et al., 2006; Porteiro et al., 2010; Besugo, 2013; Carneiro et al., 2014; Smith-Vaniz and Herrera, 2015c; Parin and Nakamura, 2016)

and Madeira[▲] (Lowe, 1833a, 1837, 1839, 1840b, 1844a, 1849; Günther, 1860; de Velloso, 1892; Goode and Bean, 1895, 1896; Brauer, 1906; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; INIP, 1980, 1982c, 1984b; Fischer *et al.*, 1981; Parin, 1986b; Lloris *et al.*, 1991; Nakamura and Parin, 1993; Severino, 2004; Carneiro *et al.*, 2014; Smith-Vaniz and Herrera, 2015c; Parin and Nakamura, 2016; Froese and Pauly, 2019).

Family Trichiuridae

***Aphanopus carbo* Lowe 1839**

Common name: **Black scabbardfish; Peixe-espada-preto**, Espada, Espada-preta, Espada-preto, Jibóia, Peixe-espada, Peixe-espada-preta.

References: mainland Portugal[▲] (de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Günther, 1887; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Goode and Bean, 1895, 1896; Bragança, 1902, 1903; Brauer, 1906; Seabra, 1911; Nobre, 1935; Fowler, 1936b; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Martins, 1981; Parin, 1986b; Nakamura and Parin, 1993; WWF, 2001; Moura, 2004; Figueiredo *et al.*, 2005; Fowler *et al.*, 2005; ICES, 2005; Fernandes *et al.*, 2007; Abecasis *et al.*, 2009; Bordalo-Machado *et al.*, 2009; Silva, 2010; Veiga *et al.*, 2013; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014; Bañón *et al.*, 2016a; Priede, 2017; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores[▲] (Wheeler, 1978; INIP, 1985a; Shcherbachov *et al.*, 1985b (PECS); Parin, 1986b; Nakamura and Parin, 1993; Martins, and Cascalho, 1990; Dias, 1991; Lloris *et al.*, 1991; Monteiro, 1991; Nakamura and Parin, 1993; Silva *et al.*, 1995; Menezes, 1996; Arruda, 1997; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Uiblein *et al.*, 1999 (PECS); Hareide and Garnes, 2001 (PECS); Morato *et al.*, 2001a; Kukuev, 2002 (PECS); Melo and Menezes, 2002; Cardigos *et al.*, 2005; Menezes *et al.*, 2006, 2009, 2012; Pakhorukov, 2008 (PESC); Bordalo-Machado *et al.*, 2009; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Parin and Nakamura, 2016 (PECS); Priede, 2017; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1839, 1840b, 1844a, 1849; NHMUK 1855.11.29.6, 1854-1861; Günther, 1860, 1887; de Velloso, 1892; Goode and Bean, 1895, 1896; Brauer, 1906; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Maul, 1948b, 1949b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Wolff, 1971; Fischer *et al.*, 1981; INIP, 1982c, 1984a, b, c; Parin, 1986b; Lloris *et al.*, 1991; Nakamura and Parin, 1993; Martins and Ferreira 1995; Severino, 2004; Bordalo-Machado *et al.*, 2009; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Parin and Nakamura, 2016; POGID, 2016; Delgado *et al.*, 2017; Porteiro *et al.*, 2017; Priede, 2017; Froese and Pauly, 2019). Two unique holotypes of *Aphanopus carbo* Lowe 1839 and *Aphanopus acus* Maul 1948 both caught off in Madeira are in NHMUK (NHMUK 1855.11.29.6) and MMF collections. *A. acus* is synonym of *A. carbo*.

***Aphanopus intermedius* Parin 1983**

Common name: **Intermediate scabbardfish; Peixe-espada-intermédio**.

References: Azores[▲] (Nakamura and Parin, 1993; Kukuev, 2002 (PECS); Porteiro *et al.*, 2010; Biscoito *et al.*, 2011; Carneiro *et al.*, 2014; Delgado, 2015c; Froese and Pauly, 2019) and Madeira (Parin and Nakamura, 1993, 2016; Biscoito *et al.*, 2011; Carneiro *et al.*, 2014; Delgado, 2015c; POGID, 2016; Froese and Pauly, 2019). This species is sometimes confused with *Aphanopus carbo* Lowe 1839 (Parin and Nakamura, 2016).

***Benthodesmus elongatus* (Clarke 1879)**

Common name: **Elongate frostfish; Espada-de-má-água-comprido**, Espada-de-má-água, Espada-branca-de-natura.

References: mainland Portugal (?) (Parin, 1986b; Saldanha *et al.*, 1995; Erzini *et al.*, 2001b; Borges *et al.*, 2002; Santos *et al.*, 2002a, b; Moura, 2004; Fernandes *et al.*, 2007; Silva, 2008, 2010, 2011; Bordalo-Machado *et al.*, 2009; Silva *et al.*, 2009; Carneiro *et al.*, 2014; Silva and Borges, 2014) Azores (?) (Morato-Gomes *et al.*, 1998; Melo and Menezes, 2002) and Madeira (?) (INIP, 1982c, 1984a, b, c; Parin, 1986b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014). The records of *B. elongatus* could probably be the subspecies *Benthodesmus elongatus simonyi* (Steindachner 1891) which was raised to species *B. simonyi*. Then, those records of *B. elongatus* should be considered as *B. simonyi*. Thus, having in mind the current southern circumglobal geographic distribution of *B. elongatus* (Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) this species is not considered in the total number of valid species.

Benthodesmus simonyi (Steindachner 1891)

Common name: **Simony's frostfish; Espada-de-má-água**, Espada, Espada-branca-de-natura, Espada-de-água-má, Peixe-espada, Peixe-espada-branca-de-natura, Peixe-espada-de-água-má, Peixe-espada-de-má-água.

References: mainland Portugal (Bragança, 1903; MNHN-IC-1905-0578, 1904; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; INIP, 1982a; Nakamura and Parin, 1993; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (Silva *et al.*, 1995; Menezes, 1996; Melo and Menezes, 2002; Santos *et al.*, 1997; Menezes *et al.*, 2006, 2009, 2012; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014) and Madeira (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Nakamura and Parin, 1993; Ferreira *et al.*, 2008; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Parin and Nakamura, 2016; POGID, 2016; Porteiro *et al.*, 2017; Froese and Pauly, 2019).

Benthodesmus tenuis (Günther 1877)

Common name: **Slender frostfish; Espada-de-má-água-elegante**, Peixe-espada.

References: mainland Portugal (INIP, 1982a). Eschmeyer *et al.* (2019) mention it as a circumglobal species in tropical through warm temperate seas. Thus, this is the northernmost occurrence in the Eastern Atlantic.

Lepidopus caudatus (Euphrasen 1788)

Common name: **Silver scabbardfish; Peixe-espada**, Anjo, Espada, Espada-branca, Espada-branco, Espada-legítimo, Lírio, Peixe-anjo, Peixe-espada-legítimo, Peixe-espada-branco, Talabarte.

References: mainland Portugal▲ (Vandelli, 1771, 1797; Holten, 1802; Shaw, 1803; Cuvier, 1829; de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1894c, 1897, 1898; Nobre, 1894, 1903, 1935; Goode and Bean, 1895, 1896; Vaillant, 1919; Seabra, 1911; Fowler, 1936b; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández *et al.*, 1981; INIP, 1981a, 1982a, b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Parin, 1986b; Fischer *et al.*, 1987; Nakamura and Parin, 1993; Saldanha *et al.*, 1995; Erzini *et al.*, 2001b; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002b, c; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Pakhorukov, 2008; Silva, 2008, 2010, 2011; Ward *et al.*, 2008; Silva *et al.*, 2009; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Christiansen *et al.*, 2014; Martins *et al.*, 2015; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Sampaio, 1904; Roule, 1919a; Vaillant, 1919; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Wood, 1974; Dias *et al.*, 1976, 1990; Maul, 1976 (PECS); Ehrich, 1977; Martins, 1982a, b; INIP, 1985a, b, c; Dias and Monteiro, 1986; Parin, 1986b; Dias, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Nakamura and Parin, 1993; Krug, 1995; Silva *et al.*, 1994, 1995; Clarke and Martins, 1995; Clarke *et al.*, 1995; Pinho and Pereira, 1995a; Silva, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Arruda, 1997; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Morato *et al.*, 2001a; Kukuev, 2002 (PECS); Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Pakhorukov, 2008 (PECS); Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2013, 2017; Besugo, 2013; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837; de Velloso, 1892; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Ehrich, 1977; INIP, 1980; Fischer *et al.*, 1981; Parin, 1986b; Nakamura and Parin, 1993; Lloris *et al.*, 1991; Severino, 2004; Pakhorukov, 2008 (PECS); Carneiro *et al.*, 2014; Parin and Nakamura, 2016; POGID, 2016; Porteiro *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of *Vandellius lusitanicus* Shaw 1803 was caught off in mainland Portugal and is deposited in the NHMUK collections. Several specimens caught off at the Meteor Bank are in the MMF collections (MMF 22357, 1967, set of eight specimens and MMF 22358, 1967, set of nineteen specimens). One specimen caught off at the Portuguese continental slope in October 2010 is in the MB collections.

Trichiurus lepturus Linnaeus 1758

Common name: **Largehead hairtail; Lírio, Chibata, Espada, Espada-lírio, Peixe-espada. Peixe-espada-lírio.**

References: mainland Portugal[▲] ([de Brito Capello, 1867a, d, 1880](#); [Gervais and Boulart, 1877b](#); [Goode and Bean, 1895, 1896](#); [Steindachner, 1867](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#); [YPM ICH 005193, 1959](#); [INIP, 1981a](#); [Martins, 1981](#); [Borges, 1984](#); [Parin, 1986b](#); [Fischer et al., 1987](#); [Nakamura and Parin, 1993](#); [Moura, 2004](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Collette et al., 2015b](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)), Azores (?) ([Osório de Castro, 1967](#)) and Madeira (?) ([Osório de Castro, 1967](#); [Fischer et al., 1981](#)). [Eschmeyer et al. \(2019\)](#) mention it as a circumglobal species in tropical and warm temperate seas. For Collette et al. (2015b) *T. lepturus* is a single, globally distributed species found in temperate and tropical waters. However, there are multiple lines of evidence showing geographically-correlated divergence and supporting the treatment of at least three species in the *T. lepturus* complex ([Hsu et al., 2009](#); [Chakraborty and Iwatsuki, 2006](#)). The Eastern Atlantic population reported as *T. lepturus* is probably an undescribed species ([Chakraborty and Iwatsuki, 2006](#)). Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores and Madeira as rare, without indicating the geographic location. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores. Thus, the occurrence records to the Azores and Madeira Islands are doubtful and this species is not considered in the total number of valid species for those islands.

Family Scombridae

Acanthocybium solandri (Cuvier 1832)

Common name: **Wahoo; Cavala-da-Índia**, Bicuda-da-Índia, Cavala-gigante, Ilhéu, Peixe-fumo, Peixe-serra, Sarra, Serra, Uaú.

References: mainland Portugal ([Carneiro et al., 2014](#); [Collette and Heessen, 2015a](#)), Azores[▲] ([Ferreira, 1937a](#); [Osório de Castro, 1967](#); [Collette, 1986b\(?\)](#); [Lloris et al., 1991](#); [Azevedo and Heemstra, 1995](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Cardigos et al., 2005](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [IMAR/DOP Underwater visual census 5592, 2014](#); [Collette and Heessen, 2015a](#); [Afonso, 2016](#); [Collette, 2016d](#)) and Madeira[▲] ([Nunes, 1953, 1974](#); [Osório de Castro, 1967](#); [Collette, 1986b\(?\)](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015a](#); [Collette, 2016d](#); [Froese and Pauly, 2019](#)). [Eschmeyer et al. \(2019\)](#) mention it as a circumglobal species in tropical and warm temperate seas. The distribution map of this species reported by Collette and Heessen (2015) partially included the south-west area of the Portuguese mainland EEZ. Sporadic catches of this species in southern Portugal (Algarve) has been reported, but the occurrence of this species needs further confirmation.

Auxis rochei (Risso 1810)

Common name: **Bullet tuna; Judeu**, Cacharra, Cachorra, Cachorreta, Chapouto, Gaiado, Serra.

References: mainland Portugal[▲] ([de Brito Capello, 1867a, 1880](#); [Steindachner, 1868a](#); [Gervais and Boulart, 1877b](#); [Almeida and Roquette, 1892](#); [Bragança, 1899](#); [Seabra, 1911](#); [Postel, 1979](#); [Collette and Nauen, 1983](#); [Collette, 1986b](#); [Fischer et al., 1987](#); [Martins et al., 1992, 2015](#); [Santos et al., 2002a, b, c](#); [ICES, 2005](#); [Gomes and Olim, 2007](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015b](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Nobre, 1924, 1930](#); [Ferreira, 1937](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Dias and Barraca, 1971](#); [Wheeler, 1978](#); [Collette, 1986b](#); [Lloris et al., 1991](#); [Collette and Aadland, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Collette, 2016d](#)) and Madeira ([Lowe, 1850, 1852](#); [Noronha and Sarmento, 1934](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Collette, 1986b](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015b](#); [Froese and Pauly, 2019](#)). This circumglobal species in tropical through temperate seas can be confused with *Auxis thazard* (Lacepède 1800) and appears to be the more common of the two species (*A. rochei* and *A. thazard*) in the Eastern Atlantic and Mediterranean (Collette, 2016d). One specimen caught off at the Portuguese continental slope in October 2010 is in the MB collections (MB06-005365).

Auxis thazard (Lacepède 1800)

Common name: **Frigate tuna; Judeu-liso**, Atum, Boneito, Bonito, Cachorra, Cachorreta, Cachorrinha, Chapouto, Gaiado, Judeu, Judia, Merma, Sarra, Serra.

References: mainland Portugal ([de Brito Capello, 1867a, 1880](#); [Bragança, 1899](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Collette and Nauen,](#)

1983; Collette *et al.*, 2011a; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (Nobre, 1924; Osório de Castro, 1967) and Madeira▲ (Lowe, 1850, 1852; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Collette *et al.*, 2011a; Carneiro *et al.*, 2014). According to Froese and Pauly (2019) many authors have used the name *Auxis thazard* as including *Auxis rochei* in the belief that there was only a single worldwide species of *Auxis*. In the Atlantic, most catches reported as *A. thazard* are probably *A. rochei* (Collette *et al.*, 2011a).

***Euthynnus alletteratus* (Rafinesque 1810)**

Common name: **Little tunny; Merma**, Albacora, Alvacora, Apluro, Atum, Barrilete, Boneito, Bonito, Cacerete, Cachorra, Cachorreta, Cachorrinha, Cachorro, Falsa-albacora, Fule-fule, Judeu, Judia, Melena, Melva, Peixe-judeu, Tonina.

References: mainland Portugal▲ (Steindachner, 1868a; de Brito Capello, 1880; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1898; Bragança, 1899; Seabra, 1911; Osório, 1917; Frade and Buen, 1932; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Collette and Nauen, 1983; Collette, 1986b; Fischer *et al.*, 1987; Erzini *et al.*, 2001a; Santos *et al.*, 2002a; Carneiro *et al.*, 2014; Collette and Heessen, 2015c; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Dias and Barraca, 1971; Arruda, 1997) and Madeira (Osório de Castro, 1967; Collette and Heessen, 2015c).

***Katsuwonus pelamis* (Linnaeus 1758)**

Common name: **Skipjack tuna; Gaiado**, Atum, Atum-bonito, Boneito, Bonito, Bonito-de-barriga-listada, Bonito-de-ventre-listado, Bonito-de-ventre-raiado, Cacerete, Cachorra, Cachorreta, Cachorrinha, Canela, Listado, Listão, Listrado, Melancia, Sarra, Sarrajão, Sarrapão, Serra, Serrajão, Trazaste, Traz-haste, Triaste.

References: mainland Portugal▲ (de Brito Capello, 1867a, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Almeida and Roquette, 1892; Vieira, 1898; Bragança, 1899; Seabra, 1911; Frade and Buen, 1932; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981, 1987; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Collette, 1986b; Santos *et al.*, 2002a; Carneiro *et al.*, 2014; Collette and Heessen, 2015d; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Sampaio, 1904; Ferreira, 1932, 1937a; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Dias and Barraca, 1971; Wheeler, 1978; Fischer *et al.*, 1981; Muzanor and Westhaus, 1981; Martins, 1982a; Pereira, 1982a, b, 1983b, 1992a, d, 1994a, 1995; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Bard, 1986; Collette, 1986b; Monteiro, 1991; Arruda, 1992, 1997; Silva, 1992; Santos *et al.*, 1994a, 1997; Pinho and Pereira, 1995a, b; Pinho *et al.*, 1995; Isidro, 1996; Morato *et al.*, 2001a; Cardigos *et al.*, 2005; Porteiro *et al.*, 2010, 2017(?); Carneiro *et al.*, 2014; Collette and Heessen, 2015d; Collette, 2016d; Fauconnet *et al.*, 2019) and Madeira▲ (Lowe, 1837; Roule, 1919a; Noronha and Sarmento, 1934; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Collette, 1981, 1986b; Fischer *et al.*, 1981; Severino, 2004; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Collette and Heessen, 2015d; Collette, 2016d; POGIS, 2016; Froese and Pauly, 2019).

***Orcynopsis unicolor* (Geoffroy St. Hilaire 1817)**

Common name: **Plain bonito; Palometa**, Atum, Boneito, Bonito, Palamita, Palmeta, Palombeta, Pelamide, Pelombeta, Plumbeta, Trazaste, Traz-haste, Triaste.

References: mainland Portugal▲ (Collette, 1981; Fischer *et al.*, 1981; Santos *et al.*, 2002a; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Lloris *et al.*, 1991) and Madeira (Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Lloris *et al.*, 1991; Collette and Heessen, 2015e). According to Eschmeyer *et al.* (2019) the current geographic distribution of *O. unicolor* is the western Baltic Sea, North Sea, Mediterranean Sea and, in the Eastern Atlantic, from southern Norway to Senegal. The distribution maps of this species reported by Fischer *et al.* (1981) and Collette and Heessen (2015e) partially included the area of Selvagens Islands (Madeira EEZ).

***Sarda sarda* (Bloch 1793)**

Common name: **Atlantic bonito; Sarrajão**, Albacôra, Atum, Boneito, Bonito, Bonito-de-dorso-listado, Bonito-de-dorso-raiado, Bonito-de-lombo-raiado, Cavalão, Cerda, Minerva, Sarrajão, Sarda, Sarragão, Sarrajão, Sarralhão, Serra, Serrajão, Serralhão, Xarda.

References: mainland Portugal[▲] (Vandelli, 1771; de Brito Capello, 1867a, 1880; Gervais and Boulart, 1877b; Posser, 1880; Osório, 1888, 1895, 1917; Baldaque da Silva, 1891; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895a, 1935; Bragança, 1899, 1903; Seabra, 1911; Frade and Buen, 1932; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Postel, 1979; Collette, 1981, 1986b; Fischer et al., 1981, 1987; INIP, 1981a; Martins, 1981; Collette and Nauen, 1983; Martins et al., 1999, 2015; Erzini et al., 2001a; WWF, 2001; Santos et al., 2002a, c; Henriques et al., 2007, 2008; Abecasis et al., 2009; Carneiro et al., 2014; OCEANA, 2014; Collette and Fernandes, 2015; Collette, 2016d; Martins and Carneiro, 2018; Baptista et al., 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Richard, 1905b; Fowler, 1936b; Ferreira, 1937a; Collins, 1954; Osório de Castro, 1967; Collett and Chao 1975; Wheeler, 1978; Harmelin and Harmelin-Vivien, 1979; Postel, 1979; Ré, 1979a; Martins, 1982a; Rey et al., 1984; Collette, 1986b; Patzner et al., 1990, 1992; Wirtz, 1990a, 1991, 1994; Lloris et al., 1991; Arruda, 1992, 1997; Saldanha, 1995; Silva and Menezes, 1995a; Isidro, 1996; Santos et al., 1997; Harmelin-Vivien et al., 2001; Morato et al., 2001a, b; Cardigos et al., 2005; Menezes et al., 2006; Porteiro et al., 2010; Carneiro et al., 2014; Collette and Fernandes, 2015; Afonso, 2016) and Madeira[▲] (Lowe, 1850, 1852; Osório de Castro, 1954, 1967; Collette, 1981, 1986b; Noronha and Sarmento, 1934; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Fischer et al., 1981; Lloris et al., 1991; Delgado, 1998; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Collette and Fernandes, 2015; Collette, 2016d; POGIS, 2016; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in March 2011 is in the MB collections (MB06-005327).

Scomber colias Gmelin 1789

Common name: **Atlantic chub mackerel; Cavala**, Cavala-comum, Cavala-corrente, Cavala-frescal, Cavala-legítima, Cavalão, Judeu, Manica, Peixe-chuva, Ruama, Sarda, Sereia, Xarda.

References: mainland Portugal[▲] (Carvalho, 1765(?); de Brito Capello, 1867a, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1895a, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; INIP, 1981a, 1982a, b; Martins, 1981; Collette, 1986b; Fischer et al., 1987; Martins et al., 1992, 1999, 2015; Erzini et al., 1996, 1997, 2001a; Gonçalves et al., 1996, 1997; Borges et al., 2002; Jorge et al., 2002; Santos et al., 2002a, c; Mendes et al., 2004; ICES, 2005; Sousa et al., 2005, 2006; Stergiou et al., 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Silva, 2008, 2010, 2011; Batista et al., 2009; Silva et al., 2009; Gonçalves et al., 2010; Veiga et al., 2010; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Christiansen et al., 2014; Costa, 2014; OCEANA, 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Baptista et al., 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Ferreira, 1937a; Fowler, 1919, 1936b; Roule, 1919a; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1978; Postel, 1979; Muzanor and Westhaus, 1981; Martins, 1982a; INIP, 1985a, b, c; Collette, 1986b; Dias and Monteiro, 1986; Isidro, 1996, 1988, 1990a, b; Dias, 1991; Lloris et al., 1991; Arruda, 1992; Pereira, 1992d, 1995; Silva, 1992; Santos et al., 1994a, 1995a, 1997; Pinho et al., 1995; Silva and Menezes, 1995a, b; Silva et al., 1995; Isidro, 1996; Menezes, 1996; Monteiro, 1996; Monteiro et al., 1996b; Morato-Gomes et al., 1998; Sousa et al., 1999; Cardigos et al., 2005; Menezes et al., 2006; Rosa et al., 2006; Pakhorukov, 2008 (PECS); Porteiro et al., 2010; Carneiro et al., 2014; Diogo and Pereira, 2014; Afonso, 2016; Fauconnet et al., 2019) and Madeira[▲] (Lowe, 1850, 1852; Günther, 1860; Von Martens, 1876; de Velloso, 1892; Noronha and Sarmento, 1934; Nobre, 1935; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; INIP, 1980, 1982c, 1984a, b; Collette, 1981, 1986b; Fischer et al., 1981; Lloris et al., 1991; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Collette, 2016d; POGID, 2016; POGIS, 2016; Biscoito et al., 2017; Froese and Pauly, 2019).

Scomber scombrus Linnaeus 1758

Common name: **Atlantic mackerel; Sarda**, Cavala, Cavala-frescal, Manica, Sereia, Xarda.

References: mainland Portugal[▲] (Carvalho, 1756; Vandelli, 1771; de Brito Capello, 1867a, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Osório, 1896; Nobre, 1894, 1895a, 1903, 1935; Bragança, 1903; Richard, 1908; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956;

Osório de Castro, 1954, 1967; Postel, 1979; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982a, b; Martins, 1981; Collette and Nauen, 1983; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Collette, 1986b; Martins *et al.*, 1992, 1999, 2015; Erzini *et al.*, 2001a, b; Gomes *et al.*, 2001; Monteiro *et al.*, 2001; Borges *et al.*, 2002; Jorge *et al.*, 2002; Santos *et al.*, 2002a, c; Prista *et al.*, 2003; Garnerot *et al.*, 2004; Mendes *et al.*, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Stergiou *et al.*, 2006; Fernandes *et al.*, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Silva, 2008, 2010; Batista *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Collette *et al.*, 2015c; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Mecklenburg *et al.*, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Sampaio, 1904; Nobre, 1924, 1930; Fowler, 1936b; Ribeiro, 1936; Ferreira, 1937a; Osório de Castro, 1967; Postel, 1979; Collette, 1981, 1986b; INIP, 1985a; Dias and Monteiro, 1986; Lloris *et al.*, 1991; Arruda, 1992, 1997; Evseenko *et al.*, 2004 (PECS); Carneiro *et al.*, 2014; Mecklenburg *et al.*, 2018) and Madeira (Lowe, 1837; Zugmayer, 1911b; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; ZMH 105383, 1968; Fischer *et al.*, 1981; Collette, 1986b, 2016c; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Collette *et al.*, 2015c; Biscoito *et al.*, 2017; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. scombrus* is the Arctic Ocean, Baltic Sea, North Sea, Mediterranean Sea, Black Sea, and the temperate waters of the North Atlantic. Santos *et al.* (1997) mention that several authors reported the occurrence of *S. scombrus* in the Azores. However, these authors have serious reservations on its occurrence since they haven't seen this species in the Azores.

***Scomberomorus tritor* (Cuvier 1832)**

Common name: **West African Spanish mackerel; Serra-branca**, Atum, Boneito, Bonito, Cavala, Cavala-bonita, Cavala-bonito, Cavala-espanhola, Cavala-moira, Cavala-moirinha, Ilhéu, Moira, Moirinha, Moura, Mourinha, Peixe-serra, Perna-de-moça, Pintada, Sarra, Serra.

References: Azores (?) (Osório de Castro, 1967) and Madeira (Osório de Castro, 1967; Fischer *et al.*, 1981; Collette *et al.*, 2015d). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. tritor* in the Eastern Atlantic is concentrated in the Gulf of Guinea from the Canary Islands south to southern Angola. Osório de Castro (1967) reported the occurrence of *Scomberomorus maculatus* (Mitchill 1815) in the fishing grounds of the Azores and Madeira, without indicating the geographic location. Collette *et al.* (2015d) include the Selvagens Islands area in the distribution map of this species. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores. *S. maculatus* is currently valid and its current distribution is in the Western Atlantic (Eschmeyer *et al.* 2019). The distribution of *S. maculatus* reported by Osório de Castro (1967) lead us to concluding that probably was referring to *S. maculatus* (non Mitchill 1815), which is the synonymous of *Scomberomorus tritor* (Cuvier 1832). Froese and Pauly (2019) mention that *S. tritor* caught off in the Eastern Atlantic is often confused with *S. maculatus* (Collette, 2016d) having been erroneously considered as a synonym of *S. tritor* by many authors. The presence of *S. tritor* in the Azores is not yet confirmed. Thus, this species is not considered in the total number of valid species for the Azores.

***Thunnus alalunga* (Bonnaterre 1788)**

Common name: **Albacore; Atum-voador**, Albacó, Albacor, Albacora, Albacore, Alvacó, Alvacor, Alvacora, Àsinha, Atum, Atum-branco, Atum-de-barbatana-comprida, Atum-de-galha-comprida, Atum-de-grelha-comprida, Atum-de-guelha-comprida, Atum-guelha-comprida, Atum-negro, Atum-verdadeiro, Atum-voador, Avoador, Àzinha, Germão, Ilhéu, Trazaste, Voador.

References: mainland Portugal▲ (de Brito Capello, 1872, 1873, 1880; Posser, 1880; Almeida and Roquette, 1892; Nobre, 1895a, 1935; Bragança, 1899; Seabra, 1911; Frade and Buen, 1932; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Postel, 1979; Collette, 1981, 1986b; Fischer *et al.*, 1981, 1987; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Bordalo-Machado *et al.*, 2009; Santos *et al.*, 2002a; Carneiro *et al.*, 2014; Collette *et al.*, 2015e; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Richard, 1900, 1910; Gain, 1913; Ferreira, 1932, 1940; Fowler, 1936b; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Clarke, 1956; Dias and Barraca, 1971; Wheeler, 1978; Postel, 1979; Bauchot and Pras, 1980; Fischer *et al.*, 1981; Muzanor and Westhaus, 1981; Garcez, 1982; Martins, 1982a; Pereira, 1982a, b, 1983b, 1992a, d, 1994a, 1995; Collette and

Nauen, 1983; Nishikawa *et al.*, 1985; Bard, 1986; Collette, 1986b, 2016c; Lloris *et al.*, 1991; Monteiro, 1991; Morin, 1992; Silva, 1992; Pinho and Pereira, 1995a, b; Pinho *et al.*, 1995; Isidro, 1996; Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Porteiro *et al.*, 2010, 2017(?); Carneiro *et al.*, 2014; Collette *et al.*, 2015e; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1849; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Collette, 1981, 1986b, 2016c; Fischer *et al.*, 1981; Collette and Nauen, 1983; Lloris *et al.*, 1991; Severino, 2004; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Collette *et al.*, 2015e; Froese and Pauly, 2019).

Thunnus albacares (Bonnaterre 1788)

Common name: **Yellowfin tuna; Atum-albacora**, Albacó, Albacor, Albacora, Albacore, Albecora, Albesco, Alvacó, Alvacor, Alvacora, Atum, Atum-amarelo, Atum-de-alheta-amarela, Atum-de-barbatana-amarela, Atum-de-cacho, Atum-de-galha, Atum-de-galha-à-ré, Atum-de-grelha-à-ré, Atum-verdadeiro, Galha-à-ré, Grelha-à-ré, Lelofino, Labacória, Lombo-lombo, Peixe-de-galha-à-ré, Peixe-de-grelha-à-ré, Peixinho-da-ilha, Rabã, Rabão, Rabo-seco, Valcória.

References: mainland Portugal▲ (Frade, 1929, 1932; Frade and Buen, 1932; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Collette, 1981, 1986b, 1990, 2016c; Fischer *et al.*, 1981; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Collette *et al.*, 2011b; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Frade, 1931b, c, 1960; Ferreira, 1932, 1937a, 1940; Nobre, 1935; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Parin, 1970; Dias and Barraca, 1971; Wheeler, 1978; Postel, 1979; Collette, 1981, 1986b, 1990, 2016c; Fischer *et al.*, 1981; Muzanor and Westhaus, 1981; Martins, 1982a; Pereira, 1982a, b, 1983b, 1986b, 1992d, 1995; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Bard, 1986; Lloris *et al.*, 1991; Pinho and Pereira, 1995a; Pinho *et al.*, 1995; Isidro, 1996; Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Porteiro *et al.*, 2010, 2017(?); Collette *et al.*, 2011b; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1860; Cunningham, 1910; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Postel, 1979; Collette, 1981, 1986b, 2016c; Fischer *et al.*, 1981; Collette and Nauen, 1983; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Collette *et al.*, 2011b; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Thunnus atlanticus (Lesson 1831)

Common name: **Blackfin tuna; Atum-de-barbatana-negra**, Albacorinha.

References: Azores (PECS) (MCZ 75623, 1984; Carneiro *et al.*, 2014).

Thunnus obesus (Lowe 1839)

Common name: **Bigeye tuna, Atum-patudo**, Albacó, Albacora, Albecora, Alvacó, Alvacor, Alvacora, Atum, Atum-de-cala, Atum-de-olhos-grandes, Atum-verdadeiro, Atum-vermelho, Cachorra, Cara-larga, Cavalinho, Patudo.

References: mainland Portugal▲ (Frade, 1929; Fischer *et al.*, 1981; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Collette, 1986b; Collette *et al.*, 2011c; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019) Azores▲ (Frade, 1929, 1931a, c, 1960; Borodin, 1930; Ferreira, 1932, 1937a, 1940; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Dias and Barraca, 1971; Collignon and Aloncle, 1973; Postel, 1979; Collette, 1981, 1986b, 1990, 2016c; Fischer *et al.*, 1981; Muzanor and Westhaus, 1981; Garcez, 1982; Martins, 1982a; Pereira, 1982a, b, 1983b, 1985a, b, 1987a, 1989a, 1992a, d, 1994a, 1995; Collette and Nauen, 1983; Nishikawa *et al.*, 1985; Bard, 1986; Silva, 1992; Lloris *et al.*, 1991; Clarke and Martins, 1995; Pinho and Pereira, 1995a, b; Pinho *et al.*, 1995; Isidro, 1996; Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005; Porteiro *et al.*, 2010, 2017(?); Collette *et al.*, 2011c; Carneiro *et al.*, 2014; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1860; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Postel, 1979; Collette, 1981, 1986b, 1990, 2016c; Fischer *et al.*, 1981; Collette and Nauen, 1983; Lloris *et al.*, 1991; Severino, 2004; Wirtz *et al.*, 2008; Collette *et al.*, 2011c; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019).

Thunnus thynnus (Linnaeus 1758)

Common name: **Atlantic bluefin tuna; Atum-rabilho**, Albacora, Alvacor, Alvacora, Atuárrio, Atuarro, Atum, Atum-azul, Atum-comum, Atum-de-barbatana-azul, Atum-de-cacho, Atum-de-cardume, Atum-de-direito, Atum-do-Algarve, Atum-do-Mediterrâneo, Atum-legítimo, Atum-preto, Atum-rabil, Atum-

rabilha, Atum-rabilo, Atum-de-recuado, Atum-de-retorno, Atum-de-revés, Atum-de-tabela, Atum-de-través, Atum-estacionário, Atum-pairante, Atum-verdadeiro, Atum-vermelho, Atum-vulgar, Cachorreta, Cachorreta, Rabã, Rabão, Rabelho, Rabil, Rabilha, Rabilho, Rabilo, Reu-reu, Tonina, Toninha, Toninho, Tonino.

References: mainland Portugal[▲] (de Brito Capello, 1867a, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1935; Vieira, 1898; Bragança 1899; Seabra, 1911; Frade, 1932; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Postel, 1979; Collette, 1981, 1986b; Fischer et al., 1981, 1987; Collette and Nauen, 1983; Nishikawa et al., 1985; Santos et al., 2002a; Carneiro et al., 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Hilgendorf, 1888a; Sampaio, 1904; Heldt, 1931; Frade, 1929, 1931c, 1960; Ferreira, 1932, 1937a, 1940; Fowler, 1936b; Poll, 1947; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Dias and Barraca, 1971; Wheeler, 1978; Postel, 1979; Garcez, 1982; Martins, 1982a; Pereira, 1982a, b, 1983b, 1987a, 1992d, 1995; Collette and Nauen, 1983; Cort and Rey, 1984; Nishikawa et al., 1985; Bard, 1986; Collette, 1986b, 2016c; Lloris et al., 1991; Arruda, 1992, 1997; Pinho and Pereira, 1995a; Pinho et al., 1995; Isidro, 1996; Santos et al., 1997; Porteiro et al., 2010, 2017(?); Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Lowe, 1837; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Collette, 1981, 1986b, 2016c; Fischer et al., 1981(?); Collette and Nauen, 1983; Wirtz et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019).

Family Scombrolabracidae

Scombrolabrax heterolepis Roule 1921

Common name: **Longfin escolar; Escolar-de-peitoral-longa**.

References: mainland Portugal (Nishikawa et al., 1985; Costa et al., 2012; Carneiro et al., 2014; Ralph, 2015m), Azores (Nishikawa et al., 1985; Santos et al., 1997; IMAR/DOP 3SMA-02-P04SCH, 2004; IMAR/DOP COLETA 164, 2004; Kukuev, 2002 (PECS); Pakhorukov, 2008; Porteiro et al., 2010; Carneiro et al., 2014; Ralph, 2015m) and Madeira (MOM-IC-0091-1226, 1915; Roule, 1921, 1922; NHMUK 2017.1.5.58, 1966; Parin and Bekker, 1979; Parin, 1986b; Lloris et al., 1991; Carneiro et al., 2014; Ralph, 2015m; Roberts et al., 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off south of Madeira is in the MOM collections (MOM-IC-0091-1226).

Family Centrolophidae

Centrolophus niger (Gmelin 1789)

Common name: **Rudderfish; Liro-preto**, Laro, Liro, Pescada, Pescada-preta, Preta.

References: mainland Portugal (de Brito Capello, 1867a, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Osório, 1896, 1909; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer et al., 1981, 1987; Haedrich, 1986a; Saldanha et al., 1995; Erzini et al., 2001b; Santos et al., 2002b; Moura, 2004; Gomes and Olim, 2007; Carneiro et al., 2014; Silva and Borges, 2014; Knudsen et al., 2015g), Azores (Cligny, n.d.; Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Guiart, 1935; Fowler, 1936b; Le Gall, 1937a; Poll, 1947; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Parin, 1970; Bauchot and Pras, 1980; Haedrich, 1986a; Lloris et al., 1991; Arruda, 1992, 1997; Santos et al., 1997; Kukuev, 2002 (PECS); Melo and Menezes, 2002; Menezes et al., 2006, 2012; Porteiro et al., 2010; Carneiro et al., 2014; Knudsen et al., 2015g) and Madeira (Cligny, n.d.; Lowe, 1839, 1840b, 1844a, 1849; NHMUK 1862.6.14.16, 1861-1890; Johnson, 1862a, 1863a; Goode and Bean, 1895, 1896; Osório de Castro, 1967; Haedrich, 1986a; Collett, 1896; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer et al., 1981; Lloris et al., 1991; Carneiro et al., 2014; Knudsen et al., 2015g; Haedrich, 2016a). The holotype (unique) of *Schedophilus elongatus* Johnson 1862, synonym of *C. niger*, caught off in Madeira is in the NHMUK collections (NHMUK 1862.6.14.16). One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005164).

***Hyperoglyphe perciformis* (Mitchill 1818)**

Common name: **Barrelfish; Liro-errante**, Laro, Liro, Pescada-preta.

References: mainland Portugal ([Osório, 1895](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Haedrich, 1986a](#); [Karrer, 1986b](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912 \(PECS\)](#); [Fowler, 1936b](#); [Osório de Castro, 1967](#); [Santos et al., 1997](#); [Carneiro et al., 2014](#); [Nielsen et al., 2015g](#); [Bañón et al., 2019b](#)) and Madeira ([Osório de Castro, 1967](#); [Nielsen et al., 2015g](#)). The holotype (unique) of *Centrolophus newtoni* Osório 1895 caught off near Matosinhos (off mainland Portugal) and preserved in the Museu Bocage was destroyed in the 1978 fire. *C. newtoni* is synonym of *H. perciformis*.

***Schedophilus maculatus* Günther 1860**

Common name: **Pelagic butterfish; Liro-manchado**, Liro.

References: Azores ([Murray and Hjort, 1912](#); [Fowler, 1936b](#); [Osório de Castro, 1967](#); [Arruda, 1997](#); [Carneiro et al., 2014](#)) and Madeira ([Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Maul, 1964](#); [Osório de Castro, 1967](#); [Carneiro et al., 2014](#)).

***Schedophilus medusophagus* (Cocco 1839)**

Common name: **Cornish blackfish; Liro-mole**, Liro, Pampo-mole.

References: mainland Portugal ([Haedrich, 1986a](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1896](#); [Murray and Hjort, 1912](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [MCZ 149614, 1963](#); [Haedrich, 1964, 1986](#); [Osório de Castro, 1967](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Porteiro et al., 2010](#); [de Casamajor and Morandeau, 2013](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)) and Madeira ([MCZ 153226, 1986](#); [Osório de Castro, 1967\(?\)](#); [Haedrich, 1986a](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [de Casamajor and Morandeau, 2013](#); [Carneiro et al., 2014](#); [Bañón et al., 2016a](#); [Froese and Pauly, 2019](#)).

***Schedophilus ovalis* (Cuvier 1833)**

Common name: **Imperial blackfish; Liro-imperial**, Choupa, Leiro, Liro, Lírio, Pâmpano.

References: mainland Portugal ([Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Fischer et al., 1981, 1987](#); [Haedrich, 1986a](#); [Carneiro et al., 2014](#)), Azores[▲] ([Cuvier and Valenciennes, 1833](#); [MNHN-IC-0000-1795, 1833](#); [Collett, 1896a](#); [Murray and Hjort, 1912](#); [Fowler, 1936b](#); [Barnard, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Haedrich, 1979, 1986, 2002](#); [Fischer et al., 1981](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Pinho and Pereira, 1995b](#); [Silva et al., 1995](#); [Menezes, 1996](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Haedrich, 2016a](#); [Brown et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Lowe, 1833a, 1837, 1838, 1839, 1840b, 1844a, 1849](#); [Günther, 1860, 1887](#); [de Brito Capello, 1871](#); [de Velloso, 1892](#); [Collett, 1896](#); [Nobre, 1935](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Maul, 1964](#); [Osório de Castro, 1954, 1967](#); [Haedrich, 1979, 1986, 2002](#); [Fischer et al., 1981](#); [INIP, 1984c](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Severino, 2004](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Haedrich, 2016a](#); [Biscoito et al., 2017](#); [Brown et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Centrolophus crassus* Cuvier & Valenciennes 1833, synonym of *S. ovalis*, caught off 125 NM west of the Azores is in the MNHN collections ([MNHN-IC-0000-1795](#)).

Family Nomeidae

***Cubiceps gracilis* (Lowe 1843)**

Common name: **Driftfish; Tirone**.

References: mainland Portugal ([Osório, 1909](#); [Seabra, 1911](#); [Norman, 1930](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Dinis, 1985](#); [Haedrich, 1986a](#); [Fischer et al., 1987](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Smith-Vaniz and Herrera, 2015d](#); [Froese and Pauly, 2019](#)), Azores ([Günther, 1860, 1887, 1889](#); [MCZ 32119, 1928](#); [Borodin, 1931](#); [Fowler, 1936b](#); [Koefoed, 1952](#); [Collins, 1954](#); [Haedrich, 1964, 1967](#); [Osório de Castro, 1967](#); [Bauchot and Pras, 1980](#); [Haedrich, 1986a](#); [Agafonova and Kukuyev, 1990](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Smith-Vaniz and Herrera, 2015d](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1843, 1844b](#); [Günther, 1860, 1887](#),

1889; Johnson, 1863b; Collett, 1896; Regan, 1902; Norman, 1930; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1967; Haedrich, 1986a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Smith-Vaniz and Herrera, 2015d; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of *Seriola gracilis* Lowe 1843 was caught off in Madeira and its whereabouts are unknown. The syntype of *Aphareus obtusirostris* Borodin 1930 was caught off in the Azores (MCZ 32119, 1928). *Seriola gracilis* and *Aphareus obtusirostris* are synonyms of *C. gracilis*.

***Nameus gronovii* (Gmelin 1789)**

Common name: **Man-of-war fish; Pequeno-argonauta**.

References: Azores ([FMNH 87602, 1963](#); Haedrich, 1986a; Arruda, 1997; Carneiro *et al.*, 2014; Ralph, 2015n) and Madeira ([Fowler, 1936b](#); [Albuquerque, 1954-1956](#); Osório de Castro, 1967; Haedrich, 1986a). According to Haedrich (1986c) this species is rare in the Eastern Atlantic being also rare in the Madeira ([Albuquerque, 1954-1956](#)).

***Psenes cyanophrys* Valenciennes 1833**

Common name: **Freckled driftfish; Peixe-pastor-amarelado**.

References: Madeira ([MCZ 79635, 1970](#); Carneiro *et al.*, 2014; Singh-Renton *et al.*, 2015). The distribution maps of this species reported by Singh-Renton *et al.* (2015) partially included in the area of Selvagens Islands (Madeira EEZ).

***Psenes maculatus* Lütken 1880**

Common name: Silver driftfish; **Peixe-pastor-prateado**.

References: Azores ([Lütken, 1880 \(PECS\)](#); [MNHN-IC-1995-0021, 1951 \(PECS\)](#); Haedrich, 1967, 1986; Arruda, 1997; Santos *et al.*, 1997; Bogutskaya, 2007c; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014). Ten syntypes of this species caught off north and south of Azores, 39°N, 25°04'W and 34°40.0' N, 27°26.0' W (in the PECS area) are in the ZMUC collections [ZMUC 30-38 (set of nine specimens) and ZMUC 83].

Family Tetragonuridae

***Tetragonurus atlanticus* Lowe 1839**

Common name: Bigeye squaretail; **Escalar-de-natura-olho-grande**, Escolar-de-natura.

References: mainland Portugal ([Haedrich, 1986a](#); Carneiro *et al.*, 2014; Ralph, 2015o), Azores ([FMNH 72534, 1920](#); Grey, 1955; Janssen and Harbison, 1981; Haedrich, 1986a; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Ralph, 2015o) and Madeira ([Lowe, 1839, 1840b, 1844a, 1949](#); Goode and Bean, 1895, 1896; Haedrich, 1979, 1986; MCZ 97044, 1986; Carneiro *et al.*, 2014; Ralph, 2015o). The holotype (unique) of this species was caught off in Madeira and its whereabouts are unknown (probably NHMUK 1914.7.14.86).

***Tetragonurus cuvieri* Risso 1810**

Common name: **Smalleye squaretail; Escolar-de-natura-olho-pequeno**, Escolar-de-natura.

References: mainland Portugal ([Haedrich, 1986a](#); Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015b), Azores ([NHMUK 1876.6.21.23, 1864-1877](#); [NHMUK 1917.7.14.86, 1912-1936](#); [FMNH 72536, 1922](#); [Borodin, 1931](#); Grey, 1955; [NHMUK 2006.9.19.7, 1966](#); Janssen and Harbison, 1981; Haedrich, 1986a; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015b) and Madeira ([Lowe, 1833a, 1839, 1840b, 1843-1860, 1844a, 1849](#); Günther, 1861a; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Haedrich, 1986a; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015b). The holotype of this species caught off in Madeira is in the NHMUK collections (NHMUK 1917.7.14.86).

Family Stromateidae

***Stromateus fiatola* Linnaeus 1758**

Common name: **Blue butterfish; Pampo-godinho**, Godinho, Pambo, Pâmpalo, Pampano, Pâmpano, Pampló, Pampo, Pando, Peixe-pombo, Pombo, Pompo, Ranholoso, Subscrito.

References: mainland Portugal▲ ([Carvalho, 1765](#); [Vandelli, 1771](#); [de Brito Capello, 1867a, 1880](#); [Steindachner, 1868a](#); [Gervais and Boulart, 1877b](#); [Almeida and Roquette, 1892](#); [Vieira, 1897, 1898](#);

Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Haedrich, 1979, 1986, 2016b; Fischer *et al.*, 1987; Borges *et al.*, 2002; Jorge *et al.*, 2002; Santos *et al.*, 2002a; ICES, 2005; Gomes and Olim, 2007; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015c; Eschmeyer *et al.*, 2019; Martins and Carneiro, 2018), Azores (Nobre, 1924, 1930; Albuquerque, 1954-1956; Arruda, 1997; Santos *et al.*, 1997(?); Carneiro *et al.*, 2014) and Madeira (INIP, 1984b; Haedrich, 1986a; Carneiro *et al.*, 2014).

Order Syngnathiformes

Family Aulostomidae

Aulostomus cf. *maculatus*

Common name: Atlantic trumpetfish; **Trombeta-salpicada**, Bom-alimento, Peixe-trombeta, Trombeta, Trombeteiro.

References: Madeira (Maul, 1959). According to Bowen *et al.* (2001) and Eschmeyer *et al.* (2019) this species is distributed in the tropical western Atlantic Ocean. However, there are occurrence records in West African coast, near Cape Verde Islands. It is possible that some specimens identified as *A. maculatus* have been misidentifications of *A. strigosus*.

Aulostomus strigosus Wheeler 1955

Common name: Atlantic cornetfish; **Trombeta**, Bom-alimento, Peixe-trombeta.

References: Madeira (Maul, 1959; Wheeler, 1979e; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Mello, 2000; Bowen *et al.*, 2001; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Poliom, 2015; Fritzsche, 2016a; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Family Fistulariidae

Fistularia petimba Lacepède 1803

Common name: **Red cornetfish**; **Corneta-rosada**, Colómu, Corneta, Pau, Peixe-pau, Peixe-tabaco, Tabaco.

References: mainland Portugal (Carneiro *et al.*, 2014) and Azores (Santos *et al.*, 1997; Azevedo *et al.*, 2004; Morato *et al.*, 2004; Wirtz, 2005; Cardigos *et al.*, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014).

Fistularia tabacaria Linnaeus 1758

Common name: **Cornetfish**; **Corneta-malhada**, Bombom, Corneta, Pau, Peixe-corneta, Peixe-pau, Peixe-tabaco, Tabaco.

References: Azores (MNHN-IC-A-7663, n.d.; Carneiro *et al.*, 2014; Carpenter *et al.*, 2015b).

Family Centriscidae

Macroramphosus gracilis (Lowe 1839)

Common name: **Slender snipefish**; **Trombeteiro-fino**, Apara-lápis, Trombeteiro.

References: mainland Portugal (Bragança, 1903; Nobre, 1935; Mohr, 1937; Albuquerque, 1954-1956; Osório de Castro, 1967; Nunes, 1974; INIP, 1981b; Fernández *et al.*, 1981; Martins, 1981; Borges, 1984; Dinis, 1985; Gomes *et al.*, 2001; Henriques *et al.*, 2008; Carneiro *et al.*, 2014), Azores (Lütken, 1880; Hilgendorf, 1888a; Vaillant, 1919; Fowler, 1936a; Mohr, 1937, Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1967; Dias *et al.*, 1976, 1990; INIP, 1985b, c; Carneiro *et al.*, 2014; Costa, 2014; Almeida and Biscoito, 2019) and Madeira (Lowe, 1837, 1839, 1840b, 1844a, 1849; Günther, 1861a; Regan, 1914; Nobre, 1935; Fowler, 1936a; Mohr, 1937; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Nunes, 1974; Wheeler, 1979e; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique and probably lost) of *Centriscus gracilis* Lowe 1839, synonym of *M. gracilis*, was caught off in Madeira.

Macroramphosus scolopax (Linnaeus 1758)

Common name: **Longspine snipefish**; **Trombeteiro**, Apara-lápis, Galinhola, Peixe-flauta, Peixe-trombeteiro, Peixe-trompeteiro, Trombeta, Vampiro.

References: mainland Portugal[▲] (de Brito Capello, 1868, 1880; Osório, 1895; Vieira, 1900; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Nunes, 1953; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; INIP, 1981a, b; Martins, 1981; Fischer et al., 1981, 1987; Borges, 1984, 2001; Dinis, 1985; Martins et al., 1999; Gomes et al., 2001; Monteiro et al., 2001; WWF, 2001; Santos et al., 2002a; Fernandes et al., 2007; Henriques et al., 2007, 2008; Gomes and Olim, 2007; Pakhorukov, 2008; Abecasis et al., 2009; Batista et al., 2009; Costa et al., 2012; Porteiro et al., 2010; Carneiro et al., 2014; Costa, 2014; OCEANA, 2014; Priede, 2017; Froese and Pauly, 2019), Azores (Lütken, 1880; Hilgendorf, 1888a; Zugmayer, 1911b; Vaillant, 1919; Fowler, 1936a; Mohr, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Mapstone and Wood, 1974, 1975; Dias et al., 1976, 1990; Maul, 1976 (PECS); Ehrich, 1977; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a; Dias, 1991; Arruda, 1992, 1997; Pierce et al., 1994; Wirtz, 1994; Azevedo, 1995; Gonçalves et al., 1995; Granadeiro et al., 1995; Pinho et al., 1995; Steiner 1995; Monteiro, 1996; Monteiro et al., 1996a, b; Santos et al., 1997; Harmelin-Vivien et al., 2001; Kukuev, 2002 (PECS); Parin and Pakhorukov, 2005 (PECS); Menezes et al., 2006; Pakhorukov, 2008 (PECS); Carneiro et al., 2014; Porteiro et al., 2017(?); Almeida and Bischoff, 2019) and Madeira (Regan, 1914; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Mohr, 1937; Nunes, 1953; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977; Fischer et al., 1981; INIP, 1984c; Abreu et al., 1995; Carneiro et al., 2014; Fritzsche, 2016b; POGID, 2016; POGIS, 2016). The distribution map of this species reported by Fischer et al. (1981) partially included the area of Selvagens Islands (Madeira EEZ). Several specimens, currently in the MMF collections, were caught off at the Josephine Bank (MMF 22462, 1967, set of six specimens and MMF 22807, 1967, one specimen) and at the Meteor Bank (MMF 22376, 1967, set of seven specimens and MMF 22353, 1967, set of eighty-six specimens).

Family Syngnathidae

Entelurus aequoreus (Linnaeus 1758)

Common name: **Snake pipefish**; **Marinha-amarela**, Cavalinho, Marinha.

References: mainland Portugal (de Brito Capello, 1880; Nobre, 1895c, 1903, 1935; Osório, 1895, 1917; Vieira, 1897; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Dawson, 1986; Arruda et al., 1988; Garnerot et al., 2004; Henriques et al., 2007, 2008; Carneiro et al., 2014; Mecklenburg et al., 2018; Froese and Pauly, 2019) and Azores (Günther, 1870; Collett, 1896a; Murray and Hjort, 1912; Duncker, 1915; Vaillant, 1919; Küenze, 1929; Fowler, 1936a; Poll, 1947; Koefoed, 1953; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1978, 1979e; Bauchot and Pras, 1980; Dawson, 1986; Lloris et al., 1991; Arruda, 1992, 1997; Santos et al., 1994a, 1995a, 1997; Floeter et al., 2008; Porteiro et al., 2010, 2017; Carneiro et al., 2014; Bañón et al., 2016a; Mecklenburg et al., 2018; Hamilton et al., 2017; Ryland and Lancaster, 2017; Froese and Pauly, 2019).

Hippocampus erectus Perry 1810

Common name: Lined Seahorse; **Cavalinho-estriado**, Cavalinho, Marinha.

References: Azores (IMAR/DOP COLETA 3120, 2004; Woodall et al., 2009, 2017; Porteiro et al., 2010; Afonso et al., 2013; Boehm et al., 2013; Carneiro et al., 2014; Lourie et al., 2016; Froese and Pauly, 2019(?)).

Hippocampus guttulatus Cuvier 1829

Common name: **Long-snouted seahorse**; **Cavalinho-de-focinho-comprido**, Cavalinho, Marinha, Cavalinho-do-mar, Cavalinho-marinho, Hipocampo, Lincorna.

References: mainland Portugal (de Brito Capello, 1872, 1873, 1880; Nobre, 1895a, 1935; Vieira, 1897; Bragança, 1902, 1903; Seabra, 1911; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1979e; Dawson, 1986; Antunes et al., 1988; Almeida, 1996; Lourie et al., 2004, 2016; Curtis and Vincent, 2005; Ribeiro et al., 2006, 2012; Gamito, 2008; Henriques et al., 2007, 2008; Neves et al., 2008; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Caldwell and Vincent, 2012; Boehm et al., 2013; Carneiro et al., 2014; Correia, 2014; Vieira et al., 2014; Correia et al., 2015, 2016; Lourie et al., 2016; Woodall et al., 2017; Froese and Pauly, 2019), Azores (Hilgendorf, 1888b; Regan, 1903a; Nobre, 1924, 1930; Fowler, 1936a; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Ré, 1979a; Dawson, 1986; Patzner et al., 1990, 1992; Azevedo et al., 1991; Lloris et al., 1991; Arruda, 1992, 1997;

Arruda *et al.*, 1992; Santos *et al.*, 1994a, 1995a, 1997; Wirtz, 1994; Azevedo, 1995; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Lourie *et al.*, 2016; Fritzsche, 2016a; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1843-1860; Günther, 1870; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1979e; Dooley *et al.*, 1985; Dawson, 1986; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Fritzsche, 2016a; Froese and Pauly, 2019(?)).

***Hippocampus hippocampus* (Linnaeus 1758)**

Common name: **Short snouted seahorse**; **Cavalo-marinho-de-focinho-curto**, Cavalinho-do-mar, Cavalinho, Cavalinho-marinho, Hipocampo, Lincorna.

References: mainland Portugal (Linnaeus 1758; Vandelli, 1771; de Brito Capello, 1869a, b, 1880; Osório, 1888; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895c, 1903, 1935; Vieira, 1897; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1979e; Dawson, 1986; Cabral, 1999; Erzini *et al.*, 2001a; Borges *et al.*, 2002; Palma *et al.*, 2003; Garnerot *et al.*, 2004; Lourie *et al.*, 2004, 2016; Curtis and Vincent, 2005; Ribeiro *et al.*, 2006, 2012; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Caldwell and Vincent, 2012; Boehm *et al.*, 2013; Carneiro *et al.*, 2014; Correia, 2014; Vieira *et al.*, 2014; Correia *et al.*, 2015, 2016; Lourie *et al.*, 2016; Woodall *et al.*, 2017; Froese and Pauly, 2019), Azores (Drouët, 1861b; Hilgendorf, 1888a; Collins, 1954; Osório de Castro, 1967; Arruda, 1992, 1997; Santos *et al.*, 1994a, 1995a; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Lourie *et al.*, 2016) and Madeira (Lowe, 1837; Drouët 1861b; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Wirtz, 1998; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Fritzsche, 2016a; Froese and Pauly, 2019). Three neotypes of *Syngnathus hippocampus* Linnaeus 1758, synonym of *H. hippocampus*, caught off in the West Coast of Spain and Portugal are in the NHMUK collections (NHMUK 1872.2.6.3-5, set of three specimens).

***Nerophis lumbriciformis* (Jenyns 1835)**

Common name: **Worm pipefish**; **Marinha-lombriga**, Cavalinho-marinho, Marinha.

References: mainland Portugal (Osório, 1888, 1895; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1979e; Dawson, 1986, 1990; Arruda, 1990; Ribeiro *et al.*, 2006, 2012; Henriques *et al.*, 2007, 2008; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Fritzsche, 2016a; Ryland and Lancaster, 2017; Froese and Pauly, 2019) and Azores (PECS) (MNHN-IC-0000-6074, n.d.; Carneiro *et al.*, 2014).

***Nerophis maculatus* Rafinesque 1810**

Common name: Spotted pipefish; **Marinha-sarapintada**, Cavalinho-marinho, Marinha.

References: mainland Portugal (Nobre, 1895a, 1895c, 1935; Vieira, 1897; Seabra, 1911; Osório de Castro, 1967; Wheeler, 1979e; Dawson, 1986; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Azores (Hilgendorf, 1888a; Fowler, 1936b; Collins, 1954; Osório de Castro, 1967; Dawson, 1986; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Nerophis ophidion* (Linnaeus 1758)**

Common name: **Straightnose pipefish**; **Marinha-fina**, Cavalinho-marinho, Marinha.

References: mainland Portugal (Vandelli, 1771; Osório, 1888, 1917; Vieira, 1897; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Muus and Dahlström, 1978; Wheeler, 1979e; Dawson, 1986; Antunes *et al.*, 1988; Garnerot *et al.*, 2004; Ribeiro *et al.*, 2006, 2008, 2012; Chaves, 2008; Henriques *et al.*, 2008; França *et al.*, 2011; Carneiro *et al.*, 2014; Vieira *et al.*, 2014; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Wirtz, 1998; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Hamilton *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Syngnathus abaster* Risso 1827**

Common name: Black-striped pipefish; **Marinha-do-rio**, Agulhinha, Agulhinha-do-rio, Cavalinho-marinho, Marinha.

References: mainland Portugal (Osório, 1888, 1895; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Dawson, 1986; Antunes *et al.*, 1988; Arruda *et al.*,

1988; Moreira *et al.*, 1992; Cabral, 1999; Cakic *et al.*, 2002; Garnerot *et al.*, 2004; Ribeiro *et al.*, 2006, 2012; Silva *et al.*, 2006; Henriques *et al.*, 2008; Veiga *et al.*, 2009; França *et al.*, 2011; Carneiro *et al.*, 2014; Vieira *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Azores (PECS) (MNHN-IC-1955-0044, 1955; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is Mediterranean Sea, Sea of Marmara, Black Sea, and from Bay of Biscay south to Portugal in the Eastern Atlantic. Carneiro *et al.* (2014) considered very doubtful the occurrence of this species in the geographic coordinates indicated in the MNHN-IC-1955-0044 record. However, this occurrence record was confirmed by information from the MNHN.

***Syngnathus acus* Linnaeus 1758**

Common name: **Greater pipefish; Marinha-comum**, Agulhinha, Cavalo-marinho, Chicote, Culebra, Culevra, Marinha, Peixe-pau.

References: mainland Portugal[▲] (Vandelli, 1771; de Brito Capello, 1869a, 1872, 1873, 1880; Osório, 1888, 1895; Nobre, 1895a, c, 1903, 1935; Vieira, 1897; Bragança, 1903; Seabra, 1911; Nobre *et al.*, 1915; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1979e; Dawson, 1986; Antunes *et al.*, 1988; Arruda *et al.*, 1988; Almeida, 1996; Cabral, 1999; Jorge *et al.*, 2002; Palma *et al.*, 2003; Garnerot *et al.*, 2004; Ribeiro *et al.*, 2006, 2012; Henriques *et al.*, 2007, 2008; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Vieira *et al.*, 2014; Martins *et al.*, 2015; Ryland and Lancaster, 2017; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a; Silva, 1893; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Wheeler, 1979e; Dooley *et al.*, 1985; Dawson, 1986; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1994b, 1997; Wirtz, 1994; Azevedo, 1995; Gonçalves *et al.*, 1995; Santos and Nash, 1995; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Günther, 1870; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Dawson, 1986; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Syngnathus fucus* Storer 1839**

Common name: Northern pipefish; **Marinha-setentrional**, Marinha.

References: Azores (Collins, 1954; Arruda, 1997). Collins (1954) and Arruda (1997) mention the occurrence of the *Syngnathus pellegrini* Fowler 1919 in the Azores. The species *S. pellegrini* is synonym of *S. fucus*. According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is the Western North Atlantic, from Canada to Florida. However, specimens are needed for confirmation in the Azores.

***Syngnathus pelagicus* Linnaeus 1758**

Common name: Sargassum pipefish; **Marinha-do-sargaço**, Cavalo-marinho, Marinha.

References: mainland Portugal (Osório, 1888; Nobre, 1935), Azores (PECS) (Murray and Hjort, 1912; Guiart, 1935(?); Osório de Castro, 1967; ROM 36947, 1969; Carneiro *et al.*, 2014) and Madeira (Osório de Castro, 1967). According to Eschmeyer *et al.* (2019) *S. pelagicus* is present in the Atlantic, associated with floating sargassum (Dawson, 1990). The historical record of this species by Osório (1888) is the northernmost occurrence in the Eastern Atlantic, if the taxonomic status of the Eastern Atlantic population will be confirmed (Froese and Pauly, 2019). Nobre (1935) indicated it as a very rare species. Zugmayer (1911b) and Fowler (1936a) reported the occurrence of this species in the Atlantic, Southwest of Azores, from 30° to 31°N, 42°W and off Azores, from 24°21'N, 34°32'W to 23°59'N, 37°59'W'. These geographic coordinates correspond to a range South and SW of Azores, but out of the Azorean PECS area. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. However, its presence in Madeira is likely due to its wide distribution and association with *Sargassum* seaweed rafts in open oceanic waters. Thus, specimens are needed for confirmation in Madeira Archipelago.

***Syngnathus phlegon* Risso 1827**

Common name: No English common name; **Marinha-espinhosa**.

References: mainland Portugal (Dawson, 1986(?); Carneiro *et al.*, 2014; Wiswedel, 2014a; Eschmeyer *et al.*, 2019) and Madeira (NHMUK 2017.3.9.13-27, 2017).

Syngnathus rostellatus Nilsson 1855

Common name: **Nilsson's pipefish; Marinha-cabeça-chata.**

References: mainland Portugal ([Almeida, 1986](#); [Henriques et al., 2008](#); [Hablützel and Wilson, 2011](#); [Carneiro et al., 2014](#); [Wiswedel, 2014b](#); [Froese and Pauly, 2019](#)).

Syngnathus tenuirostris Rathke 1836

Common name: Narrow-snouted pipefish; **Marinha-de-focinho-estreito**, Cavalo Marinho.

References: mainland Portugal (†) ([de Brito Capello, 1869a, b, 1873](#); [Osório, 1888](#); [Girard, 1892](#); [Seabra, 1911](#); [Nobre, 1935](#)). Regionally extinct, the last bibliographical reference dates from 1869 in the Lagos Bay (Algarve) and 1888 in Setúbal (de Brito Capello, 1869a, b; Osório, 1888). Thus, this species is not considered in the total number of valid species.

Syngnathus typhle Linnaeus 1758

Common name: **Broadnosed pipefish; Marinha-focinho-grosso**, Agulhinha, Cavalo-marinho, Cobra-de-água, Marinha, Peixe-pau.

References: mainland Portugal ([de Brito Capello, 1872, 1873, 1880](#); [Osório, 1888](#); [Vieira, 1893, 1897](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Muus and Dahlström, 1978](#); [Wheeler, 1979e](#); [Dinis, 1985](#); [Dawson, 1986](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Garnerot et al., 2004](#); [Henriques et al., 2007, 2008](#); [Neves et al., 2008](#); [Ribeiro et al., 2008, 2012](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Pollock, 2014](#); [Vieira et al., 2014](#); [Ryland and Lancaster, 2017](#); [Froese and Pauly, 2019](#)) and Madeira ([Osório de Castro, 1967](#)). The southernmost occurrence record of this species in the Eastern Atlantic is at Saharan Fan, south of Canary Islands (MNHN-IC-0000-9265). Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. Specimens are needed for confirmation in Madeira Archipelago.

Family Dactylopteridae

Dactylopterus volitans (Linnaeus 1758)

Common name: **Flying gurnard; Cabrinha-de-leque**, Apisto, Cabra-voadora, Ganeta, Ganete, Pássaro-ganê, Peixe-ganeta, Peixe-ganete, Peixe-pássaro, Peixe-pássaro-ganê, Peixe-voador, Ruivo, Voador, Voador-de-fundo.

References: mainland Portugal ([Posser, 1880\(?\)](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Monod, 1979d](#); [Fischer et al., 1981, 1987](#); [Dooley et al., 1985](#); [Roux, 1986b](#); [Carneiro et al., 2014](#); [Lorance et al., 2015c](#); [Smith-Vaniz, 2016a](#); [Froese and Pauly, 2019](#)), Azores ([Roux, 1986b](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Lorance et al., 2015c](#); [Smith-Vaniz, 2016a](#); [Froese and Pauly, 2019](#)) and Madeira ([Fischer et al., 1981](#); [Roux, 1986b](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#); [Wirtz et al., 2008\(?\)](#); [Carneiro et al., 2014](#); [Lorance et al., 2015c](#); [Smith-Vaniz, 2016a](#); [Froese and Pauly, 2019](#)).

Order Gobiiformes

Family Gobiidae

Aphia minuta (Risso 1810)

Common name: **Transparent goby; Caboz-transparente**, Alcaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal ([Nobre, 1895c, 1903, 1935](#); [Osório, 1895](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Miller, 1979, 1986](#); [Fischer et al., 1987](#); [Arruda et al., 1988](#); [Jorge et al., 2002](#); [Garnerot et al., 2004](#); [da Cunha and Antunes, 2008](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Kovačić and Patzner, 2011\(?\)](#); [Ribeiro et al., 2012](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Buenia affinis Iljin 1930

Common name: **De Buen's goby; Caboz-de-De Buen.**

References: Madeira ([Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#)).

Buenia jeffreysii (Günther 1867)

Common name: Jeffrey's goby; **Caboz-de-Jeffrey**.

References: mainland Portugal ([Gordo et al., 1990](#); [Henriques et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#)).

Chromogobius britoi Van Tassell 2001

Common name: Brito's goby; **Caboz-de-Brito**.

References: mainland Portugal ([Beldade et al., 2006a](#); [Kovačić and Patzner, 2011](#); [Patzner et al., 2011](#); [Carneiro et al., 2014](#); [Herler et al., 2014a](#); [Froese and Pauly, 2019](#)) and Madeira ([Wirtz et al., 1998](#), [2008](#); [Van Tassell, 2001](#); [Brito et al., 2002](#); [Floeter et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Herler et al., 2014a](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Crystallogobius linearis (Düben 1845)

Common name: **Crystal goby; Caboz-de-cristal**.

References: mainland Portugal ([Miller, 1979](#), [1986](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Kovačić et al., 2014d](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Kovačić et al., 2014d](#); [Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Deltentosteus collarianus (Risso 1820)

Common name: Toothed goby; **Caboz-dentudo**.

References: mainland Portugal ([Maul, 1976](#); [Hureau and Monod, 1979](#); [Miller, 1986](#); [Kovačić and Golani, 2007](#); [Carneiro et al., 2014](#); [Kovačić et al., 2014e](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22808, 1967).

Deltentosteus quadrimaculatus (Valenciennes 1837)

Common name: **Four-spotted goby; Caboz-de-quatro-manchas**, Alcaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal ([Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976](#); [Miller, 1979, 1986, 1990](#); [Fischer et al., 1987](#); [Rebelo 1992](#); [Borges et al., 2002](#); [Garnerot et al., 2004](#); [Gomes and Olim, 2007](#); [da Cunha and Antunes, 2008](#); [Henriques et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). Two specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22832, 1967; MMF 22828, 1967).

Gnatholepis thompsoni Jordan 1904

Common name: Goldspot goby; **Caboz-de-pinta-dourada**, Alcaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal (?) ([Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Araújo and Freitas, 2002](#); [Brito et al., 2002](#); [Freitas and Araújo, 2006](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [EMEPC, 2010](#); [Kovačić and Patzner, 2011](#); [Larson and Buckle, 2012](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Williams et al., 2015i](#); [Miller and Murdy, 2016](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)). Two specimens are in the MMF collections (MFF 34338 and MFF 35075). According to Eschmeyer et al. (2019) the actual geographic distribution of this species is the Western and Eastern Atlantic islands. The occurrence of this species in mainland Portugal reported by Froese and Pauly (2019) - List of marine fishes from Portugal - is based on Araújo and Freitas (2002). However, the latter authors only mention the presence of *G. thompsoni* in Madeira Island, indicating that is the most northerly sighting of this species in the Eastern Atlantic. The presence of this species in mainland Portugal was erroneously considered by Carneiro et al. (2014) based on the FishBase records. Thus, this species is not considered in the total number of valid species for mainland Portugal.

Gobius ater Bellotti 1888

Common name: **Bellotti's goby; Caboz-de-Bellotti**.

References: mainland Portugal ([Pombo et al., 2002](#); [Garnerot et al., 2004](#); [Carneiro et al., 2014](#)). According to Kovačić et al. (2014a) and Eschmeyer et al. (2019) this species is endemic to the Mediterranean Sea. However, [Pombo et al. \(2002\)](#) report the occurrence of *G. ater* in the Ria de Aveiro in the years of 1988, 1999 and 2000. Since [Pombo et al. \(2002\)](#) and [Garnerot et al. \(2004\)](#) no other author describes the presence of this species in mainland Portugal.

Gobius auratus Risso 1810

Common name: Golden goby; **Caboz-dourado**, Acaboz, Alcabroz, Caboz, Cabroz, Caboz-das-pedras, Cadoz.

References: mainland Portugal ([Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Miller, 1986](#); [Almeida, 1996](#); [Henriques et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#)) and Madeira ([Zugmayer, 1933](#); [Fowler, 1936b](#); [Osório de Castro, 1967](#); [Maul, 1976\(?\)](#); [Miller, 1979](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). According to [Kovačić et al. \(2014b\)](#) *G. auratus* is a Mediterranean endemic species and previous records of this species from the Atlantic Ocean, from northern Spain to Madeira, Portugal and the Canary Islands, pertain to another species. However, [Eschmeyer et al. \(2019\)](#) mention that the current geographic distribution of *G. auratus* is the Mediterranean Sea and, in the Eastern Atlantic, from Spain to Madeira and Canary islands. One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22809, 1967).

Gobius bucchichi Steindachner 1870

Common name: **Bucchich's goby**; **Caboz-mosqueado**.

References: mainland Portugal ([Miller, 1986](#); [Kovačić and Golani, 2007](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Kovačić et al., 2015a](#); [Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Gobius cobitis Pallas 1814

Common name: **Giant goby**; **Caboz-cabeçudo**, Acaboz, Alcabroz, Caboz, Cabroz, Cadoz, Escama, Peixe-escama.

References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Girard, 1892](#); [Osório, 1895, 1896](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Arruda, 1979b](#); [Gomes and Almeida, 1979](#); [Miller, 1979, 1986](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Faria and Almada, 1995](#); [Henriques et al., 1999, 2007, 2008](#); [da Cunha and Antunes, 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Golani and Fricke, 2018](#); [Eschmeyer et al., 2019](#)). Three syntypes of *Gobius algarbiensis* de Brito Capello 1880, synonym of *G. cobitis*, were caught off near Lisbon, Setúbal and Lagos (Portugal) and its whereabouts are unknown.

Gobius couchi Miller & El-Tawil 1974

Common name: Couch's goby; **Caboz-de-Couch**.

References: mainland Portugal ([Ribeiro et al., 2006, 2012](#); [Kovačić and Patzner, 2011](#)).

Gobius cruentatus Gmelin 1789

Common name: **Red-mouthed goby**; **Caboz-de-boca-vermelha**, Acaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal ([Nobre, 1903, 1935](#); [Seabra, 1911](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Miller, 1979, 1986, 1990](#); [Almeida, 1996](#); [Borges et al., 2002](#); [Ribeiro et al., 2006, 2012](#); [Henriques et al., 2007, 2008](#); [Gonçalves et al., 2010](#); [Kovačić and Patzner, 2011](#); [Baldock and Kay, 2012](#); [Carneiro et al., 2014](#); [Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Gobius gasteveni Miller 1974

Common name: Steven's goby; **Caboz-de-escama**, Acaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal ([Henriques et al., 1999, 2007, 2008](#); [Beldade et al., 2007](#); [da Cunha and Antunes, 2008](#); [Monteiro et al., 2008](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [Kovačić and Patzner, 2011](#); [Baldock and Kay, 2012](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#)) and Madeira ([Miller, 1974, 1984, 1986, 1990](#); [Hureau and Monod, 1979](#); [Lloris et al., 1991](#); [Alberto et al., 1999](#); [Ahnelt and Dorda, 2004](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). One paratype of this species is in the MMF collections (MFF 13096A).

Gobius geniporus Valenciennes 1837

Common name: **Slender goby**; **Caboz-de-bochechas porosas**.

References: mainland Portugal ([Gonçalves et al., 2010](#)). According to Miller (1986), Eschmeyer *et al.* (2019) and Froese and Pauly (2019) *G. geniporus* is an endemic species of the Mediterranean Sea. However, three specimens of this species were caught off in the Arade River estuary (Algarve) in 2010.

***Gobius niger* Linnaeus 1758**

Common name: **Black goby; Caboz-negro**, Acaboz, Alcabroz, Caboz, Cabozeira, Cabroz, Cadoz, Cardosa, Cardoza.

References: mainland Portugal ([Vandelli, 1771](#); [de Brito Capello, 1868, 1869a, b, 1880](#); [Steindachner, 1868a](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Miller, 1979, 1986](#); [Maugé, 1986](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Cabral, 1999](#); [Garnerot et al., 2004](#); [Ribeiro et al., 2006, 2012](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [da Cunha and Antunes, 2008](#); [Neves et al., 2008](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Kovačić et al., 2015b](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Drouët, 1861b](#); [Ramos, 1869](#); [Hilgendorf, 1888a](#); [Sampaio, 1904](#), [Nobre, 1924, 1930](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Arruda, 1992, 1997](#); [Kovačić and Patzner, 2011](#)) and Madeira ([Lowe, 1837](#); [Osório de Castro, 1967\(?\)](#); [EMEPC, 2010](#); [Kovačić and Patzner, 2011](#); [Almada et al., 2015](#); [Araújo and Wirtz, 2015](#); [Kovačić et al., 2015b](#); [POGIS, 2016](#)). According to Santos *et al.* (1997) the records of *G. niger* in the Azores were misidentifications of *Gobius paganellus* Linnaeus 1758. Thus, this species is not considered in the total number of valid species to the Azores.

***Gobius ophiocephalus* Pallas 1814**

Common name: **Grass goby; Caboz-cabeça-de-cobra**.

References: Azores (?) ([Sampaio, 1904](#); [Arruda, 1997](#)). Sampaio (1904) mentions the presence of this species as *Gobius iota* Cuvier 1829 and Arruda (1997) as *Zosteriessor ophiocephalus* (Pallas 1814), and both are synonyms of *G. ophiocephalus*. According to Herler *et al.* (2014b) and Eschmeyer *et al.* (2019) the current geographic distribution of *G. ophiocephalus* is the Black Sea, Sea of Azov and Mediterranean Sea (including adjacent estuaries). Thus, this species is not considered in the total number of valid species to the Azores.

***Gobius paganellus* Linnaeus 1758**

Common name: **Rock goby; Caboz-da-rocha**, Acaboz, Alcabroz, Bochecha, Caboz, Cabozeira, Cabroz, Cadoz, Escama, Joana, Paganel, Peixe-escama, Velha, Viúva.

References: mainland Portugal ([de Brito Capello, 1867a, 1868, 1869a, b, 1880](#); [Osório, 1888, 1895](#); [Girard, 1892](#); [Vieira, 1897, 1898](#); [Nobre, 1903, 1904, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Dooley et al., 1985](#); [Miller, 1979, 1986, 1990](#); [Maugé, 1986](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Arruda, 1990](#); [Faria and Almada, 1995](#); [Almeida, 1996](#); [Cabral, 1999](#); [Garnerot et al., 2004](#); [Beldade et al., 2006b](#); [Ribeiro et al., 2006, 2012](#); [Henriques et al., 2007, 2008](#); [da Cunha and Antunes, 2008](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Golani and Fricke, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Drouët, 1861b](#); [Ramos, 1869](#); [Hilgendorf, 1888a, b](#); [Regan, 1903a](#); [Sampaio, 1904](#); [Nobre, 1924, 1930](#); [Fowler, 1916, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Wood and Williams, 1974](#); [Arruda, 1979a, 1980b, 1992, 1997](#); [Miller, 1984](#); [Maugé, 1986](#); [Miller, 1986, 1990](#); [Azevedo et al., 1990, 1991, 1992, 1995](#); [Patzner, 1990](#); [Patzner and Santos, 1990a, c, e, 1993](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1994a, b, 1997](#); [Wirtz, 1990a, 1994](#); [Lloris et al., 1991](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Gonçalves et al., 1995](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Porteiro et al., 2010](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Diogo and Pereira, 2014](#); [Miller and Murdy, 2016](#); [Golani and Fricke, 2018](#); [Almeida and Biscoito, 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Cuvier and Valenciennes, 1837](#); [Lowe, 1839, 1849](#); [Günther, 1861a](#); [Von Martens, 1876](#); [Collett, 1890b](#); [Boulenger, 1916](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Augier, 1985](#); [Miller, 1986, 1990](#); [Lloris et al., 1991](#); [Falcón et al., 2000](#); [Araújo et al., 2005](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#)).

Miller and Murdy, 2016; POGIS, 2016; Golani and Fricke, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Gobius roulei de Buen 1928

Common name: **Roule's goby; Caboz-de-Roule.**

References: mainland Portugal (Maul, 1976; Hureau and Monod, 1979; Miller, 1986; Kovačić, 1995, 2001; Louisy, 2002; Kovačić and Golani, 2006; Henriques *et al.*, 2008; Liu *et al.*, 2009; Gonçalves *et al.*, 2010; Kovačić *et al.*, 2012, 2014c; Carneiro *et al.*, 2014; Miller and Murdy, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019; Kovačić and Schembri, 2019) and Madeira (?) (Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of this species is the Mediterranean Sea, and in the Eastern Atlantic, southern Portugal and Canary Islands. The presence of this species in Madeira was erroneously considered by Carneiro *et al.* (2014) and it is not included in the total number of valid species to Madeira. One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22861, 1967).

Gobius xanthocephalus Heymer & Zander 1992

Common name: **Golden goby; Caboz-de-cabeça-dourada.**

References: mainland Portugal (Almeida and Arruda, 1998; Henriques *et al.*, 2007, 2008; Gonçalves *et al.*, 2010; Kovačić and Patzner, 2011; Carneiro *et al.*, 2014; Herler *et al.*, 2014c; Froese and Pauly, 2019) and Madeira (Miller, 1986; EMEPC, 2010; Herler *et al.*, 2014c; Almada *et al.*, 2015; Miller and Murdy, 2016; POGIS, 2016; Froese and Pauly, 2019). According to Herler *et al.* (2014c) this species was previously (prior to 1992) identified as *Gobius auratus* (Risso 1810) and is easily misidentified, especially with the yellow morph of *G. auratus*.

Gobiusculus flavescens (Fabricius 1779)

Common name: **Two-spotted goby; Caboz-de-duas-manchas**, Alcaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Miller, 1986; Almeida, 1999; Moura, 2004; da Cunha and Antunes, 2008; Henriques *et al.*, 2007, 2008; Kovačić and Patzner, 2011; Carneiro *et al.*, 2014; Herler *et al.*, 2014d; Eschmeyer *et al.*, 2019).

Lebetus guilleti (Le Danois 1913)

Common name: **Guillet's goby; Caboz-de-Guillet.**

References: mainland Portugal (Miller, 1971, 1986, 1990; Ré, 1981; Henriques *et al.*, 2008; Baldock and Kay, 2012; Carneiro *et al.*, 2014; Riolo and Betti, 2015; Miller and Murdy, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Carneiro *et al.*, 2014; Herler *et al.*, 2014e; Araújo and Wirtz, 2015; Miller and Murdy, 2016).

Lebetus scorpioides (Collet 1874)

Common name: Diminutive goby; **Caboz-anão.**

References: mainland Portugal (Ré, 1981; Henriques *et al.*, 2008; Kovačić and Patzner, 2011; Baldock and Kay, 2012; Carneiro *et al.*, 2014) and Madeira (Floeter *et al.*, 2008). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2018) the current geographic distribution of *L. scorpioides* is the Eastern North Atlantic and the southern distribution limit is the Bay of Biscay. However, the occurrence of this species has already been confirmed for mainland Portugal. Rare species in Madeira.

Lesueurigobius friesii (Malm 1874)

Common name: **Fries's goby; Caboz-de-escama-grande**, Alcadoz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández *et al.*, 1981; Miller, 1986; ICES, 2005; da Cunha and Antunes, 2008; Silva *et al.*, 2009; Kovačić and Patzner, 2011; Carneiro *et al.*, 2014; Garcia, 2015g; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005181).

Lesueurigobius heterofasciatus Maul 1971

Common name: No English common name; **Caboz-listado.**

References: Madeira (Maul, 1971b, 1976; Miller, 1979, 1986, 1990; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Kovačić and Patzner, 2011; Carneiro *et al.*, 2014; Miller and Murdy, 2016;

Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype and two paratypes of this species are in the MMF collections (MMF 22818, MMF 11587 and MMF 22824, respectively).

***Lesueurigobius sanzi* (de Buen 1918)**

Common name: Sanzo's goby; **Caboz-de-Sanzo**, Alcaboz, Alcabroz, Caboz, Cabroz, Caboz-de-bandeira, Cadoz.

References: mainland Portugal (Osório de Castro, 1954; Almeida and Gomes, 1978; Hureau and Monod, 1979; INIP, 1981b; Dinis, 1985; Miller, 1986, 1990; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; da Cunha and Antunes, 2008; Chaves, 2008; Henriques *et al.*, 2008; Carneiro *et al.*, 2014; Costa, 2014; Miller and Murdy, 2016; Gerovasileiou *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (MNHN-IC-2005-2482, 1926).

***Lesueurigobius suerii* (Risso 1810)**

Common name: Lesueur's goby; **Caboz-de-Lesueur**, Alcaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal (Bragança, 1902; Nobre, 1935; MLFPI31, 2009; MLFPI32, 2009; Carneiro *et al.*, 2014), Azores (Vaillant, 1919; Fowler, 1936b; Santos *et al.*, 1997; Carneiro *et al.*, 2014) and Madeira (MNHN-IC-1883-0029, 1882; Floeter *et al.*, 2008; Carneiro *et al.*, 2014).

***Mauligobius maderensis* (Valenciennes 1837)**

Common name: No English common name; **Caboz-da-Madeira**, Caboz-de-escama.

References: Azores (Regan, 1903a; Miller, 1986(?); MCZ 13141, 1994; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019) and Madeira (Cuvier and Valenciennes, 1837; Miller, 1979, 1986, 1990; Augier, 1985; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Kovačić and Patzner, 2011; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Miller and Murdy, 2016; POGID, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Six syntypes of *Gobius maderensis* Valenciennes 1837 caught off in Madeira are in the MNHN collections (MNHN-IC-A-1164, set of three specimens and MNHN-IC- A-1160, set of three specimens). *Gobius maderensis* is synonym of *M. maderensis*.

***Neogobius fluviatilis* (Pallas 1814)**

Common name: Monkey goby; **Caboz-do-rio**.

References: mainland Portugal (?) (Osório, 1896; Seabra, 1911; Nobre, 1931). According to Eschmeyer *et al.* (2019) the habitat of this species is the freshwater, brackish and marine waters and its current geographic distribution is the Black Sea and Caspian Sea basins. Nobre (1931) referred that this species was introduced in Leça river (near Matosinhos). Since Nobre (1931) no other author described the presence of this species in mainland Portugal and its occurrence in this EEZ was not yet confirmed by more recent data. Thus, this species is not considered in the total number of valid species.

***Pomatoschistus knerii* (Steindachner 1861)**

Common name: Kner's goby; **Caboz-de-Kner**.

References: mainland Portugal (Leitão *et al.*, 2007; Henriques *et al.*, 2008; Herler *et al.*, 2014f).

***Pomatoschistus lozanoi* (de Buen 1923)**

Common name: Lozano's goby; **Caboz-de-Lozano**.

References: mainland Portugal (Miller, 1986; Arruda and Azevedo, 1987; Jorge *et al.*, 2002; da Cunha and Antunes, 2008; Henriques *et al.*, 2008; Kovačić and Patzner, 2011; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Herler *et al.*, 2014g; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Five specimens/vouchers of this species are deposited in the NCBI (GenBank JQ775030.1-33.1, set of five specimens and JQ775039.1) (see: <https://www.ncbi.nlm.nih.gov/nuccore/>).

***Pomatoschistus marmoratus* (Risso 1810)**

Common name: Marbled goby; **Caboz-marmoreado**.

References: mainland Portugal (Miller, 1979, 1986; Antunes *et al.*, 1988; Garnerot *et al.*, 2004; Henriques *et al.*, 2007, 2008; da Cunha and Antunes, 2008; Carneiro *et al.*, 2014; Herler *et al.*, 2014h; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Pomatoschistus microps* (Krøyer 1838)**

Common name: **Common goby; Caboz-comum.**

References: mainland Portugal ([Miller, 1979, 1986, 1990](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Moreira et al., 1992](#); [Cabral, 1999](#); [Vieira, et al., 2002](#); [Garnerot et al., 2004](#); [Ribeiro et al., 2006, 2012](#); [da Cunha and Antunes, 2008](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Veiga et al., 2009](#); [Freyhof, 2010](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

***Pomatoschistus minutus* (Pallas 1770)**

Common name: **Sand goby; Caboz-da-areia**, Alcaboz, Alcabroz, Caboz, Cabroz, Cadoz.

References: mainland Portugal ([MNHN-IC-1887-0186, 1883](#); [Osório, 1888](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Muus and Dahlström, 1978](#); [Miller, 1979\(?\)](#), [1986](#); [INIP, 1981a](#); [Martins, 1981](#); [Arruda et al., 1988](#); [Cabral, 1999](#); [Jorge et al., 2002](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Gomes and Olim, 2007](#); [da Cunha and Antunes, 2008](#); [Chaves, 2008](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Ribeiro et al., 2006, 2012](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [Silva, 2010](#); [França et al., 2011](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

***Pomatoschistus norvegicus* (Collett 1902)**

Common name: **Norway goby; Caboz-norueguês.**

References: mainland Portugal ([Gordo et al., 1990](#); [Kovačić and Patzner, 2011](#)). The first specimen of this species was caught off at 27 NM W off Espinho in November 20, 1984.

***Pomatoschistus pictus* (Malm 1865)**

Common name: Mediterranean painted goby; **Caboz-de-manchas**, Góbio-da-areia.

References: mainland Portugal ([Cunha, 1984](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Ribeiro et al., 2006, 2012](#); [Henriques et al., 2007, 2008](#); [da Cunha and Antunes, 2008](#); [Gonçalves et al., 2010](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#)), Azores ([Patzner and Santos, 1990a, c, e](#); [Patzner et al., 1990, 1992](#); [Ré, 1990](#); [Santos et al., 1990b, 1997](#); [Azevedo et al., 1991, 1992](#); [Nash et al., 1991](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Arruda, 1997](#); [Floeter et al., 2008](#); [Veiga et al., 2009](#); [Bertонcini et al., 2010](#); [Porteiro et al., 2010](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#)) and Madeira ([Wirtz, 1998](#); [Araújo et al., 2005](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

***Thorogobius ephippiatus* (Lowe 1839)**

Common name: Leopard-spotted goby; **Caboz-leopardo**, Bochecha-pintada, Caboz, Caboz-de-escama.

References: mainland Portugal ([Almeida and Gomes, 1978](#); [Hureau and Monod, 1979](#); [Miller, 1986](#); [Almeida and Gomes, 1978](#); [Henriques et al., 2007, 2008](#); [Abecasis et al., 2009](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Azevedo et al., 1990](#); [Patzner and Santos, 1990a, c, e](#); [Patzner et al., 1990, 1992](#); [Wirtz, 1990a, 1994](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Harmelin-Vivien et al., 2001](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#)) and Madeira ([Lowe, 1839, 1840b, 1844a, 1849](#); [Miller, 1979, 1986, 1990](#); [Lloris et al., 1991](#); [Falcón et al., 2000](#); [Araújo et al., 2005](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Miller and Murdy, 2016](#); [POGIS, 2016](#); [Kovačić and Svensen, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Gobius ephippiatus* Lowe 1839 was caught off in Madeira and its whereabouts are unknown. *Gobius ephippiatus* is synonym of *T. ephippiatus*.

***Vanneaugobius canariensis* Van Tassel, Miller & Brito 1988**

Common name: No English common name; **Caboz-das-Canárias.**

References: Madeira ([Debelius, 1998](#); [Wirtz, 1998](#); [Freitas and Araújo, 2006](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Kovačić and Patzner, 2011](#); [Carneiro et al., 2014](#); [Herler et al., 2015c](#); [Miller and Murdy, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Order Istiophoriformes

Family Xiphiidae

Xiphias gladius Linnaeus 1758

Common name: **Swordfish; Espadarte**, Agulha, Agulhão, Catana, Espada, Espadão, Peixe-agulha, Peixe-espada.

References: mainland Portugal▲ (Vandelli, 1771; de Brito Capello, 1867a, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Osório, 1888; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895a, 1903, 1935; Bragança, 1903; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979g; Fischer et al., 1981, 1987; INIP, 1981a; Martins, 1981; Dinis and Marecos, 1984; Nishikawa et al., 1985; Nakamura, 1985, 1986; Erzini et al., 2001b; Borges et al., 2002; Santos et al., 2002b, c; ICES, 2005; Gomes and Olim, 2007; Bordalo-Machado et al., 2009; Carneiro et al., 2014; OCEANA, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Sampaio, 1904; Ribeiro, 1936; Ferreira, 1937a; Collins, 1954; Osório de Castro, 1967; Parin, 1970; Ueyanagi et al., 1970; Wheeler, 1978; Fischer et al., 1981; Palko et al., 1981; Martins, 1982a; Rey, 1984; Nishikawa et al., 1985; Nakamura, 1985, 1986; Bard, 1987; Rey, 1987; Monteiro and Lopes, 1990; Lloris et al., 1991; Monteiro, 1991; Silva, 1992; Clarke and Martins, 1995; Clarke et al., 1995; Esteves et al., 1995; Pinho and Pereira, 1995a; Simões and Silva, 1995; Isidro, 1996; Menezes, 1996; Smale, 1996; Arruda, 1997; Santos et al., 1997; Cardigos et al., 2005; Menezes et al., 2006; Borges et al., 2010; Porteiro et al., 2010, 2017(?); Besugo, 2013; Martins, 2013; Carneiro et al., 2014; Nakamura, 2016; Fauconnet et al., 2019) and Madeira▲ (Lowe, 1849; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979g; Fischer et al., 1981; Nakamura, 1986, 2016; Lloris et al., 1991; Severino, 2004; Carneiro et al., 2014).

Family Istiophoridae

Istiophorus platypterus (Shaw 1792)

Common name: **Indo-Pacific sailfish; Veleiro-do-Indo-Pacífico**, Bicuda, Bicudo, Caravela, Espadarte, Espadarte-veleiro, Peco, Peito, Peixe-de-vela, Peto, Veleiro, Veleiro-americano, Veleiro-do-Atlântico, Voador.

References: mainland Portugal▲ (Fischer et al., 1981, 1987; Nishikawa et al., 1985; Nakamura, 1986; Carneiro et al., 2014; Nakamura and Collette, 2016), Azores (Fischer et al., 1981; Nishikawa et al., 1985; Martin, 1986; Nakamura, 1985, 1986; Arruda, 1997; Santos et al., 1997; Carneiro et al., 2014; Nakamura and Collette, 2016; Porteiro et al., 2017(?)) and Madeira (Fischer et al., 1981; Nakamura, 1986; Carneiro et al., 2014; Nakamura and Collette, 2016). There are records of the occurrence of *Istiophorus albicans* (Latreille 1804) in Portugal. However, according to Collette et al. (2006) the species *I. albicans* (Latreille 1804) is a synonym of *I. platypterus*. According to Nakamura and Collette (2016) the Atlantic population of the species previously known as *I. albicans* have not any morphological or genetic differences from the Indo-Pacific population of *I. platypterus*. For Eschmeyer et al. (2019) *I. platypterus* is distributed in the Atlantic ocean if *albicans* is treated as a synonym.

Kajikia albida (Poey 1860)

Common name: **Atlantic white marlin; Espadim-branco-do-Atlântico**, Agulha, Agulhão, Espadarte, Espadarte-branco, Espadim, Espadim-branco, Espadim-pequeno, Marlim-branco, Peco, Peito, Peixe-agulha, Peto.

References: mainland Portugal▲ (Rodriguez-Roda and Howard, 1962; Robins and de Sylva, 1963(?); Osório de Castro, 1967; Nishikawa et al., 1985; Nakamura, 1985, 1986; Fischer et al., 1987; Collette et al., 2011d; Carneiro et al., 2014; Nakamura and Collette, 2016; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Ueyanagi et al., 1970; Fischer et al., 1981; Nishikawa et al., 1985; Nakamura, 1985; Arruda, 1997; Santos et al., 1997; Porteiro et al., 2010; Collette et al., 2011d; Carneiro et al., 2014; Nakamura and Collette, 2016; Froese and Pauly, 2019) and Madeira▲ (Osório de Castro, 1967; Fischer et al., 1981; Nakamura, 1985, 1986; Collette et al., 2011d; Carneiro et al., 2014; Nakamura and Collette, 2016; Froese and Pauly, 2019).

Makaira nigricans Lacepède 1802

Common name: **Blue marlin; Espadim-azul-do-Atlântico**, Agulha, Agulhão, Agulha-sombra, Espadarte, Espadarte-sombra, Espadim, Espadim-azul, Peco, Peito, Peixe-agulha, Peto.

References: mainland Portugal[▲] ([Gonçalves, 1941\(?\)](#); [Rodriguez-Roda and Howard, 1962\(?\)](#); [Osório de Castro, 1954, 1967\(?\)](#); [Fischer et al., 1981](#); [Nishikawa et al., 1985](#); [Nakamura, 1985, 1986](#); [Collette et al., 2011e](#); [Santos et al., 2002a](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Osório de Castro, 1967](#); [Ueyanagi et al., 1970](#); [Fischer et al., 1981](#); [Nishikawa et al., 1985](#); [Nakamura, 1985, 1986](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017\(?\)](#); [Collette et al., 2011e](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Osório de Castro, 1954, 1967](#); [Fischer et al., 1981](#); [Nakamura, 1985, 1986](#); [Severino, 2004](#); [Collette et al., 2011e](#); [Carneiro et al., 2014](#); [Nakamura and Collette, 2016](#); [Froese and Pauly, 2019](#)).

Tetrapturus belone Rafinesque 1810

Common name: **Mediterranean spearfish; Espadim-do-Mediterrâneo**, Águia, Espadim, Espadim-água, Merlim-água, Merlim-do-Mediterrâneo, Peco, Peito, Peto, Pêto, Veleiro.

References: mainland Portugal (?) ([Gonçalves, 1941\(?\)](#); [Albuquerque, 1954-1956\(?\)](#)), Azores (?) ([Osório de Castro, 1967\(?\)](#)) and Madeira ([de Velloso, 1892](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967\(?\)](#); [IGFA 1823-15809, 1980](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015f](#)). Gonçalves (1941) reported the occurrence of one specimen of the genus *Tetrapturus* on the southern coast of Portugal without identifying the species; Albuquerque (1954-1956) included this record in the description of the *T. belone* without indicating the occurrence of this species in mainland Portugal, and probably this record could be misidentifications of a similar species occurring in the area (*Tetrapturus* or *Makaira* species). Noronha and Sarmento (1948) reported the occurrence of *T. belone* in Madeira, Albuquerque (1956) also reported it as a rare species in Madeira and Osório de Castro (1967) as a doubtful occurrence in the Azores and Madeira. Collette and Heessen (2015) mention that there is an International Game Fish Association record from Madeira (IGFA 1823-15809, June 2, 1980), outside the Mediterranean Sea (B. Collette pers. comm. 2013); identification, based on a photograph, is uncertain, but an expert opinion (B. Collette, J. Graves, J. Schratwieser pers. comm. 2013) consider that the record is likely to refer to *Tetrapturus belone*, representing a possible range extension for this species. However, according to Froese and Pauly (2019) and Eschmeyer et al. (2019) this species is endemic in the Mediterranean Sea. Thus, the occurrence of this species in the southern coast of Portugal needs further confirmation and in the Azores is questionable. This species is not considered in the total number of valid species for mainland Portugal and the Azores.

Tetrapturus georgii Lowe 1841

Common name: **Roundscale spearfish; Espadim-peto**, Marlim-peto, Peito.

References: mainland Portugal ([KUI 23773, 1961](#); [Hureau and Monod, 1979](#); [Fischer et al., 1981, 1987](#); [Nakamura, 1985, 1986](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015g](#); [Nakamura and Collette, 2016](#); [Froese and Pauly, 2019](#)), Azores ([Porteiro et al., 2010](#); [Bernarda et al., 2013 \(PECS\)](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015g](#); [Nakamura and Collette, 2016](#)) and Madeira ([Lowe, 1840a, 1841, 1849](#); [Fowler, 1936b](#); [Hureau and Monod, 1979](#); [Fischer et al., 1981](#); [Nakamura, 1985, 1986](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Collette and Heessen, 2015g](#); [Nakamura and Collette, 2016](#); [Froese and Pauly, 2019](#)). This species can be confused with *Kajikia albida* (Poey 1860) (Nakamura and Collette, 2016). The holotype (unique) of this species was caught off in Madeira and its whereabouts are unknown.

Tetrapturus pfluegeri Robins & de Sylva 1963

Common name: **Longbill spearfish; Espadim-bicudo**, Marlim-bicudo.

References: mainland Portugal ([Nishikawa et al., 1985](#); [Nakamura, 1985, 1986](#); [Collette et al., 2011f](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Ueyanagi et al., 1970](#); [Fischer et al., 1981](#); [Nishikawa et al., 1985](#); [Nakamura, 1985](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Borges et al., 2010](#); [Porteiro et al., 2010](#); [Collette et al., 2011f](#); [Carneiro et al., 2014](#)) and Madeira ([Nakamura, 1985, 1986](#); [Lloris et al., 1991](#); [Collette et al., 2011f](#); [Nakamura and Collette, 2016](#); [Froese and Pauly, 2019](#)).

Order Pleuronectiformes

Family Citharidae

***Citharus linguatula* (Linnaeus 1758)**

Common name: **Spotted flounder; Carta-de-bico**, Areeiro, Avieiro, Carta.

References: mainland Portugal▲ (Vandelli, 1771; MNHN-IC-1886-0039, 1883; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández et al., 1981; INIP, 1981b; Martins, 1981; Dinis and Marecos, 1984; Dinis, 1985; Nielsen, 1986c; Fischer et al., 1987; Martins et al., 1992, 2015; Erzini et al., 2001a; Monteiro et al., 2001; Borges et al., 2002, 2003; Santos et al., 2002a; Mendes et al., 2004; Moura, 2004; ICES, 2005; Stergiou et al., 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Silva, 2008, 2010, 2011; Batista et al., 2009; Silva et al., 2009; Teixeira and Cabral, 2009; Teixeira et al., 2010; Costa et al., 2012; Chaves and Silva, 2013; Costa, 2014; Carneiro et al., 2014; Silva and Borges, 2014; de Sola et al., 2015a; Munroe, 2016b; Martins and Carneiro, 2018; Baptista et al., 2019) and Madeira (MNHN-IC-2005-2372, n.d.; Carneiro et al., 2014; de Sola et al., 2015a; Munroe, 2016b). The distribution maps of this species reported by de Sola et al. (2015a) and Munroe (2016b) partially included the area of Selvagens Islands (Madeira EEZ).

Family Scophthalmidae

***Lepidorhombus boscii* (Risso 1810)**

Common name: **Four-spot megrim; Areeiro-de-quatro-manchas**, Areeiro, Carta, Solha.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; Vieira, 1897, 1900; Bragança, 1902, 1903; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Nielsen, 1979j, 1986c, 1990c; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a, b, 1982a, b; Martins, 1981; Dinis and Marecos, 1984; Dinis, 1985; Gordo et al., 1990; Martins et al., 1992; Figueiredo et al., 1994; Gomes et al., 2001; Monteiro et al., 2001; Borges et al., 2002, 2003; Mendes et al., 2004; Moura, 2004; ICES, 2005; Sousa et al., 2005, 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Pakhorukov, 2008; Silva, 2008, 2010, 2011; Abecasis et al., 2009; Batista et al., 2009; Silva et al., 2009; Teixeira et al., 2010; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Silva and Borges, 2014; Munroe and Chanet, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (PECS) (MNHN-IC-1956-0007, 1955; MNHN-IC-1956-0008, 1955; Pakhorukov, 2008 (PECS); Carneiro et al., 2014) and Madeira (PECS) (Shcherbachov et al., 1985a; Pakhorukov, 2008; Carneiro et al., 2014). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22799, 1967). This species can be confused with *Lepidorhombus whiffiagonis* (Walbaum 1792) (Munroe and Chanet, 2016).

***Lepidorhombus whiffiagonis* (Walbaum 1792)**

Common name: **Megrim; Areeiro**, Pregado, Solha, Solha-da-fundura, Solha-grande.

References: mainland Portugal▲ (Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nielsen, 1979j, 1986c; INIP, 1981b, 1982a, b; Fernández et al., 1981; Martins, 1981; Dinis and Marecos, 1984; Dinis, 1985; Fischer et al., 1987; Gordo et al., 1990; Figueiredo et al., 1994; Borges et al., 2002; Moura, 2004; ICES, 2005; Fernandes et al., 2007; Henriques et al., 2008; Pakhorukov, 2008; Silva, 2008, 2011; Chaves, 2009; Silva et al., 2009, 2010; Teixeira et al., 2010; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; OCEANA, 2014; Silva and Borges, 2014; Munroe and Chanet, 2016; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores▲ (Cligny, n.d.; MNHN-IC-1886-0037, 1883; Vaillant, 1888, 1919; Goode and Bean, 1895, 1896; Brauer, 1906; Norman, 1934; Fowler, 1936; Chabanaud, 1939; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Nash et al., 1991; Arruda, 1992, 1997; Silva et al., 1995; Menezes, 1996; Santos et al., 1997; Sousa et al., 1999; Menezes et al., 2006, 2009, 2012; Pakhorukov, 2008 (PECS); Porteiro et al., 2010, 2013; Carneiro et al., 2014; Almeida and Biscoito, 2019) and Madeira (Osório de Castro, 1967; Maul, 1976; Pakhorukov, 2008 (PECS)). This species can be confused with *Lepidorhombus boscii* (Risso 1810) (Munroe and Chanet, 2016).

Scophthalmus maximus (Linnaeus 1758)

Common name: **Turbot; Pregado**, Parracho, Parrocho, Redovalho, Renovalho, Rodavalho, Rodovalho, Rodovalho-pregado, Solha.

References: mainland Portugal▲ ([de Brito Capello, 1867a, 1868, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1893, 1897, 1900](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Osório, 1896](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Muus and Dahlström, 1978](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, b, 1982b](#); [Borges, 1984](#); [Dinis, 1985](#); [Nielsen, 1986c](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Erzini et al., 2001a](#); [Jorge et al., 2002](#); [Santos et al., 2002c](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Ribeiro et al., 2006](#); [Chaves, 2008](#); [Henriques et al., 2008](#); [Batista et al., 2009](#); [Garrido et al., 2009](#); [Carneiro et al., 2014](#); [Munroe and Chanet, 2016](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-2005-2424, 1926](#); [Carneiro et al., 2014](#)).

Scophthalmus rhombus (Linnaeus 1758)

Common name: **Brill; Rodovalho**, Clerigo, Clérigo, Parracho, Parrocho, Patrúcia, Pegadeira, Redovalho, Rodovalho, Rodovalho-legítimo, Renovalho, Solha-do-rio, Tapa-potes.

References: mainland Portugal▲ ([Carvalho, 1756](#); [Vandelli, 1771](#); [de Brito Capello, 1867a, 1868, 1869b, 1880](#); [Steindachner, 1868b](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1895a, 1935](#); [Vieira, 1897, 1901](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Nielsen, 1979j, 1986c](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1999, 2015](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Ribeiro et al., 2006](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Munroe and Chanet, 2016](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Zeugopterus punctatus (Bloch 1787)

Common name: **Topknot; Rodovalho-bruxa**, Rodovalho.

References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Vieira, 1897, 1900](#); [Nobre, 1903, 1904, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Antunes et al., 1988](#); [Almeida, 1996](#); [Henriques et al., 2007, 2008](#); [Batista et al., 2009](#); [Garrido et al., 2009](#); [Carneiro et al., 2014](#)).

Zeugopterus regius (Bonnaterre 1788)

Common name: Eckström's topknot; **Bruxa**.

References: mainland Portugal ([Bragança, 1903\(?\)](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Nielsen, 1979j, 1986c](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Fischer et al., 1987](#); [Martins et al., 1992](#); [Henriques et al., 2007, 2008](#); [Carneiro et al., 2014](#); [Munroe and Chanet, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Family Paralichthyidae

Syacium papillosum (Linnaeus 1758)

Common name: **Dusky flounder; Carta-sombria**.

References: Azores (PECS) ([ROM 23911, 1966](#); [Carneiro et al., 2014](#)). This species can be confused with *Syacium micrurum* (Ranzani 1842) ([Munroe, 2016c](#)). The occurrence record of the Royal Ontario Museum mentions the catch of the specimen off Florida (Atlantic Ocean). However, the geographic coordinates displayed correspond to a location off the Azores, in the PECS area (82 NM west of the Great Meteor Tablemount). According to Eschmeyer et al. (2019) the current geographic distribution of *S. papillosum* is the Western and Eastern Atlantic. The record from Azores Island requires verification.

Syacium micrurum Ranzani 1842

Common name: **Channel flounder; Carta-moira**, Carta, Linguado, Linguado-moiro, Linguado-mouro.

References: Azores (?) ([Fowler, 1936a](#); [Collins, 1954](#); [Osório de Castro, 1967](#)) and Madeira (?) ([Osório de Castro, 1967](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *S. micrurum* is

the Western Atlantic. Arruda (1997) and Santos *et al.* (1997) mention that the original record of the species *S. micrurum* by Lampe (1914) was in Porto Grande, São Vicente, Cape Verde Archipelago. The further citations of this species for the Azores by Fowler (1936a) and Collins (1954) should have been based on a partial mislabelling of the original material (Arruda, 1997; Santos *et al.*, 1997). Osório de Castro (1967) reported the occurrence of this species in the fishing grounds of the Azores and Madeira, without indicating the geographic location. The presence of this species in the Azores and Madeira Islands are probably based on misidentification of *Syacium guineensis* (Bleeker 1862) or mistaken records. Thus, this species is not considered in the total number of valid species for those islands.

Family Bothidae

Arnoglossus grohmanni (Bonaparte 1837)

Common name: No English common name; [Carta-de-Grohmann](#), Areeiro.

References: mainland Portugal (?) ([Nobre, 1903, 1935](#); [Seabra, 1911](#)). Nobre reported the occurrence of *Pleuronectes grohmanni* Bonaparte 1837 in Setúbal (mainland Portugal) in 1903 and considered it a very rare species. Some authors considered *P. grohmanni* in part in the synonymy of *Arnoglossus thori* Kyle 1913. The current geographic distribution of *A. grohmanni* is in the Mediterranean Sea and Sea of Marmara (endemic) (Eschmeyer, 2019). The occurrence of this species was not reported in mainland Portugal since Nobre (1935) and its presence in this EEZ was not yet confirmed by more recent data. Thus, this species is not considered in the total number of valid species.

Arnoglossus imperialis (Rafinesque 1810)

Common name: [Imperial sculpfish](#); [Carta-imperial](#), Areeiro.

References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Nielsen, 1979j](#); [Fernández et al., 1981](#); [Dinis 1981](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Nielsen, 1986c](#); [Fischer et al., 1987](#); [Erzini et al., 2001a](#); [Borges et al., 2002, 2003](#); [Cabral et al., 2002](#); [Prista et al., 2003](#); [Moura, 2004](#); [ICES, 2005](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Pakhorukov, 2008](#); [Abecasis et al., 2009](#); [Batista et al., 2009](#); [Borges et al., 2009](#); [Silva, 2010, 2011](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [de Sola et al., 2015b](#); [Munroe, 2016c](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1896a](#); [Kyle, 1913](#); [Norman, 1930](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Nielsen, 1986c](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Kukuev, 2002 \(PECS\)](#); [Floeter et al., 2008](#); [Pakhorukov, 2008 \(PECS\)](#); [Carneiro et al., 2014](#); [de Sola et al., 2015b](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Norman, 1934](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976 \(PECS\)](#); [Fischer et al., 1981](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Pakhorukov, 2008 \(PECS\)](#); [Carneiro et al., 2014](#); [de Sola et al., 2015b](#); [Munroe, 2016c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). Two specimens caught off in different areas are in the MMF collections: one at the coast of mainland Portugal (MMF 22830, 1967) and the other at the Josephine Bank (MMF 22805, 1967).

Arnoglossus laterna (Walbaum 1792)

Common name: [Mediterranean sculpfish](#); [Carta-do-Mediterrâneo](#), Areeiro, Carta.

References: mainland Portugal▲ ([de Brito Capello, 1868](#); [Nobre, 1903, 1904, 1935](#); [Seabra, 1911](#); [Osório de Castro, 1954, 1967](#); [Nielsen, 1979j](#); [INIP, 1982b](#); [Fernández et al., 1981](#); [Dinis 1981](#); [Martins, 1981](#); [Dinis and Marecos, 1984](#); [Nielsen, 1986c](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Martins et al., 1992, 2015](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Prista et al., 2003](#); [ICES, 2005](#); [Ribeiro et al., 2006, 2008](#); [Sousa et al., 2006](#); [Fernandes et al., 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Garrido et al., 2009](#); [Silva et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Paulo-Martins et al., 2011](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [de Sola et al., 2015c](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([de Sola et al., 2015c](#); [Munroe, 2016c](#)).

***Arnoglossus rueppellii* (Cocco 1844)**

Common name: Rüppell's scaldback; **Carta-estreita**, Areeiro.

References: mainland Portugal ([Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Martins, 1981](#); [INIP, 1982a](#); [Borges et al., 2002](#); [ICES, 2005](#); [Pakhorukov, 2008](#); [Ribeiro et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [OCEANA, 2014](#); [Martins and Carneiro, 2018](#)), Azores ([Kyle, 1913](#); [Bertin, 1932, 1936](#); [Chabanaud, 1932](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977 \(PECS\)](#); [Arruda, 1997](#); [Fock et al., 2002 \(PECS\)](#); [Kukuev, 2002 \(PECS\)](#); [IMAR/DOP 1AÇO-04-P03SOL1, 2003](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [de Sola et al., 2015d](#)) and Madeira ([Osório de Castro, 1967\(?\)](#); [Maul, 1976 \(PECS\)](#); [Pakhorukov, 2008 \(PECS\)](#); [Christiansen et al., 2009](#); [Carneiro et al., 2014](#); [de Sola et al., 2015d](#); [Eschmeyer et al., 2019](#)). Several specimens caught off in different areas are in the MMF collections: eleven specimens at the Josephine Bank (MMF 22804, 1967, set of five specimens and MMF 22850, 1967, set of six specimens) and forty six specimens at the Meteor Bank (MMF 223364, 1967, set of seven specimens, MMF 22366, 1967, set of three specimens, MMF 22406, 1967, set of six specimens, MMF 22846, 1967, set of four specimens, MMF 22854, 1967, one specimen, MMF 22871, 1967, set of seventeen specimens and MMF 22882, 1967, set of eight specimens).

***Arnoglossus thori* Kyle 1913**

Common name: **Thor's scaldfish**; **Carta-pontuada**, Areeiro.

References: mainland Portugal ([MNHN-IC-1886-0011, 1883](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Martins, 1981](#); [Nielsen, 1979j, 1986c](#); [Bauchot, 1987](#); [Fischer et al., 1987](#); [Borges et al., 2002](#); [Santos et al., 2002a](#); [Palma et al., 2003](#); [ICES, 2005](#); [Ribeiro et al., 2006, 2012](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Neves et al., 2008](#); [Silva, 2008, 2011](#); [Batista et al., 2009](#); [Chaves, 2009](#); [Gonçalves et al., 2010](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Munroe, 2016c](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-1883-0124, 1882](#); [MNHN-IC-1883-0125, 1882](#); [Carneiro et al., 2014](#); [de Sola et al., 2015e](#)).

***Bothus podas* (Delaroche 1809)**

Common name: **Wide-eyed flounder**; **Carta-de-olhos-grandes**, Careta, Carta, Carteta, Sola, Solha, Tapacu.

References: mainland Portugal[▲] ([Vieira, 1897, 1901](#); [Nobre, 1903, 1937](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Nielsen, 1986c](#); [Fischer et al., 1987](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [Palma et al., 2003](#); [Ribeiro et al., 2006, 2012](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Batista et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [de Sola et al., 2015f](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Hilgendorf, 1888a](#); [Collett, 1896a](#); [Kyle, 1913](#); [Lampe, 1914](#); [Vaillant, 1919](#); [Nobre, 1924, 1930](#); [Zugmayer, 1933](#); [Norman, 1934](#); [Fowler, 1936a, b](#); [Cadenat, 1937](#); [Chabanaud, 1939](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967\(?\)](#); [Wood and Williams, 1974](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Barreiros, 1990](#); [Nash et al., 1990, 1991, 1994a, b](#); [Patzner and Santos, 1990a, e](#); [Patzner, 1990](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1994b, 1997](#); [Wirtz, 1990a, 1994](#); [Azevedo et al., 1991, 1992](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Gonçalves et al., 1995](#); [Santos and Nash, 1995](#); [Morato et al., 2001b](#); [Menezes et al., 2006](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010, 2017](#); [Diogo and Pereira, 2013, 2014](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Munroe, 2016c](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1833a, 1837, 1838](#); [Günther, 1862](#); [de Velloso, 1892](#); [Collett, 1896](#); [Vaillant, 1919](#); [Fowler, 1923, 1936b](#); [Norman, 1934](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#); [Fischer et al., 1981](#); [Augier, 1985](#); [Dooley et al., 1985](#); [Nielsen, 1986c](#); [Lloris et al., 1991](#); [Abreu et al., 1995](#); [Andrade and Albuquerque, 1995](#); [Araújo et al., 2005](#); [Wirtz et al., 2008](#); [Falcón et al., 2000](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [de Sola et al., 2015f](#); [Munroe, 2016c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The syntypes of *Rhombus maderensis* Lowe 1834, synonym of *B. podas*, were caught off in Madeira and its whereabouts are unknown. This species can be confused with *Bothus mellissi* Norman 1931 (Munroe, 2016c).

Chascanopsetta lugubris Alcock 1894

Common name: **Pelican flounder; Carta-pelícano.**

References: mainland Portugal ([USNM 282746, 1964](#); [USNM 282748, 1964](#); [Carneiro et al., 2014](#)). The occurrence records of the National Museum of Natural History, Smithsonian Institution, are the most northerly in the Eastern Atlantic. Since these occurrences, there is no other evidence of the presence of this species in mainland Portugal. Eschmeyer *et al.* (2019) mention it as a circumglobal species in tropical and warm temperate seas. The records from mainland Portugal require verification.

Monolene microstoma Cadenat 1937

Common name: Smallmouth moonflounder; **Carta-de-boca-pequena, Carta.**

References: Madeira (PECS) ([SOC Discovery 785703_210_FIS_503001, 1972](#); [Carneiro et al., 2014](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *M. microstoma* in the Eastern Atlantic is from Senegal to Namibia. However, the northernmost occurrence records of this species in the Eastern Atlantic is a human observation of a specimen caught off at the Josephine Bank, 235 NM NNO of Porto Santo Island (Madeira) but its presence in the Portuguese ichthyofauna is not yet documented. Specimens are needed for its confirmation in Madeira.

Family Pleuronectidae

Glyptocephalus cynoglossus (Linnaeus 1758)

Common name: **Witch flounder; Solhão, Cinoglosso, Língua.**

References: mainland Portugal ([Ryland and Lancaster, 2017](#)).

Microstomus kitt (Walbaum 1792)

Common name: **Lemon sole; Solha-limão, Solha-microcéfala.**

References: mainland Portugal (?) ([Ramos, 2010](#); [Carneiro et al., 2014](#); [WoRMS Editorial Board, 2018](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *M. kitt* is the Western Baltic Sea, North Sea, eastern-north Atlantic, South-east coast of Greenland. Bañón *et al.* (2010) indicate the occurrence of this species in Galician waters. Probably the southern limit of distribution of this species is the offshore on sandy or shelly grounds of the northern Portuguese coast. Its presence among the Portuguese ichthyofauna is not yet documented. Specimens are needed for confirmation in mainland Portugal. Thus, this species is not considered in the total number of valid species.

Platichthys flesus (Linnaeus 1758)

Common name: **European flounder; Solha-das-pedras, Azevia, Dorminhoca, Patença, Patruça, Patruçadas-pedras, Patruça-de-pedras, Patrúcia, Patrussa, Patuça, Patusca, Petisca, Solha, Solha-de-pedras, Solha-marinha, Solhão, Solho.**

References: mainland Portugal[▲] ([Vandelli, 1771](#); [de Brito Capello, 1867a, 1868, 1880](#); [Vieira, 1893, 1894a, 1900](#); [Osório, 1896](#); [Nobre, 1903, 1931, 1932, 1935](#); [Seabra, 1911](#); [Nobre et al., 1915](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Nielsen, 1979j, 1986c](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Cabral, 1999](#); [Jorge et al., 2002](#); [Thiel et al., 2003](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Cabral et al., 2007](#); [Chaves, 2008](#); [Henriques et al., 2008](#); [Garrido et al., 2009](#); [Gonçalves et al., 2010](#); [Teixeira et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Munroe, 2016b](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Mecklenburg et al., 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)) and Azores (?) ([Osório de Castro, 1967](#)). Osório de Castro (1967) reported the occurrence of this species in the fishing grounds of the Azores, without indicating the geographic location. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores. Specimens are needed for confirmation in the Azores. Thus, this species is not considered in the total number of Azorean species.

Pleuronectes platessa Linnaeus 1758

Common name: **European plaice; Solha, Solha-avessa, Solha-legítima, Solhão.**

References: mainland Portugal[▲] ([de Brito Capello, 1868, 1880](#); [Baldaque da Silva, 1891](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Nielsen, 1979j, 1986c](#); [Muus and Dahlström, 1978](#); [Fernández et al., 1981](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Fischer et al., 1987](#); [Martins et al., 1992, 1999, 2015](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Henriques et al., 2008](#); [Carneiro et](#)

al., 2014; Freyhof, 2015; Munroe, 2016b; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Mecklenburg *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Azores (Sampaio, 1904; Ribeiro, 1936; Collins, 1954; Arruda, 1997; Diveboard 90772_208483_0, 2013; Carneiro *et al.*, 2014). According to Santos *et al.* (1997) this species do not occur in the Azores. However, in 2013 by human observation (Scuba diving) the presence of this species in the Pico, Faial and São Jorge islands was checked and relaunched the doubt about the occurrence in the Azores. Specimens are needed for its confirmation.

Family Soleidae

***Bathysolea profundicola* (Vaillant 1888)**

Common name: **Deep water sole; Linguado-da-fundura**, Linguado.

References: mainland Portugal (MNHN-IC-1883-0127, 1882; MNHN-IC-1883-0128, 1882; Vaillant, 1888; Goode and Bean, 1895, 1896; Brauer, 1906; Seabra, 1911; Nobre, 1935; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Torchio, 1979; Quéro *et al.*, 1986; Fischer *et al.*, 1987; Saldanha *et al.*, 1995; Desoutter and Chapleau, 1997; ICES, 2005; Chaves, 2008, 2009; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Carneiro *et al.*, 2014; Silva and Borges, 2014; Herrera *et al.*, 2015; Bañón *et al.*, 2016a; Desoutter-Meniger and Munroe, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967) and Madeira (Osório de Castro, 1967; Herrera *et al.*, 2015; Desoutter-Meniger and Munroe, 2016). This species can be confused with *Bathysolea polli* (Chabanaud 1950) (Desoutter-Meniger and Munroe, 2016). The holotype and one paratype of *Solea profundicola* Vaillant 1888, synonym of *B. profundicola*, both caught off in mainland Portugal are in the MNHN collections (MNHN-IC-1883-0128 and MNHN-IC-1883-0127, respectively). Osório de Castro (1967) reported the occurrence of this species in the fishing grounds of the Azores, without indicating the geographic location. Since Osório de Castro (1967) no other author described the presence of this species in the Azores. Moreover, the current geographic range of this benthic species is in the Eastern Atlantic continental slope. Thus, this species is not considered in the total number of valid species to the Azores.

***Buglossidium luteum* (Risso 1810)**

Common name: **Solenette; Língua-de-gato**, Língua, Linguado.

References: mainland Portugal (Bragança, 1902; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Torchio, 1979; Fernández *et al.*, 1981; INIP, 1982a; Dinis, 1985; Quéro *et al.*, 1986; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Gordo *et al.*, 1990; Martins *et al.*, 1992; Erzini *et al.*, 1997; Cabral, 1999; Borges *et al.*, 2002; Prista *et al.*, 2003; ICES, 2005; Gomes and Olim, 2007; Henriques *et al.*, 2007; Henriques *et al.*, 2008; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; de Sola *et al.*, 2015g; Desoutter-Meniger and Munroe, 2016; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (PECS) (MNHN-IC-1959-0604, 1959; Carneiro *et al.*, 2014). This species can be confused with *Microchirus boscanion* (Chabanaud 1926) (Desoutter-Meniger and Munroe, 2016).

***Dagetichthys lusitanicus* (de Brito Capello 1868)**

Common name: **Portuguese sole; Língua-de-vaca**, Azevia, Azevia-da-costa, Língua, Língua-da-costa, Linguado, Linguado-português.

References: mainland Portugal (NHMUK 1867.7.23.9, 1864-1877; de Brito Capello, 1868, 1880; MNHN-IC-0000-4804, 1868; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1903, 1935; Bragança, 1903; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Torchio, 1979; Fischer *et al.*, 1981, 1987; Cabral, 1999; Palma *et al.*, 2003; Ribeiro *et al.*, 2006; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Gonçalves *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Desoutter-Meniger and Munroe, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019). Two syntypes of *Synaptura lusitanica* de Brito Capello 1868 both caught off near of Lisbon and Setúbal (mainland Portugal) are in the NHMUK and MNHN collections (NHMUK 1867.7.23.9 and MNHN-IC-0000-4804, respectively). Six others syntypes preserved in the Museu Bocage were destroyed during the 1978 fire. *Synaptura lusitanica* is synonym of *D. lusitanicus*.

***Dicologlossa cuneata* (Moreau 1881)**

Common name: **Wedge sole; Língua, Asvião, Língua, Linguada, Linguado.**

References: mainland Portugal▲ ([Carvalho, 1765](#); [Bragança, 1902, 1903](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Torchio, 1979](#); [INIP, 1981b](#); [Martins, 1981](#); [Fischer et al., 1981, 1987](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Quéro et al., 1986](#); [Martins et al., 1992, 2015](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Monteiro et al., 2001](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Santos et al., 2002a](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Thiel et al., 2003](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Gomes and Olim, 2007](#); [Chaves, 2008](#); [Henriques et al., 2008](#); [Ribeiro et al., 2008](#); [Silva, 2008, 2011](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Monroe et al., 2015a](#); [Desoutter-Meniger and Munroe, 2016](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-1988-0651, 1925](#); [MNHN-IC-1988-1175, 1953](#); [Carneiro et al., 2014](#)).

***Microchirus azevia* (de Brito Capello 1867)**

Common name: Bastard sole; **Azevia**, Cabo-velho, Linguado, Malacueco.

References: mainland Portugal▲ ([de Brito Capello, 1867e, 1868, 1869b, 1880](#); [MNHN-IC-0000-4805, 1867](#); [Baldaque da Silva, 1891](#); [Girard, 1892](#); [Nobre, 1894, 1903, 1935](#); [Vieira, 1898, 1900](#); [Bragança, 1902, 1903](#); [Seabra, 1911](#); [Aboim, 1941](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Torchio, 1979](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Quéro et al., 1986](#); [Martins et al., 1992, 1999, 2015](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [Santos et al., 2002a, c](#); [Moura, 2004](#); [Afonso-Dias et al., 2005](#); [ICES, 2005](#); [Ribeiro et al., 2006](#); [Stergiou et al., 2006](#); [Henriques et al., 2008](#); [Ribeiro et al., 2008](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [Teixeira et al., 2010](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Keskin et al., 2015a](#); [Desoutter-Meniger and Munroe, 2016](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Keskin et al., 2015a](#); [Desoutter-Meniger and Munroe 2016](#)). Two syntypes of *Solea azevia* de Brito Capello 1867 are in the MNHN and ZMB collections (MNHN-IC-0000-4805 and ZMB 6456), a third syntype was preserved in the Museu Bocage and destroyed in the 1978 fire, all caught off near Lisbon (mainland Portugal). *Solea azevia* is synonym of *M. azevia*. According to Eschmeyer et al. (2019) the current geographic distribution of *M. azevia* is the Mediterranean Sea, Sea of Marmara and, in the Eastern Atlantic, from Portugal to Senegal; strays to British Isles. The southernmost occurrence in the Eastern Atlantic is a preserved specimen deposited in the collections of the Muséum national d'Histoire naturelle (MNHN-IC-1988-0631), caught off on July, 1979 and there are also occurrence records of this species in the Canary Islands (MNHN-IC-2006-1129). The distribution maps of this species reported by Keskin et al. (2015a) and Desoutter-Meniger and Munroe (2016) partially included the area of Selvagens Islands (Madeira EEZ). It is possible the occurrence of this species in the Selvagens Islands and specimens are needed for confirmation.

***Microchirus boscanion* (Chabanaud 1926)**

Common name: **Lusitanian sole**; **Azevia-lusitânica**, Azevia-marginada, Cascarra, Linguado.

References: mainland Portugal▲ ([Borges et al., 2002, 2003](#); [Palma et al., 2003](#); [ICES, 2005](#); [Ribeiro et al., 2006](#); [Gomes and Olim, 2007](#); [Henriques et al., 2008](#); [Chaves, 2009](#); [Silva et al., 2009](#); [Silva, 2010](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Martins and Carneiro, 2018](#)) and Madeira ([MNHN-IC-1987-1415, 1983](#); [Carneiro et al., 2014](#); [Munroe et al., 2015b](#); [Desoutter-Meniger and Munroe, 2016](#); [Baptista et al., 2019](#)). This species can be confused with *Buglossidium luteum* (Risso 1810) (Desoutter-Meniger and Munroe, 2016). The distribution maps of this species reported by Munroe et al. (2015b) and Desoutter-Meniger and Munroe (2016) included the occurrence of this species at 45 NM WSW of the Susan Seamount (Madeira EEZ).

***Microchirus hexophthalmus* (Bennett 1831)**

Common name: **Ocellated wedge sole**; **Azevia-de-olhos**, Linguado, Linguado-das-manchas, Linguado-de-manchas, Linguado-de-olhos, Linguado-moiro, Linguado-mouro.

References: mainland Portugal ([Torchio, 1979](#); [Dooley et al., 1985](#); [Quéro et al., 1986](#); [Fischer et al., 1987](#); [Cabral, 1999](#); [Martins et al., 1999](#); [Borges et al., 2002, 2003](#); [Jorge et al., 2002](#); [Sousa et al., 2006](#); [Neves et al., 2008](#); [Carneiro et al., 2014](#); [Desoutter-Meniger and Munroe, 2016](#); [Martins and Carneiro, 2018](#); [Froese and Pauly, 2019](#)) and Madeira ([Torchio, 1979](#); [Dooley et al., 1985](#); [Quéro et al., 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Munroe, 2015](#); [Froese and Pauly, 2019](#)).

***Microchirus ocellatus* (Linnaeus 1758)**

Common name: **Foureyed sole; Azevia-de-malhas**, Azevia, Linguado, Solha, Sôlha-linguado.

References: mainland Portugal (de Brito Capello, 1868, 1880; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Torchio, 1979; Dooley et al., 1985; Quéro et al., 1986; Fischer et al., 1987(?); Martins et al., 1992; Cabral, 1999; Erzini et al., 2001a; Borges et al., 2002; Mendes et al., 2004; ICES, 2005; Henriques et al., 2008; Batista et al., 2009; Carneiro et al., 2014; Costa, 2014; Keskin et al., 2015b; Desoutter-Meniger and Munroe, 2016; Martins and Carneiro, 2018) and Madeira (Günther, 1862; Johnson, 1863b; Noronha and Sarmento, 1934; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Torchio, 1979; Dooley et al., 1985; Quéro et al., 1986; Lloris et al., 1991; Wirtz et al., 2008; Carneiro et al., 2014; Keskin et al., 2015b; Desoutter-Meniger and Munroe, 2016; Froese and Pauly, 2019).

***Microchirus variegatus* (Donovan 1808)**

Common name: **Thickback sole; Azevia-raiada**, Azevia, Cascorra, Linguado, Raposo, Sóia, Solha-tapa, Tapa, Tapa-cu.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; Steindachner, 1868b; Collett, 1896; Bragança, 1902, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Torchio, 1979; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a, 1982a, b; Antunes et al., 1988; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Quéro et al., 1986; Martins et al., 1999, 2015; Erzini et al., 2001a; Gomes et al., 2001; Borges et al., 2002; Thiel et al., 2003; Mendes et al., 2004; ICES, 2005; Sousa et al. 2005, 2006; Fernandes et al., 2007; Chaves, 2008, 2009; Henriques et al., 2008; Silva, 2008, 2010, 2011; Batista et al., 2009; Garrido et al., 2009; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Silva and Borges, 2014; Keskin et al., 2015c; Martins and Carneiro, 2018; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Quéro et al., 1986; Carneiro et al., 2014; Keskin et al., 2015c; Desoutter-Meniger and Munroe, 2016; Froese and Pauly, 2019). This species can be confused with *Microchirus wittei* Chabanaud 1950 (Desoutter-Meniger and Munroe, 2016).

***Microchirus wittei* Chabanaud 1950**

Common name: **Banded sole; Azevia-raiada-africana**, Azevia, Linguado, Linguado-raiado.

References: Madeira (MNHN-IC-1987-1410, 1983; MNHN-IC- 1987-1411, 1983; Carneiro et al., 2014; Ralph, 2015p). This species can be confused with *Microchirus variegatus* (Donovan 1808) (Desoutter-Meniger and Munroe, 2016). According to Eschmeyer et al. (2019) the current geographic distribution of *M. wittei* in the Eastern Atlantic is from Mauritania to Congo. The specimens in the Ichthyology Collection of the Muséum national d'Histoire naturelle represent the extent of the northern limit of distribution of this species.

***Monochirus atlanticus* Chabanaud 1940**

Common name: **Whiskered sole; Cascarra-do-Atlântico**, Cascarra, Linguado.

References: mainland Portugal (Veiga et al., 2009; Monroe et al., 2015b; Desoutter-Meniger and Munroe, 2016; Froese and Pauly, 2019). According to Eschmeyer et al. (2019) the geographic distribution of *M. atlanticus* is the coast of West Africa and Monroe et al. (2015b) widened the distribution from the Cantabrian Sea to the Gabon, but not including the Mediterranean Sea.

***Monochirus hispidus* Rafinesque 1814**

Common name: **Whiskered sole; Cascarra**, Cascorra, Lapa-cu, Linguado, Tapa-cu.

References: mainland Portugal (de Brito Capello, 1868, 1880; Osório, 1888; Nobre, 1895a, c, 1904, 1935; Vieira, 1900; Seabra, 1911; Gonçalves, 1941; Hellings, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Torchio, 1979; Martins, 1981; Quéro et al., 1986; Fischer et al., 1987; Monteiro, 1989; Martins et al., 1992; da Cunha, 1994; Desoutter, 1997; Cabral, 1999, 2000a; Ribeiro et al., 2006, 2008, 2012; Cabral et al., 2007; Henriques et al., 2007, 2008; Chaves, 2008; Neves et al., 2008; Silva, 2008, 2010, 2011; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019). According to Eschmeyer et al. (2019) the geographic distribution of *M. hispidus* is the Mediterranean Sea, Sea of Marmara and, in the Eastern Atlantic, from Portugal south to Ghana. However, Rijnsdorp et al. (2015a) considered this species to be an endemic Mediterranean one. The previous quotations of *M. hispidus* to mainland Portugal can not be clearly

related to one of the two former taxonomic categories even after the elevation of the *atlanticus* subspecies to the species status. Thus, this subject deserves further detailed attention.

Pegusa impar (Bennett 1831)

Common name: **Adriatic sole; Linguado-do-Adriático**.

References: mainland Portugal ([Erzini et al., 2001a](#); Gomes and Olim, 2007; MLFPI 84, 2008(?); MLFPI 85, 2008(?); MLFPI 86, 2008(?); MLFPI 143, 2010(?); MLFPI 144, 2010(?); Carneiro et al., 2014).

Pegusa lascaris (Risso 1810)

Common name: **Sand sole; Linguado-da-areia**, Cascarra, Gafo, Linguado, Linguado-de-areia, Linguado-preto, Linguado-serrano, Macaca, Serranho, Serrano, Tapa-cu.

References: mainland Portugal[▲] ([Gervais and Boulart, 1877a](#); [Osório, 1888, 1905](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Bragança, 1902\(?\)](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Torchio, 1979](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Quéro, 1981](#); [Dinis, 1985](#); [Dooley et al., 1985](#); [Quéro et al., 1986](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Afonso-Dias et al., 2002](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Ribeiro et al., 2006](#); [Henriques et al., 2008](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Keskin et al., 2015d](#); [Desoutter-Meniger and Munroe, 2016](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Carneiro et al., 2014](#)) and Madeira ([Günther, 1862](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Ben-Tuvia, 1990b](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). This species was erroneously considered present in the Azores ([Carneiro et al., 2014](#)). Thus, this species is not included in the total number of valid species in the Azores.

Pegusa nasuta (Pallas 1814)

Common name: **Snouted sole; Linguado-narigudo**.

References: mainland Portugal ([Borges et al., 2002](#)). According to Eschmeyer et al. (2019) and Froese and Pauly (2019) the current geographic distribution of this species is the Northern Mediterranean Sea, Sea of Marmara, Black Sea and Sea of Azov. Borges et al. (2002) reported the occurrence of this species in the southern coast of Portugal (Algarve) and identified it as *Solea nasuta* (Pallas 1814).

Solea senegalensis Kaup 1858

Common name: **Senegalese sole; Linguado-do-Senegal**, Linguado, Linguado-branco, Linguado-da-costa, Linguado-de-fóra, Peixe-banda.

References: mainland Portugal[▲] ([Bragança, 1903](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Torchio, 1979](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Dinis, 1985](#); [Quéro et al., 1986](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1992, 2015](#); [Erzini et al., 1996, 2001a](#); [Gonçalves et al., 1996](#); [Cabral, 1999, 2000b](#); [Santos et al., 2002a](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Cabral et al., 2007](#); [Ribeiro et al., 2006, 2008](#); [Stergiou et al., 2006](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008](#); [Neves et al., 2008](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Silva, 2011](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Keskin et al., 2015e](#); [Desoutter-Meniger and Munroe, 2016](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-1999-0052, 1959](#); [Carneiro et al., 2014](#); [Keskin et al., 2015e](#)). The distribution maps of this species reported by Keskin et al. (2015e) partially included the area of Selvagens Islands (Madeira EEZ).

Solea solea (Linnaeus 1758)

Common name: **Common sole; Linguado-legítimo**, Lingoad, Linguado, Linguado-imperial, Linguado-real, Linguado-vulgar, Pescado-real.

References: mainland Portugal[▲] ([Vandelli, 1771](#); [de Brito Capello, 1867e, 1868, 1869b, 1880](#); [Steindachner, 1868b](#); [Baldaque da Silva, 1891](#); [Vieira, 1893, 1897, 1900](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Osório, 1895](#); [Bragança, 1902, 1903](#); [Seabra, 1911](#); [Murray and Hjort, 1912](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Muus and Dahlström 1978](#); [Torchio, 1979](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#);

Borges, 1984; Dinis, 1985; Dooley *et al.*, 1985; Quéro *et al.*, 1986; Arruda *et al.*, 1988; Martins *et al.*, 1992, 1999, 2015; Cabral, 1999, 2000b; Erzini *et al.*, 2001a; Borges *et al.*, 2002; Jorge *et al.*, 2002; Santos *et al.*, 2002a; Palma *et al.*, 2003; Prista *et al.*, 2003; Garnerot *et al.*, 2004; Mendes *et al.*, 2004; ICES, 2005; Stergiou *et al.*, 2006; Cabral *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.* 2007, 2008; Chaves, 2008, 2009; Neves *et al.*, 2008; Batista *et al.*, 2009; Garrido *et al.*, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Monroe *et al.*, 2015c; Desoutter-Meniger and Munroe, 2016; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Sampaio, 1904; Collins, 1954; Osório de Castro, 1967; Arruda, 1997) and Madeira (Osório de Castro, 1967; Ben-Tuvia, 1990b; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Monroe *et al.*, 2015c; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). According to Santos *et al.* (1997) this species do not occur in the Azores. Thus, this species is not considered in the total number of valid species to the Azores.

***Synapturichthys kleinii* (Risso 1827)**

Common name: Klein's sole; **Linguado-turco**, Linguado, Linguado-revesso, Zebra.

References: mainland Portugal (de Brito Capello, 1880; Seabra, 1911; Nobre, 1935; Osório de Castro, 1954, 1967(?); Fernández *et al.*, 1981; Quéro *et al.*, 1986; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Keskin *et al.*, 2015f; Martins and Carneiro, 2018) and Madeira (Keskin *et al.*, 2015f; Desoutter-Meniger and Munroe, 2016). The distribution maps of this species reported by Keskin *et al.* (2015f) and Desoutter-Meniger and Munroe (2016) partially included the area of Selvagens Islands (Madeira EEZ), but specimens are needed for confirmation.

Family Cynoglossidae

***Cynoglossus canariensis* Steindachner 1882**

Common name: Canary tonguesole; **Língua-de-cão-das-Canárias**, Língua-de-cão, Língua, Língua-de-vaca, Linguado.

References: Madeira (Osório de Castro, 1967; Fischer *et al.*, 1981; Adeofe *et al.*, 2015). The type was described by Steindachner (1882) from the Canary Islands. According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. canariensis* is the Eastern central Atlantic. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. The distribution maps of this species reported by Fischer *et al.* (1981) and Adeofe *et al.* (2015) partially included the area of Selvagens Islands (Madeira EEZ). It is possible the occurrence of this species in the Selvagens Islands, but specimens are needed for confirmation.

***Cynoglossus lineolatus* Steindachner 1867**

Common name: No English common name; **Língua-de-cão-riscada**.

References: Azores (?) ([MGYA00167975](#), 2012). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2019) the current geographic distribution of *C. lineolatus* is the Western North Pacific. MGnify (2018) reported a 18S DNA sequence reads of this species (sample TARA_N000002739) collected during the Tara Oceans expedition (2009-2013) at station TARA_151, 137 NM SSW of Pico island (36°10.866'N, 28°56.238'W). The DNA sequence underlying this identification was SSU (small sub-unit or 18S) which is not sufficiently detailed and does not allow any certainty of this occurrence record. Thus, this species is not considered in the total number of valid species to the Azores.

***Syphurus insularis* Munroe, Brito & Hernández 2000**

Common name: Macaronesian tonguesole; **Língua-da-Maraconésia**.

References: Azores ([Munroe et al.](#), 2000; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Lorance *et al.*, 2015d; [Munroe, 2016d](#); Froese and Pauly, 2019) and Madeira (Wirtz, 1994; [Munroe et al.](#), 2000; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Lorance *et al.*, 2015d; [Munroe, 2016d](#); Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). According to Munroe (2016d) this species is sometimes confused with two other species of the genus *Syphurus*, namely *S. nigrescens* Rafinesque 1810 and *S. reticulatus* Munroe 1990. One paratype of this species is in the MMF collections (MFF 22999).

***Syphurus ligulatus* (Cocco 1844)**

Common name: Elongate tonguesole; **Língua-comprida**.

References: mainland Portugal ([Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#)) and Madeira (?) ([Munroe, 1990](#)). This species can be confused with *Syphurus vanmelleae* Chabanaud 1952 ([Munroe, 2016d](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *Syphurus ligulatus* is the Mediterranean Sea and, in the Eastern Atlantic, from Morocco to Angola. Also, Munroe (2016d) does not include Madeira Island in the distribution of this species. Thus, this species is not considered in the total number of valid species to Madeira.

***Syphurus nigrescens* Rafinesque 1810**

Common name: **Spotted tonguesole; Língua-avessa**, Língua, Linguado.

References: mainland Portugal ([Gonçalves, 1941](#); [Chabanaud, 1950](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Torchio, 1979](#); [Fernández et al., 1981](#); [Martins, 1981](#); [INIP, 1982a](#); [Quéro et al., 1986](#); [Fischer et al., 1987](#); [Borges et al., 2002](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Munroe, 2016d](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Collett, 1896a](#); [Brauer, 1906](#); [Kyle, 1913](#); [Norman, 1930](#); [Chabanaud, 1933, 1950](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Torchio, 1979](#); [Quéro et al., 1986](#); [Munroe, 1990](#); [Patzner et al., 1990, 1992](#); [Lloris et al., 1991](#); [Nash et al., 1991](#); [Arruda, 1992, 1997](#); [Azevedo, 1995](#); [Santos et al., 1997](#); [Munroe et al., 2000](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Munroe, 2016d](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira (?) ([Osório de Castro, 1967\(?\)](#); [Maul, 1976](#); [Wirtz, 1994](#); [Tous et al., 2015](#)). This species can be confused with *Syphurus normani* Chabanaud 1950 ([Munroe, 2016d](#)). Three specimens caught off in the coast of mainland Portugal are in the MMF collections (MMF 22797, 1967, set of three specimens). Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira as doubtful and without indicating the geographic location. Munroe (1990) re-identified the Maul's specimen [*S. nigrescens* (not Rafinesque)] as *S. reticulatus*. Later on, Munroe (2000) reclassified Maul's specimen and Wirtz's underwater photos specimens (1994) as *Syphurus insularis* Munroe, Brito & Hernández 2000. The distribution map of this species reported by Tous *et al.* (2015) partially included the area of the Madeira EEZ. It is possible the occurrence of this species in the Madeira Islands and specimens are needed for confirmation. According to Munroe (2000) *S. insularis* is the only species of symphurine tonguefish taken in shallow-water habitats of the Macaronesian subprovince, but this author (Munroe, 2016d) mentions that this species is not frequently reported from the Azores. Thus, this species is not considered in the total number of valid species to Madeira.

***Syphurus reticulatus* Munroe 1990**

Common name: **Reticulated tonguefish; Língua-reticulada**.

References: Madeira (?) ([Munroe, 1990](#); [Brito 1991](#); [Lloris et al., 1991](#); [Wirtz, 1994](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). *S. reticulatus* is often confused with *Syphurus nigrescens* Rafinesque 1810 ([Munroe, 2016d](#)). According to Eschmeyer *et al.* (2019) and Brown *et al.* (2019) this species is endemic to Saint Helena Island. For Munroe *et al.* (2000) all records of *S. reticulatus* until 2000 are referred to *S. insularis*. Thus, this species is not considered in the total number of valid species to Madeira.

Order Atheriniformes

Family Atherinidae

***Atherina boyeri* Risso 1810**

Common name: **Big-scale sand smelt; Peixe-rei-do-Mediterrâneo**, Aterina, Bruxo, Camarão, Camarão-bruxo, Peixe-rei, Piarda, Pica, Rei, Verdugo.

References: mainland Portugal[▲] ([de Brito Capello, 1868, 1880](#); [Nobre, 1895c, 1935](#); [Vieira, 1897, 1900](#); [Seabra, 1911](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Kiener and Spillmann, 1979](#); [Quignard and Pras, 1986a](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Garnerot et al., 2004](#); [Henriques et al., 2008](#); [Francisco et al., 2011](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Gervais and Boulart, 1877b](#); [Kiener and Spillmann, 1979](#); [Quignard and Pras, 1986a](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Tito de Morais et al., 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). This species can be confused with *Atherina presbyter* Cuvier 1829 (Tito de Morais *et al.*, 2016).

Atherina hepsetus Linnaeus 1758

Common name: **Mediterranean sand smelt; Peixe-rei-do-alto**, Beligué, Bruxo, Camarão-bruxo, Guelro, Peixe-rei, Piaba, Piarda, Piava, Pica, Realengo, Rei, Rei-do-alto.

References: mainland Portugal▲ ([Osório de Castro, 1954, 1967; Almaça, 1965b; Quignard and Pras, 1986a; Fischer et al., 1987; Carneiro et al., 2014; Eschmeyer et al., 2019](#)), Azores (?) ([Osório de Castro, 1967](#)) and Madeira (MHNH-IC-1955-0013, n.d.; [Günther, 1861a; Fowler, 1923, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Kiener and Spillmann, 1979; Dooley et al., 1985; Quignard and Pras, 1986a; Lloris et al., 1991; Wirtz et al., 2008; Francisco et al., 2011; Carneiro et al., 2014; Tito de Morais et al., 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *A. hepsetus* is the Mediterranean Sea, Black Sea and, in the Eastern Atlantic, from Galicia (Spain) to Morocco, including Madeira and Canary islands. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location. The presence of this species in the Azores is not yet confirmed and specimens are needed. Thus, this species is not considered in the total number of valid species to the Azores.

Atherina presbyter Cuvier 1829

Common name: **Sand smelt; Peixe-rei**, Beligué, Breta, Bruxo, Camarão, Camarão-bruxo, Camelo, Guelro, Ligueirão, Linguarão, Lingueirão, Peixe-rei-do-alto, Piaba, Piarda, Piava, Pica, Rainha, Realengo, Rei, Rei-do-alto, Verdugo.

References: mainland Portugal▲ ([Vandelli, 1771; de Brito Capello, 1868, 1869b, 1880; Steindachner, 1868b; Posser, 1880; Osório, 1888, 1895; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1903, 1935; Vieira, 1897, 1900; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Quignard and Pras, 1986a; Fischer et al., 1987; Antunes et al., 1988; Arruda et al., 1988; Cabral, 1999; Martins et al., 1999, 2015; Borges et al., 2002; Palma et al., 2003; Garnerot et al., 2004; ICES, 2005; Fernandes et al., 2007; Henriques et al., 2007, 2008; Neves et al., 2008; Ribeiro et al., 2008, 2012; Veiga et al., 2009; Gonçalves et al., 2010; Carneiro et al., 2014; Costa, 2014; Ryland and Lancaster, 2017; Baptista et al., 2019](#)), Azores ([Osório de Castro, 1967; Wood and Williams, 1974; IMAR/DOP COLETA 22, 1990; Wirtz, 1994; Gonçalves et al., 1995; Arruda, 1997; Santos et al., 1997; Bertoncini et al., 2010; Porteiro et al., 2010; Francisco et al., 2011; Carneiro et al., 2014; IMAR/DOP Underwater visual census 5605, 2014; Tito de Morais et al., 2016](#)) and Madeira ([Lowe, 1837; Günther, 1861a; de Veloso, 1892; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Quignard and Pras, 1986a; Lloris et al., 1991; Delgado, 1998; Falcón et al., 2000; Mello, 2000; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; POGIS, 2016; Tito de Morais et al., 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019](#)). This species can be confused with *Atherina boyeri* Risso 1810 (Tito de Morais et al., 2016).

Order Beloniformes

Family Scomberesocidae

Scomberesox saurus (Walbaum 1792)

Common name: **Atlantic saury; Agulhão-do-Atlântico**, Agulhão, Agulha, Catuta, Catutinho, Catuto, Delfim, Marabumbo, Marabundo, Maribombo, Peixe-agulha, Ratinho, Ratinho-do-alto, Tira-vira, Tiravira.

References: mainland Portugal▲ ([de Brito Capello, 1876, 1880; Gervais and Boulart, 1877a; Vieira, 1901; Almeida and Roquette, 1892; Girard, 1892; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muus and Dahlström, 1978; Fischer et al., 1981, 1987; Parin, 1986a; Gordo et al., 1990; ICES, 2005; Carneiro et al., 2014; Martins and Carneiro, 2018; Mecklenburg et al., 2018; Froese and Pauly, 2019](#)), Azores ([Collett, 1896; Murray and Hjort, 1912; Roule, 1919a; Vaillant, 1919; Zugmayer, 1933; Fowler, 1936a; Belloc, 1938; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Parin, 1970, 1976, 1986; Wheeler, 1978; Hubbs and Wisner, 1980; Monteiro, 1996; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017\(?\)](#); Carneiro et al., 2014; Collette, 2016a; Mecklenburg et al., 2018; Froese and Pauly, 2019) and Madeira ([Lowe, 1837; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Parin, 1976, 1986; Fischer et al., 1981](#);

Wisner, 1981, 1990; Carneiro *et al.*, 2014; Collette, 2016a; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019).

***Scomberesox simulans* (Hubbs & Wisner 1980)**

Common name: **Dwarf saury; Agulhão-anão**.

References: mainland Portugal (NHMUK 1990.7.30.182, 1958 (PECS); Fischer *et al.*, 1981; Parin, 1986; Carneiro *et al.*, 2014; Collette, 2015b; Froese and Pauly, 2019), Azores (Murray and Hjort, 1912; Ueyanagi *et al.*, 1972; Hubbs and Wisner, 1980; Fischer *et al.*, 1981; John, 1983; Parin, 1986a; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017(?); Carneiro *et al.*, 2014; Collette, 2015b, 2016a; Froese and Pauly, 2019) and Madeira (NHMUK 1990.7.30.181, 1955; Hubbs & Wisner 1980; Fischer *et al.*, 1981; Wisner, 1981; Parin, 1986a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Collette, 2015b, 2016a; Froese and Pauly, 2019). One specimen is in the MMF collection (MMF 2866).

Family Belonidae

***Belone acus* Risso 1827**

Common name: Garfish; **Agulha-do-Mediterrâneo**, Agulha, Peixe-agulha.

References: mainland Portugal (Collette and Parin, 1970; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Collette and Parin, 1970; Lloris *et al.*, 1991; Collette, 2016a; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; Collette and Parin, 1970; Lloris *et al.*, 1991; Collette, 2016a; Froese and Pauly, 2019). Two syntypes of *Belone gracilis* Lowe 1839 were caught off Madeira, synonym of *B. acus*, and their whereabouts are unknown.

***Belone belone* (Linnaeus 1761)**

Common name: **Garfish; Agulha**, Agulhão, Agulhinha, Catuta, Forda, Peixe-agulha, Peixe-forda, Peixe-poda, Poda.

References: mainland Portugal[▲] (Carvalho, 1765; Vandelli, 1771; NHMUK 2012.10.31.1, 1854-1861; Günther, 1866; de Brito Capello, 1869a, 1880; Posser, 1880; Osório, 1888, 1895, 1896; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1901; Nobre, 1894, 1895a, c, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Parin, 1976, 1986; Borges, 1984; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Arruda *et al.*, 1988; Martins *et al.*, 1992, 1999, 2015; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Cabral, 1999; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Garnerot *et al.*, 2004; ICES, 2005; Henriques *et al.*, 2007, 2008; Gomes and Olim, 2007; Batista *et al.*, 2009; Garrido *et al.*, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Collett, 1890b; Regan, 1903a; Sampaio, 1904; Roule and Angel, 1920, 1930; Nobre, 1924; Fowler, 1936a, b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Svetovidov, 1955; Osório de Castro, 1967; Collette and Parin, 1970; Wood and Williams, 1974; Parin, 1976, 1986; Ré, 1979a; Martins, 1982a, b; Chícharo and Teodósio, 1990; Wirtz, 1990a, 1994; Arruda, 1992, 1997; Nash *et al.*, 1994; Santos *et al.*, 1994b; Pinho *et al.*, 1995; Santos and Nash, 1995; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Froese and Pauly, 2019) and Madeira (Bowdich, 1825; Lowe, 1837, 1839, 1849; Collett, 1890b; de Velloso, 1892; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Parin, 1976, 1986; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One syntype of *Belone cornidii* Günther 1866 caught off in the coast of Portugal, synonym of *B. belone*, is in the NHMUK collections (NHMUK 2012.10.31.1).

***Belone svetovidovi* Collette & Parin 1970**

Common name: Short-beaked garfish; **Agulha-do-bico-curto**, Agulha, Peixe-Agulha.

References: mainland Portugal (Collette and Parin, 1986; Parin, 1986; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Previously confused with *Belone belone* (Linnaeus 1761).

***Platybelone argalus* (Lesueur 1821)**

Common name: **Keeltail needlefish; Agulha-de-quilha**, Agulha, Peixe-Agulha.

References: Azores (Fowler, 1919, 1936b; Norman, 1935a; Collins, 1954; Collette and Parin, 1986, 1970; Parin, 1986a; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Collette, 2016a). Eschmeyer *et al.* (2019) mention *P. argalus* as a circumglobal species in tropical seas and a polytypic species with five to seven subspecies recognized by some authors.

***Tylosurus acus* (Lacepède 1803)**

Common name: **Agujon needlefish; Agulha-imperial**, Agulha, Agulheta, Peixe-Agulha.

References: mainland Portugal (?), Azores (Nobre, 1924, 1930; Fowler, 1936b; Collins, 1954; Osório de Castro, 1967; Arruda, 1997(?); Santos *et al.*, 1997(?); Carneiro *et al.*, 2014) and Madeira (Albuquerque, 1954-1956(?); Osório de Castro, 1967; Carneiro *et al.*, 2014; Collette *et al.*, 2015f; Collette, 2016a). This polytypic species with four valid subspecies (Collette *et al.*, 2015f) can be confused with *Tylosurus crocodilus* (Péron & Lesueur 1821) (Collette, 2016a). According to Santos *et al.* (1997) the identity of the species and their synonyms in this family occurring in the Azores are in need of further confirmation. In mainland Portugal it is possible the occurrence of the subspecies *Tylosurus acus imperialis* (Rafinesque, 1810). Specimens are needed for confirmation in mainland Portugal. See also the remark under *Tylosurus crocodilus*.

***Tylosurus crocodilus* (Péron & Lesueur 1821)**

Common name: **Hound needlefish; Agulha-crocodilo**, Agulha, Peixe-Agulha.

References: mainland Portugal (?). Sporadic catches of this species in southern Portugal (Algarve) has been reported, but the presence of this species along the south coast is uncertain and needs further confirmation. These reports may be misidentifications of the subspecies *Tylosurus acus imperialis* (Rafinesque, 1810). Thus, this species is not considered in the total number of valid species.

Family Hemiramphidae

***Hemiramphus balao* Lesueur 1821**

Common name: **Balao halfbeak; Meia-agulha-preta**, Meia-agulha.

References: Madeira (Fischer *et al.*, 1981; Wirtz, 1994; Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; Nieto, 2015; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019(?)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *H. balao* in the Eastern Atlantic is from Madeira and Canary islands south to Angola. The distribution map of this species reported by Fischer *et al.* (1981) partially included the area of Selvagens Islands (Madeira EEZ).

Family Exocoetidae

***Cheilopogon cyanopterus* (Valenciennes 1847)**

Common name: **Margined flyingfish; Peixe-voador-barbudo**, Voador-barbudo, Voador.

References: mainland Portugal (USNM 228994, 1957; Carneiro *et al.*, 2014; Collette *et al.*, 2015g).

***Cheilopogon exsiliens* (Linnaeus 1771)**

Common name: Bandwing flyingfish; **Peixe-voador-de-asas-listadas**, Avoador, Peixe-voador, Voador.

References: mainland Portugal (Seabra, 1911; Nobre, 1935(?); Carneiro *et al.*, 2014; Collette, 2015c), Azores (Koefoed, 1953; MCZ 149636, 1984; Parin, 1986a (PECS), 2002; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Collette, 2015c) and Madeira (Bowdich, 1825; Lowe, 1837; Lloris *et al.*, 1991; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Collette, 2015c; POGIS, 2016).

***Cheilopogon furcatus* (Mitchill 1815)**

Common name: **Spotfin flyingfish; Peixe-voador-de-barbatana-manchada**, Peixe-voador, Voador-de-barbatana-manchada, Voador.

References: Azores (Roule and Angel, 1930; USNM 299781, 1963 (PECS); MCZ 156674, 1964 (PECS); SMF 13560, 1976; Arruda, 1997; Parin, 2002; Carneiro *et al.*, 2014; Collette, 2015d) and Madeira (Lowe, 1839, 1840a, b, 1841, 1844a, 1849; Fowler, 1936a; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro *et al.*, 2014; Collette, 2015d).

***Cheilopogon heterurus* (Rafinesque 1810)**

Common name: Mediterranean flyingfish; **Peixe-voador-do-Mediterrâneo**, Leiteira, Peixe-voador, Voador, Voador-do-Atlântico.

References: mainland Portugal ([Osório de Castro, 1954, 1967](#); [Parin, 1979, 1986\(?\)](#); [Carneiro et al., 2014](#); [Collette, 2015e](#)), Azores ([Parin and Belyanina, 2000](#); [Parin, 2002](#); [Collette, 2015e](#); [Parin and Shakhovskoy, 2016](#)) and Madeira ([MNHN-IC-0000-4371, 1996](#); [Osório de Castro, 1967](#); [Carneiro et al., 2014](#); [Collette, 2015e](#); [Parin and Shakhovskoy, 2016](#)).

***Cheilopogon pinnatibarbatus* (Bennett 1831)**

Common name: **Bennett's flyingfish**; **Peixe-voador-de-Bennett**, Leiteira, Peixe-voador, Voador, Voador-de-capim.

References: mainland Portugal ([de Brito Capello, 1869a, b, 1880](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Osório, 1896](#); [Vieira, 1901](#); [Bragança, 1902](#); [Seabra, 1911](#); [Nobre, 1935](#); [Bruun, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Parin, 1979, 1986](#); [Martins et al., 1999](#); [Borges et al., 2002](#); [Carneiro et al., 2014](#); [Collette, 2015f](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a](#); [Roule and Angel, 1921, 1923](#); [Nobre, 1924, 1930, 1935](#); [Bruun, 1934, 1935](#); [Fowler, 1936b](#); [Collins, 1954](#); [Koefoed, 1953](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Parin, 1979, 1986](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Collette, 2015f](#); [Parin and Shakhovskoy, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1840a, 1841, 1849](#); [Günther, 1866](#); [Pellegrin, 1905](#); [Bruun, 1935](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Parin, 1979, 1986](#); [Lloris et al., 1991](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Collette, 2015f](#); [Parin and Shakhovskoy, 2016](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)).

***Exocoetus obtusirostris* Günther 1866**

Common name: Oceanic two-wing flyingfish; **Andorinha-do-mar-oceânica**, Andorinha-do-mar, Peixe-voador, Voador.

References: mainland Portugal ([Parin, 1979, 1986](#); [Carneiro et al., 2014](#); [Shakhovskoy, 2018](#); [Froese and Pauly, 2019](#)), Azores ([Bruun, 1935](#); [Koefoed, 1953](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Parin, 1979, 1986, 2002](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#); [Parin and Shakhovskoy, 2016](#); [Lewallen et al., 2017](#); [Froese and Pauly, 2019](#)) and Madeira ([Parin, 1986a](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Parin and Shakhovskoy, 2016](#); [Lewallen et al., 2017](#); [Froese and Pauly, 2019](#)).

***Exocoetus volitans* Linnaeus 1758**

Common name: **Tropical two-wing flyingfish**; **Andorinha-do-mar-tropical**, Andorinha-do-mar, Avoante, Peixe-voador, Voador.

References: mainland Portugal (?) ([Nunes, 1953, 1974](#); [Collette, 2015g](#)), Azores (?) ([Sampaio, 1904](#); [Zugmayer, 1911b](#); [Vaillant, 1919](#); [Nobre, 1895a](#); [Fowler, 1936a](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Parin, 1979, 1986, 2002](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Zugmayer, 1911b](#); [Vaillant, 1919](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Parin, 1979, 1986](#); [Lloris et al., 1991](#); [Parin and Shakhovskoy, 2000, 2016](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Collette, 2015g](#); [Lewallen et al., 2016, 2017](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)). Nunes (1953, 1974) mentioned the occurrence of this species in mainland Portugal and Collette (2015g) included this area in its geographic range. However, its presence in the mainland Portugal is not yet confirmed by more recent data. According to Parin and Shakhovskoy (2000) some records of wandering individuals off the Azores and Madeira may be based on misidentifications. Lewallen et al. (2016) and Parin and Shakhovskoy (2016) indicate that the current geographic distribution of *E. volitans* in the Eastern Atlantic is from 30°N to 30°S, and Eschmeyer et al. (2019) consider it as a circumglobal species in tropical through temperate seas (including Caribbean Sea and Mediterranean Sea). Thus, this species probably may occur in the southern Azores region, which eventually includes the PECS area. Under these circumstances, specimens are needed for confirmation in the Azores and mainland Portugal regions and therefore this species is not considered in the total number of valid species to those areas.

Hirundichthys rondeletii (Valenciennes 1847)

Common name: **Black wing flyingfish; Peixe-voador-de-asa-negra**, Peixe-voador, Voador.

References: mainland Portugal ([Nobre, 1935](#); Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; MCZ 156322, 1969; Parin, 1986a; Fischer et al., 1987; Parin and Belyanina, 2002; Carneiro et al., 2014; Collette, 2015h; Parin and Shakhovskoy, 2016; Froese and Pauly, 2019), Azores ([MNHN-IC-B-0771, 1827](#); Parin, 1986a; Arruda, 1997; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([MCZ 153185, 1986](#); Fischer et al., 1981; Parin, 1986a; Wirtz et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019). According to Parin and Belyanina (2002) *H. rondeletii* is considered to be a superspecies complex that includes four allopatric species found in subtropical waters of all oceans: *H. volador* in the western Atlantic; *H. rondeletii* in the Mediterranean; *H. gilberti* in the northern Pacific; and *H. rufipinnis* in the Southern Hemisphere. These authors consider that *H. rondeletii* is endemic and common in the north-eastern Atlantic, and is known only from the Mediterranean Sea and the adjacent coasts of the Iberian Peninsula and Morocco (Collette, 2015h).

Hirundichthys speculiger (Valenciennes 1847)

Common name: Mirrorwing flyingfish; **Peixe-voador-de-asa-espelhada**, Peixe-voador, Voador.

References: Azores (PECS) ([Parin, 2002](#); Collette, 2015i) and Madeira ([Parin, 1986a](#); [Carneiro et al., 2014](#); [Collette, 2015i](#); [Parin and Shakhovskoy, 2016](#); [Froese and Pauly, 2019](#)). According to Eschmeyer et al. (2019) *H. speculiger* is a circumglobal species in tropical through warm temperate sea (including Caribbean Sea). The distribution map of this species reported by Collette (2015i) includes the area of the Great Meteor Tablemount (Azorean PECS).

Hirundichthys volador (Jordan 1884)

Common name: **Atlantic blackwing flyingfish; Peixe-voador-de-asa-negra-do-Atlântico**.

References: mainland Portugal ([Parin and Belyanina, 2002](#); [Dooley et al., 2015](#); [Froese and Pauly, 2019](#)), Azores ([Parin and Belyanina, 2002](#); [Dooley et al., 2015](#); [Parin and Shakhovskoy, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Parin and Shakhovskoy, 2016](#)). This species can be confused with *Hirundichthys rondeletii* (Valenciennes 1847) ([Parin and Shakhovskoy, 2016](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *H. volador* is the Gulf of Mexico and subtropical Atlantic.

Order Mugiliformes

Family Mugilidae

Chelon labrosus (Risso 1827)

Common name: **Thicklip grey mullet; Tainha-liça**, Coca, Corveo, Córveo, Corvêo, Curveu, Corvêu, Esguiado, Fataça, Garmento, Garnento, Liça, Limento, Muge, Mugem, Muginha, Muja, Mujo, Negrão, Negreta, Negrete, Peixe-saltão, Saltão, Tainha, Tainha-beiçuda, Tainha-de-moda, Tainha-negra, Tainha-negrão, Tainha-preta, Tanha, Tinca.

References: mainland Portugal[▲] ([de Brito Capello, 1868, 1880](#); [Steindachner, 1868b](#); [Posser, 1880](#); [Osório, 1895](#); [Vieira, 1897, 1900](#); [Nobre, 1903, 1931, 1935](#); [Seabra, 1911](#); [Nobre et al., 1915](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Muus and Dahlström, 1978](#); [Trewavas, 1979b](#); [Fischer et al., 1981, 1987](#); [Thomson, 1981, 1990](#); [Dooley et al., 1985](#); [Ben-Tuvia, 1986b](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Erzini et al., 2001a](#); [Jorge et al., 2002](#); [Garnerot et al., 2004](#); [Henriques et al., 2007, 2008](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [Ribeiro et al., 2012](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Turan, 2015](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Drouët, 1861b](#); [Ramos, 1869](#); [Collett, 1897](#); [Hilgendorf, 1888a, b](#); [Sampaio, 1904](#); [Lampe, 1914](#); [Vaillant, 1919](#); [Roule and Angel, 1920, 1930](#); [Nobre, 1923, 1924, 1930](#); [Ferreira, 1934](#); [Fowler, 1936a](#); [Ribeiro, 1936](#); [Cadenat, 1937](#); [Koefoed, 1952](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Trewavas and Ingham, 1972](#); [Mapstone and Wood, 1974, 1975](#); [Wood and Williams, 1974](#); [Wheeler, 1978](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Trewavas, 1979b](#); [Martins, 1982a](#); [Dooley et al., 1985](#); [Ben-Tuvia, 1986b](#); [Barreiros, 1990, 1995](#); [Nash et al., 1990, 1994a](#); [Patzner, 1990](#); [Patzner et al., 1990, 1992](#); [Azevedo et al., 1991, 1995](#); [Lloris et al., 1991](#); [Arruda et al., 1992](#); [Patzner and Santos, 1993](#); [Santos, 1992a](#); [Wirtz, 1994](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Santos et al., 1994b, 1997](#); [Azevedo, 1995](#); [Gonçalves et al., 1995](#); [Pinho et al., 1995](#); [Santos and Nash, 1995](#); [Harmelin-Vivien et al., 2001](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#);

Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Turan, 2015; Afonso, 2016; Harrison, 2016; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Fauconnet *et al.*, 2019) and Madeira (Lowe, 1837, 1843, 1843-1860; Günther, 1861a; Boulenger, 1916; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936a; Albuquerque, 1954-1956; Osório de Castro, 1967; Trewavas, 1979b; Fischer *et al.*, 1981; Thomson, 1981; Dooley *et al.*, 1985; Ben-Tuvia, 1986b; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Falcón *et al.*, 2000; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Turan, 2015; Harrison, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). This species can be confused with *Oedalechilus labeo* (Cuvier 1829) (Harrison, 2016). The whereabouts of the holotype (unique) of *Mugil corrugatus* Lowe 1838 caught off in Madeira, synonym of *G. darwinii*, are unknown.

***Chelon ramada* (Risso 1827)**

Common name: **Thinlip grey mullet; Tainha-fataça**, Alvor, Bicudo, Cabeçudo, Coca, Corveo, Fataça, Fataça-do-Ribeiro, Ilhalvo, Liça, Maguje, Moleca, Muge, Mugem, Muginha, Oirives, Olhal, Olhalvo, Olhinho, Ourives, Peixe-saltão, Saltão, Tagana, Tagano, Tagarra, Tagona, Tainha, Tainha-branca, Tanha, Tinca.

References: mainland Portugal▲ (Carvalho, 1765; de Brito Capello, 1868, 1880; Steindachner, 1868b; Posser, 1880; Osório, 1888; Vieira, 1894a, 1897, 1900; Nobre, 1894, 1931, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Almaça, 1965b; Trewavas, 1979b; Fischer *et al.*, 1981, 1987; INIP, 1981a; Thomson, 1981, 1990; Borges, 1984; Dinis and Marecos, 1984; Ben-Tuvia, 1986b; Antunes *et al.*, 1988; Cabral, 1999; Martins *et al.*, 1999, 2015; Erzini *et al.*, 2001a; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Prista *et al.*, 2003; Garnerot *et al.*, 2004; ICES, 2005; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Batista *et al.*, 2009; Chaves, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Turan, 2015; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Arruda, 1997; Ben-Tuvia, 1986b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Turan, 2015; Harrison, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967; Fischer *et al.*, 1981; Ben-Tuvia, 1986b; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Turan, 2015; Harrison, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019(?)). This species can be confused with *Chelon dumerili* (Steindachner 1870) (Harrison, 2016).

***Chelon saliens* (Risso 1810)**

Common name: **Leaping mullet; Tainha-de-salto**, Coca, Eirigo, Eirigo-do-rio, Fataça, Garrento, Liça, Muge, Mugem, Peixe-saltão, Saltão, Tanha, Tinca.

References: mainland Portugal▲ (Rey, 1935; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Trewavas, 1979b; Ben-Tuvia, 1986b; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Borges *et al.*, 2002; Garnerot *et al.*, 2004; Freyhof, 2008a; Henriques *et al.*, 2008; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Turan, 2015; Martins *et al.*, 2015; Harrison, 2016; Freyhof and Kottelat, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Arruda, 1979a, 1980b, 1997; Santos and Almada, 1988; Santos *et al.*, 1997(?); Turan, 2015; Harrison, 2016) and Madeira (Turan, 2015). This species can be confused with *Chelon dumerili* (Steindachner 1870) (Harrison, 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. saliens* is the Mediterranean Sea, Black Sea, Sea of Azov, and in the Eastern Atlantic, from Bay of Biscay to Morocco. According to Santos *et al.* (1997) the quotations by Arruda (1979, 1980b, 1997) to *Liza saliens* (Risso 1810) are probably a misidentification of *C. labrosus* and it is highly probable that this is the only species occurring in the Azores. However, other authors after Santos *et al.* (1997) describe the presence of this species in the Azores.

***Liza aurata* (Risso 1810)**

Common name: **Golden grey mullet; Tainha-garrento**, Alvor, Barrigudo, Coca, Eirigo, Eirigo-do-rio, Eró, Fataça, Gairenta, Gairrenta, Garmento, Garnento, Garranto, Garrento, Geado, Liça, Maguje, Muge, Mugem, Muginha, Muja, Mujo, Negrão, Negreta, Negrete, Oirives, Ourives, Peixe-saltão, Saltão, Surrega, Tainha, Tainha-amarela, Tainha-da-modas, Tanha, Tinca.

References: mainland Portugal▲ (de Brito Capello, 1868, 1880; Steindachner, 1868b; Posser, 1880; Almeida and Roquette, 1892; Vieira, 1893, 1896; Nobre, 1895a, 1903, 1931, 1935; Osório, 1896; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro,

1954, 1967; Almaça, 1965b; Muusand Dahlström, 1978; Trewavas, 1979b; INIP, 1981a; Martins, 1981; Thomson, 1981, 1990; Dooley *et al.*, 1985; Ben-Tuvia, 1986b; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Arruda *et al.*, 1988; Cabral, 1999; Martins *et al.*, 1999, 2015; Erzini *et al.*, 2001a; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Palma *et al.*, 2003; Prista *et al.*, 2003; Garnerot *et al.*, 2004; ICES, 2005; Henriques *et al.*, 2007, 2008; Gomes and Olim, 2007; Chaves, 2008, 2009; Neves *et al.*, 2008; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Ribeiro *et al.*, 2012; Carneiro *et al.*, 2014; Turan, 2015; Golani and Fricke, 2018; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Fowler, 1936a; Collins, 1954, Albuquerque, 1954-1956; Osório de Castro, 1967; Trewavas, 1979b; Ben-Tuvia, 1986b; Dooley *et al.*, 1985, Barreiros, 1990, Nash *et al.*, 1990, Patzner, 1990, Ré, 1990; Lloris *et al.*, 1991; Arruda, 1997; Morato *et al.*, 2001a; Almeida *et al.*, 2009; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Turan, 2015; Golani and Fricke, 2018; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1839, 1840b; 1843, 1843-1860, 1844a, 1849; de Velloso, 1892; Boulenger, 1916; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Dooley *et al.*, 1985; Ben-Tuvia, 1986b; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Turan, 2015; Harrison, 2016; POGID, 2016; POGIS, 2016; Golani and Fricke, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). This species is sometimes confused with two other species of the genus *Chelon*, namely *C. dumerili* (Steindachner 1870) and *C. ramada* (Risso 1827) (Harrison, 2016). Two specimens caught off in the coast of Lisbon (Portugal), in 1908, are in the MB collections (MB06-004718).

***Mugil cephalus* Linnaeus 1758**

Common name: **Flathead grey mullet; Tainha-olhalvo**, Cabarraz, Cagarraz, Coca, Corvéu, Corveu, Eirigo, Eirigo-do-rio, Eró, Fataça, Ilhalvo, Garmento, Garrento, Liça, Mogueira, Muge, Mugem, Muginha, Mugeira, Mugueiru, Murgueira, Negrão, Olhal, Olhalvo, Olhinho, Papalvo, Peixe-saltão, Saltão, Tagana, Tagano, Tagarra, Tagona, Tainha, Tainha-mogueira, Tanha, Tinca.

References: mainland Portugal▲ (Vandelli, 1771; de Brito Capello, 1868, 1880; Steindachner, 1868b; Posser, 1880; Osório, 1888; Almeida and Roquette, 1892; Nobre, 1895a, 1931, 1935; Vieira, 1897, 1900; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Almaça, 1965b; Trewavas, 1979b; INIP, 1981b; Dooley *et al.*, 1985; Ben-Tuvia, 1986b; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Thomson, 1990; Martins *et al.*, 1992, 1999, 2015; Moreira *et al.*, 1992; Cabral, 1999; Erzini *et al.*, 2001a; Borges *et al.*, 2002; Jorge *et al.*, 2002; Garnerot *et al.*, 2004; ICES, 2005; Henriques *et al.*, 2007; Freyhof, 2008b; Henriques *et al.*, 2008; Neves *et al.*, 2008; Batista *et al.*, 2009; Durand *et al.*, 2013; Carneiro *et al.*, 2014; Turan, 2015; Antunes *et al.*, 2016; Camara *et al.*, 2017; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Cadenat, 1937; Osório de Castro, 1967; Ben-Tuvia, 1986b; Lloris *et al.*, 1991; Arruda, 1997; Turan, 2015) and Madeira▲ (Lowe, 1837, 1843, 1843-1880; Günther, 1861a; Gervais and Boulart, 1877b; Boulenger, 1916; Fowler, 1936a; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Ben-Tuvia, 1986b; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Falcón *et al.*, 2000; Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; Almada *et al.*, 2015; Turan, 2015; Harrison, 2016; POGIS, 2016; Froese and Pauly, 2019(?)). This species can be confused with *Mugil bananensis* (Pellegrin 1927) (Harrison, 2016). Eschmeyer *et al.* (2019) mention it as a nearly circumglobal species in temperate and tropical seas and estuaries. According to Santos *et al.* (1997) the quotation by Cadenat (1937) to *M. cephalus* is probably a misidentification of *C. labrosus* and it is highly probable that this is the only species occurring in the Azores. However, Turan (2015) describes the presence of this species in the Azores and specimens are needed for confirmation. Thus, this species is not considered in the total number of valid species of the Azores.

***Odalechilus labeo* (Cuvier 1829)**

Common name: **Boxlip mullet; Tainha-sabão**, Tainha.

References: mainland Portugal (Fischer *et al.*, 1987; Garnerot *et al.*, 2004; Turan, 2015), Azores (?) (Drouët, 1861b; Hilgendorf, 1888a; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967) and Madeira (Osório de Castro, 1967; Wirtz, 1994; Abreu *et al.*, 1995; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019). According to Trewavas (1979b) and to Santos *et al.* (1997)

the occurrence records of *O. labeo* to the Azores are misidentifications of the species *Chelon labrosus* (Risso, 1826). Thus, this species is not considered in the total number of valid species to the Azores.

Order Gobiesociformes

Family Gobiesocidae

Apletodon dentatus (Facciolà 1887)

Common name: **Small-headed clingfish; Sugador-de-cabeça-pequena**, Sugador, Pegador.

References: mainland Portugal ([Helling, 1943](#); [Albuquerque, 1954-1956](#); [Arruda, 1990](#); [Gonçalves et al., 2002](#); [Henriques et al., 2007, 2008](#); [Brandl et al., 2011](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#)) and Madeira ([EMEPC, 2010](#); [Almada et al., 2015](#); [Fricke et al., 2016a](#); [POGIS, 2016](#)).

Apletodon incognitus Hofrichter & Patzner 1997

Common name: No English common name; **Sugador-dos-ouriços**, Peixe-ventosa-dos-ouriços.

References: mainland Portugal ([Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#)), Azores ([MCZ 12940, 1860](#); [Debelius, 1997](#); [Hofrichter and Patzner, 1997](#); [Santos et al., 1997](#); [Cardigos et al., 2005](#); [Bilecenoğlu and Kaya, 2006](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Fricke et al., 2016a](#); [Froese and Pauly, 2019](#); [Kovačić and Schembri, 2019](#)) and Madeira ([MNHN-IC-1883-0042, 1882](#); [Debelius, 1997](#); [Bilecenoğlu and Kaya, 2006](#); [Floeter et al., 2008](#); [Wirtz et al., 2008\(?\)](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019\(?\)](#); [Kovačić and Schembri, 2019](#)).

Apletodon pellegrini (Chabanaud 1925)

Common name: **Chubby clingfish; Sugador-bochechudo**, Chupa-sangue, Peixe-ventosa, Sugador.

References: Azores ([Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Chabanaud, 1925](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953](#); [Albuquerque, 1954-1956](#); [Briggs, 1955, 1979, 1986, 1990](#); [Osório de Castro, 1967](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Fricke et al., 2016a](#); [POGIS, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Diplecogaster bimaculata (Bonnaterre 1788)

Common name: **Two-spotted clingfish; Sugador-de-duas-manchas**, Pegador, Sugador.

References: mainland Portugal ([Osório, 1895](#); [ZMH 20853, 1901](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Briggs, 1979, 1986](#); [Ribeiro et al., 2006, 2008, 2012](#); [Henriques et al., 2007, 2008](#); [Carneiro et al., 2014](#); [Fricke et al., 2015](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a, b](#); [Collett, 1896, 1897](#); [Guitel, 1904](#); [Roule and Angel, 1921](#); [Fage, 1935](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Lloris et al., 1991](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#)) and Madeira ([Osório de Castro, 1967\(?\)](#); [Fricke, 2007](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Diplecogaster pectoralis Briggs 1955

Common name: No English common name; **Sugador-das-Canárias**, Pegador, Peixe-ventosa, Sugador.

References: Azores ([Briggs, 1979, 1986, 1990](#); [Dooley et al., 1985](#); [Arruda, 1997](#); [Santos et al., 1995a](#); [Floeter et al., 2008](#); [Carneiro et al., 2014](#); [Fricke et al., 2015, 2016a](#); [Bilecenoğlu et al., 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Floeter et al., 2008](#); [Fricke et al., 2015, 2016a](#); [Eschmeyer et al., 2019](#)).

Lepadogaster candolii Risso 1810

Common name: **Connemara clingfish; Sugador-de-De Candolle**, Chupa-sangue, Pegador, Sugador.

References: mainland Portugal ([Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Briggs, 1979, 1986](#); [Arruda, 1990](#); [Henriques et al., 2007, 2008](#); [Carneiro et al., 2014](#); [Fricke et al., 2016a](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1837](#); [Briggs, 1990](#); [Wirtz, 1994](#); [Falcón et al., 2000](#); [Henriques et al., 2002](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Fricke et al., 2016a](#); [POGIS, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Lepadogaster lepadogaster (Bonnaterre 1788)

Common name: **Shore clingfish; Sugador-de-Gouan**, Chupa-sangue, Pegador, Sugador.

References: mainland Portugal (de Brito Capello, 1876, 1880; Nobre, 1895a, 1903, 1935, 1937; Osório, 1895, 1917; Vieira, 1895, 1897, 1901; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Henriques et al., 2002, 2008; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1861a; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Briggs, 1955, 1979, 1986; Osório de Castro, 1967; Augier, 1985; Dooley et al., 1985; Briggs, 1990; Lloris et al., 1991; Falcón et al., 2000; Henriques et al., 2002; Floeter et al., 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Fricke et al., 2016a; POGIS, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of *Lepadogaster zebrinus* Lowe 1839 caught off in Madeira, synonym of *L. lepadogaster*, are unknown.

***Lepadogaster purpurea* (Bonnaterre 1788)**

Common name: **Cornish sucker; Sugador-púrpura**, Chupa-sangue, Sugador.

References: mainland Portugal (ICM-CSIC 6916, n.d.; Briggs, 1979, 1986; Henriques et al., 2002, 2008; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (PECS) (VLIZ 318571, n.d.; Carneiro et al., 2014) and Madeira (Henriques et al., 2002; Floeter et al., 2008; Wirtz et al., 2008; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019).

***Opeatogenys gracilis* (Canestrini 1864)**

Common name: No English common name; **Sugador-do-Mediterrâneo**.

References: mainland Portugal (Gonçalves et al., 2005; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019; Kovačić and Schembri, 2019).

Order Blenniiformes Family Tripterygiidae

***Tripterygion delaisi* Cadenat & Blache 1970**

Common name: **Black-faced blenny; Caboz-de-três-dorsais-de-Delais**, Caboz, Caboz-de-três-dorsais, Cabrito.

References: mainland Portugal (Almeida and Gomes, 1978; Zander, 1986a; Wirtz, 1990b; Oliveira et al., 1992; Almeida, 1996; Gonçalves and Almada, 1998; Almada et al., 2005a; Henriques et al., 2007, 2008; Gonçalves et al., 2010; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Collett, 1897; Roule and Angel, 1920, 1930; Wood and Williams, 1974; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Barreiros, 1990, 1995; Patzner, 1990; Patzner and Santos, 1990c, d, 1993; Patzner et al., 1990, 1992; Santos et al., 1990b, 1997; Wirtz, 1990a, 1994; Azevedo et al., 1991, 1992; Arruda, 1992, 1997; Arruda et al., 1992; Oliveira et al., 1992; Santos, 1992a; Gonçalves et al., 1995; Harmelin-Vivien et al., 2001; Almada et al., 2005a; Cardigos et al., 2005; Floeter et al., 2008; Bertoncini et al., 2010; Porteiro et al., 2010; Carneiro et al., 2014; Williams, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Zander, 1986a; Lloris et al., 1991; Abreu et al., 1995; Delgado, 1998; Falcón et al., 2000; Mello, 2000; Almada et al., 2005a; Araújo et al., 2005; Ribeiro et al., 2005b; Floeter et al., 2008; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Almada et al., 2005b, 2015; Carneiro et al., 2014; Afonso, 2016; POGID, 2016; POGIS, 2016; Williams, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019).

***Tripterygion tripteronotum* (Risso 1810)**

Common name: No English common name; **Caboz-de-três-dorsais-do-Mediterrâneo**, Caboz, Cabrito.

References: Madeira (?) (Lowe, 1837, 1849; Günther, 1861a; Gervais and Boulart, 1877b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Wheeler, 1979g; MNHN-IC-1986-0460, 1985; Carneiro et al., 2014). According to Carreras-Carbonell et al. (2007) *T. tripteronotum* is endemic to the Mediterranean and Eschmeyer et al. (2019) state that the current geographic distribution of this species is the Black Sea and Mediterranean Sea. Carreras-Carbonell et al. (2007), based on genetic and morphological characteristics, split *T. tripteronotum* complex into two distinct species, *T. tripteronotum* and *T. tartessicum* Carreras-Carbonell, Pascual and Macpherson 2007. The current geographic distribution of *T. tartessicum* is the southwestern Mediterranean Sea and the adjacent coast of northern Morocco (in the Eastern Atlantic). There is only one occurrence record of this species outside of the Mediterranean Sea (Eschmeyer et al. 2019). The low larval and adult dispersal capabilities of

Tripterygion species (Carreras-Carbonell *et al.*, 2007), the existence of a natural barrier (e.g. the Gibraltar Strait) and the circulation regime that separates the northern from the southern basins (Send *et al.*, 1999; Carreras-Carbonell *et al.*, 2007) restrict the dispersal of the species outside and within the Mediterranean Sea. The Eastern Atlantic boundary in the distribution of this species is unknown (Carreras-Carbonell *et al.*, 2007). Thus, the quotes of *T. tripteronotum* to Madeira are most likely misidentifications of *T. delaisi* or are less likely *T. tartessicum*. Therefore, this species is not considered in the total number of valid species for Madeira.

Family Labrisomidae

Labrisomus nuchipinnis (Quoy & Gaimard 1824)

Common name: **Hairy blenny**; **Marachomba-peluda**, Alcabroz, Baroti, Caboz, Cabrito, Chova, Manel, Manuel, Marachomba.

References: Madeira (Dooley *et al.*, 1985; Lloris *et al.*, 1991; Oliveira *et al.*, 1992; Ribeiro, 2008; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Williams and Springer, 2016; Froese and Pauly, 2019).

Family Clinidae

Clinatrachus argentatus (Risso 1810)

Common name: **Cline**; **Marachomba-prateada**, Caboz, Macaco, Peixe-macaco.

References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; Wheeler, 1979g; Almeida, 1986; Wirtz and Zander, 1986; Arruda, 1990; Oliveira *et al.*, 1992; Henriques *et al.*, 2007, 2008; Carneiro *et al.*, 2014; Herler *et al.*, 2015a; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Family Blenniidae

Aidablennius sphynx (Valenciennes 1836)

Common name: No English common name; **Marachomba-esfinge**, Marachomba.

References: mainland Portugal (Almada *et al.*, 2001)

Blennius ocellaris Linnaeus 1758

Common name: **Butterfly blenny**; **Marachomba-borboleta**, Alcabroz, Alcabroz, Caboz, Caboz-ocelado, Cabroz, Marachomba.

References: mainland Portugal (MNHN-IC-1887-0165, 1883; Bragança, 1902, 1903; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990a; Fernández *et al.*, 1981; Zander, 1986b; Fischer *et al.*, 1987; Martins *et al.*, 1992; Oliveira *et al.*, 1992; Almada *et al.*, 2001; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; Chaves, 2008; Henriques *et al.*, 2008; Silva, 2008, 2010, 2011; Garrido *et al.*, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Williams *et al.*, 2015a; Bañón *et al.*, 2016a; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Azores (Regan, 1903a; Albuquerque, 1954-1956; Santos, 1987, 1992a; Oliveira *et al.*, 1992; Santos *et al.*, 1994a, 1995a, 1997; Wirtz, 1994; Arruda, 1997; Almada *et al.*, 2001; Almada *et al.*, 2005a; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019). One specimen caught off in the Portuguese continental slope in October 2010 is in the MB collections (MB06-005351).

Coryphoblennius galerita (Linnaeus 1758)

Common name: **Montagu's blenny**; **Marachomba-das-rochas**, Caboz, Caboz-de-crista, Marachomba.

References: mainland Portugal (Osório, 1888, 1896; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990b; Dooley *et al.*, 1985; Zander, 1986b; Antunes *et al.*, 1988; Arruda, 1990; Oliveira *et al.*, 1992; Almeida, 1996; Gonçalves and Almada, 1998; Almada *et al.*, 2001; Almada *et al.*, 2005a; Henriques *et al.*, 2007; Henriques *et al.*, 2008; Garrido *et al.*, 2009; Carneiro *et al.*, 2014; Williams *et al.*, 2015b; Eschmeyer *et al.*, 2017; Ryland and Lancaster, 2017; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a, b; Collett, 1896(?), 1897; Osório de Castro, 1967; Wood and Williams, 1974; Bath, 1977, 1990a; Arruda, 1979a, 1980b; Ré, 1979a; Almada *et al.*, 1983; Almeida and Harmelin-Vivien, 1983; Zander, 1986b; Santos, 1987, 1992a; Barreiros, 1990; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, b, 1997;

Azevedo *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Oliveira *et al.*, 1992; Almada and Santos, 1995; Azevedo, 1995; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2001; Almada *et al.*, 2005a; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Williams *et al.*, 2015b; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1836; Lowe, 1837, 1849; Günther, 1861a; Von Martens, 1876; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990b; Dooley *et al.*, 1985; Zander, 1986b; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Almada *et al.*, 2001; Almada *et al.*, 2005a, 2015; Araújo *et al.*, 2005; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Williams *et al.*, 2015b; POGIS, 2016; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of *Salarias simplicos* Hilgendorf 1888 and the syntype of *Blennius artedii* Cuvier & Valenciennes 1836, synonyms of *C. galerita*, are in the ZMB and MNHN collections (ZMB 12654, caught off in the Azores and MNHN-IC-A-1833, set of two specimens, caught off in Madeira, respectively).

***Hypseurochilus bananensis* (Poll 1959)**

Common name: No English common name; **Marachomba-banana**.

References: mainland Portugal (Almada *et al.*, 2001; Carneiro *et al.*, 2014; Williams *et al.*, 2015c).

***Hypseurochilus fissicornis* (Quoy & Gaimard 1824)**

Common name: No English common name; **Marachomba-da-toca**.

References: Azores (MNHN-IC-0000-0736, 1836 (PECS); Cuvier and Valenciennes, 1836 (PECS); Bath, 1977, 1990b, 1994; Zander, 1986b; Oliveira *et al.*, 1992; Santos *et al.*, 1997; Carneiro *et al.*, 2014; Williams and Craig, 2014(?); Froese and Pauly, 2019). The holotype (unique) of *Blennius fucorum* Cuvier & Valenciennes 1836 caught off about 240 NM South of Azores, synonym of *H. fissicornis*, is in the MNHN collections (MNHN-IC-0000-0736).

***Lipophrys pholis* (Linnaeus 1758)**

Common name: Shanny; **Marachomba-frade**, Alcaboz, Alcabroz, Cabós, Caboz, Caboz-gigante, Cabrão, Cabrito, Cabroz, Diabo, Langanhoso, Lula, Marachomba, Peixe-diabo.

References: mainland Portugal (de Brito Capello, 1868, 1880; Osório, 1888, 1895, 1896; Baldaque da Silva, 1891; Vieira, 1893, 1897; Nobre, 1894, 1895c, 1903, 1935; Bragança, 1902, 1903; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990a; Zander, 1986b; Fischer *et al.*, 1987; Arruda, 1990; Oliveira *et al.*, 1992; Almeida, 1996; Gonçalves and Almada, 1998; Almada *et al.*, 2001; Almada *et al.*, 2005a; Henriques *et al.*, 2007, 2008; Garrido *et al.*, 2009; França *et al.*, 2011; Carneiro *et al.*, 2014; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Arruda, 1979a, 1980b, 1997; Bath, 1979; Harmelin and Harmelin-Vivien, 1979; Zander, 1980; Santos, 1987, 1989, 1992a; Santos *et al.*, 1988, 1994a, 1995a, 1997; Almada *et al.*, 1990; Barreiros, 1990; Patzner *et al.*, 1990, 1992; Azevedo *et al.*, 1991; Arruda *et al.*, 1992; Oliveira *et al.*, 1992; Azevedo, 1995; Almada *et al.*, 2001, 2009; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2005a; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1839, 1843, 1844b; NHMUK 1863.9.10.12-14, 1861-1890; NHMUK 1953.11.15.1, 1937-1960; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990a; Zander, 1986b; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Almada *et al.*, 2001; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Four syntypes of *Pholis bufo* Lowe 1843 caught off in Madeira, synonym of *L. pholis*, are in the NHMUK collections (NHMUK 1863.9.10.12-14, set of three specimens and NHMUK 1953.11.15.1).

***Lipophrys trigloides* (Valenciennes 1836)**

Common name: No English common name; **Marachomba-trigloide**, Alcaboz, Alcabroz, Barriguda, Caboz, Cabroz, Diabo, Marachomba, Peixe-diabo.

References: mainland Portugal (Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990b; Zander, 1986b; Fischer *et al.*, 1987; Arruda, 1990; Oliveira *et al.*, 1992; Almeida, 1996; Almada *et al.*, 2001; Almada *et al.*, 2005a; Henriques *et al.*, 2007, 2008; Carneiro *et al.*, 2014;

Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Arruda, 1979a, 1980b, 1997; Harmelin and Harmelin-Vivien, 1979; Zander, 1980; Almeida and Harmelin-Vivien, 1983; Santos, 1987, 1992a; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994b, 1995a, 1997; Azevedo *et al.*, 1991; Arruda *et al.*, 1992; Oliveira *et al.*, 1992; Wirtz, 1994; Azevedo, 1995; Almada *et al.*, 2001; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2005a; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1836; Lowe, 1837, 1839, 1843, 1849; Günther, 1861a; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979, 1990b; Zander, 1986b; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Almada *et al.*, 2001; Almada *et al.*, 2005a; Araújo *et al.*, 2005; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Microlipophrys canevae* (Vinciguerra 1880)**

Common name: No English common name; **Marachomba-de-Caneva**.

References: mainland Portugal (Almeida and Gomes, 1978; Hureau and Monod, 1979; Zander, 1986; Bath, 1990b; Oliveira *et al.*, 1992; Almada *et al.*, 2001; Almada *et al.*, 2005a; Henriques *et al.*, 2007, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Microlipophrys dalmatinus* (Steindachner & Kolombatović 1883)**

Common name: No English common name; **Marachomba-dálmata**.

References: mainland Portugal (Almeida *et al.*, 1980; Zander, 1986b; Bath, 1990b; Oliveira *et al.*, 1992; Almada *et al.*, 2001; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Ophioblennius atlanticus* (Valenciennes 1836)**

Common name: Redlip Blenny; **Marachomba-do-Atlântico**, Caboz, Cavallo, Marachomba, Moreia-negra, Negro-moreia, Rói-anzóis, Velha.

References: Azores (Collins, 1954; Osório de Castro, 1967; Wood and Williams, 1974; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Ré and Almeida, 1981; Almeida and Harmelin-Vivien, 1983; Santos, 1987, 1992a, 1995a, b; Barreiros, 1990, 1995; Patzner, 1990; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994b, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Arruda *et al.*, 1992; Oliveira *et al.*, 1992, 1993; Santos and Barreiros, 1993; Azevedo, 1995; Gonçalves *et al.*, 1995; Arruda, 1997; Almada *et al.*, 2001; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2005a; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Afonso, 2016; Williams and Springer, 2016; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1836; Lowe, 1837, 1849; Günther, 1861a; Zugmayer, 1911b; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; MNHN-IC-A-2024, 1971; MNHN-IC-B-2510, 1971; Zander, 1986b; Bath, 1990b; Abreu *et al.*, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Almada *et al.*, 2001; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The lectotype and the paralectotype of *Salarias atlanticus* Cuvier & Valenciennes 1836, caught off in Madeira, synonyms of *O. atlanticus*, are in the MNHN collections (MNHN-IC-A-2024 and MNHN-IC-B-2510, respectively).

***Parablennius gattorugine* (Brünnich 1768)**

Common name: Tompot blenny; **Marachomba-babosa**, Alcaboz, Alcabroz, Arremata, Caboz, Cabroz, Diabo, Lula, Macaco, Marachomba, Marachona, Murfeguge, Murtefuge, Peixe-diabo, Peixe-lula, Peixe-macaco, Ranhosa.

References: mainland Portugal (de Brito Capello, 1869a, b, 1880; Posser, 1880; Osório, 1888, 1895, 1896; Girard, 1892; Vieira, 1893, 1897; Nobre, 1894, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Almeida, 1979, 1982, 1996; Bath, 1979, 1990a; Zander, 1986b; Fischer *et al.*, 1987; Arruda *et al.*, 1988; Arruda, 1990; Oliveira *et al.*, 1992; Erzini *et al.*, 1997; Cabral, 1999; Almada *et al.*, 2001, 2007; Garnerot *et al.*, 2004; Almada *et al.*, 2005a; Ribeiro *et al.*, 2006, 2008, 2012; Henriques *et al.*, 2007, 2008; Garrido *et al.*, 2009; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Regan, 1903a; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Bath, 1979; Almeida,

1982; Almeida and Harmelin-Vivien, 1983; Wood and Williams, 1974; Arruda, 1979a, 1980b, 1997; Chícharo and Teodósio, 1990; Carneiro *et al.*, 2014) and Madeira (?) (Lowe, 1849; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bath, 1979; Carneiro *et al.*, 2014). According to Oliveira *et al.* (1992) the occurrence records of *P. gattorugine* from the Azores and Madeira are misidentifications of *Parablennius ruber* (Cuvier and Valenciennes 1836). Thus, the quotes to the Azores and Madeira islands are incorrect and this species is not considered in the total number of valid species for those islands.

***Parablennius incognitus* (Bath 1968)**

Common name: No English common name; **Marachomba-das-cracas**, Caboz-das-cracas.

References: mainland Portugal (Almeida *et al.*, 1980; Zander, 1986b; Arruda, 1990; Bath, 1990b; Oliveira *et al.*, 1992; Almada *et al.*, 2001; Almada *et al.*, 2005a; Henriques *et al.*, 2007, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Nobre, 1924, 1930; Albuquerque, 1954-1956; Wood and Williams, 1974; Almeida and Harmelin-Vivien, 1983; Santos, 1987, 1992a, 1995a; Patzner, 1990; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, b, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Arruda *et al.*, 1992; Oliveira *et al.*, 1992; Almada and Santos, 1995; Azevedo, 1995; Arruda, 1997; Almada *et al.*, 2001; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2005a, b; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Zander, 1986b; Bath, 1990b; Falcón *et al.*, 2000; Almada *et al.*, 2001; Almada *et al.*, 2005a; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Parablennius parvicornis* (Valenciennes 1836)**

Common name: **Rock-pool blenny; Marachomba-das-poças**, Caboz-das-poças.

References: Azores (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Collins, 1954; Albuquerque, 1954-1956; Arruda, 1979a, 1980b, 1992; Zander, 1979, 1980, 1986; Bath, 1982, 1990a, b; Almeida and Harmelin-Vivien, 1983; Santos, 1984, 1985a, b, 1986, 1987, 1989, 1992a, 1995a; Santos and Almada, 1988; Santos *et al.*, 1989, 1994b, 1995b, 1996, 1997; Barreiros, 1990; Chícharo and Teodósio, 1990; Patzner, 1990; Patzner *et al.*, 1990, 1992; Wirtz, 1990a; Azevedo *et al.*, 1991, 1992; Lloris *et al.*, 1991; Monteiro, 1991; Arruda *et al.*, 1992; Oliveira *et al.*, 1992; Gonçalves, 1993; Gonçalves *et al.*, 1993; Santos and Barreiros, 1993; Taborsky, 1994; Almada and Santos, 1995; Azevedo, 1995; Santos and Nash, 1996; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2005a, b; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Williams and Springer, 2016; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1836; Bath, 1971, 1979, 1990b; Dooley *et al.*, 1985; Zander, 1986b; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Almada *et al.*, 2001; Almada *et al.*, 2005a, b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Several authors confused the species *P. parvicornis* and *P. sanguinolentus* since both may be found in rock pools and are of similar shape and colour (Santos *et al.*, 1997). Almada *et al.* (2005b) proved that the genetic distance between *P. parvicornis* and *P. sanguinolentus* is similar or higher than the one found for other blennius that are widely recognized as distinct species and demonstrated that the two are different species due to significant genetic distance. These results, together with the distinct geographic distribution and meristic differences, support that *P. sanguinolentus* and *P. parvicornis* should be considered as two valid species. Their current geographic distributions are: *P. parvicornis* in the Eastern Atlantic, is from Senegal to Congo, and Azores, Madeira, Cape Verde and Canary islands; and *P. sanguinolentus* in the Mediterranean Sea, Black Sea and, in the Eastern Atlantic, from France to Morocco.

***Parablennius pilicornis* (Cuvier 1829)**

Common name: Ringneck blenny; **Marachomba-cornuda**.

References: mainland Portugal (de Brito Capello, 1867a; Almeida *et al.*, 1980; Zander, 1986b; Almada *et al.*, 1987, 2001, 2005a, b; Antunes *et al.*, 1988; Bath, 1990b; Oliveira *et al.*, 1992; Almeida, 1996; Gonçalves and Almada, 1998; Palma *et al.*, 2003; Almada *et al.*, 2005a; Ribeiro *et al.*, 2006, 2008, 2012; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Abecasis *et al.*, 2009; Gonçalves *et al.*, 2010; Pastor and Francour, 2010; Carneiro *et al.*, 2014; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Williams *et al.*, 2015d). According to Eschmeyer *et al.* (2019) the current

geographic distribution of *P. pilicornis* in the Eastern Atlantic is from southern Bay of Biscay to South Africa. The distribution map of this species reported by Williams *et al.* (2015d) partially included the area of Selvagens Islands (Madeira EEZ).

***Parablennius rouxi* (Cocco 1833)**

Common name: No English common name; **Marachomba-branca**, Marachomba, Marachomba-de-risca. References: mainland Portugal (Almeida and Gomes, 1978; Hureau and Monod, 1979; Zander, 1986b; Bath, 1990b; Oliveira *et al.*, 1992; Cabral, 1999; Almada *et al.*, 2001; Henriques *et al.*, 2007, 2008; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Parablennius ruber* (Valenciennes 1836)**

Common name: Portuguese blenny; **Marachomba-portuguesa**, Caboz, Caboz-lusitano, Caboz-português, Murtefuge. References: mainland Portugal (de Brito Capello, 1869a, b; Girard, 1892; Almeida, 1979, 1982; Oliveira *et al.*, 1992; Almada *et al.*, 2001, 2007; Henriques *et al.*, 2008; Abecasis *et al.*, 2009; Carneiro *et al.*, 2014; Williams *et al.*, 2014; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Regan, 1903a; Collins, 1954; Albuquerque, 1954-1956; Wood and Williams, 1974; Arruda, 1979a, 1980b, 1997; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a; Almeida, 1982; Almeida and Harmelin-Vivien, 1983; Zander, 1986b; Santos, 1987, 1992a; Chicharo and Teodósio, 1990; Bath, 1990b; Patzner, 1990; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994b, 1995a, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Lloris *et al.*, 1991; Arruda *et al.*, 1992; Oliveira *et al.*, 1992; Azevedo, 1995; Gonçalves *et al.*, 1995; Almada *et al.*, 2001, 2007; Harmelin-Vivien *et al.*, 2001; Azevedo and Homem, 2002; Almada *et al.*, 2005a, b; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Williams *et al.*, 2014; Afonso, 2016; Williams and Springer, 2016; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1839; Zander, 1986b; Bath, 1990b; Lloris *et al.*, 1991; Oliveira *et al.*, 1992; Almada *et al.*, 2001; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Williams *et al.*, 2014; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Parablennius sanguinolentus* (Pallas 1814)**

Common name: Rusty blenny; **Marachomba-do-Mar-Negro**, Alcaboz, Alcabroz, Caboz, Caboz-das-poças, Frade, Marachomba, Peixe-frade.

References: mainland Portugal (Osório de Castro, 1954, 1967; Zander, 1986b; Fischer *et al.*, 1987; Bath, 1990b; Oliveira *et al.*, 1992; Almada *et al.*, 2001; Garnerot *et al.*, 2004; Almada *et al.*, 2005a, b; Henriques *et al.*, 2007, 2008; Ribeiro *et al.*, 2012; Carneiro *et al.*, 2014; Williams *et al.*, 2015e; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Collett, 1896a; Regan, 1903a; Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1967; Bath, 1982; Santos, 1987, 1989; Santos *et al.*, 1989; Chicharo and Teodósio, 1990; Wirtz, 1990a; Azevedo *et al.*, 1991, 1992; Arruda, 1992, 1997; Taborsky, 1994; Azevedo, 1995; Almada *et al.*, 2001; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (?) (Cuvier and Valenciennes, 1836; Lowe, 1837, 1939, 1843, 1844b, 1849; Günther, 1861a; Collett, 1890b, 1896; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Almada *et al.*, 2001; Araújo *et al.*, 2005; Carneiro *et al.*, 2014; Froese and Pauly, 2019). According to Almada *et al.* (2005b) and Eschmeyer *et al.* (2019) the current geographic distribution of *P. sanguinolentus* is the Mediterranean Sea, Black Sea and, in the Eastern Atlantic, from France to Morocco. Thus, the quotes to the Azores and Madeira islands are incorrect and this species is not considered in the total number of valid species for these archipelagos.

***Parablennius tentacularis* (Brünnich 1768)**

Common name: **Tentacled blenny**; **Marachomba-tentaculada**, Alcaboz, Alcabroz, Caboz, Cabroz, Diabo, Marachomba, Murtefuge, Peixe-diabo.

References: mainland Portugal (de Brito Capello, 1868, 1869b, 1880; Girard, 1892; Osório, 1896; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Bath, 1979, 1990b; Dooley *et al.*, 1985; Zander, 1986b; Fischer *et al.*, 1987; Oliveira *et al.*, 1992(?); Gordo and Cabral, 2001; Carneiro *et al.*, 2014; Williams and Springer, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019(?)) and Madeira (Williams *et al.*, 2015f). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. tentacularis* is the Mediterranean Sea, Black Sea and, in the Eastern Atlantic, from Bay of Biscay to

Senegal, including Canary Islands. The distribution map of this species reported by Williams *et al.* (2015f) partially included the area of Selvagens Islands (Madeira EEZ).

***Salaria basilisca* (Valenciennes 1836)**

Common name: No English common name; **Marachomba-listada**.

References: Azores (?) ([Harmelin and Harmelin-Vivien, 1979](#); [Almeida and Harmelin-Vivien, 1983](#); [Arruda, 1997](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. basilisca* is the Mediterranean Sea (endemic). Harmelin and Harmelin-Vivien (1979) cited the occurrence of *B. basiliscus* in the Azores in a preliminary report, but this misidentification was corrected in a subsequent paper (Almeida and Harmelin-Vivien, 1983). Thus, this species is not considered in the total number of valid species to the Azores.

***Salaria fluviatilis* (Asso y del Rio 1801)**

Common name: Freshwater blenny; **Marachomba-de-água-doce**, Alcaboz, Alcabroz, Caboz, Caboz-de-água-doce, Cabroz.

References: mainland Portugal (?) ([Reis, 1924](#) (cit. by Nobre (1935)); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Almaça, 1965b, 1995](#); [Garcia de Jalon *et al.*, 1989](#); [Oliveira *et al.*, 1992](#); [Collares-Pereira *et al.*, 2000](#); [Freyhof, 2006](#); [Almada *et al.*, 2009b](#)), Azores (?) ([Nobre, 1924](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#)) and Madeira (?) ([Lowe, 1837\(?\)](#)). According to Eschmeyer *et al.* (2019) this species occurs in the following habitats: freshwater, brackish and marine waters. However, *S. fluviatilis* has a preference for lakes and streams with moderate current and stone bottom (Freyhof, 2006). In streams, the deepest and fastest microhabitats are preferred; sometimes in coastal lagoons with low salinity (Freyhof, 2006). In mainland Portugal (Guadiana River basin) *S. fluviatilis* is native, resident and an exclusively freshwater species (Collares-Pereira *et al.*, 2000). The occurrence records of *Blennius inaequalis* Valenciennes 1836, synonym of *S. fluviatilis*, to the Azores and Madeira Islands are probably exclusively to freshwater. Thus, this species is not considered in the total number of valid species.

***Salaria pavo* (Risso 1810)**

Common name: Peacock blenny; **Marachomba-pavão**, Alcaboz, Alcabroz, Caboz, Cabroz, Diabo, Judia, Marachomba, Murtefuge, Peixe-diabo.

References: mainland Portugal ([de Brito Capello, 1868, 1869b, 1880](#); [Posser, 1880](#); [Girard, 1892](#); [Osório, 1896](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bath, 1979, 1990a](#); [Zander, 1986b](#); [Fischer *et al.*, 1987](#); [Antunes *et al.*, 1988](#); [Oliveira *et al.*, 1992](#); [Gonçalves and Almada, 1998](#); [Almada *et al.*, 2001, 2009a](#); [Almada *et al.*, 2005a](#); [Ribeiro *et al.*, 2006, 2008, 2012](#); [Carneiro *et al.*, 2014](#); [Williams *et al.*, 2015g](#); [Eschmeyer *et al.*, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-2005-2353, 1928](#); [Bath, 1973](#); [Carneiro *et al.*, 2014](#)).

***Scartella cristata* (Linnaeus 1758)**

Common name: **Molly miller; Marachomba-de-crista**, Alcaboz, Alcabroz, Caboz, Cabroz, Marachomba.

References: mainland Portugal ([Williams *et al.*, 2015h](#)), Azores ([Williams *et al.*, 2015h](#)) and Madeira ([Falcón *et al.*, 2000](#); [Freitas and Araújo, 2006](#); [EMEPC, 2010](#); [Almada *et al.*, 2015](#); [Williams *et al.*, 2015h](#); [POGIS, 2016](#)). According to Eschmeyer *et al.* (2019) *S. cristata* is widely distributed in the Atlantic Ocean, in the Eastern Atlantic, from Bay of Biscay to Namibia including Canary and Ascension islands, and in the western Atlantic, from Bermuda and Florida to Brazil; the current geographic distribution also includes the Mediterranean Sea. *S. cristata* is generally recognized to be a species complex and currently it is under taxonomic review (Williams *et al.*, 2015h).

Order Lophiiformes

Family Lophiidae

***Lophius budegassa* Spinola 1807**

Common name: **Blackbellied angler; Tamboril-sovaco-preto**, Enxarroco, Peixe-sapo, Penadeira, Pinadeira, Recaimão, Sapo, Tamboril, Tamboril-preto, Tamboril-ruço, Xarrocó.

References: mainland Portugal[▲] ([Osório, 1894](#); [Bragança, 1902](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Almeida and Gomes, 1978](#); [Hureau and Monod, 1979](#); [Fernández *et al.*, 1981](#); [Fischer *et al.*, 1981, 1987](#); [INIP, 1982a](#); [Dooley *et al.*, 1982](#)).

1985; Caruso, 1986, 1990; Martins *et al.*, 1992; Figueiredo *et al.*, 1994; Saldanha *et al.*, 1995; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002a; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; OCEANA, 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Azores (MHNH-IC-1980-1465, 1971; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019). This species can be confused with *Lophius piscatorius* Linnaeus 1758 (Caruso, 2016a).

***Lophius piscatorius* Linnaeus 1758**

Common name: **Angler; Tamboril**, Diabo, Embarroco, Enxarroco, Frango-do-mar, Panadeira, Peixe-diabo, Peixe-engana, Peixe-menino, Peixe-rã, Peixe-sapo, Peixe-tamboril, Penadeira, Pinadeira, Piscatório, Rã, Recaimão, Sapo, Tamboril-branco, Xarroco, Xarroco-maior.

References: mainland Portugal[▲] (Carvalho, 1756(?); Vandelli, 1771; de Brito Capello, 1868, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1903, 1935; Osório, 1895; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muus and Dahlström, 1978; Monod and Le Danois, 1979; Fernández *et al.*, 1981; INIP, 1981a, 1982a, b; Martins, 1981; Dinis, 1985; Caruso, 1986; Fischer *et al.*, 1987; Martins *et al.*, 1992; Saldanha *et al.*, 1995; Erzini *et al.*, 2001a; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002b; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005; Fernandes *et al.*, 2007; Henriques *et al.*, 2008; Pakhorukov, 2008; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Chaves and Silva, 2013; Porteiro *et al.*, 2013; Carneiro *et al.*, 2014; Costa, 2014; OCEANA, 2014; Silva and Borges, 2014; Bañón *et al.*, 2016a; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Mecklenburg *et al.*, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Vaillant, 1888; Silva, 1893; Carqueja, 1894; Goode and Bean, 1895, 1896; Regan, 1903b; Tåning, 1923; Fowler, 1936b; Ferreira, 1940; Collins, 1954; Albuquerque, 1954-1956; Haedrich, 1964; Osório de Castro, 1967; Martins, 1982a; Arruda, 1992, 1997; Krug, 1995; Silva *et al.*, 1995; Santos *et al.*, 1997; Morato *et al.*, 2001a; Melo and Menezes, 2002; Cardigos *et al.*, 2005; Menezes *et al.*, 2006, 2012; Pakhorukov, 2008 (PECS); Porteiro *et al.*, 2010, 2013, 2017; Besugo, 2013; Carneiro *et al.*, 2014) and Madeira (?) (Osório de Castro, 1967; Carneiro *et al.*, 2014). This species can be confused with *Lophius budegassa* Spinola 1807 (Caruso, 2016a). Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. According to Wirtz *et al.* (2008) this species can be found in Madeira fish markets but is not caught off in Madeira Archipelago. The specimen caught off in the Azores in 1903 in Talisman expedition and identified by Regan as *Lophius vaillanti* (MHNH-IC-1887-0223: Paralectotype of *Lophius Vaillanti* Regan, 1903) was subsequently reassessed as paralectotype of *L. piscatorius* Linnaeus 1758.

Family Antennariidae

***Antennarius multiocellatus* (Valenciennes 1837)**

Common name: Longlure frogfish; **Peixe-sapo-multiocelado**.

References: Azores (MHNH-IC-A-5265, 1827; Carneiro *et al.*, 2014). Rare species.

***Antennatus nummifer* (Cuvier 1817)**

Common name: **Spotfin frogfish; Peixe-sapo-salpicado**.

References: Azores (Pietsch, 1986a; Pietsch and Grobecker, 1987; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; NHMUK 1865.6.20.2., 1864-1877; Pietsch, 1986a; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of *Cheironectes bicornis* Lowe 1839, synonym of *A. nummifer*, caught off in Madeira is in the NHMUK collections (NHMUK 1865.6.20.2).

***Fowlerichthys radiosus* Garman 1896**

Common name: Singlespot frogfish; **Peixe-sapo-uniocelado**.

References: Azores (Arruda, 1997; Santos *et al.*, 1997; IMAR/DOP COLETA 129, 2003; IMAR/DOP COLETA 293, 2004; Porteiro and Afonso, 2007; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira (Pietsch,

1986a; Monod and Le Danois, 1979; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Fowlerichthys senegalensis Cadenat 1959

Common name: Senegalese frogfish; **Peixe-sapo-do-Senegal**.

References: Azores (Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Carneiro *et al.*, 2014) and Madeira (Floeter *et al.*, 2008).

Histrio histrio (Linnaeus 1758)

Common name: **Sargassumfish**; **Peixe-sapo-do-sargaço**, Peixe-das-algas, Peixe-rato, Peixe-sapo, Rato.

References: mainland Portugal (ZMH 20979, 1906 (PECS); Pietsch, 1986a; Carneiro *et al.*, 2014), Azores (Collett, 1896a; Ferreira, 1940; Osório de Castro, 1967; Pietsch, 1986a; Pietsch and Grobecker, 1987; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010, 2017(?); Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967(?); Pietsch, 1986a; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

Family Chaunacidae

Chaunacops coloratus (Garman 1899)

Common name: No English common name; **Peixe-sapo-colorido**.

References: Azores (?) (PECS) (MNHN-IC-1996-0225, 1995; Carneiro *et al.*, 2014). The specimen MNHN-IC-1996-0225 collected in September 2, 1995 and initially identified as *Bathychaunax roseus* (Barbour 1941) was renamed as *C. coloratus* in 2006. According to Eschmeyer *et al.* (2019) the actual geographic distribution of this species is the Eastern Pacific. Thus, this species is not considered in the total number of valid species to the Azores.

Chaunacops roseus (Barbour 1941)

Common name: No English common name; **Peixe-sapo-rosado**.

References: Azores (PENN-C 1410, n.d.; PENN-C 1411, n.d.; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014).

Chaunax pictus Lowe 1846

Common name: **Pink frogmouth**; **Peixe-sapo-mole**, Peixe-sapo, Sapo, Sapo-mole.

References: mainland Portugal (Osório, 1911, 1917; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Fernández *et al.*, 1981; Caruso and Pietsch, 1986; Monteiro *et al.*, 2001; Borges *et al.*, 2002; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Silva, 2008, 2011; Chaves, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; OCEANA, 2014; Rijnsdorp *et al.*, 2015b; Froese and Pauly, 2019), Azores (Roule, 1919a; Fowler, 1936b; Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977; Shcherbachov *et al.*, 1985b (PECS); Caruso and Pietsch, 1986; Lloris *et al.*, 1991; Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997(?); Fock *et al.*, 2002 (PECS); Melo and Menezes, 2002; Pakhorukov, 2008 (PECS); Porteiro *et al.*, 2010; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Rijnsdorp *et al.*, 2015b; Froese and Pauly, 2019) and Madeira (Lowe, 1843-1860, 1846a, b, c, 1850, 1852; NHMUK 1857.6.13.183, 1854-1861; Günther, 1861a; Goode and Bean, 1895, 1896; Brauer, 1906; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1967; Ehrich, 1977; Monod, 1979b; Caruso and Pietsch, 1986; Caruso, 1989; Lloris *et al.*, 1991; Quéro *et al.*, 1997; Carneiro *et al.*, 2014; Rijnsdorp *et al.*, 2015b; Caruso, 2016b; Froese and Pauly, 2019). The holotype (unique) of this species caught off 5-6 NM west of Funchal (Madeira) is in the NHMUK collections (NHMUK 1857.6.13.183). According to Santos *et al.* (1997) the Roule's specimen (MOM-IC-910541, year 1902, st. 1338) it probably belongs to the species *Chaunax suttkusi* Caruso 1989. Fowler (1936), Ferreira (1940), Collins (1954), Albuquerque (1954-1956) and Ehrich (1977) are all based in Roule (1919). However, some other authors refer the presence of *C. pictus* in the Azores. One specimen caught off in the Portuguese continental slope in June 2009 is in the MB collections (MB06-005301). Three specimens caught off at the Meteor Bank are in the MMF collections (MMF 22350, 1967, one specimen and MMF 22879, 1967, set of two specimens).

Chaunax suttkusi Caruso 1989

Common name: No English common name; **Peixe-sapo-de-Suttkus**, Sapo.

References: mainland Portugal ([Gonçalves, 1941](#); [Quéro et al., 1997](#); [Caruso, 2016b](#)) and Azores ([Roule, 1919a](#); [Fowler, 1936](#); [Ferreira, 1940](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Ehrich, 1977](#); [Caruso, 1989](#); [Azevedo and Heemstra, 1995](#); [Arruda, 1997](#); [Quéro et al., 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Delgado et al., 2017](#); [Almeida and Biscoito, 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Family Caulophrynidae

Caulophryne jordani Goode & Bean 1896

Common name: Fanfin angler; **Peixe-pescador-de-Jordan**.

References: Azores ([MOM-IC-0091-1529, 1912](#); [Roule and Angel, 1932, 1933](#); [Fowler, 1936b](#); [Ferreira, 1940](#); [Belloc, 1949](#); [Albuquerque, 1954-1956](#); [Maul, 1979e](#); [Pietsch, 1979, 2009](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Bertelsen, 1986a\(?\)](#); [Lloris et al., 1991](#)). The holotype (unique) of *Ceratocaulophryne regani* Roule & Angel 1932, synonym of *C. jordani*, caught off in the Azores (38°33.0'N, 34°4.20'W) is in the MOM collections ([MOM-IC-0091-1529](#)). According to Bertelsen (1986a) the species *C. jordani* is regarded as being present in SW of Madeira, but was caught off outside the EEZ and outside the proposed extension of the Portuguese continental shelf ([Carneiro et al., 2014](#)).

Caulophryne polynema Regan 1930

Common name: No English common name; **Peixe-pescador-da-Madeira**.

References: Azores ([Bertelsen, 1986a](#); [Arruda, 1997](#); [Carneiro et al., 2014](#)) and Madeira ([NHMUK 1930.2.7.1., 1912-1936](#); [Regan, 1930](#); [Regan and Trewavas, 1932a](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Maul, 1979e](#); [Bertelsen, 1986a, 1990a](#); [Lloris et al., 1991](#); [Pietsch, 2009](#); [Carneiro et al., 2014](#)). The holotype (unique) of this species caught off Funchal Bay (Madeira) is in the NHMUK collections ([NHMUK 1930.2.7.1.](#)).

Family Neoceratiidae

Neoceratias spinifer Pappenheim 1914

Common name: Spiny seadevil; **Diabo-marinho-espinhoso**.

References: Azores ([PECS](#)) ([Koefoed, 1932](#); [Pietsch, 2009](#); [Carneiro et al., 2014](#)). Eschmeyer et al. (2019) mention it as a circumglobal species in temperate and tropical seas.

Family Melanocetidae

Melanocetus johnsonii Günther 1864

Common name: **Humpback anglerfish**; **Diabo-marinho-de-Johnson**.

References: mainland Portugal ([Regan and Trewavas, 1932a](#); [Bertelsen, 1986a](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Brauer, 1906](#); [Murray and Hjort, 1912 \(PECS\)](#); [Regan, 1926](#); [Fowler, 1936a](#); [Bertelsen, 1951](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#)) and Madeira ([Günther, 1864b, 1865, 1889](#); [NHMUK 1864.7.18.6, 1864-1877](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Regan, 1926](#); [Regan and Trewavas, 1932a](#); [Fowler, 1936a](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Maul, 1961, 1962a, b, 1979e](#); [Bertelsen, 1986a, 1990a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in the sea of Madeira is in the NHMUK collections ([NHMUK 1864.7.18.6](#)).

Melanocetus murrayi Günther 1887

Common name: Deepsea blackdevil; **Diabo-marinho-de-Murray**.

References: Azores ([PECS](#)) ([ZMH 118749, 1979](#) ; [MAR-ECO - HamPelFish - TI/SF_MAR-ECO_2932, 2004](#); [MAR-ECO - HamPelFish - TI/SF_MAR-ECO_2931, 1979](#); [NHMUK 2004.11.6.88, 1981](#); [Carneiro et al., 2014](#)) and Madeira ([Maul, 1962a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#)). The previous quote by Carneiro et al. (2014) is updated to the Azorean PECS area.

Family Himantolophidae

***Himantolophus albinares* Maul 1961**

Common name: No English common name; **Peixe-farol-da-Madeira**, Farol.

References: Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF_MAR-ECO_4376, 1979](#); Carneiro *et al.*, 2014) and Madeira (Maul, 1961, 1962b; Osório de Castro, 1967; Uyeno *et al.*, 1983; Bertelsen, 1986a, 1990a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off at Câmara de Lobos (Madeira) is in the MMF collections (MMF 2598, 1961).

***Himantolophus brevirostris* (Regan 1925)**

Common name: No English common name; **Peixe-farol-de-focinho-curto**.

References: Azores (Roule and Angel, 1930; Fowler, 1936b; Albuquerque, 1954-1956; MAR-ECO - HamPelFish - TI/SF_MAR-ECO_4379, 1979 (PECS); MAR-ECO - HamPelFish - TI/SF_MAR-ECO_4380, 1979 (PECS); Bertelsen and Krefft, 1988; Carneiro *et al.*, 2014). According to Bertelsen and Krefft (1988) the species *Lipactis tumidus* Regan 1925 is in *Himantolophus brevirostris* group. Bertelsen and Krefft (1988) refer a specimen (ISH 3247/79, 35mm, caught off at 35°12.0' N, 30°9.6' W) of a metamorphosed male of the *Himantolophus brevirostris* group.

***Himantolophus compressus* (Osório 1909)**

Common name: No English common name; **Peixe-farol-sapo**, Farol, Peixe-de-farol, Peixe-farol, Peixe-sapo, Sapo, Tamboril.

References: mainland Portugal (Osório, 1909, 1912; Nobre, 1935; Osório de Castro, 1954, 1967; Bertelsen, 1986a; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Maul, 1961, 1979e; Osório de Castro, 1967; Bertelsen, 1986a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Corynophorus compressus* Osório 1909, synonym of *H. compressus*, caught off at Sesimbra (mainland Portugal) was preserved in the Museu Bocage but destroyed in the 1978 fire.

***Himantolophus groenlandicus* Reinhardt 1837**

Common name: Atlantic footballfish; **Peixe-farol-do-Atlântico**, Peixe-de-farol.

References: mainland Portugal (Girard, 1893a; Brauer, 1906; Osório, 1909, 1917; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Grey, 1956; Maul, 1979e; Bertelsen, 1986a; Carneiro *et al.*, 2014; Froese and Pauly, 2019), Azores (Clarke, 1956; Bertelsen, 1986a; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Maul, 1961; Bertelsen, 1986a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Himantolophus mauli* Bertelsen & Krefft 1988**

Common name: No English common name; **Peixe-farol-de-Maul**.

References: Madeira (Bertelsen and Krefft, 1988; Eschmeyer 1997; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype of this species caught off at Câmara de Lobos (Madeira) is in the MMF collections (MMF 18291).

Family Diceratiidae

***Bufoceratias wedli* (Pietschmann 1926)**

Common name: No English common name; **Peixe-farol-de-Wedl**.

References: mainland Portugal (Pietsch *et al.*, 2004; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Pietschmann, 1926, 1930; Regan and Trewavas, 1932a; Fowler, 1936b; Albuquerque, 1954-1956; Grey, 1956; Maul, 1962a; Hureau and Monod, 1979; Bertelsen, 1986a, 1990a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Phrynichthys wedli* Pietschmann 1926, synonym of *B. wedli*, caught off in Madeira is in the NMW collections (NMW 3524).

Family Oneirodidae

***Chaenophryne draco* Beebe 1932**

Common name: **Smooth dreamer**; **Peixe-farol-dragão**.

References: mainland Portugal ([YPM ICH 004356, 1959](#); [Bertelsen, 1986a](#); [NHMUK 2004.9.12.63, 1987](#); [Carneiro et al., 2014](#)), Azores ([Regan and Trewavas, 1932a](#); [Bertelsen, 1951, 1986a, 1990a](#); [Grey, 1956](#); [Nielsen, 1974](#); [Pietsch, 1979](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#)) and Madeira ([Maul, 1979e](#); [Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Chaenophryne melanodactylus* Regan & Trewavas 1932, synonym of *C. draco*, caught off 56 NM WSW off Santa Maria Island (Azores), 36°36.0'N, 26°14.0'W, is in the ZMUC collections (ZMUC P92116).

***Chaenophryne longiceps* Regan 1925**

Common name: Can-opener smoothdream; **Peixe-farol-cabeça-comprida**.

References: mainland Portugal ([Porteiro et al., 2017](#)), Azores ([Regan, 1925, 1926](#); [Fowler, 1936b](#); [Collins, 1954](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010](#); [Carneiro et al., 2014](#)) and Madeira ([Bertelsen, 1986a](#); [Carneiro et al., 2014](#)).

***Chaenophryne ramifera* Regan & Trewavas 1932**

Common name: No English common name; **Peixe-farol-ramudo**.

References: Azores ([Koefoed, 1944 \(PECS\)](#); [Grey, 1956\(?\)](#); [Bertelsen, 1990a \(PECS\)](#); [Carneiro et al., 2014](#)). Koefoed (1944) mentioned the presence of the species *Chaenophryne pacis* Koefoed 1944 in the Azores PECS which is synonym of *C. ramifera*.

***Ctenochirichthys longimanus* Regan & Trewavas 1932**

Common name: No English common name; **Peixe-farol-barbatana-comprida**.

References: Madeira ([Regan and Trewavas, 1932a](#); [Maul, 1979e](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#)).

***Dolopichthys allector* Garman 1899**

Common name: No English common name; **Peixe-farol-cornudo**.

References: Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF_MAR-ECO_4967, 1979](#); [ZMH 119163, 1979](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Maul, 1961](#); [Carneiro et al., 2014](#)).

***Dolopichthys danae* Regan 1926**

Common name: No English common name; **Peixe-farol-do-Dana**.

References: Madeira ([Regan, 1926](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Maul, 1961, 1979e](#); [Bertelsen, 1986a, 1990a](#); [Lloris et al., 1991](#); [Pietsch, 2009](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Dolopichthys jubatus* Regan & Trewavas 1932**

Common name: No English common name; **Peixe-farol-de-crista**.

References: Azores ([Regan, 1925, 1926](#); [Regan and Trewavas, 1932a](#); [Collins, 1954](#); [Pietsch, 1972a, b](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Ralph, 2015q](#)) and Madeira ([Bertelsen, 1986a, 1990a](#); [Lloris et al., 1991](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Ralph, 2015q](#); [Froese and Pauly, 2019](#)).

***Dolopichthys karsteni* Leipertz & Pietsch 1987**

Common name: No English common name; **Peixe-farol-de-Karsten**.

References: mainland Portugal (?) ([Carneiro et al., 2014](#)), Azores (PECS) ([MCZ 149624, 1978](#); [Carneiro et al., 2014](#)) and Madeira (?) ([Carneiro et al., 2014](#)). According to Eschmeyer et al. (2019) the actual geographic distribution of this species is the Western North Atlantic. This species was erroneously considered present in Madeira and mainland Portugal by Carneiro et al. (2014). Thus, this species is not considered in the total number of valid species for mainland Portugal and Madeira. Males unknown.

***Dolopichthys longicornis* Parr 1927**

Common name: No English common name; **Peixe-farol-diabo**.

References: Azores ([Grey, 1956](#); [MAR-ECO 4339, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017 \(PECS\)](#)) and Madeira ([Regan and Trewavas, 1932](#); [Grey, 1956](#); [Maul, 1979e](#)).

***Dolopichthys pullatus* Regan & Trewavas 1932**

Common name: Lobed dreamer; **Peixe-farol-lobado**.

References: Azores (PECS) ([ZMH 120845, 1982](#); [MAR-ECO - HamPelFish - TI/SF_MAR-ECO_4779, 1982](#); [MAR-ECO - HamPelFish - TI/SF_MAR-ECO_4780, 1982](#); [Pietsch, 2009](#); [Carneiro et al., 2014](#)).

Leptacanthichthys gracilispinis (Regan 1925)

Common name: Plainchin dreamarm; **Peixe-farol-de-queixo-direito**.

References: Azores ([MAR-ECO 7702, 2004](#); [MAR-ECO 7314, 2004](#); [NHMUK 2004.6.29.14, 2004 \(PECS\)](#); [Pietsch, 2009](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([Swinney, 1995a](#); [Pietsch, 2009](#); [Carneiro et al., 2014](#)).

Lophodolos acanthognathus Regan 1925

Common name: Whalehead dreamer; **Peixe-farol-cabeça-de-baleia**.

References: mainland Portugal ([GNM 1946-7637, 1946](#); [Bertelsen, 1986a](#); [Carneiro et al., 2014](#)), Azores ([Regan and Trewavas, 1932b](#); [ZMH 118130, 1979](#); [ZMH 119152, 1979](#); [ZMH 114081, 1975](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017 \(PECS\)](#)) and Madeira ([Porteiro et al., 2017](#)).

Microlophichthys microlophus (Regan 1925)

Common name: Short-rod anglerfish; **Peixe-farol-haste-curta**.

References: Azores ([PECS](#)) ([MAR-ECO 5707, 2004](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#)).

Oneirodes anisacanthus (Regan 1925)

Common name: No English common name; **Peixe-farol-de-bolbo-assimétrico**.

References: Azores ([NHMUK 1925.8.11.13, 1922](#)) and Madeira ([Maul, 1949c, d, 1961](#); [Hureau and Monod, 1979](#); [Bertelsen, 1986a](#), [1990a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Oneirodes carlsbergi (Regan & Trewavas 1932)

Common name: No English common name; **Peixe-farol-de-Carlsberg**.

References: Azores ([NHMUK 2002.11.17.85, 1982](#)).

Oneirodes clarkei Swinney & Pietsch 1988

Common name: No English common name; **Peixe-farol-de-Clarke**.

References: Madeira ([NMSZ 1986.005.1, 1986](#); [Swinney and Pietsch, 1988](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in Madeira (32°24.0'N, 16°51.0'W) is in the NMSZ collections (NMSZ 1986.005.1).

Oneirodes eschrichtii Lütken 1871

Common name: Bulbous dreamer; **Peixe-farol-de-Eschricht**.

References: mainland Portugal ([YPM ICH 005196, 1959](#); [Regan, 1926](#); [Fowler, 1936b](#); [Bertelsen, 1986a](#), [1990a](#); [Carneiro et al., 2014](#); [Rijnsdorp et al., 2015c](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912](#); [Fowler, 1936b](#); [ZMH 119150, 1979](#); [NHMUK 2002.11.17.84, 1982](#); [Arruda, 1997](#); [Carneiro et al., 2014](#); [Rijnsdorp et al., 2015c](#)) and Madeira ([Maul, 1949c, 1951](#); [ZMH 110101, 1971](#); [Bertelsen, 1986a](#), [1990a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Rijnsdorp et al., 2015c](#); [Froese and Pauly, 2019](#)).

Oneirodes krefftii Pietsch 1974

Common name: Krefft's dreamer; **Peixe-farol-de-Krefft**.

References: Madeira ([ZMH 110428, 1971](#)). According to Eschmeyer et al. (2019) *O. krefftii* is a cosmopolitan species in Southern Hemisphere. However, in the Ichthyology Collection of the Zoological Museum of University of Hamburg (ZMH) there is a specimen of this species captured at Madeira (32°47.0'N, 16°24.0'W).

Oneirodes macronema (Regan & Trewavas 1932)

Common name: No English common name; **Peixe-farol-de-Santa Cruz**.

References: Azores ([Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Oneirodes macrosteus Pietsch 1974

Common name: No English common name; **Peixe-farol-do-Atlântico-Norte**.

References: Azores ([MCZ 50022, 1970](#); [Melo and Menezes, 2002](#); [MAR-ECO 7300, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)).

Oneirodes myrionemus Pietsch 1974

Common name: No English common name; **Peixe-farol-multifilamentoso**.

References: Madeira ([ZMH 25185, 1971](#); [ZMH 25186, 1971](#); [Pietsch, 1974](#); [Hureau and Monod, 1979](#); [Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype and one paratype of *O. myrionemus*, caught off in Madeira (32°47.0'N, 16°24.0'W) are in the ZMH collections (ZMH 25185, 1971 and the paratype ZMH 25186, 1971).

Oneirodes posti Bertelsen & Grobecker 1980

Common name: No English common name; **Peixe-farol-de-Post**.

References: Azores (PECS) ([ZMH 25189, 1979](#); [Bertelsen and Grobecker, 1980](#); [Hartel et al., 2008](#); [Carneiro et al., 2014](#)). The holotype of *O. posti*, caught off SW off the Azores (35°24.0' N, 32°01.0' W) are in the ZMH collections (ZMH 25189, 1979).

Pentherichthys atratus (Regan & Trewavas 1932)

Common name: No English common name; **Peixe-farol-do-Panamá**.

References: Azores ([Ralph, 2015r](#)) and Madeira ([Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Ralph, 2015r](#); [Froese and Pauly, 2019](#)). *Pentherichthys venustus* (Regan and Trewavas 1932) is synonym of *P. atratus*.

Phyllorhinichthys micractis Pietsch 1969

Common name: No English common name; **Peixe-farol-de-Guadalupe**.

References: Azores ([MAR-ECO - HamPelFish - TI/SF_MAR-ECO_1419, 1982](#) (PECS); [MAR-ECO 5637, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)).

Puck pinnata Pietsch 1978

Common name: No English common name; **Peixe-farol-alado**.

References: Azores (PECS) ([MAR-ECO - HamPelFish - TI/SF_MAR-ECO_1294, 1979](#); [Carneiro et al., 2014](#)). Very rare species.

Spiniphryne gladiiferae (Beebe 1932)

Common name: Prickly dreamer; **Peixe-farol-espinhoso**.

References: Azores ([NHMUK 2014.9.26.5, 1970](#); [ZMH 117861, 1979](#); [Carneiro et al., 2014](#)). Very rare species, males unknown.

Family Thaumatichthyidae

Lasiognathus amphirhamphus Pietsch 2005

Common name: No English common name; **Peixe-ganchudo**.

References: Azores (PECS) ([NHMUK 2003.11.16.12, 1981](#); [Pietsch, 2005, 2009](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught SW off the Azores (32°22.2'N, 29°50.7'W) is in the NHMUK collections (NHMUK 2003.11.16.12).

Lasiognathus beebei Regan & Trewavas 1932

Common name: No English common name; **Peixe-ganchudo-de-Beebe**.

References: Madeira ([Bertelsen and Pietsch, 1996](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Lasiognathus saccostoma Regan 1925

Common name: No English common name; **Peixe-ganchudo-tridente**.

References: Azores (PECS) ([SOC Discovery 1126120_85_FIS_483001, 1985](#); [Carneiro et al., 2014](#)) and Madeira ([Maul, 1962a, 1979e](#); [Hureau and Monod, 1979](#); [Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Bertelsen and Pietsch, 1996](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype and one paratype of *Lasiognathus ancistrophorus* Maul 1962, synonym of *L. saccostoma*, caught off in Madeira (32°42.0'N, 16°32.0'W) are in the MMF collections (MMF 19019, 1961 and the paratype MMF 19020, 1961). The previous quote by Carneiro et al. (2014) is updated to the Azorean PECS area. Its a very rare species.

Family Centrophrynidae

Centrophryne spinulosa Regan & Trewavas 1932

Common name: Horned lantern fish; **Peixe-farol-córneo**.

References: Madeira ([Regan, 1926](#); [Albuquerque, 1954-1956](#); [Carneiro et al., 2014](#)).

Family Ceratiidae

Ceratias holboelli Krøyer 1845

Common name: **Kroyer's deep-sea angler fish; Peixe-farol-de-Holbøll.**

References: Azores ([Clarke, 1950, 1956](#); [NHMUK 2003.1.13.59, 1970](#); [Pietsch, 1986b](#); [Arruda, 1997](#); [Santos et al., 1997\(?\)](#); [MAR-ECO 5617, 2004](#); [Sutton et al., 2008](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017](#)) and Madeira ([Maul, 1962b](#)). Murray and Hjort (1912) indicate the occurrence of this species between Gran Canaria and Faial, Azores (31°21.0'N, 35°7.0'W). These geographic coordinates correspond to a position near the Atlantis Fracture Zone, but out of the PECS area. Fowler (1936b) and Albuquerque (1954-1956) mention the occurrence of this species near the Azores based on the reference made by Murray and Hjort (1912).

Cryptopsaras couesi Gill 1883

Common name: **Triplewart seadevil; Peixe-farol-de-Coues.**

References: mainland Portugal ([Kobyliansky et al., 2015d](#)), Azores ([Murray and Hjort, 1912](#); [Fowler, 1936b](#); [Bertelsen, 1951](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Pietsch, 1986b](#); [Du Buit et al., 1989](#); [Bekker et al., 1982](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 1999, 2010, 2017 \(PECS\)](#); [Melo and Menezes, 2002](#); [Menezes et al., 2012](#); [Carneiro et al., 2014](#); [Kobyliansky et al., 2015d](#); [Froese and Pauly, 2019](#)) and Madeira ([Bertelsen, 1951, 1986a](#); [NHMUK 2004.7.5.23, 1966](#); [MCZ 49995, 1970](#); [MCZ 50002, 1970](#); [MCZ 51209, 1972](#); [Pietsch, 1986b](#); [Lloris et al., 1991](#); [NHMUK 1996.2.14.24, 1995](#); [Carneiro et al., 2014](#); [Kobyliansky et al., 2015d](#); [Froese and Pauly, 2019](#)). According to Eschmeyer et al. (2019) *C. couesi* is a circumglobal species in tropical through cold temperate seas. The distribution map of this species reported by Kobyliansky et al. (2015d) included the area of mainland Portugal. Specimens are needed for confirmation in mainland Portugal. Barbour (1941) indicates the occurrence of *Cryptopsaras atlantidis* Barbour 1941, synonym of *C. couesi*, caught off north of Azores (50°00'N, 25°20.0'W). These geographic coordinates correspond to a position out of the Azores EEZ and the PECS area. This unique holotype is in the MCZ collections (MCZ 31650).

Family Gigantactinidae

Gigantactis ios Bertelsen, Pietsch & Lavenberg 1981

Common name: No English common name; **Peixe-chicote-das-Canárias.**

References: Madeira ([Pietsch, 2009](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). Eschmeyer et al. (2019) indicated the occurrence of this species [unique holotype in the NHMUK collections (NHMUK 1977.9.13.1)], SW of Madeira (29°49.0'N, 23°0.0'W), based on the reference made by Bertelsen et al., 1981. These geographic coordinates correspond to a position at Canary Basin, west Canary Islands, but out of the Madeira EEZ and the PECS area.

Gigantactis longicirra Waterman 1939

Common name: No English common name; **Peixe-chicote-filamentoso.**

References: Azores (PECS) ([NHMUK 1998.9.8.7, 1980](#); [Carneiro et al., 2014](#)).

Gigantactis vanhoeffeni Brauer 1902

Common name: No English common name; **Peixe-chicote-de-Van Höffen.**

References: Azores ([MAR-ECO 2787, 2004](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017 \(PECS\)](#)).

Family Linophrynidae

Haplophryne mollis (Brauer 1902)

Common name: Soft leafvent angler; **Peixe-sestro-macio.**

References: Azores ([Murray and Hjort, 1912\(?\)](#); [Roule and Angel, 1930\(?\)](#); [Fowler, 1936b\(?\)](#); [NHMUK 2004.8.17.48-50, 1952](#); [Albuquerque, 1954-1956](#); [Collins, 1954](#); [MCZ 68012, 1984 \(PECS\)](#); [SOC Discovery 1126126_84_FIS_382001, 1985 \(PECS\)](#); [SOC Discovery 1126168_86_FIS_382001, 1985 \(PECS\)](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Carneiro et al., 2014](#)) and Madeira ([Maul, 1961, 1962a, 1971a, 1979f](#); [Bertelsen, 1986a, 1990b](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype of

Edriolychnus roulei Regan & Trewavas 1932, synonym of *H. mollis*, caught off between Madeira and Azores islands is in the MOM collections.

***Linophryne arborifera* Regan 1925**

Common name: No English common name; **Peixe-sestro-ramoso**.

References: Madeira ([Maul, 1961, 1979f](#); [Bertelsen, 1990b](#))

***Linophryne algibarbata* Waterman 1939**

Common name: No English common name; **Peixe-sestro-de-Waterman**.

References: Azores (?) ([Porteiro et al., 2017](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *L. algibarbata* is the North Atlantic where the occurrence records are distributed between 49°N and 14°N. Porteiro *et al.* (2017) indicate that this species may probably occur in the northern MAR region, which eventually includes the Azorean Zone. Specimens are needed for confirmation its occurrence in the MAR Azores region. Thus, this species is not considered in the total number of valid species for the Azores.

***Linophryne arcturi* (Beebe 1926)**

Common name: No English common name; **Peixe-sestro-do-Arcturus**.

References: Madeira ([Bertelsen, 1990b](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Linophryne brevibarbata* Beebe 1932**

Common name: No English common name; **Peixe-sestro-de-barbilho-trirramoso**.

References: Azores ([Murray and Hjort, 1912](#); [Koefoed, 1944](#); MAR-ECO - HamPelFish - TI/SF_MAR-ECO_3463, 1979 (PECS); [NHMUK 2002.11.17.87](#), 1982; [Santos et al., 1997](#); [Bertelsen, 1986a](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([ZMH 103546](#), 1966; [Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

***Linophryne coronata* Parr 1927**

Common name: No English common name; **Peixe-sestro-coroad**.

References: mainland Portugal ([Pietsch, 2009](#); [Froese and Pauly, 2019](#)).

***Linophryne indica* (Brauer 1902)**

Common name: Headlight angler; **Peixe-sestro-do-Índico**.

References: Azores (?) ([Murray and Hjort, 1912](#); [Albuquerque, 1954-1956](#)). There are three records of *Aceratias indicus* Brauer specimens, caught off in the Atlantic ocean (west of Canaries; south-west of Azores and Azores) ([Murray and Hjort 1912](#)). However, the current geographic distribution is Indo-Pacific ([Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). Thus, this species is not considered in the total number of valid species.

***Linophryne lucifer* Collett 1886**

Common name: No English common name; **Peixe-sestro-lucifer**.

References: Azores (?) ([Porteiro et al., 2017](#)) and Madeira ([Collett, 1886](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Regan, 1926](#); [Regan and Trewavas, 1932a](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Grey, 1956](#); [Maul, 1979f](#); [Bertelsen, 1986a](#); [Lloris et al., 1991](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in Madeira (36°N, 20°W) is in the ZMUC collections (ZMUC J5560). According to Eschmeyer *et al.* (2019) the current geographic distribution of *L. lucifer* is the North Atlantic. Porteiro *et al.* (2017) indicate this species probably occurs in the northern MAR region, which may eventually includes the Azorean Zone. Specimens are needed for confirmation in the Azores region.

***Linophryne macrodon* Regan 1925**

Common name: No English common name; **Peixe-sestro-dentudo**.

References: Azores ([MAR-ECO 5655, 2004](#); [Sutton et al., 2008\(?\)](#); [Carneiro et al., 2014](#); [Porteiro et al., 2017 \(PECS\)](#)).

***Linophryne maderensis* Maul 1961**

Common name: No English common name; **Peixe-sestro-da-Madeira**.

References: Madeira ([Maul, 1961, 1979f; NHMUK 2003.3.18.27, 1962; Bertelsen, 1986a; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019](#)). The holotype of this species caught off in Madeira (stomach content) is in the MMF collections (MMF 9094, 1956).

Linophryne polypogon Regan 1925

Common name: No English common name; **Peixe-sestro-barbudo**.

References: Azores (PECS) ([NHMUK 2004.9.12.167, 1981](#)) and Madeira ([Regan, 1925; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Grey, 1956; ZMUC P92145, 1970; Maul, 1979f; Bertelsen, 1986a; Lloris et al., 1991; Carneiro et al., 2014; Froese and Pauly, 2019](#)). The holotype of this species caught off north Madeira (33°15.6'N, 16°35.4'W) is in the ZMUC collections (ZMUC P92145).

Linophryne racemifera Regan & Trewavas 1932

Common name: No English common name; **Peixe-sestro-pompom**.

References: Madeira ([Pietsch, 2009; Carneiro et al., 2014; Froese and Pauly, 2019](#)).

Linophryne sexfilis Bertelsen 1973

Common name: No English common name; **Peixe-sestro-hexafilamento**.

References: Madeira ([Bertelsen, 1986a; Lloris et al., 1991; Carneiro et al., 2014; Knudsen, 2015f; Froese and Pauly, 2019](#)).

Photocorynus spiniceps Regan 1925

Common name: No English common name; **Peixe-sestro-cabeça-espinhosa**.

References: Madeira ([Swinney, 1995b; Carneiro et al., 2014; Ralph, 2015s](#)).

Order Tetraodontiformes

Family Molidae

Masturus lanceolatus (Liénard 1840)

Common name: **Sharptail mola; Peixe-lua-rabudo**, Lua, Peixe-lua, Peixe-porco.

References: mainland Portugal ([Carneiro et al., 2014; Ralph, 2015t](#)), Azores ([Collett, 1896; Collins, 1954; Tortonese, 1979h, 1986e; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Cardigos et al., 2005; Porteiro et al., 2010; Carneiro et al., 2014; Ralph, 2015t; Matsuura, 2016b](#)) and Madeira ([Maul, 1949; Wirtz et al., 2008; Carneiro et al., 2014; Ralph, 2015t; Froese and Pauly, 2019](#)). Eschmeyer et al. (2019) mention it as a circumglobal species in tropical through temperate seas.

Mola alexandrini (Ranzani 1839)

Common name: Bump-head sunfish; **Peixe-lua-de-Ranzani**.

References: Azores ([Sawai et al., 2018; Wirtz and Biscoito, 2019](#)) and Madeira ([Wirtz and Biscoito, 2019](#)). According to Sawai et al. (2018) the adults of *M. alexandrini* can be distinguished from other species of *Mola* by the head profile (with bump and chin with bump), body scales rectangular, clavus rounded and ossicles on the rear margin. However, for Wirtz and Biscoito (2019) this species mainly differs from the other *Mola* species by the shape of the clavus and the scales, because in the Eastern Atlantic *Mola mola* (Linnaeus 1758) it can also exhibit a bump on the head, similar to the *M. alexandrini*. This species can be confused with *Mola mola* (Linnaeus 1758) (Sawai et al., 2018).

Mola mola (Linnaeus 1758)

Common name: **Ocean sunfish; Peixe-lua**, Benzedor, Bezedor, Lua, Mergulhão, Mola, Mula, Orelhão, Orelhudo, Peixe-porco, Peixe-roda, Pendão, Porco, Redim, Roda, Rodim, Rolim.

References: mainland Portugal[▲] ([Vandelli, 1771; de Brito Capello, 1869a, 1880; Vieira, 1893, 1897; Almeida and Roquette, 1892; Girard, 1894; Osório, 1896; Bragança, 1902, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979h, 1986e; Borges, 1984; Fischer et al., 1987; Erzini et al., 2001b; Borges et al., 2002; Santos et al., 2002a; ICES, 2005; Fernandes et al., 2007; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Abecasis et al., 2009; Batista et al., 2009; Chaves, 2009; Costa et al., 2012; Carneiro et al., 2014; Costa, 2014; OCEANA, 2014; Martins et al., 2015; Liu et al., 2015; Matsuura, 2016b; Sawai et al., 2018; Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a; Collett, 1896a; Sampaio, 1904; Brian, 1912; Guiart, 1935; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1978; Tortonese, 1979h, 1986e; Lloris et al., 1991; Arruda, 1992, 1997; Santos et al., 1997;](#)

Barreiros and Teves, 2005; Cardigos *et al.*, 2005; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Liu *et al.*, 2015; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Liu *et al.*, 2015; Froese and Pauly, 2019). This species can be confused with *Mola alexandrini* (Ranzani 1839) (Sawai *et al.*, 2018). One specimen is in the MMF collections (MMF 1002).

***Ranzania laevis* (Pennant 1776)**

Common name: **Slender sunfish; Peixe-lua-comprido**, Lua, Lua-comprida, Peixe-lua, Peixe-lua-oblongo. References: mainland Portugal (de Brito Capello, 1869a, 1880; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Fischer *et al.*, 1987; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Matsuura, 2016b; Froese and Pauly, 2019), Azores (Tortonese, 1986e; IMAR/DOP COLETA 203, 1996; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010, 2017(?); Carneiro *et al.*, 2014) and Madeira (Noronha and Sarmento, 1934, 1948; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005312).

Family Diodontidae

***Chilomycterus reticulatus* (Linnaeus 1758)**

Common name: **Pacific burrfish; Porco-espinho-de-pintas**, Ananaz, Baiacú, Balhacú, Bòiacú, Corum, Espinho, Inchabas, Peixe-ouríço, Peixe-porco, Porco, Porco-espinho, Sapo. References: mainland Portugal (Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Dooley *et al.*, 1985; Duron and Quéro, 1990b; Leis, 2006(?); Carneiro *et al.*, 2014; Costa and Rijnsdorp, 2015; Froese and Pauly, 2019), Azores (Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Costa and Rijnsdorp, 2015) and Madeira (Lowe, 1837; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; INIP, 1982c; Dooley *et al.*, 1985; Duron and Quéro, 1990b; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Leis, 2006; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Costa and Rijnsdorp, 2015; POGIS, 2016; Froese and Pauly, 2019).

***Chilomycterus schoepfii* (Walbaum 1792)**

Common name: Striped burrfish; **Porco-espinho-listado**. References: Azores (?) (PECS) (ROM 23914, 1966; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. schoepfii* is the Western North Atlantic. The occurrence record ROM 23914 indicates Atlantic Ocean off Florida (as locally collected) but the geographic coordinates correspond to a position off the Azores at 82 NM west of the Great Meteor Tablemount. However, there is probably a mistake in the geographic coordinates. Thus, this species is not considered in the total number of valid species.

***Chilomycterus spinosus mauretanicus* (Le Danois 1954)**

Common name: Guinean burrfish; **Porco-espinho-da-Guiné**, Baiacú, Balhacú, Bòiacú, Espinho, Peixe-porco, Porco, Porco-espinho, Porco-espinho-listado, Sapo. References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Leis, 2006(?); Jing *et al.*, 2015a), Azores (?) (Osório de Castro, 1967(?)) and Madeira (?) (Osório de Castro, 1967(?)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. spinosus* (subspecies *mauretanicus*) is the Eastern Atlantic. The specimen in the AVG collections (AVG CODC-Peixes PE/1107) probably refers to *Chilomycterus spinosus* subsp. *mauretanicus* (Le Danois 1954). Osório de Castro (1967) only mentions the doubtful occurrence of *Chilomycterus spinosus* (Linnaeus 1758) in the Azores and Madeira Archipelagos. Thus, this species is not considered in the total number of valid species in the Azores and Madeira islands.

***Chilomycterus spinosus spinosus* (Linnaeus 1758)**

Common name: No English common name; **Porco-espinho-americano**, Baiacú, Balhacú, Bòiacú, Espinho, Peixe-porco, Peixe-sapo-grande; Porco, Porco-espinho, Sapo.

References: mainland Portugal (?) ([AVG CODC-Peixes PE/1107, 1899; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967](#)), Azores (?) ([Osório de Castro, 1967\(?\)](#)) and Madeira (?) ([Osório de Castro, 1967\(?\)](#)). The current geographic distribution of this subspecies is in the Western central Atlantic and along South-west America (Leis, 2006; Simon *et al.*, 2013; Jing *et al.*, 2015a; Eschmeyer *et al.*, 2019). There is a specimen of *Chilomycterus spinosus spinosus* (Linnaeus, 1758) in the Coleção do Museu Oceanográfico do Rei D. Carlos I, Aquário Vasco da Gama, Lisbon. Catalogue number PE/1107, caught off in Nazaré, October 23, 1899. Osório de Castro (1967) only mentions the doubtful occurrence of *Chilomycterus spinosus* (Linnaeus 1758) in the Azores and Madeira Archipelagos. Probably the records of the subspecies *spinosus* referred to subspecies *mauretanicus*. Thus, this subspecies is not considered in the total number of valid species.

***Diodon eydouxii* Brisout de Barneville 1846**

Common name: Pelagic porcupinefish; [Peixe-ouriço-pelágico](#), Peixe-ouriço.

References: Azores ([Azevedo *et al.*, 2004; Leis, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Barreiros and Fricke, 2015; Ralph, 2015u; Froese and Pauly, 2019](#)) and Madeira ([Floeter *et al.*, 2008; EMEPC, 2010; Almada *et al.*, 2015; POGIS, 2016](#)).

***Diodon holocanthus* Linnaeus 1758**

Common name: **Longspined porcupinefish**; [Peixe-ouriço-de-crista](#), Espinho, Peixe-ouriço, Peixe-ouriço, Porco, Porco-espinho.

References: Azores ([Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014](#)).

***Diodon hystriculus* Linnaeus 1758**

Common name: **Spot-fin porcupinefish**; [Peixe-ouriço-ponteado](#), Espinho, Inchados, Oiriço, Oiriço-do-mar, Ouriço, Ouriço-do-mar, Peixe-balão-espinhoso, Peixe-ouriço, Peixe-porco, Peixe-porco-de-espinho, Porco, Porco-espinho, Sapo, Sapo-de-espinhos, Sapo-grande.

References: mainland Portugal ([Gonçalves, 1941; Helling, 1943; Lozano, 1952; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Carneiro *et al.*, 2014; Papaconstantinou and Rijnsdorp, 2015d](#)), Azores ([Collins, 1954; Wirtz, 1994; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Ordines *et al.*, 2018a](#)) and Madeira ([Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Lloris *et al.*, 1991; Delgado, 1998; Wirtz *et al.*, 2008\(?\)](#); Carneiro *et al.*, 2014; Ordines *et al.*, 2018a; Froese and Pauly, 2019). According to Azevedo *et al.* (2004) and Wirtz *et al.* (2008) it is possible that some of the records of *D. hystriculus* for the Azores and Madeira, respectively, are in fact a misidentification of *Diodon eydouxii* Brisout de Barneville 1846. The identity of the Azorean and Madeiran *Diodon* needs checking. Eschmeyer *et al.* (2019) mention it as a circumglobal species in tropical and warm temperate seas.

Family Tetraodontidae

***Canthigaster capistrata* (Lowe 1839)**

Common name: Macaronesian sharpnose-puffer; [Peixe-bola-da-Macaronésia](#), Baiacú, Balhacú, Bola, Peixe-sapo, Porquinho, Sapinho, Sapo.

References: mainland Portugal ([Moura, 2004; OCEANA, 2014](#)), Azores ([Moura and Castro, 2002; Cardigos *et al.*, 2006; Floeter *et al.*, 2008; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1839, 1840b, 1844a, 1849; MCZ 99309, 1970; Araújo *et al.*, 2005; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Rijnsdorp and Papaconstantinou, 2015b; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019](#)). Two syntypes of *Tetronodon capistratus* Lowe 1839, synonym of *C. capistrata*, were caught off in Madeira and their whereabouts are unknown.

***Canthigaster rostrata* (Bloch 1786)**

Common name: Sharpnose puffer; [Peixe-bola-das-Caraíbas](#), Baiacú, Balhacú, Bola, Peixe-bola, Porquinho, Sapinho, Sapo.

References: mainland Portugal ([Abecasis *et al.*, 2009; Carneiro *et al.*, 2014\(?\)](#)), Azores ([Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; IMAR/DOP Underwater visual census 5359, 2014; Afonso, 2016](#)) and Madeira (?) ([Lowe, 1839, 1849](#);

Günther, 1870; Jordan and Edwards, 1889; CAS ICH 24034, 1921; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; INIP, 1982c, 1984a, b, c; Dooley *et al.*, 1985; SAIAB FC 34751, 1990; Shipp, 1990; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; POGID, 2016; POGIS, 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. rostrata* is the Western and Eastern Atlantic. However, according to Moura and Castro (2002) all *Canthigaster* specimens so far recorded from Madeira belong to the species *Canthigaster capistrata* (Lowe 1839).

Ephippion guttifer (Bennett 1831)

Common name: **Prickly puffer**; **Peixe-bola-de-pintas**, Baiacú, Baiacú-de-pintas, Balhacú, Bola, Sapo, Tamboril, Tamboril-do-alto.

References: mainland Portugal (ICES, 2005; Silva, 2008, 2010; Carneiro *et al.*, 2014; Matsuura, 2016a; Bañón *et al.*, 2018a) and Madeira (Fischer *et al.*, 1981; MNHN-IC-A-8364, n.d.; Carneiro *et al.*, 2014). The occurrence records of the Muséum national d'Histoire naturelle indicate the presence of this species in the fishing grounds of Morocco (MNHN-IC-A-8364) but the geographic coordinates correspond to a location in the Madeira EEZ.

Lagocephalus laevigatus (Linnaeus 1766)

Common name: **Smooth puffer**; **Peixe-bola-verde**, Baiacú, Baiacú-verde, Baiaçu-verde, Balhacú, Bola, Globo, Peixe-bola, Peixe-sapo, Sapo.

References: mainland Portugal (Fischer *et al.*, 1981; Carneiro *et al.*, 2014; Matsuura, 2016a) and Madeira (Fischer *et al.*, 1981; Lloris *et al.*, 1991; Carneiro *et al.*, 2014). This species can be confused with *Lagocephalus lagocephalus* (Linnaeus 1758) (Matsuura, 2016a).

Lagocephalus lagocephalus (Linnaeus 1758)

Common name: **Oceanic puffer**; **Peixe-bola-oceânico**, Alicate, Baiacú, Balão, Balhacú, Bola, Globo, Peixe-balão, Peixe-bola, Peixe-porco, Peixe-sapo, Porco, Sapinho-do-mar, Sapo.

References: mainland Portugal (de Brito Capello, 1869a, 1880; Osório, 1888; MNHN-IC-1905-0585, 1900; Bragança, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Erzini *et al.*, 2001b; Santos *et al.*, 2002b; Garnerot *et al.*, 2004; Carneiro *et al.*, 2014; Papakonstantinou and Rijnsdorp, 2015c; Matsuura, 2016a; Encarnação *et al.*, 2019), Azores (Vaillant, 1919; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Shipp, 1974; Wheeler, 1978; Tortonese, 1979h, 1986e; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Papakonstantinou and Rijnsdorp, 2015c; Froese and Pauly, 2019) and Madeira (Nobre, 1935; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Papakonstantinou and Rijnsdorp, 2015c; Matsuura, 2016a; Froese and Pauly, 2019). This species can be confused with *Lagocephalus laevigatus* (Linnaeus 1766) (Matsuura, 2016a).

Sphoeroides marmoratus (Lowe 1838)

Common name: Guinean pufferfish; **Peixe-bola-da-Guiné**, Baiacú, Balão, Balhacú, Bola, Pampo, Peixe-balão, Peixe-bola, Peixe-coelho, Peixe-sapo, Pombo, Sapinho, Sapo, Sopapo, Urrô.

References: mainland Portugal (Osório de Castro, 1967; Dooley *et al.*, 1985; Almeida, 1986; Shipp, 1990; Henriques *et al.*, 2007, 2008; Carneiro *et al.*, 2014; Rijnsdorp and Papakonstantinou, 2015c; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a; Regan, 1903a; Sampaio, 1904; Zugmayer, 1933; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese 1973; Wood, 1973; Mapstone and Wood, 1974, 1975; Wood and Williams, 1974; Shipp, 1974; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Saldanha, 1980; Dooley *et al.*, 1985; Barreiros, 1990; Patzner, 1990; Patzner and Santos 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994b; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Nash *et al.*, 1994a; Azevedo, 1995; Gonçalves *et al.*, 1995; Santos and Nash, 1995; Harmelin-Vivien *et al.*, 2001; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Rijnsdorp and Papakonstantinou, 2015c; Afonso, 2016; Almeida and Biscoito, 2019; Froese and Pauly, 2019; Eschmeyer *et al.*, 2019) and Madeira (Bowdich, 1825; Lowe, 1837; Vinciguerra, 1883; MNHN-IC-0000-2160, 1848; Osório de Castro, 1967; Tortonese,

1979h; Dooley *et al.*, 1985; Abreu *et al.*, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Rijnsdorp and Papakonstantinou, 2015c; Matsuura, 2016a; POGID, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). This species can be confused with *Sphoeroides spengleri* (Bloch 1785) (Matsuura, 2016a). The holotype (unique) of *Tetrodon marmoratus* Lowe 1838, synonym of *S. marmoratus*, caught off in Madeira was in the NHMUK collections (NHMUK 1850.11.29.28) but it is not accessible in NHMUK data portal.

***Sphoeroides pachygaster* (Müller & Troschel 1848)**

Common name: **Blunthead puffer**; **Peixe-bola-de-focinho-achatado**, Baiacú, Balão, Balhacú, Bola, Peixe-balão, Peixe-bola, Peixe-sapo, Sapo, Sapo-do-alto.

References: mainland Portugal (Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Calvário *et al.*, 1980; INIP, 1982b; Tortonese, 1986e; Quéro *et al.*, 1997, 1998; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Silva, 2008, 2011; Ward *et al.*, 2008; Silva *et al.*, 2009; Carneiro *et al.*, 2014; Silva and Borges, 2014; Rijnsdorp and Papakonstantinou, 2015d), Azores (Fowler, 1919, 1936b; Collins, 1954; Barnard, 1927; Albuquerque, 1954-1956; Osório de Castro, 1967; Shipp, 1974; Tortonese, 1979h, 1986e; Calvário *et al.*, 1980; Brito, 1983; Dooley *et al.*, 1985; Vacchi, 1986; Lloris *et al.*, 1991; Menezes, 1996; Arruda, 1997; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Rijnsdorp and Papakonstantinou, 2015d) and Madeira (Osório de Castro, 1967; Floeter *et al.*, 2008; Carneiro *et al.*, 2014; Rijnsdorp and Papakonstantinou, 2015d; Matsuura, 2016a; POGID, 2016; Bischoff *et al.*, 2017).

***Sphoeroides spengleri* (Bloch 1785)**

Common name: **Bandtail puffer**; **Peixe-bola-de-pintas**, Sapo, Pombo.

References: mainland Portugal (Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Tortonese, 1979h, 1986e; Dooley *et al.*, 1985; Antunes *et al.*, 1988; Carneiro *et al.*, 2014(?)), Azores (Regan, 1903a; Fowler, 1936b; Albuquerque, 1954-1956; Tortonese, 1979h, 1986e; Dooley *et al.*, 1985; SAIAB FC 36087, 1990; SAIAB FC 36089, 1990; SAIAB FC 36114, 1990; Lloris *et al.*, 1991; SAIAB FC 51520, 1991; SAIAB FC 48896, 1995; Carneiro *et al.*, 2014(?)) and Madeira (Lowe, 1837; Günther, 1870; Vinciguerra, 1883; ZMH 20697, 1931; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Tortonese, 1979h, 1986e; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Carneiro *et al.*, 2014(?)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. spengleri* is the Western and Eastern Atlantic. However, some authors indicate this species is replaced by *S. marmoratus* (Müller and Troschel, 1848) in the Eastern Atlantic.

Family Ostraciidae

***Acanthostracion notacanthus* (Bleeker 1863)**

Common name: Island boxfish; **Peixe-cofre**, Cofre, Peixe-caixa, Peixe-boi.

References: Azores (USNM 42144, 1889-1890; Fowler, 1919, 1936b; Albuquerque, 1954-1956; Tortonese, 1979h, 1986e; Duron and Quéro, 1990a; Lloris *et al.*, 1991; Klassen, 1995; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Tighe, 2015d; Matsuura, 2014, 2016a; Froese and Pauly, 2019).

***Acanthostracion quadricornis* (Linnaeus 1758)**

Common name: **Scrawled cowfish**; **Peixe-cofre-touro**, Cofre, Couraceiro, Peixe-cofre, Peixe-toiro, Peixe-touro, Toiro, Touro.

References: Azores (PECS) (Fowler, 1919, 1936b; Albuquerque, 1954-1956; ROM 23915, 1966; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *A. quadricornis* is the Western Atlantic.

Family Monacanthidae

***Aluterus monoceros* (Linnaeus 1758)**

Common name: **Unicorn leatherjacket filefish**; **Peixe-porco-galhudo**, Peixe-porco, Porco.

References: mainland Portugal (Matsuura *et al.*, 2015; Papakonstantinou and Rijnsdorp, 2015a), Azores (Arruda, 1997; Santos *et al.*, 1997; Morato *et al.*, 2004; Cardigos *et al.*, 2006; Floeter *et al.*, 2008;

Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Freitas and Biscoito, 2002; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Matsuura *et al.*, 2015; Papakonstantinou and Rijnsdorp, 2015a; Matsuura, 2016a; Froese and Pauly, 2019). The distribution map of this species reported by Matsuura *et al.* (2015) included the area of mainland Portugal. Specimens are needed for confirmation in mainland Portugal.

***Aluterus scriptus* (Osbeck 1765)**

Common name: **Scribbled leatherjacket filefish**; **Peixe-porco-bicudo**, Peixe-porco, Peixe-porco-galhudo, Porco.

References: mainland Portugal (Papakonstantinou and Rijnsdorp, 2015b), Azores (IMAR/DOP COLETA 36, 2000; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Papakonstantinou and Rijnsdorp, 2015b; Froese and Pauly, 2019) and Madeira (Freitas and Biscoito, 2002; Freitas and Araújo, 2006; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Papakonstantinou and Rijnsdorp, 2015b; Froese and Pauly, 2019).

***Stephanolepis hispidus* (Linnaeus 1766)**

Common name: **Planehead filefish**; **Peixe-porco-áspero**, Peixe-gatilho-galhudo, Peixe-porco, Peixe-porco-galhudo, Porco, Porco-galhudo.

References: Azores (Wirtz, 1994; Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Lowe, 1850, 1852; Günther, 1870; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979h, 1986e; Dooley *et al.*, 1985; Harmelin-Vivien and Quéro, 1990b; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019). Several syntypes of *Monacanthus auriga* Lowe 1852, synonym of *S. hispidus*, were caught off in Madeira and their whereabouts are unknown.

***Stephanolepis setifer* (Bennett 1831)**

Common name: Pygmy filefish; **Peixe-porco-galhudo**, Peixe-porco, Porco.

References: Madeira (?) (MNHN-IC-0000-6318, 1848; Günther, 1870; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. setifer* is the Western Atlantic. However, there are seven occurrence records of this species in the eastern central Atlantic. All preserved specimens are in the fishes collection of the Muséum national d'Histoire naturelle and in the Canadian Museum of Nature. For K. Matsuura (pers comm.) the record of Madeira specimen is probably a misidentification. Therefore, we considered this record as doubtful until the specimen will be re-evaluated. Thus, this species is not considered in the total number of valid species.

Family Balistidae

***Balistes capriscus* Gmelin 1789**

Common name: **Grey triggerfish**; **Cangulo-cinzento**, Asno, Balista, Balisto, Burro, Cangul, Cangulo, Fambil, Peixe-burro, Peixe-porco, Porco, Tambor.

References: mainland Portugal▲ (de Brito Capello, 1869a, 1880; Osório, 1896; Vieira, 1897; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979h, 1986e; Fischer *et al.*, 1981, 1987; Martins, 1981; Dooley *et al.*, 1985; Almeida, 1996; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Erzini *et al.*, 1997; Martins *et al.*, 1999, 2015; Borges *et al.*, 2002; Jorge *et al.*, 2002; Santos *et al.*, 2002a, b, c; Palma *et al.*, 2003; Garnerot *et al.*, 2004; Gonçalves *et al.*, 2004; Mendes *et al.*, 2004; ICES, 2005; Stergiou *et al.*, 2006; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Abecasis *et al.*, 2009; Batista *et al.*, 2009; Gonçalves *et al.*, 2010; Veiga *et al.*, 2010; Carneiro *et al.*, 2014; Costa, 2014; OCEANA, 2014; Matsuura, 2016a; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores▲ (Hilgendorf, 1888a; Fowler, 1936b; Richard, 1912; Collins, 1954; Albuquerque, 1954-1956; Bauchot, 1967; Osório de Castro, 1967; Wood, 1973; Wood and Williams, 1974; Dias *et al.*, 1976; Wheeler, 1978; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Tortonese, 1979h, 1986e; Saldanha, 1980, 1995; Dooley *et al.*, 1985; Martins, 1982a; Dooley *et al.*, 1985; Sazonov and Galaktionova, 1987; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1997; Wirtz, 1990a, 1994; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Gonçalves, 1993; Azevedo, 1995; Gonçalves *et al.*, 1995; Pinho and Pereira, 1995b;

Silva and Menezes, 1995b; Menezes, 1996; Harmelin-Vivien *et al.*, 2001; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010, 2017; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso, 2016; Almeida and Bischoff, 2019; Fauconnet *et al.*, 2019) and Madeira[▲] (Lowe, 1837; Günther, 1870; de Brito Capello, 1872; Fowler, 1936b; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nunes, 1974; Tortonese, 1979h, 1986e; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Severino, 2004; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Matsuura, 2016a; POGID, 2016; POGIS, 2016; Froese and Pauly, 2019).

Balistes punctatus Gmelin 1789

Common name: **Bluespotted triggerfish; Cangulo-pintado**.

References: Azores (Floeter *et al.*, 2008) and Madeira (Fischer *et al.*, 1981; Dooley *et al.*, 1985; Tortonese, 1986e; Harmelin-Vivien and Quéro, 1990a; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; Matsuura, 2014, 2016a; Tighe, 2015c; Froese and Pauly, 2019(?)).

Balistes vetula Linnaeus 1758

Common name: **Queen triggerfish; Cangulo-real**, Balista, Balisto, Cangulo, Peixe-porco, Porco, Rainha.

References: mainland Portugal (Fischer *et al.*, 1981; Dooley *et al.*, 1985), Azores (Fowler, 1919, 1936b; Collins, 1954; Albuquerque, 1954-1956; Dooley *et al.*, 1985; Tortonese, 1986e; Harmelin-Vivien and Quéro, 1990a; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Bischoff, 2015h; Matsuura, 2016a; Froese and Pauly, 2019) and Madeira (Fischer *et al.*, 1981; Dooley *et al.*, 1985; Matsuura, 2016a).

Canthidermis maculata (Bloch 1786)

Common name: **Rough triggerfish; Cangulo-áspero**, Cangulo, Peixe-porco.

References: mainland Portugal (Rijnsdorp and Papaconstantinou, 2015a), Azores (Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Rijnsdorp and Papaconstantinou, 2015a) and Madeira (MCZ 86747, 1970; Rijnsdorp and Papaconstantinou, 2015a; Matsuura, 2016a).

Canthidermis sufflamen (Mitchill 1815)

Common name: **Ocean triggerfish; Cangulo-oceânico**, Cangulo.

References: Azores (PECS) (ROM 23921, 1966; Carneiro *et al.*, 2014) and Madeira (Falcón *et al.*, 2000; Freitas and Araújo, 2006; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Ralph, 2015v; POGIS, 2016; Froese and Pauly, 2019).

Xanthichthys ringens (Linnaeus 1758)

Common name: Sargassum triggerfish; **Cangulo-do-sargaço**, Balista, Balisto, Cangulo, Peixe-porco, Porco.

References: Azores (?) (Hollard, 1854; Fowler, 1936b; Albuquerque, 1954-1956; Moore, 1967; Osório de Castro, 1967; Randall *et al.*, 1978(?); Arruda, 1997(?)). The holotype (unique) of *Balistes elongatus* Hollard 1854, synonym of *X. ringens*, caught off in the Azores was in the MNHN collections but is not found. According to Randall *et al.* (1978) the Hollard's description of *B. elongatus* is not compatible with the morphological characters of *X. ringens*. Previously, Fowler (1936), Albuquerque (1954-1956) and Moore (1967) reported the species from the Azores on the assumption that *Balistes elongatus* Hollard is *X. ringens*. Eschmeyer *et al.* (2019) reported the current geographic distribution of *X. ringens* in the Western Atlantic and Ascension Island. Thus, this species is not considered in the total number of valid species.

Order Scorpaeniformes

Family Sebastidae

Helicolenus dactylopterus (Delaroche 1809)

Common name: **Blackbelly rosefish; Cantarilho-legítimo**, Acantarilho, Alcantarilha, Alcantarilho, Boca-negra, Cantaril, Cantarilha, Cantarilho, Cantariz, Cântaro, Cardeal, Carneiro, Coça, Galinha, Galinha-do-

mar, Matulha, Pai-de-gato, Papa-jaca, Peixe-carneiro, Rascasso, Requeima, Requeime, Requeimo, Requeme, Rocaz, Ronca, Roncz, Samonete, Serrano, Serrão, Vermelho.

References: mainland Portugal[▲] ([de Brito Capello, 1867a, d, 1880](#); [Steindachner, 1867](#); [Posser, 1880](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1895a, c, 1903, 1935](#); [Osório, 1895](#); [Vieira, 1897, 1898](#); [Brauer, 1906](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Blanc and Hureau, 1979](#); [INIP, 1981a, 1982a, b](#); [Fernández et al., 1981](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Hureau and Litvinenko, 1986](#); [Fischer et al., 1987](#); [Martins et al., 1992](#); [Figueiredo et al., 1994](#); [Saldanha et al., 1995](#); [Erzini et al., 2001a, b](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002b](#); [Mendes et al., 2004](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Pakhorukov, 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Porteiro et al., 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Christiansen et al., 2014](#); [OCEANA, 2014](#); [Bañón et al., 2016a](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Vaillant, 1888, 1919](#); [Collett, 1896a](#); [Regan, 1903a](#); [Sampaio, 1904](#); [Brauer, 1906](#); [Roule, 1919a](#); [Roule and Angel, 1923](#); [Nobre, 1924, 1930, 1935](#); [Norman, 1935a](#); [Fowler, 1936b](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Eschmeyer 1969](#); [Dias et al., 1976, 1990](#); [Ehrich, 1977](#); [Wheeler, 1978](#); [Blanc and Hureau, 1979](#); [Martins, 1982a](#); [INIP, 1985a, b, c](#); [Dias and Monteiro, 1986](#); [Silva, 1986a, b, 1992, 1995](#); [Hureau and Litvinenko, 1986](#); [Bauchot, 1987](#); [Isidro, 1987, 1989, 1996](#); [Eschmeyer and Demster, 1990](#); [Dias, 1991](#); [Dias and Cascalho, 1991](#); [Lloris et al., 1991](#); [Monteiro, 1991](#); [Monteiro et al., 1991](#); [Arruda, 1992, 1997](#); [Pérèz, 1992](#); [Silva and Krug, 1992](#); [Krug, 1995](#); [Santos et al., 1994a, 1995a, 1997](#); [Silva et al., 1994, 1995](#); [Silva and Menezes, 1995a, b](#); [Isidro, 1996](#); [Menezes, 1996](#); [Morato-Gomes et al., 1998](#); [Sousa et al., 1999](#); [Hareide and Garnes, 2001 \(PECS\)](#); [Morato et al., 2001a](#); [Kukuev, 2002 \(PECS\)](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010, 2013](#); [Gomes-Pereira et al., 2012, 2014](#); [Besugo, 2013](#); [Higgins et al., 2013](#); [Pham et al., 2013](#); [Carneiro et al., 2014](#); [Poss, 2016](#); [Almeida and Biscoito, 2019](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([MNHN-IC-1955-0021, n.d.](#); [Lowe, 1837](#); [Günther, 1860](#); [von Martens, 1876](#); [de Velloso, 1892](#); [Collett, 1896](#); [Goode and Bean, 1895, 1896](#); [Norman, 1935a](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [Blanc and Hureau, 1979](#); [Hureau and Litvinenko, 1986](#); [Bauchot, 1987](#); [Eschmeyer and Demster, 1990](#); [Lloris et al., 1991](#); [Ferreira et al., 2008](#); [Pakhorukov, 2008 \(PECS\)](#); [Carneiro et al., 2014](#); [Poss, 2016](#); [Froese and Pauly, 2019](#)). Several specimens caught off in different areas are in the MMF collections: one specimen at the coast of mainland Portugal (MMF 22863) and the others in Josephine Bank (MMF 22416, 1967, set of twenty two specimens and MMF 22852, 1967, one specimen).

***Sebastes norvegicus* (Ascanius 1772)**

Common name: **Golden redfish; Peixe-vermelho**, Cantarilho-do-mares-do-norte, Comunista, Peixe-fino, Sabonete, Vermelho.

References: Azores (PECS) ([Porteiro et al., 2017](#)).

***Sebastes mentella* Travin 1951**

Common name: **Beaked redfish; Peixe-vermelho-da-fundura**.

References: Azores (PECS) ([Hareide and Garnes, 2001](#))

***Trachyscorpia cristulata* (Goode & Bean 1896)**

Common name: **Atlantic thornyhead; Cantarilho-espinhosos**.

References: mainland Portugal ([Blanc and Hureau, 1979](#); [Hureau and Litvinenko, 1986](#); [Fischer et al., 1987](#); [Eschmeyer and Demster, 1990](#); [Moura, 2004](#); [ICES, 2005](#); [Silva, 2008, 2011](#); [Carneiro et al., 2014](#); [Lorance et al., 2015b](#); [Bañón et al., 2016a](#)), Azores ([Arruda, 1997](#); [Santos et al., 1997](#); [Saldanha and Biscoito, 1997](#); [Desbruyères et al., 2001](#); [Melo and Menezes, 2002](#); [Menezes et al., 2006, 2012](#); [IMAR/DOP ImagDOP videos Condor 2087, 2010](#); [Porteiro et al., 2010, 2013](#); [Carneiro et al., 2014](#); [Lorance et al., 2015b](#)) and Madeira ([Hureau and Litvinenko, 1986](#); [Carneiro et al., 2014](#); [Lorance et al., 2015b](#)).

Family Setarchidae

***Etreposebastes imus* Garman 1899**

Common name: Midwater scorpionfish; **Rascasso-da-fundura**.

References: Azores (PECS) ([Kukuev, 2002](#)) and Madeira ([Eschmeyer and Buddo, 2015](#)).

***Setarches guentheri* Johnson 1862**

Common name: **Channeled rockfish; Requeime-de-natura**, Rascasso, Requeime, Requeme, Requeme-de-natura.

References: mainland Portugal ([Marques and Saldanha, 1998](#); [Borges et al., 2002](#); [ICES, 2005](#); [Pakhorukov, 2008](#); [Velasco et al., 2010](#); [Costa, 2014](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Silva and Borges, 2014](#); [Priede, 2017](#)), Azores ([Maul, 1976 \(PECS\)](#); [Saldanha, 1977a](#); [Shcherbachov et al., 1985a](#); [IMAR/DOP COLETA 206, 1989](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Fock et al., 2002 \(PECS\)](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#)) and Madeira[▲] ([NHMUK 1862.4.22.36, 1861-1890](#); [Johnson, 1862a, 1863a](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Norman, 1935b](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Eschmeyer, 1969](#); [Maul, 1976](#); [Poss and Rama Rao, 1984](#); [Blanc and Hureau, 1979](#); [Poss and Rama Rao, 1984](#); [Hureau and Litvinenko, 1986](#); [Eschmeyer and Demster, 1990](#); [Lloris et al., 1991](#); [Marques and Saldanha, 1998](#); [Pakhorukov, 2008 \(PECS\)](#); [Velasco et al., 2010](#); [Carneiro et al., 2014](#); [POGID, 2016](#); [Poss, 2016](#); [Roberts et al., 2017](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in Madeira is in the NHMUK collections ([NHMUK 1862.4.22.36](#)). One specimen caught off at the Meteor Bank is in the MMF collections ([MMF 22883, 1967](#)).

Family Scorpaenidae

***Phenacoscopius nebris* Eschmeyer 1965**

Common name: Short-tube scorpionfish; **Rascasso-de-tubo-curto**.

References: Madeira (PECS) ([SDSC 16587, 1982](#); [Carneiro et al., 2014](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *P. nebris* is the Western Atlantic.

***Pontinus kuhlii* (Bowdich 1825)**

Common name: **Offshore rockfish; Cantarilho-requeime**, Brage, Cantareiro, Cantaril, Cantarilho, Cântaro, Cardeal, Chapado, Galhudo, Palhaço, Rascasso, Requeima, Requeime, Requeimo, Requeme, Toupeira, Vermelho.

References: mainland Portugal[▲] ([de Brito Capello, 1867a, d, 1880](#); [Steindachner, 1867](#); [Baldaque da Silva, 1891](#); [Seabra, 1911](#); [Nobre, 1935](#); [Norman, 1935b](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Eschmeyer, 1969](#); [Blanc and Hureau, 1979](#); [INIP, 1982a](#); [Martins, 1981](#); [Dooley et al., 1985](#); [Hureau and Litvinenko, 1986](#); [Fischer et al., 1987](#); [Eschmeyer and Demster, 1990](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [OCEANA, 2014](#); [Poss, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Regan, 1903a](#); [Norman, 1935b](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Eschmeyer, 1969](#); [Dias et al., 1976, 1990](#); [Maul, 1976](#); [Ehrich, 1977](#); [Blanc and Hureau, 1979](#); [Martins, 1982a](#); [Dooley et al., 1985](#); [INIP, 1985a, b, c](#); [Dias and Monteiro, 1986](#); [Hureau and Litvinenko, 1986](#); [Dias, 1991](#); [Lloris et al., 1991](#); [Monteiro, 1991](#); [Monteiro et al., 1991](#); [Silva and Krug, 1992](#); [Krug, 1995](#); [Silva et al., 1994, 1995](#); [Silva, 1995](#); [Silva and Menezes, 1995a, b](#); [Isidro, 1996](#); [Menezes, 1996](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Morato-Gomes et al., 1998](#); [Sousa et al., 1999](#); [Morato et al., 2001a](#); [Kukuev, 2002 \(PECS\)](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Floeter et al., 2008](#); [Porteiro et al., 2010, 2013](#); [Gomes-Pereira et al., 2012, 2014](#); [Besugo, 2013](#); [Higgins et al. 2013](#); [Pham et al., 2013](#); [Carneiro et al., 2014](#); [Poss, 2016](#); [Eschmeyer et al., 2019](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#) and Madeira[▲] ([Bowdich, 1825](#); [Lowe, 1837, 1843-1860](#); [Günther, 1860, 1880](#); [Sauvage, 1878](#); [de Velloso, 1892](#); [Goode and Bean, 1895, 1896](#); [Collett, 1896](#); [Zugmayer, 1911b \(PECS\)](#); [Roule, 1919a](#); [Nobre, 1935](#); [Norman, 1935a](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Eschmeyer, 1969](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [Blanc and Hureau, 1979](#); [Dooley et al., 1985](#); [Hureau and Litvinenko, 1986](#); [Lloris et al., 1991](#); [Severino, 2004](#); [Ferreira et al., 2008](#); [Floeter et al., 2008](#); [Christiansen et al., 2009](#); [Carneiro et al., 2014](#); [Poss, 2016](#); [Biscoito et al., 2016](#)).

2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The specimen from Lowe of *Scorpaena kuhlii* Bowdich 1825, synonym of *P. kuhlii*, caught off in Madeira is in the SMF collections (SMF 14).

Scorpaena azorica Eschmeyer 1969

Common name: **Azores scorpionfish; Rascasso-dos-Açores**.

References: Azores (USNM 94463, 1884; Eschmeyer, 1969; Blanc and Hureau, 1979; Hureau and Litvinenko, 1986; Lloris *et al.*, 1991; Santos *et al.*, 1994a, 1995a, 1997(?); Arruda, 1997; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Menezes 2015; Poss, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of this species caught off in Terceira Island (Azores) is in the USNM collections (USNM 94463).

Scorpaena canariensis (Sauvage 1878)

Common name: Canary scorpionfish; **Rascasso-das-Canárias**.

References: Azores (Wirtz, 1994, 2005; Motomura *et al.*, 2005; Cardigos *et al.*, 2006; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Poss, 2016; Eschmeyer *et al.*, 2019) and Madeira (Brito, 1991; Brito *et al.*, 2002; Motomura *et al.*, 2005; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Poss, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Scorpaena elongata Cadenat 1943

Common name: **Slender rockfish; Rascasso-rosado**, Galinha-do-mar, Rascasso, Rescasso, Rocaz.

References: mainland Portugal (Hureau and Litvinenko, 1986; Borges *et al.*, 2002; Carneiro *et al.*, 2014; Martins and Carneiro, 2018), Azores (Poss, 2016) and Madeira (Poss, 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. elongata* is the Mediterranean Sea and the Eastern Atlantic. The distribution map of this species reported by Poss (2016) partially included the area of Selvagens Islands (Madeira EEZ). Specimens are needed for confirmation in Madeira Archipelago.

Scorpaena laevis Troschel 1866

Common name: **Senegalese rockfish; Rascasso-do-Senegal**, Galinha-do-mar, Rascasso, Requeima, Requeime, Requeimo, Rescasso, Rocaz, Roncaz.

References: Azores (Collins, 1954; Eschmeyer, 1969; Blanc and Hureau, 1979; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Carneiro *et al.*, 2014; Biscoito, 2015e; Poss, 2016; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira (Eschmeyer, 1969; Blanc and Hureau, 1979; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Lloris *et al.*, 1991; Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; Poss, 2016; Eschmeyer *et al.*, 2019).

According to Froese and Pauly (2019) this species is absent of Madeira.

Scorpaena loppei Cadenat 1943

Common name: **Cadenat's scorpionfish; Rascasso-de-Loppé**.

References: mainland Portugal (MNHN-IC-1887-0307, 1883; Maul, 1976; Blanc and Hureau, 1979; Hureau and Litvinenko, 1986; Fischer *et al.*, 1987; Eschmeyer and Demster, 1990; Carneiro *et al.*, 2014; Poss, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (PECS) (SDSC 23056, 1967; Maul, 1976; Ehrich, 1977; SDSC 9481, 1998; Kukuev, 2002; Carneiro *et al.*, 2014) and Madeira (PECS) (SDSC 23052, 1967; SDSC 23053, 1967; Maul, 1976; Carneiro *et al.*, 2014). Several specimens caught off in different areas are in the MMF collections: four specimens at the Josephine Bank (MMF 22803, 1967, set of three specimens, and MMF 22851, 1967) and at the Meteor Bank (MMF 22841, 1967, set of five specimens, MMF 22878, 1967, one specimen and MMF 22881, 1967, set of two specimens).

Scorpaena maderensis Valenciennes 1833

Common name: **Madeira rockfish; Rascasso-da-Madeira**, Cantarilho, Coça, Papa-jaca, Rascaço, Rescasso, Rocaz.

References: mainland Portugal (de Brito Capello, 1867d; Seabra, 1911; Nobre, 1935(?); Boutière, 1958; Eschmeyer, 1969; Dooley *et al.*, 1985; Abecasis *et al.*, 2009; Carneiro *et al.*, 2014; Machado *et al.*, 2014; OCEANA, 2014; Encarnação *et al.*, 2019), Azores (Collett, 1896a, 1897; Sampaio, 1904; Brauer, 1906; Vaillant, 1919; Norman, 1935b; Fowler, 1936; Albuquerque, 1954-1956; Eschmeyer, 1969; Wood and Williams, 1974; Mapstone and Wood, 1974, 1975; Saldanha, 1977a, 1980, 1995; Blanc and Hureau, 1979; Dooley *et al.*, 1985; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Patzner, 1990;

Patzner and Santos, 1990d; Patzner et al., 1990, 1992; Ré, 1990; Santos et al., 1990b, 1997; Wirtz, 1990a, 1994; Azevedo et al., 1991, 1992; Lloris et al., 1991; Arruda, 1992, 1997; Arruda et al., 1992; Santos, 1992a; Azevedo, 1995; Gonçalves et al., 1995; Harmelin-Vivien et al., 2001; Morato et al., 2001b; Cardigos et al., 2005; Menezes et al., 2006; Floeter et al., 2008; Bertoncini et al., 2010; Botelho et al., 2010; Porteiro et al., 2010; Carneiro et al., 2014; Machado et al., 2014; Afonso, 2016; Poss, 2016; Almeida and Biscoito, 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1833; Lowe, 1837, 1840a, 1843-1860; Günther, 1860; Brauer, 1906; Roule, 1919a; Nobre, 1935; Norman, 1935b; Eschmeyer, 1969; MNHN-IC-0000-6682, 1970; MNHN-IC-0000-6683, 1970; Blanc and Hureau, 1979; Augier, 1985; Dooley et al., 1985; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Lloris et al., 1991; Abreu et al., 1995; Falcón et al., 2000; Mello, 2000; Araújo et al., 2005; Ribeiro et al., 2005b; Floeter et al., 2008; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Machado et al., 2014; POGID, 2016; POGIS, 2016; Poss, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). Eight specimens of *Scorpaena madurensis* Valenciennes 1833, synonym of *S. maderensis*, all caught off in Madeira are deposited in the MNHN collections as syntypes (MNHN-IC-0000-6682 (set of four specimens) and MNHN-IC-0000-6683 (set of four specimens)). These eight specimens were donated by the NHMUK and the collection date is much earlier than the date of determination as syntypes (1970) by Prof. Dr. Wolfgang Klausewitz.

***Scorpaena notata* Rafinesque 1810**

Common name: **Small red scorpionfish; Rascasso-escorpião**, Carneiro, Coça, Matindias, Mete-em-dias, Palhaço, Peixe-carneiro, Requeima, Requeime, Requeimo, Requeme, Rascaço, Rascasso, Rescasso, Rocaz, Roncaz, Serrão.

References: mainland Portugal (Osório, 1896, 1917; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Blanc and Hureau, 1979; Fernández et al., 1981; INIP, 1982a; Borges, 1984; Dinis and Marecos, 1984; Dooley et al., 1985; Hureau and Litvinenko, 1986; Fischer et al., 1987; Martins et al., 1992; Erzini et al., 1996, 1997, 2001a; Gonçalves et al., 1996, 1997; Cabral, 1999; Borges et al., 2002, 2003; Santos et al., 2002a; Mendes et al., 2004; ICES, 2005; Ribeiro et al., 2006, 2012; Fernandes et al., 2007; Henriques et al., 2007, 2008; Gomes and Olim, 2007; Chaves, 2008, 2009; Neves et al., 2008; Silva, 2008, 2010, 2011; Batista et al., 2009; Gonçalves et al., 2010; França et al., 2011; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Costa, 2014; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Collett, 1896a; Vaillant, 1919; Norman, 1935b; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Eschmeyer, 1969; Maul, 1976; Blanc and Hureau, 1979; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Dooley et al., 1985; INIP, 1985a, b; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Patzner, 1990; Patzner and Santos, 1990e; Patzner et al., 1990, 1992; Azevedo et al., 1991, 1992; Dias, 1991; Lloris et al., 1991; Arruda, 1992, 1997; Arruda et al., 1992; Santos, 1992a; Azevedo, 1995; Silva et al., 1995; Menezes, 1996; Santos et al., 1997; Harmelin-Vivien et al., 2001; Morato et al., 2001b; Menezes et al., 2006; Stergiou et al., 2006; Floeter et al., 2008; Bertoncini et al., 2010; Porteiro et al., 2010; Carneiro et al., 2014; Poss, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1840a, 1841, 1849; Günther, 1860; Vinciguerra, 1883; Collett, 1896; Johnson, 1863b; NHMUK 1917.7.14.87, 1912-1936; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Eschmeyer, 1969; maul, 1976; Blanc and Hureau, 1979; Dooley et al., 1985; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Lloris et al., 1991; Floeter et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; POGIS, 2016; Poss, 2016; Eschmeyer et al., 2019). Two specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22865, 1967, set of two specimens). The holotype of *Scorpaena ustulata* Lowe 1841, synonym of *S. notata*, caught off in Madeira is in the NHMUK collections (NHMUK 1917.7.14.87). Wirtz et al. (2008) do not indicate the presence of this species as a coastal fish of Madeira, but confirm its existence only in depths deeper than 100 meters. Froese and Pauly (2019) reported the absence of this species in Madeira due to a misinterpretation of the text of Wirtz et al. (2008).

***Scorpaena plumieri* Bloch 1789**

Common name: **Pacific spotted scorpionfish; Rascasso-negro**, Coça, Rascasso, Rescasso, Rocaz.

References: Azores (?) ([Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Eschmeyer and Demster, 1990](#); [Carneiro et al., 2014](#); [Poss, 2016](#)) and Madeira (?) ([Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Carneiro et al., 2014](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. plumieri* is the Western Atlantic, Ascension and Saint Helena islands. Santos *et al.* (1997) referred that the records of *S. plumieri* to the Azores are misidentifications of *Scorpaena laevis* Troschel 1866. Additionally, Poss (2016) considered that the report of this species from the Azores has been questioned and no materials exist to substantiate its presence there. Thus, this species is not considered in the total number of valid species.

***Scorpaena porcus* Linnaeus 1758**

Common name: **Black scorpionfish; Rascasso-de-pintas**, Cancarro, Carneiro, Coça, Palhaço, Peixe-carneiro, Peixe-porco, Porco, Rascasso, Rascasso-escuro, Rascasso-preto, Requeime, Requeime-preto, Requeimo, Requeme, Rescasso, Rocaz, Roncaz, Sarasco, Serrão.

References: mainland Portugal[▲] ([Vandelli, 1771](#); [Günther, 1860](#); [de Brito Capello, 1867a, d, 1869b, 1880](#); [Steindachner, 1867](#); [Posser, 1880](#); [Osório, 1888](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1895a, c, 1935](#); [Vieira, 1897, 1898](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Blanc and Hureau, 1979](#); [Halstead 1980](#); [Martins, 1981](#); [Hureau and Litvinenko, 1986](#); [Fischer et al., 1987](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [Ribeiro et al., 2006, 2012](#); [Henriques et al., 2007, 2008](#); [Batista et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Poss, 2016](#); [Fricke et al., 2018](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Seabra, 1911](#); [Roule, 1919a](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Eschmeyer, 1969](#); [Blanc and Hureau, 1979](#); [Dooley et al., 1985](#); [Hureau and Litvinenko, 1986](#); [Eschmeyer and Demster, 1990](#); [Lloris et al., 1991](#); [Wirtz, 1994](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Poss, 2016](#); [Fricke et al., 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Osório de Castro, 1954, 1967](#); [Hureau and Litvinenko, 1986](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Wirtz et al., 2008\(?\)](#); [Carneiro et al., 2014](#); [Poss, 2016](#); [Fricke et al., 2018](#); [Froese and Pauly, 2019\(?\)](#)). One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005305).

***Scorpaena scrofa* Linnaeus 1758**

Common name: **Red scorpionfish; Rascasso-vermelho**, Cancarro, Cantaril, Cantarilho, Carneiro, Coça, Palhaço, Papa-jaca, Peixe-carneiro, Porca-marinha, Rainúnculo, Ranúnculo, Rascassa-vermelha, Rascasso, Requeima, Requeime, Requeimo, Requeme, Rescasso, Rocaz, Roncaz, Serrão.

References: mainland Portugal[▲] ([Günther, 1860](#); [de Brito Capello, 1867a, d, 1880](#); [Posser, 1880](#); [Steindachner, 1867](#); [Osório, 1888, 1895, 1896](#); [Vieira, 1893, 1897, 1898](#); [Nobre, 1894, 1895a, 1903, 1935](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Fernández et al., 1981](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Dooley et al., 1985](#); [Hureau and Litvinenko, 1986](#); [Fischer et al., 1987](#); [Mendes et al., 2004](#); [Dinis, 1985](#); [Cabral, 1999](#); [Erzini et al., 2001a](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Silva, 2008](#); [Silva et al., 2009](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Poss, 2016](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Günther, 1860](#); [Drouët, 1861b](#); [Goode and Bean, 1895, 1896](#); [Ramos, 1869](#); [Hilgendorf, 1888b](#); [Roule and Angel, 1921](#); [Sampaio, 1904](#); [Nobre, 1924, 1930](#); [Duncker, 1934](#); [Fowler, 1936b](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Eschmeyer, 1969](#); [Wood, 1973](#); [Wood and Williams, 1974](#); [Wood, 1974](#); [Dias et al., 1976](#); [Maul, 1976](#); [Wheeler, 1978](#); [Blanc and Hureau, 1979](#); [Harmelin and Harmelin-Vivien, 1979](#); [Martins, 1982a](#); [Dooley et al., 1985](#); [INIP, 1985a, c](#); [Dias and Monteiro, 1986](#); [Hureau and Litvinenko, 1986](#); [Azevedo et al., 1991, 1992](#); [Dias, 1991](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Krug, 1995](#); [Silva et al., 1994, 1995](#); [Wirtz, 1994](#); [Gonçalves et al., 1995](#); [Silva and Menezes, 1995a, b](#); [Isidro, 1996](#); [Menezes, 1996](#); [Santos et al., 1997](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001a](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Pham et al., 2013](#); [Carneiro et al., 2014](#); [Diogo and Pereira, 2014](#); [Gomes-Pereira et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Lowe, 1837, 1840a, 1843-1860](#); [Günther, 1860](#); [Goode and Bean, 1895, 1896](#); [Fowler, 1923, 1936b](#); [Norman, 1935b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Eschmeyer, 1969](#); [Maul, 1976](#); [Blanc and Hureau, 1979](#); [Augier, 1985](#); [Dooley et al., 1985](#)).

al., 1985; Hureau and Litvinenko, 1986; Eschmeyer and Demster, 1990; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Ferreira *et al.*, 2008; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; POGID, 2016; Poss, 2016; Priede, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen caught off at the coast of mainland Portugal in 1967 is in the MMF collections (MMF 22467, 1967), a second specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005306).

***Scorpaena stephanica* Cadenat 1943**

Common name: **Spotted-fin rockfish; Rascasso-de-barbatana-pontuada**, Rascasso, Rascasso-pontuado, Rocaz.

References: Madeira (PECS) (NHMUK 2016.3.10.105, 1972; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. stephanica* in the Eastern Atlantic is from Mauritania south to Angola. The northernmost occurrence in the Eastern Atlantic is a preserved specimen deposited in the Natural History Museum (London) (NHMUK 2016.3.10.105) as *Scorpaena gaillardei* Roux 1954, caught off on April 11, 1972 during the research cruise R.R.S. Discovery Cruise, and represents the only historical record of this species from Madeira (in the PECS area). The closest known record occurs in Canary Islands. Probably it has a discontinuous distribution being Madeira the northern limit of its range.

***Scorpaenodes arenai* Torchio 1962**

Common name: Messina rockfish; **Rascasso-de-Messina**.

References: Azores (Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Family Triglidae

***Chelidonichthys cuculus* (Linnaeus 1758)**

Common name: **Red gurnard; Cabra-vermelha**, Antininho, Bebedo, Bêbedo, Bêbedo-ruivo, Bêbo, Bêvo, Bobo, Cabra, Cabrinha, Carriço, Peixe-de-Santo-António, Roberto, Ruivo, Santo-António.

References: mainland Portugal[▲] (Vandelli, 1771; de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Vieira, 1893, 1897, 1898; Nobre, 1894, 1903, 1935; Osório, 1895; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Blanc and Hureau, 1979; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Dooley *et al.*, 1985; Hureau, 1986b; Bauchot, 1987; Martins *et al.*, 1992, 1999, 2015; Gomes *et al.*, 2001; Borges *et al.*, 2002; Mendes *et al.*, 2004; ICES, 2005; Sousa *et al.*, 2005; Olim and Borges, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Batista *et al.*, 2009; Silva, 2011; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; OCEANA, 2014; Richard, 2016; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Regan, 1903a; Fowler, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Dias *et al.*, 1976, 1990; Maul, 1976; Blanc and Hureau, 1979; Fischer *et al.*, 1981; INIP, 1985a, b, c; Hureau, 1986b; Dias and Monteiro, 1986; Richards and Saksena, 1990; Dias, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Silva *et al.*, 1995; Menezes, 1996; Santos *et al.*, 1997; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837; Günther, 1860; Fowler, 1936b; Noronha and Sarmento, 1948; Maul, 1949; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Blanc and Hureau, 1979; Fischer *et al.*, 1981; INIP, 1984a, c; Hureau, 1986b; Richards and Saksena, 1990; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22426, 1967).

***Chelidonichthys lastoviza* (Bonnaterre 1788)**

Common name: **Streaked gurnard; Cabra-riscada**, Bêbedo, Bêbo, Bêdo, Cabra, Cabrinha, Carriço, Roberto, Ruivo.

References: mainland Portugal▲ (Günther, 1860; de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Baldaque da Silva, 1891; Girard, 1892; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Blanc and Hureau, 1979; Fernández et al., 1981; Fischer et al., 1981, 1987; Richards, 1981, 2016; Dooley et al., 1985; Hureau, 1986b; Bauchot, 1987; Richards and Saksena, 1990; Martins et al., 1992, 2015; Erzini et al., 2001a; Borges et al., 2002; Santos et al., 2002a; Palma et al., 2003; ICES, 2005; Olim and Borges, 2006; Stergiou et al., 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Neves et al., 2008; Batista et al., 2009; Chaves, 2009; Silva et al., 2009; Gonçalves et al., 2010; Silva, 2010, 2011; Costa et al., 2012; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Richards, 1968; Blanc and Hureau, 1979; Dooley et al., 1985; Hureau, 1986b; Lloris et al., 1991; Arruda, 1992, 1997; Wirtz, 1994; Gonçalves et al., 1995; Santos et al., 1997; Porteiro et al., 2010; Carneiro et al., 2014; Richard, 2016; Almeida and Bischoff, 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Günther, 1860; Vinciguerra, 1883; Fowler, 1923, 1936b; Maul, 1949; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Blanc and Hureau, 1979; Fischer et al., 1981; Dooley et al., 1985; Hureau, 1986b; Lloris et al., 1991; Abreu et al., 1995; Ribeiro, 2008; Wirtz et al., 2008; Carneiro et al., 2014; POGID, 2016; POGIS, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in April 2011 is in the MB collections (MB06-005340).

Chelidonichthys lucerna (Linnaeus 1758)

Common name: **Tub gurnard; Cabra-cabaço**, Alfondega, Antoninho, Bêbado, Bêbo, Cabação, Cabaço, Caboço, Cabra, Cabra-morena, Cabrinha, Carriço, Peixe-cabra, Peixe-de-Santo-António, Roberto, Ruivo, Santo-António, Surdo.

References: mainland Portugal▲ (Vandelli, 1771; de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895c, 1935; Osório, 1895; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muus and Dahlström, 1978; Blanc and Hureau, 1979; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a, 1982a, b; Martins, 1981; Richards, 1981; Borges, 1982; Dinis and Marecos, 1984; Dinis, 1985; Dooley et al., 1985; Hureau, 1986b; Antunes et al., 1988; Arruda et al., 1988; Richards and Saksena, 1990; Martins et al., 1992, 1999, 2015; Almeida, 1996; Cabral, 1999; Erzini et al., 2001a; Gomes et al., 2001; Borges et al., 2002, 2003; Jorge et al., 2002; Santos et al., 2002a; Palma et al., 2003; Prista et al., 2003; Garnerot et al., 2004; Mendes et al., 2004; ICES, 2005; Olim and Borges, 2006; Ribeiro et al., 2006, 2008; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Neves et al., 2008; Silva, 2008, 2010; Batista et al., 2009; Silva et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Costa et al., 2012; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Richard, 2016; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Fowler, 1923, 1936b; Osório de Castro, 1967(?); Wirtz et al., 2008; Carneiro et al., 2014; Eschmeyer et al., 2019; Froese and Pauly, 2019).

Chelidonichthys obscurus (Walbaum 1792)

Common name: **Longfin gurnard; Cabra-de-bandera**, Cabaço, Cabra, Cabrinha, Carriço, Roberto, Ruivo.

References: mainland Portugal▲ (de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Girard, 1892; Nobre, 1895a, 1903, 1935; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Blanc and Hureau, 1979; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Dooley et al., 1985; Hureau, 1986b; Richards and Saksena, 1990; Martins et al., 1992, 1999, 2015; Erzini et al., 2001a; Borges et al., 2002, 2003; Santos et al., 2002a; Prista et al., 2003; Mendes et al., 2004; ICES, 2005; Moura et al., 2005; Olim and Borges, 2006; Stergiou et al., 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Silva, 2008, 2010; Silva et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Richard, 2016; Martins and Carneiro, 2018; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (?) (Blanc and Hureau, 1979; Fischer et al., 1981(?); Dooley et al., 1985(?); Hureau, 1986b; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997(?); Carneiro et al., 2014; Richard, 2016(?); Eschmeyer et al., 2019(?); Froese and

Pauly, 2019) and Madeira (Osório de Castro, 1967(?); Blanc and Hureau, 1979; Fischer et al., 1981(?); Richards, 1981; Dooley et al., 1985; Hureau, 1986b; Lloris et al., 1991; Wirtz et al., 2008; Carneiro et al., 2014; Richard, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). The occurrence of this species in the Azores was mentioned by several authors, but the records are unconfirmed [Dooley et al. (1985); Richard (2016); Eschmeyer et al. (2019)]. Santos et al. (1997) also considered those records not valid for the Azores. Specimens are needed for confirmation in the Azores Archipelago. Thus, this species is not considered in the total number of valid species in the Azores.

Eutrigla gurnardus (Linnaeus 1758)

Common name: **Grey gurnard; Cabra-morena**, Antoninho, Bacamarte, Bêbedo, Bêbo, Bêvo, Borracho, Cabra, Cabra-moira, Cabra-moura, Carriço, Emprenhador, Peixe-de-Santo-António, Roberto, Ruivo, Santo-António.

References: mainland Portugal▲ (Vandelli, 1771; de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Vieira, 1893, 1897, 1898; Bragança, 1902; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Blanc and Hureau, 1979; Fernández et al., 1981; INIP, 1981a, 1982b; Martins, 1981; Dinis and Marecos, 1984; Dooley et al., 1985; Hureau, 1986b; Fischer et al., 1987; Martins et al., 1992; Cabral, 1999; ICES, 2005; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Olim and Borges, 2006; França et al., 2011; Silva, 2011; Costa et al., 2012; Carneiro et al., 2014; Richard, 2016; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Lloris et al., 1991) and Madeira (Blanc and Hureau, 1979; Hureau, 1986b; Wirtz et al., 2008; Carneiro et al., 2014; Richard, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019).

Lepidotrigla cavillone (Lacepède 1801)

Common name: **Large-scaled gurnard; Ruivo**, Cabra.

References: mainland Portugal (Steindachner, 1867; Bragança, 1902; Pereira, 1915; Osório, 1917; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Blanc and Hureau, 1979, 1986; Fernández et al., 1981; Martins, 1981; INIP, 1982b; Dinis and Marecos, 1984; Dinis, 1985; Bauchot, 1987; Fischer et al., 1987; Erzini et al., 2001a; Gomes et al., 2001; Borges et al., 2002; ICES, 2005; Sousa et al., 2005, 2006; Olim and Borges, 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Ribeiro et al., 2008, 2009; Silva, 2008; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019).

Lepidotrigla dieuzeidei Blanc & Hureau 1973

Common name: **Spiny gurnard; Ruivo-espinhoso**.

References: mainland Portugal (Maul, 1976; Borges et al., 2002; Olim and Borges, 2006; Chaves, 2008, 2009; Henriques et al., 2008; Batista et al., 2009; Silva et al., 2009; Silva, 2010, 2011; Costa et al., 2012; Carneiro et al., 2014; Costa, 2014; EMBL, 2014; Silva and Borges, 2014; Martins and Carneiro, 2018).

Trigla lyra Linnaeus 1758

Common name: **Piper gurnard; Cabra-lira**, Bode, Cabra, Cabra-morena, Cabrinha, Cabrinha-damourama, Carriço, Peixe-cabra, Roberto, Ruivo.

References: mainland Portugal▲ (de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Vieira, 1893, 1897, 1898; Nobre, 1894, 1903, 1935; Osório, 1895; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Blanc and Hureau, 1979; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a, 1982a, b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Hureau, 1986b; Martins et al., 1992, 1999; Borges et al., 2002, 2003; Santos et al., 2002a; Mendes et al., 2004; Moura, 2004; ICES, 2005; Olim and Borges, 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Silva, 2008, 2010, 2011; Batista et al., 2009; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (?) (Sampaio, 1904; Collins, 1954; Osório de Castro, 1967; Arruda, 1997; Santos et al., 1997(?)) and Madeira▲ (Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Blanc and Hureau, 1979; Hureau, 1986b; Richards and Saksena, 1990; Lloris et al., 1991; Wirtz et al., 2008; Carneiro et al., 2014; Richard, 2016; Eschmeyer et

al., 2019; Froese and Pauly, 2019). Santos *et al.* (1997) indicated that the record of *T. lyra* from the Azores is based on a probable misidentification of *Chelidonichthys cuculus* (Linnaeus 1758) by Sampaio (1904). The occurrence of this species in the Azores needs further documentation (Santos *et al.*, 1997; Carneiro *et al.*, 2014). Thus, this species is not considered in the total number of valid species in the Azores.

Family Peristediidae

Peristedion cataphractum (Linnaeus 1758)

Common name: **African armoured searobin**; **Cabra-de-casca**, Armado, Bargela, Bêbedo, Bergela, Berguela, Cabra, Cabra-armada, Cabra-castelhana, Cabra-de-argel, Cabra-de-casco, Cabra-espinhosa, Cabra-francesa, Cabrinha-da-moirama, Cabrinha-da-moirana, Cabrinha-da-mourama, Cabrinha-de-moirama, Cabrinha-de-mourama, Casca, Espinhosa, Peixe-cabra.

References: mainland Portugal (Vandelli, 1771; de Brito Capello, 1867a, d, 1880; Osório, 1888, 1895; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1897, 1898; Bragança, 1902; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Blanc and Hureau, 1979; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; Martins, 1981; Miller, 1981; Hureau, 1986b; Miller and Richards, 1990; Monteiro *et al.*, 2001; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; Silva, 2008, 2010, 2011; Chaves, 2009; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Garcia, 2015e; Richards and Miller, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (IMAR/DOP COLETA 219, 1992; Santos *et al.*, 1997; Carneiro *et al.*, 2014; Martins and Carneiro, 2018) and Madeira (MNHN-IC-2005-2422, 1922; Carneiro *et al.*, 2014; Garcia, 2015e).

Family Cottidae

Myoxocephalus scorpius (Linnaeus 1758)

Common name: **Shorthorn sculpin**; **Escorpião**, Charrasco, Charrôco, Diabo, Peixe-diabo, Peixe-sapo, Sapo, Saranco, Sarronca, Sarrouco, Xarroco.

References: mainland Portugal (Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro *et al.*, 2014) and Azores (?) (Osório de Castro, 1967). According to Eschmeyer *et al.* (2019) the current geographic distribution of *M. scorpius* is circumpolar, Northern Hemisphere (including White Sea, Baltic Sea, North Sea). The presence of this species off the Portuguese mainland is considered rare (Nobre, 1935) and only historical records are known. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location. The most southern occurrence record in the Atlantic ocean, out of the current geographic distribution area, occurred at 26°28.8'N, 52°50.4'W (DFO-NFLD 31984.SHSCUL, 1980). Specimens are needed for confirmation in the Azores. Thus, this species is not considered in the total number of valid species in the Azores.

Taurulus bubalis (Euphrasen 1786)

Common name: Longspined bullhead; **Escorpião-roco**, Diabo, Escorpião, Padre, Peixe-diabo, Peixe-roco, Peixe-sapo, Peixe-sapo-da-ribeira, Roco, Sapo, Sapo-da-ribeira, Saranco, Sarronca, Sarrouco, Xarroco.

References: mainland Portugal (Osório, 1888, 1895, 1917; Vieira, 1897, 1898; Nobre, 1903, 1935; Seabra, 1911; Vinciguerra, 1923; Gonçalves, 1941; Helling, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954; Neyelov, 1979; Fedorov, 1986; Fischer *et al.*, 1987; Arruda, 1990; Garnerot *et al.*, 2004; Henriques *et al.*, 2007, 2008; Carneiro *et al.*, 2014; Lorance *et al.*, 2014; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Family Psychrolutidae

Cottunculus thomsonii (Günther 1882)

Common name: Pallid sculpin; **Escorpião-pálido**.

References: mainland Portugal (Fedorov and Nelson, 1986; Nelson, 1990; Carneiro *et al.*, 2014; Florin *et al.*, 2015; Froese and Pauly, 2019).

Family Cyclopteridae

Cyclopterus lumpus Linnaeus 1758

Common name: **Lumpfish; Peixe-lapa**, Lapa.

References: mainland Portugal ([Nobre, 1935](#); [Albuquerque, 1954–1956](#); [Almaça, 1965a](#); [Osório de Castro, 1967](#); [Lindberg, 1979](#); [Stein, 1986a](#); [Vasconcelos et al., 2004](#); [Bañón et al., 2008](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. lumpus* in the Eastern Atlantic is the Baltic Sea, North Sea, North Atlantic and the Arctic Ocean. This species is of rare occurrence in southern waters and the southernmost occurrence record includes catches off southern Portugal.

Eumicrotremus spinosus (Fabricius 1776)

Common name: Atlantic spiny lump sucker; **Peixe-lapa-espinhoso**, Lapa, Peixe-lapa.

References: Azores ([Collett, 1897, 1905](#); [Arruda, 1992, 1997](#); [Santos et al., 1997\(?\)](#); [Carneiro et al., 2014](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *E. spinosus* is the Arctic Ocean and the North Atlantic.

Family Liparidae

Liparis liparis (Linnaeus 1766)

Common name: Striped seasnail; **Peixe-lesma-listado**.

References: mainland Portugal ([Chernova et al., 2004](#)).

Liparis montagui (Donovan 1804)

Common name: **Montagus seasnail; Peixe-lesma-de-Montagus**.

References: mainland Portugal ([Fage, 1918](#); [Lindberg, 1979](#); [Arruda, 1990](#); [Florin et al., 2014](#)).

Paraliparis copei Goode & Bean 1896

Common name: Blacksnout seasnail; **Peixe-lesma-de-focinho-preto**.

References: Azores ([Stein, 1986b, 2016 \(PECS\)](#); [ARC 26932, 1995](#); [Arruda, 1997](#); [Carneiro et al., 2014](#); [Mecklenburg et al., 2018](#); [Froese and Pauly, 2019](#)).

Paraliparis membranaceus Günther 1887

Common name: No English common name; **Peixe-lesma-membranoso**.

References: mainland Portugal (?) ([Günther, 1887](#); [Burke, 1930](#); [Nobre, 1935](#); [Albuquerque, 1954–1956](#); [Lindberg, 1979](#); [Stein and Able, 1986](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. membranaceus* is the Eastern South Pacific. The *P. membranaceus* record off mainland Portugal is erroneous ([Stein, 2005](#); [Carneiro et al., 2014](#)). Nobre (1935), Albuquerque (1954–1956), Lindberg (1979) and Stein and Able (1986) reported the occurrence of this species in mainland Portugal based on Günther (1887) and Burke (1930). Thus, this species is not considered in the total number of valid species.

Paraliparis nigellus Chernova & Møller 2008

Common name: No English common name; **Peixe-lesma-escuro**.

References: Azores ([Chernova and Møller, 2008](#); [MAR-ECO 2619, 2004](#); [MAR-ECO 2633, 2004](#); [Carneiro et al., 2014](#); [Biscoito, 2015f](#); [Porteiro et al., 2017](#)).

Order Centrarchiformes

Family Kyphosidae

Kyphosus bigibbus Lacepède 1801

Common name: **Brown chub; Preguiçosa-cinzenta**, Preguiçosa.

References: mainland Portugal ([Knudsen and Clements, 2013\(?\)](#); [Knudsen et al., 2015h](#)) and Madeira ([Knudsen and Clements, 2013](#); [Sakai and Nakabo, 2014, 2016](#); [Knudsen et al., 2015h](#)). According to Eschmeyer *et al.* (2019) *Kyphosus bosquii* (Lacepède 1802) is synonym of *Kyphosus bigibbus* Lacepède 1801 and its current geographic distribution is the Atlantic (if valid). However, Sakai and Nakabo (2014) have a different view of the taxonomy of the Atlantic species of *Kyphosus* and consider *K. bosquii* to occur in the Atlantic Ocean.

Kyphosus sectatrix (Linnaeus 1758)

Common name: **Bermuda sea chub; Preguiçosa-branca**, Choupa, Preguiçosa, Patruça, Salema-do-Brasil, Sopa.

References: mainland Portugal (Canas *et al.*, 2005; Francour and Mouine, 2008; de Casamajor and Morandieu, 2013; Carneiro *et al.*, 2014; OCEANA, 2014; Bañón *et al.*, 2017a; Froese and Pauly, 2019), Azores[▲] (Murray and Hjort, 1912; Fowler, 1936b; Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1967; Harmelin and Harmelin-Vivien, 1979; Patzner *et al.*, 1990, 1992; Ré, 1990; Santos *et al.*, 1990b, 1997; Wirtz, 1990a, 1994; Arruda *et al.*, 1992; Santos, 1992a; Azevedo, 1995; Gonçalves *et al.*, 1995; Arruda, 1997; Harmelin-Vivien *et al.*, 2001; Floeter *et al.*, 2008; Francour and Mouine, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Afonso, 2016; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Günther, 1859; Norman, 1935b; Fowler, 1936b; Noronha and Sarmento, 1948; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Desoutter, 1979; Fischer *et al.*, 1981; Tortonese, 1986c; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Wirtz, 1994; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; de Casamajor and Morandieu, 2013; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Friedlander *et al.*, 2017; Froese and Pauly, 2019). Knudsen and Clements (2013) reported it is possible that the records of *K. sectatrix* at the Azores (Santos *et al.*, 1997) could have included the occasional *K. bigibbus*. On the other hand, according to the former authors the specimen reported by Canas *et al.* (2005) from the Algarve coast was identified as *K. sectator* (i.e. older name *K. sectatrix*), but its photo resembles *K. bigibbus*.

Kyphosus vaigiensis (Quoy & Gaimard 1825)

Common name: **Brassy chub; Preguiçosa-bronzeada**, Preguiçosa. [Yellow sea chub; Preguiçosa-amarela, Preguiçosa, Patruça, Salema-do-Brasil (as *Kyphosus incisor* (Cuvier 1831))].

References: mainland Portugal (Knudsen *et al.*, 2015i; Bañón *et al.*, 2019b), Azores (Patzner *et al.*, 1992; Arruda, 1997; Santos *et al.*, 1997; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Knudsen *et al.*, 2015i) and Madeira (Fischer *et al.*, 1981; Tortonese, 1986c; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Diogo and Pereira, 2013; Carneiro *et al.*, 2014; Sakai and Nakabo, 2014, 2016; Knudsen *et al.*, 2015i). According to Knudsen and Clements (2013) and Eschmeyer *et al.* (2019) *Kyphosus incisor* (Cuvier 1831) (in the Atlantic) is synonym of *K. vaigiensis*. The current geographic distribution includes Atlantic Ocean (Eschmeyer *et al.*, 2019).

Order Acropomatiformes

Family Polyprionidae

Polyprion americanus (Bloch & Schneider 1801)

Common name: **Wreckfish; Cherne**, Charnota, Cherna, Cherna-de-pau, Cherna-de-pau, Cherne-americano, Cherne-galego, Chernota, Mero, Mero-francês, Pardete, Pardil, Pardilho, Romeiro.

References: mainland Portugal[▲] (Carvalho, 1765(?); de Brito Capello, 1867a, d, 1869b, 1880; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1935; Richard, 1908; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979b, 1986a; Fischer *et al.*, 1981, 1987; Erzini *et al.*, 2001b; Santos *et al.*, 2002b, c; ICES, 2005; Fernandes *et al.*, 2007; Pakhorukov, 2008; Chaves, 2009; Silva, 2011; Carneiro *et al.*, 2014; Collette *et al.*, 2015h; Bañón *et al.*, 2016a; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Sampaio, 1904; Murray and Hjort, 1912; Fowler, 1936b; Ribeiro, 1936; Cadenat, 1937; Koefoed, 1952; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood, 1974; Dias *et al.*, 1976, 1990; Wheeler, 1978; Tortonese, 1979b; Muzanor and Westhaus, 1981; Martins, 1982a; INIP, 1985a, b, c; Silva, 1986a, b, 1992; Dias, 1991; Arruda, 1992, 1997; Silva and Krug, 1992; Santos *et al.*, 1994a, 1995a, 1997; Silva *et al.*, 1994, 1995; Krug, 1995; Pinho and Pereira, 1995b; Sedberry, 1995, 2002; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Sedberry *et al.*, 1996; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Hareide and Garnes, 2001 (PECS); Morato *et al.*, 2001a; Kukuev, 2002 (PECS); Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Porteiro *et al.*, 2010,b, 2013, 2017; Besugo, 2013; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Fauconnet *et al.*, 2019) and Madeira[▲] (Günther, 1859; Lowe, 1841, 1843-1860; de Velloso, 1892; Goode and Bean, 1895, 1896; Collett, 1896; Fowler, 1936b; Noronha and

Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979b; Fischer *et al.*, 1981; INIP, 1984c; Severino, 2004; Ferreira *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Collette *et al.*, 2015h; POGID, 2016; Roberts 2016; Biscoito *et al.*, 2017; Froese and Pauly, 2019).

Family Epigonidae

Epigonus constanciae (Giglioli 1880)

Common name: Constance deepwater cardinalfish; **Olhudo-de-espinho**.

References: mainland Portugal (OCEANA, 2014), Azores (Santos *et al.*, 1997; Sutton *et al.*, 2008; Porteiro *et al.*, 2010, 2017; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015d) and Madeira (Hureau and Monod, 1979; Tortonese, 1986b; Maugé and Mayer, 1990; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015d; Gon, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Epigonus denticulatus Dieuzeide 1950

Common name: **Pencil cardinal**; **Olhudo-denteado**.

References: mainland Portugal (Saldanha *et al.*, 1995; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; Chaves, 2008; Carneiro *et al.*, 2014; Costa, 2014; Bañón *et al.*, 2016a), Azores (PECS) (Hareide and Garnes, 2001 (PECS); Carneiro *et al.*, 2014).

Epigonus telescopus (Risso 1810)

Common name: **Black cardinal fish**; **Olhudo**, Besouro, Besouro, Boca-negra, Diabo, Escamuda, Escamuda-preta, Escuro, Jordão, Olhuda, Olhudo; Peixe-diabo, Peixe-sabão, Peixe-sardão, Pimentel, Preto, Ribaldo-preto, Robaldo, Robaldo-do-alto, Robaldo-preto, Robalo, Robalo-do-alto, Robalo-preto, Sabão, Salmonete, Salmonete-da-fundura, Salmonete-de-fundura, Salmonete-preto, Sardão.

References: mainland Portugal▲ (de Brito Capello, 1867a, d, e, 1880; Steindachner, 1867; Posser, 1880; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Goode and Bean, 1895, 1896; Bragança, 1903; Osório, 1909; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979a, 1986b; Fernández *et al.*, 1981; Figueiredo *et al.*, 1994; Borges *et al.*, 2002; Moura, 2004; ICES, 2005; Fernandes *et al.*, 2007; Pakhorukov, 2008; Silva, 2008; Bordalo-Machado *et al.*, 2009; Carneiro *et al.*, 2014; OCEANA, 2014; Garcia, 2015f; Bañón *et al.*, 2016a; Martins and Carneiro, 2018), Azores▲ (Koefoed, 1952; Collins, 1954; Scherbachev *et al.*, 1985 (PECS); Santos *et al.*, 1995a, 1997; Silva *et al.*, 1995; Menezes, 1996; Saldanha and Biscoito, 1997; Arruda, 1997; Desbruyères *et al.*, 2001; Hareide and Garnes, 2001 (PECS); Kukuev, 2002 (PECS); Melo and Menezes, 2002; Menezes *et al.*, 2006, 2009, 2012; Martins *et al.*, 2006; Rosa *et al.*, 2006; Pakhorukov, 2008 (PECS); Porteiro *et al.*, 2010; Besugo, 2013; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Garcia, 2015f) and Madeira▲ (Lowe, 1837, 1843; Steindachner, 1867; de Brito Capello, 1871; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Nunes, 1953; Tortonese, 1979a; INIP, 1982c, 1984a, b; Maugé and Mayer, 1990; Lloris *et al.*, 1991; Severino, 2004; Carneiro *et al.*, 2014; Garcia, 2015f; Gon, 2016; POGID, 2016; Delgado *et al.*, 2017; Froese and Pauly, 2019).

Microichthys coccoi Rüppell 1852

Common name: Dwarf deepwater cardinalfish; **Olhudo-anão**.

References: Azores (MCZ 149629, 1978; MCZ 149630, 1978; Arruda, 1997; Porteiro *et al.*, 1999, 2010, 2017; Sutton *et al.*, 2008; Carneiro *et al.*, 2014; Gon, 2016; Priede, 2017; Ordines *et al.*, 2018b; Eschmeyer *et al.*, 2019).

Family Howellidae

Howella atlantica Post & Quéro 1991

Common name: **Pelagic basslet**; **Peixe-alado-do-Atlântico**.

References: mainland Portugal (MCZ 86138, 1972(?); Carneiro *et al.*, 2014), Azores (MCZ 149545, 1978; ZMH 25084, 1979 (PECS); ZMH 25086, 1982 (PECS); Santos *et al.*, 1997; Porteiro *et al.*, 1999, 2010, 2017; Carneiro *et al.*, 2014; Heemstra, 2016c; Froese and Pauly, 2019) and Madeira (Noronha and Sarmento, 1948; Tortonese, 1979a, 1986a; Carneiro *et al.*, 2014; Heemstra, 2016c; Froese and Pauly, 2019). The current geographic distribution of *H. atlantica* is in the Atlantic (Eschmeyer *et al.* 2019). According to

Heemstra (2016d) this species is frequently misidentified as *Howella sherborni* (non Norman 1930) or *Howella brodiei* (non Ogilby 1899). The specimen MCZ 86138 was identified as *Howella brodiei* Ogilby 1899. It is recommended the re-evaluation of these specimens identification. The records of *H. brodiei* in the northeast Atlantic should probably be considered as belonging to *H. atlantica* (Carneiro *et al.*, 2014).

***Howella brodiei* (Ogilby 1899)**

Common name: **Pelagic basslet; Peixe-alado-de-Brodie.**

References: mainland Portugal (?) (MCZ 86138, 1972; ZSM-Pis-029105 / 534316 / 385904, 1992; ZSM-Pis-029212 / 534417 / 386005, 1992), Azores (?) (ZMH 7125, 1970 (PECS); ZMH 117143, 1979 (PECS); ZMH 117276, 1979 (PECS); MAR-ECO 7276, 2004; MAR-ECO 7425, 2004; MAECO153-06, 2006; MAECO154-06, 2006; Sutton *et al.*, 2008; Porteiro *et al.*, 2010) and Madeira (?) (Noronha and Sarmento, 1948; ZMH 103325, 1966; MCZ 86142, 1969; MCZ 86162, 1972; Tortonese, 1979a). The specimens deposited in the Natural History Collections of the University of Bergen (MAR-ECO), Museum of Comparative Zoology of the Harvard University (MCZ), Zoological Museum of the Universität Hamburg (ZMH), and in the Staatliche Naturwissenschaftliche Sammlungen Bayerns (ZSM) were not identified to the subspecies level. Two specimens of the genus *Howella* caught off in Azores were identified as *H. brodiei* and their vouchers (MAECO) are in the GenBank. However, according to Eschmeyer *et al.* (2019) the current geographic distribution of *Howella brodiei* (Ogilby 1899) is the Indo-Pacific: South Africa east to Hawaiian Islands and California (U.S.A.), north to southern Japan, south to northern Australia. This geographic distribution is in contradiction with the records and references of this species in the north-west and north-east Atlantic. Thus, it is recommended the re-evaluation of these specimens' identification and consequently this species is not considered in the total number of valid species.

***Howella sherborni* (Norman 1930)**

Common name: **Pelagic basslet; Peixe-alado-de-Sherborn.**

References: Azores (PECS) (?) (NHMUK 2015.7.9.249, 1981; Carneiro *et al.*, 2014) and Madeira (?) (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Tortonese, 1986a; Lloris *et al.*, 1991; Carneiro *et al.*, 2014). The specimens reported for the Azores and Madeira could have been misidentified according to the circumglobal geographic distribution of this species in the southern temperate seas, as reported by Post and Quéro (1991), Heemstra (2016c) and Eschmeyer *et al.* (2019). However, the specimen with the occurrence record NHMUK 2015.7.9.249 was confirmed by genetic analyses as *H. sherborni*, such as other three specimens caught off in the North-western Atlantic. Thus, it is recommended the re-evaluation of this specimen identification and consequently this species is not considered in the total number of valid species.

***Howella simplex* (Parr 1933)**

Common name: No English common name; **Peixe-alado.**

References: mainland Portugal (?) (YPM ICH 005179, 1959; YPM ICH 005180, 1959; YPM ICH 005181, 1959; YPM ICH 005182, 1959; YPM ICH 005183, 1959; YPM ICH 005184, 1959; YPM ICH 005185, 1959; YPM ICH 005186, 1959). This species is sometimes confused with *Howella brodiei* Ogilby 1899 (Heemstra, 2016c). These specimens were identified as *Bathysphyrænops simplex* Parr 1933 [synonyms of *H. simplex* (Parr 1933)] and are preserved in the fishes collections in the Yale Peabody Museum's Division of Vertebrate Zoology. However, these specimens could have been misidentified according to the circumglobal geographic distribution of this species in the tropical and subtropical Western Atlantic, as reported by Heemstra (2016c). Thus, it is recommended the re-evaluation of these specimens identification and consequently this species is not considered in the total number of valid species.

Order Perciformes

Family Moronidae

***Dicentrarchus labrax* (Linnaeus 1758)**

Common name: **European seabass; Robalo-legítimo**, Cachaço, Cachote, Chalizo, Cornilha, Corninha, Corninho, Cutelo, Peixe-branco, Robailo, Robalete, Robaldinho, Robaldo, Robalete, Robalizo, Robalinho, Robalo, Robalo-marisco, Robalote, Solhado, Tareco, Xale.

References: mainland Portugal▲ ([Carvalho, 1756](#); [de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1893, 1894, 1895a, c, 1903, 1935](#); [Vieira, 1893, 1897, 1898](#); [Osório, 1895](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979b, 1986a](#); [Fischer et al., 1981, 1987](#); [Smith, 1981](#); [Bauchot, 1987](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Almeida, 1996](#); [Gonçalves et al., 1996, 1997](#); [Erzini et al., 2001a](#); [Jorge et al., 2002](#); [Palma et al., 2003](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Henriques et al., 2007, 2008](#); [Gomes and Olim, 2007](#); [Neves et al., 2008](#); [Ribeiro et al., 2008, 2012](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Pollard, 2015a](#); [Ryland and Lancaster, 2017](#); [Golani and Fricke, 2018](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Pollard, 2015](#); [Heemstra, 2016d](#)).

Dicentrarchus punctatus (Bloch 1792)

Common name: **Spotted seabass**; **Robalo-baila**, Alvaira, Avaira, Avária, Baila, Baile, Bailadeira, Balha, Balhadeira, Cachaço, Cachote, Chaliço, Cornilha, Corninha, Corninho, Cutelo, Peixe-branco, Pinta, Pintainha, Pintainho, Robaldete, Robaldinho, Robaldo, Robalete, Robaliço, Robalo, Robalo-de-pintas, Robalo-pinta, Robalote, Solhado, Tareco, Vaila, Vaira, Vairão, Varia, Vária, Variaz, Xale.

References: mainland Portugal▲ ([de Brito Capello, 1867a, d, e, 1869b, 1880; Steindachner, 1867](#); [Osório, 1888](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1895a, 1903, 1935](#); [Vieira, 1897, 1898](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Blanc et al., 1971](#); [Fischer et al., 1981, 1987](#); [Tortonese, 1986a](#); [Gonçalves et al., 1995, 1996, 1997](#); [Garnerot et al., 2004](#); [ICES, 2005](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Ribeiro et al., 2008](#); [Carneiro et al., 2014](#); [Craig, 2015a](#); [Martins et al., 2015](#); [Golani and Fricke, 2018](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Bowdich, 1825](#); [Fowler, 1936b](#); [Osório de Castro, 1967](#); [Fischer et al., 1981\(?\)](#); [Craig, 2015a](#)).

Family Serranidae

Anthias anthias (Linnaeus 1758)

Common name: **Swallowtail seaperch**; **Canário-do-mar**, Andorinha, Barbeiro, Canário, Capitão, Castanheta, Castanheta-amarela, Castanheta-do-alto, Folião, Folião-do-alto, Imperador, Imperador-do-alto, Infante, Inzempo, Papagaio-do-mar, Pedador, Pedadore, Periquito, Piriquito.

References: mainland Portugal▲ ([de Brito Capello, 1867a, d, 1880; Steindachner, 1867](#); [Osório, 1895](#); [Vieira, 1897, 1898](#); [Bragança, 1903](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Tortonese, 1979b, 1986a](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982b](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Martins et al., 1992](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [Gonçalves et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2006](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Pakhorukov, 2008](#); [Abecasis et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [OCEANA, 2014](#); [Heemstra and Anderson \(Jr.\), 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores▲ ([Drouët, 1861b](#); [Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Nobre, 1924, 1930](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Dias et al., 1976](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a](#); [Tortonese, 1979b, 1986a](#); [Saldanha, 1980, 1995](#); [Martins, 1982b](#); [Dooley et al., 1985](#); [Dias and Monteiro, 1986](#); [Barreiros, 1990, 1995](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1997](#); [Wirtz, 1990a, 1992, 1994](#); [Dias, 1991](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Martins et al., 1992](#); [Azevedo, 1995](#); [Silva and Menezes, 1995a](#); [Silva et al., 1995](#); [Menezes, 1996](#); [Morato-Gomes et al., 1998](#); [Sousa et al., 1999](#); [Harmelin-Vivien et al., 2001](#); [Kukuev, 2002 \(PECS\)](#); [Menezes et al., 2006](#); [Floeter et al., 2008](#); [Pakhorukov, 2008 \(PECS\)](#); [Porteiro et al., 2010, 2013](#); [Gomes-Pereira et al., 2012, 2014](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Heemstra and Anderson \(Jr.\), 2016](#); [Almeida and Biscoito, 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira▲ ([Bowdich, 1825](#); [Lowe, 1837, 1843-1860](#); [Günther 1859, 1880](#); [MNHN-IC-0000-4316, 1868](#); [Guichenot, 1868](#); [de Brito Capello, 1871](#); [NHMUK 1890.5.31.2-3, 1878-1892](#); [Johnson, 1890](#); [Zugmayer, 1911b](#); [Roule, 1919a \(PECS\)](#); [Fowler, 1923, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [Tortonese, 1979b](#); [Fischer et al., 1981](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#);

Wirtz *et al.*, 2008; Christiansen *et al.*, 2009; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Heemstra and Anderson (Jr.), 2016; POGID, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Two syntypes of *Aylopon canariensis* Guichenot 1868 and *Anthias mundulus* Johnson 1890, synonyms of *A. anthias*, caught off in Madeira are in the collections of the MNHN (MNHN-IC-0000-4316, set of two specimens) and of the NHMUK (NHMUK 1890.5.31.2-3, set of two specimens), respectively. One specimen caught off at the Portuguese continental slope in October 2009 is in the MB collections. Several specimens caught off at the Josephine Bank and Meteor Bank are in the MMF collections [MMF 22406, 1967, set of sixty one specimens (Josephine Bank) and MMF 22371, 1967, set of nine specimens, MMF 22840, 1967, one specimen, MMF 22853, 1967, one specimen and MMF 22873, 1967, one specimen, from the Meteor Bank].

Cephalopholis nigri (Günther 1859)

Common name: **Niger hind; Garoupa-do-Níger**; Garoupa-negra; Garoupa-preta.

References: Madeira (Heemstra & Anderson (Jr.) 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. nigri* is the Mediterranean Sea (Malta) and, in the Eastern Atlantic, from Canary Islands, Senegal to Angola, including islands in the Gulf of Guinea. The distribution map of this species reported by Heemstra & Anderson (Jr.) (2016) partially included the area of Selvagens Islands (Madeira EEZ). Specimens are needed for confirmation in Madeira Archipelago.

Epinephelus aeneus (Geoffroy St. Hilaire 1817)

Common name: **White grouper; Garoupa-legítima**, Baixa-bedaque, Cherna, Cherne, Garoupa, Garoupa-de-Portugal, Garoupa-verde, Garupa, Merlo, Mero, Murianga, Nero, Peixe-conta.

References: mainland Portugal[▲] (Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981, 1987; Tortonese, 1986a; Heemstra and Randall, 1993; Carneiro *et al.*, 2014; Craig, 2015b; Heemstra and Anderson (Jr.), 2016; Bañón *et al.*, 2017b; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Tortonese, 1979a; Lloris *et al.*, 1991).

Epinephelus adscensionis (Osbeck 1765)

Common name: **Rock hind; Garoupa-pintada-da-rocha**, Garoupa.

References: Azores (Craig *et al.*, 2011; Ferreira *et al.*, 2018; Froese and Pauly, 2019). According to Eschmeyer *et al.* (2019) the geographic distribution of *E. adscensionis* in the Eastern Atlantic is restricted to the islands in Gulf of Guinea.

Epinephelus caninus (Valenciennes 1843)

Common name: **Dogtooth grouper; Mero-gigante**, Cherna, Cherna-da-costa, Cherne, Garoupa, Garoupa-gigante, Garoupa-grande, Garoupa-negra, Garoupa-preta, Garupa, Mera, Merlo, Mero, Mero-bicudo, Negro, Nero, Pica-pau.

References: mainland Portugal (Vinciguerra, 1883; Osório de Castro, 1967; Tortonese, 1986a; Fischer *et al.*, 1987; Heemstra, 1991; Heemstra and Randall, 1993; Carneiro *et al.*, 2014; Craig and Pollard, 2015a; Froese and Pauly, 2019), Azores (Osório de Castro, 1967) and Madeira (Osório de Castro, 1967).

This species is sometimes confused with *Epinephelus alexandrinus* (Valenciennes 1828) (Heemstra & Anderson (Jr.), 2016). In mainland Portugal it is a rare species according to Osório de Castro (1967). This author also mentions the presence of this species in the fishing grounds of Madeira and the Azores without indicating its geographic location. Specimens are needed for confirmation in the Azores.

Epinephelus costae (Steindachner 1878)

Common name: **Goldblotch grouper; Garoupa-dourada**.

References: mainland Portugal (Fischer *et al.*, 1987; Heemstra and Randall, 1993; Carneiro *et al.*, 2014; Craig and Pollard, 2015b; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019); Azores (?) (Wood, 1973, 1974; Wood and Williams, 1974; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Saldanha, 1980; Wirtz, 1990a; Martins, and Santos, 1991; Patzner *et al.*, 1990; Santos, 1992b) and Madeira (?) (Heemstra and Randall, 1993; Craig and Pollard, 2015b). This species was named *Epinephelus alexandrinus* (Valenciennes 1828) from 1895 to 1988 (Craig and Pollard, 2015b). Currently *Epinephelus alexandrinus* (non Valenciennes 1828) is synonym of *E. costae* and *Epinephelus alexandrinus* (Valenciennes 1828) is synonym of *Epinephelus fasciatus* (Forsskål 1775). Several authors mention the occurrence of this species in the Azores and Madeira on the basis of misidentification of *Mycterooperca fusca* (Lowe 1838)

(Santos *et al.*, 1997; Craig and Pollard, 2015b; Froese and Pauly, 2019). Thus, this species is not considered in the total number of valid species for Madeira and the Azores.

Epinephelus goreensis (Valenciennes 1830)

Common name: **Dungat grouper; Mero-amarelo**, Canário, Cherna, Cherne, Engenhim, Garoupa, Garoupa-amarela, Garoupa-branca, Garoupa-castanha, Garoupa-de-bigode, Garoupa-negra, Garoupa-preta, Garupa, Mera, Merlo, Mero, Mero-castanho, Nero, Nero-amarelo, Nero-castanho, Ruta, Ruta-branca, Sabão.

References: mainland Portugal (†) ([de Brito Capello, 1867a, 1880](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#)). This species is sometimes confused with *Epinephelus alexandrinus* (non Valenciennes 1628) (Heemstra & Anderson (Jr.), 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *E. goreensis* is from Mauritanea to Angola, including Cape Verde Islands and islands of the Gulf of Guinea. Osório de Castro (1967) considered it a rare species and his reference is the last historical record known for mainland Portugal, and probably it is regionally extinct. Thus, this species is not considered in the total number of valid species.

Epinephelus guaza (Linnaeus 1758) - *Nomen dubium* in Mycteroptera

Some authors indicated *E. guaza* as synonym of *Epinephelus marginatus* (Lowe 1834) but others considered it as of uncertain status in Mycteroptera, Serranidae: Epinephelinae. The occurrence records of this species in PECS, mainland Portugal, Azores and Madeira are assumed to belong to the species *E. marginatus*. Thus, this species is not considered in the total number of valid species.

Epinephelus marginatus (Lowe 1834)

Common name: **Dusky grouper; Mero-legítimo**, Cachorro, Cherna, Cherna-da-costa, Cherne, Garoupa, Garoupa-negra, Garoupa-preta, Garupa, Mera, Merato, Merlo, Mero, Mero-de-Cabo-Verde, Mero-preto, Negro, Nero.

References: mainland Portugal▲ ([de Brito Capello, 1867c, d, e, 1880](#); [Posser, 1880](#); [Osório, 1888](#); [Baldaque da Silva, 1891](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Vieira, 1897, 1898](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1940, 1943](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979b\(?\)](#); [Fischer et al., 1981, 1987](#); [Heemstra and Randall, 1993](#); [Gomes and Olim, 2007](#); [Carneiro et al., 2014](#); [Harmelin-Vivien and Craig, 2015](#); [Heemstra and Anderson \(Jr.\), 2016](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores▲ ([Hilgendorf, 1888a](#); [Harmelin and Harmelin-Vivien Sampaio, 1904](#); [Fowler, 1936b](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Tortonese, 1954, 1979b\(?\)](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Wood, 1973](#); [Wood and Williams, 1974](#); [Wood, 1974](#); [Dias et al., 1976](#); [Wheeler, 1978](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Saldanha, 1980, 1992](#); [Martins, 1982a](#); [Bruslé, 1985](#); [Dooley et al., 1985](#); [INIP, 1985c](#); [Skaramuka et al., 1989](#); [Patzner, 1990](#); [Patzner and Santos, 1990c](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990a, 1994a, b, 1995a, 1997](#); [Wirtz, 1990a, 1994](#); [Azevedo et al., 1991, 1992, 1995](#); [Dias, 1991](#); [Heemstra, 1991](#); [Lloris et al., 1991](#); [Martins and Santos, 1991](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Azevedo, 1992](#); [Santos, 1992a, b](#); [Heemstra and Randall, 1993](#); [Gonçalves, 1993](#); [Heemstra and Randall, 1993](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Silva et al., 1995](#); [Isidro, 1996](#); [Menezes, 1996](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001a](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Diogo and Pereira, 2013, 2014](#); [Carneiro et al., 2014](#); [Harmelin-Vivien and Craig, 2015](#); [Afonso, 2016](#); [Heemstra and Anderson \(Jr.\), 2016](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira▲ ([Lowe, 1833a, 1837, 1838](#); [Günther, 1859](#); [de Brito Capello, 1871](#); [Steindachner, 1876](#); [Vinciguerra, 1883](#); [de Velloso, 1892](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Tortonese, 1979b\(?\)](#); [Augier, 1985](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Heemstra and Randall, 1993](#); [Abreu et al., 1995](#); [Delgado, 1998](#); [Falcón et al., 2000](#); [Mello, 2000](#); [Araújo et al., 2005](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Harmelin-Vivien and Craig, 2015](#); [Heemstra and Anderson \(Jr.\), 2016](#); [POGID, 2016](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Epinephelus gigas* (Brünnich 1768) (Heemstra & Anderson (Jr.), 2016). One specimen of *Serranus marginatus* Lowe 1834, synonym of *E. marginatus*, caught off in Madeira is in the MMF collections (MMF 3388) and labelled as neotype. The specimen identified by de Brito Capello (1867) as *Serranus cernioides* is uncertain as *Epinephelus marginatus* (Lowe 1834) (Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Hyporthodus haifensis* (Ben-Tuvia 1953)**

Common name: **Haifa grouper; Garoupa-de-Haifa.**

References: mainland Portugal ([Fischer et al., 1987\(?\)](#); [Eschmeyer et al., 2019](#)). This species is sometimes confused with *Epinephelus marginatus* (Lowe 1834) ([Heemstra & Anderson \(Jr.\), 2016](#)).

***Mycteroperca fusca* (Lowe 1838)**

Common name: **Island grouper; Garoupa-insular**, Badejo, Badeijo, Badeija.

References: Azores[▲] ([Wood, 1973](#); [Wood and Williams, 1974](#); [Wood, 1974](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a](#), [1990](#); [Saldanha, 1980](#), [1995](#); [Martins, 1982a](#); [Patzner et al., 1990](#), [1992](#); [Wirtz, 1990a](#), [1994](#); [Azevedo et al., 1991](#); [Heemstra, 1991](#); [Martins, and Santos, 1991](#); [Arruda et al., 1992](#); [Santos, 1992b](#); [Heemstra and Randall, 1993](#); [Santos et al., 1994a](#), [1995a](#), [1997](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Arruda, 1997](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001a](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Diogo and Pereira, 2013](#); [Carneiro et al., 2014](#); [Heemstra and Anderson \(Jr.\), 2016](#); [Froese and Pauly, 2019](#); [Eschmeyer et al., 2019](#) and [Madeira \(Lowe, 1837, 1838; de Brito Capello, 1871; Vinciguerra, 1883; Fowler, 1936b; Heemstra, 1991; Heemstra and Randall, 1993; Abreu et al., 1995; Delgado, 1998; Falcón et al., 2000; Mello, 2000; Araújo et al., 2005; Ribeiro et al., 2005b; Floeter et al., 2008; Ribeiro, 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Afonso, 2016; Heemstra and Anderson \(Jr.\), 2016; POGID, 2016; POGIS, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019\)](#). This species is sometimes confused with three other species, namely *Mycteroperca rubra* (Bloch 1793), *M. acutirostris* (Valenciennes 1828) and *Epinephelus alexandrinus* (Valenciennes 1628) ([Heemstra & Anderson \(Jr.\), 2016](#)). The neotype of *Serranus fuscus* Lowe 1838, synonym of *M. fusca*, caught off in Madeira is in the MMF collections (MMF 24928).

***Mycteroperca phenax* Jordan & Swain 1884**

Common name: **Scamp; Garoupa-mariola.**

References: Azores ([Morato et al., 2004](#); [Cardigos et al., 2006](#); [Porteiro et al., 2010](#); [Craig et al., 2012](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

***Mycteroperca rubra* (Bloch 1793)**

Common name: **Mottled grouper; Garoupa-chumbo**, Badeja, Badejo, Badejo-das-ilhas, Bodejo, Cherne, Garoupa, Garoupa-badejo, Garupa.

References: mainland Portugal ([Fischer et al., 1981](#), [1987](#); [Tortonese, 1986a](#); [Heemstra, 1991](#); [Heemstra and Randall, 1993](#); [Carneiro et al., 2014](#); [Heemstra and Anderson \(Jr.\), 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Osório de Castro, 1967](#); [IMAR/DOP COLETA 139, 1992](#)) and Madeira[▲] (?) ([de Velloso, 1892](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979b, 1986a](#); [Lloris et al., 1991](#); [Hoese et al., 2006](#)). This species is sometimes confused with *M. fusca* (Lowe 1838) ([Heemstra & Anderson \(Jr.\), 2016](#)). Specimens from Madeira, Azores, Canary Islands, and Cape Verde have been examined by Heemstra (1991) and were identified as *Mycteroperca fusca* (Lowe 1838). Thus, this species is not considered in the total number of valid species for the Azores and Madeira.

***Serranus atricauda* Günther 1874**

Common name: **Blacktail comber; Serrano-de-rolo**, Caroupão, Ferreiro, Garoupa, Garoupa-da-costa, Garoupa-de-natura, Garoupa-de-rolo, Garoupa-do-rolo, Garupa, Serrão.

References: mainland Portugal[▲] ([Bragança, 1903](#); [Richard, 1904, 1908](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Fischer et al., 1981, 1987](#); [Dias et al., 1990](#); [Erzini et al., 2001a](#); [WWF, 2001](#); [Gonçalves et al., 2004](#); [Henriques et al., 2007, 2008](#); [Abecasis et al., 2009](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [OCEANA, 2014](#); [Heemstra and Anderson \(Jr.\), 2016](#)), Azores[▲] ([Drouët, 1861b](#); [Günther, 1874](#); [Boulenger, 1895](#); [Collett, 1896a](#); [Ramos, 1869](#); [Hilgendorf, 1888a, b](#); [Regan, 1903a](#); [Vaillant, 1919](#); [Fowler, 1919, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bauchot and Blanc, 1961](#); [Osório de Castro, 1967](#); [Wood, 1973](#); [Mapstone and Wood, 1974, 1975](#); [Wood and Williams, 1974](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Tortonese, 1979b](#); [Saldanha, 1980](#); [Fischer et al., 1981](#); [INIP, 1985c](#); [Dias and Monteiro, 1986](#); [Silva, 1986a, b](#); [Dias et al., 1990](#); [Barreiros, 1990, 1995](#); [Patzner and Santos, 1990d, 1993](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1997](#); [Smith, 1990d](#); [Wirtz, 1990a, 1992, 1994](#); [Azevedo et al., 1991, 1992](#); [Dias, 1991](#);

Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Gonçalves, 1993; Silva *et al.*, 1994, 1995; Azevedo, 1995; Gonçalves *et al.*, 1995; Krug, 1995; Silva and Menezes, 1995a, b; Menezes, 1996; Morato-Gomes *et al.*, 1998; Sousa *et al.*, 1999; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001b; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Gomes-Pereira *et al.*, 2014; Afonso, 2016; Heemstra and Anderson (Jr.), 2016; Almeida and Biscoito, 2019; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Günther, 1859, 1874; de Velloso, 1892; Boulenger, 1895; Collett, 1896; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979b; INIP, 1980, 1982c; Fischer *et al.*, 1981; Augier, 1985; Dooley *et al.*, 1985; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Severino, 2004; Araújo *et al.*, 2005; Ferreira *et al.*, 2008; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Heemstra and Anderson (Jr.), 2016; POGID, 2016; POGIS, 2016; Friedlander *et al.*, 2017; Froese and Pauly, 2019).

***Serranus cabrilla* (Linnaeus 1758)**

Common name: **Comber; Serrano-alecrim**, Alecrim, Andorinha, Cagalhão-chumbado, Caroupa, Garoupa, Garoupa-azul, Garoupa-da-costa, Garoupa-do-alto, Garoupa-riscada, Garrupa, Garupa, Liali, Peixe-alecrim, Rainha-do-alto, Requeima, Requeime, Requeimo, Requeme, Serrano, Serrão.

References: mainland Portugal (de Brito Capello, 1867d, 1880; Steindachner, 1867; Posser, 1880; Osório, 1888, 1895, 1896; Baldaque da Silva, 1891; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895a, c, 1903, 1935; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979b, 1986a; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982b; Martins, 1981; Costa and Quéro, 1988; Martins *et al.*, 1992; Almeida, 1996; Erzini *et al.*, 1996, 1997, 2001a; Gonçalves *et al.*, 1996, 1997; Cabral, 1999; Borges *et al.*, 2002; Santos *et al.*, 2002a; Mendes *et al.*, 2004; ICES, 2005; Ribeiro *et al.*, 2006, 2008; Fernandes *et al.*, 2007; Henriques *et al.*, 2007, 2008; Batista *et al.*, 2009; Chaves, 2009; Silva *et al.*, 2009; Silva, 2010, 2011; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Silva and Borges, 2014; Golani and Fricke, 2018; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Sampaio, 1904; Nobre, 1924, 1930; Roule and Angel, 1930; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Tortonese, 1954, 1979b, 1986a; Albuquerque, 1954-1956; Osório de Castro, 1967; Wheeler, 1978; INIP, 1985a; Chícharo and Teodósio, 1990; Dias, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Wirtz, 1992; Silva and Menezes, 1995a, b; Silva *et al.*, 1995; Menezes, 1996; Santos *et al.*, 1997; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Gomes-Pereira *et al.*, 2014; Heemstra and Anderson (Jr.), 2016; Froese and Pauly, 2019) and Madeira (Lowe, 1837; Boulenger, 1895; Borodin, 1934; Fowler, 1936b; Osório de Castro, 1967; Maul, 1976; Tortonese, 1979b, 1986a; INIP, 1982c; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Heemstra and Anderson (Jr.), 2016; Golani and Fricke, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in April 2011 is in the MB collections (MB06-005339).

***Serranus hepatus* (Linnaeus 1758)**

Common name: **Brown comber; Serrano-ferreiro**, Ferreiro, Ferreiro-serrano, Garoupa, Garoupa-encarnada, Garupa.

References: mainland Portugal (de Brito Capello, 1869a, b, 1880; Bragança, 1902; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979b; INIP, 1982b; Fischer *et al.*, 1981, 1987; Martins, 1981; Dinis, 1985; Antunes *et al.*, 1988; Smith, 1990d; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Santos *et al.*, 2002a; ICES, 2005; Sousa *et al.*, 2005, 2006; Ribeiro *et al.*, 2006, 2008, 2012; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Neves *et al.*, 2008; Silva, 2008, 2010, 2011; Batista *et al.*, 2009; Silva *et al.*, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Heemstra and Anderson (Jr.), 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Serranus scriba (Linnaeus 1758)

Common name: **Painted comber; Serrano-riscado**, Caroupa, Garoupa, Garoupa-pequena, Garrupa, Garupa, Mirate, Peixe-alecrim, Requeima, Requeime, Requeimo, Requeme, Salmonete, Seima, Serrano, Serrano-pequeno, Serrão, Vaca, Vaqueta.

References: mainland Portugal (de Brito Capello, 1867a, d, 1869b, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979b, 1986a; INIP, 1982b; Fischer et al., 1981, 1987; Martins et al., 1992; Erzini et al., 1996, 2001a; Henriques et al., 2008; Carneiro et al., 2014; Heemstra and Anderson (Jr.), 2016; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Fowler, 1936b; Collins, 1954; Osório de Castro, 1967; Tortonese, 1979b, 1986a; Lloris et al., 1991; Arruda, 1992, 1997; Wirtz, 1994; Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira (Bowdich, 1825; MNHN-IC-2005-0213, 1922; Fowler, 1936b; Osório de Castro, 1967; Tortonese, 1979b; Lloris et al., 1991; Wirtz et al., 2008(?); Carneiro et al., 2014; Heemstra and Anderson (Jr.), 2016). According to Santos et al. (1997) the original report of the occurrence of *S. scriba* for the Azores by Drouët (1861b) is a misidentification of *Serranus atricauda* Günther 1874 and the former authors never found *S. scriba* in the Azores. Based on the Drouët record, the species was subsequently erroneously referred by several authors as present in the Azores. According to Wirtz et al. (2008) and Froese and Pauly (2019) the presence of this species in Madeira is also questionable. Thus, specimens from Madeira and/or the Azores are needed for confirmation the presence of this species in these areas having in mind the contradictions of these authors.

Family Callanthiidae

Callanthias ruber (Rafinesque 1810)

Common name: **Parrot seaperch; Canarinho-do-mar**, Andorinha, Castanheta, Castanheta-amarela-do-alto, Imperador, Imperador-de-natura, Imperador-do-alto, Papagaio, Papagaio-do-mar.

References: mainland Portugal (Nobre, 1895c, 1935; Vieira, 1897, 1898; Seabra, 1911; Fowler, 1936b; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Tortonese, 1979b, 1986a; Fischer et al., 1981, 1987; Martins, 1981; Dinis and Marecos, 1984; Borges et al., 2002; ICES, 2005; Chaves, 2008, 2009; Henriques et al., 2008; Pakhorukov, 2008; Carneiro et al., 2014; Christiansen et al., 2014; OCEANA, 2014; Anderson (Jr.) et al., 2015; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977 (PECS); Tortonese, 1979b, 1986a; Shcherbachov et al., 1985a; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Fock et al., 2002 (PECS); Kukuev, 2002 (PECS); Floeter et al., 2008; Pakhorukov, 2008 (PECS); Porteiro et al., 2010, 2013; Gomes-Pereira et al., 2012, 2014; Carneiro et al., 2014; Costa, 2014; Anderson (Jr.) et al., 2015; Anderson (Jr.) 2016; Almeida and Biscoito, 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Cocco 1829; Lowe, 1839, 1840b, 1843-1860, 1844a, 1849; NHMUK 1855.11.29.13., 1854-1861; Günther, 1859; Boulenger, 1895; Roule, 1919a (PECS); Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976 (PECS); Ehrich, 1977; Tortonese, 1979b, 1986a; Fischer et al., 1981; Lloris et al., 1991; Floeter et al., 2008; Pakhorukov, 2008 (PECS); Christiansen et al., 2009; Carneiro et al., 2014; Anderson (Jr.) et al., 2015; Anderson (Jr.), 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). The holotype of *Callanthias paradisaeus* Lowe 1839, synonym of *C. ruber*, caught off in Madeira is in the NHMUK collections (NHMUK 1855.11.29.13.). Several specimens caught off at the Josephine Bank and Meteor Bank are in the MMF collections (MMF 22461, 1967, set of thirty-one specimens and MMF 22839, 1967, set of five specimens, respectively).

Family Priacanthidae

Cookeolus japonicus (Cuvier 1829)

Common name: **Longfinned bullseye; Fura-vasos-alfonsim**, Fura-vasos.

References: Azores (Azevedo and Heemstra, 1995; Arruda, 1997; Santos et al., 1997; Floeter et al., 2008; Porteiro et al., 2010; Carneiro et al., 2014). This species is often misidentified as *Cookeolus boops* Forster 1801 (Froese and Pauly, 2019). However, for these authors there are no substantiated records in Eastern Atlantic with the exception of the Santa Helena Island.

Heteropriacanthus cruentatus (Lacepède 1801)

Common name: **Glasseye; Fura-vasos-da-rocha**, Alferaz, Alferraz, Alfonsim, Alfonsim-da-costa, Alfonsim-da-rocha, Alfonsim-de-costa, Alfonsim-de-rolo, Alfonsim-do-rolo, Alforaz, Boca-de-vaso, Façola, Fura-vasos, Olho-de-vidro, Realista, Vaso.

References: Azores ([Floeter et al., 2008](#)) and Madeira[▲] ([MNHN-IC-1955-0027](#), n. d.; [Bowdich, 1825](#); [Günther, 1859](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [MCZ 147838, 1970](#); [Hureau, 1979a, 1986c](#); [Fischer et al., 1981](#); [Starnes, 1981, 1988, 2016](#); [Lloris et al., 1991](#); [Araújo et al., 1995](#); [Andrade and Albuquerque, 1995](#); [Delgado, 1998](#); [Falcón et al., 2000](#); [Mello, 2000](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGID, 2016](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)). According to Starnes (2016), Eschmeyer et al. (2019) and Froese and Pauly (2019) this species is characterized by circumglobal geographic range in tropical and tropically influenced seas.

Heteropriacanthus fulgens (Lowe 1838)

Common name: Bright glasseye; **Fura-vasos-da-rocha-brilhante**, Alfarraz, Alferaz, Alferraz. Alfonsim, Alfonsim-da-costa, Alfonsim-de-costa, Alfonsim-de-rolo, Alfonsim-do-rolo, Alforaz, Boca-de-vaso, Fura-vasos, Fura-vasos-da-rocha, Olho-de-vidro, Realista, Vaso.

References: Madeira ([Bowdich, 1825](#); [Lowe, 1837, 1838](#); [Fernandez-Silva and Ho, 2017](#); [Eschmeyer et al., 2019](#)).

Priacanthus arenatus Cuvier 1829

Common name: **Atlantic bigeye; Fura-vasos-vulgar**, Alfonsim, Alfonsim-da-costa, Alfonsim-de-costa, Alfonsim-de-rolo, Alfonsim-de-rolo, Catalufa, Façola.

References: Azores ([NHMUK 1983.9.8.12, 1971-1985](#); [Wirtz, 1994](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Azevedo et al., 2004](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1841](#); [Johnson, 1862a, 1863a](#); [Boulenger, 1895](#); [Osório de Castro, 1967](#); [Hureau, 1979a, 1986c](#); [Dooley et al., 1985](#); [Starnes, 1988, 2016](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). The holotype of *Priacanthus insularum* Johnson 1862, synonym of *P. arenatus*, caught off in Madeira was in the NHMUK collections (apparently lost).

Family Apogonidae

Apogon imberbis (Linnaeus 1758)

Common name: **Cardinal fish; Alcarraz**, Afonsinhos, Alfonsinho, Alfonsinho-da-costa, Cardeal, Folião.

References: mainland Portugal ([Hureau and Monod, 1979](#); [Tortonese, 1986b](#); [Fischer et al., 1987](#); [Erzini et al., 2001a](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Guimarães, 1884](#); [Hilgendorf, 1888a](#); [Vaillant, 1919](#); [Nobre, 1924, 1930](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Wood, 1973, 1974](#); [Wood and Williams, 1974](#); [Harmelin and Harmelin-Vivien, 1979](#); [Hureau and Monod, 1979](#); [Ré, 1979a, 1990](#); [Tortonese, 1979a, 1986b](#); [Saldanha, 1980](#); [Dooley et al., 1985](#); [Barreiros, 1990, 1995](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1997](#); [Wirtz, 1990a, 1994](#); [Azevedo et al., 1991, 1992](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Gonçalves et al., 1995](#); [Harmelin-Vivien et al., 2001](#); [Cardigos et al., 2005](#); [Floeter et al., 2008](#); [Bertoni et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Diogo and Pereira, 2014](#); [Afonso, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Lowe, 1843](#); [Günther, 1859](#); [Fowler, 1923, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#); [Tortonese, 1979a](#); [Dooley et al., 1985](#); [Lloris et al., 1991](#); [Abreu et al., 1995](#); [Falcón et al., 2000](#); [Araújo et al., 2005](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Gon, 2016](#); [POGID, 2016](#); [POGIS, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Family Pomatomidae

Pomatomus saltatrix (Linnaeus 1766)

Common name: **Bluefish; Anchova**, Anxova, Chova, Dichova, Enchova, Xidama.

References: mainland Portugal[▲] (de Brito Capello, 1867a, 1880; Steindachner, 1868a; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Quéro, 1969; Monod, 1979e; Tortonese, 1986b; Fischer et al., 1987; Antunes et al., 1988; Martins et al., 1992, 2015; Erzini et al., 1996; Quéro et al., 1997, 1998; Santos et al., 2002a; ICES, 2005; Gomes and Olim, 2007; Henriques et al., 2008; Ribeiro et al., 2008; Carneiro et al., 2014; Martins and Carneiro, 2018; Baptista et al., 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Sampaio, 1904; Le Gall, 1934; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Martins, 1982a; Nash et al., 1990, 1994a; Patzner et al., 1990, 1992; Wirtz, 1990a, 1994; Arruda, 1992, 1997; Santos et al., 1994b, 1997; Azevedo, 1995; Santos and Nash, 1995; Isidro, 1996; Harmelin-Vivien et al., 2001; Morato et al., 2001a, b; Menezes et al., 2006; Bertoncini et al., 2010; Botelho et al., 2010; Porteiro et al., 2010; Diogo and Pereira, 2013, 2014; Carneiro et al., 2014; Collette, 2016b) and Madeira[▲] (Lowe, 1837; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Collette, 1981, 2016b; Fischer et al., 1981; Tortonese, 1986b; Lloris et al., 1991; Wirtz et al., 2008; Carneiro et al., 2014; Froese and Pauly, 2019).

Family Rachycentridae

Rachycentron canadum (Linnaeus 1766)

Common name: **Cobia; Fogueteiro-galego**, Bacalhau, Cobia, Filho-pródigo, Fogueteiro, Peixe-sargento, Sargento, Soldado-desconhecido.

References: Madeira (Fischer et al., 1981; Collette et al., 2015i; Collette, 2016b). According to Eschmeyer et al. (2019) *R. canadum* is a circumglobal species in tropical and subtropical seas (including Red Sea), strays to Mediterranean Sea. The distribution maps of this species reported by Fischer et al. (1981) partially included the area of Selvagens Islands (Madeira EEZ).

Family Echeneidae

Echeneis naucrates Linnaeus 1758

Common name: **Live sharksucker; Pegador-listado**, Agarrador, Apogador, Pega-coberta, Peixe-agarrador, Peixe-pegador, Peixe-piolho, Peixe-pogador, Peixe-pregador, Peixe-sapato, Peixe-ventosa, Pendura, Piolho, Piolho-do-Brasil, Pegador, Pregador, Pogador, Rémora, Rémora-maior, Ventosa.

References: mainland Portugal (Steindachner, 1868a; Vieira, 1897, 1898; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro et al., 2014; Collette, 2016b), Azores (Lachner, 1986; Arruda, 1997; Santos et al., 1997; Porteiro et al., 2010; Carneiro et al., 2014; Collette, 2016b) and Madeira (Lowe, 1837, 1839, 1840b, 1844a, 1849, 1850, 1852; Günther, 1860; Vinciguerra, 1883; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Maul, 1956a; Osório de Castro, 1967; Lachner, 1986; Wirtz et al., 2008; Carneiro et al., 2014; Collette, 2016b; Froese and Pauly, 2019).

Phtheirichthys lineatus (Menzies 1791)

Common name: **Slender suckerfish; Pegador-delgado**, Agarrador, Peixe-ventosa, Pendura, Pegador, Peixe-pegador, Peixe-ventosa, Rémora, Ventosa.

References: mainland Portugal (Bragança, 1903; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1967; Carneiro et al., 2014; Collette, 2016b), Azores (NHMUK 1903.6.27.24, 1893-1903; Regan, 1903a; Albuquerque, 1954-1956; Osório de Castro, 1967; Lachner, 1986; Lloris et al., 1991; Arruda, 1997; Santos et al., 1997; Porteiro et al., 2010; Carneiro et al., 2014; Collette, 2016b) and Madeira (Osório de Castro, 1967; Lachner, 1986(?); Carneiro et al., 2014; Collette, 2016b).

Remora albescens (Temminck & Schlegel 1850)

Common name: **White suckerfish; Pegador-branco**, Agarrador, Chupa-sangue, Pegador, Peixe-agarrador, Peixe-pegador, Peixe-ventosa, Pendura, Rémora, Sugador, Ventosa.

References: mainland Portugal (Osório, 1894; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Lachner, 1986; Carneiro et al., 2014; Collette, 2016b), Azores (Collett, 1897; Arruda, 1992, 1997; Santos et al., 1997; Cardigos et al., 2005; Porteiro et al., 2010; Carneiro et al., 2014; Collette, 2016b) and Madeira (Osório, 1894; Noronha and Sarmento, 1948; Albuquerque, 1954-1956;

Maul, 1956a; Osório de Castro, 1967; Lachner, 1986; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Collette, 2016b; Froese and Pauly, 2019).

Remora australis (Bennett 1840)

Common name: Whalesucker; **Pegador-das-baleias**.

References: mainland Portugal ([Collette *et al.*, 2015j](#); [Collette, 2016b](#)), Azores ([Collette *et al.*, 2015j](#); [Collette, 2016b](#)) and Madeira ([Collette *et al.*, 2015j](#); [Collette, 2016b](#)).

Remora brachyptera (Lowe 1839)

Common name: **Spearfish remora**; **Pegador-azul**, Agarrador, Chupa-sangue, Pegador, Peixe-agarrador, Peixe-pegador, Peixe-pegajoso, Peixe-ventosa, Pendura, Rémora, Sugador, Ventosa.

References: mainland Portugal ([Osório, 1917](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Lachner, 1986](#); [Quéro *et al.*, 2003](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Bañón *et al.*, 2017b](#)), Azores ([Collette, 1896](#); [Regan, 1903a](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Lachner, 1986](#); [Lloris *et al.*, 1991](#); [Arruda, 1992, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#)) and Madeira ([Lowe, 1839, 1840b, 1844a, 1849](#); [Günther, 1860](#); [Johnson, 1863b](#); [Collette, 1896](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Maul, 1956a](#); [Osório de Castro, 1967](#); [Lachner, 1979, 1986](#); [Lloris *et al.*, 1991](#); [Wirtz *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Roberts *et al.*, 2017](#); [Froese and Pauly, 2019](#)). The whereabouts of the holotype of *Echeneis brachyptera* Lowe 1839, synonym of *R. brachyptera*, caught off in Madeira are unknown.

Remora osteochir (Cuvier 1829)

Common name: **Marlin sucker**; **Pegador-dos-espadartes**, Agarrador, Pegador, Peixe-agarrador, Peixe-pegador, Peixe-piolho, Peixe-ventosa, Rémora, Ventosa.

References: mainland Portugal ([Girard, 1893b](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Bañón *et al.*, 2019b](#)), Azores ([Arruda, 1997](#); [Santos *et al.*, 1997](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Bañón *et al.*, 2019b](#)) and Madeira ([Lowe, 1839, 1840b, 1844a, 1849](#); [Maul, 1956a](#); [Wirtz *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Froese and Pauly, 2019](#)). The holotypes of *Echeneis jacobaea* Lowe 1839 and *Echeneis pediculus* Girard 1893, synonyms of *R. osteochir*, caught off in Madeira and coast of Portugal, respectively, are lost or probably lost.

Remora remora (Linnaeus 1758)

Common name: **Shark sucker**; **Pegador**, Agarrador, Chupa-sangue, Lapador, Pegadeira, Peixe-agarrador, Peixe-pegador, Peixe-pegajoso, Peixe-piolho, Peixe-sapato, Peixe-ventosa, Pendura, Rémora, Ventosa.

References: mainland Portugal ([de Brito Capello, 1867a, 1869b, 1880](#); [Steindachner, 1868a](#); [Gervais and Boulart, 1877b](#); [Vieira, 1898](#); [Bragança, 1903](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Osório, 1917](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Lachner, 1986](#); [Fischer *et al.*, 1987](#); [Gonçalves *et al.*, 2004](#); [Abecasis *et al.*, 2009](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#)), Azores ([Guimarães, 1882a](#); [Hilgendorf, 1888a](#); [Collett, 1897](#); [Sampaio, 1904](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Wheeler, 1978](#); [Lachner, 1986](#); [Wirtz, 1990a, 1994](#); [Arruda, 1992, 1997](#); [Santos *et al.*, 1997](#); [Cardigos *et al.*, 2005](#); [Porteiro *et al.*, 2010](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Porteiro *et al.*, 2017\(?\)](#)) and Madeira ([Lowe, 1839, 1849](#); [Günther, 1860](#); [Vinciguerra, 1883](#); [Fowler, 1923, 1936b](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Maul, 1959](#); [Osório de Castro, 1967](#); [Lachner, 1979, 1986](#); [Lloris *et al.*, 1991](#); [Mello, 2000](#); [Wirtz *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Collette, 2016b](#); [Froese and Pauly, 2019](#)).

Family Carangidae

Alectis alexandrina (Geoffroy St. Hilaire 1817)

Common name: **Alexandria pompano**; **Xareu-enxada**, Cacôa, Charro-alto, Charro-enxada, Charro-largo, Coa, Enxareu, Enxareu-enxada, Fanta, Lírio-de-serro, Lua, Macua, Palmeta, Peixe-prata, Sereia, Xareu, Xareu-corcovado, Xareu-de-natura.

References: mainland Portugal ([Smith-Vaniz, 1986](#); [Fischer *et al.*, 1987](#); [Carneiro *et al.*, 2014](#); [Smith-Vaniz and Herrera, 2015e](#); [Froese and Pauly, 2019](#)) and Madeira ([Fischer *et al.*, 1981](#); [Smith-Vaniz and Herrera, 2015e](#)).

Campogramma glaycos (Lacepède 1801)

Common name: **Vadigo; Xareu-palheta**, Doirado, Dourada, Enxareu, Enxareu-enxada, Palmeta, Pelheta, Palmoneta, Palombeta, Palombeta-campo, Palometa, Prombeta, Prumbeta, Salema, Serra, Xareu, Xareu-palmeta.

References: mainland Portugal ([Osório de Castro, 1967](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981, 1987](#); [Smith-Vaniz, 1986](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015e](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Osório de Castro, 1967](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Dooley et al., 1985](#); [Smith-Vaniz, 1986, 2016b](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015e](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Caranx cryos (Mitchill 1815)

Common name: **Blue runner; Xareu-azul**, Bonito, Enxareu, Gacesse, Gacesso, Írio-de-serra, Xarelete, Xareu, Xareu-gacesse.

References: mainland Portugal ([Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015f](#)), Azores[▲] ([Berry 1959\(?\)](#); [Wirtz, 1990a, c, 1991, 1994](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Cardigos et al., 2005](#); [Floeter et al., 2008](#); [Porteiro et al., 2010](#); [Afonso et al., 2013](#); [IMAR/DOP Underwater visual census 4818, 2013](#); [Carneiro et al., 2014](#)) and Madeira ([Falcón et al., 2000](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Herrera and Smith-Vaniz, 2015f](#); [POGIS, 2016](#); [Smith-Vaniz, 2016b](#); [Froese and Pauly, 2019](#)).

Caranx hippos (Linnaeus 1766)

Common name: **Crevalle jack; Xareu-macoa**, Charéu, Charréu, Charréu-macoa, Charro, Charro-largo, Charro-xareu, Chicharro, Coa, Enchareu-de-natura, Encharréu, Enxareu, Enxarreu, Enxarréu, Japão, Macoa, Macoleta, Macua, Peixe-Japão, Peixe-prussiano, Prussiano, Sereia, Xareu, Xaréu, Xareu-cavalão, Xareu-comum, Xareu-de-natura, Xareu-olho-de-boi, Xarreu, Xarréu, Xereu, Xeréu, Xireu, Xiréu, Xureu, Xuréu.

References: mainland Portugal (?) ([Bragança, 1902](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Palmer, 1971](#); [Hureau and Tortonese, 1979](#); [Smith-Vaniz, 1986](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores (?) ([Hilgendorf, 1888a](#); [Fowler, 1936b](#); [Arruda, 1997](#); [Carneiro et al., 2014](#)) and Madeira (?) ([ZMH 14574, 1885](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). According to Smith-Vaniz and Carpenter (2007) the *Caranx hippos* species complex comprises three extant species: *Caranx hippos* (Linnaeus 1766) from both the Western and Eastern Atlantic Ocean; *Caranx caninus* Günther 1868 from the Eastern Pacific Ocean; and *Caranx fischeri* Smith-Vaniz and Carpenter 2007, from the Eastern Atlantic, including the Mediterranean Sea and Ascension Island. *Caranx hippos* and *C. fischeri* are largely sympatric (Smith-Vaniz et al., 2015b). In the Eastern Atlantic *C. hippos* is known from Mauritania to Angola (Smith-Vaniz and Carpenter, 2007), but historical records for the mainland Portugal and Madeira are probably based on misidentifications of *C. fischeri* and *C. latus*, respectively. Smith-Vaniz and Carpenter (2007) mention that the records of the species from the Azores are unreliable and can not be confirmed. Specimens are needed to decide if both species occur at Madeira or only one of them. Thus, this species is not considered in the total number of valid species.

Caranx latus Agassiz 1831

Common name: **Horse-eye jack; Xareu-de-olho-papudo**, Xareu.

References: Madeira ([Noronha and Sarmento, 1948](#); [Maul, 1949d](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Hureau and Tortonese, 1979](#); [Smith-Vaniz, 1986, 2016b](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Caranx hippos* (Linnaeus 1766) (Smith-Vaniz, 2016b).

Caranx lugubris Poey 1860

Common name: **Black jack; Xareu-enforcado**, Charéu, Charréu, Charréu-beiçudo, Charréu-bicudo, Chicharro, Enforcado, Enxareu, Encharéu, Encharréu, Lírio, Pungo-enforcado, Xareu, Xareu-beiçudo.

References: Azores ([Zugmayer, 1911b](#); [Fowler, 1936b](#); [Osório de Castro, 1954, 1967](#); [Harmelin and Harmelin-Vivien, 1979](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986, 2016b](#);

Lloris *et al.*, 1991; Arruda, 1997; Floeter *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Günther, 1860; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Smith-Vaniz, 1986, 2016b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Caranx rhonchus* Geoffroy St. Hilaire 1817**

Common name: **False scad; Xareu-amarelo**, Cadela, Carapau-amarelo, Carapau-espanhol, Carapau-francês, Carapau-manteiga, Carapau-moiro, Charro, Charro-amarelo, Charro-branco, Chicharro-amarelo, Chicharro-espanhol, Chicharro-francês, Chicharro-moiro, Chicharro-molar, Chicharro-mouro, Xareu, Xareu-moiro.

References: mainland Portugal[▲] (Osório de Castro, 1954, 1967; Borges, 1984; Smith-Vaniz, 1986; Bauchot, 1987; Fischer *et al.*, 1987; Martins *et al.*, 1992; Erzini *et al.*, 1996; Erzini *et al.*, 1997; Santos *et al.*, 2002a, c; ICES, 2005; Gomes and Olim, 2007; Ribeiro *et al.*, 2008; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015g; Martins and Carneiro, 2018; Froese and Pauly, 2019) and Madeira (?) (Fowler, 1936b; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014(?); Froese and Pauly, 2019(?)). Fowler (1936b) mentions the existence of one adult specimen in the U.S. National Museum, obtained by Captain William Stimpson. However, the current geographic distribution of this species is the southern Mediterranean Sea and in the Eastern Atlantic, from Morocco south to Namibia, without including the Madeira Archipelago. Specimens are needed for confirmation in Madeira Archipelago. Thus, this species is not considered in the total number of valid species to Madeira.

***Caranx ruber* (Bloch 1793)**

Common name: **Bar jack; Xareu-carvoeiro**, Enxareu-carvoeiro.

References: Azores (Afonso *et al.*, 2013; Carneiro *et al.*, 2014). In the Eastern Atlantic in addition to known occurrence records from Ascension and St Helena islands, a new occurrence of this Western Atlantic species was sighted by J. Fontes while diving outside Horta Harbour breakwater in 2007 (Afonso *et al.*, 2013).

***Caranx sexfasciatus* Quoy & Gaimard 1825**

Common name: **Bigeye trevally; Charéu-de-natura**, Encharéu.

References: Madeira (?) (Maul, 1949a, d). This species was referred to as being present off Madeira by Maul (1949a) indicating its very rare occurrence. Hureau and Tortonese (1979) reported that Maul's identification might be questionable. Moreover, the current geographic distribution does not include the Atlantic Ocean (Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Specimens are needed for confirmation in Madeira Archipelago. Thus, this species is not considered in the total number of valid species.

***Decapterus macarellus* (Cuvier 1833)**

Common name: **Mackerel scad; Charro-olho-largo**, Carapau-moiro, Carapau-mouro, Cavala-preta, Cavalinha, Charro-mouro, Chicharro-moiro, Chicharro-mouro, Olho-largo.

References: Azores (Hureau and Tortonese, 1979; Smith-Vaniz, 1986, 2016b; Lloris *et al.*, 1991; Azevedo and Heemstra, 1995; Arruda, 1997; Santos *et al.*, 1997; Azevedo *et al.*, 2004; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Bañón *et al.*, 2019b; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967; Hureau and Tortonese, 1979; Smith-Vaniz, 1986, 2016b; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Bañón *et al.*, 2019b; Froese and Pauly, 2019).

***Decapterus punctatus* (Cuvier 1829)**

Common name: **Round scad; Charro-moiro**, Carapau-moiro, Carapau-mouro, Carapau-negrão, Cavala-preta, Cavalinha, Charro-mouro, Chicharro-moiro, Chicharro-mouro, Chicharro-negrão, Olho-largo, Peixe-chuva.

References: Azores (NHMUK 2017.1.5.43-54, 1966; Carneiro *et al.*, 2014) and Madeira (Noronha and Sarmento, 1948; Maul, 1949d; Albuquerque, 1954-1956; Osório de Castro, 1967(?); Fischer *et al.*, 1981; Smith-Vaniz, 1986, 2016b; Smith-Vaniz *et al.*, 1990; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Bañón *et al.*, 2019b; Froese and Pauly, 2019). The occurrence records mentioned by Carneiro *et al.* (2014) were wrong, the geographic coordinates did not correspond to the indicated location. However,

the presence of this species in the Azores is confirmed by a preserved specimen in the Collection Specimens, at the Natural History Museum (London).

Decapterus tabl Berry 1968

Common name: **Rougear scad; Charro-de-cauda-vermelha.**

References: Madeira ([Bañón et al., 2019b](#)). Three specimens caught off in Madeira of this species are in the MHNUSC collections (MHNUSC 25116-1, 2017; MHNUSC 25116-2, 2017; MHNUSC 25116-3, 2017).

D. tabl is an unusual or rare species in the Eastern Atlantic ocean where it was only known in Ascension and St Helena islands (Smith-Vaniz, 2016b).

Elagatis bipinnulata (Quoy & Gaimard 1825)

Common name: **Rainbow runner; Fogueteiro-arco-íris**, Falso-charuteiro, Fogueteiro, Salemão, Salmão, Salmão-arco-íris, Salmão-do-índico, Salemão-Índico.

References: mainland Portugal ([Ramos, 2010](#); [Carneiro et al., 2014](#)) and Azores ([Azevedo and Heemstra, 1995](#); [IMAR/DOP COLETA 132, 1995](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015h](#); [Smith-Vaniz, 2016b](#)) and Madeira ([Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015h](#); [Smith-Vaniz, 2016b](#)). Eschmeyer *et al.* (2019) mention it as a circumglobal species in tropical and warm temperate seas. The distribution map of this species reported by Smith-Vaniz (2016b) partially included the area of Selvagens Islands (Madeira EEZ). However, specimens are needed for confirmation. Apparently rare in the Eastern Atlantic.

Lichia amia (Linnaeus 1758)

Common name: **Leerfish; Palombeta**, Chambé, Colombeta, Doirada, Dourada, Palmeta, Palmoneta, Palmonete, Palombela, Palometa, Palompeta, Pelumbeta, Pombinha, Prombeta, Plumbeta, Prumbeta.

References: mainland Portugal[▲] ([de Brito Capello, 1869a, 1880](#); [Vieira, 1893, 1897, 1898](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981, 1987](#); [Smith-Vaniz, 1986, 2016b](#); [Bauchot, 1987](#); [Smith-Vaniz et al., 1990](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [de Morais et al., 2015a](#); [Smith-Vaniz and Herrera, 2015f](#); [Froese and Pauly, 2019](#)), Azores (?) ([Collett, 1896](#); [Wood, 1974](#); [Wood and Williams, 1974](#); [Arruda, 1997](#); [Carneiro et al., 2014](#)) and Madeira ([Lowe, 1850, 1852](#); [Günther, 1860](#); [Gervais and Boulart, 1877b](#); [Collett, 1896](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986, 2016b](#); [Lloris et al., 1991](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [de Morais et al., 2015a](#); [Smith-Vaniz and Herrera, 2015f](#); [Froese and Pauly, 2019](#)). According to Santos *et al.* (1997) the occurrence of this species in the Azores is a misidentification of *Seriola* spp. that occurs in the Azores. Thus, this species is not considered in the total number of valid species to the Azores.

Naucrates ductor (Linnaeus 1758)

Common name: **Pilotfish; Peixe-piloto**, Cherne-pequeno, Guia, Peixe-guia, Peixe-piolho, Peixe-piloto, Piloto, Piolho, Romeirinho, Romeiro.

References: mainland Portugal ([Carvalho, 1765](#); [de Brito Capello, 1867a, 1869b, 1880](#); [Steindachner, 1868a](#); [Gervais and Boulart, 1877b](#); [Posser, 1880](#); [Almeida and Roquette, 1892](#); [Vieira, 1893, 1897, 1898](#); [Nobre, 1894, 1935](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Berry and Smith-Vaniz, 1978](#); [Fischer et al., 1981, 1987](#); [Smith-Vaniz, 1986](#); [Santos et al., 2002b](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a](#); [Collett, 1896](#); [Vaillant, 1919](#); [Sampaio, 1904](#); [Nobre, 1924, 1930](#); [Roule and Angel, 1930](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Parin, 1970](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986](#); [Wirtz, 1990a](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Pinho and Pereira, 1995b](#); [Saldanha, 1995](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017\(?\)](#); [Carneiro et al., 2014](#); [Smith-Vaniz, 2016b](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1837, 1840a, 1849](#); [Günther, 1860](#); [de Velloso, 1892](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936b](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986, 2016b](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)).

Pseudocaranx dentex (Bloch & Schneider 1801)

Common name: **White trevally; Xareu-bicudo**, Charéu, Encharéu, Enxareu-bicudo, Xaréu, Xareu-beiçudo, Xareu-da-Novazelândia.

References: mainland Portugal ([Almeida, 1996](#); [Henriques et al., 2008](#); [Abecasis et al., 2009](#); [Herrera and Smith-Vaniz, 2015i](#)), Azores[▲] ([Hilgendorf, 1888a](#); [Regan, 1903a](#); [Sampaio, 1904](#); [Vaillant, 1919](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Berry, 1959](#); [Wood and Williams, 1974](#); [Dias et al., 1976](#); [Berry and Smith-Vaniz, 1978](#); [Hureau and Tortonese, 1979](#); [Ré, 1979a, 1990](#); [Fischer et al., 1981](#); [Martins, 1982a](#); [Dooley et al., 1985](#); [Smith-Vaniz, 1986](#); [Barreiros, 1990](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b](#), [1994a](#), [1995a](#); [Smith-Vaniz et al., 1990](#); [Wirtz, 1990a](#), c, [1991, 1994](#); [Lloris et al., 1991](#); [Arruda et al., 1992](#); [Azevedo et al., 1992](#); [Santos, 1992a](#); [Gonçalves, 1993](#); [Nash et al., 1994a](#); [Azevedo, 1995](#); [Gonçalves et al., 1995](#); [Saldanha, 1995](#); [Santos and Nash, 1995](#); [Silva and Menezes, 1995a](#); [Silva et al., 1995](#); [Isidro, 1996](#); [Menezes, 1996](#); [Arruda, 1997](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001a](#); [Menezes et al., 2006](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Diogo and Pereira, 2013, 2014](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015i](#); [Afonso, 2016](#); [Smith-Vaniz, 2016b](#); [Almeida and Biscoito, 2019](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Lowe, 1837](#); [Steindachner, 1868a](#); [Vinciguerra, 1883](#); [de Velloso, 1892](#); [Nunes, 1953, 1974](#); [Hureau and Tortonese, 1979](#); [Fischer et al., 1981](#); [Augier, 1985](#); [Dooley et al., 1985](#); [Smith-Vaniz, 1986, 2016b](#); [Smith-Vaniz et al., 1990](#); [Lloris et al., 1991](#); [Abreu et al., 1995](#); [Andrade and Albuquerque, 1995](#); [Delgado, 1998](#); [Falcón et al., 2000](#); [Mello, 2000](#); [Ribeiro et al., 2005b](#); [Afonso et al., 2008](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Herrera and Smith-Vaniz, 2015i](#); [POGID, 2016](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Caranx georgianus* Cuvier 1833 ([Smith-Vaniz, 2016b](#)).

Selar boops (Cuvier 1833)

Common name: **Oxeye scad; Charro-olho-de-boi**.

References: mainland Portugal ([Fischer et al., 1981](#); [Smith-Vaniz, 1986\(?\)](#), [2016b](#); [Smith-Vaniz et al., 1990](#); [Eschmeyer et al., 2019\(?\)](#); [Froese and Pauly, 2019\(?\)](#)). There is a single confirmed record of this species off Portugal ([Smith-Vaniz, 2016b](#)).

Selar crumenophthalmus (Bloch 1793)

Common name: **Bigeye scad; Charro-preto**.

References: mainland Portugal ([Palmer, 1971](#)). Nine specimens caught in 366-439 m at Sezimbra Bay, in 1965, lengths in SL from 174-180 mm. Since Palmer (1971) no other author describes the presence of this species in mainland Portugal and its occurrence in this EEZ was not yet confirmed by more recent data. The specimens have been in the collection of the former Instituto de Biologia Marítima (IBM) but their whereabouts are unknown.

Selene dorsalis (Gill 1863)

Common name: **African moonfish; Corcovado-africano**, Charuteiro, Charoteira, Corcovado, Limão, Mussolini, Peixe-limão.

References: mainland Portugal ([Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986, 2016b](#); [Smith-Vaniz et al., 1990](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Bowdich, 1925](#); [Fischer et al., 1981](#); [Smith-Vaniz, 1986, 2016b](#); [Smith-Vaniz et al., 1990](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015j](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with two other species, namely *Vomer setapinnis* (Mitchill 1815) and *Selene vomer* (Linnaeus 1758) ([Smith-Vaniz, 2016b](#)). The distribution map of this species reported by Herrera and Smith-Vaniz (2015j) partially included the area of Selvagens Islands (Madeira EEZ).

Selene setapinnis (Mitchill 1815)

Common name: **Atlantic moonfish; Corcovado-do-Golfo**, Corcovado.

References: mainland Portugal (?) ([Nobre, 1935](#); [Albuquerque, 1954-1956](#)) and Madeira (?) ([Bowdich, 1925](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *S. setapinnis* is the Western Atlantic. In the Eastern Atlantic is replaced by *S. dorsalis* (Froese and Pauly, 2019). Thus, this species is not considered in the total number of valid species (see *Selene dorsalis* (Gill, 1863)).

Seriola carpenteri Mather 1971

Common name: **Guinean amberjack; Charuteiro-da-Guiné.**

References: mainland Portugal ([Herrera and Smith-Vaniz, 2015k](#); [Smith-Vaniz, 2016b](#); [Eschmeyer et al., 2019](#)) and Madeira ([Floeter et al., 2008](#); [Herrera and Smith-Vaniz, 2015k](#); [Smith-Vaniz, 2016b](#)). This species is sometimes confused with *Seriola dumerili* (Risso 1810) ([Smith-Vaniz, 2016b](#)).

Seriola dumerili (Risso 1810)

Common name: **Greater amberjack; Charuteiro-catarino**, Anchova, Azeite, Bonite, Charro-catarino, Charuteiro, Charuteiro-de-serril, Charuteiro-de-serrilho, Charuteiro-de-serro, Charuteiro-do-Mediterrâneo, Eiro, Enchova, Enchoveta, Esmoregal, Limão, Lírio, Írio, Peixe-azeite, Peixe-limão, Seriola, Serze, Soldado-desconhecido.

References: mainland Portugal[▲] ([de Brito Capello, 1880](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Hureau and Tortonese, 1979](#); [Smith-Vaniz, 1986](#); [Bauchot, 1987](#); [Fischer et al., 1987](#); [Martins et al., 1992](#); [WWF, 2001](#); [Gonçalves et al., 2004](#); [Henriques et al., 2008](#); [Abecasis et al., 2009](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Fowler, 1936a, b](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Saldanha, 1980](#); [Patzner et al., 1990, 1992](#); [Wirtz, 1990a, 1991, 1994](#); [Arruda, 1992](#); [Arruda et al., 1992](#); [Saldanha, 1995](#); [Santos et al., 1995c, 1997](#); [Harmelin-Vivien et al., 2001](#); [Cardigos et al., 2005](#); [Machado and Barreiros, 2006](#); [Menezes et al., 2006](#); [Floeter et al., 2008](#); [Porteiro et al., 2010, 2017](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Froese and Pauly, 2019](#)) and Madeira ([Osório de Castro, 1967](#); [Falcón et al., 2000](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGIS, 2016](#); [Smith-Vaniz, 2016b](#); [Friedlander et al., 2017](#); [Porteiro et al., 2017](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Seriola carpenteri* Mather 1971 ([Smith-Vaniz, 2016b](#)).

Seriola fasciata (Bloch 1793)

Common name: **Lesser amberjack; Charuteiro-listado**, Anchova, Charoteiro, Charuteira, Enchova, Enchoveta, Limão, Peixe-limão, Serie, Seriola, Serze.

References: Azores ([Machado and Barreiros, 2006](#); [Porteiro et al., 2010](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Smith-Vaniz, 2016b](#); [Froese and Pauly, 2019](#)) and Madeira ([Fischer et al., 1981](#); [Smith-Vaniz, 1986, 2016b](#); [Smith-Vaniz et al., 1990](#); [Lloris et al., 1991](#); [Andrade and Albuquerque, 1995](#); [Falcón et al., 2000](#); [Floeter et al., 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [POGIS, 2016](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Seriola carpenteri* Mather 1971 ([Smith-Vaniz, 2016b](#)).

Seriola lalandi Valenciennes 1833

Common name: **Yellowtail amberjack; Charuteiro-azeite**, Anchova, Azeite, Charuteira, Charuteira-de-serrilho, Charoteiro, Charoteiro-de-serril, Charuteiro-de-serrilho, Charuteiro-de-serro, Enchova, Enchoveta, Limão, Lírio-de-serro, Peixe-azeite, Peixe-limão, Seriola, Soldado-desconhecido.

References: mainland Portugal[▲] (?) ([de Brito Capello, 1867a, 1880](#); [Almeida and Roquette, 1892](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Santos et al., 2002a](#); [Carneiro et al., 2014](#)), Azores (?) ([Osório de Castro, 1967](#); [Wood and Williams, 1974\(?\)](#); [Arruda, 1997](#)) and Madeira (?) ([Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Carneiro et al., 2014](#)). [Eschmeyer et al. \(2019\)](#) mention it as a circumglobal species in subtropical to temperate seas of Southern Hemisphere and according to [Smith-Vaniz \(2016b\)](#) in the Eastern Atlantic known only from St Helena Island and South Africa. However, historical records for the mainland Portugal, Azores and Madeira are probably based on a misidentification of one of the others *Seriola* species occurring in the north-eastern Atlantic. The records of official landings from commercial catches for the period 2015-2018[▲] can not be confirmed. Thus, this species is not considered in the total number of valid species.

Seriola rivoliana Valenciennes 1833

Common name: **Longfin yellowtail; Charuteiro-falcado**, Anchova, Azeite, Charuteira, Charoteiro, Charuteiro, Enchova, Enchoveta, Írio, Limão, Lírio, Peixe-azeite, Peixe-limão, Preguiçosa, Seriola.

References: mainland Portugal ([Fischer et al., 1981](#); [Smith-Vaniz, 1984, 1986, 2002a, 2016b](#); [Smith-Vaniz et al., 1990](#); [WWF, 2001](#); [Gonçalves et al., 2004](#); [Abecasis et al., 2009](#); [de Casamajor and Morandeau,](#)

2013; Carneiro *et al.*, 2014; OCEANA, 2014; Froese and Pauly, 2019), Azores[▲] (Saldanha, 1980; Fischer *et al.*, 1981; Smith-Vaniz, 1986, 2002a, 2016b; Smith-Vaniz *et al.*, 1990; Azevedo *et al.*, 1991; Lloris *et al.*, 1991; Wirtz, 1994; Saldanha, 1995; Santos *et al.*, 1995a, 1997; Arruda, 1997; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001b; Cardigos *et al.*, 2005; Machado and Barreiros, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; de Casamajor and Morandeau, 2013; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso, 2016; Froese and Pauly, 2019) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1860; Noronha and Sarmento, 1948; Maul, 1949d; Albuquerque, 1954-1956; Osório de Castro, 1967; Hureau and Tortonese, 1979; Fischer *et al.*, 1981; Smith-Vaniz, 1986, 2002a, 2016b; Smith-Vaniz *et al.*, 1990; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; de Casamajor and Morandeau, 2013; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of *Seriola dubia* Lowe 1839, synonym of *S. rivoliana*, caught off in Madeira are unknown.

***Seriola zonata* (Mitchill 1815)**

Common name: **Banded rudderfish; Charuteiro-olhete**, Charuteira, Charuteiro, Limão, Peixe-limão.

References: mainland Portugal (?) (Hureau and Tortonese, 1979) and Madeira (?) (Hureau and Tortonese, 1979; Lloris *et al.*, 1991). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. zonata* is the Western Atlantic. It is sometimes confused with *Seriola lalandi* Valenciennes 1833 in the South Atlantic and with *Seriola dumerili* and *Seriola fasciata* elsewhere in its range (Smith-Vaniz, 2002b). The Madeira and mainland Portugal quotes are probably referring the species *Seriola dumerili* (Risso 1810). Thus, this species is not considered in the total number of valid species.

***Trachinotus carolinus* (Linnaeus 1766)**

Common name: **Florida pompano; Sereia-da-Florida**.

References: Azores (MCZ 84185, 1964; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *T. carolinus* is the Western Atlantic.

***Trachinotus ovatus* (Linnaeus 1758)**

Common name: **Pompano; Sereia-camochilo**, Bebecá, Cabra, Camochila, Camochilo, Doirada, Dourada, Estaca, Facaio, Fogueiro, Írio, Palmeta, Palmoneta, Palombela, Palombeta, Palometa, Palomonete, Palompeta, Pâmpano, Pelombeta, Peixe-cabra, Pelumbeta, Plombeta, Pombinha, Pômpano, Prombeta, Prumbeta, Ranhosa, Ranhoso, Salema, Saléma, Sereia, Sereia-riscada, Trombeta, Tronbeta, Trumbeta.

References: mainland Portugal[▲] (Vieira, 1898; Seabra, 1911; Nobre, 1935; Osório de Castro, 1954, 1967; Hureau and Tortonese, 1979; Smith-Vaniz, 1986; Bauchot, 1987; Fischer *et al.*, 1987; Erzini *et al.*, 1996; Borges *et al.*, 2002; Palma *et al.*, 2003; Gomes and Olim, 2007; Henriques *et al.*, 2008; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Martins *et al.*, 2015; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Sampaio, 1904; Zugmayer, 1911b; Lampe, 1914; Fowler, 1919, 1936b; Ninni, 1934; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Wood, 1974; Wheeler, 1978; Harmelin and Harmelin-Vivien, 1979; Hureau and Tortonese, 1979; Ré, 1979a, 1990; Saldanha, 1980; Martins, 1982a; Smith-Vaniz, 1986; Isidro, 1988, 1990a, b; Barreiros, 1990; Nash *et al.*, 1990; Patzner, 1990; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994b, 1997; Wirtz, 1990a, 1994; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Azevedo *et al.*, 1992; Santos, 1992a; Azevedo, 1995; Gonçalves *et al.*, 1995; Pinho *et al.*, 1995; Saldanha, 1995; Santos and Nash, 1995; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a, b; Santos *et al.*, 2002a; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Afonso, 2016; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837; Günther, 1860; de Velloso, 1892; Fowler, 1919, 1936b; Borodin, 1934; Noronha and Sarmento, 1934; Nobre, 1935; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Hureau and Tortonese, 1979; Fischer *et al.*, 1981; Smith-Vaniz, 1986; Lloris *et al.*, 1991; Falcón *et al.*, 2000; Mello, 2000; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGIS, 2016; Smith-Vaniz, 2016b; Froese and Pauly, 2019). The holotype of *Trachinotus madeirensis* Borodin 1934, synonym of *T. ovatus*, caught off in Madeira was in the VMM collections (VMM 1279).

Trachurus lathami Nichols 1920

Common name: **Rough scad; Carapau-rugoso**, Carapau-negrão-austral.

References: Azores ([MCZ 166359, 1969](#); [Carneiro et al., 2014](#)). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2019) the current geographic distribution of *T. lathami* is the Western Atlantic. *T. lathami* is widely distributed in the Western Atlantic, in schools near the bottom over the continental shelf, but there are only two specimens caught off in the Azores and Cape Verde islands.

Trachurus mediterraneus (Steindachner 1868)

Common name: **Mediterranean horse mackerel; Carapau-do-Mediterrâneo**, Charro.

References: mainland Portugal[▲] ([Fischer et al., 1981, 1987](#); [Smith-Vaniz, 1986](#); [Bauchot, 1987](#); [Smith-Vaniz et al., 1990](#); [Erzini et al., 2001a](#); [Borges et al., 2002, 2003](#); [ICES, 2005](#); [Henriques et al., 2007, 2008](#); [Chaves, 2009](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([SMF 13315, n.d.](#); [Hureau and Tortonese, 1979](#); [Robins et al., 1991](#); [Froese and Pauly, 2019](#)).

Trachurus picturatus (Bowdich 1825)

Common name: **Blue jack mackerel; Carapau-negrão**, Amália, Barba-ruiva, Caneco, Carapau-azul, Carapau-de-gato, Carapau-francês, Carapau-moiro, Carapau-molar, Carapau-negro, Carapau-preto, Charrinho, Charréu, Charro, Chicharro, Chicharrinho-de-cana, Chicharro-azul, Chicharro-do-alto, Chicharro-creolo, Chicharro-criolo, Chicharro-de-gorazeira, Chicharro-de-grozeira, Chicharro-de-rolo, Chicharro-do-rolo, Chicharro-espanhol, Chicharro-francês, Chicharro-gorazeiro, Chicharro-grado, Chicharro-miúdo, Chicharro-moiro, Chicharro-negrão, Chicharro-preto, Churreu, Jaquim, Jaquizinho, Joaquim, Joaquininho, Ruama, Salmonete-saloio, Xureu.

References: mainland Portugal[▲] ([de Brito Capello, 1867a, c, 1869b, 1880](#); [Vinciguerra, 1883](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1893, 1897, 1898](#); [Nobre, 1895a, 1903, 1935](#); [Osório, 1909](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Hureau and Monod, 1979](#); [Hureau and Tortonese, 1979](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [INIP, 1982a, b](#); [Borges, 1984](#); [Dinis, 1985](#); [Smith-Vaniz, 1986](#); [Bauchot, 1987](#); [Smith-Vaniz et al., 1990](#); [Monteiro et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002a](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Fernandes et al., 2007](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Pakhorukov, 2008](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Christiansen et al., 2014](#); [Silva and Borges, 2014](#); [Martins et al., 2015](#); [Smith-Vaniz, 2016b](#); [Martins and Carneiro, 2018](#); [Moreira et al., 2018](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Drouët, 1861b](#); [Ramos, 1869](#); [Lütken, 1880](#); [Hilgendorf, 1888a](#); [Collett, 1896a](#); [Regan, 1903a](#); [Sampaio, 1904](#); [Murray and Hjort, 1912](#); [Lampe, 1914](#); [Clark 1913](#); [Nobre, 1924, 1930](#); [Fowler, 1919, 1936](#); [Ramalho, 1929c](#); [Ferreira, 1932, 1937a](#); [Ribeiro, 1936](#); [Letacounoux, 1951](#); [Koefoed, 1952](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Frade, 1960](#); [Osório de Castro, 1967](#); [Shaboneyev, 1973, 1980](#); [Wood, 1974](#); [Wood and Williams, 1974](#); [Dias et al., 1976, 1990](#); [Maul, 1976 \(PECS\)](#); [Ehrich, 1977](#); [Shaboneyev and Riazantseva, 1977](#); [Martins, 1978, 1982a, b, 1992](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a](#); [Shaboneyev and Kotlyar, 1979](#); [Fischer et al., 1981](#); [Muzanor and Westhaus, 1981](#); [INIP, 1985a, b, c](#); [Shcherbachov et al., 1985a](#); [Dias and Monteiro, 1986](#); [Smith-Vaniz, 1986](#); [Isidro, 1988, 1990a, b](#); [Barreiros, 1990](#); [Smith-Vaniz et al., 1990](#); [Azevedo et al., 1991](#); [Dias, 1991](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Pereira, 1992d](#); [Silva, 1992](#); [Nash et al., 1994a](#); [Pierce et al., 1994](#); [Santos et al., 1994a, b, 1995a, c, 1997](#); [Wirtz, 1994](#); [Azevedo, 1995](#); [Clarke and Martins, 1995](#); [Clarke et al., 1995](#); [Gonçalves et al., 1995](#); [Granadeiro et al., 1995](#); [Pinho and Pereira, 1995b](#); [Pinho et al., 1995](#); [Santos and Nash, 1995](#); [Silva and Menezes, 1995a, b](#); [Pereira, 1995](#); [Silva et al., 1995](#); [Isidro, 1996](#); [Menezes, 1996](#); [Monteiro, 1996](#); [Morato-Gomes et al., 1998](#); [Sousa et al., 1999](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001a](#); [Kukuev, 2002 \(PECS\)](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Rosa et al., 2006](#); [Pakhorukov, 2008 \(PECS\)](#); [Sutton et al., 2008](#); [Bertoni et al., 2010](#); [Porteiro et al., 2010, 2013, 2017](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#); [Diogo and Pereira, 2014](#); [Afonso, 2016](#); [Smith-Vaniz, 2016b](#); [Moreira et al., 2018](#); [Almeida and Biscoito, 2019](#); [Fauconnet et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([Bowdich, 1825](#); [Lowe, 1837](#); [Vinciguerra, 1883](#); [de Velloso, 1892](#); [Fowler, 1923, 1936](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Maul, 1976](#); [Ehrich, 1977](#); [Hureau and Monod, 1979](#); [Hureau and Tortonese, 1979](#); [INIP, 1980, 1982c, 1984a, b, c](#); [Fischer et al., 1981](#); [Dooley et al., 1981](#)).

1985; Smith-Vaniz, 1986; Bauchot, 1987; Smith-Vaniz *et al.*, 1990; Lloris *et al.*, 1991; Jesus, 1992; Ribeiro *et al.*, 2005b; Vasconcelos *et al.*, 2006; Pakhorukov, 2008 (PECS); Wirtz *et al.*, 2008; Christiansen *et al.*, 2009; Carneiro *et al.*, 2014; POGID, 2016; Smith-Vaniz, 2016b; Biscoito *et al.*, 2017; Porteiro *et al.*, 2017; Moreira *et al.*, 2018; Froese and Pauly, 2019). Several specimens caught off at the Meteor Bank are in the MMF collections (MMF 22344, 1967, one specimen, MMF 22360, 1967, one specimen, MMF 22539, 1967, set of ten specimens, and MMF 22855, 1967, one specimen).

***Trachurus trachurus* (Linnaeus 1758)**

Common name: **Atlantic horse mackerel; Carapau**, Amália, Caneco, Carapau-branco, Carrapau-de-corrida, Carapau-de-gato, Carapau-de-orelhas, Carapau-pipi, Carapau-sardinheiro, Carapau-vizinho, Carapau-vivo, Carapau-vulgar, Charrinho, Charro, Charro-do-alto, Chicharro-creolo, Chicharro-do-alto, Chicharro-branco, Chicharro-charéu, Chicharro-enzareu, Chicharro-francês, Chicharro-grado, Chicharro-grande, Chicharro-miúdo, Chicharro-vulgar, Chicharro-xaréu, Chicharro-xarréu, Chirelo, Fadista, Jaquim, Jaquizinho, Joaquim, Joaquinzinho, Pencudo, Pipi, Plim, Ruama, Salmonete-saloio, Serêlo, Sorêlo.

References: mainland Portugal▲ (Carvalho, 1765; Vandelli, 1771; de Brito Capello, 1867a, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895a, c, 1903, 1935; Osório, 1895, 1896; Seabra, 1911; Murray and Hjort, 1912; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Hureau and Tortonese, 1979; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982a, b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Smith-Vaniz, 1986; Bauchot, 1987; Antunes *et al.*, 1988; Arruda *et al.*, 1988; Martins *et al.*, 1992, 1999, 2015; Almeida, 1996; Erzini *et al.*, 1996, 1997, 2001a, b; Gonçalves *et al.*, 1996, 1997; Cabral, 1999; Gomes *et al.*, 2001; Monteiro *et al.*, 2001; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Santos *et al.*, 2002a; Palma *et al.*, 2003; Prista *et al.*, 2003; Garnerot *et al.*, 2004; Mendes *et al.*, 2004; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Stergiou *et al.*, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Neves *et al.*, 2008; Silva, 2008, 2010, 2011; Batista *et al.*, 2009; Ribeiro *et al.*, 2008; Garrido *et al.*, 2009; Silva *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Smith-Vaniz *et al.*, 2015c; Bañón *et al.*, 2016a; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Mecklenburg *et al.*, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Drouët, 1861b; Hilgendorf, 1888a(?); Regan, 1903a; Sampaio, 1904; Murray and Hjort, 1912; Lampe, 1914; Clark, 1915; Ferreira, 1932, 1937a; Collins, 1954; NHMUK 1998.8.9.13460-13469, 1966; Osório de Castro, 1967; Dias *et al.*, 1976; Dias, 1991; Arruda, 1992, 1997; Carneiro *et al.*, 2014) and Madeira (Cuvier and Valenciennes, 1833; Günther, 1860, 1862; Drouët 1861b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Smith-Vaniz, 1986, 2016b; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Smith-Vaniz *et al.*, 2015c; Mecklenburg *et al.*, 2018; Froese and Pauly, 2019). According to Santos *et al.* (1997) several authors referred erroneously the occurrence of *T. trachurus* in the Azores based on misidentification of *T. picturatus*. However, there are three specimens in museum collections of U.S. National Museum and Natural History Museum (London) classified as *T. trachurus*. Thus, these museum specimens require re-evaluation.

Family Coryphaenidae

***Coryphaena equiselis* Linnaeus 1758**

Common name: **Pompano dolphinfish; Doirado-pampo**, Delfim, Doirada, Doirada-amarela, Doirada-fêmea, Dourada, Dourada-amarela, Dourada-fêmea, Dourado, Dourado-fêmea, Dourado-pampo.

References: mainland Portugal (de Brito Capello, 1880; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Carneiro *et al.*, 2014), Azores (MNHN-IC-A-8189, 1883; Cuvier and Valenciennes, 1833; Drouët, 1861b; Steindachner, 1868a; Ramos, 1869; Hilgendorf, 1888a; Sampaio, 1904; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979c; Fischer *et al.*, 1981; Palko *et al.*, 1982; Collette, 1986a, 2016b; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1843-1860, 1849; Gervais and Boulart, 1877b(?); de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Collette, 1986a, 2016b; Lloris *et al.*, 1991; Wirtz

et al., 2008; Carneiro *et al.*, 2014; Froese and Pauly, 2019). This species is sometimes confused with *Coryphaena hippurus* Linnaeus 1758 (Collette, 2016b). The holotype of *Coryphaena azorica* Valenciennes in Cuvier & Valenciennes 1833, synonym of *C. equiselis*, caught off near Azores is in the MNHN collections (MNHN-IC-A-8189).

***Coryphaena hippurus* Linnaeus 1758**

Common name: **Common dolphinfish; Doirado**, Delfim, Doirada, Doirada-amarela, Doirada-fêmea, Doirada-de-cabeça-de-macho, Doirado-de-topete, Doirado-do-alto, Doirado-macho, Dourada, Dourada-amarela, Dourado, Dourado-de-cabeça-de-macho, Dourado-de-topete, Dourado-do-alto, Dourado-macho, Sapatorra.

References: mainland Portugal▲ (Osório de Castro, 1954, 1967; Fischer *et al.*, 1981, 1987; Collette, 1986a; Erzini *et al.*, 2001b; Santos *et al.*, 2002a; Bordalo-Machado *et al.*, 2009; Carneiro *et al.*, 2014), Azores▲ (Richard, 1905b; Bouvier, 1907; Guiart, 1935; Fowler, 1936b; Collins, 1954; Osório de Castro, 1967; Fischer *et al.*, 1981; Martins, 1982a; Collette, 1986a; Lloris *et al.*, 1991; Pinho and Pereira, 1995b; Arruda, 1997; Santos *et al.*, 1997; Morato *et al.*, 2001a; Cardigos *et al.*, 2005; Porteiro *et al.*, 2010, 2017(?); Menezes *et al.*, 2012; Carneiro *et al.*, 2014; Collette, 2016b) and Madeira▲ (Bowdich, 1825; Lowe, 1837, 1839, 1840b, 1844a, 1849; Günther, 1860; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979c; Fischer *et al.*, 1981; Collette, 1986a; Lloris *et al.*, 1991; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Collette, 2016b; POGID, 2016; Froese and Pauly, 2019). The whereabouts of the syntype of *Coryphaena nortoniana* Lowe 1839, synonym of *C. hippurus*, caught off in Madeira are unknown.

Family Bramidae

***Brama brama* (Bonnaterre 1788)**

Common name: **Atlantic pomfret; Xaputa**, Brama-rai, Chaputa, Castanheta, Freira, Palometa, Plumbeta. References: mainland Portugal▲ (de Brito Capello, 1867a, 1869b, 1880; Steindachner, 1868a; Gervais and Boulart, 1877b; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1903, 1904, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Muus and Dahlström, 1978; Mead, 1979; Fischer *et al.*, 1981, 1987; INIP, 1981a; Martins, 1981; Haedrich, 1986b; Erzini *et al.*, 2001b; Santos *et al.*, 2002b; ICES, 2005; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2008; Carneiro *et al.*, 2014; Iwamoto *et al.*, 2015; Bañón *et al.*, 2016a; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Hilgendorf, 1888a; Regan, 1903a; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Mead and Haedrich, 1965; Osório de Castro, 1967; Mead, 1972; Wheeler, 1978; Fischer *et al.*, 1981; Haedrich, 1986b; Lloris *et al.*, 1991; Arruda, 1992, 1997; Silva *et al.*, 1995; Menezes, 1996; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Besugo, 2013; Carneiro *et al.*, 2014; Iwamoto *et al.*, 2015) and Madeira▲ (Bowdich, 1825; Lowe, 1837, 1849; Günther, 1860; de Velloso, 1892; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Haedrich, 1986b; Lloris *et al.*, 1991; Severino, 2004; Carneiro *et al.*, 2014; Iwamoto *et al.*, 2015). One specimen caught off at the Portuguese continental slope in October 2010 is in the MB collections (MB06-005355).

***Pteraclis carolinus* Valenciennes 1833**

Common name: Fanfish; **Freira**, Xaputa.

References: Azores (PECS) (MCZ 76279, 1973; MCZ 76284, 1984; Carneiro *et al.*, 2014) and Madeira (Lowe, 1843, 1844b; Günther, 1860; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Mead, 1979; Haedrich, 1986b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Froese and Pauly, 2019). The holotype (unique) of *Pteraclis papilio* Lowe 1843, synonym of *P. carolinus*, caught off in Madeira is apparently lost.

***Pterycombus brama* Fries 1837**

Common name: **Atlantic fanfish; Freira-do-Atlântico**.

References: mainland Portugal (Haedrich, 1986b; Carneiro *et al.*, 2014; Collette and Smith-Vaniz, 2015; Bañón *et al.*, 2016a), Azores (Haedrich, 1986b; Arruda, 1997; Santos *et al.*, 1997; Menezes *et al.*, 2006;

Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira (Haedrich, 1986b; Carneiro *et al.*, 2014; Collette and Smith-Vaniz, 2015).

***Taractes asper* Lowe 1843**

Common name: **Rough pomfret; Xaputa-rugosa**.

References: mainland Portugal (Haedrich, 1986b; Carneiro *et al.*, 2014), Azores (Hilgendorf, 1888a; Fowler, 1936a; ZMH 117464, 1979; Haedrich, 1986b; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014) and Madeira (Lowe, 1843, 1844b; Günther, 1860; MOM-IC-0091-1731, 1927; Roule, 1929; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Maul, 1954a; Mead and Maul, 1958; Mead, 1979; Haedrich, 1986b; Lloris *et al.*, 1991; Carneiro *et al.*, 2014; Bañón *et al.*, 2016a; Roberts *et al.*, 2017; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of this species caught off in Madeira are unknown.

***Taractes rubescens* (Jordan & Evermann 1887)**

Common name: Black pomfret; **Xaputa-negra**.

References: Azores (Arruda, 1997; Santos *et al.*, 1997; Cardigos *et al.*, 2005(?); Carvalho-Filho *et al.*, 2009; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira (Mead, 1979; Haedrich, 1986b; Lloris *et al.*, 1991; Carvalho-Filho *et al.*, 2009; Carneiro *et al.*, 2014; Froese and Pauly, 2019).

***Taractichthys longipinnis* (Lowe 1843)**

Common name: **Big-scale pomfret; Xaputa-galhuda**, Cangulo, Capelo, Chaputa, Chaputão, Freira, Freira-de-natura, Freira-do-alto, Xaputa, Xaputão.

References: mainland Portugal (de Brito Capello, 1880; Osório, 1888; Almeida and Roquette, 1892; Vieira, 1897, 1898; Bragança, 1903; Seabra, 1911; Nobre, 1935; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fischer *et al.*, 1981; Haedrich, 1986b; Erzini *et al.*, 2001b; Carneiro *et al.*, 2014), Azores (Hilgendorf, 1888a(?); Fowler, 1936b; Osório de Castro, 1967; Fischer *et al.*, 1981; Haedrich, 1986b; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Carvalho-Filho *et al.*, 2009; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1843, 1844b; Günther, 1860; Johnson, 1863; de Brito Capello, 1871; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1934, 1948; Nobre, 1935; Fowler, 1936b; Nunes, 1953, 1974; Albuquerque, 1954-1956; Mead and Maul, 1958; Osório de Castro, 1954, 1967; Mead, 1979; Fischer *et al.*, 1981; Haedrich, 1986b; Lloris *et al.*, 1991; Carvalho-Filho *et al.*, 2009; Carneiro *et al.*, 2014; Roberts *et al.*, 2017; Froese and Pauly, 2019). The holotype (unique) of *Brama longipinnis* Lowe 1843, synonym of *T. longipinnis*, caught off in Madeira is lost.

Family Caristiidae

***Caristius fasciatus* (Borodin 1930)**

Common name: Greenland manefish; **Peixe-crina-da-Gronelândia**.

References: Azores (Kukuev *et al.*, 2013). *Caristius groenlandicus* Jensen 1941 is synonym of *C. fasciatus*.

***Caristius macropus* (Bellotti 1903)**

Common name: Manefish; **Peixe-crina-barbatana-comprida**.

References: Azores (PECS) (?) (ZMH 117002, 1979; ZMH 120747, 1982; ZMH 120761, 1982; Carneiro *et al.*, 2014). According to Eschmeyer *et al.* (2019) the current geographic distribution of *C. macropus* is the North Pacific. Stevenson and Kenaley (2013) in the revision of the genera *Caristius* and *Platyberyx*, include the ZMH 120747 and ZMH 120761 specimens as *Platyberyx opalescens* Zugmayer 1911. Thus, this species is not considered in the total number of valid species.

***Paracaristius maderensis* (Maul 1949)**

Common name: No English common name; **Peixe-crina-da-Madeira**, Vampiro.

References: Azores (Melo and Menezes, 2002; Porteiro *et al.*, 2010, 2017; Menezes *et al.*, 2012; Carneiro *et al.*, 2014) and Madeira (Maul, 1949c, 1954a; Albuquerque, 1954-1956; Osório de Castro, 1967; Nielson, 1979e; Lloris *et al.*, 1991; Trunov *et al.* 2006; Stevenson and Kenaley, 2011; Carneiro *et al.*, 2014; Stevenson *et al.*, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype and one

paratype of *Caristius maderensis* Maul 1949, synonym of *P. maderensis*, caught off in Madeira is in the MMF collections (MMF 2343 and MMF 2344, respectively).

Platyberyx andriashevi (Kukuev, Parin & Trunov 2012)

Common name: No English common name; **Peixe-crina-de-Andriashev**.

References: Madeira ([Stevenson and Kenaley, 2013](#); [Carpenter et al., 2015c](#); [Stevenson et al., 2016](#)).

Platyberyx mauli Kukuev, Parin & Trunov 2012

Common name: No English common name; **Peixe-crina-de-Maul**.

References: Madeira ([Maul, 1949](#); [Kukuev et al., 2012](#); [Stevenson and Kenaley, 2013](#); [Stevenson et al., 2016](#)).

Platyberyx opalescens Zugmayer 1911

Common name: No English common name; **Peixe-crina-opala**, Aia.

References: mainland Portugal ([MOM-IC-0091-1196](#), 1910; Zugmayer, 1911a, b; Fowler, 1936a; [Albuquerque, 1954-1956](#); Osório de Castro, 1967; Kukuev et al., 2012; Carneiro et al., 2014; Froese and Pauly, 2019), Azores ([Koefoed, 1953](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010, 2017](#); Carneiro et al., 2014; Froese and Pauly, 2019) and Madeira ([Noronha and Sarmento, 1948](#); [Maul, 1949c, 1954a](#); [Albuquerque, 1954-1956](#); Osório de Castro, 1967; Nielsen, 1979h; Lloris et al., 1991; Stevenson and Kenaley 2013; Carneiro et al., 2014; Stevenson et al., 2016; Porteiro et al., 2017; Froese and Pauly, 2019). The holotype (unique) of this species caught off in mainland Portugal (36°06'N, 9°W), is in the MOM collections (MOM-IC-0091-1179). Kukuev et al. (2012) made the re-description of the exemplar previously described by Maul (1949) as *P. opalescens* and re-evaluated it as *P. mauli* Kukuev, Parin & Trunov 2012.

Family Lutjanidae

Etelis carbunculus Cuvier 1828

Common name: **Deep-water red snapper**; **Vermelho-salmão**, Salmão.

References: Madeira ([Günther, 1859](#); [Maul, 1949d](#); [Albuquerque, 1954-1956](#); Osório de Castro, 1967; Carneiro et al., 2014). According to Froese and Pauly (2019) and Eschmeyer et al. (2019) the current geographic distribution of *E. carbunculus* is the Indo-Pacific region: East Africa to the Hawaiian Islands, north to southern Japan, and south to Australia. However, there are historical records on the occurrence of this species in Madeira.

Etelis oculatus (Valenciennes 1828)

Common name: **Queen snapper**; **Rosa-salmão**.

References: Madeira ([Lowe, 1843-1860](#); [Günther, 1859](#); [NHMUK 1857.6.13.112, 1854-1861](#); Carneiro et al., 2014). According to Eschmeyer et al. (2019) the current geographic distribution of *E. oculatus* is the Western Atlantic. However, there is a specimen in the NHMUK collections (NHMUK 1857.6.13.112) registered as *Serranus oculatus* Valenciennes 1828, which is a synonym of *E. oculatus*.

Lutjanus cyanopterus (Cuvier 1828)

Common name: **Cubera snapper**; **Luciano-de-ventre-vermelho**, Caranha, Caranho, Vermelho-caranho.

References: Azores ([Ribeiro et al., 2017](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *L. cyanopterus* is the Western Atlantic.

Lutjanus goreensis (Valenciennes 1830)

Common name: **Gorean snapper**; **Luciano-da-Goreia**, Bica, Castanhola-da-Goreia, Castanholas, Eburgi, Garoupa, Luciano, Mero.

References: Madeira ([Floeter et al., 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Rhomboplites aurorubens (Cuvier 1829)

Common name: **Vermilion snapper**; **Luciano-rosa**, Vermelho.

References: Azores (PECS) ([ROM 23910, 1966](#); [Carneiro et al., 2014](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *R. aurorubens* is the Western Atlantic.

Family Lobotidae

Lobotes surinamensis (Bloch 1790)

Common name: **Tripletail; Furriel**, Bentana-mar-de-fora, Lobotes, Peixe-folha.

References: mainland Portugal ([Roux, 1986c](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#); [Craig, 2015c](#); [Froese and Pauly, 2019](#)), Azores ([Roux, 1986c](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Porteiro et al., 2010](#); [Afonso et al., 2013](#); [Carneiro et al., 2014](#)) and Madeira[▲] ([Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Tortonese, 1979d](#); [Fischer et al., 1981](#); [Roux, 1986c](#); [Lloris et al., 1991](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Craig, 2015c](#); [Carpenter, 2016](#); [Froese and Pauly, 2019](#)).

Family Haemulidae

Brachydeuterus auritus (Valenciennes 1832)

Common name: **Bigeye grunt; Roncador-colo-colo**, Colo-colo, Roncador, Roncador-de-olhos-grande.

References: mainland Portugal[▲] (?). According to Eschmeyer *et al.* (2019) the current geographic distribution of *B. auritus* is the Eastern Atlantic, from Morocco to Angola. The most northern occurrence record of *B. auritus* is a specimen deposited in the National Museum of Natural History, Smithsonian Institution (USNM 400916, 1969) caught off west of Larache (Morocco). Some specimens presumably of this species caught off by an artisanal fishing vessel was reported in Quarteira auction (Algarve) in 2017. This occurrence is questionable and needs further confirmation. Thus, this species is not considered in the total number of valid species.

Parapristipoma humile (Bowdich 1825)

Common name: Guinean grunt; **Roncador-canela**, Canela, Cherna, Cherne, Mulato, Pargo, Pargo-mulato, Peixe-canela.

References: mainland Portugal[▲] ([Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Carneiro et al., 2014](#)) and Madeira ([Bowdich, 1825](#); [Günther, 1859](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Roux, 1979](#); [Eschmeyer, 1998](#); [Carneiro et al., 2014](#); [Russell, 2014a](#); [Carpenter and Johnson, 2016](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Parapristipoma octolineatum* (Valenciennes 1833) ([Carpenter and Johnson, 2016](#) and [Sadeghi, 2019](#)). The distribution map of this species reported by Russell (2014a) partially included the area of Selvagens Islands (Madeira EEZ).

Parapristipoma octolineatum (Valenciennes 1833)

Common name: **African striped grunt; Riscado**, Ajenil, Burro, Canela, Cherna, Cherne, Mulato, Pargo, Pargo-mulato, Peixe-burro, Peixe-canela, Peixe-romão, Pombo, Remão, Riscadinho, Romão, Roncador, Roncador-riscadinho, Roncador-riscado, Xaputa.

References: mainland Portugal[▲] ([Bragança, 1903](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Roux, 1979, 1981](#); [Fischer et al., 1981, 1987](#); [Ben-Tuvia and McKay, 1986](#); [Erzini et al., 2001a](#); [Santos et al., 2002a, c](#); [ICES, 2005](#); [Carneiro et al., 2014](#); [Russell, 2014](#); [Carpenter and Johnson, 2016](#); [Bañón et al., 2019b](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#); [Sadeghi, 2019](#)) and Madeira ([Noronha and Sarmento, 1934](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Fischer et al., 1981](#); [Roux, 1981](#); [Ben-Tuvia and McKay, 1986](#); [Lloris et al., 1991](#); [Falcón et al., 2000](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Russell, 2014](#); [Almada et al., 2015](#); [Carpenter and Johnson, 2016](#); [POGIS, 2016](#); [Bañón et al., 2019b](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Parapristipoma humile* (Bowdich 1825) ([Carpenter and Johnson, 2016](#) and [Sadeghi, 2019](#)).

Plectorhinchus mediterraneus (Guichenot 1850)

Common name: **Rubberlip grunt; Pombo**, Ajenil, Algenil, Angelina, Anjomil, Anjonil, Anjouil, Anjoumil, Anjounil, Asmil, Asno, Burro, Chaputa, Cherna, Cherne, Cherne-do-alto, Cherne-do-Cabo-Branco, Leopard, Leote, Matateu, Mulato, Mulato-legítimo, Pargo, Pargo-liso, Pargo-mulato, Peixe-batata, Peixe-burro, Peixe-mulato, Roncador-mulato, Roncador-pombo, Xaputa.

References: mainland Portugal[▲] ([Guimarães, 1882b](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1897, 1898](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Roux, 1979](#); [Fischer et al., 1981, 1987](#); [Roux, 1981](#); [Ben-Tuvia and McKay, 1986](#); [Martins et al., 1992](#); [Heemstra, 1995](#); [Erzini et al., 2001a](#); [Santos et al., 2002a](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#);

Russell, 2014c; Carpenter and Johnson, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019; Sadeghi, 2019) and Madeira (Osório de Castro, 1967). The holotype (unique) of *Pseudohelotes guntheri* Guimarães 1882, synonym of *P. mediterraneus*, caught off in Setúbal (mainland Portugal) is apparently destroyed. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. Since Osório de Castro (1967) no other author describes the presence of this species in Madeira and its occurrence in this EEZ was not yet confirmed by more recent data. However, its occurrence in Canary Island is confirmed.

Pomadasys incisus (Bowdich 1825)

Common name: **Bastard grunt; Roncador-bravura**, Bejigó, Besugo, Bisugo, Bolo-bolo, Bravura, Bujigú, Cacòrrecór, Cachucho-preto, Cocòrrecór, Combolobolo, Dibolo-bolo, Garoupa, Músico, Peixe-bravura, Roncador, Roncôlho, Ronqueirão, Roqueirão, Salmão.

References: mainland Portugal (Erzini *et al.*, 1997; Moura, 2004; Henriques *et al.*, 2008; Bañón *et al.*, 2014; Carneiro *et al.*, 2014; Russell, 2014d), Azores[▲] (Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Russell, 2014d) and Madeira[▲] (Bowdich, 1825; Lowe, 1837; NHMUK 1860.7.14.7, 1854-1861; Günther, 1859; de Velloso, 1892; Fowler, 1936a, b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Roux, 1979, 1981; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Ben-Tuvia and McKay, 1986; Hureau, 1991; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Delgado, 1998; Mello, 2000; Severino, 2004; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; Bañón *et al.*, 2014; Carneiro *et al.*, 2014; Russell, 2014d; Carpenter and Johnson, 2016; POGID, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The species *Anomalodon incisus* Bowdich 1825 described for Madeira Island for the first time is synonym of *P. incisus*. The holotype *Pristipoma bennettii* Lowe 1838, synonym of *P. incisus*, caught off in Madeira is in the NHMUK collections (NHMUK 1860.7.14.7).

Pomadasys perotaei (Cuvier 1830)

Common name: **Parrot grunt; Roncador-papagaio**, Camandombe, Roncador.

References: Madeira (Fischer *et al.*, 1981; Roux, 1981). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. perotaei* in the Eastern Atlantic is from Mauritania to Angola and Cape Verde Islands. This species is relatively widespread in tropical Eastern Atlantic, and is considered relatively common (de Morais *et al.*, 2015b) and probably often confused with other *Pomadasys* species (Fischer *et al.*, 1981). The distribution map of this species reported by Fischer *et al.* (1981) includes the Madeira area. However, since Fischer *et al.* (1981) no other author describes the presence of this species in Madeira and its occurrence in this EEZ was not yet confirmed by more recent data.

Family Sparidae

Boops boops (Linnaeus 1758)

Common name: **Bogue; Boga-do-mar**, Boga, Boga-legítima, Boga-pintada, Bonga, Carapau, Choupa, Pintada, Ruama.

References: mainland Portugal[▲] (Carvalho, 1765; de Brito Capello, 1867d, 1869b; Steindachner, 1867; Posser, 1880; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, c, 1903, 1935; Osório, 1896; Vieira, 1898; Seabra, 1911; Osório de Castro, 1954, 1967; Tortonese, 1979e; Dinis, 1981; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982a, b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Bauchot and Hureau, 1986; Antunes *et al.*, 1988; Hureau, 1991; Martins *et al.*, 1992, 1999, 2015; Almeida, 1996; Erzini *et al.*, 1996, 2001a, b; Gonçalves *et al.*, 1996, 1997; Cabral, 1999; Gomes *et al.*, 2001; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Santos *et al.*, 2002a; Pista *et al.*, 2003; Garnerot *et al.*, 2004; Mendes *et al.*, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Stergiou *et al.*, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Neves *et al.*, 2008; Silva, 2008, 2010, 2011; Batista *et al.*, 2009; Silva *et al.*, 2009; Gonçalves *et al.*, 2010; Costa *et al.*, 2012; Ribeiro *et al.*, 2012; Carneiro *et al.*, 2014; Silva and Borges, 2014; OCEANA, 2014; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Faria e Silva, 1892; Collett, 1896a; Regan, 1903a; Sampaio, 1904; Roule, 1919a; Vaillant, 1919; Nobre, 1924, 1930; Fowler, 1919, 1936; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood and Williams, 1974; Wheeler, 1978; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Tortonese, 1979e; Martins, 1982a, b; INIP,

1985a, c; Bauchot and Hureau, 1986; Dias and Monteiro, 1986; Isidro, 1988, 1990a, b; Nash *et al.*, 1990, 1994a; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, b, 1995a, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991; Dias, 1991; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Krug, 1995; Silva *et al.*, 1994, 1995; Azevedo, 1995; Pereira, 1995; Barreiros, 1995; Gonçalves *et al.*, 1995; Pinho *et al.*, 1995; Santos and Nash, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso, 2016; Almeida and Biscoito, 2019; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Cuvier and Valenciennes, 1830; Lowe, 1837, 1849; Gervais and Boulart, 1877b; Posser, 1880; Vinciguerra, 1883; de Velloso, 1892; Fowler, 1923; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; Augier, 1985; Bauchot and Hureau, 1986; Hureau, 1991; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Floeter *et al.*, 2008; Ribeiro *et al.*, 2005b; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Carpenter and Iwatsuki, 2016; POGID, 2016; POGIS, 2016; Friedlander *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Centracanthus cirrus Rafinesque 1810

Common name: **Curled picarel; Boqueirão**, Bocairão, Bogarão, Mãe-da-boga.

References: mainland Portugal (de Brito Capello, 1867a, d, 1880; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Tortonese *et al.*, 1979; Fischer *et al.*, 1981, 1987; Dooley *et al.*, 1985; Tortonese, 1986c; Antunes *et al.*, 1988; Heemstra, 1990; Azevedo *et al.*, 2004; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Regan, 1903; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Tortonese *et al.*, 1979; Fischer *et al.*, 1981; Heemstra, 1981, 1990; Dooley *et al.*, 1985; Shcherbachov *et al.*, 1985a; Tortonese, 1986c; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Kukuev, 2002 (PECS); Azevedo *et al.*, 2004; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Cuvier and Valenciennes, 1830; Lowe, 1837; Günther, 1859; Gervais and Boulart, 1877b; de Velloso, 1892; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Tortonese *et al.*, 1979; Fischer *et al.*, 1981; Heemstra, 1981, 1990; Dooley *et al.*, 1985; Tortonese, 1986c; Lloris *et al.*, 1991; Azevedo *et al.*, 2004; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Christiansen *et al.*, 2009; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; POGID, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Dentex canariensis Steindachner 1881

Common name: **Canary dentex; Dentão-das-Canárias**, Capatão, Chacarona, Chancarina, Chancarona, Dentão, Dentão-quissanga, Pargo, Pargo-dentão, Pargo-quissanga, Quissanga.

References: mainland Portugal (Martins *et al.*, 1992; Erzini *et al.*, 2001a; Santos *et al.*, 2002c; Carneiro *et al.*, 2014; Martins and Carneiro, 2018).

Dentex dentex (Linnaeus 1758)

Common name: **Common dentex; Capatão-legítimo**, Bombeiro, Cachucho, Capatão, Capatão-de-catula, Catula, Corcunda, Dentão, Dentelha, Dentudo, Mitrado, Pargo, Pargo-capatão, Pargo-cataludo, Pargo-catutudo, Pargo-corcunda, Pargo-cotuludo, Pargo-dentelha, Pargo-de-batata, Pargo-de-catuta, Pargo-de-corcunda, Pargo-de-cotula, Pargo-de-marreca, Pargo-de-mitra, Pargo-de-moiro, Pargo-de-morro, Pargo-de-mouro, Pargo-dentelha, Pargo-marreco, Pargo-mitrado, Pargo-moiro, Pargo-mouro, Roncador.

References: mainland Portugal[▲] (Carvalho, 1765(?); de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Posser, 1880; Osório, 1888; Baldaque da Silva, 1891; Vieira, 1897, 1898; Seabra, 1911; Nobre, 1935; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Martins, 1979; Tortonese, 1979e; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982b; Bauchot and Hureau, 1986; Bauchot, 1987; Erzini *et al.*, 2001a; Santos *et al.*, 2002a; ICES, 2005; Henriques *et al.*, 2008; Batista *et al.*, 2009; França *et al.*, 2011; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores (MCZ 10640, 1859; Cadenat, 1937; Wheeler, 1978; Arruda, 1997; Carneiro *et al.*, 2014) and Madeira (Günther, 1859; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; Dooley *et al.*, 1985;

Bauchot and Hureau, 1986, 1990; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Dentex gibbosus (Rafinesque 1810)

Common name: **Pink dentex**; **Capatão-de-bandeira**, Bandeireiro, Brasil, Capatão, Capatão-de-catuta, Capatão-de-cotula, Corcunda, Mariana, Pargo, Pargo-bandeira, Pargo-capatão, Pargo-catulo, Pargo-cotuludo, Pargo-de-bandeira, Pargo-de-bandeira-legítimo, Pargo-de-catula, Pargo-de-cotula; Pargo-capelo, Pargo-cotulado, Pargo-de-mitra, Padro-mitrado, Pargo-de-morro, Pargo-do-morro, Pargo-mitrado, Pargo-mouro, Sama, Samo.

References: mainland Portugal (de Brito Capello, 1867d, 1869b, 1880; Vinciguerra, 1883; Osório, 1888; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1935; Bragança, 1902; Seabra, 1911; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Borges, 1984; Dooley *et al.*, 1985; Bauchot and Hureau, 1986; Bauchot and Hureau, 1990; Martins *et al.*, 1992; Erzini *et al.*, 1996, 1997, 2001a; Santos *et al.*, 2002a; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Fischer *et al.*, 1981; Wirtz, 1994; Freitas and Araújo, 2006; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; POGID, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The distribution map of this species reported by Fischer *et al.* (1981) partially included the area of Selvagens Islands (Madeira EEZ).

Dentex macrophthalmus (Bloch 1791)

Common name: **Large-eye dentex**; **Cachucho**, Breca, Cachucha, Chucho, Salvador, Tambor, Vermelhofundo.

References: mainland Portugal[▲] (de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Posser, 1880; Vinciguerra, 1883; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1935; Goode and Bean, 1895, 1896; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Martins, 1979; Tortonese, 1979e; INIP, 1982b; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Borges, 1984; Dinis, 1985; Dooley *et al.*, 1985; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Erzini *et al.*, 2001a; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; Henriques *et al.*, 2008; Batista *et al.*, 2009; Carneiro *et al.*, 2014; Costa, 2014; Russell, 2014e; Carpenter and Iwatsuki, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Russell, 2014e; Carpenter and Iwatsuki, 2016). The distribution map of this species reported by Russell (2014e) and Carpenter and Iwatsuki (2016) partially included the area of Selvagens Islands (Madeira EEZ).

Dentex maroccanus Valenciennes 1830

Common name: **Morocco dentex**; **Cachucho-dentão**, Cachucho, Dentão, Dentão-de-Marrocos, Dentelha, Pargo.

References: mainland Portugal[▲] (de Brito Capello, 1867a, d, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1898; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Krefft and Bekker, 1979; Martins, 1979; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Borges, 1984; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Erzini *et al.*, 2001a; Borges *et al.*, 2002; ICES, 2005; Ribeiro *et al.*, 2006, 2012; Henriques *et al.*, 2008; Batista *et al.*, 2009; Carneiro *et al.*, 2014; Russell, 2014f; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967(?)) and Madeira (?) (Osório de Castro, 1967(?)). The unique holotype of *Dentex parvulus* de Brito Capello 1867a was preserved in the Museu Bocage and destroyed in the 1978 fire, and it is considered as synonym of *Dentex maroccanus* Valenciennes 1830. Osório de Castro (1967) mentioned the presence of this species in the fishing grounds of the Azores and Madeira as doubtful. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores or Madeira islands. Thus, this species is not considered in the total number of valid species for the Azores and Madeira.

Diplodus annularis (Linnaeus 1758)

Common name: **Annular seabream**; **Sargo-alcorraz**, Acarraz, Albricoque, Alcarraz, Alcornoque, Alcorraz, Alvar, Choupa, Choupo, Macharra, Micharra, Micharra-alvar, Mocharra, Mojarra, Mucharra, Mucharra-alvar, Mujarra, Salema, Sargo-salema, Sargueta, Sarguete.

References: mainland Portugal▲ (de Brito Capello, 1869a, b, 1880; Posser, 1880; Almeida and Roquette, 1892; Osório, 1896; Vieira, 1897, 1898; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Hureau and Monod, 1979; Tortonese, 1979e; Fischer et al., 1981, 1987; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Martins et al., 1992, 2015; Almeida, 1996; Erzini et al., 1996, 1997, 2001a; Cabral, 1999; Borges et al., 2002, 2003; Santos et al., 2002a, c; Palma et al., 2003; Garnerot et al., 2004; Ribeiro et al., 2006, 2008, 2012; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Batista et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Costa et al., 2012; Carneiro et al., 2014; Russell, 2014g; Fricke et al., 2016b; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Bauchot et al., 1981; Fischer et al., 1981; Dooley et al., 1985; Bauchot and Hureau, 1986, 1990; Lloris et al., 1991; Andrade and Albuquerque, 1995; Wirtz et al., 2008(?); Carneiro et al., 2014; Russell, 2014g; Carpenter and Iwatsuki, 2016; Fricke et al., 2016b; Eschmeyer et al., 2019; Froese and Pauly, 2019). According to Wirtz et al. (2008) the record of this species from Madeira is doubtful. The distribution map of this species reported by Russell (2014g) partially included the area of Selvagens Islands (Madeira EEZ).

***Diplodus bellottii* (Steindachner 1882)**

Common name: **Senegal seabream; Sargo-do-Senegal**, Choupa, Choupo, Mucharra, Sargo, Sargueta.

References: mainland Portugal (Martins et al., 1992, 2015; Erzini et al., 1996, 1997, 2001a; Gonçalves et al., 1996, 1997; Cabral, 1999; Borges et al., 2002; Santos et al., 2002a, c; Palma et al., 2003; Prista et al., 2003; Thiel et al., 2003; ICES, 2005; Ribeiro et al., 2006, 2008, 2012; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Neves et al., 2008; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Carneiro et al., 2014; Martins and Carneiro, 2018; Baptista et al., 2019; Froese and Pauly, 2019) and Madeira (MNHN-IC-1972-0078, 1972(?); MNHN-IC-1972-0079, 1972(?); MNHN-IC-1978-0648, 1978).

***Diplodus cadenati* de la Paz, Bauchot & Daget 1974**

Common name: White seabream; **Sargo-legítimo-do-Atlântico**, Chamoa, Mariquita, Olho-de-boi, Pintagola, Sargo, Sargo-chamôa, Sargo-comum, Sargo-legítimo, Sargo-vulgar.

References: mainland Portugal (MNHN-IC-1973-0011, n.d.; Bauchot et al., 1981; Fischer et al., 1981; Bauchot and Hureau, 1990; ICES, 2005; Carneiro et al., 2014; Fricke et al., 2016b; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Regan, 1903a; Sampaio, 1904; Lampe, 1914; Vaillant, 1919; Nobre, 1924, 1930; Fowler, 1919, 1936b; Ribeiro, 1936; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Wood, 1973; Mapstone and Wood, 1974, 1975; Wood, 1974; Wood and Williams, 1974; de la Paz et al., 1974; de la Paz, 1975; Dias et al., 1976; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Muzanor and Westhaus, 1981; Martins, 1982a; Bauchot and Hureau, 1986; Dias and Monteiro, 1986; Barreiros, 1990, 1995; Nash et al., 1990, 1994a; Patzner and Santos, 1990d, 1993; Patzner et al., 1990, 1992; Santos et al., 1990b, 1994a, b, 1995a, 1997; Wirtz, 1990a, 1994; Azevedo et al., 1991, 1992; Dias, 1991; Arruda et al., 1992; Santos, 1992a, 1995b; Gonçalves, 1993; Azevedo, 1995; Gonçalves et al., 1995; Santos and Nash, 1995; Silva and Menezes, 1995a; Silva et al., 1995; Isidro, 1996; Menezes, 1996; Harmelin-Vivien et al., 2001; Morato et al., 2001a, b; Cardigos et al., 2005; Menezes et al., 2006; Rosa et al., 2006; Floeter et al., 2008; Carneiro et al., 2014; Afonso, 2016; Fricke et al., 2016b; Almeida and Biscoito, 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Bauchot et al., 1981; Fischer et al., 1981; Dooley et al., 1985; Bauchot and Hureau, 1986; Lloris et al., 1991; Floeter et al., 2008; Carneiro et al., 2014; Carpenter and Iwatsuki, 2016; Fricke et al., 2016b; Eschmeyer et al., 2019; Froese and Pauly, 2019). This species is often confused with *Diplodus sargus* (Linnaeus 1758).

***Diplodus cervinus* (Lowe 1838)**

Common name: **Zebra seabream; Sargo-veado**, Choupa, Choupo, Mariquita, Mariquita-da-pedra, Marquita, Olho-de-boi, Peixe-zebra, Pintagola, Raiado, Riscado, Saima, Saime, Sargo, Sargo-branco, Sargo-mariquita, Sargo-preto, Sargo-raiado, Sargo-riscado, Sargueta, Sarguete, Seima, Zebra.

References: mainland Portugal▲ (de Brito Capello, 1867a, 1880; Vieira, 1897, 1898; Seabra, 1911; Nobre, 1935; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Fischer et al., 1987; Antunes et al., 1988; Martins et al., 1993, 2015; Almeida, 1996; Erzini et al., 1996, 1997, 2001a; Gonçalves et al., 1996, 1997; Santos et al., 2002a; ICES, 2005; Ribeiro et al., 2006, 2008; Henriques et al., 2007, 2008; Gomes and Olim, 2007; Gonçalves et al., 2010; Carneiro et al., 2014; Russell, 2014h; Fricke et al., 2016b; Martins and Carneiro,

2018; Froese and Pauly, 2019), Azores (?) (Collins, 1954; Arruda, 1997(?); Santos *et al.*, 1997(?)) and Madeira▲ (Lowe, 1837; Günther, 1859; de Brito Capello, 1871; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; INIP, 1980; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; Augier, 1985; Dooley *et al.*, 1985; Bauchot and Hureau, 1986; Hureau, 1991; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Severino, 2004; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Russell, 2014h; Almada *et al.*, 2015; Carpenter and Iwatsuki, 2016; Fricke *et al.*, 2016b; POGID, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Collins (1954) cites *Diplodus trifasciatus* (Rafinesque 1810), a common synonym for *D. cervinus*, based on a specimen that he observed at the Museum of Ponta Delgada (not found in the MCM collections) (Santos *et al.*, 1997). Arruda (1997) and Santos *et al.* (1997) consider doubtful the citations of Collins (1954). According to Santos *et al.* (1997) these citations are probable misidentifications of *D. cadenati*. However, Russell (2014h) considered the *Diplodus cervinus* complex needs a taxonomic review. Thus, this species is not considered in the total number of valid species for the Azores.

***Diplodus fasciatus* (Valenciennes 1830)**

Common name: **Banded seabream; Sargo-de-faixas**, Choupa, Choupo, Mariquita, Mariquita-da-pedra, Marquita, Olho-de-boi, Peixe-zebra, Pintagola, Raiado, Riscado, Saima, Saime, Sargo, Sargo-branco, Sargo-mariquita, Sargo-negro-de-Cabo-Verde, Sargo-preto, Sargo-preto-de-Cabo-Verde, Sargo-raiado, Sargo-riscado, Sargo-veado, Seima, Zebra.

References: mainland Portugal (?) (Steindachner, 1867; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Nobre, 1935; Osório de Castro, 1967), Azores (?) (Osório de Castro, 1967(?)) and Madeira (?) (Osório de Castro, 1967). This species can be confused with *Diplodus cervinus* (Lowe 1838). According to Fricke *et al.* (2016b), Carpenter and Iwatsuki (2016), Froese and Pauly (2019) and Eschmeyer *et al.* (2019) the current geographic distribution of *D. fasciatus* is the Cape Verde Islands (as endemic species). Osório de Castro (1967) mentions the presence of this species as rare in the fishing grounds of mainland Portugal and Madeira without indicating the geographic location. For the Azores, Osório de Castro (1967) indicated its occurrence as doubtful. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores or in Madeira. Thus, this species is not considered in the total number of valid species.

***Diplodus lineatus* (Valenciennes 1830)**

Common name: White seabream; **Sargo-de-Cabo-Verde**, Bicudo, Choupa, Choupo, Olho-de-boi, Sargo, Sargo-bicudo, Sargueta.

References: mainland Portugal (?) (MNHN-IC-A-8102(?), 1830; Vieira, 1897, 1898; Seabra, 1911; Nobre, 1935(?)). One specimen of *Sargus lineatus* Cuvier and Valenciennes 1830 was caught off on January, 28 1893 (southern Portugal, Faro) (Vieira, 1897, 1898). This specimen is preserved in the Zoology Gallery of the Museum of Science of the Coimbra University, catalogue number ZOO.0006093 (dry), under the identification of *Diplodus sargus* (L.), which should probably been labelled as subspecies *lineatus* (?). The subspecies *Diplodus sargus lineatus* (Valenciennes 1830) is currently valid as *Diplodus lineatus* (Valenciennes 1830). However, for taxonomic conclusion, the molecular analysis confirmation of this specimen is needed. The holotype MNHN-IC-A-8102 (unknown origin: Lisbon ?).

***Diplodus puntazzo* (Walbaum 1792)**

Common name: **Shaptnout seabream; Sargo-bicudo**, Palharete, Sargo, Sidra.

References: mainland Portugal▲ (de Brito Capello, 1869a, b, 1880; Girard, 1892; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Bauchot and Hureau, 1986, 1990; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Almeida, 1996; Erzini *et al.*, 1997, 2001a; Santos *et al.*, 2002a; ICES, 2005; Ribeiro *et al.*, 2006, 2008, 2012; Henriques *et al.*, 2007, 2008; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Russell, 2014i; Martins *et al.*, 2015; Fricke *et al.*, 2016b; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Russell, 2014i). The distribution map of this species reported by Russell (2014i) partially included the area of Selvagens Islands (Madeira EEZ).

Diplodus sargus (Linnaeus 1758)

Common name: White seabream; **Sargo-legítimo-do-Mediterrâneo**, Bicudo, Chamôa, Choupa, Choupo, Mariquita, Marquita, Olho-de-boi, Pintagola, Safia, Sargo, Sargo-bicudo, Sargo-branco, Sargo-chamôa, Sargo-comum, Sargo-legítimo, Sargo-mariquita, Sargo-marquita, Sargo-vulgar, Sargueta, Sarguêta, Sarguete, Sefia, Séfia, Seifia.

References: mainland Portugal▲ (?) (Vandelli, 1771; de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Gervais and Boulart, 1877b; Posser, 1880; Osório, 1888; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895c, 1904, 1935; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Fernández et al., 1981; INIP, 1982b; Martins, 1981; Borges, 1984; Fischer et al., 1987; Antunes et al., 1988; Martins et al., 1991, 1999, 2015; Almeida, 1996; Erzini et al., 1996, 1997, 2001a; Gonçalves et al., 1996, 1997, 2007; Cabral, 1999; Jorge et al., 2002; Santos et al., 2002a, c; Palma et al., 2003; Prista et al., 2003; Thiel et al., 2003; Garnerot et al., 2004; Ribeiro et al., 2006, 2008, 2012; Henriques et al., 2007, 2008; Neves et al., 2008; Batista et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Costa et al., 2012; Carneiro et al., 2014; Martins and Carneiro, 2018; Baptista et al., 2019; Froese and Pauly, 2019), Azores▲ (?) (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Regan, 1903; Lampe, 1914; Vaillant, 1919; Fowler, 1919, 1936b; Nobre, 1924, 1930; Ribeiro, 1936; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood, 1973, 1974; Mapstone and Wood, 1974, 1975; Wood and Williams, 1974; Dias et al., 1976; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Tortonese, 1979e; Muzanor and Westhaus, 1981; Martins, 1982a; INIP, 1985a; Dias and Monteiro, 1986; Barreiros, 1990; Nash et al., 1990; Patzner and Santos, 1990d; Patzner et al., 1990; Santos et al., 1990b, 1994a, b, 1997(?); Wirtz, 1990a; Azevedo et al., 1991, 1992; Dias, 1991; Arruda, 1992, 1997; Gonçalves, 1993, 1993; Wirtz, 1994; Azevedo, 1995; Santos and Nash, 1995; Silva and Menezes, 1995a; Silva et al., 1995; Isidro, 1996; Menezes, 1996; Morato-Gomes et al., 1998; Morato et al., 2001b; Bertoncini et al., 2010; Botelho et al., 2010; Porteiro et al., 2010; Diogo and Pereira, 2013, 2014; Carneiro et al., 2014; Afonso, 2016; Fauconnet et al., 2019; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1837, 1839, 1840b, 1844a, b, 1849; Günther, 1859; de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Augier, 1985; Hureau, 1991; Abreu et al., 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón et al., 2000; Mello, 2000; Araújo et al., 2005; Ribeiro et al., 2005b; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; POGID, 2016; POGIS, 2016; Friedlander et al., 2017; Eschmeyer et al., 2019; Froese and Pauly, 2019). According to Eschmeyer et al. (2019) the geographic distribution of *D. sargus* is the Mediterranean Sea, Black Sea and in the Eastern Atlantic (Madeira). However, Fricke et al. (2016b) mention the current geographic distribution only in the Mediterranean Sea (except Israel) and the Black Sea. Probably part of the citations of this species to the Eastern Atlantic are misidentifications of *Diplodus cadenati* de la Paz, Bauchot and Daget 1974. According to Santos et al. (1997) all the citations to the Azores are synonyms, misspellings and misidentifications of *Diplodus cadenati* de la Paz, Bauchot and Daget 1974. However, after Santos et al. (1997) several authors cite this species to the Azores and there are records from commercial catches and direct observation by underwater fish visual census. These citations are probable also synonyms, misspellings and misidentifications of *D. cadenati* as mentioned by Santos et al. (1997). This subject deserves further detailed attention.

Diplodus vulgaris (Geoffroy St. Hilaire 1817)

Common name: **Common two-banded seabream**; **Sargo-safia**, Alcorraz, Choupa, Choupo, Cruta, Macharra, Micharra, Mucharra, Mujarra, Olho-de-boi, Palmeiro, Pargo, Pintagola, Ruçada, Safia, Sargo, Sargo-amarelo, Sargo-negro, Sargo-salema, Sargueta, Sarguêta, Sefia, Séfia, Seifia.

References: mainland Portugal▲ (Carvalho, 1756; de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Osório, 1888, 1896; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1903, 1935; Vieira, 1897, 1898; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Martins, 1979; Tortonese, 1979e; Bauchot et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Bauchot and Hureau, 1986, 1990; Antunes et al., 1988; Arruda et al., 1988; Hureau, 1991; Almeida, 1996; Erzini et al., 1996, 1997, 2001a; Gonçalves et al., 1996, 1997; Cabral, 1999; Martins et al., 1999, 2015; Gomes et al., 2001; Borges et al., 2002; Jorge et al., 2002; Santos et al., 2002a, c; Palma et al., 2003; Prista et al., 2003; Garnerot et al.,

2004; Mendes *et al.*, 2004; Moura, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Ribeiro *et al.*, 2006, 2008, 2012; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Neves *et al.*, 2008; Silva, 2008; Batista *et al.*, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Fricke *et al.*, 2016b; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Sampaio, 1904; Nobre, 1919, 1924, 1930; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; MNHN-IC-2016-0611, 1971; Ré, 1979a; Tortonese, 1979e; Arruda, 1992, 1997; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Afonso, 2016) and Madeira[▲] (Lowe, 1849; Günther, 1859; de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; INIP, 1982c, 1984b; Augier, 1985; Dooley *et al.*, 1985; Bauchot and Hureau, 1986; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Afonso, 2016; Carpenter and Iwatsuki, 2016; Fricke *et al.*, 2016b; POGID, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Evynnis ehrenbergii* (Valenciennes 1830)**

Common name: **Bluespotted seabream**; **Pargo-ruço-de-Ehrenberg**, Pargo, Pargo-amarelo, Pargo-boi, Pargo-boio, Pargo-branco, Pargo-de-pintas-azuis, Pargo-enxofre, Pargo-ruço, Pargueta, Parguete, Ruço, Senapa, Senapai, Sinapa, Sinapaio, Solapa.

References: mainland Portugal (?) (Tortonese, 1979e). Cuvier and Valenciennes (1830) described this species for the first time in the Eastern Mediterranean Sea, having attributed the name *Pagrus ehrenbergii* Valenciennes 1830. Tortonese (1979e) described from mainland Portugal *E. ehrenbergii*, of which there are no further reports. Eschmeyer *et al.* (2019) considers *Evynnis ehrenbergii* (Valenciennes 1830) a valid species and indicates that the current geographic distribution is the Mediterranean Sea and west African coast (if valid). Probably, *E. ehrenbergii* is replaced by *Pagrus caeruleostictus* (Valenciennes 1830) in the West African coast. Thus, this species is not considered in the total number of valid species.

***Lithognathus mormyrus* (Linnaeus 1758)**

Common name: **Sand steenbras**; **Ferreira**, Besugo, Besugo-de-ova, Besugo-dova, Besugo-trombeiro, Besugo-trombeteiro, Besugo-trombudo, Bezugo-d'ova, Bisugo, Ferreira, Fureira, Garapau, Nacho, Peixão, Riscado, Sargo-de-areia, Trombeiro, Trombudo, Vesugo, Vesugo-da-ova, Vesugo-dova, Vesugo-trombeiro, Vesugo-trombeteiro, Vesugo-trombudo, Vezugo-trombudo.

References: mainland Portugal[▲] (Carvalho, 1765; de Brito Capello, 1967a, d, 1869b, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1903, 1935; Vieira, 1897, 1898; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Bauchot and Hureau, 1986, 1990; Martins *et al.*, 1992, 1999, 2015; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Santos *et al.*, 2002a, c; Palma *et al.*, 2003; Sousa *et al.*, 2005; Ribeiro *et al.*, 2006, 2008; Gomes and Olim, 2007; Henriques *et al.*, 2008; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Russell, 2014j; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967) and Madeira[▲] (Osório de Castro, 1954, 1967; Alves and Alves, 2002; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Russell, 2014j; Carpenter and Iwatsuki, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). According to Eschmeyer *et al.* (2019) the current geographic distribution of *L. mormyrus* in the Eastern Atlantic is from the Bay of Biscay to South Africa, including Madeira and Canary Islands. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores. Thus, this species is not considered in the total number of valid species for the Azores.

***Oblada melanura* (Linnaeus 1758)**

Common name: **Saddled seabream**; **Dobrada**, Dobrada, Galana, Judeu.

References: mainland Portugal[▲] (Vandelli, 1771; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Gonçalves *et al.*, 1996, 1997; Borges *et al.*, 2002, 2003;

Henriques *et al.*, 2007, 2008; Ribeiro *et al.*, 2008; Carneiro *et al.*, 2014; Russell, 2014k; Martins *et al.*, 2015; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837, 1849; Günther, 1859; de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; Augier, 1985; Dooley *et al.*, 1985; Bauchot and Hureau, 1986, 1990; Hureau, 1991; Lloris *et al.*, 1991; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Severino, 2004; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Russell, 2014k; Carpenter and Iwatsuki, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Pagellus acarne* (Risso 1827)**

Common name: **Axillary seabream; Besugo**, Besugo-legítimo, Besujo, Bezugo, Bisugo, Garapau, Vesugo. References: mainland Portugal[▲] (de Brito Capello, 1867d, 1869a, b, 1880; Steindachner, 1867; Posser, 1880; Nobre, 1894, 1895a, 1903, 1935; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1898; Seabra, 1911; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, b, 1982a, b; Martins, 1981; Borges, 1984; Dinis, 1985; Bauchot and Hureau, 1986, 1990; Martins *et al.*, 1992, 1999, 2015; Gonçalves *et al.*, 1996, 1997; Erzini *et al.*, 1997, 2001a; Cabral, 1999; Gomes *et al.*, 2001; Borges *et al.*, 2002, 2003; Jorge *et al.*, 2002; Santos *et al.*, 2002a, c; Prista *et al.*, 2003; Mendes *et al.*, 2004; ICES, 2005; Sousa *et al.*, 2005, 2006; Stergiou *et al.*, 2006; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Ribeiro *et al.*, 2008; Silva, 2008, 2010, 2011; Batista *et al.*, 2009; Silva *et al.*, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Russell, 2014l; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Sampaio, 1904; Roule, 1919a; Vaillant, 1919; Nobre, 1924, 1930; Fowler, 1919, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Harmelin and Harmelin-Vivien, 1979; Tortonese, 1979e; Muzanor and Westhaus, 1981; Martins, 1982a; Dooley *et al.*, 1985; INIP, 1985a, b, c; Dias and Monteiro, 1986; Dias *et al.*, 1990; Patzner *et al.*, 1990, 1992; Dias, 1991; Arruda, 1992, 1997; Santos, 1992a; Krug, 1995; Nash *et al.*, 1994a; Santos *et al.*, 1994a, 1995a, 1997; Silva *et al.*, 1994, 1995; Wirtz, 1994; Azevedo, 1995; Gonçalves *et al.*, 1995; Pinho *et al.*, 1995; Santos and Nash, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Morato-Gomes *et al.*, 1998; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a; Barreiros *et al.*, 2002; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Russell, 2014l; Afonso, 2016; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837, 1839, 1840b, 1844a, 1849; Gervais and Boulart, 1877b; de Velloso, 1892; Vaillant, 1919; Nunes, 1953, 1974; Osório de Castro, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; INIP, 1982c, 1984b, c, 1985c; Augier, 1985; Dooley *et al.*, 1985; Bauchot and Hureau, 1986, 1990; Lloris *et al.*, 1991; Ferreira *et al.*, 2008; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Russell, 2014l; Carpenter and Iwatsuki, 2016; POGID, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Pagellus bellottii* Steindachner 1882**

Common name: **Red pandora; Bica-buço**, Bica, Bica-bicuda, Bicanço, Bicudo, Breca, Breca-bica, Breca-colorada, Buço, Malaguita, Pargo-bico, Pargo-bicudo, Salvador, Tico-tico, Vermelho-fundo.

References: mainland Portugal[▲] (SAIAB FC 4316, 1957; Osório de Castro, 1967; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Santos *et al.*, 2002a; Henriques *et al.*, 2008; Carneiro *et al.*, 2014; Martins *et al.*, 2015; Baptista *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967) and Madeira (Osório de Castro, 1967; Russell, 2014m). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. bellottii* in the Eastern Atlantic is from the Gibraltar to Angola and some offshore islands. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location. Since Osório de Castro (1967) no other author describes the presence of this species in the Azores. Thus, this species is not considered in the total number of valid species for the Azores.

Pagellus bogaraveo (Brünnich 1768)

Common name: **Blackspot seabream; Goraz**, Besugo, Besugo-legítimo, Bezugo, Bisugo, Buama, Buaninha, Carapau, Garapau, Goraz-de-pinta, Massacato, Massacote, Olho-mole, Olhos-moles, Pacharro, Pachel, Pachelão, Paxelão, Pachorro, Pancho, Paxão, Peixão, Pexão, Vesugo.

References: mainland Portugal▲ (Carvalho, 1765; de Brito Capello, 1867a, d, 1869b, 1873, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895a, c, 1935; Osório, 1895, 1896; Bragança, 1902; Richard, 1908; Seabra, 1911; Murray and Hjort 1912; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Martins, 1979; Tortonese, 1979e; Bauchot et al., 1981; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a, 1982a, b; Bauchot and Hureau, 1986, 1990; Martins et al., 1992; Dinis and Marecos, 1984; Dinis, 1985; Bauchot, 1987; Gonçalves et al., 1997; Erzini et al., 2001a, b; Borges et al., 2002; Santos et al., 2002a; Garnerot et al., 2004; Mendes et al., 2004; ICES, 2005; Sousa et al., 2005, 2006; Fernandes et al., 2007; Henriques et al., 2008; Ribeiro et al., 2008, 2012; Silva, 2008, 2010, 2011; Chaves, 2009; Silva et al., 2009; Chaves and Silva, 2013; Carneiro et al., 2014; Costa, 2014; Silva and Borges, 2014; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores▲ (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Sampaio, 1904; Fowler, 1919, 1936b; Ribeiro, 1936; Cadenat, 1937; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Dias et al., 1976, 1990; Wheeler, 1978; Harmelin and Harmelin-Vivien, 1979; Tortonese, 1979e; Saldanha, 1980; Muzanor and Westhaus, 1981; Martins, 1982a, 1992; Dooley et al., 1985; INIP, 1985a, b, c; Dias and Monteiro, 1986; Krug, 1986, 1989, 1991, 1995, 1998; Silva, 1986a, b, 1992, 1995; Isidro, 1987, 1989, 1996; 1988, 1990a, b; Barreiros, 1990; Patzner, 1990; Patzner et al., 1990, 1992; Ré, 1990; Santos et al., 1990b, 1994a, 1995a, 1997; Wirtz, 1990a; Dias, 1991; Dias and Cascalho, 1991; Monteiro, 1991, 1996; Arruda, 1992, 1997; Azevedo et al., 1992; Pereira, 1992d, 1995; Santos, 1992a; Silva and Krug, 1992, 1995; Nash et al., 1994a; Silva et al., 1994, 1995; Azevedo, 1995; Clarke et al., 1995; Gonçalves et al., 1995; Pinho et al., 1995; Santos and Nash, 1995; Silva and Menezes, 1995a, b; Menezes, 1996; Monteiro et al., 1996; Morato-Gomes et al., 1998; Sousa et al., 1999; Harmelin-Vivien et al., 2001; Morato et al., 2001a; Cardigos et al., 2005; Menezes et al., 2006; Rosa et al., 2006; Floeter et al., 2008; Porteiro et al., 2010, 2013; Besugo, 2013; Pham et al., 2013; Carneiro et al., 2014; Diogo and Pereira, 2014; Afonso, 2016; Torres et al., 2016; Almeida and Biscoito, 2019; Fauconnet et al., 2019; Froese and Pauly, 2019) and Madeira▲ (Lowe, 1837; Von Martens, 1876; de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot et al., 1981; Fischer et al., 1981; INIP, 1984b; Dooley et al., 1985; Bauchot and Hureau, 1986, 1990; Lloris et al., 1991; Araújo et al., 2005; Ferreira et al., 2008; Floeter et al., 2008; Wirtz et al., 2008; Carneiro et al., 2014; Carpenter and Iwatsuki, 2016; POGID, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019).

Pagellus erythrinus (Linnaeus 1758)

Common name: **Common pandora; Bica**, Bica-bicuda, Bicanço, Bicudo, Breca, Breca-bica, Doirada, Malagueta, Salvador, Tico-tico, Vermelho-fundo.

References: mainland Portugal▲ (de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Posser, 1880; Nobre, 1884, 1895a, 1935; Osório, 1888; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954; Martins, 1979; Tortonese, 1979e; INIP, 1981a, 1982b; Bauchot et al., 1981; Fernández et al., 1981; Fischer et al., 1981, 1987; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Martins et al., 1992, 2015; Gonçalves et al., 1996, 1997; Erzini et al., 2001a; Borges et al., 2002; Santos et al., 2002a; Palma et al., 2003; Mendes et al., 2004; ICES, 2005; Sousa et al., 2005; Stergiou et al., 2006; Fernandes et al., 2007; Chaves, 2008, 2009; Henriques et al., 2008; Ribeiro et al., 2008; Batista et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Russell, 2014n; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Fowler, 1936b; Albuquerque, 1954-1956; Wheeler, 1978; IMAR/DOP 3SMI-01-P12BIC, 2012; IMAR/DOP 6FCO-03-P12BIC, 2012; IMAR/DOP 6FCO-01-P12BIC, 2012; IMAR/DOP 6FCO-04-P12BIC, 2012; Carneiro et al., 2014; Martins and Carneiro, 2018; Baptista et al., 2019) and Madeira▲ (Lowe, 1837; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954; Tortonese, 1979e; Bauchot et al., 1981; Fischer et al., 1981; Dooley et al., 1985; Bauchot and Hureau, 1986, 1990; Lloris et al., 1991; Araújo et al., 2005; Ribeiro et al., 2005b; Floeter et al., 2008;

Ribeiro, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Russell, 2014n; Carpenter and Iwatsuki, 2016; POGID, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of *Pagellus rostratus* Lowe 1838, synonym of *P. erythrinus*, caught off in Madeira are unknown.

Pagrus auriga Valenciennes 1843

Common name: **Redbanded seabream; Pargo-sêmola**, Brasino, Mariana, Pargo, Pargo-de-manchas-azuis, Pargo-de-riscas, Pargo-de-riscas-travessas, Pargo-francês, Pargo-rosa, Pargo-seima, Pargo-sémea, Pargo-tereso, Pargueta, Parguete, Seima, Sémia, Sêmola, Tereso, Traquete.

References: mainland Portugal▲ (de Brito Capello, 1872, 1873, 1880; Baldaque da Silva, 1891; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Martins *et al.*, 1992; Erzini *et al.*, 1996, 2001a; Santos *et al.*, 2002a; Palma *et al.*, 2003; Prista *et al.*, 2003; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Neves *et al.*, 2008; Bañón *et al.*, 2014; Carneiro *et al.*, 2014; Russell, 2014o; Carpenter and Iwatsuki, 2016; Martins and Carneiro, 2018; Froese and Pauly, 2019) and Madeira (Günther, 1859; Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1967; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; Bauchot and Hureau, 1986, 1990; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Bañón *et al.*, 2014; Carneiro *et al.*, 2014; Russell, 2014o; Carpenter and Iwatsuki, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). This species is sometimes confused with *Pagrus caeruleostictus* (Valenciennes 1830) (Carpenter and Iwatsuki, 2016). One specimen caught off at the Portuguese continental slope in October 2010 is in the MB collections.

Pagrus caeruleostictus (Valenciennes 1830)

Common name: **Bluespotted seabream; Pargo-ruço**, Pargo, Pargo-amargo, Pargo-boi, Pargo-boio, Pargo-cabeçudo, Pargo-cão, Pargo-de-cão, Pargo-dourado, Pargo-dourado, Pargo-enxofre, Pargo-de-manchas-negra, Pargo-legítimo-vermelho, Pargo-de-pintas-azuis, Pargo-redondo, Pargo-sêmola, Pargo-solapa, Pargo-vermelho, Pargueta, Parguete, Ruço, Sama, Sêmola, Sinapa, Solapa.

References: mainland Portugal▲ (Günther, 1861b, c; de Brito Capello, 1867a, d, 1880; Seabra, 1911; Nobre, 1935; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese, 1979e; Fischer *et al.*, 1981, 1987; Borges, 1984; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Henriques *et al.*, 2008; Carneiro *et al.*, 2014; Russell, 2014p; Carpenter and Iwatsuki, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (?) (Osório de Castro, 1967(?)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. caeruleostictus* is the Mediterranean Sea and, in the Eastern Atlantic, from Portugal to Angola. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira as doubtful, without indicating the geographic location. Since Osório de Castro (1967) no other author describes the presence of this species in Madeira. Thus, this species is not considered in the total number of valid species for Madeira.

Pagrus pagrus (Linnaeus 1758)

Common name: **Red porgy; Pargo-legítimo**, Capatão, Pargo, Pargo-comum, Pargo-de-direiro, Pargo-de-lombo-azul, Pargo-de-mancha-roxa, Pargo-de-riscas-azuis, Pargo-direito, Pargo-legítimo-azul, Pargo-verdadeiro, Pargo-vulgar, Pargueta, Parguete, Parguête.

References: mainland Portugal▲ (de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Gervais and Boulart, 1877b; Posser, 1880; Osório, 1888; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1894, 1895a, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982b; Martins, 1981; Borges, 1984; Bauchot and Hureau, 1986, 1990; Martins *et al.*, 1992, 2015; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Cabral, 1999; Borges *et al.*, 2002; Santos *et al.*, 2002a, c; Palma *et al.*, 2003; ICES, 2005; Sousa *et al.*, 2005; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Batista *et al.*, 2009; Silva *et al.*, 2009; Veiga *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Froese and Pauly, 2019), Azores▲ (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Sampaio, 1904; Nobre, 1924, 1930; Fowler, 1919, 1936; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Dias *et al.*, 1976, 1990; Wheeler, 1978; Tortonese, 1979e; Muzanor and Westhaus, 1981; Martins, 1982a; INIP, 1985a, c; Dias and Monteiro, 1986; Silva, 1986a,b, 1992; Patzner *et al.*, 1990, 1992; Wirtz, 1990a, 1994; Dias, 1991; Dias and Cascalho, 1991; Arruda, 1992, 1997; Azevedo *et al.*, 1992; Santos, 1992a; Silva and Krug, 1992; Santos *et al.*, 1994a,

1995a, 1997; Silva *et al.*, 1994, 1995; Azevedo, 1995; Clarke and Martins, 1995; Gonçalves *et al.*, 1995; Krug, 1995; Serafim and Krug, 1995; Silva and Menezes, 1995a, b; Isidro, 1996; Menezes, 1996; Morato-Gomes *et al.*, 1998; Morato *et al.*, 2001a; Cardigos *et al.*, 2005; Menezes *et al.*, 2006; Rosa *et al.*, 2006; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Pham *et al.*, 2013; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Afonso, 2016; Fauconnet *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837; Günther, 1859; de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; INIP, 1980, 1982c, 1984b, c; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Bauchot and Hureau, 1986, 1990; Lloris *et al.*, 1991; Andrade and Albuquerque, 1995; Delgado, 1998; Mello, 2000; Severino, 2004; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Ferreira *et al.*, 2008; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; POGID, 2016; POGIS, 2016; Biscoito *et al.*, 2017; Froese and Pauly, 2019).

***Sarpa salpa* (Linnaeus 1758)**

Common name: **Salema; Salema**, Boga, Choupa, Chupo, Saléma, Salêma, Salenca, Voga.

References: mainland Portugal[▲] (Carvalho, 1756; de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Posser, 1880; Baldaque da Silva, 1891; Girard, 1892; Nobre, 1895a, c, 1903, 1935; Osório, 1895, 1896; Vieira, 1897, 1898; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Martins, 1981; Borges, 1984; Bauchot and Hureau, 1986, 1990; Antunes *et al.*, 1988; Martins *et al.*, 1992, 1999, 2015; Almeida, 1996; Erzini *et al.*, 1997, 2001a; Cabral, 1999; Jorge *et al.*, 2002; Santos *et al.*, 2002a; Borges *et al.*, 2003; ICES, 2005; Ribeiro *et al.*, 2006, 2008, 2012; Fernandes *et al.*, 2007; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Chaves, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Ramos, 1869; Hilgendorf, 1888a; Faria e Silva, 1892; Collett, 1896a; Sampaio, 1904; Lampe, 1914; Roule, 1919a; Nobre, 1924, 1930; Ninni, 1933; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Wood, 1973, 1974; Wood and Williams, 1974; Wheeler, 1978; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Tortonese, 1979e; Saldanha, 1980; Martins, 1982a; Harmelin-Vivien *et al.*, 1985; Barreiros, 1990, 1995; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, 1995a, 1997; Wirtz, 1990a, 1991, 1994; Azevedo *et al.*, 1991, 1992; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Nash *et al.*, 1994a; Azevedo, 1995; Clarke and Martins, 1995; Gonçalves *et al.*, 1995; Santos and Nash, 1995; Isidro, 1996; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso, 2016) and Madeira[▲] (Bowdich, 1825; Lowe, 1837, 1849; Günther, 1859; de Velloso, 1892; Collett, 1896; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981; Bauchot and Hureau, 1986, 1990; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Carpenter and Iwatsuki, 2016; POGID, 2016; POGIS, 2016; Friedlander *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

***Sparus aurata* Linnaeus 1758**

Common name: **Gilthead seabream; Dourada**, Doirada, Doirada-fêmea, Doiradinha, Dourada-fêmea, Douradinha, Safata, Sapata.

References: mainland Portugal[▲] (Carvalho, 1765; Vandelli, 1771; de Brito Capello, 1867a, d, 1869b, 1880; Steindachner, 1867; Posser, 1880; Osório, 1888, 1895; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1898; Nobre, 1895a, 1903, 1935; Bragança, 1902; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese, 1979e; Bauchot *et al.*, 1981; Fischer *et al.*, 1981, 1987; Bauchot and Hureau, 1986, 1990; Bauchot, 1987; Antunes *et al.*, 1988; Arruda *et al.*, 1988; Hureau, 1991; Almeida, 1996; Erzini *et al.*, 1996, 2001a; Gonçalves *et al.*, 1996, 1997; Cabral, 1999; Martins *et al.*, 1999, 2015; Jorge *et al.*, 2002; Santos *et al.*, 2002a, c; Palma *et al.*, 2003; Garnerot *et al.*, 2004; ICES, 2005; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Ribeiro *et al.*, 2008, 2012; Batista *et al.*, 2009; Veiga *et al.*, 2009; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; POGID, 2016; POGIS, 2016; Biscoito *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

al., 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Carneiro *et al.*, 2014; Russell, 2014q; Golani and Fricke, 2018; Martins and Carneiro, 2018; Baptista *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Drouët, 1861b; Hilgendorf, 1888a; Fowler, 1936b; Collins, 1954; Osório de Castro, 1967; Tortonese, 1979e; Dooley *et al.*, 1985; Carneiro *et al.*, 2014) and Madeira (Osório de Castro, 1967(?); Alves and Alves, 2002(?); Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; Russell, 2014q; ICES, 2015b; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019(?)). According to Santos *et al.* (1997) the citation by Drouët (1861b), Hilgendorf (1888b), Fowler (1936b) and Collins (1954) *Chrysophrys aurata* is *Pagrus pagrus* (Linnaeus, 1758). Thus, this species is not considered in the total number of valid species for the Azores. Several specimens of *S. aurata* were observed during a underwater visual census which probably escaped from a recent open sea aquaculture unit in the area (ICES, 2015b). It is currently considered as an introduced species in Madeira.

***Spicara flexuosa* Rafinesque 1810**

Common name: No English common name; **Trombeiro-do-Mediterrâneo**, Trombeiro.

References: mainland Portugal (Nobre, 1895a; 1903, 1935; Fischer *et al.*, 1987; Sánchez 1999; Borges *et al.*, 2002, 2003; ICES, 2005; Ramos, 2010). This species is sometimes confused with *Spicara maena* (Linnaeus 1758) (Carpenter and Iwatsuki, 2016).

***Spicara maena* (Linnaeus 1758)**

Common name: **Blotched picarel; Trombeiro-choupa**, Trombeiro, Trombeiro-de-nariz-azul, Trombeta, Trombeteiro, Ferreiro.

References: mainland Portugal (de Brito Capello, 1867a, d, 1869a, b, 1880; Steindachner, 1867; Posser, 1880; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1935; Seabra, 1911; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1967; Tortonese *et al.*, 1979; Fischer *et al.*, 1981, 1987; Dooley *et al.*, 1985; Tortonese, 1986c; Heemstra, 1990; Erzini *et al.*, 2001a; Borges *et al.*, 2003; Prista *et al.*, 2003; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Chaves, 2008, 2009; Gonçalves *et al.*, 2010; Costa *et al.*, 2012; Carneiro *et al.*, 2014; Russell, 2014r; Carpenter and Iwatsuki, 2016; Saygili *et al.*, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (Fischer *et al.*, 1981; Dooley *et al.*, 1985; Russell, 2014r; Carpenter and Iwatsuki, 2016) and Madeira (Fischer *et al.*, 1981; INIP, 1984a, b, c; Russell, 2014r; Carpenter and Iwatsuki, 2016). This species is sometimes confused with *Spicara flexuosa* Rafinesque 1810 (Carpenter and Iwatsuki, 2016). The distribution map of this species reported by Russell (2014r) partially included the area of Selvagens Islands (Madeira EEZ).

***Spicara melanurus* (Valencienneas 1830)**

Common name: **Blackspot picarel; Trombeiro-malha-redonda**, Choupa, Dobrada, Taco-taco, Trombeiro.

References: Madeira (Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese *et al.*, 1979; Fischer *et al.*, 1981(?); Tortonese, 1986c; Heemstra, 1990; Lloris *et al.*, 1991; Wirtz *et al.*, 2008(?); Carneiro *et al.*, 2014; Carpenter *et al.*, 2014; Froese and Pauly, 2019(?)). Wirtz *et al.* (2008) consider the presence of this species very doubtful. However, according to Eschmeyer *et al.* (2019) the current geographic distribution of *S. melanurus* is the Eastern Atlantic. The northern extent range distribution of this species is the Canary Islands (MNHN-IC-1978-0683, 1978). Thus, specimens are needed for confirmation.

***Spicara smaris* (Linnaeus 1758)**

Common name: **Picarel; Trombeiro-boga**, Agulhão, Albatroz, Alcarraz, Alcatraz, Alcorraz, Baila, Caramelo, Ferreiro, Peixe-flauta, Trombeiro, Trombeiro-de-nariz-azul, Trombeta, Trombeteiro.

References: mainland Portugal (Vandelli, 1771; de Brito Capello, 1867a, d, 1880; Steindachner, 1867; Osório, 1888; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1897, 1898; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Tortonese *et al.*, 1979; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; Dooley *et al.*, 1985; Tortonese, 1986c; Heemstra, 1990; Martins *et al.*, 1992; Santos *et al.*, 2002a; Carneiro *et al.*, 2014; Carpenter and Iwatsuki, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Fischer *et al.*, 1981; Heemstra, 1990; Wirtz *et al.*, 2008(?); Carpenter and Iwatsuki, 2016; Froese and Pauly, 2019(?)). According to Wirtz *et al.* (2008) *S. smaris* was erroneously recorded from Madeira by Heemstra (1990). However, according to Eschmeyer *et al.* (2019) the current geographic distribution of *S. smaris* is the

Mediterranean Sea and the Black Sea, in the Eastern Atlantic: from Portugal to Canary Islands. Thus, specimens are needed for confirmation.

Spondylisoma cantharus (Linnaeus 1758)

Common name: **Black seabream; Choupa**, Ambua, Cântaro, Casaleira, Choupo, Doirada, Dourada, Loriana, Muchara, Mucharra, Oriana, Ruta, Sama, Saime, Salema, Sama, Viúva.
 References: mainland Portugal▲ ([Carvalho, 1765\(?\)](#); [de Brito Capello, 1867a, d, 1869b, 1880](#); [Steindachner, 1867](#); [Posser, 1880](#); [Osório, 1888, 1896](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Nobre, 1894, 1895a, c, 1903, 1935](#); [Vieira, 1893, 1897, 1898](#); [Bragança, 1902](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979e](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Bauchot and Hureau, 1986, 1990](#); [Bauchot, 1987](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Almeida, 1996](#); [Erzini et al., 1996, 1997, 2001a](#); [Gonçalves et al., 1996, 1997](#); [Cabral, 1999](#); [Pajuelo and Lorenzo 1999](#); [Gomes et al., 2001](#); [Borges et al., 2002, 2003](#); [Jorge et al., 2002](#); [Santos et al., 2002a, c](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Ribeiro et al., 2006, 2008, 2012](#); [Stergiou et al., 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Garrido et al., 2009](#); [Silva et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Russell, 2014s](#); [Silva and Borges, 2014](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Fowler, 1936b](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Wheeler, 1978](#); [Tortonese, 1979e](#); [Martins, 1982a](#); [Arruda, 1992, 1997](#); [Santos et al., 1997\(?\)](#); [Pajuelo and Lorenzo, 1999](#); [Morato et al., 2001a](#); [Carneiro et al., 2014](#); [Russell, 2014s](#)) and Madeira▲ ([Lowe, 1837, 1839, 1840b, 1844a, 1849](#); [Gervais and Boulart, 1877b](#); [de Velloso, 1892](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979e](#); [Fischer et al., 1981](#); [Augier, 1985](#); [Bauchot and Hureau, 1986, 1990](#); [Lloris et al., 1991](#); [Delgado, 1998](#); [Pajuelo and Lorenzo, 1999](#); [Mello, 2000](#); [Araújo et al., 2005](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Russell, 2014s](#); [Carpenter and Iwatsuki, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). According to Eschmeyer et al., (2019) and Froese and Pauly (2019) the current geographic distribution of *S. cantharus* in the Eastern Atlantic is from Norway south to the northern Namibia, including Madeira, Canary and Cape Verde islands. Santos et al. (1997) did not find any evidence of the presence of this species in the Azores. However, Morato et al. (2001a) reported catches of *S. cantharus* in the Azores for the period 1982 - 1999. Thus, this subject deserves further detailed attention and this species is not considered in the total number of valid species for the Azores.

Family Sciaenidae

Argyrosomus regius (Asso 1801)

Common name: **Meagre; Corvina-legítima**, Borregata, Corvina, Corvina-branca, Corvina-real, Corvinata, Corvina-vulgar, Rabeta, Raveta.

References: mainland Portugal▲ ([de Brito Capello, 1867a, d, 1869b, 1880](#); [Steindachner, 1867](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1893, 1897, 1898](#); [Nobre, 1894, 1895a, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Trewavas, 1979a](#); [Chao, 1981](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [Dooley et al., 1985](#); [Chao, 1986, 2014](#); [Chao and Trewavas, 1990](#); [Erzini et al., 2001a](#); [Santos et al., 2002a, c](#); [Gomes and Olim, 2007](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Batista et al., 2009](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Martins et al., 2015](#); [Golani and Fricke, 2018](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Sampaio, 1904](#); [Ribeiro, 1936](#); [Collins, 1954](#); [Osório de Castro, 1967](#); [Arruda, 1997](#); [Martins, 1982a](#); [Carneiro et al., 2014](#)) and Madeira ([Osório de Castro, 1967](#); [Fischer et al., 1981](#); [Chao, 2014a](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *A. regius* is the Western Baltic Sea, North Sea, Mediterranean Sea, Black Sea and, in the Eastern Atlantic, from Norway south to Congo; Red Sea (Mediterranean Sea immigrant). According to Santos et al. (1997) this species, or any other member of

the family Sciaenidae, does not occur in the region. Thus, this species is not considered in the total number of valid species for the Azores. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of Madeira without indicating the geographic location. The distribution maps of this species reported by Fischer *et al.* (1981) and Chao (2014a) partially included the area of Selvagens Islands (Madeira EEZ).

Cynoscion regalis (Bloch & Schneider 1801)

Common name: **Squeteague; Corvinata-real**, Corvina, Fraco, Peixe-fraco.

References: mainland Portugal[▲] ([Martins et al., 2015](#); [Béarez et al., 2016](#); [Morais and Teodósio, 2016](#); [Bañón et al., 2017c](#); [Gomes et al., 2017](#); [ICES, 2017b](#); [Morais et al., 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#)). This is an invasive species.

Pseudotolithus typus Bleeker 1863

Common name: **Longneck croaker; Rainha-branca**, Corvina, Corvina-branca, Corvina-rainha, Rainha.

References: Madeira ([Chao and Trewavas, 1990](#); [Froese and Pauly, 2019](#)). Often confused with *Pseudotolithus senegalensis* (Valenciennes 1833)

Sciaena umbra Linnaeus 1758

Common name: **Brown meagre; Roncadeira-preta**, Borregata, Calafate-preto, Corvina, Corvina-preta, Corvinata, Rabeta, Rainha, Raveta, Roncadeira, Roncador.

References: mainland Portugal ([Vandelli, 1771](#); [Vieira, 1897, 1898](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Trewavas, 1979a](#); [Chao, 1981, 1986, 2014](#); [Dooley et al., 1985](#); [Fischer et al., 1987](#); [Chao and Trewavas, 1990](#); [Erzini et al., 2001a](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Fischer et al., 1981](#); [Chao, 1986, 2014b](#); [Carneiro et al., 2014](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. umbra* is the Mediterranean Sea, Black Sea and, in the Eastern Atlantic, from the English Channel to Senegal and possibly Gabon, including Canary and Cape Verde islands. Fischer *et al.* (1981) and Chao (2014b) reported the Selvagens Islands area in the distribution maps of this species.

Umbrina canariensis Valenciennes 1843

Common name: **Canary drum; Calafate-das-Canárias**, Calafate, Corvina, Corvina-marreca, Corvinata, Formosa, Rainha, Roncadeira, Roncador.

References: mainland Portugal ([Chao, 1981, 1986, 2016](#); [Fischer et al., 1987](#); [Martins et al., 1992](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Osório de Castro, 1967](#)) and Madeira ([MNHN-IC-1912-0208, 1922](#); [MNHN-IC-1912-0212, 1922](#); [Osório de Castro, 1967](#); [Chao, 1981, 2014c, 2016](#); [Fischer et al., 1981](#); [Carneiro et al., 2014](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *U. canariensis* is the South-western Mediterranean Sea and, in the Eastern Atlantic, from Bay of Biscay to South Africa. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location and according to Santos *et al.* (1997) this species, or any other member of the family Sciaenidae, does not occur in the region. Thus, this species is not considered in the total number of valid species for the Azores. Fischer *et al.* (1981) and Chao (1981, 2014c, 2016) reported the Selvagens Islands area in the distribution maps of this species.

Umbrina cirrosa (Linnaeus 1758)

Common name: **Shi drum; Calafate-das-riscas**, Calafate, Corvina, Corvina-marreca, Corvinata, Labarda, Rainha, Roncadeira, Roncador, Viúva.

References: mainland Portugal[▲] ([de Brito Capello, 1880](#); [Seabra, 1911](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Trewavas, 1979a](#); [Chao, 1981, 1986, 2016](#); [Fischer et al., 1981, 1987](#); [Dooley et al., 1985](#); [Chao and Trewavas, 1990](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores (?) ([Osório de Castro, 1967](#)) and Madeira ([Osório de Castro, 1967](#); [Fischer et al., 1981](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *U. cirrosa* is the Mediterranean Sea, Sea of Marmara, Black Sea and, in the Eastern Atlantic, from Bay of Biscay to Gulf of Guinea. Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores and Madeira without indicating the geographic location. According to Santos *et al.* (1997) this species, or any other member of the family Sciaenidae, do not occur in the region. Fischer et

al. (1981) reported the Selvagens Islands area in the distribution map of this species. Thus, this species is not considered in the total number of valid species for the Azores.

***Umbrina ronchus* Valenciennes 1843**

Common name: **Fusca drum; Calafate-fosco**, Corvina, Roncador.

References: mainland Portugal ([Martins, 1979](#); [Chao, 1986, 2014d](#); [Fischer et al., 1987](#); [Chao and Trewavas, 1990](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#)), Azores (?) ([Osório de Castro, 1967](#)) and Madeira ([Günther, 1860](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Fischer et al., 1981](#); [Carneiro et al., 2014](#); [Chao, 2016](#)). Osório de Castro (1967) mentions the presence of this species in the fishing grounds of the Azores without indicating the geographic location. According to Santos *et al.* (1997) this species, or any other member of the family Sciaenidae, do not occur in the region. Thus, this species is not considered in the total number of valid species for the Azores. Fischer *et al.* (1981) and Chao (2016) reported the Selvagens Islands area in the distribution maps of this species.

Family Mullidae

***Mullus barbatus* Linnaeus 1758**

Common name: **Red mullet; Salmonete-da-vasa**, Salmarino, Salmonejo, Salmonete, Salmonête, Salmonete-vermelho.

References: mainland Portugal▲ ([Vandelli, 1771](#); [Steindachner, 1867](#); [de Brito Capello, 1867d, 1869b, 1880](#); [Osório, 1896](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Hureau, 1979b](#); [Ben-Tuvia, 1981, 1990a](#); [Fischer et al., 1981, 1987](#); [Martins, 1981](#); [INIP, 1982b](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Hureau, 1986d](#); [Erzini et al., 2001a](#); [Borges et al., 2002](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa, 2005](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Batista et al., 2009](#); [Silva, 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Carneiro et al., 2014](#); [Craig and Pollard, 2015c](#); [Golani, 2016](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Drouët, 1861b](#); [Ramos, 1869](#); [Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Roule, 1919a](#); [Nobre, 1924, 1930](#); [Fowler, 1936b](#); [Ribeiro, 1936](#); [Cadenat, 1937](#); [Collins, 1954](#), [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Hureau, 1979b, 1986](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Carneiro et al., 2014](#); [Craig and Pollard, 2015c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Günther, 1859](#); [Vaillant, 1919](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Ben-Tuvia, 1981](#); [Fischer et al., 1981](#); [Carneiro et al., 2014](#); [Craig and Pollard, 2015c](#)).

***Mullus surmuletus* Linnaeus 1758**

Common name: **Surmullet; Salmonete-legítimo**, Salmarino, Salmonejo, Salmonéta, Salmonete, Salmonete-da-costa, Salmonete-da-pedra, Salmonete-do-rolo, Salmonete-vermelho.

References: mainland Portugal▲ ([Carvalho, 1756\(?\)](#); [Vandelli, 1771](#); [de Brito Capello, 1867d, 1869b, 1880](#); [Posser, 1880](#); [Baldaque da Silva, 1891](#); [Almeida and Roquette, 1892](#); [Girard, 1892](#); [Vieira, 1893, 1897, 1898](#); [Nobre, 1894, 1895a, c, 1903, 1935](#); [Osório, 1895](#); [Bragança, 1902](#); [Seabra, 1911](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Hureau, 1979b](#); [Ben-Tuvia, 1981, 1990a](#); [Fernández et al., 1981](#); [Fischer et al., 1981, 1987](#); [INIP, 1981a, 1982a, b](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis, 1985](#); [Hureau, 1986d](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Martins et al., 1992, 1999, 2015](#); [Almeida, 1996](#); [Gonçalves et al., 1996, 1997](#); [Erzini et al., 1997, 2001a](#); [Cabral, 1999](#); [Borges et al., 2002](#); [Jorge et al., 2002](#); [Santos et al., 2002a](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [Moura, 2004](#); [ICES, 2005](#); [Sousa et al., 2005](#); [Ribeiro et al., 2006, 2008, 2012](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Chaves, 2008, 2009](#); [Silva, 2008, 2010, 2011](#); [Batista et al., 2009](#); [Silva et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Diogo and Pereira, 2013](#); [Carneiro et al., 2014](#); [Silva and Borges, 2014](#); [Golani, 2016](#); [Ryland and Lancaster, 2017](#); [Martins and Carneiro, 2018](#); [Baptista et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores▲ ([Drouët, 1861b](#); [Ramos, 1869, 1871](#); [Guimarães, 1882a](#); [Hilgendorf, 1888a](#); [Sampaio, 1904](#); [Lampe, 1914](#); [Roule, 1919a](#); [Vaillant, 1919](#); [Nobre, 1924, 1930](#); [Ribeiro, 1936](#); [Cadenat, 1937](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Wood, 1973](#); [Wood and Williams, 1974](#); [Wheeler, 1978](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Saldanha, 1980](#); [Martins, 1982a](#); [Chícharo and Teodósio, 1990](#); [Barreiros, 1990, 1995](#); [Nash et al., 1990, 1994a](#); [Patzner, 1990](#); [Patzner et al., 1990, 1992](#); [Santos](#)

et al., 1990b, 1994a, b, 1995a, 1997; Wirtz, 1990a, 1991, 1994; Hureau, 1991; Arruda, 1992, 1997; Azevedo *et al.*, 1992; Santos, 1992a; Azevedo, 1995; Gonçalves *et al.*, 1995; Saldanha, 1995; Santos and Nash, 1995; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a, b; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Afonso, 2016; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Bowdich, 1825; Lowe, 1837; Günther, 1859; de Velloso, 1892(?); Fowler, 1936b; Noronha and Sarmento, 1948; Maul, 1949d; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Hureau, 1979b; Wirtz, 1994; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Golani, 2016; POGID, 2016; POGIS, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Family Chaetodontidae

Chaetodon ocellatus Bloch 1787

Common name: Spotfin butterflyfish; **Borboleta-de-risca-amarela**, Lebre, Borboleta, Peixe-borboleta.

References: Azores (?) (PECS) (ROM 23917, 1966; Carneiro *et al.*, 2014). The geographic coordinates of the occurrence records of the ROM Fish Collection (ROM 23917, 1966) in the PECS Azores were rectified to off the Florida coast, on March 4, 2018. Thus, this species is not considered in the total number of valid species for the Azores.

Chaetodon sedentarius Poey 1860

Common name: Reef butterflyfish; **Borboleta-do-recife**, Lebre, Borboleta, Peixe-borboleta.

References: Azores (Santos *et al.*, 1997; Morato *et al.*, 2004; Cardigos *et al.*, 2006; Floeter *et al.*, 2008; Porteiro *et al.*, 2010; Afonso *et al.*, 2013; Carneiro *et al.*, 2014) and Madeira (Floeter *et al.*, 2008). According to Allen *et al.* (2010), Eschmeyer *et al.* (2019) and Froese and Pauly (2019) the current geographic distribution of *C. sedentarius* is the Western Atlantic. However, the occurrence of this species has already been confirmed for the Azores. Specimens are needed for confirmation the presence of this species in Madeira.

Family Pomacanthidae

Pomacanthus paru (Bloch 1787)

Common name: **French angelfish; Lebre-preta**, Lebre.

References: Madeira (?) (Bowdich, 1825; Fowler, 1936b, Albuquerque, 1954-1956; Osório de Castro, 1967). Bowdich (1825) reported the occurrence of *Chaetodon leachii* and Maugé (1990) considered this species as possible synonym of *Pomacanthus paru* (Bloch 1787) but its current status is uncertain as *P. paru* by Eschmeyer *et al.* (2019). Thus, this species is not considered in the total number of valid species.

Family Cepolidae

Cepola macrophthalma (Linnaeus 1758)

Common name: **Red bandfish; Suspensório**, Peixe-cor-de-rosa.

References: mainland Portugal[▲] (de Brito Capello, 1868, 1880; Steindachner, 1868b; Osório, 1888, 1917; Girard, 1892; Vieira, 1897, 1898; Bragança, 1902, 1903; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Monod, 1979e; Fernández *et al.*, 1981; Fischer *et al.*, 1981, 1987; INIP, 1981a, 1982b; Martins, 1981; Dinis, 1985; Tortonese, 1986b; Quero, 1990c; Borges *et al.*, 2002; ICES, 2005; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques *et al.*, 2008; Silva, 2008, 2010, 2011; Silva *et al.*, 2009; Costa *et al.*, 2012; Chaves and Silva, 2013; Carneiro *et al.*, 2014; Costa, 2014; Silva and Borges, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Family Pomacentridae

Abudefduf saxatilis (Linnaeus 1758)

Common name: **Sergeant-major; Castanheta-raiada**, Castanheta.

References: Madeira (Freitas and Araújo, 2006; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Edwards, 2016; Froese and Pauly, 2019).

Chromis chromis (Linnaeus 1758)

Common name: **Damselfish; Castanheta-báia**, Castanheta, Castanheira, Castanheira-báia, Castanheta-branca, Romeiro, Sargo.

References: mainland Portugal ([Vandelli, 1771](#); [Osório, 1917](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Saldanha, 1966c](#); [Wood, 1977](#); [Monod, 1979f](#); [Quignar and Pras, 1986](#); [Fischer et al., 1987](#); [Lloris and Rocabado, 1990](#); [Domingues et al., 2006](#); [Henriques et al., 2007, 2008](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015l](#); [Edwards, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] (?) ([Hilgendorf, 1888a, b](#); [Sampaio, 1904](#); [Nobre, 1924, 1930](#); [Fowler, 1919, 1936b](#); [Cadenat, 1937](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bauchot and Blanc, 1961](#); [Saldanha, 1966c, 1980](#); [Osório de Castro, 1967](#); [Wood, 1973, 1977](#); [Mapstone and Wood, 1974, 1975](#); [Wood and Williams, 1974, 1975](#); [Williams and Wood, 1974](#); [Arruda, 1977](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Arruda, 1980a, 1992](#); [Martins, 1982a](#); [Ré and Gomes, 1982](#); [Lloris and Rocabado, 1990](#); [Lloris et al., 1991](#); [Herrera and Smith-Vaniz, 2015l](#); [Froese and Pauly, 2019](#)) and Madeira[▲] (?) ([MNHN-IC-0000-8159, 1833](#); [Cuvier and Valenciennes, 1833](#); [Günther, 1862](#); [Vinciguerra, 1883](#); [de Velloso, 1892](#); [Osório, 1917](#); [Fowler, 1919, 1923, 1936b](#); [ZMH 16971, 1930](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [AMNH 16870, 1946](#); [Maul, 1949](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Saldanha, 1966c](#); [Quignard and Pras, 1986b](#); [SAIAB FC 22576, 1990](#); [SAIAB FC 34750, 1990](#); [Lloris and Rocabado, 1990](#); [Lloris et al., 1991](#); [Andrade and Albuquerque, 1995](#); [Carneiro et al., 2014](#); [Herrera and Smith-Vaniz, 2015l](#); [Froese and Pauly, 2019](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *C. chromis* is the Mediterranean Sea, Black Sea, Eastern Atlantic: Spain and Portugal to Gulf of Guinea and possibly Angola, including offshore islands. Santos et al. (1997) mentioned that the records of this species to the Azores are synonyms, misspellings and misidentifications of *Chromis limbata* (Valenciennes 1833). Similarly, Wirtz et al. (2008) referred that all the citations to this species to Madeira are erroneous and should be considered as misidentifications of *C. limbata* based on the study by Domingues et al. (2005). However, there are specimens identified as *C. chromis* caught off in Madeira in several museum collections (perhaps not yet re-evaluated) and bibliographic references (years 2015 and 2018) to the occurrence of *C. chromis* in Madeira and the Azores. Thus, given these contradictory opinions, it was decided do not consider in the total number of valid species of Madeira and the Azores.

Chromis limbata (Valenciennes 1833)

Common name: **Azores chromis; Castanheta-da-Madeira**, Castanheta, Castanheta-amarela, Castanheta-báia, Castanheta-branca, Sarabanete.

References: mainland Portugal ([Abecasis et al., 2009](#); [OCEANA, 2014](#)), Azores ([Hilgendorf, 1888a, b](#); [Sampaio, 1904](#); [Nobre, 1924, 1930](#); [Fowler, 1919, 1936](#); [Cadenat, 1937](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bauchot and Blanc, 1961](#); [Saldanha, 1966c, 1980](#); [Wood, 1973, 1977](#); [Mapstone and Wood, 1974, 1975](#); [Wood and Williams, 1974](#); [Willis and Wood, 1974](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Arruda, 1977, 1980a](#); [Martins, 1982a](#); [Ré and Gomes, 1982](#); [Edwards, 1986](#); [Patzner and Santos, 1990b, d, 1993](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1997](#); [Wirtz, 1990a, 1994](#); [Allen, 1991](#); [Azevedo et al., 1991, 1992](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001b](#); [Cardigos et al., 2005](#); [Domingues et al., 2005, 2006](#); [Floeter et al., 2008](#); [Leite et al., 2009](#); [Bertонcini et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Diogo and Pereira, 2013, 2014](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Edwards, 2016](#); [Froese and Pauly, 2019](#)) and Madeira[▲] ([MNHN-IC-0000-1377, 1833](#); [Cuvier and Valenciennes, 1833](#); [Lowe, 1837](#); [Wood, 1977](#); [Emery, 1981](#); [Fischer et al., 1981](#); [Augier, 1985](#); [Lloris et al., 1991](#); [Delgado, 1998](#); [Falcón et al., 2000](#); [Mello, 2000](#); [Abreu et al., 2005](#); [Araújo et al., 2005](#); [Domingues et al., 2005, 2006](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [Leite et al., 2009](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Edwards, 2016](#); [POGID, 2016](#); [Friedlander et al., 2017](#); [Froese and Pauly, 2019](#)). The holotype of *Heliases limbatus* Valenciennes [A.] in Cuvier & Valenciennes 1833, synonym of *C. limbata*, caught off in Madeira is in the MNHM collections ([MNHN-IC-0000-1377](#)). According to Domingues et al. (2005) this species is restricted to the Macaronesian islands (Azores, Madeira, and Canaries) and the western coast of North Africa.

Similiparma lurida (Cuvier 1830)

Common name: **Canary damsel; Castanheta-preta**, Castanheta, Castanheta-ferreira.

References: mainland Portugal (Allen, 1991; Gonçalves et al., 2004; Abecasis et al., 2009; Carneiro et al., 2014), Azores[▲] (Hilgendorf, 1888a, b; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Wood, 1973; Mapstone and Wood, 1974, 1975; Wood and Williams, 1974; Willis and Wood, 1974; Harmelin and Harmelin-Vivien, 1979; Hureau and Monod, 1979; Ré, 1979a, b, 1980, 1990; Saldanha, 1980; Edwards, 1986; Quignar and Pras, 1986; Lloris and Rocabado, 1990; Patzner, 1990; Patzner and Santos, 1990b, d, 1993; Patzner et al., 1990, 1992; Santos et al., 1990b, 1997; Wirtz, 1990a, 1994; Allen 1991; Azevedo et al., 1991, 1992; Lloris et al., 1991; Arruda, 1992, 1997; Arruda et al., 1992; Santos, 1992a; Santos and Barreiros, 1993; Azevedo, 1995; Barreiros, 1995; Gonçalves et al., 1995; Harmelin-Vivien et al., 2001; Morato et al., 2001b; Cardigos et al., 2005; Floeter et al., 2008; Bertoncini et al., 2010; Botelho et al., 2010; Porteiro et al., 2010; Carneiro et al., 2014; Diogo and Pereira, 2014; OCEANA, 2014; Herrera and Smith-Vaniz, 2015m; Afonso, 2016; Edwards, 2016; Almeida and Biscoito, 2019; Froese and Pauly, 2019) and Madeira[▲] (MNHN-IC-0000-5286, 1830; Cuvier and Valenciennes, 1830, 1835; Lowe, 1837; Günther, 1862; Vinciguerra, 1883; de Velloso, 1892; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954; Hureau and Monod, 1979; Quignar and Pras, 1986; Augier, 1985; Lloris and Rocabado, 1990; Lloris et al., 1991; Abreu et al., 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón et al., 2000; Mello, 2000; Araújo et al., 2005; Domingues et al., 2005; Ribeiro et al., 2005b; Floeter et al., 2008; Ribeiro, 2008; Wirtz et al., 2008; EMEPC, 2010; Carneiro et al., 2014; Almada et al., 2015; Herrera and Smith-Vaniz, 2015m; Edwards, 2016; POGID, 2016; POGIS, 2016; Friedlander et al., 2017; Froese and Pauly, 2019). The holotype (unique) of *Glyphisodon luridus* Cuvier in Cuvier & Valenciennes 1830, synonym of *S. lurida*, caught off in Madeira is in the MNHM collections (MNHN-IC-0000-5286).

Family Labridae

***Acantholabrus palloni* (Risso 1810)**

Common name: **Scale-rayed wrasse; Bodião-do-alto**, Bodião, Truta, Truta-do-alto, Vidrão.

References: mainland Portugal (Steindachner, 1868b; Osório, 1895, 1917; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Bauchot and Quignard, 1979; INIP, 1981a; Martins, 1981; Dinis and Marecos, 1984; Quignard and Pras, 1986b; Fischer et al., 1987; Gomon and Forsyth, 1990; Martins et al., 1992; Erzini et al., 2001a; Almada et al., 2002; Mendes et al., 2004; ICES, 2005; Gomes and Olim, 2007; Chaves, 2008; Henriques et al., 2008; Costa et al., 2012; Carneiro et al., 2014; OCEANA, 2014; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Collett, 1896; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Quignard, 1966; Osório de Castro, 1967; Maul, 1976; Wheeler, 1978; Bauchot and Quignard, 1979; Dooley et al., 1985; Shcherbachov et al., 1985a; Quignard and Pras, 1986b; Michel et al., 1987; Gomon and Forsyth, 1990; Lloris et al., 1991; Arruda, 1992, 1997; Porteiro et al., 1996, 2010; Santos et al., 1997; Kukuev, 2002 (PECS); Menezes et al., 2006; Floeter et al., 2008; Carneiro et al., 2014; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837, 1839, 1840b, 1843, 1844a; 1849; NHMUK 1857.6.13.103, 1854-1861; Günther, 1862; Steindachner, 1868b; Guimarães, 1884; Collett, 1896; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Bauchot and Quignard, 1979; Dooley et al., 1985; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris et al., 1991; Floeter et al., 2008; Carneiro et al., 2014; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). The holotype of *Acantholabrus imbricatus* Lowe 1839, synonym of *A. palloni*, caught off in Madeira is in the NHMUK collections (NHMUK 1857.6.13.103). One specimen caught off at the coast of mainland Portugal is in the MMF collections (MMF 22798, 1967).

***Bodianus scrofa* (Valenciennes 1839)**

Common name: **Barred hogfish; Bodião-dente-de-cão**, Bedja, Bodião, Bodinho, Cachorro, Canzala, Cão, Cão-do-mar, Cão-marinho, Gaio, Peixe-cachorro, Peixe-cão, Velha, Viola, Xaroco.

References: Azores[▲] (Guimarães, 1884; Hilgendorf, 1888a; Collins, 1954; Albuquerque, 1954-1956; Wood and Williams, 1974; Harmelin and Harmelin-Vivien, 1979; Dias et al., 1976, 1990; Ré, 1979a; Saldanha, 1980; Martins, 1982a; Dooley et al., 1985; INIP, 1985a; Gomon and Forsyth, 1990; Patzner et al., 1990, 1992; Wirtz, 1990a, 1994; Azevedo et al., 1991, 1992; Dias, 1991; Arruda, 1992, 1997; Arruda et al., 1992; Santos, 1992a; Gonçalves, 1993; Azevedo, 1995; Silva et al., 1995; Porteiro et al., 1996,

2010; Santos *et al.*, 1997; Harmelin-Vivien *et al.*, 2001; Almada *et al.*, 2002; Cardigos *et al.*, 2005; Gomon, 2006; Menezes *et al.*, 2006; Floeter *et al.*, 2008; Botelho *et al.*, 2010; Carneiro *et al.*, 2014; Diogo and Pereira, 2014; Afonso, 2016; Westneat, 2016; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837, 1839, 1840b, 1844a, 1849; Günther, 1862; de Velloso, 1892; Nunes, 1953, 1974; Osório de Castro, 1967; Bauchot and Quignard, 1979; Saldanha, 1979; INIP, 1980; Fischer *et al.*, 1981; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Hanel *et al.*, 2003; Gomon, 2006; Ferreira *et al.*, 2008; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Westneat, 2016; Friedlander *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The whereabouts of the holotype of *Crenilabrus caninus* Lowe 1839, synonym of *B. scrofa*, caught off in Madeira are unknown.

***Bodianus speciosus* (Bowdich 1825)**

Common name: **Blackbar hogfish; Bodião-cão**, Bedja, Bodião, Bodinho, Cachorro, Canzala, Cão, Cão-do-mar, Cão-marinho, Gaio, Peixa-côa, Peixe-cachorro, Peixe-cão, Peixe-côa, Peixe-quarenta-e-cinco, Veja, Velha, Viola, Xarroco.

References: mainland Portugal (?) (Osório de Castro, 1954(?)), Azores (?) (Fowler, 1936b; Albuquerque, 1954-1956; Collins, 1954; Osório de Castro, 1967; Wood and Williams, 1974; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a; Saldanha, 1980; Martins, 1981; INIP, 1985a; Dias, 1991; Arruda, 1997) and Madeira (?) (Bowdich, 1825(?); Lowe, 1837, 1839, 1849; Cuvier and Valenciennes, 1839; Günther, 1862; Guimarães, 1884; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris *et al.*, 1991; Russell and Pollard, 2010; Carneiro *et al.*, 2014; Westneat, 2016). According to Eschmeyer *et al.* (2019) the current geographic distribution of *B. speciosus* in the Eastern Atlantic is from Mauritania to Angola, including Cape Verde Islands. Osório de Castro (1954) mentioned this species as *Bodianus iagonensis* (Bowdich 1825) and considered its presence in mainland Portugal doubtful. Santos *et al.* (1997) mentioned that the presence of *B. speciosus* in the Azores has never been confirmed, so all the citations to this species or its common synonyms (e.g. *Diastodon speciosus* Bowdich 1825) are considered misidentifications of *Bodianus scrofa* (Valenciennes 1839). According to Wirtz *et al.* (2008) this species has never been found at Madeira and Eschmeyer *et al.* (2019) mention that Madeira records are erroneous. Thus, this species is not considered in the total number of valid species for mainland Portugal, the Azores and Madeira islands.

***Centrolabrus exoletus* (Linnaeus 1758)**

Common name: **Rock cook; Bodião-de-boca-pequena**.

References: mainland Portugal[▲] (Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Martins *et al.*, 1992; Almeida, 1996; Cabral, 1999; Almada *et al.*, 2002; Henriques *et al.*, 2007, 2008; Batista *et al.*, 2009; Carneiro *et al.*, 2014; Craig *et al.*, 2015a; Froese and Pauly, 2019).

***Coris julis* (Linnaeus 1758)**

Common name: **Rainbow wrasse; Judia**, Canário, Canário-do-mar, Caralete, Caralete-de-el-rei, Caraludo, Dentilha, Elefante, Freirinha, Judia, Júlia, Juliana, Lambaz, Mordedor, Mordedor, Palouco-amarelo, Peise-de-rolo, Peixe-do-rei, Peixe-piça, Peixe-pissa, Peixe-rei, Peixe-rei-amarelo, Peixe-rei-do-alto, Peixe-rosa, Pica-de-el-rei, Piça-del-rei, Pichinha-del-rei; Pico, Pissa, Pissa-de-el-rei, Pissinha-de-el-rei, Rei-amarelo, Rosa, Salmonete, Samonete, Taralhete, Toucinho, Verdelho.

References: mainland Portugal (de Brito Capello, 1868, 1869b, 1880; Posser, 1880; Osório, 1888, 1895; Baldaque da Silva, 1891; Almeida and Roquette, 1892; Girard, 1892; Vieira, 1893, 1897, 1900; Nobre, 1894, 1895a, c, 1903, 1935; Bragança, 1902, 1903; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Fischer *et al.*, 1981, 1987; Quignard and Pras, 1986; Bauchot, 1987; Gomon and Forsyth, 1990; Martins *et al.*, 1992; Almeida, 1996; Erzini *et al.*, 1996; Gonçalves *et al.*, 1996, 1997; Almada *et al.*, 2002; Gonçalves *et al.*, 2004; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Abecasis *et al.*, 2009; Batista *et al.*, 2009; Gonçalves *et al.*, 2010; Pollard and Afonso, 2010; Carneiro *et al.*, 2014; Christiansen *et al.*, 2014; OCEANA, 2014; Afonso *et al.*, 2015; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Drouët, 1861b; Günther, 1862; Ramos, 1869; Hilgendorf, 1888a; USNM 42127, 1889; Collett,

1896a; Regan, 1903a; Sampaio, 1904; Fowler, 1919, 1936b; Vaillant, 1919; Roule, 1919a; Nobre, 1924, 1930; Zugmayer, 1933; Ribeiro, 1936; Le Gall, 1937b; Collins, 1954; Albuquerque, 1954-1956; Quignard, 1966; Osório de Castro, 1967; Wood, 1973; Mapstone and Wood, 1974, 1975; Wood and Williams, 1974; Wheeler, 1978; Bauchot and Quignard, 1979; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Saldanha, 1980, 1995; Fischer *et al.*, 1981; Martins, 1982a; Quignard and Pras, 1986b; Bauchot, 1987; Laurentand Lejeune, 1988; Barreiros, 1990, 1995; Gomon and Forsyth, 1990; Patzner, 1990; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, 1995a, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a, 1995b; Azevedo, 1995; Gonçalves *et al.*, 1995; Santos *et al.*, 1995a; Isidro, 1996; Porteiro *et al.*, 1996, 2010; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a, b; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso *et al.*, 2015; Afonso, 2016; Westneat, 2016; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1836; Lowe, 1837, 1839, 1840b, 1844a, 1849; Günther, 1862; Gervais and Boulart, 1877b; Collett, 1896; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Fischer *et al.*, 1981; Augier, 1985; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Pollard and Afonso, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; Afonso *et al.*, 2015; POGID, 2016; POGIS, 2016; Westneat, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The holotype (unique) of *Julis azorensis* Fowler 1919 caught off in Horta, Faial Island (Azores), is in the USNM collections (USNM 42127) and the whereabouts of holotype of *Julis melanura* Lowe 1839 caught off in Madeira are unknown and both are synonym of *C. julis*. Five specimens caught off at the coast of Setúbal (Portugal) in 1908 are in the MB collections (MB06-004715, set of three specimens and MB06-004725, set of two specimens).

Ctenolabrus rupestris (Linnaeus 1758)

Common name: **Goldsinny-wrasse**; **Bodião-rupreste**, Bodião, Bodiano, Peixe-rei, Rei, Verdelhão, Verdelho.

References: mainland Portugal (Osório, 1896; Nobre, 1903, 1935; Seabra, 1911; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Dinis, 1985; Quignard and Pras, 1986b; Bauchot, 1987; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Martins *et al.*, 1992; Almeida, 1996; Cabral, 1999; Almada *et al.*, 2002; ICES, 2005; Ribeiro *et al.*, 2006; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Silva, 2008; Garrido *et al.*, 2009; Gonçalves *et al.*, 2010; Carneiro *et al.*, 2014; Pollard, 2015b; Ryland and Lancaster, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019).

Labrus bergylta Ascanius 1767

Common name: **Ballan wrasse**; **Bodião-reticulado**, Bodião, Bodiano, Bodião, Bodião-azul, Bodião-da-costa, Bodião-pardo, Bodião-Santo-António, Bodião-vermelho, Burrinho, Burrinho-manso, Burrinho-moiro, Burrinho-mouro, Chalrão, Chita, Gidilhão, Godião, Godião-azul, Lucinha, Lucinheira, Marabota, Maraconta, Maracoto, Maragota, Maragote, Margota, Maria-Júlia, Marmota, Peixe-rei-do-alto, Pilrão, Truta-da-costa, Truta-vermelha, Verdelho.

References: mainland Portugal[▲] (de Brito Capello, 1868, 1873(?); Steindachner, 1868b; Posser, 1880; Baldaque da Silva, 1891; Girard, 1892; Nobre, 1894, 1895a, 1903, 1935; Osório, 1895, 1896; Vieira, 1898; Richard, 1908; Fowler, 1936b; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; INIP, 1982b; Quignard and Pras, 1986b; Bauchot, 1987; Fischer *et al.*, 1987; Gomon and Forsyth, 1990; Martins *et al.*, 1992, 1999; Almeida, 1996; Erzini *et al.*, 1996, 2001a; Almada *et al.*, 2002; Garnerot *et al.*, 2004; Gonçalves *et al.*, 2004; ICES, 2005; Ribeiro *et al.*, 2006; Henriques *et al.*, 2007, 2008; Abecasis *et al.*, 2009; Garrido *et al.*, 2009; Carneiro *et al.*, 2014; OCEANA, 2014; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Vaillant, 1919; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Quignard, 1966; Osório de Castro, 1967; Wood and Williams, 1974; Wood, 1974; Bauchot and Quignard, 1979; Harmelin and Harmelin-Vivien, 1979; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Bauchot, 1987; Barreiros, 1990, 1995; Gomon and Forsyth, 1990; Patzner *et al.*, 1990, 1992; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Lloris *et al.*, 1991; Arruda *et al.*, 1992; Santos, 1992a; Gonçalves, 1993; Azevedo, 1995; Gonçalves *et al.*, 1995; Porteiro *et al.*, 1996,

2010; Arruda, 1997; Santos et al., 1997; Harmelin-Vivien et al., 2001; Morato et al., 2001b; Cardigos et al., 2005; Floeter et al., 2008; Bertoncini et al., 2010; Botelho et al., 2010; Diogo and Pereira, 2013, 2014; Carneiro et al., 2014; Afonso, 2016; Westneat, 2016; Fauconnet et al., 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1839, 1840b, 1844a, 1849; Günther, 1862; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Dooley et al., 1985; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris et al., 1991; Floeter et al., 2008; Wirtz et al., 2008; Carneiro et al., 2014; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). The holotype (unique) of *Labrus reticulatus* Lowe 1839, synonym of *L. bergylta*, caught off in Madeira is in the NHMUK collections (NHMUK 1857.6.13.111). One specimen caught off in the coast of Lisbon is in the MB collections (MB06-004825).

***Labrus merula* Linnaeus 1758**

Common name: **Brown wrasse; Bodião-fusco**, Bedião, Bodiano, Bodião-tordo, Chita, Gidilhão, Godião, Verdelho.

References: mainland Portugal (Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Fischer et al., 1987; Almeida, 1996; Erzini et al., 1996, 1997; Garnerot et al., 2004; Ribeiro et al., 2006; Henriques et al., 2008; Carneiro et al., 2014; Craig and Pollard, 2015d; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a; Sampaio, 1904; Fowler, 1936b; Ribeiro, 1936; Albuquerque, 1954-1956; Osório de Castro, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Michel et al., 1987; Patzner et al., 1990; Lloris et al., 1991; Arruda, 1992, 1997; Carneiro et al., 2014; Craig and Pollard, 2015d; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Osório de Castro, 1967). According to Eschmeyer et al. (2019) the current geographic distribution of *Labrus merula* is the Mediterranean Sea and, in the Eastern Atlantic, from Spain and Portugal to northern Morocco and the Azores. Osório de Castro (1967) mentioned the presence of this species in the fishing grounds of Madeira without indicating the geographic location. Since then no other author reports the presence of this species in Madeira to confirm its occurrence in this EEZ.

***Labrus mixtus* Linnaeus 1758**

Common name: **Cuckoo wrasse; Bodião-canário**, Bedião, Bodiano, Bodião-malola, Canário, Carvalho, Chita, Clérigo, Dentelha, Dentilha, Gidilhão, Godião, Papagaio, Peixe-carvalho, Peixe-piça, Peixe-pissa, Peixe-pissa-legítimo, Peixe-rei-do-alto, Pissa, Taberneiro, Truta-do-alto, Verdelho.

References: mainland Portugal (NHMUK 1860.4.22.44-45, 1854-1861; de Brito Capello, 1868, 1880; Steindachner, 1868b; Posser, 1880; Nobre, 1894, 1895a, 1903, 1935; Osório, 1895; Vieira, 1898, 1900; Seabra, 1911; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Dinis, 1985; Quignard and Pras, 1986b; Fischer et al., 1987; Gomon and Forsyth, 1990; Hureau, 1991; Martins et al., 1992; Almeida, 1996; Mendes et al., 2004; ICES, 2005; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Batista et al., 2009; Garrido et al., 2009; Costa et al., 2012; Carneiro et al., 2014; Ryland and Lancaster, 2017; Martins and Carneiro, 2018; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores▲ (Osório de Castro, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris et al., 1991; Harmelin-Vivien et al., 2001; Menezes et al., 2006; Porteiro et al., 2010; Carneiro et al., 2014; Gomes-Pereira et al., 2014; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1850, 1852; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Bauchot, 1987; Gomon and Forsyth, 1990; Lloris et al., 1991; Carneiro et al., 2014; Westneat, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019). Four specimens caught off in different areas are in the MB collections: three caught off at the coast of mainland Portugal (MB06-004719, 1907 at the coast of Praia da Nazaré, MB06-004723 at the coast of Matosinhos by Isaac Newton, and MB06-005359, 2010) and one specimen caught off at the coast of Pico Island (Azores) (MB06-004666).

***Labrus viridis* Linnaeus 1758**

Common name: **Green wrasse; Bodião-tordo**, Bedião, Bodiano, Bodião, Bodião-verde, Chita, Gidilhão, Godião, Lucinha, Peixe-tordo, Tordo, Verdelho.

References: mainland Portugal (de Brito Capello, 1868, 1880; Baldaque da Silva, 1891; Nobre, 1904, 1935; Seabra, 1911; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Fischer et al., 1987; Almeida, 1996; Garnerot et al., 2004; Henriques et

al., 2008; Ribeiro *et al.*, 2012; Carneiro *et al.*, 2014; Craig *et al.*, 2015b; Westneat, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Azores (?) (Harmelin and Harmelin-Vivien, 1979; Ré, 1979; Arruda, 1997). According to Eschmeyer *et al.* (2019) the current geographic distribution of *L. viridis* is the Mediterranean Sea, Sea of Marmara, Black Sea, Portugal and southern Spain south to Morocco (Eastern Atlantic). Harmelin-Vivien *et al.* (2001) verified the identification of several species using underwater photographs taken in the Azores in 1979 and re-evaluated the presence of *Labrus viridis* as a misidentified green morph of *Labrus bergylta* Ascanius 1767. Thus, this species is not considered in the total number of valid species for the Azores.

***Lappanella fasciata* (Cocco 1833)**

Common name: **Sharp-toothed wrasse; Bodião-verdelho**, Bodiano, Bodião, Verdelhão, Verdelho.

References: mainland Portugal (OCEANA, 2011, 2014; Gomes-Pereira *et al.*, 2017), Azores (Kukuev, 2002 (PECS); Gomes-Pereira, 2010, 2016; Porteiro *et al.*, 2010, 2013; Carneiro *et al.*, 2014; Pollard, 2014; Gomes-Pereira *et al.*, 2017) and Madeira (Lowe, 1840a, 1849; Günther, 1862; Gervais and Boulart, 1877b; Vaillant, 1919; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976 (PECS); Bauchot and Quignard, 1979; Hureau and Monod, 1979 (PECS); Shcherbachov *et al.*, 1985a (PECS); Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; Christiansen *et al.*, 2009; Carneiro *et al.*, 2014; Pollard, 2014; Westneat, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). One specimen caught off at the Josephine Bank is in the MMF collections (MMF 22856, 1967).

***Syphodus bailloni* (Valenciennes 1839)**

Common name: **Baillon's wrasse; Bodião-tentilhão**, Bodiano, Bodião, Canário, Serrão, Tendelhão, Tentelhão, Tentilhão, Tentilho, Verdelhão, Verdelho.

References: mainland Portugal (de Brito Capello, 1868, 1869b, 1880; Steindachner, 1868b; Osório, 1888, 1895; Baldaque da Silva, 1891; Girard, 1892; Vieira, 1900; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Quignard and Pras, 1986b; Bauchot, 1987; Fischer *et al.*, 1987; Antunes *et al.*, 1988; Arruda *et al.*, 1988; Martins *et al.*, 1992; Almeida, 1996; Erzini *et al.*, 1997, 2001a; Almada *et al.*, 2002; Jorge *et al.*, 2002; Santos *et al.*, 2002a; Hanel *et al.*, 2003; Palma *et al.*, 2003; Garnerot *et al.*, 2004; ICES, 2005; Ribeiro *et al.*, 2006; Gomes and Olim, 2007; Henriques *et al.*, 2007, 2008; Neves *et al.*, 2008; Batista *et al.*, 2009; Veiga *et al.*, 2009; Gonçalves *et al.*, 2010; França *et al.*, 2011; Ribeiro *et al.*, 2012; Carneiro *et al.*, 2014; Craig and Pollard, 2015e; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (?) (Osório de Castro, 1967) and Madeira (Osório de Castro, 1967). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. bailloni* is in the Eastern Atlantic from North Sea to Mauritania (straying to western Mediterranean Sea). Osório de Castro (1967) mentioned the presence of this species in the fishing grounds of the Azores and Madeira without indicating the geographic location. Having in mind that this species inhabits littoral and inshore coastal habitats and the occurrence records in Canary Islands (MNHN-IC-1909-0459 and MZC 26411), its presence in Madeira is possible. However, perhaps due to the geographic isolation of Azorean archipelago together with the dynamics of boundary current interactions, climatic conditions and intrinsic limitations of the species, its presence in the Azores is considered doubtful in the current work and is not included in the total number of valid species in this EEZ.

***Syphodus caeruleus* (Azevedo 1999)**

Common name: No English common name; **Bodião-dos-Açores**.

References: Azores (Azevedo, 1999; Almada *et al.*, 2002; Hanel *et al.*, 2003; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Almada *et al.*, 2002; Froese and Pauly, 2019). The holotype of *Centrolabrus caeruleus* Azevedo 1999, synonym of *S. caeruleus*, caught off Baixa da Moaçor, Ponta Delgada, São Miguel Island (Azores) is in the MCM collections (MCM 777).

***Syphodus cinereus* (Bonnaterre 1788)**

Common name: **Grey wrasse; Bodião-cinzento**, Bodiano, Bodião, Canário, Verdelhão, Verdelho.

References: mainland Portugal ([Steindachner, 1868b](#); [Vieira, 1897](#); [Osório, 1917](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967\(?\)](#); [Quignard and Pras, 1986b](#); [Bauchot, 1987](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Cabral, 1999](#); [Almada et al., 2002](#); [Palma et al., 2003](#); [Ribeiro et al., 2006, 2012](#); [Gomes and Olim, 2007](#); [Henriques et al., 2007, 2008](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Carneiro et al., 2014](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

***Syphodus mediterraneus* (Linnaeus 1758)**

Common name: **Axillary wrasse; Bodião-do-Mediterrâneo**, Abadejo, Bodiano, Bodião, Bodião-azul, Canário, Costureira, Trombeta-roxa, Trombetão, Truta-do-alto, Verdelho.

References: mainland Portugal ([Nobre, 1895c, 1903, 1935](#); [Vieira, 1897, 1900](#); [Seabra, 1911](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bauchot and Quignard, 1979](#); [Quignard and Pras, 1986b](#); [Fischer et al., 1987](#); [Gomon and Forsyth, 1990](#); [Almada et al., 2002](#); [Henriques et al., 2007, 2008](#); [Abecasis et al., 2009](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Westneat, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Vaillant, 1919](#); [Zugmayer, 1933](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Quignard, 1966](#); [Osório de Castro, 1967](#); [Collignon and Aloncle, 1973](#); [Wood and Williams, 1974](#); [Bauchot and Quignard, 1979](#); [Harmelin and Harmelin-Vivien, 1979](#); [Michel et al., 1987](#); [Gomon and Forsyth, 1990](#); [Patzner, 1990](#); [Patzner et al., 1990, 1992](#); [Ré, 1990](#); [Arruda et al., 1992](#); [Azevedo et al., 1992](#); [Santos, 1992a](#); [Wirtz, 1994](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Porteiro et al., 1996, 2010](#); [Arruda, 1997](#); [Santos et al., 1997](#); [Harmelin-Vivien et al., 2001](#); [Cardigos et al., 2005](#); [Floeter et al., 2008](#); [Bertонcini et al., 2010](#); [Diogo and Pereira, 2013](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Westneat, 2016](#); [Almeida and Biscoito, 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1837](#); [NHMUK 1855.12.26.404, 1854-1861](#); [Günther, 1862](#); [Gervais and Boulart, 1877b](#); [Fowler, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Bauchot and Quignard, 1979](#); [Gomon and Forsyth, 1990](#); [Delgado, 1998](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Westneat, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The holotype (unique) of *Crenilabrus pictus* Lowe 1838, synonym of *S. mediterraneus*, caught off in Madeira is in the NHMUK collections (NHMUK 1855.12.26.404).

***Syphodus melops* (Linnaeus 1758)**

Common name: **Corkwing wrasse; Bodião-vulgar**, Bodiano, Bodião, Canário, Maracoto, Maragota, Maragote, Margota, Melope, Serrão, Verdelho.

References: mainland Portugal ([de Brito Capello, 1869a\(?\)](#), b, [1880](#); [Steindachner, 1868b](#); [Girard, 1892](#); [Osório, 1895](#); [Vieira, 1897, 1900](#); [Nobre, 1904, 1935](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bauchot and Quignard, 1979](#); [Quignard and Pras, 1986b](#); [Fischer et al., 1987](#); [Antunes et al., 1988](#); [Arruda, 1990](#); [Almeida, 1996](#); [Cabral, 1999](#); [Almada et al., 2002](#); [Garnerot et al., 2004](#); [Ribeiro et al., 2006](#); [Henriques et al., 2007, 2008](#); [Garrido et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [Carneiro et al., 2014](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Hilgendorf, 1888a, b](#); [Vaillant, 1919](#); [Fowler, 1936b](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Bini, 1968](#); [Bauchot and Quignard, 1979](#); [Quignard and Pras, 1986b](#); [Michel et al., 1987](#); [Chícharo and Teodósio, 1990](#); [Ré, 1990](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Santos et al., 1997\(?\)](#); [Carneiro et al., 2014](#); [Westneat, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-2005-2393, 1926](#); [Osório de Castro, 1967\(?\)](#)). The two specimens preserved in the Museum national d'histoire naturelle et Reseau des Herbiers de France were caught off on October 10, 1926. Another preserved specimen of this species caught off in Canary Islands, on February 1, 1906, the southernmost occurrence record of this species in the Eastern Atlantic, is deposited in the Lund Museum of Zoology (MZLU L906/3076, 1906). Since Osório de Castro (1967) no other author described the presence of this species in Madeira. Three specimens caught off in the coast of Faro (Algarve) by Alberto Alexandre Girard in 1908 are in the MB collections (MB06-004688), another specimen caught off at the coast of Viana do Castelo, also in 1908, is in the MB collections (MB06-004714).

***Syphodus ocellatus* (Linnaeus 1758)**

Common name: No English common name; **Bodião-de-pinta**, Bodião.

References: mainland Portugal ([de Brito Capello, 1868](#); [Seabra, 1911](#); [Nobre, 1935](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Ribeiro et al., 2006](#); [Henriques et al., 2007, 2008](#); [Veiga et al., 2010](#); [Carneiro et al., 2014](#); [Froese and Pauly, 2019](#)).

Syphodus roissali (Risso 1810)

Common name: Five-spotted wrasse; **Bodião-manchado**, Bodiano, Bodião, Canário, Verdelho.

References: mainland Portugal ([Steindachner, 1868b](#); [Osório de Castro, 1954, 1967\(?\)](#); [Bauchot and Quignard, 1979](#); [Quignard and Pras, 1986b](#); [Fischer et al., 1987](#); [Almada et al., 2002](#); [Louisy, 2005](#); [Ribeiro et al., 2006, 2012](#); [Henriques et al., 2007, 2008](#); [Abecasis et al., 2009](#); [Gonçalves et al., 2010](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Pollard and Craig, 2015a](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Syphodus rostratus (Bloch 1791)

Common name: No English common name; **Bodião-das-ervas**.

References: mainland Portugal ([Almada et al., 2002](#); [Ribeiro et al., 2006](#); [Henriques et al., 2007, 2008](#); [Carneiro et al., 2014](#)).

Syphodus tinca (Linnaeus 1758)

Common name: Peacock wrasse; **Bodião-pavão**, Bodiano, Bodião, Burro, Canário, Verdelhão, Verdelho.

References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Osório, 1888](#); [Nobre, 1903, 1935](#); [Seabra, 1911](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bauchot and Quignard, 1979](#); [Quignard and Pras, 1986b](#); [Fischer et al., 1987](#); [Louisy, 2005](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Pollard and Craig, 2015b](#); [Froese and Pauly, 2019](#)). Four specimens caught off at the coast of mainland Portugal are in the MB collections: two specimens at the coast of Setúbal, in 1908 (MB06-004744, set of two specimens), one at coast of Lisbon (MB06-004726, one specimen) and another at the coast of Estoril (MB06-004727, one specimen).

Syphodus trutta (Lowe 1834)

Common name: **Emerald wrasse**; **Bodião-mosqueado**, Bodião, Bodião-verde, Maracota, Maracoto, Truta-da-costa, Truta-verde.

References: mainland Portugal ([ICM-CSIC 7751, n.d.](#); [Abecasis et al., 2009](#); [OCEANA, 2014](#)), Azores[▲] ([Drouët, 1861b](#); [Hilgendorf, 1888a, b](#); [Regan, 1903a](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Bauchot and Blanc, 1961](#); [Osório de Castro, 1967](#); [Wood, 1974](#); [Wood and Williams, 1974](#); [Arruda, 1979a, 1980b](#); [Bauchot and Quignard, 1979](#); [Harmelin and Harmelin-Vivien, 1979](#); [Ré, 1979a, 1990](#); [Dooley et al., 1985](#); [Quignard and Pras, 1986b](#); [Gomon and Forsyth, 1990](#); [Patzner et al., 1990, 1992](#); [Santos et al., 1990b, 1994b, 1997](#); [Wirtz, 1990a, 1994](#); [Azevedo et al., 1991](#); [Lloris et al., 1991](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Santos, 1992a](#); [Azevedo, 1995](#); [Barreiros, 1995](#); [Gonçalves et al., 1995](#); [Porteiro et al., 1996](#); [Harmelin-Vivien et al., 2001](#); [Carneiro et al., 2014](#); [Afonso, 2016](#); [Westneat, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Lowe, 1833a, 1837](#); [Johnson, 1862b](#); [Günther, 1862](#); [Fowler, 1923, 1936b](#); [Noronha and Sarmento, 1948](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bauchot and Quignard, 1979](#); [Dooley et al., 1985](#); [Quignard and Pras, 1986b](#); [Gomon and Forsyth, 1990](#); [Lloris et al., 1991](#); [Abreu et al., 1995](#); [Andrade and Albuquerque, 1995](#); [Delgado, 1998](#); [Falcón et al., 2000](#); [Almada et al., 2002](#); [Hanel et al., 2003](#); [Araújo et al., 2005](#); [Ribeiro et al., 2005b](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Afonso, 2015](#); [POGID, 2016](#); [POGIS, 2016](#); [Westneat, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). The whereabouts of the holotype (unique) of *Crenilabrus trutta* Lowe 1834, synonym of *S. trutta*, caught off in Madeira are unknown.

Thalassoma pavo (Linnaeus 1758)

Common name: **Ornate wrasse**; **Bodião-verde**, Bodeão, Bodião, Bodião-meredia, Bodião-pardo, Bodião-parvo, Bodião-vermelho, Cagão, Cambulo, Carnate, Judia, Júlia, Moca, Peixe-de-rolo, Peixe-rei, Peixe-verde, Rainha, Realengo, Reilangue, Torcida, Verdelhão, Verdugo.

References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Posser, 1880](#); [Seabra, 1911](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Bauchot and Quignard, 1979](#); [Quignard and Pras, 1986b](#); [Fischer et al., 1987](#); [Gomon and Forsyth, 1990](#); [Almeida, 1996](#); [Gonçalves et al., 2004](#); [Abecasis et al., 2009](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Westneat, 2016](#); [Martins and Carneiro, 2018](#); [Encarnação et al., 2019](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores[▲] ([Drouët, 1861b](#); [Ramos, 1869](#); [Hilgendorf, 1888a](#); [Regan, 1903a](#); [Vaillant, 1919](#); [Nobre, 1924, 1930](#); [Zugmayer, 1933](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Quignard, 1966](#); [Osório de Castro, 1967](#); [Wood, 1973](#); [Mapstone and Wood, 1974, 1975](#); [Wood and Williams, 1974](#); [Bauchot](#)

and Quignard, 1979; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Saldanha, 1980, 1995; Martins, 1982a; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Michel *et al.*, 1987; Barreiros, 1990, 1995; Gomon and Forsyth, 1990; Patzner, 1990; Patzner and Santos, 1990d, 1993; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1997; Wirtz, 1990a, 1991, 1994; Azevedo *et al.*, 1991, 1992, 1995; Lloris *et al.*, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a, 1995b; Van Tassel *et al.*, 1994; Azevedo, 1995; Gonçalves *et al.*, 1995; Porteiro *et al.*, 1996, 2010; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001b; Cardigos *et al.*, 2005; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso, 2016; Westneat, 2016; Almeida and Bischof, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Cuvier and Valenciennes, 1839; Lowe, 1837, 1840a, 1843-1860, 1849; Drouët, 1861b; Günther, 1862; Vinciguerra, 1883; Fowler, 1923, 1936b; Nobre, 1935; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Augier, 1985; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Gomon and Forsyth, 1990; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Westneat, 2016; Friedlander *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). This species is sometimes confused with *Thalassoma newtoni* (Osório 1891) (Westneat, 2016). The whereabouts of the syntype of the subspecies *Julis turcica lemniscata* Lowe 1843, caught off in Madeira, synonym of *T. pavo*, are unknown.

Xyrichtys novacula (Linnaeus 1758)

Common name: **Pearly razorfish; Mordedor**, Bodião-da-areia, Cão, Coelho-da-costa, Doirada, Dourada, Elefante, Papagaio, Peixe-cão, Peixe-coelho-da-costa, Peixe-padre.

References: mainland Portugal (Quignard and Pras, 1986b; Fischer *et al.*, 1987; Carneiro *et al.*, 2014; Westneat, 2016; Froese and Pauly, 2019; Nirchio *et al.*, 2019), Azores (Hilgendorf, 1888a; Regan, 1903a; Vaillant, 1919; Nobre, 1924, 1930; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Quignard, 1966; Osório de Castro, 1967; Bauchot and Blanc, 1961; Bauchot and Quignard, 1979; Dooley *et al.*, 1985; Michel *et al.*, 1987; Gomon and Forsyth, 1990; Wirtz, 1990a, 1994; Lloris *et al.*, 1991; Arruda, 1992, 1997; Porteiro *et al.*, 1996, 2010; Santos *et al.*, 1997; Floeter *et al.*, 2008; Bertoncini *et al.*, 2010; Carneiro *et al.*, 2014; Westneat, 2016; Froese and Pauly, 2019) and Madeira (Lowe, 1837; Cuvier and Valenciennes, 1839; MNHM-IC-0000-6503, 1840; Günther, 1862; Fowler, 1923, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Bauchot and Quignard, 1979; Fischer *et al.*, 1981; Augier, 1985; Dooley *et al.*, 1985; Gomon and Forsyth, 1990; Lloris *et al.*, 1991; Wirtz, 1994; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Falcón *et al.*, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Wirtz *et al.*, 2008; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Westneat, 2016; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The syntype of *Xyrichtys cultratus* Valenciennes in Cuvier & Valenciennes 1840, synonym of *X. novacula*, caught off in Madeira is in the MNHN collections (MNHM-IC-0000-6503).

Family Scaridae

Sparisoma cretense (Linnaeus 1758)

Common name: **Parrotfish; Papagaio-velho**, Badeão, Badião, Bidião, Bidião-de-boca-curta, Bodião, Bodião-de-olho, Bodião-vermelho, Peixe-papagaio, Velha, Veja, Véja.

References: mainland Portugal (Monod, 1979g; Fischer *et al.*, 1981, 1987; Randall, 1981, 1990; Quignard and Pras, 1986b; Abecasis *et al.*, 2008; Carneiro *et al.*, 2014; Westneat, 2016; Eschmeyer *et al.*, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Regan, 1903a; Sampaio, 1904; Lampe, 1914; Roule, 1919a; Vaillant, 1919; Fowler, 1936b; Ribeiro, 1936; Collins, 1954; Albuquerque, 1954-1956; Schultz, 1958; Bauchot and Blanc, 1961; Osório de Castro, 1967; Wood, 1973; Wood and Williams, 1974; Harmelin and Harmelin-Vivien, 1979; Ré, 1979a, 1990; Monod, 1979g; Saldanha, 1980; Fischer *et al.*, 1981; Randall, 1981, 1990; Martins, 1982a; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Barreiros, 1990; Patzner, 1990; Patzner and Santos, 1990c; Patzner *et al.*, 1990, 1992; Santos *et al.*, 1990b, 1994a, 1995a, 1997; Wirtz, 1990a, 1994; Azevedo *et al.*, 1991, 1992; Lloris *et al.*, 1991; Monteiro, 1991; Arruda, 1992, 1997; Arruda *et al.*, 1992; Santos, 1992a; Azevedo, 1995; Gonçalves *et al.*, 1995; Isidro, 1996; Harmelin-Vivien *et al.*, 2001; Morato *et al.*, 2001a, b; Cardigos *et al.*,

2005; Floeter *et al.*, 2008; Abecasis *et al.*, 2009b; Bertoncini *et al.*, 2010; Botelho *et al.*, 2010; Porteiro *et al.*, 2010; Diogo and Pereira, 2013, 2014; Carneiro *et al.*, 2014; Afonso, 2016; Westneat, 2016; Froese and Pauly, 2019) and Madeira[▲] (Lowe, 1837; Cuvier and Valenciennes, 1840; Günther, 1862; Vinciguerra, 1883; de Velloso, 1892; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1967; Monod, 1979g; Fischer *et al.*, 1981; Randall, 1981, 1990; INIP, 1984b; Augier, 1985; Dooley *et al.*, 1985; Quignard and Pras, 1986b; Lloris *et al.*, 1991; Abreu *et al.*, 1995; Andrade and Albuquerque, 1995; Delgado, 1998; Falcón *et al.*, 2000; Mello, 2000; Araújo *et al.*, 2005; Ribeiro *et al.*, 2005b; Floeter *et al.*, 2008; Ribeiro, 2008; Wirtz *et al.*, 2008; Abecasis *et al.*, 2009b; EMEPC, 2010; Carneiro *et al.*, 2014; Almada *et al.*, 2015; POGID, 2016; POGIS, 2016; Westneat, 2016; Friedlander *et al.*, 2017; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of *Scarus mutabilis* Lowe 1838, synonym of *S. cretense*, caught off in Madeira are unknown.

***Sparisoma chrysopterum* (Bloch & Schneider 1801)**

Common name: **Redtail parrotfish; Papagaio**, Badeão, Badião, Bidião, Bidião-salema, Bidião-verde, Bodião, Bodião-boca-verde, Bodião-pardo, Peixe-papagaio, Veja.

References: Azores (?) (Osório de Castro, 1967) and Madeira (?) (Fowler, 1936b; Albuquerque, 1954-1956; Osório de Castro, 1967). According to Eschmeyer *et al.* (2019) and Froese and Pauly (2019) the current geographic distribution of *S. chrysopterum* is the Western Atlantic. Fowler (1936b) mentioned the existence of four specimens from Funchal (Madeira) in the U.S. National Museum, with a length between 185 and 298 mm, identified by Adolfo Cesar di Noronha as *Sparisoma flavescens* (Bloch & Schneider 1801), synonymous of *S. chrysopterum*. Albuquerque (1954-1956) reported the occurrence of this species in Madeira following the quotation of Fowler. Osório de Castro (1967) mentioned the presence of this species in the fishing grounds of the Azores and Madeira without indicating the geographic location. Nevertheless, its occurrence in those EEZs is not confirmed by more recent data. Thus, this species is not considered in the total number of valid species.

Family Zoarcidae

***Lycenchelys alba* (Vaillant 1888)**

Common name: No English common name; **Enguia-beiçuda-alva**.

References: Azores (PECS) (MNHN-IC-1886-0590, 1883; Vaillant, 1888; Brauer, 1906; Andriashev, 1979, 1986; Albuquerque, 1954-1956; Grey, 1956; Andriashev, 1979, 1986(?); Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). The lectotype of *Lycodes albus* Vaillant 1888, synonym of *L. alba*, caught off between Azores and France is in the MNHM collections (MNHN-IC-1886-0590).

***Melanostigma atlanticum* Koefoed 1952**

Common name: **Atlantic soft pout; Enguia-beiçuda-do-Atlântico**.

References: mainland Portugal (Van der Land *et al.*, 2001; Carneiro *et al.*, 2014; Nielsen *et al.*, 2015h; Eschmeyer *et al.*, 2019), Azores (Nielsen *et al.*, 2015h; Porteiro *et al.*, 2017 (PECS)) and Madeira (Nielsen *et al.*, 2015h). According to Eschmeyer *et al.* (2019) the current geographic distribution of *M. atlanticum* is the Western Mediterranean Sea, and in the eastern-north Atlantic, from Scotland to Western Sahara; Western North Atlantic: Newfoundland to south-western United States. Porteiro *et al.* (2017) mention the presence of this species in the MAR at 55°N and to about 45°N (located within the Azorean PECS area) and North-East Atlantic. Nielsen *et al.* (2015h) reported the Selvagens Islands area in the distribution map of this species.

***Pachycara thermophilum* Geistdoerfer 1994**

Common name: No English common name; **Enguia-beiçuda-termofílica**.

References: Azores (Biscoito and Almeida, 2004 (PECS); Porteiro *et al.*, 2010; Baily, 2014; Carneiro *et al.*, 2014; WoRMS Editorial Board, 2018).

***Pachycara saldanhai* Biscoito & Almeida 2004**

Common name: No English common name; **Enguia-beiçuda-de-Saldanha**.

References: Azores (PECS) (Biscoito and Almeida, 2004; Stefanni *et al.*, 2007; Biscoito *et al.*, 2009; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019). The holotype of this species caught off Rainbow hydrothermal vent field, north Azores segment of Mid-Atlantic Ridge, (36°13.43'N, 33°54.08'W) is in the

MMF collections (MMF 31373). According to Stefanni *et al.* (2007) the morphological differences reported by Biscoito and Almeida (2004) may eventually be considered phenotypic expressions associated to the large geographic distance that separates these vents. Thus, the former authors suggest a revision of the nomenclature considering *P. thermophilum* (Geistdoerfer 1994) as the only *Pachycara* species inhabiting the North Atlantic vent fields. However, Eschmeyer *et al.* (2019) continue to consider them as two valid species.

Family Anarhichadidae

Anarhichas denticulatus Krøyer 1845

Common name: **Northern wolffish; Peixe-lobo-azul**; Gata; Gata-cinzenta; Gata-de-água-má; Gata-de-água-má-cinzenta.

References: Azores (PECS) ([Porteiro *et al.*, 2017](#)).

Family Pholidae

Pholis gunnellus (Linnaeus 1758)

Common name: **Rock gunnel; Peixe-manteiga**.

References: mainland Portugal ([Vandelli, 1771](#); [ICES, 2005](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *P. gunnellus* is the North Atlantic and adjacent Arctic. The only two historical records of its occurrence are separated by more than 230 years and they are probably the most southern records of distribution of this species in the European coasts. Extremely rare species.

Family Chiasmodontidae

Chiasmodon niger Johnson 1864

Common name: **Black swallower; Peixe-glutão-negro**, Glutão, Glutão-preto, Pancinha, Peixe-glutão, Sapinho-da-fundura, Sapinho-de-fundura.

References: mainland Portugal ([de Brito Capello, 1868, 1880](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Osório, 1909](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft, 1979d](#); [Johnson and Keene, 1986](#); [Melo, 2009](#); [Prokofiev and Kukuev, 2009b](#); [Carneiro *et al.*, 2014](#); [Smith-Vaniz *et al.*, 2015d](#); [Froese and Pauly, 2019](#)), Azores ([Murray and Hjort, 1912](#); [Fowler, 1936b](#); [Koefoed, 1952](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft, 1979d](#); [Johnson and Keene, 1986](#); [Lloris *et al.*, 1991](#); [Arruda, 1997](#); [Santos *et al.*, 1997](#); [Desbruyères *et al.*, 2001](#); [Melo and Menezes, 2002](#); [Menezes *et al.*, 2006, 2012](#); [Sutton *et al.*, 2008](#); [Prokofiev and Kukuev, 2009b](#); [Porteiro *et al.*, 2010, 2017](#); [Besugo, 2013](#); [Carneiro *et al.*, 2014](#); [Smith-Vaniz *et al.*, 2015d](#); [Almeida and Biscoito, 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([NHMUK 1863.12.12.4](#), 1861-1890; [Johnson, 1864a, b](#); [Günther, 1864a, 1887](#); [Carte, 1866](#); [Goode and Bean, 1895, 1896](#); [Brauer, 1906](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Fowler, 1936b](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Krefft, 1979d](#); [INIP, 1982c, 1984a, b, c](#); [Johnson and Keene, 1986](#); [Lloris *et al.*, 1991](#); [Severino, 2004](#); [Prokofiev and Kukuev, 2009b](#); [Carneiro *et al.*, 2014](#); [Smith-Vaniz *et al.*, 2015d](#); [POGID, 2016](#); [Froese and Pauly, 2019](#)). The holotype (unique) of this species caught off in Madeira is in the NHMUK collections (NHMUK 1863.12.12.4). Another holotype of *Chiasmodon bolangeri* Osório 1909, synonym of *C. niger*, caught off at Setúbal (mainland Portugal) was preserved in the Museu Bocage (MB T135 (1506)) and destroyed in the 1978 fire.

Dysalotus alcocki MacGilchrist 1905

Common name: No English common name; **Peixe-glutão-de-Alcock**.

References: Azores ([MAR-ECO 5735, 2004 \(PECS\)](#); [Wenneck *et al.*, 2008](#); [Sutton *et al.*, 2008](#); [Carneiro *et al.*, 2014](#); [Carpenter *et al.*, 2015d](#); [Porteiro *et al.*, 2017 \(PECS\)](#)) and Madeira ([Carpenter *et al.*, 2015d](#)). Eschmeyer *et al.* (2019) mention it as a nearly circumglobal species in tropical and subtropical seas, but not present in Eastern Pacific. Melo (2017) mentions it as a widely distributed species in the Atlantic, Pacific and Indian Oceans, more frequently in low latitudes. In the Atlantic Ocean, it is more frequently found throughout the South Atlantic and the western North Atlantic Ocean, including the Gulf of Mexico and Caribbean Sea.

Dysalotus oligoscolus Johnson & Cohen 1974

Common name: No English common name; **Peixe-glutão**.

References: Azores (ZMH 114082, 1975; MAR-ECO - HamPelFish - TI/SF MAR-ECO 4788, 1982; Carneiro et al., 2014; Carpenter et al., 2015e) and Madeira (Carpenter et al., 2015e). Eschmeyer et al. (2019) mention it as a circumglobal species, except western North Atlantic. Melo (2017) mentions that *D. oligoscolus* is distributed in the Atlantic, Pacific and Indian Oceans, more frequently in high latitudes. In the Atlantic Ocean, it is known from the central South Atlantic, and from the Eastern-north Atlantic.

Kali indica Lloyd 1909

Common name: No English common name; **Peixe-glutão-do-Índico**.

References: Azores (PECS) (MAR-ECO 6971, 2004; Carneiro et al., 2014). This species is distributed circumglobally in tropical and subtropical waters, to the north (Prokofiev, 2011) and in temperate through subarctic seas (Eschmeyer et al., 2019).

Kali kerberti (Weber 1913)

Common name: No English common name; **Peixe-glutão-de-Kerbert**.

References: Azores (PECS) (ZMH 120805, 1982; Carneiro et al., 2014). First identified as *Kali normani* (Parr 1931). Melo (2008) re-evaluated *K. kerberti* as valid and identified it as senior synonym of *K. normani* (Parr 1931).

Kali macrodon (Norman 1929)

Common name: No English common name; **Peixe-glutão-dentudo**.

References: Azores (ZMH 114066, 1975; MAR-ECO 7402, 2004; Sutton et al., 2008; Carneiro et al., 2014; Ralph, 2015w; Porteiro et al., 2017) and Madeira (ZMH 109831, 1971; Krefft, 1979d; Johnson and Keene, 1986; Lloris et al., 1991; Carneiro et al., 2014; Ralph, 2015w). Eschmeyer et al. (2019) mention it as a circumglobal species in temperate through subarctic seas.

Kali macrura (Parr 1933)

Common name: No English common name; **Peixe-glutão-rabudo**.

References: Azores (MAR-ECO 1995, 2004 (PECS); Sutton et al., 2008; Carneiro et al., 2014 (PECS)) and Madeira (Krefft, 1979d; Johnson and Keene, 1986; Lloris et al., 1991; Carneiro et al., 2014). Eschmeyer et al. (2019) mention it as a circumglobal species in tropical and subtropical seas, but not in the Eastern Pacific.

Kali parri Johnson & Cohen 1974

Common name: No English common name; **Peixe-glutão-de-Parr**.

References: Azores (PECS) (MAR-ECO - HamPelFish - TI/SF MAR-ECO 4016, 1979; Carneiro et al., 2014). Eschmeyer et al. (2019) mention it as a circumglobal species in tropical and subtropical seas, except western Indian Ocean and Eastern Pacific.

Pseudoscopelus altipinnis Parr 1933

Common name: No English common name; **Peixe-glutão-de-vela**.

References: mainland Portugal (GNM 1946-8375, 1946; ZMH 105437, 1968; Krefft, 1979d; Johnson and Keene, 1986; Carneiro et al., 2014; Ralph, 2015x), Azores (Collins, 1954; MCZ 49086, 1964; Arruda, 1997; Santos et al., 1997; Porteiro et al., 1999, 2010, 2017; Menezes et al., 2006; Sutton et al., 2008; Melo, 2010; Carneiro et al., 2014; Ralph, 2015x) and Madeira (Noronha and Sarmento, 1948; Albuquerque, 1954-1956; ZMH 105435, 1968; Krefft, 1979d; Johnson and Keene, 1986; Lloris et al., 1991; Carneiro et al., 2014; Ralph, 2015x).

Pseudoscopelus obtusifrons (Fowler 1934)

Common name: No English common name; **Peixe-glutão-de-focinho-curvo**.

References: Azores (ZMH 121111, 1982 (PECS); ZMH 121112, 1982 (PECS); MAR-ECO 1211, 2004; Sutton et al., 2008; Melo, 2010 (PECS); Carneiro et al., 2014; Porteiro et al., 2017) and Madeira (Porteiro et al., 2017). According to Eschmeyer et al. (2019) the current geographic distribution of *P. obtusifrons* is the tropical Atlantic, the western Pacific and south-eastern Indian Ocean.

Pseudoscopelus sagamianus Tanaka 1908

Common name: No English common name; **Peixe-glutão-de-Sagami**.

References: Azores (PECS)(?)[\(Prokofiev, 2009\)](#). According to Roa-Varón and Iwamoto (2019b) the records of this species from the Atlantic reported by Prokofiev (2009, 2011) are misidentifications of *Pseudoscopelus scriptus* and for Melo (2010) it is a misidentification of *P. cordilluminatus*. However, there is an ongoing discussion on the correct identification of this species (Roa-Varón and Iwamoto, 2019b). Thus, this species is not considered in the total number of valid species.

***Pseudoscopelus scutatus* Krefft 1971**

Common name: No English common name; [Peixe-glutão-escudado](#).

References: Azores (MAR-ECO 4918, 2004; Sutton et al., 2008; Carneiro et al., 2014; Porteiro et al., 2017 (PECS)). According to Prokofiev and Kukuev (2006) and Eschmeyer et al. (2019) the current geographic distribution of *P. scutatus* is the Atlantic and Indo-West Pacific and most of the records are from tropical and subtropical of North and South Atlantic.

Family Ammodytidae

***Ammodytes tobianus* Linnaeus 1758**

Common name: **Small sandeel; Galeota-menor**, Agulhão, Frachão, Galeota, Galiota, Ligueirão, Linguarão, Lingueirão, Pescada-bicuda, Pica, Sandil, Sandilho, Sandilho-menor.

References: mainland Portugal[▲] ([ICM-CSIC 7724](#), n.d.; Vandelli, 1771; de Brito Capello, 1876, 1880; Osório, 1888; Almeida and Roquette, 1892; Vieira, 1897, 1900; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954(?), 1967(?); Reay, 1986; Fischer et al., 1987; Antunes et al., 1988; Arruda et al., 1988; Jorge et al., 2002; Garnerot et al., 2004; ICES, 2005; Chaves, 2008; Henriques et al., 2008; Batista et al., 2009; Garrido et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; Carneiro et al., 2014; OCEANA, 2014; Martins et al., 2015; Mecklenburg et al., 2018; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019).

***Gymnammodytes cicerelus* (Rafinesque 1810)**

Common name: **Mediterranean sand eel; Galeota-da-areia**, Bicuda, Frachão, Galeota, Ligueirão, Linguarão, Lingueirão, Pescada, Pescada-bicuda.

References: mainland Portugal ([Vieira, 1893](#); Osório, 1917; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Wheeler, 1979f; Dinis, 1985; Henriques et al., 2008; Batista et al., 2009; Carneiro et al., 2014; Heessen et al., 2015a; Nizinski, 2016; Eschmeyer et al., 2019; Froese and Pauly, 2019) and Azores ([Porteiro et al., 2010](#); Afonso et al., 2013; Carneiro et al., 2014; Nizinski, 2016).

***Gymnammodytes semisquamatus* (Jourdain 1879)**

Common name: **Smooth sandeel; Galeota-frachão**, Frachão, Galeota, Ligueirão, Linguarão, Lingueirão.

References: mainland Portugal ([Duncker and Mohr, 1939](#); Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Wheeler, 1979f; INIP, 1982b; Martins, 1981; Reay, 1986; ICES, 2005; Henriques et al., 2008; Garrido et al., 2009; Carneiro et al., 2014; Collette et al., 2014; Froese and Pauly, 2019). Two specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22810, 1967, set of two specimens).

***Hyperoplus immaculatus* (Corbin 1950)**

Common name: **Greater sand-eel; Galeota-maior-de-Corbin**.

References: mainland Portugal ([Gomes et al., 2018](#)).

***Hyperoplus lanceolatus* (Le Sauvage 1824)**

Common name: **Great sandeel; Galeota-maior**, Agulhão, Bicuda, Frachão, Ligueirão, Linguarão, Lingueirão, Galeota, Galiota, Pescada, Pescada-bicuda, Sandilho-maior.

References: mainland Portugal ([Osório, 1896, 1917](#); Vieira, 1897, 1900; Reay, 1986; Nobre, 1903, 1935; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández et al., 1981; Arruda et al., 1988; Martins et al., 1999; Palma et al., 2003; ICES, 2005; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Garrido et al., 2009; Costa et al., 2012; Carneiro et al., 2014; Froese and Pauly, 2019).

Family Trachinidae

Echiichthys vipera (Cuvier 1829)

Common name: **Lesser weever; Peixe-aranha-menor**, Aranha, Aranha-comum, Aranha-do-mar, Aranha-grande, Aranhuço, Escorpião, Esquipão, Lacraia, Peixe-aranha, Peixe-aranha-comum, Peixe-escorpião, Ratinho, Víbora.

References: mainland Portugal (de Brito Capello, 1867d, 1869b, 1880; Baldaque da Silva, 1891; Girard, 1892; Nobre, 1895c, 1903, 1935; Seabra, 1911; Murray and Hjort, 1912; Osório de Castro, 1954, 1967; Wheeler, 1979f; Fernández et al., 1981; INIP, 1981a, 1982b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Tortonese, 1986d; Fischer et al., 1987; Antunes et al., 1988; Arruda et al., 1988; Martins et al., 1999, 2015; Jorge et al., 2002; Palma et al., 2003; Prista et al., 2003; Garnerot et al., 2004; ICES, 2005; Ribeiro et al., 2006; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Garrido et al., 2009; Veiga et al., 2009; Gonçalves et al., 2010; França et al., 2011; Costa et al., 2012; Carneiro et al., 2014; Ryland and Lancaster, 2017; Baptista et al., 2019; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores (Hilgendorf, 1888a; Fowler, 1936; Osório de Castro, 1967; MNHN-IC-2009-0431, 1971; MNHN-IC-2009-0449, 1971; Arruda, 1997; Morato et al., 2001b; Porteiro et al., 2010; Carneiro et al., 2014; Afonso, 2016; Almeida and Biscoito, 2019) and Madeira (Lowe, 1837; Nunes, 1953, 1974; Osório de Castro, 1954, 1967; Wheeler, 1979f; Fischer et al., 1981; Tortonese, 1986d; Roux, 1990; Lloris et al., 1991; Araújo et al., 2005; Carneiro et al., 2014; Smith, 2016g; Eschmeyer et al., 2019; Froese and Pauly, 2019).

Trachinus araneus Cuvier 1829

Common name: **Spotted weever; Peixe-aranha-pontuado**, Aranha, Aranha-grande, Aranha-pontuada, Aranhão, Aranhuço, Escorpião, Lacraia, Peixe-aranha, Peixe-escorpião.

References: mainland Portugal (Nobre, 1895a, 1903, 1935; Seabra, 1911; Osório, 1917; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Wheeler, 1979f; Fischer et al., 1981, 1987; Tortonese, 1986d; Roux, 1990; Palma et al., 2003; Carneiro et al., 2014; Collette et al., 2015k; Smith, 2016g; Froese and Pauly, 2019). This species is sometimes confused with *Trachinus radiatus* Cuvier 1829 (Smith, 2016g).

Trachinus draco Linnaeus 1758

Common name: **Greater weever; Peixe-aranha-maior**, Aranha, Aranha-do-mar, Aranha-grande, Aranhola, Aranhuço, Escorpião, Lacraia, Peixe-aranha, Peixe-aranha-grande, Peixe-aranhola, Peixe-escorpião.

References: mainland Portugal▲ (de Brito Capello, 1867a, d, 1869b, 1880; Almeida and Roquette, 1892; Girard, 1892; Nobre, 1895a, 1903, 1935; Vieira, 1897; Seabra, 1911; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Wheeler, 1979f; Fernández et al., 1981; Fischer et al., 1981, 1987; INIP, 1981a; Martins, 1981; Borges, 1984; Dinis, 1985; Tortonese, 1986d; Antunes et al., 1988(?); Roux, 1990; Martins et al., 1992, 2015; Erzini et al., 1996, 2001a; Gonçalves et al., 1996, 1997; Borges et al., 2002; Santos et al., 2002a; Palma et al., 2003; Mendes et al., 2004; ICES, 2005; Sousa et al., 2005; Ribeiro et al., 2006; Stergiou et al., 2006; Gomes and Olim, 2007; Henriques et al., 2007, 2008; Chaves, 2008, 2009; Batista et al., 2009; Costa et al., 2012; Carneiro et al., 2014; Heessen et al., 2015b; Baptista et al., 2019; Froese and Pauly, 2019) and Madeira (Lowe, 1837; Günther, 1860; Fowler, 1923, 1936b; Zugmayer, 1933; Noronha and Sarmento, 1948; Maul, 1949; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Wheeler, 1979f; Fischer et al., 1981; Roux, 1981, 1990; Augier, 1985; Dooley et al., 1985; Tortonese, 1986d; Lloris et al., 1991; Andrade and Albuquerque, 1995; Wirtz et al., 2008; Carneiro et al., 2014; Heessen et al., 2015b; POGID, 2016; Smith, 2016g; Eschmeyer et al., 2019; Froese and Pauly, 2019).

Trachinus pellegrini Cadenat 1937

Common name: **Cape Verde weever; Peixe-aranha-de-Cabo-Verde**, Aranha-de-Cabo-Verde.

References: Madeira (Fischer et al., 1981; Carpenter et al., 2015f; González Pérez, 2015b). According to Eschmeyer et al. (2019) the current geographic distribution of *T. pellegrini* is the Canary and Cape Verde islands and from the Senegal to Cameroon. Fischer et al. (1981), Carpenter et al. (2015f) and González Pérez (2015b) reported the Selvagens Islands area in the distribution maps of this species.

Trachinus radiatus Cuvier 1829

Common name: **Starry weever; Peixe-aranha-raiado**, Aranha, Aranha-grande, Aranha-preta, Aranha-raiado, Aranhuço, Escorpião, Lacraia, Peixe-aranha, Peixe-escorpião.

References: mainland Portugal ([Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Fischer et al., 1981, 1987](#); [Dooley et al., 1985](#); [Martins, 1981](#); [Palma et al., 2003](#); [Henriques et al., 2008](#); [Carneiro et al., 2014](#); [Collette et al., 2015l](#); [Smith, 2016g](#)) and Madeira ([MNHN-IC-2005-2436, 1925](#); [Fischer et al., 1981](#); [Collette et al., 2015l](#)). This species is sometimes confused with *Trachinus araneus* Cuvier 1829 ([Smith, 2016g](#)). Fischer et al. (1981) reported the Selvagens Islands area in the distribution map of this species.

Family Uranoscopidae

Uranoscopus scaber Linnaeus 1758

Common name: **Stargazer; Cabeçudo**, Aranhuço, Astrónomo, Bufo, Masca-tabaco, Papa-tabaco, Peixe-astrónomo, Peixe-cabeçudo, Peixe-sapo, Peixe-tabaqueiro, Sapo, Sapo-cabeçudo, Tabaqueiro, Terrorista.

References: mainland Portugal ([Vandelli, 1771](#); [de Brito Capello, 1867a, d, 1880](#); [Posser, 1880](#); [Osório, 1888](#); [Girard, 1892](#); [Nobre, 1895a, 1903, 1935](#); [Vieira, 1897](#); [Bragança, 1903](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Wheeler, 1979f](#); [Fernández et al., 1981](#); [Hureau, 1986e](#); [Fischer et al., 1981, 1987](#); [Dooley et al., 1985](#); [Hureau, 1986e](#); [Antunes et al., 1988](#); [Halstead et al., 1990](#); [Martins et al., 1992](#); [Erzini et al., 2001a](#); [Palma et al., 2003](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Henriques et al., 2008](#); [Batista et al., 2009](#); [Gonçalves et al., 2010](#); [Carneiro et al., 2014](#); [Smith, 2016h](#); [Golani and Fricke, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Wirtz, 1994](#); [Abreu et al., 1995](#); [Floeter et al., 2008](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [Carneiro et al., 2014](#); [Golani and Fricke, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Family Callionymidae

Callionymus lyra Linnaeus 1758

Common name: **Dragonet; Peixe-pau-lira**, Aranha, Pau, Peixe-aranha, Peixe-pau, Peixe-pimenta, Peixe-sapo, Sapo.

References: mainland Portugal[▲] ([Vandelli, 1771](#); [de Brito Capello, 1868, 1880](#); [Steindachner, 1868a](#); [Posser, 1880](#); [Nobre, 1895c, 1903, 1904, 1935](#); [Osório, 1895, 1917](#); [Goode and Bean, 1895, 1896](#); [Vieira, 1897](#); [Bragança, 1902](#); [Seabra, 1911](#); [Gonçalves, 1941](#); [Helling, 1943](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Wheeler, 1979h](#); [Costa, 1980](#); [Fernández et al., 1981](#); [INIP, 1981a, 1982b](#); [Martins, 1981](#); [Sobral, 1981](#); [Borges, 1982](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Fricke, 1986, 2016c](#); [Fischer et al., 1987](#); [Almeida, 1988](#); [Antunes et al., 1988](#); [Arruda et al., 1988](#); [Monteiro, 1989](#); [Jorge, 1991](#); [Rebelo, 1992](#); [da Cunha and Antunes, 1993](#); [da Cunha, 1994](#); [Erzini et al., 1996, 2001a](#); [Gonçalves et al., 1996, 1997, 2006](#); [Martins et al., 1992, 1999, 2015](#); [Erzini et al., 1997](#); [Cabral, 1999](#); [Henriques et al., 1999, 2007](#); [Cabral et al., 2000](#); [Gomes et al., 2001](#); [Borges et al., 2002, 2003](#); [Santos et al., 2002a](#); [Palma et al., 2003](#); [Prista et al., 2003](#); [Garnerot et al., 2004](#); [Mendes et al., 2004](#); [ICES, 2005](#); [Sousa et al., 2005, 2006](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Castro, 2008](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Neves et al., 2008](#); [Silva, 2008, 2010](#); [Batista et al., 2009](#); [Garrido et al., 2009](#); [Veiga et al., 2009](#); [Gonçalves et al., 2010](#); [França et al., 2011](#); [Costa et al., 2012](#); [da Cunha and Antunes, 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Herler et al., 2015b](#); [Fricke and Ordines, 2017](#); [Ryland and Lancaster, 2017](#); [Baptista et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([USNM 94473, 1894](#); [Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1972](#); [Wheeler, 1978, 1979h](#); [Fricke, 1986](#); [Lloris et al., 1991](#); [Arruda, 1997](#); [Carneiro et al., 2014](#); [Herler et al., 2015b](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([MNHN-IC-1883-0035, 1882](#); [MNHN-IC-2005-2418, 1922](#); [Osório de Castro, 1967\(?\)](#); [Herler et al., 2015b](#)). William Trelease on July 3, 1894 cited this species for the Azores ([USNM 94473](#)), but since then no additional occurrences have been reported ([Santos et al., 1997](#)). The above mentioned specimen was re-examined and identified as belonging to the species *Callionymus reticulatus* Valenciennes 1837 in November, 1996 and January, 1997 ([Santos et al., 1997](#)). However, according to Eschmeyer et al. (2019) in north-eastern Atlantic, the current geographic distribution of this species is from Iceland and Norway south to Mauritania, including Azores.

Callionymus maculatus Rafinesque 1810

Common name: Spotted dragonet; **Peixe-pau-malhado**, Aranha, Pau, Pau-malhado, Peixe-aranha, Peixe-pau, Peixe-pimenta, Peixe-sapo, Sapo.

References: mainland Portugal ([Bragança, 1903](#); [Nobre, 1935](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1967](#); [Maul, 1976](#); [Muus and Dahlström, 1978](#); [Wheeler, 1979h](#); [Fernández et al., 1981](#); [INIP, 1982a, b](#); [Martins, 1981](#); [Dinis, 1985](#); [Fricke, 1986, 2016c](#); [Fischer et al., 1987](#); [Almeida, 1996](#); [Cabral, 1999](#); [Martins et al., 1999](#); [Cabral et al., 2000](#); [Gordo and Cabral, 2001](#); [Borges et al., 2002](#); [Thiel et al., 2003](#); [Gonçalves et al., 2006, 2010](#); [Ribeiro et al. 2006, 2012](#); [Castro, 2008](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Silva, 2008, 2010, 2011](#); [Garrido et al., 2009](#); [Silva et al., 2009](#); [Veiga et al., 2009](#); [França et al., 2011](#); [Costa et al., 2012](#); [da Cunha and Antunes, 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [Silva and Borges, 2014](#); [Fricke and Ordines, 2017](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). Several specimens caught off at the coast of mainland Portugal are in the MMF collections (MMF 22829, 1967, set of twelve specimens).

Callionymus pusillus Delaroche 1809

Common name: Sailfin dragonet; **Peixe-pau-rabudo**.

References: mainland Portugal ([Fricke, 1986, 2016c](#); [Fischer et al., 1987](#); [Carneiro et al., 2014](#); [Herler et al., 2014i](#); [Fricke and Ordines, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Callionymus reticulatus Valenciennes 1837

Common name: Reticulated dragonet; **Peixe-pau-listado**.

References: mainland Portugal ([Wheeler, 1979h](#); [INIP, 1982b](#); [Dinis, 1985](#); [Fricke, 1986, 2016c](#); [Fischer et al., 1987](#); [da Cunha, 1994](#); [Cabral, 1999](#); [Henriques et al., 1999, 2007, 2008](#); [Cabral et al., 2000](#); [Borges et al., 2002](#); [Prista et al., 2003](#); [ICES, 2005](#); [Gonçalves et al., 2006, 2010](#); [Neves et al., 2008](#); [Batista et al., 2009](#); [Veiga et al., 2009](#); [França et al., 2011](#); [Silva, 2011](#); [da Cunha and Antunes, 2012](#); [Ribeiro et al., 2012](#); [Carneiro et al., 2014](#); [Herler et al., 2014j](#); [Ryland and Lancaster, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Azores ([Fowler, 1936b](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Maul, 1972](#); [Wheeler, 1978](#); [Santos et al., 1997](#); [Fricke, 2002](#); [Porteiro et al., 2010](#); [Carneiro et al., 2014](#); [Herler et al., 2014j](#); [Fricke and Ordines, 2017](#); [Froese and Pauly, 2019](#)).

Callionymus risso Lesueur 1814

Common name: Risso's dragonet; **Peixe-pau-pintado**, Aranha, Pau, Peixe-aranha, Peixe-pau, Peixe-pimenta, Peixe-sapo, Sapo.

References: mainland Portugal ([Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954](#); [Wheeler, 1979h](#); [Fricke, 1986, 2016c](#); [Fischer et al., 1987](#); [da Cunha, 1994](#); [Gonçalves et al., 2006, 2010](#); [Ribeiro et al., 2006](#); [Henriques et al., 2008](#); [Veiga et al., 2009](#); [da Cunha and Antunes, 2012](#); [Carneiro et al., 2014](#); [Herler et al., 2014k](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)).

Protogrammus sousai (Maul 1972)

Common name: Meteor dragonet; **Peixe-pau-de-Sousa**.

References: mainland Portugal ([Pakhorukov, 2008](#)), Azores (PECS) ([Maul, 1972a, 1976](#); [Ehrich, 1977](#); [Davis and Fricke, 1990](#); [Fock et al., 2002](#); [Kukuev, 2002](#); [Pakhorukov, 2008](#); [Carneiro et al., 2014](#); [Biscoito, 2015g](#); [Fricke, 2016](#); [Fricke and Ordines, 2017](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira (PECS) ([Pakhorukov, 2008](#)). The holotype and three paratypes of *Callionymus sousai* Maul 1972, synonym of *P. sousai*, caught off at the Great Meteor Bank are in the MMF collections (MMF 22877, 1967, one specimen, MMF 22483a and 22483b, set of two paratypes and MMF 22389, one paratype, respectively).

Synchiropus phaeton (Günther 1861)

Common name: **Phaeton dragonet**; **Peixe-pau-rosa**.

References: mainland Portugal ([Palmer, 1971](#); [Maul, 1976](#); [Wheeler, 1979h](#); [INIP, 1981a, b, 1982a, b](#); [Fernández et al., 1981](#); [Martins, 1981](#); [Borges, 1984](#); [Dinis and Marecos, 1984](#); [Dinis, 1985](#); [Fricke, 1986](#); [Fischer et al., 1987](#); [Davis and Fricke, 1990](#); [Borges et al., 2002, 2003](#); [Moura, 2004](#); [ICES, 2005](#); [Fernandes et al., 2007](#); [Gomes and Olim, 2007](#); [Chaves, 2008, 2009](#); [Henriques et al., 2008](#); [Pakhorukov, 2008](#); [Silva, 2008, 2010, 2011](#); [Silva et al., 2009](#); [Costa et al., 2012](#); [Chaves and Silva, 2013](#); [Carneiro et al., 2014](#); [Costa, 2014](#); [OCEANA, 2014](#); [Silva and Borges, 2014](#); [Fricke, 2016](#); [Bañón et al., 2018b](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([MNHN-IC-1887-0176, 1883](#); [Vaillant, 1888](#);

Goode and Bean, 1895, 1896; Brauer, 1906; Roule and Angel, 1921; Maul, 1972a (PECS), 1976; Ehrich, 1977 (PECS); Wheeler, 1979h; Fricke, 1981, 1986; Davis and Fricke, 1990; Lloris *et al.*, 1991; Arruda, 1992, 1997; Santos *et al.*, 1997; Sousa *et al.*, 1999; Fock *et al.*, 2002 (PECS); Kukuev, 2002 (PECS); Floeter *et al.*, 2008; Pakhorukov, 2008 (PECS); Porteiro *et al.*, 2013; Carneiro *et al.*, 2014; Fricke, 2016; Bañón *et al.*, 2018b; Almeida and Biscoito, 2019; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Maul, 1976 (PECS); Fricke, 1986, 2016c; Lloris *et al.*, 1991; Floeter *et al.*, 2008; Pakhorukov, 2008 (PECS); Wirtz *et al.*, 2008; Porteiro *et al.*, 2010; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Several specimens caught off in different areas are in the MMF collections: two specimens at the coast of mainland Portugal (MMF 22862, 1967), one specimen at the Gettyburg Bank (MMF 22812, 1967), one specimen at the Josephine Bank (MMF 22867) and at the Meteor Bank (MMF 22843, one specimen, MMF 22405, 1967, set of three specimens, MMF 22869, 1967, one specimen, MMF 22373, 1967, set of two specimens and MMF 22352, 1967, set of four specimens).

Family Draconettidae

Centrodraco acanthopoma (Regan 1904)

Common name: **Atlantic deepwater dragonet; Peixe-pau-da-fundura-do-Atlântico.**

References: mainland Portugal (Fricke, 1986; Marques, 1998; Carneiro *et al.*, 2014; OCEANA, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019), Azores (PECS) (Maul, 1976; Ehrich, 1977; Fricke, 1986, 2016c; Fock *et al.*, 2002; Kukuev, 2002 (PECS); Pakhorukov, 2008; Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019) and Madeira (Maul, 1976 (PECS); Ehrich, 1977; Hureau and Monod, 1979; Fricke, 1986, 2016c; Lloris *et al.*, 1991; Pakhorukov, 2008 (PESC); Carneiro *et al.*, 2014; Eschmeyer *et al.*, 2019; Froese and Pauly, 2019). Several specimens caught off in different areas are in the MMF collections: two specimens at the Josephine Bank (MMF 22868, 1967, one specimen) and at the Meteor Bank (MMF 22542, 1967, set of three specimens, MMF 22880, 1967, one specimen).

Family Luvaridae

Luvarus imperialis Rafinesque 1810

Common name: **Luvar; Boquinho**, Louvar, Peixe-sol.

References: mainland Portugal (Decamps, 1986; Carneiro *et al.*, 2014; Craig, 2015d), Azores (Roule and Angel, 1924, 1930; Fowler, 1936b; Decamps, 1986; Wheeler, 1978; Lloris *et al.*, 1991; Arruda, 1997; Santos *et al.*, 1997; Porteiro *et al.*, 2010; de Casamajor and Morandeau, 2013; Carneiro *et al.*, 2014; Craig, 2015d; Collette, 2016c; Froese and Pauly, 2019) and Madeira (Lowe, 1837, 1840a, 1843, 1844b; Günther, 1860; Gervais and Boulart, 1877b; Goode and Bean, 1895, 1896; Fowler, 1936b; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Tropp, 1979; Decamps, 1986; Lloris *et al.*, 1991; de Casamajor and Morandeau, 2013; Carneiro *et al.*, 2014; Craig, 2015d; Collette, 2016c; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of *Astroderma plumbeum* Lowe 1843, synonym of *L. imperialis*, caught off in Madeira are unknown.

Family Acanthuridae

Acanthurus monroviae Steindachner 1876

Common name: **Monrovia doctorfish; Peixe-cirurgião**, Alicate, Asno, Asno-de-seta, Barbeiro, Canivete, Castanheta, Cirurgião, Coma, Còrinhas, Faquista, Lanceta, Lanceteiro, Peixe-canivete, Peixe-coma, Unha, Unhas.

References: mainland Portugal▲ (Horta e Costa and Gonçalves, 2013; Carneiro *et al.*, 2014; Herrera and Smith-Vaniz, 2015n) and Madeira (Floeter *et al.*, 2008).

Family Sphyraenidae

Sphyraena barracuda (Edwards 1771)

Common name: **Great barracuda; Bicuda-gigante**, Barracuda, Barracuda-grande, Barrecuda, Bicuda, Bicuda-americana, Bicuda-grande, Grande-barracuda, Grande-bicuda, Lobo-do-mar, Pescada-bicuda

References: Madeira (Wirtz, 1998; Freitas and Araújo, 2006; Wirtz *et al.*, 2008; Carneiro *et al.*, 2014; Russell, 2016c; Froese and Pauly, 2019).

Sphyraena guachancho Cuvier 1829

Common name: **Guachanche barracuda**; **Bicuda-africana**, Barracuda, Barrecuda, Bicuda, Lobo-do-mar, Pescada, Pescada-bicuda, Pescadinha.

References: Madeira ([Fischer et al., 1981](#); [Dooley et al., 2015](#); [Russell, 2016c](#)). According to Eschmeyer *et al.* (2019) the current geographic distribution of *S. guachancho* is the Western and Eastern Atlantic. Russell (2016c) reported the Selvagens Islands area in the distribution map of this species.

Sphyraena sphyraena (Linnaeus 1758)

Common name: **European barracuda**; **Bicuda-europeia**, Barracuda, Barracuda-europeia, Barrecuda, Bicuda, Bicuda-das-ilhas, Lobo-do-mar, Pescada-bicuda, Pescadinha.

References: mainland Portugal▲ ([Steindachner, 1867](#); [Osório, 1896](#); [Seabra, 1911](#); [Nobre, 1935](#); [Gonçalves, 1941](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979f](#); [Fischer et al., 1981, 1987](#); [Ben-Tuvia, 1986b](#); [Ribeiro et al., 2008](#); [Carneiro et al., 2014](#); [Smith-Vaniz and Herrera, 2015g](#); [Martins and Carneiro, 2018](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)), Azores ([Drouët 1861b](#); [Collett, 1896](#); [CAS ICH 24035, 1921](#); [Fowler, 1936a](#); [Albuquerque, 1954-1956](#); [Tortonese, 1979f](#); [SAIAB FC 36172, 1990](#); [Lloris et al., 1991](#); [Isidro, 1996](#); [Arruda, 1997](#); [Carneiro et al., 2014](#); [Smith-Vaniz and Herrera, 2015g](#); [Russell, 2016c](#); [Froese and Pauly, 2019](#); [Eschmeyer et al., 2019](#)) and Madeira▲ ([Bowdich, 1825](#); [Lowe, 1837](#); [de Velloso, 1892](#); [Collett, 1896](#); [Fowler, 1923, 1936a](#); [Noronha and Sarmento, 1934, 1948](#); [Nobre, 1935](#); [Nunes, 1953, 1974](#); [Albuquerque, 1954-1956](#); [Osório de Castro, 1954, 1967](#); [Tortonese, 1979f](#); [Fischer et al., 1981](#); [Dooley et al., 1985](#); [Ben-Tuvia, 1986b](#); [Sylva, 1990](#); [Lloris et al., 1991](#); [Edwards, 1993](#); [de Casamajor and Morandeu, 2013](#); [Carneiro et al., 2014](#); [Smith-Vaniz and Herrera, 2015g](#); [Russell, 2016c](#); [Eschmeyer et al., 2019](#)). This species is sometimes confused with *Sphyraena viridensis* Cuvier 1829 ([Russell, 2016c](#)). According to Santos *et al.* (1997) the records of this species from the Azores and Russell (2016b) from the Azores, Madeira and Canary Islands, and Wirtz *et al.* (2008) from Madeira, are probably misidentifications of *Sphyraena viridensis* Cuvier 1829. However, Smith-Vaniz and Herrera (2015e), Russell (2016c) and Eschmeyer *et al.* (2019) consider the occurrence of this species in Mediterranean Sea, Sea of Marmara, Black Sea and, in the Eastern Atlantic, from Bay of Biscay south to Angola, including Azores, Madeira and Saint Paul's Rocks islands and Froese and Pauly (2019) to Canary and Azores islands.

Sphyraena viridensis Cuvier 1829

Common name: **Yellowmouth barracuda**; **Bicuda-de-Cabo-Verde**, Bacuda, Barracuda, Barracuda-boca-amarela, Barracuda-de-Cabo-Verde, Bicuda, Bicuda-boca-amarela, Pescada-bicuda, Pescadinha.

References: mainland Portugal ([Gonçalves et al., 2004](#); [Abecasis et al., 2009](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Smith-Vaniz and Herrera, 2015h](#); [Martins and Carneiro, 2018](#)), Azores▲ ([Drouët, 1861b](#); [Ramos, 1869](#); [Hilgendorf, 1888a](#); [Collett, 1896a](#); [Sampaio, 1904](#); [Zugmayer, 1933](#); [Fowler, 1936a](#); [Ribeiro, 1936](#); [Cadenat, 1937](#); [Collins, 1954](#); [Albuquerque, 1954-1956](#); [Wood, 1973](#); [Wood and Williams, 1974](#); [Ré, 1979a, 1990](#); [Saldanha, 1980, 1995](#); [Martins, 1982a](#); [Dooley et al., 1985](#); [Patzner et al., 1990, 1992](#); [Wirtz, 1990a, 1991, 1994](#); [Arruda, 1992, 1997](#); [Arruda et al., 1992](#); [Azevedo et al., 1992](#); [Santos et al., 1994a, 1995a, 1997](#); [Gonçalves et al., 1995](#); [Isidro, 1996](#); [Menezes, 1996](#); [MCZ 149607, 1999](#); [Harmelin-Vivien et al., 2001](#); [Morato et al., 2001a, b](#); [Barreiros et al., 2002](#); [Cardigos et al., 2005](#); [Menezes et al., 2006](#); [Bertoni et al., 2010](#); [Botelho et al., 2010](#); [Porteiro et al., 2010](#); [Diogo and Pereira, 2013, 2014](#); [Carneiro et al., 2014](#); [Smith-Vaniz and Herrera, 2015h](#); [Afonso, 2016](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)) and Madeira ([Fischer et al., 1981](#); [Dooley et al., 1985](#); [Wirtz, 1994, 1998](#); [Delgado, 1998](#); [Falcón et al., 2000](#); [Mello, 2000](#); [Ribeiro et al., 2005b](#); [Ribeiro, 2008](#); [Wirtz et al., 2008](#); [EMEPC, 2010](#); [Carneiro et al., 2014](#); [Almada et al., 2015](#); [Smith-Vaniz and Herrera, 2015h](#); [POGIS, 2016](#); [Russell, 2016c](#); [Eschmeyer et al., 2019](#); [Froese and Pauly, 2019](#)). This species is sometimes confused with *Sphyraena sphyraena* (non Linnaeus 1758) ([Russell, 2016c](#)).

Family Caproidae

Antigonia capros Lowe 1843

Common name: **Deepbody boarfish**; **Periquito**, Periquito-comprido, Pimpim, Tem-te-em-pé, Tinta-em-pé.

References: mainland Portugal ([Shcherbachov et al., 1985b](#); [Quéro, 1986b](#); [Zidowitz and von Westernhagen, 2004](#); [ICES, 2005](#); [Pakhorukov, 2008](#); [Carneiro et al., 2014](#); [OCEANA, 2014](#); [Garcia, 2015h](#);

Froese and Pauly, 2019), Azores (Roule, 1919a; Fowler, 1936b; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977; Krefft, 1979e; Quéro and Duron, 1980; Dooley et al., 1985; Shcherbachov et al., 1985b (PECS); Quéro, 1986b; Lloris et al., 1991; Arruda, 1997; Fock et al., 2002 (PECS); Kukuev, 2002 (PECS); Zidowitz and von Westernhagen, 2004 (PECS); Pakhorukov, 2008 (PECS); Porteiro et al., 2010, 2013; Carneiro et al., 2014; Garcia, 2015h; Heemstra, 2016f; Bañón et al., 2019b; Froese and Pauly, 2019) and Madeira (Lowe, 1843, 1844b; Günther, 1860; Goode and Bean, 1895, 1896; Noronha and Sarmento, 1948; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976; Ehrich, 1977; Hureau and Monod, 1979; Krefft, 1979e; Dooley et al., 1985; Shcherbachov et al., 1985b (PECS); Quéro, 1986b; Lloris et al., 1991; Zidowitz and von Westernhagen, 2004; Pakhorukov, 2008 (PECS); Carneiro et al., 2014; Garcia, 2015h; Heemstra, 2016f; Bañón et al., 2019b; Froese and Pauly, 2019). The whereabouts of the holotype (unique) of this species caught off in Madeira are unknown. One specimen caught off at the Portuguese continental slope in June 2009 is in the MB collections (MB06-005299). Several specimens caught off at the Meteor Bank are in the MMF collections (MMF 22341, 1967, set of three specimens, MMF 22342, 1967, set of ten specimens and MMF 22343, 1967, set of five specimens).

***Capros aper* (Linnaeus 1758)**

Common name: **Boarfish; Pim-pim**, Advim, Mini-saia, Pau, Periquito-comprido, Periquito-redondo, Peixe-pau, Rascaço, Rascasso, Tem-te-em-pé, Tinta-em-pé.

References: mainland Portugal (de Brito Capello, 1867a, 1880; Steindachner, 1868a; Posser, 1880; Osório, 1895, 1917; Vieira, 1897, 1898; Bragança, 1903; Nobre, 1903, 1935; Seabra, 1911; Roule, 1919a; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976; Krefft, 1979e; Fernández et al., 1981; INIP, 1981a, 1982a, b; Martins, 1981; Borges, 1984; Dinis and Marecos, 1984; Dinis, 1985; Quéro, 1986b; Fischer et al., 1987; Antunes et al., 1988; Karrer and Post, 1990b; Saldanha et al., 1995; Gomes et al., 2001; Monteiro et al., 2001; Borges et al., 2002, 2003; Moura, 2004; ICES, 2005; Sousa et al., 2005, 2006; Fernandes et al., 2007; Gomes and Olim, 2007; Chaves, 2008, 2009; Henriques et al., 2008; Silva, 2008, 2010, 2011; Abecasis et al., 2009; Batista et al., 2009; Silva et al., 2009; Costa et al., 2012; Chaves and Silva, 2013; Carneiro et al., 2014; Christiansen et al., 2014; Costa, 2014; Silva and Borges, 2014; OCEANA, 2014; López and Garcia, 2015; Eschmeyer et al., 2019; Froese and Pauly, 2019), Azores[▲] (Hilgendorf, 1888a; Collett, 1896a; Regan, 1903a; Murray and Hjort, 1912; Richard, 1900; Zugmayer, 1911b; Vaillant, 1919; Roule and Angel, 1923; Fowler, 1936b; Koefoed, 1952; Collins, 1954; Albuquerque, 1954-1956; Osório de Castro, 1967; Maul, 1976 (PECS); Ehrich, 1977; Martins, 1982b; INIP, 1985b, c; Quéro, 1986b; Pereira, 1987b, 1992d, 1995; Dias et al., 1990; Dias, 1991; Lloris et al., 1991; Monteiro, 1991, 1996; Arruda, 1992, 1997; Arruda et al., 1992; Hamer et al., 1994; Nash et al., 1994a; Pierce et al., 1994; Wirtz, 1994; Azevedo, 1995; Clarke and Martins, 1995; Clarke et al., 1995; Granadeiro et al., 1995; Pinho et al., 1995; Santos and Nash, 1995; Silva et al., 1995; Menezes, 1996; Monteiro et al., 1996a, b; Santos et al., 1997; Kukuev, 2002 (PECS); Menezes et al., 2006; Pakhorukov, 2008 (PECS); Sutton et al., 2008; Porteiro et al., 2010; 2017; Carneiro et al., 2014; López and Garcia, 2015; Heemstra, 2016f; Froese and Pauly, 2019) and Madeira (Lowe, 1837; Günther, 1860; Gervais and Boulart, 1877b; Goode and Bean, 1895, 1896; Roule, 1919a; Bowdich, 1825; Fowler, 1936b; Noronha and Sarmento, 1948; Nunes, 1953, 1974; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Maul, 1976 (PECS); Ehrich, 1977; Krefft, 1979e; Augier, 1985; Quéro, 1986b; Karrer and Post, 1990b; Lloris et al., 1991; Abreu et al., 1995; Delgado, 1998; Wirtz et al., 2008; Christiansen et al., 2009; Carneiro et al., 2014; López and Garcia, 2015; Heemstra, 2016f; POGID, 2016; POGIS, 2016; Bischoff et al., 2017; Porteiro et al., 2017; Eschmeyer et al., 2019; Froese and Pauly, 2019). One specimen caught off at the Portuguese continental slope in April 2011 is in the MB collections. Five specimens caught off at the Josephine Bank and Meteor Bank in 1967 are in the MMF collections (MMF 22436, 1967, set of three specimens and MMF 22842, 1967, set of two specimens, respectively).

Family Gasterosteidae

***Gasterosteus aculeatus* Linnaeus 1758**

Common name: **Three-spined stickleback; Esgana-gata**, Coroa-de-espinhos, Esgana-gata, Esgana-gato, Espinal, Espinhela, Espinho, Lixa, Peixe-espinho, Peixe-lixa.

References: mainland Portugal ([de Brito Capello, 1867a, d, 1880; Gervais and Boulart, 1876; Posser, 1880; Osório, 1888; Vieira, 1894a, 1900; Seabra, 1911; Nobre, 1931, 1935; Gonçalves, 1941; Helling, 1943; Albuquerque, 1954-1956; Muus and Dahlström, 1978; Monod, 1979c; Banister, 1986; Garnerot et al., 2004; Carneiro et al., 2014; Ryland and Lancaster, 2017; Froese and Pauly, 2019](#)).

***Spinachia spinachia* (Linnaeus 1758)**

Common name: **Sea stickleback; Esgana-gata-marinha**, Esgana-gata, Esgana-gata-do-mar, Esgana-gato, Esganagata.

References: mainland Portugal ([Vandelli, 1771; de Brito Capello, 1876, 1880; Seabra, 1911; Nobre, 1935; Gonçalves, 1941; Albuquerque, 1954-1956; Osório de Castro, 1954, 1967; Fernández-Cerdeiro, 1997; Pombo et al., 2002; Carneiro et al., 2014](#)). According to Eschmeyer et al. (2019) the current geographic distribution of *S. spinachia* is the Baltic Sea, North Sea and in the Eastern North Atlantic, from Norway to Bay of Biscay. However, Nobre (1935) reported the occurrence of a specimen in Leixões Bay, near Porto. There is also a specimen in the Science Museum of the University of Porto that was collected in Foz do Douro (December 1938) (Fernández-Cerdeiro, 1997).

Global appraisal of taxonomic diversity

The checklist comprises a total of 1,349 species, for the whole study area, where 87 were considered of doubtful occurrence.

A total of 1,262 were considered valid species, which are allocated to 3 super-classes, 4 classes, 51 orders, 219 families and 645 genera.

In 2019, there are 17,538 worldwide valid marine fish species and the Portuguese ichthyofauna account for about 7.2 % of the world's biodiversity.

The order Perciformes shows the highest diversity, with 37 families, 106 genera and 107 species. Among the families, Stomiidae (90 species), Myctophidae (69 species), Macrouridae and Gobiidae (37 species each) stand out for their higher specific richness.

The total valid species occurring in Portuguese waters or under national jurisdiction add up to 1,191 and 71 are found exclusively in the platform extension area. The largest number of species was recorded for Madeira (920), followed by the mainland (845) and the Azores (761).

A total of 148 species occurrences were included in the PECS area (140 in the Azores, 15 in Madeira and 5 in the mainland), five of which are common to the Azores and Madeira platform extensions and two are common to the mainland and the Azores platform extensions.

Only 235 (19.8 %) of all species occurring in Portuguese waters or under Portuguese jurisdiction are commercially exploited. On the mainland Portugal, 184 species are traded, which represent 21.8 % of those occurring in this area. In the Azores the number of species marketed is 102 and 97 in Madeira which correspond to 13.4 % and 10.4 % of the total species occurring in each area, respectively.

This checklist includes invasive species (e.g. *Cynoscion regalis* (Bloch & Schneider 1801) in mainland Portugal and *Sparus aurata* Linnaeus 1758 in Madeira) that have been introduced by different pathways (e.g. intentional, aquaculture, aquarophilia and ballast waters) leading to changes in biodiversity.

Avaliação global da diversidade taxonómica

A lista comprehende um total de 1349 espécies, 87 das quais foram consideradas de ocorrência duvidosa na área de estudo.

As 1262 espécies consideradas válidas distribuem-se por 3 superclasses, 4 classes, 51 ordens, 219 famílias e 645 géneros.

Em 2019, encontram-se contabilizadas, a nível mundial, 17 538 espécies de peixes marinhos consideradas válidas, representando a ictiofauna marinha portuguesa cerca de 7,2 % desta biodiversidade.

A ordem dos Perciformes apresenta a maior diversidade, com 37 famílias, 106 géneros e 107 espécies. De entre as famílias, destacam-se Stomiidae (90 espécies), Myctophidae (69 espécies), Macrouridae e Gobiidae (37 espécies cada) por apresentarem maior riqueza específica.

Das espécies consideradas como válidas, 1191 ocorrem em águas portuguesas ou sob jurisdição nacional e 71 encontram-se exclusivamente na área da extensão da plataforma. O maior número foi registado na Madeira (920), seguida pelo continente (845) e pelos Açores (761).

Na área do PECS foi assinalada a ocorrência de 148 espécies (140 nos Açores, 15 na Madeira e 5 no continente), cinco das quais são comuns às extensões da plataforma dos Açores e da Madeira e duas são comuns às extensões das plataformas do continente e dos Açores.

De todas as espécies que ocorrem em águas portuguesas ou sob jurisdição portuguesa apenas 235 (19,8 %) são alvo de exploração comercial. No continente são comercializadas 184 espécies representando 21,8 % das existentes nesta área. Nos Açores são 102 e na Madeira 97 que correspondem, respetivamente, a 13,4 % e 10,4 % das espécies que ocorrem nestas áreas.

Na lista constam espécies invasoras (por exemplo, *Cynoscion regalis* (Bloch & Schneider 1801), no continente e *Sparus aurata* Linnaeus 1758, na Madeira) que têm sido introduzidas por diferentes vias (por exemplo, intencional, aquacultura, aquarofilia e águas de balastro), podendo levar a alterações de biodiversidade.

Taking into consideration the criteria for the assignment of Portuguese common names, new 874 names are proposed. It was intended to harmonize the existing common names or to give a common name for the first time.

Significant changes in the structure and number of species included in the different taxonomic groups of the Portuguese ichthyofauna were observed relatively to previous studies. This is due to the use of a different classification taxonomic system [Nelson (2006) vs Van der Laan *et al.* (2019)], because these classification systems do not follow the same taxonomic structure.

The use of molecular biological techniques provides information that can significantly modify the natural groups which could not be done based on the methodology followed in traditional taxonomic studies.

Final comments

During the elaboration of this checklist some gaps and inaccuracies were detected, where the most significant are the following:

- Some databases include errors in either species identification or geographic coordinates and/or in the *Datum* system records and their associated error. Thus, it is necessary the improvement of databases by addition of complementary information and/or corrections.
- In some museum collections there are specimens which are misidentified and/or mislabelled and therefore a re-evaluation is needed.
- Larvae specimens of *Allothunnus fallai* Serventy 1948 and *Thunnus maccoyii* (Castelnau 1872) were collected in the Northern hemisphere (including the Azores and mainland Portugal) (Nishikawa *et al.*, 1985), but they are currently circumglobal species in the Southern hemisphere (Eschmeyer *et al.*, 2019), so they were not considered in this checklist.
- In the geographic distribution maps of some species, the authors include the Selvagens Islands, which are an integral part of the Madeira archipelago, but they do not mention their occurrence in Madeira.

Tendo em consideração os critérios para a atribuição de nomes vulgares, são propostos 874 novos nomes em português. Estes nomes visam harmonizar designações vulgares já existentes ou, pela primeira vez, atribuir um nome vulgar.

Verificaram-se alterações significativas na estrutura e no número de espécies incluídas nos diferentes grupos taxonómicos da ictiofauna portuguesa relativamente a estudos anteriores como resultado da adoção de um sistema taxonómico de classificação diferente (Nelson (2006) vs Van der Laan *et al.* (2019), sistemas estes que não adotam a mesma estrutura taxonómica).

O recurso a técnicas de biologia molecular permite obter informação que pode alterar significativamente os grupos naturais o que não era possível realizar com base na metodologia seguida em estudos taxonómicos tradicionais.

Considerações finais

No decorrer da elaboração desta lista foram detetadas algumas lacunas e incorreções, das quais importa salientar as seguintes:

- Algumas bases de dados incluem erros, quer na identificação de espécies, quer nos registos das coordenadas geográficas e/ou no sistema *Datum* e respetivo erro associado, pelo que é necessário melhorar essas bases pelo envio de informação complementar e/ou correção do erro.
- Nalgumas coleções museológicas encontram-se espécimes que apresentam erros de identificação e/ou etiquetagem e carecem de verificação, pelo que seria conveniente a sua reavaliação.
- Amostras de larvas de *Allothunnus fallai* Serventy 1948 e de *Thunnus maccoyii* (Castelnau 1872) foram coletadas no hemisfério Norte (incluindo as áreas dos Açores e Portugal continental) (Nishikawa *et al.*, 1985), mas a atual distribuição geográfica destas espécies é circumglobal no hemisfério Sul (Eschmeyer *et al.*, 2019), pelo que não foram consideradas nesta lista.
- Nos mapas de distribuição geográfica de algumas espécies, os autores incluem as Ilhas Selvagens, que fazem parte integrante do arquipélago da Madeira, mas não referem a sua ocorrência na Madeira.

Finally, this checklist highlights the specific biodiversity of the Portuguese ichthyofauna and the richness of the precious linguistic heritage registered since the last quarter of the 18th century. The denomination of new common names and the harmonization of existing Portuguese names is a proposal for the Portuguese nomenclature of marine fish.

Por último, a presente lista, evidencia a biodiversidade específica da ictiofauna portuguesa e a riqueza do valioso património linguístico coligido desde o último quartel do século XVIII. A atribuição de novos nomes vulgares e a harmonização das denominações portuguesas já existentes constituem uma proposta para a nomenclatura portuguesa dos peixes marinhos.

Acknowledgements

The authors are grateful to the following individuals and institutions for their assistance on various aspects of this study:

Bárbara Serra-Pereira (Instituto Português do Mar e da Atmosfera / Divisão de Modelação e Gestão de Recursos de Pesca); Bill Eschmeyer (Professor *Emeritus* at California Academy of Sciences); Filipe O. Costa (Departamento de Biologia da Universidade do Minho); Helena Abreu (Centro de Informação e Documentação / Biblioteca Professor Luiz Saldanha da Estação de Biologia Marinha do Funchal); Inês Tojeira (Estrutura de Missão para a Extensão da Plataforma Continental, Paço de Arcos); Ingvar Byrkjedal (University Museum of Bergen); Kei Matsuura (National Museum of Nature and Science, Tokyo); Kevin Swagel (The Field Museum of Natural History, Chicago); Magnus Lundgren (Conservation Photographer & Writer); Marcelo Melo (Departamento de Oceanografia Biológica da Universidade de São Paulo); Maria Judite Alves (Museu Bocage / Museu Nacional de História Natural); Matthias Stehmann (ICHTHYS - Ichthyological Research Laboratory and Consultant, Hamburg); Monica Landi (Centro de Biologia Molecular e Ambiental da Universidade do Minho); Paulo Vaz Pires (Departamento de Produção Aquática - ICBAS da Universidade do Porto); Richard Prum (Department of Ecology and Evolutionary Biology at Yale University / Peabody Museum of Natural History, New Haven); Ronald Fricke (Department of Zoology State Museum of Natural History Stuttgart); Zouhaira Gabsi (Muséum national d'Histoire naturelle, Paris).

We wish also to thank to Francisco Ruano for the revision and comments which contributed to improve this work.

Agradecimentos

Os autores agradecem às seguintes pessoas e instituições pelo apoio em vários aspectos deste estudo:

Desejamos igualmente agradecer a Francisco Ruano pela revisão e comentários que contribuíram para melhorar este trabalho.

Published on line, December 2019

Alphabetical indexes of scientific and common names**Índices alfabéticos dos nomes científicos e comuns****Index of scientific names / Índice dos nomes científicos**

Scientific name	English common name	Portuguese common name	page
<i>Abudeodus saxatilis</i>	Sergeant-major	Castanheta-raiada	278
<i>Abyssobrotula galatheae</i>	No English common name	Abadejo-abissal	156
<i>Acanthochaenus luetkenii</i>	Pricklefish	Peixe-de-coroa-espinhosa	153
<i>Acanthocybium solandri</i>	Wahoo	Serra-da-Índia	166
<i>Acantholabrus palloni</i>	Scale-rayed wrasse	Bodião-do-alto	280
<i>Acanthonus armatus</i>	Bony-eared assfish	Abadejo-espinhosos	156
<i>Acanthostracion notacanthus</i>	Island boxfish	Peixe-cofre	226
<i>Acanthostracion quadricornis</i>	Scrawled cowfish	Peixe-cofre-touro	226
<i>Acanthurus monroviae</i>	Monrovia doctorfish	Peixe-cirurgião	295
<i>Acipenser naccarii</i>	Adriatic sturgeon	Esturjão-do-Adriático	55
<i>Acipenser sturio</i>	Sturgeon	Esturjão	55
<i>Aetomylaeus bovinus</i>	Bull ray	Ratão-bispo	51
<i>Ahliasaurus berryi</i>	No English common name	Peixe-vigia-de-Berry	107
<i>Aidablennius sphynx</i>	No English common name	Marachomba-esfinge	207
<i>Aldrovandia affinis</i>	Pale halosaur	Peixe-lagarto-pálido	56
<i>Aldrovandia gracilis</i>	Slender halosaur	Peixe-lagarto-delgado	56
<i>Aldrovandia oleosa</i>	Oily halosaur	Peixe-lagarto-oleoso	56
<i>Aldrovandia phalacra</i>	Hawaiian halosaurid fish	Peixe-lagarto-dos-Açores	56
<i>Aldrovandia rostrata</i>	Sharpnose halosaur	Peixe-lagarto-de-bico	56
<i>Alectis alexandrina</i>	Alexandria pompano	Xareu-enxada	250
<i>Alepisaurus brevirostris</i>	Short snouted lancetfish	Lírio-ferro-de-focinho-curto	113
<i>Alepisaurus ferox</i>	Long snouted lancetfish	Lírio-ferro-de-focinho-comprido	113
<i>Alepocephalus agassizii</i>	Agassiz' slickhead	Celindra-negra-de-Agassiz	74
<i>Alepocephalus australis</i>	Small scaled brown slickhead	Celindra-austral	74
<i>Alepocephalus bairdii</i>	Baird's slickhead	Celindra-de-Baird	74
<i>Alepocephalus productus</i>	Smalleye smooth-head	Celindra-de-olho-pequeno	75
<i>Alepocephalus rostratus</i>	Risso's smooth-head	Celindra-de-Risso	75
<i>Allocyttus verrucosus</i>	Warty dory	Galo-verrugoso	129
<i>Alopias superciliosus</i>	Pelagic thresher or Thresher	Tubarão-raposo-olhudo	19
<i>Alopias vulpinus</i>	Pelagic thresher or Bigeye thresher	Tubarão-raposo	20
<i>Alosa alosa</i>	Allis shad	Sável	71
<i>Alosa fallax</i>	Twaite shad	Savelha	72
<i>Aluterus monoceros</i>	Unicorn leatherjacket filefish	Peixe-porco-galhudo	226
<i>Aluterus scriptus</i>	Scribbled leatherjacket filefish	Peixe-porco-bicudo	226
<i>Amblyraja jensenii</i>	Shorttail skate	Raia-de-cauda-curta	42
<i>Amblyraja radiata</i>	Starry ray	Raia-repregada	42
<i>Ammodytes tobianus</i>	Small sandeel	Galeota-menor	291
<i>Anarchias longicauda</i>	Pygmy moray	Moreia-de-cauda-longa	60
<i>Anarchias similis</i>	Pygmy moray	Moreia-anã	60
<i>Anarhichas denticulatus</i>	Northern wolffish	Peixe-lobo-azul	289
<i>Anguilla anguilla</i>	European eel	Enguia	69
<i>Anguilla rostrata</i>	American eel	Enguia-americana	69
<i>Anoplogaster cornuta</i>	Common fangtooth	Peixe-ogre	156
<i>Anotopterus pharao</i>	Daggertooth	Barracudina-dente-de-adaga	112
<i>Antennarius multiocellatus</i>	Longlure frogfish	Peixe-sapo-multiocelado	213
<i>Antennatus nummifer</i>	Spotfin frogfish	Peixe-sapo-salpicado	213
<i>Anthias anthias</i>	Swallowtail seaperch	Canário-do-mar	242
<i>Antigonia capros</i>	Deepbody boarfish	Periquito	296
<i>Antimora rostrata</i>	Blue antimora	Mora-azul	139
<i>Aphanopus carbo</i>	Black scabbardfish	Peixe-espada-preto	164

Scientific name	English common name	Portuguese common name	page
<i>Aphanopus intermedius</i>	Intermediate scabbardfish	Peixe-espada-intermédio	164
<i>Aphia minuta</i>	Transparent goby	Caboz-transparente	178
<i>Aphyonus gelatinosus</i>	Gelatinous blindfish	Abadejo-cego-gelatinoso	160
<i>Apletodon dentatus</i>	Small-headed clingfish	Sugador-de-cabeça-pequena	204
<i>Apletodon incognitus</i>	No English common name	Sugador-dos-ouriços	204
<i>Apletodon pellegrini</i>	Chubby clingfish	Sugador-bochechudo	205
<i>Apogon imberbis</i>	Cardinal fish	Alcarraz	248
<i>Apristurus laurussonii</i>	Madeira catshark	Tubarão-gato-da-Madeira	20
<i>Apristurus manis</i>	Ghost catshark	Tubarão-gato-fantasma	20
<i>Apristurus microps</i>	Smalleye catshark	Tubarão-gato-olhinho	20
<i>Apristurus profundorum</i>	Deep-water catshark	Tubarão-gato-abissal	21
<i>Apterichtus anguiformis</i>	Slender finless eel	Cobra-de-boca-negra	64
<i>Apterichtus caecus</i>	European finless eel	Congrinho-da-areia	64
<i>Arctozenus risso</i>	Spotted barracudina	Barracudina-pintada	109
<i>Argentina sphyraena</i>	Argentine	Argentina-branca	82
<i>Argyripricus atlanticus</i>	No English common name	Pai-velho-do-Atlântico	89
<i>Argyropelecus aculeatus</i>	Atlantic silver hatchetfish	Pai-velho-prateado-do-Atlântico	89
<i>Argyropelecus affinis</i>	Pacific hatchet fish	Pai-velho-do-Pacifico	89
<i>Argyropelecus bogachei</i>	No English common name	Pai-velho-de-Bocage	89
<i>Argyropelecus gigas</i>	Hatchetfish	Pai-velho-gigante	90
<i>Argyropelecus hemigymnus</i>	Half-naked hatchetfish	Pai-velho-manchado	90
<i>Argyropelecus olfersii</i>	Silver hatchetfish	Pai-velho-prateado	90
<i>Argyropelecus sladeni</i>	Sladen's hatchet fish	Pai-velho-de-Sladen	90
<i>Argyrosomus regius</i>	Meagre	Corvina-legítima	275
<i>Ariosoma balearicum</i>	Bandtooth conger	Congro-das-Baleares	65
<i>Aristostomias grimaldii</i>	No English common name	Peixe-dragão-de-Grimaldi	93
<i>Aristostomias lunifer</i>	No English common name	Peixe-dragão-lunar	93
<i>Aristostomias tittmanni</i>	Loosejaw	Peixe-dragão-de-Tittmann	93
<i>Aristostomias xenostoma</i>	No English common name	Peixe-dragão-das-Caraíbas	93
<i>Arnoglossus grohmanni</i>	No English common name	Carta-de-Grohmann	188
<i>Arnoglossus imperialis</i>	Imperial scaldfish	Carta-imperial	189
<i>Arnoglossus laterna</i>	Mediterranean scaldfish	Carta-do-Mediterrâneo	189
<i>Arnoglossus rueppellii</i>	Rüppell's scaldback	Carta-estreita	189
<i>Arnoglossus thori</i>	Thor's scaldfish	Carta-pontuada	190
<i>Asquamiceps caeruleus</i>	No English common name	Celindra-de-cabeça-azul	75
<i>Asquamiceps hjorti</i>	Barethroat slickhead	Celindra-de-Hjort	75
<i>Asquamiceps velaris</i>	Fanfin smooth-head	Celindra-de-leque	75
<i>Asthenomacrurus victoris</i>	Victory whiptail	Largartixa-do-mar-de-Vitória	131
<i>Astronesthes atlanticus</i>	No English common name	Peixe-dragão-do-Atlântico	93
<i>Astronesthes cyaneus</i>	No English common name	Peixe-dragão-azulado	93
<i>Astronesthes gemmifer</i>	Snaggletooth	Peixe-dragão-da-Terra Nova	94
<i>Astronesthes indicus</i>	Black snaggletooth	Peixe-dragão-do-Índico	94
<i>Astronesthes leucopogon</i>	No English common name	Peixe-dragão-de-barbilo-branco	94
<i>Astronesthes macropogon</i>	No English common name	Peixe-dragão-de-barbilo-graúdo	94
<i>Astronesthes micropogon</i>	Snaggletooth	Peixe-dragão-de-barba-curta	94
<i>Astronesthes neopogon</i>	Snaggletooth	Peixe-dragão-de-barbilo-plano	94
<i>Astronesthes niger</i>	Snaggletooth	Peixe-dragão-manchado	94
<i>Astronesthes richardsoni</i>	Richardson's snaggletooth	Peixe-dragão-de-Richardson	95
<i>Astronesthes similis</i>	No English common name	Peixe-dragão-de-bolbo-filamentoso	95
<i>Astronesthes zharovi</i>	No English common name	Peixe-dragão-de-Zharov	95
<i>Ataxolepis apus</i>	No English common name	Peixe-baleia-mosaico	153
<i>Atherina boyeri</i>	Big-scale sand smelt	Peixe-rei-do-Mediterrâneo	197
<i>Atherina hepsetus</i>	Mediterranean sand smelt	Peixe-rei-do-alto	197
<i>Atherina presbyter</i>	Sand smelt	Peixe-rei	198
<i>Aulopus filamentosus</i>	Royal flagfin	Lagarto-do-mar	104

Scientific name	English common name	Portuguese common name	page
<i>Aulostomus cf. maculatus</i>	Atlantic trumpetfish	Trombeta-salpicada	174
<i>Aulostomus strigosus</i>	Atlantic cornetfish	Trombeta	174
<i>Auxis rochei</i>	Bullet tuna	Judeu	166
<i>Auxis thazard</i>	Frigate tuna	Judeu-liso	166
<i>Avocettina infans</i>	Avocet snipe eel	Cobra-centopeia	67
<i>Bajacalifornia megalops</i>	Bigeye smooth-head	Celindra-de-olho-grande	75
<i>Balistes capricornis</i>	Grey triggerfish	Cangulo-cinzento	227
<i>Balistes punctatus</i>	Bluespotted triggerfish	Cangulo-pintado	227
<i>Balistes vetula</i>	Queen triggerfish	Cangulo-real	228
<i>Barathrites iris</i>	No English common name	Abadejo-arco-íris	156
<i>Barathronus multidens</i>	No English common name	Abadejo-cego-dentudo	160
<i>Barathronus parfaiti</i>	No English common name	Abadejo-cego-de-Vaillant	160
<i>Barathronus roulei</i>	No English common name	Abadejo-cego-de-Roule	160
<i>Barbantus curvifrons</i>	Palebelly searsid	Celindra-de-tubo-de-ventre-claro	80
<i>Barbourisia rufa</i>	Velvet whalefish	Peixe-baleia-veludo	153
<i>Bassozetus compressus</i>	Abyssal assfish	Abadejo-da-fundura	157
<i>Bassozetus levistomatus</i>	No English common name	Abadejo-boca-doce	157
<i>Bassozetus nielseni</i>	Masked assfish	Abadejo-de-Nielsen	157
<i>Bassozetus taenia</i>	No English common name	Abadejo-ténia	157
<i>Bathophilus brevis</i>	No English common name	Peixe-dragão-oval	95
<i>Bathophilus digitatus</i>	Scaleless black dragonfish	Peixe-dragão-iridescente	95
<i>Bathophilus longipinnis</i>	Longfin dragonfish	Peixe-dragão-de-barbatana-longa	95
<i>Bathophilus metallicus</i>	No English common name	Peixe-dragão-de-Welsh	95
<i>Bathophilus nigerrimus</i>	Scaleless dragonfish	Peixe-dragão-chicote	96
<i>Bathophilus pawnee</i>	Pawnee dragonfish	Peixe-dragão-de-Pawnee	96
<i>Bathophilus proximus</i>	No English common name	Peixe-dragão-das-Bermudas	96
<i>Bathophilus schizochirus</i>	No English common name	Peixe-dragão-cinzento	96
<i>Bathophilus vaillanti</i>	Scaleless dragonfish	Peixe-dragão-de-Vaillant	96
<i>Bathygadus favosus</i>	No English common name	Lagartixa-do-mar-favosa	131
<i>Bathygadus melanobranchus</i>	Vaillant's grenadier	Lagartixa-do-mar-de-Vaillant	131
<i>Bathylaco nigricans</i>	Black warrior	Celindra-negra	76
<i>Bathylagichthys greyae</i>	Grey's deepsea smelt	Argentina-da-fundura-de-Grey	84
<i>Bathylagus euryops</i>	Goiter blacksmelt	Argentina-da-fundura-negra	84
<i>Bathylichnops brachyrhynchus</i>	No English common name	Peixe-fantasma-dardo-listado	84
<i>Bathylichnops exilis</i>	Javelin spookfish	Peixe-fantasma-dardo	84
<i>Bathymicrops multispinis</i>	No English common name	Peixe-tripé-espinhoso	105
<i>Bathymicrops regis</i>	No English common name	Peixe-tripé-real	105
<i>Bathyonus laticeps</i>	No English common name	Abadejo-cabeçudo	157
<i>Bathyprion danae</i>	Fangtooth smooth-head	Celindra-de-Dana	76
<i>Bathypterois dubius</i>	Spiderfish	Peixe-tripé-filamentoso	105
<i>Bathypterois grallator</i>	Tripodfish	Peixe-tripé	106
<i>Bathypterois longipes</i>	Abyssal spiderfish	Peixe-tripé-abissal	106
<i>Bathypterois phenax</i>	Blackfin spiderfish	Peixe-tripé-de-barbatana-negra	106
<i>Bathyraja pallida</i>	Pale ray	Raia-pálida	49
<i>Bathyraja richardsoni</i>	Richardson's ray	Raia-de-Richardson	49
<i>Bathysaurus ferox</i>	Deep-sea lizardfish	Peixe-lagarto-da-fundura	109
<i>Bathysaurus mollis</i>	Highfin lizardfish	Peixe-lagarto-da-costa	109
<i>Bathysolea profundicola</i>	Deep water sole	Linguado-da-fundura	191
<i>Bathytoshia lata</i>	Brown stingray	Uge-de-cardas	49
<i>Bathytroctes macrolepis</i>	Koefoed's smooth-head	Celindra-escamuda	76
<i>Bathytroctes michaelsarsi</i>	Michael Sars smooth-head	Celindra-de-Michael-Sars	76
<i>Bathytroctes microlepis</i>	Smallscale smooth-head	Celindra-de-escama-pequena	76
<i>Bathytyphlops marionae</i>	Marion's spiderfish	Peixe-tripé-de-Marion	106
<i>Bathytyphlops sewelli</i>	No English common name	Peixe-tripé-de-Sewell	106
<i>Bathyuroconger vicinus</i>	Long-tooth conger	Congro-castanho-dentão	66

Scientific name	English common name	Portuguese common name	page
<i>Bellottia apoda</i>	No English common name	Brótula-coxa	160
<i>Belone acus</i>	Garfish	Agulha-do-Mediterrâneo	198
<i>Belone belone</i>	Garfish	Agulha	199
<i>Belone svetovidovi</i>	Short-beaked garfish	Agulha-do-bico-curto	199
<i>Benthalbella infans</i>	Zugmayer's pearleye	Olho-pérola-de-Zugmayer	106
<i>Benthocometes robustus</i>	No English common name	Abadejo-cometa	157
<i>Benthodesmus elongatus</i>	Elongate frostfish	Espada-de-má-água-comprido	164
<i>Benthodesmus simonyi</i>	Simony's frostfish	Espada-de-má-água	165
<i>Benthodesmus tenuis</i>	Slender frostfish	Espada-de-má-água-elegante	165
<i>Benthosema glaciale</i>	Glacier lantern fish	Peixe-lanterna-glacial	113
<i>Benthosema suborbitale</i>	Smallfin lanternfish	Peixe-lanterna-corcunda	114
<i>Beryx decadactylus</i>	Alfonsino	Imperador	149
<i>Beryx splendens</i>	Splendid alfonsino	Imperador-de-costa-estreita	149
<i>Blennius ocellaris</i>	Butterfly blenny	Marachomba-borboleta	207
<i>Bodianus scrofa</i>	Barred hogfish	Bodião-dente-de-cão	280
<i>Bodianus speciosus</i>	Blackbar hogfish	Bodião-cão	281
<i>Bolinichthys indicus</i>	Smoothcheek lanternfish	Peixe-lanterna-do-índico	114
<i>Bolinichthys longipes</i>	Popeye lampfish	Peixe-lanterna-barbatana-comprida	114
<i>Bolinichthys photothorax</i>	Spurcheek lanternfish	Peixe-lanterna-esporão	114
<i>Bolinichthys pyrsobolus</i>	Fiery lanternfish	Peixe-lanterna-arco-íris	114
<i>Bolinichthys supralateralis</i>	Stubby lanternfish	Peixe-lanterna-atarracado	114
<i>Bonapartia pedaliota</i>	Longray fangjaw	Boqueirão-de-raio-longo	86
<i>Boops boops</i>	Bogue	Boga-do-mar	263
<i>Borostomias antarcticus</i>	Snaggletooth	Peixe-dragão-do-Antártico	96
<i>Borostomias elucens</i>	No English common name	Peixe-dragão-brilhante	96
<i>Borostomias mononema</i>	Sickle snaggletooth	Peixe-dragão-abutre	96
<i>Bothus podas</i>	Wide-eyed flounder	Carta-de-olhos-grandes	190
<i>Brachydeuterus auritus</i>	Bigeye grunt	Roncador-colo-colo	262
<i>Brama brama</i>	Atlantic pomfret	Xaputa	259
<i>Bregmaceros atlanticus</i>	Antenna codlet	Peixe-antena-do-Atlântico	131
<i>Brosme brosme</i>	Tusk	Bolota	144
<i>Brotulotaenia brevicauda</i>	No English common name	Abadejo-de-cauda-curta	157
<i>Brotulotaenia crassa</i>	Blue cusk eel	Abadejo-grosso	158
<i>Brotulotaenia nigra</i>	Dark cusk	Abadejo-negro	158
<i>Buenia affinis</i>	De Buen's goby	Caboz-de-De Buen	178
<i>Buenia jeffreysi</i>	Jeffrey's goby	Caboz-de-Jeffrey	179
<i>Bufockeratias wedli</i>	No English common name	Peixe-farol-de-Wedl	216
<i>Buglossidium luteum</i>	Solenette	Língua-de-gato	192
<i>Callanthias ruber</i>	Parrot seaperch	Canarinho-do-mar	247
<i>Callionymus lyra</i>	Dragonet	Peixe-pau-lira	293
<i>Callionymus maculatus</i>	Spotted dragonet	Peixe-pau-malhado	293
<i>Callionymus pusillus</i>	Sailfin dragonet	Peixe-pau-rabudo	294
<i>Callionymus reticulatus</i>	Reticulated dragonet	Peixe-pau-listado	294
<i>Callionymus risso</i>	Risso's dragonet	Peixe-pau-pintado	294
<i>Campogramma glayco</i>	Vadigo	Xareu-palheta	250
<i>Canthidermis maculata</i>	Rough triggerfish	Cangulo-áspero	228
<i>Canthidermis sufflamen</i>	Ocean triggerfish	Cangulo-oceânico	228
<i>Canthigaster rostrata</i>	Sharpnose puffer	Peixe-bola-das-Caraíbas	224
<i>Canthigaster capistrata</i>	Macaronesian sharpnose-puffer	Peixe-bola-da-Macaronésia	224
<i>Capros aper</i>	Boarfish	Pim-pim	297
<i>Caranx cryos</i>	Blue runner	Xareu-azul	251
<i>Caranx hippos</i>	Crevalle jack	Xareu-macoa	251
<i>Caranx latus</i>	Horse-eye jack	Xareu-de-olho-papudo	251
<i>Caranx lugubris</i>	Black jack	Xareu-enforcado	251
<i>Caranx rhonchus</i>	False scad	Xareu-amarelo	251

Scientific name	English common name	Portuguese common name	page
<i>Caranx ruber</i>	Bar jack	Xareu-carvoeiro	252
<i>Caranx sexfasciatus</i>	Bigeye trevally	Charéu-de-natura	252
<i>Carapus acus</i>	Pearl fish	Peixe-pérola-cobrelo	159
<i>Carcharhinus altimus</i>	Bignose shark	Tubarão-baboso	24
<i>Carcharhinus brachyurus</i>	Copper shark	Tubarão-cobre	24
<i>Carcharhinus brevipinna</i>	Graceful shark	Tubarão-tecelão	25
<i>Carcharhinus falciformis</i>	Silky shark	Tubarão-luzidio	25
<i>Carcharhinus galapagensis</i>	Galapagos shark	Tubarão-dos-Galápagos	25
<i>Carcharhinus isodon</i>	Finetooth shark	Tubarão-dente-liso	25
<i>Carcharhinus leucas</i>	Pigeye shark	Tubarão-buldogue	26
<i>Carcharhinus limbatus</i>	Blacktip reef shark	Tubarão-de-pontas-negras	26
<i>Carcharhinus longimanus</i>	Oceanic whitetip shark	Tubarão-de-pontas-brancas	26
<i>Carcharhinus obscurus</i>	Dusky shark	Tubarão-faqueta	27
<i>Carcharhinus plumbeus</i>	Sandbar shark	Tubarão-corre-costa	27
<i>Carcharhinus signatus</i>	Night shark	Tubarão-da-noite	27
<i>Carcharias taurus</i>	Smalltooth sand tiger shark	Tubarão-toiro	17
<i>Carcharodon carcharias</i>	Great white shark	Tubarão-branco	18
<i>Caristius fasciatus</i>	Greenland manefish	Peixe-crina-da-Gronelândia	260
<i>Caristius macropus</i>	Manefish	Peixe-crina-barbatana-comprida	260
<i>Cataetyx allenii</i>	No English common name	Brótula-de-Allen	160
<i>Cataetyx laticeps</i>	No English common name	Brótula-de-cabeça-chata	160
<i>Caulophryne jordani</i>	Fanfin angler	Peixe-pescador-de-Jordan	214
<i>Caulophryne polynema</i>	No English common name	Peixe-pescador-da-Madeira	215
<i>Centracanthus cirrus</i>	Curled picarel	Boqueirão	264
<i>Centrobranchus andreae</i>	Andre's lanternfish	Peixe-lanterna-de-Andréas	115
<i>Centrobranchus choerocephalus</i>	No English common name	Peixe-lanterna-de-focinho-comprido	115
<i>Centrobranchus nigroocellatus</i>	Roundnose lanternfish	Peixe-lanterna-de-focinho-redondo	115
<i>Centrodraco acanthopoma</i>	Atlantic deepwater dragonet	Peixe-pau-da-fundura-do-Atlântico	295
<i>Centrolabrus exoletus</i>	Rock cook	Bodião-de-boca-pequena	281
<i>Centrolophus niger</i>	Rudderfish	Liro-preto	171
<i>Centrophorus cf. zeehaani</i>	Little gulper shark	Quelme-do-Zeehaan	37
<i>Centrophorus granulosus</i>	Gulper shark	Barroso	35
<i>Centrophorus lusitanicus</i>	Lowfin gulper shark	Tubarão-lusitano	35
<i>Centrophorus machiquensis</i>	No English common name	Quelmo	36
<i>Centrophorus squamosus</i>	Leafscale gulper shark	Lixa	36
<i>Centrophorus uyato</i>	Little gulper shark	Quelme	36
<i>Centroprhyne spinulosa</i>	Horned lantern fish	Peixe-farol-córneo	219
<i>Centroscyllium fabricii</i>	Black dogfish	Lixinha-negra	30
<i>Centroscymnus coelolepis</i>	Portuguese dogfish	Carocho	32
<i>Centroscymnus owstonii</i>	Roughskin dogfish	Xara-preta-de-natura	32
<i>Centroselachus crepidater</i>	Roughskin dogfish	Sapata-preta	33
<i>Cephalopholis nigri</i>	Niger hind	Garoupa-do-Níger	242
<i>Cepola macrophthalmia</i>	Red bandfish	Suspensório	278
<i>Ceratias holboelli</i>	Kroyer's deep-sea angler fish	Peixe-farol-de-Holbøll	219
<i>Ceratoscopelus maderensis</i>	Madeira lantern fish	Peixe-lanterna-da-Madeira	115
<i>Ceratoscopelus townsendi</i>	Dogtooth lampfish	Peixe-lanterna-de-Townsend	115
<i>Ceratoscopelus warmingii</i>	Warming's lanternfish	Peixe-lanterna-de-Warming	115
<i>Cetichthys indagator</i>	No English common name	Peixe-baleia-do-Galathea	153
<i>Cetomimus gilli</i>	Gill's flabby whalefish	Peixe-baleia-de-Gill	153
<i>Cetomimus hempeli</i>	Whalefish	Peixe-baleia-de-Hempel	154
<i>Cetonurus globiceps</i>	Globehead grenadier	Lagartixa-do-mar-de-cabeça-redonda	132
<i>Cetorhinus maximus</i>	Basking shark	Tubarão-frade	19
<i>Cetostoma regani</i>	Pink flabby whalefish	Peixe-baleia-rosa	154
<i>Chaenophryne draco</i>	Smooth dreamer	Peixe-farol-dragão	216
<i>Chaenophryne longiceps</i>	Can-opener smoothdream	Peixe-farol-cabeça-comprida	216

Scientific name	English common name	Portuguese common name	page
<i>Chaenophryne ramifera</i>	No English common name	Peixe-farol-ramudo	216
<i>Chaetodon ocellatus</i>	Spotfin butterflyfish	Borboleta-de-risca-amarela	278
<i>Chaetodon sedentarius</i>	Reef butterflyfish	Borboleta-do-recife	278
<i>Chascanopsetta lugubris</i>	Pelican flounder	Carta-pelícano	190
<i>Chauliodus danae</i>	Dana viperfish	Peixe-dragão-do-Dana	96
<i>Chauliodus sloani</i>	Sloane's viperfish	Peixe-dragão-de-Sloane	97
<i>Chaunacops coloratus</i>	No English common name	Peixe-sapo-colorido	214
<i>Chaunacops roseus</i>	No English common name	Peixe-sapo-rosado	214
<i>Chaunax pictus</i>	Pink frogmouth	Peixe-sapo-mole	214
<i>Chaunax suttkusi</i>	No English common name	Peixe-sapo-de-Suttkus	214
<i>Cheilopogon cyanopterus</i>	Margined flyingfish	Peixe-voador-barbudo	200
<i>Cheilopogon exsiliens</i>	Bandwing flyingfish	Peixe-voador-de-asas-listadas	200
<i>Cheilopogon furcatus</i>	Spotfin flyingfish	Peixe-voador-de-barbatana-manchada	200
<i>Cheilopogon heterurus</i>	Mediterranean flyingfish	Peixe-voador-do-Mediterrâneo	200
<i>Cheilopogon pinnatibarbatus</i>	Bennett's flyingfish	Peixe-voador-de-Bennett	200
<i>Chelidonichthys cuculus</i>	Red gurnard	Cabra-vermelha	234
<i>Chelidonichthys lastoviza</i>	Streaked gurnard	Cabra-riscada	234
<i>Chelidonichthys lucerna</i>	Tub gurnard	Cabra-cabaço	235
<i>Chelidonichthys obscurus</i>	Longfin gurnard	Cabra-de-bandeira	235
<i>Chelon labrosus</i>	Thicklip grey mullet	Tainha-liça	202
<i>Chelon ramada</i>	Thinlip grey mullet	Tainha-fataça	202
<i>Chelon saliens</i>	Leaping mullet	Tainha-de-salto	203
<i>Chiasmodon niger</i>	Black swalower	Peixe-glutão-negro	289
<i>Chilomycterus reticulatus</i>	Pacific burrfish	Porco-espinho-de-pintas	222
<i>Chilomycterus schoepfii</i>	Striped burrfish	Porco-espinho-listado	223
<i>Chilomycterus spinosus mauretanicus</i>	Guinean burrfish	Porco-espinho-da-Guiné	223
<i>Chilomycterus spinosus spinosus</i>	No English common name	Porco-espinho-americano	223
<i>Chimaera monstrosa</i>	Rabbit fish	Quimera-comum	53
<i>Chimaera opalescens</i>	Opal chimaera	Quimera-opala	53
<i>Chiostomias pliopterus</i>	Scaleless dragonfish	Peixe-dragão-de-bolbo-papilar	97
<i>Chlamydoselachus anguineus</i>	Frilled shark	Tubarão-cobra	16
<i>Chlopsis bicolor</i>	Bicoloured false moray	Congrinho-bicolor	63
<i>Chlopsis dentatus</i>	Mottled false moray	Congrinho-malhado	63
<i>Chlorophthalmus agassizii</i>	Shortnose greeneye	Olho-verde	105
<i>Chromis chromis</i>	Damsel fish	Castanheta-báia	278
<i>Chromis limbata</i>	Azores chromis	Castanheta-da-Madeira	279
<i>Chromogobius britoi</i>	Brito's goby	Caboz-de-Brito	179
<i>Ciliata mustela</i>	Fivebeard rockling	Laibeque-de-cinco-barbilhos	144
<i>Citharus linguatula</i>	Spotted flounder	Carta-de-bico	186
<i>Clinitrichus argentatus</i>	Cline	Marachomba-prateada	207
<i>Clupea harengus</i>	Atlantic herring	Arenque-do-Atlântico	72
<i>Coccarella atlantica</i>	Atlantic sabretooth	Dente-de-sabre-do-Atlântico	112
<i>Coelorinchus caelorhincus</i>	Hollowsnout grenadier	Lagartixa-do-mar-de-focinho-côncavo	132
<i>Coelorinchus labiatus</i>	Spearsnouted grenadier	Lagartixa-do-mar-de-focinho-comprido	132
<i>Coelorinchus occa</i>	Swordsnout grenadier	Lagartixa-do-mar-de-focinho-curto	133
<i>Coelorinchus parallelus</i>	Spiny grenadier	Lagartixa-do-mar-espinhosa	133
<i>Conger conger</i>	European conger	Congro	66
<i>Conger tripiceps</i>	Manytooth conger	Congro-dentudo	66
<i>Conocara fiontii</i>	Fiontii's smooth-head	Celindra-de-Fiontii	76
<i>Conocara macropterum</i>	Longfin smooth-head	Celindra-de-barbatana-comprida	76
<i>Conocara murrayi</i>	Murray's smooth-head	Celindra-de-Murray	77
<i>Conocara salmonicum</i>	Salmon smooth-head	Celindra-salmonada	77
<i>Conocara werneri</i>	Werner's smooth-head	Celindra-de-Werner	77
<i>Cookeolus japonicus</i>	Longfinned bullseye	Fura-vasos-alfonsim	247
<i>Coris julis</i>	Rainbow wrasse	Judia	281

Scientific name	English common name	Portuguese common name	page
<i>Corniger spinosus</i>	Spinycheek soldierfish	Peixe-soldado-espinhoso	156
<i>Coryphaena equiselis</i>	Pompano dolphinfish	Doirado-pampo	258
<i>Coryphaena hippurus</i>	Common dolphinfish	Doirado	258
<i>Coryphaenoides armatus</i>	Abyssal grenadier	Lagartixa-do-mar-abissal	133
<i>Coryphaenoides brevibarbis</i>	Shortbeard grenadier	Lagartixa-do-mar-de-barbilho-curto	133
<i>Coryphaenoides carapinus</i>	Carapine grenadier	Lagartixa-do-mar-carapina	133
<i>Coryphaenoides guentheri</i>	Günther's grenadier	Lagartixa-do-mar-de-Günther	133
<i>Coryphaenoides leptolepis</i>	Ghostly grenadier	Lagartixa-do-mar-pálida	134
<i>Coryphaenoides mediterraneus</i>	Mediterranean grenadier	Lagartixa-do-mar-do-Mediterrâneo	134
<i>Coryphaenoides profundiculus</i>	Deepwater grenadier	Lagartixa-do-mar-da-fundura	134
<i>Coryphaenoides ruddis</i>	Madeiran grenadier	Lagartixa-do-mar-cabeçuda	134
<i>Coryphaenoides rupestris</i>	Roundnose grenadier	Lagartixa-da-rocha	134
<i>Coryphaenoides thelestomus</i>	Roughlip grenadier	Lagartixa-do-mar-da-Madeira	134
<i>Coryphaenoides zaniophorus</i>	Thickbeard grenadier	Lagartixa-do-mar-de-barbilho-grosso	135
<i>Coryphoblennius galerita</i>	Montagu's blenny	Marachomba-das-rochas	207
<i>Cottunculus thomsonii</i>	Pallid sculpin	Escorpião-pálido	237
<i>Cryptopsaras couesii</i>	Triplewart seadevil	Peixe-farol-de-Coues	219
<i>Crystallogobius linearis</i>	Crystal goby	Caboz-de-crystal	179
<i>Ctenochirichthys longimanus</i>	No English common name	Peixe-farol-barbatana-comprida	217
<i>Ctenolabrus rupestris</i>	Goldsinny-wrasse	Bodião-rupestre	282
<i>Ctenoscopelus phengodes</i>	Bright lanternfish	Peixe-lanterna-brilhante	116
<i>Cubiceps gracilis</i>	Driftfish	Tirone	172
<i>Cyclopterus lumpus</i>	Lumpfish	Peixe-lapa	237
<i>Cyclothona acclinidens</i>	Benttooth bristlemouth	Boqueirão-de-dentes-curvos	86
<i>Cyclothona alba</i>	Pale bristlemouth	Boqueirão-branco	86
<i>Cyclothona braueri</i>	Garrick	Boqueirão-de-Brauer	86
<i>Cyclothona livida</i>	Bristlemouth	Boqueirão-lívido	87
<i>Cyclothona microdon</i>	Veiled anglemouth	Boqueirão-de-dentes-pequenos	87
<i>Cyclothona obscura</i>	Hidden bristlemouth	Boqueirão-sombrio	87
<i>Cyclothona pallida</i>	Tan bristlemouth	Boqueirão-bicolor	87
<i>Cyclothona parapallida</i>	Shadow bristlemouth	Boqueirão-sombreado	87
<i>Cyclothona pseudopallida</i>	Slender bristlemouth	Boqueirão-delgado	87
<i>Cyclothona signata</i>	Showy bristlemouth	Boqueirão-vistoso	87
<i>Cyema atrum</i>	Bobtail eel	Cobra-de-leme	69
<i>Cynoglossus canariensis</i>	Canary tonguesole	Língua-de-cão-das-Canárias	196
<i>Cynoglossus lineolatus</i>	No English common name	Língua-de-cão-riscada	196
<i>Cynoponticus ferox</i>	Guinean pike conger	Congro-branco-africano	65
<i>Cynoscion regalis</i>	Squeteague	Corvinata-real	275
<i>Cyttopsis rosea</i>	Red dory	Galo-de-natura	129
<i>Dactylopterus volitans</i>	Flying gurnard	Cabrinha-de-leque	178
<i>Dagetichthys lusitanicus</i>	Portuguese sole	Língua-de-vaca	192
<i>Dalatias licha</i>	Kitefin shark	Gata	29
<i>Dalophis imberbis</i>	Armless snake eel	Cobra-imberbe	64
<i>Dasyatis pastinaca</i>	Common stingray	Uge	49
<i>Dasy scopelus selenops</i>	Wisner's lantern fish	Peixe-lanterna-de-Wisner	116
<i>Deania calcea</i>	Birdbeak dogfish	Sapata	37
<i>Deania hystricosa</i>	Rough longnose dogfish	Sapata-áspera	37
<i>Deania profundorum</i>	Longsnout dogfish	Sapata-flecha	38
<i>Decapterus macarellus</i>	Mackerel scad	Charro-olho-largo	252
<i>Decapterus punctatus</i>	Round scad	Charro-moiro	252
<i>Decapterus tabl</i>	Roughear scad	Charro-de-cauda-vermelha	252
<i>Deltentosteus collarinus</i>	Toothed goby	Caboz-dentudo	179
<i>Deltentosteus quadrimaculatus</i>	Four-spotted goby	Caboz-de-quatro-manchas	179
<i>Dentex canariensis</i>	Canary dentex	Dentão-das-Canárias	264
<i>Dentex dentex</i>	Common dentex	Capatão-legítimo	264

Scientific name	English common name	Portuguese common name	page
<i>Dentex gibbosus</i>	Pink dentex	Capatão-de-bandeira	264
<i>Dentex macrourus</i>	Large-eye dentex	Cachucho	265
<i>Dentex maroccanus</i>	Morocco dentex	Cachucho-dentão	265
<i>Derichthys serpentinus</i>	Narrownecked oceanic eel	Enguia-de-pescoco-fino	63
<i>Diaphus adenomus</i>	Firebow lanternfish	Peixe-lanterna-de-Gilbert	116
<i>Diaphus bertelseni</i>	Bertelsen's lanternfish	Peixe-lanterna-de-Bertelsen	116
<i>Diaphus brachycephalus</i>	Short-headed lantern fish	Peixe-lanterna-de-cabeça-curta	116
<i>Diaphus dumerilii</i>	Dumeril's lanternfish	Peixe-lanterna-de-Duméril	117
<i>Diaphus effulgens</i>	Headlight lanternfish	Peixe-lanterna-de-focinho-luminoso	117
<i>Diaphus holti</i>	Small lantern fish	Peixe-lanterna-de-Holt	117
<i>Diaphus lucidus</i>	Spotlight lanternfish	Peixe-lanterna-cintilante	117
<i>Diaphus luetkeni</i>	Luetken's lanternfish	Peixe-lanterna-de-Lütken	117
<i>Diaphus metopoclampus</i>	Spothead lantern fish	Peixe-lanterna-de-cabeça-manchada	117
<i>Diaphus mollis</i>	Soft lanternfish	Peixe-lanterna-mole	117
<i>Diaphus perspicillatus</i>	Flatface lanternfish	Peixe-lanterna-luneta	118
<i>Diaphus rafinesquii</i>	White-spotted lantern fish	Peixe-lanterna-de-Rafinesque	118
<i>Diaphus splendidus</i>	Horned lanternfish	Peixe-lanterna-cornudo	118
<i>Diaphus subtilis</i>	Flabby lanternfish	Peixe-lanterna-flácido	118
<i>Diaphus termophilus</i>	Taaning's lantern fish	Peixe-lanterna-termófilo	118
<i>Diaphus theta</i>	California headlightfish	Peixe-lanterna-da-California	118
<i>Dicentrarchus labrax</i>	European seabass	Robalo-legítimo	241
<i>Dicentrarchus punctatus</i>	Spotted seabass	Robalo-baila	241
<i>Dicologlossa cuneata</i>	Wedge sole	Língua	192
<i>Diodon holocanthus</i>	Pelagic porcupinefish	Peixe-ouriço-pelágico	223
<i>Diodon holocanthus</i>	Longspined porcupinefish	Peixe-ouriço-de-crista	223
<i>Diodon hystrix</i>	Spot-fin porcupinefish	Peixe-ouriço-ponteado	224
<i>Diogenichthys atlanticus</i>	Atlantic lanternfish	Peixe-lanterna-do-Atlântico	118
<i>Diogenichthys laternatus</i>	Diogenes lanternfish	Peixe-lanterna-de-Diogenes	119
<i>Diplecogaster bimaculata</i>	Two-spotted clingfish	Sugador-de-duas-manchas	205
<i>Diplecogaster pectoralis</i>	No English common name	Sugador-das-Canárias	205
<i>Diplodus annularis</i>	Annular seabream	Sargo-alcorraz	265
<i>Diplodus bellottii</i>	Senegal seabream	Sargo-do-Senegal	266
<i>Diplodus cadenati</i>	White seabream	Sargo-legítimo-do-Atlântico	266
<i>Diplodus cervinus</i>	Zebra seabream	Sargo-veado	266
<i>Diplodus fasciatus</i>	Banded seabream	Sargo-de-faixas	267
<i>Diplodus lineatus</i>	White seabream	Sargo-de-Cabo-Verde	267
<i>Diplodus puntazzo</i>	Shapsnout seabream	Sargo-bicudo	267
<i>Diplodus sargus</i>	White seabream	Sargo-legítimo-do-Mediterrâneo	267
<i>Diplodus vulgaris</i>	Common two-banded seabream	Sargo-safia	268
<i>Diplophos taenia</i>	Pacific portholefish	Boqueirão-comprido	88
<i>Diplospinus multistriatus</i>	Striped escolar	Escolar-raiado	162
<i>Dipturus batis</i>	Blue skate	Raia-oirega	43
<i>Dipturus intermedius</i>	Flapper Skate	Raia-delgada	43
<i>Dipturus nidorosiensis</i>	Norwegian skate	Raia-norueguesa	43
<i>Dipturus oxyrinchus</i>	Longnosed skate	Raia-bicuda	43
<i>Dipturus sp. Stehmann</i>	Norwegian skate	Raia-norueguesa	44
<i>Diretmichthys parini</i>	Parin's spinyfish	Peixe-enfadado-de-Parin	155
<i>Diretmoides pauciradiatus</i>	Silver spinyfin	Peixe-enfadado-barbatana-comprida	155
<i>Diretmus argenteus</i>	Silver spinyfin	Peixe-enfadado-prateado	156
<i>Discoverichthys praecox</i>	No English common name	Peixe-tripé-temporão	106
<i>Ditropichthys storeri</i>	No English common name	Peixe-baleia-de-Storer	154
<i>Dolicholagus longirostris</i>	Longsnout blacksmelt	Argentina-da-fundura-de-focinho-comprido	84
<i>Dolichopteryx longipes</i>	Brown-snout spookfish	Peixe-fantasma-de-barbatanas-longas	84
<i>Dolopichthys allector</i>	No English common name	Peixe-farol-cornudo	217
<i>Dolopichthys danae</i>	No English common name	Peixe-farol-do-Dana	217

Scientific name	English common name	Portuguese common name	page
<i>Dolopichthys jubatus</i>	No English common name	Peixe-farol-de-crista	217
<i>Dolopichthys karsteni</i>	No English common name	Peixe-farol-de-Karsten	217
<i>Dolopichthys longicornis</i>	No English common name	Peixe-farol-diabo	217
<i>Dolopichthys pullatus</i>	Lobed dreamer	Peixe-farol-lobado	217
<i>Dysalotus alcocki</i>	No English common name	Peixe-glutão-de-Alcock	289
<i>Dysalotus oligoscolus</i>	No English common name	Peixe-glutão	289
<i>Dysomma brevirostre</i>	Batnose eel	Enguia-de-focinho-bolboso	58
<i>Dysommina proboscideus</i>	No English common name	Enguia-de-tromba-de-elefante	58
<i>Echelus myrus</i>	Painted eel	Cobra-de-orelhas	64
<i>Echeneis naucrates</i>	Live sharksucker	Pegador-listado	249
<i>Echiichthys vipera</i>	Lesser weever	Peixe-aranha-menor	291
<i>Echinomacrurus mollis</i>	No English common name	Lagartixa-do-mar-macia	135
<i>Echinorhinus brucus</i>	Bramble shark	Tubarão-prego	39
<i>Echiodon dentatus</i>	No English common name	Peixe-pérola-dentado	159
<i>Echiodon drummondii</i>	No English common name	Peixe-pérola-de-Drummond	160
<i>Echiostoma barbatum</i>	Threadfin dragonfish	Peixe-dragão-barbatana-filamentosa	97
<i>Ectreposebastes imus</i>	Midwater scorpionfish	Rascasso-da-fundura	229
<i>Einara edentula</i>	Toothless smooth-head	Celindra-desdentada	77
<i>Einara macrolepis</i>	Loosescale smooth-head	Celindra-de-escama-caduca	77
<i>Elagatis bipinnulata</i>	Rainbow runner	Fogueteiro-arco-íris	253
<i>Electrona risso</i>	Electric lantern fish	Peixe-lanterna-faísca	119
<i>Enchelycore anatina</i>	Fangtooth moray	Moreia-serpente	60
<i>Engraulis encrasiculus</i>	European anchovy	Biqueirão	74
<i>Entelurus aequoreus</i>	Snake pipefish	Marinha-amarela	175
<i>Ephippion guttifer</i>	Prickly puffer	Peixe-bola-de-pintas	224
<i>Epigonus constanciae</i>	Constance deepwater cardinalfish	Olhudo-de-espinho	239
<i>Epigonus denticulatus</i>	Pencil cardinal	Olhudo-denteado	239
<i>Epigonus telescopus</i>	Black cardinal fish	Olhudo	240
<i>Epinephelus adscensionis</i>	Rock hind	Garoupa-pintada-da-rocha	243
<i>Epinephelus aeneus</i>	White grouper	Garoupa-legítima	243
<i>Epinephelus caninus</i>	Dogtooth grouper	Mero-gigante	243
<i>Epinephelus costae</i>	Goldblotch grouper	Garoupa-dourada	243
<i>Epinephelus goreensis</i>	Dungat grouper	Mero-amarelo	243
<i>Epinephelus guaza</i>	<i>Nomen dubium</i> in Mycteroperca	<i>Nomen dubium</i> in Mycteroperca	243
<i>Epinephelus marginatus</i>	Dusky grouper	Mero-legítimo	244
<i>Eretmophorus kleinenbergi</i>	No English common name	Peixe-remo-de-Kleinenberg	139
<i>Etelis carbunculus</i>	Deep-water red snapper	Vermelho-salmão	261
<i>Etelis oculatus</i>	Queen snapper	Rosa-salmão	261
<i>Etmopterus princeps</i>	Great lanternshark	Lixinha-da-fundura-grada	31
<i>Etmopterus pusillus</i>	Smooth lanternshark	Xarinha-preta	31
<i>Etmopterus spinax</i>	Velvet belly	Lixinha-da-fundura	31
<i>Eumicrotremus spinosus</i>	Atlantic spiny lump sucker	Peixe-lapa-espinhoso	237
<i>Eurypharynx pelecanoides</i>	Pelican eel	Peixe-pelícano	70
<i>Eustomias aciniosus</i>	No English common name	Peixe-dragão-de-barbilo-acinoso	97
<i>Eustomias braueri</i>	No English common name	Peixe-dragão-lusitano	97
<i>Eustomias contiguus</i>	No English common name	Peixe-dragão-biboloso	97
<i>Eustomias drechseli</i>	No English common name	Peixe-dragão-de-Drechsel	98
<i>Eustomias dubius</i>	No English common name	Peixe-dragão-ambíguo	98
<i>Eustomias enbarbatus</i>	Barbate dragonfish	Peixe-dragão-barbado	98
<i>Eustomias filifer</i>	No English common name	Peixe-dragão-de-barbilo-filifero	98
<i>Eustomias fissibarbis</i>	No English common name	Peixe-dragão-de-barbilo-bifurcado	98
<i>Eustomias furcifer</i>	Scaleless dragonfish	Peixe-dragão-de-furfur	98
<i>Eustomias lipochirus</i>	No English common name	Peixe-dragão-de-Santa Cruz	98
<i>Eustomias longibarba</i>	No English common name	Peixe-dragão-de-barbilo-crescido	98
<i>Eustomias macronema</i>	Bigbarb dragonfish	Peixe-dragão-de-barbilo-grande	98

Scientific name	English common name	Portuguese common name	page
<i>Eustomias macrurus</i>	Yellowstem dragonfish	Peixe-dragão-de-barbilo-amarelo	99
<i>Eustomias melanostigma</i>	No English common name	Peixe-dragão-de-mancha-negra	99
<i>Eustomias monodactylus</i>	No English common name	Peixe-dragão-de-bolbo-monodáctilo	99
<i>Eustomias obscurus</i>	Scaleless dragonfish	Peixe-dragão-obscurio	99
<i>Eustomias parri</i>	No English common name	Peixe-dragão-de-Parr	99
<i>Eustomias radicifilis</i>	No English common name	Peixe-dragão-de-barbilo-ramificado	99
<i>Eustomias satterleei</i>	Twinray dragonfish	Peixe-dragão-de-Satterlee	99
<i>Eustomias schiffi</i>	No English common name	Peixe-dragão-de-Schiff	99
<i>Eustomias schmidti</i>	Schmidt's dragonfish	Peixe-dragão-de-Schmidt	99
<i>Eustomias simplex</i>	No English common name	Peixe-dragão-de-bolbo-simples	99
<i>Eustomias tenisoni</i>	No English common name	Peixe-dragão-de-Tenison	100
<i>Eustomias tetranema</i>	No English common name	Peixe-dragão-da-Madeira	100
<i>Eutaeniophorus festivus</i>	Festive ribbonfish	Peixe-serpentina	154
<i>Euthynnus alletteratus</i>	Little tunny	Merma	167
<i>Eutrigla gurnardus</i>	Grey gurnard	Cabra-morena	235
<i>Evermannella balbo</i>	Balbo sabretooth	Dente-de-sabre-de-Balbo	112
<i>Evermannella melanoderma</i>	Indian sabretooth	Dente-de-sabre-ponteado	112
<i>Evynnis ehrenbergii</i>	Bluespotted seabream	Pargo-ruço-de-Ebrenberg	269
<i>Exocoetus obtusirostris</i>	Oceanic two-wing flyingfish	Andorinha-do-mar-oceânica	201
<i>Exocoetus volitans</i>	Tropical two-wing flyingfish	Andorinha-do-mar-tropical	201
<i>Facciolella oxyrhynchus</i>	Facciola's sorcerer	Cobra-cabeça-de-pato	65
<i>Fistularia petimba</i>	Red cornetfish	Corneta-rosada	174
<i>Fistularia tabacaria</i>	Cornetfish	Corneta-malhada	174
<i>Flagellosomias boureei</i>	Longbarb dragonfish	Peixe-dragão-dos-Açores	100
<i>Fowlerichthys radiosus</i>	Singlespot frogfish	Peixe-sapo-uniocelado	213
<i>Fowlerichthys senegalensis</i>	Senegalese frogfish	Peixe-sapo-do-Senegal	213
<i>Gadella imberbis</i>	Beardless codling	Abrótea-imberbe	139
<i>Gadella maraldi</i>	Gadella	Abrótea-de-natura	139
<i>Gadiculus argenteus</i>	Silvery pout	Badejinho	142
<i>Gadomus arcuatus</i>	Doublethread grenadier	Lagartixa-do-mar-filamentosa	135
<i>Gadomus dispar</i>	No English common name	Lagartixa-do-mar-ímpar	135
<i>Gadomus longifilis</i>	Threadfin grenadier	Lagartixa-do-mar-de-bandeira	135
<i>Gadus morhua</i>	Atlantic cod	Bacalhau-do-Atlântico	142
<i>Gaidropsarus granti</i>	Azores rockling	Laibequer-dos-Açores	144
<i>Gaidropsarus macrophthalmus</i>	Bigeye rockling	Laibequer-do-Atlântico	145
<i>Gaidropsarus mauli</i>	No English common name	Laibequer-de-Maul	145
<i>Gaidropsarus mediterraneus</i>	Shore rockling	Laibequer-do-Mediterrâneo	145
<i>Gaidropsarus vulgaris</i>	Three-bearded rockling	Laibequer-de-três-barbilhos	145
<i>Galeichthys feliceps</i>	White barbel	Bagre-barba-branca	82
<i>Galeocerdo cuvier</i>	Tiger shark	Tubarão-tigre	27
<i>Galeorhinus galeus</i>	Tope shark	Perna-de-moça	23
<i>Galeus atlanticus</i>	Atlantic sawtail catshark	Leitão-do-Atlântico	21
<i>Galeus melastomus</i>	Blackmouth catshark	Leitão	21
<i>Galeus murinus</i>	Mouse catshark	Leitão-da-Islândia	21
<i>Gasterosteus aculeatus</i>	Three-spined stickleback	Esgana-gata	297
<i>Gempylus serpens</i>	Snake mackerel	Escolar-serpente	162
<i>Gephyroberyx darwinii</i>	Darwin's slimehead	Olho-de-vidro-imperatriz	154
<i>Gigantactis ios</i>	No English common name	Peixe-chicote-das-Canárias	220
<i>Gigantactis longicirra</i>	No English common name	Peixe-chicote-filamentoso	220
<i>Gigantactis vanhoeffeni</i>	No English common name	Peixe-chicote-de-Van Höffen	220
<i>Glaucostegus cemiculus</i>	Blackchin guitarfish	Viola-barba-negra	42
<i>Glossanodon leioglossus</i>	Smalltoothed argentine	Argentina-de-dentes-pequenos	82
<i>Glyptocephalus cynoglossus</i>	Witch flounder	Solhão	191
<i>Gnatholepis thompsoni</i>	Goldspot goby	Caboz-de-pinta-dourada	179
<i>Gnathophis codoniphorus</i>	No English common name	Congro-dos-Açores	67

Scientific name	English common name	Portuguese common name	page
<i>Gnathophis mystax</i>	Thinlip conger	Coreano	67
<i>Gobius ater</i>	Bellotti's goby	Caboz-de-Bellotti	179
<i>Gobius auratus</i>	Golden goby	Caboz-dourado	180
<i>Gobius bucchichi</i>	Buccich's goby	Caboz-mosqueado	180
<i>Gobius cobitis</i>	Giant goby	Caboz-cabeçudo	180
<i>Gobius couchi</i>	Couch's goby	Caboz-de-Couch	180
<i>Gobius cruentatus</i>	Red-mouthed goby	Caboz-de-boca-vermelha	180
<i>Gobius gasteveni</i>	Steven's goby	Caboz-de-escama	180
<i>Gobius geniporus</i>	Slender goby	Caboz-de-bochechas porosas	180
<i>Gobius niger</i>	Black goby	Caboz-negro	181
<i>Gobius ophiocephalus</i>	Grass goby	Caboz-cabeça-de-cobra	181
<i>Gobius paganellus</i>	Rock goby	Caboz-da-rocha	181
<i>Gobius roulei</i>	Roule's goby	Caboz-de-Roule	181
<i>Gobius xanthocephalus</i>	Golden goby	Caboz-de-cabeça-dourada	182
<i>Gobiusculus flavescens</i>	Two-spotted goby	Caboz-de-duas-manchas	182
<i>Gonichthysocco</i>	Cocco's lanternfish	Peixe-laterna-de-Cocco	119
<i>Gonostoma atlanticum</i>	Atlantic fangjaw	Boqueirão-do-Atlântico	88
<i>Gonostoma denudatum</i>	Bristlemouth	Boqueirão-do-Mediterrâneo	88
<i>Grammatostomias circularis</i>	No English common name	Peixe-dragão-de-mancha-redonda	100
<i>Grammatostomias dentatus</i>	No English common name	Peixe-dragão-colmilhoso	100
<i>Grammatostomias flagellibarba</i>	No English common name	Peixe-dragão-de-barbilho-filamentoso	100
<i>Grammicolepis brachiusculus</i>	Thorny tinselfish	Peixe-prata-espinhos	130
<i>Grammonus ater</i>	No English common name	Brótula-negra	161
<i>Guentherus altivelia</i>	Jellynose	Peixe-de-vela-alta	104
<i>Guttigadus latifrons</i>	No English common name	Abrótea-cabeçuda	140
<i>Gymnammodytes cicerelus</i>	Mediterranean sand eel	Galeota-da-areia	291
<i>Gymnammodytes semisquamatus</i>	Smooth sandeel	Galeota-frachão	291
<i>Gymnothorax afer</i>	Dark moray	Moreão-escuro	60
<i>Gymnothorax bacalladoi</i>	Canary moray	Moreia-das-Canárias	60
<i>Gymnothorax maderensis</i>	Sharktooth moray	Moreia-da-Madeira	61
<i>Gymnothorax miliaris</i>	Goldentail moray	Moreia-dourada	61
<i>Gymnothorax polygonius</i>	Polygon moray	Moreia-poligonal	61
<i>Gymnothorax unicolor</i>	Brown moray	Moreia-castanha	61
<i>Gymnothorax vicinus</i>	Purplemouth moray	Moreão-amarelo	61
<i>Gymnura altavela</i>	Spiny butterfly ray	Uge-manta	50
<i>Gymnura hirundo</i>	Madeira butterfly ray	Uge-da-Madeira	50
<i>Gyrinomimus myersi</i>	Myers' flabby whalefish	Peixe-baleia-de-Myers	154
<i>Halargyreus johnsonii</i>	Slender codling	Abrótea-delgada	140
<i>Halobatrachus didactylus</i>	Lusitanian toadfish	Charroco	162
<i>Halosauropsis macrochir</i>	Abyssal halosaur	Peixe-lagarto-abissal	56
<i>Halosaurus guentheri</i>	Günther's halosaur	Peixe-lagarto-de-Guenther	56
<i>Halosaurus jonhsonianus</i>	Johnson's halosaur	Peixe-lagarto-de-Johnson	56
<i>Halosaurus ovenii</i>	Oven's halosaur	Peixe-lagarto-de-Owen	57
<i>Haplophryne mollis</i>	Soft leafvent angler	Peixe-sestro-macio	220
<i>Harengula clupeola</i>	False herring	Sardinha-escamuda	72
<i>Harriotta raleighana</i>	Pacific longnose chimaera	Peixe-rato-de-Raleigh	54
<i>Helicolenus dactylopterus</i>	Blackbelly rosefish	Cantarilho-legítimo	228
<i>Hemiramphus balao</i>	Balao halfbeak	Meia-agulha-preta	200
<i>Heptranchias perlo</i>	Sharpnose sevengill shark	Tubarão-de-sete-guelras	15
<i>Herwigia krefftii</i>	Krefft's smooth-head	Celindra-de-Krefft	77
<i>Heteroconger longissimus</i>	Brown garden eel	Enguia-de-jardim	67
<i>Heterophotus ophistoma</i>	Wingfin snaggletooth	Peixe-dragão-de-boca-de-serpente	100
<i>Heteropriacanthus cruentatus</i>	Glasseseye	Fura-vasos-da-rocha	247
<i>Heteropriacanthus fulgens</i>	Bright glasseseye	Fura-vasos-da-rocha-brilhante	248
<i>Hexanchus griseus</i>	Bluntnose sixgill shark	Tubarão-albafar	16

Scientific name	English common name	Portuguese common name	page
<i>Hexanchus nakamurai</i>	Bluntnose sixgill shark	Tubarão-albafar-de-pontas-brancas	16
<i>Himantolophus albinares</i>	No English common name	Peixe-farol-da-Madeira	215
<i>Himantolophus brevirostris</i>	No English common name	Peixe-farol-de-focinho-curto	215
<i>Himantolophus compressus</i>	No English common name	Peixe-farol-sapo	216
<i>Himantolophus groenlandicus</i>	Atlantic footballfish	Peixe-farol-do-Atlântico	216
<i>Himantolophus mauli</i>	No English common name	Peixe-farol-de-Maul	216
<i>Hippocampus erectus</i>	Lined Seahorse	Cavalo-marinho-estriado	175
<i>Hippocampus guttulatus</i>	Long-snouted seahorse	Cavalo-marinho-de-focinho-comprido	175
<i>Hippocampus hippocampus</i>	Short snouted seahorse	Cavalo-marinho-de-focinho-curto	176
<i>Hirundichthys rondeletii</i>	Black wing flyingfish	Peixe-voador-de-asa-negra	201
<i>Hirundichthys speculiger</i>	Mirrorwing flyingfish	Peixe-voador-de-asa-espelhada	201
<i>Hirundichthys volador</i>	Atlantic blackwing flyingfish	Peixe-voador-de-asa-negra-do-Atlântico	202
<i>Histiobranchus australis</i>	Southern cut-throat eel	Enguia-da-fundura-austral	58
<i>Histiobranchus bathybius</i>	Deep-water arrowtooth eel	Enguia-da-fundura	58
<i>Histrio histrio</i>	Sargassumfish	Peixe-sapo-do-sargaço	213
<i>Holcomycteronus squamosus</i>	No English common name	Abadejo-escamudo	158
<i>Holtbyrnia anomala</i>	Bighead searsid	Celindra-de-tubo-cabeçuda	80
<i>Holtbyrnia innesi</i>	Teardrop tubeshoulder	Celindra-de-tubo-de-Innes	80
<i>Holtbyrnia macrops</i>	Bigeye searsid	Celindra-de-tubo-de-olhuda	80
<i>Holtbyrnia melanocephala</i>	No English common name	Celindra-de-tubo-de-cabeça-negra	80
<i>Hoplostethus atlanticus</i>	Orange roughy	Olho-de-vidro-laranja	154
<i>Hoplostethus cadenati</i>	Black slimehead	Olho-de-vidro-preto	155
<i>Hoplostethus mediterraneus</i>	Mediterranean slimehead	Olho-de-vidro-do-Mediterrâneo	155
<i>Howella atlantica</i>	Atlantic pelagic basslet	Peixe-alado-do-Atlântico	240
<i>Howella brodiei</i>	Pelagic basslet	Peixe-alado-de-Brodie	240
<i>Howella sherborni</i>	Sherborn's pelagic bass	Peixe-alado-de-Sherborn	241
<i>Howella simplex</i>	No English common name	Peixe-alado	241
<i>Hydrolagus affinis</i>	Smalleyed rabbitfish	Ratazana-da-fundura	54
<i>Hydrolagus lusitanicus</i>	No English common name	Ratazana-lusitânica	54
<i>Hydrolagus mirabilis</i>	Large-eyed rabbitfish	Ratazana-olhuda	54
<i>Hydrolagus pallidus</i>	Pale chimaera	Ratazana-pálida	54
<i>Hygophum benoiti</i>	Benoit's lanternfish	Peixe-lanterna-de-Benoit	119
<i>Hygophum hygomii</i>	Bermuda lantern fish	Peixe-lanterna-das-Bermudas	120
<i>Hygophum macrochir</i>	Largefin lanternfish	Peixe-lanterna-de-barbatanas-grandes	120
<i>Hygophum reinhardtii</i>	Reinhardt's lanternfish	Peixe-lanterna-de-Reinhardt	120
<i>Hygophum taanungi</i>	Tåning's lanternfish	Peixe-lanterna-de-Tåning	120
<i>Hymenocephalus italicus</i>	Glasshead grenadier	Lagartixa-do-mar-prateada	135
<i>Hymenogadus gracilis</i>	Graceful grenadier	Lagartixa-do-mar-elegante	135
<i>Hyperoglyphe perciformis</i>	Barrelfish	Liro-errante	172
<i>Hyperoplus immaculatus</i>	Greater sand-eel	Galeota-maior-de-Corbin	291
<i>Hyperoplus lanceolatus</i>	Great sandeel	Galeota-maior	291
<i>Hypoleurochilus bananensis</i>	No English common name	Marachomba-banana	207
<i>Hypoleurochilus fissicornis</i>	No English common name	Marachomba-da-toca	208
<i>Hyporthodus haifensis</i>	Haifa grouper	Garoupa-de-Haifa	244
<i>Ichthyococcus ovatus</i>	Ovate lightfish	Peixe-luminoso-oval	92
<i>Idiacanthus fasciola</i>	Ribbon sawtail fish	Peixe-dragão-de-cauda-serreada	100
<i>Ilyophis blachei</i>	Blache's ooze ee	Enguia-da-fundura-de-Blache	58
<i>Ilyophis brunneus</i>	Muddy arrowtooth eel	Enguia-da-fundura-castanha	58
<i>Isistius brasiliensis</i>	Cookie cutter shark	Tubarão-luminoso	30
<i>Isistius plutodus</i>	Largetooth cookiecutter shark	Tubarão-dentão	30
<i>Istiophorus platypterus</i>	Indo-Pacific sailfish	Veleiro-do-Indo-Pacífico	185
<i>Isurus oxyrinchus</i>	Longfin mako	Tubarão-anequim	18
<i>Isurus paucus</i>	Longfin mako	Tubarão-anequim-de-gadanha	18
<i>Kajikia albida</i>	Atlantic white marlin	Espadim-branco-do-Atlântico	185
<i>Kali indica</i>	No English common name	Peixe-glutão-do-Índico	290

Scientific name	English common name	Portuguese common name	page
<i>Kali kerberti</i>	No English common name	Peixe-glutão-de-Kerbert	290
<i>Kali macrodon</i>	No English common name	Peixe-glutão-dentudo	290
<i>Kali macrura</i>	No English common name	Peixe-glutão-rabudo	290
<i>Kali parri</i>	No English common name	Peixe-glutão-de-Parr	290
<i>Katsuwonus pelamis</i>	Skipjack tuna	Gaiado	167
<i>Kyphosus bigibbus</i>	Brown chub	Preguiçosa-cinzenta	238
<i>Kyphosus sectatrix</i>	Bermuda sea chub	Preguiçosa-branca	238
<i>Kyphosus vaigiensis</i>	Brassy chub	Preguiçosa-bronzeada	239
<i>Labichthys carinatus</i>	Shortgut fintail snipe eel	Cobra-de-crista	68
<i>Labrisomus nuchipinnis</i>	Hairy blenny	Marachomba-peluda	206
<i>Labrus bergylta</i>	Ballan wrasse	Bodião-reticulado	282
<i>Labrus merula</i>	Brown wrasse	Bodião-fusco	283
<i>Labrus mixtus</i>	Cuckoo wrasse	Bodião-canário	283
<i>Labrus viridi</i>	Green wrasse	Bodião-torto	283
<i>Laemonema robustum</i>	Robust cod	Abrótea-robusta	140
<i>Laemonema yarrellii</i>	Codling	Abrótea-de-Yarrell	140
<i>Lagocephalus laevigatus</i>	Smooth puffer	Peixe-bola-verde	224
<i>Lagocephalus lagocephalus</i>	Oceanic puffer	Peixe-bola-oceânico	225
<i>Lamna nasus</i>	Porbeagle	Tubarão-sardo	19
<i>Lampadена anomala</i>	Anomalous lanternfish	Peixe-lanterna-anómalo	120
<i>Lampadена atlantica</i>	Atlantic tail-light lanternfish	Peixe-lanterna-romeirinho	120
<i>Lampadена chavesi</i>	Chaves' lanternfish	Peixe-lanterna-de-Chaves	120
<i>Lampadена luminosa</i>	Luminous lanternfish	Peixe-lanterna-luminoso	121
<i>Lampadена speculigera</i>	Mirror lanternfish	Peixe-lanterna-espelhado	121
<i>Lampanyctus alatus</i>	Winged lanternfish	Peixe-lanterna-alado	121
<i>Lampanyctus ater</i>	Dusky lanternfish	Peixe-lanterna-escuro	121
<i>Lampanyctus crocodilus</i>	Jewel lanternfish	Peixe-lanterna-crocodilo	121
<i>Lampanyctus cuprarius</i>	Atlantic blackcap lanternfish	Peixe-lanterna-cobreado	121
<i>Lampanyctus festivus</i>	Festive lanternfish	Peixe-lanterna-divertido	122
<i>Lampanyctus gemmifer</i>	No English common name	Peixe-lanterna-nacarado	122
<i>Lampanyctus intricarius</i>	Intricate lanternfish	Peixe-lanterna-complexo	122
<i>Lampanyctus lineatus</i>	Lanternfish	Peixe-lanterna-riscado	122
<i>Lampanyctus macdonaldi</i>	MacDonald's lanternfish	Peixe-lanterna-de-MacDonald	122
<i>Lampanyctus niger</i>	Black lantern fish	Peixe-lanterna-preto	122
<i>Lampanyctus photonotus</i>	Dotback lanternfish	Peixe-lanterna-lobisomem	123
<i>Lampanyctus pusillus</i>	Pygmy lanternfish	Peixe-lanterna-anão	123
<i>Lampetra fluviatilis</i>	European river lamprey	Lampreia-do-rio	15
<i>Lampris guttatus</i>	Opah	Peixe-cravo	127
<i>Lampris lauta</i>	East Atlantic Opah	Peixe-cravo-elegante	127
<i>Lamprogrammus niger</i>	No English common name	Abadejo-cristado	158
<i>Lappanella fasciata</i>	Sharp-toothed wrasse	Bodião-verdelho	284
<i>Lasiognathus amphirhamphus</i>	No English common name	Peixe-ganchudo	219
<i>Lasiognathus beebei</i>	No English common name	Peixe-ganchudo-de-Beebe	219
<i>Lasiognathus saccostoma</i>	No English common name	Peixe-ganchudo-tridente	219
<i>Lebetus guilleti</i>	Guillet's goby	Caboz-de-Guillet	182
<i>Lebetus scorpioides</i>	Diminutive goby	Caboz-anão	182
<i>Lepadogaster candolii</i>	Connemarra clingfish	Sugador-de-De Candolle	205
<i>Lepadogaster lepadogaster</i>	Shore clingfish	Sugador-de-Gouan	205
<i>Lepadogaster purpurea</i>	Cornish sucker	Sugador-púrpura	205
<i>Lepidion guentheri</i>	Guenther's codling	Abrótea-de-Günther	141
<i>Lepidion lepidion</i>	Mediterranean codling	Abrótea-do-Mediterrâneo	141
<i>Lepidion schmidti</i>	Giant lepidion	Abrótea-de-Schmidt	141
<i>Lepidocybium flavobrunneum</i>	Escolar	Escolar-preto	162
<i>Lepidophanes gaussi</i>	Gauss' lanternfish	Peixe-lanterna-de-Gauss	123
<i>Lepidophanes guentheri</i>	Günther's lanternfish	Peixe-lanterna-de-Günther	123

Scientific name	English common name	Portuguese common name	page
<i>Lepidopus caudatus</i>	Silver scabbardfish	Peixe-espada	165
<i>Lepidorhombus boscii</i>	Four-spot megrim	Areeiro-de-quatro-manchas	187
<i>Lepidorhombus whiffiagonis</i>	Megrim	Areeiro	187
<i>Lepidotrigla cavillone</i>	Large-scaled gurnard	Ruivo	236
<i>Lepidotrigla dieuzeidei</i>	Spiny gurnard	Ruivo-espinhosos	236
<i>Leptacanthichthys gracilispinis</i>	Plainchin dreamarm	Peixe-farol-de-queixo-direito	217
<i>Leptocephalus giganteus</i>	Giant leptocephalus	sem nome comum	57
<i>Leptochilichthys agassizii</i>	Agassiz's tongueless smooth-head	Celindra-de-Agassiz	77
<i>Leptochilichthys pinguis</i>	Vaillant's tongueless smooth-head	Celindra-de-Vaillant	77
<i>Leptoderma macrophtalmum</i>	No English common name	Celindra-dos-Açores	78
<i>Leptoderma macrops</i>	Grenadier smooth-head	Celindra-olhuda	78
<i>Leptostomias gladiator</i>	Scaleless dragonfish	Peixe-dragão-gladiador	100
<i>Leptostomias haplocaulus</i>	No English common name	Peixe-dragão-de-barbilo-singelo	101
<i>Leptostomias leptobolus</i>	No English common name	Peixe-dragão-de-bolbo-alongado	101
<i>Leptostomias longibarba</i>	No English common name	Peixe-dragão-de-barbilo-longo	101
<i>Lestidiops affinis</i>	Barracudina	Barracudina-ruça	109
<i>Lestidiops jayakari</i>	Pacific barracudina	Barracudina-do-Pacífico	109
<i>Lestidiops pseudosphyraenoides</i>	No English common name	Barracudina-do-Atlântico-nordeste	110
<i>Lestidiops similis</i>	No English common name	Barracudina-análoga	110
<i>Lestidiops sphyreoides</i>	Barracudina	Barracudina-do-Mediterrâneo	110
<i>Lestidium atlanticum</i>	Atlantic barracudina	Barracudina-do-Atlântico	110
<i>Lesueurigobius friesii</i>	Fries's goby	Caboz-de-escama-grande	182
<i>Lesueurigobius sanzi</i>	Sanzo's goby	Caboz-de-Sanzo	182
<i>Lesueurigobius suerii</i>	Lesueur's goby	Caboz-de-Lesueur	183
<i>Lesueurigobius heterofasciatus</i>	No English common name	Caboz-listado	182
<i>Leucobrotula adipata</i>	No English common name	Brótula-falsa-branca	161
<i>Leucoraja circularis</i>	Sandy ray	Raia-de-São-Pedro	44
<i>Leucoraja fullonica</i>	Shagreen ray	Raia-pregada	44
<i>Leucoraja naevus</i>	Cuckoo ray	Raia-de-dois-olhos	45
<i>Lichia amia</i>	Leerfish	Palombeta	253
<i>Linophryne arborifera</i>	No English common name	Peixe-sestro-ramoso	220
<i>Linophryne arcturi</i>	No English common name	Peixe-sestro-do-Arcturus	220
<i>Linophryne brevibarbata</i>	No English common name	Peixe-sestro-de-barbilo-trirramoso	220
<i>Linophryne coronata</i>	No English common name	Peixe-sestro-coroad	221
<i>Linophryne indica</i>	Headlight angler	Peixe-sestro-do-Índico	221
<i>Linophryne lucifer</i>	No English common name	Peixe-sestro-lucifer	221
<i>Linophryne macrodon</i>	No English common name	Peixe-sestro-dentudo	221
<i>Linophryne maderensis</i>	No English common name	Peixe-sestro-da-Madeira	221
<i>Linophryne polypogon</i>	No English common name	Peixe-sestro-barbudo	221
<i>Linophryne racemifera</i>	No English common name	Peixe-sestro-pompom	221
<i>Linophryne sexfilis</i>	No English common name	Peixe-sestro-hexafilamentoso	221
<i>Linophryne algibarbata</i>	No English common name	Peixe-sestro-de-Waterman	220
<i>Liparis liparis</i>	Striped seasnail	Peixe-lesma-listado	237
<i>Liparis montagui</i>	Montagus seasnail	Peixe-lesma-de-Montagus	238
<i>Lipophrys pholis</i>	Shanny	Marachomba-frade	208
<i>Lipophrys trigloides</i>	No English common name	Marachomba-trigloide	208
<i>Lithognathus mormyrus</i>	Sand steenbras	Ferreira	269
<i>Liza aurata</i>	Golden grey mullet	Tainha-garrento	203
<i>Lobianchia dofleini</i>	Doflein's lantern fish	Peixe-lanterna-de-Doflein	123
<i>Lobianchia gemellarii</i>	Gemellar's lanternfish	Peixe-lanterna-de-Gemellar	124
<i>Lobotes surinamensis</i>	Tripletail	Furriel	261
<i>Lophius budegassa</i>	Blackbellied angler	Tamboril-sovaco-preto	212
<i>Lophius piscatorius</i>	Angler	Tamboril	212
<i>Lophodolos acanthognathus</i>	Whalehead dreamer	Peixe-farol-cabeça-de-baleia	217
<i>Lophotus lacepede</i>	Crested oarfish	Peixe-remo-de-crista	127

Scientific name	English common name	Portuguese common name	page
<i>Loweina interrupta</i>	Barebelly lanternfish	Peixe-lanterna-ferreiro	124
<i>Loweina rara</i>	Rare lanternfish	Peixe-lanterna-raro	124
<i>Lutjanus cyanopterus</i>	Cubera snapper	Luciano-de-ventre-vermelho	261
<i>Lutjanus goreensis</i>	Gorean snapper	Luciano-da-Goreia	261
<i>Luvarus imperialis</i>	Luvar	Boquinho	295
<i>Lycenchelys alba</i>	No English common name	Enguia-beiçuda-alva	288
<i>Lyconus brachycolus</i>	No English common name	Peixe-de-focinho-arreganhado	148
<i>Macroparalepis affinis</i>	Barracudina	Barracudina-das-Canárias	110
<i>Macroparalepis brevis</i>	No English common name	Barracudina-pequena	110
<i>Macroparalepis nigra</i>	No English common name	Barracudina-negra	110
<i>Macroramphosus gracilis</i>	Slender snipefish	Trombeteiro-fino	174
<i>Macroramphosus scolopax</i>	Longspine snipefish	Trombeteiro	174
<i>Macrourus berglax</i>	Roughhead grenadier	Lagartixa-do-mar-olho-de-cebola	136
<i>Macruronus maderensis</i>	No English common name	Peixe-orelhudo-da-Madeira	148
<i>Magnisudis atlantica</i>	Duckbill barracudina	Barracudina-bico-de-pato	110
<i>Makaira nigricans</i>	Blue marlin	Espadim-azul-do-Atlântico	185
<i>Malacocephalus laevis</i>	Softhead grenadier	Lagartixa-do-mar-redonda	136
<i>Malacocephalus occidentalis</i>	Western softhead grenadier	Lagartixa-do-mar-ocidental	136
<i>Malacoraja kreffti</i>	Krefft's ray	Raia-de-Krefft	45
<i>Malacosteus niger</i>	Stoplight loosejaw	Peixe-dragão-de-draga	101
<i>Manducus maderensis</i>	No English common name	Boqueirão-da-Madeira	88
<i>Margrethia obtusirostra</i>	Bighead portholefish	Boqueirão-cabeçudo	88
<i>Masturus lanceolatus</i>	Sharptail mola	Peixe-lua-rabudo	222
<i>Mauligobius maderensis</i>	No English common name	Caboz-da-Madeira	183
<i>Maulisia argipalla</i>	Palegold searsid	Celindra-de-tubo-dourada	80
<i>Maulisia mauli</i>	Maul's searsid	Celindra-de-tubo-de-Maul	80
<i>Maulisia microlepis</i>	Smallscale searsid	Celindra-de-tubo-de-escama-curta	80
<i>Maurolicus amethystinopunctatus</i>	Pearlsides	Peixe-luz	90
<i>Maurolicus muelleri</i>	Silvery lightfish	Peixe-luz-de-Muller	91
<i>Megalops atlanticus</i>	Tarpon	Tarpão-do-Atlântico	55
<i>Melamphaes ebelingi</i>	No English common name	Peixe-rã-de-Ebeling	150
<i>Melamphaes falsidicus</i>	No English common name	Peixe-rã-falso	150
<i>Melamphaes indicus</i>	No English common name	Peixe-rã-do-Índico	150
<i>Melamphaes longivelis</i>	Eyebrow bigscale	Peixe-rã-sobrolho	150
<i>Melamphaes microps</i>	Ridgehead	Peixe-rã-olhinho	150
<i>Melamphaes polylepis</i>	No English common name	Peixe-rã-cabecinha	151
<i>Melamphaes purnilus</i>	No English common name	Peixe-rã-anão	151
<i>Melamphaes simus</i>	Ridgehead	Peixe-rã-nariz-achatado	151
<i>Melamphaes suborbitalis</i>	Shoulderspine bigscale	Peixe-rã-de-ombro-espinhoso	151
<i>Melamphaes typhlops</i>	Ridgehead	Peixe-rã-da-Madeira	151
<i>Melanocetus johnsonii</i>	Humpback anglerfish	Diabo-marinho-de-Johnson	215
<i>Melanocetus murrayi</i>	Deepsea blackdevil	Diabo-marinho-de-Murray	215
<i>Melanolagus bericoides</i>	Bigscale deepsea smelt	Argentina-da-fundura-escamuda	84
<i>Melanonus zugmayeri</i>	Arrowtail	Peixe-cauda-de-seta	142
<i>Melanostigma atlanticum</i>	Atlantic soft pout	Enguia-beiçuda-do-Atlântico	288
<i>Melanostomias bartonbeani</i>	Scaleless black dragonfish	Peixe-dragão-de-Barton Bean	101
<i>Melanostomias biseriatus</i>	No English common name	Peixe-dragão-de-barbilho-bisseriado	101
<i>Melanostomias macrophotus</i>	No English common name	Peixe-dragão-luminoso	101
<i>Melanostomias melanopogon</i>	No English common name	Peixe-dragão-de-barbilho-negro	102
<i>Melanostomias melanops</i>	No English common name	Peixe-dragão-de-Sumatra	102
<i>Melanostomias spilorhynchus</i>	No English common name	Peixe-dragão-de-focinho-manchado	102
<i>Melanostomias tentaculatus</i>	Tentacle dragonfish	Peixe-dragão-tentaculado	102
<i>Melanostomias valdiviae</i>	Valdivia black dragon fish	Peixe-dragão-de-Valdívia	102
<i>Mentodus facilis</i>	Dark searsid	Celindra-de-tubo-escura	81
<i>Mentodus mesalirus</i>	Midridged searsid	Celindra-de-tubo-de-nervura	81

Scientific name	English common name	Portuguese common name	page
<i>Mentodus rostratus</i>	Beaked searsid	Celindra-de-tubo-bicuda	81
<i>Merlangius merlangus</i>	Whiting	Badejo	143
<i>Merluccius merluccius</i>	European hake	Pescada-branca	148
<i>Meteoria erythrops</i>	No English common name	Abadejo-cego-dos-Açores	161
<i>Meteoria longidorsalis</i>	No English common name	Abadejo-cego-de-dorsal-longa	161
<i>Microchirus azevia</i>	Bastard sole	Azevia	192
<i>Microchirus boscanion</i>	Lusitanian sole	Azevia-lusitânica	193
<i>Microchirus hexophthalmus</i>	Ocellated wedge sole	Azevia-de-olhos	193
<i>Microchirus ocellatus</i>	Foureyed sole	Azevia-de-malhas	193
<i>Microchirus variegatus</i>	Thickback sole	Azevia-raiada	193
<i>Microchirus wittei</i>	Banded sole	Azevia-raiada-africana	194
<i>Microichthys coccoi</i>	Dwarf deepwater cardinalfish	Olhudo-anão	240
<i>Microlipophrys canevae</i>	No English common name	Marachomba-de-Caneva	208
<i>Microlipophrys dalmatinus</i>	No English common name	Marachomba-dálmata	209
<i>Microlophichthys microlophus</i>	Short-rod anglerfish	Peixe-farol-haste-curta	217
<i>Micromesistius poutassou</i>	Blue whiting	Verdinho	143
<i>Microstoma microstoma</i>	Slender argentine	Argentina-fina	83
<i>Microstomus kitt</i>	Lemon sole	Solha-limão	191
<i>Mirapinna esau</i>	Hairyfish	Peixe-baleia-Esaú	154
<i>Mirognathus normani</i>	Norman's smooth-head	Celindra-de-Norman	78
<i>Mitsukurina owstoni</i>	Goblin shark	Tubarão-duede	17
<i>Mobula birostris</i>	Giant manta	Manta-gigante	52
<i>Mobula mobular</i>	Devil fish	Manta-do-Mediterrâneo	52
<i>Mobula tarapacana</i>	Chilean devil ray	Manta-cornuda	52
<i>Mobula thurstoni</i>	Smoothtail mobula	Manta-de-rabo-liso	53
<i>Mola alexandrini</i>	Bump-head sunfish	Peixe-lua-de-Ranzani	222
<i>Mola mola</i>	Ocean sunfish	Peixe-lua	222
<i>Molva dypterygia</i>	Blue ling	Maruca-azul	146
<i>Molva macrophthalma</i>	Spanish ling	Maruca-do-Mediterrâneo	146
<i>Molva molva</i>	Ling	Maruca	146
<i>Monacoa grimaldii</i>	Mirrorbelly	Peixe-sola-de-Grimaldi	85
<i>Monochirus atlanticus</i>	Whiskered sole	Cascarra-do-Atlântico	194
<i>Monochirus hispidus</i>	Whiskered sole	Cascarra	194
<i>Monognathus bertini</i>	Bertin's one-jawed eel	Enguia-da-fundura-de-Bertin	70
<i>Monognathus boehlkei</i>	Böhlke's one-jawed eel	Enguia-da-fundura-de-Böhlke	70
<i>Monognathus herringi</i>	Herring's one-jawed eel	Enguia-da-fundura-de-Herring	70
<i>Monognathus jesperseni</i>	No English common name	Enguia-da-fundura-de-Jespersen	70
<i>Monognathus nigeli</i>	Nigel's one-jawed eel	Enguia-da-fundura-de-Nigel	70
<i>Monolene microstoma</i>	Smallmouth moonflounder	Carta-de-boca-pequena	190
<i>Monomitopus metriostoma</i>	No English common name	Abadejo-da-Mauritânia	158
<i>Mora moro</i>	Common mora	Mora	141
<i>Mugil cephalus</i>	Flathead grey mullet	Tainha-olhalvo	204
<i>Mullus barbatus</i>	Red mullet	Salmonete-da-vasa	277
<i>Mullus surmuletus</i>	Surmullet	Salmonete-legítimo	277
<i>Muraena augusti</i>	White-eyed moray	Moreia-da-Macaronésia	62
<i>Muraena helena</i>	Mediterranean moray	Moreia	62
<i>Muraena melanotis</i>	Honeycomb moray	Moreia-pintada	63
<i>Mustelus asterias</i>	Starry smooth-hound	Cação-pintado	23
<i>Mustelus canis</i>	Dusky smooth-hound	Cação-cachorro	23
<i>Mustelus mustelus</i>	Dusky smooth-hound	Cação-liso	24
<i>Mustelus punctulatus</i>	Blackspotted smooth-hound	Cação-ponteado	24
<i>Mycteroperca fusca</i>	Island grouper	Garoupa-insular	244
<i>Mycteroperca phenax</i>	Scamp	Garoupa-mariola	245
<i>Mycteroperca rubra</i>	Mottled grouper	Garoupa-chumbo	245
<i>Myctophum affine</i>	Metallic lanternfish	Peixe-laterna-metálico	124

Scientific name	English common name	Portuguese common name	page
<i>Myctophum nitidulum</i>	Pearlyspotted lanternfish	Peixe-lanterna-fosforecente	124
<i>Myctophum punctatum</i>	Spotted lanternfish	Peixe-lanterna-manchado	124
<i>Myliobatis aquila</i>	Common eagle ray	Ratão-água	51
<i>Myoxocephalus scorpius</i>	Shorthorn sculpin	Escorpião	237
<i>Myrichthys pardalis</i>	Leopard eel	Cobra-leopardo	64
<i>Myrophis plumbeus</i>	Leaden worm eel	Cobra-chumbo	64
<i>Myxine glutinosa</i>	Hagfish	Enguia-de-casulo	15
<i>Nansenia atlantica</i>	No English common name	Argentina-atlântica	83
<i>Nansenia boreacrassicauda</i>	No English common name	Argentina-boreal	83
<i>Nansenia cf. pelagica</i>	No English common name	Argentina-pelágica	83
<i>Nansenia groenlandica</i>	Greenland argentine	Argentina-da-Groenlândia	83
<i>Nansenia iberica</i>	No English common name	Argentina-ibérica	83
<i>Nansenia longicauda</i>	No English common name	Argentina-de-cauda-longa	83
<i>Nansenia obliqua</i>	No English common name	Argentina-do-Mediterrâneo	83
<i>Nansenia tenera</i>	No English common name	Argentina-dourada	83
<i>Narcetes erimelas</i>	No English common name	Celindra-de-Goa	78
<i>Narcetes stomias</i>	Blackhead salmon	Celindra-de-cabeça-negra	78
<i>Naucrates ductor</i>	Pilotfish	Peixe-piloto	253
<i>Nealotus triplus</i>	Black snake mackerel	Escolar-coelho-de-natura	163
<i>Negaprion brevirostris</i>	Lemon shark	Tubarão-limão	28
<i>Nemichthys curvirostris</i>	Pale threadtail snipe eel	Cobra-pálida-ponteada	68
<i>Nemichthys scolopaceus</i>	Slender snipe eel	Cobra-de-bico	68
<i>Neoceratias spinifer</i>	Spiny seadevil	Diabo-marinho-espinhososo	215
<i>Neocyttus helgae</i>	Irish oreo	Galo-irlandês	129
<i>Neocyttus rhomboidalis</i>	Spiky oreo	Galo-espinhososo	129
<i>Neogobius fluviatilis</i>	Monkey goby	Caboz-do-rio	183
<i>Neonesthes capensis</i>	Cape snaggletooth	Peixe-dragão-do-Cabo	102
<i>Neoraja iberica</i>	Iberian pygmy skate	Raia-pigmeia-ibérica	45
<i>Neoscopelus macrolepidotus</i>	Large-scaled lantern fish	Peixe-lanterna-escamudo	113
<i>Neoscopelus microchir</i>	Shortfin neoscopelid	Peixe-lanterna-de-barbatanas-curtas	113
<i>Nerophis lumbriciformis</i>	Worm pipefish	Marinha-lombriga	176
<i>Nerophis maculatus</i>	Spotted pipefish	Marinha-sarapintada	176
<i>Nerophis ophidion</i>	Straightnose pipefish	Marinha-fina	176
<i>Nesiarchus nasutus</i>	Black gemfish	Escolar-geribé	163
<i>Nessorhamphus ingolfianus</i>	Duckbill oceanic eel	Enguia-bico-de-pato	63
<i>Nettastoma melanurum</i>	Blackfin sorcerer	Cobra-de-cauda-preta	65
<i>Nezumia aequalis</i>	Common Atlantic grenadier	Largatixa-do-mar-lisa	136
<i>Nezumia bairdii</i>	Marlin-spike grenadier	Largatixa-do-mar-de-Baird	137
<i>Nezumia longebarbata</i>	Bluntnose grenadier	Largatixa-do-mar-de-focinho-curto	137
<i>Nezumia sclerorhynchus</i>	Roughtip grenadier	Largatixa-do-mar-áspera	137
<i>Nameus gronovii</i>	Man-of-war fish	Pequeno-argonauta	173
<i>Normichthys operosus</i>	Multipore searsid	Celindra-de-tubo-multiporo	81
<i>Notacanthus bonaparte</i>	Bonaparte's spiny eel	Enguia-espinhosa-de-Bonaparte	57
<i>Notacanthus chemnitzae</i>	Chemnitz's spiny eel	Enguia-espinhosa-de-Chemnitz	57
<i>Notolichnus valdiviae</i>	Topside lanternfish	Peixe-lanterna-do-Valdívia	125
<i>Notoscopelus bolini</i>	Fanfare lanternfish	Peixe-lanterna-de-Bolin	125
<i>Notoscopelus caudispinosus</i>	Spinetail lanternfish	Peixe-lanterna-de-cauda-espinhosa	125
<i>Notoscopelus elongatus</i>	No English common name	Peixe-lanterna-do-Mediterrâneo	125
<i>Notoscopelus kroyeri</i>	Krøyer's lanternfish	Peixe-lanterna-de-Krøyer	126
<i>Notoscopelus resplendens</i>	Patchwork lanternfish	Peixe-lanterna-oliva	126
<i>Nybelinella brevianalis</i>	No English common name	Abadejo-cego-da-Madeira	161
<i>Nybelinella erikssoni</i>	No English common name	Abadejo-cego-de-Eriksson	161
<i>Oblada melanura</i>	Saddled seabream	Dobradiça	269
<i>Odontaspis ferox</i>	Sand tiger shark	Tubarão-areia	17
<i>Odontaspis noronhai</i>	Bigeye sand tiger shark	Tubarão-de-Noronha	17

Scientific name	English common name	Portuguese common name	page
<i>Odontomacrurus murrayi</i>	Roundhead grenadier	Lagartixa-do-mar-de-Murray	137
<i>Odontostomops normalops</i>	Undistinguished sabretooth	Dente-de-sabre-olho-redondo	112
<i>Oedalechilus labeo</i>	Boxlip mullet	Tainha-sabão	204
<i>Omosudis lowii</i>	Omosudid	Peixe-mandíbula-de-martelo	112
<i>Oncorhynchus mykiss</i>	Rainbow trout	Truta-arco-íris	85
<i>Oneirodes anisacanthus</i>	No English common name	Peixe-farol-de-bolbo-assimétrico	217
<i>Oneirodes carlsbergi</i>	No English common name	Peixe-farol-de-Carlsberg	218
<i>Oneirodes clarkei</i>	No English common name	Peixe-farol-de-Clarke	218
<i>Oneirodes eschrichtii</i>	Bulbous dreamer	Peixe-farol-de-Eschricht	218
<i>Oneirodes krefftii</i>	Krefft's dreamer	Peixe-farol-de-Krefft	218
<i>Oneirodes macronema</i>	No English common name	Peixe-farol-de-Santa Cruz	218
<i>Oneirodes macrosteus</i>	No English common name	Peixe-farol-do-Atlântico-Norte	218
<i>Oneirodes myrionemus</i>	No English common name	Peixe-farol-multifilamentoso	218
<i>Oneirodes posti</i>	No English common name	Peixe-farol-de-Post	218
<i>Opetogenys gracilis</i>	No English common name	Sugador-do-Mediterrâneo	206
<i>Ophidion barbatum</i>	Snake blenny	Abadejo-cobrelo	158
<i>Ophioblennius atlanticus</i>	Redlip Blenny;	Marachomba-do-Atlântico	209
<i>Ophisurus serpens</i>	Serpent eel	Cobra-do-mar	64
<i>Opisthoproctus soleatus</i>	Barrel-eye	Peixe-sola	85
<i>Orcynopsis unicolor</i>	Plain bonito	Palometá	167
<i>Oxynotus centrina</i>	Angular roughshark	Peixe-porco	34
<i>Oxynotus paradoxus</i>	Sailfin roughshark	Peixe-porco-de-vela	35
<i>Pachycara saldanhai</i>	No English common name	Enguia-beiçuda-de-Saldanha	288
<i>Pachycara thermophilum</i>	No English common name	Enguia-beiçuda-termofílica	288
<i>Pachystomias microdon</i>	Smalltooth dragonfish	Peixe-dragão-dente-curto	102
<i>Pagellus acarne</i>	Axillary seabream	Besugo	270
<i>Pagellus bellottii</i>	Red pandora	Bica-buço	270
<i>Pagellus bogaraveo</i>	Blackspot seabream	Goraz	270
<i>Pagellus erythrinus</i>	Common pandora	Bica	271
<i>Pagrus auriga</i>	Redbanded seabream	Pargo-sêmola	271
<i>Pagrus caeruleostictus</i>	Bluespotted seabream	Pargo-ruço	272
<i>Pagrus pagrus</i>	Red porgy	Pargo-legítimo	272
<i>Parablennius gattorugine</i>	Tompot blenny	Marachomba-babosa	209
<i>Parablennius incognitus</i>	No English common name	Marachomba-das-cracas	209
<i>Parablennius parvicornis</i>	Rock-pool blenny	Marachomba-das-poças	210
<i>Parablennius pilicornis</i>	Ringneck blenny	Marachomba-cornuda	210
<i>Parablennius rouxi</i>	No English common name	Marachomba-branca	210
<i>Parablennius ruber</i>	Portuguese blenny	Marachomba-portuguesa	210
<i>Parablennius sanguinolentus</i>	Rusty blenny	Marachomba-do-Mar-Negro	211
<i>Parablennius tentacularis</i>	Tentacled blenny	Marachomba-tentaculada	211
<i>Parabrotula plagiophthalma</i>	False cusk	Brótula-falsa-dos-Açores	161
<i>Paracaristius maderensis</i>	No English common name	Peixe-crina-da-Madeira	260
<i>Paraconger macrops</i>	Manypore conger	Congro-de-natura	67
<i>Paraconger notialis</i>	Guinean conger	Congro-da-Guiné	67
<i>Paralepis brevirostris</i>	No English common name	Barracudina-de-focinho-curto	110
<i>Paralepis coregonoides</i>	Lancet fish	Barracudina-lanceta	111
<i>Paralepis elongata</i>	Barracudina	Barracudina-translúcida	111
<i>Paralepis speciosa</i>	No English common name	Barracudina-manchada	111
<i>Paraliparis copei</i>	Blacksnout seasnail	Peixe-lesma-de-focinho-preto	238
<i>Paraliparis membranaceus</i>	No English common name	Peixe-lesma-membranoso	238
<i>Paraliparis nigellus</i>	No English common name	Peixe-lesma-escuro	238
<i>Parapristipoma humile</i>	Guinean grunt	Roncador-canela	262
<i>Parapristipoma octolineatum</i>	African striped grunt	Riscado	262
<i>Paraphidion vassali</i>	No English common name	Abadejo-vermelho	158
<i>Pegusa impar</i>	Adriatic sole	Linguado-do-Adriático	194

Scientific name	English common name	Portuguese common name	page
<i>Pegusa lascaris</i>	Sand sole	Linguado-da-areia	194
<i>Pegusa nasuta</i>	Snouted sole	Linguado-narigudo	195
<i>Penopus microphthalmus</i>	No English common name	Abadejo-de-olho-pequeno	158
<i>Pentherichthys atratus</i>	No English common name	Peixe-farol-do-Panamá	218
<i>Peristedion cataphractum</i>	African armoured searobin	Cabra-de-casca	236
<i>Persparsia kopua</i>	Spangled tubeshoulder	Celindra-de-tubo-estrelada	81
<i>Petromyzon marinus</i>	Sea lamprey	Lampreia-do-mar	15
<i>Phenacoscorpius nebris</i>	Short-tube scorpionfish	Rascasso-de-tubo-curto	230
<i>Pholis gunnellus</i>	Rock gunnel	Peixe-manteiga	289
<i>Photocorynus spiniceps</i>	No English common name	Peixe-sestro-cabeça-espinhosa	221
<i>Photonectes braueri</i>	Brauer's dragonfish	Peixe-dragão-de-Brauer	102
<i>Photonectes dinema</i>	No English common name	Peixe-dragão-bifilar	103
<i>Photonectes leucospilus</i>	No English common name	Peixe-dragão-de-pinta-branca	103
<i>Photonectes margarita</i>	No English common name	Peixe-dragão-boca-negra	103
<i>Photonectes mirabilis</i>	Blueband dragonfish	Peixe-dragão-maravilhoso	103
<i>Photonectes parvimanus</i>	Fleshyfin dragonfish	Peixe-dragão-barbatana-curta	103
<i>Photostomias atrox</i>	No English common name	Peixe-dragão-atroz	103
<i>Photostomias goodeyeari</i>	No English common name	Peixe-dragão-de-Goodyear	103
<i>Photostomias guernei</i>	Loosejaw	Peixe-dragão-de-Guerne	103
<i>Photostylus pycnopterus</i>	Starry smooth-head	Celindra-luminosa	78
<i>Phtheirichthys lineatus</i>	Slender suckerfish	Pegador-delgado	249
<i>Phycis blennoides</i>	Greater forkbeard	Abrótea-do-alto	146
<i>Phycis phycis</i>	Forkbeard	Abrótea-da-costa	147
<i>Phyllorhinichthys micractis</i>	No English common name	Peixe-farol-de-Guadalupe	218
<i>Physiculus dalwigki</i>	Black codling	Abrótea-de-Dalwigk	142
<i>Pisodonophis semicinctus</i>	Saddled snake eel	Cobra-malhada	65
<i>Platichthys flesus</i>	European flounder	Solha-das-pedras	191
<i>Platybelone argalus</i>	Keeltail needlefish	Agulha-de-quilha	199
<i>Platyberyx andriashevii</i>	No English common name	Peixe-crina-de-Andriashev	260
<i>Platyberyx mauli</i>	No English common name	Peixe-crina-de-Maul	260
<i>Platyberyx opalescens</i>	No English common name	Peixe-crina-opala	260
<i>Platytroctes apus</i>	Legless searsid	Celindra-de-tubo-curva	81
<i>Plectorhinchus mediterraneus</i>	Rubberlip grunt	Pombo	262
<i>Pleuronectes platessa</i>	European plaice	Solha	191
<i>Pollachius pollachius</i>	Pollack	Juliana	143
<i>Pollichthys mauli</i>	Stareye lightfish	Peixe-luminoso-de-Maul	92
<i>Polyacanthonotus challengerii</i>	Challenger's spiny eel	Enguia-espinhosa-do-Challenger	57
<i>Polyacanthonotus merretti</i>	Merrett's spiny eel	Enguia-espinhosa-de-Merrett	58
<i>Polyacanthonotus rissoanus</i>	Smallmouth spiny eel	Enguia-espinhosa-de-boca-pequena	58
<i>Polyipnus polli</i>	No English common name	Pai-velho-redondo	91
<i>Polymetme corythaeola</i>	Rendezvous fish	Peixe-luminoso-do-Índico	92
<i>Polymetme thaeocoryla</i>	No English common name	Peixe-luminoso-do-Atlântico	92
<i>Polymixia nobilis</i>	Stout beardfish	Salmonete-do-alto	149
<i>Polypriion americanus</i>	Wreckfish	Cherne	239
<i>Pomacanthus paru</i>	French angelfish	Lebre-preta	278
<i>Pomadasys incisus</i>	Bastard grunt	Roncador-bravura	263
<i>Pomadasys perotaei</i>	Parrot grunt	Roncador-papagaio	263
<i>Pomatomus saltatrix</i>	Bluefish	Anchova	248
<i>Pomatoschistus kneri</i>	Kner's goby	Caboz-de-Kner	183
<i>Pomatoschistus lozanoi</i>	Lozano's goby	Caboz-de-Lozano	183
<i>Pomatoschistus marmoratus</i>	Marbled goby	Caboz-marmoreado	183
<i>Pomatoschistus microps</i>	Common goby	Caboz-comum	183
<i>Pomatoschistus minutus</i>	Sand goby	Caboz-da-areia	184
<i>Pomatoschistus norvegicus</i>	Norway goby	Caboz-norueguês	184
<i>Pomatoschistus pictus</i>	Mediterranean painted goby	Caboz-de-manchas	184

Scientific name	English common name	Portuguese common name	page
<i>Pontinus kuhlii</i>	Offshore rockfish	Cantarielho-requeime	230
<i>Porogadus miles</i>	Slender cuskeel	Abadejo-fino	159
<i>Poromitra capito</i>	Ridgehead	Peixe-rã-de-nariz-espinhoso	151
<i>Poromitra crassiceps</i>	Crested bigscale	Peixe-rã-de-crista	152
<i>Poromitra megalops</i>	Bigscale	Peixe-rã-olhudo	152
<i>Poromitra nigriceps</i>	No English common name	Peixe-rã-de-cabeça-preta	152
<i>Priacanthus arenatus</i>	Atlantic bigeye	Fura-vasos-vulgar	248
<i>Prionace glauca</i>	Blue shark	Tintureira	28
<i>Pristis pristis</i>	Common sawfish	Espadarte-serra	42
<i>Promethichthys prometheus</i>	Roudi escolar	Escolar-branco	163
<i>Protogrammus sousai</i>	Meteor dragonet	Peixe-pau-de-Sousa	294
<i>Protomyctophum arcticum</i>	Arctic lanternfish	Peixe-lanterna-do-Ártico	126
<i>Psenes cyanophrys</i>	Freckled driftfish	Peixe-pastor-amarelado	173
<i>Psenes maculatus</i>	Silver driftfish	Peixe-pastor-prateado	173
<i>Pseudocaranx dentex</i>	White trevally	Xareu-bicudo	253
<i>Pseudonezumia flagellicauda</i>	No English common name	Lagartixa-do-mar-de-chicote	138
<i>Pseudophichthys splendens</i>	Purplemouthed conger	Congro-de-boca-púrpura	67
<i>Pseudoscopelus altipinnis</i>	No English common name	Peixe-glutão-de-vela	290
<i>Pseudoscopelus obtusifrons</i>	No English common name	Peixe-glutão-de-focinho-curvo	290
<i>Pseudoscopelus sagamianus</i>	No English common name	Peixe-glutão-de-Sagami	290
<i>Pseudoscopelus scutatus</i>	No English common name	Peixe-glutão-escudado	291
<i>Pseudotolithus typus</i>	Longneck croaker	Rainha-branca	276
<i>Pseudotriakis microdon</i>	False catshark	Tubarão-mona	22
<i>Pteraclis carolinus</i>	Fanfish	Freira	259
<i>Pteroplatytrygon violacea</i>	Pelagic stingray	Uge-violeta	50
<i>Pterothriuss gissu</i>	Japanese gissu	Gissu-japonês	55
<i>Pterycombus brama</i>	Atlantic fanfish	Freira-do-Atlântico	259
<i>Puck pinnata</i>	No English common name	Peixe-farol-alado	219
<i>Rachycentron canadum</i>	Cobia	Fogueteiro-galego	249
<i>Radiicephalus elongatus</i>	Tapetail	Peixe-remo-de-cauda-fina	128
<i>Raja asterias</i>	Mediterranean starry ray	Raia-pintada	45
<i>Raja brachyura</i>	Blonde ray	Raia-ponteada	45
<i>Raja clavata</i>	Thornback ray	Raia-lenga	45
<i>Raja maderensis</i>	Madeiran ray	Raia-da-Madeira	46
<i>Raja microocellata</i>	Small-eyed ray	Raia-zimbreira	47
<i>Raja miraletus</i>	Brown ray	Raia-de-quatro-olhos	47
<i>Raja montagui</i>	Mediterranean starry ray	Raia-manchada	47
<i>Raja undulata</i>	Undulate ray	Raia-curva	48
<i>Rajella bathyphila</i>	Deep-water ray	Raia-da-fundura	48
<i>Rajella bigelowi</i>	Bigelow's ray	Raia-de-Bigelow	48
<i>Rajella fyllae</i>	Round ray	Raia-redonda	48
<i>Rajella lintea</i>	Sailray	Raia-nevoeira	48
<i>Raniceps raninus</i>	Tadpole fish	Rainúnculo-negro	143
<i>Ranzania laevis</i>	Slender sunfish	Peixe-lua-comprido	222
<i>Regalecus glesne</i>	King of herrings	Rei-dos-Arenques	128
<i>Regalecus russelli</i>	Oarfish;	Peixe-remo-do-Índico	129
<i>Remora albescens</i>	White suckerfish	Pegador-branco	249
<i>Remora australis</i>	Whalesucker	Pegador-das-baleias	249
<i>Remora brachyptera</i>	Spearfish remora	Pegador-azul	249
<i>Remora osteochir</i>	Marlin sucker	Pegador-dos-espadartes	250
<i>Remora remora</i>	Shark sucker	Pegador	250
<i>Rhadinesthes decimus</i>	Slender snaggletooth	Peixe-dragão-delgado	104
<i>Rhincodon typus</i>	Whale shark	Tubarão-baleia	17
<i>Rhinobatos rhinobatos</i>	Common guitarfish	Viola	41
<i>Rhinochimaera atlantica</i>	Straightnose rabbitfish	Peixe-rato-narigudo	54

Scientific name	English common name	Portuguese common name	page
<i>Rhinoptera marginata</i>	Lusitanian cownose ray	Gavião-do-mar	51
<i>Rhizoprionodon acutus</i>	Milk shark	Tubarão-bicudo	28
<i>Rhomboplites aurorubens</i>	Vermilion snapper	Luciano-rosa	261
<i>Rhynchogadus hepaticus</i>	No English common name	Abrótea-amuada	142
<i>Rhynchohalus natalensis</i>	Glasshead barreleye	Peixe-sola-cabeça-de-vidro	85
<i>Rinocetes nasutus</i>	Abyssal smooth-head	Celindra-abissal	78
<i>Rondeletia bicolor</i>	Hooked redmouth whalefish	Peixe-baleia-bicolor	153
<i>Rondeletia loricata</i>	Redmouth whalefish	Peixe-baleia-de-boca-vermelha	153
<i>Rosenblattichthys hubbsi</i>	Hubb's pearleye	Olho-pérola-de-Hubb	107
<i>Rostroraja alba</i>	White skate	Raia-tairoga	48
<i>Rouleina attrita</i>	Softskin smooth-head	Celindra-macia	78
<i>Rouleina maderensis</i>	Madeiran smooth-head	Celindra-da-Madeira	79
<i>Ruvettus pretiosus</i>	Oilfish	Escolar	163
<i>Saccopharynx ampullaceus</i>	Northern swallower eel	Peixe-pelícano-do-Norte	70
<i>Saccopharynx harrisoni</i>	No English common name	Peixe-pelícano-de-Harrison	71
<i>Saccopharynx hjorti</i>	No English common name	Peixe-pelícano-de-Hjort	71
<i>Saccopharynx paucovertebratis</i>	No English common name	Peixe-pelícano-dos-Açores	71
<i>Saccopharynx ramosus</i>	Shortline swallower eel	Peixe-pelícano-de-cauda-ramosa	71
<i>Saccopharynx thalassa</i>	Hairytail swallower eel	Peixe-pelícano-de-cauda-peluda	71
<i>Saccopharynx trilobatus</i>	No English common name	Peixe-pelícano-de-cauda-trilobada	71
<i>Sagamichthys schnakenbecki</i>	Schnakenbeck's searsid	Celindra-de-tubo-de-Schnakenbeck	81
<i>Salaria basilisca</i>	No English common name	Marachomba-listada	211
<i>Salaria fluviatilis</i>	Freshwater blenny	Marachomba-de-água-doce	211
<i>Salaria pavo</i>	Peacock blenny	Marachomba-pavão	212
<i>Salmo salar</i>	Atlantic salmon	Salmão-do-Atlântico	85
<i>Salmo trutta</i>	Sea trout	Truta-marisca	86
<i>Sarda sarda</i>	Atlantic bonito	Sarrajão	167
<i>Sardina pilchardus</i>	European pilchard	Sardinha	72
<i>Sardinella aurita</i>	Round sardinella	Sardinela-lombuda	73
<i>Sardinella maderensis</i>	Madeiran sardinella	Sardinela-da-Madeira	73
<i>Sargocentron hastatum</i>	Red squirrelfish	Esquilo-real	156
<i>Sarpa salpa</i>	Salema	Salema	273
<i>Scartella cristata</i>	Molly miller	Marachomba-de-crista	212
<i>Schedophilus maculatus</i>	Pelagic butterfish	Liro-manchado	172
<i>Schedophilus medusophagus</i>	Cornish blackfish	Liro-mole	172
<i>Schedophilus ovalis</i>	Imperial blackfish	Liro-imperial	172
<i>Sciadonus pedicellaris</i>	No English common name	Abadejo-cego-do-Pacífico	161
<i>Sciaena umbra</i>	Brown meagre	Roncadeira-preta	276
<i>Scomber colias</i>	Atlantic chub mackerel	Cavala	168
<i>Scomber scombrus</i>	Atlantic mackerel	Sarda	168
<i>Scomberesox saurus</i>	Atlantic saury	Agulhão-do-Atlântico	198
<i>Scomberesox simulans</i>	Dwarf saury	Agulhão-anão	198
<i>Scomberomorus tritor</i>	West African Spanish mackerel	Serra-branca	169
<i>Scombrolabrax heterolepis</i>	Longfin escolar	Escolar-de-peitoral-longa	171
<i>Scopelarchus analis</i>	Short fin pearleye	Olho-pérola-de-barbatana-curta	107
<i>Scopelarchus guentheri</i>	Staring pearleye	Olho-pérola-de-Guenther	107
<i>Scopelarchus michaelsarsi</i>	Bigfin pearleye	Olho-pérola-de-Michael Sars	107
<i>Scopelengys tristis</i>	Pacific blackchin	Peixe-lanterna-do-Pacífico	113
<i>Scopeloberyx opisthoterurus</i>	Bigscale	Peixe-rã-delgado	152
<i>Scopeloberyx robustus</i>	Longjaw bigscale	Peixe-rã-robusto	152
<i>Scopeloberyx rubriventer</i>	Ridgehead	Peixe-rã-dos-Açores	152
<i>Scopelogadus beanii</i>	Bean's bigscale	Peixe-rã-de-Bean	153
<i>Scopelogadus mizolepis</i>	Ragged bigscale	Peixe-rã-couraça	153
<i>Scopelopsis multipunctatus</i>	Multispotted lanternfish	Peixe-lanterna-sarapintado	126
<i>Scopelosaurus argenteus</i>	Waryfish	Peixe-vigia-prateado	107

Scientific name	English common name	Portuguese common name	page
<i>Scopelosaurus lepidus</i>	Blackfin waryfish	Peixe-vigia-de-barbatana-preta	107
<i>Scopelosaurus smithii</i>	No English common name	Peixe-vigia-de-Smith	108
<i>Scophthalmus maximus</i>	Turbot	Pregado	187
<i>Scophthalmus rhombus</i>	Brill	Rodovalho	188
<i>Scorpaena azorica</i>	Azores scorpionfish	Rascasso-dos-Açores	230
<i>Scorpaena canariensis</i>	Canary scorpionfish	Rascasso-das-Canárias	230
<i>Scorpaena elongata</i>	Slender rockfish	Rascasso-rosado	230
<i>Scorpaena laevis</i>	Senegalese rockfish	Rascasso-do-Senegal	231
<i>Scorpaena loppei</i>	Cadenat's scorpionfish	Rascasso-de-Loppé	231
<i>Scorpaena maderensis</i>	Madeira rockfish	Rascasso-da-Madeira	231
<i>Scorpaena notata</i>	Small red scorpionfish	Rascasso-escorpião	232
<i>Scorpaena plumieri</i>	Pacific spotted scorpionfish	Rascasso-negro	232
<i>Scorpaena porcus</i>	Black scorpionfish	Rascasso-de-pintas	232
<i>Scorpaena scrofa</i>	Red scorpionfish	Rascasso-vermelho	233
<i>Scorpaena stephanica</i>	Spotted-fin rockfish	Rascasso-de-barbatana-pontuada	233
<i>Scorpaenodes arenai</i>	Messina rockfish	Rascasso-de-Messina	234
<i>Scyliorhinus canicula</i>	Small-spotted catshark	Pata-roxa	21
<i>Scyliorhinus stellaris</i>	Polkadot catshark	Pata-roxa-gata	22
<i>Scymnodalatias garricki</i>	Azores dogfish	Tubarão-cão-dos-Açores	33
<i>Scymnodon ringens</i>	Knifetooth dogfish	Arreganhada	33
<i>Searsia koefoedi</i>	Koefoed's searsid	Celindra-de-tubo-de-Koefoed	82
<i>Sebastes mentella</i>	Beaked redfish	Peixe-vermelho-da-fundura	229
<i>Sebastes norvegicus</i>	Golden redfish	Peixe-vermelho	229
<i>Selar boops</i>	Oxeye scad	Charro-olho-de-boi	254
<i>Selar crumenophthalmus</i>	Bigeye scad	Charro-preto	254
<i>Selene dorsalis</i>	African moonfish	Corcovado-africano	254
<i>Selene setapinnis</i>	Atlantic moonfish	Corcovado-do-Golfo	254
<i>Seriola dumerili</i>	Greater amberjack	Charuteiro-catarino	254
<i>Seriola fasciata</i>	Lesser amberjack	Charuteiro-listado	255
<i>Seriola lalandi</i>	Yellowtail amberjack	Charuteiro-azeite	255
<i>Seriola rivoliana</i>	Longfin yellowtail	Charuteiro-falcado	255
<i>Seriola zonata</i>	Banded rudderfish	Charuteiro-olhete	256
<i>Seriola carpenteri</i>	Guinean amberjack	Charuteiro-da-Guiné	254
<i>Serranus atricauda</i>	Blacktail comber	Serrano-de-rolo	245
<i>Serranus cabrilla</i>	Comber	Serrano-alecrim	246
<i>Serranus hepatus</i>	Brown comber	Serrano-ferreiro	246
<i>Serranus scriba</i>	Painted comber	Serrano-riscado	246
<i>Serrivomer beanii</i>	Bean's sawtooth eel	Cobra-couraça	68
<i>Serrivomer lanceolatoides</i>	Bronze sawtooth eel	Cobra-de-dentes-curtos	69
<i>Setarches guentheri</i>	Channeled rockfish	Requeime-de-natura	229
<i>Sigmops bathyphilus</i>	Spark anglemouth	Boqueirão-brilhante	88
<i>Sigmops elongatus</i>	Elongated bristlemouth fish	Boqueirão-dentudo	89
<i>Simenchelys parasitica</i>	Snubnosed eel	Fluta	59
<i>Similiparma lurida</i>	Canary damsel	Castanheta-preta	279
<i>Solea senegalensis</i>	Senegalese sole	Linguado-do-Senegal	195
<i>Solea solea</i>	Common sole	Linguado-legítimo	195
<i>Somniosus microcephalus</i>	Greenland shark	Tubarão-da-Gronelândia	34
<i>Somniosus rostratus</i>	Little sleeper shark	Pailona	34
<i>Paracanthurus chrysopterus</i>	Redtail parrotfish	Papagaio	288
<i>Paracanthurus cretense</i>	Parrotfish	Papagaio-velho	287
<i>Sparus aurata</i>	Gilthead seabream	Dourada	273
<i>Spectrunculus crassus</i>	No English common name	Abadejo-dos-Açores	159
<i>Spectrunculus grandis</i>	Pudgy cuskeel	Abadejo-grande	159
<i>Sphagmacrurus grenadæ</i>	Pugnose grenadier	Lagartixa-do-mar-de-nariz-arrebitado	138
<i>Sphagmacrurus hirundo</i>	Swallow grenadier	Lagartixa-do-mar-dos-Açores	138

Scientific name	English common name	Portuguese common name	page
<i>Sphoeroides marmoratus</i>	Guinean pufferfish	Peixe-bola-da-Guiné	225
<i>Sphoeroides pachygaster</i>	Blunthead puffer	Peixe-bola-de-focinho-achatado	225
<i>Sphoeroides spengleri</i>	Bandtail puffer	Peixe-bola-de-pintas	226
<i>Sphyraena gladiifera</i>	Prickly dreamer	Peixe-farol-espinhoso	219
<i>Sphyraena barracuda</i>	Great barracuda	Bicuda-gigante	295
<i>Sphyraena guachancho</i>	Guachanche barracuda	Bicuda-africana	295
<i>Sphyraena sphyraena</i>	European barracuda	Bicuda-europeia	295
<i>Sphyraena viridensis</i>	Yellowmouth barracuda	Bicuda-de-Cabo-Verde	296
<i>Sphyraena lewini</i>	Scalloped hammerhead	Tubarão-martelo-recortado	28
<i>Sphyraena mokarran</i>	Great hammerhead	Tubarão-martelo-gigante	29
<i>Sphyraena zygaena</i>	Smooth hammerhead	Tubarão-martelo	29
<i>Spicara flexuosa</i>	No English common name	Trombeiro-do-Mediterrâneo	274
<i>Spicara maena</i>	Blotched picarel	Trombeiro-choupa	274
<i>Spicara melanurus</i>	Blackspot picarel	Trombeiro-malha-redonda	274
<i>Spicara smaris</i>	Picarel	Trombeiro-boga	274
<i>Spinachia spinachia</i>	Sea stickleback	Esgana-gata-marinha	297
<i>Spondylisoma cantharus</i>	Black seabream	Choupa	274
<i>Sprattus sprattus</i>	European sprat	Espadilha	74
<i>Squaliolus laticaudus</i>	Spined pygmy shark	Tubarão-anão	30
<i>Squalogadus modifikatus</i>	Tadpole whiptail	Lagartixa-do-mar-girino	138
<i>Squalus acanthias</i>	Spotted spiny dogfish	Galhudo-malhado	38
<i>Squalus blainvillei</i>	Longnose spurdog	Galhudo	38
<i>Squalus megalops</i>	Picked spurdog	Galhudo-de-focinho-curto	39
<i>Squalus mitsukurii</i>	Shortspine spurdog	Galhudo-de-espinho-curto	39
<i>Squatina aculeata</i>	Sawback angelshark	Anjo-espinhoso	39
<i>Squatina oculata</i>	Smoothback angelshark	Anjo-de-malhas	39
<i>Squatina squatina</i>	Angelshark	Anjo	40
<i>Stephanolepis hispidus</i>	Planehead filefish	Peixe-porco-áspero	226
<i>Stephanolepis setifer</i>	Pygmy filefish	Peixe-porco-galhudo	227
<i>Sternopyx diaphana</i>	Diaphanous hatchet fish	Pai-velho-diáfano	91
<i>Sternopyx pseudobscura</i>	Highlight hatchetfish	Pai-velho-oblíquo	91
<i>Stomias boa</i>	Boa dragonfish	Peixe-dragão-jibóia	104
<i>Stomias brevibarbus</i>	No English common name	Peixe-dragão-barbilho-curto	104
<i>Stomias longibarbus</i>	Longbarb scaly dragonfish	Peixe-dragão-escamudo	104
<i>Stromateus fiatola</i>	Blue butterfish	Pampo-godinho	173
<i>Stylephorus chordatus</i>	Tube-eye	Peixe-cauda-de-fio	131
<i>Sudis atrox</i>	Fierce pike smelt	Barracudina-feroz	111
<i>Sudis hyalina</i>	Barracudina	Barracudina-imperial	111
<i>Syacium micrurum</i>	Channel flounder	Carta-moira	188
<i>Syacium papillosum</i>	Dusky flounder	Carta-sombria	188
<i>Symbolophorus rufinus</i>	Rufous lanternfish	Peixe-lanterna-de-Rufous	126
<i>Symbolophorus veranyi</i>	Large-scale lantern fish	Peixe-lanterna-de-Vérany	126
<i>Sympodus bailloni</i>	Baillon's wrasse	Bodião-tentilhão	284
<i>Sympodus caeruleus</i>	No English common name	Bodião-dos-Açores	284
<i>Sympodus cinereus</i>	Grey wrasse	Bodião-cinzento	284
<i>Sympodus mediterraneus</i>	Axillary wrasse	Bodião-do-Mediterrâneo	285
<i>Sympodus melops</i>	Corkwing wrasse	Bodião-vulgar	285
<i>Sympodus ocellatus</i>	No English common name	Bodião-de-pinta	285
<i>Sympodus roissali</i>	Five-spotted wrasse	Bodião-manchado	285
<i>Sympodus rostratus</i>	No English common name	Bodião-das-ervas	286
<i>Sympodus tinca</i>	Peacock wrasse	Bodião-pavão	286
<i>Sympodus trutta</i>	Emerald wrasse	Bodião-mosqueado	286
<i>Syphurus insularis</i>	Macaronesian tonguesole	Língua-da-Maraconésia	196
<i>Syphurus ligulatus</i>	Elongate tonguesole	Língua-comprida	196
<i>Syphurus nigrescens</i>	Spotted tonguesole	Língua-avessa	196

Scientific name	English common name	Portuguese common name	page
<i>Syphurus reticulatus</i>	Reticulated tonguefish	Língua-reticulada	197
<i>Synaphobranchus affinis</i>	Grey cutthroat eel	Moreão-do-golfo	59
<i>Synaphobranchus brevidorsalis</i>	Shortdorsal cutthroat eel	Moreão-de-dorsal-curta	59
<i>Synaphobranchus dolichorhynchus</i>	No English common name	Moreão-bicudo	59
<i>Synaphobranchus kaupii</i>	Kaup's arrowtooth eel	Moreão-de-natura	59
<i>Synapturichthys kleinii</i>	Klein's sole	Linguado-turco	195
<i>Synchiropus phaeton</i>	Phaeton dragonet	Peixe-pau-rosa	294
<i>Syngnathus abaster</i>	Black-striped pipefish	Marinha-do-rio	176
<i>Syngnathus acus</i>	Greater pipefish	Marinha-comum	177
<i>Syngnathus fuscus</i>	Northern pipefish	Marinha-setentrional	177
<i>Syngnathus pelagicus</i>	Sargassum pipefish	Marinha-do-sargaço	177
<i>Syngnathus phlegon</i>	No English common name	Marinha-espinhosa	177
<i>Syngnathus rostellatus</i>	Nilsson's pipefish	Marinha-cabeça-chata	178
<i>Syngnathus tenuirostris</i>	Narrow-snouted pipefish	Marinha-de-focinho-estreito	178
<i>Syngnathus typhle</i>	Broadnosed pipefish	Marinha-focinho-grosso	178
<i>Synodus foetens</i>	Inshore lizardfish	Peixe-lagarto-da-praia	108
<i>Synodus saurus</i>	Atlantic lizardfish	Peixe-lagarto-do-Atlântico	108
<i>Synodus synodus</i>	Redbarred lizardfish	Peixe-lagarto-de-rolo	108
<i>Taeniuichthys bathyphilus</i>	Deepwater lanternfish	Peixe-lanterna-da-fundura	126
<i>Taeniuichthys minimus</i>	Waistcoat lanternfish	Peixe-lanterna-de-Cabo-Verde	127
<i>Taeniuichthys paurolychnus</i>	Naked lanternfish	Peixe-lanterna-nu	127
<i>Taenirops grabatus</i>	Round stingray	Uge-redonda	50
<i>Talismania antillarum</i>	Antillean smooth-head	Celindra-das-Antilhas	79
<i>Talismania homoptera</i>	Hairfin smooth-head	Celindra-da-Mauritânia	79
<i>Talismania mekistonema</i>	Threadfin smooth-head	Celindra-de-barbatana-filamentosa	79
<i>Taractes asper</i>	Rough pomfret	Xaputa-rugosa	259
<i>Taractes rubescens</i>	Black pomfret	Xaputa-negra	260
<i>Taractichthys longipinnis</i>	Big-scale pomfret	Xaputa-galhuda	260
<i>Taurulus bubalis</i>	Longspined bullhead	Escorpião-roco	237
<i>Tetragonurus atlanticus</i>	Bigeye squaretail	Escolar-de-natura-olho-grande	173
<i>Tetragonurus cuvieri</i>	Smalleye squaretail	Escolar-de-natura-olho-pequeno	173
<i>Tetrapturus belone</i>	Mediterranean spearfish	Espadim-do-Mediterrâneo	186
<i>Tetrapturus georgii</i>	Roundscale spearfish	Espadim-peto	186
<i>Tetrapturus pfluegeri</i>	Longbill spearfish	Espadim-bicudo	186
<i>Tetronarce nobiliana</i>	Electric ray	Tremelga-negra	40
<i>Thalassoma pavo</i>	Ornate wrasse	Bodião-verde	286
<i>Thorogobius ephippiatus</i>	Leopard-spotted goby	Caboz-leopardo	184
<i>Thunnus alalunga</i>	Albacore	Atum-voador	169
<i>Thunnus albacares</i>	Yellowfin tuna	Atum-albacora	170
<i>Thunnus atlanticus</i>	Blackfin tuna	Atum-de-barbatana-negra	170
<i>Thunnus obesus</i>	Bigeye tuna	Atum-patudo	170
<i>Thunnus thynnus</i>	Atlantic bluefin tuna	Atum-rabilho	170
<i>Torpedo marmorata</i>	Marbled electric ray	Tremelga-marmoreada	41
<i>Torpedo torpedo</i>	Common torpedo	Tremelga-de-olhos	41
<i>Trachinocephalus myops</i>	Snakefish	Peixe-lagarto-de-focinho-rombo	109
<i>Trachinotus carolinus</i>	Florida pompano	Sereia-da-Florida	256
<i>Trachinotus ovatus</i>	Pompano	Sereia-camochilo	256
<i>Trachinus araneus</i>	Spotted weever	Peixe-aranha-pontuado	292
<i>Trachinus draco</i>	Greater weever	Peixe-aranha-maior	292
<i>Trachinus pellegrini</i>	Cape Verde weever	Peixe-aranha-de-Cabo-Verde	292
<i>Trachinus radiatus</i>	Starry weever	Peixe-aranha-raiado	292
<i>Trachipterus arcticus</i>	Dealfish	Peixe-tábuia	128
<i>Trachipterus trachypterus</i>	Mediterranean dealfish	Peixe-tábuia-do-Mediterrâneo	128
<i>Trachonurus sulcatus</i>	Bristly grenadier	Largatixa-do-mar-arrepiada	138
<i>Trachonurus villosus</i>	Furry whiptail	Lagartixa-do-mar-peluda	138

Scientific name	English common name	Portuguese common name	page
<i>Trachurus lathami</i>	Rough scad	Carapau-rugoso	256
<i>Trachurus mediterraneus</i>	Mediterranean horse mackerel	Carapau-do-Mediterrâneo	256
<i>Trachurus picturatus</i>	Blue jack mackerel	Carapau-negrão	257
<i>Trachurus trachurus</i>	Atlantic horse mackerel	Carapau	257
<i>Trachyrincus scabrus</i>	Roughsnout grenadier	Peixe-lima	139
<i>Trachyscorpia cristulata</i>	Atlantic thornyhead	Cantarielho-espinhoso	229
<i>Trichiurus lepturus</i>	Largehead hairtail	Lírio	165
<i>Trigla lyra</i>	Piper gurnard	Cabra-lira	236
<i>Trigonolampa miriceps</i>	Threelight dragonfish	Peixe-dragão-listado	104
<i>Triphoturus nigrescens</i>	Highseas lampfish	Peixe-lanterna-do-alto	127
<i>Tripterygion delaisi</i>	Black-faced blenny	Caboz-de-três-dorsais-de-Delais	206
<i>Tripterygion tripteronotum</i>	No English common name	Caboz-de-três-dorsais-do-Mediterrâneo	206
<i>Trisopterus capelanus</i>	Poor cod Mediterranean	Fançao-do-Mediterrâneo	143
<i>Trisopterus luscus</i>	Pouting	Faneca	144
<i>Trisopterus minutus</i>	Poor cod	Fanecão	144
<i>Tylosurus acus</i>	Agujon needlefish	Agulheta-imperial	199
<i>Tylosurus crocodilus</i>	Hound needlefish	Agulheta-crocodilo	200
<i>Umbrina canariensis</i>	Canary drum	Calafate-das-Canárias	276
<i>Umbrina cirrosa</i>	Shi drum	Calafate-das-riscas	276
<i>Umbrina ronchus</i>	Fusca drum	Calafate-fosco	276
<i>Uncisudis longirostra</i>	No English common name	Barracudina-da-Madeira	111
<i>Uncisudis quadrimaculata</i>	Barracudina	Barracudina-de-quatro-manchas	111
<i>Uranoscopus scaber</i>	Stargazer	Cabeçudo	293
<i>Valenciennellus tripunctulatus</i>	Constellationfish	Pai-velho-de-três-pontos	91
<i>Vanneaugobius canariensis</i>	No English common name	Caboz-das-Canárias	184
<i>Venefica proboscidea</i>	Whipsnout sorcerer	Cobra-de-bico-comprido	65
<i>Vinciguerra attenuata</i>	Slender lightfish	Peixe-luminoso-esguio	92
<i>Vinciguerra lucetia</i>	Panama lightfish	Peixe-luminoso-do-Panamá	92
<i>Vinciguerra nimbaria</i>	Oceanic lightfish	Peixe-luminoso-oceânico	93
<i>Vinciguerra poweriae</i>	Power's deep-water bristle-mouth fish	Peixe-luminoso-de-Power	93
<i>Xanthichthys ringens</i>	Sargassum triggerfish	Cangulo-do-sargaço	228
<i>Xenodermichthys copei</i>	Bluntnose smooth-head	Celindra-de-focinho-curto	79
<i>Xiphias gladius</i>	Swordfish	Espadarte	184
<i>Xyrichtys novacula</i>	Pearly razorfish	Mordedor	287
<i>Yarrella blackfordi</i>	No English common name	Peixe-luminoso-de-Blackford	93
<i>Zameus squamulosus</i>	Smallmouth knifetooth dogfish	Arreganhada-de-focinho-comprido	34
<i>Zenion hololepis</i>	Dwarf dory	Galito-rosado	130
<i>Zenopsis conchifer</i>	Silvery John dory	Galo-branco	130
<i>Zeugopterus punctatus</i>	Topknot	Rodovalho-bruxa	188
<i>Zeugopterus regius</i>	Eckström's topknot	Bruxa	188
<i>Zeus faber</i>	John dory	Galo-negro	130
<i>Zu cristatus</i>	Scalloped ribbonfish	Peixe-tábua-recortado	128

Index of the English names / Índice dos nomes Ingleses

English common name	Scientific name	Portuguese common name	page
Abyssal assfish	<i>Bassozetus compressus</i>	Abadejo-da-fundura	157
Abyssal grenadier	<i>Coryphaenoides armatus</i>	Lagartixa-do-mar-abissal	133
Abyssal halosaur	<i>Halosauropsis macrochir</i>	Peixe-lagarto-abissal	56
Abyssal smooth-head	<i>Rinoctes nasutus</i>	Celindra-abissal	78
Abyssal spiderfish	<i>Bathypterois longipes</i>	Peixe-tripé-abissal	106
Adriatic sole	<i>Pegusa impar</i>	Linguado-do-Adriático	194
Adriatic sturgeon	<i>Acipenser naccarii</i>	Esturjão-do-Adriático	55
African armoured searobin	<i>Peristedion cataphractum</i>	Cabra-de-casca	236
African moonfish	<i>Selene dorsalis</i>	Corcovado-africano	254
African striped grunt	<i>Parapristipoma octolineatum</i>	Riscado	262
Agassiz' slickhead	<i>Alepocephalus agassizii</i>	Celindra-negra-de-Agassiz	74
Agassiz's tongueless smooth-head	<i>Leptochilichthys agassizii</i>	Celindra-de-Agassiz	77
Agujon needlefish	<i>Tylosurus acus</i>	Agulheta-imperial	199
Albacore	<i>Thunnus alalunga</i>	Atum-voador	169
Alexandria pompano	<i>Alectis alexandrina</i>	Xareu-enxada	250
Alfonsino	<i>Beryx decadactylus</i>	Imperador	149
Allis shad	<i>Alosa alosa</i>	Sável	71
American eel	<i>Anguilla rostrata</i>	Enguia-americana	69
Andre's lanternfish	<i>Centrobranchus andreae</i>	Peixe-lanterna-de-Andréas	115
Angelshark	<i>Squatina squatina</i>	Anjo	40
Angler	<i>Lophius piscatorius</i>	Tamboril	212
Angular roughshark	<i>Oxynotus centrina</i>	Peixe-porco	34
Annular seabream	<i>Diplodus annularis</i>	Sargo-alcorraz	265
Anomalous lanternfish	<i>Lampadena anomala</i>	Peixe-lanterna-anómalo	120
Antenna codlet	<i>Bregmaceros atlanticus</i>	Peixe-antena-do-Atlântico	131
Antillean smooth-head	<i>Talismania antillarum</i>	Celindra-das-Antilhas	79
Arctic lanternfish	<i>Protomyctophum arcticum</i>	Peixe-lanterna-do-Ártico	126
Argentine	<i>Argentina sphyraena</i>	Argentina-branca	82
Armless snake eel	<i>Dalophis imberbis</i>	Cobra-imberbe	64
Arrowtail	<i>Melanonus zugmayeri</i>	Peixe-cauda-de-seta	142
Atlantic barracudina	<i>Lestidium atlanticum</i>	Barracudina-do-Atlântico	110
Atlantic bigeye	<i>Priacanthus arenatus</i>	Fura-vasos-vulgar	248
Atlantic blackcap lanternfish	<i>Lampanyctus cuprarius</i>	Peixe-lanterna-cobreado	121
Atlantic blackwing flyingfish	<i>Hirundichthys volador</i>	Peixe-voador-de-asas-negras-do-Atlântico	202
Atlantic bluefin tuna	<i>Thunnus thynnus</i>	Atum-rabilho	170
Atlantic bonito	<i>Sarda sarda</i>	Sarrajão	167
Atlantic chub mackerel	<i>Scomber colias</i>	Cavala	168
Atlantic cod	<i>Gadus morhua</i>	Bacalhau-do-Atlântico	142
Atlantic cornetfish	<i>Aulostomus strigosus</i>	Trombeta	174
Atlantic deepwater dragonet	<i>Centrodraco acanthopoma</i>	Peixe-pau-da-fundura-do-Atlântico	295
Atlantic fanfish	<i>Pterycombus brama</i>	Freira-do-Atlântico	259
Atlantic fangjaw	<i>Gonostoma atlanticum</i>	Boqueirão-do-Atlântico	88
Atlantic footballfish	<i>Himantolophus groenlandicus</i>	Peixe-farol-do-Atlântico	216
Atlantic herring	<i>Clupea harengus</i>	Arenque-do-Atlântico	72
Atlantic horse mackerel	<i>Trachurus trachurus</i>	Carapau	257
Atlantic lanternfish	<i>Diogenichthys atlanticus</i>	Peixe-lanterna-do-Atlântico	118
Atlantic lizardfish	<i>Synodus saurus</i>	Peixe-lagarto-do-Atlântico	108
Atlantic mackerel	<i>Scomber scombrus</i>	Sarda	168
Atlantic moonfish	<i>Selene setapinnis</i>	Corcovado-do-Golfo	254
Atlantic pelagic basslet	<i>Howella atlantica</i>	Peixe-alado-do-Atlântico	240
Atlantic pomfret	<i>Brama brama</i>	Xaputa	259
Atlantic sabretooth	<i>Coccarella atlantica</i>	Dente-de-sabre-do-Atlântico	112
Atlantic salmon	<i>Salmo salar</i>	Salmão-do-Atlântico	85

English common name	Scientific name	Portuguese common name	page
Atlantic saury	<i>Scomberesox saurus</i>	Agulhão-do-Atlântico	198
Atlantic sawtail catshark	<i>Galeus atlanticus</i>	Leitão-do-Atlântico	21
Atlantic silver hatchetfish	<i>Argyropelecus aculeatus</i>	Pai-velho-prateado-do-Atlântico	89
Atlantic soft pout	<i>Melanostigma atlanticum</i>	Enguia-beiçuda-do-Atlântico	288
Atlantic spiny lump sucker	<i>Eumicrotremus spinosus</i>	Peixe-lapa-espinhoso	237
Atlantic tail-light lanternfish	<i>Lampadena atlantica</i>	Peixe-lanterna-romeirinho	120
Atlantic thornyhead	<i>Trachyscorpia cristulata</i>	Cantarilho-espinhoso	229
Atlantic trumpetfish	<i>Aulostomus cf. maculatus</i>	Trombeta-salpicada	174
Atlantic white marlin	<i>Kajikia albida</i>	Espadim-branco-do-Atlântico	185
Avocet snipe eel	<i>Avocettina infans</i>	Cobra-centopeia	67
Axillary seabream	<i>Pagellus acarne</i>	Besugo	270
Axillary wrasse	<i>Syphodus mediterraneus</i>	Bodião-do-Mediterrâneo	285
Azores chromis	<i>Chromis limbata</i>	Castanheta-da-Madeira	279
Azores dogfish	<i>Scymnodalatias garricki</i>	Tubarão-cão-dos-Açores	33
Azores rockling	<i>Gaidropsarus granti</i>	Laibeque-dos-Açores	144
Azores scorpionfish	<i>Scorpaena azorica</i>	Rascasso-dos-Açores	230
Baillon's wrasse	<i>Syphodus bailloni</i>	Bodião-tentilhão	284
Baird's slickhead	<i>Alepocephalus bairdii</i>	Celindra-de-Baird	74
Balao halfbeak	<i>Hemiramphus balao</i>	Meia-agulha-preta	200
Balbo sabretooth	<i>Evermannella balbo</i>	Dente-de-sabre-de-Balbo	112
Ballan wrasse	<i>Labrus bergylta</i>	Bodião-reticulado	282
Banded rudderfish	<i>Seriola zonata</i>	Charuteiro-olhete	256
Banded seabream	<i>Diplodus fasciatus</i>	Sargo-de-faias	267
Banded sole	<i>Microchirus wittei</i>	Azevia-raiada-africana	194
Bandtail puffer	<i>Sphoeroides spengleri</i>	Peixe-bola-de-pintas	226
Bandtooth conger	<i>Ariosoma balearicum</i>	Congro-das-Baleares	65
Bandwing flyingfish	<i>Cheilopogon exsiliens</i>	Peixe-voador-de-asas-listadas	200
Bar jack	<i>Caranx ruber</i>	Xareu-carvoeiro	252
Barbate dragonfish	<i>Eustomias enbarbatus</i>	Peixe-dragão-barbado	98
Barebelly lanternfish	<i>Loweina interrupta</i>	Peixe-lanterna-ferreiro	124
Barethroat slickhead	<i>Asquamiceps hjorti</i>	Celindra-de-Hjort	75
Barracudina	<i>Lestidiops affinis</i>	Barracudina-ruça	109
Barracudina	<i>Lestidiops sphyrenoides</i>	Barracudina-do-Mediterrâneo	110
Barracudina	<i>Macroparalepis affinis</i>	Barracudina-das-Canárias	110
Barracudina	<i>Paralepis elongata</i>	Barracudina-translúcida	111
Barracudina	<i>Sudis hyalina</i>	Barracudina-imperial	111
Barracudina	<i>Uncisudis quadrimaculata</i>	Barracudina-de-quatro-manchas	111
Barred hogfish	<i>Bodianus scrofa</i>	Bodião-dente-de-cão	280
Barrel-eye	<i>Opisthoproctus soleatus</i>	Peixe-sola	85
Barrelfish	<i>Hyperoglyphe perciformis</i>	Liro-errante	172
Basking shark	<i>Cetorhinus maximus</i>	Tubarão-frade	19
Bastard grunt	<i>Pomadasys incisus</i>	Roncador-bravura	263
Bastard sole	<i>Microchirus azevia</i>	Azevia	192
Batnose eel	<i>Dysomma brevirostre</i>	Enguia-de-focinho-bolboso	58
Beaked redfish	<i>Sebastes mentella</i>	Peixe-vermelho-da-fundura	229
Beaked searsid	<i>Mentodus rostratus</i>	Celindra-de-tubo-bicuda	81
Bean's bigscale	<i>Scopelogadus beanii</i>	Peixe-rã-de-Bean	153
Bean's sawtooth eel	<i>Serrivomer beanii</i>	Cobra-couraça	68
Beardless codling	<i>Gadella imberbis</i>	Abrótea-imberbe	139
Bellotti's goby	<i>Gobius ater</i>	Caboz-de-Bellotti	179
Bennett's flyingfish	<i>Cheilopogon pinnatibarbatus</i>	Peixe-voador-de-Bennett	200
Benoit's lanternfish	<i>Hygophum benoiti</i>	Peixe-lanterna-de-Benoit	119
Benttooth bristlemouth	<i>Cyclothona acclinidens</i>	Boqueirão-de-dentes-curvos	86
Bermuda lantern fish	<i>Hygophum hygomii</i>	Peixe-lanterna-das-Bermudas	120
Bermuda sea chub	<i>Kyphosus sectatrix</i>	Preguiçosa-branca	238

English common name	Scientific name	Portuguese common name	page
Bertelsen's lanternfish	<i>Diaphus bertelseni</i>	Peixe-lanterna-de-Bertelsen	116
Bertin's one-jawed eel	<i>Monognathus bertini</i>	Enguia-da-fundura-de-Bertin	70
Bicoloured false moray	<i>Chlopsis bicolor</i>	Congrinho-bicolor	63
Bigbarb dragonfish	<i>Eustomias macronema</i>	Peixe-dragão-de-barbilho-grande	98
Bigelow's ray	<i>Rajella bigelowi</i>	Raia-de-Bigelow	48
Bigeye grunt	<i>Brachydeuterus auritus</i>	Roncador-colo-colo	262
Bigeye rockling	<i>Gaidropsarus macropthalmus</i>	Laibeque-do-Atlântico	145
Bigeye sand tiger shark	<i>Odontaspis noronhai</i>	Tubarão-de-Noronha	17
Bigeye scad	<i>Selar crumenophthalmus</i>	Charro-preto	254
Bigeye searsid	<i>Holtbyrnia macrops</i>	Celindra-de-tubo-de-olhuda	80
Bigeye smooth-head	<i>Bajacalifornia megalops</i>	Celindra-de-olho-grande	75
Bigeye squaretail	<i>Tetragonurus atlanticus</i>	Escolar-de-natura-olho-grande	173
Bigeye trevally	<i>Caranx sexfasciatus</i>	Charéu-de-natura	252
Bigeye tuna	<i>Thunnus obesus</i>	Atum-patudo	170
Bigfin pearleye	<i>Scopelarchus michaelisarsi</i>	Olho-pérola-de-Michael Sars	107
Bighead portholefish	<i>Margrethia obtusirostra</i>	Boqueirão-cabeçudo	88
Bighead searsid	<i>Holtbyrnia anomala</i>	Celindra-de-tubo-cabeçuda	80
Bignose shark	<i>Carcharhinus altimus</i>	Tubarão-baboso	24
Bigscale	<i>Poromitra megalops</i>	Peixe-rã-olhudo	152
Bigscale	<i>Scopeloberyx opistopterus</i>	Peixe-rã-delgado	152
Bigscale deepsea smelt	<i>Melanolagus bericoides</i>	Argentina-da-fundura-escamuda	84
Big-scale pomfret	<i>Taractichthys longipinnis</i>	Xaputa-galhuda	260
Big-scale sand smelt	<i>Atherina boyeri</i>	Peixe-rei-do-Mediterrâneo	197
Birdbeak dogfish	<i>Deania calcea</i>	Sapata	37
Blache's ooze ee	<i>Ilyophis blachei</i>	Enguia-da-fundura-de-Blache	58
Black cardinal fish	<i>Epigonous telescopus</i>	Olhudo	240
Black codling	<i>Physiculus dalwigkii</i>	Abrótea-de-Dalwigk	142
Black dogfish	<i>Centroscyllium fabricii</i>	Lixinha-negra	30
Black gemfish	<i>Nesiarchus nasutus</i>	Escolar-geribé	163
Black goby	<i>Gobius niger</i>	Caboz-negro	181
Black jack	<i>Caranx lugubris</i>	Xareu-enforcado	251
Black lantern fish	<i>Lampanyctus niger</i>	Peixe-lanterna-preto	122
Black pomfret	<i>Taractes rubescens</i>	Xaputa-negra	260
Black scabbardfish	<i>Aphanopus carbo</i>	Peixe-espada-preto	164
Black scorpionfish	<i>Scorpaena porcus</i>	Rascasso-de-pintas	232
Black seabream	<i>Spondylisoma cantharus</i>	Choupa	274
Black slimehead	<i>Hoplostethus cadenati</i>	Olho-de-vidro-preto	155
Black snaggletooth	<i>Astronesthes indicus</i>	Peixe-dragão-do-Índico	94
Black snake mackerel	<i>Nealotus tripes</i>	Escolar-coelho-de-natura	163
Black swallower	<i>Chiasmodon niger</i>	Peixe-glutão-negro	289
Black warrior	<i>Bathylaco nigricans</i>	Celindra-negra	76
Black wing flyingfish	<i>Hirundichthys rondeletii</i>	Peixe-voador-de-asa-negra	201
Blackbar hogfish	<i>Bodianus speciosus</i>	Bodião-cão	281
Blackbellied angler	<i>Lophius budegassa</i>	Tamboril-sovaco-preto	212
Blackbelly rosefish	<i>Helicolenus dactylopterus</i>	Cantarilho-legítimo	228
Blackchin guitarfish	<i>Glaucostegus cemiculus</i>	Viola-barba-negra	42
Black-faced blenny	<i>Tripterygion delaisi</i>	Caboz-de-três-dorsais-de-Delais	206
Blackfin sorcerer	<i>Nettastoma melanurum</i>	Cobra-de-cauda-preta	65
Blackfin spiderfish	<i>Bathypterois phenax</i>	Peixe-tripé-de-barbatana-negra	106
Blackfin tuna	<i>Thunnus atlanticus</i>	Atum-de-barbatana-negra	170
Blackfin wavyfish	<i>Scopelosaurus lepidus</i>	Peixe-vigia-de-barbatana-preta	107
Blackhead salmon	<i>Narcetes stomias</i>	Celindra-de-cabeça-negra	78
Blackmouth catshark	<i>Galeus melastomus</i>	Leitão	21
Blacksnout seasnail	<i>Paraliparis copei</i>	Peixe-lesma-de-focinho-preto	238
Blackspot picarel	<i>Spicara melanurus</i>	Trombeiro-malha-redonda	274

English common name	Scientific name	Portuguese common name	page
Blackspot seabream	<i>Pagellus bogaraveo</i>	Goraz	270
Blackspotted smooth-hound	<i>Mustelus punctulatus</i>	Cação-ponteado	24
Black-striped pipefish	<i>Syngnathus abaster</i>	Marinha-do-rio	176
Blacktail comber	<i>Serranus atricauda</i>	Serrano-de-rolo	245
Blacktip reef shark	<i>Carcharhinus limbatus</i>	Tubarão-de-pontas-negras	26
Blonde ray	<i>Raja brachyura</i>	Raia-ponteada	45
Blotched picarel	<i>Spicara maena</i>	Trombeiro-choupa	274
Blue antimora	<i>Antimora rostrata</i>	Mora-azul	139
Blue butterfish	<i>Stromateus fiatola</i>	Pampo-godinho	173
Blue cusk eel	<i>Brotulotaenia crassa</i>	Abadejo-grosso	158
Blue jack mackerel	<i>Trachurus picturatus</i>	Carapau-negrão	257
Blue ling	<i>Molva dypterygia</i>	Maruca-azul	146
Blue marlin	<i>Makaira nigricans</i>	Espadim-azul-do-Atlântico	185
Blue runner	<i>Caranx cryos</i>	Xareu-azul	251
Blue shark	<i>Prionace glauca</i>	Tintureira	28
Blue skate	<i>Dipturus batis</i>	Raia-oirega	43
Blue whiting	<i>Micromesistius poutassou</i>	Verdinho	143
Blueband dragonfish	<i>Photonectes mirabilis</i>	Peixe-dragão-maravilhoso	103
Bluefish	<i>Pomatomus saltatrix</i>	Anchova	248
Bluespotted seabream	<i>Evynnis ehrenbergii</i>	Pargo-ruço-de-Ebrenberg	269
Bluespotted seabream	<i>Pagrus caeruleostictus</i>	Pargo-ruço	272
Bluespotted triggerfish	<i>Balistes punctatus</i>	Cangulo-pintado	227
Blunthead puffer	<i>Sphoeroides pachygaster</i>	Peixe-bola-de-focinho-achatado	225
Bluntnose grenadier	<i>Nezumia longebarbata</i>	Largatixa-do-mar-de-focinho-curto	137
Bluntnose sixgill shark	<i>Hexanchus griseus</i>	Tubarão-albafar	16
Bluntnose sixgill shark	<i>Hexanchus nakamurai</i>	Tubarão-albafar-de-pontas-brancas	16
Bluntsnout smooth-head	<i>Xenodermichthys copei</i>	Celindra-de-focinho-curto	79
Boa dragonfish	<i>Stomias boa</i>	Peixe-dragão-jibóia	104
Boarfish	<i>Capros aper</i>	Pim-pim	297
Bobtail eel	<i>Cyema atrum</i>	Cobra-de-leme	69
Bogue	<i>Boops boops</i>	Boga-do-mar	263
Böhlke's one-jawed eel	<i>Monognathus boehlkei</i>	Enguia-da-fundura-de-Böhlke	70
Bonaparte's spiny eel	<i>Notacanthus bonaparte</i>	Enguia-espinhosa-de-Bonaparte	57
Bony-eared assfish	<i>Acanthonus armatus</i>	Abadejo-espinhososo	156
Boxlip mullet	<i>Oedalechilus labeo</i>	Tainha-sabão	204
Bramble shark	<i>Echinorhinus brucus</i>	Tubarão-prego	39
Brassy chub	<i>Kyphosus vaigiensis</i>	Preguiçosa-bronzeada	239
Brauer's dragonfish	<i>Photonectes braueri</i>	Peixe-dragão-de-Brauer	102
Bright glasseye	<i>Heteropriacanthus fulgens</i>	Fura-vasos-da-rocha-brilhante	248
Bright lanternfish	<i>Ctenoscopelus phengodes</i>	Peixe-lanterna-brilhante	116
Brill	<i>Scophthalmus rhombus</i>	Rodovalho	188
Bristlemouth	<i>Cyclotheta livida</i>	Boqueirão-lívido	87
Bristlemouth	<i>Gonostoma denudatum</i>	Boqueirão-do-Mediterrâneo	88
Bristly grenadier	<i>Trachonurus sulcatus</i>	Largatixa-do-mar-arrepiada	138
Brito's goby	<i>Chromogobius britoi</i>	Caboz-de-Brito	179
Broadnosed pipefish	<i>Syngnathus typhle</i>	Marinha-focinho-grosso	178
Bronze sawtooth eel	<i>Serrivomer lanceolatoides</i>	Cobra-de-dentes-curtos	69
Brown chub	<i>Kyphosus bigibbus</i>	Preguiçosa-cinzenta	238
Brown comber	<i>Serranus hepatus</i>	Serrano-ferreiro	246
Brown garden eel	<i>Heteroconger longissimus</i>	Enguia-de-jardim	67
Brown meagre	<i>Sciaena umbra</i>	Roncadeira-preta	276
Brown moray	<i>Gymnothorax unicolor</i>	Moreia-castanha	61
Brown ray	<i>Raja miraletus</i>	Raia-de-quatro-olhos	47
Brown stingray	<i>Bathyrajaa lata</i>	Uge-de-cardas	49
Brown wrasse	<i>Labrus merula</i>	Bodião-fusco	283

English common name	Scientific name	Portuguese common name	page
Brown-snout spookfish	<i>Dolichopteryx longipes</i>	Peixe-fantasma-de-barbatanas-longas	84
Bucchich's goby	<i>Gobius bucchichi</i>	Caboz-mosqueado	180
Bulbous dreamer	<i>Oneirodes eschrichtii</i>	Peixe-farol-de-Eschricht	218
Bull ray	<i>Aetomylaeus bovinus</i>	Ratão-bispo	51
Bullet tuna	<i>Auxis rochei</i>	Judeu	166
Bump-head sunfish	<i>Mola alexandrini</i>	Peixe-lua-de-Ranzani	222
Butterfly blenny	<i>Blennius ocellaris</i>	Marachomba-borboleta	207
Cadenat's scorpionfish	<i>Scorpaena loppei</i>	Rascasso-de-Loppé	231
California headlightfish	<i>Diaphus theta</i>	Peixe-lanterna-da-California	118
Canary damsel	<i>Similiparma lurida</i>	Castanheta-preta	279
Canary dentex	<i>Dentex canariensis</i>	Dentão-das-Canárias	264
Canary drum	<i>Umbrina canariensis</i>	Calafate-das-Canárias	276
Canary moray	<i>Gymnothorax bacalladoi</i>	Moreia-das-Canárias	60
Canary scorpionfish	<i>Scorpaena canariensis</i>	Rascasso-das-Canárias	230
Canary tonguesole	<i>Cynoglossus canariensis</i>	Língua-de-cão-das-Canárias	196
Can-opener smoothdream	<i>Chaenophryne longiceps</i>	Peixe-farol-cabeça-comprida	216
Cape snaggletooth	<i>Neonesthes capensis</i>	Peixe-dragão-do-Cabo	102
Cape Verde weever	<i>Trachinus pellegrini</i>	Peixe-aranha-de-Cabo-Verde	292
Carapine grenadier	<i>Coryphaenoides carapinus</i>	Lagartixa-do-mar-carapina	133
Cardinal fish	<i>Apogon imberbis</i>	Alcarraz	248
Challenger's spiny eel	<i>Polyacanthonotus challengerii</i>	Enguia-espinhosa-do-Challenger	57
Channel flounder	<i>Syacium micrum</i>	Carta-moira	188
Channeled rockfish	<i>Setarches guentheri</i>	Requeime-de-natura	229
Chaves' lanternfish	<i>Lampadена chavesi</i>	Peixe-lanterna-de-Chaves	120
Chemnitz's spiny eel	<i>Notacanthus chemnitzii</i>	Enguia-espinhosa-de-Chemnitz	57
Chilean devil ray	<i>Mobula tarapacana</i>	Manta-cornuda	52
Chubby clingfish	<i>Apletodon pellegrini</i>	Sugador-bochechudo	205
Cline	<i>Clinitrichus argentatus</i>	Marachomba-prateada	207
Cobia	<i>Rachycentron canadum</i>	Fogueteiro-galego	249
Cocco's lanternfish	<i>Gonichthysocco</i>	Peixe-lanterna-de-Cocco	119
Codling	<i>Laemonema yarrellii</i>	Abrótea-de-Yarrell	140
Comber	<i>Serranus cabrilla</i>	Serrano-alecrim	246
Common Atlantic grenadier	<i>Nezumia aequalis</i>	Largatixa-do-mar-lisa	136
Common dentex	<i>Dentex dentex</i>	Capatão-legítimo	264
Common dolphinfish	<i>Coryphaena hippurus</i>	Doirado	258
Common eagle ray	<i>Myliobatis aquila</i>	Ratão-água	51
Common fangtooth	<i>Anoplogaster cornuta</i>	Peixe-ogre	156
Common goby	<i>Pomatoschistus microps</i>	Caboz-comum	183
Common guitarfish	<i>Rhinobatos rhinobatos</i>	Viola	41
Common mora	<i>Mora moro</i>	Mora	141
Common pandora	<i>Pagellus erythrinus</i>	Bica	271
Common sawfish	<i>Pristis pristis</i>	Espadarte-serra	42
Common sole	<i>Solea solea</i>	Linguado-legítimo	195
Common stingray	<i>Dasyatis pastinaca</i>	Uge	49
Common torpedo	<i>Torpedo torpedo</i>	Tremelga-de-olhos	41
Common two-banded seabream	<i>Diplodus vulgaris</i>	Sargo-safia	268
Connemarra clingfish	<i>Lepadogaster candolii</i>	Sugador-de-De Candolle	205
Constance deepwater cardinalfish	<i>Epigonus constanciae</i>	Olhudo-de-espinho	239
Constellationfish	<i>Valenciennellus tripunctulatus</i>	Pai-velho-de-três-pontos	91
Cookie cutter shark	<i>Isistius brasiliensis</i>	Tubarão-luminoso	30
Copper shark	<i>Carcharhinus brachyurus</i>	Tubarão-cobre	24
Corkwing wrasse	<i>Syphodus melops</i>	Bodião-vulgar	285
Cornetfish	<i>Fistularia tabacaria</i>	Corneta-malhada	174
Cornish blackfish	<i>Schedophilus medusophagus</i>	Liro-mole	172
Cornish sucker	<i>Lepadogaster purpurea</i>	Sugador-púrpura	205

English common name	Scientific name	Portuguese common name	page
Couch's goby	<i>Gobius couchi</i>	Caboz-de-Couch	180
Crested bigscale	<i>Poromitra crassiceps</i>	Peixe-rã-de-crista	152
Crested oarfish	<i>Lophotus lacepede</i>	Peixe-remo-de-crista	127
Crevalle jack	<i>Caranx hippos</i>	Xareu-macoa	251
Crystal goby	<i>Crystallagogibius linearis</i>	Caboz-de-cristal	179
Cubera snapper	<i>Lutjanus cyanopterus</i>	Luciano-de-ventre-vermelho	261
Cuckoo ray	<i>Leucoraja naevus</i>	Raia-de-dois-olhos	45
Cuckoo wrasse	<i>Labrus mixtus</i>	Bodião-canário	283
Curled picarel	<i>Centracanthus cirrus</i>	Boqueirão	264
Daggertooth	<i>Anotopterus pharao</i>	Barracudina-dente-de-adaga	112
Damselfish	<i>Chromis chromis</i>	Castanheta-báia	278
Dana viperfish	<i>Chauliodus danae</i>	Peixe-dragão-do-Dana	96
Dark cusk	<i>Brotulaenia nigra</i>	Abadejo-negro	158
Dark moray	<i>Gymnothorax afer</i>	Moreão-escuro	60
Dark searsid	<i>Mentodus facilis</i>	Celindra-de-tubo-escura	81
Darwin's slimehead	<i>Gephyroberyx darwini</i>	Olho-de-vidro-imperatriz	154
De Buen's goby	<i>Buenia affinis</i>	Caboz-de-De Buen	178
Dealfish	<i>Trachipterus arcticus</i>	Peixe-tábua	128
Deep water sole	<i>Bathysolea profundicola</i>	Linguado-da-fundura	191
Deepbody boarfish	<i>Antigonion capros</i>	Periquito	296
Deepsea blackdevil	<i>Melanocetus murrayi</i>	Diabo-marinho-de-Murray	215
Deep-sea lizardfish	<i>Bathysaurus ferox</i>	Peixe-lagarto-da-fundura	109
Deep-water arrowtooth eel	<i>Histiobranchus bathybius</i>	Enguia-da-fundura	58
Deep-water catshark	<i>Apristurus profundorum</i>	Tubarão-gato-abissal	21
Deepwater grenadier	<i>Coryphaenoides profundiculus</i>	Lagartixa-do-mar-da-fundura	134
Deepwater lanternfish	<i>Taaningichthys bathyphilus</i>	Peixe-lanterna-da-fundura	126
Deep-water ray	<i>Rajella bathyphila</i>	Raia-da-fundura	48
Deep-water red snapper	<i>Etelis carbunculus</i>	Vermelho-salmão	261
Devil fish	<i>Mobula mobular</i>	Manta-do-Mediterrâneo	52
Diaphanous hatchet fish	<i>Sternopyx diaphana</i>	Pai-velho-diáfano	91
Diminutive goby	<i>Lebetus scorpioides</i>	Caboz-anão	182
Diogenes lanternfish	<i>Diogenichthys laternatus</i>	Peixe-lanterna-de-Diogenes	119
Doflein's lantern fish	<i>Lobianchia dofleini</i>	Peixe-lanterna-de-Doflein	123
Dogtooth grouper	<i>Epinephelus caninus</i>	Mero-gigante	243
Dogtooth lampfish	<i>Ceratoscopelus townsendi</i>	Peixe-lanterna-de-Townsend	115
Dotback lanternfish	<i>Lampanyctus photonotus</i>	Peixe-lanterna-lobisomem	123
Doublethread grenadier	<i>Gadomus arcuatus</i>	Lagartixa-do-mar-filamentosa	135
Dragonet	<i>Callionymus lyra</i>	Peixe-pau-lira	293
Driftfish	<i>Cubiceps gracilis</i>	Tirone	172
Duckbill barracudina	<i>Magnisudis atlantica</i>	Barracudina-bico-de-pato	110
Duckbill oceanic eel	<i>Nessorhamphus ingolfianus</i>	Enguia-bico-de-pato	63
Dumeril's lanternfish	<i>Diaphus dumerilii</i>	Peixe-lanterna-de-Duméril	117
Dungat grouper	<i>Epinephelus goreensis</i>	Mero-amarelo	243
Dusky flounder	<i>Syacium papillosum</i>	Carta-sombria	188
Dusky grouper	<i>Epinephelus marginatus</i>	Mero-legítimo	244
Dusky lanternfish	<i>Lampanyctus ater</i>	Peixe-lanterna-escuro	121
Dusky shark	<i>Carcharhinus obscurus</i>	Tubarão-faquetá	27
Dusky smooth-hound	<i>Mustelus canis</i>	Cação-cachorro	23
Dusky smooth-hound	<i>Mustelus mustelus</i>	Cação-liso	24
Dwarf deepwater cardinalfish	<i>Microichthys coccoi</i>	Olhudo-anão	240
Dwarf dory	<i>Zenion hololepis</i>	Galito-rosado	130
Dwarf saury	<i>Scomberesox simulans</i>	Agulhão-anão	198
East Atlantic Opah	<i>Lampris lauta</i>	Peixe-cravo-elegante	127
Eckström's topknot	<i>Zeugopterus regius</i>	Bruxa	188
Electric lantern fish	<i>Electrona risso</i>	Peixe-lanterna-faísca	119

English common name	Scientific name	Portuguese common name	page
Electric ray	<i>Tetronarce nobiliana</i>	Tremelga-negra	40
Elongate frostfish	<i>Benthodesmus elongatus</i>	Espada-de-má-água-comprido	164
Elongate tonguesole	<i>Syphurus ligulatus</i>	Língua-comprida	196
Elongated bristlemouth fish	<i>Sigmops elongatus</i>	Boqueirão-dentudo	89
Emerald wrasse	<i>Syphodus trutta</i>	Bodião-mosqueado	286
Escolar	<i>Lepidocybium flavobrunneum</i>	Escolar-preto	162
European anchovy	<i>Engraulis encrasiculus</i>	Biqueirão	74
European barracuda	<i>Sphyraena sphyraena</i>	Bicuda-europeia	295
European conger	<i>Conger conger</i>	Congro	66
European eel	<i>Anguilla anguilla</i>	Enguia	69
European finless eel	<i>Apterichtus caecus</i>	Congrinho-da-areaia	64
European flounder	<i>Platichthys flesus</i>	Solha-das-pedras	191
European hake	<i>Merluccius merluccius</i>	Pescada-branca	148
European pilchard	<i>Sardina pilchardus</i>	Sardinha	72
European plaice	<i>Pleuronectes platessa</i>	Solha	191
European river lamprey	<i>Lampetra fluviatilis</i>	Lampreia-do-rio	15
European seabass	<i>Dicentrarchus labrax</i>	Robalo-legítimo	241
European sprat	<i>Sprattus sprattus</i>	Espadilha	74
Eyebrow bigscale	<i>Melamphaes longivelis</i>	Peixe-rã-sobrolho	150
Facciola's sorcerer	<i>Facciolella oxyrhynchus</i>	Cobra-cabeça-de-pato	65
False catshark	<i>Pseudotriakis microdon</i>	Tubarão-mona	22
False cusk	<i>Parabrotula plagiophthalma</i>	Brótula-falsa-dos-Açores	161
False herring	<i>Harengula clupeola</i>	Sardinha-escamuda	72
False scad	<i>Caranx rhonchus</i>	Xareu-amarelo	251
Fanfare lanternfish	<i>Notoscopelus bolini</i>	Peixe-lanterna-de-Bolin	125
Fanfin angler	<i>Caulophryne jordani</i>	Peixe-pescador-de-Jordan	214
Fanfin smooth-head	<i>Asquamiceps velaris</i>	Celindra-de-leque	75
Fanfish	<i>Pteraclis carolinus</i>	Freira	259
Fangtooth moray	<i>Enchelycore anatina</i>	Moreia-serpente	60
Fangtooth smooth-head	<i>Bathyprion danae</i>	Celindra-de-Dana	76
Festive lanternfish	<i>Lampanyctus festivus</i>	Peixe-lanterna-divertido	122
Festive ribbonfish	<i>Eutaeniophorus festivus</i>	Peixe-serpentina	154
Fierce pike smelt	<i>Sudis atrox</i>	Barracudina-feroz	111
Fiery lanternfish	<i>Bolinichthys pyrsobolus</i>	Peixe-lanterna-arco-íris	114
Finetooth shark	<i>Carcharhinus isodon</i>	Tubarão-dente-liso	25
Fiolenti's smooth-head	<i>Conocara fiolenti</i>	Celindra-de-Fiolenti	76
Firebrow lanternfish	<i>Diaphus adenomus</i>	Peixe-lanterna-de-Gilbert	116
Fivebeard rockling	<i>Ciliata mustela</i>	Laibeque-de-cinco-barbillhos	144
Five-spotted wrasse	<i>Syphodus roissali</i>	Bodião-manchado	285
Flabby lanternfish	<i>Diaphus subtilis</i>	Peixe-lanterna-flácido	118
Flapper Skate	<i>Dipturus intermedius</i>	Raia-delgada	43
Flatface lanternfish	<i>Diaphus perspicillatus</i>	Peixe-lanterna-luneta	118
Flathead grey mullet	<i>Mugil cephalus</i>	Tainha-olhalvo	204
Fleshyfin dragonfish	<i>Photonectes parvimanus</i>	Peixe-dragão-barbatana-curta	103
Florida pompano	<i>Trachinotus carolinus</i>	Sereia-da-Florida	256
Flying gurnard	<i>Dactylopterus volitans</i>	Cabrinha-de-leque	178
Forkbeard	<i>Phycis phycis</i>	Abrótea-da-costa	147
Foureyed sole	<i>Microchirus ocellatus</i>	Azevia-de-malhas	193
Four-spot megrim	<i>Lepidorhombus boscii</i>	Areeiro-de-quatro-manchas	187
Four-spotted goby	<i>Deltentosteus quadrifasciatus</i>	Caboz-de-quatro-manchas	179
Freckled driftfish	<i>Psenes cyanophrys</i>	Peixe-pastor-amarelado	173
French angelfish	<i>Pomacanthus paru</i>	Lebre-preta	278
Freshwater blenny	<i>Salaria fluviatilis</i>	Marachomba-de-água-doce	211
Fries's goby	<i>Lesueurigobius friesii</i>	Caboz-de-escama-grande	182
Frigate tuna	<i>Auxis thazard</i>	Judeu-liso	166

English common name	Scientific name	Portuguese common name	page
Frilled shark	<i>Chlamydoselachus anguineus</i>	Tubarão-cobra	16
Furry whiptail	<i>Trachonurus villosus</i>	Lagartixa-do-mar-peluda	138
Fusca drum	<i>Umbrina ronchus</i>	Calafate-fosco	276
Gadella	<i>Gadella maraldi</i>	Abrótea-de-natura	139
Galapagos shark	<i>Carcharhinus galapagensis</i>	Tubarão-dos-Galápagos	25
Garfish	<i>Belone acus</i>	Agulha-do-Mediterrâneo	198
Garfish	<i>Belone belone</i>	Agulha	199
Gerrick	<i>Cyclothona braueri</i>	Boqueirão-de-Brauer	86
Gauss' lanternfish	<i>Lepidophanes gaussii</i>	Peixe-lanterna-de-Gauss	123
Gelatinous blindfish	<i>Aphyonus gelatinosus</i>	Abadejo-cego-gelatinoso	160
Gemellar's lanternfish	<i>Lobianchia gemellarii</i>	Peixe-lanterna-de-Gemellar	124
Ghost catshark	<i>Apristurus manis</i>	Tubarão-gato-fantasma	20
Ghostly grenadier	<i>Coryphaenoides leptolepis</i>	Lagartixa-do-mar-pálida	134
Giant goby	<i>Gobius cobitis</i>	Caboz-cabeçudo	180
Giant lepidion	<i>Lepidion schmidti</i>	Abrótea-de-Schmidt	141
Giant leptocephalus	<i>Leptocephalus giganteus</i>	sem nome comum	57
Giant manta	<i>Mobula birostris</i>	Manta-gigante	52
Gill's flabby whalefish	<i>Cetomimus gilli</i>	Peixe-baleia-de-Gill	153
Gilthead seabream	<i>Sparus aurata</i>	Dourada	273
Glacier lantern fish	<i>Benthosema glaciale</i>	Peixe-lanterna-glacial	113
Glasseye	<i>Heteropriacanthus cruentatus</i>	Fura-vasos-da-rocha	247
Glasshead barreleye	<i>Rhynchohyalus natalensis</i>	Peixe-sola-cabeça-de-vídeo	85
Glasshead grenadier	<i>Hymenocephalus italicus</i>	Lagartixa-do-mar-prateada	135
Globehead grenadier	<i>Cetonurus globiceps</i>	Lagartixa-do-mar-de-cabeça-redonda	132
Goblin shark	<i>Mitsukurina owstoni</i>	Tubarão-duede	17
Goiter blacksmelt	<i>Bathylagus euryops</i>	Argentina-da-fundura-negra	84
Goldblotch grouper	<i>Epinephelus costae</i>	Garoupa-dourada	243
Golden goby	<i>Gobius auratus</i>	Caboz-dourado	180
Golden goby	<i>Gobius xanthocephalus</i>	Caboz-de-cabeça-dourada	182
Golden grey mullet	<i>Liza aurata</i>	Tainha-garrento	203
Golden redfish	<i>Sebastes norvegicus</i>	Peixe-vermelho	229
Goldentail moray	<i>Gymnothorax miliaris</i>	Moreia-dourada	61
Goldsinny-wrasse	<i>Ctenolabrus rupestris</i>	Bodião-rupestre	282
Goldspot goby	<i>Gnatholepis thompsoni</i>	Caboz-de-pinta-dourada	179
Gorean snapper	<i>Lutjanus goreensis</i>	Luciano-da-Goreia	261
Graceful grenadier	<i>Hymenogadus gracilis</i>	Lagartixa-do-mar-elegante	135
Graceful shark	<i>Carcharhinus brevipinna</i>	Tubarão-tecelão	25
Grass goby	<i>Gobius ophiocephalus</i>	Caboz-cabeça-de-cobra	181
Great barracuda	<i>Sphyraena barracuda</i>	Bicuda-gigante	295
Great hammerhead	<i>Sphyrna mokarran</i>	Tubarão-martelo-gigante	29
Great lanternshark	<i>Etmopterus princeps</i>	Lixinha-da-fundura-grada	31
Great sandeel	<i>Hyperoplus lanceolatus</i>	Galeota-maior	291
Great white shark	<i>Carcharodon carcharias</i>	Tubarão-branco	18
Greater amberjack	<i>Seriola dumerili</i>	Charuteiro-catarino	254
Greater forkbeard	<i>Phycis blennoides</i>	Abrótea-do-alto	146
Greater pipefish	<i>Syngnathus acus</i>	Marinha-comum	177
Greater sand-eel	<i>Hyperoplus immaculatus</i>	Galeota-maior-de-Corbin	291
Greater weever	<i>Trachinus draco</i>	Peixe-aranha-maior	292
Green wrasse	<i>Labrus viridis</i>	Bodião-torto	283
Greenland argentine	<i>Nansenia groenlandica</i>	Argentina-da-Groenlândia	83
Greenland manefish	<i>Caristius fasciatus</i>	Peixe-crina-da-Gronelândia	260
Greenland shark	<i>Somniosus microcephalus</i>	Tubarão-da-Gronelândia	34
Grenadier smooth-head	<i>Leptoderma macrops</i>	Celindra-olhuda	78
Grey cutthroat eel	<i>Synaphobranchus affinis</i>	Moreão-do-golfo	59
Grey gurnard	<i>Eutrigla gurnardus</i>	Cabra-morena	235

English common name	Scientific name	Portuguese common name	page
Grey triggerfish	<i>Balistes capriscus</i>	Cangulo-cinzento	227
Grey wrasse	<i>Syphodus cinereus</i>	Bodião-cinzento	284
Grey's deepsea smelt	<i>Bathylagichthys greyae</i>	Argentina-da-fundura-de-Grey	84
Guanchane barracuda	<i>Sphyraena guachancho</i>	Bicuda-africana	295
Guenther's codling	<i>Lepidion guentheri</i>	Abrótea-de-Günther	141
Guillet's goby	<i>Lebetus guilleti</i>	Caboz-de-Guillet	182
Guinean amberjack	<i>Seriola carpenteri</i>	Charuteiro-da-Guiné	254
Guinean burrfish	<i>Chilomycterus spinosus mauretanicus</i>	Porco-espinho-da-Guiné	223
Guinean conger	<i>Paraconger notialis</i>	Congro-da-Guiné	67
Guinean grunt	<i>Parapristipoma humile</i>	Roncador-canela	262
Guinean pike conger	<i>Cynoponticus ferox</i>	Congro-branco-africano	65
Guinean pufferfish	<i>Sphoeroides marmoratus</i>	Peixe-bola-da-Guiné	225
Gulper shark	<i>Centrophorus granulosus</i>	Barroso	35
Günther's halosaur	<i>Halosaurus guentheri</i>	Peixe-lagarto-de-Guenther	56
Günther's lanternfish	<i>Lepidophanes guentheri</i>	Peixe-lanterna-de-Günther	123
Günther's grenadier	<i>Coryphaenoides guentheri</i>	Lagartixa-do-mar-de-Günther	133
Hagfish	<i>Myxine glutinosa</i>	Enguia-de-casulo	15
Haifa grouper	<i>Hyporthodus haifensis</i>	Garoupa-de-Haifa	244
Hairfin smooth-head	<i>Talismania homoptera</i>	Celindra-da-Mauritânia	79
Hairy blenny	<i>Labrisomus nuchipinnis</i>	Marachomba-peluda	206
Hairyfish	<i>Mirapinna esau</i>	Peixe-baleia-Esaú	154
Hairytail swallower eel	<i>Saccopharynx thalassa</i>	Peixe-pelícano-de-cauda-peluda	71
Half-naked hatchetfish	<i>Argyropelecus hemigymnus</i>	Pai-velho-manchado	90
Hatchetfish	<i>Argyropelecus gigas</i>	Pai-velho-gigante	90
Hawaiian halosaurid fish	<i>Aldrovandia phalacra</i>	Peixe-lagarto-dos-Açores	56
Headlight angler	<i>Linophryne indica</i>	Peixe-sestro-do-Índico	221
Headlight lanternfish	<i>Diaphus effulgens</i>	Peixe-lanterna-de-focinho-luminoso	117
Herring's one-jawed eel	<i>Monognathus herringi</i>	Enguia-da-fundura-de-Herring	70
Hidden bristlemouth	<i>Cyclothona obscura</i>	Boqueirão-sombrio	87
Highfin lizardfish	<i>Bathysaurus mollis</i>	Peixe-lagarto-da-costa	109
Highlight hatchetfish	<i>Sternopyx pseudoboscara</i>	Pai-velho-oblíquo	91
Highseas lampfish	<i>Triphoturus nigrescens</i>	Peixe-lanterna-do-alto	127
Hollowsnout grenadier	<i>Coelorinchus caelorhincus</i>	Lagartixa-do-mar-de-focinho-côncavo	132
Honeycomb moray	<i>Muraena melanotis</i>	Moreia-pintada	63
Hooked redmouth whalefish	<i>Rondeletia bicolor</i>	Peixe-baleia-bicolor	153
Horned lantern fish	<i>Centrophryne spinulosa</i>	Peixe-farol-córneo	219
Horned lanternfish	<i>Diaphus splendidus</i>	Peixe-lanterna-cornudo	118
Horse-eye jack	<i>Caranx latus</i>	Xareu-de-olho-papudo	251
Hound needlefish	<i>Tylosurus crocodilus</i>	Agulheta-crocodilo	200
Hubb's pearleye	<i>Rosenblattichthys hubbsi</i>	Olho-pérola-de-Hubb	107
Humpback anglerfish	<i>Melanocetus johnsonii</i>	Diabo-marinho-de-Johnson	215
Iberian pygmy skate	<i>Neoraja iberica</i>	Raia-pigmeia-ibérica	45
Imperial blackfish	<i>Schedophilus ovalis</i>	Liro-imperial	172
Imperial scaldfish	<i>Arnoglossus imperialis</i>	Carta-imperial	189
Indian sabertooth	<i>Evermannella melanoderma</i>	Dente-de-sabre-ponteado	112
Indo-Pacific sailfish	<i>Istiophorus platypterus</i>	Veleiro-do-Indo-Pacífico	185
Inshore lizardfish	<i>Synodus foetens</i>	Peixe-lagarto-da-praia	108
Intermediate scabbardfish	<i>Aphanopus intermedius</i>	Peixe-espada-intermédio	164
Intricate lanternfish	<i>Lampanyctus intricarius</i>	Peixe-lanterna-complexo	122
Irish oreo	<i>Neocyttus helgae</i>	Galo-irlandês	129
Island boxfish	<i>Acanthostracion notacanthus</i>	Peixe-cofre	226
Island grouper	<i>Mycteroperca fusca</i>	Garoupa-insular	244
Japanese gissu	<i>Pterothriussus gissu</i>	Gissu-japonês	55
Javelin spookfish	<i>Bathylychnops exilis</i>	Peixe-fantasma-dardo	84
Jeffrey's goby	<i>Buenia jeffreysii</i>	Caboz-de-Jeffrey	179

English common name	Scientific name	Portuguese common name	page
Jellynose	<i>Guentherus altivelis</i>	Peixe-de-vela-alta	104
Jewel lanternfish	<i>Lampanyctus crocodilus</i>	Peixe-lanterna-crocodilo	121
John dory	<i>Zeus faber</i>	Galo-negro	130
Johnson's halosaur	<i>Halosaurus johnsonianus</i>	Peixe-lagarto-de-Johnson	56
Kaup's arrowtooth eel	<i>Synaphobranchus kaupii</i>	Moreão-de-natura	59
Keeltail needlefish	<i>Platybelone argalus</i>	Aguilha-de-quilha	199
King of herrings	<i>Regalecus glesne</i>	Rei-dos-Arenques	128
Kitefin shark	<i>Dalatias licha</i>	Gata	29
Klein's sole	<i>Synapturichthys kleinii</i>	Linguado-turco	195
Kner's goby	<i>Pomatoschistus knerii</i>	Caboz-de-Kner	183
Knifetooth dogfish	<i>Scymnodon ringens</i>	Arreganhada	33
Koefoed's searsid	<i>Searsia koefoedi</i>	Celindra-de-tubo-de-Koefoed	82
Koefoed's smooth-head	<i>Bathytroctes macrolepis</i>	Celindra-escamuda	76
Krefft's dreamer	<i>Oneirodes krefftii</i>	Peixe-farol-de-Krefft	218
Krefft's ray	<i>Malacoraja krefftii</i>	Raiá-de-Krefft	45
Krefft's smooth-head	<i>Herwigia krefftii</i>	Celindra-de-Krefft	77
Krøyer's lanternfish	<i>Notoscopelus kroyeri</i>	Peixe-lanterna-de-Krøyer	126
Krøyer's deep-sea angler fish	<i>Ceratias holboelli</i>	Peixe-farol-de-Holbøll	219
Lancet fish	<i>Paralepis coregonoides</i>	Barracudina-lanceta	111
Lanternfish	<i>Lampanyctus lineatus</i>	Peixe-lanterna-riscado	122
Large-eye dentex	<i>Dentex macrophthalmus</i>	Cachicho	265
Large-eyed rabbitfish	<i>Hydrolagus mirabilis</i>	Ratazana-olhuda	54
Largefin lanternfish	<i>Hygophum macrochir</i>	Peixe-lanterna-de-barbatanas-grandes	120
Largehead hairtail	<i>Trichiurus lepturus</i>	Lírio	165
Large-scale lantern fish	<i>Symbolophorus veranyi</i>	Peixe-lanterna-de-Vérany	126
Large-scaled gurnard	<i>Lepidotrigla cavillone</i>	Ruivo	236
Large-scaled lantern fish	<i>Neoscopelus macrolepidotus</i>	Peixe-lanterna-escamudo	113
Targetooth cookiecutter shark	<i>Isistius plutodus</i>	Tubarão-dentão	30
Leaden worm eel	<i>Myrophis plumbeus</i>	Cobra-chumbo	64
Leafscale gulper shark	<i>Centrophorus squamosus</i>	Lixa	36
Leaping mullet	<i>Chelon saliens</i>	Tainha-de-salto	203
Leerfish	<i>Lichia amia</i>	Palombeta	253
Legless searsid	<i>Platytroctes apus</i>	Celindra-de-tubo-curva	81
Lemon shark	<i>Negaprion brevirostris</i>	Tubarão-limão	28
Lemon sole	<i>Microstomus kitt</i>	Solha-limão	191
Leopard eel	<i>Myrichthys pardalis</i>	Cobra-leopardo	64
Leopard-spotted goby	<i>Thorogobius ephippiatus</i>	Caboz-leopardo	184
Lesser amberjack	<i>Seriola fasciata</i>	Charuteiro-listado	255
Lesser weever	<i>Echiichthys vipera</i>	Peixe-aranha-menor	291
Lesueur's goby	<i>Lesueuriogobius suerii</i>	Caboz-de-Lesueur	183
Lined Seahorse	<i>Hippocampus erectus</i>	Cavalo-marinho-estriado	175
Ling	<i>Molva molva</i>	Maruca	146
Little gulper shark	<i>Centrophorus cf. zeehaani</i>	Quelme-do-Zeehaan	37
Little gulper shark	<i>Centrophorus uyato</i>	Quelme	36
Little sleeper shark	<i>Somniosus rostratus</i>	Pailona	34
Little tunny	<i>Euthynnus alletteratus</i>	Merma	167
Live sharksucker	<i>Echeneis naucrates</i>	Pegador-listado	249
Lobed dreamer	<i>Dolopichthys pullatus</i>	Peixe-farol-lobado	217
Long snouted lancetfish	<i>Alepisaurus ferox</i>	Lírio-ferro-de-focinho-comprido	113
Longbarb dragonfish	<i>Flagellostomias boureei</i>	Peixe-dragão-dos-Açores	100
Longbarb scaly dragonfish	<i>Stomias longibarbatus</i>	Peixe-dragão-escamudo	104
Longbill spearfish	<i>Tetrapturus pfluegeri</i>	Espadim-bicudo	186
Longfin dragonfish	<i>Bathophilus longipinnis</i>	Peixe-dragão-de-barbatana-longa	95
Longfin escolar	<i>Scombrolabrax heterolepis</i>	Escolar-de-peitoral-longa	171
Longfin gurnard	<i>Chelidonichthys obscurus</i>	Cabra-de-bandeira	235

English common name	Scientific name	Portuguese common name	page
Longfin mako	<i>Isurus oxyrinchus</i>	Tubarão-anequim	18
Longfin mako	<i>Isurus paucus</i>	Tubarão-anequim-de-gadanha	18
Longfin smooth-head	<i>Conocara macropterum</i>	Celindra-de-barbatana-comprida	76
Longfin yellowtail	<i>Seriola rivoliana</i>	Charuteiro-falcado	255
Longfinned bullseye	<i>Cookeolus japonicus</i>	Fura-vasos-alfonsim	247
Longjaw bigscale	<i>Scopeloberyx robustus</i>	Peixe-rã-robusto	152
Longlure frogfish	<i>Antennarius multiocellatus</i>	Peixe-sapo-multiocelado	213
Longneck croaker	<i>Pseudotolithus typus</i>	Rainha-branca	276
Longnose spurdog	<i>Squalus blainvillei</i>	Galhudo	38
Longnosed skate	<i>Dipturus oxyrinchus</i>	Raia-bicuda	43
Longray fangjaw	<i>Bonapartia pedaliota</i>	Boqueirão-de-raio-longo	86
Longsnout blacksmelt	<i>Dolicholagus longirostris</i>	Argentina-da-fundura-de-focinho-comprido	84
Longsnout dogfish	<i>Deania profundorum</i>	Sapata-flecha	38
Long-snouted seahorse	<i>Hippocampus guttulatus</i>	Cavalo-marinho-de-focinho-comprido	175
Longspine snipefish	<i>Macroramphosus scolopax</i>	Trombeteiro	174
Longspined bullhead	<i>Taurulus bubalis</i>	Escorpião-roco	237
Longspined porcupinefish	<i>Diodon holocanthus</i>	Peixe-ouriço-de-crista	223
Long-tooth conger	<i>Bathyuroconger vicinus</i>	Congro-castanho-dentão	66
Loosejaw	<i>Aristostomias tittmanni</i>	Peixe-dragão-de-Tittmann	93
Loosejaw	<i>Photostomias guernei</i>	Peixe-dragão-de-Guerne	103
Loosescale smooth-head	<i>Einara macrolepis</i>	Celindra-de-escama-caduca	77
Lowfin gulper shark	<i>Centrophorus lusitanicus</i>	Tubarão-lusitano	35
Lozano's goby	<i>Pomatoschistus lozanoi</i>	Caboz-de-Lozano	183
Luetken's lanternfish	<i>Diaphus luetkeni</i>	Peixe-lanterna-de-Lütken	117
Luminous lanternfish	<i>Lampadena luminosa</i>	Peixe-lanterna-luminoso	121
Lumpfish	<i>Cyclopterus lumpus</i>	Peixe-lapa	237
Lusitanian cownose ray	<i>Rhinoptera marginata</i>	Gavião-do-mar	51
Lusitanian sole	<i>Microchirus boscanion</i>	Azevia-lusitanica	193
Lusitanian toadfish	<i>Halobatrachus didactylus</i>	Charroco	162
Luvar	<i>Luvarus imperialis</i>	Boquinho	295
Macaronesian sharpnose-puffer	<i>Canthigaster capistrata</i>	Peixe-bola-da-Macaronésia	224
Macaronesian tonguesole	<i>Syphurus insularis</i>	Língua-da-Maraconésia	196
MacDonald's lanternfish	<i>Lampanyctus macdonaldi</i>	Peixe-lanterna-de-MacDonald	122
Mackerel scad	<i>Decapterus macarellus</i>	Charro-olho-largo	252
Madeira butterfly ray	<i>Gymnura hirundo</i>	Uge-da-Madeira	50
Madeira catshark	<i>Apristurus laurussonii</i>	Tubarão-gato-da-Madeira	20
Madeira lantern fish	<i>Ceratoscopelus maderensis</i>	Peixe-lanterna-da-Madeira	115
Madeira rockfish	<i>Scorpaena maderensis</i>	Rascasso-da-Madeira	231
Madeiran grenadier	<i>Coryphaenoides rutilus</i>	Lagartixa-do-mar-cabeçuda	134
Madeiran ray	<i>Raja maderensis</i>	Raia-da-Madeira	46
Madeiran sardinella	<i>Sardinella maderensis</i>	Sardinela-da-Madeira	73
Madeiran smooth-head	<i>Rouleina maderensis</i>	Celindra-da-Madeira	79
Manefish	<i>Caristius macropus</i>	Peixe-crina-barbatana-comprida	260
Man-of-war fish	<i>Nameus gronovii</i>	Pequeno-argonauta	173
Many pore conger	<i>Paraconger macrops</i>	Congro-de-natura	67
Manytooth conger	<i>Conger triporiceps</i>	Congro-dentudo	66
Marbled electric ray	<i>Torpedo marmorata</i>	Tremelga-marmoreada	41
Marbled goby	<i>Pomatoschistus marmoratus</i>	Caboz-marmoreado	183
Margined flyingfish	<i>Cheilopogon cyanopterus</i>	Peixe-voador-barbudo	200
Marion's spiderfish	<i>Bathytyphlops marionae</i>	Peixe-tripé-de-Marion	106
Marlin sucker	<i>Remora osteochir</i>	Pegador-dos-espádarpes	250
Marlin-spoke grenadier	<i>Nezumia bairdii</i>	Largatixa-do-mar-de-Baird	137
Masked assfish	<i>Bassozetus nielseni</i>	Abadejo-de-Nielsen	157
Maul's searsid	<i>Maulisia mauli</i>	Celindra-de-tubo-de-Maul	80
Meagre	<i>Argyrosomus regius</i>	Corvina-legítima	275

English common name	Scientific name	Portuguese common name	page
Mediterranean codling	<i>Lepidion lepidion</i>	Abrótea-do-Mediterrâneo	141
Mediterranean dealfish	<i>Trachipterus trachypterus</i>	Peixe-tábua-do-Mediterrâneo	128
Mediterranean flyingfish	<i>Cheilopogon heterurus</i>	Peixe-voador-do-Mediterrâneo	200
Mediterranean grenadier	<i>Coryphaenoides mediterraneus</i>	Lagartixa-do-mar-do-Mediterrâneo	134
Mediterranean horse mackerel	<i>Trachurus mediterraneus</i>	Carapau-do-Mediterrâneo	256
Mediterranean moray	<i>Muraena helena</i>	Moreia	62
Mediterranean painted goby	<i>Pomatoschistus pictus</i>	Caboz-de-manchas	184
Mediterranean sand eel	<i>Gymnammodytes cicerelus</i>	Galeota-da-areia	291
Mediterranean sand smelt	<i>Atherina hepsetus</i>	Peixe-rei-do-alto	197
Mediterranean scaldfish	<i>Arnoglossus laterna</i>	Carta-do-Mediterrâneo	189
Mediterranean slimehead	<i>Hoplostethus mediterraneus</i>	Olho-de-vidro-do-Mediterrâneo	155
Mediterranean spearfish	<i>Tetrapturus belone</i>	Espadim-do-Mediterrâneo	186
Mediterranean starry ray	<i>Raja asterias</i>	Raia-pintada	45
Mediterranean starry ray	<i>Raja montagui</i>	Raia-manchada	47
Megrim	<i>Lepidorhombus whiffagonis</i>	Areeiro	187
Merrett's spiny eel	<i>Polyacanthonotus merretti</i>	Enguia-espinhosa-de-Merrett	58
Messina rockfish	<i>Scorpaenodes arenai</i>	Rascasso-de-Messina	234
Metallic lanternfish	<i>Myctophum affine</i>	Peixe-lanterna-metálico	124
Meteor dragonet	<i>Protogrammus sousai</i>	Peixe-pau-de-Sousa	294
Michael Sars smooth-head	<i>Bathytroctes michaelisarsi</i>	Celindra-de-Michael-Sars	76
Midridged searsid	<i>Mentodus mesalinus</i>	Celindra-de-tubo-de-nervura	81
Midwater scorpionfish	<i>Ectreposebastes imus</i>	Rascasso-da-fundura	229
Milk shark	<i>Rhizoprionodon acutus</i>	Tubarão-bicudo	28
Mirror lanternfish	<i>Lampadена speculigera</i>	Peixe-lanterna-espelhado	121
Mirrorbelly	<i>Monacoa grimaldii</i>	Peixe-sola-de-Grimaldi	85
Mirrorwing flyingfish	<i>Hirundichthys speculiger</i>	Peixe-voador-de-asa-espelhada	201
Molly miller	<i>Scartella cristata</i>	Marachomba-de-crista	212
Monkey goby	<i>Neogobius fluviatilis</i>	Caboz-do-rio	183
Monrovia doctorfish	<i>Acanthurus monroviae</i>	Peixe-cirurgião	295
Montagu's blenny	<i>Coryphoblennius galerita</i>	Marachomba-das-rochas	207
Montagus seasnail	<i>Liparis montagui</i>	Peixe-lesma-de-Montagus	238
Morocco dentex	<i>Dentex maroccanus</i>	Cachucho-dentão	265
Mottled false moray	<i>Chlopsis dentatus</i>	Congrinho-malhado	63
Mottled grouper	<i>Mycteroperca rubra</i>	Garoupa-chumbo	245
Mouse catshark	<i>Galeus murinus</i>	Leitão-da-Íslândia	21
Muddy arrowtooth eel	<i>Ilyophis brunneus</i>	Enguia-da-fundura-castanha	58
Multipore searsid	<i>Normichthys operosus</i>	Celindra-de-tubo-multiporo	81
Multispotted lanternfish	<i>Scopelopsis multipunctatus</i>	Peixe-lanterna-sarapintado	126
Murray's smooth-head	<i>Conocara murrayi</i>	Celindra-de-Murray	77
Myers' flabby whalefish	<i>Gyrinomimus myersi</i>	Peixe-baleia-de-Myers	154
Naked lanternfish	<i>Taaningichthys paurolychnus</i>	Peixe-lanterna-nu	127
Narrownecked oceanic eel	<i>Derichthys serpentinus</i>	Enguia-de-pescoço-fino	63
Narrow-snouted pipefish	<i>Syngnathus tenuirostris</i>	Marinha-de-focinho-estreito	178
Nigel's one-jawed eel	<i>Monognathus nigeli</i>	Enguia-da-fundura-de-Nigel	70
Niger hind	<i>Cephalopholis nigri</i>	Garoupa-do-Niger	242
Night shark	<i>Carcharhinus signatus</i>	Tubarão-da-noite	27
Nilsson's pipefish	<i>Syngnathus rostellatus</i>	Marinha-cabeça-chata	178
No English common name	<i>Argyripricus atlanticus</i>	Pai-velho-do-Atlântico	89
No English common name	<i>Barathronus roulei</i>	Abadejo-cego-de-Roule	160
No English common name	<i>Bassozetus taenia</i>	Abadejo-ténia	157
No English common name	<i>Bathophilus metallicus</i>	Peixe-dragão-de-Welsh	95
No English common name	<i>Bathophilus proximus</i>	Peixe-dragão-das-Bermudas	96
No English common name	<i>Bathophilus schizochirus</i>	Peixe-dragão-cinzento	96
No English common name	<i>Bathylichnops brachyrhynchus</i>	Peixe-fantasma-dardo-listado	84
No English common name	<i>Bathytyphlops sewelli</i>	Peixe-tripé-de-Sewell	106

English common name	Scientific name	Portuguese common name	page
No English common name	<i>Benthocometes robustus</i>	Abadejo-cometa	157
No English common name	<i>Borostomias elucens</i>	Peixe-dragão-brilhante	96
No English common name	<i>Chilomycterus spinosus spinosus</i>	Porco-espinho-americano	223
No English common name	<i>Cynoglossus lineolatus</i>	Língua-de-cão-riscada	196
No English common name	<i>Eustomias drechseli</i>	Peixe-dragão-de-Drechsel	98
No English common name	<i>Eustomias lipochirius</i>	Peixe-dragão-de-Santa Cruz	98
No English common name	<i>Eustomias simplex</i>	Peixe-dragão-de-bolbo-simples	99
No English common name	<i>Eustomias tenisoni</i>	Peixe-dragão-de-Tenison	100
No English common name	<i>Gigantactis longicirra</i>	Peixe-chicote-filamentoso	220
No English common name	<i>Grammatostomias circularis</i>	Peixe-dragão-de-mancha-redonda	100
No English common name	<i>Hydrolagus lusitanicus</i>	Ratazana-lusitanica	54
No English common name	<i>Lampanyctus gemmifer</i>	Peixe-lanterna-nacarado	122
No English common name	<i>Lestidiops pseudosphyraenoides</i>	Barracudina-do-Atlântico-nordeste	110
No English common name	<i>Linophryne lucifer</i>	Peixe-sestro-lucifer	221
No English common name	<i>Linophryne racemifera</i>	Peixe-sestro-pompom	221
No English common name	<i>Meteoria longidorsalis</i>	Abadejo-cego-de-dorsal-longa	161
No English common name	<i>Nansenia iberica</i>	Argentina-ibérica	83
No English common name	<i>Nansenia obliterata</i>	Argentina-do-Mediterrâneo	83
No English common name	<i>Photonectes leucospilus</i>	Peixe-dragão-de-pinta-branca	103
No English common name	<i>Pseudoscopelus sagamianus</i>	Peixe-glutão-de-Sagami	290
No English common name	<i>Salaria basilisca</i>	Marachomba-listada	211
No English common name	<i>Asquamiceps caeruleus</i>	Celindra-de-cabeça-azul	75
No English common name	<i>Abyssobrotula galatheae</i>	Abadejo-abissal	156
No English common name	<i>Ahliasaurus berryi</i>	Peixe-vigia-de-Berry	107
No English common name	<i>Aidablennius sphynx</i>	Marachomba-esfinge	207
No English common name	<i>Apletodon incognitus</i>	Sugador-dos-ouricós	204
No English common name	<i>Argyropelecus bogachevi</i>	Pai-velho-de-Bocage	89
No English common name	<i>Aristostomias grimaldii</i>	Peixe-dragão-de-Grimaldi	93
No English common name	<i>Aristostomias lunifer</i>	Peixe-dragão-lunar	93
No English common name	<i>Aristostomias xenostoma</i>	Peixe-dragão-das-Caraíbas	93
No English common name	<i>Arnoglossus grohmanni</i>	Carta-de-Grohmann	188
No English common name	<i>Astronesthes atlanticus</i>	Peixe-dragão-do-Atlântico	93
No English common name	<i>Astronesthes cyaneus</i>	Peixe-dragão-azulado	93
No English common name	<i>Astronesthes leucopogon</i>	Peixe-dragão-de-barbilo-branco	94
No English common name	<i>Astronesthes macropogon</i>	Peixe-dragão-de-barbilo-graúdo	94
No English common name	<i>Astronesthes similis</i>	Peixe-dragão-de-bolbo-filamentoso	95
No English common name	<i>Astronesthes zharovi</i>	Peixe-dragão-de-Zharov	95
No English common name	<i>Ataxolepis apus</i>	Peixe-baleia-mosaico	153
No English common name	<i>Barathrites iris</i>	Abadejo-arco-íris	156
No English common name	<i>Barathronus multidens</i>	Abadejo-cego-dentudo	160
No English common name	<i>Barathronus parfaiti</i>	Abadejo-cego-de-Vaillant	160
No English common name	<i>Bassozetus levistomaticus</i>	Abadejo-boca-doce	157
No English common name	<i>Bathophilus brevis</i>	Peixe-dragão-oval	95
No English common name	<i>Bathygadus favosus</i>	Lagartixa-do-mar-favosa	131
No English common name	<i>Bathymicrops multispinis</i>	Peixe-tripé-espinhososo	105
No English common name	<i>Bathymicrops regis</i>	Peixe-tripé-real	105
No English common name	<i>Bathyonus laticeps</i>	Abadejo-cabeçudo	157
No English common name	<i>Bellottia apoda</i>	Brótula-coxa	160
No English common name	<i>Brotulotaenia brevicauda</i>	Abadejo-de-cauda-curta	157
No English common name	<i>Bufoheratias wedli</i>	Peixe-farol-de-Wedl	216
No English common name	<i>Cataetyx allenii</i>	Brótula-de-Allen	160
No English common name	<i>Cataetyx laticeps</i>	Brótula-de-cabeça-chata	160
No English common name	<i>Caulophryne polynema</i>	Peixe-pescador-da-Madeira	215
No English common name	<i>Centrobranchus choerocephalus</i>	Peixe-lanterna-de-focinho-comprido	115
No English common name	<i>Centrophorus macquensis</i>	Quelmo	36

English common name	Scientific name	Portuguese common name	page
No English common name	<i>Cetichthys indagator</i>	Peixe-baleia-do-Galathea	153
No English common name	<i>Chaenophryne ramifera</i>	Peixe-farol-ramudo	216
No English common name	<i>Chaunacops coloratus</i>	Peixe-sapo-colorido	214
No English common name	<i>Chaunacops roseus</i>	Peixe-sapo-rosado	214
No English common name	<i>Chaunax suttkusi</i>	Peixe-sapo-de-Suttkus	214
No English common name	<i>Ctenochirichthys longimanus</i>	Peixe-farol-barbatana-comprida	217
No English common name	<i>Diplecogaster pectoralis</i>	Sugador-das-Canárias	205
No English common name	<i>Discoverichthys praecox</i>	Peixe-tripé-temporão	106
No English common name	<i>Ditropichthys storeri</i>	Peixe-baleia-de-Storer	154
No English common name	<i>Dolopichthys allector</i>	Peixe-farol-cornudo	217
No English common name	<i>Dolopichthys danae</i>	Peixe-farol-do-Dana	217
No English common name	<i>Dolopichthys jubatus</i>	Peixe-farol-de-crista	217
No English common name	<i>Dolopichthys karstenii</i>	Peixe-farol-de-Karsten	217
No English common name	<i>Dolopichthys longicornis</i>	Peixe-farol-diabo	217
No English common name	<i>Dysalotus alcocki</i>	Peixe-glutão-de-Alcock	289
No English common name	<i>Dysalotus oligoscolus</i>	Peixe-glutão	289
No English common name	<i>Dysommina proboscideus</i>	Enguia-de-tromba-de-elefante	58
No English common name	<i>Echinomacrurus mollis</i>	Lagartixa-do-mar-macia	135
No English common name	<i>Echiodon dentatus</i>	Peixe-pérola-dentado	159
No English common name	<i>Echiodon drummondii</i>	Peixe-pérola-de-Drummond	160
No English common name	<i>Eretmophorus kleinenbergi</i>	Peixe-remo-de-Kleinenberg	139
No English common name	<i>Eustomias acinosus</i>	Peixe-dragão-de-barbilo-acinoso	97
No English common name	<i>Eustomias braueri</i>	Peixe-dragão-lusitano	97
No English common name	<i>Eustomias contiguus</i>	Peixe-dragão-bibolboso	97
No English common name	<i>Eustomias dubius</i>	Peixe-dragão-ambíguo	98
No English common name	<i>Eustomias filifer</i>	Peixe-dragão-de-barbilo-filífero	98
No English common name	<i>Eustomias fissibarbis</i>	Peixe-dragão-de-barbilo-bifurcado	98
No English common name	<i>Eustomias longibarba</i>	Peixe-dragão-de-barbilo-crescido	98
No English common name	<i>Eustomias melanostigma</i>	Peixe-dragão-de-mancha-negra	99
No English common name	<i>Eustomias monodactylus</i>	Peixe-dragão-de-bolbo-monodáctilo	99
No English common name	<i>Eustomias parri</i>	Peixe-dragão-de-Parr	99
No English common name	<i>Eustomias radicifilis</i>	Peixe-dragão-de-barbilo-ramificado	99
No English common name	<i>Eustomias schiffi</i>	Peixe-dragão-de-Schiff	99
No English common name	<i>Eustomias tetranema</i>	Peixe-dragão-da-Madeira	100
No English common name	<i>Gadomus dispar</i>	Lagartixa-do-mar-impar	135
No English common name	<i>Gaidropsarus mauli</i>	Laibeque-de-Maul	145
No English common name	<i>Gigantactis ios</i>	Peixe-chicote-das-Canárias	220
No English common name	<i>Gigantactis vanhoeffeni</i>	Peixe-chicote-de-Van Höffen	220
No English common name	<i>Gnathophis codoniphorus</i>	Congro-dos-Açores	67
No English common name	<i>Grammatostomias dentatus</i>	Peixe-dragão-colmillos	100
No English common name	<i>Grammatostomias flagellibarba</i>	Peixe-dragão-de-barbilo-filamentoso	100
No English common name	<i>Grammonus ater</i>	Brótula-negra	161
No English common name	<i>Guttigadus latifrons</i>	Abrótea-cabeçuda	140
No English common name	<i>Himantolophus albinares</i>	Peixe-farol-da-Madeira	215
No English common name	<i>Himantolophus brevirostris</i>	Peixe-farol-de-focinho-curto	215
No English common name	<i>Himantolophus compressus</i>	Peixe-farol-sapo	216
No English common name	<i>Himantolophus mauli</i>	Peixe-farol-de-Maul	216
No English common name	<i>Holcomycteronus squamosus</i>	Abadejo-escamudo	158
No English common name	<i>Holtbyrnia melanocephala</i>	Celindra-de-tubo-de-cabeça-negra	80
No English common name	<i>Howella simplex</i>	Peixe-alado	241
No English common name	<i>Hyleurochilus bananensis</i>	Marachomba-banana	207
No English common name	<i>Hyleurochilus fissicornis</i>	Marachomba-da-toca	208
No English common name	<i>Kali indica</i>	Peixe-glutão-do-Índico	290
No English common name	<i>Kali kerberti</i>	Peixe-glutão-de-Kerbert	290
No English common name	<i>Kali macrodon</i>	Peixe-glutão-dentudo	290

English common name	Scientific name	Portuguese common name	page
No English common name	<i>Kali macrura</i>	Peixe-glutão-rabudo	290
No English common name	<i>Kali parri</i>	Peixe-glutão-de-Parr	290
No English common name	<i>Lamprogrammus niger</i>	Abadejo-cristado	158
No English common name	<i>Lasiognathus amphirhamphus</i>	Peixe-ganchudo	219
No English common name	<i>Lasiognathus beebei</i>	Peixe-ganchudo-de-Beebe	219
No English common name	<i>Lasiognathus saccostoma</i>	Peixe-ganchudo-tridente	219
No English common name	<i>Leptoderma macrophthalmum</i>	Celindra-dos-Açores	78
No English common name	<i>Leptostomias haplocaulus</i>	Peixe-dragão-de-barbilo-singelo	101
No English common name	<i>Leptostomias leptobolus</i>	Peixe-dragão-de-bolbo-alongado	101
No English common name	<i>Leptostomias longibarba</i>	Peixe-dragão-de-barbilo-longo	101
No English common name	<i>Lestidiops similis</i>	Barracudina-análoga	110
No English common name	<i>Lesueurigobius heterofasciatus</i>	Caboz-listado	182
No English common name	<i>Leucobrotula adipata</i>	Brótula-falsa-branca	161
No English common name	<i>Linophryne arborifera</i>	Peixe-sestro-ramoso	220
No English common name	<i>Linophryne arcturi</i>	Peixe-sestro-do-Arcturus	220
No English common name	<i>Linophryne brevibarbata</i>	Peixe-sestro-de-barbilo-trirramoso	220
No English common name	<i>Linophryne coronata</i>	Peixe-sestro-coroad	221
No English common name	<i>Linophryne macrodon</i>	Peixe-sestro-dentudo	221
No English common name	<i>Linophryne maderensis</i>	Peixe-sestro-da-Madeira	221
No English common name	<i>Linophryne polypogon</i>	Peixe-sestro-barbudo	221
No English common name	<i>Linophryne sexfilis</i>	Peixe-sestro-hexafilamento	221
No English common name	<i>Linophryne algibarbata</i>	Peixe-sestro-de-Waterman	220
No English common name	<i>Lipophrys trigloides</i>	Marachomba-trigloide	208
No English common name	<i>Lycenchelys alba</i>	Enguia-beiçuda-alva	288
No English common name	<i>Lyconus brachycolus</i>	Peixe-de-focinho-arreganhado	148
No English common name	<i>Macroparalepis brevis</i>	Barracudina-pequena	110
No English common name	<i>Macroparalepis nigra</i>	Barracudina-negra	110
No English common name	<i>Macruronus maderensis</i>	Peixe-orelhudo-da-Madeira	148
No English common name	<i>Manducus maderensis</i>	Boqueirão-da-Madeira	88
No English common name	<i>Mauligobius maderensis</i>	Caboz-da-Madeira	183
No English common name	<i>Melamphaes ebelingi</i>	Peixe-rã-de-Ebeling	150
No English common name	<i>Melamphaes falsidicus</i>	Peixe-rã-falso	150
No English common name	<i>Melamphaes indicus</i>	Peixe-rã-do-Índico	150
No English common name	<i>Melamphaes polylepis</i>	Peixe-rã-cabecinha	151
No English common name	<i>Melamphaes pumilus</i>	Peixe-rã-anão	151
No English common name	<i>Melanostomias biseriatus</i>	Peixe-dragão-de-barbilo-bisseriado	101
No English common name	<i>Melanostomias macrophotus</i>	Peixe-dragão-luminoso	101
No English common name	<i>Melanostomias melanopogon</i>	Peixe-dragão-de-barbilo-negro	102
No English common name	<i>Melanostomias melanops</i>	Peixe-dragão-de-Sumatra	102
No English common name	<i>Melanostomias spilorhynchus</i>	Peixe-dragão-de-focinho-manchado	102
No English common name	<i>Meteoria erythrops</i>	Abadejo-cego-dos-Açores	161
No English common name	<i>Microlipophrys canevae</i>	Marachomba-de-Caneva	208
No English common name	<i>Microlipophrys dalmatinus</i>	Marachomba-dálmata	209
No English common name	<i>Monognathus jesperseni</i>	Enguia-da-fundura-de-Jespersen	70
No English common name	<i>Monomitopus metriostoma</i>	Abadejo-da-Mauritânia	158
No English common name	<i>Nansenia atlantica</i>	Argentina-atlântica	83
No English common name	<i>Nansenia boreacrassicauda</i>	Argentina-boreal	83
No English common name	<i>Nansenia cf. pelagica</i>	Argentina-pelágica	83
No English common name	<i>Nansenia longicauda</i>	Argentina-de-cauda-longa	83
No English common name	<i>Nansenia tenera</i>	Argentina-dourada	83
No English common name	<i>Narcetes erimelas</i>	Celindra-de-Goa	78
No English common name	<i>Notoscopelus elongatus</i>	Peixe-lanterna-do-Mediterrâneo	125
No English common name	<i>Nybelinella brevianalis</i>	Abadejo-cego-da-Madeira	161
No English common name	<i>Nybelinella erikssoni</i>	Abadejo-cego-de-Eriksson	161
No English common name	<i>Oneirodes anisacanthus</i>	Peixe-farol-de-bolbo-assimétrico	217

English common name	Scientific name	Portuguese common name	page
No English common name	<i>Oneirodes carlsbergi</i>	Peixe-farol-de-Carlsberg	218
No English common name	<i>Oneirodes clarkei</i>	Peixe-farol-de-Clarke	218
No English common name	<i>Oneirodes macronema</i>	Peixe-farol-de-Santa Cruz	218
No English common name	<i>Oneirodes macrosteus</i>	Peixe-farol-do-Atlântico-Norte	218
No English common name	<i>Oneirodes myrionemus</i>	Peixe-farol-multifilamentoso	218
No English common name	<i>Oneirodes posti</i>	Peixe-farol-de-Post	218
No English common name	<i>Opeatogenys gracilis</i>	Sugador-do-Mediterrâneo	206
No English common name	<i>Pachycara saldanhai</i>	Enguia-beiçuda-de-Saldanha	288
No English common name	<i>Pachycara thermophilum</i>	Enguia-beiçuda-termofílica	288
No English common name	<i>Parablennius incognitus</i>	Marachomba-das-cracas	209
No English common name	<i>Parablennius rouxi</i>	Marachomba-branca	210
No English common name	<i>Paracaristius maderensis</i>	Peixe-crina-da-Madeira	260
No English common name	<i>Paralepis brevirostris</i>	Barracudina-de-focinho-curto	110
No English common name	<i>Paralepis speciosa</i>	Barracudina-manchada	111
No English common name	<i>Paraliparis membranaceus</i>	Peixe-lesma-membranoso	238
No English common name	<i>Paraliparis nigellus</i>	Peixe-lesma-escuro	238
No English common name	<i>Parophidion vassali</i>	Abadejo-vermelho	158
No English common name	<i>Penopus microphthalmus</i>	Abadejo-de-olho-pequeno	158
No English common name	<i>Pentherichthys atratus</i>	Peixe-farol-do-Panamá	218
No English common name	<i>Photocorynus spiniceps</i>	Peixe-sestro-cabeça-espinhosa	221
No English common name	<i>Photonectes dinema</i>	Peixe-dragão-bifilar	103
No English common name	<i>Photonectes margarita</i>	Peixe-dragão-boca-negra	103
No English common name	<i>Photostomias atrox</i>	Peixe-dragão-atroz	103
No English common name	<i>Photostomias goodyeari</i>	Peixe-dragão-de-Goodyear	103
No English common name	<i>Phyllorhinichthys micractis</i>	Peixe-farol-de-Guadalupe	218
No English common name	<i>Platyberyx andriashevi</i>	Peixe-crina-de-Andriashev	260
No English common name	<i>Platyberyx mauli</i>	Peixe-crina-de-Maul	260
No English common name	<i>Platyberyx opalescens</i>	Peixe-crina-opala	260
No English common name	<i>Polyipnus polli</i>	Pai-velho-redondo	91
No English common name	<i>Polymetme thaeocoryla</i>	Peixe-luminoso-do-Atlântico	92
No English common name	<i>Poromitra nigriceps</i>	Peixe-rã-de-cabeça-preta	152
No English common name	<i>Pseudonezumia flagellicauda</i>	Lagartixa-do-mar-de-chicote	138
No English common name	<i>Pseudoscopelus altipinnis</i>	Peixe-glutão-de-vela	290
No English common name	<i>Pseudoscopelus obtusifrons</i>	Peixe-glutão-de-focinho-curvo	290
No English common name	<i>Pseudoscopelus scutatus</i>	Peixe-glutão-escudado	291
No English common name	<i>Puck pinnata</i>	Peixe-farol-alado	219
No English common name	<i>Rhynchogadus hepaticus</i>	Abrótea-amuada	142
No English common name	<i>Saccopharynx harrisoni</i>	Peixe-pelícano-de-Harrison	71
No English common name	<i>Saccopharynx hjorti</i>	Peixe-pelícano-de-Hjort	71
No English common name	<i>Saccopharynx paucovertebratis</i>	Peixe-pelícano-dos-Açores	71
No English common name	<i>Saccopharynx trilobatus</i>	Peixe-pelícano-de-cauda-trilobada	71
No English common name	<i>Sciadonus pedicellaris</i>	Abadejo-cego-do-Pacifico	161
No English common name	<i>Scopelosaurus smithii</i>	Peixe-vigia-de-Smith	108
No English common name	<i>Spectrunculus crassus</i>	Abadejo-dos-Açores	159
No English common name	<i>Spicara flexuosa</i>	Trombeiro-do-Mediterrâneo	274
No English common name	<i>Stomias brevibarbatus</i>	Peixe-dragão-barbilo-curto	104
No English common name	<i>Syphodus caeruleus</i>	Bodião-dos-Açores	284
No English common name	<i>Syphodus ocellatus</i>	Bodião-de-pinta	285
No English common name	<i>Syphodus rostratus</i>	Bodião-das-ervas	286
No English common name	<i>Synaphobranchus dolichorhynchus</i>	Moreão-bicudo	59
No English common name	<i>Syngnathus phlegon</i>	Marinha-espinhosa	177
No English common name	<i>Tripterygion tripteronotum</i>	Caboz-de-três-dorsais-do-Mediterrâneo	206
No English common name	<i>Uncisudis longirostra</i>	Barracudina-da-Madeira	111
No English common name	<i>Vanneaugobius canariensis</i>	Caboz-das-Canárias	184
No English common name	<i>Yarrella blackfordi</i>	Peixe-luminoso-de-Blackford	93

English common name	Scientific name	Portuguese common name	page
<i>Nomen dubium in Mycteroptera</i>	<i>Epinephelus guaza</i>	<i>Nomen dubium in Mycteroptera</i>	243
Norman's smooth-head	<i>Mirognathus normani</i>	Celindra-de-Norman	78
Northern pipefish	<i>Syngnathus fuscus</i>	Marinha-setentrional	177
Northern swallower eel	<i>Saccopharynx ampullaceus</i>	Peixe-pelícano-do-Norte	70
Northern wolffish	<i>Anarhichas denticulatus</i>	Peixe-lobo-azul	289
Norway goby	<i>Pomatoschistus norvegicus</i>	Caboz-norueguês	184
Norwegian skate	<i>Dipturus nidanensis</i>	Raia-norueguesa	43
Norwegian skate	<i>Dipturus sp. Stehmann</i>	Raia-norueguesa	44
Oarfish;	<i>Regalecus russellii</i>	Peixe-remo-do-Índico	129
Ocean sunfish	<i>Mola mola</i>	Peixe-lua	222
Ocean triggerfish	<i>Canthidermis sufflamen</i>	Cangulo-oceânico	228
Oceanic lightfish	<i>Vinciguerria nimbaria</i>	Peixe-luminoso-oceânico	93
Oceanic puffer	<i>Lagocephalus lagocephalus</i>	Peixe-bola-oceânico	225
Oceanic two-wing flyingfish	<i>Exocoetus obtusirostris</i>	Andorinha-do-mar-oceânica	201
Oceanic whitetip shark	<i>Carcharhinus longimanus</i>	Tubarão-de-pontas-brancas	26
Ocellated wedge sole	<i>Microchirus hexophthalmus</i>	Azevia-de-olhos	193
Offshore rockfish	<i>Pontinus kuhlii</i>	Cantarilho-requeime	230
Oilfish	<i>Ruvettus pretiosus</i>	Escolar	163
Oily halosaur	<i>Aldrovandia oleosa</i>	Peixe-lagarto-oleoso	56
Omosudid	<i>Omosudis lowii</i>	Peixe-mandíbula-de-martelo	112
Opah	<i>Lampris guttatus</i>	Peixe-cravo	127
Opal chimaera	<i>Chimaera opalescens</i>	Quimera-opala	53
Orange roughy	<i>Hoplostethus atlanticus</i>	Olho-de-vidro-laranja	154
Ornate wrasse	<i>Thalassoma pavo</i>	Bodião-verde	286
Ovate lightfish	<i>Ichthyococcus ovatus</i>	Peixe-luminoso-oval	92
Owen's halosaur	<i>Halosaurus oweni</i>	Peixe-lagarto-de-Owen	57
Oxeye scad	<i>Selar boops</i>	Charro-olho-de-boi	254
Pacific barracudina	<i>Lestidiops jayakari</i>	Barracudina-do-Pacífico	109
Pacific blackchin	<i>Scopelengys tristis</i>	Peixe-lanterna-do-Pacífico	113
Pacific burrfish	<i>Chilomycterus reticulatus</i>	Porco-espinho-de-pintas	222
Pacific hatchet fish	<i>Argyropelecus affinis</i>	Pai-velho-do-Pacífico	89
Pacific longnose chimaera	<i>Harriotta raleighana</i>	Peixe-rato-de-Raleigh	54
Pacific portholefish	<i>Diplophos taenia</i>	Boqueirão-comprido	88
Pacific spotted scorpionfish	<i>Scorpaena plumieri</i>	Rascasso-negro	232
Painted comber	<i>Serranus scriba</i>	Serrano-riscado	246
Painted eel	<i>Echelus myrus</i>	Cobra-de-orelhas	64
Pale bristlemouth	<i>Cyclothona alba</i>	Boqueirão-branco	86
Pale chimaera	<i>Hydrolagus pallidus</i>	Ratazana-pálida	54
Pale halosaur	<i>Aldrovandia affinis</i>	Peixe-lagarto-pálido	56
Pale ray	<i>Bathyraja pallida</i>	Raia-pálida	49
Pale threadtail snipe eel	<i>Nemichthys curvirostris</i>	Cobra-pálida-ponteada	68
Palebelly searsid	<i>Barbantus curvifrons</i>	Celindra-de-tubo-de-ventre-claro	80
Palegold searsid	<i>Maulisia argipalla</i>	Celindra-de-tubo-dourada	80
Pallid sculpin	<i>Cottunculus thomsonii</i>	Escorpião-pálido	237
Panama lightfish	<i>Vinciguerria lucetia</i>	Peixe-luminoso-do-Panamá	92
Parin's spinyfish	<i>Diretmichthys parini</i>	Peixe-enfadado-de-Parin	155
Parrot grunt	<i>Pomadasys perotaei</i>	Roncador-papagaio	263
Parrot seaperch	<i>Callanthias ruber</i>	Canarinho-do-mar	247
Parrotfish	<i>Sparisoma cretense</i>	Papagaio-velho	287
Patchwork lanternfish	<i>Notoscopelus resplendens</i>	Peixe-lanterna-oliva	126
Pawnee dragonfish	<i>Bathophilus pawnee</i>	Peixe-dragão-de-Pawnee	96
Peacock blenny	<i>Salarias pavo</i>	Marachomba-pavão	212
Peacock wrasse	<i>Syphodus tinca</i>	Bodião-pavão	286
Pearl fish	<i>Carapus acus</i>	Peixe-pérola-cobrelo	159
Pearlsides	<i>Maurolicus amethystinopunctatus</i>	Peixe-luz	90

English common name	Scientific name	Portuguese common name	page
Pearly razorfish	<i>Xyrichtys novacula</i>	Mordedor	287
Pearlyspotted lanternfish	<i>Myctophum nitidulum</i>	Peixe-lanterna-fosforescente	124
Pelagic basslet	<i>Howella brodiei</i>	Peixe-alado-de-Brodie	240
Pelagic butterfish	<i>Schedophilus maculatus</i>	Liro-manchado	172
Pelagic porcupinefish	<i>Diodon eydouxii</i>	Peixe-ouriço-pelágico	223
Pelagic stingray	<i>Pteroplatytrygon violacea</i>	Uge-violeta	50
Pelagic thresher or Bigeye thresher	<i>Alopias vulpinus</i>	Tubarão-raposo	20
Pelagic thresher or Thresher	<i>Alopias superciliosus</i>	Tubarão-raposo-olhudo	19
Pelican eel	<i>Eurypharynx pelecanoides</i>	Peixe-pelícano	70
Pelican flounder	<i>Chascanopsetta lugubris</i>	Carta-pelícano	190
Pencil cardinal	<i>Epigonus denticulatus</i>	Olhudo-denteado	239
Phaeton dragonet	<i>Synchiropus phaeton</i>	Peixe-pau-rosa	294
Picarel	<i>Spicara smaris</i>	Trombeiro-boga	274
Picked spurdog	<i>Squalus megalops</i>	Galhudo-de-focinho-curto	39
Pigeye shark	<i>Carcharhinus leucas</i>	Tubarão-buldogue	26
Pilotfish	<i>Naucrates ductor</i>	Peixe-piloto	253
Pink dentex	<i>Dentex gibbosus</i>	Capatão-de-bandeira	264
Pink flabby whalefish	<i>Cetostoma regani</i>	Peixe-baleia-rosa	154
Pink frogmouth	<i>Chaunax pictus</i>	Peixe-sapo-mole	214
Piper gurnard	<i>Trigla lyra</i>	Cabra-lira	236
Plain bonito	<i>Orcynopsis unicolor</i>	Palometa	167
Plainchin dreamarm	<i>Leptacanthichthys gracilispinis</i>	Peixe-farol-de-queixo-direito	217
Planehead filefish	<i>Stephanolepis hispidus</i>	Peixe-porco-áspero	226
Polkadot catshark	<i>Scyliorhinus stellaris</i>	Pata-roxa-gata	22
Pollack	<i>Pollachius pollachius</i>	Juliana	143
Polygon moray	<i>Gymnothorax polygonius</i>	Moreia-poligonal	61
Pompano dolphinfish	<i>Coryphaena equiselis</i>	Doirado-pampo	258
Pompano	<i>Trachinotus ovatus</i>	Sereia-camochilo	256
Poor cod	<i>Trisopterus minutus</i>	Fanecão	144
Poor cod Mediterranean	<i>Trisopterus capelanus</i>	Fanecão-do-Mediterrâneo	143
Popeye lampfish	<i>Bolinichthys longipes</i>	Peixe-lanterna-barbatana-comprida	114
Porbeagle	<i>Lamna nasus</i>	Tubarão-sardo	19
Portuguese blenny	<i>Parablennius ruber</i>	Marachomba-portuguesa	210
Portuguese dogfish	<i>Centroscymnus coelolepis</i>	Carocho	32
Portuguese sole	<i>Dagetichthys lusitanicus</i>	Língua-de-vaca	192
Pouting	<i>Trisopterus luscus</i>	Faneca	144
Power's deep-water bristle-mouth fish	<i>Vinciguerria poweriae</i>	Peixe-luminoso-de-Power	93
Pricklefish	<i>Acanthochaenus luetkenii</i>	Peixe-de-coroa-espinhosa	153
Prickly dreamer	<i>Sphryne gladifsenae</i>	Peixe-farol-espinhoso	219
Prickly puffer	<i>Ephippion guttifer</i>	Peixe-bola-de-pintas	224
Pudgy cuskeel	<i>Spectrunculus grandis</i>	Abadejo-grande	159
Pugnose grenadier	<i>Sphagemacrurus grenadæ</i>	Lagartixa-do-mar-de-nariz-arrebitado	138
Purplemouth moray	<i>Gymnothorax vicinus</i>	Moreão-amarelo	61
Purplemouthed conger	<i>Pseudophichthys splendens</i>	Congro-de-boca-púrpura	67
Pygmy filefish	<i>Stephanolepis setifer</i>	Peixe-porco-galhudo	227
Pygmy lanternfish	<i>Lampanyctus pusillus</i>	Peixe-lanterna-anão	123
Pygmy moray	<i>Anarchias longicauda</i>	Moreia-de-cauda-longa	60
Pygmy moray	<i>Anarchias similis</i>	Moreia-anã	60
Queen snapper	<i>Etelis oculatus</i>	Rosa-salmão	261
Queen triggerfish	<i>Balistes vetula</i>	Cangulo-real	228
Rabbit fish	<i>Chimaera monstrosa</i>	Quimera-comum	53
Ragged bigscale	<i>Scopelogadus mizolepis</i>	Peixe-rã-couraça	153
Rainbow runner	<i>Elagatis bipinnulata</i>	Fogueteiro-arco-íris	253
Rainbow trout	<i>Oncorhynchus mykiss</i>	Truta-arco-íris	85
Rainbow wrasse	<i>Coris julis</i>	Judia	281

English common name	Scientific name	Portuguese common name	page
Rare lanternfish	<i>Loweina rara</i>	Peixe-lanterna-raro	124
Red bandfish	<i>Cepola macrophthalma</i>	Suspensório	278
Red cornetfish	<i>Fistularia petimba</i>	Corneta-rosada	174
Red dory	<i>Cyttopsis rosea</i>	Galo-de-natura	129
Red gurnard	<i>Chelidonichthys cuculus</i>	Cabra-vermelha	234
Red mullet	<i>Mullus barbatus</i>	Salmonete-da-vasa	277
Red pandora	<i>Pagellus bellottii</i>	Bica-buço	270
Red porgy	<i>Pagrus pagrus</i>	Pargo-legítimo	272
Red scorpionfish	<i>Scorpaena scrofa</i>	Rascasso-vermelho	233
Red squirrelfish	<i>Sargocentron hastatum</i>	Esquilo-real	156
Redbanded seabream	<i>Pagrus auriga</i>	Pargo-sêmola	271
Redbarred lizardfish	<i>Synodus synodus</i>	Peixe-lagarto-de-rolo	108
Redlip Blenny;	<i>Ophioblennius atlanticus</i>	Marachomba-do-Atlântico	209
Redmouth whalefish	<i>Rondeletia loricata</i>	Peixe-baleia-de-boca-vermelha	153
Red-mouthed goby	<i>Gobius cruentatus</i>	Caboz-de-boca-vermelha	180
Redtail parrotfish	<i>Sparisoma chrysopeterum</i>	Papagaio	288
Reef butterflyfish	<i>Chaetodon sedentarius</i>	Borboleta-do-recife	278
Reinhardt's lanternfish	<i>Hygophum reinhardtii</i>	Peixe-lanterna-de-Reinhardt	120
Rendezvous fish	<i>Polymetme corythaæola</i>	Peixe-luminoso-do-Índico	92
Reticulated dragonet	<i>Callionymus reticulatus</i>	Peixe-pau-listado	294
Reticulated tonguefish	<i>Syphurus reticulatus</i>	Língua-reticulada	197
Ribbon sawtail fish	<i>Idiacanthus fasciola</i>	Peixe-dragão-de-cauda-serreada	100
Richardson's ray	<i>Bathyraja richardsoni</i>	Raia-de-Richardson	49
Richardson's snaggletooth	<i>Astronesthes richardsoni</i>	Peixe-dragão-de-Richardson	95
Ridgehead	<i>Melamphaes microps</i>	Peixe-rã-olhinho	150
Ridgehead	<i>Melamphaes simus</i>	Peixe-rã-nariz-achatado	151
Ridgehead	<i>Melamphaes typhlops</i>	Peixe-rã-da-Madeira	151
Ridgehead	<i>Poromitra capito</i>	Peixe-rã-de-nariz-espinhoso	151
Ridgehead	<i>Scopeloberyx rubriventer</i>	Peixe-rã-dos-Açores	152
Ringneck blenny	<i>Parablennius pilicornis</i>	Marachomba-cornuda	210
Risso's dragonet	<i>Callionymus risso</i>	Peixe-pau-pintado	294
Risso's smooth-head	<i>Alepocephalus rostratus</i>	Celindra-de-Risso	75
Robust cod	<i>Laemonema robustum</i>	Abrótea-robusta	140
Rock cook	<i>Centrolabrus exoletus</i>	Bodião-de-boca-pequena	281
Rock goby	<i>Gobius paganellus</i>	Caboz-da-rocha	181
Rock gunnel	<i>Pholis gunnellus</i>	Peixe-manteiga	289
Rock hind	<i>Epinephelus adscensionis</i>	Garoupa-pintada-da-rocha	243
Rock-pool blenny	<i>Parablennius parvifrons</i>	Marachomba-das-poças	210
Roudi escolar	<i>Promethichthys prometheus</i>	Escolar-branco	163
Rough longnose dogfish	<i>Deania hystricosa</i>	Sapata-áspera	37
Rough pomfret	<i>Taractes asper</i>	Xaputa-rugosa	259
Rough scad	<i>Trachurus lathami</i>	Carapau-rugoso	256
Rough triggerfish	<i>Canthidermis maculata</i>	Cangulo-áspero	228
Roughear scad	<i>Decapterus labiatus</i>	Charro-de-cauda-vermelha	252
Roughhead grenadier	<i>Macrourus berglax</i>	Lagartixa-do-mar-olho-de-cebola	136
Roughlip grenadier	<i>Coryphaenoides theleostomus</i>	Lagartixa-do-mar-da-Madeira	134
Roughskin dogfish	<i>Centroscymnus owstonii</i>	Xara-preta-de-natura	32
Roughskin dogfish	<i>Centroselachus crepidater</i>	Sapata-preta	33
Roughsnout grenadier	<i>Trachyrincus scabrus</i>	Peixe-lima	139
Roughtip grenadier	<i>Nezumia sclerorhynchus</i>	Largatixa-do-mar-áspera	137
Roule's goby	<i>Gobius roulei</i>	Caboz-de-Roule	181
Round ray	<i>Rajella fyllae</i>	Raia-redonda	48
Round sardinella	<i>Sardinella aurita</i>	Sardinela-lombuda	73
Round scad	<i>Decapterus punctatus</i>	Charro-moiro	252
Round stingray	<i>Taeniurrops grabatus</i>	Uge-redonda	50

English common name	Scientific name	Portuguese common name	page
Roundhead grenadier	<i>Odontomacrus murayi</i>	Lagartixa-do-mar-de-Murray	137
Roundnose grenadier	<i>Coryphaenoides rupestris</i>	Lagartixa-da-rocha	134
Roundnose lanternfish	<i>Centrobranchus nigroocellatus</i>	Peixe-lanterna-de-focinho-redondo	115
Roundscale spearfish	<i>Tetrapturus georgii</i>	Espadim-peto	186
Royal flagfin	<i>Aulopus filamentosus</i>	Lagarto-do-mar	104
Rubberlip grunt	<i>Plectrohinchus mediterraneus</i>	Pombo	262
Rudderfish	<i>Centrolophus niger</i>	Liro-preto	171
Rufous lanternfish	<i>Symbolophorus rufinus</i>	Peixe-lanterna-de-Rufous	126
Rüppell's scaldback	<i>Arnoglossus rueppelii</i>	Carta-estreita	189
Rusty blenny	<i>Parablennius sanguinolentus</i>	Marachomba-do-Mar-Negro	211
Saddled seabream	<i>Oblada melanura</i>	Dobradiça	269
Saddled snake eel	<i>Pisodonophis semicinctus</i>	Cobra-malhada	65
Sailfin dragonet	<i>Callionymus pusillus</i>	Peixe-pau-rabudo	294
Sailfin roughshark	<i>Oxynotus paradoxus</i>	Peixe-porco-de-vela	35
Sailray	<i>Rajella linteata</i>	Raia-nevoeira	48
Salema	<i>Sarpa salpa</i>	Salema	273
Salmon smooth-head	<i>Conocara salmonicum</i>	Celindra-salmonada	77
Sand goby	<i>Pomatoschistus minutus</i>	Caboz-da-areia	184
Sand smelt	<i>Atherina presbyter</i>	Peixe-rei	198
Sand sole	<i>Pegusa lascaris</i>	Linguado-da-areia	194
Sand steenbras	<i>Lithognathus mormyrus</i>	Ferreira	269
Sand tiger shark	<i>Odontaspis ferox</i>	Tubarão-areia	17
Sandbar shark	<i>Carcharhinus plumbeus</i>	Tubarão-corre-costa	27
Sandy ray	<i>Leucoraja circularis</i>	Raia-de-São-Pedro	44
Sanzo's goby	<i>Lesueurigobius sanzi</i>	Caboz-de-Sanzo	182
Sargassum pipefish	<i>Syngnathus pelagicus</i>	Marinha-do-sargaço	177
Sargassum triggerfish	<i>Xanthichthys ringens</i>	Cangulo-do-sargaço	228
Sargassumfish	<i>Histrio histrio</i>	Peixe-sapo-do-sargaço	213
Sawback angelshark	<i>Squatina aculeata</i>	Anjo-espinhosos	39
Scaleless black dragonfish	<i>Bathophilus digitatus</i>	Peixe-dragão-iridescente	95
Scaleless black dragonfish	<i>Melanostomias bartonbeani</i>	Peixe-dragão-de-Barton Bean	101
Scaleless dragonfish	<i>Bathophilus nigerrimus</i>	Peixe-dragão-chicote	96
Scaleless dragonfish	<i>Bathophilus vaillanti</i>	Peixe-dragão-de-Vaillant	96
Scaleless dragonfish	<i>Chiromystes plioptyerus</i>	Peixe-dragão-de-bolbo-papilar	97
Scaleless dragonfish	<i>Eustomias furcifer</i>	Peixe-dragão-de-furfur	98
Scaleless dragonfish	<i>Eustomias obscurus</i>	Peixe-dragão-obscuro	99
Scaleless dragonfish	<i>Leptostomias gladiator</i>	Peixe-dragão-gladiador	100
Scale-rayed wrasse	<i>Acantholabrus palloni</i>	Bodião-do-alto	280
Scalloped hammerhead	<i>Sphyrna lewini</i>	Tubarão-martelo-recortado	28
Scalloped ribbonfish	<i>Zu cristatus</i>	Peixe-tábua-recortado	128
Scamp	<i>Mycteroperca phenax</i>	Garoupa-mariola	245
Schmidt's dragonfish	<i>Eustomias schmidti</i>	Peixe-dragão-de-Schmidt	99
Schnakenbeck's searsid	<i>Sagamichthys schnakenbecki</i>	Celindra-de-tubo-de-Schnakenbeck	81
Scrawled cowfish	<i>Acanthostracion quadricornis</i>	Peixe-cofre-touro	226
Scribbled leatherjacket filefish	<i>Aluterus scriptus</i>	Peixe-porco-bicudo	226
Sea lamprey	<i>Petromyzon marinus</i>	Lampreia-do-mar	15
Sea stickleback	<i>Spinachia spinachia</i>	Esgana-gata-marinha	297
Sea trout	<i>Salmo trutta</i>	Truta-marisca	86
Senegal seabream	<i>Diplodus bellottii</i>	Sargo-do-Senegal	266
Senegalese frogfish	<i>Fowlerichthys senegalensis</i>	Peixe-sapo-do-Senegal	213
Senegalese rockfish	<i>Scorpaena laevis</i>	Rascasso-do-Senegal	231
Senegalese sole	<i>Solea senegalensis</i>	Linguado-do-Senegal	195
Sergeant-major	<i>Abudefduf saxatilis</i>	Castanheta-raiada	278
Serpent eel	<i>Ophisurus serpens</i>	Cobra-do-mar	64
Shadow bristlemouth	<i>Cyclothona parapallida</i>	Boqueirão-sombreado	87

English common name	Scientific name	Portuguese common name	page
Shagreen ray	<i>Leucoraja fullonica</i>	Raia-pregada	44
Shanny	<i>Lipophrys pholis</i>	Marachomba-frade	208
Shark sucker	<i>Remora remora</i>	Pegador	250
Sharktooth moray	<i>Gymnothorax maderensis</i>	Moreia-da-Madeira	61
Sharpnose halosaur	<i>Aldrovandia rostrata</i>	Peixe-lagarto-de-bico	56
Sharpnose puffer	<i>Canthigaster rostrata</i>	Peixe-bola-das-Caraíbas	224
Sharpnose sevengill shark	<i>Heptanchias perlo</i>	Tubarão-de-sete-guelras	15
Sharpsnout seabream	<i>Diplodus puntazzo</i>	Sargo-bicudo	267
Sharptail mola	<i>Masturus lanceolatus</i>	Peixe-lua-rabudo	222
Sharp-toothed wrasse	<i>Lappanella fasciata</i>	Bodião-verdelho	284
Sherborn's pelagic bass	<i>Howella sherborni</i>	Peixe-alado-de-Sherborn	241
Shi drum	<i>Umbrina cirrosa</i>	Calafate-das-riscas	276
Shore clingfish	<i>Lepadogaster lepadogaster</i>	Sugador-de-Gouan	205
Shore rockling	<i>Gaidropsarus mediterraneus</i>	Laibeque-do-Mediterrâneo	145
Short fin pearleye	<i>Scopelarchus analis</i>	Olho-pérola-de-barbatana-curta	107
Short snouted lancetfish	<i>Alepisaurus brevirostris</i>	Lírio-ferro-de-focinho-curto	113
Short snouted seahorse	<i>Hippocampus hippocampus</i>	Cavalo-marinho-de-focinho-curto	176
Short-beaked garfish	<i>Belone svetovidovi</i>	Agulha-do-bico-curto	199
Shortbeard grenadier	<i>Coryphaenoides brevibarbis</i>	Lagartixa-do-mar-de-barbilha-curto	133
Shortdorsal cutthroat eel	<i>Synaphobranchus brevidorsalis</i>	Moreão-de-dorsal-curta	59
Shortfin neoscopelid	<i>Neoscopelus microchir</i>	Peixe-lanterna-de-barbatanas-curtas	113
Shortgut fintail snipe eel	<i>Labichthys carinatus</i>	Cobra-de-crista	68
Short-headed lantern fish	<i>Diaphus brachycephalus</i>	Peixe-lanterna-de-cabeça-curta	116
Shorthorn sculpin	<i>Myoxocephalus scorpius</i>	Escorpião	237
Shortline swallower eel	<i>Saccopharynx ramosus</i>	Peixe-pelicano-de-cauda-ramosa	71
Shortnose greeneye	<i>Chlorophthalmus agassizi</i>	Olho-verde	105
Short-rod anglerfish	<i>Microlophichthys microlophus</i>	Peixe-farol-haste-curta	217
Shortspine spurdog	<i>Squalus mitsukurii</i>	Galhudo-de-espinho-curto	39
Shorttail skate	<i>Amblyraja jensenii</i>	Raia-de-cauda-curta	42
Short-tube scorpionfish	<i>Phenacoscorpius nebris</i>	Rascasso-de-tubo-curto	230
Shoulderspine bigscale	<i>Melamphaes suborbitalis</i>	Peixe-rã-de-ombro-espinhoso	151
Showy bristlemouth	<i>Cyclothona signata</i>	Boqueirão-vistoso	87
Sickle snaggletooth	<i>Borostomias mononema</i>	Peixe-dragão-abutre	96
Silky shark	<i>Carcharhinus falciformis</i>	Tubarão-luzidio	25
Silver driftfish	<i>Psenes maculatus</i>	Peixe-pastor-prateado	173
Silver hatchetfish	<i>Argyropelecus olfersii</i>	Pai-velho-prateado	90
Silver scabbardfish	<i>Lepidotus caudatus</i>	Peixe-espada	165
Silver spinyfin	<i>Diretmoides pauciradiatus</i>	Peixe-enfadado-barbatana-comprida	155
Silver spinyfin	<i>Diretmus argenteus</i>	Peixe-enfadado-prateado	156
Silvery John dory	<i>Zenopsis conchifer</i>	Galo-branco	130
Silvery lightfish	<i>Maurolicus muelleri</i>	Peixe-luz-de-Muller	91
Silvery pout	<i>Gadiculus argenteus</i>	Badejinho	142
Simony's frostfish	<i>Benthodesmus simonyi</i>	Espada-de-má-água	165
Singlespot frogfish	<i>Fowlerichthys radiosus</i>	Peixe-sapo-uniocelado	213
Skipjack tuna	<i>Katsuwonus pelamis</i>	Gaiado	167
Sladen's hatchet fish	<i>Argyropelecus sladeni</i>	Pai-velho-de-Sladen	90
Slender argentine	<i>Microstoma microstoma</i>	Argentina-fina	83
Slender bristlemouth	<i>Cyclothona pseudopallida</i>	Boqueirão-delgado	87
Slender codling	<i>Halargyreus johnsonii</i>	Abrótea-delgada	140
Slender cuskeel	<i>Porogadus miles</i>	Abadejo-fino	159
Slender finless eel	<i>Apterichtus anguiformis</i>	Cobra-de-boca-negra	64
Slender frostfish	<i>Benthodesmus tenuis</i>	Espada-de-má-água-elegante	165
Slender goby	<i>Gobius geniporus</i>	Caboz-de-bochechas porosas	180
Slender halosaur	<i>Aldrovandia gracilis</i>	Peixe-lagarto-delgado	56
Slender lightfish	<i>Vinciguerra attenuata</i>	Peixe-luminoso-esguio	92

English common name	Scientific name	Portuguese common name	page
Slender rockfish	<i>Scorpaena elongata</i>	Rascasso-rosado	230
Slender snaggletooth	<i>Rhadinesthes decimus</i>	Peixe-dragão-delgado	104
Slender snipe eel	<i>Nemichthys scolopaceus</i>	Cobra-de-bico	68
Slender snipefish	<i>Macroramphosus gracilis</i>	Trombeteiro-fino	174
Slender suckerfish	<i>Phtheirichthys lineatus</i>	Pegador-delgado	249
Slender sunfish	<i>Ranzania laevis</i>	Peixe-lua-comprido	222
Sloane's viperfish	<i>Chauliodus sloani</i>	Peixe-dragão-de-Sloane	97
Small lantern fish	<i>Diaphus holti</i>	Peixe-lanterna-de-Holt	117
Small red scorpionfish	<i>Scorpaena notata</i>	Rascasso-escorpião	232
Small sandeel	<i>Ammodytes tobianus</i>	Galeota-menor	291
Small scaled brown slickhead	<i>Alepocephalus australis</i>	Celindra-austral	74
Smalleye catshark	<i>Apristurus microps</i>	Tubarão-gato-olhinho	20
Smalleye smooth-head	<i>Alepocephalus productus</i>	Celindra-de-olho-pequeno	75
Smalleye squaretail	<i>Tetragonurus cuvieri</i>	Escolar-de-natura-olho-pequeno	173
Smalleyed rabbitfish	<i>Hydrolagus affinis</i>	Ratazana-da-fundura	54
Small-eyed ray	<i>Raja microocellata</i>	Raia-zimbreira	47
Smallfin lanternfish	<i>Benthosema suborbitale</i>	Peixe-lanterna-corcunda	114
Small-headed clingfish	<i>Apletodon dentatus</i>	Sugador-de-cabeça-pequena	204
Smallmouth knifefooth dogfish	<i>Zameus squamulosus</i>	Arreganhada-de-focinho-comprido	34
Smallmouth moonflounder	<i>Monolene microstoma</i>	Carta-de-boca-pequena	190
Smallmouth spiny eel	<i>Polyacanthonotus rissoanus</i>	Enguia-espinhosa-de-boca-pequena	58
Smallscale searsid	<i>Maulisia microlepis</i>	Celindra-de-tubo-de-escama-curta	80
Smallscale smooth-head	<i>Bathytroctes microlepis</i>	Celindra-de-escama-pequena	76
Small-spotted catshark	<i>Scyliorhinus canicula</i>	Pata-roxa	21
Smalltooth dragonfish	<i>Pachystomias microdon</i>	Peixe-dragão-dente-curto	102
Smalltooth sand tiger shark	<i>Carcharias taurus</i>	Tubarão-toiro	17
Smalltoothed argentine	<i>Glossanodon leioglossus</i>	Argentina-de-dentes-pequenos	82
Smooth dreamer	<i>Chaenophryne draco</i>	Peixe-farol-dragão	216
Smooth hammerhead	<i>Sphyraena zygaena</i>	Tubarão-martelo	29
Smooth lanternshark	<i>Etmopterus pusillus</i>	Xarinha-preta	31
Smooth puffer	<i>Lagocephalus laevigatus</i>	Peixe-bola-verde	224
Smooth sandeel	<i>Gymnammodytes semisquamatus</i>	Galeota-frachão	291
Smoothback angelshark	<i>Squatina oculata</i>	Anjo-de-malhas	39
Smoothcheek lanternfish	<i>Bolinichthys indicus</i>	Peixe-lanterna-do-Índico	114
Smoothtail mobula	<i>Mobula thurstoni</i>	Manta-de-rabo-liso	53
Snaggletooth	<i>Astronesthes neopogon</i>	Peixe-dragão-de-barbilho-plano	94
Snaggletooth	<i>Astronesthes niger</i>	Peixe-dragão-manchado	94
Snaggletooth	<i>Borostomias antarcticus</i>	Peixe-dragão-do-Antártico	96
Snaggletooth	<i>Astronesthes gemmifer</i>	Peixe-dragão-da-Terra Nova	94
Snaggletooth	<i>Astronesthes micropogon</i>	Peixe-dragão-de-barba-curta	94
Snake blenny	<i>Ophidion barbatum</i>	Abadejo-cobrelo	158
Snake mackerel	<i>Gempylus serpens</i>	Escolar-serpente	162
Snake pipefish	<i>Entelurus aequoreus</i>	Marinha-amarela	175
Snakefish	<i>Trachinocephalus myops</i>	Peixe-lagarto-de-focinho-rombo	109
Snouted sole	<i>Pegusa nasuta</i>	Linguado-narigudo	195
Snubnosed eel	<i>Simenchelys parasitica</i>	Fluta	59
Soft lanternfish	<i>Diaphus mollis</i>	Peixe-lanterna-mole	117
Soft leafvent angler	<i>Haplophryne mollis</i>	Peixe-sestro-macio	220
Softhead grenadier	<i>Malacocephalus laevis</i>	Lagartixa-do-mar-redonda	136
Softskin smooth-head	<i>Rouleina attrita</i>	Celindra-macia	78
Solenette	<i>Buglossidium luteum</i>	Língua-de-gato	192
Southern cut-throat eel	<i>Histiobranchus australis</i>	Enguia-da-fundura-austral	58
Spangled tubeshoulder	<i>Persplesia kopua</i>	Celindra-de-tubo-estrelada	81
Spanish ling	<i>Molva macrophthalmia</i>	Maruca-do-Mediterrâneo	146
Spark anglemouth	<i>Sigmops bathophilus</i>	Boqueirão-brilhante	88

English common name	Scientific name	Portuguese common name	
Spearfish remora	<i>Remora brachyptera</i>	Pegador-azul	249
Spearsnouted grenadier	<i>Coelorinchus labiatus</i>	Lagartixa-do-mar-de-focinho-comprido	132
Spiderfish	<i>Bathypterois dubius</i>	Peixe-tripé-filamentoso	105
Spiky oreo	<i>Neocyttus rhomboidalis</i>	Galo-espinhoso	129
Spined pygmy shark	<i>Squaliolus laticaudus</i>	Tubarão-anão	30
Spinetail lanternfish	<i>Notoscopelus caudispinosus</i>	Peixe-lanterna-de-cauda-espinhosa	125
Spiny butterfly ray	<i>Gymnura altavela</i>	Uge-manta	50
Spiny grenadier	<i>Coelorinchus parallelus</i>	Lagartixa-do-mar-espinhosa	133
Spiny gurnard	<i>Lepidotrigla dieuzeidei</i>	Ruivo-espinhoso	236
Spiny seadevil	<i>Neoceratias spinifer</i>	Diabo-marinho-espinhoso	215
Spinycheek soldierfish	<i>Corniger spinosus</i>	Peixe-soldado-espinhoso	156
Splendid alfonsino	<i>Beryx splendens</i>	Imperador-de-costa-estreita	149
Spotfin butterflyfish	<i>Chaetodon ocellatus</i>	Borboleta-de-risca-amarela	278
Spotfin flyingfish	<i>Cheilopogon furcatus</i>	Peixe-voador-de-barbatana-manchada	200
Spotfin frogfish	<i>Antennatus nummifer</i>	Peixe-sapo-salgicado	213
Spot-fin porcupinefish	<i>Diodon hystrix</i>	Peixe-ourizo-ponteados	224
Spothead lantern fish	<i>Diaphus metopoclampus</i>	Peixe-lanterna-de-cabeça-manchada	117
Spotlight lanternfish	<i>Diaphus lucidus</i>	Peixe-lanterna-cintilante	117
Spotted barracudina	<i>Arctozenus risso</i>	Barracudina-pintada	109
Spotted dragonet	<i>Callionymus maculatus</i>	Peixe-pau-malhado	293
Spotted flounder	<i>Citharus linguatula</i>	Carta-de-bico	186
Spotted lanternfish	<i>Myctophum punctatum</i>	Peixe-lanterna-manchado	124
Spotted pipefish	<i>Nerophis maculatus</i>	Marinha-sarapintada	176
Spotted seabass	<i>Dicentrarchus punctatus</i>	Robalo-baila	241
Spotted spiny dogfish	<i>Squalus acanthias</i>	Galhudo-malhado	38
Spotted tonguesole	<i>Sympodus nigrescens</i>	Língua-avessa	196
Spotted weever	<i>Trachinus araneus</i>	Peixe-aranha-pontuado	292
Spotted-fin rockfish	<i>Scorpaena stephanica</i>	Rascasso-de-barbatana-pontuada	233
Spurcheek lanternfish	<i>Bolinichthys photothorax</i>	Peixe-lanterna-esporão	114
Squeteague	<i>Cynoscion regalis</i>	Corvinata-real	275
Stareye lightfish	<i>Pollimyrus mauli</i>	Peixe-luminoso-de-Maul	92
Stargazer	<i>Uranoscopus scaber</i>	Cabeçudo	293
Staring pearleye	<i>Scopelarchus guentheri</i>	Olho-pérola-de-Guenther	107
Starry ray	<i>Amblyraja radiata</i>	Raia-repregada	42
Starry smooth-head	<i>Photostylus pycnopterus</i>	Celindra-luminosa	78
Starry smooth-hound	<i>Mustelus asterias</i>	Cação-pintado	23
Starry weever	<i>Trachinus radiatus</i>	Peixe-aranha-raiado	292
Steven's goby	<i>Gobius gasteveni</i>	Caboz-de-escama	180
Stoplight loosejaw	<i>Malacosteus niger</i>	Peixe-dragão-de-draga	101
Stout beardedfish	<i>Polymixia nobilis</i>	Salmonete-do-alto	149
Straightnose pipefish	<i>Nerophis ophidion</i>	Marinha-fina	176
Straightnose rabbitfish	<i>Rhinochimaera atlantica</i>	Peixe-rato-narigudo	54
Streaked gurnard	<i>Chelidonichthys lastoviza</i>	Cabra-riscada	234
Striped burrfish	<i>Chilomycterus schoepfii</i>	Porco-espinho-listado	223
Striped escolar	<i>Diplospinus multistriatus</i>	Escolar-raiado	162
Striped seasnail	<i>Liparis liparis</i>	Peixe-lesma-listado	237
Stubby lanternfish	<i>Bolinichthys supralateralis</i>	Peixe-lanterna-atarracado	114
Sturgeon	<i>Acipenser sturio</i>	Esturjão	55
Surmullet	<i>Mullus surmuletus</i>	Salmonete-legítimo	277
Swallow grenadier	<i>Sphagemoncrurus hirundo</i>	Lagartixa-do-mar-dos-Açores	138
Swallowtail seaperch	<i>Anthias anthias</i>	Canário-do-mar	242
Swordfish	<i>Xiphias gladius</i>	Espadarte	184
Swordsnout grenadier	<i>Coelorinchus occa</i>	Lagartixa-do-mar-de-focinho-curto	133
Taaning's lantern fish	<i>Diaphus termophilus</i>	Peixe-lanterna-termófilo	118
Tadpole fish	<i>Raniceps raninus</i>	Rainúnculo-negro	143

English common name	Scientific name	Portuguese common name	page
Tadpole whiptail	<i>Squalogadus modicatus</i>	Lagartixa-do-mar-girino	138
Tan bristlemouth	<i>Cyclothona pallida</i>	Boqueirão-bicolor	87
Tåning's lanternfish	<i>Hygophum taanungi</i>	Peixe-lanterna-de-Tåning	120
Tapertail	<i>Radiiceps elongatus</i>	Peixe-remo-de-cauda-fina	128
Tarpon	<i>Megalops atlanticus</i>	Tarpão-do-Atlântico	55
Teardrop tubeshoulder	<i>Holtbyrnia innesi</i>	Celindra-de-tubo-de-Innes	80
Tentacle dragonfish	<i>Melanostomias tentaculatus</i>	Peixe-dragão-tentaculado	102
Tentacled blenny	<i>Parablennius tentacularis</i>	Marachomba-tentaculada	211
Thickback sole	<i>Microchirus variegatus</i>	Azevia-raiada	193
Thickbeard grenadier	<i>Coryphaenoides zaniophorus</i>	Lagartixa-do-mar-de-barbilho-grosso	135
Thicklip grey mullet	<i>Chelon labrosus</i>	Tainha-liça	202
Thinlip conger	<i>Gnathophis mystax</i>	Coreano	67
Thinlip grey mullet	<i>Chelon ramada</i>	Tainha-fataça	202
Thornback ray	<i>Raja clavata</i>	Raia-lenga	45
Thorny tinselfish	<i>Grammicolepis brachiusculus</i>	Peixe-prata-espinhoso	130
Thor's scaldfish	<i>Arnoglossus thori</i>	Carta-pontuada	190
Threadfin dragonfish	<i>Echiostoma barbatum</i>	Peixe-dragão-barbatana-filamentosa	97
Threadfin grenadier	<i>Gadomus longifilis</i>	Lagartixa-do-mar-de-bandeira	135
Threadfin smooth-head	<i>Talismania mekistonema</i>	Celindra-de-barbatana-filamentosa	79
Three-bearded rockling	<i>Gaidropsarus vulgaris</i>	Laibeque-de-três-barbilhos	145
Threelights dragonfish	<i>Trigonolampa miriceps</i>	Peixe-dragão-listado	104
Three-spined stickleback	<i>Gasterosteus aculeatus</i>	Esgana-gata	297
Tiger shark	<i>Galeocerdo cuvier</i>	Tubarão-tigre	27
Tompot blenny	<i>Parablennius gattorugine</i>	Marachomba-babosa	209
Toothed goby	<i>Deltentosteus collarinus</i>	Caboz-dentudo	179
Toothless smooth-head	<i>Einara edentula</i>	Celindra-desdentada	77
Tope shark	<i>Galeorhinus galeus</i>	Perna-de-moça	23
Topknot	<i>Zeugopterus punctatus</i>	Rodovalho-bruxa	188
Topside lanternfish	<i>Notolychnus valdiviae</i>	Peixe-lanterna-do-Valdívia	125
Transparent goby	<i>Aphia minuta</i>	Caboz-transparente	178
Tripletail	<i>Lobotes surinamensis</i>	Furriel	261
Triplewart seadevil	<i>Cryptopsaras couesii</i>	Peixe-farol-de-Coues	219
Tripodfish	<i>Bathypterois grallator</i>	Peixe-tripé	106
Tropical two-wing flyingfish	<i>Exocoetus volitans</i>	Andorinha-do-mar-tropical	201
Tub gurnard	<i>Chelidonichthys lucerna</i>	Cabra-cabaço	235
Tube-eye	<i>Stylephorus chordatus</i>	Peixe-cauda-de-fio	131
Turbot	<i>Scophthalmus maximus</i>	Pregado	187
Tusk	<i>Brosme brosme</i>	Bolota	144
Twaite shad	<i>Alosa fallax</i>	Savelha	72
Twinray dragonfish	<i>Eustomias satterleei</i>	Peixe-dragão-de-Satterlee	99
Two-spotted clingfish	<i>Diplecogaster bimaculata</i>	Sugador-de-duas-manchas	205
Two-spotted goby	<i>Gobiusculus flavescens</i>	Caboz-de-duas-manchas	182
Undistinguished sabretooth	<i>Odontostomops normalops</i>	Dente-de-sabre-olho-redondo	112
Undulate ray	<i>Raja undulata</i>	Raia-curva	48
Unicorn leatherjacket filefish	<i>Aluterus monoceros</i>	Peixe-porco-galhudo	226
Vadigo	<i>Campogramma glayo</i>	Xareu-palheta	250
Vaillant's tongueless smooth-head	<i>Leptochilichthys pinguis</i>	Celindra-de-Vaillant	77
Vaillant's grenadier	<i>Bathygadus melanobranchus</i>	Lagartixa-do-mar-de-Vaillant	131
Valdivia black dragon fish	<i>Melanostomias valdiviae</i>	Peixe-dragão-de-Valdívia	102
Veiled anglemouth	<i>Cyclothona microdon</i>	Boqueirão-de-dentes-pequenos	87
Velvet belly	<i>Etomopterus spinax</i>	Lixinha-da-fundura	31
Velvet whalefish	<i>Barbourisia rufa</i>	Peixe-baleia-veludo	153
Vermilion snapper	<i>Rhomboplites aurorubens</i>	Luciano-rosa	261
Victory whiptail	<i>Asthenomacrurus victoris</i>	Largartixa-do-mar-de-Vitória	131
Wahoo	<i>Acanthocybium solandri</i>	Serra-da-Índia	166

English common name	Scientific name	Portuguese common name	page
Waistcoat lanternfish	<i>Taaningichthys minimus</i>	Peixe-lanterna-de-Cabo-Verde	127
Warming's lanternfish	<i>Ceratoscopelus warmingii</i>	Peixe-lanterna-de-Warming	115
Warty dory	<i>Allocyttus verrucosus</i>	Galo-verrugoso	129
Waryfish	<i>Scopelosaurus argenteus</i>	Peixe-vigia-prateado	107
Wedge sole	<i>Dicologlossa cuneata</i>	Língua	192
Werner's smooth-head	<i>Conocara wernerii</i>	Celindra-de-Werner	77
West African Spanish mackerel	<i>Scomberomorus tritor</i>	Serra-branca	169
Western softhead grenadier	<i>Malacocephalus occidentalis</i>	Lagartixa-do-mar-occidental	136
Whale shark	<i>Rhincodon typus</i>	Tubarão-baleia	17
Whalefish	<i>Cetomimus hempheli</i>	Peixe-baleia-de-Hempel	154
Whalehead dreamer	<i>Lophodolos acanthognathus</i>	Peixe-farol-cabeça-de-baleia	217
Whalesucker	<i>Remora australis</i>	Pegador-das-baleias	249
Whipsnout sorcerer	<i>Venefica proboscidea</i>	Cobra-de-bico-comprido	65
Whiskered sole	<i>Monochirius hispidus</i>	Cascarra	194
Whiskered sole	<i>Monochirius atlanticus</i>	Cascarra-do-Atlântico	194
White barbel	<i>Galeichthys feliceps</i>	Bagre-barba-branca	82
White grouper	<i>Epinephelus aeneus</i>	Garoupa-legítima	243
White seabream	<i>Diplodus caeruleus</i>	Sargo-legítimo-do-Atlântico	266
White seabream	<i>Diplodus lineatus</i>	Sargo-de-Cabo-Verde	267
White seabream	<i>Diplodus sargus</i>	Sargo-legítimo-do-Mediterrâneo	267
White skate	<i>Rostroraja alba</i>	Raia-tairoga	48
White suckerfish	<i>Remora albescens</i>	Pegador-branco	249
White trevally	<i>Pseudocaranx dentex</i>	Xareu-bicudo	253
White-eyed moray	<i>Muraena augusti</i>	Moreia-da-Macaronésia	62
White-spotted lantern fish	<i>Diaphus rafinesquii</i>	Peixe-lanterna-de-Rafinesque	118
Whiting	<i>Merlangius merlangus</i>	Badejo	143
Wide-eyed flounder	<i>Bothus podas</i>	Carta-de-olhos-grandes	190
Winged lanternfish	<i>Lampanyctus alatus</i>	Peixe-lanterna-alado	121
Wingfin snaggletooth	<i>Heterophotus ophistoma</i>	Peixe-dragão-de-boca-de-serpente	100
Wisner's lantern fish	<i>Dasy scopelus selenops</i>	Peixe-lanterna-de-Wisner	116
Witch flounder	<i>Glyptocephalus cynoglossus</i>	Solhão	191
Worm pipefish	<i>Nerophis lumbriciformis</i>	Marinha-lombriga	176
Wreckfish	<i>Polyprion americanus</i>	Cherne	239
Yellowfin tuna	<i>Thunnus albacares</i>	Atum-albacora	170
Yellowmouth barracuda	<i>Sphyraena viridensis</i>	Bicuda-de-Cabo-Verde	296
Yellowstem dragonfish	<i>Eustomias macrurus</i>	Peixe-dragão-de-barbijo-amarelo	99
Yellowtail amberjack	<i>Seriola lalandi</i>	Charuteiro-azeite	255
Zebra seabream	<i>Diplodus cervinus</i>	Sargo-veado	266
Zugmayer's pearleye	<i>Benthalbella infans</i>	Olho-pérola-de-Zugmayer	106

Index of the Portuguese names / Índice dos nomes Portugueses

Portuguese common name	Scientific name	English common name	Page
Abadejo-abissal	<i>Abyssobrotula galatheae</i>	No English common name	156
Abadejo-arco-íris	<i>Barathrites iris</i>	No English common name	156
Abadejo-boca-doce	<i>Bassozetus levistomatus</i>	No English common name	157
Abadejo-cabeçudo	<i>Bathyonus laticeps</i>	No English common name	157
Abadejo-cego-da-Madeira	<i>Nybelinella brevianalis</i>	No English common name	161
Abadejo-cego-de-dorsal-longa	<i>Meteoria longidorsalis</i>	No English common name	161
Abadejo-cego-de-Eriksson	<i>Nybelinella eriksoni</i>	No English common name	161
Abadejo-cego-dentudo	<i>Barathronus multidens</i>	No English common name	160
Abadejo-cego-de-Roule	<i>Barathronus roulei</i>	No English common name	160
Abadejo-cego-de-Vaillant	<i>Barathronus parfaiti</i>	No English common name	160
Abadejo-cego-do-Pacifico	<i>Sciadonus pedicellaris</i>	No English common name	161
Abadejo-cego-dos-Açores	<i>Meteoria erythrops</i>	No English common name	161
Abadejo-cego-gelatinoso	<i>Aphyonus gelatinosus</i>	Gelatinous blennifish	160
Abadejo-cobrelo	<i>Ophidion barbatum</i>	Snake blenny	158
Abadejo-cometa	<i>Benthocometes robustus</i>	No English common name	157
Abadejo-cristado	<i>Lamprogrammus niger</i>	No English common name	158
Abadejo-da-fundura	<i>Bassozetus compressus</i>	Abyssal assfish	157
Abadejo-da-Mauritânia	<i>Monomitopus metriostoma</i>	No English common name	158
Abadejo-de-cauda-curta	<i>Brotulotaenia brevicauda</i>	No English common name	157
Abadejo-de-Nielsen	<i>Bassozetus nielseni</i>	Masked assfish	157
Abadejo-de-olho-pequeno	<i>Penopus microphthalmus</i>	No English common name	158
Abadejo-dos-Açores	<i>Spectrunculus crassus</i>	No English common name	159
Abadejo-escamudo	<i>Holcomycteronus squamosus</i>	No English common name	158
Abadejo-espinhososo	<i>Acanthonus armatus</i>	Bony-eared assfish	156
Abadejo-fino	<i>Porogadus miles</i>	Slender cuskeel	159
Abadejo-grande	<i>Spectrunculus grandis</i>	Pudgy cuskeel	159
Abadejo-grosso	<i>Brotulotaenia crassa</i>	Blue cusk eel	158
Abadejo-negro	<i>Brotulotaenia nigra</i>	Dark cusk	158
Abadejo-ténia	<i>Bassozetus taenia</i>	No English common name	157
Abadejo-vermelho	<i>Parophidion vassali</i>	No English common name	158
Abrótea-amuada	<i>Rhynchogadus hepaticus</i>	No English common name	142
Abrótea-cabeçuda	<i>Guttiagadus latifrons</i>	No English common name	140
Abrótea-da-costa	<i>Phycis phycis</i>	Forkbeard	147
Abrótea-de-Dalwigk	<i>Physiculus dalwigki</i>	Black codling	142
Abrótea-de-Günther	<i>Lepidion guentheri</i>	Guenther's codling	141
Abrótea-delgada	<i>Halargyreus johnsonii</i>	Slender codling	140
Abrótea-de-natura	<i>Gadella maraldi</i>	Gadella	139
Abrótea-de-Schmidt	<i>Lepidion schmidti</i>	Giant lepidion	141
Abrótea-de-Yarrell	<i>Laemonema yarrellii</i>	Codling	140
Abrótea-do-alto	<i>Phycis blennoides</i>	Greater forkbeard	146
Abrótea-do-Mediterrâneo	<i>Lepidion lepidion</i>	Mediterranean codling	141
Abrótea-imberbe	<i>Gadella imberbis</i>	Beardless codling	139
Abrótea-robusta	<i>Laemonema robustum</i>	Robust cod	140
Aguilha	<i>Belone belone</i>	Garfish	199
Aguilha-de-quilha	<i>Platybelone argalus</i>	Keeltail needlefish	199
Aguilha-do-bico-curto	<i>Belone svervidovi</i>	Short-beaked garfish	199
Aguilha-do-Mediterrâneo	<i>Belone acus</i>	Garfish	198
Aguilhão-anão	<i>Scomberesox simulans</i>	Dwarf saury	198
Aguilhão-do-Atlântico	<i>Scomberesox saurus</i>	Atlantic saury	198
Aguilheta-crocodilo	<i>Tylosurus crocodilus</i>	Hound needlefish	200
Aguilheta-imperial	<i>Tylosurus acus</i>	Agujon needlefish	199
Alcarraz	<i>Apogon imberbis</i>	Cardinal fish	248
Anchova	<i>Pomatomus saltatrix</i>	Bluefish	248

Portuguese common name	Scientific name	English common name	Page
Andorinha-do-mar-oceânica	<i>Exocoetus obtusirostris</i>	Oceanic two-wing flyingfish	201
Andorinha-do-mar-tropical	<i>Exocoetus volitans</i>	Tropical two-wing flyingfish	201
Anjo	<i>Squatina squatina</i>	Angelshark	40
Anjo-de-malhas	<i>Squatina oculata</i>	Smoothback angelshark	39
Anjo-espinhoso	<i>Squatina aculeata</i>	Sawback angelshark	39
Areeiro	<i>Lepidorhombus whiffagonis</i>	Megrim	187
Areeiro-de-quatro-manchas	<i>Lepidorhombus boscii</i>	Four-spot megrim	187
Arenque-do-Atlântico	<i>Clupea harengus</i>	Atlantic herring	72
Argentina-atlântica	<i>Nansenia atlantica</i>	No English common name	83
Argentina-boreal	<i>Nansenia boreacrassicauda</i>	No English common name	83
Argentina-branca	<i>Argentina sphyraena</i>	Argentine	82
Argentina-da-fundura-de-focinho-comprido	<i>Dolicholagus longirostris</i>	Longsnout blacksmelt	84
Argentina-da-fundura-de-Grey	<i>Bathylagichthys greyae</i>	Grey's deepsea smelt	84
Argentina-da-fundura-escamuda	<i>Melanolagus bericoides</i>	Bigscale deepsea smelt	84
Argentina-da-fundura-negra	<i>Bathylagus euryops</i>	Goiter blacksmelt	84
Argentina-da-Groenlândia	<i>Nansenia groenlandica</i>	Greenland argentine	83
Argentina-de-cauda-longa	<i>Nansenia longicauda</i>	No English common name	83
Argentina-de-dentes-pequenos	<i>Glossanodon leioglossus</i>	Smalltoothed argentine	82
Argentina-do-Mediterrâneo	<i>Nansenia obliterata</i>	No English common name	83
Argentina-dourada	<i>Nansenia tenera</i>	No English common name	83
Argentina-fina	<i>Microstoma microstoma</i>	Slender argentine	83
Argentina-ibérica	<i>Nansenia iberica</i>	No English common name	83
Argentina-pelágica	<i>Nansenia cf. pelagica</i>	No English common name	83
Arreganhada	<i>Scymnodon ringens</i>	Knifetooth dogfish	33
Arreganhada-de-focinho-comprido	<i>Zameus squamulosus</i>	Smallmouth knifetooth dogfish	34
Atum-albacora	<i>Thunnus albacares</i>	Yellowfin tuna	170
Atum-de-barbatana-negra	<i>Thunnus atlanticus</i>	Blackfin tuna	170
Atum-patudo	<i>Thunnus obesus</i>	Bigeye tuna	170
Atum-rabilho	<i>Thunnus thynnus</i>	Atlantic bluefin tuna	170
Atum-voador	<i>Thunnus alalunga</i>	Albacore	169
Azevia	<i>Microchirus azevia</i>	Bastard sole	192
Azevia-de-malhas	<i>Microchirus ocellatus</i>	Foureyed sole	193
Azevia-de-olhos	<i>Microchirus hexophthalmus</i>	Ocellated wedge sole	193
Azevia-lusitanica	<i>Microchirus boscanion</i>	Lusitanian sole	193
Azevia-raiada	<i>Microchirus variegatus</i>	Thickback sole	193
Azevia-raiada-africana	<i>Microchirus wittei</i>	Banded sole	194
Bacalhau-do-Atlântico	<i>Gadus morhua</i>	Atlantic cod	142
Badejinho	<i>Gadiculus argenteus</i>	Silvery pout	142
Badejo	<i>Merlangius merlangus</i>	Whiting	143
Bagre-barba-branca	<i>Galeichthys feliceps</i>	White barbel	82
Barracudina-análoga	<i>Lestidiops similis</i>	No English common name	110
Barracudina-bico-de-pato	<i>Magnisudis atlantica</i>	Duckbill barracudina	110
Barracudina-da-Madeira	<i>Uncisudis longirostra</i>	No English common name	111
Barracudina-das-Canárias	<i>Macroparalepis affinis</i>	Barracudina	110
Barracudina-de-focinho-curto	<i>Paralepis brevirostris</i>	No English common name	110
Barracudina-dente-de-adaga	<i>Anoptopterus pharao</i>	Daggetooth	112
Barracudina-de-quatro-manchas	<i>Uncisudis quadrimaculata</i>	Barracudina	111
Barracudina-do-Atlântico	<i>Lestidium atlanticum</i>	Atlantic barracudina	110
Barracudina-do-Atlântico-nordeste	<i>Lestidiops pseudosphyraenoides</i>	No English common name	110
Barracudina-do-Mediterrâneo	<i>Lestidiops sphyrenoides</i>	Barracudina	110
Barracudina-do-Pacífico	<i>Lestidiops jayakari</i>	Pacific barracudina	109
Barracudina-feroz	<i>Sudis atrox</i>	Fierce pike smelt	111
Barracudina-imperial	<i>Sudis hyalina</i>	Barracudina	111
Barracudina-lanceta	<i>Paralepis coregonoides</i>	Lancet fish	111
Barracudina-manchada	<i>Paralepis speciosa</i>	No English common name	111

Portuguese common name	Scientific name	English common name	Page
Barracudina-negra	<i>Macroparalepis nigra</i>	No English common name	110
Barracudina-pequena	<i>Macroparalepis brevis</i>	No English common name	110
Barracudina-pintada	<i>Arctozenus risso</i>	Spotted barracudina	109
Barracudina-ruça	<i>Lestidiops affinis</i>	Barracudina	109
Barracudina-translúcida	<i>Paralepis elongata</i>	Barracudina	111
Barroso	<i>Centrophorus granulosus</i>	Gulper shark	35
Besugo	<i>Pagellus acarne</i>	Axillary seabream	270
Bica	<i>Pagellus erythrinus</i>	Common pandora	271
Bica-buço	<i>Pagellus bellottii</i>	Red pandora	270
Bicuda-africana	<i>Sphyraena guachancho</i>	Guanchane barracuda	295
Bicuda-de-Cabo-Verde	<i>Sphyraena viridensis</i>	Yellowmouth barracuda	296
Bicuda-europeia	<i>Sphyraena sphyraena</i>	European barracuda	295
Bicuda-gigante	<i>Sphyraena barracuda</i>	Great barracuda	295
Biqueirão	<i>Engraulis encrasicolus</i>	European anchovy	74
Bodião-canário	<i>Labrus mixtus</i>	Cuckoo wrasse	283
Bodião-cão	<i>Bodianus speciosus</i>	Blackbar hogfish	281
Bodião-cinzento	<i>Syphodus cinereus</i>	Grey wrasse	284
Bodião-das-ervas	<i>Syphodus rostratus</i>	No English common name	286
Bodião-de-boca-pequena	<i>Centrolabrus exoletus</i>	Rock cook	281
Bodião-dente-de-cão	<i>Bodianus scrofa</i>	Barred hogfish	280
Bodião-de-pinta	<i>Syphodus ocellatus</i>	No English common name	285
Bodião-do-alto	<i>Acantholabrus palloni</i>	Scale-rayed wrasse	280
Bodião-do-Mediterrâneo	<i>Syphodus mediterraneus</i>	Axillary wrasse	285
Bodião-dos-Açores	<i>Syphodus caeruleus</i>	No English common name	284
Bodião-fusco	<i>Labrus merula</i>	Brown wrasse	283
Bodião-manchado	<i>Syphodus roissali</i>	Five-spotted wrasse	285
Bodião-mosqueado	<i>Syphodus trutta</i>	Emerald wrasse	286
Bodião-pavão	<i>Syphodus tinca</i>	Peacock wrasse	286
Bodião-reticulado	<i>Labrus bergylta</i>	Ballan wrasse	282
Bodião-rupestre	<i>Ctenolabrus rupestris</i>	Goldsinny-wrasse	282
Bodião-tentilhão	<i>Syphodus bailloni</i>	Baillon's wrasse	284
Bodião-torto	<i>Labrus viridi</i>	Green wrasse	283
Bodião-verde	<i>Thalassoma pavo</i>	Ornate wrasse	286
Bodião-verdelho	<i>Lappanella fasciata</i>	Sharp-toothed wrasse	284
Bodião-vulgar	<i>Syphodus melops</i>	Corkwing wrasse	285
Boga-do-mar	<i>Boops boops</i>	Bogue	263
Bolota	<i>Brosme brosme</i>	Tusk	144
Boqueirão	<i>Centracanthus cirrus</i>	Curled picarel	264
Boqueirão-bicolor	<i>Cyclothona pallida</i>	Tan bristlemouth	87
Boqueirão-branco	<i>Cyclothona alba</i>	Pale bristlemouth	86
Boqueirão-brilhante	<i>Sigmops bathyphilus</i>	Spark anglemouth	88
Boqueirão-cabeçudo	<i>Margrethia obtusirostra</i>	Bighead portholefish	88
Boqueirão-comprido	<i>Diplophos taenia</i>	Pacific portholefish	88
Boqueirão-da-Madeira	<i>Manducus maderensis</i>	No English common name	88
Boqueirão-de-Brauer	<i>Cyclothona braueri</i>	Gerrick	86
Boqueirão-de-dentes-curvos	<i>Cyclothona acclinidens</i>	Benttooth bristlemouth	86
Boqueirão-de-dentes-pequenos	<i>Cyclothona microdon</i>	Veiled anglemouth	87
Boqueirão-delgado	<i>Cyclothona pseudopallida</i>	Slender bristlemouth	87
Boqueirão-dentudo	<i>Sigmops elongatus</i>	Elongated bristlemouth fish	89
Boqueirão-de-raio-longo	<i>Bonapartia pedaliota</i>	Longray fangjaw	86
Boqueirão-do-Atlântico	<i>Gonostoma atlanticum</i>	Atlantic fangjaw	88
Boqueirão-do-Mediterrâneo	<i>Gonostoma denudatum</i>	Bristlemouth	88
Boqueirão-lívido	<i>Cyclothona livida</i>	Bristlemouth	87
Boqueirão-sombreado	<i>Cyclothona parapallida</i>	Shadow bristlemouth	87
Boqueirão-sombrio	<i>Cyclothona obscura</i>	Hidden bristlemouth	87

Portuguese common name	Scientific name	English common name	Page
Boqueirão-vistoso	<i>Cyclothona signata</i>	Showy bristlemouth	87
Boquinho	<i>Luvarus imperialis</i>	Luvar	295
Borboleta-de-risca-amarela	<i>Chaetodon ocellatus</i>	Spotfin butterflyfish	278
Borboleta-do-recife	<i>Chaetodon sedentarius</i>	Reef butterflyfish	278
Brótula-coxa	<i>Bellottia apoda</i>	No English common name	160
Brótula-de-Allen	<i>Cataetyx allenii</i>	No English common name	160
Brótula-de-cabeça-chata	<i>Cataetyx laticeps</i>	No English common name	160
Brótula-falsa-branca	<i>Leucobrotula adipata</i>	No English common name	161
Brótula-falsa-dos-Açores	<i>Parabrotula plagiophthalma</i>	False cusk	161
Brótula-negra	<i>Grammonus ater</i>	No English common name	161
Bruxa	<i>Zeugopterus regius</i>	Eckström's topknot	188
Cabeçudo	<i>Uranoscopus scaber</i>	Stargazer	293
Caboz-anão	<i>Lebetus scorpioides</i>	Diminutive goby	182
Caboz-cabeça-de-cobra	<i>Gobius ophiocephalus</i>	Grass goby	181
Caboz-cabeçudo	<i>Gobius cobitis</i>	Giant goby	180
Caboz-comum	<i>Pomatoschistus microps</i>	Common goby	183
Caboz-da-areaia	<i>Pomatoschistus minutus</i>	Sand goby	184
Caboz-da-Madeira	<i>Mauligobius maderensis</i>	No English common name	183
Caboz-da-rocha	<i>Gobius paganellus</i>	Rock goby	181
Caboz-das-Canárias	<i>Vanneaugobius canariensis</i>	No English common name	184
Caboz-de-Bellotti	<i>Gobius ater</i>	Bellotti's goby	179
Caboz-de-boca-vermelha	<i>Gobius cruentatus</i>	Red-mouthed goby	180
Caboz-de-bochechas porosas	<i>Gobius geniporus</i>	Slender goby	180
Caboz-de-Brito	<i>Chromogobius britoi</i>	Brito's goby	179
Caboz-de-cabeça-dourada	<i>Gobius xanthocephalus</i>	Golden goby	182
Caboz-de-Couch	<i>Gobius couchi</i>	Couch's goby	180
Caboz-de-cristal	<i>Crystallogobius linearis</i>	Crystal goby	179
Caboz-de-De Buen	<i>Buenia affinis</i>	De Buen's goby	178
Caboz-de-duas-manchas	<i>Gobiusculus flavesiensis</i>	Two-spotted goby	182
Caboz-de-escama	<i>Gobius gasteveni</i>	Steven's goby	180
Caboz-de-escama-grande	<i>Lesueurigobius friesii</i>	Fries's goby	182
Caboz-de-GUILLET	<i>Lebetus guilleti</i>	Guillet's goby	182
Caboz-de-Jeffrey	<i>Buenia jeffreysii</i>	Jeffrey's goby	179
Caboz-de-Kner	<i>Pomatoschistus knerii</i>	Kner's goby	183
Caboz-de-Lesueur	<i>Lesueurigobius suerii</i>	Lesueur's goby	183
Caboz-de-Lozano	<i>Pomatoschistus lozanoi</i>	Lozano's goby	183
Caboz-de-manchas	<i>Pomatoschistus pictus</i>	Mediterranean painted goby	184
Caboz-dentudo	<i>Deltentosteus collaris</i>	Toothed goby	179
Caboz-de-pinta-dourada	<i>Gnatholepis thompsoni</i>	Goldspot goby	179
Caboz-de-quatro-manchas	<i>Deltentosteus quadrimaculatus</i>	Four-spotted goby	179
Caboz-de-Roule	<i>Gobius roulei</i>	Roule's goby	181
Caboz-de-Sanzo	<i>Lesueurigobius sanzi</i>	Sanzo's goby	182
Caboz-de-três-dorsais-de-Delais	<i>Tripterygion delaisi</i>	Black-faced blenny	206
Caboz-de-três-dorsais-do-Mediterrâneo	<i>Tripterygion tripteronotum</i>	No English common name	206
Caboz-do-rio	<i>Neogobius fluviatilis</i>	Monkey goby	183
Caboz-dourado	<i>Gobius auratus</i>	Golden goby	180
Caboz-leopardo	<i>Thorogobius ephippiatus</i>	Leopard-spotted goby	184
Caboz-listado	<i>Lesueurigobius heterofasciatus</i>	No English common name	182
Caboz-marmoreado	<i>Pomatoschistus marmoratus</i>	Marbled goby	183
Caboz-mosqueado	<i>Gobius buccichi</i>	Buccich's goby	180
Caboz-negro	<i>Gobius niger</i>	Black goby	181
Caboz-norueguês	<i>Pomatoschistus norvegicus</i>	Norway goby	184
Caboz-transparente	<i>Aphia minuta</i>	Transparent goby	178
Cabra-cabaço	<i>Chelidonichthys lucerna</i>	Tub gurnard	235
Cabra-de-bandeira	<i>Chelidonichthys obscurus</i>	Longfin gurnard	235

Portuguese common name	Scientific name	English common name	Page
Cabra-de-casca	<i>Peristedion cataphractum</i>	African armoured searobin	236
Cabra-lira	<i>Trigla lyra</i>	Piper gurnard	236
Cabra-morena	<i>Eutrigla gurnardus</i>	Grey gurnard	235
Cabra-riscada	<i>Chelidonichthys lastoviza</i>	Streaked gurnard	234
Cabra-vermelha	<i>Chelidonichthys cuculus</i>	Red gurnard	234
Cabrinha-de-leque	<i>Dactylopterus volitans</i>	Flying gurnard	178
Cação-cachorro	<i>Mustelus canis</i>	Dusky smooth-hound	23
Cação-liso	<i>Mustelus mustelus</i>	Dusky smooth-hound	24
Cação-pintado	<i>Mustelus asterias</i>	Starry smooth-hound	23
Cação-ponteado	<i>Mustelus punctulatus</i>	Blackspotted smooth-hound	24
Cachucho	<i>Dentex macrophthalmus</i>	Large-eye dentex	265
Cachucho-dentão	<i>Dentex maroccanus</i>	Morocco dentex	265
Calafate-das-Canárias	<i>Umbrina canariensis</i>	Canary drum	276
Calafate-das-riscas	<i>Umbrina cirrosa</i>	Shi drum	276
Calafate-fosco	<i>Umbrina ronchus</i>	Fusca drum	276
Canarinho-do-mar	<i>Callanthias ruber</i>	Parrot seaperch	247
Canário-do-mar	<i>Anthias anthias</i>	Swallowtail seaperch	242
Cangulo-áspero	<i>Canthidermis maculata</i>	Rough triggerfish	228
Cangulo-cinzento	<i>Balistes capriscus</i>	Grey triggerfish	227
Cangulo-do-sargaço	<i>Xanthichthys ringens</i>	Sargassum triggerfish	228
Cangulo-oceânico	<i>Canthidermis sufflamen</i>	Ocean triggerfish	228
Cangulo-pintado	<i>Balistes punctatus</i>	Bluespotted triggerfish	227
Cangulo-real	<i>Balistes vetula</i>	Queen triggerfish	228
Cantarilho-espinhososo	<i>Trachyscorpia cristulata</i>	Atlantic thornyhead	229
Cantarilho-legítimo	<i>Helicolenus dactylopterus</i>	Blackbelly rosefish	228
Cantarilho-requeime	<i>Pontinus kuhlii</i>	Offshore rockfish	230
Capatão-de-bandeira	<i>Dentex gibbosus</i>	Pink dentex	264
Capatão-legítimo	<i>Dentex dentex</i>	Common dentex	264
Carapau	<i>Trachurus trachurus</i>	Atlantic horse mackerel	257
Carapau-do-Mediterrâneo	<i>Trachurus mediterraneus</i>	Mediterranean horse mackerel	256
Carapau-negrão	<i>Trachurus picturatus</i>	Blue jack mackerel	257
Carapau-rugoso	<i>Trachurus lathami</i>	Rough scad	256
Carocho	<i>Centroscymnus coelolepis</i>	Portuguese dogfish	32
Carta-de-bico	<i>Citharus linguatula</i>	Spotted flounder	186
Carta-de-boca-pequena	<i>Monolene microstoma</i>	Smallmouth moonflounder	190
Carta-de-Grohmann	<i>Arnoglossus grohmanni</i>	No English common name	188
Carta-de-olhos-grandes	<i>Bothus podas</i>	Wide-eyed flounder	190
Carta-do-Mediterrâneo	<i>Arnoglossus laterna</i>	Mediterranean scaldfish	189
Carta-estreita	<i>Arnoglossus rueppelii</i>	Rüppell's scaldback	189
Carta-imperial	<i>Arnoglossus imperialis</i>	Imperial scaldfish	189
Carta-moira	<i>Syacium micrurum</i>	Channel flounder	188
Carta-pelícano	<i>Chascanopsetta lugubris</i>	Pelican flounder	190
Carta-pontuada	<i>Arnoglossus thori</i>	Thor's scaldfish	190
Carta-sombria	<i>Syacium papillosum</i>	Dusky flounder	188
Cascarra	<i>Monochirius hispidus</i>	Whiskered sole	194
Cascarra-do-Atlântico	<i>Monochirius atlanticus</i>	Whiskered sole	194
Castanheta-báia	<i>Chromis chromis</i>	Damsel fish	278
Castanheta-da-Madeira	<i>Chromis limbata</i>	Azores chromis	279
Castanheta-preta	<i>Similiiparma lurida</i>	Canary damsel	279
Castanheta-raiada	<i>Abudefduf saxatilis</i>	Sergeant-major	278
Cavala	<i>Scomber colias</i>	Atlantic chub mackerel	168
Cavalo-marinho-de-focinho-comprido	<i>Hippocampus guttulatus</i>	Long-snouted seahorse	175
Cavalo-marinho-de-focinho-curto	<i>Hippocampus hippocampus</i>	Short snouted seahorse	176
Cavalo-marinho-estriado	<i>Hippocampus erectus</i>	Lined Seahorse	175
Celindra-abissal	<i>Rinocutes nasutus</i>	Abyssal smooth-head	78

Portuguese common name	Scientific name	English common name	Page
Celindra-austral	<i>Alepocephalus australis</i>	Small scaled brown slickhead	74
Celindra-da-Madeira	<i>Rouleina maderensis</i>	Madeiran smooth-head	79
Celindra-da-Mauritânia	<i>Talismania homoptera</i>	Hairfin smooth-head	79
Celindra-das-Antilhas	<i>Talismania antillarum</i>	Antillean smooth-head	79
Celindra-de-Agassiz	<i>Leptochilichthys agassizii</i>	Agassiz's tongueless smooth-head	77
Celindra-de-Baird	<i>Alepocephalus bairdii</i>	Baird's slickhead	74
Celindra-de-barbatana-comprida	<i>Conocara macropterum</i>	Longfin smooth-head	76
Celindra-de-barbatana-filamentosa	<i>Talismania mekistonema</i>	Threadfin smooth-head	79
Celindra-de-cabeça-azul	<i>Asquamiceps caeruleus</i>	No English common name	75
Celindra-de-cabeça-negra	<i>Narcetes stomias</i>	Blackhead salmon	78
Celindra-de-Dana	<i>Bathyprion danae</i>	Fangtooth smooth-head	76
Celindra-de-escama-caduca	<i>Einara macrolepis</i>	Loosescale smooth-head	77
Celindra-de-escama-pequena	<i>Bathytroctes microlepis</i>	Smallscale smooth-head	76
Celindra-de-Fiolenti	<i>Conocara fiolenti</i>	Fiolenti's smooth-head	76
Celindra-de-focinho-curto	<i>Xenodermichthys copei</i>	Bluntsnout smooth-head	79
Celindra-de-Goa	<i>Narcetes erimelas</i>	No English common name	78
Celindra-de-Hjort	<i>Asquamiceps hjorti</i>	Barethroat slickhead	75
Celindra-de-Krefft	<i>Herwigia krefftii</i>	Krefft's smooth-head	77
Celindra-de-leque	<i>Asquamiceps velaris</i>	Fanfin smooth-head	75
Celindra-de-Michael-Sars	<i>Bathytroctes michaelsarsi</i>	Michael Sars smooth-head	76
Celindra-de-Murray	<i>Conocara murrayi</i>	Murray's smooth-head	77
Celindra-de-Norman	<i>Mirognathus normani</i>	Norman's smooth-head	78
Celindra-de-olho-grande	<i>Bajacalifornia megalops</i>	Bigeye smooth-head	75
Celindra-de-olho-pequeno	<i>Alepocephalus productus</i>	Smalleye smooth-head	75
Celindra-de-Risso	<i>Alepocephalus rostratus</i>	Risso's smooth-head	75
Celindra-desdentada	<i>Einara edentula</i>	Toothless smooth-head	77
Celindra-de-tubo-bicuda	<i>Mentodus rostratus</i>	Beaked searsid	81
Celindra-de-tubo-cabeçuda	<i>Holtbyrnia anomala</i>	Bighead searsid	80
Celindra-de-tubo-curva	<i>Platytroctes apus</i>	Legless searsid	81
Celindra-de-tubo-de-cabeça-negra	<i>Holtbyrnia melanocephala</i>	No English common name	80
Celindra-de-tubo-de-escama-curta	<i>Maulisia microlepis</i>	Smallscale searsid	80
Celindra-de-tubo-de-Innes	<i>Holtbyrnia innesi</i>	Teardrop tubeshoulder	80
Celindra-de-tubo-de-Koefoed	<i>Searsia koefoedi</i>	Koefoed's searsid	82
Celindra-de-tubo-de-Maul	<i>Maulisia mauli</i>	Maul's searsid	80
Celindra-de-tubo-de-nervura	<i>Mentodus mesalirurus</i>	Midridged searsid	81
Celindra-de-tubo-de-olhuda	<i>Holtbyrnia macrops</i>	Bigeye searsid	80
Celindra-de-tubo-de-Schnakenbeck	<i>Sagamichthys schnakenbecki</i>	Schnakenbeck's searsid	81
Celindra-de-tubo-de-ventre-claro	<i>Barbantus curvifrons</i>	Palebelly searsid	80
Celindra-de-tubo-dourada	<i>Maulisia argipalla</i>	Palegold searsid	80
Celindra-de-tubo-escura	<i>Mentodus facilis</i>	Dark searsid	81
Celindra-de-tubo-estrelada	<i>Perspersia kopua</i>	Spangled tubeshoulder	81
Celindra-de-tubo-multiporo	<i>Normichthys operosus</i>	Multipore searsid	81
Celindra-de-Vaillant	<i>Leptochilichthys pinguis</i>	Vaillant's tongueless smooth-head	77
Celindra-de-Werner	<i>Conocara werneri</i>	Werner's smooth-head	77
Celindra-dos-Açores	<i>Leptoderma macrophthalmum</i>	No English common name	78
Celindra-escamuda	<i>Bathytroctes macrolepis</i>	Koefoed's smooth-head	76
Celindra-luminosa	<i>Photostylus pycnopterus</i>	Starry smooth-head	78
Celindra-macia	<i>Rouleina attrita</i>	Softskin smooth-head	78
Celindra-negra	<i>Bathylaco nigricans</i>	Black warrior	76
Celindra-negra-de-Agassiz	<i>Alepocephalus agassizii</i>	Agassiz' slickhead	74
Celindra-olhuda	<i>Leptoderma macrops</i>	Grenadier smooth-head	78
Celindra-salmonada	<i>Conocara salmonicum</i>	Salmon smooth-head	77
Charéu-de-natura	<i>Caranx sexfasciatus</i>	Bigeye trevally	252
Charroco	<i>Halobatrachus didactylus</i>	Lusitanian toadfish	162
Charro-de-cauda-vermelha	<i>Decapterus tabl</i>	Roughearth scad	252

Portuguese common name	Scientific name	English common name	Page
Charro-moiro	<i>Decapterus punctatus</i>	Round scad	252
Charro-olho-de-boi	<i>Selar boops</i>	Oxeye scad	254
Charro-olho-largo	<i>Decapterus macarellus</i>	Mackerel scad	252
Charro-preto	<i>Selar crumenophthalmus</i>	Bigeye scad	254
Charuteiro-azeite	<i>Seriola lalandi</i>	Yellowtail amberjack	255
Charuteiro-catarino	<i>Seriola dumerili</i>	Greater amberjack	254
Charuteiro-da-Guiné	<i>Seriola carpenteri</i>	Guinean amberjack	254
Charuteiro-falcado	<i>Seriola rivoliana</i>	Longfin yellowtail	255
Charuteiro-listado	<i>Seriola fasciata</i>	Lesser amberjack	255
Charuteiro-olhete	<i>Seriola zonata</i>	Banded rudderfish	256
Cherne	<i>Polyprion americanus</i>	Wreckfish	239
Choupa	<i>Spondyliosoma cantharus</i>	Black seabream	274
Cobra-cabeça-de-pato	<i>Facciolella oxyrhynchus</i>	Facciola's sorcerer	65
Cobra-centopeia	<i>Avocettina infans</i>	Avocet snipe eel	67
Cobra-chumbo	<i>Myrophis plumbeus</i>	Leaden worm eel	64
Cobra-couraça	<i>Serrivomer beanii</i>	Bean's sawtooth eel	68
Cobra-de-bico	<i>Nemichthys scolopaceus</i>	Slender snipe eel	68
Cobra-de-bico-comprido	<i>Venefica proboscidea</i>	Whipsnout sorcerer	65
Cobra-de-boca-negra	<i>Apterichtus anguiformis</i>	Slender finless eel	64
Cobra-de-cauda-preta	<i>Nettastoma melanurum</i>	Blackfin sorcerer	65
Cobra-de-crista	<i>Labichthys carinatus</i>	Shortgut fintail snipe eel	68
Cobra-de-dentes-curtos	<i>Serrivomer lanceolatoides</i>	Bronze sawtooth eel	69
Cobra-de-leme	<i>Cyema atrum</i>	Bobtail eel	69
Cobra-de-orelhas	<i>Echelus myrus</i>	Painted eel	64
Cobra-do-mar	<i>Ophisurus serpens</i>	Serpent eel	64
Cobra-imberbe	<i>Dalophis imberbis</i>	Armless snake eel	64
Cobra-leopardo	<i>Myrichthys pardalis</i>	Leopard eel	64
Cobra-malhada	<i>Pisodonophis semicinctus</i>	Saddled snake eel	65
Cobra-pálida-ponteada	<i>Nemichthys curvirostris</i>	Pale threadtail snipe eel	68
Congrinho-bicolor	<i>Chlopsis bicolor</i>	Bicoloured false moray	63
Congrinho-da-areia	<i>Apterichtus caecus</i>	European finless eel	64
Congrinho-malhado	<i>Chlopsis dentatus</i>	Mottled false moray	63
Congo	<i>Conger conger</i>	European conger	66
Congo-branco-africano	<i>Cynoponticus ferox</i>	Guinean pike conger	65
Congo-castanho-dentão	<i>Bathyuroconger vicinus</i>	Long-tooth conger	66
Congo-da-Guiné	<i>Paraconger notialis</i>	Guinean conger	67
Congo-das-Baleares	<i>Ariosoma balearicum</i>	Bandtooth conger	65
Congo-de-boca-púrpura	<i>Pseudophichthys splendens</i>	Purplemouthed conger	67
Congo-de-natura	<i>Paraconger macrops</i>	Manypore conger	67
Congo-dentudo	<i>Conger triporiceps</i>	Manytooth conger	66
Congo-dos-Açores	<i>Gnathophis codoniphorus</i>	No English common name	67
Corcovado-africano	<i>Selene dorsalis</i>	African moonfish	254
Corcovado-do-Golfo	<i>Selene setapinnis</i>	Atlantic moonfish	254
Coreano	<i>Gnathophis mystax</i>	Thinlip conger	67
Corneta-malhada	<i>Fistularia tabacaria</i>	Cornetfish	174
Corneta-rosada	<i>Fistularia petimba</i>	Red cornetfish	174
Corvina-legítima	<i>Argyrosomus regius</i>	Meagre	275
Corvinata-real	<i>Cynoscion regalis</i>	Squeteague	275
Dentão-das-Canárias	<i>Dentex canariensis</i>	Canary dentex	264
Dente-de-sabre-de-Balbo	<i>Evermannella balbo</i>	Balbo sabretooth	112
Dente-de-sabre-do-Atlântico	<i>Coccarella atlantica</i>	Atlantic sabretooth	112
Dente-de-sabre-olho-redondo	<i>Odontostomops normalops</i>	Undistinguished sabretooth	112
Dente-de-sabre-ponteado	<i>Evermannella melanoderma</i>	Indian sabretooth	112
Diabo-marinho-de-Johnson	<i>Melanocetus johnsonii</i>	Humpback anglerfish	215
Diabo-marinho-de-Murray	<i>Melanocetus murrayi</i>	Deepsea blackdevil	215

Portuguese common name	Scientific name	English common name	Page
Diabo-marinho-espinhosos	<i>Neoceratias spinifer</i>	Spiny seadevil	215
Dobradiça	<i>Oblada melanura</i>	Saddled seabream	269
Doirado	<i>Coryphaena hippurus</i>	Common dolphinfish	258
Doirado-pampo	<i>Coryphaena equiselis</i>	Pompano dolphinfish	258
Dourada	<i>Sparus aurata</i>	Gilthead seabream	273
Enguia	<i>Anguilla anguilla</i>	European eel	69
Enguia-americana	<i>Anguilla rostrata</i>	American eel	69
Enguia-beiçuda-alva	<i>Lycenchelys alba</i>	No English common name	288
Enguia-beiçuda-de-Saldanha	<i>Pachycara saldanhai</i>	No English common name	288
Enguia-beiçuda-do-Atlântico	<i>Melanostigma atlanticum</i>	Atlantic soft pout	288
Enguia-beiçuda-termofílica	<i>Pachycara thermophilum</i>	No English common name	288
Enguia-bico-de-pato	<i>Nessorhamphus ingolfianus</i>	Duckbill oceanic eel	63
Enguia-da-fundura	<i>Histiobranchus bathybius</i>	Deep-water arrowtooth eel	58
Enguia-da-fundura-austral	<i>Histiobranchus australis</i>	Southern cut-throat eel	58
Enguia-da-fundura-castanha	<i>Ilyophis brunneus</i>	Muddy arrowtooth eel	58
Enguia-da-fundura-de-Bertin	<i>Monognathus bertini</i>	Bertin's one-jawed eel	70
Enguia-da-fundura-de-Blache	<i>Ilyophis blachei</i>	Blache's ooze ee	58
Enguia-da-fundura-de-Böhlke	<i>Monognathus boehlkei</i>	Böhlke's one-jawed eel	70
Enguia-da-fundura-de-Herring	<i>Monognathus herringi</i>	Herring's one-jawed eel	70
Enguia-da-fundura-de-Jespersen	<i>Monognathus jesperseni</i>	No English common name	70
Enguia-da-fundura-de-Nigel	<i>Monognathus nigeli</i>	Nigel's one-jawed eel	70
Enguia-de-casulo	<i>Myxine glutinosa</i>	Hagfish	15
Enguia-de-focinho-bolboso	<i>Dysomma brevirostre</i>	Batnose eel	58
Enguia-de-jardim	<i>Heterconger longissimus</i>	Brown garden eel	67
Enguia-de-pescoço-fino	<i>Derichthys serpentinus</i>	Narrownecked oceanic eel	63
Enguia-de-tromba-de-elefante	<i>Dysommina proboscideus</i>	No English common name	58
Enguia-espinhosa-de-boca-pequena	<i>Polyacanthonotus rissoanus</i>	Smallmouth spiny eel	58
Enguia-espinhosa-de-Bonaparte	<i>Notacanthus bonaparte</i>	Bonaparte's spiny eel	57
Enguia-espinhosa-de-Chemnitz	<i>Notacanthus chemnitzii</i>	Chemnitz's spiny eel	57
Enguia-espinhosa-de-Merrett	<i>Polyacanthonotus merretti</i>	Merrett's spiny eel	58
Enguia-espinhosa-do-Challenger	<i>Polyacanthonotus challengerii</i>	Challenger's spiny eel	57
Escolar	<i>Ruvettus pretiosus</i>	Oilfish	163
Escolar-branco	<i>Promethichthys prometheus</i>	Roudi escolar	163
Escolar-coelho-de-natura	<i>Nealotus tripes</i>	Black snake mackerel	163
Escolar-de-natura-olho-grande	<i>Tetragonurus atlanticus</i>	Bigeye squaretail	173
Escolar-de-natura-olho-pequeno	<i>Tetragonurus cuvieri</i>	Smalleye squaretail	173
Escolar-de-peitoral-longa	<i>Scomberolabrax heterolepis</i>	Longfin escolar	171
Escolar-geribé	<i>Nesiarchus nasutus</i>	Black gemfish	163
Escolar-preto	<i>Lepidocybium flavobrunneum</i>	Escolar	162
Escolar-raiado	<i>Diplospinus multistriatus</i>	Striped escolar	162
Escolar-serpente	<i>Gempylus serpens</i>	Snake mackerel	162
Escorpião	<i>Myoxocephalus scorpius</i>	Shorthorn sculpin	237
Escorpião-pálido	<i>Cottunculus thomsonii</i>	Pallid sculpin	237
Escorpião-roco	<i>Taurulus bubalis</i>	Longspined bullhead	237
Esgana-gata	<i>Gasterosteus aculeatus</i>	Three-spined stickleback	297
Esgana-gata-marinha	<i>Spinachia spinachia</i>	Sea stickleback	297
Espada-de-má-água	<i>Benthodesmus simonyi</i>	Simony's frostfish	165
Espada-de-má-água-comprido	<i>Benthodesmus elongatus</i>	Elongate frostfish	164
Espada-de-má-água-elegante	<i>Benthodesmus tenuis</i>	Slender frostfish	165
Espadarte	<i>Xiphias gladius</i>	Swordfish	184
Espadarte-serra	<i>Pristis pristis</i>	Common sawfish	42
Espadilha	<i>Sprattus sprattus</i>	European sprat	74
Espadim-azul-do-Atlântico	<i>Makaira nigricans</i>	Blue marlin	185
Espadim-bicudo	<i>Tetrapturus pfluegeri</i>	Longbill spearfish	186
Espadim-branco-do-Atlântico	<i>Kajikia albida</i>	Atlantic white marlin	185

Portuguese common name	Scientific name	English common name	Page
Espadim-do-Mediterrâneo	<i>Tetrapturus belone</i>	Mediterranean spearfish	186
Espadim-peto	<i>Tetrapturus georgii</i>	Roundscale spearfish	186
Esquilo-real	<i>Sargocentron hastatum</i>	Red squirrelfish	156
Esturjão	<i>Acipenser sturio</i>	Sturgeon	55
Esturjão-do-Adriático	<i>Acipenser naccarii</i>	Adriatic sturgeon	55
Faneca	<i>Trisopterus luscus</i>	Pouting	144
Fanecão	<i>Trisopterus minutus</i>	Poor cod	144
Fanecão-do-Mediterrâneo	<i>Trisopterus capelanus</i>	Poor cod Mediterranean	143
Ferreira	<i>Lithognathus mormyrus</i>	Sand steenbras	269
Fluta	<i>Simenchelys parasitica</i>	Snubnosed eel	59
Fogueteiro-arco-íris	<i>Elagatis bipinnulata</i>	Rainbow runner	253
Fogueteiro-galego	<i>Rachycentron canadum</i>	Cobia	249
Freira	<i>Pteraclis carolinus</i>	Fanfish	259
Freira-do-Atlântico	<i>Pterycombus brama</i>	Atlantic fanfish	259
Fura-vasos-alfonsim	<i>Cookeolus japonicus</i>	Longfinned bullseye	247
Fura-vasos-da-rocha	<i>Heteropriacanthus cruentatus</i>	Glasseye	247
Fura-vasos-da-rocha-brilhante	<i>Heteropriacanthus fulgens</i>	Bright glasseye	248
Fura-vasos-vulgar	<i>Priacanthus arenatus</i>	Atlantic bigeye	248
Furriel	<i>Lobotes surinamensis</i>	Tripletail	261
Gaiado	<i>Katsuwonus pelamis</i>	Skipjack tuna	167
Galeota-da-areia	<i>Gymnammodytes cicerelus</i>	Mediterranean sand eel	291
Galeota-frachão	<i>Gymnammodytes semisquamatus</i>	Smooth sandeel	291
Galeota-maior	<i>Hyperoplus lanceolatus</i>	Great sandeel	291
Galeota-maior-de-Corbin	<i>Hyperoplus immaculatus</i>	Greater sand-eel	291
Galeota-menor	<i>Ammodytes tobianus</i>	Small sandeel	291
Galhudo	<i>Squalus blainvilliei</i>	Longnose spurdog	38
Galhudo-de-espinho-curto	<i>Squalus mitsukurii</i>	Shortspine spurdog	39
Galhudo-de-focinho-curto	<i>Squalus megalops</i>	Picked spurdog	39
Galhudo-malhado	<i>Squalus acanthias</i>	Spotted spiny dogfish	38
Galito-rosado	<i>Zenion hololepis</i>	Dwarf dory	130
Galo-branco	<i>Zenopsis conchifer</i>	Silvery John dory	130
Galo-de-natura	<i>Cyttopsis rosea</i>	Red dory	129
Galo-espinhoso	<i>Neocyttus rhomboidalis</i>	Spiky oreo	129
Galo-irlandês	<i>Neocyttus helgae</i>	Irish oreo	129
Galo-negro	<i>Zeus faber</i>	John dory	130
Galo-verrugoso	<i>Allocyttus verrucosus</i>	Warty dory	129
Garoupa-chumbo	<i>Mycteroperca rubra</i>	Mottled grouper	245
Garoupa-de-Haifa	<i>Hyporthodus haifensis</i>	Haifa grouper	244
Garoupa-do-Níger	<i>Cephalopholis nigri</i>	Niger hind	242
Garoupa-dourada	<i>Epinephelus costae</i>	Goldblotch grouper	243
Garoupa-insular	<i>Mycteroperca fusca</i>	Island grouper	244
Garoupa-legítima	<i>Epinephelus aeneus</i>	White grouper	243
Garoupa-mariola	<i>Mycteroperca phenax</i>	Scamp	245
Garoupa-pintada-da-rocha	<i>Epinephelus adscensionis</i>	Rock hind	243
Gata	<i>Dalatias licha</i>	Kitefin shark	29
Gavião-do-mar	<i>Rhinoptera marginata</i>	Lusitanian cownose ray	51
Gissu-japonês	<i>Pterothrius gissu</i>	Japanese gissu	55
Goraz	<i>Pagellus bogaraveo</i>	Blackspot seabream	270
Imperador	<i>Beryx decadactylus</i>	Alfonsino	149
Imperador-de-costa-estreita	<i>Beryx splendens</i>	Splendid alfonsino	149
Judeu	<i>Auxis rochei</i>	Bullet tuna	166
Judeu-liso	<i>Auxis thazard</i>	Frigate tuna	166
Judia	<i>Coris julis</i>	Rainbow wrasse	281
Juliana	<i>Pollachius pollachius</i>	Pollack	143
Lagartixa-da-rocha	<i>Coryphaenoides rupestris</i>	Roundnose grenadier	134

Portuguese common name	Scientific name	English common name	Page
Lagartixa-do-mar-abissal	<i>Coryphaenoides armatus</i>	Abyssal grenadier	133
Lagartixa-do-mar-cabeçuda	<i>Coryphaenoides rudis</i>	Madeiran grenadier	134
Lagartixa-do-mar-carapina	<i>Coryphaenoides carapinus</i>	Carapine grenadier	133
Lagartixa-do-mar-da-fundura	<i>Coryphaenoides profundiculus</i>	Deepwater grenadier	134
Lagartixa-do-mar-da-Madeira	<i>Coryphaenoides thelestomus</i>	Roughlip grenadier	134
Lagartixa-do-mar-de-bandeira	<i>Gadomus longifilis</i>	Threadfin grenadier	135
Lagartixa-do-mar-de-barbilho-curto	<i>Coryphaenoides brevibarbis</i>	Shortbeard grenadier	133
Lagartixa-do-mar-de-barbilho-grosso	<i>Coryphaenoides zaniophorus</i>	Thickbeard grenadier	135
Lagartixa-do-mar-de-cabeça-redonda	<i>Cetonus globiceps</i>	Globehead grenadier	132
Lagartixa-do-mar-de-chicote	<i>Pseudonezumia flagellicauda</i>	No English common name	138
Lagartixa-do-mar-de-focinho-comprido	<i>Coelorinchus labiatus</i>	Spearsnouted grenadier	132
Lagartixa-do-mar-de-focinho-côncavo	<i>Coelorinchus caelorrhincus</i>	Hollowsnout grenadier	132
Lagartixa-do-mar-de-focinho-curto	<i>Coelorinchus occa</i>	Swordsnout grenadier	133
Lagartixa-do-mar-de-Günther	<i>Coryphaenoides guentheri</i>	Günther's grenadier	133
Lagartixa-do-mar-de-Murray	<i>Odontomacrus murrayi</i>	Roundhead grenadier	137
Lagartixa-do-mar-de-nariz-arrebitado	<i>Sphagmacrurus grenadae</i>	Pugnose grenadier	138
Lagartixa-do-mar-de-Vaillant	<i>Bathygadus melanobranchus</i>	Vaillant's grenadier	131
Lagartixa-do-mar-do-Mediterrâneo	<i>Coryphaenoides mediterraneus</i>	Mediterranean grenadier	134
Lagartixa-do-mar-dos-Açores	<i>Sphagmacrurus hirundo</i>	Swallow grenadier	138
Lagartixa-do-mar-elegante	<i>Hymenogadus gracilis</i>	Graceful grenadier	135
Lagartixa-do-mar-espinhosa	<i>Coelorinchus parallelus</i>	Spiny grenadier	133
Lagartixa-do-mar-favosa	<i>Bathygadus favosus</i>	No English common name	131
Lagartixa-do-mar-filamentosa	<i>Gadomus arcuatus</i>	Doublethread grenadier	135
Lagartixa-do-mar-girino	<i>Squalogadus modificatus</i>	Tadpole whiptail	138
Lagartixa-do-mar-ímpar	<i>Gadomus dispar</i>	No English common name	135
Lagartixa-do-mar-macia	<i>Echinomacrus mollis</i>	No English common name	135
Lagartixa-do-mar-ocidental	<i>Malacocephalus occidentalis</i>	Western softhead grenadier	136
Lagartixa-do-mar-olho-de-cebola	<i>Macrourus berglax</i>	Roughhead grenadier	136
Lagartixa-do-mar-pálida	<i>Coryphaenoides leptolepis</i>	Ghostly grenadier	134
Lagartixa-do-mar-peluda	<i>Trachonurus villosus</i>	Furry whiptail	138
Lagartixa-do-mar-prateada	<i>Hymenocephalus italicus</i>	Glasshead grenadier	135
Lagartixa-do-mar-redonda	<i>Malacocephalus laevis</i>	Softhead grenadier	136
Lagarto-do-mar	<i>Aulopus filamentosus</i>	Royal flagfin	104
Laibeque-de-cinco-barbilhos	<i>Ciliata mustela</i>	Fivebeard rockling	144
Laibeque-de-Maul	<i>Gaidropsarus mauli</i>	No English common name	145
Laibeque-de-três-barbilhos	<i>Gaidropsarus vulgaris</i>	Three-bearded rockling	145
Laibeque-do-Atlântico	<i>Gaidropsarus macrophthalmus</i>	Bigeye rockling	145
Laibeque-do-Mediterrâneo	<i>Gaidropsarus mediterraneus</i>	Shore rockling	145
Laibeque-dos-Açores	<i>Gaidropsarus granti</i>	Azores rockling	144
Lampreia-do-mar	<i>Petromyzon marinus</i>	Sea lamprey	15
Lampreia-do-rio	<i>Lampetra fluviatilis</i>	European river lamprey	15
Largartixa-do-mar-de-Vitória	<i>Asthenomacrus victoris</i>	Victory whiptail	131
Largatixa-do-mar-arrepiada	<i>Trachonurus sulcatus</i>	Bristly grenadier	138
Largatixa-do-mar-áspera	<i>Nezumia sclerorhynchus</i>	Roughtip grenadier	137
Largatixa-do-mar-de-Baird	<i>Nezumia bairdii</i>	Marlin-spike grenadier	137
Largatixa-do-mar-de-focinho-curto	<i>Nezumia longebarbata</i>	Bluntnose grenadier	137
Largatixa-do-mar-lisa	<i>Nezumia aequalis</i>	Common Atlantic grenadier	136
Lebre-preta	<i>Pomacanthus paru</i>	French angelfish	278
Leitão	<i>Galeus melastomus</i>	Blackmouth catshark	21
Leitão-da-Islândia	<i>Galeus murinus</i>	Mouse catshark	21
Leitão-do-Atlântico	<i>Galeus atlanticus</i>	Atlantic sawtail catshark	21
Língua	<i>Dicologlossa cuneata</i>	Wedge sole	192
Língua-avessa	<i>Syphurus nigrescens</i>	Spotted tonguesole	196
Língua-comprida	<i>Syphurus ligulatus</i>	Elongate tonguesole	196
Língua-da-Maraconésia	<i>Syphurus insularis</i>	Macaronesian tonguesole	196

Portuguese common name	Scientific name	English common name	Page
Língua-de-cão-das-Canárias	<i>Cynoglossus canariensis</i>	Canary tonguesole	196
Língua-de-cão-riscada	<i>Cynoglossus lineolatus</i>	No English common name	196
Língua-de-gato	<i>Buglossidium luteum</i>	Solenette	192
Língua-de-vaca	<i>Dagetichthys lusitanicus</i>	Portuguese sole	192
Linguado-da-areaia	<i>Pegusa lascaris</i>	Sand sole	194
Linguado-da-fundura	<i>Bathysolea profundicola</i>	Deep water sole	191
Linguado-do-Adriático	<i>Pegusa impar</i>	Adriatic sole	194
Linguado-do-Senegal	<i>Solea senegalensis</i>	Senegalese sole	195
Linguado-legítimo	<i>Solea solea</i>	Common sole	195
Linguado-narigudo	<i>Pegusa nasuta</i>	Snouted sole	195
Linguado-turco	<i>Synapturichthys kleinii</i>	Klein's sole	195
Língua-reticulada	<i>Syphurus reticulatus</i>	Reticulated tonguefish	197
Lírio	<i>Trichiurus lepturus</i>	Largehead hairtail	165
Lírio-ferro-de-focinho-comprido	<i>Alepisaurus ferox</i>	Long snouted lancetfish	113
Lírio-ferro-de-focinho-curto	<i>Alepisaurus brevirostris</i>	Short snouted lancetfish	113
Liro-errante	<i>Hyperoglyphe perciformis</i>	Barrelfish	172
Liro-imperial	<i>Schedophilus ovalis</i>	Imperial blackfish	172
Liro-manchado	<i>Schedophilus maculatus</i>	Pelagic butterfish	172
Liro-mole	<i>Schedophilus medusophagus</i>	Cornish blackfish	172
Liro-preto	<i>Centrolophus niger</i>	Rudderfish	171
Lixa	<i>Centrophorus squamosus</i>	Leafscale gulper shark	36
Lixinha-da-fundura	<i>Etmopterus spinax</i>	Velvet belly	31
Lixinha-da-fundura-grada	<i>Etmopterus princeps</i>	Great lanternshark	31
Lixinha-negra	<i>Centroscyllium fabricii</i>	Black dogfish	30
Luciano-da-Goreia	<i>Lutjanus goreensis</i>	Gorean snapper	261
Luciano-de-ventre-vermelho	<i>Lutjanus cyanopterus</i>	Cubera snapper	261
Luciano-rosa	<i>Rhomboplites aurorubens</i>	Vermilion snapper	261
Manta-cornuda	<i>Mobula tarapacana</i>	Chilean devil ray	52
Manta-de-rabo-liso	<i>Mobula thurstoni</i>	Smoothtail mobula	53
Manta-do-Mediterrâneo	<i>Mobula mobular</i>	Devil fish	52
Manta-gigante	<i>Mobula birostris</i>	Giant manta	52
Marachomba-babosa	<i>Parablennius gattorugine</i>	Tompot blenny	209
Marachomba-banana	<i>Hyleurochilus bananensis</i>	No English common name	207
Marachomba-borboleta	<i>Blennius ocellaris</i>	Butterfly blenny	207
Marachomba-branca	<i>Parablennius rouxi</i>	No English common name	210
Marachomba-cornuda	<i>Parablennius pilicornis</i>	Ringneck blenny	210
Marachomba-dálmata	<i>Microlipophrys dalmatinus</i>	No English common name	209
Marachomba-das-cracas	<i>Parablennius incognitus</i>	No English common name	209
Marachomba-das-poças	<i>Parablennius parvicornis</i>	Rock-pool blenny	210
Marachomba-das-rochas	<i>Coryphoblennius galerita</i>	Montagu's blenny	207
Marachomba-da-toca	<i>Hyleurochilus fissicornis</i>	No English common name	208
Marachomba-de-água-doce	<i>Salarias fluviatilis</i>	Freshwater blenny	211
Marachomba-de-Caneva	<i>Microlipophrys canevae</i>	No English common name	208
Marachomba-de-crista	<i>Scartella cristata</i>	Molly miller	212
Marachomba-do-Atlântico	<i>Ophioblennius atlanticus</i>	Redlip Blenny;	209
Marachomba-do-Mar-Negro	<i>Parablennius sanguinolentus</i>	Rusty blenny	211
Marachomba-esfinge	<i>Aidablennius sphynx</i>	No English common name	207
Marachomba-frade	<i>Lipophrys pholis</i>	Shanny	208
Marachomba-listada	<i>Salarias basilisca</i>	No English common name	211
Marachomba-pavão	<i>Salarias pavo</i>	Peacock blenny	212
Marachomba-peluda	<i>Labrisomus nuchipinnis</i>	Hairy blenny	206
Marachomba-portuguesa	<i>Parablennius ruber</i>	Portuguese blenny	210
Marachomba-prateada	<i>Clinitrichus argentatus</i>	Cline	207
Marachomba-tentaculada	<i>Parablennius tentacularis</i>	Tentacled blenny	211
Marachomba-trigloide	<i>Lipophrys trigloides</i>	No English common name	208

Portuguese common name	Scientific name	English common name	Page
Marinha-amarela	<i>Entelurus aequoreus</i>	Snake pipefish	175
Marinha-cabeça-chata	<i>Syngnathus rostellatus</i>	Nilsson's pipefish	178
Marinha-comum	<i>Syngnathus acus</i>	Greater pipefish	177
Marinha-de-focinho-estreito	<i>Syngnathus tenuirostris</i>	Narrow-snouted pipefish	178
Marinha-do-rio	<i>Syngnathus abaster</i>	Black-striped pipefish	176
Marinha-do-sargaço	<i>Syngnathus pelagicus</i>	Sargassum pipefish	177
Marinha-espinhosa	<i>Syngnathus phlegon</i>	No English common name	177
Marinha-fina	<i>Nerophis ophidion</i>	Straightnose pipefish	176
Marinha-focinho-grosso	<i>Syngnathus typhle</i>	Broadnosed pipefish	178
Marinha-lombriga	<i>Nerophis lumbriciformis</i>	Worm pipefish	176
Marinha-sarapintada	<i>Nerophis maculatus</i>	Spotted pipefish	176
Marinha-setentrional	<i>Syngnathus fuscus</i>	Northern pipefish	177
Maruca	<i>Molva molva</i>	Ling	146
Maruca-azul	<i>Molva dypterygia</i>	Blue ling	146
Maruca-do-Mediterrâneo	<i>Molva macrophthalmia</i>	Spanish ling	146
Meia-agulha-preta	<i>Hemiramphus balao</i>	Balao halfbeak	200
Merma	<i>Euthynnus alletteratus</i>	Little tunny	167
Mero-amarelo	<i>Epinephelus goreensis</i>	Dungat grouper	243
Mero-gigante	<i>Epinephelus caninus</i>	Dogtooth grouper	243
Mero-legítimo	<i>Epinephelus marginatus</i>	Dusky grouper	244
Mora	<i>Muraena moro</i>	Common mora	141
Mora-azul	<i>Antimora rostrata</i>	Blue antimora	139
Mordedor	<i>Xyrichtys novacula</i>	Pearly razorfish	287
Moreão-amarelo	<i>Gymnothorax vicinus</i>	Purplemouth moray	61
Moreão-bicudo	<i>Synaphobranchus dolichorhynchus</i>	No English common name	59
Moreão-de-dorsal-curta	<i>Synaphobranchus brevidorsalis</i>	Shortdorsal cutthroat eel	59
Moreão-de-natura	<i>Synaphobranchus kaupii</i>	Kaup's arrowtooth eel	59
Moreão-do-golfo	<i>Synaphobranchus affinis</i>	Grey cutthroat eel	59
Moreão-escuro	<i>Gymnothorax afer</i>	Dark moray	60
Moreia	<i>Muraena helena</i>	Mediterranean moray	62
Moreia-anã	<i>Anarchias similis</i>	Pygmy moray	60
Moreia-castanha	<i>Gymnothorax unicolor</i>	Brown moray	61
Moreia-da-Macaronésia	<i>Muraena augusti</i>	White-eyed moray	62
Moreia-da-Madeira	<i>Gymnothorax maderensis</i>	Sharktooth moray	61
Moreia-das-Canárias	<i>Gymnothorax bacalladoi</i>	Canary moray	60
Moreia-de-cauda-longa	<i>Anarchias longicauda</i>	Pygmy moray	60
Moreia-dourada	<i>Gymnothorax miliaris</i>	Goldentail moray	61
Moreia-pintada	<i>Muraena melanotis</i>	Honeycomb moray	63
Moreia-poligonal	<i>Gymnothorax polygonius</i>	Polygon moray	61
Moreia-serpente	<i>Enchelycore anatina</i>	Fangtooth moray	60
<i>Nomen dubium in Myctoperca</i>	<i>Epinephelus guaza</i>	<i>Nomen dubium in Myctoperca</i>	243
Olho-de-vidro-do-Mediterrâneo	<i>Hoplostethus mediterraneus</i>	Mediterranean slimehead	155
Olho-de-vidro-imperatriz	<i>Gephyroberyx darwinii</i>	Darwin's slimehead	154
Olho-de-vidro-laranja	<i>Hoplostethus atlanticus</i>	Orange roughy	154
Olho-de-vidro-preto	<i>Hoplostethus cadenati</i>	Black slimehead	155
Olho-pérola-de-barbatana-curta	<i>Scopelarchus analis</i>	Short fin pearleye	107
Olho-pérola-de-Guenther	<i>Scopelarchus guentheri</i>	Staring pearleye	107
Olho-pérola-de-Hubb	<i>Rosenblattichthys hubbsi</i>	Hubb's pearleye	107
Olho-pérola-de-Michael Sars	<i>Scopelarchus michaelsarsi</i>	Bigfin pearleye	107
Olho-pérola-de-Zugmayer	<i>Benthalbella infans</i>	Zugmayer's pearleye	106
Olho-verde	<i>Chlorophthalmus agassizii</i>	Shortnose greeneye	105
Olhudo	<i>Epigonus telescopus</i>	Black cardinal fish	240
Olhudo-anão	<i>Microichthys coccoi</i>	Dwarf deepwater cardinalfish	240
Olhudo-de-espinho	<i>Epigonus constanciae</i>	Constance deepwater cardinalfish	239
Olhudo-denteado	<i>Epigonus denticulatus</i>	Pencil cardinal	239

Portuguese common name	Scientific name	English common name	Page
Pailona	<i>Somniosus rostratus</i>	Little sleeper shark	34
Pai-velho-de-Bocage	<i>Argyropelecus bocagei</i>	No English common name	89
Pai-velho-de-Sladen	<i>Argyropelecus sladeni</i>	Sladen's hatchet fish	90
Pai-velho-de-três-pontos	<i>Valenciennellus tripunctulatus</i>	Constellationfish	91
Pai-velho-diáfano	<i>Sternoptyx diaphana</i>	Diaphanous hatchet fish	91
Pai-velho-do-Atlântico	<i>Argyripnus atlanticus</i>	No English common name	89
Pai-velho-do-Pacífico	<i>Argyropelecus affinis</i>	Pacific hatchet fish	89
Pai-velho-gigante	<i>Argyropelecus gigas</i>	Hatchetfish	90
Pai-velho-manchado	<i>Argyropelecus hemigymnus</i>	Half-naked hatchetfish	90
Pai-velho-oblíquo	<i>Sternoptyx pseudobscura</i>	Highlight hatchetfish	91
Pai-velho-prateado	<i>Argyropelecus olfersii</i>	Silver hatchetfish	90
Pai-velho-prateado-do-Atlântico	<i>Argyropelecus aculeatus</i>	Atlantic silver hatchetfish	89
Pai-velho-redondo	<i>Polyipnus polli</i>	No English common name	91
Palombeta	<i>Lichia amia</i>	Leerfish	253
Palometa	<i>Orcynopsis unicolor</i>	Plain bonito	167
Pampo-godinho	<i>Stromateus fiatola</i>	Blue butterfish	173
Papagaio	<i>Sparisoma chrysopterum</i>	Redtail parrotfish	288
Papagaio-velho	<i>Sparisoma cretense</i>	Parrotfish	287
Pargo-legítimo	<i>Pagrus pagrus</i>	Red porgy	272
Pargo-ruço	<i>Pagrus caeruleostictus</i>	Bluespotted seabream	272
Pargo-ruço-de-Ebrenberg	<i>Evynnis ehrenbergii</i>	Bluespotted seabream	269
Pargo-sêmola	<i>Pagrus auriga</i>	Redbanded seabream	271
Pata-roxa	<i>Scyliorhinus canicula</i>	Small-spotted catshark	21
Pata-roxa-gata	<i>Scyliorhinus stellaris</i>	Polkadot catshark	22
Pegador	<i>Remora remora</i>	Shark sucker	250
Pegador-azul	<i>Remora brachyptera</i>	Spearfish remora	249
Pegador-branco	<i>Remora albescens</i>	White suckerfish	249
Pegador-das-baleias	<i>Remora australis</i>	Whalesucker	249
Pegador-delgado	<i>Phterichthys lineatus</i>	Slender suckerfish	249
Pegador-dos-espardartes	<i>Remora osteochir</i>	Marlin sucker	250
Pegador-listado	<i>Echeneis naucrates</i>	Live sharksucker	249
Peixe-alado	<i>Howella simplex</i>	No English common name	241
Peixe-alado-de-Brodie	<i>Howella brodiei</i>	Pelagic basslet	240
Peixe-alado-de-Sherborn	<i>Howella sherborni</i>	Sherborn's pelagic bass	241
Peixe-alado-do-Atlântico	<i>Howella atlantica</i>	Atlantic pelagic basslet	240
Peixe-antena-do-Atlântico	<i>Bregmaceros atlanticus</i>	Antenna codlet	131
Peixe-aranha-de-Cabo-Verde	<i>Trachinus pellegrini</i>	Cape Verde weever	292
Peixe-aranha-maior	<i>Trachinus draco</i>	Greater weever	292
Peixe-aranha-menor	<i>Echiichthys vipera</i>	Lesser weever	291
Peixe-aranha-pontuado	<i>Trachinus araneus</i>	Spotted weever	292
Peixe-aranha-raiado	<i>Trachinus radiatus</i>	Starry weever	292
Peixe-baleia-bicolor	<i>Rondeletia bicolor</i>	Hooked redmouth whalefish	153
Peixe-baleia-de-boca-vermelha	<i>Rondeletia loricata</i>	Redmouth whalefish	153
Peixe-baleia-de-Gill	<i>Cetomimus gilli</i>	Gill's flabby whalefish	153
Peixe-baleia-de-Hempel	<i>Cetomimus hempeli</i>	Whalefish	154
Peixe-baleia-de-Myers	<i>Gyrinomimus myersi</i>	Myers' flabby whalefish	154
Peixe-baleia-de-Storer	<i>Ditropichthys storeri</i>	No English common name	154
Peixe-baleia-do-Galathea	<i>Cetichthys indagator</i>	No English common name	153
Peixe-baleia-Esaú	<i>Mirapinna esau</i>	Hairyfish	154
Peixe-baleia-mosaico	<i>Ataxolepis apus</i>	No English common name	153
Peixe-baleia-rosa	<i>Cetostoma regani</i>	Pink flabby whalefish	154
Peixe-baleia-veludo	<i>Barbourisia rufa</i>	Velvet whalefish	153
Peixe-bola-da-Guiné	<i>Sphoeroides marmoratus</i>	Guinean pufferfish	225
Peixe-bola-da-Macaronésia	<i>Canthigaster capistrata</i>	Macaronesian sharpnose-puffer	224
Peixe-bola-das-Caraíbas	<i>Canthigaster rostrata</i>	Sharpnose puffer	224

Portuguese common name	Scientific name	English common name	Page
Peixe-bola-de-focinho-achatado	<i>Sphoeroides pachygaster</i>	Blunthead puffer	225
Peixe-bola-de-pintas	<i>Sphoeroides spengleri</i>	Bandtail puffer	226
Peixe-bola-de-pintas	<i>Ephippion guttifer</i>	Prickly puffer	224
Peixe-bola-oceânico	<i>Lagocephalus lagocephalus</i>	Oceanic puffer	225
Peixe-bola-verde	<i>Lagocephalus laevis</i>	Smooth puffer	224
Peixe-cauda-de-fio	<i>Stylephorus chordatus</i>	Tube-eye	131
Peixe-cauda-de-seta	<i>Melanonus zugmayeri</i>	Arrowtail	142
Peixe-chicote-das-Canárias	<i>Gigantactis ios</i>	No English common name	220
Peixe-chicote-de-Van Höffen	<i>Gigantactis vanhoefeni</i>	No English common name	220
Peixe-chicote-filamentoso	<i>Gigantactis longicirra</i>	No English common name	220
Peixe-cirurgião	<i>Acanthurus monroviae</i>	Monrovia doctorfish	295
Peixe-cofre	<i>Acanthostracion notacanthus</i>	Island boxfish	226
Peixe-cofre-touro	<i>Acanthostracion quadricornis</i>	Scrawled cowfish	226
Peixe-cravo	<i>Lampris guttatus</i>	Opah	127
Peixe-cravo-elegante	<i>Lampris lauta</i>	East Atlantic Opah	127
Peixe-crina-barbatana-comprida	<i>Caristius macropus</i>	Manefish	260
Peixe-crina-da-Gronelândia	<i>Caristius fasciatus</i>	Greenland manefish	260
Peixe-crina-da-Madeira	<i>Paracaristius maderensis</i>	No English common name	260
Peixe-crina-de-Andriashov	<i>Platyberyx andriashovi</i>	No English common name	260
Peixe-crina-de-Maul	<i>Platyberyx mauli</i>	No English common name	260
Peixe-crina-opala	<i>Platyberyx opalescens</i>	No English common name	260
Peixe-de-coroa-espinhosa	<i>Acanthochænus luetkenii</i>	Picklefish	153
Peixe-de-focinho-arreganhado	<i>Lyconus brachycolus</i>	No English common name	148
Peixe-de-vela-alta	<i>Guentherus altivelis</i>	Jellynose	104
Peixe-dragão-abutre	<i>Borostomias mononema</i>	Sickle snaggletooth	96
Peixe-dragão-ambíguo	<i>Eustomias dubius</i>	No English common name	98
Peixe-dragão-atroz	<i>Photostomias atrox</i>	No English common name	103
Peixe-dragão-azulado	<i>Astronesthes cyaneus</i>	No English common name	93
Peixe-dragão-barbado	<i>Eustomias enbarbatus</i>	Barbate dragonfish	98
Peixe-dragão-barbatana-curta	<i>Photonectes parvimanus</i>	Fleshyfin dragonfish	103
Peixe-dragão-barbatana-filamentosa	<i>Echiostoma barbatum</i>	Threadfin dragonfish	97
Peixe-dragão-barbilho-curto	<i>Stomias brevibarbus</i>	No English common name	104
Peixe-dragão-bibolboso	<i>Eustomias contiguus</i>	No English common name	97
Peixe-dragão-bifilar	<i>Photonectes dinema</i>	No English common name	103
Peixe-dragão-boca-negra	<i>Photonectes margarita</i>	No English common name	103
Peixe-dragão-brilhante	<i>Borostomias elucens</i>	No English common name	96
Peixe-dragão-chicote	<i>Bathophilus nigerrimus</i>	Scaleless dragonfish	96
Peixe-dragão-cinzento	<i>Bathophilus schizochirus</i>	No English common name	96
Peixe-dragão-colmilhoso	<i>Grammatostomias dentatus</i>	No English common name	100
Peixe-dragão-da-Madeira	<i>Eustomias tetranema</i>	No English common name	100
Peixe-dragão-das-Bermudas	<i>Bathophilus proximus</i>	No English common name	96
Peixe-dragão-das-Caraíbas	<i>Aristostomias xenostoma</i>	No English common name	93
Peixe-dragão-da-Terra Nova	<i>Astronesthes gemmifer</i>	Snaggletooth	94
Peixe-dragão-de-barba-curta	<i>Astronesthes micropogon</i>	Snaggletooth	94
Peixe-dragão-de-barbatana-longa	<i>Bathophilus longipinnis</i>	Longfin dragonfish	95
Peixe-dragão-de-barbilho-acinoso	<i>Eustomias acinosus</i>	No English common name	97
Peixe-dragão-de-barbilho-amarelo	<i>Eustomias macrurus</i>	Yellowstem dragonfish	99
Peixe-dragão-de-barbilho-bifurcado	<i>Eustomias fissibarbis</i>	No English common name	98
Peixe-dragão-de-barbilho-bisseriado	<i>Melanostomias biseriatus</i>	No English common name	101
Peixe-dragão-de-barbilho-branco	<i>Astronesthes leucopogon</i>	No English common name	94
Peixe-dragão-de-barbilho-crescido	<i>Eustomias longibarba</i>	No English common name	98
Peixe-dragão-de-barbilho-filamentoso	<i>Grammatostomias flagellibarba</i>	No English common name	100
Peixe-dragão-de-barbilho-filifero	<i>Eustomias filifer</i>	No English common name	98
Peixe-dragão-de-barbilho-grande	<i>Eustomias macronema</i>	Bigbarb dragonfish	98
Peixe-dragão-de-barbilho-graúdo	<i>Astronesthes macropogon</i>	No English common name	94

Portuguese common name	Scientific name	English common name	Page
Peixe-dragão-de-barbilo-longo	<i>Leptostomias longibarba</i>	No English common name	101
Peixe-dragão-de-barbilo-negro	<i>Melanostomias melanopogon</i>	No English common name	102
Peixe-dragão-de-barbilo-plano	<i>Astronesthes neopogon</i>	Snaggletooth	94
Peixe-dragão-de-barbilo-ramificado	<i>Eustomias radicifilis</i>	No English common name	99
Peixe-dragão-de-barbilo-singelo	<i>Leptostomias haplocaulus</i>	No English common name	101
Peixe-dragão-de-Barton Bean	<i>Melanostomias bartonbeani</i>	Scaleless black dragonfish	101
Peixe-dragão-de-boca-de-serpente	<i>Heterophotus ophistoma</i>	Wingfin snaggletooth	100
Peixe-dragão-de-bolbo-alongado	<i>Leptostomias leptobolus</i>	No English common name	101
Peixe-dragão-de-bolbo-filamentoso	<i>Astronesthes similis</i>	No English common name	95
Peixe-dragão-de-bolbo-monodáctilo	<i>Eustomias monodactylus</i>	No English common name	99
Peixe-dragão-de-bolbo-papilar	<i>Chirostomias pliopterus</i>	Scaleless dragonfish	97
Peixe-dragão-de-bolbo-simples	<i>Eustomias simplex</i>	No English common name	99
Peixe-dragão-de-Brauer	<i>Photonectes braueri</i>	Brauer's dragonfish	102
Peixe-dragão-de-cauda-serreada	<i>Idiacanthus fasciola</i>	Ribbon sawtail fish	100
Peixe-dragão-de-draga	<i>Malacosteus niger</i>	Stoplight loosejaw	101
Peixe-dragão-de-Drechsel	<i>Eustomias drechseli</i>	No English common name	98
Peixe-dragão-de-focinho-manchado	<i>Melanostomias spilorhynchus</i>	No English common name	102
Peixe-dragão-de-furfur	<i>Eustomias furcifer</i>	Scaleless dragonfish	98
Peixe-dragão-de-Goodyear	<i>Photostomias goodeyeari</i>	No English common name	103
Peixe-dragão-de-Grimaldi	<i>Aristostomias grimaldii</i>	No English common name	93
Peixe-dragão-de-Guerne	<i>Photostomias guernei</i>	Loosejaw	103
Peixe-dragão-delgado	<i>Rhadinesthes decimus</i>	Slender snaggletooth	104
Peixe-dragão-de-mancha-negra	<i>Eustomias melanostigma</i>	No English common name	99
Peixe-dragão-de-mancha-redonda	<i>Grammatostomias circularis</i>	No English common name	100
Peixe-dragão-dente-curto	<i>Pachystomias microdon</i>	Smalltooth dragonfish	102
Peixe-dragão-de-Parr	<i>Eustomias parri</i>	No English common name	99
Peixe-dragão-de-Pawnee	<i>Bathophilus pawnee</i>	Pawnee dragonfish	96
Peixe-dragão-de-pinta-branca	<i>Photonectes leucospilus</i>	No English common name	103
Peixe-dragão-de-Richardson	<i>Astronesthes richardsoni</i>	Richardson's snaggletooth	95
Peixe-dragão-de-Santa Cruz	<i>Eustomias lipochirus</i>	No English common name	98
Peixe-dragão-de-Satterlee	<i>Eustomias satterleei</i>	Twinray dragonfish	99
Peixe-dragão-de-Schiff	<i>Eustomias schiffi</i>	No English common name	99
Peixe-dragão-de-Schmidt	<i>Eustomias schmidti</i>	Schmidt's dragonfish	99
Peixe-dragão-de-Sloane	<i>Chauliodus sloani</i>	Sloane's viperfish	97
Peixe-dragão-de-Sumatra	<i>Melanostomias melanops</i>	No English common name	102
Peixe-dragão-de-Tenison	<i>Eustomias tenisoni</i>	No English common name	100
Peixe-dragão-de-Tittmann	<i>Aristostomias tittmanni</i>	Loosejaw	93
Peixe-dragão-de-Vaillant	<i>Bathophilus vaillanti</i>	Scaleless dragonfish	96
Peixe-dragão-de-Valdívia	<i>Melanostomias valdiviae</i>	Valdivia black dragon fish	102
Peixe-dragão-de-Welsh	<i>Bathophilus metallicus</i>	No English common name	95
Peixe-dragão-de-Zharov	<i>Astronesthes zharovi</i>	No English common name	95
Peixe-dragão-do-Antártico	<i>Borostomias antarcticus</i>	Snaggletooth	96
Peixe-dragão-do-Atlântico	<i>Astronesthes atlanticus</i>	No English common name	93
Peixe-dragão-do-Cabo	<i>Neonesthes capensis</i>	Cape snaggletooth	102
Peixe-dragão-do-Dana	<i>Chauliodus danae</i>	Dana viperfish	96
Peixe-dragão-do-Índico	<i>Astronesthes indicus</i>	Black snaggletooth	94
Peixe-dragão-dos-Açores	<i>Flagellostomias boureei</i>	Longbarb dragonfish	100
Peixe-dragão-escamudo	<i>Stomias longibarbus</i>	Longbarb scaly dragonfish	104
Peixe-dragão-gladiador	<i>Leptostomias gladiator</i>	Scaleless dragonfish	100
Peixe-dragão-iridescente	<i>Bathophilus digitatus</i>	Scaleless black dragonfish	95
Peixe-dragão-jibóia	<i>Stomias boa</i>	Boa dragonfish	104
Peixe-dragão-listado	<i>Trigonolampa miriceps</i>	Threelights dragonfish	104
Peixe-dragão-luminoso	<i>Melanostomias macrophotus</i>	No English common name	101
Peixe-dragão-lunar	<i>Aristostomias lunifer</i>	No English common name	93
Peixe-dragão-lusitano	<i>Eustomias braueri</i>	No English common name	97

Portuguese common name	Scientific name	English common name	Page
Peixe-dragão-manchado	<i>Astronesthes niger</i>	Snaggletooth	94
Peixe-dragão-maravilhoso	<i>Photonectes mirabilis</i>	Blueband dragonfish	103
Peixe-dragão-obscuro	<i>Eustomias obscurus</i>	Scaleless dragonfish	99
Peixe-dragão-oval	<i>Bathophilus brevis</i>	No English common name	95
Peixe-dragão-tentaculado	<i>Melanostomias tentaculatus</i>	Tentacle dragonfish	102
Peixe-enfadado-barbatana-comprida	<i>Diretmoides pauciradiatus</i>	Silver spinyfin	155
Peixe-enfadado-de-Parin	<i>Diretmichthys parini</i>	Parin's spinyfish	155
Peixe-enfadado-prateado	<i>Diretmus argenteus</i>	Silver spinyfin	156
Peixe-espada	<i>Lepidopus caudatus</i>	Silver scabbardfish	165
Peixe-espada-intermédio	<i>Aphanopus intermedius</i>	Intermediate scabbardfish	164
Peixe-espada-preto	<i>Aphanopus carbo</i>	Black scabbardfish	164
Peixe-fantasma-dardo	<i>Bathylychnops exilis</i>	Javelin spookfish	84
Peixe-fantasma-dardo-listado	<i>Bathylychnops brachyrhynchus</i>	No English common name	84
Peixe-fantasma-de-barbatanas-longas	<i>Dolichopteryx longipes</i>	Brown-snout spookfish	84
Peixe-farol-alado	<i>Puck pinnata</i>	No English common name	219
Peixe-farol-barbatana-comprida	<i>Ctenochirichthys longimanus</i>	No English common name	217
Peixe-farol-cabeça-comprida	<i>Chaenophryne longiceps</i>	Can-opener smoothdream	216
Peixe-farol-cabeça-de-baleia	<i>Lophodolos acanthognathus</i>	Whalehead dreamer	217
Peixe-farol-córnneo	<i>Centrophryne spinulosa</i>	Horned lantern fish	219
Peixe-farol-cornudo	<i>Dolopichthys alector</i>	No English common name	217
Peixe-farol-da-Madeira	<i>Himantolophus albinares</i>	No English common name	215
Peixe-farol-de-bolbo-assimétrico	<i>Oneirodes anisacanthus</i>	No English common name	217
Peixe-farol-de-Carlsberg	<i>Oneirodes carlsbergi</i>	No English common name	218
Peixe-farol-de-Clarke	<i>Oneirodes clarkei</i>	No English common name	218
Peixe-farol-de-Coues	<i>Cryptopsaras couesii</i>	Triplewart seadevil	219
Peixe-farol-de-crista	<i>Dolopichthys jubatus</i>	No English common name	217
Peixe-farol-de-Eschricht	<i>Oneirodes eschrichtii</i>	Bulbous dreamer	218
Peixe-farol-de-focinho-curto	<i>Himantolophus brevirostris</i>	No English common name	215
Peixe-farol-de-Guadalupe	<i>Phyllorhinichthys micractis</i>	No English common name	218
Peixe-farol-de-Holbøll	<i>Ceratias holboelli</i>	Kroyer's deep-sea angler fish	219
Peixe-farol-de-Karsten	<i>Dolopichthys karsteni</i>	No English common name	217
Peixe-farol-de-Krefft	<i>Oneirodes krefftii</i>	Krefft's dreamer	218
Peixe-farol-de-Maul	<i>Himantolophus mauli</i>	No English common name	216
Peixe-farol-de-Post	<i>Oneirodes posti</i>	No English common name	218
Peixe-farol-de-queixo-direito	<i>Leptacanthichthys gracilispinis</i>	Plainchin dreamarm	217
Peixe-farol-de-Santa Cruz	<i>Oneirodes macronema</i>	No English common name	218
Peixe-farol-de-Wedl	<i>Bufockeratias wedli</i>	No English common name	216
Peixe-farol-diabo	<i>Dolopichthys longicornis</i>	No English common name	217
Peixe-farol-do-Atlântico	<i>Himantolophus groenlandicus</i>	Atlantic footballfish	216
Peixe-farol-do-Atlântico-Norte	<i>Oneirodes macrosteus</i>	No English common name	218
Peixe-farol-do-Dana	<i>Dolopichthys danae</i>	No English common name	217
Peixe-farol-do-Panamá	<i>Pentherichthys atratus</i>	No English common name	218
Peixe-farol-dragão	<i>Chaenophryne draco</i>	Smooth dreamer	216
Peixe-farol-espinhosos	<i>Sphryne gladiifera</i>	Prickly dreamer	219
Peixe-farol-haste-curta	<i>Microlophichthys microlophus</i>	Short-rod anglerfish	217
Peixe-farol-lobado	<i>Dolopichthys pullatus</i>	Lobed dreamer	217
Peixe-farol-multifilamentoso	<i>Oneirodes myrionomus</i>	No English common name	218
Peixe-farol-ramudo	<i>Chaenophryne ramifera</i>	No English common name	216
Peixe-farol-sapo	<i>Himantolophus compressus</i>	No English common name	216
Peixe-ganchudo	<i>Lasiognathus amphirhamphus</i>	No English common name	219
Peixe-ganchudo-de-Beebe	<i>Lasiognathus beebei</i>	No English common name	219
Peixe-ganchudo-tridente	<i>Lasiognathus saccostoma</i>	No English common name	219
Peixe-glutão	<i>Dysalotus oligoscolus</i>	No English common name	289
Peixe-glutão-de-Alcock	<i>Dysalotus alcocki</i>	No English common name	289
Peixe-glutão-de-focinho-curvo	<i>Pseudoscopelus obtusifrons</i>	No English common name	290

Portuguese common name	Scientific name	English common name	Page
Peixe-glutão-de-Kerbert	<i>Kali kerberti</i>	No English common name	290
Peixe-glutão-dentudo	<i>Kali macrodon</i>	No English common name	290
Peixe-glutão-de-Parr	<i>Kali parri</i>	No English common name	290
Peixe-glutão-de-Sagami	<i>Pseudoscopelus sagamianus</i>	No English common name	290
Peixe-glutão-de-vela	<i>Pseudoscopelus altipinnis</i>	No English common name	290
Peixe-glutão-do-Índico	<i>Kali indica</i>	No English common name	290
Peixe-glutão-escudado	<i>Pseudoscopelus scutatus</i>	No English common name	291
Peixe-glutão-negro	<i>Chiasmodon niger</i>	Black swallower	289
Peixe-glutão-rabudo	<i>Kali macrura</i>	No English common name	290
Peixe-lagarto-abissal	<i>Halosauropsis macrochir</i>	Abyssal halosaur	56
Peixe-lagarto-da-costa	<i>Bathysaurus mollis</i>	Highfin lizardfish	109
Peixe-lagarto-da-fundura	<i>Bathysaurus ferox</i>	Deep-sea lizardfish	109
Peixe-lagarto-da-praia	<i>Synodus foetens</i>	Inshore lizardfish	108
Peixe-lagarto-de-bico	<i>Aldrovandia rostrata</i>	Sharpnose halosaur	56
Peixe-lagarto-de-focinho-rombo	<i>Trachinocephalus myops</i>	Snakefish	109
Peixe-lagarto-de-Guenther	<i>Halosaurus guentheri</i>	Günther's halosaur	56
Peixe-lagarto-de-Johnson	<i>Halosaurus jonhsonianus</i>	Johnson's halosaur	56
Peixe-lagarto-delgado	<i>Aldrovandia gracilis</i>	Slender halosaur	56
Peixe-lagarto-de-Owen	<i>Halosaurus ovenii</i>	Owen's halosaur	57
Peixe-lagarto-de-rolo	<i>Synodus synodus</i>	Redbarred lizardfish	108
Peixe-lagarto-do-Atlântico	<i>Synodus saurus</i>	Atlantic lizardfish	108
Peixe-lagarto-dos-Açores	<i>Aldrovandia phalacra</i>	Hawaiian halosaurid fish	56
Peixe-lagarto-oleoso	<i>Aldrovandia oleosa</i>	Oily halosaur	56
Peixe-lagarto-pálido	<i>Aldrovandia affinis</i>	Pale halosaur	56
Peixe-lanterna-alado	<i>Lampanyctus alatus</i>	Winged lanternfish	121
Peixe-lanterna-anão	<i>Lampanyctus pusillus</i>	Pygmy lanternfish	123
Peixe-lanterna-anómalo	<i>Lampadена anomala</i>	Anomalous lanternfish	120
Peixe-lanterna-arco-íris	<i>Bolinichthys pyrsobolus</i>	Fiery lanternfish	114
Peixe-lanterna-atarracado	<i>Bolinichthys supralateralis</i>	Stubby lanternfish	114
Peixe-lanterna-barbatana-comprida	<i>Bolinichthys longipes</i>	Popeye lampfish	114
Peixe-lanterna-brilhante	<i>Ctenoscopelus phengodes</i>	Bright lanternfish	116
Peixe-lanterna-cintilante	<i>Diaphus lucidus</i>	Spotlight lanternfish	117
Peixe-lanterna-cobreado	<i>Lampanyctus cuprarius</i>	Atlantic blackcap lanternfish	121
Peixe-lanterna-complexo	<i>Lampanyctus intricarius</i>	Intricate lanternfish	122
Peixe-lanterna-corcunda	<i>Benthosema suborbitale</i>	Smallfin lanternfish	114
Peixe-lanterna-cornudo	<i>Diaphus splendidus</i>	Horned lanternfish	118
Peixe-lanterna-crocodilo	<i>Lampanyctus crocodilus</i>	Jewel lanternfish	121
Peixe-lanterna-da-California	<i>Diaphus theta</i>	California headlightfish	118
Peixe-lanterna-da-fundura	<i>Taenigichthys bathyphilus</i>	Deepwater lanternfish	126
Peixe-lanterna-da-Madeira	<i>Ceratoscopelus maderensis</i>	Madeira lantern fish	115
Peixe-lanterna-das-Bermudas	<i>Hygophum hygomii</i>	Bermuda lantern fish	120
Peixe-lanterna-de-Andréas	<i>Centrobranchus andreae</i>	Andre's lanternfish	115
Peixe-lanterna-de-barbatanas-curtas	<i>Neoscopelus microchir</i>	Shortfin neoscopelid	113
Peixe-lanterna-de-barbatanas-grandes	<i>Hygophum macrochir</i>	Largefin lanternfish	120
Peixe-lanterna-de-Benoit	<i>Hygophum benoiti</i>	Benoit's lanternfish	119
Peixe-lanterna-de-Bertelsen	<i>Diaphus bertelseni</i>	Bertelsen's lanternfish	116
Peixe-lanterna-de-Bolin	<i>Notoscopelus bolini</i>	Fanfare lanternfish	125
Peixe-lanterna-de-cabeça-curta	<i>Diaphus brachycephalus</i>	Short-headed lantern fish	116
Peixe-lanterna-de-cabeça-manchada	<i>Diaphus metopocampus</i>	Spothead lantern fish	117
Peixe-lanterna-de-Cabo-Verde	<i>Taenigichthys minimus</i>	Waistcoat lanternfish	127
Peixe-lanterna-de-cauda-espinhosa	<i>Notoscopelus caudispinosus</i>	Spinetail lanternfish	125
Peixe-lanterna-de-Chaves	<i>Lampadena chavesi</i>	Chaves' lanternfish	120
Peixe-lanterna-de-Cocco	<i>Gonichthysocco</i>	Cocco's lanternfish	119
Peixe-lanterna-de-Diogenes	<i>Diogenichthys laternatus</i>	Diogenes lanternfish	119
Peixe-lanterna-de-Doflein	<i>Lobianchia dofleini</i>	Dofleini's lantern fish	123

Portuguese common name	Scientific name	English common name	Page
Peixe-lantera-de-Duméril	<i>Diaphus dumerili</i>	Dumeril's lanternfish	117
Peixe-lantera-de-focinho-comprido	<i>Centrobranchus choerocephalus</i>	No English common name	115
Peixe-lantera-de-focinho-luminoso	<i>Diaphus effulgens</i>	Headlight lanternfish	117
Peixe-lantera-de-focinho-redondo	<i>Centrobranchus nigrocellatus</i>	Roundnose lanternfish	115
Peixe-lantera-de-Gauss	<i>Lepidophanes gaussi</i>	Gauss' lanternfish	123
Peixe-lantera-de-Gemellar	<i>Lobianchia gemellarii</i>	Gemellar's lanternfish	124
Peixe-lantera-de-Gilbert	<i>Diaphus adenomus</i>	Firebow lanternfish	116
Peixe-lantera-de-Günther	<i>Lepidophanes guentheri</i>	Günther's lanternfish	123
Peixe-lantera-de-Holt	<i>Diaphus holti</i>	Small lantern fish	117
Peixe-lantera-de-Krøyer	<i>Notoscopelus kroyeri</i>	Krøyer's lanternfish	126
Peixe-lantera-de-Luetken	<i>Diaphus luetkeni</i>	Luetken's lanternfish	117
Peixe-lantera-de-MacDonald	<i>Lampanyctus macdonaldi</i>	MacDonald's lanternfish	122
Peixe-lantera-de-Rafinesque	<i>Diaphus rafinesquii</i>	White-spotted lantern fish	118
Peixe-lantera-de-Reinhardt	<i>Hygophum reinhardti</i>	Reinhardt's lanternfish	120
Peixe-lantera-de-Rufous	<i>Symbolophorus rufinus</i>	Rufous lanternfish	126
Peixe-lantera-de-Tåning	<i>Hygophum taanungi</i>	Tåning's lanternfish	120
Peixe-lantera-de-Townsend	<i>Ceratoscopelus townsendi</i>	Dogtooth lampfish	115
Peixe-lantera-de-Vérany	<i>Symbolophorus veranyi</i>	Large-scale lantern fish	126
Peixe-lantera-de-Warming	<i>Ceratoscopelus warmingii</i>	Warming's lanternfish	115
Peixe-lantera-de-Wisner	<i>Dasy scopelus selenops</i>	Wisner's lantern fish	116
Peixe-lantera-divertido	<i>Lampanyctus festivus</i>	Festive lanternfish	122
Peixe-lantera-do-alto	<i>Triphoturus nigrescens</i>	Highseas lampfish	127
Peixe-lantera-do-Ártico	<i>Protomyctophum arcticum</i>	Arctic lanternfish	126
Peixe-lantera-do-Atlântico	<i>Diogenichthys atlanticus</i>	Atlantic lanternfish	118
Peixe-lantera-do-Índico	<i>Bolinichthys indicus</i>	Smoothcheek lanternfish	114
Peixe-lantera-do-Mediterrâneo	<i>Notoscopelus elongatus</i>	No English common name	125
Peixe-lantera-do-Pacífico	<i>Scopelengys tristis</i>	Pacific blackchin	113
Peixe-lantera-do-Valdívia	<i>Notolychnus valdiviae</i>	Topside lanternfish	125
Peixe-lantera-escamudo	<i>Neoscopelus macrolepidotus</i>	Large-scaled lantern fish	113
Peixe-lantera-escuro	<i>Lampanyctus ater</i>	Dusky lanternfish	121
Peixe-lantera-espelhado	<i>Lampadена speculigera</i>	Mirror lanternfish	121
Peixe-lantera-esporão	<i>Bolinichthys photothorax</i>	Spurcheek lanternfish	114
Peixe-lantera-faísca	<i>Electrona risso</i>	Electric lantern fish	119
Peixe-lantera-ferreiro	<i>Lowina interrupta</i>	Barebelly lanternfish	124
Peixe-lantera-flácido	<i>Diaphus subtilis</i>	Flabby lanternfish	118
Peixe-lantera-fosforescente	<i>Myctophum nitidulum</i>	Pearlyspotted lanternfish	124
Peixe-lantera-glacial	<i>Benthosema glaciale</i>	Glacier lantern fish	113
Peixe-lantera-lobisomem	<i>Lampanyctus photonotus</i>	Dotback lanternfish	123
Peixe-lantera-luminoso	<i>Lampadena luminosa</i>	Luminous lanternfish	121
Peixe-lantera-luneta	<i>Diaphus perspicillatus</i>	Flatface lanternfish	118
Peixe-lantera-manchado	<i>Myctophum punctatum</i>	Spotted lanternfish	124
Peixe-lantera-metálico	<i>Myctophum affine</i>	Metallic lanternfish	124
Peixe-lantera-mole	<i>Diaphus mollis</i>	Soft lanternfish	117
Peixe-lantera-nacarado	<i>Lampanyctus gemmifer</i>	No English common name	122
Peixe-lantera-nu	<i>Taaningichthys paurolychnus</i>	Naked lanternfish	127
Peixe-lantera-oliva	<i>Notoscopelus resplendens</i>	Patchwork lanternfish	126
Peixe-lantera-preto	<i>Lampanyctus niger</i>	Black lantern fish	122
Peixe-lantera-raro	<i>Lowina rara</i>	Rare lanternfish	124
Peixe-lantera-riscado	<i>Lampanyctus lineatus</i>	Lanternfish	122
Peixe-lantera-romeirinho	<i>Lampadena atlantica</i>	Atlantic tail-light lanternfish	120
Peixe-lantera-sarapintado	<i>Scopelopsis multipunctatus</i>	Multispotted lanternfish	126
Peixe-lantera-termófilo	<i>Diaphus termophilus</i>	Taaning's lantern fish	118
Peixe-lapa	<i>Cyclopterus lumpus</i>	Lumpfish	237
Peixe-lapa-espinhoso	<i>Eumicromes spinosus</i>	Atlantic spiny lump sucker	237
Peixe-lesma-de-focinho-preto	<i>Paraliparis copei</i>	Blacksnout seasnail	238

Portuguese common name	Scientific name	English common name	Page
Peixe-lesma-de-Montagus	<i>Liparis montagui</i>	Montagus seasnail	238
Peixe-lesma-escuro	<i>Paraliparis nigellus</i>	No English common name	238
Peixe-lesma-listado	<i>Liparis liparis</i>	Striped seasnail	237
Peixe-lesma-membranoso	<i>Paraliparis membranaceus</i>	No English common name	238
Peixe-lima	<i>Trachyrincus scabrus</i>	Roughsnout grenadier	139
Peixe-lobo-azul	<i>Anarhichas denticulatus</i>	Northern wolffish	289
Peixe-lua	<i>Mola mola</i>	Ocean sunfish	222
Peixe-lua-comprido	<i>Ranzania laevis</i>	Slender sunfish	222
Peixe-lua-de-Ranzani	<i>Mola alexandrini</i>	Bump-head sunfish	222
Peixe-lua-rabudo	<i>Masturus lanceolatus</i>	Sharptail mola	222
Peixe-luminoso-de-Blackford	<i>Yarrella blackfordi</i>	No English common name	93
Peixe-luminoso-de-Maul	<i>Pollichthys mauli</i>	Stareye lightfish	92
Peixe-luminoso-de-Power	<i>Vinciguerria poweriae</i>	Power's deep-water bristle-mouth fish	93
Peixe-luminoso-do-Atlântico	<i>Polymetme thaeocoryla</i>	No English common name	92
Peixe-luminoso-do-Índico	<i>Polymetme corythaenoides</i>	Rendezvous fish	92
Peixe-luminoso-do-Panamá	<i>Vinciguerria lucetia</i>	Panama lightfish	92
Peixe-luminoso-esguio	<i>Vinciguerria attenuata</i>	Slender lightfish	92
Peixe-luminoso-oceânico	<i>Vinciguerria nimbaria</i>	Oceanic lightfish	93
Peixe-luminoso-oval	<i>Ichthyococcus ovatus</i>	Ovate lightfish	92
Peixe-luz	<i>Maurolicus amethystinopunctatus</i>	Pearlsides	90
Peixe-luz-de-Muller	<i>Maurolicus muelleri</i>	Silvery lightfish	91
Peixe-mandíbula-de-martelo	<i>Omosudis lowii</i>	Omosudid	112
Peixe-manteiga	<i>Pholis gunnellus</i>	Rock gunnel	289
Peixe-ogre	<i>Anoplogaster cornuta</i>	Common fangtooth	156
Peixe-orelhudo-da-Madeira	<i>Macruronus maderensis</i>	No English common name	148
Peixe-ouriço-de-crista	<i>Diodon holocanthus</i>	Longspined porcupinefish	223
Peixe-ouriço-pelágico	<i>Diodon eydouxii</i>	Pelagic porcupinefish	223
Peixe-ouriço-ponteado	<i>Diodon hystrix</i>	Spot-fin porcupinefish	224
Peixe-pastor-amarelado	<i>Psenes cyanophrys</i>	Freckled driftfish	173
Peixe-pastor-prateado	<i>Psenes maculatus</i>	Silver driftfish	173
Peixe-pau-da-fundura-do-Atlântico	<i>Centrodraco acanthopoma</i>	Atlantic deepwater dragonet	295
Peixe-pau-de-Sousa	<i>Protogrammus sousai</i>	Meteor dragonet	294
Peixe-pau-lira	<i>Callionymus lyra</i>	Dragonet	293
Peixe-pau-listado	<i>Callionymus reticulatus</i>	Reticulated dragonet	294
Peixe-pau-malhado	<i>Callionymus maculatus</i>	Spotted dragonet	293
Peixe-pau-pintado	<i>Callionymus risso</i>	Risso's dragonet	294
Peixe-pau-rabudo	<i>Callionymus pusillus</i>	Sailfin dragonet	294
Peixe-pau-rosa	<i>Synchiropus phaeton</i>	Phaeton dragonet	294
Peixe-pelícano	<i>Eurypharynx pelecanoides</i>	Pelican eel	70
Peixe-pelícano-de-cauda-peluda	<i>Saccopharynx thalassa</i>	Hairytail swallower eel	71
Peixe-pelícano-de-cauda-ramosa	<i>Saccopharynx ramosus</i>	Shortline swallower eel	71
Peixe-pelícano-de-cauda-trilobada	<i>Saccopharynx trilobatus</i>	No English common name	71
Peixe-pelícano-de-Harrison	<i>Saccopharynx harrisoni</i>	No English common name	71
Peixe-pelícano-de-Hjort	<i>Saccopharynx hjorti</i>	No English common name	71
Peixe-pelícano-do-Norte	<i>Saccopharynx ampullaceus</i>	Northern swallower eel	70
Peixe-pelícano-dos-Açores	<i>Saccopharynx paucovertebratis</i>	No English common name	71
Peixe-pérola-cobrelo	<i>Carapus acus</i>	Pearl fish	159
Peixe-pérola-de-Drummond	<i>Echiodon drummondii</i>	No English common name	160
Peixe-pérola-dentado	<i>Echiodon dentatus</i>	No English common name	159
Peixe-pescador-da-Madeira	<i>Caulophryne polynema</i>	No English common name	215
Peixe-pescador-de-Jordan	<i>Caulophryne jordani</i>	Fanfin angler	214
Peixe-piloto	<i>Naucrates ductor</i>	Pilotfish	253
Peixe-porco	<i>Oxynotus centrina</i>	Angular roughshark	34
Peixe-porco-áspero	<i>Stephanolepis hispidus</i>	Planehead filefish	226
Peixe-porco-bicudo	<i>Aluterus scriptus</i>	Scribbled leatherjacket filefish	226

Portuguese common name	Scientific name	English common name	Page
Peixe-porco-de-vela	<i>Oxynotus paradoxus</i>	Sailfin roughshark	35
Peixe-porco-galhudo	<i>Stephanolepis setifer</i>	Pygmy filefish	227
Peixe-porco-galhudo	<i>Aluterus monoceros</i>	Unicorn leatherjacket filefish	226
Peixe-prata-espinhoso	<i>Grammicolepis brachiusculus</i>	Thorny tinselfish	130
Peixe-rã-anão	<i>Melamphaes pumilus</i>	No English common name	151
Peixe-rã-cabecinha	<i>Melamphaes polylepis</i>	No English common name	151
Peixe-rã-couraça	<i>Scopelogadus mizolepis</i>	Ragged bigscale	153
Peixe-rã-da-Madeira	<i>Melamphaes typhlops</i>	Ridgehead	151
Peixe-rã-de-Bean	<i>Scopelogadus beanii</i>	Bean's bigscale	153
Peixe-rã-de-cabeça-preta	<i>Poromitra nigriceps</i>	No English common name	152
Peixe-rã-de-crista	<i>Poromitra crassiceps</i>	Crested bigscale	152
Peixe-rã-de-Ebeling	<i>Melamphaes ebelingi</i>	No English common name	150
Peixe-rã-delgado	<i>Scopeloberyx opisthopterus</i>	Bigscale	152
Peixe-rã-de-nariz-espinhoso	<i>Poromitra capito</i>	Ridgehead	151
Peixe-rã-de-ombro-espinhoso	<i>Melamphaes suborbitalis</i>	Shoulder spine bigscale	151
Peixe-rã-do-Índico	<i>Melamphaes indicus</i>	No English common name	150
Peixe-rã-dos-Açores	<i>Scopeloberyx rubriventer</i>	Ridgehead	152
Peixe-rã-falso	<i>Melamphaes falsidicus</i>	No English common name	150
Peixe-rã-nariz-achatado	<i>Melamphaes simus</i>	Ridgehead	151
Peixe-rã-olhinho	<i>Melamphaes microps</i>	Ridgehead	150
Peixe-rã-olhudo	<i>Poromitra megalops</i>	Bigscale	152
Peixe-rã-robusto	<i>Scopeloberyx robustus</i>	Longjaw bigscale	152
Peixe-rã-sobrolho	<i>Melamphaes longivelis</i>	Eyebrow bigscale	150
Peixe-rato-de-Raleigh	<i>Harriotta raleighana</i>	Pacific longnose chimaera	54
Peixe-rato-narigudo	<i>Rhinochimaera atlantica</i>	Straightnose rabbitfish	54
Peixe-rei	<i>Atherina presbyter</i>	Sand smelt	198
Peixe-rei-do-alto	<i>Atherina hepsetus</i>	Mediterranean sand smelt	197
Peixe-rei-do-Mediterrâneo	<i>Atherina boyeri</i>	Big-scale sand smelt	197
Peixe-remo-de-cauda-fina	<i>Radiicephalus elongatus</i>	Tapetail	128
Peixe-remo-de-crista	<i>Lophotus lacepede</i>	Crested oarfish	127
Peixe-remo-de-Kleinenberg	<i>Eretmophorus kleinenbergi</i>	No English common name	139
Peixe-remo-do-Índico	<i>Regalecus russellii</i>	Oarfish;	129
Peixe-sapo-colorido	<i>Chaunacops coloratus</i>	No English common name	214
Peixe-sapo-de-Suttkus	<i>Chaunax suttkusi</i>	No English common name	214
Peixe-sapo-do-sargaço	<i>Histrio histrio</i>	Sargassumfish	213
Peixe-sapo-do-Senegal	<i>Fowlerichthys senegalensis</i>	Senegalese frogfish	213
Peixe-sapo-mole	<i>Chaunax pictus</i>	Pink frogmouth	214
Peixe-sapo-multiocelado	<i>Antennarius multiocellatus</i>	Longlure frogfish	213
Peixe-sapo-rosado	<i>Chaunacops roseus</i>	No English common name	214
Peixe-sapo-salpicado	<i>Antennatus nummifer</i>	Spotfin frogfish	213
Peixe-sapo-uniocelado	<i>Fowlerichthys radiosus</i>	Singlespot frogfish	213
Peixe-serpentina	<i>Eutaeniophorus festivus</i>	Festive ribbonfish	154
Peixe-sestro-barbudo	<i>Linophryne polypogon</i>	No English common name	221
Peixe-sestro-cabeça-espinhosa	<i>Photocorynus spiniceps</i>	No English common name	221
Peixe-sestro-coroadó	<i>Linophryne coronata</i>	No English common name	221
Peixe-sestro-da-Madeira	<i>Linophryne maderensis</i>	No English common name	221
Peixe-sestro-de-barbilho-trirramoso	<i>Linophryne brevibarbata</i>	No English common name	220
Peixe-sestro-dentudo	<i>Linophryne macrodon</i>	No English common name	221
Peixe-sestro-de-Waterman	<i>Linophryne algibarbata</i>	No English common name	220
Peixe-sestro-do-Arcturus	<i>Linophryne arcturi</i>	No English common name	220
Peixe-sestro-do-Índico	<i>Linophryne indica</i>	Headlight angler	221
Peixe-sestro-hexafilamentoso	<i>Linophryne sexfilis</i>	No English common name	221
Peixe-sestro-lucifer	<i>Linophryne lucifer</i>	No English common name	221
Peixe-sestro-macio	<i>Haplophryne mollis</i>	Soft leafvent angler	220
Peixe-sestro-pompom	<i>Linophryne racemifera</i>	No English common name	221

Portuguese common name	Scientific name	English common name	Page
Peixe-sestro-ramoso	<i>Linophryne arborifera</i>	No English common name	220
Peixe-sola	<i>Opisthoproctus soleatus</i>	Barrel-eye	85
Peixe-sola-cabeça-de-vidro	<i>Rhynchohyalus natalensis</i>	Glasshead barreleye	85
Peixe-sola-de-Grimaldi	<i>Monacoa grimaldii</i>	Mirrorbelly	85
Peixe-soldado-espinhoso	<i>Corniger spinosus</i>	Spinycheek soldierfish	156
Peixe-tábuia	<i>Trachipterus arcticus</i>	Dealfish	128
Peixe-tábuia-do-Mediterrâneo	<i>Trachipterus trachypterus</i>	Mediterranean dealfish	128
Peixe-tábuia-recortado	<i>Zu cristatus</i>	Scalloped ribbonfish	128
Peixe-tripé	<i>Bathypterois grallator</i>	Tripodfish	106
Peixe-tripé-abissal	<i>Bathypterois longipes</i>	Abyssal spiderfish	106
Peixe-tripé-de-barbatana-negra	<i>Bathypterois phenax</i>	Blackfin spiderfish	106
Peixe-tripé-de-Marion	<i>Bathytyphlops marionae</i>	Marion's spiderfish	106
Peixe-tripé-de-Sewell	<i>Bathytyphlops sewelli</i>	No English common name	106
Peixe-tripé-espinhoso	<i>Bathymicrops multispinis</i>	No English common name	105
Peixe-tripé-filamentoso	<i>Bathypterois dubius</i>	Spiderfish	105
Peixe-tripé-real	<i>Bathymicrops regis</i>	No English common name	105
Peixe-tripé-temporão	<i>Discoverichthys praecox</i>	No English common name	106
Peixe-vermelho	<i>Sebastes norvegicus</i>	Golden redfish	229
Peixe-vermelho-da-fundura	<i>Sebastes mentella</i>	Beaked redfish	229
Peixe-vigia-de-barbatana-preta	<i>Scopelosaurus lepidus</i>	Blackfin waryfish	107
Peixe-vigia-de-Berry	<i>Ahliasaurus berryi</i>	No English common name	107
Peixe-vigia-de-Smith	<i>Scopelosaurus smithii</i>	No English common name	108
Peixe-vigia-prateado	<i>Scopelosaurus argenteus</i>	Waryfish	107
Peixe-voador-barbudo	<i>Cheilopogon cyanopterus</i>	Margined flyingfish	200
Peixe-voador-de-asa-espelhada	<i>Hirundichthys speculiger</i>	Mirrorwing flyingfish	201
Peixe-voador-de-asa-negra	<i>Hirundichthys rondeletii</i>	Black wing flyingfish	201
Peixe-voador-de-asa-negra-do-Atlântico	<i>Hirundichthys volador</i>	Atlantic blackwing flyingfish	202
Peixe-voador-de-asas-listadas	<i>Cheilopogon exsiliens</i>	Bandwing flyingfish	200
Peixe-voador-de-barbatana-manchada	<i>Cheilopogon furcatus</i>	Spotfin flyingfish	200
Peixe-voador-de-Bennett	<i>Cheilopogon pinnatibarbus</i>	Bennett's flyingfish	200
Peixe-voador-do-Mediterrâneo	<i>Cheilopogon heterurus</i>	Mediterranean flyingfish	200
Pequeno-argonauta	<i>Nomeus gronovii</i>	Man-of-war fish	173
Periquito	<i>Antigonia capros</i>	Deepbody boarfish	296
Perna-de-moça	<i>Galeorhinus galeus</i>	Tope shark	23
Pescada-branca	<i>Merluccius merluccius</i>	European hake	148
Pim-pim	<i>Capros aper</i>	Boarfish	297
Pombo	<i>Plectorhinchus mediterraneus</i>	Rubberlip grunt	262
Porco-espinho-americano	<i>Chilomycterus spinosus spinosus</i>	No English common name	223
Porco-espinho-da-Guiné	<i>Chilomycterus spinosus mauretanicus</i>	Guinean burrfish	223
Porco-espinho-de-pintas	<i>Chilomycterus reticulatus</i>	Pacific burrfish	222
Porco-espinho-listado	<i>Chilomycterus schoepfii</i>	Striped burrfish	223
Pregado	<i>Scophthalmus maximus</i>	Turbot	187
Preguiçosa-branca	<i>Kyphosus sectatrix</i>	Bermuda sea chub	238
Preguiçosa-bronzeada	<i>Kyphosus vaigiensis</i>	Brassy chub	239
Preguiçosa-cinzenta	<i>Kyphosus bigibbus</i>	Brown chub	238
Quelme	<i>Centrophorus uyato</i>	Little gulper shark	36
Quelme-do-Zeehaan	<i>Centrophorus cf. zeehaani</i>	Little gulper shark	37
Quelmo	<i>Centrophorus macrourus</i>	No English common name	36
Quimera-comum	<i>Chimaera monstrosa</i>	Rabbit fish	53
Quimera-opala	<i>Chimaera opalescens</i>	Opal chimaera	53
Raia-bicuda	<i>Dipturus oxyrinchus</i>	Longnosed skate	43
Raia-curva	<i>Raja undulata</i>	Undulate ray	48
Raia-da-fundura	<i>Rajella bathyphila</i>	Deep-water ray	48
Raia-da-Madeira	<i>Raja maderensis</i>	Madeiran ray	46
Raia-de-Bigelow	<i>Rajella bigelowi</i>	Bigelow's ray	48

Portuguese common name	Scientific name	English common name	Page
Raia-de-cauda-curta	<i>Amblyraja jensenii</i>	Shorttail skate	42
Raia-de-dois-olhos	<i>Leucoraja naevus</i>	Cuckoo ray	45
Raia-de-Krefft	<i>Malacoraja krefftii</i>	Krefft's ray	45
Raia-delgada	<i>Dipturus intermedius</i>	Flapper Skate	43
Raia-de-quatro-olhos	<i>Raja miraletus</i>	Brown ray	47
Raia-de-Richardson	<i>Bathyraja richardsoni</i>	Richardson's ray	49
Raia-de-São-Pedro	<i>Leucoraja circularis</i>	Sandy ray	44
Raia-lenga	<i>Raja clavata</i>	Thornback ray	45
Raia-manchada	<i>Raja montagui</i>	Mediterranean starry ray	47
Raia-nevoeira	<i>Rajella linea</i>	Sailray	48
Raia-norueguesa	<i>Dipturus nidarosiensis</i>	Norwegian skate	43
Raia-norueguesa	<i>Dipturus sp. Stehmann</i>	Norwegian skate	44
Raia-oirega	<i>Dipturus batis</i>	Blue skate	43
Raia-pálida	<i>Bathyraja pallida</i>	Pale ray	49
Raia-pigméia-ibérica	<i>Neoraja iberica</i>	Iberian pygmy skate	45
Raia-pintada	<i>Raja asterias</i>	Mediterranean starry ray	45
Raia-ponteada	<i>Raja brachyura</i>	Blonde ray	45
Raia-pregada	<i>Leucoraja fullonica</i>	Shagreen ray	44
Raia-redonda	<i>Rajella fyllae</i>	Round ray	48
Raia-repregada	<i>Amblyraja radiata</i>	Starry ray	42
Raia-tairoga	<i>Rostroraja alba</i>	White skate	48
Raia-zimbreira	<i>Raja microocellata</i>	Small-eyed ray	47
Rainha-branca	<i>Pseudotolithus typus</i>	Longneck croaker	276
Rainúnculo-negro	<i>Raniceps raninus</i>	Tadpole fish	143
Rascasso-da-fundura	<i>Etreposebastes imus</i>	Midwater scorpionfish	229
Rascasso-da-Madeira	<i>Scorpaena maderensis</i>	Madeira rockfish	231
Rascasso-das-Canárias	<i>Scorpaena canariensis</i>	Canary scorpionfish	230
Rascasso-de-barbatana-pontuada	<i>Scorpaena stephanica</i>	Spotted-fin rockfish	233
Rascasso-de-Loppé	<i>Scorpaena loppei</i>	Cadenat's scorpionfish	231
Rascasso-de-Messina	<i>Scorpaenodes arenai</i>	Messina rockfish	234
Rascasso-de-pintas	<i>Scorpaena porcus</i>	Black scorpionfish	232
Rascasso-de-tubo-curto	<i>Phenacoscorpius nebris</i>	Short-tube scorpionfish	230
Rascasso-dos-Açores	<i>Scorpaena azorica</i>	Azores scorpionfish	230
Rascasso-do-Senegal	<i>Scorpaena laevis</i>	Senegalese rockfish	231
Rascasso-escorpião	<i>Scorpaena notata</i>	Small red scorpionfish	232
Rascasso-negro	<i>Scorpaena plumieri</i>	Pacific spotted scorpionfish	232
Rascasso-rosado	<i>Scorpaena elongata</i>	Slender rockfish	230
Rascasso-vermelho	<i>Scorpaena scrofa</i>	Red scorpionfish	233
Ratão-água	<i>Myliobatis aquila</i>	Common eagle ray	51
Ratão-bispo	<i>Aetomylaeus bovinus</i>	Bull ray	51
Ratazana-da-fundura	<i>Hydrolagus affinis</i>	Smalleyed rabbitfish	54
Ratazana-lusitânica	<i>Hydrolagus lusitanicus</i>	No English common name	54
Ratazana-olhuda	<i>Hydrolagus mirabilis</i>	Large-eyed rabbitfish	54
Ratazana-pálida	<i>Hydrolagus pallidus</i>	Pale chimaera	54
Rei-dos-Arenques	<i>Regalecus glesne</i>	King of herrings	128
Requeime-de-natura	<i>Setarches guentheri</i>	Channeled rockfish	229
Riscado	<i>Parapristipoma octolineatum</i>	African striped grunt	262
Robalo-baila	<i>Dicentrarchus punctatus</i>	Spotted seabass	241
Robalo-legítimo	<i>Dicentrarchus labrax</i>	European seabass	241
Rodovalho	<i>Scophthalmus rhombus</i>	Brill	188
Rodovalho-bruxa	<i>Zeugopterus punctatus</i>	Topknot	188
Roncadeira-preta	<i>Sciaena umbra</i>	Brown meagre	276
Roncador-bravura	<i>Pomadasys incisus</i>	Bastard grunt	263
Roncador-canela	<i>Parapristipoma humile</i>	Guinean grunt	262
Roncador-colo-colô	<i>Brachydeuterus auritus</i>	Bigeye grunt	262

Portuguese common name	Scientific name	English common name	Page
Roncador-papagaio	<i>Pomadasys perotaei</i>	Parrot grunt	263
Rosa-salmão	<i>Etelis oculatus</i>	Queen snapper	261
Ruivo	<i>Lepidotrigla cavillone</i>	Large-scaled gurnard	236
Ruivo-espinhoso	<i>Lepidotrigla dieuzeidei</i>	Spiny gurnard	236
Salema	<i>Sarpa salpa</i>	Salema	273
Salmão-do-Atlântico	<i>Salmo salar</i>	Atlantic salmon	85
Salmonete-da-vasa	<i>Mullus barbatus</i>	Red mullet	277
Salmonete-do-alto	<i>Polymixia nobilis</i>	Stout beardfish	149
Salmonete-legítimo	<i>Mullus surmuletus</i>	Surmullet	277
Sapata	<i>Deania calcea</i>	Birdbeak dogfish	37
Sapata-áspera	<i>Deania hystricosa</i>	Rough longnose dogfish	37
Sapata-flecha	<i>Deania profundorum</i>	Longsnout dogfish	38
Sapata-preta	<i>Centroscelachus crepidater</i>	Roughskin dogfish	33
Sarda	<i>Scomber scombrus</i>	Atlantic mackerel	168
Sardinela-da-Madeira	<i>Sardinella maderensis</i>	Madeiran sardinella	73
Sardinela-lombuda	<i>Sardinella aurita</i>	Round sardinella	73
Sardinha	<i>Sardina pilchardus</i>	European pilchard	72
Sardinha-escamuda	<i>Harengula clupeola</i>	False herring	72
Sargo-alcorraz	<i>Diplodus annularis</i>	Annular seabream	265
Sargo-bicudo	<i>Diplodus puntazzo</i>	Sharpsnout seabream	267
Sargo-de-Cabo-Verde	<i>Diplodus lineatus</i>	White seabream	267
Sargo-de-faixas	<i>Diplodus fasciatus</i>	Banded seabream	267
Sargo-do-Senegal	<i>Diplodus bellottii</i>	Senegal seabream	266
Sargo-legítimo-do-Atlântico	<i>Diplodus cadenati</i>	White seabream	266
Sargo-legítimo-do-Mediterrâneo	<i>Diplodus sargus</i>	White seabream	267
Sargo-safia	<i>Diplodus vulgaris</i>	Common two-banded seabream	268
Sargo-veado	<i>Diplodus cervinus</i>	Zebra seabream	266
Sarrajão	<i>Sarda sarda</i>	Atlantic bonito	167
Sável	<i>Alosa alosa</i>	Allis shad	71
Savelha	<i>Alosa fallax</i>	Twaite shad	72
sem nome comum	<i>Leptocephalus giganteus</i>	Giant leptocephalus	57
Sereia-camochilo	<i>Trachinotus ovatus</i>	Pompano	256
Sereia-da-Florida	<i>Trachinotus carolinus</i>	Florida pompano	256
Serra-branca	<i>Scomberomorus tritor</i>	West African Spanish mackerel	169
Serra-da-Índia	<i>Acanthocybium solandri</i>	Wahoo	166
Serrano-alecrim	<i>Serranus cabrilla</i>	Comber	246
Serrano-de-rolo	<i>Serranus atricauda</i>	Blacktail comber	245
Serrano-ferreiro	<i>Serranus hepatus</i>	Brown comber	246
Serrano-riscado	<i>Serranus scriba</i>	Painted comber	246
Solha	<i>Pleuronectes platessa</i>	European plaice	191
Solha-das-pedras	<i>Platichthys flesus</i>	European flounder	191
Solha-limão	<i>Microstomus kitt</i>	Lemon sole	191
Solhão	<i>Glyptocephalus cynoglossus</i>	Witch flounder	191
Sugador-bochechudo	<i>Apletodon pellegrini</i>	Chubby clingfish	205
Sugador-das-Canárias	<i>Diplecogaster pectoralis</i>	No English common name	205
Sugador-de-cabeça-pequena	<i>Apletodon dentatus</i>	Small-headed clingfish	204
Sugador-de-De Candolle	<i>Lepadogaster candolii</i>	Connemarra clingfish	205
Sugador-de-duas-manchas	<i>Diplecogaster bimaculata</i>	Two-spotted clingfish	205
Sugador-de-Gouan	<i>Lepadogaster lepadogaster</i>	Shore clingfish	205
Sugador-do-Mediterrâneo	<i>Opeanogenys gracilis</i>	No English common name	206
Sugador-dos-ouriços	<i>Apletodon incognitus</i>	No English common name	204
Sugador-púrpura	<i>Lepadogaster purpurea</i>	Cornish sucker	205
Suspensório	<i>Cepola macrophthalmia</i>	Red bandfish	278
Tainha-de-salto	<i>Chelon saliens</i>	Leaping mullet	203
Tainha-fataça	<i>Chelon ramada</i>	Thinlip grey mullet	202

Portuguese common name	Scientific name	English common name	Page
Tainha-garrento	<i>Liza aurata</i>	Golden grey mullet	203
Tainha-liça	<i>Chelon labrosus</i>	Thicklip grey mullet	202
Tainha-olhalvo	<i>Mugil cephalus</i>	Flathead grey mullet	204
Tainha-sabão	<i>Oedalechilus labeo</i>	Boxlip mullet	204
Tamboril	<i>Lophius piscatorius</i>	Angler	212
Tamboril-sovaco-preto	<i>Lophius budegassa</i>	Blackbellied angler	212
Tarpão-do-Atlântico	<i>Megalops atlanticus</i>	Tarpon	55
Tintureira	<i>Prionace glauca</i>	Blue shark	28
Tirone	<i>Cubiceps gracilis</i>	Driftfish	172
Tremelga-de-olhos	<i>Torpedo torpedo</i>	Common torpedo	41
Tremelga-marmoreada	<i>Torpedo marmorata</i>	Marbled electric ray	41
Tremelga-negra	<i>Tetronarce nobiliana</i>	Electric ray	40
Trombeiro-boga	<i>Spicara smaris</i>	Picarel	274
Trombeiro-choupa	<i>Spicara maena</i>	Blotched picarel	274
Trombeiro-do-Mediterrâneo	<i>Spicara flexuosa</i>	No English common name	274
Trombeiro-malha-redonda	<i>Spicara melanurus</i>	Blackspot picarel	274
Trombeta	<i>Aulostomus strigosus</i>	Atlantic cornetfish	174
Trombeta-salpicada	<i>Aulostomus cf. maculatus</i>	Atlantic trumpetfish	174
Trombeteiro	<i>Macroramphosus scolopax</i>	Longspine snipefish	174
Trombeteiro-fino	<i>Macroramphosus gracilis</i>	Slender snipefish	174
Truta-arco-íris	<i>Oncorhynchus mykiss</i>	Rainbow trout	85
Truta-marisca	<i>Salmo trutta</i>	Sea trout	86
Tubarão-alfafar	<i>Hexanchus griseus</i>	Bluntnose sixgill shark	16
Tubarão-alfafar-de-pontas-brancas	<i>Hexanchus nakamurai</i>	Bluntnose sixgill shark	16
Tubarão-anão	<i>Squaliolus laticaudus</i>	Spined pygmy shark	30
Tubarão-anequim	<i>Isurus oxyrinchus</i>	Longfin mako	18
Tubarão-anequim-de-gadanha	<i>Isurus paucus</i>	Longfin mako	18
Tubarão-areia	<i>Odontaspis ferox</i>	Sand tiger shark	17
Tubarão-baboso	<i>Carcharhinus altimus</i>	Bignose shark	24
Tubarão-baleia	<i>Rhincodon typus</i>	Whale shark	17
Tubarão-bicudo	<i>Rhizoprionodon acutus</i>	Milk shark	28
Tubarão-branco	<i>Carcharodon carcharias</i>	Great white shark	18
Tubarão-buldogue	<i>Carcharhinus leucas</i>	Pigeye shark	26
Tubarão-cão-dos-Açores	<i>Scymnodalatias garricki</i>	Azores dogfish	33
Tubarão-cobra	<i>Chlamydoselachus anguineus</i>	Frilled shark	16
Tubarão-cobre	<i>Carcharhinus brachyurus</i>	Copper shark	24
Tubarão-corre-costa	<i>Carcharhinus plumbeus</i>	Sandbar shark	27
Tubarão-da-Gronelândia	<i>Somniosus microcephalus</i>	Greenland shark	34
Tubarão-da-noite	<i>Carcharhinus signatus</i>	Night shark	27
Tubarão-de-Noronha	<i>Odontaspis noronhai</i>	Bigeye sand tiger shark	17
Tubarão-dentão	<i>Isistius plutodus</i>	Largetooth cookiecutter shark	30
Tubarão-dente-liso	<i>Carcharhinus isodon</i>	Finetooth shark	25
Tubarão-de-pontas-brancas	<i>Carcharhinus longimanus</i>	Oceanic whitetip shark	26
Tubarão-de-pontas-negras	<i>Carcharhinus limbatus</i>	Blacktip reef shark	26
Tubarão-de-sete-guelras	<i>Heptranchias perlo</i>	Sharpnose sevengill shark	15
Tubarão-dos-Galápagos	<i>Carcharhinus galapagensis</i>	Galapagos shark	25
Tubarão-duede	<i>Mitsukurina owstoni</i>	Goblin shark	17
Tubarão-faquetá	<i>Carcharhinus obscurus</i>	Dusky shark	27
Tubarão-frade	<i>Cetorhinus maximus</i>	Basking shark	19
Tubarão-gato-abissal	<i>Apristurus profundorum</i>	Deep-water catshark	21
Tubarão-gato-da-Madeira	<i>Apristurus laurussonii</i>	Madeira catshark	20
Tubarão-gato-fantasma	<i>Apristurus manis</i>	Ghost catshark	20
Tubarão-gato-olhinho	<i>Apristurus microps</i>	Smalleye catshark	20
Tubarão-limão	<i>Negaprion brevirostris</i>	Lemon shark	28
Tubarão-luminoso	<i>Isistius brasiliensis</i>	Cookie cutter shark	30

Portuguese common name	Scientific name	English common name	Page
Tubarão-lusitano	<i>Centrophorus lusitanicus</i>	Lowfin gulper shark	35
Tubarão-luzidio	<i>Carcharhinus falciformis</i>	Silky shark	25
Tubarão-martelo	<i>Sphyraena zygaena</i>	Smooth hammerhead	29
Tubarão-martelo-gigante	<i>Sphyraena mokarran</i>	Great hammerhead	29
Tubarão-martelo-recortado	<i>Sphyraena lewini</i>	Scalloped hammerhead	28
Tubarão-mona	<i>Pseudotriakis microdon</i>	False catshark	22
Tubarão-prego	<i>Echinorhinus brucus</i>	Bramble shark	39
Tubarão-raposo	<i>Alopias vulpinus</i>	Pelagic thresher or Bigeye thresher	20
Tubarão-raposo-olhudo	<i>Alopias superciliosus</i>	Pelagic thresher or Thresher	19
Tubarão-sardo	<i>Lamna nasus</i>	Porbeagle	19
Tubarão-tecelão	<i>Carcharhinus brevipinna</i>	Graceful shark	25
Tubarão-tigre	<i>Galeocerdo cuvier</i>	Tiger shark	27
Tubarão-toiro	<i>Carcharias taurus</i>	Smalltooth sand tiger shark	17
Uge	<i>Dasyatis pastinaca</i>	Common stingray	49
Uge-da-Madeira	<i>Gymnura hirundo</i>	Madeira butterfly ray	50
Uge-de-cardas	<i>Bathyrajia lata</i>	Brown stingray	49
Uge-manta	<i>Gymnura altavela</i>	Spiny butterfly ray	50
Uge-redonda	<i>Taeniurops grabatus</i>	Round stingray	50
Uge-violeta	<i>Pteroplatytrygon violacea</i>	Pelagic stingray	50
Veleiro-do-Indo-Pacífico	<i>Istiophorus platypterus</i>	Indo-Pacific sailfish	185
Verdinho	<i>Micromesistius poutassou</i>	Blue whiting	143
Vermelho-salmão	<i>Etelis carbunculus</i>	Deep-water red snapper	261
Viola	<i>Rhinobatos rhinobatos</i>	Common guitarfish	41
Viola-barba-negra	<i>Glaucostegus cemiculus</i>	Blackchin guitarfish	42
Xaputa	<i>Brama brama</i>	Atlantic pomfret	259
Xaputa-galhuda	<i>Taractichthys longipinnis</i>	Big-scale pomfret	260
Xaputa-negra	<i>Taractes rubescens</i>	Black pomfret	260
Xaputa-rugosa	<i>Taractes asper</i>	Rough pomfret	259
Xara-preta-de-natura	<i>Centroscymnus owstonii</i>	Roughskin dogfish	32
Xareu-amarelo	<i>Caranx rhonchus</i>	False scad	251
Xareu-azul	<i>Caranx cryos</i>	Blue runner	251
Xareu-bicudo	<i>Pseudocaranx dentex</i>	White trevally	253
Xareu-carvoeiro	<i>Caranx ruber</i>	Bar jack	252
Xareu-de-olho-papudo	<i>Caranx latus</i>	Horse-eye jack	251
Xareu-enforcado	<i>Caranx lugubris</i>	Black jack	251
Xareu-enxada	<i>Alectis alexandrina</i>	Alexandria pompano	250
Xareu-macoa	<i>Caranx hippos</i>	Crevalle jack	251
Xareu-palheta	<i>Campogramma glayco</i>	Vadigo	250
Xarinha-preta	<i>Etomopterus pusillus</i>	Smooth lanternshark	31



REPÚBLICA
PORTUGUESA

MAR

