

## Plants and People in Mooro Country

Noongar Plant Use in Yellagonga Regional Park



### Contents

Yellagonga Regional Park: A Changed Landscape	9
Aboriginal Settlement	9
Exploration and Invasion	11
Colonisation and Dispossession	13
Wineries and Market Gardens	15
A Changed Landscape – Yellagonga Regional Park Today	15
Mooro Plants and People: Yellagonga Regional Park	17
Plant Species: Noongar Uses	19
Banksia: Mungite, Piara, Pulgart, Pungura	21
Bottlebrush: Kwowdjard	25
Daisy: Yoont Djet	27
Eucalypt: Jarrah, Koodjat, Marri, Moitch, Tuart	29
Grass Tree: Balga	33
Grevillea: Berrung	37
Hakea: Berrung, Pulgur	39
Kangaroo Paw: Kurulbrang	41
Lily	43
Melaleuca: Yowarl	47
Orchid: Kararr	51
Pea Flower: Koorla, Koorpa, Koweda, Marno, Pulboorn, Puyenak, Yackal Djarr	55
Rush, Sedge: Waakal Ngarnak	59
Solanum	63
Spearwood: Kitja Boorn	65
Wattle: Coojong, Panjang, Wilyawa	67
Zamia: Bayu, Djiridji	71
Appendix 1: Species List	75
Appendix 2: Noongar Seasons	85
Appendix 3: Reference List	87




#### A Note on Historical Accuracy and Noongar Vocabulary

Information on Noongar plant use has been garnered from personal communications, historical, botanical and anthropological research, early-colonial reports, and the diaries and journals of early colonisers. It is acknowledged that such references can be problematic and, where appropriate, sources have been referenced.

In constructing this publication, the researchers have attempted to use appropriate Noongar names for plant species wherever possible. Due to dialectal differences and early inconsistencies, many species have several names and/or different spellings. A full list of Noongar plant names and their sources has been included at Appendix 1.

The researchers have aimed to be as accurate as possible with regard to the use of plants by local Noongar people. However, due to the nature of the records available, it is possible that details have been unintentionally misrepresented. A bibliographical Reference List has been provided at Appendix 3 and includes all sources examined in the development of this work.





# A Changed Landscape

Yellagonga Regional Park is located on the Swan Coastal Plain within the Cities of Joondalup and Wanneroo. The Park contains a string of wetlands and swamps which form part of the 'Linear Lakes', an important north-south link with Neerabup National Park and Yanchep National Park. Included in Yellagonga Regional Park are Lake Joondalup<sup>1</sup>, Beenyup<sup>2</sup> Swamp, Walluburnup<sup>3</sup> Swamp and Lake Goollelal<sup>4</sup>.

Yellagonga Regional Park contains a diversity of eco-systems and represents flora and fauna communities that were once widespread on the Swan Coastal Plain. The Park's wetlands represent some of the last remaining freshwater systems in the Perth Metropolitan Area (DEC 2003). In addition to the abundant plant life, these wetlands provide an important breeding ground for local birds, reptiles and amphibians (DEC 2003). The largest lake in the Park, Lake Joondalup, is listed on the Register of the National Estate and is considered to be of national significance.

#### Aboriginal Settlement:

According to archaeological evidence, Noongar<sup>5</sup> people were occupying the area around Yellagonga Regional Park for at least 40,000 years prior to European colonisation (Hallam 1989:145-147). The country surrounding the Yellagonga Regional Park area was called 'Mooro' and was frequented by several large family groups up until the early-1900s. The Regional Park itself is named after an important Noongar elder of the early colonial period, Yellagonga<sup>6</sup>. Early colonist Robert Menli Lyon, believed that "Mooro, the district of Yellowgonga...is bounded by the sea on the west; by Melville water and the Swan, on the south; by Ellen's brook, on the east; and, by the Gyngoorda, on the north." (Lyon 1833:176). Prior to large-scale European settlement, it is likely that the vicinity which is now the Perth Central Business District was the focal area of Mooro Country, with Yellagonga Regional Park playing an integral role (Hallam & Tillbrook 1990:349).

- 'Joondalup' is a Noongar word which may mean 'place of whiteness or glistening', 'place of a creature that can only move backwards' or 'place of the long, white hair'.
- <sup>2</sup> 'Beenyup' is a Noongar word which may mean 'digging place' or 'place of native yams'.
- <sup>3</sup> 'Walluburnup' is a Noongar word which may mean 'open space between two trees', 'place of fish and wallaby' or 'fish in lake'.
- <sup>4</sup> 'Goollelal' is a Noongar word which may mean 'swampy sheoak' or 'place for camp'.
- <sup>5</sup> Also written as 'Noongar', 'Nyoongar', 'Nyoongah' and 'Nyungah'.
- 6 Also written as 'Yellowgonga', 'Yalagonga', 'Yalagongga', 'Yalgongga', 'Yalgongga', 'Yallagonga', 'Yallagonga', 'Yallowgonga', 'Yallowgonga 'Yelloganga', 'Yellogonga', 'Yellowgongo', 'Barragim', 'Drooreer', 'Naraganianda' and 'Ngalgonga' (Hallam & Tillbrook 1990:348-354).



A census recorded by colonist Francis Armstrong in 1837<sup>7</sup> showed there to be 28 Mooro Noongars (Armstrong 1836:192), although it is not possible to confirm this number.

For local Noongar people, the Yellagonga Regional Park area holds considerable significance. The wetlands have been used extensively for hunting, food-gathering, social, ceremonial and recreational purposes and formed part of a north-south link of rivers, lakes and wetlands along the Swan Coastal Plain. The Yellagonga Regional Park area was particularly important during the autumn and spring months, when it was utilised as a semi-permanent camping ground. The natural north-south routes, which form the Linear Lakes, provided important access tracks to other camping areas and acted as trading routes between family and tribal groups (McGuire 1996:58; O'Connor et al 1989:27).

#### Exploration and Invasion:

The crew of the 1696 Dutch voyage under Captain Willem de Vlamingh were probably among the first Europeans to see signs of Mooro people. The ships travelled approximately 80 kilometres up the Swan River, where the crews briefly disembarked. They found the camps and burning fires of Mooro Noongars, but they did not see any people. The de Vlamingh expedition resulted in the name 'Swarte Swaene-Revier' (Black Swan River), named after the black swans they saw there. Throughout the 18th and 19th Centuries, numerous other European expeditions to the south-west coast followed. Many of these voyages noted similar signs of the local inhabitants, however, few ever saw any people.

Following the official invasion of New South Wales in 1770, and in spite of a French claim of sovereignty in 1772, a British military outpost was established at Albany in 1826. One year later, an expedition under Captain James Stirling explored the Swan River, which he described as an ideal place to establish a permanent settlement. Subsequent to lobbying in Britain, a fleet was sent and Charles Fremantle, being the first to land, declared the Swan River Colony for Britain in 1829.

With the beginning of colonisation, land in the Perth area was initially taken up around the Swan River. Most of the land however, was considered to be of very poor quality and early reports sent back to England stated that the colony was near starvation (Berryman 2002). The flow of migrants to the colony dwindled, and by 1850, the European population had only reached 5,886. The majority of colonisers settled around the south-western coastline at Albany, Augusta and Bunbury. The quest for suitable farming land eventually compelled the colonisers to explore the region north of the Swan River and various expeditions were conducted into the Yellagonga Regional Park area throughout the early 1800s.

In 1834, colonist John Butler passed Lake Joondalup whilst searching for lost cattle. During his visit, Butler met some local Mooro people whom he described as "those Wannaroo men who frequent Perth in company with the Yellowgonga Tribe". He believed the local Aboriginal people to be friendly towards Europeans and advised a surveying party be sent to the area (Butler 1834:346). The survey was completed in 1837 by Thomas Watson and the land was subsequently taken up by various colonisers. These plots of land however, were never settled by their new owners and appear to have been part of what has been described as a 'land-grab', where well-off migrants acquired extra land for speculative purposes (Russo 1998:115).

Further exploratory work in the Yellagonga Regional Park area was conducted by George Grey in 1838. Grey encountered several local Mooro people with whom he tested his newly acquired language skills. Near Lake Joondalup, Grey met the Mooro Aborigines Noogongoo8, Kurral, Jeebar9, and Dudemurry10 who fed Grey's party tortoises and talked with Grey throughout the night. Grey writes "They said that, although the lake was called Mooloore, the name of the land we were sitting on was Doondalup" (Grey 1841a). Grey concluded that the land surrounding the lakes was "of the best quality" and that there was "plenty of good feed for cattle" (Grey 1841a). Grey passed through the region again in 1839 after his ship was wrecked at

<sup>&</sup>lt;sup>7</sup> The Swan River Colony was founded 8 years earlier in 1829.

<sup>&</sup>lt;sup>8</sup> Also written as 'Byerman', 'Biarman', 'Ngogonga', 'Ngoogonga', 'Ngoogoongga', and 'Nogongo' (Hallam & Tillbrook 1990:41).

<sup>&</sup>lt;sup>9</sup> Also written as 'Djibar', 'Jibar' and 'Dukadung' (Hallam & Tillbrook 1990:165).

<sup>10</sup> Also written as 'Dutomerrar', 'Decamurry', 'Djitamarra', 'Dudamurra', 'Dutermerry', Dutomerra', and 'Dutomurra' (Hallam & Tillbrook 1990:101).



Kalbarri and the party were forced to walk back to Perth. Mooro people in the Yellagonga Regional Park area provided the starving men with frogs, tortoises and zamia nuts (Grey 1841b).

Following Grey, the Surveyor-General, John Septimus Roe, escorted Governor John Hutt to the Yellagonga Regional Park area in 1841. Roe freely used the Noongar names for the lakes, 'Joondalup', 'Goollelal', 'Needubup' and 'Nowergup', and expressed his opinion that the area was suitable for settlement (Russo 1998:115). In 1842, a group of vagrant soldiers strayed to the area and for a few years Lake Joondalup was called 'Soldiers' Lake' (Russo 2998:115). The soldiers' settlement however, was short-lived and the name was seldom used again.

#### Colonisation and Dispossession:

The first substantial effort to make contact with Mooro people was the 'Native Experimental Farm' established by the Wesleyan Reverend John Smithies. Smithies arrived in the colony in 1840 with instructions from the Methodist Church to both care for the pastoral needs of the colony's Methodists, and convert the local Aboriginal population to the Wesleyan branch of Christianity. In 1843, Smithies requested permission from London to set up an Aboriginal Mission outside the main town of Perth, and an establishment was built on the banks of Lake Goollelal soon after. The establishment was dubbed 'Mission Farm', and aside from the 'Christianisation' of the local Aborigines, the general purpose of Smithies' Mission, was to encourage local Noongar people to move off their lands and integrate into European 'civilisation'. Smithies' Mission attempted to educate local Mooro children in farming skills and housework so that they could enter into servitude for the European colonisers. It was hoped that the 'civilising' of the Mooro population would also assist in reducing the gross shortfall in labour the floundering colony was experiencing at the time. Unfortunately for Smithies, the land at Lake Goollelal was unsuitable for crop planting and the Yellagonga Regional Park area provided such a plentiful supply of food and shelter that the Mission school found it could neither attract nor retain its Noongar pupils (Cook 1966:21-22). Furthermore, a significant number of children at Mission Farm died as a consequence of introduced diseases and, as a result, the local Aboriginal population refused to allow their children to stay there (Monks 1993:10-11). After less than nine years 'Mission Farm' was abandoned by the Methodists, and Smithies moved on to better prospects in the York region. Reverend John Smithies Park, located on the banks of Lake Goollelal, has been named after the Wesleyan Missionary.

The first European to permanently occupy the Yellagonga Regional Park area was James Cockman who took up land there in 1852. James Cockman with his wife Mary built a small house close to Walluburnup Swamp and, some years later, built a larger stone house close to the corner of Woodvale Drive and Wanneroo Road (Newton 2002). With the expansion of the Perth township during the mid-19th Century, food depots were created to discourage Aboriginal people from going into the town. During this period, Mooro people retreated north to the lakes (McGuire 1996:95). When Cockman resided in the area, Aboriginal people still led traditional lives in this part of Mooro Country (Brittain 1990:56). Cockman's grandson, Cecil Malcolm Cockman, grew up in the same stone house, and remembered local Aborigines still camping by Lake Joondalup in the early-1900s when he was young (Marwick 2002:6). Following the Cockmans, the Buckingham family took up land on the eastern side of Lake Joondalup in 1860. Between the 1860s and 1880s, several more families took up land in the Yellagonga Regional Park area, including those of Okely, Lander, Backshall, Darch, Duffy, Shenton, Thompson and Leach (Brittain 1990:56; Russo 1998:120). By the end of the 19th Century, the well-established north-south link had become increasingly important to the European colonisers as well.

Various overland expeditions traversed the lakes area over this period and mining exploration was also carried out. During the late-1800s, a stock route was established with sheep, cattle and horses being watered and grazed at the lakes as they were driven northwards. A mail-carrying service to northern homesteads was established in 1853 and a small cave guano harvesting industry was attempted. As additional land was taken up around Perth, colonists would travel to the northern areas for game hunting as the animal population around the Swan River became depleted. By 1888 the entire North-West Corridor



had been taken up by pastoral leases and activity along the lakes had increased considerably. Within only 60 years the traditional land and food sources of Mooro Country had been severely diminished.

By the end of the 19th Century, various laws had been passed that significantly restricted the rights of Aboriginal people in Western Australia. The Industrial Schools Act 1874 effectively introduced the institutionalisation of Aborigines by allowing Noongar children to be placed into the care of the State. The Aborigines Protection Act 1886 enabled the newly formed Aborigines Protection Board to indenture any Aboriginal child of a 'suitable age' into an apprenticeship until the age of 21. The Act also prohibited Aboriginal people from entering or remaining in towns, including Perth. Probably the most far-reaching legislation was the Aborigines Act 1905 which established the Chief Protector of Aborigines as the legal guardian of 'every Aboriginal and half-caste child' to the age of 16 years. This Act allowed for the forced removal of anyone deemed to be an 'Aboriginal native' to a 'Native Reserve' and any child to a State institution. The effect of the Aborigines Act 1905 was to considerably limit Aboriginal access to land, water, housing, employment and education and significantly reduce the ability of Aboriginal people to secure food and income (Delmege 2005). Within this context, Mooro people found themselves driven further and further to the boundaries of their Country. 'Native Reserves' were designated to the fringe areas of Perth and, by the beginning of the 20th Century, the vast majority of Mooro Noongars had been either compelled or forcibly removed from the colonised areas of Perth.

#### Wineries and Market Gardens:

The landscape of the Yellagonga Regional Park area continued to change as it was moulded by successive farming operations. Post-World War I, there was an influx of non-British migrants to Perth, particularly from Italian, Greek and Yugoslav backgrounds. The Linear Lakes area became an important source of fruit and vegetables for the entire Perth region. Between the World Wars, migrants, including the Formiatti, Crisafulli, Ariti, and Nanovich families, took up land around Yellagonga Regional Park. Other southern European families, including the Contis, Parins and Luisinis later moved to the area and set up wineries, some of which are still in operation today.

Increased market gardening and viticulture operations resulted in a significant demand for timber. Timber mills were set up along the Yellagonga Regional Park area and a burgeoning industry was developed using locally-sourced trees. A limestone industry was also operating successfully in the Yellagonga Regional Park area at this time. Limestone quarried from around the lakes was profitably used throughout Wanneroo for construction purposes (Russo 1998:113). Lime burning was also conducted and kilns were set up from the northern end of Lake Joondalup all the way south to Lake Coogee.

Following the Second World War, Perth experienced a population boom and the City expanded rapidly north and south. Much of the land previously farmed for fruit and vegetables was re-zoned for residential purposes and the suburbs of Perth gradually spread further and further out from the City Centre. During the 1960s and 1970s, the area north of Perth experienced rapid growth and became known as the 'Mortgage Belt' (Russo 1998:137). By the early 1980s, the Shire of Wanneroo had become increasingly urbanised and the Yellagonga Regional Park area was declared public open space.

#### A Changed Landscape – Yellagonga Regional Park Today:

The relationship between Noongar culture and the environment is one which "relies upon but also sustains the natural resources" (Cherikoff 1993:22). Mooro people adapted the environment to themselves and employed sustainable cultivation practices, such as controlled burning and returning roots and seeds to the soil. With the invasion of Europeans in the early-1800s, the landscape of Yellagonga Regional Park was rapidly and considerably altered. Land was cleared for farming and development, channels were cut for irrigation and drainage, limestone was quarried, trees were cut down for timber and many foreign flora species were planted. Yellagonga Regional Park today is a changed landscape, having been shaped by its distinct and varied past.




## Mooro Plants & People Yellagonga Regional Park

The people of the Mooro Country possessed an intimate knowledge of the local ecology. Contrary to popular belief, Aboriginal people did not "wander the continent in search of food in order to survive in a harsh and desolate land" (Isaacs 2002:43). Rather, they held a meticulous knowledge of their own welldefined area, their Country. The lessons and discoveries about their Country were passed down from one generation to the next, largely by oral tradition. The way to use and care for the land was 'written' into stories and songs of the Dreaming (Cherikoff 1994:24). Knowledge from the Dreaming taught people how the spirit beings made foods and medicines from the bush as well as formed the lakes, rivers and mountains.

Plants were extremely important to Noongar people. Different plants were used to create weapons, such as spears and shields, to build shelters, for medicinal purposes, and for food. Probably the most important of these uses was food. In Mooro Country, the abundance and diversity of plant species ensured that local Noongar people utilised a substantial number of plants for a variety of purposes. The flowers, stems, leaves, bark, gum, resin and roots of many plants were all used. Many Noongar plant names are utilised in today's vernacular, including 'Jarrah', 'Marri', 'Tuart', 'Wandoo', 'Bullich', 'Yarri', 'Moonah', 'Quandong', and 'Pingle'.







## Banksia

#### Mungite, Piara, Pulgart, Pungura

#### Species found in Yellagonga Regional Park:

- Banksia attenuata
   Banksia grandis
  - Banksia ilicifolia
- Banksia littoralis
- Banksia menziesii
   Banksia prionotes
   Banksia sessilis

There are over 150 species of banksias, 90% of which occur in south-western Australia. Banksia's are characterised by their sharp, serrated leaves and large, cone-shaped flowers.

In Yellagonga Regional Park, there are at least 7 species of banksia. Banksias are very important to Aboriginal people and several Noongar names for them are known. The Candle Banksia (or Candlestick Banksia, Coast Banksia, Slender Banksia) (Banksia attenuata) is known as the Piara (or Biara, Bealwra, Peera, Piras), the Bull Banksia (or Giant Banksia, Great-Flowered Banksia) (Banksia grandis) is known as the Mungite (or Poolgarla, Bulgalla), the Swamp Banksia (or River Banksia, Seaside Banksia, Swamp Oak, Western Swamp Banksia) (Banksia littoralis) is known as the Pungura (or Boongura, Gwangia), and the Parrot Bush (Banksia sessilis) is known as the Pulgart. The Holly-Leaved Banksia (or Holly Banksia) (Banksia ilicifolia), Acorn Banksia (or Orange Banksia, Saw-Tooth Banksia (Banksia prionotes), and the Firewood Banksia (or Menzies Banksia) (Banksia menziesii) are present in Yellagonga Regional Park also.

Banksia flowers produce an abundance of honey-like nectar, which is why the early colonists called this plant the **Honeysuckle**. Noongar people drink the honey straight out of the flower cone, or soak the flower in water to produce a sweet drink. This beverage is either drunk fresh or fermented to produce Gep, an intoxicating liquor. The early colonists also used the nectar of the banksia for honey and to make sweet drinks. Early 20th Century writer, Dame Mary Gilmore, described the use of banksia drinks in the treatment of sore throats and colds.



Some banksias, such as the Piara and the Mungite are used by Noongar people as torches. When alight, the dried banksia flower cone smoulders like a torch, these were used by local Noongars to transport fire from one campsite to the next. Noongar people also kept the lighted cones under their cloaks to keep themselves warm in cold weather.

The **Pulgart** is an unusual type of banksia, as it has a squat, round flower instead of a long cone. This banksia has very spiky leaves and branches and is used by Noongar people as a broom. The Pulgart is also utilised in fishing. Noongar fishermen break off the branches and walk in a line, driving the djildjit (fish), yakan (turtle) or koonak (freshwater prawns) into the fish traps.

The wood of many banksia trees is also used as firewood. In particular the **Firewood Banksia** is known for its quick burning properties.













## Bottlebrush

#### Kwowdjard

#### Species found in Yellagonga Regional Park:

- Calothamnus quadrifidus
- Calothamnus sanguineus

The term 'bottlebrush' is used to describe plants of the Callistemon genus which are usually characterised by their bottlebrush-shaped flowers.

There are at least 2 species of bottlebrush in the Yellagonga Regional Park area, the Silky-Leaved Blood Flower (Calothamnus sanguineus) and the One-Sided Bottlebrush (Calothamnus quadrifidus). The Noongar name for the One-Sided Bottlebrush is the Kwowdjard (or Queitiat).

Similar to other flowering plants, the blossoms of the bottlebrush are useful to Noongar people as a source of honey. Noongars suck the sweet nectar straight from the flower blossoms or they soak the flowers in water to produce a sweet drink. From time to time, this drink is allowed to ferment to produce Gep, an intoxicating liquor.









## Daisy Yoont Djet

#### Species found in Yellagonga Regional Park:


Lagenophora huegelii

- Olearia axillaris
- Senecio pinnatifolius (var. maritimus)
- Waitzia suaveolens

'Daisy' is a generic term that refers to plants that belong to the Asteraceae Family. Daisy plants vary in size, but are usually herbs. Daisy flowers are generally characterised by a round, central stigma surrounded by numerous petals.

In Yellagonga Regional Park there are several species of daisies. Abundant species include the **Coarse** Lagenophora (Lagenophora huegelii), Coastal Daisybush (Olearia axillaris), Coastal Groundsel (or Variable Groundsel) (Senecio pinnatifolius [var. maritimus]), and the Fragrant Waitzia (Waitzia suaveolens). The Noongar name for the Coastal Groundsel is Yoont Djet.

It is not known whether daisies were of particular importance to traditional Noongar people. However, it is believed that the early colonisers occasionally used daisies in cooking. The crushed leaves of the Coastal Daisybush for example, have a pleasant smell and were sometimes used as a herb. As early as 1696, the crew of Willem de Vlamingh's expedition used this daisybush to add flavour to their food.









## Eucalypt

#### Jarrah, Koodjat, Marri, Moitch, Tuart

#### Species found in Yellagonga Regional Park:

- Corymbia calophylla
- Eucalyptus decipiens
- Eucalyptus gomphocephala

- Eucalyptus marginata
- Eucalyptus petrensis
- Eucalyptus rudis

Eucalyptus todtiana

Eucalypts are iconic Australian plants which vary in size from low shrubs to tall trees. Eucalypts are readily characterised by their distinctive blossoms and their seed capsules known as 'gumnuts'.

There are at least 7 species of eucalypts in Yellagonga Regional Park, the Marri (or Red Gum) (Corymbia calophylla), Redheart Moit (or Limestone Marlock (Eucalyptus decipiens), Tuart (Eucalyptus gomphocephala), Jarrah (or Swan River Mahogany) (Eucalyptus marginata), Koodjat (or Straggly Mallee) (Eucalyptus petrensis), Moitch (or Kulurda, Flooded Gum) (Eucalyptus rudis), and the Coastal Blackbutt (Eucalyptus todtiana).

Eucalypts are extremely important trees for Noongar people and the wood is used for a variety of purposes. For example, the wood of the **Koodjat** and the **Jarrah** is used to make important objects such as **doarks** (sticks for knocking the tops off **Grass Trees** [Xanthorrhoea preissii]), **kitjs** (spears), **wannas** (digging sticks), and in recent times, didgeridoos. Suitable branches from the Jarrah are also used to make spear throwers. The early colonists too, used eucalyptus wood to a great extent for such purposes as construction, fencing and furniture-making.

Eucalyptus leaves also produce eucalyptus oil which is used by Noongar people for medicinal purposes. Gum leaves are rubbed between the hands and then breathed-in to clear the nasal passages. In addition, the leaves of one species, the Moitch, are sometimes covered in small white spots of manna. Manna is the product of a small mite that gathers on the base of the leaves. Noongars lick the sugary manna directly off the leaves or gather the substance into a large, sweet lolly to suck on.



Eucalypts were called 'Gum Trees' by the early colonisers due to the large quantities of gum that exude from their trunks. Noongar people use this gum for a wide variety of medicinal purposes. Gums from the Marri, Tuart and the Jarrah are used as a mild anaesthetic. Large pieces of gum have also been used as fillings for hollow teeth and to treat diarrhoea. Gums can also be ground into powder and used as an ointment on sores or infected areas, or mixed with water as a tonic for upset stomachs.

The bark from eucalypt trees is also very important. Bark from the Marri, Tuart and Jarrah was often used by Noongar people as the roofing for mia-mias (shelters). Jarrah bark is considered the best for this purpose, as it can easily be made waterproof. The high tannin content of **Jarrah** bark also made it useful as a tanning agent and for making dye. In addition, the bark of this tree can be peeled off in one large, curved sheet. Evidence of such sheets of bark being removed can be seen today on 'scarred trees'.

Eucalypts are well-known for their distinctive blossoms. These blossoms are used by Noongar people as a source of honey, either by sucking directly from the flower, or by dipping the flower in water to create a sweet drink. Ngoowak (native bees) enjoy the nectar-rich eucalyptus blossoms also, and Noongars can often find honey in the hollows of eucalyptus branches.

As well as bees, the tall eucalypts, including the Marri, Tuart and the Jarrah, attract birds which nest in the branch hollows. Noongar people can climb the trees to catch the birds or to take the eggs to eat.

■ courtesy M. Fagg, Australian National Botanic Gardens













## Grass Tree Balga

#### Species found in Yellagonga Regional Park:

Xanthorrhoea preissii

The Grass Tree is endemic to south-western Australia and, prior to large-scale land clearing, this plant could be found across the Yellagonga Regional Park area.

In Western Australia, Grass Trees (or Blackboys) (Xanthorrhoea preissii) are also known by their Noongar name, Balga. Other recorded Noongar names for this plant include, Baaluk, Balag, Balka, Barro, Kooryoop, Paaluc, Palga and Yarrlok.

The **Balga** is an extremely important plant for Noongar people and many parts of this plant can be used. The long, thin fronds of the **Grass Tree**, called **mindarie**, can be used to cover the roof of the **mia-mia** (shelter). When it rains, the water runs along the underside of the fronds, keeping the people inside dry. The early colonisers used the mindarie in a similar way for thatch. Noongars also used the mindarie as soft bedding.

Balgas produce a resin which oozes from their trunks (especially on hot days and after burning). This resin can be used as a binding agent after being crushed in a heated stone pot with charcoal and kangaroo droppings. The molten resin produced by this process is used like a cement to bind objects together, such as stone spearheads onto wooden spear shafts.

Balga resin can also be used as a tanning agent. Noongars dissolve lumps of resin in water in a rock hole heated by hot stones. The hides of yonga (kangaroo) and koomal (possum) are scraped and softened and then placed in the rock hole to soak. The skins were then worn as **bookha** (clothes), **wogga** (blankets) or used as a coorda (carry-bag). The resin was prepared by the early colonists in a similar way to make varnish.



**Balga** resin is also highly flammable and Noongar women collect pieces of resin to use as firelighters. The burning resin is also pleasantly fragrant and, when inhaled, can be useful in clearing sinuses. The early colonists noted the highly flammable nature of the **Balga** and a great number of these trees were cut down for firewood.

In addition, **Balgas** are used by Noongar people for food. In times of shortage, the **mindarie** can be pulled out and the white, soft, new leaves eaten. These soft leaves were also eaten by the early colonisers. The centre of the **Balga** is edible too and Noongar people would chop the top off the tree and scoop out the white pulp within. This pulp is used as a medicine for upset stomachs or eaten as food in times of shortage.

In times of drought, **Balgas** were very useful to Aboriginal people in locating water. After several of the **mindarie** were pulled out, a small hollow would remain. Water would then seep into this hollow which could then be drunk.

The flower spear of the **Balga** is also important to Noongar people. The long stem of the flower can be used as a torch, particularly when moving from one camp to the next, and the shaft can be used for sparking fires by friction. The long stem is also used to make spear shafts and in the construction of the **mia-mia**.













### Grevillea Berrung

#### Species found in Yellagonga Regional Park:

- Grevillea crithmifolia
- Grevillea preissii
- Grevillea vestita

The Grevillea Genus is a diverse group of over 350 plants which range from low-lying shrubs to tall trees.

There are at least 3 species of grevillea in Yellagonga Regional Park, the Grevillea crithmifolia (no common name), Grevillea preissii (no common name), and Grevillea vestita (no common name). Low, flowering shrubs, such as grevilleas, are often called Berrung by Noongar people.

The nectar from **Berrung** plants is an important source of honey. Similar to other flowering plants, the nectar from grevilleas can be sucked directly from the flowers or soaked in water to produce a sweet drink. Sometimes the drink is allowed to ferment to produce **Gep**, an intoxicating liquor.













### Hakea Berrung, Pulgur

#### Species found in Yellagonga Regional Park:

- Hakea lissocarpha
- Hakea prostrata
- Hakea trifurcata

Hakeas are shrubs which produce attractive, nectareous flowers, highly favoured by bees and honeyeaters.

In Yellagonga Regional Park there are 3 species of hakeas, the **Honey Bush** (or **Duck and Drake Bush**) (Hakea lissocarpha), the Harsh Hakea (Hakea prostrata) and the Two-Leaf Hakea (Hakea trifurcata). Flowering shrubs, such as hakeas and grevilleas, are often called **Berrung** by Noongar people. The **Harsh Hakea** is also known as the **Pulgur** (or **Doolgur**).

Similar to other flowering plants, hakea flowers are an important source of honey for Noongars. The nectar is either sucked directly from the flowers, or the blossoms are soaked in water to produce a sweet drink. Sometimes the drink is allowed to ferment to produce **Gep**, an intoxicating liquor.

The spiky branches of the **Pulgur** are also used by Noongars in fishing. The branches are broken and used to drive fish into traps. The wood from the branches of the Pulgur can also be used to make messagesticks.

Early colonists, such as George Fletcher Moore observed that the gum from the Hakea tree was eaten by Noongar people and he believed that it formed an important part of the local diet. Hakea gum can be easily stored in cakes, and it is likely that it was transported by Noongar people from place to place. In other parts of Australia, the burnt bark of the Hakea is used in bush medicine. The ash from the bark is rubbed onto the body to relieve skin sores.



# Kangaroo Pavv Kurulbrang

#### Species found in Yellagonga Regional Park:

Anigozanthos humilis
 Anigozanthos manglesii

Kangaroo paws are iconic plants, native to Western Australia. Mangles Kangaroo Paw (Anigozanthos manglesii) is in fact the floral emblem of this State.

In Yellagonga Regional Park there are 2 species of kangaroo paws, Mangles Kangaroo Paw (or Red and Green Kangaroo Paw, Common Green Kangaroo Paw) (Anigozanthos manglesii) and Catspaw (or Common Catspaw, Dwarf Catspaw) (Anigozanthos humilis). Mangles Kangaroo Paw is known by Noongar people as Kurulbrang (or Nollamara, Yonga Marra).

As well as having attractive and unusual flowers, kangaroo paws have tuberous roots which contain significant levels of stored starch. In a similar way to orchids and some lily species, the roots of kangaroo paws are eaten by Noongar people. Prior to large-scale land clearing, it is likely that kangaroo paws were far more abundant in the Yellagonga Regional Park area than they are today. Root tubers formed an important part of the traditional Noongar diet, and it is possible that the roots of kangaroo paws were gathered in large quantities.











#### Species found in Yellagonga Regional Park:

- Burchardia congesta
- Caesia micrantha
- Dianella revoluta

- Dichopogon capillipes
- Sowerbaea laxiflora
- Thysanotus arenarius

- Thysanotus manglesianus
- Thysanotus patersonii
- Thysanotus sparteus

Thysanotus triandrus

'Lily' is a general term used for a variety of flowering plant Genera.

In Yellagonga Regional Park there are at least 10 species of plants that can be considered lilies. These include, Milkmaids (Burchardia congesta), Pale Grass Lily (Caesia micrantha), Blueberry Lily (or Black-Anther Flax Lily, Blue Flax Lily, Native Flax, Spreading Flax Lily) (Dianella revoluta), Chocolate Lily (or Purple Lily) (Dichopogon capillipes), Purple Tassels (or Vanilla Lily) (Sowerbaea laxiflora), Fringed Lily (Thysanotus manglesianus), Twining Fringed Lily (Thysanotus patersonii), **Leafless Fringed Lily** (*Thysanotus sparteus*), **Three-Stammered Fringed Lily** (*Thysanotus triandrus*) and the *Thysanotus arenarius* (no common name). The Aboriginal name *Tjunguri* (or *Tjungoori*) is often applied to the **Twining Fringe Lily**, although this is probably not a Noongar word.

Many lilies are very important to Noongar people due to their nourishing root tubers. Roots were an essential part of the diet of traditional Noongar people and various species of lilies produce an abundance of edible roots. Milkmaids for example, have fleshy white roots around 5 millimetres thick which are a good source of starch. Noongars can also obtain edible roots from the Blueberry Lily, Chocolate Lily, Purple Tassels, Fringed Lily and Twining Fringe Lily. Roots are either eaten raw, steamed in an earth oven, roasted over hot coals or rolled in hot ash. Some roots, such as those of the Twining Fringe Lily are sometimes ground into a paste and made into cakes.



Some lilies also produce an edible fruit or seed. The Blueberry Lily produces a small, blue berry which is sweet to taste. Other lilies, including the Twining Fringe Lily also have edible flowers and stems. This lily can be ground after roasting and the resulting green powder can be eaten with roots.















### Melaleuca Yowarl

#### Species found in Yellagonga Regional Park:

- Melaleuca huegelii
- Melaleuca rhaphiophylla

Melaleuca's are sometimes called 'teatrees' or 'paperbarks' and are often characterised by their flaky, layered bark.

In Yellagonga Regional Park there are at least 2 species of melaleucas, the **Chenille Honeymyrtle** (Melaleuca huegelii) and the Swamp Paperbark (or Freshwater Paperbark) (Melaleuca rhaphiophylla). The Noongar name for the **Swamp Paperbark** is **Yowarl** (or **Bibool Boorn**, **Yiembak**).

Various melaleuca species are extremely important to Aboriginal people. For local Noongars, the **Swamp** Paperbark is probably one of the most significant plant species in the region. The bark of this melaleuca is thin and papery and can be used for a variety of purposes. Long strips of the bark for example, can be used as roofing for mia-mias (shelters) and smaller pieces can be used to carry water or to hold food.

Melaleuca bark is frequently used in Noongar cooking. Meat dishes, such as kweeyar (frogs), djildjit (fish) or yonga (kangaroo), are often wrapped in the bark of the Yowarl before being placed on hot coals or in an earth oven.

The bark of the **Yowarl** can also be used as a torch. After tightly rolling long pieces of bark, one end can be set alight and the high oil content of the bark keeps the torch smouldering.

Melaleuca leaves are also used by Noongar people for medicinal purposes. The leaves are either sucked, chewed or crushed and inhaled to treat head colds and flu. Green leaves from the Swamp Paperbark and the Chenille Honeymyrtle are also used for smoking ceremonies because of the pleasant aroma the



oil in the leaves lets off. A type of tea can also be made by soaking the leaves in boiling water, which is why the early colonists used the term 'Teatree' to refer to this plant.

The flowers of the Swamp Paperbark and the Chenille Honeymyrtle are also important sources of honey. Similar to other flowering plants, the honey is either sucked directly from the flower or the blossoms are soaked in water to create a sweet drink.















### Orchid Kararr

#### Species found in Yellagonga Regional Park:

- Caladenia arenicola
- Caladenia longicauda
- Diuris corymbosa
- Pheladenia deformis
- Pyrorchis nigricans

- Caladenia flava
- Caladenia macrostylis
- Diuris longifolia
- Pterostylis recurva
- Thelymitra crinita

- Caladenia latifolia
- Caladenia marginata
- Elythranthera brunonis
- Pterostylis vittate

The Orchidaceae Family is the largest family of flowering plants, many of which are highly prized for their decorative blooms.

In Yellagonga Regional Park, there are at least 14 species of orchids which have been identified, including the Carousel Spider Orchid (Caladenia arenicola), Cowslip Orchid (or Primrose Orchid) (Caladenia flava), Pink Fairy Orchid (or Pink Fairies) (Caladenia latifolia), Common White Spider Orchid (or Yellow Spider Orchid) (Caladenia longicauda), Leaping Spider Orchid (Caladenia macrostylis), White Fairy Orchid (Caladenia marginata), Donkey Orchid (Diuris corymbosa), Common Donkey Orchid (Diuris longifolia), Purple Enamel Orchid (Elythranthera brunonis), Blue Fairy Orchid (or Blue Beard) (Pheladenia deformis), Jug Orchid (Pterostylis recurva), Banded Greenhood (Pterostylis vittate), Red Beaks (or Undertaker Orchid) (Pyrorchis nigricans) and Blue Lady Orchid (or Queen Orchid, Lily Orchid) (Thelymitra crinita). Most Noongar names for orchids are no longer known, however, the Spider **Orchids** (Caladenia sp.) are known as **Kararr** (or **Kar**).



In Australia, we have become accustomed to the 'Protected' status of orchids. However, in the past, many orchids were considered an important food source. Several early explorers and colonists, including George Grey, James Drummond, Robert Brough Smyth and George Fletcher Moore, noted the use of orchids for food. These early observers identified the root tubers of various orchids, including the **Banded Greenhood** and Red Beaks, as being highly sought after by Noongars. Roots can either be roasted or baked in hot ashes, or pounded into a paste and made into cakes.





### Pea Flower

### Koorla, Koorpa, Koweda, Marno, Pulboorn, Puyenak, Yackal Djarr

#### Species found in Yellagonga Regional Park:

- Bossiaea eriocarpa
- Hovea pungens
- Jacksonia furcellata
- Kennedia prostrata
- Daviesia divaricata
- Hovea trisperma (var. trisperma)
- Jacksonia sericea
- Templetonia retusa
- Hardenbergia comptoniana
- Isotropis cuneifolia
- Jacksonia sternbergiana
- Viminaria juncea

Pea flowers belong to the Fabaceae Family, a large group of flowering plants commonly referred to as the Legume Family or the Pea Family.

There are at least 12 pea flowers in Yellagonga Regional Park and several of the Noongar names for these plants are known. The Native Wisteria (or Wild Sarsaparilla) (Hardenbergia comptoniana) is known as the Koorla (or Koorlo), the Devil's Pins (or Needle-Leaved Hovea) (Hovea pungens) is known as the Puyenak (or Buyenak), the Green Stinkwood (Jacksonia sternbergiana) is known as the Koorpa (or Kapur, Mondil, Mondum), the Scarlet Runner (or Running Postman) (Kennedia prostrata) is known as the Pulboorn (or Pulbarn, Mirdadjet), Cockies Tongues (or Common Templetonia) (Templetonia retusa) is known as Yackal Djarr, Swishbush (or Golden Spray) (Viminaria juncea) is known as Koweda (or Kower, Kweda) and the Marno (Daviesia divaricata) is known commonly by its Noongar name. The Waldjumi (Jacksonia sericea) is also known commonly by its Aboriginal name, but 'Waldjumi' is probably not a Noongar word. Yellagonga Regional Park is also home to the Common Brown Pea (Bossiaea eriocarpa), Common Hovea (Hovea trisperma [var. trisperma]), Grey Stinkwood (Jacksonia furcellata) and Granny Bonnets (or Lamb Poison) (Isotropis cuneifolia).



Pea flowers are varyingly important to Noongar people. The twining, green stems of the **Koorla** for example, can be used as string. However, the purple flowers are never eaten.

The pea flower **Koweda**, is known to be useful to Noongar people. This plant has strong, flexible branches which are used by Noongars when building mia-mias (shelters). The branches can be used in addition to more leafy branches, such as those from **zamia** plants (Macrozamia sp.), as well as bark, from such trees as the **Swamp Paperbark** (Melaleuca rhaphiophylla).

The Pulboorn is not known to have been useful to Noongars, although the early colonists used this plant for making tea. The leaves were rolled into a ball and infused in boiling water for two or three minutes before being drunk.

The early colonists considered some of the other pea flower plants to be dangerous and went to great lengths to remove them from the land. **Granny Bonnets** were also called **Lamb Poison** as hungry sheep would be poisoned after eating them. The **Stinkwood** pea flowers (Jacksonia sp.) were also undesirable to the colonists as they let off an unpleasant odour when burnt.





### Rush, Sedge Waakal Ngarnak

#### Species found in Yellagonga Regional Park:

- Baumea articulata
- Baumea preissii
- Bolboschoenus caldwellii

- Carex appressa
- Carex fascicularis
- Ficinia nodosa

- Juncus pallidus
- Lepidosperma longitudinale
   Mesomelaena pseudostygia
- Schoenoplectus validus

Rushes and sedges are members of the Poales Order of plants and are characterised by their grass-like structure.

There are at least 10 species of rushes and sedges in Yellagonga Regional Park including the **Jointed** Rush (or Jointed Twig Rush, Jointed Twig Sedge) (Baumea articulata), Broad Twig Sedge (Baumea preissii), Marsh Club-Rush (Bolboschoenus caldwellii), Tall Sedge (Carex appressa), Tassel Sedge (or Razor Sedge) (Carex fascicularis), Knotted Club Rush (Ficinia nodosa), Pale Rush (or Giant Rush) (Juncus pallidus), Pithy Sword-Sedge (or Common Sword-Sedge) (Lepidosperma longitudinale), Semaphore Sedge (Mesomelaena pseudostygia) and Lake Club-Rush (or Lake Club-Sedge, River Club-Rush) (Schoenoplectus validus).

The Noongar names for specific species of rushes and sedges are not known. However, many are referred to as Waakal Ngarnak, named after the Waakal (or Wagul, Wagyl, Waugal, Waagal), sometimes called the Rainbow Serpent. Stories from the Noongar Dreaming tell of how pieces of the Waakal's beard fell off as he twisted and wound his way through the country. Where his beard fell off, the rushes and sedges grew. Many rushes and sedges are therefore known as Waakal Ngarnak (Waakal Beard).

Various species of rushes and sedges around Australia are utilised by Aboriginal people for their roots. It is likely that the roots of many species of rushes and sedges in the Joondalup region were eaten by Noongars,



however, only the Marsh Club Rush has been positively identified from early accounts. Explorer, John Edward Eyre, early naturalist, George French Angas, and colonist, Robert Brough Smyth, described the Marsh Club Rush as having root tubers the size of walnuts which were hard and oily. Eyre observed that the roots were prepared first by roasting and were then ground into thin, flat cakes. Numerous early colonial accounts also speak of another rush, known by the Noongar name Yanchet (or Yange, Yanjet, Yandjet). The **Yanchet** was said to be eaten raw by Noongar people in vast quantities, although it is not known exactly which species this refers to.

Many species of rushes and sedges are used by Noongar people to locate water. Noongars know that you can always find fresh water under species such as the Pithy Sword-Sedge, the Semaphore Sedge and the Knotted Club-Rush.

The leaves of rushes and sedges are also used in weaving. The leaves are woven to create nets which are used as seines to catch djildjit (fish) and yakan (turtle). Rushes and sedges can also be woven to create baskets and mats and the leaves of many species are used as string.

In addition, several species of rushes and sedges with cylindrical leaves, such as the **Jointed Rush**, are sometimes hollowed out. The resulting 'pipe' can be used as a snorkel when hunting yerderap (ducks) and other water fowl.





## Solanum

#### Species found in Yellagonga Regional Park:

Solanum symonii

The term Solanum refers to a large, diverse Genus of plants, of which the common tomato, potato and eggplant are all members.

Solanums are common in Australia, particularly in desert areas where they are known as **Bush Tomatoes**, Bush Raisins and Kangaroo Apples. There is at least 1 species of Solanum that grows in the Yellagonga Regional Park area, the **Solanum symonii** (no common name).

There are many species of Solanum which are edible, and in the central desert region, many Solanum species are staple foods. The **Solanum symonii** grows a small, edible berry that turns purple-black when ripe. This berry however, is somewhat bitter to eat and it is unlikely that traditional Noongar people prized it as a bush food.

It should be noted that there are *numerous* poisonous species of *Solanum* in Australia and it would be unwise to taste any such fruits without specialised, local knowledge.







# Spearvvood Kitja Boorn

#### Species found in Yellagonga Regional Park:

Kunzea ericifolia

Kunzea glabrescens

Spearwood plants are members of the Kunzea Genus. There are at least 2 spearwoods in Yellagonga Regional Park, the Kunzea glabrescens (usually known as Spearwood) and the Native Tea (or Spearwood, Yellow Kunzea ericifolia). Noongar people called the Native Tea plant Kitja Boorn (or Poorndil, Condil).

As its English name suggests, spearwood plants are used by Noongar people in spear-making. Spears produced from the Kitja Boorn can be used to hunt animals in small swamps and water holes.

The early colonists found spearwoods useful plants also. The Kitja Boorn was used for making tea, hence the name **Native Tea**. The tea produced by this spearwood was considered not only pleasant, but was also used as a tonic. In more recent times, the stems of the Kitja Boorn have been used in market gardens as bean-sticks as well as in the construction of crayfish pots.

■ courtesy M. Fagg, Australian National Botanic Gardens









## Wattle

### Coojong, Panjang, Wilyawa

#### Species found in Yellagonga Regional Park:

- Acacia cochlearis
- Acacia cyclops
- Acacia huegelii

- Acacia lasiocarpa
- Acacia pulchella
- Acacia saligna

Acacia willdenowiana

Wattles belong to the Acacia Genus and are well-known throughout Australia for their bright yellow flowers. The **Golden Wattle** (*Acacia pycnantha*), is in fact Australia's national flower.

There are at least 7 species of wattle in Yellagonga Regional Park and the Noongar names of 3 of these species are well known. The Red-Eyed Wattle (Acacia cyclops) is known as the Wilyawa (or Woolya Wah), the Orange Wattle (or Black Wattle) (Acacia saligna) is known as the Coojong (or Cujong, Kalyung, Kileyung, Kudjong), and the Panjang (or Pajang) (Acacia lasiocarpa) is known commonly by its Noongar name. Other species present in Yellagonga Regional Park include the Rigid Wattle (Acacia cochlearis), Prickly Moses (Acacia pulchella), Grass Wattle (Acacia willdenowiana) and the Acacia huegelii (no common name).

For Noongar people, wattles are extremely important plants. Wattle seeds for example, are a very good source of fats, protein and carbohydrates. The Wilyawa and the Coojong both have edible seeds. The seeds of the Coojong can be eaten raw, and the seeds of the Wilyawa can be ground into a flour and baked into damper.

For local Noongars, the Wilyawa is probably the most important wattle in Yellagonga Regional Park. As well as using the seeds for making damper, the green seed pods are used for a variety of purposes. For instance, a pod can be crushed in the hands to release a sticky juice which, when a little water is added, can be used as a creamy sunscreen and an insect repellent. This cream is also used to treat eczema. If a



little more water is added, the pods can be rubbed between the hands and used as a soap or cleanser. Elsewhere in Australia, similar wattles have been called 'Soap Wattles'.

The Wilyawa is also an important source of gum. The gum that exudes from the trunk is edible and can be chewed like chewing gum. Other wattle species also produce edible gums that can be sucked. These gums can act as a purgative and are used by Noongar people to alleviate constipation. Wattle gum can also be soaked in water to create a glue.

Many wattle species, including the Wilyawa, are home to grubs, sometimes known as Bardi Grubs or Witchetty Grubs. When these grubs are found in rotting wattle trees, they are roasted over hot coals or in hot ashes before eating. Many early explorers, such as George Grey and Edward John Eyre, noted that grubs were harvested from both wattle trees and Grass Trees (Xanthorrhoea preissii). Those from wattle trees are larger but not as plentiful as those from **Grass Trees**.

The wood of many species of wattle around Australia is also extensively used by Aborigines. Noongar people use Acacia wood for making spear heads, kitjs (spears), wannas (digging sticks) and shields. Wattle tree-trunks can also be used as poles for constructing mia-mias (shelters), as they grow straight and are light to carry. The early colonists used Acacia wood for such things as gunstocks and fence posts.

Wattle bark is also important for tying items together. The bark can be stripped off the tree easily and then oiled with kangaroo fat or goanna oil to make it pliable. Many species of wattle are also known to have highly astringent bark. According to the author Jennifer Hagger, the early colonists used Acacia bark to make decoctions or infusions to treat ailments such as diarrhoea and eye conditions.













### Zamia Bayu, Djiridji

#### Species found in Yellagonga Regional Park:

- Macrozamia fraseri
- Macrozamia riedliei

Zamias belong to the Cycadales Family of plants, which refer to a number of ancient species dating back to the Triassic period. Now representing only a minor part of the Plant Kingdom, Cycadales were once extremely common.

Two species of zamia exists in Yellagonga Regional Park, the **Sandplain Zamia** (Macrozamia fraseri) and the Zamia Palm (Macrozamia riedliei). Noongar people called these zamias Djiridji (or Dyergee, Girijee, Jeerajee). The bright, orange seeds of the Djiridji were called bayu (or booyoo, boya, byyu).

The **Diiridji** produce large seed pods which look somewhat like green pineapples. These pods are home to a number of orange bayu which contain significant levels of toxins. Some Aboriginal people consider bayu a delicacy and different Aboriginal groups around Australia undertake various preparatory processes before eating them. In south-western Australia the bayu are sometimes collected in a reed bag which is then soaked in running water for a period of time to leach out the toxins. The seeds are then buried underground, often for 6 months or more. After this time, Noongar people peel the seeds and eat only the orange skin. Elsewhere in Australia, it is the seeds themselves that are eaten, and these are often crushed into a porridge-like meal and then formed into cakes and roasted in ashes. Some Aboriginal groups soak the seeds in water for an extended period of time and many do not bury them. Some early botanists and colonists, including James Drummond and George Fletcher Moore believed the bayu to be a culturally significant food for traditional Noongar people.

Many early explorers and colonists ate the attractive orange zamia seeds without knowledge of their toxicity. Disastrous encounters with the seeds of the Djiridji were recorded as early as 1697 by the crew of the Dutch voyage of Willem de Vlamingh. The crew were exploring the land around the Swan River when they



ate several raw **bayu** which they described as tasting like Dutch broad beans. According to the ship's log, within three hours, they had begun to "vomit so violently that there was hardly any distinction between death and us."

The **Djiridji** also produces a cotton-like substance around the base of the plant. This native cotton is very soft and absorbent and was used by traditional Noongar women for feminine hygiene purposes. The native cotton can also be used in the **coolamon** (carrying vessel) as a soft lining for babies to sleep on as they are carried from place to place. Many early colonists also utilised this native cotton as tinder for lighting fires.

The fronds of the **Djiridji** are also useful to Noongar people. The large palm-like leaves are used for shade and occasionally in the roofing of the **mia-mia** (shelter). The long zamia leaves can also be removed from their stem and used as a strong string to tie objects together.

Many early colonists took advantage of the high starch content of zamias. Early botanist James Drummond recorded that some of the colonisers prepared quantities of starch by grating down peeled seeds and pouring water over the mass. The vegetable matter would then separate and leave a starch at the bottom, similar to arrowroot.













## Appendix 1 Species List

The following Species List has been assembled from several different resources and, as such, contains the Noongar names for plants that have been documented by various people in different parts of Noongar Country. Some names, particularly those from the Great Southern and Wheatbelt regions of Western Australia may not have been used in Mooro Country. All Noongar names have therefore been referenced, so that readers may further explore their derivation.

(most frequently used name(s) is **bolded**)

Scientific Name	Common Name(s)	Noongar Name(s)
Acacia cochlearis	Rigid Wattle	
Acacia cyclops	Coastal Wattle Red-Eyed Wattle Western Coastal Wattle	Woolya Wah <sup>1</sup> Wilyawa <sup>1</sup>
Acacia huegelii		
Acacia lasiocarpa	Panjang	Pajang <sup>2</sup> <b>Panjang</b> <sup>3</sup>
Acacia pulchella	Prickly Moses Western Prickly Moses	
Acacia saligna	Black Wattle Golden-Wreath Wattle Orange Wattle	Coojong <sup>3, 4</sup> Cujong <sup>2, 3</sup> <b>Kudjong</b> <sup>3</sup> Kileyung <sup>5</sup> Kalyung <sup>1</sup>
Acacia willdenowiana	Grass Wattle	
Alexgeorgea nitens		
Allocasuarina fraseriana	Common Sheoak Fraser's Sheoak <b>Sheoak</b>	Condil <sup>2, 3, 4</sup> <b>Kondil</b> <sup>3</sup> Kwerl <sup>1</sup> <b>Kwoorl</b> <sup>1</sup>
Allocasuarina humilis	Dwarf Sheoak	
Allocasuarina lehmanniana	Dune Sheoak	
Anigozanthos humilis	Catspaw Common Catspaw Dwarf Catspaw Small Orange Kangaroo Paw	
Anigozanthos manglesii	Common Green Kangaroo Paw <b>Mangles Kangaroo Paw</b> Red and Green Kangaroo Paw Red-Stemmed Kangaroo Paw	Koroylbardang <sup>6</sup> Krulbrang <sup>2, 3</sup> Kuroolberny <sup>1</sup> <b>Kurulbrang</b> <sup>3</sup> <b>Nollamara</b> <sup>1</sup> <b>Yonga Marra</b> <sup>3</sup>
Astartea fascicularis		

Banksia attenuata	Candle Banksia Candlestick Banksia Coast Banksia Slender Banksia	Bealwra <sup>5</sup> Beara <sup>2, 3</sup> Biara <sup>2, 3, 4, 6</sup> Birytch (flower) <sup>6</sup> Biytch (flower) <sup>6</sup> Peera <sup>2, 3</sup> <b>Piara</b> <sup>2, 3</sup> Piras <sup>6</sup>
Banksia grandis	Bull Banksia Giant Banksia Great-Flowered Banksia	Beera <sup>2, 3, 7, 8</sup> Boolgalla <sup>2, 3, 7</sup> Boorarup <sup>2, 3, 7</sup> Bulgalla <sup>6</sup> Mangaat <sup>6</sup> Mangaitch <sup>9</sup> Mangatj <sup>1</sup> Mangghoyte <sup>5</sup> Mangite <sup>3, 4, 7, 10</sup> Mangyt (flower) <sup>6, 10</sup> Manjoyte <sup>11</sup> Metjo (flower) <sup>6</sup> Metjo Kundyle (seeds) <sup>6</sup> Moncat <sup>10, 12</sup> Mungart <sup>1</sup> Mungat <sup>10, 12, 13</sup> Munghite <sup>10</sup> Mungitch <sup>7</sup> Mungite <sup>10, 14</sup> Mungit <sup>10, 14</sup> Mungyte <sup>10, 16</sup> Munyaret <sup>1</sup> Poolgarla <sup>7</sup> Pulgarla <sup>3</sup>
Banksia ilicifolia	Holly Banksia Holly-Leaved Banksia	9
Banksia littoralis	River Banksia Seaside Banksia <b>Swamp Banksia</b> Swamp Oak Western Swamp Banksia	Boongura <sup>2, 3, 8</sup> Gwangia <sup>2, 3</sup> <b>Pungura</b> <sup>3</sup>
Banksia menziesii	<b>Firewood Banksia</b> Menzies Banksia	
Banksia prionotes	Acorn Banksia Orange Banksia Saw-Tooth Banksia	
Banksia sessilis	Parrot Bush	Pulgart <sup>1</sup>
Baumea articulata	Jointed Rush Jointed Twig Rush Jointed Twig Sedge	Waakal Ngarnak (rush/sedge) 1
Baumea preissii	Broad Twig Sedge	Waakal Ngarnak (rush/sedge) 1
Bolboschoenus caldwellii	Marsh Club-Rush	Belillah <sup>17</sup> Waakal Ngarnak (rush/sedge) <sup>1</sup>
Bossiaea eriocarpa	Common Brown Pea	
Burchardia congesta	Milkmaids	
Caesia micrantha	Pale Grass-Lily	
Caladenia arenicola	Carousel Spider Orchid	Kar (Caladenia sp.) <sup>1</sup> Kararr (Caladenia sp.) <sup>1</sup>

Caladenia flava	Cowslip Orchid Primrose Orchid	Kar (Caladenia sp.) <sup>1</sup> Kararr (Caladenia sp.) <sup>1</sup>
Caladenia latifolia	Pink Fairies Pink Fairy Orchid	Kar (Caladenia sp.) <sup>1</sup> Kararr (Caladenia sp.) <sup>1</sup>
Caladenia longicauda	Common White Spider Orchid Yellow Spider Orchid	Kar (Caladenia sp.) <sup>1</sup> Kararr (Caladenia sp.) <sup>1</sup>
Caladenia macrostylis	Leaping Spider Orchid	Kar (Caladenia sp.) <sup>1</sup> Kararr (Caladenia sp.) <sup>1</sup>
Caladenia marginata	White Fairy Orchid	Kar (Caladenia sp.) <sup>1</sup> Kararr (Caladenia sp.) <sup>1</sup>
Calothamnus quadrifidus	Common One-Sided Bottlebrush One-Sided Bottlebrush	Kwowdjard <sup>3</sup> Queitjat <sup>2, 3</sup>
Calothamnus sanguineus	Silky-Leaved Blood Flower	
Carex appressa	Tall Sedge	Waakal Ngarnak (rush/sedge) 1
Carex fascicularis	Razor Sedge Tassel Sedge	Waakal Ngarnak (rush/sedge) 1
Clematis linearifolia	Slender Clematis	
Clematis pubescens	Common Clematis Common White Clematis White Clematis	
Conostephium pendulum	Drooping Cone Flower Pearl Flower Pink-Tipped Pearl	
Conostylis candicans (ssp. calcicola)	<b>Grey Cottonhead</b> White Conostylis	
Corymbia calophylla	Marri Red Gum	Cardau <sup>11</sup> Gardan <sup>2, 3, 6</sup> Grydan <sup>2, 3, 8</sup> Kardan <sup>2, 3, 5</sup> Kurrden <sup>2, 3</sup> Maree <sup>2, 3</sup> Mari <sup>3</sup> Marri <sup>2, 3</sup> Marril <sup>2, 3</sup> Mundup <sup>2, 3</sup> Nalla (resin) <sup>6</sup> Nandup <sup>2, 3</sup> Ngoombit (flower) <sup>8</sup> Ngora <sup>2, 3</sup> Ngumbat (flower) <sup>2, 3</sup> Ngumbit (flower) <sup>6, 10</sup> Numbit (flower) <sup>10, 18</sup> Numbrid (flower) <sup>6, 10</sup> Nundup <sup>3</sup>
Corynotheca micrantha	Sand Lily	
Crassula colorata	Dense Crassula  Dense Stonecrop	
Daviesia divaricata	Marno	Mano <sup>2, 3</sup> <b>Marno</b> <sup>3</sup>
Desmocladus flexuosus		

Dianella revoluta	Black-Anther Flax Lily Blue Flax Lily Blueberry Lily Flax Lily Native Flax Spreading Flax Lily	
Dichopogon capillipes	Chocolate Lily Purple Lily	
Diuris corymbosa	Donkey Orchid	
Diuris longifolia	Common Donkey Orchid Donkey Orchid	
Drosera erythrorhiza	Red Ink Sundew	
Drosera macrantha	Bridal Rainbow Bridal Rainbow Sundew Climbing Drosera Rainbow	
Elythranthera brunonis	Purple Enamel Orchid	
Eremaea pauciflora		
Eryngium pinnatifidum	Blue Devils	
Eucalyptus gomphocephala	<b>Tuart</b> White Gum	Duart <sup>3</sup> Dubta (seeds) <sup>6</sup> Mooarn <sup>3</sup> Moorun <sup>2, 3</sup> Mouarn <sup>2</sup> Tooart <sup>2, 3</sup> <b>Tuart</b> <sup>2, 3, 4</sup>
Eucalyptus decipiens	Redheart Moit Limestone Marlock	
Eucalyptus marginata	Jarrah Swan River Mahogany	Budto (bark) <sup>6</sup> Cherring <sup>2, 3</sup> Chiaragl <sup>2, 3</sup> Djara <sup>3</sup> Djarrail <sup>2, 3</sup> Djarryl <sup>2, 3, 6</sup> Djerral <sup>2, 8</sup> Djerral <sup>2, 8</sup> Gharrahel <sup>11</sup> Jarrah <sup>2, 3, 4, 19</sup> Jarrail <sup>5</sup> Jarral <sup>2, 3</sup> Jarraly <sup>2, 3</sup> Jarril <sup>2, 3</sup> Jeril <sup>2, 3</sup> Jeril <sup>2, 3</sup> Jerrail <sup>2, 3</sup>
Eucalyptus petrensis	Straggly Mallee	Koodjat <sup>1</sup>
Eucalyptus rudis	Blue Gum Desert gum Flooded Gum River Gum Swamp Gum Western Australian Flooded Gum	Colaille <sup>2, 3</sup> Gooloorda <sup>2, 3, 8</sup> Gooloorto <sup>2, 3</sup> Gulurto <sup>2, 3, 6</sup> Koolert <sup>2, 3</sup> <b>Kulurda</b> <sup>2</sup> <b>Moitch</b> <sup>1, 2, 3, 4, 19</sup> Moja <sup>2, 3</sup>

Eucalyptus todtiana	Blackbutt Coastal Blackbutt Pricklybark	
Ficinia nodosa	Knobby Club Rush Knotted Club Rush	Waakal Ngarnak (rush/sedge) 1
Gompholobium tomentosum	Hairy Yellow Pea	
Grevillea crithmifolia		Berrung (low, flowering shrub) 1
Grevillea preissii		
Grevillea vestita		
Hakea lissocarpha	Duck and Drake Bush Honey Bush	Berrung (low, flowering shrub) 1
Hakea prostrata	Harsh Hakea	Berrung (low, flowering shrub) <sup>1</sup> Doolgur <sup>1</sup> Pulgur <sup>1</sup>
Hakea trifurcata	Two-Leaf Hakea	Berrung (low, flowering shrub) 1
Hardenbergia comptoniana	Native Wisteria Wild Sarsaparilla Wild Wisteria	Koorla <sup>1</sup> Koorlo <sup>1</sup> Kurrolo <sup>6</sup>
Hemiandra linearis	Speckled Snakebush	
Hibbertia aurea		
Hibbertia hypericoides	Yellow Buttercups	
Hibbertia racemosa	Stalked Guinea Flower	
Hovea pungens	Devil's Pins Needle-Leaved Hovea	Buyenak <sup>2, 3, 6</sup> <b>Puyenak</b> <sup>3</sup>
Hovea trisperma (var. trisperma)	Common Hovea	
Hybanthus calycinus	Native Violet Wild Violet	
Hypocalymma robustum	Bush Myrtle Monkey-Blossom Pink Myrtlee Pink-All-The-Way-Up Swan River Myrtle Wild Peach	
Isotropis cuneifolia	Common Lamb Poison Granny Bonnets Lamb Poison	
Jacksonia furcellata	Grey Stinkwood	
Jacksonia sericea	Waldjumi	Waldjumei <sup>3</sup> <b>Waldjumi</b> <sup>3</sup> Waljumei <sup>2</sup>
Jacksonia sternbergiana	Green Stinkwood	Cabboor <sup>5</sup> Kabbur <sup>2, 3, 4</sup> Kapbur <sup>2, 3, 6</sup> Kapoor <sup>8</sup> <b>Kapur</b> <sup>3</sup> Koorpa <sup>2, 3</sup> <b>Koorpa</b> <sup>1</sup> Mondil <sup>2, 3</sup> Mondurn <sup>2, 3</sup>
Juncus pallidus	Giant Rush Pale Rush	Waakal Ngarnak (rush/sedge) 1

Kennedia prostrata	Red Runner	Mirdadjet <sup>1</sup>
	Running Postman	Pulbarn <sup>6</sup>
	Scarlet Coral Pea	Pulboorn <sup>1</sup>
	Scarlet Runner	
Kunzea ericifolia	Native Tea	Condil <sup>2, 3</sup>
	Spearwood	Kitja Boorn <sup>1</sup>
	Yellow Kunzea	Pondil <sup>2, 3, 4</sup>
	Tellow Rulizea	Poorndil <sup>1</sup>
		Poortidii
Kunzea glabrescens	Spearwood	
Lagenophora huegelii	Coarse Lagenophora	
Lechenaultia linarioides	Coastal Leschenaultia Yellow Leschenaultia	
Lepidosperma longitudinale	Common Sword-Sedge  Pithy Sword-Sedge	Waakal Ngarnak (rush/sedge) 1
Leucopogon propinquus		
Lobelia tenuior	Slender Lobelia	Bindji-Windjil 3, 20
Macrozamia fraseri	Sandplain Zamia	Baian (seeds) <sup>2, 3</sup>
		Baio (seeds) <sup>5, 10</sup>
		Bayio (seeds) 10, 11
		Bayou (seeds) <sup>21</sup>
		Bayu (seeds) 1
		Biu (seeds) 11
		Booyoo (seeds) <sup>25</sup>
		Boya (seeds) <sup>23</sup>
		Boyern (seeds) 2,3
		Boyo (seeds) 10
		Byyu (seeds) 6, 10, 16, 24
		Djiridji <sup>1, 3</sup>
		Djiriji <sup>2, 3, 6, 21</sup>
		Djir-jy ⁵
		Dyergee <sup>2,3,8</sup>
		Dyundo (seed kernel) <sup>6</sup>
		Gargoin (seed stone) 6
		Girijee <sup>2, 3</sup>
		Jeerajee <sup>2, 3</sup>
		Jeeriji <sup>2, 3</sup>
		Jeerja <sup>2, 3</sup>
		Jirjy <sup>5</sup>
		Koondagoor <sup>2, 3</sup>
		Kundagur <sup>2, 3, 6</sup>
		Kwinin (seeds) <sup>6</sup>
		Quinine (seeds) <sup>2, 3</sup>
		Quinning (seeds) 2,3
		Wida (seed kernal) <sup>6</sup>
Magrazamia riadiai	Zamia Palm	Djiridji 1,3
Macrozamia riedlei	∠aiilia Fallii	Dyergee <sup>2,3,8</sup>
		Girijee <sup>2,3</sup>
		Jeerajee <sup>2,3</sup>
Melaleuca huegelii	Chenille Honeymyrtle	
Melaleuca rhaphiophylla	Freshwater Paperbark	Bibool Boorn <sup>1</sup>
	Swamp Paperbark	Yeymbac <sup>5</sup>
		Yiembak <sup>11</sup>
		Yowarl <sup>1</sup>
Mesomelaena pseudostygia	Semaphore Sedge	Waakal Ngarnak (rush/sedge) 1
Myoporum caprarioides	Slender Myoporum	
Olearia axillaris	Coastal Daisybush	
Cicaria axinario	Joustal Balaysusii	

Opercularia hispidula	Dogweed Hispid Dogweed Hispid Stinkweed	<b>Djural</b> (grass-like plant) <sup>1</sup> <b>Djooral</b> (grass-like plant) <sup>1</sup>
Opercularia vaginate	Dog Weed	
Orthrosanthus laxus	Morning Iris	
Patersonia occidentalis	Purple Flag Western Patersonia	Comma <sup>1</sup> Koma <sup>3</sup> <b>Komma</b> <sup>1, 2, 3, 6</sup>
Petrophile linearis	Drumsticks Narrow-Leaved Cone Bush Pixie Mops	
Petrophile macrostachya		
Pheladenia deformis	Blue Beard Blue Fairy Orchid	
Philotheca spicata	Pepper and Salt	
Phyllanthus calycinus	False Boronia	
Pterostylis recurva	Jug Orchid	
Pterostylis vittate	Banded Greenhood Banded Greenhood Orchid Greenhood	
Ptilotus drummondii	Narrow-Leaf Mulla Mulla	Moorla-Moorla <sup>1</sup> <b>Mulla-Mulla</b> <sup>3</sup>
Ptilotus polystachyus	Bottle Washer Green Mulla-Mulla Green Spiked Mulla-Mulla Long Tails Prince of Wales Feather	Moorla-Moorla <sup>1</sup> <b>Mulla-Mulla</b> <sup>3</sup>
Pyrorchis nigricans	Red Beaks Undertaker Orchid	
Regelia ciliata	Linnaea	
Rhagodia baccata (ssp. Dioica)	Berry Saltbush Sea Berry Saltbush	
Ricinocarpos glaucus	Wedding Bush	
Schoenoplectus validus	Lake Club-Rush Lake Club-Sedge River Club-Rush	Waakal Ngarnak (rush/sedge) 1
Senecio pinnatifolius (var. maritimus)	Coastal Groundsel Variable Groundsel	Yoont Djet 1
Solanum symonii		
Sowerbaea laxiflora	<b>Purple Tassels</b> Vanilla Lily	
Spyridium globulosum	Basket Bush	
Stylidium calcaratum	<b>Book Triggerplant</b> Tripperplant	
Stylidium schoenoides	Cow Kicks Cow Licks	
Stylidium striatum	Fan-Leaved Triggerplant	
Templetonia retusa	Cockies Tongues Common Templetonia Red-Flowered Templetonia Templetonia	Yackal Djarr <sup>1</sup>
Thelymitra crinita	Blue Lady Orchid Lily Orchid Queen Orchid	



## **Endnotes:**

<sup>1</sup> N Collard, personal communication, 26 June 2009

<sup>2</sup> Abbott 1983

<sup>3</sup> Bennett 1991

<sup>4</sup> Powell 1990

<sup>5</sup> Stokes 1846 <sup>6</sup> Moore 1884

Daw et al 1997

<sup>8</sup> Lyon 1833

<sup>9</sup> Roth 1903

<sup>10</sup> Meagher 1974

<sup>11</sup> Fountain 1907

<sup>12</sup> Nind 1831

13 Collie 1834

<sup>14</sup> Armstrong 1836

15 Bird & Beeck 1899

<sup>16</sup> Grey 1841b

<sup>17</sup> Eyre 1845

<sup>18</sup> Drummond 1842-1843j

<sup>19</sup> Chippendale 1973

<sup>20</sup> Reid 1977

<sup>21</sup> CoJ nd

<sup>22</sup> Hammond 2005

<sup>23</sup> Drummond 1842-1843e

<sup>24</sup> Smyth 1878



## Appendix 2 Noongar Seasons

Across Australia, Aboriginal hunting and food-gathering practices were dictated by the cycle of the seasons (Berndt & Berndt 1999:11). The way in which the year was divided depended upon the region concerned. In south-western Australia the year was generally separated into 6 seasons: Birak, Bunuru, Djerin, Makaru, Djilba and Kambarang.

December to January	Birak	At this time of year the weather was dry and hot. During Birak, scrubland was burnt to force animals into the open for easier hunting and to encourage new plant growth. Banksias were in flower during Birak and the blossoms were gathered in Mooro Country for their honey.
February to March	Bunuru	Bunuru was the hottest time of the year and during Bunuru, people gathered around the lakes, including those in the Yellagonga Regional Park area. At this time of year, food was plentiful with frogs and reptiles in abundance. Zamia seeds were collected at this time and banksia and wattle flowers were gathered for their honey.
April to May	Djerin	As Djerin approached, the weather got cooler and people travelled down the river. Scrubland was burnt to ensure food would be plentiful for the next year. Shelters were built in Djerin and skin cloaks were sewn. At this time of year, root vegetables were plentiful and various root tubers and bulbs were collected.
June to July	Makaru	During these wetter months, the people of the Swan River Plain moved up towards the shelter of the hills where they would be protected from the south-west winds. At this time of year, rains replenished the inland water resources and large animals, such as kangaroos, emus and possums were hunted for food. Smouldering banksia cones were kept under cloaks to keep warm.
August to September	Djilba	During Djilba, the days and nights were clear and cold. During this time, root tubers were an important food source. In particular, native yams near the Swan River were dug in vast quantities. Large animals, such as kangaroos, emus and possums continued to be hunted. Plants, such as milkmaids, cottonhead, myrtle and spearwood would begin to flower.
October to November	Kambarang	As the weather become warmer in Kambarang, people camped around the lakes, including those in the Yellagonga Regional Park area. Wetlands foods, including frogs and reptiles were hunted. Birds, such as ducks, swans and wild turkeys were also plentiful. Sweet gums and resins would exude from the bark of Eucalypts.



## Appendix 3 Reference List

Abbott, I 1983, Aboriginal Names for Plant Species in South-Western Australia, Technical Paper No 5, Forests Department of Western Australia, Perth.

Ah Sam, M 2006, Mitakoodi Bush Tucker: Edible and Medicinal Plants of the Northwest Highlands and Gulf Plains of Queensland, Black Ink Press, Mt Isa.

Angas, G F 1847, Savage Life and Scenes in Australia and New Zealand: Being an Artist's Impressions of Countries and People at the Antipodes, Vol 1, Smith Elder & Co, London, retrieved from American Libraries Internet Archive.

Armstrong, F 1836, 'Manners and Habits of the Aborigines of Western Australia, From Information Collected by Mr F Armstrong, Interpreter, 1836', in Green, N J (ed) 1979, Noongar, the People: Aboriginal Customs in the Southwest of Australia, Creative Research Publishers, Perth, pp 181-206.

Arnold, H 1910, 'A Page from Our Church's History: Methodism in Western Australia 80 Years Ago', The Western Methodist, Vol 3, No 2, p 5, (Battye Library).

Austen, T 1998, A Cry in the Wind: Conflict in Western Australia, 1829-1929, Darlington Publishing Group, Darlington.

Backhouse J 1843, A Narrative of a Visit to the Australian Colonies, Hamilton Adams and Co, London, retrieved from Google Books database.

Bates 1914, 'A Few Notes on Some South-Western Australian Dialects', The Journal of the Royal Anthropological Institute of Great Britain and Ireland, Vol 44, No 1, pp 65-82, retrieved from JSTOR database.

Bates, D M 1992 (1907), Aboriginal Perth: Bibbulmun Biographies and Legends, Hesperian Books, Perth.

Beck, W, Fullager, R & White N 1988, 'Archaeology from Ethnography: The Aboriginal Use of Cycad as an Example', in Meehan, B & Jones, R (eds), Archaeology with Ethnography: An Australian Perspective, Australian National University Press, Canberra, pp 137-147.

Beckle, H 1979, A Seasonal Biogeography of Lake Joondalup, Honours Thesis, University of Western Australia, Perth, (Joondalup Library).

Bennett, E M 1991, Common and Aboriginal Names of Western Australian Plant Species, Wildflower Association of Western Australia, Perth.

Berndt, R M & Berndt, C H 1999, The World of the First Australians, Aboriginal Studies Press, Canberra.

Berryman I 2002, Swan River Letters, Swan River Press, Perth.

Bird, C & Beeck C 1988, 'Traditional Plant Foods in the Southwest of Western Australia: The Evidence from Salvage Ethnography', in Meehan, B & Jones, R (eds), Archaeology with Ethnography: An Australian Perspective, Australian National University Press, Canberra, pp 113-122.

Bougher, N L & Hart, H 2004, Yellagonga Regional Park Fungi Report 2004, Perth Urban Bushland Fungi, Perth, retrieved from www.fungiperth.org.au/Reports-all/Fungi-Surveys.html.

Bougher, N L & Syme K 1998, Fungi of Southern Australia, University of Western Australia Press, Perth.

Bougher, N L 2008, Perth Urban Bushland Fungi Field Book, Perth Urban Bushland Fungi, Perth, retrieved from www.fungiperth.org.au/Fieldbook-all/Perth-Fungi-Field-Book.

Bougher, N L, Hart, H & de Bueger, S 2006, Yellagonga Regional Park Fungi Report 2006, Perth Urban Bushland Fungi, Perth, retrieved from www.fungiperth.org.au/Reports-all/Fungi-Surveys.

Brittain, R K 1987, "Perry Paddock" Historical Village City of Wanneroo: Report for the Historical Sites Advisory Committee, Kidd and Povey Architects, City of Wanneroo, Perth.

Brittain, R K 1990. Yellagonga Regional Park, City of Wanneroo, Ethnography Position Paper, report prepared for the Department of Planning and Urban Development, Kidd and Povey Architects, City of Wanneroo, Perth.

Brittain, R K 1991, East Wanneroo Historical Sites Study, report prepared for the City of Wanneroo Historical Sites Advisory Committee, City of Wanneroo, Perth.

Brouk, B 1975, Plants Consumed by Man, Academic Press, London.

Butler, J 1834, 'Mr John Butler's Report of an Excursion to Search for Cattle 35 Miles North of Perth in March 1834', in Shoobert, J (ed) 2005, Western Australian Exploration, December 1826-December 1835: The Letter, Reports and Journals of Exploration and Discovery in Western Australia, Vol 1, Hesperian Press, Perth.

Cane, S 1989, 'Australian Aboriginal Seed Grinding and its Archaeological Record: A Case from the Western Desert', in Harris, D R & Hillman, G C (eds), Foraging and Farming: The Evolution of Plant Exploitation, Unwin Hyman, Sydney, pp 99-119.

Carter, B 2006, Nyungah Land: Records of Invasion and Theft of Aboriginal Land on the Swan River, 1829-1850, Swan Valley Nyungah Community, Perth.

Chambers, A 1991, The Pioneers: A Story of Wanneroo 1834-1985, City of Wanneroo, Perth.

Cherikoff, V 1993, The Bushfood Handbook: How to Gather, Grow, Process & Cook Australian Wild Foods, Bush Tucker Supply Australia Pty Ltd, Sydney.

Chippendale, G M 1973, Eucalypts of the Western Australian Goldfields (and the Adjacent Wheatbelt), Australian Government Publishing Services, Canberra.

City of Wanneroo (CoW) 1995, 'Traditional Aboriginal Place Names and Their Meanings', in Kauler, L B 1998, Cultural Significance of Aboriginal Sites in the Wanneroo Area, report prepared for Harry Nannup of Gnangara Aboriginal Community College, Heritage Council of WA, Perth, pp 115-116.

City of Joondalup (CoJ) nd, Joondalup Mooro Boodjar: Indigenous Culture Within Mooro Country, Perth, retrieved from www.joondalup.wa.gov.au/Libraries/Documents/Joondalup\_Mooro\_Boodjar\_Brochure.pdf.

City of Wanneroo (CoW) 2003, Lake Joondalup Nature Reserve, unpublished, Perth.

City of Wanneroo (CoW) 2006, Municipal Heritage Inventory, Perth.

City of Wanneroo (CoW) nd, History of Wanneroo: A Brief History, Perth, retrieved from www.wanneroo.wa.gov.au/3/93/1/ history\_of\_wanneroo.pm.

Clarke, P 2007, Aboriginal People and their Plants, Rosenburg Publishing, Sydney.

Colbung, K 1992, Yagan: The Swan River 'Settlement', Australia Council for the Arts, Sydney.

Collie, A 1834, 'Anecdotes and Remarks Relative to the Aborigines at King George's Sound - From an Original Manuscript by a Resident at King George's Sound', in Green, N J (ed) 1979, Noongar, the People: Aboriginal Customs in the Southwest of Australia, Creative Research Publishers, Perth, pp 56-97.

Cook J 1773, 'Departure from New South Wales. A particular Description of the Country, its Products, and People. A Specimen of the Language; and some Observations upon the Currents and Tides' in Hawkesworth J (ed) An Account of the Voyages Undertaken by the Order of His Present Majesty, for Making Discoveries in the Southern Hemisphere, and Successively Performed by Commodore Byron, Captain Wallis, Captain Carteret, and Captain Cook, in the Dolphin, the Swallow, and the Endeavour: Drawn from the Journals which were kept by the Several Commanders, and from the Papers of Joseph Banks, Esq. Vol 2, New Zealand Electronic Text Centre, retrieved from www.nzetc.org/tm/scholarly/tei-HawAcco-t1-g1-t1-body-d4-d6.html

Cook, C O L 1966, John Smithies: The Lonely Methodist, paper delivered at first meeting of the Methodist Historical Society, Perth, (Battye Library).

Cribb, A B & Cribb, J W 1981, Wild Medicine in Australia, William Collins Pty Ltd, Sydney.

Cribb, A B & Cribb, J W 1987, Wild Food in Australia, 2nd Ed, Fontana, Sydney.

Daniel, G 1979, *The Story of Wanneroo*, Shire of Wanneroo, Perth.

Daw, B, Walley, T & Keighery G 1997, Bush Tucker: Plants of the South-West, Department of Conservation and Land Management, Perth.

Delmege, S 2005, 'A Trans-Generational Effect of The Aborigines Act 1905 (WA): The Making of the Fringedwellers in the South-West of Western Australia', Murdoch University Electronic Journal of Law, Vol 12, No 1, retrieved from Australasian Legal Information Institute (AustLII) database.

Department of Environment and Conservation 2009, Flora Base: The Western Australian Herbarium, Perth, retrieved from www.florabase.calm.wa.gov.au

Department of Environmental Protection 1996, System 6 and part System I update programme: unpublished bushland plot and area records analysis, Kensington, pp 42-45.

Department of Indigenous Affairs (DIA) 2004, Lost Lands Report, Perth, retrieved from www.dia.wa.gov.au/Land/Lost-Lands.

Department of Indigenous Affairs (DIA) 2005, Consulting Citizens: Engaging with Aboriginal Western Australians, Perth, retrieved from www.dia.wa.gov.au/Our-Business/Reports--Publications.

Department of Planning and Infrastructure (DPI) 2007, Your TravelSmart Guide to Yellagonga Regional Park 2007, Perth, retrieved from www.dpi.wa.gov.au/mediaFiles/Yellagonga 5DLweb.pdf.

Department of Environment and Conservation (DEC) 2003, Yellagonga Regional Park Management Plan 2003-2013, Perth, retrieved from www.dec.wa.gov.au/hotproperty/property/regional-parks/yellagongaregional-park.html

Drummond, J 1842-1843a, 'Letter No 1, 4 May 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843b, 'Letter No 2, 11 May 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843c, 'Letter No 6, 10 August 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843d, 'Letter No 7, 17 August 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843e, 'Letter No 8, 28 September 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843f, 'Letter No 9, 5 October 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843q, 'Letter No 10, 19 October 1842', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843h, 'Letter No 13, 8 March 1843', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843i, 'Letter No 14, 5 March 1843', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843j, 'Letter No 15, 22 March 1843', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843k, 'Letter No 16, 19 April 1843', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Drummond, J 1842-1843l, 'Letter No 17, 10 May 1843', Letters on the Botany of Western Australia (To the Editor of the Inquirer), (Battye Library).

Eyre, E J 1845, 'Manners and Customs of the Aborigines of Australia', in his Journals of Expeditions of Discovery into Central Australia and Overland from Adelaide to King George's Sound in the Years 1840-1: Sent by the Colonists of South Australia, with the Sanction and Support of the Government: Including an Account of the Manners and Customs of the Aborigines and the State of their Relations with Europeans, Vol 2, Project Gutenberg, retrieved from www.gutenberg.org/etext/5345.

Facey, A B 1984, A Fortunate Life, Viking, Melbourne.

Fielder, J 1994, 'Sacred Sites and the City: Urban Aboriginality, Ambivalence, and Modernity', Boundary 2, Vol 21, No 1, pp 65-83, retrieved from JSTOR database.

Flinders, Matthew 1801-1803, Voyage to Terra Australis: Undertaken for the Purpose of Completing the Discovery of that Vast Country and Produced in the Years 1801, 1802 and 1803, Vol 1, Project Gutenberg, retrieved from www.gutenberg.org/etext/12929.

Fountain, P 1907, Rambles of an Australian Naturalist, from the Notes and Journals of Thomas Ward, John Murray, London, retrieved from American Libraries Internet Archive.

Friends of Yellagonga Regional Park (FYRP) nd, Species of Plants Grown for Yellagonga Regional Park, retrieved from www.yellagonga.org/nursery.asp.

Gardner, C A & Bennetts, H W 1956, The Toxic Plants of Western Australia, West Australian Newspapers, Perth.

Geographe Community Landcare Nursery 2005, Species: Solanum Symonii, MySouthWest.com, Busselton, retrieved from www.geographelandcarenursery.mysouthwest.com.au/2009%20Nursery%20List/Solanum% 20symonii.

Giles, E 1872-1876, Australia Twice Traversed: The Romance of Exploration, Being a Narrative Compiled from the Journals of Five Exploring Expeditions Into and Through Central South Australia and Western Australia, From 1872 to 1876, Project Gutenberg, retrieved from www.gutenberg.org/etext/4974.

Gilks, L A 1977, Wanneroo: Development of a District, Teachers Higher Certificate, Education Department of Western Australia, Perth, (Joondalup Library).

Gott, B 1982, 'Ecology of Root Use by the Aborigines of Southern Australia', Archaeology in Oceania, Vol 17, No 3, pp 59-68.

Gott, R 2006, Bush Tucker, Heinemann Library, Melbourne.

Graham, D 1996, Western Australia's Other History: A Short Guide, Western Australian Advisory Committee on Reconciliation, Perth.

Green, N J 1961, Perth and its People 1830-1840, Thesis, Claremont Teachers College, Perth, (Battye Library).

Green, N J 1962, The Natives of the Swan River Colony, unpublished, Perth, (Battye Library).

Green, N J 1984, Broken Spears: Aboriginals and Europeans in the Southwest of Australia, Focus Education Services, Perth.

Green, N J 1998, 'From Princes to Paupers: The Struggle for Control of Aborigines in Western Australia 1887-1898', Early Days, Vol 11, No 4, pp 446-462, retrieved from Informit database.

Gregory, A C 1887, 'Appendix II: Memoranda on the Aborigines of Australia', The Journal of the Anthropological Institute of Great Britain and Ireland, Vol 16, No 2, pp 131-133, retrieved from JSTOR database.

Grey, G 1841a, Journals of Two Expeditions of Discovery in North-West and Western Australia, Vol 1, Project Gutenberg, retrieved from www.gutenberg.org/etext/16027.

Grey, G 1841b, Journals of Two Expeditions of Discovery in North-West and Western Australia, Vol 2, Project Gutenberg, retrieved from http://www.gutenberg.org/etext/16145.

Haebich, A 1992, For their Own Good: Aborigines and Government in the Southwest of Western Australia, 1900-1940, University of Western Australia Press, Perth.

Hagger, J 1979, Australian Colonial Medicine, Rigby, Adelaide.

Hallam, S J & Tillbrook, L 1990, Aborigines of the Southwest Region, 1829-1840: The Bicentennial Dictionary of Western Australians, Vol 8, University of Western Australia Press, Perth.

Hallam, S J 1979, Fire and Hearth: A Study of Aboriginal Usage and European Usurpation in South-Western Australia, Prehistory and Material Culture Series No 8, Australian Institute of Aboriginal Studies, Canberra.

Hallam, S J 1989, 'Plant Usage and Management in Southwest Australian Aboriginal Societies', in Harris D R & Hillman G C (eds), Foraging and Farming: The Evolution of Plant Exploitation, Unwin Hyman, Sydney, pp 136-151.

Hallam, S J 1998a, 'Wanneroo, Joondalup, Yanchep, the Aborigines', in Kauler, L B, Cultural Significance of Aboriginal Sites in the Wanneroo Area, report prepared for Harry Nannup of Gnangara Aboriginal Community College, Heritage Council of WA, Perth, pp 76-104.

Hallam, S J 1998b, 'Paradise Lost: The Aborigines', in Gentilli J & Bekle, H (eds), Wanneroo, Joondalup, Yanchep: Planning, People, Environment, City of Wanneroo, Perth, pp 91-114.

Hammond, J E 2005 (1933), Winjan's People: The Story of the South West Australian Aborigines, Hesperian Press, Perth.

Harris D R & Gordon C H (eds) 1989, Foraging and Farming: The Evolution of Plant Exploitation, Unwin Hyman, London.

Hayes A & Hayes S 2007, Ngambunyjarri: Ngambunyjarri Thalanyjibarndi Yininyjarri (Thalanyji Plant Names and Uses), Wangka Maya Pilbara Aboriginal Language Centre, Onslow.

Hele, A E 2003, Researchers' Extension Program for the Native Foods Industry: A Report for the Rural Industries Research and Development Corporation, Rural Industries Research and Development Corporation, Canberra.

Henderson, G & Henderson K-J 1988, Unfinished Voyages: Western Australian Shipwrecks 1841-1880, University of Western Australia Press, Perth.

Heritage Council of Western Australia (HCWA) 1998, Yaberoo Budjara Heritage Trail, (brochure), Perth, retrieved from www.tourism.heritage.wa.gov.au.

Heritage Council of Western Australia (HCWA) 1999, Star Swamp Heritage Trail, (brochure), Perth. retrieved from www.tourism.heritage.wa.gov.au.

Hiddins, L 2001, Bush Tucker Field Guide, Penguin Books, Melbourne.

Isaacs, J 1996, A Companion Guide to Bush Food, Lansdowne Publishing Pty Ltd, Sydney.

Isaacs, J 2002, Bush Food: Aboriginal Food and Herbal Medicine, New Holland Publishers, Sydney.

Jones, S 2007, Sharing our Stories: Guidelines for Heritage Interpretation, The National Trust of Australia WA and Museums Australia WA, Perth, retrieved from www.ntwa.com.au/sharingourstories.

Juluwarlu Aboriginal Corporation 2003, Wanggalili: Yindjibarndi and Ngarluma Plants, Roebourne.

Kalotas, A C 1996, 'Aboriginal Knowledge and Use of Fungi', in Orchard A E, Mallett, K & Grgurinovic, C (eds), Fungi of Australia, Vol 1B, ABRS/CSIRO Australia, Canberra, pp 269-295.

Kauler, L B 1998, Cultural Significance of Aboriginal Sites in the Wanneroo Area, report prepared for Harry Nannup of Gnangara Aboriginal Community College, Heritage Council of WA, Perth.

Kenderdine, S 1995, Shipwrecks 1656-1942: A Guide to Historic Wreck Sites of Perth, Western Australian Maritime Museum, Perth.

Kennealy, S 1994, Oral Histories of Wanneroo Wetlands: Recollections of Wanneroo Pioneers, Water Authority of Western Australia, Perth.

Kings Park and Botanic Gardens (KPBG) nd, Aboriginal Life, Perth, retrieved from www.bgpa.wa.gov. au/c/bgpa/pub/stories/bgpa/docs/brochures/pdf/aboriginal\_dl.pdf.

Kirkpatrick, J 1999, A Continent Transformed: Human Impact on the Natural Vegetation of Australia, 2nd Ed, Oxford University Press, Melbourne.

Lassak E V & McCarthy T 2001, Australian Medicinal Plants, Reed New Holland, Sydney.

Latz, P 1995, Bushfires and Bushtucker: Aboriginal Plant Use in Central Australia, IAD Press, Alice Springs.

Lepp, H 2003, Aboriginal Use of Fungi, Australian National Herbarium, Canberra, retrieved from www.anbg.gov.au/fungi/aboriginal.html.

Levitt, D 1981, Plants and People: Aboriginal Uses of Plants on Groote Eylandt, Australian Institute of Aboriginal Studies, Canberra.

Lippmann, L 1994, Generations of Resistance: Mabo and Justice, 3rd Ed, Longman Cheshire Pty Ltd, Melbourne.

Low, T 1989, Bush Tucker: Australia's Wild Food Harvest, Angus & Robertson Publishers Australia, Sydney.

Low, T 1990, Bush Medicine: A Pharmacopoeia of Natural Remedies, Angus & Robertson Publishers Australia, Sydney.

Low, T 1991, Wild Herbs of Australia and New Zealand, Angus and Robertson, Sydney.

Lyon, R M 1833, 'A Glance at the Manners, and Language of the Aboriginal Inhabitants of Western Australia; With a Short Vocabulary', in Green, N (ed) 1979, Noongar, The People: Aboriginal Customs in the Southwest of Australia, Creative Research Publishers, Perth, pp 141-180.

MacIntyre Dobson and Associates Pty Ltd & O'Reilly, T 2002, Report on an Ethnographic, Ethnohistorical, Archaeological and Indigenous Environmental Survey of the Underwood Avenue Bushland Project Area, Shenton Park, paper prepared for the University of Western Australia, Perth, (Battye Library).

Major, R H (ed) 1859, 'Extract from the Journal of a Voyage made to the Unexplored South Land, by Order of the Dutch East India Company, in the Years 1696 and 1697, by the Hooker De Nyptang, the Ship De Geelvink, The Galiot de Wesel and the Return to Batavia' 1701, in Early Voyages to Terra Australis, Now Called Australia: A Collection of Documents, and Extracts from Early Manuscript Maps, Illustrative of the History of Discovery on the Coasts of that Vast Island, from the Beginning of the Sixteenth Century to the Time of Captain Cook, Project Gutenberg, retrieved from www.gutenberg.net.au/ebooks06/0600361h.html pp 120-133.

Marwick, B 2002, Stories of Old Wanneroo: As Told to Bill Marwick, Bill Marwick, Perth.

McGuire, M I 1996, The Use and Management of Wetlands by Aboriginal People on the Swan Coastal Plain, Western Australia: An Exploratory Study, Honours Thesis, Murdoch University, Perth.

McNair, W & Rumley, H 1981, Pioneer Aboriginal Mission: The Work of Wesleyan Missionary John Smithies in the Swan River Colony 1840-1855, University of Western Australia Press, Perth.

Meagher, S J 1974, 'The Food Resources of the Aborigines in the South-West of Western Australia', Records of the Western Australian Museum, Vol 3, No 1, pp 14-65.

Milligan, W 1837, 'Some Account of the New Colony of Western Australia, more especially of the Swan River District, the Natives, Settlers, Climate, Soil, Productions, etc', The Madras Journal of Literature and Science, Vol 6, July-December, pp 304-336, retrieved from Google Books database.

Monks, G 1993, Mission Farm, unpublished, Perth, (Joondalup Library).

Moore, G F 1835, 'Excursion to the Northward from the Journal of George Fletcher Moore Esq', Exploration Diaries 1834-2835, Vol 2, Western Australian Department of Lands and Surveys, Perth, pp 318-281.

Moore, G F 1884, 'A Descriptive Vocabulary of the Language in Common Use Amongst the Aborigines of Western Australia; With Copious Meanings, Embodying Much Interesting Information Regarding the Habits, Manners, and Customs of the Natives, and the Natural History of the Country', in Stannage C T (ed) 1978, Diary of Ten Years of an Early Settler in Western Australia, University of Western Australia Press, Perth.

Moore, G F 2006, The Millendon Memoirs: George Fletcher Moore's Western Australian Diaries and Letters, 1830-1841, Cameron J (ed), Hesperian Press, Perth.

Moore, G F nd, 'Excursion to a River to the Northward from the Journal of George Fletcher Moore Esq', Exploration Diaries 1834-2835, Vol 2, Western Australian Department of Lands and Surveys, Perth, pp 381-404.

Newton, D 2002, The Wood Block Road to Wanneroo: Celebrating 150 Years of Settlement, Wanneroo Heritage Art, Perth.

Nind, S 1831, 'Description of the Natives of King George's Sound (Swan River Colony) and Adjoining Country', Journal of the Royal Geographical Society of London, Vol 1, pp 21-51, retrieved from JSTOR database.

O'Connor, R, Quartermaine, G & Bodney, C 1989, Report on an Investigation into Aboriginal Significance of Wetlands and Rivers in the Perth-Bunbury Region, Western Australian Water Resources Council, WA.

Osborne, R 1985, 'One Man's Poison', Palms and Cycads, No 7, retrieved from Palms and Cycad Societies of Australia, www.pacsoa.org.au/cycads/Articles.

Pearce, D 2003, 'The Transmission of Stories, History and Cultural Knowledge with the Noongar Community is Alive and Well', Studies in Western Australian History, No 23, pp 55-60, retrieved from Informit database.

Peile, A R 1980, 'Preliminary Notes of the Ethno-Botany of the Gugadja Aborigines at Balgo, Western Australia', Western Australian Herbarium Notes, No 3, pp 59-64.

Playford, P 1998, Voyage of Discovery to Terra Australis by Willem de Vlamingh in 1696-97, Western Australian Museum, Perth.

Powell, R 1990, Leaf and Branch: Trees and Tall Shrubs of Perth, Department of Conservation and Land Management, Perth.

Reid, E 1977, The Records of Western Australian Plants Used by Aboriginals as Medicinal Agents, Western Australian Institute of Technology Pharmacy Department, Perth.

Ringer, S 2009, Some Notes of Acacia, Australian Bushfoods Magazine, Maleny, retrieved from www.ausbushfoods.com/new/index.php?option=com content&view=article&id=164:wattle-notes&catid= 13:edible&Itemid=64.

Rippey, E & Rowland, B 2004, Coastal Plants: Perth and the South-West Region, University of Western Australia Press, Perth.

Robinson, M V 1984, A Brief History of Nyoongah Culture and Administration in the South-West of Western Australia, paper prepared for the Southern Aboriginal Corporation, Katanning, (Battye Library).

Roth, W E 1903, 'Notes of Savage Life in the Early Days of West Australian Settlement', Proceedings of the Royal Society of Queensland, Vol 17, pp 45-69.

Russo, G 1998, 'Early Pioneers', in Gentilli J & Bekle, H (eds), Wanneroo, Joondalup, Yanchep: Planning, People, Environment, City of Wanneroo, Perth, pp 115-138.

Searle, S 2009, Traditional Uses of Australian Acacias, World Wide Wattle, retrieved from www.worldwidewattle.com/infogallery/utilisation/aboriginal.php.

Seddon, G 1972, Sense of Place: A Response to an Environment the Swan Coastal Plain Western Australia, University of Western Australia Press, Perth.

Seddon, G 2005, The Old Country: Australian Landscapes, Plants and People, Cambridge University Press, Melbourne.

Shipley, T 2003, Full Circle: A History of the Wesley Church, Perth, Uniting Church Synod of Western Australia, Perth.

Skinner, D 1978, Bush Medicines, (manuscript), Perth, (Battye Library).

Skyring, F 2003, 'History Wars: Debates about History in the Native Title Process', in Choo, C & Hollbach S (eds), History and Native Title, Studies in Western Australian History, No 3, University of Western Australia Press, pp 71-82.

Smith, K & Smith I, 1999, Grow your own Bushfoods, New Holland Publishers, Sydney.

Smith, M 1982, 'Late Pleistocene Zamia Exploitation in Southern Western Australia', Archaeology in Oceania, Vol 17, No 3, pp 117-121.

Smithies, J 1840, Report on the Aborigines in the Swan River Colony, (manuscript), Perth, (Battye Library).

Smyth, R B 1878, The Aborigines of Victoria With Notes Relating to the Habits of the Natives of Other Parts of Australia and Tasmania, Vol 1, John Peeres Government Printer, London, retrieved from American Libraries Internet Archive.

South West Aboriginal Land and Sea Council (SWALSC), Host, J & Owen, C 2009, It's Still in My Heart, This is My Country: The Single Noongar Claim History, UWA Publishing, Perth.

Stannage, C T 1996, Lakeside City: The Dreaming of Joondalup, University of Western Australia Press, Perth.

Stewart, K & Percival, B 1997, *Bush Foods of New South Wales: A Botanic Record and An Aboriginal Oral History*, Royal Botanic Gardens, Sydney retrieved from www.rbgsyd.nsw.gov.au/education/Resources/bush\_foods.

Stokes, J L 1846, Discoveries in Australia, With An Account Of The Coasts And Rivers Explored And Surveyed During The Voyage Of H.M.S. Beagle, In The Years 1837-38-39-40-41-42-43, By Command Of The Lords Commissioners Of The Admiralty, Also A Narrative Captain Owen Stanley's Visits To The Islands In The Arafura Sea, Vol 2, Project Gutenberg, retrieved from www.gutenberg.org/etext/12146.

Swan Catchment Council (SCC) nd, Noongar Coastal Trail, (brochure), Perth.

Symons, P & Symons, S 1994, Bush Heritage: An Introduction to the History of Plant and Animal use by Aboriginal People and Colonists in the Brisbane and Sunshine Coast Areas, Pat & Sim Symons, Nambour.

Tate, G 2008, Biodiversity in Yellagonga Regional Park, videorecording, City of Joondalup, Perth.

Tindale 1974, 'Western Australia: Tindale Tribal Boundaries', in his *Aboriginal Tribes of Australia: Their Terrain, Environmental Controls, Distribution, Limits, and Proper Names*, Department of Indigenous Affairs, Perth, retrieved from www.dia.wa.gov.au/ Information/Maps.

W G Martinick & Associates Pty Ltd, Gutterage Haskins & Davey & Briandelfs and Associates 1997, *Report - City of Wanneroo Inventory of Heritage Places*, City of Wanneroo, Perth.

Ward, D J & Lamont, B B 2000, 'Probability of Grasstrees (*Xanthorrhoea Preissii*) Flowering After Fire', *Journal of the Royal Society of Western Australia*, Vol 83, No 1, pp 13-16, retrieved from www.rswa.org.au.

White, S A 1921, 'The Native of the Everard Ranges', *Man*, Vol 21, No 67, pp 113-115, retrieved from JSTOR database.

