

10071
Ilmenite Basalt (high K)
189.5 grams



Figure 1: Photo of 10071,5 illustrating large vesicles and smooth surface, rounded by micrometeorite bombardment. NASA S76-22608. Cube is 1 cm.

Introduction

10071 is a high titanium basalt with up to 15 % ilmenite and with the new mineral armalcolite. It has prominent vesicles up to 1 mm (figure 1). The exposed surface is rounded by micrometeorite bombardment.

A very significant feature of this rock, not generally seen in Apollo samples, is the apparent igneous contact between to regions with different texture and different composition (figure 2).

10071 has been measured to be 3.68 b.y. old and has been exposed to cosmic rays for ~370 m.y.

Petrography

Schmitt et al. (1970) termed 10071 as a “fine-grained, vesicular to vuggy, plumose basalt.” James and Jackson (1970) described the texture as “intersertal”.

Drake and Weill (1971) found that there were two distinct basalt lithologies separated by a distinct boundary. The finer-grained portion has a variolitic texture with acicular intergrowths of fine clinopyroxene and plagioclase. In this fine-grained portion the pyroxene exhibits extreme Fe-enrichment all the way to pyroxferroite (figure 4). Beatty and Albee (1978) termed the fine-grained portion “Iota” and studied it separately.





Figure 2: Photomicrographs of thin section of 10071 showing distinctly different fabric (with plane-polarized light, left and crossed Nicols, right). Scale is 2.5 mm. NASA S70-17978 and 979.

Roedder et al. (1970) studied melt inclusions in minerals in 10071 and discussed liquid immiscibility.

Mineralogy

Olivine: Minor olivine occurs as cores in large pyroxene grains where it is found to be Mg-rich,

Pyroxene: Pyroxenes are chemically zoned (Drake and Weill 1971, Beatty and Albee 1978)(figures 3 and 4).

Plagioclase: Plagioclase (An₇₀₋₈₀) is found to be non-stoichiometric, with excess silica (Drake and Weill 1971, Beatty and Albee 1978). Stewart et al. (1970) and Appleman et al. (1971) determined crystallographic data.

Ilmenite: Haggerty et al. (1970) and Drake and Weill (1971) give the analysis of ilmenite (with ~2% MgO). Stewart et al. (1970) determined crystallographic data.

Armalcolite: Haggerty et al. (1970) determined an opaque phase as “magnesian ferro pseudobrookite”

and gave an analysis (table 2). The phase was determined to be a new mineral and was named armalcolite in honor of the crew of Apollo 11 (Armstrong, Aldrin and Collins). Anderson et al. (1970) fully discuss armalcolite. Several large grains of armalcolite were found mantled by ilmenite (Beatty and Albee 1978).

Apatite: Drake and Weill (1971) determined REE in apatite in 10071.

Chemistry

Tatsumoto et al. (1970) and Rosholt and Tatsumoto (1970) found 10071 had high U = 0.873 ppm, such that Compston et al. (1970) correctly assigned it to the high K group of Apollo 11 basalts. Ehmann et al. (1975) re-determined Zr and Hf (562 and 16.6 ppm respectively).

The chemical composition of “Iota” (Beatty and Albee 1978) is the same as that measured on the fine-grained portion (Drake and Weill 1971), and is distinct from

Mineralogical Mode for 10071

	James and Jackson 70	Beatty and Albee 1978	Drake and Weill 1971	Haggerty et al. 1970
Olivine	1.7	1.3	1.0	
Pyroxene	51.5	47.8	53.5	58.3
Plagioclase	20.6	23.6	23.5	21.7
Ilmenite	16.8	14.8	15.5	14.7
mesostasis	6.2	9.6	6.5	2.7
silica	2.8	2.2	0.5	0.8
troilite	0.1	0.44		
phosphate	tr.	0.24		




Figure 3: Pyroxene diagram for main part of 10071 (data replotted from Beaty and Albee 1978).




Figure 4: Pyroxene diagram for fine-grained part of 10071 (data replotted from Beaty and Albee 1978).

the coarse-grained portion of the rock. The trace element content of “Iota”, has not been determined.

Radiogenic age dating

Papanastassiou et al. (1970) determined the Rb/Sr isochron age of 3.68 b.y. for 10071. The K/Ar age of the whole rock was determined as 2.88 b.y. and for the plagioclase was 3.35 b.y. by Eberhardt et al. (1974). Stettler et al. (1973, 1974) determined 3.51 b.y. by the Ar/Ar plateau technique (figure 7) and noted that data for plagioclase gave the best plateau.

Cosmogenic isotopes and exposure ages

Marti et al. (1970) and Eberhardt et al. (1974) determined the ⁸³Kr exposure age for 10071 as 372 ± 22 and 350 ± 15 m.y. respectively. Srinivasan (1974) used the Xe data of Marti et al. (1970) and Eberhardt et al. (1974) to calculate 212 and 300 m.y. exposure ages for 10071. Stettler et al. (1973, 1974) and Guggisberg et al. (1979) reported ³⁸Ar cosmic ray exposure ages of ~380 m.y. and 370-430 m.y. respectively.

Mineralogical Mode for “Iota”

	Beaty and Albee 1978	Drake and Weill 1971
Olivine	absent	0
Pyroxene	40.7	44.5
Plagioclase	36.3	35.5
Ilmenite	6.3	4.0
mesostasis	11	8.5
silica	5	7.5
troilite	0.4	
phosphate	0.34	

Other Studies

Oxygen isotopes were reported for mineral separates of 10071 by Onuma et al. (1970).

The concentrations of Sm, Nd, Lu and Hf and the isotopic ratios of ¹⁴³Nd/¹⁴⁴Nd and ¹⁷⁶Hf/¹⁷⁷Hf were determined by Unruh et al. (1984). Eugster et al. (1970) determined the isotopic composition of Gd.

The rare gas component of 10071 was determined by Funkhouser et al. (1970), Pepin et al. (1970), Marti et al. (1970), Bogard et al. (1971) and Eberhardt et al. (1970, 1974).

Processing

Apollo 11 samples were originally described and cataloged in 1969 and “re-cataloged” by Kramer et al. (1977). In 2005, a portion of this sample was presented to Neal Armstrong as part of the Astronaut Ambassador Program for public display at a museum of his choosing. There are 5 thin sections.

List of Photo #s for 10071

- S69-47614
- S69-47288 – 289 PET B&W
- S69-47292 – 297
- S69-47304 – 309
- S69-47353 – 359
- S70-17978 – 979 TS
- S70-48981 – 982 TS
- S76-22602 – 609 ,5
- S76-23372 – 373 ,11
- S76-26082 – 083 ,13
- S76-26321 – 322 TS B&W




Figure 5: Composition of 10071 compared with that of other Apollo lunar samples.




Table 2. Chemical composition of armalcolite.

reference weight	Beaty78	Anderson70	Haggerty70	
SiO2 %	0.11			(a)
TiO2	73.15	73.4	73.4	(a)
Al2O3	1.57	1.62	1.62	(a)
FeO	15.51	15.3	15.3	(a)
MnO	0.01	0.08	0.08	(a)
MgO	7.66	7.7	7.7	(a)
CaO		0.01	0.01	(a)
Cr2O3	2.02	2.15	2.15	(a)

tech: (a) emp

Table 3	U ppm	Th ppm	K ppm	Rb ppm	Sr ppm	Nd ppm	Sm ppm	technique
Tatsumoto et al. 1970	0.873	3.43						IDMS
Rosholt and Tats 1970	0.873							
Wanless et al. 1970			2574	5.88	170.6			
Goles et al. 1970							20	INAA
Annell and Helz 1970				5.2				
Gast 1970			2770	5.93	161	64.5	22.7	IDMS
Rhodes 1980		3.4		5.6	165		19.5	INAA
Unruh et al. 1984						61.5	20.9	INAA
Papamastassiou 70				5.82	165			IDMS




Figure 6: Normalized rare-earth-element composition for high-K basalt 10071 (the line) compared with that of low-K basalt 10020 and high-K basalt 10049 (the dots) (data from Wiesmann et al. 1975).




Figure 7: Argon release curves for mineral separates from basalt 10071 (from Stettler et al. 1973). The plagioclase separate gave the "best" plateau age = 3.51 +/- 0.06 b.y.

Summary of Age Data for 10071

	Rb/Sr	Ar/Ar (plag)K/Ar
Papanastassiou et al. 1970	3.68 ± 0.02 b.y.	
Stettler et al. 1973		3.51 ± 0.06
Eberhardt et al. 1974		2.88 ± 0.06
Wanless et al. 1970		2.875

Note: Ages not corrected for new decay constants.

Table 1. Chemical composition of 10071.

<i>reference weight</i>	Gast70	Wiesmann75	Goles70	Rhodes80	Annell70	Drake 71 coarse	Drake 71 fine-grained	Beaty78 "lota"	
SiO ₂ %			40.86	(c) 40.51	(d)	42.2	46.9	(f) 47.89	(g)
TiO ₂	13.8		11.7	(c) 11.87	(d)	10.2	4.8	(f) 5.75	(g)
Al ₂ O ₃			8.22	(c) 7.91	(d)	9.7	14.5	(f) 11.29	(g)
FeO			19.17	(c) 19.65	(d)	16.5	15.4	(f) 17.56	(g)
MnO			0.21	(c) 0.25	(d) 0.29	(e)			0.22 (g)
MgO			7.29	(c) 7.51	(d)	7.7	3.3	(f) 4.69	(g)
CaO	10.9		10.07	(c) 10.48	(d)	11.3	10.8	(f) 10.87	(g)
Na ₂ O	0.52	(a) 0.52	(a) 0.49	(c) 0.53	(c)	0.64	0.66	(f) 0.8	(g)
K ₂ O	0.33	(b) 0.33	(b)	0.3	(d)	0.44	1.02	(f) 0.42	(g)
P ₂ O ₅				0.22	(d)				0.16 (g)
S %									0.25 (g)
<i>sum</i>									
Sc ppm			73.2	(c) 78	(d) 97	(e)			
V			92	(c)	78	(e)			
Cr			2170	(c) 2450	(c) 3060	(e)			
Co			27.1	(c) 25.4	(d) 33	(e)			
Ni					7	(e)			
Cu			11	(c)	14	(e)			
Zn									
Ga				8.8	(d) 4.8	(e)			
Ge ppb									
As									
Se									
Rb	5.93	(b) 5.93	(b)	5.6	(d) 5.2	(e)			
Sr	161	(b) 161	(b)	165	(d) 140	(e)			
Y				166	(d) 162	(e)			
Zr			210	(c)	644	(e)			
Nb					24	(e)			
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb									
Cd ppb									
In ppb									
Sn ppb									
Sb ppb									
Te ppb									
Cs ppm	0.17	(b) 0.17	(b)						
Ba	327	(b) 327	(b) 450	(c)	470	(e)			
La	28.8	(b) 28.8	(b) 25.8	(c) 24.1	(c) 27	(e)			
Ce	83.5	(b) 83.5	(b) 84	(c) 75	(c)				
Pr									
Nd	64.5	(b) 64.5	(b)						
Sm	22.7	(b) 22.7	(b) 20	(c) 19.5	(c)				
Eu	2.32	(b) 2.32	(b) 2.12	(c) 2.03	(c)				
Gd	29.3	(b) 29.3	(b)						
Tb			5.7	(c) 4	(c)				
Dy	33.5	(b) 33.5	(b)						
Ho			9.2	(c)					
Er	21.3	(b) 21.3	(b)						
Tm									
Yb	20.5	(b) 18.2	(b) 20.8	(c) 16.2	(c)				
Lu	2.87	(b) 2.87	(b) 3.08	(c) 2.33	(c)				
Hf			19.1	(c) 15.9	(c)				
Ta			2	(c) 2.7	(c)				
W ppb									
Re ppb									
Os ppb									
Ir ppb									
Pt ppb									
Au ppb									
Th ppm				3.4	(d)				
U ppm			0.69	(c)					

technique: (a) AA, (b) IDMS, (c) INAA, (d) XRF, (e) emission spec., (f) emp, (g) modal

References for 10071

- Anderson A.T. (1971) Exotic armalcolite and the origin of Apollo 11 ilmenite basalts. *Geochim. Cosmochim. Acta* **35**, 969-973.
- Anderson A.T. and 10 others (1970) Armalcolite: A new mineral from the Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 55-63.
- Annell C.S. and Helz A.W. (1970) Emission spectrographic determination of trace elements in lunar samples from Apollo 11. *Proc. Apollo 11 Lunar Sci. Conf.* 991-994.
- Beatty D.W. and Albee A.L. (1978) Comparative petrology and possible genetic relations among the Apollo 11 basalts. *Proc. 9th Lunar Planet. Sci. Conf.* 359-463.
- Bogard D.D., Funkhouser J.G., Schaeffer O.A. and Zahringer J. (1971) Noble gas abundances in lunar material-cosmic ray spallation products and radiation ages from the Sea of Tranquillity and the Ocean of Storms. *J. Geophys. Res.* **76**, 2757-2779.
- Compston W., Arriens P.A., Vernon M.J. and Chappell B.W. (1970a) Rubidium-strontium chronology and chemistry of lunar material. *Science* **167**, 474-476.
- Compston W., Chappell B.W., Arriens P.A. and Vernon M.J. (1970b) The chemistry and age of Apollo 11 lunar material. *Proc. Apollo 11 Lunar Sci. Conf.* 1007-1027.
- Drake M.J. and Weill D.F. (1971) Petrology of Apollo 11 sample 10071. A differentiated mini-igneous complex. *Earth Planet. Sci. Lett.* **13**, 61-70.
- Eberhardt P., Geiss J., Graf H., Grogler N., Krahenbuhl U., Schwaller H. and Stettler A. (1974b) Noble gas investigations of lunar rocks 10017 and 10071. *Geochim. Cosmochim. Acta* **38**, 97-120.
- Ehmann W.D. and Chyi L.L. (1974) Abundances of the group IVB elements, Ti, Zr, and Hf and implications of their ratios in lunar materials. *Proc. 5th Lunar Sci. Conf.* 1015-1024.
- Funkhouser J.G., Schaeffer O.A., Bogard D.D. and Zahringer J. (1970) Gas analysis of the lunar surface. *Proc. Apollo 11 Lunar Sci. Conf.* 1111-1116.
- Gast P.W., Hubbard N.J. and Wiesmann H. (1970b) Chemical composition and petrogenesis of basalts from Tranquillity Base. *Proc. Apollo 11 Lunar Sci. Conf.* 1143-1163.
- Goles G.G., Randle K., Osawa M., Lindstrom D.J., Jerome D.Y., Steinborn T.L., Beyer R.L., Martin M.R. and McKay S.M. (1970) Interpretations and speculations on elemental abundances in lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1177-1194.
- Guggisberg S., Eberhardt P., Geiss J., Grogler N., Stettler A., Brown G.M. and Peckert A. (1979) Classification of the Apollo-11 basalts according to Ar³⁹-Ar⁴⁰ ages and petrological properties. *Proc. 10th Lunar Planet. Sci. Conf.* 1-39.
- Haggerty S.E., Boyd F.R., Bell P.M., Finger L.W. and Bryan W.B. (1970) Opaque minerals and olivine in lavas and breccias from Mare Tranquillitatis. *Proc. Apollo 11 Lunar Sci. Conf.* 513-538.
- James O.B. and Jackson E.D. (1970) Petrology of the Apollo 11 ilmenite basalts. *J. Geophys. Res.* **75**, 5793-5824.
- Kramer F.E., Twedell D.B. and Walton W.J.A. (1977) **Apollo 11 Lunar Sample Information Catalogue** (revised). Curator's Office, JSC 12522
- LSPET (1969) Preliminary examination of lunar samples from Apollo 11. *Science* **165**, 1211-1227.
- Marti 70
- Onuma N., Clayton R.N. and Mayeda T. (1970) Apollo 11 rocks: Oxygen isotope fractionation between minerals and an estimate of the temperature of formation. *Proc. Apollo 11 Lunar Sci. Conf.* 1429-1434.
- Papanastassiou D.A., Wasserburg G.J. and Burnett D.S. (1970a) Rb-Sr ages of lunar rocks from the Sea of Tranquillity. *Earth Planet. Sci. Lett.* **8**, 1
- Papanastassiou D.A. and Wasserburg G.J. (1971a) Lunar chronology and evolution from Rb-Sr studies of Apollo 11 and 12 samples. *Earth Planet. Sci. Lett.* **11**, 37-62.
- Pepin R.O., Nyquist L.E., Phinney D. and Black D.C. (1970) Rare gases in Apollo 11 lunar material. *Proc. Apollo 11 Lunar Sci. Conf.* 1435-1454.

- Rhodes J.M. and Blanchard D.P. (1980) Chemistry of Apollo 11 low-K mare basalts. *Proc. 11th Lunar Planet. Sci. Conf.* 49-66.
- Roedder E. and Weiblen P.W. (1970) Lunar petrology of silicate melt inclusions, Apollo 11 rocks. *Proc. Apollo 11 Lunar Sample Conf.* 801-837.
- Rose H.J., Cuttitta F., Dwornik E.J., Carron M.K., Christian R.P., Lindsay J.R., Ligon D.T. and Larson R.R. (1970b) Semimicro X-ray fluorescence analysis of lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1493-1497.
- Rosholt J.N. and Tatsumoto M. (1970) Isotopic composition of uranium and thorium in Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1499-1502.
- Schmitt H.H., Lofgren G., Swann G.A. and Simmons G. (1970) The Apollo 11 samples: Introduction. *Proc. Apollo 11 Lunar Science Conf.* 1-54.
- Stettler A., Eberhardt P., Geiss J., Grogler N. and Maurer P. (1974) On the duration of the lava flow activity in Mare Tranquillitatis. *Proc. 5th Lunar Sci. Conf.* 1865-1888.
- Stewart D.B., Appleman D.E., Huebner J.S. and Clark J.R. (1970) Crystallography of some lunar plagioclase. *Science* **167**, 634-635.
- Sutton R.L. and Schaber G.G. (1971) Lunar locations and orientations of rock samples from Apollo missions 11 and 12. *Proc. 2nd Lunar Sci. Conf.* 17-26.
- Tatsumoto M. (1970) Age of the Moon: An isotopic study of U-Th-Pb systematics of Apollo 11 lunar samples - II. *Proc. Apollo 11 Lunar Sci. Conf.* 1595-1612.
- Unruh D.M., Stille P., Patchett P.J. and Tatsumoto M. (1984) Lu-Hf and Sm-Nd evolution in lunar mare basalts. *Proc. 14th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **88**, B459-B477.
- Wanless R.K., Loveridge W.D. and Stevens R.D. (1970) Age determinations and isotopic abundance measurements on lunar samples (Apollo 11). *Proc. Apollo 11 Lunar Sci. Conf.* 1729-1739.
- Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC