



Four new species of *Rasbora* of the Sumatrana group (Teleostei: Cyprinidae) from northern Sumatra, Indonesia

DANIEL N. LUMBANTOBING^{1,2}

¹Department of Biological Sciences, The George Washington University, 2023 G Street. NW, Lisner Hall 340, Washington, D.C., 20052. E-mail: dntobing@gwmail.gwu.edu

²Division of Fishes, Smithsonian Institution, National Museum of Natural History, Washington, D.C., 20013

Abstract

Four new species of the minnow genus *Rasbora* of the Sumatrana group, *R. arundinata*, *R. haru*, *R. maninjau*, and *R. bindumatoga*, are described from northern Sumatra. *Rasbora arundinata* is distinguished from all congeners in the Sumatrana group by the black midlateral stripe overall forming a reed-leaf-like profile. *Rasbora haru* differs from its congeners in the Sumatrana group in having the black midlateral stripe overall forming a stamen-like profile. The new species endemic to Lake Maninjau in central west Sumatra, *Rasbora maninjau*, is unique among all the congeners in the Sumatrana group in having a combination of the black midlateral stripe extending from the midhumeral region of uniform width, the prominent acutely triangular basicaudal blotch, and the oval supra-anal pigmentation. *Rasbora bindumatoga* is distinguished from all congeners in the Sumatrana group by a combination of the black rectangular subdorsal blotch, the absence of supra-anal pigmentation, and the somewhat oval basicaudal blotch. *Rasbora arundinata*, *R. maninjau*, and *R. bindumatoga* occur allopatrically in the northwestern coastal region of Sumatra, while *R. haru* is known from northeastern coastal area of Sumatra. A new diagnostic character for the Sumatrana group is described: partial exposure of the upper lip due to a submedial contact between the maxilla and the lower lip, which is marked posteriorly by a lachrymal groove.

Key words: Danioninae, *Rasbora*, new species, northern Sumatra, Sundaland

Introduction

Rasbora is a small-to-moderate-sized genus in the family Cyprinidae that lives throughout a vast geographical range within Asia, including the Indian subcontinent, southern China, and Southeast Asia (Weber & Beaufort 1916; Brittan 1954a). With currently 77 species, *Rasbora* constitutes the most species-rich genus in the cyprinid subfamily Danioninae (Eschmeyer 2013; Froese & Pauly 2013). Taxonomically, *Rasbora* has been widely considered as a catch-all group due to a lack of unique diagnostic characters (Brittan 1954a; Kottelat & Vidthayanon 1993; Liao *et al.* 2010; Tang *et al.* 2010). In the first and most comprehensive revision of *Rasbora*, Brittan (1954a) recognized three subgenera (*Rasbora*, *Rasboroides*, and *Megarasbora*), and further classified the subgenus *Rasbora* into eight species complexes: the *lateristriata*, the *sumatrana-elegans*, the *caudimaculata*, the *trifasciata*, the *argyrotaenia*, the *daniconius*, the *ethovenii*, and the *pauciperforata* complexes. Brittan's species complexes have been widely used as a practical system for the classification of the group and frequently revised by many authors (Kottelat & Vidthayanon 1993; Siebert & Guiry 1996; Kottelat 2005; Liao *et al.* 2010). In their brief revision of *Rasbora*, Kottelat & Vidthayanon (1993) replaced the category of species complex with 'species group,' which has been widely used by most of the later workers.

Authors after Brittan (1954a) have created several new genera (*Boraras*, *Brevibora*, *Horadandia*, *Kottelatia*, *Rasboroides*, *Trigonopoma*, and *Trigonostigma*) for some lineages within *Rasbora* (Kottelat & Vidthayanon 1993; Kottelat & Witte 1999; Liao *et al.* 2010). Despite these newly created genera, many workers still have recognized a larger assemblage equivalent to the concept of the genus *Rasbora sensu* Brittan (1954a), being frequently referred to as "the genus *Rasbora sensu lato*" (hereafter *Rasbora s. l.*). *Rasbora s. l.* comprises the genus *Rasbora sensu stricto* (hereafter *Rasbora s. s.*) and the new genera mentioned above (Kottelat & Vidthayanon 1993; Kottelat & Witte 1999; Conway 2005; Liao *et al.* 2010; Tang *et al.* 2010).

approximately 53 km to Painan on Padang-Painan road, 1°04'52.8"S, 100°27'26"E.

Rasbora vulgaris: BMNH 1905.5.6.4–5 (syntypes), 2, 37.7 and 34.8 mm SL, Malaysia, Selangor, Kuala Lumpur; RMNH 24858, 20, 68.1–78.1 mm SL, Malaysia, Malacca, Penang, Baya Lepas; RMNH 26373, 5, 21.7–50.4, Malaysia, Selangor, 2 km Batu Tiga to Subang Road.

Acknowledgment

I express my thanks to L. Parenti (USNM) for her support and guidance throughout the project. I gratefully acknowledge J. Burns (GWU) for his supervision and critical comments on the text. I thank R. Vari (USNM) and L. Page (UF) who critically reviewed the manuscript. I am indebted to R. Hadiaty (MZB) who guided me in the field and facilitated the permits. I am grateful to M. Kottelat and H. H. Tan (ZRC) for the insightful discussion and constructive criticisms on the taxonomy of *Rasbora*. I acknowledge the following at USNM for their invaluable assistance: J. Clayton, J. Williams, J. Finan, S. Raredon, and D. Cole (base map). I thank the following for access to material under their care: M. Sabaj Pérez (ANSP), R. Britz, O. Crimmen and P. Campbell (BMNH), M. Kottelat (CMK), K. Grossenbacher and S. Hertwig (NMBE), R. de Ruiter and M. van Oijen (RMNH), R. Robins (UF), R. Vonk and H. Praagman (ZMA), and K. Lim and H. H. Tan (ZRC). I am grateful to D. Rudaya, N. M. Ray, D. Syahril, and V. Simanjuntak for their fieldwork assistance. The 2006 ichthyofaunal survey in northern Sumatra was financially supported by the Leonard P. Schultz Fund, Division of Fishes, USNM. This article constitutes part of my dissertation submitted in partial fulfillment of the degree of Doctor of Philosophy to the Department of Biological Sciences, the George Washington University.

Literature cited

- Brittan, M.R. (1954a) *A revision of the Indo-Malayan fresh-water fish genus Rasbora*. Monographs of the Institute of Science and Technology, Manila, 224 pp + 3 maps.
- Brittan, M.R. (1954b) The cyprinid fish genus *Rasbora* in Malaya. *The Bulletin of the Raffles Museum*, 25, 129–156.
- Conway, K.W. (2005) Monophyly of the genus *Boraras* (Teleostei: Cyprinidae). *Ichthyological Exploration of Freshwaters*, 16, 249–264.
- Eschmeyer, W.N. (Ed.) (2013) Catalog of Fishes electronic version. Available from: <http://research.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (accessed 11 November 2013)
- Froese, R. & Pauly, D. (Eds.) (2013) FishBase, World Wide Web electronic publication. Available from: <http://www.fishbase.org> (accessed 1 October 2013)
- Hadiaty, R.K. (2005) Keanekaragaman jenis ikan di Suaq Balimbing dan Ketambe, Taman Nasional Gunung Leuser, Provinsi Nanggroe Aceh Darussalam. *Jurnal Biologi Indonesia*, 3, 379–388.
- Kottelat, M. (1986) A review of the nominal species of fishes described by G. Tirant. *Nouvelles Archives du Muséum d'Histoire Naturelle de Lyon*, 24, 5–24.
- Kottelat, M. (2001) Fishes of Laos. Wildlife Heritage Trust, Colombo, 198 pp.
- Kottelat, M. (2005) *Rasbora notura*, a new species of cyprinid fish from the Malay Peninsula. *Ichthyological Exploration of Freshwaters*, 16, 265–270.
- Kottelat, M., Whitten, A.J., Kartikasari, S.N. & Wirjoatmodjo, S. (1993) *Freshwater fishes of Western Indonesia and Sulawesi*. Periplus, Hong Kong, 259 pp + 84 plates.
- Kottelat, M. & Vidthayanon, C. (1993) *Boraras micros*, a new genus and species of minute freshwater fish from Thailand (Teleostei: Cyprinidae). *Ichthyological Exploration of Freshwaters*, 4, 161–176.
- Kottelat, M. & Lim, K.K. (1995) Freshwater fishes of Sarawak and Brunei Darussalam: a preliminary annotated check-list. *The Sarawak Museum Journal (New Series)*, 48, 227–256.
- Kottelat, M. & Tan, H.H. (2011) *Rasbora atranus*, a new species of fish from central Borneo (Teleostei: Cyprinidae). *Ichthyological Exploration of Freshwaters*, 22, 215–220.
- Kottelat, M. & Tan, H.H. (2012) *Rasbora cryptica*, a new species of fish from Sarawak, Borneo (Teleostei: Cyprinidae). *Ichthyological Exploration of Freshwaters*, 23, 37–44.
- Liao, T.Y., Kullander, S.O. & Fang, F. (2010) Phylogenetic analysis of the genus *Rasbora* (Teleostei: Cyprinidae). *Zoologica Scripta*, 39, 155–176.
<http://dx.doi.org/10.1111/j.1463-6409.2009.00409.x>
- Liao, T.Y., Kullander, S.O. & Fang, F. (2011) Phylogenetic position of rasborin cyprinids and monophyly of major lineages among the Danioninae, based on morphological characters (Cypriniformes: Cyprinidae). *Journal of Zoological Systematics and Evolutionary Research*, 49, 224–232.

<http://dx.doi.org/10.1111/j.1439-0469.2011.00621.x>

- Lumbantobing, D.N. (2010) Four new species of the *Rasbora trifasciata*-group (Teleostei: Cyprinidae) from Northwestern Sumatra, Indonesia. *Copeia*, 2010, 644–670.
<http://dx.doi.org/10.1643/ci-09-155>
- Ott, G. (2009) Redescription of *Homaloptera ripleyi* (Fowler, 1940) from Sumatra, Indonesia (Teleostei: Balitoridae). *Bulletin of Fish Biology*, 11, 73–86.
- Sabaj Pérez, M.H. (Ed.) (2012) Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an Online Reference. Version 3.0. American Society of Ichthyologists and Herpetologists, Washington, DC. Available from: <http://www.asih.org/> (23 February 2012)
- Siebert, D.J. & Guiry, S. (1996) *Rasbora johanna*e (Teleostei: Cyprinidae), a new species of the *R. trifasciata*-complex from Kalimantan, Indonesia. *Cybium*, 20, 395–404.
- Siebert, D.J. & Richardson, P.J. (1997) *Rasbora laticlavia*, a new cyprinid from Kalimantan, Indonesia, and lectotype designation for *R. vaillantii*. *Ichthyological Exploration of Freshwaters*, 8, 89–95.
- Springer, V.G. & Allen, G.R. (2004) *Ecsenius caeruliventris* and *E. shirleyae*, two new species of bleniid fishes from Indonesia, and new distribution records for other species of *Ecsenius*. *Zootaxa*, 791, 1–12.
- Tan, H.H. & Kottelat, M. (2009) The fishes of the Batang Hari drainage, Sumatra, with description of six new species. *Ichthyological Exploration of Freshwaters*, 20, 13–69.
- Tang, K.L., Agnew, M.K., Hirt, M.V., Sado, T., Schneider, L.M., Freyhof, J., Sulaiman, Z., Swartz, E., Vidthayanon, C., Miya, M., Saitoh, K., Simons, A.M., Wood, R.M. & Mayden, R.L. (2010) Systematics of the subfamily Danioninae (Teleostei: Cypriniformes: Cyprinidae). *Molecular Phylogenetics and Evolution*, 57, 189–214.
<http://dx.doi.org/10.1016/j.ympev.2010.05.021>
- Weber, M. & de Beaufort, L.F. (1916) *The fishes of the Indo-Australian Archipelago. III. Ostariophysii: II Cyprinoidea, Apodes, Synbranchi*. E. J. Brill, Leiden, 404 pp.
- Wirjoatmodjo, S. (1987) The river ecosystem in the forest area at Ketambe, Gunung Leuser National Park, Aceh, Indonesia. *Archiv für Hydrobiologie-ergebnisse der Limnologie*, 28, 239–246.