

A key to the West Palaearctic genera of stoneflies (Plecoptera) in the larval stage

PETER ZWICK¹

Introduction

Plecoptera constitute a numerically and ecologically significant component in freshwater ecosystems, mainly in running waters of all sizes, all over the world. The fauna of the vast Holarctic Region is relatively uniform, and its stonefly families are essentially endemic to it, only some extend into the Oriental Region. The East Palaearctic and the Nearctic subregions are particularly similar, sharing the families Pteronarcyidae, Styloperlidae, and Peltoperlidae, and a number of genera and even single species that all lack from the West Palaearctic subregion. The wingless Scopuridae are endemic to Japan and Korea. Europe and the immediately adjacent parts of Asia and Palaearctic North Africa together are relatively distinct from the other subregions, not only by the absence of the beformentioned taxa but also by shared endemic genera.

Adult stonefly taxonomy is well advanced, the terrestrial imagines can reliably be identified, although in certain regions of the world and in some genera limitations remain. However, larvae are generally less well known than adults. Given the fact that many adults can be identified to species only by genital characters, a lastingly lesser taxonomic resolution must be anticipated in larvae, even after further study. In North America, excellent comprehensive literature on larval Plecoptera exists (STEWART & STARK 2002). The genera of the Russian Far East and Siberia have been treated synoptically, although in a less detailed way (ZHILTZOVA & TESLENKO 1997). There is a preliminary key to the genera of the incompletely studied Chinese fauna (HARPER 1994).

In Europe, there are several good regional generic keys to larvae in the relatively impoverished faunas in the North and in the center of the continent. Central Europe holds an outstanding position through the work of the late JAROSLAV RAUSER (1980), who published a key to the species from former Czechoslovakia. However, all these keys are of limited use in many parts of South Europe with much more diverse faunas. These include a number of endemic genera whose larvae have been described, but not all in a comparative way, nor were they distinguished in keys.

The present paper keys all Plecoptera genera of the West Palaearctic Region. Brief comments on each genus are included. Where appropriate, suggestions or critical remarks on specific identifications of central European species are added, especially for species included in official lists of indicator organisms. There are also 2 appendices dealing with the German species of genus *Nemoura* and with selected species level taxa in genus *Leuctra*, respectively.

Geographic Range

West Palaearctic Region (i.e., Europe eastwards to Ural Mts and western shore of Caspian Sea; Anatolia, Caucasus, Lebanon and areas to the South, including Israel and Africa North of Sahara). Plecoptera are unknown from Libya and areas to the East, as well as from the Arabian Peninsula.

¹ Correspondence: Dr. PETER ZWICK, Schwarzer Stock 9, 36110 Schlitz, Germany.

Included Taxa

All genera in the above range. In genera occurring also outside the West Palaearctic Region the key is designed for West Palaearctic species, exotic species of the same genera cannot necessarily be identified with the present key. All of the West-Palaearctic families of Plecoptera belong in the suborder Arctoperlaria.

For monotypic genera, the single species is always named. For other genera, the attempt is occasionally made to identify selected species by characters listed in the text accompanying the genus.

Identifiable life stages

Characteristic character expressions of larvae develop stepwise, last instar larvae with fully formed wing-pads are optimal study objects. However, much younger larvae can also be identified with the present key but no larvae of less than *ca* 3 mm bodylength, or even first instars.

Counting of body segments

Segments of the trunk are counted from front to rear, parts of appendages from base to tip. Mind that of the 1st abdominal segment only the tergite is visible, the 1st sternite being fused to the metasternum. The first free, visible abdominal sternite is actually of segment 2. The female gonopore develops on sternite 8; its attachment scar moves backward at molts until it reaches the rear margin, interrupting the terminal hair fringe in its middle. The male gonopore is located at the end of sternite 9 but barely noticeable.

Anal gills

A collective term for gill filaments arising from the soft medial faces of paraprocts. Fine whitish filaments protruding out of the anus are no gills but gut fungi of the group Harpellales, which are common in Plecoptera. If one seizes them with forceps they can easily be pulled out of the rectum.

Nomenclature and systematic arrangement

Nomenclature follows the 4th edition of the International Code for Zoological Nomenclature (1999). The world catalogues by ILLIES (1966) and ZWICK (1973) were used as base, Recent literature, year 2002 included, was added. The system is that of ZWICK (2000).

Illustrations

Illustrations are originals by P. ZWICK, except when differently indicated. Illustrations were drawn at a variety of magnifications, from whole specimens or mounted parts, in direct or transmitted light. Illustrations are not generally to scale, absolute measurements are given where needed. Overall body length (BL) of fully grown larvae is indicated under each genus.

"Hot-keys"

A few taxa of various taxonomic rank can immediately be recognized, by reference to single key characters:

- Gill tufts on thoracic segments: Perlidae.
- 4 branched cervical gills: *Amphinemura*.
- 6 tubuliform or sausage-shaped cervical gills: *Protonemura*.
- 1 three-segmented telescoping gill filament on the inner face of each coxa; abdominal tergites with unpaired projections: *Taeniopteryx*.
- Maxillary palpus with asymmetrically inserted tiny needle-like terminal segment: Chloroperlidae.
- Head trapezoidal, extremely setose: *Isoptena*.
- Thoracic dorsum shield-like, with wide parallel wing-pads, setose; very small: *Capnopsis*.

Superfamilies

- 1 A transverse suture across upper face of head, between antennae and labrum (*L*), separating clypeus (*C*) from frons (*F*) (Fig. 1a). Labrum distinctly less than twice as wide as long. Mandibles plump, inner face with distinct mola (Fig. 2a); in lateral view, mandibles not even twice as long as high at the basal articulation. Maxillary palpi short and stout (Fig. 1a). Glossae and paraglossae subequal (Figs 5b, d). Coxae close together, thorax sternites between coxae small, the area between furcal pits and spina much smaller than a coxa (Figs 4a, b). Last tarsal segment at most as long as two basal segments together. The two basal segments differ between families but are never of equal length and structure (Figs 3b-d)Nemouroidea
- 1' No transverse suture crossing upper face of head between antennae and labrum (*L*), i.e. frontoclypeus (*F+C*) undivided (Fig. 1b). Labrum transverse, more than twice as wide as long. Mandibles flat and slender, without mola (Fig. 2b); in lateral view, mandibles are several times longer than high at the basal insertion; maxillary palpus long and slender (Fig. 1b). Glossae much shorter than paraglossae (Figs 6b, 7b). Coxae wide apart, thorax sternites between coxae extended, the area between furcal pits and spina about the size of a coxa (Figs 21b, c). Last tarsal segment about twice as long as two basal segments together; the latter are of equal size and very short (Fig. 3a) . Perloidea




Fig. 1. Head, oblique anterior view. Left antenna removed, antennal foramen striate, left mandible stippled: *Brachyptera* sp. (a), *Isoptera* sp. (b). C = Clypeus, F = frons, L = labrum, Mx = maxilla.




Fig. 2. Ventral view of left mandible of *Nemoura marginata* PICTET, 1836 and *Arcynopteryx compacta* (MCLACHLAN, 1872) (with enlarged apex of mandible).




Fig. 3. Hind tarsus of *Besdolus imhoffi* (PICTET, 1841) (a), *Rhabdiopteryx alpina* (KÜHTREIBER, 1934) (b), *Nemoura uncinata* (DESPAX, 1934) (c) and *Tyrrhenoleuctra zavattarii* (CONSIGLIO, 1956) (d).

Families of Nemouroidea

Conventional keys rely mainly on the orientation of wing-pads. In Plecoptera, first small rudiments of wing-pads appear in the antepenultimate instar (ZWICK 2003), the definite shape is attained at subsequent molts. Using wing-pad position, reliable identifications are possible only for specimens in the two last instars. However, much younger larvae can be identified by less obvious characters; the present key uses position of wing-pads only as an auxiliary character.

- 1 All tarsal segments elongate, increasing in length from 1 to 3. Segment 1 deeply excavated on upper face, receiving base of segment 2. On its lower face, segment 1 projects under 2 like a sole (Fig. 3b). Taeniopterygidae
 Habitus very different between subfamilies. Lower face of head resembles Nemouridae.
- 1' Tarsal segment 2 much shorter than segments 1 or 3. Segment 1 is tubular, with segment 2 attached at its end. Segment 2 is dorsally excavated to receive the base of segment 3. Ventrally, segment 2 projects under 3 like a sole (Figs 3c, d) 2

- 2 Stout; extended hind legs reach to tip of abdomen or beyond; wing-pads diverge obliquely from body axis (Fig. 5a); pilosity variable but always distinct. Lower face of head: bases of maxillae exposed, between them is the small soft plate of the mentum. On underside of thorax, only trochantinus well defined and sclerotized. Furcal sternites not delimited, furcal pits are tiny openings at the bottom of a poorly defined groove which is difficult to recognize between the middle coxae. All coxae projecting much from sternal face, transversely extended towards body midline, in particular front and middle legs (Fig. 4a). Ventral face of coxae flat, unsclerotized, transparent. Cervical gills, or no gillsNemouridae




Fig. 4. Cross section through prothorax behind fore coxae, seen from behind (diagrammatic), only skeletal elements shown, internal ones exposed by the cut are in black. Left, *Protonemura auberti* ILLIES, 1954, only 4 of the 6 sausage-shaped gills (numbered) partly visible in background. Right, *Leuctra inermis* KEMPNY, 1899. cx, coxa; fa, furcal arm; fe, femur; pa, pleural arm; tr, trochanter.

- 2' More slender, extended hind legs do not attain end of abdomen. Wing-pads variably positioned, mostly parallel to body axis (Abb. 5c). Thoracic sternites distinctly sclerotized, furca- and spinasternum of mesothorax together forming a well-delimited Y-shaped area between the coxae. Furcal pits well visible. Coxae of normal shape, medially not extended and not strikingly projecting from sternal surface. Never with gills . 3
- 3 Mentum plate-like, extending far sideward and forward, largely covering bases of maxillae (Fig. 5d). Wing-pads always parallel to body axis (Fig. 5c) Leuctridae
- 3' Mentum small, base of labium elongate, mouthparts completely exposed (resembling Fig. 5b). Wing-pads parallel to body axis except in one single rare genus whose wing-pads diverge (like in Nemouridae); however, the present genus lacks hairs ... Capniidae




Fig. 5. Habitus of *Protonemura* sp. (a) and *Leuctra* sp. (c; from RAUŠER 1980). Head in ventral view, mentum stippled: *Capnia bifrons* (NEWMAN, 1839) (b) and *Leuctra nigra* (OLIVIER, 1811) (d).
Ca = Cardo, G = Galea, St = Stipes.

Families of Perloidea

- 1 Thorax with lateral tufts of finely branched gills. Head very wide, its sharp lateral margin conceals the basis of maxilla from above (Fig. 6, dorsal). From below, the maxillar base is largely covered by the enlarged postmentum, only cardo and stipes partly exposed (Fig. 6b). Prolonged front corners of postmentum (f) laterally embracing the base of labium. Paraglossae large, soft, cushion-like and swollen (Fig. 6, ventral)
 Perlidae
 Larvae of this family have a distinctive habitus: they appear flat, with laterally extended flat legs with long setal fringes. Wing-pads little apparent even in the last instar, rounded on the outside and medially, separated by a wide shallow V-shaped notch.




Fig. 6: *Dinocras cephalotes* (CURTIS, 1827), head in dorsal and ventral views; the cushion-like paraglossae stippled.
ca = cardo; f = front corner of postmentum; m = mentum; pm = postmentum; occ = occipital fold; st = stipes.

- 1' No gills, head and mouthparts of different structure 2

- 2 Head and mouthparts very large, together as wide or wider than prothorax. Maxillae much enlarged, their enormous bases visible from above, to the sides of the eyes (Fig. 7a), only partly covered by postmentum from below, leaving even the membranous articulation membranes exposed (Fig. 7b). Maxillary palpi slender, last segment of normal size, symmetrically inserted in middle of penultimate segment (Figs 1b, 7a, b). Long cerci. Wing-pads roughly parallel to body axis, straight on outside, medially widely separated by U-shaped notch Perlodidae
- 2' Head normal, at most as wide as prothorax, mouthparts not enlarged, bases of maxillae hidden under head capsule. Terminal segment of maxillary palpus small, peg-like, asymmetrically inserted on penultimate segment (Figs 7c, 23a, 24a, 25e). Cerci short, distinctly conical, especially in small individuals. Wing-pads rounded on outside, medially separated by narrow V-shaped cleft (Fig. 25e) Chloroperlidae




Fig. 7. a, *Perlodes dispar* RAMBUR, 1842, head and thorax in dorsal views; cardo, stipes and lacinia of maxillae stippled; antennae removed, foramina striate. b, *Arcynopteryx compacta* (MCLACHLAN, 1872), head in slightly oblique ventral view, only right eye and mandible (black) partly visible, only basal segment of labial palpus shown; cardo and stipes of maxillae stippled. c, *Siphonoperla torrentium* (PICTET, 1841), head in dorsal view; mxp = reduced, asymmetrically inserted terminal segment at tip of maxillary palpus. sg = submental gill.

Genera of Taeniopterygidae

Genus *Oemopteryx* KLAPÁLEK is not included. A single species, *O. loewii* (ALBARDA, 1899) once occurred in large European streams; it is missing for about 100 years. A few females are preserved in museums, but no single male. The larva was never found. Larvae of the 4 North American species of *Oemopteryx* differ between themselves, inferences on *O. loewii* are therefore impossible.

- 1 A telescoping 3-segmented retractable gill protruding from inner face of each coxa (Fig. 8a). Long-legged larvae lacking hair fringes and resembling Nemouridae except that they possess unpaired processes on adominal tergites (Figs 8b, c). In most species, entire abdominal dorsum appears serrate, sometimes also thorax, especially pronotum, with prominent processes. Sternite 9 normal, paraprocts freely visible
 *Taeniopteryx* PICTET
 BL up to 13 mm. The only genus of subfamily Taeniopteryginae, unmistakable by the coxal gills. *Taeniopteryx* has a disjunct North Atlantic distribution. The numerous North American species lack tergal processes. Most of the approximately 10 Palaearctic species occur in South Europe, several of them also in central European mountains. In potamal streams of central European lowlands only *T. nebulosa* (LINNÉ, 1758) is found. It ranges from South to North Europe and East to central Asia. *T. nebulosa* can be identified by missing tergal processes on segments 8 and 9 (Fig. 8c) which the other Palaearctic species have (Fig. 8b). *T. araneoides* K LAPÁLEK, 1902 (wingless; Elbe, Danube) not found for over 100 years, larva unknown.




Fig. 8. a, *Taeniopteryx nebulosa* (LINNÉ, 1758), oblique ventro-anterior view of middle thorax, with 3-segmented telescoping gill originating from coxa. b, *T. hubaulti* AUBERT, 1946, lateral view, contour (after AUBERT, 1950). c, *T. nebulosa*, contour of abdomen, lateral view (after ILLIES, 1955).
 cx, coxa; fe, femur; fu, furcal pit; pr, pleural ridge; sp, spina; tr, trochanter; tt, trochantinus.

- 1' Gill-less sprawlers usually clinging to stones in fast current. Dorsal profile smooth, not serrate, hair fringes on legs present; ventral face soft, pale, little sclerotized. Upon disturbance larvae roll up which is often also seen in preserved specimens (Fig. 9). Sternite 9 strikingly enlarged, covering paraprocts (Fig. 10c) (Brachypterainae) 2




Fig. 9. Rolled larva of *Brachyptera* sp. (from ILLIES, 1955).

- 2 Cercus base with sparse erect hair fringe. Marginal setae of abdominal tergites very short but distinct in profile. Dorsal face with dark pigment, venter pale. Behind ocelli, branches of epicranial suture connected at a very blunt angle (Fig. 10a)² *Brachyptera* NEWPORT
 BL up to 12 mm. Endemic West Palaearctic genus. *B. risi* (MORTON, 1896) occurs in all of Europe and is the only Scandinavian species. In central European mountain streams *B. seticornis* (KLAPÁLEK, 1902) is the commonest species. Several species from potamal stream sections that had disappeared for decades were recently again found in a few places. Many species are endemic in the Mediterranean Region. The literature suggests specific colour patterns but there are examples of striking variation. Last instars can reliably be identified by structural details of sternite 9 and paraprocts illustrated by ILLIES (1955; Fig. 10c) and AUBERT (1959); allegedly distinctive colour patterns to be viewed with caution!
- 2' Cercus without hair fringe. Setae along distal tergal margins barely projecting from insertion points, almost invisible even in profile. Upper and lower body face equally pale, barely pigmented. Branches of occipital suture meet at a more acute angle (Fig. 10b) *Rhabdiopteryx* KLAPÁLEK
 BL up to 12 mm. Several species, even in central Europe; *R. alpina* KÜHTREIBER, 1934 and *R. neglecta* (ALBARDA, 1889) in the Alps are most frequently found.




Fig. 10. Dorsal view of heads of *Brachyptera seticornis* (KLAPÁLEK, 1902) (a) and *Rhabdiopteryx neglecta* (ALBARDA, 1889) (b); note angle at which branches of occipital suture meet. *Brachyptera* spp., ventral view of abdominal tip with postgenital plate of males (top) and females (bottom) (c); from ILLIES, 1955)

Genera of Nemouridae

The present key works for larvae of near 2 mm body length or larger. The two genera with cervical gills do not yet have these in the first instar and the characteristic number and shape of gills develop stepwise, at early molts. *Nemurella* is in the literature normally diagnosed by a characteristic length relation of tarsal segments; however, this relation develops only late in larval life.

- 1 With cervical gills (Amphinemurinae) 2
 1' No gills (Nemourinae) 3

² The larvae of some recently described Southern European species of *Rhabdiopteryx* are unknown, it cannot presently be tested whether the difference holds for all. However, according to AUBERT (1959), the Alpine *B. trifasciata* (PICTET, 1832) resembles genus *Rhabdiopteryx*. *B. trifasciata* occurred abundantly in Alpine rivers, However, it was entirely missing during the last 50 years but was recently again found, at few sites and in low numbers. I have seen no larvae of *B. trifasciata*.

- 2 Six sausage-shaped or tubiform cervical gills; in some species each gill has a prominent subapical constriction (Fig. 11a) *Protonemura* RIS
BL up to 10 mm. Numerous species, many Mediterranean endemics. With some practice and sufficient comparative material, central European larvae can rather reliably be identified with the key by RAUŠER (1956, also 1980).
- 2' Four branched, brush-shaped cervical gills (Fig. 11b) *Amphinemura* RIS
BL close to 7 mm. According to the (partly contradictory) literature (ILLIES 1955, RAUŠER 1963) larvae of the common central European species, *A. standfussi* RIS, 1902, *A. sulcicollis* (STEPHENS, 1835) and *A. triangularis* RIS, 1902 can be identified by cercal and femoral setation. Study of pharate males did not confirm these claims. Identification of *A. borealis* (MORTON) (after RAUŠER 1963, 1980) was not checked.




Fig. 11. *Protonemura* sp. (a) and *Amphinemura* sp. (b), ventral view of prothorax and neck showing sausage-shaped and branched gills, respectively (from RAUŠER, 1980).

- 3 Very long-legged: end of hind tibia lies clearly on coxa when folded back. Ground pilosity of femora weak, but several spreading setae in distal third exceed femur width. In late instars 1st and 3rd tarsal segments of hind legs about equal (Fig. 12c). Body normally held in a characteristic posture, with upcurved abomen and arched long antennae and cerci (Fig. 12a). Late instar male unmistakable, with enlarged upcurved paraprocts (Fig. 12b). Marginal setation of tergites consisting of only 4 large setae, others not larger than weak intercalary pilosity³ (Fig. 12d). Lateral setae longer and stronger than paramedian ones, setae on tergite 9 strongest. Setae along tibiae long, several longer than tibial width, exceptionally long erect sensory hairs, especially in small larvae. Occiput often strikingly lighter than dark purplish central portion of head. Setation of cerci sparse but strong and spreading, most developed and distinctly longer than segments around segments 5-7; distal setae fine and increasingly short. Small larvae (Fig. 12e) best identified by cerci attaining body length, unique long setae on several basal flagellar segments (Fig. 12g; disappear in larger specimens), long and strong pilosity of legs, and details of tergal and cercal setation described above

..... *Nemurella* KLAPÁLEK
Only *Nemurella pictetii* KLAPÁLEK, 1900. BL up to 11 mm. From northern Spain to central Siberia. Common at sites with few other Nemouridae, especially in swampy springs, upper courses with diffusely entering groundwater, outlets of drain water, spring basins etc., but also in strongly acid waters like outlets of peat bogs. Occasionally said to be ubiquitous, but not resistant against pollution. Voltinism varies; most central European populations are bi- to trivoltine, univoltine in cool mountain sites, North European mountain populations semivoltine.

³ The term "intercalary" designates pilosity on the surface of a sclerite, not at its edge, for example on the surface of tergites or on the surface of a cercus segment, between base and apical setal ring.

- 3' Body shape, length of legs, shape of cerci and setation very different between the numerous species; 1st tarsal segment always shorter than last. Occiput of central European species not distinctly paler than middle of head. If legs are long and slender, they lack very large spreading femoral spines. Tibiae usually with dense and strong setae along outer edges, but setae not longer than tibial width. Large setae different from intercalary tergal setation along tergal margins more numerous. Male paraprocts short, unmodified. Antennae never with long setae near base of flagellum
.....*Nemoura* LATREILLE
BL mostly 8-9 mm, some species up to 11 mm. Many species stout and more plump than *Nemurella*, cerci usually straight and shorter. See Appendix I for a preliminary key to German species.




Fig. 12. *Nemurella pictetii* KLAPÁLEK, 1900. Characteristic curved posture (a; only right appendages shown), paraprocts of last instar male (b; lateral view), hind tarsus (c) and tergite 5 (d) of last instar specimen. Small larva, habitus (e), hind tarsus (f) and base of antennal flagellum (g). Figures without scale are not to scale.

Genera of Leuctridae

- 1 Abdominal segments 5 ff ring-shaped, no pleural membrane *Leuctra* STEPHENS
BL up to 12 mm or even more, but most species are less than 10 mm long. The genus occurs on both sides of the Atlantic Ocean, in North America as well as in the West Palaearctic. One single European species, *L. fusca* (LINNÉ, 1758), extends across Siberia into the Russian Far East. Otherwise, the Easternmost outlying representative is *L. kopetdaghi* ZHILTOVA, 1972 (Kopetdagh, Iran, Libanon). Males and females of most of the over 100 European species can be well distinguished (for example, RAVIZZA & VINÇON 1999, VINÇON & RAVIZZA 2001). In contrast, larval identifications are not consistently possible. Body pilosity very variable between species, from missing to very dense and rough. See Appendix II for a few central European species and Operational Taxonomic Units that can be reliably recognized.
- 1' Pleural membranes divide also abdominal segments 5-8 into tergites and sternites 2
- 2 Resembling *Leuctra*, head rounded, pronotum normal, pilosity fine, short, inconspicuous *Tyrrhenoleuctra* CONSIGLIO
BL up to 7 mm. West Mediterranean, on fragments of the former Tyrrhenian land mass: North Africa, Iberian Peninsula, Balears, Sardinia and Corsica. Limits between the 3 nominal species doubtful, despite studies applying gel electrophoresis (SEZZI 2001).
- 2' Head trapezoidal, widest posteriorly, narrowed in almost straight line towards front (Fig. 13a). Clypeus with prominent anterior corners. Pronotum transverse, ca. 1.5 times wider than long. Long erect hairs along sides of body and legs. Antennae with normal base, rapidly thinning apically, distally very fine and slender. Cerci multi-segmented but thin, thread-like, at base thinner than antennae (Fig. 13), miserable in comparison with large (up to 11,5 mm long) massive body *Pachyleuctra* DESPAX
BL up to 11 mm. *P. benllochi* (NAVÁS, 1917) and two closely related species in the Pyrenees. Through their pilosity, half grown specimens greatly resemble *Leuctra nigra* (OLIVIER, 1811) and *L. braueri* KEMPNY, 1898; the latter has also angular front corners of clypeus. Both *Leuctra* species have normally shaped roundish heads (Appendix II, Fig. 34), and in both, pronotum, antennae and cerci are average.




Fig. 13. *Pachyleuctra* sp., head and cercus (with segment numbers) of last instar larva; all to same scale.

Genera of Capniidae

- 1 Upper face with long erect setae. Pronotum approximately heart-shaped, front much wider than rear. Abdominal segments 1-5 completely, 6 partly divided by pleural membrane. Anterior wing-pads much expanded towards median line, their medial edges closely opposed, leaving only a narrow cleft between them. Mandibles with exceptional setal pocket just in front of apical teeth and labrum with very dense and short setal seam resembling the ones found in scrapers, like *Brachyptera* spp. Brownish larva (Fig. 14) *Capnopsis* MORTON
BL only 5-7 mm. The only species, *C. schilleri* (ROSTOCK, 1892) ranges from the Caucasus to western North Africa and to the polar North of Europe. Several geographical subspecies exist; only the nominate subspecies occurs in central and North Europe. One of the smallest European Plecoptera, not often found. Larvae live concealed in loose accumulations of detritus, details of habitat preference not known.
- 1' Pilosity inconspicuous, at most individual long hairs, larvae sometimes appearing completely bare. Abdominal segments 1-8 divided by pleural membrane. Pale larvae 3
- 2 Wing-pads parallel to body axis. Antennae and cerci normal. Body with sparse long pilosity *Capnia* PICTET
BL up to 9 mm. Numerous species in Asia and North America, only a few in Europe. These belong to several distantly related species groups; nevertheless, specific larval identifications are difficult. Males of several species brachypterous to almost apterous. *C. bifrons* (NEWMAN, 1839), the most widespread of these sexually dimorphic species appears to comprise several presently unnamed biospecies (RUPPRECHT 1997).
- 2' Wing-pads diverge obliquely from body axis, thereby resembling Nemouridae. Antennae long and thin, base narrower than the plump cercus base. No macroscopic body pilosity *Capnioneura* RIS
BL up to 7 mm. Of the 12 named species in this genus, only two, *C. nemuroides* RIS, 1905 and *C. mitis* DESPAX, 1932 occur in central Europe but are seldom found; habitat preferences not known. The remaining species occur in the Mediterranean Region, from western North Africa to the Caucasus; several are completely apterous.




Fig. 14. *Capnopsis schilleri* (ROSTOCK, 1892), last instar larva, habitus (not to scale) and labrum in dorsal, left mandible in ventral view.

Genera of Perlidae

Among the West Palaearctic Plecoptera, the tufts of highly branched thoracic gills are characteristic. Each consists of 1 to 3 trunks which may fork into several branches each of which carries many straight fine filaments. These gills occur in characteristic positions to which the traditional gill nomenclature refers. It is followed here because it underlines homologies with similar gills observed also in other related families (within the infra-order Systellognatha). However, such examples do not exist among West Palaearctic representatives. The following gills are observed in Perlidae:

- a substigmatic gill below each of the 3 first spiracles;
- an anterior supracoxal gill above and in front of the fore coxa;
- a posterior supracoxal gill above the middle and hind coxa, behind the pleural fold.

- 1 Postmentum with transverse lines separating each front corner from main plate-like part (Fig. 6b); marginal setation of sternite 7 complete 2
- 1' Front corners continuous with main part of postmentum; marginal setation of sternite 7 medially incomplete, setal fringe divided⁴ 3
- 2 Transverse fold across occiput regularly curved, crossing unpaired median branch of occipital suture distinctly behind forking point of suture. Long, club-shaped setae stand in a single row along transverse fold (Fig. 15b), also laterally where the fold is curved down regularly, merging into the sharp lateral edge of head below the eyes. Abdominal tergites only with fine pilosity contrasting markedly with strong marginal spines (Fig. 15c). Small; 2 pale longitudinal bands on abdomen (Fig. 15a) *Eoperla* ILLIES
The only species is *Eoperla ochracea* (KOLBE, 1885), BL up to 17 mm. Strictly Mediterranean: North Africa, Spain, Southern France, Makedonia, Greece, Asia Minor. Absent from the Italian region.
- 2' Transverse occipital fold angled forward, medially almost touching the forking point of occipital suture (Fig. 6a). Setae along transverse fold short, barely projecting over fold when viewing intact specimens. Well visible setae occur only laterally behind eyes where fold enters into sharp lateral edge of head, and are irregularly placed instead of in an orderly row. In slide preparations of exuviae, the setae are better seen; they are stouter than in *Eoperla*, almost spatulate (Fig. 15d). Abdominal tergites with dense cover of short fairly strong spines that differ less strikingly from marginal spines (Fig. 15e). No pale longitudinal bands on abdomen *Dinocras* KLAPÁLEK
BL up to 35 mm. *D. cephalotes* (CURTIS, 1827) is widely distributed over most of Europe, males brachypterous. The very similar *D. ferreri* (PICTET, 1841) (restricted to the southern edge of the Alps and small parts of the northern Appennines) and *D. megacephala* (KLAPÁLEK, 1907) (Balkans, and locally in a strip along northern edge of Alps, to French Jura) have long-winged males. Allegedly distinctive colour patterns of larval *D. cephalotes* and *D. megacephala*, respectively (AUBERT 1946, 1959, ILLIES 1955) occur within pure populations of only *D. cephalotes*, apparently representing individual variation (A. DORN & A. WEINZIERL, personal communication); structural differences between larvae not known.
- 3 Transverse occipital fold bent angularly forward behind inner edge of eye and meeting the fold running around eye and to side of head at a distinct, forward-directed angle⁵. Sideways from this point, setation includes much longer pointed setae (Fig 16a, inset); cercus base never with seam of erect soft hairs 6
- 3' Transverse occipital fold and fold running around eye merging in a regularly curved or only slightly undulating line (Fig. 18); no striking change in setation at fusion point. Cercus base with (Fig. 17b) or without seam of erect fine hairs 4

⁴ Much used in the world generic key to larvae of Perlinae (SIVEC et al. 1988); however, may be difficult to assess when there are fine and much smaller setae in the centre of setal fringe VII than on the sides which is not exceptional.

⁵ The same is true of the closely related East Asian genus *Oyamia* KLAPÁLEK.




Fig. 15. *Eoperla ochracea* (KOLBE, 1885), habitus (a; modified from DESPAX, 1931). Setae along occipital fold (b) and detail of setation of tergite 5 (c). *Dinocras cephalotes* (CURTIS, 1827), setae along occipital fold (d) and detail of setation of tergite 5 (e). Figures b and d, and c and e, respectively, are to the same scale, respectively; mind that *Eoperla* is not even half the size of *Dinocras*!




Fig. 16. *Agnetina elegantula* K LAPÁLEK, 1905, habitus of fore body (a) and detail of occipital crest and its pilosity at bend behind left eye (specimen from Austria, Burgenland, Lafnitz); *Marthamea selysii* (PICTET, 1841), fore body of brachypterous last instar male from River Moselle (b).

- 4 Silky fine hairs on cercus form a dorsal fringe, especially on basal segments. Rings of dark spines striking already on basal cercus segments, not much longer on distal segments (Fig. 17a). Fine pilosity about half as long as segment, closely appressed to segment surface. Most species bright yellow with contrasting dark brown pattern 5
- 4' Cerci without dorsal hair fringe but semi-erect fine pilosity forms dense hair whirls all around cercus segments, especially medially; many hairs longer than corresponding segment (Fig. 17b). Rings of spines missing from basal cercus segments, appearing only from about segment 8, blond, much shorter than fine hairs, inconspicuous; spines grow progressively longer on distal segments, eventually attaining half segment length. Light brownish with indistinct pattern on head and thorax, abdominal tergites with pairs of light spots *Helenoperla* SIVEC
Larvae of the single species, *H. malickyi* SIVEC, 1997 are up to 27 mm long. The discovery of this new genus was a great surprise; the insect is presently known only from two rivers in Epirus, Greece.




Fig. 17: Right cercus (dorsal side to the right), segments 6-8, of *Perla burmeisteriana* CLAASSEN, 1936 (a) and *Helenoperla malickyi* SIVEC, 1997 (b); not to scale.

- 5 Setation along occipital fold extends only to behind eye but not forward along edge of flat expanded side of head. No wavy section of occipital fold near inner corner of eye. Mesothoracic supracoxal gill with single or double trunk, depending on species

..... *Perla* GEOFFROY
BL up to 40 mm. *Perla* is endemic to the West Palaearctic Region. Species occur from North Africa and Great Britain to Iran; Cyprus is the only Mediterranean island with a *Perla* species (*P. caucasica* GUÉRIN, 1838). The genus does not inhabit North Europe. Species identities seemed to be well sorted out until SIVEC & STARK (2002) showed convincingly that species taxonomy is chaotic. There seem to be many more species than presently recognized. The authors studied only the eggs, other life stages can presently not be identified to the same degree. Therefore, specific identifications in *Perla* are presently preliminary and as of *status quo*.

Provided the study area is rigidly restricted to Germany ⁶, *P. marginata* (PANZER, 1799) can be recognized by lack of gills on paraprocts (so-called anal gills); head pattern distinctive (Fig. 18b). *P. burmeisteriana* CLAASSEN, 1936 has stout femora (hind femora about 3 times as long as wide), head pattern also distinctive (Fig. 18c). The long-legged (hind femur ca. 4 times as long as wide) Alpine larvae would (according to SIVEC & STARK 2002) all belong to a species presently called *P. grandis* RAMBUR, 1842.

⁶ The polytypic *P. pallida* GUÉRIN, 1838 (Slovenia and eastward) and *P. madritensis* RAMBUR, 1842 (Iberian Peninsula) are siblings of *P. marginata* and easily confused with it (SIVEC & STARK, 2002). *Perla carantana* SIVEC & GRAF, 2003 from Austria and Slovenia is indistinguishable from *P. burmeisteriana*, except in the characteristic egg stage (SIVEC & GRAF, 2003)!

- 5' Setation along occipital fold continues around eye and extends far forward onto expanded flat side of head, laterally in front of eye. Occipital fold just inside from each eye distinctly wavy. Mesothoracic supracoxal gill consisting of two separate main trunks, each with many filaments *Paragnetina* KLAPÁLEK
 The only West Palaearctic species, the Caucasian-Anatolian *P. spinulifera* (ZHILTZOVA, 1967) was originally described as a separate genus, *Caucasoperla* ZHILTZOVA, 1967 but was later synonymized with *Paragnetina*. The presumed larva (conspecificity unconfirmed; BL of the only known specimen 24 mm) described by Zwick (1971; Fig. 18a) was compared with Asian *P. flavotincta* (MCLACHLAN, 1872) and Nearctic *P. immarginata* (SAY, 1823) which it resembles structurally.




Fig. 18. *Paragnetina spinulifera* ZHILTZOVA, 1967, fore body (a; from ZWICK, 1971); dorsal view of head, *Perla marginata* (PANZER, 1799) (b) and *P. burmeisteriana* CLAASSEN, 1936 (c); not to scale.

- 6 Center of meso- and metanotum with a complex brown pattern consisting mainly of a median band and mediolateral curved bands which meet it posteriorly (Fig.16a). Dark pattern on head between bases of antennae forming a broad dark band in front of median ocellus that extends far forward; the band includes 3 pale spots (Fig.16a).....
 *Aagnetina* KLAPÁLEK
 BL up to 25 mm. The genus was for a long time known by the name *Phasganophora* KLAPÁLEK; see ZWICK (1984) for the confused history and the identity of the 3 West-Palaearctic species. Unfortunately, upon discovery (SCHOENEMUND 1925) the larva of *A. elegantula* was mistaken for *Marthamea vitripennis*; for a long time this severely hindered distinction of the two genera. Only *A. elegantula* KLAPÁLEK, 1905 occurs in central Europe. It is very rare and known from very few central European localities (GRAF, 1997). *A. senilis* KLAPÁLEK, 1921 and *K. weneri* (KEMPNY, 1908) are poorly known Caucasian and Anatolian species, respectively. The dark brown dorsal pattern on yellow ground of *A. elegantula* resembles the American *A. capitata* (PICTET, 1841) much. Figures of the habitus of *A. elegantula* are in SCHOENEMUND (1925; as *Marthamea vitripennis*) and GRAF (1997; photography).

- 6' Meso- and metanotum with a brown median band in a wide pale area. Dark pattern between bases of antennae medially interrupted in front so that a pale area remains in front of anterior ocellus (Fig. 16b) *Marthamea* KLAPÁLEK BL up to 20 mm. *M. vitripennis* (BURMEISTER, 1839) occurred in rivers of central and Southeast Europe but is today largely extinct. *M. selysii* (PICTET, 1841) used to occur in the River Moselle, a complete habitus figure of a specimen is in SCHOENEMUND (1925). *M. selysii* is a West European species which was recently collected at several localities in Spain. Specific identifications of *Marthamea* larvae are presently impossible; mind that the distinctly different larva illustrated as *Marthamea vitripennis* by SCHOENEMUND (1925) is actually *Agnentina elegantula*.

Genera of Perlodidae

- 1 Paraprocts pointed. Fine dorsal hair fringe lacking from base of cerci. Distally, hair fringe not longer than whorls of setae around tips of cercal segments (Figs 19a,b). Fine procumbent black pilosity closely appressed to body surface, resembling delicate dark lines on integument. Lacinia wide, with 2 apical teeth and along inner edge with a row of strong spine-like setae which grow successively finer and shorter towards base (Fig. 20a). Transverse setal row across occiput indistinct, irregular and medially more or less widely interrupted. Only first 2 abdominal segments divided into tergite and sternite *Isoperla* BANKS BL up to 17 mm. Holarctic genus. The numerous European species are best distinguished by male genitalia and drumming signals. Identification of larvae with existing keys (ILLIES 1952, 1955; RAUŠER 1980) which rely on numbers of setae along lacinia fails because the numbers depend on body size, i.e., change during growth and differ between sexes. KONAR (2000) claims easy and safe species identification is possible in Carinthia. He relies on a combination of dorsal colour patterns with shape of tergal setae (blunt as opposed to pointed).
- 1' Paraprocts blunt, their sclerotised portion apically surrounded by soft, membranous strip; dorsal cercal hair fringe long, distinct, also at base (Figs 19c, d). Body surface bare or with different kinds of pilosity; fine procumbent hairs as seen in *Isoperla* occur only in a single Spanish species with narrow unidentate lacinia. Occipital setation variable. Variable number of abdominal segments divided into tergite and sternite 2




Fig. 19. *Isoperla* sp. (a, b) and *Dictyogenus alpinum* (PICTET, 1841) (c, d), abdominal tip of female larvae in ventral view, and lateral view of detached left paraproct and cercus. In the figure of *Dictyogenus*, stippling identifies sclerotized parts of paraprocts; the strongly bulging anal membrane is concealing both edge and setal fringe of tergite 10.

- 2 Lacinia narrow, claw-like, unidentate, lacking subterminal tooth (Fig. 20b); endemic to western part of Mediterranean area 3
- 2' Lacinia with 2 unequal tips, basally from these usually with some setae along medial edge (for example, Figs 20c, 22a, e) 4
- 3 Head and prothorax shiny, completely hairless *Afroperlodes* MIRON et ZWICK
Afroperlodes lecerfi (NAVÁS, 1929), the only species, occurs only in North Africa. BL up to 17 mm.
- 3' Head and prothorax with fine procumbent pilosity, like in *Isoperla*
..... *Hemimelaena* KLAPÁLEK
The single species, *Hemimelaena flaviventris* (PICTET, 1841), is endemic to the Iberian Peninsula, living in intermittent streams; not in the Pyrenees. BL up to 14 mm.
- 4 No transverse row of setae across occiput, between rear margins of eyes (Fig. 7a). Lacinia with 2 closely appressed teeth, from there enlarged in a curved line towards bulging base. Only occasional setae along inner edge of lacinia basally from apical teeth (Fig. 20d) 5
- 4' An (often irregular or incomplete) row of setae across occiput, between rear margins of eyes (Fig. 21a). Body surface more or less densely pilose or setose. Lacinia variable, apical teeth diverge, setation along inner edge distinct (for example, Figs 20c, 22a, e) 6
- 5 Abdominal segments 1-4 divided into tergites and sternites by hairless pleural membranes. Segments 5 and following are sclerotized rings with continuous setal fringe along rear edge. Inner margin of lacinia basally from subapical tooth smooth, no angle or knob. Body surface bare or with fine pilosity, no interspersed short strong spines. Front and rear edge of pronotum (but not the sides!) and rear edge of abdominal segments with marginal setae *Perlodes* BANKS
BL up to 26 mm. Europe, Caucasus, Asia Minor. Several species, best distinguished by structural details of their eggs. Presently, not all central European biospecies seem to be named or recognized. In central Europe, *P. microcephalus* (PICTET, 1833) and *P. dispar* RAMBUR, 1842 are widely distributed and common. Larvae are contrastively patterned in yellow and dark brown. Larvae of *P. intricatus* (PICTET, 1841) and *P. jurassicus* AUBERT, 1946 pale, pattern mainly olive green.
- 5' Only abdominal segments 1-3 divided into tergites and sternites by hairless pleural membranes, already segment 4 forming a sclerotized ring with continuous caudal setal fringe. Basally from apical lacinial tooth is a small knob with a group of small setae, or even a short row of relatively strong hairs *Filchneria* KLAPÁLEK
Filchneria is one of several closely related nominal Asian genera. One species, *Filchneria balcarica* (BALINSKY, 1950) occurs in the Caucasus; its larva is unknown. The present key is based on specimens of *F. mongolica* (KLAPÁLEK, 1903) and *F. amabilis* (JEWETT, 1958; from the Himalaya). Both have strong pilosity, distinct also along sides of pronotum. Differences in details between the two species preclude inferences on the Caucasian species. BL of the mentioned Asian species are around 25 mm.




Fig. 20. Left maxillae of several genera of Perlodidae.

- 6 On mesosternum, lines directed forward from spina meet front corners of furcal pits (Fig. 21c). Lower edge of outer mandibular tooth distinctly serrate (Fig. 2b); galea shorter than, or equal to, 3 basal segments of maxillary palpus (Fig. 22a); a short finger-shaped gill on either side of base of postmentum (Fig. 7b). Setae across occiput forming an irregular plurilinear band. Basal 3 abdominal segments divided into tergite and sternite *Arcynopteryx* KLAPÁLEK
 Several Asian species and the circumpolar *Arcynopteryx compacta* (MCLACHLAN, 1872), the only European species. Disjunct boreomontane distribution with pleistocene relict populations in central and even Southwest Europe (for example, Black Forest, Carpathians, Koralpe, Balkans, Pyrenees; not in main Alpine chain). BL up to 26 mm.
- 6' On mesosternum, lines directed forward from spina meet posterior corners of furcal pits (Fig. 21b). Lower edge of outer mandibular tooth smooth, or so finely serrate that this is easily overlooked (*Bulgaroperla*); galea longer than basal 3 palpus segments (Figs 22b, e); no gills. Only basal 2 abdominal segments divided, others ring-shaped 7
- 7 Setation across rear of head plurilinear only behind eyes, medially turning into an irregular single row ending far from middle. Setal fringe along inner edge of lacinia begins at a short distance from apical teeth (Figs 22b, f) 8
- 7' Row of setae across rear of head almost continuous, only narrowly divided in middle. Setal fringe along inner edge of wide lacinia begins immediately next to apical teeth (Figs 20c, 22c-e, g) 9
- 8 Distinct marginal setae present on pronotum and abdominal tergites. Marginal setation of lacinia begins on a small projection below subapical tooth; margin between teeth and projection concave and hairless (Fig. 22b) *Diura* BILLBERG
 Two circumpolar species occur in Europe: *D. nanseni* (KEMPNY, 1900) only in North Scandinavia, *D. bicaudata* (LINNÉ, 1758) additionally on the British Isles, plus relict populations in central European highlands North and East of the Alps, not in the Alps themselves. BL up to 25 mm.
- 8' Marginal setae on pronotum and abdominal tergites short, inconspicuous. Fine serration on outer cusp of lower mandibular tooth present. Lacinia without subterminal notch, row of setae along inner edge sparse, discontinuous, mainly on slightly bulging lacinial base (Fig. 22f) *Bulgaroperla* RAUŠER
 BL up to 23 mm. *Bulgaroperla mirabilis* RAUŠER, 1966 from the Eastern Balkans and the Caucasus.




Fig. 21. Head and ventral view of mesothorax of *Dictyogenus alpinum* (PICTET, 1841) (a, b) and ventral view of mesothorax of *Arcynopteryx compacta* (MCLACHLAN, 1872) (c). Density of stippling varies to indicate degree of sclerotisation on the largely soft ventral face of thorax. Fp = furcal pits.

- 9 Teeth of lacinia about half length of lacinia; inner margin of lacinia only with about 3 marginal setae close to subapical tooth (Fig. 22g).
 *Guadalgenus* STARK et GONZÁLEZ DEL TÁNAGO
 The single species, *G. franzi* (AUBERT, 1963), lives in intermittent streams on the Iberian Peninsula. BL up to 18 mm
- 9' Teeth of lacinia less large, about 1/3 lacinial length. Convex inner edge of lacinia with prominent marginal setation (Figs 20c, 22c, e) 10
- 10 Inner edge of lacinia below subapical tooth projecting, shoulder-like. In the same area, marginal setae form a small setal patch (Figs 22c, d) *Besdolus* RICKER
 BL up to 24 mm. *B. imhoffi* (PICTET, 1841) in the northern forefield of the Alps was very abundant for example in the Rhine at Basel around 1915. It disappeared and was missing for long, because of pollution. Recently, it reappeared in several larger streams and small rivers in South Germany and northern Austria. Together with it occurred another incompletely known species whose males were redescribed as *B. ventralis* (PICTET, 1841) but may actually be an unnamed species that is presently known only from Hungary. Additional species of *Besdolus* exist in the Balkans, in North Italy and in Spain. Setation differs much between species.
- 10' Inner edge of lacinia below subapical tooth less strongly projecting, marginal lacinial setae in a single long row (Figs 20c, 22e) 11
- 11 Long erect setae along front and rear edges of pronotum; marginal setae of abdominal tergites conspicuous, attaining 1/3 of segment length. Lacinia broad, contour slightly bulging (Fig. 22b) *Dictyogenus* KLAPÁLEK
 The distinction of the 2 nominate Alpine species (*D. alpinum* (PICTET, 1841), *D. fontium* (RIS, 1896)) is doubtful. BL up to 25 mm.
- 11' Entire circumference of pronotum with very short, easily overlooked marginal setae. Marginal setae of abdominal tergites equally inconspicuous and short, attaining less than 1/5 of segment length, except a single slightly longer paramedian seta on either side. Lacinia slender (Fig. 22e) *Isogenus* NEWMAN
 BL up to 21 mm. The only species, *I. nubecula* NEWMAN, 1833 used to be widespread and common in major European rivers, occurring "by the thousands" along the Seine in Paris (RAMBUR, 1842). It was still found in the Danube at Vienna, in 1949 (my coll.). Most contemporaneous records from central Europe are of misidentified larvae of *Isoperla*.




Fig. 22. Maxillae of several genera of Perlodidae; *Bulgaroperla* after BRAASCH & JOOST (1971).

Genera of Chloroperlidae

Family Chloroperlidae comprises two subfamilies. The 3 genera constituting the Paraperlinae are restricted to the East Asian mainland and to North America, while Chloroperlinae are of Holarctic distribution, with 5 genera in the West Palaearctic Region. Chloroperlinae are structurally very homogenous. While adults of the European species can easily and reliably be assigned to the various genera by genital characters, distinction of larvae (except *Isoptena*, which is globally unique and cannot be mistaken for any other stonefly larva, from the first instar on) is tricky, in all faunal regions.

The various European species within a given genus differ a little between them which renders generic distinctions difficult; for example. *S. montana* (PICTET, 1841) has resemblances with *Chloroperla*-species. At the same time, the various species are not sufficiently characteristic to be recognized at the specific level. In a regularly sampled population of *S. torrentium* (PICTET, 1841), ontogenetic change in all characters was found to be important, creating great problems. For example, not only segment number but also general shape and relative length of cerci change importantly during growth, from short with few segments and a strongly conical contour in early instars to relatively longer, with up to 16 segments and a much more slender overall shape in the last instar. Similarly, eyes of early instars are much smaller in relation to the head capsule than they are in later and especially in the last instar. Early instars

tend to be unicolorous and generally pale, but patterns may develop and be distinct in late instars; however, no comparative study of the patterning of the European *Siphonoperla* has so far been made. *Chloroperla* larvae tend to be generally pale.

The present key is based on study of last instar specimens. Users are warned to attempt to identify single young specimens. Only if much material collected at the same site over a longer period of time is available it may be possible to follow ontogenetic change and assign also middle sized larvae to a particular taxon with confidence.

Present illustrations are of mounted whole last instar specimens observed in transmitted light and only this way (or at best when specimens are viewed from all sides in strongly oblique illumination) can the length of the sometimes exceedingly thin and usually colourless setae be precisely recognized. In slide mounts, and consequently also in the present illustrations, not all setae are in their natural positions. Therefore the direction in which they point in the figures is not decisive. The present figures never show the very fine procumbent ground pilosity. Four pore-like sensilla on the head are visible only in transmitted light; they provide no means for distinctions.

Proportions of pronotum mentioned in the key refer to measured (not estimated!) distances between insertion points of setal rows surrounding the notum. The notal sclerite itself is frequently only partly delimited by a fine edge resembling a delicate line.

- 1 Exceedingly dense and rigid setae, especially on fore body, less on abdomen. Head trapezoidal, much enlarged behind, setation very conspicuous, especially near rear corners of head. Fossorial legs, tibiae with spine combs, mainly on front leg (Fig. 23) *Isoptena* ENDERLEIN
BL up to 11 mm. The single species, *I. serricornis* (PICTET, 1841), occurs in North and central Europe, mainly in areas with Pleistocene sands where larvae live in shifting sand at the bottom of streams, sometimes at considerable depth.
- 1' Shape normal, pilosity sparse and usually soft (for example, Figs 24, 25) 2
- 2 Eyes large and strongly convex, distinctly longer than temples (Fig. 24a); ocelli large. All of pronotal circumference pilose, lateral portions not hairless; pronotum about 1.7 times wider than long 3




Fig. 23. *Isoptena serricornis* (PICTET, 1841), dorsal view of head and prothorax; ventral view of right foreleg.

- 2' Eyes small, flat, not longer than temples (Fig. 24b, c); ocelli small. Marginal setation of pronotum laterally discontinuous, concentrated on corners of thorax (Fig. 24b); pronotum about 1.5 times wider than long 4
- 3 Body distinctly patterned, brown longitudinal bands on abdomen. Cercal pilosity short, nowhere as long as the segment over which they extend (Figs 25b, e).....
..... *Pontoperla* ZWICK
P. teberdinica (BALINSKY, 1950) (BL ca. 7 mm) and the slightly larger *P. katherinae* (BALINSKY, 1950), in the Caucasus and Anatolia.
- 3' Various patterns, no longitudinal abdominal bands ⁷. At least some cercal setae as long as or longer than the respective cercus segment (Fig. 25a) *Siphonoperla* ZWICK
BL up to 12 mm, most species slightly smaller. Endemic in the West Palaearctic Region, about 9 species. Whether the North African *S. lepineyi* (NAVÁS, 1935) belongs here is not entirely clear. *S. torrentium* (PICTET, 1841) is the most common and most widely distributed species; in Scandinavia occurs only *S. burmeisteri* (PICTET, 1841).
- 4 Small delicate species whose pronotal sclerite lacks a delimiting fold; instead, the sclerite is only delimited by hair fringes. Hair fringes along front and rear margins are sparse, lateral ones widely interrupted (Fig. 24b). Setal fringe on tergite 10 very short, comb-like. Cerci on outer and inner face with a few setae (almost) as long as the segment (Fig. 25c). Abdominal antecostae may be darker on either side of midline (not always distinct) resulting in some banding *Xanthoperla* ZWICK
The single European species, *X. apicalis* (NEWMAN, 1836), is very widely distributed; BL up to 7 mm. At least in central Europe this is a potamal species. It had disappeared for decades but returned recently; for example, it is again common in the River Neisse. Specimens illustrated in this key are from Sardinia where the present species is the only Chloroperlidae. Characters of these specimens agree with descriptions by AUBERT (1953) and BRITAIN (1983), but cannot be reconciled with LILLEHAMMER'S (1988) figures; from the strongly transverse pronotum with blunt setae he illustrated he seems to have had some different taxon before him. Additional species of the genus occur in Middle Asia and the Himalaya.
- 4' Elongate pale larvae whose pronotal sclerite is distinctly delimited in front and posteriorly (Fig. 24c). Pronotal hair fringes much denser than in preceding species (Fig. 24c), hair fringe on tergite 10 much more slender. Setation of cerci along inner face near base short, not as long as segments. However, distally individual setae along dorsal and ventral faces of cerci are very long and fine (Fig. 25d) *Chloroperla* NEWMAN
BL up to 8 mm. Endemic in central and South Europe and Anatolia. The typical species, *C. tripunctata* (SCOPOLI, 1763), is very widely distributed, preferably at moderate elevations. Regionally endemic additional species occur in the major European mountains, at higher elevations. Larvae apparently live relatively deep in stream bottom substrata, probably in the hyporheic zone. *C. tripunctata* has 5-7 very long dorsal and ventral hairs on each distal cercus segment, while *C. breviata* NAVÁS, 1918 (Pyrenees) and *C. susemicheli* ZWICK, 1967 (Alps) have only about 3 on each side (Fig. 25d).

⁷ *Xanthoperla* may have paramedian sections of the tergal antecostae darker than the rest, which may resemble banding; however, the tergite itself behind the darker portion is not pigmented!




Fig. 24. Head and pronotum of *Pontoperla teberdinica* (BALINSKY, 1950) (a), *Xanthoperla apicalis* (NEWMAN, 1836) (b) and *Chloroperla susemicheli* ZWICK, 1967; whole mounts, in transmitted light.




Fig. 25. Dorsal view of left cercus (base and segments 9 & 10) of *Siphonoperla torrentium* (PICTET, 1841), *Pontoperla teberdinica* (BALINSKY, 1950), *Xanthoperla apicalis* (NEWMAN, 1836) and *Chloroperla susemicheli* ZWICK, 1967; whole mounts in transmitted light. To the right, last instar larva of *Pontoperla teberdinica* (from ZWICK, 1971).

Appendix I:**Preliminary key to the German species of genus *Nemoura*,
with notes on a few South European species**

RAUŠER, in his masterly pioneer work (1980) provided a key to the Czechoslovakian species of *Nemoura* in which the relative length of setation on particular cercus segments is of prime importance. This approach requires that segments be accurately counted because the diagnostic proportions change along the cercus. Generally, the relative length of setae increases from the base to about the basal third of the segment and then decreases again. Although these characters are basically valid, there are two problems. **F**irst, the cercus grows during development by addition of segments near the base, similar to the base of the antennal flagellum where new segments are also added. Even in last instar larvae, this basal cercal area is usually incompletely segmented, the number of dividing sulci on the outside may exceed the number on the inside. Similarly, the most basal setal fringes at the end of segments are often only developed on the outer side. Therefore, the "first" segment after the cercus base is hard to identify. **S**econd, the angle at which setae diverge from the segment varies, which influences estimates of relative length. These details may also be perceived differently in slide mounts than in intact specimens viewed in direct light under a dissecting microscope.

An alternative preliminary key was proposed by ZWICK in a lecture handout, in 1993. Although not formally published this document was nevertheless circulated by users. The present key is a re-arranged and extended version of this earlier key, a definite key to all species can still not be presented. Because of the need for future work and also because of the wider geographic frame of the present key at generic level, a few southern European species (whose larvae were previously inadequately described, or completely unknown) are also included.

Exuviae of reared males or mature males in their larval skin of all German species were available. Intact specimens were studied under direct light and slide mounts of entire larvae (or of exuviae) in transmitted light. Magnifications ~100x are sometimes required.

- 1 Pilosity of femora fairly homogenous, regular, procumbent; even though some setae may be distinctly longer than others they remain inconspicuous. Hind femur long and slender, 3 or more times longer than wide at widest point (Figs 26a, 27a)..... 2
- 1' Femora with procumbent small setae plus distinctly larger spine-like setae, pilosity sometimes markedly heterogenous, large setae in regular (Figs 26c, 27c) or irregular arrangement (Figs 26b,d, 27b). Shape of hind femur variable, often strikingly short and thick (only about 2.5x longer than wide)..... 3
- 2 Setae along caudal margin of abdominal tergites short, the longest barely half as long as the corresponding tergite, of normal spine shape. Setae around pronotum and along wing margins pointed. Setae on cerci inconspicuous, the longest not quite as long as the respective segment. Pedicellus concolorous with scapus and antennal flagellum
..... *cinerea* (RETZIUS, 1783)
Ubiquitous stream species, tolerates some pollution, found also in acid peat bogs. Rare in typical Plecoptera habitats, but may be very common in marginal habitats that other Plecoptera fail to colonize.
- 2' Setation along caudal margin of abdominal tergites heterogenous: most setae very short and apically truncate and frazzled (Fig. 28), while several others are extremely thin, pointed and attain segment length; they are apically thinned and turn into a very fine, thin, flexible hair that is easily overlooked. Setae along pronotal and wing margins

truncate, the truncate end often a little wider than the section just basally from it. Cercal setation conspicuous, forming dense hair screens on segments 10-15 that are longer than the corresponding segments. Scape paler than both head or pedicel and flagellum

..... *avicularis* MORTON, 1894

In large and small streams and along lakeshores in Scandinavia, in North Germany typically along shores of oligotrophic lakes, in the central European highlands in rhithral streams.




Fig. 26. *Nemoura* spp., right front femur in dorsal view; not to scale.




Fig. 27. *Nemoura* spp., examples of shape and setation of hind femur.




Fig. 28. *Nemoura avicularis*, left half of tergites 4 and 5, and detail of a marginal seta with frayed apex; not to scale.

- 3 Along distal margin of abdominal tergites only densely placed short setae attaining at most ¼ of tergite length..... 4
- 3' Armature along distal margin of abdominal tergites composed of longer spine-like setae, or of a mixture of spatulate setae plus several long hairs..... 5
- 4 Setal fringe around pronotum everywhere distinct from short blunt spines on disc (Fig. 29a). Setae along abdominal tergites short, blunt, club-shaped, but not spatulate (Fig. 29b). Setal ring of distal cercus segments sparse, not longer than half segment, intercalary pilosity insignificant.....*sciurus* AUBERT, 1949
In rhithral streams, with an evident preference for waters rich in carbonate.
- 4' Setal fringe on pronotum distinct along sides, but along front and especially hind margins barely discernible from the short spines on disc; these are thick and end in a sharp transverse edge with slightly extended corners (Fig. 29c). Abdominal setae straight, very short in comparison with segment length, blunt, club-shaped, those in centre spatulate (Fig. 29d). Setal rings of distal cercal segments dense, longer than half segment, soft, forming a fine brush-like pilosity, together with well developed fine intercalary hairs*dubitans* MORTON, 1894 ⁸
Characteristic inhabitant of reedy and swampy areas near the influx of little streams, in reedy shore-line pockets of little lakes in North Germany, often also in tiny puddles around stands of *Phalaris* or *Juncus*, apparently as long as some groundwater is seeping in.
- 5 Large bristles forming a narrow ring near middle or apical third of femur (Figs 26c, 27c). Pronotal setae very long, pointed 6
- 5' Large setae on femora less orderly, shaggy on hind femur (Fig. 27b) 7
- 6 Tergal marginal setation consisting of a mixture of short spatulate spines and long, distally fine, hair-like setae. The longest ones are lateral and are much longer than the corresponding tergite (Fig. 30). On posterior tergites (T7-9), spatulate setae and fine hairs alternate almost regularly, and hairs are not longer than the respective segment. Pronotal setation straight *mortoni* RIS, 1902
Alpine and montane areas in central Europe, typically in springs and spring runs; often covered with debris.

⁸ LILLEHAMMER (1988) says the first tarsal segment to be curved which is not the case in German specimens.

- 6' Tergal marginal setation consisting exclusively of pointed and rather long sparse spines, the longest about as long as the corresponding tergite. Pronotal setation a little wavy. ...
*palliventris* AUBERT, 1953
 Widespread rhithral Italian species, from the Maritime Alps to Sicily.




Fig. 29. *Nemoura sciurus* AUBERT, 1949 (a, b) and *N. dubitans* MORTON, 1894 (c, d). Setae on disc of pronotum, near midline (and along front margin, for *N. sciurus*; a, c) and left half of tergite 5 (b, d).




Fig. 30. *Nemoura mortoni* RIS, 1902, left half of tergites 4 and 5; detail of a spatulate marginal seta not to scale.

- 7 Setae in lines on surface of wing-pads are short spines of various forms 8
 7' Wing-pads with fine, erect long hairs in linear arrangement.....*lacustris* E. PICTET, 1865
 Along tiny, slow-flowing streams in southern France and on the Iberian Peninsula.

- 8 Integument matt, rough through numerous intercalary little spines on all thoracic and abdominal tergites. Tergite margins with series of slightly curved spines that grow progressively longer from middle towards sides (Fig. 31a), lateral spine pair much longer other spines, especially central pair. Large setae on fore femur leaving an extended distal area covered with only small setae (Fig. 26b).....*marginata*-group
 Several species inhabiting springs and rhithral stream sections share a common appearance and are too similar, and at the same time too variable, to be presently separated. I examined the following: *arctica* ESBEN-PETERSEN, 1910, *cambrica* STEPHENS, 1835, *erratica* CLAASSEN, 1936, *flexuosa* AUBERT, 1949, *marginata* PICTET, 1836, *minima* AUBERT, 1946, *uncinata* DESPAX, 1934, *undulata* RIS 1902. Still more species may fall in the same group.
- 8' Integument shining. Erect marginal setae along posterior edges of tergites prominent, intercalary setation weak (Fig. 31b), very sparse on posterior abdominal segments. Length of marginal setae relatively uniform, no size cline from short paramedian to long lateral setae. Femora covered with erect spines, the largest ones are close to apex, barely any fine procumbent pilosity distally from them (Fig. 26d) 9




Fig. 31. *Nemoura uncinata* DESPAX, 1934 and *N. obtusa* RIS, 1902, left half of tergites 4 and 5; to different scales.

- 9 Setae on antero-lateral margin of pronotum acute. Tiny setae in lines on wing-pads are small pointed spines and occasional slender hairs (Fig. 32a, b)..... *obtusa* RIS, 1902
 Springs and little streamlets in the Alps. A single record from the Polish Tatra.
- 9' Setae on antero-lateral margin of pronotum almost cylindrical, only apex narrowed to a short point. Tiny setae in lines on wing-pads ovoid (Figs 32c, d) *sinuata* RIS, 1902
 Springs and little streamlets in the Alps.




Fig. 32. *Nemoura obtusa* RIS, 1902 (a, b) and *N. sinuata* RIS, 1902 (c, d), rows of tiny setae on wing-pads (a, d) and lateral marginal setae of pronotum, near anterior corner (b, c).

Appendix II:

Diagnoses of selected characteristic species and species groups of *Leuctra*

RAUŠER (1980) offered a key to the Czechoslovak species of *Leuctra*. However, in this very large and difficult genus specific identifications remain problematic, with few exceptions. Below, new diagnoses are presented for a few individual species that can be reliably identified, and for two major species groups that are proposed as Operational Taxonomic Units (OTU) until more precise identifications become possible.

- Body flattened, large. Near basal part of antennal flagellum several segments with one or two outgrowths (Fig. 33); fairly hairy..... *L. geniculata* (STEPHENS, 1835)




Fig. 33. *Leuctra geniculata* (STEPHENS, 1835), base of antenna (from RAUŠER, 1980)

- Very hairy, especially head (Fig. 34) and sides of pronotum:
 - Clypeus with prominent pointed corners (Fig. 34a); basal part of antennal flagellum with striking long setae. Lower side of head below antennal insertion normal. Found mainly in deep aerated deposits of fine detritus *L. braueri* KEMPNY, 1898
 - Clypeus normal. Setae on antennal flagellum much shorter than segments, setation inconspicuous. Lower contour of head below antennal bulging outwards (Fig. 34b); on this blunt swelling arise long setae. Body of preserved specimens tends to be completely straight. Mainly in sandy deposits *L. nigra* (OLIVIER, 1811)




Fig. 34. *Leuctra braueri* KEMPNY, 1898 (a), head in dorsal view, and *L. nigra* (OLIVIER, 1811) (b) head in combined dorsal and ventral views.

- Macroscopic pilosity of normal density, or sparse, sometimes almost absent:
 - Rear edge of thorax between front wing-pads oblique, between hind wing-pads the thorax margin forms little pointed extensions (Fig. 35a)*prima-hippopus-inermis-group*
 - Rear edge of thorax between wing-pads truncate or gently rounded (standard configuration, also in the 3 individual *Leuctra* species keyed above, as well as in the genera *Tyrrhenoleuctra* and *Pachyleuctra*) (Fig. 35b).....*fusca-group*

Notes on central European members of the *prima-hippopus-inermis-group*:

- Some members of the *prima*-subgroup (for example, *prima* KEMPNY, 1899, *autumnalis* AUBERT, 1948, *signifera* KEMPNY 1899, *pseudosignifera* AUBERT, 1954, probably also others) appear relatively hairy because hairs are stiff and tend to be well pigmented.
- *L. hippopus* KEMPNY, 1899 and the species of the *inermis*-subgroup (for example, *L. inermis* KEMPNY, 1899, *L. rauscheri* AUBERT, 1957, *L. handlirschi* KEMPNY, 1898, *L. teriolensis* KEMPNY, 1900) have paler, less easily noticed hairs but those on the abdominal segments are nevertheless about half as long as the segment length and well visible in profile.




Fig.35: *Leuctra prima* (a) and *L. albida* (b), dorsal view of fore body.

Notes on central European members of the *fusca*-group: large group with many autumnal species.

- Some are normally shaped and normally pilose like *L. albida* KEMPNY, 1899 (Fig. 36b), *L. fusca* (L., 1758) and *L. digitata* KEMPNY, 1899 (Fig. 36a; the distinction of the latter two species after RAUŠER 1980 fails!);
- Many have short hairs and are slender (like *L. aurita* NAVÁS, 1919; Fig. 36c),
- Some are almost hairless, very slender and pale. The subterranean *L. major* BRINCK, 1949 and *L. leptogaster* AUBERT, 1949 are extremes.




Fig. 36. Head and thorax of *Leuctra* spp. (from RAUŠER, 1980).

Acknowledgements

I gratefully acknowledge the gift of important material to my collection by several colleagues, especially Dr H. MALICKY (Lunz), and Dr W. GRAF (Wien), Dr W. KITTEL (Łodz), Dr E. MAUCH (Dinkelscherben), Dr I. SIVEC (Ljubljana), Dr G. VINÇON (Grenoble). Dr P. HAASE (Biebergemünd) is thanked for constructive criticism. My sincere thanks for copyright permissions are extended to PROF. DR R. ROZKOŠNY (Brno; RAUŠER, 1980), to DR U. MOLTMANN (Heidelberg; ILLIES, 1955) and to Drs Y. GONSETH (Neuchâtel) and D. BURCKHARDT (Bern; AUBERT, 1959).

References

- AUBERT, J. (1946): Les Plécoptères de la Suisse Romande. - Mitt. Schweiz. ent. Ges., Lausanne, 20: 8-128.
 AUBERT, J. (1949): Plécoptères helvétiques. Notes morphologiques et systématiques. – Mitt. Schweiz. Entomol. Ges. 22: 217-236.
 AUBERT, J. (1950): Note sur les Plécoptères européens du genre *Taeniopteryx* Pictet (*Nephelopteryx* Klapálek) et sur *Capnia vidua* Klapálek. – Mitt. Schweiz. Entomol. Ges. 23: 303-316.
 AUBERT, J. (1953): Contribution à l'étude des Plécoptères et des Ephéméroptères de la Calabre. – Ann. Ist. Mus. Zool. Univ. Napoli 5 (2): 1-35.
 AUBERT, J. (1959): Plecoptera. – Insecta Helvetica Fauna 1: 1-140.
 BRAASCH, D.; JOOST, W. (1971): Zur Plecopterenfauna Bulgariens. – Limnologica 8(2): 265-294.

- BRITAIN, J.E. (1983): The first record of the nymph of *Xanthoperla apicalis* (Newman) (Plecoptera: Chloroperlidae) from Scandinavia, with a key to the mature nymphs of the Scandinavian Chloroperlidae. – Fauna Norv., Ser. 30: 52-53.
- DESPAX, R. (1931): *Perla nigratarsis* nov. sp. Plécoptère nouveau du sud de la France. – Bull. Soc. Hist. Nat. Toulouse 62(4): 419-427.
- GRAF, W. (1997): A new record of the perlid stonefly *Agnentina elegantula* (Klapálek, 1905) in Europe. – In: LANDOLT, P.; SARTORI, M. (eds), Ephemeroptera & Plecoptera Biology-Ecology-Systematics. Fribourg: Mauron + Tinguely & Lachat SA, 205-208.
- HARPER, P.P. (1994): Plecoptera. –Pp. 176-209 in: MORSE, J.C.; YANG, L., TIAN, L. (eds): Aquatic insects of China useful for monitoring water quality. Hohai University Press, Nanjing, P. R. China.
- INTERNATIONAL CODE OF ZOOLOGICAL NOMENCLATURE (1999). – The International Trust of Zoological Nomenclature, London; 4th edition, I-XXIX; 1-306.
- ILLIES, J. (1952): Die europäischen Arten der Plecopterengattung *Isoperla* Banks (*Chloroperla* Pict.). –Beitr. Entomol. 2: 369-424.
- ILLIES, J. (1955): Steinfliegen oder Plecoptera. – In: DAHL, F (1955): Die Tierwelt Deutschlands und der angrenzenden Meeresteile 43: I-IV;1-150.
- ILLIES, J. (1966): Katalog der rezenten Plecoptera. – Das Tierreich 81: I-IXXX; 1-632; 20 Figs.
- KONAR, M. (2000): Die Larvaldetermination der Arten der Gattung *Isoperla* in Kärnten (Oesterreich). – Lauterbornia 37 (1999): 57-62.
- LILLEHAMMER, A. (1988): Plecoptera fauna of Fennoscandia and Denmark. – Fauna Entomol. scand. 21, 163 pp.
- RAMBUR, J.P. (1842): Tribu des Perlides. – Pp. 449-462 in: Histoire naturelle des insectes. Névroptères. Paris : Roret, I-XVII ; 1-534 ; 12 pls..
- RAUSER, J. (1956): Zur Kenntnis der tschechoslowakischen *Protonemura*-Larven. – Práce, Acta Acad. Sci. Čsl. 28: 449-496.
- RAUSER, J. (1963): Contribution à la connaissance des larves de genre *Amphinemura* de la Tchécoslovaquie (Plecoptera). – Čas. Česk. Spol. Entomol. 60: 32-54.
- RAUŠER, J. (1980): Rád Posvatky – Plecoptera. — Pp. 86-132 in: ROZKOSNY, R. (ed.): Klic vodnich larev hmyzu. – Prague: Česk. Akad. Ved. 521 pp. In Czech.
- RAVIZZA, C.; VINÇON, G. (1999): Les Leuctridés (Plecoptera: Leuctridae) des Alpes. – Mitt. schweiz. entomol. Ges. 71(3-4) (1998): 285-342.
- RUPPRECHT, R. (1997): An Attempt to Explain Different Drumming Signals within *Capnia bifrons*. – Pp. 93-98 in: LANDOLT, P.; SARTORI, M. (eds), Ephemeroptera & Plecoptera Biology-Ecology-Systematics. Fribourg: Mauron + Tinguely & Lachat SA; I-XI; 1-569.
- SCHOENEMUND, E. (1925): Die Larven der deutschen *Perla*-Arten. – Entomol. Mitt. 14: 113-121.
- SEZZI, E. (2001): *Biogeografia molecolare e storia evolutiva del genere Tyrrhenoleuctra (Plecoptera, Insecta) nel bacino del Mediterraneo*. PhD Thesis (Biological and Biochemical Evolution). University of Viterbo (Italy). 93 pp.
- SIVEC, I.; GRAF, W. (2003): *Perla carantana* – a new species of the genus *Perla* (Plecoptera: Perlidae) from Austria and Slovenia. – Natura Sloveniae 4: 31-38.
- SIVEC, I.; STARK, B.P. (2002): The species of *Perla* (Plecoptera: Perlidae): Evidence from egg morphology. – Scopolia 49: 1-33.
- SIVEC, I.; STARK, B.P.; UCHIDA, S. (1988): Synopsis of the world genera of Perlinae (Plecoptera: Perlidae). – Scopolia 16: 1-66.
- STEWART, K.W.; STARK, B.P. (2002): Nymphs of North American stonefly genera (Plecoptera). – Columbus, Ohio: The Caddis Press, 2nd edition, XII + 510 pp.
- VINÇON, G.; RAVIZZA, R. (2001): Leuctridae (Plecoptera) of the Pyrenees. – Anns Limnol. 37: 293-322.
- ZHILTZOVA, L.A.; TESLENKO, V.A. (1997): Order Stoneflies (Plecoptera). – Pp. 248-264, 364-398 (pls. 38-55) in: TSALOLIKHIN, S.J. (ed.): Key to Freshwater Invertebrates of Russia and adjacent lands, Vol.3. Zool. Inst. Akad. Nauk., Sankt Petersburg, 446 pp. (several unnumbered, incl. 67 pls). In Russian.
- ZWICK, P. (1971): Plecoptera aus Anatolien und benachbarten Gebieten. – Mitt. Schweiz. Entomol. Ges. 44(3/4): 233-264.
- ZWICK, P. (1973): Insecta Plecoptera. Phylogenetisches System und Katalog. – Das Tierreich 94: I-XXXII, 1-465.
- ZWICK, P. (1984): Notes on the Genus *Agnentina* (= *Phasganophora*) (Plecoptera: Perlidae). – Aquatic Insects 6: 71-79.
- ZWICK, P. (2000): Phylogenetic System and Zoogeography of the Plecoptera. – Annu. Rev. Entomol. 45: 709-746.
- ZWICK, P. (2003) : Shapes and patterns of wingpad development in the Plecoptera. – Pp. 477- 483 in: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia.