

Oregon Agricultural Conservation Easement Program (ACEP)– Wetland Reserve Easements (WRE)

Wetland Restoration Criteria and Guidelines, 2023



NRCS Oregon Wetland Easement Sites

Left side (top to bottom): Muddy Valley and Valley Junction

Right side (top to bottom): Thomas and Sevenmile

September 25, 2023

TABLE OF CONTENTS

- I. INTRODUCTION
 - A. Purpose of Wetland Restoration Criteria and Guidelines
 - B. Historical Wetlands and Wetland Loss in Oregon
- II. OREGON ECOREGIONS AND WETLAND TYPES
 - A. Oregon Ecoregions
 - B. Oregon Wetland Types and Restoration Approaches
 - 1. Freshwater Wetlands
 - 2. Estuaries and Tidal Wetlands
 - 3. Flowing Water and Riparian
- III. WRE PROGRAM ELIGIBILITIES
 - A. Eligible Land Types
 - 1. Farmed or Converted Wetlands
 - 2. Riparian Areas
 - 3. Lands in the CRP
 - 4. Wetlands Restored or Protected
 - B. Eligible Adjacent Lands
 - C. Eligible Conservation Practices
 - D. Ranking Factors and Criteria
- IV. TECHNICAL CONSIDERATIONS FOR CUAs
- V. TECHNICAL CONSIDERATIONS FOR WAIVERS
- VI. LITERATURE CITED
- VII. LIST OF RESOURCES FOR FURTHER INFORMATION

List of Appendices

- A. Oregon Conservation Strategy Ecoregion Maps
- B. Ranking Worksheet
- C. ACEP-WRE Priority Geographic Regions
- D. Wetland Restoration, Conservation Practice 657-Specification Sheet

I. INTRODUCTION

A. Purpose of Wetland Restoration Criteria and Guidelines

The Wetland Restoration Criteria and Guidelines (WRCG) for Oregon contains the technical information used to guide decision making for Agricultural Conservation Easement Program - Wetland Reserve Easements (ACEP-WRE). This includes decisions related to eligibility, ranking, selection, restoration, enhancement, and management of wetlands and associated habitats. The WRCG technical criteria and guidelines in this document have been developed in consultation with the NRCS Oregon, Oregon State Technical Advisory Committee (STAC), U.S. Fish and Wildlife Service (USFWS), Oregon Department of Fish and Wildlife (ODFW), Soil and Water Conservation Districts in Oregon, and other partners. This WRCG document is designed to serve as a basis for technical determinations and decisions related to wetland restoration activities implemented under ACEP-WRE. The WRCG is also a robust document designed to be an adaptive management resource that can be applied and updated as needed throughout the lifespan of an easement or 30-year contract.

The ACEP Manual, Wetland Restoration Definition and Principles, defines wetland restoration as the rehabilitation of degraded or lost wetland and associated habitats. Restoration reestablishes original, native vegetation and hydrology to the extent practicable or establishes a hydrologic regime and native plant community to replace the original habitat functions and values. Restoration benefits migratory waterfowl and wetland-dependent wildlife and addresses local resource concerns. (See ACEP Manual (Title 440, Part 528), Subpart N (WRE-Restoration), 528.131 A and B for further detail on above: [CPM, Title 440, Part 528, Section 528.131](#)).

B. Historical Wetlands and Wetland Loss in Oregon

A historical wetland is a wetland that existed prior to the hydrologic and vegetative manipulation generally associated with conversion to agriculture at the time of European-American colonization. While a substantial proportion of wetlands in Oregon have been manipulated or converted to uplands, some examples of historical wetlands still exist today in relatively undisturbed condition. Estimates of wetland loss in Oregon since 1850 vary by data source, area of the state, and wetland type. In Oregon overall, it is estimated that 38 percent of all wetlands have been lost (Dahl, 1990). There are 518 wetland plant communities identified by the Oregon Natural Heritage Program and 29% (151) of wetland plant communities are considered imperiled (Christy and Titus, 1997; Risser, 2000; Morlan, 2000).

Klamath Basin

The landscape around what is now Lower Klamath National Wildlife Refuge in Oregon was historically a large inland shallow lake with associated wetlands connected to the Klamath River as a source of wetland hydrology (Mayer, 2005). It is estimated that the Upper Klamath Lake area has lost approximately 30,000 acres of wetlands, and the Klamath Basin overall has lost an estimated 75% of its wetlands over the past 50 years because of altered hydrology (Akins, 1970; Fretwell et al., 1996; Morlan, 2000; Mayer, 2005).

Willamette Valley

In the Willamette Valley at the time of European colonization, the U.S. Fish and Wildlife Service (2017) estimated 340,000 acres of historical wet prairie, 240,000 acres of historical riparian forested wetlands, and 13,000 acres of historical shrub wetlands. The Temperate Pacific

Freshwater Emergent Marsh was a historic wetland type found in the Willamette Valley with groundwater hydrology sources. Christy et al. (2000) estimated 8,893 acres of freshwater emergent wetlands in the Willamette Valley. Larger historic freshwater emergent wetland areas in the Willamette Valley included littoral wetlands near Lake Labish, Lousignont Lake, and Wapato Lake (Christy and Alverson, 2011). Currently, this type only occurs as small patches in floodplains and littoral zones of ponds or lakes (Rocchio and Crawford, 2009). Estimates of historic wetland loss in the Willamette Valley due to land use change and hydrology modification are 57% overall. In addition, 98% of historic wetland prairie has been lost and 44% of the historic wetland plant communities are considered imperiled (OPB, 2000; Taft and Haig, 2003; Christy and Alverson, 2011; Fickas et al., 2016).

Tidal wetland loss

A recent study by Brophy (2019) did an analysis of historic coastal wetlands. Prior to European settlement, it is estimated that 38,052 acres of wetlands existed along the Oregon Coast, and 15 large estuaries make up approximately 96.5% of historic tidal wetlands. Historic wetlands along the Oregon Coast consisted of forested and scrub-shrub wetlands, or “tidal swamp” (58%) and emergent “tidal marsh” (42%) (Brophy, 2019). Diking has resulted in the loss of 22,034 acres (57.9%) of these historic tidal wetlands and 8,335 acres (21.9%) of these historic wetlands have been converted to another vegetation type. If you combine effects of diking that result in changes to wetland hydrology and changes to wetland vegetation, approximately 95% of forested tidal wetlands and 58.9% of tidal marsh have been lost (Brophy, 2019).

Riparian wetland loss

According to the Oregon State of the Environment Report (Risser, 2000), Oregon contains approximately 114,500 miles of streams and rivers that have associated riparian ecosystems. It is estimated that older age class, coniferous tree canopy remains in 20% of the riparian forests in the Cascades and 3% of the riparian forests in the Coast Range. Since the 1850s, in the Willamette Valley Ecoregion, it is estimated that 80% of river channel complexity and associated riparian areas along the main channel and upper tributaries of the Willamette River have been lost (Risser, 2000; OPB, 2000). Hulse et al. (2002) estimated that between 1850 and 1995, 41,000 acres of river channel and island areas have been reduced and total length of all channels have been reduced from 355 miles to 264 miles. In addition, 85% of the riparian forest along the main channel McKenzie River has been lost (Risser, 2000).

II. OREGON ECOREGIONS AND WETLAND TYPES

A. Oregon Ecoregions

This WRCG focuses on those ecoregions identified by the Oregon Department of Fish and Wildlife (ODFW), Oregon Conservation Strategy (OCS) (2016) ([see OCS ecoregions website](#)) that contain the majority of the NRCS Oregon WRE and WRP easements. These ecoregions include Blue Mountains, Coast Range, East Cascades, Northern Basin and Range, and Willamette Valley (see Figure 1).

Oregon Conservation Strategy Ecoregions




Figure 1. NRCS Oregon Wetland Easements by OCS Ecoregion (ODFW, 2016)

This section includes a general overview and conservation issues for each OCS Ecoregion (Crowe et al., 2004; Oregon Department of Fish and Wildlife, 2016). See Appendix A for maps that show NRCS wetland easements and OCS wetlands for each OCS Ecoregion. The OCS (2016) considers all wetland types a priority habitat for conservation in all ecoregions throughout the state of Oregon ([see Oregon Conservation Strategy wetlands website](#)). On page ten, Table 1 includes information on which wetland types occur in which ecoregion and a list of OCS 2016 strategy species for each ecoregion that are wetland or riparian habitat dependent. Further information on specific wetland types in Oregon is included in Section B, Oregon Wetland Types and Restoration Approaches.

Blue Mountains Ecoregion (see [Blue Mountains Ecoregion](#))

General description: The Blue Mountains Ecoregion covers approximately 23,928 square miles and ranges in elevation from 1,000 feet at Snake River to 9,838 feet at Sacajawea Peak. The elevation difference results in a wide range of temperatures and precipitation. Summers are short and dry, winters are long and cold, and snow melt is the primary source of annual precipitation. The average annual precipitation is 8.0 to 24.3 inches (snowfall 11.2 to 87.6 inches) (OCS, 2006). Drier areas like Baker Valley get 9-16 inches of precipitation per year and wetter areas like Grande Ronde Valley can get 13-24 inches of precipitation per year. The ecoregion has a wide diversity of geology and landforms, including the Ochoco, Blue, and Wallowa mountain ranges, valleys, plateaus, glacially-cut deep canyons, gorges, and mountain lakes (above paraphrased from OCS 2016, Ecoregions).

This variation in geology, elevation, and climate also drives diversity in ecosystems and plant communities. Plant community variation due to elevation can be observed generally from north to south across the ecoregion, with ponderosa pine in mid-elevations and mixed coniferous forests at high elevations. Western juniper (*Juniperus occidentalis*) is a dominant community on the western side of the ecoregion and sagebrush and grassland steppe are dominant communities on the eastern side. Pre-European settlement vegetation communities included native Ponderosa pine savanna, sagebrush steppe, grasslands, and riparian woodland (above paraphrased from OCS 2016, Ecoregions).

Conservation issues: Water availability due to dry conditions, water diversion, and resulting low water table has impacted wetlands and riparian areas within the Ecoregion (Oregon State of the Environment, Ecoregions, 2000; OCS, Wetlands 2016).

Coast Range Ecoregion (see [Coast Range Ecoregion](#)):

General description: The Coast Range Ecoregion in Oregon covers approximately 9,263 square miles and ranges in elevation from 0 at the Pacific Ocean to 4,097 feet at Mary's Peak (OCS, 2016). The average annual precipitation is 60 to 98 inches (snowfall <1 to 2 inches), making it the wettest ecoregion in the state (OCS, 2006). The cool, moist air from the ocean and high rainfall of this ecoregion result in ecosystems such as temperate coniferous rainforests, which are highly productive. Diverse habitats in this ecoregion also include deciduous riparian forest, sand dunes, tidepools, estuaries and the highest density of streams in the state (above paraphrased from OCS, 2016). Forests in this ecoregion can include tree species such as Sitka spruce (*Picea sitchensis*), Douglas-fir (*Pseudotsuga menziesii*), western red cedar (*Thuja plicata*), and western hemlock (*Tsuga heterophylla*) (Omernik, 1987).

Conservation issues: Development and land use change due to increasing population, oil spills, and altered hydrology due to historic diking have caused loss of habitat area and impacts to tidal marsh and coastal wetland functions along the Oregon Coast. Increased visitation from tourists to the Oregon Coast may result in increased disturbance to nesting shorebirds, tidepool communities, and other sensitive coastal habitats (OCS, 2016).

East Cascades Ecoregion (see [East Cascades Ecoregion](#)):

General description: The East Cascades Ecoregion in Oregon covers 10,603 square miles and ranges in elevation from 70 feet at Columbia River Gorge to 8,364 feet at Gearhart Mountain (USDA Forest Service, website accessed 8/23/2023: [Fremont Winema NF](#)). The annual average precipitation is 9.8 to 89.6 inches (snowfall 19.7 to 420 inches) (OCS, 2006), and the climate is generally dry with wide variations in temperature. The terrain of the East Cascades includes features such as buttes, lava flows and caves, deep ash deposits, and craters such as Crater Lake that are a result of historical volcanic activity. The northern two-thirds of the ecoregion are drained by the Deschutes River and the southern third is primarily drained by the Klamath River (above paraphrased from OCS, 2016).

The East Cascades Ecoregion has exceptional habitat diversity including lodgepole pine (*Pinus contorta*) forests on deep Mazama ash in drier areas, montane and foothill Ponderosa pine (*Pinus ponderosa*) forests, sagebrush flats, and alpine meadows (OCS, 2016; Crowe et al., 2004; Risser, 2000). The network of Klamath Basin lakes and wetlands are also known as nationally significant waterfowl habitat and are considered critical migratory staging areas for 80% of Pacific Flyway waterfowl, with peak counts to 1.6 million birds at Lower Klamath National Wildlife Refuge during the fall migration (Mayer, 2005). On an annual basis, over 30 species of waterfowl, > 3 million ducks, and ½ million geese use Klamath Basin as a migratory stop-over location (OCS, 2016 Wetlands; OCS, 2006; Gilmer et al., 2004).

Conservation issues: Water that historically maintained extensive shallow lake and emergent marsh systems and wet meadows has been drained and converted to support agriculture and urban development throughout the ecoregion (OCS, 2016).

Northern Basin and Range Ecoregion (see [Northern Basin and Range Ecoregion](#))

General description: The Northern Basin and Range Ecoregion in Oregon covers 23,955 square miles and ranges in elevation from 2200 feet at Snake River to 9,733 feet (9738 on topo) at Steens Mountain (OCS, 2016). This ecoregion is the driest in Oregon, with 8-12 inches of precipitation per year in the southeastern part of the ecoregion. Most areas average less than 15 inches with up to 40 inches per year in higher elevations (OCS, 2006, 2016). Mountain ranges generally run north to south and often take the form of fault blocks. Soils are volcanic in origin and are generally rocky, high in minerals, thin, and low in organic matter (Crowe et al., 2004). Evaporation and drying of lakes that formed in the Pleistocene (40,000 and 10,000 years ago) has resulted in large areas of alkali flats and associated pools, shallow lakes, and wetlands that have characteristic salt and mineral deposits (OCS, 2016). Alkali flats are important habitat for migrating shorebirds that feed on abundant invertebrates.

This ecoregion is known as sagebrush desert or high desert. Unique plant communities in this

ecoregion have adapted to the high salt content of salt-desert scrub and alkali flats communities. Plant species in these communities include inland saltgrass (*Distichlis spicata*) and spiny, salt-tolerant shrubs such as black greasewood (*Sarcobatus vermiculatus*), iodine bush (*Allenrolfea occidentalis*), and hopsage (*Grayia spinosa*). Widely-spaced sagebrush is typical vegetation that can be associated with bluebunch wheatgrass (*Pseudoroegneria spicata*), Idaho fescue (*Festuca idahoensis*), and other perennial bunchgrasses. Forested vegetation is rarer, but white fir stands occur in the Steens and Hart Mountain ranges. Aspen and mountain mahogany are more widespread and can be found in the Trout Creeks, Steens Mountain, Pueblo Mountains, Oregon Canyon Mountain, and Mahogany Mountains areas of the ecoregion (above paraphrased from OCS, 2016).

Conservation issues: Lakes and wetlands in this ecoregion provide critical habitat for nesting and migratory birds. Large networks of freshwater emergent marshes and wet meadows associated with Albert, Summer, Malheur, Harney Lake, and Warner Basins have been lost or reduced due to conversion to agriculture, water diversion, and altered hydrology for other land uses (OCS, 2016, Wetlands). Uncontrolled livestock grazing prior to the Taylor Grazing Act of 1934 has degraded sensitive ecological areas, riparian areas, and some areas of sagebrush steppe in the ecoregion. Fire suppression, juniper encroachment, and invasive non-native annual grass species (cheatgrass) have also impacted native riparian and sagebrush communities throughout the ecoregion (OCS, 2016).

Willamette Valley Ecoregion (see [Willamette Valley Ecoregion](#)):

General description: The Willamette Valley Ecoregion in Oregon covers approximately 5,560 square miles and ranges in elevation from 13 to 400 feet (Fickas et al., 2016). Annual precipitation ranges from 37 inches to 46 inches (Fickas et al. 2016), with snowfall of 1.7 to 6.0 inches (OCS, 2006). The topography of the Willamette Valley is characterized by broad alluvial floodplains associated with the Willamette River, fluvial terrace upslope of Willamette River bottomlands, low hills, and mountain foothills (Pater et al., 1998). Willamette Valley ecosystems include wetland prairie and Oregon white oak (*Quercus garryana*) savanna which offer unique habitats within the state of Oregon (USFWS, 2017). Wetland prairie plant species include tufted hairgrass (*Deschampsia caespitosa*), rushes (*Juncus* spp.), and sedges (*Carex* spp.), common camas (*Camassia quamash* ssp. *maxima*), and great camas (*Camassia leichtlinii* ssp. *suksdorfii*). Extensive riparian forested communities occur along the Willamette River corridor and connected tributaries with species such as Oregon ash (*Fraxinus latifolia*), black cottonwood (*Populus trichocarpa*), red alder (*Alnus rubra*), big-leaf maple (*Acer macrophyllum*), western red cedar (*Thuja plicata*), willows (*Salix* spp.) and Douglas-fir (*Pseudotsuga menziesii*) (Pater et al., 1998).

More than 150 years ago, around the time of Euro-American colonization, open prairies were the dominant vegetation community. Prairies were surrounded by oak woodlands, solitary oaks and pine trees, and coniferous forest in the foothills (Christy and Alverson, 2011; Johannessen et al., 1971; USFWS, 2017). Pre- Euro-American colonization, the Kalapuya people managed the prairie through frequent, low severity fire to maintain food resources (camas, berries, roots, acorns, filberts) and hunting grounds for deer (Boyd, 1986; Hamman et al., 2011; Johannessen et al., 1971; Taft and Haig, 2003). Euro-American settlers established commercial farming, ranching, and logging practices and suppressed the use of fire as a management tool. This

resulted in ecological change to a vegetation community dominated by oak and pine (Christy and Alverson, 2011; Connolly, 2000; Hamman et al., 2011; Taft and Haig, 2003).

Conservation issues: Since the 1850s, much of the Willamette Valley ecoregion has been altered by development (agricultural and urban), particularly affecting oak woodland, oak savanna, grassland, riverine, and wetland habitats. The Willamette River and wetlands throughout this ecoregion, including wet prairie, have been affected by altered water regimes, pollution, land use change, and invasive plants and animals. Several federally-listed plant species are associated with Willamette Wet Prairie, including Bradshaw's lomatium (*Lomatium bradshawii*) and Nelson's checkermallow (*Sidalcea nelsoniana*). The recovery zones for these species comprise the Priority Areas within the Willamette Valley Priority Geographic Region (see [USFWS Recovery Plan for Prairie Species 2010](#)). About 96 percent of the Willamette Valley ecoregion is privately-owned, presenting challenges to conservation efforts. Conservation strategies that focus on needs of individual at-risk species and key sites are particularly critical in this ecoregion (all above paraphrased from ODFW, 2016).

B. Oregon Wetland Types and Restoration Approaches

Classifying wetland types is an approach that conservation planners can use to organize and plan wetland restoration practices. There are numerous systems used to classify wetlands based on a variety of factors including: landscape position, geomorphology, dominant hydrology source, hydrologic regime, soil types, and vegetation. Wetland classification systems that have been used in Oregon include habitat and vegetation-based systems (Christy et al., 2017a; Kagan et al., 2007; Crowe et al., 2004; Cowardin et al., 1979), and hydrogeomorphic (HGM) based classification systems (Tiner et al., 2011; Adamus, 2001, 2006; Brinson, 1993). NRCS Oregon uses the National Wetland Inventory, Cowardin system (Cowardin et al., 1979); and the HGM based system (Brinson, 1993).

Wetland habitat type categories according to the ODFW OCS (2016) that are used by NRCS Oregon will be organized and described below as follows: Freshwater Wetlands, Estuaries/Tidal Wetlands, and Flowing Water and Riparian ([see OCS wetlands website](#)). The distribution of each wetland type is variable across ecoregions in Oregon, with some wetlands being ubiquitous around the state (e.g., freshwater marshes), and others (tidally influenced salt marshes) restricted to one or two ecoregions (see Table 1).

According to the Oregon Conservation Strategy (2016), wetland restoration approaches can focus on the following functions that wetlands can provide: habitat and native plant communities, water quantity, and water quality. Invasive species control and prevention for species that occur in wetlands is a priority for all wetland restoration and management work in Oregon. At potential wetland restoration sites, geospatial datasets can be used to determine most likely historic wetland type and vegetation community (See List of Additional Resources for data sets). Local reference wetlands of a similar type (landscape position, soil types, hydrologic regime, and plant community) can also help determine which hydrology and plant community functions are most likely to succeed in restoration efforts (See Appendix D for Wetland Restoration, 657 Conservation Practice Specifications Sheet).

Table 1. Wetland types and OCS strategy wetland and riparian-dependent wildlife by ecoregion

ODFW Ecoregion	Wetland types	OCS Strategy Species-Wetland and Riparian Dependent Wildlife
Blue Mountains	Alkaline Wetlands, Deciduous Swamps and Shrublands, Marshes, Vernal Pools, Wet Meadows, Wet Prairies	<p>Mammals: Fisher (<i>Pekania pennanti</i>)</p> <p>Birds: Greater Sage-Grouse (<i>Centrocercus urophasianus</i>); Long-billed Curlew (<i>Numenius americanus</i>); Trumpeter Swan (<i>Cygnus buccinator</i>); Upland Sandpiper (<i>Bartramia longicauda</i>)</p> <p>Amphibians: Columbia Spotted Frog (<i>Rana luteiventris</i>); Rocky Mountain Tailed Frog (<i>Ascaphus montanus</i>); Western Toad (<i>Anaxyrus boreas</i>)</p> <p>Reptiles: Western Painted Turtle (<i>Chrysemys picta bellii</i>)</p> <p>Fish: Bull Trout (<i>Salvelinus confluentus</i>); Fall and Spring Chinook (<i>Oncorhynchus tshawytscha</i>); Great Basin Redband Trout (<i>Onchorynchus mykiss newberrii</i>); Summer Steelhead/Columbia Basin Redband (<i>Oncorhynchus mykiss gairdneri</i>); Western Brook Lamprey (<i>Lampetra richardsoni</i>); Westslope Cutthroat Trout (<i>Onchorynchus clarki lewisi</i>)</p> <p>Invertebrates: Columbia Clubtail (<i>Gomphus lynnae</i>); Western Ridged Mussel (<i>Gonidea angulate</i>)</p>
Coast Range	Deciduous Swamps and Shrublands, Marshes, Saltwater Marshes, Tidal Wetlands, Wet Meadows, Wet Prairies	<p>Birds: Black Brant (<i>Branta bernicla nigricans</i>); Caspian Tern (<i>Hydroprogne caspia</i>); Harlequin Duck (<i>Histrionicus histrionicus</i>); Marbled Murrelet (<i>Brachyramphus marmoratus</i>); Tufted Puffin (<i>Fratercula cirrhata</i>); Western Snowy Plover (<i>Charadrius nivosus nivosus</i>)</p> <p>Amphibians: Coastal Tailed Frog (<i>Ascaphus truei</i>); Columbia Torrent Salamander (<i>Rhyacotriton kezeri</i>); Cope's Giant Salamander (<i>Dicamptodon copei</i>); Clouded Salamander (<i>Aneides ferreus</i>); Del Norte Salamander (<i>Plethodon elongatus</i>); Foothill Yellow-legged Frog (<i>Rana boylei</i>); Northern Red-legged Frog (<i>Rana aurora</i>); Southern Torrent Salamander (<i>Rhyacotriton variegatus</i>); Western Toad</p> <p>Reptiles: Northwestern Pond Turtle (<i>Actinemys marmorata</i>); Western Painted Turtle</p> <p>Fish: Chum Salmon (<i>Oncorhynchus keta</i>); Coastal Cutthroat Trout (<i>Oncorhynchus clarkii clarkii</i>); Coho Salmon (<i>Oncorhynchus kisutch</i>); Eulachon (<i>Thaleichthys pacificus</i>); Fall and Spring Chinook; Green Sturgeon (<i>Acipenser medirostris</i>); Millicoma Dace (<i>Rhinichthys cataractae ssp.</i>); Pacific Lamprey (<i>Entosphenus tridentatus</i>); Summer/Winter Steelhead, Coastal Rainbow; Umpqua Chub (<i>Oregonichthys kalawatseti</i>); Western Brook Lamprey; Western River lamprey (<i>Lampetra ayresii</i>); White Sturgeon (<i>Acipenser transmontanus</i>)</p> <p>Invertebrates: Black Petaltail (<i>Tanypteryx hageni</i>); Pacific Walker (<i>Pomatiopsis californica</i>); Robust Walker (<i>Pomatiopsis binneyi</i>); Sisters Hesperian (<i>Hochbergellus hirsutus</i>); Western Ridged Mussel</p>
East Cascades	Alkaline Wetlands, Deciduous Swamps and Shrublands, Marshes,	<p>Birds: American White Pelican (<i>Pelecanus erythrorhynchos</i>); Greater Sandhill Crane (<i>Antigone canadensis tabida</i>); Long-billed Curlew; Red-necked Grebe (<i>Podiceps grisegena</i>); Trumpeter Swan; Yellow Rail (<i>Coturnicops noveboracensis noveboracensis</i>)</p>

	Vernal Pools, Wet Meadows, Wet Prairies	<p>Amphibians: Cascades Frog; Caspian Tern; Cope’s Giant Salamander; Oregon Spotted Frog (<i>Rana pretiosa</i>); Western Toad</p> <p>Reptiles: Northwestern Pond Turtle; Western Painted Turtle</p> <p>Fish: Bull Trout; Coho Salmon (Klamath SMU); Goose Lake Sucker (<i>Catostomus occidentalis lacusanserinus</i>); Great Basin Redband; Lost River Sucker (<i>Deltistes luxatus</i>); Miller Lake Lamprey (<i>Entosphenus minima</i>); Modoc Sucker (<i>Catostomus microps</i>); Pacific Lamprey; Pit Sculpin (<i>Cottus pitensis</i>); Shortnose Sucker (<i>Chasmistes brevirostris</i>); Spring and Fall Chinook Salmon; Summer Steelhead</p> <p>Invertebrates: Archimedes Springsnail (<i>Pyrgulopsis archimedis</i>); Black Petaltail; Crater Lake Tightcoil (<i>Pristiloma crateris</i>); Dall’s Ramshorn (<i>Vorticifex effuse dalli</i>); Great Basin Ramshorn (<i>Helisoma newberryi newberryi</i>); Highcap Lanx (<i>Lanx alta</i>); Klamath Ramshorn (<i>Vorticifex klamathensis klamathensis</i>); Lined Ramshorn (<i>Vorticifex effusa diagonalis</i>); Scale Lanx (<i>Lanx klamathensis</i>); Scalloped Juga (<i>Juga acutifilosa</i>); Sinitisin Ramshorn (<i>Vorticifex klamathensis sinitisini</i>); Siskiyou Hesperian (<i>Vespericola sierranus</i>); Turban Pebblesnail (<i>Fluminicola turbiniformis</i>)</p>
Northern Basin and Range	Alkaline Wetlands, Deciduous Swamps and Shrublands, Marshes, Vernal Pools, Wet Meadows	<p>Birds: American White Pelican; Black-necked Stilt (<i>Himantopus mexicanus</i>); Caspian Tern; Greater Sage-Grouse; Greater Sandhill Crane; Long-billed Curlew; Snowy Egret (<i>Egretta thula</i>); Trumpeter Swan; Western Snowy Plover</p> <p>Amphibians: Columbia Spotted Frog; Western Toad</p> <p>Fish: Alvord Chub (<i>Siphateles slyvordensis</i>); Borax Lake Chub (<i>Siphateles boraxobius</i>); Bull Trout; Foskett Spring Speckled Dace (<i>Rhinichthys osculus ssp.</i>); Great Basin Redband; Hutton Spring Tui Chub (<i>Siphateles bicolor oregonensis</i>); Lahontan Cutthroat Trout (<i>Oncorhynchus clarkii henshawi</i>); Pit Sculpin; Spring Chinook Salmon; Warner Sucker (<i>Catostomus warnerensis</i>)</p> <p>Invertebrates: Borax Lake Ramshorn (<i>Planorbella oregonensis</i>); Columbia Clubtail</p>
Willamette Valley	Deciduous Swamps and Shrublands, Marshes, Tidal Wetlands, Vernal Pools, Wet Prairies	<p>Birds: Dusky Canada Goose (<i>Branta canadensis occidentalis</i>)</p> <p>Amphibians: Cascade Torrent Salamander (<i>Rhyacotriton cascadae</i>), Clouded Salamander; Columbia Torrent Salamander; Foothill Yellow-legged Frog; Northern Red-legged Frog; Oregon Slender Salamander (<i>Batrachoseps wright</i>); Southern Torrent Salamander</p> <p>Reptiles: Northwestern Pond Turtle</p> <p>Fish: Bull Trout; Chum Salmon; Coastal Cutthroat Trout; Coho Salmon; Eulachon; Fall and Spring Chinook Salmon; Pacific Lamprey; Summer/Winter Steelhead/Coastal Rainbow; Western Brook Lamprey; Western River Lamprey; White Sturgeon</p> <p>Invertebrates: California Floater (<i>Anodonta californiensis</i>); Little Stonefly (<i>Capnia kersti</i>); Western Ridged Mussel; Winged Floater (<i>Anodonta nuttalliana</i>)</p>

This section includes the following for each wetland type: a general description, habitat information, and restoration approaches. Information on wetland classification, hydrology, and typical plant species for each wetland habitat type have been compiled in Table 2 at the end of the section. Information in this section has all been compiled using information from OCS (2016) ([see OCS wetlands website](#)); Christy, 2017a and 2017b ([see Major Wetland and Riparian Types website](#)); and Nature Serve ([NaturServ Explorer](#)) unless otherwise referenced.

1. Freshwater Wetlands (alkaline wetlands, deciduous swamps and shrublands, marshes, vernal pools, wet meadows, and wet prairies)

ALKALINE WETLANDS (see also: [Inter-Mountain Basins Alkaline Closed Depressions](#))

General description: Alkaline wetlands occur in intermittently saturated areas that form in depressions where alkaline soils occur. The water level can be perched due to restrictive soil layers, such as clay hardpan. These areas also accumulate salt in soil and water due to evaporation. Vegetation in these wetland types are primarily salt-tolerant species referred to as halophytes. Years with greater amounts of precipitation may result in areas that are lower in salt content where plants that are less salt-tolerant can grow.

Habitat: This wetland type is restricted to arid areas east of the Cascades. Habitat is found in flats and edges of alkaline lakes and ponds. Alkaline wetlands in Oregon are known as breeding or foraging sites for migrating birds. Many of the same species of plants and animals occur in both interior alkaline wetlands and estuarine wetlands along the coast. Unique animals associated with this wetland type are adapted to the intermittent hydrology and may emerge only every few years depending on available water and habitat conditions.

Restoration approaches: Alkaline wetlands are especially sensitive to changes in the water table, and the relationship between water availability and salinity is a key process to restore and maintain habitat quality. Commercial availability of seed for alkaline wetlands is limited. Establishment of a desirable plant community will likely require planting of commercially available species in combination with natural regeneration.

Typical restoration activities for alkaline wetlands include removal of surface or subsurface drainage features, removal of dikes and diversions obstructing runoff, and restoration of natural microtopography. Exceptional caution must be undertaken to avoid breaking through the restrictive soil layer with any earthmoving activities. Onsite investigation by a soil scientist is recommended.

Highly disturbed alkaline wetlands may be dominated by invasive species such as cheatgrass, meadow foxtail, and perennial pepperweed. These species must be controlled through mechanical and/or chemical means prior to undertaking any vegetative restoration activities. The primary long-term maintenance for alkaline wetland restoration includes control of invasive plant species. If sites are grazed, sites should be monitored closely to ensure disturbance by grazing animals does not adversely affect the plant community.

DECIDUOUS SWAMPS AND SHRUBLANDS (see also: [North Pacific Shrub Swamp](#))

General description: Deciduous swamps and shrublands occur in depressions, adjacent to lakes and ponds, and further out on river terraces in areas that are poorly drained, where organic, muck, or mineral soils and standing water occur. Forested wetlands differ from riparian forest in their higher water tables, longer duration of soil moisture, and finer-textured soils. They are typically flooded for several weeks during the growing season (seasonal flooding). Vegetation is dominated by trees such as ash species and woody shrubs such as willows, alders, and red-osier dogwood (*Cornus sericea*). Vegetation varies by area of the state and ecoregion (see Table 2 for Eastern versus Western Oregon differences). For example, in the Willamette Valley, typical shrub species can include Hooker's or Piper's willow (*Salix hookeriana*), Sitka willow (*S. sitchensis*), Pacific river willow (*S. fluviatilis*), redosier dogwood, Douglas spiraea (*Spiraea douglasii*), and salmonberry (*Rubus spectabilis*). In the Coast Range, Oregon crabapple (*Malus fusca*), twinberry (*Lonicera involucrata*), and honeysuckle (*Lonicera* spp.) may be important components of shrub swamps.

Habitat: Habitat type is edges of rivers, streams, lakes, and ponds that are poorly drained where deciduous trees and shrubs are dominant vegetation. In Deciduous Swamps where deciduous trees and shrubs both occur, the composition of the shrub and herbaceous layers depend partly upon the degree of canopy closure. Shrublands are dominated by shrub vegetation and a diversity of sedges, grasses, rushes, and forbs may also be present in the understory.

Restoration approaches: Shrub and herbaceous species that grow under a tree canopy in deciduous swamps or herbaceous species that grow in shrublands vary depending on the hydrologic regime. Therefore, recognition of the dominant sources of hydrology is critical prior to undertaking any restoration. Depending on the source of hydrology, restoring connections to groundwater and seasonal or intermittent flooding of rivers and streams is key. Elimination of surface and subsurface drainage features, control of undesirable vegetation, and establishment of appropriate trees, shrubs, and herbaceous vegetation will be part of any restoration plan.

Sites that have been or are adjacent to disturbed areas may be dominated by the invasive species found in wetlands such as reed canary grass (*Phalaris arundinacea*), so management of invasive plants may be a priority. Mechanical and/or chemical weed control around establishing woody vegetation to reduce competition will typically be required for a period of 3-5 years or until woody plants are able to outcompete adjacent vegetation. Protection of woody seedlings from depredation by elk, deer, beavers, voles, and mice will also need to be considered.

MARSHES

General description: Freshwater marshes or emergent marshes occur in depressions and around the edges of lakes, ponds, rivers, and streams where surface water or groundwater is consistently present and the soil poorly drained. Freshwater marshes in Oregon may form from riverine processes that left abandoned oxbow channels and old side channels of nearby river and stream systems. A wide variety of vegetation can grow in freshwater marshes depending on elevation microtopography, hydrology, drainage class, and soils. Emergent freshwater marshes are dominated by emergent wetland vegetation, which refers to vegetation that grows above the water surface such as sedges, bulrushes, spike rushes, rushes, and cattails. Broadleaf herbs and

shrubs may also be present. Fringes of freshwater marshes are often dominated by emergent plants. Portions of the marsh that are under water for long periods of time may have floating-leaf or deep-rooted plants within the littoral zone.

Habitat: Freshwater marshes are particularly well known as breeding, rest, and foraging sites for birds, nursery areas for a variety of fish species, and essential breeding habitat for amphibians. Habitats are found at the edges of rivers, streams, lakes, and ponds, and can be larger expanses in basins and flats.

Restoration approaches: Freshwater marshes occur in a wide variety of landscape positions and tend to have specific vegetation communities depending on hydrologic regime. Restoration hydrology targets will depend on if the marshes are lake associated, stream associated, or groundwater hydrology based. Water depth, soil conditions, and microhabitat will determine where specific plant species will grow.

Hydrology restoration is usually achieved through the decommissioning of surface and subsurface drains. In cases where the natural topography of freshwater marshes has been filled or leveled for agricultural production, it may be necessary to remove sediment and/or restore the natural microtopography of the site. For all earthmoving activities, it is important to remove and stock-pile the topsoil prior to excavating, then spread the topsoil back over the soil surface for the final grade to facilitate establishment of native vegetation and minimize problems with invasive weed encroachment and spread. It will also help maintain higher native plant richness (number of different species) and percent cover. After excavation and grading activities have been completed, vegetation restoration involves site preparation (most often, repeated chemical treatments depending on resident invasive plants) and seeding and/or planting of appropriate herbaceous plants. Availability of seed and plugs of native emergent woody and herbaceous plants may determine species to be planted.

Management of invasive species is especially critical during the establishment phase of planted vegetation. The primary invasive species that can cause significant problems in freshwater marshes is reed canarygrass, which readily forms monocultures in formerly disturbed sites and invades natural wetlands. Other invasive species include non-native *Phragmites* (*Phragmites australis*), purple loosestrife (*Lythrum salicaria*), and yellow flag iris (*Iris pseudacorus*). For freshwater marshes associated with grassland or other open communities where fragmentation of open habitat may be an issue, management of woody vegetation will need to occur periodically to prevent woody plants from taking over fringes of the marsh.

VERNAL POOLS

See also: [Western North American Vernal Pool, Columbia Plateau Vernal Pool](#)

General description: Vernal pools are one of the rarest wetland types in Oregon. They are found in depressions within shrub-steppe and open woodland areas of intermountain valleys and volcanic scablands of the Columbia Plateau. They form on layers of hardpan or basalt bedrock. Vernal pools can range in size from a few square meters to about an acre and are seasonally wet depending on local precipitation. Slow drying of these pools is characteristic of this wetland type and some pools only hold water every few years. Unique plants and animals associated with this wetland type have adapted to this hydrology.

Habitat: Typically, vernal pools occur in a “mound-depression” complex where the mounds are grassland, and the depressions are the vernal pools (Environmental Science Associates, 2007). Depressions or pools can be various sizes within steppe, woodland, or volcanic terrain. Distinct rings of vegetation may be present depending on water availability and drainage. These habitats are important for amphibians adapted to short timeframes of water availability. Species of flying insects and rare crustaceans, including fairy shrimps (Anostraca) are uniquely adapted to vernal pool ecosystems (Zedler, 2003).

Restoration approaches: Vernal pool restoration can be associated with wet prairie or wet meadow restoration, as these wetland types can occur within a wet prairie matrix (see also Wet Meadow, Wet Prairie sections below). Wetland features and functions that can be targeted for restoration for this wetland type include depth of pools and connectivity to water within a matrix of habitat patches. Micro-topography and variability of water are important factors to consider in restoration design. Hydrological restoration may involve breaking or disconnecting drainage tile or other sources of hydrological alteration. Linear swales to connect habitat patches and support wildlife dispersal and size diversity of vernal pools will increase habitat value for restoration designs (Environmental Science Associates, 2007). Restoration of native vegetation will generally follow the same approach as the associated wet meadow or wet prairie (see wet meadow and wet prairie restoration approaches sections below) within the same matrix as the vernal pool. Vernal pool-specific plant species will be targeted for planting in restorations if available.

WET MEADOWS

General description: Wet meadows typically occur in mountain valleys and can be found near headwaters of streams, seeps, lakes, and large river bottomlands. They may have shallow standing water for part of the year. Typical vegetation can include tufted hairgrass (*Deschampsia cespitosa*), sedges (*Carex* spp.), spikesedge (*Kyllinga* spp.), rushes (*Juncus* spp.), and forbs.

Habitat: This habitat type occurs along slopes, in basins, or in flats, mostly montane to subalpine and small, low-order streams can be present. This habitat type is considered important for migrating and breeding songbirds and waterbirds and other wetland-dependent wildlife. It has also been identified as significant for sage grouse breeding in the Western United States, as studies have shown that 85% of sage grouse breeding display locations (leks) occur within approximately 6 miles of wet meadows (mesic sites) (Donnelly et al., 2016).

Restoration approaches: Restoration approaches for this wetland type typically involves reconnecting natural hydrology that has been altered or reversing impacts of unmanaged grazing. Hydrology in wet meadows in eastern Oregon is often restricted by water rights. Where water rights are secured, hydrology in wet meadows is typically restored through plugging and filling of ditches, elimination of diversions, and stream habitat restoration and improvement.

Two to three years of chemical and mechanical site preparation are typically required prior to establishing native wet meadow species. If present, introduced pasture grasses, such as meadow foxtail, creeping foxtail, and Kentucky bluegrass are likely to remain in the absence of active herbaceous weed treatments. There is good availability of trees and shrubs of most of the dominant wet meadow species across Oregon. There is generally less availability of locally

adapted seed, container and plug stocks (woody and herbaceous) of wetland species for Eastern Oregon relative to the west side of the State. Without pre-planning to develop needed plant materials (seed and/or plugs, container stocks), many desirable species needed for restoration of these habitats will be unavailable. It is possible that donor wet meadows could be used to supply plugs or sod mats for reestablishment of native vegetation. Establishing vegetation may require herbaceous weed treatment practices such as spot spraying, especially to control dominant invasive species.

WET PRAIRIE

General description: Wet prairies typically occur on shallow soils over bedrock or clay soils within flooded areas of the valley floor that form a perched water table (Christy and Alverson, 2011). Typical vegetation is dominated by tufted hairgrass (*Deschampsia caespitosa*) and other grasses, rushes (*Juncus* spp.), and sedges (*Carex* spp.) (USFWS, 2017).

Habitat: Wet prairies occur as a mosaic of marsh, grasslands, and vernal pools; and were maintained historically by low intensity fire (USFWS, 2017). Habitat patches vary according to micro-topography (flats, irregular surfaces) and hydrology. Small depressions (micro lows) within the wet prairie mosaic may form vernal pool plant communities interspersed throughout a matrix of sedges and tufted hairgrass. Other wet prairie plant species that prefer slightly drier conditions may grow on micro highs.

Listed species and species of conservation concern associated with the Willamette Valley Wet Prairie system include northern harrier (*Circus cyaneus*), short-eared owl (*Asio flammeus*), American grass bug (*Acetropis americana*), Bradshaw's lomatium (*Lomatium bradshawii*), shaggy horkelia (*Horkelia congesta*), Nelson's checker mallow (*Sidalcea nelsoniana*), racemed goldenweed (*Pyrocoma racemose* var. *racemosa*), white-topped aster (*Sericocarpus rigidus*), and Willamette Valley daisy (*Erigeron decumbens*) (USFWS, 2017).

Restoration approaches: Hydrologic restoration involves plugging drainage ditches and breaking drainage tiles. The associated clay soils generally retain a restricted drainage layer to establish a perched water table and maintain wetland hydrology. Farming and land-leveling have removed much of the natural topography from former wet prairies, so construction of microtopographic features ($\leq 6''$ height) across a site can restore topographic variation where plant diversity can establish. Natural development of tussocks by sedges and tufted hairgrass will further diversify the natural topography of a site.

Wet prairie ecosystems require regular disturbance to persist, typically on a three-to-five-year disturbance interval. Prescribed fire is the desired type of disturbance as burns help control invading trees and shrubs and invigorate the native prairie plants. If prescribed burning is not feasible then mowing, haying, or grazing will need to be used to maintain disturbance that drives wet prairie ecosystem processes.

Restoration of the plant community will be dependent upon the starting condition of the vegetation community. Sites that are relatively weed free and contain an assemblage of native wet prairie species should be spot treated for weed species followed by interseeding to rejuvenate native plants. Sites that have been subject to intensive farming will need chemical (herbicide) and mechanical (mowing, disking, harrowing) treatments for a period of 2-3 years or more to

eliminate non-native plant cover and reduce the presence of non-natives in the soil seed bank. Broadcast seeding or drilling using native plant seed mixes will establish native vegetation across a diversity of hydrologic regimes at the site. Some species are better established by planting bulbs or plugs, such as camas (*Camassia quamash*), mule's ears (*Wyethia helianthoides*), and Oregon geranium (*Geranium oregonum*). Undesired species must be controlled after planting, generally two to three treatments per year (mechanical and chemical combined) to kill weeds or prevent weeds from going to seed while desired species become established.

2. Estuaries and Tidal Wetlands

SALTWATER MARSH

General description: Saltwater marshes occur in tidal zones of coastal estuaries. The saltwater marsh wetland type is distinguished from the tidal marsh wetland type by the salinity level in the water. When the salinity drops to less than 0.5 parts per thousand (ppt) as mixing with freshwater streamflow occurs, it is considered freshwater tidal marsh. The saltwater marsh is dominated by salt tolerant grasses and other herbaceous plants, while the freshwater tidal marsh is dominated by more shrubs and trees further away from the coast. The vegetation of saltwater marshes is divided into the low salt marsh zone that is more often inundated by saltwater and the high salt marsh zone with less saltwater inundation.

Habitat: The habitat type is edges of estuaries, tidal saltwater rivers and streams, and intertidal mudflats. These areas are often significant for production of shellfish and for spawning and rearing of saltwater fish. Mudflats are rich with invertebrates and seaweeds and are important feeding areas for seabirds and other waterbirds.

Restoration approaches: Saltwater intrusion, tidal influence, and intermixing are essential for saltwater marsh function and restoration. Surface elevation is a significant factor in saltwater marsh hydrology, and elevations where saltwater inundation occurs will determine where vegetation can establish (Frenkel and Morlan, 1991). Restoration may be as simple as removal of tide gates or removing or breaching existing levees. Projects that require levees and/or water control structures to avoid off-site impacts will likely require periodic maintenance to ensure continued function. Salt-water intrusion as part of restoration design would allow for establishment of a salt-loving plant community and dispersal of native saltmarsh plant propagules. Most non-native plants, such as invasive reed canarygrass, cannot tolerate salinities that occur in saltmarsh and will be replaced by native salt-tolerant plants.

TIDAL WETLANDS

General description: Freshwater tidal wetlands occur upstream of the tidal zone of estuaries in areas where salinity is less than 0.5 ppt. Distance inshore from the tidal influence varies for this wetland type and can range from 20 miles inland to 143 miles inland along the Columbia River. Freshwater tidal wetland plant communities tend to be more diverse and have more woody shrubs and trees compared to the saltwater marsh wetland type.

Habitat: Habitat type is edges of tidal rivers and streams. These areas are known to be important for fish rearing, feeding, and spawning habitats. For juvenile salmonids, these habitats provide low tide refuge and resting pools (Miller and Sadro, 2003), abundant large woody debris

(Diefenderfer and Montgomery, 2008), and abundant prey for feeding (Davis et al., 2019).

Restoration approaches: Restoration of freshwater tidal wetlands requires connection to rivers so that tidal influence and inundation from the floodplain can drive hydrology in the system. Hydrologic restoration may involve plugging drainage ditches, regrading or reshaping berms or meanders, or breaking or disconnecting drainage tiles to reconnect a stream to its floodplain. Beaver dam analogues may be installed along incised secondary channels to encourage aggradation. Restoration of microtopography may be critical in tidally influenced wetlands. Low mounds may be capable of supporting vegetation less tolerant to inundated conditions, which can greatly increase the diversity of the plant community. Caution must be taken not to create conditions for fish entrapment in depressions.

Restoration of vegetation is like the approach for the marsh wetland type. Site preparation involves a series of mechanical and chemical treatments prior to seeding and planting of appropriate herbaceous and woody plants, depending on the extent of non-native and invasive species dominance. Availability of seed and plugs of native emergent woody and herbaceous plants may determine species to be planted. Control of competitive vegetation and invasive species is critical during the establishment phase of planted native vegetation.

3. Flowing Water and Riparian

RIPARIAN WETLANDS

General description: Riparian wetlands typically occur along river corridors on stream terraces that have surface water connections during flood events (temporary flooding). Riparian vegetation communities are often dominated by trees or shrubs. These wetland types are essential for maintaining stream hydrology, nutrient processing, water temperature, and microclimate processes.

Habitat: These wetland types are important migratory corridors for many bird, amphibian, reptile, and mammal species. These wetlands can also maintain thermal regimes of streams for coldwater fish species and other aquatic species by maintaining thermal refugia or micro-climate cooling areas provided by tree canopy (Wohl et al., 2021; Olson and Burnett, 2009).

Restoration approaches: Historically, many riparian areas in Oregon were grazed intensively by livestock and soil compaction, streambank erosion, and introduction of invasive plant species may have occurred as a result. Removal of riparian buffer vegetation may also have resulted in changes to riparian micro-climate and thermal regimes. Restoration approaches for riparian wetlands should target reconnecting streams and rivers to their floodplains to maintain hydrology. Restoration of riparian vegetation will also result in removal of invasive species, stabilization of streambanks, and maintenance of thermal regimes via riparian shade and microclimate conditions.

Hydrologic restoration may involve eliminating levees, construction of setback levees, plugging drainage ditches, grading of banks and land surfaces (to allow more flooding), disconnecting drainage tile, and construction of oxbows and/or secondary channels. Historically, beavers played a large role in the dynamics of forested riparian wetlands. Consideration should be given

Table 2. Wetland type classification, hydrology, and plant species

Organized by wetland type categories: **FW**=Freshwater Wetlands; **ETW**=Estuaries and Tidal Wetlands; **FWR**=Flowing Water and Riparian

Wetland Type	Wetland Classification (HGM)	Hydrology	Plant Species
FW -Alkaline wetlands	DEPRESSIONAL	Sources may be direct precipitation and runoff; perennially to seasonally flooded	Black greasewood (<i>Sarcobatus vermiculatus</i>), inland saltgrass (<i>Distichlis spicata</i>), alkali grasses (<i>Puccinellia lemmonii</i>), bluegrass (<i>Poa secunda</i>), <i>Muhlenbergia</i> spp., beardless wildrye (<i>Leymus triticoides</i>), alkali bulrush (<i>Bolboschoenus maritimus</i>), chairmaker's bulrush (<i>Schoenoplectus americanus</i>), seaside arrow-grass (<i>Triglochin maritima</i>), and glasswort (<i>Salicornia</i> spp.)
FW -Deciduous swamps and shrublands	Multiple HGM classes: at elevation of stream: RIVERINE; if groundwater or runoff sources, DEPRESSIONAL; if groundwater source only, SLOPE HGM; if mineral soils, precipitation source, MINERAL SOIL FLATS; if brackish water, ESTUARINE FRINGE	Sources may be groundwater or streamflow; seasonal, or temporary flooding	Western Oregon: Trees-Oregon ash (<i>Fraxinus latifolia</i>), black cottonwood (<i>Populus balsamifera ssp. trichocarpa</i>), Pacific willow (<i>Salix lasiandra</i>), red alder (<i>Alnus rubra</i>), and western redcedar (<i>Thuja plicata</i>); Shrubs- Douglas spiraea (<i>Spiraea douglasii</i>), willows (<i>Salix hookeriana</i> , <i>S. sitchensis</i> , <i>S. fluviatilis</i>), vine maple (<i>Acer circinatum</i>), red-osier dogwood (<i>Cornus sericea</i>), common snowberry (<i>Symphoricarpos albus</i>), trailing blackberry (<i>Rubus ursinus</i>), salmonberry (<i>Rubus spectabilis</i>), and roses (<i>Rosa nutkana</i> , <i>R. pisocarpa</i>); Herbaceous-slough sedge (<i>Carex obnupta</i>), lady fern (<i>Athyrium filix-femina</i>), skunk cabbage (<i>Lysichiton americanus</i>), and water parsley (<i>Oenanthe sarmentosa</i>). Eastern Oregon: Trees-black cottonwood, quaking aspen (<i>Populus tremuloides</i>), white alder (<i>Alnus rhombifolia</i>), Lodgepole pine (<i>Pinus contorta</i>), Engelmann spruce (<i>Picea engelmannii</i>); Shrubs-Lewis' mockorange (<i>Philadelphus lewisii</i>), Wood's rose (<i>Rosa woodsia</i>), common snowberry, western serviceberry (<i>Amelanchier alnifolia</i>), willows (<i>Salix</i> spp.), black hawthorn (<i>Crataegus douglasii</i>); Herbaceous-blue wildrye (<i>Elymus glaucus</i>), Canada bluejoint (<i>Calamagrostis canadensis</i>), sedges (<i>C. angustata</i> , <i>C. aquatilis</i>), cow parsnip (<i>Heracleum lanatum</i>), large-leaved avens (<i>Geum macrophyllum</i>), stinging nettle (<i>Urtica dioica</i>), and lady fern. Groundwater driven system: (<i>Carex laeviculmis</i>), mannagrasses (<i>Glyceria elata</i> , <i>G. striata</i>), skunk cabbage, and

			water parsley.
FW-Marshes	If elevation of stream, RIVERINE; if groundwater or runoff, DEPRESSIONAL; if groundwater only, SLOPE HGM; if mineral soils, MINERAL SOIL FLATS; if lakes, LACUSTRINE FRINGE	Perennially to seasonally flooded	spikerush (<i>Eleocharis spp.</i>), sedges (<i>Carex spp.</i>), bulrushes (<i>Schoenoplectus spp.</i>), bur-reeds (<i>Sparganium emersum</i> Rehmman, <i>Sparganium eurycarpum</i> Engelm., <i>Sparganium natans</i> L.), cattails (<i>Typha spp.</i>); areas that dry out: beggar ticks (<i>Bidens spp.</i>), smartweeds (<i>Polygonum spp.</i>), and curvepod yellowcress (<i>Rorippa curvisiliqua</i>); deeper areas: spatterdock or wocas (<i>Nuphar polysepala</i>), floating burred (<i>Sparganium augustifolium</i>), and wapato (<i>Sagittaria latifolia</i>).
FW-Vernal Pools	Mineral soils, precipitation source, MINERAL SOIL FLATS	Intermittently to seasonally flooded	Annual hairgrass (<i>Deschampsia danthonioides</i>), elegant calicoflower (<i>Downingia elegans</i>), waterwort (<i>Elatine spp.</i>), spikerush (<i>Eleocharis spp.</i>), denseflower willowherb (<i>Epilobium densiflorum</i>), Oregon eryngo (<i>Eryngium petiolatum</i>), inch-high rush (<i>Juncus uncialis</i>), tiny mousetail (<i>Myosurus minimus</i>), white flowered navarretia (<i>Navarretia leucocephala ssp. diffusa</i>), American pillwort (<i>Pilularia americana</i>), popcorn flower (<i>Plagiobothrys spp.</i>), dense-flowered knotweed (<i>Polygonum polygaloides ssp. confertiflorum</i>), milkwort knotweed (<i>Polygonum polygaloides ssp. polygaloides</i>), dwarf woollyheads (<i>Psilocarphus brevissimus</i>), tall woollyheads (<i>Psilocarphus elatior</i>), Oregon woollyheads (<i>Psilocarphus oregonus</i>), bowl clover (<i>Trifolium cyathiferum</i>), fool's onion (<i>Triteleia hyacinthine</i>), and purslane speedwell (<i>Veronica peregrina</i>).
FW-Wet Meadows	SLOPE	Groundwater or snowmelt typically largest source; Perennially to seasonally flooded	If saturation throughout growing season, dominated by sedges like shortbeak sedge (<i>Carex simulata</i>); if dry later in growing season, often tufted hairgrass (<i>Deschampsia caespitosa</i>), Baltic rush (<i>Juncus arcticus</i>), Nebraska sedge (<i>C. nebrascensis</i>), and widefruit sedge (<i>C. angustata</i>) in areas that have been grazed.
FW-Wet Prairie	MINERAL SOIL FLATS, DEPRESSIONAL or SLOPE	Intermittently to seasonally flooded.	Tufted hairgrass (<i>Deschampsia caespitosa</i>), spike bentgrass (<i>Agrostis exarata</i>), California oatgrass (<i>Danthonia californica</i>), dense sedge (<i>Carex densa</i>), green-sheath sedge (<i>C. feta</i>), thick-headed sedge (<i>C. pachystachya</i>), and lateral sedge (<i>C. unilateralis</i>). Toad rush (<i>Juncus bufonius</i>), western rush (<i>J. occidentalis</i>) and Bolander's rush (<i>J. bolanderi</i>),

			camas (<i>Camassia quamash</i>), slender cinquefoil (<i>Potentilla gracilis</i>), self-heal (<i>Prunella vulgaris</i>), skullcap speedwell (<i>Veronica scutellata</i>) and Oregon geranium (<i>Geranium oreganum</i>).
ETW -Saltwater Marsh	ESTUARINE FRINGE	Tidal, regularly to irregularly flooded	Low salt marsh (zone nearest ocean): pickleweed (<i>Salicornia</i> spp.), marsh jaumea (<i>Jaumea carnosa</i>), saltgrass (<i>Distichlis spicata</i>), Lyngbye's sedge (<i>Carex lyngbyei</i>), and arrowgrass (<i>Triglochin</i> spp.). High salt marsh (less frequent inundation by saltwater): tufted hairgrass (<i>Deschampsia caespitosa</i>), Baltic rush (<i>Juncus arcticus</i>), pacific silverweed (<i>Potentilla pacifica</i>), and orache (<i>Atriplex patula</i>).
ETW -Tidal Wetlands (freshwater)	ESTUARINE FRINGE	Tidal, regularly to irregularly flooded	Sitka spruce (<i>Picea sitchensis</i>), western redcedar (<i>Thuja plicata</i>), western hemlock (<i>Tsuga heterophylla</i>), Hooker's willow (<i>Salix hookeriana</i>), Oregon crabapple (<i>Malus fusca</i>), twinberry honeysuckle (<i>Lonicera involucrata</i>), red-osier dogwood (<i>Cornus sericea</i>), Douglas spiraea (<i>Spiraea douglasii</i>), western Labrador tea (<i>Ledum glandulosum</i>), Salal (<i>Gaultheria shallon</i>), willows (<i>Salix</i> spp.), slough sedge (<i>Carex obnupta</i>), and skunk cabbage (<i>Lysichiton americanus</i>).
FWR -Riparian Wetlands	RIVERINE	Streamflow, groundwater at stream elevation; Perennially to seasonally flooded	Bigleaf maple (<i>Acer macrophyllum</i>), red alder (<i>Alnus rubra</i>), black cottonwood (<i>Populus balsamifera</i> ssp. <i>trichocarpa</i>), Sitka willow (<i>Salix sitchensis</i>), Pacific willow (<i>Salix lucida</i> ssp. <i>lasiandra</i>), red-osier dogwood (<i>Cornus sericea</i>), and Oregon ash (<i>Fraxinus latifolia</i>)

to encouraging beaver activity wherever site conditions allow. This can be achieved through low technology riparian restoration techniques such as beaver dam analogs (BDAs) (see Pilliod et al., 2018; Pollock et al., 2023), or simply by allowing beaver activity to occur.

Reed canarygrass (*Phalaris arundinacea*) has been widely planted for livestock forage, and this species is particularly abundant in riparian wetland systems, as its seed is readily transported by water movement. Kentucky bluegrass (*Poa pratensis*) may form the dominant ground cover in east-side riparian wetlands that have been modified for grazing or been subject to stream degradation through beaver removal, removal of natural riparian vegetation, and/or stream channelization. This rhizomatous species takes hold in areas with lowered water tables and can be difficult to control. Meadow foxtail (*Alopecurus pratensis*) and creeping meadow foxtail (*Alopecurus arundinaceus*) have also been extensively planted for livestock forage and have the potential to persist and/or invade new wetland areas. The best method of long-term control is restoration of the natural hydrologic regime and establishment of a canopy of woody vegetation to provide competition for light. Extensive chemical and mechanical weed control may also be needed before attempting to revegetate infested areas.

III. WRE PROGRAM ELIGIBILITIES

A. Eligible Land Types

All land is evaluated and determined as eligible per the ACEP manual criteria ([Conservation Program Manual \(CPM\), Title 440, Part 528, Section 528.105](#)). Any land not meeting the eligible land criteria described in this section or upon review of National policy, that does not meet the criteria for “adjacent lands” (see Section III, B Eligible Adjacent Lands) is considered ineligible for ACEP-WRE. Land eligibility categories may vary in different areas of Oregon.

The following includes a summary of information on eligible land types that are typically enrolled in ACEP-WRE in Oregon: Farmed, Converted, Degraded Wetlands; Riparian Areas; Lands in the Conservation Reserve Program (CRP); and Wetlands Restored or Protected Under a Private, State, or Federal Program.

1. **Farmed or Converted Wetlands** (See [CPM, Title 440, Part 528, Section 528.105\(C\)](#))

Farmed wetlands: Farmed under natural conditions, farmed wetlands, prior converted cropland, commenced conversion wetlands, or farmed wetland pasture. NRCS makes this determination based on Title 180, National Food Security Act Manual (NFSAM) criteria.

Converted wetlands: Land is not eligible for enrollment if the conversion was after December 23, 1985, except as provided for in [CPM, Title 440, Part 528, Section 528.105\(D\)\(6\)](#). Enrollment may be considered if all the following are true:

- a. Enrollment is needed for administrative management of boundaries.
- b. Economic and management needs of parcels for the landowner.
- c. The landowner is willing to enroll for no compensation.
- d. The landowner is willing to complete the restoration at the landowner's expense.

Degraded wetlands: The extent of degradation and the restorability of the site are considered from a hydrology standpoint to determine eligibility. For NRCS, Oregon, the site will be considered significantly degraded and eligible if more than 75 percent of the hydrology functions (e.g., surface area, depth, frequency, timing, or duration) have been altered from its historic hydrological conditions. In addition, the site will be considered substantially restorable and eligible if more than 90 percent of the identified wetland hydrology functional impacts will be restored.

2. Riparian Areas (See [CPM, Title 440, Part 528, Section 528.105\(E\)](#))

Riparian areas are eligible if they connect wetlands less than one mile apart that are currently enrolled in an ACEP-WRE program or another resource protection program, such as a State or Federal wildlife management area. If the riparian area will link already-protected wetland areas, then no additional wetland acres are required to enroll the riparian acres. Eligible riparian areas should average up to 300 feet wide maximum on one side or 600 feet for both sides. Enrollments with widths greater than 300 feet on one side and distances greater than 1 mile between enrolled wetlands may be granted a waiver by the State Conservationist (see Section V. Technical Considerations for Waivers, Riparian Widths and Distances Waiver). Greater widths and distances for riparian area enrollments may be determined based on wildlife habitat and life cycle needs and wetland ecological functions and values (see VII. List of Resources for Further Information, Wetland-Dependent Wildlife and Wetland Functions and Values). Riparian area occurrence and the boundary of the riparian areas can be determined by using soil survey information, hydrology-based indicators, and vegetation-based indicators (see VII. List of Resources for Further Information, Riparian Areas).

3. Lands in the CRP (See [CPM, Title 440, Part 528, Section 528.105\(F\)](#))

A landowner can submit a request to the Assistant State Conservationist for Programs (ASTC-P) and State Resource Conservationist (SRC) to participate in WRE program. In general, land that is currently enrolled in CRP can be enrolled in ACEP-WRE, but eligibility should be determined on a case-by-case basis related to landowner intentions for discontinuation of agricultural production. In addition, landowner applicants should consult FSA if they wish to enroll land in ACEP-WRE that is currently enrolled in CRP to avoid paying back previous rental payments during land enrollment under CRP.

It should be noted that lands established to trees under CRP are ineligible for enrollment, whether the contract is active or closed. However, the State Conservationist may grant a waiver for ACEP-WRE eligibility if trees have not been completely established and the presence of trees conforms to ACEP-WRE restoration requirements (see [CPM, Title 440, Part 528, Section 528.106B\(2\)](#) for further detail). For information on methods to determine if trees have been established, refer to: [Title 190 Forest Inventory Methods Technical Note](#).

4. Wetlands Restored or Protected Under a Private, State, or Federal Program

As described in [CPM, Title 440, Part 528, Section 528.105\(G\)](#), land previously restored under a local, State, or Federal program are eligible that applied restoration methods meeting NRCS restoration standards and specifications to provide benefits for wetland functions.

If a deed restriction is held by another federal agency, an agreement must be reached and documented by NRCS and OGC before NRCS may proceed with enrollment of the property into the ACEP-WRE program. The enrollment must provide additional resource protection, provide critical habitat for threatened or endangered species, and/or allow an opportunity for full restoration of wetland functions because of enrollment that could not otherwise be achieved.

B. Eligible Adjacent Lands

Lands that do not fall within the categories described above that will be considered for enrollment as eligible Adjacent Lands (see [CPM, Title 440, Part 528, Section 528.105\(I\)](#)) will emphasize habitats for wetland-dependent wildlife (“Associated Habitats”) that are within dispersal distances to utilize the habitat to complete their life cycle. Other functions that wetlands provide that are required for aquatic and wetland ecosystem processes, such as water availability and hydrologic stability, will be considered for enrollment as Adjacent Lands. Eligibility will be determined using technical information for justification to ensure wetland functions and values and habitat objectives are achieved (see VII. List of Resources for Further Information, Wetland Functions and Values and Wetland-Dependent Wildlife).

An upper limit of 5:1 ratio (5 acres eligible adjacent land to 1 acre eligible land acre) will be considered for enrollment in the WRE program throughout Oregon. This ratio will be determined based on wetland type and functional targets for wetland restoration on a case-by-case basis. The following examples illustrate how enrolling adjacent lands may increase the benefits of functions that enrolled wetlands provide:

1. Wetlands in one location may provide water source and hydrology functions to support other wetlands in the area (wetland complexes depend on hydrology on adjacent lands).
2. Riparian areas may connect multiple wetlands along the same stream network.
3. Aquatic or wetland-dependent wildlife species may require connected patches of riparian or wetland habitat to complete their life cycles.

Specific Examples to illustrate the need for a 5:1 maximum ratio:

1)Example 1: The wet prairie habitat is adjacent to the oak woodland. The wet prairie (eligible) is breeding habitat for the amphibian egg and larval stage. The oak woodland (adjacent) provides habitat for the amphibian adult life stage.

2)Example 2: There are multiple freshwater marshes along a river corridor. Both marshes (eligible) are along the same river within adjacent, separate easements. The riparian area (adjacent) off easement provides connectivity between the marshes (eligible) on two separate easements.

3)Example 3: A hydrology, groundwater seep source exists off easement. A riparian wetland, floodplain bottomland (eligible) is downstream of a groundwater source off easement (adjacent). The riparian wetland needs the groundwater seep that is upstream as a hydrology source to function.

Acceptable Associated Habitats

The following table provides details on eligible Adjacent Lands that are considered “Associated Habitat” types to be considered because they may provide additional habitat functions for wetland-dependent wildlife and other aquatic species. These may be considered with written approval from the State Conservationist. This list of associated

habitats applies to new enrollments in ACEP-WRE, existing enrollments (unclosed), and closed conservation easements under ACEP-WRE and predecessor programs. Any maintenance, management, or additional restoration after initial restoration that is completed on an ACEP-WRE easement must be consistent with these associated habitats.

Acceptable Associated Habitat	Expected Contribution to Wetland Functions & Values
Grasslands	Buffer areas to wetlands, riverine habitat, perennial and intermittent streams. Provides for wildlife cover, forage, nesting, and dispersal.
Riparian Areas	Lands that occur along watercourses and water bodies (e.g., flood plains, streambanks) with unique soil and vegetation characteristics influenced by the presence of water. Typical vegetation consists of woody species that benefit multiple wildlife species and provides shade to maintain temperature. Acts as a buffer zone for riverine areas and adjacent wetlands.
Shrublands	Cover and forage areas for migratory and nesting birds.
Forestland	Cover, nesting, and forage areas for migrating birds.
Other Aquatic Priority Habitats	Connectivity to other wetland and riparian habitats, dispersal to habitats for spawning or breeding.

C. Eligible Conservation Practices

The NRCS umbrella conservation practices that are eligible to implement on wetland easements include: Wetland Restoration (657), Wetland Wildlife Habitat Management (644), Wetland Enhancement (659), and Upland Wildlife Habitat Management (645). However, any reasonable practice or activity needed to restore, manage, and/or enhance these lands is eligible if consistent with program goals and objectives.

D. Ranking Factors and Criteria

Although much of the ranking criteria are established according to the most recent Farm Bill (2018) and set Nationally, the States have some flexibility to establish their own criteria as long as they do not violate policy and provide environmental benefits per [CPM, Title 440, Part 528, Section 528.111](#). Environmental benefits from wetland restoration can be related to native plant community and habitat, such as providing greater habitat diversity, complexity, and connectivity within the landscape. Wetland restoration can also provide habitat benefits for specific groups or species, including migratory birds, wetland-dependent wildlife, threatened, endangered, and at-risk species. Environmental benefits from wetland restoration can also focus on water quality and quantity to maintain and restore hydrology (surface and groundwater) and manage flood flows. Environmental benefits of wetland functions related to carbon sequestration and climate resiliency can also be considered in ranking criteria.

Oregon NRCS determines ranking criteria on an annual basis, and a ranking worksheet is used to give each WRE application a ranking score. These scores are used to prioritize potential enrollments for funding. The only limitation on scoring is that 50% of the potential points

awarded for environmental benefits must come from hydrology restoration potential (see Appendix B, Ranking Worksheet).

Ranking criteria for NRCS Oregon that quantify environmental benefits on the Ranking Worksheet (Appendix B) result in scores for the following: altered hydrology, restored hydrology, habitat for at risk species, and restoration of native plant communities. These ranking criteria are designed to reflect the following wetland functions: maintenance of surface hydrology, maintenance of groundwater flow, maintenance of wetland and hydric soils, habitat for wetland-dependent wildlife, availability of native plant diversity, and availability of habitat. Diversity of wetland types and connectivity of wetland habitat patches within the landscape are also reflected in ranking criteria (Habitat Diversity, Adjacent Protected Habitat, and Floodplain Connectivity). Various geospatial data layers and NRCS technical staff are potential resources available for assessment of wetland functions. Further details on the technical information used for assessment of wetland functions are listed below in Table 3 and can be found in section VII. List of Resources for Further Information, Data Sources and Wetland Functions and Values.

Table 3. Ranking criteria and associated wetland functions

Ranking Criteria	Wetland functions	Technical Information Used*
Altered and Restored Hydrology	Maintain surface hydrology	NWI data, OR Wetlands data, FEMA Floodplain data, NHD data, LiDAR data
Altered and Restored Hydrology	Maintain groundwater flow	NWI data, OR Wetlands data, NHD data, LiDAR data
Altered and Restored Hydrology	Maintain wetland and hydric soils	NWI data, OR Wetlands data, WebSoilSurvey; NRCS Soil Scientists
Habitat for At Risk Species	Habitat for wetland-dependent wildlife	Presence data, Habitat assessment data, WHEG, NRCS and USFWS Biologists
Native Plant Community	Maintain native plant diversity, habitat	Presence data (plant survey); community structure assessment, NRCS Biologists

* (see VII. List of Resources for Further Information, Data Sources and Wetland Functions and Values)

In addition to information recorded on the ranking worksheet, special considerations, Priority Areas for ranking that are currently being used by NRCS Oregon include: the USFWS Recovery Zone for Bradshaw’s lomatium or Nelson’s checker mallow; and the Lower Columbia River Priority Area, Priority watersheds for Oregon Coastal Coho Salmon (see Appendix C for Priority Geographic Regions for FY 2023).

IV. TECHNICAL CONSIDERATIONS FOR CUAs

A compatible use (CUA) is a use or activity conducted on a wetland reserve easement that NRCS Oregon determines is consistent with the long-term conservation and restoration goals and objectives, and wetland functions and values of the easement area. An NRCS-CPA-52 Environmental Evaluation is required for all practices and activities in a CUA in accordance with Oregon State Instruction 300 Part 393. NRCS Oregon will consider CUAs on a case-by-case basis, and CUAs will be approved by the Assistant State Conservationist or Basin Team Leaders. Each CUA is a discreet authorization subject to requirements at the time it was issued and expiration date. Technical information and parameters used for CUA assessment will be

considered and developed by NRCS Oregon using an adaptive management approach and in partnership with USFWS, SWCDs, Oregon State agency partners, and others (see VII. List of References for Further Information, Compatible Use Authorization Management Tools).

NRCS Oregon CUAs will consider geographic area, activity, and wetland type to prescribe amount, method, location, timing, frequency, intensity, and duration of the compatible use per [CPM, Title 440, Part 528, Section 528.152\(B\)\(1\)](#). NRCS Oregon CUAs must not adversely affect habitat for migratory birds, at-risk species, or threatened or endangered species. Timing of activities associated with CUAs need to be considered on a case-by-case basis according to species and part of the state.

V. TECHNICAL CONSIDERATIONS FOR WAIVERS

The State Conservationist is authorized to issue waivers based on technical considerations. All other requested policy waivers can only be approved or denied by National Headquarters. The landowner may submit the request for a waiver in writing and must acknowledge certain conditions of the waiver. More information can be found at [CPM, Title 440, Part 528, Section 528.142\(D\)](#). Program requirements covered by the statute or the rule may not be waived. Only waiver options consistent with program policy, contained within this WRCG, or consistent with this WRCG may be requested. Existing waivers issued by the State Conservationist prior to the approval and publication of this WRCG will be allowed until expiration of such existing waiver.

Riparian Widths and Distances Waiver

[Section III. B. Eligible Land Types, Riparian Areas](#) describes the “riparian” eligible land category. Waivers for additional linkages between wetland areas, widths, or for eligible wetland areas more than 1 mile apart may be granted by the State conservationist if the riparian area can provide habitat for at-risk fish or wildlife, contribute significantly to wetland functions and values of the easement area, or improve the practical administration and management of the easement area.

Excessive Restoration Costs

Lands where the cost of restoration for the easement area will exceed the fair market value of the land are ineligible. This criterion may be waived by the State Conservationist in situations in which it is documented that the restoration may be successfully accomplished without accumulating a long-term operation and maintenance cost burden to the program. These may include habitat types that are highly degraded and are labor intensive and expensive to restore.

Early Implementation of Restoration

In general, payments are not authorized for restoration practices that are started or completed before easement recording and easement restoration agreement approval date. In very special cases and for meritorious reasons only, the State Conservationist may consider a waiver for enrollments that meet all ACEP-WRE land and landowner eligibility requirements. Meritorious reasons may include:

- Alleviation of imminent and significant environmental problems.
- Prevention of damage to life or property.
- Seasonal weather constraints.

VI. LITERATURE CITED

Adamus, P.R. 2006. Hydrogeomorphic (HGM) assessment guidebook for tidal wetlands of the Oregon Coast, part 1: rapid assessment method. Report to Coos Watershed Association, US Environmental Protection Agency, and Oregon Department of State Lands, Salem. 85 pp.

Adamus, P.R. 2001. Guidebook for hydrogeomorphic (HGM)-based assessment of Oregon wetland and riparian sites: Statewide Classification and Profiles. Oregon Division of State Lands, Salem, OR.

Akins, G.J. 1970. The effects of land use and land management on the wetlands of the upper Klamath Basin. M.S. Thesis. Western Washington State College, Bellingham, WA.

Boyd, R. 1986. Strategies of Indian burning in the Willamette Valley. *Canadian Journal of Anthropology* 5:65-86.

Brinson, M.M. 1993. A hydrogeomorphic classification for wetlands. U.S. Army Corps of Engineers. Wetlands Research Program Technical Report WRP-DE-4. Waterways Experiment Station, Vicksburg, MS. 101 pp.

Brophy, L.S. 2019. Comparing historical losses of forested, scrub-shrub, and emergent tidal wetlands on the Oregon coast, USA: A paradigm shift for estuary restoration and conservation. Prepared for the Pacific States Marine Fisheries Commission and the Pacific Marine and Estuarine Fish Habitat Partnership. Estuary Technical Group, Institute for Applied Ecology, Corvallis, Oregon, USA. 88 pp.

Christy, J.A. 2017a. Key to major wetland and riparian habitats in Oregon. Oregon Biodiversity Information Center, Institute of Natural Resources, Oregon State University, 1 page.

Christy, J.A. 2017b. Keys to wetland and riparian plant associations in Oregon. Oregon Biodiversity Information Center, Institute of Natural Resources, Oregon State University, 34 pp.

Christy, J.A. and E.R. Alverson. 2011. Historical vegetation of the Willamette Valley, Oregon, circa 1850. *Northwest Science* 85:93-107.

Christy, J.A. 2010. Oregon Biodiversity Information Center and The Wetlands Conservancy. *History of Oregon's Wetlands*. <https://oregonexplorer.info/content/history-oregons-wetlands?topic=4138&ptopic=98>

Christy, J., E. Alverson, M. Dougherty, S. Kolar, L. Ashkenas, and P. Minear. 2000. Presettlement vegetation for the Willamette Valley, Oregon, version 4.0, compiled from records of the General Land Office Surveyors (c. 1850), Oregon Natural Heritage Program, Portland, OR, USA.

Christy, J.A. and J.H. Titus. 1997. Native wetland plant communities and associated sensitive, threatened, or endangered plant and animal species in Oregon. Oregon Natural Heritage Program, Portland, OR.

Connolly, T.J. 2000. Anthropological and archaeological perspectives on native fire management of the Willamette Valley. Paper presented at the 81st Annual Meeting of the American Association for the Advancement of Science, Pacific Division (Symposium: Fire History in the Pacific Northwest: Human and Climatic Influences), June 11-14, 2000, Ashland, OR.http://www.orwww.org/Kalapuya-Amin_2006/Program/ConnollyT/Indian_Burning_20000611.pdf

Cowardin, L. M., Carter, V., Golet, F. C., and LaRoe, E.T. (1979). Classification of wetlands and deepwater habitats of the United States, U.S. Fish and Wildlife Service, Washington, DC.

Crowe, E.A., B.L. Kovalchik, and M. Kerr. 2004. Riparian and wetland vegetation of central and eastern Oregon. Oregon Natural Heritage Information Center, Institute for Natural Resources, Oregon State University. 474 pp. + appendices.

Dahl, T.E. 1990. Wetland losses in the United States, 1780s to 1980s. U.S. Fish and Wildlife Service Report to Congress, Washington, D.C.

Davis, M.J., I. Woo, C.S. Ellings, S. Hodgson, D.A. Beauchamp, G. Nakai, and S.E.W. De La Cruz. 2019. Freshwater tidal forests and estuarine wetlands may confer early life growth advantages for Delta-reared Chinook salmon. *Transactions of the American Fisheries Society* 148:289–307.

Diefenderfer, H.L. and D.R. Montgomery. 2008. Pool spacing, channel morphology, and the restoration of tidal forested wetlands of the Columbia River, U.S.A. *Restoration Ecology* 17:158-168.

Donnelly, J. P., D. E. Naugle, C. A. Hagen, and J. D. Maestas. 2016. Public lands and private waters: scarce mesic resources structure land tenure and sage-grouse distributions. *Ecosphere* 7(1):e01208. 10.1002/ecs2.1208

Environmental Science Associates. 2007. Agate Desert Vernal Pool Functional Assessment Methodology. Environmental Science Associates, Sacramento, CA.

Fickas, K. C., Cohen, W. B., and Z. Yang. 2016. Landsat-based monitoring of annual wetland change in the Willamette Valley of Oregon, USA from 1972 to 2012. *Wetlands Ecology and Management* 24: 73-92.

Fretwell, J.D., J.S. Williams, and P.J. Redman, compilers. 1996. National water summary on wetland resources. U.S. Geological Survey. Water-supply paper 2425. Washington, D.C.

Gilmer, D.S., Yee, J.L., Mauser, D.M., and Hainline, J.L., 2004, Waterfowl migration on Klamath Basin National Wildlife Refuges 1953-2001: U.S. Geological Survey, Biological

Resources Discipline Biological Science Report USGS/BRD/BSR—2003-0004, 66 pp.

Good, J.W. 2000. Summary and current status of Oregon's estuarine ecosystems. In: Oregon State of the Environment Report 2000 (Risser, P.G.). SOER Science Panel.

Hamman, S.T., P.W. Dunwiddie, J.L. Nuckols, and M. McKinley. 2011. Fire as a restoration tool in the Pacific Northwest prairies and oak woodlands: challenges, Successes, and future directions. *Northwest Science* 85:317-328.

Hulse, D., S. Gregory, and J. Baker (Eds.) 2002. Willamette River Basin planning atlas: trajectories of environmental and ecological change. 2nd edition. The Pacific Northwest Ecosystem Research Consortium. Oregon State University Press, Corvallis, OR. 192 pp.

Johannessen, C.L., W.A. Davenport, A. Millet, and S. McWilliams. 1971. The vegetation of the Willamette Valley. *Annals of the Association of American Geographers* 61:286-302.

Kagan, J.S., J.A. Christy, M.P. Murray, and J.A. Titus. 2004. Classification of native vegetation of Oregon. Oregon Natural Heritage Information Center, Institute of Natural Resources, Oregon State University, 52 pp.

Marcoe, K. and S. Pilson. 2017. Habitat change in the lower Columbia River estuary, 1870–2009. *Journal of Coastal Conservation* 21(4): 505-525.

Mayer, T. D. 2005. Water-quality impacts of wetland management in the Lower Klamath National Wildlife Refuge, Oregon and California, USA. *Wetlands* 25(3): 697-712.

Miller, B.A. and S. Sadro. 2003. Residence time and seasonal movements of juvenile coho salmon in the ecotone and lower estuary of Winchester Creek, South Slough, Oregon. *Transactions of the American Fisheries Society* 132: 546-559.

Morlan, J. C., E. F. Blok, J. Miner, and W. N. Kirchner, 2010. Wetland and land use change in the Willamette Valley, Oregon: 1994 to 2005. U.S. Fish and Wildlife Service, Portland, Oregon, and Oregon Department of State Lands, Salem, Oregon.

Morlan, J.C. 2000. Summary of current status and health of Oregon's freshwater wetlands. In: Oregon State of the Environment Report 2000. Oregon Progress Board, Salem, OR. pp. 45–52.

Olson, D.H. and K. M. Burnett. 2009. Design and management of linkage areas across headwater drainages to conserve biodiversity in forested ecosystems. *Forest Ecology and Management* 258S: S117-S126.

Omernik, J.M. 1987. Ecoregions of the conterminous United States: *Annals of the Association of American Geographers* 77 (1): 118–125.

Oregon Conservation Strategy. 2016. Oregon Department of Fish and Wildlife, Salem, Oregon.

Oregon Progress Board (OPB). 2000. Oregon State of the Environment Report 2000: Statewide Summary.

Oregon State of the Environment Report 2000 - Statewide Summary. 2000. SOER Science Panel, Paul Risser, Chair. Salem, Oregon: Oregon Progress Board.

Pater, D.E., S.A. Bryce, and T.D. Thorson. 1998. Ecoregions of Western Washington and Oregon. U.S. Geological Survey, Reston, VA.

Risser, P. G. 2000. Oregon State of the Environment Report 2000, Statewide Summary. Report prepared for Oregon Progress Board, SOER Science Panel. 81 pp.

Rocchio, J. and R. Crawford. 2009. Field guide to Washington's ecological systems. Washington Natural Heritage Program. Washington Department of Natural Resources. Olympia, WA.

Taft, O.W. and S.M. Haig. 2003. Historical wetlands in Oregon's Willamette Valley: implications for restoration of winter waterbird habitat. *Wetlands* 23:51-46

Tiner, R.W. 2011. Dichotomous keys and mapping codes for wetland landscape position, landform, water flow path, and waterbody type descriptors: Version 2.0. U.S. Fish and Wildlife Service, National Wetlands Inventory Program, Northeast Region, Hadley, MA. 51 pp.

U.S. Fish and Wildlife Service. 2017. Willamette Valley Conservation Study. Pacific Region, Portland, Oregon. 148 pp.

Wohl, E., J. Castro, B. Cluer, D. Merritts, P. Powers, B. Staab, and C. Thorne. 2021. Rediscovering, reevaluating, and restoring Lost River-wetland corridors. *Frontiers of Earth Science* 9:653623.

Zedler, P.H. 2003. Vernal pools and the concept of "isolated wetlands." *Wetlands* 23(3):597-607

VII. LIST OF RESOURCES FOR FURTHER INFORMATION

Oregon Strategic Conservation

Clarke, W.E., D. White, and A.L. Schaedel. 1991. Oregon ecological regions and subregions for water quality management. *Environmental Management* 15:847–856.

Mucken, A. and B. Bateman, (Eds.). 2017. Oregon's 2017 integrated water resources strategy. Oregon Water Resources Department. Salem, OR

U.S. Fish and Wildlife Service. 2010. Recovery plan for the prairie species of Western Oregon and Southwestern Washington. U.S. Fish and Wildlife Service, Portland, Oregon. xi + 241 pp.

Wetland Types

Aldous, A.R., M.W. Gannett, M. Keith, and J. O'Connor. 2015. Geologic and geomorphic controls on the occurrence of fens in the Oregon Cascades and implications for vulnerability and conservation. *Wetlands* 35(4): 757-767.

Brophy, L.S. 2007. Estuary assessment: Component XII of the Oregon Watershed Assessment Manual. Prepared for the Oregon Department of Land Conservation and Development, Salem, OR and the Oregon Watershed Enhancement Board, Salem, OR.

Seliskar, D.M. and J.L. Gallagher. 1983. The ecology of tidal marshes of the Pacific Northwest Coast: a community profile. U.S. Fish and Wildlife Service, Division of Wildlife Services, Washington, D.C. FWS/OBS 82/32. 65 pp.

Wetland Habitat Loss

Brophy L.S., C.M. Greene, V.C. Hare, B. Holycross, A. Lanier, W.N. Heady, et al. 2019. Insights into estuary habitat loss in the western United States using a new method for mapping maximum extent of tidal wetlands. *PLoS ONE* 14(8): e0218558.

Dahl, T.E. 2006. Status and trends of wetlands in the conterminous United States 1998 to 2004. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. 112 pp.

Marcoe, K. and S. Pilson. 2017. Habitat change in the lower Columbia River estuary, 1870-2009. *Journal of Coastal Conservation* 21: 505-525.

Mayer, T. D. 2005. Water-quality impacts of wetland management in the Lower Klamath National Wildlife Refuge, Oregon and California, USA. *Wetlands* 25(3): 697-712.

Morlan, J. C. and D. Peters. 1999. Oregon study confirms continued freshwater wetland losses. *National Wetlands Newsletter* 21(3):doi/10.1002/eco.1953

Moss, J.M., T. Divoll, J.C. Morlan, and T. O'Neill. 2022. Wetland and land use change in the Willamette Valley, Oregon: 2005 to 2020. SWCA Environmental Consultants, Portland, Oregon, and Oregon Department of State Lands, Salem, Oregon.

Wetland Functions and Values

Boyd, C. S., and T.J. Svejcar. 2012. Biomass production and net ecosystem exchange following defoliation in a wet sedge community. *Rangeland Ecology & Management* 65(4): 394-400. doi:10.2111/REM-D-11-00159.1

Donnelly, J. P., D. E. Naugle, D. P. Collins, B. D. Dugger, B. W. Allred, J. D. Tack, and V. J. Dreitz. 2019. Synchronizing conservation to seasonal wetland hydrology and waterbird migration in semi-arid landscapes. *Ecosphere* 10(6):e02758. 10.1002/ecs2.2758

Donnelly, J.P., J.N. Moore, M.L. Casazza, and S.P. Coons. 2022. Functional wetland loss drives emerging risks to waterbird migration networks. *Frontiers in Ecology and Evolution* 10:844278. doi: 10.3389/fevo.2022.844278

Nash, C.S., J.S. Selker, G.E. Grant, S.L. Lewis, and P. Noel. 2018. A physical framework for evaluating net effects of wet meadow restoration on late-summer streamflow. *Ecohydrology*: <https://doi.org/10.1002/eco.1953>

Mitsch, W.J. and J.G. Gosselink. 1993. *Wetlands* (second edition): New York, Van Nostrand Reinhold Company, Inc., 722 pp.

Wetland and Riparian Restoration Approaches

Cornu, C.E. 2005a. Restoring Cox, Dalton, & Fredrickson Creek Marshes. South Slough NERR. Coastal Resource Management Series. CRMS-2005-2. Coos Bay, Oregon.

Cornu, C.E. 2005b. Restoring Anderson Creek Marsh. South Slough NERR. Coastal Resource Management Series. CRMS-2005-3. Coos Bay, Oregon.

Cornu, C.E. 2005c. Restoring Kunz Marsh. South Slough NERR. Coastal Resource Management Series. CRMS-2005-1. Coos Bay, Oregon.

Flitcroft, R.L., W.R. Brignon, B. Staab, J.R. Bellmore, J. Burnett, P. Burns, B. Cluer, G. Giannico, J.M. Helstab, J. Jennings, C. Mayes, C. Mazzacano, L. Mork, K. Meyer, J. Munyon, B.E. Penaluna, P. Powers, D.N. Scott, and S.M. Wondzell. 2022. Rehabilitating valley floors to a stage 0 condition: a synthesis of opening outcomes. *Frontiers of Environmental Science* 10:892268.

Frenkel, R.E. and J.C. Morlan. 1990. Restoration of the Salmon River salt marshes--retrospect and perspective: U.S. Environmental Protection Agency, Region 10, 142 pp.

Frenkel, R.E. and J.C. Morln. 1991. Can we restore our salt marshes? Lessons from the Salmon River, Oregon: *Northwest Environmental Journal* 7: 119-135.

Good, J.W. and C.B. Sawyer. 1998. Recommendations for a nonregulatory wetland restoration program for Oregon. Oregon Sea Grant. Corvallis, OR.

Good, J. W., J. W. Weber, J. W. Charland, J. V. Olson, and K.A. Chapin. 1998. National coastal zone management effectiveness study: protecting estuaries and coastal wetlands. Final Report to the Office of Ocean and Coastal Resources Management, National Oceanic and Atmospheric Administration. Oregon Sea Grant Special Report PI-98-001. Corvallis, OR.

Krueger, J. J., S. T. Bois, T. N. Kaye, D. M. Steeck, and T. H. Taylor. 2014. Practical guidelines for wetland prairie restoration in the Willamette Valley, Oregon - Field- tested methods and techniques. 124 pp.

Kusler, J.A., and Kentula, M.E., eds. 1990. *Wetland creation and restoration--The status of the science*: Washington, D.C., Island Press, 591 pp.

Chapter 1: Josselyn, M., J. Zedler, and T. Griswold. *Wetland Mitigation Along the Pacific Coast of the United States*; Pages 1-36.

Chapter 15: Jensen, S.E. and W.S. Platts. Restoration of Degraded Riverine/Riparian Habitat in the Great Basin and Snake River Regions; Pages 377 to 416.

Chapter 16: Riparian Wetland Creation and Restoration in the Far West: A Compilation of Information. Pages 417 to 464.

Pilliod, D.S., A.T. Rohde, S. Charnley, R.R. Davee, J.B. Dunham, H. Gosnell, G.E. Grant, M.B. Hausner, J.L. Huntington, and C. Nash. 2018. Survey of beaver-related restoration practices in rangeland streams of the Western USA. *Environmental Management* 61: 58-68. DOI 10.1007/s00267-017-0957-6

Pollock, M.M., G.M. Lewallen, K. Woodruff, C.E. Jordan, and J.M. Castro (Editors) 2023. The Beaver restoration guidebook: working with beaver to restore streams, wetlands, and floodplains. Version 2.02. United States Fish and Wildlife Service, Portland, Oregon. 189 pp. <https://www.fws.gov/media/beaver-restoration-guidebook>

Schrager, H., J. Bragg, A. DeMarzo, B. Yednock, J. Schmitt, A. Helms, J. Tally, D. Rudd, E. Dean, R. Muse, G. Cooper, C. Cornu, T. Gaskill, and P. Wilson. 2022. South Slough National Estuarine Research Reserve Management Plan, 2017-2022. Coos County Oregon, 197 pp.

Simenstad, C.A., B.E. Feist, J. Morlan, and P.B. Williams. 1999. Assessment of potential dike-breach restoration of estuarine wetlands in Tillamook Bay, Oregon. Prepared for Tillamook Bay National Estuary Project. University of Washington, School of Fisheries. 57 pp.

U.S. Department of Agriculture, Natural Resources Conservation Service. 2021. Title 210-National Engineering Handbook, Part 650 Engineering Field handbook, National Engineering Handbook, Chapter 13--Wetland restoration, Enhancement, or Creation: Washington, D.C., U.S. Department of Agriculture, Natural Resources Conservation Service, 160 pp.

Riparian Areas:

Chopin, D.M., Beschta, R.L., and H.W. Shen. 2002. Relationships between flood frequencies and riparian plant communities in the upper Klamath Basin, Oregon. *Journal of the American Water Resources Association* 38: 603-617.

Lewis, L., L. Clark, R. Krapf, M. Manning, J. Staats, T. Subirge, L. Townsend, and B. Ypsilantis. 2003. Riparian area management: Riparian-wetland soils. Technical Reference 1737-19. Bureau of Land Management, Denver, CO. BLM/ST/ST-03/001+1737. 109 pp.

Swanson, S. S. Wyman, and C. Evans. 2015. Practical grazing management to maintain or restore riparian functions and values on rangelands. *Journal of Rangeland Applications* 2: 1-28. Access online: <https://thejra.nkn.uidaho.edu/index.php/jra/article/view/20/40>

United States Department of Agriculture, Natural Resources Conservation Service. 2009. Stream Visual Assessment Protocol Version 2. National Biology Handbook, Subpart B-Conservation

Planning, Part 614. USDA, NRCS, in cooperation with the West National Technology Support Center. 75 pp.

Wyman, S., D. Bailey, M. Borman, S. Cote, J. Eisner, W. Elmore, B. Leinard, S. Leonard, F. Reed, S. Swanson, L. Van Riper, T. Westfall, R. Wiley, and A. Winward. 2006. Riparian area management: Grazing management processes and strategies for riparian-wetland areas. Technical Reference 1737-20. BLM/ST/ST-06/002+1737. U.S. Department of the Interior, Bureau of Land Management, National Science and Technology Center, Denver, CO. 105 pp.

Wetland and Riparian Plants

Christy, J.A. 2004. Native freshwater plant associations of Northwestern Oregon. Oregon Natural Heritage Information Center. Oregon State University. 246 pp.

Crowe, E.A., B.L. Kovalchik, and M.J. Kerr. 2004. Riparian and wetland vegetation of Central and Eastern Oregon. Oregon State University, Portland, OR. 473 pp.

Darris, Dale C. 2002. Native shrubs as a supplement to the use of willows as live stakes and fascines in Western Oregon and Western Washington. USDA, NRCS Plant Materials Technical Note No. 31. Portland, Oregon, 10 pp.

Hoag, J.C. 2003. Willow clump plantings. NRCS Plant Materials Technical Note No. 36. NRCS Plant Materials Center, Aberdeen, Idaho. 8 pp.

Shaw, N.L., S.M. Lambert, and C. Hoag. 2001. Plant materials for western riparian areas. USDA, NRCS Tech Note 4., Portland, Oregon, 12 pp.

Withrow-Robinson, B., M. Bennett, and G. Ahrens. 2011. A guide to riparian tree and shrub planting in the Willamette Valley: steps to success. EM 9040. Oregon State University Extension, Corvallis, Oregon. 28 pp.

Wetland Soils:

United States Department of Agriculture, Natural Resources Conservation Service. 2018. Field indicators of hydric soils in the United States, Version 8.2. L.M. Vasilas, G.W. Hurt, and J.F. Berkowitz (eds.). USDA, NRCS, in cooperation with the National Technical Committee for Hydric Soils.

Vepraskas, M. J. 1994. Redoximorphic features for identifying aquic conditions. Tech. Bulletin 301. North Carolina Ag. Research Service, North Carolina State Univ., Raleigh, North Carolina.

Website for where to find updates to hydric soils:

<https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/soil/hydric-soils#notes>

Wetland-Dependent Wildlife and Wildlife Habitat

Altman, B. and J. L. Stephens. 2012. Land managers guide to bird habitat and populations in oak ecosystems of the Pacific Northwest. American Bird Conservancy and Klamath Bird Observatory. 82 pp.

Pilliod, D.S. and E. Wind. (editors). 2008. Habitat management guidelines for amphibians and reptiles of the Northwestern United States and Western Canada. Partners in Amphibian and Reptile Conservation, Technical Publication HMG-4, Birmingham, AL. 139 pp.

Pacific Coast Joint Venture. 2013. Habitat and population objectives-wetland birds and waterbirds, North Puget Lowlands Ecoregion. 71 pp. (see [Waterbird Habitat Pacific Coast Joint Venture](#))

Davis, K.P., D.J. Augustine, A.P. Monroe, J.D. Derner, and C.L. Aldridge. 2019. Adaptive rangeland management benefits grassland birds utilizing opposing vegetation structure in the shortgrass steppe. Ecological Applications 00(00):e02020. 10.1002/eap.2020

Mesic Habitat Planning Guide: (Working Lands for Wildlife Resource)

https://efotg.sc.egov.usda.gov/references/public/CO/Mesic_Habitat_Conservation_Planning_Guide.pdf

ODFW. 2015. Guidance for conserving Oregon's native turtles including best management practices. Oregon Dept. of Fish and Wildlife. 99 pp.

Wetland Restoration Prioritization Tools:

1) Intermountain West Joint Venture, WET Tool: <https://iwjv.org/solution-based-science/wet/>

User Guide: <https://iwjv.org/wp-content/uploads/2022/11/User-Guide-2.pdf>

Peer Reviewed Publication:

<https://www.frontiersin.org/articles/10.3389/fevo.2022.844278/full>

Data Sources, Historic Wetland Types and Vegetation:

US Department of Interior, Bureau of Land Management (BLM), Survey data, Land Status & Cadastral Survey Records: <https://www.blm.gov/or/landrecords/survey/ySrvy1.php>


US Department of Interior, BLM General Land Office Records:

<https://gloreCORDS.blm.gov/default.aspx>


Historical vegetation maps: <https://inr.oregonstate.edu/hvmp/available-maps#Oregon>

LiDAR data viewer: <https://gis.dogami.oregon.gov/maps/lidarviewer/>


APPENDIX A: OCS Blue Mountains Ecoregion




APPENDIX A: OCS Coast Range Ecoregion




APPENDIX A: OCS East Cascades Ecoregion



APPENDIX A: OCS Northern Basin and Range Ecoregion



APPENDIX A: OCS Willamette Valley Ecoregion



APPENDIX B: Ranking Worksheet

Oregon NRCS Wetland Reserve Easements Ranking Worksheet			
Project Name		Total Ranking Score	
Landowner		Date of Field Evaluation	
Ranking Team		FWS	
NRCS		ODFW	
Is there an existing easement on the application property? (y/n) If yes, type:		SWCD	
		Other(s)	

Screening Question
<p>Is the application:</p> <p>Within one of the priority areas identified on the ACEP-WRE Priority Area Map? Within an approved NRCS Wetland Restoration Conservation Implementation Strategy (CIS) area or Long Range Plan (LRP)?</p> <p>A unique property that needs protection (requires State Conservationist approval)?</p> <p>If any of the above are checked, then rank application. Otherwise, application is low priority and defer ranking.</p>

Easement Area			
Attach map showing acres of each category			
Eligible Acres	Acres	Other Eligible Acres	Acres
PC		Adjacent lands	
FW		Riparian	
FWP		CRP	
Former or degraded wetlands		Restored or protected wetlands	
Substantially altered by flooding		Hydric soil minor components	
Subtotal		Subtotal	

Other Eligible Acres
<p>How will "Other Eligible Acres" contribute to the functions of the restored wetland? (Check all that apply)</p> <p><input type="checkbox"/> Provides species-specific upland habitat for targeted species</p> <p><input type="checkbox"/> Increases habitat value by providing additional nesting habitat, etc.</p> <p><input type="checkbox"/> Other, explain:</p> <p><input type="checkbox"/> Provides buffering and filtering from surrounding land uses</p> <p><input type="checkbox"/> Reduces fragmentation of or increases the area of the wetland complex</p>

APPENDIX B: Ranking Worksheet

Summary of Ranking Score (125 point funding cutoff)		
Ranking Factor	Maximum Points	Points Received
Environmental Benefits – Subtotal	165	
Hydrology ≥50% of Environmental Benefits		
1. Altered Hydrology (Present Condition)	40	
2. Restored Hydrology (Future Condition)	45	
3. Habitat for At-Risk Species	20	
4. Native Plant Communities	30	
5. Habitat Diversity	5	
6. Adjacent Protected Habitat	10	
7. Floodplain Connectivity	5	
8. Duration of Enrollment	10	
Economic Considerations – Subtotal	65	
9. Easement Cost	10	
10. Restoration Cost	10	
11. Restoration Cost-Benefit	25	
12. Operation and Maintenance	20	
Special Considerations – Subtotal	20	
13. Priority Areas	10	
14. Project Size	10	
Total	250	

Environmental Benefits	
<p>1. Altered Hydrology (Present Condition) <i>Degree of hydrologic alteration.</i> Use Certified Wetland Determination or wetland inventory (with input from Resource Soil Scientist and Wetlands Biologist) to estimate the degree of departure from original hydrology. Choose the category representing the majority of the Eligible Acres that will have hydrology restored.</p>	40 points maximum
Original wetland hydrology is significantly degraded or modified. For example, functional ditches, dikes, diversions, and tiles are affecting the historic hydrology regime over the majority of the Eligible Acres.	5 points
Original wetland hydrology is moderately degraded or modified; or original wetland hydrology was previously restored. For example, function (or partially functional) ditches, dikes, diversions, and tiles are affecting ≤ 50% of the Eligible Acres.	20 points
Original wetland hydrology is relatively unmodified or previous hydrologic modifications have largely deteriorated such that historic hydrology is present over the majority of the Eligible Acres.	40 points
<p>2. Restored Hydrology (Future Condition) <i>Extent of hydrologic restoration relative to historic conditions.</i> Percent of the Eligible Acres on which the hydrology will be restored to historic conditions suitable for the needs of the native wetland-dependent wildlife</p>	45 points maximum

APPENDIX B: Ranking Worksheet

species that occurred in the area and appropriate for the wetland functions and values that existed prior to manipulation.	
90 – 100%	45 points
75 – 89%	25 points
50 – 74%	15 points
less than 50%	0 points

3. Habitat for At-Risk Species	20 points maximum
Offered acres have known use by State or Federally listed Threatened, Endangered, or Candidate species. <i>List species:</i>	20 points
Offered acres will restore, enhance, or create habitat for use by State or Federally listed Threatened, Endangered, or Candidate species. <i>List species:</i>	10 points
None of the above	0 points

4. Native Plant Communities	30 points maximum
Percent of the Total Easement Acres that will return to a predominance of historic native vegetation after restoration. Acreage includes existing vegetation in “Other Eligible Acres”, as well as those areas planted, seeded, or allowed to naturally re-vegetate. Consider the likelihood that easement sites will retain their habitat functions and values after the restoration is complete.	
90 – 100%	30 points
75 – 89%	20 points
50 – 74%	10 points
less than 50%	0 points

5. Habitat Diversity	5 points maximum
Post-restoration condition within the easement. Utilize the “Classification of Wetlands and Deepwater Habitats of the United States” (AKA Cowardin classes). Check all that apply: <input type="checkbox"/> PEM <input type="checkbox"/> PSS <input type="checkbox"/> PFO <input type="checkbox"/> Riparian <input type="checkbox"/> Upland <input type="checkbox"/> Estuarine <input type="checkbox"/> Open Water	
3 or more types	5 points
2 types	3 points
1 type	2 points

6. Adjacent Protected Habitat	10 points maximum
Proximity of proposed easement to an existing protected area (i.e. other conservation easement, USFWS refuge, State or locally managed wildlife areas, etc.) <i>List the protected area(s):</i>	

APPENDIX B: Ranking Worksheet

Adjacent	10 points
Less than 1 mile	5 points
1 – 5 miles	2 points
More than 5 miles	0 points

7. Floodplain Connectivity Post restoration conditions support a functioning floodplain with river/creek having access to the floodplain.	5 points maximum
Yes	5 points
No	0 points

8. Duration of Enrollment Permanence of restored habitat	10 points maximum
Permanent Easement	10 points
30-year Easement or 30-year Contract with high likelihood that the site will retain its habitat functions and values after the enrollment period ends. Provide justification of high likelihood:	5 points

Subtotal for Environmental Benefits: _____

Economic Considerations	
9. Easement Cost Voluntary landowner offer to accept a reduced easement value based on the compensation that the landowner would be entitled to for the enrollment type.	10 points maximum
70 % Fair Market Value (GARC is 85% so reduce GARC by 15 %)	10 points
75 % Fair Market Value (reduce GARC by 10%)	7 points
80% Fair Market Value (reduce GARC by 5%)	5 points
85% Fair Market Value (GARC)	0 points

10. Restoration Cost The total estimated restoration cost per acre that will be borne by NRCS per the preliminary restoration plan. Total Restoration Cost Estimate \$ _____	10 points maximum
Less than \$500 per acre	10 points
\$500 - \$1500 per acre	5 points
\$1500 - \$2500	3 points
More than \$2500 per acre	0 points
Excessive restoration or long-term costs without a State Conservationist waiver	Ineligible

11. Restoration Cost-Benefit Cost per environmental benefit ratio Restoration cost per acre / Environmental Benefits points = Cost-benefit ratio	25 points maximum
---	--------------------------

APPENDIX B: Ranking Worksheet

Less than 10	25 points
10-20	10 points
More than 20	0 points

<p>12. Operation and Maintenance The cost of O&M and management requirements needed to keep structural and vegetative practices functional (consider deterioration and damage prevention, repair, and replacement.)</p>	20 points maximum
<p>Minimal. Restoration designed to minimize O&M and management costs and requirements; practices have low replacement cost, easy access, and/or infrequent maintenance requirements. For example: tile breaks; ditch plugs or fill; low berms w/o structures; plantings with infrequent vegetation management (e.g. spraying, burning, disking, mowing, reseeding) every 5 – 10 years; water control structures without seasonal management.</p>	20 points
<p>Moderate. Restoration requires a moderate degree of O&M and management costs during establishment period, with less frequent inputs thereafter. For example: existing or proposed levees require long-term maintenance, water control structures with seasonal management; structures needing minor maintenance after flood events; annual spot treatment of invasive weeds; vegetation management (spraying, burning, disking, mowing, reseeding) every 2-5 years.</p>	5 points
<p>High. Onsite or offsite conditions require high degree of O&M and management and maintenance costs (e.g. structures requiring significant maintenance after flood events, restoration requiring frequent water management, recurring treatment needed to address erosion and/or siltation, continual noxious weed re-infestation). Site is difficult or costly to access. Extensive vegetation management (e.g. spraying, burning, disking, mowing, reseeding on less than a 2 year cycle.</p>	Ineligible without a State Conservationist Approved Waiver

Subtotal for Economic Considerations: _____

Special Considerations	
<p>13. Priority Areas Refer to Priority Geographic Regions Maps</p>	10 points maximum
Project is located within a USFWS Recovery Zone for Bradshaw’s lomatium or Nelson’s checkermallow.	10 points
Project is located within a Lower Columbia River Priority Area.	10 points
Project located within a “Highest Priority” watershed for Oregon Coastal Coho Salmon.	10 points
Project is located within a Conservation Opportunity Area (COA) where wetlands (also includes wet prairie, riparian, aquatic, and estuary) are listed as a “key habitat”, as identified in ODFW’s <i>Oregon Wetlands Conservation Strategy</i> ;	5 points


APPENDIX B: Ranking Worksheet

or project is located within a target area of an approved wetlands restoration CIS or LRP.	
--	--


14. Project Complexity Level Base on prelim WRPO, consider inventory, plan, and design time, level of permitting, and NEPA, ESA, SHPO consultation requirements.	10 points maximum
Very low	10 points
Low	7 points
Moderate	5 points
High	0 points

Subtotal for Special Considerations: _____

APPENDIX C: 2023 ACEP - WRE Priority Regions, Oregon Coastal Coho Priority Watersheds



APPENDIX C: 2023 ACEP - WRE Priority Regions Lower Columbia River Priority Area



Appendix D: Wetland Restoration

Conservation Practice 657 – Specification Sheet



Emergent wetland in Klamath County, Oregon.

Definition

The return of a wetland and its functions to a close approximation of its original condition as it existed prior to disturbance on a former or degraded wetland site.

Purpose

To restore wetland function, value, habitat, diversity, and capacity to a close approximation of the pre-disturbance conditions by restoring:

- Conditions conducive to hydric soil maintenance.
- Wetland hydrology (dominant water source, hydroperiod, and hydrodynamics).
- Native hydrophytic vegetation (including the removal of undesired species, and/or seeding or planting of desired species).
- Original fish and wildlife habitats.

Where used

This practice applies only to natural wetland sites with hydric soils which have been subject to degradation of hydrology, vegetation, or soils.

Planning for Wetland Restoration

Wetland restoration is normally applied as part of a resource management system where a landowner's objectives are to return a wetland or wetland systems to a close approximation of the original pre-disturbance conditions. In general, the planner must 1) describe the likely original wetland conditions and contributing natural processes prior to anthropogenic disturbances, 2) characterize the extent of disturbance to the original system, and 3) determine the degree to which original conditions can be restored.

A critical step in wetland restoration planning is to identify reference wetlands of the same type as the wetland to be restored. Those reference wetlands with the least amount of anthropogenic disturbance are particularly useful as they typically exhibit high levels of function and can serve as a template for restoration. This critical step ensures the most appropriate wetland characteristics and functions are restored. In addition to reference wetlands, there are several resources that can help the planner determine the likely historical conditions of a given wetland including. These include, but are not limited to: GLO survey

Appendix D: Wetland Restoration

Conservation Practice 657 – Specification Sheet

notes and plat maps, soils descriptions, ecological site descriptions, old aerial photography, natural community descriptions, nearby remnant natural communities, and landowner accounts.

Most natural communities, including wetlands, exhibit a range in characteristics that are dependent upon several factors such as disturbance history and dynamic weather patterns. As such, there are usually a handful of intergrading plant communities that may be supported by a given soil type in a specific landscape position within a given ecoregion. These include both early and late successional phases. For example, in the absence of disturbance such as severe flooding or wildfire, a wet prairie community type may shift towards a shrub dominated community and ultimately a forested community. The phase that is targeted for restoration depends on site-specific capabilities and limitations, current landscape context, landowner objectives, and regional priorities. If target objectives of the project are not aligned with one of the natural phases of the historical plant community, consider using Wetland Enhancement (659) practice standard.

The amount and duration (hydroperiod) of hydrology can vary greatly among wetland types. The hydrology of a wetland may be driven by floodwater from rivers and lakes, direct precipitation, runoff from contributing upland areas, groundwater discharge, or ocean tides. Some wetlands are inherently wetter than others. Permanently inundated depressional wetlands may retain surface water in most years while vernal pools may only have surface water for a period of weeks during the wet season then dry up for the remainder of the year. Other wetland types, such as groundwater driven fens, may rarely have surface water, but can remain saturated for extended periods during the year. It is important to identify the various hydrologic influences and their impact on the wetland hydroperiod.

When analyzing hydrology, existing water rights must also be taken into consideration. The availability of sufficient water rights and the conveyance of water rights to adjacent properties should be carefully reviewed prior to restoration.

Each wetland type has a unique set of functions and values. Wetland functions include biogeochemical, hydrologic, and habitat functions. Values are those functions deemed important by society. Functions of a particular wetland may be quantified and compared to highly functional wetlands of the same type by conducting a functional assessment. The scores of a functional assessment help determine which steps need to be taken to restore a wetland to its full functional potential. Functional assessments include general assessments, such as the Oregon Rapid Wetland Assessment Protocol (ORWAP) as well as more comprehensive Hydrogeomorphic (HGM) models, which are unique to specific wetland classes. Completion of a functional assessment is a national requirement of the wetland restoration practice standard. If no suitable Functional Assessment is available, an appropriate Wildlife Habitat Evaluation Guides (WHEGs) may be used to document existing and planned conditions.

In addition to wildlife habitat, wetlands provide many ecological functions and societal values, such as water storage and flood attenuation, nutrient cycling, and sediment retention. Wetlands are important habitat to many wildlife species, and different wetland types often support a unique suite of wildlife species. The timing, depth, duration, frequency, and flow of water through a wetland have significant influences on habitats and the wildlife species they support.

When determining the restoration measures needed to restore wetland function, effects on adjacent lands must be taken into consideration. This is especially critical with hydrology. Oftentimes, existing drainage or water delivery infrastructure affects adjacent properties. Artificial drainage or conveyance infrastructure should be decommissioned to the extent practicable without adversely impacting adjacent lands. The degree to which remaining infrastructure impacts the wetland restoration must be critically evaluated. In cases where remaining infrastructure will significantly reduce the function of the restored wetland, restoration may not be practicable.

Identification of additional practices that support the wetland restoration is also important. Restored wetlands are most effective when used in combination with conservation practices that contribute to the restoration of adjacent upland areas.

Appendix D: Wetland Restoration

Conservation Practice 657 – Specification Sheet

Plans and Specifications

Site-specific requirements are listed on the specifications sheet, and associated engineering design. These specifications are used in conjunction with the Conservation Plan and Plan Map to document existing conditions and restoration actions needed to restore wetlands to the extent practicable. Refer to Conservation Practice Standard 657 – *Wetland Restoration* for additional guidance. If associated conservation practices are included in the plan, refer to individual practice specifications, as appropriate.

Operation and Maintenance

An Operation and Maintenance (O&M) Plan will be developed that describes specific actions required to ensure normal function of installed structural components over the practice lifespan. Maintenance, management and monitoring requirements needed to ensure long-term wetland function should be included in the Operation & Maintenance Plan