

**GAMOCHAETA ARGYRINEA (ASTERACEAE)  
NATURALIZED IN CALIFORNIA**

**JOHN L. STROTHER**  
University Herbarium  
University of California  
Berkeley, California 94720-2465  
strother@berkeley.edu

**RICHARD L. MOE**  
University Herbarium  
University of California  
Berkeley, California 94720-2465  
rlmoe@berkeley.edu

**GUY L. NESOM**  
2925 Hartwood Drive  
Fort Worth, Texas 76109  
guynesom@sbcglobal.net

**ABSTRACT**

*Gamochoeta argyrinea* is naturalized in the Lake Oroville area of Butte Co., California, first documented there in 1990 and again in 2000, its persistent presence confirmed in 2012. It probably arrived there from a locality in the southeastern USA.

Curation of UC and JEPS Asteraceae collections has brought attention to a previously unreported non-native species naturalized in the California flora.

**GAMOCHAETA ARGYRINEA** G.L. Nesom, Sida. 21: 718, figs. 1-4. 2004.

**California.** Butte Co.: ca. 1/8 mi NW of the Enterprise Bridge across Lake Oroville, E of the Enterprise Boat Launch Area, growing on fine damp bare disturbed granite soil, below the high water line of Lake Oroville, Foothill Woodland (destroyed), 850 ft, normal size plants, uncommon, scattered, 25 May 1990, *Lowell Ahart 6481* (UC, Figure 1); ca. 1/2 mi SW of the Enterprise Bridge across Lake Oroville, below the high water line of Lake Oroville, Foothill Woodland (destroyed), 850 ft, growing on dry bare white disturbed granite soil, large size plants, few plants seen, 16 Jun 1990, *Lowell Ahart 6523* (UC); ca. 5 mi (air) SW of Oroville, W of Larkin Road, ca. 1 1/4 mi S of the Oroville Airport, on the margin of the Thermalito Afterbay, on the W side of the access road to the boat ramp at the Thermalito Afterbay, damp red disturbed gravelly soil, freshwater marsh, 140 ft, common, normal size plants, 13 May 2000, *Lowell Ahart 8345* (JEPS; Figure 2).

Species collected by Ahart on the same day at each of the three Lake Oroville localities are characteristic of disturbed habitats and nearly half of them are not native to California. In 1990, with *Ahart 6481*: *Agrostis avenacea*, *Chorizanthe membranacea*, *Filago californica*, *Juncus acuminatus*, *Juncus bufonius*, *Juncus tenuis*, *Lindernia dubia*, *Leontodon saxatilis*, *Polygonum hydropiper*, *Polypogon interruptus*, *Rumex acetosella*, *Silene antirrhina*, *Torilis arvensis*, and *Vicia americana*. In 1990, with *Ahart 6523*: *Agoseris heterophylla*, *Alopecurus carolinianus*, *Apocynum cannabinum*, *Centaureum muehlenbergii*, *Centaureum tenuiflorum*, *Digitaria sanguinalis*, *Eragrostis cilianensis*, *Eriodictyon californicum*, *Pseudonaphalium stramineum*, *Herniaria hirsuta*, *Kickxia elatine*, *Lonicera interrupta*, *Polygonum hydropiper*, *Rorippa palustris*, *Scleranthus annuus*, *Uropappus lindleyi*, and *Veronica peregrina* subsp. *xalapensis*. In 2000, with *Ahart 8345*: *Castilleja attenuata*,




Figure 1. *Gamochaeta argyrinea* from Butte County (UC), northwest of the Enterprise Bridge, 25 May 1990.



Figure 2. *Gamochaeta argyrinea* from Butte County (JEPS), margin of the Thermalito Afterbay, 13 May 2000.

*Centunculus minimus*, *Crucianella angustifolia*, *Eleocharis macrostachya*, *Galium tricornutum*, *Gastridium phleoides*, *Gastridium ventricosum*, *Juncus acuminatus*, *Lathyrus angulatus*, *Plagiobothrys stipitatus*, *Plantago coronopus*, *Plantago virginica*, *Pseudognaphalium stramineum*, *Ranunculus pusillus*, *Rorippa curvisiliqua*, *Trifolium cernuum*, *Trifolium dubium*, and *Triodanis biflora*. Records for these collections are from the Consortium of California Herbaria (2012).



Figure 3. *Gamochoaeta argyrinea* at Thermalito Afterbay site, 17 May 2012.

On 17 May 2012, Strother and Moe found the *Ahart 6481* site to be under water and collected *Gamochoaeta argyrinea* at the *Ahart 8345* site (Figure 3; Thermalito Afterbay, *Strother 1373*, UC, duplicates to MO and TEX). They scanned and searched roadsides along minor roads but saw no other gamochaetas.

The Enterprise Bridge localities are about 20 miles from the Thermalito Afterbay — boat ramps are at both places. Perhaps cypselae in mud on tires and/or trailers account for the transport from one boat ramp area to another as well as the original dispersal to California. The popularity of Lake Oroville for bass fishing makes this a plausible scenario.

*Gamochoaeta argyrinea* has previously been known only from the southeastern USA (the type from North Carolina; Nesom 2004, 2006, 2007), where it reaches eastern Texas and Oklahoma at the westernmost extension of its range. It is essentially restricted to disturbed habitats and appears to have become widespread in the USA only recently.

The species most similar to *Gamochaeta argyrinea* appears to be *G. ustulata* (Nutt.) Holub, which is native to the Pacific Coast of the USA and southern British Columbia. Most of the other close relatives of *G. argyrinea* are from South America and it is perhaps native there, even though its occurrence has not been reported (Freire & Iharlegui 1997). In any case, it is likely that the Butte County plants are adventive from a locality in the southeastern USA.

#### ACKNOWLEDGEMENTS

We and the botanical community are indebted to Lowell Ahart and other collectors who have contributed (and continue to contribute) over the years and decades to documentation of plant distributions and introductions in California and beyond.

#### LITERATURE CITED

- Consortium of California Herbaria. 2012. Consortium database. Data provided by the participants of the Consortium. <[ucjeps.berkeley.edu/consortium/](http://ucjeps.berkeley.edu/consortium/)>
- Freire, S.E. and L. Iharlegui. 1997. Sinopsis preliminar del género *Gamochaeta* (Asteraceae, Gnaphalieae). Bol. Soc. Argent. Bot. 33: 23–35.
- Nesom, G.L. 2004. New species of *Gamochaeta* (Asteraceae: Gnaphalieae) from the eastern United States and comments on similar species. Sida 21: 717–742.
- Nesom, G.L. 2006. *Gamochaeta* (Gnaphalieae). Pp. 431–438, In: Flora of North America Editorial Committee, eds. Flora of North America North of Mexico. Vol. 19. Oxford Univ. Press, New York and Oxford.
- Nesom, G.L. 2007. Distribution of *Gamochaeta* (Asteraceae: Gnaphalieae) in Texas, Oklahoma, Arkansas, and Louisiana. J. Bot. Res. Inst. Texas 1: 1125–1130.