

## ***SPOROBOLUS JUNCEUS* (POACEAE) IN OKLAHOMA**

**JEFFREY N. MINK**

Department of Biology  
Baylor University  
Waco, Texas 76798-7388  
J\_Mink@baylor.edu

**JASON R. SINGHURST**

Wildlife Diversity Program  
Texas Parks and Wildlife Department  
4200 Smith School Road  
Austin, Texas 78704

**WALTER C. HOLMES**

Department of Biology  
Baylor University  
Waco, Texas 76798-7388

### **ABSTRACT**

*Sporobolus junceus*, Piney Woods dropseed, a native of the contiguous southeastern states from Virginia to Texas, has been discovered in Oklahoma in sandhills bordering the Harrison/Doshier Bog in Pushmataha County.

**Key Words:** Poaceae, *Sporobolus*, Oklahoma, oak-pine savanna, sandhills

A cosmopolitan genus of at least 160 species, *Sporobolus* is widely distributed in warm-temperate, sub-tropical, and tropical habitats (Clayton & Renvoize 1986). Seventy three species are native to the Western Hemisphere (Peterson et al. 2003), with 27 species native to the United States (Peterson et al. 1997).

Fourteen species of *Sporobolus* (sensu lato) are documented as occurring in Oklahoma (McGregor & Barkley 1977, 1986; Oklahoma Biological Survey 2012; USDA/NRCS 2012). The discovery of a 15<sup>th</sup> species is reported here. Studies on the flora of sandy prairies and sandhill bogs in Pushmataha County have resulted in the collection of *Sporobolus junceus* (P. Beauv.) Kunth.

Voucher specimen: **OKLAHOMA**. Pushmataha Co.: Sandhill above Harrison/Doshier Bog, 5.16 miles W (at 266°) from Antlers; 34° 13' 34.33" N, 95° 42' 34.60" W; deep sandhills, 13 August 2006, J.R. Singhurst and E. Bridges 14408 (BAYLU). Figs. 1&2.

The sandhills surrounding Harrison and Doshier bogs contains sandy land graminoid and herbaceous species such as *Aristida desmantha*, *Triplasis purpurea*, *Phemeranthus rugospermus*, *Paronychia drummondii*, *Stillingia sylvatica* subsp. *sylvatica*, *Dalea phleoides* var. *microphylla*, *Evax candida*, *Eragrostis secundiflora*, *Croptilon divaricatum*, *Hypericum drummondii*, *Chamaecrista fasciculata*, *Cnidoscolus texanus*, *Froelichia floridana*, *Pediomelum* sp., *Scutellaria cardiophylla*, and *Matelea cynanchoides*, found in association with *Sporobolus junceus*.

*Sporobolus junceus* is distributed in the lower Atlantic and Gulf coastal plains from Virginia and Tennessee to Florida and west to eastern Texas and Arkansas (Peterson, Hatch, & Weakley 2003; USDA, NRCS 2012). It is native and occurs in openings of hardwood forests, sandy prairies, pine




Figure 1. *Sporobolus junceus*. Pushmataha Co., Oklahoma (Singhurst & Bridges 14408, BAYLU).



Figure 2. Close-up of *Sporobolus junceus* panicle (Singhurst & Bridges 14408, BAYLU).

barrens, and savannas within its distribution. The Oklahoma locality is isolated from several known distributions in other states. Within Texas, *S. junceus* is known from coastal prairie areas and interspersed in scattered locations throughout east Texas. The nearest site to the Pushmataha Co., Oklahoma, collection is Upsur Co., Texas (Turner et al 2003) approximately 155 km south. A Louisiana collection from Caddo Parish (Allen 1992) is about 190 km southeast of the Pushmataha Co. record. Locations in Bradley Co. and Drew Co., Arkansas, are about 330 km distant from the Oklahoma location.

Among the Oklahoma *Sporobolus* both *S. cryptandrus* and *S. pyramidatus* appear similar to *S. junceus*. However, *S. junceus* can be distinguished from the other species by a whorled panicle (3 or more branches per lower node), long spikelets (>2.5mm) and lack of tufted sheath apex (see Figs. 1&2). *Sporobolus cryptandrus* lacks a whorled panicle, has distinct white tufts of trichomes at the distal part of the sheath and shorter spikelets (<2.5 mm). *Sporobolus pyrimadatus* has a distinctly whorled (5 or more branches per lower node) pyramidal-shaped panicle, lacks a distinct tuft of trichomes at the sheath apex and shorter spikelets (<2.5 mm).

#### ACKNOWLEDGEMENTS

The authors appreciate the efforts of Darrell S. Vodopich (Baylor Department of Biology) for his photographic efforts and image processing of the included photos of *Sporobolus junceus*.

#### LITERATURE CITED

- Clayton, W.D. and S.A. Renvoize. 1986. *Genera Graminum: Grasses of the World*. Kew, London: Royal Botanical Gardens, Great Britain.
- McGregor, R.L. (coord.) and T.M. Barkley (ed.). 1977. *Atlas of the Flora of the Great Plains*. Iowa State Univ. Press, Ames.
- McGregor, R.L. (coord.) and T.M. Barkley (ed.). 1986. *Flora of the Great Plains*. Univ. Press of Kansas, Lawrence.
- Oklahoma Biological Survey. 2012. *Vascular Plants Database*, Univ. of Oklahoma, Norman. <<http://www.biosurvey.ou.edu/>> Accessed 19 Jan 2012.
- Peterson, P.M., R.D. Webster and J. Valdés-Reyna. 1997. *Genera of New World Eragrostideae (Poaceae: Chloridoideae)*. *Smithsonian Contr. to Bot.* 87:1–50.
- Peterson, P.M., S.L. Hatch and A.S. Weakley. 2003. *Sporobolus* R. Br. Pp. 115-139, *in* *Flora of North America* Editorial Committee, *Flora of North America*, Vol. 25. Oxford Univ. Press, New York.
- Turner, B.L., H. Nichols, G. Denny, and O. Doron. 2003. *Atlas of the Vascular Plants of Texas*, Vol.II. *Sida, Bot. Misc.* 24. Bot. Res. Inst. of Texas, Fort Worth.
- USDA, NRCS. 2012. *The PLANTS Database*. National Plant Data Team, Greensboro, North Carolina. <<http://plants.usda.gov>> Accessed 24 Jan 2012.