

Supplementary Material

Halophytes as a source of salt tolerance genes and mechanisms: a case study for the Salt Lake area, Turkey

Ceyda Ozfidan-Konakci^A, Baris Uzilday^B, Rengin Ozgur^B, Evren Yildiztugay^C, A. Hediye Sekmen^B and Ismail Turkan^{B,D}

^ADepartment of Molecular Biology and Genetics, Faculty of Science, Necmettin Erbakan University, 42 090, Meram, Konya, Turkey.

^BDepartment of Biology, Faculty of Science, Ege University, 35 100, Bornova, Izmir, Turkey.

^CDepartment of Biology, Faculty of Science, Selcuk University, 42 250, Selcuklu, Konya, Turkey.

^DCorresponding author. Email: ismail.turkan@ege.edu.tr

Fig. S1.



Eutrema parvulum (Schrenk) Al-Shehbaz & Warwick
(*Thellungiella parvula*)



Linum seljukorum P.H. Davis



Halocnemum strobilaceum (Pall.) M. Bieb.



Salicornia europaea L.



Camphorosma monspeliaca L.



Reaumuria alternifolia (Labill.) Britten



Artemisia santonicum L.



Halimione verrucifera (M. Bieb.) Aellen



Centaurea tuzgoluensis Aytaç & H.Duman



Sphaerophysa kotschyana Boiss.



Taraxacum mirabile Wagenitz



Linum ertugrulii O.Tugay, Y.Bağcı&Uysal



Salvia halophila Hedge



Silene salsuginea Hub.-Mor.



Onosma halophilum Boiss. & Heldr.



Ferula halophila Peşmen



Lepidium caespitosum Desv.



Hypericum salsugineum Robson&Hub.-Mor.



Verbascum pyroliforme (Boiss.&Heldr.)
O.Kuntze



Limonium iconicum (Boiss. &Heldr.)
O.Kuntze



Acantholimon halophilum Bokhari



Gypsophila oblancoolata Bark.



Gladiolus halophilus Boiss. & Heldr.



Allium cappadocicum Boiss.