



Revista de Gestão Costeira Integrada -
Journal of Integrated Coastal Zone
Management

E-ISSN: 1646-8872

rgci.editor@gmail.com

Associação Portuguesa dos Recursos
Hídricos

Diankha, Ousmane; Sow, Bamol Ali; Thiaw, Modou; Thierno Gaye, Amadou
Seasonal variability of sea surface temperature, chlorophyll-a and *Ethmalosa fimbriata*
abundance off the coast of Senegal
Revista de Gestão Costeira Integrada - Journal of Integrated Coastal Zone Management,
vol. 13, núm. 4, 2013, pp. 491-497
Associação Portuguesa dos Recursos Hídricos
Lisboa, Portugal

Available in: <http://www.redalyc.org/articulo.oa?id=388340143007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Seasonal variability of sea surface temperature, chlorophyll-a and *Ethmalosa fimbriata* abundance off the coast of Senegal *

*Variabilidade sazonal da temperatura da superfície do mar, clorofila-a e abundância de Ethmalosa fimbriata na Zona Costeira do Senegal ***

Ousmane Diankha^{@, 1}, Bamol Ali Sow², Modou Thiaw³, Amadou Thierno Gaye¹

ABSTRACT

The seasonal variability of *Ethmalosa fimbriata* abundance off Senegal Coastal Waters was investigated using an observation sea surface temperature (SST) and chlorophyll-a (Chl-a) concentration from PATHFINDER and SeaWiFS satellites' data of 1999-2009. The results showed that the variability of *E. fimbriata* landings in Senegalese waters is related to environmental changes. The seasonal fluctuations of *E. fimbriata* landings off Senegal Coastal Waters seemed to be governed by the combined effect of SST and Chl-a. However, SST better explains the seasonal variations of *E. fimbriata* landings than Chl-a, when they are used separately.

Keywords: Chlorophyll-a, *Ethmalosa fimbriata*, Sea surface temperature, Senegal.

RESUMO

A variabilidade sazonal da abundância de *Ethmalosa fimbriata* na Zona Costeira do Senegal foi investigada através da observação da temperatura da superfície do mar (TSM) e da concentração de clorofila-a (Chl-a) a partir de dados dos anos de 1999-2009, provenientes dos satélites PATHFINDER e SeaWiFS. Os resultados observados mostraram que a variabilidade da pesca de *E. fimbriata* em águas senegalesas aparentemente está relacionada com as mudanças ambientais. As flutuações sazonais na pesca de *E. fimbriata* na Zona Costeira do Senegal parece ser controlada pelo efeito combinado da TSM e Chl-a. Porém, a TSM explica melhor as variações sazonais na pesca de *E. fimbriata* que a Chl-a, quando estas variáveis são utilizadas separadamente.

Palavras-chave: Clorofila-a, *Ethmalosa fimbriata*, Temperatura da superfície do mar, Senegal.

@ - Corresponding author

1 - Laboratoire de Physique de l'Atmosphère et de l'Océan Siméon Fongang (LPAO-SF/ESP-UCAD, Sénégal), BP 5085 Dakar Fann. E-mail: Diankha.ousmane.diankha@ucad.edu.sn Gaye

2 - Laboratoire d'Océanographie, des Sciences de l'Environnement et du Climat (LOSEC) Département de Physique - UFR Sciences & Technologies - Université de Ziguinchor BP 523 Ziguinchor - Sénégal Email:

3 - Centre de Recherche Océanographique de Dakar-Thiaroye (CRODT, ISRA Sénégal). Email:

* Submission: 3 June 2013; Evaluation: 15 July 2013; Reception of revised manuscript: 23 July 2013; Accepted: 2 August 2013; Available on-line: 3 September 2013

** Portuguese Title and captions by Ulisses M Azeiteiro on behalf of the Journal Editorial Board

1. INTRODUCTION

Small pelagics represent more than 75 percent of the artisanal catches and 55 percent of total marine catches in Senegal (Diei-Ouadi, 2005). One of the most important species in the landings of small pelagic fish off Senegal is *Ethmalosa fimbriata*. It contributes 11.6 percent of the small pelagic fish landings (Jallow, 1994). *E. fimbriata* is a pelagic-neritic species belonging to the family clupeidae found in coastal and estuarine areas. It feeds by filtering phytoplankton, chiefly diatoms and breeds throughout the year in waters of salinities 3.5-38 ppt (Abowei, 2009).

Small pelagic fish are under increasing pressure factors related to the growth of the human population, including overfishing, global climate change, pollution, and habitat degradation (Klemas, 2012).

Almost all large marine ecosystems (LME) are warming except those of California and Humboldt (Belkin, 2009; Sherman et al., 2009). Globally, sea surface temperature (SST) has risen by 0.6 °C in the last past 100 years (IPCC, 2007). In the Canary Current Large Marine Ecosystem (CCLME) SST rose by 0.52 °C from 1957–2006 (Belkin, 2009). According to Klemas (2012), about 40 years ago, ocean productivity began declining, having reached Maximum Sustainable Yield. It is thought that the relationship between climate change and future ocean primary production is determinant for fish and fisheries production (Cushing, 1982).

The two environmental parameters used most often in investigations about relationships between the environmental changes and fish abundance are sea surface temperature (SST) and chlorophyll-a (Chl-a) concentration (phytoplankton biomass). A change of 0.1 °C could be perceived by many fish species and temperature can impact fish in many different ways (Abowei, 2009). Temperature influences fish during spawning, and at the development and survival of the eggs and larvae, as well as influencing distribution, aggregation, migration and schooling behavior of juveniles and adults (Gordoa et al., 2000; Laevastu and Hayes, 1981; Sund et al., 1981). Chlorophyll-a concentration considered as the phytoplankton biomass is the primary food source of any small pelagic species (Zainudhin et al., 2004). It has been shown that phytoplankton production is related to fish landings (Chassot et al., 2007; Ware & Thomson, 2005). According to Butler et al. (1988) Chlorophyll-a concentration above 0.2 mg.m⁻³ indicate the presence of sufficient planktonic activity to sustain a viable commercial fishery.

Understanding how climate variability may affect marine population, especially small pelagic resources, within the objective of proposing a sustainable fisheries management plan, is a challenge. In this sense, this work is to investigate how Sea Surface Temperature and Sea Surface Chlorophyll-a, and their combined effects may be controlling the catch of *Ethmalosa fimbriata* off Senegal between 1999 and 2009.

2. MATERIAL AND METHODS

Fish landed catch data

To study the variability of *Ethmalosa fimbriata* abundance in Senegalese waters, monthly landed catch data from 1999-2009 were used. These data were obtained from the statistical

section of the Direction of Maritime Fisheries (Direction des Pêches Maritimes) and constitutes the total landings in Senegal. Therefore, data from artisanal catch were used since fisheries in Senegal are conducted exclusively by artisanal fishermen, operating from canoes.

Environmental data

Monthly mean values of sea surface temperature (SST, °C) from remote-sensing data for an 11-year period from the Advanced Very High Resolution Radiometer (AVHRR Pathfinder V5) satellite data at a spatial resolution of 4 km were used.

Monthly mean values of sea surface chlorophyll-a (Chl-a, mg.m⁻³) from remote-sensing data for the same 11-year period from Sea-viewing Wide Field-of-view Sensor (SeaWiFS) satellite data at a spatial resolution of 9 km were also used.

The Data cover the time period between January 1999 and December 2009. Sea Surface Temperature (SST) and chlorophyll-a (Chl-a) were averaged from 12°8' and 16°41' N and from the coast to the 200-m isobaths (Fig.1).




Figure 1. Study area (black line: 200 m isobaths).

Figura 1. Área de estudo.

Statistical analysis between *Ethmalosa fimbriata* catch and the environmental data (SST and Chl-a)

Monthly climatology was calculated in order to analyze the seasonal abundance of *E. fimbriata* in relation to SST and Chl-a variation.

The relationships between SST and Chl-a, and catch of *E. fimbriata* were evaluated. The analysis was based on Pearson's correlation coefficient between catch of *E. fimbriata*, SST and Chl-a. Correlation coefficient has been checked against the hypothesis that it was zero with a 95% confidence level. To facilitate the visualization of the relationship between the climate descriptors and the catch of *E. fimbriata*, scatterplots was produced.

To identify a possible relationship between catch of *E. fimbriata* and SST and Chl-a, a simple linear regression model (lm) was used for describing paired data sets that are related in a linear manner. The package R (version 2.14.1) was used to carry out this study. Seasons were defined as follows: winter (January, February, March), spring (April, May, June); summer (July, August, September), and autumn (October, November, December).

In the analysis, the catch of *E. fimbriata* was considered as the response (output or y) variable in order to facilitate the analyses and visual comparison of general trends. The explanatory or independent variables used are: the SST and Chl-a. We evaluated the collinearity of these variables with catch of *E. fimbriata*. We also examined whether the explained variance is increased when they are considered together or separately. Thus, three types of models were studied: (i) combined influence of SST and Chl-a; (ii) effect of the SST solely, and (iii) finally the influence of the Chl-a by itself on catch of *E. fimbriata*. A natural logarithm was used to transform the SST, Chl-a and catch in order to reduce their variability.

3. RESULTS

Landings variability

The distribution of total landings from the different landing ports in Senegal is shown in figure 2. The landings of *E. fimbriata* are most important in Thies with 64.9%, followed by Fatick (17.9%), Ziguinchor (12.52%), St-Louis (3.21%), Dakar (1.39%), and Louga (0.07). It's worth to emphasize that there are three major landings ports in Thies which are Kayar located in the "Grande Côte" Mbour and Joal situated both in the "Petite Côte" (Fig. 1).

E. fimbriata landings reveal an interannual variability with a decreasing trend (Fig. 3). The average monthly landings between 1999 and 2009 are shown in figure 4. The seasonal pattern of *E. fimbriata* landings shows that maximum of catch, ranged from 21,093.10 to 27,988.36 and occurred from January-June with a peak in February (Fig. 4). The lowest *E. fimbriata* landing is observed in September (7,974.67 t). Figure 5 shows distribution of *E. fimbriata* landings, SST and Chl-a per season, whose 33.7%, 32%, 16.5%, and 17.8% of the total landings were performed in winter, spring, summer, and autumn, respectively.

SST variability from the coast to the 200 m isobaths

SST exhibits a gradual increase from 1999 to 2009 in Senegalese waters (Fig. 3). The monthly composite of SST




Figure 2. Distribution of total landings per region in Senegal.
Figura 2. Distribuição das descargas totais por região no Senegal.




Figure 3. Interannual trends *Ethmalosa fimbriata* landings, SST and Chl-a.

Figura 3. Padrões interanuais das descargas de *Ethmalosa fimbriata*, TSM e Chl-a.

values ranged from 17.50 to 25.86 °C with a mean value of 21.60 °C and a standard deviation of 3.22 °C. The fluctuations in SST have two patterns (Fig. 4). Maximum values were observed from July to December (<22 °C) and lowest values in January to June (>22 °C). June constitutes the transition period between the upwelling season and the warming season. The minimum is observed in February and the highest in September. The SST average per season shows that the lowest temperature is observed in winter and the highest in summer (Fig. 5).




Figure 4. Monthly climatology of *Ethmalosa fimbriata* landings, SST and Chl-a (1999-2009).

Figura 4. Climatologia mensal para as descargas de *Ethmalosa fimbriata*, TSM e Chl-a (1999-2009).

Chl-a variability from the coast to the 200 m isobaths

The interannual variability of Chl-a concentration showed a decreasing trend in the 1999-2009 period (Fig. 3). Monthly Chl-a average from the coast to the 200 m isobaths exhibited two patterns, as did SST (Fig. 4). It showed a period of high Chl-a concentration from January-June with a peak in May and a period of low Chl-a concentration from July-December. The Chl-a concentration values ranging

from 0.37 to 2.45m.gm⁻³, have a mean of 1.06m.gm⁻³, and a standard deviation of 0.63m.gm⁻³. The former indicated that Chl-a concentration was highly variable within this period. The seasonal variability of Chl-a concentration showed that the highest concentration was observed during spring and the minimum in summer (Fig. 5).

Relationship between SST, Chl-a and E. fimbriata landings

Significant and negative correlation was calculated between seasonal variability of SST and *E. fimbriata* landings ($r = -0.87, p < 0.01$) (Fig. 6, left), while Chl-a is positively and significantly correlated with *E. fimbriata* landings ($r = 0.86, p < 0.01$) (Fig. 6, right). The variation of *Ethmalosa* landings per season is strongly negatively linked to SST ($r = -0.98, p < 0.01$) (Fig. 7, left), whereas Chl-a is positively related to *E. fimbriata* landings ($r = 0.86, p < 0.05$) (Fig. 7, right).

Three types of models have been carried out: (1) combined effect of SST and Chl-a and (2) SST and (3) Chl-a effect independently. The seasonal variability of *E. fimbriata* landings is mainly due to the combined effects of SST and Chl-a, supporting 87.92% of the total variance, whereas SST solely explains 75.69% and Chl-a 73.56%. The mixed effect of SST and Chl-a plays a key role on the variability of *E. fimbriata* landings per season, explaining 99.17 % of the total variance, while SST and Chl-a contribute 94.08% and 60.38%, respectively.

4. DISCUSSION

The results of this study demonstrated that there is an environmental effect on *Ethmalosa fimbriata* abundance off coast of Senegal. This can be summarized by the variability of Chl-a through the SST driven effect. The fact that seasonal *E. fimbriata* landings are strongly related to the combined effect




Figure 5. Variability of *Ethmalosa fimbriata* landings, SST and Chl-a per season.

Figura 5. Variabilidade sazonal das descargas de *Ethmalosa fimbriata*, TSM e Chl-a.




Figure 6. Seasonal correlation between *Ethmalosa fimbriata* landings and SST (left) and *Ethmalosa fimbriata* landing and Chl-a (right).
Figura 6. Correlação sazonal entre as descargas de *Ethmalosa fimbriata* e a TSM (esquerda) e *Ethmalosa fimbriata* e Chl-a (direita).




Figure 7. Relationship between *Ethmalosa fimbriata* landings, SST and Chl-a per season.
Figura 7. Relação sazonal entre as descargas de *Ethmalosa fimbriata*, TSM e Chl-a.

of SST and Chl-a could be the consequence of their effect on *E. fimbriata* larvae survival, growth rates, and timing of the reproductive peaks.

E. fimbriata landings and Chl-a showed a decreasing trend from 1999 to 2009, while SST increased. It seems that this situation, decreasing trend of Chl-a and SST warming, was unfavorable to *E. fimbriata* abundance. Besides sea warming, and the decrease of phytoplankton biomass, the general downward trend in *E. fimbriata* landings could also have been the result of non-climatic influences such as fishing effort. However, this hypothesis can be disregarded because the fishing effort during the 1999-2008 period trends is not

correlated ($r = -0.05$) against *E. fimbriata* landings (Fig.8).

Many studies have suggested that the abundance of small pelagic resources is linked to the spatio-temporal variability of environmental factors such as SST, Chl-a, wind stress, sea surface height, among others (Cury & Roy, 1989; Fréon, 1991; Bakun, 1996; Zagaglia et al., 2004). Cold temperature observed from January to June was favorable to *Ethmalosa* abundance. According to Bakun (1996), in this period sea surface temperature is significantly and negatively correlated against upwelled waters rich in nutrient. This wind-driven phenomenon induces an increase in phytoplankton and zooplankton biomass abundance which is the base of the food

chain of small pelagic resources. It has been demonstrated that the seasonal dynamics of upwelling intensity show that spawning seasons and nursery areas of small pelagic fish are linked to enrichment processes (Demarcq & Faure, 2000; Caverivière *et al.*, 2002).

It must be emphasized that this linear model (lm) presents limitations. Since the relationships between landings and climatic terms (SST and Chl-a) seem not to be totally linear (Figs. 6 and 7). Moreover, it has been demonstrated that interaction between environment and biology is complex due to the non-linear population dynamics response (Rothschild, 2000). We think this phenomenon will be better modelled using non-linear model such as a generalized additive models (GAMs). This model provided an accurate prediction of sardine habitat and its seasonal dynamics (Zwolinski *et al.*, 2011).




Figure 8. *Ethmalosa fimbriata* landing (tonnes) versus fishing effort (Hours).

Figura 8. Descargas de *Ethmalosa fimbriata* (toneladas) versus esforço de pesca (horas).

CONCLUSIONS

Overall, this study showed that there is a tight link existing between *Ethmalosa fimbriata* abundance, sea surface temperature and phytoplankton biomass off coast of Senegal. The seasonal abundance of *E. fimbriata* is strongly governed by the combined effect of SST and Chl-a whereby an increase in SST and a decrease in phytoplankton results in a decrease in the fish abundance.

Sustainable use of marine resources requires effective monitoring and management of fish stocks and fish habitat. Remote sensing satellite data, such as sea surface temperature and ocean color (Chl-a) constitute an important tools to help scientists to develop suitable management strategies for a sustainable fisheries resources management.

However, further investigation should be done using GAMs. This model showed that a non-linear multivariate combination of environment variables could be used to define species habitat better than a range of environmental variables analysed in isolation or in linear combination (Zwolinski *et al.*, 2011).

REFERENCES

- Abowei, J.F.N. (2009) - The Morphology, Abundance, Condition Factor and Length-weight Relationship of *Ethmalosa fimbriata* (Bowdich 1825) from Nkoro River Niger Delta, Nigeria. *Advance Journal of Food Science and Technology* 1(1):51-56. Available on-line at <http://maxwellsci.com/print/ajfst/51-56.pdf>
- Bakun, A. (1996) - *Patterns in the Ocean: Ocean Processes and Marine Population Dynamics*. 323pp., University of California Sea Grant, San Diego, California, USA, in cooperation with Centro de Investigaciones Biológicas de Noroeste, La Paz, Baja California Sur, Mexico. ISBN 1-888696-018. Available on-line at <http://www.seararoundus.org/researcher/dpaul/PDF/1997/JournalArticles/ReviewOfPatternsInTheOceans.pdf>
- Belkin, I.M. (2009) - Rapid warming of Large Marine Ecosystems. *Progress in Oceanography* 81: 207–213. doi: 10.1016/j.pocan.2009.04.011
- Binet, D.; Gobert, B.; Maloueki, L. (2001) - El Nino-like warm events in the Eastern Atlantic (6°N, 20°S) and fish availability from Congo to Angola (1964–1999). *Aquatic Living Resources* 14:99–113. Available on-line at http://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_7/b_fdi_57-58/010025345.pdf
- Butler, M.J.A.; Mouchot, M.C.; Barale, V.; LeBlanc, C. (1988) - *The application of remote sensing technology to marine fisheries: an introductory manual*. 165p., FAO Fish. Tech. Pap. 295, FAO - Food and Agriculture Organization of the United Nations, Rome, Italy. ISBN: ISBN 92-5-102694&7. Available on-line at <http://www.fao.org/docrep/003/t0355e/t0355e00.HTM>
- Caverivière A.; Thiam M.; Jouffre D. (eds.) (2002) – *Le poulpe commun Octopus vulgaris. Sénégal et côtes nord-ouest africaines*. 385p., éditions IRD, Paris, France. ISBN: 2709914816. Available on-line at http://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers09-03/010029142.pdf
- Chassot, E.; Mélin, F.; Le Pape, O.; Gascuel, D. (2007) - Bottom-up control regulates fisheries production at the scale of eco-regions in European seas. *Marine Ecology Progress Series* 343: 45-55. doi: 10.3354/meps06919
- Cury, P.; Roy, C. (1989) - Optimal environmental window and pelagic fish recruitment success in upwelling areas. *Canadian Journal of Fisheries and Aquatic Sciences* 46: 670-680. Available on-line at <ftp://134.246.142.21/ifremer/ird/roy/PDF/Cury&Roy,1989.pdf>
- Cushing, D.H. (1982) - *Climate and Fisheries*. 373p., Academic Press, London & New York. ISBN: 9780121997205.
- Demarcq, H.; Faure, V. (2000) - Coastal upwelling and associated retention indices derived from satellite SST. Application to *Octopus vulgaris* recruitment. *Oceanologica Acta*, 23(4):391-408. doi: 10.1016/S0399-1784(00)01113-0

- Diei-Ouadi, Y. (2005) - *Minced sardinella fillets in fishlanding and marketing sites in Senegal*. 20p., FAO Fisheries Circular no. 999, FAO - Food and Agriculture Organization of the United Nations, Rome, Italy. Available on-line at <ftp://ftp.fao.org/docrep/fao/008/a0101e/a0101e00.pdf>
- Fréon, P. (1991) - L'introduction d'une variable climatique dans les modèles globaux de production. In: Philippe Cury & Claude Roy (eds.), *Pêcheries ouest africaines: variabilité, instabilité et changement*, pp.395-424, ORSTOM, Paris, France. ISBN: 2709910403. Available on-line at http://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers11-10/36276.pdf
- Gordoa, A.; Maso, M.; Voges, L. (2000) - Satellites and fisheries: The Namibian hake, a case study. In: D. Halpern (ed.), *Satellites, Oceanography and Society*, pp.193-205, Elsevier, Amsterdam, The Netherlands. DOI: 10.1016/S0422-9894(00)80011-2
- IPCC (2007) - Summary for Policymakers. In: S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor & H.L. Miller (eds.), *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, pp.1-18, Cambridge University Press, Cambridge, U.K. and New York, NY, U.S.A. Available on-line at <http://www.ipcc.ch/pdf/assessment-report/ar4/wg1/ar4-wg1-spm.pdf>
- Jallow, A.M. (1994) - *Utilization of bonga (Ethmalosa fimbriata) in West Africa*. 28p., FAO Fisheries Circular No. 870, FAO - Food and Agriculture Organization of the United Nations, Rome, Italy. Available on-line at <http://www.fao.org/docrep/005/t3536e/t3536e00.htm>
- Klemas, V. (2012) - Remote Sensing of Coastal and Ocean Currents: An Overview. *Journal of Coastal Research* 28(3):576-586. doi: 10.2112/JCOASTRES-D-11-00197.1
- Laevastu, T.; Hayes, M.L. (1981) - *Fisheries Oceanography and Ecology*. 199p., Fishing News (Books), Oxford, U.K. ISBN: 9780852381175.
- Rothschild, B.J. (2000) - Fish stocks and recruitment: the past thirty years. *ICES Journal of Marine Science*, 57: 191-201. doi:10.1006/jmsc.2000.0645
- Sherman, K.; Belkin, I.; Friedland, K.; O'Reilly, J.; Hyde, K. (2009) - Accelerating warming and emergent trends in fishing biomass yields of the world's Large Marine Ecosystems. *AMBIO*, 38(4):215-224. doi:10.1579/0044-7447-38.4.2.215.
- Sund, P.N.; Blackburn, M.; Williams, F. (1981) - "Tuna and Their Environment in the Pacific Ocean: A Review. Oceanogr," *Oceanography and Marine Biology: An Annual Review* 19: 443-512.
- Ware, D.M.; Thomson, R.E. (2005) - Bottom-up ecosystem trophic dynamics determine fish production in the northeast Pacific. *Science*, 308(5726):1280-1284. doi: 10.1126/science.1109049
- Zainuddin, M.; Saitoh, K.; Saitoh, S. (2004) - Detection of potential fishing ground for albacore tuna using synoptic measurements of ocean color and thermal remote sensing in the northwestern North Pacific. *Geophysical Research Letters*, 31(20):L20311. doi: 10.1029/2004GL021000
- Zagaglia, C.R.; Lorenzetti, J.; Stech, J.L. (2004) - Remote sensing data and longline catches of yellowfin tuna (*Thunnus albacores*) in the equatorial Atlantic. *Remote Sensing of Environment*, 93(1-2): 267-281. doi: 10.1016/j.rse.2004.07.015
- Zwolinski, J.P.; Emmett, R.L.; Demer, D.A. (2011) - Predicting habitat to optimize sampling of Pacific sardine (*Sardinops sagax*). *ICES Journal of Marine Science*, 68(5):867-879. doi: 10.1093/icesjms/fsr038