



Journal of the Mexican Chemical Society

ISSN: 1870-249X

editor.jmcs@gmail.com

Sociedad Química de México

México

Pérez Castorena, Ana L.; Arciniegas, Amira; Villaseñor, José Luis; Romo de Vivar, Alfonso

Furanoeremophilane Derivatives from *Psacalium beamanii*

Journal of the Mexican Chemical Society, vol. 48, núm. 1, 2004, pp. 21-23

Sociedad Química de México

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=47548105>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## Investigación

Furanoeremophilane Derivatives from *Psacalium beamanii*Ana L. Pérez-Castorena,<sup>1\*</sup> Amira Arciniegas,<sup>1</sup> José Luis Villaseñor,<sup>2</sup> Alfonso Romo de Vivar.<sup>1</sup>

<sup>1</sup>Instituto de Química, <sup>2</sup>Instituto de Biología, Universidad Nacional Autónoma de México, Circuito Exterior, Ciudad Universitaria, Coyoacán 04510, D.F., México. Tel.: (5255) 56-224412. Fax: (5255) 56-162217. E-mail: alpezerc@servidor.unam.mx

Dedicated to the memory of Dr. Raymundo Cruz Almanza

Recibido el 24 de noviembre del 2003; aceptado el 26 de febrero del 2004

**Abstract.** The phytochemical study of *Psacalium beamanii* afforded the phenylethanoid **1**, three sterols, the furanoeremophilane derivatives **2-10**, and matorone acetate (**12**), which was isolated as a natural product for the first time. Its structure was established by means of spectroscopic data and confirmed by chemical correlation. The optical rotation of decompostin (**4**) was corrected and its identity was corroborated by the X ray analysis of its 2*S*-bromo derivative (**11**).

**Keywords:** *Psacalium beamanii*, Asteraceae, Senecioneae, Tussilaginatae, furanoeremophilane derivatives.

**Resumen.** El estudio fitoquímico de *Psacalium beamanii* permitió el aislamiento del feniletanoide **1**, tres esteroides, los furanoeremofilanos **2-10**, y el acetato de matorona (**12**), el cual fue aislado por primera vez como producto natural. Su estructura se estableció por medio de sus datos espectroscópicos y se confirmó por correlación química. La rotación óptica de la decompostina (**4**) se corrigió y su identidad se corroboró por el análisis de rayos X de su derivado 2*S*-bromo (**11**).

**Palabras clave:** *Psacalium beamanii*, Asteraceae, Senecioneae, Tussilaginatae, derivados de furanoeremofilano.

## Introduction


The genus *Psacalium* (Asteraceae, Senecioneae, Tussilaginatae) comprises about 47 species distributed from southwestern United States to Guatemala [1]. Previous chemical studies have shown that eremophilanes are the main secondary metabolites elaborated by species of this genus [2-5]. *P. decompositum*, *P. palmeri*, *P. peltatum*, *P. sinuatum*, *P. radulifolium* and *Acourtia thurberi* are grouped by the Tarahumara ethnica of northern Mexico, in the so called Matarique complex, and used for the treatment of rheumatism and diabetes, as well as renal, hepatic and gastrointestinal ailments [6]. Ethnomedical information has motivated several research projects such as the evaluation of the antihyperglycemic activity of *P. decompositum* and *P. peltatum* extracts [4,7], and the antimicrobial activity of *P. radulifolium* extracts [5]. Thus, in order to continue with the phytochemical research of Mexican plants of the Senecioneae tribe [8], we studied *Psacalium beamanii* H. Rob. In the present study, we report the isolation of a phenylethanoid (**1**), three sterols, and nine furanoeremophilane derivatives (**2-10**). Additionally, the characterization of matorone acetate (**12**), isolated as a natural product, for the first time, is also described.

## Results and discussion

The phytochemical study of *Psacalium beamanii* afforded the known compounds  $\beta$ -sitosterol, stigmasterol,  $\beta$ -sitosterol glucoside [9], the phenylethanoid icaraside D<sub>2</sub> (**1**) [10], the fura-

noeremophilane derivatives matorone (**2**) [11], matorone (**3**) [2], decompostin (**4**) [12], matorinin (**5**) [2,13], matorin acetate (**6**) [2,13], cacalonol (**7/8**) [14] and peroxycacalonol (**9/10**) [14]. The steroidal compounds were identified by comparison with authentic samples, and compounds **1-10** by comparison of their physical and spectroscopic features with those described in literature.

Cacalonol and peroxycacalonol were optically inactive, suggesting the presence of 1:1 mixtures of the corresponding enantiomers (**7/8** and **9/10** respectively). Compound **4**, identified as decompostin by means of its mp and spectral data, showed a specific rotation ( $[\alpha]_D^{29} -68^\circ$ ) with opposite sign to that reported in literature [12]. Its 2 $\alpha$ -bromo derivative **11** exhibited an  $[\alpha]_D^{29} -426^\circ$ , which also differed from that reported for the same compound ( $[\alpha]_D -41.4^\circ$ ) [12]. In order to clarify this ambiguity, an x-ray crystallographic analysis of the 2*S*-bromo derivative (Fig 1) was performed, thus confirming the structure.






Fig. 1. ORTEP projection of **11**.

re and absolute configuration of decompostin (**4**) and showing that the reported specific rotations of **4** and **11** were mistaken.

In addition to the above mentioned, modified furanoere-mophilane **12** was isolated as a crystalline yellow solid, mp 150-1°. Its molecular formula  $C_{16}H_{12}O_5$  deduced from HREIMS ( $m/z$  284.068,  $[M]^+$ ) as well as its UV ( $\lambda_{max}$  246, 293 and 350 nm) and IR ( $\nu_{max}$  1670, 1585 and 1543  $cm^{-1}$ ) spectra suggested an aromatic structure with a quinoid system. This was corroborated by the two singlet signals at  $\delta$  183.54 and 173.62 (C-6 and C-9 respectively) observed in the  $^{13}C$  NMR spectrum. The presence of an acetate group was deduced by its characteristic signals ( $\delta_H$  2.14,  $\delta_C$  20.85 and  $\delta_C$  170.56) observed in the  $^1H$  and  $^{13}C$  NMR spectra. The remaining signals in both spectra showed a structural relationship with matorone (**3**). The main difference observed in  $^1H$  NMR spectrum of **12** was the downfield shift of the C-13 methylene signal ( $\delta$  5.38), in agreement with the presence of an acetate group bonded to this carbon atom. The structure and identity of **12** were confirmed when matorin acetate (**6**) was transformed into **12** by Jones oxidation. Matorone acetate was reported as an acetylation product of matorone [2]. Nevertheless, this is the first time that **12** is isolated as a natural product and fully characterized by  $^1H$ ,  $^{13}C$ , COSY, HETCOR, and COLOC NMR spectral data.

## Experimental

### General Experimental Procedures

Melting points were determined on a Fisher Jones melting point apparatus and are uncorrected. Optical rotations were determined on a JASCO DIP-360 digital polarimeter. IR spectra were recorded on a Nicolet Magna-IR 750 spectrometer. EIMS data were determined on a JEOL JMS-AX505HA mass spectrometer at 70 eV.  $^1H$  NMR and  $^{13}C$  NMR data were

obtained on a Varian Unity 300 instrument. Chemical shifts were referred to TMS ( $\delta$  0). Standard Varian programs were used for COSY spectra at 300 MHz. HETCOR experiments were obtained for  $^1J_{CH} = 140$  Hz at 75 MHz. COLOC experiments were obtained for  $^nJ_{CH} = 9$  Hz at 75 MHz. Vacuum column chromatographies (VCCs) were performed using Sil G 60 E. Merck, 70-230 mesh.

### Plant Material

*Psacalium beamanii* H. Rob. was collected in Ixtlán, Oaxaca, México, June 1997. A voucher specimen (MEXU 775149) was deposited at the Herbario Nacional, Instituto de Biología, UNAM, México.

### Extraction and Isolation

Dried and ground roots (129 g) were extracted successively with hexane and MeOH at room temperature. Hexane extract (4.47 g) was analyzed by VCC eluting with an hexane/EtOAc gradient system. Fractions eluted with hexane were combined and purified by means of VCC (hexane/EtOAc 99:1) to yield matorinone (**2**, 11.1 mg) [11], mp 169-72°. Fractions eluted with hexane/EtOAc 90:10 afforded matorone acetate (**12**, 44 mg) [2], mp 150-1°. The mother liquors of the last compound and fractions eluted with hexane/EtOAc 85:15 were combined and submitted to VCC (hexane/EtOAc 95:5) to yield decompostin (**4**, 49.4 mg) [12], mp 195-6°,  $[\alpha]_D^{29} -68^\circ$  (c 0.456,  $CHCl_3$ ), and **12** (37.9 mg). Decompostin (**4**, 19 mg) was transformed into 2*S*-bromodecompostin (**11**, 6 mg) as already described [12], mp 188-192° dec,  $[\alpha]_D^{29} -426^\circ$  (c 0.2,  $CHCl_3$ ). MeOH extract (28 g) was purified by VCC eluting with an hexane/ $Me_2CO$  gradient system. Fractions eluted with hexane were analyzed by VCC (hexane/EtOAc 99:1) affording matorin (**5**, 18.1 mg) [2, 13], mp 94-6°, **2** (4.4 mg) and a mixture of  $\beta$ -sitosterol/stigmasterol (25 mg), mp 134-6°. Fractions eluted with hexane/ $Me_2CO$  90:10 were submitted to VCC (hexane/EtOAc 95:5) to yield **4** (123.6 mg) and matorin acetate (**6**, 125.5 mg) [2, 13], mp 84-6°. Fractions eluted with hexane/ $Me_2CO$  80:20 were purified by VCC (hexane/EtOAc 85:15) affording cacalonol (**7/8**, 11.4 mg) [14], mp 205-7°,  $[\alpha]_D^{29} 0^\circ$  (c 0.26,  $CHCl_3$ ) and matorone (**3**, 3 mg) [2], mp 164-70°. Dried and ground aerial parts (476.2 g) were extracted with hexane and MeOH. Hexane extract (12.7 g) was submitted to VCC eluting with an hexane/ $Me_2CO$  gradient system. Fractions eluted with hexane/ $Me_2CO$  95:5 afforded a mixture of  $\beta$ -sitosterol/stigmasterol (25 mg). Fractions eluted with hexane/ $Me_2CO$  90:10 yielded **7/8** (33.4 mg) and peroxycacalonol (**9/10**, 34.7 mg) [14], mp 204-6° dec.,  $[\alpha]_D^{29} 0^\circ$  (c 0.185,  $CHCl_3$ ). The MeOH extract (100 g) was analyzed by VCC eluting with an hexane/ $Me_2CO$  gradient system. Fractions eluted with hexane/ $Me_2CO$  95:5 were purified by VCC (hexane/EtOAc 90:10) yielding a mixture of  $\beta$ -sitosterol/stigmasterol (28 mg), and **12** (10.6 mg). Fractions eluted with hexane/ $Me_2CO$  90:10 and 85:15 were combined and purified by VCC (hexane/EtOAc 85:15) affording **12** (2.1 mg), **7/8** (31.6

mg), and **3** (17.5 mg). Fractions eluted with hexane/Me<sub>2</sub>CO 50:50 and 40:50 were combined and analyzed by VCC (CHCl<sub>3</sub>/MeOH 95:5) to give β-sitosterol glucopyranoside (75.6 mg) [9], mp 285-90°. Fractions eluted with hexane/Me<sub>2</sub>CO 30:70 were treated with charcoal/MeOH and submitted to VCC (CHCl<sub>3</sub>/MeOH 80:20) to yield icariside D<sub>2</sub> (**1**, 148.3 mg) [10], mp 153-5°.

**Maturone acetate (12)**. Yellow crystals from hexane/EtOAc, mp 150-1° [2]. UV (MeOH): λ<sub>max</sub> (log ε) 204 (4.25), 246 (4.45), 293 (3.83), 350 nm (3.71). IR (CHCl<sub>3</sub>): ν<sub>max</sub> 1744, 1670, 1585, 1543 cm<sup>-1</sup>. EIMS *m/z* (rel. int.) 284 [M]<sup>+</sup> (10), 242 [M-42]<sup>+</sup> (15), 224 [M-HOAc]<sup>+</sup> (100), 196 [224-28]<sup>+</sup> (5), 168 [196-28]<sup>+</sup> (16), 139 [168-29]<sup>+</sup> (20), 43 [C<sub>2</sub>H<sub>3</sub>O]<sup>+</sup> (63). HREIMS *m/z* [M]<sup>+</sup> 284.0681 (calcd for C<sub>16</sub>H<sub>12</sub>O<sub>5</sub> 284.0685). <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 8.17 (1H, br d, *J* = 6.6 Hz, H-1), 7.61 (1H, t, *J* = 7.5 Hz, H-2), 7.54 (1H, br d, *J* = 7.2 Hz, H-3), 7.76 (1H, s, H-12), 5.38 (2H, s, H-13), 2.81 (3H, s, H-15), 2.14 (3H, s, Ac). <sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): δ 125.99 (C-1), 132.94 (C-2), 138.49 (C-3), 142.29 (C-4), 130.55 (C-5), 183.54 (C-6), 128.70 (C-7), 152.17 (C-8), 173.62 (C-9), 134.05 (C-10), 121.61 (C-11), 147.10 (C-12), 56.47 (C-13), 23.12 (C-15), 170.56 (C=O, Ac), 20.85 (CH<sub>3</sub>, Ac).

**Oxidation of maturin acetate (6)**. A solution of **6** (27.8 mg) in Me<sub>2</sub>CO (2 ml) at 3° was oxidized with Jones reagent in the usual manner. The residue purified by VCC (hexane/EtOAc 80:20) gave maturone acetate (**12**, 16.4 mg), mp 149-51°.

**X-ray crystallographic data of 11**. Colorless crystal obtained from CHCl<sub>3</sub>/Et<sub>2</sub>O; crystal size 0.206 × 0.144 × 0.052 mm; crystal data: C<sub>17</sub>H<sub>19</sub>BrO<sub>4</sub>, mol. wt = 367.23, monoclinic, space group *P*2<sub>1</sub>, *a* = 7.242 (1) Å, *b* = 8.321 (1) Å, *c* = 14.131 (1) Å, β = 103.535 (2)°, *V* = 827.9 (2) Å<sup>3</sup>, *Z* = 2, *D*<sub>c</sub> = 1.473 Mg/m<sup>3</sup>, *F*(000) = 376, *T* = 293 (2) K, μ = 2.497 mm<sup>-1</sup>, λ (Mo Kα) = 0.71073 Å, 6805 measured intensities (2.86 < θ < 25.03°), -8 ≤ *h* ≤ 8, -9 ≤ *k* ≤ 9, -16 ≤ *l* ≤ 16, collected Bruker Smart Apex CCD diffractometer. Refinement by the full matrix-least squares on *F*<sup>2</sup> method was performed using 2907 reflections (> 2σ (*I*)) with 203 parameters. The largest difference peak was 0.497. The final *R*-factor was 4.04 % and *wR*<sup>2</sup> was 5.6 %.

Flack's parameter = -0.05 (1). Crystallographic data have been deposited at the Cambridge Crystallographic Data Center (CCDC 232110), 12 Union Road, Cambridge, CB2 1EZ, UK.

## Acknowledgement

We are indebted to Rubén Toscano, Rubén Gaviño, María Isabel Chávez, Héctor Ríos, Angeles Peña, Rocío Patiño, Luis Velasco, Javier Pérez, and Gabriela Salcedo for technical assistance.

## References

- Barkley, T. M.; Clark, B. L.; Funston, A. M. *Proceedings of the International Compositae Conference*, Kew 1994. Hind, D. J. N., Ed., Royal Botanical Gardens: Kew, 1996; Vol. 1. p. 613-620.
- Correa, J.; Romo, J. *Tetrahedron* **1966**, *22*, 685-691.
- Joseph-Nathan, P.; Negrete, M. C.; González, M. P. *Phytochemistry* **1970**, *9*, 1623-1628.
- Inman, W. D.; Luo, J.; Joland, S. D.; King, S. T.; Cooper, R. J. *Nat. Prod.* **1999**, *62*, 1088-1092.
- Garduño-Ramírez, M. L.; Trejo, A.; Navarro, V.; Bye, R.; Linares, E.; Delgado, G. *J. Nat. Prod.* **2001**, *64*, 432-435.
- Bye, R.; Linares, E.; Estrada, E., in: *Phytochemistry of Medicinal Plants*, Arnason, J. T.; Mata, R.; Romeo, J. T. Ed., Plenum Press, New York, **1995**, 65.
- Alarcón-Aguilar, F. J.; Jiménez-Estrada, M.; Reyes-Chilpa, R.; González-Paredes, B.; Contreras-Weber, C. C.; Román-Ramos, R. *J. Ethnopharmac.* **2000**, *69*, 207-215.
- Pérez-Castorena, A. L.; Arciniegas, A.; Ramírez A. M. T.; Villaseñor, J. L.; Romo de Vivar, A. *Planta Med.* **2002**, *66*, 645-647.
- Voutquenne, L.; Lavaud, C.; Massiot, G.; Sevenet, T.; Hadi, H. A. *Phytochemistry* **1999**, *50*, 63-69.
- Miyase, T.; Ueno, A.; Takizawa, N.; Kobayashi, H.; Oguchi, H. *Phytochemistry* **1989**, *28*, 3483-3485.
- Brown, P. M.; Thomson, R. H. *J. Chem. Soc. C* **1969**, 1184-1186.
- Rodríguez-Hahn, L.; Guzmán, A.; Romo, J. *Tetrahedron* **1968**, *24*, 477-483.
- Bohlmann, F.; Zdero, C.; Grenz, M. *Chem. Ber.* **1977**, *110*, 474-486.
- Naya, K.; Miyoshi, Y.; Mori, H.; Takai, K.; Nakanishi, M. *Chem. Lett.* **1976**, 73-76.