



Darwiniana

ISSN: 0011-6793

sdenham@darwin.edu.ar

Instituto de Botánica Darwinion
Argentina

Luebert, Federico; Frohlich, Michael W.
FOUR NEW COMBINATIONS IN ARGENTINIAN HELIOTROPIACEAE
Darwiniana, vol. 4, núm. 2, diciembre, 2016, pp. 192-194
Instituto de Botánica Darwinion
Buenos Aires, Argentina

Available in: <http://www.redalyc.org/articulo.oa?id=66949983008>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



FOUR NEW COMBINATIONS IN ARGENTINIAN HELIOTROPIACEAE

Federico Luebert^{1,2} & Michael W. Frohlich³

¹Nees-Institut für Biodiversität der Pflanzen, Universität Bonn, Meckenheimer Allee 170, 53115 Bonn, Germany; fluebert@uni-bonn.de (author for correspondence).

²Departamento de Silvicultura y Conservación de la Naturaleza, Universidad de Chile, 8320000 Santiago, Chile.

³Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3DS, United Kingdom.

Abstract. Luebert, F. & M.W. Frohlich. 2016. Four new combinations in Argentinian Heliotropiaceae. *Darwiniana*, nueva serie 4(2): 192-194.

Four new combinations in the genera *Euploca* and *Myriopus* (Heliotropiaceae) are established: *Euploca aurata*, *E. catamarcense*, *Myriopus breviflorus* and *M. paniculatus* var. *austrinus*.

Keywords. Argentina; Boraginaceae; *Euploca*; *Heliotropium*; *Myriopus*; South America; *Tournefortia*.

Resumen. Luebert, F. & M.W. Frohlich. 2016. Cuatro combinaciones nuevas en Heliotropiaceae argentinas. *Darwiniana*, nueva serie 4(2): 192-194.

Se efectúan cuatro nuevas combinaciones en los géneros *Euploca* y *Myriopus* (Heliotropiaceae): *Euploca aurata*, *E. catamarcense*, *Myriopus breviflorus* y *M. paniculatus* var. *austrinus*.

Palabras clave. Argentina; Boraginaceae; *Euploca*; *Heliotropium*; *Myriopus*; Sudamérica; *Tournefortia*.

INTRODUCTION

Heliotropiaceae is a monophyletic group (Weigend et al., 2014) and is considered as a family (instead of as a subfamily of Boraginaceae s.l.) for reasons of nomenclatural stability and easiness of morphological recognition, which cannot be fulfilled when Boraginaceae is regarded in its broad sense (Luebert et al., 2016). Four genera are accepted within Heliotropiaceae (Diane et al., 2016): *Euploca* Nutt., *Heliotropium* L., *Ixorhea* Fenzl, and *Myriopus* Small, all of which are present in Argentina. *Euploca*, *Ixorhea* and *Myriopus* form a clade sister to a broadly defined *Heliotropium* (Hilger & Diane, 2003; Luebert et al., 2011, Diane et al., 2016). The latter includes *Tournefortia* L. sect. *Tournefortia* and the remaining sections traditionally ascribed to *Heliotropium*, except *H.* sect. *Orthostachys* (R.Br.) G.Don, which was transfe-

red to *Euploca* (Hilger & Diane, 2003). These phylogenetic studies also showed that the traditional *Tournefortia* is polyphyletic, with *T.* sect. *Cyphocyema* I.M.Johnst. forming a clade sister to *Euploca* and recognized under the generic name *Myriopus*. *Euploca* and *Myriopus* are morphologically well defined groups, and their generic circumscriptions are increasingly being accepted in the taxonomic literature (e.g., Degen Naumann & Melo, 2012; Feuillet, 2008, 2012, 2016; Feuillet & Halse, 2016; Hilger & Diane, 2003; Luebert et al., 2011; Melo & Semir, 2006, 2009, 2010).

During the revision of Argentinian Heliotropiaceae, we came across some species in *Heliotropium* and *Tournefortia* still in need of transfer to *Euploca* and *Myriopus*, respectively. The purpose of this contribution is to provide the necessary new combinations that can be used in

future treatments of the family in regional floras, as well as the yet incomplete treatment of Heliotropiaceae (sub Boraginaceae s.l.) of Flora Fanerogámica Argentina (Ariza-Espinar, 2006), now continued as Flora Argentina [Anton & Zuloaga]. Type specimens were examined at G, GH, MSB, P, SGO and US (Thiers, 2015).

RESULTS

Euploca Nutt., Trans. Amer. Phil. Soc. n.s. 5: 189. 1837. TYPE SPECIES: *Euploca convolvulacea* Nutt.

Heliotropium sect. *Orthostachys* (R.Br.) G.Don, Gen. Hist. 4: 361. 1838. TYPE SPECIES: *Heliotropium foliatum* R.Br. (lectotype designated by Johnston, Contr. Gray Herb. 81: 46. 1928).

Observations. Generic description found in Diane et al. (2016). *Euploca* consists of ca. 100 species of pantropical distribution, especially in semiarid zones, with centers of diversity in the Americas and Australia. Ten species have been reported to occur in Argentina [*Euploca aurata* (Phil.) Frohlich, *E. campestris* (Griseb.) Diane & Hilger, *E. catamarcense* (I.M.Johnst.) Frohlich, *E. chrysantha* (Phil.) Diane & Hilger, *E. filiformis* (Lehm.) J.I.M.Melo & Semir, *E. hassleriana* (Chodat) J.I.M.Melo & Semir, *E. mendocina* (Phil.) Diane & Hilger, *E. ocellata* (Cham.) J.I.M.Melo & Semir, *E. pallescens* (I.M.Johnst.) J.I.M.Melo & Semir, *E. procumbens* (Mill.) Diane & Hilger, and *E. salicoides* (Cham.) J.I.M.Melo & Semir; Zuloaga et al., 2008, sub *Heliotropium*].

Euploca aurata (Phil.) Frohlich, **comb. nov.** *Heliotropium auratum* Phil., Anales Mus. Nac., Santiago de Chile 8: 56. 1891. TYPE: Argentina, Catamarca, Antofagasta de la Sierra, I-1885, *F. Philippi* s.n. (lectotype SGO 54425! designated by Förther, Sendtnera 5: 180. 1998; isolectotypes SGO 42267!, GH bc GH00097639!).

Observations. This species is endemic to northwestern Argentina and was long considered as a synonym of *Euploca chrysantha* (Phil.) Diane & Hilger (e.g., Johnston, 1928; Pérez-Moreau &

Cabrera, 1983; Förther, 1998; Correa, 1999, sub *Heliotropium chrysanthum* Phil.). Di Fulvio & Ariza-Espinar (2000) consider it to be a good species, a criterion followed in subsequent publications (Zuloaga et al., 2008).

Euploca catamarcense (I.M.Johnst.) Frohlich, **comb. nov.** *Heliotropium catamarcense* I.M. Johnst., Contr. Gray. Herb. 81: 59. 1928. TYPE: Argentina, Catamarca, Pilciao, 700 m, 20-IV-1917, *P. Jörgensen 1868* (holotype GH bc GH00097604!; isotypes GH bc GH00097605!, LIL bc LIL001430 photo!, LIL bc LIL001431 photo!, SI bc SI001133 photo!, SI bc SI001134 photo!, SI bc 001135 photo!, US bc US00110847!).

Observations. This species is endemic to western Argentina (Johnston, 1928; Zuloaga et al., 2008).

Myriopus Small, Man. S.E. Fl. 1131. 1933. TYPE SPECIES: *Myriopus poliochros* (Spreng.) Small (lectotype designated by Förther, Sendtnera 5: 128. 1998).

Tournefortia sect. *Cyphocyema* I.M.Johnst., Contr. Gray Herb. 92: 72. 1930. TYPE SPECIES: *Tournefortia membranacea* (Gardner) DC.

Observations. Generic description found in Diane et al. (2016). *Myriopus* comprises around 25 species of largely Neotropical distribution from southern United States to northern Argentina with centers of diversity in Brazil and the Caribbean. Four species have been reported to occur in Argentina [*M. breviflorus* (DC.) Luebert, *M. paniculatus* (Cham.) Feuillet, *M. rubicundus* (Salzm. ex DC.) Luebert, and *M. salzmannii* (DC.) Diane & Hilger; Zuloaga et al., 2008, sub *Tournefortia*].

Myriopus breviflorus (DC.) Luebert, **comb. nov.** *Tournefortia breviflora* DC., Prodr. 9: 520. 1845. TYPE: Brazil, Rio [de Janeiro], 1834, *N. Lund* s.n. (holotype G00147013! [photo F neg 7760!, photo MSB!]; isotype P bc P03525470!).

Observations. This species occurs in Argentina, Brazil, Paraguay and Uruguay (Johnston, 1930; Zuloaga et al., 2008).

Myriopus paniculatus var. **austrinus** (I.M. Johnst.) Luebert, **comb. nov.** *Tournefortia paniculata* var. *austrina* I.M. Johnst., Contr. Gray. Herb. 92: 82. 1930. TYPE: Argentina, Tucumán, La Ramada, XII-1914, *M. Lillo 3032M* (holotype GH bc GH00096176 photo!).

Observations. This variety is distributed in northwestern Argentina and southern Bolivia (Johnston, 1930).

ACKNOWLEDGEMENTS

We thank Wilhelm Barthlott, Roberto Kiesling, Maximilian Weigend and Jun Wen for support, and the curators of the consulted herbaria for permitting access to their collections. This research received support from the SYNTHESYS project <http://www.synthesys.info/> which is financed by the European Community Research Infrastructure Action under the FP7 “Capacities” (Grant FR-TAF-1977) Program.

BIBLIOGRAPHY

- Ariza-Espinar, L. 2006. Boraginaceae, Parte 1 (Excepto *Heliotropium*), in L. Ariza-Espinar, A. Calviño, T. E. di Fulvio & N. Dottori (eds.), *Flora Fanerogámica Argentina*, fasc. 97, pp. 3-55. Córdoba: IMBIV (CONICET), Programa PROFLOTA.
- Correa, M. N. 1999. Boraginaceae, in M. N. Correa (ed.), *Flora Patagónica. Colección Científica del Instituto Nacional de Tecnología Agropecuaria* 8(6): 116-145.
- Degen Naumann, R. & J. I. M. Melo. 2012. Nuevas combinaciones en el género *Euploca* Nutt. (Boraginaceae) del Paraguay. *Candollea* 67(2): 368.
- Di Fulvio, T. E. & L. Ariza-Espinar. 2000. Rehabilitación de *Heliotropium auratum* (Boraginaceae). *Kurtziana* 28(2): 279-286.
- Diane, N.; H. H. Hilger; H. Förther; M. Weigend & F. Luebert. 2016. Heliotropiaceae, in J. W. Kadereit & V. Bittrich (eds.), *The Families and genera of vascular plants* 14: 203-211. Cham: Springer International Publishing. DOI: 10.1007/978-3-319-28534-4_17
- Feuillet, C. 2008. Folia Taxonomica 4. Conspectus of *Myriopus* (Heliotropiaceae: Boraginales) in the Guiana Shield. *Journal of the Botanical Research Institute of Texas* 2(1): 263-265.
- Feuillet, C. 2012. Boraginaceae, in P. Acevedo-Rodríguez & M. T. Strong (eds.), *Catalogue of seed plants of the West Indies. Smithsonian Contributions to Botany* 98: 157-174.
- Feuillet, C. 2016. Two new combinations in *Euploca* Nutt. (Heliotropiaceae, Boraginales) and a conspectus of the species of the Guiana Shield area. *PhytoKeys* 61: 101-124. DOI: 10.3897/phytokeys.61.6260
- Feuillet, C. & R. R. Halse. 2016. New combinations from *Heliotropium* to *Euploca* (Boraginales: Heliotropiaceae) for North American species. *Phytoneuron* 2016(5): 1-3.
- Förther, H. 1998. Die infragenerische Gliederung der Gattung *Heliotropium* L. und ihre Stellung innerhalb der subfam. Heliotropioideae (Schrad.) Arn. (Boraginaceae). *Sendtnera* 5: 35-241.
- Hilger, H. H. & N. Diane. 2003. A systematic analysis of Heliotropiaceae (Boraginales) based on trnL and ITS1 sequence data. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 125(1): 19-51. DOI: 10.1127/0006-8152/2003/0125-0019
- Johnston, I. M. 1928. Studies in the Boraginaceae VII.1. The South American species of *Heliotropium*. *Contributions from the Gray Herbarium of Harvard University* 81: 3-73.
- Johnston, I. M. 1930. Studies in the Boraginaceae VIII.1. Observations on the species of *Cordia* and *Tournefortia* known from Brazil, Paraguay, Uruguay and Argentina. *Contributions from the Gray Herbarium of Harvard University* 92: 3-89.
- Luebert, F.; G. Brokamp; J. Wen; M. Weigend & H. H. Hilger. 2011. Phylogenetic relationships and morphological diversity in Neotropical *Heliotropium* (Heliotropiaceae). *Taxon* 60(3): 663-680.
- Luebert, F.; L. Cecchi; M. W. Frohlich; M. Gottschling; C. M. Williams; H. H. Hilger; K. E. Hasenstab-Lehman; J. S. Miller; M. Mittelbach; M. Nazaire; M. Nepi; D. Nocentini; D. Ober; R. G. Olmstead; F. Selvi; M. G. Simpson; K. Sutorý; B. Valdés; G. K. Walden & M. Weigend [Boraginales Working Group]. 2016. Familial classification of Boraginales. *Taxon* 65(3): 502-522. DOI: 10.12705/653.5
- Melo, J. I. M. & J. Semir. 2006. *Euploca rodaliae* J. I. M. Melo & Semir - a new species of *Euploca* (Heliotropiaceae) from Brazil. *Candollea* 61(2): 453-456.
- Melo, J. I. M. & J. Semir. 2009. Two new Brazilian species and new combinations in *Euploca* (Heliotropiaceae). *Kew Bulletin* 64(2): 285-289.
- Melo, J. I. M. & J. Semir. 2010. Taxonomia do gênero *Euploca* Nutt. (Heliotropiaceae) no Brasil. *Acta Botanica Brasílica* 24(1): 111-132. DOI: 10.1590/S0102-33062010000100012
- Pérez-Moreau, R. L. & A. L. Cabrera. 1983. Boraginaceae, in A. L. Cabrera (ed.), *Flora de La Provincia de Jujuy, República Argentina. Colección Científica del Instituto Nacional de Tecnología Agropecuaria* 8(8): 247-291
- Thiers, B. [continuously updated, accessed 2015]. Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium; available from: <http://sweetgum.nybg.org/ih/> (accessed 2015).
- Weigend, M.; F. Luebert; M. Gottschling; T. L. P. Couvreur; H. H. Hilger & J. S. Miller. 2014. From capsules to nutlets –Phylogenetic relationships in the Boraginales. *Cladistics* 30(5): 508-518. DOI: 10.1111/cla.12061
- Zuloaga, F. O.; O. Morrone & M. J. Belgrano. 2008. Catálogo de las plantas vasculares del Cono Sur (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay). *Monographs in Systematic Botany from the Missouri Botanical Garden* 107: 1-3348.