

Metadata of the Book that will be visualized online

Book Title	Medicinal Plants of Central Asia: Uzbekistan and Kyrgyzstan
Book SubTitle	
Copyright Year	2013
Copyright Holder	the Editor

Editor	Family Name	Eisenman
	Particle	
	Given Name	Sasha
	Suffix	
	Division	College of Liberal Arts, School of Environmental Design
	Organization	Temple University
	Address	580 Meetinghouse Rd, 19122, Ambler, Pennsylvania, USA
	Email	eisenman@temple.edu
Editor	Family Name	Struwe
	Particle	
	Given Name	Lena
	Suffix	
	Division	School of Enviro and Biological Sciences, Dept of Plant Biology and Pathology
	Organization	Rutgers University
	Address	08901, New Brunswick, New Jersey, USA
	Email	struwe@aesop.rutgers.edu
Editor	Family Name	Zaurov
	Particle	
	Given Name	David
	Suffix	
	Division	School of Enviro and Biological Sciences, Dept. of Plant Biology and Pathology
	Organization	Rutgers University
	Address	59 Dudley Rd, 08901, New Brunswick, New Jersey, USA
	Email	zaurov@aesop.rutgers.edu

Medicinal Plants of Central Asia:
Uzbekistan and Kyrgyzstan

1

2

Uncorrected Proof

Uncorrected Proof

Sasha Eisenman • Lena Struwe • David Zaurov 3
Editors 4

Medicinal Plants of 5
Central Asia: Uzbekistan 6
and Kyrgyzstan 7

Uncorrected Proof

8	<i>Editors</i>		22
9	Sasha Eisenman	David Zaurov	23
10	College of Liberal Arts, School of Environmental Design	School of Enviro and Biological Sciences	24
11	Temple University	Dept. of Plant Biology and Pathology	25
12	580 Meetinghouse Road	Rutgers University	26
13	Ambler, PA 19122	59 Dudley Road	27
14	USA	New Brunswick, NJ 08901	28
15		USA	29
16	Lena Struwe		
17	School of Enviro and Biological Sciences		
18	Dept of Plant Biology and Pathology		
19	Rutgers University		
20	New Brunswick, NJ 08901		
21	USA		

30 ISBN 978-1-4614-3911-0 ISBN 978-1-4614-3912-7 (eBook)

31 DOI 10.1007/978-1-4614-3912-7
32 Springer New York Heidelberg Dordrecht London

33
34 Library of Congress Control Number: 2012939119

35
36 © The Editor 2013

37 This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of
38 translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or informa-
39 tion storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted
40 from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being
41 entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only
42 under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer.
43 Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective
44 Copyright Law.

45 The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific
46 statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

47 While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the
48 publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with
49 respect to the material contained herein.

50
51 Printed on acid-free paper

52
53 Springer is part of Springer Science+Business Media (www.springer.com)

[AU1] In ancient times, people used the gifts of nature found in their surrounding environments to 55
treat their illnesses. Medicinal plants were of great significance, and the utilization of various 56
plants in folk medicine has a very long history. As far back as 3000 BCE, herbs such as thebaic 57
poppy, rhubarb, ginseng, etc., were well known. Hippocrates listed around 200 different 58
medicinal herbs. In the first century, Dioscorides described about 400 medicinal plants, and the 59
Avesta, the holy book of the Zoroastrians, included a thousand plants. In the eleventh century, 60
Al-Beruni and Avicenna, two great scholars of Central Asia, made important contributions to 61
the science of medicinal plants. Al-Beruni conceived a new area of science concerning medi- 62
cal herbs, now called pharmacognosy, and classified and described numerous plant species. In 63
the year 1025, Avicenna gave the world *The Canon of Medical Science*, where he described the 64
herbs that were most widely researched and used in medical practice of the time. 65

[AU2] Today, many of those plants are still used in medicine in Central Asia. Many centuries of 66
herbal use has proven that plants contain substances that have healing power. The experience 67
from folk medicine has also shown that different parts of each plant often have different effects 68
and, therefore, are used for different diseases, for example, roots for one type of disease and 69
the aboveground parts for another. Similarly, leaves, flowers, fruits, and seeds may have differ- 70
ent medicinal uses. Active compounds usually accumulate in large amounts in only certain 71
parts of a plant (Wink 1999). The amounts of active substances in a plant, and consequently 72
their physiological effect when taken as a medicine, significantly fluctuate depending on the 73
season of the year, habitat, altitude, yearly climatic conditions, soil composition, and other 74
factors (Evans 2002). 75

[AU3] There are more than 20,000 plant species in the former Soviet Union. Of these, 4,500 grow 76
in Uzbekistan and 4,100 in Kyrgyzstan (Komarov 1934; Pratorov 1998; Umralina and Lazkov 77
2008). There are about 35,000–70,000 plants used in folk and scientific medicine worldwide 78
(Hamilton 2004). As of 2004, at least 200,000 phytochemicals (excluding DNA-encoded pro- 79
teins and peptides) have been characterized, but this is still thought to represent only a small 80
percentage of phytochemicals that exist in nature (Raskin and Ripoll 2004). This further indi- 81
cates the importance of drugs of herbal origin for folk and modern medicine. Currently, more 82
than 400 wild and cultivated medicinal plants in Uzbekistan have been studied and described 83
and more than 200 in Kyrgyzstan as well (Nikitina 1962). However, many medicinal plants 84
found in Uzbekistan and Kyrgyzstan have not been thoroughly scientifically evaluated for their 85
potential value in modern medicine. 86

Due to the increased interest in medicinal plants of researchers from different countries, the 87
issue of preservation of the natural environment becomes important and, in particular, the con- 88
servation of medicinal plants in their original habitat. Habitat destruction and environmental 89
pollution are factors that strongly affect medicinal plants in the wild. This complex issue is 90
the subject of international agreements, which are united under the general concept of environ- 91
mental preservation. For adequate conservation, it is important to identify the plant species that 92
are most threatened due to over-collection in the wild. These species must receive the highest 93
prioritization for preservation. It is important to bring the most utilized plants in medicine and 94
veterinary science into cultivation with the goals of increasing the content of basic active com- 95
pounds in the plants and providing a sustainable source of plant material. With the implemen- 96

97 tation of new agricultural practices, the industrial and medical sectors can be supplied with
98 necessary amounts of high-quality product without depleting wild populations. Additional
99 research is necessary to identify plants that have medicinal properties and to scientifically vali-
100 date their pharmacological activities. It is important to conduct these efforts with the involve-
101 ment of a wide circle of international researchers. Information exchange, job creation, and
102 joint conferences will undoubtedly help researchers in their work and will also increase the
103 conservation of the rich floras of Central Asian republics. A logical starting point for such
104 systematic research would be the plants that were studied by our great ancestors and have tra-
105 ditionally been used in folk medicine in the different regions of Central Asia.

106 More than 200 of the most important medicinal plants of Central Asia are listed in this
107 book, and it includes many whose medicinal uses and activities are being compiled for the first
108 time. Most of the plants described grow wild in Central Asia, and some are endemic (e.g.,
109 *Vinca erecta* and *Ajuga turkestanica*). This book is aimed at scientists engaged in research on
110 medicinal plants; physicians; as well as students of biology, pedagogy, agriculture, forestry,
111 pharmacology, and medicine. This book is also a valuable reference for biodiversity conserva-
112 tion efforts and protection of rare and endangered species of the Central Asian flora.

113 We would like to warn our readers that conducting self-treatment with herbs and herbal
114 preparations is dangerous. Medicinal plants can contain extremely strong physiologically
115 active compounds and are often very poisonous. Without the proper recommendations of a
116 medical doctor, no preparations of medicinal plants should be taken. The information in this
117 book is not to be used to diagnose or treat any medical conditions.

118
119
120
121
122

Dr. Ravshanbek D. Kurbanov
Dr. Khasan Ch. Buriev
Dr. Djamin A. Akimaliev
Dr. Ilya Raskin

[Au4]	Medicinal Plants of Central Asia:	123
	Uzbekistan and Kyrgyzstan	124

Edited by 125

Sasha W. Eisenman, Lena Struwe, and David Zaurov 126

Translations by 127

David E. Zaurov, Sasha W. Eisenman, Dilmurad A. Yunusov, and Venera Isaeva 128

These below images are reproduced under the following Creative Commons licenses: 129

Creative Commons Attribution-ShareAlike 3.0 Unported (CC BY-SA 3.0) 130

Gleditsia triacanthos: Georg Slickers, Andrew Butko 131

Lithospermum officinale and *Thalictrum foetidum*: Petr Filippov 132

Xanthium strumarium: Stan Shebs 133

Creative Commons Attribution-Share Alike 2.1 Spain 134

Gleditsia triacanthos: Luis Fernández García 135

Creative Commons Attribution-Share Alike 2.0 Generic 136

Nigella sativa: Pamela J. Eisenberg 137

138

Uncorrected Proof

Uncorrected Proof

Acknowledgments

139

[AU5]

[AU6]

This collaborative project came into being as a result of the International Cooperative Biodiversity Groups (ICBG) Central Asia Project. The ICBG program is managed by the Fogarty International Center of the National Institutes of Health (NIH) and is currently funded by the NIH, the National Science Foundation (NSF), the US Department of Agriculture (USDA) National Institute of Food and Agriculture, the Department of Energy (DOE) Office of Biological and Environmental Research, and the National Oceanographic and Atmospheric Administration (NOAA) Oceans and Human Health Program. We would like to express our great appreciation to the many photographers who have allowed us to use their images in this book. SWE would like to thank Hillary, Elijah, and Avishai for putting up with the long hours devoted to the production of this book. DZ would like to thank Dr. C. Reed Funk for his encouragement and support.

140
141
142
143
144
145
146
147
148
149
150

Uncorrected Proof

Uncorrected Proof

Contents

151

1	The Geography, Climate and Vegetation of Kyrgyzstan	1	152
	Djamin A. Akimaliev, David E. Zaurov, and Sasha W. Eisenman		153
2	The Geography, Climate and Vegetation of Uzbekistan.....	5	154
	Igor V. Belolipov, David E. Zaurov, and Sasha W. Eisenman		155
3	A Short History of Medicinal Plant Use in Central Asia	9	156
	Dr.Anvar G. Kurmukov, Anarbek A. Akimaliev		157
4	Phytochemistry of Medicinal Plants.....	13	158
	Anvar G. Kurmukov		159
5	The Medicinal Plants of Uzbekistan and Kyrgyzstan.....	15	160
	David E. Zaurov, Dr.Igor V. Belolipov, Anvar G. Kurmukov,		161
	Ishembay Sodembekov, Anarbek A. Akimaliev, and Dr.Sasha W. Eisenman		162
	Appendix 1 English-Russian Translations of Botanical and Ecological Terms.....	275	163
	Appendix 2 English-Russian Translations of Chemical Terms.....	283	164
	Appendix 3 English-Russian Translations of Medical Terms	293	165
	References	301	166
	General Index.....	321	167
	Index to Plant Species.....	335	168

[AU7]

Uncorrected Proof

Contributors

169

Anarbek A. Akimaliev Biology and Soil Science Institute of the National Academy of Science of Kyrgyzstan, Bishkek, The Kyrgyz Republic	170 171
Djamin A. Akimaliev Kyrgyz Agricultural Research Institute, Bishkek, The Kyrgyz Republic	172 173
Igor V. Belolipov Tashkent State Agrarian University, Tashkent, The Republic of Uzbekistan	174
Khasan Ch. Buriev Tashkent State Agrarian University, Tashkent, The Republic of Uzbekistan	175 176
Sasha W. Eisenman Department of Landscape Architecture and Horticulture, Temple University, College of Liberal Arts, School of Environmental Design, Ambler, PA, USA	177 178
Ravshanbek D. Kurbanov The Specialized Center for Cardiology, Tashkent, The Republic of Uzbekistan	179 180
Anvar G. Kurmukov The Specialized Center for Cardiology, Tashkent, The Republic of Uzbekistan	181 182
Ilya Raskin Department of Plant Biology and Pathology, Rutgers University – School of Environmental and Biological Sciences, New Brunswick, NJ, USA	183 184
Ishembay Sodembekov Kyrgyz Botanical Garden of the National Academy of Kyrgyzstan, Bishkek, The Kyrgyz Republic	185 186
David E. Zaurov Department of Plant Biology and Pathology, Rutgers University, School of Environmental and Biological Sciences, New Brunswick, NJ, USA	187 188

Author Queries

Chapter No.: FM 0001543047

Queries	Details Required	Author's Response
AU1	Please check if "thebaic poppy" should be changed to "thebaine poppy".	
AU2	Please check if edit to the sentence starting "Al-Beruni conceived...." is okay.	
AU3	Please check if the name of the book " <i>The Canon of Medical Science</i> " should be changed to " <i>The Canon of Medicine</i> ".	
AU4	Should this page appropriate here? Kindly confirm.	
AU5	Please check if "Office of Biological and Environmental Research" should be edited as "Office of Science, Biological and Environmental Research".	
AU6	Please check if "National Oceanographic and Atmospheric Administration (NOAA) Oceans and Human Health Program" should be edited as "National Oceanic and Atmospheric Administration (NOAA) Oceans and Human Health Initiative".	
AU7	Author Ishembay Sodembekov does not match with COP of Chap. 5. Please check.	

Uncorrected Proof

1 The Geography, Climate and 2 Vegetation of Kyrgyzstan

1

3 Djamin A. Akimaliev, David E. Zaurov,
4 and Sasha W. Eisenman

5 Kyrgyzstan is a mountainous country in the northeastern part
6 of Central Asia. The Kyrgyz Republic shares borders to the
7 south and southeast with Tajikistan and China, to the north
8 and northwest with Kazakhstan, and with Uzbekistan to the
9 west. The country covers 198,500 km² (76,621 sq miles) and
10 has a population of approximately 5.3 million. Kyrgyzstan is
11 divided into seven provinces (Fig. 1.1).

12 The highest point of elevation is in the Kakshaal-Too range,
13 along the Chinese border, where Jengish Chokusu (Pik Pobedy)
14 is the highest peak at 7,439 m (24,400 ft). The lowest point of
15 elevation, 132 m (433 ft) above sea level, occurs along the
16 Kara Darya River in the Fergana Valley. Other notable valleys
17 are the low-montane Talas and Chui valleys, the mid-montane
18 Issyk-Kul and Middle Naryn valleys, and the high-montane
19 Ak-Say and Alai valleys. Ninety-four percent of the country is
20 montane with the Tian Shan mountain system covering the
21 major portion of the country. Lake Issyk-Kul, in the north
22 western Tian Shan, is the largest lake in Kyrgyzstan and the
23 second largest mountain lake in the world.

24 The principal river in Kyrgyzstan is the Naryn, which
25 flows west through the Fergana Valley into Uzbekistan.
26 There it meets another of Kyrgyzstan's major rivers, the Kara
27 Darya. They merge to form the Syr Darya, which eventually
28 flows into the Aral Sea. Heavy water usage for irrigation in
29 Uzbekistan now causes the river to run dry before reaching
30 the sea. The Chu River also briefly flows through Kyrgyzstan
31 before entering Kazakhstan.

D.A. Akimaliev (✉)
Kyrgyz Agricultural Research Institute,
73/1, Timur Frunze St., Bishkek 720027, The Kyrgyz Republic
e-mail: krif@mail.kg

D.E. Zaurov
Department of Plant Biology and Pathology, Rutgers University,
School of Environmental and Biological Sciences,
20 Ag. Extension Way, New Brunswick, NJ 08901, USA

S.W. Eisenman
Department of Landscape Architecture and Horticulture, Temple
University, College of Liberal Arts, School of Environmental Design,
580 Meetinghouse Rd., Ambler, PA 19002, USA

32 Kyrgyzstan's climate is influenced by its position between
33 the temperate and sub-tropical zones, its high elevation and
34 its distance from oceans. These conditions cause intense sun
35 radiation, lack of precipitation and a harsh continental cli-
36 mate. The mountain relief causes altitudinal zoning of cli-
37 mate parameters such as temperature and moisture. In July
38 the average air temperature in the lowlands can range from
39 17 to 40°C (62.6–104°F), whereas at a higher elevation the
40 temperature may be much cooler. During winters frosts may
41 occur in all regions of Kyrgyzstan.

42 The southwestern Fergana Valley is dry-subtropical and
43 hot in summer, with air temperatures reaching 40°C (104°F).
44 The plains of southern and northern Kyrgyzstan have a hot
45 desert or semi-desert climate and in these areas air tempera-
46 tures can reach 35–40°C (95–104°F) during the summer
47 months as well (Mamitov 1965). The northern foothills have
48 a temperate climate and the climate in the Tian Shan moun-
49 tain system varies from dry continental to polar, depending on
50 elevation. The mountain regions have steppe, meadow-steppe,
51 meadow, and high-mountain tundra climates and the highest
52 areas are permanently snow covered (Ryazantseva 1965).

53 The yearly precipitation in Kyrgyzstan varies between
54 100 and 1,000 mm (3.9–39 in.) and is distributed unevenly
55 throughout the country. The highest levels of precipitation
56 (>900 mm; 35.4 in.) occur in the mid-belt of the southwest-
57 ern slopes of the Fergana and Chatkal ranges, the high moun-
58 tain areas of the northern slopes in the Kyrgyz Range, in the
59 Kemin valley, and in the eastern Issyk-Kul area. The Talas
60 and Chui valleys receive from 250 up to 500 mm (9.9–
61 19.7 in.) precipitation and the valley and foothills in Fergana
62 receive from 300 to 700 mm (11.8–27.6 in.) per year. Most
63 of the internal and central areas of the Tian Shan system
64 average 200–300 mm (7.9–11.8 in.) of rain annually and
65 western Issyk-Kul and portions of Fergana may have less
66 than 150 mm (5.9 in.) per year. On average, the foothills of
67 the north and the eastern Issyk-Kul basin receive 15–20 cm
68 (5.9–7.9 in.) of snow annually. The amount of snow fall in
69 the high-altitude valleys of the Tian Shan is distributed very
70 unevenly. The Ak-Shiyrak and Karakol valleys receive an



Fig. 1.1 Kyrgyzstan with provincial boundaries

71 average of 3 cm (1.2 in.) of snow whereas, on average, the
 72 Karakudjur valley receives 9 cm (3.5 in.). The mid-altitude
 73 and high-altitude belts of the Fergana range can receive
 74 upwards of 150 cm (59 in.).

75 The flora of the Kyrgyzstan contains more than 4,100 spe-
 76 cies of vascular plants (Umralina and Lazkov 2008). Around
 77 1,600 species have economic and/or useful value including
 78 species for fodder (450 species), for honey production (300
 79 species) for medicinal use (200 species), for essential oils
 80 (62 species), and for food (50 species; Nikitina 1962). The
 81 largest portion of land used for agriculture is devoted to the
 82 cultivation of grain crops. Vegetables, oil crops and cotton
 83 are also grown to a lesser extent (Abaihanova et al. 2006).

84 The distribution of the vegetation follows a pattern of ele-
 85 vation belts and is mainly influenced by land relief, climate,
 86 and soil zones. Twenty-two classes of ecosystems have been
 87 identified in Kyrgyzstan. The ecosystems are unevenly distrib-
 88 uted throughout the country. Fourteen of the ecosystems occur
 89 in middle mountain zone (2,000–3,000 m), which occupies

just 30% of the country’s area. The Western and Central Tian 90
 Shan regions have 16 and the Alai has 13 ecosystems. In the 91
 Northern Tian Shan and Issyk-Kul regions 10 ecosystems can 92
 be found. The southern Kazakhstan biogeographic region has 93
 five of the ecosystems and the Fergana valley has the fewest 94
 with only three (Ministry of Environmental Protection 1998). 95

The ecosystems include deciduous and evergreen forests, 96
 shrublands, grasslands (savannahs, meadows and steppes), 97
 deserts, various wetlands and bodies of water. The river 98
 floodplains have shrubby forests (tugai) with *Rhamnus* spp., 99
Salix spp., *Rosa* spp., etc. The valleys and foothills contain 100
 perennial herbs, ephemerals, and on stony soils, thorny herbs 101
 and succulents. In the mid-belt of the mountains, depending 102
 on precipitation levels, there are deserts, steppes, meadows 103
 and shrublands. The high elevation areas consist of glacial and 104
 subglacial areas as well as cryophylic steppes, alpine mead- 105
 ows and deserts. The majority of these deserts are *Artemisia* 106
 spp. dominated, fewer being *Salsola* spp. deserts, and a very 107
 few dominated by *Ephedra* spp. (Golovkova 1990). 108

109 In spring and in the beginning of the summer, *Astragalus*
110 spp., *Crocus* spp., *Gagea* spp., *Iris kolpakowskiana*,
111 *Ranunculus* spp. and *Tulipa* spp., as well as medicinal plants
112 like *Betonica* spp., *Salvia* spp., *Thymus* spp., *Ziziphora* spp.,
113 etc. are found in the low- and middle mountain steppes.
114 Meadows are less common than steppes, but they have a
115 diverse floral composition including *Aconitum* spp.,
116 *Androsace ovczinnikovii*, *Aster alpinus*, *Cerastium* spp.,
117 *Codonopsis clematidea*, *Delphinium* spp., *Erigeron auranti-*
118 *acus*, *Gentiana karelinii*, *Primula algida*, etc.

119 Only about 4.0% of Kyrgyzstan is covered with forests.
120 Spruce and juniper forest account for a major portion of the
121 forested area and over 350 herbaceous plant species can be
122 found in the spruce forests. In the southern part of Kyrgyzstan
123 the world's largest naturally occurring nut tree forests occupy
124 about 608,500 ha (2,350 sq miles). These forests occur
125 mainly in the Chatkal and Fergana ranges at an elevation of
126 1,000–2,200 m (3,280–7,218 ft). Many of the species in
127 these forests are wild relatives of domesticated nut and fruit
128 crops. These wild populations are important reservoirs of
129 genetic diversity, which can be utilized in breeding programs
130 to develop cultivars with cold tolerance, disease and insect
131 resistance, and other important characteristics. The main for-
132 est species is *Juglans regia* (Persian walnut), which occupies
133 about 40,000 ha (155 sq miles). Other wild fruits and nuts
134 include *Prunus amygdalus* (almond) and *Pistacia vera* (pis-
135 tachio), *Berberis oblonga*, *Cerasus mahaleb* and *C. tians-*
136 *chanica*, *Crataegus songorica* and *Cr. turkestanica*, *Malus*
137 *kirghisorum* and *M. sieversii*, *Prunus sogdiana*, *Pyrus com-*
138 *munis*, *P. korshinskyi*, and *P. regelii*.

139 Due to their extreme environment and climate, portions of
140 the country have limited or no biodiversity. These areas

141 account for around 45% of the country and consist of high
142 altitude areas (above 3,500 m [11,483 ft]) of rock and
143 glaciers, open areas of rock, gravel or clay, and deserts.

144 There are 65 plant species on the list of endangered spe-
145 cies in the *Red Data Book of Kirghiz SSR* (1985). Sultanova
146 et al. (1998) published a more up-to-date list with 386 spe-
147 cies recommended for inclusion to the red book. At the
148 present time there is a need for the establishment of orga-
149 nized medicinal plant farming and for the protection of
150 endangered species. Many of the plants used in Kyrgyz folk
151 medicine have not been studied using modern scientific
152 techniques. Pharmacological studies are necessary to char-
153 acterize the biological activity of the medicinal plants and
154 their components. Folk medicine is an invaluable source of
155 information on the properties and activities of medicinal
156 plants and for discovery of novel medicines. Further study
157 of the Kyrgyz ethno-medicine will help facilitate the
158 identification of new medicinal plants, which may possibly
159 serve as sources for new pharmaceuticals. Further expan-
160 sion of botanical and floristic research is also necessary,
161 including detailed mapping of all medicinal plant resources
162 and determination of regions for cultivation of valuable and
163 rare species.

164 Currently all ecosystems are subject to human influence.
165 The overall biodiversity of Kyrgyzstan is threatened as a
166 result of human disturbance. Over-grazing has degraded
167 many of the plant communities and over-use has greatly
168 reduced the overall size of forest ecosystems. Intensifying
169 anthropogenic influence threatens the diversity of the natural
170 resources of the country. Preservation and conservation of
171 these unique natural resources is of extreme importance for
172 future generations of Kyrgyz people.

1 **The Geography, Climate**
2 **and Vegetation of Uzbekistan**

3 Igor V. Belolipov, David E. Zaurov,
4 and Sasha W. Eisenman

5 Uzbekistan is a country in Central Asia that extends from the
6 foothills of the Tian Shan and Pamir mountains in the east to
7 just west of the Aral Sea. In the north Uzbekistan borders
8 Kazakhstan, in the east and southeast Kyrgyzstan and Tajikistan,
9 in the west Turkmenistan, and in the south Afghanistan. The
10 country covers 447,400 km² (172,742 sq miles) and has a popu-
11 lation of about 26 million. Uzbekistan is divided into 12 prov-
12 inces and 1 autonomous republic (Fig. 2.1).

13 The highest point of elevation is in the Gissar mountain
14 range at 4,643 m (15,233 ft), and the lowest point of elevation
15 is the Sarykamysh depression at 20 m (ca. 65.6 ft) below sea
16 level. About 80% of Uzbekistan's land consists of plains and
17 deserts. The vast Kyzylkum desert lies in central Uzbekistan
18 and is largely uninhabited except for mining towns.

19 There is a wide spectrum of natural environments from
20 the hot sand and gypsum deserts of Kyzylkum to the eternal
21 snows and glaciers of the Pamiro-Alai mountains. All val-
22 leys receive their water from glaciers in the Tian Shan and
23 Pamiro-Alai mountains. Uzbekistan's two most important
24 rivers, the Syr Darya and Amu Darya, flow from the Tian
25 Shan and Pamiro-Alai mountain ranges to the Aral Sea.

26 The climate of Uzbekistan is continental with predomi-
27 nance towards harsh continental. It is characterized by low
28 precipitation (70–100 mm [~2.75 to 3.94 in.] per year) in the
29 plains of the northern-western part of the country and up to
30 1,200 mm (47.25 in.) of precipitation in mountainous regions.

Over 70% of the precipitation falls in the autumn to spring
31 period, with a maximum in March and April. Summers in
32 Uzbekistan are long, dry, and hot, summer rains are very
33 rare, and summer temperatures may reach 45°C (113°F). In
34 the south the winter is mild, but sometimes with considerable
35 frosts. In the northern regions winters are cold and tempera-
36 tures may drop to –37°C (–35°F).
37

38 The flora of Uzbekistan contains more than 4,500 vascular
39 plants in 650 genera, in 115 families (Chemonics International
40 Inc. 2001). More than 4,000 species of algae and more than
41 2,000 species of fungi also occur in Uzbekistan (National
42 Biodiversity Strategy Project Steering Committee 1998). The
43 most species-rich plant families account for a large portion of
44 the flora. These families include Asteraceae (600 species),
45 Fabaceae (450 species), Poaceae (>250 species), Brassicaceae,
46 Lamiaceae, Rosaceae, Boraginaceae and Apiaceae.

47 Agriculture and cultivated crops occupy considerable
48 areas of irrigated and non-irrigated land. Some of the major
49 crops are cotton, maize (corn), alfalfa, wheat, barley, sor-
50 ghum, rice, mulberry for silkworm culture, vegetables, mel-
51 ons, fruit trees, and others. The natural vegetation of
52 Uzbekistan is a very rich source of fodder (more than 1,700
53 species), medicinal plants (600 species) and plants with
54 essential oils (>650 species), saponins (>100 species), and
55 tannins (ca. 400 species).

56 The vegetation of Uzbekistan is divided into four main eco-
57 systems. The main cause for ecosystem zonation is change in
58 hydrothermal conditions. These zones form belts which are
59 directly correlated to an increase in precipitation and eleva-
60 tion. As elevation increases there are changes in environmen-
61 tal conditions. Growing periods become shorter, temperature
62 decreases and precipitation increases. Due to the increase in
63 precipitation water is no longer a limiting factor above 2,500 m
64 (~8,200 ft). Diverse soil conditions, in combination with the
65 environmental conditions, result in a great diversity of vegeta-
66 tion. The local names “chul” (arid plain, desert), “adyr” (foot-
67 hills), “tau” (mountains), and “yailau” (alpine zone) are widely
68 used by the people of Uzbekistan and correspond to the zones
69 produced by vertical changes in the landscape.

I.V. Belolipov (✉)

Tashkent State Agrarian University, Tashkent-140,
Tashkent 100140, The Republic of Uzbekistan
e-mai: ivbelolipov@mail.ru

D.E. Zaurov

Department of Plant Biology and Pathology, Rutgers University,
School of Environmental and Biological Sciences,
20 Ag. Extension Way, New Brunswick, NJ 08901, USA

S.W. Eisenman

Department of Landscape Architecture and Horticulture,
Temple University, College of Liberal Arts, School of Environmental
Design, 580 Meetinghouse Rd., Ambler, PA 19002, USA



Fig. 2.1 Uzbekistan with provincial boundaries

70 **The chul zone (arid plain, desert):** The chul consists of
 71 the flat territory of Uzbekistan, which is usually considered
 72 desert. The chul continues up to 500–600 m (~1,640 to
 73 1,970 ft) above sea level and has a dry period of 3–6 months.
 74 The climate of the chul zone is ultra-continental and is char-
 75 acterized by low precipitation of 70–208 mm (~2.75 to 8.2 in.)
 76 per year and humidity levels that drop to as low as 1–2%. The
 77 dry period in the chul zone lasts from May to October.
 78 Summer temperatures can reach 45°C (113°F) while winter
 79 temperatures often drop below –30°C (–22°F).

80 The chul zone occupies most of the Central Asian plain
 81 (Turan) and displays four soil types: salty chul, sandy chul,
 82 gypsum (stony) chul, and clay chul (National Biodiversity
 83 Strategy Project Steering Committee 1998). Portions of the
 84 salty chul ecosystem that have extremely high salt concen-
 85 trations support no plant life. Areas of salty chul with lower
 86 salt content are dominated by *Artemisia halophila* and spe-
 87 cies in the Chenopodiaceae family such as *Halocnemum*
 88 *strobilaceum*, *Halostachys caspica*, *Haloxylon aphyllum*,
 89 *Salicornia herbacea*, *Salsola dendroides*, *Suaeda dendroides*

and *S. microphylla*. Sandy chul is dominated by
Acanthophyllum korolkowi, *Ammodendron conollyi*,
Astragalus villosissima, *Calligonum aphyllum*, *Convolvulus*
hamadae, *Ephedra strobilacea*, *Ferula foetida*, *Salsola*
arbuscula and *S. richteri*. The gypsum chul is located in the
 hills of the southwestern and central Kyzylkum desert.
Artemisia associations predominate in the gypsum chul zone.
 The most common association is *Artemisia diffusa* (less fre-
 quently *A. ferganensis*) with *Convolvulus hamadae* or with
 co-dominance of *Aellenia subaphylla*, *Anabasis eriopoda*,
Anabasis turkestanica and *Salsola arbuscula*. The species
Calligonum junceum and *Reaumuria turkestanica*, and oth-
 ers are commonly found in the gypsum chul and are charac-
 teristic for the area. *Nanophyton erinaceum* is less frequent
 and restricted mainly to the hills.

Where river valleys cut into the chul zone the increased
 humidity in the valleys facilitates the development of special
 mesophytic communities that are locally called “tugai”.
 Common species that occur in these communities are *Alhagi*
persarum, *Apocynum scabrum*, *Asparagus persicus*, *Clematis*

110 *orientalis*, *Elaeagnus orientalis*, *Erianthus purpurascens*,
 111 *Glycyrrhiza glabra*, *Halimodendron halodendron*, *Hippophae*
 112 *ramnoides*, *Karelinia caspia*, *Limonium otolepis*, *Lycium*
 113 *ruthenicum*, *Phragmites communis*, *Populus diversifolia* and
 114 *P. pruinosa*, and *Tamarix* spp.

115 **The adyr zone (lowlands and foothills):** The adyr zone
 116 is a broad belt at an elevation of around 500–1,500 m (1,640–
 117 4,921 ft). This band is found around all the mountains of
 118 Central Asia. It occupies the range between two contrasting
 119 ecological zones: the xerothermic chul (desert) and the meso-
 120 thermic tau (mountain region). The soils of the adyr zone
 121 contain less salt and more humus than the chul soils and are
 122 classified as sierozem (Makhmudov 2001). Bedrock is often
 123 found exposed on the surface.

124 The annual precipitation is between 250 (9.8 in.) and
 125 400 mm (15.7 in.) and rarely reaches 500 mm (19.7 in.). The
 126 mean monthly temperature for July is 25°C (77°F), which is
 127 3–4°C lower than in the chul and 5–6°C higher than in the tau
 128 zone. The dry period lasts from June to September. Due to its
 129 location the adyr zone is exposed to the influence of both the
 130 hot desert along its lower edge, and the cooling effects of the
 131 mountains on its upper edge. This causes the lower section of
 132 the adyr zone to be closer to the environmental conditions of
 133 the chul and the upper section to be similar to the mountain-
 134 ous environment of the tau zone. Because of this gradient the
 135 adyr is divided into subzones: the lower adyr with rolling
 136 relief and the upper adyr with broken relief.

137 Typical species found in the lower adyr area are *Amygdalus*
 138 *spinosissima*, *Artemisia sogdiana*, *Carex pachystylis*, *Mediasia*
 139 *macrophylla*, *Phlomis thapsoides*, *Pistacia vera* and *Psoralea*
 140 *drupacea*. At altitudes of 1,200–1,500 m in the upper adyr
 141 zone, typical species are *Acanthophyllum gypsophiloides*,
 142 *Agropyron trichophorum*, *Astragalus eximius*, *Bunium persi-*
 143 *cum*, *Centaurea squarrosa*, *Cousinia pulchella*, *Onobrychis*
 144 *spp.*, *Phlomis salicifolia* and *P. olgae*, *Potentilla soongarica*,
 145 *Scabiosa songarica* and *Ziziphora pamiroalaica*.

146 **The tau zone (mid-mountain zone):** The tau zone is a
 147 broad belt at an elevation of around 1,500–2,800 m (4,921–
 148 9,186 ft). The dominant soil of the tau zone is of the brown
 149 soil type. Precipitation in this zone exceeds more than 500 mm
 150 (19.7 in.) per year, with a dry period that lasts for 3 months
 151 from July to September. The growing period is in spring,
 152 summer, and autumn with a dormant period in the winter. The
 153 mean monthly temperature in July is 19°C (66°F).

154 In terms of economy, the tau zone is an important area for
 155 growing cereals and leguminous crops, for producing hay,
 156 and for use as pastures. The dominating wild and cultivated
 157 shrub and arboreal species (*Crataegus* spp., *Juglans regia*,
 158 *Malus* spp., *Prunus* spp., etc.) of the area provide the local
 159 population with fuel, building materials, and food. In the tau
 160 zone shrubs can be found in large groups or as individuals.
 161 The common species of shrubs are *Berberis oblonga*, *Cerasus*

tianshanica, *Ephedra equisetina*, *Lonicera microphylla*,
 162 *Rosa kokanica* and *Spiraea hypericifolia*. Some of the woody
 163 species found in the tau zone are gymnosperms such as
 164 *Juniperus semiglobosa*, *J. seravschanica* and *J. turkestanica*,
 165 and broad-leaved deciduous trees such as *Acer turkestan-*
 166 *icum*, *Betula tianschanica*, *Crataegus pontica* and *C. turkes-*
 167 *tanica*, *Juglans regia*, *Malus sieversii*, *Prunus sogdiana*,
 168 *Sorbus persica*, *Ziziphus jujuba* and others.

169 **The yailau zone (high mountain zone):** The yailau zone
 170 is the high-altitude, subalpine to alpine zone and extends
 171 from 2,800 to around 3,400 m (9,186–11,155 ft). This zone
 172 is characterized by environmental conditions that will not
 173 support the development of arboreal and shrub vegetation.
 174 The soil is mainly light brown and of the meadow-steppe
 175 type (Kaurichev 1989). Summer is short and rather warm,
 176 with sharp changes between day and night temperatures.
 177 Summer daytime temperature reaches up to 25°C (77°F), but
 178 can drop to 0°C (32°F) at night. In the winter the temperature
 179 may drop to –40°C (–40°F). Precipitation varies from 400
 180 (15.7 in.) to more than 600 mm (23.6 in.) per year. This zone
 181 has stony taluses, glacial valleys, glacial cirques, and glacial
 182 tongues, and fields with heavy clay soils. There are many
 183 sheer rock formations in the southwestern Tian Shan and the
 184 western Pamiro-Alai.

185 In terms of economy, the yailau region is utilized as the
 186 main summer pasture. While the Karakul sheep graze mainly
 187 in the chul, the Merinos and fat-tailed breeds of sheep
 188 (including the Gissar breed) are pastured mainly in the yailau.
 189 Other agriculture is limited by low temperatures.

190 Tallgrass meadows are an important portion of the vegeta-
 191 tion cover of the yailau. These meadows also contain
 192 *Polygonum bucharicum* and *P. hissaricum*. In western Tian
 193 Shan and the southern Pamiro-Alai there are Apiaceae-rich
 194 meadows with *Ferula tenuisecta* and *Prangos pabularia*.
 195 Meadows containing *Alopecurus*, *Artemisia*, and *Geranium*
 196 spp. are also common in the yailau zone. Shortgrass mead-
 197 ows, also known as alpine meadows, are found in small
 198 patches in the upper yailau. The high-altitude meadows are
 199 comprised of a diversity of grasses and other herbaceous
 200 genera such as *Gentiana*, *Oxytropis*, *Potentilla* and
 201 *Ranunculus*. Meadows of grasses such as *Alopecurus* spp.,
 202 *Festuca* spp., *Poa alpina* and *Phleum alpinum*, and sedges
 203 such as *Carex* and *Kobresia* are also characteristic of the
 204 upper yailau zone.

205 The great extremes of elevation, temperature, precipita-
 206 tion, and soil types found in Uzbekistan provide a wide range
 207 of habitats, which support a great diversity of vegetation.
 208 Due to human activities many of the natural areas of the
 209 planet are being disturbed or destroyed. Conservation of nat-
 210 ural environments and resources are of great importance for
 211 the future of mankind and the conservation of Uzbekistan's
 212 natural resources is no exception.
 213

1 A Short History of Medicinal Plant Use 2 in Central Asia

3 Anvar G. Kurmukov and Anarbek A. Akimaliev

4 Central Asia is a synthesis of many nations and many
5 cultures. There is a long history of using and documenting
6 medicinal plants in this region. Great contributions to the
7 knowledge of medicinal plants were made by the Greeks as
8 early as the seventh century BCE. In the sixth century BCE,
9 Central Asia was part of the Persian Empire founded by
10 Cyrus (Bobokhanova and Bekturgunava 1996). In the first
11 half of the fourth century BCE, Alexander the Great helped
12 expand the Greek empire into the regions of Bactria and
13 Sogdiana (territories that included much of present-day
14 Central Asia) and formed the Greco-Bactrian Kingdom. It is
15 known that Chinese travelers visited Central Asia and sur-
16 rounding regions since the establishment of the Great Silk
17 Road. China has a long history of herbal medicine and
18 undoubtedly had a great influence on the development of
19 Central Asian herbology. The Zoroastrian holy book, the
20 *Avesta*, written over a long period (the ninth century BCE
21 – third century CE), is a valuable source of information about
22 the social structure, medicine, and way of life in the ancient
23 societies of Central Asia, Iran, and Azerbaijan. The book
24 includes information concerning all aspects of life, including
25 natural philosophies and medical views. In the *Avesta*, medi-
26 cine is described as the art of keeping the body in health.
27 There were several kinds of treatments: (1) surgery, (2) treat-
28 ment with herbs, and (3) treatment with words. This source
29 was expanded by other researchers and physicians up to the
30 eighth century and by that time, contained information on
31 1,000 plants. The *Avesta* contains information about useful
32 and unhealthy plants, herbs that were used as sedatives, anes-
33 thetics, narcotics, restoratives, tonics, antiseptics, antidotes,
34 and as other types of remedies. In the fifth and sixth centuries

CE a large Turkic kaganate was developed as result of the
unification of diverse nomadic tribes. In the seventh century
Arabs brought Islam to Central Asia and during this time the
knowledge and science of medicinal plants grew greatly. In
1220 CE the Mongols, led by Genghis Kahn, invaded Central
Asia and there is no doubt that this invasion had an influence
on the culture of the local population.

The Central Asian scientists Abu Rayhan Muhammad ibn
Ahmad Al-Beruni (973–1048) and Abu Ali ibn Sina
(Avicenna; 980–1037) made considerable contributions to
the knowledge of medicinal plants. Both were great scholars,
and besides other sciences, studied pharmacognosy and
pharmacology. The work *Kitab-al-Saidana (Materia Medica)*
was written by Beruni towards the end of his life (1041–
1048). It describes about 750 plants and contains informa-
tion about the botanical characteristics of plants and their
geographical locality. About 400 geographical place names
from where the plants had been brought (Central Asia,
Afghanistan, Iran, Arabia, Azerbaijan, Armenia, and others
areas) are mentioned.

Abu Ali ibn Sina (Avicenna) is famous due to his works
on philosophy and medicine. Being a doctor, he studied bot-
any as well, and often used medicinal plants to treat his
patients. His most important medical work is the *Al-Qanun fi*
al-Tibb (The Canon of Medicine). The second volume of this
work is dedicated to medicinal remedies used during his
time. The book describes more than 800 pharmaceutical sub-
stances of vegetative, animal, and mineral origin. Besides
remedies produced in Central Asia and other countries of the
Near and Middle East, Avicenna described a number of drugs
brought from India, China, Greece, Africa, Mediterranean
islands, and other parts of the world. The book includes the
practices of scientific medicine as well as the traditional folk
medicine of the time. Many medicines (drugs) described by
Avicenna have entered the pharmacopoeia and are still in
use. The fifth volume of *The Canon of Medicine* represents
his pharmacopoeia. It describes how to make and use differ-
ent forms of drugs and complex medicinal formulations. In
the chapter named “Necessity of complex drugs”, Avicenna

A.G. Kurmukov (✉)
The Specialized Center for Cardiology, 4 Murtozoev St.,
Tashkent 100052, The Republic of Uzbekistan
e-mail: kurmukov_anvar@mail.ru

A.A. Akimaliev
Biology and Soil Science Institute of the National Academy
of Science of Kyrgyzstan, 265 Chuy St., Bishkek 720071,
The Kyrgyz Republic

74 recommended making complex drug formulations in order
75 to increase effects of a drug; to prevent side effects of one
76 drug by another drugs composition; to strengthen the effect
77 of the main drug by adding another one (synergy); to increase
78 penetration of one drug into tissues with the help from
79 another drug, or to slow down an effect of a drug by reducing
80 absorbability caused by a second drug and this way elongate
81 the effect of the first drug; and to use drugs for guiding deliv-
82 ery of the main active substance to a point (organ) of action.
83 Carl Linnaeus later named *Avicennia*, a genus of tropical
84 mangrove trees, in honor of Avicenna.

85 In the eleventh to twelfth centuries, Ismail al-Jurjani
86 (Ismail ibn Muhammad al-Husayn Jurjani), wrote an ency-
87 clopedic work on medicine called *Zakhirah-i Khvarazm'Shahi*
88 (*Treasure of Khorezm Shah*). Later, in the fourteenth century,
89 Mansur ibn Ilyas (Mansur ibn Muhammad ibn Ahmad ibn
90 Yusuf ibn Ilyas) published his work *Kifayah-i Mansuri*
91 (*Mansur's Sufficient Book*), which was also known as
92 *Kifayah-i Mujahidiyah* (*The Sufficient [Book] for Mujahid*).
93 Besides being summaries of medical theory and practice,
94 these works gave basic information about plant-based
95 medicine. In the eighteenth century, Muhammad Husayn
96 (Muhammad Husayn ibn Muhammad Hadi al-Aqili al-Alavi
97 al-Khurasani al-Shirazi, also known as or Hakim Muhammad
98 Hadikhan) described the therapeutic qualities of more than
99 2,000 plants, preparations of animal origin and minerals in
100 his works *Majma al-javami va-zakha'ir al-Tarakib* (*The*
101 *Assemblage of Generalities and Treasures of Compounds*)
102 and *Makhzan-al-Adviyah* (*The Storehouse of Medicaments*).
103 These works were largely based on the earlier writings of his
104 great uncle Alavi Khan and documented centuries of past
105 achievements in the field of folk medicine, the practices of
106 previous physicians, and his personal research.

107 In the past, the use of plants for medicine was not rigor-
108 ously based in science. Modern scientific techniques have
109 been used to prove the effectiveness of many plant remedies
110 used in folk medicine and prescribed by ancient physicians.
111 For instance, *Rauwolfia serpentina* has been used in Indian
112 medicine for about 2,000 years, while Europeans discovered
113 the value of this plant only in the middle of twentieth century
114 (Balick and Cox 1996; Gupta 2002). Since ancient times,
115 Africans have used *Strophanthus* seeds to make arrow poison
116 and as a cardiac remedy, but only at the end of nineteenth
117 century did *Strophanthus* enter the European pharmacopoeia
118 (Norn and Kruse 2004). To the present day, these plants are
119 considered irreplaceable cardiac remedies used to treat car-
120 dio-vascular diseases.

121 Many of the specific activities of plant remedies described
122 by Avicenna have been confirmed by modern research con-
123 ducted at the Institute of Chemistry of Vegetative Substances
124 (ICVS) of the Academy of Sciences of the Republic of
125 Uzbekistan. For example, according to Avicenna, the plant
126 *Haplophyllum perforatum* has anti-inflammatory and seda-

127 tive effects. The alkaloids perforine, evoksine, skimmianine,
128 and others have been isolated from this plant. It has been
129 found that at medium doses these alkaloids have sedative,
130 and in higher doses sleep-inducing effects. Some of these
131 alkaloids also have an anti-inflammatory action (Sadritdinov
132 and Kurmukov 1980). Another example is *Khiltit* (the gum
133 from *Ferula foetida*), which Avicenna noted could be used as
134 a treatment for malignant and fatal tumors by cutting the
135 tumor open and applying the gum. According to Avicenna
136 this gum also strengthens the libido and stimulates menstrua-
137 tion. The esters of sesquiterpene alcohols, ferutinine, ferutin
138 and others have been isolated from this species. A prepara-
139 tion from this plant, *Panoferol*, and also the individual com-
140 pounds ferutin and ferutinine, have pronounced estrogenic
141 action. *Panoferol* strengthens the libido and increases impreg-
142 nation in sheep, pigs, and cattle. A mixture of ferutin and
143 ferutinine (under the name *Tefestrol*) has been introduced to
144 obstetric-gynecologic practice as an estrogenic preparation
145 (Kurmukov and Akhmedkhodzhaeva 1994). There are many
146 more examples of corresponding effects of various plants
147 described by Avicenna and recent data gathered by modern
148 pharmacologists (Sadritdinov and Kurmukov 1980).

149 In the twentieth century, research on Central Asian
150 medicinal plants was especially productive, particularly in
151 Uzbekistan. In 1943, the Laboratory of Chemistry of
152 Alkaloids (headed by Professor S. Yu. Yunusov) was founded
153 at the Institute of Chemistry in the Uzbek branch of the
154 Academy of Sciences of the USSR. Due to the successful
155 activity of the laboratory, Yunusov created the Institute of
156 Chemistry of Plant Substances at the Academy of Sciences
157 of Uzbek Soviet Socialist Republic in 1956. The Institute
158 had laboratories devoted to distinct chemical groups includ-
159 ing alkaloids, glycosides, fats, proteins, terpenes and acids,
160 lignin, coumarins and phosphorous-containing organic com-
161 pounds, as well as botany, laboratories of pharmacology and
162 toxicology, phytotoxicology, experimental technology and
163 physical and quantitative analysis and others.

164 The Institute's scientific directions consisted of a complex
165 of investigations into plant substances. All plant parts col-
166 lected during different growth periods and from different
167 regions, were investigated. Applied laboratories had the task
168 of studying the pharmacological activity of compounds; to
169 determine the possibility of introduction into medical prac-
170 tice; to study the natural habitat of the medicinal plants; to
171 organize long-term plant collecting; to maintain the safety of
172 natural populations; and to organize the development of
173 medicinal formulations and their production. The overall
174 goal of the institute was to create medicinal products follow-
175 ing a research pipeline, which included collecting of data on
176 pharmacognosy, isolation of individual compounds, study of
177 their pharmacological activity and creation of medicinal
178 preparations up to the point of introduction into medical
179 practice.

180 Researchers in the Laboratory of Alkaloid Chemistry iso- 233
181 lated and studied many alkaloids, including a number of new 234
182 alkaloids belonging to various chemical groups. Research on 235
183 alkaloid chemistry was summarized in the monograph 236
184 *Alkaloids* by Yunusov (1974, 1981). These newly isolated 237
185 alkaloids were also studied by the pharmacology and toxicol- 238
186 ogy labs. The Glycoside Chemistry Laboratory (headed by 239
187 Professor N.K. Abubakirov) studied cardiac glycosides. This 240
188 laboratory made significant contributions to the knowledge 241
189 of the chemistry of triterpene glycosides. Among the studied 242
190 compounds, glycosides with immunomodulatory, gonado- 243
191 tropic, and hypolipidemic activities were identified. Studies 244
192 of *Astragalus* led to the isolation of methyl-steroids of the 245
193 cycloartan series. Many species of *Allium* (onions) native to 246
194 Central Asia were investigated and as a result more than 30 247
195 new compounds were isolated. One of the most important 248
196 scientific directions of the laboratory during the past years 249
197 has been investigations of phytoecdysteroids. This labora- 250
198 tory identified the structure of 25 of the 95 phytoecdysteroids 251
199 described in the literature by 1980. Studies of *Amorpha fru-* 252
200 *ticosa* led to the discovery of a new class of plant glycosides 253
201 containing rotenone derivatives as the aglycone. 254

202 The Laboratory of Lactones, Coumarins, and Terpenoids 255
203 (headed by Prof. G.P. Sidiyakin) studied various plants for lac- 256
204 tone content, particularly for the lactones leucomisine and 257
205 austricine, which were isolated from *Artemisia leucodes*. Both 258
206 lactones possess pronounced anti-inflammatory action. As a 259
207 compound possessing pronounced angio-protective, hypolipi- 260
208 demic, hypo-cholesterolemic, and anti-inflammatory actions, 261
209 leucomisine has passed medical tests and has been introduced 262
210 into medical practice under the preparation name *Oligvon*. 263
211 Since 1970, systematic studies of chemical compounds found 264
212 in various species of the genus *Ferula*, which grows in the ter- 265
213 ritory of Uzbekistan and adjacent republics, have been con- 266
214 ducted. As a result, more than 50 species of *Ferula* have been 267
215 investigated, from which more than 250 new terpenoids, cou- 268
216 marins, and esters have been isolated and their chemical struc- 269
217 tures determined. Natural esters of mono- and sesquiterpene 270
218 alcohols with aromatic acids were discovered for the first time 271
219 in this lab (Kurmukov and Akhmedkhodzhaeva 1994). 272

220 In 1957, the Pharmacology Laboratory was founded at the 273
221 Institute (headed by associate prof. I.K. Kamilov). The initial 274
222 activities of the laboratory were related to alkaloids. The 275
223 findings were mainly summarized in the books *Pharmacology* 276
224 *of Plant Alkaloids and Their use in Medicine* (Sadritdinov 277
225 and Kurmukov 1980) and *Alkaloids and Herbal Preparations* 278
226 *for Hypertensive Treatment* (Kurmukov and Zakirov 1992). 279
227 During these years preparations containing the alkaloids vin- 280
228 canine (a preparation of vincanine hydrochloride, a strych- 281
229 nine-like analeptic), vincamine (a preparation of vincametrine, 282
230 a stimulator of uterine smooth muscles), ervinine (a CNS 283
231 analeptic with primary stimulating effect on the respiratory 284
232 center) and others were introduced into medical practice. 285

Later the alkaloid lappaconitine, in the preparation *Allapenin* 233
developed by S.Yu. Yunusov and F.N. Dzhakhangirov and 234
isolated from *Aconitum soongaricum*, was introduced into 235
medical practice and was widely used as an antiarrhythmic 236
drug. The same authors developed the compound prepara- 237
tion *Aklezin* from similar alkaloids and which was also used 238
as an antiarrhythmic drug. Pharmacological investigations of 239
alkaloids from *Peganum harmala* resulted in the introduction 240
of an anticholinesterase preparation, *Desoxypeganine*, into 241
medical practice (Tulyaganov et al. 1986). The rotenoid gly- 242
coside amorphine was isolated from the plant *Amorpha fru-* 243
244 *ticosa* in the laboratory of chemistry of glycosides. 245
Pharmacological studies revealed the hypolipidemic, hypoc- 246
247 holesteremic, and angio-protecting actions of the preparation 248
(Aizikov et al. 1984; Kurmukov et al. 1982, 1984a, b, 1986). 249
After completion of clinical tests, the preparation *Glirofam* 250
(containing amorphine), was introduced as a prophylaxis and 251
treatment of atherosclerosis. 252

253 A series of studies on the pharmacology of phytoecdys- 254
255 teroids (ecdysterone, turkesterone, ciasterone, viticosterone) 256
isolated from *Rhaponticum carthamoides*, *Ajuga turkestan-* 257
258 *ica*, and various species of *Serratula* have been conducted. 259
These compounds possess tonic and anabolic actions, and 260
unlike the steranebols (nerobol) do not have androgenic 261
effects. They increase exercise performance, accelerate reha- 262
263 bilitation of lost physical capabilities, and increase an organ- 264
265 isms' ability to adapt to extreme environmental conditions 266
(Kurmukov and Syrov 1976; Syrov and Kurmukov 1975a, b, 267
268 c, 1976a, b, c, d, 1977, 1980; Kurmukov et al. 1980, 1982; 269
270 Syrov 1984, 1994; Saatov et al. 1994). The preparation 271
272 *Ecdisten* was developed from ecdysterone, and is used in 273
274 medical practice as a restorative, to improve memory, as a 275
276 prophylaxis for and treatment of myocardial infarction, and 277
278 especially for rehabilitation after cardiac infarction. 279

280 The Institute of Bioorganic Chemistry of the Academy 281
282 of Sciences of Uzbekistan was founded by academician 283
284 A.S. Sadikov. Besides various chemical laboratories, there is 285
also a Laboratory of Pharmacology at this Institute (headed by 286
Prof. S.Kh. Nasirov). In addition to natural compounds, the insti- 287
tute has studied medicinal plants, particularly alkaloids from 288
the species *Anabasis aphylla* and *A. jaxartica*, *Ammodendron* 289
290 *argenteum*, *Calligonum minimum*, *Colchicum kesselringii*, 291
292 *Merendera raddeana* and others. Other plant compounds, 293
294 including proanthocyanidins from the seeds of grapes, are stud- 295
296 ied at the institute as well (Pirniyazov et al. 2003). 297

298 Medicinal plants and their compounds are studied in the 299
300 Pharmaceutical Institute of the Ministry of Health of The 301
302 Rep. of Uzbekistan, especially in the subdepartments of 303
304 Pharmacognosy (Prof. Kh.Kh. Khalmatov and his students), 305
306 Pharmacology (Prof. Kh.U. Aliev) and Botany. Prof. 307
308 Khalmatov and his associates published a series of books 309
310 about the medicinal plants of Central Asia and Uzbekistan, 311
312 and about their use in medicine. Similar studies are conducted 313

286 in the subdepartments of the medical institutes and related
287 laboratories of the scientific research institutes. As a result of
288 the research on plant substances in the Laboratory of
289 Experimental Cardiology of the Scientific Research Institute
290 of Cardiology, now known as the Republican Specialized
291 Center for Cardiology (headed by Prof. R.D. Kurbanov), the
292 preparations *Oligvon*, *Glirofam*, *Ecdisten*, *Kavergal* and oth-
293 ers were introduced into medical practice.

294 In Kyrgyzstan scientific studies of medicinal plants began
295 in the pharmacology laboratory of the Institute of Regional
296 Medicine of the Kyrgyzstan National Academy of Science in
297 1954. Later the name of the lab was changed to laboratory of
298 pharmacognosy. The laboratory developed a tincture and the
299 preparation *Foetidin* from the aboveground parts of
300 *Thalictrum foetidum*, which was used to treat the first and
301 second stages of hypertension. Later, Dr. P.K. Alimbaeva
302 studied all species of the genus *Lagochilus* found in
303 Kyrgyzstan. These studies showed that *Lagochilus platya-*
304 *canthus* and *L. platycalyx* had the same effects on the cardio-
305 vascular system and blood coagulation as the species *L.*
306 *inebrians*. Dr. B.N. Aronova conducted pharmacognostical
307 studies of *Betonica foliosa*. As a result, a liquid extract of the
308 aboveground parts of this species was introduced into medi-
309 cal practice as a treatment for uterine diseases.

310 The department of biopharmacology (headed by
311 Academician Altimishev) was organized in 1969. This
312 department included the laboratory of pharmacology and
313 toxicology (led by Academician Altimishev), lab of resources
314 (led by Dr. A.A. Akimaliev), and the lab of pharmacognosy
315 (led by Dr. P.K. Alimbaeva). The main scientific goals of the
316 department were pharmacotoxicology studies and justification
317 for the use of natural and synthesized physiologically active
318 compounds. The preparation *Licorin* was introduced into
319 medical practice to treat bronchial and lung diseases. The
320 Ministry of Public Health Committee of the USSR
321 (Pharmacology committee) permitted the use of the linament
322 *Karagai* and *Hippophae rhamnoides* oils, in the preparation
323 *Gippol*, which were developed by scientists from the depart-
324 ment. The medicinal balsams (alcoholic plant extracts),
325 including *Arashan*, *Uccurisky*, *Kobuctan* and *Sibir*, were
326 developed and commercialized. *Arashan* was awarded a seal
327 of quality by the USSR and a gold medal at an international
328 exhibition in Leipzig in 1977.

With support from the Soviet Space Program, Drs. O.I. 329
Gorelkina, E.P. Zotov and S.N. Khabibrakhmanov of the 330
department of biopharmacology, developed and introduced 331
special adaptogens such as *Gipkos*, *Giprex*, *Gipomin*, *Daugil*, 332
etc. for use in the space program and in sports medicine. The 333
preparation *Dipsacozide*, prepared from *Dipsacus azureus* 334
roots, was developed and studied. Experiments showed that 335
this preparation increased organisms' resistance to hypoxia 336
and had hepatoprotective and antiatherosclerotic activities, 337
which were proven after clinical studies. A non-alcoholic 338
drink called *Omur*, based on the preparation *Dipsacozide*, 339
was developed and recommended as a prophylactic for ath- 340
erosclerosis. Additionally, the glycoside fraction, *Zongorozid*, 341
was isolated from the roots of *Scabiosa songorica*. In exper- 342
iments with animals the fraction significantly decreased 343
arterial blood pressure and had sedative effects (Alimbaeva 344
et al. 1986). 345

In recent years medicinal plants have been studied at the 346
laboratory of biopharmacology (led by Dr. A.A. Akimaliev) 347
at the Soil Biology Institute of the Kyrgyzstan National 348
Academy of Science. Based on edible and medicinal plants, 349
this lab developed the dietary supplement *Chabal*, which is 350
recommended to people who have been exposed to radiation 351
(such as atomic power station workers), as well as recom- 352
mended to weak patients and athletes as a general tonic. 353
Chabal has been approved by the Pharmacology and 354
Pharmacopeia Committee of the Ministry of Public Health 355
of the Kyrgyz Republic. 356

Many therapeutic syrups have been developed using medic- 357
inal plants from the flora of Kyrgyzstan. The syrup *Beykut* is 358
used as a sedative and *Glitimal* is used as an expectorant and 359
anti-inflammatory. The syrup *Akan* is used to prevent the 360
development of stones in the urinary tract and bile pathways 361
and is also recommended as a treatment for cholecystitis 362
and hepatitis. All of these syrups were approved by the 363
Pharmacology and Pharmacopeia Committee of the Ministry 364
of Public Health of the Kyrgyz Republic. At the Medical 365
Academy of Science, under the leadership of the Corresponding 366
Academician of the National Academy of Science of the 367
Kyrgyz Republic Professor A.Z. Zurdinov, a preparation 368
Immunaz, with immunomodulatory properties, was developed 369
from the leaves of *Padus grayana* and introduced into 370
medical practice. 371

Plants contain organic as well as inorganic substances that can provide therapeutic effects. Different plants may possess a wide spectrum of effects due to the presence of various groups of chemical compounds and various microelements. A preparation obtained from one plant can simultaneously be an analgesic, sedative, cardiotoxic, anti-inflammatory, and expectorant. Well-formed herbal preparations can be used protractedly when necessary, without injury to a patient, which is very important when treating chronic ailments. Medicinal plants are widely used as prophylaxis for, and treatment of, many diseases, including gastritis, stomach and duodenal ulcers, cholecystitis, colitis, enteritis, pyelonephritis, cystitis, atherosclerosis, cardiac insufficiency, and arrhythmia. They are also used for treatment of hypertensive and hypotensive neurocirculatory dystonia, neurosis and asthenia, menopausal disorders, and also to boost the body's immune system during times of disease, for rehabilitation of post-infarction conditions, as a tonic, and to increase adaptive capabilities of the organism.

Rational phytotherapy can promote recovery from dysbolism, normalize nervous system function, contribute to stabilization of blood pressure, improve coronary blood circulation and cerebral blood supply, help reduce insomnia and increase capacity for work. Herbal preparations promote excretion of toxic substances, help individuals to regain normal strength, increase energy metabolism and stop further disease progress during atherosclerosis and hypertension.

It is known that the effectiveness of medicinal plants and their pharmacotherapeutic action is due to their complex diversity of chemical compounds. Among these compounds are alkaloids, glycosides, lactones, tannins, proanthocyanidins, pigments, ecdysones, saponins and others.

Alkaloids – Alkaloids are nitrogen-containing organic bases. They are characterized by high pharmacological activity. In small doses, alkaloids represent valuable pharmaceuti-

cal substances such as lappaconitine, vincamine, reserpine, morphine, quinidine, strychnine, atropine, caffeine, ephedrine, nicotine and others. They form the main active ingredients of many medical products used for treatment of various diseases. Decoctions, infusions, extracts and others are made of alkaloid-containing plants.

Glycosides – Glycosides are organic compounds of vegetative origin, composed of a sugar component (glycoside, glycone) and a non-sugar component (aglycone, genin). The aglycone forms the main physiologically active part. Depending on their chemical nature and structure, glycosides are divided into cyanogenic glycosides (aglycones contain prussic acid), cardiac glycosides (aglycones are cardinolides and bufadienolides), saponins (aglycones are triterpene and steroid compounds), anthraglycosides (aglycones are derivatives of anthracene), phenolics (aglycones are coumarins, flavonoids, and others), and glycoalkaloids (aglycones are nitrogen-containing steroid compounds). Cardiac glycosides are used in medicine to treat cardiac disorders. They are toxic and have to be used under the supervision of a physician.

Saponins – Saponins are glycosides that make suds when shaken in water. The name comes from the Latin word “Sapo” meaning soap. Saponins are used as expectorants, diuretics, hypotensives and hypocholesterolemics. Saponins from *Aralia mandschurica*, *Echinopanax elatus*, *Eleutherococcus* spp., and *Panax* spp. have stimulating effects. Saponins cause hemolysis after intravenous introduction. Because of this, they are only introduced orally.

Anthraglycosides – Substances which belong to anthraglycosides look like red-orange crystals. Plant extracts containing anthraglycosides usually have a blood-red color. These compounds have purgative and choleric actions.

Phenol compounds – Simple phenols, coumarins, chromones, lignan, tropolones, flavonoids and their glycosides, tannins, proanthocyanidins and others are in this group. This group of substances has the most diverse pharmacological activity. Among them there are substances that have antihypoxic, antioxidant, choleric, cardio-, angio-, and hepatoprotecting and hemostatic actions.

A.G. Kurmukov (✉)
The Specialized Center for Cardiology,
4 Murtozoev St., Tashkent 100052, The Republic of Uzbekistan
e-mail: kurmukov_anvar@mail.ru

76 **Proanthocyanidins** – These are polyphenol compounds,
77 which possess pronounced antihypoxic, antioxidant and anti-
78 inflammatory actions. They have vitamin-P activity.

79 **Flavones and flavonoids** – This group includes heterocy-
80 clic compounds, uneasily dissolved in water. Flavones and
81 their derivatives have a yellow color, due to which they
82 obtained their name (flavum=yellow). These compounds
83 (rutin, quercetin, hesperidin, citrin and others) have the abil-
84 ity to decrease the permeability of vascular walls and fragil-
85 ity of capillary walls, have antispasmodic actions used for
86 spasms of vessels and smooth-muscle organs, and are used to
87 treat stomach and duodenal ulcers, and hepatitis.

88 **Coumarins and furocoumarins** – These compounds
89 increase human and animal sensitivity to ultraviolet light and
90 are used to treat vitiligo. Some have phyto-estrogenic action.
91 When eaten by sheep and other animals, plants containing
92 coumarins and furocoumarins have contraceptive action.
93 Ingestion can cause fetal death in early pregnancy as well.

94 **Tannins** – Tannins promote inhibition of pathogenic
95 microbial growth and reduce reproduction of viruses and
96 bacteria. They also have astringent, tanning and hemostatic
97 actions, and increase stability of capillary walls.

98 **Organic acids** – These acids are contained in plants in
99 free form as well as in the form of salts and esters. Among
100 them there are malic, citric, succinic, tartaric, oxalic, formic,
101 acetic and other acids. They participate actively in metabo-
102 lism, strengthen activity of salivary glands, and increase bile
103 excretion and gastric juices. Organic acids are contained in
104 lemons, apples, cranberries, currants, rosehips, sea-buck-
105 thorn berries, sorrel leaves, asparagus, greater celandine and
106 other plants. Valeric and isovaleric acids (valerian, milfoil
107 and others), and benzoic acid (in red whortleberry) have
108 medicinal effects.

109 **Esters of mono – and sesquiterpene alcohols with aro-**
110 **matic acids** – These have estrogenic, hypolipidemic, and
111 hypo-triglyceridemic activity and moderately increase blood
112 pressure.

Fatty oils and fat-like substances – Fats and oils are esters 113
of glycerin and higher fatty acids. In pure form, oils (castor, sea- 114
buckthorn and others) are used as remedies or as solvents for 115
pharmaceutical substances. Fatty oils are used in medicine to 116
make ointments, liniments and emollients, and for skin care and 117
therapeutic massage. Some oils have therapeutic action. For 118
example castor oil is used as a purgative and sea-buckthorn is 119
used internally to treat stomach and duodenal ulcers and exter- 120
nally for skin burns. Plant waxes, sterols and other substances 121
are fat-like substances. Some of them are used in medicinal 122
preparations. Unsaturated fatty acids (oleic, linoleic, linolenic, 123
palmitic and other acids) prevent development of atherosclero- 124
sis. They are contained in plant seeds (almond, sunflower, flax 125
and others), and fruits (olives and sea-buckthorn). 126

Mucilage – Consists of nitrogen-free compounds of vari- 127
ous chemical compositions, mainly polysaccharides. They 128
have coating and emollient actions, and can be found in 129
Althaea roots and flax seeds. 130

Gums – Gums are polysaccharides. They are hardened 131
fluids released out of damaged tree and shrub bark. They are 132
used as emulsifying agents and also as adhesives. Apricot, 133
cherry, plum and others are sources for gum. 134

Pectins, starch, and various sugars – Like mucilage and 135
gums, these are related to carbohydrate groups and are used 136
as additives in drug formulations. 137

All of the above mentioned groups of chemical com- 138
pounds are the main active principles of the medicinal plants 139
that are used today. However, only a very small percentage of 140
the great diversity of plant-based compounds that exist in 141
nature has been explored. Through the scientific process, 142
new compounds having other effects are currently being 143
revealed and will continue to be revealed in the future. 144

Milligram % (mg%) – A unit used to describe concentra- 145
tion. Milligrams of a specific substance contained in 100 ml 146
of a solution or in 100 g of the analyzed material. This unit 147
of measure is often used to describe vitamin content in plants 148
and foods. 149

The Medicinal Plants of Uzbekistan and Kyrgyzstan

1
2

5

[AU1]

David E. Zaurov, Igor V. Belolipov, Anvar G. Kurmukov,
Ishembay S. Sodombekov, Anarbek A. Akimaliev, and
Sasha W. Eisenman

3
4
5

Uncorrected Proof

D.E. Zaurov
Department of Plant Biology and Pathology, Rutgers University,
School of Environmental and Biological Sciences, 20 Ag.
Extension Way, New Brunswick, NJ 08901, USA

I.V. Belolipov
Tashkent State Agrarian University, Tashkent-140, Tashkent 100140,
The Republic of Uzbekistan

A.G. Kurmukov
The Specialized Center for Cardiology, 4 Murtozoev St., Tashkent
100052, The Republic of Uzbekistan

I.S. Sodombekov
Kyrgyz Botanical Garden of the National Academy of Kyrgyzstan,
1A Akhunbabaev St. Bishkek 720064, The Kyrgyz Republic

A.A. Akimaliev
Biology and Soil Science Institute, National Academy of Science of
Kyrgyzstan, 265 Chuy St., Bishkek 720071, The Kyrgyz Republic

S.W. Eisenman
Department of Landscape Architecture and Horticulture, Temple
University, College of Liberal Arts, School of Environmental Design,
580 Meetinghouse Rd., Ambler, PA 19002, USA

6 **Achillea asiatica** Serg. – Asteraceae7 **Synonyms:** *Achillea millefolium* var. *manshurica* Kitam., *Achillea setacea* ssp. *asiatica* (Serg.) Worosch.8 **English name:** Chinese yarrow, Mongolian yarrow9 **Russian name:** Тысячелистник азиатский (Tysyachelistnik aziatskiy)10 **Uzbek name:** Unknown11 **Kyrgyz name:** Азия каз тандайы (Aziya kaz tandayy)12 **Description:** Herbaceous perennial with thin, branched rhizomes. Stems few or solitary, usually 25–50 cm tall, grayish with
13 long, entangled, white hairs, often with short leafy branches in mid and upper leaf axils. Leaves bipinnatisect, usually
14 oblong, green or grayish-green, more or less densely hairy; leaves of sterile shoots up to 25 cm long, long-petiolate; lower
15 stem leaves 7–20 cm long, petiolate to subsessile; upper leaves sessile, usually 1–6 cm long. Inflorescences capitula
16 arranged in loose, convex corymbs of unequal heights. Involucre cup-shaped; involucre bracts oviform, pale yellowish-
17 green. Ray flower ligules pink, very rarely white. Fruits oblong, wedge-shaped achenes, truncated at the apex.18 **Other distinguishing features:** Bases of mid-stem leaves partially clasping to auriculate. Ray flower ligules usually 1–3 mm
19 long and 1.5–2.5 mm wide.20 **Phenology:** Flowers in August and fruits in August and September.21 **Reproduction:** By seeds and rhizomes.22 **Distribution:** Ysyk-Kol, Osh, Jalal-Abad, and Chuy Provinces of Kyrgyzstan; not found in the flora of Uzbekistan.23 **Habitat:** Found in forests, steppes, and abandoned fields.24 **Population status:** Common, forming dense groups.25 **Traditional use:** The aboveground parts are used as a hemostatic for bloody noses, bleeding gums, small wounds, abrasions,
26 scratches, lung and uterine hemorrhages, and hemorrhoidal hemorrhages. It is used to treat inflammation, metropathy, and
27 for gastrointestinal diseases, such as colitis and ulcers. It is also recommended for treating inflammation of the urinary
28 tract (Plant Resources of the USSR 1993).29 **Documented effects:** This species is used in the same manner as *Achillea millefolium*, and is anti-inflammatory, hemostatic,
30 and antibacterial (Tolmachev 1976).31 **Phytochemistry:** The plant contains alkaloids, flavonoids, sesquiterpene lactones, essential oils, vitamins C and K, resin,
32 carotene, phytoncides, and bitter and astringent substances (Plant Resources of the USSR 1993; Glasl et al. 2001).

Achillea filipendulina Lam. – Asteraceae	33
Synonyms: <i>Achillea eupatorium</i> M. Bieb.	34
English name: Fern-leaf yarrow	35
Russian name: Тысячелистник таволголистный (Tusyachelistnik tavgolistnyy)	36
Uzbek name: Dastarbosh	37
Kyrgyz name: Табылгы жалбырактуу каз тандай (Tabylgy zhalbyraktuu kaz tanday)	38
Description: Perennial herb. Stems erect, up to 60–80 cm tall, thick, striated, densely hairy, densely-leafy. Leaves pubescent, punctate glandular; basal leaves petiolate, oblanceolate, 10–20 cm long and 3–7 cm wide, pinnatipartite with acute segments; upper leaves pinnatifid with large, incised-dentate segments, sessile. Inflorescences capitula gathered into thick, unequally high, terminal corymbs. Ray flowers 1–4, yellow, trilobate. Disc flowers yellow with flattened corolla tube. Fruits oblong, wedge-shaped achenes, 2–2.25 cm long, grayish-black.	39 40 41 42 43
Other distinguishing features: Flowers have a specific pungent smell.	44
Phenology: Flowers in June-beginning of September, fruits in the end of August-September.	45
Reproduction: Reproduces abundantly by seeds and vegetatively by rhizomes.	46
Distribution: Widespread in Kyrgyzstan; Tashkent, Samarqand, Andijon, Farg'ona and Surxondaryo provinces of Uzbekistan.	47 48
Habitat: The adyr and tau zones. On stony, shallow-soiled, slopes with rocky debris, in mountain fissures, in valleys along rivers and brooks, in agricultural zones, and rarely along the banks of small irrigation canals.	49 50
Population status: Common, often found in large populations.	51
Traditional use: A decoction of the herb is used to treat gastric diseases, hemorrhoids, and as an abortifacient (Khalmatov 1964; Sadyrbekov et al. 2006a).	52 53
Documented effects: An extract of the inflorescences has anti-inflammatory activity and strongly inhibited expression of genes associated with inflammation processes (Dey et al. 2008).	54 55
Phytochemistry: The herb contains 0.07–0.26 % essential oil, alkaloid traces, asparagine, amino acids and nitrogen-containing substances. Plants growing in Uzbekistan have high variation in the amount of essential oils, which can vary from 0.04 % to 0.5 %. Around 3 % aldehydes and ketones and 0.5 % phenols are found in the oil composition. Flowering plants from Burchmulla village (Tashkent province, Uzbekistan) contained 0.2–0.27 % essential oil, which contained 10 % octylene, ~5 % pinene, 8 % camphene, 0.35 % C ₁₀ H ₁₈ O alcohol, about 30 % borneol and formic, acetic and caprylic acid (Khakimov and Tsukervanik 1948; Khalmatov 1964). Essential oil extracted from plants growing in the Botanical Garden of the Institute of Phytochemistry, Karaganda, Kazakstan, consisted mainly of santolina alcohol (29 %), 1,8-cineol (19.1 %) and borneol (27.8 %; Sadyrbekov et al. 2006a). The sesquiterpene lactone leucomisine was isolated from the aboveground parts (Konovalov and Nesterova 2003).	56 57 58 59 60 61 62 63 64

65 ***Achillea setacea* Waldst. & Kit. – Asteraceae**66 **Synonyms:** None67 **English name:** Unknown68 **Russian name:** Тысячелистник шетинистый (Tusyachelistnik shchetinisty)69 **Uzbek name:** Unknown70 **Kyrgyz name:** Катуу туктуу каз тандай (Katuu tuktuu kaz tanday)

71 **Description:** Herbaceous perennial. Stems up to 80 cm tall, whitish hairy. Leaves linear-lanceolate, 3–10 cm long, up to
72 2 cm wide, bi- or tripinnatisect, lobes linear-lanceolate; basal and lower stem leaves petiolate; upper leaves sessile.
73 Inflorescences capitula, densely arranged in convex, compound corymbs; involucre oblong-cylindrical; involucral bracts
74 greenish-yellow. Ray flowers 4–5, white, slightly 3-lobed; disc flowers 10–20, yellow, 5-lobed. Fruits oblong achenes,
75 1.8–2 mm long, light brown.

76 **Other distinguishing features:** Capitula 2.5–3 mm across, with peduncles ca. 3 mm long.77 **Phenology:** Flowers in April–June and fruits in July–August.78 **Reproduction:** By seeds and rhizomes.79 **Distribution:** Kungay Ala-Too and Terskey Ala-Too, Chuy valley, Kyrgyz Ala-Too and Alai mountain ranges of Kyrgyzstan;
80 not found in the flora of Uzbekistan.81 **Habitat:** Steppes, meadow-steppes, meadows, among shrubs, forests edges, in abandoned fields, and near roads.82 **Population status:** Common.83 **Traditional use:** Used in the same way as *Achillea millefolium* and *A. asiatica*. A decoction is used to treat internal and
84 external bleeding and hemorrhoids (Plant Resources of the USSR 1993; Alimbaeva and Shambetov 1988).85 **Documented effects:** The essential oil had antimicrobial effects against *Clostridium perfringens*, *Acinetobacter woffii*, and
86 *Candida albicans* (Unlu et al. 2002). Sesquiterpenes isolated from this species exhibited anti-inflammatory activity in the
87 croton oil ear test (Zitterl-Eglseer et al. 1991).88 **Phytochemistry:** This plant contains essential oil, alkaloids, glycosides, tannins, resins, organic acids, vitamins C and K
89 (Plant Resources of the USSR 1993). The aboveground parts contain sesquiterpenes (Zitterl-Eglseer et al. 1991). The
90 essential oil, isolated from air-dried aerial parts, contained over 51 constituents with eucalyptol (1,8-cineole) being the
91 major component (Unlu et al. 2002).

Aconitum karakolicum Rapaics. – Ranunculaceae	92
Synonyms: <i>Aconitum napellus</i> var. <i>turkestanicum</i> B. Fedtsch., <i>Aconitum soongaricum</i> Stapf. (some authors recognize this as a separate species), <i>Aconitum winkleri</i> Rapaics.	93 94
English name: Unknown	95
Russian name: Аконит каракольский (Akonit karakol'skiy)	96
Uzbek name: Karakool parpisi	97
Kyrgyz name: Исykkол уу коргошуну (Isykkol uu korgoshchunu)	98
Description: Herbaceous perennial with conical tuber-like roots. Stems up to 2 m tall, branched. Leaves appressed to stem, short-petiolate; blade circular, up to 10 cm long and 15 cm wide, palmatisect with 5 segments divided to the base; each segment pinnatifid with 2–3 linear lobes, lobes 1.5–3 mm wide. Inflorescence an dense apical raceme; pedicels with two bracteoles. Flowers irregular, with 5 petaloid sepals, dark-violet. Upper sepal hood-shaped, semispherical, with a small beak. Petals 2, each with a spur. Fruit a follicetum with 3–5 glabrous follicles.	99 100 101 102 103
Other distinguishing features: The roots form horizontal, chain-like rows. Distinguished from <i>Aconitum soongaricum</i> by having narrower leaf lobes and appressed pubescence on the inflorescence rachis and pedicels.	104 105
Phenology: Flowers in July-September and fruits in August-October.	106
Reproduction: By seeds.	107
Distribution: Ysyk-Kol province of Kyrgyzstan; not found in the flora of Uzbekistan.	108
Habitat: In meadows with diverse grass species and in spruce forests.	109
Population status: Common, forming dense groups.	110
Traditional use: In Kyrgyz folk medicine, an infusion of the tubers in fermented horse milk or water and ground tubers added to meat broth, are used to treat tuberculosis, radiculitis, and headaches. Tubers are also used to treat different types of cancer (Khalmatov et al. 1984).	111 112 113
Documented effects: An alcoholic tincture of the roots is applied externally to treat radiculitis, neuralgia, rheumatism, and as an analgesic. This tincture is a component of the preparation <i>Akofit</i> . An infusion of the tubers and the aboveground parts is used as a component of the preparation <i>Anginol</i> , which is used to treat sore throats. Because of the high toxicity the plant is not widely used in medicine (Khalmatov et al. 1984). Compounds isolated from the plant exhibited anti-tumor activity in vitro (Chodoeva et al. 2005).	114 115 116 117 118
Phytochemistry: The roots contain up to 2.35 % alkaloids and the aboveground parts up to 0.5 %. The roots contain starch and organic acids as well (Khalmatov et al. 1984). The alkaloids phenyl- β -naphthylamine, karakoline, neoline, delcosine, monticamine, songorine, napelline, acetyl napelline, isoboldine, karasamine and 1-benzoylkarasamine, etc. were found in the aboveground parts (Sultankhodzhaev et al. 1973; 1986; Sultankhodzhaev and Tadzhibaev 1976; Sultankhodzhaev 1993; Atta-ur-Rahman et al. 2005; Chodoeva et al. 2005).	119 120 121 122 123



▲ *Achillea setacea* Waldst. & Kit. Photos: Andrei Lubchenko
◀ *Achillea asiatica* Serg. Photo: Evgeny Davkaev



▲ *Achillea filipendulina* Lam. Photos: Alim Gaziev
▼ *Aconitum karakolicum* Rapaics. Photos: Alexander Naumenko



Aconitum leucostomum Worosch. – Ranunculaceae	124
Synonyms: None	125
English name: Unknown	126
Russian name: Аконит белоустый (Akonit beloustyy)	127
Uzbek name: Unknown	128
Kyrgyz name: Бурма кара, Ак темгилдуу, Үү коргошуну (Burma kara, Ak temgilduu, Uu korgoshchunu)	129
Description: Herbaceous perennial with rope-like roots. Stem 70–200 cm tall, erect. Leaves large, 10–20 cm long and 20–40 cm wide, reniform in outline, palmatisect with 5–11 lobes; basal leaves and lower stem leaves long-petiolate. Inflorescence a dense, many-flowered raceme; pedicels with 2 bracteoles. Flowers irregular, with 5 sepals. Sepals petaloid, dark violet, interior almost white; upper sepal hood-shaped, beaked. Petals 2, each with a spur. Fruit a follicetum with 3 follicles, glabrous or glandular hairy.	130 131 132 133 134
Other distinguishing features: Interior of sepals almost white.	135
Phenology: Flowers in July and August and fruits in August and September.	136
Reproduction: By seeds.	137
Distribution: Ysyk-Kol and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan.	138
Habitat: On the edges of spruce and juniper forests, forest glades; found on northern slopes.	139
Population status: Common, forming dense groups.	140
Traditional use: Unknown.	141
Documented effects: The plant has antibacterial and antiarrhythmic activity. The preparation <i>Allapinin</i> , which is prepared from the aboveground parts and contains the hydrobromic salts of lappaconitine alkaloids, is used as an antiarrhythmic-class I (Gammerman et al. 1990). The alkaloid songorine was found to enhance excitatory synaptic transmission in rat hippocampus and may act as a non-competitive antagonist at the GABA(A) receptor (Zhao et al. 2003).	142 143 144 145
Phytochemistry: The plant contains high quantities of alkaloids: roots – 0.8–4.9 %, stems – 0.3–1, leaves – 0.6–3.9 %, and flowers – 1.3–4.5 %. Lappaconitine, lappaconidine, corydine, glaunidine, N-dimethyl colletine, and others have been isolated from the aboveground parts. The alkaloids mesaconitine, aksine, acsinatine, excelsine, lappaconitine, lappaconidine have been isolated from the root. Flavonoids, coumarins, saponins, and tannins are also found in the roots (Gammerman et al. 1990; Yue et al. 1996; Zhao et al. 2003).	146 147 148 149 150

151 **Aconitum soongaricum Stapf. – Ranunculaceae**152 **Synonyms:** *Aconitum karakolicum* Rapaics. (some authors recognize this as a separate species).153 **English name:** Unknown154 **Russian name:** Аконит джунгарский (Akonit dzhungarskiy)155 **Uzbek name:** Zhoongar parpisi156 **Kyrgyz name:** Жунгар уу коргошуну (Zhungar uu korgoshchunu)157 **Description:** Herbaceous perennial with tuber-like roots. Stems 70–130 cm tall, simple or branched. Leaf blades circular-
158 cordate in outline, 5–9 cm long, 8–12 cm wide, palmatisect with 5 segments divided to the base; segments pinnatifid with
159 2 or 3 linear lobes, lobes 3–5 mm wide. Inflorescence an apical raceme. Flowers irregular. Sepals 5, petaloid, violet; upper
160 sepal hood-shaped, convex, with a long beak. Petals 2, each with a spur. Fruit a follicetum with 3 follicles. Seeds 4–5 mm
161 long.162 **Other distinguishing features:** Forms conical, horizontally segmented (chain-like) roots. Distinguished from *Aconitum*
163 *karakolicum* by having wider leaf segments and glabrous inflorescence rachis and pedicels.164 **Phenology:** Flowers in June-July, fruits in July-August.165 **Reproduction:** By seeds.166 **Distribution:** Ysyk-Kol province of Kyrgyzstan; not found in the flora of Uzbekistan.167 **Habitat:** In meadows with diverse grass species.168 **Population status:** Common, forming dense groups.169 **Traditional use:** Prepared in the same way as *Aconitum karakolicum*. Used in Kyrgyz folk medicine to treat tuberculosis,
170 radiculitis, and headaches, and also to treat different types of cancer (Khalmatov et al. 1984).171 **Documented effects:** An infusion of the tubers is a component in the preparations *Acofit (Radiculin)*, which is used to treat
172 radiculitis, neuritis, and rheumatism, and *Ehinor (Anginol)*, which is used to treat tonsillitis and malignant tumors. The
173 coumarin fraction has antitumor properties. Because of high toxicity the plant is not widely used in medicine (Tolmachev
174 1976). Alkaloids isolated from the plant have antiarrhythmic and CNS-stimulating activities (Salimov et al. 2004).175 **Phytochemistry:** Underground parts contain carbohydrates, starch, organic acids, 1.23–3.4 %, alkaloids, and coumarins up
176 to 0.3 %. Aboveground parts contain 0.56–0.7 % alkaloids and vitamin C. The inflorescence contains flavonoids, and
177 seeds contain up to 32 % fatty oil (Plant Resources of the USSR 1985; Salimov et al. 2004).

Aconitum talassicum Popov – Ranunculaceae	178
Synonyms: None	179
English name: Monkshood	180
Russian name: Аконит таласский (Akonit talasskiy)	181
Uzbek name: Ok parpi	182
Kyrgyz name: Талас уу коргошуну, Кара барпы (Talas uu korgoshchunu, Kara barpy)	183
Description: Herbaceous perennial with narrow-conical roots. Stems up to 1.5 m tall. Leaf blades circular-pentagonal, 6–11 cm long, 7–16 cm wide, palmatisect nearly to the base, with 3–5 wedge-shaped, narrow segments; each segment divided into 2–3 sharply toothed lobes; lobes broadly lanceolate. Inflorescence an apical raceme. Flowers irregular. Sepals 5, petaloid, light-blue to blue; upper sepal hood-shaped with beak; lateral sepals obovate. Petals 2, each with a spur. Fruit a follicetum with 3 follicles.	184 185 186 187 188
Other distinguishing features: Forms segmented (chain-like), horizontal roots. Leaves not as finely dissected as <i>Aconitum karakolicum</i> and <i>A. soongaricum</i> .	189 190
Phenology: Flowers in June-August and fruits in August-September.	191
Reproduction: By seeds.	192
Distribution: Talas province of Kyrgyzstan; Toshkent and Samarqand provinces of Uzbekistan.	193
Habitat: In sub-alpine zones, in wet meadows in river valleys, and among junipers. Endemic to the Tian Shan and Pamiro-Alai mountains.	194 195
Population status: Common, forming dense groups.	196
Traditional use: An infusion of the roots is used to treat rheumatism and malaria (Khalmatov 1964). In veterinary medicine an infusion is used for flesh wounds and skin ulcers (Aldashev 1979).	197 198
Documented effects: The alkaloid talatizamine has effects similar to those of curare as well as ganglio-blocking actions (Khamdamov 1972).	199 200
Phytochemistry: The aboveground parts contain 1.01 % total alkaloids and roots contain 1.92–3.63 % total alkaloids (Khalmatov 1964). Talatizamine, talatizine, talatizidine, isotalatizine, condelphine, and others compounds were isolated from the total alkaloids (Yunusov et al. 1954; Yunusov 1981; Nishanov et al. 1991; Yue et al. 1994).	201 202 203

204 **Acroptilon repens (L.) DC. – Asteraceae**205 **Synonyms:** *Acroptilon picris* (Pall.) C.A. Mey., *Centaurea repens* L.206 **English name:** Russian knarweed207 **Russian name:** Горчак ползучий (Gorchak polzuchiy)208 **Uzbek name:** Kakra209 **Kyrgyz name:** Соилоочу кекире (Soyloochu kekire)

210 **Description:** Herbaceous rhizomatous, perennial. Stems 20–60 cm tall, straight, arachnoid-hairy, with sessile glands. Leaves
211 coriaceous, grayish-green, oblong, sessile; basal and lower leaves oblong, 4–15 cm long; upper leaves oblong, linear or
212 linear-lanceolate, 1–7 cm long. Inflorescences oval capitulas, from 8 to 65, arranged in panicles. Disk flowers 1–1.5 cm
213 long, dark pink. Ray flowers absent. Fruits obovate achenes, 2–4 mm long, light in color, 8–30 in each capitula. Pappus
214 white, short-pinnate.

215 **Other distinguishing features:** Staminal filaments are free and smooth. Basal and lower leaves often withering by flowering
216 time.

217 **Phenology:** Flowers in June-July, fruits in July-August.218 **Reproduction:** Reproduces abundantly by seeds and rhizomes.219 **Distribution:** All of Uzbekistan and Kyrgyzstan.220 **Habitat:** The chul, adyr, and tau zones. Grows on stony and clay-soiled slopes, and in abandoned fields.221 **Population status:** Common, forms large populations.

222 **Traditional use:** A water infusion of the herb is used to treat malaria, epilepsy, and other diseases. The root of the herb is
223 used as an emetic (Khalmatov 1964). In the folk medicine of Central Asia, Azerbaijan, and Crimea, a water infusion of
224 the plant is used to treat malaria, and in Azerbaijan for treatment of epilepsy. Because this plant is poisonous, internal use
225 of this species must be done with caution (Makhlayuk 1992).

226 **Documented effects:** Severe poisoning in farm animals occurs when animals are fed hay containing small amounts of the
227 herb, but toxicity has only been observed when plants are in flower; plants mowed before flowering do not seem to be
228 poisonous (Ogolevitz 1951). The plant causes a nervous system disease and neural cell necrosis when consumed by
229 horses. Repin, a sesquiterpene lactone isolated from the plant, showed high toxicity to chicken embryo sensory neurons
230 (Stevens et al. 1990). Volatile oil isolated from the aboveground parts strongly inhibited the growth of the bacteria
231 *Staphylococcus saprophyticus* and *Staphylococcus epidermidis* (Norouzi-Arasi et al. 2006).

232 **Phytochemistry:** The plant contains traces of saponins, tannins and bitter substances, 4 % glycoalkaloids, 0.06 % essential
233 oils and sesquiterpene lactones (Ogolevitz 1951; Stevens et al. 1990). The main constituent of volatile oil isolated from
234 the aboveground parts was caryophyllene oxide (36.6 %; Norouzi-Arasi et al. 2006).



◀ **Aconitum leucostomum**
Worosch.

Photos: *left*: Vladimir Epiktetov; *center* and *right*: Denis Mirin

Aconitum soongaricum Stapf. ▶

Photos: Vladimir Epiktetov

▼ **Aconitum talassicum** Popov

Photos: Evgeny Davkaev



Acroptilon repens (L.) DC. ▼▶

Photos: *right*: Evgeny Davkaev;
below and *lower right*: Alim Gaziev



236 ***Agrimonia asiatica* Juz. – Rosaceae**237 **Synonyms:** *Agrimonia eupatoria* ssp. *asiatica* (Juzepczuk) Skalický238 **English name:** Agrimony239 **Russian name:** Репейничек азиатский (Repeynichek aziatskiy)240 **Uzbek name:** Sariq choiy241 **Kyrgyz name:** Азия уйгакчасы (Aziya uygakchasy)

242 **Description:** Herbaceous perennial, 30–130 cm tall, with a short, thick rhizome. Stem densely hairy with very dense, stiff,
243 horizontal hairs and fewer shorter, softer hairs. Leaves odd-pinnate, stipulate, hairy, with few small yellow glands; leaflets
244 with large-dentate margins. Inflorescence a spike-shaped raceme, reaching 40 cm during fruiting. Flowers 10–12 mm in
245 diameter, with short pedicels, petals yellow, twice as long as sepals. Fruits achenes enclosed in the hypanthium. Hypanthium
246 6–9 mm long and almost as wide, with rows of prickles towards the top.

247 **Other distinguishing features:** Wounded roots exude a fluid that quickly turns black.248 **Phenology:** Flowers in May-June, fruits in July-August.249 **Reproduction:** By seeds.250 **Distribution:** All of Uzbekistan; Chuy, Osh, Talas, Jalal-Abad, and Batken provinces of Kyrgyzstan.251 **Habitat:** The chul, adyr, and tau zones. Plains, walnut forests, fields, along small canals, along roads, in bushy thickets, and
252 shaded areas of orchards.253 **Population status:** Common, usually found as individual plants.

254 **Traditional use:** A decoction of the underground parts and dried stems and leaves is used in case of gastrointestinal diseases,
255 as an astringent, to treat rheumatism, intestinal infections, fever, edema, as diuretic, and as a mouth wash. A decoction of
256 the flowers is used to treat hemorrhoids, body rashes, and as a hemostatic (Akopov 1981).

257 **Documented effects:** An infusion and liquid extract showed hemostatic effects (Khalmatov 1964). An aqueous extraction of
258 the aboveground plant parts increases diuresis, and it has been shown that an infusion and liquid extracts have hemostatic
259 actions (Akopov 1981). An aqueous extraction of the aboveground parts of *Agrimonia eupatoria* inhibited hepatitis B
260 surface antigen production in vitro (Kwon et al. 2005).

261 **Phytochemistry:** The plant contains ursolic acid (Ibragimov and Khazanovich 1972). Above and underground parts contain
262 tannins, flavonol glycosides, B-vitamins, saponins, and trace alkaloids (Akopov 1981).

Ajuga turkestanica (Regel) Briq. – Lamiaceae	263
Synonyms: None	264
English name: Unknown	265
Russian name: Живучка туркестанская (Zhivuchka turkestanakaya)	266
Uzbek name: Kapalak kunmas	267
Kyrgyz name: Unknown	268
Description: Subshrub with a robust root. Stems 10–50 cm tall, pale brown, reddish or whitish, upper portions covered with fine, soft hairs. Leaves opposite, oblong-elliptic or obovate, 4.5–6 cm long, 1.4–1.8 cm wide, often soft-hairy, nearly sessile, margins usually entire. Flowers solitary, axillary. Calyx campanulate, hairy, with 5 narrowly lanceolate lobes. Corolla 2-lipped, bright pink-purple, with dark veins, rarely white, 2.5–4 cm long; upper lip very short; lower lip large, trilobite, the center lobe clawed and with 2 lobules; stamens 4. Fruits oblong nutlets, 7 mm long, olive-brown.	269 270 271 272 273
Other distinguishing features: Corolla tube nearly twice as long as calyx.	274
Phenology: Flowers in May- June, fruits by the end of May.	275
Reproduction: By seeds.	276
Distribution: South Pamiro-Alai: Surxondaryo Province of Uzbekistan; absent in Kyrgyzstan.	277
Habitat: The ady and tau zones. On clay-soiled and stony slopes in areas with mixed soil types and areas with gypsum and red sandstone.	278 279
Population status: Usually found in populations of 10–300 individuals, and in greater numbers in herbaceous <i>Artemisia</i> -grass complexes.	280 281
Traditional use: Plants in the genus <i>Ajuga</i> are used medicinally to treat weight deficiency, reduced hair growth, ulcers, burns, and to heal wounds. They are also used as a restorative for weakened people (Iordanov et al. 1970; Ikan and Ravid 1971; Kovaleva 1971; Zavrazhanov et al. 1972).	282 283 284
Documented effects: Biological activity of ethanolic extracts of the aboveground parts is due to the presence of phytoecdysteroid compounds. The phytoecdysterones ecdysterone, turkesterone, and cyasterone have anabolic activity. In contrast to the stero-anabolics (nerobol), androgenic action is absent in studied phytoecdysterones. In animals, these compounds have a tonic action and increase resistance to various stress factors (Syrov et al. 1975a, b; Syrov and Kurmukov 1975b, 1976b, c, d; Aizikov et al. 1978; Mamatkhanov et al. 1998). During animal tests, ecdysterone decreased the area of necrosis after experimental myocardial infarction, decreased intracellular enzyme release into the blood and accelerated enzyme reduction until normal (Ermishina et al. 1982; Kurmukov and Ermishina 1986, 1991). Ecdysterone, under the preparation name <i>Ecdysten</i> , has successfully passed clinical tests in several clinics in Russia and Uzbekistan and has been allowed for medical use in the treatment of cardiac infarction, rehabilitation of post-infarction conditions, and to treat fetal hypoxic hypotropia (Iskanderova and Sharipova 1992; Kurmukov and Ermishina 1991; Kurmukov and Kurmukova 1992; Kurmukova 2000a, b; Kurmukova and Kurbanov 1999). The preparation <i>Ayustan</i> , which contains phytoecdysterones, is also used in medicine.	285 286 287 288 289 290 291 292 293 294 295 296
Phytochemistry: This species contains the following oxysteroid compounds and phytoecdysterones: turkesterone, ecdysterone, cyasterone, and others (Saatov et al. 1977; Usmanov et al. 1975, 1977; Mamatkhanov et al. 1998; Abdukadirov et al. 2005).	297 298 299

300 **Alhagi pseudalhagi (M. Bieb.) Desv. – Fabaceae**

301 **Synonyms:** *Alhagi camelorum* Fisch. ex DC., *Alhagi maurorum* Medic., *Alhagi persarum* Boiss. & Buhse, *Hedysarum*
302 *alhagi* L., *Hedysarum pseudalhagi* M. Bieb.

303 **English name:** Camel's thorn

304 **Russian name:** Янтак ложный, Верблюжья колючка ложная (Yantak lozhnyy, Verblyuzh'ya kolyuchka lozhnaya)

305 **Uzbek name:** Yontok

306 **Kyrgyz name:** Жантак (Zhantak)

307 **Description:** Herbaceous perennial with a deep root system. Stems green, 50–120 cm tall, much branched, with upward
308 curving thorns. Leaves alternate, simple, obovate, 7–20 mm long, margins entire, apex rounded. Inflorescences axillary
309 racemes with 3–8 flowers; inflorescence rachis ending with a spine. Calyx campanulate, with or without 5 teeth. Corolla
310 papilionaceous, 8–9 mm long, pink to brownish-red. Fruit a moniliform legume, 1–3 cm long, curved or straight with 1–5
311 seeds. Seeds small, glabrous, kidney-shaped, brown.

312 **Other distinguishing features:** Stamens 10, nine of the filaments fused. Banner petal obovate, keel blunt, equal in length to
313 banner, wings shorter than keel.

314 **Phenology:** Flowers in May-September, fruits in August-October.

315 **Reproduction:** By seeds and rhizomes.

316 **Distribution:** Almost all provinces of Uzbekistan; Chuy, Osh, and Jalal-Abad provinces of Kyrgyzstan.

317 **Habitat:** The chul, adyr and tau zones. In abandoned fields, as a weed in fields, and along irrigation networks.

318 **Population status:** Common.

319 **Traditional use:** Used in Iran and other eastern countries as a laxative and antipyretic (Khalmatov 1964). An infusion of the
320 roots is used to treat liver diseases and stomach and duodenal ulcers as well as diuretic. A galenic preparation of the
321 aboveground parts is used to treat colitis, gastritis, stomach ulcers, dysentery, cervical erosion, to heal wounds, to treat
322 inflammation of the ear, nose and throat, as a choleric, to quench thirst, to reduce sweating and as an antipyretic, anti-
323 inflammatory, and cough remedy (Karimov and Shomakhmudov 1993).

324 **Documented effects:** A dry extract from the aboveground parts, as well as the total proanthocyanidins have antihypoxic,
325 antioxidant, angioprotective, and hypocholesterimic abilities (Aizikov et al. 1986). An ethanolic extract of the aboveground
326 plant parts had antiulcerogenic activity in rats (Amani et al. 2006), and a methanolic extract exhibited an antidiarrheal
327 effect (Atta and Mouneir 2004).

328 **Phytochemistry:** The herb contains 3.9–8.2 % tannins, up to 0.2 % coumarins, up to 1,000 mg%,¹ vitamin C, about 0.8 %
329 essential oils, as well as up to 3.4 % flavonoids. The roots contain alkaloids (0.17–0.19 %), glycosides, resins (up to
330 5.67 %), pigments and sugars. Proanthocyanidins have been isolated from this species (Karimov and Shomakhmudov
331 1993). The flavonoids catechin, epigallocatechin, galocatechin, leucodelphinidin, quercetin, rutin, etc. and the flavonone
332 glycosides alhagitin and alhagidin have also been isolated from the plant (Singh et al. 1999; Awaad Amani et al. 2006).

333

¹Milligram% (mg%) – A unit used to describe concentration. Milligrams of a specific substance contained in 100 ml of a solution or in 100 g of the analyzed material. Often used to describe vitamin content in plants and foods.

Allium karataviense Regel – Alliaceae	334
Synonyms: None	335
English name: Unknown	336
Russian name: Лук каратавский (Luk karatavskiy)	337
Uzbek name: Chuchka kuloq	338
Kyrgyz name: Кара Тоо пиязы (Kara Too piyazy)	339
Description: Herbaceous perennial with large, spherical bulb. Bulb 2–6 cm in diameter, with a blackish or grayish paper-like coat. Stem short, 10–25 cm tall, sometimes half buried in the soil, stems shorter than leaves. Leaves lanceolate, oblong, (3–)5–15 cm wide with smooth margins. Inflorescence a dense, many-flowered, spherical umbel. Pedicels equal in length, 3–4 times longer than perianths, lacking bracts. Flowers with 6 tepals. Tepals 5–7 mm long, linear, apex rounded, pink-violet with a dark vein. Stamens 6. Fruit an obovate capsule, 8 mm wide.	340 341 342 343 344
Other distinguishing features: Ovary with a rough surface. Leaves broader than those of related species.	345
Phenology: Flowers in April-May, fruits in May-June.	346
Reproduction: By seeds and vegetatively by lateral bulblets.	347
Distribution: Toshkent and Farg'ona provinces of Uzbekistan; Chuy and Osh provinces of Kyrgyzstan.	348
Habitat: The tau zone. On limestone taluses.	349
Population status: Common.	350
Traditional use: A decoction of the bulb is used in folk medicine to treat lung diseases and shortness of breath (Khalmatov 1964).	351 352
Documented effects: No data.	353
Phytochemistry: Most <i>Allium</i> species contain essential oils, volatile organic compounds, flavonol glycosides, phenols, vitamins, ascorbic acid, mineral salts and microelements (Khalmatov 1964), as well as steroidal saponins and sapogenins (Mimaki et al. 1999).	354 355 356



▲ Agrimonia asiatica Juz.

Photos: *far left and left:* Andrei Lubchenko; *right and far right:* Evgeny Davkaev



◀ Ajuga turkestanica (Regel)

Briq. Photos: *far left:* Authors; *left:* Diana Cheng



▼ Alhagi pseudalhagi (M. Bieb.)

Desv. Photos: *left:* Evgeny Davkaev; *right:* Alim Gaziev

▼ Allium karataviense Regel

Photos: *left:* Alim Gaziev; *middle and right:* Evgeny Davkaev



Allium suvorovii Regel – Alliaceae	358
Synonyms: None	359
English name: Unknown	360
Russian name: Лук Суворова (Luk Suvorova)	361
Uzbek name: Yowoiy piyoz	362
Kyrgyz name: Суворов пиязы (Suvorov piyazy)	363
Description: Herbaceous perennial plant to 1 m tall, with a spherical bulb. Bulb 2–3 cm in diameter, covered with grayish, cracked, almost coriaceous coat that sheathes the base of the stem. Stem 30–100 cm tall. Leaves 2–6, belt-like, much shorter than stem, 5–20 mm wide, margins rough. Inflorescence a dense, many flowered, semispherical or spherical umbel. Pedicels equal in length, 2–5 times longer than perianths, lacking bracts. Flowers with 6 tepals. Tepals 6, ~4 mm long, linear, apex rounded, pink-violet with a darker vein. Stamens 6. Fruit a capsule, broadly-ovate, 5 mm wide.	364 365 366 367 368
Other distinguishing features: It differs from closely related species by having a smooth ovary.	369
Phenology: Flowers in May, fruits in June.	370
Reproduction: By seeds and vegetatively by lateral bulblets.	371
Distribution: Toshkent, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Chuy and Osh provinces of Kyrgyzstan.	372 373
Habitat: The adyr zone. Grows in shallow soil on foothills, as a weed along canals, along the edges of plowed fields, and in orchards and cemeteries. Found in places inaccessible for pasturing and mowing.	374 375
Population status: Rare. Found sporadically as individual plants and in small populations; listed in the Red Book of Rare and Endangered Species of Uzbekistan.	376 377
Traditional use: The bulbs pickled in wine vinegar, are used to treat hemoptysis and to treat incipient tuberculosis. Also used as a phytoncidal remedy to treat various skin diseases, especially eczema and psoriasis (Khalmatov 1964).	378 379
Documented effects: No data.	380
Phytochemistry: See <i>Allium karataviense</i> . The bulbs of <i>A. suvorovii</i> contain various carbohydrates (Khodzhaeva and Turakhozhaev 1992; Khodzhaeva 1994); the seeds contain the carbohydrate stachyose (Khodzhaeva and Kondratenko 1984).	381 382 383

384 ***Allochrusa gypsophiloides* (Regel) Schischk. – Caryophyllaceae**385 **Synonyms:** *Acanthophyllum gypsophiloides* Regel.386 **English name:** Turkestan soaproot387 **Russian name:** Аллохруза качимовидная, Колючелистник качимовидный, Мыльный корень (Allokhruza kachimovid-
388 naya, Kolyuchelistnik kachimovidnyy, Myl'nyy koren')389 **Uzbek name:** Beh, Etmak, Kachimsimon etmak390 **Kyrgyz name:** Качимдай кок тикен (Kachimday kok tiken)391 **Description:** Herbaceous perennial, 30–80 cm tall, with a strong taproot reaching 6 m deep. Stems thin, branched, short-
392 pubescent or glabrous. Leaves opposite, linear or linear-lanceolate, acute, 1–2.5 cm long, glabrous, sessile. Inflorescence
393 paniculiform, loose, branched. Flowers pale-pink, with long pedicels. Fruit a capsule with 1–2 seeds. Seeds rough,
394 flattened, light-brown.395 **Other distinguishing features:** Capsule obovate or spherical, ca. 2 mm long.396 **Phenology:** Flowers in June-July, fruits in July-August.397 **Reproduction:** By seeds.398 **Distribution:** Toshkent, Jizzax, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; in the Chatkal, Talas,
399 and Pskem ranges in Kyrgyzstan.400 **Habitat:** The adyr and tau zones. Stony slopes with rocky debris.401 **Population status:** Rare, listed in the Red Book of Rare and Endangered Species of Uzbekistan.402 **Traditional use:** Roots are used to treat coughs and applied externally to heal wounds. A decoction of the root is recom-
403 mended as an expectorant for bronchitis (Khalmatov 1964). An infusion of the roots is used as a choleric, diuretic, and
404 laxative. The root is brewed in a tea and drunk to treat gastrointestinal, skin and venereal diseases, spleen, liver and kidney
405 diseases, as well as metabolism dysfunction. An infusion of the aboveground parts is used as an expectorant and laxative
406 (Khodzhimatov 1989).407 **Documented effects:** Saponins from this species increase the secretory activity of glands. Pure saponin from this species is
408 used in veterinary medicine to prepare vaccines against anthrax and brucellosis (Khodzhimatov 1989). Treatment with a
409 saponin extracted from the roots antagonized the narcotic effect of chloral hydrate, potentiated the convulsive effect of
410 strychnine, decreased the convulsive and toxic effect of Corazole, and increased diuresis in mice (Polievtssev and Sultanov
411 1971).412 **Phytochemistry:** An important source of saponins. Roots contain up to 30 % saponins with a hemolytic index of 1:1000 or
413 1:2860 and aboveground parts of the plant have saponins with an index of 1:240 (Khalmatov 1964; Yukhananov et al.
414 1972). The roots contain 10–30 % triterpene saponins from which the glycosides gypsogenin and acanthophyllosides B,
415 C and D were isolated (Putieva et al. 1970, 1975, 1979). The aboveground parts contain polysaccharides as well as many
416 saponins (Arifkhodzhaev and Kondratenko 1983; Khodzhimatov 1989).

- Althaea nudiflora Lindl. – Malvaceae** 418
- Synonyms:** *Alcea leucantha* Fisch., *Alcea nudiflora* (Lindl.) Boiss. 419
- English name:** Naked-flowered hollyhock 420
- Russian name:** Алтей голоцветный (Altey golotsvetnyy) 421
- Uzbek name:** Oq gulhairy 422
- Kyrgyz name:** Туксуз гулдуу гулкайыр (Tuksuz gulduu gulkayyr) 423
- Description:** Herbaceous biennial or perennial, to 1.5–2 m high, stellate hairy. Stems cylindrical. Leaves simple, long-petio- 424
late; blade 5–7-lobed with coarse-dentate margins, rough with crowded, stiff, stellate hairs on both sides. Inflorescences 425
terminal, racemiform. Calyx with 5 triangular-lanceolate lobes, densely stellate hairy. Corolla white, 5–8 cm in diameter; 426
petals 5, obovate. Stamens fused into a column. Fruits wheel-shaped schizocarps. Seeds 3–4 mm long, kidney-shaped, 427
red-brown. 428
- Other distinguishing features:** Foliaceous bracts absent. Style with numerous branches. 429
- Phenology:** Flowers in June-August, fruits in July-September. 430
- Reproduction:** By seeds. 431
- Distribution:** Toshkent, Andijon, Namangan, Farg'ona, Samarqand, Qashqadaryo and Surxondaryo provinces of Uzbekistan; 432
Chuy, Ysyk-Kol, Talas, Osh and Jalal-Abad provinces of Kyrgyzstan. 433
- Habitat:** The adyr and tau zones. Shallow soil and stony slopes. 434
- Population status:** Common. 435
- Traditional use:** An infusion of the dried flowers is given to children to treat diarrhea and sialorrhea. A decoction of the roots 436
and seeds is recommended as a hemostatic for post-natal bleeding. A plaster of the flower and leaf powder is used to treat 437
tumors (Khalmatov 1964). The fresh stem is cut and applied to cuts on the skin. The roots and seeds are made into a tea 438
to treat dysuria (Sezik et al. 2004). 439
- Documented effects:** None. 440
- Phytochemistry:** The plant contains mucilage. Leaves contain 165–176 mg% of vitamin C (Khalmatov 1964). The roots 441
contain lipids with cyclopropanoid fatty acids (Sagdullaev et al. 2001) and the flowers contain kaempferol glycosides 442
(Pakudina et al. 1970). 443

445 ***Althaea officinalis* L. – Malvaceae**

446 **Synonyms:** *Althaea kragujevacensis* Pančić ex Diklić & Stevan., *Althaea micrantha* Borbás, *Althaea sublobata* Stokes,
447 *Althaea taurinensis* DC., *Althaea vulgaris* Bubani, *Malva althaea* E.H.L. Krause, *Malva maritima* Salisb., *Malva officinalis*
448 (L.) Schimp. & Spenn.

449 **English name:** Common marshmallow

450 **Russian name:** Алтей лекарственный (Altey lekarstvennyy)

451 **Uzbek name:** Dorivor gulhairi

452 **Kyrgyz name:** Дары гулкан (Dary gulkan)

453 **Description:** Herbaceous perennial. Stems single or multiple, 40–150 cm tall, tomentose. Basal leaves 3–5-lobed; cauline
454 leaves with rounded or cordate base and acuminate apex, margins coarsely serrate. Flowers up to 3 cm across, clustered
455 in leaf axils. Epicalyx with 8–12 segments. Corolla pale-pink with 5 petals. Fruit a disc-shaped schizocarp, with 15–25
456 small, laterally flattened mericarps. Seeds kidney-shaped.

457 **Other distinguishing features:** Stamens many, staminal filaments connate and forming a tube.

458 **Phenology:** Flowers in June-September, fruits in June-October.

459 **Reproduction:** By seeds.

460 **Distribution:** All provinces of Kyrgyzstan; Tashkent, Farg'ona and Samarqand provinces of Uzbekistan.

461 **Habitat:** In places with a high water-table, and along rivers and canals.

462 **Population status:** Common, forming dense groups.

463 **Traditional use:** The roots, flowers and leaves are used as an anti-inflammatory and to treat flu, sore throat, hepatitis, and
464 urinary incontinence. They are also used to treat kidney stones, cystitis, prostate tumors, chronic prostatitis, and joint pain
465 (Kurochkin 1998).

466 **Documented effects:** The plant is used internally to treat eczema, psoriasis, dermatitis and to normalize metabolism. In
467 combination with other preparations, this species is used to treat gastritis, stomach and duodenal ulcers, enterocolitis,
468 food poisoning, dysentery, kidney inflammation, and urinary incontinence. The preparation *Mucaltin*, which is prepared
469 from the herb, is used as an expectorant to treat bronchitis and pneumonia (Kurochkin 1998). A methanolic extract and a
470 decoction of the roots inhibited a variety of bacteria known to cause periodontal disease (Iauk et al. 2003). An extract of
471 the root has been shown to be a potent inhibitor of calcium mobilization associated with UVB-induced pigmentation of
472 skin (Kobayashi et al. 2002a).

473 **Phytochemistry:** Roots consist of 35 % mucilage substances, about 37 % starch, 10 % sucrose, betaine, flavonoids, cou-
474 marins, phenolic acids, and fatty oil. Aboveground parts contain mucilage, carbohydrates (glucose and sucrose), essential
475 oils, vitamin C, and carotene. Seeds contain up to 12 % fatty oil, 1 % phospholipids and pectin (Khalmatov 1964;
476 Tolmachev 1976; Capek et al. 1987; Gudej 1991).



◀ **Allium suvorovii** Regel
Photos: Alim Gaziev



▲ **Althaea nudiflora** Lindl. Photos: Evgeny Davkaev



▲ **Allochrysa gypsophiloides**
(Regel) Schischk.
Photos: Evgeny Davkaev

▼ **Althaea officinalis** L.
Photos: *left*: Andrei Lubchenko;
center and right: Vadim Prokhorov



478 **Amaranthus retroflexus L. – Amaranthaceae**479 **Synonyms:** some authors consider *Amaranthus tricolor* L. a synonym.480 **English name:** Redroot amaranth481 **Russian name:** Щирица запрокинутая (Shchiritsa zaprokinutaya)482 **Uzbek name:** Gultojihuroz, Eshakshura483 **Kyrgyz name:** Кайрылган амарант (Каурылган amarant)484 **Description:** Herbaceous annual with a taproot. Stems 20–100 cm tall, pubescent. Leaves ovate-rhomboid, apex obtuse.485 Inflorescence a dense panicle; bracts lanceolate. Flowers unisexual. Pistillate flowers with 5 tepals and 3 stigmas. Staminate
486 flowers at top of inflorescence; tepals 5, stamens 3–5. Seeds shiny black, lenticular, contained in circumscissile utricles.487 **Other distinguishing features:** Pistillate tepals membranaceous with emarginate or obtuse apices.488 **Phenology:** Flowers in May-June, fruits in August.489 **Reproduction:** By seeds.490 **Distribution:** Agricultural zones in all provinces of Kyrgyzstan and Uzbekistan.491 **Habitat:** In vegetable gardens, orchards, waste places, and along the edges of fields.492 **Population status:** Common, found in dense groups.493 **Traditional use:** A water infusion of the aboveground parts is used to treat colitis, intestinal colic, and as a laxative for con-
494 stipation, as well as a hemostatic to treat hemoptysis, and menstrual and hemorrhoid hemorrhages. A water extract of the
495 dried plant collected during flowering stage is used as an antiprotist and antibacterial. A decoction of the roots is used to
496 treat guinea worm and jaundice. Young stems are used as a source of vitamins. Leaves are used as a diuretic and a decoc-
497 tion of the leaves is used to treat headaches (Zolotnitskaya 1965; Makhlayuk 1967).498 **Documented effects:** An antimicrobial peptide was isolated from the seeds and effectively inhibited the growth of multiple
499 fungi species (Lipkin et al. 2005).500 **Phytochemistry:** Roots contain the betacyanins amaranthin and isobetanin. Leaves contain nitrogenous compounds, 0.96 %
501 betaine and fatty oils, which contain the following fatty acids: miristic, palmitic, stearic, linoleic and linolenic acid. Seeds
502 contain 4.3–7 % fatty oil with the following fatty acids: palmitic (18.9 %), stearic (1.9 %), oleic (51.5 %), linoleic (27.9 %);
503 Plant Resources of the USSR 1985).

Anagallis arvensis L. – Myrsinaceae (formerly in Primulaceae)	505
Synonyms: <i>Anagallis latifolia</i> L., <i>Anagallis phoenicea</i> Scop.	506
English name: Scarlet pimpernel	507
Russian name: Очный цвет пашенный (Ochnyy tsvet pashennyy)	508
Uzbek name: Savun ut, Savunak	509
Kyrgyz name: Кызгылт анагаллис (Kyzgylt anagallis)	510
Description: Herbaceous annual or biennial with multiple branches. Stems quadrangular, glabrous, 10–25 cm long. Leaves opposite, sessile, ovate to elongate-ovate, with black dots on abaxial surface. Flowers brick-red, individual, with long pedicels. Fruit a spherical capsule, opening by a small cover. Seeds small, oval, 3-sided, black, many in each capsule.	511 512 513
Other distinguishing features: Differs from related species by having a brick-red corolla.	514
Phenology: Flowers and fruits in April-September.	515
Reproduction: Only by seeds.	516
Distribution: Toshkent, Farg'ona, Samarqand and Surxondaryo provinces of Uzbekistan; Chuy, Talas and Osh provinces of Kyrgyzstan.	517 518
Habitat: The chul, adyr and tau zones. Along banks of small canals, in river valleys, along roads, in orchards, in fields, and on loess slopes.	519 520
Population status: Common, often found in small populations.	521
Traditional use: A decoction of the herb is used to treat shortness of breath, tuberculosis, gynecological disorders, rabies, and as a diuretic in cases of edema, and is used externally for washing wounds (Khalmatov 1964). The essence from blooming plants is used in homeopathy (Ogolevitz 1951).	522 523 524
Documented effects: Plants collected in the Toshkent region contained saponins with a hemolytic index of 1:2230 (Khalmatov 1964). An aqueous extract of the plant showed significant antifungal activity against isolates of <i>Microsporum canis</i> , <i>Trichophyton mentagrophytes</i> , and <i>Trichophyton violaceum</i> (Ali-Shtayeh and Abu Ghdeib 1999). Saponins isolated from the plant exhibited strong molluscicidal activity when tested against <i>Biomphalaria glabrata</i> and <i>Oncomelania quadrasi</i> (Abdel Gawad et al. 2000). A triterpene saponin isolated from the plant inhibited the replication of herpes simplex virus type 1 and poliovirus type 2 in vitro (Amoros et al. 1987). The plant is noted as being poisonous and extracts of the plant have been shown to be highly toxic to rats (Ogolevitz 1951; Al-Sultan et al. 2003).	525 526 527 528 529 530 531
Phytochemistry: The herb contains the glucoside cyclamine, saponoids and other terpenoid saponins as well as the enzyme primveraza (Ogolevitz 1951; Amoros et al. 1987). The aboveground plant parts contain flavonoids (kaempferol, quercetin), phenylcarbonic acids (caffeic, ferulic, etc.), anthocyanins and fatty oil (MedicineLib.ru 2008).	532 533 534

536 **Anagallis foemina Mill. – Myrsinaceae (formerly in Primulaceae)**

537 **Synonyms:** *Anagallis arvensis* f. *coerulea* (Schreb.) Arechav., *Anagallis arvensis* var. *coerulea* (Schreb.) Gren. & Godr.,
538 *Anagallis arvensis* ssp. *foemina* (Mill.) Schinz & Thell., *Anagallis coerulea* Schreb.

539 **English name:** Blue pimpernel

540 **Russian name:** Очный цвет голубой (Ochnyy tsvet goluboy)

541 **Uzbek name:** Savun ut, Savunak

542 **Kyrgyz name:** Когултур анагаллис (Kogultur anagallis)

543 **Description:** The botanical description of this plant is very similar to that of *Anagallis arvensis*. This species differs by hav-
544 ing a blue corolla. Some botanists consider *Anagallis coerulea* a form or variety of *Anagallis arvensis*.

545 **Other distinguishing features:** Differs from related species by having a blue corolla with dentate lobes and no glands.

546 **Phenology:** Flowers and fruits in April-September.

547 **Reproduction:** Only by seeds.

548 **Distribution:** Surxondaryo province of Uzbekistan; Chuy, Osh and Jalal-Abad provinces of Kyrgyzstan.

549 **Habitat:** The chul, adyr and tau zones. Along banks of small canals, river valleys, along the roads, in orchards, fields, and
550 on loess slopes.

551 **Population status:** Common, found in small populations.

552 **Traditional use:** Same as *Anagallis arvensis*.

553 **Documented effects:** No data.

554 **Phytochemistry:** Similar to *Anagallis arvensis*.

555

Uncorrected Proof

<i>Anchusa azurea</i> Mill. – Boraginaceae	556
Synonyms: <i>Anchusa italica</i> Retz.	557
English name: Italian bugloss, large blue alkanet	558
Russian name: Анхуза итальянская (Ankhuza ital'yanskaya)	559
Uzbek name: Hukuz tili	560
Kyrgyz name: Италия анхузасы (Italiya ankhuzasy)	561
Description: Perennial herb to 1.5 m tall, with multiple, thick, conjoined taproots. Entire plant densely covered with bristly hairs. Stem usually single, sometimes branching, erect. Basal leaves in a rosette, oblanceolate, 10–30 cm long, petiolate; upper leaves alternate, oblong or lanceolate, sessile. Inflorescences terminal, bracteate, helicoid racemiform. Calyx lobes linear, divided nearly to the base. Corolla bright blue, funnellform, 10–15 mm in diameter, 5-lobed, with 1–1.5 cm long, bristly-hairy pedicels. Fruits gray nutlets, erect, 5–8 long mm long, 3–5 mm wide.	562 563 564 565 566
Other distinguishing features: Stamens inserted at the top of corolla tube. Fruits 3-sided.	567
Phenology: Flowers in the end of April-July, fruits in May-August.	568
Reproduction: Only by seeds.	569
Distribution: Toshkent, Farg'ona, Andijon, Samarqand, and Surxondaryo provinces of Uzbekistan; Chuy, Osh and Jalal-Abad provinces of Kyrgyzstan.	570 571
Habitat: The chul, adyr and tau zones. In fields, waste grounds, orchards, wheat fields, and oases.	572
Population status: Common, found as single individuals.	573
Traditional use: A decoction of the flowers is used to treat chest aches, neurasthenia, and asthma, and is used as a laxative, febrifuge, and cough remedy as well. An ointment, prepared by boiling the roots in cow fat, is used as hemostatic and to heal wounds (Khalmatov 1964). In Iraq, a decoction of the flowers is used as a sedative, analgesic, sudorific, and diuretic (Al-douri 2000).	574 575 576 577
Documented effects: Saponins from this species have a hemolytic index of 1:2800–1:20000 (Khalmatov 1964). An extract of the aboveground parts showed significant antibacterial effect against <i>Pseudomonas aeruginosa</i> (Bazzaz and Haririzadeh 2003).	578 579 580
Phytochemistry: All plant tissue contain saponins. The roots contain dyes (alkanin and anchusin), anchusa acid, resins, and waxes (Khalmatov 1964). Oil extracted from the seeds contains γ - and α -linolenic acid as well as stearidonic acid (Guil-Guerrero et al. 2001).	581 582 583



▲ **Amaranthus retroflexus L.**

Photos: Maxim Kucherov

▼ **Anagallis foemina Mill.** Photos: *top*: Alim Gaziev; *bottom*: Andrei Lubchenko



▲ **Anagallis arvensis L.** Photos: Sasha Eisenman



▼ **Anchusa azurea Mill.** Photos: *top left*: Evgeny Davkaev; *lower left and right*: Alim Gaziev



Artemisia absinthium L. – Asteraceae	585
Synonyms: None	586
English name: Common wormwood, absinthium, armoise absinthe	587
Russian name: По́лынь го́рькая (Polyn' gor'kaya)	588
Uzbek name: Erman, Achik erman	589
Kyrgyz name: Эрман шыбак (Erman shybak)	590
Description: Herbaceous perennial with a short taproot. Stems up to 1.5 m tall, sometimes with short, lateral, vegetative stems. Basal leaves bi- or tripinnatisect, long-petiolate; cauline leaves alternate; lower cauline leaves short-petiolate, bipinnatisect; upper cauline leaves small, almost sessile. Inflorescences many-flowered capitula with 40–70 flowers, ca. 3 mm in diameter, globose, nodding, in narrow to broadly pyramidal panicles; involucral bracts linear; receptacle convex, densely hairy. Disc flowers yellow; ray flowers absent. Fruits oblong to wedge-shaped achenes, about 1 mm long.	591 592 593 594 595
Other distinguishing features: The whole plant is gray-silver due to short, appressed hairs.	596
Phenology: Flowers in July-August, fruits in August-September.	597
Reproduction: By seeds.	598
Distribution: Almost all provinces of Kyrgyzstan and Uzbekistan.	599
Habitat: From valleys to the mid-belt of mountains. On slopes, in meadows, along rivers, and near cultivated and in abandoned fields.	600 601
Population status: Common, in some places forming dense groups.	602
Traditional use: Leaves, which are collected before and at the beginning of flowering, are used in a decoction as a carminative, a vermifuge, and to treat dyspepsia, loss of appetite, insomnia, diseases of the liver, stomach, spleen, and gall bladder, fever, hemorrhoids, malaria, and intestinal ulcers, as well as to heal wounds (Khalmatov et al. 1984).	603 604 605
Documented effects: This species is used to make preparations to treat cases of gastritis with low stomach acidity, to increase appetite, and as a choleric. In combination with other medicines, preparations are also used to treat chronic diseases of the pancreas, stomach, and intestinal tract. Because of the presence of azulene, this species is used to treat allergic reactions of the skin (Kurochkin 1998). Ethyl acetate and chloroform extracts of the whole plant inhibited a variety of microorganisms (Erdogrul 2002).	606 607 608 609 610
Phytochemistry: The herb contains 0.17–2 % essential oil which contains sesquiterpene lactones (absinthin, anabsinthin and artabasin), flavonoids (artemetin), tannins, organic acids, vitamin C and carotene (Khalmatov et al. 1984). The roots were found to contain many lignans (Greger and Hofer 1980).	611 612 613

615 **Artemisia annua L. – Asteraceae**

616 **Synonyms:** *Artemisia chamomilla* C. Winkl.

617 **English name:** Sweet sagewort, sweet wormwood, sweet annie, chinese wormwood

618 **Russian name:** Полюнь однолетняя (Polyn' odnoletnyaya)

619 **Uzbek name:** Burgan

620 **Kyrgyz name:** Бир жылдык шыбак (Bir zhyldyk shybak)

621 **Description:** Herbaceous annual. Stems often single, 15–200 cm tall, erect. Lower leaves up to 7 cm long and wide, ovate
622 in outline, bi- or tripinnatisect, petiolate; cauline leaves bipinnatisect, triangular to broadly ovate, becoming simpler and
623 smaller towards top of stem. Inflorescences globose capitula with ca. 30 flowers, in a leafy, open panicle; involucre bracts
624 linear. Disc flowers pale- or greenish-yellow; ray flowers absent. Fruits flat achenes, 0.5–1 mm long.

625 **Other distinguishing features:** Plant has a sweet aroma.

626 **Phenology:** Flowers in July and August, fruits in August and September.

627 **Reproduction:** By seeds.

628 **Distribution:** All provinces of Kyrgyzstan and Uzbekistan.

629 **Habitat:** From valleys to the mid-belt of mountains. In agricultural zones, near canals, in orchards, and vegetable gardens.

630 **Population status:** Common, forming dense groups.

631 **Traditional use:** Leaves are collected in spring and aboveground parts in autumn. The juice from fresh leaves is used to treat
632 skin diseases (scabies, abscesses, bacterial and fungal diseases, etc.). The dried leaves are used to prepare an ointment that
633 is used to treat eczema. A decoction of the aboveground parts is used to increase appetite. Traditional doctors use an infu-
634 sion of the herb to treat rheumatism and skin diseases (Khodzhimatov 1989).

635 **Documented effects:** An extract of this species inhibits the development of anthrax, by causing loss of pathogenic ability
636 and killing bacterial cells (Khodzhimatov 1989). This plant species is the source of artemisinin, which, in combination
637 with other drugs, is used as a highly effective treatment for malaria worldwide (World Health Organization 2006).
638 Artemisinin has also been shown to cause apoptosis in human cancer cells (Singh and Lai 2004).

639 **Phytochemistry:** The aboveground parts contain lactones (artemisinin and arteannuin), coumarins (scopoletin), 0.12–0.65 %
640 essential oil, tannins, alkaloids, resins, sugars, and vitamin C. Maximum essential oil content was observed during the
641 flowering period. Plants collected near Bishkek (Kyrgyzstan) contained 0.21 % essential oil and 2.44 % tannins (Khalmatov
642 1964; Khodzhimatov 1989).

643

Artemisia dracunculus L. – Asteraceae

644

Synonyms: *Artemisia aromatica* A. Nelson, *Artemisia dracunculina* S. Watson, *Artemisia dracunculoides* Pursh, *Artemisia dracunculoides* ssp. *dracunculina* (S. Watson) H. M. Hall & Clements, *Artemisia glauca* Pallas ex Willdenow, *Oligosporus dracunculus* (L.) Poljak.

645

646

647

English name: Russian tarragon, wild tarragon, estragon, silky wormwood

648

Russian name: По́лынь Эстрагон (Polyn' estragon)

649

Uzbek name: Sherolgin

650

Kyrgyz name: Шыраалжын шыбак (Shyraalzhyn shybak)

651

Description: Herbaceous, rhizomatous perennial to 50–120 cm tall, with a woody caudex and fibrous roots. Stems numerous, erect, green, yellowish or reddish brown, partially woody, glabrous. Leaves alternate, 5–8 cm long, linear-lanceolate, usually entire; lower leaves often irregularly lobed or trilobate, mostly glabrous. Inflorescences globose to ovate capitula arranged in panicles. Disk flowers pale-yellow, only peripheral flowers fertile; ray flowers absent. Fruits oblong achenes, ca. 1 mm long, brown.

652

653

654

655

656

Other distinguishing features: Receptacle where flowers are attached is naked (lacking chaff, scales, hairs, etc.). The plant has a unique smell.

657

658

Phenology: Flowers in June, fruits in September.

659

Reproduction: By seeds and rhizomes.

660

Distribution: All of Uzbekistan and Kyrgyzstan.

661

Habitat: The tau zone. On soft and rocky mountain sides. Often planted in vegetable gardens for use as a culinary herb.

662

Population status: Common, often found in dense groups.

663

Traditional use: It is used to treat edema and scurvy, dyspepsia, to improve appetite, and as a carminative. A powder of the plant is used to treat oral diseases. Tarragon from Uzbekistan has been noted to have anti-helminthic action. Leaves are also recommended as a good source of carotene (Khalmatov 1964).

664

665

666

Documented effects: After clinical tests, use of the liquid extract of tarragon was recommended to treat patients with chronic low-acid gastritis (Khalmatov et al. 1984). Essential oil isolated from the aboveground parts of *Artemisia dracunculus* "Piemontese" exhibited strong antifungal activity when tested against *Candida albicans*, *C. lusitaniae*, *C. glabrata*, and *C. tropicalis*, and weak antimicrobial effects against *Xanthomonas maltophilia* and *Proteus mirabilis* (Curini et al. 2006). An ethanolic extract of the plant significantly reduced hyperglycemia in mice with chemically induced insulin deficiency and diabetes, and reduced hyperglycemia in genetically diabetic mice (Logendra et al. 2006; Ribnicky et al. 2006).

667

668

669

670

671

672

Phytochemistry: The flowering herb contains 0.1–0.7 % essential oils, 41.8 mg% (for absolute dry weight) carotene, 190 mg% vitamin C and alkaloid traces (Khalmatov et al. 1984). The essential oils of Central Asian plants contain 65–85 % d-sabinene, about 10 % myrcene, 5 % sesquiterpene fractions, about 0.5 % methoxy-cinnaroid aldehyde, and 7–15 % resins. Central Asian tarragon oil is substantially different from Western European tarragon oil because it doesn't contain methyl-chavicol (Khalmatov 1964). The herb contains flavonoids, alkaloids, and coumarins (Mallabaev et al. 1971, 1970; Mallabaev and Sidiyakin 1976; Hofer et al. 1986; Bohm and Stuessy 2001; Saadali et al. 2001; Logendra et al. 2006).

673

674

675

676

677

678

679

680

681 **Artemisia leucodes Schrenk – Asteraceae**682 **Synonyms:** *Seriphidium leucodes* (Schrenk) Poljak.683 **English name:** Unknown684 **Russian name:** Полюнь беловатая (Polyn' belovataya)685 **Uzbek name:** Oq shuvoq686 **Kyrgyz name:** Ак шыбак (Ak shybak)

687 **Description:** Herbaceous annual or biennial, 30–90 cm tall, covered with long, white, raised hairs. Stems single or multiple,
688 erect, branched. Lower stem leaves petiolate, tripartite-pinnatisect, 3–7 cm long; cauline leaves sessile, tripartite.
689 Inflorescences 3–5-flowered capitula in panicles. Disk flowers yellow, punctate glandular; ray flowers absent. Fruits obo-
690 vate achenes, 2–2.25 mm long, olive-colored.

691 **Other distinguishing features:** Leaves produce a strong smell of camphor when rubbed.692 **Phenology:** Flowers in September, fruits in October.693 **Reproduction:** Only by seeds.694 **Distribution:** Karakalpakstan autonomous republic, Toshkent, Andijon, Farg'ona, Samarqand, Surxondaryo, and Buxoro
695 provinces of Uzbekistan; Chuy and Osh provinces of Kyrgyzstan.696 **Habitat:** The chul, adyr, and tau zones. On sandy soil, clay-soiled slopes with rocky debris, and in areas with soils containing
697 a wide diversity of minerals.698 **Population status:** Uncommon, found as small populations in *Artemisia*-ephemeral communities.699 **Traditional use:** Unknown.

700 **Documented effects:** The lactone leucomisine has strong anti-inflammatory action which is due to its antagonism of the
701 main inflammation mediators: histamine, serotonin (5-hydroxytryptamine), and prostaglandin F2 α and E2 (Kurmukov
702 1987). It reduces the volume of atherosclerotic aorta involvement in (tested) rabbits with hypercholesteremic atheroscle-
703 rosis (method of Anichkov and Holatova), reduces aorta wall permeability, has angioprotective action, and has medicinal
704 effects on experimental myocarditis (Aizikov et al. 1991; Kurmukova et al. 1997a, b; Kurmukov et al. 1991b; Prokhorova
705 et al. 1992a). Ascorbic acid strengthens the effects of leucomisine (Kurmukova and Aizikov 1997). *Oligvon*, a preparation
706 containing leucomisine, is used to prevent and treat artherosclerosis. The lactone austricine also has combined angiopro-
707 tective and hypolipidemic activity (Prokhorova et al. 1993; Aizikov et al. 1993a, b). A total lactone extract increased the
708 intensity of bile production and increased the concentration of cholesterol in the bile of normal rats, as well as in rats with
709 chemically induced hepatitis (Tursunova et al. 2002).

710 **Phytochemistry:** Leaves and inflorescences contain up to 1 % essential oils, which consist of up to 90 % levorotatory cam-
711 phor. The lactones leucomisine and austricine are obtained from the aboveground plant parts, as well as the sesquiterpe-
712 noids matricarin, anhydroaustricine, parishin B, parishin C, artelin, and artelein (Ribalko 1978; Tursunova et al. 2002).
713 The seeds contain lipids with epoxy-, monohydroxy-, and dihydroxyacids (Ul'chenko and Glushenkova 2001).



▲ **Artemisia absinthium L.** Photos: Andrei Lubchenko



◀ **Artemisia dracunculus L.**
Photos: Sasha Eisenman

▼ **Artemisia annua L.**
Photos: *left:* Kristian Peters;
right: Sasha Eisenman

▼ **Artemisia leucodes Schrenk**
Photo: Authors



715 **Artemisia scoparia Waldst. & Kit. – Asteraceae**716 **Synonyms:** *Oligosporius scoparia* (Waldst. & Kit.) Less.717 **English name:** Redstem wormwood718 **Russian name:** Полюнь метельчатая (Polyn' metyol' chataya)719 **Uzbek name:** Kizilburgan720 **Kyrgyz name:** Шыпыргы шыбак (Shyurgy shybak)

721 **Description:** Herbaceous annual or biennial plant with a thin vertical root. Stems single or few, 30–90 cm tall. Basal leaves
722 petiolate, bi- or tripinnatisect, segments linear-lanceolate, apex acute; middle cauline leaves smaller, sessile, segments
723 narrow, linear; upper cauline leaves deeply tri-lobed or entire. Inflorescences small capitula with 10–12 flowers, in a wide,
724 nodding panicle; involucral bracts brownish or pink-violet. Disc flowers yellow; ray flowers absent. Fruits achenes, ca.
725 0.6 mm long, ovate, flat with narrow ribs.

726 **Other distinguishing features:** Stems and leaves sparsely hairy or glabrous. Capitula subglobose.727 **Phenology:** Flowers in July, fruits in August-September.728 **Propagation:** By seeds.729 **Distribution:** Agricultural lands in all provinces of Kyrgyzstan and Uzbekistan.730 **Habitat:** Near and in cultivated and abandoned fields.731 **Population status:** Common, often forming dense groups.

732 **Traditional use:** Used to treat respiratory disease and rheumatism, and used as a diuretic (Gammerman et al. 1990). A tinc-
733 ture (2.5–10 %) of the plant is used to treat radiculitis. An infusion of the plant is used as a vermifuge, and to treat epilepsy
734 and irregularities in the menstruation cycle (Khodzhimatov 1989).

735 **Documented effects:** Aboveground parts are collected during the flowering stage for use as raw material. The plant is
736 slightly toxic. An infusion of the plants has diuretic properties. Its essential oil has laxative properties and is included in
737 the preparation *Artemisol*, which has antispasmodic action, increases the solubility of salts in urine and promotes the pas-
738 sage of kidney stones (Maksudov 1964). Intravenous administration of a hydro-methanolic extract of the plant produced
739 hypotensive and bradycardiac effects. Studies indicate that the plant contains Ca⁺⁺ channel-blocking constituents (Gilani
740 et al. 1994). The essential oil exhibited considerable inhibitory effects against a number of different bacteria (Cha et al.
741 2005).

742 **Phytochemistry:** Aboveground parts contain 4.35–5.57 % resins, with 1.08–1.37 % resins in roots and 7.91 % in flowers.
743 The whole plant contains organic acids (citric, malic, oxalic, acetic, propionic, and valerianic) and tannins (3.61–4.74 %
744 in aboveground parts and 2–2.5 % in the roots). The aboveground part contains essential oil of which the maximum accu-
745 mulation (0.96 %) happens during the flowering stage (Khodzhimatov 1989). The major components of the essential oils
746 are camphor, 1,8-cineole, and β -caryophyllene (Cha et al. 2005).

Artemisia viridis Willd. – Asteraceae	748
Synonyms: <i>Artemisia rupestris</i> ssp. <i>viridis</i> (Willd.) Ameljczenko, <i>Artemisia rupestris</i> var. <i>viridis</i> (Besser) A. DC.	749
English name: Wormwood	750
Russian name: По́лынь зелёная (Polyn' zelyonaya)	751
Uzbek name: Unknown	752
Kyrgyz name: Мамыр шыбак (Mamyr shybak)	753
Description: Herbaceous perennial. Stems 6–20 cm tall; vegetative stems short, densely leafy; flower-bearing stems wide, reddish-violet, rarely green, hairy. Leaves 1.5–5 cm long, 1–2.5 cm wide, wrinkled, hairy below; basal leaves with wide petioles, bipinnatisect, lower segments entire, mid- and upper segments with 3–5 narrow lobes. Inflorescences multiflorous capitula with ca. 70–80 flowers, arranged in racemes or spikes. Disc flowers reddish-brown; ray flowers absent. Fruits achenes, oblong-oviform, striated.	754 755 756 757 758
Other distinguishing features: Outer involucre bracts linear; internal involucre bracts triangular or elliptic.	759
Phenology: Flowers in July, fruits in September.	760
Reproduction: By seeds.	761
Distribution: High mountain regions of all provinces in Kyrgyzstan; not found in the flora of Uzbekistan.	762
Habitat: In high mountain steppes.	763
Population status: Common, forming dense groups.	764
Traditional use: An infusion of the herb is used to treat stomach ulcers, and diseases of the kidneys, liver, and bile ducts (Nanaeva 1960; Isakov 1969).	765 766
Documented effects: No data.	767
Phytochemistry: Aboveground parts contain essential oil, phenols, ketones, flavonoids, alkaloids, and coumarins (Plant Resources of the USSR 1993).	768 769

771 **Artemisia vulgaris L. – Asteraceae**772 **Synonyms:** *Artemisia coarctata* Forselles, *Artemisia opulenta* Pampanini.773 **English name:** Common mugwort, felon-herb, green-ginger, armoise vulgaire774 **Russian name:** Полынь обыкновенная (Polyn' obyknovennaya)775 **Uzbek name:** Oddiy erman776 **Kyrgyz name:** Кадимки куурай (Kadimki kuuray)

777 **Description:** Herbaceous perennial. Stems 30–180 cm tall. Basal leaves petiolate, 3–15 cm long, 1.5–12 cm wide, pinnatifid
778 or pinnatisect, segments lanceolate or linear; cauline leaves sessile, entire or pinnatisect. Inflorescences capitula in com-
779 pact racemiform or paniculiform clusters; involucral bracts hairy; outer bracts oblong; inner bracts elliptical. Disc flower
780 corollas brownish. Fruits ellipsoid achenes, grayish-brown, glabrous.

781 **Other distinguishing features:** Leaves densely white-haired beneath, usually glabrous above.782 **Phenology:** Flowers in July, fruits in September.783 **Reproduction:** By seeds.784 **Distribution:** In agricultural lands in all provinces of Kyrgyzstan and Uzbekistan.785 **Habitat:** From valleys to mid-belt of mountains. Near canals and in vegetable gardens, orchards and waste grounds.786 **Population status:** Common, forming dense groups.

787 **Traditional use:** Leafy tops and roots are used for medicinal purposes. Leaves are collected from plants during the flowering
788 stage and the roots are collected in the autumn. Avicenna used the herb in baths to treat kidney stones and uterine ulcers,
789 and to induce menstruation. A decoction of the herb is used to treat sinus colds (Khalmatov et al. 1984), nervous diseases,
790 epilepsy, and neurasthenia, and is also used as an anticonvulsant. The aboveground parts are used to treat poisoning,
791 inflammation of the gastrointestinal tract, tuberculosis, and to increase the appetite. It is also used externally as a lotion to
792 treat ulcers and persistent wounds (Maznev 2004).

793 **Documented effects:** Data suggest that aqueous and chloroform extracts from leaves of *A. vulgaris* have antihypertensive
794 actions (Tigno et al. 2000). Essential oils showed a broad spectrum of antimicrobial activity (Blagojevic et al. 2006). Two
795 flavonoids, eriodictyol and apigenin, found in *A. vulgaris*, exhibited estrogenic effects in vitro (Lee et al. 1998).

796 **Phytochemistry:** The herb contains essential oils (contains cineol, thujone, borneol, and aldehydes), flavonoids, alkaloids,
797 carotene, and ascorbic acid (Khalmatov et al. 1984; Lee et al. 1998).

Arum korolkowii Regel – Araceae	799
Synonyms: <i>Arum elongatum</i> Steven, <i>Biarum sewertzowii</i> Regel.	800
English name: Korolkov's arum	801
Russian name: Аронник Королькова (Aronnik Korol'kova)	802
Uzbek name: Kuchala, Chayon ut	803
Kyrgyz name: Корольков аруму (Korol'kov arumu)	804
Description: Perennial herb, 30–50 cm high, with a flat-spherical tuber that is 3–4 cm in diameter. Base of leaf petiole sheathing, petiole short to twice as long as the blade. Leaf blade cordate, acuminate (spear-shaped) or triangular. Inflorescence a spadix; peduncle longer than leaf petioles, 50–60 cm long with reddish stripes; spathe exterior green, white inside, elongate-lanceolate, narrow-cylindrical, almost 2 times longer than spadix, apex acute. Fruits red berries.	805 806 807 808
Other distinguishing features: Fruits are densely clustered on spadix.	809
Phenology: Flowers and fruits in May-June.	810
Reproduction: By seeds.	811
Distribution: All of Uzbekistan; Jalal-Abad province of Kyrgyzstan.	812
Habitat: The adyr and tau zones. Shady, wet places, in gorges, among rocks.	813
Population status: Not common, found as single individuals.	814
Traditional use: The powdered tuber is used to treat scorpion and poisonous snake bites and is mixed with honey to treat fungal skin diseases and white spots on the skin of the neck. Bread made with tuber powder and sesame oil is prescribed (to be eaten) to treat hemorrhoids (Khalmatov 1964).	815 816 817
Documented effects: All parts of the fresh plant are poisonous due the presence of saponins (Khalmatov 1964).	818
Phytochemistry: Tubers contain poisonous saponins, which produce hydrocyanic acid as a result of hydrolysis, alkaloids (possibly volatile cicutine), lipids, pectic substances, fructosans and 28–30 starch. A carotenoid, lycopene, was found in the fruits (Khalmatov 1964; Chernenko et al. 2000).	819 820 821



▲ *Artemisia scoparia* Waldst. & Kit. Photos: Andrei Lubchenko



▲ *Artemisia viridis* Willd.

Photos: Vladimir Epiktetov

▶ *Artemisia vulgaris* L.

Photos: Dmitri Oreshkin

▼ *Arum korolkowii* Regel

Photos: *left*: Evgeny Davkaev

right: Alexander Naumenko



Asparagus persicus Baker – Asparagaceae	823
Synonyms: <i>Asparagus inderiensis</i> Blume ex Ledeb., <i>A. ledebourii</i> Mishchenko.	824
English name: Persian asparagus	825
Russian name: Спаржа персидская (<i>Sparzha persidskaya</i>)	826
Uzbek name: Томirdori	827
Kyrgyz name: Персия спаржасы (<i>Persiya sparzhasy</i>)	828
Description: Perennial herb. Stems 60–120 cm high, smooth, glabrous, branched; branch angles at 90° or obtuse to the stem.	829
Cladodes 1–8 per cluster, usually 1.5–2 cm long and unequal in length, glabrous, smooth; upper and middle leaves scale-like with a sharp spur. Flowers arise from the stems and branches; female flowers 3 mm long, semispherical, campanulate, greenish-white; male flowers campanulate, 5–6 mm long. Fruit a red berry, spherical, 6–7 mm wide; on a long pedicel up to 2 cm in length.	830 831 832 833
Other distinguishing features: Stems often winding, curling, or trailing.	834
Phenology: Flowers in May, fruits in June.	835
Reproduction: By seeds and rhizomes.	836
Distribution: Toshkent, Farg'ona, and Samarqand provinces of Uzbekistan; Naryn, Ysyk-Kol and Chuy provinces of Kyrgyzstan.	837 838
Habitat: The tau zone. Among tall grasses and in the tree-shrub belt of mountains.	839
Population status: Uncommon, found as single individuals.	840
Traditional use: Of the wild species of <i>Asparagus</i> found in Uzbekistan, this is the only species used in folk medicine. In some regions of Toshkent province (Uzbekistan) it is used to treat numerous diseases (Khalmatov 1964).	841 842
Documented effects: No data.	843
Phytochemistry: All the species of <i>Asparagus</i> found in Uzbekistan contain alkaloids, essential oils, vitamins, asparagine, saponins, steroid saponins and related substances. The seeds contain fatty oils (Khalmatov 1964; Tairov 1969).	844 845

846 **Astragalus sieversianus Pall. – Fabaceae**847 **Synonyms:** *Lithoon sieversianum* (Pall.) Nevski.848 **English name:** Unknown849 **Russian name:** Астрагал Сиверса (Astragal Siversa)850 **Uzbek name:** Pakhtak851 **Kyrgyz name:** Тулку куурай (Tulku kuuray)

852 **Description:** Herbaceous perennial. Stems 60–150 cm tall, up to 2 cm in diameter, densely hairy. Leaves alternate, pinnate,
853 15–30 cm long, long-stipulate; leaflets in 8–12 pairs, from narrow-ovate to elliptic, densely hairy on undersides, margins
854 entire. Inflorescence axillary racemes with 3–9 flowers. Calyx tubular, densely hairy. Corolla papilionaceous, pale-yellow.
855 Fruits ovate-spherical legumes, 15–20 mm long, densely covered with long, entangled hairs. Seeds kidney-shaped,
856 brown.

857 **Other distinguishing features:** Stamens 10, nine filaments are fused. Legumes ovate-spherical, very hairy.858 **Phenology:** Flowers in May-June, fruits in July-August.859 **Reproduction:** By seeds.

860 **Distribution:** Farg'ona, Toshkent, Samarqand, Navoiy, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Chuy,
861 Talas, Osh and Jalal-Abad provinces of Kyrgyzstan.

862 **Habitat:** The adyr and tau zones. Stony slopes in the mountain-steppe belt and lower belt of juniper stands.863 **Population status:** Common.

864 **Traditional use:** A decoction of the fruits is drunk to remove kidney and bladder stones. The seeds, taken internally, are
865 recommended to treat hernias in children and are smoked to treat syphilis (Khalmatov 1964).

866 **Documented effects:** In experiments, the ethanol extracts of flowers showed high antioxidant, lipid-reducing, and anti-
867 aggregating activities. Preparations of the plant have sedative, antibacterial, anti-inflammatory, and bile-stimulating
868 actions. Intravenous introduction of the alkaloid smirnovine in narcotized animals, at the dose of 2 mg/kg, reduced blood
869 pressure by 32–56 % for a short time and excited breathing, which is apparently due to ganglio-blocking actions
870 (Sadritdinov and Kurmukov 1980). Saponins extracted from the roots protected the liver from induced chemical injury in
871 mice (Zhang et al. 1992).

872 **Phytochemistry:** The aboveground parts contain saponins, up to 0.1 % alkaloids (especially smirnovine), coumarins, tan-
873 nins, flavonoids (0.9 % in stems, up to 4.9 % in leaves), vitamins C, E and P, and carotene. The roots contain triterpenoids,
874 alkaloids, coumarins, and saponins (Khalmatov 1964; Svechnikova et al. 1983; Gan et al. 1986a, b).

875

Atraphaxis spinosa L. – Polygonaceae	876
Synonyms: <i>Tragopyrum spinosum</i> (L.) C. Presl.	877
English name: Goat's wheat, manna	878
Russian name: Курчавка колючая (Kurchavka kolyuchaya)	879
Uzbek name: Tuya singren	880
Kyrgyz name: Боз караган (Boz karagan)	881
Description: Woody shrub, 30–100 cm tall. Branches long, slender, apex leafless, spine-tipped. Leaves alternate, sessile or short-petiolate, 3–7 mm long, 2–5 mm wide, elliptic to ovate, coriaceous, glabrous, margins entire. Ocreae cylindric, 1–3 mm long, membranous, brown at base. Inflorescences 2–6-flowered clusters, occurring in leaf axils of current year's branchlets; pedicels ~5 mm. Tepals 4, pink. Fruits lenticularly compressed nutlets, light brown, smooth, shiny.	882 883 884 885
Other distinguishing features: Stamens 6, styles 2.	886
Phenology: Flowers and fruits in May-June.	887
Reproduction: By seeds.	888
Distribution: Karakalpakstan autonomous republic, Buxoro province, and probably other provinces of Uzbekistan; Naryn and Ysyk-Kol provinces of Kyrgyzstan.	889 890
Habitat: The tau zone. Stony slopes in lower mountain areas.	891
Population status: Common.	892
Traditional use: A decoction from the leaves and flowers are used in folk medicine to treat fever (Khalmatov 1964).	893
Documented effects: In animal studies, the total alkaloids isolated from leaves increased blood pressure (Khalmatov 1964).	894 895
Phytochemistry: The leaves contain alkaloids and tannins (Khalmatov 1964). Flavonoids has also been isolated from the plant (Chumbalov et al. 1970, 1971).	896 897

899 **Berberis integerrima Bunge – Berberidaceae**900 **Synonyms:** Some consider this species synonymous with *Berberis oblonga* (Regel) Schneid.901 **English name:** Unknown902 **Russian name:** Барбарис цельнокрайний (Barbaris tsel'noкраyniy)903 **Uzbek name:** Kizil zirk904 **Kyrgyz name:** Бёру карагат (Byoru karagat)

905 **Description:** Branched shrub, up to 4 m. Oldest branches gray, young branches reddish brown, glabrous. Branches armed
906 with 3- to 5-fid spines; spines straight, ca. 3 cm long. Leaves clustered on short shoots, petiolate, 3–3.5 cm long, 1.5–
907 1.7 cm wide, coriaceous, obovate or elongate, margins mostly entire. Inflorescences racemiform, 6–10 cm long, axillary.
908 Flowers ca. 1 cm in diameter; pedicel ca. 1 cm long. Sepals similar to 6 yellow petals. Style very short. Fruit an elongated
909 berry, purple-red, gray-glaucous, 7–8 mm long. Seeds elongated, dark brown.

910 **Other distinguishing features:** 10–12 berries per raceme. Differs from *Berberis nummularia*, which has red fruits when
911 fully ripe.

912 **Phenology:** Flowers in May-June, fruits in July-August.913 **Reproduction:** By seeds.914 **Distribution:** Tashkent, Farg'ona, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.

915 **Habitat:** The tau zone. Usually grows as a single shrub along mountain river banks, sometimes growing in small populations
916 at 1,500–1,600 m elevation.

917 **Population status:** In Uzbekistan the prevalence of this species is decreasing due to human disturbance of its natural
918 habitat.

919 **Traditional use:** Fruits are used as an antipyretic, to relieve thirst and as a spice (Khalmatov 1964). In northern Tajikistan
920 the roots are used to treat wounds, bone fractures, rheumatism, radiculitis, heart pain, and stomach aches. A decoction of
921 the leaves is used to treat kidney stones. A tea made with the flowers is used to treat lung tuberculosis, chest pains, and
922 headaches. An infusion of the fruits is used to treat constipation and wounds (Khodzhimatov 1989).

923 **Documented effects:** In experiments with animals, the alkaloid berberine lowers blood pressure, has minor ganglion block-
924 ing action, stimulates contractility of smooth muscles of the uterine horns and intestines, depresses central nervous sys-
925 tem, prevents tumors, and has a pronounced choleric action (Supek 1946; Selivanova 1954; Shvarev and Tsetlin 1972;
926 Idzumi and Conti 1962; Conti 1962). In medical practice, a preparation with berberine is prescribed to treat chronic chole-
927 cystitis. Berberine has antitumor and bacteriostatic activity, increases phagocytic activity of leucocytes, and prevents
928 animal death from septicemia. It also is effective for patients with initial pulpitis. In vitro, berberine has bactericidal action
929 against *Vibrios cholerae* (Turova et al. 1984). An extract prepared from the dried berries protected rat hepatocytes against
930 induced cytotoxicity in vitro. In vivo, pretreatment and treatment of animals with the extract protected the liver against
931 induced injuries (Jamshidzadeh and Niknahad 2006).

932 **Phytochemistry:** The plant contains many alkaloids (including berberine, columbamine, jatrorrhizine and oxyacanthine)
933 and organic acids. Leaves from plants at the fruit-bearing stage from the Chon-Kemin valley in Kyrgyzstan, contained
934 0.18 % total alkaloids and the young shoots contained 1.5 % total alkaloids. Berberine, berbamine, berbaminine, isobol-
935 dine, isocorydine, isotetrandrine, oxyacanthine, magnoflorine, palmatine, talicmidine, reticuline and others were isolated
936 from the total alkaloids (Karimov et al. 1977; Yunusov 1981; Karimov et al. 1993a, b; Khamidov et al. 1996).

Berberis oblonga (Regel) Schneid. – Berberidaceae	938
Synonyms: Some consider this species synonymous with <i>Beberis integerrima</i> Bunge.	939
English name: Unknown	940
Russian name: Барбарис продолговатый (Barbaris prodolgovatyy)	941
Uzbek name: Zirk, Kora zirk, Kora qand	942
Kyrgyz name: Созунку бёру карагат (Sozunku byoru karagat)	943
Description: Branched shrub, up to 4 m tall. Older branches dark, bark with long, shallow cracks; younger branches reddish-brown, often grayish with simple or 3-branched spines, spines ca. 1.5 cm long. Leaves clustered on short shoots in groups of 5–7, up to 6 cm long and 3 cm wide, wide-elliptic or obovate, narrow cuneate, glabrous, margins usually entire, occasionally with short, spiny teeth. Inflorescences usually racemiform, 3–4.5 cm long, with 10–30 flowers in each cluster. Flowers up to 1 cm in diameter with 6 yellow petals. Fruits ellipsoid berries, up to 1 cm long, 6 mm wide, black-purple, gray-glaucous. Seeds 2, rarely 1, dark brown.	944 945 946 947 948 949
Other distinguishing features: Young plants have 5–11-branched spines. The bark on branches and roots is bright-yellow inside.	950 951
Phenology: Flowers in May, fruits in July-August.	952
Reproduction: By seeds.	953
Distribution: Tashkent, Namangan, Andijon, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.	954 955
Habitat: The tau zone. Rocky slopes of medium and, sometimes, lower mountain zones.	956
Population status: Common, sometimes forming dense groups.	957
Traditional use: In folk medicine, the fruits are used as a heart tonic, to treat neurasthenia, as an antipyretic, to relieve thirst and as an antidiarrheal remedy. The root decoction is used to treat rheumatism, fever, eye diseases, and as oral rinse for wounds of the mouth (Khalmatov 1964). The residue from a dehydrated water extract of the root is eaten, mixed with hot water and drunk, or applied to a cloth and applied externally, to treat jaundice, stomach aches, back pain and arthralgia (Sezik et al. 2004; Pak 2005).	958 959 960 961 962
Documented effects: Giving an infusion of the plant to laboratory animals resulted in cardiogenic action and a mild decrease of blood pressure. In experiments with dogs the preparation stimulated blood coagulation (Ibragimov and Dzhumabaev 1971; Dzhumabaev 1972).	963 964 965
Phytochemistry: Roots from plants collected in Kyrgyzstan (Arslanbob) at the end of the growing season contained 4.5 % total alkaloids. Young shoots collected from flowering plants in Uzbekistan (Chingan) contained 1 % alkaloids, and the leaves contained 0.01 % total alkaloids. Berberine, berbamine, berbaminine, glaucine, isocorydine, columbamine, magnoflorine, oblongine, oxyacanthine, palmatine, thalimidine, and others have been isolated from the total alkaloids (Karimov et al. 1976, 1977; Yunusov 1981; Khamidov et al. 2003).	966 967 968 969 970



◀ **Asparagus persicus**
Baker
Photos: Alim Gaziev

▼ **Astragalus sieversianus**
Pall.
Photos: Evgeny Davkaev



▼ **Berberis integerrima** Bunge Photos: *top and lower left:* Vladimir Epiktetov; *lower right:* Alim Gaziev

▼ **Atraphaxis spinosa** L.
Photos: Avinoam Danin



Betonica foliosa Rupr.– Lamiaceae	971
Synonyms: <i>Stachys betoniciflora</i> Rupr., <i>Stachys betonicifolia</i> Regel, <i>Stachys foliosa</i> Regel.	972
English name: Unknown	973
Russian name: Буквица олиственная, Чистец буквицветный (Bukvitsa olistvennaya, Chistets bukvitsetsvetnyy)	974
Uzbek name: Тог кудуси	975
Kyrgyz name: Жалбырактуу бетоника (Zhalbyraktuu betonika)	976
Description: Herbaceous perennial with short rhizomes. Stems 60–100 cm tall, 4-sided, densely hairy below, scattered hairy above. Leaves opposite; lower leaves petiolate, 13–15 cm long, 4–5 cm wide, obovate, bases oblique, margins crenate; upper leaves sessile, ovate-lanceolate, 5–6 cm long, 2–3 cm wide, margins serrate; terminal leaves lanceolate, entire. Inflorescences 10–12-flowered verticillasters, in terminal, compact spikes. Flowers sessile. Calyx 10–15 mm long, campanulate with lanceolate teeth. Corolla 2-lipped, lilac. Fruits dark-brown nutlets.	977 978 979 980 981
Other distinguishing features: Fruits 3-sided with longitudinal grooves.	982
Phenology: Flowers in June-August, fruits in August and September.	983
Propagation: By seeds.	984
Distribution: All provinces of Kyrgyzstan; Toshkent province of Uzbekistan.	985
Habitat: Among shrubs and juniper stands, in steppes, forests, and in high mountain meadows.	986
Population status: Common, forming dense groups.	987
Traditional use: The aboveground parts are used to treat hysteria, hypertension, epilepsy, fainting, gout, jaundice, and rheumatism. A tea made from the herb is used to treat gastrointestinal pain, hemoptysis, respiratory disease, inflammation of the kidneys, and bladder, and is also used as a sedative. An infusion of the roots is used as a laxative (Khodzhimatov 1989).	988 989 990 991
Documented effects: A tincture and liquid extract of this species is used in obstetric-gynecological practices as a treatment to increase uterine muscle tonus, increase uterine contractions, and as a hemostatic (Tolmachev 1976). Preparations of the plant have anti-inflammatory, anti-asthmatic, antiseptic, analgetic, hemostatic, and choleric properties. The preparations are used as expectorants, to decrease blood pressure, to increase metabolism, to improve blood circulation and to improve digestion (Plant Resources of the USSR 1991).	992 993 994 995 996
Phytochemistry: Plants collected in Talas-Alatau (Kyrgyzstan) contained ~54 mg/% vitamin C (in the leaves). The aboveground parts contained 1.54 % flavonoids, alkaloids (up to 0.49 % stachydrine), 1 % iridoids, 3.11 % resins, 0.12 % essential oil, ~49.5 mg/% vitamin C, 2 % organic acids, 1.02 % calcium salts, 3.98 % sugars, phenolcarbonic acids, and vitamin k ₁ (Khalmatov et al. 1984).	997 998 999 1000

1002 **Bidens tripartita L. – Asteraceae**1003 **Synonyms:** *Bidens comosa* (Gray) Wieg., *Bidens orientalis* Velen.1004 **English name:** Threelobe beggarticks1005 **Russian name:** Череда трехраздельная (Chereda trekhrazdel'naya)1006 **Uzbek name:** Eteetkanak, Karakeez1007 **Kyrgyz name:** Уч болуктуу ит уйчан (Uch boluktuu it uychan)1008 **Description:** Herbaceous annual with a taproot. Stems 20–110 cm tall. Leaves opposite, lower and middle tripartite; upper
1009 leaves unlobed, lanceolate. Inflorescences capitula, single or in groups of 2–3; involucral bracts ovate or lanceolate-ovate,
1010 internal bracts shorter. Flowers yellow, usually only disc type. Fruits dark brown achenes, flattened with 4 edges, often
1011 with retrorsely barbed awns.1012 **Other distinguishing features:** 1–5 ray flowers occasionally present. Fruits usually not tuberculate.1013 **Phenology:** Flowers in June-September, fruits in July-October.1014 **Reproduction:** By seeds.1015 **Distribution:** All provinces of Kyrgyzstan and Uzbekistan.1016 **Habitat:** In wet places near ponds, lakes, bogs and canals.1017 **Population status:** Common, sometimes forming dense groups.1018 **Traditional use:** The top parts of the plant, with leaves and immature capitula, are collected for use in traditional medicine.
1019 They are used to treat respiratory diseases, scrofula, scurvy, scabies, bacterial and fungal skin diseases, poor digestion,
1020 toothaches, blood diseases (including anemia), arteriosclerosis, anthrax, and tuberculosis, and also to regulate the metabo-
1021 lism (Maznev 2004).1022 **Documented effects:** Used as an antipyretic, as a diuretic for treatment of urogenital diseases, and as a diaphoretic and anti-
1023 inflammatory (Kurochkin 1998). A bath infused with the herb is used to treat diathesis in children. An infusion of the herb
1024 is drunk to induce sweating and to treat common colds (Grinkevitch 1991). Although the content of flavonoids in the
1025 flower heads was found to be half of that found in the herb, an extract from flowers had nearly 2 times higher antioxidant
1026 activity (Wolniak et al. 2007).1027 **Phytochemistry:** The herb contains flavonoids, coumarins, ascorbic acid, carotene, tannins, mucilage, γ -lactones, and traces
1028 of essential oils (Serbin et al. 1972 a, b, c, 1975; Khalmatov et al. 1984; Maznev 2004; Wolniak et al. 2007).

1029

Biebersteinia multifida DC. – Biebersteiniaceae	1030
Synonyms: None	1031
English name: Unknown	1032
Russian name: Биберштейния многораздельная (Bibershteyniya mnogorazdel'naya)	1033
Uzbek name: Kontepar	1034
Kyrgyz name: Кёп балуктуу биеберштения (Куор baluktuu biyebershteniya)	1035
Description: Herbaceous perennial with a thick, tuberiform root. Stem 30–60 cm tall, sturdy. Plant woolly-hairy and glandular. Leaves alternate, stipulate, short-petiolate, 10–20 cm long, 2–8 cm wide, tripinnatisect, both sides spreading-hairy. Inflorescences racemiform. Flowers orange-yellow. Sepals 5. Petals 5. Stamens 10; filaments glabrous, connate at the base, forming a ring. Styles 5, connate; stigma capitate. Fruit a schizocarp. Seeds very wrinkled, coriaceous.	1036 1037 1038 1039
Other distinguishing features: Root turns pink when fractured.	1040
Phenology: Flowers and fruits in April-June.	1041
Reproduction: By seeds.	1042
Distribution: Karakalpakstan autonomous republic, Toshkent, Samarqand, Buxoro, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Chuy, Talas and Osh provinces of Kyrgyzstan.	1043 1044
Habitat: The tau zone. Gypsum soil on low mountains. Shallow-soiled and stony slopes of lower and medium mountain zones.	1045 1046
Population status: Uncommon.	1047
Traditional use: A decoction of the root is used as a hemostatic for post-natal bleeding and to treat gastric diseases (Khalmatov 1964). In Iran an ointment made of the powdered root mixed with tallow is used to treat musculoskeletal disorders and bone fractures (Farsam et al. 2000).	1048 1049 1050
Documented effects: An extract of the root had anti-inflammatory effects on induced rat paw edema and analgesic effects in tests with rats (Farsam et al. 2000). In parenteral toxicity tests in mice, an extract of the total alkaloids was classified as a moderately toxic agent. Dermal acute toxicity tests showed no sign of toxicity (Ostad et al. 2003).	1051 1052 1053
Phytochemistry: The aboveground parts have essential oils. The roots contain tannins, carbohydrates, and saponins. 0.08 % total alkaloids were obtained from the extracts of tops of plants collected at the Usturt Plateau during budding stage and vasicinone was isolated (Yunusov 1981). The roots and aboveground parts contain polysaccharides (Arifkhodzhaev et al. 1985; Arifkhodzhaev and Rakhimov 1986, 1993) as well as the flavones luteolin 7-glucoside and 7-rutinoside (Omurkamzinova et al. 1991).	1054 1055 1056 1057 1058

1060 **Bunium chaerophylloides (Regel & Schmalh.) Drude – Apiaceae**

1061 **Synonyms:** *Buniella chaerophylloides* (Regel & Schmalh.) Schischk., *Carum chaerophylloides* Regel & Schmalh., *Carum*
1062 *confusum* O. Fedtsch., *Carum sogdianum* Lipsky.

1063 **English name:** Unknown

1064 **Russian name:** Буниум бутеневый (Bunium butenevyy)

1065 **Uzbek name:** Qarga oyeq

1066 **Kyrgyz name:** Бутен зиреси (Buten ziresi)

1067 **Description:** Herbaceous, glabrous perennial, with a spherical tuber up to 2 cm in diameter. Stem 50–70 cm high, narrowly
1068 striated, cylindrical, hollow. Basal leaves long-petiolate, blade triangular-oval, tri-pinnatisect, with lanceolate lobes; upper
1069 leaves alternate, sessile or with short membranous, sheathing petioles. Inflorescences compound umbels with 10–16 radi-
1070 als, flat-topped. Calyx toothless. Petals broadly obovate, white. Fruits oblong-linear schizocarps, 4–4.5 cm long, some-
1071 times curved.

1072 **Other distinguishing features:** Bractlets lacking. The fruits are similar to those of *Bunium persicum*, but do not smell when
1073 crushed.

1074 **Phenology:** Flowers in April- May, fruits in June.

1075 **Reproduction:** By seeds.

1076 **Distribution:** Toshkent, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh province of
1077 Kyrgyzstan.

1078 **Habitat:** The tau zone. Mountainous slopes.

1079 **Population status:** Common.

1080 **Traditional use:** A powder of roasted and ground fruits mixed with honey is used in folk medicine to dissolve renal and
1081 cystic stones and to treat skin diseases (for white spots on the skin). The powdered tuber is applied to mouth injuries and
1082 for reddening of the tongue (Khalmatov 1964).

1083 **Documented effects:** No data.

1084 **Phytochemistry:** Fruits contain essential oil and roots contain around 24 % starch (Khalmatov 1964).

1085



▲ *Betonica foliosa* Rupr. Photos: Alexander Naumenko



▲ *Biebersteinia multifida* DC. Photos: *top right*: Vladimir Epiktetov; *bottom*: Evgeny Davkaev

▼ *Bunium chaerophylloides* (Regel & Schmalh.) Drude Photo: Komiljon Tojibaev



► *Bidens tripartita* L.
Photos: Dmitri Oreshkin



1086 **Bunium persicum (Boiss.) B. Fedtsch. – Apiaceae**1087 **Synonyms:** *Carum persicum* Boiss., *Carum heterophyllum* Regel & Schmalh.1088 **English name:** Black cumin, wild cumin1089 **Russian name:** Буниум персидский (*Bunium persidskiy*)1090 **Uzbek name:** Zira1091 **Kyrgyz name:** Персия зиреси (*Persiya ziresi*)

1092 **Description:** Herbaceous perennial with an irregular spherical tuber. Stem 40–60 cm high, striated, pale green, glabrous,
1093 branching from the middle to upper portion. Basal leaves with long petioles, blades wide-triangular, bi- or tripinnatisect;
1094 cauline leaves alternate, bipinnatisect with highly dissected sections, sessile, sheathing. Inflorescence a compound umbel
1095 with 15–20 rays. Petals white, ca. 1 mm long. Fruits oblong schizocarps, 3–4 mm long, dark brown, ridged, shorter than
1096 the pedicel.

1097 **Other distinguishing features:** Involucel with 2–5 linear bractlets. The ripe fruits have a very specific smell, unique to
1098 *Bunium persicum*.

1099 **Phenology:** Flowers in June, fruits in July.1100 **Reproduction:** Only by seeds.1101 **Distribution:** Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh province of Kyrgyzstan.1102 **Habitat:** The adyr zone. Slopes of loess foothills.1103 **Population status:** Due to mass collection of seeds, it now occurs infrequently.

1104 **Traditional use:** Seeds are used to prevent stomach aches and to eliminate spleen tumors. An infusion of the fruits in vinegar
1105 is used as a hemostatic to stop nose bleeds. Roasted fruits are recommended as a diuretic (Khalmatov 1964). The fruits
1106 are taken to increase appetite, and to treat kidney stones and liver diseases. A decoction and infusion is used as a diapho-
1107 retic and vermifuge, as well as to improve digestion. Roasted fruits are used to treat bladder incontinence and obesity. The
1108 fresh tubers are used to improve digestion (Khodzhimatov 1989).

1109 **Documented effects:** Modern research has shown that a decoction of the fruits stimulates gastric secretion, favors creation of
1110 complete gastric fluids with a higher acid index of pepsin and pepsinogen and also has evident cholagogic, anti-inflammatory,
1111 and antispasmodic actions. It improves disinfectant and secretory functions of the liver. Decoctions of *Bunium persicum* are
1112 recommended to treat chronic hypo- and anacidity gastritis, chronic colitis and cholecystitis. Roasted fruits have a diuretic
1113 action (Khalmatov and Kosimov 1994). Extracts of the aboveground plant parts exhibited significant antibacterial effects
1114 against *Bacillus subtilis*, as well as antifungal activity and low cytotoxicity (Sardari et al. 1998; Bazzaz and Haririzadeh
1115 2003). The essential oils has antioxidant and anti-inflammatory properties (Jassbi et al. 2005).

1116 **Phytochemistry:** Seeds contain up to 3 % essential oils (carene, cymol, terpinolene, carvone, linalool, carvacrol) and 13.6 %
1117 oils and proteins (Khalmatov and Kosimov 1994). Essential oil collected from plants in Iran contained mostly monoter-
1118 penes and phenylpropanoids, such as α -pinene, p-cymene, limonene, γ -terpinene, cuminaldehyde, cuminyl alcohol,
1119 myristicin, and dillapiole (Jassbi et al. 2005).

Campanula glomerata L. – Campanulaceae	1120
Synonyms: None	1121
English name: Clustered bell flower, Dane's blood	1122
Russian name: Колокольчик скученный (Kolokol'chik skuchenny)	1123
Uzbek name: Kungrok gul	1124
Kyrgyz name: Топтолгон конгуроо гул (Toptolgon konguroo gul)	1125
Description: Perennial herb, gray-hairy. Stems 25–70 cm high, erect, slightly angled. Leaves alternate, simple, slightly toothed; lower leaves oblanceolate, apex acute, long-petiolate; upper leaves ovate to narrowly triangular, sessile, sometimes clasping. Flowers sessile in terminal compact clusters or few in the upper leaf axils. Sepals narrowly lanceolate. Corolla campanulate, 5-lobed with ovate-triangular acute lobes, lilac to blue-violet, 1–3 cm long, hairy outside. Fruit a capsule opening by lateral pores.	1126 1127 1128 1129 1130
Other distinguishing features: Stems slightly angled.	1131
Phenology: Flowers and fruits in June-August.	1132
Reproduction: By seeds.	1133
Distribution: Toshkent, Samarqand, and Farg'ona provinces of Uzbekistan; all of Kyrgyzstan.	1134
Habitat: The tau zone. Shallow-soiled and stony slopes.	1135
Population status: Uncommon, found as single individuals.	1136
Traditional use: An infusion and decoction of the leaves are used as an oral rinse, to treat sore throat and hoarseness and are applied externally as a lotion for erysipelatos inflammations and taken internally to treat headache. A decoction of the flowering herbs is also used to treat hydrophobia, for bathing children with a fear of water and for treating people who have seizures (Khalmatov 1964).	1137 1138 1139 1140
Documented effects: An ethanolic extract of the roots has hypolipidemic and antioxidant properties (Eliseeva 2005).	1141
Phytochemistry: The plant contains small amounts of alkaloids. The leaves contain up to 1,000 mg% of vitamin C (Khalmatov 1964).	1142 1143

1145 **Capparis spinosa L. – Capparidaceae**1146 **Synonyms:** *Capparis herbacea* Willd.1147 **English name:** Caper bush1148 **Russian name:** Каперцы колючие (Kapertsy kolyuchiye)1149 **Uzbek name:** Kovul1150 **Kyrgyz name:** Тикендуу коңуз баш (Tikenduu konuz bash)

1151 **Description:** Herbaceous perennial plant with thick roots. Stems numerous, decumbent, up to 2.5 m long, woody towards
1152 the base, glabrous. Leaves alternate, ovate, obovate, or round, 3.5–6 cm long, glabrous, short-petiolate with stipular
1153 spines. Flowers single, 5–8 cm in diameter, white or sometimes cream to pinkish in color, with long pedicels. Fruits fleshy,
1154 berry-like capsules, round to obovate, 2.5–5 cm long, green, glabrous, smooth. Seeds 3–3.5 mm long, round-elliptic or
1155 kidney-shaped, brown.

1156 **Other distinguishing features:** Flowers have strong aroma similar to honey.1157 **Phenology:** Flowers in May-June, fruits in July-August.1158 **Reproduction:** By seeds.1159 **Distribution:** All of Uzbekistan; Chuy, Talas, Osh and Jalal-Abad provinces of Kyrgyzstan.

1160 **Habitat:** The chul and adyr zones. A weed and ruderal; found on hills, among unirrigated winter wheat fields, along roads,
1161 in dry river-beds and on stony slopes of the lower mountains.

1162 **Population status:** Common, forms large populations.

1163 **Traditional use:** One of the oldest folk remedies and used to treat a variety of diseases. Avicenna recommended caper bush
1164 as an analgesic and vermifuge, for healing wounds, and to treat asthma and gastrointestinal diseases. A decoction of the
1165 roots is used to treat hepatitis, and the root bark is smoked to treat syphilis. Juice from the flowers are used as a treatment
1166 for scrofula. A decoction of the fruit is used for hemorrhoids and toothaches, and to strengthen the gums (Akopov
1167 1981).

1168 **Documented effects:** 25 % root extract in 96 % ethanol and 25 % root decoction accelerate blood coagulation (Akopov
1169 1981). A tincture of the root increased the number of thrombocytes in blood. In experiments with guinea pigs, treatment
1170 with a decoction of the roots caused desensitization to animal and plant allergens. The fresh juice from the fruits was clini-
1171 cally tested and recommended for the treatment of exophthalmic goiters (Khodzhimatov 1989). A methanol extract of the
1172 flower buds exhibited strong antioxidant activity (Germano et al. 2002). p-Methoxy benzoic acid isolated from an extract
1173 of this plant species was found to possess significant activity against induced hepatotoxicity in vivo and in vitro (Gadgoli
1174 and Mishra 1999).

1175 **Phytochemistry:** Flowers and buds contain rutin, quercetin, vitamin C, saponins, pigments and glycosides. Seeds contain
1176 25–35 % semi-drying oils, 25 % oleic and 33 % linoleic acids. The aboveground plant parts contain 0.32 % rutin and
1177 quercetin, up to 100 mg% vitamin C, stachydrine, and thioglycosides. Fruits contain up to 36 % sugar, 25–35%mg vita-
1178 min C, flavonoids, and thioglycosides. Roots contain 1.2 % alkaloids (stachydrine), 0.44 % flavonoids, 4.5 % sugars,
1179 coumarins, and other substances (Khalmatov and Kosimov 1994).

- Capsella bursa-pastoris (L.) Medik. – Brassicaceae** 1181
- Synonyms:** *Bursa bursa-pastoris* (L.) Britton, *Bursa pastoris* Weber ex F.H. Wigg., *Capsella hyrcana* Grossh., *Crucifera capsella* E.H.L. Krause, *Iberis bursa-pastoris* (L.) Crantz, *Thlapsi bursa-pastoris* L. 1182
1183
- English name:** Shepherd's purse 1184
- Russian name:** Пастушья сумка обыкновенная (Pastush'ya sumka obyknovennaya) 1185
- Uzbek name:** Ochambiti, zhag-zhag 1186
- Kyrgyz name:** Койчу баштык (Koychu bashtyk) 1187
- Description:** Herbaceous annual. Stems 5–60 cm tall. Basal leaves in a rosette, petiolate, oblanceolate entire to pinnatifid with triangular lobes; cauline leaves elongate, upper leaves almost linear with sagittate bases. Inflorescence an apical raceme. Flowers small, pedicellate with 4 white petals. Fruit a silicle, triangular to heart-shaped. Seeds small, oval, slightly flattened, yellow-brownish. 1188
1189
1190
1191
- Other distinguishing features:** Stamens 6 (4 long, 2 short). 1192
- Phenology:** Flowers in April-June, fruits in June-July. 1193
- Reproduction:** By seeds. 1194
- Distribution:** All provinces of Kyrgyzstan and Uzbekistan. 1195
- Habitat:** From the foothills to high mountains, on waste grounds, abandoned fields, near houses, along roads and canals. 1196
- Population status:** Common, forming dense groups. 1197
- Traditional use:** The aboveground parts are used in Kyrgyz folk medicine to treat uterine bleeding, malignant ulcers, stomach cancer, dysentery, gastritis, tuberculosis, and venereal diseases. In Chinese medicine the roots are used to treat dysentery and eye diseases, and in Tibet they are used as an antiemetic (Plant Resources of the USSR 1986). 1198
1199
1200
- Documented effects:** The herb strengthens the tonus of uterine muscles and narrows the peripheral veins (Maznev 2004). 1201
The peptides shepherdin I and shepherdin II, isolated from the roots, exhibited antimicrobial activity against gram-negative bacteria and fungi (Park et al. 2000). 1202
1203
- Phytochemistry:** The herb contains rhamnoglucosides (including hyssopin), choline, acetylcholine, tyramine, inosine, tannins, bursic, fumaric, malic, tartaric and citric acids, vitamins A, B, C and K, saponins, phytoncides, and essential oils (Kurochkin 1998). 1204
1205
1206




▲ *Campanula glomerata* L. Photos: *left and center*: Alexander Naumenko; *right*: Ilya Raskin



▲ *Capparis spinosa* L. Photos: *left and center*: John B. Taft; *right*: Alexander Naumenko

▼ *Capsella bursa-pastoris* (L.) Medik. Photos: Sergey Appolonov

▼ *Bunium persicum* (Boiss.)
B. Fedtsch. Photo: Authors



Carum carvi L. – Apiaceae	1208
Synonyms: <i>Carum gracile</i> Lindl., <i>Carum rosellum</i> Woronow.	1209
English name: Caraway	1210
Russian name: Тмин обыкновенный (Tmin obyknovennyy)	1211
Uzbek name: Korazira	1212
Kyrgyz name: Кадимки карум (Kadimki karum)	1213
Description: Herbaceous biennial, occasionally annual or perennial. Stems 30–90 cm tall. Leaves bi- or tripinnatisect; basal leaves long-petiolate, segments linear-lanceolate; cauline leaves short-petiolate. Inflorescence a compound umbel with 8–16 rays. Flowers small with 5 petals, white or pink. Fruit a 2-seeded schizocarp, brown, 3–5 mm long, 1–2 mm wide, sides flattened.	1214 1215 1216 1217
Other distinguishing features: Fruits have a distinct aroma.	1218
Phenology: Flowers in June, fruits in July.	1219
Reproduction: By seeds.	1220
Distribution: All provinces of Kyrgyzstan; Toshkent, Samarqand and Surxondaryo provinces of Uzbekistan.	1221
Habitat: From valleys to high mountains. Found in meadows, along canals and river floodplains, near bogs and in forest glades.	1222 1223
Population status: Common, forming dense groups.	1224
Traditional use: Fruits, which are collected in July and August, are used for medicinal purposes. They are used as a sedative, expectorant, diuretic, and is included in a preparation used as a carminative, laxative, sedative, and to increase appetite (Turova and Sapozhnikova 1984).	1225 1226 1227
Documented effects: Caraway strengthens the appetite, promotes digestion, reduces spasms in smooth muscles (intestinal, uterine, and urethral), increases diuresis, and promotes expelling of phlegm and sputum (Turova and Sapozhnikova 1984). The essential oil isolated from the fruits exhibited antibacterial activity against a variety of gram-positive and gram-negative bacteria (Iacobellis et al. 2005).	1228 1229 1230 1231
Phytochemistry: Fruits contain 3–7 % essential oil, 14–22 % fatty oil, and tannins. The essential oil contains limonene, carvacrol, carvone, and other compounds. The flavonoids quercetin, camphorol, isorhamnetin and polyenes were isolated from the aboveground parts and flowers that were collected during the flowering stage (Khalmatov et al. 1984; Turova and Sapozhnikova 1984; Iacobellis et al. 2005).	1232 1233 1234 1235

1237 **Centaurea depressa M. Bieb. – Asteraceae**

1238 **Synonyms:** *Cyanus depressus* (M. Bieb.) Soják.

1239 **English name:** Low cornflower, Iranian knapweed

1240 **Russian name:** Василек придавленный (Vasilek pridavlenny)

1241 **Uzbek name:** Butakuz

1242 **Kyrgyz name:** Жагалак кёп башы (Zhagalak kyop bashy)

1243 **Description:** Herbaceous annual plant, to 15–60 cm tall. Stems multiple, abundantly branched, gray-tomentose, foliaceous
1244 from the base. Basal and lower leaves simple, petiolate, oblong, 5–10 cm long, entire to pinnatifid, gray-tomentose; upper
1245 leaves sessile, linear-lanceolate, entire. Inflorescences pedunculate capitula, arranged solitarily; involucre bracts coriaceous,
1246 silvery fimbriate along the edges. Ray flowers bright blue or blue-violet; disk flowers violet. Fruits obovate achenes,
1247 mostly smooth, shiny, with pappus.

1248 **Other distinguishing features:** Outer pappi of stiff bristles, unequal, up to 8 mm long.

1249 **Phenology:** Flowers in May-June, fruits in June-July.

1250 **Reproduction:** Only by seeds.

1251 **Distribution:** All of Uzbekistan; Chuy, Talas, Osh and Jalal-Abad provinces of Kyrgyzstan.

1252 **Habitat:** The adyr and tau zones. Clay-soiled slopes with rocky debris. Often occurs as a weed in wheat fields.

1253 **Population status:** Common, found in small populations.

1254 **Traditional use:** A decoction of the flowers is used for melancholy, neurasthenia, eye diseases, and as a cholagogue for
1255 hepatitis (Khalmatov 1964).

1256 **Documented effects:** An extract of the aboveground parts had antibacterial effects against *Bacillus subtilis*, *Escherichia coli*,
1257 *Proteus mirabilis*, and *Pseudomonas aeruginosa* (Bazzaz and Haririzadeh 2003; Arif et al. 2004). A hexane extract of the
1258 plant showed antifungal activity against *Candida krusei* (Karamenderes et al. 2006).

1259 **Phytochemistry:** The herb contains flavonoids, phenolic compounds and small amounts of alkaloids (Khalmatov 1964;
1260 Bandyukova et al. 1969; Hosseinimehr et al. 2007). The main components of the essential oil isolated from plants in Iran,
1261 were piperitone and elemol (Esmaeili et al. 2005).

1262

Ceratocephala testiculata (Crantz) Roth – Ranunculaceae	1263
Synonyms: <i>Ceratocephala orthoceras</i> DC., <i>Ceratocephala reflexa</i> Steven, <i>Ranunculus testiculatus</i> Crantz.	1264
English name: Bur buttercup, tubercled crowsfoot, curvseed butterwort	1265
Russian name: Рогоглавник яичковидный, Рогоглавник пряморогий (Rogoglavnik yaichkovidnyy, Rogoglavnik pryamorogiy)	1266 1267
Uzbek name: Uchma, Kuitikan	1268
Kyrgyz name: Unknown	1269
Description: Herbaceous annual. Stems erect or suberect, tomentose. Basal leaves petiolate, blades broadly spatulate in outline, 1–3-dissected, segments linear. Flowers solitary, terminal. Sepals 5, 2–8 mm long, spreading, tomentose. Petals 5, yellow. Fruits achenes in ovoid clusters. Achenes 4.5–6.5 mm long, with a straight, sharp beak.	1270 1271 1272
Other distinguishing features: Early flowering ephemeral. Sepals persistent in fruit.	1273
Phenology: Flowers in March-April, fruits in April-May.	1274
Reproduction: By seeds.	1275
Distribution: Widespread throughout Uzbekistan and Kyrgyzstan.	1276
Habitat: The chul, adyr, and tau zones. Clayey and sandy soils, pebbly, dry slopes, and very salty areas.	1277
Population status: Common, weedy.	1278
Traditional use: The plant is used to treat wounds, injuries, eczema, and other skin diseases (Khalmatov 1964).	1279
Documented effects: The plant is used as a treatment for pyoderma and furunculosis. An ointment made of the dried herb is used to treat sores. Oil extracts of the fresh herb can accelerate reduction of inflammatory edema and stimulate steady increase in tissue granulation and wound epithelization (Khalmatov 1964). Studies showed that plants growing in Utah (USA) were toxic to sheep with a minimum lethal dosage of 11 g (wet weight) of green plant material/kg. Signs of poisoning are weakness, depression, diarrhea, labored breathing, anorexia, and occasional fever (Nachman and Olsen 1983).	1280 1281 1282 1283 1284
Phytochemistry: Contains anemonin, uronic acids, resins, carotene, and sugars (Khalmatov 1964). Analyses of plants collected in Utah (USA), revealed that the “early flower” stage contained the highest concentration of the toxic compound ranunculin (Nachman and Olsen 1983).	1285 1286 1287

1289 **Cichorium intybus L. – Asteraceae**1290 **Synonyms:** None1291 **English name:** Chicory1292 **Russian name:** Цикорий обыкновенный (Tsikoriy obyknovennyy)1293 **Uzbek name:** Sachratki1294 **Kyrgyz name:** Кадимки дарчын (Kadimki darchyn)

1295 **Description:** Herbaceous perennial with a large taproot. Stems 15–120 cm tall. Basal leaves long-elliptic, 10–28 cm long,
1296 2–6 cm wide, pinnatifid to pinnatisect, base tapering to the petiole; lower cauline leaves oblong-ovate to broadly lanceo-
1297 late, large dentate; upper cauline leaves small, linear to lanceolate, margins almost entire. Inflorescences capitula, axillary,
1298 in groups of 1–3; involucre bracts in 2 rows. Flowers only ligulate; ligules blue, 5-toothed. Fruits 3–6-sided achenes,
1299 2–3 mm long, brown; pappus of very short scales.

1300 **Other distinguishing features:** Anthers bluish, connate around style. Sap milky.1301 **Phenology:** Flowers in June-July, fruits in July-August.1302 **Reproduction:** By seeds.

1303 **Distribution:** All provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Tashkent, Samarqand, Farg'ona, Buxoro
1304 and Surxondaryo provinces of Uzbekistan.

1305 **Habitat:** On waste grounds, near roads, in fallow meadows and dry, stoney or clayey waterways, in vegetable gardens and
1306 in cultivated fields.

1307 **Population status:** Common, forming dense groups.

1308 **Traditional use:** The roots and inflorescences are used to prepare folk medicines. The roots are used to increase the appetite
1309 and improve digestion. Inflorescences are used to treat inflammation of the stomach lining, diseases of the large and small
1310 intestines, gall bladder, and kidneys, as well as to treat kidney and gallstones. The inflorescences are also used swelling
1311 due to various heart conditions (Ladigina and Morozova 1987; Nogaller et al. 1987).

1312 **Documented effects:** According to experiments, an infusion of the inflorescences has sedative effects on the central nervous
1313 system and strengthens the heart function (Akopov 1990). A decoction of the aboveground parts has diuretic, astringent,
1314 and antimicrobial effects. Liquid extracts from the roots reduced the blood sugar content in people with early stage dia-
1315 betes (Khalmatov et al. 1984). Extracts of the plant have been shown to affect cholesterol uptake, tumor development,
1316 prevent immunotoxicity induced by ethanol, and have anti-inflammatory properties (Schmidt et al. 2007). Experiments
1317 with mice and rats showed that an extract of the root, rich in sesquiterpene lactones, significantly reduced inflammation,
1318 by down-regulating pro-inflammation gene expression and reducing nitric oxide production (Ripoll et al. 2007).

1319 **Phytochemistry:** Roots contain up to 65 % inulin, the glycoside intibin, alkaloids, organic acids, and vitamin B and C. The
1320 flowers contain the glycoside cichoriin, coumarins, flavonoids, and tannins. The plant sap contains lactucin, lactucopicrin,
1321 and taraxasterol (Khalmatov et al. 1984; Schmidt et al. 2007).

1322



▲ **Carum carvi L.** Photos: *left and center*: Sergey Appolonov; *right* : Dmitri Oreshkin



▲ **Centaurea depressa M. Bieb.** Photos: *left*: Alim Gaziev; *right*: Evgeny Davkaev

▼ **Cichorium intybus L.**

Photos: *top*: Evgeny Davkaev; *bottom*: Sasha Eisenman



◀ **Ceratocephala testiculata (Crantz) Roth** Photos: Maxim Kucherov



1323 **Clematis orientalis L. – Ranunculaceae**

1324 **Synonyms:** *Clematis grata* Wall., *Clematis incisodentata* Rich., *Clematis orveniae* Harvey & Sonder, *Clematis petersiana*
1325 Klotzsch, *Clematis thunbergii* Steud., *Clematis triloba* Thunb., *Clematis viridiflora* Bertol., *Viticella orientalis* (L.) W.A.
1326 Weber.

1327 **English name:** Oriental virginsbower

1328 **Russian name:** Ломонос восточный, Клематис восточный (Lomonos vostochnyy, Klematis vostochnyy)

1329 **Uzbek name:** Пан чуп

1330 **Kyrgyz name:** Чыгыш жебелгеси (Chygysh zhebelgesi)

1331 **Description:** Perennial, semi-woody climbing vine. Stems 2–8 m long, sometimes reddish, glabrous or densely short-hairy.
1332 Leaves pinnately compound; leaflets (3–)5–7, lanceolate to ovate, slightly lobed, entire or coarsely dentate, with short
1333 appressed hairs or nearly glabrous. Flowers pedicellate, solitary or in axillary cymes. Sepals 4, greenish-yellow, recurved,
1334 often hairy. Petals absent. Fruits hairy achenes, 2 mm long with a long beak (3–8 cm).

1335 **Other distinguishing features:** Plant climbs using tendril-like petioles and leaf-rachises. Staminal filaments hairy towards
1336 base.

1337 **Phenology:** Flowers in June-September, fruits in July-October.

1338 **Reproduction:** By seeds.

1339 **Distribution:** Widespread throughout all of Uzbekistan and Kyrgyzstan.

1340 **Habitat:** The chul, adyr, and tau zones. Along river banks and irrigation canals, along fences and among bushes.

1341 **Population status:** Common.

1342 **Traditional use:** The fresh herb is used as a source for antivenom to treat snake bites. A powder and decoction has strong
1343 insecticidal properties (Khalmatov 1964). In Central Asia the plant is used to treat tuberculosis (Ogolevitz 1951).
1344 Preparations are applied externally to treat chronic eczema with itching. In Chinese medicine preparations are used as a
1345 sedative, analgesic, diuretic, diaphoretic, to treat cystitis and as an anti-inflammatory to treat rheumatism, gout, and
1346 chronic gonorrhea. Other species, particularly *Clematis hexapetala*, are used as an antivenom remedy to treat snake bites
1347 and as an analgesic (Ibragimov and Ibragimova 1960).

1348 **Documented effects:** Extract of the leaves have strong bactericidal and fungicidal actions, possibly due to the presence of
1349 anemonin. The fresh herb is considered poisonous, probably because of anemonin, which disappears after drying
1350 (Ogolevitz 1951). In experiments, an extract of the herb had antibacterial actions on gram-positive microbes (Khalmatov
1351 1964).

1352 **Phytochemistry:** The aboveground parts contain the alkaloid clematine, green resin with melissic acid, myricyl alcohol, and
1353 caulosapogenin glycoside. The roots contain alkaloids (Khalmatov 1964).

1354

- Cnicus benedictus L. – Asteraceae** 1355
- Synonyms:** *Centaurea benedicta* (L.) L., *Hierapicra benedicta* (L.) Kuntze. 1356
- English name:** Blessed thistle 1357
- Russian name:** Кникус благословенный, Волчец кудрявый (Knikus blagoslovennyy, Volchets kudryavyy) 1358
- Uzbek name:** Saryq gul, Kushkunmas 1359
- Kyrgyz name:** Тармал кникус (Tarmal knikus) 1360
- Description:** Herbaceous annual, 5–70 cm high with taproot. Stems prostrate to erect, usually branching, slightly striated, often reddish, loosely hairy. Basal leaves elongate, pinnatipartite, up to 20 cm long, margins spiny-toothed, base of leaf tapering to winged petiole; stem leaves alternate, sessile; upper leaves simple, up to 5.5 cm long, sinuate with small spiny teeth. Inflorescences wide-ovate capitula, each solitary at the ends of branches; involucre bracts pinnate with spiny tips. Disk flowers yellow; ray flowers few, very slender. Fruits achenes 6–10 mm long, yellow-brown, with 20 ribs and pappi consisting of 2 rows of awns. 1361–1366
- Other distinguishing features:** Leaves glandular and slightly or densely hairy. Ray flowers sterile, 3-lobed. Achenes slightly curved. 1367–1368
- Phenology:** Flowers in May-June, fruits in June-July. 1369
- Reproduction:** Only by seeds. 1370
- Distribution:** Tashkent, Samarqand, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan. 1371
- Habitat:** The chul and adyr zones. A weed along roads and in waste places. 1372
- Population status:** Common, does not grow in large populations. 1373
- Traditional use:** The plant has a long history of use in folk medicine. Preparations were used as a cancer remedy. A decoction of the upper plant parts (capitula and leaves) is used to treat constipation, jaundice, liver diseases, hypochondria, respiratory tract catarrh, intermittent fever, gastrointestinal atonia, gout, ulcers, kidney diseases, urination disorders, and indigestion and is also used as an emetic (Ogolevitz 1951; Khalmatov et al. 1984; Khodzhimatov 1989). 1374–1377
- Documented effects:** In modern medicine, infusions and extracts of the plant are used to stimulate the appetite and improve digestion (Khalmatov et al. 1984). The herb increases the flow of gastric juices, which relieves dyspepsia, indigestion, and headaches associated with liver congestion. Extracts of the plants, including cnicin, essential oil, and polyacetylenes, have antibiotic properties. The compound cnicin has been shown to have antitumor, antimicrobial, cytotoxic, and anti-inflammatory activities. The lignans arctiin and arctigenin act as platelet activating factor (PAF) antagonists and have exhibited anti-HIV activity, as well as cytotoxic activity in vitro and antitumor activity in vivo (Tamayo et al. 2000). 1378–1383
- Phytochemistry:** The herb contains bitter substances, cnicin (a sesquiterpene lactone), resin, mucilage, sterins, tannins, essential oils, and vitamin C. The essential oil contains n-paraffin, aromatic aldehydes (cinnamaldehyde, benzaldehyde, cuminaldehyde), and monoterpenes (citronellol, fenchone, p-cymene, etc.). The seeds contain 24–28 % semi-drying fatty oil and lignans (including arctiin and arctigenin), some of which are phytoestrogen precursors for mammalian lignans (Khalmatov 1964; Khalmatov et al. 1984; Tamayo et al. 2000). 1384–1388

1390 **Codonopsis clematidea (Schrenk) C.B. Clarke – Campanulaceae**1391 **Synonyms:** *Glossocomia clematidea* (Schrenk) Fisch., *Wahlenbergia clematidea* Schrenk.1392 **English name:** Asian bellflower1393 **Russian name:** Кодонопсис ломоносовидный (*Kodonopsis lomonosovidnyy*)1394 **Uzbek name:** Qoraqurt, Dogboyut1395 **Kyrgyz name:** Конгуроодой сасык гул (*Konguroodoy sasyk gul*)

1396 **Description:** Perennial herb, 50–80 cm tall. Root fusiform, vigorous. Stem erect or winding, densely branching from the
1397 base, pubescent or glabrous, deep-green. Leaves alternate or sub-opposite, oval, acute, petiolate except at top of plant,
1398 margins entire, short-hairy. Calyx with 5 deep lobes, glabrous or pubescent; lobes up to 2 cm long, triangular, oblong or
1399 ovate-lanceolate, during flowering becoming recurved. Corolla 2–3 cm long, widely campanulate, with 5 short lobes,
1400 whitish or bluish with darker blue veins. Fruit a compressed capsule, obconical or oval, acute. Seeds oblong, shining or
1401 dull, wingless.

1402 **Other distinguishing features:** Flowering plant with strong, objectionable odor.1403 **Phenology:** Flowers and fruits in June-August.1404 **Reproduction:** By seeds.1405 **Distribution:** Toshkent, Samarqand, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.1406 **Habitat:** The tau zone. Wet taluses, along canyon bottoms and along mountain streams.1407 **Population status:** Fairly often, does not grow in dense populations.

1408 **Traditional use:** The aboveground parts are widely used in medicine as a cholagogue for hepatitis and cholecystitis
1409 (*Rakhimov et al. 2003*).

1410 **Documented effects:** This plant is a part of cholagogue collection by professor Khodzhimatov. Effective doses of the alka-
1411 loid codonopsin provoked general depression in mice. Codonopsin reduced blood pressure in acute experiments on cats
1412 and caused premature ventricular beats when applied intravenously to rabbits (*Khanov et al. 1971*). Codonopsinine and
1413 codonopsine have antibiotic properties and exhibit hypotensive activity with no observed effects on the central nervous
1414 system in animal tests (*Haddad and Larchevêque 2003*).

1415 **Phytochemistry:** The aboveground plant parts contain the alkaloids codonopsin and codonopsinin (*Yunusov 1974*;
1416 *Tashkhodzhaev et al. 2004*).

1417

Conium maculatum L. – Apiaceae

- Synonyms:** *Cicuta major* Lam., *Cicuta officinalis* Crantz, *Conium cicuta* Neck., *Conium maculosum* Pall., *Coriandrum cicutum* Crantz, *Coriandrum maculatum* (L.) Roth, *Selinum conium* (Vest) E.L. Krause, *Sium conium* Vest. 1418-1420
- English name:** Poison hemlock 1421
- Russian name:** Болиголов пятнистый (Boligolov pyatnistyy) 1422
- Uzbek name:** Sasik alaf 1423
- Kyrgyz name:** Уу балдыркан (Uu baldyrkan) 1424
- Description:** Herbaceous biennial. Stems 60–200 cm with red-brownish spots on lower portion. Basal leaves triangular in outline, petiolate, 30–60 cm long, tripinnatisect; primary and secondary segments petiolulate, tertiary segments sessile, oblong-ovate, pinnatifid. Inflorescence a compound umbel with 10–20 rays; bracts 4, lanceolate, acute; bracteoles 3–7, connate at the base. Petals 5, obcordate, white. Fruit a 2-seeded schizocarp, 3–3.5 mm long, nearly orbicular to ovate, with wavy ribs. 1425-1429
- Other distinguishing features:** Flowers have sharp smell. 1430
- Phenology:** Flowers in June, fruits in July. 1431
- Reproduction:** By seeds. 1432
- Distribution:** All provinces of Kyrgyzstan; Tashkent, Samarqand, and Surxondaryo provinces of Uzbekistan. 1433
- Habitat:** In forest glades, long-used animal corrals in the tallgrass-meadow belt of mountains, and in valleys. 1434
- Population status:** Common, forming dense groups. 1435
- Traditional use:** Aboveground parts of the plant and seeds are used. The plant is poisonous. It is used as a sedative, anticonvulsant, and analgesic, and to treat chorea, epilepsy, pertussis, migraine headaches, cancer, and uterine fibroids (Khalmatov 1964). 1436-1438
- Documented effects:** Extracts and plasters from this plant are rarely used externally as analgesics (Khalmatov 1964). The plant contains piperidine alkaloids that are toxic to humans and animals. These alkaloids have also been shown to cause congenital birth defects in goats and pigs (Panter et al. 1985a, b). These alkaloids have 2 modes of action. The first is similar to curare, which effects neuromuscular function and can cause respiratory failure. The second action effects the autonomic ganglia and can cause salivation, mydriasis, and tachycardia, followed by bradycardia and occasionally rhabdomyolysis and acute tubular necrosis (Frank et al. 1995; Lopez et al. 1999). 1439-1444
- Phytochemistry:** The roots contain up to 0.042 % total alkaloids, with the stems up to 0.065 % and leaves up to 0.1 %. The fruits contain up to 1 % total alkaloids, but sometimes unripe fruits contain up to 2 % (with 50 % of it being coniine). Other alkaloids include conhydrine, pseudoconhydrine, γ -coniceine, and methyl-coniine. The above parts also contain essential oils (mainly terpenes), vitamin C, carotene, and caffeic acid. Quercetin and kaempferol have been isolated from the flowers (Khalmatov 1964; Lopez et al. 1999). 1445-1449



▲ **Codonopsis clematidea**
(Schrenk) C.B. Clarke
Photos: Sasha Eisenman



► **Conium maculatum** L.
Photos: *top*: Clinton Shock;
bottom: Alim Gaziev

▼ **Cnicus benedictus** L.
Photos: Bazar Dovletov



Clematis orientalis L. Photos: Alexander Naumenko ►

- Convolvulus arvensis L. – Convolvulaceae** 1451
- Synonyms:** *Convolvulus chinensis* Ker Gawl., *Convolvulus sagittifolius* (Fisch.) T. Liou & Ling. 1452
- English name:** Field bindweed 1453
- Russian name:** Вьюнок полевой (V'yunok polevoy) 1454
- Uzbek name:** Куу печак 1455
- Kyrgyz name:** Чырмак (Chyrmok) 1456
- Description:** Herbaceous perennial. Stems twining or prostrate. Leaves alternate, long-petiolate, ovate to oblong; apex acute, obtuse to rounded; base usually hastate. Inflorescences axillary cymes, 2–3-flowered, or flowers solitary; long-pedunculate. Corolla funnelform, up to 3 cm in diameter, pink or white. Stamens 5. Stigmas 2. Fruit a smooth, spherical capsule. Seeds 3–4 mm long, brownish or black. 1457
- Other distinguishing features:** Flowers plicate when young. 1458
- Phenology:** Flowers in May–September, fruits in June–September. 1459
- Reproduction:** By seeds and rhizomes. 1460
- Distribution:** In all provinces of Kyrgyzstan and Uzbekistan. 1461
- Habitat:** In abandoned fields and waste grounds. 1462
- Population status:** Common, forming dense groups. 1463
- Traditional use:** The roots, leaves, stems, and flowers are used in folk medicine. The powdered leaves are used to treat wounds, cuts, and bruises. Juice from the leaves mixed with cow fat is used to treat lung and ear diseases. The root is used as a laxative. A decoction of the herb is used to wash wounds and to treat skin ulcers, fungal skin diseases, and scabies. Avicenna used this species to treat asthma, lung disease, chest pains, liver and spleen diseases, and as a choleric remedy (Khalmatov 1964). 1464
- Documented effects:** In experiment on animals, this plant species had hypotensive, antispasmodic, anti-inflammatory, and styptic properties (Plant Resources of the USSR 1985). A methanol extract of the plant induced a dose-dependent relaxation of duodenal smooth muscle in rabbits (Atta and Mouneir 2004). Mice fed high doses of the plant died or had severe hepatic necrosis and gastritis after 4–7 days. Mice fed low doses of the plant had no clinical disease or large lesions, but developed mild multifocal hepatitis and gastritis (Schultheiss et al. 1995). 1465
- Phytochemistry:** All parts of the plant contain alkaloids. The roots contain up to 5 % resins. The resins contain convolvine, jalapine, convolvuline, and caffeic acid. The aboveground parts contain flavonoids (quercetin and kaempferol) and caffeic acid. The leaves contain carotene and vitamin C (Khalmatov 1964). Plants from a pasture in Colorado (USA) were found to contain the tropane alkaloids tropine, pseudotropine, and tropinone, and the pyrrolidine alkaloids cuscohygrine and hygrine (Todd et al. 1995). 1466

1483 **Convolvulus subhirsutus Regel & Schmalh. – Convolvulaceae**

1484 **Synonyms:** *Convolvulus chondrilloides* Boiss. var. *sericeus* Kuntze, *Convolvulus dorychium* ssp. *subhirsutus* (Regel &
1485 Schmalh.) Saad, *Convolvulus tschimganicus* Popov.

1486 **English name:** Unknown

1487 **Russian name:** Вьюнок жёстковолосистый, Вьюнок шерстистый (V'yunok zhyostkovolosisty, V'yunok sherstisty)

1488 **Uzbek name:** Mingbosh

1489 **Kyrgyz name:** Туктуу чырмоок (Tuktuu chymook)

1490 **Description:** Hairy perennial herb to 40–100 cm with a thick taproot. Stems multiple, erect, with spreading branches. Leaves
1491 alternate, simple, elliptic, elongate-lanceolate or obovate, 1.5–10 cm long, narrowly acute, margins entire; petioles
1492 0.5–2 cm long. Inflorescence a long-branched dichasium. Flowers pink, lilac, or rarely white. Corolla funnelform,
1493 1–2.3 cm long. Fruit an ovoid capsule, 4–7 mm long, 1-seeded, glabrous. Seeds oviform or flat-elliptical, brown.

1494 **Other distinguishing features:** Sepals are recurved when plant is fruiting. Seeds velutinous.

1495 **Phenology:** Flowers in June–July, fruits in July–August.

1496 **Reproduction:** By seeds.

1497 **Distribution:** Toshkent, Andijon, Farg'ona, Samarqand, Buxoro, and Surxondaryo provinces of Uzbekistan; all of
1498 Kyrgyzstan.

1499 **Habitat:** The adyr and tau zones. Shallow soil, more rarely on shallow-soiled, stony slopes.

1500 **Population status:** Common, usually found in small populations.

1501 **Traditional use:** In the folk medicine of Tajikistan, a decoction of the seeds is drunk to treat gastrointestinal diseases. An
1502 infusion of the herb is used as an analgesic, anticonvulsant, to heal wounds, and to treat asthma and lung tuberculosis
1503 (Khalmatov et al. 1984; Khodzhimatov 1989).

1504 **Documented effects:** The alkaloids convolvine and convolamine have anesthetic properties, produce irritation of the mucous
1505 membranes of the eyes, and large doses can paralyze the central nervous system. Due to high toxicity they are not used in
1506 medical practice. Their derivatives, convocaine and tropacin, were developed as preparations for medical use. *Tropacin* is
1507 used to treat Parkinson's disease, spastic paresis and other diseases following muscle tone increase. *Tropacin* is also rec-
1508 ommended to treat ulcers, bronchial asthma, other cases of spasms of the smooth muscular system, and poisoning with
1509 phosphorganic compounds (Mashkovskii 1953; Khalmatov et al. 1984; Khodzhimatov 1989).

1510 **Phytochemistry:** The herb contains alkaloids (convolvine, convolamine, convolidine, phyllalbine, phyllalbine N-oxide, etc.)
1511 and the aminoalcohol nortropine (Razzakov and Aripova 2004; Gapparov et al. 2007). Roots collected at the end of the
1512 growing season in a Toshkent suburb (Kaplanbek) contained 4.1 % total alkaloids. The aboveground parts collected in the
1513 beginning of the growing season (March) contained 2.08 % total alkaloids (Yunusov 1974).

Cousinia lappacea Schrenk – Asteraceae	1515
Synonyms: None	1516
English name: Unknown	1517
Russian name: Кузиния репейниковидная (Kuziniya repеynikovidnaya)	1518
Uzbek name: Unknown	1519
Kyrgyz name: Уйгактай кокуй тикен (Uygaktay kokuy tiken)	1520
Description: Herbaceous perennial. Stems numerous, erect, 40–70 cm tall, up to 2 cm in diameter at base, usually pinkish or purple. Leaves soft, gray-hairy; basal leaves petiolate, obovate, margins spinose-toothed; cauline leaves oblanceolate, finely prickly-toothed, sessile, densely arranged. Inflorescences oviform capitula, 12–13 mm long, 5–6 mm wide, with 4–5 flowers; involucre bracts 25–30; outer bracts closely appressed, ovate, apices acuminate-hooked. Disk flowers purple; ray flowers lacking. Fruits obovoid achenes, 6 mm long, 4 mm wide, smooth.	1521 1522 1523 1524 1525
Other distinguishing features: Inner involucre bracts purple at apices, ending with a thin hooked spine. Receptacle with smooth bristles.	1526 1527
Phenology: Flowers in June, fruits in July.	1528
Reproduction: By seeds.	1529
Distribution: Chuy, Naryn, and Jalal-Abad provinces of Kyrgyzstan; Samarqand and Farg'ona provinces of Uzbekistan.	1530
Habitat: On stony slopes of the middle mountain belt.	1531
Population status: Common, found as single individuals.	1532
Traditional use: An infusion of the herb and roots is used to prevent tumor growth and to treat gastrointestinal ulcers (Plekhanova et al. 1985).	1533 1534
Documented effects: No data.	1535
Phytochemistry: The aboveground parts contain monosaccharides (glucose and fructose), oligosaccharides (5.1–5.5 %), and pectic substances. The roots contain water-soluble polysaccharides (2.18–2.78 %; Plekhanova et al. 1985).	1536 1537

1539 **Cousinia umbrosa Bunge – Asteraceae**1540 **Synonyms:** None1541 **English name:** Unknown1542 **Russian name:** Кузиния теневая (Kuziniya tenevaya)1543 **Uzbek name:** Okboshikon1544 **Kyrgyz name:** Колоко кокуй тикен (Koloko kokuy tiken)

1545 **Description:** Herbaceous perennial. Stems numerous, 60–120 cm tall, deeply grooved, branched above. Leaves green and
1546 glabrous on adaxial side, abaxial side grayish felted; basal leaves very large, petiolate, obovate, cordate, margins irregu-
1547 larly dentate; cauline leaves similar but becoming gradually reduced towards apex. Inflorescences ovoid capitula arranged
1548 in a panicle; involucre bracts oblong, bases appressed, spreading above and tapering to incurved hooks. Disc flowers
1549 10–12, pink; ray flowers lacking. Fruits obovoid achenes, 6 mm long, 3 mm wide, light brown with dark spots.

1550 **Other distinguishing features:** Outer involucre bracts with 1 or 2 pairs of glandular hairs; inner bracts linear, apex acumi-
1551 nate. Receptacles with smooth bristles.

1552 **Phenology:** Flowers in June-July, fruits in July-August.1553 **Reproduction:** By seeds.1554 **Distribution:** Chuy and Jalal-Abad provinces of Kyrgyzstan; Toshkent and Qashqadaryo provinces of Uzbekistan.1555 **Habitat:** In shady places in the foothills and the lower mountain belt.1556 **Population status:** Common, found in dense groups.

1557 **Traditional use:** An infusion and decoction of the aboveground and underground parts is applied to treat stomach ulcers and
1558 hypoxia in mountainous conditions, and is also used as a general tonic during recovery from a variety of diseases
1559 (Turdumambetov 1995).

1560 **Documented effects:** No data.

1561 **Phytochemistry:** The roots and aboveground plant parts contain oligo- and polysaccharides (fructan), pectic substances, and
1562 hemicellulose (Turdumambetov 1995; Turdumambetov et al. 2007). The fruits contain lipids, hydrocarbons, triterpene
1563 alcohols, sterols, mono- and diacylglycerides, etc. The predominant fatty acids are 16:0, 18:1, and 18:2 (Ul'chenko et al.
1564 1999)

1565



▲ **Convolvulus arvensis L.**
Photos: *top*: Sasha Eisenman;
bottom: Mary Backlund



▲ **Convolvulus subhirsutus Regel & Schmalh.**
Photos: Evgeny Davkaev



▲ **Cousinia umbrosa Bunge**
Photos: Evgeny Davkaev



◀ **Cousinia lappacea Schrenk**
Photos: Georgy Lazkov

1566 **Crambe kotschyana Boiss. – Brassicaceae**

1567 **Synonyms:** *Crambe cordifolia* ssp. *kotschyana* (Boiss.) Jafri, *Crambe cordifolia* var. *kotschyana* (Boiss.) O.E. Schulz,
1568 *Crambe palmatifida* Regel & Schmulh., *Crambe sewerzowii* Regel.

1569 **English name:** Colewort

1570 **Russian name:** Катран (Katran)

1571 **Uzbek name:** Katron

1572 **Kyrgyz name:** Кочи катраны (Kochi katrany)

1573 **Description:** Perennial herb with a thick, fleshy root. Stems 50–150(–250) cm tall, single or multiple, coarse-hairy; branches
1574 spreading. Basal leaves with long petioles up to 30 cm long, blades cordate-reniform to ovate-oblong, up to 50 cm wide,
1575 roughly lobed, coarsely toothed, coriaceous, covered with rough prominent hairs; upper leaves alternate, ca. 1 cm long.
1576 Inflorescences racemes arranged in large panicles. Petals 4, obovate, white. Stamens 6, tetradynamous, the longer 4
1577 toothed. Fruit an elongate-spherical silique, 6–7 mm long.

1578 **Other distinguishing features:** Seeds 3–4 mm in diameter, pale brown.

1579 **Phenology:** Flowers and fruits from the end of March until mid-June.

1580 **Reproduction:** Propagates by seeds, it can be easily cultivated.

1581 **Distribution:** Karakalpakstan autonomous republic, Toshkent, Andijon, Farg'ona, Jizzax, Samarqand, Qashqadaryo, and
1582 Surxondaryo provinces of Uzbekistan; Chuy, Talas, Osh and Jalal-Abad provinces of Kyrgyzstan.

1583 **Habitat:** The adyr and tau zones. On soils with a high diversity of minerals and shallow-soiled slopes with rocky debris.

1584 **Population status:** Common.

1585 **Traditional use:** Seeds are used to treat respiratory tract catarrh. Roots are used by veterinarians to treat gastric diseases in
1586 camels. The roots, baked or boiled, are used for food by people (Khalmatov 1964).

1587 **Documented effects:** No data.

1588 **Phytochemistry:** Roots contain 18–19 % sugar (monosaccharides 10.54 %, disaccharide 9.2 %), 39.62 % starch. Seeds
1589 contain up to 40 % oils (Khalmatov 1964). The aboveground parts contain a variety of lipids of which a high proportion
1590 is palmitic acid. The seeds have high erucic and linolenic acid contents (Bekker et al. 2003).

1591

Crataegus altaica (Loudon) Lange – Rosaceae

- Synonyms:** *Crataegus chlorocarpa* Lenne & K. Koch, *Crataegus korolkowii* L. Henry, *Crataegus purpurea* var. *altaica* Loudon, *Crataegus sanguinea* var. *incisa* Regel, *Crataegus sanguinea* var. *inermis* Kar. & Kir., *Crataegus wattiana* var. *incisa* C.K. Schneid. 1592–1595
- English name:** Altai hawthorn, Altai mountain thorn 1596
- Russian name:** Боярышник алтайский (Boyaryshnik altayskiy) 1597
- Uzbek name:** Dulana zardak, Sarik dulana 1598
- Kyrgyz name:** Алтай долоносу, Сары долоно (Altay dolonosu, Sary dolono) 1599
- Description:** Shrubby tree with multiple trunks, up to 3–5 m tall. Bark smooth, mostly gray; 1-year old stems red-brown or green-brown, smooth, with many white lenticels; older branches gray-orange with large lenticels; some branches with short (1–1.5 cm), thick spines in the leaf axils. Leaves petiolate, broadly triangular, oval or circular, entire or 3–7-lobed, coarsely toothed. Inflorescence corymbiform with 10–30 flowers. Flowers up to 1.9 cm in diameter, with 5 white petals. Fruits spherical pomes, yellow or dark-brown with 3–5 seeds. 1600–1604
- Other distinguishing features:** Fruit 8–10 mm in diameter. Leaves glabrous or slightly pubescent. 1605
- Phenology:** Flowers in May-June, fruits in August-September. 1606
- Reproduction:** By seeds. 1607
- Distribution:** All provinces of Kyrgyzstan; Toshkent, Farg'ona and Qashqadaryo provinces of Uzbekistan. 1608
- Habitat:** Along forest edges, in the understory, and in meadows. 1609
- Population status:** Common, found as single plants. 1610
- Traditional use:** Flowers, fruits, leaves and bark are used in folk medicine for treatment of various illnesses. A decoction of leaves and tea from dried flowers and fruits are taken to treat hypertension, dizziness, tachycardia, insomnia, heart diseases, and common colds. Fresh fruits are recommended as a laxative (Khodzhimatov 1989). 1611–1613
- Documented effects:** Extracts from fruits or tinctures from flowers are used to treat cardiovascular diseases and hypertension, to tone cardiac muscles, and as a sedative. This hawthorn extract is part of *Cardiovalen*, which is used to treat rheumatic heart disease, cardiosclerosis, stenocardia, and vegetative neurosis (Khodzhimatov 1989). 1614–1616
- Phytochemistry:** The bark contains tannins and the fruits contain up to 260 mg% vitamin C (Zapryagaeva 1964). Flowers contain flavonoids (hyperoside, quercetin, vitexin, and vitexin-ramnoside), triterpene saponins (ursolic and oleanolic acids) and essential oil. The fruits contain flavonoids, saponins, tannins, polysaccharides, fatty oil, and phenolcarboxylic acids (chlorogenic and caffeic; Khodzhimatov 1989). 1617–1620

1622 **Crataegus songarica** K. Koch. – Rosaceae

1623 **Synonyms:** *Crataegus fischeri* C.K. Schneid.

1624 **English name:** Dzhungarskei hawthorn

1625 **Russian name:** Боярышник сонгорский (Boyaryshnik songorskiy)

1626 **Uzbek name:** Dulana

1627 **Kyrgyz name:** Сонгор долоносу (Songor dolonosu)

1628 **Description:** Shrubby tree, up to 4–5 m tall with multiple trunks, each 5–9 cm in diameter. Bark reddish-gray to blackish
1629 with small cracks; young twigs green, glabrous or slightly hairy with spines up to 1.7 cm long; 1-year old twigs reddish-
1630 brown; older branches brown with smooth bark. Leaves petiolate, 6–8 cm long, 5–6 cm wide, broadly triangular to almost
1631 circular, 5–7-lobed, margins dentate. Inflorescence corymbiform with 28–35 flowers. Flowers up to 1.8 cm in diameter
1632 with 5 white petals. Fruits round pomes, dark red, with 2–3 seeds.

1633 **Other distinguishing features:** Differs from other *Crataegus* species by having smooth, brown older branches and reddish-
1634 brown year-old twigs.

1635 **Phenology:** Flowers in May-June, fruits in August-September.

1636 **Reproduction:** By seeds.

1637 **Distribution:** Chuy, Jalal-Abad, and Osh provinces of Kyrgyzstan; Toshkent, Samarqand, Qashqadaryo and Surxondaryo
1638 provinces of Uzbekistan.

1639 **Habitat:** Along river valleys and on mountain slopes.

1640 **Population status:** Common, found in loosely arranged groups.

1641 **Traditional use:** A tea made of dried flowers and infusions of dried fruits are used to treat heart pain, dyspnea, hypertension,
1642 and gastrointestinal diseases (Khodzhimatov 1989).

1643 **Documented effects:** Clinical tests of tincture of this hawthorn have yielded positive results when used against the active
1644 form of rheumatism (Kuchin 1955). An extract from the fruits and tinctures from the flowers are also used to treat cardio-
1645 vascular diseases by strengthening the heart muscle, as a sedative, and to treat hypertension (Khodzhimatov 1989).

1646 **Phytochemistry:** Fruits contain vitamin C, carotene, tannins (0.53–0.85 %), and the catechins epicatechin and leucoantho-
1647 cyanidin (Petrova 1972).

1648

- Dactylorhiza umbrosa (Kar. & Kir.) Nevski – Orchidaceae** 1649
- Synonyms:** *Dactylorchis umbrosa* (Kar. & Kir.) Wendelbo, *Orchis magna* Czerniak, *Orchis orientalis* ssp. *turkestanica* 1650
Klinge, *Orchis umbrosa* Kar. & Kir. 1651
- English name:** Unknown 1652
- Russian name:** Дактилориза теневая, Яртышник теневой (Daktiloriza tenevaya, Yartyshnik tenevoy) 1653
- Uzbek name:** Saleeb 1654
- Kyrgyz name:** Колоколуу арала (Kolokoluy arala) 1655
- Description:** Perennial herb, 30–50(–80) cm high, with a cluster of 1–6 finger-like tubers. Stems erect, thick, hollow. Leaves 1656
usually 6–7 in number, lanceolate to linear-lanceolate, apex acute, parallel veined; basal leaves 10–20 cm long, 2–5 cm 1657
wide; upper leaves smaller and more narrow, usually positioned up to the base of spike. Inflorescence spiciform, 5–18 cm 1658
long, from elongate-oviform to short- or long-cylindrical, dense and many-flowered. Flowers zygomorphic, with 6 tepals 1659
in 2 whorls, lilac- or violet-purple; lower inner tepal forming a large lip with a white “ω”-shaped blotch, spurred. Fruit a 1660
capsule with very small seeds. 1661
- Other distinguishing features:** Leaves lack spots. Bracts narrow-lanceolate, green or violet. 1662
- Phenology:** Flowers from May to the end of July. Bears fruit in July-August. 1663
- Reproduction:** By seeds. 1664
- Distribution:** Toshkent and Samarqand provinces of Uzbekistan; Chuy and Ferghana valleys and Talas province of 1665
Kyrgyzstan. 1666
- Habitat:** The adyr and tau zones. Wet, boggy places, meadows, edges of brooks, river banks, tugai, wet slopes, and ravines 1667
from 700 to 2.800 m above sea level. 1668
- Population status:** Uncommon, found as single individuals. 1669
- Traditional use:** Avicenna recommended a decoction of the tubers to treat gout, paralysis, convulsions, and joint pain. In 1670
Russian folk medicine the plant is used as a diuretic, as well as to treat fevers and gynecological diseases. The crushed 1671
tubers are mixed with lard and applied to abscesses. Fresh tuber is applied to the teeth to treat toothaches and is used 1672
to stimulate hair growth. Tadjiks use a decoction of the tubers to treat hand convulsions, paralysis, stomach catarrh, kidney 1673
stones, and to stimulate blood production. The boiled roots are used to rejuvenate the elderly and people with lung tuber- 1674
culosis. A tea made of the fried, crushed tubers is used to treat coughs, inflammation of the respiratory tract, to increase 1675
energy and to calm nerves. The roots boiled in milk is used to treat coughs, impotence, and gastrointestinal tract weak- 1676
ness. A powder of the tubers mixed with honey is used as a tonic (Khodzhimatov 1989). 1677
- Documented effects:** A decoction of the tubers is used in modern medicine to coat the digestive tract as a treatment for 1678
gastritis, enterocolitis, and other gastrointestinal diseases. It is also used as an enema to treat diarrhea in children. 1679
Experiments have shown that this plant has anti-inflammatory activities (Khalmatov et al. 1984; Khodzhimatov 1989). 1680
- Phytochemistry:** The main component of the tubers is a water soluble mucilage, which contains starch, sugars, mineral 1681
salts, bitters and proteins, essential oils, etc. During the fruit bearing stage, polysaccharide content in tubers reaches 1682
68.48 % (Khalmatov et al. 1984). The aboveground parts of many species in the genus *Dactylorhiza* contain the glycoside 1683
loroglossine. This species was found to contain traces of alkaloids and saponins as well as lactone compounds in the 1684
leaves (Khodzhimatov 1989). 1685



◀ **Crataegus kotschyana Boiss.**
Photos: Alexander Naumenko

Crataegus altaica (Loudon)
Lange Photos: Alexander Naumenko ▶

▼ **Crataegus songarica K. Koch.**
Photos: Alexander Naumenko



▼ **Dactylorhiza umbrosa**
(Kar. & Kir.) Nevski
Photos: *top*: Stefano Doglio;
bottom: Ilya Raskin



Datura stramonium L. – Solanaceae	1687
Synonyms: <i>Datura tatula</i> L., <i>Stramonium spinosum</i> Lam.	1688
English name: Jimsonweed	1689
Russian name: Дурман обыкновенный, Дурман вонючий (Durman obyknovennyy, Durman vonyuchiy)	1690
Uzbek name: Bangi divana	1691
Kyrgyz name: Кадимки чочко жангак (Kadimki chochko zhangak)	1692
Description: Herbaceous annual, up to 1 m tall. Stems erect, branching. Leaves alternate, petiolate, 8–20 cm long, up to 15 cm wide, ovate, apex acuminate, slightly lobed, margins roughly dentate. Flowers singular, in leaf and branch axils.	1693
Calyx tubular, 5-sided, up 6 cm long. Corolla white, up to 12 cm long, tubular-funnelform, 5-sided. Fruit an oviform capsule up to 5 cm long and 2.5 cm wide, densely covered with hard prickles up to 1 cm long. Seeds up to 0.5 cm long, kidney-shaped, nearly black, finely tuberculate.	1694 1695 1696 1697
Other distinguishing features: Capsule splits open into 4 valves when ripe and can contain up to 800 seeds.	1698
Phenology: Flowers in June-September, fruits in July-September.	1699
Reproduction: By seeds.	1700
Distribution: Agriculture zones in all provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Toshkent, Samarqand, Andijon, Sirdaryo, Jizzax and Surxondaryo provinces of Uzbekistan.	1701 1702
Habitat: In vegetable gardens, in orchards and waste places, near forested areas, and along rivers and canals.	1703
Population status: Common, sometimes forming dense groups.	1704
Traditional use: This species has been widely used since ancient times. Avicenna said that this plant makes you drunk, is too dangerous for the brain, and is the enemy of the heart. Beruni wrote that half a gram of the seeds can make you drunk and 4.2 g can kill you. A decoction of the seeds is used as a gargle for people with tooth- and headaches, as a painkiller and sedative, and to treat fevers, neuralgia, rheumatism, and radiculitis (Khodzhimatov 1989). Oil from the seeds is used to treat hemorrhoids and the leaves are laid over the eyes to treat eye aches (Khalmatov et al. 1984).	1705 1706 1707 1708 1709
Documented effects: The plant is poisonous. Preparations of this species is used as an antispasmodic, and mainly used to treat bronchial asthma, neuralgia, and convulsions. The leaves are a component of the preparations <i>Asthmatin</i> and <i>Asthmatol</i> . A liquid extract of the leaves is used in the preparation <i>Solutan</i> , which is used to treat bronchial asthma and bronchitis (Khodzhimatov 1989). In general, preparations of this species are used as an antispasmodic to treat bronchial asthma, stomach ulcers, cholecystitis, colitis, spastic constipation, cardio-vascular diseases, and bradycardia. Preparations derived from the plant also used as a preventive treatment for sea and air sickness (Turova and Sapozhnikova 1984). The active compounds are hyoscyamine and scopolamine. The basic pharmacological action of hyoscyamine is to block N-cholinoreceptors. Hyoscyamine increases heart rate, but decreases saliva secretion, gastric and sweat production, secretions of the pancreas, and the tonus of smooth muscles in the bronchial tubes and abdominal cavity. Hyoscyamine causes prolonged mydriasis. It tones and increases the activity of the respiratory center. In experiments with frogs a tincture reduced heart beat amplitude (Turova and Sapozhnikova 1984).	1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720
Phytochemistry: The whole plant contains alkaloids, with the main alkaloids being hyoscyamine, atropine, and scopolamine (Khalmatov et al. 1984). The leaves and stems also contain coumarins (scopoletin, esculetin, and esculin), tannins, essential oils, and carotene. Seeds contain up to 25 % fatty oil, containing linoleic, oleic, palmitic, stearic, and lignoceric acids (Khodzhimatov 1989).	1721 1722 1723 1724

1726 **Daucus carota L. – Apiaceae**

1727 **Synonyms:** *Daucus bactrianus* Bunge, *Daucus exarmatus* Korovin, *Daucus pulcherrimus* (Willd.) Koch ex DC., *Carota*
1728 *sativa* Rupr., *Carota sylvestris* (Mill.) Rupr., *Caucalis carota* (L.) Crantz, *Caucalis daucus* Crantz.

1729 **English name:** Wild carrot, Queen Anne's lace

1730 **Russian name:** Морковь дикая (Morkov' dikaya)

1731 **Uzbek name:** Yovoyi sabzi

1732 **Kyrgyz name:** Жапайы сабиз (Zhapayy sabiz)

1733 **Description:** Herbaceous biennial with a thin taproot. Stem up to 1 m high, branching, rough due to scattered, stiff hairs.
1734 Basal leaves petiolate, bipinnatisect with narrow, lanceolate or linear sections, sometimes glabrous on adaxial side, abax-
1735 ial side hairy along veins; stem leaves alternate, becoming sessile and sheathing. Inflorescence a compound umbel with
1736 many rays, up to 10 cm wide, subtended by pinnate bracts. Petals white. Fruits schizocarps with 2 one-seeded mericarps,
1737 oval or oblong, flattened, covered with short bristles along and between the ribs.

1738 **Other distinguishing features:** Umbel curving inwards in fruit and becoming spherical.

1739 **Phenology:** Flowers in May-June, fruits in July.

1740 **Reproduction:** Propagates by seeds. One plant bears up to 4,000 seeds.

1741 **Distribution:** All regions of Uzbekistan and Kyrgyzstan.

1742 **Habitat:** The chul, adyr and tau zones. Occurs fairly often as a weed in irrigated regions, predominantly in shaded places.

1743 **Population status:** Common, found as single individuals.

1744 **Traditional use:** Avicenna used wild carrot fruits as a diuretic. For a long time, the essential oils of wild carrot fruits have
1745 been used in medicine for making astringent and spicy extracts. The plant extract has been used as vermifuge and purga-
1746 tive (Khalmatov et al. 1984).

1747 **Documented effects:** In the past, flavonoids isolated from the fruits were made into a preparation called *Daukarin*. This was
1748 in used in cardiology to improve coronary blood circulation as well as chronic coronary disease (Khalmatov et al. 1984).
1749 Presently the preparation is not made because more active alternatives have been developed. An extract of the fruits of
1750 wild carrot, contained in the preparation *Urolesan*, is used in medicine. This preparation has been approved for the treat-
1751 ment of liver and kidney diseases, for acute and chronic cholestasis, and different kinds of kidney and gallbladder stones
1752 (Gammerman et al. 1990). An extract of carrot root exhibited hepatoprotective activity in mice (Bishayee et al. 1995).
1753 Compounds isolated from the seeds showed significant inhibition of cyclooxygenase (COX) enzymes (Momin et al.
1754 2003).

1755 **Phytochemistry:** Wild carrot seeds contained up to 2.5 % essential oils, which consisted of 17 substances including
1756 1- α -pinene, myrcene, bergamotene, β -bisabolene, caratol and asarone. Besides essential oil, the seeds contained flavonoids,
1757 coumarins, steroidal compounds and fatty oil. The roots also contain essential oils, which consisted of asarone, caratol,
1758 bisabolene and 5–9 % carotene. Large amounts of pyrrolidine and daucene were found in essential oils extracted from the
1759 herb. In the herb and flowers flavonoids, coumarins, anthocyanins, as well as large amounts of carotenoids, vitamins in
1760 the groups B and C, pantothenic acid, anthocyanidin, essential oils, umbelliferone and sugars were found (Gammerman
1761 et al. 1990).

Delphinium confusum Попов – Ranunculaceae	1763
Synonyms: None	1764
English name: Larkspur	1765
Russian name: Живоко́сть спутанная (Zhivokost' sputannaya)	1766
Uzbek name: Isfarak	1767
Kyrgyz name: Татыш туктуу бутуо (Tatysh tuktuu butoo)	1768
Description: Herbaceous perennial. Stems 30–70 cm tall, tomentose; leaves aggregated on the lower half of stem. Leaves petiolate, the blade circular to kidney-shaped in outline, 5–13 cm long, 7–20 cm wide, 3-lobed with sinuses half-way into leaf blade; middle lobe elongate-obovate, with 3–5 lobules; lateral lobes of leaf with 2–3 lobules; all lobes with unequal triangular-lanceolate teeth. Inflorescence a multi-flowered raceme; bracts broadly lanceolate. Flowers with 5 dark-violet tepals, upper tepal with a spur at the base. Fruit a follicetum with 3 follicles.	1769 1770 1771 1772 1773
Other distinguishing features: Spur straight, positioned almost horizontally, curving at the end.	1774
Phenology: Flowers in June-August, fruits in July-September.	1775
Reproduction: By seeds.	1776
Distribution: Chuy, Naryn, Talas, and Ysyk-Kol provinces of Kyrgyzstan; in the western Tian-Shan (Akhangaran region) and the Alai Range in Uzbekistan.	1777 1778
Habitat: In alpine meadows, spruce forests, juniper stands, on stony slopes of mountains, in feather-grass steppes, and in grassy meadows.	1779 1780
Population status: Common, found in small groups.	1781
Traditional use: A decoction of this plant is used to treat intestinal disorders, to increase muscle tone, and as an antiparasitic treatment for cows (Altimishev 1991).	1782 1783
Documented effects: In modern medicine a decoction of the stems and leaves is used in medicine to disinfect animals. Finely ground seeds mixed with butter is used to treat pediculosis (lice infestation). Tablets of condelphine are used to treat psycho-neurological diseases. The compounds delsemine and mellictine are used as anesthesia during surgical procedures (Altimishev 1991). The alkaloid condelphine has an activity similar to curare. Physicians use tablets of 0.025 g to treat conditions of excess skeletal muscle contraction, Parkinson's disease, multiple sclerosis, spastic and traumatic paralysis, etc. This preparation cannot be used by patients with conditions of reduced muscle contraction, liver and kidney diseases, or heart decompensation (Khalmatov et al. 1984).	1784 1785 1786 1787 1788 1789 1790
Phytochemistry: Flowers, stems, and leaves contain condelphine, delphiline, delatine, delsine, delsoline, isobaldine, etc., as well as aconitic acid. During the bud stage buds contain up to 0.8 % alkaloids, and during the flowering stage flowers contain up to 2 % alkaloids. The roots contain up to 1 % alkaloids (Dzhakupova 1968; Vaisov and Yunusov 1987; Narsullaev et al. 1989; Altimishev 1991).	1791 1792 1793 1794

1796 **Delphinium semibarbatum** Bienert. – **Ranunculaceae**

1797 **Synonyms:** *Delphinium bitternatum* var. *leiocarpum* Freyn., *Delphinium hybridum* var. *sulphureum* Regel, *Delphinium zalil*
 1798 Aitch.

1799 **English name:** Unknown

1800 **Russian name:** Живоко́сть полуборода́тая (Zhivokost' poluborodataya)

1801 **Uzbek name:** Isfarak

1802 **Kyrgyz name:** Жарым сакалчалуу бутёё (Zharym sakalchaluu butyoyo)

1803 **Description:** Perennial herb, 35–70 cm high. Stem unbranched or branched, glabrous or short-pubescent on lower part of
 1804 stem. Basal leaves long-petiolate; leaf blades palmatifid with 5 segments with petiolules; segments tripartite with long,
 1805 narrow-linear, glabrous or slightly villose lobes. Inflorescence racemiform. Flowers with 5 bright yellow, obovate tepals,
 1806 the upper tepal with a spur at the base. Fruits with 3 follicles, glabrous. Seeds tiny, 3-edged, paleaceous.

1807 **Other distinguishing features:** Pedicels are glabrous up to the calyx. Follicle with 3 sharply protruding longitudinal ribs
 1808 and slightly-protruded ribs between them.

1809 **Phenology:** Flowers in May-June, fruits in July.

1810 **Reproduction:** Reproduces by seeds.

1811 **Distribution:** Toshkent, Namangan, Andijon, Farg'ona, Samarqand, Qashqadaryo, Buxoro and Surxondaryo provinces of
 1812 Uzbekistan; Chuy, Osh and Jalal-Abad provinces of Kyrgyzstan.

1813 **Habitat:** The adyr and tau zones. On dry, mixed-grass steppes and loess slopes. It is a typical component of the plant associa-
 1814 tions in these zones.

1815 **Population status:** Common.

1816 **Traditional use:** A decoction of the herb mixed with barley flour is used to treat various tumors. The ashes are applied to
 1817 treat eczema and scabies (Khalmatov 1964). A decoction of the herb is used to treat fever, flu, sore throat, pertussis,
 1818 stomach diseases, burns, and used as an anticonvulsive. It is also used as an insecticide to kill flies and cockroaches. It is
 1819 important to note that since the plant is very toxic, it should be used with extreme caution (Kulikov 1975; Khalmatov
 1820 et al. 1984; Khalmatov and Kosimov 1994).

1821 **Documented effects:** The preparation *Delsemine* has a muscle relaxing effect and has been used during surgery (Khodzhimatov
 1822 1989). Intravenous injection of the alkaloid delsemine at 0.5–2 mg/kg, was used to cause relaxation during surgery. At
 1823 doses of 5–6 mg/kg, delsemine was used to stop autonomous breathing. Presently, delsemine is not used in medical prac-
 1824 tice. Intravenous injection of the alkaloid licoctonine caused general calming with muscle relaxation, respiratory depres-
 1825 sion, and decrease of blood pressure while pulse rate remained the same (Tulyaganov et al. 1976). Intravenous introduction
 1826 of the alkaloid methyllycaconitine had a curariform effect on narcotized animals. Methyllycaconitine provoked brief
 1827 hypotensive effects. The alkaloid is used for spastic paresis of pyramidal character, postencephalitic arachnoencephalitis,
 1828 and spinal arachnoiditis (Dozortseva 1958, 1959). A preparation of methyllycaconitine, *Mellictin*, is used to treat
 1829 Parkinson's diseases and cerebral palsy (Khalmatov et al. 1984).

1830 **Phytochemistry:** All plant parts contain alkaloids. Before flowering, the top portion of the plant contained 0.25 % total
 1831 alkaloids and at flowering, 0.09 % total alkaloids. Delsemine, licoctonine, delphirine, methyllycaconitine, and anthra-
 1832 nolylycoctonine were isolated from the total alkaloids (Yunusov 1974). The flowers contained up to 4 % pigments. From
 1833 this, the flavonoids isorhamnetin, quercetin, and their glycosides were isolated (Khalmatov et al. 1984).



▲ *Datura stramonium* L. Photos: *left and center*: Vadim Prokhorov; *right*: Alim Gaziev

▼ *Daucus carota* L. Photos: *top*: Sasha Eisenman; *center*: Benjamin Zwitter; *bottom*: Evgeny Davkaev



▲ *Delphinium confusum* Popov
Photo: Stefano Doglio

► *Delphinium semibarbatum* Bienert. Photos: *top right*: Evgeny Davkaev; *lower right*: John B. Taft; *lower left*: Bazar Dovlet



1835 **Descurainia sophia (L.) Webb ex Prantl – Brassicaceae**

1836 **Synonyms:** *Hesperis sophia* (L.) Kuntze, *Sisymbrium sophia* L., *Sisymbrium tenuissimum* Kar. & Kir., *Sophia lobelii* Rupr.

1837 **English name:** Flixweed

1838 **Russian name:** Дескурения Софьи (Deskureniya Sof'i)

1839 **Uzbek name:** Shuvaran, sassyk кара

1840 **Kyrgyz name:** София дескурениясы (Sofiya deskureniyasy)

1841 **Description:** Herbaceous annual, grayish pubescent. Stem 10–90 cm tall, erect, unbranched or with spreading branches.

1842 Stem leaves alternate, bi- or tripinnatisect with linear, acute lobes; basal and lower leaves petiolate; upper leaves sessile.

1843 Inflorescence many-flowered raceme. Flowers 3–5 mm long with 4 sepals, 4 yellow petals, and 6 stamens, pedicellate.

1844 Fruit a silique, erect, slightly arcuate. Seeds light brown, 1–1.5 mm long, 0.5–0.75 mm wide.

1845 **Other distinguishing features:** The silique has a prominent vein along the septum.

1846 **Phenology:** Flowers and fruits from the beginning of April until the end of May.

1847 **Reproduction:** By seeds.

1848 **Distribution:** All regions of Uzbekistan and Kyrgyzstan.

1849 **Habitat:** The chul, adyr, and tau zones. As a weed, near roads, in fields, and in pastures.

1850 **Population status:** Common.

1851 **Traditional use:** A decoction of the plant is recommended as a febrifuge to treat laryngeal diseases, measles, and smallpox
1852 and is also used as a hemostatic. Fresh leaves are used to heal flesh wounds and are thought to have antibacterial action
1853 (Khalmatov 1964). The herb is often used to treat diarrhea and dysentery. Essence from the fresh flowering plant is used
1854 in homeopathy (Ogolevitz 1951). A decoction of the aboveground parts is used to treat throat diseases and as an anti-
1855 pyretic for smallpox and measles. In veterinary medicine, a decoction of the roots is used for diarrhea and helminthosis
1856 in cattle and horses. In Tibetan medicine, the roots are used for treatment of anthrax and ergotism. A tincture is used as a
1857 antihelmintic, diuretic, and hemostatic for internal hemorrhages (Bekker et al. 2005). A decoction of the herb is promoted
1858 as and considered a stimulant in the Russian Far East (Mamedov 2005).

1859 **Documented effects:** Experiments showed that a galenic preparation of this species reduced hypotension (Khalmatov 1964).

1860 An alcoholic extract of the seeds increases the tonus of muscles responsible for intestinal contractions. This extract is also
1861 used as a laxative to treat constipation (Khodzhimatov 1989).

1862 **Phytochemistry:** Leaves contain 10 mg% carotene, seeds contain 27–30 % oils and 1.5 % sinigrin glycoside, which produce
1863 0.8–0.9 % mustard essential oil after enzymic hydrolysis. The latter consists of 60 % benzyl isothiocyanate, 15 % allyl-
1864 isothiocyanate, and 5 % propenyl isothiocyanate (Khalmatov 1964). The seeds contain glucosides of quercetin, kaemp-
1865 ferol, and isorhamnetin as well as sinapic acid, ethyl ester, and trimethoxyl-cinnamic acid (Wang et al. 2004a), as well as
1866 lipids consisting of hydrocarbons, esters of fatty acids and cyclic alcohols, triacylglycerides, epoxyacylglycerides, free
1867 fatty acids, triterpenols, sterols, diacylglycerides and monoacylglycerides. Linolenic, linoleic, arachic, and erucic acids
1868 were the main components of the total lipids and triglycerides (Bekker et al. 2005).

1869

- Dianthus superbus L. – Caryophyllaceae** 1870
- Synonyms:** Some consider Central Asian populations to be a distinct species, *Dianthus hoeltzeri* Winkl. 1871
- English name:** Unknown 1872
- Russian name:** Гвоздика Гельцера (Gvozdika Gel'tsera) 1873
- Uzbek name:** Unknown 1874
- Kyrgyz name:** Гельцер чеге гул ("Gel'tser chege gul") 1875
- Description:** Herbaceous perennial with rhizomes. Stems of 2 kinds: non-elongated vegetative (non-reproductive) and 1876
reproductive, 15–60 cm tall. Leaves opposite, linear-lanceolate, 4–6 cm long, 2–4 mm wide, opposite blades connate at 1877
base, sheathing the stem. Sheath 2–4 mm long. Flowers solitary or in groups of 2–4. Calyx cylindrical, violet-tinged. 1878
Petals 5, light pink to dark pink, deeply fringed. Fruit a cylindrical capsule. 1879
- Other distinguishing features:** Capsule longer than calyx. 1880
- Phenology:** Flowers in June-August, fruits in July-September. 1881
- Reproduction:** By seeds. 1882
- Distribution:** All provinces of Kyrgyzstan; Toshkent Province of Uzbekistan. 1883
- Habitat:** In the tallgrass-meadow and forest belts, in glades and meadow slopes, and in subalpine meadows. 1884
- Population status:** Common, found as single plants. 1885
- Traditional use:** In folk medicine a decoction of the aboveground parts is drunk to treat heart diseases, gastrointestinal 1886
diseases, and uterine bleeding. The herb is also used to treat people bitten by rabid animals (Alimbaeva and Goncharova 1887
1971). A decoction of the aboveground parts and roots is used to treat various uterine diseases (Khalmatov 1964). 1888
- Documented effects:** No data. 1889
- Phytochemistry:** All plant parts contain saponins (triterpenes), alkaloids, tannins, flavonoids, and lipids (Boguslavskaya 1890
et al. 1983; Plant Resources of the USSR 1985). Phytoecdysteroids have been isolated from the plant (Saatov et al. 1891
1990). 1892

1894 **Dictamnus angustifolius G. Don fil. ex Sweet – Rutaceae**1895 **Synonyms:** *Dictamnus albus* ssp. *turkestanicus* Wint.1896 **English name:** Burning bush1897 **Russian name:** Ясенец узколистный (Yasenets uzkolistnyy)1898 **Uzbek name:** Togturbid1899 **Kyrgyz name:** Ичке жалбырактуу диктамнус (Ichke zhalbyraktuu diktamnus)

1900 **Description:** Herbaceous perennial. Stems 50–100 cm high, long- or short-hairy, but often glabrous. Leaves odd-pinnate
1901 with 5–6 pairs of large oblong or elongated-elliptic leaflets; leaflets with serrate margins and narrow-winged petiolules.
1902 Inflorescence racemose-paniculate, strongly glandular. Corollas lilac-purple, 3.5–4.5 cm long. Staminal filaments hairy,
1903 style glabrous. Ovary hairy. Seeds 4–5 mm long, black, shiny, smooth.

1904 **Other distinguishing features:** During dry weather, when the flowering plant is exposed to fire, it flares up, but the plant
1905 remains intact. Therefore people had named it “burning bush”.

1906 **Phenology:** Flowers in May-June, fruits in June-July.1907 **Reproduction:** By seeds.1908 **Distribution:** Toshkent and Farg’ona provinces of Uzbekistan; all of Kyrgyzstan.1909 **Habitat:** Among shrubs in the tau zone.1910 **Population status:** Uncommon, found as individuals.

1911 **Traditional use:** Avicenna used the plant as a purgative, but noted that it was poisonous and when taken in large doses
1912 (6–7 g), caused nausea, vomiting, and even death due to excessive vomiting. Avicenna also noted that some physicians
1913 prescribed the plant decoction to treat paralysis. An infusion of the leaves and flowers is used to treat rheumatism in
1914 Central Asian folk medicine. An aromatic water is made out of the flowers, which is used in cosmetics for facial skin care
1915 (Khalmatov 1964). *Dictamnus angustifolius*, growing in the Xin Jiang province of China, has been used as an alternative
1916 for *D. dasycarpus* in the treatment of rheumatism, bleeding, itching, jaundice, chronic hepatitis, and skin diseases, and as
1917 an anti-inflammatory agent, febrifugal, and detoxicant drug (Wu et al. 1999a).

1918 **Documented effects:** The root bark has antispasmodic, vermifugal, and antihysterical actions (Khalmatov 1964). The
1919 alkaloids dubinidine, evoxin, and skimmianine caused central nervous system depression resulting in sleep and then
1920 narcosis, when introduced in active doses into animals, and also had hypothermic action and increased pain threshold
1921 (Berezhinskaya and Trutneva 1959; Polievstev 1962a, b, Polievstev 1965; Polievstev et al. 1967; Sadritdinov 1968).
1922 In clinical tests at doses 0.6–0.8 g/day the alkaloid dubinidine had good sedative effect, especially on patients with
1923 severe insomnia. However, it was not recommended for clinical use (Polievstev 1965; Evdokimova and Kurmukov
1924 1972). The alkaloid dictamine was toxic when injected intravenously at 0.05–0.055 mg, evoked convulsions of rear
1925 extremities, decreased respiration, and eventually caused death of animals from asphyxia (Kovalenko 1946). Flavonoids
1926 from this species showed choleric, anti-inflammatory, and capillary strengthening activity (Komissarenko and
1927 Levashova 1988, 1989). A methanolic extract of the root bark of *Dictamnus angustifolius* showed significant vascular
1928 relaxing activity (Wu et al. 1999a).

1929 **Phytochemistry:** Roots contain 0.21 % and seeds contain 0.025 % total alkaloids. Skimmianine, dictamine, dubamine,
1930 dubinidine (roots), evoxin (haploperin), and other alkaloids were isolated from the total alkaloids (Yunusov 1974).
1931 Limonoids and coumarins have been isolated from the root bark (Wu et al. 1999a). The plant also contains essential oils
1932 and seeds contain 18–21 % drying oils (Khalmatov 1964).

- Dipsacus dipsacoides (Kar. & Kir.) Botsch. – Dipsacaceae** 1934
- Synonyms:** *Cephalaria dipsacoides* Kar. & Kir., *Dipsacus azureus* Schrenk. 1935
- English name:** Teasel 1936
- Russian name:** Ворсянка лазоревая (Vorsyanka lazorevaya) 1937
- Uzbek name:** Unknown 1938
- Kyrgyz name:** Когултур топчу баш (Kogultur topchu bash) 1939
- Description:** Herbaceous biennial or perennial. Stems 60–140 cm tall, branching above. Basal leaves lanceolate, up to 40 cm 1940
long, margins entire; lower leaves sessile, 15–30 cm long, 4–5 cm wide, pinnatifid towards leaf base; uppermost leaves 1941
smaller, becoming lanceolate or linear, sometimes pinnatisect. Inflorescences nearly spherical heads; involucre bracts 1942
awn-like, stiff, prickly. Corolla tubular, 4-lobed, bright blue, pubescent on the outside, each with a stiff green bract. Fruits 1943
4-sided achenes. 1944
- Other distinguishing features:** Inflorescence heads 2.5–4 cm long, with involucre bracts only slightly shorter. 1945
- Phenology:** Flowers in July, fruits in August-September. 1946
- Reproduction:** By seeds. 1947
- Distribution:** Chuy, Jalal-Abad, and Osh provinces of Kyrgyzstan; Toshkent, Andijon, Farg'ona, Samarqand and Surxondaryo 1948
provinces of Uzbekistan. 1949
- Habitat:** Among diverse grass assemblages on foothills, grassy steppes, and more often in bushy places. 1950
- Population status:** Common, forming dense groups. 1951
- Traditional use:** Infusions and decoctions of the herb are used to treat acute rheumatism, ulcers, and stomach cancer 1952
(Alimbaeva and Goncharova 1971). 1953
- Documented effects:** In experiment on animals (mice and rats), the preparation *Dipsacozide* (total plant saponins) was non- 1954
toxic and caused short-term decrease in arterial pressure. It noticeably raised the animals' tolerance to hypoxia, as found 1955
in foothill and high mountain conditions. In lipid metabolism *Dipsacozide* caused results similar to the known preparation 1956
Polysponin, and it also had hepatoprotective abilities (Alimbaeva et al. 1986). 1957
- Phytochemistry:** Roots contains glucose, lactose, organic acids, triterpene glycosides (18.9–31.8 %, hederagenin deriva- 1958
tives), alkaloids, vitamin C, phenolcarboxylic acids, coumarins, and flavonoids (2.18 %). The aboveground parts contain 1959
organic acid saponins (4.51–18.3 %, hederagenin derivatives), alkaloids (gentianine), phenolcarboxylic acids, coumarins, 1960
and flavonoids (0.5 %; Mukhamedziev and Alimbaeva 1969; Rakhmatullaev and Yunusov 1972a; Alimbaeva et al. 1986; 1961
Akimaliev et al. 1989; Putieva and Mukhamedziev 1998). 1962
- 1963



▲ **Dictamnus angustifolius G.**
Don fil. ex Sweet
Photos: Alim Gaziev



▲ **Dipsacus dipsacoides**
(Kar. & Kir.) Botsch.
Photos: Evgeny Davkaev



▲ **Descurainia sophia (L.)**
Webb ex Prantl Photos:
top: Denis Mirin;
bottom: Alim Gaziev



► **Dianthus superbus L.**
Photo: Sergey Appolonov

Dipsacus laciniatus L. – Dipsacaceae	1964
Synonyms: None	1965
English name: Cutleaf teasel	1966
Russian name: Ворсянка разрезная (Vorsyanka razreznaya)	1967
Uzbek name: Tungiztarok	1968
Kyrgyz name: Кесиктуу ворсянка (Kesiktuu vorsyanka)	1969
Description: Herbaceous biennial. Stems 50–150 cm tall; stems covered by prickles. Basal leaves elongate-obovate, toothed or pinnatilobate; cauline leaves opposite, up to 30 cm long, up to 15 cm wide; opposite leaves connate at base, forming a cup-shaped sheath; underside of midvein with prickles. Inflorescence a dense, elongate-oviform head; involucral bracts linear-lanceolate, coarse, prickly. Corolla tubular, 4 lobed, pale-blue to white, each with a stiff green bract. Fruits gray-brown achenes.	1970 1971 1972 1973 1974
Other distinguishing features: Inflorescence heads 5–8 cm long. Involucral bracts usually shorter than inflorescence head.	1975 1976
Phenology: Flowers in July, fruits in August-September.	1977
Reproduction: By seeds.	1978
Distribution: Chuy and Jalal-Abad provinces of Kyrgyzstan; Toshkent, Andijon, Farg'ona, Samarqand, and Surxondaryo provinces of Uzbekistan.	1979 1980
Habitat: In wet meadows and near canals.	1981
Population status: Common, found as single plants.	1982
Traditional use: An infusion of the roots is applied to treat tuberculosis and syphilis. A decoction, ointment, or paste is used as an anesthetic for hemorrhoids, calluses, and snake bites. Preparations of the aboveground parts are used as an anti-inflammatory and diuretic, and to stimulate respiratory function, cardiovascular function, and blood circulation. A decoction is used to treat fevers, ulcers, and stomach cancer, and is applied as compresses to treat skin cancer. A decoction of the inflorescence is used to treat rheumatism (Alimbaeva et al. 1986; Plant Resources of the USSR 1990).	1983 1984 1985 1986 1987
Documented effects: In experiment on animals, a preparation of the total saponins showed low toxicity and reduced arterial pressure for a short time (Alimbaeva and Goncharova 1971).	1988 1989
Phytochemistry: The roots contain triterpene glycosides (10 %), iridoids, and alkaloids (0.24 %). The aboveground parts contain triterpene glycosides (8.5 %), alkaloids (0.4 %), iridoid and phenolic glucosides, and flavonoids. Fruits contain iridoids (Alimbaeva et al. 1986; Abdallah 1991; Kocsis et al. 1993).	1990 1991 1992

1994 **Dodartia orientalis L. – Phrymaceae (formerly in Scrophulariaceae)**

1995 **Synonyms:** *Dodartia atro-coerulea* Pavlov.

1996 **English name:** Unknown

1997 **Russian name:** Додартия восточная (*Dodartiya vostochnaya*)

1998 **Uzbek name:** Takasoqol

1999 **Kyrgyz name:** Чыгыш теке сакалы (*Chygysh teke sakaly*)

2000 **Description:** Herbaceous perennial. Nearly leafless, lower leaves opposite. Stems multiple, erect, multi-branched, 25–40 cm
2001 high, younger shoots with curly hairs, becoming glabrous with age. Inflorescence a loose raceme. Flowers sessile, dark
2002 purple-violet. Calyx campanulate, 5-lobed. Corolla 1.6–2.2 cm long, 2-lipped, glabrous outside, bearded in the throat;
2003 lower lip 3-lobed, longer and broader than upper lip; upper lip short, erect. Stamens 4. Fruit a spherical capsule. Seeds
2004 multiple, oviform, deep-brown, 0.5–0.75 mm long, 0.5 mm wide.

2005 **Other distinguishing features:** Multi-branched, green, almost leafless herb with dark violet flowers. Middle lobe of lower
2006 lip smaller than lateral lobes.

2007 **Phenology:** Flowers in May-July, fruits in July-August.

2008 **Reproduction:** By seeds and rhizomes.

2009 **Distribution:** All of Uzbekistan and Kyrgyzstan.

2010 **Habitat:** The adyr and tau zones. Plains and slopes of hills.

2011 **Population status:** More often found as single individuals, but due to intensive vegetative propagation by rhizomes it grows
2012 as small but dense populations.

2013 **Traditional use:** A decoction made of the aboveground plant parts is used as a purgative and to treat syphilis (Khalmatov
2014 1964).

2015 **Documented effects:** The plant extract has slight purgative action, which is strengthened when mixed with other drugs
2016 (Ogolevitz 1951).

2017 **Phytochemistry:** The plant has barely been investigated chemically. The aboveground plant parts contain alkaloids and pos-
2018 sibly saponins (Khalmatov 1964). Mussaneoside [mussaenoside] has been isolated from this species (Umarova et al.
2019 1988).

2020

Eminium regelii Vved. – Araceae	2021
Synonyms: Some consider <i>E. regelii</i> a synonym of <i>Eminium lehmannii</i> Kuntze.	2022
English name: Unknown	2023
Russian name: ЭМИНИУМ Регеля (<i>Eminium Regelya</i>)	2024
Uzbek name: It kuchala, Korakulak	2025
Kyrgyz name: Тамыр кучала (<i>Tamyr kuchala</i>)	2026
Description: Perennial herb to 15–40 cm high, with a flat-spherical tuber, 3 cm in diameter. Leaves basal, light green, entire, oblanceolate to elliptic, the base wide-cuneate, sheathing, petiolate. Inflorescence a spadix; spathe tube 4–7 cm long, spathe blade ovate or oblong, inside velvety black-violet; spadix appendix 5–7 cm long, cylindrical, black-blue. Fruits subglobose berries, 1–2-seeded.	2027 2028 2029 2030
Other distinguishing features: Inflorescence produces the odor of rotten meat.	2031
Phenology: Flowering and fruits in April-May.	2032
Reproduction: By seeds and tubers.	2033
Distribution: Toshkent, Farg'ona, and Samarqand provinces of Uzbekistan; Osh province of Kyrgyzstan.	2034
Habitat: The adyr zone. Loess slopes of hills, on dry, shallow-soiled slopes with rocky debris.	2035
Population status: Not common, found as single individuals.	2036
Traditional use: The powdered tubers are used as an analgesic to treat rheumatism (Khalmatov 1964). The powdered tuber is also used internally to treat stomach aches, abdominal pain, internal diseases, and dysentery (Sezik et al. 2004; Pak 2005).	2037 2038 2039
Documented effects: An extract of the tubers had strophanthine-like action on the heart (Khalmatov 1964).	2040
Phytochemistry: The tubers contain poisonous saponins, traces of alkaloids and starch. The spathe contains pigments (Khalmatov 1964). The leaves and tuber contain a number of different lipids. The leaves contain carotinoids: neoxanthine and carotene (Chernenko et al. 2005).	2041 2042 2043
	2044

2045 **Ephedra equisetina Bunge – Ephedraceae**2046 **Synonyms:** *E. procera* var. *typica* Regel.2047 **English name:** Ephedra, ma huang2048 **Russian name:** Хвойник хвощевой (Khvoynik khvoshchevoy)2049 **Uzbek name:** Zogoza, Kizilcha2050 **Kyrgyz name:** Кырк муундай чекенде (Kyrk muunday chekende)

2051 **Description:** Large dioecious shrub, to 1.5(–2.5) m high, usually with a single, thick stem (occasionally multiple). Bark gray
2052 or brown, cracking, spongy; older branches thick, woody, erect; young branches green, opposite or whorled on older
2053 branches. Leaves opposite, scale-like, paleaceous, triangular, 2.5–3.5 mm long. Male inflorescences consist of pollen
2054 cones, solitary or in clusters of 2–4 at the nodes. Female cones usually opposite at nodes, each cone composed of overlap-
2055 ping bracts. Mature female cone berry-like, 6–7 mm long, spherical, fleshy, red. Seeds 4–6 mm long, elongate-ovoid, dark
2056 brown.

2057 **Other distinguishing features:** Leaves reddish to deep brown in color, connate for three fourth of their length.2058 **Phenology:** Flowers in May-June, fruits in July-August.2059 **Reproduction:** Propagates by seeds and rhizomes.2060 **Distribution:** Toshkent, Farg'ona, Samarqand, Qashqadaryo, Buxoro, and Surxondaryo provinces of Uzbekistan; all of
2061 Kyrgyzstan.2062 **Habitat:** The tau zone. Stony slopes in tree-shrub belt of mountains.2063 **Population status:** Common, sometimes forming dense groups.2064 **Traditional use:** For a long time, an infusion of the green shoots has been used for acute rheumatism, scabies, malaria, ulcers
2065 and other gastric diseases, altitude sickness, fever and heart diseases (Khalmatov 1964; Khalmatov et al. 1984). The
2066 stems, inflorescences, and berries are used as a treatment for bronchial asthma (Mamedov and Craker 2001).

2067 **Documented effects:** This plant is one of the main sources of ephedrine. The alkaloid d-pseudoephedrine (0.5 mg/kg intra-
2068 venously) evokes pressor action in narcotized animals. Repeated injections of the alkaloid usually cause tachyphylaxis.
2069 Pseudoephedrine has some properties of sympathomimetics of indirect action (Cession-Fossion 1967; de Meyts and
2070 Cession-Fossion 1966, 1967, 1968). In dogs, the alkaloid (1–2 mg/kg) provoked heartbeat deceleration, increased the
2071 blood pressure, and it also increased oxygen content in the blood of the coronary sinus. In general, pseudoephedrine has
2072 positive effects as a vasoconstrictor for rhinitis, tracheitis, and pharyngitis. Unlike L-ephedrine, pseudoephedrine has little
2073 or no effect on hemodynamics (Rowe et al. 1965). d-pseudoephedrine has direct stimulating effects on β -adrenoreceptors,
2074 and L-ephedrine has indirect stimulating effects (Tye et al. 1967).

2075 **Phytochemistry:** All plant parts contain alkaloids. Young shoots have up to 3.5 % alkaloids (ephedrine and pseudoephed-
2076 rine), tannins, vitamin C, and pigments. The stems contain up to 14 % tannins. The core of the wood stems contained from
2077 30 % to 65 % tannins. Seeds contained 4 pigments of the flavone series (Khalmatov 1964; Khalmatov et al. 1984).

2078




▲ **Dipsacus laciniatus L.** Photos: *left:* Andrei Lubchenko; *center and right:* Evgeny Davkaev




▲ **Ephedra equisetina Bunge**
Photo: Evgeny Davkaev

▼ **Dodartia orientalis L.** Photos: *left:* Evgeny Davkaev; *right:* Alim Gaziev



▼ **Eminium regelii Vved.** Photos: Evgeny Davkaev



2079 ***Ephedra intermedia* Schrenk & C.A. Mey. – Ephedraceae**

2080 **Synonyms:** *Ephedra ferganensis* V. Nikitin, *Ephedra glauca* Regel, *Ephedra microsperma* V. Nikitin, *Ephedra persica*
2081 (Stapf) V. Nikitin, *Ephedra tesquorum* V. Nikitin, *Ephedra tibetica* (Stapf) V. Nikitin, *Ephedra valida* V. Nikitin.

2082 **English name:** Ephedra

2083 **Russian name:** Хвойник средний, Эфедра средняя, Пустынная Эфедра (Kpвоуник sredniy, Efedra srednyaya, Pustynnaya
2084 efedra)

2085 **Uzbek name:** Kizilcha, Chul kizilcha

2086 **Kyrgyz name:** Орточо чекенде (Ortocho chekende)

2087 **Description:** Perennial, evergreen bush up to 1 m tall. Branches dense, erect, opposite or whorled, segmented, gray-green,
2088 glabrous; bark gray, fibrous. Leaves reduced, triangular to scale-shaped, opposite or in whorls, up to 3.5 mm long, leaves
2089 partially connate. Male cones usually clustered at nodes, subtended by circular or ovate bracts, connate at the base. Mature
2090 female cones berry-like, spherical, juicy, up to 6 mm long.

2091 **Other distinguishing features:** Has longer micropylar tubes than other *Ephedra* species; cones with 2–3 seeds.

2092 **Phenology:** Flowers in May and fruits in June-July.

2093 **Reproduction:** By seeds.

2094 **Distribution:** Surrounding Ysyk-Kol lake and in the Boom gorge of Kyrgyzstan; Toshkent, Farg'ona, Jizzax and Samarqand
2095 provinces of Uzbekistan.

2096 **Habitat:** On stony slopes in the lower and middle belt of mountains, and on rocks.

2097 **Population status:** Common, forming dense groups.

2098 **Traditional use:** This is one of the three species of *Ephedra* that is officially used in the Chinese Pharmacopoeia as a source
2099 of ma huang, a stimulant and antiasthmatic that has been used for at least 2,000 years (Abourashed et al. 2003).

2100 **Documented effects:** Ephedrine is obtained from the herb and is issued in the form of ephedrine chloride. It is widely used
2101 to treatment allergies (bronchial asthma, rashes, rhinitis, etc.). It acts by stimulating the central nervous system. In cases
2102 of morphine, scopolamine, and ganglioplegic poisoning, a preparation of ephedrine is used to raise arterial pressure, ren-
2103 der positive inotropic action on the heart, increase heart rate and to tone peripheral vessels, relax smooth muscles of
2104 bronchial tubes, and to stimulate breathing. The basic mechanism of ephedrine's activity is its ability to cause the libera-
2105 tion of noradrenaline from its reserves in nervous fibers and inhibit the return of noradrenaline to nervous fibers. In addi-
2106 tion, it protects noradrenaline and adrenaline from decomposition and strengthens their effects (Dobrokhotova and
2107 Chudinov 1966; Gammerman 1967). An extract of the plant exhibited antibacterial effects against *Micrococcus luteus* and
2108 *Klebsiella pneumoniae* (Shahidi Bonjar 2004).

2109 **Phytochemistry:** The thin, green stems contain up to 2.2 % total alkaloids, flavonoids, pigments, and up to 8 % tannins. Of
2110 the total alkaloids, up to 75 % is pseudoephedrine, with the rest being ephedrine and others (Khalmatov et al. 1984; Kim
2111 et al. 2005). The main constituent (12.80 %) of the essential oil isolated from the dried stems was 1,4-cineole (Ji et al.
2112 1997).

Epilobium hirsutum L. – Onagraceae	2114
Synonyms: <i>Chamaenerion hirsutum</i> (L.) Scop., <i>Epilobium tomentosum</i> Vent., <i>Epilobium velutinum</i> Nevski, <i>Epilobium villosum</i> Thunb.	2115 2116
English name: Codlins and cream, great willowherb, great hairy willowherb	2117
Russian name: Кипрей мохнатый, Кипрей волосистый (Kiprey mokhnatyy, Kiprey volosisty)	2118
Uzbek name: Kizilkon	2119
Kyrgyz name: Сапсагай кипрей (Sapsagay kiprey)	2120
Description: Herbaceous perennial. Stems up to 1.5 m tall, densely gray-hairy. Leaves sessile, oblanceolate, 4–10 cm long, 1–2 cm wide, margins serrulate, upper and lower surfaces densely pubescent. Flowers in a raceme. Calyx campanulate, lobes lanceolate, pubescent. Corolla lilac-purple, deeply lobed. Stigma deeply 4-lobed, recurved. Fruit a capsule, 4–10 cm long, pubescent. Seeds brown or light-brown, papillate.	2121 2122 2123 2124
Other distinguishing features: Leaves clasping the stem.	2125
Phenology: Flowers in June-August, fruits in August-September.	2126
Reproduction: By seeds and rhizomes.	2127
Distribution: All provinces of Kyrgyzstan; the Karakalpakstan autonomous republic (delta of the Amu-Darya river) and Toshkent province of Uzbekistan.	2128 2129
Habitat: In wet and marshy places near rivers and canals.	2130
Population status: Common, forming dense groups.	2131
Traditional use: The aboveground parts are used as a hemostatic, astringent, and anti-inflammatory (Vandisheva et al. 1977).	2132 2133
Documented effects: In experiments on animals, a water infusion caused reduced heart rate, increased amplitude of heartbeats, and caused diuresis (Appolonova 1956). Extracts of the plant exhibited a significant inhibitory effect on the reproduction of influenza viruses (Ivancheva et al. 1992), and prolonged the lifespan of mice with 2 types of tumorous cancers (Voynova et al. 1991).	2134 2135 2136 2137
Phytochemistry: The aboveground parts contain saponins, phenols, phenolcarboxylic acids, tannins, flavonoids (hyperoside, rutin, etc.), trace alkaloids, vitamin C, and coumarins (Plant Resources of the USSR 1987; Barakat et al. 1997).	2138 2139

2141 **Equisetum arvense L. – Equisetaceae**2142 **Synonyms:** *Equisetum boreale* Bong., *Equisetum calderi* B.Boivin, *Equisetum saxicola* Suksd.2143 **English name:** Field horsetail2144 **Russian name:** Хвощ полевой (Khvoshch polevoy)2145 **Uzbek name:** Kirk bugim2146 **Kyrgyz name:** Талаа кырк мууну (Talaa kyrk muunu)

2147 **Description:** Herbaceous, rhizomatous perennial, bearing spores. Stems dimorphic. Vegetative stems (developing later than
2148 sporebearing stems), 10–15 cm high, green, with 6–12 ribs, segmented with whorls of branches, with reduced leaves,
2149 forming a toothed sheath. Sporebearing stems (appearing in spring and die back after spores ripen), up to 40 cm tall,
2150 fleshy, reddish, brown or brownish-yellow, unbranched, topped with conical-cylindrical spore-bearing cones; sheathes
2151 longer than on vegetative stems.

2152 **Other distinguishing features:** Spores green, spherical.2153 **Phenology:** Spores ripen in April-May.2154 **Reproduction:** By spores and rhizomes.2155 **Distribution:** All provinces of Kyrgyzstan and Uzbekistan.2156 **Habitat:** In wet places and sandy meadows in valleys as well as in the lower and mid mountain belt.2157 **Population status:** Common, forming dense groups.

2158 **Traditonal use:** Preparations of the herb are used as a diuretic, anti-inflammatory, astringent, hemostatic, and disinfectant,
2159 and also to increase metabolism and treat skin wounds (Altimishev 1991). The herb is used to treat kidney and bladder
2160 diseases, edema, rheumatism, and stomach and intestinal growths (Kurochkin 1998).

2161 **Documented effects:** Preparations of this plant have been approved as a medicinal remedy. An infusion or liquid extract as
2162 a component of a tea mixture is used as a very strong diuretic for patients with kidney or heart diseases, to treat inflammation
2163 of the bladder and urinary tract, to stop stomach, intestinal, hemorrhoidal, and uterine bleeding, and as a treatment for
2164 pleurisy and some types of tuberculosis (Khalmatov et al. 1984). The compounds onitin-9-*O*-glucoside and luteolin, iso-
2165 lated from the plant, exhibited hepatoprotective activity in vitro, as well as strong superoxide scavenging effects (Oh et al.
2166 2004).

2167 **Phytochemistry:** The herb contains flavonoids (equisetrine, luteolin and glycosides of luteolin, quercetin, kaempferol, etc.),
2168 up to 5 % saponins (equisetonin), alkaloids (equisetin and nicotine) resins, organic acids (malic, aconitic, and oxalic), up
2169 to 25 % silicic acid, carotene, vitamin C, tannins, etc. (Khalmatov et al. 1984; Oh et al. 2004).

2170

- Eremurus regelii Vved. – Asphodelaceae (formerly in Liliaceae)** 2171
- Synonyms:** *Eremurus spectabilis* ssp. *regelii* (Vved.) Wendelbo. 2172
- English name:** Fox tail lily 2173
- Russian name:** Шириш Регеля (Shirish Regelya) 2174
- Uzbek name:** Shirach 2175
- Kyrgyz name:** Регель чырашы (Regel' chyrashy) 2176
- Description:** Herbaceous perennial with fusiform-incrassate roots. Stem glabrous, 80–180 cm high. Leaves wide-linear, 2.5–5 cm wide, 20–40 cm long, fluted, keeled, blue-gray, glabrous. Inflorescence a dense, multiflorous raceme, mostly erect while flowering. Flowers with 6 perianth segments, pale-pink, each with wide brownish-purple stripe. Fruits spherical capsules, latitudinally wrinkled, 6–8 mm in diameter. Seeds narrow-winged. 2177–2180
- Other distinguishing features:** When fruiting, pedicels arcuate, capsules crowded around inflorescence axis. 2181
- Phenology:** Flowers in May in the foothills, in June in the mountains. Fruits accordingly in June and August. 2182
- Reproduction:** Propagates by seeds and rhizomes. 2183
- Distribution:** Toshkent, Samarqand, Jizzax, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh and Jalal-Abad provinces of Kyrgyzstan. 2184–2185
- Habitat:** The adyr and tau zones, on gentle slopes. 2186
- Population status:** Common. 2187
- Traditional use:** In folk medicine the leaves are used as a carminative. Roots are used to treat gastrointestinal diseases and to coat the digestive tract. The roots are a source of native mannose. The polysaccharide eremuran is used to produced glucose and mannose by acid hydrolysis. A high quality glue is produced from the roots (Khalmatov and Kosimov 1992). The powdered rhizomes are used to treat psoriasis (Mamedov et al. 2004). 2188–2191
- Documented effects:** In acute tests on narcotized animals, the alkaloid hordenine, at doses of 0.5–1 mg/kg and higher, provoked rapid breathing. These effects are due to sympathomimetic (adrenomimetic) activity; it also has moderate vasoconstrictive action (Aliev et al. 1967; Khalmatov and Kosimov 1992). A polysaccharide, isolated from the roots, was found to increase the survival rate of rabbits subjected to hemorrhagic shock and had activity similar to that of reopoliglukin (Rakhimov 1997). 2192–2196
- Phytochemistry:** The roots contain the polysaccharide eremuran. The total alkaloid content of the roots was 0.035 % out of which 0.012 % was the alkaloid eremursine. The leaves contain vitamin C and carotene (Khalmatov 1964). The leaves and roots contain polysaccharides (Yuldasheva et al. 1993; Rakhimov 1997). 2197–2199

▶ **Equisetum arvense L.**
Photos: *left and center left:*
Vladimir Epiktetov; *center*
right: Dmitri Oreshkin;
right: Vadim Prokhorov



Eremurus regelii Vved. Photos: Evgeny Davkaev ▼



▲ **Epilobium hirsutum L.** Photos: Dmitri Oreshkin

▼ **Ephedra intermedia Schrenk & C.A. Mey.** Photos: John B. Taft



- Erodium cicutarium* (L.) L'Her. ex Aiton – Geraniaceae** 2201
- Synonyms:** *Erodium pulchellum* Karel. ex Ledeb., *Geranium cicutarium* L. 2202
- English name:** Redstem stork's bill, Redstem filaree 2203
- Russian name:** Аистник обыкновенный (Aistnik obyknovennyy) 2204
- Uzbek name:** Lailac tumshuk, qora mashaq 2205
- Kyrgyz name:** Цикутадай турна тумшук (Tsikutaday turna tumshchuk) 2206
- Description:** Ephemeral annual, with shortened vegetation cycle. Stems 10–60 cm tall, prostrate or upright, loosely villous. 2207
Leaves oblong, pinnatisect, segments pinnatipartite, stipulate. Inflorescences umbelliform, axillary. Sepals 5. Petals 5, 2208
4–6 mm long, purple-pink. Fruit a schizocarp, splitting into 5 mericarps, each attached to the styler column by a terminal 2209
awn. 2210
- Other distinguishing features:** Sepals apiculate. Fertile stamens 5, alternating with 5 staminodes. While drying, the awn 2211
twists spirally and separates from receptacle. If the soil has enough moisture in it, the terminal awn of the mericarp will 2212
penetrate into it. 2213
- Phenology:** Flowers and fruits in April-June. 2214
- Reproduction:** Abundantly propagates by seeds. 2215
- Distribution:** It is widespread in the irrigated farming zones of Uzbekistan; all of Kyrgyzstan. 2216
- Habitat:** The chul and adyr zones. Among wheat and alfalfa fields, vegetable gardens, orchards, on small hills, waste places 2217
and dry slopes. 2218
- Population status:** In some areas fairly dense populations can be found; especially in wheat farming areas, vegetable gar- 2219
dens and *Artemisia*-rich ephemeral complexes of the adyr zone. 2220
- Traditional use:** Central Asian folk medicine uses powdered leaves, as well as powder mixed with melted lamb fat to treat 2221
abscesses and as wound healing remedy. In the past the plant was widely used but its current use is limited (Khalmatov 2222
1964). In Iraq, a decoction of the whole plant is used for treatment of anasarca and metrorrhagia (Al-douri 2000). In 2223
Turkey, a decoction of the whole plant is used externally to treat pains (Simsek et al. 2004). 2224
- Documented effects:** A decoction is recommended as a hemostatic for internal uterine bleeding. An acetone-alcohol extract 2225
of the herb has been introduced as a hemostatic treatment (Aliev et al. 1972). Zavrazhanov et al. (1977) stated this species 2226
has astringent, anti-inflammatory, hemostatic, and sedative properties. The decoction of the herb is used to treat internal 2227
and uterine bleeding (Aliev et al. 1972) and also as an anticonvulsant (Fruentov 1972; Akopov 1981). A water extract, as 2228
well as a methanol extract and its fractions, were found to have antiviral effect on myxoviruses, herpes virus type 1, 2229
vesicular stomatitis and vaccinia virus (Zielinska-Jencylyk et al. 1987). In vivo, a methanol extract injected intravenously 2230
induced interferon in mice (Zielinska-Jencylyk et al. 1988). In vitro, low concentrations of a polyphenolic fraction from 2231
an extract of the plant stimulated free radical activity of human granulocytes, whereas high concentrations inhibited the 2232
activity (Fecka et al. 1997). 2233
- Phytochemistry:** The herb contains bitters, 2.1 % resins, tannins, acetylcholine, 55 mg% carotene, up to 4.94 % sugar, 1.9 % 2234
general titratable organic acids, 37.5–91.85 mg% vitamin C, and 0.64 mg% vitamin K, and 12–14 % ash, which includes 2235
up to 47 % K₂O (Akopov 1981). The aboveground parts contain a variety of tannins and flavonoids (geraniin, didehydrog- 2236
eraniin, corilagin, rutin, hyperin, quercetin, isoquercitrin, kaempferol, myricetin, polyphenolic acids, etc.) (Fecka and 2237
Cisowski 2005). 2238

2240 **Eryngium biebersteinianum Nevski – Apiaceae**

2241 **Synonyms:** *Eryngium caucasicum* Trautv. (some consider this the correct name), *Eryngium coeruleum* M. Bieb., *Eryngium*
2242 *pskemense* Pavlov

2243 **English name:** Bieberstein's sea-holly

2244 **Russian name:** Синеголовник Биберштейна (Sinogolovnik Bibershteyna)

2245 **Uzbek name:** Кок тыкан, koz tykan

2246 **Kyrgyz name:** Биберштейн тикен башы (Bibershteyn tiken bashy)

2247 **Description:** Glabrous perennial, bluish in color with widely fusiform root. Stems up to 1 m tall, often solitary. Branches
2248 emanating from above middle of stem and branching again, forming a wide, corymbose top. Basal leaves long-petiolate,
2249 thin-coriaceous, blue-gray, the blades oblong-oval with a cordate base; upper leaves sessile, deeply divided, margins
2250 spiny-dentate. Inflorescences subglobose heads up to 10 mm in diameter. Involucral bracts stiff, spiny, 2–4 times longer
2251 than heads. Petals blue, ca. 2 mm long. Fruits composed of obovate mericarps; mericarps angular, covered with long, nar-
2252 row, lanceolate scales along the edges.

2253 **Other distinguishing features:** First basal leaves have smooth margins and senesce early.

2254 **Phenology:** Flowers in May, fruits in July.

2255 **Reproduction:** Only by seeds.

2256 **Distribution:** Toshkent, Samarqand, Buxoro, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh and Jalal-Abad
2257 provinces of Kyrgyzstan.

2258 **Habitat:** The adyr zone. As a weed in orchards, vegetable gardens, unirrigated, cultivated fields and abandoned and long-
2259 fallow fields.

2260 **Population status:** Common, as single individuals.

2261 **Traditional use:** Infusions of the roots of other *Eryngium* species are used as a blood cleanser and as a sedative. The infu-
2262 sions are also used to treat edema, scrofula, gonorrhea, headaches, heart pain, and various tumors, and are used as a treat-
2263 ment for pertussis, anti-convulsant for epileptics, and as cough medicine, diaphoretic and diuretic. The roots are used to
2264 treat mushroom poisoning and bites from venomous animals. The herb is recommended for anemia (Khalmatov 1964;
2265 Minayeva 1991).

2266 **Documented effects:** This species has been shown to have expectorant action (Minayeva 1991).

2267 **Phytochemistry:** The plant contains essential oils, saponins, and tannids (Minayeva 1991; Khalmatov and Kosimov 1992).

2268

- Erysimum diffusum Ehrh. – Brassicaceae** 2269
- Synonyms:** *Erysimum canescens* Roth., *Erysimum andrzejowskianum* DC. 2270
- English name:** Diffuse wallflower 2271
- Russian name:** Желтушник рассеянный, Желтушник серый, Желтушник раскидистый (Zheltushnik rasseyannyu, Zheltushnik seryy, Zheltushnik raskidisty) 2272
- Uzbek name:** Kulrang zhyoltushnik 2274
- Kyrgyz name:** Чачырак даргын (Chachyrak dargyn) 2275
- Description:** Herbaceous biennial. Stems erect, single or few, 30–80 cm tall, sometimes branched. Basal rosette leaves petiolate, linear-lanceolate, margins entire; lower cauline leaves short-petiolate; upper cauline leaves sessile, margins entire. Inflorescence a few-flowered raceme. Flowers small, perfect, pedicellate. Petals 4, yellow. Stamens 6 (tetradynamous), erect. Fruits 4-sided siliques, thin, 3–10 cm long, 1–1.5 mm wide, whitish, hairy. Seeds ellipsoid, yellow-brown, up to 1.5 mm long. 2276–2280
- Other distinguishing features:** Fruit smooth, with 4 lines of white trichomes. 2281
- Phenology:** Flowers in June, fruits in July. 2282
- Reproduction:** By seeds. 2283
- Distribution:** All provinces of Kyrgyzstan; Toshkent, Namangan and Farg'ona provinces of Uzbekistan. 2284
- Habitat:** On steppes and dry stony exposures. 2285
- Population status:** Common, found as single plants. 2286
- Traditional use:** In Kyrgyzstan, an infusion of the herb is used as a diuretic, sedative and anti-depressant, and to treat heart problems. It is said to be one of the best treatments for edema (Altimishev 1991). In the folk medicine of Tajikistan, the aboveground parts are used to make a tea used as a diuretic and laxative, and to treat heart weakness, tachycardia, and hypertension (Khodzhimatov 1989). 2287–2290
- Documented effects:** The preparations *Erysimine*, *Erysimoside*, *Coreside*, liquid extracts, and *Cardiovalen* (a complex preparation) are used to treat mitral failure, hypertension, and arteriosclerotic cardiosclerosis (Khalmatov et al. 1984). 2291–2292
- Phytochemistry:** All plant parts contain cardiac glycosides. The greatest quantity is found in flowers and seeds (2–6 %). More than 10 cardiac glycosides have been isolated, including erysimine, erysimoside, and others. Seeds contains up to 30–40 % fatty oil (Tadzhibaev et al. 1977; Khalmatov et al. 1984). 2293–2295

2297 **Euphorbia jaxartica Prokh. – Euphorbiaceae**

2298 **Synonyms:** *Euphorbia virgata* Waldst. & Kit. ssp. *jaxartica* (Prokh.) Prokh., *Euphorbia waldsteinii* (Sojak) A. Radcliffe-
 2299 Smith ssp. *jaxartica* (Prokh.) Oudejans, *Tithymalus graminifolius* (Vill.) Sojak ssp. *jaxarticus* (Prokh.) Sojak.

2300 **English name:** Unknown

2301 **Russian name:** Молочай сырдарьинский (Molochay syrdar'inskiy)

2302 **Uzbek name:** Sultama

2303 **Kyrgyz name:** Сыр-Дарыя суттуу чобу (Syr-Daryya suttuu chobu)

2304 **Description:** Herbaceous perennial, 30–100 cm tall, blue-gray. Root thick, vertical or obliquely descending. Stems many or
 2305 few, erect, pubescent on lower portions, branching in upper parts with vegetative branches below the flowering branches.
 2306 Leaves alternate, nearly sessile, oblong-linear, 4–13 cm long, 2–7 cm wide, margins entire. Inflorescences cyathia, on
 2307 upper axile branches and on terminal peduncles arranged in umbels with 8–12 rays; bracts subtending inflorescences
 2308 opposite, partially connate, kidney-shaped or ovate-triangular, 6–20 mm long, 8–22 mm wide; cyathia campanulate with
 2309 ciliate lobes. Styles 2–3 mm long, connate nearly to the middle. Fruit an ovoid schizocarp, 3.5–4.5 mm long, 4–5 mm
 2310 wide, trisulcate. Seeds oval, 2.5 mm long, whitish-gray, smooth, with a small scarious appendage.

2311 **Other distinguishing features:** Leaves on vegetative branches densely arranged. Nectaries yellowish, crescent-shaped,
 2312 2-horned.

2313 **Phenology:** Flowers and fruits in April-September.

2314 **Reproduction:** Reproduces by rhizomes and seeds.

2315 **Distribution:** Toshkent, Namangan, Andijon, and Farg'ona provinces of Uzbekistan; Naryn Ysyk-Kol, Chuy and Talas
 2316 provinces of Kyrgyzstan.

2317 **Habitat:** The adyr and tau zones. In river valleys, along canals, in long-cultivated fields; often found along ravines into the
 2318 mountains, where it grows on stony slopes with rocky debris and in wet meadows.

2319 **Population status:** Common, found in small populations.

2320 **Traditional use:** The powdered root is used to treat wounds and syphilis. The latex is used to treat fungal skin diseases and
 2321 scabies, and to remove corns and warts (Khalmatov 1964).

2322 **Documented effects:** No data.

2323 **Phytochemistry:** The plant contains a large amount of latex, which contains 1.5 % caoutchouc, resins, and euphorbin
 2324 (Pavlov 1947). The plant contains triterpenes and polyphenols (quercetin-3-galactoside, kaempferol, gallic acid, etc.;
 2325 Azimov and Nazirov 1969, 1970; Abdulladzhanova et al. 2003).

2326



▲ Erodium cicutarium (L.) L'Her. ex Aiton Photos: *left*: Alim Gaziev; *center and right*: Evgeny Davkaev



◀ Eryngium biebersteinianum Nevski Photos: *left*: Komiljon Tojibaev; *right*: Vadim Prokhorov



▼ Euphorbia jaxartica Prokh. Photos: *left and center*: Alim Gaziev; *right*: Evgeny Davkaev



◀ Erysimum diffusum Ehrh. Photos: Maxim Zaitsev

2327 **Euphorbia rapulum Kar. & Kir. – Euphorbiaceae**

2328 **Synonyms:** *Tithymalus rapulum* (Kar. & Kir.) Klotzsch & Garcke.

2329 **English name:** Unknown

2330 **Russian name:** Молочай репчатый (Molochay repchatyy)

2331 **Uzbek name:** Ikhrozh

2332 **Kyrgyz name:** Туймоктуу суттуу чоп (Tuymoktuu suttuu chop)

2333 **Description:** Herbaceous perennial up to 25(–35) cm tall with a spherical, sometimes branching tuber, 3–6 cm in diameter.

2334 Stems erect, thicker towards the base; top of plant wide-paniculiform with bifurcating, flowering branches. Basal leaves
2335 ovate, sheathed; cauline leaves alternate, 3–4 cm long, 1–2 mm wide, spatulate or lanceolate-elliptic, sometimes cordate
2336 at the base, entire, short-petiolate to sessile. Inflorescences broadly campanulate cyathia, 2–3 mm in diameter, margin of
2337 lobes densely ciliate. Styles 1–1.5 mm long, connate at the base and forked at the top. Fruit an ovoid schizocarp, 4.5–
2338 5.5 mm long, 4–5 mm wide, trisulcate, glabrous, shiny. Seeds flattened-oblong, smooth, brownish, with a short-stalked
2339 conical appendage.

2340 **Other distinguishing features:** Upper flowering branches sometimes trifurcated.

2341 **Phenology:** Flowers and fruits in March–June.

2342 **Reproduction:** By seeds.

2343 **Distribution:** Toshkent, Samarqand, and Buxoro provinces of Uzbekistan; Chuy and Jalal-Abad provinces of Kyrgyzstan.

2344 **Habitat:** The adyr and tau zones. Clayey, stony slopes and slopes with red sandstone.

2345 **Population status:** Common, found as single individuals or in groups with 2–3 plants.

2346 **Traditional use:** Powdered root is used as a strong purgative and also for tuberculosis (Khalmatov 1964).

2347 **Documented effects:** In experiments, ethanolic extracts of the aboveground parts showed high antioxidant action (Eliseeva
2348 2005).

2349 **Phytochemistry:** The tuber contains upto 5 % resins and 0.5–0.6 % caoutchouc. Caoutchouc can also be found in the stems
2350 (up to 0.24 %) and in the fruits (up to 1.4 %). The resin contains the poisonous chemical euphorbin (Khalmatov 1964).

2351

Ferula foetida (Bunge) Regel – Apiaceae

Synonyms: *Ferula assa-foetida* L., *Peucedanum asa-foetida* (L.) Baill., *Scorodosma foetidum* Bunge.

English name: Giant fennel

Russian name: Ферула вонючая (*Ferula vonyuchaya*)

Uzbek name: Sassyk kavrak, kovrak, sassyk kurayi

Kyrgyz name: Жыттуу ала гул (*Zhyttuu ala gul*)

Description: Herbaceous, monocarpic perennial, with a large, oval, fleshy root up to 15 cm in diameter. Stem thick, up to 1–1.2 m high, upper portion branching and forming a dense globose panicle. Leaves mostly glabrous above, more or less soft-villous beneath, senescing early; basal leaves short-petiolate with broad blade, ternate with bipinnatisect lobes, lobules decurrent, 15 cm long, 5 cm wide; lower leaves alternate; upper leaves smaller and becoming reduced to sheaths. Inflorescences compound umbels; terminal umbel sessile or on a reduced peduncle, spherical, 15–20 cm wide; lateral umbels on long peduncles. Petals light-yellow, almost cream colored. Fruit a schizocarp with 2 one-seeded mericarps; mericarps flattened, pubescent, 1.6–2.2 cm long and wide.

Other distinguishing features: Ovary and fruit pubescent. Seeds have an extremely objectionable, persistent odor.

Phenology: Flowers in March-April, fruits in April-May.

Reproduction: By seeds.

Distribution: Desert areas of Karakalpakstan autonomous republic, Toshkent, Samarqand, Buxoro, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; not found in Kyrgyzstan.

Habitat: The adyr zone. Plains in foothills, on stony-clay soils.

Population status: Common, found as single individuals.

Traditional use: The plant has been used in Central Asian folk medicine since ancient times, as an anticonvulsant, vermifuge, and to treat some nervous diseases. The gum-resin is used in Chinese medicine as a restorative and tonic for hysterics, neurasthenia and vegetative neurosis, and to treat some skin diseases and common colds, as an expectorant and anticonvulsant, and mixed with other drug substances to treat lung tuberculosis, exudative diathesis, lymphadenitis, and osteitis. Avicenna used this plant to treat tumors, jaundice, and other liver diseases, as well as stomach, kidney, and spleen diseases, and as a diuretic and hemostatic for uterine bleeding (Khalmatov and Khabibov 1976; Kurmukov and Akhmedkhodzhaeva 1994).

Documented effects: Extracts of various *Ferula* species and individual compounds isolated from the extract exhibit phytoestrogenic activities. Based on these compounds 2 phytoestrogenic preparations, *Tefestrol* and *Panoferol*, were developed (Prokhorova and Kurmukov 1997; Prokhorova et al. 1992b; Kurmukov and Akhmedkhodzhaeva 1994). Infusions, pills, and emulsions of the gum-resin are used as an antispasmodic asthma treatment, to treat hysteria and other nerve diseases, and as an anticonvulsant. An infusion of *Ferula*, injected intravenously, reduced blood pressure. Its hypotensive activity is due to antispasmodic action on blood vessels (Sarkisyants 1969a, 1972). Dried resin of the roots reduced platelet adhesiveness and aggregative properties, depressed blood thromboplastic activity and elongated time and intensity of bleeding (Mansurov 1967). An infusion and decoction of *Ferula foetida* stimulated stomach secretory activity, and also had an impact on activity of the gastrointestinal tract. Thirty to thirty-five days of treatment with a water infusion (10–20 % by weight) of the herb, at 0.5–1 g/kg animal mass, prevented animal death from anaphylactic shock and development of Arthus-Sakharov phenomenon, i.e. it shows anti-allergic affect (Isakov 1969; Sarkisyants 1969b; Sarkisyants and Azizova 1971; Kurmukov and Akhmedkhodzhaeva 1994). In Kuwait this species is used as an treatment for diabetes and has hypolipidemic activity (Al-Awadi and Shoukry 1988).

Phytochemistry: In the early 1930s coumarins and organic sulfides were isolated from *Ferula* spp. (Tsukervanik et al. 1935; Kurmukov and Akhmedkhodzhaeva 1994). Later, umbelliferone, ferulic and galbanic acids and coumarins were isolated from the resins (Kurmukov and Akhmedkhodzhaeva 1994). In a systematic study of 50 species of *Ferula* in Uzbekistan, more than 250 terpenoids were isolated. It was shown for the first time that plants of this genus contained complex esters of terpenoid alcohols with aliphatic and aromatic acids. The structure and stereochemical abilities of more than 150 new terpenoids were determined (Saidkhodzhaev and Nikinov 1973, 1974; Sagitdinova and Saidkhodzhaev 1977; Sagitdinova et al. 1978). The compounds isolated from species in the genus *Ferula* can be divided into 3 groups: (1) coumarins, (2) compound esters of terpenoids and sesquiterpenoid alcohols with aromatic acids, and (3) sesquiterpenoid lactones (Bagirov et al. 1978). The roots of all the species found in Central Asia have similar chemical compounds to that of *F. foetida* and contain resins, essential oil, gums, high amounts of starch, and other compounds (Kurmukov and Akhmedkhodzhaeva 1994; Khalmatov and Kosimov 1994). The gum-resin of *F. foetida* contains 4–28 % essential oils: disulfide, hexenyl-disulfides, paraoxycoumarins, 0.68 % free asaresinotannol, asaresinol and their ether with ferulic acid, umbelliferone (which is formed from ferulic acid), asaresin A, farnesferol A, B, C, and other substances (Kurmukov and Akhmedkhodzhaeva 1994).

2352

2353

2354

2355

2356

2357

2358

2359

2360

2361

2362

2363

2364

2365

2366

2367

2368

2369

2370

2371

2372

2373

2374

2375

2376

2377

2378

2379

2380

2381

2382

2383

2384

2385

2386

2387

2388

2389

2390

2391

2392

2393

2394

2395

2396

2397

2398

2399

2400

2401

2402

2403

2405 **Ferula kuhistanica Korovin – Apiaceae**2406 **Synonyms:** *Ferula jaeschkeana* Vatke.2407 **English name:** Unknown2408 **Russian name:** Ферула кухи́станская (*Ferula kukhistanskaya*)2409 **Uzbek name:** Chair2410 **Kyrgyz name:** Unknown

2411 **Description:** Herbaceous perennial, monocarpic, with thick oviform root. Stem thick, stocky, up to 1 m high, upper third
2412 branching into thick, oviform panicle. Leaves quickly senescing, abaxial side glabrous, hairy beneath; leaf blades are
2413 wide-triangular in outline, ternate with bipinnatisect lobes, lobules oblanceolate. Inflorescences compound umbels;
2414 umbels of 2 kinds: the terminal umbel nearly sessile, with 20–25 rays, up to 12 cm wide; lateral umbels long-pedunculate,
2415 in clusters of 3, exceeding the terminal umbel. Petals yellow. Fruit a schizocarp with 2 one-seeded mericarps; mericarps
2416 flattened, oval, 2–3,2 cm long and 1–2,2 cm wide, reddish-brown.

2417 **Other distinguishing features:** Ovary and fruit glabrous.2418 **Phenology:** Flowers in May-June, fruits in July-August.2419 **Reproduction:** By seeds.

2420 **Distribution:** Andijon, Namangan, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Naryn
2421 and Ysyk-Kol Provinces of Kyrgyzstan.

2422 **Habitat:** The tau and yailau zones. Gentle mountain slopes of the tree-shrub belt.2423 **Population status:** Rare, forming dense groups.

2424 **Traditional use:** The plant's resin, boiled with milk, is used to treat syphilis. It is applied externally as a treatment for per-
2425 sistent wounds, tumors, and other diseases (Khalmatov and Kosimov 1992; Kurmukov and Akhmedkhodzhaeva 1994).

2426 **Documented effects:** Similar to *Ferula foetida*. Compounds isolated from the fruits were toxic against gram-positive bacte-
2427 ria, including methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* (Tamemoto et al. 2001).

2428 **Phytochemistry:** Refer to *Ferula foetida* for general information on the genus *Ferula*. All plant parts of *F. kuhistanica* con-
2429 tain resin and essential oils. The fresh roots contain 0.42–0.72 % essential oils, the fruits 0.54 % and the fresh leaves
2430 0.08 %. From steam distillation, 11.7–14.8 % green-colored, strong smelling essential oils were extracted. Leaf oil con-
2431 tains 85 % d-pinene. Roots contain up to 28 % and fruits 10–11 % resins. The resin contains n-carbolic acid (12.5 %),
2432 anisic and angelic acids, and umbelliferone (Khalmatov 1964). Daucane-type sesquiterpenes and daucane esters have
2433 been isolated from the roots and stems (Chen et al. 2000).

2434

- Ferula moschata* (Reinsch.) Koso-Pol. – Apiaceae** 2435
- Synonyms:** *Ferula pseudo-oreoselinum* (Regel & Schmalh.) Koso-Pol., *Ferula sumbul* (Kaufm.) Hook. f., *Ferula urceolata* 2436
Korov. 2437
- English name:** Musk fennel 2438
- Russian name:** Ферула сумбул (*Ferula sumbul*) 2439
- Uzbek name:** Sumbul 2440
- Kyrgyz name:** Unknown 2441
- Description:** Herbaceous perennial, with multiple, thick, conjoined taproots. Stems few, up to 50 cm high, slender, pubes- 2442
cent becoming subglabrous, corymbiform branching above. Leaves stiff, persisting long into the growing season, abaxial 2443
side slightly hairy, long-petiolate; basal leaves oval-triangular in outline, blade tripinnatisect, leaf segments lanceolate or 2444
oblong, 20–30 mm long, 10–15 mm wide, entire or deeply dissected; cauline leaves becoming smaller, upper leaves 2445
reduced to sheaths. Inflorescences compound umbels; umbels variable; terminal with 6–10 rays, 4–6 cm wide; lateral 2446
umbels single or in pairs, distinctly below level of terminal umbel. Petals yellow. Fruit a schizocarp with 2 one-seeded 2447
mericarps; mericarps 7 mm long, twice as long as the pedicels, flattened, with filiform ribs. 2448
- Other distinguishing features:** Umbellets 10–15-flowered. Fractured roots produce a specific pleasant smell. 2449
- Phenology:** Flowers in June, fruits in July. 2450
- Reproduction:** By seeds. 2451
- Distribution:** Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Jalal-Abad province of Kyrgyzstan. 2452
- Habitat:** The tau and yailau zones. Stony open slopes among shrubs. 2453
- Population status:** Rare, found as single individuals; listed in the Red Book of Rare and Endangered Species of 2454
Uzbekistan. 2455
- Traditional use:** Used as a folk medicine in Eastern and European countries. The resin was often used as a tonic and as a 2456
stimulatory remedy for gastric pneumatosis, pertussis, cholera, and other diseases. Avicenna applied it to treat tumors, 2457
jaundice and other diseases of the liver, stomach, kidneys, and spleen, and he also used it as a diuretic and hemostatic for 2458
uterine bleeding (Ogolevitz 1951; Kurmukov and Akhmedkhodzhaeva 1994; Khalmatov and Kosimov 1994). 2459
- Documented effects:** Similar to *Ferula foetida*. Compounds isolated from a methanol extract of the dried roots of *Ferula* 2460
sumbul showed anti-HIV activity (Zhou et al. 2000). 2461
- Phytochemistry:** See *Ferula foetida* for general information on the genus *Ferula*. The roots contain up to 21.5 % resins 2462
containing phytosterin, vanillic acid, umbelliferone, fatty acids (including isovaleric), up to 4 % essential oil, consisting 2463
of linalyl acetate, citronellyl acetate, ferulene, sesquiterpenes, doremon, doremol and its acetic ester, as well as the ses- 2464
quiterpene sambulene and up to 24.41 % total sugars (Tsukurvanik and Simkhaev 1948; Khalmatov 1964). The dried 2465
roots contained many different coumarins and sesquiterpene lactones (Zhou et al. 2000; El-Razek et al. 2001). 2466



▲ *Euphorbia rapulum* Kar. & Kir. Photos: Evgeny Davkaev



▲ *Ferula foetida* (Bunge) Regel Photos: Alim Gaziev



◀ *Ferula moschata* (Reinsch.) Koso-Pol.
Photo: Red Book of Uzbekistan



▲ *Ferula kuhistanica* Korovin
Photos: Alim Gaziev

- Fragaria vesca L. – Rosaceae** 2468
- Synonyms:** *Fragaria chinensis* Losinsk., *Fragaria concolor* Kitag., *Potentilla vesca* (L.) Scop. 2469
- English name:** Woodland strawberry 2470
- Russian name:** Земляника лесная (Zemlyanika lesnaya) 2471
- Uzbek name:** Yavoiy klubnay 2472
- Kyrgyz name:** Токой кожогаты (Tokoy kozhogaty) 2473
- Description:** Herbaceous, stoloniferous perennial, 5–30 cm tall. Stems compressed, hairy. Leaves in basal rosette, petiolate, trifoliolate, margins sharply toothed, lateral leaflets sessile, middle leaflet often short petiolulate. Inflorescence cymose, on an elongated stem. Sepals 5, appressed hairy, margins entire. Petals 5, white. Stamens many. Fruits small achenes, attached to surface of swollen receptacle. Receptacle berry-like, bright red, fleshy, 0.7–2 cm in diameter, elongated or nearly spherical. 2474–2478
- Other distinguishing features:** Runners develop in the axils of leaves. Pedicels appressed hairy. 2479
- Phenology:** Flowers in May, fruits in June. 2480
- Reproduction:** By seeds and runners. 2481
- Distribution:** All provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 2482
- Habitat:** In the forest belt of mountains, spruce-fir forests, and glades. 2483
- Population status:** Common, found in small groups. 2484
- Traditional use:** In folk medicine infusions of fruits and leaves are used to treat sore throats, jaundice, hemorrhoids, fatigue, uterine bleeding, and children with diarrhea. Fresh leaves are applied to old skin ulcers (Akopov 1990). Fresh fruits are used to treat kidney stones, inflammation of the gall bladder and bile duct, gout, stomach catarrh, constipation, hypertension, and arteriosclerosis, and is used as a vermifuge. Crushed fruits are applied to the skin to treat eczema. A decoction of the dried fruits is used as a diaphoretic and of the leaves as a diaphoretic and diuretic. A decoction of the roots is used as a hemostatic (Altimishev 1991). 2485–2490
- Documented effects:** Berries of wild strawberry possess tonic, anti-inflammatory, diuretic, antiscorbutic, choleric, and hypoglycemic properties (Maznev 2004). An alcoholic extract of the aboveground parts increased the cellular mass of the spleen and thymus, protected the mucus membrane of the stomach and decreased stress in cyclophosphane-treated mice, as well as exhibited antiulcer and stress-protective effects (Aksinenko et al. 2003; Klimentova et al. 2005). 2491–2494
- Phytochemistry:** Fruits contain pectins, sugars, citric, malic and phosphoric acids, carotene, vitamin C, essential oils, and over 10 microelements, including iron. Leaves contain of vitamin C (high amounts), carotene, tannins, flavonoids, many different organic acids, essential oils, and up to 20 micro- and macroelements. The roots are rich in tannins and iron salts (Poludenny and Zhuravlev 2000). 2495–2498

2500 **Fumaria vaillantii Loisel. – Fumariaceae (Papaveraceae)**2501 **Synonyms:** *Fumaria vaillantii* var. *schrammii* (Asch.) Velen.2502 **English name:** Earthsmoke2503 **Russian name:** Дымянка Вайяна (Думуанка Ваууана)2504 **Uzbek name:** Shotara2505 **Kyrgyz name:** Вайлант фумариясы (Vaylant fumariyası)

2506 **Description:** Annual herb with taproot. Stems 10–35 cm tall, erect or reclining, branching from the base. Leaves alternate,
2507 long-petiolate, tri-pinnatisect, segments linear or linear-lanceolate, margins entire. Inflorescences terminal or leaf-opposed
2508 racemes. Flowers zygomorphic with 2 small sepals. Corolla pink-violet, darker towards the apex, 5–6 mm long. Petals 4,
2509 in 2 whorls, 1 outer petal with a short spur. Fruits indehiscent capsules, subglobose, 1.5–2.5 mm in diameter, tuberculate-
2510 wrinkled, 1-seeded.

2511 **Other distinguishing features:** Staminal filaments connate into 2 groups.2512 **Phenology:** Flowering and fruits in March-July.2513 **Reproduction:** By seeds.2514 **Distribution:** In all regions of Uzbekistan and Kyrgyzstan.2515 **Habitat:** The adyr and tau zones. Mainly as a weed amongst crops, in orchards and in abandoned fields.2516 **Population status:** Common, especially in abandoned fields.

2517 **Traditional use:** A decoction of the herb is taken as a blood-cleanser and diuretic, to treat coughs, jaundice, headache, fever,
2518 gonorrhoea, uterine bleeding, erysipelas, and for clearing the intestines. It is also used externally in a bath to treat itching,
2519 rashes, and pimples (Khalmatov et al. 1984).

2520 **Documented effects:** The alkaloid protopine (fumarine) caused narcosis in amphibians and, in mammals, caused paralysis
2521 of sensory nerve endings and increased reflex excitability. The alkaloid slightly increased the effects of analeptics and
2522 induced catalepsy (Chen-Gu 1957; Cheney 1963). In acute experiments with animals under narcosis, reduced heart rate
2523 and increased heartbeat amplitude occurred and, for a short time, decreased blood pressure was observed. Protopine has
2524 antiarrhythmic action with better effects than novocainamide and quinidine (Sadritdinov and Kurmukov 1980). In a
2525 screen to determine effects on platelet aggregation, extracts of this species showed complete inhibition of aggregation.
2526 This result was found to be caused by protopine (Sener 1994). Extracts of the dried plant displayed high rates of inhibition
2527 against the enzymes acetylcholinesterase and butyrylcholinesterase, which are associated with Alzheimer's disease
2528 (Orhan et al. 2004).

2529 **Phytochemistry:** The aboveground parts contain alkaloids (protopine, vaillantine, parfumine, fumaridine, fumvailline, etc.),
2530 sugars, resins, pigments, fumaric acid, traces of essential oil, and vitamins C and K₁ (Ibragimova et al. 1974; Khalmatov
2531 et al. 1984; Khodzhimatov 1989). The seeds contain phospholipids (Gazizov and Glushenkova 1997).

2532

Galium septentrionale Roem. & Schult. – Rubiaceae	2533
Synonyms: some consider <i>G. septentrionale</i> a subspecies of <i>G. boreale</i> [<i>G. boreale</i> ssp. <i>septentrionale</i> (Roem. & J. A. Schult.) H. Hara].	2534
English name: Northern bedstraw	2535
Russian name: Подмаренник северный (Podmarennik severnyy)	2536
Uzbek name: Chakamoog	2537
Kyrgyz name: Тундук галиум (Tunduk galium)	2538
Description: Herbaceous perennial, with rhizomes. Stems 30–70 cm tall. Leaves in whorls of 4, 4.5–5.5 cm long, 0.7–0.8 cm wide, elongate-lanceolate, 3-nerved. Inflorescence a dense, many-flowered, terminal panicle. Flowers small, white. Corolla rotate, 4-lobed. Fruits bristly nutlets.	2539
Other distinguishing features: Stems glabrous.	2540
Phenology: Flowering in June-August, fruits in July-September.	2541
Reproduction: By seeds.	2542
Distribution: All provinces of Kyrgyzstan; not found in the flora of Uzbekistan.	2543
Habitat: In glades, among shrubs, and on river banks.	2544
Population status: Common, forming dense groups.	2545
Traditional use: Infusions and decoctions of the aboveground parts are used to treat deafness, malignant tumors, and applied to eyes to treat conjunctivitis. In Tibetan medicine a decoction of the aboveground parts is used to treat heart diseases, gastritis, and gynecological diseases. The rhizomes are used to treat pneumonia and gynecological diseases (Plant Resources of the USSR 1990).	2546
Documented effects: In experiments with frogs a tincture decreased heart beat amplitude (Turova and Nikolskaya 1954).	2547
Phytochemistry: The roots contain steroid saponins, tannins, flavonoids, coumarins, and anthraquinones. The aboveground parts contain essential oils, triterpene acids, iridoids, steroid saponins, alkaloids, tannins, coumarins, anthraquinones, and vitamin C (Revina and Shustova 1982).	2548
	2549
	2550
	2551
	2552
	2553
	2554
	2555
	2556
	2557

2558 **Galium verum L. – Rubiaceae**2559 **Synonyms:** *Galium glabratum* Klokov.2560 **English name:** Yellow spring bedstraw, Lady's Bedstraw2561 **Russian name:** Подмаренник настоящий (Podmarennik nastoyashchiy)2562 **Uzbek name:** Tilkisoomai2563 **Kyrgyz name:** Кадимки галиум (Kadimki galium)

2564 **Description:** Herbaceous perennial, with branched rhizomes. Stems 30–125 cm tall, thin. Leaves in whorls of 8–12, narrow,
2565 linear, 1–4 cm long, 0.5–3 mm wide, 1-nerved, apex acute, margins sometimes recurved. Inflorescence a long, dense-
2566 flowered panicle. Flowers bright-yellow. Corolla rotate, 4-lobed. Fruits 2-parted.

2567 **Other distinguishing features:** Flowers smell like honey.2568 **Phenology:** Flowers in June-July, fruits in July-August.2569 **Reproduction:** By seeds.

2570 **Distribution:** Ysyk-Kol and Chuy provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Toshkent, Samarqand,
2571 Farg'ona, Andijon and Surxondaryo provinces of Uzbekistan.

2572 **Habitat:** In steppes and meadow-steppes.2573 **Population status:** Common, forming dense groups.

2574 **Traditional use:** An infusion of the fresh aboveground parts is drunk as a hemostatic, analgesic, sedative, and diuretic for
2575 people with swelling associated with heart or kidney diseases. A bath or compresses soaked with the infusion are used to
2576 treat rheumatism, various skin diseases, scrofula, and furunculosis. The rhizomes are used as to strengthen the libido. In
2577 Tibetan medicine, the rhizomes are used to treat pneumonia and liver diseases (Shreter 1975; Akopov 1990).

2578 **Documented effects:** In vitro, ethanolic extracts of the plant showed low to moderate cytotoxic activity in human lympho-
2579 blastoid Raji cells (Spiridonov et al. 2005). Rubiadin exhibited antifungal and antituberculosis activity, as well as cyto-
2580 toxicity to BC and NCI-H187 cancer cell lines (Kanokmedhakul et al. 2005). Asperuloside has laxative effects
2581 (Milkowska-Leyck et al. 1999).

2582 **Phytochemistry:** The herb contains anthraglycosides and anthraquinones (galiosin, rubiadin, asperuloside, etc.), flavonoids,
2583 traces of essential oils, tannins, and dyeing substances. Rhizomes contain iridoids, steroid glycosides, coumarins, and
2584 flavonoids (Akopov 1990; Muzychkina 2000; Demirezer et al. 2006; Tamas et al. 2006; Zhao et al. 2008). Cultivated cal-
2585 lous tissue produced a variety of different anthraquinones (Banthorpe and White 1995).

2586



◀ **Fragaria vesca L.**

Photos: Sergey Appolonov



▲ **Fumaria vaillantii Loisel.**

Photos: Alexander Naumenko



▼ **Galium septentrionale**

Roem. & Schult. Photos:

Vadim Prokhorov



▶ **Galium verum L.** Photos:

top and center: Alim Gaziev;

bottom: Natalie Schultz



2587 **Gentiana olivieri Griseb. – Gentianaceae**2588 **Synonyms:** *Gentiana regeliana* Gand., *Gentiana weschniakowii* Regel.2589 **English name:** Unknown2590 **Russian name:** Горечавка Оливье (Gorechavka Oliv'ye)2591 **Uzbek name:** Gazakut, erbahasi2592 **Kyrgyz name:** Оливье кок базини (Oliv'ye kok bazini)

2593 **Description:** Herbaceous perennial with thin rhizomes. Stems several, erect, 10–40 cm high, glabrous, smooth, rounded,
2594 sheathed at the base with the older basal leaves. Basal leaves in a rosette, elongate-lanceolate to elongate-spatulate,
2595 2–12 cm long and 0.4–1 cm wide, green on both sides, glabrous; cauline leaves opposite, 2–3 pairs, lanceolate or narrow-
2596 lanceolate. Inflorescences terminal corymbiform cymes with 1–6 flowers. Corolla conical with 5 lobes, bluish-violet, dark
2597 blue or pale blue, rarely white, 2–4 cm long. Fruit an oblong capsule, 1–2 cm long, 2-valved. Seeds many, small, wingless,
2598 seed coat thick, surface reticulately patterned.

2599 **Other distinguishing features:** Stamens inserted just below middle of corolla tube. This species has multiple forms distin-
2600 guished by their pedicel lengths.

2601 **Phenology:** Flowers in May-June, fruits in June-July.2602 **Reproduction:** By seeds and vegetative rhizomes.

2603 **Distribution:** Toshkent, Samarqand, Jizzax, Buxoro, Qashqadaryo, Farg'ona, Andijon, and Surxondaryo provinces of
2604 Uzbekistan; Naryn, Osh, Chuy and Talas provinces of Kyrgyzstan.

2605 **Habitat:** The adyr and tau zones. On dry slopes with rocky debris.2606 **Population status:** Common, usually found in small populations.

2607 **Traditional use:** Decoction of the flowering herb is used for gastric diseases, malaria, toothaches, bleeding gums, and as an
2608 oral rinse, as well as is applied externally to treat abscesses and tumors. Syrup, made by boiling gentian and barberry roots
2609 for a long time, is recommended for side pains, rheumatic pain and chest pains (Khalmatov 1964).

2610 **Documented effects:** The alkaloid gentianine, at doses of 50 mg/kg and higher, had sedative effects. At doses of 150–
2611 200 mg/kg it had central muscle relaxant action. At 10–25 mg/kg, the alkaloid noticeably prolonged the activity of
2612 soporifics in experiments on mice, eliminated aggressive reaction in rats, provoked a hypothermic effect, depressed devel-
2613 oped conditioned reflexes and decreased stimulant action of caffeine and benzedrine (i.e., it has sedative and tranquilizing
2614 effect; Tulyaganov and Sadritdinov 1968; Tulyaganov et al. 1971; Danilevskii et al. 1972; Sadritdinov and Kurmukov
2615 1980). The alkaloids gentianadine, gentianamine, and oliverine had anti-inflammatory action in rabbits and rats (Sadritdinov
2616 and Tulyaganov 1967, 1972; Sadritdinov 1971a). In experiments with rats, extracts of the plant exhibited hepatoprotective
2617 effects (Orhan et al. 2003). Methanolic extracts of the plant exhibited significant hypoglycemic effects on hyperglycemic
2618 rats (Sezik et al. 2005).

2619 **Phytochemistry:** The aboveground plant parts contain alkaloids and bitter glycosides (Ersoz and Calis 1991; Orhan et al.
2620 2003; Sezik et al. 2005). The plants around Toshkent had the following bitter index: leaves 1:20,000, flowers 1:5,000,
2621 stems 1:2,500, and the total aboveground parts 1:5,000. From the aboveground parts collected in the Toshkent province
2622 (village of Kaplanbek), 0.35 % total alkaloids were isolated and these included gentianine, gentiananine, gentianine,
2623 gentianadine, gentioflavine, gentiotibetine, oliverine, oliveridine, oliveramine, and others (Rakhmatullaev and Yunusov
2624 1972b; Yunusov 1974).

Geranium collinum Steph. ex Willd. – Geraniaceae	2626
Synonyms: <i>Geranium minutum</i> Ikonn., <i>Geranium saxatile</i> Kar. & Kir., <i>Geranium wakhanicum</i> (Pauls.) Ikonn.	2627
English name: Geranium	2628
Russian name: Герань холмовая (Geran' kholmovaya)	2629
Uzbek name: Anzhabor	2630
Kyrgyz name: Шалбай каз таманы (Shalbay kaz tamany)	2631
Description: Herbaceous perennial. Stems 15–55 cm tall. Leaves hairy, palmate, circular in outline with 3–7 lobes divided more than three fourth to the base, lobes nearly rhomboid, pinnatifid with 3–5 lobules. Flowers in small groups at end of axillary branches. Sepals 5, oblong-elliptic, 4–10 mm long. Petals 5, obovate, 10–19 mm long, pink-violet, apex rounded, claw ciliate. Fruit an elongated, beaked capsule covered with short hairs.	2632 2633 2634 2635
Other distinguishing features: Beak up to 3 cm long.	2636
Phenology: Flowers in May-July, fruits in June-August.	2637
Reproduction: By seeds.	2638
Distribution: All provinces of Kyrgyzstan; Tashkent, Andijon, Farg'ona, Samarqand and Surxondaryo provinces of Uzbekistan.	2639 2640
Habitat: In steppes, in wet meadows in the forest-meadow mountain belt, along canals, in orchards, and in boggy places.	2641
Population status: Common, forming dense groups.	2642
Traditional use: The underground parts are used as an astringent and hemostatic. A decoction is used to treat gastric diseases, hemoptysis, and diarrhea, and also as a hemostatic after childbirth (Khalmatov 1964). In folk medicine it used internally to treat malignant tumors, broken bones, and fever (Amirov 1974).	2643 2644 2645
Documented effects: In experiments on animals preparations from the leaves inhibited malignant tumors (Amirov 1974).	2646
Phytochemistry: Underground parts of the plant contains tannins and phenols (pyrogallol and pyrocatechin). The aboveground parts contain flavone glycosides, saponins, alkaloids, and tannins (12–27.2 %). The whole plant is rich in tannins (Chumbalov et al. 1968; Chumbalov and Bikbulatova 1970; Plant Resources of the USSR 1988).	2647 2648 2649

2651 **Geum rivale L. – Rosaceae**2652 **Synonyms:** None2653 **English name:** Purple avens, water avens2654 **Russian name:** Гравилат речной (Gravilat rechnoy)2655 **Uzbek name:** Shirchai2656 **Kyrgyz name:** Ийилген гулду геум (Iyilgen guldu geum)

2657 **Description:** Herbaceous perennial, with thick rhizomes. Stems 20–70 cm tall. Leaves in basal rosette, petiolate, sparsely
2658 hairy, lyrate-pinnately compound, with 3–7 leaflets, terminal leaflet largest and lobed; cauline leaves smaller, simple to
2659 3-lobed, stipulate. Inflorescence terminal, corymbiform, 2–4-flowered, often nodding. Flowers 5-merous, pedicels pubes-
2660 cent. Sepals reddish-purple. Petals yellow with reddish brown-purple veins. Stamens and carpels numerous, styles plu-
2661 mose. Fruits long-beaked achenes in a globose aggregate; achenes fusiform, 3–4 mm long, yellow villous.

2662 **Other distinguishing features:** Achenes have a hooked style to aid in dispersal.2663 **Phenology:** Flowers in June, fruits in July.2664 **Reproduction:** By seeds.2665 **Distribution:** All provinces of Kyrgyzstan; Toshkent province of Uzbekistan.2666 **Habitat:** In forest glades, in long-used animal corrals in the tallgrass-meadow belt of mountains, and in valleys and along
2667 brooks.2668 **Population status:** Common, forming dense groups.

2669 **Traditional use:** Infusions and decoctions of the herb are used to treat paradontosis, stomatitis, laryngitis, stomach catarrh,
2670 dysentery, vomiting, and intestinal colitis. Infusions and decoctions of the rhizomes are used to treat headaches, insomnia,
2671 eye diseases, rheumatism, and hemorrhoids, and is effective against snake venom (Krilov 1972).

2672 **Documented effects:** Extracts of this plant showed anti-inflammatory activity in vitro (Tunon et al. 1995).

2673 **Phytochemistry:** Rhizomes contain carbohydrates (glucose, arabinose, and ketose), pectins, organic acids (6.46 %), essen-
2674 tial oils, saponins, alkaloids, vitamin C, and tannins. The leaves contain vitamin C, carotene, and tannins. The flowers
2675 contain tannins (7.35 %) and the fruits contain carbohydrates (Blinova 1957; Aliev et al. 1961). The roots were found to
2676 contain small amounts of proanthocyanidins and high amounts of ellagic acid (Oszmianski et al. 2007).

2677

- Glaucium fimbrilligerum Boiss. – Papaveraceae** 2678
- Synonyms:** *Dicranostigma iliense* C.Y. Wu & H. Chuang, *Glaucium luteum* var. *fimbrilligerum* (Boiss.) Trautv. 2679
- English name:** Unknown 2680
- Russian name:** Глауциум бахромчатый (*Glaucium bakhromchaty*) 2681
- Uzbek name:** Urmon kora 2682
- Kyrgyz name:** Туктуу саргалдак (*Tuktuu sargaldak*) 2683
- Description:** Annual or biennial herb. Stems 8–65 cm tall, branched, leafy. Lower leaves lyrate-pinnatisect, 5–30 cm long; upper leaves clasping, many-lobed. Flowers solitary, axillary. Buds 15–20 mm long, glabrous. Petals bright yellow, lacking spots, wide-obovate or round, 1.5–3.7 cm long, 2.5–3.5 cm wide, margin wavy. Fruit a silique-like capsule, 10–25 cm long, up to 0.4 cm wide, dehiscent from the top nearly to the base, straight or arching, sparsely covered with appressed bristles. Seeds kidney-shaped, 1.5–2 mm long, brown. 2684–2688
- Other distinguishing features:** Pedicels up to 20 mm long in fruit. Capsules with 2 horns at the tip. 2689
- Phenology:** Flowers and fruits in May–July. 2690
- Reproduction:** Seeds. 2691
- Distribution:** Toshkent, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh, Chuy and Talas provinces of Kyrgyzstan. 2692–2693
- Habitat:** The adyr and tau zones. On clay bluffs, stony slopes and in dry river beds with rocky debris. 2694
- Population status:** Rare. 2695
- Traditional use:** The crushed, roasted seeds are recommended as a hemostatic and tonic for women after childbirth. The oil has the same abilities. A decoction of the leaves and flowers, as a tea, is given as a tonic and stimulant for people recovering from diseases. Large doses have emetic and soporific effects, but can cause asphyxiation. The seeds are considered a strong laxative (Khodzhimatov 1989). 2696–2699
- Documented effects:** In acute experiments with animals under anesthesia, reduced heart rate and increased heartbeat amplitude occurred and, for a short time, decreased blood pressure was observed. Protopine has antiarrhythmic action with better effects than novocainamide and quinidine (Sadritdinov and Kurmukov 1980). The alkaloid corydine has a general activity similar to that of bulbocarpine, and like bulbocarpine can cause catalepsy (Berezhinskaya et al. 1968). In acute experiments with animals, the alkaloid chelerythrine produced 2 phases of action on arterial pressure: hypertensive effects due to the alkaloids influence on the vasomotor center and hypotensive effects due to the alkaloids influence on the muscle walls of vessels. Chelerythrine has analgesic activities, potentiates analgetic action of morphine and elongates sleep produced sleeping preparations (Chelombito and Muravyova 1971). 2700–2706
- Phytochemistry:** The entire plant contains alkaloids (protopine, corydine, sanguinarine, corytuberine, glauvine, glaunine, norcorydine, isoboldine, etc.). The seeds contain up to 30 % drying fatty oil (Yunusov et al. 1973; Yunusov and Israilov 1974; Karimova et al. 1980, 1983; Khodzhimatov 1989; Shafiee et al. 1998). 2708–2710



▲ **Gentiana olivieri** Griseb.
Photo: Stefano Doglio



▲ **Glaucium fimbriigerum** Boiss. Photos: Evgeny Davkaev
▼ **Geum rivale** L. Photos: Alexander Naumenko



▼ **Geranium collinum** Steph. ex Willd. Photos: *left*: Alim Gaziev; *center*: Sasha W. Eisenman;
right: John B. Taft



Gleditsia triacanthos L. – Fabaceae

- Synonyms:** *Acacia americana* Cat. Long. ex Stokes, *Acacia triacanthos* (L.) Gron., *Caesalpiniodes triacanthum* (L.) Kuntze, *Gleditsia brachycarpa* (Michx.) Pursh, *Gleditsia bujotii* Neumann, *Gleditsia elegans* Salisb., *Gleditsia hebecarpa* S. McCoy, *Gleditsia heterophylla* Raf., *Gleditsia horrida* Salisb., *Gleditsia inermis* L., *Gleditsia meliloba* Walter, *Gleditsia micracantha* Loddiges ex Steudel, *Gleditsia polysperma* Stokes, *Gleditsia spinosa* Marsh, *Gleditsia triacanthus* (L.) Mill., *Melilobus heterophylla* Raf. 2712-2717
- English name:** Honey-locust 2718
- Russian name:** Гледичия обыкновенная (*Gledichiya obyknovennaya*) 2719
- Uzbek name:** Tikandaraxht 2720
- Kyrgyz name:** Кадимки гледичия (*Kadimki gledichiya*) 2721
- Description:** Deciduous tree, 12–20 m tall. Trunk and branches with long, simple or branched, reddish-brown, 2–8 cm long thorns. Leaves of 2 kinds: pinnate and bipinnate, petioles pubescent; once-pinnate leaves on short lateral spurs; bipinnate leaves on long shoots; leaflets oblanceolate. Inflorescences perfect or staminate, in separate racemes arising from the short lateral spurs. Flowers yellow-green, calyx and petals pubescent. Fruit a dark brown legume, flat, often slightly twisted, up to 40 cm long. Seeds elongate-elliptic, up to 15 mm long. 2722-2726
- Other distinguishing features:** Has sweet pulp between the seeds. Thornless cultivars exist and are used as an ornamental plant. 2727-2728
- Phenology:** Flowers in May, fruits in September. 2729
- Reproduction:** By seeds. 2730
- Distribution:** Native to North America. Cultivated nearly everywhere in Kyrgyzstan and Uzbekistan. 2731
- Habitat:** Only cultivated. 2732
- Population status:** Common. 2733
- Traditional use:** The plant is used to treat spastic colitis, chronic cholecystitis, stomach ulcers, and bronchial asthma (Rakhmanberdyeva et al. 2002). 2734-2735
- Documented effects:** In experiments the alkaloid triacanthine showed hypotensive and antispasmodic activity. The antispasmodic actions occurred in the bronchial smooth muscles and the intestines. The saponin triacanthocide showed anti-arrhythmic action in experiments (Khalmatov et al. 1984). A preparation of triacanthine is used to treat digestive system problems (Altimishev 1991). 2736-2739
- Phytochemistry:** Young leaves contain up to 1 % of the alkaloid triacanthine and flowers contain up to 0.3 %. Leaves contain up to 400 mg% ascorbic acid. Fruits contain olmelin, fustin, and no less than 10 triterpene glycosides. The fruit walls contain around 2.6 % anthraglycosides, 3.1 % tannins, and traces of essential oil. The pulp of the fruits contain up to 29 % sugars, and the seeds contain up to 39 % mucilage, carbohydrates, lipids, fatty acids (palmitic, oleic, linoleic, and linolenic), carotinoids, etc. (Khalmatov et al. 1984; Rakhmanberdyeva et al. 2002). 2740-2744

2746 **Glycyrrhiza glabra L. – Fabaceae**2747 **Synonyms:** *Glycyrrhiza glandulifera* Waldst. & Kit., *Glycyrrhiza hirsuta* Pall., *Glycyrrhiza violacea* Boiss. & Noë.2748 **English name:** Common licorice2749 **Russian name:** Солодка голая (Solodka golaya)2750 **Uzbek name:** Kizilmiya, Chuchuk miya, Shirin miya2751 **Kyrgyz name:** Тукуз кызыл мыя (Tukuz kyzyl myya)

2752 **Description:** Herbaceous perennial with deep root system down to 5 m. Stems erect, simple or branched, 45–120 cm high,
2753 sparsely short-hairy with scattered glands or glandular prickles. Leaves alternate, odd-pinnate, 5–20 cm long with (2–)3–9
2754 pairs of leaflets; leaflets oblong, ovate or lanceolate, 2–4 cm long, 0.8–2 cm wide, with glands on abaxial side. Inflorescences
2755 loose racemes, 5–12 cm long. Flowers 8–12 mm long. Calyx 5-lobed, upper 2 lobes half as long as lower 3. Corolla pap-
2756 ilionaceous, whitish-violet. Fruit a legume, 2–7-seeded, straight or slightly curved, glabrous or with dense glandular
2757 prickles. Seeds small, 3 mm in diameter, almost round, smooth, deep-brown.

2758 **Other distinguishing features:** Stamens 10 (9 united). Interior of root is lemon-yellow and has a specific sweet taste.2759 **Phenology:** Flowers in April–July, fruits in May–June.2760 **Reproduction:** By seeds and rhizomes.2761 **Distribution:** All regions of Uzbekistan and Kyrgyzstan.2762 **Habitat:** The chul and adyr zones. River banks, embankments, along canals, salty-soiled areas (salanchaks), tugai, on gentle
2763 slopes of mountains and foothills, and in melon and cotton fields as a weed.2764 **Population status:** Common, sometimes found in large groups.

2765 **Traditional use:** The plant root has been used to treat various diseases since ancient times. Decoctions and extracts of the
2766 root are used as a diaphoretic and purgative and also to treat cough, chest pains, and other diseases. Avicenna recom-
2767 mended the roots to treat renal, lung, and bladder diseases, as well as gastritis, fever, and other diseases. The root decoc-
2768 tion is used for throat dryness and spasms and as an expectorant for coughs and respiratory tract catarrh (Khalmatov et al.
2769 1984).

2770 **Documented effects:** Modern medicine uses preparations made of the roots (syrup, thick and dry extracts), as well as in
2771 combinations with other substances such as a mixture of powders, as a slight laxative, expectorant, and to coat the stom-
2772 ach. The powder, thick and dry extracts, and root syrup are widely used in pharmaceutical practice to make pills, improve
2773 mixture taste, and for other purposes. It was established that the active ingredients of the roots (glycyrrhizic and glycyrr-
2774 hetinic acids) have antispasmodic and antihistamine activities, similar to adrenal hormones (deoxycorticosterone and
2775 hydrocortisone) and are recommended to treat skin diseases and inflammatory processes (Mashkovskii 1984). The prepa-
2776 rations have tonic and adaptogenic activities and are useful for recovery of general health and memory improvement
2777 (Kurmukov 1976). Licochalcone-A, an estrogenic flavonoid found in licorice root has been shown to effectively inhibit
2778 proliferation of prostate cancer cells (Fu et al. 2004). Isoliquiritigenin inhibited platelet aggregation and aldose reductase
2779 activity in vivo (Aida et al. 1990; Tawata et al. 1992), and in vitro, inhibited proliferation and induced apoptosis in prostate
2780 cancer cell lines (Kanazawa et al. 2003; Jung et al. 2006a, b).

2781 **Phytochemistry:** Underground organs contain 4.6–23 % glycyrrhizin, up to 10.5 % sugars, up to 8.1 % bitters (glycyrrama-
2782 rin), flavonoids (liquiritin, liquirazide, liquitigenin and 2'-4,4'-trihydroxychalcone and its glycoside isoliquiritigenin),
2783 glabric acid, 0.035 % essential oil, β -sitosterol, extriol, 1–4 % asparagines, dyes, and other substances (Kurmukov 1976;
2784 Mashkovskii 1984).

Glycyrrhiza uralensis Fisch. ex DC. – Fabaceae	2786
Synonyms: <i>Glycyrrhiza asperrima</i> var. <i>desertorum</i> Regel, <i>Glycyrrhiza asperrima</i> var. <i>uralensis</i> (Fisch. ex DC.) Regel, <i>Glycyrrhiza glandulifera</i> Ledeb.	2787 2788
English name: Chinese licorice	2789
Russian Name: Солодка уральская (Solodka ural'skaya)	2790
Uzbek name: Shirinmiya	2791
Kyrgyz name: Урал кызыл мыясы (Ural kyzyly myyasy)	2792
Description: Herbaceous perennial, with large rhizome. Stems simple or branched, 40–70 cm high, short pubescent with punctuate glands or raised glands. Leaves alternate, odd-pinnate, 10–25 cm long; leaflets 3–8 pairs, 2–6 cm long, 1.5–3.5 cm wide, obovate or elliptic. Inflorescences densely flowered, axillary racemes. Flowers 1.5–2.5 cm long. Calyx 8–14 mm long, toothed, pubescent. Corolla papilionaceous; petals violet, banner petal rounded (cupped) or sinuate. Fruits crescent-shaped legumes, 2–4 cm long, in dense, tangled clusters. Seeds round to kidney-shaped, brown, smooth.	2793 2794 2795 2796 2797
Other distinguishing features: Has a more dense-flowered raceme and larger flowers than <i>Glycyrrhiza glabra</i> .	2798
Phenology: Flowers in May-June, fruits in August-September.	2799
Reproduction: By seeds and rhizomes.	2800
Distribution: Southern and eastern parts of Kyrgyzstan; Surxondaryo province of Uzbekistan.	2801
Habitat: In meadows with relatively high water tables and along canals and rivers.	2802
Population status: Common, found in dense groups.	2803
Traditional use: The underground parts are used as a diuretic, laxative, and carminative, and to treat pneumonia, bronchitis, asthma, and ulcers, and also as a remedy for poisoning (Khodzhimatov 1989).	2804 2805
Documented effects: Similar to <i>Glycyrrhiza glabra</i> . Because of the high flavonoid content in the above and below ground parts of <i>Glycyrrhiza uralensis</i> , it is used as raw material for antispasmodic and anti-ulcer preparations (Khalmatov et al. 1984). Extracts of the root exhibited apoptotic effects on human breast cancer cells (Jo et al. 2004).	2806 2807 2808
Phytochemistry: The aboveground parts have up to 3.3 % total flavonoids. The below ground parts have up to 4.3 % total flavonoids (glycyrrhizic acid, glycyrrhetic acid, fermononetin, isoliquiritigenin, etc.; Tolmachev 1976; Nakanishi et al. 1985; Wang et al. 2004b).	2809 2810 2811

2813 **Haplophyllum acutifolium (DC.) G. Don. f. – Rutaceae**

2814 **Synonyms:** *Haplophyllum flexuosa* Boiss., *Haplophyllum perforatum* (M. Bieb.) Kar. & Kir., *Haplophyllum sieversii* Fisch.,
 2815 *Ruta acutifolia* DC., *Ruta flexuosa* (Boiss.) Engl., *Ruta perforata* M. Bieb., *Ruta sieversii* (Fisch.) F. Fedtsch.

2816 **English name:** Unknown

2817 **Russian name:** Цельнолистник остролистный (Tsel' nolistnik ostrolistnyy)

2818 **Uzbek name:** Toshbakatol, Tashbakftol

2819 **Kyrgyz name:** Unknown

2820 **Description:** Herbaceous perennial, covered with punctate glands. Stem erect, corymbiform- branching, glabrous, 20–70 cm
 2821 high. Leaves alternate, simple, broadly-oblong to narrowly-lanceolate, entire, glabrous, short-petiolate. Inflorescence
 2822 paniculate-corymbiform, multiflorous. Calyx lobes 5, ovate-triangular, acute, glabrous or slightly pubescent. Petals 5,
 2823 yellow, 3.5–5 mm long, tapering to a claw. Stamens 10. Style glabrous; ovary sessile. Fruit a capsule with indehiscent
 2824 deciduous segments, on a very short stipe, densely tuberculate.

2825 **Other distinguishing features:** Pellucid dots on leaves observable when held up to the light. Leaves produce a specific
 2826 objectionable odor when bruised.

2827 **Phenology:** Flowers in May-June, fruits in July-August.

2828 **Reproduction:** By seeds and rhizomes.

2829 **Distribution:** Toshkent, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; not found in
 2830 Kyrgyzstan.

2831 **Habitat:** The adyr and tau zones. Shallow soils, rarely on stony slopes.

2832 **Population status:** Common.

2833 **Traditional use:** Preparations of the plant are used in folk medicine as a sedative for anxiety and cardiac neurosis, as well
 2834 as for hysterics, epilepsy, gastric spasms, and menstrual period disturbance (Kovaleva 1971). A leaf decoction is used to
 2835 treat toothaches, chest and stomach diseases, and for bloated abdomens. A decoction and infusion of the herb, together
 2836 with decoction of common wormwood (*Artemisia absinthium*), is used as a bath to treat various skin diseases (Khalmatov
 2837 1964).

2838 **Documented effects:** The plant contains the alkaloids perforine, haplofolin, and haplofilidine, which in small doses produce
 2839 sedative effects, in medium doses produce soporific effects, and in large doses, causes narcosis. These alkaloids are
 2840 strongly pronounced antagonists against some analeptics (camphor, strychnine, and caffeine) and reinforce actions of
 2841 some soporifics and narcotics. Haplofilidine eliminated fear in tested rats, but perforine did not have such an action
 2842 (Akhmedhodjaeva and Polievtsev 1963; Danilevskii et al. 1972; Akhmedhodjaeva 1978). The majority of alkaloids con-
 2843 tained in the plant have estrogenic activity (Akhmedhodjaeva 1978). When tested for in vitro cytotoxicity, extracts of the
 2844 aboveground plant parts had strong cytotoxic activity against multiple types of cancer cell lines (Varamini et al. 2007).

2845 **Phytochemistry:** Plants collected in the Qashqadaryo province of Uzbekistan contained varying amounts of total alkaloids
 2846 during different phenologic periods: alkaloid content of leaves during flower bud stage was 0.1 %; alkaloid content of
 2847 leaves during flowering stage was 0.4 %; alkaloid content of leaves during seed maturation was 1.2 %, alkaloid content in
 2848 stems was 0.075–0.14 %; alkaloid content at the stage of full fruit maturity, in roots was 0.025 % and in seeds was 1.6 %.
 2849 From different parts of plants growing in several regions of Uzbekistan, 25 alkaloids were isolated, including evoxin,
 2850 skimmianine, haplofilidine, perforine, haplopine, flindersine, glycoferine, methyl-evoxin, evodine, evoxidine,
 2851 haplofidine, anhydroperforine, perfamine, foliosidine, dubinidine, etc., and the lignan eudesmine (Razzakova et al. 1973,
 2852 1986; Yunusov 1981). Kusunokinin, β -sitosterol, oleanolic acid, cholesterol and hexadecanoic acid, as well as the alka-
 2853 loids haplophytin-A and B, were isolated from the plant (Ali et al. 2001).



▲ *Gleditsia triacanthos* L. Photos: *left*: Georg Slickers; *center*: Andrew Butko; *right*: Luis Fernández García ▼ *Glycyrrhiza uralensis* Fisch. ex DC. Photos: Alexander Naumenko



▼ *Haplophyllum acutifolium* (DC.) G. Don. f. Photo: Evgeny Davkaev



▲ *Glycyrrhiza glabra* L. Photos: Maxim Kucherov

2855 **Helichrysum maracandicum Popov ex Kirp. – Asteraceae**2856 **Synonyms:** None2857 **English name:** Unknown2858 **Russian name:** Цмин самаркандский, Бессмертник самаркандский (Tsmin samarkandskiy, Bessmertnik samarkandskiy)2859 **Uzbek name:** Samarqand buznoch2860 **Kyrgyz name:** Самарканд очпос гулу (Samarqand ochpos gulu)

2861 **Description:** Herbaceous perennial. Stems 15–75 cm tall, striated, hairy. Leaves alternate, greenish, gray-green, to yellow-
2862 green, densely hairy; basal and cauline leaves linear to linear-lanceolate, apex very acute, margins entire, base partly sheath-
2863 ing stem. Inflorescences capitula, 5 mm wide, spherical, semispherical, campanulate, or elliptic; capitula in groups of 20–80
2864 and forming dense clusters or compact corymbiform structures. Involucral bracts 40–60 in 5 rows, stiff-membranaceous,
2865 yellow. Flowers 50–80 per capitulum; corollas yellow. Fruits dark-brown achenes with pappus of 20–25 very thin, whitish-
2866 yellow bristles.

2867 **Other distinguishing features:** The entire plant is densely hairy. Outer involucral bracts lanceolate to elliptical, more
2868 numerous than inner bracts. Inner bracts spatulate, glabrous, shiny.

2869 **Phenology:** Flowers in June, fruits in September-October.2870 **Reproduction:** By seeds.

2871 **Distribution:** Naryn, Chuy, Talas, and Jalal-Abad provinces of Kyrgyzstan; Toshkent, Andijon, Farg'ona, Samarqand and
2872 Surxondaryo provinces of Uzbekistan.

2873 **Habitat:** In steppes, on stony slopes, and among bushes.2874 **Population status:** Common, forming dense groups.

2875 **Traditional use:** This species is used as a replacement for *Helichrysum arenarium*. The inflorescences are collected at the
2876 beginning of flowering and are used to make a tea, which is taken to treat liver diseases, jaundice, gall and kidney stones,
2877 edema, and tuberculosis. It is also used as a hemostatic for hemorrhoidal bleeding, as a vermifuge (particularly for ascari-
2878 dosis), as a common cold remedy, and as a diuretic (Khodzhimatov 1989; Sezik et al. 2004).

2879 **Documented effects:** In modern medicine, water decoctions and infusions, liquid extracts, and dry concentrates of the
2880 inflorescences, as well as the preparation *Flamin*, are used as a choleric for treating liver disease, cholecystitis and hepa-
2881 tocholecystitis (Khodzhimatov 1989). An ethanolic extract of the flowers and the compound naringenin chalcone (iso-
2882 lated from the extract) showed antiproliferative activity against mouse skin tumor cells in vitro. Application of
2883 isosalipurposide, isolated from the flowers, delayed formation of papillomas in an in vivo assay of carcinogenesis on
2884 mouse skin (Yagura et al. 2008).

2885 **Phytochemistry:** The flowers contain flavonoids, glycosides, diterpenes, coumarins, sterins, vitamin K, essential oil, gum,
2886 dyeing substances, fatty acids, etc. (Khodzhimatov 1989; Baimukhamedov and Komissarenko 1990; Ul'chenko et al.
2887 2000; Yagura et al. 2008).

2888

Herniaria glabra L. – Caryophyllaceae	2889
Synonyms: <i>Herniaria suavis</i> Klokov.	2890
English name: Rupturewort, smooth rupturewort	2891
Russian name: Грыжник голый (Gryzhnik golyy)	2892
Uzbek name: Tuksuz saminchop	2893
Kyrgyz name: Туксуз самын чоп (Tuksuz samyn chop)	2894
Description: Yellowish-green perennial herb with woody taproot. Stems prostrate, sometimes ascending, 5–25 cm long, strongly branched from the base, glabrous or slightly hairy. Leaves mostly opposite, simple, elliptic to obovate, 2–7 mm long, 1–3 mm wide, short-petiolate, usually glabrous or sometimes short-ciliate. Inflorescences axillary clusters or capitate-spiceform, usually leaf opposed. Flowers sessile. Calyx 5-lobed, whitish-green, lanceolate to oblong, glabrous. Petals absent. Stamens 5. Styles 2, lower 1/3 connate. Fruit a utricle, 1–1.3 mm, usually longer than calyx.	2895 2896 2897 2898 2899
Other distinguishing features: Differs from closely related species by having herbaceous stems (sometimes woody at base) and mostly glabrous leaves.	2900 2901
Phenology: Flowers and fruits in June-August.	2902
Reproduction: By seeds.	2903
Distribution: Toshkent, Farg'ona, Samarqand, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.	2904
Habitat: The adyr and tau zones. In open, dry, sandy, stony places, along rivers, near roads, and on mountain slopes.	2905
Population status: Not common, found as single individuals.	2906
Traditional use: Used as diuretic to treat edema and acute catarrh of the bladder, as an astringent, to treat syphilis, pulmonary and other diseases, as well as for kidney inflammation and jaundice (Khalmatov et al. 1984).	2907 2908
Documented effects: Preparations have antispasmodic and diuretic activities, and are especially effective for urinary bladder inflammation. In medicine, an infusion is used for renal pain, inflammation of the renal pelvis, ureteritis, and to help excrete stones from kidneys and the urinary bladder (Khalmatov et al. 1984). A water extract from the aboveground parts increased diuresis in rats by 73 % (Khodzhimatov 1989). Treating hypertensive rats with saponins from <i>Herniaria glabra</i> resulted in a significant decrease in blood pressure (Rhiouani et al. 1999; Rhiouani et al. 2001).	2909 2910 2911 2912 2913
Phytochemistry: The herb contains coumarins (umbelliferone and herniarine) and their derivatives, flavonoids (quercetin, rutin, quercetin triglycoside, quercetin arabinoside, quercetin galactoside, rhamnoglycoside, isorhamnetin triglycoside, etc.), triterpene saponins, essential oil, and traces of alkaloids (Khodzhimatov 1989; Akopov 1990; Schröder et al. 1993).	2914 2915 2916 2917

2919 **Hibiscus trionum L. – Malvaceae**

2920 **Synonyms:** *Hibiscus ternatus* Cav.

2921 **English name:** Flower of an-hour

2922 **Russian name:** Гибискус тройчатый (Gibiskus troychatyy)

2923 **Uzbek name:** Burytaroq

2924 **Kyrgyz name:** Уч айчыктуу гибиск (Uch aychyktuu gibisk)

2925 **Description:** Herbaceous annual, 5–75 cm tall. Stems erect, mostly branched, lower branches elongated, stems with scattered stiff, forked and stellate hairs,. Leaves alternate, petiolate, stipulate, adaxial surface of leaf nearly glabrous, abaxial side with scattered stellate-hairs; stem leaves palmatilobate, usually with 3 oblong, pinnatilobate segments; uppermost leaves unlobed to slightly lobed. Flowers solitary in leaf axils, with long pedicels; epicalyx with 7–13 bractlets, linear, ciliate-bristly. Calyx campanulate, 5-lobed, with purple veins, bristly and stellate hairy. Corolla very deeply 5-lobed, pale-yellow with reddish-purple center, 1.5–3.5 cm wide. Staminal column 3–4 mm long. Stigmas 5, reddish-purple. Fruit a black capsule, hairy. Seeds 2.5 mm long, kidney-shaped or irregular.

2932 **Other distinguishing features:** Stipules 2–7 mm long, long-ciliate. Flowers quickly fading. Calyx becomes inflated in fruit.

2934 **Phenology:** Flowers and fruits in July-September.

2935 **Reproduction:** Only by seeds.

2936 **Distribution:** All regions of Uzbekistan and Kyrgyzstan.

2937 **Habitat:** The chul zone. As a weed in cotton and melon fields, vegetable gardens, and all irrigated farming areas.

2938 **Population status:** Common.

2939 **Traditional use:** An infusion of the leaves is used as an expectorant to treat catarrh in the upper respiratory tract. In Romania the plant is used as a diuretic (Khalmatov 1964).

2941 **Documented effects:** An infusion and extract made from different plant parts have a diuretic effect. Special diuretic properties were documented from preparations of the leaves (Khalmatov 1964). Extracts of the plant exhibit antimicrobial activity (Szabo et al. 2006).

2944 **Phytochemistry:** Gossypol has been isolated from the seeds (Schmidt and Wells 1990). The main fatty acids isolated from the seed oil were linoleic acid (63.61 %), hexadecanoic acid (16.72 %), oleic acid (12.30 %), stearic acid (2.23 %), and the total content of the unsaturated fatty acids was 79.11 % (Hu et al. 2006).

2947

Hippophae rhamnoides L. – Elaeagnaceae

2948

Synonyms: *Elaeagnus rhamnoides* (L.) A. Nelson, *Hippophae angustifolia* Lodd. ex Dippel, *Hippophae littoralis* Salisb.,
Hippophae rhamnoideum Saint-Lager, *Hippophae sibirica* Hort. ex Steud., *Osyris rhamnoides* Scop., *Rhamnoides hippo-*
phae Moench.

2949

2950

2951

English name: Sea buckthorn, seaberry

2952

Russian name: Облепиха крушиновая (Oblepikha krushinovaya)

2953

Uzbek name: Chakanda

2954

Kyrgyz name: Кадимки чычырканак (Kadimki chychyrkanak)

2955

Description: Deciduous, dioecious shrub or small tree, 1.5–11 m tall. Branches with 2–7 cm long spines. Leaves alternate,
short-petiolate, linear-lanceolate, 2–8 cm long, 2–8 mm wide, adaxial side gray-green, abaxial side brownish-silver due
to scales and stellate hairs. Flowers unisexual. Staminate flowers in short spikes; flowers 5–8 mm long, 4–6 mm wide,
outside covered with brown and white scales. Pistillate flowers covered with scales, very short-pedicellate, in groups of
2–5 in branch and thorn axils. Fruit a juicy, orange, red or yellow ellipsoidal drupe, 0.5–1 cm long, 3.8 mm wide. Seeds
dark-brown, shiny.

2956

2957

2958

2959

2960

2961

Other distinguishing features: Fruits have a peculiar flavor and aroma.

2962

Phenology: Flowers in April-May, fruits in August.

2963

Reproduction: By seeds.

2964

Distribution: All provinces of Kyrgyzstan; Toshkent, Farg'ona and Samarqand provinces of Uzbekistan.

2965

Habitat: Along stream and river banks in valleys and into the mountains.

2966

Population status: Common, found in dense groups.

2967

Traditional use: The fruits are used as an analgesic, as a remedy for stomach pain, to improve digestion, and to treat scurvy.

2968

A decoction of the fruits is drunk to treat ulcers and is added to baths to prevent skin diseases. Fresh fruits are used to
moisturize the skin, to help heal small wounds and burns, and to treat skin diseases associated with poor metabolism. An
infusion of the leaves is drunk or the leaves are directly applied to the body to treat rheumatism. A decoction of the seeds
is used as a laxative (Khalmatov et al. 1984; Khodzhimatov 1989).

2969

2970

2971

2972

Documented effects: Fruits are a rich source of polyvitamins. Oil from the fruits is used as an analgesic and to treat burns,
frostbite, eczema, persistent wounds, as well as stomach and duodenal ulcers. The oil is used during radiation treatment
for esophageal cancer (Tolmachev 1976). A study of the radioprotective action of a preparation of this species resulted in
an 82 % survival rate in mice that received the treatment compared to no survival in irradiated control (Goel et al. 2002).
Alcoholic extracts of leaves and fruits of sea buckthorn were found to inhibit chromium-induced free radical production,
apoptosis, and DNA fragmentation, and restored the anti-oxidant status to that of control cells. These extracts also were
able to arrest the chromium-induced inhibition of lymphocyte proliferation (Geetha et al. 2002). Flavonoids isolated from
the plant are reported to have antioxidant, anti-ulcerogenic, and hepato-protective properties (Yue et al. 2004).

2973

2974

2975

2976

2977

2978

2979

2980

Phytochemistry: The fruits contain carotene, vitamins C, E, B₁ and B₂, folic acid, sugars, organic acids, quercetin, isorham-
netin, tannins, and semi-drying fatty oil. The leaves contain tannins, vitamin C, and polyphenols (kaempferol, quercetin,
isorhamnetin and myristicine; Khalmatov et al. 1984; Yue et al. 2004).

2981


2982

2983

2984



▲ **Helichrysum maracandicum Popov ex Kirp.** Photos: *left:* Evgeny Davkaev; *center and right:* Alexander Naumenko



▲ **Herniaria glabra L.**

Photos: Maxim Kucherov

◀ **Hippophae rhamnoides L.**

Photos: *top:* Vadim Prokhorov;

center: Rostislav Lezhoev;

bottom: Sergey Mayorov

▶ **Hibiscus trionum L.**

Photos: *bottom:* Alim Gaziev;

center and top: Sasha Eisenman

Hyoscyamus niger L. – Solanaceae	2985
Synonyms: <i>Hyoscyamus agrestis</i> Kit. ex Schult., <i>Hyoscyamus bohemicus</i> F.W. Schmidt.	2986
English name: Black henbane	2987
Russian name: Белена чёрная (Belena chyornaya)	2988
Uzbek name: Ming divana	2989
Kyrgyz name: Кара мeндубана (Kara mendubana)	2990
Description: Herbaceous biennial with taproot. Stems green, 15–150 cm, villous. Leaves alternate, simple, dull green from above, gray-green below with long hairs; basal rosette leaves long-petiolate, elliptic, pinnatifid; cauline leaves sessile, elongate-lanceolate, with triangular lobes. Flowers solitary in axils or in scorpioid spikes. Calyx tubular-campanulate, with 5 broadly triangular lobes. Corolla funnelform with 5 lobes, greenish-yellow with purple reticulate veins. Fruit a bilocular capsule, circumscissile, 15–18 mm long. Seeds up to 500 per capsule, brownish-gray.	2991 2992 2993 2994 2995
Other distinguishing features: The entire plant is densely hairy and has an unpleasant aroma.	2996
Phenology: Flowers in May-August, fruits in July-September.	2997
Reproduction: By seeds.	2998
Distribution: Almost all provinces of Kyrgyzstan and Uzbekistan.	2999
Habitat: In waste places, near houses, in vegetable gardens, and cultivated and fallow fields.	3000
Population status: Common, forming dense groups.	3001
Traditional use: Avicenna recommended the juice of the leaves to treat eye, ear, tooth, and uterine pain and as a hemostatic for uterine bleeding. He also suggested that a paste made with the leaves and seeds be used as an analgesic for pain associated with gout. In current folk medicine this plant is still used as an analgesic. The leaf juice is used to treat tumors and earaches. A water infusion of the seeds is used to treat convulsions and smoke from the burning seeds is used to treat toothaches. A plaster of the leaves is put on swollen abscesses to draw out pus (Khalmatov et al. 1984).	3002 3003 3004 3005 3006
Documented effects: The plant is highly toxic. Preparations from this species are mostly used as antispasmodic and analgesic medicines. Atropine is used to treat bile ducts, stomach and duodenal ulcers, intestinal spasms, and bronchial asthma, and is used in ophthalmology as a mydriatic. Scopolamine is used as a depressant of the central nervous system in surgery and psychiatry. Oil from the leaves is used as analgesic to treat rheumatism and neurological pains. The leaves are used to prepare antiasthmatic medicines (<i>Asthmatin</i> ; Tolmachev 1976).	3007 3008 3009 3010 3011
Phytochemistry: The entire plant contains alkaloids including hyoscyamine (isomer of atropine), scopolamine, and glycosides. The seeds contain essential oils. The leaves are rich in flavonoids such as rutin (Tolmachev 1976; Gammerman et al. 1990).	3012 3013 3014

3016 **Hypericum perforatum L. –Hypericaceae**3017 **Synonyms:** *Hypericum komorovii* Gorschk., *Hypericum nachitschevanicum* Grossh.3018 **English name:** Common St. Johnswort, St. Johnswort, Klamath weed, goat weed3019 **Russian name:** Зверобой продырявленный (Zveroboy prodyryavlennyy)3020 **Uzbek name:** Kizil-poicha3021 **Kyrgyz name:** Козонокчолуу сары чай чоп (Kozonokcholu sary chay chop)

3022 **Description:** Herbaceous perennial plant with a much-branched taproot. Stems one to many, erect, 20–100 cm tall, the upper
3023 portions branched. Leaves simple, opposite, sessile, entire, elliptic or elongate-obovate, dotted with light-colored translu-
3024 cent and black (along margins) glands. Inflorescences cymes or corymbiform. Flowers 1.5–2.5 cm wide. Sepals 5, lanceo-
3025 late to oblong. Petals 5, yellow, twice as long as sepals with marginal black dots. Fruit a capsule, 5–9 mm long,
3026 elongate-ovoid. Seeds small, elongate, brown.

3027 **Other distinguishing features:** Stamens united at base into 3–5 fascicles. Stems ridged below leaves.3028 **Phenology:** Flowers in June–July, fruits in July–August.3029 **Reproduction:** By seeds.3030 **Distribution:** All provinces of Kyrgyzstan and Uzbekistan.3031 **Habitat:** In meadow-steppes, meadows, stony slopes on foothills, along canals, and in fallow fields.3032 **Population status:** Common, not found in very large groups.

3033 **Traditional use:** One of the most commonly used herbs in Central Asia. A decoction of the herb is used as an astringent,
3034 anti-inflammatory, antiseptic, tonic, and hemostatic, and is used to treat kidney diseases, heart diseases, diarrhea, and
3035 hemoptysis. The decoction is applied externally to treat wounds (Khalmatov et al. 1984).

3036 **Documented effects:** Preparations of this species are used externally as an astringent, disinfectant, and anti-inflammatory,
3037 and used internally to treat gastrointestinal diseases and acute and chronic colitis of non-bacterial origin. Oil from the
3038 plant is used to treat gingivitis and stomatitis. A tincture of the herb is used to rinse the mouth and is drunk to treat colitis,
3039 gallstones, and cystitis. The antibacterial preparation *Novoimanin* is used against gram-positive bacteria, including peni-
3040 cillin-resistant *Staphylococcus*. Externally, it is applied to infected wounds, carbuncles, paronychia, and furuncles.
3041 *Novoimanin* is used to treat mastitis and the cracked nipples of lactating women, in stomatology, to treat stomatitis ulcers,
3042 and in otolaryngology to treat acute rhinitis, pharyngitis, laryngitis, highmoritis, chronic tonsillitis, and chronic and acute
3043 otitis (Maznev 2004). The extracts of the plant has been shown to have antidepressant, antiviral, and antibacterial effects.
3044 The flavonoid hyperforin has been identified as one of the major constituents responsible for antidepressant activity
3045 (Barnes et al. 2001).

3046 **Phytochemistry:** The herb contains anthocyanins (hypericin, pseudohypericin, protopseudohypericin, frangula-emodin
3047 anthronol, etc.) and flavonoids (hyperoside, rutin, quercetrin, isoquercetrin, and quercetin). The herb also contains essen-
3048 tial oil with terpenes, sesquiterpenes, and complex esters of isovalerianic acids, tannins, carotene, ceryl alcohol, choline,
3049 and traces of alkaloids (Khodzimatov 1989; Nahrstedt and Butterweck 1997).

3050

- Hypericum scabrum L. – Hypericaceae** 3051
- Synonyms:** *Drosanthe scabra* (L.) Spach, *Hypericum asperum* Ledeb. 3052
- English name:** None 3053
- Russian name:** Зверобой шероховатый (Zveroboy sherokhovaty) 3054
- Uzbek name:** Dalachoi, Choichoop 3055
- Kyrgyz name:** Бодурлуу сары чай чоп (Bodurluu sary chay chop) 3056
- Description:** Herbaceous perennial. Stems numerous, 20–70 cm tall, brown or reddish, covered with small, rigid papillae. 3057
 Leaves opposite, sessile, oblong to lanceolate or elongate-linear, apex rounded or mucronate, covered with glands. 3058
 Inflorescence a dense, corymbiform cyme. Sepals 5, partially connate. Petals 5, yellow with marginal black glands. Fruit 3059
 a brown, ovoid to elongate-elliptical capsule. Seeds 1.5 mm long, brown. 3060
- Other distinguishing features:** Stems rough, stamens in 3 fascicles. 3061
- Phenology:** Flowers in May-June, fruits in July-September. 3062
- Reproduction:** By seeds. 3063
- Distribution:** Jalal-Abad, Naryn, Talas, and Chuy provinces of Kyrgyzstan; Toshkent, Farg'ona, Samarqand, Qashqadaryo, 3064
 and Surxondaryo provinces of Uzbekistan. 3065
- Habitat:** On dry, stony mountain slopes and in dry stream beds. 3066
- Population status:** Common, found in small groups. 3067
- Traditional use:** In folk medicine *Hypericum scabrum* is used in a similar manner as *H. perforatum* (Khalmatov 1964). The 3068
 aboveground parts are collected during flowering before the appearance of unripe fruits and are used to treat coughs and 3069
 liver, heart, stomach, intestinal, and bladder diseases. An infusion of the flowers is used to treat jaundice (Khodzhimatov 3070
 1989). 3071
- Documented effects:** Crude extracts of *Hypericum scabrum* showed antimicrobial activity in vitro against *Bacillus cereus*, 3072
E. coli, *Staphylococcus aureus*, *Branhamella catarrhalis*, *Clostridium perfringens* and *Candida albicans* (Sokmen et al. 3073
 1999). Experiments demonstrated that an aqueous extract of the plant, given orally to rats, showed significant antiulcero- 3074
 genic activity (Yesilada et al. 1993). Also in vitro, compounds isolated from the plant had moderate cytotoxicity against 3075
 human tumor cells and mild antibacterial activity against methicillin-resistance *Staphylococcus aureus* (MRSA) and 3076
 methicillin-sensitive *Staphylococcus aureus* (MSSA; Matsuhisa et al. 2002; Tanaka et al. 2004). 3077
- Phytochemistry:** The total flavonoids isolated from this species are nearly identical to those of *Hypericum perforatum*. 3078
 Xanthonenes, vitamin C, carotene, anthocyanins, essential oil, sugars, mucilage, resins, organic acids, and saponins and 3079
 others have also been isolated from the plant (Plant Resources of the USSR 1986; Matsuhisa et al. 2002; Tanaka et al. 3080
 2004). 3081

3083 **Hyssopus seravschanicus (Dub.) Pazij – Lamiaceae**3084 **Synonyms:** *Hyssopus tianschanicus* Boriss., *Hyssopus ferganensis* Boriss.3085 **English name:** Tian Shan hyssop3086 **Russian name:** Иссоп тянь-шанский (Issop tyan'-shanskiy)3087 **Uzbek name:** Dorivor kukut3088 **Kyrgyz name:** Тянь-Шань иссобу (Tyan'-Shan' issobu)

3089 **Description:** Subshrub. Stems 40–50 cm tall, twig-like, 4-sided, glabrous. Leaves opposite, linear, 1–3.5 cm long, 1–3 mm
3090 wide, almost glabrous, margins curled. Inflorescences 4–6-flowered verticillasters, found in narrow spikes. Calyx 5–6 mm
3091 long, blue, with sharp triangular teeth. Corolla blue-violet, about 10 mm long, 2-lipped, upper lip ovate, lower lip 3-lobed.
3092 Fruits oblong nutlets, 2 mm long, 1 mm wide, glabrous.

3093 **Other distinguishing features:** Stamens 4, two equal to length of corolla and two longer. Style exserted.3094 **Phenology:** Flowers in July-August, fruits in September.3095 **Reproduction:** By seeds.

3096 **Distribution:** Jalal-Abad and Talas provinces of Kyrgyzstan; Toshkent, Andijon, Samarqand, and Surxondaryo provinces of
3097 Uzbekistan.

3098 **Habitat:** On stony slopes, on rocky and pebbly soils, on steppes.3099 **Population status:** Common, found in small groups.

3100 **Traditional use:** An infusion is used as an expectorant, anti-inflammatory, astringent, tonic, antihelminthic, to heal wounds,
3101 and to treat bronchial asthma, gastrointestinal diseases, dyspepsia, rheumatism, anemia, stenocardia, neurosis, scrophula,
3102 meteorism and hyperhydrosis. It applied to the mouth to treat stomatitis and bad breath, and externally to heal persistent
3103 wounds. In Indian medicine it is used to treat bronchial asthma and acute respiratory infections (Zotov 1975; Dzhumaev
3104 1980).

3105 **Documented effects:** The plant has antiprotist, antibacterial, and antifungal activities, as well as lactogenic properties. The
3106 essential oil and phytoncides have antibacterial actions. In veterinary science an infusion is used to treat inflammation of
3107 the gastrointestinal tract in calves (Zotov et al. 1977).

3108 **Phytochemistry:** The plant contains steroids (β -sitosterin), flavonoids (diosmine), essential oil (containing camphene,
3109 β -pinene, pinocamphone, 1,8-cineol, linalool, α -terpenyl-acetate, bornyl acetate, myrcene, limonene, etc.), triterpenoids
3110 (ursolic and oleanolic acids), vitamins B₁, B₂ and C, and phenolcarbonic acids and their derivatives. The seeds contain
3111 fatty oil including palmitic, stearic, oleinic, linoleic, and linolenic acids (Zotov 1975).



▲ *Hyssopus seravschanicus* (Dub.) ▲ *Hyoscyamus niger* L. Photos: Alim Gaziev
Pazij Photo: Dr. Petr Kocna, www1.lf1.cuni.cz/~kocnaflowr_myflow_gb.htm.jpg

▼ *Hypericum perforatum* L. Photos: *left*: Mary Backlund; *right and center*: Sergey Appolonov



▼ *Hypericum scabrum* L. Photos: *left*: John B. Taft; *right*: Vladimir Epiktetov



3112 ***Impatiens parviflora* DC. – Balsaminaceae**3113 **Synonyms:** *Impatiens brachycentra* Kar. & Kir.3114 **English name:** Small balsam, small flower touch-me-not3115 **Russian name:** Недотрога мелкоцветная (*Nedotroga melkotsvetnaya*)3116 **Uzbek name:** Hinagina, Chupkhina3117 **Kyrgyz name:** Майда гулду кына (*Mayda guldu kyna*)

3118 **Description:** Herbaceous annual with fibrous roots. Stems erect, 30–70 cm tall, succulent, glabrous. Leaves alternate,
3119 8–17 cm long, 4–8 cm wide, elliptic or ovate, apex acuminate, margins serrate-dentate, gradually tapering to 1–2 cm long
3120 petiole. Inflorescences loose axillary racemes, with 4–12 flowers; peduncles similar in length to the leaves; pedicels thin,
3121 1.5–2 cm long. Flowers irregular, up to 1 cm long. Sepals 3, 2 lateral sepals small, ovate; lower sepal petaloid with
3122 4–5 mm long spur. Petals 5, lateral petals connate in pairs, 3-lobed, yellow with red spots in the throat; fifth petal sub-
3123 orbicular. Fruit an oblong capsule, 2 cm long, 3–4 mm wide, explosively dehiscent along raised longitudinal seams. Seeds
3124 oval, almost round.

3125 **Other distinguishing features:** Leaf teeth glandular. Flowers directed upward or aside, not drooping. Ripe fruits burst when
3126 touched.

3127 **Phenology:** Flowers in June–July, fruits in July–August.3128 **Reproduction:** By seeds.

3129 **Distribution:** Tashkent, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Chuy and Jalal-Abad prov-
3130 inces of Kyrgyzstan.

3131 **Habitat:** The tau zone. Grows in wet, shady places, walnut forests, in oases, and can be found in shaded areas of orchards,
3132 as a weed.

3133 **Population status:** Common.3134 **Traditional use:** Used in folk medicine as a hemostatic and as a treatment for various uterine diseases (Khalmatov 1964).

3135 **Documented effects:** An alcohol extract of this species had highly significant hemostatic activity. Extracts of the herb con-
3136 tain antibacterial substances (Khalmatov 1964). An infusion of the herb in alcohol had sedative and hypotensive effects,
3137 regulated the menstrual cycle, and accelerated childbirth delivery (Ibragimov and Ibragimova 1960). An experiment with
3138 an aqueous extract of the plant, to determine cyclooxygenase inhibition, showed negative inhibition, indicating an enzyme-
3139 stimulating effect (Tunon et al. 1995).

3140 **Phytochemistry:** Aboveground parts contained flavonoid glycosides (0.43 %), alkaloids (0.016 %), resins (3.53 %), vitamin
3141 C (7.2 mg%), and traces of carotene. The compounds N-oxy-benzoic acid, vanillic, gentisinic, ferulic acid, N-coumarinic
3142 and caffeic acids, as well as 2-methoxy-1,4 naphthoquinone have been isolated from the leaves (Khalmatov 1964). Oil
3143 from the seeds contains parinaric acid (Tsevegsuren et al. 1998).

3144

- Inula britannica* L. – Asteraceae** 3145
- Synonyms:** *Conyza britannica* (L.) Moris ex Rupr., *Inula serrata* Gilib., *Inula tymiensis* Kudô. 3146
- English name:** British yellowhead, British elecampane, meadow fleabane, yellow starwort 3147
- Russian name:** Девясил британский (Devyasil britanskiy) 3148
- Uzbek name:** Chachalbosh 3149
- Kyrgyz name:** Сары баш карындыз (Sary bash karyndyz) 3150
- Description:** Herbaceous perennial, with thin creeping rhizomes. Stems mostly erect, often villous or with orange glands, 10–70 cm tall. Basal leaves elliptic, lanceolate or ovate, 3–13 cm long, 1–3.2 cm wide; cauline leaves alternate, sessile, elongate-lanceolate to lanceolate. Inflorescence a capitulum, 3–5 cm wide, single or in corymbiform groups; involucrel bracts linear, 4–6 mm long, in 2 rows. Ray flowers many (ca. 40–70), 1–1.5 cm long, yellow, twice as long as bracts; disc flowers 4–6 mm long, yellow. Fruits linear-oblong achenes, ribbed, brown, with gray-white pappus. 3151–3155
- Other distinguishing features:** Pappus consists of 15–25 simple hairs. 3156
- Phenology:** Flowers and fruits in May-September. 3157
- Reproduction:** By seeds. 3158
- Distribution:** All provinces of Kyrgyzstan; Toshkent, Farg'ona, Andijon, Jizzax and Surxondaryo provinces of Uzbekistan. 3159–3160
- Habitat:** In semi-desert areas, steppes, and meadows, along the edges of rivers and lakes, and among bushes. 3161
- Population status:** Common, found in small groups. 3162
- Traditional use:** This species is gathered in autumn or early spring. An infusion or decoction of the underground parts is used to treat cystitis, diabetes, jaundice, respiratory catarrh, bone tuberculosis, rheumatism, and hemorrhoids, and is used as a vermifuge, hemostatic for uterine bleeding, and to improve the appetite. An infusion of the leaves is drunk as an anti-inflammatory and astringent remedy (Khodzhimatov 1989). 3163–3166
- Documented effects:** In modern medicine *Inula britannica* is used the same way as *Inula helenium* (Khodzhimatov 1989). Flavonoids isolated from this species were shown to protect cultured rat cortical cells from cell death caused by oxidative stress (Kim et al. 2002). Results from experiments with mice suggest an aqueous extract from the flowers of *Inula britannica* ssp. *japonica* Kitam. has a preventative effect on autoimmune diabetes by regulating cytokine production (Kobayashi et al. 2002b). The sesquiterpene lactone ergolide has anti-inflammatory activity (Han et al. 2001). 3167–3171
- Phytochemistry:** The aboveground parts contain flavonoids, essential oil, tannins, ergolide, britanin, and other sesquiterpene lactones. The underground parts contain essential oil (with alantolactone and isoalantolactone), alkaloids, and inulin. The leaves contain vitamin C (Khalmatov et al. 1984; Zhou et al. 1993; Han et al. 2001; Kim et al. 2002). 3172–3174

3176 **Inula grandis Schrenk ex Fisch. & C.A. Mey. – Asteraceae**

3177 **Synonyms:** *Codonocephalum grande* (Schrenk ex Fisch. & C.A. Mey) B. Fedtsch., *Inula macrophylla* Kar. & Kir.

3178 **English name:** Large-leaved elecampane

3179 **Russian name:** Девясил крупнолистный (Devyasil krupnolistnyy)

3180 **Uzbek name:** Sari andiz, Ok andiz

3181 **Kyrgyz name:** Unknown

3182 **Description:** Herbaceous perennial, 50–200 cm high with a thick, vigorous, branched root. Stem erect, branching towards
3183 the top. Leaves coriaceous, shiny, adaxial side glabrous, abaxial side glandular, margins serrate-dentate. Basal leaves
3184 widely elliptic, up to 25–85 cm long and 18–32 cm wide, petioles 10–20 cm; stem leaves elongate-elliptical, 20–37 cm
3185 long, 8–15 cm wide, sessile; upper leaves lanceolate, 3–10 cm long, 1–4.5 cm wide. Inflorescences capitula, 2–5 arranged
3186 in a corymbiform raceme; capitula 4.5–6 cm in diameter with ray and disk flowers. Ray flowers yellow, 1–3 cm long. Fruit
3187 a cylindrical achene, brown, with multiple longitudinal ribs and yellowish pappus.

3188 **Other distinguishing features:** Leaves stiff, odorous, with vaguely sinuate edges and distinctly reticulate veins. Involucral
3189 bracts lanceolate, acute, coriaceous.

3190 **Phenology:** Flowers in May-July, fruits in July-August.

3191 **Reproduction:** By seeds.

3192 **Distribution:** Toshkent, Samarqand, Andijon, Farg'ona, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.

3193 **Habitat:** The adyr and tau zones. Shallow soils and gentle slopes.

3194 **Population status:** Common, usually found in small populations.

3195 **Traditional use:** A decoction made of underground organs of this plant, and related species, are used to treat brucellosis,
3196 tuberculosis, gastrointestinal diseases, and as a vermifuge. Young juicy stems, with the bark removed, are used as a restor-
3197 ative and to treat phthisis (Khalmatov and Kosimov 1992).

3198 **Documented effects:** Preparations are used to treat ulcers and gastric catarrh, as well as duodenal ulcers (Khalmatov 1964).
3199 Compounds isolated from this species showed unique anti-oxidant activity (Kogure et al. 2004).

3200 **Phytochemistry:** The roots contain essential oils (up to 3 %), the main portion of which is sesquiterpene lactones: alantolac-
3201 tones and isoalantolactones, proazulen and alantone. The roots also contain saponins, inulin (up to 44 %), resins, traces
3202 of alkaloids, and the sesquiterpene lactones carabron and granilin (Kulikov 1975; Akopov 1981; Khalmatov and Kosimov
3203 1992). The bark contains many mono- and sesquiterpene lactones (Fu et al. 2001; Su et al. 2000, 2001).

3204

- Inula helenium L. – Asteraceae** 3205
- Synonyms:** *Aster helenium* (L.) Scop., *Corvisartia helenium* (L.) Mérat, *Helenium grandiflorum* Gilib. 3206
- English name:** Elecampane 3207
- Russian name:** Девясил высокий (Devyasil vysokiy) 3208
- Uzbek name:** Kora andiz 3209
- Kyrgyz name:** Бийик карындыз (Biyik karyndyz) 3210
- Description:** Herbaceous perennial, with thick, fragrant rhizomes. Stem erect, 0.5–2.5 m tall, white-hairy. Lower leaves 3211
alternate, long-petiolate, elliptic, irregularly shallow-dentate, up to 50 cm long, up to 10–20 cm wide, densely hairy on 3212
abaxial side; stem leaves elongate-ovate, becoming sessile towards the top. Inflorescence a capitulum, 3–8 cm wide, in 3213
groups forming loose racemes or corymbs. Ray flowers golden-yellow, numerous (ca. 50–100), ca. 3–4 cm long, thin; disc 3214
flowers 9–11 mm long. Fruits brown achenes, quadrangular, with light colored pappus. 3215
- Other distinguishing features:** Basal leaves very broad, up to 20 cm wide. 3216
- Phenology:** Flowers in June-August, fruits August-September. 3217
- Reproduction:** By seeds. 3218
- Distribution:** In the Chuy valley (North slopes of Kyrgyz Alatau and Talas Alatau) and Ferghana Range (Arslanbob and 3219
Kara-Alma areas) in Kyrgyzstan; Tashkent, Samarqand, Jizzax, Andijon and Farg'ona provinces of Uzbekistan. 3220
- Habitat:** Along rivers and streams, in the lower and mid belt of mountains. 3221
- Population status:** Common, found in small groups. 3222
- Traditional use:** Avicenna stated that this plant belongs to a group with the ability to act as a tonic and invigorate and 3223
strengthen the heart. It is useful to treat inflammation of the sciatic nerve and joint pain. Mixed with honey it is used as an 3224
expectorant. A decoction of the rhizomes, especially a syrup made from it, works as a diuretic and promotes menstruation. 3225
The rhizomes are still used to treat gastrointestinal diseases, malaria, cystitis, bone tuberculosis, rheumatism, radiculitis, 3226
diabetes, jaundice, edema, and respiratory catarrh. An ointment or water infusion is applied to treat eczema and scabies. 3227
A tincture of roots (in vodka) is drunk to treat gastritis, stomach and duodenal ulcers, tuberculosis, nervous diseases, 3228
goiters, heart diseases, and hypertension and is used as an expectorant for treatment of chronic respiratory diseases (tra- 3229
cheitis, lung tuberculosis, and bronchitis). It is also used to treat gastroenteritis and diarrhea of non-infectious origins 3230
(Khodzhimatov 1989). 3231
- Documented effects:** In modern medicine, a decoction of the underground parts is recommended to treat respiratory and 3232
gastrointestinal diseases. The preparation *Alanton* is used to treat ulcers (Poludenny and Zhuravlev 2000). Experimental 3233
results show that a decoction of the underground parts act as an expectorant, weak diuretic, choleric, and weak hemo- 3234
static, and normalizes the function of the gastrointestinal tract. The essential oil acts as a very strong vermifuge, especially 3235
against *Ascaris* worms and pork and beef tapeworms. A preparation from this plant is used externally to treat skin diseases 3236
such as eczema, scabies, and neurodermatitis (Akopov 1990). Extracts of the roots, and isolated sesquiterpene lactones, 3237
showed significant inhibitory activity against a variety of cancer cell lines in vitro as well as against *Mycobacterium* 3238
tuberculosis (Cantrell et al. 1999; Konishi et al. 2002). 3239
- Phytochemistry:** The underground parts contain 1–3 % essential oil (including sesquiterpene lactones such as alantolactone, 3240
isoalantolactone, dihydroalantolactone, etc.), up to 44 % inulin and other sugars, pigments, gums, mucilage, alkaloids, 3241
and acetic and benzoic acid. The aboveground parts contain alkaloids, essential oil, alantopicrine, and folic acid (Khalmatov 3242
et al. 1984; Khodzhimatov 1989; Akopov 1990; Cantrell et al. 1999; Konishi et al. 2002). 3243



▲ *Impatiens parviflora* DC. Photos: Sergey Appolonov



▲ *Inula grandis* Schrenk ex Fisch. & C.A. Mey.

Photos: Alexander Naumenko



▲ *Inula britannica* L. Photos: Sergey Appolonov

▼ *Inula helenium* L. Photos: Sergey Mayorov



- Juglans regia L. – Juglandaceae** 3245
- Synonyms:** *Juglans duclouxiana* Dode, *Juglans fallax* Dode, *Juglans kamaonia* (C. DC.) Dode, *Juglans orientis* Dode, *Juglans sinensis* (C. DC.) Dode. 3246
- English name:** Persian walnut, English walnut 3248
- Russian name:** Грецкий орех (Gretskiy orekh) 3249
- Uzbek name:** Yöng'oq 3250
- Kyrgyz name:** Грек жангагы (Grek zhangagy) 3251
- Description:** Large monoecious tree with a wide, dense crown, 15–35 m tall. Trunk diameter up to 1.5(–2.5) m wide; young trees with slightly cracked, light-gray bark; older trees have darker, strongly cracked bark. Leaves alternate, 19–54 cm long, 15–40 cm wide, dark-green, odd-pinnate with 3–5 pairs of leaflets; leaflets ovate, coriaceous, glabrous with entire margins. Male flowers arranged in catkins, each flower with 8–40 stamens. Female flowers in groups of 1–3 on ends of young branches. Fruit drupe-like, spherical, pericarp green, drying when ripe; endocarp or “shell” light brown, hard. Seed covered with thin yellow papery layer. 3252–3257
- Other distinguishing features:** The pith of young branches is chambered. Leaves produce a specific, pungent smell when crushed. 3258–3259
- Phenology:** Flowers in April-May, fruits in September. 3260
- Reproduction:** By seeds. 3261
- Distribution:** All regions of Uzbekistan; Osh and Jalal-Abad provinces of Kyrgyzstan. 3262
- Habitat:** The tau zone. Mountain river banks, rarely on slopes, as single trees or groups. 3263
- Population status:** As individual trees and groups, creates walnut forests. 3264
- Traditional use:** Young, green fruits are used to prepare a concentrate of vitamins. A decoction of the nuts is drunk to treat high arterial pressure, cardiac diseases, and to rinse the mouth to treat gum disease. Juice from the fruit husk is used as an ointment to treat different kinds of external ulcers, eczema, and other cases of itchy dermatosis. A tea of the leaves is drunk to treat diabetes and decrease sugar content in the urine. The leaves are used as a vermifuge and to treat skin diseases, venereal diseases, catarrh of the gastrointestinal tract, and tuberculosis. A decoction of the leaves is drunk to treat scrofula and rickets. The bark from the roots is used to make a very mild laxative (Akopov 1981). 3265–3270
- Documented effects:** The leaves and the fruit husks are used to make the preparation *Juglon*. It is used externally to treat skin tuberculosis and *Staphylococcus* and *Streptococcus* lesions. It has antimicrobial and anti-inflammatory actions and is used to heal wounds. Unsaturated fatty acids from the nuts help to prevent arteriosclerosis (Nuraliev 1989). Clinical studies showed that a water extract of the leaves was effective in treating some forms of skin tuberculosis, tuberculosis lymphadenitis, and tuberculosis of the larynx (Altimishev 1991). In experiments with mice, extracts of *Juglans regia* improved glucose tolerance in hypoglycemic activity screens (Neef et al. 1995). In vitro, an extract of the nuts inhibited the oxidation of human plasma and low density lipoproteins (Anderson et al. 2001). Polyphenols, isolated from the nuts, exhibited antioxidative properties (Fukudu et al. 2003). 3271–3278
- Phytochemistry:** The leaves contain α -hydrojuglone, which easily oxidizes to juglone, β -hydrojuglone, flavonoids (hyperoside, quercetin-3-arabinoside), ascorbic acid, vitamin P, B₁, tannins, carotene, pigments and essential oil. The fruit husk contains ascorbic acid, tannins, and α - and β -hydrojuglone. The seeds contain carotene, vitamins C, B₁, E, and P, and fatty oil, which contains glycerides of linoleic and oleic acids. β -sitosterol and its glycoside have been isolated from the papery layer surrounding the seed. The bark contains tannins, pigments, and gallic and ellagic acids (Khodzhimatov 1989; Colaric et al. 2005). 3279–3284

3286 **Juniperus sabina L. – Cupressaceae**3287 **Synonyms:** *Sabina vulgaris* Antoine3288 **English name:** Savin juniper, Savin3289 **Russian name:** Можжевельник казацкий (Mozhzhewel'nik kazatskiy)3290 **Uzbek name:** Archa3291 **Kyrgyz name:** Кара арча (Kara archa)

3292 **Description:** Dioecious, evergreen, more or less prostrate shrub, or occasionally an erect small tree, to 5 m high. Bark red-
3293 dish-gray. Branchlets slender, densely arranged. Leaves of 2 kinds, needle-like and scale-like; needle-like leaves present
3294 on young plants and sterile branches only, 3–7 mm long, appressed; scale-like leaves 1–3 mm long with a oval shaped
3295 gland on the back. Pollen (male) cones ellipsoid or oblong, 3–4 mm long. Seed (female) cones berry-like, brown-black,
3296 pruinose, round-oval, 2–5-seeded. Seeds brown, 4–5 mm long.

3297 **Other distinguishing features:** Leaves have a characteristic smell when ground. Prostrate branches sometimes form roots.3298 **Phenology:** Flowers in April, fruits ripen in the fall or the spring of the following year.3299 **Reproduction:** By seeds.3300 **Distribution:** The Chuy and Naryn provinces of Kyrgyzstan; not found in the flora of Uzbekistan.3301 **Habitat:** In the steppe and forest belts and on stony slopes of hills and low mountains.3302 **Population status:** Common, forming dense groups.

3303 **Traditional use:** An infusion of the berries is used to treat urogenital diseases. A decoction of the berries is drunk to treat
3304 kidneys and bladder illnesses, kidney stones, liver diseases, rheumatism, scurvy, jaundice, and to improve choleric
3305 action. The berries are eaten to treat stomach ulcers and to improve appetite. A decoction of the roots is used for stomach
3306 ulcers, bronchitis, tuberculosis, kidney stones, and skin illnesses. A tincture of the bark and roots is drunk to treat arthritis.
3307 A decoction of the bark is used to raise the libido. The cones and green branches are also used in baths to treat rheumatism
3308 (Makhlayuk 1992).

3309 **Documented effects:** Extracts from the fruits and branches have cytotoxic effects on cancer cell lines in vitro (Jafarian-
3310 Dehkordi et al. 2004). Results from experiments with mice indicate that the abortifacient effect of essential oil from
3311 *Juniperus sabina* is related to an implantation inhibiting effect induced by sabinyl acetate (Pages et al. 1996). Cyclolignans,
3312 isolated from the leaves, exhibits anti-cancer and anti-viral activity (San Feliciano et al. 1993).

3313 **Phytochemistry:** The plant contains podophyllotoxin and other cyclolignans. Fresh branches, leaves, bark and fruits contain
3314 essential oil which contains pinene, cadinene, terpinene, terpinolene, camphene, cedrol, etc. The bark and stems contain
3315 tannins and the leaves contain vitamin C. The leaves, bark, and unripe cones contain pigments. The fruits contain sugar,
3316 juniperin, resins, pentosan, and organic acids (Makhlayuk 1992; San Feliciano et al. 1993).

3317

Juniperus semiglobosa Regel – Cupressaceae	3318
Synonyms: <i>Juniperus jarkendensis</i> Kom., <i>Juniperus sabina</i> var. <i>jarkendensis</i> (Kom.) Silba, <i>Juniperus schunganica</i> Kom., <i>Juniperus tianshanica</i> Sumnev., <i>Sabina vulgaris</i> var. <i>jarkendensis</i> (Kom.) Cheng-yuan Yang.	3319 3320
English name: Russian Juniper	3321
Russian name: Можжевельник полушаровидный (Mozhzhevel'nik polusharovidnyy)	3322
Uzbek name: Saur archa	3323
Kyrgyz name: Саур-арча (Saur-archa)	3324
Description: Dioecious or occasionally monoecious evergreen tree or shrub, up to 10 m tall. Branchlets thick, straight, loosely arranged. Leaves of 2 kinds, needle-like and scale-like; needle-like leaves usually present on young plants, rarely on adult plants, 3–7 mm long; scale-like leaves closely appressed, rhomboid-ovate, 0.9–2.5 mm long. Pollen (male) cones ellipsoid, 3–5 mm long. Seed (female) cones berry-like, 4–8 mm long, 5–10 mm wide, globose to semispherical, green- brown when unripe, black when ripe, pruinose, 2–4-seeded. Seeds up to 6 mm long, up to 3.5 mm wide, brown.	3325 3326 3327 3328 3329
Other distinguishing features: Branchlets thick.	3330
Phenology: Flowers in March-May, fruits ripen the following year.	3331
Reproduction: By seeds.	3332
Distribution: Chuy, Naryn, and Jalal-Abad provinces of Kyrgyzstan; Toshkent, Farg'ona and Samarqand provinces of Uzbekistan.	3333 3334
Habitat: In the vegetation of the tree-shrub belt, at elevations of 1,500–2,700 m on dry and stony slopes.	3335
Population status: Common, found growing as single plants.	3336
Traditional use: An infusion or decoction of the fruits is used to induce appetite, aid in better digestion, to increase urina- tion, as a disinfectant of the urinary system, an anti-inflammatory to treat pneumonia, and as an analgesic and expectorant (Altimishev 1991). The green branches are burned in homes to provide a pleasant odor (Khodzhimatov 1989). The essen- tial oil from the needles and fruits is used to treat skin conditions (Mamedov et al. 2004).	3337 3338 3339 3340
Documented effects: In contemporary medicine a tincture of the fruits is applied externally to treat rheumatism and gout. A decoction of the fruits is use to treat hypoacidic gastritis, cholecystitis, as a disinfectant of the bladder and to increase urination. The ground fruits are spelled to treat strong headaches (Altimishev 1991).	3341 3342 3343
Phytochemistry: The wood contains 0.02 % essential oil, whereas unripe fruits and green branches contains 0.64–1.6 %. The essential oil contains up to 53 % sabinene, up to 21 % cedrol, and some sesquiterpenes and aldehydes. The fruits contain sugar and pigments. An extract from fresh branches collected in the Gissar mountains contained 0.38–0.54 % essential oil including pinene (up to 76 %), myrcene (5.4 %), cedrol (7 %), and few aldehydes (Khodzhimatov 1989).	3344 3345 3346 3347

3348 **Juniperus seravschanica Kom. – Cupressaceae**

3349 **Synonyms:** *Juniperus excelsa* var. *macrocarpa* Regel, *Juniperus kulsaica* Dmitr., *Juniperus polycarpus* var. *seravschanica*
 3350 (Kom.) Kitam., *Juniperus polysperma* Dmitr., *Juniperus pseudosabina* var. *typica* Regel, *Juniperus sabina* var. *globosa*
 3351 Regel, *Juniperus sabina* var. *macrocarpa* Regel, *Juniperus taurica* Lipsky, *Juniperus zaaminica* Dmitr., *Sabina seravschan-*
 3352 *ica* (Kom.) Nevski.

3353 **English name:** Unknown

3354 **Russian name:** Можжевельник зеравшанский (Mozhzhevel'nik zeravshanskiy)

3355 **Uzbek name:** Qora archa

3356 **Kyrgyz name:** Кызыл арча, Кара арча (Kyzyl archa, Kara archa)

3357 **Description:** Dioecious, evergreen tree up to 5–25 m tall, or sometimes a stocky bush with a dense oval or conical crown.
 3358 Bark reddish or brick-brown in color. Branches spreading, relatively short with many smaller branches. Leaves scale-like,
 3359 elongate-oval, apex acute, with long vein and gland on the lower surface. Female cone berry- or drupe-like, 9–12 mm
 3360 long, globular; young cones green, mature cones deep-brown, heavily covered with a gray coating, contains 2–3(–4)
 3361 seeds. Seeds 5–7.5 mm long, vaguely triquetrous-oval, curved, with longitudinal furrows on the sides.

3362 **Other distinguishing features:** Seedlings with needle-like leaves in whorls of 3. Seeds white when immature, brown when
 3363 ripe.

3364 **Phenology:** Pollen released in March–April. Cones ripen in September–October of the following year.

3365 **Reproduction:** By seeds.

3366 **Distribution:** Toshkent, Farg'ona, Samarqand, and Buxoro provinces of Uzbekistan; Naryn, Osh, Jalal-Abad and Batken
 3367 provinces of Kyrgyzstan.

3368 **Habitat:** The upper adyr and tau zones. Stony, shallow-soiled mountain slopes with rocky debris.

3369 **Population status:** Common, found as individual trees and also in small groves.

3370 **Traditional use:** Smoke from burning branches is used in Central Asian folk medicine to treat rheumatism. The fruits, mixed
 3371 with sesame oil, are applied to treat deafness. The powdered plant is sniffed to treat headaches. The essential oil is used
 3372 to treat wounds and skin diseases (Khalmatov et al. 1984; Mamedov et al. 2004). An infusion of the dry fruits is used to
 3373 treat the urogenital path, and a decoction is drunk to treat scurvy, liver disease and rheumatism. The fruits are also used
 3374 to treat edema and nervous disorders. Fresh fruits are eaten to treat stomach ulcers and to increase the appetite, and a
 3375 decoction is drunk as a choleric and to treat jaundice. A decoction of the roots is recommended to treat stomach ulcers,
 3376 bronchitis, lung tuberculosis, kidney stones, and skin diseases. An infusion of the roots and bark is used to treat arthritis.
 3377 A decoction of the bark is drunk to treat impotency. A decoction of the fruits and green branches is used in a bath to treat
 3378 rheumatism (Khodzhimatov 1989).

3379 **Documented effects:** A 5 % solution of cedrol (isolated from the essential oil) in castor oil is used as a treatment for fester-
 3380 ing and slowly healing wounds and chilblain ulcers, and is applied as a salve on bandages or dressings. For osteomyelitis,
 3381 this solution is poured into bone cavities (Gammerman 1960; Khalmatov et al. 1984). Essential oil from the leaves is used
 3382 to treat trichomoniasis (Khodzhimatov 1989). Some terpenoids isolated from the fruits showed moderate antimalarial
 3383 activity (Okasaka et al. 2006).

3384 **Phytochemistry:** Fresh branches contain 0.45–0.75 % essential oils with d-pinene, d-camphene, myrcene, cedrene, and
 3385 other sesquiterpenes. The bark, young branches, and unripe fruits contain 7–8 % tannins. Ripe fruits contain yellow pig-
 3386 ments and up to 18.6 % sugar. Leaf samples from Tajikistan contained 120–140 mg% vitamin C (Khalmatov et al. 1984).
 3387 A variety of compounds, including diterpenes and sesquiterpenes, has been isolated from the dried fruits (Okasaka et al.
 3388 2006).

3389

Juniperus turkestanica Kom. – Cupressaceae	3390
Synonyms: <i>Juniperus intermedia</i> V.P. Drobow, <i>Juniperus pseudosabina</i> Fisch. & C.A. Mey., <i>Juniperus pseudosabina</i> var. <i>turkestanica</i> (Kom.) Silba, <i>Juniperus pseudosabina</i> var. <i>typica</i> Regel.	3391 3392
English name: Unknown	3393
Russian name: Можжевельник туркестанский (Mozhzhevel'nik turkestanskiy)	3394
Uzbek name: Urik archa, Balik archa	3395
Kyrgyz name: Орук арча, Жапалак арча (Oruk archa, Zhapalak archa)	3396
Description: Dioecious evergreen tree to 18(–25) m tall or a stocky shrub up to 2 m high, with a dense crown. Bark brownish-gray. Branches spreading, ascending or horizontal. Leaves scale-like, about 2 mm long, ovate or rhombic, bright green, slightly pointed with a prominent gland on the back or with a prominent keel. Fruit a berry-like or drupe-like cone, 10–15 mm long and 8–10 mm wide, juicy, globular, sometimes oblong, with a single seed. Young cones green, mature cones black, shiny with a light gray coating. Seeds 6–10 mm long, oblong, pointed on the base, striated on the edges, with dark stripe on upper half; seed-coats thick, woody.	3397 3398 3399 3400 3401 3402
Other distinguishing features: Seedlings with needle-like leaves in whorls of 3. Cones taste sweet.	3403
Phenology: Pollen released in April-June, fruits the next year in September-November.	3404
Reproduction: By seeds.	3405
Distribution: Toshkent, Samarqand, and Buxoro provinces of Uzbekistan; Naryn, Osh, Jalal-Abad and Talas provinces of Kyrgyzstan.	3406 3407
Habitat: The tau and yailau zones. Stony and shallow-soiled slopes with rocky debris.	3408
Population status: Common, found as single individuals and in groups.	3409
Traditional use: Fruit decoction is recommended by folk medicine as mouth-wash to treat gingivitis (Khalmatov et al. 1984). A decoction and ointment made with the fruits are used to treat eczema, tuberculosis, skin diseases and as a diuretic (Khodzhimatov 1989).	3410 3411 3412
Documented effects: The fruits are used as a diuretic, for swelling due to kidney ailments, to treat kidney stones and are combined with other preparations to treat chronic respiratory disease and as an expectorant. The cedrol fraction, from the essential oil of young branches, together with castor oil is used as a remedy for persistent wounds and ulcers (Minayeva 1991).	3413 3414 3415 3416
Phytochemistry: The fruits contain essential oil which has up to 100 components including pinene, camphene, borneol, caphor, and other terpenes, the fruits also contain 40 % sugars, resins, flavonoids, pectic substances, etc. (Minayeva 1991).	3417 3418



▲ **Juglans regia L.** Photos: *left:* Thomas Molnar; *right:* Alim Gaziev
center: Dimitri Oreshkin

▼ **Juniperus sabina L.** Photos: Vladimir Epiktetov



▼ **Juniperus seravschanica Kom.** Photos: Evgeny Davkaev



Korolkowia sewerzowii (Regel) Regel – Liliaceae	3419
Synonyms: <i>Fritillaria sewerzowii</i> Regel.	3420
English name: Unknown	3421
Russian name: Корольковия Северцова (Korol'koviya Severtsova)	3422
Uzbek name: Olgi	3423
Kyrgyz name: Северцов алгысы (Severtsov algysy)	3424
Description: Herbaceous perennial with a spherical bulb, 3–7 cm wide. Stem thick, glabrous, erect, 30–60 cm high. Leaves simple; lower leaves opposite, wide-lanceolate to ovate, up to 20 cm long; upper leaves alternate. Inflorescence a loose terminal raceme. Flowers funnelform-campanulate, with 6 lobes, nodding; lobes greenish-brown or reddish-brown. Stamens 6, slightly shorter than the perianth. Fruit a capsule, 3–5 cm high and wide. Seeds flat, light-brown.	3425 3426 3427 3428
Other distinguishing features: Capsules erect, winged.	3429
Phenology: Flowers in April-July, fruits in May-August, depending on altitude of location.	3430
Reproduction: By seeds.	3431
Distribution: Toshkent, Farg'ona, and Samarqand provinces of Uzbekistan; Osh, Chuy and Jalal-Abad provinces of Kyrgyzstan.	3432 3433
Habitat: The adyr and tau zones. Clay-soiled slopes.	3434
Population status: Common, found as single individuals.	3435
Traditional use: In folk medicine the bulb is used as a strong diaphoretic. The bulbs contain high amounts of starch and are used as food (Khalmatov 1964).	3436 3437
Documented effects: The alkaloid alginine has local anesthetic effects, and is 4 times less toxic than cocaine. The hydrochloric salt of alginine and the total alkaloids of the plant in the form of hydrochloric salt are recommended as a local anesthetic to be used in medical practice (Khalmatov 1964). Alginine, like novocaine, acts as a conduction anesthesia and a 3–4 % solution causes widening of the pupils (Sadritdinov and Kurmukov 1980). In experiments with narcotized animals, the alkaloid alginidine (3–20 mg/kg) decreased arterial pressure and ganglioblocking effects were observed (Ishmukhamedov and Sultanov 1965).	3438 3439 3440 3441 3442 3443
Phytochemistry: Bulbs collected in the Chatkal Valley (Uzbekistan), when aboveground parts had nearly senesced, contained 0.8–0.92 % total alkaloids. Plants collected near Toshkent, during the flowering stage contained 2.3 % alkaloids in the aboveground parts and 1.4 % in the bulbs. More than 20 alkaloids have been isolated from the total alkaloids, including alginine, korseverinine, alginidine, korseveramine and korseveridine, etc. (Yunusov 1981; Samikov et al. 1989; Harrison 1990; Abdullaeva and Shakirov 2006).	3444 3445 3446 3447 3448

3450 **Lachnophyllum gossypinum Bunge – Asteraceae**3451 **Synonyms:** Unknown3452 **English name:** Unknown3453 **Russian name:** Шерстистолистник хлопковидный (Sherstistolistik khlopkovidnyy)3454 **Uzbek name:** Момуқ, Оқ момуқ3455 **Kyrgyz name:** Пахтадай лахнофиллум (Pakhtaday lakhnofillum)

3456 **Description:** Annual, 10–50 cm high, densely covered with soft, grayish, felted hairs, with abundant glands. Stems erect,
3457 often heavily branched. Lowest leaves obovate, apex obtuse or rounded, 1.5–4.5 cm long, 0.5–1.7 cm wide, leaf narrow-
3458 ing to petiole; middle leaves sessile, slightly amplexicaul with auricles on the base; upper leaves acute, narrow. Flowers
3459 in thick pubescent capitulum, heterogamous; marginal flowers ligulate, female, lilac-bluish; disk flowers bisexual, yellow.
3460 Fruits oblanceolate achenes, 2–3 mm long, flat, villous.

3461 **Other distinguishing features:** The whole plant smells nice, like ripe melon.3462 **Phenology:** Flowers and fruits in July-October.3463 **Reproduction:** By seeds.3464 **Distribution:** All regions of Uzbekistan; Osh, Chuy and Jalal-Abad provinces of Kyrgyzstan.3465 **Habitat:** The chul (bordering adyr), adyr and tau zones. Stony slopes with rocky debris.3466 **Population status:** Uncommon, found as single individuals.

3467 **Traditional use:** Fresh leaf juice is used to heal wounds. The juice is heated and brought to a thicker consistency and this is
3468 applied over the surface of old, slow-healing wounds and furuncles (Khalmatov 1964).

3469 **Documented effects:** Small doses of the crystalline material from the essential oil has strong effects on the sympathetic
3470 nervous system (Khalmatov 1964).

3471 **Phytochemistry:** A crystalline material, C₁₁H₁₂O₂, has been isolated from the essential oil (Khalmatov 1964). The principal
3472 components of the essential oil, from plants collected in the Moynkumy desert of southern Kazakhstan, were methyl
3473 lachnophyllate (80.1 %), β-pinene (4.8 %) and caryophyllene (1 %). Other compounds isolated from the essential oil were
3474 α-pinene, β-myrcene, limonene, camphor, caryophyllene oxide, etc. (Sadyrbekov et al. 2006b).

3475

Lagochilus gypsaceus Vved. – Lamiaceae	3476
Synonyms: None	3477
English name: Unknown	3478
Russian name: Заячья губа гипсовая (Zayach'ya guba gipsovaya)	3479
Uzbek name: Bozulbang	3480
Kyrgyz name: Unknown	3481
Description: Subshrub. Stems 30–40 cm high, woody at the very base, erect, often branched, with white, shiny bark, villous, subsequently becoming glabrous. Leaves petiolate, villous, rhomboid or wide-ovate in outline, 3–5-lobed, base cuneate; lobes oval or large-dentate. Inflorescences verticillasters with 4–6 flowers. Bracteoles awl-shaped, 3-sided, stiff. Flowers sessile. Calyx campanulate with spinescent lobes. Corolla 2-lipped, white or pink, with brown veins, 20–25 mm long. Fruits glabrous nutlets, 4–5 mm long.	3482 3483 3484 3485 3486
Other distinguishing features: Corolla 1–1.5 times longer than the calyx. Differs from related species by having villous stems.	3487 3488
Phenology: Flowers in May-August, fruits in June-September.	3489
Reproduction: Only by seeds.	3490
Distribution: Endemic plant of Qashqadaryo and Surxondaryo provinces of Uzbekistan; not found in Kyrgyzstan.	3491
Habitat: The adyr zone. Slopes of foothills with rocky debris and areas with a high diversity of soil minerals including gypsum.	3492 3493
Population status: Uncommon, sometimes in small populations.	3494
Traditional use: Infusions and decoctions of the plant are used to stop bleeding and as a sedative tea (Akopov 1981).	3495
Documented effects: In experiments with animals, an intravenous injection of a 10 % infusion of the plant extract accelerated coagulation of the blood by 30 % in 30 min, decreased the time of recalcification by 38 %, increased toleration of plasma to hepatitis by 35 %, and decreased blood pressure by 7 %. Preparations (infusion and tincture) made of the above-ground parts have hemostatic and sedative effects and decrease blood pressure. This plant is used in modern medicine as a preventive and therapeutic agents for various kinds of hemorrhage (traumatic, uterine, hemorrhoidal, pulmonary, lung, and nasal), and also to treat hemophilia and Werlhof's disease (Akopov 1981).	3496 3497 3498 3499 3500 3501
Phytochemistry: Leaves contain the diterpene alcohol lagochilin, 0.03 % essential oil, 11–14 % tannins, 7–10 mg% carotene, vitamin C, organic acids, calcium, and iron salts, and 0.6–0.7 % flavonoid glycosides. Lagochilin (1.98 %), tannins (2–2.7 %), ascorbic acid (106.29 mg%), carotene (4.39 mg%), and essential oils (0.083 %) were isolated from air dried plants (Akopov 1981).	3502 3503 3504 3505

3506 **Lagochilus platyacanthus Rupr. – Lamiaceae**3507 **Synonyms:** *Lagochilus iliensis* C.Y. Wu & S.J. Hsuan, *Lagochilus keminensis* Isakov, *Lagochilus macrodontus* Knorr.3508 **English name:** Unknown3509 **Russian name:** Зайцегуб плоскоколючий (Zaytsegub ploskokolyuchiy)3510 **Uzbek name:** Unknown3511 **Kyrgyz name:** Жалпак тикендуу ак тикен (Zhalpak tikenduu ak tiken)

3512 **Description:** Herbaceous perennial. Stems 15–45 cm, branching from the base, covered with bristly hairs. Leaves pinnatisect with linear or ovate lobes, ciliate-margined. Lower leaves rhomboid, winged-petiolate; upper leaves more rounded. 3513
3514 Inflorescence a verticillaster with 4–8 flowers; bracteoles lanceolate to linear-lanceolate, stiff-spinescent, densely covered with glandular hairs. Calyx narrowly campanulate, tomentose, with ovate or triangular teeth. Corolla pale pink, 2-lipped, 3515
3516 twice as long as calyx, upper lip 2 or 3 lobed. Fruits brown nutlets.

3517 **Other distinguishing features:** Bracteoles 7–12 mm long.3518 **Phenology:** Flowers in June–July, fruits in August.3519 **Reproduction:** By seeds.3520 **Distribution:** Ysyk-Kol, Chuy, and Naryn provinces of Kyrgyzstan; not found in the flora of Uzbekistan.3521 **Habitat:** In steppes and tree-shrub belts, and on pebbly to stony slopes.3522 **Population status:** Common, found in small groups.3523 **Traditional use:** No data.

3524 **Documented effects:** In experiments, an infusion of the aboveground parts showed low toxicity and hemostatic and sedative 3525
3526 effects equal to, and hypotensive effects surpassing, those of *Lagochilus inebrians* Bunge. An infusion promoted blood coagulation and possessed antibacterial activity (Rakhimova and Pulatova 1972).

3527 **Phytochemistry:** The aboveground parts contain flavonoids, organic acids, essential oil, diterpenoids (lagochilin), alkaloids 3528
(stachydrine), vitamin C, tannins, coumarins, lipids, etc. (Plant Resources of the USSR 1991).

3529

- Lagochilus platycalyx Schrenk ex Fisch. & Mey. – Lamiaceae** 3530
- Synonyms:** *Chlainanthus platycalyx* (Schrenk ex Fisch. & Mey.) Briq. 3531
- English name:** Unknown 3532
- Russian name:** Зайцегуб плоскочашечный, Заячья губа широкочашечная (Zaytsegub ploskochashechnyy, Zayach'ya guba shirokochashechnaya) 3533
- Uzbek name:** Unknown 3535
- Kyrgyz name:** Жазы чейчокчолуу ак тикен (Zhazy cheychokcholuu ak tiken) 3536
- Description:** Perennial subshrub with woody roots. Stems herbaceous, erect, simple or branched, 20–50 cm tall, densely covered with fine hairs. Leaves opposite, with winged petioles, rhomboid in outline, pinnatisect nearly to mid-vein, scattered-hairy; lobes ovate, elongate or linear; upper leaves spiny or awl-like. Inflorescences verticillasters with 4–6 flowers. Calyx narrow-campanulate, with short triangular (sometimes merged) lobes, appressed-hairy. Corolla pale pink, with dark veins, 2-lipped, upper lip with 2 short lobes, lower lip with 3 wide lobes, lateral lobes elongate, oblong. Fruits nutlets, glabrous. 3537–3542
- Other distinguishing features:** Bracts 3–7 mm long, hairy. Corolla as long or 1.5 times as long as the calyx. 3543
- Phenology:** Flowers in May-June, fruits in July. 3544
- Reproduction:** By seeds. 3545
- Distribution:** Jalal-Abad, Chuy, and Talas provinces of Kyrgyzstan; Toshkent province of Uzbekistan. 3546
- Habitat:** On pebbly to stony slopes of foothills, in dry steppes and on exposures. 3547
- Population status:** Common, found in small groups. 3548
- Traditional use:** No data. 3549
- Documented effects:** Effects on the cardio-vascular system and blood coagulability is equal to that of *Lagochilus inebrians* Bunge (Abdurakhmanov 1962). A tincture of the aboveground parts had low toxicity, hemostatic and sedative properties similar to those of *L. inebrians*, and hypotensive effects which surpassed those of *L. inebrians*. Clinical tests established the efficacy of the tincture for the treatment of hypertonic illness and as a hemostatic. A tincture of leaves and flowers exhibited hypotensive and sedative effects, and increased the speed of blood coagulation without increase of the prothrombin time (Alimbaeva 1961). 3550–3555
- Phytochemistry:** The plant contains essential oil, alkaloids (stachydrine, etc.), organic acids (chlorogenic, caffeic, hydroxycinnamic, and citric), flavonoids, diterpenoids (lagochilin), vitamin C, and tannins (Plant Resources of the USSR 1991; Nasrullaev and Makhsudova 1991; Zainutdinova et al. 1994; Kotenko et al. 1994). 3556–3558



▲ **Lachnophyllum gossypinum**
Bunge Photo: Alim Gaziev

◀ **Lagochilus platycalyx**
Schrenk ex Fisch. & Mey.
Photos: *top*: Evgeny Davkaev;
bottom left and bottom right:
Alexander Naumenko



◀ **Korolkowia sewerzowii**
(Regel) Regel Photos: *top and bottom*: Evgeny Davkaev;
center: Alexander Naumenko



▼ **Lagochilus platyacanthus** Rupr. Photos: Vladimir Epiktetov



Lallemantia royleana (Benth.) Benth. – Lamiaceae	3560
Synonyms: <i>Dracocephalum inderiense</i> Less. ex Kar. & Kir., <i>Dracocephalum royleanum</i> Benth., <i>Nepeta erodiifolia</i> Boiss.	3561
English name: Unknown	3562
Russian name: Лаллеманция Ройла (Lallemantsiya Royla)	3563
Uzbek name: Mallachoi	3564
Kyrgyz name: Ройл лаллеманциясы (Royl lallemantsiyasy)	3565
Description: Annual herb covered with dense, short pubescence. Stems simple or branching, 5–30 cm tall. Lower leaves petiolate, ovate, 1.5–4 cm long, 0.8–2.5 cm wide, margins crenate; upper leaves smaller, subsessile. Flowers in whorls of 4–6, arranged in erect, interrupted, terminal, spiciform inflorescences. Bracteoles up to 1.5 cm long, with 2–4 awned teeth. Calyx tubular, prominently nerved with short obtuse lobes. Corolla 2-lipped, 6.5–9 mm long, azure, outside pubescent and glandular. Fruits oblinear nutlets, 2.5–3 mm long, trigonous, glabrous, smooth, dark-brown.	3566 3567 3568 3569 3570
Other distinguishing features: Corolla slightly exceeding calyx in length. The leaves produce a distinct smell when crushed.	3571 3572
Phenology: Flowers in April-July, fruits in May-July.	3573
Reproduction: Only by seeds.	3574
Distribution: All regions of Uzbekistan; Osh, Chuy, Jalal-Abad and Talas provinces of Kyrgyzstan.	3575
Habitat: The chul, adyr, and tau zones.	3576
Population status: Common, as part of the ephemeral vegetation of foothills in the adyr zone.	3577
Traditional use: A decoction of the fruits is used in folk medicine as a diuretic and expectorant and to treat gastric diseases and asthenia. An infusion of the herb is recommended for coughs and gastric pains (Khalmatov 1964).	3578 3579
Documented effects: Pharmacological studies have proved the diuretic action of the aboveground plant parts. A water extract of the herb increased diuresis in rats to 52 %, but the plant was toxic, and caused the death of 20 % of the tested animals (Khalmatov 1964).	3580 3581 3582
Phytochemistry: Forty-six compounds were detected in the essential oil from the aboveground parts of <i>Lallemantia royleana</i> . Among them, verbenone and <i>trans</i> -carveol were found to be the major components of the oil (Ghannadi and Zolfaghari 2003). Plants collected in the Toshkent region contained traces of essential oils. The seeds of another related species, <i>Lallemantia iberica</i> , contain 27–35 % semi-drying oil. This oil is used for industrial purposes, as a food, and to produce soap (Ogolevitz 1951).	3583 3584 3585 3586 3587

3589 **Leontice ewersmanni Bunge – Berberidaceae**3590 **Synonyms:** *Leontice leontopetalum* ssp. *ewersmannii* (Bunge) Coode.3591 **English name:** Unknown3592 **Russian name:** Леонтица Эверсмана (*Leontitsa Eversmana*)3593 **Uzbek name:** Yersovun3594 **Kyrgyz name:** Эверсман леонтичасы (*Eversman leontitsasy*)

3595 **Description:** Herbaceous perennial with a large, ovate tuber, 5–15 cm wide. Stem 20–60 cm tall, with subterranean part
3596 nearly as long. Basal leaves 1 or 2, with 3 petioluled lobes; each lobe trisected, middle lobe tripartite with sessile, bisected
3597 lateral lobes; upper leaves 3–5, lower 2 similar to basal leaves, the most upper leaves smaller and less divided or entire.
3598 Inflorescence apical, paniculiform, formed of racemes with 20–40 flowers. Flowers on long, horizontally spreading pedi-
3599 cels. Sepals yellow, petaloid. Petals 6, reduced, yellow. Stamens 6. Fruit an inflated capsule, ca. 15 mm in diameter. Seeds
3600 1–2 per fruit, 5 mm wide, spherical, smooth.

3601 **Other distinguishing features:** Petioles of basal leaves originating below ground. Tubers can grow 15–40 cm under the soil
3602 surface and weigh more than 1 kg (2.2 lbs).

3603 **Phenology:** Flowers in March, fruits in April.3604 **Reproduction:** By seeds.

3605 **Distribution:** Toshkent, Jizzax, Samarqand, Namangan, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Chuy
3606 province of Kyrgyzstan.

3607 **Habitat:** The chul and adyr zones. Sandy and clay deserts, loess hills in the foothills.3608 **Population status:** Uncommon.

3609 **Traditional use:** The powdered tuber is used in folk medicine to treat wounds and is smoked to treat syphilis. An infusion
3610 of the tuber is drunk for treating delayed menstruation and bladder stones (Khalmatov 1964).

3611 **Documented effects:** In acute tests on animals with the alkaloids pachycarpine and d-lupanine, doses of 2–10 mg/kg
3612 decreased arterial blood pressure. Starting from the dose of 1–3 mg/kg it depressed and at the dose of 5 mg/kg it com-
3613 pletely blocked parasympathetic cardiac ganglions. At the same dose, it potentiated hypotensive effect of acetylcholine
3614 and hypertensive effect of adrenaline and decreased reaction of cat's third eyelid, arterial pressure and respiration caused
3615 by cytisine introduction. Seventy to eighty percent of d-lupanine is excreted from the body: 50–70 % with urination,
3616 10–14 % through defecation and 30–40 % turned into oxylupanine (Wittenburg and Nehring 1965). The alkaloid had
3617 minor tonic action on uterine muscles and had anticholinesterase action (Trutneva and Berezhinskaya 1960).

3618 **Phytochemistry:** Roots contain tannins, up to 1.5 % total alkaloids (leontidine, leontine, leontamine, pachycarpine, and
3619 d-lupanine) and up to 30 % starch. The plant contains saponins with a hemolytic index of 1:240 in the aboveground por-
3620 tion of the plant and 1:6,000 in the tubers. Taspine, methylcytidine, and isoleontine were isolated from aboveground por-
3621 tion of the plant (Yunusov 1981).

3622

Leonurus turkestanicus V. Krecz & Kuprian. – Lamiaceae	3623
Synonyms: <i>Leonurus cardiaca</i> ssp. <i>turkestanicus</i> (V. Krecz. & Kuprian.) Rech.	3624
English name: Turkestan motherwort	3625
Russian name: Пустырник туркестанский (Pustyrnik turkestanskiy)	3626
Uzbek name: Arslon kuirug	3627
Kyrgyz name: Туркстан дулой чалканы (Turkstan duloy chalkany)	3628
Description: Perennial herb with woody rhizome. Stems 50–150 cm tall, purple-red, branched, pubescent or glabrous.	3629
Leaves opposite, petiolate, wide-ovate to nearly circular in outline, palmatipartite; lobes pinnately divided into broadly lanceolate lobules. Flowers sessile, 15–20 per verticillaster, forming spiciform inflorescences. Bracts awl-like, pubescent.	3630
Calyx 8–9 mm long, funnelform, short-pubescent, with triangular spinescent lobes. Corolla pink-lilac, ca. 1 cm long, 2-lipped, villous outside. Fruits triquetrous nutlets, light brown.	3631
	3632
	3633
Other distinguishing features: Upper corolla lip obovate, lower lip with 3 lobes, middle lobe larger than lateral lobes.	3634
Phenology: Flowers in June-July, fruits in July-August.	3635
Reproduction: By seeds.	3636
Distribution: Toshkent, Samarqand, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.	3637
Habitat: The tau zone. On stony, shallow-soiled slopes, among trees and shrubs.	3638
Population status: Uncommon.	3639
Traditional use: A decoction of the aboveground parts is used to treat heart, stomach and nervous system diseases (Khalmatov 1964). A tea and an infusion of the aboveground parts are used to treat nervous disorders, hypertension, hysteria, epilepsy, tachycardia, gastrointestinal, and female diseases, and are used as soporific, anti-inflammatory, diaphoretic, and laxative remedies (Khodzhimatov 1989).	3640
	3641
	3642
	3643
Documented effects: Studies show that a tincture of the herb has a sedative effect, which is twice as strong as the effect of a valerian tincture. The tincture also causes decreased arterial pressure and strengthens the contraction of uterus muscles (Khalmatov 1964). Stachydrine exhibited protective effects when given to rats with experimental myocardial ischemia-reperfusion injury (Ma and Yang 2006).	3644
	3645
	3646
	3647
Phytochemistry: The aboveground parts contain alkaloids (stachydrine), flavonoids, essential oils, tannins, saponins, resins, bitter substances and other compounds (Khalmatov 1964; Pulatova 1969; Khalmatov and Kosimov 1994).	3648
	3649
	3650

3651 **Lepidium perfoliatum L. – Brassicaceae**3652 **Synonyms:** *Crucifera diversifolia* E.H.L. Krause, *Nasturtium perfoliatum* (L.) Besser, *Nasturtium perfoliatum* (L.) Kuntze.3653 **English name:** Claspig pepper-grass3654 **Russian name:** Клоповник пронзенный (Клоровник пронзеннуу)3655 **Uzbek name:** Unknown3656 **Kyrgyz name:** Кучакталган сасык кычы (Kuchaktalghan sasyk kychy)3657 **Description:** Herbaceous biennial. Stems up to 20–25 cm tall, erect, branched, hairy at the bottom, glabrous towards the top.3658 Leaves alternate, dimorphic; basal leaves (in rosette) and lower cauline leaves lanceolate, bi- or tripinnatisect with acute,
3659 simple or trilobate segments, hairy; upper cauline leaves sessile, ovate, cordate or nearly round, acute, amplexicaul, gla-
3660 brous. Inflorescence racemose. Sepals 4. Petals 4, ca. 1.5 mm long, pale-yellow. Stamens 6. Fruits glabrous siliques,
3661 orbicular or rhombic, thin, 4–5 mm long. Seeds 1.5–2.5 mm long, 0.75–1.5 mm wide, dark-brown.3662 **Other distinguishing features:** Seeds have a narrow wing around their entire edge.3663 **Phenology:** Flowers in March-May, fruits in April-June.3664 **Reproduction:** Only by seeds.3665 **Distribution:** All regions of Uzbekistan and Kyrgyzstan.3666 **Habitat:** The chul and adyr zones.3667 **Population status:** Common, found in small populations as a part of ephemeral associations.3668 **Traditional use:** A decoction of the herb is taken to treat headaches, and ground seeds mixed with other pharmaceuticals are
3669 recommended for treatment of general weakness and to reinforce the nervous system. Avicenna applied the plant as a
3670 dressing or ointment with honey to treat “hard” and malignant tumors, as well as podagra, and used as an expectorant
3671 mixed with other drugs (Khalmatov 1964).3672 **Documented effects:** No data.3673 **Phytochemistry:** The herb contains glycosides, which are produced after the enzymic hydrolysis of mustard essential oil.3674 Seeds contain 12–19 % drying oil. There is a possibility for the presence of prussic acid in young plants in the spring
3675 (Khalmatov 1964). The plant contains the flavonoid lepidoside (Fursa and Litvinenko 1970). The seeds were found to
3676 contain quercetin derivatives, as well as 18.71 % oil, in which alpha-linolenic, oleic, erucic and eicosenoic acid were the
3677 most abundant (Dolya et al. 1973a, b).


3678

Lepidolopsis turkestanica (Regel & Schmalh.) Poljakov – Asteraceae	3679
Synonyms: <i>Chrysanthemum turkestanicum</i> (Regel & Schmalh.) Gilli, <i>Crossostephium turkestanicum</i> Regel & Schmalh.	3680
English name: Unknown	3681
Russian name: Лепидолопис туркестанский (Lepidolopsis turkestanskiy)	3682
Uzbek name: Zarkuloq	3683
Kyrgyz name: Туркстан лепидолописи (Turkstan lepidolopsisi)	3684
Description: Perennial herb with thick rhizome. Stems solitary or few, 40–100 cm high, erect, leafy, with long and short hairs, later becoming glabrous. Basal leaves and lower stem leaves up to 10–15 cm long and 3.5 cm wide, petiolate; blades blue-gray-green, sparsely hairy, oblanceolate in outline, bi- or tripinnatisect with narrow-linear segments, terminal segments with short, cartilaginous tips; upper leaves reduced, sessile. Inflorescences composed of many small capitula arranged in compressed spicate-panicles, 15–30 cm long; involucre 4–6 mm in diameter, often golden-tinged. Flowers all tubular disk florets, yellow. Fruits achenes, 1.5–1.75 mm long, angled on top, with a paleaceous corona that is deeply divided into 8–12 narrow teeth.	3685 3686 3687 3688 3689 3690 3691
Other distinguishing features: Basal leaves senesce early. Receptacle glabrous.	3692
Phenology: Flowers in June-August, fruits in July-September.	3693
Reproduction: Only by seeds.	3694
Distribution: Toshkent, Samarqand, Farg'ona, Andijon, and Surxondaryo provinces of Uzbekistan; Osh, Jalal-Abad and Batken provinces of Kyrgyzstan.	3695 3696
Habitat: Loess foothills, most often in the adyr zone.	3697
Population status: Uncommon. Found in small populations but more often as single individuals.	3698
Traditional use: A decoction of flower heads is used in folk medicine to treat chest pains, heavy breathing, malaria, and delayed menstruation, and is also used as a vermifuge and diuretic remedy (Khalmatov 1964).	3699 3700
Documented effects: Pharmacological investigations of the plant extract showed effects on uterine activity (Khalmatov 1964).	3701 3702
Phytochemistry: This species contained traces of alkaloids (Khalmatov 1964), and at flowering it contained 0.12–0.13 % essential oil (Kudryashev 1932).	3703 3704



▲ **Lallemantia royleana (Benth.) Benth.** Photo: Alexander Naumenko

▼ **Lepidium perfoliatum L.**
Photos: *top*: Maxim Kucherov;
bottom: Clinton Shock



▲ **Leonurus turkestanicus V. Krecz & Kuprian.**
Photos: Evgeny Davkaev



▼ **Leontice ewersmanni Bunge** Photos: Alexander Naumenko



Lithospermum officinale L. – Boraginaceae	3705
Synonyms: None	3706
English name: European stoneseed, common gromwell	3707
Russian name: Воробейник лекарственный (Vorobeynik lekarstvennyy)	3708
Uzbek name: Пончоор	3709
Kyrgyz name: Дары таранчы чоп (Dary taranchy chop)	3710
Description: Herbaceous perennial, with stout rhizome. Stems single to many, 30–100 cm tall, branched above. Leaves	3711
opposite, nearly sessile, lanceolate to oblanceolate, 3–8 cm long, 5–15 mm wide, leaves on upper portion of stems	3712
crowded. Inflorescences dense cymes, in upper leaf axils. Calyx lobes 5, oblong-linear. Corolla 3–6 mm long, tubular with	3713
5 lobes, yellowish or greenish-white. Stamens inserted at middle of corolla tube. Stigma capitate. Fruits ovoid nutlets,	3714
about 4 mm long, white or light brown, shiny.	3715
Other distinguishing features: Stems and leaves are scabrid-hairy.	3716
Phenology: Flowers in May-June, fruits in June-July.	3717
Reproduction: By seeds.	3718
Distribution: Ysyk-Kol, Chuy, and Osh provinces of Kyrgyzstan; Tashkent, Farg'ona, Andijon, Samarqand, Buxoro and	3719
Surxondaryo provinces of Uzbekistan.	3720
Habitat: In the tallgrass-meadow belt, in meadows, river floodplains, and among bushes.	3721
Population status: Common, found growing as single plants.	3722
Traditional use: The freshly ground plant is applied to heal bruises and cuts (Khalmatov 1964).	3723
Documented effects: In preclinical tests an extract of the plant showed satisfactory results for treatment of hyperpituitarism	3724
and displayed antihormonal properties (Vyazovskaya 1963). When administered together with thyroid stimulating hor-	3725
mone (TSH), an extract of the plant blocked the TSH-induced increase in endocytotic activity of the thyroid glands fol-	3726
lowed by a strong decline of thyroid hormone levels. When the extract was injected alone it caused a decline in endogenous	3727
TSH-levels as well as in thyroidal secretion and thyroid hormone levels (Winterhoff et al. 1983). A decoction of the	3728
aboveground parts is used to treat the gastrointestinal tract (Utkin 1931). A water extract possesses antigonadotropic,	3729
contraceptive, and spermatocidal properties. A decoction of the ground fruits is parturifacient, and is used to treat dys-	3730
menorrhea, kidney diseases, kidney stones, and dyspepsia. Roots show protistocidal activity (Dilman et al. 1968).	3731
Phytochemistry: All plant parts contain cyclitols, organic acids (citric, malic, maleic, succinic, and fumaric), steroids, phe-	3732
nylcarbonic acids and their derivatives, tannins, and flavonoids. The underground parts contain carbohydrates (glucose,	3733
saccharose, glucofructose, and fructose), cyanogenic compounds, phenylcarbonic acids, naphthoquinones, and fatty acids.	3734
The aboveground parts contain organic acids, flavonoids, and phenylcarbonic acids. Fruits contain cyclitols, aliphatic	3735
alcohols, steroids, vitamin E, phenylcarbonic acids and their derivatives, tannins, fatty oil, fatty acids, and pyrrolizidine	3736
alkaloids. Seeds contain carbohydrates, aliphatic alcohols, steroids, fatty oil and fatty acids (Dilman et al. 1968; Krenn	3737
et al. 1994).	3738

3740 **Lycopus europaeus L. – Lamiaceae**3741 **Synonyms:** None3742 **English name:** Gypsywort3743 **Russian name:** Зюзник европейский (Zyuznik evropeyskiy)3744 **Uzbek name:** Khorok, Tadzh3745 **Kyrgyz name:** Европа ликопусу (Evropa likopusu)

3746 **Description:** Herbaceous perennial, with rhizomes and stolons. Stems single or many, 20–90 cm tall, erect. Leaves opposite,
3747 3–9 cm long, 1–4 cm wide, oblong-elliptic to lanceolate-elliptic, coarsely dentate, base attenuate, apex acuminate.
3748 Inflorescences dense verticillasters, 18–20-flowered; bracteoles linear-subulate. Calyx with 4 or 5 triangular-lanceolate
3749 lobes. Corolla 2-lipped, 3 mm long, white with reddish-purple spots. Fruits oblong nutlets, glabrous.

3750 **Other distinguishing features:** Two exerted and two reduced stamens. Upper leaves coarsely dentate.3751 **Phenology:** Flowers in June–July, fruits in August–September.3752 **Reproduction:** By seeds and rhizomes.

3753 **Distribution:** Jalal-Abad, Chuy, and Osh provinces of Kyrgyzstan. Karakalpakstan autonomous republic, Toshkent, Andijon,
3754 Farg'ona, Samarqand, Buxoro, Surxondaryo and Xorazm provinces of Uzbekistan.

3755 **Habitat:** On along rivers and in wet meadows, sometimes in water.3756 **Population status:** Common, sometimes found in small groups.

3757 **Traditional use:** The herb is used to reduce swelling and as a hemostatic for uterine bleeding (Akopov 1990). A decoction
3758 and infusion of the aboveground parts is used to normalize increased heart rate due to stress, as a sedative and an anti-
3759 pyretic, and to treat uterine bleeding (Khalmatov 1964). Extracts from the plant are traditionally used to treat mild forms
3760 of hyperthyroidism (Vönhoff et al. 2006).

3761 **Documented effects:** Clinical studies showed that the herb normalizes the function of the thyroid gland, has sedative and
3762 hypotensive effects, and dilates the coronary arteries. An infusion or tincture is recommended for the above effects
3763 (Akopov 1990). In Azerbaijan an infusion and decoction is used to treat heart diseases and lung tuberculosis. In Bulgaria
3764 a decoction is used to treat rheumatis. Based on preclinical tests, an alcoholic solution (of the polyphenols) and ointment
3765 accelerated healing of wounds and were effective in treating purulent otitis. Preparations of this species are proposed for
3766 treatment of atherosclerosis, hypertension, and coronary insufficiency. In experiments, a water infusion normalized pro-
3767 duction of thyroxine, slowed development of goiters, lowered metabolism in cases of exophthalmic goiters, showed low
3768 toxicity, and was recommended for clinical studies as a treatment for thyroidtoxicosis. A liquid extract possessed antithy-
3769 roid activity, normalized the gas content of blood, reduced the ability of the thyroid gland to accumulate iodine, and posi-
3770 tively influenced lactation. An ether extract also showed antibacterial and antifungal activity (Plant Resources of the
3771 USSR 1991). High doses of an extract of the plant caused a reduction of thyroid hormone levels in animal experiments,
3772 whereas in hyperthyroid patients treated with low doses, an improvement of cardiac symptoms was reported without
3773 major changes in thyroid hormone concentrations. Extracts diminished thyroidal secretion, reduced the plasma concentra-
3774 tion of thyroxine (T_4), triiodothyronine (T_3), and inhibited the conversion of thyroxine to triiodothyronine. *Lycopus* extract
3775 also reduced heart rate and blood pressure and alleviated cardiac hypertrophy (Vönhoff et al. 2006). Two diterpenes iso-
3776 lated from the plant caused twofold potentiation of the activities of tetracycline and erythromycin against two strains of
3777 multi-drug resistant *Staphylococcus aureus* (Gibbons et al. 2003).

3778 **Phytochemistry:** The plant contains organic acids (tartaric, citric, and malic), essential oil, diterpenoids (phytol), triterpe-
3779 noids, steroids, saponins, alkaloids, choline, vitamin C, carotene, phenylcarbonic acids (rosmarinic acid), tannins, cou-
3780 marins, flavonoids, cardiac glycosides, carbohydrates and anthocyanins (cyanin and pelargonin; Akopov 1990; Plant
3781 Resources of the USSR 1991).

- Marrubium anisodon K. Koch. – Lamiaceae** 3783
- Synonyms:** *Marrubium alternidens* Rech. f., *Marrubium kusnezowii* N.P. Popov. 3784
- English name:** Horehound 3785
- Russian name:** Шандра очереднозубая, Шандра очереднозубчатая (Shandra ocheryodnozubaya, Shandra ocheryodnozubchataya) 3786
- Uzbek name:** Devoltegiuit 3788
- Kyrgyz name:** Ар тшчелуу маррубium (Ar tishcheluu marrubium) 3789
- Description:** Herbaceous perennial. Stems simple or branched, woolly-hairy, 30–100 cm high. Leaves opposite; lower leaves elliptic, orbicular or almost oblong, 5 cm long, 4 cm wide, base wide-cuneate, serrate-crenate, petiolate; upper leaves similar to lower leaves but smaller, wrinkled, densely pubescent. Inflorescences axillary verticillasters, multiflorous. Calyx hairy, with 10 teeth, 5 long alternating with 5 short. Corolla 2-lipped, 9–11 mm long, pale pink, pale yellow or white, stellate-hairy on the outside. Fruits obovoid nutlets, triquetrous, 1.5 mm long, glabrous, dark-brown or black. 3790–3794
- Other distinguishing features:** Corolla 1.5 times as long as calyx. 3795
- Phenology:** Flowers and fruits in May-September. 3796
- Reproduction:** By seeds. 3797
- Distribution:** All regions of Uzbekistan and Kyrgyzstan. 3798
- Habitat:** The chul, adyr, and tau zones. A weed, mainly a ruderal. 3799
- Population status:** Common. 3800
- Traditional use:** A decoction of the herb is used in folk medicine to treat chronic catarrh of the respiratory tract and throat diseases; also used as a mouthwash to treat toothaches (Khalmatov and Kosimov 1994). 3801–3802
- Documented effects:** Used as a sedative, it exceeds twice the action of valerian tincture; decreased blood pressure and increased tone of uterine muscles. The alkaloid stachydrine is slightly toxic. At doses of 5–100 mg/kg there was almost no effect on blood pressure. At the dose of 5 mg/kg there was a positive chronotropic action on the heart, where as at doses 10–100 mg/kg, there was a negative chronotropic action on the heart. Injected intravenously, starting from the dose of 5 mg/kg, stachydrine stimulated blood coagulation in dogs (Akopov et al. 1958). Research showed that marrubinic acid has choleric effects (Khalmatov 1964). An extract of the plant showed significant antibacterial activity against *Escherichia coli* (Bazzaz and Haririzadeh 2003). 3803–3809
- Phytochemistry:** The aboveground plant parts contained 0.4 % essential oils, 1.12 % flavonoids, the alkaloid stachydrine, resins, 116.57 mg% vitamin C, and other compounds (Khalmatov and Kosimov 1994). Two diterpenoids, vulgarol and marubiin, have been isolated from the plant. (Sagitdinova et al. 1996). The total amino acids, polysaccharides, tanning agents, acids, flavonoids, phenols, essential oils, coumarins, saponins and alkaloids, from plants at the budding, flowering, and fruiting stages, were quantified by Kurbatova et al. (2003). 3810–3814

3816 **Mediasia macrophylla (Regel & Schmalh.) Pimenov – Apiaceae**

3817 **Synonyms:** *Athamanta macrophylla* (Regel & Schmalh.) Korovin, *Seseli macrophyllum* Regel & Schmalh.

3818 **English name:** Unknown

3819 **Russian name:** Медиазия крупнолистная (Mediaziya krupnolistnaya)

3820 **Uzbek name:** Hunich, Alkor

3821 **Kyrgyz name:** Unknown

3822 **Description:** Herbaceous perennial with a 1.5–3 cm thick root. Stems several, 0.5–1.5 m tall, hollow, round, striated, covered
3823 with thick, short hairs. Leaves alternate, broadly ovate, 20–60 cm long and 20–40 cm wide, bipinnatipartite, coated with
3824 short hairs, long-petiolate; leaflets 4.5–12 cm long, 3.5–10 cm wide, bases cordate. Inflorescences compound umbels,
3825 apical, 5–10 cm wide, with 13–23 unequal rays; umbellets 5–6 mm wide, ca. 20-flowered. Flowers white or greenish-
3826 yellow, very hairy outside. Fruit a schizocarp with 2 mericarps; mericarps flattened, oval in outline, 5–6 mm long, hairy.

3827 **Other distinguishing features:** Leaflets of upper leaves often trilobate. Leaves produce a pleasant odor when crushed.

3828 **Phenology:** Flowers in June-July, fruits in August.

3829 **Reproduction:** By seeds.

3830 **Distribution:** Toshkent, Samarqand, Farg'ona, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh, Chuy and
3831 Jalal-Abad provinces of Kyrgyzstan.

3832 **Habitat:** Tau zone. Stony slopes and large-fragmental taluses in the tree-shrub belt.

3833 **Population status:** Common, found as individual plants.

3834 **Traditional use:** A decoction of the roots is taken as a hemostatic. Local people use the fruit as a spice (Khalmatov 1964).

3835 **Documented effects:** No data.

3836 **Phytochemistry:** This species contained 1.67 % essential oils, the composition of which included aldehydes (Khalmatov
3837 1964). According to Baser et al. (1997b), a total of 33 compounds were found in the essential oil, of which the principal
3838 components were *p*-cymene (27.2 %), thymol (15.1 %), carvacrol (12.5 %), and palmitic acid (2.9 %). A variety of neu-
3839 tral-, glyco- and phospho-lipids were isolated and identified from the leaves, and various free and bound fatty acids and
3840 carotenoids were quantified (Chernenko et al. 2002).

3841



▲ **Lithospermum officinale** L. Photos: *left*: Sergey Mayorov; *right*: Petr Filippov

▼ **Lycopus europaeus** L. Photos: *top left* and *lower right*: Rostislav Lezhoyev; *top right*: Sergey Appolonov



▲ **Marrubium anisodon**
K. Koch. Photos:
Vladimir Epiktetov



▼ **Mediasia macrophylla**
(Regel & Schmalh.) Pimenov
Photos: Alim Gaziev



3842 **Melilotus officinalis (L.) Pall. – Fabaceae**3843 **Synonyms:** *Melilotus graveolens* Bunge, *Melilotus suaveolens* Ledeb., *Trifolium officinale* L.3844 **English name:** Yellow sweetclover3845 **Russian name:** Донник лекарственный (Donnik lekarstvennyy)3846 **Uzbek name:** Kashkar beda3847 **Kyrgyz name:** Дары кашка беде (Dary kashka bede)

3848 **Description:** Herbaceous biennial, with branching taproot. Stems one to many, up to 2 m tall. Leaves alternate, trifoliate,
3849 petiolate, stipulate; leaflets oblanceolate to obovate, serrulate, terminal leaflet stalked. Inflorescences axillary racemes,
3850 5–15 cm long. Flowers 5–7 mm long. Calyx 2–2.5 mm long, toothed. Corolla papilionaceous, yellow. Fruit an oval
3851 legume, 3–5 mm long, with a beaked tip, 1–2-seeded. Seeds greenish-yellow.

3852 **Other distinguishing features:** Stipules lanceolate, entire.3853 **Phenology:** Flowers in May-June, fruits in July-August.3854 **Reproduction:** By seeds.3855 **Distribution:** Almost all provinces of Kyrgyzstan and Uzbekistan.3856 **Habitat:** In meadows, fallow fields, along rivers and roads, and in cultivated fields.3857 **Population status:** Common, sometimes found in dense groups.

3858 **Traditional use:** An infusion of the herb is drunk to treat chronic catarrh of the bronchial tubes, migraines, and hypertension,
3859 for bladder and kidney pain, and is used during menopause. It is applied externally in the form of compresses, plasters,
3860 and as a wash, which are used as an emollient and analgesic to treat furuncles, carbuncles, purulent wounds and inflamed,
3861 pus-producing infections of the middle ear (Maznev 2004).

3862 **Documented effects:** Coumarins from this species suppress the central nervous system and possess anticonvulsive and nar-
3863 cotic properties. After radiation treatments leucopenia patients treated with coumarins had increased leucocytes. The
3864 preparation *Dicumalin* (*Dicumarol*) has the ability to inhibit blood coagulation and is widely used as an anticoagulant and
3865 anti-vitamin K₁ to treat thrombophlebitis and heart attacks. This preparation must be used very carefully, not only because
3866 of inhibition of blood coagulation, but also because of increased permeability of capillaries and prolonged bleeding
3867 (Khalmatov et al. 1984). An extract of the plant had anti-inflammatory effects due to the activation of circulating phago-
3868 cytes and lowered citrulline production (Plesca-Manea et al. 2002). A coumarinic extract from the plant was effective in
3869 reducing lymphedema in 79 % of patients with chronic lymphedema of the upper arm, caused by lymphadenectomy for
3870 breast cancer (Pastura et al. 1999).

3871 **Phytochemistry:** The herb contains coumarins, melilotin, melilotic acid, melilotocide, purine derivatives, essential oil, vita-
3872 mins C and E, carotene, protein, and lipids. The seeds contain protein, fatty oils, starch, and alkaloids (Tolmachev 1976;
3873 Khalmatov et al. 1984; Chikov 1989; Martino et al. 2006).

3874

- Melissa officinalis L. – Lamiaceae** 3875
- Synonyms:** *Melissa bicornis* Klokov. 3876
- English name:** Lemon balm 3877
- Russian name:** Мелисса лекарственная (Melissa lekarstvennaya) 3878
- Uzbek name:** Limonuit 3879
- Kyrgyz name:** Дары мелиссасы (Dary melissasy) 3880
- Description:** Herbaceous perennial. Stems up to 120 cm tall, 4-sided, hairy. Leaves opposite, ovate, up to 7.5 cm long and 2–4 cm wide, serrate-crenate, hairy above, nearly glabrous below, long-petiolate. Inflorescences axillary verticillasters, 2–14-flowered. Calyx 2-lipped, angular, lobes about 2/3 as long as tube. Corolla white, yellowish or pinkish, 2-lipped, upper lip 2-lobed, lower lip 3-lobed. Fruits obovoid nutlets, brown. 3881
3882
3883
3884
- Other distinguishing features:** Crushed leaves have a lemon scent. 3885
- Phenology:** Flowers in June-July, fruits August-September. 3886
- Reproduction:** By seeds. 3887
- Distribution:** Jalal-Abad, Osh, and Chuy provinces of Kyrgyzstan; Toshkent and Surxondaryo provinces of Uzbekistan. 3888
- Habitat:** In Persian walnut forests, deciduous forests, on shady slopes, and among shrubs. 3889
- Population status:** Common, forming dense groups. 3890
- Traditional use:** This herb is widely used to treat migraines, insomnia, gynecological diseases, gout, dizziness, and anemia (Poludenny and Zhuravlev 2000). It is used as an antispasmodic for cardiovascular disease, as an analgesic, sedative, hypotensive, diuretic, and to improve digestion and to treat tympanites and pregnancy toxicosis (Kurochkin 1998). 3891
3892
3893
- Documented effects:** Preparations of this species are used as a sedative, anticonvulsive, analgesic, and anti-flu medicine. 3894
It is used as a cardiac remedy and acts by slowing down the rate of breaths and heartbeats and by reducing tachycardia, palpitations, shortness of breath, and chest pain (Maznev 2004). An extract of the plant given orally produced a significantly better outcome in cognitive function in patients with mild to moderate Alzheimer's disease, than a placebo given to the control group (Akhondzadeh et al. 2003). Healthy people who received an extract of the herb orally exhibited a reduction in negative effects of laboratory induced stress and, at a higher dose, significantly increased the speed of mathematical processing with no reduction in accuracy (Kennedy et al. 2004). The essential oil has anti-tumor and anti-oxidant activities (de Sousa et al. 2004). 3895
3896
3897
3898
3899
3900
3901
- Phytochemistry:** The leaves contain tannins, caffeic, oleanolic, ursolic acids, and essential oil (including citral, citronellol, myrcene, and geraniol). The aboveground parts contain ascorbic acid, potassium, calcium, magnesium, iron, manganese, copper, zinc, molybdenum, chromium, aluminium, barium, tungsten, silicon, nickel, sulfur, lead, and selenium. The seeds contain fatty oil (Volinsky et al. 1983; Carnat et al. 1998; de Sousa et al. 2004; Maznev 2004). 3902
3903
3904
3905

3907 **Melo agrestis (Naudin) Pang. – Cucurbitaceae**3908 **Synonyms:** *Cucumis agrestis* (Naudin) Grebensc., *Cucumis melo* var. *agrestis* Naudin.3909 **English name:** Unknown3910 **Russian name:** Дыня полевая (Дупуа polevaya)3911 **Uzbek name:** It qavun3912 **Kyrgyz name:** Жапайы коон (Zhapayy koon)

3913 **Description:** Herbaceous taprooted annual with stiff, rough hairs. Stems prostrate, multiple, spreading, branched, slightly
3914 edged, 30–100 cm long. Leaves alternate, oblong or oblong-oval, 4–6 cm wide, slightly 3–5-lobed, seldom deeply notched,
3915 bristly-hairy, petiolate. Flowers uni- or bisexual. Staminate flowers in umbelliform inflorescences; pistillate flowers soli-
3916 tary. Calyx campanulate, 5-lobed, densely hairy. Corolla yellow, broadly funnellform, usually with 5 deep lobes. Fruit an
3917 oval-oblong berry (pepo), 2–5 cm long, usually green, yellowing at maturity, almost no aroma; rind rough, with a pattern
3918 in the form of deep-green, solid or interrupted, longitudinal stripes. Seeds small, white-yellowish, oval.

3919 **Other distinguishing features:** Male flowers have 5 stamens, 4 in pairs and the fifth free. The pulp of the fruit tastes sour or
3920 bitter and is greenish-white with a large amount of cucumber-like placenta.

3921 **Phenology:** Flowers in June-September, fruits in August-September.3922 **Reproduction:** Only by seeds.3923 **Distribution:** All regions of Uzbekistan; Osh, Chuy and Jalal-Abad provinces of Kyrgyzstan.3924 **Habitat:** The chul zone. As a weed in cotton and melon fields, rarely along canals and river banks.3925 **Population status:** Not common, found as single individuals.3926 **Traditional use:** A root decoction is used to treat edema and jaundice, and is used as mouthwash to treat bumps in the mouth.

3927 A fruit decoction is prescribed externally to treat eczema (Khalmatov 1964).

3928 **Documented effects:** Cultivated forms of this species are used as a food with medicinal value to treat asthenia, constipation,
3929 and hepatitis and are used as a diuretic and prophylaxis to prevent arteriosclerosis and anemia (Karimov and Shomakhmudov
3930 1993).

3931 **Phytochemistry:** Roots collected in the Toshkent region contained 1.16 % tannins and up to 2 % sugars. Stems contained
3932 0.87 % tannins, up to 4 % sugars, 0.4 % titratable organic acids, and alkaloid traces. Leaves contain 1.74 % tannins,
3933 0.53 % titratable organic acids, and alkaloid traces; fruits contain up to 2 % sugars, 1.07 % titratable organic acids, and
3934 alkaloid traces (Khalmatov 1964).

3935

- Mentha asiatica Boriss. – Lamiaceae** 3936
- Synonyms:** *Mentha kopetdaghensis* Boriss., *Mentha longifolia* (L.) Huds. var. *asiatica* (Boriss.) Rech. f., *Mentha vagans* Boriss. 3937
- English name:** Asian mint 3938
- Russian name:** Мята лесная (Myata lesnaya) 3939
- Uzbek name:** Yalpeez 3940
- Kyrgyz name:** Жалбыз (Zhalbyz) 3941
- Description:** Herbaceous perennial, with rhizomes. Stems erect, 40–100 cm tall, 4-sided, branched, finely hairy. Leaves short-petiolate, ovate, elliptic or oblanceolate, margins serrate-dentate, both sides finely hairy, very glandular on underside; upper leaves sessile. Inflorescences verticillasters in terminal, cylindrical spikes; bracts awl-shaped, equal in length to the calyx. Calyx campanulate with linear teeth, densely hairy. Corolla 4–5 mm long, lilac, funnellform. Fruits ovoid nutlets. 3942
- Other distinguishing features:** Leaves in inflorescence tapering to a point and extending past the verticillasters. 3943
- Phenology:** Flowers in July-September, fruits September-October. 3944
- Reproduction:** By seeds and rhizomes. 3945
- Distribution:** Jalal-Abad, Osh, and Chuy provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Andijon, Farg'ona, Samarqand, Buxoro and Surxondaryo provinces of Uzbekistan. 3946
- Habitat:** In wet places and near canals, springs and streams. 3947
- Population status:** Common, forming dense groups. 3948
- Traditional use:** In Tibetan, Chinese, Mongolian, Indian and Central Asian folk medicine an infusion and decoction of this plant is used as an anti-inflammatory, hemostatic, and is used to treat wounds, gastritis, dysentery, diarrhea, colitis, gastralgia, tuberculosis, respiratory infections, pertussis, and toothaches. An infusion of the leaves and inflorescences is used as a choleric and to treat gall bladder diseases (Minayeva 1991). 3949
- Documented effects:** In an evaluation for antimicrobial activity, essential oils from the related species *Mentha longifolia* ssp. *longifolia* and *Mentha sylvestris* L. exhibited activity against 30 different microorganisms including *Bacillus subtilis*, *Micrococcus luteus*, *Escherichia coli*, *Serratia marcescens* and *Aspergillus oryzae* (Carvalho et al. 1999; Gulluce et al. 2007). 3950
- Phytochemistry:** Aboveground parts contain essential oil with menthol, menthone, carvacrol, and pulegone (Khalmatov 1964; Gulluce et al. 2007). Thirty-seven compounds were characterized representing 97 % of the total components detected. The major constituents of the oil were trans-piperitone oxide (64.51 %) and piperitenone oxide (12.34 %; Baser et al. 1997a). The seeds contain a variety of different fatty acids (Gusakova et al. 1976). 3951



▲ **Melilotus officinalis (L.) Pall.** Photos: *left and right:* Evgeny Davkaev; *center:* Rostislav Lezhoyev

▼ **Melo agrestis (Naudin) Pang.** Photos: Sergey Zelentsov



▼ **Melissa officinalis L.**
Photos: *top:* Andrei Lubchenko;
bottom: Sasha Eisenman



▼ **Mentha asiatica Boriss.**
Photos: *left:* Alim Gaziev;
top right and bottom right:
Evgeny Davkaev



Morus alba L. – Moraceae	3968
Synonyms: None	3969
English name: White mulberry	3970
Russian name: Шелковица белая (Shelkovitsa belaya)	3971
Uzbek name: Oq toot	3972
Kyrgyz name: Ак тыт (Ak tyt)	3973
Description: Monoecious or dioecious tree, up to 15(–18) m tall. Bark light-brown with shallow furrows; branches gray or gray-brown, young branches pubescent. Leaves alternate, ovate or rarely oblong-oval, 8–14 cm long, 5–10 cm wide, often lobed with 2–5 lobes, base rounded or slightly cordate, sometimes uneven, apex acute, margins entire or serrate, long-petiolate. Inflorescences catkins; male catkins cylindrical; female catkins oviform, densely-flowered. Flowers unisexual, sessile, glabrous. Fruits small, drupelet-like, arranged a syncarp; syncarp 0.5–2.5 cm long, white, pink, or red.	3974 3975 3976 3977 3978
Other distinguishing features: The leaves are often glossy and the sap is milky. The stigmas are covered with papillae.	3979
Phenology: Flowers in April, fruits in May-June.	3980
Reproduction: By seeds, cuttings, and grafts.	3981
Distribution: Cultivated throughout all of Uzbekistan, especially in the plains and lower mountain zones, often becoming naturalized; in agricultural zones of all provinces of Kyrgyzstan.	3982 3983
Habitat: The chul, adyr, and tau zones. Cultivated lands near canals and backyards.	3984
Population status: Common.	3985
Traditional use: In the folk medicine of Central Asia, mulberry leaves are used to treat angina. Fresh leaf juice is used to treat toothaches, and fruits and fruit juice are used to treat oral and throat bumps, dysentery, anemia, and as a diuretic and hemostatic for uterine bleeding, rashes, and scarlet fever. Fresh fruits are used to treat ulcers and the duodenum, and as a blood purifier, as well as an antipyretic and diuretic to improve heart function for cases of myodystrophy (Khalmatov 1964; Gammerman et al. 1990).	3986 3987 3988 3989 3990
Documented effects: Resins from the leaves decrease blood pressure. An infusion of the leaves was shown to slightly reduce blood sugar levels (Gammerman et al. 1990). The flavonoid leachianone G was isolated from the root bark and showed potent antiviral activity against herpes simplex type 1 virus (Du et al. 2003). Two flavonoids isolated from the leaves significantly inhibited the growth of a human leukemia cell line (Kim et al. 2000). Flavonol glycosides, isolated from an extract of the leaves, showed some inhibition of low-density lipoprotein (LDL) oxidation (Katsube et al. 2004, 2006).	3991 3992 3993 3994 3995
Phytochemistry: The leaves contained tannins (3.2–3.7 %), flavonoids (up to 1 %), coumarins, organic acids, resins, and small amounts of essential oils (0.03–0.04 %). Rutin, hyperoside, and quercetin were isolated from the total flavonoids and ostchol was isolated from the coumarins. The fruits contained up to 12 % sugars (occasionally up to 23 %), flavonoids, carotene, pectin, organic acids, small amounts of vitamin C, and tannins (Gammerman et al. 1990).	3996 3997 3998 3999

4001 ***Nepeta pannonica* L. – Lamiaceae**

4002 **Synonyms:** *Nepeta nuda* L.

4003 **English name:** Unknown

4004 **Russian name:** Котовник венгерский (Kotovnik vengerskiy)

4005 **Uzbek name:** Zofo

4006 **Kyrgyz name:** Венгер непетасы (Venger nepetasy)

4007 **Description:** Herbaceous perennial. Stems erect, up to 120 cm tall. Leaves 3.5–6.5 cm long, 1.5–2.5 cm wide, oblong-ovate
4008 to lanceolate, above green, nearly glabrous, pale beneath, pubescent, margin crenate or serrate. Inflorescences terminal
4009 paniculiform cymes, bracts narrow-linear. Calyx tubular, pubescent. Corolla pale-violet, pink or white, 2-lipped, upper lip
4010 2-lobed, lower lip 3-lobed. Fruits oblong, brown nutlets.

4011 **Other distinguishing features:** Plants branching above middle of stem, inflorescences long and narrow.

4012 **Phenology:** Flowers in June-August, fruits in August-September.

4013 **Reproduction:** By seeds.

4014 **Distribution:** All provinces of Kyrgyzstan; Toshkent, Farg'ona and Surxondaryo provinces of Uzbekistan.

4015 **Habitat:** In tallgrass meadows, meadow-steppes and steppes, and forest belts.

4016 **Population status:** Common, found growing as single plants.

4017 **Traditional use:** An infusion of the herb is used to treat asthenia and syphilis. The essential oil is used in perfumery (Plant
4018 Resources of the USSR 1991).

4019 **Documented effects:** The entire plant shows antibacterial activity (Plant Resources of the USSR 1991).

4020 **Phytochemistry:** Aboveground plant parts contains essential oil with 60 components with cineole and nepetalactone as the
4021 major constituents (Kobaisy et al. 2005), iridoids, steroidal saponins, flavonoids, alkaloids, and tannins. Seeds contain
4022 fatty oil, steroids, sterols, and sterol esters (Stepanenko et al. 1980; Plant Resources of the USSR 1991).

4023

- Nigella sativa* L. – Ranunculaceae** 4024
- Synonyms:** *Nigella indica* Roxb. ex Flem., *Nigella truncata* Viv. 4025
- English name:** Black cumin, fennel-flower, love-in-a mist 4026
- Russian name:** Чернушка посевная (Chernushka posevnaya) 4027
- Uzbek name:** Sedana 4028
- Kyrgyz name:** Сейдана ундоосу (Seydana undoosu) 4029
- Description:** Herbaceous annual. Stem 20–75 cm high, striated, oval, simple, slightly glandular-hairy. Leaves 1.5–3 cm long, bi- or tripinnatisect with linear, acute lobules; lower leaves petiolate, early-senescing; upper leaves sessile, similar to lower leaves. Flowers solitary, terminal or in leaf axes, 10–15 mm long, 15 mm wide, short-pubescent. Sepals 5, 1–1.5 cm long, petaloid with a short stalk, blue. Petals developed into 2-lipped nectaries. Stamens many. Fruits composed of 5 inflated follicles, ~1.5 cm long, connate nearly to apices, with erect, ribbed beaks. Seeds triquetrous, wrinkly-tuberculate, light brown. 4030
4031
4032
4033
4034
4035
- Other distinguishing features:** Follicles granular-tuberculate. Seeds have a specific bitter taste. 4036
- Phenology:** Flowers and fruits in May-June. 4037
- Reproduction:** Only by seeds. 4038
- Distribution:** Toshkent and Samarqand provinces of Uzbekistan; all of Kyrgyzstan. 4039
- Habitat:** The chul and adyr zones. Among crops in cultivated areas. 4040
- Population status:** Rare. 4041
- Traditional use:** Avicenna used this plant to treat headaches, facial paralysis, and eye cataracts, and when mixed together with honey in hot water to remove bladder and kidney stones. An infusion of the seeds is used to treat toothaches, gastric and intestinal diseases and chest pains, and is used as a diuretic, soporific, and vermifuge for children (seeds in vinegar), as well as to treat angina and stimulate milk production in women (Karimov and Shomakhmudov 1993). 4042
4043
4044
4045
- Documented effects:** An infusion of the seeds had positive inotropic and negative chronotropic action, and reduced heart function due to increased cardiac output (Ogolevitz 1951). In a variety of experiments, extracts of the plant exhibited antibacterial activity, and in animals, increased bile and uric acid secretion, protected against histamine induced broncho-spasm, shortened bleeding time, and inhibited fibrinolytic activity. Volatile oil from the seeds caused a dose-dependent increase in respiratory rate and intracranial pressure in anesthetized guinea pigs and reduced heart rate and blood pressure in anesthetized rats. Ingestion of the seeds caused reduction in cholesterol and blood glucose levels in humans. The seeds were also found to enhance immunity and had anti-cancer activity against malignant cells in mice and in humans. An aqueous extract of the seeds had anti-inflammatory and analgesic activity (Al-Ghamdi 2001). 4046
4047
4048
4049
4050
4051
4052
4053
- Phytochemistry:** Seeds contain 0.4–1.5 % essential oil (with a pleasant aroma), up to 40 % fatty oil, vitamin C, flavonoids (quercetin and camphorol), steroid alkaloids, coumarins, quinones, saponins, mineral salts, etc. (Karimov and Shomakhmudov 1993; Ali and Blunden 2003). The major components of the essential oil were thymoquinone, ρ -cymene, carvacrol, trans-anethole, 4-terpineol and longifolin (Ali and Blunden 2003). 4054
4055
4056
4057

4059 **Onopordum acanthium L. – Asteraceae**4060 **Synonyms:** *Acanos spina* Scop.4061 **English name:** Scotch thistle4062 **Russian name:** Татарник обыкновенный (Tatarnik obyknovennyy)4063 **Uzbek name:** Okkarrak4064 **Kyrgyz name:** Кадимки коко тикен (Kadimki koko tiken)

4065 **Description:** Herbaceous biennial, white-gray tomentose. Stems erect, 35–300 cm tall, spiny-winged. Leaves sinuate-lobed
4066 or toothed, teeth and lobes tipped with sharp spines; basal leaves up to 30 cm long, petiole winged; cauline leaves sessile.
4067 Inflorescences ovoid-spherical capitula, single or in corymbiform groups; involucre bracts linear, arranged in many rows,
4068 ending in sharp spines. Disc flowers purple, many; ray flowers absent. Fruits achenes, elongate-obovate, dark-gray with
4069 brown spots, pappi brownish.

4070 **Other distinguishing features:** Leaves and stems white-gray tomentose. Involucre bracts are linear.4071 **Phenology:** Flowers in June-July, fruits in July-August.4072 **Reproduction:** By seeds.

4073 **Distribution:** All provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Tashkent, Samarqand, Jizzax, Sirdaryo and
4074 Surxondaryo provinces of Uzbekistan.

4075 **Habitat:** As weed in waste grounds, fallow fields, and pastures.4076 **Population status:** Common, often forming dense groups.

4077 **Traditional use:** The inflorescences, roots, seeds, and late developing leaves (with spines removed), are harvested for use in
4078 folk medicine. They are used internally to treat inflammation of the bladder and urinary system, bronchial asthma, pertus-
4079 sis, scrofula, hypostasis of various origins, common colds, hemorrhoids, as a blood cleanser, and for treating skin dis-
4080 eases. The plant is used externally in the form of compresses, lotions, and fresh juice, which is especially effective, to treat
4081 skin diseases, purulent wounds, ulcers, and furuncles. An infusion of the top of the stem collected during flowering is
4082 drunk to treat nervous breakdowns, common colds, and inflammation of the respiratory system, and is put in baths for
4083 frightened children (Turova and Sapozhnikova 1984; Maznev 2004).

4084 **Documented effects:** Experiments have shown that preparations of this species has very low toxicity and even after long
4085 periods of use show no side effects. It possesses cardiogenic, hemostatic, styptic, diuretic, and bacteriocidal properties and
4086 raises arterial pressure and causes narrowing of the blood vessels. In small doses preparations of this plants work as a
4087 tonic, and in larger doses, depress the central nervous system (Khalmatov et al. 1984). In some countries the herb is used
4088 to treat skin cancer and as a prophylactic after removal of a tumor (Akopov 1990). An aqueous extract of the plant exhib-
4089 ited anti-tumor activity in vitro (AbuHarfeil et al. 2000).

4090 **Phytochemistry:** Leaves contain alkaloids, glycosides, bitter substances, sesquiterpene lactones (arctiopicrin and onopor-
4091opicrin), vitamin C and K₁, resins, titratable acids, sugars, tannins, terpenoids (taraxasteryl acetate), etc. Seeds contain
4092 alkaloids, acetates of lupeol and amyirin, and drying fatty oil (Khalmatov et al. 1984; Ul'chenko et al. 1993; Khalilova
4093 et al. 2004).

4094



▲ *Nepeta pannonica* L. Photos: Andrei Lubchenko



▲ *Morus alba* L. Photo: Bruce Hamilton



▲ *Nigella sativa* L. Photo: Pamela J. Eisenberg

▼ *Onopordum acanthium* L. Photos: *lower left and center:* Rostislav Lezhoyev; *top left and right:* Sasha Eisenman



4095 **Origanum tyttanthum Gontsch. – Lamiaceae**

4096 **Synonyms:** *Origanum vulgare* var. *genuinum* O. Fedtsch., *Origanum vulgare* var. *prismaticum* Gaudin, *Origanum vulgare*
4097 var. *viride* (Boiss.) Hayek.

4098 **English name:** Кыргыз oregano

4099 **Russian name:** Душица мелкоцветная (Dushitsa melkotsvetnaya)

4100 **Uzbek name:** Тоғ райхон, жамбил

4101 **Kyrgyz name:** Майда гулдуу кок чай чоп (Mayda gulduu kok chay chop)

4102 **Description:** Herbaceous perennial, rhizomatous. Stems many, 20–90 cm tall, erect, branched, square, villous. Leaves oppo-
4103 site, oval or oblong, 0.5–3 cm long, adaxial side nearly glabrous, abaxial side villous along veins, covered with punctuate
4104 glands, short-petiolate. Inflorescence a complex panicle, 10–30 cm long. Flowers nearly sessile. Calyx campanulate,
4105 3 mm long, short pubescent. Corolla pale-pink, 5 mm long. Fruits nutlets, deep brown, less than 1 mm long.

4106 **Other distinguishing features:** It differs from *Origanum vulgare* by having a more narrow inflorescence and smaller
4107 flowers.

4108 **Phenology:** Flowers in June-August, fruits in July-September.

4109 **Reproduction:** By seeds and rhizomes.

4110 **Distribution:** Toshkent, Samarqand, Andijon, Farg'ona, and Surxondaryo provinces of Uzbekistan; Osh, Jalal-Abad, and
4111 Batken provinces of Kyrgyzstan.

4112 **Habitat:** The adyr and tau zones. Grows on rocky and pebbly slopes.

4113 **Population status:** Common, often makes large populations.

4114 **Traditional use:** A decoction of the herb is used in folk medicine to stimulate the appetite and to improve digestion, to treat
4115 inflammation of mucous membranes in the upper respiratory tract, and decrease nervous excitability. Infusions and decoc-
4116 tions are applied externally as compresses for abscesses, and are also used in a bath to treat children who have rickets or
4117 scrofula. Water extractions of the aboveground plant parts are used to treat acute and chronic gastritis, bronchitis, chole-
4118 cystitis, pneumonia, and urolithiasis and are also used as a cholagogue. A tea is used to treat tympanites, laryngitis, stoma-
4119 titis, and angina, and as an oral and throat rinse (Khalmatov et al. 1984; Khodzhimatov 1989).

4120 **Documented effects:** It is an effective remedy to treat hypertension, atherosclerosis, kidney, liver, and epilepsy (Kovaleva
4121 1971). It is a sedative for excitement of the central nervous system (Turova 1974). A decoction of the dried leaves and
4122 flowers is used to treat intestinal atonia and as an expectorant. The plant is a component of a diaphoretic tea and is added
4123 to baths. The leaves are used as a spice and in liquor production (Tsitsina 1962). The essential oil has shown antimicro-
4124 bial, hypocholesteremic, and hypolipidemic activity (Nuraliev and Zubaidova 1994; Takeda et al. 2008).

4125 **Phytochemistry:** The flowering plant contain 0.17–0.6 % essential oil, which includes 35–66 % phenols (mostly thymol and
4126 carvacrol). The seeds contain up to 25 % fatty oils (Khalmatov 1964; Khalmatov et al. 1984). The plant contains phenolic
4127 glycosides, lipids and coumarins (Takeda et al. 2008).

- Origanum vulgare L. – Lamiaceae** 4129
- Synonyms:** *Micromeria formosana* C. Marquand, *Origanum creticum* Lour., *Origanum dilatatum* Klokov, *Origanum normale* D. Don, *Origanum puberulum* (G. Beck) Klokov. 4130
- English name:** Oregano, wild marjoram, Greek oregano 4131
- Russian name:** Душица обыкновенная (Dushitsa obyknovennaya) 4132
- Uzbek name:** Jambil 4133
- Kyrgyz name:** Кадемки кок чай чоп (Kademki kok chay chop) 4134
- Description:** Herbaceous perennial, with rhizomes. Stems purplish, erect or prostrate, pubescent, 20–60 cm tall. Leaves 4135
opposite, petiolate, broadly ovate to oblong, 1–4 cm long and 0.5–1.5 cm wide, densely hairy below. Inflorescences spikes 4136
in corymbiform or paniculiform clusters; bracts ovate-elliptic, green or purple. Calyx with triangular-lanceolate teeth, 4137
dark-purple. Corolla 5–7 mm long, light-purple, lilac-pink or white, 2-lipped; upper lip erect, apex 2-lobed; lower lip 4138
3-lobed. Fruits nutlets, orbicular, bluntly 3-sided, brown. 4139
- Other distinguishing features:** Inflorescence wider, and flowers bigger, than the closely related species, *Oreganum tyttan-* 4140
thum Gontsch. 4141
- Phenology:** Flowers in June-July, fruits in August-September. 4142
- Reproduction:** By seeds. 4143
- Distribution:** Ysyk-Kol and Chuy Provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 4144
- Habitat:** On northern slopes in tallgrass-meadow belts and forests, among bushes, and along forest edges. 4145
- Population status:** Common, found in dense groups. 4146
- Traditional use:** An infusion is used to treat stomach ailments, common colds, and gynecological problems. It is used exter- 4147
nally as a lotion, compress, and in baths for the treatment of eczema, infected skin diseases, and to wash wounds 4148
(Gammerman et al. 1990). 4149
- Documented effects:** An infusion of the herb is used to treat insomnia, hypo- and anacidic gastritis, and atonia of the intes- 4150
tines. It is also used as an expectorant for bronchitis and bronchiectasis, as well as to increase appetite (Turova and 4151
Sapozhnikova 1984). An extract of the herb is used as a component in the preparation *Urolesan*, which is used as an anti- 4152
spasmodic, antiseptic, and anti-inflammatory for the urinary tract, as well as to help eliminate ureter stones, and to 4153
increase bile production. The preparation increases diuresis and improves blood circulation through the liver (Gammerman 4154
et al. 1990). The essential oil of *Origanum vulgare* ssp. *vulgare* exhibited significant antimicrobial activity against 10 4155
species of bacteria and 15 fungal species (Sahin et al. 2004). Essential oils from *Origanum vulgare* ssp. *hirtum* exhibited 4156
high levels of antimicrobial activity against 8 strains of gram-positive and gram-negative bacteria. The essential oil also 4157
exhibited high levels of cytotoxicity against 4 permanent animal cell lines including 2 derived from human cancers 4158
(Sivropoulou et al. 1996). Aqueous and methanolic extracts of oregano have been shown to have effective antioxidant 4159
properties (Cervato et al. 2000). 4160
- Phytochemistry:** The herb contains essential oil (with phenols such as thymol and isomers of carvacrol), bi and tricyclic 4161
sesquiterpenes, free alcohols, tannins, ascorbic acids, and flavonoids (Akopov 1990; Sivropoulou et al. 1996). A total 62 4162
constituents were identified from the essential oil of *Origanum vulgare* ssp. *vulgare*, with the main constituents being 4163
caryophyllene, spathulenol, germacrene-D, and terpineol (Sahin et al. 2004). 4164

4167 **Orthurus kokanicus (Regel & Schmalh.) Juz. – Rosaceae**4168 **Synonyms:** *Geum kokanicum* Regel & Schmalh.4169 **English name:** Unknown4170 **Russian name:** Прямохвостник кокандский (Pryamokhlostnik kokandskiy)4171 **Uzbek name:** Yerchoy, Shirchoy4172 **Kyrgyz name:** Unknown

4173 **Description:** Herbaceous perennial. Stems erect, 15–50 cm high, hairy. Basal leaves in dense rosette, lyrate-pinnatisect, lateral segments many, large and small, usually in opposite or sub-opposite pairs, ovate-rhomboid, single- or bi-dentate, terminal segment larger, slightly lobed, bidentate; cauline leaves few, small, short-petiolate, oval to nearly round, trilobed, large-dentate. Inflorescences cyme-like with 2–7 flowers, crowded, but becoming loose with age, stiff-hairy. Hypanthium widely campanulate. Outer and inner sepals 5. Petals 5, yellow, about as long as sepals. Style erect with 2 parts; upper glabrous and deciduous; lower part persistent, longer than fruitlet, covered with retrorse bristles. Fruitlets 4–10, stiff-hairy.

4180 **Other distinguishing features:** When fractured, the roots produce pleasant a eugenol smell.4181 Sepals wider and gynophore shorter than *Orthurus heterocarpus*.4182 **Phenology:** Flowers in June–July, fruits in July–August.4183 **Reproduction:** By seeds.4184 **Distribution:** Toshkent, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; not found in Kyrgyzstan.4185 **Habitat:** The tau zone. Stony mountain slopes in juniper stands.4186 **Population status:** Uncommon.4187 **Traditional use:** A decoction of the roots is used in folk medicine internally for chest pains and as an astringent. A decoction of roots and leaves is used to rinse the mouth and throat. The roots are also used a tea substitute (Khalmatov 1964).

4188 **Documented effects:** The essential oil, which is rich with eugenol, is used in dentistry instead of imported clove oil. A decoction or infusion of the roots are recommended as an astringent for gastrointestinal diseases (Khalmatov 1964). Essential oil, isolated from the plant, had strong antibiotic effects against *Shigella dysenteriae*, *Bacillus subtilis*, and *Aspergillus flavus* (Faramarzi et al. 2008).

4193 **Phytochemistry:** Roots contain 22–25 % tannins, 10–13 % sugars, essential oils (up to 0.45 % eugenol), resins, and organic acids (Khalmatov 1964). The major compounds in the essential oil, distilled from underground parts, were eugenol (80.9 %) and myrtenol (5.2 %) (Faramarzi et al. 2008).

Padus avium Mill. – Rosaceae	4196
Synonyms: <i>Padus racemosa</i> (Lam.) Gilib., <i>Prunus padus</i> L., <i>Prunus racemosa</i> Lam.	4197
English name: Bird cherry	4198
Russian name: Черёмуха обыкновенная (Cheryomukha obyknovennaya)	4199
Uzbek name: Unknown	4200
Kyrgyz name: Кадемки моюл (Kademki moyul)	4201
Description: Deciduous tree, 2–10 m tall. Bark light tan to black-gray, cracked; young branches brown with white-yellow lenticels. Leaves simple, alternate, short-petiolate, glabrous, elliptic, margins serrate. Inflorescences hanging racemes, 8–12 cm long. Hypanthium cup-shaped, glabrous outside, hairy inside, with 5 short sepals. Petals 5, obovate, white. Fruits black, spherical drupes, 8–10 mm in diameter.	4202 4203 4204 4205
Other distinguishing features: Differs from <i>Padus asiatica</i> Kom. by having glabrous young branches, shorter racemes, and smaller corollas.	4206 4207
Phenology: Flowers in May to the beginning of June, fruits in July-August.	4208
Reproduction: By seeds and rhizomes.	4209
Distribution: Osh and Chuy provinces of Kyrgyzstan; cultivated in Uzbekistan.	4210
Habitat: On floodplains.	4211
Population status: Rare, found in small groups.	4212
Traditional use: Bark, leaves, flowers, and mature fruits are used in folk medicine. Fruits are used as an astringent remedy to treat diarrhea of non-infectious origins and other intestinal disorders, as well as a secondary treatment for infectious colitis and diarrhea (Kurochkin 1998).	4213 4214 4215
Documented effects: Mature fruits are used as a bactericide, anti-inflammatory, to normalize intestine and stomach function, as a source of vitamins, and as a tonic. The bark is used as a diaphoretic, antipyretic, and diuretic. Leaves are used to treat diarrhea and as a source of vitamins. Flowers are used as an anti-inflammatory. Preparations from this species are counter-indicated during pregnancy (Maznev 2004). An extract of the seeds had antibacterial activity against 5 different species, including methicillin-resistant <i>Staphylococcus aureus</i> , and had strong anti-oxidant activity (Kumarasamy et al. 2002, 2007).	4216 4217 4218 4219 4220 4221
Phytochemistry: Leaves, flowers, bark and seeds contain glycosides (amygdalin, prulaurasin, and prunasin). Prussic acid is found in the bark and leaves. Fruits contain malic and citric acids, sugar, astringent substances, ascorbic acid, and flavonoids (Maznev 2004; Deineka et al. 2004).	4222 4223 4224



▲ **Origanum vulgare L.**
Photo: Sasha Eisenman

◀ **Origanum tyttanthum Gontsch.**
Photos: *top left*: Evgeny Davkaev;
top right: John B. Taft; *bottom*:
Alim Gaziev



▼ **Padus avium Mill.** Photos: Sergey Appolonov



- Papaver pavoninum Schrenk – Papaveraceae** 4226
- Synonyms:** *Papaver ocellatum* Woronow. 4227
- English name:** Peacock poppy 4228
- Russian name:** Мак павлиний (Мак pavliniy) 4229
- Uzbek name:** Lola qizg'aldak 4230
- Kyrgyz name:** Кызгалдак апиийми (Kyzgaldak apiyimi) 4231
- Description:** Herbaceous annual. Stem simple or branched from the base, 10–50 cm high, densely coated with bristles. 4232
 Leaves multiple, bipinnatisect; segments oval-oblong, sessile, with bristles on adaxial side. Flowers often in groups of 3; 4233
 buds rounded or oval, 8–15 mm long. Calyx coated with long whitish or reddish bristles, with 2 long, hollow, prominent 4234
 apical horns. Petals ca. 2.5 cm long and 4 cm wide, bright red with an arching black spot at the base. Fruit an ovoid cap- 4235
 sule, roundish, 5–10 mm long, 8 mm wide, ribbed, coated with stiff bristles. Seeds 1 mm long, light brown-gray, 4236
 oblong. 4237
- Other distinguishing features:** Cauline leaves often have an elongated terminal lobe. 4238
- Phenology:** Flowers and fruits in March-June. 4239
- Reproduction:** By seeds. 4240
- Distribution:** Toshkent, Buxoro, Samarqand, Andijon, Farg'ona, and Surxondaryo provinces of Uzbekistan; Chuy, Ysyk- 4241
 Kol, Talas, Osh, Jalal-Abad and Batken provinces of Kyrgyzstan. 4242
- Habitat:** The chul and adyr zones. Clay deserts, on loess or stony slopes and in unirrigated winter wheat fields. 4243
- Population status:** Common. 4244
- Traditional use:** Juice from the petals is used as a drink to treat heatstroke (in children) and as a rinse for eye ailments. The 4245
 dried petals of related species, *Papaver rhoeas*, *Papaver orientale*, and *Papaver bracteatum*, are used to prepare a tea to 4246
 treat coughs (Khalmatov 1964; Seredin and Sokolov 1969). 4247
- Documented effects:** Protopine has been shown to strongly inhibit induced platelet aggregation in vitro. In vivo, pretreat- 4248
 ment with protopine protected rabbits from the lethal effects of specific platelet aggregation agonists. Protopine also 4249
 inhibited carrageenan-induced rat paw edema and had 3 times the potency of aspirin (Saeed et al. 1997). 4250
- Phytochemistry:** The herb contains 0.1 % total alkaloids, from which a-allocryptopine, protopine, and roemeridine have 4251
 been isolated (Khalmatov 1964). In another study less than 0.05 % total alkaloids were found in the plant and a β -carbo- 4252
 line was the dominant alkaloid (Taborska et al. 1988). 4253

4255 ***Patrinia intermedia* (Hornem.) Roem. & Schult. – Valerianaceae**

4256 **Synonyms:** *Fedia intermedia* Hornem., *Fedia rupestris* var. *intermedia* (Hornem.) Vahl, *Patrinia nudiuscula* Fisch.

4257 **English name:** Unknown

4258 **Russian name:** Патриния средняя (Patriniya srednyaya)

4259 **Uzbek name:** Unknown

4260 **Kyrgyz name:** Орто патриния (Orto patriniya)

4261 **Description:** Herbaceous perennial, with taproot that branches towards the top. Stems single or few, 20–75 cm tall, up to
4262 3 cm in diameter, short-haired. Leaves opposite, 4–18 cm long, 2–5 cm wide, gray-green, glabrous; basal leaves long-
4263 petiolate, elongate-oblong, strongly dentate, pinnatilobate or pinnatisect; lower stem leaves sessile, in 2–5 pairs, pinnati-
4264 sect; upper stem leaves lanceolate-ovate, 3-nerved. Inflorescence corymbiform-paniculate. Corolla bright yellow,
4265 campanulate, 5-lobed. Fruits slightly hairy achenes.

4266 **Other distinguishing features:** Flowers have 4 stamens.

4267 **Phenology:** Flowers in June-July, fruits in July-September.

4268 **Reproduction:** By seeds.

4269 **Distribution:** All provinces of Kyrgyzstan; not found in the flora of Uzbekistan.

4270 **Habitat:** On stony floodplains of mountain rivers, on stony slopes and in steppe and forest-meadow belts of mountains.

4271 **Population status:** Common, found in loosely arranged groups.

4272 **Traditional use:** An infusion or decoction of the roots is used like valerian to treat nervous excitement and cardiac neurosis
4273 (Turova and Sapozhnikova 1984).

4274 **Documented effects:** The biological activity is due to the presence of saponins, and removal of the saponins from the tinc-
4275 ture leads to complete loss of the pharmacological properties (Ivanova 1963). The sedative effect of this species is nearly
4276 twice as strong as that of *Valeriana* (Tolmachev 1976). The roots of this species reduce excitability of the nervous system.
4277 Clinical tests showed that application of an alcohol infusion stopped or noticeably reduced chest pain as well as nervous
4278 and cardiovascular excitation caused by hypodermic introduction of caffeine (Akopov 1990).

4279 **Phytochemistry:** Roots contain triterpene saponins (patrinoside A, B, C and interoside B), inulin, organic acids, tannins, and
4280 essential oils. The seeds contain alkaloids (Khodzhimatov 1989).

4281

- Peganum harmala L. – Zygophyllaceae** 4282
- Synonyms:** None 4283
- English name:** Syrian rue 4284
- Russian name:** Гармала обыкновенная (Garmala obyknovennaya) 4285
- Uzbek name:** Isiriq 4286
- Kyrgyz name:** Адыршаман (Adyrshaman) 4287
- Description:** Herbaceous perennial, with thick woody taproot. Stems few, 20–80 cm tall, heavily branched, glabrous, slightly 4288
grooved. Leaves 3–8 cm long, sessile to short-petiolate, irregularly pinnatisect with linear-lanceolate segments, stipulate. 4289
Flowers in groups of 1–3, pedicellate, terminal on branches. Calyx deeply divided into 5 linear lobes, 1.5–2 cm long. 4290
Petals 5, white or pale yellow, elliptic, 1.5–2 cm long. Fruit a globular capsule, ca. 1 cm in diameter, 3-valved, splitting 4291
when ripe. Seeds many, triquetrous, dark brown. 4292
- Other distinguishing features:** Calyx persistent in fruit. Dry leaves and plants with fruits have a specific smell when 4293
burned. 4294
- Phenology:** Flowers and fruits in May-September. 4295
- Reproduction:** By seeds. 4296
- Distribution:** All regions of Uzbekistan and Kyrgyzstan. 4297
- Habitat:** The chul and adyr zones. Waste places, abandoned fields, around the periphery of wells and in villages. In clayey 4298
and sandy soils, rich with nitrates. 4299
- Population status:** Common, found in small populations. 4300
- Traditional use:** This is a well-known herb to all Central Asian people. Avicenna used the plant as an analgesic for patients 4301
with sciatic nerve inflammation. In folk medicine the herb is used in baths to treat rheumatism, scabies, and other skin 4302
diseases. A decoction or infusion of the plant is drunk to treat common colds, malaria, fever, syphilis, neurasthenia, and 4303
epilepsy, and is also used as a mouth wash to treat gum disease. The smoke of the burning herb is good for headaches; for 4304
epileptic diseases the patient's room is filled with the smoke. A decoction of the seeds mixed with flax seeds is recom- 4305
mended for asthma and breathlessness, it is mixed with chili pepper to treat syphilis, and it is used as a diuretic and dia- 4306
phoretic (Khalmatov et al. 1984; Gammerman et al. 1990). In Tajikistan smoke from the plant is used to treat paralytics. 4307
The leaves are used as a poultice to treat swelling (Khodzhimatov 1989). 4308
- Documented effects:** An infusion of the roots and a preparation, *Salyanokisli garmine*, are used for Parkinson's disease after 4309
lethargic encephalitis (von Economo's disease), for epilepsy, and as a soporific. The alkaloid harmine is a reversible 4310
inhibitor of monoamine oxidase (Gorkin 1964; Coates and Cox 1972). The strong impact of harmine on the central ner- 4311
vous system often causes major mental disorders. Due to these effects, it is classified as a psychomimetic substance of 4312
adrenergic action (Sadritdinov and Kurmukov 1980). The alkaloid peganine depresses acetyl-cholinesterase as well as 4313
butyryl-cholinesterase (Sharapov 1959). In acute tests on cats and chronic tests on dogs the alkaloid increases bile flow 4314
up to 40–100 %, at the dose of 5 mg/kg. At the same time, secretion of bilirubin also increases (Rabinovich et al. 1966). 4315
Deoxypeganine exhibited strong anticholinesterase activity in vitro and in vivo (Tulyaganov et al. 1986). In in vitro tests, 4316
the alkaloids peganol and peganidine have inhibitory action on activity of acetyl- and butyryl-cholinesterase of the blood 4317
and brain (Rustamov et al. 1974). 4318
- Phytochemistry:** The aboveground parts of plants collected during the early the vegetative stage in the Buxoro and 4319
Surxondaryo provinces contained 2.1 % total alkaloids, with the young roots containing 3.32 % and older roots 1.68 % 4320
total alkaloids. At the bud stage, the aboveground parts contained 2–2.3 % total alkaloids, which decreased at the stage of 4321
flowering to 1.86–1.95 %. More than 15 alkaloids were isolated from the total alkaloids including harmine, harmaline 4322
harmalol, peganine, vasicinone, deoxypeganine, peganine, peganidine, peganol, and dipegene, etc. (Yunusov 1981). 4323

4325 **Perovskia abrotanoides Kar. – Lamiaceae**

4326 **Synonyms:** *Perovskia artemisioides* Boiss.

4327 **English name:** Russian sage, Caspian Russian sage, Caspian Perovskia

4328 **Russian name:** Перовския полынная (Perovskiya polynnaya)

4329 **Uzbek name:** Khapri, Abrik

4330 **Kyrgyz name:** Шыбактай коен томук (Shybaktay koyen tomuk)

4331 **Description:** Perennial subshrub. Stems up to 100 cm tall, bases woody, white-hairy. Leaves petiolate, oblong-ovate, 2–7 cm
4332 long, 1–3 cm wide, bi-pinnatipartite. Inflorescences verticillasters, found in loose panicles; bracts lanceolate-linear. Calyx
4333 around 4.5 mm long, tubular-campanulate, 2-lipped, violet, often densely hairy. Corolla violet, funnelform, 2-lipped;
4334 upper lip 4-lobed, middle 2 lobes smaller; lower lip entire. Fruits smooth, brown nutlets.

4335 **Other distinguishing features:** Differs from *Perovskia atriplicifolia* Benthams by having bi-pinnatipartite leaves.

4336 **Phenology:** Flowers in June–August, fruits in August.

4337 **Reproduction:** By seeds and rhizomes.

4338 **Distribution:** Ysyk-Kol, Naryn, Osh, and Chuy Provinces of Kyrgyzstan; not found in the flora of Uzbekistan.

4339 **Habitat:** In pebbly, dry stream beds and on dry, stony places in the mountains.

4340 **Population status:** Common.

4341 **Traditional use:** The aboveground parts are used to heal wounds. A decoction is used to treat scabies (Massagetov 1932).

4342 The plant is used externally to treat leishmaniasis in Iran (Moallem and Niapour 2008).

4343 **Documented effects:** Compounds isolated from the roots exhibited leishmanicidal activity in vitro and inhibited growth of
4344 cultured malaria parasites, human lymphocytes, and human carcinoma cell lines (Sairafianpour et al. 2001). Compounds
4345 isolated from the aboveground parts exhibited cytotoxic activity against leukemia cells (Aoyagi et al. 2006).

4346 **Phytochemistry:** The plant contains tanshinones (Sairafianpour et al. 2001). Water-distilled essential oils from leaves col-
4347 lected in Arslonbob (Kyrgyzstan) contained cineole, pinene, epi-13-manool, bornyl acetate, camphene, camphor, caryo-
4348 phyllene, caryophyllene oxide, humulene, caryophylladienol, borneol, and other compounds (Basher et al. 1997).

4349



▲ **Papaver pavoninum** Schrenk Photos: *left and right:* Alim Gaziev; *center:* Alexander Naumenko



▲ **Patrinia intermedia** (Hornem.) Roem. & Schult. Photos: Lina Valdshmit



► **Peganum harmala** L. Photos: *top:* John B. Taft; *center:* Alim Gaziev; *bottom:* Evgeny Davkaev

▼ **Perovskia abrotanoides** Kar.

Photos: *left:* John B. Taft; *right:* Bazar Dovletov



4350 ***Picea schrenkiana* Fisch. & C.A. Mey – Pinaceae**

4351 **Synonyms:** *Abies schrenkiana* (Fisch. & C.A. Mey) Lindl. & Gordon, *Picea morinda* ssp. *tianschanica* (Rupr.) Berezin,
4352 *Picea obovata* Ledeb. var. *schrenkiana* (Fisch. & C.A. Mey) Carrière, *Picea prostrata* Isakov, *Picea robertii* P. Vipper, *Picea*
4353 *tianschanica* Rupr.

4354 **English name:** Schrenk's spruce

4355 **Russian name:** Ель Шренка (El' Shrenka)

4356 **Uzbek name:** Неизвестно

4357 **Kyrgyz name:** Archa

4358 **Description:** Evergreen tree, up to 40 m tall, with narrow conical crown. Bark grayish-brown with thick plates. Leaves
4359 (needles) arranged radially, 20–25 mm long, linear, 4-sided, apex acute. Seed (female) cones ellipsoid-cylindric, 6–15 cm
4360 long, up to 3.5 cm wide. Seed scales triangular-ovate, apex rounded, brown. Seeds up to 4 mm long, flat-ovoid to fusiform,
4361 winged, brown.

4362 **Other distinguishing features:** Branchlets pendulous, pale yellow.

4363 **Phenology:** Seeds ripen in September.

4364 **Reproduction:** By seeds.

4365 **Distribution:** Ysyk-Kol, Naryn, Talas, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan.

4366 **Habitat:** On mountain slopes from 1,000 to 3,500 m.

4367 **Population status:** Common, forming forests.

4368 **Traditional use:** An infusion of needles from spring branchlets and cones are drunk to treat persistent common colds and is
4369 added to baths to treat rheumatism. An infusion of young branchlets in vodka is used to treat lung tuberculosis. The
4370 ground bark, mixed with wax and butter or lard, is applied in the form of a plaster to treat furuncles. The needles are used
4371 to prevent and treat scurvy and as a source of vitamins (Bykov 1950; Gan 1970).

4372 **Documented effects:** None.

4373 **Phytochemistry:** Needles and young branches contain vitamin C, essential oil (with up to 40 components such as camphene,
4374 myrcene, bornyl acetate, and others), flavonoids, and microelements (iron, manganese, chromium, aluminium and copper;
4375 Bykov 1950). Thirty-eight diterpenoids have been identified in the oleoresin (Raldugin et al. 1991). Sesquiterpenoids,
4376 diterpenoids, triterpenoids, steroids, and tocopherol were isolated from needles and twigs. Dehydroabietol, patchouli
4377 alcohol, guaial, β -sitosterol, and campesterol were the main components of the unsaponifiable matter (Zhou 2001).

4378

Plantago lanceolata L. – Plantaginaceae	4379
Synonyms: None	4380
English name: Narrowleaf plantain	4381
Russian name: Подорожник ланцетолистный (Podorozhnik lantsetolistnyy)	4382
Uzbek name: Nishtarsimon bargizub, Zabturum	4383
Kyrgyz name: Бака жалбырак (Baka zhalbyrak)	4384
Description: Herbaceous perennial. Leaves in basal rosettes, narrow-elliptic to lanceolate-elliptic, 7.5–35 cm long, 0.5–3.5 cm wide, with 3–5 parallel veins, apex acute, narrow petiolate. Inflorescence a dense, erect, cylindrical spike, 1.5–8 cm tall; peduncles 15–60 cm tall, with 5 ribs; bracts ovate, acute. Corolla 4-lobed. Fruit a 2-seeded, circumscissile capsule. Seeds elongate-oval.	4385 4386 4387 4388
Other distinguishing features: Leaves narrow-elliptic to lanceolate-elliptic. Stamens exerted.	4389
Phenology: Flowers in June-August, fruits in August-September.	4390
Reproduction: By seeds.	4391
Distribution: All provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Toshkent, Samarqand, Farg'ona, Buxoro, Andijon, Namangan, Surxondaryo and Xorazm provinces of Uzbekistan.	4392 4393
Habitat: Along canals and roads and in fallow fields.	4394
Population status: Common, found in small groups.	4395
Traditional use: A decoction of the leaves is used as a diuretic and as a treatment for cystitis, gastric diseases, lung tuberculosis, headaches, and to detoxify snake bites. Decoctions, infusions, extracts, and juice are used as a bacteriostatic, anti-inflammatory, antispasmodic, and expectorant, and also to treat enterocolitis, stomach ulcers, liver diseases, malaria, bronchitis, pertussis, bronchial asthma, allergic conjunctivitis, as well as to heal wounds, furuncles, skin ulcers, and purulent wounds (Plant Resources of the USSR 1990).	4396 4397 4398 4399 4400
Documented effects: Preparation from this species are used as a hemostatic (Zemlinsky 1958) and to treat chronic bronchitis (Nosal and Nosal 1959). Compounds in the herb showed inhibitory effects on mouse ear edema (Murai et al. 1995). Results of experimental research confirmed anti-inflammatory, antispasmodic, and immunostimulatory actions (Wegener and Kraft 1999).	4401 4402 4403 4404
Phytochemistry: Roots contain steroids (sitosterin, stigmasterin, cholesterol, and campesterin) and higher fatty acids. The aboveground parts contain iridoids, phenolcarbonic acids, flavonoids, carbohydrates, organic acids, and protocatechins. Seeds contain iridoids, carbohydrates, muscilage, and fatty oil (Plant Resources of the USSR 1990; Murai et al. 1995).	4405 4406 4407

4409 **Plantago major L. – Plantaginaceae**

4410 **Synonyms:** *Plantago borysthenica* (Rogow.) Wissjul., *Plantago dregeana* Decne., *Plantago latifolia* Salisb., *Plantago*
4411 *officinarum* Crantz.

4412 **English name:** Common plantain, broadleaf plantain

4413 **Russian name:** Подорожник большой (Podorozhnik bol'shoy)

4414 **Uzbek name:** Zupturoom, Buzchi, Bakayaprok

4415 **Kyrgyz name:** Чон бака жалбырак (Chon baka zhalbyrak)

4416 **Description:** Herbaceous perennial. Leaves in basal rosettes, broad-elliptic to broad-ovate, 4–21 cm long, 3–14 cm wide,
4417 3–9 parallel veins, sheathing petiolate. Inflorescences dense, erect, narrow-cylindric spikes, 5–15 cm tall; peduncles
4418 15–70 cm tall; bracts ovate, acute. Corolla greenish or yellowish white with 4 reflexed lobes. Fruit a 2-seeded, circumscis-
4419 sile capsule. Seeds 1–1.5 mm long, densely reticulate.

4420 **Other distinguishing features:** Differs from *Plantago rugelii* Decne. by having fruits dehisce near the middle rather than
4421 far below the middle.

4422 **Phenology:** Flowers in June-August, fruits in August-September.

4423 **Reproduction:** By seeds.

4424 **Distribution:** Almost all provinces of Kyrgyzstan and Uzbekistan.

4425 **Habitat:** In meadows, along streams and canals, and in orchards and wet places.

4426 **Population status:** Common, forming dense groups.

4427 **Traditional use:** This species has been used for centuries. Avicenna used the leaves as a hemostatic, to heal wounds, tumors,
4428 eye inflammation, chronic skin ulcers, and elephantitis as well as for liver and kidney diseases. In more recent times the
4429 plant has been used to treat lung tuberculosis, pertussis, stomach catarrh with low acidity, acute gastritis, enterocolitis,
4430 stomach and duodenum ulcers, and as a hemostatic (Khalmatov et al. 1984). A tea made from the dried leaves is used to
4431 treat coughing, diarrhea, dysentery (with tea from seeds is most effective), inflammation of the bladder, and malaria, and
4432 as an expectorant (Altimishev 1991).

4433 **Documented effects:** Experiments with animals showed that a 20 % extract of leaves healed wounds, decreased pus volume,
4434 stimulated epithelialization of the wound surface, had sedative and soporific effects, and reduced blood pressure (Aliev
4435 1945). The triterpenoid, ursolic acid, and isolated from the plant showed significant COX-2 inhibitory activity (Ringbom
4436 et al. 1998). Five compounds, including caffeic and chlorogenic acids, isolated from extracts of the plant exhibited potent
4437 antiviral activity (Chiang et al. 2002). The preparation *Plantaglucid*, made from a water extract of the plant, is used as an
4438 anti-ulcer treatment and to heal wounds (Poludenny and Zhuravlev 2000). A preparation of the plant is used to treat respi-
4439 ratory tract diseases, pertussis, lung tuberculosis, and chronic nephritis (Khodzhimatov 1989).

4440 **Phytochemistry:** Leaves contain the glycoside aucubin, phenolic compounds (caffeic acid, chlorogenic acid, ferulic acid,
4441 and p-coumaric acid), triterpenes (oleanolic acid and ursolic acid), bitter substances, tannins, carotene, vitamin C and K,
4442 high amounts of potassium, mucilage, organic acids, saponins, essential oil, flavonoids (baicalein, scutellarin, apigenin,
4443 etc.), and small amounts of alkaloids. The seeds contain mucilage, fatty oil, carbohydrates, saponins, etc. (Khalmatov et
4444 al. 1984; Poludenny and Zhuravlev 2000; Chiang et al. 2002).

- Polemonium caucasicum N. Busch – Polemoniaceae** 4446
- Synonyms:** *Polemonium caeruleum* ssp. *caucasicum* (N. Busch) V.E. Avet. 4447
- English name:** Unknown 4448
- Russian name:** Синюха кавказская (Sinyukha kavkazskaya) 4449
- Uzbek name:** Unknown 4450
- Kyrgyz name:** Кавказ полемону (Kavkaz polemonu) 4451
- Description:** Herbaceous perennial, with rhizomes. Stems 50–100 cm tall. Leaves alternate, 7–20 cm long, odd-pinnatisect with 5–21 pairs of segments; segments lanceolate, sessile. Inflorescence many-flowered, corymbiform. Calyx 6–8 mm long, glandular-hairy. Corolla rotate, 8–15 mm long, blue or seldom white, 5-lobed. Stamens 5. Fruit a capsule, almost spherical, 5–7 mm long. Seeds brown, 3–3.5 mm long. 4452
4453
4454
4455
- Other distinguishing features:** Stamens exerted. Seeds angular and rugose. 4456
- Phenology:** Flowers in June-July, fruits in July-August. 4457
- Reproduction:** By seeds. 4458
- Distribution:** Ysyk-Kol, Naryn, and Chuy Provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 4459
- Habitat:** In tallgrass meadows, subalpine meadows, and meadow-steppes. 4460
- Population status:** Common, found in small groups. 4461
- Traditional use:** An infusion of the underground parts is used to treat fevers. A decoction is used as a bath to treat spazmophilia. An infusion of the leaves is used as a sedative and to treat syphilis. An infusion of the flowers is used internally to treat leucorrhoea (Plant Resources of the USSR 1990). 4462
4463
4464
- Documented effects:** The closely related species *Polemonium caeruleum* L. contains saponins that act as an expectorant. The plant has also been shown to have hemostatic effects and acts as a highly effective sedative (8–10 times that of *Valeriana*), but can be fatal to experimental animals at high doses. Preparations are used as expectorants, sedatives, treatments for stomach and duodenum ulcers, epilepsy, and chronic and acute bronchitis (Akopov 1990). 4465
4466
4467
4468
- Phytochemistry:** The entire plant of the closely related species, *Polemonium caeruleum*, contains triterpene saponins, triterpene glycosides, resins, organic acids, essential oils, fatty oils, and many macro- and micro-elements (Akopov 1990; Kurochkin 1998). 4469
4470
4471



▲ *Picea schrenkiana* Fisch. & C.A. Mey Photos: *left* and *right*: Vladimir Epiktetov; *center*: Evgeny Davkaev

▼ *Plantago lanceolata* L. Photos: *left*: Dmitri Oreshkin; *right*: Sergey Appolonov



▼ *Polemonium caucasicum* N. Busch Photo: Vladimir Epiktetov



► *Plantago major* L.

Photos: *left*: Andrei Lubchenko; *right*: Vadim Prokhorov



- Polygala hybrida DC.– Polygalaceae** 4473
- Synonyms:** *Polygala comosa* var. *altaica* Chodat, *Polygala comosa* Schkuhr var. *hybrida* (DC.) Petelin, (some consider *P. hybrida* a synonym of *P. comosa* Schkuhr). 4474
- English name:** Milkwort 4475
- Russian name:** Истод гибридный (Istod gibridnyy) 4476
- Uzbek name:** Unknown 4477
- Kyrgyz name:** Аргын истод (Argyn istod) 4478
- Description:** Herbaceous perennial. Stems 15–40 cm tall, short-hairy. Leaves sessile, 1.5–4.5 cm long, 2–4 cm wide, elliptic or lanceolate, margins entire. Inflorescences densely flowered, terminal racemes. Calyx with 3 outer, elliptic-lanceolate sepals, and 2 inner, large petaloid, elliptic sepals. Corolla with 3 petals, purple or pink, keel shorter than lateral petals. Fruits winged capsules, 6 mm long. Seeds densely covered with appressed hairs. 4479
- Other distinguishing features:** Filaments connate for the entire length. Capsules oblong. Seeds arillate. 4480
- Phenology:** Flowers in June-July, fruits in July-August. 4481
- Reproduction:** By seeds. 4482
- Distribution:** All provinces of Kyrgyzstan; Tashkent, Farg'ona and Samarqand provinces of Uzbekistan. 4483
- Habitat:** In subalpine and alpine meadows and meadow-steppes. 4484
- Population status:** Common, found growing as individual plants. 4485
- Traditional use:** An infusion of the herb is used to treat heart and gastrointestinal illnesses and rabid dog bites. The herb is also used to treat croupous pneumonia, coughs, asphyxia, fainting, sore throats, and oral diseases. In Mongolian medicine it is used as an expectorant to treat tuberculosis, purulent pleuritis, and as a hemostatic to treat uterine bleeding. In the Tibetan and Mongolian medicine the seeds are used to treat myopathy, obesity, tumors and wounds, and as a hemostatic (Plant Resources of the USSR 1988). 4486
- Documented effects:** Other species of *Polygala* have been shown to contain biologically active saponins that exhibit significant immunological properties in vitro (Desbène et al. 1999; Estrada et al. 2000). 4487
- Phytochemistry:** The roots contain carbohydrates, saponins, tannins, and fatty oil. The aboveground parts contain alkaloids (Turova and Sapozhnikova 1984; Lugmanova et al. 2007). 4488

4500 **Polygonum aviculare L. – Polygonaceae**

4501 **Synonyms:** *Polygonum aequale* Lindm., *Polygonum agreste* Sumner, *Polygonum aphyllum* Krock., *Polygonum araraticum*
4502 Kom., *Polygonum arenastrum* Boreau, *Polygonum berteroi* Phil., *Polygonum heterophyllum* Lindm., *Polygonum retinerve*
4503 Vorosch., *Polygonum striatum* K. Koch, *Polygonum uruguense* H. Gross.

4504 **English name:** Prostrate knotweed, Yard knotweed

4505 **Russian name:** Горец птичий (Gorets ptichiy)

4506 **Uzbek name:** Kiziltasma

4507 **Kyrgyz name:** Тошолгон кымыздык (Tosholgon kymyzdyk)

4508 **Description:** Herbaceous annual, with a slightly-branched taproot. Stems prostrate or suberect, 7–60 cm long. Leaves alter-
4509 nate, of 2 sizes; early leaves lanceolate, 2.5–6 cm long, 4–15 mm wide; later leaves much reduced; ocreae 4–8 mm long,
4510 membranaceous, lacerate. Flowers many, very small, in groups of 2–5 at nodes. Tepals 5, partially connate, white, green-
4511 ish or pink-red. Fruits triquetrous, dark-brown achenes.

4512 **Other distinguishing features:** The fruits are equal to or slightly exserted past the tepals.

4513 **Phenology:** Flowers and fruits in May-October.

4514 **Reproduction:** By seeds.

4515 **Distribution:** Almost all provinces of Kyrgyzstan and Uzbekistan.

4516 **Habitat:** In meadows and fallow or cultivated fields, from foothills up to the alpine belt of the mountains.

4517 **Population status:** Common, found in dense groups.

4518 **Traditional use:** A decoction and infusion of the herb is used to treat stomach spasms, intestinal infections, diarrhea and as
4519 a tonic, hemostatic and diuretic. The plant is used in a bath to treat bacterial and fungal skin diseases and rashes. The fresh
4520 herb is put on tumors, wounds, and skin ulcers (Khalmatov et al. 1984). An infusion of the herb is used to wash the head
4521 to increase the health of hair and encourage hair growth. A decoction of the herb in milk is taken to treat convulsions
4522 (Poludenny and Zhuravlev 2000).

4523 **Documented effects:** Water and alcohol extracts of the plant have been shown to increase the rate of blood coagulation,
4524 decrease blood pressure, increase inhalation volume, improve lung function, tone uterine muscles, and increase diuresis.
4525 The preparation *Avicularen* is used in gynecological practice as a hemostatic (Khalmatov et al. 1984). The infusion of the
4526 herb is recommended to improve metabolism and treat diabetes (Kurochkin 1998). Experiments indicate that a methanol
4527 extract of the plant has anti-fibrotic effects on rats with induced liver fibrosis (Nan et al. 2000).

4528 **Phytochemistry:** The plant contains essential oils, vitamin K₁, sugars, saponins, coumarins, mucilage, anthraglycosides, etc.
4529 (Khalmatov et al. 1984). Leaves contains tannins, flavonoids (avicularin), vitamin C, carotene, and silicic acid compounds
4530 (Tolmachev 1976).

4531

- Polygonum coriarium Grig. – Polygonaceae** 4532
- Synonyms:** *Aconogonon bucharicum* (Grig.) Holub, *Aconogonon coriarium* (Grig.) Soják, *Aconogonon coriarium* ssp. 4533
bucharicum (Grig.) Soják, *Pleuropteropyrum bucharicum* (Grig.) Nevski, *Polygonum bucharicum* Grig. 4534
- English name:** Unknown 4535
- Russian name:** Горлец дубильный (Gorlets dubil'nyy) 4536
- Uzbek name:** Taran dubil'nyy 4537
- Kyrgyz name:** Ашаткыч кымыздык (Ashatkych kymyzdyk) 4538
- Description:** Herbaceous perennial, with a large rhizome (up to 5–8 kg). Stems up to 1–1.5 m tall, abundantly branched, 4539
glabrous. Leaves alternate, short-petiolate, ovate or ovate-lanceolate, 6–10 cm long, 2.5–5 cm wide, base wide-cuneate or 4540
rounded, abaxial side (sometimes both sides) densely hairy, rarely glabrous, margin bristly-ciliate. Ocreae membranous, 4541
tubular, 1.5–2.5 cm long, brown, not persisting. Inflorescence large panicle, branched, dense, up to 35 cm long and 25 cm 4542
wide. Perianth usually 2.5–3.5 cm long, with 5 white tepals. Fruit a triquetrous nutlet with sharp edges, 3–4.5 mm long, 4543
shiny, slightly exserted from perianth. 4544
- Other distinguishing features:** Stamens 8, styles 3. Branches of inflorescence nodding in fruit. 4545
- Phenology:** Flowers and fruits in June-August. 4546
- Reproduction:** By seeds and rhizomes. 4547
- Distribution:** Toshkent, Samarqand, and Surxondaryo provinces of Uzbekistan; Talas, Osh and Jalal-Abad provinces of 4548
Kyrgyzstan. 4549
- Habitat:** The tau zone. Stony, shallow soil on wet slopes of mountains. 4550
- Population status:** Common. 4551
- Traditional use:** A decoction of the underground plant parts is used in folk medicine as an astringent for treatment of diar- 4552
rhea with and without blood (Khalmatov 1964). 4553
- Documented effects:** Astringent preparations effective for gastrointestinal tract diseases of alimentary origin were prepared 4554
from root powder in combination with protein (called *Taranalbin*) or formaldehyde (called *Taranform*). An extract 4555
obtained from the plant roots in the dose of 20 mg/kg increases stability for exercise stress (physical activity) and elon- 4556
gates swimming time of mice up to 61%. Proanthocyanidin and catacin (katacine) have distinct antihypoxic action and 4557
decrease the oxygen-need of tissues (Kurmukov et al. 1991b), which is connected to its influence on energy metabolism 4558
(Nazrullaev et al. 1990). 4559
- Phytochemistry:** Underground organs contain up to 28–35 % tannins, mainly of the pyrocatechin group (proanthocyanidin; 4560
Ogolevitz 1951). Many proanthocyanidins have been isolated from the roots (Makhmatkulov et al. 1992, 1994; Keneshov 4561
et al. 1997a, b). The leaves contain flavonoids (Chumbalov and Omurkamzinova 1968). 4562

4564 **Portulaca oleracea L. – Portulacaceae**

4565 **Synonyms:** *Portulaca consanguinea* Schldtl., *Portulaca intermedia* Link ex Schldtl., *Portulaca marginata* Kunth, *Portulaca*
4566 *mundula* I.M. Johnst., *Portulaca neglecta* Mack. & Bush, *Portulaca pilosa* L., *Portulaca pusilla* Kunth, *Portulaca retusa*
4567 Engelm.

4568 **English name:** Purslane, Little hogweed

4569 **Russian name:** Портулак огородный (Portulak ogorodnyy)

4570 **Uzbek name:** Semiz ut

4571 **Kyrgyz name:** Огород портулагы (Ogorod portulagy)

4572 **Description:** Herbaceous annual. Stem 10–35 cm long, glabrous, fleshy, prostrate, spreading, branched from the base. Leaves
4573 alternate or sub-opposite, obovate or spatulate, 4–28 × 2–13 mm, apex rounded to obtuse, fleshy, sessile. Flowers 3–10 mm
4574 wide, solitary or in small clusters of 2–3 in branch and leaf axils. Sepals 2, deciduous. Petals usually 5, yellow, obovate.
4575 Fruit a circumscissile capsule, ovoid, 5–8 mm long, many-seeded. Seeds orbiculate or elongate, flattened, surface covered
4576 with tubercles, black to dark brown, slightly shining, 0.7–1 mm long, 0.25 mm wide.

4577 **Other distinguishing features:** Stamens usually 8–15. Stigmas 3–6.

4578 **Phenology:** Flowers and fruits in June-October.

4579 **Reproduction:** By seeds.

4580 **Distribution:** All regions of Uzbekistan and Kyrgyzstan.

4581 **Habitat:** The chul, adyr and tau zones. A weed of irrigated agricultural areas.

4582 **Population status:** Common.

4583 **Traditional use:** A decoction of the herb is used in Chinese medicine to alleviate pain and swelling, as an anti-inflammatory
4584 and diuretic, and for gonorrhea, kidney and liver diseases, bacterial dysentery, syphilitic arthritis, and palsies of infectious
4585 origin. In Central Asia this herb is used as a choleric, an antipyretic for fevers associated with hepatitis, nephritis, and
4586 cystitis, and as a treatment for intestinal ulcers and bloody diarrhea. The plant is also used to treat intestinal infections
4587 (Khalmatov 1964; Chen et al. 2003).

4588 **Documented effects:** An extract of this herb sharply increases blood pressure, due to its high noradrenaline content.
4589 Hemostatic action for internal hemorrhaging has been documented (Khalmatov 1964). In experiments with mice and rats,
4590 an ethanolic extract of the dried aboveground parts showed significant antiinflammatory and analgesic effects after intra-
4591 peritoneal and topical, but not oral, administration (Chan et al. 2000). Studies indicated that the consumption of the plant
4592 may help to reduce the occurrence of cancer and heart diseases. Catecholamines (noradrenaline and dopamine) contained
4593 in the plant are generally considered to be the effective component in the treatment of shock. Studies have also shown that
4594 noradrenaline is a modulator of the immune system and may have anti-cancer properties (Chen et al. 2003).

4595 **Phytochemistry:** The herb contains alkaloids, glycosides, traces of saponins, and bitter substances (Khalmatov 1964).
4596 250 mg% noradrenaline has been obtained from the fresh herb (Khalmatov 1964). The plant contains an abundance of the
4597 catecholamines noradrenaline and dopamine, free oxalic acids, alkaloids, coumarins, flavonoids, cardiac and anthraqui-
4598 none glycosides, proteins, high amounts of beta-carotenes and has a higher content of omega-3 fatty acids (especially
4599 α -linolenic acid) than many other vegetables (Guil-Guerrero and Rodríguez-García 1999; Chen et al. 2003; Fontana et al.
4600 2006).



▲ *Polygala hybrida* DC. Photos: Lina Valdshmit

▼ *Polygonum coriarium* Grig. Photos: Vladimir Epiktetov



▼ *Polygonum aviculare* L. Photos: *top right* and *bottom right*: Vadim Prokhorov; *bottom left*: Denis Mirin



▲ *Portulaca oleracea* L.

Photos: *top*: Andrei Lubchenko;
bottom: Mary Backlund



4602 **Potentilla canescens Bess. – Rosaceae**4603 **Synonyms:** *Potentilla adscendens* Waldst. & Kit. ex Willd., *Potentilla inclinata* Vill.4604 **English name:** Hoary cinquefoil, ashy cinquefoil4605 **Russian name:** Лапчатка седоватая (Lapchatka sedovataya)4606 **Uzbek name:** Unknown4607 **Kyrgyz name:** Агыш туктуу казтаман (Agysh tuktuu kaztaman)

4608 **Description:** Herbaceous perennial. Stems few, erect, 10–50 cm tall, pubescent. Leaves palmately compound with 5–7
4609 leaflets, stipulate, petioles pubescent; leaflets obovate or obovate-lanceolate, pubescent, margins coarse serrate.
4610 Inflorescence many-flowered, corymbiform or cymose-paniculiform. Flowers pedicellate, ca. 10 mm in diameter. Sepals
4611 5, epicalyx segments 5, alternating with sepals. Petals 5, yellow, ovate, slightly longer than sepals. Fruits wrinkled
4612 achenes.

4613 **Other distinguishing features:** Lower side of leaflets tomentose. Base of style thickened.4614 **Phenology:** Flowers in June, fruits in August.4615 **Reproduction:** By seeds.4616 **Distribution:** Naryn, Talas, and Chuy provinces of Kyrgyzstan; Tashkent, Andijon, Namangan, and Farg'ona provinces of
4617 Uzbekistan.4618 **Habitat:** In the foothills and steppe belt of mountains, along roads, fallow fields and in lowland steppes.4619 **Population status:** Common, found in small groups.4620 **Traditional use:** An infusion of the underground parts is used to treat menorrhagia, diarrhea, and hematuria. An infusion of
4621 the aboveground parts is used to treat laryngitis (Plant Resources of the USSR 1987).4622 **Documented effects:** No data.4623 **Phytochemistry:** The rhizomes and flowers contain traces of alkaloids. Leaves and flowers contain vitamin C (Plant
4624 Resources of the USSR 1987).

4625

Prangos pabularia Lindl. – Apiaceae

Synonyms: *Hippomarathrum sarawschanicum* Regel & Schmalh., *Hyalolaena sewerzowii* Regel & Herd., *Koelzella pabularia* (Lindl.) Hiroe, *Prangos cylindrocarpa* Korovin, *Prangos hissarica* Korovin, *Prangos lamellata* Korovin, *Prangos seravschanica* (Regel & Schmalh.) Korovin.

English name: Unknown

Russian name: Прангос кормовой (Prangos kormovoy)

Uzbek name: Tulky kuyruq

Kyrgyz name: Тоют аюу чачы (Toyut ayuu chachy)

Description: Herbaceous perennial with thick taproot. Stems several, up to 0.6–2 m tall, angular-striated, strongly branching from the middle, nearly glabrous. Basal leaves densely clustered, pointing upward, long-petiolate; blades 30–70 cm long, 6–12 cm wide, elliptic or oblong in outline, 4–5-pinnate with filiform or narrow-linear segments. Inflorescence an irregular compound umbel, 8–20 rays; umbellets 10–15-flowered. Sepals triangular, acute. Petals obovate, ca.1.5 mm long, yellow. Fruit a schizocarp with 2 mericarps; mericarps oblong-cylindrical, 15–18 mm long, often violet in color with prominent ribs, grooves between ribs narrow, lined with tubercles.

Other distinguishing features: Leaves cause strong burns and photosensitivity when touched. Leaves quickly senescing after which the stem is covered with leaf remnants. Flowers along the outer margin of the umbellets are bisexual; flowers in the center are male.

Phenology: Flowers in May-June, fruits in June-July.

Reproduction: By seeds.

Distribution: Toshkent, Samarqand, Jizzax, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Osh and Jalal-Abad provinces of Kyrgyzstan.

Habitat: The tau and yailau zones. Clayey and clayey-stony mountain slopes.

Population status: Common, often found in large populations.

Traditional use: Root decoctions and root tinctures, sometimes mixed with tinctures of iodine and St. John's wort (*Hypericum*), are used to treat scabies in humans (Ogolevitz 1951). A decoction of the roots is used to disinfect the mouth and to kill ticks, fleas, and bed bugs on farm animals. The roots are put on hot ashes and after 2–3 h are then put on surface wounds. The above and underground parts are used in a bath to treat skin diseases (scabies, fungal, etc.). A decoction of the aboveground parts is used as a mouth wash to treat toothaches (Khodzhimatov 1989).

Documented effects: In experiments the coumarin osthol increased blood pressure, pulse rhythm, stimulated respiratory activity, weakened acetylcholine effect, and had vermifugal activity. (Ogolevitz 1951; Jamwal et al. 1962). A butenyl coumarin isolated from the plant had analeptic activity on respiration and the heart, stimulated brain functions, and exhibited antiacetylcholinic and antihistaminic action (Chicco 1966). It is also used as antidote in the poisoning due to hypnotics. The coumarin osthol showed antibiotic activity against *Staphylococcus aureus* (MRSA) and *Pseudomonas aeruginosa*. Twelve compounds isolated from this plant had immunosuppressive activity (Tada et al. 2002).

Phytochemistry: Leaves, fruits, roots and root resin contain coumarins. The coumarin osthol and the furocoumarins oxypeucedanin, imperatorin, prangenin, prangenidin, and others have been isolated from the total coumarins. Seeds contain 0.2–0.3 % alkaloids from which the alkaloid prangosine has been isolated (Ogolevitz 1951; Yunusov 1981; Khodzhimatov 1989). This plant is a rich source of coumarins, coumarin derivatives, and terpenoids with 29 different compounds being identified (Tada et al. 2002). One hundred twenty-eight compounds were characterized from the volatile constituents of the fruits. The major constituents of the essential oil were α -humulene, bicyclogermacrene, spathulenol, germacrene D, and α -pinene (Ozek et al. 2007).

4626
4627
4628
4629
4630
4631
4632
4633
4634
4635
4636
4637
4638
4639
4640
4641
4642
4643
4644
4645
4646
4647
4648
4649
4650
4651
4652
4653
4654
4655
4656
4657
4658
4659
4660
4661
4662
4663
4664
4665
4666

4667

4668 **Prunus sogdiana Vassilcz. – Rosaceae**

4669 **Synonyms:** *Prunus cerasifera* ssp. *sogdiana* (Vassilcz.) Cinovskis, *Prunus cerasifera* var. *orientalis* Popov, *Prunus mirabilis*
4670 Sumner, *Prunus orientalis* (Popov) Kudr.

4671 **English name:** Sogdian plum

4672 **Russian name:** Слива согдийская (Sliva sogdiyskaya)

4673 **Uzbek name:** Togolcha

4674 **Kyrgyz name:** Жапайы алча (Zhapayy alcha)

4675 **Description:** Tree or shrub from 2.5 to 7 m tall, with multiple trunks. Older bark dark-gray, cracked; young branches
4676 brownish-green to red-brown. Leaves alternate, petiolate, elliptic, ovate or obovate, 4.5–5.6 cm long, 2.2–4 cm wide,
4677 glabrous above, light in color and pubescent along midvein below, margins serrate or serrate-crenate. Flowers ca. 2 cm in
4678 diameter, pedicillate. Sepals 5, glabrous. Petals 5, ovate, white or with purple base. Fruit a dark purple drupe, spherical to
4679 slightly elongated, 1–2 cm in diameter, often glaucous.

4680 **Other distinguishing features:** Stamens 15–30, in 2 whorls, filaments unequal in height.

4681 **Phenology:** Flowers in May, fruits in July–October.

4682 **Reproduction:** By seeds.

4683 **Distribution:** Jalal-Abad, Osh, and Chuy provinces of Kyrgyzstan; Toshkent and Surxondaryo provinces of Uzbekistan.

4684 **Habitat:** On the edges, and in the understory of deciduous forests, and among bushes.

4685 **Population status:** Common, forming dense groups.

4686 **Traditional use:** The fruits, leaves, flowers, bark and gum are used in folk medicine. An infusion of the leaves and flowers
4687 is used as a light laxative. A decoction of the dried fruits is used to increase appetite, to aid in digestion, and as an expect-
4688 torant. The gum is used as a treatment for coughs. A water extract of the bark and roots is used as a diaphoretic, anti-
4689 pyretic, and anti-inflammatory (Nuraliev 1989). In Pamir-Alai it is used to treat acute respiratory diseases (Zapryagaeva
4690 1964).

4691 **Documented effects:** No data.

4692 **Phytochemistry:** Fruits contain sugars, organic acids (malic and citric), vitamin C, provitamin A, pectins, tannins, minerals,
4693 and fatty oil (Nuraliev 1989).

4694

- Pseudosophora alopecuroides (L.) Sweet – Fabaceae** 4695
- Synonyms:** *Goebelia alopecuroides* (L.) Bunge, *Sophora alopecuroides* L., *Vexibia alopecuroides* (L.) Yakovl. 4696
- English name:** Unknown 4697
- Russian Name:** Вексибия лисохвостная, Талхак обыкновенный (Veksibiya lisokhlostnaya, Talkhak obyknovenny) 4698
- Uzbek name:** Akmia, Achikmia 4699
- Kyrgyz name:** Ак мяя (Ak myya) 4700
- Description:** Herbaceous perennial. Stems 40–70 cm tall. Leaves alternate, odd-pinnate, with 5–12 pairs of oval or elongate-ovate leaflets; leaflets 1.2–3 cm long, 4–12 mm wide. Inflorescence an densely-flowered, apical raceme. Calyx widely campanulate with 5 uneven teeth. Corolla papilionaceous, white or slightly yellowish. Fruit a legume, 5–12 cm long, constricted between the seeds, with extended tip at the end. Seeds spherical, light-brown, smooth. 4701–4704
- Other distinguishing features:** The whole plant is gray-green hairy. 4705
- Phenology:** Flowers in May-June, fruits in June-August. 4706
- Reproduction:** By seeds and rhizomes. 4707
- Distribution:** All provinces of Kyrgyzstan; Tashkent, Farg'ona, Samarqand and Bukhara provinces of Uzbekistan. 4708
- Habitat:** In foothills and in abandoned and cultivated fields. 4709
- Population status:** Common, found in dense groups. 4710
- Traditional use:** In Tibetan and Mongolian medicine the roots are used to treat diseases of the heart, aorta, and vascular system, diphtheria, and rheumatism, and are used as an antipyretic and restorative as well (Khaidav 1965). The ground seeds are used to treat poor digestion and loss of appetite (Khalmatov 1964). 4711–4713
- Documented effects:** In modern medicine preparations of this plant are used to increase respiratory function (Khalmatov 1964). In experiments, large doses of sophocarpine act as a weak ganglioblocker and cause contraction of the myometrium. Matrine, sophoridine, sophocarpine, and aloperine have stimulating activities, but aloperine causes short-term hypotension. Sophocarpine and sophoridine cause narrowing of the peripheral vessels, and in small doses strengthen peristalsis and intestinal tonus, paralyze skeletal muscles, and have ganglioblocking properties (Georgadze 1938; Kruglikova-Livova 1952). Quinolizidine alkaloids *isolated from the plant* have very weak antiviral activities (Ma et al. 2002a). 4714–4720
- Phytochemistry:** The roots contain alkaloids (spartein, sophoridine, and sophocarpine) and flavonoids (quercetin and rutoside), anthraquinones (aloemodin, anthraquinone sennosides, etc.). The aboveground parts contain alkaloids (sophoridine, cytosine, neosophoramine, sophoramine, sophocarpine and aloperine; Yusupova et al. 1984; Plant Resources of the USSR 1987; Iinuma et al. 1995). The alkaloids oxymatrin, oxysophocarpine, cytosine, matrine, sophocarpine, sophoridine, and nicotine have been isolated from the seeds (Zhang et al. 1997). 4721–4725



▲ Potentilla canescens Bess.

Photo: Andrei Lubchenko



▲ Pseudosophora alopecuroides (L.) Sweet

Photos: Lina Valdshmit



▲ Prunus sogdiana Vassilcz. Photos: *left*: John B. Taft; *right*: Vladimir Epikhetov



▼ Prangos pabularia Lindl. Photos: Evgeny Davkaev



Psoralea drupacea Bunge – Fabaceae	4727
Synonyms: <i>Cullen drupacea</i> (Bunge) C.H. Stirt., <i>Lotodes drupaceum</i> (Bunge) Kuntze.	4728
English name: Scurfy-pea	4729
Russian name: Псоралея костянковая (<i>Psoraleya kostyankovaya</i>)	4730
Uzbek name: Ok kuraiy	4731
Kyrgyz name: Соокчёлуу ак куурай (<i>Sookchyoluu ak kuuray</i>)	4732
Description: Herbaceous perennial to 40–150 cm tall, with vigorous, thick, woody roots. Stems erect, branched, densely hairy, glandular. Leaves alternate, simple or ternate, short-petiolate; leaflets nearly round, 1.5–5 cm long, 2–6 cm wide, densely hairy beneath with glands on both sides, margins coarsely dentate; stipules linear-lanceolate, 0.5–1.5 cm long, hairy, glandular. Inflorescences in loose axillary racemes. Flowers 4–7 mm long, on very short pedicels. Calyx tubular-campanulate with unequal teeth, densely hairy with glands. Corolla white-lilac. Fruit a 1-seeded legume, suborbicular, densely hairy, ca. 5 mm long, 2.5–3.5 mm wide. Seed very small, adnate to the fruit wall.	4733 4734 4735 4736 4737 4738
Other distinguishing features: The root has a yellow color inside. The fruit is indehiscent and beakless.	4739
Phenology: Flowers in May-June-July, fruits in June-September.	4740
Reproduction: By seeds.	4741
Distribution: Toshkent, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Chuy, Talas and Jalal-Abad provinces of Kyrgyzstan.	4742 4743
Habitat: The chul and adyr zones. Found in combination with ephemeroïd vegetation in shallow, loamy, sierozem soil, rocky-clay loams, and light-clay loams.	4744 4745
Population status: Common, found in large populations.	4746
Traditional use: Leaf powder is used in folk medicine to treat abscesses (furuncles and carbuncles), vitiligo, eczema, and hair loss (Shimanov 1972). The essential oil from the fruits and galenical preparations of the legumes and roots are used to treat skin problems (Mamedov et al. 2004).	4747 4748 4749
Documented effects: The chemical psoralen has photosensitizing, estrogenic, contraceptive, and embryotoxic actions (Shimanov 1972; Kurmukov and Akhmedkhodzhaeva 1975; Kurmukov et al. 1976). A medical preparation <i>Psoralen</i> (a mixture of furocoumarins from the fruits) is used to treat vitiligo and patchy, irregular hair loss. Treatment with the preparation <i>Drupanol</i> , at the dose of 10 mg/kg for 10 days, has anabolic and androgenic effects. In tests on chickens, the androgenic effect of drupanol caused a significant growth stimulation of the crest. The testing groups' crest increased by 1.6 times compared to control group (Syrov et al. 1976; Akopov 1990).	4750 4751 4752 4753 4754 4755
Phytochemistry: The aboveground parts contain the antibiotic bakuchiol and the coumarin umbelliferol. The fruits and roots contain psoralen, isopsoralen, tannins, and semi-solid essential oil, and the fruits contain fatty oil, the alkaloid drupacine, and the phenol drupanol (Golovina and Nikonov 1973). The mature fruits have the highest coumarin content (0.1 %) and the roots contains tannins (Akopov 1990).	4756 4757 4758 4759

4761 ***Pulicaria salviifolia* Bunge – Asteraceae**

4762 **Synonyms:** *Pulicaria afghanica* Kitam., *Pulicaria lachnophylla* C. Winkl., *Pulicaria olivascens* Rech. f., *Pulicaria sublepi-*
4763 *dota* Rech. f.

4764 **English name:** Sage-leaf fleabane

4765 **Russian name:** Блошница шалфеелистная (Bloshnitsa shalfeelistnaya)

4766 **Uzbek name:** Gulband

4767 **Kyrgyz name:** Шалфей жалбырактуу пуликария (Shalfey zhalbyraktuu pulikariya)

4768 **Description:** Herbaceous perennial, 40–60 cm tall, densely covered with woolly hairs. Stem single or few, each with many
4769 straight branches from near the base. Leaves simple, spatulate, apex obtuse, gradually tapering to petiole, margins sinuate;
4770 lower leaves much larger than upper. Inflorescences capitula, numerous, arranged in loose racemes. Involucre 9–15 mm
4771 in diameter; involucre bracts usually in 2 rows, lanceolate, acute with membranaceous margin, grayish-hairy; inner bracts
4772 much more narrow and membranaceous. Flowers yellow; ray flowers with short, obovate ligules; disk flowers narrow,
4773 5–8 mm long. Fruits achenes, 2–2.7 mm long, sparsely hairy, glandular; inner row of pappus plumose-barbed, whitish
4774 hairs, 5–8 mm long.

4775 **Other distinguishing features:** Plant strongly sweet-scented, smelling like honey. Varieties of this species differ in the color
4776 and amount of pubescence.

4777 **Phenology:** Flowers in July-August, fruits in August-September.

4778 **Reproduction:** By seeds.

4779 **Distribution:** Toshkent, Farg'ona, Samarqand, and Surxondaryo provinces of Uzbekistan; Osh and Jalal-Abad provinces of
4780 Kyrgyzstan.

4781 **Habitat:** The adyr zone. Dry stony, slopes with rocky debris and pebbly, gypsum-soiled foothills.

4782 **Population status:** Common, often forming dense groups.

4783 **Traditional use:** A decoction of the herb is used for decreasing blood sugar content for treatment of diabetes (Abdunazarov
4784 2000).

4785 **Documented effects:** The flavonoid pulicarin exhibited hypolipidemic effects in experiments with rats (Sagitdinova et al.
4786 1992). When administered orally, salvin, salvicin and salvifolin, show significant hypoglycemic activity in rats
4787 (Tashmukhamedova et al. 1992).

4788 **Phytochemistry:** The plant contains terpenoids and diterpenoids (salvin, salvifolin, salvicin, salvicinin, salvicinolide, and
4789 salvicinolin, etc.), as well as flavonoids (rutin, etc.), triterpenoids, and sterols (Nurmukhamedova et al. 1985, 1986;
4790 Sagitdinova et al. 1992, 1994; Eshbakova et al. 1997; Eshbakova and Saidkhodzhaev 2001).

4791

Reseda luteola L. – Resedaceae	4792
Synonyms: None	4793
English name: Weld, Yellow dye, Dyer's rocket	4794
Russian name: Резеда жёлтенькая (Rezeda zhyolten'kaya)	4795
Uzbek name: Sayok	4796
Kyrgyz name: Сары резеда (Sary rezeda)	4797
Description: Herbaceous biennial, up to 80 cm high. Stem single, erect, glabrous, densely foliaceous. Leaves alternate, oblanceolate to linear, 3–9 cm long, 5–12 mm wide, glabrous, sessile, margins entire. Inflorescences spiciform racemes, 15–45 cm long, erect. Flowers with 4 rounded sepals and 4 yellowish, irregularly lobed petals. Stamens 20–25. Fruit a subglobose capsule, 3-parted. Seeds ca.1 mm long, brown-black, glabrous, shiny, smooth.	4798 4799 4800 4801
Other distinguishing features: The bracts, calyx, and filaments are persistent during fruiting.	4802
Phenology: Flowers and fruits in May-August.	4803
Reproduction: Only by seeds.	4804
Distribution: Toshkent, Samarqand, and Surxondaryo provinces of Uzbekistan; Jalal-Abad province of Kyrgyzstan.	4805
Habitat: The adyr and tau zones. Dry hills, along roads, river valleys, mountain slopes and near field crops.	4806
Population status: Uncommon, found in small populations of 4–8 individuals.	4807
Traditional use: A decoction of the root is taken as a vermifuge (Khalmatov 1964).	4808
Documented effects: The glycoside glucobarbarin has antithyroid action (Khalmatov 1964). In an inhibition assay, an extract of the plant inhibited the activity of trypsin by 97 % (Johansson et al. 2002). Luteolin showed anti-inflammatory activity in a variety of different in vivo assays and has also exhibited anti-cancer activity (Chowdhury et al. 2002; Ziyani et al. 2007).	4809 4810 4811 4812
Phytochemistry: The aboveground parts of the herb contain 1–3 % dyeing substances (luteolin), mustard essential oil (mostly in roots). 32–34 % fatty oil was extracted from the seeds. The leaves, inflorescence and the seeds include the glycosides glucocapparin and glucobarbarin (Khalmatov 1964). The aboveground parts contain cinnamamide and alkaloids (Lutfullin et al. 1976, 1977). The plant was also found to contain phenyl- β -naphthylamine (Sultankhodzhaev and Tadzhibaev 1976).	4813 4814 4815 4816 4817

4819 **Rhamnus cathartica L. – Rhamnaceae**4820 **Synonyms:** None4821 **English name:** Common buckthorn4822 **Russian name:** Жостер слабительный (Zhoster slabitel'nyy)4823 **Uzbek name:** Togzhumroot4824 **Kyrgyz name:** Ич алдыргыч карк моюл (Ich aldyrgych kark moyul)

4825 **Description:** Bush or small tree, up to 3–8 m tall, usually dioecious. Old bark nearly black, rough, exfoliating; young bark
4826 red-brown; some branches ending in short spines. Leaves opposite, ovate to elliptic, 2–8 cm long, 1.5–5 cm wide, base
4827 round-cuneate, margins crenate-serrate. Flowers perfect or unisexual, in leaf axils. Sepals 4, twice as long as petals. Petals
4828 erect, lanceolate, 1–1.5 mm long in staminate flowers, ca. 0.5 mm in pistillate flowers. Fruits black drupes, 6–8 mm in
4829 diameter, juicy, round, shiny.

4830 **Other distinguishing features:** Lateral leaf veins strongly upcurved. Fruits 4-seeded.4831 **Phenology:** Flowers in May, fruits in August.4832 **Reproduction:** By seeds.4833 **Distribution:** All provinces of Kyrgyzstan; Toshkent, Farg'ona and Samarqand provinces of Uzbekistan.4834 **Habitat:** On mountain slopes, along rivers, among bushes and in forest plantations.4835 **Population status:** Common, found in small groups.

4836 **Traditional use:** Thin branches, bark and fruits are used in folk medicine. Fruits are used as a laxative to treat chronic con-
4837 stipation, and the treatment starts to take effect 8–10 h after ingestion. A decoction of the fruits, with the addition of dairy
4838 whey, oil, and baking soda is used as a laxative for atonic and spastic constipation, and is used as a stool softener for cases
4839 of hemorrhoids and wounds to the colon. An infusion of the fruits in vodka is used externally to treat rheumatism.
4840 A decoction of the branches is used internally to treat ulcers and externally, as a compress to heal wounds. A decoction of
4841 the bark is used to treat stomach catarrh, low acidity of the stomach, and Polish plait (Maznev 2004).

4842 **Documented effects:** Preparations of this species have laxative properties that are associated with the presence of anthragly-
4843 cosides and related compounds (mainly emodin), which act by stimulating the walls of the large intestines and moderately
4844 strengthen wave-and pendulum-like movements (Maznev 2004). An ethanolic extract of this species had high antimyco-
4845 bacterial activity but also purgative effects (Newton et al. 2000).

4846 **Phytochemistry:** Fruits contain anthraglycosides (glucofrangulin, frangulin, frangula-emodin, etc.), flavonol glycosides,
4847 sugars, organic acids, pectins, etc. Leaves contain ascorbic acid. Bark contains chrysophanic acid, anthraglycosides, and
4848 high amounts of tannins (Khalmatov et al. 1984).

4849



▲ *Rhamnus cathartica* L. Photos: Dmitri Oreshkin



▲ *Reseda luteola* L. Photos: Bazar Dovletov

▶ *Psoralea drupacea* Bunge Photos: *left:* Evgeny Davkaev; *right:* Alim Gaziev



▼ *Pulicaria salviifolia* Bunge
Photos: Alim Gaziev



4850 **Rheum maximowiczii Losinsk. – Polygonaceae**4851 **Synonyms:** *Rheum emodi* Wall., *Rheum megalocarpon* Maxim.4852 **English name:** Unknown4853 **Russian name:** Ревень Максимовича (Reven' Maksimovicha)4854 **Uzbek name:** Rovach, Chukhra4855 **Kyrgyz name:** Чукурук (Chukuruk)

4856 **Description:** Herbaceous perennial with a thick rhizome. Stem 40–100 cm tall, up to 2 cm thick, leafless, branched, tough,
4857 reddish, rough due to tiny papillae, rarely smooth. Ocreae rust-colored, tightly surrounding the stem above each leaf axil.
4858 Leaves basal, round to kidney-shaped, 18–50 cm long, 20–60 cm wide, with 3 veins and short, flat petioles. Inflorescence
4859 paniculiform, pyramidal; flowers clustered on long peduncles. Tepals 6, each 3 mm long, 1 mm wide, greenish. Fruits
4860 achenes, 2 cm long, 1.5 cm wide, winged, lilac-reddish.

4861 **Other distinguishing features:** The adaxial sides of the leaves are glabrous, but the abaxial sides are rough due to papillae
4862 near the veins.

4863 **Phenology:** Flowers and fruits in May-June.4864 **Reproduction:** By seeds.

4865 **Distribution:** Toshkent, Samarqand, Qashqadaryo, and Andijon provinces of Uzbekistan; Osh and Jalal-Abad provinces of
4866 Kyrgyzstan.

4867 **Habitat:** The tau zone. Grassy slopes or slopes with rocky debris.4868 **Population status:** Common.

4869 **Traditional use:** A decoction of the root and fresh juice from the leaves are used to treat diarrhea. Juice from the petioles is
4870 recommended as a treatment for malaria (Khalmatov 1964). The young petioles and stems, and the fresh juice or compote
4871 made from them are used as a tonic, antipyretic and hypotensive to prevent anemia and to detoxify. The plant is used to
4872 increase the appetite and to treat gastritis with low acidity, liver (hepatitis) and gallbladder diseases, tuberculosis, hemor-
4873 rhoids, chronic constipation, polyarthritis, and fevers (Nuraliev 1989).

4874 **Documented effects:** The powdered root has astringent properties (Khalmatov 1964). This species improves the liver's abil-
4875 ity to detoxify, helps patients with moderately high blood pressure, has diuretic actions, and is good for treatment of
4876 constipation and fevers (Nuraliev 1989). Compounds isolated from plants collected in Uzbekistan exhibited antioxidative
4877 activity (Kogure et al. 2004). (+)-rhododendrol and epi-rhododendrin isolated from *Acer nikoense* Maxim. suppressed
4878 nitric oxide (NO) production in mouse peritoneal macrophages in vivo (Fushiya et al. 1998).

4879 **Phytochemistry:** The roots contains tannins (catechins, gallic acid, pyrogallol, and pyrocatechin), carbohydrates, and gly-
4880 cosides (Khalmatov 1964). The aboveground parts contain vitamins C, A, E, B₁, B₂, B₆, B₁₅, organic acids (malic and
4881 oxalic), sugars, fibers, hemicellulose, pectin, and macroelements (Nuraliev 1989). Roots collected in Uzbekistan con-
4882 tained new phenylbutanoid and stilbene derivatives as well as the known compounds rhododendrol, epi-rhododendrin,
4883 lindleyin, torachryson, etc. (Shikishima et al. 2001).

4884

Rhodiola linearifolia Boriss. – Crassulaceae

- Synonyms:** *Rhodiola kirilowii* (Regel) Maxim., *Rhodiola longicaulis* (Praeger) S.H. Fu, *Rhodiola macrolepis* (Franch.) S.H. Fu, *Rhodiola robusta* (Praeger) S.H. Fu, *Sedum kirilowii* Regel, *Sedum longicaule* Praeger, *Sedum macrolepis* Franch.,
Sedum robustum Praeger.
- English name:** Unknown
- Russian name:** Родиола линейнолистная (*Rodiola lineynolistnaya*)
- Uzbek name:** Unknown
- Kyrgyz name:** Сызгыч чегендир (*Syzgych chegendir*)
- Description:** Herbaceous perennial, with a thick caudex. Stems 10–60 cm tall, densely leafy. Caudex leaves scale like; stem leaves alternate, sessile, linear-lanceolate, 2–6 cm long, 3–15 mm wide, slightly serrate. Inflorescences cymose, dense, compact. Flowers unisexual or occasionally bisexual. Sepals linear or triangular, 1.5–3 mm long. Petals 3–4 mm long, brownish-red, pink or yellow. Fruits paired elongate follicles with curved apical beaks.
- Other distinguishing features:** Stem leaves linear-lanceolate. Stamens 8–10, yellow.
- Phenology:** Flowers in May-July, fruits in July-September.
- Reproduction:** By seeds.
- Distribution:** Jalal-Abad, Naryn, Osh, Ysyk-Kol, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan.
- Habitat:** In wet places, forest meadows, and high mountains.
- Population status:** Common, found growing as individual plants.
- Traditional use:** An infusion of the underground parts is used to treat weariness, neurotic conditions, and decreased ability to work (Krasnov and Demidenko 1981).
- Documented effects:** The total flavonoids isolated from the underground parts show antitumor activity (Krasnov and Demidenko 1981). Salidroside has protective effects against oxidative stress-induced cell apoptosis and has been shown to enhance the ability of hemoglobin to carry oxygen and protect neuronal cells against hypoxia/reoxygenation injury in vitro (Chen et al. 2007; Yang et al. 2007; Zhang et al. 2007). Daucosterol was shown to have an inhibitory effect on the viral enzyme reverse transcriptase (Kimura et al. 2003).
- Phytochemistry:** Underground parts contain organic acids (citric and malic), triterpene glycoside derivatives of oleanolic acids, phenols, phenolcarbonic acids, coumarins, flavonoids, tannins, tyrosol, daucosterol, lotaustralin, salidroside, and sucrose. The aboveground parts contain coumarins and tannins (Krasnov et al. 1979; Kurlin and Zapesochnaya 1986; Peng et al. 1994; Kolesnikov and Gins 2001).

4915 **Ribes nigrum L. – Grossulariaceae (formerly in Saxifragaceae)**4916 **Synonyms:** *Botrycarpum nigrum* (L.) A. Rich., *Grossularia nigra* (L.) Rupr., *Ribes cyathiforme* Pojark., *Ribes olidum*
4917 Moench, *Ribes pauciflorum* Turcz. ex Ledeb.4918 **English name:** Black currant4919 **Russian name:** Смородина чёрная (Smorodina chyornaya)4920 **Uzbek name:** Kora smorodina, korakat4921 **Kyrgyz name:** Чыны карагат (Chyny karagat)4922 **Description:** Shrub up to 1.5 m tall. Young branches dull yellow, hairy; older branches brownish, almost glabrous. Leaves
4923 3–5-lobed, up to 12 cm wide, dotted with yellow glands beneath; lobes wide-triangular, margins serrate-dentate.
4924 Inflorescences drooping racemes, 3–8 cm long, 4–12-flowered. Flowers perfect, usually 5-merous, 5–7 mm wide, pedicel-
4925 late. Hypanthium campanulate, pink, greenish-red or greenish-yellow, pubescent, glandular. Calyx lobes reflexed, 3–4 mm
4926 long. Petals ovate, 2–3 mm long. Fruit a many-seeded, black berry, ca. 10 mm in diameter.4927 **Other distinguishing features:** Ovary inferior. Stamens inserted below rim of hypanthium, alternating with petals.4928 **Phenology:** Flowers in May–July, fruits in July.4929 **Reproduction:** By seeds and vegetatively.4930 **Distribution:** Ysyk-Kol and Naryn provinces of Kyrgyzstan; cultivated in Uzbekistan.4931 **Habitat:** In gorges and mountain river valleys up to 3,000 m elevation.4932 **The Population status:** Common, forming dense groups.4933 **Traditional use:** Fruits and leaves are used to treat anemia and edema, and as a light laxative. Fresh fruits are used to
4934 decrease blood pressure, to treat heart and liver diseases and atherosclerosis. A decoction of the young branches is drunk
4935 to treat children's diabetes and skin tuberculosis (Poludenny and Zhuravlev 2000). A decoction of the fruits is used as a
4936 diaphoretic, anti-inflammatory, and diuretic. The fresh juice is used to treat stomach and duodenum ulcers and gastritis
4937 with low stomach acidity; mixed with honey it is used to treat respiratory diseases. Leaves are used in a tea to treat skin
4938 and bladder diseases, kidney stones, rheumatism, common colds, and also as a diuretic (Khalmatov et al. 1984).4939 **Documented effects:** This species is used to treat infectious diseases, hemorrhagic diathesis, gastritis (particularly with low
4940 acidity), and is used as a tonic to treat the cardio-vascular system (Akopov 1990). An extract of the fruits has shown anti-
4941 viral activity against herpes and influenza A and B viruses (Knox et al. 2001, 2003; Suzutani et al. 2003). Proanthocyanidins
4942 isolated from the leaves exhibit anti-inflammatory effects in rats (Garbacki et al. 2004).4943 **Phytochemistry:** The fruits contain vitamins (ascorbic acid, B₁, and carotene), sugars, organic acids (citric and malic), pec-
4944 tins, anthocyanin compounds (cyanidin and delphinindin) and their glycosides, as well as quercetin and isoquercetin.
4945 Buds contain essential oil with d-pinene, l- and d-sabinene, d-caryophyllene, alcohol, and phenols. The leaves contain
4946 essential oil and ascorbic acid (Akopov 1990; Knox et al. 2001).

4947

Roemeria refracta DC. – Papaveraceae**Synonyms:** *Glaucium refractum* Steven ex DC., *Papaver refractum* (DC.) K.F. Gunther.**English name:** Spotted Asian poppy**Russian name:** Рёмерия отогнутая (Ryomeriya otognutaya)**Uzbek name:** Kizgaldok**Kyrgyz name:** Ийилген кызгалдак (Iyilgen kyzgaldak)**Description:** Annual herb, slightly hairy. Stem usually branched, rarely simple, 8–60 cm tall. Leaves bi- or tripinnatisect; basal and lower stem leaves petiolate; upper leaves alternate, sessile, pinnatisect. Flowers solitary, axillary and terminal. Petals 4, bright red, with a black spot at the base, broadly obovate, 2–4 cm long, 1.5–3.5 cm wide. Fruit a capsule, 4–6 cm long, 2–3 mm wide, glabrous. Seeds kidney-shaped, gray, pitted or reticulated.**Other distinguishing features:** Pedicel 10–12 cm long when in fruit. Fruits have 3–4 awn-like projections on the top, each 3–5 mm long.**Phenology:** Flowers in April-May, fruits in May-June.**Reproduction:** By seeds.**Distribution:** All regions of Uzbekistan and Kyrgyzstan.**Habitat:** The chul and adyr zones. On clay slopes of foothills and as a weed in crop fields and orchards.**Population status:** Common.**Traditional use:** The juice from the petals are used in a drink to treat children with sunstroke and as a wash to treat eye problems. A decoction of the plant is used to treat smallpox and fevers, and is applied externally to treat skin rashes (Khalmatov 1964). The dried petals are used as a sedative to treat cardiac and digestive organ pains (Khodzhimatov 1989).**Documented effects:** Pharmacological studies of the alkaloid roemerine showed that it has curare-like actions and an overdose can cause convulsions. The derivatives of roemerine also have curare-like and ganglio-blocking actions, but only for a short time. Roemerine has the ability to potentiate the effects of the analeptics corazol, cardiamine, caffeine, and strychnine. The alkaloid, and one of its derivatives, have strong antibacterial action against pathogenic microorganisms (Khodzhimatov 1989). (–)-roemerine isolated from the leaves of *Annona senegalensis*, was found to enhance the cytotoxic response mediated by vinblastine with multidrug-resistant human cancer cells in vitro (You et al. 1995).**Phytochemistry:** At the time of flowering the plants contain 0.2 % total alkaloids (roemerine, l-isoremerin, anonaine, liri-odenine, remrefidine, remrefine, l-ephedrine, d-pseudoephedrine, and l-mecambroline). The plant has also been found to contain a variety of additional alkaloids (Gozler et al. 1988, 1990a, b). The aboveground parts also contain tannins, organic acids, vitamin C, and sugar. The seeds contain a significant amount of fatty oils (Yunusov 1981; Khodzhimatov 1989).

4948

4949

4950

4951

4952

4953

4954

4955

4956

4957

4958

4959

4960

4961

4962

4963

4964

4965

4966

4967

4968

4969

4970

4971

4972

4973

4974

4975

4976

4977

4978

4979

4980



▲ **Ribes nigrum L.** Photos: *left*: Sergey Mayorov; *center and right*: Denis Mirin



◀ **Rheum maximowiczii Losinsk.** Photos: Evgeny Davkaev



◀ **Rhodiola linearifolia Boriss.** Photos: *left*: John B. Taft; *center and right*: Vladimir Epiktetov

▼ **Roemeria refracta DC.** Photos: *left*: Evgeny Davkaev; *center*: Alim Gaziev; *right*: Authors



- Rosa canina L. – Rosaceae** 4981
- Synonyms:** *Rosa ciliatosepala* Blocki, *Rosa sosnovskyi* Chrshan. 4982
- English name:** Dog rose 4983
- Russian name:** Шиповник собачий, Роза собачья (Shipovnik sobachiy, Roza sobach'ya) 4984
- Uzbek name:** Itburun 4985
- Kyrgyz name:** Ит мурун (It murun) 4986
- Description:** Shrub, up to 3 m tall. Stems arching with stout, flattened, hooked or rarely straight prickles. Leaves alternate, stipulate, pinnately compound with 5–7 leaflets; leaflets glabrous, elliptic, apex acute, margins sharply serrate. Inflorescence a corymb or rarely single flowered. Flowers 2–8 cm wide. Sepals 5, usually glabrous, reflexed, deciduous. Petals 5, bright pink, pale pink or white. Stamens many. Fruit a large hip (1.5–2.6 cm long), wide-ovoid or elongate-ovoid, smooth, bright or light-red, containing stony achenes. 4987–4991
- Other distinguishing features:** Stipules adnate to petiole for more than half their length. Outer sepals pinnatifid. 4992
- Phenology:** Flowers in June, fruits in August. 4993
- Reproduction:** By seeds. 4994
- Distribution:** Jalal-Abad and Osh provinces of Kyrgyzstan; Toshkent, Farg'ona, Samarqand, Qashqadaryo and Surxondaryo provinces of Uzbekistan. 4995–4996
- Habitat:** Along rivers and streams, on edges of deciduous forests, and in juniper stands. 4997
- Population status:** Common, found in small groups. 4998
- Traditional use:** A decoction of the petals, leaves, branches and roots is used to treat rheumatism, radiculitis, and stomach and heart ailments (Poludenny and Zhuravlev 2000). Decoction or tea of the fruits is used to treat scurvy, common colds, and as a diuretic. A decoction of the roots is used to treat liver and gastrointestinal tract diseases (Khalmatov et al. 1984; Khodzhimatov 1989). A decoction and infusion of the fruits is taken as an astringent (particularly for regular and bloody diarrhea), to treat fevers, intestinal infections, as a hemostatic for uterine bleeding, to improve the metabolism, and as a mouth wash for gum disease. The seeds are used as a diuretic and to treat kidney diseases. The powdered leaves are used to treat wounds and skin ulcers (Khalmatov et al. 1984). 4999–5005
- Documented effects:** Fruits are used as raw material for the preparation *Kholosas*, which has choleric activity and is used to treat cholecystitis and hepatitis (Khalmatov et al. 1984). Extracts of the fresh fruits exhibit high anti-ulcerogenic activity in rats (Gurbuz et al. 2003). A galactolipid, which is found in this species, has been shown to possess antitumor-promoting properties, as well as anti-inflammatory effects (Larsen et al. 2003). In a clinical trial, treatment with a standardized rose-hip powder showed significant reduction of symptoms associated with osteoarthritis (Warholm et al. 2003). 5006–5010
- Phytochemistry:** Fruits contain vitamin C, sugars, tannins, flavonoids (cyanidin-3-*O*-glucoside, phloridzin, isoquercitrin and glycosides of kaempferol, quercetin, taxifolin, and eriodictyol), conjugates of methyl gallate, pigments (carotene, lycopene, xanthophyll, etc.), pectins, pentosan and vitamins K₁, B₂, P and E. The seeds contain fatty oils and the flowers contain essential oil (Tolmachev 1976; Khalmatov et al. 1984; Hvattum 2002). 5011–5014

5016 **Rosa fedtschenkoana Regel – Rosaceae**

5017 **Synonyms:** *Rosa caraganifolia* Sumn., *Rosa coeruleifolia* Sumn., *Rosa epipsila* Sumn., *Rosa lavrenkoi* Sumn., *Rosa lip-*
5018 *schitzii* Sumn., *Rosa minusculifolia* Sumn., *Rosa oligosperma* Sumn.

5019 **English name:** Fedtschenko's rose

5020 **Russian name:** Шиповник Федченко, Роза Федченко (Shipovnik Fedchenko, Roza Fedchenko)

5021 **Uzbek name:** Namatak

5022 **Kyrgyz name:** Федченко ит мурун (Fedchenko it murun)

5023 **Description:** Shrub, 2–3(–6) m tall. Branches prickly; prickles yellowish, firm, straight, expanded at the base, up to 13 mm
5024 long. Leaves alternate, stipulate, pinnately compound with 5–9 leaflets, 3–4.5 cm long; leaflets 1–2.5 cm long, ovate to
5025 elongate-ovate, glabrous, margins serrate. Flowers 3–9 cm in diameter, solitary or in groups of 3–4. Sepals 5, lanceolate,
5026 pubescent above, glandular below. Petals 5, white or pink, broad-obovate. Fruit a fleshy, red hip, 2–5 cm long, elongate-
5027 ovoid, glandular-bristly, with persistent sepals, and containing stony achenes.

5028 **Other distinguishing features:** Leaflets glabrous. Hip to 5 cm long, densely glandular-bristly.

5029 **Phenology:** Flowers in June-August, fruits in July-September.

5030 **Reproduction:** By seeds.

5031 **Distribution:** All provinces of Kyrgyzstan; Tashkent, Farg'ona, Samarqand, Qashqadaryo and Surxondaryo provinces of
5032 Uzbekistan.

5033 **Habitat:** In forest glades, among bushes.

5034 **Population status:** Common, found in small groups.

5035 **Traditional use:** Fruits are widely used as a tonic, choleric, and as a remedy for scurvy. An infusion of the fruits with
5036 honey is used to treat upper respiratory colds and coughs. A decoction of the roots is drunk to treat diarrhea. A decoction
5037 of the leaves is taken to treat dysentery and as a diuretic (Khodzhimatov 1989). The hips from this and related species are
5038 used to prevent scurvy and avitaminosis, to treat arteriosclerosis, cholecystitis, hepatitis, and gastrointestinal diseases,
5039 particularly with reduced bile production (Altimishev 1991). Oil of rose is used externally to treat cracked and injured
5040 nipples of breast feeding women, bedsores, trophic ulcers of the shins, and dermatosis (Muravyova 1978). The fruits are
5041 used to treat lung tuberculosis, diphtheria, scarlet fever, flu, and sore throat (Zakordonets 1953).

5042 **Documented effects:** The fruits of this species are an official source of polyvitamins. Preparations (extracts, syrups, candies,
5043 pills, etc.) are used to treat hypo- and avitaminosis (particularly vitamin C deficiency) as well to treat diseases related to
5044 vitamin deficiency. The fruits are used as a component in an anti-asthmatic mixture. Oil from the seeds is used to treat
5045 burns, dermatosis, and radiation exposure. Ascorbic acid and an oil extract *Karotolin* (containing carotenoids, vitamin E,
5046 and linolic acid) are isolated from the pericarp. *Karotolin* is used to treat trophic skin ulcers, eczema, erythrodermia, and
5047 other skin diseases (Khalmatov et al. 1984).

5048 **Phytochemistry:** Fruits contain vitamins C, E, P, B₂, K₁, carotene, organic acids (malic and citric), sugars, flavonoids, pec-
5049 tins, and tannins. Seeds contain up to 37 % fatty oil (Tolmachev 1976; Khodzhimatov 1989). The flowers were found to
5050 contain glycosides of quercetin, kaempferol, cyanidin, and peonidin (Mikanagi et al. 1995).

5051

- Rubia tinctorum L. – Rubiaceae** 5052
- Synonyms:** *Rubia iberica* (Fisch. ex DC.) K. Koch. 5053
- English name:** Madder, common madder 5054
- Russian name:** Марена красильная (Marena krasil'naya) 5055
- Uzbek name:** Ruʻyan 5056
- Kyrgyz name:** Боечу марена (Boyechu marena) 5057
- Description:** Herbaceous perennial, with taproot and horizontal rhizomes. Stems 0.5–2 m tall, prostrate or climbing, 4-sided, 5058
with curved prickles. Leaves in whorls of 4 or 6, up to 9 cm long, up to 3 cm wide, narrow-ovate, apex acute. Inflorescences 5059
spreading complex panicles. Flowers small. Corollas yellow, 1–1.5 mm in diameter, 5-lobed. Stamens 5. Fruits berry-like, 5060
juicy, black with 2 hemispherical seeds. 5061
- Other distinguishing features:** Abaxial midvein and margins of leaves with curved prickles. 5062
- Phenology:** Flowers in June-August, fruits in August-September. 5063
- Reproduction:** By seeds and rhizomes. 5064
- Distribution:** The Osh province of Kyrgyzstan; Toshkent, Andijon, Farg'ona, Samarqand and Surxondaryo provinces of 5065
Uzbekistan. 5066
- Habitat:** Along canals, near springs, in orchards, and near rivers in tree-shrub forests. 5067
- Population status:** Common, found in small groups. 5068
- Traditional use:** In the past the underground parts of this species were used to treat rickets, constipation, jaundice, joint ail- 5069
ments, rheumatic back aches, and other diseases. Avicenna used a water extract as a strong diuretic to purify the liver and 5070
spleen and to treat spleen tumors. The extract, mixed with honey, was drunk to treat the sciatic nerves and paralysis. The 5071
underground parts, mixed with vinegar, were applied to treat fungal skin diseases and to remove skin spots. In recent times 5072
the underground parts have been used to treat kidney stones, gallstones, and gout, and also as a diuretic and laxative. In 5073
Central Asia, the roots mixed with honey is used to treat jaundice, to improve memory, and as a diuretic (Khalmatov et al. 5074
1984; Grinkevich 1991). 5075
- Documented effects:** Alcohol and water extracts of the roots inhibited the growth of *Aeromonas hydrophila*, *Bacillus mega-* 5076
terium, *Corynebacterium xenosis*, *Pseudomonas aeruginosa*, *Micrococcus luteus*, *Enterococcus faecalis*, and 5077
Staphylococcus aureus, but was not an effective inhibitor of *Escherichia coli* (Golcu et al. 2002). In experiments with rats 5078
that ate fresh roots decreased bladder and kidney stone formation was observed, but increased death rates were exhibited. 5079
In experiments with rabbits that were given root extracts orally, decreased calcium oxalate crystal formation in the kid- 5080
neys and hepatotoxicity was observed. Genotoxic effects were observed in bacterial and mammalian cell systems 5081
(Blumenthal 1998). 5082
- Phytochemistry:** The underground parts contain anthraglycosides and anthraquinone derivatives (ruberythric acid, galiosin, 5083
purpurin, rubiadin, mollugin, lucidin, etc.), organic acids (citric, malic, and tartaric), sugars, and traces of alkaloids. The 5084
young shoots contain the glycoside asperuloside. (Khalmatov et al. 1984; Gammerman et al. 1990; Kawasaki et al. 1992; 5085
Derksen et al. 2002). 5086

5088 **Rubus caesius L. – Rosaceae**5089 **Synonyms:** *Rubus psilophyllus* Nevski, *Rubus turkestanicus* (Regel) Pavlov.5090 **English name:** European dewberry5091 **Russian name:** Ежевика сизая (Ezhevika sizaya)5092 **Uzbek name:** Parmanchak, Maimunzhon5093 **Kyrgyz name:** Когултур кара булдукон (Kogultur kara buldurkon)5094 **Description:** Shrub, up to 1 m tall. Primocanes arching, glaucous, with stout, hooked prickles, rooting at the tip. Leaves
5095 trifoliate (basal leaves sometimes 5-foliate), stipulate, petiole prickly; leaflets broad-ovate, margins unevenly dentate,
5096 soft-pubescent beneath. Inflorescence racemiform or paniculiform. Flowers with 5 sepals and 5 white petals. Stamens and
5097 pistils many. Fruit an aggregate of drupelets, black or red, glaucous.5098 **Other distinguishing features:** Fruits separating from the stem with receptacle.5099 **Phenology:** Flowers in June-July, fruits in July-August.5100 **Reproduction:** By seeds and vegetatively.5101 **Distribution:** All provinces of Kyrgyzstan; Tashkent, Namangan, Farg'ona, Qashqadaryo and Surxondaryo provinces of
5102 Uzbekistan.5103 **Habitat:** Among shrubs, in forests and deforested areas, and along rivers and canals.5104 **Population status:** Common, found in dense groups.5105 **Traditional use:** Fresh fruits, infusion of the dried fruits, syrup or jam, or taken with tea, are widely used to quench the thirst,
5106 as a tonic, diaphoretic, diuretic, laxative, and sedative, as a remedy to increase the appetite, and to treat chronic gastritis
5107 and enterocolitis, stomach and duodenum ulcers, liver diseases, the flu, sore throats, pneumonia, stomatitis, dysentery,
5108 typhoid and fever. Water extracts, infusions or tea of the leaves and roots, is commonly used to treat stomach ulcers,
5109 chronic gastritis, and kidney stones (Nuraliev 1989). A decoction of the fruits, leaves, and branches is taken to treat cysti-
5110 titis, pyelitis, bronchitis, diabetes, urinary incontinence, eczema, vitiligo, psoriasis, fungal skin diseases, hair loss, and
5111 during menopause (Kurochkin 1998).5112 **Documented effects:** A decoction of the fruits is used as a source of vitamins, to improve digestion, and as a laxative and
5113 diaphoretic (Nuraliev 1989).5114 **Phytochemistry:** Fruits contain sugars, pectins, organic acids (citric, tartaric, malic, and salicylic), fiber, tannins, rutin, nico-
5115 tinic acid, flavonoids, and vitamins C, P, B₁, A, PP, E, and K. The leaves and branches contain flavonoids, tannins and
5116 ascorbic, malic, oxalic and lactic acids (Nuraliev 1989; Kurochkin 1998; Gudej and Tomczyk 2004).

5117



▲ **Rosa canina L.** Photos: *top left:* Alim Gaziev; *top right, center and bottom:* Maxim Kucherov



▲ **Rosa fedtschenkoana Regel** Photos: *top:* Alim Gaziev; *bottom:* Vladimir Epiktetov



▲ **Rubia tinctorum L.** Photos: Maxim Kucherov

▼ **Rubus caesius L.** Photos: *left and right:* Maxim Kucherov; *center:* Sergey Appolonov



5118 **Rubus idaeus L. – Rosaceae**5119 **Synonyms:** None5120 **English name:** Red raspberry5121 **Russian name:** Малина обыкновенная (Malina obyknovennaya)5122 **Uzbek name:** Parmanchak, Malina5123 **Kyrgyz name:** Кадимки дан куурай (Kadimki dan kuuray)

5124 **Description:** Shrub, up to 80–200 cm tall. Primocanes green, glaucous, with thin, straight prickles. Floricanes yellowish or
5125 green, slightly woody. Leaves odd-pinnate with 3–5(–7) leaflets, stipules thread-like; leaflets white tomentose below,
5126 margin unevenly serrate. Flowers in few-flowered racemes in corymbiform-paniculate inflorescences. Sepals 5, reflexed,
5127 grayish-green. Petals 5, white. Stamens and pistils many. Fruit a red (raspberry) aggregate of drupelets .

5128 **Other distinguishing features:** Fruits separating from the receptacle.5129 **Phenology:** Flowers in June, fruits in July-August.5130 **Reproduction:** By seeds and vegetatively.5131 **Distribution:** Jalal-Abad, Ysyk-Kol, Osh, and Chuy provinces of Kyrgyzstan; cultivated in Uzbekistan.5132 **Habitat:** In meadows, along rivers, and in deforested areas in the shrub and forest belt of mountains.5133 **Population status:** Common, found in dense groups.

5134 **Traditional use:** Fruits are used as a diaphoretic and antipyretic. Leaves are used as an astringent and hemostatic, and to treat
5135 diarrhea. A decoction and infusion of the leaves is recommended as a cough remedy, and is gargled to treat sore throats.
5136 An infusion of the leaves and flowers is used to treat hemorrhoids and gynecological conditions. A paste of the fresh
5137 leaves is used to treat acne and rashes. A decoction of the flowers is used as a wash for acne, erysipelas, and conjunctivitis
5138 (Khalmatov et al. 1984). The fresh fruits are considered to have sobering effects for drunkenness (Kurochkin 1998).

5139 **Documented effects:** Preparations from raspberries improve stomach and intestine function, have antiseptic, analgesic,
5140 antipyretic, expectorant, anti-inflammatory, and anti-emetic properties, and improve metabolism (Maznev 2004). Extracts
5141 of the fruits have antioxidant effects and exhibit antimicrobial properties (Kahkonen et al. 1999; Rauha et al. 2000;
5142 Puupponen-Pimia et al. 2001). Components of raspberry leaf extract exhibited relaxant activity in an in vitro gastrointes-
5143 tinal tissue (Rojas-Vera et al. 2002).

5144 **Phytochemistry:** Fruits contain vitamins (C, B₁, B₂, B₆, PP, E, and A), organic acids (citric, malic, salicylic, tartaric, formic,
5145 and capronic), ellagic acid and its derivatives, sugars, pectins, minerals, essential oil, anthocyanins, flavonoids, and tannins.
5146 Seeds contain fatty oils, sitosterin, tocopherols, neutral lipids, phospholipids, and free fatty acids. The main fatty acids of
5147 crude oil were 18:2 (54.5 %), 18:3 (29.1 %), 18:1 (12 %), and 16:0 (2.7 %; Tolmachev 1976; Khalmatov et al. 1984;
5148 Kurochkin 1998; Oomah et al. 2000; Zafrilla et al. 2001).

5149

- Rumex confertus Willd. – Polygonaceae** 5150
- Synonyms:** *Rumex alpinus* L. var. *subcalligerus* Boiss. 5151
- English name:** Russian dock 5152
- Russian name:** Щавель конский (Shchavel' konskiy) 5153
- Uzbek name:** Ot quloq 5154
- Kyrgyz name:** Ат кулак (At kulak) 5155
- Description:** Herbaceous perennial, 60–150 cm tall, with a thick root. Stems erect, striated. Blades of basal leaves elongated triangular-oval, 15–25 cm long, 6–12 cm wide, apex obtuse, cordate, margins sinuate, abaxial side with stiff hairs towards veins; petiole equal to or exceeding the length of the blade. Cauline leaves smaller, acute, oval-lanceolate, short-lanceolate. Inflorescence terminal, narrow-cylindrical or wide-paniculiform, composed of pedicellate flowers densely arranged in multiflorous whorls. Perianth with 6 tepals, 6–9 mm long, 6–11 mm wide. Fruit a triquetrous achene, 3–5 mm long, 1.5–2.5 mm wide. 5156–5161
- Other distinguishing features:** Ocreae mostly deciduous. 5162
- Phenology:** Flowers and fruits in May-June. 5163
- Reproduction:** By seeds. 5164
- Distribution:** Toshkent and Qashqadaryo provinces of Uzbekistan; not found in Kyrgyzstan. 5165
- Habitat:** The adyr and tau zones. River banks and grassy slopes and as a weed in cultivated fields. 5166
- Population status:** Common, sometimes makes dense populations. 5167
- Traditional use:** This plant has been used for treatment of multiple diseases such as scabies and scurvy, and as an astringent for diarrhea. A decoction of roots and leaves is used to treat skin disorders (fungal skin diseases and rashes), ulcers, and wounds (Seredin and Sokolov 1969; Khalmatov et al. 1984). 5168–5170
- Documented effects:** Small doses of preparations (infusions and extracts) have astringent effects and in big doses, purgative effects. Currently they are recommended to improve intestinal function. They are also used for anemia with simultaneous gastrointestinal tract dysfunction, and for colitis, hemorrhagic enterocolitis, hemorrhagic colitis, and child's diarrhea (Seredin and Sokolov 1969). Experiments show that a preparation of this species acts as a vermifuge and has hemostatic and hypotensive ability (Sokolov and Zamotaev 1985). An extract of the plant exhibited cytotoxic effects against human lymphoblastoid cells in vitro (Spiridonov et al. 2005). Chrysophanic acid, isolated from *Dianella longifolia*, has been found to inhibit the replication of poliovirus types 2 and 3 in vitro (Semple et al. 2001). In vitro, emodin inhibits tyrosine kinase, an enzyme overexpressed in certain breast cancer cells. The combination of emodin and paclitaxel synergistically inhibited tumor growth and prolonged survival in mice (Zhang et al. 1999). 5171–5179
- Phytochemistry:** Underground organs contain tannins (ellagic acid, phloroglucinol, and caffeic acid), flavonoids (nepodin, chrysophanic acid, emodin, etc.), resins, essential oils, and calcium oxalate. Leaves contain flavone glycosides (hyperoside and rutin), carotene, vitamin C, and calcium oxalate (Seredin and Sokolov 1969; Mukhamed'yarova and Chumbalov 1979). 5180–5183

5185 **Rumex tianschanicus** Losinsk. – Polygonaceae5186 **Synonyms:** None5187 **English name:** Unknown5188 **Russian name:** Щавель Тяньшанский (Shchavel' tyan'shanskiy)5189 **Uzbek name:** Unknown5190 **Kyrgyz name:** Ат кулак (At kulak)

5191 **Description:** Herbaceous perennial. Stem single, up to 2 m tall, thick, branched, hollow, largely striated. Basal leaves wide-
5192 ovate, 17–25 cm long, up to 15 cm wide, apex acute, base cordate, margin undulate, short-petiolate; stem leaves smaller;
5193 ocreae membranous, falling off early. Inflorescence paniculate. Flowers with 6 tepals arranged in 2 whorls. Fruits 3-sided
5194 achenes, pointed, light-brown, 2 mm long.

5195 **Other distinguishing features:** Each tepal with a prominent vein.5196 **Phenology:** Flowering and fruits in May-June.5197 **Reproduction:** By seeds.5198 **Distribution:** Chuy province of Kyrgyzstan; not found in the flora of Uzbekistan.5199 **Habitat:** In rivers valleys and orchards.5200 **Population status:** Common, found in small groups.5201 **Traditional use:** In Uzbekistan, a fresh leaf is applied externally to an abscess to provoke maturation (Sezik et al. 2004).

5202 **Documented effects:** The underground parts have slight antitumor activity. An infusion and alcohol extract are used to treat
5203 pellagra and dyspepsia. Fruits are used to treat dyspepsia in children (Belodubrovskaya et al. 2002).

5204 **Phytochemistry:** All parts of the plant contain phenolcarboxylic acids, flavonoids, and catechins. The seeds contain fatty oil
5205 (Plant Resources of the USSR 1985). The roots also contain sugars, inulin, organic acids, tannins, anthraquinones, and
5206 leucoanthocyanides. The leaves contain vitamins (C, P, K), carotenoids, and tannins (Belodubrovskaya et al. 2002;
5207 Kharlamova 2007).

5208

- Salvia deserta Schangin – Lamiaceae** 5209
- Synonyms:** *Salvia jailicola* Klokov, *Salvia moldavica* Klokov, some considered *S. deserta* a synonym of *Salvia nemorosa* L. 5210
- English name:** Unknown 5211
- Russian name:** Шалфей пустынный (Shalfey pustynnyy) 5212
- Uzbek name:** Mavrak 5213
- Kyrgyz name:** Чол шалфейи (Chol shalfeyi) 5214
- Description:** Perennial subshrub. Stems erect, simple or branched, densely curly pubescent, 60–80 cm tall. Leaves opposite, ovate to ovate-lanceolate, apex acute to acuminate, base cordate, adaxial side dark green, abaxial side gray-pubescent, margin crenate-serrate, petiolate. Inflorescences terminal, racemiform, composed of verticillasters with 4–6 flowers, pubescent. Bracts broadly ovate, 4–6 mm long, purple-red. Flowers short pedicellate. Calyx 5–6 mm long, 2-lipped. Corolla 9–10 mm long, 2-lipped, blue-purple to violet. Nutlets rounded-triangular, black, 1.5 mm long. 5215–5219
- Other distinguishing features:** Upper lip of calyx shorter than lower lip. 5220
- Phenology:** Flowers in May–August, fruits in June–September. 5221
- Reproduction:** Only by seeds. 5222
- Distribution:** Toshkent, Farg’ona, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan. 5223
- Habitat:** The adyr and tau zones. As a weed in orchards, on grassy slopes and as a weed in oases. 5224
- Population status:** Common, often found as single individuals. 5225
- Traditional use:** A decoction of leaves and flowers is used for cardiac neurosis and neurasthenia, to increase appetite, as a gargle to treat sore throat, and for intestinal infections and fever. A powder of roasted seeds is recommended for dysentery and heart palpitations. Ground fruits mixed with oil are used to heal wounds. Preparations of *Salvia deserta* are used in the same way as the preparations of the aboveground parts of *Salvia sclarea* (Khalmatov and Kosimov 1992; Gammerman et al. 1990). 5226–5230
- Documented effects:** Essential oils from this species are used in the pharmaceutical industry to add an aroma to drugs and in the fragrance industry as an aroma fixative (Khalmatov and Kosimov 1992; Gammerman et al. 1990). Water and MeOH extracts of the plant strongly inhibited aldose reductase activity, an enzyme associated with diabetic complications (Kasimu et al. 1998). Compounds from the plant were found to inhibit prolyl endopeptidase (PEP), an enzyme thought to be involved with learning and memory processes, and the inhibition of which may produce anti-amnesic effects (Tezuka et al. 1999). 5231–5236
- Phytochemistry:** Flowering plant tops contain 0.01–0.04 % essential oil (similar to *Salvia sclarea* in composition) with a pleasant aroma. In leaves there were 47 mg% vitamin C, and seeds contained up to 19 % oil (Khalmatov 1964). Triterpenoids, including ursane, oleanane, and lupane derivatives were isolated from the aboveground parts (Savona et al. 1987). The roots were found to contain a number of caffeic acid derivatives (rosmarinic acid, lithospermic acid B, etc.), diterpenes (royleanone, ferruginol, taxodione, etc.), and the steroid daucosterol (Tezuka et al. 1998). 5237–5241

▼ **Salvia deserta Schangin** Photos: Evgeny Davkaev



▼ **Rumex tianschanicus** Losinsk. Photos: Vladimir Epiktetov



▼ **Rubus idaeus L.** Photos: *top and bottom:* Dmitri Oreshkin; *center:* Sergey Mayorov



▼ **Rumex confertus Willd.** Photos: *top:* Sergey Appolonov; *bottom:* Anatoly Lisitzyn



- Salvia sclarea L. – Lamiaceae** 5243
- Synonyms:** *Salvia asperata* Falc. ex Benth., *Salvia pamirica* Gand. 5244
- English name:** Clary, Clary sage 5245
- Russian name:** Шалфей мускатный (Shalfey muskatnyy) 5246
- Uzbek name:** Mavrak, Marmarak, Khutan 5247
- Kyrgyz name:** Мускат шалфейи (Muskat shalfeyi) 5248
- Description:** Herbaceous perennial, with taproot. Stems few, erect, 20–150 cm tall, 4-sided, hairy, branched, upper portions glandular. Leaves opposite, simple, 7–30 cm long, 3–22 cm wide, rugose, ovate or oblong-ovate, margins unevenly dentate. Inflorescences verticillasters in panicles. Bracts round-ovate, 1–3 cm long, often whitish with red-purple tips. Calyx tubular, 2-lipped, upper lip 3-lobed, lower lip 2-lobed. Corolla 2-lipped, pink, lilac or white. Fruits are ellipsoid nutlets, brown, 2–3 mm long. 5249
5250
5251
5252
5253
- Other distinguishing features:** Upper lip of corolla arching, longer than tube and extending past lower lip. 5254
- Phenology:** Flowers in June-August, fruits in August-September. 5255
- Reproduction:** By seeds. 5256
- Distribution:** Osh, Jalal-Abad, Talas, and Chuy provinces of Kyrgyzstan; Tashkent, Andijon, Farg'ona, Samarqand and Surxondaryo provinces of Uzbekistan. 5257
5258
- Habitat:** On the slopes of mountains, along high mountain rivers, and in fallow fields and orchards. 5259
- Population status:** Common, forming dense groups. 5260
- Traditional use:** The aboveground parts are used to treat fevers, stomach ulcers, headaches, epilepsy, to improvement digestion, and as an antiseptic. It is used in bathes to treat bladder diseases, polyarthritis, osteomyelitis, deforming arthrosis, and trophic ulcers. The leaves are used as a antispasmodic and anti-inflammatory. A decoction of the leaves is used as a mouth wash for acute respiratory diseases and throat illnesses, periostitis and is applied externally to purulent wounds and furuncles. The decoction of the leaves and inflorescences are used to treat tachycardia and asthenia (Plant Resources of the USSR 1991). 5261
5262
5263
5264
5265
5266
- Documented effects:** Clinical studies showed that an ointment (with 5–20 % plant extract) was highly effective in treating psoriasis (Khalmatov et al. 1984). An emulsion of the oil was successfully used to treat osteomyelitis, varicose veins, paronychia, burns, and other diseases (Sklarovsky 1972). Extracts of the roots show antibacterial activity and are used in antibacterial preparations (Gammerman et al. 1990). In experiments, a tincture of the herb increased respiration and arterial pressure and had diuretic properties. The tincture affected an isolated frog heart in a similar manner as camphor. An infusion of the herb is used in stomatology to treat caries, pulpitis, periodontitis, and catarrhal gingivitis (Plant Resources of the USSR 1991). A number of the diterpenoids and sesquiterpenes isolated from the plant were found to be active against *Staphylococcus aureus* and *Candida albicans* and caryophyllene oxide showed activity against *Proteus mirabilis* (Ulubelen et al. 1994). 5267
5268
5269
5270
5271
5272
5273
5274
5275
- Phytochemistry:** Aboveground parts contain essential oil (linalyl-acetate, linalool, ocimene, myrcene, cedrene, nerolidol, sclareol, etc.), coumarins, flavonoids, saponins, and trace alkaloids. Seeds contains drying fatty oil which contains oleo-lic, linoleic, linolenic, arachidic, behenic, lignoceric, and cerotinic acids, pigments (carotene and chlorophyll), and stear- ins. The roots contains quinones (tanshinone, isotanshinone, oxytanshinone, etc.; Khalmatov et al. 1984; Khodzhimatov 1989; Gammerman et al. 1990). An extract of the whole plant contained flavonoids (apigenin, luteolin and their deriva- tives, etc.), diterpenes (sclareol, manool, ferruginol, etc.), sesquiterpenes (caryophyllene oxide and spathulenol), alpha- myrin, and β -sitosterol (Ulubelen et al. 1994). 5276
5277
5278
5279
5280
5281
5282

5284 **Sanguisorba officinalis L. – Rosaceae**5285 **Synonyms:** *Sanguisorba glandulosa* Kom.5286 **English name:** Great burnet, Official burnet5287 **Russian name:** Кровохлёбка аптечная (Krovokhlyobka aptechnaya)5288 **Uzbek name:** Sangvizorba, Dorivor kukat, Dorivor krovoklebka5289 **Kyrgyz name:** Дары кансоргуч (Dary kansorguch)5290 **Description:** Herbaceous perennial, with thick rhizome. Stems up to 1 m tall, single or few, hollow, ribbed, branched above.5291 Leaves alternate, odd-pinnately compound, glabrous, stipulate; leaflets elongate-ovate, margins serrate; lower leaves
5292 large, long-petiolate; upper leaves sessile. Inflorescences ellipsoid to cylindrical heads, 1–3 cm long. Sepals 4, petaloid,
5293 purple-brown. Petals lacking. Fruit a brown achene.5294 **Other distinguishing features:** Stamens 4, equal in length to sepals.5295 **Phenology:** Flowers in June, fruits in August.5296 **Reproduction:** By seeds.5297 **Distribution:** Chuy province of Kyrgyzstan; not found in the flora of Uzbekistan.5298 **Habitat:** Along rivers, in the forest-meadow mountain belt, among shrubs, on grassy slopes, and in bogs.5299 **Population status:** Rare, found in small groups.5300 **Traditional use:** Used in folk medicine as an astringent and hemostatic. Used to treat gastrointestinal diseases, tuberculosis,
5301 hemoptysis, and uterine bleeding. Used externally to heal wounds (Khalmatov et al. 1984; Akopov 1990; Grinkevitch
5302 1991).5303 **Documented effects:** The species is used to treat upper respiratory illnesses, hemorrhoids, scurvy and gingivitis, and as an
5304 expectorant and astringent (Kovaleva 1971). The decoction of this species is used externally to treat wounds and skin
5305 ulcers and as a douche to treat cervical erosion (Akopov 1990). A decoction of the roots has antimicrobial effects against
5306 *Trichomonas*, *Candida* sp., and *Giardia lamblia* (Zavrazhanov et al. 1977). Two triterpene glycosides isolated from the
5307 roots were found to have cytotoxic activity against human carcinoma cells in vitro (Mimaki et al. 2001). Both in vitro and
5308 in vivo, a triterpene glycoside isolated from the roots diminished tumor necrosis factor-alpha production (Cho et al.
5309 2006).5310 **Phytochemistry:** Underground parts contain tannins (pyrogallol groups), saponins, sterols, acids (gallic, ellagic, oxalic,
5311 and ascorbic), a number of triterpenes and triterpene glycosides, gallotannins, carotene, starch, pigments, phytoncides,
5312 essential oil, and micro- and macroelements (Kurochkin 1998; Mimaki et al. 2001; Liu et al. 2005; Cho et al. 2006).

5313

Scabiosa songarica Schrenk – Dipsacaceae	5314
Synonyms: <i>Trochocephalus songaricus</i> (Schrenk) Á. Löve & D. Löve.	5315
English name: Unknown	5316
Russian name: Скабиоза джунгарская (Skabioza dzhungarskaya)	5317
Uzbek name: Zhoongor scabiozasi	5318
Kyrgyz name: Жунгар бешилик чобу (Zhungar beshilik chobu)	5319
Description: Herbaceous perennial, with woody roots. Stems 20–65(–100) cm tall, with short hairs. Basal and lower stem leaves petiolate; lower stem leaves lanceolate, entire or slightly pinnate; upper stem leaves opposite, lanceolate, hairy, pinnatifid with a larger apical lobe. Inflorescence a head, 2.5–3 cm in diameter; involucre bracts narrow-lanceolate, densely long-bristled; involucre expanded above into corona. Marginal flowers up to 2 cm long. Corolla yellow-violet, hairy outside. Fruits bristly achenes, adnate to the involucre and crowned by the calyx.	5320 5321 5322 5323 5324
Other distinguishing features: Calyx teeth twice as long as corolla.	5325
Phenology: Flowers in June-July, fruits in August-September.	5326
Reproduction: By seeds and vegetatively.	5327
Distribution: All provinces of Kyrgyzstan; Tashkent, Andijan, Farg'ona, Samarqand, and Surxondaryo provinces of Uzbekistan.	5328 5329
Habitat: On foothills, in steppes with a wide diversity of grass species.	5330
Population status: Common, forming dense groups.	5331
Traditional use: A decoction of the herb is used to treat respiratory infections and common colds (Sinitsin 1959).	5332
Documented effects: In experiments on animals, the total saponins isolated from the roots and the preparation <i>Zongorozid</i> caused a significant decrease in arterial pressure, increased resistance to hypoxia, and had sedative effects. In experiments with dogs, the preparation <i>Zongorozid</i> increased the sodium in erythrocytes and reduced potassium in blood plasma as well as in erythrocytes. A one time dose of the preparation has blood coagulating effects but multiple applications, over 5–7 days, have better effects. The effects include an increase in tolerance to heparin, reduction of prothrombin time and fibrinolytic activity, increase in fibrinogen content (up to 45 %), and an increase of the adhesion index with an increase in blood coagulation potential (Alimbaeva et al. 1986).	5333 5334 5335 5336 5337 5338 5339
Phytochemistry: Roots contain organic acids, saponins (17 triterpene glycosides and oleanolic acid derivatives), steroids, alkaloids, vitamin C, flavonoids, coumarins, and tannins. The aboveground parts contain organic acids, saponins, alkaloids, phenolcarboxylic acids, coumarins, and flavonoids (Alimbaeva and Akimaliev 1975).	5340 5341 5342

5344 ***Serratula sogdiana* Bunge – Asteraceae**

5345 **Synonyms:** *Serratula alata* C.A. Mey. ex Rupr., *Serratula dissecta* var. *asperula* Regel & Herder, *Serratula trautvetteri-*
5346 *ana* Regel & Schmalh.

5347 **English name:** Unknown

5348 **Russian name:** Серпуха согдийская (*Serpukha sogdiyskaya*)

5349 **Uzbek name:** Unknown

5350 **Kyrgyz name:** Согдия чогойносу (*Sogdiya chogoynosu*)

5351 **Description:** Herbaceous perennial with a thick, woody, branching rhizome and string-like roots. Stems erect, 25–55 cm tall,
5352 ribbed, foliaceous, with long, appressed, straight, thin branches. Basal and lower leaves thin-coriaceous, oblong-lyrate,
5353 ca. 8 cm long, lower half of blade incised-toothed, upper half entire, petiolate with stipule-like auricles at the base; middle
5354 leaves and leaves on branches linear-lanceolate, ca. 3 cm long, some deeply incised, toothed; upper leaves becoming very
5355 reduced and spinescent. Inflorescences capitula, mostly solitary, 12–15 mm wide, 25 mm long; peduncles with several
5356 small spiny leaves; involucre bracts coriaceous, yellowish-green, imbricate, short-hairy on the outside, gradually tapering
5357 into pointed tip. Corollas pink or purple, ca. 1.5 cm long with linear lobes, protruding well past involucre bracts. Fruits
5358 oblong achenes, ca. 5 mm long, dentate-edged on top, reddish-brown; pappus with dense, plumose bristles, deciduous.

5359 **Other distinguishing features:** Receptacle with smooth bristles that are ca. 1 cm long.

5360 **Phenology:** Flowers in June–August, fruits in August–September.

5361 **Reproduction:** By seeds.

5362 **Distribution:** Farg'ona province of Uzbekistan; lower belt of Alai mountain range (mountains-Kiziltau, Galtin, Mashalang,
5363 Katrantau, Hurdjuntau); Osh, Chuy and Jalal-Abad provinces of Kyrgyzstan.

5364 **Habitat:** The lower tau zone. On gentle, stony slopes.

5365 **Population status:** Uncommon, sometimes found in small populations.

5366 **Traditional use:** Decoctions and infusions of plants in the genus *Serratula* are used to heal wounds, to treat anemia, as a
5367 restorative for weakness due to fever and as a treatment for liver diseases (Zavrazhanov et al. 1972).

5368 **Documented effects:** Ecdysterone and extracts of *Serratula sogdiana* L. have anabolic activity as well as the ability to keep
5369 nitrogenous compounds in the organism and assist in acceleration of protein synthesis (Syrov and Kurmukov 1975a, b, c;
5370 Saatov et al. 1999).

5371 **Phytochemistry:** The phytoecdysteroids ecdysterone, viticosterone, and sogdisterone were identified in extracts of the
5372 inflorescences (Zatsny et al. 1971, 1973a, b; Saatov et al. 1999).

5373



▲ *Sanguisorba officinalis* L. Photos: *left*: Denis A. Davydov; *center and right*: Rostislav Lezhoyev



▲ *Scabiosa songarica* Schrenk

Photos: *left and center*: Evgeny Davkaev; *right*: Alim Gaziev



◀ *Salvia sclarea* L.

Photos: *left and center*: Alim Gaziev; *right*: Evgeny Davkaev

▼ *Serratula sogdiana* Bunge

Photos: Evgeny Davkaev



5374 **Silybum marianum (L.) Gaertn. – Asteraceae**5375 **Synonyms:** *Carduus marianus* L.5376 **English name:** Blessed milk thistle, Milk thistle5377 **Russian name:** Расторопша пятнистая (Rastoropsha pyatnistaya)5378 **Uzbek name:** Unknown5379 **Kyrgyz name:** Unknown

5380 **Description:** Annual or biennial 0.3–3 m, usually ~1.5 m tall. Stem erect, usually branched, grooved, farinose, thinly arach-
5381 noid-hairy, foliaceous. Leaves green with large white spots and veins, oblanceolate to elliptical, pinnately lobed, petiolate;
5382 lobes prickly or prickly toothed; basal leaves up to 80 cm long and 30 cm wide, forming a rosette; upper leaves reduced,
5383 sessile, clasping, prickly lobed. Inflorescences terminal capitula, nodding, oblong or globose, 3–6 cm in diameter, solitary
5384 with slender peduncles. Involucral bracts imbricate; outer and middle bracts up to 3 cm long, spreading, stiff, erect,
5385 spinescent with 4–6 spines on the margin. Flowers discoid, 2.5–3.5 cm long, pink, purple or white, numerous; tube long,
5386 slender, throat abruptly wider, corolla lobes linear. Fruits elliptical or obovate achenes, ca. 6 mm long, 3 mm wide, slightly
5387 flattened, brownish-black and sometimes white spotted, glabrous; pappus composed of a deciduous ring of minutely
5388 barbed bristles, ca. 2 cm long.

5389 **Other distinguishing features:** Receptacles flat and covered with whitish bristles.5390 **Phenology:** Flowers in April-May, fruits May-June.5391 **Reproduction:** Only by seeds.5392 **Distribution:** Qashqadaryo and Surxondaryo provinces of Uzbekistan; not found in Kyrgyzstan.5393 **Habitat:** The chul and adyr zones. A weed growing along roads and edges of agricultural fields.5394 **Population status:** Uncommon, sometimes makes dense populations numbering up to 40 individuals.

5395 **Traditional use:** The seeds are used to treat jaundice, hepatitis, chronic coughing and hemoptysis, gall-stones and
5396 inflammation of the gall bladder and bile duct, liver and spleen diseases, fevers, hemorrhoids, and other diseases. Juice
5397 from the leaves is drunk as a choleric and diuretic and to treat colitis and constipation. A decoction of the root is drunk
5398 to treat stomach catarrh. Currently, an alcohol-water extraction of the seeds is used to treat liver diseases (Khalmatov
5399 1964; Khodzhimatov 1989).

5400 **Documented effects:** Preparations such as *Karsil*, *Legalon*, and *Silimarin* are used in modern medicine to restore liver mem-
5401 branes and to treat bile-duct and gall-bladder diseases (Gammerman et al. 1990). A variety of experiments have shown
5402 that silymarin increases liver regeneration after damage caused by liver diseases. Similar effects were found in kidney
5403 cells in vitro (Sonnenbichler et al. 1999).

5404 **Phytochemistry:** The aboveground parts contain flavonoids and fumaric acid. Seeds contain 0.08 % essential oil, vitamin K,
5405 mucilage, resins, biogenic amines (thiramine and histamine), trace alkaloids, saponins, flavonoids, and flavolignans
5406 (isosilibinin, silibinin, silicristin and silidianin; Khodzhimatov 1989; Gammerman et al. 1990; Kurochkin 1998;
5407 Sonnenbichler et al. 1999). The seed oil is rich in linoleic and oleic acids and contains 5 major triacylglycerols. Campesterol,
5408 5-stigmasterol, β -sitosterol, 7-stigmasterol, avenasterol, and spinasterol were also detected in the seed oil (El-Mallah
5409 et al. 2003).

Sorbus tianschanica Rupr. – Rosaceae	5411
Synonyms: <i>Pyrus tianschanica</i> (Rupr.) Franch.	5412
English name: Tian Shan mountain ash, Tian Shan rowan	5413
Russian name: Рябина тьяншанская (Ryabina tyan'shanskaya)	5414
Uzbek name: Kizilchetan	5415
Kyrgyz name: Тяньшань четини (Tyan'shan' chetini)	5416
Description: Tree, 3–5 m tall. Branches brown, with lenticels; young shoots reddish-brown. Leaves alternate, odd-pinnately compound with 13–15 leaflets, 12–18 cm long (including rachis), stipules membranaceous; leaflets ovate-lanceolate, glabrous, margins serrate. Inflorescences loose clusters, many-flowered. Flowers 1.5–2 cm wide, hypanthium campanulate. Sepals 5, triangular. Petals 5, ovate or elliptic, white. Stamens 15–20. Styles 3–5. Fruits nearly round pomes, 10–12 mm wide, scarlet to dark-red, glaucous.	5417 5418 5419 5420 5421
Other distinguishing features: Buds white, pubescent.	5422
Phenology: Flowers in June-July, fruits in August-September.	5423
Reproduction: By seeds.	5424
Distribution: All provinces of Kyrgyzstan; Toshkent and Samarqand provinces of Uzbekistan.	5425
Habitat: In the upper forest-shrub belt of mountains (2,000–3,200 m).	5426
Population status: Common, found growing as single plants.	5427
Traditional use: The fruits of this species are used to treat hepatitis and cholecystitis (Sumnevich 1942).	5428
Documented effects: Fruits and seeds have antibacterial properties. An alcohol extract and fatty oil are used to treat paratyphoid fever (Aitbaeva 1972).	5429 5430
Phytochemistry: All parts of plant contain phenolcarbonic acids, flavonoids, and catechins. The seeds contain fatty oil. Fruits contain ascorbic acid, vitamin A, tannins, and carotene (Zapesochnaya et al. 1973; Dzhangaliev et al. 2003).	5431 5432

5434 **Sphaerophysa salsula (Pall.) DC. – Fabaceae**5435 **Synonyms:** *Phaca salsula* Pall.5436 **English name:** Alkali swainsonpea, Austrian Peaweed5437 **Russian name:** Сферофиза солонцовая, Круглоплодник солончаковый (Sferofiza solontsovaya, Krugloplodnik
5438 solonchakovyy)5439 **Uzbek name:** Shildir bosh5440 **Kyrgyz name:** Шорчул сферофиза (Shorchul sferofiza)5441 **Description:** Herbaceous perennial with scattered short, appressed hairs. Stems erect, 30–70 cm tall, with appressed
5442 branches. Leaves alternate, odd-pinnate, 4–9.5 cm long; leaflets in 6–10 pairs, elliptic to oblong-elliptic, mucronulate.
5443 Inflorescences racemes, 4–10 cm long. Flowers numerous, short-pedicellate. Calyx campanulate, 4–5 mm long with 5
5444 short teeth. Corolla papilionaceous, brick-red. Stamens diadelphous. Fruits swollen legumes, wide-oblong, 2.5–3.5 cm
5445 long, 1.8–2 cm wide, papery-membranous, glabrous or with scattered hairs, stipitate. Seeds ~1.5 mm long, round to kid-
5446 ney-shaped, brown, dull.5447 **Other distinguishing features:** Legume many-seeded, indehiscent.5448 **Phenology:** Flowers in May-June, fruits in July-August.5449 **Reproduction:** By seeds and vegetatively by rhizomes.5450 **Distribution:** Nearly all provinces of Uzbekistan; Talas and Jalal-Abad provinces of Kyrgyzstan.5451 **Habitat:** The chul zone. Wet, salty places, river banks, and tugais.5452 **Population status:** Uncommon, usually found as small populations.5453 **Traditional use:** An infusion of the herb is used as a hemostatic after childbirth and to treat uterine atonia (Akopov 1981).

5454 It is used for the treatment of hypertension in China (Ma et al. 2002b).

5455 **Documented effects:** The alkaloid spherophysine, which was isolated from the aboveground plant parts, has hypotensive
5456 activity and effects uterine action. In the form of a benzoic-acid salt, it is used for essential hypertension of the first and
5457 second degrees. Spherophysine is used for arterial hypertension, weak birthing activity (labor difficulties), and post natal
5458 bleeding (Sokolov and Zamotaev 1989). A stilbene isolated from the plant was synthesized and tested for antioxidant
5459 activity and showed superior antioxidative activity when compared to the well-known antioxidants resveratrol, vitamin C
5460 and butylated hydroxyanisole (BHA) (Venkateswarlu et al. 2003). Additionally, synthesized stilbenes based on naturally
5461 occurring compounds were active against leukemia and lymphoma cell lines (Tolomeo et al. 2005).5462 **Phytochemistry:** Aboveground plant parts contain 0.4 % total alkaloids, from which spherophysine, spherosine, and
5463 saponins with hemolytic index of 1:40 have been isolated (Sokolov and Zamotaev 1989). Isoflavans, lignans, coumarins,
5464 flavonoids, and sterols have also been isolated from the plant (Ma et al. 2002b, 2003, 2004a, b; Hou et al. 2005).

5465

<i>Spinacia turkestanica</i> Iljin – Amaranthaceae (formerly in Chenopodiaceae)	5466
Synonyms: <i>Spinachia tetrandra</i> Steven ex M. Bieb.	5467
English name: Turkestan spinach, Wild spinach	5468
Russian name: Шпинат туркестанский (Shpinat turkestanskiy)	5469
Uzbek name: Chuchka tikan	5470
Kyrgyz name: Туркстан шпинаты (Turkstan shpinaty)	5471
Description: Dioecious, herbaceous annual, glabrous or with slight farinaceous bloom. Stem 10–60 cm tall, unbranched or sometimes with elongated lower branches. Basal leaves and lower stem leaves runcinate, with a large triangular-hastate terminal lobe and oblong or linear lateral lobes, long-petiolate; upper stem leaves triangular-hastate with shorter petioles. Male inflorescences interrupted spikes, axillary and terminal, nearly-leafless. Female flowers clustered in leaf axils. Fruits consist of 4–6 flowers accreted in to a spiny aggregate (3–8 mm long) with thorny horns. Surface of aggregate and horns smooth or wrinkled; horns usually oblong- pyramidal, triangular in cross-section.	5472 5473 5474 5475 5476 5477
Other distinguishing features: Staminate flowers with 4 perianth segments and very exserted stamens. Fresh leaves have an alkaline flavor.	5478 5479
Phenology: Flowers and fruits in April-June.	5480
Reproduction: By seeds.	5481
Distribution: Toshkent, Samarqand, Buxoro, and Surxondaryo provinces of Uzbekistan; Osh and Chuy provinces of Kyrgyzstan.	5482 5483
Habitat: The adyr and tau zones. A weed of irrigated and unirrigated fields and foothill pastures.	5484
Population status: Common.	5485
Traditional use: The leaves are used as a carminative. It is recommended as a poly-vitamin for treatment of anemia and rickets (Khalmatov 1964).	5486 5487
Documented effects: Spinach is a valuable food crop due its high iodine, calcium, iron, vitamin, protein, and fat content. Spinach can compete with milk with its protein content; the protein is mainly contained in the leaves (Bakiev and Makhkamov 1987). The spinach protein, secretin, is used in medicine like pilocarpine, as a therapeutic agent to stimulate the mucus coating of the stomach lining and the pancreatic glands (Khalmatov 1964).	5488 5489 5490 5491
Phytochemistry: The leaves contain 80 mg% carotene, 64 units/100 g of vitamin B1, up to 40 units of vitamin B2, 16 mg% of vitamin C, and peculiar proteins (Khalmatov 1964).	5492 5493



▲ *Silybum marianum* (L.) Gaertn. Photos: Evgeny Davkaev



◀ *Sphaerophysa salsula* (Pall.) DC. Photos: Alexander Ivanov

▼ *Spinacia turkestanica* Iljin Photos: Chris Kik



▼ *Sorbus tianschanica* Rupr. Photos: Vladimir Epiktetov



- Tanacetum vulgare L. – Asteraceae** 5495
- Synonyms:** *Chrysanthemum tanacetum* Vis., *Chrysanthemum vulgare* (L.) Bernh., *Pyrethrum vulgare* (L.) Boiss., *Tanacetum boreale* Fisch. ex DC., *Tanacetum crispum* Steud., *Tanacetum umbellatum* Gilib. 5496
- English name:** Common tansy 5497
- Russian name:** Пижма обыкновенная (Pizhma obyknovennaya) 5498
- Uzbek name:** Oddi dastarbosh 5499
- Kyrgyz name:** Кадимки танацетум (Kadimki tanatsetum) 5500
- Description:** Herbaceous perennial, with long, woody rhizomes. Stems many, erect, 50–150 cm tall, branched in upper part. 5501
Leaves alternate, up to 20 cm long, 3–10 cm wide, bipinnatisect, elongate-ovate; basal leaves petiolate; stem leaves sessile; lobes pinnatifid or dentate. Inflorescences capitula in flat-topped corymbs; capitula semispherical, compact, 5–10 cm wide, with up to 200 flowers. Disc flowers yellow, 2–3 mm long, 5-lobed, peripheral flowers ca. 20, 3–4-lobed; ray flowers absent. Fruits elongate achenes, often ribbed. 5502
- Other distinguishing features:** Leaves nearly or completely glabrous. 5503
- Phenology:** Flowers and fruits in July-October. 5504
- Reproduction:** By seeds and vegetatively. 5505
- Distribution:** Naryn, Ysyk-Kol, Osh, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 5506
- Habitat:** On steppes, grassy slopes, in meadows, among shrubs, along rivers and roads, and in spruce forests. 5507
- Population status:** Common, found in small groups. 5508
- Traditional use:** Inflorescences are used as a vermifuge, carminative, and choleric, to heal wounds, and to treat intoxication due to lung tuberculosis, fevers, gastrointestinal diseases, and low acidity (Khalmatov et al. 1984). An infusion of the inflorescences is used in Russian folk medicine to increase appetite, bile and sweat production, blood pressure, and decrease heart rates. The infusion is also used as an antipyretic, antispasmodic, anti-inflammatory, analgesic, vermifuge, insecticide, and anti-microbial, and to heal wounds. In the folk medicine of North Caucasus a decoction of the herb is used to treat headaches, and is used externally to treat rheumatism. A decoction of the inflorescences is used to treat skin cancer (Altimishev 1991). 5509
- Documented effects:** A decoction of the inflorescences is used as a vermifuge (for ascarides and pinworm), to treat liver diseases (hepatitis and angiocholitis), gall bladder diseases, and acute gastrointestinal diseases. A water infusion of the inflorescences has shown to be an effective treatment for enterocolitis and other intestinal diseases. An infusion of the inflorescences and leaves is used externally as a bath and compress as a pain killer, to treat gout, rheumatism, joint pain, sprains, bruises and to heal wounds. The infusion of this plant is prohibited for pregnant women (Altimishev 1991). In experiments with animals, an infusion of the inflorescences increased heart beat amplitude and blood pressure, decreased heart rate, increased choleresis, tonified the gastrointestinal tract, and increased its secretions (Akopov 1990). An extract of the plant and isolated compounds have been shown to have anti-inflammatory properties in vivo (Schinella et al. 1998). 5510
- Phytochemistry:** Leaves and inflorescences contain essential oil (α -thujone, β -thujone, L-camphor, thujol, borneol, pinene, etc.), flavonoids (luteolin, quercetin, apigenin, diosmetin, etc.), tannins, bitter substances, and alkaloids (Khalmatov et al. 1984; Akopov 1990; Schinella et al. 1998; Williams et al. 1999). 5511

5533 **Taraxacum officinale F.H. Wigg. – Asteraceae**

5534 **Synonyms:** *Leontodon taraxacum* L., *Taraxacum dens-leonis* Desf., *Taraxacum retroflexum* H. Lindb., *Taraxacum sylvani-*
5535 *cum* R. Doll.

5536 **English name:** Common dandelion

5537 **Russian name:** Одуванчик лекарственный (Oduvanchik lekarstvennyy)

5538 **Uzbek name:** Koki, Momakaimok, Gulkoku

5539 **Kyrgyz name:** Дары какымы (Dary kakumu)

5540 **Description:** Herbaceous perennial, with taproot. Leaves in basal rosette, numerous, oblanceolate, 10–25 cm long, 1.5–5 cm
5541 wide, pinnifid or wide-triangularly toothed. Inflorescences capitula, with hollow peduncles up to 50 cm tall; involucrel
5542 bracts in 2 series. Flowers all ligulate, yellow. Fruits light brown achenes, 3–4 mm long, with a long, thin beak, bearing
5543 white pappus.

5544 **Other distinguishing features:** Mature inflorescences with mature fruits look spherical due to large pappi.

5545 **Phenology:** Flowers in April-May, fruits in May-June.

5546 **Reproduction:** By seeds.

5547 **Distribution:** Almost all provinces of Kyrgyzstan and Uzbekistan.

5548 **Habitat:** In meadows, forest glades, in orchards and parks, near roads and in populated areas as a weed.

5549 **Population status:** Common, forming dense groups.

5550 **Traditional use:** Fresh juice from the leaves is recommended as a laxative and to treat anemia and general weakness. Milky
5551 juice, from the roots, is used to eliminate warts and a galenical preparation of the roots is used to treat skin conditions
5552 (Khalmatov et al. 1984; Mamedov et al. 2004). The roots are collected in autumn and the leaves in spring before flowering.
5553 In Chinese medicine all parts of the plant are used as a antipyretic (diaphoretic), and leaves are used to strengthen the
5554 function of mammary glands (Akopov 1990). The plant is used as a remedy for jaundice, liver and gallbladder disorders,
5555 and as a treatment for water retention and breast and uterus cancer (Koo et al. 2004).

5556 **Documented effects:** A methanol extract of the flowers inhibited inflammation in induced mouse ear edema experiments
5557 (Yasukawa et al. 1998). In scientific medicine a decoction or extract is used to increase the appetite, to aid in function of
5558 the digestive tract and is used as a choleric and laxative. A powder from the roots is used in a complex remedy to treat
5559 arteriosclerosis (Khalmatov et al. 1984). Flower extracts have shown antioxidant activity in vitro (Hu and Kitts 2004). An
5560 aqueous extract of the plant has exhibited anti-tumor actions and was shown to induce apoptosis of human carcinoma cells
5561 in vitro. Taraxasterol has also been shown to have anticarcinogenic activity (Koo et al. 2004).

5562 **Phytochemistry:** Roots contain sesquiterpene lactones, triterpene compounds (taraxerol, taraxasterol, pseudotaraxasterol,
5563 β -sitosterin, and stigmasterin), taraxol, inulin, caoutchouc, and fatty oil, which contains glycerides of palmitic, oleic,
5564 linoleic, melissic, and cerotinic acid. The inflorescence and leaves contain coumarins (cichoriin and aesculin), flavonoids
5565 and flavonoid glycosides (chrysoeriol [3'-methoxyluteolin], luteolin, luteolin 7-glucoside and its derivatives), carotinoids
5566 (taraxanthin, flavoxanthin, and lutein), triterpene alcohols (arnidiol and faradiol) and vitamin B2. The leaves contain
5567 ascorbic and chicoric acid. Monocaffeoyltartaric and chlorogenic acid have been found throughout the plant (Tolmachev
5568 1976; Akopov 1990; Williams et al. 1996; Kisiel and Barszcz 2000).

- Thalictrum foetidum L. – Ranunculaceae** 5570
- Synonyms:** *Thalictrum minus* var. *foetidum* (L.) Hook. f. & Thomson 5571
- English name:** Foetid meadow rue 5572
- Russian name:** Василистник вонючий (Vasilistnik vonyuchiy) 5573
- Uzbek name:** Sassiik sanchikoot 5574
- Kyrgyz name:** Сасык тармал чоп (Sasyk tarmal chop) 5575
- Description:** Herbaceous perennial. Stems 15–100 cm tall, simple or branched, gray-green, often violet. Leaves petiolate, wide-triangular, tri- or quadripinnate; leaflets 4–15 mm long, 2–15 mm wide, broad-ovate, 3-lobed; lobes entire or with 2–3 rounded teeth. Inflorescence a loose, spreading panicle. Sepals 5, 2.5–4 mm long, ovate, violet. Petals absent. Stamens many, with yellow anthers. Fruits ovoid achenes, 3–5 mm long. 5576
5577
5578
5579
- Other distinguishing features:** The entire plant, especially the lower surfaces of the leaves, are covered with hairs and small glands. 5580
5581
- Phenology:** Flowers in June-July, fruits in July-August. 5582
- Reproduction:** By seeds. 5583
- Distribution:** Ysyk-Kol, Jalal-Abad, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 5584
- Habitat:** On stony slopes with rocky debris, in subalpine and alpine mountain belts. 5585
- Population status:** Common, found in large groups. 5586
- Traditional use:** The plant is used as an antiemetic. In Tibetan medicine this species is used to treat edema and gynecological diseases (Akopov 1990). Decoction or tea of the aboveground parts is recommended to treat epilepsy, jaundice, edema, lung tuberculosis, nose bleeds, gastrointestinal ailments, common colds, and gynecological diseases, as well as a general tonic. The herb is used in a poultice to treat bruises, wounds, abscesses, and rheumatism. A decoction of the roots is drunk to treat diarrhea, ulcers, and liver and kidney diseases. A tea of the seeds and herb is drunk to treat side pains, headache, dizziness, and bronchitis (Khodzhimatov 1989; Mamedov et al. 2004). 5587
5588
5589
5590
5591
5592
- Documented effects:** An infusion is used as a treatment for early stage hypertension, stenocardia, and poor blood circulation. The alkaloid foetidin has anti-inflammatory and antiedemic action (Gammerman et al. 1990; Grinkevich 1991). Glycosides isolated from the plant reduced cholesterol in the blood serum and showed anti tumor activity as well as contraceptive effects (Khamidullina et al. 2006). 5593
5594
5595
5596
- Phytochemistry:** The aboveground parts contains alkaloids (foetidine, magnoflorine, thalfine, and thalfinine), flavonoids (rutin, glycosides, etc.), coumarins, triterpene glycosides, tannins, organic acids, and resins. The underground parts contain alkaloids (Ganenko et al. 1986; Rakhimov et al. 1987; Akopov 1990). 5597
5598
5599

5601 **Thalictrum isopyroides C.A. Mey. – Ranunculaceae**5602 **Синонимы:** Unknown5603 **English name:** Unknown5604 **Russian name:** Василистник изопириодный (Vasilistnik izopiroidnyy)5605 **Unbek name:** Sanchikut5606 **Kyrgyz name:** Терен кесиктуу тармал чоп (Teren kesiktuu tarmal chop)

5607 **Description:** Herbaceous perennial with fibrous roots. Stem 8–45 cm tall, simple or branched, glabrous. Leaves tri- or quad-
5608 ripinnatisect, gray, glabrous, with short petioles, concentrated at the base of stem, usually in groups of 2–3; leaflets
5609 broadly rhomboid, 3-lobed, thick; lobes lanceolate-linear or oblanceolate; terminal lobe divided up to the middle or to the
5610 base into 2–3 lanceolate segments with smooth margins. Inflorescence a very loose panicle. Sepals greenish, ca. 2 mm
5611 long. Petals absent. Stamens 5–8. Fruits narrow-ovoid achenes, 4–5 mm long, 1–2 mm wide.

5612 **Other distinguishing features:** Stigma triangular-winged, persistent in fruit.5613 **Phenology:** Flowers in April- early May, fruits in May-June.5614 **Reproduction:** By seeds.

5615 **Distribution:** Toshkent, Andijon, Farg'ona, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; Talas,
5616 Osh and Jalal-Abad provinces of Kyrgyzstan.

5617 **Habitat:** The adyr and tau zones. On shallow-soiled, stony slopes with rocky debris on hills and mountains.5618 **Population status:** Rare, found as single individuals.

5619 **Traditional use:** In Tajikistan a tea made from the aboveground parts is used to treat fever, chest pain, and as an anticonvul-
5620 sive. A decoction of the herb is drunk to treat epilepsy, jaundice, tachycardia, nose bleeds, lung tuberculosis, gastrointes-
5621 tinal, and feminine diseases. A decoction of the roots is drunk to treat stomach ulcers, liver and kidney disease, and high
5622 blood pressure. A tea of the seeds is recommended to treat dizziness, high blood pressure, bronchitis, and edema
5623 (Khodzhimatov 1989). An infusion of the plant is used to treat diarrhea, jaundice, malaria, epilepsy and lung tuberculosis,
5624 and is used externally to treat skin diseases (Khalmatov 1964).

5625 **Documented effects:** In experiments with animals, the alkaloid thalisopine acted as a sedative and had pronounced anticon-
5626 vulsant activity which surpassed that of phenytoin and trimetin (Tashbaev and Sultanov 1962, 1965). When injected
5627 intravenously, it had distinct antiarrhythmic action on experimental models (Akbarov et al. 1972). The alkaloid cryptopine
5628 stimulated uterine smooth muscles, had vasoconstrictive action, and increased arterial pressure in narcotized animals. The
5629 alkaloid magnoflorine reduced blood pressure due to its ganglio-blocking action (Fakhrutdinov 1971; Fakhrutdinov and
5630 Sultanov 1972). In anesthetized animals, intravenous injections of the alkaloid thalicminine caused short-term reduction
5631 of blood pressure and heart rate (Abdalla et al. 1991).

5632 **Phytochemistry:** Plants studied were found to contain 3.22 % total alkaloids. Thaliosopine, thalisopidine, dehydrothalic-
5633 mine, thalicmine, thalicminine, cryptopine, magnoflorine, and others were isolated from the total alkaloids (Yunusov
5634 1974; Abduzhabbarova et al. 1978).

5635



▲ *Thalictum isopyroides* C.A. Mey. Photos: Alim Gaziev

▼ *Taraxacum officinale* F.H. Wigg. Photos: *top*: Dmitri Oreshkin; *center and bottom*: Sergey Appolonov



▲ *Thalictum foetidum* L. Photos: Petr Filippov



◀ *Tanacetum vulgare* L.
Photos: *left*: Mary Backlund;
right: Sergey Appolonov



5636 **Thalictrum minus L. – Ranunculaceae**5637 **Синонимы:** Unknown5638 **English name:** Small meadow-rue, lesser meadow-rue5639 **Russian name:** Василистник малый (Vasilistnik malyy)5640 **Uzbek name:** Sanchuq ut5641 **Kyrgyz name:** Кичинекей тармал чоп (Kichinekey tarmal chop)

5642 **Description:** Herbaceous perennial. Stems 30–100 cm high, glabrous, smooth, erect or irregularly bending, evenly folia-
5643 ceous. Leaves alternate, tri- or quadripinnatisect, wide-triangular in outline, greenish-gray, petiolate (upper leaves ses-
5644 sile); leaflets almost round, 0.8–4 cm long and wide, irregularly lobed. Inflorescence an oval or pyramidal panicle. Sepals
5645 ovate, 3–4 mm long, 2 mm wide, yellowish-green. Stamens 10–15. Fruits ovoid achenes, 4–5 mm long, 2 mm wide,
5646 ribbed, with an erect or slightly bent tip, sessile.

5647 **Other distinguishing features:** Inflorescence spreading and much branched. Leaves deflected from the stem.5648 **Phenology:** Flowers in June–July, fruits in June–August.5649 **Reproduction:** By seeds.5650 **Distribution:** Toshkent, Jizzax, Samarqand, Qashqadaryo, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan.5651 **Habitat:** The adyr and tau zones. In the valleys of lowland and mountain rivers.5652 **Population status:** Uncommon, found as single individuals.

5653 **Traditional use:** An infusion of the herb is used to treat various diseases: skin, diarrhea, hepatitis, malaria, epilepsy, tuber-
5654 culosis, fevers, and is also used as a hemostatic (Khalmatov et al. 1984).

5655 **Documented effects:** Extracts are used as a hemostatic, for hypotonia to increase blood pressure, and as an antibacterial
5656 against gram-positive bacteria. In various animals using different modes of application, the alkaloids thalicmine and thal-
5657 icmidine caused depression of the central nervous system and elongated effects of soporifics. In higher doses they pro-
5658 duced catalepsy (Zabirov and Kasmaliev 1962; Sadritdinov et al. 1971; Sadritdinov 1973; Sadritdinov and Khamdamov
5659 1975). The alkaloid thalictimine had ganglion blocking action (cardiac ganglion n. vagus) and inhibited the cough reflex
5660 (Sadritdinov and Kurmukov 1980). Thalictimine and thalmine have sedative and short-term hypotensive effects; thalmine
5661 also had anti-inflammatory, analgesic, and antipyretic action (Sadritdinov and Sultanov 1971; Fakhrutdinov and Sultanov
5662 1972; Sadritdinov 1971b, 1973; Abdalla et al. 1991). The alkaloids O-methyl-thalicberine, thalisopine, and thalmine had
5663 antiarrhythmic action. Thalispine exceeded the activity of quinidine and procainamide-hydrochloride (Akbarov et al.
5664 1978). Experiments have shown that a number of the alkaloids isolated from the plant have antimicrobial activity against
5665 *Mycobacterium smegmatis* (Liao et al. 1978). The alkaloid thaliblastine exhibited activity against various types of cancer
5666 (Mircheva and Stoychkov 1976; Ilarionova et al. 1980; Stoychkov and Miloushev 1980; Todorov and Zeller 1992; Chen
5667 et al. 1992).

5668 **Phytochemistry:** The aboveground parts contained up to 1 % total alkaloids (thalmine and thalictimine), flavonoids (1.64 %),
5669 saponins (3.1 %), vitamin C (175.7–761.7 mg%), organic acids, tannins, bitter, and other substances; the roots contained
5670 1.1 % total alkaloids (thalictimine, thalictidine, thalictitrine, tolmetin, argemonine and others). The seeds contained
5671 22.9–28.4 % fatty oil (Yunusov 1981; Khalmatov et al. 1984; Sidjimov et al. 1998).

Thermopsis alterniflora Regel & Schmalh. – Fabaceae	5673
Synonyms: <i>Thermopsis rigida</i> Vassilcz.	5674
English name: Unknown	5675
Russian name: Термопсис очередноцветковый (Termopsis ocherednotsvetkovyy)	5676
Uzbek name: Afsonak (Афсонак)	5677
Kyrgyz name: Кезек гулдуу сары мья (Kezek gulduu sary myya)	5678
Description: Herbaceous perennial with vigorous rhizomes. Stems erect, up to 100 cm tall, branched, middle and upper portion with varying amounts of hairs. Leaves alternate, trifoliolate, petiolate with lanceolate stipules; leaflets oblong-elliptic, 2.5–5 cm long, 1–2 cm wide, acuminate, adaxial side glabrous, abaxial side slightly hairy. Inflorescence a loose apical raceme, 9–20 cm long, with oblanceolate bracts. Flowers alternate. Calyx 10–20 mm long, densely covered with silky hairs. Corolla papilionaceous, yellow. Fruits oblong-elliptic legumes, 3–6 cm long, 1–1.6 cm wide, covered with short, appressed hairs, few-seeded. Seeds kidney-shaped, 5–6 mm long, 3–4 mm wide, brownish-red-greenish, glabrous.	5679 5680 5681 5682 5683 5684
Other distinguishing features: Calyx teeth one third to one half as long as tube. All 10 stamens free.	5685
Phenology: Flowers in May-June, fruits in June-August.	5686
Reproduction: By seeds and rhizomes.	5687
Distribution: Tashkent province of Uzbekistan, in the Western Tien Shan; Osh and Jalal-Abad provinces of Kyrgyzstan.	5688
Habitat: The tau zone. Shallow-soiled slopes and mountain brook valleys; as a weed among unirrigated cereal crops.	5689
Population status: Uncommon, sometimes in populations with up to 50 individuals.	5690
Traditional use: A galenical preparation of the stems, leaves, flowers and fruits is used to treat bronchial asthma (Mamedov and Craker 2001).	5691 5692
Documented effects: Used as an expectorant and vermifuge (Khalmatov 1964). The alkaloid cytisine is used to prepare a 0.15 % solution, called <i>cytion</i> , which is used to increase respiration in cases of respiratory standstill, such as during operations and traumas, from infectious diseases, shocks, various intoxication (such as poisoning by carbon oxide, prussic acid, and narcotics), asphyxia of newborns, and others. Pachycarpine increases uterine contractility and is used in obstetrical practice to stimulate contractions for weak labors, and also to stop bleeding during the post-natal period (Mashkovskii 1984).	5693 5694 5695 5696 5697 5698
Phytochemistry: At the beginning of flowering, 3.5 % total alkaloids were obtained from the aboveground parts. Cytisine (>50 % of total alkaloids), pachycarpine, methylcytisine, thermopsine, anagirine, argentine, alteramine, dimethylamine, and other alkaloids were isolated from the total alkaloids. The flavonoids cinaroside, luteolin, chrysoeriol, thermopsocide, genistein, and genistin were also isolated from the aboveground parts. The aboveground portion also contained 4.88 % titrated organic acids, up to 4.8 % sugars, and 5.08 % resins. Roots contained 0.81 % and seeds contained up to 3.34 % total alkaloids (Khalmatov et al. 1984).	5699 5700 5701 5702 5703 5704

5706 **Thermopsis lanceolata R. Br. – Fabaceae**

5707 **Synonyms:** *Sophora lupinoides* L., *Thermopsis dahurica* Czefr., *Thermopsis glabra* Czefr., *Thermopsis lupinoides* (L.)
5708 Link., *Thermopsis sibirica* Czefr.

5709 **English name:** Lanceleaf thermopsis

5710 **Russian name:** Термопсис ланцетный, Мышатник (*Thermopsis lantsetnyu*, *Myshatnik*)

5711 **Uzbek name:** Lantsetcemon termopsis, Lantsetcemon afsonak

5712 **Kyrgyz name:** Ланцетный сары мыя (*Lantsetnyu sary myya*)

5713 **Description:** Herbaceous perennial, with a deep main root and lateral rhizomes. Stems up to 40 cm tall, many, erect, branched,
5714 striated, hairy. Leaves alternate, petiolate, trifoliolate; leaflets elongate- or oblanceolate, 2.3–7.6 cm long, 0.8–2.3 cm wide,
5715 glabrous above, hairy below. Flowers in whorls forming terminal racemose inflorescences. Calyx campanulate, with 5
5716 lanceolate lobes. Corolla papilionaceous, yellow. Fruit a narrow-linear legume, straight or slightly arched, 4–8.8 cm long,
5717 0.7–1.2 cm wide, short hairy. Seeds nearly round, dark olive or nearly black, glaucous.

5718 **Other distinguishing features:** Stamens 10, all distinct. Legumes not flattened, sharply tapering at the end.

5719 **Phenology:** Flowers in May–June, fruits in July–August.

5720 **Reproduction:** By seeds and rhizomes.

5721 **Distribution:** Ysyk-Kol, Naryn, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan.

5722 **Habitat:** Found among *Achnatherum splendens* (Trin.) Nevski on the coast of Lake Ysyk-Kol, on solonchic soils, and in
5723 fallow and cultivated fields. Not found high into the mountains.

5724 **Population status:** Common, found in dense groups.

5725 **Traditional use:** The aboveground parts are harvested before flowering. Decoctions of the aboveground parts are used to
5726 treat respiratory catarrh, flu, bronchitis, pneumonia, and headaches (Akopov 1990; Mamedov and Craker 2001).

5727 **Documented effects:** An infusion of the herb is used as an expectorant to treat chronic bronchitis and residual pneumonia.
5728 The preparation *Cyiton*, which contains the alkaloid cytosine isolated from the seeds, is used to stimulate respiratory
5729 function and improve blood circulation. The preparation is used to treat asphyxia in newborns and when a person stops
5730 breathing during surgical procedures or from trauma (Khalmatov et al. 1984). The alkaloid pachycarpine, isolated from
5731 this plant, is used to treat peripheral vessel spasms and to induce labor when necessary (Akopov 1990).

5732 **Phytochemistry:** The herb contains alkaloids (thermopsine, homothermopsine, methylcytisine, pachycarpine, and anagyrene),
5733 saponins, tannins, resins, mucilage, traces of essential oil, and ascorbic acid. The seeds contain alkaloids, mainly cytosine
5734 (Tolmachev 1976; Akopov 1990).

5735

Thermopsis turkestanica Gand. – Fabaceae	5736
Synonyms: <i>Thermopsis kaxgarica</i> Chang Y. Yang, <i>Thermopsis lanceolata</i> ssp. <i>turkestanica</i> (Gand.) Gubanov.	5737
English name: Unknown	5738
Russian name: Термопсис туркестанский (<i>Thermopsis turkestanica</i>)	5739
Uzbek name: Unknown	5740
Kyrgyz name: Туркестан сары мяясы (<i>Turkestan sary myyasy</i>)	5741
Description: Herbaceous perennial, with a deep main root and lateral rhizomes. Stems many, erect, 30–50 cm tall, striated, branched; branches appressed to main stems. Leaves alternate, petiolate, trifoliate; leaflets 3.5–8 cm long, 0.5–1 cm wide, narrowly lanceolate. Flowers in whorls forming terminal racemose inflorescences. Calyx campanulate; lobes 5, lanceolate. Corolla papilionaceous, yellow. Fruit an elongate-linear legume, 4.5–7 cm long, 0.8–1 cm wide, light-brown, densely covered with short hairs. Seeds ellipsoid, dark green.	5742 5743 5744 5745 5746
Other distinguishing features: Stamens 10, all distinct. Legumes flattened and slowly tapering to the end.	5747
Phenology: Flowers in June-July, fruits in July-August.	5748
Reproduction: By seeds and rhizomes.	5749
Distribution: Ysyk-Kol, Naryn, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan.	5750
Habitat: In fallow fields and on solonchic soils among <i>Achnatherum splendens</i> (Trin.) Nevski.. Found in high mountain valleys.	5751 5752
Population status: Common, forming dense groups.	5753
Traditional use: A decoction of the herb is used to treat low blood pressure and as an expectorant (Teslov 1960).	5754
Documented effects: An alcoholic extract of the aboveground parts strengthens respiratory function and raises blood pressure (Chefranova 1954).	5755 5756
Phytochemistry: The aboveground parts contain alkaloids (cytisine, thermopsine, N-methylcytisine, anagyrine, and sparteine; Plant Resources of the USSR 1987).	5757 5758



▲ Thermopsis alterniflora Regel & Schmalh. Photos: Alim Gaziev



▲ Thalictrum minus L. Photos: Alim Gaziev

▼ Thermopsis turkestanica Gand.

▼ Thermopsis lanceolata R. Br. Photo: Klazina Witteveen

Photo: Vladimir Epiktetov



Thymus marschallianus Willd. – Lamiaceae

- Synonyms:** *Thymus amictus* Klok., *Thymus latifolius* (Bess.) Andrz., *Thymus pannonicus* All., *Thymus pannonicus* ssp. *marschallianus* (Willd.) Soó, *Thymus platyphyllus* Klok., *Thymus pseudopannonicus* Klok., *Thymus stepposus* Klok. & Shost. 5760-5763
- English name:** Unknown 5764
- Russian name:** Тимьян Маршаллов (Tim'yan Marshallov) 5765
- Uzbek name:** Kaklikoot, Toshchop 5766
- Kyrgyz name:** Кадимки кийик оту (Kadimki kiyik otu) 5767
- Description:** Perennial subshrub. Stems short, much branched, upper-half retrorse-pubescent, flower-bearing branches 12–37 cm tall. Leaves opposite, sessile, oblanceolate or elongate-elliptic, 12.5–30 mm long, 2.5–7.5 mm wide, abaxially glandular, margin entire or slightly serrulate. Inflorescences verticillasters in apical spikes; pedicels densely pubescent. Calyx tubular-campanulate, 2-lipped; upper lip 3-toothed; lower lip 2-toothed. Corolla red-purple, lilac or white, pubescent, 2-lipped; lower lip 3-lobed. Fruits ovoid nutlets. 5768-5772
- Other distinguishing features:** Plants gynodioecious. 5773
- Phenology:** Flowers in May, fruits in August. 5774
- Reproduction:** By seeds. 5775
- Distribution:** Ysyk-Kol and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 5776
- Habitat:** On foothills, in meadow-steppes, meadow slopes, on the edges of spruce forests, among juniper stands. 5777
- Population status:** Common. 5778
- Traditional use:** An infusion of the herb is used to treat stomatitis and toothaches. A decoction in milk is used to treat acute respiratory infections and amenorrhea. In Bulgaria the herb is used to heal wounds and a decoction is used to treat stomach ulcers and bad breath. An infusion is used in the Altai region to treat fevers and headaches and in the Middle Volga region as an expectorant for acute respiratory infections and pertussis (Plant Resources of the USSR 1991). 5779-5782
- Documented effects:** In modern medicine the herb is used in a similar manner as *Thymus serpyllum*. A decoction and liquid extract is recommended for internal use as an expectorant to treat bronchitis and other upper respiratory illnesses. It is used externally in compresses and baths as an analgesic to treat radiculitis and neuritis (Tolmachev 1976). The complex preparation *Pertussin*, containing this herb, is used as an expectorant and cough suppressant and to treat bronchitis and other upper respiratory illnesses (Kurochkin 1998). Ethanolic extracts of *Thymus marschallianus* exhibited antioxidative activity (Budincevic et al. 1995). Volatile oils isolated from the plant exhibited antibacterial activity against the gram-positive bacterium *Diplococcus pneumoniae* (Oprean et al. 2007). 5783-5788
- Phytochemistry:** Aboveground parts contain phenolcarboxylic acids and their derivatives (caffeic, rosmarinic, 1-caffeoylquinic, and 5-caffeoylquinic acids), flavonoids (luteolin, apigenin, scutellarein, and anthocyanins) and essential oil (containing thymol, carvacrol, α -pinene, camphene, sabinene, n-thymol, isoborneol, borneol, undecanoic acid, and amyl alcohol; Plant Resources of the USSR 1991; Kolesnikov and Gins 2001; Stahl-Biskup 2002). 5790-5793

5795 **Tribulus terrestris L. – Zygophyllaceae**5796 **Synonyms:** *Tribulus bicornutus* Fisch. & Mey.5797 **English name:** Puncturevine, Caltrop5798 **Russian name:** Якорцы стелющиеся (Yakortsy stelyushchiesya)5799 **Uzbek name:** Temirtikan5800 **Kyrgyz name:** Тошолмо мык тикен (Tosholmo myk tiken)

5801 **Description:** Herbaceous annual with a thin taproot. Stems 20–80 cm long, branched, spreading, prostrate and rising only at
 5802 tips, usually hairy. Leaves opposite, even-pinnate, 3–6 cm long, short-petiolate, with small stipules; leaflets in 6–8 pairs,
 5803 oblong, 4–10 mm long, adaxial side glabrous, abaxial side hairy. Flowers solitary in leaf axils, 1–1.2 cm in diameter,
 5804 pedicels 4–10 mm long. Sepals 5. Petals 5, yellowish. Stamens 10. Style 1. Fruits schizocarpic, flattened, star-shaped;
 5805 mericarps 5, dry, angular, tuberculate with 2 or 4 divergent spines.

5806 **Other distinguishing features:** Plant often appears glaucescent.5807 **Phenology:** Flowers and fruits in May-August.5808 **Reproduction:** By seeds.5809 **Distribution:** All of Uzbekistan and Kyrgyzstan.5810 **Habitat:** The arid zone. Waste places, oases, unirrigated fields, near roads, dry slopes, and slopes along rivers and brooks.5811 **Population status:** Common, not forming dense groups.

5812 **Traditional use:** This plant has been used since ancient times for various diseases. Avicenna recommended caltrop for
 5813 tumors and ulcers, especially for festering ulcers of the gums, as a diuretic, and to remove kidney and bladder stones. Folk
 5814 medicine in the East uses decoctions and infusions of the herb as a purgative, diuretic and tonic, for gonorrhoea, headaches
 5815 and eye inflammations, and for strong side pains. Cleaned roots are boiled in milk and used for chronic malaria and as a
 5816 energizing remedy (Seredin and Sokolov 1969; Khalmatov et al. 1984). In Western countries it is used to increase the
 5817 libido, and as a tonic, astringent, and diuretic (Gammerman et al. 1990).

5818 **Documented effects:** A liquid extract of this species (collected during flowering) is used to treat people with low levels of
 5819 stomach acidity due to hypo- and anacidic gastritis and as a diuretic to treat swelling. Extracts made from the plant (col-
 5820 lected during fruiting period) are also used as a diuretic. A preparation from the leaves, *Tribusponin*, which contains ste-
 5821 roid glycosides, is used as an antisclerotic treatment (Seredin and Sokolov 1969; Gammerman et al. 1990). Two compounds
 5822 isolated from the plant, tribulosin and β -sitosterol-D-glucoside, exhibited antihelmintic activity (Deepak et al. 2002).
 5823 Steroidal saponins, isolated from the plant, exhibited antifungal activity against *Candida albicans* and *Cryptococcus*
 5824 *neoformans* and anti-cancer activity against a variety of cancer cell lines (Bedir et al. 2002). Rats that were given an oral
 5825 extract of the fruits exhibited weight gain and improvement in sexual behavior parameters (Gauthaman et al. 2003). The
 5826 systolic blood pressure of hypertensive rats that were fed an extract of the fruits was significantly decreased compared to
 5827 unfed hypertensive rats. The ACE (angiotensin-converting enzyme) activity in all tissues of extract fed hypertensive rats
 5828 was significantly lower than that of the control rats (Sharifi et al. 2003).

5829 **Phytochemistry:** The plant contains flavonoids, alkaloids (harman, etc.), amides, and steroidal saponins (diosgenin dehy-
 5830 dration products including crystalline diosgenin, gitogenin, ruscogenin, and 25-D-spirosta-3,5-diene), and saponins with
 5831 a hemolytic index of 1:240. The leaves contain up to 160 mg% vitamin C. The seeds contain alkaloids and the fruits con-
 5832 tain around 5 % tannins and fatty drying oil (Seredin and Sokolov 1969; Gammerman et al. 1990; Wang et al. 1997; Wu
 5833 et al. 1999b; Deepak et al. 2002).

Trichodesma incanum (Bunge) A. DC. – Boraginaceae	5835
Synonyms: <i>Friedrichsthalia incana</i> Bunge.	5836
English name: Unknown	5837
Russian name: Триходесма седая (<i>Trikhodesma sedaya</i>)	5838
Uzbek name: Kampir chopon	5839
Kyrgyz name: Боз триходесма (<i>Boz trikhodesma</i>)	5840
Description: Rhizomatous perennial up to 30–100 cm tall. Stems ascending, branched, herbaceous, densely covered with short, gray pubescence, becoming shiny, woody and glabrescent below. Leaves alternate or subopposite, ovate to oblanceolate, 3–8 cm long, 1.3–2.8 cm wide, apex acute, margins entire, sessile, both sides silky gray-pubescent. Inflorescences loose, narrow-paniculate, composed of terminal scorpioid cymes. Flowers pedicellate, drooping. Calyx ovate-campanulate, gray-tomentose with 5 deeply divided oblanceolate lobes. Corolla ca. 2 cm in diameter, with a short tube and 5 broad, triangular-ovate lobes with tail-like appendages; at the beginning of flowering the tube is white and lobes light-blue, later the tube turns pink and lobes dark-blue. Anthers yellow, forming an exerted cone. Fruits ovoid nutlets, 6–8 mm long, grayish-brown, dull and covered with tiny wrinkles and tubercles, edges slightly uneven or toothed.	5841 5842 5843 5844 5845 5846 5847 5848
Other distinguishing features: Anthers with spirally-twisted awn-like appendages. Calyx enlarged in fruit, becoming disk-shaped and membranous.	5849 5850
Phenology: Flowers and fruits from May to November.	5851
Reproduction: By seeds and rhizomes.	5852
Distribution: Karakalpakstan autonomous republic, Toshkent, Andijon, Farg'ona, Samarqand, and Surxondaryo provinces of Uzbekistan; Talas, Batken and Osh provinces of Kyrgyzstan.	5853 5854
Habitat: The adyr and tau zones. Loess slopes of hills, stony slopes with rocky debris, as well as unirrigated and abandoned fields.	5855 5856
Population status: Uncommon, found as single individuals.	5857
Traditional use: The plant roots (as a root-powder plaster) are used to heal persistent wounds and furunculosis. A decoction of the roots and leaves is recommended for scabies and is applied on infected skin areas (Khalmatov 1964).	5858 5859
Documented effects: This plant is highly poisonous. The alkaloids contained in this species act as neurovascular toxins. The alkaloids depress blood production, destroy erythrocytes, induce hypoxia in tissues, and increase vein wall permeability. These toxins have the ability to accumulate in the body (Vilner 1974). The alkaloid incanine (the N-oxide [amine oxide]) and the alkaloid trichodesmine lower arterial pressure and have antispasmodic action (Mashkovskii 1983).	5860 5861 5862 5863
Phytochemistry: All plant parts contain alkaloids. Immature fruits contain up to 1.5 %, mature fruits 2.7 %, and the above-ground parts, before flowering, up to 1 % alkaloids. The flowering herb contains only 0.3 % total alkaloids. The alkaloids incanine (1.5 % in seeds), N-oxide form of incanine, trichodesmine, and N-oxide form of trichodesmine have been isolated from the total alkaloids. At flowering period the plant top contains up to 70 % trichodesmine from the total alkaloid content (Yunusov 1981).	5864 5865 5866 5867 5868

5870 **Trifolium pratense L. – Fabaceae**5871 **Synonyms:** *Trifolium ucrainicum* Opperm. ex Wissjul.5872 **English name:** Red clover

5873 Russian name: Клевер луговой, Клевер красный (Klever lugovoy, Klever krasnyy)

5874 **Uzbek name:** Sebarga5875 **Kyrgyz name:** Шалбаа уй бедеси (Shalbaa uy bedesi)5876 **Description:** Herbaceous perennial, slightly hairy, with taproot. Stems to 80 cm tall, erect or suberect, simple or branched.

5877 Leaves trifoliate, stipulate; lower leaves long-petiolate; upper leaves short-petiolate or sessile; leaflets obovate or elliptical, usually sinuate, rarely serrulate, often with a white triangular blotch. Inflorescence head-like, ovoid or globose.

5878 Flowers 1.3–2 cm long, in globose or ovoid heads. Calyx tubular-campanulate, with 5 teeth (1 longer, 4 shorter). Corolla papilionaceous, light-pink to dark-red. Fruits small legumes nearly enclosed by calyx.

5881 **Other distinguishing features:** Stamens 10 (9 united). Heads on top of stems and lateral branches.5882 **Phenology:** Flowers in May-September, fruits in June-October.5883 **Reproduction:** By seeds.5884 **Distribution:** All provinces of Kyrgyzstan; Tashkent, Farg'ona, Samarqand, Qashqadaryo and Surxondaryo provinces of Uzbekistan.5886 **Habitat:** Along rivers, in damp meadows and valleys, in the high-mountain meadow and forest belt, and in tallgrass meadows.5888 **Population status:** Common, found in dense groups.5889 **Traditional use:** An infusion or tea of the flowers is used as an antiseptic, expectorant, diuretic, anti-inflammatory, and analgesic, and to treat lung and upper respiratory tract diseases, sore throat, bronchial asthma, pertussis, malaria, rheumatism, hypertension, stenocardia, anemia, uterine bleeding, leucorrhea, shortness of breath, coughs, and painful menstruation. Freshly ground leaves or fresh juice from the plant are applied externally to treat infected wounds and skin ulcers, burns, and rubella (Nuraliev 1989; Akopov 1990). The seeds are used to increase the libido and are used to treat prolonged fevers (Khodzhimatov 1989).5895 **Documented effects:** A tincture of the plant is used to treat arteriosclerosis in patients with normal blood pressure (Nuraliev 1989). In modern medicine this species is used as an expectorant, diuretic, and antiseptic (Khodzhimatov 1989). Metabolites of isoflavones found in the plant were found to protect against UV radiation-induced inflammation and immunosuppression (Widyarini et al. 2001). Isoflavones found in red clover inhibited COX enzyme activity in certain cancer cell types (Lam et al. 2004). Extracts of red clover and individual flavonoid constituents exhibited estrogenic activity in a variety of in vitro assays (Overk et al. 2005).5901 **Phytochemistry:** The herb contains many flavonoids (trifolin, isotrifolin, trifoside, etc.), asparagine, tyrosine, coumarinic and salicylic acids, alkaloids, fatty oil, carotene, B vitamins, and vitamin C. The roots contain coumarins (Khalmatov 1964; Akopov 1990; Lin et al. 2000; Klejdus et al. 2001).



▲ **Trifolium pratense L.** Photos: *left*: Dmitri Oreshkin; *center*: Sasha Eisenman; *right*: Sergey Appolonov

▼ **Thymus marschallianus Willd.** Photos: Andrei Lubchenko



◀ **Tribulus terrestris L.** Photos: *top*: Sergey Mayorov; *center and bottom*: Sasha Eisenman

▼ **Trichodesma incanum (Bunge) A. DC.** Photos: Alim Gaziev



5905 **Tussilago farfara L. – Asteraceae**5906 **Synonyms:** None5907 **English name:** Coltsfoot5908 **Russian name:** Мать-и-мачеха обыкновенная (Mat'-i-machekha obyknovennaya)5909 **Uzbek name:** Okkaldirmok5910 **Kyrgyz name:** Кадимки огой Эне (Kadimki ogoy ene)

5911 **Description:** Herbaceous perennial, with creeping rhizomes. Flowering stems 7–45 cm tall, felted-hairy, with small, alter-
5912 nate bracts. Leaves basal, long-petiolate, orbicular-cordate, 8–15 cm long, up to 12 cm wide, glabrous above, felted-hairy
5913 beneath, shallowly lobed, margins unequally dentate. Stem leaves scale-like, ovate-lanceolate, 0.6–1.5 cm long, 0.3–
5914 0.8 cm wide, sessile, purple-violet, appressed to stem. Inflorescences capitula. Ray flowers 100–300, golden yellow; disc
5915 flowers 20–40, yellowish. Fruits linear achenes, 3–4.5 mm long; pappus white, longer than achene.

5916 **Other distinguishing features:** The plant blossoms before the leaves appear.5917 **Phenology:** Flowers in May–September, fruits in June–October.5918 **Distribution:** All provinces of Kyrgyzstan; Toshkent, Farg'ona, Samarqand and Surxondaryo provinces of Uzbekistan.5919 **Habitat:** On damp lawns, along mountain rivers and streams, and in areas with water-eroded soils and alluvial deposits.5920 **Population status:** Common, found in dense groups.

5921 **Traditional use:** An infusion and decoction of the leaves and flowers are used as an expectorant and cough suppressant, to
5922 treat bronchial asthma, as well as a diuretic to treat edema and scrofula. It is applied externally as a poultice or wash to
5923 treat tumors, abscesses, and furuncles. Juice from fresh leaves and roots is used to treat tuberculosis and malaria, and as
5924 a choleric and diaphoretic (Khalmatov et al. 1984). Leaves are used to treat acute and chronic bronchitis, catarrh of the
5925 upper respiratory system, pneumonia, laryngitis, bronchial pneumonia, and a hoarse voice. Preparations of coltsfoot are
5926 used to treat tracheitis, kidney and bladder diseases, the gastrointestinal tract, loss of appetite, fever, erysipelous skin
5927 inflammation, scrofula, hair loss, and abscesses. Fresh juice from the leaves is inhaled into the nostrils to eliminate sinus
5928 colds. The juice of leaves is also mixed with powdered sugar to treat tuberculosis (Maznev 2004).

5929 **Documented effects:** An infusion of the leaves and flowers is used as an expectorant and demulcent. A tea made from the
5930 leaves is used to treat bronchitis, laryngitis, bronchiectasis, abscesses, and gangrene of the lungs. A poultice is used exter-
5931 nally as a demulcent, disinfectant, and anti-inflammatory (Tolmachev 1976; Khalmatov et al. 1984). A sesquiterpene
5932 isolated from extracts of the buds was found to have anti-inflammatory effects in vitro and reduced induced rat foot edema
5933 (Hwang et al. 1987). Extracts of both the aboveground parts and rhizomes showed antimicrobial activity against *Bacillus*
5934 *cereus* and *Staphylococcus aureus* (Kokoska et al. 2002). Flavonoids isolated from the flower buds exhibited antioxidative
5935 activity (Kim et al. 2006). Various compounds isolated from the plant induced cardiovascular and respiratory stimulation
5936 and have been shown to have anti-inflammatory activities by inhibiting arachidonic acid metabolism, platelet-activating
5937 factor receptors, and the activity of nitric oxide synthesis (Ryu et al. 1999). The ethyl acetate fraction of the plant extract
5938 had neuroprotective and antioxidant effects in vitro (Cho et al. 2005).

5939 **Phytochemistry:** Leaves contain bitter glycosides, carotenoids, alkaloids, flavonoids, coumarins, saponins, mucilage, tan-
5940 nins, organic acids, cytosterin, inulin, tussilaglin, and vitamin C. Flowers contain rutin, arnidiol, faradiol, taraxanthin,
5941 stigmaterin, cytosterin, phytosterins, n-heptacosane, tannins, etc. (Tolmachev 1976; Khodzhimatov 1989; Ryu et al.
5942 1999).

- Ungernia victoris Vved. ex Artjushenko – Amaryllidaceae** 5944
- Synonyms:** None 5945
- English name:** Unknown 5946
- Russian name:** Унгерния Виктора (Ungerniya Viktora) 5947
- Uzbek name:** Omonqora 5948
- Kyrgyz name:** Unknown 5949
- Description:** Herbaceous perennial up to 20 cm tall. Bulbs ovoid, small, 4–7 cm wide, with multiple black-brown, papery coats. Stem bearing inflorescence flattened, 5–10 cm long. Leaves 7–10 in number, in 2 rows, linear, 20–25 cm long and 2–3 cm wide, light blue-gray, smooth. Inflorescence an umbel bearing 4–7 flowers. Flowers funnelform, 5–6 mm wide, yellowish to yellow-pink. Stamens 6. Fruit a capsule, 2–3 cm wide, with 3 wide heart-shaped valves. Seeds flat, black. 5950–5953
- Other distinguishing features:** Flowers after leaves have senesced. 5954
- Phenology:** Flowers in August, fruits in September. 5955
- Reproduction:** By seeds and vegetatively by bulbs. 5956
- Distribution:** Gissar mountain range, Chulbair mountains in Surxondaryo province of Uzbekistan; not found in Kyrgyzstan. 5957–5958
- Habitat:** The tau zone. Shallow-soiled slopes and ravines. 5959
- Population status:** Rare endemic of southwest Pamiro-Alai; listed in the Red Book of Rare and Endangered Species of Uzbekistan. 5960–5961
- Traditional use:** Baked bulbs are used to heal wounds; they are also applied to furuncles to remove pus. According to Avicenna the herb and its seeds are the best treatment for diarrhea. If the seeds are taken with water or wine it helps to heal stomach ulcers and improve digestion. Wine infused with the seeds are used to treat kidney stones (Karimov and Shomakhmudov 1993). 5962–5966
- Documented effects:** This species is recommended as the raw material to obtain the alkaloid galanthamine. Hydrobromic salt of galanthamine is widely used in medical practice to treat myasthenia, myopathia, and for post-poliomyelitis, radiculitis, and polyneuritis palsies, as well as traumatic injuries of sensory and motor nerves. The alkaloid narwedine has anti-narcotic action and facilitates transfer of nervous excitation to H- and M-cholinergic synapses. A preparation was recommended for clinical trials as an anti-narcotic drug. The alkaloid pancratine lowers blood pressure, has sedative action, and increases activity of soporifics. In acute tests, hordenine shows adrenomimetic action. It is used for intestinal peristalsis inhibition in diarrhea. Licorine has anti-inflammatory, analgesic, and antipyretic action and strengthens the hypothermic action of amidopyrine. This alkaloid also strengthens the secretion of intestines and lung-bronchial glands in dogs and cats (Sadritdinov and Kurmukov 1980). An extract derived from cultured plant cells exhibited antimutagenic properties (Dvornyk et al. 2002). 5967–5975
- Phytochemistry:** Leaves contain 0.35–1 %, bulbs 0.8–0.9 %, and roots 2.5 % total alkaloids. Galanthamine, pancratine, narwedine, hordenine, and licorine have been isolated from the leaves. Similarly, galanthamine, licorine, pancratine, tatsetine, and hippeastrine have been isolated from the bulbs (Yunusov 1981). 5976–5978

5980 **Urtica dioica L. – Urticaceae**5981 **Synonyms:** None5982 **English name:** Stinging nettle5983 **Russian name:** Крапива двудомная (Krapiva dvudomnaya)5984 **Uzbek name:** Gazanda, Kichitki oot, Chayan oot5985 **Kyrgyz name:** Чалкан (Chalkan)5986 **Description:** Dioecious, herbaceous perennial plant, with creeping rhizomes. Stems and leaves covered with stinging hairs.

5987 Stems erect, 30–170 cm tall. Leaves opposite, simple, ovate-lanceolate, 8–17 cm long, 2–8 cm wide, apex acuminate,
5988 margins large dentate. Inflorescences axillary panicles. Flowers unisexual, small, green. Staminate flowers with 4 equal
5989 tepals; stamens 4. Pistillate flowers with 4 tepals, inner 2 equal to achene, outer 2 smaller. Fruits ovoid or elliptic achenes,
5990 1–1.5 mm long.

5991 **Other distinguishing features:** Staminate inflorescences ascending, pistillate lax or recurved in fruit. Achenes smooth.5992 **Phenology:** Flowers and fruits in May–September.5993 **Reproduction:** By seeds and rhizomes.

5994 **Distribution:** All provinces of Kyrgyzstan; Karakalpakstan autonomous republic, Toshkent, Andijon, Namangan, Farg'ona,
5995 Samarqand, Qashqadaryo and Xorazm provinces of Uzbekistan.

5996 **Habitat:** Found as a weed in settlements, along canals, in woods, and among bushes.5997 **Population status:** Common, forming dense groups.

5998 **Traditional use:** A decoction and powder of leaves is used as a diuretic, laxative, expectorant, vasoconstrictor, and hemo-
5999 static to treat internal bleeding and hemorrhoids, and also to treat rheumatism, stomach diseases, diabetes, and chronic
6000 ulcers. It is used in a bath to treat various types of swelling. The roots and fruits are used to treat diarrhea. The leaves are
6001 used in a wash to treat hair loss. A water extract of the leaves, along with extracts of different plant species are used to
6002 prepare a cream with cow bone marrow which is used to wash and encourage hair growth (Khalmatov et al. 1984;
6003 Khodzhimatov 1989).

6004 **Documented effects:** Preparations from this plant species are used internally as a hemostatic, to increase uterine contrac-
6005 tions and to increase blood coagulation. The preparations are effective in treating climacteric menopause and hemor-
6006 rhoids. Extracts of this species are used to normalize the menstrual cycle. A preparation of the herb has pressor action on
6007 internal organ vessels (Tolmachev 1976). Preparations of this species decrease blood cholesterol content and have chol-
6008 eretic and anti-inflammatory activities. In studies with diabetic patients, a decoction of leaves decreased blood and urine
6009 sugar levels (Kurochkin 1998). Preparations of nettle normalize metabolism and blood sugar content, increase blood
6010 coagulability, increase milk production in feeding mothers, normalize lipid metabolism, increase hemoglobin and eryth-
6011 rocyte content, increase intestine and cardiovascular tonus and stimulate epithelization of wounded tissues. This species
6012 helps to treat liver illnesses, joint rheumatism, and gastrointestinal and bladder diseases (Maznev 2004). A water extract
6013 of the plant had antioxidant and analgesic activity, showed antimicrobial activity against 9 microorganisms, and exhibited
6014 antiulcer activity against ethanol-induced ulcerogenesis (Gulcin et al. 2004). A methanolic extract of the roots exhibited
6015 antiproliferative effects on human prostate cancer cells in vivo and in vitro (Konrad et al. 2000). A fraction from the
6016 extract of the leaves caused a marked increase in insulin secretion by the pancreatic islets of Langerhans in normal and
6017 induced diabetic rats (Farzami et al. 2003).

6018 **Phytochemistry:** Leaves contain carotene and other carotenoids, organic acids (formic, pantothenic, caffeic, p-coumaric,
6019 and ferulic), glycosides (urticin), sitosterin, phytoncides, quercetin, acetylcholine, histamine, tannins, mineral salts, vita-
6020 mins C, K and group B, resin, protoporphyrin, koproporphyrin, and 5-hydroxytryptamine (Tolmachev 1976; Chikov
6021 1989; Kurochkin 1998).

6022

- Vaccaria hispanica (Mill.) Rauschert – Caryophyllaceae** 6023
- Synonyms:** *Saponaria segetalis* Neck, *Saponaria vaccaria* L., *Vaccaria parviflora* Moench., *Vaccaria pyramidata* Medik., 6024
Vaccaria segetalis (Neck) Garke ex. Asch., *Vaccaria vulgaris* Host. 6025
- English name:** Cow cockle, Cow herb 6026
- Russian name:** Тысячеголов пирамидальный, Тысячеголов посевной (Tysyachegolov piramidal'nyy, Tysyachegolov 6027
posevnoy) 6028
- Uzbek name:** Qora mug 6029
- Kyrgyz name:** Айдама мин баш (Aydama min bash) 6030
- Description:** Herbaceous annual, glabrous, glaucous. Stem erect, 30–70 cm tall, heavily branched towards top. Leaves oppo- 6031
site, simple, sessile, ovate-lanceolate to oblong-ovate, 2–9 cm long, blue-gray, apex acute, base almost cordate and slightly 6032
connate. Inflorescence a paniculiform-cyme, pedicels 1–6 cm long. Calyx 1.3–1.5 cm long, consists of 5 connate sepals, 6033
yellowish-green. Petals 5, with linear claws, pink. Stamens 10. Styles 2. Fruit a capsule, wide-ovoid, shorter than the 6034
calyx. Seeds black, globose, tuberculate, 1.5 mm wide. 6035
- Other distinguishing features:** Calyx with 5 raised longitudinal ribs. When fruiting, calyx swollen at the base, the top very 6036
narrowed. 6037
- Phenology:** Flowers and fruits in April-July. 6038
- Reproduction:** By seeds. 6039
- Distribution:** All provinces of Uzbekistan; in agricultural zones of all provinces of Kyrgyzstan. 6040
- Habitat:** The chul and adyr zones. A weed of cultivated fields, especially in unirrigated wheat fields. 6041
- Population status:** Common. 6042
- Traditional use:** In Chinese medicine, the seeds are used as an analgesic, to stimulate milk let-down, to promote diuresis, to 6043
activate blood circulation, relieve carbuncles, and to treat amenorrhea and breast infections. It is also used in ointments, 6044
which are used for treating skin diseases (eczema and psoriasis). In Central Asia, a plaster of the herb is used to treat 6045
tumors and as an analgesic (Khalmatov 1964; Morita et al. 1997b; Sang et al. 2000). 6046
- Documented effects:** Hemolytic index of the herb is equal to 1:1450, of the roots 1:4000, and the seeds contain 3.18 % of a 6047
poisonous saponin with a high hemolytic index (1:50,000 in human blood and 1:25,000 in dog's blood). Convolvine and 6048
convolamine act as local anesthetics. However, because they are highly toxic and not very effective, they are not used for 6049
this purpose. After modification, a derivative of convolamine, convocaine, was introduced for use in hospitals (Ogolevitz 6050
1951). Peptides isolated from the seeds exhibited estrogen-like activity and caused uterine contractions in vitro (Morita et 6051
al. 1997a, b). 6052
- Phytochemistry:** A wide assortment of chemical compounds have been isolated from the seeds including triterpene saponins, 6053
alkaloids (up to 0.5 %, convolvine and convolamine), cyclic peptides, phenolic acid, flavonoids, and steroids. Roots con- 6054
tain 5 % saponins, sugars, saporubin, and saporubinic acid. Leaves contain the glycoside saponarin (Ogolevitz 1951; 6055
Morita et al. 1997a,b; Sang et al. 2000, 2003). 6056



▲ *Tussilago farfara* L. Photos: *left and right*: Alexander Naumenko; *center*: Evgeny Davkaev

▼ *Ungernia victoris* Vved. ex Artjushenko Photos: Alim Gaziev



▲ *Urtica dioica* L. Photos: Sergey Appolonov

▼ *Vaccaria hispanica* (Mill.) Rauschert
Photos: Andrei Lubchenko



Valeriana officinalis L. – Valerianaceae**Synonyms:** *Valeriana baltica* Pleijel, *Valeriana exaltata* Mikan fil., *Valeriana palustris* Kreyer.**English name:** Valerian, Garden valerian, Garden heliotrope**Russian name:** Валериана лекарственная (*Valeriana lekarstvennaya*)**Uzbek name:** Asaroon**Kyrgyz name:** Дары мышык тамыр (*Dary myshyk tamyr*)**Description:** Herbaceous perennial, with short rhizomes. Stems single or few, 50–150 cm tall, hollow, furrowed. Leaves opposite, 7–25 cm long, odd-pinnately compound with 6–8 pairs of leaflets, lower leaves petiolate; leaflets ovate-lanceolate or almost linear, entire to dentate. Inflorescence corymbiform or paniculiform, apical. Flowers perfect. Corolla funnelform with 5 lobes, white or pale-lilac. Stamens 3. Fruits flattened achenes, 2–3 mm long.**Other distinguishing features:** Calyx initially small, later enlarged with plumose, pappus-like segments. Roots have a strong, specific smell.**Phenology:** Flowers in May, fruits in July.**Reproduction:** By seeds and division of rhizomes.**Distribution:** All provinces of Kyrgyzstan; cultivated in Uzbekistan.**Habitat:** Cultivated.**Population status:** Common.**Traditional use:** Valerian is used as a sedative, carminative, and vermifuge, as an aid in digestion, and to treat hypercondria, psychological traumas, hysteria, migraines, convulsive pains, heart pains, heart failure, epilepsy, insomnia, and anxiety (Turova and Sapozhnikova 1984; Altimishev 1991). A decoction or tincture is used as a heart remedy, a sedative to treat nervous disorders, as well as to treat headaches, and cancer, and to improve the appetite. It is used in a bath to relax hyperactive children so they sleep well and to treat hysteria, convulsions, acute typhus, epilepsy, and internal aches (Kurochkin 1998).**Documented effects:** Preparations of valerian influence the nervous system and have sedative effects as well as antispasmodic actions (Kurochkin 1998). They are used to treat insomnia, neurosis of the cardiovascular system and to treat spasms of the gastrointestinal tract. In acute and chronic experiments with dogs given valerian infusion intravenously and orally, arterial pressure was decreased (only when applied intravenously) and the speed of blood coagulation was increased (Akopov 1990). Valepotriates suppress aggression, have anticonvulsant effects against pentylenetetrazol- and strychnine-induced seizures, increase thiopental-induced sleeping time, reduce motility and have dose-dependent sedative effects. The sesquiterpenes reduce locomotion and increase pentobarbital and hexobarbital-induced sleeping time of mice. Some sesquiterpenes, especially valerenic acid, influence serotonin and noradrenaline levels (Ortiz et al. 1999).**Phytochemistry:** Underground parts contain essential oil with sesquiterpenes, iridoids, etc. (including bornyl-isovalerianate, dihydrovaltrate, valtrate, acevaltrate, isovaltrate, valerenic and iso-valerianic acid, borneol, myrtenol, myrtenyl isovalerianate, camphene, α -pinene, d-terpeneol, limonene, alcohols, etc.), alkaloids (valerine, chatinene, etc.), glycosides (valeride), tannins, sugars, acids (formic, acetic, malic, stearic, palmitinic, etc.) and macro- and micro-elements (Akopov 1990; Bos et al. 1998; Kurochkin 1998; Ortiz et al. 1999).

6058

6059

6060

6061

6062

6063

6064

6065

6066

6067

6068

6069

6070

6071

6072

6073

6074

6075

6076

6077

6078

6079

6080

6081

6082

6083

6084

6085

6086

6087

6088

6089

6090

6091

6092

6093

6094

6095 **Veratrum lobelianum Bernh. – Melanthiaceae**

6096 **Synonyms:** *Veratrum album* ssp. *lobelianum* (Bernh.) Schuebl. & Martens, *Veratrum album* ssp. *virescens* (Gaudin) Jav. &
6097 Soo, *Veratrum album* var. *lobelianum* (Bernh.) Koch, *Veratrum album* var. *virescens* Gaudin.

6098 **English name:** Unknown

6099 **Russian name:** Чемерица Лобеля (Chemeritsa Lobelya)

6100 **Uzbek name:** Maralkulok

6101 **Kyrgyz name:** Лобел марал кулагы (Lobel maral kulagy)

6102 **Description:** Herbaceous perennial, with short rhizomes. Stem single, erect, 2–3 cm in diameter, 70–170 cm tall. Leaves
6103 cauline, alternate, simple, sheathing the stem, prominently veined, margins entire; lower leaves wide-elliptic, 15–25 cm
6104 long, 10–15 cm wide; upper leaves smaller, lanceolate. Inflorescence an apical panicle, 20–60 cm tall. Flowers with 6
6105 white-green tepals and 6 stamens. Fruit an ovoid capsule, 3-lobed. Seeds flat, elliptical, broad-winged, 6–10 mm long.

6106 **Other distinguishing features:** Ovary superior.

6107 **Phenology:** Flowers in June-July, fruits in July-September.

6108 **Reproduction:** By seeds and rhizomes.

6109 **Distribution:** Ysyk-Kol province of Kyrgyzstan; not found in the flora of Uzbekistan.

6110 **Habitat:** In wet meadows, bogs and along rivers.

6111 **Population status:** Common, forming dense groups.

6112 **Traditional use:** The underground parts infused in cream is recommended to treat eczema (Khalmatov et al. 1984). This
6113 plant is used to treat mental illness and is used externally to treat joint rheumatism and neuralgia (Altimishev 1991).
6114 A tincture of the rhizome is used as a hypotensive in Bulgarian folk medicine (Ivancheva and Stantcheva 2000).

6115 **Documented effects:** A preparation of this species has insecticidal activity. In medicine an alcoholic or water infusion is
6116 used externally on skin parasites and scabies. An alcohol infusion, decoction, and ointment prepared with the under-
6117 ground parts are used as an analgesic to treat neuralgia, arthritis, rheumatism, and common colds. Alkaloids isolated from
6118 this species have hypotensive and analgesic abilities. Because of the high toxicity of the alkaloids they are not widely used
6119 in medicine (Khalmatov et al. 1984; Kurochkin 1998). The alkaloid jervine isolated from this plant effected fibroblasts
6120 and isolated animal organs in vitro in a similar manner as serotonin (Suladze et al. 2006).

6121 **Phytochemistry:** The plant contains alkaloids (jervine, pseudojervine, rubijervine, isorubijervine, etc.), tannins, resins, sug-
6122 ars, and pigments (Bondarenko 1972; Khashimov et al. 1970; Tolmachev 1976; Shakirov et al. 1995; Suladze et al.
6123 2006).

6124

- Verbascum songaricum Schrenk – Scrophulariaceae** 6125
- Synonyms:** *Verbascum khorassanicum* Boiss., *Verbascum lychnitis* L., *Verbascum polystachyum* Kar. & Kir. 6126
- English name:** Songar mullein 6127
- Russian name:** Коровяк джунгарский (Korovyak dzhungarskiy) 6128
- Uzbek name:** Sigir kuuruq 6129
- Kyrgyz name:** Жунгар аюу кулагы (Zhungar ayuu kulagy) 6130
- Description:** Herbaceous biennial, densely stellate hairy. Stem 40–150 cm high, erect, foliaceous, branched on top. Basal 6131
leaves lanceolate to oblanceolate, 15–40 cm long, 4–12 cm wide, base attenuate, margin nearly entire, grayish-hairy on 6132
both sides, nearly sessile to petiolate; stem leaves lanceolate to oblong, sessile; base of upper leaves subcordate. 6133
Inflorescence pyramidal-paniculate, 20–40 cm long. Flowers in bunches of 4–7, pedicellate. Calyx 4–10 mm long with 5 6134
deep linear-lanceolated lobes, whitish-hairy. Corolla yellow, 1.5–3 cm in diameter, 5-lobed. Stamens 5. Fruit a wide-ovoid 6135
capsule, 5–8 mm long, densely hairy. Seeds tiny, obconic-prismatic, 0.7–0.9 mm long, 0.6 mm wide, linearly pitted. 6136
- Other distinguishing features:** Staminal filaments are coated with whitish hairs. 6137
- Phenology:** Flowers in June-August, fruits in July-September. 6138
- Reproduction:** By seeds. 6139
- Distribution:** Toshkent, Farg'ona, Samarqand, and Surxondaryo provinces of Uzbekistan; all of Kyrgyzstan. 6140
- Habitat:** The adyr and tau zones. Dry slopes of foothills. 6141
- Population status:** Common, found as single individuals. 6142
- Traditional use:** The plant is used to heal wounds. Thoroughly boiled leaves are put on burns, tumors and wounds, and the 6143
fresh leaf juice is applied on the surface of wounds. It is also used for toothaches, eye inflammations, and as an expecto- 6144
rant to relieve chronic cough. A decoction of the flowers is used to treat stomach and intestinal catarrh and gall bladder 6145
and liver inflammation (Seredin and Sokolov 1969). 6146
- Documented effects:** An infusion of the flowers in water is used as an expectorant. Decoctions of the leaves and flowers of 6147
this species, as well as the related species *Verbascum thapsus*, *V. phlomoides*, and *V. thapsiforme* are used as an expecto- 6148
rant and to coat and sooth the mouth and throat to reduce effects of catarrh and coughs (Seredin and Sokolov 1969). 6149
Phenylethanoid glycosides isolated from a methanolic extract of the plant inhibited mammalian DNA polymerases (Iida 6150
et al. 2003). 6151
- Phytochemistry:** The whole plant contains alkaloids, including anabasine and plantagonine, saponins, triterpenoid saponins, 6152
and vitamin C (Khodzhimatov 1989; Seifert et al. 1991; Hartleb and Seifert 1995). The aboveground parts contains 6153
saponins with a hemolytic index of 1:250 (Khalmatov 1964). 6154

6156 **Verbascum thapsus L. – Scrophulariaceae**6157 **Synonyms:** None6158 **English name:** Common mullein6159 **Russian name:** Коровяк обыкновенный (Korovyak obyknovennyu)6160 **Uzbek name:** Unknown6161 **Kyrgyz name:** Аю кулак (Ayu kulak)

6162 **Description:** Herbaceous biennial, densely felted-hairy. Stem thick, leafy, up to 2 m tall. Lower leaves in a basal rosette, peti-
6163 olate, oblong or oblanceolate, up to 30 cm long, up to 5–10 cm wide, usually entire; upper leaves alternate, becoming
6164 smaller, sessile, decurrent on stem to next leaf below. Inflorescence a dense, apical, spiciform raceme, appearing in the
6165 second year. Calyx deeply 5-lobed. Corolla yellow, 1–2.5 cm in diameter, 5-lobed, the lower 3 lobes slightly longer than
6166 the upper 2. Fruit a septicidal capsule with 2 valves. Seeds small, furrowed.

6167 **Other distinguishing features:** Stamens 5, upper 3 shorter than the lower 2.6168 **Phenology:** Flowers in May-June, fruits in July-August.6169 **Reproduction:** By seeds.6170 **Distribution:** Jalal-Abad, Osh, and Chuy Provinces of Kyrgyzstan; not found in the flora of Uzbekistan.6171 **Habitat:** In fallow fields and pastures and along canals.6172 **Population status:** Common, found in loosely arranged groups.

6173 **Traditional use:** A decoction of the herb is used to treat neurosis and epilepsy, as a diuretic to treat kidney stones, and gout
6174 and swelling due to kidney and heart problems. It is used externally to treat throat diseases, neuralgia of facial nerves, in
6175 a bath to treat hemorrhoids, scrofula, and rickets, and as a compress or lotion to heal wounds and treat eye diseases. A
6176 decoction of the roots and leaves is used to treat diarrhea. An infusion and decoction of the leaves and flowers is used as
6177 an expectorant, anti-inflammatory, demulcent and coating to treat acute respiratory diseases, pneumonia, bronchial
6178 asthma, gastritis and liver and gall bladder diseases (Plant Resources of the USSR 1990).

6179 **Documented effects:** Extracts of the plant exhibited varying antibacterial activity against *Klebsiella pneumonia*,
6180 *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Escherichia coli*, inhibited *Agrobacterium tumefaciens*-induced
6181 tumors in vitro, and had antiviral activity (McCutcheon et al. 1995; Turker and Camper 2002).

6182 **Phytochemistry:** The plant contain polysaccharides, iridoid glycosides (harpagoside, harpagide, and aucubin) flavonoids
6183 (3-methylquercetin, hesperidin, and verbascoside) saponins, essential oil, steroids, mucilage, etc. (Turker and Gurel
6184 2005).

6185



▲ **Valeriana officinalis L.**
Photos: Vadim Prokhorov



▲ **Veratrum lobelianum Bernh.**
Photos: *top, upper center and lower center:* Vadim Prokhorov;
bottom: Sergey Appolonov



▲ **Verbascum songaricum**
Schrenk Photos: Alim Gaziev

▼ **Verbascum thapsus L.**
Photo: Sasha Eisenman



6186 **Vexibia pachycarpa (Schrenk ex C.A. Mey.) Yakovlev – Fabaceae**

6187 **Synonyms:** *Goebelia pachycarpa* (Schrenk ex C.A. Mey.) Bunge ex Boiss., *Sophora pachycarpa* Schrenk ex C.A. Mey.

6188 **English name:** Unknown

6189 **Russian name:** Вексибия толстоплодная (Veksibiya tolstoplodnaya)

6190 **Uzbek name:** Achykmiya

6191 **Kyrgyz name:** Unknown

6192 **Description:** Herbaceous perennial, 30–60 cm tall, stems branching from the base, densely covered with short, white hairs.

6193 Leaves alternate, compound, odd-pinnate, 10–18 cm long; leaflets in 6–12 pairs, elliptic or oblong, 1.5–2 cm long, 3–8 mm
6194 wide, both sides white-hairy. Inflorescences cylindrical, apical racemes. Calyx wide-campanulate, with wide-triangular
6195 teeth, densely hairy. Corolla papilionaceous, white to creamy-yellow colored, up to 1.5 cm long. Fruits club-shaped
6196 legumes, 3–6 cm long, 7–9 mm wide, with varying amounts of pubescence, legumes oriented vertically. Seeds slightly
6197 kidney-shaped to oval, deep-brown, glabrous.

6198 **Other distinguishing features:** Legumes slightly constricted between seeds, with elongated, conical tip.

6199 **Phenology:** Flowers in April-May, fruits in June-July.

6200 **Reproduction:** By seeds and rhizomes.

6201 **Distribution:** Toshkent, Farg'ona, Samarqand, and Buxoro provinces of Uzbekistan; found in some provinces of
6202 Kyrgyzstan.

6203 **Habitat:** The chul and adyr zones. On river banks, in long-fallow fields, on loess hills, sandy soils, and as a weed in unirri-
6204 gated wheat fields.

6205 **Population status:** Common, usually occurs in small populations or as single individuals.

6206 **Traditional use:** The ground seeds are recommended for loss of appetite. An decoction of the aboveground plant parts is
6207 used to treat skin diseases (eczema, fungal, and scabies) and as a spasmolytic, analgesic, and vermifuge (Khalmatov et al.
6208 1984; Khodzhimatov 1989; Mamedov et al. 2004).

6209 **Documented effects:** Only pachycarpine is used in medical practice. Pachycarpine is a ganglionic blocking agent and is
6210 used internally for hypertension strokes, peripheral vessels spasms (endarteritis, intermittent claudication), for myopathy
6211 and to stimulate labor during child birth. Dermatologists use pachycarpine preparations for scleroderma, idiopathic skin
6212 atrophy, and to treat chronic eczema (Mashkovskii 1984).

6213 **Phytochemistry:** The aboveground parts contain up to 3 %, and seeds up to 2.2 %, total alkaloids. Plants from Kenimesh
6214 massif (Buxoro province, Uzbekistan) contained 3.90–6.4 % (aboveground) and 1.5–2.98 % (roots) total alkaloids. The
6215 main alkaloids are pachycarpine, sophocarpine, matrine, and sophoramine. Pachycarpidine, quercetin, kaempferol, and
6216 genistein and its xyloglucoside have also been isolated. The roots contained 9–12 % (and the root bark 22–25 %) phenolic
6217 pigments, flavonoids, steroid glucosides, etc. (Yunusov 1981; Botirov et al. 2006; Muminova et al. 2006; Emami et al.
6218 2007).

6219

- Vicia cracca L. – Fabaceae** 6220
- Synonyms:** *Vicia hiteropus* Freyn, *Vicia lilacina* sensu B. Fedtsch., *Vicia macrophylla* (Maxim.) B. Fedtsch. 6221
- English name:** Bird vetch, cow vetch, tufted vetch 6222
- Russian name:** Горошек мышиный (Goposhek myshinyy) 6223
- Uzbek name:** Unknown 6224
- Kyrgyz name:** Жапайы жер буурчак (Zharayu zher buurchak) 6225
- Description:** Herbaceous perennial vine. Stems trailing or climbing, pubescent. Leaves pinnate, usually with 5–11 pairs of 6226
 leaflets, a tendril replacing the terminal leaflet; leaflets linear-lanceolate to narrowly oblong, 1.5–3 cm long, 4–8 mm 6227
 wide, apex mucronate. Inflorescence a long-peduncled one-sided raceme, many-flowered. Calyx campanulate, pink or 6228
 bluish, lobes unequal. Corolla papilionaceous, blue-violet or rarely white. Fruits elongate-lanceolate legumes, 15–20 mm 6229
 long. Seeds dark-brown, spherical. 6230
- Other distinguishing features:** Legumes glabrous. 6231
- Phenology:** Flowers in June-July, fruits in August-September. 6232
- Reproduction:** By seeds. 6233
- Distribution:** Ysyk-Kol, Naryn, Talas, and Chuy provinces of Kyrgyzstan; not found in the flora of Uzbekistan. 6234
- Habitat:** In tall-grass meadows, among bushes, in forests, along canals and in floodplains. 6235
- Population status:** Common, found in small groups. 6236
- Traditional use:** The aboveground parts are used as a demulcent, hemostatic and to heal wounds. A tincture is used to treat 6237
 diarrhea and as a diuretic. A poultice is used to treat rectal prolapse and prolapsed hemorrhoids. The crushed, dry or fresh 6238
 herb is applied as a compress to treat abscesses. In the Bryansk area of Russia a decoction of the roots is used to treat hepa- 6239
 titis. In the Tibetan medicine, the aboveground parts are used to treat swelling, ascites, and as a hemostatic (Plant Resources 6240
 of the USSR 1987). 6241
- Documented effects:** Lectins, isolated from this species, show bonding specificity with human blood type A (Sharon and 6242
 Lis 2004). In experiment on animals, an infusion and decoction of the plant had antibacterial activity (Plant Resources of 6243
 the USSR 1987). 6244
- Phytochemistry:** The aboveground parts contains the non-protein, amino acid canavanine (Enneking 1995), cyanogenic 6245
 glycosides (vicianine), hydrocyanic acid, and vitamin C. The leaves contain vitamins C, P and carotene. The flowers 6246
 contain vitamins, flavonoids and anthocyanins (Savoskin et al. 1971; Shreter 1975). 6247

6249 **Vinca erecta Regel & Schmalh. – Apocynaceae**6250 **Synonyms:** None6251 **English name:** Unknown6252 **Russian name:** Барвинок прямостоящий, Барвинок прямой (Barvinok pryamostoyashchiy, Barvinok pryamoy)6253 **Uzbek name:** Burygul6254 **Kyrgyz name:** Туз бору гул (Tuz boru gul)

6255 **Description:** Herbaceous perennial with horizontal, woody, scale-covered rhizomes. Stems many, erect, 15–50 cm high,
 6256 unbranched, glabrous or hairy. Leaves opposite, densely arranged, sessile; lower leaves simple, glabrous or pubescent,
 6257 1–2 cm long, up to 7 mm wide, apex obtuse or rounded; upper leaves ovate to wide-lanceolate, 2.5–5.5 cm long, 1.2–3 cm
 6258 wide, apex acute. Flowers single, axillary, pedicellate. Corolla 2–2.5 cm long, pale lilac outside, white inside with dark-
 6259 violet tube, glabrous. Fruits composed of 2 linear-cylindrical follicles, 3–6.5 cm long, brown, densely covered with large
 6260 tubercles. Seeds 1.1–1.6 cm long, 2–3 mm wide, light-brown.

6261 **Other distinguishing features:** Follicles have 1–7 seeds, usually 3–4.6262 **Phenology:** Flowers and fruits in May–August.6263 **Reproduction:** By seeds and rhizomes.6264 **Distribution:** Toshkent, Farg'ona, and Surxondaryo provinces of Uzbekistan; Osh and Jalal-Abad provinces of
 6265 Kyrgyzstan.6266 **Habitat:** The adyr and tau zones. Stony slopes, shale taluses in foothills, on rocks.6267 **Population status:** Uncommon, found in small populations.

6268 **Traditional use:** In mountain zones where these plants are growing, local populations use decoctions and infusions of the
 6269 aboveground parts to treat febrile diseases. A decoction of the roots is used as an emetic (Kurmukov 1970). The leaves
 6270 are used in a tea to treat diarrhea and gastrointestinal disorders, headaches and dizziness, and as a mouthwash for tooth-
 6271 aches. The fresh leaves are applied to wounds. A powder is used externally as an astringent and to heal wounds (Khalmatov
 6272 et al. 1984; Khodzhimatov 1989).

6273 **Documented effects:** The total alkaloids of the aboveground parts have different actions at different doses. In low and
 6274 middle doses they act as a sedative and at major doses they have a stimulating effect on the central nervous system. The
 6275 preparation, *Vinsumine*, has antispasmodic, adrenolytic and ganglion blocking (ganglions of cardiac branches of vagus
 6276 nerves) actions, it changes signals from the carotid and sciatic nerves which influence arterial pressure, and it releases and
 6277 prevents cardiac arrhythmia caused by electric heart stimulation and by intravenous introduction of 10 % calcium chloride
 6278 solution. All effects of *Vinsumine* are due to the alkaloids it contains (Kurmukov and Sultanov 1965). The alkaloids aku-
 6279 amidine, tombozine, and ervine have α -adrenolytic action. Ervine shows pronounced anti-fibrillating action for cardiac
 6280 arrhythmia (Kurmukov 1968b, 1970, 1975, 1978). The alkaloid ervinine is an analeptic of the central nervous system with
 6281 primary influence on respiration and stimulates reticular formations of the medulla oblongata and midbrain due to stimu-
 6282 lation of adrenergic structures (Saidkasimov 1960; Kurmukov and Saidkasimov 1968; Kurmukov and Saidkasimov 1969;
 6283 Kurmukov 1970). The alkaloid vincamine has stimulatory action on uterine unstriped muscles and stimulates contractions
 6284 in weak labors. It was used in obstetrics under the preparation name *Vikametrin*. The alkaloid vincarine has anti-arrhyth-
 6285 mic action and is not inferior to aimaline (Khanov et al. 1968, 1972; Kurmukov 1968a, 1970; Kurmukov and Sultanov
 6286 1971). The alkaloid vincanine is a strychnine-like spasmodic and analeptic of the central nervous system (Sultanov
 6287 1959b; Shamansurov and Sultanov 1967). The main effect of the alkaloid vincanidine is an apomorphine-like emetic
 6288 action (Sultanov 1959a, 1960).

6289 **Phytochemistry:** The following alkaloids have been obtained from the aboveground plant parts collected in different areas
 6290 of Uzbekistan: vincamine, ervamine, ervinine, ervine, vinervine, vinervinine, akuamine, akuamidine, reserpinine, isores-
 6291 erpine, and vincamine. Alkaloids, including vincanine and vincanidine, have been isolated from the roots. More than 60
 6292 other alkaloids have been isolated from this species (Yunusov 1981; Yagudaev et al. 1983).

6293

Viola suavis M. Bieb. – Violaceae	6294
Synonyms: <i>Viola pontica</i> W. Beck.	6295
English name: Russian violet	6296
Russian name: Фиалка приятная (Fialka priyatnaya)	6297
Uzbek name: Gunafsha	6298
Kyrgyz name: Жагымдуу ала гул (Zhagymduu ala gul)	6299
Description: Herbaceous perennial, with a short rhizome and short, stout stolons. Leaves in a rosette, obovate to broad-ovate, base cordate, long-petiolate; spring leaves 3–8 cm long; summer leaves up to 20 cm long, margins dentate. Flowers solitary. Sepals 5. Sepals 5, violet with white throat; lower petal with spur. Fruit a spherical capsule, glabrous or pubescent. Seeds with conspicuous elaiosomes.	6300 6301 6302 6303
Other distinguishing features: Stipules free, lanceolate, long-fimbriate. This species also produces cleistogamous flowers.	6304 6305
Phenology: Flowers in April, fruits in May.	6306
Reproduction: By seeds and stolons.	6307
Distribution: Cultivated in Kyrgyzstan and Uzbekistan.	6308
Habitat: Found escaped into the wild.	6309
Population status: Common.	6310
Traditional use: A syrup made from the aboveground parts is used as a diuretic, anti-inflammatory, expectorant, diaphoretic, and choleric. A decoction is used to treat coughs, sinus colds and illnesses of the eyes, throat, and stomach. The roots are used as an emetic and laxative. In Turkmenistan and the Caucasus a decoction of the flowers with sugar is used to treat heart illnesses (Plant Resources of the USSR 1986).	6311 6312 6313 6314
Documented effects: Unknown.	6315
Phytochemistry: Aboveground parts have essential oil and vitamin C (Plant Resources of the USSR 1986).	6316



▲ **Vexibia pachycarpa**
(Schrenk ex C.A. Mey.)
Yakovlev
Photos: Evgeny Davkaev



▲ **Vicia cracca** L.
Photos: *top*: Evgeny Davkaev;
center: Rostislav Lezhoyev;
bottom: Radu Chibzii



▲ **Viola suavis** M. Bieb.
Photos: Maxim Zaitsev
▼ **Vinca erecta** Regel &
Schmalh. Photo: Authors



Xanthium strumarium L. – Asteraceae

- Synonyms:** *Xanthium americanum* Walter, *Xanthium cavanillesii* Schouw, *Xanthium chasei* Fernald, *Xanthium chinense* Mill., *Xanthium curvescens* Millsp. & Sherff, *Xanthium echinatum* Murray, *Xanthium echinellum* Greene ex Rydb., *Xanthium globosum* C. Shull, *Xanthium inflexum* Mack. & Bush, *Xanthium italicum* Moretti, *Xanthium natalense* Widder, *Xanthium orientale* L., *Xanthium oviforme* Wallr., *Xanthium pensylvanicum* Wallr., *Xanthium pungens* Wallr., *Xanthium speciosum* Kearney, *Xanthium varians* Greene, *Xanthium wootonii* Cockerell. 6318-6323
- English name:** Common cocklebur, Rough cocklebur 6324
- Russian name:** Дурнишник обыкновенный (Durnishnik obyknovennyy) 6325
- Uzbek name:** Guzatkon, Patanak 6326
- Kyrgyz name:** Кадимки манкоо (Kadimki manko) 6327
- Description:** Herbaceous annual, with taproot. Stems 20–200 cm tall, branched, appressed hairy or subglabrous. Leaves alternate, long-petiolate, broadly ovate to suborbicular, shallowly 3–5-lobed, irregularly dentate. Inflorescences unisexual heads. Staminate heads many-flowered with highly reduced involucre, heads in a terminal cluster. Pistillate heads in short axillary clusters, heads cylindric to ovoid, 1–3.5 cm long, 2-flowered, enclosed by involucre forming a bur (false-fruit) with curved prickles. Fruits thick achenes with no pappus. 6328-6332
- Other distinguishing features:** Leaves broad, no spines in the axils. 6333
- Phenology:** Flowers in June-July, fruits in July-September. 6334
- Reproduction:** By seeds. 6335
- Distribution:** All provinces of Kyrgyzstan and Uzbekistan. 6336
- Habitat:** Near roads, canals, waste places, sandy riverbanks, and in agricultural fields. 6337
- Population status:** Common. 6338
- Traditional use:** In folk medicine a decoction of the seeds and roots is used to treat dysentery, scrofula, and bladder diseases. A tincture of the entire plant in vodka is drunk to treat goiters, rheumatism and common colds, and inflammatory diseases and is also used as a diaphoretic, antipyretic, and sedative. A tea made from the entire plant is used to treat cancer. An infusion is used to stimulate digestion, to treat intestinal atonia, stomach spasms, liver inflammation, jaundice, acute and chronic bronchitis, pertussis, painful menstruation, kidney stones, goiters, cancer, and to reduce sexual excitability. It is used externally in dry and damp compresses and aromatic baths. Fruits and seeds are used to treat eczema, itchy dermatosis, insect stings, and paralysis. A decoction of the root is used externally to treat skin diseases and furunculosis. A decoction of the entire plant is applied to the face after shaving, especially on pimples and fungal skin diseases (Maznev 2004). 6339-6346
- Documented effects:** An extract of the leaves exhibited trypanocidal activity in vitro and in vivo (Talakal et al. 1995). Rodents treated with an extract of the plant exhibited alterations in behavior patterns that suggested the extract had significant depressing activity on the central nervous system (Mandal et al. 2001). An extract of the plant showed slight activity against *Candida albicans* (Murillo-Alvarez et al. 2001). Caffeic acid isolated from the fruits induced a dose-dependent decrease of plasma glucose in streptozotocin-induced and insulin-resistant diabetic rats (Hsu et al. 2000). 6348-6352
- Phytochemistry:** The entire plant contains iodine. The leaves contain alkaloids, ascorbic acid, essential oil (with limonene, carveol, and α -ionone being the major constituents), sesquiterpenoids (xanthanine, xanthanol, xanthosine, xanthamine, xanthinine, xanthumanol, and xanthinosin), phenolic acids (caffeic), chalcones, tannins, steroids, (β - and ϵ -sitosterin) saponins and carotenoids. The fruits contain drying fatty oil, resins, flavonoids, alkaloids and the glycoside xanthostrumarin (Khodzhimatov 1989; Marco et al. 1993; Belodubrovskaya et al. 2002). 6353-6357

6359 **Ziziphora bungeana Juz. – Lamiaceae**

6360 **Synonyms:** Some authors consider this species synonymous with *Ziziphora clinopodioides* Lam., *Ziziphora clinopodioides*
6361 ssp. *bungeana* (Juz.) Rech. f.

6362 **English name:** Unknown

6363 **Russian name:** Зизифора Бунге (Zizifora Bunge)

6364 **Uzbek name:** Kiyik ut

6365 **Kyrgyz name:** Кокомерен (Kokomeren)

6366 **Description:** Perennial subshrub, with woody roots. Stems many, 12–30 cm tall, branched, bases woody, densely retrorse
6367 pubescent towards apex. Leaves opposite, simple, short-petiolate, 5–15 mm long, 1.5–6 mm wide, narrowly lanceolate to
6368 ovate-lanceolate, glandular, margins entire. Inflorescences verticillasters, crowded into semiglobose, terminal heads.
6369 Calyx tubular. Corolla pink, 2-lipped; upper lip entire; lower lip 3-lobed. Fruits smooth, ovoid nutlets.

6370 **Other distinguishing features:** Plant has a strong smell. Two longer, fertile stamens, reaching upper corolla lip, and two
6371 reduced or absent stamens.

6372 **Phenology:** Flowers in July, fruits in August.

6373 **Reproduction:** By seeds.

6374 **Distribution:** Jalal-Abad, Ysyk-Kol, Naryn, and Chuy provinces of Kyrgyzstan; Tashkent province of Uzbekistan.

6375 **Habitat:** On stony slopes.

6376 **Population status:** Common, found in small groups.

6377 **Traditional use:** Extracts and infusions of the aboveground parts are recommended for hypertonia, for cardiac and climac-
6378 teric neurosis, rheumacarditis with poor blood circulation, and rheumatic endomyocarditis of children in the active phase
6379 of illness. A decoction of the leaves is used to treat gastric colic, nausea, to stimulate the appetite, and as a diuretic. It is
6380 used externally to treat throat illnesses in children. An extract of the flowers is used to treat gastritis, frequent vomiting,
6381 and meteorism (Dobrokhotova and Chudinov 1966; Plant Resources of the USSR 1991).

6382 **Documented effects:** In experiments on animals, an extract, infusion and decoction possessed hemostatic properties, raised
6383 the activity of respiratory enzymes during hypoxia, had positive influence on collateral coronary blood flow and showed
6384 prophylactic activity for, and effective treatment of, myocardial infarctions and myocarditis. In experiments, the total
6385 alkaloids showed cardiotoxic properties. The preparation *Ziziphorine* has antiarrhythmic properties on model ventricular
6386 arrhythmia in dogs, and has cardiotoxic and hypotensive actions (Plant Resources of the USSR 1991). The essential oil
6387 exhibited antibacterial activity against *Staphylococcus epidermidis*, *S. aureus*, *Escherichia coli*, and *Bacillus subtilis*
6388 (Sonboli et al. 2006).

6389 **Phytochemistry:** The roots contain organic acids, essential oils, saponins, alkaloids, vitamin C, flavonoids, and tannins.
6390 Aboveground parts contain essential oils, triterpenoids, alkaloids, flavonoids, and tannins (Dobrokhotova and Chudinov
6391 1966). The essential oil contains over 32 components with pulegone, isomenthone, 1,8-cineole and piperitenone as the
6392 main constituents (Sonboli et al. 2006).

Ziziphora clinopodioides Lam. – Lamiaceae

Synonyms: *Ziziphora afghanica* Rech. f., *Ziziphora borzhomica* Juz., *Ziziphora brevicalyx* Juz., *Ziziphora bungeana* Juz.,
Ziziphora clinopodioides ssp. *afghanica* (Rech. f.) Rech. f., *Ziziphora clinopodioides* ssp. *bungeana* (Juz.) Rech. f., *Ziziphora*
denticulata Juz., *Ziziphora dzhavakhshvili* Juz., *Ziziphora turcomaica* Juz.

English name: Unknown

Russian name: Зизифора пахучковидная (*Zizifora pakhuchkovidnaya*)

Uzbek name: Kiyik ut

Kyrgyz name: Кокомерен (*Kokomeren*)

Description: Perennial subshrub, with woody roots. Stems many, 8–40 cm tall, bases woody, rarely branched, densely ret-
 orse pubescent towards top. Leaves opposite, simple, petiolate, 6–25 mm long, 3–12 mm wide, broadly elliptic, ovate or
 elongate-ovate, glandular, margins entire or slightly toothed. Inflorescences are verticillasters, crowded into semiglobose,
 terminal heads. Calyx tubular. Corolla lilac, 2-lipped; upper lip entire; lower lip 3-lobed. Fruits smooth, ovoid nutlets.

Other distinguishing features: Plant has a strong smell. Two longer, fertile stamens, reaching upper corolla lip, and two
 reduced or absent stamens.

Phenology: Flowers in June, fruits in August.

Reproduction: By seeds.

Distribution: Naryn and Chuy provinces of Kyrgyzstan; Toshkent, Jizzax, Samarqand, Qashqadaryo, and Surxondaryo
 provinces of Uzbekistan.

Habitat: On stony slopes of mountains and gorges in spruce forests and the subalpine zone.

Population status: Common, found in small groups.

Traditional use: In Kyrgyzstan, an infusion and decoction is used to treat tachycardia, gastralgia, and heart illnesses with
 swelling. Juice from the plant is used as a vermifuge for pinworm in children (Alimbaeva and Goncharova 1971). In the
 Altai region of Russia, a tincture is used to treat common colds, rheumatism, and scrofula and it is used externally to treat
 toothaches. In Indian medicine an infusion of the leaves is used as an antipyretic and a decoction is used to treat typhoid
 fever (Plant Resources of the USSR 1991).

Documented effects: A tincture of the herb possesses hypotensive, cardiogenic, and antihelminthic properties. An 8 and
 10 % water solution of the total flavonoids possesses hypotensive properties (Alimbaeva and Goncharova 1971). The
 essential oil shows antibacterial and fungicidal activity (Delova and Guskova 1974). In experiments with mice, pretreat-
 ments with extracts of the plant reduced the biochemical, macro-, and microscopic effects of induced inflammatory bowel
 disease (Ghafari et al. 2006). Extracts the plant showed significant antibacterial activity against *Staphylococcus aureus*
 and *Pseudomonas aeruginosa*. The essential oil showed antibacterial activity against *Staphylococcus epidermidis*, *S.*
saprophyticus, *Escherichia coli*, *Shigella flexneri*, and *Salmonella typhi* (Bazzaz and Haririzadeh 2003; Tabatabaei-
 Anaraki et al. 2007).

Phytochemistry: The aboveground parts, collected during flowering stage, contained essential oil with limonene, menthone,
 isomethone, isomenthol, and pulegone as the main constituents (Korolyuk et al. 2002). Twenty-six compounds were iso-
 lated from the essential oil of plant material collected in Iran. The major components were pulegone and piperitenone
 (Mohammadreza 2008). In other collections from Iran the main compounds were: thymol, p-cymene and carvacrol, or
 1,8-cineole and terpinen-4-ol (Tabatabaei-Anaraki et al. 2007). The aboveground parts also contain saponins, coumarins,
 and flavonoids. The seeds contain fatty oil (palmitic, oleic, stearic, linoleic, and linolenic; Plant Resources of the USSR
 1991).

6393
6394
6395
6396
6397
6398
6399
6400
6401
6402
6403
6404
6405
6406
6407
6408
6409
6410
6411
6412
6413
6414
6415
6416
6417
6418
6419
6420
6421
6422
6423
6424
6425
6426
6427
6428
6429
6430
6431
6432
6433

6434 **Ziziphora pedicellata Pazij & Vved. – Lamiaceae**6435 **Synonyms:** None6436 **English name:** Unknown6437 **Russian name:** Зизифора цветоножечная (Zizifora tsvetonozhchnaya)6438 **Uzbek name:** Kiik ut6439 **Kyrgyz name:** Гулсапчалуу кокомерен (Gulsapchaluu kokomeren)

6440 **Description:** Perennial herb. Stems 20–40 cm tall, numerous, slightly winding. Leaves opposite, short-petiolate, lanceolate,
6441 glabrous or with short, spreading hairs. Flowers with long, hairy pedicels, in verticillasters crowded into head-like
6442 inflorescences. Calyx tubular, hairy; slightly 2-lipped, upper lip with 3 teeth, lower lip with 2 teeth. Corolla light-violet,
6443 2-lipped, upper lip entire, lower lip 3-lobed, tube surpassing the calyx. Fruits smooth nutlets, almost brown.

6444 **Other distinguishing features:** When rubbed, leaves and flowers produce a strong menthol aroma.6445 **Phenology:** Flowers in June-August and fruit July-September.6446 **Reproduction:** By seeds.6447 **Distribution:** Toshkent province of Uzbekistan; Western Tien-Shan; not found in Kyrgyzstan.6448 **Habitat:** The tau zone. Stony slopes with rocky debris.6449 **Population status:** Common.

6450 **Traditional use:** In Central Asia and Kazakhstan a tincture and decoction of the aboveground parts are used as a diuretic and
6451 the fresh ground plant is used to heal wounds. In Uzbekistan an infusion of the herb, taken as a tea, is used as a hypoten-
6452 sive and to treat headaches (Khalmatov 1964; Gusakova and Khomova 1997; Sezik et al. 2004).

6453 **Documented effects:** In pharmacological studies, infusions, tinctures and liquid extracts of this plant had positive effects on
6454 myocarditis and myocardial infarction. The same preparations acted as a cardiotonic, decreased arterial pressure, and
6455 increased diuresis (Khalmatov 1964).

6456 **Phytochemistry:** The plant contains essential oil composed of pulegone, pinene, menthol, menthone, isomenthone, alco-
6457 hols, and other substances. The seeds and leaves contain carotenoids and lipids. The flowers contain terpenes (Khalmatov
6458 1964; Gusakova and Khomova 1997).

6459



► **Xanthium strumarium L.**
Photos: *center*: Stan Shebs;
left and right: Clinton Shock
▼ **Ziziphora
clinopodioides Lam.**
Photos: Evgeny Davkaev



▼ **Ziziphora pedicellata Pazij & Vved.**
Photos: Alim Gaziev



6460 **Ziziphora tenuior L. – Lamiaceae**

6461 **Synonyms:** *Faldermannia parviflora* Trautv.

6462 **English name:** Unknown

6463 **Russian name:** Зизифора тонкая (*Zizifora tonkaya*)

6464 **Uzbek name:** Chul yalpiz

6465 **Kyrgyz name:** Ичке кокомерен (*Ichke kokomeren*)

6466 **Description:** Annual herb. Stems erect, unbranched or branching from the base, 5–30 cm tall, curly-hairy. Leaves opposite, linear-lanceolate to lanceolate, apex acuminate, base attenuate, the edges and abaxial side along veins curly-hairy, margin entire, short-petiolate; upper leaves ciliate. Inflorescences axillary verticillasters, usually 2–6-flowered, arranged into spikes. Calyx tubular, slightly curving downwards, obscurely 2-lipped, spreading-hairy, upper lip 3-toothed, lower lip 2-toothed. Corolla light violet, 2-lipped; upper lip entire; lower lip 3-lobed, spreading; tube noticeably protruding past the calyx. Fruits oblong-linear nutlets, 1.5 mm long, 3-edged, brown.

6472 **Other distinguishing features:** Leaves in the inflorescences much longer than the flowers. Plant produces a strong menthol aroma when crushed.

6474 **Phenology:** Flowers in May-June, fruits in June-August.

6475 **Reproduction:** Only by seeds.

6476 **Distribution:** All provinces of Uzbekistan and Kyrgyzstan.

6477 **Habitat:** The chul, adyr, and tau zones. Clay and stony soils.

6478 **Population status:** Common in *Artemisia*-ephemeral complexes, sometimes makes small populations.

6479 **Traditional use:** A decoction of the herb is recommended for intestinal diseases, diarrhea, children's colitis, neurasthenia and for maintaining cardiac activity (Khalmatov 1964).

6481 **Documented effects:** Pulegone, which is isolated from the essential oil, is reduced to produce menthol (Ogolevitz 1951). Extracts of the plant exhibited antifungal and antibacterial activity (Sardari et al. 1998; Tajadod and Majd 2007).

6483 **Phytochemistry:** Plants contain 0.3–1 % essential oils which consist of 75–87.1 % pulegone (Khalmatov 1964; Salehi et al. 2005).

6485

Ziziphus jujuba Mill. – Rhamnaceae**Synonyms:** *Rhamnus zizyphus* L., *Ziziphus sativa* Gaertn., *Ziziphus vulgaris* Lam.**English name:** Jujube, Chinese date**Russian name:** Унаби (Unabi)**Uzbek name:** Unaby, Chylon jiida**Kyrgyz name:** Кадимки унаби (Kadimki unabi)

Description: Shrub or small tree usually to 3–4(–10) m high, with or without spines. Bark brown or gray-brown. New branches purple-red or gray-brown, flexuose, with 2 stipular spines or not; long spines erect, stout, to 3 cm; short spines recurved; annual branchlets pendulous, green, resembling compound leaves, solitary or 2–7-fascicled on short shoots. Leaves alternate, short-petiolate with small spinose stipules at the base, oblong-ovate to broadly-lanceolate, rounded or slightly cordate and unequal at the base, prominently 3-veined, coriaceous, glabrous, dark green and shiny above, pale green below, margins crenate-serrate. Inflorescences axillary cymes on very short peduncles. Flowers 3–4 mm in diameter, with fleshy disk. Sepals 5, ovate-triangular. Petals 5, greenish-yellow, obovate, clawed at base. Stamens 5. Fruit a drupe, globular or oblong, reddish-orange to red-purple.

Other distinguishing features: Leaves have anesthetic effect when chewed, and causes inability to taste sugar, salt and pepper for 1–2 min.

Phenology: Flowers and fruits in July-September.**Reproduction:** By seeds and rhizomes.

Distribution: Naturally occurs in Surxondaryo province, but is cultivated throughout Uzbekistan; Jalal-Abad province of Kyrgyzstan.

Habitat: The tau zone. Dry slopes with rocky debris.**Population status:** Uncommon, occurs in small populations and as solitary individuals.

Traditional use: Fruits are used for catarrh of the upper airways, fevers, and to treat intestinal infections. The root bark is used as a stimulant and the fruits have antibacterial action. In Central Asia, a decoction of the fruit is used for anemia, chest pains, asthma, coughs, smallpox, diarrhea, and as an analgesic for diseases of the liver, kidneys, and intestines and also as hypotensive drug. In China, the preparation *landutzao* is made by processing the fruits of this species with steam from water in which *Aconitum leucostomum* L. has been boiled. This preparation is used to treat tuberculosis, lymph nodes, bones, skin, eyes, and lungs (Sakhobiddinov 1948; Gammerman et al. 1990).

Documented effects: As a result of pharmacological studies, fruits and leaves in a 10 % infusion were recommended as a medical treatment for its hypotensive and diuretic effects. In the therapeutic clinic of Samarqand Medical Institute, preparations of jujube fruits have shown positive results for the treatment of hypertensive patients (Akopov 1981; Gammerman et al. 1990). Betulinic acid and a fatty acid mixture of linoleic, oleic and stearic acids isolated from extracts of the seeds showed moderate and significant levels of cyclooxygenase-2 inhibition respectively (Su et al. 2002). The flavonoids spinosin and swertish, isolated from the seeds, exhibited significant sedative effects (Cheng et al. 2000). In vivo experiments with mice showed that an extract of the seeds possessed anxiolytic effects at lower dose and sedative effects at higher dose (Peng et al. 2000). Triterpenoids isolated from the fruit exhibited high cytotoxic activity against a number of different tumor cell lines (Lee et al. 2003).

Phytochemistry: Triterpenoid saponins, triterpenoids, flavonoids, and alkaloids have been isolated from species in this genus (Li et al. 2005). Leaves contain 27–30 % tannins (including 15 % pure tannin), tetra-saccharide, methyl ether of gallic acid, and free gallic acid. The leaves also contain myricitrin and other flavonoids, up to 0.01 % of essential oils, up to 122 mg% of vitamin C, and pigments. The fruit coat contains tannins. The fruits contain micro and macro-elements (iron, iodine, zinc, copper, cobalt, and others), as well as triterpenoids (Akopov 1981; Gammerman et al. 1990; Lee et al. 2003). Over 22 different compounds have been isolated from the seeds including flavonoids, phenyl glycosides, triterpenes, and alkaloids (Cheng et al. 2000; Li et al. 2005).

6486

6487

6488

6489

6490

6491

6492

6493

6494

6495

6496

6497

6498

6499

6500

6501

6502

6503

6504

6505

6506

6507

6508

6509

6510

6511

6512

6513

6514

6515

6516

6517

6518

6519

6520

6521

6522

6523

6524

6525

6526

6527

6528

6529

6530

6531 **Zygophyllum oxianum** Boriss. – **Zygophyllaceae**6532 **Synonyms:** *Zygophyllum fabago* L. var. *oxianum* (Boriss.) Kitam.6533 **English name:** Beancaper6534 **Russian name:** Парнолистник амударьинский (Parnolistnik amudar'inskiy)6535 **Uzbek name:** Tujatovan, It tovon6536 **Kyrgyz name:** Unknown

6537 **Description:** Herbaceous perennial with a thick, woody, vertical root. Stems few, erect, 30–70 cm tall, divaricate-branched
6538 above, thick, striated, glabrous. Leaves opposite, compound, with 1 pair of leaflets; stipules 4–7 mm long; leaflets obliquely
6539 ovate to orbicular, flat, fleshy, up to 3–4 cm long. Flowers singular or paired in upper leaf axils, pedicels 1–1.2 cm long.
6540 Sepals 5. Petals 5, oblong, ca. 1 cm long, lower half orange-red, top white, apex rounded. Stamens 10, orange. Fruits
6541 oblong-cylindrical capsules, 1.5–2 cm long, sharp-angular with 5 ribs and 5 grooves, erect. Seeds 5–8 mm long, glabrous,
6542 gray.

6543 **Other distinguishing features:** Differs from *Zygophyllum fabago* which has longer, drooping fruits.6544 **Phenology:** Flowers and fruits in May-August.6545 **Reproduction:** Most often by seeds, and rarely by rhizomes.6546 **Distribution:** All provinces of Uzbekistan; not found in Kyrgyzstan.6547 **Habitat:** The chul and adyr zones. Primary habitats are river floodplains, on slightly salty soils, and in oases of desert and
6548 semi-desert zones.6549 **Population status:** Common in typical habitats, mostly as a solitary individuals.

6550 **Traditional use:** Plasters made of fresh leaves are used to treat abscesses, as well as to heal wounds. An infusion of the
6551 leaves is used as a vermifuge and to treat fatigue and weak heart function. An extract made of this species has bacteria-
6552 cidal properties. A decoction of the root is used as a wash to treat rheumatism, wounds and carbuncles. An ointment,
6553 prepared by mixing powdered root with sheep fat, is used to treat wounds (Seredin and Sokolov 1969). In China the plant
6554 is used as a cough suppressant, expectorant, anti-inflammatory, and analgesic (Feng et al. 2007).

6555 **Documented effects:** An extract of the closely related species *Zygophyllum fabago* exhibited low activity as an acetylcho-
6556 linesterase inhibitor and exhibited much higher inhibitory activity against butyrylcholinesterase (Orhan et al. 2004).
6557 Extracts also showed very strong antifungal activity against *Candida albicans* and significant antibacterial activity against
6558 *Escherichia coli* and *Bacillus subtilis* (Zaidi and Crow 2005). Compounds isolated from a bark extract exhibited anti-tu-
6559 mor activity (Feng et al. 2007).

6560 **Phytochemistry:** The whole plant contains up to 2 % alkaloids, the main ones being zygofabagine, harmine, and others.
6561 Leaves contain 15.7–70 mg% and fruits up to 10 mg% of vitamin C (Seredin and Sokolov 1969). The bark contains trite-
6562 penoid glycosides, quinovic acid and its derivatives as well as a cincholic acid derivative (Feng et al. 2007).



▲ *Ziziphora tenuior* L. Photos: Evgeny Davkaev

◀ *Ziziphus jujuba* Mill. Photos: Alim Gaziev

▼ *Zygophyllum oxianum* Boriss.

Photos: Alim Gaziev



Author Query

Chapter No.: 5 Eisenman_Ch05

Query	Details Required	Author's Response
AU1	"Karimov et al. 1976" and "Karimov et al. 1977" are cited in text, but not provided in reference list. Kindly advice.	

Uncorrected Proof

1 Appendix 1

2 **English-Russian Translations of Botanical**
3 **and Ecological Terms**

t1.1	Abandoned field	зброшенная пашня (залеж)
t1.2	Abundant	обильный
t1.3	Achene	семянка
t1.4	Acuminate	заострённый
t1.5	Acute	острый, заострённый
t1.6	Adnate	сросшийся
t1.7	Aggregate fruit	сложный плод
t1.8	Alluvial deposits	аллювиальные наносы
t1.9	Along	вдоль
t1.10	Alpine	альпийский
t1.11	Alternate	очерёдный
t1.12	Amplexicaul	стеблеобъемлющий (лист)
t1.13	Angled	гранистый
t1.14	Angular	угловатый
t1.15	Annual	однолетний
t1.16	Anther	пыльник
t1.17	Apex	верхушка
t1.18	Apical	верхушечный
t1.19	Apiculate	с коротким узкозаострённым
t1.20		концом
t1.21	Appendage	придаток
t1.22	Appressed	прижатый
t1.23	Arachnoid-hairy	паутинисто-опушённый
t1.24	Arching, arcuate	дуговидный
t1.25	Arcuate, arching	дуговидный
t1.26	Aril	присемянник
t1.27	Artemisia-grass complex	полынно-разнотравный фитоценоз
t1.28	Ascending	приподнимающиеся
t1.29	Attenuate	суженный
t1.30	Auricles	ушки (листа)
t1.31	Auriculate	ушковидный
t1.32	Awl-shaped	шиловидный
t1.33	Awn	ость
t1.34	Axil	пазуха
t1.35	Axillary	пазушные
t1.36	Banner petal	флаг
t1.37	Barb	шип

Bark	кора
Basal leaves	прикорневые листья
Beak	носик (клюв)
Belt-like	ремневидный
Beneath	снизу
Berry	ягода
Biennial	двухлетний
Bifurcating	раздвоенный
Bipinnate	двуперистый
Bipinnatipartite	двуперистораздельный
Bipinnatisect	дважды перисторассечённый
Bisexual	обоеполюй
Biternate	двоякотройчатый
Bitter	горький
Blade	пластинка
Blunt	тупой
Bog	болото
Bract	прицветник
Bracteate	с прицветниками
Bracteole	прицветник на вторичной оси
Bractlet	прицветничек
Branch	ветка
Branched	ветвистый
Branchlet	веточка
Bristly	щетинистый
Bristles, setae	щетины
Broadly-oblong	широко-продолговатый
Brook, stream	сай, ручей, маленькая речка
Brook, stream	ручей, сай, маленькая речка
Buds (flower)	бутоны
Bulb	луковица
Bulblet	луковичка
Burst	лопаться
Bush	кустарник
Calyx	чашечка (цветка)
Campanulate	колокольчатый
Canyon bottoms	дно ущелий
Capitate	головчатый
Capitulum	корзинка (форма соцветия)
Capsule	коробочка (форма соцветия)
Carpel	плодолистик

t1.38	Cartilaginous	хрящеватый	Cup-shaped	бокальчатый	t1.94
t1.39	Catkin	серёжка (форма соцветия)	Curling	вьющийся	t1.95
t1.40	Caudex	каудекс	Curly	курчавый	t1.96
t1.41	Cemetery	кладбище	Curly-hairy	курчаво-опушённый	t1.97
t1.42	Chain	цепочка	Curved	изогнутый	t1.98
t1.43	Chamber	камера	Cyathium	циатий	t1.99
t1.44	Ciliate	бахромчатый, ресничатый	Cylindrical	цилиндрический	t1.100
t1.45	Ciliate	ресничатый, бахромчатый,	Cyme	полузонттик (соцветие)	t1.101
t1.46	Circumscissile	открывается по круговой линии	Deciduous	опадающий	t1.102
t1.47	City	город	Deciduous	теряющий на зиму листву	t1.103
t1.48	Cladodes	кладодий	Decumbent	приподнимающийся	t1.104
t1.49	Clasping	охватывающий, стеблеобъемлющий	Decurrent	нисходящий	t1.105
t1.50	Clasping	стеблеобъемлющий, охватывающий	Deep	глубокий	t1.106
t1.51	Claw	ноготок	Deeply dissected	глубоко рассечённый	t1.107
t1.52	Clay bluff	глинистый обрыв	Dehiscent (fruits)	растрескивающийся плод	t1.108
t1.53	Clayey	глинистый	Densely	густой	t1.109
t1.54	Clay-soiled	глинистая почва	Dentate	зубчатый	t1.110
t1.55	Cleistogamous	клеистогамный	Desert	пустыня	t1.111
t1.56	Climbing	цепляющиеся	Diadelphous	двубратственный	t1.112
t1.57	Club-shaped	булавовидный	Diameter	диаметр	t1.113
t1.58	Cluster	гроздь	Dichasium	дихазий	t1.114
t1.59	Cluster (flowers)	кисть (соцветие)	Dimorphic	диморфный	t1.115
t1.60	Coarse-dentate	крупно-зубчатый	Dioecious	двудомное (растение)	t1.116
t1.61	Coat	оболочка	Disc flowers	дисковые цветки	t1.117
t1.62	Compact	скупенный	Disk flower	трубчатый цветок	t1.118
t1.63	Compacted (soil)	хрящеватая (почва)	Disk-shaped	дискообразный	t1.119
t1.64	Compound leaf	сложный лист	Dispersal of seeds	распространение семян	t1.120
t1.65	Compound umbel	сложный зонтик	Dissected	рассечённый	t1.121
t1.66	Compressed	сжатая	Divaricate-branched	вильчато-ветвистый	t1.122
t1.67	Cone (berry-like)	шишкоягода (плод)	Dots	точки	t1.123
t1.68	Cone [pine type]	шишка	Drooping	поникающий	t1.124
t1.69	Cone [shape]	конус	Drupe	костянка (плод)	t1.125
t1.70	Conical	конический	Drupelet	костяночка	t1.126
t1.71	Conjoined	многоглавый	Dry	сухой	t1.127
t1.72	Connate	сросшийся	Dry river-bed	сухое русло реки	t1.128
t1.73	Constricted	перетянутый	Dull	матовый	t1.129
t1.74	Convex	выпуклый	Elaiosome	элайосома	t1.130
t1.75	Cordate	сердцевидный	Elliptic	эллиптический	t1.131
t1.76	Coriaceous	кожистый	Elongate	удлинённый, продолговатый	t1.132
t1.77	Corolla	венчик	Emarginate	выемчатый	t1.133
t1.78	Corona	корона	Embankment	насыпь	t1.134
t1.79	Corymb	щиток	Endocarp	эндокарпий	t1.135
t1.80	Corymbiform	щитковидный	Entire (margin)	цельный	t1.136
t1.81	Corymbiform cyme	щитковидный ползотик	Ephemeral	эфемерный	t1.137
t1.82	Cotton field	хлопковое поле	Ephemeroid	эфемероидный	t1.138
t1.83	Crack	трещина	Epicalyx	наружная чашечка	t1.139
t1.84	Cracked	трещиноватый	Erect	прямо стоящий	t1.140
t1.85	Creeping	ползучий	Even-pinnate	парноперистый	t1.141
t1.86	Creeping roots	корнеотпрысковый	Even-pinnate	парноперистый	t1.142
t1.87	Crenate	городчатый (лист)	Evergreen	вечнозелёный	t1.143
t1.88	Crescent-shaped	серповидный	Explosively dehiscing	вскрывается	t1.144
t1.89	Crowded	скупенный	Exserted	выступающий	t1.145
t1.90	Crown	крона	Farinose	покрыт мучнистым налётом	t1.146
t1.91	Cultivated	культивируемый	Fascicle	пучок	t1.147
t1.92	Cultivated fields	посевы	Feather-grass steppes	ковыльные степи	t1.148
g1.93	Cuneate	клиновидный	Felted	войлочный	g1.149

t1.150	Felted-hairy	войлочно опушённый	Hood	шлем (часть цветка)	t1.206
t1.151	Female flower	женский цветок	Hooked	крюкообразный	t1.207
t1.152	Fibrous roots	мочковатые корни	Horizontal	горизонтальный	t1.208
t1.153	Field	поле	Horn	рожок	t1.209
t1.154	Filiform	нитевидный	Horn-like	роговидный	t1.210
t1.155	Fimbriate, fringed	бахромчатый	Hypanthium	гипантий	t1.211
t1.156	Finger-like	пальчатообразный	Imbricate	чешуйчатый	t1.212
t1.157	Fir	пихта	Incised	надрезанный	t1.213
t1.158	Fissure	трещина	Incrassate	утолщённый	t1.214
t1.159	Flat	плоский	Incurved	внутри изогнутый	t1.215
t1.160	Flattened	сплюснутый	Indehiscent	нераскрывающиеся (плоды)	t1.216
t1.161	Fleshy	мясистый	Inflated	вздутый	t1.217
t1.162	Flexuose	извилистый	Inflorescence	соцветие	t1.218
t1.163	Floodplain	пойма (реки)	Inflorescence axis	ось соцветия	t1.219
t1.164	Fluted	желобчатый	Inserted	расположенный	t1.220
t1.165	Foliaceous	листовидный	Interrupted	прерывистый	t1.221
t1.166	Follicle	листовка (плод)	Involucel	вторичная обёртка (цветка)	t1.222
t1.167	Foothills	предгорье	Involucral bract	листочек обёртки	t1.223
t1.168	Forest	лес	Involucre	обёртка соцветия	t1.224
t1.169	Forest edges	опушка леса	Irregular	неправильный	t1.225
t1.170	Forked	вилчатый	Irrigated	орошаемый	t1.226
t1.171	Four-sided (4-sided)	четырёхгранный	Irrigation canal	арык	t1.227
t1.172	Fragrant	душистый	Juicy	сочный	t1.228
t1.173	Fringed, fimbriate	бахромчатый	Juniper	арча	t1.229
t1.174	From the base	от основания	Juniper stand	арчëвник	t1.230
t1.175	Fruit	плод	Keel	лодочка, киль	t1.231
t1.176	Funnelform	воронковидный	Keel	киль, лодочка	t1.232
t1.177	Furrowed	бороздчатый	Keeled	килевидный	t1.233
t1.178	Fusiform	веретеновидный	Kidney-shaped	почковидный	t1.234
t1.179	Flabrous	голый	Lake	озеро	t1.235
t1.180	Glade	поляна	Lanceolate	ланцетный	t1.236
t1.181	Glands	железки	Large-dentate	крупнозубчатый	t1.237
t1.182	Glandular	железистый	Lateral	боковой	t1.238
t1.183	Glandular prickles	шиповидные железки	Lawn	лужайка	t1.239
t1.184	Glaucous	покрытый налётом	Leaf	лист	t1.240
t1.185	Globular	шаровидный, сферический	Leaflet	листочек	t1.241
t1.186	Glossy, shiny	блестящий	Leafy	облиственный	t1.242
t1.187	Gorge	ущелье	Legume	боб	t1.243
t1.188	Gradually	постепенно	Lenticel	чечевичка	t1.244
t1.189	Granular	гранулярный	Lenticular	чечевицеобразный	t1.245
t1.190	Groove	борозда	Limestone talus	известняковая осыпь	t1.246
t1.191	Gum, resin, pitch	камедь, смола	Linear	линейный	t1.247
t1.192	Gypsum	гипс	Lip	губа	t1.248
t1.193	Habitat	местообитание	Loam	суглинок	t1.249
t1.194	Hairs	волоски	Lobe	доля, лопасть	t1.250
t1.195	Hairy	опушённый	Lobe	лопасть, доля	t1.251
t1.196	Hanging	пониклый	Lobed	лопастной, дольчатый	t1.252
t1.197	Hastate	стреловидный	Lobed	дольчатый, лопастной	t1.253
t1.198	Head (inflorescence)	головка (соцветия)	Lobule	долька	t1.254
t1.199	Head-like	головчатовидный	Loess	лëсс	t1.255
t1.200	Helicoid	спиралеобразный	Long	длина	t1.256
t1.201	Herbaceous plant	травянистое растение	Long-fallow field	перелог	t1.257
t1.202	Heterogamous	гетерогамный	Longitudinal	продольный	t1.258
t1.203	Hill	холм	Loose	рыхлый	t1.259
t1.204	Hollow	полый	Lyrate	лировидный	t1.260
t1.205	Honey	мёд	Male flower	мужской цветок	t1.261

t1.262	Male inflorescence	мужское соцветие	Papilla	сосочек	t1.318
t1.263	Many-flowered	многоцветковый	Papillate	бородавчатый	t1.319
t1.264	Many-lobed	многолопастной	Pappus	хохолок	t1.320
t1.265	Margin entire	цельнокройный	Parallel	паралельный	t1.321
t1.266	Marginal flowers	краевые цветки	Pasture	пастбище	t1.322
t1.267	Meadow	луг	Pebbly	галечниковый	t1.323
t1.268	Melon	дыня	Pedicel	плодоножка, цветоножка	t1.324
t1.269	Melon field	бахчёвое поле	Pedicel	цветоножка, плодоножка	t1.325
t1.270	Membranaceous	плёнчатый	Peduncle	цветонос	t1.326
t1.271	Membranous	перепончатый	Pellucid dots	исколотый	t1.327
t1.272	Mericaip	мерикаип	Pendulous	плакучий	t1.328
t1.273	Milky	молочный	Pepo	тыквина	t1.329
t1.274	Milky sap	млечный сок	Perennial	многолетний	t1.330
t1.275	Moniliform	чёткообразный	Perfect (flower)	обоепольный (цветок)	t1.331
t1.276	Monocarpic	монокарпический	Perianth	околоцветник	t1.332
t1.277	Monoecious	однодомное (растение)	Pericarp	околоплодник	t1.333
t1.278	Mountain	гора	Petal	лепесток	t1.334
t1.279	Mucilage (plant)	слизи (растений)	Petaloid	лепестковидный	t1.335
t1.280	Mucronulate	маленькое острое окончание	Petiolate	черешковый	t1.336
t1.281		(листа)	Petiole	черешок	t1.337
t1.282	Narrow	узкий	Pinnate	перистый (лист)	t1.338
t1.283	Nectary	нектарник	Pinnatifid	перистонадрезанный (лист)	t1.339
t1.284	Needle-like	игловидный	Pinnatilobate	перистолопастной (лист)	t1.340
t1.285	Nerve, vein	жилка (растения)	Pinnatipartite	перисторазделный (лист)	t1.341
t1.286	Nodding	пониклый	Pinnatisect	перисторассечённый (лист)	t1.342
t1.287	Nutlet	орешек (плод)	Pistil	пестик (цветка)	t1.343
t1.288	Oasis	оазис	Pistillate flower	женский цветок	t1.344
t1.289	Obconical	обратноконический	Pitch, gum, resin	камедь, смола	t1.345
t1.290	Oblanceolate	обратноланцетный	Pith	сердцевина (стебля)	t1.346
t1.291	Obliquely descending	скошенный	Pitted	ямчатый	t1.347
t1.292	Oblong, elongate	продолговатый	Placenta	плацента	t1.348
t1.293	Obovate	обратнояйцевидный	Plain	равнина	t1.349
t1.294	Obtuse	притупленный	Plate	пластинка	t1.350
t1.295	Ocrea	раструб	Plicate	складчатый	t1.351
t1.296	Odd-pinnate	непарнопирестый	Plowed field	пашня	t1.352
t1.297	Odorous	пахучий	Plumose	перистый	t1.353
t1.298	Opposite	супротивный	Plumose-barbed	перисто-зазубренный	t1.354
t1.299	Orchard, garden	сад	Pod	стручок	t1.355
t1.300	Oriented	ориентированный	Pomaceous	яблокообразный	t1.356
t1.301	Ovary	завязь	Pore	дырочка	t1.357
t1.302	Ovate	яйцевидный, овальный	Prickle	шип	t1.358
t1.303	Ovate	овальный, яйцевидный	Prickly	шиповатый	t1.359
t1.304	Ovule	семяпочка	Prismatic	призматический	t1.360
t1.305	Paleaceous	чешуйчатый, плёнчатый	Projection	вырост	t1.361
t1.306	Paleaceous	плёнчатый, чешуйчатый	Prominent	выдающийся	t1.362
t1.307	Palmate	пальчатый	Prostrate	стелящийся	t1.363
t1.308	Palmately compound	пальчатосложный	Pubescent	волосистый	t1.364
t1.309	Palmatifid	дланевидно-надрезный	Pulp	мякоть (плодов)	t1.365
t1.310	Palmatilobate	пальмовидно-лопастный	Punctate glandular	точечные железки	t1.366
t1.311	Palmatipartite	пальчато-лопастный	Pyramidal	пирамидальный	t1.367
t1.312	Palmatisect	пальчато-рассечённый	Quadrangular	четырёхгранный	t1.368
t1.313	Panicle	метёлка (соцветие)	Quadripinnate	четыреждыперистый	t1.369
t1.314	Paniculate	метёлчатый	Raceme (cluster)	кисть (соцветие)	t1.370
t1.315	Paniculiform	метёлковидное	Racemiform	кистевидный	t1.371
t1.316	Paper-like, papery	бумагообразный	Rachis	ось	t1.372
g.317	Papilionaceous	мотыльковый (цветок)	Raised	выступающий	t1.373

t1.374	Raised gland	железистый шипик	Semispherical	полушаровидный, полусферический	t1.430
t1.375	Ray	луч			t1.431
t1.376	Ray flower	язычковый цветок	Semi-woody	полудеревянистый	t1.432
t1.377	Receptacle	цветоложе	Senescing	скороувядающий (лист)	t1.433
t1.378	Reclining	приподнимающийся	Sepal	чашелистик	t1.434
t1.379	Recurved	отогнутый вниз	Septicidal	растрескивающийся по перегородкам (плода)	t1.435
t1.380	Red sandstone	краснопесчаник			t1.436
t1.381	Reduced	редуцированный, уменьшенный	Septum	перегородка	t1.437
t1.382	Reduced	уменьшенный, редуцированный	Serrate	пильчатый	t1.438
t1.383	Reflexed	отогнутый вниз	Serrulate	мелкозубчатый	t1.439
t1.384	Resin, gum, pitch	камедь, смола	Sessile	сидячий	t1.440
t1.385	Resin, gum, pitch	смола, камедь	Setae, bristles	щетинки	t1.441
t1.386	Reticulate	сетчатый	Shady	тенистый	t1.442
t1.387	Reticulate veined	сетка жилок	Shale, slate	сланец	t1.443
t1.388	Retrorse	направленный вниз	Shallow	неглубокий	t1.444
t1.389	Rhizome	корневище, корневой отпрыск	Shallow soil	мелкоземистая почва	t1.445
t1.390	Rhobic	ромбический	Sheath	влагалище (листа)	t1.446
t1.391	Rib	ребро	Shell	скорлупа	t1.447
t1.392	Ribbed	ребристый	Shiny, glossy	блестящий	t1.448
t1.393	Ridge	рубчик	Short	короткий	t1.449
t1.394	Rind	корка	Short shoot	короткая веточка	t1.450
t1.395	Ring	кольцо	Short-petiolate	короткий черешок	t1.451
t1.396	Ripe	созревший, спелый	Shrub	кустарник	t1.452
t1.397	Ripe	спелый, созревший	Silicle	короткий стручок	t1.453
t1.398	River	река	Silique	стручок	t1.454
t1.399	River valley	долина реки	Silique-like	стручковидный	t1.455
t1.400	Road	дорога	Silky	шелковистый	t1.456
t1.401	Rocky debris	щебнистый	Simple	простой	t1.457
t1.402	Root	корень	Single, solitary	одиночный	t1.458
t1.403	Root crown	корневая шейка	Sinuate	выемчатый	t1.459
t1.404	Root system	корневая система	Slightly	слегка	t1.460
t1.405	Rosette	розетка	Slope	склон	t1.461
t1.406	Rotate	колесовидный	Small groves	небольшая роща	t1.462
t1.407	Rough	шероховатый	Smell	запах	t1.463
t1.408	Round	круглый	Smooth	гладкий	t1.464
t1.409	Row	ряд	Soft	мягкий	t1.465
t1.410	Ruderal	рудеральный	Soil	почва	t1.466
t1.411	Rugose	морщинистый	Solitary, single	одиночный	t1.467
t1.412	Runcinate	обращённые назад доли (листа)	Solonchic	солонцеватый	t1.468
t1.413	Sagittate	стреловидное основание (листа)	Sour	кислый	t1.469
t1.414	Salty area (very)	солончак	Spadix	початок	t1.470
t1.415	Sandstone	песчаник	Spathe	обвёртка	t1.471
t1.416	Sandy	песчаный	Spatulate	лопатчатый, лопатовидный	t1.472
t1.417	Sap	сок	Spatulate	лопатовидный, лопатчатый	t1.473
t1.418	Scabrid	шершавый	Spear-shaped	копьевидный	t1.474
t1.419	Scale	чешуя	Spherical	сферический, шаровидный	t1.475
t1.420	Scale-like	чешуевидный	Spherical	шаровидный, сферический	t1.476
t1.421	Scarious	пластинчатый	Spiciform	колосовидный	t1.477
t1.422	Scattered	разбросанный	Spike	колос (соцветие)	t1.478
t1.423	Schizocarp	распадающийся плод, дробный	Spine	колочка	t1.479
t1.424	Schizocarp	дробный, распадающийся плод	Spine, thorn	колочка	t1.480
t1.425	Scorpioid cyme	завиток (соцветие)	Spinescent	колочий	t1.481
t1.426	Seam	шов	Spine-tipped	оканчивающийся колочкой	t1.482
t1.427	Segment	сегмент	Spiny-toothed	колочче-зубчатый	t1.483
t1.428	Segmented	сегментированный	Spirally	винтообразный	t1.484
t1.429	Semi-desert	полупустыня	Spirally-twisted	спиралезакрученный	t1.485

t1.486	Spongy	мочалистый	Trigonus, triquetrous	трёхгранный	t1.541
t1.487	Spore	спора	Tripartite	трёхраздельный (лист)	t1.542
t1.488	Spot	пятно	Tripartite-pinnatisect	тройчато-перисторассечённый	t1.543
t1.489	Spreading branchy	оттопыренно-ветвистый	Tripinnate	трижды перистый	t1.544
t1.490	Spring	родник	Tripinnatisect	трижды перисторассечённый (лист)	t1.545
t1.491	Spruce forest	еловый лес	Triquetrous, trigonous	трёхгранный	t1.546
t1.492	Spur	шпора	Trisulcate	трёхборзчатый	t1.547
t1.493	Stamen	тычинка	Truncated	усечённый	t1.548
t1.494	Staminal column	тычиночная колонка	Trunk	ствол (растения)	t1.549
t1.495	Staminal filament	тычиночная нить	Tube	трубка	t1.550
t1.496	Staminate flowers	мужские (тычиночные) цветки	Tuber	клубень	t1.551
t1.497	Staminode	бесплодая тычинка	Tubercle	бугорок	t1.552
t1.498	Stellate	звёзчатый	Tuberculate	бугорчатый	t1.553
t1.499	Stellate-hairy	звёздно-опушённый	Tuberiform	клубневидный	t1.554
t1.500	Stem	стебель	Tubular	трубчатый	t1.555
t1.501	Steppe	степь	Tugai	тугай	t1.556
t1.502	Stiff	жёсткий	Twig-like	прутьевидный	t1.557
t1.503	Stigma	рыльце (цветка)	Twining	вьющийся	t1.558
t1.504	Stinging hairs	жгучие волоски	Two-horned	двурогий	t1.559
t1.505	Stipulate	снабжённый прилистниками	Two-lipped (2-lipped)	двугубый	t1.560
t1.506	Stipule	прилистник	Two-valved (2-valved)	двустворчатый	t1.561
t1.507	Stocky	коренастый	Umbel	зонтик (соцветие)	t1.562
t1.508	Stolon	столон	Umbellet	вторичный зонтик	t1.563
t1.509	Stony	каменистый	Umbelliform	зонтиковидный	t1.564
t1.510	Straight	прямой	Understory	подлесок	t1.565
t1.511	Stream bed	русло ручья, сая, маленькой речки	Undulate	волнистый	t1.566
t1.512	Stream, brook	ручей, сай, маленькая речка	Unequal	неравный	t1.567
t1.513	Striated	бороздчатый	Unirrigated	богара (не орошаемая зона)	t1.568
t1.514	String-like	шнуровидный	Unisexual	однополый	t1.569
t1.515	Style	столбик (цветка)	Upper	верхний	t1.570
t1.516	Suberect	приподнятый	Utricle	мешочек (плод)	t1.571
t1.517	Subshrub	полукустарник	Valley	долина	t1.572
t1.518	Succulent	сочный	Valve	створка	t1.573
t1.519	Syncarp	синкарпий	Vegetable garden	огород	t1.574
t1.520	Tail-like	хвостоподобный	Vegetation	растительность	t1.575
t1.521	Talus	осьпь	Vegetatively	вегетативный	t1.576
t1.522	Tangled	спутанный	Vein	жилка	t1.577
t1.523	Tapering	суженный	Velutinous, velvety	бархатистый	t1.578
t1.524	Tarroot	стержневой корень	Velvety, velutinous	бархатистый	t1.579
t1.525	Tendril	усик (растения)	Vertical	вертикальный	t1.580
t1.526	Tepal	листочек околоцветника	Verticillaster	полумутовка (соцветие)	t1.581
t1.527	Terminal	верхушечный	Vigorous	мощный	t1.582
t1.528	Ternate	тройчатый (лист)	Village	посёлок	t1.583
t1.529	Thick	толстый	Villous, tomentose, hairy	опушённый	t1.584
t1.530	Thin	тонкий	Vine	цепляющееся растение	t1.585
t1.531	Thorn, spine	колючка	Walnut forests	ореховый лес	t1.586
t1.532	Thread-like	нитевидный	Waste place	мусорное место	t1.587
t1.533	Three follicles	трёхлистовка (плод)	Water-eroded	смытый	t1.588
t1.534	Tip	носик (тонкий конец)	Wavy	выемчатый	t1.589
t1.535	Tomentose	опушённый	Wedge-shaped	клиновидный	t1.590
t1.536	Trailing	стелющийся (растение)	Weed	сорняк	t1.591
t1.537	Triangular	треугольный	Well	колодец	t1.592
t1.538	Trichome	трихома	Wet	сырой	t1.593
t1.539	Trifoliolate	тройчатый	Wheat field	посевы пшеницы	t1.594
t1.540	Trifurcated	трёхветвистый	Whorl	мутовка	t1.595

t1.596	Wide	шириной (в ширину)	Withering	отмирают	t1.601
t1.597	Winding	извилистый	Woody	одревенелый	t1.602
t1.598	Wing	крыло	Woolly-hairy	шерстисто опушённый	t1.603
t1.599	Winged petiole	крылатый черешок	Wrinkled	морщинистый	t1.604
t1.600	Wingless	бескрылый	Zygomorphic	зигоморфный	t1.605

Uncorrected Proof

19 **English-Russian Translations of Chemical Terms**

t2.1	25-d-spirosta-3,5-diene	25-d-спирост-3,5-диен	Amino acid	аминокисота
t2.2	2-methoxy-1,4 naphthoquinone	2-метокси-1,4-нафтохинон	Aminoalcohol	аминоспирт
t2.3	Absinthin	абсинтин	Amygdalin	амигдалин
t2.4	Acanthophylloside	аконтофиллазид	Amyl alcohol	амиловый спирт
t2.5	Acetic acid	уксусная кислота	Amyrin	амирин
t2.6	Acetylcholine	ацетилхолин	Anabasine	анабазин
t2.7	AcetylNapelline	ацетилнапеллин	Anabsinthin	анабсинтин
t2.8	Acevaltrate	ацевалтрат	Anagirine	анагирин
t2.9	Aconitic acid	акотиновая кислота	Anagyrene	анагирин
t2.10	Acsinatine	аксинатин	Anchusa acid	анхузовая кислота
t2.11	Aesculin	эскулин	Anchusin	анхизин
t2.12	Aglycone	агликон	Anemonin	анемонин
t2.13	Aksine	аксин	Anethole	анетол
t2.14	Akuamidine	акуаммидин	Angelic acid	анисовая кислота
t2.15	Akuamine	акуамин	Anhydroaustricine	ангидроаустрицин
t2.16	Alantolactone	алантолактон	Anhydroperforine	ангидроперфорин
t2.17	Alantone	алантон	Anisic acid	ангеликовая кислота
t2.18	Alantopicrine	алантопикрин	Anonaine	аноанин
t2.19	Alcohol	спирт	Anthocyan	антоциан
t2.20	Aldehyde	альдегид	Anthocyanidin	антоцианидин
t2.21	Alginidine	алгинидин	Anthocyanin	антоцианин
t2.22	Alginine	алгинин	Anthracene	антрацен
t2.23	Alhagidin	алхагидин	Anthraglycoside	антрагликозид
t2.24	Alhagitin	алхагитин	Anthranoyllycoctonine	антраноилликоктонин
t2.25	Aliphatic alcohol	алифатический спирт	Anthraquinone	антрахинон
t2.26	Alkaloid	алкалоид	Apigenin	апигенин
t2.27	Alkamide	алкамид	Arabinose	арабиноза
t2.28	Alkanin	алканин	Arachic acid	арахиновая кислота
t2.29	Allocryptopine	аллокриптопин	Arachidic acid	арахидиновая кислота
t2.30	Allyl-isothiocyanate	аллилизотиоцианат	Arctigenin	арктигенин
t2.31	Aloemodin	алоэмодин	Arctiin	арктиин
t2.32	Aloperine	алоперин	Arctiopicrin	арктиопикрин
t2.33	Alpha-amyrin	а-амирин	Argemonine	аргемонин
t2.34	Alteramine	альтерамин	Argentine	аргентин
t2.35	Aluminium	алюминий	Arnidiol	арнидиол
t2.36	Amaranthin	амарантин	Aromatic acid	ароматическая кислота
t2.37	Amide	амид	Aromatic aldehyde	ароматический альдигид
			Artabasin	артабсин
			Arteannuin	артеаннуин
			Artelein	артелеин

t2.38	Artelin	артелин	Carabron	караброн	t2.93
t2.39	Artemetin	артеметин	Carbohydrates	углеводы	t2.94
t2.40	Artemisinin	артемизинин	Carbolic acid	n-оксибензойная кислота	t2.95
t2.41	Asaresin	асарезен	Cardiac glycoside	сердечный гликозид	t2.96
t2.42	Asaresinol	асарезинол	Cardinolide	карденолид	t2.97
t2.43	Asaresinotannol	асарезинотанол	Carene	карен	t2.98
t2.44	Asarone	азарон	Carotene	каротин	t2.99
t2.45	Ascorbic acid	аскорбиновая кислота	Carotenoid	каротиноид	t2.100
t2.46	Ash	зола	Carvacrol	карвакрол	t2.101
t2.47	Asparagine	аспарагин	Carvone	карвон	t2.102
t2.48	Asperuloside	асперулозид	Caryophylladienol	кариофилла-диен-ол	t2.103
t2.49	Astringent substances	вяжущие вещества	Caryophyllene	кариофиллин	t2.104
t2.50	Atropine	атропин	Caryophyllene oxide	окись кариофиллина	t2.105
t2.51	Aucubin	аукубин	Catechin	катехин	t2.106
t2.52	Austricine	аустрицин	Catecholamine	катехоламин	t2.107
t2.53	Avenasterol	авенастерол	Caulosapogenin glycoside	каулосапогенин-гликозид	t2.108
t2.54	Avicularin	авикуларин	Cedrene	цедрен	t2.109
t2.55	Bactericidal	бактерицидный	Cedrol	цедрол	t2.110
t2.56	Baicalein	байкалеин	Cerotic acid	церотиновая кислота	t2.111
t2.57	Bakuchiol	бакучиол	Ceryl-alcohol	церильевый спирт	t2.112
t2.58	Barium	барий	Chatinene	хатинин	t2.113
t2.59	Behenic acid	бегеновая кислота	Chemical compound	химическое соединение	t2.114
t2.60	Benzaldehyde	бензойный альдегид	Chicoric acid	цикоревая кислота	t2.115
t2.61	Benzoic acid	бензойная кислота	Chlorogenic acid	хлорогеновая кислота	t2.116
t2.62	Benzyl isothiocyanate	бензилотиоцианат	Chlorophyll	хлорофилл	t2.117
t2.63	Berbamine	бербамин	Cholesterol	холестерин	t2.118
t2.64	Berbamunine	бербамунин	Choline	холин	t2.119
t2.65	Berberine	берберин	Chromium	хром	t2.120
t2.66	Bergamotene	бергамотен	Chromone	хромон	t2.121
t2.67	Betaine	бетаин	Chrysoeriol	хризозериол	t2.122
t2.68	Bicyclic	бициклический	Chrysophanic acid	хризофановая кислота	t2.123
t2.69	Bicyclogermacrene	бициклогермакрен	Cichoriin	цикорин	t2.124
t2.70	Biogenic amine	биогенный амин	Cicutine, coniine	коницин	t2.125
t2.71	Bitter substances	горькие вещества	Cinaroside	цинarosин	t2.126
t2.72	Bitters	горечи	Cincholic acid	хинолиновая кислота	t2.127
t2.73	Borneol	борнеол	Cineol	цинеола	t2.128
t2.74	Bornyl acetate	борнил ацетат	Cinnamaldehyde	коричный альдегид	t2.129
t2.75	Bornyl-isovalerianate	борнил-изовалерианат	Cinnamamide	циннамамид	t2.130
t2.76	Britanin	британин	Citral	цитраль	t2.131
t2.77	Bufadienolide	буфадиенолид	Citric acid	лимонная кислота	t2.132
t2.78	Bursic acid	бурсовая кислота	Citrin	цитрин	t2.133
t2.79	Cadinene	кадинен	Citronellol	цитронеллол	t2.134
t2.80	Caffeic acid	кофейная кислота	Citronellyl acetate	цитронеллиацетат	t2.135
t2.81	Caffeine	кофеин	Clematine	клематин	t2.136
t2.82	Caffeoylquinic acid	кофеоилхинная кислота	Cnicin	кницин	t2.137
t2.83	Calcium oxalate	оксалат кальция	Cobalt	кобальт	t2.138
t2.84	Campesterin	кампестерин	Codonopsin	кодонопсин	t2.139
t2.85	Campesterol	кампестерол	Codonopsinin	кодонопсинин	t2.140
t2.86	Camphene	камфена	Columbamine	колумбамин	t2.141
t2.87	Camphene	камфена	Compound	соединение	t2.142
t2.88	Camphorol	кемпферол	Condorphine	кондельфин	t2.143
t2.89	Canavanine	канаванин	Conhydrine	конгидрином	t2.144
t2.90	Caoutchouc	каучук	Convolamine	конволамин	t2.145
t2.91	Capronic acid	капроновая кислота	Convalidine	конволидин	t2.146
t2.92	Caprylic acid	каприловая кислота	Convolvine	конвольвин	t2.147

t2.148	Convolvuline	конвольвулин	Diosmine	диосмин	t2.203
t2.149	Copper	медь	Dipegene	дипегин	t2.204
t2.150	Corydine	коридин	Disaccharide	дисахаридов	t2.205
t2.151	Corytuberine	коритуберин	Disulfide	дисульфид	t2.206
t2.152	Coumaric acid	кумаровая кислота	Dopamine	допамин	t2.207
t2.153	Coumarin	кумарин	Doremol	доремол	t2.208
t2.154	Coumarinic acid	кумариновая кислота	Doremon	доремон	t2.209
t2.155	Cryptopine	криптопин	Drupacine	друпацин	t2.210
t2.156	Crystalline	кристаллический	Drupanol	друпанол	t2.211
t2.157	Crystals	кристаллы	Drying fatty oil	жирное высыхающее масло	t2.212
t2.158	Cuminaldehyde	куминовый альдегид	Dubamine	дубамин	t2.213
t2.159	Cuminyl alcohol	куминилловый спирт	Dubinidine	дубинидин	t2.214
t2.160	Cuscohygrine	кускогигрин	Ecdysone	экизон	t2.215
t2.161	Cyanidin	цианидин	Ecdysterone	экидистерон	t2.216
t2.162	Cyanidin-3-glucoside	цианидин-3-глюкозид	Eicosenoic acid	гадолеиновая кислота	t2.217
t2.163	Cyanin	цианин	Elemol	элебол	t2.218
t2.164	Cyanogenic compound	цианогенное соединение	Ellagic acid	эллаговая кислота	t2.219
t2.165	Cyanogenic glycoside	цианогенный гликозид	Emodin	эмодин	t2.220
t2.166	Cyasterone	циастерон	Enzyme	фермент	t2.221
t2.167	Cyclamine	цикламин	Enzymic hydrolysis	ферментативный гидролиз	t2.222
t2.168	Cyclic alcohol	циклический спирт	Ephedrine	эфедрин	t2.223
t2.169	Cyclic peptide	циклиновый пептид	Epi-13-manool	13-эпиманоол	t2.224
t2.170	Cyclitols	циклитолы	Epicatechin	эпикатехин	t2.225
t2.171	Cyclolignan	циклолигнан	Epigallocatechin	эпигалокатехин	t2.226
t2.172	Cyclopropenoid fatty acid	циклопропаноидная жирная кислота	Epi-rhododendrin	эпирододендрин	t2.227
t2.173			Epoxyacylglyceride	эпоксинацилглицерид	t2.228
t2.174	Cymene	цимен	Equisetin	эквишетин	t2.229
t2.175	Cymol	цимол	Equisetonin	эквишетонин	t2.230
t2.176	Cytisine	цитизин	Equisetrine	эквишетрин	t2.231
t2.177	Cytosterin	ситостерин	Eremuran	эремуран	t2.232
t2.178	Daucane esters	дауциновые эфиры	Eremursine	эремурсин	t2.233
t2.179	Daucane-type sesquiterpene	сесквитерпен, типа дауцин	Ergolide	эрголид	t2.234
t2.180	Daucene	дауцин	Eriodictyol	эриодиктиол	t2.235
t2.181	Daucosterol	даукостерол	Erucic acid	эруковая кислота	t2.236
t2.182	Dehydroabietol	дигидроабиетол	Ervamine	эрвамин	t2.237
t2.183	Dehydrothalicmine	дегидроталикмин	Ervine	эрвин	t2.238
t2.184	Delatine	делатин	Ervinine	эрвинин	t2.239
t2.185	Delphinine	дельфелин	Erysimine	эризимин	t2.240
t2.186	Delphinindin	дельфинидин	Erysimoside	эризимозид	t2.241
t2.187	Delphirine	дельпирин	Esculetin	эскулетин	t2.242
t2.188	Delsemine	дельсемин	Esculin	эскулин	t2.243
t2.189	Delsine	дельсин	Essential oil	эфирное масло	t2.244
t2.190	Delsoline	дельсолин	Ester	сложный эфир	t2.245
t2.191	Delsosine	делькозин	Ether	эфир	t2.246
t2.192	Deoxypeganane	дезоксипеганин	Ethyl	этил	t2.247
t2.193	Derivatives	производные	Ethyl ester	этиловый эфир	t2.248
t2.194	Diacylglyceride	диацилглицерид	Eucalyptol	эвкалиптол	t2.249
t2.195	Dictamine	диктамин	Eudesmine	эудесмин	t2.250
t2.196	Didrovaltrate	дидровалтрат	Eugenol	евгенол	t2.251
t2.197	Dihydroalantolactone	дигидроалантолактон	Euphorbin	эуфорбин	t2.252
t2.198	Dihydroxyacids	диоксикислота	Evodine	эводин	t2.253
t2.199	Dillapiole	диллапиол	Evoxin (haploperin)	эвоксин (хаплоперин)	t2.254
t2.200	Dimethamine	диметамин	Evoxidine	эвоксоидин	t2.255
t2.201	Diosgenin	диосгенина	Excelsine	эксцельзин	t2.256
t2.202	Diosmetin	диосметин	Extriol	экстриол	t2.257

t2.258	Faradiol	фарадиол	Glucocapparin	глюкокаппарин	t2.313
t2.259	Farnesferol	фарнезиферол	Glucofrangulin	глюкофрангулин	t2.314
t2.260	Fat-like substances	жироподобные вещества	Glucofructose	глюкофруктоза	t2.315
t2.261	Fatty acid	жирная кислота	Glucose	глюкоза	t2.316
t2.262	Fatty oil	жирное масло	Glucoside	глюкозид	t2.317
t2.263	Fenchone	фенхон	Glycerin	глицерина	t2.318
t2.264	Fermononetin	формононетин	Glycoalkaloid	гликоалкалоид	t2.319
t2.265	Ferruginol	ферругинол	Glycone	гликон	t2.320
t2.266	Ferulic acid	феруловая кислота	Glycoperine	гликоперин	t2.321
t2.267	Fiber	клетчатка	Glycoside	гликозид	t2.322
t2.268	Flavanone glycoside	флаванонгликозид	Glycyrramarin	глицирамарин	t2.323
t2.269	Flavolignan	флаволигнан	Glycyrrhetic acid	глицирретиниковая кислота	t2.324
t2.270	Flavone	флавоон	Glycyrrhizic acid	глицирризиновая кислота	t2.325
t2.271	Flavonoid	флавоноид	Glycyrrhizin	глицирризин	t2.326
t2.272	Flavonol glycoside	флавоновый гликозид	Gossypol	госсипол	t2.327
t2.273	Flavoxanthin	флавоксантин	Granilin	гранилин	t2.328
t2.274	Flindersine	флиндерсин	Guaiol	гвайол	t2.329
t2.275	Foetidine	фетидин	Gum	камедь	t2.330
t2.276	Folic acid	фоливая кислота	Haplofidine	хаплофидин	t2.331
t2.277	Foliosidine	фолиозидин	Haplofilidine	хаплофилидин	t2.332
t2.278	Formic acid	муравьиная кислота	Haplophytin	хаплофитин	t2.333
t2.279	Fractions	фракции	Haplopine	хаплопин	t2.334
t2.280	Frangula-emodin	франгулаэмодин	Harmaline	гармалин	t2.335
t2.281	Frangula-emodin anthronol	франгулаэмодинантранол	Harmalol	гармалол	t2.336
t2.282	Frangulin	франгулин	Harman	гарман	t2.337
t2.283	Free fatty acid	свободная жирная кислота	Harmine	гармин	t2.338
t2.284	Fructose	фруктоза	Harpagide	гарпагид	t2.339
t2.285	Fumaric acid	фумаровая кислота	Harpagoside	гарпагосид	t2.340
t2.286	Fumaridine	фумаридин	Hederagenin derivatives	производные хедерагенина	t2.341
t2.287	Fumvailline	фумвайлин	Hemicellulose	гемицеллюлоза	t2.342
t2.288	Furocoumarin	фурукумарин	Hemolytic index	гемолитический индекс	t2.343
t2.289	Fustin	фустин	Herniarine	герниарин	t2.344
t2.290	Galanthamine	галантамин	Hesperidin	гесперидин	t2.345
t2.291	Galiosin	галиозин	Heterocyclic	гетероциклический	t2.346
t2.292	Gallic acid	галловая кислота	Hexadecanoic acid (palmitic acid)	пальмитиновая кислота (гексадекановая кислота)	t2.347
t2.293	Gallocatechin	галокатехин	Hexynyl disulfide	гексенилдисульфид	t2.348
t2.294	Gallotannin	галлотанид	Hippeastrine	гиппеастрин	t2.349
t2.295	Genistein	генистеин	Histamine	гистамин	t2.350
t2.296	Genistin	генистин	Homothermopsine	гомотермопсин	t2.351
t2.297	Gentianadine	генцианадин	Hordenine	горденин	t2.352
t2.298	Gentiananine	генциананин	Humulene	хумулен	t2.353
t2.299	Gentianine	генцианин	Hydrocarbons	углеводороды	t2.354
t2.300	Gentioflavine	генциофлавин	Hydrocyanic acid	цианистоводородная кислота	t2.355
t2.301	Gentiotibetine	генциотибетин	Hydroxycinnamic acid	гидроксикоричная кислота	t2.356
t2.302	Gentisinic acid	гептизиновая кислота	Hydroxytryptamine	гидрокситриптамин	t2.357
t2.303	Gentianaine	генцианаин	Hygrine	гигрин	t2.358
t2.304	Geraniol	гераниол	Hyoscyamine	гиосциамин	t2.359
t2.305	Germacrene	гермакрин	Hypericin	гиперицин	t2.360
t2.306	Gitogenin	гитогенина	Hyperoside	гиперозид	t2.361
t2.307	Glabric acid	глабровая кислота	Hypsogenin	гипсогенин	t2.362
t2.308	Glaucine	глауцин	Hyssopin	гиссопин	t2.363
t2.309	Glaunidine	глаунидин	Imperatorin	императорин	t2.364
t2.310	Glaunine	глаунин	Incanine	инканин	t2.365
t2.311	Glauvine	глауфин	Inorganic	неорганический	t2.366
t2.312	Glucobarbarin	глюкобарбарин			t2.367

t2.368	Inosine	инозин	Leontamine	леонтамин	t2.424
t2.369	Interoside	интерозид	Leontidine	леонтидин	t2.425
t2.370	Intibin	интибин	Leontine	леонтин	t2.426
t2.371	Inulin	инулин	Lepidoside	лепитоцид	t2.427
t2.372	Iodine	йод	Leucoanthocyanide	лейкоантоцианид	t2.428
t2.373	Iridoid glucoside	иридоидный гликозид	Leucoanthocyanidin	лейкоантоцианидин	t2.429
t2.374	Iridoids	иридоиды	Leucodelphinidin	лукоделфинидин	t2.430
t2.375	Iron	железо	Leucomisine	леукомизин	t2.431
t2.376	Isoalantolactone	изоалантолактон	Levorotatory	левовращающийся	t2.432
t2.377	Isobaldine	изоболдин	Licoctonine	ликкоктонин	t2.433
t2.378	Isobetanine	изобетанин	Licorine	ликорин	t2.434
t2.379	Isocorydine	изокоридин	Lignan	лигнан	t2.435
t2.380	Isoflavan	изофлаван	Lignin	лигнин	t2.436
t2.381	Isoleontine	изолеонтин	Lignoceric acid	лигиоцериновая кислота	t2.437
t2.382	Isoliquiritigenin	изоликвиритигенин	Limonene	лимонен	t2.438
t2.383	Isomenthone	изоментон	Limonoid	лимоноид	t2.439
t2.384	Isopsoralen	изопсорален	Linalool	линалоол	t2.440
t2.385	Isoquercitrin	изокверцитрин	Linalyl acetate	линалилацетат	t2.441
t2.386	Isoremerin	изорёмерин	Lindleyin	линдлеин	t2.442
t2.387	Isoreserpiline	изорезерпилин	Linoleic acid	линолевая кислота	t2.443
t2.388	Isorhamnetin	изорамнетин	Linolenic acid	линоленовая кислота	t2.444
t2.389	Isorubijervine	изорубийервин	Lipid	липид	t2.445
t2.390	Isosalipurposide	изосалипурпозид	Liquirazide	ликвиритозид	t2.446
t2.391	Isotalatizine	изоталатизин	Liquiritin	ликвиритин	t2.447
t2.392	Isotanshinone	изотаншинон	Liquitigenin	ликвитигенин	t2.448
t2.393	Isotetrandrine	изотетрандрин	Liriodenine	лириоденин	t2.449
t2.394	Isotrifolin	изотрифолин	Lithospermic acid	литоспермовая кислота	t2.450
t2.395	Isovalerianic acid, isovaleric acid	изовалериановая кислота	Longifolin	лонгифолен	t2.451
t2.396	Isovalerianic acid	изовалериановая кислота	Loroglossine	лороглоссин	t2.452
t2.397	Isovalerianic acid, isovalerianic acid	изовалериановая кислота	Lotaustralin	лотаустралин	t2.453
t2.398	Isovaltrate	изовалтрат	Lucidin	луцидин	t2.454
t2.400	Jalapine	ялапин	Lupane	лупан	t2.455
t2.401	Jatrorrhizine	ятроррицин	Lupanine	лупанин	t2.456
t2.402	Jervine	йервин	Lupeol	лупеол	t2.457
t2.403	Juglone	юглон	Lutein	лютеин	t2.458
t2.404	Juniperin	юниперин	Luteolin	лютеолин	t2.459
t2.405	Kaempferol	кампферол	Luteolin 7-glucoside	лютиолин-7-глюкозид	t2.460
t2.406	Karakoline	караколин	Luteolin 7-rutinoside	лютиолин-7-рутинозид	t2.461
t2.407	Karasamine	карасамин	Lycopene	ликопин	t2.462
t2.408	Ketone	кетон	Macroelement	макроэлемент	t2.463
t2.409	Ketose	кетосахар	Magnesium	магний	t2.464
t2.410	Koproporphyrin	копропорфирин	Magnofflorine	магнофлорин	t2.465
t2.411	Korseveramine	корсеверамин	Maleic acid	малеиновая кислота	t2.466
t2.412	Korseveridine	корсеверидин	Malic acid	яблочная кислота	t2.467
t2.413	Korseverinine	корсеверинин	Manganese	марганец	t2.468
t2.414	Kusunokinin	кусунокинин	Manool	маноол	t2.469
t2.415	Lactic acid	молочная кислота	Marubiin	марубиин	t2.470
t2.416	Lactone	лактон	Matricarin	матрикарин	t2.471
t2.417	Lactose	лактоза	Matrine	матрин	t2.472
t2.418	Lactucin	лактуцин	Mecambroline	мекамбролин	t2.473
t2.419	Lactucopicrin	лактупекрин	Melilotic acid	мелилотиновая кислота	t2.474
t2.420	Lagochilin	лагохиллин	Melilotin	мелилотин	t2.475
t2.421	Lappaconidine	лаппаконидин	Melilotocide	мелилотозид	t2.476
t2.422	Lappaconitine	лаппаконитин	Melissic acid	мелиссовая кислота	t2.477
t2.423	Lead	свинец	Menthol	ментол	t2.478

t2.479	Menthone	ментон	Ocimene	оцимен	t2.534
t2.480	Mesaconitine	мезаконитин	Octylene	октилен	t2.535
t2.481	Methoxy-cinnarolic aldehyde	метоксикоричный альдегид	Oil	масло	t2.536
t2.482	Methyl gallate	метилгаллат	Oleanane	<i>олеанан</i>	t2.537
t2.483	Methyl lachnophyllate	метил лакнофиллат	Oleanolic acid	олеаноловая кислота	t2.538
t2.484	Methyl-chavicol	метилхавикол	Oleic acid	олеиновая кислота	t2.539
t2.485	Methyl-coniine	метилconiин	Oleoresin	олеорезин	t2.540
t2.486	Methylcytidine	метилцитизин	Oligosaccharide	олигосахарид	t2.541
t2.487	Methylcytisine	метилцитизин	Oliveramine	оливерамин	t2.542
t2.488	Methyl-evoxin	метилэвоксин	Oliveridine	оливеридин	t2.543
t2.489	Methyllycaconitine	метилликаконитин	Oliverine	оливерин	t2.544
t2.490	Methylquercetin	метилкверцетин	Olmelin	олмелин	t2.545
t2.491	Microelements	микроэлементы	Omega-3 fatty acid	омега-3 жирная кислота	t2.546
t2.492	Mineral salts	минеральные соли	Onopordopicrin	онопордопикрин	t2.547
t2.493	Mollugin	моллугин	Organic	органический	t2.548
t2.494	Molybdenum	молибден	Organic acid	органическая кислота	t2.549
t2.495	Monoacylglyceride	моноацилглицерид	Osthol	остхол	t2.550
t2.496	Monocaffeoyltartaric acid	<i>монокофейл</i> -винная кислота	Oxalic acid	щавелевая кислота	t2.551
t2.497	Monohydroxyacid	монооксикислота	Oxyacanthine	оксиакантин	t2.552
t2.498	Monosaccharide	моносахарид	Oxymatrin	оксиматрин	t2.553
t2.499	Monoterpene	монотерпен	Oxypeucedanin	оксипейцеданин	t2.554
t2.500	Monticamine	монтикамин	Oxysophocarpine	окись софокарпина	t2.555
t2.501	Morphine	морфин	Oxysteroid	оксистероид	t2.556
t2.502	Mucilage	слизи	Oxytanshinone	окситаншинон	t2.557
t2.503	Mussaenoside	муссаенозид	Pachycarpine	пахикарпин	t2.558
t2.504	Mustard essential oil	горчичное эфирное масло	Palmatine	пальматин	t2.559
t2.505	Myrcene	мирцен	Palmitic acid (hexadecanoic acid)	пальмитиновая кислота	t2.560
t2.506	Myricitrin	мирицитрин	Pancratine	панкратин	t2.561
t2.507	Myricyl alcohol	мирициловый спирт	Pantotenic acid	пантотеновая кислота	t2.562
t2.508	Myristic acid	миристиновая кислота	Paraffin	парафин	t2.563
t2.509	Myristicin	миристицин	Paraoxycoumarin	параоксикумарин	t2.564
t2.510	Myrtenol	миртенол	Parfumine	парфумин	t2.565
t2.511	Myrtenyl isovalerianate	миртенил изовалерианат	Parinaric acid	паринариновая кислота	t2.566
t2.512	Napellin	напеллин	Parishin	паришин	t2.567
t2.513	Naphthoquinone	нафтохинон	Patchouli alcohol	пачулиевый <i>спирт</i>	t2.568
t2.514	Naringenin chalcone	нарингенин халькон	Patrinocide	патринозид	t2.569
t2.515	Narwedine	нарведин	Pectic substances	пектиновые вещества	t2.570
t2.516	n-dimethyl colletine	п-диметилколлетин	Pectins	пектины	t2.571
t2.517	Neoline	неолин	Pegamine	пегамин	t2.572
t2.518	Neosophoramine	неософорамин	Peganidine	пеганидин	t2.573
t2.519	Neoxanthine	неоксантин	Peganine	пеганин	t2.574
t2.520	Nepetalactone	непеталактон	Peganol	пеганол	t2.575
t2.521	Nepodin	неподин	Pelargonin	пелларгонин	t2.576
t2.522	Nerolidol	неролидол	Pentosan	пентозаны	t2.577
t2.523	n-heptacosane	п-гептакозан	Peonidin	пеонидин	t2.578
t2.524	Nickel	никель	Perfamine	перфамин	t2.579
t2.525	Nicotine	никотин	Perforine	перфорин	t2.580
t2.526	Nicotinic acid	никотиновая кислота	Phenol	фенол	t2.581
t2.527	Nitrogen	азот	Phenolcarbonic acid	фенолкарбоновая кислота	t2.582
t2.528	Nitrogenous compounds	азотсодержащие соединения	Phenolic acid	феноловая кислота	t2.583
t2.529	Noradrenaline	норадреналин	Phenolic glucoside	феноловый гликозид	t2.584
t2.530	Norcorydine	норкоридин	Phenyl glycoside	фенил гликозид	t2.585
t2.531	Nortropine	нортропин	Phenylbutanoid	фенилбутаноид	t2.586
t2.532	n-oxy-benzoic acid	п-оксибензойная кислота	Phenylpropanoid	фенилпропаноид	t2.587
t2.533	Oblongine	облонгин			t2.588

t2.589	Phenyl- β -naphthylamine	фенил- β -нафтиламин	Quercetin arabinoside	арабинозид кверцетина	t2.644
t2.590	Phloridzin	флоридзин	Quercetin galactoside	галактозид кверцетина	t2.645
t2.591	Phloroglucinol	флороглуцин	Quercetin triglycoside	тригликозид кверцетина	t2.646
t2.592	Phospholipid	фосфолипид	Quercetin-3-arabinoside	3-арабинозид кверцетина	t2.647
t2.593	Phosphoric acid	фосфорная кислота	Quercetin-3-galactoside	кверцетин-3-галактозид	t2.648
t2.594	Phyllalbine	филлальбин	Quercetrin	кверцитрин	t2.649
t2.595	Phytoecdysone	фитоэктизон	Quinidine	хинидин	t2.650
t2.596	Phytoecdysteroid	фитоэктистероид	Quinone	хинон	t2.651
t2.597	Phytoestrogen	фитоэстроген	Quinovic acid	хинновая кислота	t2.652
t2.598	Phytol	фитол	Ranunculin	ранункулин	t2.653
t2.599	Phytoncid	фитонцид	Remrefidine	ремрефидин	t2.654
t2.600	Phytosterin	фитостерин	Remrefine	ремрефин	t2.655
t2.601	Pigments, dyeing substances	красящие вещества	Reserpine	резерпин	t2.656
t2.602	Pilocarpine	пилокарпин	Reserpinine	резерпинин	t2.657
t2.603	Pinene	пинен	Reticuline	ретикулин	t2.658
t2.604	Pinocamphone	пинокамфон	Rhamnoglucoside	рамноглюкозид	t2.659
t2.605	Piperitone	пиперитон	Rhamnoglycoside	рамногликозид	t2.660
t2.606	Plantagonine	плантагонин	Rhododendrol	рододендрол	t2.661
t2.607	Podophyllotoxin	подофилотоксин	Roemeridine	ремеридин	t2.662
t2.608	Poisonous	ядовитый	Roemerine	рёмерин	t2.663
t2.609	Polyenes	полиены	Rosmarinic acid	розмариновая кислота	t2.664
t2.610	Polyphenol	полифенол	Royleanone	ройлеанон	t2.665
t2.611	Polysaccharide	полисахарид	Ruberythric acid	руберитриновая кислота	t2.666
t2.612	Potassium	калий	Rubiadin	рубиадин	t2.667
t2.613	Prangenidin	прангенидин	Rubijervine	рубийервин	t2.668
t2.614	Prangenin	прангенин	Rubi dine	рубиодин	t2.669
t2.615	Prangosine	прангосин	Rus ogenin	рускогенин	t2.670
t2.616	Primveraza	примвераз	Rutin	рутин	t2.671
t2.617	Proanthocyanidin	проантоцианидин	Rutinoside	рутиназид	t2.672
t2.618	Proazulen	проазулен	Sabinene	сабинена	t2.673
t2.619	Propenyl isothiocyanate	пропенил изотиоцианат	Salicylic acid	салициловая кислота	t2.674
t2.620	Propionic acid	пропионовая кислота	Salidoside	салидросид	t2.675
t2.621	Protein	белок	Salvicin	сальвицин	t2.676
t2.622	Protocatechin	протокатехин	Salvicinin	сальвицинин	t2.677
t2.623	Protopine	протопин	Salvicinolide	сальвицинолид	t2.678
t2.624	Protoporphyrin	протопорфирин	Salvicinolin	сальвицинолин	t2.679
t2.625	Protopseudohypericin	протопсевдогиперицин	Salvifolin	сальвифолин	t2.680
t2.626	Prulaurasin	прулауразин	Salvin	сальвин	t2.681
t2.627	Prunasin	пруназин	Sambulene	самбулен	t2.682
t2.628	Prussic acid	синильная кислота	Sanguinarine	сангвинарин	t2.683
t2.629	Pseudoconhydrine	псевдоконгидрином	Santolina alcohol	сантолиновый спирт	t2.684
t2.630	Pseudoephedrine	псевдоэфедрин	Sapogenin	сапогенин	t2.685
t2.631	Pseudohypericin	псевдогиперицин	Saponarin	сапонарин	t2.686
t2.632	Pseudojervine	псевдойервин	Saponin	сапонин	t2.687
t2.633	Pseudotaraxasterol	псевдотаракастерол	Saporubin	сапорубин	t2.688
t2.634	Pseudotropine	псевдотропин	Saporubinic acid	сапорубиновая кислота	t2.689
t2.635	Psoralen	псорален	Sclareol	склареол	t2.690
t2.636	Pulegone	пулегон	Scopolamine	скополамин	t2.691
t2.637	Purine derivatives	производные пурина	Scopoletin	скополетин	t2.692
t2.638	Purpurin	пурпурин	Scutellarin	скутеллярин	t2.693
t2.639	Pyrocatechin	пирокатехин	Selenium	селен	t2.694
t2.640	Pyrogallol	пирогаллол	Semi-drying oil	полувьсышающее масло	t2.695
t2.641	Pyrrolidine	пирролидин	Sesquiterpene	сесквитерпен	t2.696
t2.642	Pyrrolidine alkaloid	пирролидиновый алкалоид	Sesquiterpene alcohol	сесквитерпеновый спирт	t2.697
t2.643	Quercetin	кверцетин	Sesquiterpene lactone	сесквитерпеновый лактон	t2.698

t2.699	Shepherin	шеперин	Terpinen-4-ol	терпинен-4-ол	t2.754
t2.700	Silicic acid	кремневая кислота	Trpinene	терпенен	t2.755
t2.701	Silicon	кремний	Terpineol	терпенеол	t2.756
t2.702	Silicristin	силикрестин	Terpinolene	терпинолен	t2.757
t2.703	Silimarin	силимарин	Thalfine	тальфин	t2.758
t2.704	Silybin	силибин	Thalfinite	тальфинин	t2.759
t2.705	Sinapic acid	синапиновая кислота	Thalicmidine	таликмидин	t2.760
t2.706	Sinigrin	синигрин	Thalicmine	таликмин	t2.761
t2.707	Skimmianine	скиммианин	Thalicminine	таликминин	t2.762
t2.708	Smirnovine	смирновин	Thalicmitrine	таликмитрин	t2.763
t2.709	Sogdisterone	согдистерон	Thalisopidine	тализопидин	t2.764
t2.710	Songorine	сонгорин	Thalisopine	тализопин	t2.765
t2.711	Sophocarpine	софокарпин	Thalmine	тальмин	t2.766
t2.712	Sophoramine	софорамин	Thalminine	тальминин	t2.767
t2.713	Sophoridine	софоридин	Thermopsine	термопсин	t2.768
t2.714	Sparteine	спартеин	Thermopsocide	термопсозид	t2.769
t2.715	Sparteine	спартеин	Thioglycoside	тиогликозид	t2.770
t2.716	Spathulenol	спатуленол	Thiramine	тирамин	t2.771
t2.717	Spherophysine	сферофизин	Thujone	туйон	t2.772
t2.718	Spherosine	сферозин	Thymol	тимол	t2.773
t2.719	Spinasterol	спинастерол	Thymoquinone	timoхинон	t2.774
t2.720	Stachydrine	стахидрин	Titrateable organic acids	титруемые органические кислоты	t2.775
t2.721	Stachyose	стахиоза	Tocopherol	токоферол	t2.776
t2.722	Starch	крахмала	Tolmetin	таьметин	t2.777
t2.723	Stearic acid	стеариновая кислота	Torachrysone	торахризон	t2.778
t2.724	Stearidonic acid	стеаридоновая кислота	Toxic	токсичный	t2.779
t2.725	Sterin	стерин	Trace	следы	t2.780
t2.726	Steroid	стероид	<i>trans</i> -carveol	транс-карвеол	t2.781
t2.727	Steroidal saponin	стероидный сапонин	Triacanthine	триакантин	t2.782
t2.728	Sterol	стерол	Triacylglyceride	триацилглицерид	t2.783
t2.729	Stigmasterin	стигмастерин	Triacylglycerol	триацилглицерол	t2.784
t2.730	Stigmasterol	стигмастерол	Trichodesmine	триходесмин	t2.785
t2.731	Stilbene derivatives	производные стибена	Tricyclic	трициклический	t2.786
t2.732	Strychnine	стрихнин	Trifolin	трифолин	t2.787
t2.733	Substance	вещество	Trifoside	трифозид	t2.788
t2.734	Succinic acid	янтарная кислота	Triglyceride	триглицерид	t2.789
t2.735	Sucrose	сахароза	Triglycoside isoramnetin	тригликозид изорамнетина	t2.790
t2.736	Sugar	сахар	Trihydroxychalcone	тригидрооксихалкон	t2.791
t2.737	Sulfur	сера	Trimethoxyl-cinnamic acid	триметоксил коричная кислота	t2.792
t2.738	Talatzamine	талатизамин	Triterpene	тритерпен	t2.793
t2.739	Talatzidine	талатизидин	Triterpene alcohol	тритерпеновый спирт	t2.794
t2.740	Talatzine	талатизин	Triterpene glycoside	тритерпеновый гликозид	t2.795
t2.741	Talicymidine	таликмидин	Triterpene saponin	тритерпеновый сапонин	t2.796
t2.742	Tannins	дубильные вещества	Triterpenoid saponin	тритерпеноидный сапонин	t2.797
t2.743	Tanshinone	таншинон	Triterpenol	тритерпенол	t2.798
t2.744	Taraxanthin	тараксантин	Tropane alkaloid	тропановый алкалоид	t2.799
t2.745	Taraxasterol	тараксастерол	Tropine	тропин	t2.800
t2.746	Taraxerol	тараксерол	Tropinone	тропинон	t2.801
t2.747	Taraxol	тараксол	Tropolone	трополон	t2.802
t2.748	Tartaric acid	винная кислота	Tungsten	вольфрам	t2.803
t2.749	Taspine	таспин	Turkesterone	туркестерон	t2.804
t2.750	Tatsetine	тацеттин	Tussilagin	туссиягин	t2.805
t2.751	Taxifolin	таксифолин	Tyramine	тирамин	t2.806
t2.752	Taxodione	таксодион	Tyrosine	тирозин	t2.807
t2.753	Terpene	терпен			t2.808

t2.809	Tyrosol	тиросол	Viticosterone	витикостерон	t2.836
t2.810	Umbelliferol	умбеллиферол	Volatile	летучий	t2.837
t2.811	Umbelliferone	умбеллиферон	Vulgarol	вулгарол	t2.838
t2.812	Undecanoic acid	ундекановая кислота	Waxes	воск	t2.839
t2.813	Unsaturated fatty acid	ненасыщенная жирная кислота	Xanthamine	ксантумин	t2.840
t2.814	Uronic acid	уроновая кислота	Xanthanine	ксантоксин	t2.841
t2.815	Ursane	<i>урсан</i>	Xanthanol	ксантанол	t2.842
t2.816	Ursolic acid	уроловая кислота	Xanthinine	ксантинин	t2.843
t2.817	Urticin	уртицин	Xanthinosin	ксантинозин	t2.844
t2.818	Vaillantine	вайлантин	Xanthone	ксантон	t2.845
t2.819	Valerianic acid	валериановая кислота	Xanthophyll	ксантофилл	t2.846
t2.820	Valeric acid	валериановая кислота	Xanthosine	ксантоксин	t2.847
t2.821	Valeride	валерид	Xanthostrumarin	ксантострумарин	t2.848
t2.822	Valerine	валерин	Xanthumanol	ксантуманол	t2.849
t2.823	Valtrate	валтрат	Xyloglucoside	ксилоглюкозид	t2.850
t2.824	Vanillic acid	ванилиновая кислота	Zinc	цинк	t2.851
t2.825	Vasicinone	вазицинон	Zygodabagine	зигофабагин	t2.852
t2.826	Verbascoside	вербаскозид	α -humulene	α -хумулен	t2.853
t2.827	Verbenone	вербенон	α -hydrojuglone	α -гидроюглон	t2.854
t2.828	Vincanidine	винканидин	α -linolenic acid	α -линолиновая кислота	t2.855
t2.829	Vincanine	винканин	α -terpenyl-acetate	α -терпенилацетат	t2.856
t2.830	Vinervine	винервин	β -bisabolene	β -бизаболен	t2.857
t2.831	Vinervinine	винервинин	β -carboline	β -карболин	t2.858
t2.832	Vin amine	винкамин	β -hydrojuglone	β -гидроюглон	t2.859
t2.833	Vitamin	витамин	β -sitosterin	β -ситостерин	t2.860
t2.834	Vitexin	витексин	γ -coniceine	γ -коницеином	t2.861
t2.835	Vitexin-ramnoside	витексинрамнозид	γ -terpinene	γ -терпинен	t2.862

Uncorrected Proof

39 English-Russian Translations of Medical Terms

t3.1	Abdomen	живот
t3.2	Abdominal cavity	брюшная полость
t3.3	Abortifacient	абортирующее (средство)
t3.4	Abrasion	ссадина
t3.5	Abscess	нарыв, гнойник
t3.6	Acetylcholine	ацетилхолин
t3.7	Acetylcholinesterase	ацетилхолинэстераза
t3.8	Ache	боль, угръ
t3.9	Acute tests on animals	острые опыты на животных
t3.10	Adaptagen	адаптоген
t3.11	Adrenal gland	надпочечник
t3.12	Adrenaline	адреналин
t3.13	Adrenergic	адренергический
t3.14	Aggregative properties	агрегационные свойства
t3.15	Aimaline	аймалин
t3.16	Air sickness	воздушная болезнь
t3.17	Aldose reductase	альдоз редуктаза
t3.18	Alimentary	алиментарный
t3.19	Allergen	аллерген
t3.20	Allergy	аллергия
t3.21	Altitude sickness	горная болезнь
t3.22	Amenorrhea	амеоррея
t3.23	Anabolic activity	анаболическая активность
t3.24	Anacidic gastritis	анацидный гастрит
t3.25	Anacidity	анацидный
t3.26	Analeptic	аналептик
t3.27	Analgesic	болеутоляющее, анальгезирующее (средство)
t3.28		
t3.29	Anaphylactic shock	анафилактический шок
t3.30	Anasarca	анасарка
t3.31	Androgenic action	андрогенное действие
t3.32	Anemia	малокровие, анемия
t3.33	Anemonin	анемонин
t3.34	Anesthesia	анестезия
t3.35	Anesthetic	анестезирующий
t3.36	Angiocholitis	ангиохолит
t3.37	Angioprotector	ангиопротектор

Anorexia	анорексия
Antagonist	антагонист
Anthrax	сибирская язва
Anti-aggregant	антиагрегант
Anti-amnesic	антиамнезийный
Antiarrhythmic	антиаритмический
Anti-asthmatic	противоастмати-ческий
Antibacterial	антибактериальный, противобактериаль-ный
Antibiotic	антибиотик
Anticarcinogenic	антикарциногенный
Anticonvulsive	противосудорожный
Anti-cough	противокашлевой
Antidiarrheal	противоноосный
Antiedemic	противоотёчный
Antiemetic	противорвотный
Anti-fibrillant	антифибрилят
Antifungal	антимикотический, противогрибковый
Antigonadotropic	антигонадотропный
Antihelminthic	антигельминтный
Antihistamine	антигистамин
Anti-HIV	анти-ВИЧ
Antihypertensive	противогипертонический
Antihypoxic	антигипоксический
Antihysterical	противоистерич-ческий
Anti-inflammatory	противовоспалитель-ный
Antimicrobial	антимикробный
Antioxidant	антиоксидант
Antiparasitic	противопаразитар-ный
Antiproliferative	антипролифератив-ный
Antiprotist	протистоцидный
Antipyretic	жаропонижающий
Antisclerotic	антисклеротический
Antiseptic	противосептический
Antispasmodic	противоспазматический
Antithyroid	антитиреоидный
Antitumor	противоопухольевый
Antiulcerogenic	противоязвенный
Antivenom	противоядие
Aorta	аорта

т3.38	Apoptosis	апоптоз	Carcinogenesis	канцерогенез	т3.94
т3.39	Apoptotic effect	апоптический эффект	Cardiosclerosis	кардиосклероз	т3.95
т3.40	Appetite	аппетит	Cardiotonic	кардиотонический	т3.96
т3.41	Arachno-encephalitis	арахноэнцефалит	Caries	кариес	т3.97
т3.42	Arachnoiditis	арахноидит	Carminative	ветрогонное (средство)	т3.98
т3.43	Arrhythmia	аритмия	Carotid nerve	каротидный нерв	т3.99
т3.44	Arterial pressure	артериальное давление	Catalepsy	каталепсия	т3.100
т3.45	Arteriosclerosis	атеросклероз	Cataract	катаракта	т3.101
т3.46	Arthralgia	артралгия	Cell	клетка	т3.102
т3.47	Arthritis	артрит	Central nervous system	центральная нервная система	т3.103
т3.48	Arthrosis	артроз	Cerebral cortex	головной мозг	т3.104
т3.49	Ascariasis	аскаридоз	Cervical erosion	эрозия шейки матки	т3.105
т3.50	Asphyxia	асфиксия	Cervix	шейка матки	т3.106
т3.51	Asthenia	астения	Chest	грудь	т3.107
т3.52	Asthma	астма	Cholagogue, choloretic	желчегонное (средство)	т3.108
т3.53	Astringent	вяжущее (средство)	Cholecystitis	холецистит	т3.109
т3.54	Atonia	атония	Cholera	холера	т3.110
т3.55	Autoimmune	аутоиммунный	Choleresis	желчеотделение	т3.111
т3.56	Autonomic ganglia	автономные ганглии	Choleric action	желчегонное действие	т3.112
т3.57	Avitaminosis	авитаминоз	Choleric, cholagogue	желчегонное (средство)	т3.113
т3.58	Back pain	боли в пояснице	Cholesterol	холестерин	т3.114
т3.59	Bacteria	бактерии	Cholinergic	холинергический	т3.115
т3.60	Bactericidal	бактерицидный	Cholinesterase	холинэстеразный	т3.116
т3.61	Bacteriostatic	бактериостати-ческий	Chorea	хорея	т3.117
т3.62	Bed bug	клоп	Chronic	хронический	т3.118
т3.63	Bedsores	пролежни	Chronotropic	хронотропный	т3.119
т3.64	Bile	желчь	Coagulation	свёртывание	т3.120
т3.65	Bile duct	желчный путь	Coating (remedy)	обволакивающее (средство)	т3.121
т3.66	Bile-stimulant	желчестимулирующий	Colitis	колит	т3.122
т3.67	Bilirubin	билирубин	Common cold	простуда	т3.123
т3.68	Bleeding, hemorrhaging	кровотечение	Compress	компресс	т3.124
т3.69	Bloated	вздутие	Congenital defect	врождённый дефект	т3.125
т3.70	Blood	кровь	Congestion	закупорка	т3.126
т3.71	Blood circulation	кровообращение	Conjunctivitis	конъюнктивит	т3.127
т3.72	Blood cleanser	кровоочистительный	Constipation	запор	т3.128
т3.73	Blood coagulation	свёртывание крови	Contraceptive	контрацептивный, противозачаточный	т3.129
т3.74	Blood sugar	сахар крови	Contractility	сократимость	т3.130
т3.75	Bone	кость	Contraction	сокращение	т3.131
т3.76	Bone fracture	перелом кости	Convulsions	судорги	т3.132
т3.77	Bone marrow	костный мозг	Coronary	коронарный	т3.133
т3.78	Bradycardia	брадикардия	Cough	кашель	т3.134
т3.79	Bronchiectasis	bronхоэктаз	Croupous pneumonia	крупозная пневмония	т3.135
т3.80	Bronchitis	бронхит	Curare	кураре	т3.136
т3.81	Bronchospasm	бронхоспазм	Cuts	порезы	т3.137
т3.82	Bronchus	бронх	Cyclooxygenase (OX) enzyme	циклооксигеназный фермент (ЦОГ)	т3.138
т3.83	Brucellosis	бруцеллёз	Cystitis	цистит	т3.139
т3.84	Bruise	ушиб	Cytotoxicity	цитотоксичность	т3.140
т3.85	Burn	ожог	Deafness	глухота	т3.141
т3.86	Butyrylcholinesterase	бутирилхолин-эстераза	Decoction	отвар	т3.142
т3.87	Ca ⁺⁺ channel	кальциевые каналы	Decompensation	декомпенсация	т3.143
т3.88	Ballus	мозоль	Demulcent	мягчительное (средство)	т3.144
т3.89	Calm	успокоение	Depression (emotional)	депрессия (эмоциональная)	т3.145
т3.90	Cancer	рак (болезнь)	Depression (physical, physiological)	угнетение	т3.146
т3.91	Capillary	капилляр			т3.147
т3.92	Capillary strengthening	капилляроукрепляющее			т3.148
т3.93	Carbuncle	карбункул			т3.149

t3.150	Dermatitis	дерматит	Faint	обморок	t3.206
t3.151	Dermatosis	дерматоз	Fatigue	упадок сил	t3.207
t3.152	Desensitization	десенсибилизация	Febrifuge	жаропонижающее (средство)	t3.208
t3.153	Detoxicant	детоксикант	Festering wounds	гнойные раны	t3.209
t3.154	Detoxify	детоксицировать	Fetal hypoxic hypotropia	гипоксическая гипотрофия плода	t3.210
t3.155	Diabetes	диабет	Fever	лихорадка	t3.211
t3.156	Diaphoretic, sudorific	потогонное (средство)	Fibrinolytic	фибринолитический	t3.212
t3.157	Diarrhea	понос, диарея	Fibroblast	фибробласт	t3.213
t3.158	Diathesis	диатез	Fibrosis	фиброз	t3.214
t3.159	Digestion	пищеварение	Flea	блоха	t3.215
t3.160	Digestive organs	пищеварительные органы	Food poisoning	пищевое отравление	t3.216
t3.161	Digestive system	пищеварительная система	Fragmentation	фрагментация	t3.217
t3.162	Digestive tract	пищеварительный тракт	Free radicals	свободные радикалы	t3.218
t3.163	Diphtheria	дифтерия	Frostbite	обморожение	t3.219
t3.164	Disease	заболевание	Fungal skin disease	лишай	t3.220
t3.165	Disinfectant	обеззараживающий	Fungicide	фунгицид	t3.221
t3.166	Disinfection	дезинфекция	Fungus	грибок	t3.222
t3.167	Dissolve	растворение	Furuncle	фурункул	t3.223
t3.168	Diuresis	диурез	Furunculosis	фурункулёз	t3.224
t3.169	Diuretic	мочегонное (средство)	Galenic preparation	галеновый препарат	t3.225
t3.170	Dizziness	головокружение	Gallbladder	желчный пузырь	t3.226
t3.171	DNA	ДНК	Gallstone	желчный камень	t3.227
t3.172	Duodenum	двенадцатиперстная кишка	Ganglion-blocking	ганглиоблокиро-вание	t3.228
t3.173	Dysentery	дизентерия	Gangrene	гангрена	t3.229
t3.174	Dysmenorrhea	дисменорея	Gargle	полоскание горла	t3.230
t3.175	Dyspepsia	диспепсия, плохое пищеварение	Gastralgia	гастралгия	t3.231
t3.176	Dyspnea	удушьё	Gastric disease	желудочное заболевание	t3.232
t3.177	Dysuria	дизурия	Gastric fluid	желудочный сок	t3.233
t3.178	Ear	ухо	Gastric pneumatosis	пневматоз желудка	t3.234
t3.179	Eczema	экзема	Gastritis	гастрит	t3.235
t3.180	Edema	водянка, отёк	Gastroenteritis	гастроэнтерит	t3.236
t3.181	Elephantitis	слоновая болезнь	Gastrointestinal tract	желудочнокишечный тракт	t3.237
t3.182	Emetic	рвотное (средство)	Gene expression	экспрессия ген	t3.238
t3.183	Emollient	мягчительное (средство)	General tonic	общеукрепляющее (средство)	t3.239
t3.184	Enteritis	энтерит	Genotoxic	генотоксический	t3.240
t3.185	Endocytotic activity	эндоцитозная активаность	Gingivitis	гингивит	t3.241
t3.186	Endogenous	эндогенный	Gland	железа	t3.242
t3.187	Endomyocarditis	эндомиокардит	Goiter	зоб	t3.243
t3.188	Enterocolitis	энтероколит	Gonorrhea	гонорея	t3.244
t3.189	Enzyme	фермент	Gout	подагра	t3.245
t3.190	Enzyme-stimulating	ферментостимулирование	Gram-negative bacteria	грамотрицательные бактерии	t3.246
t3.191	Epilepsy	эпилепсия	Gram-positive bacteria	грамположительные бактерии	t3.247
t3.192	Epithelization	эпителизация	Granulation	грануляция	t3.248
t3.193	Ergotism	эрготизм	Guinea pig	морская свинка	t3.249
t3.194	Erysipelas	рожа (болезнь)	Guinea worm	ришта, гвинейский червь	t3.250
t3.195	Erysipelatous inflammation	рожистое воспаление	Gum	десна	t3.251
t3.196	Erythrocyte	эритроцит	Gynecological disorders	женские заболевания	t3.252
t3.197	Erythrodermia	эритродермия	Headache	головная боль	t3.253
t3.198	Esophagus	пищевод	Heart	сердце	t3.254
t3.199	Estrogen	эстроген	Heart failure	порок сердца	t3.255
t3.200	Exophthalmic goiter	базедовая болезнь	Helminthosis	гельментоз	t3.256
t3.201	Expectorant	отхаркивающее (средство)	Hematuria	гематурия	t3.257
t3.202	External bleeding	наружное кровотечение	Hemodynamics	гемодинамика	t3.258
t3.203	Exudative diathesis	экссудативный диатез	Hemoglobin	гемоглобин	t3.259
t3.204	Eye	глаз	Hemolytic index	гемолитический индекс	t3.260
t3.205	Facial paralysis	паралич лицевого нерва	Hemoptysis	кровохаркание	t3.261

t3.262	Hemorrhagic shock	геморрагический шок	Intestinal colic	кишечные колики	t3.318
t3.263	Hemorrhaging, bleeding	кровотечение	Intestinal disorder	растройство кишечника	t3.319
t3.264	Hemorrhoidal hemorrhage	геморроидальное кровотечение	Intestines	кишки, кишечник,	t3.320
t3.265	Hemorrhoids	геморрой	Intoxication	интоксикация	t3.321
t3.266	Hemostatic	гемостатический,	Intracellular	внутриклеточный	t3.322
t3.267		кровоостанавливающий	Intracranial	внутричерепной	t3.323
t3.268	Hepatitis	гепатит	Intravenous injection	внутривенное вливание	t3.324
t3.269	Hepatochole-cystitis	гепатохолецистит	Itch	зуд	t3.325
t3.270	Hepatoprotector	гепатопротектор	Jaundice	желтуха	t3.326
t3.271	Hepatotoxic	гепатотоксичный	Joint	сустав	t3.327
t3.272	Hernia	грыжа	Kidney	почка	t3.328
t3.273	Herpes	герпес	Kidney stone	почечный камень	t3.329
t3.274	Herpes simplex type 1	герпес симплекс типа 1	Lactation	лактация	t3.330
t3.275	Highmoritis	гайморит	Lactogenic	лактогенный	t3.331
t3.276	Hippocampus	гиппокамп	Laryngitis	ларингит	t3.332
t3.277	Histamine	гистамин	Larynx	гортань	t3.333
t3.278	Hoarseness	охриплость	Laxative	слабительное (средство)	t3.334
t3.279	Homeopathy	гомеопатия	Leishmaniasis	лейшманиоз	t3.335
t3.280	Hydrophobia	водобоязнь	Lethargic encephalitis	летаргический энцефалит	t3.336
t3.281	Hyperglycemia	гипергликемия	Leucocytes	лейкоциты	t3.337
t3.282	Hyperhydrosis	гипергидроз	Leucomycin	леукомизин	t3.338
t3.283	Hyperpituitarism	гиперпитуитаризм	Leucopenia	лейкопения	t3.339
t3.284	Hypertension	гипертония	Leukemia	лейкемия	t3.340
t3.285	Hypertensive	гипертензивный	Leukorrhea	бели (болезнь)	t3.341
t3.286	Hyperthyroidism	гипертиреоз	Libido	либидо	t3.342
t3.287	Hypertrophy	гипертрофия	Liver	печень	t3.343
t3.288	Hypoacidic gastritis	гипоацидный гастрит	Local anesthesia	местная анестезия	t3.344
t3.289	Hypochondria	ипохондрия	Lotion	примочка	t3.345
t3.290	Hypogastritis	гипогастрит	Low stomach acidity	пониженная кислотность желудка	t3.346
t3.291	Hypoglycemic	гипогликемический	Lungs	лёгкие	t3.347
t3.292	Hypolipidemic	гиполипидемический	Lupus	волчанка (болезнь)	t3.348
t3.293	Hypotension	гипотензия	Lymph nodes	лимфатические узлы	t3.349
t3.294	Hypotensive	гипотензивный	Lymphaden-ectomy	лимфаденэктомия	t3.350
t3.295	Hypothermic	гипотермический	Lymphadenitis	лимфаденит	t3.351
t3.296	Hypotonia	гипотония	Lymphedema	лимфодемия	t3.352
t3.297	Hypoxia	гипоксия	Lymphoblastoid	лимфобластид	t3.353
t3.298	Hysteria	истерия	Macrophage	макрофаг	t3.354
t3.299	Idiopathic skin atrophy	идиопатическая атрофия кожи	Malaria	малярия	t3.355
t3.300	Immunological	иммунологический	Malignant	злокачественный	t3.356
t3.301	Immunosuppression	иммуносупрессия	Mammalian	млекопитающий	t3.357
t3.302	Immunotoxicity	иммунная токсикация	Mannose	манноза	t3.358
t3.303	Implantation	имплантация	Mastitis	мастит	t3.359
t3.304	Impotence	импотенция	Measles	корь	t3.360
t3.305	Infected	инфицированный	Mediator	медиатор	t3.361
t3.306	Infection	инфекция	Medulla oblongata	продолговатый мозг	t3.362
t3.307	Infectious diseases	инфекционные заболевания	Melancholy	меланхолия	t3.363
t3.308	Inflammation	воспалительный процесс	Menopause	климакс	t3.364
t3.309	Influenza	грипп	Menorrhagia	меноррагия	t3.365
t3.310	Infusion	настой	Menstruation	менструация	t3.366
t3.311	Inhibition	ингибирование, задерживание, затормаживание	Menstruation cycle	менструальный цикл	t3.367
t3.312			Metabolism	обмен веществ	t3.368
t3.313	Inotropic action	инотропное действие	Methicillin-resistance	метицилин-резистентный	t3.369
t3.314	Insecticide	инсектицид	Methicillin-sensitive	метицилин-чувствительный	t3.370
t3.315	Insomnia	бессонница	Metropathy	метропатия	t3.371
t3.316	Insulin	инсулин			t3.372
t3.317	Internal	внутренний			48

t3.373	Metrorrhagia	метроррагия	Paronychia	панариций	t3.428
t3.374	Midbrain	средний мозг	Parturifacient	родовспомогатель-ный	t3.429
t3.375	Migraine	мигрень	Paste	паста	t3.430
t3.376	Mitral failure	митральный порок	Pathogenic	потогенность	t3.431
t3.377	Molluscicidal	моллюскицидный	Pediculosis (lice infestation)	педикулёз	t3.432
t3.378	Mouse	мышь	Pellagra	пеллагра	t3.433
t3.379	Mouth wash	полоскание рта	Penicillin	пеницилин	t3.434
t3.380	Mucous membrane	слизистая оболочка	Pepsin	пепсин	t3.435
t3.381	Mucus	слизи (носоглотки)	Pepsinogen	пепсиноген	t3.436
t3.382	Multiple sclerosis	рассеянный склероз	Periodontal disease	пародантоз	t3.437
t3.383	Muscle	мускул	Periodontitis	периодонтит	t3.438
t3.384	Musculoskeletal	скелетно-мускульный	Periostitis	периостит	t3.439
t3.385	Myasthenia	миастения	Peripheral	периферический	t3.440
t3.386	Mydriasis	мидриатический	Peristalsis	перистальтика	t3.441
t3.387	Mydriatic	мидриатик	Peritoneal	перитонеальный	t3.442
t3.388	Myocardial infarction	инфаркт миокарда	Permeability	проницаемость	t3.443
t3.389	Myocarditis	миокардит	Pertussis	коклюш	t3.444
t3.390	Myodystrophy	миодистрофия	Phagocyte	фагоцит	t3.445
t3.391	Myometrium	миометрия	Phagocytic	фагоцитарный	t3.446
t3.392	Myopathy	миопатия	Pharmacological	фармакологический	t3.447
t3.393	Narcosis	наркоз	Pharyngitis	фарингит	t3.448
t3.394	Narcotic	наркотик	Phthisis	чахотка	t3.449
t3.395	Narcotized	наркотизированный	Phytoestrogen	фитоэстроген	t3.450
t3.396	Nausea	тошнота	Pimple	прыщик	t3.451
t3.397	Necrosis	некроз	Pinworm	острица	t3.452
t3.398	Nephritis	нефрит	Plague, pestilence	чума	t3.453
t3.399	Neural	невральный	Plasma	плазма	t3.454
t3.400	Neuralgia	невралгия	Platelet activating factor (PAF)	фактор активации тромбоцитов (ФАТ)	t3.455
t3.401	Neurasthenia	неврастения	Pleurisy	плеврит	t3.456
t3.402	Neuritis	неврит	Pneumonia	пневмония, воспаление лёгких	t3.457
t3.403	Neurodermatitis	нейродермотит	Poison	яд, отравы	t3.458
t3.404	Neuromuscular	нервно-мышечный	Poliomyelitis	полиомиелит	t3.459
t3.405	Neuron	нейрон	Poliovirus	полиовирус	t3.460
t3.406	Neuroprotective	нейрозащитный	Polyarthritis	полиартрит	t3.461
t3.407	Neurosis	невроз	Polyp	полип	t3.462
t3.408	Noradrenaline	норадреналин	Polyvitamin	поливитамин	t3.463
t3.409	Nose	нос	Postencephalitic	постэнцефалитический	t3.464
t3.410	Obesity	ожирение	Poultice	припарка	t3.465
t3.411	Obstetric-gynecological	акушерско-гинекологический	Powder	порошок	t3.466
t3.412	Ointment	мазь	Pressor action	прессорное действие	t3.467
t3.413	Osteitis	остит	Proliferation	пролиферация	t3.468
t3.414	Osteoarthritis	остеоартрит	Prophylactic	профилактика	t3.469
t3.415	Osteomyelitis	остеомиелит	Prostaglandin	простагландин	t3.470
t3.416	Otitis	отит	Prostate	предстательная железа (простата)	t3.471
t3.417	Otolaryngology	отоларингология	Prostitis	простатит	t3.472
t3.418	Pancreas	поджелудочная железа	Prothrombin	протромбин	t3.473
t3.419	Papilloma	папиллома	Psoriasis	псориаз	t3.474
t3.420	Paradontosis	парадонтоз	Psychiatry	психиатрия	t3.475
t3.421	Paralysis	паралич	Psychomimetic	психомиметический	t3.476
t3.422	Paralytic	паралитик	Psycho-neurological diseases	психоневрологические заболевания	t3.477
t3.423	Parasite	паразит	Pulpitis	пульпит	t3.478
t3.424	Parasympathetic ganglions	парасимпатические ганглии	Pupil	зрачок	t3.479
t3.425	Paratyphoid	паратиф	Purgative	рвотный	t3.480
t3.426	Parenteral	парэнтеральный	Pus	гной	t3.481
t3.427	Parkinson's disease	болезнь Паркинсона			t3.482

†3.484	Pyelitis	пиелит	Skin ulcer	язва кожи	†3.540
†3.485	Pyoderma	пиодермия	Sleep	сон	†3.541
†3.486	Quench thirst	жаждоутоляющее	Smallpox	оспа	†3.542
†3.487	Rabies	бешенство	Smooth muscles	гладкая мускулатура	†3.543
†3.488	Radiculitis	радикулит	Snake bites	укус змеи	†3.544
†3.489	Rash	крапивница, сыпь	Snake venom	змеиный яд	†3.545
†3.490	Rat	крыса	Soporific	снотворное средство	†3.546
†3.491	Recalcification	рекальцификация	Sore throat	ангина	†3.547
†3.492	Receptor	рецептор	Spasm	спазм	†3.548
†3.493	Rectal prolapse	выпадение прямой кишки	Spastic	спастический	†3.549
†3.494	Reduced hair growth	плохой рост волос	Spastic paresis	спастический парез	†3.550
†3.495	Reflex excitability	рефлекторная возбудимость	Spazmophilia	спазмофилия	†3.551
†3.496	Relax	расслабляться	Spermatocidic	спермацидный	†3.552
†3.497	Remedy	средство	Spleen	селезёнка	†3.553
†3.498	Renal pelvis	почечная лоханка	Sprain	вывих	†3.554
†3.499	Reoxygenation	реокисление	<i>Staphylococcus</i>	стафилокок	†3.555
†3.500	Residual	остаточный	Stenocardia	стенокардия	†3.556
†3.501	Resistance	резистентность, устойчивость	Stimulate	стимулировать, возбуждать	†3.557
†3.502	Respiration	дыхание	Stomach	желудок	†3.558
†3.503	Respiratory	респираторный	Stomach ache	боли желудка	†3.559
†3.504	Respiratory disease	респираторные заболевания,	Stomach catarrh	катар желудка	†3.560
†3.505		болезнь органов дыхания	Stomatitis	стоматит	†3.561
†3.506	Restorative	общеукрепляющее	<i>Streptococcus</i>	стрептокок	†3.562
†3.507	Reticular	ретикулярный	Streptozotocin-induced	стрептозотоцин-индуцированный	†3.563
†3.508	Rhabdomyolysis	рабдомиолиз	Stress	стресс	†3.564
†3.509	Rheumacarditis	ревмакардит	Stress factor	стрессфактор	†3.565
†3.510	Rheumatic pain	ломота	Sudorific, diaphoretic	потогонное (средство)	†3.566
†3.511	Rheumatism	ревматизм	Sunstroke	солнечный удар	†3.567
†3.512	Rhinitis	рени́т	Surgery	хирургия	†3.568
†3.513	Rickets	рахит	Swelling	опухоль, отёк	†3.569
†3.514	Rubella	краснуха	Sympathetic nervous system	симпатическая нервная система	†3.570
†3.515	Salivation	слюноотечение	Sympatho-mimetic	симпатомиметик	†3.571
†3.516	Scabies	чесотка	Synapse	синапс	†3.572
†3.517	Scarlet fever	скарлатина	Synergistic	синергичный	†3.573
†3.518	Sciatic nerve	седалищный нерв	Syphilis	сифилис	†3.574
†3.519	Scleroderma	склеродермия	Systolic	систолический	†3.575
†3.520	Scratch	царапина	Tachycardia	тахикардия	†3.576
†3.521	Scrofula	скрофулёз, золотуха	Tachyphylaxis	тахифилаксия	†3.577
†3.522	Scurvy	цинга	Tapeworm	солитёр	†3.578
†3.523	Sea sickness	морская болезнь	Tetanus	столбняк (болезнь)	†3.579
†3.524	Secretion	секреция	Throat	горло	†3.580
†3.525	Secretory activity	секреторная деятельность	Thrombocytes	тромбоциты	†3.581
†3.526	Secretory function	выделительная функция	Thrombophlebitis	тромбофлебит	†3.582
†3.527	Sedative	седативный, успокаивающее	Thromboplastic activity	тромбопластическая активность	†3.583
†3.528		(средство)	Thymus	тимус	†3.584
†3.529	Seizure	припадок	Thyroid gland	щитовидная железа	†3.585
†3.530	Sensory	сенсорный	Thyroid stimulating hormone (TSH)	тиреотропный гормон (ТТГ)	†3.586
†3.531	Septicemia	септицемия	Thyroidotoxicosis	тиреотоксикоз	†3.588
†3.532	Serotonin	серотонин	Thyroxine (T ₄)	тироксин (Т ₄)	†3.589
†3.533	Shin	голень	Tick	клещь	†3.590
†3.534	Shortness of breath	одышка	Tincture	тинктура, настойка	†3.591
†3.535	Sialorrhea	слюноотечение	Tissue	ткань (клетки)	†3.592
†3.536	Sinus	синус	Tongue	язык	†3.593
†3.537	Sinus cold	насморк	Tonic	тонизирующее, укрепляющее (средство)	†3.594
†3.538	Skin	кожа			†3.595
†3.539	Skin diseases	заболевания кожи			

t3.596	Tonic action	тонизирующее действие	Uterine fibroids	фибриома матки	t3.623
t3.597	Tonsillitis	тонзиллит	Uterine hemorrhages	маточное кровотечение	t3.624
t3.598	Tonus	тонус	Uterine horn	рог матки	t3.625
t3.599	Tooth	зуб	Uterine ulcers	язва матки	t3.626
t3.600	Toothache	зубная боль	Uterus	матка	t3.627
t3.601	Toxic	токсичный	Vaccine	вакцина	t3.628
t3.602	Tracheitis	трахеит	Vagus nerve	блуждающий нерв	t3.629
t3.603	Tranquilizing effect	транквилизирующий эффект	Vasoconstrictor	сосудосуживающий	t3.630
t3.604	Trichomoniasis	трихомониаз	Vasomotor center	сосудодвигательный центр	t3.631
t3.605	Triiodothyronine (T ₃)	трийодтиронин (Т3)	Vegetative neurosis	вегетативный невроз	t3.632
t3.606	Trophic ulcer	трофическая язва	Vein	вена	t3.633
t3.607	Trypanocidal	трипаноцидный	Venereal diseases	венерические болезни	t3.634
t3.608	Tuberculosis	туберкулёз	Vermifuge	глистогонное (средство)	t3.635
t3.609	Tubular necrosis	некроз трубчатых клеток почек	Vessel	сосуд	t3.636
t3.610	Tumor	опухоль	Veterinary medicine	ветеринария (ветеринарная медицина)	t3.637
t3.611	Tympanites	метеоризм	Virus	вирус	t3.638
t3.612	Typhoid fever	брюшной тиф	Vitiligo	витилиго	t3.640
t3.613	Ulcer	язва	Vomit	рвота	t3.641
t3.614	Ulcer disease	язвенная болезнь	Wart	бородавка	t3.642
t3.615	Upper respiratory	верхние дыхательные пути	Weakened	ослабленный	t3.643
t3.616	Ureter	мочеточник	Weakness	слабость	t3.644
t3.617	Urethra	уретра	Weariness	усталость	t3.645
t3.618	Urinary incontinence	недержание мочи	Weight deficiency	низкий вес	t3.646
t3.619	Urinary tract	мочевыводящий путь	Wound	рана	t3.647
t3.620	Urination disorders	расстройство мочеотделения	Wound healing	ранозаживляющее (средство)	t3.648
t3.621	Urogenital	мочеполовой			
t3.622	Uterine atonia	антония матки			

- 55 eFloras (2008) Published on the Internet. <http://www.efloras.org>.
 56 Accessed 2007–2008. Missouri Botanical Garden/Harvard
 57 University Herbaria, St. Louis/Cambridge
- [AU5] Abaihanova Z, Alpiev M, Bogdetsky V, Bortsova S, Vorobyev G,
 59 Djangaracheva M, Ilyasov M, Ilyasov S, Kabanova N, Kashilov A,
 60 Koshoev M, Lailieva M, Makeyev T, Mateev U, Meng S, Penkina L,
 61 Torgashova L, Filkova T, Hasanov R, Uzakbaeva Z, Shevchenko V,
 62 Shukurov E (2006) Kyrgyzstan: environment and natural resources
 63 for sustainable development. In: Shukurov E, Makeev T, Koshoev M
 64 (eds) CARNet digital informational network on environment and
 65 sustainable development in Central Asia and Russia
- 66 Abdalla S, Al-Khalil S, Afifi F (1991) Observations on the pharmacol-
 67 ogy of thalicminine, an oxyaporphine alkaloid from *Thalictrum*
 68 *isopyroides*. Gen Pharmacol 22(2):253–257
- 69 Abdallah OM (1991) Phenolic glucoside and other constituents of
 70 *Dipsacus laciniatus*. Phytochemistry 30(8):2805–2806
- 71 Abdel Gawad MM, El Amin SM, Ohigashi H, Watanabe Y, Takeda N,
 72 Sugiyama H, Kawanaka M (2000) Molluscicidal saponins from
 73 *Anagallis arvensis* against schistosome intermediate hosts. Jpn J
 74 Infect Dis 53(1):17–19
- 75 Abdulkadirov IT, Yakubova MR, Nuriddinov KhR, Mamatkhanov AU,
 76 Turakhozhaev MT (2005) Ecdysterone and turkesterone in *Ajuga*
 77 *turkestanica* determined by HPLC. Chem Nat Compd
 78 41(4):475–476
- 79 Abdulladzhanova NG, Mavlyanov SM, Dalimov DN (2003) Polyphenols
 80 of certain plants of the Euphorbiaceae family. Chem Nat Compd
 81 39(4):399–400
- 82 Abdullaeva DU, Shakirov R (2006) Alkaloids from *Korolkowia sewert-*
 83 *zovii*. Sevedinine N-oxide. Chem Nat Compd 42(5):623–624
- 84 Abdunazarov SKh (2000) Plants of the Fergana valley – *Pulicaria*
 85 Gaertn. Bioecological peculiarities. Synopsis of dissertation for the
 86 degree of Ph.D. in biology, Kokand (in Uzbek)
- 87 Abdurakhmanov TR (1962) About the pharmacology of some species
 88 of *Lagochilus*. Synopsis of dissertation for the degree of Ph.D. in
 89 medicine, Karaganda (in Russian)
- 90 Abduzhabbarova S, Maekh SKh, Yunusov SYu, Yagudaev MR,
 91 Kurbakov D (1978) Alkaloids of *Thalictrum isopyroides*. Chem Nat
 92 Compd 14(4):400–402
- 93 Abourashed E, El-Alfy A, Khan I, Walker L (2003) Ephedra in perspec-
 94 tive—a current review. Phytother Res 17(7):703–712
- 95 Abuharfeil NM, Maraqa A, Von Kleist S (2000) Augmentation of natu-
 96 ral killer cell activity in vitro against tumor cells by wild plants from
 97 Jordan. J Ethnopharmacol 71(1–2):55–63
- 98 Aida K, Tawata M, Shindo H, Onaya T, Sasaki H, Yamaguchi T, Chin
 99 M, Mitsushashi H (1990) Isoliquiritigenin: a new aldose reductase
 100 inhibitor from glycyrrhizae radix. Planta Med 56(3):254–258
- Aitbaeva RKh (1972) Antibacterial activities of a preparation of Tian
 Shan mountain ash. Zdravookhraneniye Kazakhstana (1):76–77 (in
 Russian) 102
- Aizikov MI, Kurmukov AG, Sirov VN (1978) Physiological activity
 and correlative changes in protein, carbohydrate digestions and lipid
 exchange under influence of ecdizones and nerobols. In: Sadritdinov
 BS (ed) Pharmacology of natural substances. FAN, Tashkent, pp
 107–125 (in Russian) 108
- Aizikov MI, Rasulova SA, Kurmukov AG (1984) Hypocoagulatory
 action of amorphin in experimental atherosclerosis. Dokl Akad
 Nauk UzSSR (10):49–50 111
- Aizikov MI, Syrov VN, Kurmukov AG, Khushbaktova ZA, Rasulova
 SA, Dzhukharova MKh, Malikov VM, Batyrov EKh (1986) Effect
 of polyphenols on lipid metabolism in hyperlipemia and atheroscle-
 rosis. Meditsinskii Zh Uzbekistana (12):45–48 (in Russian) 115
- Aizikov MI, Kurmukov AG, Rasulova SA et al (1991) Angioprotecting
 and hypolipidemic activity of leucomisine on experimental athero-
 sclerosis. Farmakol Toksikol (3):35–37 (in Russian) 118
- Aizikov MI, Kurmukov AG, Rasulova SA (1993a) Influence of leu-
 comisine on the regression of experimental atherosclerosis.
 Meditsinskii Zh Uzbekistana (4):72–75 (in Russian) 121
- Aizikov MI, Prohorova IR, Kurmukov AG (1993b) Angioprotecting,
 hypolipidemic and atherosclerotic activity of austicine. In: Tezisui
 dokladov – 1 konferentsiyakardiologov Moldovui (Proceedings of the
 1st conference of cardiologists of Moldova), Kishinev, pp 150–151 125
- Akbarov ZS, Aliev KhU, Sultanov MB (1972) Antiarrhythmic action of
 thalisopine, novocainamide, and quinidine on experimental arrhyth-
 mia models. Farmakol. In: Sultanov MB (ed) Farmakologiya
 Alkaloidov i Ikh Proizvodnikh (Pharmacology of alkaloids and their
 derivatives). FAN, Tashkent, pp 129–133 (in Russian) 130
- Akbarov ZS, Aliev KhU, Sultanov MB (1978) Antiarrhythmic action of
 alkaloids of bisbenzil, isohinolin and criptopine series. In: Sadritdinov
 BS (ed) Farmakologiya prirodnikh veshchestv (Pharmacology of
 natural substances). FAN, Tashkent, pp 11–29 (in Russian) 134
- Akhmedhodjaeva HS (1978) The influence of the alkaloids of
Haplophyllum acutifolium on sex cycle and reproductive function of
 rats. In: Sadritdinov BS (ed) Farmakologiya prirodnikh veshchestv
 (Pharmacology of natural substances). FAN, Tashkent, pp 51–56 (in
 Russian) 139
- Akhmedhodjaeva HS, Kurmukov AG (1975) Haplamine pharmaco-
 logy. Dokl Akad Nauk UzSSR (8):36 (in Russian) 141
- Akhmedhodjaeva HS, Polievtev NP (1963) Sedative properties of the
 alkaloid performine and its synergy with soporifics and narcotics. In:
 Voprosy biologii i krayevoi meditsinui (Issues of Biology and
 Regional Medicine), no 4. Izdatelstvo Akademii Nauk UzSSR,
 Tashkent (in Russian) 146

- 147 Akhmedzhanova VI, Bessonova IA, Yunusov SYu (1975) Haplamine-a
148 new alkaloid from *Haplophyllum perforatum*. Chem Nat Prod
149 10(1):121–122
- 150 Akhondzadeh S, Noroozian M, Mohammadi M, Ohadinia S, Jamshidi
151 AH, Khani M (2003) *Melissa officinalis* extract in the treatment of
152 patients with mild to moderate Alzheimer's disease: a double blind,
153 randomised, placebo controlled trial. J Neurol Neurosurg Psychiatry
154 74(7):863–866
- 155 Akimaliev AA, Putieva ZhM, Alimbaeva PK, Abubakirov NK (1989)
156 Triterpene glucosides of *Dipsacus azureus*. III. Structure of dipsa-
157 cobioside. Chem Nat Compd 25(3):174–176
- 158 Akopov IE (1981) Hemostatic plants. Meditsina, Tashkent (in Russian)
- 159 Akopov IE (1990) The most important native medicinal plants and their
160 uses. Meditsina, Tashkent (in Russian)
- 161 Akopov IE, Konovalova VA, Mansurov MM (1958). Pharmacology of
162 stachydrine hydrochloride. Farmakol Toksikol 21: 44–46
- 163 Aksinenko SG, Klimentova DA, Pashinsky VG (2003) Cytoprotective
164 effects of alcohol extract from above-ground part of *Fragaria vesca*
165 L. under the conditions of intoxication by cyclophosphane.
166 Rastitel'nye Resursy 39(4):130–134 (in Russian)
- 167 Al-Awadi F, Shoukry M (1988) The lipid lowering effect of an anti-
168 diabetic plant extract. Acta Diabetol 25(1):1–5
- 169 Aldashev AA (1979) *Aconitum*: pharmacology, toxicology and uses.
170 Ilim, Frunze (in Russian)
- 171 Al-douri NA (2000) A survey of medicinal plants and their traditional
172 uses in Iraq. Pharm Biol 38(1):74–79
- 173 Al-Ghamdi MS (2001) The anti-inflammatory, analgesic and antipyretic
174 activity of *Nigella sativa*. J Ethnopharmacol 76(1):45–48
- 175 Ali BH, Blunden G (2003) Pharmacological and toxicological proper-
176 ties of *Nigella sativa*. Phytother Res 17(4):299–305
- 177 Ali MS, Pervez MK, Saleem M, Tareen RB (2001) Haplophytin-A and
178 B: the alkaloidal constituents of *Haplophyllum acutifolium*.
179 Phytochemistry 57(8):1277–1280
- 180 Aliev RK (1945) A wound-healing preparation of the leaves of broad-leaf
181 plantain. Farmatsiya (2):28–29, Meditsina, Moscow (in Russian)
- 182 Aliev KhU, Kamilov IK (1967) Pharmacology of protopine and its
183 derivatives. In: Kamilov IK (ed) Farmakologiya Alkaloidov i
184 Glikozidov (Pharmacology of alkaloids and glycosides). FAN,
185 Tashkent, pp 176–178 (in Russian)
- 186 Aliev RK, Aliev ND, Rakhimova AK (1961) Materials about the study
187 of the rhizomes of *Geum rivale*. Dokl Akad Nauk AzSSR 17(6):519–
188 524 (in Russian)
- 189 Aliev KhU, Zakirov UB, Kamilov IK (1967) Pharmacology of the alka-
190 loid hordenine and its derivatives. In: Kamilov IK (ed) Farmakologiya
191 Alkaloidov i Glikozidov (Pharmacology of alkaloids and glyco-
192 sides). FAN, Tashkent, pp 114–117
- 193 Aliev RK, Prilipko LI, Damirov IA, Islamova NA, Guseinov DYa
194 (1972) In: Guseinov DYa (ed) Lekarstvenniye rasteniya
195 Azerbaidzhana: spravochnik (Medicinal plants of Azerbaijan: man-
196 ual). Azerbaidzhanskoye Knizhnoye Izdatelstvo, Baku
- 197 Alimbaeva PK (1961) About the pharmacology of *Lagochilus* growing
198 in Kyrgyzstan, vol 15. Trudi Kyrgyzskogo Meditsinskogo Instituta,
199 Frunze (in Russian)
- 200 Alimbaeva PK, Akimaliev AA (1975) Phytochemical study of *Scabiosa*
201 *songarica* Schrenk. Farmatsiya 24(3):33–36, Meditsina, Moscow
202 (in Russian)
- 203 Alimbaeva PK, Goncharova AV (1971) Dikorastushchiye lekarstven-
204 niye rasteniya v Kirgizii (farmakolognosticheskiye issledovaniye)
205 [Wild-growing medicinal plants of Kyrgyzstan (pharmacognostic
206 studies)]. Kyrgyzstan, Frunze (in Russian)
- 207 Alimbaeva PK, Shambetov TM (1988) Current status of harvesting
208 medicinal plant raw material in the Kyrgyz SSR. Rastitel'nye
209 Resursy 24(2) (in Russian)
- 210 Alimbaeva PK, Mukhamedziev MM, Akimaliev AA (1986)
211 Lekarstvenniye rasteniya semeistva vorcyankovuiikh flori Kirgizii
(Medicinal plants of the family Dipsacaceae from the flora of
Kyrgyzstan). Ilim, Frunze (in Russian)
- 212 Ali-Shtayeh MS, Abu Ghdeib SI (1999) Antifungal activity of plant
213 extracts against dermatophytes. Mycoses 42(11–12):665–672
- 214 Al-Sultan SI, Hussein YA, Hegazy A (2003) Toxicity of *Anagallis*
215 *arvensis* plant. Pak J Nutr 2(3):116–122
- 216 Altimishev AA (1991) Prirodniye tselebnuiye sredstva, izd. 2 (Natural
217 medicinal remedies), 2nd edn. Kyrgyzstan, Bishkek (in Russian)
- 218 Amirov MN (1974) A study of the connection between anti-tumor
219 activity and the chemical structures of plant tannins.
220 Zdravookhraneniye Kazakhstana (8):36–38 (in Russian)
- 221 Amoros M, Fauconnier B, Girre RL (1987) In vitro antiviral activity of
222 a saponin from *Anagallis arvensis*, Primulaceae, against herpes sim-
223 plex virus and poliovirus. Antiviral Res 8(1):13–25
- 224 Anderson KJ, Teuber SS, Gobeille A, Cremin P, Waterhouse AL,
225 Steinberg FM (2001) Walnut polyphenolics inhibit in vitro human
226 plasma and LDL oxidation. J Nutr 131(11):2837–2842
- 227 Aoyagi Y, Takahashi Y, Satake Y, Takeya K, Aiyama R, Matsuzaki T,
228 Hashimoto S, Kurihara T (2006) Cytotoxicity of abietane diterpe-
229 noids from *Perovskia abrotanoides* and of their semisynthetic ana-
230 logues. Bioorg Med Chem 14(15):5285–5291
- 231 Appolonova IB (1956) Influences of *Petrosimonia*, *Berteroa incana* and
232 *Epilobium hirsutum* on the activity of the heart. Synopsis of disser-
233 tation for the degree of Ph.D. in medicine, Ivanovo (in Russian)
- 234 Arif R, Küpeli E, Ergun F (2004) The biological activity of *Centaurea*
235 L. species. GU J Sci 17(4):149–164
- 236 Arifkhozhaev AO, Rakhimov DA (1986) Polysaccharides of saponin-
237 bearing plants. III. Polysaccharides of the epigeal organs of
238 *Biebersteinia multifida*. Chem Nat Compd 22(6):719–720
- 239 Arifkhozhaev AO, Rakhimov DA (1993) Polysaccharides of saponin-
240 bearing plants. IV. Structure of glucans A, B, and C of *Biebersteinia*
241 *multifida*. Chem Nat Compd 29(2):151–153
- 242 Arifkhozhaev AO, Kondratenko ES (1983) Polysaccharides of
243 saponin-bearing plants. I. Characteristics of the polysaccharides of
244 *Allochrysa gypsophiloides*. Chem Nat Compd 19:222–223
- 245 Arifkhozhaev AO, Arifkhozhaev KhA, Kondratenko ES (1985)
246 Polysaccharides of saponin-bearing plants. II. Isolation and charac-
247 terization of the polysaccharides of *Biebersteinia multifida*. Chem
248 Nat Compd 21(6):714–716
- 249 Atta AH, Mounier SM (2004) Antidiarrhoeal activity of some Egyptian
250 medicinal plant extracts. J Ethnopharmacol 92(2–3):303–309
- 251 Atta-ur-Rahman, Atia-tul-Wahab, Sultankhozhaev MN, Teshebaeva
252 UT, Choudhary MI (2005) Norditerpenoid alkaloids from *Aconitum*
253 *karakolicum* Rapais. Nat Prod Res 19(7):713–718
- 254 Awaad Amani AS, Maitland DJ, Soliman GA (2006) Antiulcerogenic
255 activity of *Alhagi maurorum*. Pharm Biol 44(4):292–296
- 256 Azimov MA, Nazirov ZN (1969s) A study of the triterpenes of
257 *Euphorbia jaxartica*. Chem Nat Compd 5(6):517
- 258 Azimov MA, Nazirov ZN (1970) Flavonoids of *Euphorbia jaxartica*
259 and *E. lamprocarpa*. Chem Nat Compd 6(2):268
- 260 Azizov KN, Aizikov MI, Akhmatova NF et al (1992) The influence of
261 leucomisine on thrombocyte aggregation in healthy volunteers and
262 in CHD patients. Meditsinskii Zh Uzbekistana (8):24–27 (in
263 Russian)
- 264 Bagirov VY, Sheichenko VI, Gasanova RY, Pimenov MG (1978) The
265 lactones of *Ferula malacophylla*. Chem Nat Compd 14(4):377–380
- 266 Baimukhamedov MA, Komissarenko NF (1990) Coumarins of
267 *Helichrysum maracandicum*. Chem Nat Compd 25(5):617
- 268 Bakiev A, Makhkamov S (1987) Zelyonie i Malo Rasprostanenniye
269 Ovoshchniye Kulturui (Green and not widely used vegetable crops).
270 Mekhnat, Tashkent (in Russian)
- 271 Balick MJ, Cox PA (1996) Plants, people, and culture: the science of
272 ethnobotany. W.H. Freeman/Scientific American Library, New York
- 273 Bandyukova VA, Khalmatov KhKh, Alimov KhI (1969) Flavonoids of
274 *Centaurea depressa*. Chem Nat Compd 5(4):274–275

- 277 Banthorpe DV, White JJ (1995) Novel anthraquinones from undifferentiated cell cultures of *Galium verum*. *Phytochemistry* 38(1):107–111
- 278 Barakat HH, Hussein SAM, Marzouk MS, Merfort I, Linscheid M, Nawwar MAM (1997) Polyphenolic metabolites of *Epilobium hirsutum*. *Phytochemistry* 46(5):935–941
- 281 Barnes J, Anderson LA, Phillipson DJ (2001) St John's wort (*Hypericum perforatum* L.): a review of its chemistry, pharmacology and clinical properties. *J Pharm Pharmacol* 53(5):583–600
- 282 Baser KHC, Nuriddinov KR, Nigmatullaev AM, Aripov KN (1997a) Essential oil of *Mentha asiatica* Boriss. from Uzbekistan. *Plant Physiol Biochem* 9(4):453–454
- 283 Baser KHC, Ozek T, Nuriddinov KhR, Nigmatullaev AM, Khadzimatov KKh, Aripov KhN (1997b) Essential oils of *Mediasia macrophylla* (Regel et Schmalh.) Pimen. and *Foeniculum vulgare* Mill. from Uzbekistan. *J Essent Oil Res* 9(2):249–250
- 284 Basher KHC, Ozek T, Demirchakmak B, Abduganiev BYo, Nuriddinov KhR, Aripov KhN, Doriev AS, Karataeva ChSh (1997) Essential oil of *Perovskia angustifolia* from Kyrgyzstan. *Chem Nat Compd* 33(3):296–298
- 285 Bazzaz BSF, Haririzadeh G (2003) Screening of Iranian plants for antimicrobial activity. *Pharm Biol* 41(8):573–583
- 286 Bedir E, Khan IA, Walker LA (2002) Biologically active steroidal glycosides from *Tribulus terrestris*. *Pharmazie* 57(7):491–493
- 287 Bekker NP, Ul'chenko NT, Glushenkova AI (2003) Lipids from the aerial part of *Crambe kotschyana*. *Chem Nat Compd* 39(2):218
- 288 Bekker NP, Ul'chenko NT, Glushenkova AI (2005) Lipids from *Descurainia sophia* seeds. *Chem Nat Compd* 41(3):346–347
- 289 Belodubrovskaya GA et al (2002) Encyclopedic dictionary of medicinal plants. <http://www.spcpa.ru/slr/preview/articles/1.html>. Accessed 22 Feb 2007 (in Russian)
- 290 Berezhinskaya VV, Trutneva EA (1959) Pharmacology of evoxin. *Farmakol Toksikol* 22(2):117–122 (in Russian)
- 291 Berezhinskaya VV, Aleshinskaya EE, Aleskina YaA (1968) Comparative pharmacological study of some alkaloids of aporphine group. *Farmakol Toksikol* 31(1):44–46 (in Russian)
- 292 Bishayee A, Sarkar A, Chatterjee M (1995) Hepatoprotective activity of carrot (*Daucus carota* L.) against carbon tetrachloride intoxication in mouse liver. *J Ethnopharmacol* 47(2):69–74
- 293 Blagojevic P, Radulovic N, Palic R, Stojanovic G (2006) Chemical composition of the essential oils of Serbian wild-growing *Artemisia absinthium* and *Artemisia vulgaris*. *J Agric Food Chem* 54(13):4780–4789
- 294 Blinova KF (1957) *Geum* species as tannin producing plants. *Trudy Leningradskogo Khimiko-farmatsevticheskogo Instituta* (2):80–90 (in Russian)
- 295 Blumenthal M (Senior ed) (1998) The complete German commission E monographs – therapeutic guide to herbal medicine. American Botanical Council, Austin
- 296 Bobokhanova LT, Bekturgunava SD (1996) Uzbekistan i Uzbeki (Uzbekistan and Uzbeks). Uzbekiston, Tashkent (in Russian)
- 297 Boguslavskaya LI, Dem'yanenko SI, Salam DK (1983) Flavonoids of some species of the genus *Dianthus*. *Chem Nat Compd* 19(3):366
- 298 Bohm BA, Stuessy TF (2001) Flavonoids of the sunflower family (Asteraceae). Springer, Vienna
- 299 Bondarenko NV (1972) Alkaloids of *Veratrum lobelianum*. IV. *Chem Nat Compd* 8(6):800–801
- 300 Bos R, Woerdenbag HJ, van Putten FM, Hendriks H, Scheffer JJ (1998) Seasonal variation of the essential oil, valerianic acid and derivatives, and velopotriates in *Valeriana officinalis* roots and rhizomes, and the selection of plants suitable for phytomedicines. *Planta Med* 64(2):143–147
- 301 Botirov EK, Drenin AA, Makarova AV (2006) A chemical study of flavanoids from medicinal and edible plants. *Khimiya rastitel'nogo syrya* (1):45–48 (in Russian)
- 302 Budincevic M, Vrbaski Z, Turkulov J, Dimic E (1995) Antioxidant activity of *Oenothera biennis* L. *Fett* 97(7–8):277–280
- 303 Bykov BA (1950) Elovyye lesa Tian'-Shanya ikh istoriya, osobennosti i tekhnologiya (Spruce forests of the Tian-Shan: their history, uniqueness and technology). *Isdatel'stvo Akademii Nauk, Alma-Ata*, pp 21–27 (in Russian)
- 304 Cantrell CL, Abate L, Fronczek FR, Franzblau SG, Quijano L, Fischer NH (1999) Antimycobacterial eudesmanolides from *Inula helenium* and *Rudbeckia subtomentosa*. *Planta Med* 65(4):351–355
- 305 Capek P, Rosik J, Kardosova A, Toman R (1987) Polysaccharides from the roots of the marsh mallow (*Althaea officinalis* L., var *Rhobusta*): structural features of an acidic polysaccharide. *Carbohydr Res* 164:443–452
- 306 Carnat AP, Carnat A, Fraisse D, Lamaison JL (1998) The aromatic and polyphenolic composition of lemon balm (*Melissa officinalis* L. subsp. *officinalis*) tea. *Pharm Acta Helv* 72(5):301–305
- 307 Carvalho JCT, Vignoli VV, de Souza GHB, Ujikawa K, Neto JJ (1999) Antimicrobial activity of essential oils from plants used in Brazilian popular medicine. *Acta Hort* 501:77–81
- 308 Cervato G, Carabelli M, Gervasio S, Cittera A, Cazzola R, Cestaro B (2000) Anti-oxidant properties of oregano (*Origanum vulgare*) leaf extracts. *J Food Biochem* 24(6):453–456
- 309 Cession-Fossion A (1967) Action of sympathicomimetic amines on the adrenal medulla of the rat perfused in vitro. *Arch Int Pharmacodyn Ther* 167(2):486–487 (in French)
- 310 Cha JD, Jeong MR, Jeong SI, Moon SE, Kim JY, Kil BS, Song YH (2005) Chemical composition and antimicrobial activity of the essential oils of *Artemisia scoparia* and *A. capillaris*. *Planta Med* 71(2):186–190
- 311 Chan K, Islam MW, Kamil M, Radhakrishnan R, Zakaria MNM, Habibullah M, Attas A (2000) The analgesic and anti-inflammatory effects of *Portulaca oleracea* L. subsp. *sativa* (Haw.) Celak. *J Ethnopharmacol* 73(3):445–451
- 312 Chefranova (1954) The genus *Thermopsis* R. Br. and its industrial importance. Synopsis of dissertation for the degree of Ph.D. in biology, Leningrad (in Russian)
- 313 Chelombito VA, Muravyova DA (1971) About the pharmacology of the alkaloid chelerythrine isolated from bokkonii melkoplodnoi (*Bocconia microcarpa* Maxim). In: *Aktualniye Problemy Farmakologii i Farmatsii* (Actual problems in pharmacology and pharmacy), Moscow, pp 183–188 (in Russian)
- 314 Chemonics International Inc (2001) Biodiversity assessment for Uzbekistan, task order under the biodiversity & sustainable forestry. IQC (BIOFOR) USAID contract number: LAG-I-00-99-00014-00, submitted to: USAID Central Asian Republics Mission, Almaty. http://pdf.usaid.gov/pdf_docs/PNACN475.pdf. Accessed 22 Feb 2007
- 315 Chen G, Todorov DK, Zeller WJ (1992) Antitumor activity of thaliblastine (NSC-68075) in experimental ovarian tumor cell lines sensitive and resistant to cisplatin. *Cancer Lett* 62(2):173–177
- 316 Chen B, Teranishi R, Kawazoe K, Takaishi Y, Honda G, Itoh M, Takeda Y, Kodzhimatov OK (2000) Sesquiterpenoids from *Ferula kuhistanica*. *Phytochemistry* 54(7):717–722
- 317 Chen J, Shi Y, Liu J (2003) Determination of noradrenaline and dopamine in Chinese herbal extracts from *Portulaca oleracea* L. by high-performance liquid chromatography. *J Chromatogr A* 1003(1–2):127–132
- 318 Chen Z, Lu Y, Wang Z, Tao X, Wei D (2007) Protective effects of salidroside on hypoxia/reoxygenation injury by sodium hydrosulfite in pc12 cells. *Pharm Biol* 45(8):604–612
- 319 Cheney RH (1963) Therapeutic potential of *Eschscholtzia californica* herba. *Pharm Biol* 3(3):413–416
- 320 Cheng G, Bai Y, Zhao Y, Tao J, Liu Y, Tu G, Ma L, Liao N, Xu X (2000) Flavonoids from *Ziziphus jujuba* Mill var. *spinosa*. *Tetrahedron* 56(45):8915–8920
- 321 Chen-Gu Ch (1957) Certain alkaloids from Chinese medicinal plants. *Farmakol Toksikol* 20(1):49–52 (in Russian)
- 322 Chernenko TV, Glushenkova AI, Kotenko LD (2000) Lipids of *Arum korolkowii* tubers. *Chem Nat Compd* 36(6):565–567

- 409 Chernenko TV, Glushenkova AI, Nigmatullaev AM (2002) Lipids from
410 *Mediasia macrophylla* leaves. Chem Nat Compd 38(4):307–309
- 411 Chernenko TV, Gazizov FYu, Glushenkova AI, Nigmatullaev AM
412 (2005) Lipids of *Eminium lehmannii* leaves and tubers. Chem Nat
413 Compd 41(2):141–143
- 414 Chiang LC, Chiang W, Chang MY, Ng LT, Lin CC (2002) Antiviral
415 activity of *Plantago major* extracts and related compounds in vitro.
416 Antiviral Res 55(1):53–62
- 417 Chicco A (1966) *Prangos pabularia*, an Indian medicinal plant. Corriere
418 del Farmacista 21(17):440–441
- 419 Chikov PS (1989) Лекарственные растения: справочник, 2-е изд,
420 Лекарственные растения: справочник, изд 2 (Medicinal plants:
421 manual, 2nd edn). Agropromizdat, Moscow [in Russian]
- 422 Cho J, Kim HM, Ryu JH, Jeong YS, Lee YS, Jin C (2005) Neuroprotective
423 and antioxidant effects of the ethyl acetate fraction prepared from
424 *Tussilago farfara* L. Biol Pharm Bull 28(3):455–460
- 425 Cho JY, Yoo ES, Cha BC, Park HJ, Rhee MH, Han YN (2006) The
426 inhibitory effect of triterpenoid glycosides originating from
427 *Sanguisorba officinalis* on tissue factor activity and the production
428 of TNF- α . Planta Med 72(14):1279–1284
- 429 Chodoeva A, Bosc JJ, Guillon J, Decendit A, Petraud M, Absalon C,
430 Vitry C, Jarry C, Robert J (2005) 8-O-Azeloyl-14-benzoylaconine: a
431 new alkaloid from the roots of *Aconitum karacolicum* Rapcs. and its
432 antiproliferative activities. Bioorg Med Chem 13(23):6493–6501
- 433 Chowdhury AR, Sharma S, Mandal S, Goswami A, Mukhopadhyay S,
434 Majumder HK (2002) Luteolin, an emerging anti-cancer flavonoid,
435 poisons eukaryotic DNA topoisomerase I. Biochem J 366(Pt
436 2):653–661
- 437 Chumbalov TK, Bikbulatova TN (1970) Polyphenolic compounds of
438 *Geranium collinum*. IV. Chem Nat Compd 6(4):523–524
- 439 Chumbalov TK, Omurkamzinova VB (1969) Flavonoids of the leaves
440 of *Polygonum coriarium*. I. Chem Nat Compd 4(5):72
- 441 Chumbalov TK, Bikbulatova TN, Chanysheva IS (1968) Flavone gly-
442 cosides of *Geranium collinum*. III. Chem Nat Compd 4(6):323
- 443 Chumbalov TK, Mukhamed'yarova MM, Chanysheva IS (1970)
444 Flavonoids of *Atraphaxis spinosa*. I. Chem Nat Compd 6(5):639
- 445 Chumbalov TK, Mukhamed'yarova MM, Chanysheva IS, Il'yasova
446 MM (1971) Flavonoids of *Atraphaxis pyrifolia* and *spinosa*. II.
447 Chem Nat Compd 7(4):504–505
- 448 Coates GH, Cox B (1972) Harmine tremor after brain monoamine oxi-
449 dase inhibition in the mouse. Eur J Pharmacol 18(2):284–286
- 450 Colaric M, Veberic R, Solar A, Hudina M, Stampar F (2005) Phenolic
451 acids, syringaldehyde, and juglone in fruits of different cultivars of
452 *Juglans regia* L. J Agric Food Chem 53(16):6390–6396
- 453 Conti A (1962) Study of the antitumor activity of berberine, particularly
454 of the protoberberine alkaloid group. Referativni Zh Biokhimiya
455 (11):1464 (in Russian)
- 456 Curini M, Epifano F, Genovese S, Tammaro F, Menghini L (2006)
457 Composition and antimicrobial activity of the essential oil of
458 *Artemisia dracunculus* "Piemontese" from Italy. Chem Nat Compd
459 42(6):738–739
- 460 Danilevskii BL, Tulyaganov NT, Sadritdinov FS (1972) Ataractic
461 actions of alkaloids haplofilidine, perforine, gentianine, and tetrahy-
462 droberberine compared to meprostan and aminazin. Dokl Akad Nauk
463 UzSSR 29(3):37–38 (in Russian)
- 464 De Combarieu E, Fuzzati N, Lovati M, Mercalli E (2003) Furostanol
465 saponins from *Tribulus terrestris*. Fitoterapia 74(6):583–591
- 466 de Meyts P, Cession-Fossion A (1966) Ephedrine as a sympathicom-
467 imetic amine with indirect action in the rat. Comptes Rendus des
468 Séances de la Société de Biologie et de Ses Filiales 160(11):2224–
469 2227 (in French)
- 470 de Meyts P, Cession-Fossion A (1967) D(-)-ephedrine and L(+)-psi-
471 ephedrine as sympathomimetic amines with indirect action in the
472 rat. Comptes Rendus des Séances de la Société de Biologie et de Ses
473 Filiales 161(5):1162–1164 (in French)
- de Meyts P, Cession-Fossion A (1968) Vasomotor action of D(-)- 474
ephedrine and L(+)-psi-ephedrine in rats. Arch Int Pharmacodyn 475
Ther 174(1):233–242 (in French) 476
- de Sousa AC, Alviano DS, Blank AF, Alves PB, Alviano CS, Gattass 477
CR (2004) *Melissa officinalis* L. essential oil: antitumoral and anti- 478
oxidant activities. J Pharm Pharmacol 56(5):677–681 479
- Deepak M, Dipankar G, Prashanth D, Asha MK, Amit A, Venkataraman 480
BV (2002) Tribulosin and β -sitosterol-D-glucoside, the anthelmintic 481
principles of *Tribulus terrestris*. Phytomedicine 9(8): 753–756 482
- Deineka VI, Grigor'ev AM, Borzenko ON, Staroverov VM, Trubitsyn MA 483
(2004) HPLC analysis of anthocyanins: cyanidine glycosides from 484
fruits of plants of the *Prunus* genus. Pharm Chem J 38(8):437–440 485
- Delova GB, Guskova IN (1974) Antibacterial and antifungal effects of 486
essential oil from some species of *Ziziphora*. In: Sobolevskaya KA 487
(ed) Complex studies of the useful plants of Siberia. Nauka, 488
Novosibirsk, pp 131–146 (in Russian) 489
- Demirezer LO, Gurbuz F, Guvenalp Z, Stroch K, Zeeck A (2006) 490
Iridoids, flavonoids and monoterpene glycosides from *Galium* 491
verum subsp. *verum*. Turk J Chem 30(4):525–534 492
- Derksen GC, Niederlander HA, van Beek TA (2002) Analysis of 493
anthraquinones in *Rubia tinctorum* L. by liquid chromatography 494
coupled with diode-array UV and mass spectrometric detection. J 495
Chromatogr A 978(1–2):119–127 496
- Desbène S, Hanquet B, Shoyama Y, Wagner H, Lacaille-Dubois M 497
(1999) Biologically active triterpene saponins from callus tissue of 498
Polygala amarella. J Nat Prod 62(6):923–926 499
- Dey M, Ripoll C, Pouleva R, Dorn R, Aranovich I, Zurov D, Kurmukov 500
A, Eliseyeva M, Belolipov I, Akimaliev A, Sodobekov I, 501
Akimaliev D, Lila MA, Raskin I (2008) Plant extracts from central 502
Asia showing antiinflammatory activities in gene expression assays. 503
Phytother Res 22(7):929–934 504
- Dilman VM, Kozhina IC, Klyuchnikova LA, Kibalchich PN, Kusov 505
Yu Yu (1968) Study of the abilities and constituents of *Lithospermum* 506
officinale: perspectives on use in oncology. Vopr Onkol 14(7):86–91 507
(in Russian) 508
- Dobrokhotova K, Chudinov V (1966) Medicinal plants of Kazakhstan. 509
Kazakhstan, Alma-Ata (in Russian) 510
- Dolya VS, Fursa NS, Litvinenko VI (1973a) Flavonoids of the seeds 511
of some species of the family Cruciferae. Chem Nat Compd 512
9(6):766 513
- Dolya VS, Shkurupii EN, Fursa NS (1973b) The oils of four plants. 514
Chem Nat Compd 9(5):578–579 515
- Dozortseva PM (1958) Mellictine – a new preparation for treatment of 516
spastic paralysis. Meditsinskaya promuihlnennost SSSR (11):54 (in 517
Russian) 518
- Dozortseva PM (1959) The pharmacology of the alkaloid methyllyca- 519
conitine (mellictine). Farmakol Toksikol 22(1):34–38 (in Russian) 520
- Du J, He ZD, Jiang RW, Ye WC, Xu HX, But PPH (2003) Antiviral 521
flavonoids from the root bark of *Morus alba* L. Phytochemistry 522
62(8):1235–1238 523
- Dvomyk AS, Pererva TP, Kukakh VA (2002) Screening substances 524
derived from cultures of medicinal plants for antimutagenic activity 525
in the *Escherichia coli*-bacteriophage lambda system. Tsitologii i 526
Genetika 36(2):3–10 (in Ukrainian) 527
- Dzhakupova NU (1968) The medicinal plant *Delphinium confusum* in 528
the western Tian Shan. Vestnik Akademii Nauk KazSSR (11):66–68 529
(in Russian) 530
- Dzhangaliev AD, Salova TN, Turekhanova PM (2003) The wild fruit 531
and nut plants of Kazakhstan. Horticult Rev 29 532
- Dzhumabaev TZ (1972) The pharmacology of *Berberis oblonga* and 533
Berberis nummularia. Synopsis of dissertation for the degree of 534
Ph.D., Ryazan 535
- Dzhumabaev TZ, Ibragimov II, Lutfullin KL (1970) Effect of the sum of 536
alkaloids in the root and bark of the stem of the oblong barberry on the 537
blood coagulation system. Meditsinskii Zh Uzbekistana (8):22–25 538

- 539 Dzharmaev K (1980) The most prospective wild-growing aromatic
540 plants of the Surkhan-Sherabad valley and their economic value. In:
541 Aktualnuiye voprosy izucheniya I ispolzobaniya efirnikh masel
542 (Proceedings of 3rd symposium "Important questions in the study
543 of using essential oil bearing plants and essential oil"). Simferopol,
544 24–26 Sept 1980, pp 96–97 (in Russian)
- 545 Eliseeva M (2005) Internation cooperative biodiversity groups (ICBG)
546 Central Asia 2005 progress report (unpublished report)
- 547 El-Mallah MH, El-Shami SM, Hassanein MM (2003) Detailed studies
548 on some lipids of *Silybum marianum* (L.) seed oil. *Grasas y Aceites*
549 54(4):397–402
- 550 El-Razek MHA, Ohta S, Ahmed AA, Hirata T (2001) Monoterpene
551 coumarins from *Ferula ferulago*. *Phytochemistry* 57(8):1201–1203
- 552 Emami SA, Amin-Ar-Ramimeh E, Ahi A, Kashy MRB, Schneider B,
553 Iranshahi M (2007) Prenylated flavonoids and flavonostilbenes from
554 *Sophora pachycarpa* roots. *Pharm Biol* 45(6):453–457
- 555 Enneking D (1995) The toxicity of *Vicia* species and their utilisation as
556 grain legumes, vol 6, Centre for legumes in mediterranean agricul-
557 ture (CLIMA) occasional publication. University of Western
558 Australia, Nedlands
- 559 Erdogru OT (2002) Antibacterial activities of some plant extracts used
560 in folk medicine. *Pharm Biol* 40(4):269–273
- 561 Ermishina OA, Kurmukov AG, Salikhova RE, Aizikov MI (1982)
562 Effect of ecdysterone on enzyme levels during experimental myo-
563 cardial infarction. *Meditsinskii Zh Uzbekistana* (11):90–91 (in
564 Russian)
- 565 Ersoz T, Calis I (1991) C-glucosylflavones from *Gentiana olivieri*.
566 Hacettepe Universitesi Eczacilik Fakultesi Dergisi 11:29–36
- 567 Eshbakova KA, Saidkhodzhaev AI (2001) Triterpenoids and sterols
568 from three species of *Pulicaria*. *Chem Nat Compd* 37(2):196–197
- 569 Eshbakova KA, Sagitdinova GV, Levkovich MG, Rasulev BF, Abdullaev
570 ND, Malikov VM (1997) Diterpenoids of *Pulicaria salviifolia* IV.
571 Structures of salvicinolide and salvicinolin. *Chem Nat Compd*
572 33(4):458–461
- 573 Esmaeili A, Rustaiyan A, Nadimi M, Masoudi S, Tadavon F, Sedaghat
574 S, Ebrahimpur N, Haiyzadeh E (2005) Volatile constituents of
575 *Centaurea depressa* M.B. and *Carduus pycnocephalus* L. two com-
576 positae herbs growing wild in Iran. *J Essent Oil Res* 17(5):539–541
- 577 Estrada A, Katselis GS, Laarveld B, Barl B (2000) Isolation and evalua-
578 tion of immunological adjuvant activities of saponins from *Polygala*
579 *senega* L. *Comp Immunol Microbiol Infect Dis* 23(1):27–43
- 580 Evans WC (2002) Production of crude drugs. In: Evans WC (ed) *Trease*
581 and *Evans pharmacognosy*, 15th edn. W.B. Saunders, Edinburgh, pp
582 61–66
- 583 Evdokimova NI, Kurmukov AG (1972) Effect of skimmianine in com-
584 parison with meprotran on a change in the EEG (electroencephalo-
585 gram) of rabbits. *Meditsinskii Zh Uzbekistana* (5):61 (in Russian)
- 586 Fakhrutdinov SF (1971) Comparative pharmacological study of the
587 methiodides of corytuberine alkaloids and iodides of magnoflorine
588 and O, O-diacetylmagnoflorine. In: Sultanov MB (ed) *Farmakologiya*
589 *Alkaloidov i Serdechnykh Glikozidov*. FAN, Tashkent, pp 155–158
590 (in Russian)
- 591 Fakhrutdinov SF, Sultanov MB (1972) Pharmacology of lirioidenine,
592 thalicminine, corydine and isocorydine hydrochlorides. In: Sultanov
593 MB (ed) *Farmakologiya Alkaloidov Serdechnykh Glikozidov*. FAN,
594 Tashkent (in Russian)
- 595 Faramarzi MA, Moghimi M, Monsef-Esfahani HR, Shahverdi AR,
596 Khodae S (2008) Chemical composition and antimicrobial activity
597 of essential oils from *Geum kokanicum*. *Chem Nat Compd*
598 44(6):811–813
- 599 Farsam H, Amanlou M, Dehpour AR, Jahaniani F (2000) Anti-
600 inflammatory and analgesic activity of *Biebersteinia multifida* DC.
601 root extract. *J Ethnopharmacol* 71(3):443–447
- 602 Farzami B, Ahmadvand D, Vardasbi S, Majin FJ, Khaghani Sh (2003)
603 Induction of insulin secretion by a component of *Urtica dioica* leave
604 extract in perfused Islets of Langerhans and its in vivo effects in nor-
605 mal and streptozotocin diabetic rats. *J Ethnopharmacol* 89(1):47–53
- Fazly Bazzaz BS, Haririzadeh G (2003) Screening of Iranian plants for
606 antimicrobial activity. *Pharm Biol* 41(8):573–583
- 607 Fecka I, Cisowski W (2005) Tannins and flavonoids from the *Erodium*
608 *cicutarium* herb. *Z Naturforsch B: Chem Sci* 60(5):555–560
- 609 Fecka I, Gasiorowski K, Brokos B (1997) The phytochemical analysis
610 and immunotropic activity evaluation of polyphenolic fraction from
611 the common storks bill herb (*Erodium cicutarium* (L.) L'Herit.).
612 *Herba Pol* 43(3):214–221
- 613 Feng YL, Li HR, Xu LZ, Yang SL (2007) 27-Nor-triterpenoid glyco-
614 sides from the barks of *Zygophyllum fabago* L. *J Asian Nat Prod*
615 *Res* 9(6):505–510
- 616 Flora of Kirghiz SSR. 1952–1965, vols 1–11. Izdatelstvo Akademii
617 Nauk Kirghizskoi SSR, Frunze (in Russian)
- 618 Flora of North America Editorial Committee (ed) (1993+) *Flora of*
619 *North America north of Mexico*, vol 12+. Oxford University Press,
620 New York/Oxford
- 621 Flora of Uzbekistan. Vol 1–6. Izdatelstvo Akademii Nauk Uzbekskoi
622 SSR, Tashkent (in Russian)
- 623 Fontana E, Hoeberechts J, Nicola S, Cros V, Battista Palmegiano G,
624 Peiretti PG (2006) Nitrogen concentration and nitrate/ammonium
625 ratio affect yield and change the oxalic acid concentration and fatty
626 acid profile of purslane (*Portulaca oleracea* L.) grown in a soilless
627 culture system. *J Sci Food Agric* 86(14):2417–2424
- 628 Frank BS, Michelson WB, Panter KE, Gardner DR (1995) Ingestion of
629 poison hemlock (*Conium maculatum*). *West J Med* 163(6):573–574
- 630 Fruentov NK (1972) Lekarstvennuiye rasteniya Dalnego Vostoka
631 (Medicinal plants of the Far East). Khabarovskoye knizhnoye izda-
632 telstvo, Khabarovsk (in Russian)
- 633 Fu B, Su BN, Takaishi Y, Honda G, Ito M, Takeda Y, Kodzhimatov OK,
634 Ashurmetov O (2001) A bis-sesquiterpene and sesquiterpenolides
635 from *Inula macrophylla*. *Phytochemistry* 58(7):1121–1128
- 636 Fu Y, Hsieh TC, Guo J, Kunicki J, Lee MY, Darzynkiewicz Z, Wu JM
637 (2004) Licochalcone-A, a novel flavonoid isolated from licorice
638 root (*Glycyrrhiza glabra*), causes G2 and late-G1 arrests in andro-
639 gen-independent PC-3 prostate cancer cells. *Biochem Biophys Res*
640 *Commun* 322(1):263–270
- 641 Fukudu T, Ito H, Yoshida T (2003) Antioxidative polyphenols from
642 walnuts (*Juglans regia* L.). *Phytochemistry* 63(7):795–801
- 643 Fursa NS, Litvinenko VI (1970) Lepidoside from *Lepidium perfolia-*
644 *tum*. *Chem Nat Compd* 6(1):119
- 645 Fushiya S, Kabe Y, Ikegaya Y, Takano F (1998) (+)-rhododendrol and
646 epi-rhododendrin suppress the NO production by activated mac-
647 rophages in vivo. *Planta Med* 64(7):598–602
- 648 Gadgoli C, Mishra SH (1999) Antihepatotoxic activity of p-methoxy
649 benzoic acid from *Capparis spinosa*. *J Ethnopharmacol* 66(2):
650 187–192
- 651 Gammerman AF (1960) *Kurs Farmakognozii* (Pharmacognosy course),
652 Medgiz (in Russian)
- 653 Gammerman AF (1967) *Kurs Farmakognozii*, izd. 6 (Pharmacognosy
654 course), 6th edn. Meditsina, Leningrad (in Russian)
- 655 Gammerman AF, Kadaev GN, Yatsenko-Hmelevsky AA (1990)
656 *Lekarstvennuiye Rasteniya* (Medicinal plants). Vuishaya shkola,
657 Moscow (in Russian)
- 658 Gan PA (1970) *Ekologicheskoye osnovuiy introduktsii i lesorazvedeniya*
659 *v poyasye yelovuiikh lesov Tyan-Shan* (Basic ecology of introduc-
660 tion and reforestation in the Spruce Belt of the Tian-Shan). Ilim,
661 Frunze (in Russian)
- 662 Gan LX, Han XB, Chen YQ (1986a) Astrasieversianins IX, XI and XV,
663 cycloartane derived saponins from *Astragalus sieversianus*.
664 *Phytochemistry* 25(6):1437–1441
- 665 Gan LX, Han XB, Chen YQ (1986b) The structures of thirteen
666 astrasieversianins from *Astragalus sieversianus*. *Phytochemistry*
667 25(10):2389–2393
- 668

- 669 Ganenko TV, Isaev MI, Gromova AS, Abdullaev ND, Lutskii VI, Larin
670 MF, Semenov AA, Gorovits MB, Abubakirov NK (1986) Triterpene
671 glycosides and their genins from *Thalictrum foetidum*. IV. Structure
672 of cyclofoetigenin B. *Chem Nat Compd* 22(3):288–294
- 673 Gapparov AM, Razzakov NA, Aripova SF (2007) Alkaloids of
674 *Convolvulus subhirsutus* from Uzbekistan. *Chem Nat Compd*
675 43(3):291–292
- 676 Garbacki N, Tits M, Angenot L, Damas J (2004) Inhibitory effects of
677 proanthocyanidins from *Ribes nigrum* leaves on carrageenin acute
678 inflammatory reactions induced in rats. *BMC Pharmacol* 4(1):25
- 679 Gauthaman K, Ganesan AP, Prasad RN (2003) Sexual effects of punc-
680 turevine (*Tribulus terrestris*) extract (protodioscin): an evaluation
681 using a rat model. *J Altern Complement Med* 9(2):257–265
- 682 Gazizov FYu, Glushenkova AI (1997) Phospholipids of the seeds of
683 *Fumaria vaillantii*. *Chem Nat Compd* 33(6):627–629
- 684 Geetha S, Sai Ram M, Singh V, Ilavazhagan G, Sawhney RC (2002)
685 Anti-oxidant and immunomodulatory properties of seabuckthorn
686 (*Hippophae rhamnoides*)—an in vitro study. *J Ethnopharmacol*
687 79(3):373–378
- 688 Georgadze VH (1938) Pharmacology of the alkaloids from *Sophora*.
689 *Fiziologicheskii Zh SSSR* 25(1–2):179–195 (in Russian)
- 690 Germano MP, De Pasquale R, D'Angelo V, Catania S, Silvani V, Costa
691 C (2002) Evaluation of extracts and isolated fraction from *Capparis*
692 *spinosa* L. buds as an antioxidant source. *J Agric Food Chem*
693 50(5):1168–1171
- 694 Ghafari H, Yasa N, Mohammadirad A, Dehghan G, Zamani MJ,
695 Nikfar S, Khorasani R, Minaie B, Abdollahi M (2006) Protection
696 by *Ziziphora clinopoides* of acetic acid-induced toxic bowel
697 inflammation through reduction of cellular lipid peroxidation
698 and myeloperoxidase activity. *Hum Exp Toxicol* 25(6):
699 325–332
- 700 Ghannadi A, Zolfaghari B (2003) Compositional analysis of the essen-
701 tial oil of *Lallemantia royleana* (Benth. in Wall.) Benth. from Iran.
702 *Flavour Frag J* 18(3):237–239
- 703 Gibbons S, Oluwatuyi M, Veitch NC, Gray AI (2003) Bacterial resis-
704 tance modifying agents from *Lycopus europaeus*. *Phytochemistry*
705 62(1):83–87
- 706 Gilani AH, Janbaz KH, Lateef A, Zaman M (1994) Ca⁺⁺ channel
707 blocking activity of *Artemisia scoparia* extract. *Phytother Res*
708 8(3):161–165
- 709 Glasl S, Presser A, Gunbilig D, Werner I, Narantuya S, Haslinger E,
710 Jurenitsch J, Kubelka W (2001) Highly hydroxylated guaianolides
711 of *Achillea asiatica* and Middle European *Achillea* species.
712 *Phytochemistry* 58(8):1189–1194
- 713 Goel HC, Prasad J, Singh S, Sagar RK, Prem Kumar I, Sinha AK (2002)
714 Radioprotection by a herbal preparation of *Hippophae rhamnoides*,
715 RH-3, against whole body lethal irradiation in mice. *Phytomedicine*
716 9(1):15–25
- 717 Golcu A, Dolaz M, Diğrak M, Serin S (2002) The biological activity of
718 dyer's madder (*Rubia tinctorium* L.). In: Yayli N, Kucuk M (eds)
719 Proceedings of the 1st international congress on the chemistry of
720 natural products. Karadeniz University, Trabzon, pp 255–258
- 721 Golovina LA, Nikonov GK (1973) The structure of drupanol – a new
722 phenol from *Psoralea drupacea*. *Chem Nat Compd* 9(1):7–9
- 723 Golovkova AG (1990) Rastitelnost Kirgizii (ratsionalnoye ispolzo-
724 vaniye i okhrana) [Vegetation of Kyrgyzstan (rational use and pro-
725 tection)]. Ilim, Frunze (in Russian)
- 726 Gorkin VZ (1964) Current progress in the study of the nature and physi-
727 ological role of mitochondrial monoamine oxidase. *Voprosy med-*
728 *itsinskoi khimii* (10):115–134 (in Russian)
- 729 Gozler B, Gozler T, Freer AJ, Shamma M (1988) Some unusual pavine
730 and isopavine alkaloids from *Roemeria refracta*. *J Nat Prod*
731 51(4):760–764
- 732 Gozler B, Kivçak B, Gozler T, Shamma M (1990a) Two new benzyltetra-
733 rahydroisoquinoline alkaloids from *Roemeria refracta*. *J Nat Prod*
734 53(3):666–668
- Gozler B, Ozic P, Freyer AJ, Shamma M (1990b) Morphinandienone
alkaloids from *Roemeria refracta*. *J Nat Prod* 53(4):986–988
- Greger H, Hofer O (1980) New unsymmetrically substituted tetrahy-
drofurofuran lignans from *Artemisia absinthium*. Assignment of the
relative stereochemistry by lanthanide induced chemical shifts.
Tetrahedron 36(24):3551–3558
- Grinkevich NE (Senior ed) (1991) Lekarstvennuiye Rasteniy: Sprav.
Posobiye (Medicinal plants: infomative manual). Vuishaya shkola,
Moscow (in Russian)
- Gudej J (1991) Flavonoids, phenolic acids and coumarins from the
roots of *Althaea officinalis*. *Planta Med* 57(3):284–285
- Gudej J, Tomczyk M (2004) Determination of flavonoids, tannins and
ellagic acid in leaves from *Rubus* L. species. *Arch Pharm Res*
27(11):1114–1119
- Guil-Guerrero JL, Rodríguez-García I (1999) Lipids classes, fatty acids
and caroteners of the leaves of six edible wild plants. *Eur Food Res*
Technol 209(5):313–316
- Guil-Guerrero JL, García Maroto FF, Giménez A (2001) Fatty acid
profiles from forty-nine plant species that are potential new sources
of γ -linolenic acid. *J Am Oil Chem Soc* 78(7):677–684
- Gulcin I, Kufrevioglu OI, Oktay M, Buyukokuroglu ME (2004)
Antioxidant, antimicrobial, antiulcer and analgesic activities of
nettle (*Urtica dioica* L.). *J Ethnopharmacol* 90(2–3):205–215
- Gulluce M, Sahin F, Sokmen M, Ozer H, Daferera D, Sokmen A,
Polissiou M, Adiguzel A, Ozkan H (2007) Antimicrobial and anti-
oxidant properties of the essential oils and methanol extract from
Mentha longifolia L. ssp. *Longifolia*. *Food Chem* 103(4):
1449–1456
- Gupta SPK (2002) Rustom Jal Vakil (1911–1974) father of modern car-
diology – a profile. *J Indian Acad Clin Med* 3(1):100–104
- Gurbuz I, Ustun O, Yesilada E, Sezik E, Kutsal O (2003) Anti-
ulcerogenic activity of some plants used as folk remedy in Turkey. *J*
Ethnopharmacol 88(1):93–97
- Gusakova SD, Khomova TV (1997) Lipids of *Ziziphora pedicellata*.
Chem Nat Compd 33(6):633–635
- Gusakova SD, Kholmatov KhK, Umarov AU (1976) Composition of
the seed oils of *Origanum tyttanthum* and *Mentha asiatica*. *Chem*
Nat Compd 12(2):134–139
- Gusinin IA (1962) Toksikologiya Yadovituikh Rasteniy (Toxicology
of poisonous plants). Izdaniye selkhozliteraturi zhurnalov i plaka-
tov, Moscow (in Russian)
- Haddad M, Larchevêque M (2003) Diastereocontrolled synthesis of
(–)-codonopsinine. *Synlett* 2:274–276
- Hamilton AC (2004) Medicinal plants, conservation and livelihoods.
Biodivers Conserv 13:1477–1517
- Han JW, Lee BG, Kim YK, Yoon JW, Jin HK, Hong S, Lee HY, Lee
KR, Lee HW (2001) Ergolide, sesquiterpene lactone from *Inula bri-*
tannica, inhibits inducible nitric oxide synthase and cyclo-oxyge-
nase-2 expression in RAW 264.7 macrophages through the
inactivation of NF- κ B. *Br J Pharmacol* 133(4):503–512
- Harrison DM (1990) Steroidal alkaloids. *Nat Prod Rep* 7(2):139–147
- Hartleb I, Seifert K (1995) Triterpenoid saponins from *Verbascum son-*
garicum. *Phytochemistry* 38(1):221–224
- Hofer O, Szabó G, Greger H (1986) 2-Hydroxy-4-methoxy-trans-
cinnamic acid as a precursor of herniarin in *Artemisia dracunculus*.
Org Chem Biochem 117(10):1219–1222
- Hosseinnimehr SJ, Pourmorad F, Shahabimajid N, Shahrbandy K,
Hosseinzadeh R (2007) In vitro antioxidant activity of *Polygonium*
hyrcanicum, *Centaurea depressa*, *Sambucus ebulus*, *Mentha spicata*
and *Phytolacca americana*. *Pak J Biol Sci* 10(4):637–640
- Hou BL, Li ZL, Li X (2005) A new flavonoid glycoside from the roots
and stems of *Sphaerophysa salsula*. *Yao Xue Xue Bao* 40(6):533–
535 (in Chinese)
- Hsu FL, Chen YC, Cheng JT (2000) Caffeic acid as active principle
from the fruit of *Xanthium strumarium* to lower plasma glucose in
diabetic rats. *Planta Med* 66(3):228–230

- 801 Hu C, Kitts DD (2004) Luteolin and luteolin-7-O-glucoside from dan- 866
 802 dellion flower suppress iNOS and COX-2 in RAW264.7 cells. *Mol* 867
 803 *Cell Biochem* 265(1–2):107–113
- 804 Hu KF, Hou XY, Gong YT, Abbas A (2006) Analysis of physical-chem- 868
 805 ical properties and fatty acid compositions of *Hibiscus trionum* L. 869
 806 seed oil. *Food Sci* 27(11):455–457 (in Chinese)
- 807 Hvattum E (2002) Determination of phenolic compounds in rose hip 870
 808 (*Rosa canina*) using liquid chromatography coupled to electrospray 871
 809 ionisation tandem mass spectrometry and diode-array detection. 872
 810 *Rapid Commun Mass Spectrom* 16(7):655–662
- 811 Hwang SB, Chang MN, Garcia ML, Han QQ, Huang L, King VF, 873
 812 Kaczorowski GJ, Winqvist RJ (1987) L-652,469 – a dual receptor 874
 813 antagonist of platelet activating factor and dihydropyridines from 875
 814 *Tussilago farfara* L. *Eur J Pharmacol* 141(2):269–281
- 815 Iacobellis NS, Cantore PL, Capasso F, Senatore F (2005) Antibacterial 876
 816 activity of *Cuminum cyminum* L. and *Carum carvi* L. essential oils. 877
 817 *J Agric Food Chem* 53(1):57–61
- 818 Iauk L, Lo Bue AM, Milazzo I, Rapisarda A, Blandino G (2003) 878
 819 Antibacterial activity of medicinal plant extracts against periodon- 879
 820 topathic bacteria. *Phytother Res* 17(6):599–604
- 821 Ibragimov FI, Ibragimova VS (1960) Osnovnuiye Lekarstvennuiye 880
 822 Sredstva Kitaiskoi Meditsinui (The basic medicinal remedies of 881
 823 Chinese medicine). Medgiz, Moscow (in Russian)
- 824 Ibragimov AY, Khazanovich RL (1972) Ursolic acid from *Agrimonia* 882
 825 *asiatica*. *Chem Nat Compd* 8(3):390–391
- 826 Ibragimov II, Dzhumabaev TZ (1971) Influence of *Berberis oblonga* on 883
 827 blood coagulation. In: *Problemy Svyortuivaniya Krovi i Gemostaza* 884
 828 (Problems of blood coagulation and hemostasis), vol 33. Nauchni 885
 829 Trudy Kubanskogo Med Instituta (Scientific Works of Kuban 886
 830 Medical Institute), pp 110–114
- 831 Ibragimova MU, Israilov IA, Yunusov MS, Yunusov SYu (1974) 887
 832 Alkaloids of *Fumaria vaillantii* structure of vaillantine. *Chem Nat* 888
 833 *Compd* 10(4):481–482
- 834 Idzumi I, Conti A (1962) Study of the activity of the alkaloid berberine 889
 835 on the intestines and uterus in mice. *Referativnii Zh Biokhimii* 890
 836 (12):1504 (in Russian)
- 837 Iida A, Kiuchi F, Ito M, Honda G, Mizushina Y, Yoshida H, Sarsenbaev 891
 838 K (2003) Phenylethanoid glycosides targeting mammalian DNA 892
 839 polymerases. *Nat Med* 5(4):146–149
- 840 Iinuma M, Ohyama M, Tanaka T (1995) Six flavonostilbenes and a 893
 841 flavanone in roots of *Sophora alopecuroides*. *Phytochemistry* 894
 842 38(2):519–525
- 843 Ikan R, Ravid U (1971) The isolation and identification of ecdysterone 895
 844 from *Ajuga iva*. *Planta Med* 20(1):33–35
- 845 Ilarionova M, Maneva K, Todorov D (1980) Pathomorphological stud- 896
 846 ies on animals bearing experimental tumors treated with thaliblas- 897
 847 tine. *Dokl Bolgarskoi Akad Nauk* 33(5):719–722
- 848 Iordanov D, Nikolov P, Boichinov A (1970) Fitoterapiya. Lecheniye 898
 849 Lekarstvennuiimi Travami. Vtoroye dopolnennoye Russkoye 899
 850 Izdaniye. (Phytotherapy: treatment with medicinal herbs, 2nd 900
 851 Russian ed). Meditsina i Fizkultura, Sofia (in Russian)
- 852 Isakov SHI (1969) Study of medicinal plant material used by Kyrgyz 901
 853 people. Synopsis of dissertation for the degree of Ph.D. in medicine, 902
 854 Ashkabad (in Russian)
- 855 Ishmukhamedov NA, Sultanov MB (1965) Pharmacology of the new 903
 856 alkaloid alginidine. In: *Farmakologiya Alkaloidov* (Pharmacology 904
 857 of alkaloids). Nauka, Tashkent, pp 342–346 (in Russian)
- 858 Iskanderova SD, Sharipova UC (1992) Influence of *Ecdysten* on physi- 905
 859 cal working capacity of patients, after myocardial infarction. 906
 860 *Meditsinskii Zh Uzbekistana* (3):14–16 (in Russian)
- 861 Ivancheva S, Stantcheva B (2000) Ethnobotanical inventory of medici- 907
 862 nal plants in Bulgaria. *J Ethnopharmacol* 69(2):165–172
- 863 Ivancheva S, Manolova N, Serkedjieva J, Dimov V, Ivanovska N (1992) 908
 864 Polyphenols from Bulgarian medicinal plants with anti-infectious 909
 865 activity. *Basic Life Sci* 59:717–728
- Ivanova VM (1963) Study of the roots and rhizomes of *Patrinia interme-* 866
 867 *dia* as a source for new medicinal preparations. Synopsis of disserta- 868
 869 tion for the degree of Ph.D. in Pharmacology, Moscow (in Russian)
- Jafarian-Dehkordi A, Emami SA, Saeidi M, Sadeghi H (2004) 870
 871 Cytotoxicologic studies of the extracts of Iranian *Juniperus sabina* 872
 873 and *Platycladus orientalis* on cancer cells. *J Res Med Sci* 9(5):7–11
- Jamshidzadeh A, Niknahad H (2006) Hepatoprotective activity of 874
 875 *Berberis integerrima* Bge extract in rats treated with CCl₄: in vitro 876
 877 and in vivo studies. *Toxicol Lett* 164(Suppl 1):S310
- Jamwal KS, Anand KK, Chopra IC (1962) Pharmacological properties 878
 879 of a crystalline substance (osthol) isolated from *Prangos pabularia*. 880
 881 *Arch Int Pharmacodyn Ther* 138:400–411
- Jassbi AR, Mehrdad M, Soleimani M, Mirzaei M, Sonboli A (2005) 882
 883 Chemical composition of the essential oils of *Bunium elegans* and 884
 885 *Bunium caroides*. *Chem Nat Compd* 41(4):415–417
- Ji L, Xu Z, Pan G, Yang G (1997) GC-MS analysis of constituents of 886
 887 essential oils from stems of *Ephedra sinica* Stapf, *E. intermedia* 888
 889 Schrenk et C.A. Mey. and *E. equisetina* Bge. *Zhongguo Zhong Yao* 890
 891 *Za Zhi* 22(8):489–492 (in Chinese)
- Jo EH, Hong HD, Ahn NC, Jung JW, Yang SR, Park JS, Kim SH, Lee 892
 893 YS, Kang KS (2004) Modulations of the Bcl-2/Bax family were 894
 895 involved in the chemopreventive effects of licorice root (*Glycyrrhiza* 896
 897 *uralensis Fisch*) in MCF-7 human breast cancer cell. *J Agric Food* 898
 899 *Chem* 52(6):1715–1719
- Johansson S, Göransson U, Lujendijk T, Backlund A, Claesson P, 900
 901 Bohlin L (2002) A neutrophil multitarget functional bioassay to 902
 903 detect anti-inflammatory natural products. *J Nat Prod* 65(1):32–41
- Jung JI, Chung E, Seon MR, Shin HK, Kim EJ, Lim SS, Chung WY, 904
 905 Park KK, Park JH (2006a) Isoliquiritigenin (ISL) inhibits ErbB3 906
 907 signaling in prostate cancer cells. *Biofactors* 28(3–4):159–168
- Jung JI, Lim SS, Choi HJ, Cho HJ, Shin HK, Kim EJ, Chung WY, Park 908
 909 KK, Park JH (2006b) Isoliquiritigenin induces apoptosis by depo- 909
 910 larizing mitochondrial membranes in prostate cancer cells. *J Nutr* 910
 911 *Biochem* 17(10):689–696
- Kahkonen MP, Hopia AI, Vuorela HJ, Rauha J, Pihlaja K, Kujala TS, 912
 913 Heinonen M (1999) Antioxidant activity of plant extracts contain- 914
 915 ing phenolic compounds. *J Agric Food Chem* 47(10):3954–3962
- Kanazawa M, Satomi Y, Mizutani Y, Ukimura O, Kawauchi A, Sakai T, 916
 917 Baba M, Okuyama T, Nishino H, Miki T (2003) Isoliquiritigenin 918
 919 inhibits the growth of prostate cancer. *Eur Urol* 43(5):580–586
- Kanokmedhakul K, Kanokmedhakul S, Phatchana R (2005) Biological 920
 921 activity of anthraquinones and triterpenoids from *Prismatomeris* 921
 922 *fragrans*. *J Ethnopharmacol* 100(3):284–288
- Karamenderes C, Khan S, Tekwani BL, Jacob MR, Ikhlas IA (2006) 923
 924 Antiprotazoal and antimicrobial activities of *Centaurea* species 924
 925 growing in Turkey. *Pharm Biol* 44(7):534–539
- Karimov VA, Shomakhmudov A (1993) Khalk tabobati va zamonavii 926
 927 imli tilda kullaniladigan shifobakhsh usimliklar. Avicenna, Toshkent 927
 928 (in Uzbek)
- Karimov A, Telezhenetskaya MV, Lutfullin KL, Yunusov SYu (1975) 928
 929 Alkaloids of *Echium vulgare* and *Berberis oblonga*. *Chem Nat* 929
 930 *Compd* 11(3):452
- Karimov A, Abdullaev ND, Telezhenetskaya MV, Lutfullin K, Yunusov 931
 932 SYu (1977a) Oblongine – a new quaternary alkaloid from *Berberis* 932
 933 *oblonga*. *Chem Nat Compd* 12(1):111–112
- Karimov A, Telezhenetskaya MV, Lutfullin KL, Yunusov SYu (1977b) 934
 935 Berberis alkaloids. The new alkaloid oblongamine. *Chem Nat* 935
 936 *Compd* 13(1):68–70
- Karimov A, Tashkhodzhaev B, Rashkes YV, Makhmudov MK, 937
 938 Mil'grom EG (1993a) *Berberis* alkaloids. XXI. Intebrine – a new 938
 939 N-benzylisoquinoline alkaloid from *Berberis integerrima*. *Chem* 939
 940 *Nat Compd* 29(1):53–57
- Karimov A, Vinogradova VI, Shakirov R (1993b) *Berberis* alkaloids. 940
 941 XXII. Interbrinine and interbrimine – new alkaloids from *Berberis* 941
 942 *integerrima*. *Chem Nat Compd* 29(1):57–60

- 931 Karimova SU, Israilov IA, Yunusov MS, Yunusov SYu (1980) Alkaloids
932 of *Glaucium fimbriigerum*. Chem Nat Compd 16(2):177–180
- 933 Karimova SU, Israilov IA, Vezhnik F, Yunusov MS, Slavik Yu, Yunusov
934 SYu (1983) Alkaloids of *Glaucium fimbriigerum*. II. Chem Nat
935 Compd 19(4):464–467
- 936 Kasimu R, Tanaka K, Tezuka Y, Gong ZN, Li JX, Basnet P, Namba T,
937 Kadota S (1998) Comparative study of seventeen *Salvia* plants:
938 aldose reductase inhibitory activity of water and MeOH extracts and
939 liquid chromatography-mass spectrometry (LC-MS) analysis of
940 water extracts. Chem Pharm Bull 46(3):500–504
- 941 Katsube T, Tabata H, Ohta Y, Yamasaki Y, Anuurad E, Shiwaku K,
942 Yamane Y (2004) Screening for antioxidant activity in edible plant
943 products: comparison of low-density lipoprotein oxidation assay,
944 DPPH radical scavenging assay, and Folin-Ciocalteu assay. J Agric
945 Food Chem 52(8):2391–2396
- 946 Katsube T, Imawaka N, Kawano Y, Yamazaki Y, Shiwaku K, Yamane Y
947 (2006) Antioxidant flavonol glycosides in mulberry (*Morus alba* L.)
948 leaves isolated based on LDL antioxidant activity. Food Chem
949 97(1):25–31
- 950 Kaurichev IC (Senior ed) (1989) Pochvovedeniye, 4 izd (Soil science),
951 4th edn. Agropromizdat, Moscow (in Russian)
- 952 Kawasaki Y, Goda Y, Yoshihira K (1992) The mutagenic constituents of
953 *Rubia tinctorum*. Chem Pharm Bull 40(6):1504–1509
- 954 Keneshov BM, Kuliev ZA, Vdovin AD, Abdullaev ND, Makhmatkulov
955 AB, Nishanov AA (1997a) Proanthocyanidins of *Polygonum cori-*
956 *arium*. III. Structures of proanthocyanidins T1 and T2. Chem Nat
957 Compd 33(4):453–457
- 958 Keneshov BM, Kuliev ZA, Vdovin AD, Abdullaev ND, Nishanov AA
959 (1997b) Proanthocyanidins of *Polygonum coriarium*. IV. Structures
960 of proanthocyanidins T3 and T4. Chem Nat Compd 33(5):
961 548–553
- 962 Kennedy DO, Little W, Scholey AB (2004) Attenuation of laboratory-
963 induced stress in humans after acute administration of *Melissa*
964 *officinalis* (Lemon Balm). Psychosom Med 66:607–613
- 965 Khaidav Ts (1965) Medicinal plants used in Mongolian Folk medicine.
966 In: Tez. Doklad. 41-I Itog, Nauch. Konf. Chenovits. Med. Inst.
967 (Proceedings of the 41st scientific conference, Chernovitsa Medical
968 Institute), pp 41–45 (in Russian)
- 969 Khakimov MKh, Tsukervanik IP (1948) Essential oil of *Achillea*
970 *filipendulina* Lam., vol 1. Trudy Instituta Khimii, Tashkent (in
971 Russian)
- 972 Khalilova AZ, Litvinov IA, Beskrovnyi DV, Gubaidullin AT, Shakurova
973 ER, Nuriev IR, Khalilov LM, Dzhemilev UM (2004) Isolation and
974 crystal structure of taraxasteryl acetate from *Onopordum acan-*
975 *thium*. Chem Nat Compd 40(3):254–257
- 976 Khalmatov KhKh (1964) Dikorastushchiye Lekarstvenniye Rasteniya
977 Uzbekistana (Wild-growing medicinal plants of Uzbekistan).
978 Meditsina, Tashkent (in Russian)
- 979 Khalmatov KhKh, Khabibov ZKh (1976) Kovrak – Ferula vonyuchaya
980 – Ferula assafoetida L. Sp Pl. In: Uzbekistonning shifobakhsh usim-
981 liklar. Meditsina, Tashkent, pp 49–50 (in Uzbek)
- 982 Khalmatov KhKh, Kosimov AI (1992) Russian-Latin-Uzbek dictionary
983 of medicinal plants (herbal plants). Ibn Sino, Tashkent (in Russian)
- 984 Khalmatov KhKh, Kosimov AI (1994) Dorivor Usimliklar
985 (Malumotnoma) [Medicinal plants (references)]. Ibn Sino,
986 Tashkent (in Uzbek)
- 987 Khalmatov KhKh, Kharlamov IA, Alimbayeva PK, Karriev MO,
988 Khaetov IH (1984) Osnovniye Lekarstvenniye Rasteniya Srednei
989 Azii (The main medicinal plants of Central Asia). Meditsina,
990 Tashkent (in Russian)
- 991 Khamdamov I (1972) In: Sultanov MB (ed) Farmakologiya Alkaloidov
992 i Ikh Proizvodnikov (Pharmacology of alkaloids and their derivatives).
993 FAN, Tashkent (in Russian)
- 994 Khamidov II, Tashkhodzhaev B, Aripova SF, Telezhenetskaya MV,
995 Karimov AK (1996) *Berberis* alkaloids. XXXVII. Investigation of
996 the alkaloids of *B. oblongata* and *B. integerrima*. Crystal structure
of 8-trichloromethylidhydroberberine. Chem Nat Compd 32(6):876–879
- Khamidov II, Aripova SF, Karimov AK (2003) *Berberis* alkaloids. XLI.
Alkaloids from leaves of cultivated *Berberis oblonga*. Chem Nat
Compd 39(4):407
- Khamidullina EA, Gromova AS, Lutsky VI, Owen NL (2006) Natural
products from medicinal plants: non-alkaloidal natural constituents
of the *Thalictrum* species. Nat Prod Rep 23(1):117–129
- Khanov M, Kurmukov AG, Sultanov MB, Akhmedkhodzhaeva KhS
(1968) Influence of vincarin on central nervous system. Farmakol
Toksikol 31(5):538–541 (in Russian)
- Khanov M, Sultanov MB, Egorova TA (1971) Pharmacological study of
the alkaloid cononopsine extracted from *Codonopsis clematidea*.
Farmakol. Alkoloidov Serdech Glikoyidov, pp 210–212 (in Russian)
- Khanov M, Kurmukov AG, Akhmedkhodzhaeva KhS, Sultanov MB
(1972a) Effect of vincarine on experimental heart arrhythmias. In:
Sultanov MB (ed) Farmakologiya Alkaloidov i Ikh Proizvodnikov
(Pharmacology of alkaloids and their derivatives). FAN, Tashkent,
pp 22–25 (in Russian)
- Khanov M, Kurmukov AG, Akhmedkhodzhaeva HS (1972b) Influence
of vincarin on experimental cardiac arrhythmia. In: Pharmacology
of alkaloids and their derivatives. FAN, Tashkent (in Russian)
- Kharlamova TV (2007) Reaction of frangula-emodin with α -bromoal-
kylmethylketones. Chem Nat Compd 43(4):391–394
- Khashimov AM, Shakirov R, Yunusov SYu (1970) A study of the alka-
loids of *Veratrum lobelianum*. Chem Nat Compd 6(3):338–340
- Khodzhaeva MA (1994) Carbohydrates of *Allium*. XIII. Glucofructans
of *Allium suvorovii*. Chem Nat Compd 30(2):162–165
- Khodzhaeva MA, Kondratenko ES (1984) Carbohydrates of *Allium*. VI.
Stachyose from the seeds of *Allium suvorovii*. Chem Nat Compd
20(6):745–746
- Khodzhaeva MA, Turakhozhayev MT (1992) Carbohydrates of *Allium* IX.
Polysaccharides of *A. suvorovii*. Chem Nat Compd 28(6):540–541
- Khodzhimatov M (1989) Dikorastushchiye Lekarstvenniye Rasteniya
Tadzhikistana (Wild-growing medicinal plants of Tadzhikistan).
Glavnaya nauchnaya redaktsiya Tadzhikskoi Sovetskoi entsiklope-
dii, Dushanbe (in Russian)
- Kim SY, Gao JJ, Kang HK (2000) Two flavonoids from the leaves of
Morus alba induce differentiation of the human promyelocytic leu-
kemia (HL-60) cell line. Biol Pharm Bull 23(4):451–455
- Kim SR, Park MJ, Lee MK, Sung SH, Park EJ, Kim J (2002) Flavonoids
of *Inula britannica* protect cultured cortical cells from necrotic cell
death induced by glutamate. Free Radic Biol Med 32(7):596–604
- Kim HK, Choi YH, Erkelens C, Lefeber AW, Verpoorte R (2005)
Metabolic fingerprinting of *Ephedra* species using ¹H-NMR spec-
troscopy and principal component analysis. Chem Pharm Bull
53(1):105–109
- Kim MR, Lee YJ, Lee HH, Aryal DK, Kim YG, Kim SK, Woo ER,
Kang KW (2006) Antioxidative effects of quercetin-glycosides iso-
lated from the flower buds of *Tussilago farfara* L. Food Chem
Toxicol 44(8):1299–1307
- Kimura T, Jyo M, Nakamura N, Komatsu K, Hattori M, Shimotohno K,
Shimotohno K, Kakiuchi N (2003) Inhibitory effect of Tibetan medic-
inal plants on viral polymerases. J Tradit Med 20(6):243–250
- Kisiel W, Barszcz B (2000) Further sesquiterpenoids and phenolics
from *Taraxacum officinale*. Fitoterapia 71(3):269–273
- Klejduš B, Vitamvasova-Sterbova D, Kuban V (2001) Identification of
isoflavone conjugates in red clover (*Trifolium pratense*) by liquid
chromatography-mass spectrometry after two-dimensional solid-
phase extraction. Anal Chim Acta 450(1–2):81–97
- Klimentova DA, Aksinenko SG, Gorbacheva AV, Nesterova YuV,
Pashinskiy VG (2005) Antiulcerous properties of *Fragaria vesca*
(Rosaceae) aboveground part infusion. Rastitel'nyye Sursy
41(2):129–133 (in Russian)
- Knox YM, Hayashi K, Suzutani T, Ogasawara M, Yoshida I, Shiina R,
Tsukui A, Terahara N, Azuma M (2001) Activity of anthocyanins

- 1063 from fruit extract of *Ribes nigrum* L. against influenza A and B
1064 viruses. *Acta Virol* 45(4):209–215
- 1065 Knox YM, Suzutani T, Yoshida I, Azuma M (2003) Anti-influenza
1066 virus activity of crude extract of *Ribes nigrum* L. *Phytother Res*
1067 17(2):120–122
- 1068 Kobaisy M, Tellez MR, Dayan FE, Mamonov LK, Mukanova GS,
1069 Sitpaeva GT, Gemejjeva NG (2005) Composition and phytotoxic
1070 activity of *Nepeta pannonica* L essential oil. *J Essent Oil Res*
1071 17(6):704–707
- 1072 Kobayashi A, Hachiya A, Ohuchi A, Kitahara T, Takema Y (2002a)
1073 Inhibitory mechanism of an extract of *Althaea officinalis* L. on endothe-
1074 lin-1-induced melanocyte activation. *Biol Pharm Bull* 25(2):229
- 1075 Kobayashi T, Song QH, Hong T, Kitamura H, Cyong JC (2002b)
1076 Preventative effects of the flowers of *Inula britannica* on autoim-
1077 mune diabetes in C57BL/KsJ mice induced by multiple low doses
1078 of streptozotocin. *Phytother Res* 16(4):377–382
- 1079 Kocsis A, Szabo LF, Podanyi B (1993) New bis-iridoids from *Dipsacus*
1080 *laciniatus*. *J Nat Prod* 56(9):1486–1499
- 1081 Kogure K, Yamauchi I, Tokumura A, Kondou K, Tanaka N, Takaishi Y,
1082 Fukuzawa K (2004) Novel antioxidants isolated from plants of the
1083 genera *Ferula*, *Inula*, *Prangos* and *Rheum* collected in Uzbekistan.
1084 *Phytomedicine* 11(7–8):645–651
- 1085 Kokoska L, Polesny Z, Rada V, Nepovim A, Vanek T (2002) Screening
1086 of some Siberian medicinal plants for antimicrobial activity. *J*
1087 *Ethnopharmacol* 82(1):51–53
- 1088 Kolesnikov MP, Gins VK (2001) Phenolic substances in medicinal
1089 plants. *Appl Biochem Microbiol* 37(4):392–399
- 1090 Komarov VL (Senior ed) (1934) Flora of the USSR, vols 1–30.
1091 *Izdatelstvo Akademii Nauk SSSR, Leningrad* (in Russian)
- 1092 Komissarenko NF, Levashova IG (1988) Phototoxic compounds of the
1093 genus *Dictamnus*. In: *Vtoraya Resp. Konf. Po Med. Bot. – Kiev*
1094 (Proceedings of the 2nd Republican conference on medicinal botany,
1095 Kiev), Nauka, Kiev, pp 258–259 (in Russian)
- 1096 Komissarenko NF, Levashova IG (1989) Flavonoids and coumarins of
1097 some plant from Siberia and the far East. In: *Novuiye Lekarstvennuiye*
1098 *Preparatui iz Rastenii Sibiri i Dalnego Vostoka* (New medical prepa-
1099 *rations from plants of Siberia and the far East*), vol 2, Tomsk, pp
1100 81–82 (in Russian)
- 1101 Konishi T, Shimada Y, Nagao T, Okabe H, Konoshima T (2002)
1102 Antiproliferative sesquiterpene lactones from the roots of *Inula*
1103 *helenium*. *Biol Pharm Bull* 25(10):1370–1372
- 1104 Konovalov DA, Nesterova TA (2003). Phytochemical study of some
1105 Caucasian species of Asteraceae and the possibility of their aller-
1106 genic properties. *Vestnik VGU. Seriya: Khimiya. Biologiya.*
1107 *Farmatsiya* (2):205–207 (in Russian)
- 1108 Konrad L, Müller HH, Lenz C, Laubinger H, Aumüller G, Lichius JJ
1109 (2000) Antiproliferative effect on human prostate cancer cells by a
1110 stinging nettle root (*Urtica dioica*) extract. *Planta Med* 66(1):44–47
- 1111 Koo HN, Hong SH, Song BK, Kim CH, Yoo YH, Kim HM (2004)
1112 *Taraxacum officinale* induces cytotoxicity through TNF- α and
1113 IL-1 α secretion in Hep G2 cells. *Life Sci* 74(9):1149–1157
- 1114 Korolyuk EA, Konig V, Tkachev AV (2002) Compounds in the essen-
1115 tial oil of *Ziziphora clinopodioides* Lam. from the Altai region and
1116 Altai Republic. *Khim Rastitel'nogo Sirya* (1):49–52 (in Russian)
- 1117 Kotenko LD, Yakubova MYa, Tselishcheva NA, Turakhozhayev MT,
1118 Badalbaeva TA (1994) Quantitative determination of the total iridoids
1119 in plants of the genus *Lagochilus*. *Chem Nat Compd* 30(6):669–672
- 1120 Kovalenko VN (1946) About the pharmacological and pharmaco-
1121 logical characteristics of the alkaloid dictamnine from *Dictamnus*
1122 *albus turkestanicus*. *Farmatsiya* 9(5):20
- 1123 Kovaleva NG (1971) Treatment with plants. *Meditsina, Moscow* (in
1124 Russian)
- 1125 Krasnov EA, Demidenko LA (1981) Herbacetin glycosides in plants of
1126 the family Crassulaceae and their biological activity. In: *Materialui*
1127 *4-go Vseros. Sezda Farmatsevtov* (Proceedings of the 4th all-union
1128 conference in pharmacology), Voronezh, USSR, pp 439–440
- Krasnov EA, Saratikov AS, Surov YuP (1979) Semeistva 1129
Tolstyankovuih. Plants of the family Crassulaceae. Tomskii gos- 1130
darstvenii universitet, Tomsk (in Russian) 1131
- Krenn L, Wiedenfeld H, Roeder E (1994) Pyrrolizidine alkaloids from 1132
Lithospermum officinale. *Phytochemistry* 37(1):275–277 1133
- Krillov GV (1972) *Travui Zhizni i Ikh Iskateli* (Herbs for life and their 1134
seekers). Zapad-Sibirskoye Knezhnoye Izdatelstvo, Novosibirsk (in 1135
Russian) 1136
- Kruglikova-Livova RP (1952) About the pharmacology of sophocarpine. 1137
Farmakol Toksikol 15(6):51–55 (in Russian) 1138
- Kuchin NN (1955) About using an infusion of the flowers of *Crataegus* 1139
songarica for the treatment of patients with rheumatism. Synopsis 1140
of dissertation for the degree of Ph.D. in Medical Sciences, Alma- 1141
Ata (in Russian) 1142
- Kudryashev S (1932) Essential oils from wild plants of the central 1143
region of the Gissar mountains. *Parfums de France* 12:98–101 1144
- Kulikov VV (1975) *Lekarstvennuiye Rasteniyi Altaiskogo Kraya, 2-I* 1145
izd (Medicinal plants of the Altai territory), 2nd edn. Altaickoye 1146
khizhnoye isdatelstvo, Barnaul (in Russian) 1147
- Kumarasamy Y, Cox PJ, Jaspars M, Nahar L, Sarker SD (2002) 1148
Screening seeds of Scottish plants for antibacterial activity. *J* 1149
Ethnopharmacol 83(1–2):73–77 1150
- Kumarasamy Y, Byres M, Cox PJ, Jaspars M, Nahar L, Sarker SD 1151
(2007) Screening seeds of some Scottish plants for free radical scav- 1152
enging activity. *Phytother Res* 21(7):615–621 1153
- Kurbatova NV, Muzychkin RA, Mukhitdinov NM, Parshina GN 1154
(2003) Comparative phytochemical investigation of the composi- 1155
tion and content of biologically active substances in *Marrubium* 1156
vulgare and *M. alternidens*. *Chem Nat Compd* 39(5):501–502 1157
- Kurlin VA, Zapesochnaya GG (1986) Terpenoids of the rhizomes of 1158
Rhodiola linearifolia. *Chem Nat Compd* 22(5):607 1159
- Kurmukov AG (1956) Pharmacological investigations of some 1160
Cruciferous plants of Uzbekistan for the presence of cardiac glyco- 1161
sides. In: *Nauchnuiye rabotui studentov Tash. Gos. Med. Inst.* 1162
(Scientific works by students of Tashkent State Medical Institute). 1163
Izdatelstvo Akademii Nauk UzSSR, Tashkent, pp 59–65 (in Russian) 1164
- Kurmukov AG (1967a) Electroencephalogram analysis of influences of 1165
reserpine, vincamine, ervamine and ervamidine on CNS. In: *Kamilov* 1166
IK (ed) *Farmakologiya Alkaloidov i Glikozidov* (Pharmacology of 1167
alkaloids and glycosides). FAN, Tashkent, pp 74–79 (in Russian) 1168
- Kurmukov AG (1967b) On the pharmacology of alkaloid vinervinine. 1169
In: *Kamilov IK* (ed) *Farmakologiya Alkaloidov i Glikozidov* 1170
(Pharmacology of alkaloids and glycosides). FAN, Tashkent, pp 1171
79–82 (in Russian) 1172
- Kurmukov AG (1968a) Alkaloids as stimulants of uterine activity. *Dokl* 1173
Akad Nauk UzSSR 25(12):23–24 (in Russian) 1174
- Kurmukov AG (1968b) On the pharmacology of the alkaloid ervami- 1175
dine (akuamidine). *Farmakol Toksikol* 31(1):47 (in Russian) 1176
- Kurmukov AG (1970) Pharmacological investigations of some indole 1177
alkaloids. Synopsis of dissertation for the degree of doctor of medi- 1178
cine, Tashkent (in Russian) 1179
- Kurmukov AG (1975) The pharmacology of the alkaloid ervine. 1180
Farmakol Toksikol 38(1):82–84 (in Russian) 1181
- Kurmukov AG (1976) Pharmacology of licorice and its preparations 1182
(Influences on the hypothalamo-hypophyseal-adrenal systems). In: 1183
Sultanov MB (ed) *Farmakologiya Rastitelnuikh Veshchestv* 1184
(Pharmacology of plant substances). FAN, Tashkent, pp 174–187 1185
(in Russian) 1186
- Kurmukov AG (1978) The pharmacology of the indole alkaloids 1187
akuamine, aimaline, vinervine and the derivatives ervamine and 1188
pseudocopsinine. In: *Sadritdinov BS* (ed) *Farmakologiya Prirodnikh* 1189
Veshchestv (Pharmacology of natural substances). FAN, Tashkent, 1190
pp 32–51 (in Russian) 1191
- Kurmukov AG (1987) About the anti-inflammatory action of the lac- 1192
tone leucomisine. *Meditsinskii Zh Uzbekistana* (9):72–75 (in 1193
Russian) 1194

- 1195 Kurmukov AG (2000) Hypoxic hypotropia of a fetus and its correction
1196 by introduction of ecdysten. *Pathology* (2):24–27 (in Russian)
- 1197 Kurmukov AG, Akhmedkhodzhaeva KhS (1975) The experimental
1198 influence of psoralen and infekundin on the sexual cycle and
1199 embryogenesis. *Meditinskii Zh Uzbekistana* (8):35 (in Russian)
- 1200 Kurmukov AG, Akhmedkhodzhaeva KhS (1994) Estrogennuiye leka-
1201 rstvennuiye preparatui iz rastenii roda *Ferula* (Estrogenic medicinal
1202 preparations of plants in the Genus *Ferula*). Ibn Sino, Tashkent (in
1203 Russian)
- 1204 Kurmukov AG, Ermishina OA (1991) The effect of ecdysterone on
1205 experimental arrhythmias and changes in the hemodynamics and
1206 myocardial contractility induced by coronary artery occlusion.
1207 *Farmakol Toksikol* 54(1):27–29 (in Russian)
- 1208 Kurmukov AG, Kurmukova NA (1992) Influence of ecdysterone on
1209 prenatal hypertrophy, caused by infringement of placental-uterine
1210 circulation. In: *Eksperimentalnaya farmalogiya klinishi: "Tezisui*
1211 *dokladov"*. Vinnytsia, Kiev, pp 42–43
- 1212 Kurmukov AG, Ermishina OA (1986) Influence of ecdysterone on the
1213 size of the necrotic zone after experimental myocardial infarction.
1214 In: *Otsenka Razmera i Taktika Lecheniya Infarkta Miokarda*
1215 *(Estimation of the size, and methods for treating, myocardial infarc-*
1216 *tion)*, Tomsk, pp 62–63 (in Russian)
- 1217 Kurmukov AG, Rasulova SA (1983) About anti-sclerotic and cardio-
1218 protecting actions of leucomisine. In: *Trudy Tash. Med. Inst. "Ishemicheskaya*
1219 *Blezn Serdtsa"* (Scientific works of Tashkent State Medical Institute
1220 "Ischemic Heart-disease"). Tashkent, pp 55–58 (in Russian)
- 1221 Kurmukov AG, Saidkasymov T (1968) Mechanism of the effect of ervinine
1222 on the central nervous system. *Meditinskii Zh Uzbekistana* 9:6–9
- 1223 Kurmukov AG, Sultanov MB (1965) Pharmacology of *Vinsumine*. In:
1224 *Farmakologiya Alkaloidov (Pharmacology of alkaloids)*. Nauka,
1225 Tashkent, pp 143–157 (in Russian)
- 1226 Kurmukov AG, Sultanov MB (1971) Vinkametrin – a new stimulator of
1227 child delivery. In: Sultanov MB (ed) *Farmakologiya Alkaloidov i*
1228 *Serdechnikh Glikozidov (Pharmacology of alkaloids and cardiac*
1229 *glycosides)*. FAN, Tashkent (in Russian)
- 1230 Kurmukov AG, Saidkasimov TK (1969) About the main mechanism of
1231 action of ervinine. *Meditinskii Zh Uzbekistana* (3):6–9 (in Russian)
- 1232 Kurmukov AG, Syrov VN (1976) About the anabolic activity of the
1233 phytoecdysone ecdysterone isolated from *Rhaponticum cartham-*
1234 *oides*. *Farmakol Toksikol* (6):690–693 (in Russian)
- 1235 Kurmukov AG, Zakirov UB (1992) Alkaloidui i Preparatui Tselebnuih
1236 Trav dlya Lecheniya Gipertenzivnuikh Sostoyanii (Alkaloids and
1237 preparations of medicinal herbs for treating hypertensive condi-
1238 tions). Ibn Sino (Avicenna), Tashkent (in Russian)
- 1239 Kurmukov AG, Akhmedkhodzhaeva KhS, Sirov VN, Sidiyakin VN
1240 (1976) Phytoestrogens from Central Asian plants. *Rastitel'nye*
1241 *Resursy* (4):515–525 (in Russian)
- 1242 Kurmukov AG, Syrov VN, Akhmedkhodzhaeva KhS, Aizikov MI
1243 (1977) Hormonal activity of plant substances. In: *Tezisui Dokladov*
1244 *Vsesoyuznoi Konferentsii. Leningrad-Pushkin, 17–19 Sept 1977, p*
1245 *225* (in Russian)
- 1246 Kurmukov AG, Syrov VN, Ermishina OA (1980) Effects of
1247 Phytoecdysones and sapogenin 6-ketoderivatives on cardiac muscle
1248 metabolism. *J Mol Cell Cardiol* (12):86
- 1249 Kurmukov AG, Aizikov MI, Rasulova SA (1984a) Hypolipidemic and
1250 antisclerotic activities of fruticine in experiments and clinics. In:
1251 *Demiologiya, diagnostika, klinika, techeniye I reabilitatsiya ser-*
1252 *dechno-sosudistuih zabolevaniy (Demology, diagnostics, clinics,*
1253 *progression and rehabilitation of cardiovascular diseases)*. Kanaus,
1254 pp 288–289 (in Russian)
- 1255 Kurmukov AG, Aizikov MI, Rasulova SA (1984b) Hypocoagulative
1256 effects of amorphine for experimental atherosclerosis. *Dokl Akad*
1257 *Nauk UzSSR* (10):49–50 (in Russian)
- 1258 Kurmukov AG, Aizikov MI, Rasulova SA, Makhmudov OKh (1986)
1259 Hypolipidemic, hypocholesterolemic and antisclerotic effects of the
1260 rotenoid glycoside amorphine. *Farmakol Toksikol* 49(6):54–57
- Kurmukov AG, Aizikov MI, Rasulova SA, Sidiyakin GP, Sham'ianov
ID, Malikov VM (1991a) The angioprotector and hypolipidemic
activity of leucomisine on experimental atherosclerosis. *Farmakol*
Toksikol 54(3):35–37 (in Russian)
- Kurmukov AG, Nazrullaev SS, Abdullahadjaeva DG (1991b)
Pharmacological treatment of hypoxic conditions with plant proan-
thocyanidins. In: *Farmakologicheskaya korrektsiya gipoksicheskikh*
sostoyanii. Chast II. (Pharmacological treatment of hypoxic condi-
tions, part 2) Grodno (in Russian)
- Kurmukova NA (2000a) Fetal hypoxic hypotrophy and its correction by
ecdysten introduction. *Patologiya* (2):24–27 (in Russian)
- Kurmukova NA (2000b) Seasonal stability (resistance) of fetus against
uteroplacental hemodynamics disturbance and its correction by
ecdysten introduction. *Chem Nat Compd (special edn):148–150* (in
Russian)
- Kurmukova NA, Kurbanov SD (1999) Evaluation of immune state con-
dition of women with "Inuterine Delayed Embro Development"
during the treatment with ecdysterone. *Pediatrics (special edn):250*
(in Russian)
- Kurmukova ShR, Aizikov MI (1997) Studies of the anti-inflammatory
action of a mixture of leucomisine and ascorbic acid. In: *Trudy Tash*
Med Inst "Farmakoterapiya patalogicheskikh sostoyanii" (Scientific
works of Tashkent State Medical Institute, "Pharmacotherapy of
pathological conditions"), vol 1. Tashkent, pp 65–68 (in Russian)
- Kurmukova ShR, Aizikov MI, Zakirov NU (1997a) Study of the effec-
tivity of leucomisine and leucomisine mixed with ascorbic acid on
experimental izadrinic myocarditis in rats. In: *III Kongress*
Assotsiatsii Kardiologov Stran Tsentralnoi Azii (III congress of
association of cardiologists in Middle Asia), Tashkent, pp. 173 (in
Russian)
- Kurmukova ShR, Zakirov NU, Aizikov MI (1997b) Cardio-protecting
action of leucomisine L-AK on experimental myocarditis. *Kimye va*
farmatsiya (3):37–40 (in Russian)
- Kurmukov AG, Makhmudov O, Aizikov MI (1982) The influence of
fruticine on the content of major compounds for lipid metabolism in
rabbits with experimental atherosclerosis. In: *Materialov yubileinoi*
nauchno-prakticheskoi konferentsii posvyashchonnoi 50 lyetiyu
sanatoriya "Uzbekistan" v gorode Kislovodskye po problemeye
"Etapnoye lecheniye IBS", Tashkent
- Kurochkin EI (1998) *Lekarstvennuiye Rasteniya (Medicinal plants)*.
Parus, Samara (in Russian)
- Kwon DH, Kwon HY, Kim HJ, Chang EJ, Kim MB, Yoon SK, Song
EY, Yoon DY, Lee YH, Choi IS, Choi YK (2005) Inhibition of hepa-
titis B virus by an aqueous extract of *Agrimonia eupatoria* L.
Phytother Res 19(4):355–358
- Ladigina EA, Morozova RS (1987) *Fitoterapiya (Phytotherapy)*.
Meditcina, Leningrad (in Russian)
- Lam ANC, Demasi M, James MJ, Husband AJ, Walker C (2004) Effect
of red clover isoflavones on cox-2 activity in murine and human
monocyte/macrophage cells. *Nutr Cancer* 49(1):89–93
- Larsen E, Kharazmi A, Christensen LP, Christensen SB (2003) An anti-
inflammatory galactolipid from rose hip (*Rosa canina*) that inhibits
chemotaxis of human peripheral blood neutrophils in vitro. *J Nat*
Prod 66(7):994–995
- Lee SJ, Chung HY, Maier CGA, Wood AR, Dixon RA, Mabry TJ
(1998) Estrogenic flavonoids from *Artemisia vulgaris* L. *J Agric*
Food Chem 46(8):3325–3329
- Lee SM, Min BS, Lee CG, Kim KS, Kho YH (2003) Cytotoxic triterpenoids
from the fruits of *Zizyphus jujuba*. *Planta Med* 69(11):1051–1054
- Li LM, Liao X, Peng SL, Ding LS (2005) Chemical constituents from
the seeds of *Zizyphus jujuba* var. *spinosa* (Bunge) Hu. *J Integr Plant*
Biol 47(4):494–498
- Li L, Tsao R, Yang R, Kramer JKG, Hernandez M (2007) Fatty acid
profiles, tocopherol contents, and antioxidant activities of heartnut
(*Juglans ailanthifolia* var. *cordiformis*) and Persian walnut (*Juglans*
regia L.). *J Agric Food Chem* 55(4):1164–1169

- 1327 Liao WT, Beal JL, Wu WN, Duskotch RW (1978) Alkaloids of
1328 *Thalictrum* XXVII. New hypotensive aporphine-benzylisoquinoline
1329 derived dimeric alkaloids from *Thalictrum minus* race B. *Lloydia*
1330 41(3):271–276
- 1331 Lin LZ, He XG, Lindenmaier M, Yang J, Cleary M, Qiu SX, Cordell
1332 GA (2000) LC-ESI-MS study of the flavonoid glycoside malonates
1333 of red clover (*Trifolium pratense*). *J Agric Food Chem*
1334 48(2):354–365
- 1335 Lipkin A, Anisimova V, Nikonorova A, Babakov A, Krause E, Bienert
1336 M, Grishin E, Egorov T (2005) An antimicrobial peptide Ar-AMP
1337 from amaranth (*Amaranthus retroflexus* L.) seeds. *Phytochemistry*
1338 66(20):2426–2431
- 1339 Liu X, Cui Y, Yu Q, Yu B (2005) Triterpenoids from *Sanguisorba*
1340 *officinalis*. *Phytochemistry* 66(14):1671–1679
- 1341 Logendra S, Ribnicki DM, Yang H, Poulev A, Ma J, Kennelly EJ,
1342 Raskin I (2006) Bioassay-guided isolation of aldose reductase
1343 inhibitors from *Artemisia dracunculus*. *Phytochemistry*
1344 67(14):1539–1546
- 1345 Lopez TA, Cid MS, Bianchini ML (1999) Biochemistry of hemlock
1346 (*Conium maculatum* L.) alkaloids and their acute and chronic toxic-
1347 ity in livestock. A review. *Toxicol* 37(6):841–865
- 1348 Lugmanova MR, Muzafarov AM, Mikhailenko OI, Fedorov NI (2007)
1349 Results revealing alkaloid-bearing species in the flora of the moun-
1350 tain-forest-steppe zones in the southern Urals. *Vestnik Bashkirskogo*
1351 *Universiteta* (Bashkirsk University) (1) (in Russian)
- 1352 Lutfullin KL, Tadzhibaev MM, Abdullaev UA, Malikov VM, Yunusov
1353 SYu (1977) Alkaloids of *Reseda luteola*. *Chem Nat Compd*
1354 12(5):559–563
- 1355 Lutfullin KL, Tadzhibaev MM, Malikov VM, Abdullaev UA,
1356 Rakhmankulov U (1976) Dynamics of the accumulation of the alka-
1357 loids of *Reseda luteola* structure of lutine and lutinine. *Chem Nat*
1358 *Compd* 13(6):696–699
- 1359 Ma Yu-H, Yang J-R (2006) Protective effect of stachydrine on myocar-
1360 dial ischemia-reperfusion injury in rats. *Zhongguo Shiyan Fangjixue*
1361 *Zazhi* 12(5):40–42 (in Chinese)
- 1362 Ma SC, Du J, But PPH, Deng XL, Zhang YW, Ooi VEC, Xu HX, Lee
1363 SHS, Lee SF (2002a) Antiviral Chinese medicinal herbs against
1364 respiratory syncytial virus. *J Ethnopharmacol* 79(2):205–211
- 1365 Ma ZJ, Li X, Li N, Wang JH (2002b) Stilbenes from *Sphaerophysa*
1366 *salsula*. *Fitoterapia* 73(4):313–315
- 1367 Ma ZJ, Tian J, Fujii I, Ebizuka Y, Li X (2003) A new lignan from the
1368 seeds of *Sphaerophysa salsula*. *Nat Prod Res* 17(6):423–425
- 1369 Ma ZJ, Fujii I, Ebizuka Y, Li X (2004a) A new lignan from the seeds of
1370 *Sphaerophysa salsula*. *J Asian Nat Prod Res* 6(4):265–269
- 1371 Ma ZJ, Fujii I, Ebizuka Y, Li X, Shimomaki S, Sakano Y (2004b)
1372 Flavonoids from the seeds of *Sphaerophysa salsula*. *J Asian Nat*
1373 *Prod Res* 6(1):69–73
- 1374 Makhlayuk VP (1967) *Lekarstvennuiye Rasteniya v Narodnoi*
1375 *Meditisinye*, izd. 2 (Medicinal plants in folk medicine), 2nd edn.
1376 *Privol'skoye Knezhnoye Izdatel'stvo*, Saratov (in Russian)
- 1377 Makhlayuk VP (1992) *Lekarstvennuiye Rasteniya v Narodnoi*
1378 *Meditisinye* (Medicinal plants in folk medicine). *Niva Rossii*,
1379 Moscow (in Russian)
- 1380 Makhmatkulov AB, Kuliev ZA, Vdovin AD, Yagudaev MR, Malikov
1381 VM (1992) Proanthocyanidins of *Polygonum coriarium*. I. *Chem*
1382 *Nat Compd* 28(1):48–54
- 1383 Makhmatkulov AB, Kuliev ZA, Vdovin AD, Malikov VM (1994)
1384 Proanthocyanidins of *Polygonum coriarium*. II. *Chem Nat Compd*
1385 30(2):214–222
- 1386 Makhmudov MM (2001) Country pasture/forage resource profiles:
1387 Uzbekistan. [http://www.fao.org/ag/AGP/AGPC/doc/Counprof/](http://www.fao.org/ag/AGP/AGPC/doc/Counprof/Uzbekistan/uzbekistan.htm)
1388 [Uzbekistan/uzbekistan.htm](http://www.fao.org/ag/AGP/AGPC/doc/Counprof/Uzbekistan/uzbekistan.htm). Accessed 12 Jan 2008
- 1389 Maksudov NKH (1964) Poluchenie Efirnykh Masel (Terpenov) i Ikh
1390 Primenenie pri Mochekamennoi Bolezni [Production of essential
1391 oils (Terpenes) and their use for the treatment of Urolithic diseases].
1392 *Meditisina*, Tashkent (in Russian)
- Mallabaev A, Sidiyakin GP (1976) Artemidinol – a new isocoumarin 1393
from *Artemisia dracunculus*. *Chem Nat Compd* 12(6):729 1394
- Mallabaev A, Saitbaeva IM, Sidiyakin GP (1971) The isocoumarin 1395
artemidinal from *Artemisia dracunculus*. *Chem Nat Compd* 1396
7(3):248–249 1397
- Mallabaev A, Yagudaev MR, Saitbaeva IM, Sidiyakin GP (1970) The 1398
isocoumarin artemidin from *Artemisia dracunculus*. *Chem Nat* 1399
Compd 6(4):479 1400
- Mamatkhanov AU, Yakubova MR, Syrov VN (1998) Isolation of 1401
turkesterone from the epigeal part of *Ajuga turkestanica* and its ana- 1402
bolic activity. *Chem Nat Compd* 34(2):150–154 1403
- Mamedov N (2005) Adaptogenic, geriatric, stimulant and antidepres- 1404
sant plants of Russian Far East. *J Cell Mol Biol* 4:71–75 1405
- Mamedov N, Craker LE (2001) Medicinal plants used for the treatment 1406
of bronchial asthma in Russia and Central Asia. *J Herbs Spices Med* 1407
Plant 8(2/3):91–118 1408
- Mamedov N, Gardner Z, Craker LE (2004) Medicinal plants used in 1409
Russia and Central Asia for the treatment of selected skin condi- 1410
tions. *J Herbs Spices Med Plant* 11(1/2):191–222 1411
- Mamitov AM (1965) Classification, vertical belts and provinces of 1412
Kyrgyz soils. In: Kovda VA, Lobova EV (eds) *Geography and* 1413
classification of Asian soils. Nauka, Moscow, pp 50–51 (in 1414
Russian) 1415
- Mandal SC, Dhara AK, Ashok Kumar CK, Maiti BC (2001) 1416
Neuropharmacological activity of *Xanthium strumarium* Linn 1417
extract. *J Herbs Spices Med plant* 8(1):69–77 1418
- Mansurov MM (1967) Effect of *Ferula asafetida* on blood coagulabil- 1419
ity. *Meditinskii Zh Uzbekistana* (6):46–49 (in Russian) 1420
- Marco JM, Sanz-Cervera JF, Corral J, Carda M, Jakupovic J (1993) 1421
Xanthanolides from *Xanthium*: absolute configuration of xanthanol, 1422
isoxanthanol and their C-4 epimers. *Plant Physiol Biochem* 1423
34(6):1569–1576 1424
- Martino E, Ramaola I, Urbano M, Bracco F, Collina S (2006) 1425
Microwave-assisted extraction of coumarin and related compounds 1426
from *Melilotus officinalis* (L.) Pallas as an alternative to Soxhlet and 1427
ultrasound-assisted extraction. *J Chromatogr A* 1125(2):147–151 1428
- Mashkovskii MD (1953) The pharmacology of tropacin. *Farmakol* 1429
Toksikol (5):3–8 (in Russian) 1430
- Mashkovskii MD (1984) *Lekarstvennuiye Sredstva: Posobiye po* 1431
Farmakoterapii Dlya Vrachei, 9 izd (Medicinal remedies: manual 1432
on pharmacotherapy for doctors), 9th edn. *Meditisina*, Moscow (in 1433
Russian) 1434
- Massagetov PC (1932) Tian Shan expedition of NIKhFI. *Khim.* 1435
Farmatsev, Promishlennost (4–5): .134–c.142 (in Russian) 1436
- Matsuhisa M, Shikishima Y, Takaishi Y, Honda G, Ito M, Takeda Y,
1437 Shibata H, Higuti T, Kodzhimatov OK, Ashurmetov O (2002) 1438
Benzoylphloroglucinol derivatives from *Hypericum scabrum*. *J Nat* 1439
Prod 65(3):290–294 1440
- Maznev N (2004) *Bolshaya Entsiklopediya Narodnoi Meditsinui* (Big 1441
encyclopedia of traditional medicines). *Martin*, Moscow (in 1442
Russian) 1443
- McCutcheon AR, Roberts TE, Gibbons E, Ellis SM, Babiuk LA, 1444
Hancock RE, Towers GH (1995) Antiviral screening of British 1445
Columbian medicinal plants. *J Ethnopharmacol* 49(2):101–110 1446
- MedicineLib.ru. Medicinal plants. Ochnui tsvet pashennui. [http://medi-](http://medicinelib.ru/travi/O/travi158.html) 1447
[cinelib.ru/travi/O/travi158.html](http://medicinelib.ru/travi/O/travi158.html). Accessed 12 Jan 2008 (in 1448
Russian) 1449
- Mikanagi Y, Yokoi M, Ueda Y, Saito N (1995) Flower flavonol and 1450
anthocyanin distribution in subgenus *Rosa*. *Biochem Syst Ecol* 1451
23(2):183–200 1452
- Milkowska-Leyck K, Borkowski B, Rozanski M (1999) Selected poly- 1453
phenolic compounds Part III. Biological properties of iridoids. 1454
Herba Pol 45(3):232–241 1455
- Mimaki Y, Kuroda M, Fukasawa T, Sashida Y (1999) Steroidal saponins 1456
from the bulbs of *Allium karataviense*. *Chem Pharm Bull* 1457
47:738–743 1458

- 1459 Mimaki Y, Fukushima M, Yokosuka A, Sashida Y, Furuya S, Sakagami
1460 H (2001) Triterpene glycosides from the roots of *Sanguisorba*
1461 *officinalis*. *Phytochemistry* 57(5):773–779
- 1462 Minayeva VG (1991) Lekarstvennuiye Rasteniya Sibiri, 5 izd (Medicinal
1463 plants of Siberia), 5th edn. Nauka, Novosibirsk (in Russian)
- 1464 Ministry of Environmental Protection (1998) Kyrgyz Republic biodi-
1465 versity strategy and action plan. Republic of Kyrgyzstan Ministry of
1466 Environmental Protection, Bishkek
- 1467 Mircheva J, Stoychkov J (1976) Effect of thaliblastine on trans-plant-
1468 able tumors in mice. *Biomed Express* 25:280–281
- 1469 Moallem SA, Niapour M (2008) Study of embryotoxicity of *Perovskia*
1470 *abrotanoides*, an adulterant in folk-medicine, during organogenesis
1471 in mice. *J Ethnopharmacol* 117(1):108–114
- 1472 Mohammadreza VR (2008) Essential oil composition and biological
1473 activity of *Ziziphora clinopodioides* Lam. from Iran. *Res J Pharmacol*
1474 2(2):17–19
- 1475 Momin RA, De Witt DL, Nair MG (2003) Inhibition of cyclooxygenase
1476 (COX) enzymes by compounds from *Daucus carota* L. seeds.
1477 *Phytother Res* 17(8):976–979
- 1478 Morita H, Yun YS, Takeya K, Itokawa H, Shirota O (1997a) Thionation
1479 of segetalins A and B, cyclic peptides with estrogen-like activity
1480 from seeds of *Vaccaria segetalis*. *Bioorg Med Chem* 5(3):631–636
- 1481 Morita H, Yun YS, Takeya K, Itokawa H, Yamada K, Shirota O (1997b)
1482 Vaccaroid A, a new triterpenoid saponin with contractility of rat uter-
1483 ine from *Vaccaria segetalis*. *Bioorg Med Chem Lett* 7(8):1095–1096
- 1484 Mukhamed'yarova MM, Chumbalov TK (1979) Flavonoids of *Rumex*
1485 *confertus*. *Chem Nat Compd* 15(6):758
- 1486 Mukhamedziev MM, Alimbaeva PK (1969) Triterpene glycosides of
1487 *Dipsacus azureus*. *Chem Nat Compd* 5(5):376–377
- 1488 Muminova BA, Batirov EK, Yuldashev MP, Inamova ZG (2006)
1489 Flavonoids from *Sophora griffithii* and *Goebelia pachycarpa*. *Chem*
1490 *Nat Compd* 42(1):108–109
- 1491 Murai M, Tamayama Y, Nishibe S (1995) Phenylethanoids in the herb
1492 of *Plantago lanceolata* and inhibitory effect on arachidonic acid-
1493 induced mouse ear edema. *Planta Med* 61(5):479–480
- 1494 Muravyova DA (1978) Farmakognoziya (Pharmacognosy). *Meditcina*,
1495 Moscow (in Russian)
- 1496 Murillo-Alvarez JI, Encarnación DR, Franzblau SG (2001)
1497 Antimicrobial and cytotoxic activity of some medicinal plants from
1498 Baja California Sur (Mexico). *Pharm Biol* 39(6):445–449
- 1499 Muzychkina RA (2000) Certain Kazakhstan *Galium* species. I. *Chem*
1500 *Nat Compd* 36(3):321–322
- 1501 Nachman RJ, Olsen JD (1983) Ranunculin: a toxic constituent of the
1502 poisonous range plant bur buttercup (*Ceratocephalus testiculatus*).
1503 *J Agric Food Chem* 31(6):1358–1360
- 1504 Nahrstedt A, Butterweck V (1997) Biologically active and other chemi-
1505 cal constituents of the herb of *Hypericum perforatum* L.
1506 *Pharmacopsychiatry* 30:129–134
- 1507 Naidovich LP, Trumneva EA, Tolkachev ON, Vasileva VD (1976)
1508 Chemical compounds of local species in the family Berberidaceae.
1509 *Farmatsiia* 25(4):33–38 (in Russian)
- 1510 Nakanishi T, Inada A, Kambayashi K, Yoneda K (1985) Flavonoid gly-
1511 cosides of the roots of *Glycyrrhiza uralensis*. *Phytochemistry*
1512 24(2):339–341
- 1513 Nan JX, Park EJ, Kim HJ, Ko G, Sohn DH (2000) Antifibrotic effects
1514 of the methanol extract of *Polygonum aviculare* in fibrotic rats
1515 induced by bile duct ligation and scission. *Biol Pharm Bull*
1516 23(2):240–243
- 1517 Nanaeva MT (1960) About the pharmacology of *Artemisia viridis*. In:
1518 2-ya Konf. Fiziologov, Biokhimikov i farmakologov Srednei Azii i
1519 Kazakhstana (Proceedings of the 2nd conference of physiologists,
1520 biochemists and pharmacologists of Central Asia and Kazakhstan),
1521 Frunze, pp 239–240 (in Russian)
- 1522 Nasrullaev AS, Yunusov MS, Matveev VM, Sabirov SS (1989)
1523 14-Acetylnudicaulidine and 18-deoxyglycoctonine – new diterpene
1524 alkaloids from *Delphinium confusum*. *Chem Nat Compd* 25(1):41–42
- Nasirov SH (1973) The pharmacology of the alkaloid methyl ether of
1525 aphyllinic acid. In: *Farmakologiya rastitel'nykh Veshestv Nauchnuiye*
1526 *Trudy, Vuipusk 457* [Pharmacology of plant substances (Scientific
1527 works, edn 457)]. FAN, Tashkent, pp 60–63 (in Russian)
- Nasrullaev FD, Makhsudova BT (1991) Flavonoids of *Lagochilus*
1528 *platycalyx*. *Chem Nat Compd* 27(4):511–512
- National Biodiversity Strategy Project Steering Committee (1998)
1529 Republic of Uzbekistan biodiversity conservation: national strategy
1530 and action plan, Tashkent. http://bbsp-neca.brim.ac.cn/books/act-pln_uzbek/index.html. Accessed online 22 Feb 2007
- Nazrullaev CC, Akhmerov RN, Kurmukov AG (1990) Effects of an
1531 antihypoxant-catechin on the energy of myocardial metabolism.
1532 *Meditinskii Zh Uzbekistana* 11:7–9 (in Russian)
- Neef H, Declercq P, Laekeman G (1995) Hypoglycaemic activity of
1533 selected European plants. *Phytother Res* 9(1):45–48
- Newton SM, Lau C, Wright CW (2000) A review of antimycobacterial
1534 natural products. *Phytother Res* 14(5):303–322
- Nikitina EV (1962) Flora i Rastitelnost Pastbishch i Senokosov Khrebt
1535 Kirgizskii Ala-Too (The flora and vegetation of pastures and ranges
1536 in the Kyrgyz Ala-Too). Izdatel'stvo Akademii Nauk KyrgSSR,
1537 Frunze (in Russian)
- Nishanov AA, Sultankhodzhaev MN, Yunosov MS, Yusupova IM,
1538 Tashkhozhaev B (1991) Alkaloids of *Aconitum talassicum*. *Chem*
1539 *Nat Compd* 27(1):82–86
- Nogaller AM, Nikolaeva VG, Alexeyev VF (1987) Fitoterapiya zabol-
1540 evanii pishchevareniya (Phytotherapy of digestive system diseases).
1541 Pavlov Medical Institute, Ryazan (in Russian)
- Norn S, Kruse PR (2004) Cardiac glycosides: from ancient history
1542 through Withering's foxglove to endogenous cardiac glycosides.
1543 *Dansk Medicinhistorisk Arbog*:119–132 (in Danish)
- Norouzi-Arasi H, Yavari I, Chalabian F, Kiarostami V, Ghaffarzadeh F,
1544 Nasirian A (2006) Chemical constituents and antimicrobial activi-
1545 ties of the essential oil of *Acroptilon repens* (L.) DC. *Flavour Frag J*
1546 21(2):247–249
- Nosal MA, Nosal IM (1959) Lekarstvennuiye Rasteniya i Sposobui ikh
1547 Primniya v Narodye (Medicinal plants and the ways that they are
1548 used by people). *Meditcina, Kiev* (in Russian)
- Nuraliev Yu (1989) Lekarstvennuiye Rasteniya, 2-ye izd (Medicinal
1549 plants), 2nd edn. MAORIF, Dushanbe (in Russian)
- Nuraliev YuN, Zubaidova TM (1994) Hypocholesterinemic and hypo-
1550 lipidemic effects of ether oil of *Origanum tyttanthum* Gontsch.
1551 *Khim Farm Zh* 28(10):63–64 (in Russian)
- Nurmukhamedova MR, Kasymov ShZ, Abdullaev ND, Sidiyakin GP,
1552 Yagudayev MR (1985) Diterpenoids of *Pulicaria salviifolia* I.
1553 Structures of salvin and salvinin. *Chem Nat Compd* 21(2):188–191
- Nurmukhamedova MR, Abdullaev ND, Sidiyakin GP (1986)
1554 Diterpenoids of *Pulicaria salviifolia*. II. Structure of salvicin. *Chem*
1555 *Nat Compd* 22(3):277–279
- Ogolevitz GS (1951) Entsiklopedicheskii Slovar Lekarstvennuikh,
1556 Efiromaslichnuikh i Yadovituikh Rastenii (Encyclopedic dictionary
1557 of medicinal, essential oil and poisonous plants). Gos Izdatel'stvo
1558 selskokhozyaistvennoi literatury, Moscow (in Russian)
- Oh H, Kim DH, Cho JH, Kim YC (2004) Hepatoprotective and free radi-
1559 cal scavenging activities of phenolic petrosins and flavonoids iso-
1560 lated from *Equisetum arvense*. *J Ethnopharmacol* 95(2–3):421–424
- Okasaka M, Takaishi Y, Kashiwada Y, Kodzhimatov OK, Ashurmetov O,
1561 Lin AJ, Consentino LM, Lee KH (2006) Terpenoids from *Juniperus*
1562 *polycarpus* var. *seravschanica*. *Phytochemistry* 67(24):2635–2640
- Omurkamzinova VB, Maurel' ND, Bikbulatova TN (1991) Flavonoids
1563 of *Biebersteinia multifida*. *Chem Nat Compd* 27(5):636–637
- Oomah BD, Ladet S, Godfrey DV, Liang J, Girard B (2000)
1564 Characteristics of raspberry (*Rubus idaeus* L.) seed oil. *Food Chem*
1565 69(2):187–193
- Oprean L, Mărculescu A, Cristea V, Gaspar E (2007) The antibacterial
1566 activity of the volatile oils extracted from medicinal plants. *Acta*
1567 *Aliment Hung* 36(3):303–309

- 1591 Orhan DD, Aslan M, Aktay G, Ergun E, Yesilada E, Ergun F (2003)
1592 Evaluation of hepatoprotective effect of *Gentiana olivieri* herbs on
1593 subacute administration and isolation of active principle. *Life Sci*
1594 72(20):2273–2283
- 1595 Orhan I, Sener B, Choudhary MI, Khalid A (2004) Acetylcholinesterase
1596 and butyrylcholinesterase inhibitory activity of some Turkish
1597 medicinal plants. *J Ethnopharmacol* 91(1):57–60
- 1598 Ortiz JG, Nieves-Natal J, Chavez P (1999) Effects of *Valeriana*
1599 *officinalis* extracts on [3h]flunitrazepam binding, synaptosomal [³H]
1600 GABA uptake, and hippocampal [³H]GABA release. *Neurochem*
1601 *Res* 24(11):1373–1378
- 1602 Ostad SN, Montazeran H, Minaee B, Monsef HR (2003) The study of
1603 acute toxicity of *Biebersteinia multifida* total extract in mice. *Toxicol*
1604 *Lett* 144(Suppl 1):s63
- 1605 Oszmianski J, Wojdylo A, Lamer-Zarawska E, Swiader K (2007)
1606 Antioxidant tannins from Rosaceae plant roots. *Food Chem*
1607 100(2):579–583
- 1608 Overk CR, Yao P, Chadwick LR, Nikolic D, Sun Y, Cuendet MA, Deng
1609 Y, Hedayat AS, Pauli GF, Farnsworth NR, van Breemen RB, Bolton
1610 JL (2005) Comparison of the in vitro estrogenic activities of com-
1611 pounds from hops (*Humulus lupulus*) and red clover (*Trifolium*
1612 *pratense*). *J Agric Food Chem* 53(16):6246–6253
- 1613 Ozek G, Ozek T, Iscan G, Baser KHC, Hamzaoglu E, Duran A (2007)
1614 Comparison of hydrodistillation and microdistillation methods for
1615 the analysis of fruit volatiles of *Prangos pabularia* Lindl., and eval-
1616 uation of its antimicrobial activity. *S Afr J Bot* 73(4):563–569
- 1617 Pages N, Fournier G, Baduel C, Tur N, Rusnac M (1996) Sabinyl acetate,
1618 the main component of *Juniperus sabina* L'Hérit. essential oil, is
1619 responsible for antiimplantation effect. *Phytother Res* 10(5):438–440
- 1620 Pak VV (2005) Medicine plants of folk medicine used for treatment of
1621 gastro-intestinal problems in Fergana valley. (*Food Tech*), Korean
1622 Food Reseach Institute 18(4):150–157
- 1623 Pakudina ZP, Sadykov AS, Zuparov A (1970) Kaempferol glycosides
1624 from the flowers of *Alcea nudiflora*. *Chem Nat Compd*
1625 6(5):643–644
- 1626 Panter KE, Keeler RF, Buck WB (1985a) Induction of cleft palate in
1627 newborn pigs by maternal ingestion of poison hemlock. *Am J Vet*
1628 *Res* 46(6):1368–1371
- 1629 Panter KE, Keeler RF, Buck WB (1985b) Congenital skeletal malfor-
1630 mations induced by maternal ingestion of *Conium maculatum* (poi-
1631 son hemlock) in newborn pigs. *Am J Vet Res* 46(10): 2064–2066
- 1632 Park CJ, Park CB, Hong SS, Lee HS, Lee SY, Kim SC (2000)
1633 Characterization and cDNA cloning of two glycine- and histidine-
1634 rich antimicrobial peptides from the roots of shepherd's purse,
1635 *Capsella bursa-pastoris*. *Plant Mol Biol* 44(2):187–197
- 1636 Pastura G, Mesiti M, Saitta M, Romeo D, Settineri N, Maisano R, Petix
1637 M, Giudice A (1999) Lymphedema of the upper extremity in patients
1638 operated for carcinoma of the breast: clinical experience with cou-
1639 marinic extract from *Melilotus officinalis*. *La Clinica Terapeutica*
1640 150(6):403–408 (in Italian)
- 1641 Pavlov NV (1947) Rastitelnoye Suiryo Kazakhstana (Rasteniya; ikh
1642 Veshchestva i Ispolzovaniye) [Plant raw material of Kazakhstan
1643 (Plants: their substances and uses)]. Izdatelstvo Akademii Nauk
1644 SSSR, Moscow (in Russian)
- 1645 Peng JN, Ma CY, Ge YC (1994) Chemical constituents of *Rhodiola*
1646 *kirilowii* (Regel) Regel. *Zhongguo Zhong Yao Za Zhi* 19(11):676–
1647 677 (in Chinese)
- 1648 Peng WH, Hsieh MT, Lee YS, Lin YC, Liao J (2000) Anxiolytic effect
1649 of seed of *Ziziphus jujuba* in mouse models of anxiety. *J*
1650 *Ethnopharmacol* 72(3):435–441
- 1651 Petrova VP (1972) Catechins and leucoanthocyanins from the fruits of
1652 wild hawthorn species (*Crataegus* L.). *Ukr Botanichnii Zh*
1653 29(2):144–147 (in Ukrainian)
- 1654 Pirniyazov AZ, Abdulladzhanova NG, Mavlyanov SM, Kamaev FG,
1655 Dalimov DN (2003) Polyphenols from *Vitis vinifera* seeds. *Chem*
1656 *Nat Compd* 39(4):349–355
- Plant Resources of the USSR (1985) Fam. Magnoliaceae – Limoniaceae, 1657
vol 1. Nauka, Leningrad (in Russian) 1658
- Plant Resources of the USSR (1986) Fam. Paeoniaceae – Thymelaeaceae, 1659
vol 2. Nauka, Leningrad (in Russian) 1660
- Plant Resources of the USSR (1987) Fam. Hydrangeaceae – 1661
Haloragaceae, vol 3. Nauka, Leningrad (in Russian) 1662
- Plant Resources of the USSR (1988) Fam. Rutaceae – Elaeagnaceae, 1663
vol 4. Nauka, Leningrad (in Russian) 1664
- Plant Resources of the USSR (1990) Fam. Caprifoliaceae – 1665
Plantaginaceae, vol 5. Nauka, Leningrad (in Russian) 1666
- Plant Resources of the USSR (1991) Fam Hippuridaceae – Lobeliaceae, 1667
vol 6. Nauka, Leningrad (in Russian) 1668
- Plant Resources of the USSR (1993) Fam. Asteraceae, vol 7. Nauka, 1669
Saint Petersburg (in Russian) 1670
- Plekhanova NV, Turdumambetov KT, Fedorchenko GP (1985)
1671 Lechebnui Sakhar iz Sornuikh Rastanii Kirgizii (Medicinal sugar
1672 from weedy plants of Kyrgyzstan). Ilim, Frunze 1673
- Plesca-Manea L, Pärvu AE, Pärvu M, Taămaş M, Buia R, Puia M 1674
(2002) Effects of *Melilotus officinalis* on acute inflammation. 1675
Phytother Res 16(4):316–319 1676
- Polievtev NP (1962a) The synergism of an alkaloid, dubinidine, with 1677
hypnotics and narcotics. *Dokl Akad Nauk UzSSR* 19(5):50–54 (in 1678
Russian) 1679
- Polievtev NP (1962b) About some analgesic properties of the alka- 1680
loid dubinidine. In: *Farmakologiya Alkaloidov* (Pharmacology of 1681
alkaloids), vol 1. Izdatelstvo Akademii Nauk UzSSR, Tashkent, 1682
pp 138–144 (in Russian) 1683
- Polievtev NP (1965) The influence of dubinidine on the central ner- 1684
vous system. In: *Farmakologiya Alkaloidov* (Pharmacology of alka- 1685
loids). Nauka, Tashkent, pp 60–80 (in Russian) 1686
- Polievtev NP, Sultanov MB (1971) Pharmacology of a saponin 1687
extracted from the plant *Acanthophyllus gypsophiloides*. In: 1688
Sultanov MB (ed) *Farmakologiya Alkaloidov i Serdechnuikh* 1689
Glikozidov (Pharmacology of alkaloids and cardiac glycosides). 1690
FAN, Tashkent, pp 240–243 (in Russian) 1691
- Polievtev NP, Kamilov IK, Evdokimova NI (1967) Pharmacology of 1692
the alkaloid skimmianine. In: Kamilov IK (ed) *Farmakologiya* 1693
Alkaloidov i Glikozidov (Pharmacology of alkaloids and glyco- 1694
sides). FAN, Tashkent, pp 35–42 (in Russian) 1695
- Poludenny LV, Zhuravlev YuP (2000) Zagotovka, Vuirashchivanie I 1696
pererabotka lekarstvennuikh rastanii (Preparation, management and 1697
processing of medicinal plants). Moscow Agricultural Academy 1698
(MCXA), Moscow (in Russian) 1699
- Pratov U (ed) (1998) Red data book of the Republic of Uzbekistan: rare 1700
and endangered species of plants and animals, vol 1, Plants. Chinor 1701
ENK, Tashkent (in Russian) 1702
- Prokhorova IR, Kurmukov AG (1997) Pre-clinical research of angio- 1703
protecting and hypolipidemic activity of the phytoestrogenic prepa- 1704
ration *Tefestrol*. In: III Kongress Kardiologov Stran Tsentralnoi Azii 1705
(3rd congress of cardiologists of Central Asian countries), Tashkent, 1706
p 175 (in Russian) 1707
- Prokhorova IR, Aizikov MI, Kurmukov AG (1992a) Antisclerotic trend 1708
of lactone A. In: *Eksperimentalnaya Farmakologiya Kliniky* (Tezisui 1709
Dokladov Simposiuma-soveshchaniya s Mezhdunarodnuim uchasti- 1710
yem Vinnitskogo Med Inst) [Clinical pharmacology experiments. 1711
(Proceedings of international symposium, Vinnitsa Medical Institute)] 1712
Vinnitsa Medical Institute, Vinnitsa-Kiev, pp 70–71 (in Russian) 1713
- Prokhorova IR, Aizikov MI, Kurmukov AG (1992b) Hypolipidemic 1714
activity of ferutinin. *Meditsinskii Zh Uzbekistana* (6):63 (in Russian) 1715
- Prokhorova IR, Aizikov MI, Kurmukov AG (1993) Angioprotective, 1716
hypolipidemic and antiatherosclerotic activity of austriacine. In: 1717
Tezisui Dokladov: 1-aya Konferentsiya Kardiologov Moldovui 1718
(Proceedings of the 1st conference of Moldavian cardiologists), 1719
Kishinev, pp 150–151 1720
- Pulatova TP (1969) Presence of alkaloids in some plants of the family 1721
Labiatae. *Chem Nat Compd* 5(1):55 1722

- 1723 Putieva ZhM, Mukhamedziev MM (1998) Triterpene glycosides of
1724 *Dipsacus azureus*. Chem Nat Compd 34(3):341–342
- 1725 Putieva ZhM, Mzhel'skaya LG, Kondratenko ES, Abubakirov NK
1726 (1970) Triterpenoid glycosides of *Acanthophyllum gypsophiloides*.
1727 Chem Nat Compd 6(4):506–507
- 1728 Putieva ZhM, Mzhel'skaya LG, Gorovits TT, Kondratenko ES,
1729 Abubakirov NK (1975) Triterpenoid glycosides of *Acanthophyllum*
1730 *gypsophiloides*. IV. Structure of acanthophyllosides B and C. Chem
1731 Nat Compd 11(6):756–761
- 1732 Putieva ZhM, Gorovits TT, Kondratenko ES, Abubakirov NK (1979)
1733 Triterpene glycosides of *Acanthophyllum gypsophiloides*. VI.
1734 Structure of acanthophylloside D. Chem Nat Compd 15(2):148–151
- 1735 Puupponen-Pimia R, Nohynek L, Meier C, Kahkonen M, Heinonen M,
1736 Hopia A, Oksman-Caldentey KM (2001) Antimicrobial properties
1737 of phenolic compounds from berries. J Appl Microbiol
1738 90(4):494–507
- 1739 Rabinovich MI, Leskov AI, Gladkikh AS (1966) In: Mater Konf Fiziol,
1740 Biokhim, Farmakol. Uchastiem Prakt Vrachei, Ufa, pp 181–184 (in
1741 Russian)
- 1742 Rakhimov DA (1997) Polysaccharides of *Eremurus*. XXX. Method of
1743 isolating a polysaccharide and its pharmacological activity. Chem
1744 Nat Compd 33(2):207
- 1745 Rakhimov KD, Vermenichev SM, Lutskii VI, Gromova AS, Ganenko
1746 TV, Semenov AA (1987) Triterpene alkaloids of *Thalictrum foet-*
1747 *idum* L. and *Thalictrum minus* L. and their antitumorigenic activity.
1748 Pharm Chem J 21(12):848–850
- 1749 Rakhimov DA, Salimova ESh, Zakirova RP (2003) Polysaccharides of
1750 plant tissue culture. III. Polysaccharides of *Codonopsis clematidea*
1751 callus culture. Chem Nat Compd 39(1):46–47
- 1752 Rakhimova IV, Pulatova TP (1972) Study of the antimicrobial proper-
1753 ties of some plants in the family Lamiaceae. In: Phytocids: results,
1754 prospects and research problems, Kiev, pp 94–96 (in Russian)
- 1755 Rakhmanberdyeva RK, Talipova M, Gazizov F, Rakhimov DA (2002)
1756 Carbohydrates and lipids of *Gleditsia triacanthos* seeds. Chem Nat
1757 Compd 38(1):24–26
- 1758 Rakhmatullaev TU, Yunusov SYu (1972a) Alkaloids of *Dipsacus*
1759 *azureus*. Chem Nat Compd 8(3):401
- 1760 Rakhmatullaev TU, Yunusov SYu (1972b) Alkaloids of *Gentiana*
1761 *olivieri*. Chem Nat Compd 9(1):56–58
- 1762 Raldugin VA, Demenkova LI, Pentegova VA (1991) Terpenoids of the
1763 oleoresin of the Schrenk spruce. Chem Nat Compd 27(3):279–282
- 1764 Raskin I, Ripoll C (2004) Can an apple a day keep the doctor away?
1765 Curr Pharm Des 10(27):3419–3429
- 1766 Rauha JP, Remes S, Heinonen M, Hopia A, Kähkönen M, Kujala T,
1767 Pihlaja K, Vuorela H, Vuorela P (2000) Antimicrobial effects of
1768 Finnish plant extracts containing flavonoids and other phenolic
1769 compounds. Int J Food Microbiol 56(1):3–12
- 1770 Razzakov NA, Aripova SF (2004) Confolidine, a new alkaloid from the
1771 aerial part of *Convolvulus subhirsutus*. Chem Nat Compd 40(1):54–55
- 1772 Razzakova DM, Bessonova IA, Yunusov SYu (1973) Alkaloids of
1773 *Haplophyllum acutifolium*. Chem Nat Compd 9(2):199–202
- 1774 Razzakova DM, Bessonova IA, Yunusov SYu (1986) Components of
1775 *Haplophyllum acutifolium*. Chem Nat Compd 22(3):363–364
- 1776 Red Data Book of Kirghiz SSR (1985) Redkiye i Nakhodyashchiesya
1777 Pod Ugrozoi Ischeznoveniya Vidui Shuvotmikh i Rastenii (The rare
1778 and endangered flora and fauna species). Izdatelstvo Akademii
1779 Nauk KyrgSSR, Frunze (in Russian)
- 1780 Revina TA, Shustova TN (1982) Study of *Galium* species in connection
1781 with their prospective medicinal use. In: Rastitel'nuiye Resursui
1782 Yuzhnoi Sibiri, ikh Ratsionalnoye Ispolzovaniye i Okhrana (Plant
1783 resources of Southern Siberia, their rational use and conservation),
1784 Tomsk, pp 91–94 (in Russian)
- 1785 Rhiouani H, Settaf A, Lyoussi B, Cherrah Y, Lacaille-Dubois MA,
1786 Hassar M (1999) Effects of saponins from *Herniaria glabra* on
1787 blood pressure and renal function in spontaneously hypertensive
1788 rats. Therapie 54(6):735–739
- Rhiouani H, Lyoussi B, Settaf A, Cherrah Y, Hassar M (2001) 1789
Antihypertensive effect of *Herniaria glabra* saponins in the sponta- 1790
neously hypertensive rats. Ann Pharm Fr 59(3):211–214 (in French) 1791
- Ribalko KS (1978) Natural sesquiterpene lactones. Meditsina, Moscow 1792
(in Russian) 1793
- Ribina EV, Kurmukov AG, Reshetov TA et al (1982) Certificate of 1794
authorship “A remedy for increasing egg-laying productivity in hens” 1795
#948365 (USSR), 7 Apr 1982, p 7, Information bulletin, no. 29. MKI 1796
A 23 KI/165. # 948365, application #3290796, p 1 (in Russian) 1797
- Ribina EV, Salikhov GV, Kurmukov AG (1983) Increasing the rate of 1798
maturity of pullets by using the phytoestrogen *Panoferol*. In: Trudy 1799
UzNIIZH (Scientific works of The Uzbek Research Institute for 1800
Livestock), 38th edn, Tekhnologiya proizvodstva produktov zhivot- 1801
novodctva [“Production technology of livestock products”]. The 1802
Uzbek Research Institute for Livestock, Tashkent, pp 113–117 (in 1803
Russian) 1804
- Ribnicky DM, Poulev A, Watford M, Cefalu WT, Raskin I (2006) 1805
Antihyperglycemic activity of Tarralin, an ethanolic extract of 1806
Artemisia dracunculus L. Phytomedicine 13(8):550–557 1807
- Ringbom T, Segura L, Noreen Y, Perera P, Bohlin L (1998) Ursolic acid 1808
from *Plantago major*, a selective inhibitor of cyclooxygenase-2 cata- 1809
lyzed prostaglandin biosynthesis. J Nat Prod 61(10):1212–1215 1810
- Ripoll C, Schmidt B, Ilić N, Poulev A, Dey M, Kurmukov AG, Raskin 1811
I (2007) Anti-inflammatory effects of a sesquiterpene lactone extract 1812
from chicory (*Cichorium intybus* L.) roots. Nat Prod Commun 1813
2(7):717–722 1814
- Rojas-Vera J, Patel AV, Dacke CG (2002) Relaxant activity of raspberry 1815
(*Rubus idaeus*) leaf extract in guinea-pig ileum in vitro. Phytother 1816
Res 16(7):665–688 1817
- Rowe GG, Castillo CA, Crumpton CW, Maxwell GM (1965) Systemic 1818
and coronary hemodynamic effects of pseudoephedrine. Proc Soc 1819
Exp Biol Med 118:682–684 1820
- Rustamov B, Tulyaganov N, Sadritdinov F (1974) Pharmacology of the 1821
alkaloid peganol. Meditsinskii Zh Uzbekistana (12):22–23 (in 1822
Russian) 1823
- Ryazantseva Z (Senior ed) (1965) Klimat Kirgizskoi SSR (Climate of 1824
the Kirghiz SSR). Ilim, Frunze (in Russian) 1825
- Ryu JH, Jeong YS, Sohn DH (1999) A new bisabolene epoxide from 1826
Tussilago farfara, and inhibition of nitric oxide synthesis in LPS- 1827
activated macrophages. J Nat Prod 62(10):1437–1438 1828
- Saadali B, Boriky D, Blaghen M, Vanhaelen M, Talbi M (2001) 1829
Alkamides from *Artemisia dracunculus*. Phytochemistry 58(7): 1830
1083–1086 1831
- Saatov Z, Syrov VN, Mamatkhanov AU, Abubakirov NK (1994) 1832
Phytoecdysteroids of *Ajuga* and their biological activity. 1. 1833
Distribution and chemical structures of the compounds isolated. 1834
Chem Nat Compd, (30)2:138–145 1835
- Saatov Z, Usmanov BZ, Abubakirov NK (1977) Phytoecdyzones of 1836
Ajuga turkestanica. IV. Chem Nat Compd 13(3):359 1837
- Saatov Z, Gorovits MB, Abubakirov NK (1990) Phytoecdysteroids of 1838
the plant *Dianthus hoeltzeri*. Chem Nat Compd 26(6):717–718 1839
- Saatov Z, Agzamkhodzhaeva DA, Syrov VN (1999) Distribution of 1840
phytoecdysteroids in plants of Uzbekistan and the possibility of 1841
using drugs based on them in neurological practice. Chem Nat 1842
Compd 35(2):186–191 1843
- Sadritdinov FS (1968) Effect of the alkaloids, dubinidine and haplophi- 1844
lidine, on higher neural activity. Meditsinskii Zh Uzbekistana 1845
(6):71–73 (in Russian) 1846
- Sadritdinov FS (1971a) Comparative study of the antiinflammatory 1847
properties of alkaloids from *Gentiana* plants. In: Sultanov MB (ed) 1848
Farmakologiya Alkaloidov i Serdechnykh Glikozidov (Pharmacology 1849
of alkaloids and cardiac glycosides). FAN, Tashkent, pp 146–148 1850
(in Russian) 1851
- Sadritdinov FS (1971b) Antiinflammatory activity of 1852
bis(benzylisoquinoline) alkaloids from *Thalictrum* plants in rela- 1853
tion to their chemical structure. In: Sultanov MB (ed) Farmakologiya 1854

- 1855 Alkaloidov i Serdechnykh Glikozidov (Pharmacology of alkaloids
1856 and cardiac glycosides). FAN, Tashkent, pp 122–125 (in Russian)
- 1857 Sadritdinov FS (1973) Effects of the alkaloids thalamicine and thalimi-
1858 dine on the central nervous system. Meditsinskii Zh Uzbekistana
1859 12:40–43 (in Russian)
- 1860 Sadritdinov FS, Khamdamov I (1975) Pharmacological properties of
1861 the alkaloid thalamicine. Farmakol Toksikol 38(4):490–491
- 1862 Sadritdinov FS, Kurmukov AG (1980) Farmakologiya Rastitelnuikh
1863 Alkaloidov i Ikh Primeneniye v Meditsinye (Pharmacology of plant
1864 alkaloids and their use in medicine). Meditsina, Tashkent (in Russian)
- 1865 Sadritdinov FS, Sultanov MB (1971) Antiinflammatory activity of the
1866 alkaloids thalmine and O-methylthalicberine. In: Sultanov MB (ed)
1867 Farmakologiya Alkaloidov i Serdechnykh Glikozidov (Pharmacology
1868 of alkaloids and cardiac glycosides). FAN, Tashkent, pp 120–122
1869 (in Russian)
- 1870 Sadritdinov FS, Tulyaganov N (1967) Pharmacology of the alkaloids
1871 gentianine and gentianadine. In: Kamilov IK (ed) Farmakologiya
1872 Alkaloidov i Glikozidov (Pharmacology of alkaloids and glyco-
1873 sides). FAN, Tashkent, pp 128–137 (in Russian)
- 1874 Sadritdinov FS, Tulyaganov N (1972) Pharmacology of the new alka-
1875 loid gentianadine. In: Sultanov MB (ed) Farmakologiya Alkaloidov
1876 i Ikh Proizvodnikh (Pharmacology of alkaloids and their deriva-
1877 tives). FAN, Tashkent, pp 152–154 (in Russian)
- 1878 Sadritdinov FS, Sultanov MB, Ismailov ZF (1971) Effect of some apor-
1879 phine alkaloids on the behavior of animals. In: Sultanov MB (ed)
1880 Farmakologiya Alkaloidov i Serdechnykh Glikozidov (Pharmacology
1881 of alkaloids and cardiac glycosides). FAN, Tashkent, pp 127–132
1882 (in Russian)
- 1883 Sadyrbekov DT, Suleimenov EM, Tikhonova EV, Atazhanova GA,
1884 Tkachev AV, Adekenov SM (2006a) Component composition of
1885 essential oils from four species of the genus *Achillea* growing in
1886 Kazakhstan. Chem Nat Compd 42(3):294–297
- 1887 Sadyrbekov DT, Ryazantsev OG, Kenesov BK, Raldugin VA,
1888 Atazhanova GA, Adekenov SM (2006b) Spectral properties of
1889 methyl-2-cis-lachnophyllate. Chem Nat Compd 42(4):493–494
- 1890 Saeed SA, Gilani AH, Majoo RU, Shah BH (1997) Anti-thrombotic and
1891 anti-inflammatory activities of protopine. Pharmacol Res 36(1):1–7
- 1892 Sagdullaev BT, Shakhidoyatov RKh, Khodzhaeva MA, Chernenko TV,
1893 Turakhozhaev MT, Abduazimova MA (2001) Lipids and carbohy-
1894 drates from *Althaea nudiflora* and *A. armeniaca* Roots. Chem Nat
1895 Compd 37(3):208–212
- 1896 Sagitdinova GV, Saidkhodzhaev AI (1977) The structures of juniferin
1897 and juniferinin. Chem Nat Compd 13(6):665–670
- 1898 Sagitdinova GV, Saidkhodzhaev AI, Malikov VM (1978) The struc-
1899 tures of juferin and juniferidin. Chem Nat Compd 14(6):693–694
- 1900 Sagitdinova GV, Eshbakova KA, Khushbaktova ZA, Malikov VM,
1901 Olimov V (1992) Flavonoid pulicarpin from *Pulicaria salviifolia*
1902 and its hypolipidemic activity. Chem Nat Compd 28(3–4):286–288
- 1903 Sagitdinova GV, Eshbakova KA, Malikov VM (1994) Diterpenoids of
1904 *Pulicaria salviifolia*. III. The structure of salvicinin. Chem Nat
1905 Compd 30(2):226–228
- 1906 Sagitdinova GV, Makhmudov MK, Tashkhodzhaev B, Mal'tsev II
1907 (1996) Labdanoids of *Marrubium anisodon*. Chem Nat Compd
1908 32(1):43–46
- 1909 Sahin F, Gulluce M, Daferera D, Sokmen A, Sokmen M, Polissiou M,
1910 Agar G, Ozer H (2004) Biological activities of the essential oils and
1911 methanol extract of *Origanum vulgare* ssp. *vulgare* in the Eastern
1912 Anatolia region of Turkey. Food Control 15(7):549–557
- 1913 Saidkasimov TK (1960) About the pharmacology of the alkaloid ervin-
1914 ine. In: Voprosy Biologii i Krayevoi Meditsinui (Issues of biology
1915 and regional medicine). Izdatelstvo Akademii Nauk UzSSR,
1916 Tashkent, pp 242–247 (in Russian)
- 1917 Saidkhodzhaev AI, Nikinov GN (1973) Structure of ferutin. Chem
1918 Nat Compd 9(1):25–26
- 1919 Saidkhodzhaev AI, Nikinov GN (1974) Structure of ferutinol. Chem
1920 Nat Compd 10(2):178–181
- Sairafianpour M, Christensen J, Staerk D, Budnik BA, Kharazmi A, 1921
Bagherzadeh K, Jaroszewski JW (2001) Leishmanicidal, antiplas- 1922
modial, and cytotoxic activity of novel diterpenoid 1,2-quinones 1923
from *Perovskia abrotanoides*: new source of tanshinones. J Nat Prod 1924
64(11):1398–1403 1925
- Sakhobiddinov SS (1948) Dikorastushchiye Lekarstvennuiye Rasteniya 1926
Srednei Azii (Wild-growing medicinal plants of Central Asia). 1927
Gosudarstvennoye Izdatelstvo UzSSR, Tashkent (in Russian) 1928
- Salehi P, Sonboli A, Eftekhari F, Nejad-Ebrahimi S, Yousefzadi M 1929
(2005) Essential oil composition, antibacterial and antioxidant 1930
activity of the oil and various extracts of *Ziziphora clinopodioides* 1931
subsp. *rigida* (BOISS.) RECH. f. from Iran. Biol Pharm Bull 1932
28(10):1892–1896 1933
- Salimov BT, Turgunov KK, Tashkhodzhaev B, Dzhakhangirov FN 1934
(2004) Structure and antiarrhythmic activity of 12-acetyl-12-epi- 1935
nappelline, a new diterpenoid alkaloid from *Aconitum soongoricum*. 1936
Chem Nat Compd 40(2):151–155 1937
- Samikov K, Shakirov R, Yunusov SYu (1989) Alkaloids of *Korolkowia* 1938
sewerzowii. Structure of korselidine. Chem Nat Compd 25(1):28–30 1939
- San Feliciano A, Gordialza M, Miguel del Corral JM, Castro MA, 1940
García-Grávalos MD, Ruiz-Lázaro P (1993) Antineoplastic and anti- 1941
viral activities of some cyclo lignans. Planta Med 59(3):246–249 1942
- Sang SM, Lao AN, Leng Y, Gu ZP, Chen ZL, Uzawa J, Fujimoto Y 1943
(2000) Segetoside F a new triterpenoid saponin with inhibition of 1944
luteal cell from the seeds of *Vaccaria segetalis*. Tetrahedron Lett 1945
41(48):9205–9207 1946
- Sang SM, Xia Z, Lao A, Cao L, Chen Z, Uzawa J, Fujimoto Y (2003) 1947
Studies on the constituents of the seeds of *Vaccaria segetalis*. 1948
Heterocycles 59(2):811–821 1949
- Sardari S, Amin G, Micetich RG, Daneshlab M (1998) 1950
Phytopharmaceuticals. Part 1. Antifungal activity of selected Iranian 1951
and Canadian plants. Pharm Biol 36(3):180–188 1952
- Sarkisyants RG (1969a) Influence of *Ferula* of experimental arterial 1953
pressure. Meditsinskii Zh Uzbekistana (9):23–24 1954
- Sarkisyants RG (1969b) Influence of *Ferula* on the fuction of the gas- 1955
tro-intestinal tract of experimental animals. In: Tezisui 1 Nauchnoi 1956
Konferentsii Tsentralnoi Nauchno-Issledovatel'skoi Laboratorii 1957
(Proceedings of the 1st conference of the Central Research Scientific 1958
Laboratory), Samarkand, pp 19–20 1959
- Sarkisyants RG (1972) Pharmacological characterization of *Ferula* 1960
(galenic forms, asafetids and sodium salts of galbanic acids. Synopsis 1961
of dissertation for the degree of Ph.D. in medicine (in Russian) 1962
- Sarkisyants RG, Azizova OM (1971) Influence of *Ferula* on progres- 1963
sion and development of the Arthus reaction. Meditsinskii Zh 1964
Uzbekistana (9):46–47 1965
- Savona G, Bruno M, Rodriguez B, Marco JL (1987) Triterpenoids from 1966
Salvia deserta. Phytochemistry 26(12):3305–3308 1967
- Savoskin IP, Pekker EG, Kadirova RB (1971) Amino acids and vitamin 1968
potential of some plant species in the bean family (Leguminosae) 1969
found growing the the south-east Altai. In: Rastitelnuie Bogatstva 1970
Sibiri (Plant riches of Siberia). Nauka, Novosibirsk, pp 104–109 (in 1971
Russian) 1972
- Schinella GR, Giner RM, Recio MC, Mordujovich de Buschiazio P, Ríos 1973
JL, Mániz S (1998) Anti-inflammatory effects of South American 1974
Tanacetum vulgare. J Pharm Pharmacol 50(9):1069–1074 1975
- Schmidt JH, Wells R (1990) Evidence for the presence of gossypol in 1976
malvaceous plants other than those in the “cotton tribe”. J Agric 1977
Food Chem 38(2):505–508 1978
- Schmidt BM, Ilic N, Poulev A, Raskin I (2007) Toxicological evaluation 1979
of a chicory root extract. Food Chem Toxicol 45(7):1131–1139 1980
- Schröder H, Schubert-Zsilavecz M, Reznicek G, Cart J, Jurenitsch J, 1981
Haslinger E (1993) A triterpene saponin from *Herniaria glabra*. 1982
Phytochemistry 34(6):1609–1613 1983
- Schultheiss PC, Knight AP, Traub-Dargatz JL, Todd FG, Stermitz FR 1984
(1995) Toxicity of field bindweed (*Convolvulus arvensis*) to mice. 1985
Vet Hum Toxicol 37(5):452–454 1986

- 1987 Seifert K, Preiss A, John S, Schmidt J, Lien NT, Lavaud C, Massiot G
1988 (1991) Triterpene saponins from *Verbascum songaricum*.
1989 Phytochemistry 30(10):3395–3400
- 1990 Selivanova LN (1954) Pharmacological study of the alkaloids allocrip-
1991 topine, protopine and berberine. Synopsis of dissertation for the
1992 degree of Ph.D., Moscow (in Russian)
- 1993 Semple SJ, Pyke SM, Reynolds GD, Flower RL (2001) In vitro antiviral
1994 activity of the anthraquinone chrysophanic acid against poliovirus.
1995 Antiviral Res 49(3):169–178
- 1996 Sener B (1994) Recent results in the search for bioactive compounds from
1997 Turkish medicinal plants. Pure Appl Chem 66(10/11): 295–2298
- 1998 Serbin AG, Borisov MI, Chernobai VT (1972a) Flavonoids of *Bidens*
1999 *tripartita*. I. Chem Nat Compd 8(1):119–120
- 2000 Serbin AG, Borisov MI, Chernobai VT (1972b) Flavonoids of *Bidens*
2001 *tripartita*. II. Chem Nat Compd 8(4):439–441
- 2002 Serbin AG, Borisov MI, Chernobai VT (1972c) Coumarins of *Bidens*
2003 *tripartita*. Chem Nat Compd 8(5):652
- 2004 Serbin AG, Borisov MI, Chernobai VT, Kovalev IP, Gordienko VG
2005 (1975) Flavonoids of *Bidens tripartita*. III. Chem Nat Compd
2006 11(2):160–162
- 2007 Seredin RM, Sokolov SD (1969) Lekarstvenniye Rasteniya i Ikh
2008 Primneniye, izd 2 (Medicinal plants and their uses), 2nd edn.
2009 Stavropolskoye knizhnoye izdatelstvo, Stavropol (in Russian)
- 2010 Sezik E, Yesilada E, Shadidoyatov H, Kulivey Z, Nigmatullaev AM,
2011 Aripov HN, Takaishi Y, Takeda Y, Honda G (2004) Folk medicine
2012 in Uzbekistan. I. Toshkent, Djizzax, and Samarqand provinces. J
2013 Ethnopharmacol 92(2–3):197–207
- 2014 Sezik E, Aslan M, Yesilada E, Ito S (2005) Hypoglycaemic activity of
2015 *Gentiana olivieri* and isolation of the active constituent through bio-
2016 assay- directed fractionation techniques. Life Sci 76(11):1223–1238
- 2017 Shafiee A, Morteza-Semnani K, Amini M (1998) (+)-Bulbocapnine-beta-
2018 N-oxide from *Glaucium fimbriigerum*. J Nat Prod 61(12):1564–1565
- 2019 Shahidi Bonjar GH (2004) Evaluation of antibacterial properties of
2020 Iranian medicinal-plants against *Micrococcus luteus*, *Serratia mar-*
2021 *escens*, *Klebsiella pneumoniae* and *Bordetella bronchoseptica*.
2022 Asian J Plant Sci 3(1):82–86
- 2023 Shakirov R, Kul'kova VV, Nakhatov I (1995) Alkaloids of *Veratrum*
2024 *lobelianum* verdinine and 3,15-DI-O-(2-methylbutyroyl)germine.
2025 Chem Nat Compd 31(1):79–82
- 2026 Shamansurov ShSh, Sultanov MB (1967) Use of vincanine hydrochlor-
2027 ide in neurologic practice. Zh Nevropatol Psikh 67(12):1807–1809
2028 (in Russian)
- 2029 Sharapov IM (1959) Pharmacology of peganine. Farmakol Toksikol
2030 22(1):69–73 (in Russian)
- 2031 Sharifi AM, Darabi R, Akbarloo N (2003) Study of antihypertensive
2032 mechanism of *Tribulus terrestris* in 2K1C hypertensive rats: role of
2033 tissue ACE activity. Life Sci 73(23):2963–2971
- 2034 Sharon N, Lis H (2004) History of lectins: from hemagglutinins to bio-
2035 logical recognition molecules. Glycobiology 14(11):53R–62R
- 2036 Shikishima Y, Takaishi Y, Honda G, Ito M, Takeda Y, Kodzhimatov
2037 OK, Ashurmetov O (2001) Phenylbutanoids and stilbene derivatives
2038 of *Rheum maximowiczii*. Phytochemistry 56(4):377–381
- 2039 Shimanov VG (1972) Gormonalnaya Aktivnost Pastbishchniikh
2040 Rastanii I vliyaniye ikh na Plodovitost Karakulskikh Ovets
2041 (Hormonal activity of pasture plants and their influence on the ferti-
2042 lity of karakul sheep). FAN, Tashkent (in Russian)
- 2043 Shreter AI (1975) Lekarstvennaya Flora Sovetskogo Dalnego Vostoka
2044 (Medicinal flora of the Soviet Far East). Meditsina, Moscow (in
2045 Russian)
- 2046 Shvarev IF, Tsetlin AL (1972) Anti-blastic properties of berberine and
2047 its derivatives. Farmakol Toksikol 35:73–75
- 2048 Sidjimov AK, Tawara JN, Stermitz FR, Rithner CD (1998) An isopavine
2049 alkaloid from *Thalictrum minus*. Phytochemistry 48(2):403–405
- 2050 Simsek I, Aytekin F, Yesilada E, Yildirimli S (2004) An ethnobotanical
2051 survey of the Beypazari, Ayas, and GÜdül district towns of Ankara
2052 province (Turkey). Econ Bot 58(4):705–720
- Singh NP, Lai HC (2004) Artemisinin induces apoptosis in human can-
2053 cer cells. Anticancer Res 24(4):2277–2280 2054
- Singh VP, Yadav B, Pandey VB (1999) Flavanone glycosides from
2055 *Alhagi pseudalhagi*. Phytochemistry 51(4):587–590 2056
- Sinitzin GC (1959) About the use of medicinal plants in traditional and
2057 veterinary medicine in the western Tian-Shan. Zhurnal Vestnik
2058 Akademii Nauk KazakSSR Alma-Ata (4):92–94 (in Russian) 2059
- Sivropoulou A, Papanikolaou E, Nikolaou C, Kokkini S, Lanaras T,
2060 Arsenakis M (1996) Antimicrobial and cytotoxic activities of
2061 *Origanum* essential oils. J Agr Food Chem 44(5):1202–1205 2062
- Sklarovsky LY (1972) Tselebniye Svoistva Pishchevuih Rastanii
2063 (Healing properties of edible plants). Rosselkhozuzdat, Moscow (in
2064 Russian) 2065
- Sokmen A, Jones BM, Erturk M (1999) Antimicrobial activity of
2066 extracts from the cell cultures of some Turkish medicinal plants.
2067 Phytother Res 13(4):355–357 2068
- Sokolov SYa, Zamotaev IP (1985) Spravochnik Po Lekarstvennuim
2069 Rastenyam. Fitoterapiya. (Guide to medicinal plants. Phytotherapy).
2070 Meditsina, Moscow (in Russian) 2071
- Sokolov SYa, Zamotaev IP (1989) Spravochnik Po Lekarstvennuim
2072 Rastenyam. Fitoterapiya, izd. 2 (Guide to medicinal plants.
2073 Phytotherapy), 2nd edn. Nedra, Moscow (in Russian) 2074
- Sonboli A, Mirjalili MH, Hadian J, Ebrahimi SN, Yousefzadi M (2006)
2075 Antibacterial activity and composition of the essential oil of
2076 *Ziziphora clinopodioides* subsp. *bungeana* (Juz.) Rech. f. from Iran.
2077 Z Naturforsch 61(9–10):677–680 2078
- Sonnenbichler J, Scalera F, Sonnenbichler I, Weyhenmeyer R (1999)
2079 Stimulatory effects of silibinin and silicristin from the milk thistle
2080 *Silybum marianum* on kidney cells. J Pharmacol Exp Ther
2081 290(3):1375–1383 2082
- Spiridonov NA, Kononov DA, Arkhipov VV (2005) Cytotoxicity of
2083 some Russian ethnomedicinal plants and plant compounds.
2084 Phytother Res 19(5):428–432 2085
- Stahl-Biskup E (2002) Essential oil chemistry of the genus thymus – a
2086 global view. In: Stahl-Biskup E, Saez F (eds) Thyme: the genus thy-
2087 mus, vol 24, Medicinal and aromatic plants – industrial profiles.
2088 Taylor & Francis, London, pp 75–124 2089
- Stepanenko GA, Gusakova SD, Umarov AU (1980) Composition of the
2090 coats and kernels of the seeds of *Nepeta pannonica* and *Lavandula*
2091 *vera*. Chem Nat Compd 16(5):434–439 2092
- Stevens KL, Riopelle RJ, Wong RY (1990) Repin, a sesquiterpene lac-
2093 tone from *Acroptilon repens* possessing exceptional biological
2094 activity. J Nat Prod 53(1):218–221 2095
- Stoychkov J, Miloushev AL (1980) Enhancement of antitumor effect of
2096 cyclophosphamide by thaliblastine in Lewis lung carcinoma and
2097 L1210 leukemia in mice. Biomedicine 33(2):42–44 2098
- Su BN, Takaishi Y, Tori M, Takaoka S, Honda G, Ito M, Takeda Y,
2099 Kodzhimatov OK, Ashurmetov O (2000) Macrophyllidimers A and
2100 B, two novel sesquiterpene dimers from the bark of *Inula macro-*
2101 *phylla*. Tetrahedron Lett 41(9):1475–1479 2102
- Su BN, Takaishi Y, Yabuuchi T, Kusumi T, Tori M, Takaoka S, Honda
2103 G, Ito M, Takeda Y, Kodzhimatov OK, Ashurmetov O (2001)
2104 Sesquiterpenes and monoterpenes from the bark of *Inula macro-*
2105 *phylla*. J Nat Prod 64(4):466–471 2106
- Su BN, Cuendet M, Farnsworth NR, Fong HHS, Pezzuto JM, Kinghorn
2107 AD (2002) Activity-guided fractionation of the seeds of *Ziziphus*
2108 *jujuba* using a cyclooxygenase-2 inhibitory assay. Planta Med
2109 68(12):1125–1128 2110
- Suladze TSh, Vachnadze VYu, Tsakadze DM, Gedevanishvili MD,
2111 Tsutsunava LE, Malazoniya NA (2006) Alkaloid accumulation
2112 dynamics in *Veratrum lobelianum* growing in Georgia and biologi-
2113 cal activity of jervine. Chem Nat Compd 42(1):71–74 2114
- Sultankhodzaev MN (1993) Karakanine – a new alkaloid from the epi-
2115 geal part of *Aconitum karakolicum*. Chem Nat Compd 29(1):51–52 2116
- Sultankhodzaev MN, Tadzhibaev MM (1976) Phenyl-β-naphthylamine
2117 from three species of plants. Chem Nat Compd 12(3):361 2118

- 2119 Sultankhodzhaev MN, Yunusov MS, Yunusov SYu (1973) Alkaloids of
2120 *Aconitum karakolicum*. Chem Nat Compd 9(1):129–130
- 2121 Sultankhodzhaev MN, Yunusov MS, Yunusov SYu (1986) Alkaloids of
2122 the epigeal part of *Aconitum karakolicum* structure of karasamine
2123 and of 1-benzoylkarasamine. Chem Nat Compd 22(2):192–195
- 2124 Sultanov MB (1959a) Pharmacology of the alkaloid of vincanidine.
2125 Izvestiya Akademii Nauk UzSSR (Seriya Meditsinskaya) (1):29–37
2126 (in Russian)
- 2127 Sultanov MB (1959b) The strychnine-like action of the alkaloid vinca-
2128 nine. Izvestiya Akad Nauk UzSSR (Seriya Meditsinskaya) (3):38–40
2129 (in Russian)
- 2130 Sultanov MB (1960) About the emetic action of the alkaloid vincani-
2131 dine. Meditsinskii Zh Uzbekistana (6):30 (in Russian)
- 2132 Sultanov MB, Kurmukov AG (1965) Pharmacology of the alkaloid
2133 ervamine. In: Farmakologiya Alkaloidov (Pharmacology of alka-
2134 loids). FAN, Tashkent, pp 128–142 (in Russian)
- 2135 Sultanova BA, Lazkov GA, Lebedeva LP, Ionov RN (1998) Preview list
2136 of vascular plants for protection and entering in the Red Data Book.
2137 Nauka i Novye Technologii (2):119–127
- 2138 Sumnevich GP (1942) Wild-growing fruit trees and shrubs. In:
2139 Suirevuiye resursii Uzbekistana (Raw material resources of
2140 Uzbekistan), vol 2(1). Tashkent, pp 10–26 (in Russian)
- 2141 Supek Z (1946) Effect of berberine on the uterus in test animals.
2142 Farmakol Toksikol 9(6):12–14 (in Russian)
- 2143 Suzutani T, Ogasawara M, Yoshida I, Azuma M, Knox YM (2003)
2144 Anti-herpesvirus activity of an extract of *Ribes nigrum* L. Phytother
2145 Res 17(6):609–613
- 2146 Svechnikova A, Umarova RU, Gorovits MB, Abdullaev ND, Abubakirov
2147 NK (1983) Triterpene glycosides of *Astragalus* and their genins. XI.
2148 Cyclosiversioside G – a triglycoside of *Astragalus sieversianus*.
2149 Chem Nat Compd 19(3):296–299
- 2150 Syrov VN (1984) Mechanism of the anabolic action of phytoecdysteroids
2151 in mammals. Biologicheskiiye Nauki (11):16–20 (in Russian)
- 2152 Syrov VN (1994) Phytoecdysteroids: their biological effects in the body
2153 of higher animals and the outlook for their use in medicine.
2154 Eksperimentalnaya i Klinicheskaya farmakologiya 57(5):61–66 (in
2155 Russian)
- 2156 Syrov VN, Kurmukov AG (1975) On the anabolic action of viticoster-
2157 one E. Dokl Akad Nauk UzSSR 32(6):31–32 (in Russian)
- 2158 Syrov VN, Kurmukov AG (1975a) The anabolic and androgenic prop-
2159 erties of ecdysterone and methyl-androstenediol. In: Voprosy
2160 Farmakologii i Toksikologii (Issues in pharmacology and toxicol-
2161 ogy). Meditsina, Tashkent, pp 22–25 (in Russian)
- 2162 Syrov VN, Kurmukov AG (1975c) Effect of Viticosterone E on the
2163 weight of organs and their protein content in rats of different ages.
2164 Dokl Akad Nauk UzSSR (9):40–41 (in Russian)
- 2165 Syrov VN, Kurmukov AG (1976a) Anabolic activity of phytoecdysone-
2166 ecdysterone isolated from *Rhaponticum carthamoides* (Willd.) Iljin.
2167 Farmakol Toksikol 39(6):690–693 (in Russian)
- 2168 Syrov VN, Kurmukov AG (1976b) The anabolic properties of turkester-
2169 one phytoecdysone and turkesterone tetraacetate in experiments on
2170 male rats. Problemy Endokrinologii 22(3):107–112 (in Russian)
- 2171 Syrov VN, Kurmukov AG (1976c) About the biological activity of
2172 cyasterone in tests on male rats. Biologicheskiiye Nauka (5):72–74 (in
2173 Russian)
- 2174 Syrov VN, Kurmukov AG (1976d) Influence of ecdysterones on haem-
2175 atopoiesis of experimental posthemorrhagic anemia in lab animals.
2176 In: Aktualniye Voprosy Akusherstva i Ginekologii (Current issues in
2177 obstetrics and gynecology), Tashkent, pp 62–63 (in Russian)
- 2178 Syrov VN, Kurmukov AG (1977) Tonic properties of ecdysterone iso-
2179 lated from *Rhaponticum carthamoides*. Dokl Akad Nauk UzSSR
2180 (12):27–30 (in Russian)
- 2181 Syrov VN, Kurmukov AG (1980) Influence of phytoecdysones, their
2182 6-ketoanalogs, and nerobol on anabolic processes in heart muscle
2183 (experimental data). In: Materialui IV Nauchno-Prakticheskoi
2184 Konferentsii IV Glavnogo Upravleniya pri Min. Zdravye UzSSR
(Proceedings of the 4th scientific-practioners conference of 4th
General Directorate of the Ministry of Health of the UzSSR), pp
119–122 (in Russian)
- Syrov VN, Aizikov MI, Kurmukov AG (1975a) Effect of ecdysterone
on the content of protein, glycogen, and fat in white rat liver, heart,
and muscle. Dokl Akad Nauk UzSSR (8):37–38 (in Russian)
- Syrov VN, Kurmukov AG, UsmanovBZ (1975b) About the anabolic
action of turkesterone and tetra-acetate of turkesterone. Dokl Akad
Nauk UzSSR (2):32–34 (in Russian)
- Syrov VN, Kurmukov AG, Akhmedkhodzhaeva KhS (1976) About the
anabolic and androgenic actions of drupanol. Dokl Akad Nauk
UzSSR (1):34–36 (in Russian)
- Syrov VN, Matveev SB, Kurmukov AG, IslambekovUS (1986) Effect
of ecdysterone and nerobol on the healing of experimental bone
fractures. Meditsinskii Zh Uzbekistana (3):67–69 (in Russian)
- Szabo I, Vonzaz G, Pallag A, Nemeth S (2006) In vitro specific anti-
microbial properties of *Hibiscus trionum* L. extractive solutions. In:
Proceedings of 4th conference of the association for medicinal and
aromatic plants of South-East European countries. Ia i, România,
Alma Mater, Sibiu, 28–31 May 2006
- Tabatabaei-Anaraki M, Chalabian F, Masoudi S, Rustaiyan A (2007) The
chemical composition and in vitro antibacterial activities of the oil of
Ziziphora clinopodioides Lam. from Iran. Planta Med 73(9):94
- Taborska E, Dostal J, Bochorakova H, Veznik F (1988) Alkaloids of
Papaver argemone L. and *Papaver pavoninum* Fisch. et Mey. from
the Argemonorhoeades FEDDE section. Collect of Czech Chem
Commun 53(8):1845–1850
- Tada Y, Shikishima Y, Takaishi Y, Shibata H, Higuti T, Honda G, Ito M,
Takeda Y, Kodzhimatov OK, Ashurmetov O, Ohmoto Y (2002)
Coumarins and γ -pyrone derivatives from *Prangos pabularia*: anti-
bacterial activity and inhibition of cytokine release. Phytochemistry
59(6):649–654
- Tadzhibaev YuA, Mukhamedova KhS, Akramov ST (1977)
Phospholipids of the seeds of two species of *Erysimum*. Chem Nat
Compd 13(5):517–520
- Tairov T (1969) Steroid sapogenins of *Asparagus persicus*. Chem Nat
Compd 5(4):277
- Tajadod G, Majd A (2007) Antibacterial effects of alcoholic extracts of
different parts of *Mentha longifolia* and *Ziziphora tenuior*. Planta
Med 73(9):176
- Takeda Y, Tomonari M, Arimoto S, Masuda T, Otsuka H, Matsunami K,
Honda G, Ito M, Takaishi Y, Kiuchi F, Khodzhimatov OK,
Ashurmetov OA (2008) A new phenolic glucoside from an Uzbek
medicinal plant, *Origanum tyttanthum*. J Nat Med 62(1):71–74
- Talalal TS, Dwivedi SK, Sharma SR (1995) In vitro and in vivo
antitrypanosomal activity of *Xanthium strumarium* leaves.
J Ethnopharmacol 49(3):141–145
- Tamas M, Doina S, Simina T (2006) Comparative phytochemical
research of *Galium verum* L. and *G. mollugo* L. Notulae Botanicae
Horti Agrobotanici Cluj-Napoca 34(1):18–20
- Tamayo C, Richardson MA, Diamond S, Skoda I (2000) The chemistry
and biological activity of herbs used in Flor-Essence™ herbal
tonic and Essiac. Phytother Res 14(1):1–14
- Tamemoto K, Takaishi Y, Chen B, Kawazoe K, Shibata H, Higuti T,
Honda G, Ito M, Takeda Y, Kodzhimatov OK, Ashurmetov O (2001)
Sesquiterpenoids from the fruits of *Ferula kuhistanica* and antibac-
terial activity of the constituents of *F. kuhistanica*. Phytochemistry
58(5):763–767
- Tanaka N, Takaishi Y, Shikishima Y, Nakanishi Y, Bastow K, Lee KH,
Honda G, Ito M, Takeda Y, Kodzhimatov OK, Ashurmetov O (2004)
Prenylated benzophenones and xanthenes from *Hypericum scabrum*.
J Nat Prod 67(11):1870–1875
- Tashbaev KH, Sultanov MB (1965) Antiepileptic and sedative proper-
ties of thalozopine dihydriodide. Farmakologiya Alkaloidov,
Akademii Nauk UzSSR, Inst Khim Rast Veshchestv (2):197–212
(in Russian)

- 2251 Tashbaev KhI, Sultanov MB (1962) Anticonvulsive action of the alka- 2317
 2252 loid talizopine. *Farmakologiya Alkaloidov, Akademii Nauk UzSSR,* 2318
 2253 *Inst Khim Rast Veshchestv* (1):210–219 (in Russian) 2319
- 2254 Tashkhodzhaev B, Aripova SF, Turgunov KK, Abdilalimov O (2004) 2320
 2255 Stereochemistry of the alkaloids codonopsine and codonopsinine. 2321
 2256 *Chem Nat Compd* 40(6):618–619 2322
- 2257 Tashmukhamedova MA, Mukhina OA, Syrov VN, Khushbaktova ZA, 2323
 2258 Katkova SP, Kosovskii MI (1992) Hypoglycemic activity of bicyclic 2324
 2259 diterpenoids of the clerodane series as compared to adebit and 2325
 2260 maninil. *Problemy Endokrinologii* 38(1):48–50 (in Russian) 2326
- 2261 Tawata M, Aida K, Noguchi T, Ozaki Y, Kume S, Sasaki H, Chin M, 2327
 2262 Onaya T (1992) Anti-platelet action of isoliquiritigenin, an aldose 2328
 2263 reductase inhibitor in licorice. *Eur J Pharmacol* 212(1):87–92 2329
- 2264 Teslov SV (1960) Pharmacognostic studies of some Central Asian spe- 2330
 2265 cies in the genus *Thermopsis* R.Br. Synopsis of dissertation for the 2331
 2266 degree of Ph.D. in pharmacology, University of Tartu, Tartu (in 2332
 2267 Russian) 2333
- 2268 Tezuka Y, Kasimu R, Li JX, Basnet P, Tanaka K, Namba T, Kadota S 2334
 2269 (1998) Constituents of roots of *Salvia deserta* Schang. (Xinjiang- 2335
 2270 Danshen). *Chem Pharm Bull* 6(1):107–112 2336
- 2271 Tezuka Y, Fan W, Kasimu R, Kadota S (1999) Screening of crude drug 2337
 2272 extracts for prolyl endopeptidase inhibitory activity. *Phytochemistry* 2338
 2273 6(3):197–203 2339
- 2274 Tigno XT, de Guzman F, Flora AM (2000) Phytochemical analysis and 2340
 2275 hemodynamic actions of *Artemisia vulgaris* L. *Clin Hemorheol* 2341
 2276 *Microcirc* 23(2–4):167–175 2342
- 2277 Todd FG, Stermitz FR, Schultheis P, Knight AP, Traub-Dargatz J (1995) 2343
 2278 Tropane alkaloids and toxicity of *Convolvulus arvensis*. 2344
 2279 *Phytochemistry* 39(2):301–303 2345
- 2280 Todorov DK, Zeller WJ (1992) Antiproliferative activity of the non- 2346
 2281 myelotoxic antitumor agent of plant origin thaliblastine on two 2347
 2282 human glioma cell lines. *J Cancer Res Clin Oncol* 118(2):83–86 2348
- 2283 Tolmachev AI (Senior ed) (1976) Atlas Arealov I Resursov 2349
 2284 Lekarstvennuikh Rastenii SSSR (Atlas of distributions and resources 2350
 2285 of medicinal plants of the USSR), Moscow (in Russian) 2351
- 2286 Tolomeo M, Grimaudo S, Di Cristina A, Roberti M, Pizzirani D, Meli 2352
 2287 M, Dusonchet L, Gebbia N, Abbadessa V, Crosta L, Barucchetto R, 2353
 2288 Grisolia G, Invidiata F, Simoni D (2005) Pterostilbene and 2354
 2289 3'-hydroxypterostilbene are effective apoptosis-inducing agents in 2355
 2290 MDR and BCR-ABL-expressing leukemia cells. *Int J Biochem Cell* 2356
 2291 *Biol* 37(8):1709–1726 2357
- 2292 Trutneva EA, Berezhinskaya VV (1960) Pharmacology of Lupinan 2358
 2293 group of alkaloids. *Farmakol Toksikol* (5):445–449 2359
- 2294 Tsevegsuren N, Christie WW, Lösel D (1998) *Tanacetum* 2360
 2295 (*Chrysanthemum corymbosum* seed oil—a rich source of a novel 2361
 2296 conjugated acetylenic acid. *Lipids* 33(7):723–727 2362
- 2297 Tsitsina NB (Senior ed) (1962) Atlas Lekarstvennuikh Rastenii SSSR 2363
 2298 (Atlas of medicinal plants of the USSR). Izdatelstvo Meditsinskaya 2364
 2299 literatura, Moscow (in Russian) 2365
- 2300 Tsukervanik IP, Bersutskii VP, Burtseva CV, Aizikovich BI (1935) 2366
 2301 Composition of (B) roots of *Ferula pseudoreosolinum* Lipsky. *Bull* 2367
 2302 *Univ Asie Centrale* (21):55–68 (in Russian) 2368
- 2303 Tsukurvanik IP, Simkhaev NG (1948) Study of *Ferula sumbul* Hooker 2369
 2304 f. roots. In: *Trudy Instituta Khimii Akademii Nauk UzSSR:* 2370
 2305 *Khimicheskiye Issledovaniya Dikikh Rastenii Srednei Azii, vuip. I* 2371
 2306 (Works of the institute of chemistry of the academy of science of 2372
 2307 Uzbekistan: chemical studies of the wild plants of Central Asia), vol 2373
 2308 1. Izdatelstvo Akademii Nauk UzSSR, Tashkent, pp 24–26 (in 2374
 2309 Russian) 2375
- 2310 Tulyaganov NT (1976) Cholinergic properties of quinazoline alkaloids 2376
 2311 from *Peganum harmala*. In: Sultanov MB (ed) *Farmakologiya* 2377
 2312 *Rastitelnuikh Veshestv* (Pharmacology of plant substances). FAN, 2378
 2313 Tashkent, pp 101–110 (in Russian) 2379
- 2314 Tulyaganov NT, Sadritdinov FS (1968) Effects of gentianine on the 2380
 2315 central nervous system. *Meditsinskii Zh Uzbekistana* (11):51–53 (in 2381
 2316 Russian) 2382
- Tulyaganov NT, Danilevskii BL, Sadritdinov FS (1971) Comparative 2317
 sedative activity of gentianine, gentianadine, mepropan, and ami- 2318
 nazine. In: Sultanov MB (ed) *Farmakologiya Alkaloidov i* 2319
Serdechnykh Glikozidov (Pharmacology of alkaloids and cardiac 2320
 glycosides). FAN, Tashkent, pp 148–151 (in Russian) 2321
- Tulyaganov NT, Dzhakhangirov FN, Sadritdinov FS, Khamdamov I 2322
 (1976) Pharmacology of aconite alkaloids. In: Sultanov MB (ed) 2323
Farmakologiya Rastitelnuikh Veshestv (Pharmacology of plant sub- 2324
 stances). FAN, Tashkent, pp 76–91 (in Russian) 2325
- Tulyaganov N, Sadritdinov FS, Sule manova GA (1986) Pharmacological 2326
 characteristics of desoxyepganine hydrochloride. *Farmakol Toksikol* 2327
 49(3):37–40 (in Russian) 2328
- Tunon H, Olavsdotter C, Bohlin L (1995) Evaluation of anti- 2329
 inflammatory activity of some Swedish medicinal plants. Inhibition 2330
 of prostaglandin biosynthesis and PAF-induced exocytosis. 2331
J Ethnopharmacol 48(2):61–76 2332
- Turdumambetov K (1995) Carbohydrates from plants in the genus 2333
Cousinia. Synopsis of dissertation for the degree of Ph.D. in chem- 2334
 istry, Bishkek (in Russian) 2335
- Turdumambetov K, Rakhimov DA, Malikova MKh (2007) Oligo- and 2336
 polysaccharides from *Cousinia umbrosa*. *Chem Nat Compd* 2337
 43(3):308–309 2338
- Turker AU, Camper ND (2002) Biological activity of common mullein, 2339
 a medicinal plant. *J Ethnopharmacol* 82(2–3):117–125 2340
- Turker AU, Gurel E (2005) Common mullein (*Verbascum thapsus* L.): 2341
 recent advances in research. *Phytother Res* 19(9):733–739 2342
- Turova AD (1974) Lekarstvennuye Rasteniya SSSR i ikh Premeneniye 2343
 (Medicinal plants of the USSR and their uses). *Meditsina, Moscow* 2344
 (in Russian) 2345
- Turova AD, Nikolskaya BS (1954) Short notes on the medicinal flora of 2346
 Moscow Province. *Farmakol Toksikol* 17(1):54–58 (in Russian) 2347
- Turova AD, Sapozhnikova EN (1984) Lekarstvennuye Rasteniya SSSR 2348
 i ikh Premeneniye (Medicinal plants of the USSR and their uses), 2349
 4th edn. *Meditsina, Moscow* (in Russian) 2350
- Tursunova SA (1973) Pharmacology of aphylline hydrochloride. 2351
Nauchnye Trudy – Tashkentskii Gosudarstvennyi Universitet im. V. 2352
I. Lenina (457):64–68 (in Russian) 2353
- Tursunova NV, Syrov VN, Khushbaktova ZA, Mukhamatkhanova RF, 2354
 Sham'yanov ID, Nabiev AN (2002) Effect of the total sesquiterpe- 2355
 noid extract from *Artemisia leucodes* on cholepoietic processes in 2356
 rat liver in the normal state and during experimental hepatitis. *Pharm* 2357
Chem J 36(2):37–39 2358
- Tye A, Baldesberger R, Lapidus JB, Patil PN (1967) Steric aspects of 2359
 adrenergic drugs. VI. Beta adrenergic effects of ephedrine isomers. 2360
J Pharmacol Exp Ther 157:356–362 2361
- Ul'chenko NT, Glushenkova AI (2001) Oxygenated fatty acids from 2362
 neutral lipids of *Artemisia leucodes* seeds. *Chem Nat Compd* 2363
 37(5):406–408 2364
- Ul'chenko NT, Gusakova SD, Glushenkova AI (1993) Oxygenated tria- 2365
 clylglycerols of the lipids of *Onopordum acanthium* seeds. *Chem* 2366
Nat Compd 29(5):578–581 2367
- Ul'chenko NT, Mukhamedova KhS, Glushenkova AI (1999) Lipids from 2368
 seeds of two species of *Cousinia*. *Chem Nat Compd* 35(6): 680–681 2369
- Ul'chenko NT, Mukhamedova KhS, Glushenkova AI (2000) Lipids of 2370
Helichrysum maracandicum flowers. *Chem Nat Compd* 36(1): 2371
 93–94 2372
- Ulubelen A, Topcu G, Eris C, Sonmez U, Kartal M, Kurucu S, Bozok- 2373
 Johansson C (1994) Terpenoids from *Salvia sclarea*. *Phytochemistry* 2374
 36(4):971–974 2375
- Umarova RU, Gorovits MB, Abubakirov NK (1988) Mussaneoside 2376
 from *Dodartia orientalis*. *Chem Nat Compd* 24(2):259 2377
- Umralina AR, Lazkov GA (2008) Endemic and rare plant species of 2378
 Kyrgyzstan (Atlas). Biotechnology Institute, National Academy of 2379
 Sciences Kyrgyz Republic, Bishkek 2380
- Unlu M, Daferera D, Donmez E, Polissiou M, Tepe B, Sokmen A (2002) 2381
 Compositions and the in vitro antimicrobial activities of the essential 2382

- 2383 oils of *Achillea setacea* and *Achillea teretifolia* (Compositae). J
2384 Ethnopharmacol 83(1–2):117–121
- [AL2005] Ur-Rehman J, Ahmad M, Saeed MA, Younas M (1991) J Chem Soc Pak
2386 13:56
- 2387 Usmanov BZ, Gorovits MB, Abubakirov NK (1975) Phytoecdysones
2388 of *Ajuga turkestanica*. III. The structure of turkesterone. Chem Nat
2389 Compd 11(4):484–487
- 2390 Usmanov BZ, Saatov Z, Abubakirov NK (1977) Phytoecdysones of
2391 *Ajuga turkestanica*. V. Chem Nat Compd 13(5):595
- 2392 Utkin LA (1931) Traditional medicinal plants of Siberia. Trudy
2393 Nauchno-Issledovatel'skogo Khimiko-farmatsevticheskogo Instituta
2394 24:1–133 (in Russian)
- 2395 Vaisov ZM, Yunusov MS (1987) Methylisotalatisidine – a new alkaloid
2396 from *Delphinium confusum*. Chem Nat Compd 23(6):725–727
- 2397 Vandisheva BI, Yusupova AA, Alimbaeva PK, Nuralieva ZhC (1977)
2398 Lekarstvennuiye Rasteniyi Kirgizii i Perspektivui Ikh Ispolzovaniya
2399 (Medicinal plants of Kyrgyzstan and perspectives on their uses).
2400 Ilim, Frunze (in Russian)
- 2401 Varamini P, Doroudchi M, Mohagheghzadeh A, Soltani M, Ghaderi A
2402 (2007) Cytotoxic evaluation of four *Haplophyllum* species with
2403 various tumor cell lines. Pharm Biol 45(4):299–302
- 2404 Venkateswarlu S, Satyanarayana B, Sureshbabu CV, Subbaraju GV
2405 (2003) Synthesis and antioxidant activity of 4-[2-(3,5-dimethoxy-
2406 phenyl)ethenyl]-1,2-benzenediol, a metabolite of *Sphaerophysa sal-
2407 sula*. Biosci Biotechnol Biochem 67(11):2463–2466
- 2408 Vilner AM (1974) Kormoviye otravleniya (Poisonous fodder plants),
2409 5th edn. Kolos, Leningrad (in Russian)
- 2410 Volinsky BT, Bender BT, Friedman CL (1983) Rasteniyi v Meditsinye
2411 (Plants in medicine). Izdatelstvo Saratovskogo Universiteta (Saratov
2412 University), Saratov (in Russian)
- 2413 Vonhoff C, Baumgartner A, Hegger M, Korte B, Biller A, Winterhoff H
2414 (2006) Extract of *Lycopus europaeus* L. reduces cardiac signs of
2415 hyperthyroidism in rats. Life Sci 78(10):1063–1070
- 2416 Voynova E, Dimitrova S, Naydenova E, Karadjov P (1991) Inhibitory
2417 action of extracts of *Maclura aurantiaca* and *Epilobium hirsutum* on
2418 tumour models in mice. Acta Physiol Pharmacol Bulg 17(4):50–52
- 2419 Vyazovskaya RD (1963) Depressive actions of plant substances on
2420 some endocrine functions. Uspekhi sovremennoi biologii 56(2):265–
2421 273 (in Russian)
- 2422 Wang Y, Ohtani K, Kasai R, Yamasaki K (1997) Steroidal saponins
2423 from fruits of *Tribulus terrestris*. Phytochemistry 45(4):811–817
- 2424 Wang AQ, Wang XK, Li JL, Cui XY (2004a) Isolation and structure
2425 identification of chemical constituents from the seeds of *Descurainia
2426 sophia* (L.) Webb ex Prantl. Yao Xue Xue Bao 39(1):46–51 (in
2427 Chinese)
- 2428 Wang QE, Ma S, Fu B, Lee FSC, Wang X (2004b) Development of
2429 multi-stage countercurrent extraction technology for the extraction
2430 of glycyrrhizic acid (GA) from licorice (*Glycyrrhiza uralensis
2431 Fisch*). Biochem Eng J 21(3):285–292
- 2432 Warholom O, Skaar S, Hedman E, Molmen HM, Eik L (2003) The
2433 effects of a standardized herbal remedy made from a subtype of
2434 *Rosa canina* in patients with osteoarthritis: a double-blind, random-
2435 ized, placebo-controlled clinical trial. Curr Ther Res 64(1):21–31
- 2436 Wegener T, Kraft K (1999) Plantain (*Plantago lanceolata* L.): anti-
2437 inflammatory action in upper respiratory tract infections. Wien Med
2438 Wochenschr 149(8–10):211–216 (in German)
- 2439 Widyarani S, Spinks N, Husband AJ, Reeve VE (2001) Isoflavonoid
2440 compounds from red clover (*Trifolium pratense*) protect from
2441 inflammation and immune suppression induced by UV radiation.
2442 Photochem Photobiol 74(3):465–470
- 2443 Williams CA, Goldstone F, Greenham J (1996) Flavonoids, cinnamic
2444 acids and coumarins from the different tissues and medicinal prepara-
2445 tions of *Taraxacum officinale*. Phytochemistry 42(1):121–127
- 2446 Williams CA, Harborne JB, Geiger H, Hoult JR (1999) The flavonoids
2447 of *Tanacetum parthenium* and *T. vulgare* and their anti-inflammatory
2448 properties. Phytochemistry 51(3):417–423
- Wink M (1999) Introduction: biochemistry, role and biotechnology of 2449
secondary metabolites. In: Wink M (ed) Biochemistry of plant sec- 2450
ondary metabolism, vol 2, Annual plant reviews. CRC Press, Boca 2451
Raton, pp 1–16 2452
- Winterhoff H, Sourgens H, Kemper FH (1983) Antihormonal effects of 2453
plant extracts. Pharmacodynamic effects of lithospermum officinale 2454
on the thyroid gland of rats; comparison with the effects of iodide. 2455
Horm Metab Res 15(10):503–507 2456
- Wittenburg H, Nehring K (1965) Studies on the effect of pure lupinine 2457
alkaloids on the animal body. The effect of lupinine on rats. 2458
Pharmazie 20:156–158 (in German) 2459
- Wolniak M, Tomczykowa M, Tomczyk M, Gudej J, Wawer I (2007) 2460
Antioxidant activity of extracts and flavonoids from *Bidens tripar-
2461 tita*. Acta Pol Pharm 64(5):441–447 2462
- World Health Organization (2006) Guidelines for the treatment of malaria. ii. 2463
<http://www.who.int/malaria/docs/TreatmentGuidelines2006.pdf>. 2464
Accessed 27 Mar 2007 2465
- Wu J, Beal JL, Wu WN, Doskotch RW (1977) Alkaloids of *Thalictrum* 2466
XXVI. New hypotensive and other alkaloids from *Thalictrum minus* 2467
race B. Lloydia 41(3):593 2468
- Wu TS, Li CY, Leu YL, Hu CQ (1999a) Limonoids and alkaloids of the 2469
root bark of *Dictamnus angustifolius*. Phytochemistry 50(3):509–512 2470
- Wu TS, Shi LS, Kuo SC (1999b) Alkaloids and other constituents from 2471
Tribulus terrestris. Phytochemistry 50(8):1411–1415 2472
- Yagudaev MR, Khalmirzaev MM, Yunusov SYu (1983) NMR study of 2473
alkaloids. V. ¹³C NMR spectra and reconsideration of the structures 2474
of 11- and 10-hydroxypleiocarpamines and the structures of new 2475
alkaloids from *Vinca erecta*. Chem Nat Compd 19(4):454–458 2476
- Yagura T, Motomiya T, Ito M, Honda G, Iida A, Kiuchi F, Tokuda H, 2477
Nishino H (2008) Anticarcinogenic compounds in the Uzbek 2478
medicinal plant, *Helichrysum maracandicum*. J Nat Med 2479
62(2):174–178 2480
- Yan W, Ohtani K, Kasai R, Yamasaki K (1997) Steroidal saponins from 2481
fruits of *Tribulus terrestris*. Phytochemistry 45(4):811–817 2482
- Yang R, Gao G, Liu T, Liu S, Li G (2007) Enhanced ability of hemoglobin 2483
to carry oxygen by salidroside. Electrochem Commun 9(1):94–96 2484
- Yasukawa K, Akihisa T, Inoue Y, Tamura T, Yamanouchi S, Takido M 2485
(1998) Inhibitory effect of the methanol extracts from compositae 2486
plants on 12-O-tetradecanoylphorbol-13-acetate-induced ear 2487
oedema in mice. Phytother Res 12(7):484–487 2488
- Yesilada E, Sezic E, Fujita T, Tanaka S, Tabata M (1993) Screening of 2489
some Turkish medicinal plants for their antiulcerogenic activities. 2490
Phytother Res 7(3):263–265 2491
- You M, Wickramaratne DB, Silva GL, Chai H, Chagwedera TE, 2492
Farnsworth NR, Cordell GA, Kinghorn AD, Pezzuto JM (1995) 2493
(–)-Roemerine, an aporphine alkaloid from *Annona senegalensis* 2494
that reverses the multidrug-resistance phenotype with cultured cells. 2495
J Nat Prod 58(4):598–604 2496
- Yue J, Xu J, Chen Y, Chen SN (1994) Diterpenoid alkaloids from 2497
Aconitum talassicum. Phytochemistry 37(5):1467–1470 2498
- Yue J, Xu J, Zhao Q, Sun H, Chen Y (1996) Diterpenoid alkaloids from 2499
Aconitum leucostomum. J Nat Prod 59(3):277–279 2500
- Yue ME, Jiang TF, Shi YP (2004) Fast determination of flavonoids in 2501
Hippophae rhamnoides and its medicinal preparation by capillary 2502
zone electrophoresis using dimethyl-β-cyclodextrin as modifier. 2503
Talanta 62(4):695–699 2504
- Yukhananov DKh, Yatsyno AI, Turova AD, Kuvaev VB (1972) 2505
Saponins of the Caryophyllaceae family and their hemolytic activ- 2506
ity. Rastitel'nye Resursy 8(3):374–377 (in Russian) 2507
- Yuldasheva NP, Rakhimov DA, Turakhozaev MT (1993) Polysaccharides 2508
of *Eremurus*. XXVI. Study of the structure of the pectin from the 2509
leaves of *Eremurus regelii*. Chem Nat Compd 29(2):154–156 2510
- Yunusov SYu (1974) Alkaloidui izd. 2 (Alkaloids), 2nd edn. FAN, 2511
Tashkent (in Russian) 2512
- Yunusov SYu (1981) Alkaloidui, izd. 3 (Alkaloids), 3rd edn. FAN, 2513
Tashkent (in Russian) 2514

- 2515 Yunusov AKh, Israilov IA (1974) A study of the alkaloids of *Glaucium*
2516 *fimbrilligerum*. Chem Nat Compd 10(4):555
- 2517 Yunusov SYu, Sichkova EV, Potemkin GF (1954) Alkaloids of *Aconitum*
2518 *talassicum* Pop. Zh Obshch Khim 24(12):2237
- 2519 Yunusov AKh, Israilov IA, Yunusov MS (1973) Alkaloids of *Glaucium*
2520 *fimbrilligerum*. Chem Nat Compd 9(5):658
- 2521 Yusupova SS, Batirov EKh, Abdullaev ShV, Malikov VM (1984)
2522 Flavonoids of *Vexibia alopecuroides*. Chem Nat Compd 20(2):233
- 2523 Zabirotv ISh, Kasmaliev DK (1962) About the pharmacology of the alka-
2524 loids thalicmine and thalimidine. In: Materialui X Vsesoyuznoi
2525 Konferentsii Farmakologov (Proceedings of the tenth All-Union con-
2526 ference of pharmacologists), Volgograd, pp 116–118 (in Russian)
- 2527 Zafrilla P, Ferreres F, Tomás-Barberán FA (2001) Effect of processing
2528 and storage on the antioxidant ellagic acid derivatives and flavonoids
2529 of red raspberry (*Rubus idaeus*) jams. J Agric Food Chem
2530 49(8):3651–3655
- 2531 Zaidi MA, Crow SA Jr (2005) Biologically active traditional medicinal
2532 herbs from Balochistan, Pakistan. J Ethnopharmacol
2533 96(1–2):331–334
- 2534 Zainutdinova UN, Pulatova MP, Badalbaeva TA, Umarova RU,
2535 Mavlyankulova ZI, Pulatova TP, Aslanova KhA (1994) Diterpene
2536 lactones and iridoid glycosides of plants of the genus *Lagochilus*.
2537 Chem Nat Compd 30(1):27–30
- 2538 Zakordonets AI (1953) Kultura Vuicokovitaminuikh Shipovnikov
2539 (Culturing *Rosa* with high vitamin content), Kiev (in Russian)
- 2540 Zapesochnaya GG, Aitbaeva RKh, Bankovsky AI (1973) Flavonoids
2541 and chlorogenic acid in *Sorbus tianschanica*. II. Chem Nat Compd
2542 9(1):112–113
- 2543 Zapryagaeva VI (1964) Dikorastushchiye Plodovuiye Tadzshikistana
2544 (Wild growing fruit-bearing species of Tajikistan). Nauka, Moscow-
2545 Leningrad (in Russian)
- 2546 Zatsny IL, Gorovits MB, Abubakirov NK (1971) Ecdysterone from
2547 *Serratula sogdiana*. Chem Nat Compd 7(6):822
- 2548 Zatsny IL, Gorovits MB, Abubakirov NK (1973a) Arbuitin from
2549 *Serratula sogdiana*. Chem Nat Compd 9(3):415
- 2550 Zatsny IL, Gorovits MB, Abubakirov NK (1973b) Phytoecdysones of
2551 *Serratula* II. Viticosterone E from *Serratula sogdiana* and its partial
2552 synthesis. Chem Nat Compd 9(2):170–173
- 2553 Zavrazhanov VI, Kitaeva RI, Khmelev KF (1972) Lekarstvennuiye
2554 Rasteniya Tsentralnogo Chernozemya (Medicinal plants of Central
2555 Chernozem). Izdatelstvo Voronezh Inst, Voronezh (in Russian)
- 2556 Zavrazhanov VI, Kitaeva RI, Khmelev KF (1977) Lekarstvennuiye
2557 Rasteniya Tsentralnogo Chernozemya (Medicinal plants of Central
2558 Chernozem). Izdatelstvo Voronezh Inst, Voronezh (in Russian)
- 2559 Zemlinsky SYa (1958) Lekarstvennuiye Rasteniya SSSR, 3-ye izd
2560 (Medicinal plants of the USSR), 3rd edn. Medgiz, Moscow (in
2561 Russian)
- 2562 Zhang YD, Shen JP, Zhu SH, Huang DK, Ding Y, Zhang XL (1992)
2563 Effects of *Astragalus* (ASI, SK) on experimental liver injury. Yao
2564 Xue Xue Bao 27(6):401–406 (in Chinese)
- 2565 Zhang L, Li G, Houghton PJ, Jackson S, Twentyman PR (1997)
2566 Alkaloids in *Sophora alopecuroides* seed and relevant tests for activ-
2567 ity. Zhongguo Zhong Yao Za Zhi 22(12):740–764 (in Chinese)
- Zhang L, Lau YK, Xia W, Hortobagyi GN, Hung MC (1999) Tyrosine
kinase inhibitor emodin suppresses growth of HER-2/neu-overex-
pressing breast cancer cells in athymic mice and sensitizes these
cells to the inhibitory effect of paclitaxel. Clin Cancer Res
5(2):343–353
- Zhang L, Yu H, Sun Y, Lin X, Chen B, Tan C, Cao G, Wang Z (2007)
Protective effects of salidroside on hydrogen peroxide-induced
apoptosis in SH-SY5Y human neuroblastoma cells. Eur J Pharmacol
564(1–3):18–25
- Zhao XY, Wang Y, Li Y, Chen XQ, Yang HH, Yue JM, Hu GY (2003)
Songorine, a diterpenoid alkaloid of the genus *Aconitum*, is a novel
GABA(A) receptor antagonist in rat brain. Neurosci Lett
337(1):33–36
- Zhao C-C, Shao J-H, Li X, Kang X-D, Zhang Y-W, Meng D-L, Li N
(2008) Flavonoids from *Galium verum* L. J Asian Nat Prod Res
10(7):611–615
- Zhou WC (2001) Study on chemical constituents of unsaponifiable
matter in chlorophyll-carotene paste from needles and twigs of
Picea schrenkiana. Chem Ind Forest Prod 21(2):53–56 (in
Chinese)
- Zhou BN, Bai NS, Lin LZ, Cordell GA (1993) Sesquiterpene lactones
from *Inula Britannica*. Phytochemistry 34(1):249–252
- Zhou P, Takaishi Y, Duan H, Chen B, Honda G, Itoh M, Takeda Y,
Kodzhimatov OK, Lee K-H (2000) Coumarins and bicoumarin from
Ferula sumbul: anti-HIV activity and inhibition of cytokine release.
Phytochemistry 53(6):689–697
- Zielinska-Jenczylik J, Sypula A, Budko E, Rzadkowska-Bodalska H
(1987) Interferonogenic and antiviral effect of extracts from *Erodium*
cicutarium. Arch Immunol Ther Exp 35(2):211–220
- Zielinska-Jenczylik J, Sypula A, Budko E, Rzadkowska-Bodalska H
(1988) Interferonogenic and antiviral effect of extracts from *Erodium*
cicutarium. II. Modulatory activity of *Erodium cicutarium* extracts.
Arch Immunol Ther Exp 36(5):527–536
- Zitterl-Eglseer K, Jurenitsch J, Korhammer S, Haslinger E, Sosa S,
Della Loggia R, Kubelka W, Franz C (1991) Sesquiterpenelactones
of *Achillea setacea* with antiphlogistic activity. Planta Med
57(5):444–446
- Ziyan L, Yongmei Z, Nan Z, Ning T, Baolin L (2007) Evaluation of the
anti-inflammatory activity of luteolin in experimental animal mod-
els. Planta Med 73(3):221–226
- Zolotnitskaya SY (1965) Lekarstvennuiye Rasteniya Florui Armenii T.
2 (Medicinal plants in the flora of Armenia), vol 2. Izdatelstvo
Akademii Nauk Arm. SSR, Yerevan (in Russian)
- Zotov EP (1975) Phytochemical study of *Hyssopus seravshanicus* and
ferganensis and production of medicinal forms. Synopsis of disserta-
tion for the degree of Ph.D. in pharmacology, Frunze-Tartu (in
Russian)
- Zotov EP, Kolodina LK, Khazanovich RL (1977) Antibacterial activi-
ties of preparations of *Hyssopus seravshanicus* and *ferganensis*.
In: Sovremennuiye Voprosy Mikrobiologii I Immunologii
Tuberkuloz (Current issues in the microbiology and immunology
of tuberculosis). Kyrgyz Medical Institute, Frunze, pp 101–108 (in
Russian)

Author Queries

Chapter No.: 0001543048

Queries	Details Required	Author's Response
AU1	Please cite the following references in text: Akhmedhodjaeva and Kurmukov (1975), Akhmedzhanova et al. (1975), Azizov et al. (1992), De Combarieu et al. (2003), Dzhumabaev et al. (1970), 'eFloras (2008), Fazly Bazzaz and Haririzadeh (2003), Flora of Kirghiz SSR, Flora of North America Editorial Committee (1993+), Flora of Uzbekistan, Gusinin (1962), Karimov et al. (1975, 1977a, b), Kurmukov (1956, 1967a, b, 2000), Kurmukov and Rasulova (1983), Kurmukov and Syrov (1977, 1980), Kurmukov a et al. (1991a), Li et al. (2007), Naidovich et al. (1976), Nasirov (1973), Norn and Kruse (2004), Ribina et al. (1982, 1983), Sultanov and Kurmukov (1965), Syrov et al. (1986), Tulyaganov (1976), Tursunova (1973), Ur–Rehman et al. (1991), Wu et al. (1977), Yan et al. (1997)”.	
AU2	Please provide conference location for Abaihanova et al. (2006).	
AU3	Please provide volume number for Aitbaeva (1972), Aizikov et al. (1984, 1986, 1991, 1993a), Akhmedhodjaeva and Kurmukov (1975), Aliev (1945), Amirov (1974), Azizov et al (1992), Blinova (1957) , Botirov et al. (2006), Conti (1962), Dozortseva (1958), Dzhakupova (1968), Dzhumabaev et al. (1970), Ermishina et al (1982), Evdokimova and Kurmukov (1972), Gorkin (1964), Idzumi and Conti (1962), Iskanderova and Sharipova (1992), Kononov and Nesterova (2003), Korolyuk et al. (2002), Kurmukov (1987, 2000), Kurmukov and Akhmedkhodzhaeva (1975), Kurmukov and Saidkasimov (1969), Kurmukov and Syrov (1976), Kurmukov et al. (1976, 1980, 1984b), Kurmukova (2000a, b), Kurmukova and Kurbanov (1999), Kurmukovа et al. (1997b), Lugmanova et al. (2007), Mansurov (1967), Mashkovskii (1953), Massagetov (1932), Nazrullaev et al. (1990), Norn and Kruse (2004), Prokhorova et al. (1992b), Rustamov et al. (1974), Sadritdinov (1968), Sarkisyants (1969a), Sarkisyants and Azizova (1971), Sinitsin (1959), Sultanov (1959a, b, 1960), Sultanova et al (1998), Syrov (1984), Syrov and Kurmukov (1975c, 1976c, 1977), Syrov et al. (1975a, b, 1976, 1986), Tashbaev and Sultanov (1962, 1965), Trutneva and Berezhinskaya (1960), Tsukervanik et al (1935), Tulyaganov and Sadritdinov (1968), Tursunova (1973).	
AU4	Please provide chapter title for Aliev et al. (1972), Khamdamov (1972).	
AU5	Please provide page range for Dzhangaliev et al (2003).	
AU6	Please provide publisher location for Ibragimov and Dzhumabaev (1971)	
AU7	Please provide complete bibliography details for Khanov et al. (1971).	
AU8	Please provide missing parenthesis in Sarkisyants (1972)	
AU9	Please check if updated publisher name and location to Stahl–Biskup (2002) is okay.	
AU10	Please provide article title for Ur–Rehman et al. (1991).	
AU11	Please check if updated volume number and page range to Zhou (2001) is okay.	

General Index

- A**
- Abortifacient, 17, 148
- Abrasion, 16
- Abscess, 42, 85, 107, 122, 137, 180, 205, 222, 237, 250, 261, 272
- Absinthin, 41
- Acanthophylloside, 32
- Acetic acid, 14, 17, 46, 145, 255
- Acetylcholine, 65, 107, 160, 201, 252
- Acetylcholinesterase, 118, 187, 272
- Acetylcapelline, 19
- Acevaltrate, 255
- Aches, 39, 54, 55, 62, 87, 99, 255
- Aconitic acid, 89, 104
- Acsinatine, 21
- [AU1] Adaptagen, 128
- [AU2] Adrenal gland,
- Adrenaline, 102, 160
- Adrenergic, 187, 262
- Aesculin, 236
- Aglycone, 11, 13
- Aimaline, 262
- Air sickness, 87
- Aksine, 21
- Akuamidine, 262
- Akuamine, 262
- Alantolactone, 143–145
- Alantone, 144
- Alantopicrine, 145
- Aldose reductase, 128, 223
- Alginidine, 153
- Alginine, 153
- Alhagidin, 28
- Alhagitin, 28
- Alimentary, 197
- Aliphatic alcohol, 165
- Alkamide, 43
- Alkanin, 39
- Allergen, allergy, 41, 64, 102, 191
- Allocriptopine, 185
- Allyl-isothiocyanate, 92
- Aloemodin, 203
- Aloperine, 203
- Alpha-amyrin, 225
- α -humulene, 201
- α -hydrojuglone, 147
- α -linolenic acid, 39, 198
- α -terpenyl-acetate, 140
- Alteramine, 241
- Altitude sickness, 100
- Amaranthin, 36
- Amenorrhea, 245, 253
- Amygdalin, 183
- Amyl alcohol, 245
- Amyrin, 178
- Anabasine, 257
- Anabolic activity, 27, 228
- Anabsinthin, 41
- Anacidic gastritis, anacidity, 62, 181, 246
- Anagirine, 241
- Anagyrene, 242, 243
- Analeptic, 11, 118, 130, 201, 213, 262
- Analgesic, 13, 19, 39, 59, 64, 72, 75, 78, 99, 120, 125, 135, 137, 149, 170, 171, 177, 187, 198, 220, 235, 240, 245, 248, 251–253, 256, 260, 271, 272
- Anaphylactic shock, 113
- Anasarca, 107
- Anchusa acid, 39
- Anchusin, 39
- Androgenic action, 27
- Anemia, 58, 108, 140, 171, 172, 175, 210, 212, 221, 228, 233, 236, 248, 271
- Anemonin, 69, 72
- Anesthesia, 89, 125, 153
- Anesthetic, 9, 78, 97, 153, 253, 271
- Anethole, 177 [AU3]
- Angelic acid, 114
- Angiocholitis, 235
- Angioprotector, 28, 44 [AU4]
- Anhydroaustricine, 44
- Anhydroperforine, 130
- Anisic acid, 114
- Anonaine, 213
- Anorexia, 69
- Antagonist, 21, 73, 130
- Anthocyan, 37, 138, 220, 245, 261
- Anthocyanidin, 88
- Anthocyanin, 88, 139, 166, 212
- Anthracene, 13
- Anthraglycoside, 13, 120, 127, 196, 208, 217
- Anthranoyllycoctonine, 90
- Anthraquinone, 119, 120, 198, 203, 217, 222
- Anthrax, 32, 42, 58, 92
- Anti-aggregant, 52 [AU5]
- Anti-amnesic, 223
- Antiarrhythmic, 11, 21, 22, 118, 125, 127, 238, 240, 262, 266
- Anti-asthmatic, 57, 102, 137, 216

- Antibacterial, 16, 21, 36, 39, 52, 62, 67, 68, 72, 92, 102, 138–140, 142, 156, 166, 167, 176, 177, 183, 213, 225, 231, 240, 245, 258, 261, 266, 267, 270–272
- Antibiotic, 73, 74, 182, 201, 205
- Anticarcinogenic, 236
- Anticonvulsive, 90, 170, 171, 238
- [AU6] Anti-cough,
- Antidiarrheal, 28, 55
- Antiedemic, 237
- Antiemetic, 65, 220, 237
- [AU7] Anti-fibrillant, 262
- Antifungal, 65, 220, 237
- Antigonadotropic, 165
- Antihelminthic, 43, 140, 267
- Antihistamine, 128
- Anti-HIV, 73, 115
- Antihypertensive, 48
- Antihypoxic, 13, 14, 28, 197
- Antihysterical, 94
- Anti-inflammatory, 10–14, 16–18, 28, 34, 44, 52, 57–59, 62, 70, 72, 73, 77, 85, 94, 97, 103, 104, 107, 117, 122, 124, 138, 140, 143, 147, 149, 161, 170, 173, 177, 181, 183, 191, 198, 202, 207, 212, 215, 220, 235, 237, 240, 248, 250–252, 258, 263, 272
- Antimicrobial, 18, 36, 43, 48, 65, 70, 73, 134, 139, 147, 173, 180, 181, 220, 226, 235, 240, 250, 252
- Antioxidant, 13, 14, 28, 52, 58, 62–64, 112, 135, 144, 171, 181, 183, 220, 232, 236, 250, 252
- Antiparasitic, 89
- Antiproliferative, 132, 252
- Antiprotist, 36, 140
- Antipyretic, 28, 54, 55, 58, 92, 166, 175, 183, 198, 202, 203, 210, 220, 235, 236, 240, 251, 265, 267
- Antisclerotic, 246
- Antiseptic, 57, 138, 181, 220, 225, 248
- Antispasmodic, 14, 46, 62, 77, 87, 94, 113, 127–129, 133, 137, 171, 181, 191, 225, 235, 247, 255, 262
- Antithyroid, 166, 207
- Antitumor, 19, 22, 54, 73, 171, 178, 211, 215, 222, 236, 272
- Antulcerogenic, 28, 135, 139, 215
- Antivenom, 72
- Aorta, 44, 203
- Apigenin, 48, 192, 225, 235, 245
- Apoptosis, apoptotic effect, 42, 128, 129, 135, 211, 236
- Appetite, 41–43, 48, 62, 67, 70, 73, 143, 148–150, 180, 181, 202, 203, 210, 218, 223, 235, 236, 250, 255, 260, 266
- Arabinose, 124
- Arachic acid, 92
- Arachidic acid, 225
- Arachno-encephalitis, 90
- Arachnoiditis, 90
- Arctigenin, 73
- Arctiin, 73
- Arctiopicrin, 178
- Argemone, 240
- Argentine, 241
- Arnidiol, 236, 250
- Arrhythmia, 13, 262, 266
- Artabasin, 41
- Arteannuin, 42
- Artelein, 44
- Artelin, 44
- Artemetin, 41
- Artemisinin, 42
- Arterial pressure, 12, 95, 97, 102, 125, 147, 153, 160, 161, 178, 225, 227, 238, 247, 255, 262, 268
- Arteriosclerosis, 58, 117, 147, 172, 216, 236, 248
- Arthralgia, 55
- Arthritis, 148, 150, 198, 256
- Arthrosis, 225
- Asaresin, 113
- Asaresinol, 113
- Asaresinotannol, 113
- Asarone, 88
- Ascariasis, 132
- Ascorbic acid, 29, 44, 48, 58, 127, 147, 155, 171, 181, 183, 208, 212, 216, 218, 226, 231, 236, 242, 265
- Asparagine, 17, 51, 248
- Asperuloside, 120, 217
- Asphyxia, 94, 195, 241, 242
- Asthenia, 13, 113, 159, 172, 176, 225
- Asthma, 39, 64, 77, 78, 87, 100, 102, 113, 127, 129, 137, 140, 178, 187, 191, 241, 248, 250, 258, 271
- Astringent, 14, 26, 70, 88, 103, 104, 107, 123, 133, 138, 140, 143, 182, 183, 197, 210, 215, 220, 221, 226, 246, 262
- Astringent substances, 16, 183
- Atonia, 73, 180, 181, 265
- Atropine, 13, 87, 137
- Aucubin, 192, 258
- Austricine, 11, 44
- Autoimmune, 143
- Autonomic ganglia, 75
- Avenasterol, 230
- Avicularin, 196
- Avitaminosis, 216
- B**
- Back pain, 55
- Bacteria, 14, 24, 34, 46, 65, 67, 114, 138, 181, 240, 272
- Bactericidal, 54, 72
- Bacteriostatic, 54, 191
- Baicalin, 192
- Bakuchiol, 205
- Barium, 171
- Bed bug, 201
- Bedsore, 216
- Behenic acid, 225
- Benzaldehyde, 73
- Benzoic acid, 14, 64, 145, 232
- Benzyl isothiocyanate, 92
- Berbamine, 54, 55
- Berbamunine, 54, 55
- Berberine, 54, 55
- Bergamotene, 88
- β -bisabolene, 88
- β -carbolone, 185
- β -hydrojuglone, 147
- Betaine, 34, 36
- β -sitosterin, 140, 236, 265
- Bicyclogermacrene, 201
- Bile, 12, 14, 44, 52, 177, 181, 187, 216, 235
- Bile duct, 47, 117, 137, 230
- Bile-stimulant, 52
- Bilirubin, 187
- Biogenic amine, 230
- Bitter substances, bitters, 16, 24, 73, 161, 178, 192, 198, 235, 240
- Bleeding, hemorrhaging, 16, 18, 33, 59, 65, 93, 94, 104, 107, 113, 115, 117, 118, 122, 132, 137, 143, 155, 166, 170, 175, 177, 195, 198, 215, 226, 232, 241, 248, 252
- Bloated, 130

- Blood, 64, 166, 187, 197, 253, 261
 circulation, 13, 57, 88, 97, 181, 237, 242, 266
 cleanser, 108, 118, 178
 coagulation, 12, 55, 64, 155–157, 167, 170, 196, 227, 252, 253, 255
 sugar, 70, 175, 206, 252
- Bone, 123, 143, 145, 150, 271
 fracture, 54, 59
 marrow, 252
- Borneol, 17, 48, 151, 188, 235, 255
- Bornyl acetate, 140, 188, 190
- Bornyl-isovalerianate, 255
- Bradycardia, 75, 87
- Britanin, 143
- Bronchiectasis, 181, 250
- Bronchitis, 32, 34, 87, 129, 145, 148, 150, 180, 181, 191, 193, 218, 237, 238, 242, 245, 250, 265
- Bronchospasm, 177
- [AU9] Bronchus,
 Bruellosis, 32, 144
 Bruise, 77, 165
 Bufadienolide, 13
 Burn, 14, 27, 90, 135, 201, 216, 225, 248, 257
 Bursic acid, 65
 Butyrylcholinesterase, 118, 187, 272
- C**
- Ca⁺⁺ channel, 46
- Cadinene, 148
- Caffeic acid, 37, 75, 77, 83, 142, 157, 171, 192, 221, 223, 245, 252, 265
- Caffeine, 13, 122, 130, 186, 213
- Caffeoylquinic acid, 245
- Calcium oxalate, 217, 221
- Callus, 97, 120
- Calm, calming, 85, 90. *See also* Relax
- Campesterin, 191
- Campesterol, 190, 230
- Camphene, 17, 140, 148, 151, 188, 190, 245, 255
- Camphorol, 67, 177
- Canavanine, 261
- Cancer, 19, 22, 42, 65, 73, 75, 95, 97, 103, 120, 128–130, 135, 145, 148, 170, 177, 178, 181, 198, 213, 221, 235, 236, 240, 246, 248, 252, 255, 265
- Caoutchouc, 110, 112, 236
- Capillary, capillary strengthening, 14, 94
- Capronic acid, 220
- Caprylic acid, 17
- Carabron, 144
- Carbohydrates, 14, 22, 31, 34, 59, 124, 127, 165, 166, 191, 192, 195, 210
- Carbolic acid, 114
- Carbuncle, 138, 170, 205, 253
- Carcinogenesis, 132
- Cardiac glycoside, 11, 13, 109, 166, 198
- Cardinolide, 13
- Cardiosclerosis, 83, 109
- Cardiotonic, 13, 55, 178, 266–268
- Carene, 62
- Carminative, 41, 43, 67, 105, 129, 233, 235, 255
- Carotene, 16, 34, 41, 43, 48, 52, 58, 69, 75, 77, 84, 87, 88, 92, 99, 104, 105, 107, 117, 124, 135, 138, 139, 142, 147, 155, 166, 170, 175, 192, 196, 212, 215, 216, 221, 225, 226, 231, 233, 248, 252, 261
- Carotenoid, 49, 88, 168, 216, 222, 250, 252, 265, 268
- Carotid nerve, 262
- Carvacrol, 62, 67, 168, 173, 177, 180, 181, 245, 267
- Carvone, 62, 67
- Caryophylladienol, 188
- Caryophyllene, 154, 181, 188
- Caryophyllene oxide, 24, 154, 188, 225
- Catalepsy, 118, 125, 240
- Cataract, 177
- Catechin, 28, 84, 210, 222, 231
- Catecholamine, 198
- Caulosapogenin glycoside, 72
- Cedrene, 150, 225
- Cedrol, 148–151
- Central nervous system, 54, 70, 74, 78, 94, 102, 137, 170, 178, 180, 187, 240, 262, 265
- Cerebral cortex, [AU10]
- Cerotic acid, 225, 236
- Cervical erosion, 28, 226
- Cervix, [AU11]
- Ceryl-alcohol, 138
- Chatinene, 255
- Chemical compound, 11, 13, 14, 113, 253
- Chicoric acid, 236
- Chlorogenic acid, 83, 157, 192, 236
- Chlorophyll, 225
- Cholagogue, choleric, 13, 28, 32, 41, 54, 57, 68, 74, 77, 94, 117, 132, 145, 148, 150, 167, 173, 180, 198, 215, 216, 230, 235, 236, 250, 252, 263
- Cholecystitis, 12, 13, 54, 62, 74, 87, 127, 132, 149, 180, 215, 216, 231
- Cholera, 115
- Choleresis, 235
- Choleric action, 13, 54, 148
- Choleric, cholagogue, 13, 28, 32, 41, 54, 57, 68, 74, 77, 94, 117, 132, 145, 148, 150, 167, 173, 180, 198, 215, 216, 230, 235, 236, 250, 252, 263
- Cholesterol, 44, 70, 130, 177, 191, 237, 252
- Choline, 65, 138, 166
- Cholinergic, 251
- Cholinesterase, [AU12]
- Chorea, 75
- Chromium, 171, 190
- Chromone, 13
- Chronotropic, 167, 177
- Chrysoeriol, 236, 241
- Chrysophanic acid, 208, 221
- Cichoriin, 70, 236
- Cicutine, conicine, 49, 75
- Cinaroside, 241
- Cincholic acid, 272
- Cineol, 48, 176, 188
- Cinnamaldehyde, 73
- Cinnamamide, 207
- Citral, 171
- Citric acid, 65, 183
- Citrin, 14
- Citronellol, 73, 171
- Citronellyl acetate, 115
- Clematine, 72
- Cnicin, 73
- Coagulation, 12, 55, 64, 155–157, 167, 170, 196, 227, 252, 255
- Coating (remedy), 14, 151, 258 [AU13]
- Cobalt, 271
- Codonopsin, 74
- Codonopsinin, 74
- Colitis, 13, 16, 28, 36, 62, 87, 124, 127, 138, 173, 183, 221, 230, 270
- Columbamine, 54, 55

- Common cold, 58, 83, 113, 132, 178, 181, 187, 190, 212, 215, 227, 237, 256, 265, 267
- Compound, 10–14, 18, 19, 23, 27, 29, 36, 60, 62, 67–69, 72, 73, 75, 78, 85, 87–89, 104, 113–115, 124, 132, 139, 142, 144, 150, 154, 159, 161, 165, 167, 168, 173, 182, 188, 191, 192, 196, 200, 201, 208, 210, 212, 215, 216, 223, 226, 228, 231, 232, 235, 236, 246, 250, 253, 255, 260, 267, 271, 272
- Compress, 97, 120, 170, 178, 180, 181, 208, 235, 245, 258, 261, 265
- Condolphine, 23, 89
- Congenital defect, 75
- Congestion, 73
- Conhydrine, 75
- Conjunctivitis, 119, 191, 220
- Constipation, 36, 54, 73, 87, 92, 117, 172, 208, 210, 217, 230
- Contraceptive, 14, 165, 205, 237
- Contractility, 54, 241
- Contraction, 57, 89, 92, 161, 203, 241, 252, 253, 262
- Convolamine, 78, 253
- Convalidine, 78
- Convolvine, 77, 78, 253
- Convolvuline, 77
- Convulsions, 85, 87, 94, 137, 196, 213, 255
- Coronary, 13, 88, 100, 166, 266
- Corydine, 21, 125
- Corytuberine, 125
- Cough, 28, 32, 39, 85, 108, 118, 128, 139, 159, 185, 195, 202, 216, 220, 240, 245, 248, 250, 257, 263, 271, 272
- Coumaric acid, 192, 252
- Coumarin, 10, 11, 13, 14, 21, 22, 28, 34, 42, 43, 47, 52, 58, 64, 70, 87, 88, 94, 95, 103, 113, 115, 119, 120, 132, 156, 166, 167, 170, 175, 177, 180, 196, 198, 201, 205, 211, 225, 227, 232, 236, 237, 248, 250, 267
- Coumarinic acid, 248
- Cryptopine, 238
- Crystalline, 154, 246
- Cuminaldehyde, 62, 73
- Cuminyl alcohol, 62
- Curare, 23, 75, 89, 213
- Cuscohygrine, 77
- Cuts, 6, 33, 77, 165
- Cyanidin, 84, 212, 215, 216
- Cyanidin-3-*O*-glucoside, 215
- Cyanin, 166
- Cyanogenic compound, 165
- Cyanogenic glycoside, 13, 261
- Cyasterone, 27
- Cyclamine, 37
- Cyclic alcohol, 92
- Cyclic peptide, 253
- Cyclitols, 165
- Cyclolignan, 148
- Cyclooxygenase (COX) enzyme, 88
- Cyclopropenoid fatty acid, 33
- Cymene, 62, 73, 168, 177, 267
- Cymol, 62
- Cystitis, 13, 34, 54, 72, 138, 143, 145, 180, 191, 198, 218
- Cytisine, 203, 241–243
- Cytosterin, 250
- Cytotoxicity, 54, 62, 120, 130, 139, 181
- D**
- Daucane esters, 114
- Daucane-type sesquiterpene, 114
- Daucene, 88
- Daucosterol, 211, 223
- Deafness, 119, 150
- Decompensation, 89
- Dehydroabietol, 190
- Dehydrothalicmine, 238
- Delatine, 89
- Delphinine, 89
- Delphinindin, 212
- Delphirine, 90
- Delsemine, 89, 90
- Delsine, 89
- Delsoline, 89
- Delsosine, 89 [AU14]
- Demulcent, 250, 258, 261
- Dental caries, 225 [AU15]
- Deoxyepanin, 187
- Depression [AU16]
- emotional, 69, 74
- physical/physiological, 90, 94, 240
- Dermatitis, 34
- Dermatosis, 147, 216, 265
- Desensitization, 64
- Detoxicant, detoxify, 94, 191, 210
- Diabetes, 43, 70, 113, 143, 145, 147, 196, 206, 212, 218, 252
- Diacylglyceride, 80, 92
- Diaphoretic, sudorific, 39, 58, 62, 72, 108, 117, 128, 153, 161, 180, 183, 187, 202, 212, 218, 220, 236, 250, 263, 265
- Diarrhea, 33, 69, 85, 92, 117, 123, 138, 145, 173, 183, 192, 196–198, 200, 210, 215, 216, 220, 221, 237, 238, 240, 251, 252, 258, 261, 262, 270, 271
- Diathesis, 58, 113, 212
- Dictamine, 94
- Didrovaltrate, 255
- Digestion, 57, 58, 62, 67, 70, 73, 135, 149, 171, 180, 202, 203, 218, 225, 251, 255, 265
- Digestive organs, 213
- Digestive system, 127
- Digestive tract, 85, 105, 236
- Dihydroalantolactone, 145
- Dihydroxyacids, 44
- Dillapiole, 62
- Dimethamine, 241
- Diosgenin, 246
- Diosmetin, 246
- Diosmine, 140
- Dipegene, 187
- Diphtheria, 203, 216
- Disaccharide, 82
- Disinfectant, disinfection, 62, 104, 138, 149, 250
- Disulfide, 113
- Diuresis, 26, 67, 103, 133, 159, 181, 196, 253, 268
- Diuretic, 13, 26, 28, 32, 36, 37, 39, 46, 58, 62, 67, 70, 72, 85, 88, 92, 97, 104, 108, 109, 113, 115, 117, 118, 120, 129, 132–134, 145, 151, 159, 163, 171, 172, 175, 177, 178, 183, 187, 191, 196, 198, 210, 212, 215–218, 225, 230, 246, 248, 250, 252, 258, 261, 263, 266, 268, 271
- Dizziness, 83, 171, 237, 238, 262
- DNA, 135, 257
- Dopamine, 198
- Doremol, 115
- Doremon, 115
- Drupacine, 205
- Drupanol, 205
- Drying fatty oil, 73, 125, 135, 178, 225, 246, 265
- 25-D-spirosta-3,5-diene, 246
- Dubamine, 94
- Dubinidine, 94, 130

- Duodenum, 175, 192, 193, 212, 218
 Dyeing substances, 120, 132, 207
 Dysentery, 28, 34, 65, 92, 99, 124, 175, 192, 198, 216, 218, 223, 265
 Dysmenorrhea, 165
 Dyspepsia, 41, 43, 73, 140, 165, 222
 Dyspnea, 84
 Dysuria, 33
- E**
 Ear, 18, 28, 77, 137, 170, 191, 236
 Ecdysone, 13
 Ecdysterone, 11, 27, 228
 Eczema, 31, 34, 42, 69, 72, 90, 117, 135, 145, 147, 151, 172, 181, 205, 216, 218, 253, 256, 260, 265
 Edema, 26, 37, 43, 59, 69, 104, 108, 109, 132, 133, 145, 150, 172, 185, 191, 212, 236–238, 250
 Eicosenoic acid, 162
 Elemol, 68
 Elephantitis, 192
 Ellagic acid, 124, 220, 221
 Emetic, 24, 73, 125, 262, 263
 Emodin, 208, 221
 Emollient, 14, 170
 Endarteritis, 260
 Endocytotic activity, 165
 Endogenous, 165
 Endomyocarditis, 266
 Enterocolitis, 34, 85, 191, 192, 218, 221, 235
 Enzyme, 27, 37, 142, 211, 221, 223, 246, 248
 Enzyme-stimulating, 142
 Ephedrine, 13, 100, 102, 213
 Epicatechin, 84
 Epigallocatechin, 28
 Epilepsy, 24, 46, 48, 57, 75, 130, 161, 180, 187, 193, 225, 237, 238, 240, 255, 258
 Epi-13-manool, 188
 Epi-rhododendrin, 210
 Epithelization, 69, 252
 Epoxyacylglyceride, 92
 Equisetin, 104
 Equisetonin, 104
 Equisetrine, 104
 Eremuran, 105
 Eremursine, 105
 Ergolide, 143
 Ergotism, 92
 Eriodictyol, 48, 215
 Erucic acid, 92
 Ervamine, 262
 Ervine, 262
 Ervinine, 11, 262
 Erysimine, 109
 Erysimoside, 109
 Erysipelas, 118, 220
 Erysipelatous inflammation, 63, 250
 Erythrocyte, 252
 Erythrodermia, 216
 Esculetin, 87
 Esculin, 87
 Esophagus, 135
 Essential oil, 2, 5, 16–18, 24, 28, 29, 34, 41–44, 46–48, 51, 57–60, 62, 65, 67, 68, 73, 75, 83, 85, 87, 88, 92, 94, 102, 108, 113–115, 117–120, 124, 127, 128, 132, 133, 137–140, 143–145, 147–151, 154–157, 159, 161–163, 166–168, 170, 171, 173, 175–177, 180–182, 186, 188, 190, 192, 193, 196, 201, 205, 207, 212, 215, 220, 221, 223, 225, 226, 230, 235, 245, 255, 258, 263, 265–268, 270, 271
 Estrogen, 253
 Ethyl ester, 92
 Eucalyptol, 18
 Eudesmine, 130
 Eugenol, 182
 Euphorbin, 110, 112
 Evodine, 130
 Evoxin (haploperin), 94, 130
 Evoxidine, 130
 Excelsine, 21
 Exophthalmic goiter, 64, 166
 Expectorant, 12, 13, 32, 34, 57, 67, 108, 113, 128, 134, 140, 145, 149, 151, 159, 162, 180, 181, 191–193, 195, 202, 220, 226, 241–243, 245, 248, 250, 252, 257, 258, 263, 272
 External bleeding, 18
 Extriol, 128
 Exudative diathesis, 113
 Eye, 78, 87, 119, 137, 177, 185, 192, 213, 246, 257, 258, 263, 271
- F**
 Facial paralysis, 177
 Faint, 57, 195
 Faradiol, 236, 250
 Farnesferol, 113
 Fatigue, 117, 272
 Fatty oil, 14, 22, 34, 36, 37, 51, 67, 73, 83, 87, 88, 109, 125, 135, 140, 147, 165, 170, 171, 176–178, 180, 191–193, 195, 202, 205, 207, 213, 215, 216, 220, 222, 225, 231, 236, 240, 248, 265, 267
 Febrifuge, 39, 92
 Fenchone, 73
 Fermononetin, 129
 Ferruginol, 223, 225
 Ferulic acid, 113, 142, 192
 Festering wounds, 150
 Fetal hypoxic hypotropia, 27
 Fever, 26, 41, 53, 55, 69, 73, 85, 87, 90, 97, 100, 118, 123, 128, 175, 187, 193, 198, 210, 213, 215, 216, 218, 223, 225, 228, 230, 231, 235, 238, 240, 245, 248, 250, 267, 271
 Fiber, 102, 210, 218
 Fibrinolytic, 177, 227
 Fibroblast, 256
 Fibrosis, 196
 Flavanone glycoside, [AU19]
 Flavolignan, 230
 Flavone, 14, 59, 100, 123, 221, 248
 Flavonoid, 13, 14, 16, 21, 22, 28, 34, 37, 41, 43, 47, 48, 52, 53, 57, 58, 64, 67, 68, 70, 77, 83, 88, 90, 93–95, 97, 102–104, 107, 117, 119, 120, 128, 129, 132, 133, 135, 137–140, 142, 143, 147, 151, 155–157, 161, 162, 165–167, 175–177, 181, 183, 190–192, 196–198, 203, 206, 211, 215, 216, 218, 220–222, 225, 227, 230–232, 235–237, 240, 241, 245, 246, 248, 250, 253, 258, 260, 261, 265–267, 271
 Flavonol glycoside, 26, 29, 175, 208
 Flavoxanthin, 236
 Flea, 201
 Flindersine, 130
 Foetidine, 237
 Folic acid, 135, 145
 Foliosidine, 130
 Food poisoning, 34
 Formic acid, 220, 252, 255
 Fragmentation, 135

[AU17]

[AU18]

[AU19]

Frangula-emodin, 138, 208
 Frangulin, 208
 Free radicals, 107, 135
 Frostbite, 135
 Fructose, 79, 165
 Fumaric acid, 118, 230
 Fumaridine, 118
 Fumvailline, 118
 Fungal skin disease, 49, 58, 77, 110, 196, 217, 218, 221, 265
 [AU20] Fungicide, 72, 267
 [AU21] Fungus, 5, 36, 65
 Furocoumarin, 14, 205
 Furuncle, 138, 154, 170, 178, 190, 191, 205, 225, 250, 251
 Furunculosis, 69, 120, 247, 265
 Fustin, 127

G

Galanthamine, 251
 Galiosin, 120, 217
 Gallbladder, 88, 210, 236
 Gallic acid, 110, 210, 271
 Gallocatechin, 28
 Gallotannin, 226
 Gallstone, 70, 138, 217
 γ -coniceine, 75
 γ -terpinene, 62
 Ganglion-blocking, 54, 240, 260, 262
 Gangrene, 250
 Gargle, 87, 220, 223
 Gastralgia, 173, 267
 Gastric disease, 17, 59, 82, 100, 122, 123, 159, 191
 Gastric fluid, 62
 Gastric pneumatosis, 115
 Gastritis, 13, 28, 34, 41, 43, 62, 65, 77, 85, 119, 128, 145, 149, 173, 180, 181, 192, 210, 212, 218, 246, 258, 266
 Gastroenteritis, 145
 Gastrointestinal tract, 48, 85, 113, 140, 145, 147, 165, 197, 215, 221, 235, 250, 255
 Gene expression, 70
 General tonic, 12, 80, 237
 Genistein, 241, 260
 Genistin, 241
 Genotoxic, 217
 Gentianadine, 122
 Gentianaine, 122
 Gentiananine, 122
 Gentianine, 95, 122
 Gentioflavine, 122
 Gentiobetaine, 122
 Gentsinic acid, 142
 Geraniol, 171
 Germacrene, 181, 201
 Gingivitis, 138, 151, 225, 226
 Gitogenin, 246
 Glabric acid, 128
 Gland, 14, 24, 26, 32, 38, 128–130, 138, 139, 143, 148, 150, 151, 154, 165, 166, 180, 205, 212, 233, 236, 237, 251
 Glaucine, 55
 Glaunidine, 21
 Glaunine, 125
 Glauvine, 125
 Glucobarbarin, 207
 Glucocapparin, 207
 Glucofrangulin, 208
 Glucofructose, 165

Glucose, 34, 79, 95, 105, 124, 147, 165, 177, 265
 Glucoside, 37, 59, 92, 97, 104, 215, 236, 246, 260
 Glycerin, 14
 Glycoalkaloid, 13, 24
 Glycone, 13
 Glycoperine, 130
 Glycoside, 11–13, 70, 72, 85, 92, 128, 147, 192, 207, 211, 217, 226, 253, 265
 Glycyrramarin, 128
 Glycyrrhetic acid, 128, 129
 Glycyrrhizic acid, 129
 Glycyrrhizin, 128
 Goiter, 64, 145, 166, 265
 Gonorrhoea, 72, 108, 118, 198, 246
 Gossypol, 134
 Gout, 57, 72, 73, 85, 117, 137, 149, 171, 217, 235, 258
 Gram-negative bacteria, 65, 67, 181
 Gram-positive bacteria, 114, 138, 240
 Granilin, 144
 Granulation, 69
 Guaiol, 190
 Guinea worm, 36
 Gum, 10, 14, 16, 64, 113, 122, 132, 145, 147, 187, 202, 215, 246
 Gynecological disorders, 37

H

Hair growth, 27, 85, 196, 252
 Haplofidine, 130
 Haplofilidine, 130
 Haplophytin, 130
 Haplopine, 130
 Harmaline, 187
 Harmalol, 187
 Harman, 246
 Harmine, 187, 272
 Harpagide, 258
 Harpagoside, 258
 Headache, 19, 22, 36, 54, 63, 73, 75, 87, 108, 118, 124, 149, 150, 162, 177, 187, 191, 225, 235, 237, 242, 245, 246, 255, 262, 268
 Heart, 54, 55, 65, 70, 83, 84, 87, 89, 93, 99, 100, 102–104, 108, 109, 118–120, 125, 138, 139, 145, 161, 166, 167, 170, 175, 177, 195, 198, 201, 203, 212, 215, 223, 225, 235, 238, 251, 255, 258, 262, 263, 267, 272
 Heart failure, 255
 Hederagenin, 95
 Helminthosis, 92
 Hematuria, 200
 Hemicellulose, 80, 210
 Hemodynamics, 100
 Hemoglobin, 211, 252
 Hemolytic index, 32, 37, 39, 160, 232, 246, 253, 257
 Hemoptysis, 31, 36, 57, 123, 138, 226, 230
 Hemorrhagic shock, 105
 Hemorrhaging, bleeding, 16, 18, 33, 59, 65, 93, 94, 104, 107, 113, 115, 117, 118, 122, 132, 137, 143, 155, 166, 170, 175, 177, 195, 198, 215, 226, 232, 241, 248, 252
 Hemorrhoidal hemorrhage, 16
 Hemorrhoids, 17, 18, 26, 41, 49, 64, 87, 97, 117, 124, 143, 178, 208, 210, 220, 226, 230, 252, 258, 261
 Hemostatic, 13, 14, 16, 26, 33, 36, 39, 57, 59, 62, 92, 103, 104, 107, 113, 115, 117, 120, 123, 125, 132, 137, 138, 142, 143, 145, 155–157, 166, 168, 173, 175, 178, 191–193, 195, 196, 198, 215, 220, 221, 226, 232, 240, 252, 261, 266
 Hepatitis, 12, 14, 26, 34, 44, 64, 68, 74, 77, 94, 155, 172, 198, 210, 215, 216, 230, 231, 235, 240, 261

- [AU22] Hepatocholecystitis, 132
 [AU22] Hepatoprotector, 12, 88, 95, 104, 122
 [AU23] Hepatotoxic, 64, 217
 Hernia, 52, 133
 Herniarine, 133
 Herpes, 37, 107, 175, 212
 Herpes simplex type 1, 175
 Hesperidin, 14, 258
 Heterocyclic, 14
 Hexadecanoic acid (palmitic acid), 82, 130, 134, 168
 Hexynyl disulfide, 113
 Highmoritis, 138
 Hippeastrine, 251
 Hippocampus, 21
 Histamine, 44, 177, 230, 252
 Hoarseness, 63
 Homeopathy, 37, 92
 Homothermopsine, 242
 Hordenine, 105, 251
 Humulene, 188, 201
 Hydrocyanic acid, 49, 261
 Hydrophobia, 63
 Hydroxycinnamic acid, 157
 Hydroxytryptamine, 44, 252
 Hygrine, 77
 Hyoscyamine, 87, 137
 Hyperglycemia, 43
 Hyperhydrosis, 140
 Hypericin, 138
 Hyperoside, 83, 103, 138, 147, 175, 221
 Hyperpituitarism, 165
 Hypertension, 12, 13, 57, 83, 84, 109, 117, 145, 161, 166, 170, 180, 232, 237, 248, 260
 Hypertensive, 11, 13, 125, 133, 160, 246, 271
 Hyperthyroidism, 166
 Hypertrophy, 166
 Hypoacidic gastritis, 149
 Hypochondria, 73
 [AU24] Hypogastritis,
 Hypoglycemic, 117, 122, 147, 206
 Hypolipidemic, 11, 14, 44, 63, 113, 180, 206
 Hypotension, 92, 203
 Hypotensive, 13, 46, 74, 77, 90, 113, 125, 127, 142, 156, 157, 160, 166, 171, 210, 221, 232, 240, 256, 266–268, 271
 Hypothermic, 94, 122, 251
 Hypotonia, 240
 Hypoxia, 12, 80, 95, 211, 227, 247, 266
 [AU25] Hypsogenin,
 Hyssopin, 65
 Hysteria, 57, 113, 161, 255
- I**
 Idiopathic skin atrophy, 260
 Immunological, 195
 Immunosuppression, 248
 Immunotoxicity, 70
 Imperatorin, 201
 Implantation, 148
 Impotence, 85
 Incanine, 247
 Infected, 138, 181, 247, 248
 Infection, 26, 140, 170, 173, 196, 198, 215, 223, 227, 245, 253, 271
 Infectious diseases, 212, 241
 Inflammation, 16, 17, 28, 34, 44, 48, 57, 63, 70, 85, 104, 117, 133, 140, 145, 178, 180, 187, 192, 230, 236, 246, 248, 250, 257, 265
- Influenza, 103, 212
 Inhibition, 14, 88, 118, 135, 142, 170, 175, 207, 223, 251, 271
 Inorganic, 13
 Inosine, 65
 Inotropic action, 102
 Insecticide, 90, 235
 Insomnia, 13, 41, 83, 94, 124, 171, 181, 255
 Insulin, 43, 252, 265
 Internal, 1, 18, 24, 47, 58, 92, 99, 107, 198, 245, 252, 255
 Interoside, 186
 Intestinal colic, 36
 Intestinal disorder, 89, 183, 262
 Intestines, 54, 70, 118, 127, 181, 208, 251, 271
 Intibin, 70
 Intoxication, 235, 241
 Intracellular, 27
 Intracranial, 177
 Intravenous injection, 90, 155, 238
 Inulin, 70, 143–145, 186, 222, 236, 250
 Iodine, 166, 201, 233, 265, 271
 Iridoid glucoside, 258
 Iridoids, 57, 97, 119, 120, 176, 191, 255
 Iron, 117, 155, 171, 190, 233, 271
 Isoalantolactone, 143–145
 Isobaldine, 89
 Isobetanin, 36
 Isocorydine, 54, 55
 Isoflavan, 232
 Isoleontine, 160
 Isoliquiritigenin, 128, 129
 Isomenthone, 266, 268
 Isopsoralen, 205
 Isoquercitrin, 107, 215
 Isoremerin, 213
 Isoreserpiline, 262
 Isorhamnetin, 67, 90, 92, 133, 135
 Isorubijervine, 256
 Isosalipurposide, 132
 Isotalatizine, 23
 Isotanshinone, 225
 Isotetrandrine, 54
 Isotriofolin, 248
 Isovalerianic acid, 138
 Isovaleric acid, 14
 Isovaltrate, 255
 Itch, 72, 94, 118
- J**
 Jalapine, 77
 Jatrorrhizine, 54
 Jaundice, 36, 55, 57, 73, 94, 113, 115, 117, 118, 132, 133, 139, 143, 145, 148, 150, 172, 217, 230, 236–238, 265
 Jervine, 256
 Joint, 34, 85, 145, 217, 235, 252, 256
 Juglone, 147
 Juniperin, 148
- K**
 Kaempferol, 33, 37, 75, 77, 92, 104, 107, 110, 135, 215, 216, 260
 Karakoline, 19
 Karasamine, 19
 Ketone, 17, 47
 Ketose, 124
- [AU26]

- Kidney, 28, 32–34, 47, 52, 57, 64, 70, 73, 87–89, 104, 110, 113, 115, 120, 125, 129, 133, 134, 138, 148, 151, 165, 170, 180, 192, 198, 210, 213, 215, 217, 230, 232, 237, 238, 241, 246, 250, 258, 260, 271
- Kidney stone, 34, 46, 48, 54, 62, 85, 117, 132, 148, 150, 151, 165, 177, 212, 217, 218, 251, 258, 265
- Koproporphyrin, 252
- Korseveramine, 153
- Korseveridine, 153
- Korseverinine, 153
- Kusunokinin, 130
- L**
- Lactation, 166
- Lactic acid, 218
- Lactogenic, 140
- Lactone, 11, 13, 16, 17, 24, 41, 42, 44, 58, 70, 73, 85, 113, 115, 143–145, 178, 236
- Lactose, 95
- Lactucin, 70
- Lactucopicrin, 70
- Lagochilin, 155–157
- Lappaconidine, 21
- Lappaconitine, 11, 13, 21
- Laryngitis, 124, 138, 180, 200, 250
- Larynx, 147
- Laxative, 28, 32, 36, 39, 46, 57, 67, 77, 83, 92, 109, 120, 125, 128, 129, 135, 147, 161, 202, 208, 212, 217, 218, 236, 252, 263
- Leishmaniasis, 188
- Leontamine, 160
- Leontidine, 160
- Leontine, 160
- Lepidoside, 162
- Lethargic encephalitis, 187
- Leucoanthocyanide, 222
- Leucoanthocyanidin, 84
- Leucocytes, 54, 170
- Leucodelphinidin, 28
- Leucomisine, 11, 17, 44
- [AU27] Leucomycin, 11, 17, 44
- Leucopenia, 170
- Leukemia, 175, 188, 232
- [AU28] Leukorrhea, 44
- Levorotatory, 44
- Libido, 10, 120, 148, 246, 248
- Licoctonine, 90
- Licorine, 251
- Lignan, 13, 41, 73, 130, 232
- Lignin, 10
- Lignoceric acid, 87
- Limonene, 62, 67, 140, 154, 255, 265, 267
- Limonoid, 94
- Linalool, 62, 140, 225
- Linalyl acetate, 115, 225
- Lindleyin, 210
- Linoleic acid, 64, 134
- Linolenic acid, 36, 39, 82, 140, 198
- Lipid, 33, 44, 49, 52, 80, 82, 92, 93, 95, 99, 127, 156, 170, 180, 220, 252, 268
- Liquirazide, 128
- Liquiritin, 128
- Liquitigenin, 128
- Liriodenine, 213
- Lithospermic acid, 223
- Liver, 28, 32, 41, 47, 52, 54, 62, 73, 77, 88, 89, 113, 115, 120, 132, 139, 148, 150, 180, 181, 191, 192, 196, 198, 210, 212, 215, 217, 218, 228, 230, 235–238, 252, 257, 258, 265, 271
- Local anesthesia, 153, 253 [AU29]
- Longifolin, 177
- Loroglossine, 85
- Lotaustralin, 211
- Lotion, 48, 63, 178, 181, 258
- Low stomach acidity, 41, 212
- Lucidin, 217
- Lungs, 250, 271
- Lupane, 223
- Lupanine, 160
- Lupeol, 178
- Lupus, [AU30]
- Lutein, 236
- Luteolin, 59, 104, 207, 225, 235, 236, 241, 245
- Luteolin 7-glucoside, 59, 236
- Luteolin 7-rutinoside, 59
- Lycopene, 49, 215
- Lymphadenectomy, 170
- Lymphadenitis, 113, 147
- Lymphedema, 170
- Lymph nodes, 271
- Lymphoblastoid, 120, 221
- M**
- Macroelement, 117, 210, 226, 271
- Macrophage, 210
- Magnoflorine, 54, 55, 237, 238
- Malaria, 23, 24, 41, 42, 100, 122, 145, 163, 187, 188, 191, 192, 210, 238, 240, 246, 248, 250
- Maleic acid, 165
- Malic acid, 14, 46, 65, 104, 117, 165, 166, 183, 202, 210–212, 216–218, 220, 255
- Malignant, 10, 22, 65, 119, 123, 162, 177
- Mammalian, 73, 217, 257
- Manool, 188, 225
- Marubiin, 167
- Mastitis, 138
- Matricarin, 44
- Matrine, 203, 260
- Measles, 92
- Mecambrolone, 213
- Mediator, 44
- Medulla oblongata, 262
- Melancholy, 68
- Melilotic acid, 170
- Melilotin, 170
- Melilotocide, 170
- Melissic acid, 72
- Menopause, 170, 218, 252
- Menorrhagia, 200
- Menstruation, 10, 46, 48, 145, 160, 163, 248, 265
- Menthol, 173, 268, 270
- Menthone, 173, 267, 268
- Mesaconitine, 21
- Metabolism, 13, 14, 32, 34, 57, 58, 95, 104, 135, 166, 196, 197, 215, 220, 250, 252
- Methicillin, 114, 139, 183
- Methoxy-cinnaric aldehyde, 43
- 2-Methoxy-1,4 naphthoquinone, 142
- Methyl-chavicol, 43
- Methyl-coniine, 75
- Methylcytidine, 160

Methylcytosine, 241–243
 Methyl-evoxin, 130
 Methyl gallate, 215
 Methyl lachnophyllate, 154
 Methyllycaconitine, 90
 Methylquercetin, 258
 Metropathy, 16
 Metrorrhagia, 107
 Microelements, 13, 29, 117, 190, 193, 255
 Midbrain, 262
 Migraine, 75, 170, 171, 255
 Mineral salts, 29, 85, 177, 252
 Mitral failure, 109
 Mollugin, 217
 Molluscicidal, 37
 Molybdenum, 171
 Monoacylglyceride, 92
 Monocaffeoyltartaric acid, 236
 Monohydroxyacid, 44
 Monosaccharide, 79, 82
 Monoterpene, 62, 73
 Monticamine, 19
 Morphine, 13, 102, 125
 Mouse, 132, 191, 210, 236
 Mouth wash, 26, 151, 167, 172, 187, 201, 215, 225, 262
 Mucilage, 14, 33, 34, 58, 73, 85, 127, 139, 145, 192, 196, 230, 242, 250, 258
 Mucous membrane, 78, 180
 Mucus, 117, 233
 Multiple sclerosis, 89
 Muscle, 57, 78, 84, 89, 90, 122, 125
 Musculoskeletal, 59
 Mussaenoside, 98
 Mustard essential oil, 92, 162, 207
 Myasthenia, 251
 Mydriasis, 75, 87
 Mydriatic, 137
 Myocardial infarction, 11, 27, 268
 Myocarditis, 44, 266, 268
 Myodystrophy, 175
 Myometrium, 203
 Myopathy, 195, 260
 Myrcene, 43, 88, 140, 149, 150, 154, 171, 190, 225
 Myricitrin, 271
 Myricyl alcohol, 72
 Myristic acid,
 Myristicin, 62, 135
 Myrtenol, 182, 255
 Myrtenyl isovalerianate, 255

N

Napelline, 19
 Naphthoquinone, 142, 165
 Narcosis, 94, 118, 130
 Narcotic, 9, 32, 130, 170, 241, 251
 Narcotized, 52, 90, 100, 105, 153, 238
 Naringenin chalcone, 132
 Narwedine, 251
 Nausea, 94, 266
 N-dimethyl colletine, 21
 Necrosis, 24, 27, 75, 77, 226
 Neoline, 19
 Neosophoramine, 203
 Neoxanthine, 99
 Nepetalactone, 176

Nephritis, 192, 198
 Nepodin, 221
 Nerolidol, 225
 Neural, 24
 Neuralgia, 19, 87, 256, 258
 Neurasthenia, 39, 48, 55, 68, 113, 187, 223, 270
 Neuritis, 22, 245, 251
 Neurodermatitis, 145
 Neuromuscular, 75
 Neuron, 24
 Neuroprotective, 250
 Neurosis, 13, 83, 113, 130, 140, 186, 223, 255, 258, 266
 N-heptacosane, 250
 Nicotine, 13, 104, 203
 Nicotinic acid, 218
 Nitrogenous compounds, 36, 228
 Noradrenaline, 102, 198, 255
 Norcorydine, 125
 Nortropine, 78
 Nose, 16, 28, 62, 237, 238
 N-oxy-benzoic acid, 142

O

Obesity, 62, 195
 Oblongine, 55
 Obstetric-gynecological, 57
 Ocimene, 225
 Octylene, 17
 Ointment, 14, 39, 42, 59, 69, 97, 145, 147, 151, 162, 166, 225, 253, 255, 272
 Oleanane, 223
 Oleanolic acid, 83, 130, 140, 192, 211, 227
 Oleic acid, 134, 147, 230
 Oleoresin, 190
 Oligosaccharide, 79
 Oliveramine, 122
 Oliveridine, 122
 Oliverine, 122
 Olmelin, 127
 Omega-3 fatty acid, 198
 Onopordopicrin, 178
 Organic acid, 14, 18, 19, 22, 41, 46, 54, 57, 70, 95, 104, 107, 117, 124, 135, 139, 148, 155–157, 165, 166, 172, 175, 182, 186, 191–193, 202, 208, 210–213, 216–218, 220, 222, 227, 237, 240, 241, 250, 252, 266
 Osteitis, 113
 Osteoarthritis, 215
 Osteomyelitis, 150, 225
 Osthol, 201
 Otitis, 138, 166
 Otolaryngology, 138
 Oxalic acid, 198
 Oxycanthine, 54, 55
 Oxymatrin, 203
 Oxypeucedanin, 201
 Oxyphosphocarpine, 203
 Oxysteroid, 27
 Oxytanshinone, 225

P

Pachycarpine, 160, 241, 242, 260
 Palmatine, 54, 55
 Palmitic acid (hexadecanoic acid), 82, 130, 134, 168
 Pancratine, 251

- Pancreas, 41, 87
 Pantotenic acid, 88, 252
 Papilloma, 132
 Parodontosis, 124
 Paraffin, 73
 Paralysis, 85, 89, 94, 118, 177, 217, 265
 Paralytic, 187
 Paraoxycoumarin, 113
 Parasite, 188, 286
 [AU32] Parasympathetic ganglions, 160
 Paratyphoid, 231
 Parenteral, 59
 Parfumine, 118
 Parinaric acid, 142
 Parishin, 44
 Parkinson's disease, 78, 89, 90, 187
 Paronychia, 138, 225
 Parturifacient, 165
 Patchouli alcohol, 190
 Pathogenic, 14, 42, 213
 Patrinoside, 186
 Pectic substances, 49, 79, 80, 151
 Pectins, 14, 117, 124, 202, 208, 212, 215, 216, 218, 220
 Pediculosis (lice infestation), 89
 Pegamine, 187
 Peganidine, 187
 Peganine, 11, 187
 Peganol, 187
 Pelargonin, 166
 Pellagra, 222
 Penicillin, 138
 Pentosan, 148, 215
 Peonidin, 216
 Pepsin, 62
 Pepsinogen, 62
 Perfamine, 130
 Perforine, 10, 130
 Periodontal disease, 34
 Periodontitis, 225
 Perioritis, 225
 Peripheral, 43, 65, 102, 203, 235, 242, 260
 Peristalsis, 203, 251
 Peritoneal, 198, 210
 Permeability, 14, 44, 170, 247
 Pertussis, 75, 90, 108, 115, 173, 178, 191, 192, 245, 248, 265
 Phagocyte, 170
 Phagocytic, 54
 Pharmacological, 3, 10, 11, 13, 87, 159, 163, 186, 213, 268, 271
 Pharyngitis, 100, 138
 Phenol, 13, 17, 29, 47, 123, 167, 180, 181, 211, 212
 Phenolcarbonic acid, 57, 83, 95, 103, 140, 191, 211, 222, 227, 231, 245
 Phenolic acid, 34, 107, 253, 265
 Phenolic glucoside, 97
 Phenyl- β -naphthylamine, 19, 207
 Phenylbutanoid, 210
 Phenyl glycoside, 271
 Phenylpropanoid, 62
 Phloridzin, 215
 Phloroglucinol, 221
 Phospholipid, 34, 118, 168, 220
 Phosphoric acid, 117
 Phthisis, 144
 Phyllalbine, 78
 Phytoecdysone, 27
 Phytoecdysteroid, 11, 27, 93, 228
 Phytoestrogen, 14, 73, 113
 Phytol, 166
 Phytoncid, 16, 31, 65, 140, 226, 252
 Phytosterin, 115, 250
 Pigments, 13, 28, 64, 90, 99, 100, 102, 118, 145, 147–150, 215, 225, 226, 256, 260, 271
 Pilocarpine, 233
 Pimple, 118, 265
 Pinene, 17, 62, 88, 114, 140, 148–151, 154, 188, 201, 212, 235, 245, 255, 268
 Pinocamphone, 140
 Pinworm, 235, 267
 Piperitone, 68, 173
 Plague, pestilence, [AU33]
 Plantagonine, 257
 Plasma, 147, 155, 166, 227, 265
 Platelet activating factor (PAF), 73
 Pleurisy, 104
 Pneumonia, 34, 102, 119, 120, 129, 149, 180, 195, 218, 242, 245, 250, 258
 Podophyllotoxin, 148
 Poison, 10, 75
 Poisonous, 24, 37, 49, 72, 75, 87, 94, 99, 112, 247, 253
 Poliomyelitis, 251
 Poliovirus, 37, 221
 Polyarthritis, 210, 225
 Polyenes, 67
 Polyp, [AU34]
 Polyphenol, 14, 107, 110, 135, 147, 166
 Polysaccharide, 14, 32, 59, 79, 80, 83, 85, 105, 167, 258
 Polyvitamin, 135, 216, 233
 Postencephalitic, 90
 Potassium, 171, 192, 227
 Poulitice, 187, 237, 250, 261
 Prangenidin, 201
 Prangenin, 201
 Prangosine, 201
 Pressor action, 100, 252
 Primveraza, 37
 Proanthocyanidin, 11, 13, 14, 28, 124, 197, 212
 Proazulen, 144
 Proliferation, 128, 135
 Propenyl isothiocyanate, 92
 Prophylactic, 12, 178, 266
 Propionic acid, 46
 Prostaglandin, 44
 Prostate, 34, 128, 252
 Prostatitis, 34
 Protein, 10, 62, 85, 147, 170, 175, 197, 198, 228, 233, 261
 Prothrombin, 157, 227
 Protocatechin, 191
 Protopine, 118, 125, 185
 Protoporphyrin, 252
 Protopseudohypericin, 138
 Prulaurasin, 183
 Prunasin, 183
 Prussic acid, 13, 162, 183, 261
 Pseudoconhydrine, 75
 Pseudoephedrine, 100, 102, 213
 Pseudohypericin, 138
 Pseudojervine, 256
 Pseudotaraxasterol, 236
 Pseudotropine, 77
 Psoralen, 205
 Psoriasis, 31, 34, 218, 225, 253
 Psychiatry, 137
 Psychomimetic, 187

Psycho-neurological diseases, 89
 Pulegone, 173, 266–268, 270
 Pulpitis, 54, 225
 Purgative, 13, 14, 88, 94, 98, 112, 128, 208, 221, 246
 Purine derivatives, 170
 Purpurin, 217
 Pus, 137, 170, 192, 251
 Pyelitis, 218
 Pyoderma, 69, 105
 Pyrocatechin, 123, 197, 210
 Pyrogallol, 123, 210
 Pyrrolidine, 77, 88
 Pyrrolidine alkaloid, 77

Q

Quercetin, 14, 28, 37, 64, 67, 75, 77, 83, 90, 92, 104, 107, 110, 133, 135, 138, 147, 162, 175, 177, 203, 212, 215, 216, 235, 252, 258, 260
 Quercetin arabinoside, 133
 Quercetin-3-arabinoside, 147
 Quercetin galactoside, 133
 Quercetin-3-galactoside, 110
 Quercetin triglycoside, 133
 Quercetrin, 138
 Quinidine, 13, 118, 125, 240
 Quinone, 177, 225
 Quinovic acid, 272

R

Rabies, 37
 Radiculitis, 19, 22, 46, 54, 87, 145, 215, 245, 251
 Ranunculin, 69
 Rash, 26, 102, 118, 175, 196, 213, 220, 221
 Recalcification, 155
 Receptor, 21, 87, 100, 250
 Rectal prolapse, 261
 Reflex excitability, 118
 Relax, 102, 255. *See also* Calm
 Remedy, 10, 28, 31, 39, 55, 72, 73, 77, 104, 107, 115, 129, 132, 135, 143, 151, 163, 171, 180, 183, 216, 218, 220, 236, 246, 255
 Remrefidine, 213
 Remrefine, 213
 Renal pelvis, 133
 Reoxygenation, 211
 Reserpine, 13
 Reserpinine, 262
 Resistance, 3, 12, 27, 139, 227
 Respiration, 94, 160, 201, 225, 241, 262
 Respiratory, 11, 46, 57, 58, 73, 75, 82, 85, 87., 90, 97, 128, 134, 140, 143, 145, 151, 167, 173, 177, 178, 180, 192, 201–203, 212, 216, 225–227, 241–243, 245, 248, 250, 258, 266
 Respiratory disease, 46, 57, 58, 145, 151, 202, 212, 225, 258
 Restorative, 9, 11, 27, 113, 144, 203, 228
 Reticular, 262
 Reticuline, 54
 Rhabdomyolysis, 75
 Rhamnoglucoside, 65
 Rhamnoglycoside, 133
 Rheumacarditis, 266
 Rheumatic pain, 122
 Rheumatism, 19, 22, 23, 26, 42, 46, 54, 55, 57, 72, 84, 87, 94, 95, 97, 99, 100, 104, 120, 124, 135, 137, 140, 143, 145, 148–150, 187, 190, 203, 208, 212, 215, 235, 237, 248, 252, 256, 265, 267, 272

Rhinitis, 100, 102, 138
 Rhododendrol, 210
 Ricketts, 147, 180, 217, 233, 258
 Roemeridine, 185
 Roemerine, 213
 Rosmarinic acid, 166, 223
 Royleanone, 223
 Rubella, 248
 Ruberythric acid, 217
 Rubiadin, 120, 217
 Rubiadine,
 Rubijervine, 256
 Ruscogenin, 246
 Rutin, 14, 28, 64, 107, 133, 137, 138, 175, 206, 218, 221, 237, 250
 Rutinoside, 59, 103

S

Sabinene, 43, 149, 212, 245
 Salicylic acid, 248
 Salidroside, 211
 Salivation, 75
 Salvicin, 206
 Salvicinin, 206
 Salvicinolate, 206
 Salvicinolin, 206
 Salvifolin, 206
 Salvin, 206
 Sambulene, 115
 Sanguinarine, 125
 Santolina alcohol, 17
 Sapogenin, 29, 51, 72
 Saponarin, 253
 Saponin, 5, 13, 21, 24, 26, 29, 32, 37, 39, 49, 51, 52, 59, 64, 65, 83, 85, 93, 95, 97–99, 103, 104, 108, 119, 123, 124, 127, 133, 139, 144, 160, 161, 166, 167, 176, 177, 186, 192, 193, 195, 196, 198, 225–227, 230, 232, 240, 242, 246, 250, 253, 257, 258, 265–267, 271
 Saporubin, 253
 Saporubinic acid, 253
 Scabies, 42, 58, 77, 90, 100, 110, 145, 187, 188, 201, 221, 247, 256, 260
 Scarlet fever, 175, 216
 Sciatic nerve, 145, 187, 217, 262
 Sclareol, 225
 Scleroderma, 260
 Scopolamine, 87, 102, 137
 Scopoletin, 42, 87
 Scratch, 16
 Scrofula, 58, 64, 108, 120, 147, 178, 180, 250, 258, 265, 267
 Scurvy, 43, 58, 135, 148, 150, 190, 215, 216, 221, 226
 Scutellarin, 192
 Sea sickness, 87
 Secretion, 62, 87, 165, 166, 177, 187, 235, 251, 252
 Secretory activity, 32, 113
 Secretory function, 62
 Sedative, 9, 10, 12, 13, 39, 52, 57, 67, 70, 72, 75, 83, 84, 87, 94, 107–109, 120, 122, 130, 142, 155–157, 161, 166, 167, 171, 180, 186, 192, 193, 213, 218, 227, 238, 240, 251, 255, 262, 265, 271
 Seizure, 63, 255
 Selenium, 171
 Semi-drying oil, 64, 159
 Sensory, 24, 118, 251
 Septicemia, 54
 Serotonin, 44, 255

[AU36]

- Sesquiterpene, 10, 11, 14, 16–18, 24, 41, 43, 70, 73, 114, 115, 138, 143–145, 149, 150, 178, 181, 225, 236, 250, 255
- Sesquiterpene alcohol, 10, 11, 14
- Sesquiterpene lactone, 16, 17, 24, 41, 70, 73, 115, 143–145, 178, 236
- Shepherin, 65
- Shortness of breath, 29, 37, 171, 248
- Sialorrhea, 33
- Silicic acid, 104, 196
- Silicon, 171
- Silicristin, 230
- Silimarin, 230
- [AU37] Silybin, 230
- Sinapic acid, 92
- Sinigrin, 92
- Sinus, 48, 89, 100, 250, 263
- Sinus cold, 48, 250, 263
- Skimmianine, 10, 94, 130
- Skin, 14, 23, 31–34, 41, 42, 49, 58, 60, 69, 77, 94, 97, 104, 110, 113, 117, 120, 128, 130, 132, 135, 142, 147–151, 178, 181, 187, 191, 192, 196, 201, 205, 212, 213, 215–218, 221, 226, 235, 236, 238, 240, 247, 248, 250, 253, 256, 260, 261, 265, 271
- Skin ulcer, 23, 77, 117, 191, 192, 196, 215, 216, 226, 248
- Sleep, 10, 94, 125, 255
- Smallpox, 92, 213, 271
- Smirnovine, 52
- Smooth muscles, 11, 54, 67, 87, 102, 127, 238
- Snake bites, 49, 72, 97, 191
- Snake venom, 124
- Sogdisterone, 228
- Songorine, 19, 21
- Sophocarpine, 203, 260
- Sophoramine, 203, 260
- Sophoridine, 203
- Soporific, 122, 125, 130, 161, 177, 187, 192, 240, 251
- Sore throat, 19, 34, 63, 90, 117, 195, 216, 218, 220, 223, 248
- Sparteine, 203, 243
- Sparteine, 243
- Spasm, 14, 67, 78, 128, 130, 137, 177, 196, 242, 255, 260, 265
- Spastic, 78, 87, 89, 90, 127, 208
- Spastic paresis, 78, 90
- Spathulenol, 181, 201, 225
- Spazmophilia, 193
- Spermatocidic, 165
- Spherophysine, 232
- Spherosine, 232
- Spinasterol, 230
- Spleen, 32, 41, 62, 77, 113, 115, 117, 217, 230
- Sprain, 235
- Stachydrine, 57, 64, 156, 157, 161, 167
- Stachyose, 31
- Staphylococcus, 24, 114, 138, 139, 147, 166, 183, 201, 217, 225, 250, 258, 266, 267
- Starch, 14, 19, 22, 34, 49, 60, 82, 85, 99, 113, 153, 160, 170, 226
- Stearic acid, 134, 271
- Stearidonic acid, 39
- Stenocardia, 83, 140, 237, 248
- Sterin, 73, 132
- Steroid, 119, 120, 140, 165, 166, 176, 177, 190, 191, 223, 227, 246, 253, 258, 260, 265
- Steroid saponin, 29, 176, 246
- Sterol, 14, 80, 92, 176, 206, 232
- Stigmasterin, 191, 236, 250
- Stigmasterol, 230
- Stilbene derivatives, 210
- Stimulate, 10, 54, 62, 69, 73, 85, 97, 102, 177, 180, 233, 241, 242, 252, 253, 260, 262, 265, 266
- Stomach, 13, 14, 28, 34, 41, 47, 65, 70, 80, 85, 87, 90, 95, 97, 104, 113, 115, 117, 124, 127, 128, 130, 135, 137, 139, 145, 148, 150, 161, 181, 183, 191–193, 196, 208, 212, 215, 218, 220, 225, 230, 233, 238, 245, 246, 251, 252, 257, 263, 265
- Stomach ache, 54, 55, 62, 99
- Stomach catarrh, 85, 117, 124, 192, 208, 230
- Stomatitis, 107, 124, 138, 140, 180, 218, 245
- Streptococcus, 147
- Streptozotocin-induced, 265
- Stress, 27, 117, 143, 166, 171, 197, 211
- Stress factor, 27
- Strychnine, 11, 13, 32, 130, 213, 255, 262
- Substance, 9, 10, 12–14, 16, 17, 24, 34, 49, 51, 64, 73, 79, 80, 88, 113, 120, 128, 132, 142, 151, 161, 178, 183, 187, 192, 198, 207, 235, 240, 268
- Succinic acid, 14, 165
- Sucrose, 34, 211
- Sudorific, diaphoretic, 39, 58, 62, 72, 108, 117, 128, 153, 161, 180, 183, 187, 202, 212, 218, 220, 236, 250, 263, 265
- Sugar, 13, 14, 28, 42, 57, 64, 69, 70, 82, 85, 88, 107, 115, 117, 118, 127, 128, 135, 139, 145, 147–151, 172, 175, 178, 182, 183, 196, 202, 206, 208, 210, 212, 213, 215–218, 220, 222, 241, 250, 252, 253, 255, 256, 263, 271
- Sulfur, 171
- Sunstroke, 213
- Surgery, 9, 90, 137
- Swelling, 70, 120, 151, 166, 187, 198, 246, 252, 258, 261, 267
- Sympathetic nervous system, 154
- Sympathomimetic, 100, 105
- Synapse, 251
- Synergistic, 221
- Syphilis, 52, 64, 97, 98, 110, 114, 133, 160, 176, 187, 193
- Systolic, 246
- T**
- Tachycardia, 75, 83, 109, 161, 171, 225, 238, 267
- Tachyphylaxis, 100
- Talatizamine, 23
- Talatizidine, 23
- Talatizine, 23
- Talimidine, 54
- Tannins, 5, 13, 14, 18, 21, 24, 26, 28, 41, 42, 46, 52, 53, 58, 59, 65, 67, 70, 73, 83, 84, 87, 93, 102–104, 107, 117, 119, 120, 123, 124, 127, 135, 138, 143, 147, 148, 150, 155–157, 160, 161, 165, 166, 171, 172, 175, 176, 178, 181, 182, 186, 192, 195–197, 202, 205, 208, 210, 211, 213, 215, 216, 218, 220–222, 226, 227, 231, 235, 237, 240, 242, 246, 250, 252, 255, 256, 265, 266, 271
- Tanshinone, 188, 225
- Tapeworm, 145
- Taraxanthin, 236, 250
- Taraxasterol, 70, 236
- Taraxerol, 236
- Taraxol, 236
- Tartaric acid, 14, 65, 166, 217, 218, 220, 236
- Taspine, 160
- Tatsetine, 251
- Taxifolin, 215
- Taxodione, 223
- Terpene, 10, 75, 138, 151, 268
- Terpinene, 62, 148
- Terpinen-4-ol, 267
- Terpineol, 177, 181
- Terpinolene, 62, 148
- Tetanus,

[AU38]

[AU39]

- Thalfine, 237
 Thalfinine, 237
 Thalicmidine, 55, 240
 Thalicmine, 238, 240
 Thalicminine, 238, 240
 Thalicmitrine, 240
 Thalispodine, 238
 Thalispopine, 238, 240
 Thalmine, 240
 Thalminine, 240
 Thermopsine, 241–243
 Thermopsocide, 241
 Thioglycoside, 64
 Thiramine, 230
 Thirst, 28, 54, 55, 218
 Throat, 19, 28, 34, 63, 90, 92, 98, 117, 128, 142, 167, 175, 180, 182, 195, 216, 218, 220, 223, 225, 230, 248, 257, 258, 263, 266
 Thrombocytes, 64
 Thrombophlebitis, 170
 Thromboplastic activity, 113
 Thujone, 48, 235
 Thymol, 168, 180, 181, 245, 267
 Thymoquinone, 177
 Thymus, 3, 117, 245
 Thyroid gland, 165, 166
 Thyroid stimulating hormone (TSH), 165
 Thyroidtoxicosis, 166
 Thyroxine (T4), 166
 Tick, 201
 Tincture, 12, 19, 46, 57, 64, 83, 84, 87, 92, 119, 138, 145, 148, 149, 155, 157, 161, 166, 167, 186, 201, 225, 248, 255, 256, 261, 265, 267, 268
 Tissue, 10, 39, 69, 120, 197, 220, 246, 247, 252
 Titratable organic acids, 107, 172
 Tocopherol, 190, 220
 Tolmetin, 240
 Tongue, 7, 60
 Tonic, 11–13, 27, 55, 80, 85, 113, 115, 117, 125, 128, 138, 140, 145, 160, 178, 183, 196, 210, 212, 216, 218, 237, 246
 Tonsillitis, 22
 Tonus, 57, 65, 87, 92, 203, 252
 Tooth, 87, 137
 Toothache, 262
 Torachryson, 210
 Toxic, 13, 32, 37, 46, 59, 69, 75, 90, 94, 95, 114, 137, 153, 159, 167, 213, 253
 Tracheitis, 100, 145, 250
 Tranquilizing effect, 122
 Trans-carveol, 159
 Triacanthine, 127
 Triacylglyceride, 92
 Triacylglycerol, 230
 Trichodesmine, 247
 Trichomoniasis, 150
 Tricyclic, 181
 Trifolin, 248
 Trifoside, 248
 Triglyceride, 92
 Triglycoside isorhamnetin, 133
 Trihydroxychalcone, 128
 Triiodothyronine (T3), 166
 Trimethoxyl-cinnamic acid, 92
 Triterpene, 11, 13, 32, 37, 80, 83, 93, 95, 97, 110, 119, 127, 133, 186, 192, 193, 211, 226, 227, 236, 237, 253, 271
 Triterpene alcohol, 80, 236
 Triterpene glycoside, 11, 95, 97, 127, 193, 211, 226, 227, 237
 Triterpene saponin, 32, 37, 83, 93, 133, 186, 193, 253
 Triterpenoid saponin, 257, 271
 Triterpenol, 92
 Tropane alkaloid, 77
 Trophic ulcer, 216, 225
 Tropine, 77
 Tropinone, 77
 Tropolone, 13
 Trypanocidal, 265
 TSH.. *See* Thyroid stimulating hormone (TSH)
 Tuberculosis, 19, 22, 31, 37, 48, 54, 58, 65, 72, 78, 85, 97, 104, 112, 113, 120, 132, 143–145, 147, 148, 150, 151, 166, 173, 190–192, 195, 210, 212, 216, 226, 235, 237, 238, 240, 250, 271
 Tubular necrosis, 75
 Tumor, 10, 22, 33, 34, 54, 62, 70, 79, 90, 103, 108, 113–115, 119, 122, 123, 132, 137, 139, 162, 178, 192, 195, 196, 217, 221, 226, 246, 250, 253, 257, 258, 271
 Turkesterone, 11, 27
 Tussilagins, 250
 Tympanites, 171, 180
 Typhoid fever, 231, 267
 Tyramine, 65
 Tyrosine, 221, 248
 Tyrosol, 211
- U**
- Ulcer, 13, 14, 16, 23, 27, 28, 34, 41, 47, 48, 65, 73, 77–80, 87, 95, 97, 100, 117, 127, 129, 135, 138, 144, 145, 147, 148, 150, 151, 175, 178, 191–193, 196, 198, 208, 212, 215, 216, 218, 221, 225, 226, 237, 238, 245, 246, 248, 251, 252
 Umbelliferol, 205
 Umbelliferone, 88, 113–115, 133
 Undecanoic acid, 245
 Unsaturated fatty acid, 14, 134, 147
 Upper respiratory, 134, 180, 216, 226, 245, 248, 250
 Ureter, 133, 181
 Urethra, 67
 Urinary incontinence, 34, 218
 Urinary tract, 12, 16, 104, 181
 Urination disorders, 73
 Urogenital, 58, 148, 150
 Uronic acid, 69
 Ursane, 223
 Ursolic acid, 26, 171, 192
 Urticin, 252
 Uterine atonia, 232
 Uterine fibroids, 75
 Uterine hemorrhages, 16
 Uterine horn, 54
 Uterine ulcers, 48
 Uterus, 161, 236
- V**
- Vaccine, 32
 Vagus nerve, 262
 Vaillantins, 118
 Valerianic acid, 46
 Valeric acid, 14
 Valeride, 255
 Valerine, 255
 Valtrate, 255
 Vanillic acid, 115, 142
 Vasicinone, 59, 187

Vasoconstrictor, 100, 252
Vasomotor center, 125
Vegetative neurosis, 83, 113
Vein, 27, 29, 31, 65, 74, 85, 88, 92, 124, 134, 137, 144, 150, 155, 157, 180, 191, 192, 208, 210, 221, 222, 225, 230, 247, 256, 270, 271
Venereal diseases, 32, 65, 147
Verbascoside, 258
Verbenone, 159
Vermifuge, 41, 46, 62, 64, 88, 113, 117, 132, 143–145, 147, 163, 177, 207, 221, 235, 241, 255, 260, 267, 272
Vessel, 14, 102, 113, 125, 178, 203, 242, 252, 260
Veterinary medicine, 23, 32, 92
Vincamine, 11, 13, 262
Vincandine, 262
Vincanine, 11, 262
Vinervine, 262
Vinervinine, 262
Virus, 14, 37, 103, 107, 175, 212, 221
Vitamin, 14, 16, 18, 22, 26, 28, 29, 33, 34, 36, 41–43, 51, 52, 57, 63–65, 70, 73, 75, 77, 83, 84, 88, 95, 100, 103–105, 107, 117–119, 124, 132, 135, 139, 140, 142, 143, 147, 148, 150, 155–157, 165–167, 170, 175, 177, 178, 183, 190, 192, 196, 200, 202, 210, 212, 213, 215, 216, 218, 220–223, 227, 230–233, 236, 240, 246, 248, 250, 252, 257, 261, 263, 266, 271, 272
Vitexin, 83
Vitexin-ramnoside, 83
Viticosterone, 11, 228
Vitiligo, 14, 205, 218
Vomit, 94, 124, 266
Vulgarol, 167

W

Wart, 110, 236
Waxes, 14, 39
Weakened, 27, 201
Weakness, 69, 85, 109, 162, 228, 236
Weariness, 211
Weight deficiency, 27
Wound, 16, 23, 26–28, 32, 37, 39, 41, 48, 54, 55, 64, 69, 77, 78, 92, 104, 107, 110, 114, 135, 138, 140, 147, 150, 151, 154, 160, 166, 170, 173, 178, 181, 188, 191, 192, 195, 196, 201, 208, 215, 221, 223, 225, 226, 228, 235, 237, 245, 247, 248, 251, 252, 257, 258, 261, 262, 268, 272
Wound healing, 64, 107, 150, 154

X

Xanthamine, 265
Xanthanine, 265
Xanthanol, 265
Xanthinine, 265
Xanthinosin, 265
Xanthone, 139
Xanthophyll, 215
Xanthosine, 265
Xanthostrumarin, 265
Xanthumanol, 265
Xyloglucoside, 260

Z

Zygodabagine, 272

Author Queries

Chapter No.: 0625532

Query	Details Required	Author's Response
AU1	Found as Adaptagenic. Please check.	
AU2	Not found in text. Please check.	
AU3	Found as Trans-anethole. Please check.	
AU4	Found as Angioprotective. Please check.	
AU5	Found as Anti aggregating. Please check.	
AU6	Not found in text. Please check.	
AU7	Found as Anti-fibrillating. Please check.	
AU8	Found as Ascariidosis. Please check.	
AU9	Not found in text. Please check.	
AU10	Not found in text. Please check.	
AU11	Not found in text. Please check.	
AU12	Not found in text. Please check.	
AU13	We have referred the page numbers only where coating is related to remedy. Please check if this is ok.	
AU14	Found as Delsoline. Please check.	
AU15	Not found exactly. Please check.	
AU16	We have referred the page numbers only where depression is related to emotional, physical/physiological. Please check if this is ok.	
AU17	Found as Esophageal. Please check.	
AU18	Found as Fainting. Please check.	
AU19	Not found in text. Please check.	
AU20	Found as Fungicidal. Please check.	
AU21	Found as Fungi. Please check.	
AU22	Found as Hepatoprotective. Please check.	
AU23	Found as Hepatotoxicity. Please check.	
AU24	Not found in text. Please check.	
AU25	Not found in text. Please check.	
AU26	Found as Itching. Please check.	
AU27	Found as Leucomisine. Please check.	
AU28	Not found in text. Please check.	
AU29	Found as Local anesthetic. Please check.	
AU30	Not found in text. Please check.	
AU31	Not found in text. Please check.	
AU32	Found as Parasympathetic cardiac ganglions. Please check.	
AU33	Not found in text. Please check.	
AU34	Not found in text. Please check.	
AU35	Found as Propionic acid. Please check.	
AU36	Not found in text. Please check.	
AU37	Found as Silybum. Please ccheck.	
AU38	Found as Synergistically. Please check.	
AU39	Not found in text. Please check.	

1 Index to Plant Species

2	A		
3	<i>Abies schrenkiana</i> , 190	<i>Anchusa azurea</i> , 39	54
4	<i>Acacia americana</i> , 127	<i>Anchusa italica</i> , 39	55
5	<i>Acacia triacanthos</i> , 127	<i>Artemisia absinthium</i> , 41, 130	56
6	<i>Acanos spina</i> , 178	<i>Artemisia annua</i> , 42	57
7	<i>Acanthophyllum gypsophiloides</i> , 7, 32	<i>Artemisia aromatica</i> , 43	58
8	<i>Achillea asiatica</i> , 16	<i>Artemisia chamomilla</i> , 42	59
9	<i>Achillea eupatorium</i> , 17	<i>Artemisia coarctata</i> , 48	60
10	<i>Achillea filipendulina</i> , 17	<i>Artemisia dracunculina</i> , 43	61
11	<i>Achillea millefolium</i> var. <i>manshurica</i> , 16	<i>Artemisia dracunculoides</i> , 43	62
12	<i>Achillea setacea</i> , 16, 18	<i>Artemisia dracunculoides</i> ssp. <i>dracunculina</i> , 43	63
13	<i>Achillea setacea</i> ssp. <i>asiatica</i> , 16	<i>Artemisia dracunculus</i> , 43	64
14	<i>Aconitum karakolicum</i> , 19, 22, 23	<i>Artemisia glauca</i> , 43	65
15	<i>Aconitum leucostomum</i> , 21, 271	<i>Artemisia leucodes</i> , 11, 44	66
16	<i>Aconitum napellus</i> var. <i>turkestanicum</i> , 19	<i>Artemisia opulenta</i> , 48	67
17	<i>Aconitum soongaricum</i> , 11, 19, 22	<i>Artemisia rupestris</i> ssp. <i>viridis</i> , 47	68
18	<i>Aconitum talassicum</i> , 23	<i>Artemisia rupestris</i> var. <i>viridis</i> , 47	69
19	<i>Aconitum winkleri</i> , 19	<i>Artemisia scoparia</i> , 46	70
20	<i>Aconogonon bucharicum</i> , 197	<i>Artemisia viridis</i> , 47	71
21	<i>Aconogonon coriarium</i> , 197	<i>Artemisia vulgaris</i> , 48	72
22	<i>Aconogonon coriarium</i> ssp. <i>bucharicum</i> , 197	<i>Arum elongatum</i> , 49	73
23	<i>Acroptilon picris</i> , 24	<i>Arum korolkowii</i> , 49	74
24	<i>Acroptilon repens</i> , 24	<i>Asparagus inderiensis</i> , 51	75
25	<i>Agrimonia asiatica</i> , 26	<i>Asparagus ledebourii</i> , 51	76
26	<i>Agrimonia eupatoria</i> ssp. <i>asiatica</i> , 26	<i>Asparagus persicus</i> , 6, 51	77
27	<i>Ajuga turkestanica</i> , 11, 27	<i>Aster helenium</i> , 145	78
28	<i>Alcea leucantha</i> , 33	<i>Astragalus sieversianus</i> , 52	79
29	<i>Alcea nudiflora</i> , 33	<i>Athamanta macrophylla</i> , 168	80
30	<i>Alhagi camelorum</i> , 28	<i>Atraphaxis spinosa</i> , 53	81
31	<i>Alhagi maurorum</i> , 28		
32	<i>Alhagi persarum</i> , 6, 28	B	82
33	<i>Alhagi pseudalhagi</i> , 28	<i>Berberis integerrima</i> , 54	83
34	<i>Allium karataviense</i> , 29, 31	<i>Berberis oblonga</i> , 3, 7, 54, 55	84
35	<i>Allium suvorovii</i> , 31	<i>Betonica foliosa</i> , 12, 57	85
36	<i>Allochrysa gypsophiloides</i> , 32	<i>Biarum sewertzowii</i> , 49	86
37	<i>Althaea kragujevacensis</i> , 34	<i>Bidens comosa</i> , 58	87
38	<i>Althaea micrantha</i> , 34	<i>Bidens orientalis</i> , 58	88
39	<i>Althaea nudiflora</i> , 33	<i>Bidens tripartita</i> , 58	89
40	<i>Althaea officinalis</i> , 34	<i>Biebersteinia multifida</i> , 59	90
41	<i>Althaea sublobata</i> , 34	<i>Botrycarpum nigrum</i> , 212	91
42	<i>Althaea taurinensis</i> , 34	<i>Buniella chaerophylloides</i> , 60	92
43	<i>Althaea vulgaris</i> , 34	<i>Bunium chaerophylloides</i> , 60	93
44	<i>Amaranthus retroflexus</i> , 36	<i>Bunium persicum</i> , 7, 60, 62	94
45	<i>Amaranthus tricolor</i> , 36	<i>Bursa bursa-pastoris</i> , 65	95
46	<i>Anagallis arvensis</i> , 37, 38	<i>Bursa pastoris</i> , 65	96
47	<i>Anagallis arvensis</i> f. <i>coerulea</i> , 38		
48	<i>Anagallis arvensis</i> ssp. <i>foemina</i> , 38	C	97
49	<i>Anagallis arvensis</i> var. <i>coerulea</i> , 38	<i>Caesalpiniodes triacanthum</i> , 127	98
50	<i>Anagallis coerulea</i> , 38	<i>Campanula glomerata</i> , 63	99
51	<i>Anagallis foemina</i> , 38	<i>Capparis herbacea</i> , 64	100
52	<i>Anagallis latifolia</i> , 37	<i>Capparis spinosa</i> , 64	101
53	<i>Anagallis phoenicea</i> , 37		

- 102 *Capsella bursa-pastoris*, 65
 103 *Capsella hircana*, 65
 104 *Carduus marianus*, 230
 105 *Carota sativa*, 88
 106 *Carota sylvestris*, 88
 107 *Carum carvi*, 67
 108 *Carum chaerophylloides*, 60
 109 *Carum confusum*, 60
 110 *Carum gracile*, 67
 111 *Carum heterophyllum*, 62
 112 *Carum persicum*, 62
 113 *Carum rosellum*, 67
 114 *Carum sogdianum*, 60
 115 *Caucalis carota*, 88
 116 *Caucalis daucus*, 88
 117 *Centaurea benedicta*, 73
 118 *Centaurea depressa*, 68
 119 *Centaurea repens*, 24
 120 *Cephalaria dipsacoides*, 95
 121 *Ceratocephala orthoceras*, 69
 122 *Ceratocephala reflexa*, 69
 123 *Ceratocephala testiculata*, 69
 124 *Chamaenerion hirsutum*, 103
 125 *Chlainanthus platycalyx*, 157
 126 *Chrysanthemum tanacetum*, 235
 127 *Chrysanthemum turkestanicum*, 163
 128 *Chrysanthemum vulgare*, 235
 129 *Cichorium intybus*, 70
 130 *Cicuta major*, 75
 131 *Cicuta officinalis*, 75
 132 *Clematis grata*, 72
 133 *Clematis incisedentata*, 72
 134 *Clematis orientalis*, 72
 135 *Clematis orveniae*, 72
 136 *Clematis petersiana*, 72
 137 *Clematis thunbergii*, 72
 138 *Clematis triloba*, 72
 139 *Clematis viridiflora*, 72
 140 *Cnicus benedictus*, 73
 141 *Codonocephalum grande*, 144
 142 *Codonopsis clematidea*, 3, 74
 143 *Conium cicuta*, 75
 144 *Conium maculatum*, 75
 145 *Conium maculosum*, 75
 146 *Convolvulus arvensis*, 77
 147 *Convolvulus chinensis*, 77
 148 *Convolvulus chondrilloides*, 78
 149 *Convolvulus dorychium* ssp. *subhirsutus*, 78
 150 *Convolvulus sagittifolius*, 77
 151 *Convolvulus subhirsutus*, 78
 152 *Convolvulus tschimganicus*, 78
 153 *Conyza britannica*, 143
 154 *Coriandrum cicutum*, 75
 155 *Coriandrum maculatum*, 75
 156 *Corvisartia helenium*, 145
 157 *Cousinia lappacea*, 77
 158 *Cousinia umbrosa*, 80
 159 *Crambe cordifolia* ssp. *kotschyana*, 82
 160 *Crambe cordifolia* var. *kotschyana*, 82
 161 *Crambe kotschyana*, 82
 162 *Crambe palmatifida*, 82
 163 *Crambe sewerzowii*, 82
 164 *Crataegus altaica*, 83
 165 *Crataegus chlorocarpa*, 83
 166 *Crataegus fischeri*, 84
 167 *Crataegus korolkowii*, 83
 168 *Crataegus purpurea* var. *altaica*, 83
Crataegus sanguinea var. *incisa*, 83 169
Crataegus sanguinea var. *inermis*, 83 170
Crataegus songarica, 84 171
Crataegus wattiana var. *incisa*, 83 172
Crossostephium turkestanicum, 163 173
Crucifera capsella, 65 174
Crucifera diversifolia, 162 175
Cucumis agrestis, 172 176
Cucumis melo var. *agrestis*, 172 177
Cullen drupacea, 205 178
Cyanus depressus, 68 179
- D** 180
Dactylorhiza umbrosa, 85 181
Dactylorhiza umbrosa, 85 182
Datura stramonium, 87 183
Datura tatula, 87 184
Daucus bactrianus, 88 185
Daucus carota, 88 186
Daucus exarmatus, 88 187
Daucus pulcherrimus, 88 188
Delphinium bitternatum var. *leiocarpum*, 90 189
Delphinium confusum, 89 190
Delphinium hybridum var. *sulphureum*, 90 191
Delphinium semibarbatum, 90 192
Delphinium zalil, 90 193
Descurainia sophia, 92 194
Dianthus hoeltzeri, 93 195
Dianthus superbus, 93 196
Dicranostigma iliense, 125 197
Dictamnus albus ssp. *turkestanicus*, 94 198
Dictamnus angustifolius, 94 199
Dipsacus azureus, 12, 95 200
Dipsacus dipsacoides, 95 201
Dipsacus laciniatus, 97 202
Dodartia atro-coerulea, 98 203
Dodartia orientalis, 98 204
Dracocephalum inderiense, 159 205
Dracocephalum royleanum, 159 206
Drosanthe scabra, 139 207
- E** 208
Elaeagnus rhamnoides, 135 209
Eminium lehmannii, 99 210
Eminium regelii, 99 211
Ephedra equisetina Bunge, 100 212
Ephedra ferganensis, 102 213
Ephedra glauca, 102 214
Ephedra intermedia, 102 215
Ephedra microsperma, 102 216
Ephedra persica, 102 217
Ephedra procera var. *typica*, 100 218
Ephedra tesquorum, 102 219
Ephedra tibetica, 102 220
Ephedra valida, 102 221
Epilobium hirsutum, 103 222
Epilobium tomentosum, 103 223
Epilobium velutinum, 103 224
Epilobium villosum, 103 225
Equisetum arvense, 104 226
Equisetum boreale, 104 227
Equisetum calderi, 104 228
Equisetum saxicola, 104 229
Eremurus regelii, 105 230
Eremurus spectabilis ssp. *regelii*, 105 231

- 232 *Erodium cicutarium*, 107
- 233 *Erodium pulchellum*, 107
- 234 *Eryngium biebersteinianum*, 108
- 235 *Eryngium caucasicum*, 108
- 236 *Eryngium coeruleum*, 107
- 237 *Eryngium pskemense*, 108
- 238 *Erysimum andrzejowskianum*, 109
- 239 *Erysimum canescens*, 109
- 240 *Erysimum diffusum*, 109
- 241 *Euphorbia jaxartica*, 110
- 242 *Euphorbia rapulum*, 112
- 243 *Euphorbia virgata* ssp. *jaxartica*, 110
- 244 *Euphorbia walsteinii* ssp. *jaxartica*, 110
- 245 **F**
- 246 *Faldermannia parviflora*, 270
- 247 *Fedia intermedia*, 186
- 248 *Fedia rupestris* var. *intermedia*, 186
- 249 *Ferula assa-foetida*, 113
- 250 *Ferula foetida*, 6, 10, 113–115
- 251 *Ferula jaeschkeana*, 114
- 252 *Ferula kuhistanica*, 114
- 253 *Ferula moschata*, 115
- 254 *Ferula pseudo-oreoselinum*, 115
- 255 *Ferula sumbul*, 115
- 256 *Ferula urceolata*, 115
- 257 *Fragaria chinensis*, 117
- 258 *Fragaria concolor*, 117
- 259 *Fragaria vesca*, 117
- 260 *Friedrichsthalia incana*, 247
- 261 *Fritillaria sewerzowii*, 153
- 262 *Fumaria vaillantii* Loisel, 118
- 263 *Fumaria vaillantii* var. *schrammii*, 118
- 264 **G**
- 265 *Galium boreale* ssp. *septentrionale*, 119
- 266 *Galium glabratum*, 120
- 267 *Galium septentrionale*, 119
- 268 *Galium verum*, 120
- 269 *Gentiana olivieri*, 122
- 270 *Gentiana regeliana*, 122
- 271 *Gentiana weschniakowii*, 122
- 272 *Geranium cicutarium*, 107
- 273 *Geranium collinum*, 123
- 274 *Geranium minutum*, 123
- 275 *Geranium saxatile*, 123
- 276 *Geranium wakhanicum*, 123
- 277 *Geum kokanicum*, 182
- 278 *Geum rivale*, 124
- 279 *Glaucium fimbriigerum*, 125
- 280 *Glaucium luteum* var. *fimbriigerum*, 125
- 281 *Glaucium refractum*, 213
- 282 *Gleditsia brachycarpa*, 127
- 283 *Gleditsia bujotii*, 127
- 284 *Gleditsia elegans*, 127
- 285 *Gleditsia hebecarpa*, 127
- 286 *Gleditsia heterophylla*, 127
- 287 *Gleditsia horrida*, 127
- 288 *Gleditsia inermis*, 127
- 289 *Gleditsia meliloba*, 127
- 290 *Gleditsia micracantha*, 127
- 291 *Gleditsia polysperma*, 127
- 292 *Gleditsia spinosa*, 127
- 293 *Gleditsia triacanthos*, 127
- 294 *Gleditsia triacanthus*, 127
- Glossocomia clematidea*, 74 295
- Glycyrrhiza asperrima* var. *desertorum*, 129 296
- Glycyrrhiza asperrima* var. *uralensis*, 129 297
- Glycyrrhiza glabra*, 7, 128, 129 298
- Glycyrrhiza glandulifera*, 128, 129 299
- Glycyrrhiza hirsuta*, 128 300
- Glycyrrhiza uralensis*, 129 301
- Glycyrrhiza violacea*, 128 302
- Goebelia alopecuroides*, 203 303
- Goebelia pachycarpa*, 260 304
- Grossularia nigra*, 212 305
- H** 306
- Haplophyllum acutifolium*, 130 307
- Haplophyllum flexuosa*, 130 308
- Haplophyllum perforatum*, 10, 130 309
- Haplophyllum sieversii*, 130 310
- Hedysarum alhagi*, 28 311
- Hedysarum pseudalhagi*, 28 312
- Helenium grandiflorum*, 145 313
- Helichrysum maracandicum*, 132 314
- Herniaria glabra*, 133 315
- Herniaria suavis*, 133 316
- Hesperis sophia*, 92 317
- Hibiscus ternatus*, 134 318
- Hibiscus trionum*, 134 319
- Hierapicra benedicta*, 73 320
- Hippomarathrum sarawschanicum*, 201 321
- Hippophae angustifolia*, 135 322
- Hippophae littoralis*, 135 323
- Hippophae rhamnoides*, 7, 12, 135 324
- Hippophae rhamnoides*, 135 325
- Hippophae sibirica*, 135 326
- Hyalolaena sewerzowii*, 201 327
- Hyoscyamus agrestis*, 137 328
- Hyoscyamus bohemicus*, 137 329
- Hyoscyamus niger*, 137 330
- Hypericum asperum*, 139 331
- Hypericum komorovii*, 137 332
- Hypericum nachitschevanicum*, 138 333
- Hypericum perforatum*, 138, 139 334
- Hypericum scabrum*, 139 335
- Hyssopus ferganensis*, 140 336
- Hyssopus seravschanicus*, 140 337
- Hyssopus tianschanicus*, 140 338
- I** 339
- Iberis bursa-pastoris*, 65 340
- Impatiens brachycentra*, 142 341
- Impatiens parviflora*, 142 342
- Inula britannica*, 143 343
- Inula grandis*, 144 344
- Inula helenium*, 143, 145 345
- Inula macrophylla*, 144 346
- Inula serrata*, 143 347
- Inula tymiensis*, 143 348
- J** 349
- Juglans duclouxiana*, 147 350
- Juglans fallax*, 147 351
- Juglans kamaonia*, 147 352
- Juglans orientis*, 147 353
- Juglans regia*, 3, 7, 147 354
- Juglans sinensis*, 147 355

- 356 *Juniperus excelsa* var. *macrocarpa*, 150
 357 *Juniperus intermedia*, 151
 358 *Juniperus jarkendensis*, 149
 359 *Juniperus kulsai*, 150
 360 *Juniperus polycarpos* var. *seravschanica*, 150
 361 *Juniperus polysperma*, 150
 362 *Juniperus pseudosabina*, 150, 151
 363 *Juniperus pseudosabina* var. *turkestanica*, 151
 364 *Juniperus pseudosabina* var. *typica*, 150, 151
 365 *Juniperus sabina*, 148–150
 366 *Juniperus sabina* var. *globosa*, 150
 367 *Juniperus sabina* var. *jarkendensis*, 149
 368 *Juniperus sabina* var. *macrocarpa*, 150
 369 *Juniperus schunganica*, 149
 370 *Juniperus semiglobosa*, 7, 149
 371 *Juniperus seravschanica*, 150
 372 *Juniperus taurica*, 150
 373 *Juniperus tianshanica*, 149
 374 *Juniperus turkestanica*, 151
 375 *Juniperus zaaminica*, 150
- 376 **K**
 377 *Koelzella pabularia*, 201
 378 *Korolkowia sewerzowii*, 153
- 379 **L**
 380 *Lachnophyllum gossypinum*, 154
 381 *Lagochilus gypsaceus*, 155
 382 *Lagochilus iliensis*, 156
 383 *Lagochilus keminensis*, 156
 384 *Lagochilus macrodontus*, 156
 385 *Lagochilus platyacanthus*, 12, 156
 386 *Lagochilus platycalyx*, 12, 157
 387 *Lallemantia royleana*, 159
 388 *Leontice ewersmannii*, 160
 389 *Leontice leontopetalum* ssp. *ewersmannii*, 160
 390 *Leontodon taraxacum*, 236
 391 *Leonurus cardiaca* ssp. *turkestanicus*, 161
 392 *Leonurus turkestanicus*, 161
 393 *Lepidium perfoliatum*, 162
 394 *Lepidolopsis turkestanica*, 163
 395 *Lithoon sieversianum*, 52
 396 *Lithospermum officinale*, 165
 397 *Lotodes drupaceum*, 205
 398 *Lycopus europaeus*, 166
- 399 **M**
 400 *Malva althaea*, 34
 401 *Malva maritima*, 34
 402 *Malva officinalis*, 34
 403 *Marrubium alternidens*, 167
 404 *Marrubium anisodon*, 167
 405 *Marrubium kusnezowii*, 167
 406 *Mediasia macrophylla*, 7, 168
 407 *Melilobus heterophylla*, 127
 408 *Melilotus graveolens*, 170
 409 *Melilotus officinalis*, 170
 410 *Melilotus suaveolens*, 170
 411 *Melissa bicornis*, 171
 412 *Melissa officinalis*, 171
 413 *Melo agrestis*, 172
 414 *Mentha asiatica*, 173
 415 *Mentha kopetdaghensis*, 173
 416 *Mentha longifolia*, 173
Mentha vagans, 173
Micromeria formosana, 181
Morus alba, 175
- N**
Nasturtium perfoliatum, 162
Nepeta erodiifolia, 159
Nepeta nuda, 176
Nepeta pannonica, 176
Nigella indica, 177
Nigella sativa, 177
Nigella truncata, 177
- O**
Oligosporus scoparia, 46
Oligosporus dracuncululus, 43
Onopordum acanthium, 178
Orchis magna, 85
Orchis orientalis ssp. *turkestanica*, 85
Orchis umbrosa, 85
Origanum creticum, 181
Origanum dilatatum, 181
Origanum normale, 181
Origanum puberulum, 181
Origanum tyttanthum, 180
Origanum vulgare, 181
Origanum vulgare var. *genuinum*, 180
Origanum vulgare var. *prismaticum*, 180
Origanum vulgare var. *viride*, 180
Orthurus kokanicus, 182
Osyris rhamnoides, 135
- P**
Padus avium, 183
Padus racemosa, 183
Papaver ocellatum, 185
Papaver pavoninum, 185
Papaver refractum, 213
Patrinia intermedia, 186
Patrinia nudiuscula, 186
Peganum harmala, 11, 187
Perovskia abrotanoides, 188
Perovskia artemisioides, 188
Peucedanum asa-foetida, 113
Phaca salsula, 232
Picea morinda ssp. *tianschanica*, 190
Picea obovata Ledeb. var. *schrenkiana*, 190
Picea prostrata, 190
Picea robertii, 190
Picea schrenkiana, 190
Picea tianschanica, 190
Plantago borysthena, 192
Plantago dregeana, 192
Plantago lanceolata, 191
Plantago latifolia, 192
Plantago major, 192
Plantago officinarum, 192
Pleuropterypyrum bucharicum, 197
Polemonium caeruleum ssp. *caucasicum*, 193
Polemonium caucasicum, 193
Polygala comosa, 195
Polygala comosa var. *altaica*, 195
Polygala comosa var. *hybrida*, 195
Polygala hybrida, 197

- 478 *Polygonum aequale*, 196
 479 *Polygonum agreste*, 196
 480 *Polygonum aphyllum*, 196
 481 *Polygonum araraticum*, 196
 482 *Polygonum arenastrum*, 196
 483 *Polygonum aviculare*, 196
 484 *Polygonum berteroi*, 196
 485 *Polygonum bucharicum*, 7, 197
 486 *Polygonum coriarium*, 197
 487 *Polygonum heterophyllum*, 196
 488 *Polygonum retinerve*, 196
 489 *Polygonum striatum*, 196
 490 *Polygonum uruguense*, 196
 491 *Portulaca consanguinea*, 198
 492 *Portulaca intermedia*, 198
 493 *Portulaca marginata*, 198
 494 *Portulaca mundula*, 198
 495 *Portulaca neglecta*, 198
 496 *Portulaca oleracea*, 198
 497 *Portulaca pilosa*, 198
 498 *Portulaca pusilla*, 198
 499 *Portulaca retusa*, 198
 500 *Potentilla adscendens*, 200
 501 *Potentilla canescens*, 200
 502 *Potentilla inclinata*, 200
 503 *Potentilla vesca*, 117
 504 *Prangos cylindrocarpa*, 201
 505 *Prangos hissarica*, 201
 506 *Prangos lamellata*, 201
 507 *Prangos pabularia*, 7, 201
 508 *Prangos seravschanica*, 201
 509 *Prunus cerasifera* ssp. *sogdiana*, 202
 510 *Prunus cerasifera* var. *orientalis*, 202
 511 *Prunus mirabilis*, 202
 512 *Prunus orientalis*, 202
 513 *Prunus padus*, 183
 514 *Prunus racemosa*, 183
 515 *Prunus sogdiana*, 3, 7, 202
 516 *Pseudosiphora alopecuroides*, 203
 517 *Psoralea drupacea*, 7, 205
 518 *Pulicaria afghanica*, 206
 519 *Pulicaria lachnophylla*, 206
 520 *Pulicaria olivascens*, 206
 521 *Pulicaria salviifolia*, 206
 522 *Pulicaria sublepidota*, 206
 523 *Pyrethrum vulgare*, 235
 524 *Pyrus tianschanica*, 231
- R**
- 525 *Ranunculus testiculatus*, 69
 527 *Reseda luteola*, 207
 528 *Rhamnoides hippophae*, 7, 12, 135
 529 *Rhamnus cathartica*, 208
 530 *Rhamnus zizyphus*, 271
 531 *Rheum emodi*, 210
 532 *Rheum maximowiczii*, 210
 533 *Rheum megalocarpon*, 210
 534 *Rhodiola kirilowii*, 211
 535 *Rhodiola linearifolia*, 211
 536 *Rhodiola longicaulis*, 211
 537 *Rhodiola macrolepis*, 211
 538 *Rhodiola robusta*, 211
 539 *Ribes cyathiforme*, 212
 540 *Ribes nigrum*, 212
 541 *Ribes olidum*, 212
 542 *Ribes pauciflorum*, 212
- Roemeria refracta*, 213
Rosa canina, 215
Rosa caraganifolia, 216
Rosa ciliatosepala, 215
Rosa coeruleifolia, 216
Rosa epipsila, 216
Rosa fedtschenkoana, 216
Rosa lavrenkoi, 216
Rosa lipschützii, 216
Rosa minusculifolia, 216
Rosa oligosperma, 216
Rosa sosnovskyi, 215
Rubia iberica, 217
Rubia tinctorum, 217
Rubus caesius, 218
Rubus idaeus, 220
Rubus psilophyllus, 218
Rubus turkestanicus, 218
Rumex alpinus var. *subcalligerus*, 221
Rumex confertus, 221
Rumex tianschanicus, 222
Ruta acutifolia, 130
Ruta flexuosa, 130
Ruta perforata, 130
Ruta sieversii, 130
- S**
- Sabina seravschanica*, 150
Sabina vulgaris, 148, 149
Sabina vulgaris var. *jarkendensis*, 149
Salvia asperata, 225
Salvia deserta, 223
Salvia jailicola, 223
Salvia moldavica, 223
Salvia nemorosa, 223
Salvia pamirica, 225
Salvia sclarea, 223, 225
Sanguisorba glandulosa, 226
Sanguisorba officinalis, 226
Saponaria segetalis, 253
Saponaria vaccaria, 253
Scabiosa songarica, 7, 227
Scorodosma foetidum, 113
Sedum kirilowii, 211
Sedum longicaule, 211
Sedum macrolepis, 211
Sedum robustum, 211
Selinum conium, 75
Seriphidium leucodes, 44
Serratula alata, 228
Serratula dissecta var. *asperula*, 228
Serratula sogdiana, 228
Serratula trautvetteriana, 228
Seseli macrophyllum, 168
Silybum marianum, 230
Sisymbrium sophia, 92
Sisymbrium tenuissimum, 92
Sium conium, 75
Sophia lobelii, 92
Sophora alopecuroides, 203
Sophora lupinoides, 242
Sophora pachycarpa, 260
Sorbus tianschanica, 231
Sphaerophysa salsula, 232
Spinachia tetrandra, 233
Spinacia turkestanica, 233
- 543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
- 568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607

- 608 *Stachys betoniciflora*, 57
609 *Stachys betonicifolia*, 57
610 *Stachys foliosa*, 57
611 *Stramonium spinosum*, 87
- T**
- 612 *Tanacetum boreale*, 235
614 *Tanacetum crispum*, 235
615 *Tanacetum umbellatum*, 235
616 *Tanacetum vulgare*, 235
617 *Taraxacum dens-leonis*, 236
618 *Taraxacum officinale*, 236
619 *Taraxacum retroflexum*, 236
620 *Taraxacum sylvanicum*, 236
621 *Thalictrum foetidum*, 12, 237
622 *Thalictrum isopyroides*, 238
623 *Thalictrum minus*, 237, 240
624 *Thalictrum minus* var. *foetidum*, 237
625 *Thermopsis alterniflora*, 241
626 *Thermopsis dahurica*, 242
627 *Thermopsis glabra*, 242
628 *Thermopsis kaxgarica*, 243
629 *Thermopsis lanceolata*, 242, 243
630 *Thermopsis lanceolata* ssp. *turkestanica*, 243
631 *Thermopsis lupinoides*, 242
632 *Thermopsis rigida*, 241
633 *Thermopsis sibirica*, 242
634 *Thermopsis turkestanica*, 243
635 *Thlapsi bursa-pastoris*, 65
636 *Thymus amictus*, 245
637 *Thymus latifolius*, 245
638 *Thymus marschallianus*, 245
639 *Thymus pannonicus*, 245
640 *Thymus pannonicus* ssp. *marschallianus*, 245
641 *Thymus platyphyllus*, 245
642 *Thymus pseudopannonicus*, 245
643 *Thymus stepposus*, 245
644 *Tithymalus graminifolius* ssp. *jaxarticus*, 110
645 *Tithymalus rapulum*, 112
646 *Tragopyrum spinosum*, 53
647 *Tribulus bicornutus*, 246
648 *Tribulus terrestris*, 246
649 *Trichodesma incanum*, 247
650 *Trifolium officinale*, 170
651 *Trifolium pratense*, 248
652 *Trifolium ucrainicum*, 248
653 *Trochocephalus songaricus*, 227
654 *Tussilago farfara*, 250
- U**
- 655 *Ungernia victoris*, 251
657 *Urtica dioica*, 252
- V**
- 658 *Vaccaria hispanica*, 253
660 *Vaccaria parviflora*, 253
661 *Vaccaria pyramidata*, 253
662 *Vaccaria segetalis*, 253
663 *Vaccaria vulgaris*, 253
664 *Valeriana baltica*, 255
665 *Valeriana exaltata*, 255
666 *Valeriana officinalis*, 255
667 *Valeriana palustris*, 255
- Veratrum album* ssp. *lobelianum*, 256
Veratrum album ssp. *virescens*, 256
Veratrum album var. *lobelianum*, 256
Veratrum album var. *virescens*, 256
Veratrum lobelianum, 256
Verbascum khorassanicum, 257
Verbascum lychnitis, 257
Verbascum polystachyum, 257
Verbascum songaricum, 257
Verbascum thapsus, 257, 258
Vexibia alopecuroides, 203
Vexibia pachycarpa, 260
Vicia cracca, 261
Vicia hiteropus, 261
Vicia lilacina, 261
Vicia macrophylla, 261
Vinca erecta, 262
Viola pontica, 263
Viola suavis, 263
Viticella orientalis, 72
- W**
- Wahlenbergia clematidea*, 74
- X**
- Xanthium americanum*, 265
Xanthium cavanillesii, 265
Xanthium chasei, 265
Xanthium chinense, 265
Xanthium curvscens, 265
Xanthium echinatum, 265
Xanthium echinellum, 265
Xanthium globosum, 265
Xanthium inflexum, 265
Xanthium italicum, 265
Xanthium natalense, 265
Xanthium orientale, 265
Xanthium oviforme, 265
Xanthium pennsylvanicum, 265
Xanthium pungens, 265
Xanthium speciosum, 265
Xanthium strumarium, 265
Xanthium varians, 265
Xanthium wootonii, 265
- Z**
- Ziziphora afghanica*, 267
Ziziphora borzhomica, 267
Ziziphora brevicalyx, 267
Ziziphora bungeana, 266, 267
Ziziphora clinopodioides, 266, 267
Ziziphora clinopodioides ssp. *afghanica*, 267
Ziziphora clinopodioides ssp. *bungeana*, 266, 267
Ziziphora denticulata, 267
Ziziphora dzhavakhishvili, 267
Ziziphora pedicellata, 268
Ziziphora tenuior, 270
Ziziphora turcomaica, 267
Ziziphus jujuba, 7, 271
Ziziphus sativa, 271
Ziziphus vulgaris, 271
Zygophyllum fabago var. *oxianum*, 272
Zygophyllum oxianum, 272