

Changes to the nomenclature of the skates (Chondrichthyes: Rajiformes)

PETER R. LAST¹, SIMON WEIGMANN² & LEI YANG³

¹CSIRO National Research Collections Australia, Australian National Fish Collection, Castray Esplanade, Hobart, TAS, 7001, AUSTRALIA. E-mail: peter.last@csiro.au

²Elasmo-Lab, Elasmobranch Research Laboratory, Schlägertwiete 5b, 21335 Lüneburg, Germany. E-mail: simon.weigmann@elasmo-lab.de

³Department of Biology, College of Charleston, Charleston, SC 29401, USA. E-mail: leiyangslu@gmail.com

<http://zoobank.org/urn:lsid:zoobank.org:act:2F49A24B-812C-4008-BF2E-A7692337C214>

<http://zoobank.org/urn:lsid:zoobank.org:act:B6434AFE-AD1E-4D32-81C6-E9936F25157D>

Abstract

In the course of the NSF-funded project “Jaws and Backbone: Chondrichthyan Phylogeny and a Spine for the Vertebrate Tree of Life”, morphological and molecular data were collected for a huge number of species (including type specimens). Molecular studies using mitochondrial and nuclear markers with dense taxon sampling corroborate that the skates consist of four main family-level groups, i.e. Anacanthobatidae, Arhynchobatidae, Gurgesiellidae and Rajidae. The *Rays of the World* book followed this subdivision of skates resulting in several nomenclatural decisions at both supraspecific and species levels, which are described and discussed in the present paper. These nomenclatural changes include: 1) resurrection of the family Gurgesiellidae, comprising all eight species of *Cruriraja*, eight species of *Fenestraja* and three species of *Gurgesiella*; 2) supraspecific changes to anacanthobatid nomenclature, i.e. elevation of subgenus *Schroederobatis* to generic level and resurrection of *Springeria* from subgeneric rank as a valid genus-level taxon; 3) provisional assignment of members of two undefined genus-level taxa, the “North Pacific Assemblage” and the “Amphi-American Assemblage”; 4) reassignment of species to the genus *Dentiraja*; 5) resurrection of *Dipturus intermedius* as a valid species from synonymy with *D. batis*; 6) resurrection of the tribe Pavorajini McEachran, 1984; and 7) erection of two new tribes, Bathyrājini (type genus *Bathyrāja*) and Crurirājini (type genus *Cruriraja*). Furthermore, an annotated checklist of rajiform species is provided to explain major nomenclatural changes and place the list in context with other contemporary lists.

Key words: Rajiformes, skates, nomenclatural changes, genera, species, checklist.

Introduction

A key objective of a 5-year, NSF-funded project (“Jaws and Backbone: Chondrichthyan Phylogeny and a Spine for the Vertebrate Tree of Life”) was to investigate diversity in the chondrichthyan fishes and provide an inventory of species. Possibly the most challenging part of this task involved an investigation of the skates (Rajiformes), which comprise almost half of all batoid fishes. As part of the Tree of Life project, one of us (PL) travelled to museums globally to re-examine types and key specimens, and obtain images, for the compilation of skate chapters in the *Rays of the World* book. These data were supplemented by long-term datasets acquired by the co-authors (J. McEachran, B. Séret, M. Stehmann, and PL) and summarised in the skate chapters. The Tree of Life project also involved molecular analyses (NADH2 gene and mitochondrial genome) including data for more than half of the world’s extant skate species (Yang *et al.*, unpubl.). These

projects yielded new information at all taxonomic levels and these findings are combined herein with those of a comprehensive study of the group conducted concurrently by Weigmann (2016), with an update by Weigmann (in press). These investigations, along with those conducted by other contemporary leaders in the field (M. de Carvalho, D. Ebert, H. Ishihara, C. Jeong, J. McEachran, B. Séret, M. Stehmann) have led to the description of several new species in recent times, including 4 specifically for the ray book.

Studies by Naylor *et al.* (2012a,b) using broad taxon coverage for the NADH2 gene, Aschliman *et al.* (2012) using both mitochondrial and nuclear genes, and more recent studies by Naylor *et al.* (in press) and Yang *et al.* (unpubl.), using the mitochondrial genome and nuclear markers for even denser taxon sampling, corroborate that the skates consist of four main family-level groups. The *Rays of the World* (Last *et al.*, in press a) family contributions, i.e. Anacanthobatidae (Séret

et al., in press), Arhynchobatidae (Last *et al.*, in press b), Gurgesiellidae (Weigmann *et al.*, in press) and Rajidae (Last *et al.*, in press c), followed this subdivision of the skates resulting in the nomenclatural decisions at both supraspecific and species levels discussed below. An annotated checklist of species is provided to explain major nomenclature changes and place the list in context with other contemporary lists (i.e. Compagno, 2005; Weigmann, 2016).

Supraspecific changes to skate nomenclature

Resurrection of the family Gurgesiellidae de Buen, 1959

Familial structure of the skates (order Rajiformes *sensu* Naylor *et al.*, 2012a) has been the subject of much discussion in recent decades. Based on a morphological analysis, the group was considered to consist of a single family by McEachran & Dunn (1998) comprised of 5 tribes: Amblyrajini, Aryhchobatini, Gurgesiellini, Rajini and Riorajini. In preliminary work for a proposed catalogue of rays of the world, Compagno (1999) divided the order into three families: Anacanthobatidae, Arhynchobatidae and Rajidae. Naylor *et al.* (2012a,b), using a molecular analysis

based on the NADH2 gene, recovered the families Arhynchobatidae and Rajidae, but *Cruriraja* Bigelow & Schroeder, 1948 fell outside these groups and was well divergent from the Anacanthobatidae (see Naylor *et al.*, 2012a: Figure 2.9; Naylor *et al.*, 2012b: Figure 77). *Cruriraja* grouped as the sister to the Arhynchobatidae, and *Sinobatis* Hulley, 1973 grouped as the sister to a larger group containing the rest of these skates. In the past, the phylogenetic classification of *Cruriraja* (from latin *crus* = leg or limb and skate genus *Raja* Linnaeus; type species *Cruriraja atlantis* Bigelow & Schroeder, 1948) has changed frequently. *Cruriraja* was originally placed in the family Rajidae by Bigelow & Schroeder (1948). This classification was adopted by, e.g. Wallace (1967), Hulley (1986) and Carvalho *et al.* (2006). However, *Cruriraja* was placed in its own family Crurirajidae by Hulley (1972a) based on characters of the pelvic girdle and clasper structure. Later, species of the families Anacanthobatidae and Crurirajidae were combined in the family Anacanthobatidae without a reason given by Compagno (1999). This placement was followed by, e.g. Ebert & Compagno (2007), Last & Séret (2008) and Last & Stevens (2009), who justified the placement by the joint possession of leg-like anterior pelvic-fin lobes in all species. However, skates of the genus *Cruriraja* morphologically differ greatly from anacanthobatid legskates (Figure 1).




Figure 1: Comparison of the external morphology of *Cruriraja* and anacanthobatids. Adult female 532 mm TL (ZMH 105118) of *Cruriraja hulleyi* Aschliman, Ebert & Compagno, 2010 in (A) dorsal and (B) ventral views, as well as juvenile female 234.5 mm TL (ZMH 25928) of *Indobatis ori* (Wallace, 1967) in (C) dorsal and (D) ventral views.

The differences include firm, thick skin in *Cruriraja* (*vs.* soft, thin skin in anacanthobatids), a solid tail (*vs.* a thin, cord-like tail), the presence of two dorsal fins, except for one aberrant specimen of ZMH 122862 with only one dorsal fin (*vs.* no dorsal fins), and numerous thorns on the dorsal surface (*vs.* largely naked dorsally). The absence of thorns in the Anacanthobatidae was expressed also in the etymology of the family name (Weigmann *et al.*, 2014a). Furthermore, in *Cruriraja*

species the disc has somewhat angular outer corners (*vs.* rounded outer corners in anacanthobatids), the snout is rather short and stout (*vs.* at least moderately long and delicate), the mesopterygia of the pectoral skeleton are clearly better developed, the scapulocoracoid has fewer and larger postventral fenestrae, and the pelvic girdle has much shorter prepelvic processes (Figure 2; see also Weigmann *et al.*, 2014a).




Figure 2: Comparison of the skeletal morphology of *Cruriraja* and anacanthobatids. Radiographs in dorsal total views: (A) adult female 532 mm TL (ZMH 105118) of *Cruriraja hulleyi* Aschliman, Ebert & Compagno, 2010; (B) adult male 425 mm TL (ZMH 25926) of *Indobatis ori*.

Based on the numerous differences, Weigmann *et al.* (2014a) removed *Cruriraja* from the Anacanthobatidae, placing it within its own family Crurirajidae following Hulley (1972a). This classification was adopted by van der Laan *et al.* (2014), Ebert & van Hees (2015) and Weigmann (2016). The latter reference, however, indicated that unpublished data from the Tree of Life project suggested placement of *Cruriraja* in the family Gurgesiellidae de Buen, 1959, together with the genera *Fenestraraja* McEachran & Compagno, 1982 and *Gurgesiella* de Buen, 1959. The external morphologies of the latter two genera are shown in Figure 3.

Very recent molecular studies (Naylor *et al.*, in press; Yang *et al.*, unpubl.) have also provided compelling data supporting the resurrection of the family Gurgesiellidae. However, McEachran & Compagno (1979) had earlier discussed the validity of the monotypic families Gurgesiellidae and Pseudorajidae Bigelow & Schroeder, 1954 and revealed that genera *Gurgesiella* and *Pseudoraja* Bigelow & Schroeder, 1954

share five derived characters unique within the order (as suborder Rajoidei). Nevertheless, the Gurgesiellidae and Pseudorajidae had previously been recognized to be distinct families based on the structure of the pelvic girdle, neurocranium and hyomandibular cartilage (Hulley, 1972b). McEachran & Dunn (1998) later again placed these genera in separate families: i.e. *Gurgesiella* (Rajidae) and *Pseudoraja* (Arhynchobatidae). Although McEachran & Compagno's proposal that these genera should be placed together in the Pseudorajidae has credibility based on their original findings and reassignment of *Gurgesiella*, no fresh tissues are available for *Pseudoraja*, nor is an adult male available for clasper examination (J. McEachran, pers. comm.). In the absence of such critical information, Gurgesiellidae is provisionally resurrected as the name of this family-level taxon pending definitive information on *Pseudoraja*. The family comprises 8 species of *Cruriraja*, 8 species of *Fenestraraja*, and 3 species of *Gurgesiella* (Weigmann, 2016; Weigmann *et al.*, in press).




Figure 3: External morphology of *Fenestraraja* and *Gurgesiella*. Adult male 232 mm TL (ZMH 119851) of *Fenestraraja plutonia* (Garman, 1881) in (A) dorsal and (B) ventral views, as well as adult male holotype 424 mm TL (ZMH 25046) of *Gurgesiella dorsalis* McEachran & Compagno, 1980 in (C) dorsal and (D) ventral views.

Supraspecific changes to anacanthobatid nomenclature

The family Anacanthobatidae and the type genus *Anacanthobatis* were erected by von Bonde & Swart (1923) for *Leiobatis marmoratus* (type species, designated with lectotype by Hulley, 1973) and *L. dubius* n. spp. from South Africa. Subsequently, a second anacanthobatid genus was described, i.e. *Springeria* Bigelow & Schroeder, 1951 with the newly described type species *S. foliostris*. *Springeria* was later redefined by Hulley (1973) as a subgenus of *Anacanthobatis*, along with the erection of two new subgenera, *Schroederobatis* and *Sinobatis*, and the definition of subgenus *Anacanthobatis* von Bonde & Swart, 1923.

All four subgenera were defined based on differences in external and skeletal clasper morphology. The differences were described in detail by Hulley (1973) and updated by Weigmann *et al.* (2014a) and Stehmann & Weigmann (2016). *Sinobatis* was elevated to generic level by Last & Séret (2008) based on differences in clasper morphology. As such differences are considered to be diagnostic of genera also in other rajiform families, elevation of subgenera *Schroederobatis* and *Springeria* to generic level was recommended by Last & Séret (2008), but not formally performed. Subsequently, Weigmann *et al.* (2014a) erected a new genus, *Indobatis* Weigmann, Stehmann & Thiel, 2014, for *Anacanthobatis ori* (Wallace, 1967) due to strong differences in the formerly unknown external and skeletal clasper morphology, clearly distinguishing this genus from all other described anacanthobatid genera and subgenera.

Considering the extent of differences in external and skeletal clasper characters between all anacanthobatid genera and subgenera (Hulley, 1973; Weigmann *et al.*, 2014a; Stehmann & Weigmann, 2016), the subgenus *Schroederobatis* is herein elevated to generic level and *Springeria* is resurrected from subgeneric rank as a valid genus-level taxon.

Provisional assignment of members of two undefined genus-level taxa, the “North Pacific Assemblage” and the “Amphi-American Assemblage”

In a classification of the Rajidae based on a morphology-based phylogenetic analysis, McEachran & Dunn (1998) recognised 8 genus-level taxa within the tribe Rajini, including two un-named taxa, *Raja* “North Pacific Assemblage” and *Raja* “Amphi-American Assemblage”.

The *Raja* “North Pacific Assemblage” contained 6 species: *R.* “North Pacific Assemblage” *binoculata* Girard, 1855; *R.* “North Pacific Assemblage” *rhina* Jordan & Gilbert, 1880; *R.* “North Pacific Assemblage” *inornata* Jordan & Gilbert, 1881; *R.* “North Pacific Assemblage” *stellulata* Jordan & Gilbert, 1880; *R.* “North Pacific Assemblage” *pulchra* Liu, 1932; and *R.* “North Pacific Assemblage” *cortezensis* McEachran &

Miyake, 1988. The genus *Beringraja* was later erected by Ishihara *et al.* (2012) for *R. binoculata* and *R. pulchra* based on egg capsule morphology. Molecular analyses for the Chondrichthyan Tree of Life project (Yang *et al.*, unpubl.) have provided strong support for recognition of a supraspecific group including all 6 species of the “North Pacific Assemblage” (no molecular data are available for *R. cortezensis*) in the genus *Beringraja* within the tribe Rajini.

The *Raja* “Amphi-American Assemblage” is a morphologically variable group comprising 7 species: *R.* “Amphi-American Assemblage” *eglanteria* Bosc, 1800; *R.* “Amphi-American Assemblage” *ackleyi* Garman, 1881; *R.* “Amphi-American Assemblage” *equatorialis* Jordan & Bollman, 1890; *R.* “Amphi-American Assemblage” *texana* Chandler, 1921; *R.* “Amphi-American Assemblage” *cervigoni* Bigelow & Schroeder, 1964; *R.* “Amphi-American Assemblage” *bahamensis* Bigelow & Schroeder, 1965; and *R.* “Amphi-American Assemblage” *velezi* Chirichigno, 1973. Molecular analyses (Yang *et al.*, unpubl.) indicate that two of these species, *R. eglanteria* and *R. velezi* (no data for the other taxa), group with another of McEachran & Dunn’s (1998) rajine taxa, *Rostroraja* (i.e. *Rostroraja alba* (Lacepède, 1803)), within the newly erected tribe Rostrorajini (Chiquillo *et al.*, 2014). Hence, retention of *Raja* (in the Rajini) for this group is untenable so the *Raja* “Amphi-American Assemblage” is provisionally assigned to *Rostroraja* following the recommendation of Naylor *et al.* (2012a,b). Nevertheless, *Rostroraja alba* differs morphologically from members of the *Raja* “Amphi-American Assemblage” and is a massive skate occurring in the Eastern Atlantic and South-West Indian Ocean (rather than the Western Atlantic and Eastern Pacific). More research is needed to verify this placement.

Reassignment of species to the genus *Dentiraja* Whitley, 1940

The genus *Dentiraja* was originally described as a subgenus of *Raja* Linnaeus by Whitley (1940), with *Raja dentata* Klunzinger, 1872 as type species. The species was later found to be a junior synonym of *Raja lemprieri* Richardson, 1845 (Paxton *et al.*, 1989; Gomon *et al.*, 1994; Last & Stevens, 1994; Hoese *et al.*, 2006), which has subsequently been referred to as *Raja lemprieri* (Paxton *et al.*, 1989; Gomon *et al.*, 1994), *Raja* (*Dipturus*) *lemprieri* (Last & Stevens, 1994), *Dipturus lemprieri* (Last & Yearsley, 2002; Hoese *et al.*, 2006; Jeong *et al.*, 2007) and *Okamejei lemprieri* (McEachran & Dunn, 1998; Compagno, 1999; Ebert & Compagno, 2007).

The subgenus *Dentiraja* was resurrected and assigned to *Dipturus* Rafinesque, 1810 by Last & Yearsley (2002). Last & Gledhill (2008) subsequently elevated *Dentiraja* to full generic rank, based particularly on clasper morphology.

Recently, all species of the family Rajidae have been reviewed for the rajid chapter (Last *et al.*, in press c) in the *Rays of the World* (Last *et al.*, in press a). As pointed out in the remarks column of Table II for the respective species in Weigmann (2016), unpublished molecular results by Naylor (pers. comm.) indicated that several further species of *Dipturus* in fact belong to *Dentiraja*, i.e. *Dipturus australis* (Macleay, 1884), *Dipturus cerva* (Whitley, 1939), *Dipturus confusus* Last, 2008, *Dipturus endeavouri* Last, 2008 and *Dipturus polyommata* (Ogilby, 1910). Further data collected for the *Rays of the World* indicate that *Dipturus falloargus* Last, 2008, *Dipturus healdi* Last, White & Pogonoski, 2008 and *Dipturus oculus* Last, 2008 belong to *Dentiraja* as well. Accordingly, *Dentiraja* now comprises 10 valid species, i.e. *Dentiraja australis* (Macleay, 1884), *Dentiraja cerva* (Whitley, 1939), *Dentiraja confusa* (Last, 2008), *Dentiraja endeavouri* (Last, 2008), *Dentiraja falloarga* (Last, 2008), *Dentiraja flindersi* Last & Gledhill, 2008, *Dentiraja healdi* (Last, White & Pogonoski, 2008), *Dentiraja lemprieri* (Richardson, 1845), *Dentiraja oculata* (Last, 2008) and *Dentiraja polyommata* (Ogilby, 1910).

Species of *Dentiraja* differ from *Dipturus* species in the small (*vs.* large) size, short to moderately elongate (*vs.* elongate to very elongate) snout, and large internarial width usually exceeding 60% of prenasal snout length in adults (*vs.* usually less than 60% in adults). *Dentiraja* species resemble species of *Okamejei* Ishiyama, 1958 in the small size, but differ in the short (*vs.* long) inter- and postdorsal tail sections, and the absence (*vs.* presence), except for *Okamejei ornata* Weigmann, Stehmann & Thiel, 2015) of an external clasper component funnel. For clasper details of *Okamejei ornata* see Weigmann *et al.* (2015).

Annotated checklist for the order Rajiformes

The order Rajiformes consists of at least 291 species in four families: 14 species belong to the family Anacanthobatidae, 104 to the Arhynchobatidae, 19 to the Gurgesiellidae, and 154 to the Rajidae. The checklist is annotated and ordered by tribes within families (Table 1, see page 18). Assignments are supported by a combination of a molecular analysis of the mitochondrial genome (Yang *et al.*, unpubl.) and morphological data (Last, Séret & Weigmann, unpubl.). Molecular data provide support for the retention of the 5 tribes of McEachran & Dunn (1998): Amblyrajini, Gurgesiellini, Rajini, as well as the Rostrorajini (Chiquillo *et al.*, 2014) in the subfamily Rajinae; and Arhynchobatini and Riorajini in the Arhynchobatinae. There is also strong evidence for the resurrection of the Pavorajini McEachran, 1984 (modified generic composition) and inclusion of two taxa that are newly erected herein, Bathyrajini (type genus *Bathyraja* Ishiyama, 1958) and Crurirajini (type genus *Cruriraja* Bigelow & Schroeder, 1948).

The family Rajidae now comprises 3 tribes. The Amblyrajini (*sensu* McEachran & Dunn) contained 5 genera (*Amblyraja*, *Breviraja*, *Dactylobatus*, *Leucoraja* and *Rajella*). While no samples of *Dactylobatus* were available, there is strong support for the inclusion of the remaining taxa in this tribe of the Rajidae. The tribe Rajini (*sensu* McEachran & Dunn) of the subfamily Rajinae contained 8 genera (*Anacanthobatis*, *Cruriraja*, *Dipturus*, *Okamejei*, *Raja*, *Rostroraja*, and two un-named taxa). Based on molecular data (Yang *et al.*, unpubl.), the tribe now contains *Dipturus*, *Okamejei*, and *Raja*, as well as 5 additional taxa, *Beringraja*, *Dentiraja*, *Hongeo*, *Spiniraja* and *Zearaja* (all supported by morphological and molecular data). *Dipturus intermedius* (Parnell, 1837) is resurrected from the synonymy of *D. batis* (Linnaeus, 1758) based on Iglésias *et al.* (2010). *Anacanthobatis* and *Cruriraja* are transferred to other families (i.e. Anacanthobatidae and Gurgesiellidae respectively), and *Rostroraja* is transferred to the Rostrorajini. The third tribe of the Rajidae, the Rostrorajini, includes four genera *Malacoraja*, *Neoraja*, *Orbiraja* and *Rostroraja*. *Malacoraja* and *Neoraja* were formerly assigned to the Gurgesiellini (in the Rajinae) by McEachran & Dunn.

The family Arhynchobatidae now comprises 4 tribes. The Arhynchobatini (*sensu* McEachran & Dunn) consisted of 9 genera (*Arhynchobatis*, *Bathyraja*, *Irolita*, *Notoraja*, *Psammobatis*, *Pavoraja*, *Pseudoraja*, *Rhinoraja* and *Sympterygia*). It is now divided to include only the 4 genera *Arhynchobatis*, *Irolita*, *Psammobatis* and *Sympterygia* (all supported by molecular data). The Riorajini (*sensu* McEachran & Dunn) contained *Atlantoraja* and *Rioraja* and this association is well supported. An Indo-Pacific group defined by McEachran (1984) as the Pavorajini is resurrected herein and contains *Brochiraja*, *Insentiraja*, *Notoraja* and *Pavoraja* with very strong molecular support. *Pseudoraja* is provisionally retained in this group, but most likely will be assigned to the family Gurgesiellidae when fresh material is made available for study. *Bathyraja* and *Rhinoraja* are assigned to a newly erected tribe, the Bathyrajini.

The family Gurgesiellidae contains 2 strongly divergent groups based on both molecular and morphological data. The Gurgesiellini contains a single genus (*Gurgesiella*) and 3 species. A newly erected tribe Crurirajini contains *Cruriraja* (assigned to the Rajini by McEachran & Dunn) and *Fenestraja* (assigned to the Gurgesiellini of the subfamily Rajinae by McEachran & Dunn).

Anacanthobatis was assigned to the Rajini (Rajinae) by McEachran & Dunn. The family Anacanthobatidae now includes *Anacanthobatis* and species reassigned to 4 other genera (*Indobatis*, *Schroederobatis*, *Sinobatis* and *Springeria*) based on recent morphological research (Hulley, 1973; Weigmann *et al.*, 2014a; Stehmann & Weigmann, 2016).

Some nominal species included (except for *Raja arctowskii*) as questionable species in Weigmann (2016) could not be definitely assigned to valid taxa. Therefore, the following species are provisionally considered invalid and need further evaluation:

- *Raja alia* Garman, 1899 is considered a *nomen dubium* (Weigmann, 2016) and may be a junior synonym of *Rostroraja ackleyi* (Garman, 1881).
- *Raja arctowskii* Dollo, 1904, based on empty egg capsules, is probably valid and possibly identical to *Bathyraja* sp. 2 *sensu* Stehmann (1985) following Weigmann (2016).
- *Raja rondeleti* Bougis, 1959 is questionably valid (Weigmann, 2016). Considered doubtful according to Compagno (1999, 2005), but valid following Capapé *et al.* (2006). Probably synonymous with either *Raja clavata* (Stehmann & Bürkel, 1984) or *Leucoraja fullonica* (Serena, 2005). See Weigmann (2016) for more information.
- *Anacanthobatis donghaiensis* (Deng, Xiong & Zhan, 1983) is a likely junior synonym of *Sinobatis borneensis* (Séret, 1986). See Weigmann (2016) for more information.
- *Anacanthobatis nanhaiensis* (Meng & Li in Chu *et al.*, 1981) is a likely junior synonym of *Sinobatis borneensis* (Ishihara, 1984; Last & Compagno, 1999; Last & Séret, 2008; White & Last, 2013). See Weigmann (2016) for more information.

Table 1. Annotated checklist for the order Rajiformes

Rajidae

Amblyrajini

<i>Amblyraja doellojuradoi</i> (Pozzi, 1935)
<i>Amblyraja frerichsi</i> (Krefft, 1968) Possibly a junior synonym of <i>A. hyperborea</i> due to strong morphological similarity according to Ebert & Stehmann (2013) and Weigmann (2016).
<i>Amblyraja georgiana</i> (Norman, 1938) Morphologically distinct but confused with <i>A. hyperborea</i> leading to possible misidentification of preliminary molecular data (Last, unpubl.).
<i>Amblyraja hyperborea</i> (Collett, 1879) <i>Raja badia</i> Garman, 1899 and <i>Raja robertsi</i> Hulley, 1970 are considered synonyms (Weigmann, 2016; Last, unpubl.). The literature also contains examples of several valid <i>Amblyraja</i> that have been confused with this species.
<i>Amblyraja jenseni</i> (Bigelow & Schroeder, 1950) Provisionally retained but treated as junior synonym of <i>A. hyperborea</i> by Weigmann (2016) due to strong morphological similarity and preliminary molecular data. Further work is needed to resolve this issue.
<i>Amblyraja radiata</i> (Donovan, 1808)
<i>Amblyraja reversa</i> (Lloyd, 1906) Retained in <i>Amblyraja</i> but generic placement questionable (Last, unpubl.) and needs further investigation.
<i>Amblyraja taaf</i> (Meisner, 1987) Possibly a junior synonym of <i>A. hyperborea</i> due to strong morphological similarity according to Ebert & Stehmann (2013) and Weigmann (2016).
<i>Breviraja claramaculata</i> McEachran & Matheson, 1985
<i>Breviraja colesi</i> Bigelow & Schroeder, 1948
<i>Breviraja mouldi</i> McEachran & Matheson, 1995
<i>Breviraja nigriventralis</i> McEachran & Matheson, 1985
<i>Breviraja spinosa</i> Bigelow & Schroeder, 1950
<i>Dactylobatus armatus</i> Bean & Weed, 1909 Placement of <i>Dactylobatus</i> not yet assessed for molecular data.
<i>Dactylobatus clarkii</i> (Bigelow & Schroeder, 1958)
<i>Leucoraja circularis</i> (Couch, 1838)
<i>Leucoraja compagnoi</i> (Stehmann, 1995)
<i>Leucoraja erinacea</i> (Mitchell, 1825)
<i>Leucoraja fullonica</i> (Linnaeus, 1758)
<i>Leucoraja garmani</i> (Whitley, 1939) <i>Leucoraja garmani</i> contains two valid subspecies: <i>Leucoraja garmani caribbaea</i> (McEachran, 1977) and <i>Leucoraja garmani virginica</i> (McEachran, 1977) (Schmitter-Soto <i>et al.</i> , 2000; McEachran 2002; Moore <i>et al.</i> , 2003; Weigmann, 2016).
<i>Leucoraja lentiginosa</i> (Bigelow & Schroeder, 1951)
<i>Leucoraja leucosticta</i> (Stehmann, 1971)
<i>Leucoraja melitensis</i> (Clark, 1926)

<i>Leucoraja naevus</i> (Müller & Henle, 1841)
<i>Leucoraja ocellata</i> (Mitchill, 1815)
<i>Leucoraja pristispina</i> Last, Stehmann & Séret, 2008
<i>Leucoraja wallacei</i> (Hulley, 1970)
<i>Leucoraja yucatanensis</i> (Bigelow & Schroeder, 1950)
<i>Rajella annandalei</i> (Weber, 1913)
<i>Rajella barnardi</i> (Norman, 1935)
<i>Rajella bathyphila</i> (Holt & Byrne, 1908)
<i>Rajella bigelowi</i> (Stehmann, 1978)
<i>Rajella caudaspinosa</i> (von Bonde & Swart, 1923)
<i>Rajella challengerii</i> Last & Stehmann, 2008
<i>Rajella dissimilis</i> (Hulley, 1970)
<i>Rajella eisenhardti</i> Long & McCosker, 1999 Not included in Compagno (2005).
<i>Rajella fuliginea</i> (Bigelow & Schroeder, 1954)
<i>Rajella fyllae</i> (Lütken, 1887) <i>Breviraja marklei</i> McEachran & Miyake, 1987 is a junior synonym (Stehmann <i>et al.</i> , 2008; Weigmann, 2016).
<i>Rajella kukujevi</i> (Dolganov, 1985)
<i>Rajella leoparda</i> (von Bonde & Swart, 1923) Formerly as <i>R. leopardus</i> but amended to <i>R. leoparda</i> in accordance with ICZN Article 34.2. by Weigmann <i>et al.</i> (2014b). Still spelt <i>R. leopardus</i> by Compagno (2005).
<i>Rajella lintea</i> (Fries, 1838) Generally listed as published in 1838 (Eschmeyer <i>et al.</i> , 2016), but as 1839 in Stehmann (2012) and Weigmann (2016). As <i>Dipturus? linteus</i> by Compagno (2005).
<i>Rajella nigerrima</i> (de Buen, 1960)
<i>Rajella paucispinosa</i> Weigmann, Stehmann & Thiel, 2014
<i>Rajella purpuriventralis</i> (Bigelow & Schroeder, 1962) Placed erroneously in <i>Dipturus</i> Rafinesque, 1810 based on material from the Gulf of Mexico (Weigmann, 2016) following Cotton & Stehmann (pers. comm), but this was based on a misidentification.
<i>Rajella ravidula</i> (Hulley, 1970)
<i>Rajella sadowskii</i> (Kreffft & Stehmann, 1974) Spelt <i>R. sadowskyii</i> by Compagno (2005).

Rajini

<p><i>Beringraja binoculata</i> (Girard, 1855)</p> <p>Initially placed in <i>Beringraja</i> Ishihara, Treloar, Bor, Senou & Jeong, 2012 by Ishihara <i>et al.</i> (2012) with very strong molecular support (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Beringraja cortezensis</i> (McEachran & Miyake, 1988)</p> <p>Newly assigned to <i>Beringraja</i> with strong molecular support (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Beringraja inornata</i> (Jordan & Gilbert, 1881)</p> <p>Newly assigned to <i>Beringraja</i> with strong molecular support (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Beringraja pulchra</i> (Liu, 1932)</p> <p>Type of <i>Beringraja</i>. As <i>Raja 'pulchra'</i>, junior homonym of <i>Raja pulchra</i> Schaffhaeutl, 1863 for fossil dermal tubercles from the Eocene of Bavaria by Compagno (2005).</p>
<p><i>Beringraja rhina</i> (Jordan & Gilbert, 1880)</p> <p>Newly assigned to <i>Beringraja</i> with strong molecular support (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Beringraja stellulata</i> (Jordan & Gilbert, 1880)</p> <p>Newly assigned to <i>Beringraja</i> with strong molecular support (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Dentiraja australis</i> (Macleay, 1884)</p> <p>Recently placed in <i>Dipturus</i> by Last & Stevens (2009) and Weigmann (2016) but now assigned to <i>Dentiraja</i> Whitley, 1940 following new molecular data (Yang <i>et al.</i>, unpubl.). As <i>Okamejei australis</i> by Compagno (2005).</p>
<p><i>Dentiraja cerva</i> (Whitley, 1939)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.). As <i>Okamejei cerva</i> by Compagno (2005).</p>
<p><i>Dentiraja confusa</i> (Last, 2008)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Dentiraja endeavouri</i> (Last, 2008)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Dentiraja falloarga</i> (Last, 2008)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Dentiraja flindersi</i> Last & Gledhill, 2008</p> <p>Reassigned from <i>Dipturus</i> to <i>Dentiraja</i> by Last & Gledhill (2008).</p>
<p><i>Dentiraja healdi</i> (Last, White & Pogonoski, 2008)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Dentiraja lemprieri</i> (Richardson, 1845)</p> <p>Reassigned from <i>Dipturus</i> to <i>Dentiraja</i> by Last & Gledhill (2008). As <i>Okamejei lemprieri</i> by Compagno (2005).</p>
<p><i>Dentiraja oculata</i> (Last, 2008)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.).</p>
<p><i>Dentiraja polyommata</i> (Ogilby, 1910)</p> <p>Newly assigned to <i>Dentiraja</i> based on molecular data (Yang <i>et al.</i>, unpubl.). As <i>Raja polyommata</i> by Compagno (2005).</p>
<p><i>Dipturus acrobelus</i> Last, White & Pogonoski, 2008</p>
<p><i>Dipturus amphispinus</i> Last & Alava, 2013</p>
<p><i>Dipturus apricus</i> Last, White & Pogonoski, 2008</p>
<p><i>Dipturus batis</i> (Linnaeus, 1758)</p> <p>Still treated as a composite species in Weigmann (2016) but with remark that <i>D. batis</i> comprises a small (<i>D. cf. flossada</i>) and a large (<i>D. cf. intermedia</i>) species following Iglésias <i>et al.</i> (2010). As a single species, <i>D. batis</i>, by Compagno (2005) and used herein for (<i>D. cf. flossada</i>).</p>

<i>Dipturus bullisi</i> (Bigelow & Schroeder, 1962)
<i>Dipturus campbelli</i> (Wallace, 1967)
<i>Dipturus canutus</i> Last, 2008 Molecular data (Yang <i>et al.</i> , unpubl.) supports placement in <i>Dipturus</i> rather than <i>Dentiraja</i> .
<i>Dipturus chinensis</i> (Basilewsky, 1855) Considered valid by Cheng & Zhou (1997) in the genus <i>Raja</i> but questionably valid by Weigmann (2016). New data confirm its validity and indicate its assignment to <i>Dipturus</i> (Last, unpubl.). Not included in Compagno (2005).
<i>Dipturus crosnieri</i> (Séret, 1989)
<i>Dipturus doutrei</i> (Cadenat, 1960)
<i>Dipturus ecuadoriensis</i> (Beebe & Tee-Van, 1941)
<i>Dipturus garricki</i> (Bigelow & Schroeder, 1958)
<i>Dipturus gigas</i> (Ishiyama, 1958)
<i>Dipturus grahami</i> Last, 2008
<i>Dipturus gudgeri</i> (Whitley, 1940) As <i>Dipturus?</i> <i>gudgeri</i> by Compagno (2005).
<i>Dipturus innominatus</i> (Garrick & Paul, 1974) As <i>Dipturus?</i> <i>innominatus</i> by Compagno (2005).
<i>Dipturus intermedius</i> (Parnell, 1837) Newly resurrected from the synonymy of <i>Dipturus batis</i> (Linnaeus, 1758) based on Iglésias <i>et al.</i> (2010). Not in Compagno (2005) and still treated as a composite species in <i>D. batis</i> by Weigmann (2016).
<i>Dipturus johannisdavisi</i> (Alcock, 1899) Spelt <i>D. johannisdavesi</i> by Compagno (2005).
<i>Dipturus kwangtungensis</i> (Chu, 1960) Confused with <i>Dipturus chinensis</i> (Basilewsky, 1855) in recent literature.
<i>Dipturus laevis</i> (Mitchill, 1818)
<i>Dipturus lanceostratus</i> (Wallace, 1967) Spelt <i>D. lanceostrata</i> by Compagno (2005).
<i>Dipturus leptocaudus</i> (Krefft & Stehmann, 1975) Spelt <i>D. leptocauda</i> by Compagno (2005).
<i>Dipturus macrocaudus</i> (Ishiyama, 1955) Spelt <i>D. macrocauda</i> by Compagno (2005).
<i>Dipturus melanospilus</i> Last, White & Pogonoski, 2008
<i>Dipturus mennii</i> Gomes & Paragó, 2001 <i>Dipturus diehi</i> Soto & Mincarone 2001 is a junior synonym of <i>D. mennii</i> (Moreira <i>et al.</i> , 2011; Weigmann, 2016).
<i>Dipturus nidarosiensis</i> (Storm, 1881) Probably a composite species in the eastern North Atlantic according to Ebert & Stehmann (2013) and Weigmann (2016). However, at least one form appears widespread with populations from Ireland and South Africa identical based on molecular data (Yang <i>et al.</i> , unpubl.). Authorship indicated as (Collett, 1880) by Compagno (2005).
<i>Dipturus olseni</i> (Bigelow & Schroeder, 1951)
<i>Dipturus oregoni</i> (Bigelow & Schroeder, 1958)

<p><i>Dipturus oxyrinchus</i> (Linnaeus, 1758)</p> <p>Probably a composite species comprising a small and a large species according to Ebert & Stehmann (2013) and Weigmann (2016). Spelt <i>D. oxyrhynchus</i> by Compagno (2005).</p>
<p><i>Dipturus pullopunctatus</i> (Smith, 1964)</p> <p>Spelt <i>D. pullopunctata</i> by Compagno (2005).</p>
<p><i>Dipturus queenslandicus</i> Last, White & Pogonoski, 2008</p>
<p><i>Dipturus springeri</i> (Wallace, 1967)</p>
<p><i>Dipturus stenorhynchus</i> (Wallace, 1967)</p>
<p><i>Dipturus teevani</i> (Bigelow & Schroeder, 1951)</p>
<p><i>Dipturus tengu</i> (Jordan & Fowler, 1903)</p>
<p><i>Dipturus trachydermus</i> (Krefft & Stehmann, 1974)</p> <p>Spelt <i>D. trachyderma</i> by Compagno (2005).</p>
<p><i>Dipturus wengi</i> Séret & Last, 2008</p>
<p><i>Dipturus wuhanlingi</i> Jeong & Nakabo, 2008</p>
<p><i>Hongoe koreana</i> (Jeong & Nakabo, 1997)</p> <p>Distinct morphologically and based on molecular data (Yang <i>et al.</i>, unpubl.) from other rajine skates. As <i>Okamejei? koreana</i> by Compagno (2005).</p>
<p><i>Okamejei acutispina</i> (Ishiyama, 1958)</p>
<p><i>Okamejei arafurensis</i> Last & Gledhill, 2008</p>
<p><i>Okamejei boesemani</i> (Ishihara, 1987)</p>
<p><i>Okamejei cairae</i> Last, Fahmi & Ishihara, 2010</p>
<p><i>Okamejei heemstrai</i> (McEachran & Fechhelm, 1982)</p>
<p><i>Okamejei hollandi</i> (Jordan & Richardson, 1909)</p>
<p><i>Okamejei kenojei</i> (Müller & Henle, 1841)</p>
<p><i>Okamejei leptoura</i> Last & Gledhill, 2008</p>
<p><i>Okamejei meerdervoortii</i> (Bleeker, 1860)</p>
<p><i>Okamejei mengae</i> Jeong, Nakabo & Wu, 2007</p> <p>Possibly a junior synonym of <i>O. hollandi</i> (Weigmann, 2016).</p>
<p><i>Okamejei ornata</i> Weigmann, Stehmann & Thiel, 2015</p>
<p><i>Okamejei schmidti</i> (Ishiyama, 1958)</p>
<p><i>Raja africana</i> Capapé, 1977</p> <p>Listed as valid by McEachran & Séret (2016) but questionable following Weigmann (2016) with the holotype supposedly representing an aberrant <i>R. miraletus</i> or <i>R. straeleni</i>, and a homonym, preoccupied by <i>Raja africana</i> Bloch & Schneider, 1801, which is now considered a junior synonym of a dasyatid <i>Urogymnus asperrimus</i>. Also treated as a questionably valid species by Compagno (2005).</p>
<p><i>Raja asterias</i> Delaroche, 1809</p>
<p><i>Raja brachyura</i> Lafont, 1873</p>
<p><i>Raja clavata</i> Linnaeus, 1758</p>
<p><i>Raja herwigii</i> Krefft, 1965</p>

<i>Raja maderensis</i> Lowe, 1838 Recent molecular results indicate that <i>R. maderensis</i> is possibly conspecific with <i>R. clavata</i> (Ball <i>et al.</i> , 2016).
<i>Raja microocellata</i> Montagu, 1818
<i>Raja miraletus</i> Linnaeus, 1758 Confused with <i>Raja ocellifera</i> Regan, 1906 and <i>Raja parva</i> Last & Séret, 2016 by McEachran & Séret 2016 and others (Last & Séret 2016).
<i>Raja montagui</i> Fowler, 1910
<i>Raja ocellifera</i> Regan, 1906 Frequently confused with <i>Raja miraletus</i> Linnaeus, 1758 but newly resurrected as a valid species by Last & Séret (2016).
<i>Raja parva</i> Last & Séret, 2016 Newly described species frequently confused with <i>Raja miraletus</i> .
<i>Raja pita</i> Fricke & Al-Hassan, 1995 Placed as <i>incertae sedis</i> in the general genus <i>Raja</i> , probably belonging to either <i>Leucoraja</i> Malm 1877 or <i>Rajella</i> Stehmann 1970 (Weigmann <i>et al.</i> , 2015; Weigmann, 2016). As <i>Okamejei pita</i> by Compagno (2005).
<i>Raja polystigma</i> Regan, 1923
<i>Raja radula</i> Delaroche, 1809
<i>Raja straeleni</i> Poll, 1951
<i>Raja undulata</i> Lacepède, 1802
<i>Spiniraja whiteyi</i> (Iredale, 1938) Support for this genus-level taxon based on Last (unpubl.) and Yang <i>et al.</i> (unpubl.).
<i>Zearaja argentinensis</i> (de Astarloa, Mabragana, Hanner & Figueroa, 2008) Placed in <i>Zearaja</i> Whitley, 1939 instead of <i>Dipturus</i> Rafinesque, 1810, based on Yang <i>et al.</i> (unpubl.); see also Weigmann (2016).
<i>Zearaja chilensis</i> (Guichenot, 1848) Possibly a composite species based on molecular analyses (Naylor <i>et al.</i> , 2012b; Vargas-Caro <i>et al.</i> , 2014; Weigmann, 2016). As <i>Dipturus? chilensis</i> by Compagno (2005).
<i>Zearaja maugeana</i> Last & Gledhill, 2007
<i>Zearaja nasuta</i> (Müller & Henle, 1841) As <i>Dipturus? nasutus</i> by Compagno (2005).

Rostrorajini

<i>Malacoraja krefftii</i> (Stehmann, 1978)
<i>Malacoraja obscura</i> Carvalho, Gomes & Gadig, 2005
<i>Malacoraja senta</i> (Garman, 1885)
<i>Malacoraja spinacidermis</i> (Barnard, 1923)
<i>Neoraja africana</i> (Stehmann & Séret, 1983)
<i>Neoraja caerulea</i> (Stehmann, 1976)
<i>Neoraja carolinensis</i> McEachran & Stehmann, 1984
<i>Neoraja iberica</i> Stehmann, Séret, Costa & Baro, 2008 Morphologically distinct from <i>Neoraja caerulea</i> (Stehmann, 1976) but the species are indistinguishable based on their <i>NADH2 gene sequences</i> (Yang <i>et al.</i> , unpubl.).

<i>Neoraja stehmanni</i> (Hulley, 1972)
<i>Orbiraja jensenae</i> (Last & Lim, 2010) Relocated to newly described genus <i>Orbiraja</i> Last, Weigmann & Dumale, 2016 (Last <i>et al.</i> , 2016).
<i>Orbiraja philipi</i> (Lloyd, 1906) Now placed in <i>Orbiraja</i> (Last <i>et al.</i> , 2016) but possibly a junior synonym of <i>O. powelli</i> (Weigmann <i>et al.</i> , 2015; Weigmann, 2016).
<i>Orbiraja powelli</i> (Alcock, 1898) Now placed in <i>Orbiraja</i> (Last <i>et al.</i> , 2016).
<i>Rostroraja ackleyi</i> (Garman, 1881) Provisionally assigned to <i>Rostroraja</i> based on molecular data (Yang <i>et al.</i> , unpubl). Placed as <i>incertae sedis</i> in the general genus <i>Raja</i> by Compagno (2005) and Weigmann (2016).
<i>Rostroraja alba</i> (Lacepède 1803)
<i>Rostroraja bahamensis</i> (Bigelow & Schroeder, 1965) Provisionally assigned to <i>Rostroraja</i> based on molecular data (Yang <i>et al.</i> , unpubl). Placed as <i>incertae sedis</i> in the general genus <i>Raja</i> by Compagno (2005) and Weigmann (2016).
<i>Rostroraja cervigoni</i> (Bigelow & Schroeder, 1964) See comment above for <i>Rostroraja bahamensis</i> .
<i>Rostroraja eglanteria</i> (Bosc, 1800) See comment above for <i>Rostroraja bahamensis</i> .
<i>Rostroraja equatorialis</i> (Jordan & Bollman, 1890) See comment above for <i>Rostroraja bahamensis</i> .
<i>Rostroraja texana</i> (Chandler, 1921) See comment above for <i>Rostroraja bahamensis</i> .
<i>Rostroraja velezi</i> (Chirichigno, 1973) See comment above for <i>Rostroraja bahamensis</i> .

Arhynchobatidae

Arhynchobatini

<i>Arhynchobatis asperrimus</i> Waite, 1909
<i>Irolita waitii</i> (McCulloch, 1911) Spelt <i>I. waitei</i> by Compagno (2005).
<i>Irolita westraliensis</i> Last & Gledhill, 2008
<i>Psammobatis bergi</i> Marini, 1932
<i>Psammobatis extenta</i> (Garman, 1913)
<i>Psammobatis lentiginosa</i> McEachran, 1983
<i>Psammobatis normani</i> McEachran, 1983
<i>Psammobatis parvacauda</i> McEachran, 1983 Unpublished data support the validity of this rare species (Weigmann, 2016).
<i>Psammobatis rudis</i> Günther, 1870
<i>Psammobatis rutrum</i> Jordan, 1891
<i>Psammobatis scobina</i> (Philippi, 1857)

<i>Sympterygia acuta</i> Garman, 1877
<i>Sympterygia bonapartii</i> Müller & Henle, 1841 Spelt <i>S. bonapartei</i> by Compagno (2005).
<i>Sympterygia brevicaudata</i> (Cope, 1877)
<i>Sympterygia lima</i> (Poëppig, 1835)

Bathyrājini

<i>Bathyrāja abyssicola</i> (Gilbert, 1896)
<i>Bathyrāja aguja</i> (Kendall & Radcliffe, 1912)
<i>Bathyrāja albomaculata</i> (Norman, 1937) Compagno (1999) reallocated this species from <i>Bathyrāja</i> Ishiyama, 1958 to <i>Rhinorāja</i> Ishiyama, 1952, but the validity of this move is questionable (Weigmann, 2016; Last, unpubl.).
<i>Bathyrāja aleutica</i> (Gilbert, 1896)
<i>Bathyrāja andriashevi</i> Dolganov, 1983
<i>Bathyrāja bergi</i> Dolganov, 1983
<i>Bathyrāja brachyurops</i> (Fowler, 1910) Probably belongs to the subgenus <i>Arctorāja</i> Ishiyama, 1958 based on clasper characters (Stehmann, pers. comm. in Weigmann, 2016).
<i>Bathyrāja cousseauae</i> Diaz de Astarloa & Mabrágana, 2004 Spelt <i>B. cousseau</i> by Compagno (2005).
<i>Bathyrāja diplotaenia</i> (Ishiyama, 1952)
<i>Bathyrāja eatonii</i> (Günther, 1876)
<i>Bathyrāja fedorovi</i> Dolganov, 1983
<i>Bathyrāja griseocauda</i> (Norman, 1937)
<i>Bathyrāja hesperaficana</i> Stehmann, 1995
<i>Bathyrāja interrupta</i> (Gill & Townsend, 1897) Following Weigmann (2016), as <i>Rhinorāja interrupta</i> by Raschi & McEachran (1991), Compagno (1999, 2005), Dolganov (1999), Sheiko & Fedorov (2000), Hoff (2002) and Orlov (2003), but as <i>Bathyrāja interrupta</i> by Stehmann (1986), McEachran & Dunn (1998), Mecklenburg <i>et al.</i> (2002), Spies <i>et al.</i> (2011) and Page <i>et al.</i> (2013). Spies <i>et al.</i> (2011) provide morphological and molecular support for the placement in <i>Bathyrāja</i> instead of <i>Rhinorāja</i> . <i>Bathyrāja interrupta</i> possibly represents a species complex (Orr <i>et al.</i> , 2011; Spies <i>et al.</i> , 2011; Weigmann, 2016).
<i>Bathyrāja irrasa</i> Hureau & Ozouf-Costaz, 1980
<i>Bathyrāja isiharai</i> Stehmann, 2005
<i>Bathyrāja isotrachys</i> (Günther, 1877)
<i>Bathyrāja kincaidii</i> (Garman, 1908) Treated as junior synonym of <i>B. interrupta</i> by Weigmann (2016) following Ishihara & Ishiyama (1985), Stehmann (1986), Sheiko & Fedorov (2000) and Ishihara (pers. comm.), but as valid species spelt <i>B. kincaidi</i> by Compagno (2005). Further work is needed to resolve this issue.
<i>Bathyrāja leucomelanos</i> Iglésias & Lévy-Hartmann, 2012
<i>Bathyrāja lindbergi</i> Ishiyama & Ishihara, 1977 Valid according to Compagno (1999, 2005), Weigmann (2016) and Last (unpubl.), but a junior synonym of <i>B. matsubarae</i> according to Sheiko & Fedorov (2000).

<i>Bathyrāja longicauda</i> (de Buen, 1959)
<i>Bathyrāja maccaini</i> Springer, 1971
<i>Bathyrāja macloviana</i> (Norman, 1937) Compagno (1999) reallocated this species from <i>Bathyrāja</i> to <i>Rhinoraja</i> , but the validity of this move is questionable (Weigmann, 2016; Last, unpubl.).
<i>Bathyrāja maculata</i> Ishiyama & Ishihara, 1977
<i>Bathyrāja magellanica</i> (Philippi, 1902) Compagno (1999) reallocated this species from <i>Bathyrāja</i> to <i>Rhinoraja</i> , but the validity of this move is questionable (Weigmann, 2016; Last, unpubl.).
<i>Bathyrāja mariposa</i> Stevenson, Orr, Hoff & McEachran, 2004
<i>Bathyrāja matsubarae</i> (Ishiyama, 1952) <i>Bathyrāja caeluronigricans</i> Ishiyama & Ishihara, 1977 is a junior synonym (Ishihara, 1990; Weigmann, 2016; Last, unpubl.).
<i>Bathyrāja meridionalis</i> Stehmann, 1987
<i>Bathyrāja microtrachys</i> (Osburn & Nichols, 1916) Treated as junior synonym of <i>B. trachura</i> by Weigmann (2016) following Castro-Aguirre & Espinosa Pérez (1996) and Mecklenburg <i>et al.</i> (2002), but as valid species by Compagno (2005). Further work is needed to resolve this issue.
<i>Bathyrāja minispinosa</i> Ishiyama & Ishihara, 1977
<i>Bathyrāja multispinis</i> (Norman, 1937) Compagno (1999) reallocated this species from <i>Bathyrāja</i> to <i>Rhinoraja</i> , but the validity of this move is questionable (Weigmann, 2016; Last, unpubl.).
<i>Bathyrāja murrayi</i> (Günther, 1880) Compagno (1999) reallocated this species from <i>Bathyrāja</i> to <i>Rhinoraja</i> , but the validity of this move is questionable (Weigmann, 2016; Last, unpubl.).
<i>Bathyrāja notoroensis</i> Ishiyama & Ishihara, 1977 Treated as junior synonym of <i>B. matsubarae</i> by Weigmann (2016) following Ishihara (1990), but as valid species by Compagno (2005). Further work is needed to resolve this issue (Last, unpubl.).
<i>Bathyrāja pacifica</i> Last, Stewart & Séret, 2016 Newly described species.
<i>Bathyrāja pallida</i> (Forster, 1967)
<i>Bathyrāja panthera</i> Orr, Stevenson, Hoff, Spies & McEachran, 2011 Belongs to the subgenus <i>Arctoraja</i> (Orr <i>et al.</i> , 2011; Weigmann, 2016).
<i>Bathyrāja papilionifera</i> Stehmann, 1985
<i>Bathyrāja parmifera</i> (Bean, 1881) Belongs to the subgenus <i>Arctoraja</i> (Orr <i>et al.</i> , 2011; Weigmann, 2016). <i>Rhinoraja obtusa</i> (Gill & Townsend, 1897) and <i>Rhinoraja rosispinis</i> (Gill & Townsend, 1897) are junior synonyms (Stehmann 1986; Mecklenburg <i>et al.</i> , 2002; Orr <i>et al.</i> , 2011; Weigmann, 2016).
<i>Bathyrāja peruana</i> McEachran & Miyake, 1984
<i>Bathyrāja richardsoni</i> (Garrick, 1961)
<i>Bathyrāja scaphiops</i> (Norman, 1937)
<i>Bathyrāja schroederi</i> (Krefft, 1968)
<i>Bathyrāja shuntovi</i> Dolganov, 1985

<i>Bathyrāja simoterus</i> (Ishiyama, 1967) Belongs to the subgenus <i>Arctoraja</i> (Orr <i>et al.</i> , 2011; Weigmann, 2016). According to Weigmann (2016), a junior synonym of <i>B. parmifera</i> following Dolganov (2001) and Dolganov & Korolev (2006), but valid according to Stehmann (1986), McEachran & Dunn (1998), Compagno (1999, 2005), Nakabo (2002, 2013), Orr <i>et al.</i> (2011) and Spies <i>et al.</i> (2011).
<i>Bathyrāja smirmovi</i> (Soldatov & Pavlenko, 1915) See comment above for <i>Bathyrāja simoterus</i> .
<i>Bathyrāja smithii</i> (Müller & Henle, 1841)
<i>Bathyrāja spinicauda</i> (Jensen, 1914)
<i>Bathyrāja spinosissima</i> (Beebe & Tee-Van, 1941)
<i>Bathyrāja taranetzi</i> (Dolganov, 1983) As <i>Rhinoraja taranetzi</i> by Weigmann (2016). Following Weigmann (2016), as <i>Bathyrāja taranetzi</i> by Mecklenburg <i>et al.</i> (2002) and Page <i>et al.</i> (2013), but as <i>Rhinoraja taranetzi</i> by Raschi & McEachran (1991), McEachran & Dunn (1998), Compagno (1999, 2005), Dolganov (1999), Sheiko & Fedorov (2000), Stevenson <i>et al.</i> (2004), Spies <i>et al.</i> (2011) and Ishihara (pers. comm.). <i>Rhinoraja longi</i> Raschi & McEachran, 1991 is a junior synonym (Mecklenburg <i>et al.</i> , 2002; Stevenson <i>et al.</i> , 2004; Weigmann, 2016).
<i>Bathyrāja trachouros</i> (Ishiyama, 1958)
<i>Bathyrāja trachura</i> (Gilbert, 1892)
<i>Bathyrāja tunae</i> Stehmann, 2005 Possibly conspecific with the wide-ranging <i>Bathyrāja richardsoni</i> (Garrick, 1961) (Last, unpubl.).
<i>Bathyrāja tzinovskii</i> Dolganov, 1983
<i>Bathyrāja violacea</i> (Suvorov, 1935)
<i>Rhinoraja kujiensis</i> (Tanaka, 1916) <i>Bathyrāja</i> and <i>Rhinoraja</i> were considered among the most problematic of the chondrichthyan genera by Naylor <i>et al.</i> (2012a). All species assigned to <i>Rhinoraja</i> were found to be interspersed among the 20 species of <i>Bathyrāja</i> included in their molecular analysis. However, none of the species originally assigned to <i>Rhinoraja</i> by Ishiyama (1952, 58) was included. The separation of <i>Rhinoraja</i> from <i>Bathyrāja</i> is still doubtful as the only diagnostic character is the presence or absence of a segmented rostral cartilage (Ishihara, pers. comm. in Weigmann, 2016). For the time being, <i>Rhinoraja</i> is retained pending further investigation.
<i>Rhinoraja longicauda</i> Ishiyama, 1952 See comment above.
<i>Rhinoraja odai</i> Ishiyama, 1958 See comment above.

Riorajini

<i>Atlantoraja castelnaui</i> (Miranda Ribeiro, 1907) Author name indicated as Ribeiro by Compagno (2005).
<i>Atlantoraja cyclophora</i> (Regan, 1903)
<i>Atlantoraja platana</i> (Günther, 1880)
<i>Rioraja agassizii</i> (Müller & Henle, 1841)

Pavorajini

<i>Brochiraja aenigma</i> Last & McEachran, 2006
<i>Brochiraja albilabiata</i> Last & McEachran, 2006
<i>Brochiraja asperula</i> (Garrick & Paul, 1974) Still as <i>Notoraja asperula</i> by Compagno (2005).
<i>Brochiraja heuresa</i> Last & Séret, 2012
<i>Brochiraja leviveneta</i> Last & McEachran, 2006
<i>Brochiraja microspinifera</i> Last & McEachran, 2006
<i>Brochiraja spinifera</i> (Garrick & Paul, 1974) Still as <i>Notoraja spinifera</i> by Compagno (2005).
<i>Brochiraja vittacauda</i> Last & Séret, 2012
<i>Insentiraja laxipella</i> Yearsley & Last, 1992 As <i>Notoraja laxipella</i> by Compagno (2005). Status of <i>Insentiraja</i> needs further investigation (Last, unpubl.).
<i>Insentiraja subtilispinosa</i> (Stehmann, 1989) As <i>Notoraja subtilispinosa</i> with date of the original publication given as 1985 by Compagno (2005).
<i>Notoraja alisae</i> Séret & Last, 2012
<i>Notoraja azurea</i> McEachran & Last, 2008
<i>Notoraja fijiensis</i> Séret & Last, 2012
<i>Notoraja hirticauda</i> Last & McEachran, 2006
<i>Notoraja inusitata</i> Séret & Last, 2012
<i>Notoraja lira</i> McEachran & Last, 2008
<i>Notoraja longiventralis</i> Séret & Last, 2012
<i>Notoraja ochroderma</i> McEachran & Last, 1994
<i>Notoraja sapphira</i> Séret & Last, 2009
<i>Notoraja sticta</i> McEachran & Last, 2008
<i>Notoraja tobitukai</i> (Hiyama, 1940)
<i>Pavoraja alleni</i> McEachran & Fechhelm, 1982
<i>Pavoraja arenaria</i> Last, Mallick & Yearsley, 2008
<i>Pavoraja mosaica</i> Last, Mallick & Yearsley, 2008
<i>Pavoraja nitida</i> (Günther, 1880)
<i>Pavoraja pseudonitida</i> Last, Mallick & Yearsley, 2008
<i>Pavoraja umbrosa</i> Last, Mallick & Yearsley, 2008
? <i>Pseudoraja fischeri</i> Bigelow & Schroeder, 1954

Gurgesiellidae

Crurirajini

<i>Cruriraja andamanica</i> (Lloyd, 1909)
<i>Cruriraja atlantis</i> Bigelow & Schroeder, 1948
<i>Cruriraja cadenati</i> Bigelow & Schroeder, 1962
<i>Cruriraja durbanensis</i> (von Bonde & Swart, 1923)
<i>Cruriraja hulleyi</i> Aschliman, Ebert & Compagno, 2010
<i>Cruriraja parcomaculata</i> (von Bonde & Swart, 1923) <i>Cruriraja triangularis</i> Smith, 1964 is a junior synonym (Compagno & Ebert, 2007; Aschliman <i>et al.</i> , 2010; Weigmann, 2016).
<i>Cruriraja poeyi</i> Bigelow & Schroeder, 1948
<i>Cruriraja rugosa</i> Bigelow & Schroeder, 1958
<i>Fenestraja atripinna</i> (Bigelow & Schroeder, 1950)
<i>Fenestraja cubensis</i> (Bigelow & Schroeder, 1950)
<i>Fenestraja ishiyamai</i> (Bigelow & Schroeder, 1962)
<i>Fenestraja maceachrani</i> (Séret, 1989)
<i>Fenestraja mamillidens</i> (Alcock, 1889)
<i>Fenestraja plutonia</i> (Garman, 1881)
<i>Fenestraja sibogae</i> (Weber, 1913)
<i>Fenestraja sinusmexicanus</i> (Bigelow & Schroeder, 1950)

Gurgesiellini

<i>Gurgesiella atlantica</i> (Bigelow & Schroeder, 1962)
<i>Gurgesiella dorsalifera</i> McEachran & Compagno, 1980
<i>Gurgesiella furvescens</i> de Buen, 1959

Anacanthobatidae

Anacanthobatini

<i>Anacanthobatis marmorata</i> (von Bonde & Swart, 1923) Formerly as <i>A. marmoratus</i> but amended to <i>A. marmorata</i> in accordance with ICZN Article 34.2. by Weigmann <i>et al.</i> (2014a). Date of the original description indicated as 1924 by Compagno (2005).
<i>Indobatis ori</i> (Wallace, 1967) Still placed in <i>Anacanthobatis</i> by Compagno (2005).
<i>Schroederobatis americana</i> (Bigelow & Schroeder, 1962) As <i>Anacanthobatis americanus</i> by Compagno (2005). Still placed in the subgenus <i>Schroederobatis</i> Hulley, 1973 of genus <i>Anacanthobatis</i> von Bonde & Swart, 1923 by Weigmann <i>et al.</i> (2014a), Weigmann (2016), Weigmann & Stehmann (2016) and Stehmann & Weigmann (2016). Specific name amended from <i>americanus</i> to <i>americana</i> in accordance with ICZN Article 34.2. by Weigmann <i>et al.</i> (2014a).
<i>Sinobatis andamanensis</i> Last & Bussarawit, 2016 Newly described species.
<i>Sinobatis borneensis</i> (Chan, 1965) As <i>Anacanthobatis borneensis</i> by Compagno (2005), but assigned to <i>Sinobatis</i> by Last & Séret (2008).

<p><i>Sinobatis brevicauda</i> Weigmann & Stehmann, 2016 Newly described species.</p>
<p><i>Sinobatis bulbicauda</i> Last & Séret, 2008</p>
<p><i>Sinobatis caerulea</i> Last & Séret, 2008</p>
<p><i>Sinobatis filicauda</i> Last & Séret, 2008</p>
<p><i>Sinobatis kotlyari</i> Stehmann & Weigmann, 2016 Newly described species, not included in the <i>Rays of the World</i> family contribution Anacanthobatidae (Séret <i>et al.</i>, in press).</p>
<p><i>Sinobatis melanosoma</i> (Chan, 1965) As <i>Anacanthobatis melanosoma</i> by Compagno (2005), but provisionally assigned to <i>Sinobatis</i> by Last & Séret (2008), which was supported by the examination of the claspers of a subadult male by Weigmann <i>et al.</i> (2014a).</p>
<p><i>Sinobatis stenosoma</i> (Li & Hu, in Chu, Meng, Hu & Li, 1982) As <i>Anacanthobatis stenosoma</i> by Compagno (2005). As questionably valid species in the general genus <i>Anacanthobatis</i> by Weigmann (2016) based on Ishihara & Ishiyama (1986), Séret (1986), and Last & Compagno (1999), who treated the species as possibly invalid, and White & Last (2013), who treated it as questionably valid.</p>
<p><i>Springeria folirostris</i> Bigelow & Schroeder, 1951 As <i>Anacanthobatis folirostris</i> by Compagno (2005). Still placed in the subgenus <i>Springeria</i> Bigelow & Schroeder, 1951 of genus <i>Anacanthobatis</i> by Weigmann <i>et al.</i> (2014a), Weigmann (2016), Weigmann & Stehmann (2016) and Stehmann & Weigmann (2016).</p>
<p><i>Springeria longirostris</i> (Bigelow & Schroeder, 1962) As <i>Anacanthobatis longirostris</i> by Compagno (2005). Still placed in the subgenus <i>Springeria</i> of genus <i>Anacanthobatis</i> by Weigmann <i>et al.</i> (2014a), Weigmann (2016), Weigmann & Stehmann (2016) and Stehmann & Weigmann (2016).</p>

Acknowledgements

This paper is a supplement to the skate component of the *Rays of the World* book. Major input to these chapters was provided by B. Séret, M. Stehmann and J. McEachran. The project was co-funded by a National Science Foundation (NSF; <http://www.nsf.gov>) grant (Jaws and Backbone: Chondrichthyan Phylogeny and a Spine for the Vertebrate Tree of Life (CTOL); DEB-01132229), and was also supported by CSIRO's Wealth from Oceans (WFO) Flagship and CSIRO National Facilities and Collections. Technical papers and unpublished data generated by the CTOL team provided valuable taxonomic and nomenclatural insights and we thank G. Naylor and S. Corrigan (College of Charleston) for their input. PL thanks D. Gledhill, L. Joseph and A. Young (CSIRO National Facilities and Collections), and D. Smith (CSIRO Oceans & Atmosphere), for providing support for this research. C. Devine, J. Pogonoski and A. Graham (CSIRO Australian National Fish Collection, Hobart) all provided assistance to facets of this project. SW is grateful to R. Thiel and I. Eidus (ZMH) for granting access to ZMH specimens and photo- and radiography facilities at ZMH, as well as help with radiography and collection database. The authors thank W. Eschmeyer (CAS), R. Fricke (SMNS) and R. van der Laan (Hogeschool Utrecht), for making available a wealth of valuable data in their Catalog of Fishes (Eschmeyer *et al.*, 2016).

References

- Aschliman, N.C., Ebert, D.A. & Compagno, L.J.V. (2010) A New Legskate (Rajoidei: Genus *Cruriraja*) from Southern Africa. *Copeia*, 2010 (3), 364–372.
- Aschliman, N.C., Nishida, M., Miya, M., Inoue, J.G., Rosana, K.M. & Naylor, G.J.P. (2012) Body plan convergence in the evolution of skates and rays (Chondrichthyes: Batoidea). *Molecular Phylogenetics and Evolution*, 63 (1), 28–42.
- Ball, R.E., Serra-Pereira, B., Ellis, J., Genner, M.J., Iglésias, S., Johnson, A., Jones, C.S., Leslie, R., Lewis, J., Mariani, S., Menezes, G., Neat, F.C., Noble, L.R., Sims, D.W. & Griffiths, A.M. (2016) Resolving taxonomic uncertainty in vulnerable elasmobranchs: are the Madeira skate (*Raja maderensis*) and the thornback ray (*Raja clavata*) distinct species? *Conservation Genetics*, 17 (3), 565–576.
- Bigelow, H.B. & Schroeder, W.C. (1948) New genera and species of batoid fishes. *Journal of Marine Research*, 7, 543–566.
- von Bonde, C. & Swart, D.B. (1923) The Platosomia (Skates and Rays) collected by the S.S. "Pickle". *Reports of the Fisheries and Marine Biological Survey of the Union of South Africa*, 3, 1–22.
- Capapé, C., Guélorget, O., Vergne, Y., Marqués, A. & Quignard, J.P. (2006) Skates and rays (Chondrichthyes) from waters off the Languedocian coast (southern France, northern Mediterranean): a historical survey and present status. *Annales, Series historia naturalis*, 16 (2), 165–178.
- Carvalho, M.R. de, Gomes, U.L. & McEachran, J.D. (2006) First report and description of a legskate, genus *Cruriraja*, from the southwestern Atlantic Ocean (Chondrichthyes: Rajidae). *Cybium* 30 (4), 343–354.
- Castro-Aguirre, J.L. & Espinosa Pérez, H. (1996) *Listados Faunísticos de México. VII. Catálogo Sistemático de las Rayas y Especies Afines de México (Chondrichthyes: Elasmobranchii: Rajiformes: Batoideiomorpha)*. Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, 75 pp.
- Cheng, Q.T., Zhou, C.W., Wang, J.X., Yang, Q. & Lin, H.Y. (1997) *The Fishes of Shandong Province*. Shandong Science and Technology Press, Jinan, 549 pp. [in Chinese]
- Chiquillo, K.L., Ebert, D.A., Slager, C.J., Crow, K.D. (2014) The secret of the mermaid's purse: phylogenetic affinities within the Rajidae and the evolution of a novel reproductive strategy in skates. *Molecular Phylogenetics and Evolution*, 75, 245–251.
- Compagno, L.J.V. (1999) Checklist of Living Elasmobranchs. In: Hamlett, W.C. (Ed.) *Sharks, Skates, and Rays – The Biology of Elasmobranch Fishes*. The Johns Hopkins University Press, Baltimore, pp. 471–498.
- Compagno, L.J.V. (2005) Checklist of Living Chondrichthyes. In: Hamlett, W.C. (Ed.) *Reproductive Biology and Phylogeny of Chondrichthyes – Sharks, Batoids and Chimaeras*. Science Publishers, Inc., Enfield, pp. 503–548.
- Compagno, L.J.V. & Ebert, D.A. (2007) Southern African skate biodiversity and distribution. *Environmental Biology of Fishes*, 80 (2–3), 125–145.
- Dolganov, V.N. (1999) Geographic and bathymetric distribution of the skates of the family Rajidae in the far eastern seas of Russia and the adjacent waters. *Voprosy Ikhtiologii*, 39 (3), 428–430. [In Russian, English translation in *Journal of Ichthyology*, 39 (4), 340–342]
- Dolganov, V.N. (2001) Origin and distribution of skates of the suborder Rajoidei of the Far East seas of Russia. *Voprosy Ikhtiologii*, 41 (3), 304–311. [In Russian, English translation in *Journal of Ichthyology*, 41 (5), 354–361]
- Dolganov, V.N. & Korolev, M.R. (2006) On validity of the skate species from *parmifera* group of the genus *Bathyraja* (Rajidae, Rajoidei). *Izvestiya Tikhookeanskogo nauchno-issledovatel'skogo instituta rybnogo khozyajstva i okeanografii*, 147, 179–182. [In Russian with English abstract]
- Ebert, D.A. & Compagno, L.J.V. (2007) Biodiversity and systematics of skates (Chondrichthyes: Rajiformes: Rajoidei). *Environmental Biology of Fishes*, 80 (2/3), 111–124.

- Ebert, D.A. & van Hees, K.E. (2015) Beyond *Jaws*: rediscovering the 'lost sharks' of southern Africa. *African Journal of Marine Science*, 37 (2), 141–156.
- Ebert, D.A. & Stehmann, M.F.W. (2013) *Sharks, batoids, and chimaeras of the North Atlantic*. FAO Species Catalogue for Fishery Purposes, 7. FAO, Rome, 523 pp.
- Eschmeyer, W.N., Fricke, R. & van der Laan, R. (Eds.) (2016) Catalog of Fishes: genera, species, references. Available at <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (last accessed 12 November 2016).
- Gomon, M.F., Glover, J.C.M. & Kuitert, R.H. (1994) *The Fishes of Australia's South Coast*. State Print, Adelaide, 992 pp.
- Hoese, D.F., Bray, D.J., Allen, G.R., Paxton, J.R., Wells, A. & Beesley, P.L. (Eds.) (2006) *Zoological Catalogue of Australia Vol. 35, Parts 1–3. Fishes*. Zoological Catalogue of Australia Series. 3 vols. 2248 pp.
- Hoff, G.R. (2002) New records of the Aleutian skate, *Bathyraja aleutica* from northern California. *California Fish and Game*, 88 (3), 145–148.
- Hulley, P.A. (1972a) The origin, interrelationships and distribution of southern African Rajidae (Chondrichthyes, Batoidei). *Annals of the South African Museum*, 60, 1–103.
- Hulley, P.A. (1972b) The family Gurgesiellidae (Chondrichthyes, Batoidei), with reference to *Pseudoraja atlantica* Bigelow and Schroeder. *Copeia*, 1972 (2), 356–359.
- Hulley, P.A. (1973) Interrelationships within the Anacanthobatidae (Chondrichthyes, Rajoidea), with a description of the lectotype of *Anacanthobatis marmoratus* von Bonde & Swart, 1923. *Annals of the South African Museum*, 62(4), 131–158.
- Hulley, P.A. (1986) Rajidae. In: Smith, M.M. & Heemstra, P.C. (Eds.) *Smiths' Sea Fishes*. First edition. Springer-Verlag, Berlin, pp. 115–127.
- Iglésias, S.P., Toulhoat, L. & Sellos, D.Y. (2010) Taxonomic confusion and market mislabelling of threatened skates: important consequences for their conservation status. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 20 (3), 319–333.
- Ishihara, H. (1984) Second record of the rare skate *Anacanthobatis borneensis* from the East China Sea. *Japanese Journal of Ichthyology*, 30 (4), 448–451.
- Ishihara, H. (1990) *Studies on the systematics and resources of the skates (Rajidae) in the North Pacific Ocean*. Unpublished PhD Thesis, Tokyo University.
- Ishihara, H. & Ishiyama, R. (1985) Two New North Pacific Skates (Rajidae) and a Revised Key to *Bathyraja* in the Area. *Japanese Journal of Ichthyology*, 32 (2), 143–179.
- Ishihara, H. & Ishiyama, R. (1986) Systematics and distribution of the skates of the North Pacific (Chondrichthyes, Rajoidei). In: Uyeno, T., Arai, R., Taniuchi, T. & Matsuura, K. (Eds.) *Indo-Pacific fish biology: proceedings of the Second International Conference on Indo-Pacific Fishes*. Ichthyological Society of Japan, Tokyo, pp. 269–280.
- Ishihara, H., Treloar, M., Bor, P.H.F., Senou, H. & Jeong, C.H. (2012) The Comparative Morphology of Skate Egg Capsules (Chondrichthyes: Elasmobranchii: Rajiformes). *Bulletin of the Kanagawa Prefectural Museum (Natural Science)*, 41, 9–25.
- Ishiyama, R. (1952) Studies on the rays and skates belonging to the family Rajidae, found in Japan and adjacent regions. 4. A revision of three genera of Japanese rajids with descriptions of one new genus and four new species mostly occurred in northern Japan. *Journal of the Shimonoseki College of Fisheries*, 2 (2), 1–34.
- Ishiyama, R. (1958) Studies on the rajid fishes (Rajidae) found in the waters around Japan. *Journal of the Shimonoseki College of Fisheries*, 7 (2/3), 193–394.
- Jeong, C.H., Nakabo, T. & Wu, H.L. (2007) A new species of skate (Chondrichthyes: Rajidae), *Okamejei mengae* from the South China Sea. *Korean Journal of Ichthyology*, 19 (1), 57–65.
- van der Laan, R., Eschmeyer, W.N. & Fricke, R. (2014) Family-group names of Recent fishes. *Zootaxa*, 3882 (1), 1–230.
- Last, P.R. & Compagno, L.J.V. (1999) Anacanthobatidae, Legskates. In: Carpenter, K.E. & Niem, V.H. (Eds.) *The living marine resources of the Western Central Pacific. Vol. 3: Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae)*. FAO Species Identification Guide for Fishery Purposes. FAO, Rome, pp. 1462–1466.
- Last, P.R. & Gledhill, D.C. (2008) A new skate of the genus *Dentiraja* (Rajoidei: Rajidae) from southern Australia, pp. 109–118. In: Last, P.R., White, W.T., Pogonoski, J.J. & Gledhill, D.C. (Eds.) *Descriptions of New Australian Skates (Batoidea: Rajoidei)*. CSIRO Marine & Atmospheric Research Paper, 021, 181 pp.
- Last, P.R. & Séret, B. (2008) Three new legskates of the genus *Sinobatis* (Rajoidei: Anacanthobatidae) from the Indo-West Pacific. *Zootaxa*, 1671, 33–58.
- Last, P.R. & Séret, B. (2016) A new Eastern Central Atlantic skate *Raja parva* sp. nov. (Rajoidei: Rajidae) belonging to the *Raja miraletus* species complex. *Zootaxa*, 4147 (4), 477–489.
- Last, P.R. & Stevens, J.D. (1994) *Sharks and Rays of Australia*. CSIRO Publishing, Melbourne, 513 pp.
- Last, P.R. & Stevens, J.D. (2009) *Sharks and Rays of Australia*. Second edition. CSIRO Publishing, Melbourne, 644 pp.
- Last, P.R. & Yearsley, G.K. (2002) Zoogeography and relationships of Australasian skates (Chondrichthyes: Rajidae). *Journal of Biogeography*, 29 (12), 1627–1641.

- Last, P.R., Weigmann, S. & Dumale, D. (2016) A new skate genus *Orbiraja* (Rajiformes: Rajidae) from the Indo-West Pacific. *Zootaxa*, 4184 (1). 52–62.
- Last, P.R., White, W.T., Carvalho, M.R. de, Séret, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.) (in press a) *Rays of the World*. CSIRO Publishing, Melbourne.
- Last, P.R., Stehmann, M.F.W., Séret, B. & Weigmann, S. (in press b) Softnose Skates, Family Arhynchobatidae. In: Last, P.R., White, W.T., Carvalho, M.R. de, Séret, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.) *Rays of the World*. CSIRO Publishing, Melbourne, pp. 364–472.
- Last, P.R., Séret, B., Stehmann, M.F.W. & Weigmann, S. (in press c) Skates, Family Rajidae. In: Last, P.R., White, W.T., Carvalho, M.R. de, Séret, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.) *Rays of the World*. CSIRO Publishing, Melbourne, pp. 204–363.
- McEachran, J.D. (1984) Anatomical Investigations of the New Zealand Skates *Bathyraja asperula* and *B. spinifera*, with an Evaluation of their Classification within the Rajoidei (Chondrichthyes). *Copeia*, 1984 (1), 45–58.
- McEachran, J.D. (2002) Skates. Family Rajidae. In: Collette, B.B. & Klein-MacPhee, G. (Eds.) *Bigelow and Schroeder's fishes of the Gulf of Maine*. Smithsonian Institution Press, Washington, D.C., pp. 60–75.
- McEachran, J.D. & Compagno, L.J.V. (1979) A further description of *Gurgesiella furvescens* with comments on the interrelationships of Gurgesiellidae and Pseudorajidae (Pisces, Rajoidei). *Bulletin of Marine Science*, 29 (4), 530–553.
- McEachran, J.D. & Dunn, K.A. (1998) Phylogenetic analysis of skates, a morphologically conservative clade of elasmobranchs (Chondrichthyes: Rajidae). *Copeia*, 1998 (2), 271–290.
- McEachran, J.D. & Séret, B. (2016) Rajidae, Skates. In: Carpenter, K.E. & Angelis, N. de (Eds.) *The living marine resources of the Eastern Central Atlantic. Vol. 2: Bivalves, gastropods, hagfishes, sharks, batoid fishes, and chimaeras*. FAO Species Identification Guide for Fishery Purposes. FAO, Rome, pp. 1379–1403.
- Mecklenburg, C.W., Mecklenburg, T.A. & Thorsteinson, L.K. (2002) *Fishes of Alaska*. American Fisheries Society, Bethesda, 1037 pp.
- Moore, J.A., Hartel, K.E., Craddock, J.E. & Galbraith, J.K. (2003) An annotated list of deepwater fishes from off the New England region, with new area records. *Northeastern Naturalist*, 10 (2), 159–248.
- Moreira, R.A., Gomes, U.L. & Carvalho, M.R. de (2011) Morphological description of *Dipturus mennii* (Chondrichthyes: Elasmobranchii: Rajidae) and its differentiation from *Dipturus trachyderma*. *Zoologia*, 28 (1), 97–111.
- Nakabo, T. (Ed.) (2002) *Fishes of Japan with pictorial keys to the species, English edition. Vol. 1*. Tokai University Press, Hadano-shi, 866 pp.
- Nakabo, T. (Ed.) (2013) *Fishes of Japan with pictorial keys to the species, Vol. 1, 3rd edition*. Tokai University Press, Hadano-shi, 864 pp. [in Japanese]
- Naylor, G.J.P., Caira, J.N., Jensen, K., Rosana, K.A.M., Straube, N. & Lakner, C. (2012a) Elasmobranch Phylogeny: A Mitochondrial Estimate Based on 595 Species. In: Carrier, J.C., Musick, J.A. & Heithaus, M.R. (Eds.) *The Biology of Sharks and Their Relatives*. CRC Press, Boca Raton, pp. 31–56.
- Naylor, G.J.P., Caira, J.N., Jensen, K., Rosana, K.A.M., White, W.T. & Last, P.R. (2012b) A DNA Sequence-Based Approach to the Identification of Shark and Ray Species and its Implications for Global Elasmobranch Diversity and Parasitology. *Bulletin of the American Museum of Natural History*, 367, 1–262.
- Naylor, G.J.P., Yang, L., Corrigan, S. & Carvalho, M.R. de (in press) 2. Phylogeny and Classification of Rays. In: Last, P.R., White, W.T., Carvalho, M.R. de, Séret, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.) *Rays of the World*, CSIRO Publishing, Melbourne.
- Orlov, A.M. (2003) Diets, feeding habits, and trophic relations of six deep-benthic skates (Rajidae) in the western Bering Sea. *Aqua, International Journal of Ichthyology*, 7 (2), 45–60.
- Orr, J.W., Stevenson, D.E., Hoff, G.R., Spies, I. & McEachran, J.D. (2011) *Bathyraja panthera*, a new species of skate (Rajidae: Arhynchobatinae) from the western Aleutian Islands, and resurrection of the subgenus *Arctoraja* Ishiyama. *NOAA Professional Paper NMFS*, 11, 1–50.
- Page, L.M., Espinosa-Pérez, H., Findley, L.D., Gilbert, C.R., Lea, R.N., Mandrak, N.E., Mayden, R.L. & Nelson, J.S. (2013) *Common and Scientific Names of Fishes from the United States, Canada, and Mexico. 7th Edition*. American Fisheries Society Special Publication 34, 243 pp.
- Paxton, J.R., Hoese, D.F., Allen, G.R. & Hanley, J.E. (1989) Zoological Catalogue of Australia, Vol. 7: Pisces, Petromyzontidae to Carangidae. Australian Government Publishing Service, Canberra. *Zoological Catalogue of Australia*, 7, 1–665.
- Raschi, W.G. & McEachran, J.D. (1991) *Rhinoraja longi*, a new species of skate from the outer Aleutian Islands, with comments on the status of *Rhinoraja* (Chondrichthyes, Rajoidei). *Canadian Journal of Zoology*, 69 (7), 1889–1903.
- Schmitter-Soto, J.J., Vásquez-Yeomans, L., Aguilar-Perera, A., Curiel-Mondragón, C., Caballero-Vázquez, J.A. (2000) Lista de peces marinos del Caribe mexicano. *Anales del Instituto de Biología Universidad Nacional Autónoma de México (Serie Zoológica)*, 71 (2), 143–177.

- Serena, F. (2005) *Field identification guide to the sharks and rays of the Mediterranean and Black Sea*. FAO Species Identification Guide for Fishery Purposes. FAO, Rome, 97 pp.
- Séret, B. (1986) Deep water skates of Madagascar. Part I. Anacanthobatidae (Pisces, Chondrichthyes, Batoidea), second record of the skate *Anacanthobatis ori* (Wallace, 1967) from off Madagascar. *Cybium*, 10 (4), 307–326.
- Séret, B., Last, P.R., Weigmann, S. & Stehmann, M.F.W. (in press) Legskates, Family Anacanthobatidae. In: Last, P.R., White, W.T., Carvalho, M.R. de, Séret, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.) *Rays of the World*. CSIRO Publishing, Melbourne, pp. 494–508.
- Sheiko, B.A. & Fedorov, V.V. (2000) Part 1. In: Artuykhin, Y.B. & Sheyko, B.A. (Eds.) *Catalog of the Vertebrates of Kamchatka and Adjacent Waters*. Rossiyskaya Akademiya Nauk, Moscow, pp. 7–69. [In Russian with English abstract]
- Spies, I.B., Stevenson, D.E., Orr, J.W. & Hoff, G.R. (2011) Molecular systematics of the skate subgenus *Arctoraja* (*Bathyraja*: Rajidae) and support for an undescribed species, the leopard skate, with comments on the phylogenetics of *Bathyraja*. *Ichthyological Research*, 58 (1), 77–83.
- Stehmann, M.F.W. (1985) Ergebnisse der Forschungsreisen des FFS “Walter Herwig” nach Südamerika. LXIV. *Bathyraja papilionifera* sp. n. (Pisces, Batoidea, Rajidae), eine weitere neue Rochenart aus dem Südwestatlantik vom nordargentinischen Kontinentalabhang. *Archiv für Fischereiwissenschaft*, 36 (1/2), 195–211.
- Stehmann, M.F.W. (1986) Notes on the systematics of the Rajid genus *Bathyraja* and its distribution in the world oceans. In: Uyeno, T., Arai, R., Taniuchi, T. & Matsuura, K. (Eds.) *Indo-Pacific fish biology: proceedings of the Second International Conference on Indo-Pacific Fishes*. Ichthyological Society of Japan, Tokyo, pp. 261–268.
- Stehmann, M.F.W. (2012) Complementary redescription of *Raja lintea* Fries, 1839 (Elasmobranchii, Rajidae) and its revised generic assignment. *Zootaxa*, 3331, 44–68.
- Stehmann, M. & Bürkel, D.L. (1984) Rajidae. In: Whitehead, P.J.P., Bauchot, M.-L., Hureau, J.-C., Nielsen, J. & Tortonese, E. (Eds.) *Fishes of the North-Eastern Atlantic and the Mediterranean, Vol. 1*. UNESCO, Paris, pp. 163–196.
- Stehmann, M.F.W. & Weigmann, S. (2016) A new deepwater legskate, *Sinobatis kotlyari* n. sp. (Rajiformes, Anacanthobatidae) from the southeastern Indian Ocean on Broken Ridge. *Zootaxa*, 4189 (2), 327–347.
- Stehmann, M.F.W., Séret, B., Costa, M.E. & Baro, J. (2008) *Neoraja iberica* n. sp., a new species of pygmy skate (Elasmobranchii, Rajidae) from the southern upper slope of the Iberian Peninsula (Eastern North Atlantic). *Cybium*, 32 (1), 51–71.
- Stevenson, D.E., Orr, J.W., Hoff, G.R. & McEachran, J.D. (2004) *Bathyraja mariposa*: A New Species of Skate (Rajidae: Arhynchobatinae) from the Aleutian Islands. *Copeia*, 2004 (2), 305–314.
- Vargas-Caro, C., Bustamante, C., Bennett, M.B. & Ovenden, J.R. (2014) The complete validated mitochondrial genome of the yellownose skate *Zearaja chilensis* (Guichenot 1848) (Rajiformes, Rajidae). *Mitochondrial DNA*.
- Wallace, J.H. (1967) The Batoid Fishes of the East Coast of Southern Africa. Part III: Skates and Electric Rays. *Investigational Report of the Oceanographic Research Institute*, 17, 1–62.
- Weigmann, S. (2016) Annotated checklist of the living sharks, batoids and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity. *Journal of Fish Biology*, 88 (3), 837–1037.
- Weigmann, S. (in press) Reply to Borsa: “Molecular genetics as a valid contribution to the taxonomy of species complexes in stingrays: comment on Weigmann’s global checklist of chondrichthyan fishes”. *Journal of Fish Biology*. 10.1111/jfb.13234.
- Weigmann, S. & Stehmann, M.F.W. (2016) *Sinobatis brevicauda* n. sp., a new deep-water legskate (Rajiformes, Anacanthobatidae) and first generic record from the western Indian Ocean. *Zootaxa*, 4137 (4), 478–500.
- Weigmann, S., Stehmann, M.F.W. & Thiel, R. (2014a) Complementary redescription of *Anacanthobatis ori* (Wallace, 1967) and its assignment to *Indobatis* n. g. (Elasmobranchii, Anacanthobatidae), with comments on other legskates. *Zootaxa*, 3779 (2), 101–132.
- Weigmann, S., Stehmann, M.F.W. & Thiel, R. (2014b) *Rajella paucispinosa* n. sp., a new deep-water skate (Elasmobranchii, Rajidae) from the western Indian Ocean off South Mozambique, and a revised generic diagnosis. *Zootaxa*, 3847 (3), 359–387.
- Weigmann, S., Stehmann, M.F.W. & Thiel, R. (2015) *Okamejei ornata* n. sp., a new deep-water skate (Elasmobranchii, Rajidae) from the northwestern Indian Ocean off Socotra Islands. *Deep-Sea Research II*, 115, 18–29.
- Weigmann, S., Séret, B., Last, P.R. & McEachran, J.D. (in press) Pygmy Skates, Family Gurgesiellidae. In: Last, P.R., White, W.T., Carvalho, M.R. de, Séret, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.) *Rays of the World*. CSIRO Publishing, Melbourne, pp. 473–493.
- White, W.T. & Last, P.R. (2013) Notes on shark and ray types at the South China Sea Fisheries Research Institute (SCSFRI) in Guangzhou, China. *Zootaxa*, 3752 (1), 228–248.
- Whitley, G.P. (1940) *The fishes of Australia. Part I. The sharks, rays, devil-fish, and other primitive fishes of Australia and New Zealand*. Royal Zoological Society of New South Wales, Sydney, 280 pp.