## **SCAMIT Provisional Species Voucher Sheet**

**PROVISIONAL NAME**: *Lumbrineris* sp. E **TAXON**: Annelida, Eunicida, Lumbrineridae

SCAMIT CODE: noneDATE EXAMINED: 12 March 2001VOUCHER BY: Larry Lovell & Kelvin Barwick

**SYNONYMY**: *Lumbrineris* nr. *limicola* (used during SCAMIT meeting presentation 12 March 2001)

## **DIAGNOSTIC CHARACTERS:**

- 1. Composite hooded hooks present, begin setiger 1.
- 2. Yellow acicula.
- 3. Median parapodia with pre- and post-setal lobes, pre-setal lobes shorter than setae, postsetal lobes equal to length of setae. (See Fig. 1)
- 4. Posterior parapodia with pre-setal lobes absent, post-setal lobes rounded at tip, shorter than length of setae. (See Fig. 2)

## **RELATED SPECIES AND CHARACTER DIFFERENCES:**

- 1. *Lumbrineris cruzensis*: pre- and post-setal lobes prolonged in median and posterior segments.
- 2. *Lumbrineris latreilli*: pre-setal lobes absent and post-setal lobes short, blunt with rounded lobe in median segments.
- 3. *Lumbrineris limicola*: post-setal lobes in posterior segments prolonged extending beyond the length of the setae.

**DEPTH RANGE**: 44-80 m (sediment: 10 - 36% fine sand)

**DISTRIBUTION:** Northern Channel Islands.




Figure 1. Anterior segments.




Figure 2. Posterior segments.