

Biogeographical affinities of fish associated to the shrimp trawl fishery in the Gulf of Tehuantepec, Mexico

Marco A. Martínez-Muñoz¹, Domènec Lloris², Adolfo Gracia¹, Ricardo Ramírez-Murillo³, Saul Sarmiento-Nafáte⁴, Sebastián Ramos-Cruz⁴ & Felipe Fernández⁵

1. Unidad Académica de Ecología Marina, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, PO Box 70-305, México, D.F., 04510; marcoam@unam.mx, gracia@unam.mx
2. Institut de Ciències del Mar (CMIMA-CSIC), Pasage Marítim de la Barceloneta 37-49 Barcelona, Spain; ictios@gmail.com
3. Instituto de Educación Media Superior del Distrito Federal (IEMS-DF). Plantel Tlalpan I. Av. División del Norte # 906, Col. Narvarte Poniente, 03020, Del. Benito Juárez México, D.F.; hippoglossina@hotmail.com
4. Instituto Nacional de Pesca, Centro Regional de Investigación Pesquera (CRIP) Salina Cruz, Prol. Playa Abierta S/N, Col. Miramar, 70680. Salina Cruz, Oaxaca, México; nafatess@yahoo.com.mx, ramoscruz@yahoo.com
5. Departament d'Ecologia, Facultat de Biologia, Universitat de Barcelona, Diagonal 645, 08028 Barcelona, Spain; ffernandez@ub.edu

Received 05-V-2015. Corrected 20-X-2015. Accepted 20-XI-2015.

Abstract: Fish by-catch of shrimp fishery from the Gulf of Tehuantepec is composed of several species that are mainly discarded. In this study, fish by-catch species composition, distribution and biogeographical affinities were analyzed. For this, a total of 15 cruises were carried out on the continental shelf, at depths from 15 to 64 m, during 2003, 2004, 2005 and 2013. Results showed that fish by-catch was represented by 58 families, 129 genera and 242 species. The families Haemulidae, Sciaenidae, Paralichthyidae, Gerreidae and Carangidae accounted for > 70 % of the catch. *Haemulopsis axillaris*, *Syacium ovale*, *Selene peruviana*, *Dipterus peruvianus*, *Larimus acclivis* and *Stellifer erycimba* were the most frequent species at < 40 m depth (inner shelf), and *Prionotus stephanophrys*, *Scorpaena russula*, *Porichthys analis* and *Synodus scituliceps* were dominant at 40-60 m depth (outer shelf). Analysis of biogeographical affinities showed that 36.1 % of species had a wide distribution, from San Diego Province to the Panamic Province, while 13.2 % had a restricted distribution in the Mexican and Panamic Provinces. The ichthyofaunal composition was markedly influenced by the local environment and seasonal conditions. Rev. Biol. Trop. 64 (2): 683-700. Epub 2016 June 01.

Key words: biogeography, ichthyofauna, shrimp, fisheries, by-catch, Gulf of Tehuantepec.

The Gulf of Tehuantepec is influenced by an array of environmental factors like the current pattern, seasonal wind regimes, seasonal upwellings, rivers runoff and extensive coastal lagoon systems, that makes this region a very productive one in terms of fisheries (Tapia-García, 1998).

A valuable trawl shrimp fishery is conducted in this area mainly based on four species: *Litopenaeus vannamei* (White shrimp), *Farfantepenaeus californiensis* (Yellowleg shrimp), *Litopenaeus stylirostris* (Blue shrimp) and *Farfantepenaeus brevirostris* (Crystal shrimp)

(Cervantes-Hernández, Gallardo-Berumen, Ramos-Cruz, Gómez-Ponce & Gracia, 2008). Gulf of Tehuantepec shrimp catch contributes nearly 5 % (1880 m) to the total annual catch of the Mexican Pacific (37600 m). Brown and white shrimp are the most important species as they represent 90 % of the total shrimp catch in the area.

Shrimp is the target species in trawling operations, but is associated with a highly diverse fish fauna, which is usually discarded. Both, the large catches discarded and the high ichthyofaunal diversity have caused concern



and prompted attempts to improve its use and management (Allsop, 1985; Andrew & Pepperell, 1992; Hendrickson & Griffin, 1993; Griffiths, Larson, & Courtney, 2004; Sarmiento-Náfate, Gil-López, & Arroyo, 2007; Stobutzki, Miller, Jones, & Salini, 2001).

Studies on the Gulf of Tehuantepec demersal community caught in trawl nets include that of Acal & Arias (1990), who pointed out the high diversity in fish communities in the region. Bianchi (1991) studied the demersal faunal assemblages (fish, crustaceans and cephalopods) of the shelf and upper slope in a wide area of the Gulf of Tehuantepec. Tapia-García, García-Abad, González-Medina, Macuitl-Montes, and Cerdaneras-Ladrón de Guevara (1994) recorded composition and abundance of fish reporting 178 species, with the greatest diversity located in front of the lagoon-estuarine systems of Oaxaca and Chiapas coasts. The dominant species were: *Syacium latifrons*, *S. ovale*, *Eucinostomus gracilis*, *Bothus constellatus*, *Orthopristis chalceus* and *Pomadasys nitidus*. Siqueiros-Beltrones and De la Cruz-Agüero (2004) stood out the importance of taxonomic lists and the analysis of biogeographical distribution as essential tools for fishery management. According to this, several studies have recently assumed increased interest in the Southeastern Mexican Pacific Ocean (Aguilar-Palomino, Mariscal, González, & Rodríguez, 1996; Madrid-Vera, Ruiz, & Rosado, 1998; Moncayo-Estrada, Castro-Aguirre, & De la Cruz-Agüero, 2006). In spite of the ecological and fisheries importance of the Gulf of Tehuantepec, there are only a few by-catch studies on the demersal species from shrimp trawl fishery in this region (Tapia-García, 1998; Tapia-García & García-Abad, 1998).

Up to now, studies are scarce, spatially and temporally scattered with little comprehensive integration and mainly focused on the fishery perspective. This study presents an updated systematic list of by-catch fishes from shrimp fishery, its abundance and analyzes of their biogeographical relationships.

MATERIALS AND METHODS

By-catch sampling was carried out during the shrimp closed seasons (April-August) in the Southern Pacific region of Mexico. A total of 15 exploratory cruises were developed during 2003 (five), 2004 (two), 2005 (four) and then resumed in 2013 (four) in a wide section of the continental shelf of the Gulf of Tehuantepec between Punta Chipehua (Oaxaca) and Puerto Madero (Chiapas) ($14^{\circ}43' - 16^{\circ}01'$ N - $92^{\circ}53' - 95^{\circ}22'$ W). A sampling design of 62 stations was established for all cruises based on previous studies done in the area (Reyna Cabrera & Ramos-Cruz, 1998). Total number of locations sometimes varied due to weather conditions; however, this number was never above 25 % (Fig. 1).

Samples were obtained day and night with twin shrimp nets (24.4 mm horizontal mouth opening, 40.0 mm stretched mesh body, 38.1 mm stretched mesh cod-end) towed at each side of the vessel. The net had a turtle excluder device (TED) fitted in front of the codend. Each trawl had one-hour duration at an average speed of 1.5-2.5 knots; 344 hauls were done in a 15-64 m depth range; 206 hauls (60 %) were made during the day and 138 (40 %) at night; 234 hauls (68 %) were made below 40 m depth and 110 trawls (32 %) were carried above 40 m depth.

The catch from each trawl was cast on deck, and total weight per trawl was estimated filling crates or boxes of 40 kg. A sample of 20 kg was taken; previous analysis showed that this was enough to reflect adequately the species composition. Each sample was kept in the freezer at -30° C.

The biological material was identified and processed in the laboratories of the Regional Centre of Fisheries Research, Salina Cruz (CRIP-SC). The standard length of each fish was measured with a standard ichthyometer of 50 cm length and a precision of 0.05 cm. Large individual fish were weighted with a scale (25 g of precision) and small fishes were weighted with an OHAUS digital scale of 2.6 kg of capacity and a precision of 0.05 g. In the






Fig. 1. Study area and sampling stations in the Gulf of Tehuantepec, Mexico.

Fig. 1. Área de estudio y estaciones de muestreo en el Golfo de Tehuantepec, México.

laboratory, fish were separated, washed, re-labelled and stored in 70 % isopropyl alcohol.

Fish identification followed the keys of Castro-Aguirre (1978) Eschmeyer (1998), Bussing and López (1993), Fischer, Krupp, Schneider, Sommer, Carpenter and Niem (1995), Robertson and Allen (2002), and Amezcu-Linares (1996). Specialized literature for some specific groups was also consulted: rays, Castro-Aguirre and Espinosa-Pérez (1996); sharks, Espinosa-Pérez, Castro-Aguirre and Huidobro-Campos (2004); Sciaenidae, McPhail (1958); Pleuronectiformes, Norman (1934), and Ginsburg (1958); *Diplectrum*, Roseblatt and Johnson (1974); and *Porichthys*, Walter and Roseblatt (1988).

Systematic ordination followed the criteria of Nelson (2006). Genera and their species are presented in alphabetical order. Common names were taken from Robertson and Allen (2002); Nelson et al. (2004); and Love, Mecklenburg, Mecklenburg and Thorsteinson (2005).

The relative abundance of each species was estimated from the number of individuals within the total of individuals sampled. The species were grouped in four categories: (A) Abundant, relative abundance > 1 %; (F)

Frequent, 0.1-0.99 %; (C) Common, 0.01-0.099 %; and (R) Rare, < 0.01 %. Differences in biomass values between day and night as well as the depth intervals were calculated through variance analysis (ANOVA) and t-test (Zar, 1999).

Analysis of zoogeographic affinities followed the basic scheme of Briggs (1974, 1995) and Walker (1960), with modifications proposed by Boschi (2000), Galván-Magaña, Gutiérrez-Sánchez, Abitia-Cárdenas and Rodríguez-Romero (2000), Hastings (2000), Robertson & Allen (2002), Robertson, Grove and McCoster (2004), and Horn, Allen and Lea (2006), considering the following provinces: A) Oregonian: from Puget Sound to Punta Concepción in California. The Southern section is considered a transition zone between this province and that of San Diego. B) San Diego: from Punta Concepción to Bahía Magdalena in the South of Baja California, with temperate-warm waters. C) Cortés: Southern Bahía Magdalena and the Gulf of California. D) Mexican: from Mazatlán, Sinaloa to Tehuantepec (Mexico). E) Panamic: from the South of Salvador to Cabo Blanco (Peru). F) Tropical Eastern Pacific: includes the West coast of the American



continent between 25° N in the Southern part of Bahía Magdalena and 5 °S at Cabo Blanco, Northern Peru. Differences in species biogeographic affinities between the years 2003 and 2013 were calculated by analysis of variance (ANOVA) (Zar, 1999).

The analysis also considered species with wide biogeographic distribution, such as the Circumtropical, Trans-Pacific and amphi-American species of the Central American Isthmus, the Tropical Eastern Pacific and the Western Atlantic.

Classification of the abundant, frequent, common and rare species used the Olmstead-Tukey method modified by Sokal and Rohlf

(1995). The data matrices were expressed as frequency of occurrence within each sampling period. Then the frequency of occurrence per cruise was calculated (mean \pm standard deviation). Richness was considered as the number of fish species in the study area.

RESULTS

The total number of samplings from all cruises was 344, with a total weight of 2818 kg. The 66 908 individual fishes represented two Classes, 20 Orders, 58 families, 129 genera and 242 species (Table 1). Previous studies in the area showed a lower number of demersal

TABLE 1
Fishes of the Gulf of Tehuantepec, Mexico

CUADRO 1
Peces del Golfo de Tehuantepec, México

Species	A	B	C	D
Phylum Chordata				
Class Chondrichthyes				
Subclass Elasmobranchii				
Subdivision Selachii				
Order Carcharhiniformes				
Family Triakidae				
<i>Mustelus lunulatus</i> Jordan & Gilbert, 1883	Sicklefin smooth-hound	94-108	R	PS-PP
Family Carcharhinidae				
<i>Rhizoprionodon longurio</i> (Jordan & Gilbert 1882)	Pacific sharpnose shark	230	R	PS-PP
Family Sphyrnidae				
<i>Sphyraena lewini</i> (Griffith & Smith, 1834)	Scalloped hammerhead	176-350	R	CT
<i>Sphyraena tiburo</i> (Linnaeus, 1758)	Bonnethead	260	R	PS-PP
Subdivision Batoidea				
Order Torpediniformes				
Family Narcinidae				
<i>Narcine entemedor</i> Jordan & Starks, 1895	Giant electric ray	40-399	C	PC-PP
<i>Narcine vermiculatus</i> Breder, 1928	Vermiculate electric ray	44-444	F	POT
Order Rajiformes				
Family Rhinobatidae				
<i>Rhinobatos glaucoptigma</i> Jordan & Gilbert, 1883	Speckled guitarfish	60-335	F	POT
<i>Rhinobatos leucorhynchus</i> Günther, 1867	Whitesnout guitarfish	68-571	C	PC-PP
<i>Zapteryx exasperata</i> (Jordan & Gilbert, 1880)	Banded guitarfish	95-230	C	PS-PP
<i>Zapteryx xyster</i> Jordan & Evermann, 1896	Witch guitarfish	406	R	PC-PP
Order Myliobatiformes				
Family Urolophidae				
<i>Urolophus halleri</i> Cooper, 1863	Haller's round ray	170	R	POT



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
<i>Urolophus maculatus</i> (Garman, 1913)	Spotted round ray	110-125	R	PS-PC
<i>Urotrygon aspidura</i> (Jordan & Gilbert, 1882)	Spiny-tail round ray	41-395	F	PM-PP
<i>Urotrygon chilensis</i> (Günther, 1872)	Chilean round ray	35-364	A	PC-PP
<i>Urotrygon munda</i> Gill, 1863	Munda round ray	42-230	F	PM-PP
<i>Urotrygon nana</i> Miyake & McEachran, 1988	Dwarf round ray	30-235	F	PM-PP
<i>Urotrygon reticulata</i> Miyake & McEachran, 1988	Reticulate round ray	90-151	R	PM-PP
<i>Urotrygon rogersi</i> (Jordan & Starks, 1895)	Roger's round ray	64-411	C	PC-PP
Family Dasyatidae				
<i>Dasyatis brevis</i> (Garman, 1880)	Whiptail stingray	146	R	PO-PP
<i>Dasyatis longa</i> (Garman, 1880)	Longtail stingray	118-227	R	PO-PP
Family Gymnuridae				
<i>Gymnura marmorata</i> (Cooper, 1864)	California butterfly ray	200-326	C	PS-PP
Family Myliobatidae				
<i>Aetobatus narinari</i> (Euphrasen, 1790)	Spotted eagle ray	165-195	C	CT
<i>Rhinoptera steindachneri</i> Evermann & Jenkins, 1891	Pacific cownose ray	175-406	R	PS-PP
Class Actinopterygii				
Subclass Neopterygii				
Division Teleostei				
Order Albuliformes				
Family Albulidae				
<i>Albula nemoptera</i> (Fowler, 1911)	Threadfin bonefish	134-276	C	POT
<i>Albula vulpes</i> (Linnaeus, 1758)	Bonefish	175-231	C	POT
Order Anguilliformes				
Suborder Muraenoidei				
Family Muraenidae				
<i>Gymnothorax equatorialis</i> (Hildenbrand, 1946)	Spotted-tail moray	213-681	C	PC-PP
<i>Gymnothorax panamensis</i> (Steindachner, 1876)	Panamic moray	531-569	C	PS-PP
Suborder Congroidei				
Family Ophichthidae				
<i>Myrophis vafer</i> Jordan & Gilbert, 1883	Pacific worm eel	247	R	PS-PP
<i>Ophichthus zophochir</i> Jordan & Gilbert, 1882	Yellow snake eel	205-580	C	PS-PP
<i>Pseudomyrophis micropinna</i> Wade, 1946	Smallfin worm eel	169	R	PM-PP
Family Congridae				
<i>Ariosoma gilberti</i> (Ogilby, 1898)	Gilbert's garden eel	109-165	C	PC
<i>Paraconger californiensis</i> Kanazawa, 1961	California conger	483	R	PS-PP
Order Clupeiformes				
Suborder Clupeoidei				
Family Engraulidae				
<i>Anchoa argentivittata</i> (Regan, 1904)	Regan's anchovy	50-65	R	PC-PP
<i>Anchoa helleri</i> (Hubbs, 1921)	Heller's anchovy	38-109	C	PC-PM
<i>Anchoa ischana</i> (Jordan & Gilbert, 1882)	Slender anchovy	40-119	F	PC-PP
<i>Anchoa lucida</i> (Jordan & Gilbert, 1882)	Bright anchovy	46-193	F	PC-PP
<i>Anchoa mundeola</i> (Gilbert & Pierson, 1898)	False Panama anchovy	50-72	C	PC-PP
<i>Anchoa nasus</i> (Kner & Steindachner, 1867)	Longnose anchovy	60-119	C	PC-PP
<i>Anchoa starksii</i> (Gilbert & Pierson, 1898)	Black-tail anchovy	84	R	PP
<i>Anchoa walkeri</i> Baldwin & Chang, 1970	Walker's anchovy	40-145	C	PC-PP
<i>Anchovia macrolepidota</i> (Kner, 1863)	Bigscale anchovy	47-177	F	PC-PM



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
<i>Cetengraulis mysticetus</i> (Günther, 1867)	Pacific anchoveta	36-134	C	PS-PP
Family Clupeidae				
<i>Harengula thrissina</i> Jordan & Gilbert, 1882	Pacific flatiron herring	117-180	C	PC-PP
<i>Lile gracilis</i> Castro-Aguirre & Vivero, 1990	Graceful piquitinga	80-190	C	PC-PP
<i>Lile stolifera</i> (Jordan & Gilbert, 1882)	Pacific piquitinga	118	R	PC-PP
<i>Opisthonema libertate</i> (Günther, 1867)	Pacific thread herring	73-195	F	PS-PP
<i>Opisthonema medirastre</i> Berry & Barrett, 1963	Middling thread herring	118-166	C	PC-PP
<i>Sardinops sagax</i> (Jenyns, 1842)	South American pilchard	207	R	PA-PC
Family Pristigasteridae				
<i>Opisthoterpes dovii</i> (Günther, 1868)	Dove's longfin herring	59-180	F	PS-PP
<i>Pliosteostoma lutipinnis</i> (Jordan & Gilbert, 1882)	Yellowfin herring	45-482	F	PM-PP
Order Siluriformes				
Family Ariidae				
<i>Ariopsis guatemalensis</i> (Günther, 1864)	Blue sea catfish	66-195	F	PC-PP
<i>Ariopsis seemanni</i> (Günther, 1864)	Tete sea catfish	41-236	F	PC-PP
<i>Bagre panamensis</i> (Gill, 1863)	Chilhuil sea catfish	64-410	F	PC-PP
<i>Bagre pinnimaculatus</i> (Steindachner, 1876)	Red sea catfish	152-325	R	PC-PP
<i>Cathorops dasycephalus</i> (Günther, 1864)	Big-bellied sea catfish	38-172	C	PM
<i>Cathorops fuerthii</i> (Steindachner, 1876)	Congo sea catfish	40-205	F	PC-PP
<i>Cathorops steindachneri</i> (Gilbert & Starks, 1904)	Steindachner's sea catfish	119-191	C	PM-PP
<i>Cathorops taylori</i> (Hildebrand, 1925)	Sea catfish	176-213	R	PP
<i>Galeichthys peruvianus</i> Lütken, 1874	Peruvian sea catfish	140-241	C	PM-PP
<i>Notarius kessleri</i> (Steindachner, 1876)	Sculptured sea catfish	70-255	C	PM-PP
<i>Notarius planiceps</i> (Steindachner, 1876)	Flathead sea catfish	68-247	C	PM-PP
<i>Notarius troschelii</i> (Gill, 1863)	Chili sea catfish	28-275	F	PM-PP
<i>Occidentarius platypogon</i> (Günther, 1864)	Cominate sea catfish	57-290	F	PC-PP
<i>Sciaes dowii</i> (Gill, 1863)	Brown sea catfish	170-190	R	PP
Order Aulopiformes				
Suborder Synodontoidei				
Family Synodontidae				
<i>Synodus evermanni</i> Jordan & Bollman, 1890	Inotted lizardfish	84-312	F	PS-PP
<i>Synodus scituliceps</i> Jordan & Gilbert, 1882	Shorthead lizardfish	91-420	F	PS-PP
Order Gadiformes				
Family Merluccidae				
<i>Merluccius angustimanus</i> Garman, 1899	Panama hake	68-155	C	PS-PP
<i>Merluccius productus</i> (Ayres, 1855)	North Pacific hake	46-180	C	PS-PP
Order Ophidiiformes				
Suborder Ophidioidei				
Family Ophidiidae				
<i>Brotula clarkae</i> Hubbs, 1944	Pacific bearded brotula	154-315	C	PC-PP
<i>Chilara taylori</i> (Girard, 1858)	Spotted cusk-eel	178	R	PO-PP
<i>Lepophidium pardale</i> (Gilbert, 1890)	Leopard cusk eel	120-169	R	POT
<i>Lepophidium prorates</i> (Jordan & Bollman, 1890)	Prowspine cusk eel	60-254	C	POT
<i>Ophidion galeoides</i> (Gilbert, 1890)	Striped cusk eel	175	R	PM-PP
<i>Otophidium indefatigabile</i> Jordan & Bollman, 1890	Bighead cusk eel	169	R	PS-PP
Order Batrachoidiformes				
Family Batrachoididae				
<i>Batrachoides bouleengeri</i> Gilbert & Starks, 1904	Boulenger's toadfish	87-210	C	PM-PP



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
<i>Batrachoides waltersi</i> Collette & Russo, 1981	Walter's toadfish	40-292	F	PM-PP
<i>Porichthys analis</i> Hubbs & Schultz, 1939	Darkedge midshipman	10-174	A	PS-PP
<i>Porichthys marginatus</i> (Richardson, 1844)	Daisy midshipman	52-160	F	POT
Suborder Antennariidae				
Family Antennariidae				
<i>Fowlerichthys avalonis</i> (Jordan & Starks, 1907)	Roughbar frogfish	50-156	C	PS-PP
Order Mugiliformes				
Family Mugilidae				
<i>Mugil cephalus</i> Linnaeus, 1758	Flathead grey mullet	167-188	R	PS-PP
<i>Mugil curema</i> Valenciennens, 1836	White mullet	160-210	C	PC-PP
<i>Mugil setosus</i> Gilbert, 1892	Liseta mullet	155	R	PC-PP
Suborder Belonoidei				
Family Exocoetidae				
<i>Cheilopogon papilio</i> (Clark, 1936)	Butterfly flyingfish	117-127	R	PM-PP
Order Beloniformes				
Family Hemiramphidae				
<i>Cypselurus callopterus</i> (Günther, 1866)	Ornamented flyingfish	185	R	PS-PP
<i>Hemiramphus saltator</i> Gilbert & Starks, 1904	Longfin halfbeak	240-250	R	PS-PP
Suborder Cyprinodontoidei				
Order Cyprinodontiformes				
Family Poeciliidae				
<i>Poecilia butleri</i> Jordan, 1889	Pacific molly	55	R	PM
<i>Poeciliopsis fasciata</i> (Meek, 1904)	San Jeronimo livebearer	52	R	PS-PP
Order Gasterosteiformes				
Suborder Syngnathoidei				
Family Syngnathidae				
<i>Hippocampus ingens</i> Girard, 1858	Pacific seahorse	117-197	R	PO-PP
Family Fistularidae				
<i>Fistularia commersonii</i> Rüppell, 1838	Bluespotted cornetfish	170	R	PC-PP
<i>Fistularia corneta</i> Gilbert & Starks, 1904	Pacific cornetfish	164-381	C	PO-PP
Order Scorpaeniformes				
Suborder Scorpaenoidei				
Family Scorpaenidae				
<i>Pontinus sierra</i> (Gilbert, 1890)	Speckled scorpionfish	67	R	PC-PM
<i>Scorpaena histrio</i> Jenyns, 1840	Player scorpionfish	35-88	F	PS-PP
<i>Scorpaena mystes</i> Jordan & Starks, 1895	Pacific spotted scorpionfish	92-145	R	PS-PP
<i>Scorpaena russula</i> Jordan & Bollman, 1890	Reddish scorpionfish	27-130	A	PS-PP
Suborder Platycephaloidei				
Family Triglidae				
<i>Bellator gymnostethus</i> (Gilbert, 1892)	Naked-belly searobin	35-239	R	PC-PP
<i>Bellator loxias</i> (Jordan, 1897)	Barred searobin	65-277	C	POT
<i>Bellator xenisma</i> (Jordan & Bollman, 1889)	Splitnose searobin	32-282	F	PS-PP
<i>Prionotus albirostris</i> Jordan & Bollman, 1890	Whitesnout searobin	39-110	C	PC-PP
<i>Prionotus birostatus</i> Richardson, 1844	Two-beaked searobin	55-198	C	PC-PP
<i>Prionotus horrens</i> Richardson, 1844	Bristly searobin	30-125	F	POT
<i>Prionotus ruscarius</i> Gilbert & Starks, 1904	Common searobin	53-245	F	PS-PP
<i>Prionotus stephanophrys</i> Lockington, 1881	Lumptail searobin	30-200	A	PM-PP



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
Order Perciformes				
Suborder Percoidei				
Family Centropomidae				
<i>Centropomus armatus</i> Gill, 1863	Armed snook	170	R	PM-PP
<i>Centropomus medioides</i> Günther, 1864	Blackfin snook	93-140	C	PS-PP
<i>Centropomus robalito</i> Jordan & Gilbert, 1882	Yellowfin snook	58-190	F	PC-PP
<i>Centropomus unionensis</i> Bocourt, 1868	Union snook	69-162	C	PM-PP
Family Serranidae				
<i>Alphestes immaculatus</i> Breder, 1936	Pacific mutton hamlet	65-110	C	PC-PP
<i>Alphestes multiguttatus</i> (Günther, 1867)	Rivulated mutton hamlet	72-172	C	PC-PP
<i>Cephalopholis panamensis</i> (Steindachner, 1876)	Pacific graysby	180	R	PC-PP
<i>Diplectrum eumelum</i> Rosemblatt & Johnson, 1974	Orange-spotted sand perch	50-162	F	POT
<i>Diplectrum euryplectrum</i> Jordan & Bollman, 1890	Bighead sand perch	59-155	F	PC-PP
<i>Diplectrum labarum</i> Rosemblatt & Johnson, 1974	Highfin sand perch	60-158	F	PS-PP
<i>Diplectrum macropoma</i> (Günther, 1864)	Mexican sand perch	65-140	C	POT
<i>Diplectrum pacificum</i> Meek & Hildebrand, 1925	Inshore sand perch	43-195	F	PS-PP
<i>Epinephelus analogus</i> Gill, 1863	Spotted grouper	90-308	C	PS-PP
<i>Epinephelus labriformis</i> (Jenyns, 1840)	Starry grouper	104-104	R	PC-PP
<i>Hyporthodus acanthistius</i> (Gilbert, 1892)	Rooster hind	74-510	C	PS-PP
<i>Hyporthodus exsul</i> (Fowler, 1944)	Tenspine grouper	200-570	C	PC-PP
<i>Hyporthodus niphobles</i> (Gilbert & Starks, 1897)	Star-studded grouper	61-138	C	PS-PP
<i>Rypticus nigripinnis</i> Gill, 1861	Blackfin soapfish	98-160	C	PS-PP
Family Priacanthidae				
<i>Pristigenys serrula</i> (Gilbert, 1891)	Popeye catalufa	45-128	F	PO-PP
Family Carangidae				
<i>Carangoides otrynter</i> (Jordan & Gilbert, 1883)	Threadfin jack	36-468	C	PS-PP
<i>Caranx caballus</i> Günther, 1868	Green jack	132-190	R	PS-PP
<i>Caranx vinctus</i> Jordan & Gilbert, 1882	Cocinero	32-175	F	PS-PP
<i>Caranx caninus</i> Günther, 1867	Pacific crevalle jack	91-118	R	PS-PP
<i>Chloroscombrus orqueta</i> Jordan & Gilbert, 1883	Pacific bumper	48-325	A	PS-PP
<i>Decapterus macarellus</i> (Cuvier, 1833)	Mackerel scad	165-184	R	CT
<i>Decapterus macrosoma</i> Bleeker, 1851	Shortfin scad	92-192	C	CT
<i>Decapterus muroidesi</i> (Temminck & Schlegel, 1844)	Amberstripe scad	115-325	R	CT
<i>Hemicaranx leucurus</i> (Günther, 1864)	Yellowfin jack	56-190	C	POT
<i>Hemicaranx zelotes</i> Gilbert, 1898	Blackfin jack	39-195	F	PC-PP
<i>Oligoplites refulgens</i> Gilbert & Starks, 1904	Shortjaw leatherjack	113	R	PM-PP
<i>Oligoplites saurus</i> (Bloch & Schneider, 1801)	Leatherjack	144	R	PC-PP
<i>Selar crumenophthalmus</i> (Bloch, 1793)	Bigeye scad	54-195	C	PC-PP
<i>Selene brevoortii</i> (Gill, 1863)	Mexican lookdown	40-205	F	PC-PP
<i>Selene orstedii</i> Lutken, 1880	Mexican moonfish	42-78	C	PC-PP
<i>Selene peruviana</i> (Guichenot, 1866)	Pacific moonfish	31-238	A	PS-PP
Family Lutjanidae				
<i>Lutjanus argentiventris</i> (Peters, 1869)	Yellow snapper	70	R	PS-PP
<i>Lutjanus guttatus</i> (Steindachner, 1869)	Spotted rose snapper	42-405	F	PC-PP
<i>Lutjanus peru</i> (Nichols & Murphy, 1922)	Pacific red snapper	39-295	F	PC-PP
Family Gerreidae				
<i>Diapterus aureolus</i> Jordan & Gilbert 1882	Golden mojarra	52-157	F	PC-PP



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
<i>Diapterus peruvianus</i> (Cuvier, 1830)	Peruvian mojarra	10-222	A	PC-PP
<i>Eucinostomus argenteus</i> Baird & Girard, 1855	Silver mojarra	78-109	R	PS-PP
<i>Eucinostomus currani</i> Zahuranec, 1980	Pacific flagfin mojarra	51-194	A	PC-PP
<i>Eucinostomus dowii</i> (Gill, 1863)	Dow's mojarra	80-145	R	PC-PP
<i>Eucinostomus entomelas</i> Zahuranec, 1980	Dark-spot mojarra	72-145	C	PC-PP
<i>Eucinostomus gracilis</i> (Gill, 1862)	Graceful mojarra	48-170	A	PC-PP
<i>Gerres cinereus</i> (Walbaum, 1792)	Yellow fin mojarra	72-92	R	PC-PP
Family Haemulidae				
<i>Anisotremus interruptus</i> (Gill, 1862)	Burrito grunt	132	R	PC-PP
<i>Conodon serrifer</i> Jordan & Gilbert, 1882	Armed grunt	64-165	F	PS-PP
<i>Haemulon scudderii</i> Gill, 1862	Grey grunt	42-200	A	PC-PP
<i>Haemulon steindachneri</i> (Jordan & Gilbert, 1882)	Chere-chere grunt	82-205	C	POT
<i>Haemulopsis axillaris</i> (Steindachner, 1869)	Yellowstripe grunt	31-297	A	PC-PP
<i>Haemulopsis elongatus</i> (Steindachner, 1879)	Elongate grunt	132	R	PM-PP
<i>Haemulopsis leuciscus</i> (Günther, 1864)	White grunt	31-252	F	PS-PP
<i>Haemulopsis nitidus</i> (Steindachner, 1869)	Shining grunt	47-285	F	PC-PP
<i>Microlepidotus brevipinnis</i> (Steindachner, 1869)	Humpback grunt	115	R	PC-PP
<i>Orthopristis chalceus</i> (Günther, 1864)	Brassy grunt	72-215	F	PM
<i>Orthopristis reddingi</i> Jordan & Richardson, 1895	Bronze-striped grunt	98-148	C	PC-PP
<i>Pomadasys bayanus</i> Jordan & Evermann, 1898	Purplemouth grunt	55-176	C	PC-PP
<i>Pomadasys branickii</i> (Steindachner, 1879)	Sand grunt	111-182	C	PC-PP
<i>Pomadasys panamensis</i> (Steindachner, 1876)	Panama grunt	32-314	F	PC-PP
<i>Xenichthys xanti</i> Gill, 1863	Longfin salema	64-176	F	PS-PP
<i>Xenistius californiensis</i> (Steindachner, 1876)	Californian salema	102-130	R	PS-PP
Family Polynemidae				
<i>Polydactylus approximans</i> (Lay & Bennet, 1839)	Blue bobo	56-207	A	PS-PP
<i>Polydactylus opercularis</i> (Gill, 1863)	Yellow bobo	70-260	F	PC-PP
Family Sciaenidae				
<i>Aplodinotus grunniens</i> Rafinesque, 1819	Freshwater drum	95-199	C	PM
<i>Bairdiella armata</i> Gill, 1863	Armed croaker	68-185	C	PC-PP
<i>Bairdiella ensifera</i> (Jordan & Gilbert, 1862)	Swordspine croaker	130-156	C	PM-PP
<i>Cynoscion phoxocephalus</i> Jordan & Gilbert, 1882	Cachema weakfish	108	R	PC-PP
<i>Cynoscion reticulatus</i> (Günther, 1864)	Striped weakfish	160-205	R	PC-PP
<i>Cynoscion stolzmanni</i> (Steindachner, 1879)	Stolzmann's weakfish	83-184	R	PC-PP
<i>Elattarchus archidium</i> (Jordan & Gilbert 1882)	Bluestreak drum	110-155	C	PC-PP
<i>Isopisthus remifer</i> Jordan & Gilbert, 1882	Silver weakfish	50-215	F	PS-PP
<i>Larimus acclivis</i> Jordan & Bristol, 1898	Steepledrum	34-211	A	PC-PP
<i>Larimus argenteus</i> (Gill, 1863)	Silver drum	37-195	A	PC-PP
<i>Larimus effulgens</i> Gilbert, 1898	Shining drum	68-170	F	PC-PP
<i>Larimus pacificus</i> Jordan & Bollman, 1890	Pacific drum	39-155	F	POT
<i>Menticirrhus elongatus</i> (Günther, 1864)	Pacific kingcroaker	85-280	C	PC-PP
<i>Menticirrhus nasus</i> (Günther, 1868)	Highfin kingcroaker	74-215	F	PC-PP
<i>Menticirrhus panamensis</i> (Steindachner, 1876)	Panama kingcroaker	54-192	C	PC-PP
<i>Menticirrhus undulatus</i> (Girard, 1854)	California kingcroaker	87-185	C	PC-PP
<i>Micropogonias altipinnis</i> (Günther, 1864)	Tallfin croaker	59-280	F	PC-PP
<i>Micropogonias ectenes</i> (Jordan & Gilbert 1882)	Slender croaker	189-190	R	PS-PM
<i>Nebris occidentalis</i> Vaillant, 1897	Pacific smalleye croaker	54-240	F	PM-PP



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
<i>Ophioscion imiceps</i> (Jordan & Gilbert 1882)	Blinkard croaker	50-194	F	PC-PP
<i>Ophioscion strabo</i> Gilbert, 1897	Squint-eyed croaker	69-178	C	PM-PP
<i>Ophioscion typicus</i> Gill, 1863	Point-nosed croaker	70-475	C	PS-PP
<i>Paralonchurus goodei</i> Gilbert, 1898	Goode croaker	100-182	C	PM-PP
<i>Paralonchurus rathbuni</i> (Jordan & Bollman 1890)	Bearded banded croaker	112-201	C	PM-PP
<i>Stellifer chrysoleuca</i> (Günther, 1867)	Shortnose stardrum	44-246	C	PS-PM
<i>Stellifer ericymba</i> (Jordan & Gilbert, 1882)	Hollow stardrum	40-300	A	PS-PM
<i>Stellifer fuerthii</i> (Steindachner, 1876)	White stardrum	54-186	F	PP
<i>Stellifer illecebrosus</i> Gilbert, 1898	Silver stardrum	59-185	F	PS-PP
<i>Umbrina xanti</i> Gill, 1862	Polla drum	45-313	F	PM-PP
Family Mullidae				
<i>Pseudupeneus grandisquamis</i> (Gill, 1863)	Bigscale goatfish	56-172	A	PS-PP
Family Chaetodontidae				
<i>Chaetodon humeralis</i> Günther, 1860	Threebanded butterflyfish	37-112	F	PS-PP
Suborder Labroidei				
Family Labridae				
<i>Halichoeres chierchiai</i> Di Caporiacco, 1948	Wounded wrasse	128-152	R	PC-PP
Suborder Trachinoidei				
Family Ephippidae				
<i>Chaetodipterus zonatus</i> (Girard, 1858)	Pacific sleeper	125-140	R	PC-PP
<i>Parapsettus panamensis</i> (Steindachner, 1876)	Pennant goby	59-90	C	PC-PM
Family Uranoscopidae				
<i>Astroscopus zephyreus</i> Gilbert & Starks 1897	Pacific stargazer	193	R	PS-PP
Suborder Gobioidei				
Family Eleotridae				
<i>Gobiomorus maculatus</i> (Günther, 1859)	Tailspot goby	66-72	R	PC-PP
Family Gobiidae				
<i>Bollmannia ocellata</i> Gilbert, 1892	Pacific spadefish	32-176	F	PC-PP
<i>Bollmannia stigmatura</i> Gilbert, 1892	Panama spadefish	33-286	F	POT
Suborder Scombroidei				
Family Trichiuridae				
<i>Lepidopus fitchi</i> Rosenblatt & Wilson, 1987	Pacific scabbardfish	400	R	PO-PP
<i>Trichiurus lepturus</i> Linnaeus, 1758	Largehead hairtail	436	C	CT
Family Scombridae				
<i>Scomberomorus sierra</i> Jordan & Starks, 1895	Pacific sierra	135-290	C	PS-PP
Family Sphyraenidae				
<i>Sphyraena ensis</i> Jordan & Gilbert, 1882	Mexican barracuda	16-304	F	PC-PP
Suborder Stromateoidei				
Family Stromateidae				
<i>Peprilus medius</i> (Peters, 1869)	Pacific harvestfish	110-222	C	POT
<i>Peprilus ovatus</i> Horn, 1970	Shining butterfish	115-157	R	PS-PP
<i>Peprilus snyderi</i> Gilbert & Starks, 1904	Salema butterfish	49-179	C	PM-PP
Order Pleuronectiformes				
Suborder Pleuronectoidei				
Family Paralichthyidae				
<i>Ancyloplitta dendritica</i> Gilbert, 1890	Three-spot flounder	105-222	R	PM-PP
<i>Citharichthys platophrys</i> Gilbert, 1891	Small sandab	33-124	F	PM-PP



TABLE 1 (Continued) / CUADRO 1 (Continuación)

Species	A	B	C	D
<i>Cyclopsetta panamensis</i> (Steindachner, 1876)	God's flounder	45-306	A	PC-PP
<i>Cyclopsetta querna</i> (Jordan & Bollman, 1890)	Toothed flounder	56-385	F	PC-PP
<i>Etropus crossotus</i> Jordan & Gilbert, 1882	Fringed flounder	89-170	F	PS-PP
<i>Paralichthys woolmani</i> Jordan & Williams, 1897	Speckled flounder	225-290	R	PC-PP
<i>Syacium latifrons</i> (Jordan & Gilbert, 1882)	Beach flounder	40-237	A	POT
<i>Syacium longidorsale</i> Murakami & Amaoka, 1992	Longfin flounder	96	R	PM-PP
<i>Syacium ovale</i> (Günther, 1864)	Oval flounder	88-128	A	POT
Family Bothidae				
<i>Bothus constellatus</i> (Jordan, 1889)	Pacific eyed flounder	35-303	A	PS-PP
<i>Engyophrys sanctilaurentii</i> Jordan & Bollman, 1890	Speckled-tail flounder	88-90	R	PC-PP
<i>Monolepis asaedai</i> Clark, 1936	Asaedeas flounder	39-107	F	PS-PP
Family Cynoglossidae				
<i>Symphurus atramentatus</i> Jordan & Bollman, 1890	Inkspot tonguefish	130-135	R	PC-PP
<i>Symphurus atricaudus</i> (Jordan & Gilbert, 1880)	California tonguefish	126-164	C	PO-PP
<i>Symphurus elongatus</i> (Günther, 1868)	Elongate tonguefish	72-266	F	PM-PP
<i>Symphurus melanurus</i> Clark, 1936	Drab tonguefish	135	R	POT
Family Achiridae				
<i>Achirus mazatlanus</i> (Steindachner, 1869)	Mazatlan sole	34-146	C	PS-PP
<i>Achirus scutum</i> (Günther, 1862)	Network sole	51-205	F	PM
<i>Achirus zebrinus</i> Clark, 1936	Tehuantepec sole	50-182	F	PS-PP
<i>Trinectes fimbriatus</i> (Günther, 1862)	Fringed sole	57-75	R	PS-PP
<i>Trinectes fonsecensis</i> (Günther, 1862)	Spottedfin sole	58-190	C	PS-PP
Order Tetraodontiformes				
Suborder Balistoidei				
Family Balistidae				
<i>Balistes polylepis</i> Steindachner, 1876	Finescale triggerfish	45-350	C	PO-PP
<i>Pseudobalistes naufragium</i> (Jordan & Starks, 1895)	Stone triggerfish	39-65	C	PS-PP
Family Monacanthidae				
<i>Aluterus monoceros</i> (Linnaeus, 1758)	Unicorn leatherjacket filefish	280-441	R	CT
Suborder Tetraodontoidei				
Family Tetraodontidae				
<i>Sphoeroides annulatus</i> (Jenyns, 1842)	Bullseye puffer	30-264	F	PM-PP
<i>Sphoeroides kendalli</i> Meek & Hildebrand, 1928	Slick puffer	65-87	F	PS-PP
<i>Sphoeroides lobatus</i> (Steindachner, 1870)	Longnose puffer	46-210	R	PO-PP
<i>Sphoeroides sechurae</i> Hildebrand, 1946	Peruvian puffer	44-169	F	POT
Family Diodontidae				
<i>Chilomycterus reticulatus</i> (Linnaeus, 1758)	Spotfin burrfish	60-190	F	CT
<i>Diodon holocanthus</i> Linnaeus, 1758	Long-spine porcupinefish	109-275	R	CT
<i>Diodon hystrix</i> Linnaeus, 1758	Spot-fin porcupinefish	70-290	C	CT

A: Common Name in English; B: Size range (standard length, mm); C: Relative abundance; D: Province and biogeographical region (PO, Oregonian; PS, San Diego Province; PC, Cortés Province; PM, Mexican Province; PP, Panamic Province; POT, Tropical Eastern Pacific; CT, Circumtropical species; T, Trans-Pacific species). Species abundance: (A) abundant; (F) frequent; (C) common; (R) rare. Systematic arrangement according to Nelson (2006). Revision of species according to Eschmeyer (1998) and Froese & Pauly (2013).

A: Nombre común en inglés, B: Intervalo de tallas (longitud estándar, mm), C: Abundancia relativa, D: Provincias y regiones biogeográficas (PO: Provincia Oregoniana, PS: Provincia de San Diego, PC: Provincia de Cortés, PM: Provincia Mexicana, PP: Provincia Panámica, POT: Pacífico Oriental Tropical, CT: Especies Circumtropicales, y T: Especies Transpacíficas). Especie abundante (A), especie frecuente (F), especie común (C) y especie rara (R). Clasificación sistemática según Nelson (2006). Revisión de especies según Eschmeyer (1998), Froese y Pauly (2013).



TABLE 2
Comparison of previous studies regarding demersal fish lists in the continental shelf of the Gulf of Tehuantepec

CUADRO 2
Comparación de estudios anteriores con respecto a las listas de peces demersales en
la plataforma continental del Golfo de Tehuantepec

Author	Cruises	Year	Nº of Trawls	Depth Range (m)	Nº of Families	Nº of Genus	Nº of Species
Secretaría de Marina (1978)	1	1977	11	27-60	32	49	61
Acal-Arias (1990)	4	1987	161	15-200	53	115	195
Tapia-García (1998)	5	1989-1990	85	15-80	56	123	173
Present study	15	2003-2005, 2013	344	15-64	58	129	242

fish species with respect to this study (Table 2). The Order Perciformes was the most diverse, with 20 families, 55 genera and 110 species, which is typical of the ichthyofaunal groups in the intertropical regions. The families best represented in terms of number of species were: Sciaenidae (29), Haemulidae (16), Carangidae (16), and Serranidae (14). The main genera were: *Anchoa*, *Eucinostomus*, *Epinephelus* and *Urotrygon*, each with six species, followed by *Prionotus* and *Diplectrum*, each with five species. Sizes ranged from 27 mm in the red-dish scorpionfish (*Scorpaena russula*) to 681 mm in the Spotted-tail moray (*Gymnothorax equatorialis*). The species with the highest relative abundance were: *Haemulopsis axillaris* (Steindachner, 1869), *Syacium ovale* (Günther, 1864), *Selene peruviana* (Guichenot, 1866), *Bothus constellatus* (Jordan, 1889), *Diapterus peruvianus* (Cuvier and Valenciennes, 1830), *Syacium latifrons* (Jordan and Gilbert, 1882), *Scorpaena russula* (Jordan and Bollman, 1889), *Eucinostomus currani* (Zauranec, 1967), *Haemulon scudderii* (Gill, 1863), *Prionotus stephanophrys* (Lockington, 1881), and *Larimus acclivis* (Jordan and Bristol, 1898). *H. axillaris*, *S. ovale*, *S. peruviana*, *D. peruvianus*, *L. acclivis* and *Stellifer erycimba* were the most frequent species in the inner shelf (< 40 m depth), whereas *P. stephanophrys*, *S. russula*, *Porichthys analis* and *Synodus scituliceps* were dominant in the outer shelf (40-60 m depth).

Individual fish size and weight showed a similar general pattern along the four years. Most of the fishes (40 %) presented a standard

length range of 75 to 107 mm with a mean of 103 ± 31.8 . Individual weight including 70 % of the fishes varied between 1.25 to 32.5 g with a mean of 32.5 ± 32.7 . Large fishes (> 50 cm SL and 2000 g) were also found in shrimp by-catch, although their abundance may be underestimated due to reduced catchability related to trawl speed and net selectivity. Most of these fishes were elasmobranch of commercial interest like *Narcine vermiculatus*, *Gymnura marmorata* and *Rhinobatos glaucoptigma*, and teleosts like *Epinephelus multiguttatus*, *Hyporthodus exsul*, *H. acanthisthius*, *Ophioscion typicus*, *Carangoides otrynter*, *Lutjanus guttatus*, *L. peru*, *Paralichthys woolmani*, *Cyclopsetta querna*, *Trichiurus lepturus*, *Centropomus robalito*, *Micropogonias altipinnis* and *Cynoscion phoxocephalus*, which represented 25 % of the biomass. Fish catches during day and night showed significant differences (t-test = 1.93, P < 0.05). Also catches recorded below 40 m depth presented a significant difference (t-test = -5.67, P < 0.05) with those obtained deeper than 40 m.

The community was composed mainly by species with subtropical and tropical affinities (Table 1). Subtropical and tropical fish represented 34.4 % of the catch with a wide geographical distribution from the Cortés to the Panamic Province; 27.0 % were eury-thermic species from the San Diego to the Panamic Province, 13.3 % were restricted to the Mexican and Panamic Provinces, and 9.3 % were species with a distribution in the Eastern Tropical Pacific (Fig. 2). The fish community




Fig. 2. Ichthyogeographic affinities (%) in Provinces along the Eastern Pacific coast (PO, Oregonian; PS, San Diego; PC, Cortés; PM, Mexican; PP, Panamic; POT, Tropical Eastern Pacific; CT, Circumtropical species; T, Trans-Pacific species).

Fig. 2. Afinidad ictiogeográfica (%) en las provincias a lo largo de la costa oeste del Pacífico (PO: Provincia Oregoniana, PS: Provincia de San Diego, PC: Provincia de Cortés, PM: Provincia Mexicana, PP: Provincia Panámica, POT: Pacífico Oriental Tropical, CT: Especies Circumtropicales, y T: Especies Transpacíficas).

was constituted by 32.8 % of common species, whereas 28.6 % are frequent species, 30.3 % are rare species and the remaining 8.3 % were abundant.

New records for the area were: *Notarius planiceps*, *Ophioscion typicus*, *Mugil setosus*, *Alphestes immaculatus*, *Chilara taylori*, *Decapterus muroadsi*, *Lepidopus fitchi* and *Stellifer chrysoleuca*. The recorded distribution within the area was extended for *Urotrygon reticulata*, *Anchoa helleri*, *Cathorops steindachneri*, *Bollmannia stigmatura* and *Sphoeroides kendalli*.

In terms of abundance, the most important families were Haemulidae, Sciaenidae, Carangidae, Ariidae and Serranidae, which also contributed with the greatest number of species and abundance. These species are central to the understanding of the structure and function of the community of demersal fishes of the area. Its euryhaline capacity explains its abundance in front of the lagoons of the Huave, Mar Muerto, La Joya-Buenavista, Carretas-Pereyra and Chantuto-Panzacola systems.

Of the 242 species recorded, 70 (30 %) made incursions into the coastal lagoons for various purposes such as spawning and feeding. Notable for their abundance were those in the Huave (Oaxaca) and Chantuto Panzacola (Chiapas) systems: *Lile stolifera*, *L. gracilis*, *Diapterus peruvianus*, *Micropanchax altipinnis*, *Achirus zebrinus*, *A. scutum*, *Anchoa nasus*, *A. mundeola*, *Centropomus robalito*, *Cathorops fuerthii*, *Eucinostomus currani* and *Cyclopsetta panamensis*.

Three species (1.3 %) were typically freshwater species: *Poecilia butleri*, *Poecilopsis fasciata* and *Gobiomorus maculatus*. They were caught near the outlet of the Mar Muerto lagoon at 16-18 m depth, possibly due to the influence of a plume of brackish water intruding the continental shelf.

The total number of species registered in 2003 (131) was higher than the one found in 2013 (116), whereas species biogeographical affinities did not present a significant difference ($F_{12,9} = 4.41$, d.f. = 21.0, $P > 0.447$) among 2003 and 2013.



DISCUSSION

The species richness of demersal fishes recorded as by-catch from the shrimp fishery in the Gulf of Tehuantepec, with 242 species, is typical of the Eastern Pacific (Robertson & Cramer, 2009). Only 34 species (15 %) were frequent or abundant while the others were common and rare. Some of the rare species are epipelagic species (*Opisthonema libertate*, *Scomberomorus sierra*, *Decapterus macarellus* and *Cypselurus callopterus*), which probably may have been caught while the net was being recovered.

Fish by-catch individual size and weight were consistently small, with 92 % under 50 g weight and 20 cm SL, respectively. A similar composition has been previously reported for the Gulf of California (Pérez Mellado & Findley, 1985). These authors pointed out that about 1 % were fishes weighting more than 100 g whereas in this study the percentage was remarkably higher (3.4 %). These fishes, usually higher than 350 g, were frequent in by-catch and are mainly used for human consumption.

On previously compiled lists of species for the demersal fishery on the continental shelf in the Gulf of Tehuantepec, recorded 32 families, 49 genera and 61 species (Secretaría de Marina, 1978). Acal and Arias (1990) registered 292 species, but that list contained 97 inconsistencies; 50 species were not identified but simply assigned to genus, 24 others were only assigned to family, six constituted synonyms within the list, and others were clearly out of context since they were species recorded for the Atlantic Ocean; the record has accordingly been adjusted.

Of the species in previous lists from the Gulf of Tehuantepec, 167 (74 %) were also found in the present study but 99 were not. Compiling these studies we obtained a total of 331 species, which supports the fact that recorded fish species in the Mexican South Pacific are higher than those for the Western shelf of Baja California Sur, the Gulf of California, the Jalisco and Colima shelf, and the shelves of Nayarit, Michoacán and Guerrero.

This number may be greater if species that belong to the rocky, pelagic and slope zones, and to the lagoons and estuaries, are considered.

From a biogeographical point of view, the assignment of this ichthyofauna to the classical faunistic provinces (Cortés, Panamic and Mexican) presents difficulties, particularly because it is constituted by species with tropical affinities that overlap with others with subtropical or temperate affinities. In addition, it must be considered that the conjunction of provinces (Cortés-Panamic) corresponds to 12 zoogeographic divisions. This marks a distinct change in association, frequency and diversity of species from South to North, reflecting the great variety of habitats.

In the Northern area, the species were temperate ones: *Bellator xenisma*, *Diplectrum pacificum*, *Epinephelus analogus*, *Sphoeroides lobatus*, *Synodus evermanni*, *G. marmorata*, *Ophidion galeoides*, *Opisthonema medirastre* and *Merluccius productus*. To the South, the predominant species are typical of the Tropical Eastern Pacific, such as *Bellator loxias*, *Porichthys analis*, *Ancylopsetta dendritica*, *Aluterus monoceros*, *Trinectes fimbriatus*, *Diplectrum macropoma*, *Urotrygon munda* and *Paralonchurus rathbuni*. Horn et al. (2006) and Rodríguez-Romero, Hernández-Vázquez and López-Martínez (2009) noted similar tendencies.

The high diversity recorded in the Gulf of Tehuantepec, while demonstrating the importance and uniqueness of the area, reflects the concurrence of a series of environmental factors that interact, characterizing the region as a dynamic frontier or transition ecosystem with a broad thermic regime and a variety of habitats (Díaz-Ruiz, Cano-Quiroga, Aguirre-León, & Ortega-Bernal, 2004; Tapia-García, García-Abad, Carranza-Edwards & Vázquez-Gutiérrez, 2007; Velázquez-Velázquez, Vega-Cendejas & Navarro-Alberto, 2008). This uniqueness is supported not only by its geographic position with tropical and temperate characteristics, but also by: 1) the contribution of river runoff, 2) the presence of extensive coastal lagoon systems with high productivity,



3) the prevailing winds and seasonal rainfall from land to the sea, and 4) a wide continental shelf influenced by a complex oceanic current system (the sub-surface equatorial, the North-equatorial countercurrent, the California current, and the South-equatorial or coastal current of Costa Rica that flows northwards and favors the presence of important upwellings) (Galván-Magaña et al., 2000; López-Martínez, Herrera-Valdivia, Rodríguez-Romero, & Hernández-Vázquez, 2010; Mora & Robertson, 2005; Rodríguez-Romero et al. 1998; Rodríguez-Romero, Palacios-Salgado, López-Martínez, Hernández-Vázquez, & Ponce-Díaz, 2008; Zapata & Robertson, 2007).

In addition, the species of the Tropical Eastern Pacific receive strong immigration on its coasts, particularly during events such as ENSO (El Niño - Southern Oscillation), which reach the East Pacific coasts when it is of great magnitude and intensity (Lea & Rosenblatt, 2000). Their periodicity and magnitude contribute significantly to dispersion, since most fishes have life cycles with pelagic larval stages (Auster, 1988; Caddy & Sharp, 1986; Galzin & Legendre, 1987; Grossman, Freeman, Moyle, & Whitaker, 1985), which affect the distribution patterns since long time. It is considered that changes in abundance and composition of the fish community frequently occur as a result of migratory movements between neighbouring geographic areas driven by climatic and oceanographic processes of medium or long-term duration (Safran, 1990), as well as other factors such as changes in the rates of hatching and mortality, owing to abiotic environmental factors (temperature, oxygen) or biotic factors (predation, competition, food availability).

The results show the uniqueness and importance of the region in relation to the fish species richness of great value for industrial and artisanal fisheries. In this context, it would be important to include, in the actual local policies, rules that could regulate the technical aspects of the fishing methods used in the Gulf of Tehuantepec in order to preserve this ecosystem richness. Consequently, scientific research appropriate to the particular needs of

this area of the Tropical Eastern Pacific should increase, so long as the information is scarce and temporarily discontinuous.

ACKNOWLEDGMENTS

To the University of Barcelona for a collaborative grant, in particular to the Department of Ecology for the help in its computer section, library and laboratories; without these facilities it would have been difficult to proceed with the doctoral study of the first author. Thanks to the Institute of Marine Sciences of Barcelona, particularly the Department of Renewable Marine Resources of CSIC for granting the support and infrastructure necessary for this study. To the Regional Centre of Fisheries Research, Salina Cruz, of the Mexican National Institute of Fisheries and its personnel, for the opportunity to participate in the fishing expeditions in 2003 and 2005, and for the help in the laboratory and hospitality in their premises. Thanks to the Postgraduate Department of the Institute of Marine Sciences and Limnology, UNAM, for the assistance during a postdoctoral residency and for the translation of the manuscript. This study was partially supported by the Dirección General de Asuntos del Personal Académico, Universidad Nacional Autónoma de México (PAPIIT), Grant IN211214-2.

RESUMEN

En la pesquería del camarón del Golfo de Tehuantepec un gran número de especies de peces se capturan como fauna de acompañamiento y son descartadas. La composición, distribución y afinidades biogeográficas de la ictiofauna acompañante del camarón fue analizada mediante 15 cruceros desarrollados en la plataforma continental entre 15-64 m de profundidad durante 2003, 2004, 2005 y 2013. La ictiofauna descartada estuvo representada por 58 familias, 129 géneros y 242 especies. Las familias, Haemulidae, Sciaenidae, Paralichthyidae, Gerreidae y Carangidae aportaron más del 70 % de la captura. *Haemulopsis axillaris*, *Syacium ovale*, *Selene peruviana*, *Dipterus peruvianus*, *Larimus acclivis* y *Stellifer erycimba* fueron las especies más frecuentes en profundidades menores de 40 m (plataforma interna), mientras que *Prionotus stephanophrys*, *Scorpaena russula*, *Porichthys analis* y *Synodus scituliceps*, fueron dominantes entre 40-60 m de profundidad



(plataforma externa). El análisis de las afinidades biogeográficas mostró que el 36.1 % de las especies son de amplia distribución desde la provincia de San Diego a la Panámica, mientras que el 13.2 %, presentó una distribución restringida entre la provincia Mexicana y la Panámica. La composición de la ictiofauna estuvo marcadamente influenciada por las condiciones ambientales locales y sus variaciones estacionales.

Palabras clave: biogeografía, ictiofauna, camarón, pesca, fauna de acompañamiento, Golfo de Tehuantepec.

REFERENCES

- Acal, E. D. & Arias, A. (1990). Evaluación de los recursos demerso-pelágicos vulnerables a redes de arrastre de fondo en el sur del Pacífico de México. *Ciencias Marinas*, 16, 93-129.
- Aguilar-Palomino, B., Mariscal, R. J., González, S. G., & Rodríguez, I. L. E. (1996). Lista sistemática de la ictiofauna demersal de fondos blandos de la plataforma continental de Jalisco y Colima México, durante mayo y junio de 1995. *Ciencias Marinas*, 22, 469-481.
- Allsop, W. H. L. (1985). La fauna acompañante del camarón: perspectivas y manejo. En A. Yáñez-Arancibia (Ed.). *Recursos pesqueros potenciales de México: La pesca acompañante del camarón. Programa Universitario de Alimentos* (pp. 635-643). México D.F.: Instituto de Ciencias del Mar y Limnología, Instituto Nacional de la Pesca, UNAM.
- Amezcu-Linares, F. (1996). *Peces demersales de la Plataforma Continental de México*. México: UNAM/ICMYL/CONABIO.
- Andrew, N. L., & Pepperell, J. G. (1992). The by-catch of shrimp trawl fisheries. *Oceanography and Marine Biology: An Annual Review*, 30, 527-565.
- Auster, P. J. (1988). A review of present state of understanding of marine fish communities. *Journal of Northwest Atlantic Fisheries Science*, 8, 67-75.
- Bianchi, G. (1991). Demersal assemblages of the continental shelf and slope edge between the Gulf of Tehuantepec (México) and Gulf of Papagayo (Costa Rica). *Marine Ecology Progress Series*, 73, 121-140.
- Boschi, E. E. (2000). Species of decapod crustaceans and their distribution in the American marine zoogeographic provinces. *Revista de Investigación y Desarrollo Pesquero*, 13, 7-136.
- Briggs, J. C. (1974). *Marine Zoogeography*. New York: McGraw-Hill.
- Briggs, J. C. (1995). *Global biogeography*. Amsterdam, Holland: Elsevier.
- Bussing, W., & López, M. (1993). Peces demersales y pelágicos costeros del Pacífico de Centro América Meridional. *Revista de Biología Tropical, Publicación Especial*, 1, 1-164.
- Caddy, J. F., & Sharp, G. D. (1986). An ecological framework for marine fishery investigations. *FAO Fisheries Technical Paper*, 283, 152.
- Castro-Aguirre, J. L. (1978). *Catálogo sistemático de los peces marinos que penetran a las aguas continentales de México, con aspectos zoogeográficos y ecológicos*. México: Instituto Nacional de la Pesca, Depto. Pesca.
- Castro-Aguirre, J. L. & Espinosa Pérez, H. (1996). *Listados faunísticos de México. VII. Catálogo sistemático de las rayas y especies afines de México (Chondrichthyes: Elasmobranchii: Rajiformes: Batoideiomorpha)*. México, D.F: Instituto de Biología, UNAM.
- Cervantes-Hernández, P., Gallardo-Berumen, M. I., Ramos-Cruz, S., Gómez-Ponce, M. A., & Gracia, A. (2008). Análisis de las temporadas de veda en la explotación marina de camarones del Golfo de Tehuantepec, México. *Revista de Biología Marina y Oceanografía*, 43, 285-294.
- Díaz-Ruiz, S., Cano-Quiroga, E., Aguirre-León, A., & Ortega-Bernal, R. (2004). Diversidad, abundancia y conjuntos ictiofaunísticos del sistema lagunar-estuarino Chantuto-Panzacola, Chiapas, México. *Revista de Biología Tropical*, 52, 187-199.
- Eschmeyer, W. N. (1998). *Catalog of fishes. Special Publication*, 1. San Francisco: California Academy of Sciences.
- Espinosa-Pérez, H. J., Castro-Aguirre, L., & Huidobro-Campos, L. (2004). *Listados faunísticos de México. IX. Catálogo sistemático de tiburones (Elasmobranchii: Selachimorpha)*. México, D.F., México: Instituto de Biología UNAM.
- Fischer, W., Krupp, F., Schneider, W., Sommer, C., Carpenter, K. E., & Niem, V. H. (1995). *Guía FAO para la identificación de especies para los fines de la pesca Pacífico centro-oriental* (Volumen III. Vertebrados, Parte 2). Roma: FAO.
- Froese, R., & Pauly, D. (2013). *FishBase*. Retrieved from <http://www.fishbase.org>.
- Galván-Magaña, F., Gutiérrez-Sánchez, F. J., Abitia-Cárdenas, L. A., & Rodríguez-Romero, J. (2000). The distribution and affinities of the shore fishes of Baja California Sur lagoons. In M. Munawar, S. G. Lawrence, I. F. Munawar, & D. F. Malley (Eds.). *Aquatic ecosystems of Mexico* (pp. 383-398). Leiden: Ecovision world monograph series. Backhuys Publishers.
- Galzin, R., & Legendre, P. (1987). The fish communities of a coral reef transect. *Pacific Science*, 41, 158-165.



- Ginsburg, I. (1958). Flounders of the genus *Paralichthys* and related genera in American waters. *U.S. Fisheries and Wildlife Service, Fishery Bulletin*, 52, 267-351.
- Griffiths, S., Larson, H., & Courtney, T. (2004). *Trawl bycatch species in National Oceans Office, Description of key species groups in the northern planning area*. Hobart, Australia: National Oceans Office.
- Grossman, G. D., Freeman, M. C., Moyle, P. B., & Whittaker Jr., J. O. (1985). Stochasticity and assemblage organization in an Indiana stream fish assemblage. *American Naturalist*, 126, 275-285.
- Hastings, P. A. (2000). Biogeography of the tropical eastern Pacific: distribution and phylogeny of chaenopsid fishes. *Zoological Journal of Linnean Society*, 128, 319-335.
- Hendrickson, H. M., & Griffin, W. L. (1993). An analysis of management policies for reducing shrimp by-catch in the Gulf of Mexico. *North American Journal of Fisheries Management*, 13, 686-697.
- Horn, M. H., Allen, L. G., & Lea, R. N. (2006). Biogeography. In L. G. Allen, D. J. Pondella, & M. H. Horn (Eds.). *The ecology of marine fishes, California and adjacent waters* (pp. 3-25). California, U.S.A.: University of California Press, Berkeley.
- Lea, R. N., & Rosenblat, R. H. (2000). Observations on fishes associated with the 1997-98 El Niño off California. *CalCOFI Report*, 41, 117-129.
- López-Martínez, J., Herrera-Valdivia, E., Rodríguez-Romero, J., & Hernández-Vázquez, S. (2010). Composición taxonómica de peces integrantes de la fauna de acompañamiento de la pesca industrial de camarón del Golfo de California, México. *Revista de Biología Tropical*, 58, 925-942.
- Love, M. S., Mecklenburg, C. W., Mecklenburg, T. A., & Thorsteinson, L. K. (2005). *Resource Inventory of Marine and Estuarine Fishes of the West Coast and Alaska: A Checklist of North Pacific and Arctic Ocean Species from Baja California to the Alaska-Yukon Border*. Washington: U. S. Department of the Interior, U. S. Geological Survey, Biological Resources Division, Seattle, Washington.
- McPhail, J. D. (1958). Key to the croakers (Sciaenidae) of the eastern Pacific. University of British Columbia. *Institute of Fisheries, Museum Contributions*, 2, 1-20.
- Madrid-Vera, J., Ruiz, A. L., & Rosado, B. I. (1998). Peces de la plataforma continental de Michoacán y sus relaciones en el Pacífico mexicano. *Revista de Biología Tropical*, 46, 267-276.
- Moncayo-Estrada, R., Castro-Aguirre, J. L., & De La Cruz-Agüero, J. (2006). Lista sistemática de la ictiofauna de Bahía de Banderas, México. *Revista Mexicana de Biodiversidad*, 77, 67-80.
- Mora, C., & Robertson, D. R. (2005). Causes of latitudinal gradients in species richness: a test with the endemic shorefishes of the Tropical Eastern Pacific. *Ecology*, 86, 1771-1782.
- Nelson, J. S. (2006). *Fishes of the world*. New Jersey, USA: John Wiley & Sons.
- Nelson, J. S., Crossman, E. J., Espinosa-Pérez, H., Findley, L. T., Gilbert, C. R., Lea, R. N., & Williams, J. D. (2004). Common and scientific names of fishes from the United States, Canada, and Mexico. *American Fisheries Society Special Publication*, 29, Maryland, EUA.
- Norman, J. R. (1934). *A systematic monograph of the flatfishes (Heterosomata) Vol. I. Psettodidae, Bothidae, Pleuronectidae*. London: British Museum Natural History.
- Pérez-Mellado, J., & Findley, L. T. (1985). Evaluación de la ictiofauna acompañante del camarón capturado en la costa de Sonora y norte de Sinaloa, México. En A. Yáñez-Arancibia (Ed.), *Recursos pesqueros potenciales de México: La pesca acompañante del camarón* (pp. 201-254). México: PUA, ICMYL, UNAM/INP.
- Reyna Cabrera, I. E., & Ramos-Cruz, S. (1998). La pesquería de camarón de alta mar. En M. Tapia-García (Ed.), *El Golfo de Tehuantepec: el ecosistema y sus recursos* (pp. 163-178). México: Universidad Autónoma Metropolitana-Iztapalapa.
- Robertson, D. R., & Allen, G. R. (2002). *Shore fishes of the Tropical Eastern Pacific: an Information System. CD-ROM*. Panamá: Smithsonian Tropical Research Institute, Balboa, Panamá.
- Robertson, D. R., & Cramer, K. L. (2009). Shore fishes and biogeographic subdivisions of the Tropical Eastern Pacific. *Marine Ecology Progress Series*, 380, 1-17.
- Robertson, D. R., Grove, J. S., & McCoster, J. S. (2004). Tropical trans-Pacific shore fishes. *Pacific Science*, 58, 507-565.
- Rodríguez-Romero, J., Abitia, C. L., Galván, M. F., Gutiérrez, S., Aguilar, P. B., & Arvizú, M. J. (1998). Ecology of fish communities from the soft bottoms of Bahía Concepción, México. *Archive of Fishery Marine Research*, 46, 61-76.
- Rodríguez-Romero, J., Palacios-Salgado, D. S., López-Martínez, J., Hernández-Vázquez, S., & Ponce-Díaz, G. (2008). Composición y zoogeografía de los peces demersales de la costa occidental de Baja California Sur, México. *Revista de Biología Tropical*, 56, 1765-1783.
- Rodríguez-Romero, J., Hernández-Vázquez, S., & López-Martínez, J. (2009). Desarrollo potencial de peces desaprovechados. *Ciencia y Desarrollo*, 35, 45-51.



- Rosemblatt, R. H., & Johnson, G. D. (1974). Two new species of sea basses of the genus *Diplectrum* with a key to the Pacific species. *California Fish and Game*, 60, 178-191.
- Safran, P. (1990). Spatio-temporal variability in the structure of a nectobenthic fish nursery: A descriptive study. *Oceanologica Acta*, 13, 97-106.
- Sarmiento-Náfate, S., Gil-López, H. A., & Arroyo, D. (2007). Shrimp by-catch reduction using a short funnel net, in the Gulf of Tehuantepec, South Pacific, Mexico. *Revista Biología Tropical*, 55, 889-897.
- Secretaría de Marina. (1978). *Estudio oceanográfico del Golfo de Tehuantepec. Tomos I-III. (Biología, Física, y Química; Geología)*. México: Secretaría de Marina. Dirección General de Oceanografía. Talleres de la Nación, México.
- Siqueiros-Beltrones, D. A., & De la Cruz Agüero, J. (2004). Examen filosófico de las listas sistemáticas como informes científicos y porqué deben ser publicados. *Océanides*, 19, 1-9.
- Sokal, R. R., & Rohlf, F. J., (1995). *Biometry*. New York, USA: W.H. Freeman and Co.
- Stobutzki, I. C., Miller, M. J., Jones, P., & Salini, J. P. (2001). Bycatch diversity and variation in a tropical Australian peneaeid fishery; the implications for monitoring. *Fisheries Research*, 53, 283-301.
- Tapia-García, M., García-Abad, M. C., González-Medina, G., Macuitl-Montes, M. C., & Cerdáñoles-Ladrón de Guevara, G. (1994). Composición, distribución y abundancia de la comunidad de peces demersales del Golfo de Tehuantepec. *Tropical Ecology*, 35, 229-252.
- Tapia-García, M. (1998). Evaluación ecológica de la fauna demersal. In M. Tapia-García (Ed.), *El Golfo de Tehuantepec: el ecosistema y sus recursos* (pp. 129-148). México, D. F: Universidad Autónoma Metropolitana-Iztapalapa.
- Tapia-García, M., & García-Abad, M. C. (1998). Los peces acompañantes del camarón y su potencial como recurso en las costas de Oaxaca y Chiapas. In M. Tapia-García (Ed.), *El Golfo de Tehuantepec: el ecosistema y sus recursos* (pp. 179-196). México, D.F: Universidad Autónoma Metropolitana-Iztapalapa.
- Tapia-García, M., García-Abad, A., Carranza-Edwards, A., & Vázquez-Gutiérrez, F. (2007). Environmental characterization of the continental shelf of the gulf of Tehuantepec, Mexico. *Geofísica Internacional*, 7, 249-260.
- Velázquez-Velázquez, E., Vega-Cendejas, M. E., & Navarro-Alberto, J. (2008). Spatial and temporal variation of fish assemblages in a coastal lagoon of the Biosphere Reserve La Encrucijada, Chiapas, Mexico. *Revista de Biología Tropical*, 56, 557-574.
- Walker, B. W. (1960). The distribution and affinities of the marine fish fauna of the Gulf of California. *Systematic Zoology*, 9, 123-133.
- Walter, H. J., & Rosemblatt, R. H. (1988). Pacific toadfishes of the genus *Porichthys* (Batrachoididae) with description of three new species. *Copeia*, 4, 887-904.
- Zapata, F. A., & Robertson, D. R. (2007). How many species of shore fishes are there in the Tropical Eastern Pacific? *Journal of Biogeography*, 34, 38-51.
- Zar, J. H. (1999). *Biostatistical analysis*. New York, USA: Prentice Hall.