

DNA barcodes reveal 63 overlooked species of Canadian beetles (Insecta, Coleoptera)

Mikko Pentinsaari¹, Robert Anderson², Lech Borowiec³, Patrice Bouchard⁴, Adam Brunke⁴, Hume Douglas⁴, Andrew B.T. Smith², Paul D. N. Hebert¹

1 Centre for Biodiversity Genomics, 50 Stone Road East University of Guelph, Guelph, ON, N1G 2W1, Canada **2** Canadian Museum of Nature, P.O. Box 3443, Station D, Ottawa, ON, K1P 6P4, Canada **3** Department of Biodiversity and Evolutionary Taxonomy, University of Wrocław, Przybyszewskiego 65, 51-148 Wrocław, Poland **4** Canadian National Collection of Insects, Arachnids and Nematodes, Agriculture and Agri-Food Canada, 960 Carling Avenue, Ottawa, ON, K1A 0C6, Canada

Corresponding author: Mikko Pentinsaari (mpentins@uoguelph.ca)

Academic editor: A. Smith | Received 5 July 2019 | Accepted 25 October 2019 | Published 4 December 2019

<http://zoobank.org/D11503CA-5A57-4067-8179-04E0C8C162C8>

Citation: Pentinsaari M, Anderson R, Borowiec L, Bouchard P, Brunke A, Douglas H, Smith ABT, Hebert PDN (2019) DNA barcodes reveal 63 overlooked species of Canadian beetles (Insecta, Coleoptera). ZooKeys 894: 53–150. <https://doi.org/10.3897/zookeys.894.37862>

Abstract

This study demonstrates the power of DNA barcoding to detect overlooked and newly arrived taxa. Sixty-three species of Coleoptera representing 25 families are studied based on DNA barcode data and morphological analysis of the barcoded specimens. Three of the species involve synonymies or previous taxonomic confusion in North America, while the first Canadian records are published for 60 species. Forty-two species are adventive in North America, and 40 of these adventive species originate from the Palaearctic region. Three genera are recorded from the Nearctic region for the first time: *Coelostoma* Brullé, 1835 (Hydrophilidae), *Scydmorephes* Reitter, 1891 (Staphylinidae), and *Lythraria* Bedel, 1897 (Chrysomelidae). Two new synonymies are established: *Mycetoporus triangulatus* Campbell, 1991 (Staphylinidae) is a junior synonym of *Mycetoporus reichei* Pandellé, 1869, **syn. nov.** while *Bledius philadelphicus* Fall, 1919 (Staphylinidae) is a junior synonym of *Bledius gallicus* (Gravenhorst, 1806), **syn. nov.** The previously suggested move of *Ctenicera tigrina* (Fall, 1901) to the genus *Pseudanostirus* Dolin, 1964 (Elateridae) is formalized, resulting in *Pseudanostirus tigrinus* (Fall, 1901), **comb. nov.**

Keywords

DNA barcoding, new species records, adventive species

Table of contents

Introduction.....	55
Materials and methods	56
Material	56
Tissue sampling and DNA barcode sequencing.....	57
Identification and validation of the new Canadian species records.....	58
Species accounts.....	58
Availability of data.....	59
Results.....	59
Gyrinidae.....	62
Carabidae.....	62
Hydrophilidae.....	62
Leiodidae.....	64
Staphylinidae.....	66
Scarabaeidae.....	86
Clambidae.....	87
Scirtidae.....	89
Throscidae.....	92
Elateridae.....	96
Cantharidae.....	97
Dermestidae.....	102
Ptinidae.....	102
Erotylidae.....	103
Cryptophagidae.....	105
Phalacridae.....	105
Nitidulidae.....	108
Coccinellidae.....	109
Corylophidae.....	114
Mycetophagidae.....	116
Ciidae.....	116
Mordellidae.....	119
Zopheridae.....	119
Tenebrionidae.....	120
Chrysomelidae.....	122
Curculionidae.....	127
Discussion.....	134
Acknowledgements.....	136
References.....	137
Supplementary material 1.....	150

Introduction

Since being proposed as a standardized approach for identifying unknown specimens to species-level (Hebert et al. 2003) DNA barcoding has become a global research effort. By May 2019, the Barcode of Life Data Systems (BOLD, <http://boldsystems.org/>; Ratnasingham and Hebert 2007)) contained more than 7 million DNA barcode records. The utility of DNA barcodes for biosecurity and for the detection of invasive species was recognized soon after their initial proposition (Armstrong and Ball 2005). In New Zealand, DNA barcoding has been adopted as a routine tool for screening for new potential pest species in certain insect taxa (Armstrong 2010). Invasive species are one of the most important threats to biodiversity worldwide (Bellard et al. 2016; Maxwell et al. 2016). They are also responsible for major economic losses to agriculture and forestry; invasive species are estimated to cause between \$100–200 billion in losses annually in the United States of America alone (Pimentel 2011). Timely detection and accurate identification of new adventive species is important for efficient monitoring and management of potential pests. However, if an adventive species belongs to a taxonomically difficult or neglected group of morphologically similar species, it can remain undetected for decades (deWaard et al. 2010; Jendek et al. 2015).

Among the 8,302 species of Coleoptera known from Canada, 639 are adventive (Brunke et al. 2019). A series of publications by Klimaszewski et al. (2010, 2012, 2013, 2015, 2017) provides both diagnostic features and overviews of the ecology and known North American distribution of these species. While Europe has been and still is a major source of these species because of the intensive trade and ship traffic across the Atlantic, the proportion of Asian species has recently increased as a consequence of increased trade with this region (Klimaszewski et al. 2012). The establishment of Palaearctic species in North America is likely facilitated by the climatic similarity of the regions (Klimaszewski et al. 2013).

The beetle fauna of North and Central Europe has recently been DNA barcoded extensively (Pentinsaari et al. 2014; Hendrich et al. 2015; Rougerie et al. 2015). These barcode reference libraries provide an efficient tool for the detection of adventive European beetle species in other parts of the world, particularly for those taxonomic groups where the native fauna is poorly known. Detection of species shared between continents is greatly facilitated by the Barcode Index Number (BIN) system (Ratnasingham and Hebert 2013). The BIN system was created primarily as an interim taxonomic framework for the COI barcode records on BOLD which lack species level identifications. However, comparison between BINs and Linnaean species names has proved highly useful in uncovering misidentifications, synonymies, and overlooked species diversity (e.g., Landry et al. 2013). In European Coleoptera, BINs showed a perfect 1:1 correspondence to known species in ca. 90% of the studied species (Pentinsaari et al. 2014; Hendrich et al. 2015).

The Canadian arthropod fauna has been extensively sampled for DNA barcoding over the past decade, both in the field and in natural history collections (see e.g., Gwiazdowski et al. 2015, Hebert et al. 2016, Bouchard et al. 2017, Steinke et al. 2017). A joint analysis of all available European and Canadian beetle data allows rap-

id screening for species shared between continents on a wide taxonomic scale. Such screening in equivalent barcode libraries of Lepidoptera has revealed multiple new species records and synonymies (Landry et al. 2013), as well as previously overlooked species lineages (Mutanen et al. 2012).

This paper reports the first Canadian records for 60 species of beetles, which were initially detected based on DNA barcoded specimens, and resolves previous taxonomic confusion in three more species. Twenty-one species represent native North American taxa recently arrived or previously overlooked in Canada. Forty-two species are adventive, and at least four are potential pests. Two species described from North America were found to be synonyms of Palearctic species and hence are now properly recognized as adventive to the Nearctic region. We provide morphological diagnoses and illustrations for all adventive species, and for those 12 native North American species for which they are not readily available elsewhere.

Materials and methods

Material

This publication is based on the analysis of more than 130,000 DNA barcode records from Europe and Canada. The combined dataset of European and Canadian Coleoptera was screened for intercontinentally shared species. As part of the cleaning and validation process of a barcode reference library for Canadian Coleoptera, representative specimens of Canadian Barcode Index Number (BIN) clusters lacking species-level identifications were retrieved for morphological analysis. After identification and validation of new species records and synonyms, 1168 DNA barcode records (sequence length ≥ 400 bp) representing 63 species were selected for publication. Most (1147) of these records derive from freshly collected specimens obtained through projects coordinated by the Centre for Biodiversity Genomics, University of Guelph (CBG) such as the Canadian National Parks Malaise Program (<http://biodiversitygenomics.net/projects/cnp/>), the School Malaise Trap Program (Steinke et al. 2017, <https://malaiseprogram.com/>), and BIObus collecting trips across Canada (<https://biobus.ca/>). As these specimens are stored pinned or in ethanol in the CBG voucher archive, they were available for morphological study and species assignment once barcode sequences were available. The three specimens of *Attagenus smirnovi* Zhantiev, 1973 were submitted for DNA barcoding through the LifeScanner citizen science initiative (<http://www.lifescanner.net/>) and are stored in the CBG voucher specimen archive. One of the DNA barcoded specimens of *Contacyphon kongsbergensis* (Munster, 1924) is stored in the Wallis-Roughley Museum of Entomology (JBWM). As part of our effort to construct a DNA barcode reference library for Canadian beetles, we analyzed 15,811 specimens of beetles held in the Canadian National Collection of Insects, Arachnids, and Nematodes (CNC). Although sequence recovery from the CNC samples was lower than with freshly collected material, their analysis provided a wide set of well-identified reference specimens (Bouchard

et al. 2017). Seventeen of the successfully sequenced CNC specimens were found to represent new species for Canada, and are included in the dataset published here.

In addition to the barcoded material, we examined 303 specimens without DNA barcode data to obtain a more detailed understanding of the Canadian distribution of some of the newly detected species. Of these additional specimens, 257 are deposited in CNC, five in the University of Guelph Insect Collection (**DEBU**), four in the Canadian Museum of Nature (**CMNC**), and two in the Field Museum of Natural History, Chicago. Thirty-two additional records of *Notaris scirpi* (Fabricius, 1793) are from specimens deposited in the private insect collections of Claude Chantal (**CCCH**), Stéphane Dumont (**CSDU**), Pierre de Tonnancour (**CPTO**), and Robert Vigneault (**CRVI**). Three additional records of *Carpelimus elongatulus* (Erichson, 1839) are from specimens in the private insect collection of Reginald Webster (**RWC**).

Tissue sampling and DNA barcode sequencing

The tissue sampling protocol varied according to the origin of the material and size of the specimen. A single leg was detached from each CNC specimen and it was placed in a well in a 96-well microplate pre-filled with 10 µl of 96% ethanol. Each CNC specimen was also photographed, and the resultant image was uploaded to BOLD along with the label data. The specimens archived at the CBG were processed in two ways. Small specimens (body length < 6 mm) were placed into a well in a 96 well microplate for DNA extraction. Following DNA extraction, the microplates were refilled with ethanol and the specimens were stored in the microplates in the CBG voucher specimen archive. Larger specimens were either pinned or preserved in ethanol, and a single leg was used for DNA extraction. Photography of each specimen is not a standard element in the workflow because a million specimens are processed yearly at CBG. Instead, representative specimens of new Barcode Index Numbers (BINs, Ratnasingham and Hebert 2013) are retrieved from the archive for photography once specimens have been assigned to a BIN.

DNA extraction, PCR amplification, and Sanger sequencing of the COI barcode region were performed for all specimens at the Centre for Biodiversity Genomics, using standard protocols optimized for large-scale generation of COI barcode data. For detailed descriptions of the protocols, see Ivanova et al. (2006) and deWaard et al. (2008, 2018). For most specimens, a cocktail of the Folmer primers (Folmer et al. 1994) and LepF1 & LepR1 (Hebert et al. 2004) was used in the first PCR amplification attempt. When resources allowed it, specimens for which the initial PCR amplification failed were reanalyzed using primer sets that targeted 307 bp and 407 bp amplicons (Hajibabaei et al. 2006). Bidirectional sequencing was the rule prior to 2013, but subsequent sequences were often obtained through unidirectional analysis. Details on the PCR and sequencing protocol for each specimen are provided in the public BOLD dataset (see Availability of data below).

Identification and validation of the new Canadian species records

All COI barcode sequences on BOLD which fulfill quality criteria (< 1% ambiguous bases; no reading frame shifts, chimeras or obvious contaminations) are automatically assigned into BINs. The founding member sequence of a new BIN cluster must be at least 500 bp long, but shorter sequences (min. 300 bp) can be assigned into existing clusters. A detailed description of the clustering algorithm and the associated informatics workflow is provided by Ratnasingham and Hebert (2013). In short, all sequences are initially clustered based on a fixed threshold of 2.2% divergence (uncorrected p-distance). These initial clusters are then refined by Markov clustering to generate the final BINs. The clustering algorithm is run regularly on BOLD, and new BINs are generated and existing BINs may be split or merged as new sequence data are added.

The new adventive species were initially detected because Canadian specimens shared a BIN assignment with their European counterparts. When available, at least five Canadian voucher specimens were then morphologically examined to confirm the identification. Most of the extensions in the known range of native North American species were detected and validated in the same way, i.e., Canadian specimens were found to share BINs with identified specimens from the United States. A few taxa were encountered during the validation of a DNA barcode reference library for Canadian Coleoptera when representative specimens from BINs lacking a species-level identification were retrieved for morphological analysis. Only those species for which voucher specimens were available and could be reliably validated are included in this paper.

Species accounts

The brief sections on diagnostic information in this paper detail only the most relevant morphological characters for distinguishing each newly detected species from its closest relatives in North America. Due to the variety of beetle taxa covered, these sections cannot employ a completely uniform format. To provide some consistency, the terminology employed and the order in which the characters are presented follows Lawrence et al. (2011). For native North American species with good diagnoses readily available in existing literature, we simply provide references to those diagnoses. In addition to the diagnostic information, we briefly summarize the current Canadian records as well as the distribution of the species outside Canada. Finally, we provide brief ecological data, and consider collecting methods for each species. The order of the families and genera in the species treatments follows Bousquet et al. (2013). Species are ordered alphabetically within genera. We follow Wheeler and Hoebeke (2009) in our use of terms related to non-native species. All distance measures between DNA barcode sequences are reported as uncorrected p-distance (i.e., the proportion of differing nucleotides in pairwise comparisons of sequences).

Availability of data

Detailed collection information for each specimen, including both DNA barcoded material and other specimen records, as well as GenBank accession numbers for the barcode sequences, are provided in the Suppl. material 1: Table S1. All sequences, details on PCR and sequencing primers, photographs (if available) and full collection data for the DNA barcoded specimens are available through a public dataset on BOLD (DS-NEWCOL18, <https://doi.org/10.5883/DS-NEWCOL18>). The public BIN pages for each species can be accessed through the BOLD dataset, or by entering the BIN URIs provided in each species account (e.g., BOLD:AAP7843) in the search field of the public BIN portal: http://boldsystems.org/index.php/Public_BarcodeIndexNumber_Home

Results

The higher classification of the new species, and the research projects and collections from which the specimens originate, are summarized in Table 1. Detailed accounts of each species are provided below. The original distribution, habitat preferences, and possible pest status of the adventive species are summarized in Table 2.

Table 1. Summary of the higher classification of the studied species and the projects and institutions from which the specimens originate. Abbreviations: SMP: School Malaise Program; CNP & OPPMP: Canadian National Parks & Ontario Provincial Parks Malaise programs; CNC: Canadian National Collection of Insects, Arachnids, and Nematodes.

Family	Subfamily	Species	SMP	CNP& OPPMP	BioBus	Other CBG projects	CNC	Other public & private collections
Gyrinidae	Enhydrinae	<i>Dineutus emarginatus</i>			x			
Carabidae	Harpalinae	<i>Anisodactylus caenus</i>					x	
Hydrophilidae	Sphaeridiinae	<i>Coelostoma orbiculare</i>				x		
Leiodidae	Leiodinae	<i>Leiodes polita</i>				x		
Staphylinidae	Pselaphinae	<i>Biblopectus minutissimus</i>				x		
	Tachyporinae	<i>Mycetoporus reichei</i>	x	x				
		<i>Tachyporus atriceps</i>	x	x	x	x		
	Aleocharinae	<i>Amischa decipiens</i>	x			x		
		<i>Atheta vaga</i>				x		
	Aleocharinae	<i>Myllaena infuscata</i>				x		
	Oxytelinae	<i>Bledius gallicus</i>		x		x		
		<i>Carpelimus elongatulus</i>			x	x		x
	Scydmaeninae	<i>Stenichnus collaris</i>			x	x		
		<i>Stenichnus scutellaris</i>	x	x	x	x		
		<i>Scydmorephes minutus</i>		x		x		
		<i>Scydmaenus rufus</i>				x		
	Paederinae	<i>Lathrobium geminum</i>			x			
		<i>Lathrobium lineatocolle</i>		x		x		
		<i>Medon apicalis</i>				x		
		<i>Medon ripicola</i>		x				
		<i>Pseudomedon obscurellus</i>		x				
Scarabaeidae	Melolonthinae	<i>Phyllophaga implicita</i>			x			

Family	Subfamily	Species	SMP	CNP& OPPMP	BioBus	Other CBG projects	CNC	Other public & private collections
Clambidae	Calyptomerinae	<i>Calyptomerus dubius</i>	x			x		
	Clambinae	<i>Clambus simsoni</i>	x					
Scirtidae	Scirtinae	<i>Contacyphon kongsbergensis</i>			x	x		x
		<i>Contacyphon obscurellus</i>		x	x			
		<i>Contacyphon fuscescens</i>		x	x	x		
Throscidae	Throscinae	<i>Aulonothroscus distans</i>	x	x		x		
		<i>Trixagus carinifrons</i>		x		x		
		<i>Trixagus meybohmi</i>	x			x		
Elateridae	Dendrometrinae	<i>Pseudanostirus tigrinus</i>		x				
Cantharidae	Cantharinae	<i>Dichelotarsus lapponicus</i>		x				
	Malthinae	<i>Malthodes pumilus</i>		x	x			
Dermestidae	Attageninae	<i>Attagenus smirnovi</i>				x		
Ptinidae	Dorcatominae	<i>Petalium incisum</i>				x		
Erotylidae	Cryptophilinae	<i>Cryptophilus obliterated</i>	x			x		
		<i>Cryptophilus propinquus</i>				x		
Cryptophagidae	Cryptophaginae	<i>Henoticus mycetoecus</i>		x				
Phalacridae	Phalacrinae	<i>Acylomus ergati</i>	x	x	x	x		
		<i>Olibrus liquidus</i>	x					
Nitidulidae	Eपुरaeinae	<i>Eपुरaea unicolor</i>			x	x		
Coccinellidae	Chilocorinae	<i>Chilocorus renipustulatus</i>	x					
	Coccinellinae	<i>Nephus bisignatus</i>					x	
		<i>Scymnus rubromaculatus</i>	x			x		
Corylophidae	Corylophinae	<i>Orthoperus corticalis</i>				x		
Mycetophagidae	Mycetophaginae	<i>Litargus connexus</i>	x					
Ciidae	Ciinae	<i>Cis boleti</i>				x		x
		<i>Cis glabratus</i>			x			
Mordellidae	Mordellinae	<i>Mordellistena militaris</i>		x				
Zopheridae	Colydiinae	<i>Lasconotus subcostulatus</i>		x			x	
Tenebrionidae	Alleculinae	<i>Isomira angusta</i>			x			
Chrysomelidae	Galerucinae	<i>Chaetocnema hortensis</i>	x	x		x	x	
		<i>Longitarsus lewisii</i>	x					
		<i>Lythvaria salicariae</i>					x	
		<i>Scelolyperus liriophilus</i>		x				
Curculionidae	Brachycerinae	<i>Notaris scirpi</i>					x	x
	Baridinae	<i>Ampelogypter sesostris</i>				x	x	
		<i>Centrinopus helvinus</i>	x					
	Ceutorhynchinae	<i>Ceutorhynchus inaffectatus</i>				x		
		<i>Ceutorhynchus mutabilis</i>			x		x	
Cryptorhynchinae	<i>Peracalles pectoralis</i>			x				
Entiminae	<i>Exomias trichopterus</i>				x			
Scolytinae	<i>Ambrosiodmus rubricollis</i>			x	x			

Table 2. Summary of the original distribution, habitats, and possible pest status of the adventive species.

Family	Species	Original distribution	Habitat	Possible pest status
Cantharidae	<i>Malthodes pumilus</i>	Palaeartic	Dry meadows, warm forest edges, larvae probably in dead wood as predators	–
Chrysomelidae	<i>Chaetocnema hortensis</i>	Palaeartic	On various species of Poaceae	Recorded as a minor pest of wheat and barley in Europe
	<i>Longitarsus lewisii</i>	Palaeartic	On <i>Plantago</i> spp., especially <i>P. major</i>	–
	<i>Lythvaria salicariae</i>	Palaeartic	In wetlands on <i>Lysimachia</i> spp.	–
Ciidae	<i>Cis boleti</i>	Palaeartic	On polypore fungi, mainly <i>Trametes</i> spp.	–
	<i>Cis glabratus</i>	Palaeartic	On polypore fungi, main host: <i>Fomitopsis pinicola</i>	–
Clambidae	<i>Calyptomerus dubius</i>	Palaeartic	In decaying plant material	–
	<i>Clambus simsoni</i>	Australian	In decaying plant material	–

Family	Species	Original distribution	Habitat	Possible pest status
Coccinellidae	<i>Chilocorus renipustulatus</i>	Palaeartic	Deciduous forests, feeds on scale insects	–
	<i>Scymnus rubromaculatus</i>	Palaeartic	Dry, warm habitats, mainly found on Brassicaceae, feeds on aphids	–
Corylophidae	<i>Orthoperus corticalis</i>	Palaeartic	Deciduous forests, in fungus-infested dead wood	–
Curculionidae	<i>Notaris scirpi</i>	Palaeartic	On <i>Scirpus</i> and <i>Carex</i> in various wet habitats	–
	<i>Exomias trichopterus</i>	Palaeartic	Eurytopic, polyphagous, often found in orchards and gardens	Potential pest of berry crops. Recorded as a pest of strawberry, raspberry and black chokeberry in Europe.
	<i>Ceutorhynchus inaeffectatus</i>	Palaeartic	On <i>Hesperis matronalis</i> (also <i>H. tristis</i> in Europe)	–
	<i>Ambrosiodmus rubricollis</i>	Palaeartic	Ambrosia feeder, polyphagous	Potential pest of various deciduous and coniferous trees.
Dermestidae	<i>Attagenus smirnovi</i>	Afrotropical	Larval development mainly (exclusively?) indoors in temperate areas	Pest of various organic materials of animal origin
Erotylidae	<i>Cryptophilus obliteratedus</i>	Palaeartic	In decaying plant material	–
	<i>Cryptophilus propinquus</i>	Palaeartic	In decaying plant material	–
Hydrophilidae	<i>Coelostoma orbiculare</i>	Palaeartic	Aquatic, mainly in eutrophic ponds	–
Leiodidae	<i>Leiodes polita</i>	Palaeartic	Eurytopic, in forests, heaths, gardens, etc., larval development probably in subterranean fungi	–
Mycetophagidae	<i>Litargus connexus</i>	Palaeartic	In deciduous and mixed forests in fungus-infested dead wood	–
Nitidulidae	<i>Epuraea unicolor</i>	Palaeartic	Decaying and fermenting fruit, tree sap, fungal fruiting bodies, etc.	–
Phalacridae	<i>Olibrus liquidus</i>	Palaeartic	Prefers dry, warm habitats, larval development in flowers of Asteraceae, exact host plants preferences not known	–
Staphylinidae	<i>Amischa decipiens</i>	Palaeartic	Eurytopic, in various, usually moist habitats	–
	<i>Abeta vaga</i>	Palaeartic	Eurytopic, in decaying organic material, often in bird nests	–
	<i>Myllaena infuscata</i>	Palaeartic	At margins of standing and running water	–
	<i>Bledius gallicus</i>	Palaeartic	Moist soil at river banks, edges of fields, etc.	–
	<i>Carpelimus elongatus</i>	Palaeartic	Various moist habitats	–
	<i>Latrobium geminum</i>	Palaeartic	Moist, open habitats	–
	<i>Latrobium lineatocolle</i>	Palaeartic	Riparian habitats and wet meadows	–
	<i>Medon apicalis</i>	Palaeartic	Exact breeding habitat unknown, collected from various habitats	–
	Staphylinidae	<i>Medon ripicola</i>	Palaeartic	Unknown, possibly deep litter or mammal burrows
<i>Pseudomedon obscurellus</i>		Palaeartic	Wetlands, in decaying organic matter	–
<i>Biblepsectus minutissimus</i>		Palaeartic	Collected in flood debris and along river banks in Europe, exact microhabitat unknown	–
<i>Scydmaenus rufus</i>		Palaeartic	Eurytopic, found in forest edges, parks, gardens, floodplains and fields	–
<i>Scydmorephes minutus</i>		Palaeartic	Associated with ants	–
<i>Stenichmus collaris</i>		Palaeartic	Various moist habitats	–
<i>Stenichmus scutellaris</i>		Palaeartic	Forests, forest edges, gardens	–
<i>Mycetoporus reichei</i>		Palaeartic	Eurytopic, in various terrestrial habitats	–
<i>Tachyporus atriceps</i>		Palaeartic	Eurytopic, mainly in disturbed habitats in Canada	–
Throscidae	<i>Trixagus carinifrons</i>	Palaeartic	Prefers dry, warm habitats, larval development probably in soil on fungi (possibly mycorrhizal fungi)	–
	<i>Trixagus meybohmi</i>	Palaeartic	Larval development probably in soil on fungi (possibly mycorrhizal fungi)	–

Gyrinidae

Dineutus emarginatus (Say, 1825)

BOLD:ABX0942

Distribution. Native to the Nearctic region. Widespread in the eastern United States (Gustafson and Miller 2015).

Canadian records. ONTARIO: Charleston Lake Provincial Park, 22-Jun-2015 (2 exx, CBG); Charleston Lake Provincial Park, 25-Jun-2015 (2 exx, CBG).

Diagnostic information. See Gustafson and Miller (2015).

Bionomic notes. Recorded from a variety of lotic and lentic freshwater habitats (Gustafson and Miller 2015). The Canadian adult specimens were hand-collected at a boat launch site on the shore of Charleston Lake, and two larvae matched to the adults by barcode sequences were collected with a dip net at a different site in the same lake.

Carabidae

Harpalinae

Harpalini

Anisodactylus caenus (Say, 1823)

BOLD:AAZ2908

Distribution. Native to the Nearctic region. Widespread in the United States, recorded from all states bordering southern Ontario (Bousquet 2012).

Canadian records. ONTARIO: Point Pelee National Park, 08-Jun-2000 (1 ex, CNC).

Diagnostic information. See Lindroth (1968) or Bousquet (2010).

Bionomic notes. This species occurs in deciduous forests on moist soil (Larochelle and Larivière 2003). The Canadian specimen was caught with a UV light trap on a forest trail in Point Pelee National Park.

Comments. As only a single specimen was captured, it is uncertain whether this species is truly established in Canada.

Hydrophilidae

Sphaeridiinae

Coelostomatini

Coelostoma orbiculare (Fabricius, 1775)

BOLD:AAF3894

Figure 1



Figure 1. *Coelostoma orbiculare* (Fabricius), habitus, L. Borowiec. Scale bar: 1.0 mm.

Distribution. Native to the Palearctic region. Widespread and common in Europe, distributed across Eurasia to the Russian Far East and Japan (Hansen 1987, 2004). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Cambridge, 01-Jun-2015 (1 ex, CBG); Hartington, 18-Apr-2017 (1 ex, CBG).

Diagnostic information (based on Hansen 1987). Body length 4.0–4.8 mm. Habitus short and wide, convex, as in Fig. 1. Black, with the pronotal margins sometimes narrowly red-brown. Antennae with nine antennomeres and a loosely built club with three antennomeres. Base of antennae concealed in dorsal view by the expanded lateral margin of the head. Eyes emarginate. Elytra with sharply impressed sutural striae reaching from apex at least to middle. Tarsomere 1 of meso- and metatarsi longer than tarsomere 2. Abdominal ventrite 1 without medial carina.

Bionomic notes. This species is found in stagnant fresh water. It prefers eutrophic ponds with dense vegetation, and mainly occurs in shallow water at the edges (Hansen 1987). One of the two Canadian specimens was collected as a larva in a leaf litter sample from a wetland, the other (an adult) was sifted from leaf litter close to a lake shore.

Comments. This is the first record of the genus *Coelostoma* Brullé, 1835 in the Nearctic region. *Coelostoma orbiculare* leads to couplet 28 in Van Tassell's (2001) key to North American genera of Hydrophilidae together with the genera *Dactylosternum* Wollaston, 1854 and *Phaenonotum* Sharp, 1882. It can be distinguished from *Dactylosternum* by the absence of a longitudinal carina on the first abdominal ventrite (present in *Dactylosternum*), and from *Phaenonotum* by the presence of distinct sutural striae on the elytra (absent in *Phaenonotum*).

Leiodidae

Leiodinae

Leiodini

Leiodes polita (Marsham, 1802)

BOLD:AAO1258

Figure 2

Distribution. Native to the Palaearctic region. Widespread in Europe, also recorded from North Africa and Caucasus (Daffner 1983; Perreau 2004). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Puslinch Township, 15-Aug-2010 to 22-Aug-2010 (1 ex, CBG); Guelph, 18-Aug-2010 (1 ex, CBG).

Diagnostic information (based on Daffner 1983). Body length 2.2–4 mm. Habitus as in Fig. 2A. Red-brown or yellow-brown, head, pronotum and the sutural and lateral margins of elytra sometimes darkened. Antennae long, with a strongly transverse, darkened club, last antennomere narrower than antennomere 10. Head normally with four punctures in transverse series. Basal margin of pronotum sinuate laterally. Mesoventrite with a long, low and evenly curved medial carina, not reaching the transverse carina and without excavation anteriorly. Metaventrite approximately as long as abdominal ventrites 1 and 2 combined. Elytra with regular, strongly and densely punctate striae, interstitial punctures sparse. Elytra not transversely strigose or strongly microsculptured. Elytral stria 9 separated from side margin at basal third, forming a subhumeral row of punctures. Protibiae only moderately widened towards apex. Metafemora in both sexes with an apical projection at both inner and outer margins, projections stronger in males (Fig. 2B). Male metatibiae bent inward starting from the basal third. Male genitalia as in Fig. 2C, D.




Figure 2. *Leiodes polita* (Marsham) **A** habitus L. Borowiec **B** male left metafemur, dorsal view **C** aedeagus, ventral view **D** aedeagus, lateral view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B**), 0.2 mm (**C, D**).

Bionomic notes. This eurytopic species is found in forests, forest edges, heaths, gardens etc. in Europe (Koch 1989b). The Canadian specimens were collected with Malaise traps in suburban residential areas in southern Ontario.

Comments. *Leiodes polita* leads to *L. quebecensis* Baranowski, 1993 in the key to North American species of *Leiodes* (Baranowski 1993). It can be distinguished by the sinuate basal margin of the pronotum (straight in *L. quebecensis* and related species), differently formed projections of the metafemora, and the male genitalia.

Staphylinidae
Pselaphinae
Trichonychini

***Biblopectus minutissimus* (Aubé, 1833)**

BOLD:AAO0615

Figure 3

Distribution. Native to the West Palaearctic region, widespread in Europe (Schülke and Smetana 2015). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Peterborough, 24-May-2015 to 30-May-2015 (1 ex, CBG); Markham, 24-Jun-2017 to 25-Jun-2017 (1 ex, CBG).

Diagnostic information. Body length 0.9–1.1 mm. Habitus as in Fig. 3A. Female apical tergite with distinctive projection (Fig. 3A). Aedeagus as in Fig. 3B (Löbl 1960).

Bionomic notes. Pearce (1957) writes that this species can be collected from the base of grasses and under stones along sandy river banks and in flood debris in Britain. Unlike other European (Pearce 1957) or North American (Owens and Carlton 2017) species of the genus, it does not occur in deep leaf litter or damp moss. Both Pearce (1957) and Besuchet (1955) state that the species is infrequently collected but this may be due to the extremely small size and unknown microhabitat requirements. One of the Canadian specimens was collected with a Malaise trap on farmland, the other was extracted from soil and leaf litter from a mixed habitat of farmland and forest.

Comments. Only female specimens were available from the Nearctic, but they share identical barcode haplotypes with a specimen of *Biblopectus minutissimus* sampled from Germany. They were also morphologically consistent with the diagnostic characters listed above. In the Palaearctic fauna, females of this species can be recognized by a combination of small size, pale body, temples clearly longer than eyes, and apical tergite produced into a long spine (Besuchet 1955). In the Nearctic region, males are needed for an accurate morphological identification (see Chandler 1990; Owens and Carlton 2017) as many undescribed species are still expected.

Tachyporinae
Mycetoporini


***Mycetoporus reichei* Pandellé, 1869**

BOLD:ABW4933

Figure 4

= *Mycetoporus triangulatus* Campbell, 1991, syn. nov.

Distribution. Native to the West Palaearctic region and broadly distributed (Schülke and Smetana 2015). Adventive in the Nearctic region (Massachusetts, New Hamp-



Figures 3, 4. *Biblopectus minutissimus* (Aubé) **3A** habitus **3B** aedeagus, ventral view, re-drawn from Löbl (1960) **4** *Mycetoporus reichei* Pandellé **4A** habitus **4B** aedeagus, ventral and lateral view, M. Kocian. Scale bars: 0.25 mm (**3A**), 0.1 mm (**3B**), 1.0 mm (**4A**), 0.2 mm (**4B**).

shire, and Wisconsin, United States, and Ontario, Quebec, and New Brunswick, Canada) (Campbell 1991, as *M. triangulatus*).

Canadian records (DNA barcoded specimens). ONTARIO: Orangeville, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Owen Sound, 21-Aug-2014 to 04-Sep-2014 (1 ex, CBG).

Additional Canadian records. See Campbell (1991) and Brunke et al. (2014) for details of earlier records from Canada and United States (as *M. triangulatus*).

Diagnostic information. Body length: 3.1–4.3 mm. Habitus as in Fig. 4A. Ocular puncture of head located at inner edge of eye. Discal pronotal punctures absent. Elytral disc with only one row of punctures, and elytral microsculpture only distinct in apical half. Aedeagus as in Fig. 4B.

Bionomic notes. In the Nearctic, Campbell (1991) reported this species from a vole nest, spruce litter, car net, flood debris, and from a deciduous forest. Brunke et al. (2014) collected this species (as *M. triangulatus*) from soybean fields and their adjacent hedgerows in Ontario, Canada. The barcoded Canadian specimens were collected with Malaise traps, one in a suburban residential area and the other in grassland habitat.

Comments. Campbell (1991) described *Mycetoporus triangulatus* and stated that it is “almost certainly introduced” but was unable to match it to Palearctic species available for study. The Palearctic species of *Mycetoporus* were only recently revised to

include the complex sclerites of the internal sac (e.g., Schülke 2012b) and the Nearctic fauna had not been reviewed since. The Finnish specimens in the BIN are identified as *M. clavicornis* (Stephens, 1832), a close relative of *M. reichei*. These specimens may be misidentified (they were identified by MP before the presence of *M. reichei* in Finland was detected) and need to be re-examined.

Tachyporinae

Tachyporini

Tachyporus atriceps Stephens, 1832

BOLD:ABX2484, BOLD:ACL8080


Figure 5

Distribution. Native to the Palaearctic region, where it is widespread (Schülke 2012b). Adventive in the Nearctic region (British Columbia, Ontario, Quebec, Nova Scotia, New Brunswick, and Prince Edward Island, Canada).

Canadian records (DNA barcoded specimens). BRITISH COLUMBIA: Burnaby, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG). ONTARIO: Ancaster, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG); Ausable-Bayfield Conservation Authority, 30-Jun-2015 (1 ex, CBG); Cambridge, 22-Sep-2014 to 03-Oct-2014 (2 exx, CBG); Cambridge, 24-Apr-2015 to 01-May-2015 (49 exx, CBG); Carillon Park, 06-May-1982 (2 ex, CNC); Courtice, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG); Guelph, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Guelph, 23-Sep-2013 to 04-Oct-2013 (2 exx, CBG); Guelph, 26-Sep-2014 to 29-Sep-2014 (1 ex, CBG); Hartington, 04-Oct-2017 (1 ex, CBG); Mississauga, 24-May-2016 to 26-May-2016 (1 ex, CBG); Point Pelee National Park, 06-Jul-2015 (1 ex, CBG); Puslinch Township, 19-Sep-2010 to 27-Sep-2010 (2 exx, CBG); Puslinch Township, 24-Oct-2010 to 31-Oct-2010 (1 ex, CBG); Red Rock, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG); Rondeau Provincial Park, 09-Jul-2015 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (13 exx, CBG); Rouge National Urban Park, 15-Sep-2013 (1 ex, CBG); Stayner, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG). QUEBEC: Forillon National Park, 16-Sep-2013 to 23-Sep-2013 (2 exx, CBG). NEW BRUNSWICK: Florenceville-Bristol, 22-Sep-2014 to 03-Oct-2014 (3 exx, CBG); Fredericton, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG). NOVA SCOTIA: Cape Breton Highlands National Park, 13-Sep-2013 to 20-Sep-2013 (1 ex, CBG); Hubbards, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG). PRINCE EDWARD ISLAND: Prince Edward Island National Park, 11-Sep-2013 to 18-Sep-2013 (1 ex, CBG).

Additional Canadian records. ONTARIO: Carillon Park, 06-May-1982 (2 exx, CNC); Renfrew County, 4 km SE Cobden, 15-Sep-1980 (1 ex, CNC); Wentworth County, Stoney Creek, 03-Mar-1973 (1 ex, CNC). QUEBEC: Montreal, 07-Sep-1984 (1 ex, CNC). NOVA SCOTIA: Halifax, 1988 (1 ex, CNC).

Diagnostic information. Body length 2.4–3.6 mm. Habitus as in Fig. 5A. Head black, strongly contrasting with pronotum, elytra with either medial and ovoid lateral markings across disc or disc nearly entirely darkened (Fig. 5B). Elytral chaetotaxy as



Figures 5, 6. *5* *Tachyporus atriceps* Stephens **5A** habitus **5B** elytral chaetotaxy (re-drawn from Assing and Schülke 2012) and pattern variability **5C** aedeagus, ventral view, M. Schülke, modified. Abbreviations: OD outer discal row, MD middle discal row, ID inner discal row, S sutural row **6** *Amischa decipiens* (Sharp) **6A** habitus **6B** spermatheca, re-drawn from Muona (1990). Scale bars: 1.0 mm (**5A**; **6A**), 0.5 mm (**5B**, **C**), 0.1 mm (**6B**).

in Fig. 5B, with sutural row of punctures and three rows of discal punctures, with two punctures in inner discal row. Aedeagus as in Fig. 5C.

Bionomic notes. This species occurs in a variety of moist to very dry microhabitats (Schülke 2012b). In the Nearctic, this species occurs in disturbed habitats. It was collected by Brunke et al. (2014) in soybean fields and adjacent hedgerows in Ontario (misidentified as *T. canadensis* Campbell, 1979, to which it is similar). The barcoded specimens were collected in suburban residential areas and protected land adjacent to cities, mainly with Malaise or pitfall traps. Some of the barcoded specimens are larvae extracted from soil or leaf litter.

Comments. *Tachyporus atriceps* has the same elytral chaetotaxy as *T. borealis* Campbell, 1979, *T. nimbicola* Campbell, 1979, and *T. canadensis* Campbell, 1979 but can be separated from the first two by the elytra with discal markings. *Tachyporus canadensis* has a dark red-brown head, bright yellow pronotum, and either a pair of narrow linear lateral dark markings (and medial darkening) or entirely immaculate elytra, while *T. atriceps* has a deep black head, slightly darkened (dingy yellow-orange) pronotum and lateral elytral markings that are ovoid or entirely fused with the medial marking to form a broad darkened area over much of the elytra. The internal sac sclerite of *T. atriceps* is similarly shaped to *T. nimbicola* and *T. borealis* (cane-shaped, Fig. 5C) while *T. canadensis* possesses a characteristic arc-shaped sclerite that is not hooked. *Tachyporus atriceps* appears to be common in at least southern Canada and has been present in the Nearctic since at least 1973. This species was probably recently introduced, or has only recently become widespread and common, as it was not detected by Campbell (1979). *Tachyporus atriceps* is split into two closely clustered BINs which show no differences in morphology, including male genitalia.

Aleocharinae

Athetini

Amischa decipiens (Sharp, 1869)

BOLD:ABA6362

Figure 6

Distribution. Native to the Palaearctic region, occurring broadly in Europe and also reported from the Canary Islands, Tunisia, Turkey, and Mongolia (Schülke and Smetana 2015). Adventive in the Nearctic region (British Columbia and Ontario, Canada).

Canadian records. BRITISH COLUMBIA: Abbotsford, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Burnaby, 21-Sep-2015 to 02-Oct-2015 (2 exx, CBG); Port Coquitlam, 20-Apr-2015 to 08-May-2015 (1 ex, CBG); Surrey, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG). ONTARIO: Aylmer, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG); Brantford, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Cambridge, 20-Apr-2015 to 08-May-2015 (1 ex, CBG); Chesley, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Ethel, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Georgian Bay Islands National Park, 28-

Apr-2013 to 03-May-2013 (1 ex, CBG); Hagersville, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Hagersville, 23-Sep-2013 to 04-Oct-2013 (1 ex, CBG); Little Britain, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG); London, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG); London, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Manitowaning, 21-Sep-2015 to 02-Oct-2015 (5 exx, CBG); Milverton, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Napanee, 22-Sep-2014 to 03-Oct-2014 (2 exx, CBG); Perth East, 23-Sep-2013 to 04-Oct-2013 (1 ex, CBG); St. Thomas, 22-Sep-2014 to 03-Oct-2014 (10 exx, CBG); Stayner, 21-Sep-2015 to 02-Oct-2015 (2 exx, CBG); Teeswater, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Teeswater, 23-Sep-2013 to 04-Oct-2013 (1 ex, CBG); Walkerton, 22-Apr-2013 to 03-May-2013 (2 exx, CBG); Walkerton, 22-Sep-2014 to 03-Oct-2014 (2 exx, CBG); Whitby, 23-Sep-2013 to 04-Oct-2013 (1 ex, CBG).

Diagnostic information. Body length: 2.0–2.2 mm. Habitus as in Fig. 6A. Tergite VII in both sexes without a distinct notch. Spermatheca as in Fig. 6B.

Bionomic notes. This eurytopic species is usually found in moist microhabitats such as leaf litter and moldy hay (Koch 1989a). Good (1995) reported this species from agricultural fields and grasslands in Ireland. Most of the barcoded Canadian specimens were collected with Malaise traps in suburban residential areas.

Comments. One of the most distinctive species of this difficult genus, *A. decipiens* can be recognized by tergite VII lacking a notch in both sexes and by the distinctive spermatheca that bears an elongate capsule (Fig. 6B) (Muona 1990). In at least some parts of its European range (e.g., Ireland), the species is considered to be parthenogenetic (Williams 1969; Good 1995). All examined voucher specimens from the Nearctic were females, suggesting that this species is also parthenogenetic in North America. The genus *Amischa* in North America is unrevised and all Nearctic specimens that cannot be matched to Palearctic *A. analis* (Gravenhorst, 1802) or *A. decipiens* should be treated as unidentifiable pending a comprehensive study. An examination of all North American types was outside the scope of this study and should ideally be accompanied by further DNA sequencing work of both Nearctic and Palearctic *Amischa*.

Atheta vaga (Heer, 1839)


BOLD: AAP6199

Figure 7

Distribution. Native to the Palearctic region, widespread in Europe and reported from Algeria, Tunisia, East and West Siberia, and Mongolia (Schülke and Smetana 2015). Adventive in the Nearctic region (California, United States, and Nova Scotia, Canada).

Canadian records. NOVA SCOTIA: Halifax, 30-May-2013 to 06-Jul-2013 (2 exx, CBG).

Diagnostic information. Body length 2.5–2.8 mm. Habitus as in Fig. 7A. Aedeagus as in Fig. 7B. Spermatheca as in *Atheta fanaticus* Casey (Fig 7D).



Figures 7, 8. **7A** *Atheta vaga* (Heer), habitus **7B** *Atheta vaga*, aedeagus, lateral view **7C** *Atheta fanatica* Casey, aedeagus, lateral view **7D** *Atheta fanatica*, spermatheca **8** *Myllaena infuscata* Kraatz **8A** habitus **8B** aedeagus, ventral view **8C** aedeagus, lateral view **8D** spermatheca **8B–D** V. Assing. Scale bars: 1.0 mm (**7A**), 0.2 mm (**7B–D**), 0.5 mm (**8A**), 0.1 mm (**8B–D**).

Bionomic notes. Palm (1970) wrote that the species is common at sap runs on trees, on carrion, in fungi, in compost, and in the nests of birds, including ravens and birds of prey. It consistently occurs in a wide variety of bird nests in Europe (Hicks 1959). Its sister species, native Nearctic *A. fanatica* Casey, 1910, apparently lives in the same way (Klimaszewski et al. 2018) and has been collected in artificial owl nest boxes (Majka et al. 2006, Webster et al. 2009). The Canadian specimens were collected with a Malaise trap in a forested part of Point Pleasant Park in Halifax, Nova Scotia.

Comments. First reported from North America by Muona (1984) from California (without specimen data including date). This is the first record from eastern North America and for Canada. Populations in eastern and western North America may represent separate introductions, and dissection and sequencing of further material may reveal a more detailed introduction history. With the exception of its native sister species, *A. fanatica*, *A. vaga* can be easily recognized by the shape of the median lobe in lateral view and spermatheca. It can be distinguished from *A. fanatica* by the less strongly sinuate tubus of the median lobe in lateral view (compare with Fig. 7C). The spermathecae of the two species are identical. The close relationship and separate species status of the two species is confirmed by two well-separated BINs. Although the two examined vouchers from Canada are females, we are confident of their identity based on identical DNA barcode haplotypes shared with European material of *A. vaga*. *Atheta fanatica* forms a separate BIN cluster (BOLD:ACL9881) which shows ca. 10% divergence from *A. vaga*.

Aleocharinae

Myllaenini

Myllaena infuscata Kraatz, 1853

BOLD:AAX2901

Figure 8

Distribution. Native to the western Palearctic region, widely distributed in Europe but rare in the north (Palm 1968) and also reported from India (Kashmir) (Schülke and Smetana 2015), though many of these records need confirmation (Assing 2018). It has previously been reported as occurring in the Nearctic region by Schülke and Smetana (2015) but this is probably based on an erroneous synonymy of North American *M. immunda* Casey with this species in older literature. Klimaszewski (1982) corrected the synonymy, and *M. immunda* is now considered a synonym of *M. arcana* Casey, 1911. The true *M. infuscata* is reported here as adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Rouge National Urban Park, 15-Sep-2013 (1 ex, CBG).

Diagnostic information. Body length 1.2–1.5 mm. Habitus as in Fig. 8A. Aedeagus as in Fig. 8B, C. Spermatheca as in Fig. 8D.

Bionomic notes. *Myllaena infuscata* occurs in both exposed and shaded microhabitats along the margins of still and running water (Reid 1991). The Canadian specimen was extracted from soil and leaf litter collected near the mouth of Rouge River.

Comments. *Myllaena infuscata* is distinctive in the Nearctic fauna for its spermathecal shape, which forms concentric circular coils (Fig. 8D). The median lobe of the aedeagus in lateral view is also distinctive among species in North America (Fig. 8B, C). Within the western Palaearctic fauna, only *M. minuta* has similar genitalia but differs in the shape of the median lobe in lateral view. The single female from Ontario bears the characteristic spermatheca and its corresponding barcode sequence falls within the BIN associated with *M. infuscata*, rather than *M. minuta*, its similar sister species. It is unknown whether this is a recent introduction to North America or if its small size has impeded its detection.

Oxytelinae

Blediini

Bledius gallicus (Gravenhorst, 1806)

BOLD:AAH0142

Figure 9

= *Bledius philadelphicus* Fall, 1919, syn. nov.

Distribution. Native to the Palaearctic region, trans-Palaearctic (Schülke 2012a). Adventive in the Nearctic region (Maryland, Massachusetts, New Jersey, New York, and Pennsylvania, United States, and Ontario, Quebec, New Brunswick, and Newfoundland, Canada (Herman 1972, Bousquet et al. 2013, as *B. philadelphicus*).


Canadian records (DNA barcoded specimens). ONTARIO: Georgian Bay Islands National Park, 19-Aug-2013 to 27-Aug-2013 (1 ex, CBG); Grundy Lake Provincial Park, 13-Jul-1995 (1 ex, CNC); Hamilton, 21-Jul-2017 (3 exx, CBG). QUEBEC: Montreal, 19-Aug-1981 (1 ex, CNC).

Additional Canadian records. See Herman (1972) for a list of earlier records from Canada and the United States (as *B. philadelphicus*).

Diagnostic information. Body length: 3.7–4.8 mm. Habitus as in Fig. 9A, B. Male sternite VII as in Fig. 9D. Aedeagus as in Fig. 9C.

Bionomic notes. Palm (1961) (as synonym *B. fracticornis*) states that this species can be found in half-moist sand, gravel, clay or mineral soil mixed with humus, with or without vegetation cover. In Central Europe, this species occurs on sandy to muddy river banks, and also in damp field edges (Schülke 2012a). Three of the CBG specimens were collected at a UV light at a forest edge, one was caught in a Malaise trap in a forested peninsula.

Comments. *Bledius gallicus* can be recognized within Herman's (1972) 'semiferugineus group' using the following combination of characters: last segment of meta-



Figures 9, 10. **9** *Bledius gallicus* (Gravenhorst) **9A** habitus, black elytra, L. Borowiec **9B** habitus, red-brown elytra, L. Borowiec **9C** aedeagus, ventral view, VL = ventral lamella M. Schülke **9D** male sternite VII **10** *Carpelimus elongatulus* (Erichson) **10A** habitus **10B** aedeagus, ventral view, H. Schillhammer. Scale bars: 1.0 mm (**9A, B; 10A**), 0.2 mm (**9C**), 0.2 mm (**9D**), 0.25 mm (**10B**).

tarsus in dorsal view gradually expanded to apex, male sternite VII emarginate, with membranous lobe but emargination not bordered by a pair of spines, pronotum with midlongitudinal groove. The species will key easily to *B. philadelphicus* Fall, 1919 in Herman's (1972) key and we here consider these two species synonyms. Specimens in

the CNC identified as *B. philadelphicus* by Lee Herman and included in his revision of the ‘*semiferrugineus* group’ (Herman 1972) were dissected and revealed to be *B. gallicus*. The description of *Bledius philadelphicus* in Herman (1972) corresponds to that of *B. gallicus* in Schülke (2012a), including the characteristic male sternite VII (though the membranous part is slightly deeper in both Nearctic and Palearctic populations than indicated by the illustration). *Bledius gallicus* is closely related to the Palearctic *B. femoralis* (Gyllenhal, 1827) (Schülke 2012a). The two species have extremely similar aedeagi, differing only in the apex of the ventral lamella (Fig. 9C) (acute in *B. femoralis* and broadly truncate in *B. gallicus*). These two species are more easily separated by the shape of male sternite VII (Schülke 2012a).

Based on the specimens available at the CNC and reported by Herman (1972), *B. gallicus* has been in North America for quite a long time, since at least as early as 1910, when Fall (1910) first described *B. philadelphicus* as *B. dissimilis* (not Erichson 1840, preoccupied name replaced by Fall (1919)). The earliest Canadian specimens are from the 1920s.

Oxytelinae

Oxytelini

Carpelimus elongatulus (Erichson, 1839)

BOLD:AAW4293

Figure 10

Distribution. Native to the Palearctic region, distributed from western Europe to the Baikal region of Russia (Schülke 2012a). Adventive in the Nearctic region (Ontario, Quebec, and New Brunswick, Canada).

Canadian records (DNA barcoded specimens). ONTARIO: Ausable-Bayfield Conservation Authority, 30-Jun-2015 (1 ex, CBG); Indian Point Provincial Park, 28-Jul-2015 (1 ex, CBG); Puslinch Township, 09-May-2010 (1 ex, CBG).

Additional Canadian records. ONTARIO: Guelph, 10-Apr-2009, (2 exx, DEBU); Minesing Swamp, 26-Jan-2008 (1 ex, DEBU). QUEBEC: Dorval, 10-Oct-1975, (1 ex, FMNH); Sainte-Foy, 27-May-1976, (1 ex, FMNH). NEW BRUNSWICK: Charlotte County, 05-Jun-2008 (1 ex, RWC); Jackson falls, 22-May-2010, (1 ex, RWC); Musquash, 07-May-2006 (1 ex, RWC).

Diagnostic information. Body length: 2.0–2.6 mm. Habitus as in Fig. 10A. Aedeagus as in Fig. 10B.

Bionomic notes. This species occurs on banks of waterways, wet meadows, agricultural fields and in damp leaf litter (Schülke 2012a). The Canadian specimens were collected with Malaise traps in forests and extracted from leaf litter from a wetland and a river bank.

Comments. As the Nearctic *Carpelimus* have not been revised in modern times, it is currently necessary to dissect males to match with published illustrations of the aedeagus (see Webster et al. 2016). Although only female voucher specimens from the Nearctic were available for study, they easily key to *C. elongatulus* in Schülke

(2012a) and two of the barcoded Canadian specimens share identical haplotypes with European specimens of *C. elongatulus*. Similar but much smaller Palaearctic species such as *C. subtilis* are represented in BOLD and form separate BINs deeply divergent from *C. elongatulus*.

Scydmaeninae Cyrtoscydmini

Stenichnus collaris (Müller & Kunze, 1822)

BOLD:ABA1519

Figure 11

Distribution. Native to the western Palaearctic region, widely distributed in Europe (Schülke and Smetana 2015). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Peterborough, 31-May-2015 to 06-Jun-2015 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (2 exx, CBG).

Diagnostic information. Body size: 1.55–1.70 mm. Habitus as in Fig. 11A. Aedeagus as in Fig. 11B.

Bionomic notes. Koch (1989a) reports that this species is found in moist forests, forest edges and gardens among leaves and fungi, in dead wood, and at sap flows. One of the Canadian specimens was collected with a Malaise trap on farmland, the other two with pitfall traps in grassland and a river bank.

Comments. As the Nearctic *Stenichnus* fauna remains unrevised, it is only possible to associate Nearctic specimens with Palaearctic species through dissected males or barcodes. The Canadian specimens share identical barcode haplotypes with European material, and the identification was verified by examination of the male genitalia.


Stenichnus scutellaris (Müller & Kunze, 1822)

BOLD:AAN9916

Figure 12

Distribution. Native to the western Palaearctic region, widespread in Europe (Schülke and Smetana 2015). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Cambridge, 07-May-2015 to 14-May-2015 (5 exx, CBG); Cambridge, 15-Jul-2017 (15 exx, CBG); Cambridge, 25-May-2015 to 01-Jun-2015 (1 ex, CBG); Cambridge, 25-May-2015 to 31-May-2015 (4 exx, CBG); Cambridge, 29-Apr-2015 to 07-May-2015 (1 ex, CBG); Guelph, 13-May-2017 (1 ex, CBG); Guelph, 22-Apr-2013 to 03-May-2013 (2 exx, CBG); Guelph, 22-Apr-2017 (1 ex, CBG); Kitchener, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Mississauga, 15-Sep-2015 to 17-Sep-2015 (3 exx, CBG); Owen Sound, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Pickering, 24-Jun-2017 to 25-Jun-2017 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (7 exx, CBG); Rouge National Urban Park,



Figures 11, 12. 11 *Stenichnus collaris* (Müller & Kunze) 11A habitus, L. Borowiec 11B aedeagus, dorsal view 12 *Stenichnus scutellaris* (Müller & Kunze) 12A habitus, L. Borowiec 12B aedeagus, dorsal view. Scale bars: 0.5 mm (11A; 12A), 0.1 mm (11B; 12B).

11-Jun-2013 to 18-Jun-2013 (1 ex, CBG); Rouge National Urban Park, 15-Sep-2013 (1 ex, CBG); Rouge National Urban Park, 24-Jun-2017 (1 ex, CBG); Rouge National Urban Park, 29-Apr-2013 to 03-May-2013 (1 ex, CBG); Warsaw, 05-May-2014 to 23-May-2013 (1 ex, CBG); Whitby, 22-Apr-2013 to 03-May-2013 (1 ex, CBG).

Diagnostic information. Body length: 1.4–1.5 mm. Habitus as in Fig. 12A. Male profemur widened apicad to form an abrupt 90° angle in dorsal view. Aedeagus as in Fig. 12B.

Bionomic notes. This species lives in leaf litter and dead wood (Koch 1989a). It is mostly collected in forests or at forest edges, occasionally in wetlands and grasslands

(Koch 1989a). Most Canadian specimens were collected using Malaise traps, pitfall traps, or by sifting leaf litter. Most specimens were collected in disturbed forest fragments but some were from grasslands and wetlands.

Comments. As the Nearctic *Stenichmus* fauna remains unrevised, it is only possible to associate Nearctic specimens with Palaearctic species through dissected males or barcodes. The Canadian specimens share identical barcode haplotypes with European material, and the identification was verified by examination of the male genitalia. The modified male profemur of *S. scutellaris* is unique among the Central European fauna (Franz and Besuchet 1971). Without a revision of the Nearctic fauna, it is not possible to know whether other North American species also possess this character.

Scydmaeninae

Scydmaenini

Scydmaenus rufus Müller & Kunze, 1822

BOLD:AAO4392

Figure 13

Distribution. Native to the western Palaearctic region, widespread in Europe and also reported from Algeria, Tunisia, and Lebanon (Schülke and Smetana 2015). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Guelph, 17-Sep-2017 (6 exx, CBG).

Diagnostic information. Body length: 1.4 mm. Habitus as in Fig. 13A. Aedeagus as in Fig. 13B.

Bionomic notes. This eurytopic species occurs along forest edges and in parks, gardens, floodplains, and fields (Koch 1989a). The Canadian specimens were sifted from a compost heap.

Comments. As the Nearctic *Scydmaenus* fauna remains unrevised, it is only possible to associate Nearctic specimens with Palaearctic species through dissected males or barcodes. Three of the Canadian specimens share identical barcode haplotypes with European material, and the identification was verified by examination of the male genitalia.

Scydmaeninae


Glandulariini

Scydmorephes minutus (Chaudoir, 1845)

BOLD:AAO3241

Figure 14

Distribution. Native to the Palaearctic region, widespread in Europe and also reported from the Russian Far East (Schülke and Smetana 2015). Adventive in the Nearctic region (Ontario, Canada).



Figures 13, 14. **13** *Scydmaenus rufus* Müller & Kunze **13A** habitus, L. Borowiec **13B** aedeagus, ventral view **14** *Scydmoraphes minutus* (Chaudoir) **14A** habitus **14B** aedeagus, ventral view. Scale bars: 0.5 mm (**13A**; **14A**), 0.1 mm (**13B**; **14B**).

Canadian records. ONTARIO:Georgian Bay Islands National Park, 30-Jul-2013 to 06-Aug-2013 (1 ex, CBG); Peterborough, 30-Jul-2015 (1 ex, CBG).

Diagnostic information. Body length: 1.15–1.30 mm. Habitus as in Fig. 14A. Head without supraantennal notches or frontal impression, submentum with lateral sutures broadly separated. Pronotum setose, laterally margined, with a transverse groove at the base. Aedeagus as in Fig. 14B.

Bionomic notes. This species is associated with ants, especially species of the *Formica rufa* Linnaeus, 1761 group, and *Lasius fuliginosus* (Latreille, 1798) and *L. brunneus* (Latreille, 1798) in Europe (Franz and Besuchet 1971). Koch (1989a) reports *S. minutus* with *Lasius* ants in hollow trees, under loose bark and in fallen logs. The Canadian specimens were collected with Malaise traps, one in a forested peninsula and the other on farmland.

Comments. The genus *Scydmoraphes* Reitter, 1891 is here reported for the first time from North America. It was distinguished recently from the similar Nearctic genus *Parascydmus* Casey, 1897 (Jałoszynski 2019), and it does not appear to be an obvious synonym of the other similar Nearctic genus *Brachycepsis* Brendel, 1889. A detailed systematic study of the Nearctic glandulariine genera is warranted. The genus *Scydmoraphes* (with a single species in the Nearctic region) may be recognized within the Nearctic fauna of Glandulariini by the unique combination of a transverse groove on the base of the pronotum, which is margined laterally (Fig. 14A), submentum with lateral sutures broadly separated, and head dorsally lacking frontal impression and supraantennal notches (Jałoszynski 2019). In habitus, *Scydmoraphes* is similar to *Brachycepsis* and *Parascydmus* but can be easily recognized by the transverse pronotal groove.

The following couplets from O’Keefe (2001) were modified to include *Scydmoraphes*:

- 19a (18) Base of pronotum with transverse groove (Fig. 14A)
Scydmoraphes minutus
- Base of pronotum with only impressed basal foveae **19b**
- 19b (19a) Pronotum with 4 basal foveae (fig. 42.20); scutellum large; often light to dark brown in color ***Brachycepsis***
- Pronotum with 6 basal foveae (fig. 43.20); scutellum minute; often black in color ***Parascydmus***

Paederinae

Paederini

***Lathrobium geminum* Kraatz, 1857**

BOLD:AAX0181

Figure 15


Distribution. Native to the Palaearctic region, distributed from Europe to the Far East of Russia (Schülke and Smetana 2015). Adventive in the Nearctic region (British Columbia, Canada).

Canadian records. BRITISH COLUMBIA: Gulf Islands National Park Reserve, 17-Jun-2014 to 22-Jun-2014 (2 exx, CBG).

Diagnostic information. Body length: 8.0–9.0 mm. Habitus as in Fig. 15A. Female sternite VIII elongate and truncate apically, as in Fig. 15D. Aedeagus distinctive in lateral view, as in Fig. 15B, C.

Bionomic notes. In Central Europe, this is a common species in unforested, humid microhabitats such as wetlands, shorelines, agricultural fields, gardens, and heath (Assing 2012). The Canadian specimens were collected in a wetland adjacent to a lake, one with pitfall traps and the other by Berlese funnel extraction.

Comments. The voucher specimens from North America are, unfortunately, females but share identical barcode haplotypes with Palaearctic specimens of *L. geminum*



Figures 15, 16. **15** *Lathrobium geminum* Kraatz **15A** habitus, L. Borowiec **15B** aedeagus, lateral view V. Assing **15C** aedeagus, ventral view V. Assing **15D** female sternite VIII, modified from Assing and Schülke (2012) **16** *Lathrobium lineatocolle* Scriba **16A** aedeagus, lateral view, V. Assing **16B** female sternite VIII, V. Assing. Scale bars: 2.0 mm (**15A**), 0.5 mm (**15B–D**), 0.5 mm (**16A, B**).

from Germany and Finland. North American vouchers key to *L. geminum* in Assing (2012) and female sternite VIII is consistent with the shape described for this species. As the Nearctic fauna of *Lathrobium* is unrevised, comparisons with North American species are not yet possible.

Lathrobium lineatocolle Scriba, 1859

BOLD:ACL0909

Figure 16

Distribution. Native to the Palearctic region, widespread in Europe and reported from China, Iran, Turkey, and the Russian Far East (Schülke and Smetana 2015). Adventive in Canada (Ontario).

Canadian records. ONTARIO: Rouge National Urban Park, 29-Apr-2013 to 03-May-2013 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (1 ex, CBG).

Diagnostic information. Female sternite VIII elongate to narrow, scarcely emarginate apex, as in Fig. 16B. Aedeagus distinctive in lateral view, as in Fig. 16A.

Bionomic notes. In Central Europe, this species occurs mostly in riparian habitats and in wet meadows (Assing 2012). One of the Canadian specimens was collected with a Malaise trap in a forest patch; the other was caught in a riverside pitfall trap.

Comments. As the Nearctic fauna of *Lathrobium* is unrevised, comparisons with North American species are not yet possible.

Medon apicalis (Kraatz, 1857)

BOLD:ACC4101

Figure 17


Distribution. Native to the western Palaearctic region, widespread in Europe and also reported from Algeria, Morocco, Turkey, the Canary Islands, and Madeira (Schülke and Smetana 2015). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Guelph, 30-Jun-2018 (1 ex, CBG).

Diagnostic information. Body length: 3.8–4.6 mm. Habitus as in Fig. 17A. Male sternite VII as in Fig. 17C. Aedeagus as in Fig. 17B.

Bionomic notes. This species has been collected in a variety of habitats in Europe, but the breeding habitat requirements are unknown (Assing 2006). Most specimens have been collected in flight (car nets, flight interception traps) (Assing 2006). Specimens have also been collected from stream edges, haystacks, woodland and at light (Assing 2006). This species is less likely to occur in the nests of small mammals than other species of the genus (Assing 2006). The Canadian specimen was collected at a UV light in a mixed forest.

Comments. A single female voucher from Canada was available for study and, while males would normally be necessary to confirm a positive identification in *Medon*, its barcode sequence is identical to German and Austrian specimens of *M. apicalis*. All similar, widespread Palaearctic species that could be confused with *M. apicalis* (*M. ripicola* (Kraatz, 1854), *M. brunneus* (Erichson, 1839), *M. fuscus* (Mannerheim, 1830)) are represented in BOLD and form distinct BIN clusters. The female voucher was also morphologically compared to representatives of all widespread western Palaearctic *Medon* species and was consistent with the variability of body proportions, punctuation and color of *M. apicalis*. Four species known from the southwestern Palaearctic are closely related to *M. apicalis* and cannot be reliably distinguished by external characters: *M. perniger* Coiffait, 1978 (Italy and extreme southern parts of France and Switzerland); *M. maronitus* (Saulcy, 1864) (Greece to Turkmenistan); *M. sericellus* Fairmaire, 1860 (North Africa) and *M. beydaghensis* Fagel, 1969 (Turkey) (Assing 2004, 2006). None of these species are currently rep-



Figures 17–19. 17 *Medon apicalis* (Kraatz) 17A habitus 17B aedeagus, ventral and lateral view, V. Assing 17C male sternite VII, V. Assing 18 *Medon ripicola* (Kraatz) 18A habitus 18B aedeagus, ventral and lateral view, V. Assing 18C male sternite VII, V. Assing 19 *Pseudomedon obscurellus* (Erichson) 19A habitus 19B aedeagus, lateral view, V. Assing 19C male sternite VIII, V. Assing. Scale bars: 1.0 mm (17A; 18A; 19A), 0.2 mm (17B, C; 18B, C; 19B, C).

resented on BOLD. Although one or more of these species might share a BIN with *M. apicalis*, the Ontario specimen has an identical DNA barcode haplotype to specimens from Germany and Austria where *M. apicalis* is the only known representative of this species group. As the Nearctic fauna of *Medon* is unrevised, comparisons with North American species are not yet possible. Recognizing this species in the Nearctic region requires dissected males or DNA barcoding.

***Medon ripicola* (Kraatz, 1854)**

BOLD:AAO2230

Figure 18

Distribution. Native to the western Palaearctic region, widespread in Europe and also reported from Algeria, Morocco, Turkey, and Madeira (Schülke and Smetana 2015). Adventive in the Nearctic region (Nova Scotia, Canada).

Canadian records. NOVA SCOTIA: Cape Breton Highlands National Park, 10-May-2013 to 21-May-2013 (1 ex, CBG).

Diagnostic information. Body length: 3.7–4.2 mm. Habitus as in Fig. 18A. Male sternite VII as in Fig. 18C. Aedeagus as in Fig. 18B.

Bionomic notes. This species is rarely collected in the Palaearctic, with its breeding microhabitat unknown (probably in deeper litter or mammal burrows). In Central Europe, specimens have been collected mostly in wetlands (floodplains, ponds), in flood debris, mole nests, and deeper deciduous leaf litter (Assing 2012). Palm (1963) wrote that this species was rarely collected in Scandinavia: once in Sweden under pebbles on the seashore and in Denmark under seaweed. Its occasional but typical appearance near water suggests that heavy rains may flood out the breeding microhabitat and deposit the beetles elsewhere (e.g., flood debris). The collection of *M. ripicola* on northern European seashores suggests a potential mechanism for introduction to the Canadian Maritimes through ocean commerce. The Canadian specimen was collected with a Malaise trap in a riverside forest.

Comments. A single female voucher from Canada was available for study and, while males would normally be necessary to confirm a positive identification in *Medon* by morphology, its barcode sequence clustered within the European material of *M. ripicola* with only two nucleotide sites differing from the nearest European specimen. All similar Palaearctic species that could be confused with *M. ripicola* (*M. apicalis* (Kraatz, 1857), *M. brunneus* (Erichson, 1839), *M. fuscus* (Mannerheim, 1830)) are represented in BOLD in separate BIN clusters. The female voucher was also morphologically compared to representatives of all Palaearctic *Medon* species and was consistent with the body proportions, punctuation and color of *M. ripicola*. As the Nearctic fauna of *Medon* is unrevised, useful comparisons with North American species are not yet possible. Recognizing this species in the Nearctic region is reliably accomplished, at present, using dissected males or DNA barcoding.

***Pseudomedon obscurellus* (Erichson, 1840)**

BOLD:AAO1043

Figure 19

Distribution. Native to the western Palaearctic region, widespread in Europe and also reported from Algeria, Morocco, Madeira, Tunisia, Cyprus, and Turkey (Schülke and Smetana 2015). Adventive in the Neotropical region (Chile; Assing 2009) and the Nearctic region (Nova Scotia, Canada).

Canadian records. NOVA SCOTIA: Cape Breton Highlands National Park, 07-Jun-2013 to 24-Jun-2013 (1 ex, CBG).

Diagnostic information. Body length: 3.0–3.4 mm. Habitus as in Fig. 19A. Male sternite VIII as in Fig. 19C. Aedeagus as in Fig. 19B.

Bionomic notes. This species inhabits wetlands and can be collected from rotting organic matter (Assing 2012). The Canadian specimen was collected at the same site and in the same Malaise trap as the *M. ripicola* specimen.

Comments. A single female voucher from Canada was available for study and, while males would normally be necessary to confirm a positive identification in *Pseudomedon*, its barcode sequence is identical to German specimens of *P. obscurellus*. The morphologically similar Palaearctic species *P. obsoletus* forms a separate BIN cluster (BOLD:ABY0636). The female voucher from Canada also was consistent with the typical coloration of *P. obscurellus* given by Assing (2012). As the Nearctic fauna of *Pseudomedon* is unrevised, comparisons with North American species are not yet possible. Recognizing this species in the Nearctic region is reliably accomplished, at present, using dissected males or DNA barcoding.

Due to taxonomic confusion until the 1970s, reports of *Pseudomedon obscurellus* and *P. obsoletus* from regions outside of the Palaearctic need re-confirmation (Assing 2009, Klimaszewski et al. 2013). The record of *P. obsoletus* from British Columbia from Hatch (1957) is doubtful and likely refers to *P. obscurellus* as it was described as being partly dark rufous, a color more typically associated with this species (Assing 2012). To our knowledge, this is the first verified record of any Palaearctic *Pseudomedon* species from the Nearctic.

Scarabaeidae
Melolonthinae
Melolonthini***Phyllophaga implicita* (Horn, 1887)**

BOLD:AAC5184

Distribution. Native to North America. Occurs across most of the Mississippi River drainage basin in the United States (Luginbill and Painter 1953).

Canadian records. ONTARIO: Point Pelee National Park, 05-Jun-2008 (1 ex, CBG).

Diagnostic information (partially based on Luginbill and Painter 1953). Body length 14.5–17.0 mm. Dorsal surface pale to dark brown, moderately shiny (not pruinose or iridescent), glabrous, and without scales. Clypeus emarginate. Antennae with nine antennomeres. See Luginbill and Painter (1953) for images of habitus and genitalia.

Bionomic notes. Adults have been observed on numerous plants including *Tilia* L., *Fagus* L., *Betula* L., *Ulmus* L., *Lonicera* L., *Acer* L., *Platanus* L., *Rosa* L., *Juglans* L., *Salix* L. and cultivated legumes (Luginbill and Painter 1953). The Canadian specimen was collected in a mixed forest close to the shore of Lake Erie using an ultraviolet light.

Comments. It is not surprising to find a range extension of this species into Canada considering the widespread distribution in eastern North America and the apparent broad range of host plants. Since only a single specimen was collected in Canada it is difficult to assess how firmly established this species is. There are hundreds of species of *Phyllophaga* with a similar overall appearance; therefore, it is crucial to use the male or female genitalia for morphological species identifications.

Clambidae

Calyptomerinae

Calyptomerus dubius (Marsham, 1802)

BOLD:AAP9145


Figure 20

Distribution. Native to the Palaearctic region, widespread in Central Europe and around the Mediterranean (Endrödy-Younga 1961, Löbl 2006). Adventive in the Afrotropical region (South Africa), the Australian region (Australia) (Endrödy-Younga 1974) and in the Nearctic region (British Columbia, Canada).

Canadian records. BRITISH COLUMBIA: Abbotsford, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Vancouver, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Victoria, 03-Sep-2014 to 10-Sep-2014 (1 ex, CBG).

Diagnostic information (based on Endrödy-Younga 1961). Body length 1.1–1.6 mm. Habitus as in Fig. 20A. Red-brown, with the edges of pronotum and elytra paler. The antennal groove forms a continuous and even curve with the side of the frons in front of the eyes. Lateral edges of pronotum bluntly angled. Pubescence on the dorsal surface long and sparse. Elytra angled apicolaterally and truncate at the hind margin. Aedeagus as in Fig. 20B.

Bionomic notes. This species is known from decaying plant material. It has been collected from dead, fungus-infested logs of deciduous trees, leaf litter, composts, moldy hay, etc. (Koch 1989b). The Canadian specimens were collected with Malaise traps in residential areas.



Figures 20, 21. **20** *Calyptomerus dubius* (Marsham) **20A** habitus, L. Borowiec **20B** aedeagus, dorsal view
21 *Clambus simsoni* Blackburn **21A** habitus **21B** aedeagus, ventral view. Scale bars: 0.5 mm (**20A**; **21A**),
 0.2 mm (**20B**), 0.1 mm (**21B**).

Comments. *Calyptomerus oblongulus* (Mannerheim, 1853) is the only other representative of this genus known from North America. It is larger than *C. dubius* (body length 1.8–2.0 mm), with a rounded angle between the antennal groove and the lateral margin of frons, rounded lateral edges of pronotum, evenly curved (not truncate) elytral hind margins, shorter and denser pubescence on the dorsal surface, and different male genitalia (Endrödy-Younga 1961).

Clambinae

Clambus simsoni Blackburn, 1902

BOLD:AAO0037

Figure 21

Distribution. Native to the Australian region. Described from Australia, where the species is widespread and common (Endrödy-Younga 1990). Also recorded from New Zealand (Johnson 1997). Adventive in the Afrotropical region (South Africa), the Palaearctic region (British Isles and Central Europe; Endrödy-Younga 1990, Johnson 1997, Meybohm 2004, Löbl 2006), and the Nearctic region (British Columbia, Canada).

Canadian records. BRITISH COLUMBIA: West Vancouver, 20-Apr-2015 to 08-May-2015 (1 ex, CBG).

Diagnostic information (based on Endrödy-Younga 1990 and Johnson 1997). Body length 1.0–1.2 mm. Habitus as in Fig. 21A. Pale red-brown, with head and anterior part of pronotum darkened. Lateral angles of head narrowly rounded, rectangular, a line drawn between the angles level with the posterior margin of eyes. Dorsal surface without microsculpture. Apical part of elytra with large punctures. Pubescence of elytra relatively long and sparse, individual setae only a little longer than distance between seta-bearing punctures. Aedeagus as in Fig. 21B, penis finely serrate laterally in the apical quarter.

Bionomic notes. This species is known from decaying plant material. It has been collected from heaps of cut grass, heaps of shredded bark, and (in New Zealand) from tree fungi (Johnson 1997). The Canadian specimen was collected with a Malaise trap in a suburban residential area.

Comments. Morphologically, *Clambus simsoni* is most reliably identified by its characteristic male genitalia. The Canadian specimen is a male which shares an identical barcode sequence with a specimen sampled from Germany. In Endrödy-Younga's (1981) key to the New World species of *Clambus*, *C. simsoni* leads to *C. spangleri* Endrödy-Younga in couplet 14. *Clambus simsoni* is slightly larger (*C. spangleri* is 0.8–0.9 mm according to Endrödy-Younga), and the pubescence on the dorsal surface is sparser than in *C. spangleri*.

Scirtidae

Scirtinae

Contacyphon fuscescens (Klausnitzer, 1976)

BOLD:ACX3673

Figure 22

Distribution. Native to North America. Described from New York State (Ithaca) (Klausnitzer 1976).

Canadian records. YUKON TERRITORY: Kluane National Park and Reserve, 15-Jul-2014 to 24-Jul-2014 (1 ex, CBG). BRITISH COLUMBIA: Naikoon Provincial Park, 24-Jun-2014 to 03-Jul-2014 (4 exx, CBG); Naikoon Provincial Park, 13-Jul-2014 to 31-Jul-2014 (5 exx, CBG); Naikoon Provincial Park, 08-Aug-2014 to 15-Aug-2014 (4 exx, CBG). ONTARIO: Puslinch Township, 12-Jun-2010 to 19-Jun-2010 (1 ex, CBG); Short Hills Provincial Park, 26-May-2014 to 09-Jun-2014 (4 exx, CBG); Short Hills Provincial Park, 23-Jun-2014 to 07-Jul-2014 (1 ex, CBG).

Diagnostic information (based on Klausnitzer 1976). Body length 2.5–3.0 m. Dark brown, basal margin of the pronotum and elytral suture a little paler. Head with granulate punctation, pronotal punctation fine, elytra a little more coarsely punctate than pronotum. Elytra with longitudinal ribs. Male tergite VIII and IX as in Fig. 22C. Aedeagus as in Fig. 22A, B.

Bionomic notes. The Canadian specimens were collected with Malaise traps in wetlands and close to open water in forests and farmland.

Comments. *Contacyphon fuscescens* belongs to the *C. coarctatus* group of species. It is most reliably identified by the male genitalia. The identification of the Canadian specimens is based on dissected male representatives of the BIN.

Contacyphon kongsbergensis (Munster, 1924)

BOLD:AAJ3307

Figure 23


Distribution. Holarctic. Recorded from across the Palaearctic region, from Western Europe to the Russian Far East (Klausnitzer 2006). In North America, apparently previously known only from Alaska (Kalsin Bay, Kodiak (Nyholm 1972)). This record was overlooked by Bousquet et al. (2013).

Canadian records. BRITISH COLUMBIA: Smithers, 28-Jul-2014 to 05-Aug-2014 (4 exx, CBG). ALBERTA: Jasper National Park, 02-Aug-2010 to 05-Aug-2010 (1 ex, CBG). MANITOBA: Churchill, 05-Aug-2005 (1 ex, CBG); Churchill, 18-Aug-2006 (1 ex, JBWM); Churchill, 21-Jul-2007 (2 exx, CBG); Riding Mountain National Park, 16-Jul-2012 to 23-Jul-2012 (1 ex, CBG).

Diagnostic information (based on Nyholm 1972 and Lohse 1979). Body length 2.7–3.2 mm. Reddish or yellowish brown, head usually darker than pronotum and elytra. Head and pronotum with dense, coarse, granulate punctation. Elytra densely punctate. Male sternite IX and accessory sclerite as in Fig. 23C. Aedeagus as in Fig. 23A, B.

Bionomic notes. This species is known from acidic *Sphagnum* bogs (Nyholm 1972). Most Canadian specimens were collected with Malaise traps; also collected with light traps and by sweep netting.

Comments. *Contacyphon kongsbergensis* is morphologically most reliably identified by its genitalia. The lack of a modern revision of North American *Contacyphon* prevents detailed comparison with related species.



Figures 22–24. **22** *Contacyphon fuscescens* (Klausnitzer) **22A** aedeagus, dorsal plate **22B** aedeagus, ventral plate **22C** male tergite 8 and 9 **23** *Contacyphon kongsbergensis* (Munster) **23A** aedeagus, dorsal plate **23B** aedeagus, ventral plate **23C** male sternite 9 and accessory sclerite **24** *Contacyphon obscurellus* (Klausnitzer) **24A** aedeagus, dorsal plate **24B** aedeagus, ventral plate **24C** male sternite 9 **24D** male tergite 9. Scale bars: 0.2 mm (**22A–C**; **23A–C**), 0.5 mm (**24A–D**).

***Contacyphon obscurellus* (Klausnitzer, 1976)**

BOLD:AAG7260, BOLD:ACP0631, BOLD:ACP0632

Figure 24

Distribution. Native to North America. Described from New York State (Adirondack, Long Lake) (Klausnitzer 1976).

Canadian records. ONTARIO: Georgian Bay Islands National Park, 30-Jul-2013 to 06-Aug-2013 (1 ex, CBG); Guelph, 30-Jun-2018 (1 ex, CBG); Perth, 03-Jul-2014 to 17-Jul-2014 (1 ex, CBG); Warsaw, 04-Jul-2014 to 18-Jul-2014 (1 ex, CBG). NEW BRUNSWICK: Kouchibouguac National Park, 19-Aug-2009 (1 ex, CBG). NOVA SCOTIA: Cape Breton Highlands National Park, 14-Jul-2013 to 19-Jul-2013 (1 ex, CBG); Cape Breton Highlands National Park, 19-Jul-2013 to 26-Jul-2013 (1 ex, CBG); Cape Breton Highlands National Park, 02-Aug-2013 to 09-Aug-2013 (1 ex, CBG); Kejimikujik National Park, 08-Aug-2013 to 22-Aug-2013 (1 ex, CBG). NEWFOUNDLAND: Gros Morne National Park, 25-Jun-2013 to 02-Jul-2013 (1 ex, CBG); Gros Morne National Park, 09-Jul-2013 to 16-Jul-2013 (1 ex, CBG); Gros Morne National Park, 09-Jul-2013 to 20-Jul-2013 (9 exx, CBG); Gros Morne National Park, 10-Jul-2013 to 20-Jul-2013 (47 exx, CBG); Gros Morne National Park, 11-Jul-2013 (1 ex, CBG); Gros Morne National Park, 12-Jul-2013 (2 exx, CBG); Gros Morne National Park, 15-Jul-2013 (1 ex, CBG); Gros Morne National Park, 17-Jul-2013 (1 ex, CBG); Gros Morne National Park, 22-Aug-2013 to 27-Aug-2013 (1 ex, CBG); Terra Nova National Park, 24-Jul-2013 to 30-Jul-2013 (3 exx, CBG).

Diagnostic information (based on Klausnitzer 1976). Body length 2.4 mm. Brown, elytral suture pale brown, antennae and legs yellow-brown. Male sternite IX and tergite IX as in Fig. 24C, D. Aedeagus as in Fig. 24A, B.

Bionomic notes. The Canadian specimens were collected in conifer and mixed forests, mainly with Malaise traps.

Comments. *Contacyphon obscurellus* belongs to the *C. variabilis* group of species. It is most reliably identified by its genitalia. The species is split into three closely clustered BINs, which show no obvious morphological differences. The identification of the Canadian specimens is based on dissected male representatives of these BINs.

Throscidae***Aulonothroscus distans* Blanchard, 1917**

BOLD:ABA9083

Distribution. Native to North America. Blanchard (1917) listed records from Massachusetts, New York and North Carolina.

Canadian records. ONTARIO: Balsam Lake Provincial Park, 02-Jun-2014 to 16-Jun-2014 (1 ex, CBG); Cambridge, 21-May-2015 to 27-May-2015 (1 ex,

CBG); Cambridge, 25-May-2015 to 31-May-2015 (1 ex, CBG); Cambridge, 04-Jun-2015 to 11-Jun-2015 (3 exx, CBG); Cambridge, 18-Jun-2015 to 24-Jun-2015 (1 ex, CBG); Elizabethtown-Kitley, 14-May-2010 to 18-May-2010 (1 ex, CBG); Elizabethtown-Kitley, 28-May-2010 to 30-May-2010 (2 exx, CBG); Georgian Bay Islands National Park, 04-Jun-2013 to 11-Jun-2013 (1 ex, CBG); Georgian Bay Islands National Park, 11-Jun-2013 to 18-Jun-2013 (1 ex, CBG); Georgian Bay Islands National Park, 18-Jun-2013 to 26-Jun-2013 (3 exx, CBG); Georgian Bay Islands National Park, 26-Jun-2013 to 30-Jun-2013 (2 exx, CBG); Georgian Bay Islands National Park, 30-Jul-2013 to 06-Aug-2013 (2 exx, CBG); Georgian Bay Islands National Park, 06-Aug-2013 to 19-Aug-2013 (1 ex, CBG); Georgian Bay Islands National Park, 03-Sep-2013 to 10-Sep-2013 (2 exx, CBG); Guelph, 17-Oct-2013 (1 ex, CBG); Hanover, 30-Jul-2014 to 13-Aug-2014 (1 ex, CBG); Rouge National Urban Park, 04-Jun-2013 to 11-Jun-2013 (1 ex, CBG); Rouge National Urban Park, 11-Jun-2013 to 18-Jun-2013 (6 exx, CBG); Rouge National Urban Park, 02-Jul-2013 to 09-Jul-2013 (1 ex, CBG); Rouge National Urban Park, 23-Jul-2013 to 30-Jul-2013 (1 ex, CBG); Rouge National Urban Park, 20-Aug-2013 to 27-Aug-2013 (1 ex, CBG); Rouge National Urban Park, 26-May-2014 to 03-Jun-2014 (3 exx, CBG); Rouge National Urban Park, 03-Jun-2014 to 10-Jun-2014 (2 exx, CBG); Rouge National Urban Park, 10-Jun-2014 to 17-Jun-2014 (3 exx, CBG); Rouge National Urban Park, 17-Jun-2014 to 24-Jun-2014 (3 exx, CBG); Rouge National Urban Park, 25-May-2014 to 01-Jul-2014 (1 ex, CBG); Rouge National Urban Park, 01-Jul-2014 to 08-Jul-2014 (1 ex, CBG); Rouge National Urban Park, 08-Jul-2014 to 15-Jul-2014 (1 ex, CBG); Rouge National Urban Park, 22-Jul-2014 to 29-Jul-2014 (2 exx, CBG); Rouge National Urban Park, 29-Jul-2014 to 05-Aug-2014 (1 ex, CBG); Rouge National Urban Park, 12-Aug-2014 to 19-Aug-2014 (1 ex, CBG); Rouge National Urban Park, 19-Aug-2014 to 26-Aug-2014 (1 ex, CBG); Rouge National Urban Park, 16-Sep-2014 to 23-Sep-2014 (1 ex, CBG); Rouge National Urban Park, 30-Sep-2014 to 07-Oct-2014 (1 ex, CBG); Thousand Islands National Park, 01-Jun-2012 to 08-Jun-2012 (2 exx, CBG); Thousand Islands National Park, 07-Jul-2012 to 13-Jul-2012 (1 ex, CBG); Thousand Islands National Park, 08-Jun-2012 to 15-Jun-2012 (1 ex, CBG); Thousand Islands National Park, 22-Jun-2012 to 29-Jun-2012 (1 ex, CBG); Thousand Islands National Park, 25-May-2012 to 01-Jun-2012 (1 ex, CBG); Waterloo, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG).

Diagnostic information (based on Blanchard 1917). Body length 2.6–3.0 mm. Piceous brown. Eyes with a small semicircular emargination. Head with longitudinal carinae separate and slightly diverging on the frons. Prosternal striae entire.

Bionomic notes. The Canadian specimens were collected with Malaise traps, mostly in deciduous or mixed forests.

Comments. The shallow semicircular emargination of the eyes separates this species from the other known North American species of *Aulonothroscus* (Blanchard 1917).

***Trixagus carinifrons* (Bonvouloir, 1859)**

BOLD:ABW4548

Figure 25

Distribution. Native to the Palearctic region. Widespread in Europe, also recorded from the Russian Far East (Leseigneur 2007). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Guelph, 01-Aug-2013 to 08-Aug-2013 (3 exx, CBG); Guelph, 15-Aug-2013 to 22-Aug-2013 (1 ex, CBG); Guelph, 29-Aug-2013 to 05-Sep-2013 (1 ex, CBG); Guelph, 30-Sep-2013 to 04-Oct-2013 (1 ex, CBG); Rouge National Urban Park, 28-May-2013 to 04-Jun-2013 (1 ex, CBG); Rouge National Urban Park, 18-Jun-2013 to 25-Jun-2013 (4 exx, CBG); Rouge National Urban Park, 27-Aug-2013 to 03-Sep-2013 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2014 to 10-Jun-2014 (1 ex, CBG); Rouge National Urban Park, 24-Jun-2014 to 01-Jul-2014 (2 exx, CBG); Rouge National Urban Park, 29-Jul-2014 to 05-Aug-2014 (2 exx, CBG); Rouge National Urban Park, 05-Aug-2014 to 12-Aug-2014 (4 exx, CBG).

Diagnostic information (based on Leseigneur 1998, 2005). Body length 2.5–3.0 mm. Habitus as in Fig. 25A. Eyes deeply emarginate. Head with longitudinal carinae which converge towards the vertex. Sides of pronotum sinuate in males, slightly or not sinuate in females. Elytral apex in males with a fringe of hairs longer than the lateral pubescence, often hidden under the elytra and not visible without removal of the abdomen. Aedeagus as in Fig. 25B.

Bionomic notes. In Europe, this species is usually found in dry, warm habitats: heaths, forest edges, gravel pits, etc. (Koch 1989b). The Canadian specimens were collected in Malaise traps in a patch of forest and on a lawn between buildings on the University of Guelph campus.


Comments. The genus *Trixagus* includes several overlooked and probably undescribed species in Canada based on DNA barcode data and initial studies of male genitalia of the barcoded material (Bouchard et al. 2017, MP unpublished data). Until the genus is revised, the two Palearctic *Trixagus* species reported here are most reliably identified using DNA barcodes or male genitalia.

***Trixagus meyhohmi* Leseigneur, 2005**

BOLD:ACC5051

Figure 26

Distribution. Recently described, distribution not yet thoroughly known. Apparently widespread in Europe (Leseigneur 2005, Mertlik and Leseigneur 2007, Silfverberg 2010, Rassi et al. 2015). Adventive in the Nearctic region (British Columbia, Ontario, Quebec, and Nova Scotia, Canada).



Figures 25, 26. **25** *Trixagus carinifrons* (Bonvouloir) **25A** habitus, L. Borowiec **25B** aedeagus **26** *Trixagus meybohmi* Leseigneur **26A** habitus **26B** aedeagus. Scale bars: 1.0 mm (**25A**; **26A**), 0.5 mm (**25B**; **26B**).

Canadian records. BRITISH COLUMBIA: North Vancouver, 21-Sep-2015 to 02-Oct-2015 (2 exx, CBG); Vancouver, 20-May-2014 to 26-May-2014 (2 exx, CBG); Vancouver, 03-Jun-2014 to 10-Jun-2014 (1 ex, CBG); Vancouver, 12-Aug-2014 to 19-Aug-2014 (2 exx, CBG); Vancouver, 26-Aug-2014 to 02-Sep-2014 (1 ex, CBG); Vancouver, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); West Vancouver, 21-Sep-2015 to 02-Oct-2015 (12 exx, CBG). ONTARIO: Dundas, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Toronto, 21-Sep-2015 to 02-Oct-2015 (3 exx, CBG). QUEBEC: Montreal, 19-Sep-2014 to 26-Sep-2014 (1 ex, CBG). NOVA SCOTIA: Point Pleasant Park, 25-May-2013 to 01-Jun-2013 (1 ex, CBG); Point Pleasant Park, 14-Jun-2013 to 22-Jun-2013 (1 ex, CBG); Point Pleasant Park, 06-Jul-2013 to 13-Jul-2013 (2 exx, CBG); Point Pleasant Park, 03-Aug-2013 to 10-Aug-2013 (9 exx, CBG); Point Pleasant Park, 10-Aug-2013 to 17-Aug-2013 (7 exx, CBG); Point Pleasant Park, 17-Aug-2013 to 24-Aug-2013 (6 exx, CBG); Point Pleasant Park, 24-Aug-2013 to 31-Aug-2013 (4 exx, CBG); Point Pleasant Park, 31-Aug-2013 to 07-Sep-2013 (1 ex, CBG); Point Pleasant Park, 07-Sep-2013 to 14-Sep-2013 (2 exx, CBG).

Diagnostic information (based on Leseigneur 2005). Body length 2.4–3.2 mm. Habitus as in Fig. 26A. Eyes deeply emarginate. Head with slender longitudinal carinae, which are parallel or slightly converging towards the vertex. Sides of pronotum not or only slightly sinuate, slightly angulate in males, rounded or weakly angulate in females. Elytral apex in males with a fringe of long setae (setae as long as antennomere 11), which can be tucked under the elytra and may not be visible without removal of the abdomen. Aedeagus as in Fig. 26B.

Bionomic notes. The Canadian specimens were collected with Malaise traps in city parks and suburban residential areas.

Comments. Until the North American species of *Trixagus* are revised, *T. meyhohmi* is most reliably identified using DNA barcodes or male genitalia.

Elateridae

Dendrometrinae

Prosternini

Pseudanostirus tigrinus (Fall, 1901), comb. nov.

BOLD: ACU2924

Distribution. Native to North America. Previously known only from the United States, where the species is known from areas near Lake Tahoe in California (Fall 1901). CNC has additional specimens collected in northern Oregon and near Lake Tahoe in Nevada.

Canadian record (DNA barcoded specimen). BRITISH COLUMBIA: Gulf Islands National Park Reserve, 30-May-2014 to 08-Jun-2014 (1 ex, CBG).

Additional Canadian record. BRITISH COLUMBIA: Parksville, 11-Apr-2018 (1 ex, CNC).

Diagnostic information (based on Brown 1936). Body length 9.0–11.2 mm. Antennae with antennomeres 3 and 5 of equal length. Pronotum black, with pubescence pale except for two to circular patches of dark setae on each side. Elytra red-brown with pale setae, with band of darker setae surrounding scutellar shield and three angulate transverse bands of dark setae extending from suture to epipleura.

Bionomic notes. *Pseudanostirus tigrinus* has been collected by beating *Pseudotsuga* Carrière on a grassy hillside with *Quercus garryana* Douglas ex Hook. trees. Other specimens have been collected in Malaise and funnel traps also in semi-open woodland with *Arbutus* L. and *Pseudotsuga* trees in warm-summer Mediterranean climate areas. The barcoded specimen was collected with a Malaise trap in a coastal mixed forest.

Comments. This species was described as *Corymbites tigrinus* Fall, 1901. Brown (1936) placed this species in *Ludius* Berthold, 1827 as *Ludius tigrinus* (Fall, 1901), part of the *L. triundulatus* species group. Lane (1948) found that *Ludius* Eschscholtz, 1829 was a synonym of *Elater* Linnaeus, 1758 and transferred all North American *Ludius* to *Ctenicera* Latreille, 1829. Johnson (2002) indicated that all species of Brown's *triundulatus* group should be transferred to *Pseudanostirus* but did not formally present any new combinations. This combination has not been used previously in the scientific literature. Therefore the resulting combination *Pseudanostirus tigrinus* (Fall, 1901) is used here for the first time.

Pseudanostirus tigrinus is similar to *P. nebraskensis* (Bland, 1863). Its independent placement in a separate BIN cluster supports the validity of *P. tigrinus*.

Cantharidae

Cantharinae

Podabrini

Dichelotarsus lapponicus (Gyllenhal, 1810)

BOLD:ACD1483

Figure 27

Distribution. Previously only recorded from the Palaearctic region. A northern species, found in Norway, Sweden, and Finland, and across the northern Palaearctic to the Russian Far East and Japan (Hokkaido) (Kazantsev and Brancucci 2007; Silfverberg 2010; Rassi et al. 2015). Probably Holarctic and previously overlooked in North America.

Canadian records. YUKON TERRITORY: Ivvavik National Park, 17-Jun-2014 to 23-Jun-2014 (3 exx CBG); Ivvavik National Park, 23-Jun-2014 to 29-Jun-2014 (16 exx, CBG).

Diagnostic information (based on Kazantsev 1998). Body length 7–10 mm. Habitus as in Fig. 27A. Dark brown to black, basal antennomeres, mandibles and usually clypeus (at least at the margins) yellow. Legs variably yellow, usually at least the profemora yellow. Third antennomere in males ca. 1.5 times as long as the second. Pronotum as wide as long or slightly wider than long, with sides concave before acute



Figures 27, 28. **27** *Dichelotarsus lapponicus* (Gyllenhal) **27A** habitus **27B** aedeagus, ventral view (everted endophallus removed) **27C** aedeagus, lateral view (everted endophallus removed) **27D** endophallus, lateral view **28** *Malthodes pumilus* (Brébisson), habitus, L. Borowiec. Scale bars: 2.0 mm (**27A**), 0.5 mm (**27B–D**), 0.3 mm (**28**).

hind angles. All tarsal claws in both sexes with a broad, blunt basal tooth, no claws deeply cleft. Aedeagus as in Fig. 27B–D, with dorsal plate with apical notch.

Bionomic notes. In Northern Finland, this species is found both above and below the treeline, usually in wetlands (MP, pers. obs.). The Canadian specimens were collected with a Malaise trap on tundra close to the Arctic treeline.

Comments. The remote arctic collecting locality suggests that this species is more likely to be Holarctic than adventive from the Palearctic region. The legs and basal antennomeres of the Canadian specimens are darker and the body length is slightly smaller compared to North European material we examined (including the DNA bar-coded Finnish specimens with which the Canadian specimens share the BIN cluster). The male genitalia and shape of the pronotum show no differences between the European and Canadian specimens. Based on the identification keys, descriptions and figures by Fall (1927) and Fender (1961), *D. lapponicus* closely resembles *D. piniphilus* (Eschscholtz, 1830). The tarsal claw formula is the same and the shape of the pronotum is very similar in both species. The dorsal plate of the aedeagus has an apical notch in *D. lapponicus* (as in Fig. 27B), whereas in *D. piniphilus* it is apically truncate or subtruncate. The clypeus of *D. lapponicus* is usually yellow at least at the margins. The yellow color is more extensive in males in the material we have seen, and only faint red-brown spots are visible on the clypeus of some female specimens. The clypeus is black in *D. piniphilus*. Pelletier and Hébert (2014) state that *D. lapponicus* resembles *D. perplexus* (W.J. Brown, 1940), which is known from across boreal and arctic Canada, but *D. perplexus* is smaller (body length 5.0–6.5 mm) and has a different tarsal claw formula.

Malthininae

Malthodini

Malthodes pumilus (Brébisson, 1835)

BOLD: AAP7843

Figure 28

Distribution. Native to the Palearctic region. Widespread in Europe, also recorded from Iran and Turkey (Kazantsev and Brancucci 2007). Adventive in the Nearctic region (British Columbia, Alberta, Saskatchewan, Manitoba, Ontario, Quebec, New Brunswick, Nova Scotia, Prince Edward Island, and Newfoundland, Canada).

Canadian records. BRITISH COLUMBIA: Mount Revelstoke National Park, 19-Jun-2014 to 26-Jun-2014 (15 exx, CBG); Mount Revelstoke National Park, 04-Jul-2014 to 09-Jul-2014 (22 exx, CBG); New Afton Mine, 06-Jun-2013 to 13-Jun-2013 (4 exx, CBG); New Afton Mine, 20-Jun-2013 to 27-Jun-2013 (1 ex, CBG); Vancouver, 03-Jun-2014 to 10-Jun-2014 (4 exx, CBG); Vancouver, 17-Jun-2014 to 24-Jun-2014 (1 ex, CBG); Yoho National Park, 25-Jun-2014 to 07-Jul-2014 (1 ex, CBG). ALBERTA: Banff National Park, 17-Jun-2012 (1 ex, CBG); Banff National Park,

19-Jun-2012 (1 ex, CBG); Banff National Park, 15-Jun-2012 to 20-Jun-2012 (1 ex, CBG); Elk Island National Park, 22-Jun-2012 to 29-Jun-2012 (1 ex, CBG); Elk Island National Park, 01-Jul-2012 (1 ex, CBG); Elk Island National Park, 02-Jul-2012 (1 ex, CBG); Elk Island National Park, 29-Jun-2012 to 03-Jul-2012 (17 exx, CBG); Elk Island National Park, 06-Jul-2012 to 13-Jul-2012 (8 exx, CBG); Jasper National Park, 04-Jul-2012 to 11-Jul-2012 (1 ex, CBG); Waterton Lakes National Park, 27-Jun-2012 to 04-Jul-2012 (3 exx, CBG); Waterton Lakes National Park, 10-Jul-2012 to 17-Jul-2012 (4 exx, CBG); Waterton Lakes National Park, 17-Jul-2012 to 24-Jul-2012 (3 exx, CBG). SASKATCHEWAN: Grasslands National Park, 29-Jun-2014 to 08-Jul-2014 (3 exx, CBG). MANITOBA: Riding Mountain National Park, 02-Jul-2012 to 09-Jul-2012 (10 exx, CBG). ONTARIO: Algonquin Provincial Park, 01-Jul-2014 to 15-Jul-2014 (2 exx, CBG); Bayview Escarpment Provincial Park, 29-May-2014 to 12-Jun-2014 (6 exx, CBG); Bayview Escarpment Provincial Park, 26-Jun-2014 to 10-Jul-2014 (1 ex, CBG); Bruce Peninsula National Park, 21-Jun-2012 to 28-Jun-2012 (2 exx, CBG); Bruce Peninsula National Park, 28-Jun-2012 to 07-Jul-2012 (3 exx, CBG); Bruce Peninsula National Park, 05-Jul-2012 to 12-Jul-2012 (3 exx, CBG); Bruce Peninsula National Park, 12-Jul-2012 to 18-Jul-2012 (2 exx, CBG); Elizabethtown-Kitley, 02-Jun-2010 to 04-Jun-2010 (1 ex, CBG); Elizabethtown-Kitley, 04-Jun-2010 to 06-Jun-2010 (1 ex, CBG); Elizabethtown-Kitley, 06-Jun-2010 to 08-Jun-2010 (1 ex, CBG); Elizabethtown-Kitley, 12-Jun-2010 to 14-Jun-2010 (1 ex, CBG); Elizabethtown-Kitley, 15-Jun-2011 to 20-Jun-2011 (2 exx, CBG); Frontenac Provincial Park, 05-Jun-2014 to 19-Jun-2014 (3 exx, CBG); Georgian Bay Islands National Park, 04-Jun-2013 to 11-Jun-2013 (1 ex, CBG); Georgian Bay Islands National Park, 11-Jun-2013 to 18-Jun-2013 (7 exx, CBG); Georgian Bay Islands National Park, 18-Jun-2013 to 26-Jun-2013 (6 exx, CBG); Guelph, 19-Jun-2015 to 26-Jun-2015 (1 ex, CBG); Hartington, 30-May-2017 (1 ex, CBG); Hartington, 13-Jun-2017 (2 exx, CBG); Hartington, 28-Jun-2017 (1 ex, CBG); Inverhuron Provincial Park, 25-Jun-2014 to 09-Jul-2014 (4 exx, CBG); Lion's Head Provincial Park, 26-Jun-2014 to 10-Jul-2014 (1 ex, CBG); Lower Madawaska River Provincial Park, 02-Jul-2014 to 16-Jul-2014 (2 exx, CBG); Murphy's Point Provincial Park, 05-Jun-2014 to 19-Jun-2014 (1 ex, CBG); Peterborough, 24-May-2015 to 30-May-2015 (1 ex, CBG); Peterborough, 07-Jun-2015 to 13-Jun-2015 (2 exx, CBG); Pinery Provincial Park, 25-Jun-2014 to 09-Jul-2014 (1 ex, CBG); Puslinch Township, 05-Jun-2010 to 12-Jun-2010 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (2 exx, CBG); Sandbanks Provincial Park, 05-Jun-2014 to 19-Jun-2014 (1 ex, CBG); Short Hills Provincial Park, 26-May-2014 to 09-Jun-2014 (1 ex, CBG); Sudbury, 08-Jun-2010 (1 ex, CBG); Thousand Islands National Park, 01-Jun-2012 to 08-Jun-2012 (1 ex, CBG); Thousand Islands National Park, 08-Jun-2012 to 15-Jun-2012 (5 exx, CBG); Thousand Islands National Park, 15-Jun-2012 to 22-Jun-2012 (3 exx, CBG). QUEBEC: Forillon National Park, 05-Jul-2013 to 15-Jul-2013 (4 exx, CBG); Forillon National Park, 15-Jul-2013 to 22-Jul-2013 (6 exx, CBG); Forillon National Park, 22-Jul-2013 to 30-Jul-2013 (4 exx, CBG); Forillon National Park, 28-Jun-2013

to 05-Jul-2013 (1 ex, CBG); Mingan Archipelago National Park Reserve, 02-Jul-2013 to 09-Jul-2013 (1 ex, CBG). NEW BRUNSWICK: Fundy National Park, 18-Jun-2013 to 25-Jun-2013 (1 ex, CBG); Fundy National Park, 02-Jul-2013 to 09-Jul-2013 (22 exx, CBG); Fundy National Park, 09-Jul-2013 to 16-Jul-2013 (33 exx, CBG); Fundy National Park, 16-Jul-2013 to 23-Jul-2013 (8 exx, CBG). NOVA SCOTIA: Cape Breton Highlands National Park, 14-Jul-2013 to 19-Jul-2013 (1 ex, CBG); Cape Breton Highlands National Park, 19-Jul-2013 to 26-Jul-2013 (1 ex, CBG); Kejimikujik National Park, 13-Jun-2013 to 20-Jun-2013 (1 ex, CBG); Kejimikujik National Park, 27-Jun-2013 to 05-Jul-2013 (1 ex, CBG); Sable Island National Park Reserve, 01-Jul-2014 to 14-Jul-2014 (3 exx, CBG). PRINCE EDWARD ISLAND: Prince Edward Island National Park, 03-Jul-2013 to 10-Jul-2013 (1 ex, CBG). NEWFOUNDLAND: Gros Morne National Park, 12-Jul-2013 (1 ex, CBG); Gros Morne National Park, 09-Jul-2013 to 16-Jul-2013 (109 exx, CBG); Gros Morne National Park, 09-Jul-2013 to 20-Jul-2013 (24 exx, CBG); Gros Morne National Park, 10-Jul-2013 to 20-Jul-2013 (6 exx, CBG); Gros Morne National Park, 23-Jul-2013 to 30-Jul-2013 (25 exx, CBG); Gros Morne National Park, 06-Aug-2013 to 13-Aug-2013 (2 exx, CBG); Terra Nova National Park, 25-Jun-2013 to 02-Jul-2013 (4 exx, CBG); Terra Nova National Park, 09-Jul-2013 to 16-Jul-2013 (37 exx, CBG); Terra Nova National Park, 24-Jul-2013 to 30-Jul-2013 (3 exx, CBG).

Diagnostic information (based on Wittmer 1979). Body length 1.3–1.5 mm. Habitus as in Fig. 28. Unicolorous dark brown to black, tarsi slightly paler. Mandibles with a finely serrate additional tooth on the inner surface. Male with the last ventrite long and narrow, deeply emarginate (almost to the middle), last visible tergite likewise deeply emarginate.

Bionomic notes. In Europe, this eurytopic species is usually found in dry, warm habitats such as exposed forest edges, dry meadows etc. (Koch 1989b). The larvae probably live in dead wood as predators (Koch 1989b). The Canadian specimens were collected in a variety of habitats, mainly forests, and mainly in Malaise traps.

Comments. The minute size distinguishes this species from all other Canadian species of *Malthodes* except for *M. parvulus* (LeConte, 1851) (Fender 1951, Pelletier and Hébert 2014). *Malthodes parvulus* is paler, with the first two antennomeres, pronotum, elytra and legs yellow (Pelletier and Hébert 2014). The structure of the terminal abdominal segments in males is also quite different between these species (see Fender (1951) or Pelletier and Hébert (2014) for figures of *M. parvulus*). However, all morphologically examined Canadian specimens of *M. pumilus* were females. It is probably a mainly parthenogenetic species, as males are rare in Europe as well (Wittmer 1979). The genitalia and modifications of the terminal abdominal segments of males are often crucial for morphological identification of *Malthodes* species (Fender 1951; Wittmer 1970). The extreme scarcity of males for morphological diagnosis combined with the lack of recent taxonomic work on the genus in North America probably explains why *M. pumilus* has remained undetected despite being apparently widespread and common across Canada.

Dermestidae
Attageninae
Attagenini

***Attagenus smirnovi* Zhantiev, 1973)**

BOLD:AAI7919

Figure 29

Distribution. Native to the Afrotropical region. Adventive in the Palaearctic region, first recorded from Europe in the 1960s (misidentified under various species names), distribution expanded in recent decades (Stenggaard Hansen et al. 2012). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Toronto, 19-Jul-2016 (3 exx, CBG).

Diagnostic information (based on Peacock 1979, Halstead 1981, and Kalik 1992). Body length 2.3–4.0 mm. Habitus as in Fig. 29A, B. Dark brown to black with yellow pubescence, elytra red-brown in males, usually paler yellow-brown in females. Antennae and legs red-brown or yellow-brown. Male antennomere 11 slightly curved at the base, ca. four times longer than wide and ca. four times as long as the combined length of antennomeres 9 and 10. Female antennal club elongate, last antennomere not modified, ovoid. Propleurotrochantin exposed. Anterior ventral carina of mesofemur prominent and sharp, posterior carina weakly developed. Metacoxa reaching metepimeron.

Bionomic notes. This species is recorded from the nests of the Little swift (*Apus affinis* (J.E. Gray, 1830)) in Kenya (Peacock 1979). It is an indoor pest of various materials of animal origin in Europe (Stenggaard Hansen et al. 2012). The Canadian specimens (two larvae and one adult female) were collected in an apartment in Toronto.

Comments. Vernacularly known as the brown carpet beetle. The coloration makes this species quite distinctive among *Attagenus* species recorded from Canada. Presence of adults and larvae in a home suggest establishment in Canada. It is unknown how large or viable Canadian populations of this species are.

Ptinidae
Dorcatominae

***Petalium incisum* Ford, 1973**

BOLD:ACX8800

Distribution. Native to North America. Widespread in eastern United States (White 1982; Arango and Young 2012).

Canadian records. ONTARIO: Burlington, 07-Aug-2014 to 20-Aug-2014 (1 ex, CBG); Cambridge, 15-Jul-2017 (4 exx, CBG); Guelph, 30-Jul-2017 (1 ex, CBG).

Diagnostic information. See Ford (1973) and Arango and Young (2012).




Figure 29. *Attagenus smirnovi* Zhantiev, habitus, L. Borowiec **A** male **B** female. Scale bar: 1.0 mm.

Bionomic notes. Ford (1973) reared this species from *Rhus toxicodendron* L. and *Robinia pseudoacacia* L. (probably from dead dry wood). One Canadian specimen was caught with a Malaise trap; the rest were collected by beating vegetation in deciduous and mixed forests.

Erotylidae
Cryptophilinae
Cryptophilini

Cryptophilus obliteratus Reitter, 1874
 BOLD:AAP6170

Distribution. Native to the Palearctic region (Asia). Adventive in the western Palearctic region, recorded in Europe at least from Austria, Denmark, Germany, and France (Denux and Zagatti 2010). Adventive in the Nearctic region (Arkansas, Massachusetts, Maryland, Minnesota, and Wisconsin, United States (Esser 2017), and Ontario, Canada).

Canadian records. ONTARIO: Ailsa Craig, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Dunnville, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); London, 22-Apr-2013 to 03-May-2013 (2 exx, CBG); Orillia, 20-Apr-2015 to 08-May-2015 (1 ex, CBG); Peterborough, 05-Jul-2015 to 11-Jul-2015 (1 ex, CBG); Port Rowan, 22-Apr-2013

to 03-May-2013 (2 exx, CBG); Scarborough, 20-Apr-2015 to 08-May-2015 (1 ex, CBG); Whitby, 22-Apr-2013 to 03-May-2013 (5 exx, CBG).

Diagnostic information (based on Esser 2016). Body length ca. 2.5 mm. Color red-brown with more-or-less extensive black markings on the elytra, usually a transverse black band or black lateral spots. Lateral margins of pronotum rounded, more abruptly narrowed basally, concave just in front of the sharp, approximately right-angled hind angles. Meso- and metatibiae bent inwards in males. Parameres relatively shorter than in the other two species recorded from North America. For habitus and genital figures, see Esser (2016) and Jens Esser's homepage: <http://cryptophagidae.de/Cryptophilus-Erotylidae-pleasing-fungus-beetles/>

Bionomic notes. This species has been collected from heaps of compost and garden waste in Germany (Franzen 1991). The Canadian specimens were collected with Malaise traps, mainly from suburban residential areas.

Comments. Esser (2017) reported *Cryptophilus obliteratus* from the United States and synonymized *Cryptophilus seriatus* Casey, 1924 (described from Massachusetts) with it. This species has obviously been present in North America for a long time, and Canadian records older than those reported here may well be found in collections.

Cryptophilus propinquus Reitter, 1874 BOLD: AAY6550

Distribution. This species was confused with *Cryptophilus angustus* (Rosenhauer, 1856) under the name *Cryptophilus integer* (Heer, 1841) until recently. Therefore, its distribution is not yet very well known. *Cryptophilus propinquus* was described from Japan, and has been recorded at least from Germany, India, Italy, Turkey, and the United States (Esser 2016, 2017). Adventive in the Nearctic region (Maryland, Minnesota, Mississippi, and Wisconsin, United States (Esser 2017), and British Columbia and Ontario, Canada).

Canadian records. BRITISH COLUMBIA: Victoria, 25-Jun-2014 to 02-Jul-2014 (1 ex, CBG); Victoria, 23-Jul-2014 to 30-Jul-2014 (2 exx, CBG). ONTARIO: Cambridge, 04-Jun-2015 to 11-Jun-2015 (1 ex, CBG); Guelph, 15-Jul-2010 to 01-Aug-2010 (1 ex, CBG); Guelph, 17-Sep-2017 (2 exx, CBG); Rouge National Urban Park, 15-Sep-2013 (1 ex, CBG).

Diagnostic information (based on Esser 2016). Body length ca. 2 mm. Red-brown without black markings on the elytra. Sides of pronotum evenly rounded in dorsal view, no concavity before blunt hind angles. Meso- and metatibiae dilated distally, sometimes with a blunt angle on the dorsal edge in males, but not bent ventrad in either sex. Parameres relatively longer than in *C. obliteratus*, but shorter than in *C. angustus*. For habitus and genital figures, see Esser (2016) and Jens Esser's homepage: <http://cryptophagidae.de/Cryptophilus-Erotylidae-pleasing-fungus-beetles/>

Bionomic notes. This species is found in decaying plant material, e.g., in compost heaps (Koch 1989b, Ruta et al. 2011). The DNA barcoded Canadian specimens were

collected in Malaise traps in suburban residential areas, in pitfall traps in forest fragments, and by sifting compost heaps.

Comments. *Cryptophilus integer* (Heer, 1841) is listed as occurring in Canada by (Bousquet et al. 2013), but Esser (2016, 2017) discovered that the name is not valid and refers specimens identified as *C. integer* to two different species: *C. propinquus* Reitter, 1874 and *C. angustus* (Rosenhauer, 1856) (= *C. simplex* (Wollaston, 1857)). Esser (2017) reported that both of these species occur in the Nearctic region but listed no records from Canada for either species. We found only *C. propinquus* among the DNA barcoded Canadian specimens, but not *C. angustus*.

Cryptophagidae

Cryptophaginae

Cryptophagini

Henoticus mycetoecus (Park, 1929)

BOLD:ACK1696

Distribution. Native to North America. Described from Illinois (Park 1929), also recorded from Iowa (Downie and Arnett 1996).

Canadian records. ONTARIO: Rouge National Urban Park, 18-Jun-2013 to 25-Jun-2013 (1 ex, CBG).

Diagnostic information (based on Park (1929)). Body length 1.8–2.0 mm. More or less uniformly red-brown, with legs, antennae, and medial part of elytra paler. Lateral margins of pronotum serrate, sublateral carinae absent. Posterior of pronotum with two deep foveae connected by a distinct basal groove.

Bionomic notes. Park (1929) collected the type specimens from decaying fruiting bodies of the polypore fungus *Climacodon septentrionalis* (Fr.) P. Karst. in a sugar maple forest. The Canadian specimen was caught with a Malaise trap in a patch of forest.

Phalacridae

Phalacrinae


Acylomus ergoti Casey, 1890

BOLD:AAH0135

Figure 30

Distribution. United States (Gimmel 2013). *Tinodemus grouvellei* Guillebeau, 1894, synonymized under *A. ergoti* by Gimmel (2013), was described from Michigan.

Canadian records. ONTARIO: Brantford, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG); Breslau, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Cambridge, 07-May-2015 to 14-May-2015 (3 exx, CBG); Cambridge, 14-May-2015 to 21-May-2015



Figures 30, 31. **30** *Acylomus ergoti* Casey **30A** male metatibia, ventral view **30B** aedeagus, dorsal view **30C** tegmen, dorsal view **30D** tegmen, lateral view **31** *Olibrus liquidus* Erichson, habitus, L. Borowiec. Scale bars: 0.2 mm (**30A–D**), 1.0 mm (**31**).

(1 ex, CBG); Cambridge, 25-May-2015 to 31-May-2015 (1 ex, CBG); Elizabethtown-Kitley, 30-Apr-2010 to 02-May-2010 (1 ex, CBG); Elizabethtown-Kitley, 09-May-2010 to 14-May-2010 (1 ex, CBG); Elizabethtown-Kitley, 14-May-2010 to 18-May-2010 (1 ex, CBG); Elizabethtown-Kitley, 30-May-2010 to 02-Jun-2010 (1 ex, CBG); Embro, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Georgian Bay Islands National Park, 06-May-2013 to 12-May-2013 (1 ex, CBG); Georgian Bay Islands National Park, 12-May-2013 to 23-May-2013 (9 exx, CBG); Georgian Bay Islands National Park, 30-Jul-2013 to 06-Aug-2013 (1 ex, CBG); Hagersville, 23-Sep-2013 to 04-Oct-2013 (1 ex, CBG); Hartington, 28-Jun-2017 (1 ex, CBG); London, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Mississauga, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG); Orangeville, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Peterborough, 31-May-2015 to 06-Jun-2015 (1 ex, CBG); Peterborough, 29-Jun-2015 to 02-Jul-2015 (1 ex, CBG); Point Pelee National Park, 02-May-2012 to 09-May-2012 (8 exx, CBG); Point Pelee National Park, 16-May-2012 to 23-May-2012 (10 exx, CBG); Point Pelee National Park, 27-Jun-2012 to 04-Jul-2012 (1 ex, CBG); Point Pelee National Park, 05-Sep-2012 to 12-Sep-2012 (1 ex, CBG); Rouge National Urban Park, 07-Jun-2013 (2 exx, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (10 exx, CBG); Rouge National Urban Park, 17-

Jun-2014 to 24-Jun-2014 (1 ex, CBG); Rouge National Urban Park, 08-Jul-2014 to 15-Jul-2014 (1 ex, CBG); Williamstown, 22-Sep-2014 to 03-Oct-2014 (2 exx, CBG). QUEBEC: Montreal, 24-Jul-2014 to 02-Aug-2014 (1 ex, CBG); Montreal, 19-Sep-2014 to 26-Sep-2014 (1 ex, CBG). NEW BRUNSWICK: Springfield, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG).

Diagnostic information (based on the redescription of *Tinodemus grouvellei* by Švec 2002). Body length 2.0 mm. Oval, dark brown, elytral suture and base of pronotum laterally paler. Legs, antennae and mouthparts red, ventral side orange. Head and pronotum without microsculpture. Scutellum and elytra finely and densely transversely strigose. Sutural stria of elytra present in the apical 5/8. Male metatibiae short and widened apically, twice as wide at apex as proximally, medio-apical spine ca. twice as long as metatarsomere 1, curved at apex (Fig. 30A). Female metatibiae not modified. Male genitalia as in Fig. 30B, C.

Bionomic notes. The Canadian specimens were collected in various habitats (grasslands, forests, wetlands, residential areas etc.), mainly with Malaise traps.

Comments. The lack of a modern species-level revision of *Acylomus* prevents detailed comparison to most other Nearctic species of *Acylomus*. The only other species of *Acylomus* previously known from Canada, *A. pugetanus* Casey, 1916, was redescribed by Steiner and Singh (1987). It is darker, especially ventrally, and has different male genitalia and no apparent sexual dimorphism in metathoracic leg structure. At least one more species of *Acylomus* (BOLD:ACM7465) occurs in Canada according to DNA barcode data and initial morphological analysis of the barcoded specimens, but we have not been able to identify it to species level.

***Olibrus liquidus* Erichson, 1845**

BOLD:AAX0584

Figure 31

Distribution. Native to the Palaearctic region, widespread in Europe and North Africa (Švec 2007). Adventive in the Nearctic region (British Columbia, Canada).

Canadian records. BRITISH COLUMBIA: Burnaby, 20-Apr-2015 to 08-May-2015 (1 ex, CBG).

Diagnostic information (based on Thompson 1958; Vogt 1967b). Body length 1.9–2.6 mm. Habitus (Fig. 31) elongate-oval, narrower towards the elytral apex. Dark brown, elytra paler towards the apex, antennae, palpi and legs yellow-brown. Head and pronotum without microsculpture. Elytra fully covered by fine, net-like microsculpture in females, anterior third without microsculpture in males. Elytra with two sutural striae, which converge and usually meet towards the apex. Metaventricle densely and coarsely punctate, especially laterally. See Thompson (1958) for figures of the male genitalia (tegmen) and female ovipositor.

Bionomic notes. This species feeds on a variety of Asteraceae genera, usually in dry and warm habitats in Europe (Koch 1989b). The Canadian specimen was collected with a Malaise trap in a suburban residential area.

Comments. Lack of a modern revision of North American *Olibrus* prevents detailed comparison of *O. liquidus* to the native species. It is most reliably identified using male genitalia or DNA barcodes. Good illustrations of the genitalia are provided in volume 5, part 5b of the Handbooks for the Identification of British Insects (Thompson 1958).

Nitidulidae

Epuraeinae

Epuraeini

Epuraea unicolor (Olivier, 1790)

BOLD:AAX8537

Figure 32

Distribution. Native and widespread in the Palaearctic region. Recorded from North Africa and all of Europe to the Russian Far East and Japan (Audisio 1993; Jelínek and Audisio 2007) One of the most common and abundant species of the genus in Europe (Audisio 1993). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Guelph, 01-Nov-2009 (1 ex, CBG); Guelph, 22-Apr-2017 (1 ex, CBG); Guelph, 06-Jun-2018 (3 exx, CBG); Guelph, 30-Jun-2018 (1 ex, CBG); Rouge National Urban Park, 03-Jun-2013 to 09-Jun-2013 (1 ex, CBG).

Diagnostic information (based on Audisio 1993). Body length 2.3–3.2 mm. Habitus elongate, subparallel, rather flattened (Fig. 32A). Color variable, body, legs, and antennae usually yellowish or red-brown, pronotum and elytra often laterally paler, antennal club usually darkened. Elytra variably darkened, with a rounded dark spot on each elytron at the apical third, or with more extensive, irregular but symmetric dark patterns. Antennae with club ca. 1.5 times as long as wide. Head with subcircular, moderately impressed punctation, punctures approximately the size of the ommatidia, separated by 0.5–0.6 times their diameter, interspace with fine microsculpture. Punctures on pronotum and elytra slightly larger, but with similar microsculpture and relative distance between them. Pronotum 1.45–1.65 times as wide as long, broadest in the basal third, abruptly narrowed towards the protruding hind angles, anterior edge with a deep, wide, trapezoidal emargination. Elytral apices separately broadly rounded. Metaventrite with a wide V-shaped emargination at the hind edge. Male mesotibia distally slightly widened, with a small tooth at the inner margin (Fig. 32B). Female mesotibia unmodified. Male genitalia as in Fig. 32C–E.

Bionomic notes. This species occurs in decaying and fermenting organic material (e.g., fruit, fruiting bodies of fungi, tree sap), under the bark of dead trees etc., probably feeding on the microbes decomposing these materials (Audisio 1993). Often found in anthropogenic habitats such as orchards, cultivated fields, and garbage dumps (Audisio 1993). The Canadian specimens were collected by sifting a compost heap in a suburban backyard, in a Malaise trap in a residential area, and in pitfall traps at a riverside in Rouge National Urban Park.




Figure 32. *Epuraea unicolor* (Olivier) **A** habitus, L. Borowiec **B** male mesotibia **C** penis **D** parameres, ventral view **E** parameres, lateral view. Scale bars: 1.0 mm (**A**), 0.2 mm (**B–E**).

Comments. The lack of a modern revision of North American *Epuraea* prevents detailed comparison to other Canadian species at the moment. *Epuraea unicolor* can be reliably separated by DNA barcodes from all other Palearctic and Nearctic *Epuraea* species sampled so far. The diagnostic information above, in particular the male mesotibia and genitalia, should allow morphological identification.


Coccinellidae
Coccinellinae
Chilocorini

***Chilocorus renipustulatus* (Scriba, 1791)**

BOLD:AAO1521

Figure 33

Distribution. Native to the Palearctic region. Widespread in Europe, also recorded from Siberia and the Russian Far East (Kovář 2007). Adventive in the Nearctic region (Ontario, Canada).



Figures 33, 34. **33** *Chilocorus renipustulatus* (Scriba) **33A** habitus, L. Borowiec **33B** median lobe (penis guide) and parameres, ventral view **33C** penis (siphon), tip in lateral view **34** *Nephus bisignatus* (Boheman) **34A** habitus, S. Karjalainen **34B** median lobe (penis guide) and parameres, lateral view **34C** penis (siphon), tip in lateral view. Scale bars: 1.0 mm (**33A**), 0.5 mm (**33B**; **34A**), 0.2 mm (**33C**; **34B**), 0.1 mm (**34C**).

Canadian records. ONTARIO: Hamilton, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG); Mississauga, 19-Sep-2016 to 30-Sep-2016 (3 exx, CBG); Windsor, 22-Sep-2014 to 03-Oct-2014 (1 ex, CBG).

Diagnostic information (based on Fürsch 1967 and Gordon 1985). Body length 4–5 mm. Habitus as in Fig. 33A. Black, shiny, with a single rounded or slightly transverse orange-red macula on each elytron, abdomen laterally and apically orange, medial part of first ventrite black. Pronotum without distinct microsculpture on disc. Male genitalia as in Fig. 33B, C.

Bionomic notes. The main habitat in Europe is broadleaf forest, and the preferred prey are scale insects, in particular *Chionaspis salicis* (Linnaeus, 1758) (Koch 1989b). The Canadian specimens were collected with Malaise traps in suburban residential areas.

Comments. *Chilocorus kuwanae* Silvestri, 1909, an East Asian species introduced to the United States and recorded from across the country (Gordon 1985; Hendrickson et al. 1991), was recently synonymized with *C. renipustulatus* by Bieńkowski (2018). According to Bieńkowski, male genitalia are similar throughout the distribution areas of both species. However, Bieńkowski did not study any type material. One of the Canadian specimens shares an identical barcode haplotype with specimens of *C. renipustulatus* from Germany and Finland, others are slightly divergent (p-distance to European material varies from 0.006 to 0.015). Unfortunately, no barcode data are available for *C. kuwanae*. No Canadian records have been previously published under either name.

Chilocorus renipustulatus is externally very similar to *Chilocorus stigma* (Say, 1835) and its closest relatives. It can be distinguished using the male genitalia and microsculpture of the pronotum. In *C. stigma* and allied species, the interspace between punctures on the disc of the pronotum is covered by finely engraved, netlike microsculpture. In *C. renipustulatus*, the interspace is smooth and shiny, with no visible microsculpture on disc. The orange maculae on the elytra are more transverse in *C. renipustulatus* than in *C. stigma* in the examined DNA barcoded Canadian material of these species, but the maculae are known to vary in size and shape in *C. renipustulatus* (Bieńkowski, 2018).

Coccinellinae

Scymnini

Nephus bisignatus (Boheman, 1850)

BOLD:ACD2027

Figure 34

Distribution. Previously known only from Europe, where the species is more common in the north and rather sporadic in the central and southern parts (Fürsch 1965; Kovář 2007; Silfverberg 2010; Rassi et al. 2015). Probably Holarctic and previously overlooked in North America.

Canadian records. NUNAVUT: Kugluktuk, 25-Jun-2010 (1 ex, CNC); Kugluktuk, 01-Jul-2010 (1 ex, CNC); Kugluktuk, 11-Jul-2010 (1 ex, CNC); Kugluktuk, 13-Jul-2010 (1 ex, CNC).

Diagnostic information (based on Fürsch 1965, 1967, 1987). Body length 1.5–2.0 mm. Habitus elongate-oval (Fig. 34A). Black, with the anterior edge of the pronotum and often the apical edge of the elytra narrowly brown. Each elytron with a single small, obscurely delimited red-brown spot close to the apex, sometimes very faintly visible. Antennae with nine antennomeres. Pronotum very finely punctate, with strong, netlike microsculpture. Postcoxal lines on first abdominal ventrite briefly parallel to the hind margin of the ventrite at the middle, with the apices curved forward laterally. Male genitalia as in Fig. 34B, C.

Bionomic notes. *Nephus bisignatus* prefers open, usually sandy habitats in Europe (Koch 1989b). The Canadian specimens were collected in mesic tundra with yellow pan and pitfall traps.

Comments. *Nephus bisignatus* belongs to subgenus *Bipunctatus* Fürsch, 1987, which is characterized by having only nine antennomeres (Fürsch 1987). All the previously recorded Canadian species have either ten or eleven antennomeres (Gordon 1976, 1985). The remote collecting localities in the arctic tundra indicate that this species is probably Holarctic and previously overlooked in North America rather than adventive from the Palearctic region. Two subspecies are known from Europe (Kovář 2007), but we refrain from assigning the Canadian specimens to any subspecies. Among the Nearctic fauna, *N. bisignatus* resembles *N. georgei* (Weise, 1929), but has a narrower body outline and usually smaller and less conspicuous elytral spots. *Nephus georgei* also has ten antennomeres instead of nine. Gordon (1976) notes that specimens of *N. georgei* from the northern parts of the Northwest Territories are smaller and narrower compared to specimens from southern Canada and northern United States, and that the pale color pattern of the elytra is reduced in the northern specimens. Based on these notes, the arctic specimens of *N. georgei* may actually represent *N. bisignatus* and need to be re-examined.

***Scymnus rubromaculatus* (Goeze, 1777)**

BOLD:AAN9250

Figure 35

Distribution. Native to the Palearctic region, widespread across Eurasia from western Europe to the Russian Far East (Kovář 2007). Adventive in the Nearctic region (Ontario and Nova Scotia, Canada).

Canadian records. ONTARIO: Barrie, 22-Apr-2013 to 03-May-2013 (1 ex, CBG); Burlington, 21-Jul-2017 (1 ex, CBG); Guelph, 06-Jun-2013 to 13-Jun-2013 (1 ex, CBG); Guelph, 20-Jun-2013 to 27-Jun-2013 (2 exx, CBG); Guelph, 01-Aug-2013 to 08-Aug-2013 (1 ex, CBG); Guelph, 15-Aug-2013 to 22-Aug-2013 (1 ex, CBG); Mississauga, 15-Sep-2015 to 17-Sep-2015 (1 ex, CBG); Mississauga, 19-Sep-2016 to 30-Sep-2016 (3 exx, CBG); Toronto, 19-Jun-2017 to 27-Jun-2017 (1 ex, CBG). NOVA SCOTIA: Berwick, 20-Apr-2015 to 08-May-2015 (1 ex, CBG).




Figure 35. *Scymnus rubromaculatus* (Goeze) **A** female habitus, L. Borowiec **B** male habitus **C** median lobe (penis guide) and parameres, ventral view **D** median lobe (penis guide) and parameres, lateral view **E** penis (siphon), lateral view. Scale bars: 1.0 mm (**A, B**), 0.2 mm (**C, D**), 0.5 mm (**E**).

Diagnostic information (based on Fürsch 1967). Body length 1.8–2.3 mm. Habitus as in Fig. 35A, B. Color sexually dimorphic. Male: head, pronotum (apart from a black mediobasal spot), and legs yellow, otherwise black. Female: almost completely black with only the mouthparts, labrum and legs yellow. Femora often darkened. Post-coxal lines on 1st abdominal ventrite reaching the hind edge of the ventrite. Male genitalia as in Fig. 35C–E.

Bionomic notes. This species prefers dry, warm habitats in Europe and is found mainly on Brassicaceae, occasionally on trees and bushes (Koch 1989b). Most Canadian specimens were collected with Malaise traps in suburban areas.

Comments. *Scymnus rubromaculatus* leads to the couplets separating *Scymnus americanus* Mulsant, 1850, *S. apicanus* Chapin, 1973 and *S. paracanus* Chapin, 1973 in Gordon's keys to North American *Scymnus* (Gordon 1976, 1985). In *S. rubromaculatus*, the dorsal surface is more densely punctate and pubescent than in those three species, the hind margin of the elytra is slightly or not paler than the elytral disc, and the female has no pale markings on head (apart from labrum and mouthparts) or pronotum. The male genitalia differ: the apical hook of the penis (sipho), which is typical for *S. americanus* and related species, is absent in *S. rubromaculatus*.

Corylophidae

Corylophinae

Orthoperini

Orthoperus corticalis (Redtenbacher, 1845)

BOLD:ACC5439

Figure 36


Distribution. Native to the Palearctic region. Widely distributed from Western Europe to Siberia (Bowstead 1999, 2007). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Cambridge, 29-Apr-2015 to 07-May-2015 (1 ex, CBG); Cambridge, 07-May-2015 to 14-May-2015 (1 ex, CBG); Cambridge, 21-May-2015 to 27-May-2015 (1 ex, CBG).

Diagnostic information (based on Bowstead 1999). Body length 0.8–1.0 mm. Habitus slightly elongate oval, strongly convex (Fig. 36A). Dark brown to black, antennal base and legs pale, five apical antennomeres dark brown. Pronotum finely punctate, often with a transverse row of larger punctures medioposteriorly, with isodiametric microsculpture throughout. Elytral punctation fine, punctures larger basally, interspaces with similar microsculpture as the pronotum, the microsculpture forming wavy transverse rows of cells especially basally. Sutural striae of elytra present only at the apex. Male metaventricle with an elongate depression medially, and a short median keel behind the depression, distance of the keel from hind edge of metaventricle ca. 1/12 of the length of the metaventricle (Fig. 36B). Aedeagus as in Fig. 36C, D.

Bionomic notes. This species is mainly known from deciduous forests. It has been collected from a variety of fungus species growing on dead logs, and under the bark of fungus-infested logs (Bowstead 1999, Rutanen 2015). The Canadian specimens were collected with a Malaise trap at the edge of a forest.

Comments. This is the second species of *Orthoperus* recorded as adventive in Canada: the Palearctic *O. atomus* (Gyllenhal, 1808) is known from British Columbia in



Figures 36, 37. **36** *Orthoperus corticalis* (Redtenbacher) **36A** habitus, L. Borowiec **36B** male metaven-trite **36C** aedeagus, ventral view **36D** aedeagus, lateral view **37** *Litargus connexus* (Geoffroy) **37A** habitus, L. Borowiec **37B** antenna. Scale bars: 0.25 mm (**36A**), 0.2 mm (**36B–D**), 0.5 mm (**37A**), 0.2 mm (**37B**).

Canada, and Washington and Oregon in the United States (Klimaszewski et al. 2015). *Orthoperus corticalis* is darker and slightly larger than *O. atomus*, with stronger punctation on the pronotum and elytral base and denser and more strongly impressed microsculpture (Bowstead 1999). Two native North American species are currently known from Canada: *O. scutellaris* LeConte, 1878 has small V-shaped scratches on the elytra instead of punctures, and *O. suturalis* LeConte, 1878 has fine but distinctly impressed sutural striae (only faintly visible close to the elytral apex in *O. corticalis*) (LeConte 1878, Downie and Arnett 1996).

Mycetophagidae
Mycetophaginae
Mycetophagini

***Litargus connexus* (Geoffroy, 1785)**

BOLD:AAK8818

Figure 37

Distribution. Native to the Palaearctic region. Widespread in Europe, also recorded from North Africa, and across the region to the Russian Far East and Japan (Nikitsky 2008). Adventive in the Nearctic region (British Columbia, Canada).

Canadian records. BRITISH COLUMBIA: Burnaby, 20-Apr-2015 to 08-May-2015 (1 ex, CBG).

Diagnostic information (based on Vogt 1967a). Body length 2.4–2.8 mm. Dorsal habitus elongate, sides of elytra almost parallel (Fig. 37A). Black, elytra with transverse, undulating yellow bands at the base and just beyond midlength, and a yellow sutural spot close to the elytral apex. Yellow markings variable, basal band frequently broken into separate spots. Antennomere 8 somewhat wider than long, terminal antennomere approximately as long as wide, structure of the antennal club as in Fig. 37B. Elytra with epipleura concave, descending towards the lateral edge.

Bionomic notes. This species is found in deciduous and mixed forests in fungus-infested dead wood (Koch 1989b). The Canadian specimen was collected with a Malaise trap in a suburban residential area.

Comments. The combination of the elongate and nearly parallel-sided body, color pattern of the elytra and structure of the antennae will distinguish *L. connexus* from all other *Litargus* species known from Canada. Parsons (1975) provides diagnoses and illustrations of the native North American species.

Ciidae
Ciinae
Ciini


***Cis boleti* (Scopoli, 1763)**

BOLD:AAJ3268

Figure 38

Distribution. Native to the Palaearctic region. Recorded across Eurasia from western Europe to the Russian Far East and Japan (Jelínek 2008). Adventive in the Nearctic region (Ontario, Canada).

Canadian records (barcoded specimens). ONTARIO: Guelph, 21-Aug-2017 (2 exx, CBG); Guelph, 21-Oct-2017 (2 exx, CBG);



Figures 38, 39. **38** *Cis boleti* (Scopoli) **38A** habitus, L. Borowiec **38B** pronotum **39A** *Cis glabratus* Mellié, female habitus, L. Borowiec **39B** *Cis glabratus*, male head and pronotum **39C** *Cis glabratus*, aedeagus **39D** *Cis levettei* (Casey), aedeagus. Scale bars: 0.5 mm (**38A, B; 39A, B**), 0.2 mm (**39C, D**).

Additional Canadian records. ONTARIO: Horning's Mills, 08-Nov-2015 (1 ex, DEBU); Milton, 28-May-2002 (1 ex, DEBU).

Diagnostic information (based on Lohse 1967). Body length 2.8–4.0 mm. Habitus as in Fig. 38A. Dark brown to black (usually darker than in Fig. 38A). Pronotum more than 1.3 times wider than long, with concavities on disc (Fig. 38B). Lateral mar-

gins of pronotum widely deplanate and visible throughout in dorsal view. Elytra with dual punctuation: larger punctures arranged in irregular rows, with the intervals finely and densely punctate. Elytral vestiture consists of short, stout bristles.

Bionomic notes. This species feeds on polypore fungi, mainly *Trametes* species growing on deciduous trees (Koch 1989b, Reibnitz 1999). The barcoded Canadian specimens were collected from polypore fruiting bodies in a mixed forest.

Comments. *Cis submicans* Abeille de Perrin, 1874 (= *C. pistoria* Casey, 1898) is the only other representative of the mainly Palaearctic *C. boleti* species group known from North America (Lopes-Andrade et al. 2016). *Cis boleti* is a robust species, broader and on average larger than *C. submicans*, with the pronotum at least 1.3 times wider than long. The pronotal disc of *C. boleti* has indentations on both sides of the midline. These indentations are very shallow in *C. submicans*. The pronotum is also more densely punctate in *C. boleti* than in *C. submicans*.

Cis glabratus Mellié, 1848

BOLD:AAJ3241

Figure 39

Distribution. Native to the Palaearctic region, widespread in Europe (Jelínek 2008). Common in northern Europe, mainly found in higher elevations in Central Europe (Reibnitz 1999, Rassi et al. 2015). Adventive in the Nearctic region (Nova Scotia, Canada).

Canadian records. NOVA SCOTIA: Cape Breton Highlands National Park, 21-Jul-2009 (1 ex, CBG).

Diagnostic information (based on Lohse 1967). Body length 1.5–2.0 mm. Red-brown to dark brown, habitus as in Fig. 39A. Clypeus in male with two large, broad teeth (Fig. 39B). Pronotum widest behind middle, distinctly tapering towards the front angles. Vestiture on pronotum fine and pale. Outer edge of protibia serrated. Male with a large abdominal fovea on 1st abdominal ventrite. Aedeagus as in Fig. 39C.

Bionomic notes. The main host fungus in Europe is *Fomitopsis pinicola* (Sw.) P. Karst. (Reibnitz 1999). The Canadian specimen was collected in a jack pine forest in Cape Breton Highlands National Park.

Comments. *Cis glabratus* is externally very similar to *C. levettei* (Casey, 1898) and leads to that species in the key to North American species (Lawrence 1971). The microscopic vestiture of the pronotum is longer and more conspicuous in *C. glabratus*, but the most reliable morphological differences are in the male genitalia (Fig. 39C, D). *Cis levettei* forms a separate BIN (BOLD:ACA7530) which is more closely clustered to other Palaearctic members of the *C. nitidus* species group (*C. castaneus* (Herbst, 1793), *C. Jacquemartii* Mellié, 1848 and *C. lineatocribratus* Mellié, 1848) than to *C. glabratus*.

Mordellidae
Mordellinae
Mordellistenini

***Mordellistena militaris* LeConte, 1862**
BOLD:ACE3572

Distribution. Native to the Nearctic region. Previously recorded at least from Indiana, New York, North Carolina, and Ohio in the United States (Liljeblad 1945; Downie and Arnett 1996).

Canadian records. ONTARIO: Point Pelee National Park, 27-Jun-2012 to 04-Jul-2012 (5 exx, CBG).

Diagnostic information. See Liljeblad (1945).

Bionomic notes. The Canadian specimens were collected with a Malaise trap in a savanna with *Opuntia* cacti and sparse woody vegetation.

Comments. The coloration and the ridges of the hind legs of the Canadian specimens match both the Liljeblad (1945) diagnosis and the photographs of LeConte's type specimen in the type database of the Museum of Comparative Zoology at Harvard University. Therefore, we consider this record reliable despite the lack of a modern revision of the North American *Mordellistena*.

Zopheridae
Colydiinae
Synchitini

***Lasconotus subcostulatus* Kraus, 1912**
BOLD:ACU9668

Distribution. Native to the Nearctic region. Previously known from California, Idaho, Nevada, Oregon, Washington, South Dakota, Montana, and Nebraska in the United States (Lord et al. 2011–2013).

Canadian records (DNA barcoded specimen). SASKATCHEWAN: Grasslands National Park 21-May-2014 to 29-May-2014 (1 ex, CBG).

Additional Canadian records. BRITISH COLUMBIA: Aspen Grove, 20-Oct-1936 (5 exx, CNC); Merritt, 04-Jun-1922 (1 ex, CNC); Merritt, 08-Jun-1922 (2 exx, CNC); Merritt, 09-Jun-1922 (6 exx, CNC); Merritt, 10-Jun-1922 (1 ex, CNC); Merritt, 15-Jun-1922 (2 exx, CNC); Merritt, 18-Jun-1922 (1 ex, CNC); Merritt, 14-Sep-1923 (1 ex, CNC); Merritt, 03-Jun-1924 (1 ex, CNC); Merritt, 13-May-1925 (1 ex, CNC); Merritt, 25-Jul-1925 (1 ex, CNC); Olivier, 24-May-1958 (1 ex, CNC); Olivier, 12-Jun-1958 (10 exx, CNC); Olivier, 14-Jun-1958 (6 exx, CNC); Peachland, 19-Jul-1912

(1 ex, CNC); Peachland, 13-Jul-1919 (1 ex, CNC); Summerland, 25-Mar-1932 (16 exx, CNC); Summerland, 24-Sep-1932 (5 exx, CNC); Summerland, 7-Oct-1932 (1 ex, CNC); Summerland, 10-Oct-1932 (116 exx, CNC); Summerland, 11-Oct-1932 (5 exx, CNC); Summerland, 11-Nov-1932 (51 exx, CNC); Exact locality unknown, Sep-1923 (1 ex, CNC). SASKATCHEWAN: Crane Valley, 06-Oct-1914 (1 ex, CNC). MANITOBA: Aweme, 25-Jul-1919 (5 exx, CNC); Aweme, 31-Oct-1921 (1 ex, CNC); Onah, 24-Jul-1919 (5 exx, CNC).

Diagnostic information (based on Stephan 1989 and Lord et al. 2011–2013). Body length 2.5–2.8 mm. Pronotum with a central concave area covering 1/3 to 1/2 total width of pronotum, concave area bordered laterally by longitudinal raised ridges. Pronotum carinate anteriorly with double “U” shaped anterior margin. Elytral interstriae 5 more raised than other interstriae, forming a median concave area of the elytra typically on posterior half only. See Lord et al. (2011–2013) for a habitus photograph.

Bionomic notes. Hackwell (1973) reported that this species is associated with galleries of several species of bark beetles where it feeds on both fungi and bark beetles during larval development. Many of the CNC specimens were collected from pine trees (*Pinus contorta* Douglas ex Loudon, *P. monticola* Douglas ex D. Don, *P. ponderosa* Douglas ex C. Lawson). The DNA barcoded Canadian specimen was collected with a Malaise trap in a grassland.

Comments. The single DNA barcoded specimen from Saskatchewan (the only member of its BIN, with no closely clustered neighbors) was compared with specimens of this little-studied genus in the CNC. The identification of this specimen using data in Lord et al. (2011–2013) led to the further identification of several other Canadian specimens from British Columbia, Saskatchewan, and Manitoba. Examination of specimens collected 100 years ago in three provinces suggests that this species has long been part of the Canadian fauna.

Tenebrionidae

Alleculinae

Alleculini

Isomira angusta (Casey, 1891)

BOLD:AAH0400

Figure 40

Distribution. Native to the Nearctic region. Previously known from Georgia and South Carolina in the United States (Bousquet et al. 2018).

Canadian records. ONTARIO: Point Pelee National Park, 23-Jun-2010 (2 exx, CBG).

Diagnostic information (based on Aalbu et al. 2002). Body length 5.5–6.0 mm. Ventral surface of tarsi densely, finely pubescent (Fig. 40A). Male with sternite VIII deeply bilobed apically (Fig. 40B), extending beyond posterior edge of abdominal ventrite 5.




Figure 40. *Isomira angusta* (Casey) **A** anterior tarsal pubescence **B** male sternite 8. Scale bars: 0.5 mm (**A**), 1.0 mm (**B**).

Bionomic notes. The Canadian specimens were collected with a UV light trap in a meadow patch in deciduous forest.

Comments. This species was originally described as the only member of the new genus *Tedinus* by Casey (1891). *Tedinus* was included as valid in the key to the genera of Alleculini by Aalbu et al. (2002) where it was separated from species of *Isomira* Mulsant, 1856 based on the characters listed above. In addition to the new Canadian record, three new U.S. state records were found among the DNA barcoded specimens:

FLORIDA: Destin, 25-Mar-1980 (1 ex, CNC). OKLAHOMA: Willis, 15-Apr-2009 (6 exx, CBG & CNC); Willis, 18-Apr-2009 (1 ex, CNC). ILLINOIS: Pine Hills Field Station, 22-May-1967 (1 ex, CNC).

Chrysomelidae

Galerucinae

Alticini

Chaetocnema hortensis (Geoffroy, 1785)

BOLD:AAM7650

Figure 41

Distribution. Native to the Palearctic region, widespread across the region and common in many parts (Döberl 2010; Konstantinov et al. 2011). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. BRITISH COLUMBIA: Kelowna, 22-Sep-2014 to 03-Oct-2014 (2 exx, CBG); Revelstoke, 21-Sep-2015 to 02-Oct-2015 (2 exx, CBG). ONTARIO: Brampton, 19-Sep-2016 to 30-Sep-2016 (2 exx, CBG); Mississauga, 24-May-2016 to 26-May-2016 (1 ex, CBG); Mississauga, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG). NOVA SCOTIA: Cape Breton Highlands National Park, 23-Jun-2013 to 29-Jun-2013 (1 ex, CBG); Elmwood, 01-Nov-2005 (1 ex, CBG); Kejimikujik National Park, 31-Jul-2009 (1 ex, CBG); Truro, 21-Sep-2015 to 02-Oct-2015 (2 exx, CBG). LABRADOR: Happy Valley-Goose Bay, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG). NEWFOUNDLAND: Terra Nova National Park, 04-Jul-2009 (1 ex, CBG).

Diagnostic information (based on Konstantinov et al. 2011). Body length (excluding head) 1.8–2.1 mm. Habitus as in Fig. 41A, B. Pronotum and elytra with a bronze or green metallic lustre. Four basal antennomeres yellow, antennomere 2 sometimes partly brown, femora brown, tibiae yellow. Pronotal punctures separated by approximately their own diameter. The two innermost elytral rows of punctures on basal half confused, third through fifth rows confused or regular, sixth row confused. Elytral humeral callus well developed. Aedeagus as in Fig. 41C, D.

Bionomic notes. *Chaetocnema hortensis* has a wide range of host plants. It mainly feeds on various grasses (Poaceae), including cereal crop species (Koch 1992; Konstantinov et al. 2011). It has been recorded as a minor pest of wheat and barley in Europe (Pavlov 1960' Vappula 1965). Most of the barcoded Canadian specimens were collected with Malaise traps in suburban environments. A few records are from grassland and forest habitats in Canadian national parks.

Comments. *Chaetocnema hortensis* has previously been confused with *C. borealis* R. White, 1996 in Canada. We found that most Canadian specimens in CNC identified as *C. borealis* actually represent *C. hortensis*. The elytral punctuation of the two species is similarly irregular basally. In *C. borealis*, the basal antennomeres are brown rather than pale yellow, and the dorsal surface has a blue rather than bronze or green lustre. The aedeagus is differently shaped in the two species (Fig. 41D, E).




Figure 41. *Chaetocnema hortensis* (Geoffroy) **A** male habitus, L. Borowiec **B** female habitus, L. Borowiec **C** *C. hortensis*, aedeagus, ventral view **D** *C. hortensis*, aedeagus, lateral view **E** *C. borealis* R. White, aedeagus, lateral view. Scale bars: 1.0 mm (**A, B**), 0.5 mm (**C–E**).

Based on comparison of the type specimens of *C. borealis* (deposited in CNC) with the diagnoses and figures in the recent revision of Palaearctic *Chaetocnema* species (Konstantinov et al. 2011), *C. borealis* is very similar to (and possibly synonymous with) the Palaearctic *C. sahlbergii* (Gyllenhal, 1827). Both species inhabit bogs and other types of wetlands (Koch 1992; White 1996). Records of *C. borealis* from agricultural fields and other drier habitats reported e.g., by Majka and LeSage (2010) probably represent *C. hortensis*.

***Longitarsus lewisii* Baly, 1874**

BOLD:ACI5614

Figure 42

Distribution. Native to the Palearctic region. Widespread in Europe, recorded throughout Eurasia to China and the Russian Far East (Döberl 2010). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Cornwall, 19-Sep-2016 to 30-Sep-2016 (5 exx, CBG).

Diagnostic information (based on Warchalowski 1996 and Rutanen and Martikainen 2014). Body length 1.7–2.3 mm. Habitus as in Fig. 42A, B, convex in dorsal view, sides of elytra rounded. Head brown, pronotum and elytra yellow-brown, elytral suture usually narrowly dark at least near midlength, legs pale, metafemora darker. Ventral side red-brown to black. Pronotum ca. 1.5 times wider than long, finely punctate. Elytra densely and finely punctate, punctures slightly larger around scutellum. Male with a narrow longitudinal impression at the middle of the last ventrite, ending in a small, sharply delimited round pit. Last ventrite of females unmodified or with a very weak impression. Penis in lateral view strongly bent towards dorsum at the apex (Fig. 42C, D).

Bionomic notes. This species feeds on *Plantago* species, especially *P. major* L. (Koch 1992, Rutanen and Martikainen 2014). In Finland, it is most often collected in dry, barren habitats (Rutanen and Martikainen 2014). The Canadian specimens were collected with a Malaise trap in a suburban residential area.

Comments. *Longitarsus lewisii* is closely related to *L. pratensis* (Panzer, 1794), another adventive species from the Palearctic region (Warchalowski 1996, Rutanen and Martikainen 2014). *Longitarsus lewisii* is more rounded and convex, and on average slightly larger than *L. pratensis* (1.7–2.3 mm vs. 1.4–2.1 mm) (Warchalowski 1996). The elytral suture is not darkened in *L. pratensis*, and the hind femora are paler. However, color is variable in this species group, and the male genitalia and the modifications of the last ventrite are the best distinguishing characters. In males of *L. pratensis*, the impression of the last ventrite is broad, circular and less sharply delimited than in males of *L. lewisii*. The penis of *L. pratensis* is shorter than that of *L. lewisii*, and less strongly bent. Females of *L. pratensis* have an elongate-oval, shallow medial impression on the last ventrite. The preferred host plant of *L. pratensis* is *Plantago lanceolata* L., but both species use several species of *Plantago* (Koch 1992; Döberl 1994; Rutanen and Martikainen 2014).

***Lythreria salicariae* (Paykull, 1800)**

BOLD:AAO3219

Figure 43

Distribution. Native to the Palearctic region. Widespread in Europe, scattered records in Asia to East Siberia and Japan (Döberl 2010). Adventive in the Nearctic region (Ontario, Canada).




Figure 42. *Longitarsus lewisii* Baly **A** male habitus, L. Borowiec **B** female habitus, L. Borowiec **C** aedeagus, ventral view **D** aedeagus, lateral view. Scale bars: 1.0 mm (**A, B**), 0.5 mm (**C, D**).

Canadian records. ONTARIO: Cambridge, 25-May-2015 to 31-May-2015 (1 ex, CBG); Pickering, 24-Jun-2017 to 25-Jun-2017 (1 ex, CBG).

Diagnostic information (based on Mohr 1966). Body length 1.8–2.3 mm. Habitus elongate-oval (Fig. 43A, B). Yellow-brown or red-brown, apical antennomeres and ventral side darkened, sometimes also head, pronotum, and elytral suture darker brown. Base of pronotum without lateral furrows or a transverse impression. Procoxal cavities closed behind. Elytral punctures arranged in regular striae. Metatibia without a subapical dilation or tooth on the outer margin.




Figure 43. *Lythroria salicariae* (Paykull), habitus, L. Borowiec **A** male **B** female. Scale bar: 1.0 mm.

Bionomic notes. *Lythroria salicariae* is found in various wetland and marshy shoreline habitats as well as in forest depressions (Koch 1992). The larvae develop on *Lysimachia* species, and the adults occasionally feed also on *Lythrum salicaria* L. (Koch 1992, Dolgovskaya et al. 2004). The Canadian specimens were collected with pan traps in a grassy wetland and a mixed habitat of agricultural fields and forest.

Comments. *Lythroria* Bedel, 1897 is a monotypic genus reported here for the first time from North America. *Lythroria salicariae* would be identified as *Pseudorthygia* Csiki, 1940 (couplet 75) using the key to genera of Galerucinae in Riley et al. (2002) based on its closed procoxal cavities, but *L. salicariae* is not as convex in lateral profile and has a more elongate body outline. Among previously recorded Canadian leaf beetles, the habitus of *L. salicariae* is somewhat similar to *Glyptina brunnea* Horn, 1889, but the procoxal cavities are open behind in *Glyptina*.

Galerucinae

Luperini

Scelolyperus liriophilus Wilcox, 1965

BOLD:ABW1434

Distribution. Native to the Nearctic region. Widespread in eastern United States (Clark 1996).

Canadian records. QUEBEC: Forillon National Park, 05-Jul-2013 to 15-Jul-2013 (2 exx, CBG).

Diagnostic information. See Clark (1996).

Bionomic notes. This species has been collected from a wide variety of plant species (Clark 1996). The Canadian specimens were collected with a Malaise trap along a forest trail in Forillon National Park.

Curculionidae

Brachycerinae

Eriirhinini

Notaris scirpi (Fabricius, 1793)

BOLD:AAX5634


Figure 44

Distribution. Native to the Palearctic region. Widespread in Europe, with scattered records in Asia to the Russian Far East and Japan (Alonso-Zarazaga et al. 2017). Adventive in the Nearctic region (Quebec, Canada).

Canadian records (DNA barcoded specimen). QUEBEC: Laval, 11-Jun-1997 (1 ex, CNC).

Additional Canadian records. QUEBEC: Gatineau, 25-May-2012 (1 ex, CPTO); Henryville, 14-Jun-2015 (1 ex, CNC); Henryville, 07-Jun-2017 (1 ex, CMNC); Henryville, 07-Jun-2017 (9 exx, CCCH); Henryville, 14-Jun-2017 (1 ex, CMNC); Laval, 5-Jun-2004 (2 exx, CSDU); Laval, 19-Apr-2013 (1 ex, CSDU); Laval, 22-Jun-2013 (1 ex, CSDU); Longueuil, 21-May-2016 (1 ex, CPTO); Oka, 01-Jul-2004 (1 ex, CRVI); Oka, 06-Jun-2011 (1 ex, CRVI); Oka, 19-Aug-2012 (1 ex, CMNC); Oka, 26-Aug-2012 (1 ex, CPTO); Oka, 01-Jun-2014 (1 ex, CRVI); Oka, 21-Jun-2016 (1 ex, CRVI); Oka, 13-Jul-2016 to 20-Jul-2016 (2 exx, CRVI); Oka, 22-Jul-2016 (1 ex, CRVI); Oka, 15-May-2018 to 31-May-2018 (1 ex, CRVI); Saint-Côme, 13-Jul-2013 (1 ex, CSDU); Saint-Lazare, 20-Jun-2012 (1 ex, CPTO); Saint-Lazare, 17-Sep-2012 (1 ex, CPTO); Saint-Lazare, 14-Jun-2013 (1 ex, CPTO); Saint-Lazare, 18-Jul-2017 (1 ex, CSDU); Terrasse-Vaudreuil, 01-Jun-2013 (1 ex, CNC); Terrasse-Vaudreuil, 11-Jun-2007 (1 ex, CPTO); Terrasse-Vaudreuil, 30-Jun-2014 (1 ex, CPTO); Varennes, 07-Jun-2011 (1 ex, CCCH).

Diagnostic information (based on Hoffmann 1958). Body length: 4.7–7.0 mm. Habitus as in Fig. 44. Oblong-oval, black or brown, dorsal pubescence of small pili-form scales more or less regularly distributed, with a speckled color pattern formed by patches of paler scales. Rostrum elongate, narrow, curved, punctate-striate and carinate. Prothorax approximately as long as wide, sides rounded, punctation dense and deep, with median line slightly elevated anteriorly. Elytra rounded at humeri in dorsal view, sides subparallel until slightly beyond middle. Ventrally with lateral portions of abdomen, metanepisternum, metanepimeron, and lateral portion of metaventricle with dense cream-colored scales.



Figures 44, 45. **44** *Notaris scirpi* (Fabricius), habitus, L. Borowiec **45** *Centrinopus helvinus* Casey, habitus. Scale bar: 2.0 mm (**44**), 1.0 mm (**45**).

Bionomic notes. *Notaris scirpi* is oligophagous on *Scirpus* and *Carex* species in wet habitats (Koch 1992). Hoffmann (1958) notes that in France the species develops in the collar of *Carex acutiformis* Ehrh. and that adults can be collected in litter around wet areas.

Comments. These are the first records of *Notaris scirpi* from the Nearctic region. After the identification of the DNA barcoded specimen deposited in CNC, 37 additional specimens from various localities in Quebec were found in other collections. The earliest record is from 1997, and the species seems to be firmly established in Quebec. *Notaris scirpi* is easily distinguished from *Tournotaris bimaculatus* (Fabricius, 1787) and *Notaris puncticollis* (LeConte, 1876), the two most similar species already known from North America, by the dense cream-colored scales on the lateral portions of the abdomen, metanepisternum, metanepimeron, and lateral portion of the metaventricle.

Baridinae**Madarini*****Ampelogypter sesostris* (LeConte, 1876)**

BOLD:ADH7970

Distribution. Native to the Nearctic region. Previously recorded from Indiana, Michigan, Ohio, Pennsylvania, Illinois, Florida, and Missouri (O'Brien and Wibmer 1982; Saunders and Tobin 2000) but likely more widespread in eastern and midwestern United States (Riedl and Taschenberg 1984).

Canadian records (DNA barcoded specimens). ONTARIO: Pelee Island, 06-Jun-1982 (1 ex, CNC); Rouge National Urban Park, 25-Jun-2017 (1 ex, CBG).

Additional Canadian records. ONTARIO: Pelee Island, 26-Jun-1940 (1 ex, CNC); Pelee Island, 27-Jun-1940 (1 ex, CNC); Windsor, 30-May-2002 (1 ex, CMNC).

Diagnostic information (based on Blatchley and Leng 1916 and Anderson 2002). Body length: 2.7–3.0 mm. Body glabrous, shiny, elongate-oval, pale reddish brown throughout. Elytral interstriae flat. Femora not toothed. Tarsus with two claws connate at base.

Bionomic notes. This species feeds on *Vitis* L. species, and it is considered a minor pest in vineyards (Bouchard et al. 2005). The female oviposits above a stem node and hollows out additional cavities along the longitudinal axis of shoots of the host plants. The larva develops and feeds on tissues inside the shoot, causing it to swell and thereby inducing gall formation (Lasnier et al. 2019).

Comments. The red-brown *Ampelogypter sesostris*, known commonly as the grape cane gallmaker, can be separated from the other two species in this genus in the United States and Canada by color: *A. ampelopsis* (Riley, 1869) and *A. longipennis* Casey, 1892 have a black integument.

Baridinae**Apostasimerini*****Centrinopus helvinus* Casey, 1892**

BOLD:ACX7906

Figure 45

Distribution. Native to the Nearctic and Neotropical regions. Recorded from eastern and north central United States, and southward to Nicaragua (O'Brien and Wibmer 1982).

Canadian records. ONTARIO: Waterloo, 21-Sep-2015 to 02-Oct-2015 (1 ex, CBG); Waterloo, 19-Sep-2016 to 30-Sep-2016 (1 ex, CBG).

Diagnostic information (based on Casey 1920 and Anderson 2002). Body length: 2.0–2.7 mm. Body dark red-brown to black, oval, covered dorsally with pale scales, oriented perpendicularly to body axis on pronotum, oriented longitudinally on elytra

(Fig. 45). Scales somewhat denser on elytral interstriae 3, 5, 7. Prothorax only slightly narrower than elytra in dorsal view. Scutellum densely covered with scales. Female with sharply defined longitudinal sulcus anterior to procoxae. Each procoxa in male with one anteriorly projecting spine-like process in front.

Bionomic notes. Kissinger (1964) mentioned that adults in this genus are found on flowers of Asteraceae. According to Blatchley and Leng (1916) *Centrinopus helvinus* was taken on sweetscented joe pye weed, *Eutrochium purpureum* (L.) E.E. Lamont. We are not aware of any additional biological information published on this species. The barcoded Canadian specimens were collected with a Malaise trap on farmland.

Comments. The genus *Centrinopus* Casey, 1892, which is in need of a taxonomic revision (Anderson 2002), contains six species in the eastern United States and is recorded here from Canada for the first time.

Ceutorhynchinae

Ceutorhynchini

Ceutorhynchus inaeffectatus Gyllenhal, 1837

BOLD:ACB0098

Figure 46


Distribution. Native to the Palearctic region. Widespread in Europe, recorded east to Kazakhstan and West Siberia (Alonso-Zarazaga et al. 2017). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Guelph, 02-Jun-2018 (1 ex, CBG).

Diagnostic information (based on Lohse 1983). Body length 2.3–3.9 mm. Habitus as in Fig. 46A, appearing grey at low magnification due to the pale scales sparsely covering the black integument. Antennal funicle with seven antennomeres. Pronotum densely punctate, lateral tubercles absent. Elytra without apical calli, interstriae with narrow scales arranged in two or three longitudinal rows. Meso- and metafemora with small teeth. All tarsal claws with small basal tooth. Aedeagus as in Fig. 46B.

Bionomic notes. This species feeds on *Hesperis matronalis* L. and *H. tristis* L. (Brassicaceae) (Koch 1992). The larvae develop in the seed pods, the adults feed on leaves and other parts of the host plants (Koch 1992, Larsen et al. 1992). The Canadian specimen was collected by sweep netting vegetation along a recreational trail where *H. matronalis* is abundant.

Comments. *Ceutorhynchus inaeffectatus* is similar in habitus to two other Palearctic species adventive in North America: *C. obstrictus* (Marsham, 1802) and *C. rapae* Gyllenhal, 1837. These species have lateral pronotal tubercles, which are absent in *C. inaeffectatus*. The combination of toothed femora, antennal funicle with seven antennomeres, pronotum lacking lateral tubercles and toothed tarsal claws will separate *C. inaeffectatus* from native species of *Ceutorhynchus*. *Hesperis matrona-*



Figures 46–48. **46** *Ceutorhynchus inaeffectatus* Gyllenhal **46A** habitus, L. Borowiec **46B** aedeagus **47** *Ceutorhynchus mutabilis* Dietz, habitus **48** *Exomias trichopterus* (Gautier des Cottés) **48A** male habitus, L. Borowiec **48B** female habitus, L. Borowiec. Scale bars: 1.0 mm (**46A**; **47**; **48**), 0.5 mm (**46B**).

lis (dame's rocket or purple rocket) is an invasive weed in North America (Francis et al. 2009), and the strictly specialized *C. inaeffectatus* could potentially be useful in its biological control.

***Ceutorhynchus mutabilis* Dietz, 1896**

BOLD:AAZ4085

Figure 47

Distribution. Native to the Nearctic region. This species is reported from Baja California, California, Oregon, Washington, Colorado, Kansas, and North Dakota in the United States (O'Brien and Wibmer 1982; Balsbaugh and Aarhus 1990).

Canadian records. BRITISH COLUMBIA: Radium, 24-Aug-1982 (2 exx, CNC); New Afton Mine, 20-Jun-2013 to 27-Jun-2013 (1 ex, CBG). ALBERTA: Calgary, 22-Jul-1976 (1 ex, CNC); Exact locality not specified, 20-Jun-1985 (1 ex, CNC); Exact locality not specified, 09-Jun-1990 (1 ex, CNC). SASKATCHEWAN: Grasslands National Park, 19-Jul-2012 to 26-Jul-2012 (1 ex, CBG). MANITOBA: Exact locality not specified, 24-Jul-1995 (1 ex, CNC).

Diagnostic information (based on Scheibner 1963). Body length: 2.4 mm. Habitus as in Fig. 47, body with black integument covered with white to pale brown scales. Antennal funicle with seven antennomeres. Combination of two types of scales on pronotum and elytra, some broadly oval, others expanding from base with a truncate apex. Elytra with dense patch of appressed oval scales posterior to scutellum. Metafemora lacking tooth. Tarsal claws each with a small basal tooth.

Bionomic notes. The natural history and host preferences of this little-studied species are unknown (Colonnelli 2004).

Comments. Although this genus is in need of a revision, the combination of character states listed above, in combination with the habitus photograph (Fig. 47), should lead to the correct identification. Studies describing the biology of this and other native species of *Ceutorhynchus* Germar, 1824 are badly needed.

***Peracalles pectoralis* (LeConte, 1876)**

BOLD:ACY2911

Distribution. Native to the Nearctic region. Anderson (2002) reports this species from Illinois, Indiana, Ohio, Kentucky, and Missouri in the United States.

Canadian records. ONTARIO: Point Pelee National Park, 06-Jul-2015 (1 ex, CBG); Point Pelee National Park, 16-Jun-2014 to 22-Jun-2014 (3 exx, CBG).

Diagnostic information (based on Anderson 2002). Body length: 3.0–3.3 mm. Body black, covered with broad, flat, appressed dark brown to pale scales, broadly oval in dorsal view. Antennal funicle with seven articles. Prosternum with deep longitudinal sulcus for reception of rostrum, sulcus extending posteriorly to anterior edge of mesoventrite. Elytra strongly convex in lateral view, with a single row of flat, apically truncate, erect scales on each interstria.

Bionomic notes. Adults in this genus occur in leaf litter (Anderson 2002). The Canadian specimens were collected from a marsh and a swampy forest using pan traps, pitfall traps and Berlese funnel extraction.

Comments. The genus *Peracalles* Kissinger, 1964 contains two species in the United States (Anderson 2002) and is recorded here in Canada for the first time.

Entiminae Sciaphilini

Exomias trichopterus (Gautier des Cottes, 1863)

BOLD:ACZ1179

Figure 48

Distribution. Native to the Palearctic region. Widespread in Central Europe (Alonso-Zarazaga et al. 2017). Adventive in the Nearctic region (Ontario, Canada).

Canadian records. ONTARIO: Rouge National Urban Park, 24-Jun-2017 to 25-Jun-2017 (1 ex, CBG).

Diagnostic information (based on Rheinheimer and Hassler 2013). Body length: 2.7–3.4 mm. Habitus as in Fig. 48A, B. Body brown to black, covered with fine semi-erect to erect setae, legs pale yellow to red-brown. Rostrum with shallow longitudinal depression dorsally. Globose elytra with humeral angles obsolete. Elytra lacking row of long setae near suture on posterior half.

Bionomic notes. This common European species is polyphagous on herbaceous plants (Balalaikins 2011) and could become a new pest of berry crops in Canada (see Kolov and Korotyaev 2017).

Comments. *Exomias trichopterus* is very similar in appearance to *E. pellucidus pellucidus* (Boheman, 1834), another adventive Palearctic species which is common and widespread in North America. Both species were previously placed in the genus *Barypeithes* Jacquelin du Val, 1854. The former subgenus *Exomias* was elevated to the generic level by Borovec (2013). *Exomias pellucidus pellucidus* can be diagnosed primarily by the noticeably denser setae on their elytra, especially near the apex where an additional row of long setae is present along the elytral suture.

Scolytinae Xyleborini

Ambrosiodmus rubricollis (Eichhoff, 1875)

BOLD:AAR3532

Distribution. Native to the eastern Palearctic and Oriental regions (Knížek 2011). Adventive in Europe, Australia, and the Nearctic region (widespread in the United States; Ontario, Canada) (Knížek 2011; Gomez et al. 2018).

Canadian records. ONTARIO: Point Pelee National Park, 11-Jul-2012 to 18-Jul-2012 (1 ex, CBG); Point Pelee National Park, 16-Jun-2014 to 22-Jun-2014 (2 exx, CBG).

Diagnostic information (based on Gomez et al. 2018). Body length 2.4–2.6 mm. Pronotum with asperities covering entire surface. Elytral declivity with tubercles on interstriae 2 as large as those on interstriae 1 and 3.

Bionomic notes. *Ambrosiodmus rubricollis* uses symbiotic fungi to attack many genera of gymnosperm and dicot trees including species in the following Canadian genera: *Abies* Mill., *Aesculus* L., *Alnus* Mill., *Carya* Nutt., *Cornus* L., *Fraxinus* L., *Ilex* L., *Juglans* L., *Morus* L., *Pinus* L., *Populus* L., *Prunus* L., *Quercus* L., *Rhus* L. (Faccoli et al. 2009). One of the Canadian specimens was collected with a Malaise trap in a savanna, the two others were caught with pitfall traps in a swampy forest.

Comments. This is the only *Ambrosiodmus* species known from Canada, although two larger-bodied species are known from states bordering southern Ontario (Gomez et al. 2018). *Ambrosiodmus lewisi* (Blandford, 1984), and *A. tachygraphus* (Zimmermann, 1868) can be distinguished from *A. rubricollis* by their greater body lengths (3.6 to 4.0 mm).

Discussion

This study adds 60 new species to the Canadian beetle fauna and resolves taxonomic confusion in another three species. Among the 42 adventive species covered, 40 are native to the Palearctic region. The remaining two species, *Clambus simsoni* and *Attagenus smirnovi*, are native to the Australian and Afrotropical regions respectively, but also occur in the Palearctic as adventive species. *Nephus bisignatus* and *Dichelotarsus lapponicus* were previously known only from the Palearctic region, but because they were collected in remote arctic localities in Canada, we consider it likely that they are Holarctic taxa whose occurrence in North America was previously overlooked. The remaining 19 new species are native to North America, and represent either previously overlooked occurrences in Canada, or recent range expansions. The fact that many new records of native species were of species that are difficult to identify by morphological methods suggests that most of these species have been long present in Canada but overlooked. Six species were found at Point Pelee, a forest and wetland area isolated from similar habitats in both United States and Canada, further suggesting that recent range extensions are an unlikely explanation for new Canadian records of these species. The fact that 54 of the 60 new species for Canada were found in general survey samples for insects clearly indicates that much more work is needed using specialized, taxon-specific collecting techniques to achieve a full inventory of the Coleoptera diversity in Canada. We also expect that increased insect survey activity in United States would recover records for many of the same adventive species there, plus additional species as-yet unrecorded from North America.

Species that are adapted to disturbed or ruderal habitats are more likely to be accidentally transported through human activities than species that require non-synanthropic habitats (Lockwood et al. 2013). Many of the adventive beetle species established in North America are strongly synanthropic and occur mainly in human-

disturbed habitats and settlements (Klimaszewski et al. 2010, 2012). Not surprisingly, most of the new adventive species we report here were found mainly or exclusively in Southern Ontario, which is Canada's most densely populated and biodiverse region, and in the Greater Vancouver area, the third largest metropolitan area in Canada and home to the busiest port in the country (also with high native insect diversity). Some of these adventive species have likely been present in Canada for a long time, but have been overlooked due to difficulties in morphological identification. *Stenichnus scutellaris* and *Amischa decipiens* are widespread and common in southern Ontario, while *A. decipiens* is also found in the Greater Vancouver area. *Malthodes pumilus* occurs from East to West in both suburban and natural environments. All three species represent genera that have received little or no taxonomic investigation in North America in recent decades. Others, such as *Calyptromerus dubius*, *Clambus simsoni*, *Litargus connexus*, and *Olibrus liquidus*, may have arrived more recently as they have only been found in one or a few urban localities, and some are only represented by singleton specimens in the Canadian DNA barcode data. Verifying that these species are well-established in Canada will require further monitoring and study of material in existing collections.

Most of the species newly recorded here that are shared between Europe and North America probably arrived into North America from Europe because they were discovered there first. Relatively few North American beetle species are known to occur as adventive in Europe, but more may well be uncovered especially in families where the Nearctic fauna is poorly known. Adventive insect species are sometimes described as new to science after arriving in a new area (Wheeler and Hoebeke 2009), as exemplified by the two new synonymies in Staphylinidae we establish in this study. New species are less likely to escape notice in Europe where the beetle fauna, including taxonomically challenging families such as Staphylinidae, is generally better known and more intensively studied compared to North America. Our analysis of European and Canadian DNA barcode data has uncovered at least one native North American species of Staphylinidae occurring as adventive in Europe and described as new from there. This synonymy will be formally established in a future publication. A geographically well represented DNA barcode dataset can provide information on the biogeography and distributional status (native vs. adventive) of species, and potentially identify the geographic origin of adventive or expansive species (Valade et al. 2009; Lees et al. 2011). A detailed analysis of the spatial genetic variation in all the species covered here is beyond the scope of the present paper but will be a subject of future studies.

It is noteworthy that 57 of our 60 new species records for Canada were discovered, in whole or in part, using material recently collected by the Centre for Biodiversity Genomics. In fact, only two of the new species were discovered based solely on specimens from the CNC (see Table 1). The CNC was once the primary source of new data on Canadian insect species, but it is no longer the depository for most specimens from general survey and inventory work across the country. For example, the 1,085,146 Canadian insect specimens analyzed by Hebert et al. (2016) are stored in the CBG voucher archive. To further illustrate this change in specimen accumulation, a complete inventory of the Canadian Scarabaeoidea in the CNC (ABTS, unpublished data)

revealed that only 7% of the specimens in that institution were collected during the past 30 years. In contrast, 37% of the specimens in CNC were collected during the previous 30-year period (1959–1988). Inventories of Canadian Scarabaeoidea in most of the major entomological collections across the country show a similar overall pattern. Although general survey and inventory work is badly needed in Canada to detect the full diversity of the Coleoptera fauna, collecting efforts have significantly declined over the past 30 years. This leaves invasive species undetected and range expansions undiscovered for years longer than would have been the case when there were large-scale survey and inventory efforts (e.g., Northern Insect Survey; Lonsdale and Huber 2011). With the growing threat of invasive species through increasing global trade and the northward expansion of species due to the changing climate, ongoing collaborative survey and inventory efforts are needed to detect new species as they appear in Canada.

Bousquet et al. (2013) recorded 8237 Coleoptera species in Canada, an increase of almost 10% over a similar checklist published 22 years earlier (Bousquet 1991). The recent summary by Brunke et al. (2019) increased the number of Canadian beetle species to 8302. These increases are mainly due to progress in taxonomic research on Coleoptera species already present in Canada, but species recently establishing themselves in Canada also increased the count. The 60 species we report here as new for Canada increase the number of known beetle species in the country by another 0.7% compared to Brunke et al. (2019). Of these, the 40 new adventive species add 6.3% to the number of non-native Coleoptera known from Canada. Further new Canadian records and new synonymies in European and North American Coleoptera detected through DNA barcode data are currently being validated. Our study shows that DNA barcoding, combined with morphological validation of the voucher specimens, is a powerful tool for detecting and identifying overlooked or recently arrived species, both native and adventive (see also deWaard et al. 2009, Landry et al. 2013). Even though the species coverage of the European and Canadian DNA barcode reference libraries of beetles is still far from complete, our results undeniably demonstrate their usefulness for cataloguing regional biodiversity.

Acknowledgements

Josie Smith (Canadian Food Inspection Agency) provided the additional specimen of *Pseudanostirus tigrinus*. Collecting permits granted by the City of Guelph (Martin Neumann, Parks Operations and Forestry) to M. Pentinsaari led to the discovery of *Cis boleti* and *Medon apicalis*. Marko Mutanen (University of Oulu, Finland) provided European material of several species for comparison. Serge Laplante (Agriculture and Agri-Food Canada, Ottawa) helped in validating some of the identifications. Volker Assing (Hannover, Germany), Sami Karjalainen (Kirkkonummi, Finland), Matúš Kocian (Prague, Czech Republic), Harald Schillhammer (Natural History Museum Vienna, Austria), and Michael Schülke (Berlin, Germany) kindly allowed us to use their images in this publication. Anthony Davies (Agriculture and Agri-Food Canada, Ottawa) imaged many of the specimens and compiled the figure plates. Steve Paiero provided access to the University of Guelph Insect Collection. Claude Chantal (Varenes, Quebec), Stéphane Dumont (Montreal,

Quebec), Pierre de Tonnancour (Terrasse-Vaudreuil, Quebec), and Robert Vigneault (Oka, Quebec) provided additional specimen records of *Notaris scirpi* from their private collections. Reginald Webster (Charters Settlement, New Brunswick) and György Makranczy (Hungary) provided additional verified specimen records of *Carpelimus elongatulus*. All of the sequence analysis and processing of voucher specimens, as well as the CBG's collection program, were supported by grants from the Ontario Ministry of Research and Innovation, and from Genome Canada through Ontario Genomics in support of the International Barcode of Life (iBOL) project. Subsequent work to identify specimens and analyze data was enabled by the Canada First Research Excellence Fund through its support for the Food From Thought project at the University of Guelph. Michael Caterino and two anonymous reviewers provided helpful comments on a previous version of this manuscript.

References

- Aalbu RL, Triplehorn CA, Campbell JM, Brown KW, Somerby RE, Thomas DB (2002) Tenebrionidae Latreille, 1802. In: Arnett RH, Thomas MC, Skelley PE, Frank JH (Eds) American Beetles (Vol. 2). Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, 463–509. <https://doi.org/10.1201/9781420041231.ch7>
- Alonso-Zarazaga MA, Barrios H, Borovec R, Bouchard P, Caldara R, Colonnelli E, Gültekin L, Hlaváč P, Korotyaev B, Lyal CHC, Machado A, Meregalli M, Pierotti H, Ren L, Sánchez-Ruiz M, Sforzi A, Silfverberg H, Skuhrovec J, Trýzna M, Velázquez de Castro AJ, Yunakov NN (2017) Cooperative Catalogue of Palaearctic Coleoptera Curculionoidea. Monografias electrónicas SEA 8. Sociedad Entomológica Aragonesa S.E.A., Zaragoza, 729 pp. <http://graellsia.revistas.csic.es/index.php/graellsia/article/view/537/636>
- Anderson RS (2002) Curculionidae Latreille, 1802. In: Arnett RH, Thomas MC, Skelley PE, Frank JH (Eds) American Beetles (Vol. 2). Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, 722–815.
- Arango RA, Young DK (2012) Death-watch and spider beetles of Wisconsin – Coleoptera: Ptinidae. General Technical Report FPL-GTR-209. US Department of Agriculture, Forest Service, Forest Products Laboratory, Madison. <https://doi.org/10.2737/FPL-GTR-209>
- Armstrong K (2010) DNA barcoding: a new module in New Zealand's plant biosecurity diagnostic toolbox. EPPO Bulletin 40: 91–100. <https://doi.org/10.1111/j.1365-2338.2009.02358.x>
- Armstrong KF, Ball SL (2005) DNA barcodes for biosecurity: invasive species identification. Philosophical Transactions of the Royal Society of London. Series B, Biological sciences 360: 1813–1823. <https://doi.org/10.1098/rstb.2005.1713>
- Assing V (2004) A Revision of the *Medon* Species of the Eastern Mediterranean and Adjacent Regions (Insecta: Coleoptera: Staphylinidae: Paederinae). Bonner zoologische Beiträge 52: 33–82.
- Assing V (2006) A Revision of the Western Palaearctic *Medon*: the Species of the Atlantic Islands, the Western Mediterranean, and Europe, Except for the Southeast. Bonner zoologische Beiträge 54: 25–95.
- Assing V (2009) On the *Pseudomedon* species of the Palaearctic region (Coleoptera: Staphylinidae: Paederinae). Linzer biologische Beiträge 41: 1175–1189.

- Assing V (2012) Paederinae. In: Assing V, Schülke M (Eds) Die Käfer Mitteleuropas, Band 4 (2. Auflage). Spektrum Akademischer Verlag, Heidelberg, 380–383.
- Assing V (2018) On some *Myllaena* species in the East Mediterranean and Caucasus regions (Coleoptera: Staphylinidae: Aleocharinae). Linzer biologische Beiträge 50: 1015–1032.
- Audisio P (1993) Fauna d'Italia (Vol. 32): Coleoptera Nitidulidae – Kateretidae. Edizione Calderini, Bologna, 971 pp.
- Balalaikins M (2011) On Latvian Entiminae (Coleoptera: Curculionidae): 2. Tribes Trachyploeini Lacordaire, 1863 and Sciaphilini Sharp, 1891. Acta Zoologica Lithuanica 21: 253–262. <https://doi.org/10.2478/v10043-011-0031-1>
- Balsbaugh EU, Aarhus DG (1990) Checklist and new state records of Curculionidae (broad sense) (Coleoptera) for North Dakota. Journal of the Kansas Entomological Society 63: 227–236. <https://www.jstor.org/stable/25085171>
- Baranowski R (1993) Revision of the genus *Leiodes* Latreille of North and Central America. Entomologica Scandinavica Supplementum 41: 1–149.
- Bellard C, Cassey P, Blackburn TM (2016) Alien species as a driver of recent extinctions. Biology letters 12: 20150623. <https://doi.org/10.1098/rsbl.2015.0623>
- Besuchet C (1955) Monographie des *Biblopectus* et *Pseudoplectus* paléarctiques. Bulletin de la Société Entomologique Suisse 28: 153–209.
- Bieńkowski AO (2018) Key for identification of the ladybirds (Coleoptera: Coccinellidae) of European Russia and the Russian Caucasus (native and alien species). Zootaxa 4472: 233–260. [/doi.org/10.11646/zootaxa.4472.2.2](https://doi.org/10.11646/zootaxa.4472.2.2)
- Blanchard F (1917) Revision of the Throscidae of North America (Coleoptera). Transactions of the American Entomological Society 43: 1–26.
- Blatchley WS, Leng CW (1916) Rhynchophora or Weevils of North Eastern America. The Nature Publishing Company, Indianapolis, 682 pp. <https://doi.org/10.5962/bhl.title.1557>
- Borovec R (2013) Sciaphilini. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 8). Brill, Leiden, 83, 377–386.
- Bouchard P, Lesage L, Goulet H, Bostanian NJ, Vincent C, Zmudzinska A, Lasnier J (2005) Weevil (Coleoptera: Curculionoidea) diversity and abundance in two Quebec vineyards. Annals of the Entomological Society of America 98: 565–574. [https://doi.org/10.1603/0013-8746\(2005\)098\[0565:WCCDAA\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2005)098[0565:WCCDAA]2.0.CO;2)
- Bouchard P, Smith ABT, Douglas H, Gimmel ML, Brunke AJ, Kanda K (2017) Biodiversity of Coleoptera. In: Footitt RG, Adler PH (Eds) Insect Biodiversity: Science and Society. John Wiley & Sons Ltd, 337–417. <https://doi.org/10.1002/9781444308211.ch11>
- Bousquet Y (1991) Checklist of beetles of Canada and Alaska. Agriculture Canada, Ottawa, 430 pp.
- Bousquet Y (2010) Illustrated Identification Guide to Adults and Larvae of Northeastern North American Ground Beetles (Coleoptera, Carabidae). Pensoft Publishers, Sofia, 562 pp.
- Bousquet Y (2012) Catalogue of Geadephaga (Coleoptera: Adephaga) of America, north of Mexico. ZooKeys 245: 1–1722. <https://doi.org/10.3897/zookeys.245.3416>
- Bousquet Y, Bouchard P, Davies A, Sikes D (2013) Checklist of beetles (Coleoptera) of Canada and Alaska. Second edition. ZooKeys 360: 1–44. <https://doi.org/10.3897/zookeys.360.4742>

- Bousquet Y, Thomas DB, Bouchard P, Smith AD, Aalbu RL, Johnston MA, Steiner Jr. WE (2018) Catalogue of Tenebrionidae (Coleoptera) of North America. *ZooKeys* 728: 1–455. <https://doi.org/10.3897/zookeys.728.20602>
- Bowstead S (1999) A revision of the Corylophidae (Coleoptera) of the West Palearctic Region. Muséum d'histoire naturelle, Genève, Geneva, 203 pp.
- Bowstead S (2007) Corylophidae. In: Löbl I, Smetana A (Eds) Catalogue of Palearctic Coleoptera (Vol. 4). Apollo Books, Stenstrup, 631–635.
- Brown WJ (1936) American species of *Ludius*; The *fallax* and *triundulatus* groups. *The Canadian Entomologist* 68: 99–107. <https://doi.org/10.4039/Ent6899-5>
- Brunke AJ, Bahlai CA, Klimaszewski J, Hallett RH (2014) Rove beetles (Coleoptera: Staphylinidae) in Ontario, Canada soybean agroecosystems: assemblage diversity, composition, seasonality, and habitat use. *The Canadian Entomologist* 146: 652–670. <https://doi.org/10.4039/tce.2014.19>
- Brunke AJ, Bouchard P, Douglas HB, Pentinsaari M (2019) Coleoptera of Canada. *ZooKeys* 819: 361–376. <https://doi.org/10.3897/zookeys.819.24724>
- Campbell JM (1979) A revision of the genus *Tachyporus* Gravenhorst (Coleoptera: Staphylinidae) of North America. *Memoirs of the Entomological Society of Canada* 111: 1–95. <https://doi.org/10.4039/entm111109fv>
- Campbell JM (1991) A revision of the genera *Mycetoporus* Mannerheim and *Ischnosoma* Stephens (Coleoptera: Staphylinidae: Tachyporinae) of North and Central America. *Memoirs of the Entomological Society of Canada* 123: 1–169. <https://doi.org/10.4039/entm123156fv>
- Casey TL (1891) Coleopterological notices, III. *Annals of the New York Academy of Sciences* 6: 9–214. <https://doi.org/10.1111/j.1749-6632.1892.tb55403.x>
- Casey TL (1920) *Memoirs on the Coleoptera*. The New Era Printing Company, Lancaster, 529 pp.
- Chandler DS (1990) The Pselaphidae (Coleoptera) of Latimer county, Oklahoma, with revisions of four genera from eastern North America. Part I. Faroninae and Euplectinae. *Transactions of the American Entomological Society* 115: 503–529.
- Clark SM (1996) The genus *Scelolyperus* Crotch in North America (Coleoptera: Chrysomelidae: Galerucinae). *Insecta Mundi* 10: 261–280. <http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1005&context=insectamundi>.
- Colonnelli E (2004) Catalogue of Ceutorhynchinae of the world, with a key to genera (Insecta: Coleoptera: Curculionidae). Argania Editio, Barcelona, 124 pp.
- Daffner H (1983) Revision der paläarktischen Arten der Tribus Leiodini Leach (Coleoptera, Leiodidae). *Folia Entomologica Hungarica* 44: 9–163.
- Denux O, Zagatti P (2010) Coleoptera families other than Cerambycidae, Curculionidae sensu lato, Chrysomelidae sensu lato and Coccinelidae. Chapter 8.5. *BioRisk* 4: 315–406. <https://doi.org/10.3897/biorisk.4.61>
- deWaard JR, Humble LM, Schmidt BC (2010) DNA barcoding identifies the first North American records of the Eurasian moth, *Eupithecia pusillata* (Lepidoptera: Geometridae). *Journal of the Entomological Society of British Columbia* 107: 25–31. <https://journal.entsocbc.ca/index.php/journal/article/view/11>

- deWaard JR, Ivanova NV, Hajibabaei M, Hebert PDN (2008) Assembling DNA Barcodes. In: Martin CC (Ed.) *Methods in Molecular Biology* (Vol. 410): Environmental Genomics. Humana Press, 275–294. https://doi.org/10.1007/978-1-59745-548-0_15
- deWaard JR, Landry J-F, Schmidt BC, Derhousoff J, McLean JA, Humble LM (2009) In the dark in a large urban park: DNA barcodes illuminate cryptic and introduced moth species. *Biodiversity and Conservation* 18: 3825–3839. <https://doi.org/10.1007/s10531-009-9682-7>
- deWaard JR, Levesque-Beaudin V, DeWaard SL, Ivanova NV, McKeown JTA, Miskie R, Naik S, Perez KHJ, Ratnasingham S, Sobel CN, Sones JE, Steinke C, Telfer AC, Young AD, Young MR, Zakharov EV, Hebert PDN (2018) Expedited assessment of terrestrial arthropod diversity by coupling Malaise traps with DNA barcoding. *Genome* 62(3): 85–95. <https://doi.org/10.1139/gen-2018-0093>
- Döberl M (1994) Alticinae. In: Lohse GA, Lucht WH (Eds) *Die Käfer Mitteleuropas*, Band 14 (3. Supplementband). Goecke & Evers, Krefeld, 92–141.
- Döberl M (2010) Alticinae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 6). Apollo Books, Stenstrup, 491–563.
- Dolgovskaya MY, Konstantinov AS, Reznik SY, Spencer NR, Volkovitsh MG (2004) Flea beetles (Coleoptera: Chrysomelidae) associated with purple loosestrife, *Lythrum salicaria*, in Russia. In: Cullen JM, Briese DT, Kriticos DJ, Lonsdale WM, Morin L, Scott JK (Eds) *Proceedings of the XI International Symposium on Biological Control of Weeds*. CSIRO Entomology, Canberra, 96–101.
- Downie NM, Arnett RH (1996) *The Beetles of Northeastern North America* (Vol. 1, 2). Sandhill Crane Press, Gainesville, 1721 pp.
- Endrödy-Younga S (1961) Revision der Gattung *Calyptomerus* Redtb. (Coleoptera: Clambidae). *Acta Zoologica Academiae Scientiarum Hungaricae* 7: 401–412.
- Endrödy-Younga S (1974) A revision of the described Australian and New Zealand species of the family Clambidae (Coleoptera) with description of a new species. *Records of the South Australian Museum* 17: 1–10.
- Endrödy-Younga S (1981) The American species of the familia Clambidae (Coleoptera: Eucinetoidae). *Entomologia Generalis* 7: 33–67.
- Endrödy-Younga S (1990) A revision of the Australian Clambidae (Coleoptera: Eucinetoidae). *Invertebrate Taxonomy* 4: 247–280. <https://doi.org/10.1071/IT9900247>
- Esser J (2016) Über die Identität von *Cryptophilus integer* (Heer, 1841) (Coleoptera, Erotylidae). *Entomologische Nachrichten und Berichte* 60: 213–218.
- Esser J (2017) On the Nearctic *Cryptophilus* Reitter, 1874 (Coleoptera: Erotylidae). *Linzer biologische Beiträge* 49: 1133–1137.
- Faccoli M, Frigimelica G, Mori N, Petrucco Toffolo E, Vettorazzo M, Simonato M (2009) First record of *Ambrosiodmus* (Hopkins, 1915) (Coleoptera: Curculionidae, Scolytinae) in Europe. *Zootaxa* 2303: 57–60. <https://doi.org/10.11646/zootaxa.2303.1.4>
- Fall HC (1901) Notes on *Dichelonycha* and *Cantharis*, with descriptions of new species in oth-
era [sic] genera. *Transactions of the American Entomological Society* 27: 277–310.
- Fall HC (1910) Miscellaneous notes and descriptions of North American Coleoptera. *Transactions of the American Entomological Society* 36: 89–197.
- Fall HC (1919) A Change of Names (Coleoptera). *Entomological News* 30: 1–26.

- Fall HC (1927) A review of the North American species of *Podabrus*. *Entomologica Americana* 8: 65–103.
- Fender KM (1951) The Malthini of North America (Coleoptera-Cantharidae). *The American Midland Naturalist* 46: 513–629. <https://doi.org/10.2307/2421804>
- Fender KM (1961) Family Cantharidae. In: Hatch MH (Ed.) *Beetles of the Pacific Northwest. Part III: Pselaphidae and Diversicornia I*. University of Washington Press, Seattle, 44–68.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Ford EJJ (1973) A revision of the genus *Petalium* LeConte in the United States, Greater Antilles, and the Bahamas (Coleoptera: Anobiidae). *United States Department of Agriculture Technical Bulletin* 1476: 1–40.
- Francis A, Cavers PB, Warwick SI (2009) The Biology of Canadian Weeds. 140. *Hesperis matronalis* L. *Canadian Journal of Plant Science* 89: 191–206. <https://doi.org/10.4141/CJPS08094>
- Franz H, Besuchet C (1971) Scydmaenidae. In: Freude H, Harde KW, Lohse GA (Eds) *Die Käfer Mitteleuropas, Band 3*. Goecke & Evers, Krefeld, 271–303.
- Franzen B (1991) Vorläufige Mitteilung über eine *Cryptophilus*-Art neu für Mitteleuropa (Coleoptera: Languriidae). *Mitteilungen der Arbeitsgemeinschaft Rheinischer Koleopterologen* 1: 59–63.
- Fürsch H (1965) Die palaearktischen Arten der *Scymnus-bipunctatus*-Gruppe und die europäischen Vertreter der Untergattung *Sidis* (Col. Cocc.). *Mitteilungen der Münchner Entomologischen Gesellschaft* 55: 178–213.
- Fürsch H (1967) Coccinellidae. In: Freude H, Harde KW, Lohse GA (Eds) *Die Käfer Mitteleuropas, Band 7*. Goecke & Evers, Krefeld, 227–278.
- Fürsch H (1987) Übersicht über die genera und Subgenera der Scymnini mit besonderer Berücksichtigung der Westpalaearktis (Insecta, Coleoptera, Coccinellidae). *Entomologische Abhandlungen (Dresden)* 51: 57–74.
- Gimmel ML (2013) Genus-level revision of the family Phalacridae (Coleoptera: Cucujoidea). *Zootaxa* 3605: 1–147. <https://doi.org/10.11646/zootaxa.3605.1.1>
- Gomez DF, Rabaglia RJ, Fairbanks KEO, Hulcr J (2018) North American Xyleborini north of Mexico: a review and key to genera and species (Coleoptera, Curculionidae, Scolytinae). *ZooKeys* 768: 19–68. <https://doi.org/10.3897/zookeys.768.24697>
- Good JA (1995) Habitat records of *Amischa* C.G. Thomson (Staphylinidae: Aleocharinae) from Ireland. *Coleopterist* 3: 77–79.
- Gordon RD (1976) The Scymnini (Coleoptera: Coccinellidae) of the United States and Canada: Key to genera and revision of *Scymnus*, *Nephus* and *Diomus*. *Bulletin of the Buffalo Society of Natural Sciences* 28: 1–362.
- Gordon RD (1985) The Coccinellidae (Coleoptera) of America north of Mexico. *Journal of the New York Entomological Society* 93: 1–912.
- Gustafson GT, Miller KB (2015) The New World whirligig beetles of the genus *Dineutus* Macleay, 1825 (Coleoptera, Gyrinidae, Gyrininae, Dineutini). *ZooKeys* 476: 1–135. <https://doi.org/10.3897/zookeys.476.8630>

- Gwiazdowski RA, Foottit RG, Maw HEL, Hebert PDN, Imondi R, LaRocca G (2015) The Hemiptera (Insecta) of Canada: Constructing a reference library of DNA barcodes. *PLoS ONE* 10: e0125635. <https://doi.org/10.1371/journal.pone.0125635>
- Hackwell GA (1973) Biology of *Lasconotus subcostulatus* (Coleoptera: Colydiidae) with special reference to feeding behavior. *Annals of the Entomological Society of America* 66: 62–65. <https://doi.org/10.1093/aesa/66.1.62>
- Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN (2006) DNA barcodes distinguish species of tropical Lepidoptera. *Proceedings of the National Academy of Sciences of the United States of America* 103: 968–971. <https://doi.org/10.1073/pnas.0510466103>
- Halstead DGH (1981) Taxonomic notes on some *Attagenus* spp associated with stored products, including a new black species from Africa (Coleoptera: Dermestidae). *Journal of Stored Product Research* 17: 91–99. [https://doi.org/10.1016/0022-474X\(81\)90007-2](https://doi.org/10.1016/0022-474X(81)90007-2)
- Hansen M (1987) 18 Fauna Entomologica Scandinavica The Hydrophiloidea (Coleoptera) of Fennoscandia and Denmark. EJ Brill, Copenhagen, 254 pp.
- Hansen M (2004) Hydrophilidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 2). Apollo Books, Stenstrup, 44–68.
- Hatch MH (1957) *The beetles of the Pacific Northwest. Part II: Staphyliniformia*. University of Washington Press, Seattle, 384 pp.
- Hebert PDN, Cywinska A, Ball SL, deWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 270: 313–321. <https://doi.org/10.1098/rspb.2002.2218>
- Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W (2004) Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences of the United States of America* 101: 14812–14817. <https://doi.org/10.1073/pnas.0406166101>
- Hebert PDN, Ratnasingham S, Zakharov EV, Telfer AC, Levesque-Beaudin V, Milton MA, Pedersen S, Jannetta P, DeWaard JR (2016) Counting animal species with DNA barcodes: Canadian insects. *Philosophical Transactions of the Royal Society of London. Series B, Biological sciences* 371: e1001127. <https://doi.org/10.1098/rstb.2015.0333>
- Hendrich L, Morinière J, Haszprunar G, Hebert PDN, Hausmann A, Köhler F, Balke M (2015) A comprehensive DNA barcode database for Central European beetles with a focus on Germany: adding more than 3,500 identified species to BOLD. *Molecular Ecology Resources* 15: 795–818. <https://doi.org/10.1111/1755-0998.12354>
- Hendrickson RMJ, Drea JJ, Rose M (1991) A distribution and establishment program for *Chilocorus kuwanae* (Silvestri) (Coleoptera: Coccinellidae) in the United States. *Proceedings of the Entomological Society of Washington* 93: 197–200.
- Herman L (1972) Revision of *Bledius* and related genera. Part I. The *aequatorialis*, *mandibularis*, and *semiferrugineus* groups and two new genera (Coleoptera, Staphylinidae, Oxytelinae). *Bulletin of the American Museum of Natural History* 149: 111–254.
- Hicks EA (1959) *Check-list and Bibliography on the Occurrence of Insects in Birds' Nests*. Iowa State College Press, Ames, 696 pp. <https://doi.org/10.5962/bhl.title.6819>
- Hoffmann A (1958) *Faune de France 62. Coléoptères Curculionides. 3^{ème} partie*. Fédération Française des Sociétés de Sciences Naturelles, Paris, 632 pp.

- Ivanova NV, deWaard JR, Hebert PDN (2006) An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Molecular Ecology Notes* 6: 998–1002. <https://doi.org/10.1111/j.1471-8286.2006.01428.x>
- Jałoszynski P (2019) Notes on Nearctic genera *Catalinus* Casey and *Parascydmus* Casey (Coleoptera: Staphylinidae: Scydmaeninae). *Zootaxa* 4603: 145–158. <https://doi.org/10.11646/zootaxa.4603.1.7>
- Jelínek J (2008) Ciidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 5). Apollo Books, Stenstrup, 55–62.
- Jelínek J, Audisio P (2007) Nitidulidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 4). Apollo Books, Stenstrup, 459–491.
- Jendek E, Grebennikov VV, Bocak L (2015) Undetected for a century: Palaearctic *Agrilus ribesi* Schaefer (Coleoptera: Buprestidae) on currant in North America, with adult morphology, larval biology and DNA barcode. *Zootaxa* 4034: 1–112. <https://doi.org/10.11646/zootaxa.4034.1.5>
- Johnson C (1997) *Clambus simsoni* Blackburn (Col., Clambidae) new to Britain with notes on its wider distribution. *Entomologist's Monthly Magazine* 133: 161–164.
- Johnson P (2002) Elateridae Leach, 1815. In: Arnett RHJ, Thomas MC, Skelley PE, Frank JH (Eds) *American Beetles* (Vol. 2). Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, 160–173.
- Kalik V (1992) Dermestidae. In: Lohse GA, Lucht WH (Eds) *Die Käfer Mitteleuropas*, Band 13 (2. Supplementband). Goecke & Evers, Krefeld, 83–86.
- Kazantsev SV (1998) НОВЫЕ РОССИЙСКИЕ ВИДЫ *Dichelotarsus* (Coleoptera, Cantharidae) [New Russian species of *Dichelotarsus* (Coleoptera, Cantharidae)]. *Zoologicheskii zhurnal* 77: 572–575.
- Kazantsev SV, Brancucci M (2007) Cantharidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 4). Apollo Books, Stenstrup, 234–298.
- Kissinger DG (1964) Curculionidae of America north of Mexico. A key to genera. Taxonomic Publications, South Lancaster, 143 pp.
- Klausnitzer B (1976) Zur Kennits der Nordamerikanischen Arten der Gattung *Cyphon* Paykull (Col., Helodidae) (40. Beitrag zur Kennits der Helodidae). *Polskie Pismo Entomologiczne* 46: 439–453.
- Klausnitzer B (2006) Scirtidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 3). Apollo Books, Stenstrup, 316–323.
- Klimaszewski J (1982) Studies of Myllaenini (Coleoptera: Staphylinidae, Aleocharinae). 1. Systematics, phylogeny, and zoogeography of Nearctic *Myllaena* Erichson. *The Canadian Entomologist* 114: 181–242. <https://doi.org/10.4039/Ent114181-3>
- Klimaszewski J, Brunke AJ, Assing V, Langor DW, Newton AF, Bourdon C, Pelletier GA, Webster RP, Herman L, Perdereau L, Davies A, Smetana A, Chandler DS, Majka CG, Scudder GGE (2013) Synopsis of Adventive Species of Coleoptera (Insecta) Recorded from Canada. Part 2: Staphylinidae. Pensoft Publishers, Sofia, 360 pp.
- Klimaszewski J, Langor D, Batista R, Duval J-A, Majka CG, Scudder GGE, Bousquet Y (2012) Synopsis of Adventive Species of Coleoptera (Insecta) Recorded from Canada. Part 1: Carabidae. Pensoft Publishers, Sofia, 96 pp.

- Klimaszewski J, Langor D, Majka CG, Bouchard P, Bousquet Y, LeSage L, Smetana A, Sylvestre P, Pelletier G, Davies A, DesRochers P, Goulet H, Webster R, Sweeney J (2010) Review of Adventive Species of Coleoptera (Insecta) Recorded from Eastern Canada. Pensoft Series Faunistica No. 94. Pensoft Publishers, Sofia, 272 pp.
- Klimaszewski J, Langor DW, Hammond HEJ, Pelletier GA, Bousquet Y, Bourdon C, Webster RP, Borowiec L, Scudder GGE, Majka CG (2015) Synopsis of Adventive Species of Coleoptera (Insecta) Recorded from Canada. Part 3: Cucujoidea. Pensoft Publishers, Sofia, 172 pp.
- Klimaszewski J, Langor DW, Smith ABT, Hoebeke ER, Davies A, Pelletier GA, Douglas H, Webster RP, Bourdon C, Borowiec L, Scudder GGE (2017) Synopsis of Adventive Species of Coleoptera (Insecta) Recorded from Canada. Part 4: Scarabaeoidea, Scirtoidea, Buprestoidea, Byrrhoidea, Elateroidea, Derodontoidea, Bostrichoidea, and Cleroidea. Pensoft Publishers, Sofia, 215 pp.
- Klimaszewski J, Webster R, Langor D, Brunke AJ, Davies A, Bourdon C, Labrecque M, Newton AF, Dorval J-A, Frank JH (2018) Aleocharine Rove Beetles of Eastern Canada (Coleoptera, Staphylinidae, Aleocharinae): a Glimpse of Megadiversity. Springer, Cham, 879 pp.
- Knížek M (2011) Scolytinae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 7). Apollo Books, Stenstrup, 204–251.
- Koch K (1989a) Die Käfer Mitteleuropas, Ökologie (Vol. 1). Goecke & Evers Verlag, Krefeld, 440 pp.
- Koch K (1989b) Die Käfer Mitteleuropas, Ökologie (Vol. 2). Goecke & Evers Verlag, Krefeld, 382 pp.
- Koch K (1992) Die Käfer Mitteleuropas, Ökologie (Vol. 3). Goecke & Evers Verlag, Krefeld, 389 pp.
- Kolov SV, Korotyayev BA (2017) On establishing of two weevils (Coleoptera, Curculionidae: Entiminae) damaging fruit and berry crops in Southeastern Kazakhstan. Entomological Review 97: 404–405. <https://doi.org/10.1134/S0013873817030149>
- Konstantinov AS, Baselga A, Grebennikov VV, Prena J, Lingafelter SW (2011) Revision of the Palearctic *Chaetocnema* Species (Coleoptera: Chrysomelidae: Galerucinae: Alticini). Pensoft Publishers, Sofia, 363 pp.
- Kovář I (2007) Coccinellidae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 4). Apollo Books, Stenstrup, 568–631.
- Landry J-F, Nazari V, DeWaard JR, Mutanen M, Lopez-Vaamonde C, Huemer P, Hebert PDN (2013) Shared but overlooked: 30 species of Holarctic Microlepidoptera revealed by DNA barcodes and morphology. Zootaxa 3749(1): 1–93. <https://doi.org/10.11646/zootaxa.3749.1.1>
- Lane MC (1948) Some generic corrections in the Elateridae- I. (Coleoptera). Proceedings of the Entomological Society of Washington 50: 179–182.
- Larochelle A, Larivière M-C (2003) A Natural History of the Ground-beetles (Coleoptera: Carabidae) of America North of Mexico. Pensoft Publishers, Sofia, 584 pp.
- Larsen LM, Nielsen JK, Sørensen H (1992) Host plant recognition in monophagous weevils: Specialization of *Ceutorhynchus inaeffectatus* to glucosinolates from its host plant *Hesperis matronalis*. Entomologia Experimentalis et Applicata 64: 49–55. <https://doi.org/10.1111/j.1570-7458.1992.tb01593.x>

- Lasnier C, McFadden-Smith W, Moreau D, Bouchard P, Vincent C (2019) Guide to the key arthropods of vineyards of Eastern Canada. Ottawa.
- Lawrence JF (1971) Revision of the North American Ciidae (Coleoptera). Bulletin of the Museum of Comparative Zoology at Harvard College 142: 419–522.
- Lawrence JF, Slipinski A, Seago AE, Thayer MK, Newton AF, Marvaldi AE (2011) Phylogeny of the Coleoptera based on morphological characters of adults and larvae. *Annales Zoologici* 61: 1–217. <https://doi.org/10.3161/000345411X576725>
- LeConte JL (1878) Descriptions of new species. In: Hubbard HG, Schwarz EA (Eds) The Coleoptera of Michigan. Proceedings of the American Philosophical Society 17: 593–626.
- Lees DC, Lack HW, Rougerie R, Hernandez-Lopez A, Raus T, Avtzis ND, Augustin S, Lopez-Vaamonde C (2011) Tracking origins of invasive herbivores through herbaria and archival DNA: the case of the horse-chestnut leaf miner. *Frontiers in Ecology and the Environment* 9: 322–328. <https://doi.org/10.1890/100098>
- Leseigneur L (1998) Throscidae. In: Lucht W, Klausnitzer B (Eds) Die Käfer Mitteleuropas, Band 15 (4. Supplementband). Spektrum Akademischer Verlag, Heidelberg, 222–231.
- Leseigneur L (2005) Description de *Trixagus meyhohmi* n. sp. et note sur la morphologie des *Trixagus* du groupe *carinifrons* (Coleoptera, Throscidae). *Bulletin de la Société entomologique de France* 110: 89–96.
- Leseigneur L (2007) Throscidae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 4). Apollo Books, Stenstrup, 87–89.
- Liljeblad E (1945) Monograph of the family Mordellidae (Coleoptera) of North America, north of Mexico. Miscellaneous Publications, Museum of Zoology, University of Michigan, No. 62. University of Michigan Press, Ann Arbor, 229 pp.
- Lindroth CH (1968) The ground-beetles (Carabidae, excl. Cicindelinae) of Canada and Alaska. Part 5. *Opuscula Entomologica Supplementum* 33: 649–944.
- Löbl I (1960) Die tschechoslowakischen Arten der Gattung *Biblopectus* Reitter (Col. Pselaphidae). *Acta Societatis Entomologicae Čechosloveniae* 57: 58–67.
- Löbl I (2006) Clambidae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 3). Apollo Books, Stenstrup, 314–316.
- Lockwood JL, Hoopes MF, Marchetti MP (2013) *Invasion Ecology* (2nd ed.). Wiley-Blackwell, Chichester, 444 pp.
- Lohse GA (1967) Cisidae. In: Freude H, Harde KW, Lohse GA (Eds) Die Käfer Mitteleuropas, Band 7. Goecke & Evers, Krefeld, 280–295.
- Lohse GA (1979) Helodidae. In: Freude H, Harde KW, Lohse GA (Eds) Die Käfer Mitteleuropas, Band 6. Goecke & Evers, Krefeld, 249–263.
- Lohse GA (1983) Ceutorhynchinae. In: Freude H, Harde KW, Lohse GA (Eds) Die Käfer Mitteleuropas, Band 11. Goecke & Evers, Krefeld, 180–253.
- Lonsdale O, Huber JT (2011) Insect Collections of Canada Series: Canadian National Collection of Insects, Arachnids & Nematodes, Ottawa. *Newsletter of the Biological Survey of Canada* 30: 15–40.
- Lopes-Andrade C, Webster RP, Webster VL, Alderson CA, Hughes CC, Sweeney JD (2016) The Ciidae (Coleoptera) of New Brunswick, Canada: New records and new synonyms. *ZooKeys* 573: 339–366. <https://doi.org/10.3897/zookeys.573.7445>

- Lord NP, Nearn EH, Miller KB (2011–2013) Ironclad ID: Tool for Diagnosing Ironclad and Cylindrical Bark Beetles (Coleoptera: Zopheridae) of North America north of Mexico. The University of New Mexico and Center for Plant Health Science and Technology, USDA, APHIS, PPQ. <http://cerambycids.com/ironcladid/index.html>
- Luginbill P, Painter HR (1953) May beetles of the United States and Canada. United States Department of Agriculture Technical Bulletin 1060: 1–102.
- Majka CG, Klimaszewski J, Lauff RF (2006) New Coleoptera records from owl nests in Nova Scotia, Canada. *Zootaxa* 1194: 33–47.
- Majka CG, LeSage L (2010) *Chaetocnema* flea beetles (Coleoptera: Chrysomelidae, Alticini) of the Maritime provinces of Canada. *Journal of the Acadian Entomological Society* 6: 34–38.
- Maxwell SL, Fuller RA, Brooks TM, Watson JEM (2016) Biodiversity: The ravages of guns, nets and bulldozers. *Nature* 536: 143–145. <https://doi.org/10.1038/536143a>
- Mertlik J, Leseigneur L (2007) The species of the family Throscidae (Coleoptera: Elateroidea) Czech and Slovak Republics. *Elateridarium* 1: 1–55.
- Meybohm H (2004) *Clambus lohsei* n. sp. aus der Umgebung von Hamburg (Coleoptera, Clambidae). *Entomologische Blätter* 100: 13–18.
- Mohr K-H (1966) Chrysomelidae. In: Freude H, Harde KW, Lohse GA (Eds) *Die Käfer Mitteleuropas*, Band 9. Goecke & Evers, Krefeld, 95–280.
- Muona J (1984) Review of the Palearctic Aleocharinae also occurring in North America (Coleoptera: Staphylinidae). *Entomologica Scandinavica* 15: 227–231. <https://doi.org/10.1163/187631284X00190>
- Muona J (1990) The Fennoscandian and Danish species of the genus *Amischa* Thomson (Coleoptera, Staphylinidae). *Entomologisk Tidsskrift* 111: 17–24.
- Mutanen M, Hausmann A, Hebert PDN, Landry J-F, de Waard JR, Huemer P (2012) Allopatry as a Gordian knot for taxonomists: patterns of DNA barcode divergence in arctic-alpine Lepidoptera. *PLoS ONE* 7: e47214. <https://doi.org/10.1371/journal.pone.0047214>
- Nikitsky NB (2008) Mycetophagidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 5). Apollo Books, Stenstrup, 51–55.
- Nyholm T (1972) Die nordeuropäischen Arten der Gattung *Cyphon* Paykull (Coleoptera). *Entomologica Scandinavica Supplementum* 3: 1–100.
- O'Brien CW, Wibmer GJ (1982) Annotated checklist of the weevils (Curculionidae sensu lato) of North America, Central America, and the West Indies (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 34: 1–382.
- O'Keefe ST (2001) Scydmaenidae Leach, 1815. In: Arnett RH, Thomas MC (Eds) *American Beetles* (Vol. 1). Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia. CRC Press, Boca Raton, 259–267.
- Owens BE, Carlton CE (2017) Seven new species of *Biblopectus* Reitter (Coleoptera: Staphylinidae: Pselaphinae) from Eastern United States with notes on distributions of previously described species. *Zootaxa* 4243: 139–152. <https://doi.org/10.11646/zootaxa.4243.1.6>
- Palm T (1961) *Svensk Insektfauna* 9. Skalbagger, Coleoptera. Kortvingar: Fam. Staphylinidae, Häfte 2. Underfam. Oxytelinae, Oxyporinae, Steninae, Euaesthetinae. Entomologiska Föreningen i Stockholm, Stockholm, 126 pp.

- Palm T (1963) Svensk Insektfauna 9. Skalbaggar, Coleoptera. Kortvingar: Fam. Staphylinidae, Häfte 3. Underfam. Paederinae, Staphylininae. Entomologiska Föreningen i Stockholm, Stockholm, 168 pp.
- Palm T (1968) Svensk Insektfauna 9. Skalbaggar. Coleoptera, Kortvingar: Fam. Staphylinidae, Häfte 5. Underfam. Aleocharinae (*Deinopsis-Trichomicra*). Entomologiska Föreningen i Stockholm, Stockholm, 112 pp.
- Palm T (1970) Svensk Insektfauna 9. Skalbaggar. Coleoptera, Kortvingar: Fam. Staphylinidae, Häfte 6. Underfam. Aleocharinae (*Atheta*). Entomologiska Föreningen i Stockholm, Stockholm, 180 pp.
- Park O (1929) Taxonomic studies in Coleoptera, with notes upon certain species of beetles in the Chicago area, I. Journal of the New York Entomological Society 37: 429–436.
- Parsons CT (1975) Revision of Nearctic Mycetophagidae (Coleoptera). The Coleopterists Bulletin 29: 93–108.
- Pavlov IE (1960) Ecology of cereal-stem flea-beetles (Coleoptera, Chrysomelidae, Halticinae) and measures for their control [In Russian]. Entomologicheskoe Obozrenie 39: 775–795.
- Peacock ER (1979) *Attagenus smirnovi* Zhantiev (Coleoptera: Dermestidae) a species new to Britain, with keys to the adults and larvae of British *Attagenus*. Entomologist's Gazette 30: 131–136.
- Pearce EJ (1957) Coleoptera Pselaphidae. Handbooks for the identification of British insects (Vol. IV), part 9. Royal Entomological Society, London, 32 pp. http://www.royensoc.co.uk/sites/default/files/Vol04_Part09.pdf
- Pelletier GA, Hébert C (2014) The Cantharidae of Eastern Canada and Northeastern United States. Canadian Journal of Arthropod Identification 25: 1–246. <https://doi.org/10.3752/cjai.2014.25>
- Pentinsaari M, Hebert PDN, Mutanen M (2014) Barcoding Beetles: A regional survey of 1872 species reveals high identification success and unusually deep interspecific divergences. PLoS ONE 9: e108651. <https://doi.org/10.1371/journal.pone.0108651>
- Perreau M (2004) Leiodidae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 2). Apollo Books, Stenstrup, 133–203.
- Pimentel D (2011) Environmental and economic costs associated with alien invasive species in the United States. In: Pimentel D (Ed.) Biological Invasions: Economic and Environmental Costs of Alien Plant, Animal, and Microbe Species. CRC Press, Boca Raton, 411–430. <https://doi.org/10.1201/b10938-26>
- Rassi P, Karjalainen S, Clayhills T, Helve E, Hyvärinen E, Laurinharju E, Malmberg S, Mannerkoski I, Martikainen P, Mattila J, Muona J, Pentinsaari M, Rutanen I, Salokannel J, Siitonen J, Silfverberg H (2015) Kovakuoriaisten maakuntaluettelo 2015 [Provincial List of Finnish Coleoptera 2015]. Sahlbergia 21: 1–164. http://www.luomus.fi/sites/default/files/sahlbergia/sahlbergia_21_S1_2015.pdf
- Ratnasingham S, Hebert PDN (2007) BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). Molecular Ecology Notes 7: 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Ratnasingham S, Hebert PDN (2013) A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. PLoS ONE 8: e66213. <https://doi.org/10.1371/journal.pone.0066213>

- Reibnitz J (1999) Verbreitung und Lebensräume der Baumschwammfresser Südwestdeutschlands (Coleoptera: Cisidae). *Mitteilungen des Entomologischen Vereins Stuttgart* 34: 3–74.
- Reid CAM (1991) Notes on distribution of *Myllaena* spp. (Col., Staphylinidae) in Buckinghamshire and north-east England. *Entomologist's Monthly Magazine* 127: 1–24.
- Rheinheimer J, Hassler M (2013) Die Rüsselkäfer Baden-Württembergs (2nd ed.). Landesanstalt für Umwelt, Messungen und Naturschutz Baden-Württemberg (LUBW), Ubstadt-Weiher, 944 pp.
- Riedl H, Taschenberg EF (1984) Grape cane gallmaker *Ampelogypter sesostris* (LeConte). Grape IPM. Insect Identification Sheet No. 2. New York State IPM Program. Cornell University fact sheet. <https://ecommons.cornell.edu/bitstream/handle/1813/43098/grape-cane-gallmaker-FS-NYSIPM.pdf>
- Riley EG, Clark SM, Flowers RW, Gilbert AJ (2002) Chrysomelidae Latreille, 1802. In: Arnett RH, Thomas MC, Skelley PE, Frank JH (Eds) *American Beetles* (Vol. 2). Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, 617–691.
- Rougerie R, Lopez-Vaamonde C, Barnouin T, Delnatte J, Moulin N, Noblecourt T, Nusillard B, Parmain G, Soldati F, Bouget C (2015) PASSIFOR: A reference library of DNA barcodes for French saproxylic beetles (Insecta, Coleoptera). *Biodiversity Data Journal* 3: e4078. <https://doi.org/10.3897/BDJ.3.e4078>
- Ruta R, Jałoszynski P, Sienkiewicz P, Konwerski S (2011) Erotylidae (Insecta, Coleoptera) of Poland – problematic taxa, updated keys and new records. *ZooKeys* 134: 1–13. <https://doi.org/10.3897/zookeys.134.1673>
- Rutanen I (2015) *Orthoperus* Stephens -suvun kovakuoriaisten esiintymisestä Suomessa (Coleoptera: Corylophidae) [On the occurrence of the species of *Orthoperus* Stephens in Finland (Coleoptera, Corylophidae)]. *Sahlbergia* 21: 40–48.
- Rutanen I, Martikainen P (2014) *Longitarsus pratensis* (Panzer) -ryhmän lajit Suomessa [*Longitarsus pratensis* (Panzer) species group in Finland]. *Sahlbergia* 20: 28–35. https://www.luomus.fi/sites/default/files/sahlbergia/2014/Sahlbergia_2014_1-Rutanen_Martikainen.pdf
- Saunders MC, Tobin PC (2000) Grape Cane Gallmaker (Coleoptera: Curculionidae) and its impact on cultivated grapes. *Journal of Economic Entomology* 93: 795–799. <https://doi.org/10.1603/0022-0493-93.3.795>
- Scheibner RA (1963) A taxonomic study of the genus *Ceutorhynchus* (Coleoptera, Curculionidae). PhD Thesis, Michigan State University, Department of Entomology, 156 pp.
- Schülke M (2012a) Oxytelinae. In: Assing V, Schülke M (Eds) *Die Käfer Mitteleuropas*, Band 4 (2. Auflage). Spektrum Akademischer Verlag, Heidelberg, 283–284.
- Schülke M (2012b) Tachyporinae. In: Assing V, Schülke M (Eds) *Die Käfer Mitteleuropas*, Band 4 (2. Auflage). Spektrum Akademischer Verlag, Heidelberg, 130–199.
- Schülke M, Smetana A (2015) Staphylinidae. In: Löbl I, Löbl D (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 2). Brill, Leiden, 901–1134. <https://doi.org/10.1163/9789004296855>
- Silfverberg H (2010) Enumeratio renovata Coleopterorum Fennoscandiae, Daniae et Baltiae. *Sahlbergia* 16: 1–144.
- Steiner WEJ, Singh BP (1987) Redescription of an ergot beetle, *Acyломus pugetanus* Casey, with immature stages and biology (Coleoptera: Phalacridae). *Proceedings of the Entomological Society of Washington* 89: 744–758.

- Steinke D, Breton V, Berzitis E, Hebert PDN (2017) The School Malaise Trap Program: Coupling educational outreach with scientific discovery. *PLoS Biology* 15: e2001829. <https://doi.org/10.1371/journal.pbio.2001829>
- Stengaard Hansen L, Åkerlund M, Grøntoft T, Ryhl-Svendsen M, Schmidt AL, Bergh JE, Vagn Jensen KM (2012) Future pest status of an insect pest in museums, *Attagenus smirnovi*: Distribution and food consumption in relation to climate change. *Journal of Cultural Heritage* 13: 22–27. <https://doi.org/10.1016/j.culher.2011.05.005>
- Stephan KH (1989) The Bothrideridae and Colydiidae of America North of Mexico (Coleoptera: Clavicornia and Heteromera). *Occasional Papers of the Florida State Collection of Arthropods* 6: 1–65.
- Švec Z (2002) Revision of the African species of the genus *Tinodemus* Guillebeau (Coleoptera, Phalacridae). Results of the entomological expeditions of the Museum of Natural History, Berlin to Africa. 76th contribution. *Zoosystematics and Evolution* 78: 217–256. <https://doi.org/10.1002/mmz.20020780204>
- Švec Z (2007) Phalacridae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera* (Vol. 4). Apollo Books, Stenstrup, 506–513.
- Van Tassell ER (2001) Hydrophilidae Latreille, 1802. In: Arnett RH, Thomas MC (Eds) *American Beetles* (Vol. 1). Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia. CRC Press, Boca Raton, 187–208. <https://doi.org/10.1201/9781482274325>
- Thompson RT (1958) Coleoptera Phalacridae. *Handbooks for the identification of British insects* (Vol. V), part 5(b). Royal Entomological Society, London, 17 pp. http://www.royensoc.co.uk/sites/default/files/Vol05_Part05b.pdf
- Valade R, Kenis M, Hernandez-Lopez A, Augustin S, Mari Mena N, Magnoux E, Rougerie R, Lakatos F, Roques A, Lopez-Vaamonde C (2009) Mitochondrial and microsatellite DNA markers reveal a Balkan origin for the highly invasive horse-chestnut leaf miner *Cameraria ohridella* (Lepidoptera, Gracillariidae). *Molecular Ecology* 18: 3458–3470. <https://doi.org/10.1111/j.1365-294X.2009.04290.x>
- Vappula NA (1965) Pests of cultivated plants in Finland. *Acta Entomologica Fennica* 19: 1–239.
- Vogt H (1967a) Mycetophagidae. In: Freude H, Harde KW, Lohse GA (Eds) *Die Käfer Mitteleuropas*, Band 7. Goecke & Evers, Krefeld, 191–196.
- Vogt H (1967b) Phalacridae. In: Freude H, Harde KW, Lohse GA (Eds) *Die Käfer Mitteleuropas*, Band 7. Goecke & Evers, Krefeld, 158–166.
- Warchalowski A (1996) Übersicht der Westpaläarktischen Arten der Gattung *Longitarsus* Berthold, 1827 (Coleoptera: Chrysomelidae: Halticinae). *Genus* (Supplement). Wrocław, 266 pp.
- Webster RP, Davies AE, Klimaszewski J, Bourdon C (2016) Further contributions to the staphylinid fauna of New Brunswick, Canada, and the USA, with descriptions of two new *Proteinus* species (Coleoptera, Staphylinidae). *ZooKeys* 573: 31–83. <https://doi.org/10.3897/zookeys.573.7830>
- Webster RP, Klimaszewski J, Pelletier G, Savard K (2009) New Staphylinidae (Coleoptera) records with new collection data from New Brunswick, Canada. I. Aleocharinae. *ZooKeys* 22: 171–248. <https://doi.org/10.3897/zookeys.22.152>

- Wheeler AG, Hoebeke ER (2009) Adventive (Non-Native) Insects: Importance to Science and Society. In: Foottit RG, Adler PH (Eds) Insect Biodiversity: Science and Society. Wiley-Blackwell, Oxford, 475–521. <https://doi.org/10.1002/9781444308211.ch21>
- White RE (1982) A Catalog of the Coleoptera of America North of Mexico. Family: Anobiidae. Agriculture Handbook Number 529–70. United States Department of Agriculture, Agricultural Research Service, Washington, 59 pp.
- White RE (1996) A revision of the genus *Chaetocnema* of America north of Mexico (Coleoptera: Chrysomelidae). Contributions of the American Entomological Institute 29: 1–158.
- Williams SA (1969) The British species of the genus *Amischa* (Thomson) (Col., Staphylinidae), including *A. soror* Kraatz, an addition to the list. Entomologist's Monthly Magazine 113: 1–250.
- Wittmer W (1970) Zur Kenntnis der Gattung *Malthodes* Kies. (Col., Cantharidae). Entomologische Arbeiten aus dem Museum G. Frey 21: 13–107.
- Wittmer W (1979) Malthinini. In: Freude H, Harde KW, Lohse GA (Eds) Die Käfer Mitteleuropas, Band 6. Goecke & Evers, Krefeld, 40–51.

Supplementary material I

Table S1. Detailed collection data, GenBank accession numbers, and institutional storage information of the studied specimens

Authors: Mikko Pentinsaari, Robert Anderson, Lech Borowiec, Patrice Bouchard, Adam Brunke, Hume Douglas, Andrew B.T. Smith, Paul D. N. Hebert

Data type: species data

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.894.37862.suppl1>