

The complete mitochondrial genomes of five lichenized fungi in the genus *Usnea* (Ascomycota: Parmeliaceae)

Erik R. Funk^{a*}, Alexander N. Adams^{a*}, Sarah M. Spotten^{a*}, Roxanne A. Van Hove^a, Kristina T. Whittington^a, Kyle G. Keepers^a, Cloe S. Pogoda^b, James C. Lendemer^c, Erin A. Tripp^{a,d} and Nolan C. Kane^a

^aDepartment of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, USA; ^bDepartment of Molecular, Cellular and Developmental Biology, University of Colorado, Boulder, CO, USA; ^cInstitute of Systematic Botany, The New York Botanical Garden, Bronx, NY, USA; ^dMuseum of Natural History, University of Colorado, Boulder, CO, USA

ABSTRACT

Known colloquially as ‘Old Man’s Beard’, *Usnea* is a genus of lichenized Ascomycete fungi characterized by having a fruticose growth form and cartilaginous central axis. The complete mitochondrial genomes of *Usnea halei*, *U. mutabilis*, *U. subfusca*, *U. subgracilis*, and *U. subscabrosa* were sequenced using Illumina data and then assembled *de novo*. These mitogenomes ranged in size from 52,486 bp (*U. subfusca*) to 94,464 bp (*U. subgracilis*). All were characterized by having high levels of intronic and intergenic variation, such as ORFs that encode proteins with homology to two homing endonuclease types, LAGLIDADG and GIY-YIG. Genes annotated within these mitogenomes include 14 protein-coding genes, the large and small ribosomal subunits (LSU and SSU), and 23–26 tRNAs. Notably, the *atp9* gene was absent from each genome. Genomic synteny was highly conserved across the five species. Five conserved mitochondrial genes (*nad2*, *nad4*, *cox1*, *cox2*, and *cox3*) were used to infer a best estimate maximum likelihood phylogeny among these five *Usnea* and other relatives, which yielded relationships consistent with prior published phylogenies.

ARTICLE HISTORY

Received 12 February 2018
Accepted 13 February 2018

KEYWORDS

Symbiosis; *Usnea halei*;
Usnea mutabilis; *Usnea subfusca*; *Usnea subgracilis*;
Usnea subscabrosa

Introduction

Usnea is a genus of lichenized Ascomycete fungi that represents one of the most species-rich genera in Parmeliaceae, which is a family of more than 2000 species and 77 genera (Crespo et al. 2007; Lücking et al. 2016). *Usnea* has a cosmopolitan distribution, being found on every continent (Halonen 2000), and contains both saxicolous (rock-dwelling) and corticolous (bark-dwelling) species (Clerc and Herrera-Campos 1997; Clerc 1998; Halonen 2000). The genus on the whole is characterized by having a fruticose growth form and a cartilaginous central axis (Clerc 1998). Known colloquially as ‘Old Man’s Beard’, species of *Usnea* are charismatic and highly recognizable lichens across the globe (Brodo et al. 2001). Species are characteristically yellowish-green in colour due to the production of usnic acid, which is thought to have a multitude of potential functions including having antibiotic, antimycotic, and antiprotozoal properties (Lauterwein et al. 1995; Fournet et al. 1997; Giordano et al. 1997; Cocchietto et al. 2002). Here, we present newly sequenced and annotated mitochondrial genomes from five species of *Usnea* that occur in the southern Appalachian lichen biodiversity hotspot, located in eastern North America (Lendemer et al. 2013; Tripp and Lendemer in press; Tripp et al. in review). Two of these species (*U. halei* and *U. subfusca*) are endemic to eastern North America and have centres of distributions in the

region (Clerc and Herrera-Campos 1997; Lendemer et al. 2013).

Methods

Samples of *Usnea halei* (voucher Lendemer 46374; Lat/Long: 35.1356,–83.1911; NCBI accession MG722979), *U. mutabilis* (voucher Lendemer 49260; Lat/long: 34.3679,–85.6299; NCBI accession MG920803), *U. subfusca* (voucher Lendemer 46309; Lat/Long: 35.61,–83.4467; NCBI accession MG720812), *U. subgracilis* (voucher Lendemer 48717; Lat/Long: 35.1036,–83.2075; NCBI accession MG720066), and *U. subscabrosa* (voucher Lendemer 46747; Lat/long: 35.4372,–83.7481; NCBI accession MG720452) were collected by J.C. Lendemer, E.A. Tripp, K.G. Keepers and K.H. White from the Southern Appalachian Mountains in the United States between 2015 and 2017. Genomic DNA was extracted using a Qiagen DNeasy 96 plant kit, with the protocol modified to include a 10 min 65 °C incubation step for ground material in lysis buffer and a 100% ethanol wash before final drying of the membrane prior to elution. Genomic libraries were prepared using Nextera[®] XT DNA library prep kits (Illumina[®]) and each sample was barcoded using the unique dual index adapters Nextera[®] i5 and i7. Samples that passed QC were processed for paired-end 150 base pair read sequencing on the Illumina

CONTACT Kyle G. Keepers  kylekeepers@gmail.com  900 Pleasant Street, Ramaley Bldg Rm N395, Boulder, CO 80309, USA

*These authors contributed equally to this work.

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

NextSeq[®] at University of Colorado's BioFrontiers Institute Next-Generation Sequencing Facility in Boulder, Colorado. Genomic reads were trimmed using Trimmomatic-0.36 with the parameters 'ILLUMINACLIP:NexteraPE-PE.fa:2:20:10 MINLEN:140 LEADING:20 TRAILING:20' (Bolger et al. 2014). Trimmed reads were assembled *de novo* using SPAdes v.3.9, with the parameters '-meta -k 35,55,85' (Bankevich et al. 2012). Fungal mitochondrial contigs were identified using a command-line BLAST against a set of fungal mitochondrial proteins derived from the lichenized fungus *Peltigera dolichorrhiza* (NCBI accession KT946595), which were then circularized and error-corrected. Annotations of genomic features were initiated using DOGMA (Wyman et al. 2004) and completed in NCBI's Sequin 15.10 (Bethesda, MD). Genome assemblies and annotations were performed by undergraduate and graduate students enrolled in N. Kane's Genomics class at the University of Colorado, Boulder.

Phylogenetic analysis

A maximum likelihood phylogenetic hypothesis was constructed using sequence data from five loci (*cox3*, *nad2*, *nad4*, *cox1*, and *cox2*), which were concatenated prior to analysis. Outgroup sequences were obtained from NCBI: *O. vulgata* [KY315997], *P. ostiolata* [KY346830], *G. americanus* [NC_034790], *H. speciosa* [KY328643], *L. hirsutum* [NC_034928], *P. corallina* [NC_034779], *C. rangiferina* [KY460674], *C. uncialis* [KY352404], *P. stipiteum* [KY362439], *I. aleurites* [KY352227], *H. vittata* [KY362374], *M. subsimilis* [KY352491], *A. fallacina* [MG711470], *U. cornuta* [KY100278], *U. ceratina* [KX987159], and *U. pensylvanica* [KY321923]. Sequences were aligned in MEGA v.7.0.26 (Kumar et al. 2016) using the MUSCLE aligner (Edgar 2004). The resulting alignment was manually manipulated to exclude regions of the alignment that contained excessive missing data (available at Dryad Zenodo, record 1185346). Gaps were treated as missing data. The ML tree was inferred using RAxML v.8.2.10 (Stamatakis 2014). To assess support among relationships, we conducted 250 bootstrap replicates using the autoMRE criterion in RAxML and from this distribution, a consensus topology was generated. Bootstrap support values were mapped onto the most likely tree deriving from the above search. The tree was rooted using *Opegrapha vulgata*.

Results and discussion

Mitochondrial genome content and organization

The lengths of *Usnea* mitochondrial genomes in this study varied from 52,486 bp (*U. subfusca*) to 94,464 bp (*U. subgracilis*). The mitochondrial genomes of the other three species were of intermediate lengths: 61,314 bp (*U. mutabilis*), 78,464 bp (*U. subscabrosa*), and 82,851 bp (*U. halei*). The genomes presented here contained a conserved set of 14 protein coding genes (*cob*, *cox1*, *cox2*, *cox3*, *nad1*, *nad2*, *nad3*, *nad4*, *nad4L*, *nad5*, *nad6*, *atp6*, *atp8*, and *rps3*). Notably, all lacked a key mitochondrial gene (*atp9*), which has previously been shown to have been lost evolutionarily from three different lineages of lichens, one of which includes Parmeliaceae (Pogoda et al. 2018).

Synteny of protein-coding genes in the mitogenomes was highly conserved across the five species of *Usnea*, with no differences in overall gene order and no inversions detected.

Introns and intergenic spacer regions

Among the five samples, non-coding DNA contributed to overall genome size: *U. subfusca* has the smallest mitochondrial genome and also the least amount of intronic sequence. The three largest genomes (*U. subscabrosa*, *U. halei*, and *U. subgracilis*) contained variable lengths of intronic sequences, but all had approximately two to four times more total intergenic sequence compared to the other genomes. Notably, the three largest genomes also contained a large intergenic region between *atp8* and *cob*, which contained multiple fragments of a DNA polymerase (*dpo*) and ranged in size from approximately 4 kb in *U. subgracilis* to 17.9 kb in *U. subscabrosa*. All five genomes contained a large intergenic gap between the LSU and *nad2* genes, ranging in size from 4.8 to 7.7 kb and containing 7–13 tRNA sequences. *Usnea subscabrosa* also contained an ORF with homology to a GIY-YIG homing endonuclease in this region.

Open reading frames

Usnea mutabilis had multiple ORFs throughout its mitogenome. The longest ORF (1193 bp) was found preceding *nad5* and is likely co-transcribed with a 3' ORF that shares the stop codon of the protein coding gene. Two hypothetical proteins were located between *nad4* and *rps3* and between *cox1* and *nad4*, having lengths of 311 bp and 758 bp, respectively. We found that *U. mutabilis* possessed three ORFs with homology to homing endonucleases located in the introns of the *cox1* and LSU genes. *Usnea subfusca* was found to have two ORFs, one with homology to a GIY-YIG homing endonuclease that was 1445 bp and was located at the 3' end of the *nad1* gene.

Phylogenetic history

All eight species of *Usnea* (the five presented here, along with three previously sequenced species obtained from NCBI) were recovered as monophyletic with 100% bootstrap support (Figure 1). Monophyly of Parmeliaceae as a whole was also well supported (BS = 100%), with *Imshaugia*, *Hypogymnia*, *Menegazzia*, and *Alectoria* recovered as sister to *Usnea*. This topology is in general agreement with previously published phylogenies (e.g. Crespo et al. 2010; Miadlikowska et al. 2014).

Conclusions

The five newly sequenced, assembled, and annotated *Usnea* mitogenomes exhibited a large range in size, with intronic and intergenic regions of variable lengths associated with changes in genome size. These mitochondrial genomes coded for a variable set of ORFs, but all contained the same set of 14 protein-coding genes. Results from our study confirm earlier findings (Pogoda et al. 2018) that members of Parmeliaceae have lost a key gene involved in energy




Figure 1. Phylogenetic relationships among *Usnea* and other lichens inferred using maximum likelihood (see Methods section). Numbers next to nodes represent bootstrap support mapped onto the most likely tree ($-\ln L = 68803.11$). The long branch leading to *Usnea* was shortened (hashes) for the figure.

production, *atp9*, lending genomic evidence of the obligate nature of this symbiotic relationship. We thus infer that these five species of *Usnea* are likely dependent on their photosynthetic partner(s) for energy production. Finally, these data provide an important resource for further study of the evolution of parasitic, selfish elements such as LAGLIDADG and GIY-YIG type homing endonucleases in *Usnea* and related lichens. The present study yields new resources for further study of lichen mitochondrial genome evolution and co-evolution with obligate symbiotic partners.

Acknowledgements

Special thanks to the students of University of Colorado Boulder's Genomics classes of 2016 and 2017 for assembling and annotating the genomes used in this study. Publication of this chapter was funded by the University of Colorado Boulder Libraries Open Access Fund.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content of this paper.

Funding

This work was supported by a grant from the National Science Foundation's Dimensions of Biodiversity Program [award #1542639], University of Colorado award [#1432629] and New York Botanical Garden.

References

- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 19:455–477.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 30:2114–2120.
- Brodo IM, Sharnoff SD, Sharnoff S. 2001. *Lichens of North America*. Conn, New Haven: Yale University Press.
- Clerc P, Herrera-Campos MA. 1997. Saxicolous species of *Usnea* subgenus *Usnea* (lichenized ascomycetes) in North America. *Bryologist.* 281–301.
- Clerc P. 1998. Species concepts in the genus *Usnea* (lichenized Ascomycetes). *Lichenologist.* 30:321–340.
- Cocchietto M, Skert N, Nimis P, Sava G. 2002. A review on usnic acid, an interesting natural compound. *Naturwissenschaften.* 89:137–146.
- Crespo A, Kauff F, Divakar PK, del Prado R, Pérez-Ortega S, de Paz GA, Ferencova Z, Blanco O, Roca-Valiente B, Núñez-Zapata J, et al. 2010. Phylogenetic generic classification of parmelioid lichens (Parmeliaceae, Ascomycota) based on molecular, morphological and chemical evidence. *Taxon.* 59:1735–1753.
- Crespo A, Lumbsch HT, Mattsson JE, Blanco O, Divakar PK, Articus K, Wiklund E, Bawingan PA, Wedin M. 2007. Testing morphology-based hypotheses of phylogenetic relationships in Parmeliaceae (Ascomycota) using three ribosomal markers and the nuclear RPB1 gene. *Mol Phylogenet Evol.* 44:812–824.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32:1792–1797.
- Fournet A, Ferreira ME, de Arias AR, de Ortiz ST, Inchausti A, Yalaff G, Quilhot W, Fernandez E, Hidalgo ME. 1997. Activity of compounds

- isolated from Chilean lichens against experimental cutaneous leishmaniasis. *Comp Biochem Physiol C: Pharmacol Toxicol Endocrinol.* 116:51–54.
- Giordano S, Basile A, Lanzetta R, Corsaro MM, Spagnuolo V, Castaldo Cobianchi R. 1997. Potential allelochemicals from the lichen *Cladonia foliacea* and their in vitro effects on the development of mosses. *Allelopathy J.* 4:89–100.
- Halonen P. 2000. Studies on the lichen genus *Usnea* in East Fennoscandia and Pacific North America (PDF). Oulu, Finland: Oulu University Library; p. 13. ISBN 9514255232. ISSN 0355-3191.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 33:1870–1874.
- Lauterwein M, Oethinger M, Belsner K, Peters T, Marre R. 1995. In vitro activities of the lichen secondary metabolites vulpinic acid, (+)-usnic acid, and (–)-usnic acid against aerobic and anaerobic microorganisms. *Antimicrob Agents Chemother.* 39:2541–2543.
- Lendemer JC, Harris RC, Tripp EA. 2013. The lichens and allied fungi of Great Smoky Mountains National Park. *Memoires NY Bot Gard.* 104:1–152.
- Lücking R, Hodkinson BP, Leavitt SD. 2016. The 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota – approaching one thousand genera. *Bryologist.* 119:361–416.
- Miadlikowska J, Kauff F, Högnabba F, Oliver JC, Molnár K, Fraker E, Gaya E, Hafellner J, Hofstetter V, Gueidan C, et al. 2014. A multigene phylogenetic synthesis for the class Lecanoromycetes (Ascomycota): 1307 fungi representing 1139 infrageneric taxa, 317 genera and 66 families. *Mol Phylogenet Evol.* 79:132–168.
- Pogoda CS, Keepers KG, Lendemer JC, Kane NC, Tripp EA. 2018. Reductions in complexity of mitochondrial genomes in Lichen-Forming fungi shed light on genome architecture of obligate symbioses. *Mol Ecol.* doi:10.1111/mec.14519.
- Stamatakis A. 2014. RAxML Version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 30:1312–1313.
- Tripp E, Lendemer JC, McCain CM. in review. Impacts of disturbance on lichens – obligate symbiotic organisms—in a temperate biodiversity hotspot. *Divers Distrib.*
- Tripp EA, Lendemer JC. in press. Field guide to the Lichens of Great Smoky Mountains National Park. Knoxville (TN): University of Tennessee Press.
- Wyman SK, Jansen RK, Boore JL. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics.* 20:3252–3255.