



**Dr. C. Srinivasan**

---

**Studies in  
The Mosses of  
South India**

# STUDIES IN THE MOSSES OF SOUTH INDIA

**Dr. C. SRINIVASAN**, M. Sc., Ph. D.  
Reader, Department of Botany  
Annamalai University



**ANNAMALAI UNIVERSITY**  
**ANNAMALAINAGAR**

# STUDIES IN THE MOSSES OF SOUTH INDIA



**Dr. C. SRINIVASAN**, M.Sc., Ph.D.  
Reader, Department of Botany  
Annamalai University



**ANNAMALAI UNIVERSITY**  
**ANNAMALAINAGAR**

First Edition: March, 1974  
Annamalai University Publication  
Published under U. G. C. Grant

Thesis submitted to the Annamalai University  
by the author for the degree of Ph.D. in 1962

Printed at  
Sivakami Printers, Annamalainagar.

## CONTENTS

I. PREFACE	iii
II. INTRODUCTION	1
III. MATERIALS AND METHODS	4
IV. MORPHOLOGICAL OBSERVATIONS:	
1. Systematic description	8
2. General considerations of:	
(a) The leaf	39
(b) The axis	47
(c) The sporophyte	48
(d) Vegetative reproduction and regeneration	72
V. ANATOMICAL OBSERVATION	75
1. Description	76
2. Results	90
VI. PHYSIOLOGICAL OBSERVATIONS	100
VII. ECOLOGICAL OBSERVATIONS	112
VIII. DISCUSSION:	
1. Morphological and Physiological peculiarities of mosses in relation to their ecology:	
(a) Vegetative adaptation	132
(b) The sporophyte	134
(c) Osmotic concentration	137
(d) Ecological factors and their significance	138

2. Taxonomical considerations on the basis of anatomical observations:	
(a) Structure of the moss axis	139
(b) Systems of classification	140
IX. SUMMARY	142
X. ACKNOWLEDGEMENTS	146
XI. INDEX	146
XII. LITERATURE	153
XIII. EXPLANATION OF PLATES	167

---

## PREFACE

Studies in the morphology, anatomy, physiology and ecology of 64 different species of South Indian Mosses have been made. The contribution recorded herein have been made for the first time.

Systematic description of 64 species of more common South Indian Mosses has been attempted in detail. Critical study reveals that the structure of the leaves appears to be well-adapted for the retention of water. The foot of the sporophyte has been found to be of 8 different forms. The L/B (Length/Breadth) ratio of the stomata observed on the apophysis of 8 species appears to be a constant feature. The stomatal index has been found to be specific.

The transverse sections of the axes of most of the species (53) have been described in a systematic manner. On a close study of the internal structure of the moss axes, it has been possible to classify them. The proposed system of classification though based purely on anatomical peculiarities of the gametophyte is in conformity with the prevailing systems of classification, in some respects.

The sporophytes of 10 species of mosses have been induced to produce rhizoids from the base of their foot by treatment with hormones and gametophytic-extract just to show that the moss sporophytes under proper cultural conditions could lead an independent existence for a short period. Secondly the osmotic concentration of the cell-sap of the cells of the sporophytes of

7 species has been found to be always higher than that of the gametophytes. Vegetative reproduction and regeneration in 11 species have been studied and recorded.

Ecological studies in the physiographic, edaphic and biotic factors in respect of the South Indian mosses and their formations and associations and relative frequency of some of the species have been made.

It is in these and in other details that the present investigations seek to be an attempt towards advancement of knowledge in Bryology.



FRONTISPICE: ASSOCIATION OF PAPILLARIA CRUCEA, FLORIBUNDARIA  
FLORIBUNDA AND METEOROPSIS RECLINATA FORMING FENCORIS

# Introduction

The study of the South Indian mosses was undertaken on the kind suggestion of Professor T.C.N. Singh, Head of the Department of Botany of the Department of Botany of the Annamalai University. While he had been working on the vegetative reproduction in certain mosses (Singh, 1930) under the guidance of Professor Birbal Sahni, F.R.S. of the Lucknow University, it transpires that the learned Professor had expressed a desire that this long neglected field of Botany, specially Indian mosses, should be carefully studied. Perhaps, it is almost the posthumous desire of that great Professor that has fructified in a very small measure in the wake of the present investigation. The present study is based on 64 species, comprising of 55 genera spread over 24 families.

In the first instance I was introduced to the census of Indian mosses published by Brühl in 1931\*. He has rightly pointed out the limitations of an Indian teacher of the plains for a work of this kind in the study and collection of mosses which grow in the mountainous regions.

The present work is obviously an attempt: (i) to describe the more common mosses of this part of the

---

\* In 1930, Professor T.C.N. Singh had placed his entire large collection of Mosses of Simla Hills (North-Western Himalayas) at the disposal of Dr. P. Brühl.

country in detail; (2) to relate as far as possible their ecological peculiarities and their physiological behaviour; and (3) to submit a thought for the classification of the mosses on the basis of anatomical details

There is hardly any detailed work on the South Indian mosses. The earliest reference on these plants is by Brotherus (1899). He, in his report on a collection of mosses made by Walker in Coorg during the cold weather of 1897-1898, says that "the Bryological Flora of South India is very little known." He has enlisted 98 species of mosses of which 20 were described as new species. Cardot (1911), according to Foreau (1930), estimated the moss flora of the Palni Hills to consist of 191 species of which more than 50, he considered, were new to science. Dixon (1914) in his report on the mosses collected by Fischer from South India, has enumerated 58 species of which 6 were described as new species. Of these mosses 11 species find a place in the present investigation. Dixon (1923) has also enlisted 43 species of mosses collected in 1921 from the Kanara district of South India by Sedgewick and also of those received from Mons. Cardot. He has stated that he had in his herbarium, "numerous unpublished species of mosses from various parts of India including the types of a considerable number of new species mostly from the Madura district of South India." It is these mosses that form the bulk of the present treatise. Only 3 species out of his report are described in the present work. Subsequently Potier de la Varde (1922), (1925), (1928 a), (1928 b); Cardot and Varde (1922); Dixon and Varde (1927, 1928) have listed the South Indian mosses and

described some new species. It is interesting to note that Dixon and Varde (1927), (1928) have erected 2 new genera of mosses from this part of India, represented by *Trigonodyction indicum* Dix. et Vard. and *Foreauella indica* Dix. et Vard.\* Foreau (1930) has published an interesting and very useful note on the Bryological Geography of the Palni Hills and enumerated the mosses found there. Brühl (1931) in his census of Indian Mosses brought out a very comprehensive list of mosses occurring not only in India but also in Ceylon and Malaya Peninsula. His analytical key to the genera and their description were of some help in identification. Sedgewick (1947-1950) published a list of mosses of the Bombay presidency into which region I have not extended this study. Very recently Foreau (1961) has published a list of 368 species of moss flora of the Palni Hills with the exact places and dates of collection along with the Greek and Latin derivations of the generic names of the mosses.


There is, therefore, hardly any work on the South Indian mosses and the observations presented in this communication comprise of details being recorded for the first time.

---

\* This genus has been named after Rev. Fr. G. Foreau who has been a very painstaking muscologist who in spite of his old age not only showed me the natural home of the mosses but also helped me greatly in collecting them during my periodical excursions to the Kodaikanal Hills.

## Materials And Methods

About 100 different species of mosses were collected from the hills of Palni, Ginjee, Yercaud and Tirupati and from the mountainous regions of Kodaikanal (Madura district) and Ootacamund (Nilgiri district). A few were obtained from the hills in Trivandrum and Mysore. Out of these, only 64 species spread over 55 genera have been included in the present study. The collection was made in different seasons such as wintry days of November and December and the summer days of April and May. Some mosses were collected during September and October also when they were observed with their capsules. The plants were preserved dry in small paper packets. Herbaria of these plants were prepared for facilitating identification. For detailed study, the leaves were carefully removed under the stereo-microscope and mounted in a special medium consisting of 90 c.c. of 30% glycerine, 2 c.c. of 1% aqueous safranin and 8 c.c. of F. A. A. (5 : 5 : 90 of 50%). This mixture keeps the material not only well spread and stained suitably but also free from the attack of insects and fungus. Similarly, entire plants with the capsule wherever available, were mounted. The ripe capsule was ripped open and the peristome with the capsular wall, the entire apophysis region, the spores and the rhizoids were all mounted in a similar manner.



MAP OF SOUTH INDIA

The foot of the seta was carefully removed from the apex of the gametophore with a pair of very fine needles in living condition and mounted thereafter.

The usual process of fixing in F. A. A. and dehydration in chloroform and sectioning with rotary microtome was adopted for the sections of the axes and leaves. They were cut to thickness of  $10\mu$  to  $12\mu$ . Double embedding in celloidin and later in paraffin was also done specially for sectioning longitudinally the foot of the seta while still attached to the gametophore. Considerable amount of difficulties were experienced in the matter of infiltration and microtomy. The leaves shrink in absolute alcohol and the microtome sections when they become dry, shrivel and roll upwards from the paraffin wax and fall off during the processes of staining. But this difficulty could be obviated by cutting out the material towards one end with a fine sharp blade and thus exposing it while still in the block and then dipping the block either in water or in 50% glycerine for a considerable time and thus softening the material. Double staining in safranin and fast green gave contrasting differentiation. Free hand sections stained suitably and mounted in the glycerine mixture mentioned above were largely used. Staining in phlo - roglucin, methylene blue were also tried. The axes of some of the mosses were treated with Schultze's maceration fluid and the cells were subjected to micro-chemical tests.

Certain growth hormones such as I. B. A.; I. A. A.; I. N. A.; and 2, 4-D were used in different concentrations of .5 p. p. m. to 20 p. p. m. in order to see whether the sporophytes and leaves (gametophyte) produce rhizoids.

These experiments were conducted at both the hill stations of Kodaikanal and at Ootacamund in two different seasons (October and April). The same growth hormones excepting I. N. A. were applied to the local moss (*Barbula indica*) which grows very luxuriantly during winter in the plains and also on the terrace of the laboratory. For these mosses buffer solutions of different pH namely 4 to 9 were sprayed in their natural habitat to find out whether these substances promote or affect the growth of the mosses.

Osmotic concentrations within the gametophyte in the young leaves and transverse sections of axis and those of the sporophyte in the wall cells of the apophysis region of the capsule and also of the seta were determined for some of the mosses by the de Vries' plasmolytic method using the solutions of Sodium chloride, Potassium nitrate and Dextrose in different concentrations. The pH of the peaty soil formed by the local moss, *Barbula indica*, on the terrace of buildings and the soil of some of the soil mosses were determined with the help of La Motte Morgan Soil Testing set. The nitrogen content of a rock moss was examined in our Bio-chemical Section of the department. The results are given in the chapter on physiological observations.

Some of the results of these investigations have, from time to time been contributed as abstracts, being read in the Botany Section of the Indian Science Congress Association (Srinivasan, 1958, 1960a, 1960b, 1961).

The mosses without sporophytes (described in the chapter on morphological observations) are indicated with asterisks. All the diagrams and photographs have

been numbered in a serial manner. The photographs of the mosses studied, have been arranged in plates (Figs. 477-535). A few species of mosses were also photographed in their natural habitat and included (Frontispiece and Figs. 538-549). Photomicrographs in support of some of the diagrams have been appended (Figs. 550-584). The diagrams have been drawn with Zeiss Winkel camera lucida using Zeiss Winkel Research Microscope. The magnifications are noted against each. Wherever the laminar cells are enlarged, exact region of the leaf is pointed out with an arrow. Whenever the cells are different in size and shape in one and the same leaf and when they are enlarged as separate diagrams, those different regions of the leaf, are indicated by means of arrows and marked 'a' and 'b'.

# Morphological Observations

## Fissidentaceae

1. *Fissidens schimidii* C. M.: Plants grow in small tufts on clayey ground in the moist woods of Shembaganur (Kodaikanal) at an altitude of 2000 metres or on wet soil-covered-rocks in shaded regions of Trivandrum 300 metres forming thin mats, bright to dark green. Soil acidic, pH being 5.4 to 5.6 Plants (Figs. 1 & 2): very small slender, 8 mm to 10 mm high. Leaves: bifarious, stem-clasping; .88 mm to 1.21 mm long and .09mm to 13mm broad, dorsal lamella strongly developed; apex acute; leaf-cells (Fig.4) rectangular to hexagonal, 8 cells in a square of  $22\mu$ ; margin bordered by one row of elongated rectangular cells; mid-rib thin. Sporophyte: terminal or lateral; seta 8 mm to 10 mm long; Spore-capsule sub-erect to inclined, .9 mm to 1.2 mm long and .3 mm to .4 mm broad; peristome (Fig. 6) deeply bipartite  $264\mu$  long and  $22\mu$  broad at base operculum (Fig. 357) long beaked; spores (Fig. 386) spherical  $11\mu$  to  $13\mu$  in diameter.

## Ditrichaceae

2. *Ditrichum amoenum* (Thw. and Mitt.) Par.: Plants grow crowded on moist clayey soil on road-side near Observatory (Kodaikanal) 2300 metres: or in shaded cave sides at Fernhill (Ootacamund) 2300 metres. Soil

acidic, pH being 5.2 to 5.4. Plants (Figs. 7 & 8) : bright yellowish green, fruiting abundantly ; stem erect and short 10 mm to 15 mm with capsule ; gametophyte 3 mm to 5 mm ; leaves (Fig. 9) hairy 3.8 mm to 6 mm long and .3 mm to .4 mm broad at base ; apex attenuate ; leaf-cells ( Fig. 10 ) elongated rectangular, 3 cells in a square of 22 . Capsules (Fig. 11) 2 mm to 2.3 mm long and .4 mm to 5 mm in diameter ; operculum (Fig. 358)  $17\mu$  to  $190\mu$  long ; peristomial teeth (Fig. 12) slender thread like  $232\mu$  long with minute marginal papillae ; spores (Fig. 387) spherical,  $15\mu$  to  $17\mu$  in diameter.

3. *Trematodon ceylonensis* C. M. : Plants grow in swampy regions of Perumal hill shola (Kodaikanal) 1600 metres: plants (Fig. 13&14): 20 mm to 30 mm in height with the sporophyte, greenish yellow and shiny. Leaves (Fig. 15): 2.5 mm to 4 mm long and .4 mm to .5 mm broad at base, lanceolate subulate; apex revolute; margin slightly serrulate; leaf-cells (Fig. 16) rectangular, cells of the costa narrowly rectangular. Sporophyte: 20 mm to 25 mm long; seta erect yellowish and shiny; capsule (Fig. 17) 7 mm to 9 mm  $\times$  .5 mm to .7 mm with much longer neck; operculum (Fig. 359) and calyptra hooded peristome teeth (Fig. 19) entire; apophysis provided with stomata (Fig 18) resembling those of dicots but with numerous subsidiary cells varying from 7 to 8; spores (Fig. 388) spherical,  $22\mu$  to  $24\mu$  in diameter.

#### Dicranaceae

4. *Campylopus nodiflorus* (C. M.) Jaeg. : Plants grow on the bark of tree trunks in thick forests

of Silvercaskade (Kodaikanal) 2200 metres, Thoddabedda (Ootacamund) 2800 metres. Plants (Fig 20) 40 mm to 45 mm tall. Leaves 3 mm to 5 mm long .3 mm to .4 mm broad at base (Fig. 21), ovate and attenuated into straight long tip; apex (Fig. 22) with serrated margin; leaf-cells (Fig. 23)  $50\mu$  to  $55\mu \times 20\mu$  to  $22\mu$ , elongated; alar cells (Fig. 24) rectangular, marginal cells narrow and elongated. Seta unusually short and bent variously 8 mm to 10 mm long; capsule (Fig. 25) 1.4 mm to 1.75 mm long and .8 mm to 1 mm broad, spherical and drooping with long beaked operculum (Fig. 360); peristome (Fig. 27) teeth bipartite, hair-like, united at base, annulus (Fig. 26) specialised ring of vesicular cells flask-shaped; spores (Fig 389)  $15\mu$  to  $17\mu$  in diameter.

5. *Thysanomitrium depallieri* Card. \* : Plants grow on flat rocky, water-oozing substratum, in bright sunshine in the Priests' Walk (Kodaikanal, 2000 metres. Plants (Fig. 33) dark green, gametophytes 9 cm to 12 cm long. Leaves (Fig. 34): 6 mm to 9 mm long, 1.5 mm to 2 mm broad, slightly concave passing into a lanceolate upper part ending in a hair point; apex (Fig. 30) with fimbriate margin; leaf-cells (Fig. 32) almost rectangular  $33\mu$  to  $35\mu \times 16\mu$  to  $18\mu$ ; marginal and mid-rib cells narrower; alar cells quadratic (Fig. 31). Profuse unbranched rhizoids (Fig. 34) found at the base of leaves.

6. *Thysanomitrium lioneurum* (Ther. and Vard.) Broth. :\* Plants grow in dense patches on ground or rock near Thoddabedda (Ootacamund) 2600 metres and on waysides of Tigershola (Kodaikanal) 2100 metres.

Plants (Fig. 28) 4 cm, to 4.5 cm tall. Leaves (Fig. 29): 3 mm to 5 mm long and .5 mm to .75 mm broad at base, oblong passing into a lanceolate upper part ending in a blunt point (Fig. 35); leaf-cells (Fig. 36) elongated rhomboid; cell-walls thickened; alar cells differentiated; marginal cells thin in three rows; mid rib broad, cells typically rectangular and thinner.

7. *Brothera leana* (Sull.) C. m. \* : Plants grow closely on rotting tree trunks at Perumal hill road (Kodaikanal) 1675 metres, 7 mm to 10 mm tall (Fig. 38). Leaves (Fig. 39): lanceolate 3 mm to 5 mm long and .25 mm to .4 mm broad at base; margin entire; apex (Fig. 40) pointed, leaf cells (Fig. 41) hyaline, rectangular.

8. *Dicranodontium fragile* (Hook.) Broth. : Plants grow on shaded rocks and on decaying wood in the woods of Priests' Walk (Kodaikanal) 1700 metres in dark green tufts and small patches, 4 cm to 4.5 cm tall (Fig. 42). Leaves (Fig. 43) form a broad concave base, linear oblong 1 mm to 3 mm long and .6 mm to .8 mm broad at base; costa in narrow seam; cells of the lamina (Fig. 45) rectangular in linear stereids; cell wall thickened; alar cells broad rectangular, differentiated, forming 6 to 7 rows; apex (Fig. 44) prolonged with serrated margin. Seta at first decurved, finally erect 10 mm to 12 mm long; capsule (Fig. 46) 5 mm to 6 mm  $\times$  .8 mm to 1 mm, cylindric; operculum (Fig. 361) long beaked; peristome (Fig. 47) teeth 16 each with 9 to 10 basal partitions and bipartite above; spores (Fig. 39 ) spherical,  $25\mu$  to  $27\mu$  in diameter.

## Leucobryaceae

9. *Leucobryum neilgherrense* C. M. : Plants grow in small pale green clusters on tree trunks, rocks and shady ground in various parts of South India; Jinge hills (South Arcot) 250 metres, Old Ghat Road, (Kodaikanal) 1870 metres, Simp's Park (Ootacamund) 2500 metres. Soil acidic, pH being 5.6 to 5.8 plants (Fig. 48) 15 mm. to 18 mm. tall, green. Leaves (Figs 48 & 49) thick, narrow linear 3.5 mm to 5 mm long .5 mm to .75 mm broad. whitish when dry; costa differentiated; apex (Fig. 50) acute; laminar cells (Fig. 52) slightly elongated rectangular, chloroplasts light and pale green. Rhizoids found to arise from the basal, marginal and apical cells of the leaf also. Sporophyte terminal; seta short 4 mm to 5 mm long; capsule (Fig. 53) erect or inclined slightly, sub-ellipsoid 2.5 mm to 4 mm  $\times$  .7 mm to .9 mm; lid conical (Fig. 362); peristome teeth (Fig. 54) subulately bifid; spores (Fig. 390) spherical  $18\mu$  to  $20\mu$  in diameter.

10. *Leucobryum scalare* C.M. ; Plants grow in light green tufts on tree trunks in the forest of the Priests' Walk (Kodaikanal) 1700 metres and in the Sim's Park (Ootacamund) 2000 metres. Gametophytes (Fig. 55) 3 cm to 4 cm tall, branching from base in clusters of 4 to 6 fastigiate. Leaves (Fig. 56) thick, long and narrow 3.5 mm to 5 mm long and 1.5 mm to 2 mm broad; apex (Fig. 57) acute with serrulate margin; laminar cells (Fig. 59) rectangular  $44\mu \times 30$  costa not differentiated; marginal cells thin elongated.

## Pottiaceae

11. *Tortula muralis* Hedw.: Plants grow on rocks and road sides near the wood house (Ootacamund) 2750 metres in small dark green cushions about 1 cm. tall (Figs. 60 & 61). Leaves (Fig. 62) 2.5 mm to 3.5 mm long and .4 mm to .5 mm broad, keeled and ligulate; margins incurved; costa (Fig. 63) terete, prolonged beyond the leaf blade as a hyaline hair  $440_{\mu}$  to  $450_{\mu}$  long; leaf-cells (Fig. 64) rectangular, upper leaf cells smaller, cells of the costa narrow and long. Seta purple and erect 11 mm to 13 mm long; capsule cylindric (Fig. 65) 4 mm to 4.2 mm  $\times$  .5 mm to .6 mm; operculum (Fig. 363) long conical about 2 mm long beyond the capsule head; peristome (Fig. 67) with 32 filiform long twisted teeth (Fig. 112); spores (Fig. 394) spherical, sparsely granulose  $9_{\mu}$  to  $10_{\mu}$  in diameter.

12 *Barbula indica* (Schw.) Brid: Common in acidic soil, the pH being 6.2 to 6.4; cosmopolitan in distribution in the plains and in the hills; growing gregariously on old brick-walls, terrace of buildings and on moist rocks. Hill plants grow taller but usually do not fruit. Plants (Figs. 68&69) yellowish green 1 cm to 2 cm tall; rhizoids branched with oblique cross walls. Leaves (Fig. 70): spirally alternate, oblong .8 mm to 1.5 mm long and .5 mm to .75mm broad; apex (Fig. 71) blunt; laminar cells (Fig. 72) rectangular; cells of costa narrower. Sometimes club-shaped gemmae (Fig. 73) produced in clusters in the axils of upper leaves; obovate multicellular with a basal cell,  $30_{\mu}$  to  $50_{\mu}$  long and  $14_{\mu}$  to  $18_{\mu}$ . broad. Sporophyte terminal or lateral; seta erect 5 mm to 7 mm long; capsule

(Fig. 74) ovoid 2 mm  $\times$  .15 mm; peristome (Fig. 77&78) filiform with fimbriate margin; operculum (Fig. 364) conical; stomata present on apophysis, guard cells two (Fig. 75) sometimes the guard cells of the stomata in the lower regions near the seta divided by cross striations (Fig. 76); spores (Fig. 393) greenish, spherical,  $15\mu$  to  $16\mu$  in diameter.

13. *Trichostomum cylindricum* (Bruch.) C. M. : Plants bright green, grow on ground and on moist rocks of Shenbaganur (Kodaikanal) 1800 metres; 20 mm to 25 mm tall (Figs. 79 & 80). Leaves (Fig. 81) 3 mm to 4 mm long and .7 mm to .8 mm broad at base, keeled ventrally concave, lanceolate, upper portion secund; apex (Fig. 82) sharply acute; alar cells (Fig. 84) elongated rectangular to quadratic  $44\mu$  long and  $29\mu$  broad, upper cells  $5\mu$  to  $6\mu \times 2\mu$  to  $3\mu$  only, cells towards the mid-rib still smaller, cells of costa narrow and elongated rectangular. Seta 8 mm to 10 mm long; capsule (Fig. 84) slightly inclined, cylindrical, 2.5 mm to 3 mm  $\times$  .5 mm to .7 mm operculum (Fig. 365) conical; peristome teeth (Fig. 85) entire; spores (Fig. 393) spherical, brownish, thinly granulose  $15\mu$  to  $16\mu$  in diameter.

14. *Weisia macrospora* Card. : Plants grow in mats on soil and in cracks of rocks in Pillar Rocks (Kodaikanal) 2000 metres, 10 mm to 15 mm tall (Figs. 86&87). Upper leaves (Fig. 88) usually larger than the lower, erect, spreading, much elongated, lanceolate; apex (Fig. 89) prolonged, acute; mid-rib well developed, lamina folded or canaliculate; leaf-cells (Fig. 90) ovate, cells of costa narrowly elongated. Seta erect, moderately long 8 mm

to 10 mm; capsule (Fig. 91) ovoid to ellipsoidal .5 mm to .75 mm  $\times$  .5mm; operculum (Fig. 366) long beaked; peristome (Fig. 92) inner teeth short in two rows, outer peristome not developed; spores (Fig. 406) spherical, 12 $\mu$  to 14 $\mu$  in diameter.

15. *Hyophila involuta* (Hook.) Jaeg. : Plants grow on walls, and on wet rocks in Perumal hills 2000 metres Shenbaganur (Kodaikanal); short 2 cm to 2.5 cm tall (Figs. 92 & 93). Axial column scarcely differentiated. Leaves (Fig. 94) oblong, 2 mm to 3 mm long and .5 mm to .7 mm broad; apex (Fig. 97) acute with serrate margin; leaf-cells (Fig. 98) 8 to 9 in a square of 22 $\mu$ , hexagonal near mid-rib, rectangular towards margin; margin differentiated by one row of ovate cells, cells of costa thin and elongated rectangular. Seta brownish 10 mm to 12 mm long; capsule (Fig. 95) cylindrical, 2.7 mm to 3 mm  $\times$  .5 mm to .6 mm; lid (Fig. 367) conical; annulus (Fig. 96) in two rows; spores (Fig. 403) roundish, 9 $\mu$  to 10 $\mu$  in diameter.

16. *Hyophila validinervis* Card. and Vard.\*. Plants grow in small tufts on soil and on rock in Priests' Walk (Kodaikanal) 2400 metres; 1.5 cm to 2 cm tall (Figs. 99 & 100). Leaves (Fig. 101) oblong; apex (Fig. 102) obtuse; laminar cells (Fig. 103) rectangular to quadratic towards the midrib, those towards the margin hexagonal; margin not much differentiated, cells quadrate; mid-rib prominent, cells linear. Branched rhizoids produced at the base of the leaves too.

## Grimmiaceae

17. *Rhacomitrium javanicum* Bryol.\* : Plants grow with creeping axis in rigid tufts on moist rocks of Priests' Walk (Kodaikanal) 1900 metres. Plants (Fig. 104) dark green, 5 cm to 8 cm tall. Leaves (Fig. 105) 2 mm to 2.2 mm long and .7 mm to .9 mm broad at base lanceolate; apex bluntly acute, leaf-cells (Fig. 106) long and narrow, alar cells with much thickened cell-wall, wavy longitudinal and thinner, transverse thickening characteristic; the margins usually recurved; costa prominent and broad.

## Funariaceae

18. *Funaria hygrometrica* Hedw.: Plants grow close together in small patches on ground in the woods of Tiger hill 2800 metres and in the open swampy soil at Kern hill (Ootacamund) 2300 metres. "Its presence marks the site of a recent fire". Plants pale green, gametophytes 1 cm to 2 cm tall but with sporophyte 3 cm to even 8 cm tall (Fig. 108). Leaves (Fig. 109) spatulate 2 mm to 2.2 mm long and 1 mm to 1.5 mm broad; margins often incurved and slightly wavy; leaf-cells (Fig. 110) large almost hexagonal. Sporophyte terminal: seta long greenish yellow when young, reddish brown later, curved and twisted 45 mm long; capsule (Fig. 111) pear shaped, held horizontally 3.5 mm to 4 mm  $\times$  .8 mm to 1.5 mm furrowed when dry, mouth oblique bearing inner and outer peristome (Figs. 111 & 113), the latter curved and united by their 16 fimbriate tips (Fig. 114) to a minute central disc; operculum convex with its characteristic spirally

arranged cells (Fig. 368); calyptra covering the capsule hooded at base, long and pointed above; spores spherical, thick-walled, brownish (Fig. 395)  $9\mu$  to  $10\mu$  in diameter.

19. *Funaria submarginata* Card. and Vard. : Plants grow in clayey soil 1 cm to 1.5 cm tall; (Figs. 115 & 116) in loose clusters on way side at Tiger hill (Ootacamund) 2700 metres. Leaves (Fig. 117) ovate elliptic, 1.8 mm to 2 mm long and .8 mm to 1.1 mm broad; apex acute; laminar cells rectangular to pentagonal; (Fig. 118) cells of costa thin and elongated, narrow cells constituting the margin. Sporophyte terminal: seta 12 mm to 14 mm long brownish and erect; capsule ovate, (Fig. 119); 1.2 mm to 1.5 mm  $\times$  .8 mm to 1 mm; operculum (Fig. 369) slightly convex and round; peristome wanting; stomata (Fig. 121) on the wall of the apophysis; spores (Fig. 396) large spherical, granulose,  $25\mu$  to  $27\mu$  in diameter.

### Bryaceae

20. *Pohlia flexuosa* (Mitt.) Broth : Plants grow in carpets on rocks in moist and shady places of Shempaganur (Kodaikanal) 1800 metres, (Figs. 122 & 123); 2.5 cm to 3 cm tall. Leaves (Fig. 124) lanceolate 2 mm long and .3 mm to .4 mm broad; apex (Fig. 125) attenuated with marginal serration and prolonged up to a fine point; leaf-cells (Fig. 126) elongated,  $99\mu$  to  $111\mu$ . Sporophyte terminal: seta inclined reddish 15 mm to 17 mm. long; capsule (Fig. 134) ovate 7 mm to 7.1 mm  $\times$  .9 mm to 1 mm; operculum (Fig. 370) convex ending in a sharp point; inner peristome  $60\mu$ , with outer peristome

also (Fig. 127); spores (Fig. 397) spherical,  $15\mu$  to  $17\mu$  in diameter.

21. *Brachymenium exile* (Doz. and Molk.) Bryol.: Plants dirty green; grow on soil on way sides to Perumal hill (Kodaikanal) 1600 metres or at Gudalur 2300 metres, or on the way to Thoddabedda (Ootacamund) 2800 metres. Soil acidic, pH being 6.2 to 6.4. Plants form loose tufts, 2 cm to 3 cm tall (Figs. 128 & 129). Leaves ovate (Fig. 130) 5 mm to .6 mm long and .15 mm to .17 mm broad; apex prolonged into a hair point, costa excurrent; laminar cells (Fig. 131) hexagonal, basal cells rectangular. Sporophyte lateral or terminal: seta 12 mm to 14 mm long and erect; capsule 2 mm to 2.1 mm  $\times$  .17 mm; operculum (Fig. 371) cupola shaped and apiculate peristome (Fig. 132) hyaline; spores (Fig. 398) spherical, small  $12\mu$  to  $14\mu$  in diameter, thinly granulose. Multicellular hooked gemmae (Fig. 418) were found in the axils of leaves.

22. *Brachymenium leptostomoides* (C. M.) Shimp.\*: Plants grow on tree trunks in the moist shaded forest of Priests' walk (Kodaikanal) 2250 metres. The gametophytes brownish green, .5 cm to 1 cm tall (Fig. 133 & 134). Leaves (Fig. 135). 6 mm to .7 mm long and .3 mm to .4 mm broad, oblong and spatulate with a marginal rim, costa strong and excurrent; apex (Fig. 136) prolonged and acuminate; laminar cells (Fig. 137) irregularly rhomboid  $33\mu$  to  $11\mu$ , marginal cells elongated and ovoid.

23 *Anomobryum subneditum* Card. and Vard.: Plants grow in the clefts of wet rocks and on the soil near

Pillar rocks (Kodaikanal) 2000 metres or at Golf-ground shola (Ootacamund) 2600 metres. Plants yellowish green 3 mm to 4 mm tall (Figs. 138 & 139). Leaves (Fig. 140) julaceous on the stem, oblong elliptic .5 mm to .6 mm long and .4 mm to .5 mm broad; mid-rib strongly developed, apex acuminate; leaf-cells elongated rhomboid and narrow (Fig. 142). Sporophyte lateral or terminal: seta red, 15 mm to 17 mm long and inclined; capsule ovate and elliptic 2.5 mm to 2.75 mm  $\times$  1 mm to 1.1 mm.; operculum (Fig. 372) apiculate slightly beaked; peristome teeth (Fig. 141) distinctly red with filiform upper portion; spores (Fig. 399) granulose, 15 $\mu$  to 17 $\mu$  in diameter.

24, *Bryum argenteum* Brid: These silvery mosses grow in small patches between bricks of walls or on culverts at Ootacamund and Kodaikanal 2000 to 2800 metres. Plants 2 cm to 3 cm tall cosmopolitan (Figs. 143 & 114). Leaves (Fig. 145) silvery grey in appearance due to lack of chloroplasts in the upper half, slightly concave, ovate, crowded and closely overlapping making the stem prettily julaceous; apex prolonged and bristle-like; leaf-cells (Fig. 146) quadratic to irregular rhomboid. Sporophyte terminal or lateral. Seta red, inclined above, 12 mm to 14 mm long; capsule pendulous, drooping, pear-shaped 3 mm to 3.1 mm.  $\times$  1.2 mm to 1.3 mm., when matured dark red, obovoid or sub-cylindrical, neck distinct; apophysis provided with stomata; operculum apiculate with blunt end (Fig. 373); peristome (Fig. 147) prominent 220 $\mu$  to 240 $\mu$ ; inner peristome slender and numerous; spores (Fig. 400) spherical with thick coat 12 $\mu$  to 14 $\mu$  in diameter, sparsely granu ose.

25. *Rhodobryum giganteum* (Hook.) Par. : Plants grow in moist shady forests of Silver Cascade (Kodaikanal) 1700 metres or in Kern hill Reserve Forest Ootacamund) 2500 metres, with a creeping stem on the soil and rosette of broad and spreading leaves above (Fig. 148); 4 cm to 7 cm tall. Leaves (Fig. 149) obovate 8 mm to 12 mm long and 4.5 mm to 5 mm broad, upper margin serrulate; apex (Fig. 150) acute with a hair-like projection; leaf-cells (Fig. 152) much elongated rhomboid with thickened walls,  $88\mu$  to  $22\mu$ , costa well developed excurrent. Sporophyte terminal: seta long erect 50 mm to 55 mm long; capsule (Fig. 151) cylindrical, horizontal to pendent 8 mm to 9 mm  $\times$  1.6 mm to 1.75 mm; operculum (Fig. 374) umbilicate, cupola shaped; peristome with many narrow partitions free and close (Fig. 153); spores spherical (Fig. 401)  $12\mu$  to  $14\mu$  in diameter.

#### Mniaceae

26. *Mnium coriaceum* Griff. : Plants grow sometimes singly or mixed with other mosses in moist and shaded slopes of soil covered rocks and on loamy soil at circular road (Ootacamund) 2600 metres or in Bear Shola (Kodaikanal) 2300 metres. Plants dark green and prostrate (Fig. 154). Leaves (Fig. 155) tongue-like, oblong 6.5 mm to 7 mm long and 2.5 mm to 3 mm broad with a thick undulate margin and strong costa; apex (Fig. 156) obtuse, nearly mucronate above; leaf-cells roundish 4 to 5 in a square of  $88\mu$  (Fig. 157). Sporophyte terminal or lateral; seta erect and elongated 40 mm to 42 mm long; capsule (Fig. 158) inclined 10.5 mm to 11 mm  $\times$  1.2 mm to 1.3 mm, lid conical; peristome teeth 16 free;

spores (Fig. 402) spherical with a thick coat, small,  $12\mu$  to  $14\mu$  in diameter.

Kabiersch (1937) treated this as a variety of *Mnium longirostrum* Brid. (*M. rostratum*). H. Ando (1961) says that this is closely related to Japanese *M. maximowiczii* Lindb.

### Rhizogoniaceae

27. *Rhizogonium spiniforme* (Hedw.) Bruch. : Plants grow on tree trunks in close tufts in the Priests' Walk (Kodaikanal) 2250 metres. Plants bright green, 3 cm to 4 cm tall (Fig. 160). Leaves (Fig. 161) narrow lanceolate 6 mm to 7 mm long and .4 mm to .5 mm broad at base, margin serrate; apex (Fig. 162) acuminate; leaf-cells thick walled, rectangular near the mid-rib and round or ovate towards the margin; 4 to 5 cells in a square of  $22\mu$  (Fig. 163). Sporophyte terminal or lateral; seta erect 35 mm to 37 mm long, reddish brown; capsule (Fig. 164) inclined or horizontal 6 mm to 6.2 mm  $\times$  .9 mm to 1 mm, elongate-ellipsoidal; operculum (Fig. 375) conical, long beaked; peristome double, outer phase of peristome characteristic of Diplolepideae (Fig. 165), inner phase of peristome with fimbriae-like margin (Fig. 166) and horizontal striations; spores spherical  $15\mu$  to  $17\mu$  in diameter.

### Bartramiaceae

28. *Bartramia madurensis* Dix. and Vard.\*: Plants pale green, grow well in cushions to a height of 2.5 cm to 3 cm without sporophyte in the soil covered rocks of the municipal Park of Kodaikanal 2800 metres

(Fig. 167). Soil acidic, pH being 4.5 to 4.6. Leaves (Fig. 168) from a broad base abruptly narrow into a long subulate tip; 4 mm to 4.5 mm. long and .5 mm to .7 mm broad at base; apex (Fig. 169) drawn out to a very fine point with serrate margin; leaf-cells mamillate elongated and thick walled (Fig. 170), costa very broad, cells narrow and closely packed, marginal rows of cells rectangular.

29. *Philonotis anisoclada* Card. and Vard.: Plants yellowish green, grow in dense tufts 3 cm to 3.5 cm tall (Fig. 171) in the swampy soil near mountain streams of Shembaganur (Kodaikanal) 2150 metres or at Thod-dabedda shola, (Ootacamund) 2800 metres. Stem erect 2 cm to 2.5 cm and branches profusely (5 to 6) from the same point. Rhizoidal felt dense extending from the base upwards on the stem. Leaves (Fig. 172) lanceolate .8 mm to 1.2 mm long and .3 mm to .4 mm broad at base, costa strong extending upto the tip; apex (Fig. 173) prolonged into a hair point, margin towards the apex serrated; laminar cell (Fig. 174) collenchymatous, thick walled and elongated; cells of the costa and margin much thick walled. Sporophyte terminal seta reddish 30 mm to 32 mm long; capsule (Fig. 176) erect or inclined, sub-spherical 2.5 mm to 2.6 mm  $\times$  1.5 mm to 1.7 mm when dry longitudinally furrowed; operculum sharp conical (Fig. 376); peristome (Fig. 177) free  $75\mu$  to  $88\mu$  long and  $30\mu$  broad at base; spores (Fig. 405) large spherical, granulose  $20\mu$  to  $22\mu$  in diameter.

#### Orthotrichaceae

30. *Hypnodon perpusillus* (Thw. and Mitt.) C. M.: Plants very small, with sporophyte less than 1 cm tall

(Figs. 173&179); grow on barks of rotting tree trunks at Perumal hill (Kodaikanal) 1600 metres. Leaves (Fig. 180) from oblong or spatulate base passing into an upper elliptic part; costa ending at some distance from the leaf tip; lamina cells (Fig. 181) rectangular  $55\mu$  to  $22\mu$  and isodiametric and roundish above; apex (Fig. 182) acute. Sporophyte terminal; seta 2 mm to 2.5 mm long; capsule ovoid, 8-ribbed 1 mm to 1.2 mm  $\times$  .4 mm to .5 mm, perichaetium almost inseparable from sporophytic foot; operculum conical and beaked (Fig. 377); peristome teeth connate in pairs (Fig. 183)  $176\mu$  long and  $66\mu$  broad at base; spores (Fig. 407) large, spherical with a thick coat  $19\mu$  to  $22\mu$  in diameter.

31. *Schlotheimia grevilleana* Mitt. : Plants growing predominantly on rock in Priests' Walk (Kodaikanal) 2250 metres forming thick carpet with bright yellowish younger parts. Axis creeping, branching profusely: 2 cm to 2.5 cm tall fastigate, densely foliose, main axis and branch stem below covered with felt of rhizoids (Figs. 184 & 185). Leaves (Fig. 186) elongate, oblong ending in a short point, 2.2 mm to 2.5 mm long and .6 mm to .75 mm broad; apex acute (Fig. 187); leaf-cells (Fig. 188) elliptic rectangular at base; ovate in the middle, cell walls much thickened, 5 to 6 cells in a square of  $22\mu$  (Fig. 189); isodiametric and sclerenchyma-like at apex (Fig. 190); rhizoids formed also from the base of the leaf. Sporophyte terminal on the branches; seta erect and short 8 mm to 9 mm long; capsule (Fig. 191) cylindrical 2 mm to 2.1 mm  $\times$  .9 mm to 1 mm; calyptra long conical enveloping the capsule, margin lobed; peristome free, basal cells rectangular arranged in two rows, teeth

132 $\mu$  long 15 $\mu$  to 18 $\mu$  broad at base (Fig. 192); spores (Fig. 408) spherical, 16 $\mu$  to 18 $\mu$  in diameter.

### Trachypodaceae

32. *Trachypus bicolor* Rein. and Hornsch.\*: Plants grow in dense tufts in the forests of Priests' Walk (Kodaikanal) 2800 metres in the shaded regions. Stem creeping on tree trunks branching profusely (Fig. 193), deep green in the older regions and yellowish green in the younger parts. Leaves (Fig. 194) lanceolate 1.5 mm to 2 mm long and .4 mm to .7 mm broad at base; apex acuminate (Fig. 195); leaf cells spindle-shaped and elongated (Fig. 197), cells of costa narrower and thick-walled (Fig. 196).

33. *Trachypus humilis* Lindb. var. *humilis* Zanten.\*: Plants grow in close tufts on trees in the sholas of Shembaganur (Kodaikanal) 2000 metres. Axis of this hairy moss very thin, branchlets many (Fig. 198). Leaves (Fig. 199) small, lanceolate with broad base .9 mm to 1.1 mm long and .25 mm to .3 mm broad at base; apex drawn out into a long narrow hair point (Fig. 200), cells linear and thick-walled above (Fig. 203), lumen of the cells very narrow; alar cells irregularly shaped (Fig. 202); no mid-rib seen in the leaf, except that at the base where the cells diverge out above.

34. *Trachypodopsis serrulata* (Beauv.) Fleisch. var. *crispatula* (Hook.) Zanten.\*: Plants grow on trees in tufts in the Priests' Walk (Kodaikanal) 2300 metres and on forest ground in Kern hill (Ootacamund)

265) metres. Secondary stems numerous, flexuately ascending with closely arranged leaves (Fig. 204). Leaves (Fig. 205) 3 mm to 3.75 mm long and .5 mm to .75 mm broad lanceolate and elongate with serrate margin; apex ending in a narrow acuminate (Fig. 206); leaf-cells (Figs. 207 & 208) elongated elliptic with much thickened cell walls; costa well defined; the lumen of the alar cells elongated oval (Fig. 209) and spindle shaped just above.

### Myuriaceae

35. *Myurium warburgii* (C. M.) Fleisch.\* : Plants compactly tufted, growing in the shaded and moist forests of Tiger shola (Kodaikanal) 2000 metres, branching irregularly with crowded brownish green leaves (Fig. 210). Leaves (Fig. 211) narrow lanceolate 3 mm. to 4 mm long .3 mm to .75 mm broad at base; upper margin serrulate; apex (Fig. 212) acuminate and drawn out into a fine point; leaf-cells (Fig. 213) narrow and irregularly spindle-shaped above and broadly rhomboid below (Fig. 214).

### Pterobryaceae

36. *Pterobryopsis orientalis* (C. M.) Fleisch.\* : Plants grow on tree trunks and on the branches in the moist woods of Silver cascade (Kodaikanal) 2000 metres. Primary stem creeping and spreading, branchlets yellowish green (Fig. 215). Leaves (Fig. 216) ovate oblong, shortly narrowed up, 1 mm to 2 mm long and .8 mm to 1 mm broad; leaf cells (Fig. 217) spindle-shaped 10 to 12 cells in a square of  $88\mu$ , towards the middle the cells narrowly rhomboid  $70\mu$  to  $90\mu \times 5\mu$  to  $6\mu$ ; alar cells quadratic

to ovate (Fig. 218); costa extending only up to half the leaf.

### Meteoriaceae

37. *Papillaria crocea* (Hamp.) Jaeg.: Plants pale green, thread-like growing on branches of trees in the shaded moist forests of Four-road shola (Ootacamund) 2650 metres, Tiger shola (Kodaikanal) 2500 metres. Primary stem thin and hanging in the form of festoons, secondary branches not very long but numerous, irregularly pinnate (Fig. 219 and frontispiece). Leaves closely adpressed on the stem, ovate auriculate (Fig. 220) .8 mm to 1.2 mm long and .5 mm to .7 mm broad; apex gradually acute (Fig. 221); costa strongly developed; laminar cells with roundish ovate lumen, 6 cells in a square of  $22\mu$  (Fig. 222). Sporophyte lateral, drooping from the branches: seta short 5 mm to 6 mm long; capsule cylindric 2.3 mm to 2.5 mm long and 1 mm to 1.2 mm in diameter; operculum (Fig. 378) conical and beaked; peristome teeth slender; spores spherical with a thick coat (Fig. 409)  $16\mu$  to  $19\mu$  in diameter.

38. *Aerobryopsis longissima* (Doz. and Molk.) Fleisch. \*: Plants grow on tree trunks and on rocks, irregularly branching, secondary stems foliose (Fig. 223) in the Tiger shola (Kodaikanal) 2100 metres. Leaves (Fig. 224) broadly lanceolate 3 mm to 5 mm long and .6 mm to .75 mm broad; apex (Fig. 225) very long, gradually attenuated into a fine point; leaf-cells (Fig. 226) rhomboid 6 to 7 in a square of  $22\mu$ .

39. *Barbella tenax* (C. M.) Broth. \*: Plants mostly growing on branches in drooping tufts in the Tiger shola (Kodaikanal) 2100 metres or in the Four-road shola (Ootacamund) 2650 metres; branches pinnate and long, younger portions yellowish green and less foliose (Fig. 227). Leaves (Fig. 229) ovate oblong 2 mm to 2.5 mm long and .5 mm to .75 mm broad, lower margins incurved; costa absent, point of attachment with a thickening; apex (Fig. 228) short acuminate with serrate margin; laminar cells (Fig. 230) very narrow and elongated, spindle-shaped,  $65\mu$  to  $85\mu$  long and  $5\mu$  to  $7\mu$  wide.

40 *Barbella determessi* (Rein. and Card.) Fleisch. \*: Plants grow on the branches of trees and hang down in the forests of Silver cascade (Kodaikanal) 2000 metres or in Lovedale shola (Ootacamund) 2300 metres. Primary stem very long, pinnately branching (Fig. 231). Leaves (Fig. 232) of the branches spreading, lanceolate 3.5 mm to 4 mm long and .4 mm to .5 mm broad at base; stem leaves adpressed to the axis, narrowly linear lanceolate; apex (Fig. 233) prolonged acuminate with slightly toothed margin; mid-rib absent; laminar cells (Fig. 234) narrowly linear and spindle-shaped.

41. *Barbella pendula* (Sull.) Fleisch.\*: Plants creep along the bark of tree trunks and branches in the forest of Priests' Walk (Kodaikanal) 2300 metres. Primary stem fimbriate with numerous branches, long and hairy (Fig. 235). The pH of the sediment in the crevices of the bark is 6.0. Leaves (Fig. 236) spreading only on the primary stem and lower regions of the branches, lanceolate 1.5 mm to 2 mm long and .5 mm to .7mm

broad; margin serrate; mid-rib wanting, narrow and thick walled cells closely packed at the point of attachment, indicative of costa; apex (Fig. 237) prolonged, acuminate, ending bluntly; laminar cells (Fig. 238) broad, linear rhomboid  $44\mu$  to  $46\mu$  long and  $7\mu$  to  $8\mu$  wide.

42. *Floribundaria floribunda* (Doz. and Molk.) Fleisch.\*: Plants grow with other mosses in the thick forests of Lovedale shola, 2300 metres, Four road shola 2650 metres (Ootacamund) and in the Park-shola (Kodaikanal) 2350 metres. Primary axis threadlike, branches and secondary branches bear spreading leaves (Fig. 239) Leaves (Fig. 240) broad at base and slightly auricled, lanceolate 1.6 mm to 2mm long and .5 mm to .75mm broad; apex (Fig. 241) prolonged acuminate laminar cells (Fig. 242) narrowly linear, cell walls very much thickened.

43. *Meteoriopsis squarrosa* (Hook.) Fleisch \* : Plants grow with other mosses in the thick forests of Lovedale shola, 2300 metres, Four-road shola 2650 metres (Ootacamund) and in the Park-shola (Kodaikanal) 2350 metres. Primary stem creeps on the bark of tree trunks and on the branches; secondary stems numerous and drooping (Fig. 243). Leaves (Fig. 244) stem clasping squarrose, crowded, basal part broad, sub-uniform 3 mm to 3.5 mm long and 1 mm to 1.5 mm broad; mid-rib absent; apex (Fig. 245) shortly acuminate with sparsely-toothed margin; laminar cells (Fig. 246) long spindle-shaped, cell walls thickened.

44. *Meteoriopsis reclinata* (C.M.) Fleisch.: Plants grow in dense tufts on branches of moist forest trees

in the Priests' Walk (Kodaikanal) 2200 metres, and Four-road Shola (Ootacamund) 2300 metres. Primary axis long, creeping or drooping in the form of festoons. Secondary pinnate branches short and numerous with spreading leaves (Fig. 247 and frontispiece). Leaves 2.5 mm to 3 mm long and 1 mm to 1.2 mm broad at base, recurved, falcately secund, stem-clasping (Fig. 248); mid-rib absent; apex acute with toothed margin (Fig. 249); laminar cells (Fig. 250) narrow but long elliptic with tapering ends, cell walls incrassate.

#### Neckeraceae

45. *Homaliodendron exiguum* (Bryol. Jav.) Fleisch.\*: Plants grow on the barks of decaying tree trunks in the damp forests of Priests' Walk (Kodaikanal) 2300 metres. Axis slender 3 cm to 5 cm tall (Fig. 251). Leaves (Fig. 252) bifarious ovate oblong. .8 mm to .1 mm long .25 mm to .3 mm broad. apex orbicular obtuse; mid-rib wanting; laminar cells (Fig. 253) quadratic to roundish, 4 to 5 cells in a square of  $22\mu$ .

46. *Homaliodendron flabellatum* (Dix.Sm.) Fleisch.\*: Plants sturdy and dark green growing on the trunks of trees and on moist rocks, rupestral and spreading horizontally. Axis strong, without leaves, branches two-ranked, shoot remarkably flattened (Fig. 254). Leaves bifarious on the branches, oblong, mid-rib not extending up-to tip; apex broad, upper margin irregularly toothed (Fig. 255); leaf-cells (Fig. 256) quadratic to roundish 3 to 4 cells in a square of  $22\mu$ . Gemmae bud like in the leaf axils.

47. *Distichophyllum succulentum* (Mitt.) Broth. \* : Plants grow in densely shaded and moist shelters of forest ground of Perumal hill (Kodaikanal) 2000 metres and of Kotagiri (Ootacamund) 2000 metres. Plants 2.5 cm to 3 cm tall and dark green (Fig. 257). Leaves (Fig. 258) more than twice long as broad, 2 mm to 2.5 mm long and .6 mm to .75 mm broad; costa excurrent slightly; marginal layer formed apex (Fig. 259) sharply retuse; laminar cells (Fig. 260) isodiametric and pentagonal, each cell  $22\mu \times 22\mu$ .

### Hookeriaceae

48. *Hookeria acutifolia* Hook. \*: Plants grow in moist and shaded forest ground of Thoddabedda slope (Ootacamund) 2800 metres. Plants pale green; leaves arranged in 4 or 5 rows (Fig. 261). Leaves (Fig. 262) ovate, broad and elliptic 3.5 mm to 4 mm long and .7 mm to 1 mm wide; marginal seam well formed; mid-rib absent; apex (Fig. 263) angularly acute; leaf cells rhomboid  $33\mu \times 11\mu$ .

### Hypopterygiaceae

49. *Hypopterygium tenellum* Lac. = *H. ceylanicum* Mitt. \* : Plants grow on rotting tree trunks in moist and shaded forests of Priests' Walk (Kodaikanal) 2500 metres. Primary stem creeps in the crevices of rocks and bark of trees; secondary branches profuse, flat and spreading (Fig. 264). Leaves (Fig. 265) ovate oblique, .9 mm to 1.1 mm long and .5 mm to .7 mm wide; mid-rib ending at some distance below the leaf-tip and running on one side dividing the lamina into unequal

halves; upper margin serrulate; laminar cells (Fig. 266) rhomboid hexagonal 5 to 6 cells in a square of  $88\mu$ .

### Leskeaceae

50. *Rhegmatodon orthostegius* Mont. : Plants grow on the twigs of small trees in moist forests of Priests Walk (Kodaikanal) 2500 metres and in the Wood-house shola (Ootacamund) 2600 metres. Secondary stems irregularly branched, branches ascending to erect (Fig. 263). Leaves (Fig. 269) ovate 1 mm to 1.5 mm long .4 mm to .6 mm wide; apex acute (Fig. 267); mid-rib not extending up-to the tip; leaf-cells (Fig. 270) quadratic at base, ovate in the upper portion, lumen much reduced on account of cell wall thickening, 3 cells in a square of  $22\mu$  in the middle of the lamina. Sporophyte terminal or lateral on the branches : seta reddish 6 mm to 8 mm long and erect; capsule (Fig. 271) 2 mm to 2.2 mm  $\times$  .9 mm to 1 mm cylindrical; operculum conical (Fig. 379); peristome (Fig. 274) free, inner peristome slender and long (Fig. 273); spores (Fig. 410) large, spherical  $19\mu$  to  $21\mu$  in diameter. Stomata found on the wall in the apophysis region of the capsule (Fig. 272).

### Thuidiaceae

51. *Herpetineurum toccoae* (Sull. and Lesq.) Card: \* Plants grow on tree trunks or soil covered rocks in the woods of Bombay shola (Kodaikanal) 2400 metres. Primary stem creeps along horizontally and secondary stems erect 4 cm to 5 cm tall, branches spreading

(Fig. 275). Leaves (Fig. 276) lanceolate 2.5 mm to 3 mm long and .7 mm to 1 mm wide at base; upper margin serrate (Fig. 277); leaf cells (Fig. 278) very small and sub-quadratic to round or ovate, 6 to 7 cells in square of  $22\mu$ .

52. *Thuidium tamariscellum* (C.M) Bryol. : Plants grow very well in the damp shady forest ground of Bear Shola (Kodaikanal) 2300 metres in matted tufts. Primary stem hardy, secondary stems branch thrice. This tri-pinnate branching and flattened frond like form characteristic of this moss (Fig. 279). Branch leaves (Figs. 280 & 281) ovate .15 mm to .2 mm long and .08 mm to .1 mm broad; stem leaves larger, cordate (Figs. 282 & 283) .8 mm to 1 mm long and .4 mm to .5 mm broad; both leaves with finely toothed upper margin; branch leaf-cells (Fig. 284) 3 to 4 in a square of  $22\mu$ , papillose, thick walled; stem leaf-cells rhomboid at alar region (Fig. 287). Sporophyte lateral on branches : seta reddish, erect or inclined, smooth, 15 mm to 17 mm long; capsule 2.7 mm to 3 mm  $\times$  .7 mm to .9 mm ellipsoidal (Fig. 285); operculum (Fig. 380) cupula-shaped and beaked; peristome free with longitudinal striations (Fig. 286); spores (Fig. 411) spherical  $12\mu$  to  $14\mu$  in diameter.

#### Brachytheciaceae

53. *Pleuropus nilghiriense* (Mont.) Toyama: Plants grow on tree trunks in the damp shaded Tiger Shola (Kodaikanal) 2150 metres. Plants (Fig. 288) brownish green, secondary branches numerous and spreading 4 cm

to 5 cm tall. Leaves (Fig. 289) lanceolately acuminate 1.2 mm to 2 mm long and .4 mm to .5 mm broad at base; upper margin (Fig. 290) sharply serrate; laminar cells (Fig. 291) elongated and narrow,  $40\mu$  to  $45\mu$  long and  $4\mu$  to  $6\mu$  wide. Sporophyte lateral: seta long, red, and erect 8 mm to 10 mm long; capsule cylindrical (Fig. 292) 2 mm to 2.2 mm  $\times$  .5 mm; operculum conical with long beak; peristome (Fig. 293)  $120\mu$  long and blunt  $45\mu$  broad at base with a middle striation; spores spherical,  $15\mu$  to  $17\mu$  in diameter.

54. *Rhynchostegium javanicum* (Bel.) Besch. : Plants 3.5 cm to 4.5 cm tall, light green, grow on moist ledges where earth has accumulated and in the Tiger Shola 2100 metres and in the Priests' Walk (Kodaikanal) 2300 metres. Secondary stems branching (Fig. 294). Leaves (Fig. 295) apparently two-ranked 2 mm to 2.5 mm long and .7 mm to 1 mm broad, ovate lanceolate, upper margin serrulate apex (Fig. 296) acuminate; laminar cells (Fig. 297) narrow and much elongated, upper cells spindle-shaped. Sporophyte lateral: seta 18 mm to 20 mm long; capsule (Fig. 298) horizontal cylindrical 4 mm to 4.2 mm  $\times$  .7 mm to .9 mm; lid arched conical (Fig. 381); peristomial teeth very long 2.7 mm upper part bent inwards (Fig. 299); spores (Fig. 412) small spherical,  $8\mu$  to  $10\mu$  in diameter.

#### Entodontaceae

55. *Erythrodontium julaceum* (Hook.) Par. : Plants grow horizontally on trees and shrubs, or on moist rocks in the Priests' Walk (Kodaikanal) 2350 metres. Plants long, secondary branches numerous 4 cm to 6 cm

tall (Fig. 300). Rhizoids unbranched, numerous in clusters from the axis in different regions. Leaves broad, ovate (Fig. 301), slightly concave 1 mm to 1.5 mm long and .7 mm to .9 mm broad; apex sub-abruptly acuminate (Fig. 302), upper margin toothed; alar cells (Fig. 303) particularly marginal and sub-marginal ones ovate to quadratic, hardly wider than the inner cells which are elongated and rhomboid; upper cells spindle-shaped with thick walls (Fig. 304). Sporophyte lateral; seta 18 mm to 25 mm long; capsule (Fig. 305) erect, prolate, sphaeroidal 1.5 mm to 1.7 mm  $\times$  .9 mm to 1.2 mm; lid conical; peristome (Fig. 306) sharp 66 $\mu$  long and 16 $\mu$  broad at base; spores large, spherical 18 $\mu$  to 21 $\mu$  in diameter.

### Sematophyllaceae

56. *Foreauella indica* Dix. and Vard.\* : Plants grow on soil covered rocks or tree trunks in Shenbaganur (Kodaikanal) 2000 metres. Primary stem grows horizontally and numerous branches 4 cm to 5 cm long (Fig. 307) grow vertically. Stem leaves (Fig. 308) adpressed, obovate, trapezoidal; apex acuminate .9 mm to 1.2 mm long and .4 mm to .6 mm broad; alar cells oblong upper leaf cells (Fig. 309) elongate rhomboid 32 $\mu$   $\times$  22 $\mu$ .

---

\* The specific name of this plant (56) in current use is *Foreauella orthothecia* (Schw.) Dix. and Vard. according to E. B. Bartram. H. Ando points out that *Foreauella indica* is only a synonym of *Foreauella orthothecia*. This is referred to as a South Indian genus after the name of Rev. Father G. Foreau, who was kind enough to show the author this plant in its natural habitat in the Sacred Heart College Garden at Shenbaganur, Kodaikanal in 1959.

57. *Sematophllum subhumile* (C. M.) Fleisch. : Plants grow on moist rock in shaded forests of Shenbaganur (Kodaikanal) 2000 metres. Primary stem creeps horizontally 5 cm to 6 cm long; irregularly pinnate branches spreading (Fig. 311). Leaves (Fig. 312) not adpressed even when dry, long lanceolate 1.5 mm to 2 mm long and .4 mm to .5 mm broad; apex acuminate ending in a fine point; leaf cells narrow, elongated, spindle-shaped  $60\mu$  to  $64\mu \times 6\mu$  to  $7\mu$  (Fig. 313), upper cells narrow linear (Fig. 314). Sporophyte lateral. seta 10 mm to 12 mm long; capsule (Fig. 315) inclined elongate ellipsoidal 1.5 mm to 1.7 mm  $\times$  .3 mm to .5 mm; operculum (Fig. 382) cupula-shaped, beak long; peristome (Fig. 316) 1.75 mm long; spores (Fig. 413) small spherical  $9\mu$  to  $10\mu$  in diameter, with a thick coat. Gemmae in the form of vegetative buds found in the leaf axil.

58. *Warburgiella leptorrhynchoides*(Mont.)Fleisch \*: Plants: pale yellowish green, grow on the rocks in the forest of Tiger shola (Kodaikanal) 2000 metres. (Fig. 317) spreading horizontally with numerous branches. Leaves (Fig. 318) ovate oblong 1.2 mm to 2 mm long and .2 mm to .4 mm; apex (Fig. 319) ending in a narrow acumen; margin of the leaf incurved and entire at base, but obscurely denticulate near the tip; laminar cells (Fig. 320) narrowly linear  $66\mu$  long and  $7\mu$  to  $8\mu$  broad: alar cells inflated.

### Hypnaceae

59. *Hypnum cupressiforme* Hedw.\*: Plants prostrate, on forest ground or on decaying wood in Bear Shola

(Kodaikanal) 2300 metres (Fig. 321) branching pinnately and forming flat carpets. Leaves (Fig. 322) ovate lanceolate 2mm to 2.5mm long and .4mm to .6 mm broad at base; apex reflexed and secund (Fig. 223) acuminate to a point; mid-rib absent; leaf-cells (Fig. 324) narrow and elongated  $50\mu$  to  $66\mu \times 5\mu$  to  $6\mu$ , cells towards the middle of the leaf slightly more elongated with tapering ends cell walls much thickened. Simple rhizoids found to arise from the base of the leaves also.

60. *Taxiphyllum taxirameum* (Mitt.) Fleisch. : Plants grow on the bark of trees in moist forests of Priests' Walk (Kodaikanal) 2350 metres. Primary axis creeps on the bark and branches vertical on upper side (Fig. 325). Leaves (Fig. 326) ovate oblong 1.25 mm to 1.75 mm long and .3mm to .5mm wide; apex (Fig. 327) long acuminate; leaf cells isodiametric, 3 cells in square of  $22\mu$  (Fig. 328) lumen of upper leaf cells very small.

61. *Ctenidium lychnites* (Mitt.) Broth. : Plants grow on moist ground or on tree trunks in the woods of Priests' Walk (Kodaikanal) 2350 meters. Primary stem prostrate, branching pinnately 2 cm to 2.5 cm tall (Fig. 329). Stem leaves cordate ovate, rapidly narrowed into a lanceolate upper part. Branch leaves narrower, elliptic (Fig. 330) 1 mm to 1.5mm long and .4 mm to .5 mm broad; apex (Fig.331) acuminate; margin serrate; leaf-cells linear, elongated and narrow (Fig. 332). Sporophyte lateral: seta erect and inclined above 25mm to 27mm long; capsule (Fig. 333) elongated ovoid, horizontal 6.5 mm to 7mm.  $\times$  1.3 mm to 1.5 mm; lid conical with sharp beak (Fig. 383); peristome (Figs. 334 and 335) 2.75 mm

to 3 mm long with a central dividing line up-to some distance from the base. spores spherical  $12\mu$  to  $13\mu$  in diameter.

### Polytrichaceae

62. *Atrichum aculeatum* Card. and Vard.: Plants grow in dark green dense tufts and cushions on forest ground especially under the shade near tree trunks in the damp forests of Thoddabedda (Ootacamund) 2500 metres, or near St. Mary's Hill 2600 metres, or in rich humus soil of Bombay Shola 2800 metres (Kodaikanal), to a height of 2 cm to 3 cm (Fig 336). Soil acidic pH being 5.6 to 5.8; Leaves (Fig. 337) keeled, linear 10 mm to 11 mm long and 1.2 mm to 1.5mm broad with a marginal rim, upper margin serrate; apex acute; basal leaf-cells isodiametric to hexagonal  $22\mu$  to  $33\mu \times 22\mu$  (Fig. 338); upper leaf cells roundish (Fig. 339); costa prominent with narrow linear and elongated cells. Sporophyte terminal or on the branches: seta long, reddish, erect 25 mm to 27 mm; capsule inclined 5 mm to 6.5 mm  $\times$  1.2 mm to 1.5 mm cylindrical, slightly curved; (Fig 340) operculum (Fig. 384) short, conical and beaked peristome (Fig. 341) teeth 16, blunt,  $110\mu$  long and  $44\mu$  broad; spores (Fig. 414) spherical  $15\mu$  to  $17\mu$  in diameter.

63. *Pogonatum microstomum* (R. Br.) Brid. : Plants gregarious, strong and sturdy growing on the crevices of culverts, or on stone walls or on moist grounds, in Kodaikanal and in Ootacamund at all altitudes between 2000 metres to 2700 metres. Gametophyte alone very tall 10 cm to 15 cm lower part of the stem rhizomatoid (Fig. 342). Leaves (Fig. 343) linear 8 mm to 11 mm long and 1 mm to 1.5 mm broad and  $44\mu$  thick; apex (Fig. 344) convex, margin toothed;

cells of the margin quadratic, those of the leaf sheath rectangular (Figs. 345 & 349); mid-rib cells narrow linear; leaf in transverse section resembling that of *Marchantia* thallus (Fig. 349), many cells thick, lamellate.; Sporophyte terminal or lateral: seta erect 65 mm to 70mm long; capsule 5 mm to 5.5mm  $\times$  1.8 mm to 2 mm roundish, ellipsoidal: operculum broad, cupula-shaped and short beaked; calyptra hood shaped producing a dense felt of hairs enveloping the capsule; peristome teeth 32 with blunt end,  $286\mu \times 110\mu$  (Figs. 347 & 348); spores (Fig. 415) greenish, spherical  $12\mu$  to  $14\mu$  in diameter.

64. *Pogonatum neesii* (C.M.) Mitt. : Plants grow in dense tufts on the road-side cuttings to Thoddabedda (Ootacamund) 2800 metres or around St. Mary's Hill (Kodaikanal) 2600 metres. Soil acidic, pH being 5.6 to 5.8. Gametophytes 2 cm to 5 cm tall, not as tall as *Pogonatum microstomum*. Plants gregarious 4 cm to 7 cm tall (Fig. 350). Leaves (Fig. 351) whorled on the stem, linear 4 mm to 8 mm long 1 mm to 1.5 mm broad and  $40\mu$  to  $44\mu$  thick; apex acute with serrate margin (Fig. 352); leaf sheath cells (Fig. 353) quadratic at base, roundish above, cells of the costa elongated and narrow; leaf in transverse section resembles that of the thallus of *Marchantia* (Fig. 354) multicellular in thickness, lamellate. Sporophyte terminal: seta erect 35 mm to 40 mm long; capsule (Fig. 355) round, erect, elliptic 4 mm to 4.5 mm  $\times$  1.5 mm to 2 mm; lid conical, beaked (Fig. 385); calyptra hood-shaped giving rise to long hairs forming a felt; subjacent this hair-cap calyptra short thin cuculate inner calyptra also present; peristome with 32 teeth (Fig. 356); spores (Fig. 416) spherical, greenish  $9\mu$  to  $12\mu$  in diameter.

## General considerations

After Hedwig (1864) who has left magnificent illustrations of mosses, several Bryological Floras and Manuals have appeared from different parts of the world. Steere (1950, 1952) has reviewed some of these treatments. Outstanding contributions on Moss Flora have been made by Brotherus(1924). Dixon (1904), Grout (1924, 1928-1940), Bartram (1939, 1949), Jennings (1951), Dunham (1951), Jensen (1939), Krusenstijerna (1945) and Watson (1954) have produced monumental works. These however, are mainly systematic descriptions of the mosses occurring in different regions of the world. Ruhland (1924) has given a detailed account of the life-history of the mosses in general.

The general considerations on the external morphology of the South Indian mosses dealt with in this study, are based upon original observations both in the field and the laboratory. In most cases they are in conformity with the observations of previous workers.

### a) *The Leaf*

Most of the Bryologists have taken into account the character of the leaves as one of the most important keys for the identification of the mosses (Dixon, 1904),

(Dunham, 1951). It has been considered very important even in the determination of sterile mosses. Full grown and well developed leaves alone were taken up for detailed study.

### *The phyllotaxy*

The mode of arrangement of the leaves on the stem is also equally important. The usual arrangement in most cases is spirally alternate. The leaves are distichous in *Fissidens* (Figs. 1 & 2). In *Hypnum* (Fig. 322) the leaves are curved and turned in opposite directions giving a braided appearance, and in *Bryum* (Fig. 145) they are adpressed to the stem, so compactly dove-tailed that the lower leaves almost encircle the upper ones. The number and size of the leaves on the main and secondary axes vary considerably in the pleurocarpous mosses. In *Homaliodendron flabellatum* (Fig. 254) the leaves occur only on the branches but not on the main axis when they are old. In *Thuidium tamariscellum* (Figs. 280 & 281) the leaves of the main axis are larger than those of the branches. Watson's (1954) statement that the leaves of the finer branches are often different in shape from those of the main stem, is true of this moss also.

### *The shape*

The shape of the moss leaf varies greatly in different species. Some are long and narrow as in *Ditrichum amoenum* (Fig. 9), lanceolate as in *Philonotis anisoclada* (Fig. 172), ligulate in *Mnium coriaceum* (Fig. 155), falcate in *Meteoriopsis squarrosa* (Fig. 244), linear in *Myurium*

and *Thysanomitrium* (Figs. 211 & 29) and nearly orbicular in *Homaliodendron exiguum* (Fig. 252). Sometimes the shape is such as to combine a few terms for describing them such as linear-lanceolate in *Rhizogonium spiniforme* (Fig. 161) or ovate-oblique in *Hypopterygium tenellum* (Fig. 265). The mid-rib is usually present in many of the mosses but it is characteristically absent in some such as *Barbella tenax* (Fig. 229), *Floribundaria floribunda* (Fig. 240), *Meteoriopsis squarrosa* (Fig. 244). In a few cases the mid-rib ends blindly half way as in *Distichophyllum succulentum* (Fig. 258) and in *Hypopterygium tenellum* (Fig. 265), whereas in certain others it is continued up almost to the apex as in *Thuidium tamariscellum* (Figs. 280 & 232) and *Pleuropus nilghiriense* (Fig. 239). Lamella on leaves is characteristic of Polytrichaceae.

#### *The margin*

The margin of the leaves varies very much. This diagnostic character is of immense value in the identification of the species. There is entire margin in *Campylopus nodiflorus* (Fig. 21), undulate in *Mnium coriaceum* (Fig. 155), serrate in *Rhizogonium spiniforme* (Fig. 161) serrulate in *Trachypodopsis serrulata* (Fig. 205) and denticulate in the upper part of *Mnium coriaceum* (Fig. 157). There are also cases in which the lower part may be entire and the upper part toothed as in *Rhodobryum giganteum* (Fig. 152).

#### *The apex*

The apex of the leaf is also of equal importance in mosses for specific determination. It may be acumi-

-nate as in *Trachypus humilis* (Fig. 200), acute in *Hookeria acutifolia* (Fig. 263), and terete and excurrent in *Tortula muralis* (Fig. 63) and apiculate-retuse in *Distichophyllum succulentum* (Fig. 258).

### *The leaf cells*

The cellular configuration and their arrangement in the leaves of the mosses is characteristic of the species. They are isodiametric in *Taxiphyllum taxirameum* (Fig. 328), rectangular in *Leucobryum* species (Figs. 52 & 59), rhomboid in *Anomobryum subnuditum* (Fig. 142), papillose in *Thuidium tamariscellum* (Fig. 284) and narrowly elongate in *Sematophyllum subhumile* (Fig. 313). The alar cells are different from the cells of upper portions of the lamina not only in size but in form as well. The shape of the cells in different parts of the lamina is also variable in *Trachypus humilis* (Fig. 199, 201 to 203). The cells of the costa are usually different in size and shape from those of the lamina. They are elongated and narrow according to nature and thickness of the costa. The cell-walls in some mosses are thickened. In some they are so much thickened that the lumen is small or much reduced. Generally all the pleurocarpous mosses studied here, have long narrow cells throughout the lamina and in the acrocarpous mosses they are short and / or isodiametric in the upper portions and elongated rectangular at the base.

The nature of distribution of the chloroplasts is different in some mosses. The lower part of the leaf of *Bryum argenteum* is studded with chloroplasts while

the upper portion is so completely devoid of chloroplasts that it appears silver white and hence the specific name *argenteum* (Fig. 145). In *Leucobryum* the leaves are multilayered pale green and turn white when dry. The laminar cells are large and elongate some lacking chloroplasts. The leaves of the mosses in most cases are made up of only a single layer of cells except in the costa where it is multi-layered and also at the margin whenever it is thickened. The leaf of *Leucobryum* is thick and is more than one cell in thickness (Figs. 51 & 58); The leaves of *Pogonatum* species are thick or are perhaps the thickest of the lot. Longitudinal strips, the lamellae are closely arranged parallel to one another along the leaf. Dunham (1951) describes this costa as a wide mid-rib. The anatomical details of these leaves are dealt with in the chapter on the Anatomical Observations: (Figs. 51, 58, 349 & 354).

In some mosses there is variation to a great extent in the size of the leaves in the same species. The leaves of the lower part of the axis and hence the old, and the well developed young leaves in the tender region are very much different in size though there is no variation in shape as judged from Length / Breadth ratio. The observations are tabulated as under:

TABLE - I

Name	Length of leaf in mm (L)	Breadth of Leaf in mm (B)	L/B ratio
1	2	3	4
<i>Barbula indica</i>	0.80	.50	1.6
	1.20	.75	1.6
	1.00	.65	1.6
	1.30	.80	1.6
	1.15	.70	1.6
	1.25	.75	1.6
	1.09	.67	1.6
	0.90	.54	1.6
	0.85	.50	1.6
	1.50	.90	1.6
	1.32	.82	1.6
	0.88	.55	1.6
	<i>Pogonatum microstomum</i>	09.8	1.2
08.0		1.0	8.0
10.5		1.3	8.0
08.8		1.1	8.0
09.6		1.2	8.0
12.0		1.5	8.0
09.5		1.2	8.0
09.7		1.2	8.0
12.0		1.5	8.0
11.3		1.3	8.0
10.4		1.3	8.0
11.2		1.4	8.0

TABLE I (Contd.)

1	2	3	4
<i>Pogonatum</i>			
<i>neesii</i>	5.0	1.0	5.0
	6.0	1.2	5.0
	5.5	1.1	5.0
	4.0	0.8	5.0
	8.0	1.6	5.0
	7.5	1.5	5.0
	7.5	1.5	5.0
	6.6	1.3	5.0
	6.5	1.3	5.0
	6.0	1.2	5.0
	5.5	1.1	5.0
	6.5	1.3	5.0
<i>Rhodobryum</i>			
<i>giganteum</i>	08.8	4.0	2.2
	10.0	4.5	2.2
	10.2	4.6	2.2
	09.5	4.2	2.2
	09.9	4.5	2.2
	07.2	3.2	2.2
	09.7	4.1	2.2
	07.8	3.5	2.2
	08.4	3.8	2.2
	11.0	4.8	2.2
	08.2	3.7	2.2
	06.7	3.0	2.2

It may thus be seen that in spite of variation in the size of the leaves in these mosses, the L/B ratio of each species is a constant factor.

However, in the following two species the L/B ratio of the leaves is variable :

TABLE - II

Name	Length of Leaf in mm (L)	Breadth of Leaf in mm (B)	L/B ratio
<i>Fissidens</i>			
<i>schimidii</i>	0.88	.09	9.7
	0.99	.10	9.9
	1.10	.11	10.0
	0.93	.10	9.3
	1.15	.11	10.4
	1.21	.13	9.3
	0.99	.12	8.3
	1.21	.13	9.3
	1.10	.12	9.1
	1.14	.13	8.9
<i>Ditrichum</i>			
<i>amoeum</i>	4.4	0.44	10.0
	4.8	0.32	15.0
	3.8	0.30	12.6
	4.3	0.30	14.3
	5.0	0.40	12.5
	5.3	0.42	12.6
	4.6	0.38	12.1
	4.1	0.40	10.2
	5.4	0.45	12.0
	6.0	0.45	13.3

(b) *The axis :*

The mosses are classified into two groups, the acrocarpi and the pleurocarpi. Although this division is not generally regarded as a natural one, according to Watson (1954), still it is convenient to retain it. In the former the capsule is normally borne at the apex of the primary axis or on its branch. The capsule arises laterally on the axis or on a short side branch in the pleurocarpous mosses. The length of the axis of the acrocarpous members is generally not very long, but that of the pleurocarpi is very long and may grow even to a maximum length of 2 metres, being found in *Papillaria crocea*. The smallest moss under this study is *Hypnodon perpusillus* which has been referred to as "a pretty little moss" by Bartram, its maximum height without sporophyte being 1.5 mm. The axis creeps inside the soil in some mosses as in the case of *Rhacomitrium javanicum*. It is very hardy in the fan moss, *Homatiodendron flabellatum*. Even the sturdy *Pogonatum microstomum* is not as hardy as this fan moss. There are also some mosses in which an axis "worth the name" is not organised as in *Ditrichum amoenum* and *Hypnodon perpusillus*.

The branching in the mosses is very variable. In the acrocarpi where the branches are not many, they arise from the primary stem as in *Bryum argenteum*, *Brachymerium exile* and *Pogonatum microstomum*. Profuse branching from the primary stem and growing above in close tufts can be seen in *Schlotheimia grevilleana*. Irregular but profuse branching are seen in *Myurium warburgii*, *Barbella tenax* and others.

The rhizoids usually arise from the base of the axis. There are also some mosses in which they arise from

any part of the primary axis or from the branches or from base or margin of the leaves. As Narayanaswamy (1957) has shown, they are produced from the seta or even from the base of the capsule. They are simple without any thickening and brownish in colour in most of the mosses. They are profusely branched forming a felt in *Sclotheimia grevilleana* and are reddish in colour in this moss and also in *Rhizogonium spiniforme*.

It may be pointed out that axis of different types of mosses such as the terrestrial mosses, the rock mosses, and the tree mosses are typical in nature of their growth. The branching in the axis and the production of rhizoids in certain mosses are interesting

### c *The sporophyte.*

Brotherus (1899) and Cardot and Varde (1925) have referred to a few of these mosses and have described the sporophytes. Again in his excellent contribution in *Die Natürlichen Pflanzenfamilien*, Brotherus, (1924) like a few other Bryologists, has taken into account the character of the sporophyte as the most important diagnostic feature in the identification of mosses.

### *The foot*

In all the available literature on the mosses, there seems to be no detailed account on the shape and nature of articulation of foot of the sporophyte to the gametophyte. In the present study sporophytes were available only in 34 species of mosses. The morphology of the foot has been brought to light for the first time for the these mosses. At present they have tentatively been classed under eight groups.

The fertilised egg develops into a sporophyte. This is differentiated into the foot, the seta and the capsule. The foot buried in the tip of the gametophore, could be separated with a pair of fine needles without causing any injury to it. However, it comes off easily when the perichaetial bract and the gametophore tissue are ripped open. Though the shape of the foot is generally attenuated in most of the cases they show the following 8 different forms:

1. Narrowly elongated to a blunt tip : e.g. in

*Hyophila involuta* (Fig. 422)

*Ditrichum amoenum*

*Funaria hygrometrica*

*Anomobryum subnuditum*

*Papillaria crocea*

*Sematophyllum subhumile*

2. Broadly narrowed down to a blunt tip: e.g. in

*Tortula muralis* (Fig. 423)

*Funaria submarginata*

*Pohlia flexuosa*

*Rhodobryum giganteum*

3. Conical and elongated : e.g. in

*Bryum argenteum* (Fig. 424)

*Regmatodon orthostegius*

*Pleuropus nilghiriense*

*Erythrodonium julaceum*

4. Sharply attenuated to a sharp point: e.g. in  
*Barbula indica* (Fig. 425)  
*Trematodon ceylonensis*  
*Brachymenium exile*  
*Thuidium tamariscellum*
5. Gradually narrowed to blunt and mamillate tip: e.g. in  
*Rhizogonium spiniforme* (Fig. 426)  
*Campylopus nodiflorus*  
*Dicranodontium fragile*  
*Leucobryum neilgherrense*  
*Trichostomum cylindricum*  
*Weisia macrospora*
6. Flatly narrowed to blunt and undulate tip: e.g. in  
*Rhynchostegium javanicum* (Fig. 427)  
*Fissidens schimidii*  
*Philonotis anisoclada*  
*Hypnodon perpusillus*
7. Bulging and napiform below: e.g. in  
*Atrichum aculeatum* (Fig. 428)  
*Mnium coriaceum*  
*Schlotheimia grevilleana*
8. Abruptly narrowed to short tip: e.g. in  
*Pogonatum microstomum* (Fig. 429)  
*Pogonatum neesii*  
*Ctenidium lychnites*

## The Seta


The seta of the sporophyte is short in some and long in others, There appears to be no correlation or proportion between the length of the seta and the length of the axis or its branch. It is short but variously curled as in *Campylopus nodiflorus* where the length of axis is considerably long. In *Funaria hygrometrica* the seta is long, where the axis is short comparatively. The nature of looping of the seta is again very or extremely variable. There are different degrees\* of looping of the seta. They may be grouped under two heads.

A. In this group the angle of looping is  $135^\circ$  and above. The capsules of these mosses are either slightly inclined or erect. They are:

*Ditrichum amoenum* ( $180^\circ$ )

*Barbula indica* ( $180^\circ$ )

---



*Schlotheimia grevilleana* (180°)  
*Erythrodontium julaceum* (180°)  
*Fissidens schimidii* (180°)  
*Trematodon ceylonensis* (155°)  
*Campylopus nodiflorus* (140°)  
*Leucobryum neilgherrense* (155°)  
*Trichostomum cylindricum* (145°)  
*Weisia macrospora* (50°)  
*Hyophila involuta* (160°)  
*Tortula muralis* (150°)  
*Funaria submarginata* (170°)  
*Pohlia flexuosa* (170°)  
*Brachymenium exile* (140°)  
*Papillaria crocea* (160°)  
*Philonotis anisoclada* (140°)  
*Thuidium tamariscellum* (160°)  
*Pleuropus nilghiriense* (150°)  
*Rhegmatodon orthostegius* (160°)

**B.** In this group the angle of looping is below 135°. The capsules of these mosses are much inclined or even dropping down. They are:

*Rhodobryum giganteum* (120°)  
*Mnium coriaceum* (110°)  
*Rhizogonium spiniforme* (100°)  
*Ctenidium lychnites* (120°)  
*Pogonatum microstomum* (105°)  
*Pogonatum neesii* (125°)  
*Anomobryum subneditum* (110°)  
*Sematophyllum subhumile* (90°)  
*Pohlia flexuosa* (50°)  
*Bryum argenteum* (45°)

- Rhynchostegium javanicum* (55°)  
*Atrichum aculeatum* (5°)  
*Philonotis anisoclada* (20°)  
*Funaria hygrometrica* (25°)  
*Dicranodontium fragile* (10°)

The different degrees of looping of the seta in the two groups appear to have some bearing on the dispersal of spores. Spore dispersal is obviously regulated by the peristome. On closer examination, however, it has been found that the structure of the peristome in group A\* is not as elaborate as the structure of the peristome in group B. This feature seems to conform to the prophetic statement of Jennings (1951): "Species whose capsules stand vertically are not so likely to have well-developed peristome as are species whose capsules are inclined or vertical, this variation corresponding to the need for the regulation of the spore dispersal."

As the degree of looping varies according to the age of the sporophyte, only well matured sporophytes were taken into consideration. Several observations were made in each case in the foregoing classification. Further measurements of the length of the foot, seta and of the capsule were taken in the following manner: the length of the foot was measured from the point of attachment of the perichaetial bract downwards. The length of the seta excluding the length of the foot buried in the gametophore and that of the capsule excluding the length of the operculum were measured. The average of six readings in each case is tabulated (Table - III).

---

\* In fact in *Funaria submarginata* the peristome is even wanting.

TABLE III

Name of moss	Length of foot in mm	Length of seta in mm	Length of capsule in mm	Breadth of capsule in mm	Diameter of spores in $\mu$
1	2	3	4	5	6
<i>Fissidens schimidii</i>	0.49	9.0	2.8	0.45	12
<i>Ditrichum amoenum</i>	0.83	8.5	2.3	0.45	16
<i>Trematodon ceylonensis</i>	0.99	22.5	8.0	0.62	23
<i>Campylopus nodiflorus</i>	0.91	9.0	1.6	0.92	16
<i>Dicranodontium fragile</i>	1.08	11.0	5.5	0.92	26
<i>Leucobryum neilgherrense</i>	0.66	4.5	3.2	0.84	19
<i>Weisia macrospora</i>	0.46	11.0	6.1	0.52	10
<i>Trichostomum cylindricum</i>	0.99	9.2	2.8	0.61	15
<i>Hyophila involuta</i>	0.99	11.0	2.8	0.62	9
<i>Barbula indica</i>	0.54	10.0	1.8	0.72	16
<i>Tortula muralis</i>	0.79	12.5	4.2	0.52	10

TABLE III (Contd.)

Name of moss	Length of foot in mm	Length of seta in mm	Length of capsule in mm	Breadth of capsule in mm	Diameter of spores in $\mu$
1	2	3	4	5	6
<i>Funaria hygrometrica</i>	0.83	48.0	4.0	0.92	10
<i>Funaria submarginata</i>	0.79	26.5	1.3	0.90	26
<i>Pohlia flexuosa</i>	0.66	32.0	7.0	0.92	16
<i>Brachymenium exile</i>	0.66	31.0	2.0	0.75	13
<i>Anomobryum subnuditum</i>	0.66	30.5	2.6	1.12	16
<i>Bryum argenteum</i>	0.62	25.0	3.0	1.32	13
<i>Rhodobryum giganteum</i>	1.50	51.5	8.5	1.52	13
<i>Mnium coriaceum</i>	1.50	41.3	10.8	1.25	13
<i>Rhizogonium spiniforme</i>	1.42	36.0	6.1	0.92	16
<i>Philonotis anisoclada</i>	1.24	31.0	2.5	1.65	21
<i>Hypnodon perpusillus</i>	0.83	2.3	1.1	0.45	20
<i>Schlotheimia grevilleana</i>	0.87	8.5	2.1	0.92	17

TABLE III (Contd.)

Name of moss	Length of foot of seta		Length of capsule		Breadth of capsule		Diameter of spores	
	in mm	in mm	in mm	in mm	in mm	in mm	in $\mu$	$\mu$
1	2	3	4	5	6	7	8	9
<i>Papillaria crocea</i>	0.79	5.5	2.4	1.12	16			
<i>Rhegmatoxon orthostegius</i>	0.91	7.0	2.1	0.92	20			
<i>Thuidium tamariscellum</i>	0.70	17.0	3.2	0.85	13			
<i>Pleuropus nilghiriense</i>	0.91	9.0	2.1	0.52	16			
<i>Rhynchostegium javanicum</i>	0.83	19.0	4.1	0.80	9			
<i>Erythrodonium julaceum</i>	0.54	19.5	1.6	1.00	20			
<i>Sematophyllum subhumile</i>	0.66	11.0	1.6	0.45	10			
<i>Ctenidium lychnites</i>	1.07	26.5	5.2	1.32	12			
<i>Atrichum aculeatum</i>	1.70	39.5	6.8	1.45	16			
<i>Pogonatum microstomum</i>	1.66	67.0	5.3	1.92	13			
<i>Pogonatum neesii</i>	1.49	38.0	4.3	1.75	10			

It may be seen from a perusal of Table-III that there appears to be no correlation between the size of the capsule and the diameter of the spores. Nor does it appear that there exists any relation between the length of the seta and the length of the capsule. But when length of the foot and the length of the capsule are taken into consideration there is a corresponding increase in the length of the foot *pari-passu* with the increase in the length of the capsule.

It was observed that the seta is short in the following mosses which grow on the bark and branches of trees and on bare rocks:

*Hypnodon perpusillus*  
*Leucobryum neilgherrense*  
*Pleuropus nilghiriense*  
*Papillaria crocea*  
*Schlotheimia grevilleana*

This is in confirmity with the observation of Goebel (1905).

#### **The capsule**

The columella in the capsule of the mosses studied is prolonged up-to the peristomial region. The wall of the capsule in the region of the apophysis is provided with stomata in some species. It has been reported by Haberlandt (1895) that the stomata are confined to the sporogonium where they often occur in large numbers on the apophysis which is the principal organ of photosynthesis and to a small extent on the capsule wall where

it is green. He has sketched the stomata of six species of mosses and also stated that a single capsule of *Polytrichum* may bear three different types of stomata namely (1) two celled stomata; (2) one celled stomata with fused guard cells and (3) four-celled stomata in which each of the two primary guard cells is divided into two by a median transverse wall. Of the three kinds, the second type was not observed in this study. Campbell (1930), Ruhland (1924), and Grout (1924) have reproduced the illustrations of Haberlandt (1895). Van Der Wijk (1932) has reported the presence of stomata in *Orthotrichum*. Though Berkeley (1863) has stated that "stomata are not uncommon on the surface of the sporangia\* in which case they resemble very closely those of Phaenarogams," they are not found on all the capsules in this study. Out of the 34 capsules, the stomata were observed only in the following 8 species:

1. *Trematodon ceylonensis* (Fig. 18)
2. *Barbula indica* (Figs. 75 & 76)
3. *Funaria hygrometrica* (Fig. 420)
4. *Funaria submarginata* (Fig. 121)
5. *Pohlia flexuosa* (Fig. 421)
6. *Hypnodon perpusillus*
7. *Rhegmatodon orthostegius* (Fig. 272)
8. *Rhynchostegium javanicum*

In all cases, the stomata are made up of two guard cells but the number of subsidiary cells is variable: In

---

\* This term "sporangia" has been used by Berkeley (1863) obviously for "sporogonia."

*Trematodon ceylonensis* the cells of capsular wall are much oval oblong or polygonal and ovoid resembling the cells of the leaf. A few stomata occurring in the lower surface of the apophysis in *Barbula indica* are provided with four guard cells (Fig. 76.)

The frequency of the stomata was calculated by taking a unit area of one hundred micrometer squares of the Netz micrometer eye-piece. The stomatal indices of the 8 species have been calculated (Table IV). Stomatal index according to Salisbury quoted by Trease (1945) is an expression of the percentage proportion of the ultimate divisions of the dermatogen of the capsule which have been converted into stomata. The formula for stomatal index is  $I = \frac{S}{E+S}$  where S is the number of stomata per unit area and E, the number of epidermal cells in the same area.

TABLE-IV

Name of the moss	No. of cells per unit area (E)	No. of stomata per (same) unit area (S)	Stomatal index (I) $I = \frac{S}{E+S} \times 100$
1	2	3	4
<i>Trematodon ceylonensis</i>	36	2	$\frac{2}{36+2} \times 100 = 5.26$
	35	2	$\frac{2}{35+2} \times 100 = 5.32$
	37	2	$\frac{2}{37+2} \times 100 = 5.13$
	36	2	$\frac{2}{36+2} \times 100 = 5.26$
	35	2	$\frac{2}{35+2} \times 100 = 5.32$
	38	2	$\frac{2}{38+2} \times 100 = 5.00$
			<u>Average 5.21</u>

TABLE-IV (Contd.)

1	2	3	4	5
<i>Barbula indica</i>	84	2	$\frac{2}{84+2}$	$\times 100 = 2.32$
	85	2	$\frac{2}{85+2}$	$\times 100 = 2.29$
	84	2	$\frac{2}{84+2}$	$\times 100 = 2.32$
	82	2	$\frac{2}{82+2}$	$100 = 2.38$
	80	2	$\frac{2}{80+2}$	$\times 100 = 2.43$
	86	2	$\frac{2}{86+2}$	$100 = 2.26$
				<u>    </u> Average <u>2.33</u> <u>    </u>

TABLE.- IV (Contd.)

1	2	3	4	5
<i>Funaria hygrometrica</i>				
	112	5	$\frac{5}{112+5}$	$\times 100 = 4.33$
	114	5	$\frac{5}{114+5}$	$\times 100 = 4.20$
	110	5	$\frac{5}{110+5}$	$\times 100 = 4.34$
	112	5	$\frac{5}{112+5}$	$\times 100 = 4.33$
	110	5	$\frac{5}{110+5}$	$\times 100 = 4.34$
	111	5	$\frac{5}{111+5}$	$\times 100 = 4.31$
			Average	4.30

TABLE IV (Contd.)

1	2	3	4	5
<i>Funaria</i> <i>submarginata</i>	96	3	$\frac{3}{96+3}$	$\times 100 = 3.03$
	97	3	$\frac{3}{97+3}$	$\times 100 = 3.00$
	95	3	$\frac{3}{95+3}$	$\times 100 = 3.06$
	97	3	$\frac{3}{97+3}$	$\times 100 = 3.00$
	96	3	$\frac{3}{96+3}$	$\times 100 = 3.03$
	94	3	$\frac{3}{94+3}$	$\times 100 = 3.09$
				<u>Average 3.04</u>

TABLE - IV (Contd.)

1	2	3	4	5
<i>Pohlia flexuosa</i>	62	3	$\frac{3}{62+3} \times 100 = 4.61$	
	60	3	$\frac{3}{60+3} \times 100 = 4.76$	
	62	3	$\frac{3}{62+3} \times 100 = 4.61$	
	63	3	$\frac{3}{63+3} \times 100 = 4.54$	
	61	3	$\frac{3}{61+3} \times 100 = 4.68$	
	63	3	$\frac{3}{63+3} \times 100 = 4.54$	
				<u>Average 4.62</u>

TABLE-IV (Contd.)

1	2	3	4	5
<i>Hypnodon perpusillus</i>	60	2	$\frac{2}{60+2} \times 100 = 3.23$	
	62	2	$\frac{2}{62+2} \times 100 = 3.12$	
	63	2	$\frac{2}{63+2} \times 100 = 3.07$	
	60	2	$\frac{2}{60+2} \times 100 = 3.23$	
	58	2	$\frac{2}{58+2} \times 100 = 3.33$	
	59	2	$\frac{2}{59+2} \times 100 = 3.27$	
			Average	3.20

TABLE - IV (Contd.)

1	2	3	4	5
<i>Rhegmatodon</i>	78	3	$\frac{3}{78+3}$	$\times 100 = 3.70$
<i>orthostegius</i>	77	3	$\frac{3}{77+3}$	$\times 100 = 3.75$
	80	3	$\frac{3}{80+3}$	$\times 100 = 3.61$
	78	3	$\frac{3}{78+3}$	$\times 100 = 3.70$
	79	3	$\frac{3}{79+3}$	$\times 100 = 3.65$
	77	3	$\frac{3}{77+3}$	$\times 100 = 3.75$
				Average $\frac{3.69}{}$

TABLE-IV (Contd.)

1	2	3	4	5
<i>Rhynchosstegium javanicum</i>	104	4	$\frac{4}{104+4}$	$\times 100 = 3.75$
	102	4	$\frac{4}{102+4}$	$\times 100 = 3.77$
	106	4	$\frac{4}{106+4}$	$\times 100 = 3.63$
	102	4	$\frac{4}{102+4}$	$\times 100 = 3.77$
	105	4	$\frac{4}{105+4}$	$\times 100 = 3.66$
	104	4	$\frac{4}{104+4}$	$\times 100 = 3.75$
				Average 3.72

It may be seen that the stomatal indices of different species of mosses are specific as they are in the angiosperms (Trease, 1945), (Rangaswamy, 1959).

Further, measurements of the length and breadth of the stomata were made. An average of six readings in each case has been taken and tabulated as below:

TABLE-V

Name of the Mosses	No. of subsidiary cells per stoma	Length of the stoma (L) in $\mu$	Breadth of the stoma (B) in $\mu$	L/B Ratio
<i>Trematodon</i>				
<i>ceylonensis</i>	8	66	55	1.2
<i>Barbula indica</i>	10	33	27	1.2
<i>Funaria</i>				
<i>hygrometrica</i>	13	33	27	1.2
<i>Funaria</i>				
<i>submarginata</i>	10	55	44	1.2
<i>Pohlia flexuosa</i>	10	44	36	1.2
<i>Hypnodon</i>				
<i>perpusillus</i>	8	55	44	1.2
<i>Rhegmatodon</i>				
<i>orthostegius</i>	9	55	44	1.2
<i>Rhynchostegium</i>				
<i>javanicum</i>	10	49	39	1.2

It may be seen, therefore, that the length and breadth ratio of the stomata of the mosses seems to be uniformly similar.

### The operculum

The opercula of the capsules were drawn separately (*Plate Nos. XXII & XXIII*) to gain a comparative idea of their shape and size. They have been classified into four groups as under;

1. Convex as in *Funaria hygrometrica* (Fig. 368)
2. Cupula shaped as in *Anomobryum subnuditum* (Fig. 372)
3. Short beaked as in *Thuidium tamariscellum* (Fig. 380)
4. Long beaked as in *Rhynchostegium javanicum* (Fig. 381)

These groupings are in accord with the classification erected by Dunham (1951) for the opercula of the mosses of Northeastern United States.

The operculum of *Funaria hygrometrica* is slightly convex and cells are arranged in concentric rings (Fig. 369). It may persist even after maturity of the capsule though in most cases they fall off when the sporogonium ripens. There seems to be no correlation between the length of the capsule and the length of the operculum.

### The peristome

The peristome of the mosses is considered to be of very great significance in the identification of the

mosses. The description of the peristome in this treatise is in conformity with the findings of Dixon (1914), Cardot (1922), Conard (1954) and Watson (1954). The peristome in the haplolepideae is made up of one ring of teeth. Each tooth is composed of fused walls of two adjacent circles of cells. As the outer cell-wall extends the entire width of the tooth, the outer surface appears to have horizontal partitions. As the inner side of the tooth is derived from two boundaries, the surface shows a longitudinal line marking the boundaries of these cells. The outer surface of the peristome is termed as exostome and the inner endostome. The peristomial teeth of this group of mosses are often forked above as in *Fissidens schimidii* (Fig. 6).

The peristome is present in two rings in the diplolepideae. The inner peristome is more delicate and fragile than the outer and is usually lighter in colour as shown in *Pohlia flexuosa* (Fig. 127). The structure of the peristome teeth of this group of mosses is just the reverse of those of the haplolepideae. They have a zigzag line on the outer face as seen in *Rhizogonium spiniforme* (Fig. 166). The peristome is lacking in some mosses like *Funaria submarginata* (Fig. 120).

The colour of the peristome as seen in these mosses is generally yellow and reddish brown. Bright crimson colour as has been stated by Richards (1950) has not been met with so far.

### **The spore**

The study of the spores in mosses was made for the first time by Schimper (1836) though superficially.

MacClymont (1955) has studied the morphology of the spores of a large number\* of species in relation to their significance in systematics. Although these South Indian mosses have been described years ago, there is no complete description of the spores. Hence detailed information regarding the spores of these species in a comparative manner is recorded in this work

The spores of 4 species were studied following the general acetolysis method of Erdman (1952). The rest of the spores could not be studied in the same manner for want of adequate quantity of spore materials. So, temporary mounting of the spores in the safranin mixture, already referred to, was done. As it stains the spores suitably, observations and measurements could be made satisfactorily. Some spores were simply mounted in the dry condition. The camera lucida diagrams of the spores are found in Plate No. XXIV (Figs. 386-416). As recorded already in the chapter on morphological observations and in Table - III, the diameter of the spores varies from  $9\mu$  to  $27\mu$ , the smallest being observed in *Hyophila involuta* and the largest in *Funaria submarginata*.

It is interesting to note that the smallest moss in this study viz. *Hypnodon perpusillus* produces very large spores. Most of them were found to be spherical in shape. The colour of the spores in general, is brown. The sculpturing and the thickness of the Coat varies differently in different species.

---

\* 653 species.

In the present study of the sporophyte, the following observations may be noticed:

In the sporophytes of 34 species, 8 different forms of foot have been observed for the first time.

Measurements of the foot and capsule reveal that there is corresponding increase in length of the foot *pari-passu* with the increase in length of the capsule.

Though the nature of looping of the seta appears to have doubtful bearing on the structure of the peristome corresponding to the need for the regulation of spore dispersal their correlation is possible.

Stomata were observed in 8 species. The stomatal index of each one of these mosses, appears to be specific. The L/B ratio of the stomata is a constant factor.

The opercula are classified under 4 groups conforming to the classifications of Dunham (1951).

The spores have been studied in detail and in a comparative manner.

#### **Vegetative reproduction and regeneration:**

The vegetative reproduction in mosses is such a common phenomenon that Professor Goebel (1905) has expressed the opinion "almost every living cell of a moss can grow out into protonema and may produce gemmae of different kinds". The most important contributions have been made by Braithwaite (1395), Sainsbury (1935, 1938, 1952) and Correns (1899). So far as Indian mosses are concerned, Singh (1930) has contributed that the bulbils and gemmae are found in two species of mosses from

Mussoorie. Regeneration in mosses has been worked out by Meyer (1940), Narayanaswamy & Lal (1957), Kachroo (1954), Chopra & Sharma (1956) and Banerji & Sen (1957).

There seems to be no work on the South Indian mosses in this line. Vegetative reproduction and natural regeneration have been observed in several species of mosses as indicated below:

Bulbils and leafy buds were found in the leaf axils of *Campylopus nodiflorus*, *Brothera leana* (Fig. 417) and *Rhynchostegium javanicum* (Fig. 419). In these mosses the leafy buds possess 5 to 6 leaves. The leaves are all united together forming a bud from the base of which number of rhizoids arise. These buds get detached from the plant and under favourable conditions grow independently.

Multicellular club-shaped gemmae have been found in the axils of the leaves of *Barbula indica* (Fig. 73), *Bryum argenteum* and *Brachymenium exile* (Fig. 418). The gemmae of *Barbula indica* are found in clusters of 3 to 6 in the axils of apical leaves. In the axil, there is a small fleshy knob-like receptacle to which the gemmae are attached. They are in the form of club-shaped multicellular structures. To start with they are green in colour but later they become brownish. The gemmae in *Bryum argenteum* are also club-shaped but the head is narrower. They are found (6 to 10) in the axils of leaf. They resemble the brood bodies noticed in *Tortula papillosa* by Van Der Wijk (1932). The gemmae of *Brachymenium exile* exactly resemble those of *Bryum hemisphaericarpum* reported by Singh (1930). They are

also multicellular club-shaped bodies the larger head of which resembling the mandibles of insects.

Regeneration of mosses by the formation of rhizoids from detached and attached leaves has been found in *Leucobryum neilgherrense*. The rhizoids are formed from the apex of the leaf (Photomicrograph Fig. 564) and from the margin of the leaf (Photomicrograph Fig. 562) in this plant. Production of rhizoids from the base of the leaf was noticed in *Taxiphyllum taxirameum* (Photomicrograph Fig. 558) and in *Schlotheimia grevilleana*, *Sematophyllum subhumile* and *Pterobryopsis orientalis*. Profusely branched rhizoids were found to arise even from the young regions of the axis of *Leucobryum neilgherrense* and *Aerobryopsis longissima*.

Regeneration was also noticed by the formation of special asexual reproductive bodies in the form of tri- or cross-like structures named as "deciduous rhizoprotonemata" by Grout (1947). These bodies are formed when the young plants of *Barbula indica* are kept growing in petri-dishes on wet filter paper. After 3 to 4 days protonematal outgrowths arising from any part of plant are ending into "rhizoprotonemata". They are green in colour from the first appearance. Actual developmental studies were not made. It is reported that these bodies are formed when the plants of *Barbula gregaria* were treated by 2, 4-D (Narayanasamy & Lal, 1957). But no such herbicide was used in the present investigation.

## Anatomical Observations

Anatomy of the mosses in general has been worked out by Lorch (1931). Van Der Wijk (1932) has given an account of the morphology and anatomy of the mosses. Ruhland (1924) and Brotherus (1924) have left some information regarding the anatomy of the mosses. Tansley and Chick (1911) while describing the conducting system in *Polytrichaceae*, have referred to the contribution of Haberlandt (1895) to Moss Anatomy in glowing terms. However, the anatomical studies of the South Indian mosses have never before been attempted by any author.

Transverse sections of axis of 53 species of mosses and longitudinal sections of some of them have been investigated and the observations thereof, have been presented in this chapter. The diameter of the axis varies very much in different mosses. The structure of the moss axis in its cross section is not the same in all cases. The different parts of the tissue have been described with different terminology by different authors from time to time. The present investigation deals with the axis.

## 1) Description :

1.\* *Fissidens schimidii* C. M. Transverse section of axis ovate in outline  $142\mu \times 90\mu$ . Epidermis thick-walled not distinguishable from outer cortex; both forming hypodermal rind of  $18\mu$  to  $20\mu$  thickness; cells very small and cell walls heavily thickened. Inner cortex composed of two layers of very large and thin-walled cells. Central strand made up of pith; cells very small and thin-walled (Fig. 458).

2. *Ditrichum amoenum* (Thw. & Mitt.) Par. Transverse section of axis bluntly triangular in outline,  $120\mu$  in diameter. Epidermis not differentiated from outer cortex forming hypodermal rind of  $15\mu$  to  $20\mu$  thickness. Cells uniform and walls heavily thickened. Inner cortex with much thick-walled cells of 3 to 4 layers innermost layer of cortex with slightly larger cells. Central strand made up of pith, cells very small and thin-walled (Fig. 459).

4. *Campylopus nodiflorus* (C. M.) Jaeg. Transverse section of axis ovate in outline. Epidermis not distinguishable from the outer cortex; both layers forming rind of  $22\mu$  thickness. Cells heavily thickened, cell lumen elongated. Inner cortex of 4 to 5 layers of not very much thickened cell walls. Central axis hollow (Fig. 463) in the older regions. Younger axis with central narrow pith of thin-walled cells, the outer cortex and epidermis remaining the same.

5. *Thysanomitrium depallieri* Card. Transverse section of axis slightly ovate in outline  $198\mu \times 154\mu$ .

---

\* Number indicates the same serial number of mosses as referred to in the Morphological Observations.

Epidermis not distinguishable from outer cortex; both forming hypodermal rind of  $20\mu$  to  $25\mu$  thickness with heavily thickened cell walls. Inner cortex made up of 6 to 7 layers of polygonal cells gradually thickened outwards. Centre of the young axis with thin-walled pith cells; older regions with a hollow cavity (Fig. 464).

6. *Thysanomitrium lioneurum* (Ther. & Vard.) Broth. Transverse section of axis more or less round. Epidermis not distinguishable from outer cortex; cells heavily thickened. Inner cortex not differentiated; cells large; thin-walled, but becoming thicker walled outwards. Central strand made up of pith in the young condition. Centre of older axis hollow forming a small cavity.

8. *Dicranodontium fragile* (Hook.) Broth. Transverse section of axis irregularly ovate  $220\mu \times 195\mu$ . Epidermis not discernible from outer cortex; both forming rind of  $15\mu$  thickness; cells heavily thick-walled. Inner cortex not differentiated; cells large and polygonal, thin-walled but gradually becoming thicker-walled outwards. Central portion in older regions of axis hollow.

9. *Leucobryum neilgherrense* C. M. Transverse section of axis more or less roundish  $160\mu \times 175\mu$ , epidermis thick-walled. Rind conspicuous by absence. All cells beneath epidermis uniformly distributed without any differentiation of cortex or central strand. Cell-walls not much thickened (Fig. 432).

10. *Leucobryum scalare* C. M. Transverse section of axis more or less roundish  $350\mu$  to  $360\mu$  in diameter.

Epidermis thick-walled; outer cortex 3 to 4 layers of thick-walled cells forming rind. Entire tissue inside the rind made up of uniform and polygonal cells without any differentiation of cortex or central strand; cell-walls not much thickened. In older axis rind well laid. Epidermis not discernible. Inner tissue with uniformly thick-walled cells (Fig. 440).

12. *Barbula indica* (Schw.) Brid. Transverse section of axis roundish,  $154\mu$  in diameter. Epidermis not distinguishable from outercortex; both forming rind of  $30\mu$  thickness; cell walls heavily thickened. Inner cortex not differentiated; cells thin-walled. Central strand made up of pith of thin-walled cells.

16. *Hyophila validinervis* Card. and Vard. Transverse section of axis small ovate,  $220\mu$  to  $260\mu$  in outline. Epidermis not detectable from outer cortex; both forming rind of  $88\mu$  thickness; cells heavily thick-walled and elongated. Inner cortex not differentiated; cells uniformly polygonal, thin-walled but thick-walled in the outer layers. Centre hollowed into a cavity in the old axis (Fig. 465). In young axis centre made up of pith cells.

17. *Rhacomitrium javanicum* Bryol. Transverse section of axis roundish  $220\mu$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of  $25\mu$  thickness; cell walls heavily thickened. No differentiation of tissue inside rind, cells thick-walled, uniformly distributed.

18. *Funaria hygrometrica* Hedw. Transverse section of axis roundish  $264\mu$  in diameter. Epidermis thin-walled. Outer cortex 3 to 4 layers slightly thick-walled forming rind. Inside rind cells isodiametric, thin-walled uniformly distributed. In older axis epidermis and rind discernible but cell-walls thickened. Inner cortex thin-walled. Central strand made up of pith; cells very thin-walled and smaller.

19. *Funaria submarginata* Card. and Vard. Transverse section of axis small and roundish  $100\mu$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of  $22\mu$  thickness. Cell-walls very heavily thickened; cell lumen obliterated. Inside rind, axis hollowed (Fig. 456).

20. *Pohlia flexuosa*(Mitt.)Broth. Transverse section of axis slightly ovate  $160\mu \times 144\mu$ . Epidermis not distinguishable from outer cortical rind; cell-walls much thickened. Inner cortex with much thin-walled layers of cells. Central strand made up of pith; cells smaller and thin-walled.

25. *Rhodobryum giganteum*(Hook.)Par. Transverse section of axis roundish  $880\mu$  in diameter. Epidermis with thick-walled small cells, provided with epidermal hairs. Outer cortical rind made up of 4 to 5 layers of thick-walled cells. Inner cortex differentiated into 2 zones; the outer zone just beneath the rind made up of many layers of larger and thick-walled cells; the inner zone of inner cortex composed of 2 to 3 layers of much thick-walled smaller cells. The central strand in the form of

pith in younger axis; in older regions centre of the axis hollow (Fig. 467) forming a cavity.

26. *Mnium coriaceum* Griff. Transverse section of younger axis (Fig. 457) roundish  $500\mu$  in diameter. Epidermis not detectable from outer cortex; both forming rind of  $42\mu$  thickness made up of 3 to 4 layers of heavily thickened cell walls. Inner cortex not differentiated; 5 to 6 rows of thin-walled cells becoming thick-walled outwards. Central strand composed of collenchyma-like cells with angular thickening. In older axis central strand well formed with heavily thick-walled cells; innermost layer of inner cortex made up of larger cells (Fig. 470).

27. *Rhizogonium spiniforme* (L.) Bruch. Transverse section of axis irregularly pentagonal in outline  $242\mu \times 262\mu$ . Epidermis not distinguishable from outer cortex; both forming rind  $25\mu$  thickness with heavily thick-walled cells. Inner cortex not differentiated; cells thin-walled but gradually thick-walled outwards. Central strand consists of thin-walled small cells forming a pith.

28. *Bartramia madurensis* Dix. & Vard. Transverse section of axis irregular in shape  $220\mu \times 262\mu$ . Epidermis not distinguishable from outer cortex both forming rind of  $45\mu$  thickness; cells heavily thick-walled. Tissue inside rind not differentiated, made up of thick-walled cells (Fig. 442).

29. *Philonotis anisoclada* Card. and Vard. Transverse section of axis small and slightly roundish  $220\mu$

> 200 $\mu$ . Epidermis thick-walled; outer cortex of 2 layers of cells forming rind; cells narrow and thick-walled. Inner cortex extended to centre of axis; cells thick-walled. Centre of axis without strand or core but with a small cell surrounded by 4 to 5 thick-walled narrower cells (Figs. 449 and 450).

31. *Schlotheimia grevilleana* Mitt. Transverse section of axis ovate 262 $\mu$   $\times$  330 $\mu$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of 22 $\mu$  thickness; cells heavily thickened but lignification slightly different. Safranin stain not deep red as in others but light; hence thickening stippled. Tissue inside rind not differentiated; cell walls heavily thickened (Fig. 441).

32. *Trachypus bicolor* Rein. & Hornsch. Transverse section of axis ovately flattened 220 $\mu$   $\times$  330 $\mu$ . Epidermis not distinguishable from outer cortex; both forming 25 $\mu$  thick rind; cell walls very heavily thickened. Tissue inside rind made up of several layers of thick-walled narrow cells surrounding a very small narrow cell (Figs. 451 & 452).

33 *Trachypus humilis* Lindb. Var *humilis* Zanten. Transverse section of axis small. Epidermis not distinguishable from outer cortex; both forming rind of 22 $\mu$  thickness; cell walls heavily thickened. Inner tissue of axis composed of thick-walled large cells extending up-to centre of axis, possessing 10 to 12 layers of thick-walled narrow cells surrounding a single central cell.

34. *Trachypodopsis serrulata* (Beav.) Fleisch. Var. *crispatula* (Hook.) Zanten. Transverse section of axis

fairly large and irregularly rounding  $396\mu \times 462\mu$ . Epidermis not distinguishable from outer cortex; both forming  $45\mu$  thick rind; cell walls very heavily thickened. Inside rind several layers of large cells uniformly thick-walled. Centre of axis consisting of a rind of narrow 5 to 6 thick-walled cells enclosing two slightly elongated cells.

35. *Myurium warburgii* (C. M.) Fleisch. Transverse section of axis small ovate in outline  $155\mu \times 220\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $30\mu$  thickness; cell walls heavily thickened. Tissue inside rind not differentiated; cell-walls much thickened and in several concentric layers (Fig. 443).

36. *Pterobryopsis orientalis* (C. M.) Fleisch. Transverse section of axis ovate in outline  $420\mu \times 484\mu$ . Epidermis and outer cortex forming rind of  $25\mu$  thickness; cell walls so heavily thickened that cell lumen not visible. Inner cortex differentiated into outer zone and inner zone; outer zone consisting of 6-7 layers of thick-walled cells; inner zone made up of 10 to 12 layers of slightly thick-walled cells. Central strand organised with a group of very narrow and much thick-walled cells (Fig. 472).

37. *Papillaria crocea* (Hamp.) Jaeg. Axis flattened; transverse section of axis ovate in outline  $220\mu \times 310\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $42\mu$  thickness; cellwalls very heavily thickened. Inner undifferentiated tissue composed of uniformly thick-walled cells (Fig. 444).

38. *Aerobryopsis longissima* (Doz. & Molk.) Fleisch. Axis flattened; transverse section narrowly ovate  $528\mu \times 296\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $33\mu$  thickness. Tissue inside rind not differentiated; cells uniformly thick walled and distributed around a single small cell situated in the centre.

39. *Barbella tenax* (C. M.) Broth. Axis flattened; transverse section irregularly ovate  $126\mu \times 180\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $20\mu$  thickness; cell walls heavily thickened; cell lumen scarcely seen. Inside rind, axis hollowed out into large cavity (Fig. 468).

40. *Barbella determessi* (Rein & Card.) Fleisch. Transverse section of axis irregularly roundish  $176\mu$  in diameter. Epidermis and outer cortex forming rind of  $25\mu$  thickness; cell-walls heavily thickened. Cells inside rind much thick-walled. Central most cell slightly larger surrounded by a ring of 7 - 8 smaller cells.

41. *Barbella pendula* (Sull.) Fleisch. Transverse section of axis ovate in outline  $176\mu \times 143\mu$ . Epidermis and outer cortex forming rind of  $25\mu$  thickness; cell walls very heavily thickened. Tissue inside rind uniformly thick-walled; cell lumen narrow almost spindle shaped (Fig. 445).

42. *Floribundaria floribunda* (Doz. and Molk.) Fleisch. Transverse section of axis irregularly roundish  $242\mu$  in diameter. Single layer of epidermis distinguishable from outer cortex of three layers of

narrow thick-walled cells forming rind of  $22\mu$  thickness. Tissue inside rind containing uniformly thin-walled cells excepting the central region of 6 - 7 smaller cells.

43. *Meteoriopsis squarrosa* (Hook ) Fleisch. Axis flattened; transverse section of axis ovate  $396\mu \times 220\mu$  (Fig. 461). Epidermis not distinguishable from outer cortex both forming rind of  $20\mu$  thickness. Inner cortex differentiated into three zones, outer zone of thick-walled narrow cells, mid zone of larger cells and inner zone of thick-walled cells with narrow lumen. Central axis compressed\*. Lignification not very much.

44. *Meteoriopsis reclinata* (C. M.) Fleisch. Transverse section of axis ovate in outline  $440\mu \times 262\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $30\mu$  to  $40\mu$  thickness; cell walls heavily thickened. Inner cortex not differentiated; cells thin-walled but gradually becoming thick-walled outwards. Central strand organised + by pith-like thin-walled smaller cells (Fig. 460).

45. *Homaliodendron exiguum* (Bryol Jav.) Fleisch. Transverse section of axis ovate in outline  $242 \times 176\mu$ . One layer of cells of epidermis and 3 - 4 layers of outer cortex distinguishable; both forming rind of  $33\mu$  thickness; cell walls very much thickened. Tissue inside rind without any differentiation and uniformly thin-walle.

---

\* Similar feature recorded by Lorch (1931) in the Transverse section of seta of *Cinclidium stygium* Schw.

+ Similar structure observed by Lorch (1931) in the axis of *Mnium cinclidioides* Hueb.

46. *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. Transverse section of axis ovate in outline  $506_{\mu} \times 396_{\mu}$ . Epidermis not distinguishable from 6 - 7 layers of outer cortex both forming rind of  $55_{\mu}$  thickness; branch organised. Tissue inside rind not differentiated; cells isodiametric and uniformly thick-walled (Figs. 446 & 447).

47. *Distichophyllum succulentum* (Mitt.) Broth. Transverse section of axis irregularly roundish in outline  $220_{\mu}$  in diameter. Epidermis not distinguishable from outer cortex; both forming heavily thickwalled rind of  $30_{\mu}$  thickness. Inner cortex also thick-walled but differentiated by safranin, not deeply red as outer cortex, hence stippled (Fig. 434). Beneath inner cortex central strand of uniformly thin-walled and sinuated cells forming pith-like tissue.

48. *Hookeria acutifolia* Hook. Transverse section of axis ovate  $272_{\mu} \times 392_{\mu}$ . Epidermis made up of single layer of thick-walled cells; lumen smaller and distinct from the lumen of the cells inside. No differentiation of cortex; no rind; no strand. Tissue of axis below epidermis consists of isodiametric pentagonal cells. Cell-wall not much thickened, turns light red by safranin (Figs. 430 & 431).

49. *Hypopterygium tenellum* Lac. = *H. ceylanicum* Mitt. Transverse section of axis roundish  $308_{\mu}$  in diameter. Epidermis and cortex distinguishable. Cell walls very much thickened but turning light red by phloroglucin and safranin. Cortex large not differentiated. Central strand made up of pith of uniform and isodiametric very thin-walled cells (Fig. 433).

50. *Rhegmatodon orthostegius* Mont. Transverse section of axis irregularly ovate  $396_{\mu} \times 286_{\mu}$ . Epidermis not distinguishable from outer cortex; both form rind of  $25_{\mu}$  thickness; cell walls heavily thickened. Tissue inside rind not differentiated uniformly thick-walled.

51. *Herpetineurum toccoe* (Sull. & Lesq.) Card. Transverse section of axis irregularly ovate  $374_{\mu} \times 310_{\mu}$ . Epidermis and outer cortex not distinguishable. Both form rind of  $44_{\mu}$  thickness; cell walls heavily thickened. Inner cortex not differentiated; cells uniformly thick-walled. Central strand present in the form of pith of very thin walled cells.

52. *Thuidium tamariscellum* (C. M.) Bryol. Jav. Transverse section of primary axis roundish  $440_{\mu}$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of  $33_{\mu}$  thickness. Tissue inside rind not differentiated; cells uniformly thin-walled.

53. *Pleuropus nilghiriense* (Mont.) Toyama. Transverse section of axis slightly ovate  $330_{\mu} \times 310_{\mu}$ . Epidermis not distinguishable from outer cortex; both forming rind of  $40_{\mu}$  thickness; cells heavily thickened; Inner cortex differentiated into inner and outer zones; cells of the inner zone very thin-walled and cells of the outer zone uniform, thick-walled. Central axis hollowed into a narrow cavity.

54. *Rhynchostegium javanicum* (Be'.) Besch. Axis flattened. Transverse section of axis  $418_{\mu} \times 286_{\mu}$  ovate

(Fig. 462). Epidermis not distinguishable from outer cortex; both forming rind of  $18\mu$  to  $20\mu$  thickness. Tissue inside rind composed of uniformly thin-walled cells. Cells of central axis compressed forming long streak.\*

55. *Eryt'irodontium julaceum* (Hook.) Par. Transverse section of axis irregularly ovate  $286\mu \times 242\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $20\mu$  to  $40\mu$  thickness; cell walls heavily thickened; cell-lumen very small. Cells inside rind uniformly thick-walled; cell-lumen larger except in central axis; 3 to 4 cells with narrow lumen found in the centre (Fig. 453).

56. *Foreauella indica* Dix. & Vard. Transverse section of axis irregularly ovate  $264\mu \times 176\mu$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of  $45\mu$  thickness; cell walls heavily thickened. Tissue inside rind composed of uniformly thick-walled cells. Centrally-placed small single cell around a ring of seven radiating cells observable (Fig. 448).

57. *Sematophyllum subhumile* (C.M.) Fleisch. Axis very narrow; transverse section very irregular in outline  $110\mu \times 70\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of  $12\mu$  thickness; cell walls heavily thickened. Tissue inside rind composed of thick-walled cells; inner cortical cells very small (Fig. 439).

58. *Warburgiella leptorrhynchoides* (Mont.) Fleisch. Axis flattened; transverse section irregularly ovate

---

\*Similar feature recorded by Lorch (1931) in the transverse section of seta of *Cinclidium stygium* Schw.

154 $\mu$   $\times$  88 $\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of 10 $\mu$  thickness. Tissue inside rind not differentiated; cells uniformly thin-walled (Fig. 438).

59. *Hypnum cupressiforme* Hedw. Transverse section of axis irregularly ovate 176 $\mu$   $\times$  132 $\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of 12 $\mu$  thickness; cell walls heavily thickened; inner cortex differentiated with peripheral thick-walled cells and central thin-walled cells. Very small narrow lumen in the central most region of axis with 7 radially elongated cells observable (Fig. 456).

60. *Taxiphyllum taxirameum* (Mitt.) Fleisch. Transverse section of axis almost roundish, 188 $\mu$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of 15 $\mu$  thickness; cell walls heavily thickened; tissue inside rind not differentiated, cells thin-walled. Centrally placed single cell surrounded by a ring of 7 - 8 elongated and radiating cells as in (Fig. 456).

61. *Ctenidium lychnites* (Mitt.) Broth. Axis flattened; transverse section irregularly ovate 264 $\mu$   $\times$  154 $\mu$ . Epidermis not distinguishable from outer cortex; both forming rind of 15 $\mu$  thickness; cells inside rind thin-walled but peripheral layers slightly thick-walled. Centrally placed single narrow very small cell possessing ring of 5 - 6 radiating and elongated cells (Fig. 455).

62. *Atrichum aculeatum* Card. & Vard. Transverse section of axis almost roundish 264 $\mu$  in diameter.

Epidermis not distinguishable from outer cortex; both forming rind of  $10\mu$  to  $15\mu$  thickness; cell walls heavily thickened, cell lumen very small. Inner cortex differentiated into three zones; outer zone composed of two incomplete rows of thick-walled cells; mid zone made up of 3-5 rows of thick-walled cells of smaller size; inner zone composed of two rows of thin-walled cells with angular thickening. Inside inner zone of inner cortex central strand of thick-walled cells found in older axis (Fig. 474) But in younger axis central strand formed by very thin walled pith cells ( Fig 473).

63. *Pogonatum microstomum* (R. Br ) Brid. Transverse section of axis very large, roundish  $440\mu$  in diameter: Epidermis not distinguishable from outer cortex; both forming rind of  $50\mu$  thickness; cell walls very heavily thickened. Inner cortex differentiated into two zones: outer zone beneath the rind made up of several layers of highly thick-walled smaller cells; inner zone composed of several layers of larger and thin-walled collenchyma-like cells. Central core with angular projections forming the central strand composed of very heavily thickened very small and narrow cells.

64. *Pogonatum neesii* (C.M.) Mitt. Transverse section of axis very irregular in outline  $264\mu$  in diameter. Epidermis not distinguishable from outer cortex; both forming rind of  $45\mu$  thickness; cell walls so heavily thickened that cell-lumen hardly seen. Inner cortex differentiated into outer zone and inner zone;\* outer zone made up of

---

\* Similar structure of the axis has been reported by Frye and Ferguson (1943) in *Pogonatum liebmannianum* C. Muell. but without angular projections of the central strand.

6-7 layers of very much thick-walled cells, cell lumen very narrow; inner zone composed of 4 or 5 concentric layers of collenchyma-like cells. Central axis made up of strand with 5 or 6 angular projections; cells small walls very much thickened, cell lumina of strand very narrow in peripheral 5-6 layers and slightly larger in the central region (Fig. 475).

## 2) Results :

Transverse sections of the axes of 53 species of mosses were taken. The axes of the rest of the 11 species such as: *Trematodon ceylonensis*, *Brothera leana*, *Tortula muralis*, *Trichostomum cylindricum*, *Weisia macrospora*, *Hyophila involuta*, *Brachymenium exile*, *Brachymenium leptostomoides*, *Anomobryum subnuditum*, *Bryum argenteum*, and *Hypnodon perpusillus* could not be sectioned. The structure of the transverse section of the axis may be differentiated into three regions: the epidermis, the cortex and the central strand. The outer most is the epidermis. The cortex is often very much thickened in the outer portion forming a rind. The cells of the rind are very heavily thick-walled. The inner cortex may or may not be differentiated. Inside the cortex, there is a central strand or core of the axis. This is made up of either thin-walled or thick-walled cells.

There is a good deal of variation in the structure of the axis. They may be outlined as under:-

1. The epidermis and the rind may not be distinguishable but may be uniform (Fig. 431).

2. The rind may be conspicuous by its absence (Fig. 433).

3. The cortex may not be differentiated (Fig. 433) and cell-walls of pith be sinuate (Fig. 435).

4. There may absolutely be no differentiation of the tissue beneath the rind (Fig. 436). The cells may either be thin-walled (Fig. 438) or thick-walled (Fig. 440).

5. The differentiation of the tissue inside the rind may be initiated by the presence of a centrally situated small cell with an outer ring of radiating cells (Figs. 449 to 456).

6. Where a central strand is present it may be composed of thin-walled pith cells (Fig. 457) in the younger regions of the axis. In the older regions the centre of the axis may be made up of very heavily thick-walled cells forming this strand. The strand is differentiated in *Pogonatum microstomum* and *Pogonatum neesii*.

7. Instead of a pith in the centre a hollow cavity may be present (Figs. 463–469).

The anatomical features so observed have afforded a critical basis for the classification of the mosses.

Longitudinal sections of the axes of the following mosses were taken:—

*Pogonatum microstomum*

*Pogonatum neesii*

*Rhacomitrium javanicum*

*Rhizogonium spiniforme*

*Homaliodendron flabellatum*

*Thuidium tamariscellum*

It has been found that the cells of the epidermis and those of the rind are scarcely discernible as the thickening of the cell-wall is often enormous. The cells of the inner cortex are elongated and the cross walls are oblique. They are not so much thickened as the cells of the rind. The cell wall of the central strand are also equally thickened and they take up deep Safranin stain. The double staining process with Safranin and Fast Green or Haematoxylin and Eosin differentiates the central strand very clearly from the outer layer of large cells.

The cells of the axis of certain mosses were separated with the help of Schultze's maceration fluid and stained suitably with Phloroglucin. It is found that lignification has taken place in the cells of the rind and of the central strand much more than in other regions.

The anatomy of the leaves of a few mosses was studied. The transverse section of the leaf of *Fissidens schimidii* reveals the presence of dorsal lamella as an outgrowth (Fig. 5).

The cross sections of the leaf of *Leucobryum scalare* and *Leucobryum neilgherrense* (Figs. 51 & 58) have been found to possess multicellular structure. The former shows a row of collenchyma-like corner thickened cells in the centre. Similar observations were made by Brotherus (1925) in *Leucobryum sanctum*, and by Van Der Wijk (1932) and Conard (1956) in *Leucobryum glaucum*. The cross section of the leaves of the two species of *Pogonatum* is interesting. It resembles the transverse section of the thallus of Marchantiales. There are

longitudinal rows of cells arranged one above the other containing chloroplasts. They form the lamellae. The uppermost rows of cells in *Pogonatum neesii* are different from those of *Pogonatum microstomum*. In all other cases the leaves are made up of single row of cells. In the region of the mid rib thick walled cells in two or three rows are found.

It is interesting to note that the structure of the axis in the young region is considerably different from that of the axis in the older regions. For example the axis of *Funaria hygrometrica* according to Campbell (1930) possesses a central cylinder of thin-walled cells when young which gradually gets converted into a thick-walled central strand at maturity. This may be true of some mosses as *Polytrichum commune* (Tansley and Chick 1901); but not in all cases.

Lorch (1931) has sketched transverse section of the axis of *Campylopus polytrichoides* both in the young and old conditions. The former shows a well defined pith made up of thin-walled parenchyma-like cells and the latter has a hollow cavity in the centre. This feature is exhibited in some of the Nilgiri mosses such as :-

*Campylopus nodiflorus* (Fig. 463)

*Thysanomitrium depallieri* (Fig. 464)

*Thysanomitrium lioneurum*

*Dicranodontium fragile*

*Hyophila validinervis* (Fig 465)

*Funaria submarginata* (Fig 466)

*Rhodobryum giganteum* (Fig 467)

*Barbella tenax* (Fig. 468)  
*Pleuropus nilghiriense* and  
*Entodon plicatus* (Fig. 469.)

The second outstanding observation is the presence of a rind consisting of 4 to 7 rows of very much thick-walled cells in most cases. Lorch (1931) has drawn attention to this fact by showing the different patterns of thickening of the rind along with the epidemis. This rind is found to be formed and well-laid even in the young condition. It is conspicuous by its absence in a few cases such as *Distichophyllum succulentum* and *Hookeria acutifolia*. In these two species the epidermis can be clearly distinguished as being made up of barrel-shaped cells (Fig. 430). Brotherus (1924) has also observed this feature in *Sarconeurum glaciale* (Hook fil. et Wils.) Card. et Bryhn. and in *Hookeriopsis incurva* (Hook et Arev.) Broth.

In some moss axes, it has been noticed that just inside the rind, the entire central portion of the axis is made up of undifferentiated and uniformly thin-walled cells as in the following:

*Leucobryum scalare* (Fig. 440)  
*Rhacomitrium javanicum*  
*Schlotheimia grevilleana* (Young, Fig. 441.)  
*Schlotheimia grevilleana* (Old, Fig. 442.)  
*Myurium warburgii* (Fig. 443)  
*Papillaria crocea* (Fig. 444)  
*Barbella pendula* (Fig. 445)

*Hcmaliodendron flatellatum* (Fig. 446)  
*Rhegmatodon orthostegius* and  
*Herpetineurum toccoeae*.

Lorch (1931) has observed a similar structure in the axis of *Pilotrichidium antillarum*. The following mosses possess yet another kind of central structure in the axis. Instead of the thin-walled or thick-walled cell distribution in the centre there occurs a centrally situated cell having a narrow or large lumen often surrounded by a ring of cells radiating outwards.

*Philonotis anisoclada* (Fig. 449)

*Trachypus bicolor* (Fig. 451)

*Aerobryopsis longissima*

*Floribundaria floribunda*

*Erythrodontium julaceum* (Fig. 454)

*Hypnum cupressiforme* (Fig. 456) and

*Ctenidium lychnites* (Fig. 455).

The central strand (Zenral-strang) which has been referred to by some authors as the central core of conducting strand is found in the two species of *Pogonatum* belonging to the *Polytrichaceae*. Though *Polytrichum* has not been so far found to grow in South India, its relatives,\* viz. *Pogonatum microstomum* and *Pogonatum neesii*, are abundant in the South Indian

---

\* The other available species *Pogonatum aloides*, *P. hexagonium*, *P. inflexum* and *P. junghuhnianum* were not included in this work due to difference of opinion in the specific determination.

Hill stations. It has been found that this genus exhibits almost the same structure as *Polytrichum*. The central strand is well developed. The strand possesses four to six angular projections abutting on the cortex. Outside this central strand there is a layer of comparatively thin-walled, sometimes large cells followed by a few layers of thin-walled cells lying beneath the rind (Figs. 474 & 475).


Having these facts in view the 53 species of mosses belonging to 26 families have been classified on the basis of the following principles :—

1. The presence or absence of the rind.
2. The differentiation of the tissue inside the rind or the same without differentiation.
3. The presence or absence of the central strand.
4. The differentiation of the central strand or the same without differentiation.
5. The presence of a pith or cavity in the centre.
6. The presence of a small cell in the centre of the axis surrounded by a ring of cells.

The detailed classification proposed in this treatise is presented in Table-VI.

TABLE VI

SYSTEM OF CLASSIFICATION OF THE MOSSES  
 PROPOSED HERE ON THE BASIS OF ANATOMICAL STRUCTURE \*



\* Very interesting! A. J. Sharp.

It may be observed further:

1. that the pith in the young axis has become the central strand in the following :

*Mnium coriaceum* (Figs. 457 & 470)

*Artichum aculeatum* (Figs. 473 & 474)

2. that the pith in the centre of the young axis has developed a hollow cavity in the following:

*Campylopus nodiflorus* (Fig. 463)

*Thysanomitrium depallieri* (Fig. 464)

*Hyophila validinervis* (Fig. 465)

*Funaria submarginata* (Fig. 466)

*Rhodobryum giganteum* (Fig. 467)

3. that the uniformly thick walled cells in the centre of the young axis have developed a single central cell in the old axis in the following :

*Schlotheimia grevilleana* (Fig. 441)

*Philonotis anisoclada* (Figs. 449 & 450)

4. that the centrally situated single cell in the young axis has developed a hollow cavity in the following :

*Barbella tenax* (Fig. 448)

## Physiological Observations

Different aspects of physiology of mosses have been dealt with by several authors, most important among them being Haberlandt (1886), Lorch (1925), Goebel (1905), Oltmanns, (1884) Stelmach (1926) and Brilliant (1927). This wide and vast field has many facets but in this investigation a few experiments on certain aspects alone were under-taken. They are (1) on the effect of pH and hormones on the growth of moss plant; (2) the production of rhizoids from the foot of the sporophyte of mosses; and (3) comparative studies in the osmotic concentration of the gametophytic and sporophytic tissues of mosses.

There have been many limitations in the pursuit of this kind of study. Most of the experiments had to be conducted in the hill stations - the very home of the mosses. In many cases the sporophytes are not usually produced and when they are produced they are short-lived. As Garjeanne (1932) has pointed out, "a reliable method for cultivating the bryophytes on their natural substrata in the laboratory would however be of great value to the physiologist." As per the suggestion, certain experiments were carried on in the same manner in the University Laboratory. Some tufts of mosses with sufficient soil were uprooted and brought from the hills.

They were growing alright for sometime but in the long run they died even when they were enjoying normal conditions of life of the Hills.

The effect of pH on the growth of mosses:

Reference has already been made (P. 8) to the effect that the soil of the following mosses has been found to be acidic:

Name	pH
<i>Fissidens schimidii</i>	5.4 to 5.6
<i>Ditrichum amoenum</i>	5.2 to 5.4
<i>Leucobryum neilgherrense</i>	5.6 to 5.8
<i>Barbula indica</i>	6.2 to 6.4
<i>Bartramia madurensis</i>	4.5 to 4.6
<i>Barbella pendula</i>	6.0
<i>Pogonatum neesii</i>	5.6 to 5.8

These findings only confirm the observations of Olsen (1923) that "each species had a different optimum pH, corresponding more or less with the pH of its natural habitat."

One of the mosses referred to above, *Barbula indica* grows well on the open terrace of the laboratory. The plants were treated with buffer solutions of different concentrations in the following manner.

The terrace is plastered with lime-mortar. This moss grows during winter in shade cast by the parapet wall. In selected plots, small bunds of 50 cm × 30 cm

were constructed. The plots were protected from sun and rain with thatches of palm leaves. Daily towards evening the plots were irrigated with equal quantity of buffer solutions of different concentrations of varying pH from 4.6 to 9. The experiments were carried on for fifteen days. The control plot of the same dimensions was irrigated with same quantity of distilled water. The initial height of the plants in all the plots was 5 mm. Observations were taken once in every five days. The results are tabulated as under.

TABLE-VII

1	2	3	4	5	6
Height of plant	pH 4.6	pH 6	pH 8	pH 9	Control
Fifth day	7 mm	6 mm	5.5 mm	showed signs of wilting	6 mm
Tenth day	9.5 mm	8 mm	showed signs of wilting	plants died	7.5 mm
Fifteenth day	12 mm	10 mm	Plants died	Plants died	9 mm

It may be seen that increase in pH (on acidic side) is directly proportional to the growth of the stem.

In the same manner growth hormones such as Indole Acetic Acid (I A A), Indole-Butyric Acid (I B A), and 2,4-D were irrigated to investigate the nature of growth

of *Barbula indica* in a few other plots. The concentration of the hormones used was 1%. The control plot was irrigated with same quantity of distilled water. The observations are recorded in Table No. VIII.

TABLE-VIII

1	2	3	4	5
Height of Plant	I A A	I B A	2,4-D	Control
Fifth day	6 mm	6 mm	6 mm	6 mm
Tenth day	8 mm	9 mm	7.5 mm	7.5 mm
Fifteenth day	9.5 mm	10.5 mm	9 mm	9 mm
Twentieth day	12 mm	15.5 mm	12.5 mm	12 mm
Twenty-fifth day	13 mm	18 mm	16 mm	15 mm

It may be seen that I B A induces greater growth of the plants than the other hormones. Microscopic examination of the plants treated with 2,4-D reveals that the leaves upon the shoots were generally spaced farther apart though no appreciable increase in height could be noticed. This finding is in confirmation of the observations of Prior (1952). Patterson (1957) found out that I A A. accelerated early protonematal growth and increased the percentage of spore germination. But in the experiment on *Barbula indica* I A A. seems to have very little effect.

### Induced production of rhizoids:

Several experiments were conducted in the hill stations of Kodaikanal and Ootacamund in two different seasons of the year, to induce the production of rhizoids from the foot of the sporophyte using the growth hormones.

The following species of mosses were available with sporophyte :

*Ditrichum amoenum*

*Funaria hygrometrica*

*Bryum argenteum*

*Pohlia flexuosa*

*Anomobryum subnuditum*

*Schlotheimia grevilleana*

*Thuidium tamariscellum*

*Atrichum aculeatum*

*Pogonatum microstomum*

*Pogonatum neesii*

Fairly young sporophytes of the foregoing mosses were carefully removed from the tip of the gametophore and transferred into small vials containing the growth hormones in different concentrations. The hormones used were: (i) Indole-3- Acetic Acid; (2) Indole-Butyric Acid and (3) 2,4-D. Various concentrations of the hormones: .5 ppm., 1 ppm, 5 ppm., and 10 ppm. were prepared in distilled water. Besides, an extract of the gametophyte was also used. This was prepared by pounding the gametophytic part of the mosses in glass mortar after removing the sporophyte. The pounded material with

distilled water was used as such as an extract. A control in distilled water was kept for each experiment. Observations were taken every sixth hour for three days. The experiments were repeated every time with 10 sporophytes in each case. As rhizoids were formed only in I B A between 12 hours and 30 hours, the results have been tabulated accordingly as under:

TABLE-IX

		No. of sporophytes producing rhizoids out of 10			
Name of Gameto- moss and phytic duration extract		Indole-Butyric Acid			
		.5 ppm.	1 ppm.	5 ppm.	10 ppm.
1	2	3	4	5	6
<i>Ditrichum amoenum</i>					
12 hrs.	—	—	—	—	—
18 hrs.	2	—	—	—	—
24 hrs.	—	—	—	3	—
30 hrs.	—	—	—	—	—
<i>Funaria hygrometrica</i>					
12 hrs.	—	—	—	—	—
18 hrs.	—	—	—	5	—
24 hrs.	3	—	—	—	—
30 hrs.	—	—	—	—	—

TABLE-IX (Contd.)

	1	2	3	4	5	6
<i>Bryum</i>						
<i>argenteum</i>						
12 hrs.	1	—	—	—	—	—
18 hrs.	1	—	—	—	—	—
24 hrs.	—	—	—	2	—	—
30 hrs.	—	—	—	—	—	—
<i>Pohlia</i>						
<i>flexuosa</i>						
12 hrs.	2	—	—	—	3	—
18 hrs.	—	—	—	—	—	—
24 hrs.	—	—	—	—	—	—
30 hrs.	—	—	—	—	—	—
<i>Anomobryum</i>						
<i>subnuditum</i>						
12 hrs.	—	—	—	—	—	—
18 hrs.	—	—	—	—	2	—
24 hrs.	4	—	—	—	—	—
30 hrs.	—	—	—	—	—	1
<i>Schlotheimia</i>						
<i>grevilleana</i>						
12 hrs.	—	—	—	—	—	—
18 hrs.	2	—	—	—	—	—
34 hrs.	—	—	—	1	—	—
30 hrs.	—	—	—	—	—	1

TABLE-IX (Contd.)

	1	2	3	4	5	6
<i>Thuidium</i>						
<i>tamariscellum</i>						
12 hrs.	2	—	—	—	—	—
18 hrs.	—	—	—	—	3	—
24 hrs.	—	—	—	—	—	—
30 hrs.	—	—	—	—	—	—
<i>Atrichum</i>						
<i>aculeatum</i>						
12 hrs.	—	—	—	—	—	—
18 hrs.	2	—	—	—	—	—
24 hrs.	—	—	—	—	—	2
30 hrs.	—	—	—	—	—	—
<i>Pogonatum</i>						
<i>microstomum</i>						
12 hrs.	—	—	—	—	—	—
18 hrs.	—	—	—	—	—	—
24 hrs.	2	—	—	—	—	1
30 hrs.	—	—	—	2	—	—
<i>Pogonatum</i>						
<i>neesii</i>						
12 hrs.	—	—	—	—	—	—
18 hrs.	—	—	—	—	—	—
24 hrs.	3	—	—	—	—	—
30 hrs.	—	—	—	—	—	3

It may be observed that rhizoids have been produced in the extract of the gametophyte in all cases. Moreover, 5 ppm. of I B A has better effect on the formation of rhizoids from the tip of foot than other concentrations of the same hormone. Rhizoids are generally produced in larger number from the perichaetial region of the of the sporophyte than from the tip of the foot. The observations are in conformity with the findings of La Rue (1942), so far as the production of rhizoids are concerned. But, he stimulated rhizoidal production with I A A in lanolin and I B A in lanolin on detached leaves and sporophytes of several species of mosses. It is to be noted that in the present experiment I B A in distilled water itself induces the formation of rhizoids. It is not necessary that solution of I B A should be made in lanolin.

#### **Osmotic concentration of gametophyte and sporophyte:**

It is well known that the sporophyte of moss depends partly on the gametophyte for its sustenance. The osmotic concentration of the cells of the gametophyte and of the sporophyte was studied with a view to find the difference in osmotic pressures in them. Plasmolytic method of de Vries was adopted using different concentrations of sucrose solution. Well developed leaves of mosses and thin free hand sections of the axis were also employed as the gametophytic tissue. Peels of the wall of the capsule specially from the apophysis region and sections of the seta near the capsule represented the sporophytic part. As Stocking (1956), Bonner and Galston (1951) have pointed out the concentration of the solution at incipient plasmolysis in about 50% of the cells, was taken as isotonic. The leaf cells and the

cells of the axis did not show any difference of osmotic concentration. So also, the difference in osmotic concentration of the cells of the capsule wall and the cells of the seta was negligible in most cases. The atmospheric temperature at Kodaikanal where these experiments were conducted was 20 C. Data of A. Ursprung, and G. Blum (1916) as quoted by Meyer, Anderson and Swanson (1955) were taken for the estimation of atmospheric pressures to different molarity. Only seven mosses were available with young sporophytes in sufficient quantities. The experiments were repeated several times to ascertain the proper isotonic solution and the observations are tabulated as under:

TABLE-X

Plasmolysis by Sucrose solution						
Name of moss	LEAF (Gametophyte)			CAPSULE (Sporophyte)		
	Concentration of sucrose %	Molarity	Osmotic pressure (O.p) at 20° Atm.	Concentration of sucrose %	Molarity	Osmotic pressure (O.p) at 20° Atm,
1	2	3	4	5	6	7
<i>Bardula indica</i>	5	.15	4	20	.58	17.1
<i>Funaria hygrometrica</i>	5	.15	4	25	.73	22.7
<i>Anomobryum subnuditum</i>	15	.45	12.7	30	.88	28.8

TABLE-X (Contd.)

1	2	3	4	5	6	7
<i>Bryum</i>						
<i>argenteum</i>	30	.88	28.8	55	1.6	73.9
<i>Thuidium</i>						
<i>tama-</i>						
<i>riscellum</i>						
(branch						
leaf)	10	.29	7.8	20	.58	17.1
<i>Pogonatum</i>						
<i>micro-</i>						
<i>stomum</i>	45	1.31	52.2	55	1.6	73.9
<i>Pogonatum</i>						
<i>neesii</i>	40	1.71	43.7	50	1.46	62.8

It may be seen that in all cases the osmotic pressure of the sporophyte is always higher than that of the gametophyte.

The ionising substances such as Sodium chloride and Potassium nitrate were also used for the determination of osmotic concentration in the same plasmolytic method. The isotonic percentages of the solutions which brought about incipient plasmolysis in the gametophytic and sporophytic tissues are tabulated. (Table-XI).

TABLE-XI

Name of mosses	Potassium nitrate (isotonic percentage)			
	Gametophyte Transverse section of axis	Leaf	Sporophyte T. S. of seta	Capsule
1	2	3	4	5
<i>Barbula indica</i>	—	5	30	30
<i>Funaria hygrometrica</i>	—	5	—	30
<i>Anomobryum Subneditum</i>	—	10	20	25
<i>Bryum argenteum</i>	—	7.5	—	25
<i>Thuidium tamariscellum</i>	10	10	20	20
<i>Schlotheimia grevilleana</i>	—	30	40	40
<i>Rhegmatodon orthostegius</i>	—	30	—	40
<i>Pogonatum neesii</i>	30	30	40	40

It may be seen that the cells of the sporophyte have higher osmotic concentration in all cases than the cells of the gametophyte.

Apart from the foregoing, biochemical tests were done in a rock moss *Schlotheimia grevilleana*. Kjeldahl's method of Nitrogen estimation was followed. Five grams (dry weight) of this plant were taken up. It was found after repeated experiments that 0.6260% of Nitrogen is present in this moss.

## Ecological observations

Warming (1909) recognised the mosses as muscoid growth-forms and as a separate entity. For a proper ecological study of these forms, exact measurements of ecological factors, as rightly pointed out by Richards (1932) are needed. Though much more detailed study of the factors of the habitat of these mosses have to be made for entering into the inner realms of ecology, an attempt has been made to present the various factors that come into play in the study of the moss plants in their peculiarly complex environment.

Along with the external morphological details, the places of occurrence of the moss plants have been given in the preceding chapter. They are growing in the mountainous regions within the belt of  $9^{\circ}$  to  $12^{\circ}$  latitude and  $76^{\circ}$  to  $79^{\circ}$  longitude. (For map see elsewhere in this book). Within this area the most important moss growing places are the Kodaikanal (Kodai) Hills usually referred to as Upper Palnis in the Madura District and The Nilgiri Hills (Ootacamund or Ooty) in the Nilgiri District of South India.

*Altitude* : The altitude of the hills ranges between 1000 metres to 2900 metres. These hills are favoured with rain by two monsoon winds, the south west in

the months of May and June, the north east during October and November. The latter monsoon actually brings the little rainfall for this area. The monthly average rainfall and the maximum and minimum temperature of these two regions for three years consecutively during which period plants were collected, have been tabulated. (Table-XII).

**TABLE - XII**  
**MONTHLY AVERAGE TEMPERATURE AND TOTAL MONTHLY RAIN FALL AT**  
**Kodaikanal,**

Month	1956			1957			1958		
	Temperature		Rain fall	Temperature		Rain fall	Temperature		Rain fall
	Maxi- mum oF.	Mini- mum oF.	in mm	Maxi- mum oF.	Mini- mum oF.	in mm	Maxi- mum oF.	Mini- mum oF.	in mm
1	2	3	4	5	6	7	8	9	10
January	61	45	49	63	45	2	63	48	23
February	62	45	5	65	47	29	64	49	12
March	68	51	29	66	49	33	66	50	186
April	66	53	159	67	52	160	67	52	164
May	68	54	67	67	54	120	68	55	240
June	64	53	128	66	55	106	69	55	53
July	62	51	25	63	53	135	63	52	60

TABLE-XII (Contd.)

	1	2	3	4	5	6	7	8	9	10
August		63	52	187	63	53	142	64	53	121
September		63	52	124	64	52	112	65	52	114
October		61	51	348	63	52	285	63	52	177
November		60	50	343	61	50	349	63	50	211
December		61	46	57	61	49	197	61	46	48
Yearly total				152.1			167.0			140.9
Rain fall										

TABLE-XII (Contd.)

Ootacamund		1956					1957					1958					
		Tempe- rature		Rain fall		1	Temperature		Rain fall		7	Tempera- ture		Rain fall		9	10
		Maxi- mum oF.	Mini/ mum oF.	in mm	in mm		Maxi- mum oF.	Mini mum oF.	in mm	in mm		Maxi- mum oF.	Maxi- mum oF.	in mm	in mm		
Month	2	3	4	5	6	8	9	10	11	12	13	14	15	16	17	18	
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
January	73	37	7	77	33	0	73	40	2								
February	72	38	0	76	39	9	74	41	24								
March	79	45	0	72	41	38	72	45	136								
April	79	48	136	75	47	46	75	47	45								
May	77	49	83	71	50	670	78	51	187								
June	72	50	169	69	51	153	72	52	75								
July	67	49	80	66	50	100	66	51	213								

TABLE-XII (Contd.)

1	2	3	4	5	6	7	8	9	10
August	70	48	63	71	51	77	70	50	198
September	71	49	59	74	45	25	71	46	80
October	71	48	389	71	50	257	71	48	102
November	70	46	138	70	49	233	71	44	228
December	72	38	14	71	44	91	70	40	7
Yearly total			114.7			169.9			129.7
Rain fall									

## Temperature

It may be seen that the range of temperature is 33 F, to 79° F. and yearly rainfall is between 100 cc. to 170 cc. There is no snow fall but during the winter months of December and January, the hills are covered by mist specially towards the evenings. It is interesting to note that many species of mosses such as *Bryum argenteum*, and *Funaria hygrometrica* of the temperate climate grow fairly well in these regions. The tropical moss families, are well represented in this belt such as *Pterobryaceae* with 10 species, *Meteoricaceae* with 20 species, *Hookeriaceae* with 6 species and *Sematophyllaceae* with 27 species according to Foreau's list (1961). Only a few species in each of these families could be collected and still fewer have been dealt with in this study.

Apart from their growing in the hills, *Barbula indica* (Fig. 488) and *Leucobryum neilgiriense* (Fig. 485) are growing on the plains very luxuriantly.

The Dry and Wet Bulb Thermometer was used to record the range of temperature in the field conditions. Daubenmire (1977) advocates the use of this kind of thermometers, for obtaining the temperature as it is easy to work out the atmospheric pressure and the relative humidity of the locality with these readings.

The readings were noted on the high mountainous localities where the mosses are growing well, every half-hour of the day between 10 A.M. to 3 P.M. in two different seasons just to gain an idea of the climatic conditions of the moss area. The readings are tabulated (Table-XIII).

TABLE-XIII

Name of the mosses	Locality	Weather condition								
		October				May				
		Altitude in metres	Thermometer reading	Nature of weather						
1	2	3	4	5	6	7	8	9		
<i>Bryum argenteum</i>	Nilgiris									
<i>Petrobryopsis frondosa</i>	Thodda-bedda peak	2900	60-64	52-58			70-76	70-74		Bright sun-shine
<i>Rhodobryum giganteum</i>										
<i>Pleuropus nilgiriense</i>	Kodi									
<i>Barbella tenax</i>	Silver cascade	2000	54-58	52-54	Clouded	70-72	68-70	Bright		

TABLE XIII (Contd.)

	1	2	3	4	5	6	7	8	9
<i>Pogonatum</i>									
<i>neesii</i>		Nilgiris							
<i>Mnium</i>									
<i>longirostrum</i>		Fern hill	2500	56-62	52-56	Misty	70-78	68-72	Clouded
<i>Homaliodendron</i>		Kodai							Moist
<i>flabellatum</i>									windy
<i>Entodon</i>									
<i>plicatus</i>		Priest's							
<i>popillaria</i>		walk							
<i>crocea</i>			2300	60-62	56-58	Drizzling	70-74	72-74	
<i>Barbula indica</i>		Annamalai-	10	80-86	70-78	Bright day	84-90	80-88	Dry
		nagar							

There are variations, as may be seen, at different altitudes at different places during the same hours of the day. The change in temperature was also due to the wind which blows rather violently often carrying clouds with them specially at Doddabetta, the highest peak in South India.

When the temperature is at its maximum (104°F.) the local moss *Barbula indica* is almost dead or dormant waiting for the onset of rains or sprouting out when water is supplied artificially.

### **Humidity**

To measure the humidity, the psychometric tables (Weather Bureau U.S. Commerce Department 1941) were employed as indicated by Daubenmire (1947). There is difficulty in the calculation of humidity as it varies differently in different hours of the day and night and also at different seasons.

Wind profoundly affects the vegetation of mosses. The forest trees like *Lasianthus venulosus* Wt. are clothed with many moss plants (Frontispiece). Richards (1932) has observed that the "effects are particularly obvious on epiphytes." There are also a few species growing on the top of high mountains exposed to the wind.

Pure formations of the local moss *Barbula indica* are found on the terrace of the botanical laboratory during winter season of November and December. The relative humidity during the day varies from 65 to 90. It is at this period that this moss produces capsules. This observation is in conformity with the findings

of Schimper (1903) "that high atmospheric humidity promotes growth and depresses transpiration" Though this plant is abundant both in the plains and found in the hills, it has never been observed to produce capsules in the hills for several years.

### Light

Richards (1932) has observed that "the study of light factor presents special difficulties. The usual methods for measuring its intensity are admittedly unsatisfactory and its effects are very hard to disentangle from those of humidity and temperature" Many moss communities such as *Bryum argenteum*, *Brachymenium exile*, *Pogonatum microstomum*, *Schlotheimia grevilleana*, *Barbula indica* and others are exposed to rather high light intensities. *Ditrichum amoenum* and *Homaliodendron flabellatum* were found to grow away from sunshine inside the caves. The following mosses such as *Rhizogonium spiniforme* and *Rhodobryum giganteum* grow in shady habitats.

### Water

Apart from the intermittent or periodical showers of rain, water keeps oozing from the substratum so well that the mosses growing on the slopes receive a continual supply.

The atmospheric moisture is another source of water for the mosses in the hills. The epiphytic mosses such as *Floribundaria*, *Meteoriopsis* and *Papillaria* which are exposed to dessication "are able to seize the momentary opportunity and in their dry condition can instantly absorb moisture over their whole surface, e.g. algae,

mosses” Warming (1909). While speaking about the muscoid and lichenoid growth forms, he says, “their powers of enduring extreme loss of water and of rapidly replacing this by means of absorption over the whole free surface are ecologically very important ”

### Edaphic factor

The soil is of various kinds in this area, such as rocky, black loamy and humus. The following mosses are growing on tree trunks forming a carpet; *Sematophyllum subhumile*, *Pterobryopsis orientalis*, *Pleuropus nilghiriense* and *Aerobryopsis longissima*.

The following mosses are hanging from the branches of trees in the form of festoons: *Papillaria crocea* (Frontis piece) *Floribundaria floribunda*, *Barbella pendulla*, *Meteoriopsis reclinata*, *Rhizogonium spiniforme* and *Leucobryum scalare* grow in small patches on dead tree trunks.

The behaviour of growth of *Barbula indica* is interesting. It occurs on the open terraces of old buildings, on brick walls exposed to sun and rain. It is surprising to comprehend the perennation of such a delicate moss in the tropical isolation of Annamalainagar. When an entire patch of germinating moss plants was examined, it was found that there is always a tangle of *Oscillatoria* forming a mat beneath. Such patches do not form at all on the same terrace where there is cement plastering, as no *Cyanophyceae* develops there. The statement of Warming (1909) that algae may form the lower stratum for the formation of mosses is in support of this observation. Chapman (1950) has

pointed out that filamentous Cyanophyceae are capable of retaining moisture. Therefore, the occurrence of *Barbula indica* in such a hot climate may be correlated to the presence of filamentous *Oscillatoria* offering protection to the germinating spores and gemmae of the moss plant. Further, it is but appropriate to quote the temperature tolerance of the same genus of moss and blue-green alga from the hand book of Biological data edited by Spector (1956).

#### Tolerance to Extremes of Heat: Moss

Moss	Temperature °C.
<i>Barbula gracillis</i>	110-115

#### Temperature tolerance Extremes: Algae

Species	Maximum for habitat °C.	Maximum tolerated °C.
<i>Oscillatoria amphibia</i>	50	-
<i>Oscillatoria filiformis</i>	85.2	85.2
<i>Oscillatoria formosa</i>	50	-
<i>Oscillatoria geminata</i>	45	-
<i>Oscillatoria okeni</i>	44	-

However, Desikachary (1959) while referring to thermal algae, has quoted Elenkin (1914) who stated that the Cyanophyceae tolerates a maximum temperature of

85°C. It may be seen, therefore, that *Barbula indica* in the tangle of *Ocillatoria* filaments has been able to be tided over the maximum temperature of 104°F. (40°C) in the tropical isolation of Annamalainagar.

Quadrat studies on this moss were also made. Two sets of 10 c.m. quadrats were isolated each set with a control side by side. One of the quadrats of each set was denuded and observations were taken for about a month. On the third day the *Cyanophyceae* made its appearance in the denuded quadrats and on the 10th day the moss protonema was found to develop and after 20 days they were full of young moss plants. These plants were so small that they were in close aggregation. Actual counting of the 10 cm. quadrat reveals that there are 25,000 to 26,000 moss plants in pure formation.

The following mosses grow on rocky substratum: *Homaliodendron flabellatum*, *Schlotheimia grevilleana*, *Thysanomitrium depallieri*. Of these, *Homaliodendron flabellatum* known as the fan moss grows in a horizontal manner directly from moist rocks in thick forests. *Schlotheimia grevilleana* (Fig. 543) forms a thick and beautiful carpet covering rocks exposed to the sun. The plant is provided with reddish branched rhizoids in large numbers. *Thysanomitrium depallieri* is dark green growing in the sun on soil covered rocky slopes where water keeps oozing.

The two species of *Pogonatum* (Figs. 548 & 549) grow on soil in pure formation specially on the cuttings of the road side at higher altitudes extending several

metres. *Mnium coriaceum*, *Rhodobryum giganteum* (Fig. 540) and *Hypopterygium tenellum* grow on loose soil in damp and shaded forests. *Bryum argenteum* and *Anomobryum subnuditum* grow on the crevices of rocks and culverts as Chasmophytes. *Brachymerium exile* and *Fissidens schimidii* (Fig. 538) grow in black loamy soil in shaded situations. *Funaria hygrometrica* likes to grow in the damp regions where there was a forest fire. This observation has also been made by Watson (1954) and Dunham (1951). Therefore, it may be pointed out that this sort of soil is a kind of plant indicator.

When all these physiographic and edaphic factors are present and while many of the moss plants are producing the capsules, there still remains a mystery that some mosses do not seem to produce capsules at all in spite of their very luxuriant growth. It is yet to be known whether the absence of capsule in such mosses suggests that the gametophytic phase has come to be established once and for all as permanent feature without any prospect of giving rise to the sporophytic phase, or, whether, this is just a temporary means of propagating the gametophytic phase for a prolonged period.

The association of some of the mosses with other plants was studied in the manner indicated by Gams (1932) in order to understand the relative frequency of the species. The observations are tabulated as follows:

TABLE-XIV

	<i>Bryum</i> <i>argenteum</i>	<i>Mnium</i> <i>coniaceum</i>	<i>Papillaria</i> <i>crocea</i>	<i>Thuidium</i> <i>tamariscellum</i>	<i>Meteoriopsis</i> <i>reclinata</i>	<i>Pleuropus</i> <i>nilghiriense</i>	<i>Entodon</i> <i>plicatus</i>
<i>Brachymenium</i> <i>exile</i>	+ +	+					
<i>Rhodobryum</i> <i>giganteum</i>	+	+ +					
<i>Pterobryopsis</i> <i>frondosa</i>			+ +		+		
<i>Trachypus</i> <i>bicolor</i>			+	+ +			
<i>Barbella</i> <i>tenax</i>				+ +	+ +		+ +
<i>Floribundaria</i> <i>floribunda</i>					+	+ +	
<i>Sematophyllum</i> <i>subhumile</i>							+ +

TABLE-XV

Mosses with other bryophyta or lichen  
or on angiosperms

	<i>Cladonia</i>	<i>Pellia</i>	<i>Usnea</i>	<i>Coleus</i>	<i>Lasianthus venulosus</i>	<i>Syzygium arnottianum</i>	<i>Rhododendron nilagiricum</i>	<i>Plectranthes weghtii</i>
<i>Bryum argenteum</i>	++							
<i>Hypopterygium tenellum</i>		++						
<i>Pogonatum microstomum</i>	+			++				
<i>Papillaria crocea</i>			+		++			
<i>Sematophyllum subhumile</i>						++		
<i>Hypnodon perpusillus</i>								++
( <i>Rhegmatodon</i> )		1						
<i>Pterobryopsis frondosa</i>							++	

### **Biotic factor :**

This may include, besides the foregoing, the zoological organisms which take shelter under the moss formations. Some millipedes, *Xenobolus acuticonus* Att. were found under the thick carpet of *Schlotheimia grevilleana*. The profusely branched rhizoids afford suitable accommodation for these creatures. The crevices of the bark of forest trees like *Syzygium arnottianum*, *Lasianthus venulosus*, *Rhododendron nilagiricum* and *Plectranthes wightii* in which some of the epiphytic mosses like *Barbella* and *Rhegmatodon* spread their rhizoids, form natural abode of many insects like *Thyroglutus* sp. The leeches, *Haemadipsa* spp., are found in abundance generally with *Rhodobryum giganteum* especially in the Tiger Shola of Kodaikanal. In fact the leech is indicative of the presence of this moss. Similarly in the tuft of *Thuidium*, another millipede, *Sphaerotherium* sp. is always found.

Further, moss carpets of *Pogonatum microstomum* have been found to provide seed beds for flowering plants like *Coleus ovatus* (Fig. 548). It may probably be due to the water retaining capacity of the mosses as Richards (1932) has pointed out.

The South Indian mosses under this study have been classified ecologically in the scheme of classification suggested by Gams (1932).

#### **1. ADNATA**

Mosses on firm ground are.—

*Bryum*

*Fissidens*

*Pogonatum*

1. ADNATA (Contd.)

*Dicranum*

*Ditrichum*

2. RADICANTIA

Mosses on loose ground are:-

*Rhodobryum*

*Mnium*

*Philonotis*

*Funaria*

3. AMPHINEREDIA

Mosses periodically submerged are:-

*Hookeria*

*Thysanomitrium*

4. EPIPETRIA

Mosses confined to rocks are:-

*Schlotheimia*

*Barbula*

*Pleuropus*

*Homaliodendron*

5. EPIPHYTIA

Mosses on the bark of trees are:-

*Rhizogonium*

*Hypnodon*

*Pterobryopsis*

*Meteoriopsis*

*Papillaria*

*Leucobryum*

*Barbella*

6. HELOPHYTIA

Mosses that are peat-forming on very wet soil are:

*Thysanomitrium*

*Herpetineurum*

7. AMPHIPHYTIA

Mosses with periodically inundated soil are

*Rhodobryum*

*Mnium*

*Hypopterygium*

8. CHASMOPHYTIA

Mosses on mineral soil are:-

*Bryum*

*Barbula*

9. BRYOCHAMAEPHYTIA

Mosses on humus soil are:-

*Thuidium*

*Atrichum*

10. SYMBIOPHYTIA

Mosses with other plants are:-

*Rhegmatodon*

*Sematophyllum*

*Erythrodontium*

The other two groups of Gams (1932) such as *Nereidia* and *Epiphyllia* could not be observed under this collection.

# Discussion

## 1) Morphological and Physiological peculiarities of mosses in relation to their ecology

The morphological details of the 64 species of South Indian mosses are generally in accordance with the findings of the earlier bryologists (Cardot & Varde) (1922), Brotherrus (1899), Dixon & Varde (1927). However, the height of the plant, the length and breadth of the leaves, the length of seta and the dimensions of the capsule are in certain cases, slightly variable. This may be due to regional and / or seasonal changes.

### a) Vegetative adaptation:

The presence of dorsal lamellae in the leaves of *Fissidens* (Fig. 5), the papillose cells in the leaves of *Thuidium tamariscellum* (Fig. 284) and the occurrence of rows (lamellae) of cells as outgrowth on the upper leaf surface of *Pogonatum* (Figs. 349 & 354) have been considered to be of significance for the retention of water. Further, the leaf of *Leucobryum* which is mostly *multicellular* contains chloroplasts only in a few cells and the rest are without chloroplasts. This sort of spongy construction appears to be a device for the retention of water as pointed out by Goebel (1905).

The laminar cells of *Trachypodopsis serrulata* (Figs. 208 & 209), *Barbella tenax* (Fig. 238), *Floribundaria floribunda* (Fig. 242), *Erythrodonium julaceum* (Fig. 304) are very much thick-walled. The thickening of the cell-wall in these species, may be due to their xerophyllous habit. It is probable that the thickening of cell-walls serves for holding water obtained from mist, dew and rain by the process of imbibition. Warming (1909) while referring to epiphytes has stated that "many are able to seize the momentary opportunity and in their dry condition can instantly absorb moisture over their whole surface e.g. Algae, mosses, lichens."

*Bryum argenteum* derives its name from the silvery appearance of its shoots. The upper half of the leaves is responsible for this appearance. Goebel (1905) has stated that "these dead upper parts invest as with a mantle, the bud of the stem and must check the outgo of water."

The apex of leaf in the following species of mosses is hair-like: *Brachymenium exile* (Fig. 130), *Brachymenium leptostomoides* (Fig. 136), *Bryum argenteum* (Fig. 145), *Trachypus humilis* (Fig. 200), *Aerobryopsis longissima* (Fig. 225), *Barbella determessi* (Fig. 233), *Barbella pendula* (Fig. 237), *Warburgiella leptorhynchoides* (Fig. 319), *Hypnum cupressiforme* (Fig. 323), and *Taxiphyllum taxirameum* (Fig. 327). The dense aggregation of the hair point in the younger leaves may be another safeguard against drought.

The rhizoidal felt of the mosses may play an important role in the conservation of moisture. This is more so

specially in the lythophytic mosses such as *Homaliodendron flabellatum* and *Schlotheimia grevilleana*. This observation was made by Warming (1909).

It may be seen from a perusal of the anatomical description of the mosses (pp. 86 - 105) that in most of the species, the axis is provided with a peripheral rind of thick-walled cells. The tubular structure of the axis so constructed is perhaps to afford mechanical strength to the otherwise flimsy and delicate mosses. It is likely that the rind is also but a device to retain the water by the process of imbibition apart from its being a protective layer.

**b) The sporophyte:**

Out of 64 species studied, sporophytes were available in 34 mosses only and they have been examined in some detail. The nature of looping of the seta may have a bearing on the "need for the regulation of spore dispersal." Eight different forms of foot of the sporophyte have been observed in the different species of mosses.

Stomata were found only in 8 species. It may be seen from Tables IV and V that the stomata are characteristic of the species and they resemble those of the angiosperms. L B (Length/Breadth) ratio of the stomata is constant and the stomatal index is specific of the mosses. The number of subsidiary cells of the stomata in mosses is variable being 8 to 13. The epidermal cells of the sporophyte of *Trematodon ceylonensis* are much elongated, resembling those of the sporophyte of *Anthoceros pearsoni*. These observations

are in conformity with Bower (1934) who has compared the photosynthetic system of the Psilophytales with that of the Anthocerotales. He has also quoted Goebel as saying that "there is no doubt that many moss capsules show clearly traces of a reduction of the photosynthetic system of the stomata closely related to it - all this indicates that moss sporangia originally possessed a structure similar to that of *Anthoceros*." Further, Goebel (1905) while speaking about the stomata of the mosses has said that "it is remarkable that in the Bryophyta the formation of stomata repeatedly appears as for example in *Anthoceros* and in the different series of the Musci and they in every way correspond with the stomata of the spermatophyta."

Lorch (1926) discovered protoplasmic links between the cells of the foot and those of seta in some 300 mosses. The foot of the sporophyte is always buried more or less deeply in the tissue of the gametophore. Goebel (1905) has observed that in *Eriopus remotifolius* rhizoids are developed at the point where "the sporogonium sits within the ruffle-like vaginula." He has further recorded that "the only example I know of a sporangium rooting by rhizoids." La Rue (1942) successfully stimulated rhizoidal formation all over the detached sporophytes by wounding and by using wound and growth hormones dissolved in lanolin. In the present investigation the sporophyte of 10 species of mosses were induced to develop rhizoids by hormones and thereafter they could lead independent existence. The idea behind these experiments was to find out whether the moss sporophyte could be induced to live indepen-

dently for some length of time. It is interesting to note that under the influence of hormones and gametophytic extract they do produce rhizoids from the base of the foot and they are able to exist independently without the gametophyte though for a short duration of 36 hours. The only other report of a bryophyte known to lead an independent existence is by Campbell (1925) in the case of *Anthoceros fusiformis*, but so far no sporophyte of a moss is known to behave similarly.

The sporophyte of moss with rhizoids therefore appears to have attained a "Psilophytalean status" in its external morphology. It superficially resembles *Horneophyton* with its columellate sporangium. The lacuna in this correlation is the absence of vascular elements in the sporophyte of moss. It is not claimed however, that the gap between the Bryales and the Psilophytales has been bridged by the independent existence of the moss sporophyte. When it can secure this independent existence of 36 hours by Indole-Butyric Acid it may be possible by better culture methods that the moss sporophyte may be induced to become independent for a longer duration.

It may also be pointed out that this observation is an added support to the view of Scott (1923) who referring to Psilophytales, has said that "a certain affinity with the Bryophyta has also been recognised and it has even been held that the Rhyniaceae should be assigned to this sub-kingdom. The Sphagnum-like structure of the columellate sporangium or sporogonium of *Hornea* and *Sporogonites* may be regarded as

ssupporting the Bryophytic attribution which may them readily be extended to *Rhynia*.”

c) Osmotic concentration:

The osmotic concentration of the cell-sap of the gametophyte and that of the sporophyte of 7\* mosses' has been determined for the first time by de Vries, Plasmolytic method (*Table X*). The experiments were made in Kodaikanal hills during the month of October 1960.

TABLE XVI

Osmotic perssure of sporophyte and gametophyte

Name of moss	Sporophyte	Gametophyte	Difference
	O.P. in atm.	O.P. in atm.	in atm.
1	2	3	4
<i>Barbula indica</i>	17.1	4.0	13.1
<i>Funaria hygrometrica</i>	22.7	4.0	18.7
<i>Anomobryum subnuditum</i>	28.8	12.7	16.1
<i>Bryum argenteum</i>	73.9	28.8	45.1
<i>Thuidium tamariscellum</i>	17.1	7.8	9.3
<i>Pogonatum microstomum</i>	73.9	52.2	21.7
<i>Pogonatum neesii</i>	62.8	43.7	19.1

\* Only 7 species were available with sporophytes at that time.

It may be seen from Table XVI. that the osmotic concentration of the sporophyte of each species is always higher than the osmotic concentration of the corresponding gametophyte. The range of difference of the osmotic pressure between the sporophyte and the gametophyte varies from 9.3 atmospheres (*Thuidium tamariscellum*) to 45.1 atmospheres (*Bryum argenteum*). The osmotic concentration of the sporophyte is very high. It may be due to the fact that the food material has to be transported from the gametophyte to the sporophyte.

d) Ecological factors and their significance

The South Indian mosses grow at an altitude of 1000 metres to 2900 metres in the tropical belt of 9° to 12° north latitude and 76° to 79° east longitude. The atmospheric temperature varies from 33°F to 79°F and annual rainfall is between 100 c.c. to 170 c.c. The moss flora is generally rich on the slopes of steep mountains and in the moist regions of the downs. The mosses therefore enjoy a peculiar limited climate known as the "local climate" as Tansley (1946) would put it. He has stated that "Altitude, slope and exposure determine what is called "local climate," affecting the temperature, rainfall, air moisture and insolation to which a given piece of vegetation is exposed and therefore to a large extent the particular species of plant that form it."

The occurrence of tropical moss families in South India in large number of species may be due to the various climatic conditions. As has been pointed out already (p. 118) it is significant that many species of

temperate climate are growing fairly well in this area at high altitudes.

Low temperature at high altitude in the tropics seems to account for the luxuriant growth of the rich vegetation of mosses in this region as the moss plants are endowed with "powers of assimilating apparently at very low temperatures more readily than can spermatophyta" (Warming, 1909).

Some species of South Indian mosses have been observed to grow in two different "discontinuous areas" (Wolff, 1924) of almost similar altitudes. The following mosses grow nearly at the same altitude of both Kodaikanal hills and Ootacamund hills though they are about 300 miles apart from one another :

<i>Ditrichum amoenum</i>	2300 metres
<i>Meteoriopsis reclinata</i>	2200-2300 metres
<i>Papillaria crocea</i>	2500-2600 metres
<i>Distichophyllum succulentum</i>	2000 metres
<i>Atrichum aculeatum</i>	2500-2600 metres
<i>Pogonatum neesii</i>	2600-2800 metres

## 2) Taxonomical considerations on the basis of anatomical observations

### (a) Structure of the moss axis :

The transverse sections of the axis of the 53 species of South Indian mosses studied here, exhibit various peculiarities. As has been pointed out already (p. 96) the following anatomical features may be noticed :

- (i) The axis of most of the mosses, is provided with a peripheral rind.
- (ii) The axis of a few of them are not provided with this rind.
- iii) The axis may possess a well-defined central strand in the centre which may or may not be differentiated.
- iv) The axis may not possess a central strand but the tissue inside the rind may be differentiated into a pith or cavity or there may be a small cell situated in the centre surrounded by a ring of radiating cells. Sometimes the tissue inside the rind may not be differentiated and it may be made up of uniformly thin or thick-walled cells.

Having these facts in view, a classification of the mosses (53 in number) on the basis of the anatomical structure of the gametophyte, has been erected. This classification is merely an attempt to group the various mosses and it does not aim at phylogeny or evolution.

**b) Systems of classification:**

Natural system of classification was proposed by Hedwig (1801) and Jaeger and Sauerbeck (1870-78) followed it. Later Brotherus (1901) erected a classification which according to Dixon (1932) "does not greatly differ from that of Jaeger in general arrangement" and it has been based on the classification of Fleischer (1906-1908) in most respects. In 1932, Dixon proposed his classification. Phylogenetic arrangement of the true mosses was proposed by Schaffner (1938). He has stated that "Brotherus

(1924-1925) is much better than Dixon in his arrangement of the general progressive sequence of the groups from the lower to the higher evolutionary levels." He has added that, "Archidales are plainly the lowest group and the Polytrichales the highest. The Archidales do not represent reduced or degenerate species but are truly primitive in their characteristics."

The different systems of classification of mosses according to Brotherus (1924), Dixon (1932), Schaffner (1939), Conard (1956) and Watson (1954) represented diagrammatically in Plate LXII. All these systems are based mainly on the characters of the peristome. As the classification of Grout (1924) resembles that of Conard (1956) in the main, it has not been represented here.

It may be seen that Schaffner (1939) and Brotherus (1924) separated off the Polytrichales from the rest of the mosses while others place the Polytrichales in the middle of the scheme. Thus the systematic position of the Polytrichales is highly disputed. Cavers (1911) in his inter-relationships of the higher Bryophyta, has separated the Polytrichales away from the Bryales.

The anatomical observations of this group in the present investigation are only meagre as three species alone were examined. However, these three species of the Polytrichaceae stand out as a peculiarly different type of mosses from the rest. The axis has a well defined central strand unlike the other mosses. The leaves contain many rows of lamellae-a feature not found in other mosses.

# Summary

Studies in the morphology, anatomy, physiology and ecology of 64 different species of more common South Indian mosses have been made in some detail.

## Morphology:

Many of South Indian mosses have been reported to occur in other countries also and they were identified and taxonomically described long ago by Cardot (1911), Dixon (1914), Cardot & Varde (1922), Varde (1922-1925), Hedwig (1864), Dixon & Varde (1927), Fleischer (1906-1908), Brotherus (1899), and others. Some of the species were collected even from South India and described by these authors. A few others have been redescribed and specific names corrected. Excepting these sporadic morphological descriptions, there has hardly been any comprehensive work on the South Indian mosses. The present investigations on South Indian mosses, therefore, have been made for the first time in a detailed manner.

On closer examination of the mosses in detail, it has been found that the morphological observations such as the shape, margin, apex and phyllotaxy of the leaves are in conformity with the original description. The lamellar structure of the leaves, the papillose nature of the laminar cells, the absence of chloroplasts in some cells of certain leaves, the excurrent costa and

acuminate apex of leaves and the rhizoidal felt may be adaptations for retention of water. These observations conform to the observations by Goebel (1905) on mosses of other regions.

Certain observations on the sporophyte such as the following are reported for the first time:

(a) Out of 34 available sporophytes examined, 8 different forms of foot have been recognised so far.

(b) Linear measurements of the capsule and foot reveal that there is corresponding increase in length of the foot *pari-passu* with the increase in length of the capsule. The peristome and the spores have been studied in detail.

(c) The opercula have been classified under four groups conforming to the classification of Dunham (1951).

(d) The L/B (Length/Breadth) ratio of the stomata found on the apophysis of 8 species of mosses appears to be a constant factor. The stomatal index of each one of the mosses also appears to be specific.

Vegetative reproduction has been recorded: by means of bud-like bulbils in 3 species, by means of multicellular gemmae in 3 species, regeneration by the formation of rhizoids from detached or attached leaf or stem in 4 species and by the production of "rhizoprotonemata" in *Barbula indica*.

#### **Anatomy:**

Transverse sections of axes of 53 South Indian species have been described for the first time. This anatomo-

mical observation has led to a classification of these mosses on the basis of their structure. The proposed system of classification though based purely upon anatomical details of the gametophyte, is in conformity with the prevailing systems of classification of Brothier (1924) and Schaffner (1938) in some respects.

#### Physiology:

Physiological observations have exhibited that the sporophyte of the moss can be induced to produce rhizoids by treatment with hormones and thereafter it may be induced to lead an independent existence for a short time. The moss sporophyte, thus with its rhizoids arising from the base of the foot, appears to have attained a "psilophytalean" status in this regard. It may be compared in its external morphology to *Hornophyton* with its columellate sporangium, a comparison at best superficial as the moss sporophyte is not provided with any vascular element.

The osmotic concentration of the sporophyte of 10 species and that of their corresponding gametophytes have been determined for the first time. It has been found that osmotic pressure of the sporophytes is always higher than that of the gametophytes. This may be explained on the basis of transport of food from gametophyte to the sporophyte.

#### Ecology:

The natural home of the South Indian mosses is situated within the square of 9° - 12° North Latitude and 76° - 79° East Longitude (Map elsewhere). The range of temperature is between 33°F and 79°F and the annual rainfall between 100 cc. and 170 c.c.

The soil of occurrence of *Funaria hygrometrica* is a plant indicator. It has been found that it grows in regions where there was a forest-fire. This observation is in support of the findings of Dunham (1951) and Watson (1954).

The perennation of *Barbula indica* in the tropical isolation of Annamalainagar has been found to be due to the presence of *Oscillatoria* sp. forming the substratum for the mosses. Both the genera are capable of tolerating high temperature of 104 F. (40°C). This is in conformity with the report of Spector (1956).

The presence of *Rhodobryum giganteum* is indicative of the presence of a leech *Haemadipsa* sp. in the habitat and conversely the leech is indicative of the presence of this moss.

The association of mosses with other mosses and mosses with other plants have been recorded (pp. 127-128) in respect of South Indian mosses. Further, these are classified into 10 groups in the scheme of classification of Gams (pp. 129-131).

## Acknowledgments

It is a source of immense pleasure to place on record my deep sense of gratitude to professor T. C. N. Singh, M Sc., D. Sc., D.Ph., F.B.S , P.L., Professor and Head of the Department of Botany for suggesting the problem and directing the present investigation. I am further much indebted to him for his helpful criticisms, ideas and guidance and also for offering enormous facilities and continued encouragement throughout the course of this work.

I take this opportunity to express my most heartfelt thanks to Rev. Father G. Foreau, Sacred Heart College, Shembaganur, South India for his generous and continuous help in the collection of the mosses and references.

I acknowledge my indebtedness to Professor E. B. Bartram, of Pennsylvania, to Dr. Hyoji Suzuki, Head of Botanical Institute, Hiroshima University and to Dr. H. Ando, Botanical Institute, Hiroshima University for identification of the species and for their stimulating encouragement.

I am much thankful to Professor P. W. Richards, Cambridge University (England) and to Dr. E. V. Watson, University of Reading (England) for supplying reprints, and offering me encouragement; also to the Director of

Observatory, Kodaikanal and the Curator of Botanical Gardens, Ootacamund for kindly supplying meteorological data of the respective regions.

My sincere thanks are due to Shri Sanjiban Biswas for his excellent photography, to Dr. P. Govindan, M.A., Ph.D., Professor of Zoology, Annamalai University for microphotography and to Dr. V. Sriramalu, M.Sc., Ph.D., Department of Zoology of the Annamalai University for help in taking photomicrographs and to Sri R. Sadasivan of M.Sc., class and to Sri Subharayalu of the B.Sc., class for certain drawings of the herbarium. Finally, I earnestly thank all the members of the Botany Department Research Club of the Annamalai University for their useful discussion and helpful criticisms during the weekly meetings of the club.

# Index

Name of the mosses	Page No.
<i>Aerobryopsis longissiima</i>	26
<i>Anomobryum subnuditum</i>	18
<i>Atrichum aculeatum</i>	37
<i>Barbella determessi</i>	27
„ <i>pendula</i>	27
„ <i>tenax</i>	27
<i>Barbula indica</i>	13
<i>Bartramia madurensis</i>	21
<i>Brachymenium exile</i>	18
„ <i>leptostomoides</i>	18
<i>Brothera leana</i>	11
<i>Bryum argenteum</i>	19
<i>Campylopus nodiflorus</i>	9
<i>Ctenidium lychnites</i>	36
<i>Dicranodontium fragile</i>	11
<i>Distichophyllum succulentum</i>	30
<i>Ditrichum amoenum</i>	8
<i>Erythrodontium julaceum</i>	33
<i>Fissidens schimidii</i>	8
<i>Floribundaria floribunda</i>	28

Name of the mosses	Page No.
<i>Foreauella indica</i>	34
<i>Funaria hygrometrica</i>	16
„ <i>submarginata</i>	17
<i>Herpetineurum toccoae</i>	31
<i>Homaliodendron exiguum</i>	29
„ <i>flabellatum</i>	29
<i>Hookeria acutifolia</i>	30
<i>Hyophila involuta</i>	15
„ <i>validinervis</i>	15
<i>Hypnodon perpusillus</i>	22
<i>Hypnum cupressiforme</i>	35
<i>Hypopterygium tenellum</i>	30
<i>Leucobryum neilgherrense</i>	12
„ <i>scalare</i>	12
<i>Meteoriopsis reclinata</i>	23
„ <i>squarrosa</i>	28
<i>Mnium coriaceum</i>	20
<i>Myurium warburgii</i>	25
<i>Papillaria crocea</i>	26
<i>Philonotis anisoclada</i>	22
<i>Pleuropus nilghiriense</i>	32
<i>Pogonatum microstomum</i>	37
„ <i>neesii</i>	38
<i>Pohlia flexuosa</i>	17
<i>Pterobryopsis orientalis</i>	25

Name of the mosses	Page No.
<i>Rhacomitrium javanicum</i>	16
<i>Rhegmatodon orthostegius</i>	31
<i>Rhizogonum spiniforme</i>	21
<i>Rhodobryum giganteum</i>	20
<i>Rhynchostegium javanicum</i>	33
<i>Schlotheimia grevilleana</i>	23
<i>Sematophyllum subhumile</i>	35
<i>Taxiphyllum taxirameum</i>	36
<i>Thuidium tamariscellum</i>	32
<i>Tortula muralis</i>	13
<i>Trachypodopsis serrulata</i>	24
<i>Trachypus bicolor</i>	24
„ <i>humilis</i>	24
<i>Trematodon ceylonensis</i>	9
<i>Trichostomum cylindricum</i>	14
<i>Thysanomitrium depallieri</i>	10
„ <i>lioneurum</i>	10
<i>Warburgiella leptorrhynchoides</i>	35
<i>Weisia macrospora</i>	14

## Literature

1. ARNOLD, C.A. (1947): An Introduction to Palaeobotany. *McGraw-Hill Book Company, Inc., New York.*
2. BANERJI, T. and SEN, S. (1957): Life history of *Barbula indica*. *Proc. Ind. Sci. Congr. Part III Abstracts*, p. 228.
3. BARTRAM, E.B. (1939): Mosses of the Philipines. *Philip. Journ. Sci.* 68 (1-4): pp. 1-423.
4. ——— (1949): Mosses of Guatemala, *Fieldiana. Bot.* 25, pp. 1-442.
5. BERKELEY, M.J. (1863): Hand book of British Mosses. *Lovell Reeve & Company, London.*
6. BONNER, J. and GALSTON, A.W. (1951): Principles of Plant Physiology. *W. H. Freeman & Company, Sanfrancisco.*
7. \*BRAITHWAITE, R. (1895); The British Moss Flora, pp. 162-163, 207-208 (1888 - 1895).
8. \*BRILLIANT, B. (1927): Les formes de la plasmolyse produites par des solutions con centrees de sucres et de sels dans les cellules de *Mnium* et de *catharinaea*. *C.R. Acad. Sci. U.S.S.R.*, pp. 155-160.

---

\* Original not seen by me.

9. BRITTON, E. (1907): *Rhacopilum tomentosum* (Sw.)  
Brid. *The Bryologist*, Vol. X, pp. 31-33.
10. BROTHERUS, V.F. (1899): Contributions to the  
Bryological Flora of Southern India. *Rec. Bot.  
Sur. India*, Vol. 1, pp. 311-329.
11. \*— — — — (1893-1909): Musci In Engler and Prantl,  
Pflanzenfamilien 1st. Edition.
12. — — — — (1924): Musci, In Engler and Prantl,  
Naturlichen Pflanzenfamilien. Band 10 Leipzig,  
1924, 2nd. Edition and Band 11. *Duncker & Humblot,*  
*Berlin*, 1960.
13. BRÜHL, P. (.931): A census of Indian Mosses. *Rec.  
Bot. Sur. India*, Vol. XIII, pp. 1-135, and Vol.  
XIII, 2. pp. 1 to 152.
14. BOWER, F.O. (1934): Primitive Land Plants. *Hafner  
Publishing Company*, New York, 1959. pp. 70-122.
15. \*CAMPBELL, D.H. (1925): The relationships of the  
Anthocerotaceae. (Flora. Festschr.f.Goebel).
16. — — — — (1930): The Structure and development of  
Mosses and Ferns. *Macmillan Company Ltd.,  
London*.
17. \*CARDOT, M. (1925): *Rev. Bryol.* 52: p. 38.
18. \*— — — — (1927): *Arch. Bot.* 1: p.179.
19. \*— — — — and VARDE, P. de la (1922): *Rev. Bryol.*  
Vol. 49-50.
20. \*— — — — and — — — — (1923): *Arch. Bot.* Vol.1.

---

\* Original not seen by me.

21. CAVERS, F. (1911): Inter-relationships of the Higher Bryophyta. *New Phyt.* Vol. X, pp. 37-46.
22. CHOPRA, R.S. and KUMAR, S.S. (1961): Preliminary study of the Anatomy of Mosses. *Proc. Ind. Sci. Congr. Assn. Abstract*, 51, p.263.
23. ——— and SHARMA, P. D. (1956). Natural regeneration in *Pogonatum palis*. *Jour. Ind. Bot Soc.* Vol. 25, pp. 177-180.
24. CONARD, H. S. (1956): How to know the Mosses and Liverworts. *W.M.C. Brown Company*, Iowa. Revised Edition.
25. \*CORRENS, C. (1899): Untersuchungen über die Vermehrung der Laub-moose. Jena.
26. CRUM, H. (1956): Notes on *Hypnodon*, A Genus of Orthotrichaceae New to North America. *The Bryologist* : Vol. 59, pp.26-34.
27. DAUBENMIRE, R.F. (1947): Plants and Environment. *John Wiley & Sons. Inc. New York.* p.87.
28. DEB, D. B. (1955): Mosses of Manipur, Assam. *Bull. Bot. Soc. Bengal.* Vol. 9 : 2.
29. DESIKACHARY, T. V. (1959): Cyanophyta. *Ind. Council. Agri. Research, New Delhi.* pp. 51-52.
30. DIXON, H.N. and JAMESON, H.G. (1904) : Students Hand Book of British Mosses. Second Edition. *V.T. Sumfield, London.*
31. \*DIXON, H.N. (1909); Mosses from the Western Ghats. *Jour. of Bot.* Vol. 47, pp. 157-164.

---

\* Original not seen by me.

32. ——— (1914): Report on the mosses collected by Mr. C.E.C. Fisher and others from South India and Ceylon. *Rec. Bot. Surv. Ind. Vol. VI* (3): pp. 55-89.
33. ——— (1923): On a collection of mosses from the Kanara District. *Jour. Ind. Bot. Vol.2*, pp. 174-188.
34. ——— (1932): Classification of Mosses in Verdoorn's Manual of Bryology. *The Hague Martinus Nijhoff*. pp. 397-412.
35. \* ——— and VARDE, P. de la.R. 1927 : Contribution a la Flore bryologique de l'Inde Meridionale. *Arch. Bot.* 1 : pp. 161-184.
36. DUNHAM, E.M. (1951) : How to know the Mosses 2nd. Edition Mosher. Press, Mass.
37. \*ELENKIN, A.A. (1936): Monographic algarum cyanophycearum aquidutcium et terrestrium in finibus URSS inventarum, Vol. 1. pp. 675. *Acad. Nank. Moscow*. In Desikachari : Cyanophyta.
38. ERDTMAN, G. (1952) ; Pollen Morphology and Plant Taxonomy-Angiosperms. 2nd. Edition. *The Chronica Botanica Company, U. S. A.*
39. \*FLEISCHER, M (1906 -1908) : Flore de Buitenzorg ve partie, Les Muscinees Leiden.
40. FOREAU, G. Rev. (1930) : Notes on Bryological Geography for the Presidency of Madras. *Jour. of Mad. Univ.* 1930.

---

\* Original not seen by me,

1. FOREAU, G. Rev. (1961) : The Moss Flora of the Palni Hills. *Jour. of Bombay Nat. Hist. Soc.* Vol. 58 : pp. 1-47.
42. FRY, E. (1908) : British mosses. 2nd. Edition. *Witherby & Company, London.*
43. FRYE, T. C. and FERGUSON, E. (1943) : *Pogonatum liebmannianum*. *The Bryologist* : Vol. 46, pp. 141-146.
44. GAMS, H. (1932) : Bryo-cenology (Moss-Societies). In Verdoorn-Manual of Bryology. *Martinus Nijhoff, The Hague*, pp. 323-366.
45. GARJEANNE. A.J.M. (1932) : Physiology. In Verdoorn-Manual of Bryology. *Martinus Nijhoff, The Hague*, pp. 207-232.
46. GOEBEL, K. (1905) : Organography of Plants. *Authorised English Edition, Part II, Clarendon Press, Oxford*, pp. 1-167.
47. GROUT, A. J. (1908) : Some relations between the habitats of mosses and their structure. *The Bryologist* : Vol. VII, pp. 97-100.
48. ——— (1913) : Inter-relationships of the Bryophyta (Review). *The Bryologist* Vol.16, pp. 9-11.
49. ——— (1924, 1947) : Mosses with a hand-lens 3rd. & 4th. Editions. *Intelligence Printing Company. Lancaster, Pa.*
50. ——— (1928) : Classification of the Musci. *The Bryologist*, Vol. 31, pp. 56-61.

51. ——— (1924-1940) : Moss Flora of North America, Vols. 1-11, Newfare, Vermont.
52. \*HABERLANDT, G. (1886) : Beitr. Z. Anatomie u. Physiol.d. Laubmoose. *Jahrb. j. Wiss. Bot.*
53. \*—————(1896) : Physiologische Pflanzenanatomie Ed. 2. Leipzig.
54. ——— (1928) : Physiological Plant Anatomy, Translated from the fourth German Edition. *Macmillan Company London.*
55. \*HEDWIG, J. (1787-97) : Descripto et adumbratio microscopico-analytica muscorum frondosorum, Leipzig.
56. \*HEDWIG (1801) : Species Muscorum. In Dixon's classification in Verdoorn's Manual of Bryology. *Martinus Nijhoff. The Hague, p.397.*
57. \*JAEGER and SAUERBECK (1870-78) : Adumbratio floriae muscorum. In Dixon's classification in Verdoorn's Manual of Bryology. *Martinus Nijhoff: The Hague, p. 397.*
58. JENNINGS, O.E. (1951) : The American Midland Naturalist. Monograph No. 6. A manual of the mosses of Western Pennsylvania and adjacent regions 2nd. Edition. *Univ- of Notre Dame Press.*
59. \*JENSEN, C. (1939) : Skandinaviens Bladmossflora (Mosses of Scandinavia) *Ejnar Munksgaard, Copenhagen.*

---

\* Original not seen by me.

60. KACHROO P. (1954) : Spore Germination and Regeneration in *Physcomitrium pyriforme*. *Jour. Ind. Bot. Soc. Vol. 33*, pp. 262-267.
61. KRUSENSTJERNA, E. Von (1945) : Bladmossvegetation och Bladmossflora i UppsalaTrakten (Moss flora and Moss vegetation in the neighbourhood of uppsala). *Acta Phytogeog. Suecica (Svenska Vaxtgeogr. Sallsk.) 19* : 1-250 (With a short English summary).
62. LA RUE, C.D. (1942) : The Effect of Wounding of Wound Hormones and of Growth Hormones on Rhizoid Formation in Mosses and Liverworts. *The Bryologist : Vol.45*, pp.35-39.
63. \*LORCH, W. (1894) : Beiträge Zur Anatomie und Biologie der Laubmoose. Flora In R. Van Der Wijk - Morphologie und Anatomie Der Musci in Verdoorn's Manual of Bryology. *Martinus NijHoff. The Hague*, pp. 21.
64. \*LORCH, U.D. (1925) : Haustorialschläuche am Fusze d. Laubmoose. (Ber.d. deutschen bot. Gesellsch. 43.1925). In Verdoorn's Manual of Bryology. *Martinus NijHoff. The Hague* p.216.
65. \*———(1925)a : Saugzellen im Fusze etc. bei den Laubmoosen. Ibid. In Verdoorn's Manual of Bryology. *Martinus. The Hague*, p. 216.
66. LORCH, W. (1931) : Handbuch der Pflanzenanatomie. Band VII/1 Anatomie der Laubmoose. Verlag von Gebrüder, Berlin.

---

\* Original not seen by me.

67. MAC CLYMONT, J.W. (1955) : Spore studies in the Musci with special reference to the genus *Bruchia*. *The Bryologist* : Vol- 58, pp.287-3ii.
68. MEYER, S.L. (1940) : Physiological studies on Mosses I. The development of leafy gametophyte in liquid media. *Amer. Jour. Bot.* 27. pp. 221-225.
- 69 \*——— (1954) : Fifteen years of research in moss physiology : 1938-1953. *VIII<sup>o</sup> Gongr. Inter. Bot. Rapp. et Conn. Sect. 16* : pp. 125-126.
- 70 MEYER, B.S. ANDERSON, D.B., and SWANSON C.A. (1955) : *Laboratoy Plant Physiology*. Third Edition. D. Van Nostrand Company Inc. New York.
- 71 MILLER, H.A. (1954) : Field Observations on Associations of Hawaiian Mosses. *The Bryologist*: Vol. 7 : 2. pp. 167-172.
- 72 \*MITTEN, W. (1859) : Musci Indiae Orientalis : an enumeration of the mosses of the East Indies (1858). *Linn. Soc. Journ.* III, 1859 (Bot.) Suppl.
- 73 NARAYANASWAMY, S. and LAL, M. (1957) : Regeneration in some Indian mosses. *Phytomorphology*, Vol. 7 : 2, pp. 244-252.
- 74 \*OLTMANN, (1884) : Ü. d. Wasserbew in der Mossflanze Inaug. In Verdoorn's Manual of Bryology. *Martinus Nijhoff. The Hague*, p.384.
- 75\* OLSEN, C. (1923) : C. R. Lab. Calsberg, 15, 1923, pp. 107-113. In Verdoorn's Manual of Bryology. *Martinus Nijhoff. The Hague*, p. 384.

---

\* Original not seen by me

76. PASCHR, A. (1931): Die Susswasser Flora Mitteleuropas. Haft 14: Bryophyta. *Verlag Von Gustav Fischer Jena*.
77. PATTERSON, P.M. (1957): The Effect of Indole-3 Acetic acid on certain growth phases in Bryophytes. *The Bryologist: Vol. 60*. pp.277-283.
78. PRIOR, P.V. (1952): The Effect of 2,4-D on two Mosses. *The Bryologist, Vol.55*, pp. 48-50.
79. RANGASWAMI, K. (1959): Studies in the Cyto-Morphology and Cyto-taxonomy of South Indian Cucurbitaceae. *Ph. D. Thesis. Ann. Uni.* 1972.
80. RUHLAND, W. (1924): Musci. In Engler and Prantl Pflanzen-familien, Band 10, Verlag, Von Wilhelm Engelmann, Leipzig. 2nd. Edition, pp. 1-100.
81. RICHARDS, P.W. (1932): Ecology. In Verdoorn's Manual of Bryology. *Martinus Nijhoff. The Hague*, pp. 367-396.
82. ———(1950); A book of mosses. Penguin Books, London.
83. \*SAINSBURY, G.O.K. (1935): Vegetative Reproduction in New-zealand Mosses. *Journ. and Proc. Royal Soc. of New South Wales. Vol.69*: pp. 86-105.
84. ———(1938): Vegetative Reproduction in New Zealand Mosses II. *The Bryologist: Vol. 41*, pp. 11-18.

---

\* Original not seen by me

85. ——— (1952) : Ibid. Vol. 55, pp. 71-79.
86. SCHAFFNER. J.H. (1938) The Natural Orders of the True Mosses. *The Bryologist: Vol 41*, pp. 57-63.
87. SCHIMPER, A.F W. (1903) : Plant Geography upon a physio-logical Basis. Authorised English Translation-Revised, *Clarendon Press*.
88. SCHIMPER (1836-1855) : Bryologia Europaea. In McClymont's Spore studies in the Musci *The Bryologist: Vol.58.*, p.287.
89. SCOTT, D.H. (1923) : Studies in Fossil Botany, Vol.II. Third Edition, *A & C. Black Ltd., London* pp. 386-398.
90. \*SEDGEWICK (1932?) : Mosses of the Bombay Presidency. *Jour. of Botany, Vols. 47-50*.
91. SEWARD, A.C. (1931) : Plant Life Through the Ages. Reprint of 2nd. Edition. *Hafner Publishing Company, New York, 1959*.
92. SINGH, T. C. N. (1930) : Notes on vegetative reproduction in two mosses from Mussoorie. *New Phyt. Vol.29*, pp. 355-360.
93. SMITH, G.M. (1955) : Cryptogamic Botany Vol.II Bryo-phytes and Pteridophytes. *McGraw Hill Book Company, Inc. New York, 1955*.
94. SPECTOR, W.S. (1956) : Hand book of Biological Data. *W.B. Saunders Company, U.S.A.*, p.448.

---

\*. Original not seen by me

95. SRINIVASAN, C. (1958), On the Ecological Survey of the South Indian Mosses. *Proc. Ind. Sci. Congr. Assn. (1958)*, Abstract No.33, pp. 283-284.
96. ——— (1960,a): Vegetative Reproduction and Regeneration in South Indian Mosses I. *Ibid (1960)*. Abstract No. 60, pp. 338-339.
97. ——— (1960,b): On the Anatomy of the axis of the South Indian Mosses. *Ibid (1960)*. Abstract No.61, p. 339.
98. ——— (1961): Stomata on the Moss capsule. *Ibid (1961)*. Abstract No.50, pp.262.263.
99. STEERE, W.C. (1950): Reviews. *The Bryologist*, Vol.53, pp.77-79.
100. ——— (1952): Notices of some Recent Bryological Floras and Manuals. *Ibid (1952)*, Vol.52, pp.228-238
101. \*STELMACH (1926): Die Regulation d. Substratizidität durch zwei Torfmoose. *Bul. Acad. Pol.* 1926. In Verdoorn's Manual of Bryology. *Martinus Nijhoff. The Hague*, p.219.
102. STOCKING, C.R. (1956): Osmotic pressure or Osmotic value. In Ruhland's Encyclopedia of Plant Physiology Vol. II. General Physiology of the plant cell. *Springer-verlag Berlin*, 1956, pp.57-70.
103. TANSLEY, A. G.: (1946): Introduction to Plant Ecology, 2nd. Edition. Reprinted, 1949, *George Allen & Unwin Ltd., London* p.156.

---

\* Original not seen by me

104. Tansley and CHICK, E. (1901): Notes on the conducting Tissue-system in Bryophyta. *Annals of Bot.* Vol. XV, pp.1-38.
105. TREASE, G.E. (1945): A Text Book of Pharmacognancy, 4th. Edition.
106. URSPRUNG, A and BLUM, G. (1916): Osmotic pressures of Volume Molar Sucrose Solutions at 20 C. *Ber. Deutsch. Bot. Ges.* 34 : 525-554. In Meyer, Anderson & Swanson - Laboratory Plant Physiology Third Edition. *D van Nostrand Company; Inc. New York*, 1955, p.34.
107. \*VARDE POTIER DE LA, R. (1922): Musci Madurenses India Meridionalis exsiccati. *Revue Bryologique* 49 : 33-44, 50 : 17-27, 72-79 : 51 : 10-14. Cahan, 1922-1924.
108. \*——— (1925). Nouvelles herborisations dans le Sud le l'Inde. *Ibid.* (1925), 52 : 37-43.
109. \*——— (1928,a) : *Musc novi Indici.* *Ann. Crypt. Exot.* 1 : 37-47.
110. \*——— (1928,b) : Mousses nouvelles de l'Inde meridionale. *Ibid.* 1 : 279-283.
111. VAN DER WIJK, R (1932); Morphologie und Anatomie Der Musci. In Verdoorn' Manual of Bryology. *Martinus Nijhoff. The Hague*, pp. 1-40.
112. VERDOORN, FR. (1932): Manual of Bryology (Edited). *Martinus Nijhoff. The Hague*, 1932.

---

\* Original not seen by me.

113. WARMING, EUG. (1909): *Oecology of Plants*, 2nd. Impression, 1925. *Oxford Uni.Press*.
114. WATSON, E.V. (1954) : *British Mosses and Liverworts*. *Cambridge Uni. Press*. 1955.
115. WATSON, W. (1909): *The Distribution of Bryophytes in the Woodlands of Somerset*, *New Phytologist*, Vol. VIII, pp. 90-96. Reproduced *W.M. Dawson & Sons Ltd, London*, 1955.
116. WOLFF, E.V. (1950): *An Introduction to Historical Plant Geography*. Authorised English Translation, *Chronica Botanica Company, Mass, U.S.A.*, p.80.

# PLATES

## PLATE I

### *Fissidens schimidii* C. M.

- Figure 1. Showing plant with leaves and sporophyte. X 1  
Figure 2. Showing plant with leaves and sporophyte. X 10  
Figure 3. Leaf showing the characteristic appendage. X 75  
Figure 4. Cells of leaf. X 285  
Figure 5. Transverse section of leaf showing the dorsal lamella the appendage as outgrowth. X 285  
Figure 6. Peristome. X 258


### *Ditrichum amoenum* (Thw. and Mitt.) Par.

- Figure 7. Showing plant with leaf and sporophyte. X 1  
Figure 8. Showing plant with leaves of gametophyte. X 10  
Figure 9. Leaf showing shape. X 20  
Figure 10. Cells of leaf. X 285  
Figure 11. Capsule showing shape. X 20  
Figure 12. Peristome. X 285

### *Trematodon ceylonensis* C. M.

- Figure 13. Showing plant with leaf and sporophyte. X 1  
Figure 14. Showing plant with leaf and sporophyte. X 10  
Figure 15. Leaf showing shape and apex. X 75  
Figure 16. Cells of leaf. X 285  
Figure 17. Capsule showing shape. X 20  
Figure 18. Stomata on the wall of capsule near apophysis. X 490  
Figure 19. Peristome. X 285

PLATE I



## PLATE II

### *Campylopus nodiflorus* (C. M.) Jaeg.

- Figure 20. Showing plant with leaves and sporophyte X 2  
Figure 21. Leaf showing shape. X 20  
Figure 22. Leaf showing apex. X 285  
Figure 23. Cells of leaf in the middle of lamina. X 285  
Figure 24. Alar cells. X 285  
Figure 25. Capsule showing shape. X 20  
Figure 26. Annulus. X 285  
Figure 27. Peristome. X 285

### *Thysanomitrium depallieri* Card.

- Figure 28. Showing plant with leaves. X 2  
Figure 29. Leaf showing shape and margin. X 20

### *Thysanomitrium lioneurum* (Ther. and Vard.) Broth.

- Figure 30. Leaf showing apex. X 285  
Figure 31. Alar cells of leaf. X 490  
Figure 32. Cells of leaf. X 75  
Figure 33. Showing plant with leaves. X 2  
Figure 34. Leaf showing shape. X 15

### *Thysanomitrium depallieri* Card.

- Figure 35. Showing apex of leaf. X 285

### *Thysanomitrium lioneurum* (Ther. and Vard.) Broth.

- Figure 36. Cells of leaf. X 75  
Figure 37. Cells of leaf. X 285


### *Brothera leana* (Sull.) C. M.

- Figure 38. Showing plant with leaf. X 2  
Figure 39. Leaf showing shape. X 30  
Figure 40. Leaf showing apex. X 285  
Figure 41. Cells of leaf. X 285

PLATE II



20



22




23



25



26



24



27




28



29




31



32



33



34



35



36




38



39



36



37



39

### PLATE III

*Dcranodontium fragile* (Hook.) Broth.

- Figure 42. Showing plant with leaf and sporophyte.  
X 2
- Figure 43. Leaf showing shape. X 75
- Figure 44. Leaf showing apex. X 285
- Figure 45. Alar cells of leaf. X 285
- Figure 46. Showing form of capsule. X 20
- Figure 47. Peristome. X 285


*Leucobryum neilgherrense* C. M.

- Figure 48. Showing plant with leaves and sporophyte.  
X 2
- Figure 49. Leaf showing shape. X 40
- Figure 50. Leaf showing apex. X 285
- Figure 51. Transverse section of leaf. X 75
- Figure 52. Cells of leaf. X 285
- Figure 53. Capsule showing shape. X 20
- Figure 54. Peristome. X 285.

*Leucobryum scalare* C. M

- Figure 55. Showing plant with leaves. X 2
- Figure 56. Leaf showing shape. X 40
- Figure 57. Leaf showing apex. X 285
- Figure 58. Transverse section of leaf. X 75
- Figure 59. Cells of leaf. X 285

PLATE III



## PLATE IV

### *Tortula muralis* Hedw.

- Figure 60. Showing plant with leaves and sporophyte.  
X 1
- Figure 61. Showing plant with leaves and sporophyte.  
X 10
- Figure 62. Leaf showing shape. X 20
- Figure 63. Leaf showing apex. X 285
- Figure 64. Cells of leaf. X 285
- Figure 65. Capsule showing shape. X 20

### *Funaria hygrometrica* Hedw.

- Figure 66. Annulus. X 285


### *Tortula muralis* Hedw.

- Figure 67. Peristome. X 285 (Base and tip only shown)

### *Barbula indica* Schw.

- Figure 68. Showing plant with leaves and sporophyte.  
X 2
- Figure 69. Showing plant with leaves and sporophyte.  
X 10
- Figure 70. Leaf showing shape. X 40
- Figure 71. Leaf showing apex. X 285
- Figure 72. Cells of leaf. X 285
- Figure 73. Gemmae detached. X 490
- Figure 74. Capsule showing shape. X 20
- Figure 75. Stoma with two guard cells. X 285
- Figure 76. Guard cells of stoma divided by cross wall.  
X 285
- Figure 77. Peristome. X 40
- Figure 78. Peristome. X 60

PLATE IV



## PLATE V

### *Trichostomum cylindricum* (Bruch.) C.M.

- Figure 79. Showing plant with leaves and sporophyte.  
X 2
- Figure 80. Showing plant with leaves and sporophyte.  
X 10
- Figure 81. Leaf showing shape. X 40
- Figure 82. Leaf showing apex. X 285
- Figure 83. Cells of leaf. X 285
- Figure 84. Capsule showing shape. X 20
- Figure 85. With annulus. X 285


### *Weisia macrospora* Card.

- Figure 86. Showing plant with leaves and sporophyte.
- Figure 87. Showing plant with leaves and sporophyte.
- Figure 88. Leaf showing shape. X 40
- Figure 89. Leaf showing apex. X 285
- Figure 90. Cells of leaf. X 285
- Figure 91. Capsule showing shape. X 20

### *Hyophila involuta* (Hook.) Jaeg.

- Figure 92. Showing plant with leaves and sporophyte.  
X 2
- Figure 93. Showing plant with leaves and sporophyte.  
X 5
- Figure 94. Leaf showing shape. X 40
- Figure 95. Capsule showing shape X 20
- Figure 96. Annulus in two rows. X 285
- Figure 97. Leaf showing apex X 285
- Figure 98. Cells of leaf. X 200

PLATE V



## PLATE VI

### *Hyophila valtinervis* Card. and Vard.

- Figure 99. Showing plant with leaves. X 2  
Figure 100. Showing plant with leaves. X 10  
Figure 101. Leaf showing shape. X 40  
Figure 102. Leaf showing apex. X 285  
Figure 103. Cells of leaf. X 40

### *Rhacomitrium javanicum* Bryol.

- Figure 104. Showing plant with leaves. X 2  
Figure 105. Leaf showing apex. X 7  
Figure 106. Leaf showing apex. X 285  
Figure 107. Cells of leaf. X 285

### *Funaria hygrometrica* Hedw.

- Figure 108. Showing plant with leaves and sporophytes  
X 1  
Figure 109. Leaf showing shape and margin. X 75  
Figure 110. Leaf showing apex. X 75  
Figure 111. Capsule showing apex. X 10


### *Tortula muralis* Hedw.

- Figure 112. Showing peristomial teeth twisted. X 128

### *Funaria hygrometrica* Hedw.

- Figure 113. Peristome. X 75  
Figure 114. Peristome. X 285

PLATE VI



## PLATE VII

*Funaria submarginata* Card. and Vard.

Figure 115. Showing plant with leaves and sporophyte.  
X 2

Figure 116. Showing plant with leaves and sporophyte.  
X 10

Figure 117. Leaf showing shape. X 20

Figure 118. Cells of leaf.. X 285

Figure 119. Capsule showing shape. X 20

Figure 120. Part of upper portion of capsule without  
peristome X 75

*Pohlia flexuosa* (Mitt.) Broth.

Figure 122. Showing plant with leaves and sporophyte.  
X 2

Figure 123. Showing plant with leaves and sporophyte.  
X 10




Figure 124. Leaf showing shape. X 40

Figure 125. Leaf showing apex. 285

Figure 126. Cells of leaf. X 285

Figure 127. Outer and inner peristome. X 285

PLATE VII



## PLATE VIII

*Brachymenium exile* (Doz. and Molk.) Bryol.

Figure 128. Showing plant with leaves and sporophyte  
X 2

Figure 129. Showing plant with leaves and sporophyte.  
X 10

Figure 130. Leaf showing shape X 40

Figure 131. Cells of leaf. X 285

Figure 132. Peristome. X 128

*Brachymenium leptostomoides* (C. M.) Shimp.

Figure 133. Showing plant with leaves. X 2

Figure 134. Showing plant with leaves. X 10

Figure 135. Leaf showing shape. X 75

Figure 136. Leaf showing apex X 285

Figure 137. Cells of leaf. 285

*Anomobryum subneditum* Card. and Vard.

Figure 138. Showing plant with leaves and sporophyte.  
X 2

Figure 139. Showing plant with leaves and sporophyte.  
X 10

Figure 140. Leaf showing shape. X 75

Figure 141. Peristome. X 285

Figure 142. Cells of leaf. X 285

*Bryum argenteum* Linn.

Figure 143. Showing plant with leaves and sporophyte.  
X 2


Figure 144. Showing plant with leaves and sporophyte.  
X 10

Figure 145. Leaf showing shape. X 75

Figure 146. Cells of leaf. X 285

Figure 147. Peristome. X 285

PLATE VIII



## PLATE IX

*Rhodobryum giganteum* (Hook.) Par.

Figure 148: Showing plant with leaves and sporophyte,  
X 2

Figure 149. Leaf showing shape. X 10

Figure 150. Leaf showing apex. X 75

Figure 151. Capsule showing shape. X 5

Figure 152. Cells of leaf. X 285

Figure 153. Peristome X 285

*Mnium coriaceum* Griff.

Figure 154. Showing plant with leaves and sporophyte  
X 2

Figure 155. Leaf showing shape. X 20

Figure 156. Leaf showing apex. X 75

Figure 157. Cells of leaf. X 285

Figure 158. Capsule shape. X 20

Figure 159. Peristome. X 285

PLATE IX




48



49




152



153



154




150




158



155



156



157



159

## PLATE X

### *Rhizogonium spiniforme* (Hedw.) Bruch.

Figure 160. Showing plant with leaves and sporophyte.  
X 2

Figure 161. Leaf showing shape. X 20

Figure 162. Leaf showing apex. X 285

Figure 163. Cells of leaf. X 285

Figure 164. Capsule shape. X 20

Figure 165. Peristome. X 75

Figure 166. Inner peristome. X 75

### *Bartramia madurensis* Dix. and Vard.

Figure 167. Showing plant with leaves. X 2

Figure 168. Leaf showing shape. X 40

Figure 169. Leaf showing apex. X 285

Figure 170. Cells of leaf. X 285

### *Philonotis anisoclada* Card. and Vard.

Figure 171. Showing plant with leaves and sporophyte.  
X 2

Figure 172. Leaf showing shape. X 75

Figure 173. Leaf showing apex. X 285


Figure 174. Cells of leaf. X 285

Figure 175. Cells of leaf. X 750 (Note cellenchymatous thickening)

Figure 176. Capsule showing shape. X 20

Figure 177. Peristome. X 75

PLATE X



## PLATE XI

*Hypnodon perpusillus* (Thw. and Mitt.) C.M.

Figure 178. Showing plant with leaves and sporophyte.  
X 2

Figure 179. Showing plant with leaves and sporophyte.  
X 25

Figure 180. Leaf showing shape. X 40

Figure 181. Cells of leaf. X 285

Figure 182. Leaf showing apex. X 75

Figure 183. Peristome. X 285

*Schlotheimia grevilleana* Mitt.

Figure 184. Showing plant with leaves and sporophyte.  
X 2

Figure 185. Showing plant with leaves and sporophyte.  
X 5

Figure 186. Leaf showing shape. X 75

Figure 187. Leaf showing apex. 75

Figure 188. Alar cells of leaf. X 285


Figure 189. Cells of leaf. X 285

Figure 190. Cells of upper leaf. X 285

Figure 191. Capsule showing shape. X 20

Figure 192. Peristome. X 285

PLATE XI



## PLATE XII

### *Trachypus bicolor* Rein. and Horns.

Figure 193. Showing plant with leaves. X 2

Figure 194. Leaf showing shape. X 75

Figure 195. Leaf showing apex. X 285

Figure 196. Cells of leaf. X 285

Figure 197. Cells of leaf. X 750.

### *Trachypus humilis* Lind. var *humilis* Zanten.

Figure 198. Showing plant with leaves. X 2

Figure 199. Leaf showing shape, without costa. X 75

Figure 200. Leaf showing apex. X 285

Figure 201. Cells of leaf. X 750

Figure 202. Alar cells of leaf. X 285

Figure 203. Cells of upper leaf. X 285

### *Trachypodopsis serrulata* (Beauv.) Fleis. var. *crispatula* (Hook.) Zanten.

Figure 204. Showing plant with leaves. X 2

Figure 205. Leaf showing apex. X 30

Figure 206. Leaf showing apex. X 285

Figure 207. Cells of leaf. X 285

Figure 208. Cells of leaf. X 750

Figure 209. Alar cells of leaf. X 750


PLATE XII



193



194



195




196



---a




198



200



---c 201




199



200




202



203



204



205 ---b



206



207



208

## PLATE XIII

### *Myurium warburgii* (C. M.) Fleis.

- Figure 210. Showing plant with leaves. X 2  
Figure 211. Leaf showing shape. X 30  
Figure 212. Leaf showing apex. X 285  
Figure 213. Cells of upper leaf. X 285  
Figure 214. Cells of lower leaf. X 285

### *Pterobryopsis orientalis* (C. M.) Fleis.

- Figure 15. Showing plant with leaves. X 2  
Figure 216. Leaf showing shape. X 75  
Figure 217. Leaf showing apex. X 285  
Figure 218. Alar cells of leaf. X 285


### *Papillaria crocea* (Hamp.) Jaeg.

- Figure 219. Showing plant with leaves. X 2  
Figure 220. Leaf showing shape. X 75  
Figure 221. Leaf showing apex. X 285  
Figure 222. Cells of leaf. X 285

### *Aerobryopsis longissima* (Doz. and Molk.) Fleisch.

- Figure 223. Showing plant with leaves. X 2  
Figure 224. Leaf showing shape. X 20  
Figure 225. Leaf apex. X 75  
Figure 226. Cells of leaf. X 285

PLATE XIII



## PLATE XIV

*Barbella tenax* (C. M. ) Broth.

Figure 227. Showing plant with leaves. X 2

Figure 228. Leaf showing apex. X 285

Figure 229. Leaf showing shape. X 75

Figure 230. Cells of leaf. X 750

*Barbella determessi* (Rein. and Card. ) Fleish.

Figure 231. Showing plant with leaves. X 2

Figure 232. Leaf showing shape. X 40

Figure 233. Leaf showing apex. X 285

Figure 234. Cells of leaf. X 750

*Barbella pendula* (Sull. ) Fleisch.

Figure 235. Showing plant with leaves. X 2

Figure 236. Leaf showing shape. X 75

Figure 237. Leaf showing apex. X 285

Figure 238. Cells of leaf. X 750

*Floribundaria floribunda* (Doz. and Molk.) Fleisch.

Figure 239. Showing plant with leaves. X 2

Figure 240. Leaf showing shape. X 75

Figure 241. Leaf showing apex. X 285

Figure 242. Cells of leaf. X 750.


PLATE XIV



227



231




235



239




232



240




228



236




241




229



237



238a



230



234



247

## PLATE XV

*Meteoriopsis squarrosa* (Hook .) Fleisch.

Figure 243. Showing plant with leaves. X 2

Figure 244. Leaf showing shape. X 20

Figure 245. Leaf showing apex. X 285

Figure 246. Cells of leaf. X285

*Meteoriopsis reclinata* (C. M. ) Fleisch.

Figure 247. Showing plant with leaves. X 2

Figure 248. Leaf showing shape. X 75

Figure 249. Leaf showing apex. X 285

Figure 250. Cells of leaf. X 750

*Homaliodedrn exiguum* (Bryol. Jav. ) Fleisch.

Figure 251. Showing plant with leaves. X 2

Figure 252. Leaf showing shape. X 75

Figure 253. Cells of leaf. X 750

*Homaliodendron flabellatum* (Dix. Sm. ) Fleisch.

Figure 254. Showing plants with leave. X 2

Figure 255. Leaf showing shape and apex. X 75

Figure 256. Cells of leaf. X 750

PLATE XV



243



244



245



246



247



252



248



249



250




251



253



254



255



256

## PLATE XVI

*Distichophyllum succulentum* (Mitt.) Broth.

Figure 257. Showing plant with leaves. X 2

Figure 258. Leaf showing shape. X 75

Figure 259. Leaf showing apex X 285

Figure 260. Cells of leaf. X 285

*Hookeria acutifolia* Hook.

Figure 261. Showing plant with leaves. X 2

Figure 262. Leaf showing shape. X 40

Figure 263. Leaf showing apex. X 285

*Hypopterygium tenellum* Lac. = *H. ceylonicum* Mitt.

Figure 264. Showing plant with leaves. X 2

Figure 265. Leaf showing shape. X 75

Figure 266. Cells of leaf. X 285

*Rhegmatodon orthostegius* Mont.

Figure 267. Leaf showing apex. X 285

Figure 268. Showing plant with leaves and sporophyte.  
X2

Figure 269. Leaf showing shape. X 75

Figure 271. Capsule showing shape. X 20

Figure 270. Cells of leaf. X 750

Figure 272. Apophysis showing stomata. X 275


Figure 273. Inner peristome. X285

Figure 274- Peristome. X 285

PLATE XVI




257




258



259




260



263




261



265



262




268




267



264



269



270




271




273



266



272



274

## PLATE XVII

*Herpetineurum toccoae* (Sufl. and Lesq.) Card.

Figure 275. Showing plant with leaves. X 2

Figure 276. Leaf showing shape. X 75

Figure 277. Leaf showing apex. X 750

Figure 278. Cells of leaf. X 750

*Thuidium tamariscellum* (C. M) Bryol. Jav.

Figure 279. Showing branch with leaves and sporophyte. X 2

Figure 280. Branch leaf showing shape. X 75

Figure 281. Branch leaf showing apex. X 285

Figure 282. Stem leaf showing shape. X 75

Figure 283. Stem leaf showing apex. 285

Figure 284. Cells of branch leaf. X 750

Figure 285. Capsule showing shape. X 20

Figure 286. Peristome. X 285

Figure 287. Alar cells of stem leaf. X 750

*Pleuropus nilghiriense* (Mont.) Toyama

Figure 288. Showing branch with leaves and sporophyte. X 2

Figure 289. Leaf showing shape. X 75


Figure 290. Leaf showing apex. X 285

Figure 291. Cells of leaf. X 750

Figure 292. Capsule showing shape. X 20

Figure 293. Peristome. X 285

PLATE XVII



## PLATE XVIII

### *Rhynchostegium javanicum* (Bel.) Eesch.

Figure 294. Showing branch with leaves and sporophyte.  
X 2.

Figure 295. Leaf showing shape. X 20

Figure 296. Leaf showing apex. X 285

Figure 297. Cells of leaf. X 750

Figure 298. Capsule showing shape. X 20

Figure 299. Peristome. X 285

### *Erythrodontium julaceum* (Hook.) Par.

Figure 300. Showing branch with leaves and sporophyte.

Figure 301. Leaf showing shape. X 75

Figure 302. Leaf showing apex. X 285

Figure 303. Cells of lower leaf. X 750

Figure 304. Cells of upper leaf. X 750

Figure 305. Capsule showing shape. X 20

Figure 306. Peristome. X 285

PLATE XVIII



294



295




296



298



299



300



302




297



305




301



303



304



306

## PLATE XIX

### *Foreauella indica* Dix. and Vard.

Figure 307. Showing branch of plant with leaves. x 2.

Figure 308. Leaf showing shape. X 75

Figure 309. Alar cells of leaf. X 750

Figure 310. Leaf showing apex. X 285

### *Sematophyllum subhumile* (C.M.) Fleisch.

Figure 311. Showing branch of plant with leaves and sporophyte. X 2.

Figure 312. Leaf showing shape. X 75

Figure 313. Cells of leaf. X 750

Figure 314. Cells of upper leaf. X 750

Figure 315. Capsule showing shape. X 20

Figure 316. Peristome. X 285

### *Warburgiella leptorrhynchoides* (Mont.) Fleisch.

Figure 317. Showing plant with leaves. X 2

Figure 318. Leaf showing shape. X 75

Figure 319. Leaf showing apex. X 285

Figure 320. Cells of leaf. X 750

### *Hypnum cupressiforme* Hedw.


Figure 321. Showing plant with leaves. X 2

Figure 322. Leaf showing shape. X 285

Figure 323. Leaf showing apex. X 285

Figure 324. Cells of leaf. X 750

PLATE XIX



## PLATE XX

*Taxiphyllum taxirameum* (Mitt.) Fleisch.

Figure 325. Showing plant with leaves. X 2

Figure 326. Leaf showing shape. X 75

Figure 327. Leaf showing apex. X 285

Figure 328. Cells of leaf. X 750

*Ctenidium lychnites* (Mitt.) Broth.

Figure Showing branch of plant with leaves and sporophyte. X 2

Figure 330. Leaf showing apex. X 75

Figure 331. Leaf showing apex. X 285

Figure 332. Cells of leaf. X 750

Figure 333. Capsule showing shape. X 20

Figure 334. Peristome. X 75

Figure 335. Peristome. X 285

*Artichum aculeatum* Card. and Vard.

Figure 336. Showing plant with leaves and sporophyte. X 2

Figure 337. Leaf showing shape. X 20

Figure 338. Cells of leaf. X 285

Figure 339. Cells of upper leaf. X 750

Figure 340. Capsule showing shape. X 20

Figure 341. Peristome. X 285

PLATE XX




PLATE XXI

*Pogonatum microstomum* (R. Br.) Brid.

- Figure 342. Showing plant with leaves and sporophyte. X 2  
Figure 343. Leaf showing shape. X 15  
Figure 344. Leaf showing apex X 285  
Figure 345. Alar cells of leaf. X 285  
Figure 346. Capsule showing shape. X 20  
Figure 347. Peristome. X75  
Figure 348. Peristome. X 285  
Figure 349. Transverse section of leaf. X 75

*Pogonatum neesii* (C. M.) Mitt.

- Figure 350. Showing plant with leaves and sporophyte. X 2  
Figure 351. Leaf showing shape. X 15  
Figure 352. Leaf showing apex. X 285  
Figure 353. Alar cells of leaf. X 285  
Figure 354. Transverse section of leaf. X 285  
Figure 355. Capsule showing shape. X 20  
Figure 356. Peristome. X 285

PLATE XXI





PLATE XXII

- Figure 357. Operculum of *Fissidens schimidii*  
C. M. X 75
- Figure 358. Operculum of *Ditrichum amoenum*  
(Thw. and Mitt.) Par. X 75
- Figure 359. Operculum of *Trematodon ceylonensis*  
C. M. X 75
- Figure 360. Operculum of *Campylopus nodiflorus*  
(C. M.) Jaeg. X 75
- Figure 361. Operculum *Dicranodontium fragile*  
(Hook.) Broth. X 75
- Figure 362. Operculum of *Leucobryum neilgherrense*  
C. M. X 75
- Figure 363. Operculum of *Tortula muralis* Hedw.  
X 75
- Figure 364. Operculum of *Barbula indica* (Schw.)  
Brid. X 75
- Figure 365. Operculum of *Trichostomum cylindricum*  
(Bruch.) C. M. X 75
- Figure 366. Operculum of *Weisia macrospora* Card.  
X 75
- Figure 367. Operculum of *Hyophila involuta* (Hook.)  
Jaeg. X 75
- Figure 368. Operculum of *Funaria hygrometrica* Hedw.  
X 75
- Figure 369. Operculum of *Funaria submarginata* Card.  
and Vard. X 75
- Figure 370. Operculum of *Pohlia flexuosa* (Mitt.)  
Broth. X 75
- Figure 371. Operculum of *Brachymenium exile* (Doz.  
and Molk.) Bryol. X 75
- Figure 372. Operculum of *Anomobryum subnuditum*  
Card and Vard. X 75
- Figure 373. Operculum of *Bryum argenteum* Linn.  
X 75


PLATE XXII



359




359




359



360



361



362



363



364



365




366



367



368



369



371



370



372



373


PLATE XXIII

- Figure 374. Operculum of *Rhodobryum giganteum* (Hook.) Par. X 75
- Figure 375. Operculum of *Rhizogonium spiniforme* (Hedw). Bruch. X 75
- Figure 376. Operculum of *Philonotis anisoclada* Card. and Vard. X 75
- Figure 377. Operculum of *Hypnodon perpusillus* (Thw. and Mitt. C.M. X 75
- Figure 378. Operculum of *Papillaria crocea* (Hamp.) Jaeg. X 75
- Figure 379. Operculum of *Rhegmatodon orthostegius* Mont. X 75
- Figure 380. Operculum of *Thuidium tamariscellum* (C.M.) Bryol. Jav. X 75
- Figure 381. Operculum of *Rhynchostegium javanicum* (Bel.) Besch. X 75
- Figure 382. Operculum of *Sematophyllum subhumile* (C.M.) Fleisch. X 75
- Figure 383. Operculum of *Ctenidium lychnites* (Mitt.)
- Figure 384. Operculum of *Atrichum aculeatum* Card and Vard. X 75
- Figure 385. Operculum of *Pogonatum neesii* (C.M.) Mitt. X 75

PLATE XXIII



374



375



376.



377



378



379



380



381



382



383




384

PLATE XXIV

- Figure 386. Spores of *Fissidens schimidii* C.M. X 750  
 Figure 387. Spores of *Ditrichum amoenum* (Thw. and Mitt.)  
 Par. X 750  
 Figure 388. Spores of *Trematodon ceylonensis* C.M. X 750  
 Figure 389. Spores of *Campylopus nodiflorus* (C.M.) Jaeg. X 750  
 Figure 390. Spores of *Leucobryum neilgherrense* C.M. X 750  
 Figure 391. Spores of *Dicranodontium fragile* (Hook.) Broth.  
 X 750  
 Figure 392. Spores of *Trichostomum cylindricum* (Bruch.) C.M.  
 X 750  
 Figure 393. Spores of *Barbula indica* (Schw.) Brid. X 750  
 Figure 394. Spores of *Tortula muralis* Hedw. X 750  
 Figure 395. Spores of *Funaria hygrometrica* Hedw. X 750  
 Figure 396. Spores of *Funaria submarginata* Card. and Vard.  
 X 750  
 Figure 397. Spores of *Pohlia flexuosa* (Mitt.) Broth. X 750  
 Figure 398. Spores of *Brachymenium exile* (Doz. and Molk.)  
 Bryol. X 750  
 Figure 399. Spores of *Anomobryum subnuditum* Card. and Vard.  
 X 750  
 Figure 400. Spores of *Bryum argenteum* Linn. X 750  
 Figure 401. Spores of *Rhodobryum giganteum* (Hook.) Par. X 750  
 Figure 402. Spores of *Mnium coriaceum* Griff. X 750  
 Figure 403. Spores of *Hyophila involuta* (Hook.) Jaeg. X 750  
 Figure 404. Spores of *Ctenidium lychnites* (Mitt.) Broth. X 750  
 Figure 405. Spores of *Philonotis anisoclada* Card. and Vard.  
 X 750  
 Figure 406. Spores of *Weisia macrospora* Card. X 750  
 Figure 407. Spores of *Hypnodon perpusillus* (Thw. and Mitt.)  
 C.M. X 750  
 Figure 408. Spores of *Schlotheimia grevilleana* Mitt. X 750  
 Figure 409. Spores of *Papillaria crocea* (Hamp.) Jaeg. X 750  
 Figure 410. Spores of *Rhegmatodon orthostegious* Mont X 750  
 Figure 411. Spores of *Thuidium tamariscellum* (C.M.) Bryol.  
 Jav. X 750  
 Figure 412. Spores of *Rhynchostegium javanicum* (Bel.) Besch.  
 X 750  
 Figure 413. Spores of *Sematophyllum subhumile* (C.M.) Fleisch.  
 X 750  
 Figure 414. Spores of *Atrichum aculeatum* Card. and Vard. X 750  
 Figure 415. Spores of *Pogonatum microstomum* (R.Br.) Brid. X 750  
 Figure 416. Spores of *Pogonatum neesii* (C.M.) Mitt. X 750

PLATE XXIV



386




387



388




389



390



391



392



393




394



395




396



397



398



399



400




401



402



403



404



405



406



407



408



409




410



411



412



413



414



415



416



417

PLATE XXV

- Figure 417. Vegetative bud of *Brothera leana* (Sull.)  
C. M. X 75
- Figure 418. Multicellular hooked gemmae of *Brachy-  
nium exile* (Doz. & Molk.) Bryol X 490
- Figure 419. Vegetative bud on leaf of *Rhynchostegium  
javanicum* (Bel.) Fleisch. X 75
- Figure 420. Stomata on apophysis of *Funaria hygrome-  
trica* Hedw. X 285
- Figure 421. Stomata on apophysis of *Pohlia flexuosa*  
(Mitt.) Broth. X 285
- Figure 422. Foot of the sporophyte of *Hyophila  
involuta* (Hook) Jaeg. X 285
- Figure 423. Foot of the sporophyte of *Tortula muralis*  
Hedw. X 285 (Perichaetial region stippled)
- Figure 424. Foot of the sporophyte of *Bryum argen-  
teum* Hedw. X 285
- Figure 425. Foot of the sporophyte of *Barbula indica*  
(Schw.) Brid. X 285
- Figure 426. Foot of the sporophyte of *Rhizogonium  
spiniforme* (Hedw.) Bruch. X 285
- Figure 427. Foot of the sporophyte of *Rhynchostegium  
javanicum* (Bel.) Fleisch. X 285
- Figure 428. Foot of the sporophyte of *Atrichum aculea-  
tum* Card. & Varde X 285
- Figure 429. Foot of the sporophyte of *Pogonatum  
microstomum* (R.Br.) Brid. X 285

PLATE XXV









PLATE-XXVI

- Figure 430. Transverse section of axis of *Hookeria acutifolia* Hook. X 75.
- Figure 431. Transverse section of axis of *Hookeria acutifolia* Hook. Sector passing through centre of axis X 285.
- Figure 432. Transverse section of axis of *Leucobryum neilgherrense* C. M. X 285.
- Figure 433. Transverse section of axis of *Hypopterygium tenellum* X 750.
- Figure 434. Transverse section of axis *Distichophyllum succulentum* (Mitt.) Broth. X 285.
- Figure 435. Transverse section of axis of *Distichophyllum succulentum* (Mitt.) Broth. Centre of axis enlarged. X 750.
- Figure 436. Transverse section of axis of *Floribundaria floribunda* (Doz. & Molk.) Fleisch. X 285.
- Figure 437. Transverse section of axis of *Aerobryopsis longissima* (Doz. & Molk.) Fleisch. Half of section X 285.
- Figure 438. Transverse section of axis of *Warburgiella leptorhynchoides* (Mont.) Fleisch. X 285.
- Figure 439. Transverse section of axis of *Sematophyllum subhumile* (C. M.) Fleisch. X 285


PLATE XXVI




432




433



434





435



436



437




438


## PLATE XXVII

- Figure 440. Transverse section of axis of *Leucobryum scalare* C. M. X 285.
- Figure 441. Transverse section of axis of *Schlotheimia grevilleana* Mitt. X 285.
- Figure 442. Transverse section of axis of *Bartramia madurensis* Dix. & Vard. X 490.
- Figure 443. Transverse section of axis of *Myurium warburgii* (C. Mull.) Fleisch. X 285.
- Figure 444. Transverse section of axis of *Papillaria crocea* (Hamp.) Jaeg X 285.
- Figure 445. Transverse section of axis of *Barbella pendula* (Sull) Fleisch. X 285.
- Figure 446. Transverse section of axis of *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. X 75.
- Figure 447. Transverse section of axis of *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. (Sector- X 285.
- Figure 448. Transverse section of axis of *Foreauella indica* Dix. & Vard. X 285.


PLATE XXVII




440




441




442



443




444




445



448



446




447


## PLATE XXVIII

- Figure 449. Transverse section of axis of *Philonotis anisoclada* Card. & Vard. X 75.
- Figure 450. Transverse section of axis of *Philonotis anisoclada* Card. & Vard. Centre enlarged X 750.
- Figure 451. Transverse section of axis *Trachypus bicolor* Rein. & Hornsch. X 285.
- Figure 452. Transverse section of axis *Trachypus bicolor* Rein. & Hornsch. Centre enlarged X 750.
- Figure 453. Transverse section of axis of *Erythrodontium julaceum* (Hook.) Par. X 285.
- Figure 454. Transverse section of axis of *Erythrodontium julaceum* (Hook.) Par. Centre enlarged X 750.
- Figure 455. Transverse section of axis of *Ctenidium lychnites* (Mitt.) Broth. X 285.
- Figure 456. Transverse section of axis of *Hypnum cupressiforme* Hedw. X 285.
- Figure 457. Transverse section of young axis of *Mnium coriaceum* Griff. X 285
- Figure 458. Transverse section of axis of *Fissidens schimidii* X 285.
- Figure 459. Transverse section of axis of *Ditrichum amoenum* (Thw. & Mitt.) Par. 285


PLATE XXVIII




449




450




451




452




453




454




455



457



456



458




459


## PLATE-XXIX

- Figure 460. • Transverse section of axis of *Meteoriopsis reclinata* (C. M.) Fleisch. X 285
- Figure 461. Transverse section of axis of *Meteoriopsis squarrosa* (Hook.) Fleisch. sector passing through centre of axis X 285
- Figure 462. Transverse section of axis of *Rhynchostegium javanicum* (Bel.) Fleisch. X 285
- Figure 463. Transverse section of axis of *Campylopus nodiflorus* (C. M.) Jaeg. X 285
- Figure 464. Transverse section of axis of *Thysanomitrium depallieri* Card. X 285
- Figure 465. Transverse section of axis of *Hyophila validinervis* Card & Vard. X 285
- Figure 466. Transverse section of axis *Funaria submarginata* X 285
- Figure 467. Transverse section of axis of *Rhodobryum giganteum* (Hook.) par. X 75
- Figure 468. Transverse section of axis of *Barbella tenax* (C. M.) Broth. X 285
- Figure 469. Transverse section of axis of *Fleuropus nilghiriense* (Mont.) Toyama X 285


PLATE XXIX




460




461




462



463




464




465




466



467



468




469


## PLATE-XXX

- Figure 470. Transverse section of old axis of *Mnium coriaceum* Griff. Centre enlarged. X 750
- Figure 471. Transverse section of axis of *Rhegmatodon orthostegius* Mont. X 285
- Figure 472. Transverse section of axis of *Pterobryopsis orientalis* (C. M.) Fleisch. Sector passing through centre of axis. X 750
- Figure 473. Transverse section of young axis of *Atrichum aculeatum* Card. & Vard. Centre of axis enlarged X 750
- Figure 474. Transverse section of old axis of *Atrichum aculeatum* Card. & Vard. X 285
- Figure 475. Transverse section of old axis of *Pogonatum neesii* (C. M.) Mitt. X 285
- Figure 476. Transverse section of seta (at its base) of *Pogonatum neesii* (C. M.) Mitt. X 285


PLATE XXX




470




471




472




473



474



475




476

## PLATE - XXXI

(Photographs of the Herbarium)

- Figure 477. *Fissidens schimidii* C. M. plants X 2  
Figure 478. Plants of *Ditrichum amoenum* (Thw. & Mitt.) par. X 3  
Figure 479. Plants of *Trematodon ceylonensis* C. M. X 1  
Figure 480. Plants of *Campylopus nodiflorus* (C. M.) Jaeg. X 1  
Figure 481. Plants of *Thysanomitrium depalieri* Card. X 3 4  
Figure 482. Plants of *Thysanomitrium lioneurum* (Ther. & Vard.) Broth. X 1-1/4  
Figure 483. Plants of *Brothera leana* (Sul.) C. M. X 1-1/2  
Figure 484. Plants of *Dicranodontium fragile* (Hook.) Broth. X 1-1/4  
Figure 485. Plants of *Leucobryum neilgherrense* C. M. X 2  
Figure 486. Plants of *Leucobryum scalare* C. M. X 1.5  
Figure 487. Plants of *Tortula muralis* Hedw. X 4  
Figure 488. Plants of *Barbula indica* (Schw.) Brid X 2  
Figure 489. Plants of *Trichostomum cylindricum* (Bruch.) C. M. X 1.5  
Figure 490. Plants of *Weisia macrospora* Card. X 1  
Figure 491. Plants of *Hyophila involuta* (Hook.) Jage X 2

PLATE XXXI



## PLATE XXXII

(Photographs of the Herbarium)

- Figure 492. Plants of *Hyophyla validinervis* Card. & Vard. X 2-1/2
- Figure 493. Plants of *Rhacomitrium javanicum* Bryol. X 1
- Figure 494. Plants of *Funaria hygrometrica* Hew. X 1
- Figure 495. Plants of *Funaria submarginata* Carp. & Vard. X 2-1/2
- Figure 496. Plants of *Pholia flexuosa* (Mitt.) Broath. X 3
- Figure 497. Plants of *Brachymenium leptostomoides* (C. M) Schimp. X 1-1/2
- Figure 498. Plants of *Anomobryum subnuditum* Card. & Vard. X 1-1/2
- Figure 499. Plants of *Bryum argenteum* Linn. X 1
- Figure 500. Plants of *Rhodobryum giganteum* (Hook.) Par. X 1
- Figure 501. Branch of *Mnium coriaceum* Griff. 1
- Figure 502. Plants of *Rhizogonium spiniforme* (Hedw.) Bruch. X 1-1/2
- Figure 503. Plants of *Bartramia madurensis* Dix. & Vard. X 1-1/2
- Figure 504. Plants of *Philonotis anisoclada* Card. & Vard. X 1-1/4
- Figure 505. Plants of *Hypnodon perpusillus* Thw. & Mitt. X 1-3/4

PLATE XX XII





PLATE - XXXIII

- Figure 506. Branches of *Schlotheimia grevilleana* Mitt.  
X 2-1/4
- Figure 507. Branch of *Trachypus bicolor* Rein. &  
Hornsch. X 1-3/4
- Figure 508. Branch of *Trachypus humilis* Lindb. var.  
*humilis* Zanten. X 2
- Figure 509. Branch of *Trachypodopsis serrulata* (Beauv.)  
Fleisch. var. *crispatula* Hook. Zanten. X 1
- Figure 510. Branch of *Myurium warburgii* (C. M.)  
Fleisch. X 1-1/2
- Figure 511. Branch of *Petrobryopsis orientlis* (C. M.)  
Fleisch. X 1
- Figure 512. Branch of *Papillaria crocea* (Hamp.) Jaeg.  
X 3/4
- Figure 513. Branch of *Aarobryopsis longissima* Diz. &  
Molk.) Fleisch. X 1
- Figure 514. Plant of *Barbella tenax* (C. M.) Broth. X 1
- Figure 515. Branch of *Barbella determessi* (Rein. &  
Card.) Fleisch. X 3/4
- Figure 516. Branch of *Barbella pendula* (Sull.) Fleisch.  
X 7/8

PLATE XXXIII



## PLATE - XXXIV


(Photographs of the Herbarium)

- Figure 517. Branch of *Floribundaria floribunda* (Doz. & Molk.) Fleisch. X 1
- Figure 518. Branch of *Meteoriopsis squarrosa* (Hook.) Fleisch. X 1
- Figure 519i. Branch of *Meteoriopsis reclinata* (C. M.) Fleisch. X 1
- Figure 520. Plants of *Homaliodendron exiguum* (Bryol. & Jav.) Fleisch. X 1-1/4
- Figure 521. Plant of *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. X 1
- Figure 522. Plants of *Distichophyllum succulentum* (Mitt.) Broth. X 1-1/2
- Figure 523. Plants *Hookeria acutifolia* Hook. X 2
- Figure 524. Branch of *Hypopterygium tenellum* Lac. X 2
- Figure 525. Branch of *Rhegmatodon orthostegius* Mont.
- Figure 526. Plants of *Herpetineurm tocoae* (Sull. & Lesq.) X 1
- Figure 527. Plant of *Rhynchostegium javanicum* (Bel.) Besh. X 1


PLATE XXXIV




519



521




526




518




523




520



522



524



525



527




528

## PLATE - XXXV

(Photographs of the Herbarium)

- Figure 527 - 1 Plants of *Thuidium tamariscellum* (C.M.)  
Bryol. Jav. X 2
- Figure 528. Branches of *Erythrodontium julaceum*  
(Hook.)Pae. X 1-1/2
- Figure 528 - 1 Plant of *Pleuropus nilghiriense* (Mont.)  
Toyama X 1
- Figure 529. Branch of *Foreauella indica* Dix. & Vard.  
X 1-1/2
- Figure 530. Branch of *Sematophyllum subhumile* (C. M.)  
Fleisch. X 2
- Figure 531. Branch of *Warburgiella leptorrhynchoides*  
(Mont.) Fleisch. X 2
- Figure 532. Plant of *Hypnum cupressiforme* Hedw. X 1
- Figure 533. Plant of *Taxiphyllum taxirameum* (Mitt.)  
Fleisch. X 1
- Figure 534. Branch of *Ctenidium lychnites* (Mitt.) Broth.  
X 2-1/4
- Figure 535. Plants of *Atrichum aculeatum* Card. &  
Vard. X 2
- Figure 536. Plants of *Pogonatum microstomum* (R. Br.)  
Brid. X 2/3
- Figure 537. Plants of *Pogonatum neesii* (C. M.) Mitt.  
X 1-1/4

PLATE XXXV




## PLATE - XXXVI


(Photographs of the mosses in nature)

- Figure 538. Plants of *Fissidens schimidii* C. M. in natural habitat at Trivandrum on loose soil. X 1/5
- Figure 539. plants of *Campylopus nodiflorus* (C. M.) Jaeg. in small tufts at Ootacamund. X 1/4
- Figure 540. Plants of *Rhodobryum giganteum* (Hook.) Par. Flash photograph in natural habitat at Kodaikanal. X 1/6
- Figure 541. Plant carpet of *Schlotheimia grevilleana* Mitt. on rock at Kodaikanal. X 1/5
- Figure 542. Plants of *Meteorioposis reclinata* (C. M.) Fleisch. at Kodaikanal. X 1/5
- Figure 543. Plants of *Trachypus bicolor* Rein & Card. on rock at Kodaikanal. X 5


PLATE XXXVI




538




539



540



541



542



543

## PLATE - XXXVII


(Photographs of the mosses in nature)

- Figure 514. Plants of *Thuidium tamariscellum* (C. M.)  
Bryol Jav. at Kodaikanal. X 1/5
- Figure 545. Plants of *Floribundaria floribunda* (Doz. &  
Molk.) Fleisch. at Kodaikanal. X 1/5
- Figure 546. Plants of *Rhegmatodon orthostegius* Mont.  
at Kodaikanal. X 1/8
- Figure 547. Plants of *Barbella determessi* (Rein. & Card.)  
Fleisch. X 1/7
- Figure 548. Plants of *Pogonatum microstomum* (R Brid)  
at Kodaikanal. X 3/5
- Figure 549. Plants of *Pogonatum neesii* (C. M.) Mitt.  
at Ootacamund. X 1/4


PLATE XXXVII




544




543




545



546



548



549

PLATE XXXVIII

(Photomicrographs)

- Figure 550. Part of leaf of *Anomobryum subnuditum* Card. & Vard. Note rhomboid large cells and prominent costa. X 750
- Figure 551. Part of leaf of *Sematophyllum subhumile* (C.M.) Fleish. Note narrow elongated cells and absence of mid-rib. X 490
- Figure 552. Part of leaf of *Mnium coriaceum* Griff. Note margin made up of thick-walled rows of elongated cells. X 750
- Figure 553. Part of leaf of *Rhodubryum giganteum* (Hook.) Par. Note serrulate margin and heavily thickened laminar cells. X 750
- Figure 554. Part of leaf of *Rhegmatodon orthostegius* Mont. Note costa ending in the middle of the leaf. X 490
- Figure 555. Part of leaf of *Dicranodontium fragile* (Hook.) Broth. Note broad costa and rectangular laminar cells. X 490
- Figure 556. Part of leaf of *Fissidens schimidii* C. M. Note the dorsal lamella X 490
- Figure 557. Part of leaf of *Ctenidium lychnites* (Mitt.) Broth. Note elongated and narrow cells of the lamina and the margin. X 490
- Figure 558. Base of leaf of *Taxiphyllum taxirameum* (Mitt.) Fleisch. Note rhizoids arising from leaf base. X 490
- Figure 559. Part of leaf of *Hookeria acutifolia* Hook. Note large cells of lamina. X 750
- Figure 560. Part of leaf of *Hypnum cupressiforme* Hedw. Note larger alar cells X 490
- Figure 561. Part of leaf of *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. Note different shapes of cells. X 490
- Figure 562. Part of leaf of *Leucobryum neilgherrense* C. M. Note rhizoids arising from the left margin. X 490

PLATE XXXVIII





PLATE - XXXIX


(Photomicrographs)

- Figure 563. Part of leaf of *Anomobryum subnuditum* Card & Vard. Note excurrent costa. X 490
- Figure 564. Apex of leaf of *Leucobryum neilgherrense* C. M. Note production of rhizoids from leaf apex. X 490
- Figure 565. Spores of *Dicranodontium fragile* (Hook.) Broth. Note thickening of costa. X 490
- Figure 566. Part of leaf of *Funaria submarginata* Card. & Vard. Note leaf base with prominent costa. X 490
- Figure 567. Part of leaf of *Rhynchostegium javanicum* (Bel.) Fleisch. Note alar cells quadratic to rectangular X 490
- Figure 568. Peristomial teeth in *Dicranodontium fragile* (Hook.) Broth. Note Haplolepideae structure of peristome inner face. X 200
- Figure 569. Part of leaf of *Barbella deterrmessi* (Rein & Card.) Fleisch. Note absence of costa. 490


PLATE XXXIX




563




564




565



566



567



568




569

## PLATE - XL


### (Photomicrographs)

- Figure 570. Stomata on the apophysis of *Trematodon ceylonensis* C. M. Note elongated epidermal cells. X 750
- Figure 571. Stoma on the apophysis of *Funaria hygrometrica* Hedw. Note numerous subsidiary cells of stoma. X 750
- Figure 572. Transverse section of axis of *Barbella determessi* (Rein. & Card.) Fleisch. Note central small cell surrounded by a ring of cells. X 200
- Figure 573. Stomata on the apophysis of *Funaria hygrometrica* Hedw. Note frequency of stomata. X 200
- Figure 574. Transverse section of axis of *Thysanotrium depallieri* Card. Note central axis hollowed. X 200
- Figure 575. Part of Transverse section of axis of *Barbella determessi* (Rein. & Card) Fleisch. Note organisation of cavity. X 400
- Figure 576. Transverse section of axis of *Thuidium tamariscellum* (C.M.) Bryol. Note undifferentiated tissue of the axis inside the rind. X 200


PLATE XL




570




571




572




573



574



575



576

## PLATE - XLI

### (Photomicrographs)

- Figure 577. Longitudinal section of axis of *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. Note oblique or straight cross walls. X 490
- Figure 578. Part of Transverse section of *Mnium coriaceum* Griff. Note presence of pith in the centre X 490
- Figure 579. Part of Transverse section of axis of *Ctenidium lychnites* (Mitt.) Broth. Note presence of single cell with a ring of radiating and elongated cells in the centre of axis. X 490
- Figure 580. Transverse section of axis of *Mnium coriaceum* Griff. centre. enlarged. Note pith cells having angular thickening. X 750
- Figure 581. Transverse section of axis of *Homaliodendron flabellatum* (Dix. Sm.) Fleisch. Note undifferentiated uniformly thickened cells. inside rind X 200
- Figure 582. Transverse section of axis of *Hypnum cupressiforme* Hedw. centre enlarged. Note organisation of cavity. X 750
- Figure 583. Transverse section of axis of *Barbella tenax* (C.M.) Broth. Note the large hollow cavity of axis. X 490
- Figure 584. Transverse section of axis of *Pogonatum neesii* (C.M.) Mitt. Note the angular projection of the central stand and differentiation of cortex and also of strand. X 200

PLATE XLI













PLATE XLII (Contd.)

SCHEFFNER (1938)	COMB NO (1984)	PUNTI MALACZAR HYPMACZAR
POLYTRICHALES	PLEUHOCAPI	RHIZOCHLACZAR FLANACZAR
SPILACRHALES	ACROCARPI	BUI SACNMACZAR SPILACSNMACZAR
STYMALES	HETEROLOPILEAF	DISCELLACZAR FISSELENTACZAR
BOOSENALLES	RAPLOLEPIDEAF	POLYTRICZACZAR TETRAPEHIDACZAR
LEUCOLOFALLES	TERRIA	ANDREARACZAR
DICNARALLES	HEMTOLOPTEAF	SPERACHACZAR
ANCHILALLES	ADREAROBRYA	
ODOFOCARPAL	SPERACHOBRYA	

PLATE XLII (Contd.)

