

17P INSTRUCTION CHECKLIST

PATIENT: _____ **DOB:** _____

Injection Teaching

Basic overview of medication
17P (17 alpha hydroxy progesterone caproate/ delalutin)
250mg/1cc IM q week with 21G 1 ½ "needle and 3 cc syringe

- Roll the medication bottle between your palms
- Clean top of bottle with rubbing alcohol
- Open package with syringe and needle
- Pull syringe plunger back until top of black plunger is even with 1 cc black line
- Take off needle cap
- Put the needle into the top part of the bottle and inject 1 cc of air into bottle
- With needle still inside of bottle, turn bottle upside down and pull plunger back
- Fill syringe with 1 cc medication
- Re-cap needle
- Confirm correct place for shot :

- Clean area with rubbing alcohol
- Insert needle at 90 degree angle
- Pull back plunger and watch for blood
 - If you see blood, throw away in the red box and start again
- If no blood, push plunger and inject all of medication
- Dispose of needle and syringe in appropriate biohazard box

- Person taught to give injection _____
- Person taught return demo

- Lancet Disposal / Biohazard Box (office / home disposal)

Referral to for follow up

County Home Health Nurse: _____

Spoke with _____

Appointment planned _____

Referring Provider _____

Spoke with _____

Appointment planned _____

PROVIDER SIGNATURE: _____ **DATE:** _____