






# THEOREM OF THE DAY



**Netto's Conjecture (Dixon's Theorem)** *Two random random permutations of  $\{1, \dots, n\}$  generate either the symmetric group  $S_n$  or the alternating group  $A_n$  with probability tending to 1 as  $n$  tends to infinity.*



- Indexed by even permutation pair generating all of  $A_n$
- Indexed by permutation pair generating all of  $S_n$
- Indexed by permutation pair fixing a point in common

The symmetric group  $S_n$  consists of all  $n!$  permutations of  $\{1, \dots, n\}$ . For example, each  $S_n$ , for  $n \geq 5$ , contains the permutation  $(13)(254)$  (written in so-called 'disjoint cycle notation'): it permutes two numbers, 1 and 3 (this called is a transposition) and cycles through 2, 5 and 4. Each permutation has a parity, even or odd, which may be computed as the parity of the number of gaps between numbers:  $(13)(254)$  has three gaps (i.e. 1-3, 2-5 and 5-4) and is an odd permutation, whereas  $(13254)$  has four gaps and is even. The alternating group  $A_n$  is the subgroup of all  $n!/2$  even permutations. The tables above record those pairs of permutations which generate (via combining permutations by multiplication) all of  $A_n$  or  $S_n$ . This is immediately impossible for pairs which fix a common point: such pairs are also recorded, in yellow. In the limit the white and yellow cells become a vanishingly small proportion.

Eugen Netto's 1882 conjecture waited nearly a century for a proof. In 1969 John D. Dixon proved that, when  $n$  is sufficiently large, the proportion of pairs of permutations of  $n$  elements that generate either  $S_n$  or  $A_n$  exceeds  $1 - 2/(\log \log n)^2$ . This number approaches 1 very slowly (e.g. staying below 0.99 until  $n \approx 10^{10^6}$ ); Dixon conjectured that the actual proportion would be  $1 - 1/n + O(1/n^2)$ , with the  $1/n$  term being the contribution of pairs of permutations having a common fixed point (the yellow squares in our illustration). Dixon's conjecture was proved in 1989 by László Babai, assuming the classification of the finite simple groups.

**Web link:** [cameroncounts.wordpress.com/2011/04/09](http://cameroncounts.wordpress.com/2011/04/09)

**Further reading:** [Permutation Groups](#) by John D. Dixon and Brian Mortimer, Springer, 1996.

