

Aspidosperma Species as Sources of Antimalarials. Part III. A Review of Traditional Use and Antimalarial Activity*

Authors

Renata Cristina de Paula^{1**}, Maria Fâni Dolabela^{2**}, Alaíde Braga de Oliveira^{1**}

Affiliations

¹ Departamento de Produtos Farmacêuticos, Faculdade de Farmácia, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

² Departamento de Farmácia, ICS, Universidade Federal do Pará, Belém, PA, Brazil

Key words

- *Aspidosperma* spp.
- Apocynaceae
- malaria
- *Plasmodium falciparum*
- antimalarial activity

Abstract

Several plant species belonging to the genus *Aspidosperma* are traditionally used in Brazil and other Meso- and South American countries for the treatment of malaria and fevers. These traditional uses were motivation for this review. A literature survey completed for this review has identified scientific bibliographical references to the use of 24 *Aspidosperma* species to treat malaria/fevers and to 19 species that have had their extracts

and/or alkaloids evaluated, with good results, for *in vitro* and/or *in vivo* antimalarial activity. Indole alkaloids are typical constituents of *Aspidosperma* species. However, only 20 out of more than 200 known indole alkaloids isolated from this genus have been assayed for antimalarial activity. These data support the potential of *Aspidosperma* species as sources of antimalarials and the importance of research aimed at validating their use in the treatment of human malaria.

Introduction

Natural products and malaria

Malaria remains one of the most prevalent infectious diseases worldwide and is, therefore, a global health problem despite substantial efforts to control the disease over the past few decades. Approximately 3.3 billion people are at risk, and 250 million cases each year were reported in the period 2006–2008, primarily in Africa [1]. In the Americas, malaria transmission occurs in 21 countries. *P. vivax* caused 77% of all cases reported in 2008, but *P. falciparum* was responsible for almost 100% of all cases in Haiti and the Dominican Republic [1]. Brazil reported the highest number of malaria cases (603 532) in the region in 2005, primarily in the Brazilian Legal Amazon Region, where 10–15% of the population is at risk. Brazil was among the 30 highest-burden countries for malaria [2]. However, a decrease of approximately 25% in the number of reported cases has been recorded since 2006 [3,4].

Historically, plants have had a remarkable role in therapeutics and were the principal source of drugs until the 19th century. Quinine, isolated in 1820, from *Cinchona* species (Rubiaceae), was the

first antimalarial drug introduced in chemotherapy and remained the only clinical weapon until the 1940s, when chloroquine, a synthetic 4-aminoquinoline, became available. Efficient and inexpensive, chloroquine was widely used until the 1960s, when resistance to the drug by *P. falciparum* became widespread in the malaria-endemic countries, causing a strong increase in mortality rates. The antimalarial drugs in current use are artemisinin, the active compound from *Artemisia annua* L. (Asteraceae), a traditional plant used for millennia in China, and its semisynthetic derivatives artemether, artesunate, and arteether [5]. Artemisinins are currently the most effective drugs for antimalarial chemotherapy and have been globally adopted for the treatment of *P. falciparum* malaria. The most recently introduced antimalarial drug is atovaquone, a synthetic naphthoquinone based on lapachol. Lapachol, a prenylnaphthoquinone, was first isolated from *Tabeuia impetiginosa* (Mart. ex DC.) Standl. (synon. *T. avellanedeae* Lor. ex Griseb.), a South American representative of the Bignoniaceae [6].

The emergence of *P. falciparum* strains resistant to artemisinin and its derivatives would cause a resurgence of human malaria to high levels in many

received August 18, 2013
revised January 10, 2014
accepted January 11, 2014

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1368168>
Published online March 3, 2014
Planta Med 2014; 80: 378–386
© Georg Thieme Verlag KG
Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Alaíde Braga de Oliveira
Departamento de Produtos Farmacêuticos
Faculdade de Farmácia
Universidade Federal de Minas Gerais
Av. Antônio Carlos, 6.627
31270–901 Belo Horizonte, MG
Brazil
Phone: + 55 31 34096950
Fax: + 55 31 3441 5575
alaide.braga@pq.cnpq.br

* For Part I see [48], for Part II see [36]. Part of RCP Doctorate Thesis at PPGCF, UFMG, Belo Horizonte, MG, Brazil.

** All authors contributed equally to this article.

countries. For this reason, R&D on new antimalarial drugs is urgent. Plants continue to represent a valuable source of drugs. A review of all small molecules that have been approved as pharmaceutical agents within the 25-year period from 1/1981 to 6/2006 has demonstrated that approximately 50% of these molecules originated from natural products [7]. Investigations that begin by screening plants used in traditional medicines are particularly valuable. In the past decade, a substantial number of publications have focused on the screening of either extracts or natural products for antimalarial activity [8]. However, as we have noted previously, very few highly active antimalarial natural products have been evaluated for cytotoxicity and in *in vivo* assays. This lack of previous study limits the potential of these products as bases for the development of new antimalarial drugs [5,9]. The relevance of these results should not, however, be underestimated. Active naturally occurring antimalarial compounds might be useful as bases for semisynthetic derivatives, as agents for direct use if their structures are too complex for an economically and/or technologically viable total synthesis, or as templates for the total synthesis of structurally related compounds [5]. Moreover, bioactive natural products from medicinal plants might be useful as biological markers in the development of efficient and safe phytomedicines [5,9], a new approach of growing interest that would provide malaria-endemic countries with good-quality herbal medicines of low cost, that would be locally and sustainably produced [8]. High-tech methods are available to standardize phytopreparations [10,11], and new molecular biological assays can serve to screen extracts and plant constituents as well as to evaluate their pharmacological profiles, elucidate the synergistic effects of the constituents of an extract and, thus, gain a better understanding of the various mechanisms underlying their pharmacological effects [11–14]. Several *Aspidosperma* species have a history of medicinal uses, including the treatment of human malaria and/or fevers in Brazil as well as in other Meso- and South American countries [15–19]. The present review reports the results of a literature survey on *Aspidosperma* species traditionally used to treat malaria and/or fevers as well as data on those previously evaluated for antimalarial activity.

The Genus *Aspidosperma*

Taxonomy and geographic distribution

The genus *Aspidosperma* Mart. & Zucc. belongs to the family Apocynaceae, one of the ten largest angiosperm families [20]. The Apocynaceae s.l. belong to the order Gentianales and have a primarily pantropical/subtropical distribution with few temperate representatives [21].

Aspidosperma species are of outstanding economic interest as sources of valuable timber. They are found from Mexico to Argentina. The number of species in this genus is controversial. Woodson [22] recognizes 52 species, whereas Marcondes-Ferreira and Kinoshita [23] have proposed an infra-generic division for this genus with 43 species, 32 of them occurring in Brazil. However, in the “Lista de Espécies da Flora do Brasil (LEFB)”, 51 species are listed with accepted names; 21 of these species are endemic to Brazil, and 90 synonyms are reported [24].

Traditional use in treating malaria

Data obtained by searching for *Aspidosperma* species with reported traditional uses as antimalarials and/or febrifuges as well as those species that have been experimentally evaluated for antimalarial activity are shown in **Table 1**. The table includes the accepted taxonomic nomenclature, synonyms, and occurrence, according to Koch and collaborators [20], LEFB [24], Lorenzi (1992) [15], TROPICOS® Specimen Data Base [25], and The Plant List [26]. The reported uses for treating malaria/fevers are based on the scientific literature. Medicinal uses described on commercial Internet sites, as well as uses other than for malaria and/or fevers, have not been included. Reported antimalarial activity may refer to *in vitro* assays with different *Plasmodium* species, *in vivo* assays in different animals with different *Plasmodium* species, or clinical assays in humans. Details of these findings are shown in **Table 2**.

Our literature survey has identified 24 *Aspidosperma* species (**Table 1**) used to treat malaria/fevers, representing approximately 50% of the representatives of this taxon [24]. In addition to these 24 species, **Table 1** includes two other species, *A. cylindrocapon* and *A. macrocarpon*, for which no reports have been found on their use to treat malaria/fever but that have been evaluated for antimalarial activity [27,28].

The “Dicionário das plantas úteis do Brasil e das exóticas cultivadas” (Dictionary of useful plants from Brazil and of the exotics cultivated), by Pio-Corrêa (1874–1934) [17], was our first source of information on the antimalarial use of plants belonging to the genus *Aspidosperma*. More than 50 *Aspidosperma* species are listed within this six-volume collection, with taxonomic data and vernacular names. Most of the applications described there are as sources of timber, whereas only three species, *A. discolor*, *A. polyneuron*, and *A. gomezianum*, are identified as plants used to treat malaria and/or fevers.

The botanical names shown for the 26 *Aspidosperma* species in **Table 1** are those originally reported in the literature. However, it should be emphasized that these are not the presently accepted names in certain cases. This is the situation for *A. marcgravianum* Woodson (*A. marcgravianum*), which is not presently an accepted name [24] but a synonym of *A. excelsum* Benth. together with *A. nitidum* Benth. ex Müll. Arg., according to Koch and collaborators [20] and LEFB [24], but is described as a synonym for *A. excelsum* only by the The Plant List [26]. In the same source [26], however, *A. nitidum* Benth. ex Müll. Arg. is an accepted name, with *A. aquaticum* Ducke as a synonym. Another contradiction is found for *A. album* (Vahl) Benoist ex Pichon and *A. desmanthum* Benth. ex Müll. Arg., cited as accepted names for distinct species [20,24,26], whereas the second species represents a synonym for the first one [25]. Furthermore, *A. parvifolium* A. DC. is the accepted name of a species that has *A. tambopatense* A.H. Gentry and *A. vargasii* A. DC. as synonyms [20,24], whereas the latter synonym is also cited as a distinct species [25,26]. A similar situation is observed for *A. tomentosum* Mart., an accepted name having *A. gomezianum* A. DC. as a synonym [25,26], although both of them are also reported as distinct species [20,26]. These observations show that several controversies remain in the taxonomy of *Aspidosperma* species.

Phytochemistry and antimalarial activity

Chemically, *Aspidosperma* species are characterized by the presence of alkaloids and have yielded more than 200 indole alkaloids. The phytochemical investigation of several Brazilian *Aspidosperma* species, primarily in the period between 1960 and

Table 1 Literature survey on *Aspidosperma* species, their traditional use to treat malaria and/or fevers, and reported antimalarial activity.

<i>Aspidosperma</i> species	Local names in Brazil References: [15, 17, 20, 24]	Occurrence References: [15, 17, 20, 24–26]	Reported use to treat malaria/fever	Reported anti-malarial activity
<i>A. album</i> (Vahl) Benoist ex Pichon Syn. <i>A. desmanthum</i> Benth. ex Müll. Arg., <i>A. centrale</i> Markgr. [20, 24, 26]	Cabeça-de-arara, Piquiá-marfim, Araracanga, Aracanga-preta, Maparana	Bolívia, Colômbia, French Guiana, Guyana, Suriname, Venezuela, Brazil: Amazonia	Yes [62, 63]	No
<i>A. auriculatum</i> Markgr. [20, 26]	Carapanaúba Carapanaúba-amarela	Brazil: Endemic (PA)	Yes [64]	No
<i>A. cuspa</i> (Kunth) S. F. Blake ex Pittier Syn. <i>A. decipiens</i> Müll. Arg. (+ 11 synonym.) [20, 26]	Guatambuzinho, Guatambú-branco, Amargoso, Peroba-de-Goiás	Bolívia, Colombia, Ecuador, Haiti, Venezuela, Paraguay, Peru, Republica Dominicana, Trinidad Tobago, Venezuela, Brazil: Amazonia, Caatinga, Cerrado, Atlantic Rainforest	Yes [65]	No
<i>A. cylindrocarpon</i> Müll. Arg. Syn. <i>A. brevifolia</i> Rusby [20, 26]	Peroba-iquira, Peroba de Lagoa Santa, Peroba de Minas, Peroba rosa	Bolívia, Paraguay Peru, Brazil: Amazônia, Cerrado, Atlantic Rainforest	No	Yes See Table 2
<i>A. desmanthum</i> Benth. ex Müll. Arg Accepted name Syn. <i>A. cruentum</i> Woodson, <i>A. chiapense</i> Matuda, <i>A. matudae</i> Lundell [20, 24, 26] Syn. <i>A. album</i> (Vahl) Benoist ex Pichon [25]	Araraíba, Peroba, Arraracanga, Amargoso, Pequiá-marfim	Belize, Bolívia, Colombia, Ecuador, French Guiana, Honduras, Mexico, Panama, Nicaragua, Brazil: Amazonia, Atlantic Forest	Yes [17, 63, 66]	Yes See Table 2
<i>A. discolor</i> A. DC. Accepted name Syn. <i>A. francisii</i> A. DC. [20, 26]	Cabo-de-machado, Quina, Carapanaúba-amarela, Pau-pereira, Peroba-de-rego, Peroba	Bolívia, French Guiana, Guyana, Peru, Suriname, Venezuela, Brazil: Amazonia, Caatinga, Cerrado, Atlantic Forest	Yes [17]	No
<i>A. excelsum</i> Benth. Accepted name Syn. <i>A. marcgravianum</i> Woodson, <i>A. nitidum</i> Benth. ex Müll. Arg. [20, 25, 26]	Carapanaúba, Carapanaúba-preta, Sapopema, Sapopemba	Costa Rica, Panama, Colombia, French Guiana, Guyana, Ecuador, Peru, Bolívia, Venezuela, Brazil: N, NE, SE, CW	Yes [67–70]	Yes See Table 2
<i>A. gomezianum</i> A. DC. Accepted name [20, 24] Non-cons., syn. of <i>A. tomentosum</i> Mart. [25]	Peroba-amarela, Pequiá-de-pedra, Pau-cetim, Guatambu, Ipê-peroba	Brazil: NE (Bahia), SE/Minas Gerais, Espírito Santo, Rio de Janeiro	Yes [17, 28]	No
<i>A. macrocarpon</i> Mart. Accepted name Syn. <i>A. macrocarpum</i> Mart., <i>A. duckei</i> Huber ex Ducke, <i>A. gardneri</i> Müll. Arg. (+ 6 other syn.) [20, 26]	Guatambú, Guatambú-do-cerrado, Peroba-do-campo, Pau-pereira, Pereiro	Bolívia, Ecuador, Peru, Brazil: Amazonia, Cerrado	No	Yes See Table 2
<i>A. marcgravianum</i> Woodson This name is a synonym of <i>A. excelsum</i> Benth and <i>A. nitidum</i> Benth. ex Müll. Arg. [24, 26]	Carapanaúba, Carapanaúba-preta	Bolívia, Colômbia, French Guiana, Guyana, Ecuador, Panamá, Peru, Suriname, Venezuela	Yes [71]	Yes See Table 2
<i>A. megalocarpon</i> Müll. Arg. Accepted name Syn. <i>A. lundellianum</i> Woodson, <i>A. saguinale</i> Bartl., <i>A. stegomeris</i> (Woodson) Woodson [22, 26]	Pequiá-marfim, Araracanga, Carapanaúba, Carapanaúba-preta	Central America from México to Colombia [25], Brazil: Endemic (Acre) [24]	Yes [36]	Yes See Table 2
<i>A. nitidum</i> Benth ex Müll. Arg. Accepted name Syn. <i>A. aquaticum</i> Ducke, Syn. of <i>A. excelsum</i> Benth. [24–26]	Carapanaúba, Maparaná	Peru [25], Brazil: Amazonas [25], Roraima [24]	Yes [72]	Yes See Table 2
<i>A. oblongum</i> A. DC. Accepted name Syn. <i>A. khulmannii</i> Markgr. [26]	Carapanaúba, Carapanaúba-amarela	French Guiana, Guyana, Suriname, Venezuela, Brazil: Pará, Amazonas, Maranhão, Goiás	Yes [73] (<i>A. oblongum</i>)	Yes See Table 2
<i>A. olivaceum</i> Müll. Arg. Accepted name Syn. <i>A. bello-horizontinum</i> Silveira [20, 26]	Guatambú, Guatambu-mirim	Brazil: Endemic to Atlantic Rainforest (NE, SE, S) [24], Argentina [23, 25]	Yes [67]	Yes See Table 2
<i>A. parvifolium</i> A. DC. Accepted name Syn. <i>A. ingratum</i> K. Schum., <i>A. vargasii</i> A. DC., <i>A. tambopatense</i> A. H. Gentry [20, 25, 26]	Pau-Pereira, Peroba, Guatambú, Guatambú-rosa, Guatambú-marfim Amarelão	Argentina, Bolívia, Guyana, Paraguay, Peru, Venezuela, Brazil: Amazonia, Caatinga, Cerrado, Atlantic Rainforest	Yes [74]	Yes See Table 2

continued

Table 1 Continued

<i>Aspidosperma</i> species	Local names in Brazil References: [15, 17, 20, 24]	Occurrence References: [15, 17, 20, 24–26]	Reported use to treat malaria/fever	Reported anti-malarial activity
<i>A. polyneuron</i> Müll. Arg. Accepted name Syn. <i>A. dugandii</i> Standl., <i>A. peroba</i> Allemão ex Saldanha, <i>A. venosum</i> Müll. Arg. [20, 24, 26]	Peroba-rosa, Sobro, Peroba-amargosa	Argentina, Bolívia, Colombia, Paraguay, Peru, Venezuela, Brazil: Caatinga, Cerrado, Atlantic Rainforest	Yes [17, 75]	Yes See Table 2
<i>A. pyriforme</i> Mart. Accepted name Syn. <i>A. bicolor</i> Mart. (+ several others) [26]	Pereiro-do-sertão, Pereiro-preto, Pau-de-coam, Pequiá-da-restinga	Argentina, Bolívia, Paraguay, Brazil: Caatinga, Cerrado	Yes [62]	Yes See Table 2
<i>A. quebracho blanco</i> Schldt. Accepted name Syn. <i>A. quebracho</i> Griseb (+ several others) [26]	Quebracho-branco (Brazil), Quebracho-branco (Argentina)	Argentina, Bolívia, Paraguay, Uruguay, Brazil: Mato Grosso	Yes [68, 72, 76, 77]	Yes See Table 2
<i>A. ramiflorum</i> Müll. Arg. Accepted name Syn. <i>Geissospermum ramiflorum</i> (Müll. Arg.) Miers [26]	Pau-pereira, Peroba, Guatambú-amarelo	Bolívia, Brazil: Amazonia Cerrado, Atlantic Rainforest	Yes [22]	Yes See Table 2
<i>A. rigidum</i> Rusby Accepted name Syn. <i>A. acreanum</i> Markgr. (+ several others) [20, 26]	Carapanaúba, Aracaranga	Bolívia, Colombia, Costa Rica, Ecuador, Panamá, Peru, Venezuela [25], Brazil: Endemic (Amazonia, Cerrado) [24]	Yes [39]	Yes See Table 2
<i>A. sandwithianum</i> Markgr. Accepted name No synonyms [20, 26]	Carapanaúba	French Guiana, Guyana, Suriname, Brazil: Pará, Amazonas	Yes [71]	No
<i>A. schultesii</i> Woodson Accepted name No synonyms [20, 26]	Maku	French Guiana, Peru, Venezuela Brazil: Amazonas, Roraima, Mato Grosso	Yes [66, 71]	No
<i>A. spruceanum</i> Benth. ex Müll. Arg. Accepted name Syn. <i>A. igapoanum</i> Markgr., <i>A. melanocalix</i> Müll. Arg. (+ several others) [20, 26]	Amargoso, Aracaranga, Guatambu, Peroba, Quina da mata, Pequiá marfim	Bolívia, Colombia, Ecuador, French Guiana, Mexico, Panama, Peru, Venezuela, Brazil: Amazonia, Cerrado, Atlantic Rainforest	Yes [78]	Yes See Table 2
<i>A. tomentosum</i> Mart Accepted name Syn. <i>A. camporum</i> Müll. Arg., <i>A. dasycarpon</i> A. DC. and others [20, 24, 26] Non-accepted name, syn. for <i>A. gomezianum</i> A. DC. fide Woodson, Jr. R. E., 1951 [25]	Guatambú-do-cerrado, Pereiro-do-campo	Bolívia, Paraguay, Brazil: Amazonia, Caatinga, Cerrado	Yes [79]	No
<i>A. ulei</i> Markgr. Accepted name Syn. <i>A. occidentale</i> Markgr. [20, 26]	Pitiá	Guyana, Suriname, Venezuela, Brasil: Amazonia, Atlantic Forest	Yes [22]	Yes See Table 2
<i>A. vargasii</i> A. DC. Accepted name [25, 26] Non-accepted name, syn. for <i>A. parvifolium</i> A. DC. [20, 24]	Amarelão	Bolívia, Colombia, Guyana, Peru, Suriname, Venezuela	Yes [62, 66, 71]	Yes See Table 2

1980, is highly valuable and can be appreciated in a 2007 review [29]. However, there are relatively few reports on the biological activities of these alkaloids [5, 30].

Of the 26 *Aspidosperma* species included in [Table 1](#), 19 have had their crude extracts and/or alkaloidal extracts assayed for antimalarial activity. Detailed data are shown in [Table 2](#). Evaluations were performed as early as 1932, and several experiments were conducted before the methodology for the *P. falciparum* erythrocytic culture became available [31]. These experiments include *in vivo* assays in ducklings infected with *P. lophurae* [32], *in vitro* assays with *P. cathemerium* [33], and clinical assays in humans [34, 35] ([Table 2](#)). Good activity ($IC_{50} < 10 \mu\text{g/mL}$) has been reported for extracts from *A. macrocarpon* [28], *A. megalocarpon* [36], *A. oblongum* [37], *A. olivaceum* [27], *A. quebracho blanco* [38], and *A. rigidum* [39]. A hydroethanolic extract from *A. pyriforme* stem bark was inactive against the *P. falciparum* F32 strain [40]. However, a series of indole alkaloids isolated from the same species were shown to be active against *P. falciparum* FcM29 and Nigerian strains [41, 42]. Most likely, this controversy can be explained by a low content of alkaloids in the inactive crude extract. Extracts from *A. cylindrocarpon* [27, 43] and *A. macrocarpon* [28], the two species with no reported uses in treating malaria, have shown *in vitro* activity against three different *P. falciparum* clones (W2, 3D7, FcB1), with a selectivity index > 10 ($SI = CC_{50}/IC_{50}$) in every case ([Table 2](#)). These results demonstrate the validity of the taxonomic approach for the selection

de Paula RC et al. *Aspidosperma* Species as... Planta Med 2014; 80: 378–386

Table 2 Antimalarial activity of *Aspidosperma* extracts and alkaloids.

<i>Aspidosperma</i> species	Country of collection	Part of the plant	Extracts Compounds	Bioassay <i>Plasmodium</i> species/ strains	Results	References
<i>A. cylindrocarpon</i>	Brazil	Trunkwood	EtOH	<i>In vitro</i> / <i>P. falciparum</i> W2 and 3D7 strains	W2: IC ₅₀ = 44 µg/mL 3D7: IC ₅₀ = 39 µg/mL Cytotoxicity Vero cells CC ₅₀ > 500 µg/mL SI: W2 = 11.4 SI: 3D7 = 12.8	[27]
<i>A. desmanthum</i>	Brazil	Trunk bark	Aspidocarpine (2)	<i>In vitro</i> / <i>P. falciparum</i> K1	IC ₅₀ = 0.019 µM (0.07 µg/mL)	[45]
<i>A. excelsum</i>	Peru (Remo caspi)	Trunk bark	EtOH	<i>In vitro</i> / <i>P. falciparum</i> 3D7	IC ₅₀ = 42 µg/mL SI human lymphocyte inhibition IC ₅₀ > 100 µg/mL	[39]
<i>A. macrocarpon</i>	Brazil	Root bark	EtOH	<i>In vitro</i> / <i>P. falciparum</i> FcB1	FcB1: IC ₅₀ = 4.9 µg/mL Cytotoxicity MRC-5 CC ₅₀ = 79.2 µg/mL SI = 16.2	[28]
<i>A. marcgravianum</i>	Guyane	Leaves	Tetrahydro-usambaresine (175)	<i>In vitro</i> / <i>P. falciparum</i> FcM29	0.26 µg/mL (0.59 µM)	[32]
<i>A. megalocarpon</i>	Colombia	Trunk bark	MeOH	<i>In vitro</i> / <i>P. falciparum</i> D2 and F32 strains	MeOH extract D2 strain: IC ₅₀ = 8 µg/mL F32 strain: IC ₅₀ = 25 µg/mL Cytotoxicity U-937 human promonocytic cells: CC ₅₀ = 0.4 µg/mL SI: Pf/D2 = 0.05 SI: Pf/F32 = 0.02	[36]
<i>A. megalocarpon</i>	Colombia		Fendlerine (13), Aspidoalbine (14), Aspidolimidine (15)	<i>In vitro</i> / <i>P. falciparum</i> FcM29 and Nigerian strains	FcM29: IC ₅₀ = 25.6 to 59.2 µg/mL Nigerian strain: IC ₅₀ = 28.0 to 57.3 µg/mL	[41, 53]
<i>A. nitidum</i>	Brazil	–	–	<i>In vivo</i> Duckling/ <i>P. lophurae</i>	Inactive	[32, 80]
<i>A. nitidum</i>	Brazil	Trunk bark	H ₂ O extract	<i>In vivo</i> / <i>P. berghei</i>	Inactive	[44]
<i>A. oblongum</i>	Brazil	–	EtOH	<i>In vitro</i> / <i>P. falciparum</i> W2 and D6	W2: IC ₅₀ = 4742.5 ng/mL D6: IC ₅₀ = 847.4 ng/mL	[37]
<i>A. olivaceum</i>	Brazil	–	EtOH Hexane-AcOEt	<i>In vivo</i> / <i>P. berghei</i>	Inactive	[67]
<i>A. olivaceum</i>	Brazil	Leaves, trunk wood, trunk bark	DCM and EtOH	<i>In vitro</i> / <i>P. falciparum</i> W2 and 3D7	W2: IC ₅₀ = 5.0 to 7 µg/mL 3D7: IC ₅₀ = 5.0 to 25.5 µg/mL	[27]
<i>A. olivaceum</i>	Brazil	–	Olivacine (4)	<i>In vitro</i> / <i>P. falciparum</i> 3D7 and K1	K1: IC ₅₀ = 1.4 µM 3D7: IC ₅₀ = 1.2 µM Citotoxicity murine macrophages: CC ₅₀ > 4.1 × 10 ² µg/mL SI: K1 > 2.9 × 10 ² 3D7 > 3.4 × 10 ²	[48]
<i>A. parvifolium</i>	Brazil	Trunk bark	EtOH extract, Uleine (3)	<i>In vivo</i> / <i>P. berghei</i> <i>In vitro</i> / <i>P. falciparum</i> W2 and 3D7 Uleine W2 and 3D7	<i>In vivo</i> : active (50 mg/kg/day) W2: IC ₅₀ = 32.8 µg/mL 3D7: IC ₅₀ = 20.5 µg/mL W2: IC ₅₀ = 0.75 µg/mL (2.81 µM) 3D7: IC ₅₀ = 11.90 µg/mL (32.69 µM)	[27]
<i>A. polyneuron</i>	Brazil	Trunk bark	Total alkaloids	<i>In vitro</i> / <i>P. cathemerium</i>	Active	[33]
<i>A. pyrifolium</i>	Bolivia	Stem bark	EtOH-H ₂ O (7:3)	<i>In vitro</i> / <i>P. falciparum</i> F32 strain FBIT	Inactive	[40, 81]
<i>A. pyrifolium</i>	Bolivia	Stem bark	*Eight alkaloids	<i>In vitro</i> / <i>P. falciparum</i> FcM29 and Nigerian strain	FcM29: IC ₅₀ = 3.2 to 28.5 µM Nigerian strain: IC ₅₀ = 5.1 to 22.6 µM	[42] [41]
<i>A. quebrachoblanco</i>	Argentina	Trunk bark	Total alkaloids	Human	Active	[35, 80]
<i>A. quebrachoblanco</i>	Argentina	Trunk bark	EtOH	Human	Inactive	[34, 80]
<i>A. quebrachoblanco</i>	Bolivia	Leaves, trunk bark	EtOH-H ₂ O (7:3)	<i>In vitro</i> / <i>P. falciparum</i> F32 strain FBIT	Trunk bark extract F32: IC ₅₀ = 3.9 µg/mL FBIT: IC ₅₀ = 1.22 mg/mL Leaf extract: inactive	[38] <i>continued</i>

Table 2 Continued

Aspidosperma species	Country of collection	Part of the plant	Extracts Compounds	Bioassay Plasmodium species/ strains	Results	References
<i>A. ramiflorum</i>	Brazil	Leaves, trunk wood, trunk bark	DCM and EtOH	<i>In vitro/P. falciparum</i> W2 and 3D7	W2: IC ₅₀ = 19.7 to 36.5 µg/mL 3D7: IC ₅₀ = 1.0 to 48.0 µg/mL	[27]
<i>A. ramiflorum</i>	Brazil	–	–	<i>In vitro/P. falciparum</i> W1 <i>In vivo/P. berghei</i>	W2: IC ₅₀ = 11 to 40 µg/mL <i>In vivo</i> : partial activity	[82]
<i>A. rigidum</i>	Peru (Remo caspi)	Trunk bark	EtOH	<i>In vitro/P. falciparum</i> 3D7	IC ₅₀ < 10 µg/mL SI human lymphocyte inhibition 75%	[39]
<i>A. spruceanum</i>	Brazil	Leaves, trunk wood, trunk bark	DCM and EtOH	<i>In vitro/P. falciparum</i> W2 and 3D7	W2: IC ₅₀ = < 6.0 to 65.0 µg/mL 3D7: IC ₅₀ = < 6.0 to > 100 µg/mL	[27]
<i>A. tomentosum</i>	Brazil	Leaves, trunk wood, seeds, fruits	EtOH	<i>In vitro/P. falciparum</i> W2 and 3D7	W2: IC ₅₀ = 20.5 to 26.5 µg/mL 3D7: IC ₅₀ = 3.0 to 38.5 µg/mL	[27]
<i>A. ulei</i>	Brazil	Leaf, bark, trunk wood, root wood, root bark	Five alkaloids**	<i>In vitro/P. falciparum</i> K1	K1: IC ₅₀ = 16.7 to > 176.0 µM Cytotoxicity NIH3T3 murine fibroblasts	[47]
<i>A. vargasii</i>	Brazil	Trunk bark	Ellipticine (1)	<i>In vitro/P. falciparum</i> K1	IC ₅₀ = 0.073 µM (0.018 µg/mL)	[45]
<i>A. vargasii</i>	Brazil	Bark	Ellipticine (1)	<i>In vitro/P. falciparum</i> 3D7 and K1	K1: IC ₅₀ = 0.81 µM 3D7: IC ₅₀ = 0.35 µM Cytotoxicity murine macrophages: CC ₅₀ > 4.1 × 10 ² µg/mL SI: K1 > 5.0 × 10 ² 3D7 > 1.2 × 10 ³	[48]
				<i>In vivo/P. berghei</i>	<i>In vivo</i> : active (50 mg/kg/day)	

* Aspidospermidine (6), 10-methoxy-aspidospermidine (7), N-formil-aspidospermidine (8), vallesine (9), (-)-aspidospermine (10), demethoxyaspidospermine (11), palosine (12), haplocine (13). ** 20-*epi*-dasycarpidone (3), 3,4,5,6-tetrahydro-β-yohimbine, 20(*E*)-*nor*-subincanadine E, 19*E*-hunteracine, 12-hydroxy-*N*-acetyl-21(*N*)-dehydroplumeran-18-oic acid

of plant species for antimalarial screening. Negative results were described for *in vivo* assays of *A. nitidum* crude extracts [32,44] (Table 2). These results are surprising as *A. nitidum* is one of the most frequently cited *Aspidosperma* species of those used to treat malaria/fevers in Brazil. Therefore, this species deserves further investigation with a focus on the alkaloids.

Interestingly, a strong interest in *Aspidosperma* species has resumed in the past two decades, and several species whose phytochemistry was intensively investigated during 1960–1980 have been reexamined, leading to the isolation of several known indole alkaloids. Several of these alkaloids have been evaluated for various biological/pharmacological effects [30]. A total of 20 *Aspidosperma* alkaloids have been assayed for antimalarial activity and have been isolated by Brazilian researchers, one from *A. desmanthum* [45], one from *A. vargasii* [45], one from *A. parvifolium* [27,46], five from *A. ulei* [47], and one from *A. olivaceum* [48]; additionally, a French-Bolivian group has evaluated 11 alkaloids from *A. megalocarpon* and *A. pyrifolium* [41].

Ellipticine (1) and aspidocarpine (2) (Fig. 1) have been isolated from the trunk bark of *A. vargasii* and *A. desmanthum*, respectively. Both of these species have been collected at the Ducke Reserve in Manaus, Amazonas state, Brazil, and they have shown remarkable *in vitro* activity against the multidrug-resistant K1 strain of *P. falciparum* (IC₅₀ = 73 nM and 19 nM, respectively) [45]. Ellipticine (1), a pyridocarbazol alkaloid, was originally isolated from *Ochrosia elliptica* (an Australian evergreen shrub) and occurs in other genera of the family Apocynaceae, such as *Aspidosperma* and *Bleekeria* [49–51]. Ellipticine (1) and several related syn-

thetic derivatives are highly cytotoxic to human cancer cell lines, and elliptinium acetate (9-OH-NME, Celliptium®) is an antineoplastic agent currently used in the treatment of metastatic breast cancer [49–51]. Recently, ellipticine (1) and olivacine (5) (Fig. 1) were assayed against *P. falciparum* strains K1 and 3D7. Olivacine disclosed lower *in vitro* antimalarial activity than ellipticine, and low cytotoxicity for both agents was observed against murine macrophages. Ellipticine was also more active than olivacine in an evaluation of the effects in *P. berghei*-infected mice. Remarkably, 100% parasitemia suppression was observed at an oral dose of 50 mg/kg/day (four days), and no mortality or other signs of toxicity were reported [48]. Ellipticine (1) and its derivatives were described as the most active compounds, with IC₅₀ values < 1 µg/mL (range 0.08 to 0.47 µg/mL), in a series of 184 randomly selected compounds belonging to several classes of natural products, either from plants or from marine organisms, or prepared as intermediates during the synthesis or semisynthesis of isolated products that were assayed against the *P. falciparum* FcM29 strain [52].

A total of 12 indole alkaloids were recently isolated from the EtOH extracts of various parts of *A. ulei* Markgr. Only five of these alkaloids were assayed for antimalarial activity against the multidrug-resistant K1 strain of *P. falciparum*. 20-*Epi*-dasycarpidone (3) (Fig. 1) showed moderate activity (IC₅₀ 4.5 µg/mL, 16.7 µM), whereas 3,4,5,6-tetrahydro-β-yohimbine, 19*E*-hunteracine, 20(*E*)-*nor*-subincanadine E, and 12-hydroxy-*N*-acetyl-21(*N*)-dehydroplumeran-18-oic acid were inactive. Two of the known alkaloids isolated, uleine and olivacine, have previously




Fig. 1 Chemical structures of antimalarial alkaloids from *Aspidosperma* species: ellipticine from *A. vargasii* [45], aspidocarpine from *A. desmanthum* [45], uleine from *A. parvifolium* [46], olivacine from *A. olivaceum* [47] and 20-Epi-dasycarpidone from *A. ulei* [47].

been shown to be active [46,48]. The compounds were evaluated against NIH3T3 murine fibroblasts and showed no cytotoxicity up to a concentration of 50 $\mu\text{g}/\text{mL}$ [47].

Eleven known aspidospermane alkaloids (● Fig. 2) were isolated from *A. pyrifolium* and *A. megalocarpon*, both collected in Bolivia [42,53] and both also occurring in Brazil [24–26]. Three of the active compounds, aspidospermine (6), 10-methoxy-aspidospermidine (7), and *N*-formyl-aspidospermine (8) (● Fig. 2), were shown to be less cytotoxic to NIH 3 T3 cells (human fibroblasts), with calculated SI values of 22.7, 15.6, and 8.3, respectively [41]. The *in vitro* antiplasmodial activity of *A. megalocarpon* bark extract against D2 and F32 strains of *P. falciparum* has been reported; subsequently, a very low SI (< 1) was reported in relation to cytotoxicity assays in U-937 human promonocytic cells [36]. Uleine (4) (● Fig. 1) was first isolated from *A. ulei* in 1959 [54,55]. It is frequently found in species, e.g., *A. australe*, *A. dasycarpon*, *A. eburneum*, *A. excelsum*, *A. formosanum*, *A. gilbertii*, *A. gomezianum*, *A. multiflorum*, *A. nigricans*, *A. olivaceum*, and *A. parvifolium* [29], as well as in other apocynaceous representatives such as *Himathanthus lancifolius* [56].

A recent phytochemical reinvestigation of *A. parvifolium* trunk bark afforded four known indole alkaloids, uleine (4) (● Fig. 1), *epi*-uleine, apparicine, and *N*-demethyluleine [57]. A bioguided fractionation of the trunk bark alkaloidal fraction demonstrated the *in vitro* activity of uleine against chloroquine-resistant *P. falciparum* (W2), with an IC_{50} at the ng level, at least against the W2 clone (● Table 2) [27,46]. The antiplasmodial activity of uleine (4), an aspidospermane indole alkaloid, may be related to the inhibition of heme polymerization to give hemozoin, as demonstrated by its *in vivo* effect in the food vacuole of chloroquine-resistant *P. falciparum* (W2 clone) monitored by confocal microscopy [46].

Uleine (4) (● Fig. 1) has been evaluated for cytotoxicity against the NCI 60 cancer cell line panel and was inactive in all of them [58], reinforcing its high potential as a leading antimalarial compound.

In summary, crude extracts, alkaloidal extracts, and isolated alkaloids from 19 *Aspidosperma* species have been evaluated for antimalarial activity, and positive results have been observed for most of them. These results disclose the high potential of *Aspidosperma* species as sources of antimalarial alkaloids. A substantial literature search, approximately 1031 articles, describes the antiplasmodial activity of plant extracts, but very few of these re-




Fig. 2 Antimalarial alkaloids from *A. megalocarpon* and *A. pyrifolium* [41].

ports include the effect of pure constituents [14], a requirement for the development of new drugs. The development of new drugs is lengthy and extremely costly. However, if effective, safe, and locally produced phytomedicines are the goal, a more direct and less expensive route might certainly be pursued [5,59,60].

Conclusions

Our literature survey has identified 24 *Aspidosperma* species reported to be used to treat malaria/fevers. Of these, a total of 19 species have had extracts and/or isolated alkaloids evaluated for antimalarial activity by different assays, showing positive results. Only 20 of more than 200 known indole alkaloids from *Aspidosperma* species have been assayed for antimalarial activity, and variable levels of parasite inhibition have been observed. Among the assayed *Aspidosperma* alkaloids, uleine (4) (● Fig. 1) appears the most promising as an antimalarial because it has shown good *in vitro* activity [27,46] and no cytotoxicity in several human cancer cell lines [58]. These findings have motivated a patent in Brazil [61]. However, recent results on the *in vivo* evaluation of ellipticine (1) (● Fig. 1) appear to place this alkaloid in the leading position among natural antimalarials because it has, remarkably, shown 100% parasitemia suppression at an oral dose of 50 mg/kg/day and no signs of mortality or toxicity [48]. Our review also demonstrates a need for molecular genetic studies to facilitate the identification and the differentiation of *Aspidosperma* species and, therefore, to clarify the controversial question of synonyms. The high chemical diversity of alkaloids from *Aspidosperma* species and the small number of these alkaloids that have been as-

sayed for antimalarial activity coupled with the traditional use of several species of this taxon to treat malaria in Brazil as well as in other Meso- and South American countries make further investigations of the plants in this genus of great interest in the quest for natural antimalarial products. There is a need to isolate plant constituents to evaluate their pharmacological profile, to further pursue studies of synergistic effects, and to make these constituents available for use as markers in the standardization of extracts to be used in the development of effective, safe, and low-cost phytomedicines. These phytomedicines would then be available to Amazonian people, who inhabit the region where many of the *Aspidosperma* species occur and that have the highest incidence of malaria in Meso- and South America.

Acknowledgements

▼ We thank CNPq and FAPEMIG for financial support (PRONEX CNPq Process 555655/2009–1 and FAPEMIG Process CDS APQ 01129–10) and a doctoral fellowship (CNPq to RCP). Additionally, we thank Dr. Luzia H. Carvalho, CPqRR/FIOCRUZ, Belo Horizonte, Brazil, for helpful discussions.

Conflict of Interest

▼ The authors declare that they do not have any conflict of interest in reference to the content of this article.

References

- 1 World Health Organization. World Malaria Report 2009. Available at http://whqlibdoc.who.int/publications/2009/9789241563901_eng.pdf. Accessed November 25, 2009.
- 2 World Health Organization. World Malaria Report 2008. Available at http://whqlibdoc.who.int/publications/2008/9789241563697_eng.pdf. Accessed November 22, 2009.
- 3 Oliveira-Ferreira J, Lacerda MVG, Brasil P, Ladislau JLB, Tauil PL, Daniel-Ribeiro CT. Malaria in Brazil: an overview. *Malar J* 2010; 9: 115–130
- 4 World Health Organization. World Malaria Report 2011. Available at http://www.who.int/malaria/world_malaria_report_2011/9789241564403_eng.pdf. Accessed January 13, 2011.
- 5 Oliveira AB, Dolabela MF, Braga FC, Jacome RL, Varotti FP, Povoá MM. Plant-derived antimalarial agents: new leads and efficient phytomedicines. Part I. Alkaloids. *An Acad Bras Cienc* 2009; 81: 715–740
- 6 Castellanos JRG, Prieto JM, Heinrich M. Red Lapacho (*Tabebuia impetiginosa*)—a global ethnopharmacological commodity? *J Ethnopharmacol* 2009; 121: 1–13
- 7 Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25 years. *J Nat Prod* 2007; 70: 461–477
- 8 Ginsburg H, Deharo E. A call for using natural compounds in the development of new antimalarial treatments – an introduction. *Malar J* 2011; 10 (Suppl. 1): S1
- 9 Batista R, Junior AJS, Oliveira AB. Plant-derived antimalarial agents: new leads and efficient phytomedicines. Part II. Non-alkaloidal natural products. *Molecules* 2009; 14: 3037–3072
- 10 Verpoorte R, Choi Y, Mustafa N, Kim H. Metabolomics: back to basics. *Phytochem Rev* 2008; 7: 525–537
- 11 Wagner H. Natural products chemistry and phytomedicine research in the new millennium: new developments and challenges. *Arkivoc* 2004; VII: 277–284
- 12 Ulrich-Merzenich G, Zeitler H, Jobst D, Panek D, Vetter H, Wagner H. Application of the “-Omic-” technologies in phytomedicine. *Phytomedicine* 2007; 14: 70–82
- 13 Verpoorte R, Choi YH, Kim HK. Ethnopharmacology and systems biology: a perfect holistic match. *J Ethnopharmacol* 2005; 100: 53–56
- 14 Deharo E, Ginsburg H. Analysis of additivity and synergism in the antiplasmodial effect of purified compounds from plant extracts. *Malar J* 2011; 10 (Suppl. 1): S5
- 15 Lorenzi H. Árvores Brasileiras: manual de identificação e cultivo de plantas arbóreas nativas do Brasil. 2nd edition. São Paulo: Plantarum; 1998: 365
- 16 Perez-Arbeláez E. Plantas útiles de Colombia. 5th edition. Bogotá: Fondo Fen Colombia; 1996: 831
- 17 Pio-Corrêa M. Dicionário de plantas úteis do Brasil e das exóticas cultivadas. Rio de Janeiro: Ministério da Agricultura; 1978: 687
- 18 Rizzini CT. Árvores e madeiras úteis do Brasil: manual de dendrologia brasileira. 2nd edition. São Paulo: E. Blücher; 2000: 312
- 19 Schultes RE, Raffauf RF. The healing forest: medicinal and toxic plants of the northwest Amazonia. 1st edtion. Portland: Dioscorides Press; 2003: 500
- 20 Koch I, Rapini A, Kinoshita LS, Simões AO, Spina AP. Apocynaceae. Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. Available at <http://reflora.jbrj.gov.br/jabot/floradobrasil/FB21892>. Accessed December 8, 2013.
- 21 Sennblad B, Bremer B. Classification of Apocynaceae s.l. according to a new approach combining Linnaean and phylogenetic taxonomy. *Syst Biol* 2002; 51: 389–409
- 22 Woodson RE. Studies in the Apocynaceae. VIII. An Interim Revision of the Genus *Aspidosperma* Mart. & Zucc. *Ann Mo Bot Gard* 1951; 38: 119–206
- 23 Marcondes-Ferreira W, Kinoshita L. Uma nova divisão infragenerica para *Aspidosperma* Mart. (Apocynaceae). [A new infrageneric division on *Aspidosperma* Mart. (Apocynaceae)]. *Rev Brasil Bot* 1996; 19: 203–214
- 24 Lista de Espécies da Flora do Brasil. Available at <http://floradobrasil.jbrj.gov.br/jabot/listaBrasil/PrincipalUC/PrincipalUC.do>. Accessed November 29, 2011.
- 25 Tropicos. Missouri Botanical Garden – Tropicos Specimen Data Base. Available at <http://www.tropicos.org/NameSearch.aspx?name=Aspidosperma&common name=>. Accessed November 30, 2011.
- 26 The Plant List. (2010). Version 1. Available at <http://www.theplantlist.org/tp1/search?q=Aspidosperma>. Accessed November 29, 2011.
- 27 Dolabela MF. Atividade antiplasmodica e citotoxicidade de *Esenbeckia febrifuga* (A.St-Hil.) Juss. ex Mart. (Rutaceae) e de espécies do gênero *Aspidosperma* (Apocynaceae) [dissertation]. Belo Horizonte: Universidade Federal de Minas Gerais; 2007
- 28 Mesquita M, Grellier P, Mambu L, De Paula J, Espindola L. In vitro antiplasmodial activity of Brazilian Cerrado plants used as traditional remedies. *J Ethnopharmacol* 2007; 110: 165–170
- 29 Pereira MM, Jácome R, Alcântara A, Alves RB, Raslan DS. Alcalóides indólicos isolados de espécies do gênero *Aspidosperma* (Apocynaceae). *Quim Nova* 2007; 30: 970–983
- 30 Oliveira V, Freitas M, Mathias L, Braz-Filho R, Vieira I. Atividade biológica e alcalóides indólicos do gênero *Aspidosperma* (Apocynaceae): uma revisão. *Rev Bras Pl Med* 2009; 11: 92–99
- 31 Trager W, Jensen JB. Human malaria parasites in continuous culture. *Science* 1976; 193: 673–675
- 32 Becker ER. Report on 35 drugs and three plant materials tested against *Plasmodium lophurae* in the white Pekin duck. *Iowa State Coll J Sci* 1949; 23: 189–194
- 33 Wasicky R, Unti O, Barbieri E. Quinine and alkaloids in Brazil. *An Fac Farm Odontol Univ São Paulo* 1942; 3: 137
- 34 Domínguez J. Malaria treatment with *Aspidosperma quebracho blanco*. *Rev Farm (Buenos Aires)* 1932; 73: 82
- 35 Loizaga N, Sagastume L. Malaria treatment with Quechuol-Domínguez. *Sem Med Buenos Aires* 1935; 2: 562–566
- 36 Weniger B, Robledo S, Arango GJ, Deharo E, Aragon R, Muñoz V, Callapa J, Lobstein A, Anton R. Antiprotozoal activities of Colombian plants. *J Ethnopharmacol* 2001; 78: 193–200
- 37 Cabral JA, McChesney JD, Milhous WK. A new antimalarial quassinoid from *Simaba guianensis*. *J Nat Prod* 1993; 56: 1954–1961
- 38 Bourdy G, Oporto P, Gimenez A, Deharo E. A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part VI. Evaluation of the antimalarial activity of plants used by Isoceno-Guarani Indians. *J Ethnopharmacol* 2004; 93: 269–277
- 39 Kvist LP, Christensen SB, Rasmussen HB, Mejia K, Gonzalez A. Identification and evaluation of Peruvian plants used to treat malaria and leishmaniasis. *J Ethnopharmacol* 2006; 106: 390–402
- 40 Brandão MG, Grandi TS, Rocha EM, Sawyer DR, Krettlí AU. Survey of medicinal plants used as antimalarials in the Amazon. *J Ethnopharmacol* 1992; 36: 175–182

- 41 *Mitaine-Offer AC, Sauvain M, Valentin A, Callapa J, Mallie M, Zeches-Hanrot M.* Antiplasmodial activity of *Aspidosperma* indole alkaloids. *Phytomedicine* 2002; 9: 142–145
- 42 *Mitaine AC, Mesbah K, Richard B, Petermann C, Arrazola S, Moretti C, Zeches-Hanrot M, Men-Olivier LL.* Alkaloids from *Aspidosperma* species from Bolivia. *Planta Med* 1996; 62: 458–461
- 43 *Dolabela MF, Oliveira SG, Peres JM, Nascimento JM, Póvoa MM, Oliveira AB.* *In vitro* antimalarial activity of six *Aspidosperma* species from the state of Minas Gerais (Brazil). *An Acad Bras Cienc* 2012; 84: 899–910
- 44 *Carvalho LH, Brandão MG, Santos-Filho D, Lopes JL, Krettli AU.* Antimalarial activity of crude extracts from Brazilian plants studied *in vivo* in *Plasmodium berghei*-infected mice and *in vitro* against *Plasmodium falciparum* in culture. *Braz J Med Biol Res* 1991; 24: 1113–1123
- 45 *Andrade-Neto VF, Pohlit AM, Pinto AC, Silva EC, Nogueira KL, Melo MR, Henrique MC, Amorim RC, Silva LF, Costa MR, Nunomura RC, Nunomura SM, Alecrim WD, Alecrim MG, Chaves FC, Vieira PP.* *In vitro* inhibition of *Plasmodium falciparum* by substances isolated from Amazonian antimalarial plants. *Mem Inst Oswaldo Cruz* 2007; 102: 359–365
- 46 *Oliveira AB, Dolabela MF, Póvoa MM, Santos CAM, Varotti FP.* Antimalarial activity of uleïn and proof of its action on the *Plasmodium falciparum* digestive vacuole. *Mal J* 2010; 9 (Suppl. 2): O9
- 47 *Torres ZES, Silveira ER, Rocha e Silva LF, Lima ES, Vasconcellos MC, Andrade Uchoa DE, Filho RB, Pohlit AM.* Chemical composition of *Aspidosperma ulei* Markgr. and antiplasmodial activity of selected indole alkaloids. *Molecules* 2013; 18: 6281–6297
- 48 *Rocha e Silva LF, Montoia A, Amorim RC, Melo MR, Henrique MC, Nunomura SM, Costa MR, Andrade Neto VF, Costa DS, Dantas G, Lavrado J, Moreira R, Paulo A, Pinto AC, Tadei WP, Zacardi RS, Eberlin MN, Pohlit AMLF.* Comparative *in vitro* and *in vivo* antimalarial activity of the indole alkaloids ellipticine, olivacine, cryptolepine and a synthetic cryptolepine analog. *Phytomedicine* 2012; 20: 71–76
- 49 *Cirino JJV, Belletato P, Dantas SO, Ribeiro L, Ferreira GR, Santos HF.* Estrutura e propriedades de elipticinas. *Quim Nova* 2005; 28: 30–36
- 50 *Cragg GM, Newman DJ.* Plants as a source of anti-cancer agents. *J Ethnopharmacol* 2005; 100: 72–79
- 51 *Mans DR, da Rocha AB, Schwartsmann G.* Anti-cancer drug discovery and development in Brazil: targeted plant collection as a rational strategy to acquire candidate anti-cancer compounds. *Oncologist* 2000; 5: 185–198
- 52 *Passeram C, Salery M, Soh PN, Linas MD, Ahond A, Poupat C, Benoit-Vical F.* Indole and aminoimidazole moieties appear as key structural units in antiplasmodial molecules. *Phytomedicine* 2011; 18: 1118–1125
- 53 *Mitaine AC, Weniger B, Sauvain M, Lucumi E, Aragon R, Zeches-Hanrot M.* Indole alkaloids from the trunk bark of *Aspidosperma megalocarpon*. *Planta Med* 1998; 64: 487
- 54 *Büchi G, Warnhoff E.* The structure of uleine. *J Am Chem Soc* 1959; 81: 4433–4434
- 55 *Joule J.* The uleine-ellipticine-vallesamine group. In: Saxton JE, editor. *The chemistry of heterocyclic compounds, indoles: the monoterpene indole alkaloids*. 1st edition. Cleveland: John Wiley & Sons; 2009: 886
- 56 *Nardin JM, de Souza WM, Lopes JF, Florao A, de Moraes Santos CA, Welfort-Santos AM.* Effects of *Himatanthus lancifolius* on human leukocyte chemotaxis and their adhesion to integrins. *Planta Med* 2008; 74: 1253–1258
- 57 *Jácome R, Oliveira A, Raslan DS, Wagner H.* Estudo químico e perfil cromatográfico das cascas de *Aspidosperma parvifolium* A. DC. (“pau-perreira”). *Quim Nova* 2004; 27: 897–900
- 58 *PubChem Substance.* Uleine-Substance Summary. Available at <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?sid=115962>. Accessed August 11, 2013.
- 59 *Willcox ML, Bodeker G.* Traditional herbal medicines for malaria. *BMJ* 2004; 329: 1156–1159
- 60 *Wright CW.* Recent developments in research on terrestrial plants used for the treatment of malaria. *Nat Prod Rep* 2010; 27: 961–968
- 61 *Oliveira AB, Dolabela MF, Rocha FD, Gomes FMA, Neiva RMT, Jácome RLRP.* Extrato e fração padronizados de cascas de *Aspidosperma parvifolium* e/ou uleine e sua composição farmacêutica. Filing date: Dec 23, 2009. PI0905584–3. Assignee: Universidade Federal de Minas Gerais – UFMG. Belo Horizonte, MG, Brazil
- 62 *Araújo jr. JX, Antheaume C, Trindade RCP, Schmitt M, Bourguignon JJ, Sant’Ana AEG.* Isolation and characterisation of the monoterpene indole alkaloids of *Aspidosperma pyrifolium*. *Phytochem Rev* 2007; 6: 183–188
- 63 *Grenand P, Moretti C, Jacquemin H, Prévost M.* Pharmacopées traditionnelles en Guyane Créoles, Polikur, Wayapi. 1st edition. Montpellier: IRD Éditions; 2006: 816
- 64 *Barbosa WLR, Tavares ICC, Soares DC.* Alcaloides de *Aspidosperma auriculatum* Standl. *Braz J Pharmacogn* 2003; 13: 6–8
- 65 *Pittier H.* Manual de las plantas usuales de Venezuela y su suplemento. 2nd edition. Caracas: Fundación Eugenio Mendoza; 1978: 620
- 66 *Quignard ELJ, Martin Pohlit A, Nunomura SM, Da Silva Pinto AC, Mustafa dos Santos EV, Reis de Moraes SK, Alecrim AM, Da Silva Pedroso AC, Brito Cyrino BR, Melo CS.* Screening of plants found in Amazonas state for lethality towards brine shrimp *Artemia franciscana*. *Acta Amazonica* 2003; 33: 93–104
- 67 *Brandão MGL, Botelho MGA, Krettli AU.* Quimioterapia experimental antimalárica com produtos naturais: I. Uma abordagem mais racional?/Experimental antimalarial chemotherapy with natural products: I. A more rational approach? *Cienc Cult* 1985; 37: 1152–1163
- 68 *Milliken W.* Traditional anti-malarial medicine in Roraima, Brazil. *Econ Bot* 1997; 51: 212–237
- 69 *Mejía CK, Rengifo SE.* Plantas medicinales de uso popular en la Amazonía Peruana. 1st edition. Lima: Tarea Asociación Gráfica Educativa; 1995: 286
- 70 *Pérez D.* Etnobotánica medicinal y biocidas para malaria en la región Ucayali. *Folia Amazonica* 2002; 13: 2
- 71 *Duke JA, Vasquez R.* Amazonian ethnobotanical dictionary. 1st edition. Florida: CRC Press; 1994: 224
- 72 *Milliken W, Albert B.* The use of medicinal plants by the Yanomami Indians of Brazil. *Econ Bot* 1996; 50: 10–25
- 73 *DeFilippis RA, Maina SL, Crepin J.* Medicinal Plants of the Guianas (Guyana, Surinam, French Guiana). Part 1 Acanthaceae – Combretaceae. Available at http://botany.si.edu/bdg/medicinal/Medicinal_plants_master.pdf. Accessed July 28, 2011.
- 74 *Almeida MR, Lima JA, dos Santos NP, Pinto AC.* Pereirina: o primeiro alcalóide isolado no Brasil? *Braz J Pharmacogn* 2009; 19: 942–952
- 75 *Botsaris AS.* Plants used traditionally to treat malaria in Brazil: the archives of Flora Medicinal. *J Ethnobiol Ethnomed* 2007; 3: 18–19
- 76 *Milliken W, Albert B.* The use of medicinal plants by the Yanomami Indians of Brazil, Part II. *Econ Bot* 1997; 51: 264–278
- 77 *Del Vitto LA, Petenatti E, Petenatti M.* Recursos herbolarios de San Luis (República Argentina). Primera parte: Plantas nativas. *Multequina* 1997; 6: 49–66
- 78 *Freitas ADD.* Aspectos tecnológicos e morfoanatómicos de sementes maduras, plântulas e plantas jovens de *Aspidosperma spruceanum* Benth. Ex Mull. Arg. (Apocynaceae) [dissertation]. Belém: Universidade Federal Rural da Amazônia e Museu Paraense Emílio Goeldi; 2008
- 79 *Di Stasi LC.* Plantas medicinais na Amazônia e na Mata Atlântica. 2nd edition. São Paulo: UNESP; 2002: 592
- 80 *Mariath IR, Falcão HS, Barbosa-Filho JM, Sousa LCF, Tomaz ACA, Batista LM, Diniz MFFM, Athayde-Filho PF, Tavares JF, Silva MS.* Plants of the American continent with antimalarial activity. *Braz J Pharmacogn* 2009; 19: 158–191
- 81 *Muñoz V, Sauvain M, Bourdy G, Callapa J, Bergeron S, Rojas I, Bravo JA, Balderrama L, Ortiz B, Gimenez A, Deharo E.* A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part I. Evaluation of the antimalarial activity of plants used by the Chacobo Indians. *J Ethnopharmacol* 2000; 69: 127–137
- 82 *Krettli LG, de Paula RC, Varotti FP, Pereira MM, Oliveira AJB, Krettli AU.* Antimalarial activity of extracts from *Aspidosperma* plants tested *in vitro* against *Plasmodium falciparum* drug resistant parasites and in animal models. XXIV Meeting of the SBPz. Águas de Lindóia, São Paulo: Brazilian Society of Protozoology; 2008