Duck River

Conservation Opportunity Area



The Duck River flows 284 miles from its headwaters in the Eastern Highland Rim, draining much of middle Tennessee, before entering the Tennessee River. The Duck River has a high diversity of freshwater animals, with 151 species of fish, 60 species of freshwater mussels, and 22 species of aquatic snails. Several of these species are considered rare and endangered.

The Tennessee Chapter of The Nature Conservancy has developed and implemented numerous habitat improvement projects along the Duck River in collaboration with TWRA and many other partners. These projects include stabilizing stream banks through the planting of trees and shrubs; providing landowner incentive funds to construct cattle fencing, water stations, and creek crossings; and assisting state and federal agencies with freshwater mussel recovery efforts.

The highlands surrounding the headwaters of the Duck River, as well as the forested areas remaining along the river, are important for maintaining water quality and temperature. Stream bank erosion can impact water quality and degrade habitat. The Duck River provides drinking water for over 250,000 people today, and parts of the watershed are experiencing urbanization pressures. Collaborative land and water use planning will be critical to protecting the Duck River in coming years.


In addition, large areas of cedar glades and limestone barren habitats occur throughout this COA. These glades and barrens are susceptible to loss due to natural succession from lack of natural fire and management. Many of these glades are home to rare plant species and also associated with caves and sinkholes. These habitats are all susceptible to illegal dumping and incompatible recreational activities.

Public Lands within the COA

Bedford Lake, Columbia Greenway, Duck River State Scenic River, Henry Horton State Park, Normandy Fish Hatchery, Wilson School Road Forest & Cedar Glade, Yanahli WMA



Draining much of middle Tennessee, the Duck River is considered to be one of the most biologically diverse rivers in North America.



Top to bottom: Mussel diversity, Dan Hubbs; Big Falls, Duck River - Brian Stansberry/next page: The Nashville Breadroot only found in Cedar Glades habitats. - Jason Ferrell

Duck River Highlands and Headwaters Conservation Opportunity Area


Partnerships

TWRA will continue its collaborations with The Nature Conservancy and others to conserve the natural resources of the Duck River. TWRA will develop and foster partnerships with the U.S. Fish and Wildlife Service, Natural Resources Conservation Service, U.S. Army Corps of Engineers, Tennessee Valley Authority, Tennessee Dept. of Environment and Conservation, universities, and private landowners.



| Desired Change | Units |
|---|----------|
| Use government-funded incentive programs to encourage private landowners to restore habitats and protect water quality using Best Management Practices | number |
| Restore pastures and agricultural lands to native ecosystems, such as forests and grasslands. | acres |
| Restore riparian habitat and physical stream habitat in priority locations throughout the watershed | miles |
| Conduct research on life history and ecological needs of target species to establish guidelines for determining population viability. | projects |
| Conduct scientific surveys to discern the geographical distributions of lesser known target species. | projects |
| Increase acres of habitat under permanent conservation management. | acres |
| Improve population viability of native freshwater mussel species. | animals |
| Increase the number of water withdrawal plans created and implemented by appropriate state and local agencies and utilities to protect and improve instream flow for native species | plans |
| Improve forest habitat through development of new forestry Best Management Practice Standards. | assists |
| Improve and protect ecologically important instream flows by engaging in reservoir operational reviews with TVA. | reviews |

To learn more about the Duck River Watershed Conservation Opportunity Area, please contact: Josh Campbell, Wildlife Diversity Coordinator, Region 2 (615) 781-6626



| | DUCK RIVER COA GCN SPECIES (TN) |
|--------------|---|
| Taxa Group | Scientific & Common names |
| Amphibian | Ambystoma barbouri (Streamside Salamander) |
| Bat | Myotis grisescens (Gray Bat) |
| Bat | Myotis septentrionalis (Northern Myotis) |
| Bat | Myotis sodalis (Indiana Bat) |
| Bat | Perimyotis subflavus (Tri-colored Bat) |
| Bird | Accipiter striatus (Sharp-shinned Hawk) |
| Bird | Ammodramus henslowii (Henslow's Sparrow) |
| Bird | Ammodramus savannarum (Grasshopper Sparrow) |
| Bird | Anas rubripes (American Black Duck) |
| Bird | Aquila chrysaetos (Golden Eagle) |
| Bird | Asio flammeus (Short-eared Owl) |
| Bird | Bartramia longicauda (Upland Sandpiper) |
| Bird | Botaurus lentiginosus (American Bittern) |
| Bird | Calidris mauri (Western Sandpiper) |
| Bird | Calidris pusilla (Semipalmated Sandpiper) |
| Bird | Caprimulgus carolinensis (Chuck-will's-widow) |
| Bird | Caprimulgus vociferus (Whip-poor-will) |
| Bird | Chaetura pelagica (Chimney Swift) |
| Bird | Chordeiles minor (Common Nighthawk) |
| Bird | Circus cyaneus (Northern Harrier) |
| Bird | Cistothorus platensis (Sedge Wren) |
| Bird | Colinus virginianus (Northern Bobwhite) |
| Bird | Egretta caerulea (Little Blue Heron) |
| Bird | Empidonax minimus (Least Flycatcher) |
| Bird | Empidonax traillii (Willow Flycatcher) |
| Bird | Euphagus carolinus (Rusty Blackbird) |
| Bird | Falco peregrinus (Peregrine Falcon) |
| Bird | Geothlypis Formosa (Kentucky Warbler) |
| Bird | Grus americana (Whooping Crane) |
| Bird | Haliaeetus leucocephalus (Bald Eagle) |
| Bird | Helmitheros vermivorum (Worm-eating Warbler) |
| Bird | Hylocichla mustelina (Wood Thrush) |
| Bird | Icteria virens (Yellow-breasted Chat) |
| Bird | Icterus spurius (Orchard Oriole) |
| Bird | Ictinia mississippiensis (Mississippi Kite) |
| Bird | Ixobrychus exilis (Least Bittern) |
| Bird | Lanius Iudovicianus (Loggerhead Shrike) |
| Bird | Limnothlypis swainsonii (Swainson's Warbler) |
| Bird | Melanerpes erythrocephalus (Red-headed Woodpecker) |
| Bird | Nycticorax nycticorax (Black-crowned Night-heron) |
| Bird | Parkesia motacilla (Louisiana Waterthrush) |
| Bird | Pluvialis dominica (American Golden Plover) |
| Bird | Protonotaria citrea (Prothonotary Warbler) |
| Bird | Riparia riparia (Bank Swallow) |
| Bird | Scolopax minor (American Woodcock) |
| Bird | Setophaga cerulea (Cerulean Warbler) |
| Bird | Setophaga discolor (Prairie Warbler) |
| Bird | Setophaga discolor (Traine Warbler) Setophaga fusca (Blackburnian Warbler) |
| Bird Bird | Sturnella magna (Eastern Meadowlark) |
| Bird Bird | Thryomanes bewickii (Bewick's Wren) |
| Bird Bird | Tyto alba (Barn Owl) |
| Bird | Vermivora chrysoptera (Golden-winged Warbler) |
| Bird | Vermivora chrysoptera (Golden-winged Warbler) Vermivora cyanoptera (Blue-winged Warbler) |

| | DUCK RIVER COA GCN SPECIES (TN) |
|--------------------|---|
| Taxa Group | Scientific & Common names |
| Bivalve | Epioblasma capsaeformis (Oyster Mussel) |
| Bivalve | Epioblasma triquetra (Snuffbox) |
| Bivalve | Lemiox rimosus (Birdwing Pearlymussel) |
| Bivalve | Medionidus conradicus (Cumberland Moccasinshell) |
| Bivalve | Obovaria subrotunda (Round Hickorynut) |
| Bivalve Bivalve | Pleurobema oviforme (Tennessee Clubshell) |
| Bivalve Bivalve | · |
| Bivalve Bivalve | Pleurobema rubrum (Pyramid Pigtoe) Pleuronaia barnesiana (Tennessee Pigtoe) |
| Bivalve Bivalve | |
| Bivalve Bivalve | Pleuronaia dolabelloides (Slabside Pearlymussel) |
| Bivalve Bivalve | Ptychobranchus subtentum (Fluted Kidneyshell) |
| | Quadrula cylindrica cylindrica (Rabbitsfoot) |
| Bivalve | Quadrula intermedia (Cumberland Monkeyface) |
| Bivalve | Toxolasma lividum (Purple Lilliput) |
| Bivalve | Villosa iris (Rainbow) |
| Bivalve | Villosa taeniata (Painted Creekshell) |
| Bivalve | Villosa vanuxemensis (Mountain Creekshell) |
| Fish | Carpiodes velifer (Highfin Carpsucker) |
| Fish | Erimystax dissimilis (Streamline Chub) |
| Fish | Erimystax insignis (Blotched Chub) |
| Fish | Etheostoma aquali (Coppercheek Darter) |
| Fish | Etheostoma cinereum (Ashy Darter) |
| Fish | Etheostoma denoncourti (Golden Darter) |
| Fish | Etheostoma luteovinctum (Redband Darter) |
| Fish | Etheostoma striatulum (Striated Darter) |
| Fish | Etheostoma tippecanoe (Tippecanoe Darter) |
| Fish | Fundulus julisia (Barrens Topminnow) |
| Fish | Hemitremia flammea (Flame Chub) |
| Fish | Ichthyomyzon gagei (Southern Brook Lamprey) |
| Fish | Notropis asperifrons (Burrhead Shiner) |
| Fish | Notropis rubellus (Rosyface Shiner) |
| Fish | Percina phoxocephala (Slenderhead Darter) |
| Gastropod | Lithasia armigera (Armored Rocksnail) |
| Gastropod | Lithasia duttoniana (Helmet Rocksnail) |
| Gastropod | Lithasia geniculata fuliginosa (Geniculate River Snail) |
| Insect | Gomphus sandrius (Tennessee Clubtail) |
| Mammal | Neotoma magister (Allegheny Woodrat) |
| Mammal | Neotoma magister (Allegheny Woodrat) |
| Plant | Allium tricoccum (Ramps) |
| Plant | Ammoselinum popei (Pope's Sand-parsley) |
| Plant | Amsonia tabernaemontana var. gattingeri (Limestone Blue Star) |
| Plant | Anemone caroliniana (Carolina Anemone) |
| Plant | Arabis hirsuta (Western Hairy Rockcress) |
| Plant | Arenaria lanuginosa (Wooly Sandwort) |
| Plant | Arnoglossum plantagineum (Fen Indian-plantain) |
| Plant | Astragalus tennesseensis (Tennessee Milk-vetch) |
| Plant | Boechera perstellata (Braun's Rockcress) |
| Plant | Carex davisii (Davis' Sedge) |
| Plant | Castanea dentata (American Chestnut) |
| Plant | Cimicifuga rubifolia (Appalachian Bugbane) |
| Plant | Dalea foliosa (Leafy Prairie-clover) |
| Plant | Drosera brevifolia (Dwarf Sundew) |
| Plant | Echinacea tennesseensis (Tennessee Coneflower) |
| Plant | Gaylussacia dumosa (Dwarf Huckleberry) |

| | DUCK RIVER COA GCN SPECIES (TN) |
|---------------------------|--|
| Taxa Group | Scientific & Common names |
| Plant | Isoetes melanopoda (Blackfoot Quillwort) |
| Plant | Juglans cinerea (Butternut) |
| Plant | Lilium michiganense (Michigan Lily) |
| Plant | Lycopodiella alopecuroides (Foxtail Clubmoss) |
| Plant | Neviusia alabamensis (Alabama Snow-wreath) |
| Plant | Onosmodium molle ssp. subsetosum (Smooth False Gromwell) |
| Plant | Panax quinquefolius (American Ginseng) |
| Plant | Paysonia densipila (Duck River Bladderpod) |
| Plant | Phemeranthus calcaricus (Limestone Fame-flower) |
| Plant | Polygala boykinii (Boykin's Milkwort) |
| Plant | Schoenolirion croceum (Yellow Sunnybell) |
| Plant | Stellaria fontinalis (Water Stitchwort) |
| Plant | Trifolium calcaricum (Running Glade Clover) |
| Plant | Zanthoxylum americanum (Northern Prickly-ash) |
| Plant | Zigadenus leimanthoides (Death-camas) |
| Reptile | Crotalus horridus (Timber Rattlesnake) |
| Reptile | Pituophis melanoleucus melanoleucus (Northern Pinesnake) |
| Reptile | Terrapene carolina (Eastern Box Turtle) |
| Subterranean - Amphibian | Gyrinophilus palleucus (Tennessee Cave Salamander) |
| Subterranean - Amphibian | Gyrinophilus palleucus necturoides (Big Mouth Cave Salamander) |
| Subterranean - Arachnid | Hesperochernes mirabilis (Southeastern Cave Pseudoscorpion) |
| Subterranean - Crustacean | Caecidotea bicrenata ((a cave obligate isopod)) |
| Subterranean - Crustacean | Caecidotea bicrenata bicrenata ((a cave obligate isopod)) |
| Subterranean - Crustacean | |
| Subterranean - Crustacean | Orconectes australis (Southern Cave Crayfish) |
| Subterranean - Crustacean | Stygobromus exilis (Central Kentucky Cave Obligate) |
| Subterranean - Fish | Typhlichthys subterraneus (Southern Cavefish) |
| Subterranean - Flatworm | Sphalloplana percoeca ((a cave obligate planarian)) |
| Subterranean - Gastropod | Glyphyalinia specus (Hollow Glyph) |
| Subterranean - Insect | Batrisodes ferulifer ((a cave obligate beetle)) |
| Subterranean - Insect | Litocampa cherokeensis ((a bristletail)) |
| Subterranean - Insect | Litocampa cookei (Cooke's Cave Dipluran) |
| Subterranean - Insect | Litocampa davisi ((a bristletail)) |
| Subterranean - Insect | Pseudanophthalmus bendermani (Benderman's Cave Beetle) |
| Subterranean - Insect | Pseudanophthalmus tullahoma (Duck River Cave Beetle) |
| Subterranean - Insect | Pseudosinella hirsuta (Hirsute Cave Springtail) |
| Subterranean - Millipede | Scoterpes jackdanieli ((a cave obligate millipede)) |
| Subterranean - Millipede | Scoterpes musicarustica ((a cave obligate millipede)) |
| Subterranean - Millipede | Scoterpes tricorner ((a cave obligate millipede)) |

| DUCK RIVER COA HABITATS (TN) | |
|--|--|
| Terrestrial Habitat Type | Total acres of very high, high, and medium ranked habitat |
| Natural habitats | |
| Nashville Basin Limestone Glade and Woodland | 25855 |
| Southern Interior Low Plateau Dry-Mesic Oak Forest | 25688 |
| South-Central Interior Mesophytic Forest | 7696 |
| Eastern Highland Rim Prairie and Barrens | 1720 |
| South-Central Interior Large Floodplain | 749 |
| South-Central Interior Small Stream and Riparian | 214 |
| Semi-natural habitats | |
| Pasture | 14447 |
| Old Field / Successional | 4612 |
| Cropland | 125 |
| | |
| Aquatic Habitat Type | Total stream miles of very high, high, and medium ranked habitat |
| Headwaters and Streams | 131 |
| Small River | 70 |
| Medium River | 77 |