

# Dogwood Anthracnose

***Dr. Fulya Baysal-Gurel, Md Niamul Kabir and  
Angelo Randaci***

***Otis L. Floyd Nursery Research Center  
College of Agriculture  
Tennessee State University  
[fbaysalg@tnstate.edu](mailto:fbaysalg@tnstate.edu)***

**ANR-PATH-8-2017**

Dogwood anthracnose (*Discula destructiva*) is often described as the most serious disease that affects dogwood species, specifically flowering (*Cornus florida*) and mountain (*C. nuttallii*) dogwood species in the United States. The disease was first noticed on the flowering dogwoods in New York and Connecticut in 1978 (1) and on mountain dogwoods in Washington in 1979 (2). Since the first reports in the late 1970's, infection of mountain dogwoods has been reported in Oregon, Idaho, British Columbia and Northern California; and reported on flowering dogwoods in Massachusetts, New Jersey, Pennsylvania, Delaware, Maryland, Virginia, West Virginia, North Carolina, South Carolina, Tennessee, Georgia, Kentucky, Alabama, Ohio, the District of Columbia, New Hampshire, Rhode Island, Indiana, Vermont, Michigan and Missouri (3). Dogwood anthracnose threatens forest ecosystems and causes substantial economic losses for the ornamental nursery industry.

## **Symptoms**

Symptoms of dogwood anthracnose include necrosis of bracts, leaf spots, necrotic blotches (Figure 1), wilted and curled foliage, dieback of twigs, leaf blight and cankers on trunks (4). Initial symptoms develop on the lower leaves especially on the leaf margins and flower bracts. Tan spots on leaves with purple rims are often visible. The spots then develop into large, dead areas and as the leaves become necrotic they either dropoff or remain on the tree throughout the fall and winter seasons. Infections from the leaves spread through the

leaf petioles and produce cankers. Dogwood flower bracts are normally prone to infections and display reddish, brownish spots, or blotches. If rainy conditions occur during the flowering period, infected bracts are prevalent. Twig dieback may occur due to cankers at the leaf nodes. Succulent shoots often proliferate on the main branches and lower portions of the trunk on affected trees. Succulent branches are very susceptible to infection. If the disease spreads to the main stem of dogwood plants, cankers with swelling and splitting bark are observed. This can cause girdling of individual branches or death of the entire tree (2, 5 and 6).



Figure 1. Necrotic blotches caused by dogwood anthracnose.

### **Disease Cycle**

Anthrachnose fungus may persist in a dormant stage for extended periods in leaves, twigs, leaf debris, and branches. During spring, eruption of fruiting structures (acervuli) occurs on the lower side of the spotted leaves and large amounts of spores (conidia) are released through infected twigs. The spores are spread to newly developed leaves, flower bracts and green shoots by rain, air currents, and sometimes by birds on. Germination takes place when the spores come in contact with water on epicormic sprouts and leaves. In the infected leaves and twigs this fungus can overwinter and can resume growth in the spring to kill new shoots, twigs and buds. Cool temperatures (65°-75°F), along with moist weather are favorable conditions for infection and spread of the disease.

### **Disease Management**

Efficient control of dogwood anthracnose is achieved with cultural and chemical control techniques. Since dogwood anthracnose can be introduced via infected plants, careful inspection should be conducted prior to and after the purchase of plant material. During dry weather conditions it is necessary to cut out all infected twigs, limbs and suckers from infected trunks or branches to reduce the inoculum level. During the fall season, fallen leaves must be removed because those leaves are a great source of overwintering type of disease inoculum. It is important to utilize proper watering and mulching during drought to ensure tree vigor. Sufficient spacing is important for air circulation which can help to

reduce infection. Excessive use of fertilizers must be avoided, because it may increase new succulent branches on plants, rendering them more susceptible to dogwood anthracnose. If this disease is a recurring problem, resistant dogwood cultivar or hybrids need to be considered for planting. See list below of resistant dogwoods (7, 8).

***Anthracnose Resistant Dogwoods***

***C. florida*** cultivar ‘Appalachian Spring’

***C. kousa*** cultivars ‘Big Apple’, ‘China Girl’, ‘Elizabeth Lustgarten’, ‘Gay Head’, ‘Greensleeves’, ‘Julian’, ‘Milky Way’, ‘Steeple’ and ‘Temple Jewel’

***C. florida* x *C. kousa*** hybrids ‘Aurora’, ‘Celestial’, ‘Constellation’, ‘Ruth Ellen’, ‘Star Dust’, ‘Stellar Pink’, ‘Empire’, ‘Red Steeple’, ‘Pam’s Mountain’ and ‘Bouquet’

Fungicides can be used for dogwood anthracnose management in conjunction with other management strategies previously mentioned (Table 1). When there is a risk of dogwood anthracnose occurring, repeated applications of fungicides or biorational products may be necessary.

**Table 1.** Selected lists of fungicide and biorational product that can be used to prevent dogwood anthracnose.

Active Ingredient	FRAC Code	Notes
<b>Azoxystrobin</b>	11	Spray at 7 to 28 day intervals when conditions favor disease development.
<b><i>Bacillus subtilis</i> QST173</b>	44	Spray at 7 day intervals when conditions favor disease development.
<b>Chlorothalonil</b>	M5	Spray at 7 to 14 day intervals when conditions favor disease development.
<b>Chlorothalonil + thiophanate-methyl</b>	M5 + 1	Spray at 7 day intervals when conditions favor disease development.
<b>Copper hydroxide</b>	M1	Begin application at the first sign of the disease and repeat at 7 to 14 day intervals.
<b>Copper hydroxide + mancozeb</b>	M1 + M3	Spray at 10 to 14 day intervals.
<b>Copper sulfate</b>	M1	Begin application at the first sign of the disease and repeat at 7 to 10day intervals.
<b>Fluoxastrobin</b>	11	Under low disease pressure spray at 7 to 28day

<b>Active Ingredient</b>	<b>FRAC Code</b>	<b>Notes</b>
		intervals. Under high disease pressure spray at 7 to 14 day intervals.
<b>Kresoxim-methyl</b>	11	Spray strictly as a protective treatment at 7 to 10 day intervals.
<b>Mancozeb</b>	M3	Begin application at bud break in early spring.
<b>Mancozeb + myclobutanil</b>	M3 + 3	Begin application at the first sign of the disease and repeat at 7 to 10 day intervals.
<b>Myclobutanil</b>	3	Spray at 7 to 14 day intervals.
<b>Potassium bicarbonate</b>	NC	Spray at 7 to 14 day intervals.
<b>Propiconazole</b>	3	Spray at 14 to 28 day intervals when conditions favor disease development.
<b>Propiconazole + chlorothalonil</b>	3 + M5	Spray at 21 day intervals. Application is not recommended in greenhouses and to the blooms.
<b>Pyraclostrobin + boscalid</b>	11 + 7	Begin application prior to the first sign of the disease and repeat at 7 to 14 day intervals.
<b>Tebuconazole</b>	3	Begin application with spring bud break at 14 to 28 day intervals.
<b>Thiophanate-methyl</b>	1	Spray at 14 to 21 day intervals when conditions favor disease development.
<b>Thiophanate-methyl + mancozeb</b>	1 + M3	Spray at 7 day intervals.
<b>Trifloxystrobin</b>	11	Begin application at the first sign of the disease and repeat at 7 to 14 day intervals.
<b>Trifloxystrobin + triadimefon</b>	11 + 3	Begin application at the first sign of the disease and repeat at 7 to 14 day intervals.

**NOTE:** Before applying ANY disease management product, be sure to: 1) read the label to be sure that the product is allowed for the crop and the disease you intend to control; 2) read and understand the safety precautions and application restrictions.

## References

1. Pirone, P.P. 1980. Parasitic fungus affects region's dogwood. New York: New York Times; Feb. 24, pp. 34, 37.
2. Byther, R.S. and Davidson, R.M. 1979. Dogwood anthracnose. Ornamental North West Newsletter; 3:20-21.
3. U.S. Department of Agriculture Forest Service, Forest Health Protection and its partners. 2013. <http://www.fs.fed.us/nrs/tools/afpe/maps/pdf/DWA.pdf>

4. Zhang, N., and Blackwell, M. 2002. Population structure of dogwood anthracnose fungus. *Phytopathology* 92:1276-1283.
5. Daughtrey, M.L., C.R. Hibben and G.W. Hudler. 1988. Cause and control of dogwood anthracnose in northeastern United States. *Journal of Arboriculture* 14(2): 55.
6. Hibben, C.R., M.L. Daughtrey. 1988. Dogwood anthracnose in northeastern United States. *Plant Disease* 72:199-203.
7. Gillman, D. 2011. Dogwood anthracnose. <https://ag.umass.edu/landscape/fact-sheets/dogwood-anthracnose>.
8. Smith, S. and Carson, J. 2016. Anthracnose diseases of dogwood <https://www.uaex.edu/publications/pdf/FSA-7564.pdf>

**For additional information, contact your local nursery specialist office at:**

**Tennessee State University**  
College of Agriculture  
3500 John A. Merritt Blvd., Box 9635 Nashville, TN 3720-1561  
<http://www.tnstate.edu/extension>

**Tennessee State University, Otis L. Floyd Nursery Research Center**  
472 Cadillac Lane McMinnville, TN 37110 <http://www.tnstate.edu/agriculture/nrc/>

#### **Precautionary Statement**

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label.

#### **Disclaimer**

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication. Use of trade, brand, or active ingredient names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar and suitable composition, nor does it guarantee or warrant the standard of the product. The author(s) and Tennessee State University assume no liability resulting from the use of these recommendations.

Dr. Chandra Reddy, Dean, Tennessee State University, College of Agriculture  
Dr. Latif Lighari, Associate Dean Extension, Tennessee State University, College of Agriculture  
Dr. Nick Gawel, Superintendent, Otis L. Floyd Nursery Research Center, Tennessee State University, College of Agriculture

---

**TSU-18-0039(B)-15i-61065** Tennessee State University does not discriminate against students, employees, or applicants for admission or employment on the basis of race, color, religion, creed, national origin, sex, sexual orientation, gender identity/expression, disability, age, status as a protected veteran, genetic information, or any other legally protected class with respect to all employment, programs and activities sponsored by Tennessee State University. The following person has been designated to handle inquiries regarding non-discrimination policies: Rita Williams Seay, Office of Equity and Inclusion, [rseay@tnstate.edu](mailto:rseay@tnstate.edu), 3500 John Merritt Blvd., McWhorter Administration Building, Suite 260, Nashville, TN 37209, 615-963-7438. The Tennessee State University policy on nondiscrimination can be found at [www.tnstate.edu/nondiscrimination](http://www.tnstate.edu/nondiscrimination).