

Exotic butterflies and moths (Lepidoptera) in botanical gardens – potential for education and research

MARTIN SUVÁK

Botanical Garden of Pavol Jozef Šafárik University in Košice, Mánesova 23, 043 52
Košice, martin.suvak@upjs.sk

Suvák M. (2015): Exotic butterflies and moths (Lepidoptera) in botanical gardens – potential for education and research. – Thaiszia – J. Bot. 25 (Suppl. 1): 81-147. – ISSN 1210-0420.

Abstract: Exhibitions of live butterflies and moths can have very high educational potential. In natural environments, lepidopteran and plant taxa closely interact and many characteristics of representatives of both these groups are results of such relationships. Therefore, free flying butterflies in botanical gardens enable demonstrations of a wide variety of natural phenomena studied within entomology, ecology, botany, evolutionary theory etc. The Victoria greenhouse of Botanical Garden of P. J. Šafárik University in Košice provides excellent conditions for such observations. Combination of tropical and subtropical plants with live exotic butterflies and moths, in many cases from the same regions as the plant taxa, gives a unique opportunity to see phenomena usually naturally available only far away from Central Europe. During annual seasonal “butterfly shows” in the years 2008 – 2015, the visitors of this greenhouse could see at least 182 different lepidopteran taxa mostly from Central and South America, Africa and South East Asia. Next to the presented species diversity, some observations on food preferences, behavior, mimetic forms, natural enemies and other ecological relationships are discussed here.

Keywords: Lepidoptera, butterflies, moths, education, ecology, botanical garden, greenhouses.

Introduction

Exhibitions of live insects and other arthropods in insect zoos or insectaria have enormous educational value (SAUL-GERSHENZ 2009). Free-flying butterflies are among the most popular subjects there. Next to the specialized butterfly houses, presentations of live butterflies and moths have spread worldwide in a number of other sites, at least seasonally. International cooperation of butterfly breeders (usually farmers in tropical and subtropical regions) with distributors of pupae (sometimes also with their own pupae production) and rapid air mail enable to satisfy customer requirements practically everywhere. In the world, butterfly exhibits are held in various facilities of different sizes but with otherwise suitable conditions (temperature, humidity, light, access for visitors etc.), usually in greenhouses. Butterfly farmers and distributors can organize their own exhibitions but such actions are seasonally undertaken also by other subjects with appropriate spaces, using pupae from commercial suppliers. Since lepidopterans and plants have close relationships, botanical gardens can provide very suitable conditions for such presentations. Free-flying butterflies between wide variety of plants and proper explanatory boards may increase educative value of such exhibitions. However, this potential is not always fully used.

Since 2008, a live butterfly show is annually organized also in Botanical Garden of Pavol Jozef Šafárik University in Košice (hereinafter BG PJŠU). It became one of the most popular seasonal event with plenty of visitors of all ages every year (Tab. 1). Though many of them just wants to see flying strange creatures with colorful wings, there are much more to see also for more demanding visitors, students and teachers of biological disciplines, professional entomologists and others.

Tab. 1. Summary data on butterfly exhibitions in BG PJŠU

Year	Duration of exhibitions			Number of visitors	No. of pupae	No. of different butterfly species
	from	to	no. of days			
2008	15.5.2008	15.6.2008	32	19682	1055	46
2009	22.5.2009	28.6.2009	38	25463	1800	66
2010	21.5.2010	30.6.2010	41	18349	2742	55
2011	20.5.2011	3.7.2011	45	20679	2727	83
2012	18.5.2012	30.6.2012	44	15438	3295	99
2013	17.5.2013	30.6.2013	45	17337	3087	86
2014	16.5.2014	30.6.2014	46	14069	3035	68
2015	22.5.2015	30.6.2015	40	20614	3110	89
Total			331	151631	20851	182

Material and methods

Butterfly show in BG PJŠU is usually presented for visitors within a time span of cca 40 days, in May-June (Tab. 1). Some pupae are ordered in advance before opening the exhibition and next orders are realized later once a week to guarantee enough eclosed butterflies during the specified period. The pupae are

ordering from specialized suppliers either as individual species or as a special mixes according to their geographical origin or taxonomic relatedness.

The butterfly exhibition itself takes place in the Victoria greenhouse (Fig. 1) of BG PJŠU. It covers an area of 310 m² (25 m x 12.5 m), its maximum height is 4.5 m and the total volume is about 1150 m³. Two water basins are in the centre and a circular trail for visitors runs along the side walls of this greenhouse. The plants (Tab. 2) grow in all free spaces between basins and a pathway, including lianas and epiphytes on the walls and arranged tree trunks. The average temperature here is about 25°C but occasionally it can vary from 18°C to 40 °C (in hot summer days). Humidity can be relatively high (up to 95 %) with respect to the water basins.

The pupae are placed to special open boxes within the greenhouses, so enclosed butterfly are free to fly to the greenhouse space.

Next to the plants with long term fixed locations, some other plants are used to be placed there temporarily – some flowering potted plants or their cuttings arranged around basins as a source of nectar for flying butterflies. Other food sources are provided on special feeders (mostly fresh or decaying fruits).

During last 8 years of annually undertaken butterfly exhibitions, a lot of interesting phenomena was registered and many photographs were taken here (all selected illustrative figures below). Some specimens were also stored, e.g. several dead butterflies were collected, individuals of other insects (e.g. parasitoids) and spiders were sampled for future analyses. Selected data on species diversity (Fig. 2, Tab. 3) and observations on other interesting phenomena are summarised below.



Fig. 1. Victoria greenhouse in Botanical Garden of P. J. Šafárik University in Košice.

Tab. 2. Plants in the Victoria greenhouse of BG PJŠU in Košice.

Plant taxa ¹	Distribution ²
Pteridophyta	
Aspleniaceae	
<i>Asplenium nidus</i> L.	tropical SE Asia, E Australia, Hawaii, Polynesia, Christmas Island, India, E Africa
Cibotiaceae	
* <i>Cibotium</i> Kaulfuss sp.	
Dryopteridaceae	
<i>Cyrtomium falcatum</i> (L. fil.) Presl	China, Taiwan, North Korea, South Korea, Japan, Vietnam
<i>Tectaria cicutaria</i> (L.) Copel.	Puerto Rico, Cuba, Hispaniola, Jamaica
Lomariopsidaceae	
<i>Nephrolepis exaltata</i> (L.) Schott	N South America, Mexico, Central America, Florida, West Indies, Polynesia, Africa
Lygodiaceae	
<i>Lygodium japonicum</i> (Thunb.) Sw.	China, South Korea, Sri Lanka, Philippines, Moluccas, Sunda Islands, New Guinea, Peninsular Malaysia, Vietnam, Laos, Cambodia, Thailand, India Nepal
Polypodiaceae	
<i>Platycterium bifurcatum</i> (Cav.) C.Chr.	Java, New Guinea, SE Australia
<i>Platycterium</i> Desv. sp.	tropical and temperate areas of South America, Africa, SE Asia, Australia, New Guinea
Pteridaceae	
<i>Acrostichum aureum</i> L.	tropical and sub-tropical areas around the world (mangrove swamps)
<i>Adiantum raddianum</i> C.Presl	tropical and subtropical South America.
Cycadophyta	
Cycadaceae	
<i>Cycas circinalis</i> L.	Sri Lanka
* <i>Cycas rumphii</i> Miq.	Indonesia, New Guinea, Christmas Island
Zamiaceae	
<i>Encephalartos villosus</i> Lem.	S Africa
<i>Encephalartos ferox</i> (G.Bertol) Lehm.	SE Africa
Magnoliophyta	
Araliaceae	
* <i>Polyscias</i> J.R.Forst. & G.Forst. sp.	tropical areas
Acanthaceae	
<i>Fittonia verschaffeltii</i> (Lem.) Van Houtte	Peru
<i>Hypoestes sanguinolenta</i> (Van Houtte) Hook. f.	Madagascar
<i>Ruellia amoena</i> Nees	Argentina, Cerrado vegetation of Brazil, Mexico
<i>Ruellia portellae</i> Hook. fil.	Brazil
* <i>Thunbergia affinis</i> S.Moore	E Africa
Amaranthaceae	
* <i>Celosia argentea</i> L.	India
<i>Alternanthera</i> Forssk.	mostly in tropical Americas, other species in Asia, Africa, and Australia
*unidentified sp.	

Tab. 2. – cont.

Plant taxa ¹	Distribution ²
Anacardiaceae	
* <i>Mangifera indica</i> L.	India
Apocynaceae	
<i>Allamanda cathartica</i> L.	Brazil
<i>Dipladenia splendens</i> (Hook.f.) A.DC.	Brazil
<i>Stephanotis floribunda</i> Brongn.	Madagascar
Araceae	
<i>Alocasia macrorrhizos</i> (L.) G.Don	from Malaysia to Queensland
<i>Alocasia</i> (Schott) G.Don	tropical and subtropical Asia to E Australia
<i>Anthurium andraeanum</i> Linden ex André	Colombia, Ecuador
<i>Anthurium digitatum</i> (Jacq.) Schott	Venezuela
<i>Anthurium hookeri</i> Kunth	Central and South America
<i>Anthurium magnificum</i> Linden	Colombia
<i>Anthurium pedatoradiatum</i> Schott	Mexico
<i>Anthurium scherzerianum</i> Schott	Costa Rica
<i>Anthurium tetragonum</i> Hook. ex Schott	Belize, Guatemala, Honduras, Mexico, Panama
<i>Anubias</i> Schott sp.	tropical central and W Africa
<i>Caladium bicolor</i> (Aiton) Vent.	from Costa Rica to N Argentina
<i>Caladium</i> Vent. sp.	South America, Central America
<i>Colocasia esculenta</i> (L.) Schott	Malaysia
<i>Cryptocoryne pontederiifolia</i> Schott	Sumatra
<i>Cryptocoryne walkeri</i> Schott 1857	Sri Lanka
<i>Monstera decursiva</i> (Roxb.) Schott	China, Indian subcontinent, Indochina
<i>Monstera deliciosa</i> Liebm.	from S Mexico to Panama
<i>Monstera karwinskyi</i> Schott	from Mexico to N South America
<i>Monstera obliqua</i> Miq.	Brazil
<i>Philodendron melanochrysum</i> Linden & André	from Costa Rica to Columbia into Ecuador and Peru
<i>Philodendron bipinnatifidum</i> Schott ex Endl.	South America, namely Brazil, Bolivia, Argentina, and Paraguay
<i>Philodendron erubescens</i> K.Koch & Augustin	Colombia
<i>Philodendron panduriforme</i> (Kunth) Kunth	N and W areas of the Amazon basin: Peru, Ecuador, Brazil, Colombia, Venezuela
<i>Philodendron wendlandii</i> Schott	from Nicaragua to Panama
<i>Philodendron x corsinianum</i> Hort.	
<i>Philodendron xanadu</i> Croat, Mayo & J.Boos, 2002 publ. 2003	Brazil
<i>Pistia stratiotes</i> L.	pantropical, probably originated in Africa
<i>Scindapsus aureus</i> (Linden & André) Engl., 1908	French Polynesia
<i>Scindapsus pictus</i> Hassk.	Bangladesh, Thailand, Peninsular Malaysia, Borneo, Java, Sumatra, Sulawesi, Philippines
<i>Scindapsus</i> Schott sp.	SE Asia, New Guinea, Queensland, W Pacific islands
<i>Spathiphyllum wallisii</i> Regel	Central America
<i>Spathiphyllum</i> Schott sp.	tropical regions of the Americas and SE Asia
<i>Syngonium</i> Schott sp.	S Mexico, West Indies, Central and South America
* <i>Zamioculcas zamiifolia</i> (Lodd.) Engl.	E Africa
unidentified sp.	

Tab. 2. – cont.

Plant taxa ¹	Distribution ²
Araliaceae	
* <i>Dizygotheca elegantissima</i> (Veitch ex Mast.) R. Vig. & Guillaumin	New Caledonia
<i>Schefflera digitata</i> J.R.Forst. et G.Forst.	New Zealand
Areaceae	
<i>Chamaedorea</i> Willd. sp.	subtropical and tropical regions of the Americas
<i>Chamaedorea elegans</i> Mart.	S Mexico, Guatemala
<i>Chamaedorea oblongata</i> Mart.	Belize, Guatemala, Honduras, Mexico, Nicaragua
<i>Pritchardia hillebrandii</i> (Kuntze) Becc.	Hawaii
Aristolochiaceae	
<i>Aristolochia gigantea</i> Mart. & Zucc.	Brazil
* <i>Aristolochia</i> L. sp.	widespread in different climates of the world
Asparagaceae	
<i>Asparagus myriocladus</i> Baker	S Africa
* <i>Asparagus</i> L. sp.	from rainforest to semi-desert habitats of the world
<i>Cordylina terminalis</i> Kunth	tropical SE Asia, Papua New Guinea, Melanesia, NE Australia, Polynesia
<i>Dracaena</i> Vand. ex L. sp.	Africa, S Asia, Central America
<i>Ledebouria kirkii</i> (Baker) Stedje & Thulin	tropical Africa
Asteraceae	
* <i>Argyranthemum frutescens</i> (L.) Sch.Bip.	Canary Islands
Bignoniaceae	
<i>Tecomaria capensis</i> (Thunb.) Spach.	S Africa
<i>Tecoma</i> Juss. sp.	the Americas, Africa
Boraginaceae	
* <i>Heliotropium</i> L. sp.	subtropical regions of the world
Bromeliaceae	
x <i>Cryptbergia</i> (<i>Cryptanthus bahianus</i> x <i>Billbergia nutans</i>)	
<i>Aechmea bracteata</i> (Swartz) Grisebach	Central America, Mexico, Colombia, Venezuela
<i>Aechmea brasiliensis</i> Regel	Cerrado vegetation in Brazil, N Argentina, Bolivia, Paraguay, Uruguay
<i>Aechmea caudata</i> Lindman	SE Brazil
<i>Aechmea distichantha</i> Lem.	Cerrado vegetation in Brazil, N Argentina, Bolivia, Paraguay, Uruguay
<i>Aechmea fasciata</i> (Lindl.) Baker	Brazil
<i>Aechmea fillicaulis</i> (Griseb.) Mez	Venezuela
<i>Aechmea gamosepala</i> Wittm.	S Brazil
<i>Aechmea chantinii</i> (Carrière) Baker	Amazon rainforest vegetation in Brazil, Venezuela, Colombia, Ecuador and Peru
<i>Aechmea chlorophylla</i> L.B.Sm.	E Brazil
<i>Aechmea lueddemanniana</i> (K. Koch) Mez in Engler	Central America
<i>Aechmea mexicana</i> Baker	Mexico, Central America, Colombia, Ecuador
<i>Aechmea organensis</i> Wawra	SE Brazil
<i>Aechmea orlandiana</i> L.B.Sm.	Brazil
<i>Aechmea pubescens</i> Baker	Costa Rica, Honduras, Nicaragua, Panama, Colombia, Venezuela

Tab. 2. – cont.

Plant taxa ¹	Distribution ²
<i>Aechmea racinae</i> L.B. Smith	Brazil
<i>Aechmea recurvata</i> (Klotzsch) L.B. Smith	S Brazil, Paraguay, Uruguay, N Argentina
<i>Aechmea tillandsioides</i> (Mart. ex Schult. & Schult.f.) Baker	S Mexico, Central America, and N South America
<i>Aechmea weilbachii</i> Didrichsen	E Brazil
<i>Aechmea</i> Ruiz & Pav. sp.	from Mexico through South America
<i>Ananas comosus</i> (L.) Merr.	South America, Central America
<i>Billbergia decora</i> Poeppig & Endlicher	Peru, Bolivia, Brazil
<i>Billbergia euphemiae</i> E. Morren	Brazil
<i>Billbergia horrida</i> Regel	Brazil
<i>Billbergia</i> "Hoelscheriana" (<i>B. nutans</i> x <i>Saundersii</i>)	
<i>Billbergia nutans</i> H.Wendl.	Brazil, Paraguay, Uruguay, Argentina
<i>Billbergia vittata</i> Brongniart	Brazil
<i>Billbergia</i> x <i>windii</i> (<i>Billbergia nutans</i> x <i>Bilbergia decora</i>)	
<i>Billbergia</i> Thunb. sp.	S Mexico, West Indies, Central America, South America
<i>Cryptanthus bivittatus</i> Rgl.	E Brazil
<i>Edmundoa lindenii</i> (Regel) Leme	Brazil
<i>Guzmania melinonis</i> Regel	Bolivia, Peru, Colombia, the Guianas, Venezuela, Brazil, Ecuador
<i>Guzmania minor</i> Mez	Central America, N and central South America, S Mexico and the West Indies
<i>Guzmania monostachia</i> (L.) Rusby ex Mez	South America (Bolivia, Brazil, Colombia, Ecuador, Peru, Venezuela), Central America, the West Indies, Florida
<i>Guzmania</i> Ruiz & Pav. sp.	Florida, the West Indies, S Mexico, Central America, N and W South America
<i>Hohenbergia stellata</i> Schult. & Schult.f.	Trinidad and Tobago, Martinique, Netherlands Antilles, Venezuela, NE Brazil
<i>Neoregelia carolinae</i> (Beer) L.B.Sm.	Brazil
<i>Neoregelia marmorata</i> (Baker) L.B. Smith	Brazil
<i>Neoregelia pineliana</i> (Lemaire) L.B. Smith	Brazil
<i>Neoregelia spectabilis</i> (T. Moore) L.B. Smith	S Brazil
<i>Neoregelia tristis</i> (Beer) L.B. Smith	Brazil
<i>Nidularium purpureum</i> Beer	Brazil
<i>Pitcairnia carnea</i> Beer	Panama
<i>Pitcairnia corallina</i> Linden & André	Brazil, Colombia, Peru.
<i>Pitcairnia xanthocalyx</i> Martius	Mexico
<i>Quesnelia liboniana</i> (De Jonghe) Mez	Brazil
<i>Tillandsia</i> L. sp. 1	Central and South America, S United States, West Indies
<i>Tillandsia</i> L. sp. 2	Central and South America, S United States, West Indies
<i>Tillandsia stricta</i> Solander	South America, Trinidad
<i>Tillandsia usneoides</i> (L.) L.	SE United States, Mexico, Bermuda, the Bahamas, Central America, South America, West Indies
<i>Vriesea</i> hybr. (<i>Vriesea glutinosa</i> x <i>Vriesea splendens</i>)	

Tab. 2. – cont.

Plant taxa ¹	Distribution ²
<i>Vriesea scalaris</i> E. Morren	Brazil, Venezuela
<i>Vriesea</i> Lindl. sp.	Mexico, Central America, South America, West Indies
<i>Vriesea splendens</i> (Brongn.) Lem.	Trinidad, E Venezuela, the Guianas
<i>Wittrockia amazonica</i> (Baker) L.B.Sm.	Brazil
unidentified sp. 1	
unidentified sp. 2	
unidentified sp. 3	
Cactaceae	
<i>Lepismium</i> Pfeiff. sp.	tropical South America
<i>Rhipsalis baccifera</i> (J.S. Mueller) Stearn	Central and South America, the Caribbean, Florida
<i>Rhipsalis</i> Gaertn. sp.	Central America, Caribbean, N and central South America, tropical Africa, Madagascar, Sri Lanka
<i>Selenicereus chrysocardium</i> (Alexander) Kimmach	Mexico
<i>Schlumbergera</i> Lem. sp.	Brazil
Clusiaceae	
<i>Clusia rosea</i> Jacq.	Caribbean, Florida
Commelinaceae	
<i>Tradescantia</i> Ruppis ex L. sp.	New World from S Canada south to N Argentina including the West Indies
Costaceae	
<i>Costus lucanusianus</i> J.Braun & K.Schum.	tropical Africa
<i>Costus malortieanus</i> H.Wendl.	Nicaragua, Costa Rica
<i>Cheilocostus speciosus</i> (J.Konig) C.Specht	SE Asia, China, Queensland
Cucurbitaceae	
<i>Neoalsomitra sarcophylla</i> (Wall.) Hutchinson	Myanmar, Thailand, Laos, Philippines, Palawan, Cambodia, Vietnam, Sulawesi
Cyperaceae	
<i>Cyperus altermifolius</i> L.	Madagascar
* <i>Cyperus papyrus</i> L.	Africa
Erythroxylaceae	
<i>Erythroxylum coca</i> Lam.	rain forests of the Andes
Euphorbiaceae	
<i>Codiaeum</i> A.Juss. sp.	insular SE Asia, N Australia, Papua New Guinea
<i>Codiaeum variegatum</i> (L.) A.Juss.	Indonesia, Malaysia, Australia, W Pacific Ocean islands
Geraniaceae	
+ <i>Pelargonium</i> L'Hér. sp.	temperate and tropical regions of the world, many species in S Africa.
Gesneriaceae	
<i>Aeschynanthus radicans</i> Jack	humid tropics of the Malay Peninsula south to Java
<i>Columnnea schiedeana</i> Schlechtend.	E Mexico
Heliconiaceae	
<i>Heliconia bihai</i> (L.) L.	N South America, West Indies
<i>Heliconia humilis</i> (Aubl.) Jacq.	N South America, West Indies
<i>Heliconia</i> L. sp.	tropical Americas, certain islands of W Pacific, Maluku

Tab. 2. – cont.

Plant taxa ¹	Distribution ²
Hypoxidaceae	
<i>Curculigo latifolia</i> Dryand. ex W.T.Aiton	E Asia: S China, Bangladesh, Andaman Islands, Myanmar, Thailand, Vietnam, Malaysia, Indonesia, Philippines
Lamiaceae	
* <i>Salvia splendens</i> Sellow ex J.A. Schultes	Brazil
Malvaceae	
unidentified sp.	
Marantaceae	
<i>Calathea ornata</i> (Linden ex Lem.) Körn.	Colombia, Venezuela
<i>Calathea zebrina</i> (Sims) Lindl.	SE Brazil
<i>Ctenanthe oppenheimiana</i> (E.Morr) K.Schum	Brazil
<i>Maranta leuconeura</i> E.Morren	Brazil
<i>Maranta</i> L. sp.	tropical Central and South America, West Indies
Melastomataceae	
* <i>Medinilla</i> Gaudich. sp.	tropical regions of Africa, Madagascar, S Asia, W Pacific Ocean islands
* <i>Medinilla magnifica</i> Lindl.	Philippines
Moraceae	
* <i>Artocarpus</i> J.R.Forster & G.Forster sp.	SE Asia, Pacific islands
<i>Artocarpus heterophyllus</i> Lam.	parts of S and SE Asia
* <i>Ficus elastica</i> Roxb. ex Hornem. 1819	NE India, Nepal, Bhutan, Burma, China, Malaysia, Indonesia
<i>Ficus pumila</i> L.	E Asia (China, Japan, Vietnam)
Musaceae	
<i>Musa velutina</i> H.Wendl. & Drude	India, Myanmar
Nelumbonaceae	
* <i>Nelumbo nucifera</i> Gaertn.	tropical Asia and N Australia
Nymphaeaceae	
<i>Nymphaea</i> L. x hybrida	cosmopolitan distribution
<i>Nymphaea zanzibariensis</i> Casp.	Galapagos
* <i>Victoria regia</i> Lindl.	Amazon River basin
Orchidaceae	
<i>Cattleya</i> Lindl. sp.	from Costa Rica and the Lesser Antilles south to Argentina
<i>Dendrobium</i> Sw. sp.	SE Asia, Philippines, Indonesia, Australia, New Guinea, Vietnam, Pacific islands
<i>Paphiopedilum</i> Pfitzer sp.	SE Asia, the Indian Subcontinent, S China, New Guinea, the Solomon and Bismarck Islands
<i>Phalaenopsis</i> Blume sp.	S China, India, SE Asia, New Guinea, the Bismarck Archipelago, Queensland
* <i>Stanhopea oculata</i> (Lodd.) Lindl.	from Mexico to Colombia and SE Brazil
<i>Vanda</i> Gaud. ex Pfitzer sp. 1	E Asia, SE Asia, New Guinea, Queensland, W Pacific islands
<i>Vanda</i> Gaud. ex Pfitzer sp. 2	E Asia, SE Asia, New Guinea, Queensland, W Pacific islands
<i>Vanilla aphylla</i> Blume	SE Asia
<i>Vanilla planifolia</i> Jacks. ex Andrews	Mexico

Tab. 2. – cont.

Plant taxa ¹	Distribution ²
Oxalidaceae	
<i>Averrhoa carambola</i> L.	SE Asia, Indian Subcontinent
Pandanaceae	
* <i>Pandanus tectorius</i> Parkinson ex Du Roi	SE Asia, E Australia, the Pacific Islands
Passifloraceae	
<i>Passiflora caerulea</i> L.	South America (Argentina, Chile, Paraguay, Uruguay and Brazil)
* <i>Passiflora quadrangularis</i> L.	Neotropics
* <i>Passiflora racemosa</i> Brot.	Brazil
<i>Passiflora</i> L. sp.	Central and South America, SE Asia, New Guinea, Australia, New Zealand
Piperaceae	
<i>Peperomia</i> Ruiz & Pav. sp.	tropical and subtropical regions of the world
<i>Piper ornatum</i> N.E.Br.	Indonesia
Plantaginaceae	
<i>Russelia equisetiformis</i> Schlecht. & Cham.	Mexico, Guatemala
Poaceae	
<i>Oryza sativa</i> L.	China
<i>Saccharum officinarum</i> L.	SE Asia
Polygonaceae	
<i>Antigonon leptopus</i> Hook. & Arn.	Mexico
Pontederiaceae	
* <i>Eichhornia crassipes</i> (Mart.) Solms	Amazon basin
Scrophulariaceae	
+ <i>Buddleja davidii</i> Franch.	central China, Japan
Solanaceae	
* <i>Nicotiana tabacum</i> L.	tropical and subtropical America
Sterculiaceae	
* <i>Theobroma cacao</i> L.	Central and South America
Urticaceae	
<i>Pellionia pulchra</i> N.E. Br.	S Vietnam
<i>Soleirolia soleirolii</i> (Req.) Dandy	N Mediterranean region around Italy and nearby islands
Verbenaceae	
<i>Clerodendrum thomsoniae</i> Balf.	tropical W Africa
+ <i>Lantana camara</i> L.	American tropics
Vitaceae	
<i>Cissus discolor</i> Blume	temperate and tropical SE Asia
Zingiberaceae	
<i>Alpinia purpurata</i> K.Schum.	Malaysia
* <i>Hedychium</i> J.Koenig sp.	SE Asia

¹Some plants (+) are only temporarily placed to the Victoria greenhouse of BG PJŠU in time of butterfly exhibitions. These plants (potted ones or their cuttings with flowers) should serve as additional food sources (nectar) for adult butterflies. Other plants (majority) have been within permanent compositions. However some changes took place during the history of living butterfly exhibits (some plant taxa have been removed (*) and other ones have been additionally planted). This table is a complete list of all registered plant taxa registered in this greenhouse between the years 2008 and 2015. Actual situation (2015) in the Victoria greenhouse is represented by remaining plant taxa without marks (+, *).

²Data on distribution of individual taxa in the wild are adapted according to HASSLER (2015a, 2015b)




Fig. 2. All 182 lepidopteran species in exhibits in BG PJŠU. The order (from top left to bottom right, row by row) is the same as in Tab. 3.

Tab. 3. Butterflies and moths in BG PJŠU during annual exhibitions from 2008 to 2015.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Geometridae			
Geometrinae			
Dysphaniini			
1. <i>Dysphania malayanus</i> (Guérin-Méneville, 1843)	Peninsular Malaysia, Sumatra, Borneo, Palawan	Rhizophoraceae (<i>Carallia</i>)	HOLLOWAY (1996)
2. <i>Dysphania militaris</i> (Linnaeus, 1758)	tropical regions of S and SE Asia: India, S China, Thailand, Sumatra, Borneo, Bali	Myrtaceae (<i>Rhodomyrtus</i>), Rhizophoraceae (<i>Carallia</i> , <i>Kandelia</i>)	HOLLOWAY (1996), ROBINSON et al (2010), SVELA (2015)
3. <i>Dysphania transducta</i> (Walker, 1861)	Myanmar, Peninsular Malaysia, Sumatra, Borneo	Rhizophoraceae (<i>Carallia</i>)	HOLLOWAY (1996)
Lycaenidae			
Lycaeninae			
Eumaeini			
4. <i>Eumaeus minyas</i> (Hübner, 1809)	Colombia, Peru, Bolivia, C Brazil	Zamiaceae (<i>Zamia</i>)	PYLE (1981), WARREN et al (2013), SVELA (2015)
Nymphalidae			
Biblidinae			
Ageroniini			
5. <i>Hamadryas amphinome</i> (Linnaeus, 1767)	Mexico, Guatemala, Costa Rica, Panama, Colombia, Brazil, Guianas, Bolivia, Peru, Cuba	Aristolochiaceae (<i>Aristolochia</i>), Asteraceae (<i>Eupatorium</i>), Euphorbiaceae (<i>Dalechampia</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
6. <i>Hamadryas feronia</i> (Linnaeus, 1758)	S USA, Mexico, Guatemala, Honduras, Costa Rica, Venezuela, Colombia, Suriname, Ecuador, Peru, Brazil, Paraguay, Trinidad	Arecaceae (<i>Syagrus</i>), Euphorbiaceae (<i>Dalechampia</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
7. <i>Hamadryas laodamia</i> (Cramer, 1777)	Mexico, Honduras, Costa Rica, Panama, Colombia, Venezuela, Suriname, Bolivia, Brazil, Peru, Trinidad	Euphorbiaceae (<i>Dalechampia</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Biblidini			
8. <i>Biblis hyperia</i> (Cramer, 1779)	SW USA, Mexico, El Salvador, Costa Rica, Ecuador, Peru, Brazil, Paraguay, Caribbean islands (Hispaniola, Mona, Puerto Rico, Culebra, Virgin Islands, St. Lucia, Montserrat, Guadeloupe)	Euphorbiaceae (<i>Acidoton</i> , <i>Tragia</i>), Urticaceae (<i>Laportea</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
Epicaliini			
9. <i>Catonephele numilia</i> (Cramer, 1775)	Mexico, El Salvador, Costa Rica, Panama, Colombia, Venezuela, Suriname, Ecuador, Peru, Brazil, Argentina, Paraguay, Trinidad	Euphorbiaceae (<i>Alchornea</i> , <i>Aparisthmium</i> , <i>Conceveiba</i>), Lauraceae (<i>Nectandra</i>), Verbenaceae (<i>Citharexylum</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
10. <i>Catonephele orites</i> Stichel, 1899	Costa Rica, Panama, Colombia, Ecuador	Euphorbiaceae (<i>Alchornea</i>)	ROBINSON et al (2010), WARREN et al (2013)
11. <i>Myscelia cyaniris</i> Doubleday, 1848	Mexico, Honduras, Costa Rica, Panama, Venezuela, Ecuador, Peru	Euphorbiaceae (<i>Adelia</i> , <i>Dalechampia</i>)	SAVELA (2015)
12. <i>Myscelia ethusa</i> (Doyère, 1840)	S Texas, Mexico, Guatemala, Honduras, Nicaragua, Costa Rica	Euphorbiaceae (<i>Dalechampia</i>)	RIES (2012), WARREN et al (2013)
13. <i>Nessaea aglaura</i> (Doubleday, 1848)	Mexico, Costa Rica, Panama, Colombia, Venezuela, Ecuador	Asteraceae (<i>Mikania</i>), Euphorbiaceae (<i>Alchornea</i> , <i>Dalechampia</i> , <i>Plukenetia</i>), Rhamnaceae (<i>Gouania</i>)	ROBINSON et al (2010), WARREN et al (2013), MORISSON (2014), SAVELA (2015)
Danainae			
Danaini			
14. <i>Danaus chrysippus</i> (Linnaeus, 1758)	Tunisia, Algeria, Morocco, Egypt, Mauritania, Senegal, Gambia, Guinea-Bissau, Guinea, Sierra Leone, Liberia, Mali, Ivory Coast, Ghana, Burkina Faso, Togo, Benin, Nigeria, Niger, Cameroon, Equatorial Guinea, Gabon, Congo, Angola, Democratic Republic of Congo, Central African Republic, Sudan, Uganda, Ethiopia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa,	Apocynaceae (<i>Apocynum</i> , <i>Araujia</i> , <i>Asclepias</i> , <i>Aspidoglossum</i> , <i>Calotropis</i> , <i>Caralluma</i> , <i>Ceropegia</i> , <i>Cryptolepis</i> , <i>Cynanchum</i> , <i>Gomphocarpus</i> , <i>Huernia</i> , <i>Ischnostemma</i> , <i>Kanahia</i> , <i>Leichardtia</i> , <i>Leptadenia</i> , <i>Marsdenia</i> , <i>Metaplexis</i> , <i>Orbea</i> , <i>Oxystelma</i> , <i>Pachycarpus</i> , <i>Pentarrhinum</i> , <i>Pentatropis</i> , <i>Pergularia</i> , <i>Periploca</i> , <i>Pleurostelma</i> , <i>Raphistemma</i> , <i>Secamone</i> , <i>Schizoglossum</i> , <i>Stapelia</i> , <i>Stathmostelma</i> , <i>Tylophora</i> , <i>Xysmalobium</i>), Convolvulaceae (<i>Ipomoea</i>),	WILLIAMS (2008), VAN DER HEYDEN (2010), ROBINSON et al (2010), GOSWAMI (2013), INAYOSHI (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
	Lesotho, Swaziland, Madagascar, Comoro Islands, Mauritius, Rodrigues, Reunion, Bourbon, Seychelles, Aldabra, St Helena; Arabia (Yemen, Saudi Arabia, Oman, United Arab Emirates), Iran, Iraq, Palestine, Lebanon, Turkey, India, Sri Lanka, Andaman Islands, Pakistan, Afghanistan, China, Taiwan, Japan, Malaysia, Thailand, Laos, Myanmar, Vietnam, Philippines, Borneo, Java, Sulawesi, Lesser Sunda Islands, Timor, New Guinea, Australia, Fiji, New Hebrides, Cyprus, Malta, Greece, Italy, Spain, Cape Verde Islands, Canary Islands,	Euphorbiaceae (<i>Euphorbia</i>), Malvaceae, Moraceae (<i>Ficus</i>), Plumbaginaceae (<i>Dyerophytum</i>), Poaceae, Rosaceae (<i>Rosa</i>), Sapindaceae (<i>Lepisanthes</i>), Scrophulariaceae (<i>Antirrhinum</i>).	
15. <i>Danaus plexippus</i> (Linnaeus, 1758)	native to North America (Canada, USA, Mexico), Central America (Nicaragua), South America (Venezuela, French Guiana, Suriname, Guyana, Ecuador, Peru, Brazil), Trinidad, Tobago, Puerto Rico, Cuba, Hispaniola, Bahamas, Virgin Islands, Caymans, Galapagos; expanded to Australia, New Zealand, many Pacific islands (e.g. Hawaii, Samoa, Fiji, New Caledonia), Indian ocean islands (Mauritius, Reunion) and a few places in Europe (Portugal, Spain, Canary Islands) and Africa (Morocco)	Apocynaceae (<i>Apocynum</i> , <i>Araujia</i> , <i>Asclepias</i> , <i>Calotropis</i> , <i>Ceropegia</i> , <i>Cynanchum</i> , <i>Gomphocarpus</i> , <i>Gonolobus</i> , <i>Marsdenia</i> , <i>Matelea</i> , <i>Orbea</i> , <i>Oxypetalum</i> , <i>Oxystelma</i> , <i>Raphistemma</i> , <i>Sarcostemma</i> , <i>Stapelia</i> , <i>Stephanotis</i>), Convolvulaceae (<i>Ipomoea</i>), Euphorbiaceae (<i>Euphorbia</i>), Malvaceae (<i>Gossypium</i>), Rutaceae (<i>Citrus</i>), Sapotaceae (<i>Manilkara</i>)	BROWER (2007), WILLIAMS (2008), ROBINSON et al (2010), WARREN et al (2013), ZHAN et al (2014), SAVELA (2015)
16. <i>Euploea core</i> (Cramer, 1780)	India, Sri Lanka, Nepal, S China, Taiwan, Malaysia, Myanmar, Thailand, Laos, Cambodia, Vietnam, Sumatra, Borneo, Sulawesi, Java, Bali, Bismarck Archipelago, Solomon Islands, Andaman Islands, Nicobars, New Guinea, Australia	Apocynaceae (<i>Adenium</i> , <i>Allamanda</i> , <i>Anodendron</i> , <i>Apocynum</i> , <i>Asclepias</i> , <i>Brachystelma</i> , <i>Calotropis</i> , <i>Carissa</i> , <i>Cerbera</i> , <i>Ceropegia</i> , <i>Cryptolepis</i> , <i>Cryptostegia</i> , <i>Cynanchum</i> , <i>Gomphocarpus</i> , <i>Gunnessia</i> , <i>Gymnanthera</i> , <i>Hemidesmus</i> , <i>Holarthena</i> , <i>Hoya</i> , <i>Ichnocarpus</i> , <i>Leichardtia</i> , <i>Mandevilla</i> , <i>Marsdenia</i> , <i>Nerium</i> , <i>Parsonsia</i> , <i>Plumeria</i> , <i>Rhynchodia</i> , <i>Sarcolobus</i> , <i>Sarcostemma</i> , <i>Secamone</i> , <i>Stephanotis</i> , <i>Toxocarpus</i> , <i>Trachelospermum</i> , <i>Tylophora</i>), Cannabidaceae (<i>Aphananthe</i>), Moraceae (<i>Ficus</i> , <i>Streblus</i>), Rubiaceae (<i>Morinda</i>), Ulmaceae (<i>Aphananthe</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
17. <i>Euploea mulciber</i> (Cramer, 1777)	India, Bhutan, Myanmar, Thailand, Laos, Cambodia, Vietnam, S China, Taiwan, Philippines, Malaysia, Sumatra, Borneo, Sulawesi, Java, Bali	Apocynaceae (<i>Ichnocarpus</i> , <i>Marsdenia</i> , <i>Nerium</i> , <i>Pottsia</i> , <i>Strophanthus</i> , <i>Toxocarpus</i>), Aristolochiaceae (<i>Aristolochia</i>), Convolvulaceae (<i>Argyreia</i>), Moraceae (<i>Ficus</i> , <i>Streblus</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015),
18. <i>Euploea phaenareta</i> (Schaller, 1785)	Sri Lanka, India, Burma, Thailand, Vietnam, the Malay Peninsula, Sumatra, Java, Borneo, Philippines, Moluccas, New Guinea, Australia, Bismarck Islands	Apocynaceae (<i>Cerbera</i> , <i>Plumeria</i>), Moraceae (<i>Ficus</i>)	ROBINSON et al (2010), DAY (2015), SAVELA (2015)
19. <i>Idea leuconoe</i> Erichson, 1834	W Malaysia, Thailand, Taiwan, Japan, Sumatra, Borneo, Java, Singapore, Philippines	Apocynaceae (<i>Cynanchum</i> , <i>Parsonsia</i> , <i>Tylophora</i>)	ROBINSON et al (2010), HOSKINS (2015), SAVELA (2015)
20. <i>Ideopsis juvena</i> (Cramer, 1777)	Malay Peninsula, Sumatra, Java, Borneo, Sulawesi, Lesser Sunda Islands, Philippines, Moluccas, New Guinea, Bismarck Archipelago, Solomon Islands	Apocynaceae (<i>Cynanchum</i> , <i>Gymnema</i> , <i>Heterostemma</i> , <i>Parsonsia</i> , <i>Pergularia</i> , <i>Telosma</i>), Piperaceae (<i>Piper</i>)	VANE-WRIGHT & DE JONG (2003), SARI et al (2015), SAVELA (2015)
21. <i>Ideopsis similis</i> (Linnaeus, 1758)	Sri Lanka, Vietnam, Thailand, Laos, Cambodia, Malaysia, Myanmar, Nicobars, Sumatra, S China, Taiwan, Japan	Apocynaceae (<i>Cryptolepis</i> , <i>Cynanchum</i> , <i>Gymnema</i> , <i>Marsdenia</i> , <i>Parsonsia</i> , <i>Tylophora</i> , <i>Vincetoxicum</i>), Menispermaceae (<i>Cocculus</i> , <i>Diploclisia</i>)	ROBINSON et al (2010), DAY (2015), INAYOSHI (2015), SAVELA (2015)
22. <i>Lycorea halia</i> (Hübner, 1816)	S USA, Mexico, El Salvador, Belize, Colombia, Venezuela, Suriname, Ecuador, Peru, Brazil, larger Caribbean islands (Cuba, Hispaniola, Jamaica, Puerto Rico, St. Lucia, Trinidad)	Araliaceae (<i>Hedera</i>), Apocynaceae (<i>Asclepias</i>), Caricaceae (<i>Carica</i> , <i>Jacaratia</i>), Hemerocallidaceae (<i>Hemerocallis</i>), Moraceae (<i>Ficus</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
23. <i>Tirumala septentrionis</i> (Butler, 1874)	India, Sri Lanka, Nepal, Bhutan, China, Taiwan, Myanmar, Laos, Cambodia, Vietnam, Thailand, Malaysia, Singapore, Java, Borneo, Lesser Sunda Islands,, Philippines,	Apocynaceae (<i>Cosmostigma</i> , <i>Dregea</i> , <i>Heterostemma</i> , <i>Parsonsia</i> , <i>Vallis</i> , <i>Tylophora</i>), Menispermaceae (<i>Cocculus</i>)	NAIR (2005), ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
Ithomiini			
24. <i>Godyrus nero</i> (Hewitson, 1855)	Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Ecuador	Solanaceae (<i>Cestrum</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
25. <i>Mechanitis polymnia</i> (Linnaeus, 1758)	Mexico, Guatemala, Honduras, Belize, Costa Rica, Panama, Colombia, Venezuela, Suriname, Ecuador, Bolivia, Peru, Brazil, Trinidad	Apocynaceae, Solanaceae (<i>Brugmansia</i> , <i>Brunfelsia</i> , <i>Cyphomandra</i> , <i>Lycopersicon</i> , <i>Solanum</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
26. <i>Methona confusa</i> Butler, 1873	Panama, Colombia, Venezuela, Guayana, Ecuador, Peru, Bolivia, Brazil, Argentina	Solanaceae (<i>Brunfelsia</i>)	BECCALONI (1995), ROBINSON et al (2010), SAVELA (2015)
27. <i>Tithorea harmonia</i> (Cramer, 1777)	Mexico, Honduras, El Salvador, Guatemala, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Guayana, Suriname, French Guiana, Ecuador, Brazil, Peru, Bolivia, Trinidad	Apocynaceae (<i>Echites</i> , <i>Mandevilla</i> , <i>Mesechites</i> , <i>Prestonia</i>)	FREITAS et al (2001), ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
28. <i>Tithorea tarricina</i> Hewitson, 1858	Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia	Apocynaceae (<i>Echites</i> , <i>Prestonia</i>), Solanaceae (<i>Markea</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
Heliconiinae			
Acraeini			
29. <i>Acraea violae</i> (Fabricius, 1793)	India, Sri Lanka, Bangladesh, Myanmar, Thailand, Laos, Cambodia, Vietnam	Cucurbitaceae, Loganiaceae, Malvaceae (<i>Hibiscus</i>), Moraceae (<i>Castilla</i>), Passifloraceae (<i>Adenia</i> , <i>Passiflora</i>), Turneraceae (<i>Turnera</i>), Verbenaceae (<i>Vitex</i>), Violaceae (<i>Hybanthus</i> , <i>Viola</i>)	DAS et al (2010), ROBINSON et al (2010), KHOT & GAIKWAD (2011), HOSKINS (2015), INAYOSHI (2015),
30. <i>Cethosia biblis</i> (Drury, 1773)	India, Bhutan, Nepal, China, Thailand, Laos, Vietnam, Myanmar, Malaysia, Philippines, Sumatra, Borneo, Java, Sulawesi, Bali, Ambon, Serang, Moluccas, Andaman Islands	Passifloraceae (<i>Adenia</i> , <i>Passiflora</i>)	VANE-WRIGHT & DE JONG, ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
31. <i>Cethosia cyane</i> (Drury, 1773)	India, Bhutan, Myanmar, S China, Thailand, Laos, Cambodia, Vietnam	Commelinaceae (<i>Commelina</i>), Convolvulaceae (<i>Ipomoea</i>), Euphorbiaceae (<i>Ricinus</i>), Passifloraceae (<i>Adenia</i> , <i>Passiflora</i>), Verbenaceae (<i>Duranta</i>)	ROBINSON et al (2010), INAYOSHI (2015), NEUBAUER (2015), SAVELA (2015)
Heliconiini			
32. <i>Agraulis vanillae</i> (Linnaeus, 1758)	S USA, Mexico, Costa Rica, Panama, Venezuela, Suriname, French Guiana, Peru, Brazil, Paraguay, Uruguay, Argentina, Antilles, Bahamas, Galapagos, Hawaii	Caprifoliaceae (<i>Lonicera</i>), Passifloraceae (<i>Passiflora</i>), Tiliaceae (<i>Corchorus</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
33. <i>Dione juno</i> (Cramer, 1779)	Mexico, Guatemala, Nicaragua, Costa Rica, Panama, Suriname, Ecuador, Peru, Paraguay, Antilles	Passifloraceae (<i>Passiflora</i>).	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
34. <i>Dryadula phaetusa</i> (Linnaeus, 1758)	S USA, Mexico, Nicaragua, Costa Rica, Panama, Suriname, French Guiana, Brazil, Bolivia	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015)
35. <i>Dryas iulia</i> (Fabricius, 1775)	S USA, Mexico, Honduras, Costa Rica, Suriname, Ecuador, Bolivia, Caribbean Islands (Cuba, Hispaniola, Jamaica, Puerto Rico, St. Thomas, Montserrat, Guadeloupe, St. Lucia, S. Vincent, St. Kitts, Grenada, Martinique, Bahamas, Cayman Islands and others)	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SVELA (2015)
36. <i>Eueides isabella</i> (Stoll, 1781)	S USA, Mexico, Honduras, Guatemala, Nicaragua, Panama, Colombia, Venezuela, Suriname, Ecuador, Peru, Brazil, West Indies (Cuba, Puerto Rico, Hispaniola)	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
97 37. <i>Heliconius antiochus</i> (Linnaeus, 1767)	Panama, Colombia, Venezuela, Guayana, Suriname, French Guiana, Peru, Bolivia, Brazil	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015)
38. <i>Heliconius atthis</i> Doubleday, 1847	Ecuador	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015)
39. <i>Heliconius charithonia</i> (Linnaeus, 1767)	S USA, Mexico, Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Caribbean Islands (Cuba, Hispaniola, Jamaica, Bahamas, Puerto Rico, Virgin Islands, St. Kitts, Antigua and others)	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), CAST (2015)
40. <i>Heliconius cydno</i> (Doubleday, 1847)	Mexico, Guatemala, Belize, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015)
41. <i>Heliconius erato</i> (Linnaeus, 1758)	S USA, Mexico, Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia, Guayana, Suriname, French Guiana, Brazil, Paraguay, Argentina, Uruguay, Trinidad, Tobago	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), CAST (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
42. <i>Heliconius hecale</i> (Fabricius, 1776)	Mexico, Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia, Guayana, Suriname, French Guiana, Brazil	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015)
43. <i>Heliconius himera</i> Hewitson, 1867	Ecuador, Peru	Passifloraceae (<i>Passiflora</i>)	JIGGINS (2008), CAST (2015)
44. <i>Heliconius ismenius</i> Latreille, 1817	Mexico, Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, French Guiana	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), CAST (2015)
45. <i>Heliconius melpomene</i> (Linnaeus, 1758)	Mexico?, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia, Guayana, Suriname, French Guiana, Brazil, Trinidad, Tobago	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015), SAVELA (2015)
46. <i>Heliconius numata</i> (Cramer, 1780)	Guatemala?, Colombia, Venezuela, Ecuador, Peru, Bolivia, Guayana, Suriname, French Guiana, Brazil	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015), SAVELA (2015)
47. <i>Heliconius sara</i> (Fabricius, 1793)	Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia, Guayana, Suriname, French Guiana, Brazil, Argentina, Trinidad	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), CAST (2015), SAVELA (2015)
48. <i>Laparus doris</i> (Linnaeus, 1771)	Mexico, Honduras, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Guayana, Suriname, French Guiana, Peru, Brazil, Bolivia, Trinidad	Passifloraceae (<i>Passiflora</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
49. <i>Philaethria dido</i> (Linnaeus, 1763)	Mexico, Nicaragua, Panama, Colombia, Guianas?, Brazil	Passifloraceae (<i>Passiflora</i>)	GODMAN & SALVIN (1879-1901), ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
Vagrantini			
50. <i>Phalanta phalantha</i> (Drury, 1773)	SW Arabia, Socotra, India, Sri Lanka, S China, Taiwan, Myanmar, Malaysia, Japan, Sumatra, Borneo, Sulawesi, Java, Lesser Sunda Islands,	Acanthaceae (<i>Barleria</i>), Anacardiaceae (<i>Mangifera</i>), Asteraceae (<i>Tridax</i>), Bixaceae, Celastraceae (<i>Maytenus</i>), Flacourtiaceae (<i>Dovyalis</i> , <i>Flacourtia</i> ,	VANE-WRIGHT & DE JONG (2003), WILLIAMS (2008),

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources	
	Philippines, N Australia, Senegal, Gambia, Guinea-Bissau, Guinea, Sierra Leone, Burkina Faso, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Gabon, Congo, Central African Republic, Angola, Democratic Republic of Congo, Sudan, Ethiopia, Uganda, Rwanda, Burundi, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Lesotho, Madagascar, Seychelles, Aldabra, Comoro Islands, Mauritius, Reunion	<i>Oncoba</i> , <i>Trimeria</i> , <i>Xylosma</i>), Loranthaceae (<i>Loranthus</i>), Myrtaceae (<i>Melaleuca</i>), Picrodendraceae (<i>Petalostigma</i>), Primulaceae (<i>Androsace</i>), Rubiaceae (<i>Canthium</i> , <i>Coffea</i> , <i>Ixora</i>), Salicaceae (<i>Populus</i> , <i>Salix</i> , <i>Scolopia</i>), Smilacaceae (<i>Smilax</i>), Violaceae (<i>Viola</i>)	ROBINSON et al (2010), HOSKINS (2015), SVELA (2015)	
51. <i>Vindula dejone</i> (Erichson, 1834)	Thailand, Laos, Cambodia, Vietnam, Malaysia, Philippines, Sumatra, Java, Borneo, Sulawesi, Lesser Sunda Islands, Moluccas	Passifloraceae (<i>Adenia</i> , <i>Passiflora</i>)	VANE-WRIGHT & DE JONG (2003), ROBINSON et al (2010), INAYOSHI (2015), SVELA (2015)	
<hr/>				
66	Charaxinae			
	Anaeini			
	52. <i>Consul electra</i> Westwood, 1850	Mexico, Belize, Guatemala, Nicaragua, Costa Rica, Panama	Piperaceae (<i>Piper</i>)	GODMAN & SALVIN (1879-1901), MUYSHONDT (1976), SVELA (2015)
	53. <i>Consul fabius</i> (Cramer, 1776)	Mexico, Costa Rica, Panama, Colombia, Venezuela, Suriname, French Guiana, Ecuador, Brazil, Peru, Bolivia, Trinidad	Piperaceae (<i>Piper</i>)	MUYSHONDT (1974), WARREN et al (2013), SVELA (2015)
	54. <i>Fountainea eurypyle</i> (C. & R. Felder, 1862)	Mexico, Guatemala, Nicaragua, Costa Rica, Panama, Bolivia, Peru	Euphorbiaceae (<i>Croton</i>), Flacourtiaceae (<i>Casearia</i>)	ROBINSON et al (2010), SVELA (2015)
	55. <i>Fountainea nobilis</i> (Bates, 1864)	Mexico, Guatemala, El Salvador, Costa Rica, Panama, Colombia, Venezuela, Peru	Euphorbiaceae (<i>Croton</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
56. <i>Hypna clytemnestra</i> (Cramer, 1777)	Mexico, Guatemala, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Suriname, Peru, Bolivia, Brazil, Argentina, Cuba	Euphorbiaceae (<i>Croton</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SVELA (2015)	

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Charaxini			
57. <i>Charaxes brutus</i> (Cramer, 1779)	Senegal, Guinea-Bissau, Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Nigeria, Cameroon, Equatorial Guinea, Central African Republic, Gabon, Congo, Angola, Democratic Republic of Congo, Uganda, Sudan, Kenya, Rwanda, Burundi, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland	Euphorbiaceae (<i>Flueggea</i> , <i>Securinega</i>), Meliaceae (<i>Ekebergia</i> , <i>Entandrophragma</i> , <i>Khaya</i> , <i>Lepidotrichilia</i> , <i>Melia</i> , <i>Trichilia</i> , <i>Turraea</i>), Melianthaceae (<i>Bersama</i>), Sapindaceae (<i>Allophylus</i> , <i>Blighia</i>), Tiliaceae (<i>Grewia</i>)	WILLIAMS (2008), ROBINSON et al (2010)
58. <i>Charaxes candiope</i> (Godart, 1824)	Senegal, Gambia, Guinea, Sierra Leone, Ivory Coast, Ghana, Nigeria, Cameroon, Ethiopia, Kenya, Tanzania, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland	Euphorbiaceae (<i>Croton</i>), Poaceae (<i>Pennisetum</i>)	WILLIAMS (2008), ROBINSON et al (2010)
59. <i>Charaxes castor</i> (Cramer, 1775)	Senegal, Gambia, Guinea-Bissau, Guinea, Burkina Faso, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Gabon, Congo, Angola, Central African Republic, Democratic Republic of Congo, Uganda, Ethiopia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, South Africa, Swaziland, Comoro Islands	Celastraceae (<i>Cassine</i> , <i>Elaeodendron</i> , <i>Gymnosporia</i> , <i>Maytenus</i> , <i>Pleurostylia</i>), Euphorbiaceae (<i>Bridelia</i> , <i>Croton</i> , <i>Tragia</i>), Fabaceae (<i>Afzelia</i> , <i>Bauhinia</i> , <i>Brachystegia</i> , <i>Cassia</i> , <i>Copaifera</i> , <i>Entada</i> , <i>Erythrina</i> , <i>Senna</i> , <i>Schotia</i>), Iridaceae (<i>Iris</i>), Lamiaceae (<i>Gmelina</i>), Malvaceae (<i>Hibiscus</i>), Poaceae (<i>Sorghum</i>), Ulmaceae (<i>Chaetachme</i> , <i>Trema</i>)	DICKSON & KROON (1978), WILLIAMS (2008), ROBINSON et al (2010)
60. <i>Charaxes cithaeron</i> C. & R. Felder, 1859	Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, South Africa, Swaziland	Acanthaceae (<i>Chaetacanthus</i> , <i>Crabbea</i>), Celastraceae (<i>Gymnosporia</i> , <i>Hippocratea</i> , <i>Maytenus</i>), Fabaceae (<i>Acacia</i> , <i>Afzelia</i> , <i>Albizia</i> , <i>Baphia</i> , <i>Craibia</i> , <i>Dalbergia</i> , <i>Leptoderris</i> , <i>Millettia</i> , <i>Philenoptera</i> , <i>Schotia</i>), Linaceae (<i>Hugonia</i>), Sapindaceae (<i>Deinbollia</i>), Sterculiaceae (<i>Cola</i>), Tiliaceae (<i>Grewia</i>), Ulmaceae (<i>Celtis</i> , <i>Chaetachme</i> , <i>Trema</i>)	WILLIAMS (2008), ROBINSON et al (2010)
61. <i>Charaxes etesipe</i> (Godart, 1824)	Senegal, Guinea, Sierra Leone, Liberia, Benin, Ivory Coast, Ghana, Togo, Nigeria, Cameroon, Gabon, Congo, Angola, Central African Republic,	Bombacaceae (<i>Bombax</i> , <i>Ceiba</i>), Euphorbiaceae (<i>Croton</i> , <i>Ricinus</i> , <i>Tragia</i>), Erythroxylaceae (<i>Erythroxylum</i>), Fabaceae (<i>Afzelia</i> , <i>Cassia</i> , <i>Dalbergia</i> ,	WOODHALL (2005), WILLIAMS (2008), ROBINSON et al (2010),

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
	Chad, Democratic Republic of Congo, Uganda, Sudan, Rwanda, Tanzania, Ethiopia, Kenya, Somalia, Democratic Republic of Congo, Angola, Mozambique, Malawi, Zimbabwe, Swaziland, South Africa, Madagascar	<i>Entada</i> , Phyllanthaceae (<i>Margaritaria</i> , <i>Phyllanthus</i> , <i>Securinega</i>), Polygalaceae (<i>Securidaca</i>), Rhamnaceae (<i>Maesopsis</i> , <i>Ventilago</i>)	SAVELA (2015)
62. <i>Charaxes protoclea</i> Feisthamel, 1850	Senegal, Gambia, Guinea-Bissau, Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Nigeria, Cameroon, Equatorial Guinea, Central African Republic, Gabon, Congo, Angola, Democratic Republic of Congo, Uganda, Kenya, Tanzania, Zambia, Malawi, Zimbabwe, Mozambique, South Africa	Fabaceae (<i>Azelia</i> , <i>Berlinia</i> , <i>Brachystegia</i> , <i>Bussea</i> , <i>Cassia</i> , <i>Julbernardia</i>), Myrtaceae (<i>Eugenia</i> , <i>Syzygium</i>), Poaceae (<i>Oxytenanthera</i>)	WILLIAMS (2008), ROBINSON et al (2010)
63. <i>Charaxes varanes</i> (Cramer, 1764)	Senegal, Gambia, Guinea-Bissau, Guinea, Burkina Faso, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Equatorial Guinea, Central African Republic, Sudan, Ethiopia, Yemen, Oman, Kenya, Tanzania, Zambia, Angola, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland	Anacardiaceae (<i>Rhus</i>), Sapindaceae (<i>Allophylus</i> , <i>Cardiospermum</i> , <i>Schmidelia</i>)	WILLIAMS (2008), ROBINSON et al (2010)
64. <i>Charaxes violetta</i> Grose-Smith, 1885	Kenya, Tanzania, Malawi, Zimbabwe, Mozambique, South Africa	Fabaceae (<i>Azelia</i> , <i>Brachystegia</i>), Sapindaceae (<i>Blighia</i> , <i>Deinbollia</i> , <i>Paullinia</i>)	WILLIAMS (2008)
Euxanthini			
65. <i>Euxanthe wakefieldi</i> (Ward, 1873)	Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, South Africa, Swaziland	Fabaceae (<i>Azelia</i>), Sapindaceae (<i>Blighia</i> , <i>Deinbollia</i> , <i>Lecaniodiscus</i> , <i>Phialodiscus</i> , <i>Sapindus</i>)	LARSEN (1996), WILLIAMS (2008), ROBINSON et al (2010)
Preponini			
66. <i>Archaeoprepona demophon</i> (Linnaeus, 1758)	USA (Texas), Mexico, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Ecuador, Suriname, Bolivia, Brazil, Paraguay, Cuba, Hispaniola	Annonaceae (<i>Annona</i> , <i>Rollinia</i>), Connaraceae (<i>Rourea</i>), Convolvulaceae (<i>Maripa</i>), Lacistemataceae (<i>Lacistema</i>), Lauraceae (<i>Cinnamomum</i> , <i>Nectandra</i> , <i>Ocotea</i> , <i>Persea</i>), Malpighiaceae (<i>Malpighia</i>), Meliaceae (<i>Guarea</i>), Monimiaceae (<i>Mollinedia</i> , <i>Siparuna</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
67. <i>Prepona laertes</i> (Hübner, 1811)	USA (Florida), Mexico, Guatemala, Honduras, Costa Rica, Panama, Colombia, Venezuela, Peru, Ecuador, Suriname, French Guiana, Bolivia, Brazil, Paraguay, Cuba, Trinidad	Fabaceae (<i>Andira</i> , <i>Inga</i>), Sapindaceae (<i>Melicococcus</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
Limnitiidinae			
Adoliadini			
68. <i>Dophla evelina</i> (Stoll 1790)	India, Sri Lanka, Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Singapore, China; Java, Sumatra, Borneo, Celebes, Sulawesi, Philippines	Anacardiaceae (<i>Anacardium</i>), Ebenaceae (<i>Diospyros</i>), Euphorbiaceae (<i>Antidesma</i>), Fagaceae (<i>Lithocarpus</i>)	ROBINSON et al (2010), INAYOSHI (2015), SAVELA (2015)
69. <i>Euthalia aconthea</i> (Hewitson, 1874)	India, Sri Lanka, Thailand, Laos, Vietnam, Malaysia, S China, Philippines; Sumatra, Borneo, Java, Sulawesi, Bali, Lombok, Andamans	Anacardiaceae (<i>Anacardium</i> , <i>Mangifera</i>), Cucurbitaceae (<i>Bryonia</i>), Loranthaceae (<i>Loranthus</i> , <i>Scurrula</i>), Moraceae (<i>Morus</i> , <i>Streblus</i> , <i>Trophis</i>), Rosaceae (<i>Rosa</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
70. <i>Euthalia monina</i> (Moore, 1859)	India, S China, Thailand, Laos, Cambodia, Vietnam, Myanmar, Malaysia, Singapore, Sumatra, Borneo, Java, Bali, Lombok, Philippines	Dipterocarpaceae (<i>Shorea</i>), Ebenaceae (<i>Diospyros</i>), Euphorbiaceae (<i>Macaranga</i> , <i>Mallotus</i>), Melastomataceae (<i>Clidemia</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
71. <i>Lexias pardalis</i> Moore, 1878	China, Laos, Vietnam, Thailand, Myanmar, India, Singapore, Philippines, Sumatra, Borneo	Hypericaceae (<i>Cratoxylum</i>)	TAN (2010), SAVELA (2015)
72. <i>Tanaecia flora</i> Butler, 1873	Thailand, Myanmar, Peninsular Malaysia	Lecythidaceae (<i>Careya</i>), Melastomataceae (<i>Melastoma</i>)	REIMAN GARDENS (2015), SAVELA (2015)
73. <i>Tanaecia julii</i> Bougainville, 1837	India, Bhutan, Myanmar, Cambodia, Thailand, Laos, Vietnam, China, W Malaysia, Sumatra	Sapotaceae (<i>Diploknema</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
Limnitiidini			
74. <i>Athyma perius</i> (Linnaeus, 1758)	India, S China, Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Java, Sumatra, Lombok	Phyllanthaceae (<i>Glochidion</i> , <i>Phyllanthus</i>)	ROBINSON et al (2010), INAYOSHI (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Parthenini			
75. <i>Parthenos sylvia</i> (Cramer, 1776)	Sri Lanka, India, China, Myanmar, Thailand, Malaysia, Borneo, Sumatra, Sulawesi, Java, Bali, Philippines, Moluccas, New Guinea, Solomon Islands, Bismarck Archipelago,	Cucurbitaceae, Passifloraceae (<i>Adenia</i> , <i>Passiflora</i>), Mennispermaceae (<i>Tinospora</i>)	VANE-WRIGHT & DE JONG (2003), ROBINSON et al (2010), HOSKINS (2015)
Nymphalinae			
Coeini			
76. <i>Historis odius</i> Fabricius, 1775	S USA, Mexico, Guatemala, Belize, Costa Rica, Suriname, Peru, Bolivia, N Argentina, Greater Antilles (Hispaniola, Puerto Rico), Windward and Leeward Islands, Guadeloupe	Bignoniaceae (<i>Tabebuia</i>), Fabaceae (<i>Inga</i>), Ulmaceae (<i>Celtis</i> , <i>Trema</i>), Urticaceae (<i>Cecropia</i> , <i>Urtica</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SVELA (2015)
Junoniini			
77. <i>Hypolimnias antevorta</i> (Distant, 1880)	NE Tanzania	Urticaceae (<i>Laportea</i> , <i>Urera</i>)	WILLIAMS (2008), ROBINSON et al (2010)
78. <i>Hypolimnias bolina</i> (Linnaeus, 1758)	S Arabia, Socotra, India, Sri Lanka, S China, Taiwan, Japan, Myanmar, Thailand, Laos, Vietnam, Malaysia, Sumatra, Java, Borneo, Sulawesi, Lesser Sunda Islands, Timor, Moluccas, Philippines, New Guinea, Bismarck Archipelago, Solomon Islands, Christmas Island, Australia, New Zealand, New Caledonia, Fiji, French Polynesia, Madagascar	Apocynaceae, Acanthaceae (<i>Asystasia</i> , <i>Blechum</i> , <i>Eranthemum</i> , <i>Justicia</i> , <i>Pseuderanthemum</i> , <i>Rostellularia</i> , <i>Ruellia</i>), Amaranthaceae (<i>Achyranthes</i> , <i>Alternanthera</i>), Asteraceae (<i>Eclipta</i> , <i>Synedrella</i>), Commelinaceae (<i>Commelina</i>), Convolvulaceae (<i>Ipomoea</i> , <i>Merremia</i>), Fabaceae (<i>Desmodium</i> , <i>Phaseolus</i> , <i>Vigna</i>), Lamiaceae (<i>Perilla</i>), Malvaceae (<i>Abutilon</i> , <i>Malvastrum</i> , <i>Sida</i>), Moraceae (<i>Ficus</i>), Polygonaceae (<i>Persicaria</i> , <i>Polygonum</i>), Portulacaceae (<i>Portulaca</i>), Rubiaceae (<i>Richardia</i>), Tiliaceae (<i>Triumfetta</i>), Urticaceae (<i>Boehmeria</i> , <i>Elatostema</i> , <i>Laportea</i> , <i>Pipturus</i> , <i>Urtica</i>)	VANE-WRIGHT & DE JONG (2003), RAJAGOPALAN (2005), ROBINSON et al (2010), INAYOSHI (2015), SVELA (2015)
79. <i>Hypolimnias misippus</i> (Linnaeus, 1764)	Senegal, Gambia, Guinea-Bissau, Guinea, Mali, Sierra Leone, Liberia, Ivory Coast, Burkina Faso, Ghana, Togo, Benin, Nigeria, Niger, Chad, Cameroon, Equatorial Guinea, Gabon, Congo,	Acanthaceae (<i>Asystasia</i> , <i>Blepharis</i> , <i>Dyschoriste</i> , <i>Justicia</i> , <i>Pseuderanthemum</i> , <i>Ruellia</i>), Amaranthaceae (<i>Amaranthus</i>), Arecaceae (<i>Elaeis</i>), Convolvulaceae (<i>Ipomea</i>), Crassulaceae (<i>Sedum</i>),	VANE-WRIGHT & DE JONG (2003), WILLIAMS (2008), ROBINSON et al (2010), SVELA (2015),

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources	
	Central African Republic, Angola, Democratic Republic of Congo, Sudan, Uganda, Rwanda, Burundi, Ethiopia, Somalia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Lesotho, Arabia (Yemen, Saudi Arabia, Oman, United Arab Emirates), Madagascar, Comoro Islands, Mauritius, Rodrigues, Reunion, Seychelles; India, Sri Lanka, Myanmar, Thailand, Laos, Vietnam, Taiwan, Japan, Indonesia (Java, Sulawesi, Moluccas), Australia; colonised Canary Islands, USA (Florida), West Indies, French Guiana, NE Brazil; migration into Egypt, Lebanon, Turkey	Fabaceae (<i>Vigna</i>), Malvaceae (<i>Abelmoschus</i> , <i>Abutilon</i> , <i>Gossypium</i> , <i>Hibiscus</i>), Moraceae (<i>Ficus</i>), Plantaginaceae (<i>Plantago</i>), Portulacaceae (<i>Portulaca</i> , <i>Talinum</i>), Rosaceae (<i>Prunus</i>), Urticaceae (<i>Elatostema</i>)	HERBISON-EVANS & CROSSLEY (2015), HOSKINS (2015), INAYOSHI (2015)	
104	80. <i>Hypolimnas salmacis</i> (Drury, 1773)	Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Gabon, Congo, Angola, Equatorial Guinea, São Tomé and Príncipe, Democratic Republic of Congo, Uganda, Sudan, Ethiopia, Kenya, Tanzania	Urticaceae (<i>Laportea</i> , <i>Urera</i>)	WILLIAMS (2008), ROBINSON et al (2010)
	81. <i>Junonia almana</i> (Linnaeus, 1758)	India, Sri Lanka, Thailand, Laos, Cambodia, Vietnam, Andaman and Nicobar Islands, Myanmar, Malaysia, Sumatra, Borneo, Sulawesi, Java, Philippines, S China, Taiwan, Japan	Acanthaceae (<i>Acanthus</i> , <i>Barleria</i> , <i>Blechnum</i> , <i>Dyschoriste</i> , <i>Hemigraphis</i> , <i>Hygrophila</i> , <i>Ruellia</i> , <i>Strobilanthes</i>), Amaranthaceae (<i>Alternanthera</i>), Balsaminaceae (<i>Hydrocera</i>), Gesneriaceae (<i>Gloxinia</i>), Fabaceae (<i>Mimosa</i>), Linderniaceae (<i>Lindernia</i>), Melastomataceae (<i>Osbeckia</i>), Onagraceae (<i>Ludwigia</i>), Plantaginaceae (<i>Plantago</i>), Poaceae (<i>Oryza</i> , <i>Pennisetum</i>), Scrophulariaceae (<i>Antirrhinum</i> , <i>Mimulus</i>), Verbenaceae (<i>Lippia</i> , <i>Phyla</i> , <i>Stachytarpheta</i>)	VANE-WRIGHT & DE JONG (2003), ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
	82. <i>Junonia atlites</i> (Linnaeus, 1763)	India, Sri Lanka, Nepal, China, Thailand, Laos, Cambodia, Vietnam, Myanmar, W Malaysia, Sumatra, Java, Borneo, Sulawesi, Lesser Sunda Islands, Moluccas	Acanthaceae (<i>Barleria</i> , <i>Blechnum</i> , <i>Hygrophila</i> , <i>Justicia</i> , <i>Lepidagathis</i> , <i>Nelsonia</i> , <i>Pseuderanthemum</i> , <i>Ruellia</i> , <i>Strobilanthes</i>), Amaranthaceae (<i>Achyranthes</i> , <i>Alternanthera</i>), Poaceae (<i>Oryza</i>), Linderniaceae (<i>Lindernia</i>), Verbenaceae (<i>Phyla</i>)	VANE-WRIGHT & DE JONG (2003), ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
83. <i>Junonia iphita</i> (Cramer, 1779)	Sri Lanka, India, China, Taiwan, Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Sumatra, Borneo, Java, Bali, Lesser Sunda Islands, Philippines, Maldives	Acanthaceae (<i>Hygrophila</i> , <i>Justicia</i> , <i>Lepidagathis</i> , <i>Strobilanthes</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SVELA (2015)
84. <i>Junonia oenone</i> (Linnaeus, 1758)	Senegal, Gambia, Guinea-Bissau, Guinea, Mali, Sierra Leone, Liberia, Ivory Coast, Burkina Faso, Ghana, Togo, Benin, Nigeria, Niger, Cameroon, Ethiopia, Tanzania, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Lesotho, S Arabia, Madagascar, Comoro Islands, Seychelles	Acanthaceae (<i>Adhatoda</i> , <i>Asystasia</i> , <i>Barleria</i> , <i>Brillantaisia</i> , <i>Dicliptera</i> , <i>Eremomastax</i> , <i>Hypoestes</i> , <i>Isoglossa</i> , <i>Justicia</i> , <i>Mackaya</i> , <i>Phaulopsis</i> , <i>Ruellia</i>)	LARSEN (1996), WILLIAMS (2008), ROBINSON et al (2010)
85. <i>Junonia orithya</i> (Linnaeus, 1758)	Senegal, Gambia, Guinea-Bissau, Guinea, Mali, Sierra Leone, Liberia, Ivory Coast, Burkina Faso, Ghana, Togo, Benin, Nigeria, Niger, Tanzania, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Lesotho, Arabia, Madagascar, Seychelles, India, Sri Lanka, S China, Myanmar, Thailand, Malaysia, Singapore, Sumatra, Java, Borneo, Sulawesi, Lesser Sunda Islands, Timor, Philippines, Moluccas, New Guinea, New Britain, N Australia	Acanthaceae (<i>Acanthus</i> , <i>Asystasia</i> , <i>Barleria</i> , <i>Brunoniella</i> , <i>Hygrophila</i> , <i>Hypoestes</i> , <i>Justicia</i> , <i>Lepidagathis</i> , <i>Pseuderanthemum</i> , <i>Rostellularia</i> , <i>Thunbergia</i>), Annonaceae (<i>Annona</i>), Convolvulaceae (<i>Convolvulus</i> , <i>Ipomoea</i>), Lamiaceae (<i>Englerastrum</i> , <i>Plectranthus</i>), Orobanchaceae (<i>Graderia</i> , <i>Cycnium</i> , <i>Lindenbergia</i>), Plantaginaceae (<i>Plantago</i>), Scrophulariaceae (<i>Angelonia</i> , <i>Antirrhinum</i> , <i>Buchnera</i> , <i>Limosella</i> , <i>Misopates</i> , <i>Scrophularia</i> , <i>Striga</i>), Verbenaceae (<i>Lippia</i> , <i>Phyla</i> , <i>Stachytarpheta</i>), Violaceae (<i>Viola</i>)	WILLIAMS (2008), ROBINSON et al (2010), SVELA (2015)
86. <i>Precis octavia</i> (Cramer, 1777)	Senegal, Guinea, Burkina Faso, Sierra Leone, Liberia, Ghana, Togo, Benin, Nigeria, Cameroon, Gabon, Congo, Central African Republic, Sudan, Ethiopia, Somalia, Uganda, Angola, Democratic Republic of Congo, Kenya, to Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland	Fabaceae (<i>Eriosema</i>), Lamiaceae (<i>Coleus</i> , <i>Basilicum</i> , <i>Englerastrum</i> , <i>Isodon</i> , <i>Platostoma</i> , <i>Plectranthus</i> , <i>Pycnostachys</i> , <i>Rabdosiella</i> , <i>Solenostemon</i> , <i>Tetradenia</i>)	WILLIAMS (2008), ROBINSON et al (2010)
87. <i>Protogoniomorpha anacardii</i> (Linnaeus, 1758)	Sierra Leone, Ivory Coast, Ghana, Togo, Nigeria, Central African Republic, Democratic Republic of Congo, Ethiopia, Kenya, Tanzania, Zambia, Mozambique, Zimbabwe, Botswana, South Africa, Swaziland, Yemen, Madagascar	Acanthaceae (<i>Asystasia</i> , <i>Brillantaisia</i> , <i>Eremomastax</i> , <i>Hypoestes</i> , <i>Isoglossa</i> , <i>Justicia</i> , <i>Mimulopsis</i> , <i>Paulowilhelmia</i> , <i>Phaulopsis</i>)	LARSEN (1996), WILLIAMS (2008), ROBINSON et al (2010)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
88. <i>Protogoniomorpha parhassus</i> (Drury, 1782)	Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Gabon, Congo, Central African Republic, Angola, Democratic Republic of Congo, Sudan, Uganda, Rwanda, Burundi, Ethiopia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Namibia, South Africa, Swaziland	Acanthaceae (<i>Asystasia</i> , <i>Brillantaisia</i> , <i>Dicliptera</i> , <i>Eremomastax</i> , <i>Isoglossa</i> , <i>Justicia</i> , <i>Mimulopsis</i> , <i>Paulowilhelmia</i>)	LARSEN (1996), WILLIAMS (2008), ROBINSON et al (2010)
89. <i>Salamis cacta</i> (Fabricius, 1793)	Senegal, Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Gabon, Congo, Central African Republic, Angola, Democratic Republic of Congo, Uganda, Rwanda, Ethiopia, Kenya, Tanzania, Malawi, Mozambique, Zimbabwe	Acanthaceae (<i>Asystasia</i>), Urticaceae (<i>Ureca</i>)	WILLIAMS (2008), ROBINSON et al (2010)
Kallimini			
106 90. <i>Doleschallia bisaltide</i> (Cramer, 1777)	India, Sri Lanka, S China, Myanmar, Thailand, Laos, Vietnam, W Malaysia, Sumatra, Borneo, Palawan, Sulawesi, New Guinea, NE Australia, New Hebrides, Bismarck Archipelago, Solomon Islands, New Caledonia	Acanthaceae (<i>Asystasia</i> , <i>Blechum</i> , <i>Carlownrightia</i> , <i>Eranthemum</i> , <i>Graptophyllum</i> , <i>Phaulopsis</i> , <i>Pseuderanthemum</i> , <i>Ruellia</i> , <i>Strobilanthes</i>), Calycanthaceae (<i>Calycanthus</i>), Fabaceae (<i>Erythrina</i>), Moraceae (<i>Artocarpus</i>), Urticaceae (<i>Girardinia</i> , <i>Urtica</i>)	ROBINSON et al (2010), DAY (2015), HOSKINS (2015), INAYOSHI (2015), SVELA (2015)
91. <i>Kallima inachus</i> (Doyère, 1840)	India, Myanmar, Laos, Thailand, Vietnam, W Malaysia, China, Taiwan, Japan	Acanthaceae (<i>Acanthus</i> , <i>Dicliptera</i> , <i>Hygrophila</i> , <i>Lepidagathis</i> , <i>Rostellularia</i> , <i>Strobilanthes</i>), Polygonaceae (<i>Polygonum</i>), Rosaceae (<i>Prunus</i>), Urticaceae (<i>Girardinia</i>)	YANG et al (2005), ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), SVELA (2015)
92. <i>Kallima paralekta</i> (Horsfield, 1829)	Java, Sumatra	Acanthaceae (<i>Pseuderanthemum</i> , <i>Strobilanthes</i>)	ROBINSON et al (2010), SVELA (2015)
Nymphalini			
93. <i>Colobura dirce</i> (Linnaeus, 1758)	Mexico, El Salvador, Guatemala, Costa Rica, Colombia, Venezuela, Guyana, French Guiana, Suriname, Ecuador, Peru, Bolivia, Brazil, Argentina, Paraguay, Puerto Rico, Cuba, Jamaica, Hispaniola, Trinidad	Cecropiaceae (<i>Cecropia</i>), Fabaceae (<i>Cassia</i>), Meliaceae (<i>Cabralea</i>), Musaceae (<i>Musa</i>), Myrtaceae (<i>Eucalyptus</i>), Rubiaceae (<i>Coffea</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Victorini			
94. <i>Anartia amathea</i> (Linnaeus, 1758)	Belize, Panama, Colombia, Venezuela, N Chile, Suriname, Bolivia, Brazil, Argentina, Paraguay, Uruguay, Lesser Antilles (Grenada, Barbados, Antigua, Trinidad, Tobago)	Acanthaceae (<i>Blechnum</i> , <i>Dicliptera</i> , <i>Justicia</i> , <i>Ruellia</i>), Lamiaceae (<i>Melissa</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SVELA (2015)
95. <i>Siproeta epaphus</i> (Latreille, 1813)	S USA (New Mexico, Texas), Mexico, Guatemala, Costa Rica, Venezuela, Ecuador, Peru, Brazil, Argentina	Acanthaceae (<i>Blechnum</i> , <i>Ruellia</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015)
96. <i>Siproeta stelenes</i> (Linnaeus, 1758)	USA (Florida, Texas), Mexico, Honduras, Belize, Costa Rica, Panama, Ecuador, Peru, Brazil, Argentina, Bolivia, Antilles (Cuba, Jamaica, Hispaniola, Puerto Rico, St. Kitts, St. Croix), Cayman Islands	Acanthaceae (<i>Blechnum</i> , <i>Justicia</i> , <i>Pseuderanthemum</i> , <i>Ruellia</i> , <i>Strobilanthes</i>), Fabaceae (<i>Calliandra</i> , <i>Pithecellobium</i>), Lamiaceae (<i>Salvia</i>), Plantaginaceae (<i>Plantago</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SVELA (2015)
Satyrinae			
Amathusiini			
107 97. <i>Amathusia phidippus</i> (Linnaeus, 1763)	India, Myanmar, Peninsular Malaysia, Thailand, Cambodia, Vietnam, Indonesia (Sulawesi, Java, Sumatra, Borneo, Bali, Lombok), Philippines	Arecaceae (<i>Areca</i> , <i>Borassus</i> , <i>Cocos</i> , <i>Corypha</i> , <i>Cyrtostachys</i> , <i>Elaeis</i> , <i>Metroxylon</i> , <i>Nypa</i>), Musaceae (<i>Musa</i>)	ROBINSON et al (2010), INAYOSHI (2015), SVELA (2015)
Brassolini			
98. <i>Brassolis isthmia</i> Bates, 1864	Guatemala, Panama, Colombia, Ecuador, Bolivia	Arecaceae (<i>Cocos</i> , <i>Chamaedorea</i>)	YOUNG (1985), WARREN et al (2013), SVELA (2015)
99. <i>Caligo atreus</i> Kollar, 1850	Mexico, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru	Arecaceae (<i>Asterogyne</i> , <i>Geonoma</i> , <i>Reinhardtia</i>), Bromeliaceae (<i>Guzmania</i>), Costaceae (<i>Costus</i>), Cyclanthaceae, Heliconiaceae (<i>Heliconia</i>), Marantaceae (<i>Calathea</i>), Musaceae (<i>Musa</i>), Zingiberaceae (<i>Renealmia</i>)	PENZ et al (2000), WARREN et al (2013), SVELA (2015)
100. <i>Caligo eurilochus</i> (Cramer, 1775)	Mexico, Panama, Colombia, Venezuela, Suriname, Peru, Bolivia, Brazil	Arecaceae (<i>Euterpe</i>), Heliconiaceae (<i>Heliconia</i>), Marantaceae (<i>Calathea</i>), Musaceae (<i>Musa</i>), Zingiberaceae (<i>Hedychium</i>)	PENZ et al (2000), WARREN et al (2013), SVELA (2015)
101. <i>Caligo idomeneus</i> (Linnaeus, 1758)	Panama, Colombia, Suriname, Brazil, Peru, Bolivia	Marantaceae, Monimiaceae (<i>Palmeria</i>), Musaceae (<i>Musa</i>)	PENZ et al (2000), ROBINSON et al (2010), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
102. <i>Caligo illioneus</i> (Cramer, 1775)	Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Suriname, Bolivia, Peru, Brazil, Paraguay, Trinidad, Saint Vincent	Cyperaceae (<i>Scleria</i>), Heliconiaceae (<i>Heliconia</i>), Musaceae (<i>Musa</i>), Poaceae (<i>Saccharum</i>), Zingiberaceae (<i>Hedychium</i>)	PENZ et al (2000), WARREN et al (2013), SAVELA (2015)
103. <i>Caligo telamonius</i> (C. Felder & R. Felder, 1862)	Mexico, Guatemala, Belize, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Surinam	Cannaceae (<i>Canna</i>), Heliconiaceae (<i>Heliconia</i>), Marantaceae (<i>Calathea</i>), Musaceae (<i>Musa</i>), Rubiaceae (<i>Coffea</i>)?, Zingiberaceae (<i>Alpinia</i>)	PENZ et al (2000), WARREN et al (2013), SAVELA (2015)
104. <i>Eryphanis automedon</i> (Cramer, 1775)	Guatemala, Nicaragua, Costa Rica, Colombia, Suriname, Ecuador, Bolivia, Brazil, Peru	Poaceae (<i>Bambusa</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
105. <i>Opsiphanes tamarindi</i> C. & R. Felder, 1861	Mexico, Honduras, Costa Rica, Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia	Araceae (<i>Arum</i>), Arecaceae (<i>Chamaedorea</i> , <i>Cocos</i>), Cannaceae (<i>Canna</i>), Heliconiaceae (<i>Heliconia</i>), Marantaceae (<i>Calathea</i>), Musaceae (<i>Musa</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
Elymniini			
106. <i>Elymnias hypermnestra</i> (Linnaeus, 1763)	India, Sri Lanka, S China, Taiwan, Myanmar, Laos, S Vietnam, Thailand, Cambodia, W Malaysia, Philippines, Timor, Taiwan, Sumatra, Borneo, Java, Bali	Arecaceae (<i>Archontophoenix</i> , <i>Areca</i> , <i>Arenga</i> , <i>Calamus</i> , <i>Chrysalidocarpus</i> , <i>Cocos</i> , <i>Cyrtostachys</i> , <i>Elaeis</i> , <i>Licuala</i> , <i>Livistona</i> , <i>Metroxylon</i> , <i>Phoenix</i> , <i>Ptychosperma</i> , <i>Rhapis</i> , <i>Roystonea</i> , <i>Trachycarpus</i>), Fabaceae (<i>Butea</i>), Poaceae, Verbenaceae (<i>Lantana</i>)	ROBINSON et al (2010), HOSKINS (2015), SAVELA (2015)
Melanitini			
107. <i>Melanitis leda</i> (Linnaeus, 1758)	Senegal, Gambia, Guinea-Bissau, Guinea, Mali, Burkina Faso, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Kenya, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Saudi Arabia, Yemen, Oman, Cape Verde Islands, Sao Tome and Principe, Equatorial Guinea, Mauritius, Rodrigues, Reunion, Seychelles, Mafia Island, Comoro Islands, Madagascar, Sri Lanka, India, Thailand, Laos, Cambodia, Vietnam, China, Malay Peninsula, Sumatra, Java, Borneo, Sulawesi, Bali, Lesser Sunda Islands, Philippines, New Guinea, N Australia	Arecaceae (<i>Cocos</i> , <i>Elaeis</i>), Cyperaceae (<i>Cladium</i>), Poaceae (<i>Andropogon</i> , <i>Apluda</i> , <i>Axonopus</i> , <i>Bambusa</i> , <i>Brachiaria</i> , <i>Capillipedium</i> , <i>Chrysopogon</i> , <i>Coix</i> , <i>Cynodon</i> , <i>Cyrtococcum</i> , <i>Digitaria</i> , <i>Eleusine</i> , <i>Ehrharta</i> , <i>Heteropogon</i> , <i>Imperata</i> , <i>Leersia</i> , <i>Melinis</i> , <i>Microstegium</i> , <i>Miscanthus</i> , <i>Ophiuros</i> , <i>Oplismenus</i> , <i>Oryza</i> , <i>Panicum</i> , <i>Paspalum</i> , <i>Pennisetum</i> , <i>Poa</i> , <i>Prospyrtochloa</i> , <i>Pseudechinolaena</i> , <i>Rottboellia</i> , <i>Saccharum</i> , <i>Setaria</i> , <i>Sorghum</i> , <i>Sporobolus</i> , <i>Stenotaphrum</i> , <i>Themeda</i> , <i>Thysanolaena</i> , <i>Zea</i> , <i>Zizania</i>)	WILLIAMS (2008), ROBINSON et al (2010), INAYOSHI (2015), HOSKINS (2015), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Morphini			
108. <i>Morpho achilles</i> (Linnaeus, 1758)	Colombia, Venezuela, Guyana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil, Paraguay, Uruguay	Fabaceae (<i>Dalbergia</i> , <i>Inga</i> , <i>Machaerium</i> , <i>Myrocarpus</i> , <i>Platymiscium</i> , <i>Pterocarpus</i>)	ROBINSON et al (2010), HOSKINS (2015), SAVELA (2015)
109. <i>Morpho helenor</i> (Cramer, 1776)	Mexico, Honduras, Guatemala, Belize, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Guyana, Suriname, French Guiana, Venezuela, Ecuador, Peru, Brazil, Bolivia, Argentina, Paraguay, Trinidad, Tobago	Bignoniaceae (<i>Paragonia</i>), Fabaceae (<i>Arachis</i> , <i>Dalbergia</i> , <i>Dioclea</i> , <i>Inga</i> , <i>Lonchocarpus</i> , <i>Machaerium</i> , <i>Medicago</i> , <i>Mucuna</i> , <i>Pithecellobium</i> , <i>Platymiscium</i> , <i>Pterocarpus</i> , <i>Swartzia</i>)	ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
110. <i>Morpho polyphemus</i> Westwood, 1850	Mexico, Honduras, Guatemala, El Salvador, Nicaragua, Costa Rica	Fabaceae (<i>Inga</i> , <i>Pithecellobium</i>), Sapindaceae (<i>Paullinia</i>)	WARREN et al (2013), SAVELA (2015)
Satyriini			
111. <i>Mycalesis intermedia</i> Moore, 1892	Myanmar, Thailand, Laos, Cambodia, Vietnam, Peninsular Malaysia, S China, India	Poaceae	ROBINSON et al (2010), INAYOSHI (2015), SAVELA (2015)
Papilionidae			
Papilioninae			
Leptocircini			
112. <i>Graphium agamemnon</i> (Linnaeus, 1758)	China, India, Sri Lanka, China, Malaysia, Myanmar, Thailand, Laos, Cambodia, Vietnam, Sumatra, Java, Borneo, Sulawesi, Bali, Philippines, Moluccas, New Guinea, Solomon Islands, Australia	Annonaceae (<i>Anaxagorea</i> , <i>Ancana</i> , <i>Annona</i> , <i>Artabotrys</i> , <i>Cyathostemma</i> , <i>Desmos</i> , <i>Fitzalania</i> , <i>Friesodielsia</i> , <i>Goniothalamus</i> , <i>Gutteria</i> , <i>Haplostichanthus</i> , <i>Melodorum</i> , <i>Miliusa</i> , <i>Mitrephora</i> , <i>Oncodostigma</i> , <i>Polyalthia</i> , <i>Pseuduvaria</i> , <i>Rollinia</i> , <i>Uvaria</i> , <i>Xylophia</i>), Bombacaceae (<i>Durio</i>), Dioscoreaceae (<i>Dioscorea</i>), Fabaceae (<i>Cassia</i>), Lauraceae (<i>Cinnamomum</i>), Magnoliaceae (<i>Magnolia</i> , <i>Michelia</i>), Piperaceae (<i>Piper</i>), Rutaceae (<i>Citrus</i>)	DUNN & DUNN (1991), VANE-WRIGHT & DE JONG (2003), ROBINSON et al (2010) HOSKINS (2015), INAYOSHI (2015), SAVELA (2015)
113. <i>Graphium angolanus</i> (Goeze, 1779)	Senegal, Gambia, Guinea-Bissau, Guinea, Sierra Leone, Ivory Coast, Mali, Burkina Faso, Ghana, Togo, Benin, Niger, Nigeria, Cameroon, Equatorial Guinea, Sao Tome and Principe,	Annonaceae (<i>Annona</i> , <i>Artabotrys</i> , <i>Monodora</i> , <i>Uvaria</i>), Apocynaceae (<i>Landolphia</i>), Malpighiaceae (<i>Sphedamnocarpus</i>)	WILLIAMS (2008), ROBINSON et al (2010)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
	Gabon, Congo, Democratic Republic of Congo, Central African Republic, Chad, Sudan, Uganda, Rwanda, Burundi, Ethiopia, Kenya, Tanzania, Angola, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Comoro Islands		
114. <i>Graphium antheus</i> (Cramer, 1779)	Senegal, Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Gabon, Congo, Angola, Central African Republic, Democratic Republic of Congo, Sudan, Uganda, Rwanda, Burundi, Ethiopia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland	Annonaceae (<i>Annona</i> , <i>Artabotrys</i> , <i>Cleistochlamys</i> , <i>Hexalobus</i> , <i>Monanthes</i> , <i>Monodora</i> , <i>Uvaria</i>), Apocynaceae (<i>Landolphia</i>)	WILLIAMS (2008)
115. <i>Graphium colonna</i> (Ward, 1873)	Liberia, Sierra Leone, Ghana, Ivory Coast, Togo, Equatorial Guinea, Gabon, Congo, Democratic Republic of Congo, Central African Republic, Sudan, Ethiopia, Somalia, Uganda, Kenya, Tanzania, Malawi, Mozambique, Zimbabwe, South Africa, Swaziland, Madagascar	Annonaceae (<i>Annona</i> , <i>Artabotrys</i> , <i>Monodora</i> , <i>Uvaria</i>)	WILLIAMS (2008), ROBINSON et al (2010)
116. <i>Graphium doson</i> C&R Felder, 1864	India, Sri Lanka, Myanmar, Laos, Cambodia, Thailand, China, Taiwan, Korea, Vietman, Japan, Philippines, W Malaysia, Sumatra, Borneo, Sulawesi, Java	Annonaceae (<i>Annona</i> , <i>Desmos</i> , <i>Miliusa</i> , <i>Polyalthia</i> , <i>Uvaria</i>), Apocynaceae (<i>Hunteria</i>), Callophylacaceae (<i>Mesua</i>), Lauraceae (<i>Cinnamomum</i> , <i>Litsea</i>), Magnoliaceae (<i>Magnolia</i> , <i>Michelia</i>)	ROBINSON et al (2010), HOSKINS (2015), SVELA (2015)
117. <i>Graphium polistratus</i> (Grose-Smith, 1889)	Nigeria, Democratic Republic of Congo, Kenya, Tanzania, Malawi, Mozambique	Annonaceae (<i>Annona</i> , <i>Artabotrys</i> , <i>Uvaria</i>)	WILLIAMS (2008), ROBINSON et al (2010),
118. <i>Mimoides ilus</i> (Fabricius, 1793)	Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela	Annonaceae (<i>Annona</i> , <i>Rollinia</i>)	WARREN et al (2013), SVELA (2015)
119. <i>Mimoides thymbraeus</i> (Boisduval, 1836)	Mexico, Honduras, El Salvador, Guatemala	Annonaceae (<i>Annona</i>)	WARREN et al (2013), MURILLO-HILLER (2013), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
Papilionini			
120. <i>Papilio anchisiades</i> Esper, 1788	USA (Texas), Mexico, Honduras, Guatemala, El Salvador, Belize, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Guayana, Suriname, French Guiana, Ecuador, Peru, Bolivia, Brazil, Argentina, Paraguay, Trinidad, Tobago	Rutaceae (<i>Amyris</i> , <i>Casimiroa</i> , <i>Choisya</i> , <i>Citrus</i> , <i>Esenbeckia</i> , <i>Fortunella</i> , <i>Ruta</i> , <i>Skimmia</i> , <i>Swinglea</i> , <i>Zanthoxylum</i>)	LEWIS (2010), ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
121. <i>Papilio androgeus</i> Cramer, 1775	USA (Florida), Mexico, Guatemala, El Salvador, Belize, Costa Rica, Panama, Suriname, Colombia, Ecuador, Bolivia, Peru, Brazil, Argentina, Paraguay, Hispaniola, Cuba, Puerto Rico, Santa Lucia, Trinidad, Tobago	Rutaceae (<i>Amyris</i> , <i>Casimiroa</i> , <i>Choisya</i> , <i>Citrus</i> , <i>Esenbeckia</i> , <i>Swinglea</i> , <i>Zanthoxylum</i>)	LEWIS (2010), ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
122. <i>Papilio astyalus</i> Godart, 1819	USA (Texas, Arizona), Mexico, Guatemala, El Salvador, Costa Rica, Panama, Colombia, Venezuela, French Guiana, Ecuador, Peru, Bolivia, Brazil, Argentina, Paraguay, Cuba, Trinidad	Rutaceae (<i>Amyris</i> , <i>Balfourodendron</i> , <i>Casimiroa</i> , <i>Choisya</i> , <i>Citrus</i> , <i>Esenbeckia</i> , <i>Galipea</i> , <i>Pilocarpus</i> , <i>Ruta</i> , <i>Skimmia</i> , <i>Zanthoxylum</i>)	LEWIS (2010), ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
123. <i>Papilio clytia</i> Linnaeus, 1758	India, Sri Lanka, Andaman Islands, China, Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Sumatra, Borneo, Sulawesi, Philippines, Timor	Lauraceae (<i>Alseodaphne</i> , <i>Cinnamomum</i> , <i>Cryptocarya</i> , <i>Litsea</i> , <i>Persea</i> , <i>Phoebe</i>), Sapotaceae (<i>Sarcosperma</i>)	ROBINSON et al (2010), HOSKINS (2015), SVELA (2015)
124. <i>Papilio constantinus</i> Ward, 1871	Congo, Democratic Republic of Congo, Ethiopia, Somalia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, South Africa, Swaziland	Rutaceae (<i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Vepris</i> , <i>Zanthoxylum</i>)	WILLIAMS (2008), ROBINSON et al (2010)
125. <i>Papilio crespontes</i> Cramer, 1777	Canada, USA, Mexico, Honduras, Guatemala, El Salvador, Nicaragua, Costa Rica, Panama, Colombia; introduced to Cuba	Cannabaceae (<i>Humulus</i>), Cornaceae (<i>Nyssa</i>), Lauraceae (<i>Persea</i>), Piperaceae (<i>Piper</i>), Rutaceae (<i>Amyris</i> , <i>Casimiroa</i> , <i>Choisya</i> , <i>Citrus</i> , <i>Dictamnus</i> , <i>Esenbeckia</i> , <i>Helietta</i> , <i>Murraya</i> , <i>Phellodendron</i> , <i>Ptelea</i> , <i>Ruta</i> , <i>Zanthoxylum</i>), Salicaceae (<i>Populus</i>), Staphyleaceae (<i>Staphylea</i>)	LEWIS (2010), ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
126. <i>Papilio dardanus</i> Brown, 1776	Senegal, Guinea-Bissau, Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Burkina Faso, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea,	Monimiaceae (<i>Xymalos</i>), Rutaceae (<i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Orcia</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Vepris</i> , <i>Zanthoxylum</i>)	WILLIAMS (2008), ROBINSON et al (2010)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
	Sao Tome and Principe, Gabon, Congo, Angola, Central African Republic, Democratic Republic of Congo, Sudan, Uganda, Kenya, Ethiopia, Eritrea, Somalia, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, South Africa, Swaziland, Comoro Islands, Madagascar		
127. <i>Papilio deiphobus</i> Linnaeus, 1758	Sumatra, Sulawesi, New Guinea, Philippines, Moluccas	Rutaceae (<i>Atalantia</i> , <i>Citrus</i>)	ROBINSON et al (2010), SVELA (2015)
128. <i>Papilio demodocus</i> Esper, 1798	Mauritania, Senegal, Gambia, Guinea-Bissau, Guinea, Sierra Leone, Liberia, Ivory Coast, Mali, Burkina Faso, Ghana, Togo, Benin, Niger, Nigeria, Cameroon, Uganda, Kenya, Tanzania, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Lesotho, Saudi Arabia, Yemen, Cape Verde Islands, Comoro Islands, Madagascar, Mauritius, Reunion, Socotra	Anacardiaceae (<i>Pseudospondias</i>), Apiaceae (<i>Angelica</i> , <i>Athamanta</i> , <i>Deverra</i> , <i>Foeniculum</i> , <i>Notobubon</i> , <i>Peucedanum</i> , <i>Pituranthos</i>), Euphorbiaceae (<i>Euphorbia</i>), Fabaceae (<i>Acacia</i> , <i>Psoralea</i>), Rhamnaceae (<i>Ziziphus</i>), Rutaceae (<i>Aegle</i> , <i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Fagaropsis</i> , <i>Glycosmis</i> , <i>Haplophyllum</i> , <i>Murraya</i> , <i>Oricia</i> , <i>Phellodendron</i> , <i>Poncirus</i> , <i>Ptelea</i> , <i>Ruta</i> , <i>Skimmia</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Vepris</i> , <i>Zanthoxylum</i>), Ptaeroxylaceae (<i>Ptaeroxylon</i>), Sapindaceae (<i>Hippobromus</i>)	WILLIAMS (2008), ROBINSON et al (2010), SVELA (2015)
129. <i>Papilio demoleus</i> Linnaeus, 1758	Saudi Arabia, Iran, India, Sri Lanka, Afghanistan, Nepal, Myanmar, Thailand, Laos, Cambodia, Vietnam, S. China, Taiwan, Japan, Philippines, Malaysia, Sumatra, Java, Borneo, Lesser Sunda Islands, Timor, New Guinea, Australia, introduced to Central America (Dominican Republic) and the Caribbean (Puerto Rico, Jamaica)	Fabaceae (<i>Cullen</i> , <i>Psoralea</i>), Magnoliaceae (<i>Michelia</i>), Loganiaceae (<i>Fagraea</i>), Rhamnaceae (<i>Ziziphus</i>), Rutaceae (<i>Acronychia</i> , <i>Aegle</i> , <i>Atalantia</i> , <i>Chloroxylon</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Flindersia</i> , <i>Fortunella</i> , <i>Glycosmis</i> , <i>Limonia</i> , <i>Microcitrus</i> , <i>Micromelum</i> , <i>Murraya</i> , <i>Ruta</i> , <i>Toddalia</i> , <i>Triphasia</i> , <i>Zanthoxylum</i>), Tiliaceae (<i>Tilia</i>)	VANE-WRIGHT & DE JONG (2003), LEWIS (2009), ROBINSON et al (2010), HOSKINS (2015), SVELA (2015)
130. <i>Papilio echerioides</i> Trimen, 1868	Cameroon, Equatorial Guinea, Angola, Democratic Republic of Congo, Sudan, Ethiopia, Uganda, Rwanda, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, South Africa, Swaziland	Rutaceae (<i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Clausena</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Vepris</i> , <i>Zanthoxylum</i>)	WILLIAMS (2008), ROBINSON et al (2010)
131. <i>Papilio erostratus</i> Westwood, 1847	Mexico, Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica	Rutaceae (<i>Casimiroa</i> , <i>Citrus</i>)	MAES (2006), LEWIS (2010), WARREN et al (2013), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
132. <i>Papilio garamas</i> (Geyer, 1829)	Mexico, Guatemala, Honduras, El Salvador, Costa Rica, Panama	Lauraceae (<i>Persea</i> , <i>Sassafras</i>), Magnoliaceae (<i>Liriodendron</i> , <i>Magnolia</i>)	SCRIBER et al (1990), WARREN et al (2013), SVELA (2015)
133. <i>Papilio helenus</i> Linnaeus, 1758	India, Sri Lanka, Bhutan, Myanmar, Thailand, Cambodia, Laos, Vietnam, China, Taiwan, Japan, Philippines, Malaysia, Sumatra, Borneo, Sulawesi, Java	Rutaceae (<i>Citrus</i> , <i>Euodia</i> , <i>Fortunella</i> , <i>Phellodendron</i> , <i>Poncirus</i> , <i>Toddalia</i> , <i>Zanthoxylum</i>)	ROBINSON et al (2010), HOSKINS (2015), SVELA (2015)
134. <i>Papilio hipponous</i> C & R. Felder, 1862	Philippines, Sulawesi	Rutaceae (<i>Citrus</i>)	PAGE et al (2003), VANE-WRIGHT & DE JONG (2003), SVELA (2015)
135. <i>Papilio lowi</i> Druce, 1873	Philippines, Borneo	Rutaceae (<i>Citrus</i>)	REIMAN GARDENS (2015), SVELA (2015)
136. <i>Papilio maackii</i> Ménétries, 1859	SE Russia, Sakhalin, Kurile Islands, E China, Japan (Hokkaido), North Korea	Rutaceae (<i>Euodia</i> , <i>Orixa</i> , <i>Phellodendron</i> , <i>Zanthoxylum</i>)	ROBINSON et al (2010), SVELA (2015)
137. <i>Papilio memnon</i> Linnaeus, 1758	India, Nepal, Myanmar, Thailand, Laos, Cambodia, Vietnam, Peninsular Malaysia, China, Taiwan, Japan, Borneo, Sumatra, Java, Bali, Philippines	Aristolochiaceae (<i>Aristolochia</i>), Magnoliaceae (<i>Magnolia</i> , <i>Michelia</i>), Rutaceae (<i>Atalantia</i> , <i>Citrus</i> , <i>Fortunella</i> , <i>Paramignya</i> , <i>Poncirus</i> , <i>Severinia</i> , <i>Toddalia</i> , <i>Zanthoxylum</i>)	ROBINSON et al (2010), HOSKINS (2015), INAYOSHI (2015), (SVELA (2015)
138. <i>Papilio nephelus</i> Boisduval, 1836	India, Nepal, Bhutan, Myanmar, Thailand, Laos, Cambodia, Vietnam, China (Yunnan), Taiwan, Malaysia, Sumatra, Borneo, Sulawesi, Java	Rutaceae (<i>Citrus</i> , <i>Clausena</i> , <i>Euodia</i> , <i>Toddalia</i> , <i>Zanthoxylum</i>)	ROBINSON et al (2010), HOSKINS (2015), SVELA (2015)
139. <i>Papilio nireus</i> Linnaeus, 1758	Senegal, Gambia, Guinea-Bissau, Guinea, Burkina Faso, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Gabon, Congo, Central African Republic, Angola, Democratic Republic of Congo, Sudan, Uganda, Ethiopia, Eritrea, Somalia, Kenya, Tanzania, Malawi, Zambia, Mozambique, Zimbabwe, Botswana, Namibia, South Africa, Swaziland, Comoro Islands	Rutaceae (<i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Orixa</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Vepris</i> , <i>Zanthoxylum</i>)	WILLIAMS (2008), ROBINSON et al (2010), SVELA (2015)
140. <i>Papilio ophidicephalus</i> Oberthür, 1878	Kenya, Tanzania, Zambia, Democratic Republic of Congo, Malawi, Mozambique, Zimbabwe, South Africa, Swaziland	Rutaceae (<i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Zanthoxylum</i>)	WILLIAMS (2008), ROBINSON et al (2010)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
141. <i>Papilio palinurus</i> Fabricius, 1787	Thailand, Myanmar, Peninsular Malaysia, Indonesia (Sumatra, Borneo), Philippines	Rutaceae (<i>Clausena</i> , <i>Euodia</i> , <i>Micromelum</i>)	ROBINSON et al (2010), SVELA (2015)
142. <i>Papilio peranthus</i> Fabricius, 1787	Java, Sulawesi, Lesser Sunda Islands	Rutaceae (<i>Micromelum</i>)	ROBINSON et al (2010), SVELA (2015)
143. <i>Papilio phorcas</i> Cramer, 1775	Guinea, Sierra Leone, Liberia, Ivory Coast, Ghana, Togo, Benin, Nigeria, Cameroon, Equatorial Guinea, Bioko, Gabon, Angola, Congo, Democratic Republic of Congo, Central African Republic, Sudan, Uganda, Kenya, Tanzania, Malawi, Zambia	Rutaceae (<i>Calodendrum</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Macrostylis</i> , <i>Oriciopsis</i> , <i>Teclea</i> , <i>Toddalia</i> , <i>Vepris</i> , <i>Zanthoxylum</i>)	WILLIAMS (2008), ROBINSON et al (2010)
144. <i>Papilio pilumnus</i> Boisduval, 1836	USA (Texas), Mexico, Guatemala,	Lauraceae (<i>Litsea</i>)	ROBINSON et al (2010), SVELA (2015)
145. <i>Papilio polytes</i> Linnaeus, 1758	India, Sri Lanka, Himalayas, China, Taiwan, Japan, Myanmar, Thailand, Cambodia, Laos, Vietnam, W Malaysia, Singapore, Andaman and Nicobar Islands, Sumatra, Borneo, Java, Sulawesi, Bali, Timor, Philippines	Rutaceae (<i>Aegle</i> , <i>Atalantia</i> , <i>Citrus</i> , <i>Clausena</i> , <i>Correa</i> , <i>Euodia</i> , <i>Fortunella</i> , <i>Glycosmis</i> , <i>Micromelum</i> , <i>Murraya</i> , <i>Paramignya</i> , <i>Poncirus</i> , <i>Severinia</i> , <i>Toddalia</i> , <i>Triphasia</i> , <i>Zanthoxylum</i>)	ROBINSON et al (2010), INAYOSHI (2015), SVELA (2015)
146. <i>Papilio polyxenes</i> Fabricius, 1775	Canada, USA, Mexico, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Peru, Ecuador, Cuba	Apiaceae (<i>Aegopodium</i> , <i>Anethum</i> , <i>Angelica</i> , <i>Apium</i> , <i>Berula</i> , <i>Carum</i> , <i>Ciclospermum</i> , <i>Cicuta</i> , <i>Conium</i> , <i>Cryptotaenia</i> , <i>Cymopterus</i> , <i>Daucus</i> , <i>Eryngium</i> , <i>Foeniculum</i> , <i>Harbouria</i> , <i>Heracleum</i> , <i>Levisticum</i> , <i>Ligusticum</i> , <i>Lomatium</i> , <i>Musineon</i> , <i>Osmorhiza</i> , <i>Oxypolis</i> , <i>Pastinaca</i> , <i>Petroselinum</i> , <i>Ptilimnium</i> , <i>Sanicula</i> , <i>Sium</i> , <i>Spananthe</i> , <i>Spermolepis</i> , <i>Taenidia</i> , <i>Tauschia</i> , <i>Thaspium</i> , <i>Torilis</i> , <i>Zizia</i>), Asteraceae (<i>Cosmos</i> , <i>Solidago</i>), Brassicaceae (<i>Brassica</i>), Magnoliaceae (<i>Magnolia</i>), Rutaceae (<i>Dictamnus</i> , <i>Ruta</i> , <i>Thamnosma</i> , <i>Zanthoxylum</i>)	ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
147. <i>Papilio thoas</i> Linnaeus, 1771	USA (Texas), Mexico, Honduras, Guatemala, El Salvador, Belize, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Guyana, French Guiana, Suriname, Brazil, Ecuador, Peru, Bolivia, Argentina, Paraguay, Uruguay, Cuba, Trinidad, Tobago	Araceae (<i>Syngonium</i>), Rutaceae (<i>Citrus</i> , <i>Ertela?</i> , <i>Esenbeckia</i> , <i>Fortunella</i> , <i>Monnieria</i> , <i>Ptelea</i> , <i>Ruta</i> , <i>Zanthoxylum</i>), Piperaceae (<i>Piper</i>)	LEWIS (2010), ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
148. <i>Papilio torquatus</i> Cramer, 1777	Mexico, El Salvador, Guatemala, Costa Rica, Panama, Colombia, Venezuela, Suriname, French Guiana, Brazil, Ecuador, Peru, Bolivia, Argentina, Paraguay, Trinidad	Rutaceae (<i>Angostura</i> , <i>Choisya</i> , <i>Citrus</i> , <i>Ertela</i> , <i>Erythrochiton</i> , <i>Esenbeckia</i> , <i>Euxylophora</i> , <i>Galipea</i> , <i>Zanthoxylum</i>)	LEWIS (2010), ROBINSON et al (2010), WARREN et al (2013), SVELA (2015)
Troidini			
149. <i>Battus belus</i> (Cramer, 1777)	Mexico, Guatemala, Venezuela, Colombia, Peru, Bolivia, Brazil, Guyana, Surinam	Aristolochiaceae (<i>Aristolochia</i>)	ROBINSON et al (2010), SVELA (2015)
150. <i>Battus polydamas</i> Linnaeus, 1758	SE USA, Mexico, Honduras, Costa Rica, Panama, Colombia, Venezuela, Guayna, Suriname, Ecuador, Bolivia, Chile, Argentina, Peru, Brazil, Caribbean islands (Cuba, Hispaniola, Jamaica, Puerto Rico, Virgin Islands, St. Kitts, Montserrat, Guadeloupe, Trinidad, Tobago, Dominica, Martinique, St. Lucia, St. Vincent, Grenada, Bahamas and others)	Aristolochiaceae (<i>Aristolochia</i>), Rutaceae (<i>Citrus</i>)	ROTHSCHILD & JORDAN (1906), ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SVELA (2015)
115 151. <i>Atrophaneura antiphus</i> (Fabricius, 1793)	Sumatra, Borneo, Java, Philippines	Aristolochiaceae (<i>Aristolochia</i>)	ROTHSCHILD (1895), PAGE et al (2003), SVELA (2015)
152. <i>Atrophaneura kotzebuea</i> (Eschscholtz, 1821)	Philippines	Aristolochiaceae (<i>Aristolochia</i>)	PAGE et al (2003), SVELA (2015)
153. <i>Parides childrenae</i> Gray, 1832	Mexico, Guatemala, Nicaragua, Costa Rica, Panama, Colombia, Ecuador	Aristolochiaceae (<i>Aristolochia</i>)	WARREN et al (2013), SVELA (2015)
154. <i>Parides eurimedes</i> (Stoll, 1782)	Mexico, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Ecuador	Aristolochiaceae (<i>Aristolochia</i>)	ROBINSON et al (2010), SVELA (2015)
155. <i>Parides iphidamas</i> (Fabricius, 1793)	Mexico, Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Peru, Ecuador	Aristolochiaceae (<i>Aristolochia</i>)	MÖHN (2006), SVELA (2015)
156. <i>Parides lysander</i> (Cramer, 1775)	Suriname, French Guiana, Venezuela, Colombia, Ecuador, Peru, Brazil	Aristolochiaceae (<i>Aristolochia</i>), Rutaceae (<i>Citrus</i>)	ROBINSON et al (2010), SVELA (2015)
157. <i>Parides montezuma</i> (Westwood, 1842)	Mexico, Belize, El Salvador, Honduras, Nicaragua, Costa Rica, Panama	Aristolochiaceae (<i>Aristolochia</i>)	MÖHN (2006), WARREN et al (2013), SVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
158. <i>Parides photinus</i> (Doubleday, 1844)	Mexico, Honduras, Belize, Nicaragua, Costa Rica	Aristolochiaceae (<i>Aristolochia</i>)	MÖHN (2006), SAVELA (2015)
159. <i>Troides rhadamantus</i> (Lucas, 1835)	Philippines	Aristolochiaceae (<i>Aristolochia</i>)	ROBINSON et al (2010), SAVELA (2015)
Pieridae			
Coliadinae			
160. <i>Anteos clorinde</i> (Godart, 1824)	USA (Texas, Florida), Mexico, Honduras, Belize, Venezuela, Peru, Bolivia, Brazil, Argentina, Paraguay, Cuba, Hispaniola, Jamaica, Bahamas	Fabaceae (<i>Cassia</i> , <i>Pithecellobium</i> , <i>Senna</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
161. <i>Catopsilia pomona</i> Fabricius, 1775	India, Sri Lanka, Taiwan, Bangladesh, Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Philippines, Sumatra, Java, Borneo, Sulawesi, Lesser Sunda Islands, Timor, Moluccas, Solomon Islands, New Guinea, New Caledonia, Vanuatu, Australia, New Zealand	Fabaceae (<i>Bauhinia</i> , <i>Brownea</i> , <i>Butea</i> , <i>Cassia</i> , <i>Pterocarpus</i> , <i>Senna</i> , <i>Sesbania</i>)	VANE-WRIGHT & DE JONG (2003), ROBINSON et al (2010), INAYOSHI (2015), SAVELA (2015)
162. <i>Phoebis argante</i> (Fabricius, 1775)	Mexico, El Salvador, Honduras, Costa Rica, Panama, Guyana, Suriname, Peru, Brazil, Argentina, Uruguay, Trinidad, Puerto Rico, Cuba, Hispaniola, Jamaica, Hispaniola	Capparaceae, Fabaceae (<i>Caesalpinia</i> , <i>Cassia</i> , <i>Inga</i> , <i>Pentaclethra</i> , <i>Pithecellobium</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
163. <i>Phoebis philea</i> (Linnaeus, 1763)	S USA, Mexico, El Salvador, Costa Rica, Suriname, Peru, Brazil, Cuba, Hispaniola, Puerto Rico	Fabaceae (<i>Caesalpinia</i> , <i>Cassia</i> , <i>Pithecellobium</i> , <i>Senna</i>)	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)
164. <i>Phoebis sennae</i> (Linnaeus, 1758)	USA, Mexico, Honduras, Costa Rica, Panama, Suriname, French Guiana, Peru, Chile, Bolivia, Brazil, Chile, Argentina, Uruguay, Galapagos, Cuba, Hispaniola, St. Vincent, Grenada, Jamaica	Brassicaceae, Flacourtiaceae (<i>Casearia</i>), Fabaceae (<i>Cassia</i> , <i>Chamaecrista</i> , <i>Crotalaria</i> , <i>Phaseolus</i> , <i>Senna</i> , <i>Trifolium</i>)	PYLE (1981), ROBINSON et al (2010), WARREN et al (2013), HOSKINS (2015), SAVELA (2015)
Pierinae			
Pierini			
165. <i>Ascia monuste</i> (Linnaeus, 1764)	USA, Mexico, Nicaragua, Costa Rica, Suriname, Brazil, Argentina, Peru, Antilles (Cuba, Hispaniola, Jamaica, Puerto Rico, St. Vincent,	Amaryllidaceae (<i>Allium</i>), Bataceae (<i>Batis</i>), Brassicaceae (<i>Armoracia</i> , <i>Brassica</i> , <i>Cakile</i> , <i>Capsella</i> , <i>Lepidium</i> , <i>Lobularia</i> , <i>Raphanus</i> , <i>Rorippa</i> , <i>Sinapis</i>),	ROBINSON et al (2010), WARREN et al (2013), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
	Virgin Islands, St. Kitts, Antigua, St. Barthélemy, St. Martin), Bahamas	Capparaceae (<i>Capparis</i> , <i>Crateva</i>), Cleomaceae (<i>Cleome</i> , <i>Polanisia</i>), Fabaceae (<i>Cassia</i>), Tropaeolaceae (<i>Tropaeolum</i>)	
Colotini			
166. <i>Hebomoia glaucippe</i> (Linnaeus, 1758)	India, Sri Lanka, Nepal, China, Taiwan, Myanmar, Thailand, Cambodia, Laos, Vietnam, Malaysia, Singapore, Japan, Borneo, Sumatra, Java, Sulawesi, Philippines, Solomon Islands, Andaman Islands	Brassicaceae (<i>Brassica</i>), Capparaceae (<i>Capparis</i> , <i>Cleome</i> , <i>Crateva</i>)	ROBINSON et al (2010), INAYOSHI (2015), SABELA (2015)
167. <i>Pareronia valeria</i> (Cramer, 1776)	China (Hainan), India, Myanmar, Thailand, Peninsular Malaysia, Singapore, Sumatra, Java, Borneo, Philippines	Capparaceae (<i>Capparis</i>)	ROBINSON et al (2010), INAYOSHI (2015), SABELA (2015)
Saturniidae			
Saturniinae			
Attacini			
168. <i>Attacus atlas</i> (Linnaeus, 1758)	India, Sri Lanka, China, Thailand, Malaysia, Singapore, Java, Sumatra, Borneo, Sulawesi, Philippines, Timor, New Guinea	Anacardiaceae (<i>Anacardium</i> , <i>Lannea</i> , <i>Mangifera</i>), Annonaceae (<i>Annona</i> , <i>Artabotrys</i> , <i>Cananga</i>), Apocynaceae (<i>Alstonia</i> , <i>Nerium</i> , <i>Sarcostemma</i>), Aquifoliaceae (<i>Ilex</i>), Araliaceae (<i>Schefflera</i>), Asteraceae (<i>Artemisia</i>), Avicenniaceae (<i>Avicennia</i>), Berberidaceae (<i>Berberis</i>), Betulaceae (<i>Betula</i>), Bignoniaceae (<i>Spathodea</i>), Bombacaceae (<i>Ceiba</i>), Burseraceae (<i>Canarium</i>), Caprifoliaceae (<i>Lonicera</i>), Combretaceae (<i>Quisqualis</i> , <i>Terminalia</i>), Convolvulaceae (<i>Ipomoea</i>), Corylaceae (<i>Carpinus</i>), Crypteroniaceae (<i>Crypteronia</i>), Dilleniaceae (<i>Dillenia</i>), Dipterocarpaceae (<i>Shorea</i>), Ericaceae (<i>Kalmia</i> , <i>Rhododendron</i>), Euphorbiaceae (<i>Aleurites</i> , <i>Bischofia</i> , <i>Glochidion</i> , <i>Hevea</i> , <i>Phyllanthus</i> , <i>Ricinus</i> , <i>Sapium</i> , <i>Stillingia</i>), Fabaceae (<i>Albizia</i> , <i>Codariocalyx</i> , <i>Erythrina</i> , <i>Parkia</i> , <i>Sesbania</i>), Fagaceae, Goodeniaceae (<i>Scaevola</i>), Lamiaceae (<i>Leucosceptrum</i>), Lauraceae (<i>Cinnamomum</i> , <i>Persea</i> , <i>Sassafras</i>), Loganiaceae	ROBINSON et al (2010), HOSKINS (2015), SABELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
		(<i>Fagraea</i>), Lythraceae (<i>Lagerstroemia</i>), Magnoliaceae (<i>Michelia</i>), Malvaceae (<i>Hibiscus</i>), Melastomataceae (<i>Clidemia</i> , <i>Melastoma</i> , <i>Memecylon</i>), Meliaceae (<i>Sandoricum</i> , <i>Swietenia</i>), Moraceae (<i>Ficus</i> , <i>Morus</i>), Musaceae (<i>Musa</i>), Myrsinaceae (<i>Ardisia</i> , <i>Embelia</i> , <i>Maesa</i>), Myrtaceae (<i>Eucalyptus</i> , <i>Eugenia</i> , <i>Psidium</i>), Oleaceae (<i>Fraxinus</i> , <i>Jasminum</i> , <i>Ligustrum</i> , <i>Syringa</i>), Oxalidaceae (<i>Averrhoa</i>), Piperaceae (<i>Piper</i>), Poaceae (<i>Setaria</i>), Rhizophoraceae (<i>Bruguiera</i>), Rosaceae (<i>Malus</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Rosa</i>), Rubiaceae (<i>Cinchona</i> , <i>Coffea</i> , <i>Mussaenda</i> , <i>Nauclea</i> , <i>Uncaria</i> , <i>Vangueria</i>), Rutaceae (<i>Citrus</i>), Salicaceae (<i>Populus</i> , <i>Salix</i>), Sapindaceae (<i>Dimocarpus</i> , <i>Litchi</i> , <i>Nephelium</i> , <i>Schleichera</i>), Scrophulariaceae (<i>Paulownia</i>), Simaroubaceae (<i>Ailanthus</i> , <i>Brucea</i>), Sterculiaceae (<i>Theobroma</i>), Symplocaceae (<i>Symplocos</i>), Theaceae (<i>Camellia</i>), Tiliaceae (<i>Muntingia</i>), Verbenaceae (<i>Clerodendrum</i> , <i>Gmelina</i> , <i>Lantana</i> , <i>Stachytarpheta</i> , <i>Vitex</i>), Zingiberaceae (<i>Curcuma</i> , <i>Etlingera</i>)	
169. <i>Attacus caesar</i> Maassen, 1873	Philippines	Annonaceae (<i>Annona</i> , <i>Cananga</i>), Fabaceae (<i>Erythrina</i>), Lauraceae (<i>Persea</i>), Meliaceae (<i>Sandoricum</i> , <i>Swietenia</i>), Moraceae (<i>Artocarpus</i>), Myricaceae (<i>Myrica</i>), Oleaceae (<i>Syringa</i>), Rutaceae (<i>Citrus</i>), Salicaceae (<i>Salix</i>)	PAUKSTADT & PAUKSTADT (2002), ROBINSON et al (2010)
170. <i>Epiphora mythimnia</i> (Westwood, 1849)	Botswana, Kenya, Mozambique, South Africa, Tanzania, Zambia, Zimbabwe	Bignoniaceae (<i>Fernandoa</i>), Euphorbiaceae (<i>Croton</i>), Rhamnaceae (<i>Frangula</i> , <i>Helinus</i> , <i>Ziziphus</i>)	ROBINSON et al (2010), DE PRINS & DE PRINS (2015)
171. <i>Rothschildia lebeau</i> (Guérin-Méneville, 1868)	USA (Texas), Mexico, Costa Rica, Panama, Venezuela, Ecuador, Peru	Anacardiaceae (<i>Malosma</i> , <i>Schinus</i> , <i>Spondias</i>), Euphorbiaceae (<i>Ricinus</i>), Flacourtiaceae (<i>Casearia</i> , <i>Zuelania</i>), Fabaceae (<i>Acacia</i>), Lythraceae (<i>Lythrum</i>), Oleaceae (<i>Fraxinus</i> , <i>Ligustrum</i> , <i>Syringa</i>), Rosaceae (<i>Prunus</i>), Rubiaceae (<i>Cephalanthus</i> , <i>Chiococca</i> , <i>Exostema</i>), Rutaceae (<i>Citrus</i> , <i>Zanthoxylum</i>), Salicaceae (<i>Salix</i>), Simaroubaceae (<i>Ailanthus</i>)	ROBINSON et al (2010), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
172. <i>Samia cynthia</i> (Drury, 1773)	China, Korea; introduced to Japan, India, Australia, Canada, USA, Venezuela, Uruguay, Brazil, Tunisia, France, Austria, Switzerland, Germany, Spain, Bulgaria, Italy	Aceraceae (<i>Acer</i>), Adoxaceae (<i>Sambucus</i>), Apiaceae (<i>Apium</i>), Asteraceae (<i>Ageratina</i>), Anacardiaceae (<i>Rhus</i>), Apocynaceae (<i>Plumeria</i>), Aquifoliaceae (<i>Ilex</i>), Araliaceae (<i>Heteropanax</i>), Berberidaceae (<i>Berberis</i>), Burseraceae (<i>Canarium</i>), Caprifoliaceae (<i>Viburnum</i>), Caricaceae (<i>Carica</i>), Celastraceae (<i>Celastrus</i> , <i>Euonymus</i>), Coriariaceae (<i>Coriaria</i>), Cornaceae (<i>Camptotheca</i> , <i>Cornus</i>), Corylaceae (<i>Carpinus</i>), Euphorbiaceae (<i>Glochidion</i> , <i>Jatropha</i> , <i>Manihot</i> , <i>Ricinus</i>), Fabaceae (<i>Cassia</i> , <i>Laburnum</i>), Fagaceae (<i>Quercus</i>), Hamamelidaceae (<i>Liquidambar</i>), Juglandaceae (<i>Juglans</i>), Lauraceae (<i>Laurus</i> , <i>Lindera</i> , <i>Sassafras</i>), Lythraceae (<i>Lagerstroemia</i> , <i>Lawsonia</i>), Magnoliaceae (<i>Liriodendron</i> , <i>Magnolia</i> , <i>Michelia</i>), Malvaceae (<i>Alcea</i>), Meliaceae (<i>Azadirachta</i>), Myrtaceae (<i>Psidium</i>), Oleaceae (<i>Forsythia</i> , <i>Fraxinus</i> , <i>Ligustrum</i> , <i>Syringa</i>), Platanaceae (<i>Platanus</i>), Primulaceae (<i>Anagallis</i>), Rhamnaceae (<i>Ziziphus</i>), Rosaceae (<i>Malus</i> , <i>Prunus</i> , <i>Pyrus</i> , <i>Sorbus</i> , <i>Spiraea</i>), Rubiaceae (<i>Coffea</i>), Rutaceae (<i>Euodia</i> , <i>Ptelea</i> , <i>Zanthoxylum</i>), Salicaceae (<i>Salix</i>), Simaroubaceae (<i>Ailanthus</i>), Solanaceae (<i>Solanum</i>), Sterculiaceae (<i>Firmiana</i>), Tiliaceae (<i>Tilia</i>), Verbenaceae (<i>Gmelina</i>)	PEIGLER & NAUMANN (2003), ROBINSON et al (2010), SAVELA (2015)
173. <i>Samia ricini</i> (Donovan, 1798)	India, Indonesia	Euphorbiaceae (<i>Ricinus</i>), Lauraceae (<i>Cinnamomum</i>), Rubiaceae (<i>Cinchona</i>)	ROBINSON et al (2010)
Bunaeini			
174. <i>Bunaea alcinoe</i> (Stoll, 1780)	Angola, Benin, Cameroon, Chad, Democratic Republic of Congo, Egypt, Eritrea, Ethiopia, Gabon, Kenya, Lesotho, Madagascar, Malawi, Mozambique, South Africa, Sudan, São Tomé & Príncipe, Tanzania, Uganda, Zimbabwe	Anacardiaceae (<i>Harpephyllum</i> , <i>Mangifera</i> , <i>Rhus</i> , <i>Schinus</i>), Annonaceae (<i>Cananga</i>), Apocynaceae (<i>Diplorhynchus</i>), Araliaceae (<i>Cussonia</i> , <i>Panax</i>), Arecaceae (<i>Elaeis</i>), Balanitaceae (<i>Balanites</i>), Burseraceae (<i>Dacryodes</i>), Caprifoliaceae (<i>Lonicera</i>), Celastraceae (<i>Maytenus</i>), Combretaceae (<i>Terminalia</i>), Ebenaceae (<i>Diospyros</i>), Euphorbiaceae	ROBINSON et al (2010), DE PRINS & DE PRINS (2015), LATHAM (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
		(<i>Croton</i> , <i>Manihot</i> , <i>Sapium</i>), Fabaceae (<i>Acacia</i> , <i>Bauhinia</i> , <i>Erythrina</i>), Fagaceae (<i>Quercus</i>), Gentianaceae (<i>Anthocleista</i>), Lauraceae (<i>Persea</i>), Meliaceae (<i>Ekebergia</i> , <i>Khaya</i>), Myrsinaceae (<i>Maesa</i>), Myrtaceae (<i>Eucalyptus</i> , <i>Psidium</i>), Oleaceae (<i>Ligustrum</i>), Piperaceae (<i>Piper</i>), Rosaceae (<i>Crataegus</i> , <i>Malus</i> , <i>Prunus</i> , <i>Rosa</i>), Rubiaceae (<i>Coffea</i> , <i>Crossopteryx</i> , <i>Sarcocephalus</i>), Ulmaceae (<i>Celtis</i>), Zingiberaceae (<i>Aframomum</i>)	
175. <i>Gonimbrasia zambesina</i> (Walker, 1865)	Angola, Botswana, Kenya, Malawi, Mozambique, South Africa, Tanzania, Zambia, Zimbabwe	Anacardiaceae (<i>Mangifera</i> , <i>Ozoroa</i> , <i>Rhus</i> , <i>Schinus</i>), Apocynaceae (<i>Nerium</i>), Ebenaceae (<i>Diospyros</i>), Fabaceae (<i>Delonix</i>), Fagaceae (<i>Fagus</i>), Lecythidaceae (<i>Barringtonia</i>), Rosaceae (<i>Prunus</i>), Sapindaceae (<i>Deinbollia</i>), Verbenaceae (<i>Petrea</i>)	ROBINSON et al (2010), DE PRINS & DE PRINS (2015)
176. <i>Gynanisa maja</i> (Klug, 1836)	Angola, Botswana, Chad, Democratic Republic of Congo, Eritrea, Kenya, Malawi, Mozambique, Namibia, Nigeria, Senegal, Somalia, South Africa, Tanzania, Zambia, Zimbabwe	Fabaceae (<i>Acacia</i> , <i>Berlinia</i> , <i>Brachystegia</i> , <i>Cassia</i> , <i>Colophospermum</i> , <i>Elephantorrhiza</i> , <i>Julbernardia</i> , <i>Laburnum</i> , <i>Peltophorum</i> , <i>Pterocarpus</i> , <i>Robinia</i> , <i>Schotia</i>), Fagaceae (<i>Quercus</i>), Rosaceae (<i>Prunus</i>)	ROBINSON et al (2010), DE PRINS & DE PRINS (2015)
Saturniini			
177. <i>Actias selene</i> (Hübner, 1807)	Afghanistan, India, China, Korea, Sundaland, Andaman Islands	Anacardiaceae (<i>Lannea</i> , <i>Malosma</i> , <i>Mangifera</i> , <i>Schinus</i>), Betulaceae (<i>Alnus</i> , <i>Betula</i>), Combretaceae (<i>Terminalia</i>), Coriariaceae (<i>Coriaria</i>), Corylaceae (<i>Corylus</i>), Ericaceae (<i>Andromeda</i> , <i>Lyonia</i> , <i>Rhododendron</i>), Euphorbiaceae (<i>Sapium</i>), Fagaceae (<i>Castanea</i> , <i>Quercus</i>), Hamamelidaceae (<i>Liquidambar</i>), Juglandaceae (<i>Juglans</i> , <i>Pterocarya</i>), Lauraceae (<i>Cinnamomum</i>), Lythraceae (<i>Lagerstroemia</i> , <i>Lawsonia</i>), Malvaceae (<i>Hibiscus</i>), Meliaceae (<i>Azadirachta</i> , <i>Cedrela</i>), Moringaceae (<i>Moringa</i>), Oleaceae (<i>Ligustrum</i> , <i>Syringa</i>), Rhamnaceae (<i>Frangula</i> , <i>Rhamnus</i>), Rosaceae (<i>Crataegus</i> , <i>Malus</i> , <i>Photinia</i> , <i>Prunus</i> , <i>Pyrus</i>), Rutaceae (<i>Zanthoxylum</i>), Salicaceae (<i>Salix</i>), Symplocaceae (<i>Symplocos</i>)	ROBINSON et al (2010), IOPPOLO (2011), SAVELA (2015)

Tab. 3. – cont.

Lepidopteran taxa	Distribution	Host plants (families and genera) for caterpillars	Data sources
178. <i>Antheraea paphia</i> Linnaeus, 1758	India, Sri Lanka, Thailand	Anacardiaceae (<i>Anacardium</i>), Apocynaceae (<i>Carissa</i>), Bombacaceae (<i>Bombax</i>), Celastraceae (<i>Celastrus</i>), Combretaceae (<i>Anogeissus</i> , <i>Terminalia</i>), Corylaceae (<i>Carpinus</i>), Dipterocarpaceae (<i>Shorea</i>), Ericaceae (<i>Erica</i>), Euphorbiaceae (<i>Hevea</i> , <i>Ricinus</i>), Fabaceae (<i>Bauhinia</i> , <i>Chamaecrista</i> , <i>Dalbergia</i>), Fagaceae (<i>Fagus</i> , <i>Quercus</i>), Lecythidaceae (<i>Careya</i> , <i>Planchonia</i>), Lythraceae (<i>Lagerstroemia</i> , <i>Sonneratia</i>), Meliaceae (<i>Cipadessa</i>), Moraceae (<i>Ficus</i>), Myrtaceae (<i>Eucalyptus</i> , <i>Eugenia</i>), Rhamnaceae (<i>Ziziphus</i>), Rosaceae (<i>Prunus</i>), Rubiaceae (<i>Canthium</i>), Rutaceae (<i>Chloroxylon</i>), Salicaceae (<i>Salix</i>), Sapotaceae (<i>Madhuca</i>), Theaceae (<i>Camellia</i>), Sapindaceae (<i>Dodonaea</i>), Thymeleaceae (<i>Gnidia</i>), Verbenaceae (<i>Tectona</i>)	ROBINSON et al (2010)
179. <i>Argema mimosae</i> (Boisduval, 1847)	Angola, Democratic Republic of Congo, Ethiopia, Kenya, Malawi, Mozambique, Namibia, South Africa, Tanzania, Zambia, Zimbabwe,	Anacardiaceae (<i>Malosma</i> , <i>Sclerocarya</i>), Burseraceae (<i>Commiphora</i>), Euphorbiaceae (<i>Excoecaria</i> , <i>Spirostachys</i>), Hamamelidaceae (<i>Liquidambar</i>), Juglandaceae (<i>Juglans</i>), Myrtaceae (<i>Eucalyptus</i>)	ROBINSON et al (2010), DE PRINS & DE PRINS (2015),
180. <i>Argema mittrei</i> (Guérin-Méneville, 1847)	Madagascar	Altingiaceae (<i>Liquidambar</i>), Anacardiaceae (<i>Cotinus</i> , <i>Malosma</i> , <i>Pistacia</i> , <i>Rhus</i> , <i>Sclerocarya</i> , <i>Schinus</i> , <i>Toxicodendron</i>), Cunoniaceae (<i>Weinmannia</i>), Euphorbiaceae (<i>Uapaca</i>), Fabaceae (<i>Mimosa</i>), Juglandaceae (<i>Juglans</i>), Myrtaceae (<i>Eucalyptus</i> , <i>Eugenia</i>)	OPIE (2015), ROBINSON et al (2010)
181. <i>Ceranchia apollina</i> Butler, 1879	Madagascar	Apocynaceae (<i>Stephanotis</i>)	FISCHER (2014)
182. <i>Cricula agria</i> Jordan, 1909	India	?Fagaceae (<i>Quercus</i>), Lauraceae (<i>Persea</i>), Rosaceae (<i>Prunus</i>)	JORDAN (1909)

Selected phenomena seen during butterfly exhibitions in BG PJŠU

Biodiversity

Though only a small part of world lepidopteran fauna can be seen during exhibitions, the presented species are conspicuous representatives from tropical and subtropical regions from all over the world. They are usually belonging to the families Lycaenidae, Nymphalidae, Papilionidae, Pieridae and Saturnidae.

Approximately 200 species were available to visitors in BG PJŠU, 182 of them were registered as active ones in the greenhouse and dealt with in more detail (Fig. 2, Tab. 3). Identity of species was assigned by suppliers, though not in all cases (missing individual names in mixes of species). Sometimes misidentifications in provided lists also occurred. Polymorphic species were often represented by more than one form.

Reproduction and life stages

In some cases, mating behaviour (Fig. 3), seeking and checking potential host plants (Tab. 2) by females, egg laying and even development of caterpillars (Fig. 4) could be observed.



Fig. 3. Group of *Papilio demoleus* individuals during mating on *Antigonon leptopus*.



Fig. 4. Caterpillars of (from top left to bottom right) *Caligo* sp., *Eryphanis automedon*, *Attacus atlas* and *Siproeta stelenes* from the Victoria greenhouse in BG PJŠU.

Therefore, the visitors could meet not only adult butterflies but also all other stages of their developmental cycle. The details of imago morphology can be seen from the close view even at live butterflies, as many of the presented species are not timid. Eggs and caterpillars were available especially in *Caligo* sp. (on plants of the genera *Musa*, *Heliconia*, *Calathea*, *Maranta*, *Alpinia* and others), more rarely in other lepidopteran species. Various types of pupae from suppliers can be seen arranged in boxes.

Types and functions of wing colorations

Wide variety of wing colorations in butterflies and moths is one of the main reasons of attractiveness of these insects for visitors. Such colours and patterns evolved for different purposes, enhancing survival probability in natural environments. Different types of such butterfly wings are discussed e.g. by HOSKINS (2015).

Hiding from predators

There are many means which are used by butterflies to hide themselves from predators. HOSKINS (2015) shows the following types:

- camouflage as colour, pattern and texture adaptations that enables to blend against background
- disguise - appearance similar to another natural object, e.g. leaf (examples of imitations of dead brown leaves in Fig. 5)
- disruptive coloration
- transparent wings

Aposematic coloration

Aposematic coloration means distinct colours and patterns (Fig. 6) for warning predators that a potential prey species is unpalatable, toxic or otherwise dangerous. According to HOSKINS (2015), all vertebrates including insectivorous birds associate greens and blues with safety, and inherently regard red, orange, yellow and white as signs of danger. Next to it, birds can remember unpleasant experiences associated with their attempts to eat unpalatable butterflies. In the future, such birds are avoiding butterflies with coloration associated with bad taste. However, such learning process could be associated with high mortality of butterfly individuals, it depends also on species of specialized insectivorous birds (e. g. PINHEIRO 1996, 2004).

Diematic patterns

According to HOSKINS (2015), diematic patterns are defensive markings which have the effect of startling or frightening potential predators. This type is often represented by a pair of false-eyes on hind wings of some Saturnidae moths (Fig. 7). Snake head-like forms at tips of fore wings in some other moths (*Attacus* sp. or *Rothschildia* sp.) are also supposed to have this effect and can be seen in live butterfly exhibitions in BG PJSŮ.



Fig. 5. Wings imitating dead leaves, from top left to bottom right: *Consul fabius*, *Hypna clytemnestra*, *Elymnias hypermnestra*, *Doleschalia bisaltide*, *Fountainea nobilis*, *F. eurypyle*, *Salamis cacta*, *Kallima* sp.



Fig. 6. *Idea leuconoe* - both adult and pupae possess aposematic coloration showing their unpalatability.



Fig. 7. Eye spots on hind wings in African moths *Bunea alcinoe* (left) and *Gyanisa maja* (right) (both Saturnidae) manifested after irritation to deter potential predators.

Diverting patterns

It is supposed that effect of eye spots can be deterrent only temporarily and later, on the contrary, these spots can be targets of attack. For that reasons, ocelli on the underside of the wings in many butterfly species (Fig. 8) can possibly serve in diverting attention of predators away from the butterfly's body. After attack directed to false eyes on the border of the wings, a butterfly can gain time to escape.



Fig. 8. Eyespots on underside of the wings, from top left to bottom right: *Morpho helenor*, *Caligo telamonius*, *Opsiphanes tamarindi*, *Eryphanis automedon*, *Mycalesis intermedia*, *Archaeoprepona demophon*, *Salamis parhassus*, *Morpho polyphemus*.

Probably for similar reasons, some butterflies possess streaks on the underside of the wings (Fig. 9) to lead attention to the opposite end of the body. It is sometimes accompanied with a spot like a false eye, or even with tail-like protuberances simulating antennae apart from the vulnerable main body parts (HOSKINS 2015).



Fig. 9. *Amathusia phidippus* (left) and *Colobura dirce* (right) with streaks leading attention of potential attacker to the rear end of butterfly wings.

Other possible functions of wing coloration

There can be different wing patterns serving in communication between individuals of the same butterfly species. For example intensive blue flash coloration on the upperside of *Morpho* species is supposed to be of the mentioned purpose, e.g. during mating behavior. But sudden closing the wings can cause also sudden disappearing of the bright butterfly coloration from predators and exposing its eye spots on underside of the wings to them.

Dark colours of wings in some butterflies can serve for soaking up heat in the sun which can be important for their thermoregulation (HOSKINS 2015).

Mimicry

Similar colour patterns of butterfly wings in different species from the same region are usually regarded as examples of mimicry. Previously discussed aposematic coloration and unpalatability are basic assumptions for mimicry cases which can be often visible also in live exhibitions. Unpalatability is usually a result of **toxicity of food plants** used by caterpillars and toxic properties are consequently transferred to pupae and bodies of imagines. Eventually, toxic compounds can be sequestered also by adult butterflies by feeding on some plant species (BROWER et al. 2010, CARDOSO & GILBERT 2013, HOSKINS 2015).

Mimetic phenomena are widely occurred within lepidopteran insects (KOMÁREK 2004), the same colorations can be shared not only by adult butterflies or moths

but also by caterpillars of different species. Though chemical mimicry is also described in insects (including Lepidoptera), most of known cases is connected just with expressive aposematic colorations.

True mimicry with different species sharing the same coloration evolved due to the pressure from visually oriented predators, especially birds (e.g. PINHEIRO 2004). Usually one of participants in mimicry relationship is a model, the other one is a mimic which is visually adapted to model. Batesian mimicry represent relationship in which a model is unpalatable and a palatable mimic is parasitising on model species (by increasing its own protection from predator attacks and decreasing it in a model). In Müllerian mimicry type both participants are unpalatable and sharing similar wing coloration and patterns, it symbiotically reduces the risk of attack from unexperienced predators.

Some of butterfly species in live exhibitions are co-occurring also in their natural habitats and several of them forming mimetic pairs (of Batesian or Müllerian type) or more numerous mimicry rings. Mimicry ring is a group of sympatric species (or their forms) sharing a common warning pattern (JORON & MALLET 1998). Usually it is a complex of unpalatable species, together with some palatable ones, that have converged on the same colour pattern (MALLET & GILBERT 1995).

Visitors of BG PJSŮ could see the participants in mimicry relationships between butterflies from different parts of the world. In the following cases at least 2 species from the same mimicry ring or mimetic pairs (and usually corresponding subspecies or forms from the same region) have been presented:

Central and South America:

- 'Tiger' mimicry ring (MALLET & GILBERT 1995): unpalatable butterflies (some forms of *Heliconius ismenius*, *H. hecale*, *H. numata*, *Mechanitis polymnia*, *Lycorea halia*, *Tithorea harmonia*, *T. tarricina*, *Eueides isabella*) and palatable species (*Consul fabius*) (Fig. 10).
- 'Orange' mimicry ring (MALLET & GILBERT 1995): unpalatable *Dryas julia*, *Dione juno*, *Dryadula phaetusa*, *Agraulis vanillae* (Fig. 11).
- 'Blue' mimicry ring (MALLET & GILBERT 1995): unpalatable *H. sara*, *Laparus doris* (Fig. 12)
- Several representatives from at least 29 pairs of corresponding geographical forms of *Heliconius erato* and *H. melpomene* (Fig. 13) with almost identical color pattern in each pair (HOSKINS 2015). This includes also 'red' mimicry ring according to MALLET & GILBERT (1995), where some forms of these two unpalatable species (and some other *Heliconius* species) with similar red patterns are classified.
- Male of sexually dimorphic palatable *Papilio erostratus* species mimics unpalatable *Battus polydamas*, while its female mimics different unpalatable model - *Parides photinus* (KUNTE 2009). Palatable *Euritides thymbraeus* mimics *P. photinus* too (COLLINS & MORRIS 1985), while palatable females of some forms of *P. astyalus* and *P. androgeus* mimic unpalatable *B. polydamas* (SPADE et al. 1988) (Fig. 14).



Fig. 10. Representatives of 'tiger' mimicry ring (from top left to bottom right): *Tithorea tarricina*, *Heliconius hecale*, *H. ismenius metaphorus*, *Eueides isabella*, *Mechanitis polymnia*, *Lycorea halia*, *Tithorea harmonia*, *H. numata*, *Consul fabius*, *H. ismenius telchinia*.



Fig. 11. Orange mimetic ring represented by *Dryas julia* (top left), *Dione juno* (top right), *Agraulis vanillae* (bottom left) and *Dryadula phaetusa* (bottom right).



Fig. 12. Mimetic pair of *Lparagus doris* - blue form (top) and *Heliconius sara* (bottom), an example of Müllerian mimicry.



Fig. 13. Different forms of *Heliconius erato* (top row) and *H. melpomene* (bottom row). Corresponding mimetic pairs from the same region are in columns, each such pair represents an example of Müllerian mimicry.



Fig. 14. Top row: *Battus polydamas* (top left) and its mimics *Papilio erostratus* ♂ (top middle) and *Papilio astyalus* ♀ (top right). Bottom row: *Parides photinus* (bottom left) and its mimics *P. erostratus* ♀ (bottom middle) and *Mimoides thymbraeus* (bottom right). *P. erostratus* is here as an example of dimorphic mimicry where male mimics different model than its female.

- Palatable *P. anchisiades*, female of *P. torquatus* and unpalatable *Parides* sp. mentioned by HOSKINS (2015) and palatable *Eurytides ilus branchus* and unpalatable *Parides eurimedes* according to COLLINS & MORRIS (1985) deal similar wing patterns, at least in some of their forms (Fig. 15).
- Palatable *Siproeta stelenes* and unpalatable *Philaethria dido* (HOSKINS 2015) (Fig. 16).



Fig. 15. Non mimetic ♂ *Papilio torquatus* (top left) and its ♀ (top middle) mimicking unpalatable *Parides* sp., like *P. iphidamas* (♀, bottom middle) and *P. eurimedes* (♀, bottom right). *Papilio anchisiades* (top right) and *Mimoides ilus branchus* (bottom left, captured in a spider's web) are also mimicking these *Parides* models.



Fig. 16. Mimetic pair of palatable *Siproeta stelenes* (left) and unpalatable *Philaethria dido* (right), an example of Batesian mimicry.

Africa

- Female of palatable *Hypolimnas misippus* and corresponding model form of unpalatable *Danaus chrysippus* (PUNNETT 1915) (Fig. 17), both species are widespread also in Asia and Australia.
- Palatable *Papilio dardanus* and unpalatable *Danaus chrysippus* (COLLINS & MORRIS 1985), though only some not corresponding mimetic forms of both species could be seen in BG PJSU.



Fig. 17. Non-mimetic ♂ of *Hypolimnas missipus* (left column) and its ♀ (middle column) mimicking *Danaus chrysippus* (right column)

South-Eastern Asia

- Unpalatable *Euploea* sp. (*E. core*) are supposed to be models for some visually similar mimicking forms of palatable *Papilio clytia* and females of *Hypolimnas bolina* (PUNNETT 1915) (Fig. 18). Some other form of *P. clytia* (not presented in BG PJŠU) mimics also other available distasteful species – *Tirumala septentrionis* (COLLINS & MORRIS 1985).
- Unpalatable *D. chrysippus* mentioned above among African species is very widespread and in Asia it is mimicked also by other species, e.g.: *Cethosia cyane* (male) and *Elymnias hypermnestra* (STOLBERG 2014) (though only not corresponding form occurred in BG PJŠU).
- Unpalatable *T. septentrionis* is a model also for females of palatable *Pareronia valeria* (KUNTE 2008), though only non-mimetic male of this species was seen in BG PJŠU.



Fig. 18. *Euploea core* (left), one of *Euploea* sp. mimicked by some forms of *Papilio clytia* (middle) and by ♀♀ of *Hypolimnas bolina* (right).

There could be seen also individual species from other mimetic groups in BG PJSU, though without the respective counterparts. There were diverse forms of palatable mimics (*Euxanthe wakefieldi*, *Papilio echerioides*, *P. memnon*, *P. polytes*, *P. polyxenes*, *Phalanta phalantha*) and distasteful individual model species from not corresponding groups (*Godyris nero*, some *Heliconius* sp. not mentioned above, *Idea leuconoe*, various *Ideopsis* species, *Methona confusa*) (ROGERS 1911, COLLINS & MORRIS 1985, BECCALONI 1995, BROWER 2010, HERBISON-EVANS & CROSSLEY 2015, JORON 2015, NEUBAUER 2015, SAVELA 2015).

In many cases, visitors of such live butterfly exhibitions can see also differences in flight characteristics in individual representatives of mimetic groups. Palatable species with erratic and substantially faster flight are usually in contrast to unpalatable butterflies which tend to fly more slowly and regularly (DUDLEY 2000).

However, it should be mentioned, that not all cases with similarly looking wing patterns must represent mimicry relationships. Convergent evolution caused by some other environmental factors (except predation pressure mentioned above) should also be taken into account. Therefore, these phenomena are still under investigation and individual experts can differ in their opinions regarding mimetic relations between individual lepidopteran species (some differences are discussed e.g. by KUNTE 2008).

Butterfly – plant interactions

Interactions between lepidopteran and plant taxa are of principal importance in ecology of the respective species (their overall characters, way of life, behavior and distribution in nature). Especially it is the case of host plants for caterpillars but in many species feeding of adults can be also of principal importance. Mimicry relationships discussed above are a good example of interconnections between plant characteristics and butterflies and moths as primary consumers formed by selective pressure from predators, all in feedbacks.

In the Victoria greenhouse of BG PJSU, many potential host plants for presented butterflies are available (Tab. 2). Only in some cases these plants are really used for laying eggs as a food source for caterpillars (Fig. 4) but it is enough for a good demonstration. Considering plants from other greenhouses, there are much more potential hostplants (MÁRTONFIOVÁ et al. 2010) but of course, most of these plants are rare individuals growing in BG PJSU and they are not intended for such experiments. Nevertheless, visitors could be properly informed that enormous richness of higher plant taxa in nature is considerably influenced by similarly variable lepidopterans and vice-versa. Live butterfly show is a great opportunity for effective education on these selected aspects in real ecology, not through abstract constructions.

As for imagines, some of the species can use food elements from living plants (mostly nectar from flowers, Fig. 19), plant products (fresh or rotting fruits offered on special feeder devices) or other sources. The observed cases of using different food by individual butterfly species in BG PJSU are summarised in Tab. 4.



Fig. 19. *Papilio demodocus* and *P. memnon* feeding from flowers of *Musa velutina* (top) and a group of *Caligo telamonius memnon* butterflies feeding on rotting bananas on the same inflorescence about 2 months later (bottom) in the Victoria greenhouse of BG PJŠU.

Natural enemies

The overall illusion of natural tropical environment with different ecological relations between butterflies and plants can be completed also by actions of some natural enemies. Though most of the presented lepidopteran species are naturally short-lived in imaginal stage, some of them can be killed also untimely. Of course, this is undesirable from the viewpoint of organizers with their usual efforts to show as much healthy butterflies and moths as possible to visitors. On the other hand, such occasional incidents can indicate what are everyday natural threats the butterfly and moths must withstand in their specific ecosystems all over the world.

Since 2006 there were used no pesticides in most of exposition greenhouses (including the Victoria greenhouse) in BG PJSU and only biological control measures against plant pests were applied. It forms suitable, non-toxic conditions for butterflies and other animals kept in greenhouses for visitors (e.g. fishes in the basins, birds in cages etc.). Predatory and parasitic insects and mites and enthomopatogenic fungi and bacteria are not only these types intentionally released to suppress plant pests but there are also other organisms widely occurred spontaneously within greenhouses. Some of them are able to attack also exotic butterflies of all stages (Fig. 20).



Fig. 20. Some of the natural enemies of butterflies seen in the Victoria greenhouse of BG PJSU: pupa of Papilionidae parasitised with *Pteromalus* wasps (left); *Papilio anchisiades* captured by a tangle-web spider *Parasteatoda tepidariorum* (top middle); *Morpho helenor peleides* fallen into the water basin and attacked by fishes *Carassius auratus* (bottom middle); caterpillar of *Caligo* sp. infected probably with bacterial pathogens (right).

Conclusions

Live butterfly exhibitions provides unusual experiences for different categories of people spending time at such events. Everyone there can gain new experiences and inspiration, from childrens, through common general public, to more demanding students, amateurs or professionals in various areas of natural sciences. Just contact with such interesting living forms has apparent **educational effects**. This influence can be properly reinforced by suitable organization, illustrative tables, propagative materials, articles and web pages to allow visitors to see wider context. In this sense, the exhibition spaces with wide range of plants utilizable by butterflies are exceptionally important for demonstrations of ecological relationships. Botanical gardens are institutions best able to provide such environment outside the areas of natural occurrence of both lepidopteran and plant species.

Particular attention in BG PJŠU is paid especially to the aspects usually not dealt with (or only superficially) in similar exhibitions elsewhere. Above all, various lepidopteran-plant interactions and other phenomena more or less directly related to these interactions (like mimicry, influencing natural enemies, etc.) are discussed and presented in suitable form. However, much more can be seen there by visitors. Last but not least, these exhibitions could help people to recognize the immense diversity of living forms, while 182 lepidopteran species with their host plant genera in Tab. 3 are only a small sample of real richness and possible relationships of these groups in the world.

Next to the educational effects in the sense of new knowledge in entomology, botany, ecology, evolutionary theory, etc., such exhibitions could be very important also from the viewpoint of **nature protection** awareness. Biodiversity of lepidopteran species is strongly depended on healthy ecosystems in the respective countries. In this regard, local economy in tropical areas based on pupae production or tourism connected to butterfly species richness could be acceptable alternative to the destructive exploitation of tropical forests through logging, mining or transformation to agricultural land (SAUL-GERSHENZ 2009).

Under certain conditions, spaces in botanical gardens can be used also for dealing with specific **research problems** related (not only) to lepidopteran-plant interactions. It is possible to form specific qualitative and quantitative composition of plants in an area with controlled abiotic factors (temperature, humidity, light, soil properties). In such space, selected populations of butterflies can live (either from pupae delivered by external suppliers, or from own rearing in separate spaces, if possible). The range of research tasks that could be solved on such basis could be limited only by available technique or laboratory equipment in botanical gardens or their allied institutions.

There are many important groups of insects (and other organisms) but most of them are usually not such favourite like butterflies in wide public. This primary interest in Lepidoptera should be used, e.g. through live butterfly exhibitions, to increase people's knowledge on nature in its complex. Consecutively, such stimulated people could decide to study it in more detail or at least better perceive and eventually participate in solving rising environmental problems.

References

- BECCALONI G.W. (1995): Studies on the ecology and evolution of neotropical Ithomiine butterflies (Nymphalidae: Ithomiinae). A thesis submitted for the degree of Doctor of Philosophy of the University of London, 251 pp.
- BROWER A.V.Z. (2007): *Danaus plexippus* (L. 1758). The Monarch Butterfly. Version 02 August 2007 (under construction). World Wide Web electronic publication, http://tolweb.org/Danaus_plexippus/76926/2007.08.02 in The Tree of Life Web Project, <http://tolweb.org/> (accessed 20-November-2015)
- BROWER A.V.Z. (2010): *Ideopsis* Horsfield 1858. Glassy Tigers. Version 21 June 2010 (under construction). World Wide Web electronic publication, <http://tolweb.org/Ideopsis/70709/2010.06.21> in The Tree of Life Web Project, <http://tolweb.org/> (accessed 28-November-2015)
- BROWER, A. V. Z., WAHLBERG N., OGAWA J. R., BOPPRÉ M. & VANE-WRIGHT R.I. (2010): Phylogenetic relationships among genera of danaine butterflies (Lepidoptera: Nymphalidae) as implied by morphology and DNA sequences. *Systematics and Biodiversity* 8(1): 75–89.
- CARDOSO M.Z. & GILBERT L.E. (2013): Pollen feeding, resource allocation and the evolution of chemical defence in passion vine butterflies. *Journal of Evolutionary Biology* 26: 1254-1260.
- CAST M. (2015): La Diversité des Heliconius. World Wide Web electronic publication, <http://cliniquevetodax.com/Heliconius/index.html> (accessed 20-November-2015)
- COLLINS N.M. & MORRIS M.G. (1985): Threatened Swallowtail Butterflies of the World: the IUCN Red Data Book. IUCN, Gland, Switzerland and Cambridge, UK, 401 pp. + 8 cl plates.
- DAS R.P., ROY A.B., POLLEY R. AND SAHA G. (2010): "A new record of larval host plant of tawny coster *Acraea violae* (Fabricius)". *J. Bombay Nat. Hist. Soc.* 107 (1): 63.
- DAY L. (2015a): Samui Butterflies. World Wide Web electronic publication, www.samuibutterflies.com/ (accessed 20-November-2015)
- DE PRINS J. & DE PRINS W. (2015): Afromoths, online database of Afrotropical moth species (Lepidoptera). World Wide Web electronic publication, www.afromoths.net (accessed 20-November-2015)
- DICKSON C. G. C. & KROON D. M. (eds.) (1978): Pennington's Butterflies of Southern Africa. Johannesburg : Ad. Donker, 670 pp.
- DUDLEY R. (2000): The Biomechanics of Insect Flight: Form, Function, Evolution. Princeton: Princeton University Press, 476 pp.
- DUNN K.L. & DUNN L. E. (1991): Review of Australian butterflies: distribution, life history and taxonomy. Part 1. Introduction, Papilionidae, Pieridae and adult temporal data. Privately published by the authors, Melbourne. pp.1-196.
- FISCHER M. (2014): World of Silkmoths. World Wide Web electronic publication, <http://silkmoths.eu/index.html> (accessed 28-November-2015)
- FREITAS A.V.L., VASCONCELLOS-NETO J., VANINI F., TRIGO J. R., BROWN K.S. jr. (2001): Populations studies of *Aeria olena* and *Tithorea harmonia* (Nymphalidae, Ithomiinae) in southeastern Brazil. *Journal of the Lepidopterists' Society* 55(4): 150-157.
- GODMAN F. D. & SALVIN O. (1879-1901): *Insecta. Lepidoptera-Rhopalocera*. Vol. I., *Biologia Centrali Americana*, R.H. Porter, London, 487 pp.
- GOSWAMI R. (2013): "*Danaus chrysippus* (Linnaeus 1758): Trophic Strategy". *Encyclopedia of Life*. World Wide Web electronic publication, http://eol.org/data_objects/26326900 (accessed 28-November-2015)

- HASSLER M. (2015a): World Ferns: Checklist of Ferns and Lycophytes of the World (version Sep 2015). In: ROSKOV Y., ABUCAY L., ORRELL T., NICOLSON D., KUNZE T., FLANN C., BAILLY N., KIRK P., BOURGOIN T., DEWALT R.E., DECOCK W., DE WEVER A., (eds.): Species 2000 & ITIS Catalogue of Life, 27th October 2015. Digital resource at www.catalogueoflife.org/col. Species 2000: Naturalis, Leiden, the Netherlands. ISSN 2405-8858. (accessed 28-November-2015)
- HASSLER M. (2015b): World Plants: Synonymic Checklists of the Vascular Plants of the World (version Sep 2015). In: ROSKOV Y., ABUCAY L., ORRELL T., NICOLSON D., KUNZE T., FLANN C., BAILLY N., KIRK P., BOURGOIN T., DEWALT R.E., DECOCK W., DE WEVER A., (eds.): Species 2000 & ITIS Catalogue of Life, 27th October 2015. Digital resource at www.catalogueoflife.org/col. Species 2000: Naturalis, Leiden, the Netherlands. ISSN 2405-8858. (accessed 28-November-2015)
- HENDERSON C.L. (2002): Field guide to the Wildlife of Costa Rica. Austin, Texas: University of Texas Press, 559 pp., ISBN 0-292-73459-X.
- HERBISON-EVANS D. & CROSSLEY S. (2015): Australian Caterpillars and their Butterflies and Moths. World Wide Web electronic publication, <http://lepidoptera.butterflyhouse.com.au/> (accessed 28-November-2015)
- HOLLOWAY J.D. (1987): The Moths of Borneo, Part 3; Superfamily Bombycoidea: families Lasiocampidae, Eupterotidae, Bombycidae, Brahmaeidae, Saturniidae, Sphingidae. Kuala Lumpur: Southdene, 199 pp.
- HOLLOWAY J.D. (1996): The Moths of Borneo, Part 9; Family Geometridae, Subfamilies Oenochrominae, Desmobjathrinae and Geometrinae. Malayan Nat. J. 49: 147-326.
- HOSKINS A. (2015): Learn about butterflies. The complete guide to the world of butterflies and moths. World Wide Web electronic publication, <http://www.learnaboutbutterflies.com/> (accessed 6-November-2015)
- INAYOSHI, Y. (2015): A Check List of Butterflies in Indo-China (Chiefly from Thailand, Laos & Vietnam). World Wide Web electronic publication, <http://yutaka.it-n.jp/index.html> (accessed 28-November-2015)
- IOPPOLO M. (2011): Scheda di allevamento di *Actias selene*. 8 pp. World Wide Web electronic publication, <http://www.farfalledalmondo.it/schede/schedaselene.pdf> (accessed 28-November-2015)
- JIGGINS CH. (2008): *Heliconius himera* Hewitson 1867. Version 12 August 2008 (under construction). http://tolweb.org/Heliconius_himera/72241/2008.08.12 in The Tree of Life Web Project, <http://tolweb.org/> (accessed 28-November-2015)
- JORDAN K. (1909): On the Species of *Cricula*, a Genus of Saturniidae. Novitates Zoologicae 16(1): 300-306.
- JORON M. & MALLETT J.L.B. (1998): Diversity in mimicry: paradox or paradigm? Trends in Ecology and Evolution 13(11): 461-466.
- JORON M. (2015): Butterfly Ecology and Evolution. World Wide Web electronic publication, <http://isyeb.mnhn.fr/joron/index.html> (accessed 28-November-2015)
- KHOT R. & GAIKWAD K. (2011): An additional record of larval host plant of Tawny Coster *Acraea violae* (Fabricius 1775). J. Bombay Nat. Hist. Soc. 108 (2): 140.
- KOMÁREK S. (2004): Mimikry, aposematizmus a příbuzné jevy. Mimetismus v přírodě a vývoj jeho poznání. Dokořán, 189 pp.
- KUNTE K. (2008): Evolution of sex-limited mimicry in swallowtail butterflies. Dissertation presented to the Faculty of the Graduate School of The University of Texas at Austin, 132 pp.
- KUNTE K. (2009): The diversity and evolution of Batesian mimicry in *Papilio* swallowtail butterflies. Evolution 63(10): 2707-2716.
- LARSEN T.B. (1996): The Butterflies of Kenya and their natural history. Oxford University Press, 500 pp.

- LATHAM P. (2015): Edible caterpillars and their food plants in Bas-Congo Province, Democratic Republic of Congo. 3rd edition, 44 pp., ISBN: 9780992898625
- LEWIS D.S. (2009): Lime Swallowtail *Papilio demoleus* Linnaeus (Insecta: Lipidoptera: Papilionidae). EENY 444, one of a series of the Entomology and Nematology Department, UF/IFAS Extension, University of Florida, 5 pp.
- LEWIS D.S.A. (2010): Phylogeny and revision of the genus *Heraclides* Hübner, 1819 (Lepidoptera: Papilionidae: Papilioninae: Papilionini). A dissertation presented to the Graduate School of the University of Florida in partial fulfillment of the requirements for the degree of doctor of philosophy. University of Florida, 301 pp.
- MAES J.-M. (2006): Papilionidae (Lepidoptera) de Nicaragua. Rev. Nica. Ent., 66 (2006), Suplemento 3, 241 pp.
- MALLET J. & GILBERT L.E. (1995): Why are there so many mimicry rings? Correlations between habitat, behaviour and mimicry in *Heliconius* butterflies. Biological Journal of the Linnean Society, 55: 159–180.
- MÁRTONFIOVÁ L., MOCHNACKÝ S., FRIDMAN P., REPČÁKOVÁ K., KELBEL P., GREGOREK R. (2010): Catalogue of plant collections no. 5, P.J.Šafárik University Botanical Garden. PJSU Košice, 96 pp.
- MÖHN E. (2002): Papilionidae VIII. *Baronia*, *Euryades*, *Protographium*, *Neographium*, *Eurytides*. In: BAUER E. & FRANKENBACH T. (eds.): Butterflies of the World. Keltern : Goecke & Evers ; Canterbury : Hillside Books. Part 14, 36 plates + 12 pp., ISBN 978-3-931374-87-7
- MÖHN E. (2006): Papilionidae XIII. *Parides*. Butterflies of the World. Part 26. Antiquariat Goecke & Evers, Keltern, Wohmbrechts, Germany. 84 pp.
- MORISSON C. (2014): *Nessaea aglaura aglaura* (Doubleday, 1848). Parasitoid-Caterpillar-Plant Interactions in the Americas. World Wide Web electronic publication, <http://caterpillars.myspecies.info/node/2236#> (accessed 22-November-2015)
- MURILLO-HILLER L.R. (2013): Early Stages and Natural History of *Mimoides pausanias prasinus* and *M. euryleon clusoculis* (Lepidoptera, Papilionidae) from Costa Rica. Annual Review & Research in Biology 3(1): 22-30.
- MUYSHONDT A. (1974): Notes on the life cycle and natural history of butterflies of El Salvador. III. *Anaea (Consul) fabius* J. Lepid. Soc. 28: 81-89.
- MUYSHONDT A. (1976): Notes on the life cycle and natural history of butterflies of El Salvador. VIII. *Archaeoprepona antimache gulima*, *Siderone marthesia*, *Zaretis callidryas* and *Consul electra* (Nymphalidae). J. Lepid. Soc. 30: 159-168.
- NAIR P. V. (2005): *Cosmostigma racemosa*, a new host plant record of the Dark Blue Tiger Butterfly *Tirumala septentrionis* from Kerala. Journal of the Bombay Natural History Society. 102(3): 354.
- NEUBAUER T. (2015): Butterflycorner.net. World Wide Web electronic publication, <http://www.butterflycorner.net/> (accessed 28-November-2015)
- OPIE (2015): Papillon-comète de Madagascar (*Argema mittrei*). Office pour les Insectes et leur Environnement. World Wide Web electronic publication, <http://www.insectes.org/argema-mittrei/papillon-comete-de-madagascar.html> (accessed 28-November-2015)
- PAGE M.G.P., TREADAWAY C.G., BAUER E., FRANKENBACH T. (2003): Schmetterlinge der Erde. Butterflies of the world. Part 17, Papilionidae IX.: Papilionidae of the Philippine Islands. Keltern: Goecke & Evers; Canterbury: Hillside Books, pp. 12 + 58 col. Plates; ISBN 978-3-931374-45-7
- PEIGLER R.S. & NAUMANN S. (2003): A revision of the Silkmoth Genus *Samia*. Texas, San Antonio, University of the Incarnate Word. 230 pp., 10 maps, 228 figs., ISBN 0-9728266-0-2

- PAUKSTADT U. & PAUKSTADT L.H. (2002): Beitrag zur Kenntnis der Morphologie von *Attacus caesar* Maassen, 1873 von den Philippinen (Lepidoptera: Saturniidae). *Galathea, Berichte des Kreises Nürnberger Entomologen* 18(1): 44-51.
- PENZ C.M., AIELLO A. & SRYGLEY R.B. (2000): Early stages of *Caligo illioneus* and *C. idomeneus* (Nymphalidae, Brassoliniæ) from Panama, with remarks on larval food plants for the subfamily. *J. Lep. Soc.* 53 (4): 142-152
- PINHEIRO C.E.G. (1996): Palatability and escaping ability in Neotropical butterflies: tests with wild kingbirds (*Tyrannus melancholicus*, Tyrannidae). *Biological Journal of the Linnean Society* 59: 351-365.
- PINHEIRO C.E.G. (2004): Jacamars (Aves, Galbulidae) as selective agents of mimicry in neotropical butterflies. *Ararajuba* 12 (2): 137-139.
- PUNNETT R.C. (1915): *Mimicry in Butterflies*. Cambridge University Press, 185 pp.
- PYLE R.M. (1981): *The National Audubon Society field guide to North American butterflies*. A.A. Knopf, New York. 924 pp., ISBN 0-394-51914-0
- RAJAGOPALAN A. (2005): "A new food plant of the Great Eggfly". *Journal of the Bombay Natural History Society* 102 (3): 355.
- REIMAN GARDENS (2015): *Reiman Gardens Butterflies*, Iowa State University. World Wide Web electronic publication, <http://rgbutterflyapp.com/> (accessed 28-November-2015)
- RIES L. (2012): *Myscelia ethusa*, Mexican Bluewing. North American Ecology (US and Canada). *Encyclopaedia of Life*. World Wide Web electronic publication, <http://eol.org/pages/162223/overview> (accessed 22-November-2015)
- ROBINSON, G. S., P. R. ACKERY, I. J. KITCHING, G. W. BECCALONI & L. M. HERNÁNDEZ (2010): HOSTS - A Database of the World's Lepidopteran Hostplants. Natural History Museum, London. World Wide Web electronic publication, <http://www.nhm.ac.uk/hosts> (accessed: 17 November 2015).
- ROGERS K.S.A. (1911): Mimicry in East african butterflies with special reference to Danaine models. *Journal of the East Africa Society London*, 1: 81-92.
- ROTHSCHILD W. (1895): A Revision of the Papilios of the Eastern Hemisphere, exclusive of Africa. *Novitates Zoologicae* 2(3): 167-463.
- ROTHSCHILD W. & JORDAN K. (1906): A Revision of the American Papilios. *Novitates Zoologicae* 13(3): 411-752.
- SARI H.P.E., RAHAYU S.E., MASJHUDI, FAUZIAH F. (2015): Study of butterfly species at Kondang Merak beach southern Malang area. The 3rd International Conference on Biological Science 2013. *KnE Life Sciences* 2: 579-582. ISSN 2413-0877
- SAUL-GERSHENZ L. (2009): Insect zoos. Pp. 516-523 in: RESH V.G. & CARDÉ R.T. (eds.): *Encyclopedia of insects*, 2nd edition, Academic Press, 1168 pp.
- SAVELA M. (2015): Lepidoptera and some other life forms. World Wide Web electronic publication, <http://www.nic.funet.fi/pub/sci/bio/life/intro.html> (accessed 28-November-2015)
- SCRIBER J.M., LEDERHOUSE R.C. & BROWN K.S. JR. (1990/91): Hybridization of the Brazilian *Papilio* (*Pyrrhosticta*) (Section V) with the North American *Papilio* (*Pterourus*) (Section III). *Journal of Research on the Lepidoptera* 29(1-2): 21-32.
- SPADE P., TYLER H. & BROWN J.W. (1988): The biology of seven Troidine swallowtail butterflies (Papilionidae) in Colima, Mexico. *Journal of Research on the Lepidoptera* 26(1-4): 13-26.
- STOLBERG V.B. (2014): Butterflies. P. 112-114 in: LEVINE T.R. (ed.): *Encyclopedia of Deception*, 1st Edition, SAGE Publications, Inc., 1104 pp.
- TAN H. (2010): Life History of the Archduke (*Lexias pardalis dirteana*). Butterflies of Singapore. World Wide Web electronic publication, <http://butterflycircle.blogspot.sk/2010/02/life-history-of-archduke.html> (accessed: 22 November 2015)

- VAN DER HEYDEN T. (2010): *Orbea variegata* (L.) Haworth, 1812 (Apocynaceae, Asclepiadoideae) als Futterpflanze der Larven von *Danaus chrysippus* (Linnaeus, 1758) auf den Kanarischen Inseln (Spanien) (Lepidoptera: Nymphalidae, Danainae). SHILAP Revista de lepidopterología 38(149): 107-110.
- VANE-WRIGHT, R.I. & DE JONG R. (2003): The butterflies of Sulawesi: annotated checklist for a critical island fauna. Zool. Verh. Leiden 343: 3-267, ISSN 0024-1652/ISBN 90-73239-87-7
- WARREN, A. D., DAVIS K. J., STANGELAND E. M., PELHAM J. P. & GRISHIN N. V. (2013): Illustrated Lists of American Butterflies. World Wide Web electronic publication, <http://www.butterfliesofamerica.com/> (accessed: 27 November 2015)
- WILLIAMS M.C. (2008): Butterflies and skippers of the Afrotropical region (Papilionoidea and Hesperioidea). An encyclopaedia, the 7th edition. World Wide Web electronic publication, <http://atbutterflies.com/> (accessed: 22 November 2015)
- WOODHALL S. (2005): Field Guide to Butterflies of South Africa. New Holland Publishers (NZ) Limited, 440 pp.
- YANG P., QI B., DENG H.-L., CHEN J., LIU Q. (2005): The biology of *Kallima inachis* and its rearing. Journal of Southwest Agricultural University 27 (1): 44–49.
- YOUNG A. M. (1985): Natural history notes on *Brassolis isthmia* Bates (Lepidoptera: Nymphalidae: Brassolinae) in northeastern Costa Rica. J. Res. Lep. 24:385–392.
- ZHAN S., ZHANG W., NIITEPÖLD K., HSU J., HAEGER J.F., ZALUCKI M.P., ALTIZER S., DE ROODE J.C., REPERT S.M., KRONFORST M.C. (2014): The genetics of monarch butterfly migration and warning coloration. Nature, 514(7522): 317–321, doi:10.1038/nature13812.

Received: November 25th 2015
 Revised: December 14th 2015
 Accepted: December 15th 2015