Wyoming Plant and Animal Species of Concern

November 2003

CONTENTS

| Introduction1 |
|---|
| Changes from Previous Lists 1 |
| Codes and Definitions2 |
| Plants ListsP-1 |
| Species of Concern (P-1); Potential Concern (P-30); Dropped (P-33); Uncertain (P-33) |
| Animal ListsA-1 |
| Birds (A-1); Fish (A-6); Amphibians (A-8); Reptiles (A-9); Mammals (A-10); Potential Concern (A-13); Dropped (A-20) |






Douglas Keinath, Zoologist Bonnie Heidel, Botanist Gary Beauvais, Director


Wyoming Natural Diversity Database (WYNDD)

University of Wyoming
Department 3381 // 1000 E. University Avenue
Laramie, Wyoming 82071

PHONE: 307 766-3023 FAX: 307 766-3026



Survey Forms Inside Covers





EMAIL: wndd@uwyo.edu HOMEPAGE: www.uwyo.edu/WYNDD

INTRODUCTION

Interest in rare species has increased substantially over the past 40 years, and currently there is broad support for the conservation of rare plants and animals in North America. Natural resource managers, policy makers, and the public require an understanding of the identity, distribution, and abundance of rare species in order to develop effective conservation strategies. Such information is especially vital to management plans that strive to integrate the conservation of rare species with development of natural resources.

This publication provides the most complete information available on the status of rare vertebrate species and vascular plant species in Wyoming. It updates and replaces previous lists (Fertig and Beauvais 1999, Fertig and Heidel 2002), and documents 473 plants and 125 vertebrates of conservation and management concern in Wyoming. For each species, a summary of factors used in weighing species' status including distribution, abundance, trends, and intrinsic vulnerability, is also provided.

The Wyoming Natural Diversity Database (WYNDD) at the University of Wyoming serves as the information clearinghouse on species of concern in Wyoming. Professional biologists at WYNDD conduct primary field research, literature reviews, and information summaries to maintain a comprehensive database of information on rare plants and animals. Readers are encouraged to help in this effort and improve this list by submitting data on rare species to WYNDD. Rare species observation forms are included in this publication, and are also posted at [www.uwyo.edu/wyndd].

This document, supporting information, and staff contacts are available at: www.uwyo.edu/wyndd
All parties are encouraged to contact WYNDD with comments, new data, or requests for the most current status information.

CHANGES FROM PREVIOUS LISTS

The ranks, statuses, and known distributions of rare species are dynamic due to ongoing research and re-examination of existing data. Differences in these items relative to previous lists reflect WYNDD's ongoing efforts to maintain comprehensive and current information. New species added since 1999 (animals) or 2002 (plants), and any changes in ranks, counties, or managed areas, are indicated in bold. Brackets enclosing species names indicate recent synonyms.

The animal portion of this list represents a major improvement over previous versions. It employs a more systematic and replicable ranking procedure, which is explained in detail in Keinath and Beauvais (2003). A

similar ranking procedure was employed in the 2002 plant list (Fertig and Heidel 2002) with minor revisions in this 2003 application.

The enclosed animal and plant lists introduce the Wyoming Contribution Rank, which provides more precise information on how Wyoming populations of a given taxon contribute to the rangewide persistence of that taxon. Details of this ranking system are in Keinath and Beauvais (2003) and Fertig and Heidel (2003).

The list format has been changed to provide readers with more species-specific information while still maintaining a concise layout. For each species we have included information on range context, abundance, vulnerability, and other factors that contribute to the status ranks. We also provide a more complete listing of species' statuses as determined by other organizations. For example, we now denote which species have been recently petitioned for listing under the Endangered Species Act.

Both the plant and animal lists are organized into a series of sub-lists:

Species of Concern: These are taxa that WYNDD has determined to be of conservation concern within the state of Wyoming. These taxa exhibit a combination of biological factors that predispose them to decline or potential extirpation from the state. Many of these taxa are also given special statuses by federal and state agencies.

Species of Potential Concern: These are taxa for which there is some evidence suggesting a moderate degree conservation concern, including species that are locally common but occupy a restricted range that is vulnerable to landscape changes. For animals, this list also includes species for which basic biological information is anecdotal, incomplete, or otherwise lacking in the detail needed for a reliable assessment and ranking. Also, some animals on this list may also have taxonomic questions yet to be resolved. WYNDD often collects data opportunistically for species of potential concern, but focuses most of its efforts on species of known concern.

Species of Uncertain Status: These are plant taxa that may warrant conservation attention but for which there are substantial questions regarding taxonomy, identification, or continued occurrence in the state.

Dropped: These taxa appeared on previous Species of Concern or Species of Potential Concern Lists, but have been dropped from this 2003 version. Taxa are usually dropped when new data or re-examination of older data indicates a lessening in their degree of conservation concern.

CODES AND DEFINITIONS

Heritage Rank: WYNDD uses a standardized ranking system originally developed by The Nature Conservancy and its network of natural heritage programs (now coordinated by NatureServe [Arlington, Virginia]) to indicate the probability of extinction, at both the global and state scales, of each plant and animal taxon. The following letters denote the spatial scale at which a taxon's status is scored:

- G = Global rank: refers to the range-wide probability of extinction for a species
- T = Trinomial rank: refers to the range-wide probability of extinction for a subspecies or variety
- S = State rank: refers to probability of extinction from WY for a given taxon

These letters are each followed by a numeric, 1-5 score:

- 1 = Critically imperiled because of extreme rarity (often <5 extant occurrences) or because some factor makes it highly vulnerable to extinction
- 2 = Imperiled because of rarity (often 6-20 extant occurrences) or because of factors making it vulnerable to extinction
- 3 = Rare or local throughout its range or found locally in a restricted range (often 21-100 known occurrences)
- 4 = Apparently secure, although it may be quite rare in parts of its range, especially at the periphery
- 5 = Demonstrably secure, although it may be rare in parts of its range, especially at the periphery

Some taxa receive non-numeric scores, indicating special situations:

- H = Known only from historical records (typically pre-1970; varies by taxon)
- A = Accidental or vagrant: taxon appears irregularly and infrequently; *typically applied only to animals*
- X = Believed to be extinct
- U = Uncertain status: taxon possibly in peril but more information is needed

Some taxa may also receive rank modifiers, indicating other special situations:

- B = Breeding rank: indicates the status of a migratory species during the breeding season; applied only to animals
- N = Non-breeding rank: indicates the status of a migratory species during the non-breeding season; applied only to animals

- Z = Rank not applicable: indicates that a migratory species is essentially absent or unrankable for the season in question (used with "B" or "N"); applied only to animals
- Q = Questions exist regarding the taxonomic validity of a species, subspecies, or variety
- ? = Questions exist regarding the assigned numeric score

Wyoming Contribution Rank: WYNDD has developed a ranking system that indicates the contribution of WY populations of a taxon to the rangewide persistence of that taxon.

Very High = WY populations contribute greatly to the taxon's rangewide persistence. Typically applies to native, resident taxa with small continental ranges that are centered on WY

High = In combination with populations in adjacent states, WY populations contribute substantially to the taxon's rangewide persistence. Typically applies to native, resident taxa that are more secure in WY than elsewhere and have a moderate proportion of their continental range in WY

Medium = WY populations contribute to the taxon's rangewide persistence, but are apparently no more critical than populations in other states. Typically applies to native, resident taxa that are as secure in WY as elsewhere and have a moderate proportion of their continental range in WY

Low = WY populations contribute minimally to the taxon's rangewide persistence. Typically applies to taxa that are more prevalent and secure in other states, and occur only intermittently or peripherally in WY

Federal Status:

<u>US Fish and Wildlife Service (USFWS)</u>: The Endangered Species Act of 1973 directs the USFWS to identify and protect rare plants and animals. The following categories are currently used to rank listed and candidate species. The candidate system was revised in 1996, eliminating the old categories of C2 and 3C.

Endangered: In danger of extinction. For plants this applies to all of a taxon's range; for animals this may apply to only a portion of a taxon's range

Threatened: Likely to become Endangered in the near future. For plants this applies to all of a taxon's range; for animals this may apply to only a portion of a taxon's range

Proposed De-list: Formally proposed for de-listing from Endangered or Threatened status; awaiting final rule

Proposed: Formally proposed for listing as Endangered or Threatened; awaiting final rule

Candidate (formerly C1): Existing biological information supports a formal proposal to list as Endangered or Threatened, but developing a listing regulation is precluded by higher priority listing activities.

Recent De-list: Recently de-listed from Endangered or Threatened status

Listing Denied: A formal proposal for listing as Endangered or Threatened was recently denied

Petitioned: A petition to list as Endangered or Threatened has been received by USFWS; awaiting USFWS review

<u>USDA Forest Service (USFS)</u>: USFS has adopted policies to ensure that agency actions do not drive rare taxa towards listing as Threatened or Endangered under the Endangered Species Act.

USFS Region 2 and Region 4 have developed lists of "Sensitive Species" that warrant special attention on USFS lands (USDA FS 1991, 1994, 2003). Sensitive species are defined as "plant and animal species identified by the Regional Forester for which population viability is a concern as evidenced by: (a) significant current or predicted downward trends in population numbers or density, and/or (b) significant current or predicted downward trends in habitat capability that would reduce a species' existing distribution." Sensitive species are usually designated for an entire region, but independent "Forest Sensitive" lists are maintained by some individual National Forests. The following codes are used to represent sensitive species designations by USFS:

USFS R2 Sensitive: Sensitive in Bighorn, Black Hills, Medicine Bow, and Shoshone National Forests, and the Thunder Basin National Grassland

USFS R4 Sensitive: Sensitive in Bridger-Teton, Caribou, Targhee, Wasatch-Cache, and Ashley (including Flaming Gorge National Recreation Area) National Forests

<u>USDI Bureau of Land Management (BLM):</u> BLM has adopted policies to ensure that agency actions do not drive rare taxa towards listing as Threatened or Endangered under the Endangered Species Act.

The BLM Wyoming State Office has established a list of "Sensitive Species" warranting special attention on BLM lands (USDI BLM 2001, 2002). BLM Sensitive species are "species that could easily become endangered or extinct in the state", including (a) species under status review by the FWS/ National Marine and Fisheries Service, (b) species whose numbers are declining so rapidly that Federal listing may become necessary, (c) species with typically small or fragmented populations, and (d) species inhabiting specialized refugia or other unique habitats. The following code is used to represent sensitive species designations by BLM:

WY BLM Sensitive: Sensitive in all Wyoming BLM Field Offices

State Status: The text below applies only to animals. At this time there is no system used by the state of WY to indicate the status of rare native plants.

Wyoming Game and Fish Department (WGF): The WGF uses a 1-4 "Native Species Status" score to rank the status of vertebrate species:

NSS1 = Populations greatly restricted or declining, extirpation possible -OR- Ongoing significant loss of habitat

NSS2 = Populations declining, extirpation possible; habitat restricted or vulnerable but no recent or ongoing significant loss; species likely sensitive to human disturbance -OR- Populations declining or restricted in numbers or distribution, extirpation not imminent; ongoing significant loss of habitat

NSS3 = Populations greatly restricted or declining, extirpation possible; habitat not restricted, vulnerable but no loss; species not sensitive to human disturbance -OR- Populations declining or restricted in numbers or distribution, extirpation not imminent; habitat restricted or vulnerable but no recent or ongoing significant loss; species likely sensitive to human disturbance -OR- Species widely distributed; population status or trends unknown but suspected to be stable; on-going significant loss of habitat

NSS4 = Populations greatly restricted or declining, extirpation possible; habitat stable and not restricted -OR- Populations declining or restricted in numbers or distribution, extirpation not imminent; habitat not restricted, vulnerable but no loss; species not sensitive to human disturbance -OR- Species widely distributed, population status or trends unknown but suspected to be stable; habitat restricted or vulnerable but no recent or on-going significant loss; species likely sensitive to human disturbance -OR-

Populations stable or increasing and not restricted in numbers or distribution; on-going significant loss of habitat

County: WY counties in which rare species occur. Vague or unsubstantiated county records are indicated by "?". County names in brackets indicate former range of now extinct taxa. Records for Yellowstone Nat'l Park are listed by Park or Teton County.

| ALB | Albany | NAT | Natrona |
|-----|-------------|--------|------------|
| BIG | Big Horn | NIO | Niobrara |
| CAM | Campbell | PAR | Park |
| CAR | Carbon | PLA | Platte |
| CON | Converse | SHE | Sheridan |
| CRO | Crook | SUB | Sublette |
| FRE | Fremont | SWE | Sweetwater |
| GOS | Goshen | TET | Teton |
| HOT | Hot Springs | UIN | Uinta |
| JOH | Johnson | WAS | Washakie |
| LAR | Laramie WES | Westor | 1 |
| LIN | Lincoln | | |

Managed Area: Codes show federal and state managed areas in which rare plants are known to occur. Blank entries indicate that a species is known only from other state or privately owned lands. Vague or unsubstantiated records are indicated by "?". Codes are as follows:

| AFB = | Air Force Base |
|-------|---------------------------|
| BLM = | Bureau of Land Management |
| NF = | National Forest |
| NG = | National Grassland |
| NHS = | National Historic Site |
| NM = | National Monument |
| NP = | National Park |
| NRA = | National Recreation Area |
| NWR = | National Wildlife Refuge |
| SP = | State Park |

In addition, records reported from Wind River Indian Reservation are indicated by:

IR = Indian Reservation

Range context: The extent and position of a taxon's continental range relative to WY.

Local endemic: continental range is no larger (approx.) than Natrona Co., WY (72 X 72 mi)

Regional endemic: continental range is no larger (approx.) than the state of WY

Widespread: continental range is much larger than the state of WY

Core: WY encompasses >50% of the continental range

Edge: WY encompasses <50% of the continental range

Disjunct: WY populations are widely isolated (by about 300 mi or more) from the main, contiguous portion of the continental range

Number of Occurrences: Number of extant (documented since 1970) and discrete reproductive populations of a rare plant in WY.

Very Low = 1-5 occurrences
Low = 6-20 occurrences
Moderate = 21-75 occurrences
High = >75 occurrences
Unknown

Abundance: Number of individuals, or area of occupied habitat, of a rare plant in WY.

Rare = <5000 individuals or <500 occupied ac.

Uncommon = 5000-50000 individuals or 500-5000 occupied ac.

Unknown

Trend: Change in number of individuals, occurrences, or area of occupied habitat. *Historical* refers roughly to the period 1850 - 1950. *Recent* generally refers to the period since 1950.


Large decline = >50% decline in numbers, occurrences, or area

Moderate decline = <50% decline in numbers, occurrences, or area

Stable = no substantial decline or increase

Increase = any substantial increase in individuals, occurrences, or area

Unknown



Intrinsic Vulnerability: The susceptibility of a taxon to decline due to inherent biological factors. Such factors include fecundity, population density, dispersal ability, pollinator limitations, competitive ability, likelihood of hybridization, and habitat specialization. Intrinsic vulnerability is scored with an emphasis on conditions in Wyoming that may represent a subset of rangewide conditions.

High = Taxon restricted to 1 rare habitat type -ORexhibits at least 2 of the following characteristics: very low population density, very low dispersal ability, very low fecundity, pollinator limitations, predisposition to disease

Moderate = Taxon restricted to 2-4 habitat types -ORexhibits at least 1 of the following characteristics: very low population density, very low dispersal ability, very low fecundity, pollinator limitations, predisposition to disease

Low = Taxon occurs in a variety of habitat types (usually all within 1 biome) -OR- has no life history traits that predisposes it to declines resulting from moderate environmental changes

ACKNOWLEDGEMENTS - PLANTS

Much of the original data for this report comes from specimens deposited at the Rocky Mountain Herbarium. Additional information and suggestions were provided over the course of this and previous list revisions by Larry Apple, Melanie Arnett, Darcie Bacon, Frank Blomquist, Bernie Bornong, Janet Britt, Greg Brown, Sabine Mellmann-Brown, Beth Burkhart, Jeff Carroll, Reed Crook, Jean Daly, Charmaine Delmatier, Robert and Jane Dorn, Katy Duffy, Erwin Evert, Walter Fertig, Ben Franklin, Jim Glennon, Wendy Haas, Ron Hartman, Don Hazlett, Pat and Noel Holmgren, Kent Houston, Kelly Hughes, Ann Humphrey, George Jones, Greg Karow, John Kartesz, Lynn Kinter, Andy Kratz, Clay Kyte, Steve Laster, Claire Leon, Michael Mancuso, Stuart Markow, Hollis Marriott, Jerry Mastel, Bob Moseley, Mary Neighbours, B. Ernie Nelson, Dave Ode, Kevin O'Dea, Jim Ozenberger, Barb Packer, John Proctor, Deana Reyher, Richard and Bev Scott, Phil Shephard, Carolyn Sieg, Susan Spackman, Gerry Steinauer, Amy Taylor, Laura Welp, Jennifer Whipple, and Kathy Zacharkevics.

ACKNOWLEDGEMENTS - ANIMALS

Information and suggestions were provided over the course of list revision by Stan Anderson, Merav Ben-David, Jason Bennett, Frank Blomquist, Tim Byers, Steven Buskirk, Jeff Carroll, Andrea Cerovski, Darby Dark-Smiley, Robert Dorn, Eric Everett, Martin Grenier, Jeff Gruver, Greg Hayward, Mary Jennings, Laine Johnson, Steve Koslowski, Robert Luce, Larry Master, Matthew McGee, Carron Meaney, Bob Oakleaf, Deb Patla, Charles Peterson, Regan Plumb, Tom Rinkes, Brian Smith, Patricia Sweanor, Michelle VanVleet, and Jon Warder.

ACKNOWLEDGEMENTS - ARTWORK

Columbian sharp-tailed grouse, Colorado River cutthroat trout, and Canada lynx by *Summers Scholl*. Yellow lady's-slipper by *Walter Fertig*. Desert yellowhead by *Jane Dorn*.

LITERATURE CITED

- Fertig, W. and G.P. Beauvais. 1999. Wyoming Plant and Animal Species of Special Concern. Wyoming Natural Diversity Database, University of Wyoming, Laramie.
- Fertig, W. and B. Heidel. 2002. Wyoming Plant Species of Special Concern. Wyoming Natural Diversity Database, University of Wyoming, Laramie.
- Fertig, W. and B. Heidel. 2003. Wyoming plant element ranking guidelines- working document. Wyoming Natural Diversity Database, University of Wyoming, Laramie.
- Keinath, D.K. and G.P. Beauvais. 2003. Wyoming animal element ranking guidelines. Wyoming Natural Diversity Database, University of Wyoming, Laramie.
- USDA Forest Service. 1991. Threatened, Endangered, and Sensitive Species of the Intermountain Region. USDA Forest Service, Region 4, Ogden, Utah.
- USDA Forest Service. 1994. Forest Service Manual, Title 2600 - wildlife, fish and sensitive plant habitat management; Region 2 supplement 2600-94-2, Section 2672.11a, Exhibit 1. USDA Forest Service, Region 2, Lakewood, Colorado.
- USDA Forest Service. 2003. R2 Regional Forester's Sensitive Species: information memorandum 3 November 2003. USDA Forest Service, Region 2, Lakewood, Colorado.

- USDI Bureau of Land Management. 2001. Wyoming Bureau of Land Management Sensitive Species policy and list: instruction memorandum no. WY-2001-040. USDI Bureau of Land Management Wyoming State Office, Cheyenne.
- USDI Bureau of Land Management. 2002. Wyoming Bureau of Land Management update of Sensitive Species list 2002: information bulletin no. WY-2003 0001. USDI Bureau of Land Management Wyoming State Office, Cheyenne.

GENERAL REFERENCES-WY PLANTS

- Dorn, R.D. 2001. Vascular plants of Wyoming. 3rd edition. Mountain West Publishing, Cheyenne, Wyoming.
- Fertig, W., C. Refsdal, and J. Whipple. 1994. Wyoming rare plant field guide. Wyoming Rare Plant Technical Committee, Cheyenne.
- Rocky Mountain Herbarium www.rmh.uwyo.edu
- NatureServe Explorer: www.natureserve.org/explorer
- Wyoming Natural Diversity Database: www.uwyo.edu/WYNDD

GENERAL REFERENCES-WY ANIMALS

- Clark, T.W. and M. Stromberg. 1987. Mammals in Wyoming. University of Kansas Museum of Natural History, Lawrence.
- Dorn, R.D. and J.L. Dorn. 1999. Wyoming Birds. 2nd edition. Mountain West Publishing, Cheyenne, Wyoming
- Baxter, G.T and M.D. Stone. 1995. Fishes of Wyoming. Wyoming Game and Fish Department, Cheyenne.
- Baxter, G.T. and M.D. Stone. 1985. Amphibians and Reptiles of Wyoming. Wyoming Game and Fish Department, Cheyenne.
- NatureServe Explorer: www.natureserve.org/explorer/
- Wyoming Natural Diversity Database: www.uwyo.edu/WYNDD

Plant Species of Concern

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|-------------------------------|---|---------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Abies concolor | G5/S1 | | SWE, UIN? | Rock Springs BLM | Widespread/ | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| White fir | | | , | Wasatch-Cache NF? | Edge | | | | | | |
| Abronia ammophila Yellowstone sand verbena | G1/S1 | | PAR, TET | Yellowstone NP | Local Endemic/ Core | Very Low | Rare | Large Decline | Moderate Decline? | High | Very High |
| Achnatherum nevadense | G4/S1 | | CAR, FRE | Lander BLM | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| [Stipa nevadensis] Nevada needlegrass | | | , | Rawlins BLM Rock Springs BLM | 3 | | | | | | C |
| Achnatherum scribneri [Stipa scribneri] Scribner needlegrass | G4/S1 | | LAR | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Achnatherum swallenii [Oryzopsis swallenii] Swallen mountain -ricegrass | G4/S2 | | LIN, SUB | Kemmerer BLM Pinedale BLM Rock Springs BLM | Regional Endemic/ Edge | Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Adiantum aleuticum [Adiantum pedatum] Aleutian maidenhair | G5?/ S1 | | TET | Grand Teton NP Yellowstone NP | Disjunct | Very Low | Uncommon | Unknown | Stable? | Low | High |
| Adoxa moschatellina Moschatel | G5/ S2 | | ALB, CRO, PAR, SHE, WAS | Bighorn NF Black Hills NF Medicine Bow NF? Shoshone NF Yellowstone NP | Disjunct | Low | Rare | Unknown | Stable? | High | High |
| Agalinis tenuifolia var. parviflora Slender false-foxglove | G5T4/ S1 | | GOS, NIO, PLA | Ft. Laramie NHS | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |
| Agrostis mertensii [Agrostis borealis] Northern bentgrass | G5/S1 | | ALB, PAR | Medicine Bow NF Shoshone NF Yellowstone NP | Disjunct | Very Low | Unknown | Unknown | Stable? | Moderate | High |
| Agrostis oregonensis Oregon bentgrass | G4/ S1? | | PAR, TET | Bridger-Teton NF? Grand Teton NP? Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Agrostis rossiae Ross bentgrass | G1/S1 | | TET | Yellowstone NP | Local Endemic/ Core | Very Low | Rare | Moderate Decline | Moderate Decline? | High | Very High |
| Aletes humilis Larimer aletes | G2G3 / SH | | ALB | | Regional Endemic/ Edge | Very Low | Unknown | Unknown | Unknown | High | High |
| Amerorchis rotundifolia [Orchis rotundifolia] Round-leaved orchid | G5/S1 | USFS R2 Sensitive | PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Moderate Decline? | High | High |
| Ammannia robusta Robust toothcup | G5/S1 | | WES | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Androsace chamaejasme ssp. carinata [A. lehmanniana] Sweet-flowered rock-jasmine | G5T4/ S1S2 | USFS R4 Sensitive | FRE, HOT, PAR, TET | Cody BLM? Shoshone NF Targhee NF Yellowstone NP? | Widespread/ Edge | Low | Uncommon | Unknown | Stable | Moderate | Low |
| Androstephium breviflorum Purple funnel-lily | G5/S1 | | CAR, SWE | Rawlins BLM | Widespread/ Edge | Very Low | Uncommon | Unknown | Unknown | Moderate | Low |
| Anemone lyallii | G5/S1 | | TET | Targhee NF | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Lyall's wood anemone | | | | | | | | | | ? | |
| Anemone narcissiflora ssp. zephyra Zephyr windflower | G5T4/ S1 | | BIG, JOH, SHE | Bighorn NF | Regional Endemic/ Edge | Very Low | Uncommon | Stable? | Stable | Moderate | Medium |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|---|-------------------------------|---|------------------------------|-----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Antennaria arcuata Meadow pussytoes | G2/S2 | USFS R4 Sensitive WY BLM Sensitive | FRE, SUB | Lander BLM Pinedale BLM Rock Springs BLM | Regional Endemic/ Core | Moderate | Uncommon | Moderate Decline | Stable | Moderate | Very High |
| Antennaria flagellaris Stoloniferous pussytoes | G5?/ S1S2 | | PAR, TET | Shoshone NF Yellowstone NP | Disjunct | Low | Unknown | Unknown | Unknown | Moderate | High |
| Antennaria monocephala Single-head pussytoes | G4G5 / S1 | | FRE, JOH, PAR, SUB, TET | Bighorn NF Bridger-Teton NF Shoshone NF Targhee NF | Widespread/Ed ge | Low | Rare | Stable? | Stable | Moderate | Low |
| Aquilegia brevistyla Small-flower columbine | G5/S1 | USFS R2 Sensitive | PAR?, WES | Black Hills NF Newcastle BLM Shoshone NF? | Disjunct | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Medium |
| Aquilegia formosa var. formosa Crimson columbine | G5T5/ SH | | PAR? | Shoshone NF? Yellowstone NP? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Aquilegia laramiensis Laramie columbine | G2/S2 | USFS R2 Sensitive WY BLM Sensitive | ALB, CON | Medicine Bow NF Rawlins BLM? | Local Endemic/ Core | Moderate | Rare | Stable? | Stable? | High | Very High |
| Arceuthobium douglasii Douglas-fir dwarf-mistletoe | G5/S1 | | LIN, TET | Bridger-Teton NF Targhee NF | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | High | Low |
| Arctous rubra [Arctostaphylos rubra] Red manzanita | G5/S1 | | PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Argyrochosma fendleri [Notholaena fendleri] Fendler's false cloak-fern | G3/S1 | | LAR | | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Aristida curtissii [Aristida basiramea var. curtissii] Curtis' three-awn | G5T5/ S1 | | CRO?, WES | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Aristida oligantha Prairie three-awn | G5/S1 | | WES | Thunder Basin NG | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Arnica angustifolia var. tomentosa [Arnica alpina var. tomentosa] Alpine arnica | G5T5/ S1 | | FRE | Shoshone NF | Widespread/ Edge | Very Low | Rare | Stable? | Stable | Moderate | Low |
| Arnica lonchophylla Northern arnica | G4/S2 | | BIG, JOH, SHE | Bighorn NF Buffalo BLM | Disjunct | Moderate | Uncommon | Stable? | Stable? | Moderate | High |
| Artemisia biennis var. diffusa Mystery wormwood | G5T1 Q/ S1 | WY BLM Sensitive | SWE | | Regional Endemic/ core? | Very Low | Rare | Unknown | Large Decline | High | Very High |
| Artemisia porteri Porter's sage brush | G2/S2 | WY BLM Sensitive | FRE, JOH, NAT | Buffalo BLM Casper BLM Lander BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | High | Very High |
| Asclepias arenaria Sand milkweed | G5/S1 | | GOS | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High | Low |
| Asclepias engelmanniana Engelmann's milkweed | G5/S1 | | GOS | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|--|---|------------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Asclepias hallii Hall's milkweed | G3/ SH | | ALB | Rawlins BLM? | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | High |
| Asclepias subverticillata Bedstraw milkweed | G4G5 / SH | | CAR | Rawlins BLM? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Low | Low |
| Asclepias uncialis Dwarf milkweed | G3?/ SH | USFS R2 Sensitive | SWE? | Rock Springs BLM? Seedskadee NWR? | Disjunct | Very Low | Unknown | Moderate Decline? | Unknown | High | High |
| Asclepias verticillata Whorled milkweed | G5/S1 | | CRO | Devils Tower NM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Aspidotis densa [Cryptogramma densa] Pod-fem | G5/S1 | | TET | Grand Teton NP | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | High | Low |
| Asplenium trichomanes Maidenhair spleenwort | G5/S1 | | ALB, NAT | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | High | Low |
| Asplenium trichomanes-ramosum [Asplenium viride] Green spleenwort | G4/S2 | | CAR, SHE, SUB, TET, WAS | Bighorn NF Bridger-Teton NF Medicine Bow NF Targhee NF | Disjunct | Low | Rare | Unknown | Stable? | Moderate | High |
| Astragalus barrii [Orophaca barrii] Barr's milkvetch | G3/S3 | USFS R2 Sensitive | CAM, CON, JOH, NAT, NIO, SHE, WES | Buffalo BLM Casper BLM Newcastle BLM Thunder Basin NG | Regional Endemic/ Core | Moderate | Uncommon | Moderat e Decline | Moderate Decline? | Moderate | High |
| Astragalus bisulcatus var. haydenianus Hayden's milkvetch | G5T4 ?/ S1? | | CAR, FRE?, LIN?, SWE, UIN | Lander BLM Kemmerer BLM Rawlins BLM Rock Springs BLM | Widespread/ Edge | Low? | Unknown | Unknown | Unknown | Moderate | Low |
| Astragalus calycosus var. calycosus King's milkvetch | G5T4 ?/ S1S2 | | LIN, SWE, UIN | Kemmerer BLM | Widespread/ Edge | Low | Rare | Unknown | Stable? | Moderate | Low |
| Astragalus coltonii var. moabensis Moab milkvetch | G4T3 % S2 | | SWE, UIN | Kemmerer BLM Rock Springs BLM | Regional Endemic/ Edge | Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Astragalus diversifolius Meadow milkvetch | G3/ SH | USFS R4 Sensitive | SWE? | Rock Springs BLM? Seedskadee NWR? | Regional Endemic/ Edge | Unknown | Unknown | Large Decline? | Unknown | High | High |
| Astragalus gilviflorus var. purpureus [Orophaca triphylla var. purpurea] Dubois milkvetch | G5T2/ S2 | WY BLM Sensitive | FRE | Lander BLM Shoshone NF Wind River IR | Local Endemic/ Core | Low | Uncommon | Unknown | Stable | High | Very High |
| Astragalus jejunus var. articulatus Hyattville milkvetch | G3T1/ S1 | WY BLM Sensitive | BIG | Worland BLM | Local Endemic/ Core | Very Low | Rare | Unknown | Moderate Decline? | High | Very High |
| Astragalus lentiginosus var. salinus Sodaville milkvetch | G5T5/ S1 | | LIN, UIN | Fossil Butte NM Kemmerer BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | High | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|--|-----------------------|---|------------------------------|-----------|-----------|----------------------|---------------------|---------------------------------|------------------------------------|
| Astragalus leptaleus Park milkvetch | G4/ SH | USFS R2 Sensitive Targhee NF Sensitive | CAR | Medicine Bow NF | Regional Endemic/Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Medium |
| Astragalus paysonii Payson's milkvetch | G3/S2 | USFS R4 Sensitive | LIN, SUB, TET | Bridger-Teton NF Targhee NF | Regional Endemic/ Core | Moderate | Uncommon | Moderate Decline | Stable? | High | High |
| Astragalus platytropis Broad-keeled milkvetch | G5/S1 | | PAR | Cody BLM | Disjunct | Very Low | Unknown | Unknown | Unknown | High? | High |
| Astragalus proimanthus [Orophaca proimantha] Precocious milkvetch | G1/S1 | WY BLM Sensitive | SWE | Rock Springs BLM | Local Endemic/ Core | Very Low | Rare | Moderate Decline? | Moderate Decline | High | Very High |
| Astragalus racemosus var. longisetus Long-stalk ed racemose milkvetch | G5T4/ S1 | | NIO | Newcastle BLM? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Astragalus racemosus var. racemosus Racemose milkvetch | G5T5/ S1 | | NIO, WES | Newcastle BLM Thunder Basin NG | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Astragalus racemosus var. treleasei Trelease's racemose milkvetch | G5T2/ S2 | WY BLM Sensitive | SUB, UIN | Kemmerer BLM Pinedale BLM | Regional Endemic/ Core | Low | Rare | Moderate Decline | Unknown | Moderate | Very High |
| Astragalus robbinsii var. minor Robbins' milkvetch | G5T5/ S1 | | LIN, SUB, TET | Bridger-Teton NF | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Astragalus terminalis Railhead milkvetch | G3/S1 | | TET | Bridger-Teton NF Grand Teton NP National Elk Refuge | Regional Endemic/ Edge | Very Low | Uncommon | Unknown | Stable? | Moderate | High |
| Athyrium distentifolium var. americanum [Athyrium alpestre var. americanum] American alpine lady fern | G4G5 T4T5/ S1 | | ALB, TET | Grand Teton NP Medicine Bow NF Targhee NF Yellowstone NP | Widespread/ Edge | Low | Rare | Moderate Decline? | Stable? | High | Low |
| Atriplex falcata [Atriplex gardneri var.falcata] Sickle saltbush | G4Q/ S1 | | SUB, SWE, UIN | Kemmerer BLM Pinedale BLM Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Atriplex wolfii [Atriplex tenuissima] Wolf's orache | G3G4 / S1 | | CAR, SWE | Rawlins BLM Rock Springs BLM | Regional Endemic/Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Medium |
| Bacopa rotundifolia Roundleaf water-hyssop | G5/S1 | | CRO, LAR | | Widespread/ Edge | Very Low | Unknown | Unknown | Increase? | Moderate | Low |
| Bahia dissecta Dissected bahia | G5/ S1S2 | | ALB, CAR, CON, LAR | Casper BLM? Curt Gowdy SP Medicine Bow NF Rawlins BLM? | Widespread/ Edge | Low | Rare | Unknown | Stable? | Moderate | Low |
| Besseya alpina Alpine kittentails | G4/S1 | | ALB, CAR | Medicine Bow NF | Regional Endemic/ Edge | Very Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Besseya plantaginea White River kittentails | G? /S1 | | ALB | Medicine Bow NF | Regional Endemic/ Edge | Very Low | Uncommon | Moderate Decline? | Stable | Moderate | Medium |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|-----------------------|--|------------------------------|-----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Boechera crandallii [Arabis crandallii] Crandall's rockcress | G2/S1 | | CAR, SWE | Ashley NF [FGNRA] Rawlins BLM Rock Springs BLM | Regional Endemic/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Very High |
| Boechera perennans [Arabis perennans] Perennial rockcress | G5/S1 | | ALB | | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |
| Boechera pusilla [Arabis pusilla] Small rockcress | G1/S1 | WY BLM Sensitive | FRE | Rock Springs BLM | Local Endemic/ Core | Very Low | Rare | Unknown | Stable | High | Very High |
| Boechera selbyi [Arabis selbyi] Selby's rockcress | G4?Q/ S1 | | SWE | Ashley NF [FGNRA] Rawlins BLM Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Botrychium ascendens Upward-lobe moonwort | G2G3 / S1 | USFS R4 Sensitive | FRE, JOH. PAR | Bighorn NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | High | High |
| Botrychium campestre Prairie moonwort | G3G4 /S1 | USFS R2 Sensitive | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High | High |
| Botrychium lanceolatum var lanceolatum Lance-leaved moonwort | G5T4 /S1 | | CAR, FRE, JOH | Bighorn NF Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High | Low |
| Botrychium minganense [B. lunaria var. onondagense] Mingan moonwort | G4/S1 | | FRE, JOH, PAR, SHE | Bighorn NF Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High | Low |
| Botrychium multifidum Leathery grape-fern | G5/S2 | USFS R2 Sensitive | CAR, SUB, TET | Bridger-Teton NF Grand Teton NP Medicine Bow NF Yellowstone NP | Widespread/ Edge | Low | Rare | Unknown | Unknown | High | Low |
| Botrychium virginianum Rattlesnake fern | G5/S2 | | CRO, PAR, SHE, TET | Bighorn NF Black Hills NF Grand Teton NP Shoshone NF Targhee NF Yellowstone NP | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Bouteloua simplex Mat grama | G5/S1 | | LAR | FE Warren AFB | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Braya glabella Arctic braya | G5/S1 | USFS R2 Sensitive | SUB, TET | Bridger-Teton NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Braya humilis Low braya | G5/S1 | | FRE | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Brickellia microphylla var. scabra Little-leaved brickell-bush | G4G5 T4?/ S1 | | FRE, NAT, SWE | Ashley NF (FGNRA) Casper BLM Lander BLM Rock Springs BLM | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Bromus pubescens Hairy wood brome | G5/ SH | | CON | | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Campanula aparinoides Marsh bellflower | G5/S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Carex alopecoidea Foxtail sedge | G5/S2 | USFS R2 Sensitive | CRO | Black Hills NF | Disjunct | Low | Rare | Moderate Decline? | Unknown | Moderate | Medium |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|-------------------------------|--|---------------------------|--------------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Carex concinna Beautiful sedge | G4G5 / S1 | | PAR, SUB | Black Hills NF Bridger-Teton NF Shoshone NF Yellowstone NP | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Carex crawei Crawe's sedge | G5/S1 | | GOS, LAR | FE Warren AFB | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High? | Low |
| Carex cusickii Cusick's sedge | G5/ S2 | | PAR, TET | Grand Teton NP Yellowstone NP | Widespread/ Edge | Low | Uncommon | Unknown | Stable? | Moderate | Low |
| Carex deweyana var. bolanderi [Carex bolanderi] Bolander's sedge | G5T5/ S1 | | LIN, PAR, TET | Bridger-Teton NF Grand Teton NP Shoshone NF Targhee NF Yellowstone NP | Widespread/ Edge | Very Low? | Unknown | Unknown | Unknown | Moderate | Low |
| Carex diandra Lesser panicled sedge | G5/ S1S2 | USFS R2 Sensitive | ALB, PAR, TET | Grand Teton NP Medicine Bow NF Shoshone NF Yellowstone NP | Disjunct | Low | Rare | Unknown | Unknown | Moderate | High |
| Carex eburnea Ebony sedge | G5/S1 | | CRO | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Carex echinata ssp. echinata [Carex muricata] Little prickly sedge | G5/S1 | | PAR?, TET | Grand Teton NP Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Carex egglestonii Eggleston's sedge | G4/S1 | | ALB, CAR | Medicine Bow NF | Regional Endemic/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Medium |
| Carex emoryi Emory's sedge | G5/S1 | | CRO, PLA? | Devils Tower NM | Widespread/ Edge | Very Low | Uncommon | Unknown | Unknown | Moderate | Low |
| Carex flava Yellow sedge | G5/S1 | | PAR, TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Carex granularis var. haleana Meadow sedge | G5T4/ S2 | | CRO | Black Hills NF | Disjunct | Low | Unknown | Unknown | Unknown | Moderate | High |
| Carex incurviformis var. danaensis [Carex maritima] Incurved sedge | G4G5 T3/ S2 | USFS R4 Sensitive | FRE, PAR, SUB | Bridger-Teton NF Shoshone NF | Disjunct | Low | Rare | Unknown | Stable? | Moderate | Medium |
| Carex laeviculmis Smooth-stemmed sedge | G5/S1 | | TET | Grand Teton NP JDR Parkway | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Carex leptalea Bristly -stalk sedge | G5/S2 | USFS R2 Sensitive | ALB, CRO, PAR, TET | Black Hills NF Grand Teton NP Medicine Bow NF Shoshone NF Targhee NF Yellowstone NP | Disjunct | Low | Rare | Unknown | Stable? | Moderate | High |
| Carex limosa Mud sedge | G5/S2 | | BIG, CAR, PAR, SUB, TET | Bighorn NF Bridger-Teton NF Medicine Bow NF Shoshone NF Targhee NF Yellowstone NP | Disjunct | Low | Uncommon | Unknown | Stable? | Moderate | High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|------------------------------------|--|---------------------------|-----------|-----------|----------------------|---------------------|---------------------------------|------------------------------------|
| Carex livida Livid sedge | G5/S2 | USFS R2 Sensitive | PAR, TET | Shoshone NF Yellowstone NP | Disjunct | Low | Uncommon | Unknown | Unknown | High | High |
| Carex luzulina var. atropurpurea Black and purple sedge | G5T3/ S2 | USFS R4 Sensitive | FRE, SUB, TET | Bridger-Teton NF Shoshone NF | Regional Endemic/ Core | Low | Rare? | Unknown | Unknown | Moderate | High |
| Carex microglochin False uncinia sedge | G5?/ S1 | | FRE, PAR, SUB | Bridger-Teton NF Shoshone NF Yellowstone NP | Disjunct | Very Low | Uncommon | Unknown | Unknown | High | High |
| Carex misandra [Carex fuliginosa var. misandra] Short-leaf sedge | G5/S1 | | BIG, FRE, JOH, PAR | Bighorn NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Carex nelsonii Nelson's sedge | G3?/ S2 | | ALB, CAR, FRE, PAR | Medicine Bow NF Shoshone NF | Regional Endemic/ Core | Low | Uncommon | Moderate Decline? | Unknown | Moderate | High |
| Carex occidentalis Western sedge | G4/S1 | | ALB, FRE?, PAR, TET? | Cody BLM Medicine Bow NF Rawlins BLM? Shoshone NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Carex oreocharis Mountain -loving sedge | G3/S1 | | ALB, LAR | Curt Gowdy SP Medicine Bow NF | Regional Endemic/Edge | Very Low | Uncommon | Moderate Decline? | Stable? | Moderate | High |
| Carex parryana var. unica [Carex parryana ssp. hallii; C. hallii] Hall's sedge | G4?Q/ S1 | | ALB, CAR, LAR, PLA | Medicine Bow NF Rawlins BLM | Widespread/ Edge | Very Low | Uncommon | Moderate Decline | Unknown | Moderate | Low |
| Carex proposita Smoky Mountain sedge | G4/ SH | | TET | Bridger-Teton NF? Grand Teton NP? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Carex richardsonii Richardson sedge | G4/S1 | | CRO | Black Hills NF | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Carex rosea Rosy sedge | G5/S1 | | CRO | Black Hills NF | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Carex sartwellii var. sartwellii Sartwell's sedge | G4G5 T4T5/ S1 | | ALB, FRE, JOH, PAR, SHE, TET | Grand Teton NP Medicine Bow NF National Elk Refuge Wind River IR? Yellowstone NP | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |
| Carex scirpoidea var. scirpiformis Canadian single-spike sedge | G5T4 Q/ S1 | | PAR, TET | Grand Teton NP Nation Elk Refuge Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High | Low |
| Carex stricta Strict sedge | G5/S1 | | NIO | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Ceanothus herbaceus Prairie redroot | G5/S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Moderate Decline | Moderate | Low |
| Ceanothus martinii Utah mountain lilac | G4/S1 | | LIN, SWE | Fossil Butte NM Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Moderat e Decline? | Moderate | Low |
| Celtis occidentalis Common hackberry | G5/S1 | | GOS, PLA, SHE | Bighorn NF Casper BLM? | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Centunculus minimus Chaffweed | G5/S1 | | CRO, PLA | Black Hills NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cercocarpus ledifolius var. intricatus [Cercocarpus intricatus] Dwarf mountain mahogany | G5T4/ S1 | | SWE | Ashley NF (FGNRA) Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---------------------|-----------------------|--|---------------------------|-----------|-----------|--------------------|----------------------|---------------------------------|------------------------------------|
| Chamaechaenactis scaposa Fullstem | G4/ S1S2 | | SWE | Kemmerer BLM Rock Springs BLM | Regional Endemic/ Edge | Low? | Rare | Unknown | Unknown | Moderate | Medium |
| Chamaesyce geyeri [Euphorbia geyeri] Geyer's spurge | G5/S1 | | GOS | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Chamaesyce serpens [Euphorbia serpens] Matted broom-spurge | G5/S1 | | CRO | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Low | Low |
| Chenopodium incanum var. incanum Mealy goosefoot | G5T?/ S1 | | GOS | | Widespread/ Edge | Very Low | Uncommon | Unknown | Unknown | Low | Low |
| Chenopodium watsonii Watson goosefoot | G5/S1 | | GOS, LAR, WES | Bighorn NF? Buffalo BLM Casper BLM Worland BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Chionophila jamesii Rocky Mountain snowlover | G4?/ S1 | | ALB, CAR | Medicine Bow NF | Regional Endemic/Edge | Very Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Chrysothamnus greenei Greene rabbitbrush | G5/ S1? | | SWE | Rawlins BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cicuta bulbifera Bulb-bearing water-hemlock | G5/S1 | | TET | Targhee NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Circaea lutetiana var. canadensis Canadian enchanter's nightshade | G5T5/ S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Cirsium aridum Cedar Rim thistle | G2Q/ S2 | WY BLM Sensitive | CAR, FRE, SUB, SWE | Lander BLM Pinedale BLM Rawlins BLM Rock Springs BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Moderate Decline? | Moderate | Very High |
| Cirsium canovirens Gray-green thistle | G4G5 / SH | | PAR | Yellowstone NP? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cirsium foliosum Leafy thistle | G5/S1 | | PAR, SHE | Bighorn NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cirsium ownbeyi Ownbey's thistle | G3/S2 | WY BLM Sensitive | SWE | Ashley NF (FGNRA) Rock Springs BLM | Regional Endemic/Edge | Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Clarkia pulchella Large-flower clarkia | G5?/ SH | | TET | Bridger-Teton NF? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Cleome multicaulis Many-stemmed spider-flower | G2G3 / S1 | WY BLM Sensitive | NAT | Casper BLM Pathfinder NWR | Disjunct | Very Low | Uncommon | Moderate Decline? | Stable? | High | High |
| Collomia grandiflora Large-flower collomia | G5/ SH | | LIN, SWE? | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Crassula solieri [Crassula aquatica, Tillaea aquatica - misapplied] Py gmyweed | G4G5 / S1 | | FRE, PAR | Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cryptantha gracilis Slender cryptantha | G5/S1 | | SWE | Ashley NF (FGNRA) Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Cryptantha rollinsii Rollins' cat's-eye | G3/S1 | | SWE | Ashley NF? (FGNRA?) Rock Springs BLM Seedskadee NWR | Regional Endemic/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | #Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|-------------------------------|---|----------------------------|----------|-----------|----------------------|---------------------|---------------------------------|------------------------------------|
| Cryptantha subcapitata Owl Creek miner's candle | G2/S2 | WY BLM Sensitive | FRE | Boysen St. Park Lander BLM Wind River IR | Local Endemic/ Core | Very Low | Uncommon | Unknown | Stable? | High | Very High |
| Cryptogramma stelleri Fragile rockbrake | G5/S1 | | FRE, PAR, SHE, TET | Bighorn NF Bridger-Teton NF Shoshone NF Targhee NF Yellowstone NP | Disjunct | Low | Rare | Unknown | Unknown | Moderate | High |
| Cuscuta indecora var. neuropetala Pretty dodder | G5T5/ S1 | | GOS, PLA, SHE | Casper BLM? Glendo St. Park | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cuscuta megalocarpa [Cuscuta umbrosa] Big-fruited dodder | G5/ SH | | PLA | | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Cuscuta occidentalis Western dodder | G5/S1 | | LIN, SWE | Fossil Butte NM Rock Springs BLM? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cuscuta plattensis Wyoming dodder | G1?Q/ S1 | | CON, GOS, PLA | Casper BLM Glendo St. Park | Regional Endemic/ Core? | Very Low | Unknown | Large Decline? | Unknown | Moderate ? | Very High |
| Cymopterus alpinus [Oreoxis alpina] Alpine oreoxis | G4/S1 | | ALB | Medicine Bow NF Rawlins BLM? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Cymopterus evertii Evert's waferparsnip | G2G3 / S2S3 | WY BLM Sensitive | HOT, PAR | Cody BLM Shoshone NF | Local Endemic / Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Cymopterus williamsii Williams' waferparsnip | G2G3 / S2S3 | WY BLM Sensitive | BIG, JOH, NAT, WAS | Bighorn NF? Buffalo BLM Casper BLM Worland BLM | Local Endemic/ Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Cyperus acuminatus Short -point flatsedge | G5/S1 | | CAM, GOS | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cyperus bipartitus [Cyperus rivularis] Shining flatsedge | G5/S1 | | GOS, PLA | Guernsey St. Park Fort Laramie NHS | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cyperus erythrorhizos Red-root flatsedge | G5/S1 | | CRO, PLA | Keyhole St. Park Greyrocks WMA | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Cypripedium fasciculatum Clustered lady-slipper | G4/ S2S3 | USFS R4 Sensitive | ALB, CAR | Medicine Bow NF | Disjunct | Moderate | Uncommon | Moderate Decline? | Stable? | Moderate | High |
| Cypripedium montanum Mountain lady-slipper | G4G5 / S1 | USFS R2 Sensitive | JOH, SHE | Bighorn NF | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Moderate Decline | High | Low |
| Cypripedium parviflorum var. pubescens [Cypripedium calceolus var. pubescens] Large yellow lady-slipper | G5T?/ S1S2 | USFS R2 Sensitive | BIG, CRO, FRE, SHE, WAS | Bighorn NF Black Hills NF? Wind River IR | Disjunct | Low | Rare | Moderate Decline? | Moderate Decline | High | High |
| Dalea aurea Golden prairie-clover | G5/S1 | | GOS, PLA | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Dalea cylindriceps Andean prairie-clover | G3?/ S1 | | ALB, GOS, PLA | Casper BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Dalea enneandra Nine-anther prairie-clover | G5/S1 | | CRO, WES | Newcastle BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---|--------------------------------|--|----------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Dalea villosa var. villosa Silky prairie-clover | G5T5/ S1 | | GOS, LAR | Casper BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Delphinium ramosum Mountain larkspur | G4/S1 | | LAR | | Regional Endemic/ | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Deschampsia danthonioides Annual hairgrass | G5/S1 | | PAR | Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Moderate Decline? | Moderate | Low |
| Descurainia pinnata var. paysonii [Descurainia incisa ssp. paysonii] Payson's tansymustard | G5T3 % S2 | | CAR, SWE | Ashley NF (FGNRA) Rawlins BLM Rock Springs BLM | Widespread/ Edge | Low | Uncommon | Unknown | Unknown | High | Medium |
| Descurainia torulosa Wyoming tansymustard | G1/S1 | USFS R2 Sensitive USFS R4 Sensitive WY BLM Sensitive | FRE, PAR, SWE, TET | Bridger-Teton NF Cody BLM? Rock Springs BLM Shoshone NF | Local Endemic/ Core | Low | Rare | Unknown | Unknown | High | Very High |
| Dichanthelium linearifolium [Panicum linearifolium] Slim-leaf witchgrass | G5/S1 | | ALB, CRO, WES | Black Hills NF Medicine Bow NF | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Diphasiastrum complanatum [Lycopodium complanatum] Ground cedar | G5/S1 | USFS R2 Sensitive | CRO | Black Hills NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Dodecatheon jeffreyi Jeffrey's shootingstar | G5/S1 | | TET | Grand Teton NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Unknown | Low |
| Downingia laeta Great Basin downingia | G5/S1 | | ALB, CAR, UIN | Kemmerer BLM? Rawlins BLM | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Draba borealis Boreal draba | G4/S2 | Bridger- Teton NF Sensitive | LIN, PAR, SUB, TET | Bridger-Teton NF Grand Teton NP National Elk Refuge Shoshone NF Yellowstone NP | Disjunct | Low | Rare | Moderate Decline? | Stable? | Moderate | High |
| Draba fladnizensis var. pattersonii White arctic whitlow-grass | G4T2 T3/ S2 | | FRE, JOH, LIN, PAR, TET? | Bighorn NF Bridger Teton NF Grand Teton NP Shoshone NF Targhee NF | Regional Endemic/Edge | Very Low | Unknown | Unknown | Stable? | Moderate | Medium |
| Draba glabella Rock whitlow-grass | G4G5 / SH | | PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Draba juniperina [Draba oligosperma var. juniperina] Uinta draba | G3Q/ S2 | | SWE, UIN | Ashley NF (FGNRA) Rock Springs BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable | Moderate | High |
| Draba paysonii var. paysonii Payson's draba | G5T3/ S2 | | FRE, LIN, PAR, SUB | Bridger-Teton NF Shoshone NF Yellowstone NP? | Regional Endemic/ Core? | Low | Rare | Unknown | Unknown | Moderate | High |
| Draba paysonii var. treleasii Trelease's draba | G5T4 ?/ S1 | | FRE, PAR | Shoshone NF Yellowstone NP? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|-----------------------|--|---------------------------|-----------|-----------|----------------------|-------------------|---------------------------------|------------------------------------|
| Draba pectinipila [Draba oligosperma var. pectinipila] Comb-hair whitlow-grass | G1Q/ S1 | | PAR | Shoshone NF | Regional Endemic/ Core | Very Low | Rare | Unknown | Stable? | High | Very High |
| Draba porsildii [Includes vars. brevicula and porsildii] Porsild's whit low-grass | G3G4 / S1 | | FRE, HOT, PAR, SUB | Bridger-Teton NF Shoshone NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| <i>Draba spectabilis</i> var. <i>oxyloba</i> Showy draba | G3?T 3Q/ SH | | CAR | Medicine Bow NF | Regional Endemic/ Edge | Very Low | Unknown | Unknown | Large Decline? | Moderate | High |
| <i>Drosera anglica</i> English sundew | G5/S2 | USFS R2 Sensitive | PAR, TET | Grand Teton NP JDR Parkway Shoshone NF Yellowstone NP | Disjunct | Low | Rare | Moderate Decline? | Unknown | High | High |
| Dryopteris expansa [Dryopteris assimilis; Dryopteris carthusiana] Spreading woodfern | G5/S1 | | PAR, TET | Grand Teton NP Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Dulichium arundinaceum Three-way sedge | G5/S1 | | TET | Yellowstone NP | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Eleocharis bella Delicate spikerush | G5/S1 | | TET | Grand Teton NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Eleocharis flavescensvar. thermalis Warm Springs spikerush | G5T2 T3Q/ S2 | | PAR, TET | JDR Parkway Grand Teton NP Yellowstone NP | Disjunct | Low | Rare | Moderate Decline? | Stable | Moderate | High |
| Eleocharis ovata Ovate spike-rush | G5/S1 | | CRO | Black Hills NF Grand Teton NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Eleocharis parvula var. anachaeta Small spike-rush | G5/S1 | | GOS | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Eleocharis tenuis var. borealis [E. elliptica] Boreal spikerush | G5T5/ S1 | USFS R2 Sensitive | PLA, TET | Medicine Bow NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Elymus simplex var. luxurians [Leymus simplex] Long-awned alkali wild-rye | G4?Q T?/ S1 | | SWE | Rock Springs BLM | Local Endemic / Core | Very Low | Rare | Unknown | Unknown | Moderate | Very High |
| Elymus triticoides [Leymus triticoides] Creeping wildrye | G4G5 / SH | | ALB | Hutton Lake NWR | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Elymus villosus Hairy wild-rye | G5/S1 | | CRO | Black Hills NF Devils Tower NM Newcastle BLM | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Ephedra viridis var. viridis Green Mormon tea | G5T5/ S1 | | SWE | Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Epipactis gigantea Giant helleborine | G4/S1 | USFS R2 Sensitive | BIG, TET | Grand Teton NP Yellowstone NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Moderate Decline? | High | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | #Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|------------------------------|--|---------------------------|----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Equisetum fluviatile Water horsetail | G5/S1 | | PAR, TET | Grand Teton NP Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Large Decline? | Unknown | Moderate | Low |
| Equisetum scirpoides Dwarf scouring rush | G5/S1 | | CRO | Black Hills NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Equisetum sylvaticum Woodland horsetail | G5/S1 | | CRO, SHE | Bighorn NF Black Hills NF | Disjunct | Very Low | Rare | Unknown | Stable | Moderate | High |
| Eragrostis hypnoides Teal love grass | G5/S1 | | GOS, SHE | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Eriastrum wilcoxii [Eriastrum sparsiflorum var. wilcoxii] Wilcox eriastrum | G5/ S1S2 | | FRE, SWE | Lander BLM Rock Springs BLM Seedskadee NWR | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Erigeron compactus var. consimilis [Erigeron consimilis] San Rafael daisy | G4G5 / S1 | | SWE | Rawlins BLM Rock Springs BLM | Regional Endemic/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Erigeron elatior Tall fleabane | G4/S2 | | ALB, CAR | Medicine Bow NF Rawlins BLM? | Regional Endemic/ Edge | Low? | Uncommon | Moderate Decline? | Unknown | Moderate | Medium |
| Erigeron humilis Low fleabane | G4/S2 | | BIG, PAR, SUB | Bighorn NF Bridger-Teton NF Shoshone NF | Widespread/ Edge | Low | Rare | Unknown | Unknown | High | Low |
| Erigeron lanatus Woolly fleabane | G3G4 / S1 | USFS R4 Sensitive | SUB | Bridger-Teton NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Erigeron pinnatisectus Pinnate fleabane | G4/S2 | | ALB, CAR | Medicine Bow NF | Regional Endemic/Edge | Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Erigeron radicatus Taprooted fleabane | G3/S2 | | FRE, PAR, SUB, TET | Bridger-Teton NF Shoshone NF | Regional Endemic/ Edge | Low | Rare | Unknown | Unknown | Moderate | Medium |
| Eriogonum corymbosum var. corymbosum Crisp-leaf wild buckwheat | G5T5/ S1 | | SWE | Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Eriogonum divaricatum Divergent wild buckwheat | G4G5 / S1 | | LIN, SUB, SWE, UIN | Kemmerer BLM Pinedale BLM Rock Springs BLM | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |
| Eriogonum exilifolium Slender-leaved buckwheat | G3/S2 | USFS R2 Sensitive | ALB, CAR | Medicine Bow NF Rawlins BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Unknown | Moderate | High |
| Eriogonum hookeri Hooker wild buckwheat | G5/S1 | | CAR, SWE | Rawlins BLM? Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Eriogonum mancum Mancos wild buckwheat | G4/S1 | | BIG | Cody BLM | Regional Endemic/Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Medium |
| Eriogonum pauciflorum var. nebraskense Nebraska buckwheat | G5T? Q/S1 | | PLA, GOS | | Regional Endemic/Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Medium |
| Eriophorum callitrix var. callitrix Sheathed cotton-grass | G5T5/ S1 | | FRE, PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Eriophorum chamissonis Russet cotton-grass | G5/ S2 | USFS R2 Sensitive | PAR, SHE, TET | Bighorn NF Shoshone NF Yellowstone NP | Widespread/ Edge | Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|-------------------|---|---------------------------|-----------|-----------|-----------------------|---------------------|---------------------------------|------------------------------------|
| Eriophorum gracile Slender cotton-grass | G5/S2 | USFS R2 Sensitive | ALB, PAR, TET | Bridger-Teton NF? Grand Teton NP Medicine Bow NF Shoshone NF Targhee NF Yellowstone NP | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Eriophorum scheuchzeri Scheuchzer cotton-grass | G5/S1 | | FRE, PAR, SUB | Bridger-Teton NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Eriophorum viridicarinatum Green keeled cotton-grass | G5/ S1S2 | | PAR, TET | Grand Teton NP National Elk Refuge Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Eritrichum howardii Howard forget-me-not | G4/ S2? | | JOH, PAR, SHE | Bighorn NF Buffalo BLM Cody BLM Shoshone NF | Regional Endemic/Edge | Low | Uncommon | Unknown | Unknown | Moderate | Medium |
| Euphorbia exstipulata var. exstipulata Square-seeded spurge | G5T5/ SH | | PLA | Casper BLM? | Disjunct | Very Low | Unknown | Large Declin e? | Unknown | Moderate | High |
| Euphorbia hexagona Six-angle spurge | G5/S1 | | GOS, PLA | Casper BLM Guernsey SP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Eustoma grandiflorum [Eustoma russellianum] Showy prairie-gentian | G5/S1 | | GOS, NAT, PLA | | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | High | Low |
| Euthamia graminifolia var. major [Solidago graminifolia var. major] Flat-top fragrant goldenrod | G5T5/ S1 | | ALB, PLA | Casper BLM Medicine Bow NF | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Festuca hallii [Festuca altaica ssp. hallii; included in F. scabrella by some authors] Hall's fescue | G4/S2 | USFS R2 Sensitive | ALB, JOH?, PAR | Bighorn NF? Cody BLM Medicine Bow NF Shoshone NF | Widespread/ Edge | Low | Uncommon | Moderat e Decline? | Unknown | Moderate | Low |
| Filago prolifera [Evax prolifera] Bighead pygmycudweed | G5/S1 | | PLA | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Froelichia gracilis Slender cottonweed | G5/ SH | | WES | | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Galium coloradoense [Galium multiflorum var. coloradoense] Colorado bedstraw | G4/S1 | | SWE | Rawlins BLM Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Gaura neomexicana ssp. coloradensis Colorado butterfly plant | G3T2/ S2 | USFWS Threaten. | LAR, PLA | FE Warren AFB | Regional Endemic/ Core | Low | Uncommon | Large Decline? | Moderate Decline | High | Very High |
| Gentiana affinis var. bigelovii Bigelow's prairie gentian | G5T4/ S1 | | ALB, LAR | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Gentianella propinqua Four-part gentian | G5/S1 | | PAR | Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | #Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---|------------------|---|---------------------------|--------------|-----------|--------------------|----------------------|---------------------------------|------------------------------------|
| Gentianopsis simplex Hiker's gentian | G4/S1 | | TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Glandularia bipinnatifida Dakota vervain | G5/S1 | | CRO, FRE | Devils Tower NM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Glossopetalon spinescens var. meionandrum [Forsellesia meionandra] Utah greasebush | G5T3/ S1 | | SWE | Ashley NF (FGNRA) Rock Springs BLM | Regional Endemic/ Edge | Very Low | Rare | Unknown | Stable | Moderate | Medium |
| Gymnocarpium dryopteris [incl. Gymnocarpium disjunctum & G. x brittonianum] Oak fern | G5/S2 | | CAR, CRO, TET | Black Hills NF Grand Teton NP Medicine Bow NF | Widespread/Ed ge | Low | Rare | Unknown | Moderate Decline? | Moderate | Low |
| Helianthemum bicknellii [Crocanthemum bicknellii] Plains frostweed | G5/S1 | | CRO | Black Hills NF Devils Tower NM Newcastle BLM | Widespread/ Edge | Very Low | Rare | Unknown | Stable | Moderate | Low |
| Helictotrichon mortonianum Alpine oatgrass | G4/S1 | | FRE | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Hesperochiron californicus California hesperochiron | G4G5 / S1 | | SUB, UIN | Pinedale BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Hesperostipa neomexica na [Stipa neomexicana] New Mexico needlegrass | G4G5 / S1 | | PLA | Casper BLM Guernsey St. Park | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Heterocodon rariflorus Western pearl-flower | G5/ SH | | TET | Targhee NF | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Heterotheca pumila Little golden-aster | G4/S1 | | ALB, CAR | Medicine Bow NF | Regional Endemic/Edge | Very Low? | Rare? | Unknown | Unknown | Moderate | Medium |
| Horkelia fusca var. parviflora Pine woods horkelia | G5T?/ SH | | TET | Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Huperzia haleakalae [Lycopodium selago, Huperzia selago - misapplied] Fir clubmoss | G5/ SH | | TET | Grand Teton NP | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Hymenopappus tenuifolius Chalk -hill woollywhite | G5/ SH | | CRO | Black Hills NF? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Ionactis alpina [Aster scopulorum] Lava aster | G5/S1 | | PAR | Yellowstone NP | Widespread/ Edge | Very Low | Uncommon | Unknown | Stable? | Moderate | Low |
| Ipomopsis aggregata var. tenuituba [Ipomopsis tenuituba] Slender-trumpet ipomopsis | G5T4 T5/S1 | | ALB, CAR | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Ipomopsis aggregata var. weberi Weber's scarlet -gilia | G5T2/ S1 | USFS R2 Sensitive WY BLM Sensitive | CAR | Medicine Bow NF Rawlins BLM | Regional Endemic/Edge | Very Low | Rare | Unknown | Large Decline? | Moderate | High |
| Ipomopsis longiflora [Gilia longiflora] Longflowered gilia | G5/S1 | | GOS | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|-------------------------|---|----------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Ipomopsis polycladon [Gilia polycladon] Lavender ipomopsis | G4/S1 | | SWE | Ashley NF (FGNRA) | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Ipomopsis spicata var. robruthii [Ipomopsis spicata var. robruthiorum] Kirkpatrick's ipomopsis | G4?T 2/ S2 | | HOT, PAR | Shoshone NF | Local Endemic/ Core | Low | Rare | Unknown | Stable? | Moderate | Very High |
| Isoetes occidentalis [Isoetes lacustris] Western quillwort | G4G5 / S1 | | PAR, TET | Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Juncus filiformis Thread rush | G5/S1 | | CAR, PAR, TET | Grand Teton NP Medicine Bow NF Targhee NF Yellowstone NP | Widespread/ Edge | Low? | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Juncus triglumis var. albescens [Juncus albescens] Northern white rush | G5T5/ S1 | | ALB, FRE, PAR, SUB | Bridger-Teton NF Medicine Bow NF Shoshone NF | Disjunct | Very Low | Rare | Moderate Decline? | Stable? | Moderate | High |
| Juncus triglumis var. triglumis Three-flower rush | G5T5/ S1 | | JOH, PAR | Bighorn NF Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Juncus vaseyi Vasey rush | G5?/ S1 | | ALB, FRE, SUB | Bridger-Teton NF Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Kelloggia galioides Milk kelloggia | G5/S1 | | FRE?, LIN?, PAR, TET | Bridger-Teton NF Grand Teton NP Shoshone NF Yellowstone NP | Widespread/ Edge | Low? | Rare | Unknown | Unknown | Moderate | Low |
| Kobresia sibirica [Kobresia macrocarpa; Kobresia schoenoides] Siberian kobresia | G5/S1 | | PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Kobresia simpliciuscula Simple kobresia | G5/S1 | USFS R2 Sensitive | FRE?, PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Koenigia islandica Koenigia | G4/S1 | | FRE, PAR | Shoshone NF | Disjunct | Low? | Rare | Unknown | Unknown | Moderate | High |
| Lathyrus eucosmus Seemly sweetpea | G4G5 / S1 | | FRE | Lander BLM | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Lathyrus lanszwertii var. lanszwertii Nevada sweetpea | G4G5 T4/ S1 | | FRE?, HOT, UIN | Kemmerer BLM Wasatch-Cache NF Worland BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Lechea intermedia Narrowleaf pinweed | G5/S1 | | CRO | Black Hills NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| <i>Lemna valdiviana</i> Pale duckweed | G5/S1 | | TET | Grand Teton NP Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Lepidium integrifolium var. integrifolium Entire-leaved peppergrass | G1G2 T1?/ S1 | WY BLM Sensitive | LIN, UIN | Fossil Butte NM Kemmerer BLM | Regional Endemic/ Core? | Very Low | Rare | Large Decline? | Moderate Decline? | High | Very High |
| Lepidium montanum var. alyssoides [Lepidium alyssoides var. a.] Mountain peppergrass | G5T5 % S1 | | SUB | Pinedale BLM | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---|-----------------------|--|------------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Lesquerella arenosa var. argillosa Sidesaddle bladderpod | G5T3/ S1 | WY BLM Sensitive | NIO | Newcastle BLM | Regional Endemic/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Lesquerella carinata var. carinata Keeled bladderpod | G3G4 T3T4/ S1 | | FRE?, TET | Bridger-Teton NF Grand Teton NP National Elk Refuge Targhee NF | Regional Endemic/Edge | Low | Uncommon | Unknown | Stable | Moderate | Medium |
| Lesquerella fremontii Fremont bladderpod | G2/S2 | USFS R2 Sensitive WY BLM Sensitive | FRE | Lander BLM Shoshone NF | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Lesquerella macrocarpa Large-fruited bladderpod | G2/S2 | WY BLM Sensitive | FRE, LIN, SUB, SWE | Kemmerer BLM Pinedale BLM Rock Springs BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Moderate Decline? | High | Very High |
| Lesquerella multiceps Western bladderpod | G3/S1 | WY BLM Sensitive | LIN | Kemmerer BLM? Targhee NF | Regional Endemic/Edge | Very Low | Rare | Moderate Decline? | Stable? | Moderate | Medium |
| Lesquerella parvula [Lesquerella alpina var. parvula] Narrowleaved bladderpod | G5T3 ?/ S1 | | CAR, SWE, UIN | Kemmerer BLM Medicine Bow NF? | Regional Endemic/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Medium |
| Lesquerella prostrata Prostrate bladderpod | G3/S1 | WY BLM Sensitive | LIN, UIN | Kemmerer BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | High |
| Liatris lancifolia Lance-leaf gay-feather | G4/S1 | | GOS | | Widespread/ Edge | Very Low | Rare | Unknown | Moderate Decline | Moderate | Low |
| Ligularia bigelovii var. hallii [Senecio bigelovii var. hallii] Bigelow's groundsel | G4?T 3T4/ S1 | | ALB, CAR | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Ligusticum tenuifolium Slender-leaved lovage | G5/S1 ? | | CAR | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Linanthus watsonii [Leptodactylon watsonii] Watson's prickly-phlox | G3G5 / S1 | | FRE, HOT, SWE, WAS | Ashley NF (FGNRA) Bighorn NF Cody BLM Wind River IR | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Lipocarpha drummondii [Hemicarpha drummondii] Dwarf bulrush | G4G5 / S1 | | ALB, GOS, PLA, WES | Rawlins BLM? Rawhide WMA | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Listera convallarioides Broad-leaved twayblade | G5/S1 | | ALB, CON, SHE, TET | Bighorn NF Grand Teton NP Medicine Bow NF Rawlins BLM Targhee NF Yellowstone NP | Widespread/ Edge | Low | Rare | Moderate Decline? | Unknown | High | Low |
| Lithospermum multiflorum Many-flowered gromwell | G4/ SH | | LAR | | Disjunct | Very Low | Unknown | Large Decline? | Unknown | Moderate | High |
| Lobelia siphilitica Great blue lobelia | G5/S1 | | GOS, NIO | Ft. Laramie NHS | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Loeflingia squarrosa var. artemisiarum Sage-like loeflingia | G5T2 T3/SH | | SWE | Rock Springs BLM? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Medium |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|------------------|---|-------------------------------|------------|-----------|----------------------|-------------------|---------------------------------|------------------------------------|
| Loeflingia squarrosa var. texana Texas spreading loeflingia | G5T4 ?/ S1 | | CRO, WES | Newcastle BLM? | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Lomatium attenuatum Absaroka biscuitroot | G3/S2 | | PAR | Cody BLM Shoshone NF | Regional Endemic/ Core | Low | Uncommon | Unknown | Moderate Decline? | Moderate | Very High |
| Lomatium triternatum var. anomalum Ternate desert-parsley | G5T?/ S1 | | LIN | Fossil Butte NM Kemmerer BLM | Regional Endemic/ Edge? | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Lomatogonium rotatum Marsh felwort | G5/ S2 | | ALB, CAR, LAR | Medicine Bow NF | Disjunct | Low | Rare | Moderate Decline? | Unknown | Moderate | High |
| Luzula glabrata var. hitchcockii [Luzula hitchcockii] Smooth wood-rush | G5T4/ S1 | | PAR, TET | Bridger-Teton NF Grand Teton NP Shoshone NF Targhee NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Lycopodium dendroideum | G5/ | | CRO | Black Hills NF | Disjunct | Low | Uncommon | Unknown | Unknown | Moderate | High |
| Treelike clubmoss | S1S2 | | CDO TET | Dia ala Hella NIE | W: 11/ | Manual and | T T1 | I Indonesia | I I1 | Madausta | T |
| Lycopus uniflorus Northern bugleweed | G5/S1 | | CRO, TET | Black Hills NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Lysimachia thyrsiflora Swamp loosestrife | G5/S1 | | ALB, TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | High | Low |
| Lythrum alatum var. alatum Winged loosestrife | G5/S1 | | CRO, NIO, PLA | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Machaeranthera bigelovii var. bigelovii Bigelow's tansy -aster | G4G5 T? /S1 | | ALB | Medicine Bow NF | Regional Endemic/Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Medium |
| Machaeranthera coloradoensis var. coloradoensis [Haplopappus coloradoensis] Colorado tansy-aster | G2T2 ?/ S2 | USFS R2 Sensitive | ALB, CAR | Medicine Bow NF Rawlins BLM | Regional Endemic/ Core? | Low | Rare | Moderate Decline? | Unknown | Moderate | High |
| Marsilea vestita var. oligospora [Marsilea oligospora] Pepperwort | G5/S1 | | SUB, TET | Bridger-Teton NF? Grand Teton NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Melica smithii Smith melic grass | G4/ SH | | TET | Grand Tet on NP? Targhee NF? Yellowstone NP? | Widespread/ Edge | Very Low | Rare | Large Decline? | Unknown | Moderate | Low |
| Mentzelia oligosperma Few-seed stickleaf | G4/S1 | | CRO, PLA, WES | Casper BLM Guernsey St. Park | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Mentzelia rusbyi | G4?/ | | ALB, CAR | Medicine Bow NF | Widespread/ | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Rusby's stickleaf | S1 | | | Rawlins BLM? | Edge | | | | | | |
| Mentzelia speciosa Jeweled blazingstar | G3?/ S1 | | LAR | | Regional Endemic/Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Mimulus nanus | G5/S1 | | TET | Yellowstone NP | Widespread/ | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Dwarf purple monkeyflower | 03/51 | | 121 | 1 Showstone 1 41 | Edge | , cry Low | Ture | JIKIOWII | Subic: | Moderate | Low |
| Mimulus rubellu s Ciliolate-toothed monkeyflower | G5/S1 | | CAR, NAT, PAR | Casper BLM Rawlins BLM Yellowstone NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Stable | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|-------------------|--------------------------------|---|---------------------------|-----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Minuartia filiorum [Arenaria filiorum; A. rubella var. filiorum] Thread-branch stitchwort | G3G4 / S1 | | SUB, TET | Bridger-Teton NF | Regional Endemic/Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Monarda pectinata Plains lemon bee-balm | G5/S1 | | GOS | Casper BLM | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Monardella odoratissimavar. glauca Mountain wild-mint | G4G5 T?/S1 | | LIN | Bridger-Teton NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Monolepis pusilla Red poverty-weed | G5/S1 | | SWE, WAS | Rock Springs BLM Worland BLM? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Montiastrum lineare Linearleaf miner's-lettuce | G5/S1 | | PAR, TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Muhlenbergia glomerata [incl. in Muhlenbergia racemosa by some authors] Marsh muhly | G5/ S2 | | CRO, GOS?, PAR, SUB, TET | Black Hills NF National Elk Refuge Pinedale BLM? Shoshone NF Yellowstone NP | Widespread/ Edge | Low | Rare | Moderate Decline? | Stable? | Moderate | Low |
| Muhlenbergia montana Mountain muhly | G5/S1 | | CAR, LAR | Curt Gowdy SP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Muhlenbergia torreyi Ring muhly | G4/S1 | | LAR | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Musineon vaginatum Sheathed musineon | G3?/ S2 | | BIG, SHE | Bighorn NF Cody BLM | Regional Endemic/ Edge | Low | Rare | Unknown | Unknown | Moderate | Medium |
| Myosotis verna Spring forget-me-not | G5/S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Myriophyllum quitense Andean water-milfoil | G4?/ S1 | | TET | Yellowstone NP | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Myriophyllum verticillatum Whorled water-milfoil | G5/S1 | | PAR, TET | Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Najas guadalupensis Southern naiad | G5/S1 | | SUB?, TET | Bridger-Teton NF? Grand Teton NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Nothocalais troximoides False agoseris | G5/S1 | | PAR, SHE | Cody BLM Grand Teton NP Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Oenothera canescens Spotted evening-primrose | G4G5 / S1 | | GOS | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Oenothera howardii Howard's evening primrose | G3G4 / S1 | | LAR | | Regional Endemic/Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Onoclea sensibilis Sensitive fern | G5/S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Ophioglossum pusillum [Ophioglossum vulgatum] Adder's-tongue | G5/S1 | | TET | Yellowstone NP | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Opuntia erinacea var. utahensis Utah old man prickly -pear | G5T4/ S1? | | SWE | Rock Springs BLM? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|---------------------------------|--|-------------------------------|-----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Opuntia polyacantha var. juniperina Juniper prickly -pear | G5T3 ?Q/ S1 | | SUB, SWE | Ashley NF (FGNRA) Kemmerer BLM Pinedale BLM Seedskadee NWR | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Opuntia polyacantha var. rufispina Rufous-spine prickly-pear | G5T5/ S2 | | LIN, SWE | Ashley NF (FGNRA) Kemmerer BLM Rawlins BLM Rock Springs BLM | Widespread/ Edge | Low | Unknown | Unknown | Unknown | Moderate | Low |
| Orobanche corymbosa var. corymbosa [Orobanche californica var. corymbosa] Flat-top broomrape | G4T4/ S1 | | PAR, SUB, TET | Bridger-Teton NF Grand Teton NP Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Orobanche ludoviciana var. arenosa Louisiana broomrape | G5T5/ SH | | TET | Grand Teton NP Yellowstone NP? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Oxytheca dendroidea Tree-like oxytheca | G4/ SH | | FRE, HOT, SUB?, SWE | Lander BLM? Rock Springs BLM Worland BLM? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Oxytropis besseyi var. obnapiformis [Oxytropis obnapiformis] Maybell locoweed | G5T2/ S1 | | FRE, NAT, SWE, UIN | Lander BLM Kemmerer BLM | Regional Endemic/ Edge? | Low | Rare | Unknown | Unknown | Moderate | High |
| Packera crocata [Senecio crocatus] Saffron groundsel | G3/S1 ? | | ALB, SWE | Medicine Bow NF Rock Springs BLM | Regional Endemic/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | High |
| Packera indecora [Senecio indecorus] Plains ragwort | G5/S1 | | PAR | Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Packera pseudaurea var. flavula [Senecio pseudaureus var. flavulus] Streambank groundsel | G5T?/ S1 | | ALB, CAR, LAR | FE Warren AFB Medicine Bow NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Paeonia brownii Brown's peony | G5/S1 | | TET | Grand Teton NP Targhee NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Palafoxia rosea var. macrolepis Rosy palafoxia | G5T3 T4/S1 | | CON, GOS | Thunder Basin NG | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Papaver kluanense [Papaver lapponicum var. occidentale P. radicatum ssp. kluanensis] Alpine poppy | G3G4 / S2 | | BIG, FRE?, JOH?, PAR, SUB | Bighorn NF Bridger-Teton NF Cody BLM Shoshone NF | Disjunct | Low | Rare | Unknown | Stable? | Moderate | High |
| Parnassia kotzebuei Kotzebuei's grass-of-parnassus | G4/S2 | USFS R2 Sensitive | JOH, PAR, SUB, TET | Bighorn NF Bridger-Teton NF Grand Teton NP Shoshone NF | Widespread/ Edge | Low | Unknown | Unknown | Stable? | Moderate | Low |
| Paronychia jamesii James nailwort | G4/S1 | | ALB, GOS | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|---|-----------------------|--|----------------------------------|--------------|-----------|----------------------|---------------------|---------------------------------|------------------------------------|
| Paronychia pulvinata Rocky Mountain nailwort | G3/S1 | | ALB, CAR | Medicine Bow NF Rawlins BLM | Regional Endemic/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Parrya nudicaulis Naked-stemmed parrya | G5/S2 | USFS R4 Sensitive | FRE, PAR, SUB, TET | Bridger-Teton NF Shoshone NF | Disjunct | Low | Uncommon | Unknown | Stable | Moderate | High |
| Pectis angustifolia Crown-seed fetid-marigold | G4G5 / S1 | | CON, GOS, PLA | Casper BLM? Thunder Basin NG | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Pedicularis contorta var. ctenophora Coil-beaked lousewort | G5T3/ S2 | | BIG, JOH, SHE | Bighorn NF Buffalo BLM | Regional Endemic/ Edge | Low | Uncommon | Unknown | Unknown | Moderate | Medium |
| Pedicularis oederi Oeder's lousewort | G5/ S2 | | PAR | Shoshone NF | Disjunct | Very Low | Uncommon | Unknown | Stable? | Moderate | High |
| Pedicularis parryi ssp. mogollonica Mogollon lousewort | G5T2 T4Q/ S1 | | BIG?, UIN | Bighorn NF? Wasatch-Cache NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Pediomelum digitatum [Psoralea digitata] Palm-leaved scurfpea | G5/S1 | | GOS | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Pediomelum linearifolium [Psoralea linearifolia] Narrowleaf scurfpea | G4?/ S1 | | LAR | | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Pellaea atropurpurea [WY material may all be Pellaea gastonyi] Purple-stem cliff-brake | G5/S1 ? | | CRO | Black Hills NF Newcastle BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Pellaea gastonyi Gastony's cliff-brake | G2G4 / S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low? | Rare | Unknown | Unknown | Moderate | Low |
| Pellaea suksdorfiana [Pellaea glabella var. simplex] Smooth cliff-brake | G5T4 % S1 | | CRO, PLA, TET | Black Hills NF Targhee NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Penstemon absarokensis Absaroka beardtongue | G2/S2 | USFS R2 Sensitive WY BLM Sensitive | FRE, PAR | Cody BLM Shoshone NF Wind River IR | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Penstemon acaulis var. acaulis Stemless beardtongue | G3T2/ S1 | USFS R4 Sensitive WY BLM Sensitive | SWE | Ashley NF (FGNRA) Rock Springs BLM | Regional Endemic/ Core | Very Low | Rare | Unknown | Moderate Decline | High | Very High |
| Penstemon angustifolius var. caudatus Narrowleaf beardtongue | G5T4 % S1 | | GOS | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Penstemon caryi Cary's beardtongue | G3/S3 | USFS R2 Sensitive | BIG, SHE, WAS | Bighorn NF Cody BLM Worland BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | High |
| Penstemon cyathophorus North Park beardtongue | G3/ S1S2 | | CAR | Medicine Bow NF Rawlins BLM | Regional Endemic/Edge | Low | Rare | Unknown | Stable? | Moderate | Medium |
| Penstemon gibbensii Gibbens' beardtongue | G1/S1 | WY BLM Sensitive | CAR, SWE | Rawlins BLM | Regional Endemic/ Core | Very Low | Rare | Unknown | Moderate Decline? | High | Very High |
| Penstemon haydenii Blowout penstemon | G1/S1 | USFWS Endanger. | CAR | Rawlins BLM | Regional Endemic/Edge | Very Low | Rare | Unknown | Stable? | High | Very High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | #Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|-------------------------------|--|---|----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Penstemon laricifolius ssp. exilifolius White beardtongue | G4T2 Q/S2 | USFS R2 Sensitive | LAR, NAT, NIO, PAR, SUB | Medicine Bow NF, Rawlins BLM | Regional Endemic/ Core | Low | Rare | Stable? | Unknown | Moderate | High |
| Penstemon scariosusvar. garrettii Garrett's beardtongue | G4T3/ S1 | | SWE, UIN | Rock Springs BLM | Regional Endemic/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Penstemon watsonii Watson beardtongue | G5/ SH | | UIN | Kemmerer BLM? | Regional Endemic/Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Medium |
| Phacelia alba [Phacelia neomexicana var. alba] White scorpion-weed | G4G5 / S1 | | ALB, LAR | Curt Gowdy SP Medicine Bow NF Rawlins BLM | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Phacelia demissa [includes vars. demissa & knightii] Intermountain phacelia | G5/S1 | | FRE, SWE | Rock Springs BLM | Regional Endemic/ Edge [If recognized, var. knightii is a local endemic/core] | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Phacelia denticulata Rocky Mountain phacelia | G3?/ S2 | | ALB, LAR | Curt Gowdy SP Medicine Bow NF Rawlins BLM | Regional Endemic/ Edge | Low | Rare | Unknown | Moderate Decline? | Moderate | Medium |
| Phacelia glandulosa var. deserta Desert glandular phacelia | G4T1 T2Q/ S1? | | LIN, SUB, SWE | Ashley NF (FGNRA) Kemmerer BLM Pinedale BLM Rawlins BLM Rock Springs BLM | Local Endemic/ Core | Low | Rare | Unknown | Unknown | Moderate | Very High |
| Phacelia incana Western phacelia | G3G4 / S1 | | CAR, SWE | Ashley NF (FGNRA) Rawlins BLM? Rock Springs BLM | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Phacelia salina Nelson phacelia | G3?Q/ S2 | | CAR, LIN, SUB, SWE, UIN | Pinedale BLM Rawlins BLM Rock Springs BLM | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Phacelia tetramera Tiny phacelia | G4/S1 | | FRE, SWE | Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Philadelphus microphyllus var. occidentalis Little-leaf mock-orange | G5?T 3T4/ S1 | | SWE | Ashley NF (FGNRA) Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Phippsia algida Ice grass | G5/S1 | | FRE?, PAR, SUB | Bridger-Teton NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Phlox albomarginata White-margined phlox | G4/S1 | | LIN | Kemmerer BLM | Regional Endemic/Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Medium |
| Phlox diffusa ssp. scleranthifolia [included in Phlox hoodii by some authors] Diffuse phlox | G5T?/ S1 | | PAR | Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Phlox pungens Beaver Rim phlox | G2/S2 | WY BLM Sensitive | FRE, LIN, SUB | Lander BLM Pinedale BLM Rock Springs BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Phryma leptostachya Lopseed | G5/S1 | | CRO | _ | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | #Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|---|---|---------------------------|----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Physalis virginiana var. virginiana Virginia ground-cherry | G5/S1 | | CRO, WES | Black Hills NF Newcastle BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Physaria condensata Tufted twinpod | G2/S2 | WY BLM Sensitive | LIN, SUB, UIN | Fossil Butte NM Kemmerer BLM Pinedale BLM Rock Springs BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Stable | Moderate | Very High |
| Physaria dornii Dorn's twinpod | G1/S1 | WY BLM Sensitive | LIN, UIN | Kemmerer BLM | Local Endemic/ Core | Very Low | Uncommon | Unknown | Stable | High | Very High |
| Physaria lanata [Physaria didymocarpa var. lanata] Woolly twinpod | G5T2/ S2 | USFS R2 Sensitive | BIG, CAM, HOT?, JOH, PAR, SHE, WAS | Bighorn NF Buffalo BLM Worland BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable | Moderate | Very High |
| Physaria saximontana var. saximontana Rocky Mountain twinpod | G3T2/ S2 | WY BLM Sensitive | CAR?, FRE, HOT, PAR | Lander BLM Shoshone NF Wind River IR Worland BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Physocarpus alternans Dwarf ninebark | G4/S1 | | SWE | Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Platanthera orbiculata [Habenaria orbiculata] Round-leaved orchid | G5?/ S1 | USFS R2 Sensitive | CRO | Black Hills NF | Disjunct | Very Low | Rare | Unknown | Unknown | High | High |
| Polemonium micranthum Annual polemonium | G5/ SH | | PAR | Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Polygala verticillata Whorled milkwort | G5/S1 | | CAM, CRO | Black Hills NF? Newcastle BLM Thunder Basin NG | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Polygonatum biflorum Common Solomon's-seal | G5/S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Polygonum spergulariiforme Fall knotweed | G5T4 % S1 | | JOH?, PAR, SHE, WAS | Worland BLM | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Polypodium saximontanum [Polypodium vulgare var. columbianum] Rocky Mountain polypody | G3?/ S1 | | ALB, CAR, CON | Medicine Bow NF | Regional Endemic/Edge | Very Low | Rare | Unknown | Stable? | Moderate | Medium |
| Polystichum scopulinum Rock holly -fern | G5/ SH | | TET | Targhee NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Populus deltoides var. wislizeni [Populus fremontii var. wislizeni] Fremont cottonwood | G5T4/ S1 | | CAR, SWE | Rawlins BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Stable | Moderate | Low |
| Porterella carnosula Western porterella | G4/S1 | | TET, UIN | Grand Teton NP Wasatch-Cache NF Yellowstone NP | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |
| Potamogeton amplifolius Large-leaved pondweed | G5/S1 | | CAR, FRE, JOH, PAR, SUB | Bighorn NF Bridger-Teton NF Medicine Bow NF Shoshone NF | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Potamogeton diversifolius Water-thread pondweed | G5/S1 | | CRO | Black Hills NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---|--------------------------------|--|------------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Potamogeton friesii Fries pondweed | G4/S1 | | CAR, PAR, TET | Bridger-Teton NF Grand Teton NP Medicine Bow NF | Widespread/Ed ge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Potamogeton illinoensis Illinois pondweed | G5/S1 | | ALB, NAT, PAR, SUB | Bridger-Teton NF Casper BLM? Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Potamogeton nodosus Longleaf pondweed | G5/S1 | | CAR, FRE, PLA, TET | Rawlins BLM? Wind River IR Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Potamogeton obtusifolius Blunt-leaf pondweed | G5/ SH | | TET | Grand Teton NP JDR Parkway | Disjunct | Very Low | Unknown | Large Decline? | Unknown | Moderate | High |
| Potamogeton praelongus White-stem pondweed | G5/S1 | | ALB, PAR | Medicine Bow NF Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Potamogeton robbinsii Flatleaf pondweed | G5/S1 | | CAR, FRE, SUB, TET | Bridger-Teton NF Medicine Bow NF Yellowstone NP | Disjunct | Very Low | Unknown | Unknown | Large Decline? | Moderate | High |
| Potamogeton strictifolius Strict -leaved pondweed | G5/ S1? | | ALB, SUB, SWE?,TET | Bridger-Teton NF Rawlins BLM Rock Springs BLM? Yellowstone NP | Disjunct | Very Low | Unknown | Unknown | Moderate Decline? | Moderate | High |
| Potamogeton zosteriformis Flatstem pondweed | G5/ SH | | TET | Grand Teton NP | Widespread/ Edge | Very Low | Unknown | Unknown | Moderate Decline? | Moderate | Low |
| Potentilla ambigens Giant cinquefoil | G3/ SH | | ALB | Rawlins BLM? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Medium |
| Potentilla hyparctica [Potentilla nana] Arctic cinquefoil | G4G5 / S1 | | FRE, SUB | Bridger-Teton NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Potentilla multisecta [Potentilla diversifolia var. multisecta] Deep Creek cinquefoil | G3G4 Q/ S1 | | SWE, UIN | Kemmerer BLM Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Potentilla subjuga Twinleaf cinquefoil | G4/S2 | | FRE, HOT, PAR | Bridger-Teton NF Cody BLM Shoshone NF Worland BLM | Regional Endemic/ Edge | Low | Rare | Unknown | Unknown | Moderate | Medium |
| Potentilla uniflora One-flower cinquefoil | G5/S2 | | FRE, HOT?, PAR, SUB, TET | Bridger-Teton NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Prenanthes racemosa ssp. multiflora Many-flowered rattlesnake-root | G5T4 % S1 | | ALB, CRO | Black Hills NF Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Primula egaliksensis Greenland primrose | G4/S1 | USFS R2 Sensitive Bridger- Teton NF Sensitive | PAR, SUB | Bridger-Teton NF Shoshone NF | Disjunct | Very Low | Rare | Unknown | Stable? | High | High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|-------------------------------|---|---------------------------|-----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Prosartes hookeri Drops of gold (Hooker's fairybell) | G5T4 T5/S2 | | CRO | Black Hills NF | Disjunct | Low | Rare | Unknown | Unknown | Moderate | High |
| Pseudognaphalium microcephalum var. thermale [Gnaphalium microcephalum var. thermale] White cudweed | G5T4 Q/ S1 | | PAR, TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Psilocarphus brevissimus Dwarf woolly-heads | G4/ S2 | | CAM, FRE, NAT, SUB | Casper BLM Lander BLM Pinedale BLM Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Puccinellia cusickii [Incl. in P. nuttalliana by some authors] Cusick's alkali-grass | G3G4 Q/ S1 | | ALB <u>.</u> CRO, JOH, NAT | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Pyrrocoma carthamoides var. subsquarrosa [Haplopappus carthamoides var. subsquarrosus] Absaroka goldenweed | G4G5 T2T3/ S2 | USFS R2 Sensitive | PAR | Cody BLM Shoshone NF | Regional Endemic/ Core | Low | Rare | Unknown | Unknown | Moderate | Very High |
| Pyrrocoma clementis var. clementis [Haplopappus clementis] Tranquil goldenweed | G3G4 T?/S1 | | CAR, FRE | Medicine Bow NF? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Pyrrocoma clementis var. villosa [Haplopappus clementis] Hairy tranquil goldenweed | G3G4 T?/S1 | USFS R2 Sensitive | BIG, FRE, JOH, SHE, WAS | Bighorn NF Wind River IR Worland BLM | Regional Endemic/ Core | Very Low | Rare | Moderate Decline? | Unknown | High | Very High |
| Pyrrocoma crocea [Haplopappus croceus] Western goldenweed | G4?/ S1 | | CAR | Medicine Bow NF Rawlins BLM | Widespread/ Edge | Very Low | Uncommon | Moderate Decline? | Unknown | Moderate | Low |
| Pyrrocoma integrifolia [Haplopappus integrifolius] Entire-leaved goldenweed | G3?/ S1 | USFS R2 Sensitive | BIG, FRE, PAR, WAS | Shoshone NF Worland BLM? Yellowstone NP | Regional Endemic/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Medium |
| Ranunculus flabellaris Yellow water-crowfoot | G5/ SH | | UIN | Kemmerer BLM? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Ranunculus gelidus [Ranunculus verecundus] Timberline buttercup | G5/S1 | | FRE, PAR, TET | Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Rayjacksonia annua [Haplopappus annuus] Viscid tansy -aster | G4G5 / S1 | | GOS, WES | Thunder Basin NG | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|--|--|---------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Rorippa calycina Persistent sepal yellowcress | G3/ S2S3 | WY BLM Sensitive | ALB, BIG, CAR, FRE, PAR, SWE, WAS | Ashley NF (FGNRA) Bighorn Canyon NRA Boysen St. Park Buffalo Bill St. Park Cody BLM Lander BLM Pathfinder NWR Rawlins BLM Seminoe St. Park Worland BLM | Regional Endemic/ Core | Moderate | Uncommon | Moderate Decline? | Increase? | Moderate | High |
| Rorippa truncata [Rorippa curvipes var. truncata] Wild yellowcress | G5/S1 | | FRE, GOS, NAT, PAR | Pathfinder NWR | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Rubus acaulis [Rubus arcticus ssp. acaulis] Northern blackberry [Nagoonberry] | G5/S1 | USFS R2 Sensitive | JOH, TET | Bighorn NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | High | Low |
| Sagittaria latifolia var. latifolia Broadleaf arrowhead | G5T5/ S1 | | GOS | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Salix barrattiana Barratt willow | G5/S1 | USFS R2 Sensitive | PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Stable | High | High |
| Salix candida Hoary willow | G5/S2 | USFS R2 Sensitive | ALB, PAR, SUB, TET | Bridger-Teton NF Medicine Bow NF Pinedale BLM? National Elk Refuge Shoshone NF Yellowstone NP | Widespread/ Edge | Low | Rare | Moderate Decline? | Moderate Decline? | High | Low |
| Salix myrtillifolia var. myrtillifolia Myrtleleaf willow | G5T5/ S1 | USFS R2 Sensitive | PAR | Shoshone NF | Disjunct | Very Low | Rare | Unknown | Unknown | High | High |
| Salix serissima Autumn willow | G4/S1 | USFS R2 Sensitive | ALB | Medicine Bow NF | Disjunct | Very Low | Rare | Unknown | Stable | Moderate | High |
| Sambucus cerulea Blue elderberry | G5?/ S1? | | JOH, SHE, SWE | Bighorn NF? Rock Springs BLM | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Sanicula graveolens Sierra sanicle | G4/ S1S2 | | PAR, TET | Shoshone NF Yellowstone NP | Disjunct | Low | Rare | Unknown | Stable? | Moderate | High |
| Saussurea weberi [Incl. in S. densa by some authors] Weber's saw-wort | G2G3 / S2 | USFS R4 Sensitive | FRE, SUB, TET | Bridger-Teton NF Shoshone NF | Regional Endemic/Edge | Low | Uncommon | Unknown | Stable? | Moderate | Medium |
| Saxifraga serpyllifolia var. chrysantha [Saxfraga chrysantha] Golden saxifrage | G4/S1 | | ALB, CAR, FRE, PAR, SUB | Bridger-Teton NF Medicine Bow NF Shoshone NF | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Scheuchzeria palustris Pod-grass | G5/S1 | | TET | Yellowstone NP | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | #Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|---|-----------------------|---|---------------------------|----------|-----------|--------------------|-------------------|---------------------------------|------------------------------------|
| Schoenoplectus americanus [Scirpus americanus] American bulrush | G5/S2 | | BIG, FRE, PAR, TET | Yellowstone NP JDR Parkway | Widespread/ Edge | Low | Unknown | Unknown | Stable? | Moderate | Low |
| Schoenoplectus heterochaetus [Scirpus heterochaetus] Slender bulrush | G5/S1 | | CAM, CRO, SHE | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Schoenoplectus saximontanus [Scirpus saximontanus] Rocky Mountain bulrush | G5/S1 | | GOS | Casper BLM? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Schoenoplectus subterminalis [Scirpus subterminalis] Water bulrush | G4G5 / S1 | | TET | Yellowstone NP | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Scirpus cyperinus Cottongrass bulrush | G5/S2 | | CRO, YEL | Black Hills NF Yellowstone NP | Widespread/Ed ge | Low | Uncommon | Unknown | Stable? | Moderate | Low |
| Scolochloa festucacea Sprangle-top | G5/S1 | | PAR | Yellowstone NP | Disjunct | Very Low | Rare? | Unknown | Moderate Decline? | Moderate | High |
| Selaginella mutica Blunt-leaf spike-moss | G4G5 / S1 | | CAR, LAR, SWE | Ashley NF (FGNRA) Curt Gowdy SP Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Selaginella rupestris Ledge spike-moss | G5/S1 | | CRO, WES | Black Hills NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Selaginella selaginoides Low spike-moss | G5/S1 | USFS R2 Sensitive | SUB, TET | Bridger-Teton NF Pinedale BLM Targhee NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Selaginella underwoodii Underwood's spike-moss | G5?/ S1 | | ALB, LAR, PLA | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Selaginella watsonii Watson's spike-moss | G4/S1 | | PAR | Shoshone NF | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Senecio hydrophiloides Sweet-marsh butterweed | G5/S1 | | TET | Grand Teton NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Senecio spartioides var. multicapitatus Many-headed broom groundsel | G5T4/ S1 | | CAR, SWE | Rawlins BLM Rock Springs BLM | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Sesuvium verrucosum Sea purslane | G5/ S1? | | CAM | Thunder Basin NG? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Shinnersoseris rostrata Annual skeleton-weed | G5?/ S1 | | CAR, GOS, LAR, PLA | Casper BLM Rawlins BLM | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Shoshonea pulvinata Shoshonea | G2G3 / S2 | USFS R2 Sensitive WY BLM Sensitive | FRE, HOT, PAR | Cody BLM Shoshone NF Wind River IR Worland BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Silene douglasii Douglas' campion | G4/S1 | | LIN | Kemmerer BLM | Widespread/ Edge | Low | Rare | Unknown | Unknown | Moderate | Low |
| Silene king ii "var. kingii" King's campion | G2G4 QT?/ S1 | | FRE, HOT, PAR | Cody BLM Shoshone NF | Disjunct | Low | Rare | Unknown | Unknown | Moderate | High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---|------------------------------------|--|---------------------------|-----------|-----------|----------------------|----------------------|---------------------------------|------------------------------------|
| Silene kingii var. novum new variety of King's campion | G2G4 QT?/ S1 | | PAR | Shoshone NF | Local Endemic/ Core | Very Low | Rare | Unknown | Unknown | Moderate | Very High |
| Silene repens var. australe Creeping campion | G5T3/ S1 | | SUB, TET | Bridger-Teton NF | Regional Endemic/Edge | Very Low | Rare | Unknown | Unknown | Moderate | Medium |
| Silphium integrifolium var. laeve Rosinweed | G4G5 T4?/ S1 | | LAR | FE Warren AFB | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Sisyrinchium idahoense var. idahoense Idaho blue-eyed grass | G5T4/ S1 | | PAR, TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Sorghastrum nutans Indian grass | G5/S1 | | CRO, GOS, PLA | Black Hills NF Fort Laramie NHS | Widespread/ Edge | Very Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Sparganium eurycarpum Large bur-reed | G5/S1 | | CRO, NIO, SHE | Black Hills NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Sparganium natans [Sparganium minimum] Small bur-reed | G5/S1 | | CAR, FRE, PAR, TET | Grand Teton NP Medicine Bow NF Shoshone NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Sphaeromeria simplex Laramie false sagebrush | G2/S2 | WY BLM Sensitive | ALB, CAR, CON, NAT | Rawlins BLM | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | High | Very High |
| Spiranthes diluvialis Ute ladies' tresses | G3/S1 | USFWS Threaten. | CON, GOS, LAR, NIO | Casper BLM | Widespread/ Edge | Very Low | Rare | Unknown | Moderate Decline? | High | High |
| Spirodela polyrrhiza Common water-flaxseed | G5/ SH | | TET | Grand Teton NP | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Sporobolus compositus [Sporobolus asper] Longleaf dropseed | G5T5/ S1 | | CAM, CRO, GOS | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Sporobolus heterolepis Northern dropseed | G5/S1 | | CRO, WES | | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Sporobolus neglectus Small dropseed | G5/ SH | | BIG, PAR | Cody BLM? | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Stellaria crispa Crimped stitchwort | G5/S1 | | PAR, TET | Grand Teton NP Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Stephanomeria fluminea Teton wire-lettuce | G2/S2 | | PAR, SUB, TET | Bridger-Teton NF Grand Teton NP Shoshone NF | Local Endemic/ Core | Low | Uncommon | Unknown | Unknown | Moderate | Very High |
| Stephanomeria pauciflora Prairie skeletonplant | G5/S1 | | PLA | Casper BLM | Widespread/ Edge | Low | Unknown | Unknown | Unknown | Moderate | Low |
| Symphyotrichum porteri [Aster porteri] Porter's aster | G3G4 / S1 | | ALB, CAR?, LAR | Curt Gowdy SP Medicine Bow NF | Regional Endemic/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Medium |
| Talinum parviflorum Small-flowered fame-flower | G5/S2 | | ALB, CAM, CON, LAR, PLA, WES | Buffalo BLM? Medicine Bow NF Thunder Basin NG? | Widespread/ Edge | Low | Unknown | Unknown | Unknown | Moderate | Low |
| Thelesperma caespitosum Green River greenthread | G1/S1 | USFS R4 Sensitive WY BLM Sensitive | SWE | Ashley NF (FGNRA) Rock Springs BLM | Regional Endemic/ Core | Very Low | Uncommon | Unknown | Moderate Decline? | High | Very High |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|---|-----------------------|---|------------------------|-----------|-----------|--------------------|-----------------------|---------------------------------|------------------------------------|
| Thelesperma pubescens Uinta greenthread | G1/S1 | USFS R4 Sensitive WY BLM Sensitive | SWE, UIN | Rock Springs BLM Wasatch-Cache NF | Local Endemic/ Core | Very Low | Uncommon | Unknown | Stable? | High | Very High |
| Tonestus pygmaeus [Haplopappus pygmaeus] Pygmy goldenweed | G4/S1 | | ALB, CAR | Medicine Bow NF | Widespread/ Edge | Very Low | Rare | Unknown | Stable? | Moderate | Low |
| Torreyochloa pallida var. fernaldii [Puccinellia fernaldii] Fernald alkali-grass | G5?T 4Q/ S1 | | TET | Grand Teton NP | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Townsendia condensata var. anomala [Townsendia anomala] North Fork Easter-daisy | G4T2/ S2 | USFS R2 Sensitive | PAR | Shoshone NF | Local Endemic/ Core | Low | Uncommon | Unknown | Stable? | High | Very High |
| Townsendia florifera [Townsendia florifer] Showy Easter-daisy | G5/ SH | | TET | Grand Teton NP? | Widespread/ Edge | Very Low | Unknown | Large Decline? | Unknown | Moderate | Low |
| Townsendia microcephala Cedar Mountain Easter-daisy | G1/S1 | WY BLM Sensitive | SWE, UIN | Rock Springs BLM | Local Endemic/ Core | Very Low | Rare | Unknown | Stable? | High | Very High |
| Tradescantia bracteata Long-bract spiderwort | G5/S1 | | CRO | Black Hills NF? | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Trautvetteria caroliniensis Carolina tassel-rue | G5/S1 | | TET | Yellowstone NP | Disjunct | Very Low | Rare | Unknown | Unknown | Moderate | High |
| Trichophorum pumilum [Scirpus rollandii, S. pumilus] Pygmy bulrush | G3Q/ S1 | | ALB, PAR, TET | Bridger-Teton NF Medicine Bow NF National Elk Refuge Shoshone NF | Disjunct | Very Low | Rare | Unknown | Stable? | Moderate | High |
| Trifolium barnebyi [Trifolium haydenii var.barnebyi] Barneby's clover | G1/S1 | WY BLM Sensitive | FRE | Lander BLM | Local Endemic/ Core | Very Low | Uncommon | Unknown | Stable? | High | Very High |
| Trillium ovatum Western trillium | G5/S2 | | CAR | Medicine Bow NF Rawlins BLM | Disjunct | Low | Rare | Unknown | Unknown | Moderate | High |
| Triodanis holzingeri Holzinger Venus' looking-glass | G4/S1 | | GOS, PLA | Casper BLM? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Triodanis leptocarpa Slim-pod Venus' looking glass | G5?/ S1 | | CAM, CRO, JOH, SHE | Bighorn NF? Black Hills NF | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Trisetum canescens Tall trisetum | G5/S1 | | TET | Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Triteleia grandiflora [<i>Brodiaea grandiflora</i>] Large flower triteleia | G4G5 / S2 | USFS R2 Sensitive | LIN, TET | Bridger-Teton NF? Caribou NF Grand Teton NP Targhee NF Yellowstone NP | Widespread/ Edge | Low | Rare | Unknown | Moderat e Decline? | Moderate | Low |
| Utricularia intermedia Flat-leaf bladderwort | G5/S1 | | TET | National Elk Refuge | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|----------------------|-----------------------|--|------------------------|-----------|-----------|----------------------|------------------|---------------------------------|------------------------------------|
| Utricularia minor Lesser bladderwort | G5/S2 | USFS R2 Sensitive | ALB, PAR, TET, WAS | Bighorn NF Grand Teton NP Medicine Bow NF Shoshone NF Yellowstone NP | Widespread/ Edge | Low | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Vaccinium myrtillus var. oreophilum Dwarf bilberry | G5T?/ S1 | | ALB, CAR, TET | Medicine Bow NF Yellowstone NP | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Viburnum edule Squashberry | G5/S1 | | ALB, CRO, PAR | Black Hills NF Medicine Bow NF Shoshone NF Yellowstone NP | Widespread/ Edge | Low? | Rare | Moderate Decline? | Unknown | Moderate | Low |
| Viburnum opulusvar. americanum Highbush cranberry | G5T5/ S1 | USFS R2 Sensitive | CRO | Black Hills NF | Disjunct | Low | Rare | Unknown | Unknown | Moderate | High |
| Viola orbiculata Western rough-leaved violet | G4/S1 | | LIN, FRE, TET | Grand Teton NP JDR Parkway Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Moderate Decline? | Unknown | Moderate | Low |
| Viola pedatifida Prairie violet | G5/S1 | | CRO | Black Hills NF Devils Tower NM Newcastle BLM? | Widespread/ Edge | Very Low | Rare | Unknown | Unknown | Moderate | Low |
| Viola renifolia var. brainerdii Kidney leaf white violet | G5T5/ S1 | | CRO, TET | Black Hills NF Bridger-Teton NF? Targhee NF | Disjunct | Very Low | Unknown | Unknown | Unknown | Moderate | High |
| Xerophyllum tenax Western beargrass | G4G5 / S1 | | TET | Grand Teton NP JDR Parkway Targhee NF Yellowstone NP | Widespread/ Edge | Very Low | Unknown | Unknown | Unknown | Moderate | Low |
| Yermo xanthocephalus Desert yellowhead | G1/S1 | USFWS Threaten. | FRE | Lander BLM | Local Endemic/ Core | Very Low | Rare | Unknown | Stable? | High | Very High |

Plant Species of Potential Concern (continued)

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|---|-----------------------|-------------------|--|---|------------------------------|-----------|-----------|---------------------|---------------------|---------------------------------|------------------------------------|
| Antennaria aromatica Aromatic pussytoes | G3G4 /S2 | | BIG, FRE, LIN, PAR, SUB, TET | Bighorn NF Bridger-Teton NF Cody BLM Grand Teton NP Shoshone NF Yellowstone NP | Regional Endemic/Edge | Moderate | Uncommon | Unknown | Stable? | Moderate | High |
| Astragalus drabelliformis Big Piney milkvetch | G2G3 / S2S3 | | LIN?, SUB | Bridger-Teton NF Pinedale BLM | Local Endemic/Core | Moderate | Abundant | Moderate Decline | Moderate Decline | Low | Very High |
| Astragalus nelsonianus [Astragalus pectinatus var. platyphyllus] Nelson's milkvetch | G3/S3 | | FRE, NAT, SWE | Ashley NF (FGNRA) Casper BLM Lander BLM Rawlins BLM Rock Springs BLM | Regional Endemic/ Core | Moderate | Uncommon | Stable? | Stable | Low | Very High |
| Astragalus shultziorum [Astragalus molybdenus, A. m. var. shultziorum] Shultz's milkvetch | G3Q/ S3 | | LIN, SUB, TET | Bridger-Teton NF Grand Teton NP Targhee NF | Local Endemic/Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Astragalus tridactylicus Three-fingered milkvetch | G4/S2 | | ALB, LAR, PLA? | Rawlins BLM? | Regional Endemic/Edge | Low | Uncommon | Unknown | Unknown | Moderate | High |
| Boechera pendulina var. russeola [Arabis pendulina var. russeola] Daggett rock cress | G5T3/ S3 | | ALB, CAR, FRE, NAT, SUB?, SWE, UIN? | Casper BLM Lander BLM Rawlins BLM Rock Springs BLM | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Boechera williamsii var. williamsii [Arabis williamsii var. williamsii] Williams' rock cress | G3QT 3/ S3 | | FRE, PAR, SUB | Bridger-Teton NF Rock Springs BLM Shoshone NF | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Unknown | Moderate | Very High |
| Castilleja crista -galli Cock's-comb paintbrush | G3/S2 | | PAR | Shoshone NF Yellowstone NP | Regional Endemic/Edge | Low | Unknown | Unknown | Stable? | Moderate | High |
| Castilleja nivea Snow paintbrush | G3/S2 | | FRE, HOT, PAR | Buffalo Bill St. Park Cody BLM Shoshone NF | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Castilleja nivea Snow paintbrush | G3/S2 | | FRE, HOT, PAR | Buffalo Bill St. Park Cody BLM Shoshone NF | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Cryptantha stricta Erect cryptantha | G3/S2 | | CAR, FRE, NAT, SWE, UIN | Ashley NF (BCNRA) Casper BLM Kemmerer BLM Lander BLM Rawlins BLM Rock Springs BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable | Moderate | Very High |
| Draba crassa Thick-leaf whitlow-grass | G3/S2 | | FRE, PAR, SUB, TET | Bridger-Teton NF Grand Teton NP Shoshone NF | Regional Endemic/ Edge | Low | Uncommon | Unknown | Stable? | Moderate | High |

Plant Species of Potential Concern (continued)

| Scientific Name Common Name | Heri- tage Rank | Federal Status | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic Trends | Recent Trends | Intrinsic Vulner- ability | Wyoming Contribu - tion Rank |
|--|-----------------------|----------------------|--|--|---------------------------|-----------|-----------|--------------------|------------------|---------------------------------|------------------------------------|
| Draba globosa [Draba apiculata var. apiculata; D. densifolia var. apiculata] Rockcress draba | G3/S2 | USFS R4 Sensitive | ALB, FRE, LIN, PAR, SUB, TET | Bridger-Teton NF Grand Teton NP Medicine Bow NF Shoshone NF Targhee NF | Regional Endemic/ Edge | Low | Uncommon | Unknown | Stable | Moderate | High |
| Ericameria discoidea var. linearis [Haplopappus macronema var. linearis; H. m. var. canescens] Narrowleaf goldenweed | G4G5 T3/ S2 | USFS R4 Sensitive | FRE, LIN, PAR, SUB, TET | Bridger-Teton NF Kemmerer BLM Shoshone NF Wind River IR Yellowstone NP | Regional Endemic/Edge | Moderate | Uncommon? | Unknown | Stable? | Moderate | High |
| Erigeron allocotus Bighorn fleabane | G3/S2 S3 | | BIG, SHE, WAS | Bighorn Canyon NRA Bighorn NF Cody BLM Worland BLM | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Stable | Moderate | Very High |
| Erigeron flabellifolius Fan-leaved fleabane | G3G4 /S2 | | HOT, PAR | Cody BLM Shoshone NF Yellowstone NP | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Eriogonum brevicaule var. canum [Eriogonum lagopus] Rabbit buckwheat | G3/S2 | | BIG, PAR, SHE | Bighorn Canyon NRA Cody BLM | Regional Endemic/ Core | Low | Abundant | Unknown | Stable | Moderate | Very High |
| Ipomopsis crebrifolia [Ipomopsis congesta ssp. crebrifolia] Compact gilia | G3/S2 S3 | | FRE, LIN, SUB, SWE, TET | Bridger-Teton NF Kemmerer BLM Pinedale BLM Rock Springs BLM | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Stable? | Moderate | Very High |
| Lesquerella paysonii Payson's bladderpod | G3/S3 | USFS R4 Sensitive | LIN, SUB, TET | Bridger-Teton NF Grand Teton NP Pinedale BLM Targhee NF | Regional Endemic/ Core | Moderat e | Uncommon | Unknown | Stable? | Moderate | Very High |
| Lomatium bicolor var. bicolor Wasatch biscuitroot | G4T3/ S2 | | LIN, SUB, TET, UIN | Bridger-Teton NF Rock Springs BLM Targhee NF | Regional Endemic/Edge | Low | Uncommon | Unknown | Unknown | Moderate | High |
| Mentzelia sinuata Wavy-leaf stickleaf | G3?/S 2 | | ALB, LAR | Rawlins BLM? | Regional Endemic/Edge | Low | Unknown | Unknown | Unknown | Moderate | High |
| Oonopsis wardii [Haplopappus wardii] Ward's goldenweed | G3/S3 | | ALB, CAR, NAT, PLA | Casper BLM Hutton Lake NWR Rawlins BLM | Local Endemic/Core | Moderate | Uncommon | Unknown | Increase? | Moderate | Very High |
| Pedicularis pulchella Mountain lousewort | G3/S2 | | BIG, FRE, HOT, JOH, PAR, SUB, TET | Bighorn NF Bridger-Teton NF Shoshone NF | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Penstemon caryi Cary beardtongue | G3/S3 | USFS R2 Sensitive | BIG, SHE, WAS | Bighorn NF Cody BLM Worland BLM | Regional Endemic/ Core | Low | Uncommon | Unknown | Stable? | Moderate | Very High |
| Penstemon pachyphyllus var. mucronatus [P. mucronatus] Sheep Creek beardtongue | G5T4/ S2 | | CAR, SWE | Ashley NF (FGNRA) Rawlins BLM Rock Springs BLM | Regional Endemic/Edge | Low | Uncommon | Unknown | Stable? | Moderate | High |

Plant Species of Potential Concern (continued)

| Scientific Name | Heri- | Federal | County | Managed Areas | Range Context | # Occurs. | Abundance | Historic | Recent | Intrinsic | Wyoming |
|--|----------------|--|------------------------------------|---|-------------------------------|-----------|-----------|----------|----------------------|--------------------|-------------------------|
| Common Name | tage Rank | Status | | - | | | | Trends | Trends | Vulner- ability | Contribu - tion Rank |
| Phlox opalensis Opal phlox | G3/S3 | | LIN, SUB, SWE, UIN | Ashley NF (FGNRA) Kemmerer BLM Pinedale BLM Rock Springs BLM Seedskadee NWR | Regional Endemic/ Core | Moderate | Abundant | Unknown | Stable | Moderate | Very High |
| Physaria eburniflora Devil's Gate twinpod | G3/S3 | | CAR, FRE, NAT | Casper BLM Lander BLM Rawlins BLM | Local Endemic/Core | Moderate | Uncommon | Unknown | Stable | Moderate | Very High |
| Physaria vitulifera Rydberg twinpod | G4/S2 | | ALB, CAR, NAT | Medicine Bow NF Rawlins BLM | Regional Endemic/Edge | Low | Rare | Unknown | Stable | Moderate | High |
| Sisyrinchium pallidum Pale blue-eyed grass | G2G3 / S2S3 | | ALB, CAR | Hutton Lake NWR? Mortensen Lake NWR Rawlins BLM | Regional Endemic/ Edge? | Low | Abundant | Unknown | Increase? | Moderate | High |
| Stanleya tomentosa var. tomentosa Hairy prince's-plume | G4T3/ S2 | | BIG, FRE, HOT, PAR | Bighorn Canyon NRA Bighorn NF Cody BLM Lander BLM Worland BLM | Regional Endemic/ Core | Low | Rare | Unknown | Moderate Decline? | Moderate | Very High |
| Sullivantia hapemanii var. hapemanii Hapeman's sullivantia | G3T3/ S3 | USFS R2 Sensitive | BIG, HOT, JOH, NAT, SHE, WAS | Bighorn NF Buffalo BLM Cody BLM Lander BLM? Worland BLM | Regional Endemic/ Core | Moderate | Uncommon | Unknown | Stable | Moderate | Very High |
| Symphyotrichum molle [Aster mollis] Soft aster | G3/S3 | USFS R2 Sensitive USFS R4 Sensitive | BIG, JOH, NAT, SHE, SUB, WAS | Bighorn NF Bridger-Teton NF? Buffalo BLM Casper BLM Worland BLM | Regional Endemic/Core | Moderate | Abundant | Unknown | Stable | Moderate | Very High |

Taxa Dropped From Previous Lists: The following species have been dropped from the 2002 WYNDD list of plant species of special concern. Unless otherwise noted, all have been found to be more common in the state than previously recognized and are no longer considered vulnerable to extirpation.

Adenocaulon bicolor (American trailplant) Agrimonia gryposepala (Common agrimony) Anemone virginiana (Virginia anemone) Antennaria neglecta (Field pussytoes) Astragalus nelsonianus

(Nelson's milkvetch; moved to potential spp. of concern list)

Carex paupercula (Bog sedge) Carex preslii

(Presl sedge; all specimens have been redetermined)

Cypripedium fasciculatum (Clustered lady's-slipper;

moved to potential spp. of

concern list)

Elymus diversiglumis

(Interrupted wild-rye)

Epilobium oregonense (Oregon willow-herb)

Juncus brevicaudatus

(Tweedy's rush)

Lonicera caerulea var. caurina (Western honev-suckle)

Ostrya virginiana var. virginiana

(Hop-hornbeam)

Piptantherum pungens

(Slender mountain-ricegrass) Solidago speciosa var. pallida (Showy goldenrod) **Status Uncertain:** This list includes species that may warrant conservation attention, but which are not currently listed due to uncertainties regarding taxonomy, identification, or confirmed occurrence in the state.

Allium drummondii (reported)

Alnus viridis var. crispa (status not known)

Arabis fructicosa (taxonomic questions)

Arnica amplexicaulis (status not known)

Artemisia pattersonii (reported)

Astragalus undescribed species (status not known)

Botrychium boreale (reported)

Botrychium hesperium (reported)

Botrychium pinnatum (reported)

Bouteloua eriopoda (reported)

Callirhoe involucrata (status not known)

Carex stenoptila (identification problems)

Cirsium barnebyi (reported)

Claytonia tuberosa ssp. rosea (status not known)

Dichanthelium depauperatum (reported)

Draba longipes (reported)

Erigeron coulteri (reported)

Erigeron pumilus var. gracilior (taxonomic questions)

Eriogonum inflatum (reported)

Festuca thurberi (status not known)

Gayophytum decipiens (identification problems)

Gentiana nutans (taxonomic questions)

Gilia ophthalmoides (taxonomic questions)

Hieracium fendleri (status not known)

Hymenopappus filifolius var. megacephalus (taxonomic questions)

Isoetes echinospora (taxonomic questions)

Juncus articulatus (status not known)

Lesquerella subumbellata (reported)

Listera borealis (status not known)

Lysichiton americanus (reported)

Mertensia arizonica (identification problems)

Mimulus tilingii var. tilingii (status not known)

Muhlenbergia mexicana (status not known)

Orthocarpus tenuifolius (reported)

Oryzopsis canadensis (reported)

Polystichum braunii (reported)

Potamogeton epihydrous (status not known)

Potamogeton fibrillosus (taxonomic questions)

Ribes hirtellum (reported)
Salix barclayi (status not known)
Stephanomeria exigua (reported)
Stipa thurberiana (reported)
Telesonix jamesii (reported)
Thalictrum revolutum (taxonomic questions)
Utricularia ochroleuca (reported, probably U. intermedia)

Animal Species of Concern – BIRDS

| Common Name | Heritage | | WY Cont. | | | |
|---|------------|--|----------|--|--|--|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| Common loon (Gavia immer) | G5/S1B/S2N | Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. SN Rank based on migratory concerns. | High | Low portion of its range in Wyoming, Restricted continental distribution, and equal security in Wyoming relative to other areas. | USFS R4 Sensitive, WYGF NSS1 | ALB, BIG, CAR, CRO, FRE, HOT, JOH, LIN, NAT, PAR, PLA, SHE, SUB, SWE, TET, UIN |
| Clark's grebe (Aechmophorus clarkii [Achemophorus occidentalis]) | G5/S1B | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS4 | ALB, FRE, LIN, NAT, PAR, TET |
| American white pelican (Breeding colonies) (Pelecanus erythrorhynchos) | G3/S1B | Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | ALB, CAR?, FRE?, NAT, PAR, TET |
| American bittern (Botaurus lentiginosus) | G4/S3B | Large WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderat e External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | ALB, BIG, CAR, CRO, FRE, GOS, LIN, PAR, SWE, TET, UIN |
| White-faced ibis (Plegadis chihi) | G5/S1B | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental dist ribution, and Uncertain security in Wyoming relative to other areas. | Wyoming BLM Sensitive, WYGF NSS3 | ALB, CAR, FRE, GOS, HOT, LAR, LIN, SUB, SWE, TET, UIN |
| Trumpeter swan (Cygnus buccinator) | G4/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | CAR, FRE, LIN, NAT, PAR, SUB, SWE, TET |
| Harlequin duck (Histrionicus histrionicus) | G4/S1B | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive, WYGF NSS3 | BIG?, FRE, HOT, LIN, PAR, SUB, TET |
| Bald eagle (Haliaeetus leucocephalus) | G4/S3B/S5N | Moderate WY Range. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS Threatened (proposed for de-listing), WYGF NSS2 | ALB, BIG, CAM, CAR, CON, FRE, GOS, HOT, JOH, LAR, LIN, NAT, NIO, PAR, PLA, SHE, SUB, SWE, TET, UIN, WAS? |
| Northern goshawk (Accipiter gentilis) | G5/S3 | Moderate WY Range. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS4 | ALB, BIG, CAR, CRO, FRE, JOH, LAR, LIN, NAT, PAR, SHE, SUB, SWE, TET, UIN, WAS |
| Ferruginous hawk (Buteo regalis) | G4/S4B/S5N | Large WY Range. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and More security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS3 | all counties |
| American peregrine falcon (Falco peregrinus anatum) | G4/T3/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | Recently de-listed by USFWS, USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS3 | ALB, CAR, JOH, PAR, SUB, SWE, TET |

| Common Name | Heritage | | WY Cont. | | | |
|---|----------------|--|-----------|--|--|---|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| White-tailed ptarmigan (Lagopus leucurus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | USFS R2 Sensitive | ALB, CAR?, FRE?, LIN, PAR, SWE?, TET |
| Columbian sharp-tailed grouse (Tympanuchus phasianellus columbianus) | G4/T3/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive | CAR |
| Greater sage grouse (Centrocercus urophasianus) | G4/S4 | Large WY Range. Moderate Biological Vulnerability. Moderate External Threat s. Patchy Distribution. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS ESA Petitioned, USFS R2 Sensitive, Wyoming BLM Sensitive | ALB?, BIG, CAM, CAR, CON, CRO, FRE, HOT, JOH, LIN, NAT, PAR, PLA, SHE, SUB, SWE, TET, UIN, WAS, WES |
| Whooping crane (Grus americana) | G1/SAB/ S1N | Accidental. Low Range Occupation. Low Abundance. Declining Abundance. High Biological Vulnerability. High External Threats. Patchy Distribution. | Low | Accidental or occasional in Wyoming | USFWS Endangered | FRE, LIN, PAR, SUB, SWE, TET, UIN |
| Snowy plover (Charadrius alexandrinus) | G4/SA | Accidental. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive | ALB, CAR, CRO, FRE, NAT, SWE |
| Mountain plover (Charadrius montanus) | G2/S2 | Moderate WY Range. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, WYGF NSS4 | ALB, BIG, CAM, CAR, CON, CRO, FRE, GOS, HOT, JOH, LAR, LIN, NAT, NIO, PAR, PLA, SUB, SWE, WAS, WES |
| Long-billed curlew (Numenius americanus) | G5/S3B | Moderate WY Range. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS3 | ALB, CAM, CAR, CON, CRO, FRE, GOS, HOT, LAR, LIN, NAT, NIO, PAR, PLA, SHE?, SUB, SWE, TET, UIN, WES |
| Caspian tern (Sterna caspia) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS3 | ALB, CAR, FRE, LIN, PAR, TET |
| Forster's tern (Sterna forsteri) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | ALB, CAR?, FRE, LIN |
| Black tern (Breeding colonies) (Chlidonias niger) | G4/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. Moderate Biological Vulnerability. High External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | ALB, LAR?, LIN?, PAR |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|---|------------------|--|------------------|--|--|--|
| (Beleiterie Tume) | Ium | Terrange State Faith Reasons | Ittilii | Typoning Contribution Runn Reasons | 1 ederar and State Status | County of Occurrence |
| Black-billed cuckoo (Coccyzus erythropthalmus) | G5/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | | ALB, BIG, CAM, CAR, CRO, FRE, GOS, HOT?, JOH, NAT, SHE, TET, WAS? |
| Yellow-billed cuckoo (Coccyzus americanus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | USFWS ESA Candidate (Western subspecies), USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | ALB, BIG, CAR, CON, CRO, FRE, GOS, HOT, JOH, NAT, PAR, PLA, SHE, SWE, TET, UIN |
| Northern pygmy-owl (Glaucidium gnoma) | G5/S2 | Small WY Range. Low Range Occupation. Low Abundance. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS4 | FRE, LIN?, PAR, TET, UIN? |
| Burrowing owl (Athene cunicularia [Speotyto cunicularia]) | G4/S3 | Moderate WY Range. Moderate Biological Vulnerability. High External Threats. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS4 | all counties |
| Great gray owl (Strix nebulosa) | G5/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R4 Sensitive, WYGF NSS4 | CAR, FRE, PAR, SUB, TET |
| Short -eared owl (Asio flammeus) | G5/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive | all counties |
| Boreal owl (Aegolius funereus) | G5/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive, WYGF NSS4 | ALB, CAR, FRE, JOH, LIN, NAT, PAR, PLA, SHE, TET, UIN |
| Calliope hummingbird (Stellula calliope) | G5/S3 | Moderate WY Range. Low Abundance. Uncertain Abundance Trends. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, CAR, LIN, PAR, SHE, SUB, TET |
| Lewis' woodpecker (Melanerpes lewis) | G4/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | ALB, CON, CRO, FRE, LAR, NAT, PAR, PLA, TET, UIN, WES |
| Williamson's sapsucker (Sphyrapicus thyroideus) | G5/S2 | Moderate WY Range . Low Range Occupation. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, BIG?, CAR, CON, FRE, JOH, LAR, LIN?, PAR, SHE, SUB, SWE?, TET, UIN? |
| Three-toed woodpecker (Picoides tridactylus) | G5/S3 | Moderate WY Range. Low Range Occupation. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive | ALB, BIG, CAR, CRO, FRE, HOT?, JOH, LIN?, NAT, PAR, SHE, SUB, SWE?, TET, UIN, WAS, WES? |
| Black-backed woodpecker (Picoides arcticus) | G5/S1 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS4 | CRO, FRÉ?, LIN, NAT?, PAR, TET, WES? |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|--|------------------|--|---|---|
| | | | | | | • |
| Loggerhead shrike (Lanius ludovicianus) | G4/S3 | Large WY Range. Low Range Occupation. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive | all counties |
| Western scrub-jay (Aphelocoma californica [Aphelocoma coerulescens]) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | CAR, LAR, NAT, SWE |
| Juniper titmouse (Baeolophus ridgwayi [Baeolophus griseus]) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS3 | FRE, SWE, UIN? |
| Bushtit (Psaltriparus minimus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS3 | NAT, SWE, UIN |
| Pygmy nuthatch (Sitta pygmaea) | G5/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS4 | ALB, BIG, CAM, CON, CRO, JOH, LAR, LIN, NAT, PAR, SHE, SUB, TET |
| Virginia's warbler (Vermivora virginiae) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Patchy Distribution. | High | Low port ion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, CAR?, CON?, CRO?, GOS, JOH?, LIN, SUB?, SWE, UIN |
| Black-throated gray warbler (Dendroica nigrescens) | G5/S2 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, FRE, NAT, SWE, UIN? |
| Sage sparrow (Amphispiza belli) | G5/S3 | Moderate WY Range. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | High portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive | ALB, BIG, CAM, CAR, CON, FRE, GOS, HOT, JOH, LIN, NAT, NIO, PAR, PLA, SUB, SWE, UIN, WAS, WES |
| Baird's sparrow (Ammodramus bairdii) | G4/S1B?/ SZN | Uncertain WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | Low | Occasional in Wyoming. | Wyoming BLM Sensitive, WYGF NSS4 | ALB?, CAM, CON, GOS?, LAR?, NIO?, PLA?, UIN |
| McCown's longspur (Calcarius mccownii) | G5/S2 | Moderate WY Range. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS4 | ALB, CAM, CAR, CON, CRO, FRE, GOS, JOH, LAR, NAT, NIO, PAR, PLA, SWE, WES |
| Chestnut-collared longspur (Calcarius ornatus) | G5/S1 | Small WY Range. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS4 | ALB, CAM, CAR, CON, CRO, FRE, GOS, LAR, NIO, PAR, WES |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|--|------------------|--|--------------------------|--|
| | | | | | | |
| Bobolink (Dolichonyx oryzivorus) | G5/S2 | Moderate WY Range. Low Range Occupation. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS4 | CAM, CAR, CRO, FRE, NAT, PAR, SHE |
| | | Patchy Distribution. | | | | |
| Scott's oriole (Icterus parisorum) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS3 | SWE |
| Brown-capped rosy finch [Rosy finch] (Leucosticte australis [Leucosticte arctoa]) | G4/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. High External Threats. Regional Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | | ALB, CAR? |
| Black-rosy finch [Rosy finch] (Leucosticte atrata [Leucosticte artcoa]) | G4/S1B/S2N | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. High External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, FRE, LIN, NAT, PAR, SUB, SWE, TET, UIN |

Species of Concern – FISH

| Common Name | Heritage | | WY Cont. | | | |
|--|----------|---|-----------|--|--|--|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| Shovelnose sturgeon (Scaphirhynchus platorynchus) | G4/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS2 | CAM, JOH, SHE |
| Goldeye (Hiodon alosoides) | G5/S2 | Moderate WY Range. Low Range Occupation. Unknown Abundance. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS2 | ЈОН |
| Leatherside chub (Gila copei [Snyderichthys copei]) | G3G4/S1 | Very Small WY Range. Low Range Occupation. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | Wyoming BLM Sensitive, WYGF NSS1 | LIN, TET, UIN |
| Humpback chub (Gila cypha) | G1/SX | Extirpated. Local Endemic. | Low | Accidental or occasional in Wyoming | USFWS Endangered | (SWE) |
| Roundtail chub [Bonytail] (Gila robusta) | G3/S3 | Small WY Range. Low Range Occupation. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS1 | CAR, LIN, SUB, SWE, UIN |
| Bonytail chub (Gila elegans) | G1/SX | Extirpated. Patchy Distribution. | Low | Accidental or occasional in Wyoming | USFWS Endangered | (SWE) |
| Western silvery minnow (Hybognathus argyritis [Hybognathus nuchalis]) | G4/S2 | Very Small WY Range. Low Range Occupation. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS1 | BIG, CAM, CRO, HOT, JOH, PAR, SHE, WAS |
| Sturgeon chub (Macrhybopsis gelida [Hybopsis gelida]) | G3/S1 | Small WY Range. Low Range Occupation. Unknown Abundance. High Biological Vulnerability. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, WYGF NSS1 | BIG, CAM, JOH, SHE, WAS |
| Pearl dace [Northern dace] (Margariscus margarita [Semotilus margarita]) | G5/S1 | Very Small WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS1 | NIO? |
| Hornyhead chub (Nocomis biguttatus) | G5/S1 | Very Small WY Range. Low Range Occupation. Moderate Biological Vulnerability. Moderate External Threats. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS1 | ALB, CAR, CON?, FRE, GOS, LAR?, NAT?, PLA |
| Suckermouth minnow (Phenacobius mirabilis) | G5/S2 | Very Small WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS1 | GOS, PLA |
| Finescale dace (Phoxinus neogaeus) | G5/S2 | Very Small WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS1 | CRO, NIO, WES? |
| Colorado River squawfish (Ptychocheilus lucius) | G1/SX | Extirpated. Local Endemic. | Low | Accidental or occasional in Wyoming | USFWS Endangered | (SWE) |
| Kendall Warm Springs dace (Rhinichthys osculus thermalis) | G5/T1/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | WYGF NSS2, USFWS Endangered | SUB |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|--|------------------|--|---|---|
| (Scientific Ivallie) | Kalik | Heritage State Kalik Reasons | Kalik | wyoning Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| D1 1 1 1 | G4/G2 | G HWAYD I D O | T T | N. 1 | Hand by a second second | CAD AD CUD ONE TEX |
| Bluehead sucker (Catostomus discobolus) | G4/S3 | Small WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS1 | CAR, LIN, SUB, SWE, TET, UIN |
| Flannelmouth sucker (Catostomus latipinnis) | G3G4/S3 | Small WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS1 | CAR, LIN, SUB, SWE, UIN |
| Snake River Sucker (Chasmistes muriei [Chasmistes liorus]) | GX/SX | Extirpated. Low Range Occupation. | Low | Accidental or occasional in Wyoming | | (TET) |
| Razorback sucker [Humpback sucker] (Xyrauchen texanus) | G1/SX | Extirpated. Low Range Occupation. | Low | Accidental or occasional in Wyoming | USFWS Endangered | (SWE) |
| Yellowstone cutthroat trout (Native pops.) (Oncorhynchus clarki bouvieri) | G4/T2/S2 | Moderate WY Range. Low Range Occupation. High Biological Vulnerability. Regional Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS3 | BIG, CAR?, FRE, HOT, PAR, SHE, SUB, TET, WAS |
| Westslope cutthroat trout (Oncorhynchus clarki lewisi) | G4/T3/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. High Biological Vulnerability. Regional Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R4 Sensitive | PAR, TET |
| Fine-spotted Snake River cutthroat trout (Native populations) (Oncorhynchus clarki ssp. 2 (undescribed subspecies)) | G4/T1Q/S1 | Small WY Range. Low Range Occupation. Low Abundance. High Biological Vulnerability. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS4 | FRE, LIN, SUB, TET |
| Colorado River cutthroat trout (Native populations) (Oncorhynchus clarki pleuriticus) | G4/T2/S1 | Small WY Range. Low Range Occupation. Low Abundance. High Biological Vulnerability. Regional Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | USFWS Petitioned for ESA listing, USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | CAR, LIN, SUB, SWE, UIN |
| Bonneville cutthroat trout [Bear River cutthroat trout] (Native populations) (Oncorhynchus clarki utah) | G4/T2/S1 | Very Small WY Range. Low Range Occupation. High Biological Vulnerability. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | LIN, UIN |
| Greenback cutthroat trout (Oncorhynchus clarki stomias) | G4/T2/SX | Extirpated. High Biological Vulnerability. Regional Endemic. | Low | Accidental or occasional in Wyoming | USFWS Threatened | (ALB, LAR) |
| Arctic grayling (Upper Missouri River fluvial population) (Thymallus arcticus) | G5/T2/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS4, USFWS ESA Candidate | PAR, TET |
| Orangethroat darter (Etheostoma spectabile) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS2 | LAR |

$Species\ of\ Concern-AMPHIBIANS$

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|---|------------------|--|------------------|---|--|---------------------------------------|
| | | | | · · · · · · | | • |
| Boreal western toad (Northern Rocky Mountain population) (Bufo boreas boreas (undescribed taxon)) | G4/T4/S1 | Small WY Range. Low Range Occupation. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | FRE, LIN, PAR, SUB, TET, UIN, WAS? |
| Boreal western toad (Southern Rocky Mountain population) (Bufo boreas boreas (undescribed taxon)) | G4/T1/S1 | Recognized taxon for management purposes. Very Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. High Biological Vulnerability. High External Threats. Regional Endemic. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | USFWS ESA Candidate, USFS R2 Sensitive, WYGF NSS2 | ALB, CAR, CON, LAR |
| Wyoming toad (Bufo baxteri [Bufo hemiophrys baxteri]) | G1/T1/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. High Biological Vulnerability. High External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFWS Endangered, WYGF NSS1 | ALB |
| Great Basin spadefoot toad (Spea intermontana [Scaphiopus intermontanus]) | G5/S3 | Moderate WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | Wyoming BLM Sensitive, WYGF NSS4 | CAR?, FRE, LIN, NAT, SWE, UIN |
| Northern leopard frog (Rana pipiens) | G5/S3 | Large WY Range. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS4 | all counties |
| Columbia spotted frog [Spotted frog] (statewide) (Rana luteiventris [Rana pretiosa]) | G4/S3 | Moderate WY Range. Low Range Occupation. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS5 | FRE, LIN, PAR, SHE, SUB, TET |
| Wood frog (statewide) (Rana sylvatica) | G5/S1 | Small WY Range. Low Range Occupation. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | ALB, BIG, CAR, JOH, SHE |

Species of Concern – REPTILES

| Common Name | Heritage | | WY Cont. | | | |
|--|----------|--|-----------|--|--------------------------|--|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| Ornate box turtle (Terrapene ornata ornata) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | GOS |
| Six-lined racerunner [Prairie linedracerunner] (Cnemidophorus sexlineatus viridis) | G5/S2 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | GOS?, PLA |
| Northern many-lined skink (Eumeces multivirgatus multivirgatus) | G5/S1 | Very Small WY Range. Low Range Occupation. Unknown Abundance and Trends. Regional Endemic. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | GOS, LAR, NIO, PLA |
| Northern plateau lizard (Sceloporus undulatus elongatus) | G5/T5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAR, SWE |
| Red-lipped prairie lizard [Orange-lipped plateau lizard] (Sceloporus undulatus erythrocheilus) | G5/T5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, CON, LAR?, PLA |
| Northern prairie lizard (Sceloporus undulatus garmani) | G5/T5/S1 | Very Small WY Range. Low Range Occupation. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | GOS, LAR?, NIO?, PLA |
| Tree lizard (Urosaurus ornat us) | G5/S2 | Very Small WY Range. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | SWE |
| Lesser earless lizard (Holbrookia maculata) | G5/S2 | Very Small WY Range. Low Range Occupation. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | | GOS?, LAR, PLA |
| Rubber boa (Charina bottae) | G5/S2 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | BIG?, FRE, LIN, PAR, SHE, SUB, TET, WAS |
| Smooth green snake (eastern and western) (Liochlorophis vernalis [Lv. vernalis, L.v. blanchardi; Opheodrys v. vernalis, O.v. blanchardi]) | G5/S2 | Small WY Range. Low Range Occupation. Uncertain Abundance Trends. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, CAR, CON, CRO, NAT, WES |
| Black Hills redbelly snake (Storeria occipitomaculata pahasapae) | G5/T3/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFS R2 Sensitive | CRO, WES? |
| Midget faded rattlesnake (Crotalus viridis concolor) | G5/T3/S1 | Very Small WY Range. Uncertain Abundance Trends. High External Threats. Regional Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | Wyoming BLM Sensitive | SWE |

$Species\ of\ Concern-MAMMALS$

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|---|------------------|--|--|---|
| (Serentific 1 (units) | 244 | Tiende Suite Tunii Teensons | | , your government remains reasons | 1 ederar and search status | County of Securitories |
| Hayden's shrew (Sorex haydeni) | G4/S3 | Small WY Range. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, CAR, CON, CRO, JOH, SHE, WES |
| Southern Rocky Mountain pygmy shrew (Sorex hoyi montanus) | G?/S1 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. Patchy Distribution. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS2 | ALB, CAR |
| Preble's shrew (Sorex preblei) | G4/S1 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | LIN, PAR?, TET, UIN |
| Fringed myotis (statewide) (Myotis thysanodes) | G4G5/S2 | Large WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. | Low | Medium portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | ALB, BIG, CAR, CRO?, FRE?, GOS, JOH?, LAR?, NAT?, PAR, PLA?, SHE, SUB, SWE?, WAS, WES |
| Black Hills fringed myotis (Myotis thysanodes pahasapensis) | G?/T1/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. Local Endemic. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | CRO?, WES |
| Yuma myotis (Myotis yumanensis) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | BIG?, CAR, FRE?, JOH?, NAT?, PAR, SHE, SWE, UIN?, WAS? |
| Spotted bat (Euderma maculatum) | G4/S3 | Large WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | ALB?, BIG, FRE?, HOT?, JOH?, LIN?, PAR, SUB?, SWE, TET?, UIN?, WAS |
| Townsend's big-eared bat (Corynorhinus townsendii [Plecotus townsendii]) | G4/S2 | Large WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. High External Threats. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS2 | ALB?, BIG, CAM?, CAR?, CON?, CRO?, FRE?, GOS?, HOT?, JOH?, LAR?, LIN?, NAT?, NIO?, PAR, PLA?, SHE?, SUB?, SWE, TET?, UIN?, WAS, WES? |
| Pallid bat (Antrozous pallidus) | G5/S1 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS2 | BIG?, CAR?, GOS, HOT, SWE, WAS |
| Bighorn Mountain pika (Ochotona princeps obscura) | G5/T1/S1 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. High External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | | BIG, JOH?, SHE, WAS? |

Animal Species of Concern – MAMMALS (continued)

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|---|------------------|---|--|---|
| (Belefitine Ttaine) | Kank | Tierrage State Rank Reasons | Kuiik | Wyoming Contribution Rank Reasons | 1 cuci ai anu state status | County of Occurrence |
| Pygmy rabbit (Brachylagus idahoensis [Sylvilagus idahoensis]) | G4/S1 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS Petitioned for ESA listing, Action Pending; Wyoming BLM Sensitive; WYGF NSS3 | FRE, LIN, SUB, SWE, UIN |
| Bighorn Mountain snowshoe hare (Lepus americanus seclusus) | G5/T3/S1 | Very Small WY Range. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | | BIG?, JOH?, SHE?, WAS? |
| Cliff chipmunk (Tamias dorsalis utahensis [Eutamias dorsalis]) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. High External Threats. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | SWE |
| Black Hills yellow- bellied marmot (Marmota flaviventris dacota) | G?/S2 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | | CRO?, WES? |
| Allen's thirteen-lined ground squirrel (Spermophilus tridecemlineatus alleni) | G5/T1/S1 | Very Small WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive | BIG?, FRE, HOT, JOH, SUB, WAS? |
| Black-tailed prairie dog (Large towns) (Cynomys ludovicianus) | G4/S2 | Moderate WY Range. Low Range Occupation. Declining Abundance. High Biological Vulnerability. High External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS ESA Candidate, USFS R2 Sensitive, WYGF NSS3 | ALB, CAM, CON, CRO, GOS?, JOH, LAR, NAT, NIO, PAR, PLA, SHE, WAS?, WES |
| White-tailed prairie dog (Large towns) (Cynomys leucurus) | G4/S3 | Large WY Range. Low Range Occupation. High Biological Vulnerability. High External Threats. Regional Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS Petitioned for ESA listing, Action Pending; USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS3 | ALB, BIG, CAR, FRE, HOT, JOH, LIN, NAT, PAR, SUB, SWE, UIN, WAS |
| Black Hills red squirrel (Tamiasciurus hudsonicus dakotensis) | G5/T3/S2 | Very Small WY Range. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | | CRO?, WES? |
| Wyoming pocket gopher (Thomomys clusius [Thomomys talpoides]) | G2/S2 | Small WY Range. Low Range Occupation. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive | CAR, SWE |
| Idaho pocket gopher (Thomomys idahoensis [Thomomys talpoides]) | G4/S2 | Small WY Range. Low Range Occupation. Uncertain Abundance Trends. Regional Endemic. | Very High | High port ion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | Wyoming BLM Sensitive, WYGF NSS3 | LIN, SUB, UIN |
| Canyon mouse (Peromyscus crinitus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | SWE |
| Pinon mouse (Peromyscus truei) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | WYGF NSS3 | SWE |

Animal Species of Concern – MAMMALS (continued)

| Common Name | Heritage | W. W. G. A. D. J. D. | WY Cont. | W. I. G. (D. I. D. I. D. | F. 1. 1. 16/4 6/4 | G 4 60 |
|---|----------|--|-----------|---|---|--|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| Water vole (statewide) (Microtus richardsoni [Arvicola richardsoni]) | G5/S2 | Moderate WY Range. Low Range Occupation. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to | USFS R2 Sensitive, WYGF NSS3 | BIG, FRE, HOT?, JOH, LIN, PAR, SHE, SUB, TET, UIN, WAS |
| Black Hills southern red-backed vole (Clethrionomys gapperi brevicaudus) | G5/T3/S2 | Very Small WY Range. Uncertain Abundance Trends. Local Endemic. | High | other areas. Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | CRO?, WES? |
| Bear Lodge meadow jumping mouse (Zapus hudsonius campestris) | G5/T3/S1 | Small WY Range. Low Range Occupation. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Regional Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | CRO, WES |
| Preble's meadow jumping mouse (Zapus hudsonius preblei) | G5/T2/S1 | Small WY Range. Low Range Occupation. Low Abundance. High Biological Vulnerability. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFWS Threatened | ALB, CAR?, CON, GOS?, LAR, PLA |
| Gray wolf (Canis lupus) | G4/S1 | Small WY Range. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFWS Endangered | FRE, PAR, TET, SUB? |
| Swift fox (Vulpes velox) | G3/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Low portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, Wyoming BLM Sensitive, WYGF NSS3 | ALB, CAM, CAR, CON, CRO?, FRE, GOS, JOH?, LAR, NAT, NIO, PLA, SWE, WES |
| Grizzly bear (Ursus arctos horribilis) | G4/S1 | Small WY Range. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Patchy Dist. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Equal security in Wyoming relative to other areas. | USFWS Threatened | FRE, PAR, SUB, TET |
| Fisher (Martes pennanti) | G5/S1 | Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | USFS R4 Sensitive | ALB?, CAR?, FRE?, JOH?, PAR, SUB, TET |
| Black-footed ferret (Mustela nigripes) | G1/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. High Biological Vulnerability. High External Threats. Regional Endemic. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Less security in Wyoming relative to other areas. | USFWS Endangered, WYGF NSS1 | ALB, CAR |
| North American Wolverine (Gulo gulo luscus [Gulo luscus]) | G4/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | USFWS ESA Listing Denied, USFS R2 Sensitive, USFS R4 Sensitive, WYGF NSS3 | ALB?, BIG?, CAR?, FRE, LIN, PAR, SHE?, SUB, TET |
| River otter (Lontra canadensis [Lutra Canadensis]) | G5/S3 | Moderate WY Range. Low Range Occupation. Patchy Distribution. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive | ALB, BIG?, CAR?, FRE, HOT?, JOH, LIN, NAT, PAR, SHE?, SUB, SWE, TET, UIN? |
| Canada lynx [North American lynx] (Lynx canadensis [Felis lynx]) | G5/S1 | Small WY Range. Low Range Occupation. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | USFWS Threatened, WYGF NSS1 | ALB?, BIG, CAR?, FRE, JOH, LIN, PAR, SHE, SUB, TET, UIN |
| American bison (Free- ranging herds) (Bos bison [Bison bison]) | G4/S1 | Very Small WY Range. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and More security in Wyoming relative to other areas. | USFWS Petitioned for ESA listing (Yellowstone DPS) | PAR, TET |

Animal Species of Potential Concern – ALL TAXA

Please note that not all species on the potential concern list have been fully incorporated into the new ranking system. S-ranks given in square brackets (e.g., [S3B]) are carried over from the 1999 list and S-rank reasons in square brackets are from the range comments section of the 1999 list. A designation of Not Ranked (NR) in the Wyoming contribution rank columns means that these species have not yet been assigned a Wyoming contribution rank. All these fields will be updated in future lists, at which time decisions may be made to elevate or remove certain species from this list.

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|---|------------------|---|------------------|--|---|--|
| | | | | | | |
| BIRDS | | | | | | |
| Least bittern (Ixobrychus exilis) | G5/S1B | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Accidental or occasional in Wyoming | | ALB? |
| Snowy egret (Egretta thula) | G5/[S3B] | [Breed: N periph.] | NR | NR | WYGF NSS3 | ALB, CAR, FRE, GOS, HOT, JOH, LAR, LIN, NAT, PAR, SHE, SUB, SWE, TET, UIN |
| Black-crowned night- heron (Nycticorax nycticorax) | G5/[S3B] | [Breed: interior periph.] | NR | NR | WYGF NSS3 | all counties? |
| Ring-necked duck (Aythya collaris) | G5/S4B | Moderate WY Range. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Equal security in Wyoming relative to other areas. | | ALB, BIG, CAR, FRE, JOH, LIN?, NAT, PAR, SUB, TET, UIN |
| Bufflehead (Bucephala albeola) | G5/S2B | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. High External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAR, FRE, JOH, LIN, NAT, PAR, SUB, SWE, TET, UIN |
| Tundra swan (Cygnus columbianus) | G5/S2N | Uncertain WY Range. | Low | Accidental or occasional in Wyoming | | all except NIO? |
| Common goldeneye (Bucephala clangula) | G5/[S3B] | [Breed: S periph. Winter: core] | NR | NR | | all except CRO, WES |
| Golden eagle (Aquila chrysaetos) | G5/[S3B] | [Year-round: core] | NR | NR | | all counties |
| Osprey (Pandion haliaetus) | G5/[S3B] | [Breed: S periph.] | NR | NR | | all counties |
| Merlin (Falco columbarius) | G5/S4 | Large WY Range. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | ALB, BIG, CAM, CAR, CON, CRO, FRE, HOT, JOH, LAR, LIN, NAT, PAR, SHE, SUB, SWE, TET, UIN, WAS, WES |
| Arctic peregrine falcon (Falco peregrinus tundrius) | G4/T3/SU | Uncertain WY Range. | Low | Accidental or occasional in Wyoming | Recently de-listed by USFWS, Wyoming BLM Sensitive, WYGF NSS3 | all counties? |
| Greater prairie-chicken (Tympanuchus cupido) | G4/SA | Accidental. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. High Biological Vulnerability. High External Threats. Regional Endemic. | Low | Accidental or occasional in Wyoming | USFS R2 Sensitive | LAR? |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|---|------------------|--|------------------|--|--------------------------|--|
| (Belefitine Ivaine) | Kank | Heritage State Kank Keasons | Kank | wyoning Contribution Rank Reasons | rederar and State Status | County of Occurrence |
| Virginia rail (Rallus limicola) | G5/[S3B] | [Breed: core] | NR | NR | | ALB, CAR, CON, GOS, JOH?, LAR, LIN, NAT, PAR, PLA, SHE, TET, UIN |
| Sandhill crane (Grus canadensis) | G5/S3B/S5N | Small WY Range. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | all counties |
| Piping plover (Charadrius melodus) | G3/SA | Accidental. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | Low | Accidental or occasional in Wyoming | USFWS Threatened | CRO?, GOS?, LAR, NAT? |
| Black-necked stilt (Himantopus mexicanus) | G5/[S3B] | [Breed: N periph.] | NR | NR | | ALB, BIG, CAR, FRE, GOS, JOH, LIN, NAT, PAR, SUB, SWE, TET, UIN |
| American avocet (Recurvirostra americana) | G5/[S3B] | [Breed: core] | NR | NR | | all counties |
| Red-necked phalarope (Phalaropus lobatus) | G5/[S3N] | [Migratory only?] | NR | NR | | CAR, NAT, PAR, SWE, TET |
| Herring gull (Breeding colonies) (Larus argentatus) | G5/SA | Accidental. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Accidental or occasional in Wyoming | | ALB |
| Ring-billed gull (Breeding colonies) (Larus delawarensis) | G5/S2 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, FRE, PAR?, TET? |
| Eastern phoebe (Sayornis phoebe) | G5/SA | Accidental. Low Abundance. Uncertain Abundance Trends. | Low | Accidental or occasional in Wyoming | | CRO, JOH, SHE, SWE |
| Hammond's flycatcher (Empidonax hammondii) | G5/S4 | Moderate WY Range. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, FRE, GOS, LIN, PAR, SHE, SUB, SWE, TET, UIN |
| Winter wren (Troglodytes troglodytes) | G5/SA | Accidental. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Accidental or occasional in Wyoming | | FRE, LIN, NAT, SHE, TET |
| American Dipper (Cinclus mexicanus) | G5/S4 | Moderate WY Range. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | BIG?, JOH?, LIN?, PAR?, SHE? SUB?, TET?, WAS? |
| Golden-crowned kinglet (Regulus satrapa) | G5/S3B/S4N | Moderate WY Range. Low Range Occupation. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Equal security in Wyoming relative to other areas. | USFS R2 Sensitive | ALB, BIG, CAM, CAR, CRO, FRE, JOH, LAR, LIN, NAT, PAR, SHE, SUB, TET |
| Eastern bluebird (Sialia sialis) | G5/S2 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | | ALB, CON, CRO, FRE, GOS, JOH, LAR, PLA, WES |
| Townsend's warbler (Dendroica townsendi) | G5/SA | Accidental. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Accidental or occasional in Wyoming | | FRE, HOT, PAR, SUB, SWE, TET |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|---|------------------|--|---|--|
| (2000000) | | | | _ · · · J • · · · · · · · · · · · · · · · | | , 000000, 00000000000000000000000000000 |
| Rose-breasted grosbeak (Pheucticus ludovicianus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Accidental or occasional in Wyoming | | ALB, NAT |
| Dickcissel (Spiza americana) | G5/S1 | Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAM, CRO, FRE, GOS, LAR, SHE, WES |
| Cassin's sparrow (Aimophila cassinii) | G5/SA | Accidental. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Accidental or occasional in Wyoming | USFS R2 Sensitive | GOS |
| Grasshopper sparrow (Ammodramus savannarum) | G5/S4 | Moderate WY Range. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS4 | ALB, BIG, CAM, CAR, CON, CRO, FRE, GOS, HOT, JOH, LAR, LIN, NAT, NIO, PAR, PLA, SHE, SUB, SWE, UIN, WAS, WES |
| Brewer's sparrow (Spizella breweri) | G5/S5 | Large WY Range. Declining Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Medium | High portion of its range in Wyoming, Wide continental distribution, and More security in Wyoming relative to other areas. | USFS R2 Sensitive, Wyoming BLM Sensitive | all counties |
| White-winged junco (Junco hyemalis aikeni) | G5/T4/S3 | Small WY Range. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAM, CRO, LAR, NAT, NIO, PAR, SHE, WES |
| Northern bobwhite (Native populations) (Colinus virginianus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | GOS, LAR, NIO, PLA |
| California gull (Breeding colonies) (Larus californicus) | G5/[S2B] | [Breed: S periph.] | NR | NR | | ALB, CAR, FRE, NAT?, PAR, TET |
| Common tern (Sterna hirundo) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | | ALB, FRE |
| Barn owl (Tyto alba) | G5/S2 | Uncertain WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAM?, CON?, GOS?, LAR?, NIO?, PLA?, SWE?, UIN?, WES? |
| Western screech-owl (Otus kennicottii) | G5/S2 | Very Small WY Range. Low Range Occupation. Low Abundance. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | LIN?, SUB, TET, UIN? |
| Eastern screech-owl (Otus asio) | G5/[S3] | [Year-round: W periph.] | NR | NR | | ALB, BIG, CAM, CAR, CON, CRO, FRE?, GOS, HOT, JOH, LAR, NAT, NIO, PLA, PAR?, SHE, WAS, WES |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|---|------------------|---|------------------|---|---|---|
| | | | | | | |
| Flammulated owl (Otus flammeolus) | G4/S1 ? | Uncertain WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Patchy Distribution. | High | Uncertain portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive | ALB, BIG, FRE, NAT, TET |
| Chimney swift (Chaetura pelagica) | G5/[S3B] | [Breed: W periph.] | NR | NR | | CAR, CON, GOS, LAR, NAT, NIO?, PLA, SHE |
| Ash-throated flycatcher (Myiarchus cinerascens) | G5/[S3B] | [Breed: N periph.] | NR | NR | WYGF NSS3 | CAR, NAT, SWE, UIN |
| Canyon wren (Catherpes mexicanus) | G5/[S2S3] | [Year-round: NE periph.] | NR | NR | | BIG, CON, CRO, FRE, JOH, NAT, PAR, PLA, SHE, SWE, TET, WAS |
| Sage thrasher (Oreoscoptes montanus) | G5/S5 | Large WY Range. Moderate Biological Vulnerability. Moderate External Threats. | Medium | High portion of its range in Wyoming, Wide continental distribution, and More security in Wyoming relative to other areas. | Wyoming BLM Sensitive | ALB, BIG, CAM, CAR, CRO?, CON, FRE, GOS, HOT, JOH, LAR, LIN, NAT, NIO, PAR, PLA?, SHE, SUB, SWE, TET, UIN, WAS, WES |
| Red-eyed vireo (Vireo olivaceous) | G5/[S3B] | [Breed: SW periph.] | NR | NR | | ALB, CON, CRO, FRE, JOH, NAT, PAR, PLA, SHE, SWE, TET, WES |
| Blue grosbeak (Guiraca caerulea) | G5/[S3B] | [Breed: N periph.] | NR | NR | | BIG, CAR, FRE, GOS, LAR, NAT, PLA, SHE, SWE |
| White-winged crossbill (Loxia leucoptera) | G5/S2 | Moderate WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Equal security in Wyoming relative to other areas. | | ALB, CRO, FRE, LAR, PAR, SWE, TET, WES |
| Clay-colored sparrow (Spizella pallida) | G5/[S3B] | [Breed: S periph.] | NR | NR | | ALB, CAM, CON?, CRO, FRE, GOS, JOH, LAR, NAT, PAR, SHE, TET, WAS |
| Indigo bunting (Passerina cyanea) | G5/[S3B] | [Breed: WNW periph.] | NR | NR | WYGF NSS4 | CRO, WES |
| FISH | | | | | | |
| Common shiner (Luxilus cornutus) | G5/[S3S4] | [W periph. Disjunct] | NR | NR | WYGF NSS3 | ALB, CAR, CON, FRE?, GOS, LAR, NAT, PLA |
| Iowa darter (Etheostoma exile) | G5/[S3S4] | [SW periph. Disjunct] | NR | NR | WYGF NSS4 | ALB, CAR, CON, FRE?, GOS, LAR, NAT, NIO, PLA |
| Burbot [Ling] (Lota lota) | G5/[S3S4] | [S periph.] | NR | NR | WYGF NSS2 | BIG, FRE, HOT, PAR, WAS |
| Sauger (Stizostedion canadense) | G5/[S3S4] | [W periph.] | NR | NR | WYGF NSS2 | BIG, CAM, FRE, HOT, JOH, PAR, SHE, WAS |
| AMPHIBIANS | | | | | | |
| Tiger salamander (Ambystoma tigrinum) | G5/S4 | Large WY Range. Uncertain Abundance Trends. Moderate Biological Vulnerability. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS4 | all counties |
| Great Plains toad (Bufo cognatus) | G5/[S3] | [W periph.] | NR | NR | WYGF NSS4 | CAM, CRO?, NIO? WES |

| Common Name | Heritage | | WY Cont. | | | |
|---|------------|--|-----------|--|--|---|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| Columbia spotted frog [Spotted frog] (Bighorn Mountain Spotted Frog) (Rana luteiventris [Rana pretiosa] (undescribed taxon)) | G4/T1Q/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFS R2 Sensitive, USFS R4 Sensitive, Wyoming BLM Sensitive, WYGF NSS5 | SHE |
| Wood frog (Southern Rockies Wood Frog) (Rana sylvatica (undescribed taxon)) | G5/T2Q/S1 | Small WY Range. Low Range Occupation. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | ALB, CAR |
| Wood frog (Bighorn Mountain Wood Frog) (Rana sylvatica (undescribed taxon)) | G5/T1Q/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | BIG, JOH, SHE |
| REPTILES | | | | | | |
| Milk snake (Lampropeltis triangulum) | G5/S3 | Moderate WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | 0 |
| Great Basin gopher snake (Pituophis catenifer deserticola [P. melanoleucus deserticola]) | G5/T5/S3 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | SWE |
| Western plains garter snake (Thamnophis radix haydenii) | G5/T5/[S4] | [W periph.] | NR | NR | | CAM, CRO?, GOS, LAR, NIO, PLA?, WES? |
| Eastern yellowbelly racer (Coluber constrictor flaviventris) | G5/T5/[S4] | [W periph.] | NR | NR | | ALB, BIG?, CAR, CAM, CON, CRO, GOS, JOH, NAT, NIO?, PLA, SHE, WAS?, WES |
| Spiny softshell turtle (Apalone spinfera [Trionyx spiniferus]) | G5/[S4] | [W periph.] | NR | NR | | BIG, CAM, CRO?, GOS, HOT, JOH, NIO, PLA, SHE, WAS, WES? |
| MAMMALS | | | | | | |
| Dwarf shrew (Sorex nanus) | G4/S4 | Large WY Range. Low Range Occupation. Uncertain Abundance Trends. Patchy Distribution. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | WYGF NSS3 | ALB, CAM, CAR, CON?, FRE?, HOT?, JOH, LAR?, LIN, NAT?, PAR, PLA?, SUB?, SWE?, TET, UIN |
| Eastern mole (Scalopus aquaticus) | G5/S2 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | GOS, LAR?, PLA |
| Hoary bat (Lasiurus cinereus) | G5/S4 | Large WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB, BIG?, CAM, CAR, CON, CRO, FRE?, GOS?, HOT?, JOH?, LAR?, LIN, NAT?, NIO?, PAR, PLA?, SHE?, SUB?, SWE, TET, UIN?, WAS, WES? |

| Common Name | Heritage | | WY Cont. | | | |
|--|-----------|--|-----------|--|-------------------------------------|--|
| (Scientific Name) | Rank | Heritage State Rank Reasons | Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
| | | | | | | |
| Silver-haired bat (Lasionycteris noctivagans) | G5/[S3B] | [Breed: core] | NR | NR | | all except SUB, TET |
| Western small-footed myotis (Myotis ciliolabrum [Myotis leibii]) | G5/[S3B] | [Breed: core Winter: N periph.] | NR | NR | WYGF NSS3 | ALB, BIG, CAR, CON, CRO, FRE, GOS, HOT, JOH, LAR, LIN, NAT, PAR, PLA, SHE, SWE, WAS |
| Long-legged myotis (Myotis volans) | G5/[S3B] | [Breed: E periph.] | NR | NR | WYGF NSS2 | all except CAM, NIO |
| Long-eared myotis (Myotis evotis) | G5/S4 | Large WY Range. Low Range Occupation. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | Wyoming BLM Sensitive, WYGF NSS2 | ALB?, BIG, CAM?, CAR, CON?, CRO, FRE, GOS?, HOT, JOH, LAR?, LIN?, NAT?, NIO?, PAR, PLA?, SHE, SUB, SWE, TET, UIN, WAS, WES? |
| Eastern cottontail (Sylvilagus floridanus) | G5/S3 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. | Low | Accidental or occasional in Wyoming | | GOS, LAR, PLA? |
| Wyoming ground squirrel (Spermophilus elegans) | G5/[S3S4] | [Reg. endm.] | NR | NR | | ALB, CAR, CON, FRE, HOT, LAR, LIN, NAT, PAR, PLA?, SWE, SUB, TET, UIN, WAS |
| Uinta ground squirrel (Spermophilus amatus) | G5/[S3S4] | [Reg. endm.] | NR | NR | | FRE, LIN, PAR, SUB, SWE, TET, UIN |
| Abert's squirrel (Sciurus aberti) | G5/S1 | Very Small WY Range. Low Abundance. Uncertain Abundance Trends. Regional Endemic. | Low | Accidental or occasional in Wyoming | | ALB, LAR |
| Red squirrel (Bighorn Mountain population) (Tamiasciurus hudsonicus (undescribed taxon)) | G5/T3Q/S3 | Very Small WY Range. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | | BIG?, JOH?, SHE?, WAS? |
| Northern flying squirrel (Black Hills population) (Glaucomys sabrinus (undescribed taxon)) | G5/T2Q/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | CRO, WES |
| Olive-backed pocket mouse (Perognathus fasciatus) | G5/[S4] | [Core] | NR | NR | WYGF NSS3 | ALB, BIG, CAR, CAM, CON, CRO?, FRE, GOS, HOT?, JOH, NAT, NIO, LAR, PAR?, PLA, SHE, SWE, WAS, WES? |
| White-footed mouse (Peromyscus leucopus) | G5/S3 | Small WY Range. Low Range Occupation. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Less security in Wyoming relative to other areas. | | CAM, CRO, JOH?, NIO, SHE |
| Southern red-backed vole (Bighorn Mountain population) (Clethrionomys gapperi (undescribed taxon)) | G5/T3Q/S3 | Very Small WY Range. Uncertain Abundance Trends. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | | BIG?, JOH?, SHE?, WAS? |

| Common Name (Scientific Name) | Heritage Rank | Heritage State Rank Reasons | WY Cont. Rank | Wyoming Contribution Rank Reasons | Federal and State Status | County of Occurrence |
|--|------------------|---|------------------|--|---------------------------------|---|
| (Scientific Ivame) | Kalik | Heritage State Kank Reasons | Kalik | wyoning Contribution Rank Reasons | reuer at and State Status | County of Occurrence |
| Water vole (Bighorn Mountain population) (Microtus richardsoni [Arvicola richardsoni] (undescribed taxon)) | G5/T2Q/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive, WYGF NSS3 | BIG, JOH, SHE, WAS |
| Native Rocky Mountain red fox (Vulpes vulpes macroura) | G?/T1Q/S1 | Small WY Range. Low Range Occupation. Low Abundance. Declining Abundance. Moderate Biological Vulnerability. High External Threats. Patchy Distribution. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | | ALB?, CAR?, FRE?, HOT?, LIN?, PAR?, SUB?, TET?, UIN? |
| Common gray fox (Urocyon cinereoargenteus) | G5/S2 | Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAM?, CON?, CRO?, GOS?, JOH?, NAT?, NIO?, PLA?, WES |
| Ringtail (Bassariscus astutus) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. | Low | Accidental or occasional in Wyoming | | ALB, CAR?, LIN, SWE, UIN? |
| American marten [Pine marten] (Bighorn Mountain population) (Martes americana (undescribed taxon)) | G5/T2Q/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | Very High | High portion of its range in Wyoming, Restricted continental distribution, and Equal security in Wyoming relative to other areas. | USFS R2 Sensitive | BIG, JOH, SHE, WAS |
| American marten [Pine marten] (Black Hills population) (Martes americana (undescribed taxon)) | G5/T2Q/S1 | Reintroduced Taxon. Very Small WY Range. Low Range Occupation. Low Abundance. Moderate Biological Vulnerability. Moderate External Threats. Local Endemic. | High | Medium portion of its range in Wyoming, Restricted continental distribution, and Uncertain security in Wyoming relative to other areas. | USFS R2 Sensitive | CRO?, WES? |
| Least weasel (Mustela nivalis [Mustela rixosa]) | G5/S1 | Very Small WY Range. Low Range Occupation. Low Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Accidental or occasional in Wyoming | WYGF NSS3 | CAM, JOH?, SHE |
| Plains (eastern) spotted skunk (Spilogale put orius interruptua) | G5/T4/S3 | Moderate WY Range. Low Range Occupation. Unknown Abundance. Uncertain Abundance Trends. Moderate Biological Vulnerability. Moderate External Threats. | Low | Low portion of its range in Wyoming, Wide continental distribution, and Uncertain security in Wyoming relative to other areas. | | CAM?, CON?, CRO?, GOS?, JOH?, LAR?, NIO?, PLA?, SHE?, WES? |
| Bighorn sheep (Ovis canadensis) | G4/[S3S4] | [Historic: core Extant: disjunct] | NR | NR | | ALB, BIG, CAR, CON?, FRE, HOT, LIN, NAT?, PAR, SUB, TET, WAS? |

Species Dropped from Previous Lists – All Taxa

| Common Name | |
|---|---------------------------------|
| (Scientific Name) | Action Taken |
| Milk snake (Lampropeltis triangulum) | Moved to Potential Concern List |
| Great Basin gopher snake (Pituophis catenifer deserticola [P. | Moved to Potential Concern List |
| melanoleucus deserticola]) | |
| Ring-necked duck (Aythya collaris) | Moved to Potential Concern List |
| Bufflehead (Bucephala albeola) | Moved to Potential Concern List |
| Merlin (Falco colum barius) | Moved to Potential Concern List |
| Arctic peregrine falcon (Falco peregrinus tundrius) | Moved to Potential Concern List |
| Greater prairie-chicken (Tympanuchus cupido) | Moved to Potential Concern List |
| Northern bobwhite (Native populations only) | Moved to Potential Concern List |
| (Colinus virginianus) | |
| Piping plover (Charadrius melodus) | Moved to Potential Concern List |
| Upland sandpiper (Bartramia longicauda) | Dropped from Concern List |
| Wilson's phalarope (Phalaropus tricolor) | Dropped from Concern List |
| Ring-billed gull (Breeding colonies only) (Larus | Moved to Potential Concern List |
| delawarensis) | |
| Herring gull (Breeding colonies only) (Larus argentatus) | Moved to Potential Concern List |
| Common tern (Sterna hirundo) | Moved to Potential Concern List |
| Barn owl (Tyto alba) | Moved to Potential Concern List |
| Western screech-owl (Otus kennicottii) | Moved to Potential Concern List |
| Black-chinned hummingbird (Archilochus alexandri) | Dropped from Concern List |
| Rufous hummingbird (Selasphorus rufus) | Dropped from Concern List |
| Eastern phoebe (Sayornis phoebe) | Moved to Potential Concern List |
| Hammond's flycatcher (Empidonax hammondii) | Moved to Potential Concern List |
| Cassin's kingbird (Tyrannus vociferans) | Dropped from Concern List |
| Winter wren (Troglodytes troglodytes) | Moved to Potential Concern List |
| Golden-crowned kinglet (Regulus satrapa) | Moved to Potential Concern List |
| Eastern bluebird (Sialia sialis) | Moved to Potential Concern List |
| Townsend's warbler (Dendroica townsendi) | Moved to Potential Concern List |
| Rose-breasted grosbeak (Pheucticus ludovicianus) | Moved to Potential Concern List |
| Dickcissel (Spiza americana) | Moved to Potential Concern List |
| Cassin's sparrow (Aimophila cassinii) | Moved to Potential Concern List |
| Grasshopper sparrow (Ammodramus savannarum) | Moved to Potential Concern List |
| Brewer's sparrow (Spizella breweri) | Moved to Potential Concern List |
| | |

| Common Name | |
|--|-----------------------------------|
| | A |
| (Scientific Name) | Action Taken |
| White-winged junco (Junco hyemalis aikeni) | Moved to Potential Concern List |
| White-winged crossbill (Loxia leucoptera) | Moved to Potential Concern List |
| Dwarf shrew (Sorex nanus) | Moved to Potential Concern List |
| Eastern mole (Scalopus aquaticus) | Moved to Potential Concern List |
| Long-eared myotis (Myotis evotis) | Moved to Potential Concern List |
| Hoary bat (Lasiurus cinereus) | Moved to Potential Concern List |
| Eastern cottontail (Sylvilagus floridanus) | Moved to Potential Concern List |
| Spotted ground squirrel (Spermophilus spilosma) | Dropped from Concern List |
| Northern flying squirrel (Black Hills population) | Moved to Potential Concern List |
| (Glaucomys sabrinus (undescribed taxon)) | |
| Abert's squirrel (Sciurus aberti) | Moved to Potential Concern List |
| Plains pocket mouse (Perognathus flavescens) | Dropped from Concern List |
| Silky pocket mouse (Perognathus flavus) | Dropped from Concern List |
| Great Basin pocket mouse (Perognathus parvus) | Dropped from Concern List |
| Hispid pocket mouse (Chaetodipus hispidus [Perognathus hispidus]) | Dropped from Concern List |
| Plains harvest mouse (Reithrodontomys montanus) | Dropped from Concern List |
| White-footed mouse (Peromyscus leucopus) | Moved to Potential Concern List |
| Water vole (Bighorn Mountain population) (Microtus | Moved to Potential Concern List |
| richardsoni [Arvicola richardsoni] (undescribed taxon)) | Tho ved to I stellar Concern East |
| Common gray fox (Urocyon cinereoargenteus) | Moved to Potential Concern List |
| Ringtail (Bassariscus astutus) | Moved to Potential Concern List |
| American marten [Pine marten] (Bighorn Mountain population) (Martes americana (undescribed taxon)) | Moved to Potential Concern List |
| Least weasel (Mustela nivalis [Mustela rixosa]) | Moved to Potential Concern List |
| Plains (eastern) spotted skunk (Spilogale putorius | Moved to Potential Concern List |
| interruptua) | |
| Western spotted skunk (Spilogale gracilis [Spilogale | Dropped from Concern List |
| putorius]) | |
| Swainson's hawk (Buteo swainsoni) | Dropped from Watch List |
| Prairie falcon (Falco mexicanus) | Dropped from Watch List |
| Red-naped sapsucker (Sphyrapicus nuchalis) | Dropped from Watch List |
| Willow flycatcher (Empidonax traillii) | Dropped from Watch List |
| Cordilleran flycatcher [Western flycatcher] (Empidonax | Dropped from Watch List |
| occidentalis [Empidonax difficilis]) | |
| Marsh wren (Cist othorus palustris) | Dropped from Watch List |
| Lark bunting (Calamospiza melanocorys) | Dropped from Watch List |
| American marten [Pine marten] (statewide) (Martes | Dropped from Watch List |
| americana) | |

RARE ANIMAL OBSERVATION FORM WYNDD, Univ. Wyoming - Dept. 3381, Laramie, WY 82071

| SPECIES NAM | E | | | | | | | |
|-------------------|------------------------------------|--------------------|----------------------|------------------------|-----------------|--|--|--|
| Date of observa | ation | Observers | | | | | | |
| Obs. affiliation/ | contact info | | | | | | | |
| LOCATION: (in | f possible, please attach o | copy of map with | location noted) | | | | | |
| USGS quad nam | USGS quad name County | | | | | | | |
| Township | /nship Range Sect | | ection | 1⁄4 of sec | 1⁄4 of 1⁄4 sec | | | |
| Latitude | Longitude | Datum | UTM-N | UTM- | -E | | | |
| Description of l | location | | | | | | | |
| Jurisdiction (e.g | g., USFS District, BLM Re | esource Area) | | | | | | |
| BIOLOGY: | (males) | | adults (m) (f) | (m) | ubadults | | | |
| Life-history not | es | | | | | | | |
| Reproductive n | otes | | | | | | | |
| Behavioral note | es_ | | | | | | | |
| HABITAT: | Elevation | | | er | | | | |
| | te | | | | | | | |
| | urroundings | | | | | | | |
| | 3 | | | | | | | |
| | (e.g., pond, riffle, pool)_ | | | | | | | |
| DOCUMENTA | TION: Visual | Tracks | Scat | Song/call | Roadkill_ | | | |
| Other | | | | | | | | |
| Specimen collec | cted? (collector, coll. #, w | here located) | | | | | | |
| Photo taken? (p | photographer, photo #, wh | nere located) | | | | | | |
| Identification co | onfirmed?any sheets necessary to e | laborate or diagra | m any information | n associated with this | s observation)) | | | |

| DATE OF SURVET. / / OBSERVER(S). | DATE OF SURVEY: | / / | OBSERVER(S): | |
|----------------------------------|-----------------|-----|--------------|--|
|----------------------------------|-----------------|-----|--------------|--|

PLANT SPECIES OF CONCERN SURVEY FORM WYNDD, Univ. Wyoming - Dept. 3381, Laramie, WY 82071

| SPECIES NAME: | | COUNTY: | | | | |
|---|-------------------------------------|-----------------|--|--|--|--|
| LOCATION: (attach annotated topo map copies with population location or GIS electronic files if available USGS QUAD: ELEVATION RANGE: | | | | | | |
| | | | | | | |
| NATIONAL FOREST DISTR | ICT/BLM FIELD OFFICE/OTHER: | | | | | |
| • | fer to physiographic region, roads, | · | | | | |
| | | | | | | |
| SIZE (ACRES) AND DISTRI | B.: | | | | | |
| % FLOWERING | % FRUITING | % VEGETATIVE | | | | |
| VEGETATION TYPE/ DOMI | NANT SPECIES: | | | | | |
| ASSOCIATED SPECIES: _ | | | | | | |
| | | | | | | |
| | | | | | | |
| GENERAL SETTING: | | | | | | |
| | | | | | | |
| | | | | | | |
| ASPECT (S, SE, NNW, etc. |): % SLOPE: | SLOPE POSITION: | | | | |
| SLOPE SHAPE: | LIGHT EXPOSURE: | MOISTURE: | | | | |
| SOIL TEXTURE: | SOIL S | SERIES/GEOLOGY: | | | | |
| LAND USE/ SIGNS OF DIST | TURBANCE: | | | | | |
| PHOTOGRAPH TAKEN?: | VE | GETATION PLOT?: | | | | |
| SPECIMEN (COLLECTOR/ | COLL. #/REPOSITORY): | | | | | |
| COMMENTS: | | | | | | |
| | | | | | | |
| | | | | | | |

Please use the reverse side for elaborating or diagramming any of the information on this form