

ST163: Bufexamac CAS#: 2438-72-4

Patient Information

Your patch test results indicate that you have a contact allergy to bufexamac. This contact allergy may cause your skin to react when it is exposed to this substance although it may take several days for the symptoms to appear. Typical symptoms include redness, swelling, itching, and fluid-filled blisters.

Where is bufexamac found?

Bufexamac is an anti-inflammatory analgesic and is administered orally, topically, or rectally.

How can you avoid contact with bufexamac?

Avoid products that list any of the following names in the ingredients:

- 4-Butoxy-Nhydroxybenzeneacetamide
- 4-Butoxyphenylacetohydroxamic acid
- Acide p
 - butoxyphenylacethydroxamique
- Acide pbutoxyphenylacethydroxamique [French]
- Anderm
- BRN 2646848
- Bufessamac
- Bufessamac [DCIT]
- Bufexamac
- Bufexamaco
- Bufexamaco [INN-Spanish]
- Bufexamacum
- Bufexamacum [INN-Latin]

- Bufexamic acid
 - CP 1044
- CP 1044 J3
- Droxaryl
- EINECS 219-451-1
- Feximac
- Flogicid
- Flogocid N plastigel
- **J**3
- Malipuran
- Mofenar
- Norfemac
- Parfenac
- Parfenal
- p-Butoxyphenylacetohydroxamic acid

What are some products that may contain bufexamac? Medications:

- Creams
- Oral tablets

*For additional information about products that might contain **bufexamac**, go to the Household Product Database online (http:/householdproducts.nlm.nih.gov) at the United States National Library of Medicine. These lists are brief and provide just a few examples. They are not comprehensive. Product formulations also change frequently. Read product labels carefully and talk to your doctor if you have any questions. These are general guidelines. Talk to your doctor for more specific instructions.

