



Monograph

# Polychaetes of Greece: an updated and annotated checklist

Sarah Faulwetter<sup>‡</sup>, Nomiki Simboura<sup>§</sup>, Nikolaos Katsiaras<sup>§,||</sup>, Giorgos Chatzigeorgiou<sup>¶</sup>, Christos Arvanitidis<sup>¶</sup>

<sup>‡</sup> University of Patras, Department of Zoology, Section of Marine Biology, Patras, Greece

<sup>§</sup> Institute of Oceanography, Hellenic Centre for Marine Research, Anavyssos, Greece

| Department of Marine Sciences, University of the Aegean, Mytilini, Greece

<sup>¶</sup> Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine Research, Heraklion Crete, Greece

Corresponding author: Sarah Faulwetter ([sarahfaulwetter@gmail.com](mailto:sarahfaulwetter@gmail.com))

Academic editor: Vasilis Gerovasileiou

Received: 14 Sep 2017 | Accepted: 13 Dec 2017 | Published: 22 Dec 2017

Citation: Faulwetter S, Simboura N, Katsiaras N, Chatzigeorgiou G, Arvanitidis C (2017) Polychaetes of Greece: an updated and annotated checklist. Biodiversity Data Journal 5: e20997. <https://doi.org/10.3897/BDJ.5.e20997>

## Abstract

### Background

The last annotated checklist of marine polychaetes in Greece was published in 2001. Since then, global taxonomic progress, combined with many new species records for Greece, required a thorough review of the taxonomic, nomenclatural and biogeographic status of the national species list. This checklist revises the status of all extant polychaete species reported from the Greek Exclusive Economic Zone since 1832. The work was undertaken as part of the efforts on compiling a national species inventory (Greek Taxon Information System initiative) in the framework of the LifeWatchGreece Research Infrastructure.

### New information

This checklist comprises an updated and annotated inventory of polychaete species in Greek waters, compiled from literature reports, online databases, museum collections and unpublished datasets. The list provides information on 836 species-level taxa from Greece, of which 142 are considered questionable. An additional 84 species reported in the past

are currently considered absent from Greece; reasons for the exclusion of each species are given. Fourteen species are reported here for the first time from Greek waters. At least 52 species in the present list constitute in fact a complex of cryptic or pseudo-cryptic species. Forty-seven species are considered non-native to the area. In addition to the species-level taxa reported in this checklist, eleven genera have been recorded from Greece with no representatives identified to species level. One replacement name is introduced. For each species, a comprehensive bibliographic list of occurrence records in Greece and the synonyms used in these publications are provided as supplementary material. Where necessary, the taxonomic, nomenclatural or biogeographic status is discussed. Finally, the findings are discussed in the wider context of Mediterranean polychaete biogeography, taxonomic practice and worldwide research progress.

## Keywords

Eastern Mediterranean, Polychaeta, Annelida, inventory, biodiversity, new records, replacement name

## Introduction

### Polychaete research in Greece

The history of polychaete studies in Greece can be traced back to classical antiquity. Aristotle, although he tended to neglect polychaetes in his works on natural history (Voultsiadou et al. 2017) gave the first description of a “scolopendra thalassia” (referring to *Hermodice carunculata* Pallas, 1776): “*There are also marine scolopendras, similar to the terrestrial ones, but smaller in size; they live on the rocky substrates. Their colour is richer red and they bear more and much slender feet than the terrestrial scolopendras. They do not occur in great depths, as do the marine snakes*” ([History of Animals](#), 4<sup>th</sup> century B.C.). Modern systematic polychaete research in the country started in the 19<sup>th</sup> century (e.g. Brullé 1832, Forbes 1842, von Marenzeller 1893). Since then, most research has been directed towards the study of macrobenthic coastal soft bottom polychaete communities in an ecological context. To date, the only taxonomic work studying exclusively planktonic polychaetes in Greece is by Wesenberg-Lund (1939a), based on samples taken during the Danish Oceanographic Expedition of the research vessel “Thor” from a few stations in Greek waters (Mavraki et al. 2016). Hard substrates, meiobenthic communities and deep sea habitats are understudied in Greece (e.g. Tselepides et al. 2004) while no terrestrial or fresh water polychaetes are known from the country. Specialised taxonomic or faunistic works from Greece are rare and the large body of ecological studies comprises the most important source for species records and thus for the creation of national checklists.

## Rationale for the checklist

The first polychaete checklists covering the Greek waters were those of Arvanitidis (2000a) for the Aegean Sea (including the Turkish Aegean and the Sea of Marmara) under the broader framework of the entire Mediterranean inventory and of Simboura and Nicolaidou (2001) for all Greek seas; both checklists included only benthic species. In the framework of the European Union project PESI (Pan-European Species directories Infrastructures), Faulwetter (2010) combined the above lists and additional sources into an updated checklist, but this was simply a computer-based compilation of existing sources without further verification or annotation of the records.

After 2001, marine research, including benthic surveys (carried out by the [Hellenic Centre for Marine Research](#) (HCMR) and several universities) was intensified. Many new species records – often not only new for Greece, but also for the Mediterranean Sea – resulting from these studies, as well as the worldwide progress of taxonomic knowledge, required a thorough review of the taxonomic, nomenclatural and biogeographic status of many polychaete species recorded from Greece.

The construction of the Greek Taxonomic Information System (Bailly et al. 2016), which forms the taxonomic backbone of the [LifeWatchGreece](#) Research Infrastructure, provided the framework for the present study: the compilation of a comprehensive, annotated and updated checklist of the Greek polychaete fauna, expanding and updating former works (i.e. Arvanitidis 2000a, Simboura and Nicolaidou 2001). In addition, this publication complements recent checklists from other areas of the Mediterranean and Black Sea (e.g. Çınar 2005, Castelli et al. 2008, Ayari et al. 2009, Zaâbi et al. 2009, Martín and Gil 2010, Zaâbi et al. 2012, Çınar et al. 2014, Mikac 2015) and thus serves as an information resource for future polychaete studies in this biogeographic region.

## Taxonomic scope

Several recent studies indicate the paraphyletic status of polychaetes (e.g. Rouse and Pleijel 2001, de Oliveira Almeida et al. 2003, Weigert and Bleidorn 2016) and the higher classification of Annelida has been in a constant flux over the last years (e.g. Zrzavý et al. 2001, Helm et al. 2012, Weigert and Bleidorn 2016). As the phylogenetic position of many groups within the Annelida is not yet resolved, we chose to restrict the "polychaetes" to their traditional definition in this checklist and include those clades covered by Rouse and Pleijel (2001). We anticipate this approach to be more helpful to the majority of users of this checklist than following the most recent phylogeny which would exclude certain "traditional" polychaete families (see Weigert and Bleidorn 2016).

## Geographic scope and characteristics of the region

This checklist includes all polychaete taxa reported from the waters of the Greek Exclusive Economic Zone (EEZ) (as delimited by the [Marine Regions database](#), see also Bailly et al. 2016). It does not cover a biogeographic area but constitutes an inventory of the national

polychaete fauna. Thus, species reported from neighbouring countries but not (yet) from Greece are not included in this list, despite the fact that they are highly likely to occur in Greek water too.

Greece is located centrally in the Eastern Mediterranean Sea and thus lies in one of the world's marine biodiversity hotspots (Bianchi and Morri 2000a, Coll et al. 2010). Its coastline stretches over 18,000 km, showing a variety of coastal landforms, from cliffs and rocky shores to beaches, lagoons and river deltas and encompassing around 9,800 islands and rocky islets (Issaris and Salomidi 2012). The Greek EEZ comprises parts of three biogeographically distinct water bodies: the Aegean Sea in the east, the Ionian Sea in the west and the Libyan Sea in the south. The Aegean Sea exchanges waters in the north with the Black Sea through the Bosphorus Strait, the Marmara Sea and the Strait of Dardanelles. Extensive faulting and vertical tectonic movements have resulted in the fracturing of the Hellenic mountain chains and the creation of deeply incised embayments and semi-enclosed basins, whereas active underwater volcanoes and shallow-water hydrothermal vents can be found in several places in the Aegean Sea (Dando et al. 1999; see also Sini et al. (2017) for a detailed overview of the ecology of the Aegean Sea). The island of Crete forms the southern border of the Aegean and separates it from the Libyan Sea. The Libyan Sea is characterised by deep sea features such as trenches, extensive deep sea plateaus, hydrothermal vents, cold seeps and mud volcanoes. It is part of the Levantine basin and linked to the Red Sea via the Suez canal which forms the point of entry for Lessepsian migrant species. In the west, the Ionian Sea communicates with the Western Mediterranean and the Adriatic Sea. The Greek part of the Ionian Sea (East Ionian Sea) is characterised by the existence of the deep Hellenic Trench with depths exceeding 5,000 m – the deepest part of the Mediterranean Sea. Primary production in the Mediterranean decreases towards the eastern basin and thus the Greek waters are highly oligotrophic (SoHeIME 2005), characterised by a limited availability of nutrients, primarily phosphorus (Thingstad et al. 2005). However, within this oligotrophic environment relatively eutrophic areas do exist (e.g. the North-West Aegean is highly productive in terms of nutrients and biological production compared to the Ionian, the South Aegean and the Levantine basin (e.g. Lykousis et al. 2002)). Knowledge on the polychaete fauna in each of these regional seas is unequally distributed, as the overwhelming majority of benthic research (around 80% of studies) has been performed in the Aegean Sea, whereas the deep sea habitats characterising the Libyan Sea are almost completely unexplored.

## **Materials and methods**

### **Literature resources**

Existing checklists on Greek polychaetes were combined, complemented, verified and annotated with information on species occurrences in Greece, using over 260 bibliographic and online resources, covering almost 200 years of research (1832–2017). A comprehensive list of all references from Greece for each species is contained in Suppl. material 2. These sources include publications from scientific journals, book chapters, theses, conference proceedings and biogeographic and museum collection databases. In

addition, unpublished records generated in the framework of research projects conducted within the Hellenic Centre of Marine Research (HCMR) after 2001 were used to assess the occurrence status of many controversial and non-native species (Suppl. material 1). We deliberately chose to include grey literature and database records, as many of these have been used in secondary works and we prefer to draw attention to doubtful records rather than ignoring them. The additions to the previous checklists comprise polychaete records from 486 sampling stations which were compiled from a) literature and data sources from after 2001 and b) literature and data sources from before 2001 if these had not been covered by previous checklists (Fig. 1). The majority of these stations were monitored regularly or occasionally. Samples were mainly collected from shallow soft-bottom habitats, though a few samples from pelagic, deep-sea and hard substrate habitats are also included.




Figure 1. [doi](#)

Sampling stations from which polychaetes were recorded, compiled from literature sources and databases which were not included in previous checklists.

The final checklist was submitted to the [World Register of Marine Species](#) and the [Pan-European Species directories Infrastructure](#) to replace the latest version by Faulwetter (2010).

### How to read this checklist

The nomenclature used here follows the [World Register of Marine Species](#) (WoRMS) and the most recent taxonomic literature. Taxa are arranged alphabetically in hierarchical order (genera and species within families); suprafamilial classifications are omitted. The inclusion of species follows a conservative approach: species with a questionable presence in Greek waters are included in the list and clearly indicated as such; we excluded only species that are either considered taxonomically indeterminable according to the most recent taxonomic works or where Greek records were shown to be misidentifications. These are presented separately at the end of the manuscript.

For each species, one or more of the following types of information may be provided:

## 1. Nomenclature

*If no information is provided in the "Nomenclature" field:* Species has only been recorded under its currently accepted name.

*If information is provided in the "Nomenclature" field:* Not a comprehensive list of synonyms (for this, the reader is advised to consult [WoRMS](#)), but all names under which the species has been reported from Greece (respective literature resources for each name in Suppl. material 2). Subjective and objective synonyms (subjective: two different type specimens exist and original descriptions exist, synonymy is based on expert opinion; objective: different names exist as a result of assigning a species to a different genus or rank but are based on the same type specimen) as well as different rank assignments (e.g. subgeneric vs. generic level) are listed. Misspellings and spelling variations (e.g. species epithets ending in -a vs. -us) are excluded. Misidentifications (i.e. specific specimens which were in the past identified as belonging to another species but then re-examined) are also omitted to avoid confusion; instead, a remark is given in the Notes field and the record is annotated in Suppl. material 2.

## 2. Native Status

*If no information is provided in the "Native status" field:* Species is considered native to the Mediterranean.

*If information is provided in the "Native status" field:* Contains information on the native status of the species in Greece, following the terminology and definitions by Zenetos et al. (2011):

- **non-native (established):** Species with at least two independent published records from Greece, separated in time and space;
- **non-native (casual):** Reported only once in the scientific or grey literature from Greece;
- **non-native (questionable):** Non-native species reported from Greece whose presence in the country is questionable and needs to be confirmed by an expert, or which have an unresolved taxonomic status. In this paper, doubtful records of species with an alien origin are not indicated as non-native if they have never been reported before as alien from the Mediterranean, whereas species previously reported as alien from the Mediterranean Sea but shown here to be questionable are retained as "non-native (questionable)" to highlight their status for further assessment by experts;
- **cryptogenic:** Species of unknown origin that could be either native or introduced, clear evidence for either origin being absent.

### 3. Notes

*If no information is provided in the "Notes" field:* Species has an uncontroversial taxonomic and biogeographic status and regular records from Greece and the Mediterranean exist. A full reference list of all reports from Greece for each species can be found in Suppl. material 2.

*If information is provided in the "Notes" field:* May contain information on:

- the taxonomic history, nomenclatural or occurrence status where these are deemed useful for assessing the taxonomic or occurrence status of the species in Greece.
- literature references reporting the species from Greece and additional information on the species' distribution: only in cases where the species has been reported fewer than three times from Greece (excluding checklists and other secondary works).
- whether the species has originally been described from the Mediterranean.

*Distribution information:*

Where statements are made on the occurrence (or absence) of the species in other areas of the Mediterranean, this information was derived from a combined assessment of:

- the checklist of Mediterranean polychaetes by Martín and Gil (2010);
- recent regional checklists of polychaetes in the Mediterranean (e.g. Çınar 2005, Castelli et al. 2008, Ayari et al. 2009, Zaâbi et al. 2009, Çınar et al. 2014, Mikac 2015);
- extensive unpublished notes by C. Arvanitidis covering the literature on Mediterranean polychaetes until ca. 1994;
- literature research.

Information on species distributions outside the Mediterranean are derived from a combined assessment of literature sources and distribution records in the [Ocean Biogeographic Information System \(OBIS\)](#) (unless a specific source is cited). Distribution information is not intended to list all known distributions of the species but rather to provide an estimate of the species' likelihood to occur in the Mediterranean.

*Questionable species:*

Species are indicated as "questionable" if one or more of the following criteria are met:

- the species concept is not clearly defined or outdated and in need of revision, rendering literature records highly uncertain;
- species is indicated as possibly absent from the Mediterranean in a recent taxonomic work (preferably a review) based on examination of Mediterranean specimens;
- species is in the Mediterranean reported exclusively (or almost exclusively) from Greece and has otherwise a distinctly different distribution range;

- species has been reported only once in grey literature before 1993 (chosen as a cut-off date, as thorough investigations of the Greek polychaete fauna followed afterwards)

Species reported as "cf." in primary records are not indicated as questionable, as the modifier "cf." already implies uncertainties in the identification and often a possible deviation from the nominal species.

*Species complexes:*

Species known to constitute a complex of cryptic (morphologically indistinguishable but genetically different) or pseudo-cryptic (morphologically very similar) species are here treated as *sensu lato*, following *Nygren and Pleijel (2011)*. Thus, clarifications pending through molecular analyses, we assume the presence of at least one species of the complex in Greece. The keyword "Species complex" in the Notes field indicates that the name has been applied to a species complex as defined above.

## **Acoetidae Kinberg, 1856**

### ***Eupanthalis kinbergi* McIntosh, 1876**

**Notes:** Many records of *Eupanthalis kinbergi* in the Mediterranean actually belong to *Euarche tubifex* Ehlers, 1887, including the description by Fauvel (1923) (Pettibone 1989). Arvanitidis (2000a) reports both *Euarche tubifex* and *Eupanthalis kinbergi* in his checklist of Aegean polychaetes, re-assigning all records of *Eupanthalis kinbergi* that were probably identified using the key by Fauvel (1923) to *Euarche tubifex*, but no primary records of *Euarche tubifex* from Greece exist.

### ***Panthalis oerstedii* Kinberg, 1856**

### ***Polyodontes frons* Hartman, 1939**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Satsmadjis 1983, Bogdanos and Satsmadjis 1987, NCMR 1989). Commonly distributed in the Greater Caribbean Region.

### ***Polyodontes maxillosus* (Ranzani, 1817)**

**Notes:** Type locality: Mediterranean.



## Acrocirridae Banse, 1969

### *Acrocirrus frontifilis* (Grube, 1860)

Notes: Type locality: Mediterranean (Croatia).

### *Flabelligena mediterranea* (Kolmer, 1985)

Nomenclature: *Flabelligella mediterranea* Kolmer, 1985

Notes: Originally described from the Hellenic Trench (Matapan Deep; 35°49'48"N, 22°20'42"E; 4690 m depth), no other records from Greece.

### *Macrochaeta clavicornis* (M. Sars, 1835)

## Ampharetidae Malmgren, 1866

### *Adercodon pleijeli* Mackie, 1994

Notes: Reported from Greece by Simboura and Zenetos (2005). Type locality: Mediterranean (Banyuls-sur-Mer, France).

### *Alkmaria romijni* Horst, 1919

Notes: Questionable status. In the Mediterranean, only reported from Greece (Kisseleva 1961). Record doubted by Arvanitidis (1994) and Çinar et al. (2014) based on the species' ecology: *Alkmaria romijni* is typically a brackish water species but has been reported from a depth of 65–100 m in the open waters of the Aegean Sea.

### *Amage adpersa* (Grube, 1863)

Notes: Type locality: Mediterranean (Adriatic).

### *Ampharete acutifrons* (Grube, 1860)

Notes: Questionable status. Mediterranean records of *Ampharete acutifrons* in the Mediterranean are doubted by several authors (e.g. Parapar et al. 1993a, Mikac 2015) based on the species' boreal distribution. Specimens of *Ampharete acutifrons* from the Adriatic were shown to belong to *Ampharete lindstroemi* Hessle, 1917 (Mikac 2015), thus *Ampharete acutifrons* may actually be absent from the region. Jirkov (2001) considers *Ampharete acutifrons* to probably constitute a species complex, based on observed differences between Atlantic and Pacific species.

***Ampharete grubei* Malmgren, 1865**

**Notes:** *Ampharete grubei* had long been considered a junior synonym of *Ampharete acutifrons* (Grube, 1860) and was reinstated by Jirkov (2001). Primary literature records from Greece under the name *Ampharete grubei* exist but were subsequently reported by Arvanitidis (2000a) and Simboura and Nicolaidou (2001) as *Ampharete acutifrons*.

***Ampharete octocirrata* (Sars, 1835)**

**Nomenclature:** *Sabellides octocirrata* (M. Sars, 1835)

***Amphicteis gunneri* (M. Sars, 1835)*****Anobothrus gracilis* (Malmgren, 1866)**

**Notes:** Reported from worldwide locations but probably restricted to circumpolar areas; specimens from other regions may belong to other species (Jirkov 2008).

***Auchenoplax crinita* Ehlers, 1887*****Lysippe labiata* Malmgren, 1866*****Melinna monoceroides* Fauvel, 1936*****Melinna palmata* Grube, 1870*****Neosabellides oceanica* (Fauvel, 1909)**

**Notes:** Questionable status. Few records from the Mediterranean outside Greece (e.g. Spain: Nebra et al. 2011; France: Guérin and Massé 1976) otherwise known from the European Atlantic coasts. Arvanitidis (1994) regards its presence in the Mediterranean as doubtful.

***Sosane sulcata* Malmgren, 1866*****Ushakovius enigmaticus* Laubier, 1973**

**Notes:** Originally described from the Hellenic Trench (Matapan Deep; 36°01'48"N, 22° 24'36"E; 3174 m depth), no other records from Greece. Considered endemic to the type locality (Paterson et al. 2009).

## Amphinomidae Lamarck, 1818

### *Chloeia candida* Kinberg, 1857

**Nomenclature:** *Chloeia viridis* Schmarda, 1861

**Notes:** Reported from Greece by Arvanitidis (1994). In the Mediterranean also reported from the western basin (Fauvel and Rullier 1959), otherwise distributed in the Caribbean and tropical Atlantic. *Chloeia viridis* is the name prevailing in literature, with *Chloeia candida* as a synonym (referred to synonymy by Hartman (1948), although Barroso and Paiva (2011) stress that this synonymy should be re-evaluated). The publication year of Kinberg's species is sometimes cited as 1910, but the species was in fact described in 1857 (Kinberg 1857) and thus the name *Chloeia candida* takes priority.

### *Chloeia venusta* Quatrefages, 1866

**Notes:** Type locality: Mediterranean (Sicily).

### *Chloenopsis atlantica* (McIntosh, 1885)

**Notes:** Böggemann (2009) examined the syntypes of *Chloenopsis atlantica* and postulated a possible synonymy of *Chloenopsis atlantica* with *Bathychloeia sibogae* Horst, 1910. However, Kudenov and Borda (2013), having studied the types of the monotypic genera *Chloenopsis* and *Bathychloeia*, consider both genera distinct.

### *Eurythoe complanata* (Pallas, 1766)

**Native status:** Non-native (casual)

**Notes:** Species complex. Reported from Greece by Chatzigeorgiou et al. (2016) based on a single specimen (identified using Day (1967) and Barroso et al. (2010)). *Eurythoe complanata* is a complex of cryptic and pseudo-cryptic species (Barroso et al. 2010, Arias et al. 2013a), containing two cryptic forms of *Eurythoe complanata* and the species *Eurythoe laevisetis* Fauvel, 1914, which differs from *Eurythoe complanata* sensu stricto by the absence of harpoon-shaped notochaetae (Arias et al. 2013a). In the Mediterranean, two of the species in this species complex occur: *Eurythoe laevisetis* in the central and western basin, and *Eurythoe complanata* in the eastern basin and the Alboran Sea. Both species are native to the Caribbean and tropical Atlantic and non-native to the Mediterranean, but their exact pathways of introduction to the Mediterranean and thus the identity of the species occurring in the eastern basin are yet unknown (Arias et al. 2013a).

### ***Hermodice carunculata* (Pallas, 1766)**

**Nomenclature:** *Amphinome savignyi* Brullé, 1832 | *Hermodice carunculata* (Pallas, 1766)

**Notes:** The identity of the circumtropical species *Hermodice carunculata* has been the subject of several investigations during the last years. Yáñez-Rivera and Salazar-Vallejo (2011), based on morphological investigations of fixed and living specimens, resurrected *Hermodice nigrolineata* Baird, 1868 for the Eastern Atlantic and Mediterranean populations of the species. Ahrens et al. (2013) found no genetic differences between *Hermodice carunculata* populations of the Caribbean and the Eastern Atlantic and Mediterranean and placed *Hermodice nigrolineata* again into synonymy with *Hermodice carunculata*. The findings of Chatzigeorgiou et al. (2014) concerning the phylogenetic relationships of the eastern Mediterranean populations of *Hermodice carunculata* support the results of Ahrens et al. (2013).

### ***Linopherus canariensis* Langerhans, 1881**

**Native status:** Non-native (casual).

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016). Atlantic species, considered alien to the Mediterranean (Çınar 2009). Known from Turkey and Cyprus (Çınar 2009), Italy (Cosentino and Giacobbe 2010) and Egypt (Dorgham et al. 2014).

### ***Notopygos megalops* McIntosh, 1885**

**Notes:** Questionable status. The Greek records of *Notopygos megalops* probably belong to a different species. *Notopygos megalops* was originally described based on a juvenile specimen from the Caribbean Sea and used to be considered a synonym of *Notopygos crinita* (Grube, 1855) before it was re-described by Yáñez-Rivera and Carrera-Parra (2012) from the Caribbean Sea based on adult characters. This re-description of *Notopygos megalops* differs from the description by Fauvel (1923) (which was probably used for the identification of the Greek specimens) in the following characters: Branchiae from chaetiger 6 (Yáñez-Rivera and Carrera-Parra 2012) instead of chaetiger 7 (Fauvel 1923); anterior branchiae with main stem and seven branchial filaments, in median chaetigers branching into four stems with five to seven filaments each (Yáñez-Rivera and Carrera-Parra 2012) vs. small tufts of 2–4 filaments anteriorly and only a single papilliform filament anteriorly Fauvel (1923). Yáñez-Rivera and Carrera-Parra (2012) define the start of the branchiae on chaetiger 6 as characteristic for the species; this seems to be invariable in juvenile individuals, too (McIntosh 1895). Thus, the species described by Fauvel (1923) and any specimens identified based on his description probably belong to a different species.

***Pareurythoe borealis* (M. Sars, 1862)**

**Notes:** Questionable status. Reported from Greece by Kisseleva (1983). Distributed in the boreal Atlantic, in the Mediterranean reported from the western basin (e.g. Jacquotte 1963, Bellan 1964a). The species' presence in Greece is doubtful, as it has not been reported from the Mediterranean for over three decades and is usually distributed in colder waters.

**Aphroditidae Malmgren, 1867*****Aphrodita aculeata* Linnaeus, 1758**

**Notes:** Specimens of *Aphrodita aculeata* from the Adriatic and Aegean may have been confused with *Aphrodita alta* Kinberg, 1865 or *Aphrodita perarmata* Roule, 1898 (Barnich and Fiege 2000b). In addition, the description of *Aphrodita aculeata* by Fauvel (1923) contains elements of both *Aphrodita aculeata* and *Aphrodita alta*, thus specimens identified using Fauvel's key may belong to either species.

***Aphrodita perarmata* Roule, 1898**

**Notes:** Questionable status. Reported from Greece by Marinov (1959), otherwise only known from the type locality in the North-East Atlantic. *Aphrodita perarmata*, *Aphrodita aculeata* Linnaeus, 1758 and *Aphrodita alta* Kinberg, 1865 are easily confused and all records require verification (Barnich and Fiege 2000b).

***Laetmonice filicornis* Kinberg, 1856**

**Notes:** Reported from Greece by Arvanitidis (2000a) based on a single specimen; in the Mediterranean also known from France (Bellan 1964a), Spain (Campoy 1982), Cyprus (Çınar 2005), Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015), otherwise distributed in the North Atlantic.

***Laetmonice hystrix* (Savigny in Lamarck, 1818)**

**Nomenclature:** *Hermione hystrix* (Savigny in Lamarck, 1818) | *Laetmonice hystrix* (Savigny in Lamarck, 1818)

**Notes:** Type locality: Mediterranean.

***Pontogenia chrysocoma* (Baird, 1865)**

**Notes:** Type locality: Mediterranean.

## Arenicolidae Johnston, 1835

### *Abarenicola claparedi* (Levinsen, 1884)

**Nomenclature:** *Abarenicola claparedi* (Levinsen, 1884) | *Arenicola claparedi* Levinsen, 1884

**Notes:** Type locality: Mediterranean.

### *Arenicola marina* (Linnaeus, 1758)

**Notes:** Questionable status. Reported from Greece by Antoniadou et al. (2006) from rocky shores. The record is considered questionable on the basis of habitat and geographic distribution (the species occurs in tidal mudflats of the European Atlantic coast and North Sea).

### *Branchiomaldane vincenti* Langerhans, 1881

**Notes:** Reported from Greece by Arvanitidis (1994) and Chatzigeorgiou et al. (2016). In the Mediterranean also known from Israel (Ben-Eliahu 1976b), Tunisia (Cantone et al. 1978), Spain (Alós 1990), Italy (Castelli et al. 2008), Egypt (Abd-Elnaby 2009), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed in the East Atlantic (Gravina 1991).

## Asteriomyzostomidae Jägersten, 1940

### *Asteriomyzostomum asteriae* (Marenzeller, 1895)

**Nomenclature:** *Myzostoma asteriae* Marenzeller, 1895

**Notes:** Originally described from Greece as parasitic on the starfish species *Sclerasterias richardi* (Perrier in Milne-Edwards 1882) and *Sclerasterias neglecta* (Perrier, 1891) collected near Santorini, Kythira and Samos; no other records from Greece. Almost nothing is known of the species' distribution; the extensive description of its anatomy by Stummer-Traunfels (1903) is based on specimens provided by Marenzeller, presumably from the same collection as the type material.

## Capitellidae Grube, 1862

### *Capitella capitata* (Fabricius, 1780)

**Nomenclature:** *Capitella capitata* (Fabricius, 1780) | *Capitella capitata capitata* (Fabricius, 1780)

**Notes:** Questionable status. Species complex. Several sibling species have been separated from *Capitella capitata* sensu lato worldwide since Grassle and Grassle (1976) showed it to constitute a complex of species. Blake (2009) restricts *Capitella capitata* sensu stricto to Arctic areas and considers Mediterranean specimens, including the description by Fauvel (1927), to show significant differences to *Capitella capitata* and thus to belong to other species. In Greece, there is evidence so far of at least another sibling species population (Gamenick et al. 1998). However, because of the high morphological similarity of the various sibling species, they cannot be reliably distinguished in routine investigations (Blake 2000). See also notes under *Capitella teleta* Blake, Grassle & Eckelbarger, 2009.

### *Capitella giardi* (Mesnil, 1897)

**Nomenclature:** *Capitella giardi* (Mesnil, 1897) | *Capitellides giardi* Mesnil, 1897

### *Capitella minima* Langerhans, 1881

**Nomenclature:** *Capitella minima* Langerhans, 1881 | *Capitomastus minimus* (Langerhans, 1881)

### *Capitella teleta* Blake, Grassle & Eckelbarger, 2009

**Notes:** Reported from Greece by Maidanou et al. (2017). *Capitella teleta* is one of the sibling species which used to be referred to as *Capitella* sp. I in laboratory and genetic studies (Blake et al. 2009). In the Mediterranean also known from France (Blake et al. 2009) and Turkey (Çınar et al. 2017). Present in the adjacent Sea of Marmara (Çınar et al. 2015) and Black Sea (Kurt-Şahin et al. 2017). Otherwise distributed in the North Atlantic and Pacific Ocean.

### *Capitellethus dispar* (Ehlers, 1907)

**Notes:** Questionable status. Listed by Arvanitidis (2000a), subsequently included in the checklist by Simboura and Nicolaidou (2001) but in fact based on a record from the Turkish Aegean, which is considered questionable by Çınar et al. (2014). The actual first record of the species from Greece is by Akoumianaki (2004). However, this species can be confused with several species of *Notomastus* Sars, 1851, based only on the thoracic chaetal formula which was traditionally used for identification. Green (2002)

clarified some inaccuracies in previous descriptions of *Capitellethus* and used the teeth pattern in hooks as a reliable differentiating character between these two genera. If the presence of this species in Greece can be confirmed, it should be regarded as a non-native species. Originally described from New Zealand.

***Dasybranchus caducus* (Grube, 1846)**

**Notes:** Type locality: Mediterranean.

***Dasybranchus gajolae* Eisig, 1887**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Decamastus gracilis* Hartman, 1963**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Akoumianaki 2004). Distributed in the Eastern Pacific, the Greater Caribbean and questionably in the Red Sea (Wehe and Fiege 2002).

***Heteromastus filiformis* (Claparède, 1864)**

**Notes:** Frequently reported from Greece and other parts of the Mediterranean; type locality: Mediterranean (Port-Vendres, France). Green (2002) clarified confusions that had arisen from typographic errors in the genus designation made by Hutchings and Rainer (1981). However, a number of problems concerning the correct identification of the species still exist. Juvenile specimens can be confused with adults of *Mediomastus* species, since chaetigers four and five initially carry hooks which are replaced by capillaries with age (Capaccioni-Azzati and El-Haddad 2015). In addition, different opinions regarding the presence and location of abdominal gills exist in literature (Capaccioni-Azzati and El-Haddad 2015).

***Leiocapitella dollfusi* (Fauvel, 1936)**

**Nomenclature:** *Leiocapitella dollfusi* (Fauvel, 1936) | *Leiocapitella glabra* Hartman, 1947

**Notes:** Bellan (1964a) proposed the synonymy of *Leiocapitella glabra* (type locality California) with *Leiocapitella dollfusi* (originally described from the Moroccan coasts) based on the overlapping variability in the chaetal formula of the two species. Ben-Eliahu and Fiege (1995), probably unaware of Bellan (1964a), proposed a possible synonymy of *Leiocapitella dollfusi* with *Leiocapitella glabra* on the basis of the identical hook dentition, in addition to the chaetal formula (Capaccioni-Azzati and El-Haddad 2015). However, it is noteworthy that the presence of branchiae in posterior segments, a character reported by Fauvel (1936a) for *Leiocapitella dollfusi*, has not been confirmed in material previously identified as *Leiocapitella glabra* (Hartman 1947, Ben-



Eliahu and Fiege 1995), due to lack of posterior parts. In the present study, in agreement with Bellan (1964a), and Capaccioni-Azzati and El-Haddad (2015), we consider the older name *Leiocapitella dollfusi* as having priority over *Leiocapitella glabra*.

### ***Leiocapitellides analis* Hartmann-Schröder, 1960**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Akoumianaki (2004) and Neofitou et al. (2014). In the Mediterranean also known from Egypt (Abd-Elnaby 2009). Originally described from the Red Sea.

### ***Leiochrus alutaceus* Ehlers, 1908**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Akoumianaki 2004). Commonly distributed in the Pacific. Worldwide, records of this genus are rare and uncertain (e.g. Hilbig and Blake 2006, Probert et al. 2009). However, as species of *Leiochrides* can key out as *Leiochrus* in Fauchald (1977), the presence of the former genus in Greece is possible if the key by Fauchald has been used.

### ***Mastobranthus trinchessii* Eisig, 1887**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Mediomastus capensis* Day, 1961**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Maidanou et al. (2017). Originally from South Africa, in the Mediterranean also known from Italy (Castelli et al. 2008), Cyprus (Çınar 2005), Spain (Capaccioni-Azzati and El-Haddad 2015) and the Adriatic (Mikac 2015). Zenetos et al. (2005) and Zenetos et al. (2010) question the identification of the species in the Mediterranean, but without providing specific information.

### ***Mediomastus fragilis* Rasmussen, 1973**

### ***Neoheteromastus lineus* Hartman, 1960**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Akoumianaki 2004). Commonly distributed in the East Pacific.

***Neomediomastus glabrus* (Hartman, 1960)**

**Native status:** Non-native (casual)

**Notes:** In the Mediterranean only reported from Greece, by Dando et al. (1995) from shallow hydrothermal vents and by Simboura and Zenetos (2005) from a depth of 1250 m. Commonly distributed in deep sea habitats of the East Pacific, the Chukchi Sea and possibly the Bay of Biscay (Simboura and Zenetos 2005), thus the record by Dando et al. (1995) is considered questionable as it is from a shallow-water habitat.

***Neopseudocapitella brasiliensis* Rullier & Amoureux, 1979**

**Native status:** Non-native (questionable)

**Notes:** Questionable status. Reported from Greece by NCMR (1992), Simboura (1996) and Katsiaras and Simboura (2015). In the Mediterranean also known from Italy (Gravina and Somaschini 1990), Spain (Capaccioni-Azzati et al. 1992), Cyprus (Çınar 2005), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015). Originally described from Brazil. However, Capaccioni-Azzati and El-Haddad (2015) list some noteworthy inconsistencies amongst Mediterranean records and the original description regarding the presence of chaetae on the peristomium and the last setiger with capillaries present. Therefore, it is likely that the Mediterranean records belong to an undescribed species.

***Notomastus aberans* Day, 1957**

**Native status:** Non-native (established)

**Notes:** Natively distributed in East Africa and the Red Sea, first Mediterranean record from Greece (Harmelin 1968), nowadays distributed throughout the Mediterranean (Capaccioni-Azzati and El-Haddad 2015).

***Notomastus formianus* Eisig, 1887**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Notomastus latericeus* Sars, 1851**

**Notes:** Morphologically very similar to *Notomastus profundus* (Eisig, 1887), therefore some authors have suggested a possible synonymy between the two species (Capaccioni-Azzati and El-Haddad 2015; references within).

***Notomastus lineatus* Claparède, 1869**

**Notes:** Reported from Greece by Koukouras (1979) and Papazacharias (1991). Type locality: Mediterranean (Gulf of Naples).

***Notomastus profundus* (Eisig, 1887)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Peresiella clymenoides* Harmelin, 1968**

**Notes:** Originally described from specimens collected in Marseilles and the Greek islands of Crete and Santorini.

***Pseudocapitella incerta* Fauvel, 1913**

**Notes:** Type locality: Mediterranean (French Riviera).

***Pseudoleiocapitella fauveli* Harmelin, 1964**

**Notes:** Type locality: Mediterranean (Gulf of Lion).

**Chaetopteridae Audouin & Milne Edwards, 1833*****Chaetopterus variopedatus* (Renier, 1804)**

**Notes:** Type locality: Mediterranean.

***Mesochaetopterus sagittarius* (Claparède, 1870)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Phyllochaetopterus gracilis* Grube, 1863**

**Notes:** Reported from Greece by Pérès (1959). Type locality: Mediterranean (Adriatic).

***Phyllochaetopterus socialis* Claparède, 1869**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Spiochaetopterus costarum* (Claparède, 1869)**

**Nomenclature:** *Spiochaetopterus costarum* (Claparède, 1869) | *Telepsavus costarum* Claparède 1869

**Notes:** Species complex. *Spiochaetopterus costarum* was long believed to be a cosmopolitan species but in fact the name has been applied to a number of pseudo-cryptic species (e.g. Bhaud 1998, Bhaud et al. 2003). Mediterranean specimens are currently considered to belong to *Spiochaetopterus costarum* sensu stricto, as the type locality of the species is in the Gulf of Naples.

### ***Spiochaetopterus solitarius* (Rioja, 1917)**

**Nomenclature:** *Phyllochaetopterus solitarius* Rioja, 1917 | *Spiochaetopterus solitarius* (Rioja, 1917)

### ***Spiochaetopterus typicus* M. Sars, 1856**

**Notes:** Reported from Greece by Makra and Nicolaidou (2000) and Tselepides (1992). In the Mediterranean also reported from France (Guille and Laubier 1966) and the Adriatic (Mikac 2015). Bhaud (1998) restricts the species' distribution to the boreal Atlantic and considers it absent from the temperate waters of the Mediterranean, but the report by Tselepides is from deep waters (500 m and 700 m) off Crete, where environmental conditions could be more favourable for the species. Other boreal species have been shown to occur in deep Mediterranean waters (Fiege et al. 2000a).

## **Chrysopetalidae Ehlers, 1864**

### ***Arichlidon reyssi* (Katzmann, Laubier & Ramos, 1974)**

**Nomenclature:** *Arichlidon reyssi* (Katzmann, Laubier & Ramos, 1974) | *Bhawania reyssi* Katzmann, Laubier & Ramos, 1974

**Notes:** Type locality: Mediterranean (Adriatic).

### ***Chrysopetalum debile* (Grube, 1855)**

**Nomenclature:** *Chrysopetalum debile* (Grube, 1855) | *Paleanotus debile* (Grube, 1855)

**Notes:** Type locality: Mediterranean (Villfranche-sur-Mer, France).

***Hyalopale bispinosa* Perkins, 1985**

**Notes:** Recently identified from Greece (Watson and Chatzigeorgiou, submitted) based on two specimens from Crete (for details see [Record 1](#); [Record 2](#)).

***Paleanotus chrysolepis* Schmarda, 1861****Cirratulidae Carus, 1863*****Aphelochaeta filiformis* (Keferstein, 1862)**

**Nomenclature:** *Aphelochaeta filiformis* (Keferstein, 1862) | *Cirratulus filiformis* Keferstein, 1862

***Aphelochaeta marioni* (Saint-Joseph, 1894)**

**Nomenclature:** *Aphelochaeta marioni* (Saint-Joseph, 1894) | *Tharyx marioni* (Saint-Joseph, 1894)

***Aphelochaeta cf. monilaris* (Hartman, 1960)**

**Notes:** In the Mediterranean only reported from Greece (Simboura 1996, as *Aphelochaeta cf. monilaris*). Commonly distributed along the Pacific coast of North America. The Greek material differs from Hartman's original description mainly in the number of thoracic chaetigers (11 vs. 15 in the species described by Hartman 1960) and in the absence of eyespots (two eyespots in Hartman's material) and may belong to a different taxon (Simboura 1996).

***Aphelochaeta multibranchis* (Grube, 1863)**

**Nomenclature:** *Aphelochaeta multibranchis* (Grube, 1863) | *Tharyx multibranchis* (Grube, 1863)

**Notes:** Type locality: Mediterranean.

***Caulleriella alata* (Southern, 1914)**

**Nomenclature:** *Caulleriella alata* (Southern, 1914) | *Heterocirrus alatus* (Southern, 1914)

**Notes:** Mediterranean records could also comprise specimens of the recently described *Caulleriella mediterranea* Lezzi, 2017 which differs from *Caulleriella alata* mainly in having a biannulate instead of triannulate prostomium (Lezzi 2017).

### ***Caulleriella bioculata* (Keferstein, 1862)**

**Nomenclature:** *Caulleriella bioculata* (Keferstein, 1862) | *Heterocirrus bioculatus* (Keferstein, 1862)

### ***Caulleriella viridis* (Langerhans, 1881)**

**Notes:** Reported from Greece by Maidanou et al. (2017). The only other Mediterranean record is an unconfirmed record from Corsica (Ifremer 2016) and neither of the two Mediterranean reports is accompanied by a taxonomic description. Originally described from Madeira, little is known on its distribution range. If its presence is confirmed, the species should probably be considered non-native to the Mediterranean.

### ***Chaetozone caputesocis* (Saint-Joseph, 1894)**

**Nomenclature:** *Caulleriella caputesocis* (Saint-Joseph, 1894) | *Chaetozone caputesocis* (Saint-Joseph, 1894)

### ***Chaetozone corona* Berkeley & Berkeley, 1941**

**Nomenclature:** *Chaetozone corona* Berkeley & Berkeley, 1941

**Native status:** Non-native (established)

**Notes:** Reported from Greece by Simboura et al. (2010), found several times afterwards (HCMR, unpublished data). In the Mediterranean also known from Turkey (Çınar et al. 2014) and the Adriatic Sea (Munari et al. 2017). Chambers et al. (2011) doubt the validity of the Mediterranean records and the presence of the species in the Mediterranean (mainly on the basis of its Pacific distribution) but Le Garrec et al. (2016) present evidence of a wide distribution of the species in the Bay of Biscay, indicating that *Chaetozone corona* is an established alien species along the coasts of Europe and was probably introduced via shipping from its native distribution.

### ***Chaetozone gibber* Woodham & Chambers, 1994**

**Notes:** Reported from Greece by Simboura et al. (2010) and Katsiaras and Simboura (2015). In the Mediterranean also known from Tunisia (Zaâbi et al. 2012), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015); otherwise distributed around the British Isles and along the European Atlantic coasts. Chambers et al. (2011) remark that records of *Chaetozone gibber* from the Mediterranean could in fact belong to other species, as the Mediterranean environment differs significantly from conditions in the species' native distribution range.

***Chaetozone setosa* Malmgren, 1867**

**Notes:** Questionable status. Species complex. *Chaetozone setosa* is a complex of pseudo-cryptic species, with at least five species in northern Europe alone (Chambers and Woodham 2003). Several authors (e.g. Simboursa 1996, Çınar and Ergen 2007, Mikac 2015) doubt the validity of Mediterranean records, as the species has likely been confused with *Chaetozone gibber* Woodham & Chambers, 1994, *Chaetozone corona* Berkeley & Berkeley, 1941 or *Chaetozone carpenteri* McIntosh, 1911 and probably does not occur in the Mediterranean.

***Chaetozone zetlandica* McIntosh, 1911**

**Nomenclature:** *Caulleriella zetlandica* (McIntosh, 1911)

**Notes:** Chambers et al. (2011) remark that records of *Chaetozone zetlandica* might be misidentifications, as the Mediterranean environment differs significantly from conditions in the species' native distribution range in the North Atlantic.

***Cirratulus cirratus* (O. F. Müller, 1776)**

**Notes:** Species complex. Often considered cosmopolitan, but ecological, morphological and genetic differences have been found in specimens from different geographical locations (Petersen 1999, Carr et al. 2011, Weidhase et al. 2014).

***Cirriformia tentaculata* (Montagu, 1808)**

**Nomenclature:** *Audouinia tentaculata* (Montagu, 1808) | *Cirriformia tentaculata* (Montagu, 1808)

***Dodecaceria capensis* Day, 1961**

**Native status:** Non-native (established)

**Notes:** In the Mediterranean only reported from Greece (e.g. Bogdanos and Satsmadjjs 1983, Bogdanos and Satsmadjjs 1987, Simboursa 1996; full reference list in Suppl. material 2). Known from South Africa. Identification of several Greek specimens confirmed by P. H. Gibson (pers. comm. in Simboursa 1996).

***Dodecaceria concharum* Ørsted, 1843**

**Notes:** Questionable status. Castelli et al. (2008) and Mikac (2015) consider the presence of the species in the Mediterranean uncertain as it may have been confused with *Dodecaceria saxicola* (Grube, 1855), *Dodecaceria sextentaculata* (Delle Chiaje, 1822), *Dodecaceria fimbriata* (Verrill, 1880) or *Dodecaceria joubini* Gravier, 1905. Several Greek specimens were examined by P.H. Gibson and differ from the northern

European species; instead, different species (e.g. *Dodecaceria joubini*) could be present in Greece (P.H. Gibson, pers. comm. in Simboura 1996). The identification of *Dodecaceria* species depends on reproductive characteristics (Gibson 1978) but it is unknown which form was described by Ørsted. Thus, identity of *Dodecaceria concharum* is confused and Ørsted's original species is probably not valid due to an insufficient original description and loss of type material (Gibson 2015). Neotypes were designated by Gibson and Heppell (1995), but it is unclear whether these correspond to the original species of Ørsted.

### ***Dodecaceria saxicola* (Grube, 1855)**

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016). Type locality: Mediterranean (Villefranche-sur-Mer, France). As the species had long been treated as a synonym of *Dodecaceria concharum* Ørsted, 1843 it may be underreported.

### ***Kirkegaardia dorsobranchialis* (Kirkegaard, 1959)**

**Nomenclature:** *Monticellina dorsobranchialis* (Kirkegaard, 1959) | *Tharyx dorsobranchialis* (Kirkegaard, 1959)

**Notes:** A species reported from worldwide locations, partly caused by inaccurate descriptions in the past; therefore Blake (2016) considers the presence of *Kirkegaardia dorsobranchialis* in the Mediterranean uncertain and possibly restricted to West and South Africa. Many Greek records of *Kirkegaardia dorsobranchialis* belong in fact to *Kirkegaardia heterochaeta* (Laubier, 1961), as Blake (1991) had synonymised *Monticellina heterochaeta* with *Monticellina dorsobranchialis*. The distinction between the two species is difficult. *Kirkegaardia heterochaeta* has a longer peristomium than *Kirkegaardia dorsobranchialis* and possesses a mid-dorsal ridge within the dorsal channel which is absent in *Kirkegaardia dorsobranchialis*. The most reliable factor for their separation is the methyl green staining pattern (Blake 2016).

### ***Kirkegaardia heterochaeta* (Laubier, 1961)**

**Nomenclature:** *Monticellina heterochaeta* Laubier, 1961 | *Tharyx heterochaeta* (Laubier, 1961)

**Notes:** Possibly underreported, as the species was considered a synonym of *Kirkegaardia dorsobranchialis* (Kirkegaard, 1959) by Blake (1991) and reported under the latter name until Blake (1996) revoked the synonymy. Type locality: Mediterranean (France).

### ***Kirkegaardia tesselata* (Hartman, 1960)**

**Nomenclature:** *Monticellina tesselata* (Hartman, 1960)



**Notes:** Questionable status. Reported from Greece by Koulouri et al. (2015). In the Mediterranean also reported from Spain (Desbruyères et al. 1973) and Turkey (Çınar et al. 2014). Çınar (2005) reports morphological differences in chaetal features between specimens from Cyprus and the original description of *Kirkegaardia tessellata*, concluding that the material might belong to an undescribed species. Blake (2016), having examined a specimen of *Kirkegaardia* from France, refers this specimen and the literature references of *Monticellina tessellata* by Çınar et al. (2014) and Çınar (2005) to an undescribed *Kirkegaardia* species (*Kirkegaardia* sp. A). He restricts the distribution of *Kirkegaardia tessellata* to California. Greek specimens possibly do not belong to *Kirkegaardia tessellata* but the material would require re-examination to confirm its identity.

### ***Protocirrineris chrysoderma* (Claparède, 1868)**

**Nomenclature:** *Cirratulus chrysoderma* Claparède, 1868

**Notes:** Questionable status; species complex. Reported from Greece by NCMR (1989). Reported from worldwide locations but the species constitutes a complex of pseudo-cryptic species (Petersen 1991). Originally described from the Gulf of Naples and the only known species of the genus in the Mediterranean until the recent description of *Protocirrineris purgamentorum* Lezzi, Çınar & Giangrande, 2016. While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in almost three decades, therefore its presence in Greece is here considered questionable.

### ***Tharyx killariensis* (Southern, 1914)**

**Nomenclature:** *Cauleriella killariensis* (Southern, 1914) | *Heterocirrus killariensis* (Southern, 1914) | *Tharyx killariensis* (Southern, 1914)

### ***Timarete filigera* (Delle Chiaje, 1828)**

**Nomenclature:** *Cirriformia filigera* (Delle Chiaje, 1828) | *Timarete filigera* (Delle Chiaje, 1828)

**Notes:** Species complex. *Timarete filigera* has been recorded from worldwide locations, but many of these records probably belong to different species. Specimens from West Atlantic localities identified as *Timarete filigera* were re-examined morphologically and genetically by Magalhães et al. (2014) and shown to belong to undescribed species or to species previously treated as synonyms of *Timarete filigera*. Mediterranean specimens are currently considered to belong to *Timarete filigera* sensu stricto, as the type locality of the species is in the Gulf of Naples.

***Timarete punctata* (Grube, 1859)**

**Native status:** Non-native (casual)

**Notes:** Species complex. *Timarete punctata*, originally described from the Caribbean, constitutes a complex of cryptic species. Molecular analyses of specimens morphologically identified as *Timarete punctata* by Seixas et al. (2017) revealed that samples from the West Atlantic and the Pacific comprise two cryptic species, one of them widely distributed and exhibiting a low genetic diversity, the other restricted to a single location. The authors hypothesise that the low genetic diversity of the widely distributed cryptic species could be attributed to a recent human-mediated introduction at these sites. In the Mediterranean reported from Greece (Maidanou et al. 2017) and Turkey (Çınar et al. 2014).

**Cossuridae Day, 1963*****Cossura coasta* Kitamori, 1960**

**Native status:** Non-native (questionable)

**Notes:** Questionable status. Arvanitidis (1994) questions the presence of the species in the area, as the description of the specimen found by Bogdanos and Fredj (1983) shows differences both to the species described by Kitamori (1960) and to *Cossura soyeri* Laubier, 1964. Without re-examination of the Mediterranean specimens referred to *Cossura coasta*, the status of the species in the area remains doubtful.

***Cossura soyeri* Laubier, 1964**

**Notes:** Type locality: Mediterranean (Banyuls-sur-Mer, France)

**Ctenodrilidae Kennel, 1882*****Ctenodrilus serratus* (Schmidt, 1857)**

**Notes:** Reported from Greece by Makra and Nicolaidou (2000), also found by Papageorgiou et al. (2006) (unpublished dataset). Type locality: Mediterranean.

## Dorvilleidae Chamberlin, 1919

### *Dorvillea rubrovittata* (Grube, 1855)

**Nomenclature:** *Dorvillea rubrovittata* (Grube, 1855) | *Schistomeringos rubrovittata* (Grube, 1855) | *Staurocephalus rubrovittatus* Grube, 1855

**Notes:** Type locality: Mediterranean (Adriatic). Considered cosmopolitan. However, Jumars (1974) notes differences between populations and suggests the possibility of more than one species existing under the name of *Dorvillea rubrovittata*.

### *Dorvillea similis* (Crossland, 1924)

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Corsini-Foka et al. (2015). In the Mediterranean also known from Turkey (Çinar 2009), natively distributed in the Indo-Pacific Ocean and Red Sea.

### *Ophryotrocha labronica* La Greca & Bacci, 1961

**Notes:** Species complex. Reported from Greece by Åkesson (1984). Neotype from the Gulf of Naples. At least 15 sibling species have been reported under the *Ophryotrocha labronica* group worldwide (Åkesson and Paxton 2005), four of which have been recently described from areas including the Mediterranean by Paxton and Åkesson (2010): *Ophryotrocha japonica*, *Ophryotrocha macrovifera*, *Ophryotrocha robusta* and *Ophryotrocha rubra*.

### *Ophryotrocha puerilis* Claparède & Mecznirow, 1869

**Nomenclature:** *Ophryotrocha puerilis* Claparède & Mecznirow, 1869 | *Ophryotrocha puerilis puerilis* Claparède & Mecznirow, 1869

**Notes:** Species complex. Neotype from Genoa, Italy. Simbora and Nicolaidou (2001) report *Ophryotrocha puerilis* from Greece at species level, as it is reported on other parts of the Mediterranean as well (Castelli et al. 2008, Çinar et al. 2014, Núñez et al. 2013, Mikac 2015). However, Taboada et al. (2017) provide molecular evidence of at least two cryptic species under the name *Ophryotrocha puerilis* in the Mediterranean Sea.

### *Parougia caeca* (Webster & Benedict, 1884)

**Nomenclature:** *Schistomeringos caeca* (Webster & Benedict, 1884)

**Notes:** Reported from Greece by Eleftheriou et al. (1990) and Dando et al. (1995). In the Mediterranean also known from Italy (Gambi et al. 1998) and Turkey (Çınar et al. 2014); otherwise distributed in the North Atlantic and the North-East Pacific.

### ***Pettiboneia urciensis* Campoy & San Martín, 1980**

**Notes:** Type locality: Mediterranean (Águilas, Spain).

### ***Protodorvillea atlantica* (McIntosh, 1885)**

**Nomenclature:** *Protodorvillea atlantica* (McIntosh, 1885) | *Schistomeringos atlanticus* (McIntosh, 1885)

**Notes:** Questionable status. Reported from Greece by Vamvakopoulou (1991). In the Mediterranean also known from Tunisia (Zaâbi et al. 2009) and the Adriatic (Mikac 2015). Originally described from deep waters off the Azores (Atlantic); as the Greek record is from a lagoonal habitat, it must be considered questionable.

### ***Protodorvillea biarticulata* Day, 1963**

**Native status:** Non-native (established)

**Notes:** In the Mediterranean reported from Greece (Bogdanos and Satsmadjis 1983, Bogdanos and Diapoulis 1984, Simboura et al. 1995b) and Egypt (Abd-Elnaby and Gab-Alla 2007), otherwise distributed in the Indo-Pacific, South Africa and Brazil.

### ***Protodorvillea kefersteini* (McIntosh, 1869)**

**Nomenclature:** *Protodorvillea kefersteini* (McIntosh, 1869) | *Schistomeringos kefersteini* (McIntosh, 1869) | *Staurocephalus kefersteini* McIntosh, 1896

### ***Schistomeringos neglecta* (Fauvel, 1923)**

**Nomenclature:** *Dorvillea neglecta* (Fauvel, 1932) | *Schistomeringos neglecta* (Fauvel, 1923)

### ***Schistomeringos rudolphi* (Delle Chiaje, 1828)**

**Nomenclature:** *Dorvillea rudolphi* (Delle Chiaje, 1828) | *Schistomeringos rudolphi* (Delle Chiaje, 1828) | *Staurocephalus rudolphi* (Delle Chiaje, 1828)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

## Eulepethidae Chamberlin, 1919

### *Grubeulepis augeneri* Pettibone, 1969

**Notes:** One specimen from Greece in the collections of the Senckenberg Museum (SMF 15256, 35°52'58.2"N, 27°42'49.2"E, 115 m depth, coll. date 2005-09-05, det. D. Fiege). Type locality: Mediterranean (Adriatic).

### *Grubeulepis katzmanni* Pettibone, 1986

**Notes:** Questionable status. Reported from Greece by Simboura (1987) and Zenetos et al. (1997). Type locality: Mediterranean (Adriatic). The original description by Pettibone (1986) is based on juveniles and needs to be completed with adult characters; *Grubeulepis katzmanni* may turn out to be a juvenile of *Grubeulepis augeneri* Pettibone, 1969 (Pettibone 1986).

## Eunicidae Berthold, 1827

### *Eunice aphroditois* (Pallas, 1788)

**Notes:** Questionable status. Fauvel (1923) recorded *Eunice roussaei* Quatrefages, 1866 in the Adriatic Sea, but the species was later synonymised with *Eunice aphroditois* by Hartman (1944). Subsequently, several records of large-sized eunicids in the Mediterranean were assigned to the latter. After the resurrection of *Eunice roussaei* by Fauchald (1992), many authors suggested that literature records of *Eunice aphroditois* in the Mediterranean actually belong to *Eunice roussaei* (Parapar and Harto 2001, Zanol and Bettoso 2006, Çınar et al. 2014, Mikac 2015). Two specimens of *Eunice aphroditois* from the collections of the Aristotle University of Thessaloniki were re-examined by the first author and identified as *Eunice roussaei* (Faulwetter et al. 2013); it is likely that all Greek records belong to *Eunice roussaei*.

### *Eunice floridana* (Portalès, 1867)

**Native status:** Non-native (questionable)

**Notes:** Questionable status. Reported from Greece by Kisseleva (1983). In the Mediterranean also known from the western basin (e.g. Bellan 1959, Fredj 1974), Cyprus (Argyrou et al. 1999) and the Adriatic (Casellato and Stefanon 2008). Otherwise distributed in the Greater Caribbean (Fauchald et al. 2009). Commonly occurs in coralligenous habitats (Casellato and Stefanon 2008), which are little studied in Greece. However, no clear description based on Mediterranean material exists. Fauvel (1923), considers *Leodice gunneri* (Storm, 1880) a synonym of *Eunice floridana*. Day (1967) lists *Eunice floridana* sensu Fauvel (1923) as a synonym of *Eunice norvegica*

(Linnaeus, 1767) and Fauchald (1992) considers *Leodice gunneri* a synonym of the latter. Clear differentiating characters between *Eunice floridana* and *Eunice norvegica* can be found in Fauchald (1992) and Carrera-Parra and Salazar-Vallejo (1998) (which were, however, not published at the time the species was recorded from Greece).

### ***Eunice norvegica* (Linnaeus, 1767)**

**Notes:** Questionable status. Reported from Greece by NCMR (1989). In the Mediterranean also known from the western basin (e.g. Danovaro et al. 2010, Taviani et al. 2015), Tunisia (Ayari et al. 2009), the Italian coasts of the Ionian Sea (e.g. Tursi et al. 2004, Vertino et al. 2010) and the Adriatic (D'Onghia et al. 2015), otherwise distributed in the North Atlantic. Often associated with deep-water corals (Danovaro et al. 2010), which are little studied in Greece. The species has so far only been reported in grey literature and has not been found in almost three decades, therefore its presence in Greece is pending confirmation. See also notes under *Eunice floridana* on the confusion regarding the species concept in early works and keys.

### ***Eunice pennata* (Müller, 1776)**

### ***Eunice purpurea* Grube, 1866**

**Notes:** Reported from Greece by Arvanitidis (1994) and Kitsos (2003). Type locality: Mediterranean (Adriatic). Arvanitidis (1994) notes some differences in the individuals he examined compared to the description provided by Fauchald (1992). Fauvel (1917) regards *Eunice purpurea* as junior synonym of *Eunice roussaei* Quatrefages, 1866 and the species has since then been part of the confusion concerning large eunicids in the Mediterranean (see remarks under *Eunice aphroditois* Pallas, 1788). Fauchald (1992) considers it a valid species by investigating material from the Adriatic and Zanol and Bettoso (2006) highlight specific differences that do not occur in *Eunice roussaei*.

### ***Eunice roussaei* Quatrefages, 1866**

**Notes:** Reported from Greece by Kisseleva (1983). Probably underreported, as records of *Eunice aphroditois* (Pallas, 1788) from Greece likely belong to *Eunice roussaei* (see notes under *Eunice aphroditois*).

### ***Eunice schizobranchia* Claparède, 1870**

**Notes:** Questionable status. Reported from Greece by NCMR (1992). Type locality: Mediterranean (Gulf of Naples). Commonly occurs in coralligenous habitats (Casellato and Stefanon 2008), which are little studied in Greece. While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over two decades, therefore its presence in Greece is here considered questionable.

***Eunice vittata* (Delle Chiaje, 1828)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Leodice harassii* (Audouin & Milne Edwards, 1833)**

**Nomenclature:** *Eunice harassii* Audouin & Milne Edwards, 1833

**Notes:** Records of *Leodice harassii* could include specimens of *Eunice rubrocincta* Ehlers, 1868, a Mediterranean species which Fauvel (1923) considered synonymous with *Leodice harassii*. *Leodice rubrocincta* was resurrected by Fauchald (1992) and is included as a valid species in the key by Carrera-Parra and Salazar-Vallejo (1998).

***Leodice torquata* (Quatrefages, 1866)**

**Nomenclature:** *Eunice torquata* Quatrefages, 1866 | *Leodice torquata* (Quatrefages, 1866)

**Notes:** The species resembles *Leodice laurillardii* (Quatrefages, 1866) in a number of diagnostic characters. The latter was synonymised with *Leodice torquata* by Pruvot and Racovitza (1895); this was accepted by Fauvel (1923) and later authors. However, Fauchald (1992) resurrected *Leodice laurillardii* as a valid species and Arias et al. (2015) re-described the species based on type material and new specimens from the Mediterranean. They suggest that *Leodice laurillardii* could be much more widespread than previously thought but was possibly being overlooked in the Mediterranean due to the confusion with *Leodice torquata*.

***Lysidice collaris* Grube, 1870**

**Native status:** Non-native (questionable)

**Notes:** Questionable status. Frequently reported from Greece and other parts of the Mediterranean (Martín 1987, Sardá 1991, Giangrande et al. 2003, Mikac 2015). However, Çınar (2005) and Kurt Şahin and Çınar (2009) suggest that the species closely resembles the native species *Lysidice margaritacea* Claparède, 1868 and question the presence of *Lysidice collaris* in the Eastern Mediterranean. Type locality in the Red Sea, but reported from circumtropical locations.

***Lysidice ninetta* Audouin & Milne Edwards, 1833**

**Notes:** Species complex. Frequently reported from Greece. However, Iannotta et al. (2006) and Iannotta et al. (2009) investigated molecular markers and provided evidence of sibling species within the *Lysidice ninetta* complex. Two morphotypes (dark and light), differentiated by colour patterns in anterior body and acicular chaetae, were identified. Although the dark type matches well the original description, it is still

unresolved whether one or both morphotypes are new species or belong to previously described species.

### ***Lysidice unicornis* (Grube, 1840)**

**Nomenclature:** *Lysidice unicornis* (Grube, 1840) | *Nematonereis unicornis* (Grube, 1840)

**Notes:** The genus *Nematonereis* Schmarda, 1860 was synonymised with *Lysidice* Lamarck, 1818 by Zanol et al. (2013). Type locality: Mediterranean (Adriatic).

### ***Marphysa adenensis* Gravier, 1900**

**Native status:** Cryptogenic

**Notes:** In the Mediterranean only reported from Greece (Katsiaras et al. 2014) from *Posidonia oceanica* meadows, a little studied habitat in Greece. Distributed in the Indo-Pacific. The authors note that data are currently insufficient to draw conclusions on the species' native or alien distribution.

### ***Marphysa bellii* (Audouin & Milne Edwards, 1833)**

### ***Marphysa cinari* Kurt-Şahin, 2014**

**Notes:** Several Greek specimens reported by Simboura et al. (2010) as *Marphysa disjuncta* Hartman, 1961 were re-examined by Kurt Şahin (2014) and assigned to *Marphysa cinari*. Type locality: Sea of Marmara.

### ***Marphysa fallax* Marion & Bobretzky, 1875**

**Notes:** Type locality: Mediterranean (Marseille).

### ***Marphysa kinbergi* McIntosh, 1910**

### ***Marphysa sanguinea* (Montagu, 1813)**

**Notes:** Species complex. Considered cosmopolitan in the past, but *Marphysa sanguinea* actually comprises a complex of numerous pseudo-cryptic species (Zanol et al. 2016). Hutchings and Karageorgopoulos (2003) designated a neotype based on pectinate chaetae distribution instead of branchiae and Hutchings et al. (2012) encourage re-examination of all material outside the English Channel and North Sea. Recent studies have described several new species similar to *Marphysa sanguinea* or resurrected species from synonymy with the latter (Lewis and Karageorgopoulos 2008, Molina-Acevedo and Carrera-Parra 2015, Zanol et al. 2016, Liu et al. 2017). Wijnhoven



and Dekker (2010) question the presence of *Marphysa sanguinea* in the Mediterranean Sea and encourage comparisons of specimens with the neotype.

### ***Palola siciliensis* (Grube, 1840)**

**Nomenclature:** *Eunice siciliensis* Grube, 1840 | *Palola siciliensis* (Grube, 1840)

**Notes:** Type locality: Mediterranean (Palermo, Sicily, Italy).

## **Euphrosinidae Williams, 1852**

### ***Euphrosine foliosa* Audouin & Milne Edwards, 1833**

### ***Euphrosine myrtosa* Savigny in Lamarck, 1818**

**Notes:** Questionable status. Reported from Greece by Bogdanos and Satsmadjjs (1983). In the Mediterranean also reported from the Levantine Basin (Ben-Eliahu 1995) and questionably from the Adriatic (Mikac 2015), otherwise distributed in the Indo-Pacific. It might either be an overlooked alien species in the area or a misidentification.

## **Fabriciidae Rioja, 1923**

### ***Fabricia stellaris stellaris* (Müller, 1774)**

**Nomenclature:** *Fabricia sabella* (Ehrenberg, 1836) | *Fabricia stellaris stellaris* (O. F. Müller, 1774) |

### ***Manayunkia aestuarina* (Bourne, 1883)**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Kisseleva 1961, Diapoulis and Bogdanos 1983). Both Greek records are from a fully marine environment and therefore considered questionable by Arvanitidis (1994), as the species usually occurs in brackish waters. However, Harris (1970) reports the species from non-brackish environments in the British Isles and assumes that it can tolerate full salinity at least during certain stages of its life cycle. Based on the geographic range of *Manayunkia aestuarina*, the Greek records must nevertheless be considered uncertain. Undescribed species of *Manayunkia* exist in Greece (Chintiroglou et al. 2004) and the genus needs further study in the region.

### ***Novafabricia posidoniae* Licciano & Giangrande, 2006**

**Notes:** New record for Greece. Eleven specimens, Lesvos Island, 38°59'54"N, 26°32'26"E, 6 m depth, *Posidonia oceanica* meadow, collected in the framework of the PhD of N. Katsiaras. Literature used for identification: Licciano and Giangrande (2006). Type locality: Mediterranean (Italy).

### ***Pseudofabricia aberrans* Cantone, 1972**

**Notes:** Reported from Greece by Simboura and Zenetos (2005). Type locality: Mediterranean (Sicily).

### ***Pseudofabriciola analis* Fitzhugh, Giangrande & Simboura, 1994**

**Notes:** Specimens of *Fabricia filamentosa* Day 1963 from Greece (Simboura 1990) and the Adriatic (Giangrande and Castelli 1986) were re-examined by Fitzhugh et al. (1994) who described two new species based on this material: *Pseudofabriciola analis* from the Adriatic and *Pseudofabriciola longipyga* Fitzhugh, Giangrande & Simboura, 1994 from Greece. *Pseudofabriciola analis* was subsequently reported from Greece by Simboura (1996).

### ***Pseudofabriciola longipyga* Fitzhugh, Giangrande & Simboura, 1994**

**Notes:** Originally described from Greece by Fitzhugh et al. (1994). See remarks under *Pseudofabriciola analis* Fitzhugh, Giangrande & Simboura, 1994.

## **Flabelligeridae de Saint-Joseph, 1894**

### ***Bradabyssa villosa* (Rathke, 1843)**

**Nomenclature:** *Brada villosa* (Rathke, 1843)

**Notes:** Questionable status. Species complex. Salazar-Vallejo (2017) revised the genus *Brada* and transferred a large number of species (including *Brada villosa*) to *Bradabyssa* Hartman, 1967. Previous descriptions of *Bradabyssa villosa* by e.g. Fauvel (1927), comprise a wide variety of characters and probably comprise more than one species under this name (Salazar-Vallejo 2017). Likewise, molecular analyses by Carr et al. (2011) revealed at least two putative cryptic species in Canadian waters (one in the Arctic, another in the Pacific Ocean) which were initially identified as *Bradabyssa villosa*. Salazar-Vallejo (2017) restricts *Bradabyssa villosa* sensu stricto to the North Atlantic Ocean and European and Russian Arctic and re-instates *Bradabyssa parthenopeia* (Lo Bianco, 1893) from the Mediterranean (Gulf of Naples) – a species previously considered a synonym of *Bradabyssa villosa*. It is highly likely that

*Bradabyssa villosa* is absent from the Mediterranean and specimens from this region belong to other species such as *Bradabyssa parthenopeia*.

### ***Diplocirrus glaucus* (Malmgren, 1867)**

### ***Diplocirrus hirsutus* (Hansen, 1878)**

**Notes:** Questionable status. One specimen from Greece in the collections of the Senckenberg Museum (SMF 11285, 39°09'46.8"N, 19°26'21.6"E, 1008 m depth, coll. date 1993-05-19, det. D. Fiege). In the Mediterranean also reported from the Adriatic, but records are considered doubtful by Mikac (2015) based on the Arctic and Subarctic distribution of the species.

### ***Flabelligera affinis* M. Sars, 1829**

**Notes:** Species complex. Consists of at least two different putative cryptic species in the Arctic Ocean (Carr et al. 2011), specimens from other locations could belong to either of these or to a different cryptic species. Salazar-Vallejo (2012) considers the distribution of *Flabelligera affinis* to be restricted to "Arctic to cold and temperate, boreal localities" and questions the identity of specimens reported under this name from warm water localities such as Africa or Panama. Whether the Mediterranean is considered here temperate or warm water and whether the species occurs in the Mediterranean is unknown; it is considered here to be present in the Greek waters pending further information.

### ***Flabelligera diplochaitus* (Otto, 1820)**

**Notes:** Type locality: Mediterranean.

### ***Pherusa plumosa* (Müller, 1776)**

**Nomenclature:** *Pherusa plumosa* (Müller, 1776) | *Stylarioides plumosa* (O.F. Müller, 1776)

**Notes:** Questionable status. Salazar-Vallejo (2014) revised the genus *Pherusa* and reinstated a number of species, many of which were previously considered synonyms of *Pherusa plumosa*. He restricts *Pherusa plumosa* in its diagnosis and distribution; currently it is believed to occur in the Arctic and temperate Atlantic Ocean and to be absent from the Mediterranean. The three currently known species of *Pherusa* occurring in the Mediterranean Sea – previously all grouped under the name *Pherusa plumosa* – are *Pherusa incrustans* Quatrefages, 1866, *Pherusa mikacae* Salazar-Vallejo, 2014 and *Pherusa obscura* Quatrefages, 1849. As the description by Fauvel (1927) is very general and includes a large range of characters which correspond to *Pherusa plumosa* sensu stricto, *Pherusa obscura* and *Pherusa incrustans* (Salazar-Vallejo 2014), all specimens identified using Fauvel's key are questionable.

***Piromis eruca* (Claparède, 1869)**

**Nomenclature:** *Pherusa eruca* (Claparède, 1869) | *Piromis eruca* (Claparède, 1869) | *Stylarioides eruca* (Claparède, 1869)

**Notes:** Type locality: Mediterranean (Gulf of Naples)

***Stylarioides monilifer* Delle Chiaje, 1841**

**Nomenclature:** *Pherusa monilifera* (Delle Chiaje, 1841)

**Notes:** Salazar-Vallejo (2011) revised the genus *Stylarioides* and reinstated *Stylarioides hirsutus* Lo Bianco, 1893, which Fauvel (1927) had synonymised with *Stylarioides monilifer*. Thus, specimens identified using the publication by Fauvel (1927) or works based on it (e.g. Castelli 1990) could belong to either of the two species. Type locality: Mediterranean (Gulf of Naples).

***Therochaeta flabellata* (Sars in Sars, 1872)**

**Nomenclature:** *Pherusa flabellata* (M. Sars in G.O. Sars, 1872) | *Therochaeta flabellata* (M. Sars in G.O. Sars, 1872)

**Notes:** Questionable status. Salazar-Vallejo (2013) restricts *Therochaeta flabellata* to the North-East Atlantic Ocean and Arctic waters and assigns Mediterranean material, including the description by Fauvel (1927) to an undescribed species (informally named *Therochaeta* cf. *flabellata* “Mediterranean”, formal description pending). The Mediterranean form differs from the original description mainly in the size of the much larger sediment particles and in the lack of fusion of the first two chaetigers.

**Glyceridae Grube, 1850*****Glyceria alba* (O.F. Müller, 1776)**

**Nomenclature:** *Glyceria alba* (O.F. Müller, 1776) | *Glyceria minuta* (Bobretzky, 1870)

**Notes:** Böggemann (2002) distinguishes the species from *Glyceria tridactyla* Schmarda, 1861, mainly on the basis of the shape of the proboscival papillae and ailerons; although the separation was judged problematic by Worsfold (2007). Parapar and Moreira (2015) agree in separating the two species on the basis of the above two characters.

***Glycera capitata* Ørsted, 1843**

**Notes:** Frequently reported from the Mediterranean (e.g. Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015), but Böggemann (2002) and Böggemann (2015) restricts its distribution to Arctic, Antarctic, cold temperate areas and deep sea down to 5655m (Böggemann 2015). The Mediterranean material could possibly be *Glycera lapidum* Quatrefages, 1866 or *Glycera noelae* Böggemann, Bienhold & Gaudron, 2012 (Böggemann, pers. comm.)

***Glycera celtica* O'Connor, 1987**

**Notes:** Böggemann (2002) synonymised *Glycera dayi* O' Connor, 1987 with *Glycera celtica* but this synonymy has not been accepted unanimously (Worsfold 2007, Parapar and Moreira 2015).

***Glycera fallax* Quatrefages, 1850**

**Nomenclature:** *Glycera gigantea* Quatrefages, 1866

***Glycera lapidum* Quatrefages, 1866**

**Notes:** O'Connor (1987) reports at least four varieties amongst the examined material from the North-East Atlantic, although these correspond in fact to different species (Böggemann, pers. comm.).

***Glycera rouxii* Audouin & Milne Edwards, 1833**

**Notes:** Böggemann (2002) considers *Glycera rouxii* a synonym of *Glycera unicornis* Lamarck, 1818 (based on comparisons with original description as the holotype is lost). He considers the shape of the retractable branchiae (reported as bifid in *Glycera unicornis*) not to be a diagnostic character, since variations were found in the same populations (Böggeman, pers. comm.). However, Worsfold (2007) and Parapar and Moreira (2015) debate this synonymy and highlight differences in parapodial lobes besides those of the branchiae, although parapodial lobes were found to change between juvenile and adult specimens by Böggemann (2002). They propose that the branchial shape remains a diagnostic character, which should however be used with caution due to retractability. Despite the available evidence for the synonymy, two recent works consider both species as valid. Therefore, the past records of *Glycera rouxii* in Greece (see Supplementary material) are currently retained as separate, until a response to the arguments of Parapar and Moreira (2015) is available. Type locality: Mediterranean (Marseille).

***Glycera tessellata* Grube, 1863**

**Notes:** Type locality: Mediterranean (Croatia).

### ***Glycera tridactyla* Schmarda, 1861**

**Nomenclature:** *Glycera convoluta* Keferstein, 1862 | *Glycera tridactyla* Schmarda, 1861

**Notes:** Böggemann (2002) considers *Glycera convoluta* a synonym of *Glycera tridactyla* based on comparisons with the original description and illustrations. A number of authors support this synonymy (e.g. Worsfold 2007, Parapar and Moreira 2015).

### ***Glycera unicornis* Lamarck, 1818**

**Notes:** See notes under *Glycera rouxii* Audouin & Milne Edwards, 1833.

## **Goniadidae Kinberg, 1865**

### ***Glycinde bonhourei* Gravier, 1904**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Simboura (2008). In the Mediterranean also known from Israel (Böggemann 2005), Egypt (Ben-Eliahu 1972a) and Turkey (Çınar et al. 2014). Originally described from Bonhoure (Djibuti).

### ***Glycinde nordmanni* (Malmgren, 1866)**

**Nomenclature:** *Eone nordmanni* Malmgren, 1866 | *Glycinde nordmanni* (Malmgren, 1866)

### ***Goniada emerita* Audouin & Milne Edwards, 1833**

**Notes:** Type locality: Mediterranean (Nice, France).

### ***Goniada gigantea* (Verrill, 1885)**

**Notes:** Reported from Greece by Böggemann (2005) from material collected by E. Marenzeller in 1894 on the Cyclades plateau (as *Goniada emerita* Audouin & Milne Edwards, 1833). In the Mediterranean also known from Spain and France (Böggemann 2005). Otherwise reported from the West and East Atlantic and North-East Pacific (Böggemann 2005).

***Goniada hexadentes* Böggemann & Eibye-Jacobsen, 2002**

**Notes:** Reported from Greece by Böggemann (2005) from the abyssal zone (3848 m) off the Peloponnese. In the Mediterranean also known from Spain, Italy and Israel (Böggemann 2005), otherwise reported from the Eastern Atlantic and the Indian Ocean (Böggemann 2005).

***Goniada maculata* Ørsted, 1843*****Goniada norvegica* Ørsted, 1845*****Goniadella bobrezkii* (Annenkova, 1929)**

**Notes:** In the Mediterranean only reported from Greece (Arvanitidis 2000a). Commonly distributed in the Black Sea (Annenkova 1929) and around the North-East Atlantic (Wolff and Stegenga 1975, Kirkegaard 2001).

***Goniadella gracilis* (Verrill, 1873)**

**Notes:** Reported from Greece by Bogdanos and Satsmadjis (1983) and Bogdanos and Satsmadjis (1987). In the Mediterranean also known from Israel (Böggemann 2005), otherwise reported from the North-West Atlantic, East Atlantic and South-West Indian Ocean (Böggemann 2005).

**Hesionidae Grube, 1850*****Gyptis propinqua* Marion & Bobretzky, 1875**

**Nomenclature:** *Gyptis propinqua* Marion & Bobretzky, 1875 | *Oxydromus propinquus* (Marion & Bobretzky, 1875)

**Notes:** Type locality: Mediterranean (France).

***Hesione splendida* Savigny in Lamarck, 1818**

**Nomenclature:** *Hesione pantherina* Risso, 1826 | *Hesione splendida* Savigny in Lamarck, 1818

**Notes:** *Hesione* species display a high colour variability and it is currently unknown whether these are colour morphs of the same species or belong to different species (Mikac 2015, Jimi et al. 2017). For this reason, the status of *Hesione pantherina*, *Hesione splendida* and similar species is still unclear; here we follow the opinion of F.

Pleijel (pers. comm. to Mikac 2015) and list occurrences of *Hesione pantherina* under *Hesione splendida*.

### ***Hesionides arenaria* Friedrich, 1937**

**Notes:** Schmidt and Westheide (2000) assess the genetic differentiation of worldwide *Hesionides arenaria* populations. Specimens from Crete group with those from near the type locality in the North Sea, indicating the presence of *Hesionides arenaria* sensu stricto in Greece.

### ***Hesionides gohari* Hartmann-Schröder, 1960**

**Notes:** Species complex. Schmidt and Westheide (1999) recover three genetic lineages from worldwide localities which could correspond to cryptic species. Specimens from Crete group with those from near the type locality in the Red Sea, indicating the presence of *Hesionides gohari* sensu stricto in Greece.

### ***Hesiospina aurantiaca* (M. Sars, 1862)**

**Nomenclature:** *Hesiospina aurantiaca* (M. Sars, 1862) | *Hesiospina similis* (Hessle, 1925)

**Notes:** Frequently reported from Greece, considered cosmopolitan (Pleijel 2004). Simboura (1996) notes morphological differences between specimens of *Hesiospina similis* from the Mediterranean and from the native range (Japan). Specifically, Mediterranean material exhibits a stronger serration of the blades of the compound chaetae and neuropodial aciculae with rounded instead of tapering tips.

### ***Leocrates atlanticus* (McIntosh, 1885)**

**Notes:** Reported by von Marenzeller (1902) from deep waters (808 m) off Milos. Marenzeller's two specimens (Natural History Museum Vienna, Inv. No. 599, Acq. No. 15503) were examined by S. Faulwetter. One specimen was confirmed as *Leocrates atlanticus*; in the second specimen the jaws could not be observed without dissection and its identity is not confirmed.

### ***Leocrates claparedii* (Costa in Claparède, 1868)**

**Notes:** Synonymised by Hartman (1965a) with *Leocrates chinensis* Kinberg, 1866, *Leocrates claparedii* was reported under the former name for several years (see also remarks in Table 1), therefore listed in inventories of non-native species in the Mediterranean (e.g. Zenetos et al. 2010). Pleijel (1998), in his revision of the Hesionidae, considers the two species distinct. Type locality: Mediterranean (Gulf of Naples).



***Microphthalmus aberrans* (Webster & Benedict, 1887)**

**Notes:** Questionable status. Reported from Greece by Akoumianaki and Hughes (1997), also found by Papageorgiou et al. (2006) (unpublished data), but *Microphthalmus aberrans* is a taxonomically confused species. Riser (2000) revised the species and found the slides of the type material to be comprised of two different species. He assigned specimens previously assigned to *Microphthalmus aberrans* to three species: *Microphthalmus aberrans*, *Microphthalmus pettiboneae* Riser, 2000 and *Microphthalmus aggregatus* Riser, 2000. However, the species re-described as *Microphthalmus aberrans* by Riser does not resemble previously available descriptions of the species, as those were mostly copied from a re-description by Southern (1914), which was based on the wrong microscope slide. The specimen on this slide was re-described as *Microphthalmus pettiboneae* by Riser, causing confusion in the application of the name *Microphthalmus aberrans*. All specimens reported under this name would need to be re-investigated to clarify their identity.

***Microphthalmus fragilis* Bobretzky, 1870**

**Notes:** Reported from Greece by Koukouras (1979), also found by Papageorgiou et al. (2006) (unpublished data). In the Mediterranean also known from Italy (Castelli et al. 2008), otherwise distributed in the Black and North Seas.

***Microphthalmus pseudoaberrans* Campoy & Vieitez, 1982**

**Notes:** Reported from Greece by NCMR (1995) and NCMR (2000a). In the Mediterranean also known from Spain (Capaccioni-Azzati and Torres-Gavila 1989), otherwise distributed along the Atlantic coast of the Iberian Peninsula.

***Microphthalmus szcelkowiei* Metschnikow, 1865**

**Notes:** Reported from Greece by Dando et al. (1995). In the Mediterranean also known from Egypt (Dorgham et al. 2013), otherwise distributed in the Black Sea, North Sea and East coast of North America.

***Microphthalmus tyrrhenicus* Zunareli-Vandini, 1967**

**Notes:** New record for Greece. One specimen, Elafonisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Zunarelli (1967), Sordino (1990). Type locality: Mediterranean (Leghorn coast, Italy).

***Neogyptis mediterranea* (Pleijel, 1993)**

**Nomenclature:** *Gyptis mediterranea* Pleijel, 1993

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016). Type locality: Mediterranean (Banyuls-sur-Mer, France).

### ***Nereimyra punctata* (Müller, 1788)**

**Notes:** Questionable status. Pleijel et al. (2012) consider the status of the species in the Mediterranean doubtful, as no specimens could be recovered from typical habitats despite intense sampling. Based on examination of other specimens, the authors conclude that the species is probably restricted to boreal regions. Records from the Adriatic and the Sea of Marmara are likewise considered questionable by Mikac (2015) and Çınar et al. (2014) respectively.

### ***Oxydromus agilis* (Ehlers, 1864)**

**Nomenclature:** *Ophiodromus agilis* (Ehlers, 1864)

**Notes:** Questionable status. Reported from Greece by Nicolaidou et al. (1990). Type locality: Mediterranean (Martinscica, Adriatic). While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.

### ***Oxydromus flexuosus* (Delle Chiaje, 1827)**

**Nomenclature:** *Ophiodromus flexuosus* (Delle Chiaje, 1827)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Oxydromus pallidus* Claparède, 1864**

**Nomenclature:** *Ophiodromus pallidus* (Claparède, 1864) | *Podarke pallida* (Claparède, 1864)

**Notes:** Type locality: Mediterranean (Port-Vendres, France).

### ***Podarkeopsis capensis* (Day, 1963)**

**Nomenclature:** *Gyptis capensis* (Day, 1963) | *Podarkeopsis capensis* (Day, 1963)

**Notes:** May have been confused with *Podarkeopsis galangai* Laubier 1961 or *Podarkeopsis arenicolus* (La Greca, 1946), both native to the area but not included in any keys for the region and therefore probably underreported (Pleijel 2005). Reported from the Mediterranean, European Atlantic coasts and South Africa, considered questionable in Turkey by Çınar et al. (2014).

***Podarkeopsis galangai* Laubier, 1961**

**Notes:** Reported from Greece by Arvanitidis (1994) based on a single specimen. Type locality: Mediterranean (Banyuls-sur-Mer, France). Pleijel (1998) considers the species a junior synonym of *Podarkeopsis arenicolus* (La Greca, 1946) but does not provide details and does not formalise the synonymy. As Rizzo and Salazar-Vallejo (2014) list both species in their key and recognise difference in the size of the eyes, length of the median antenna and chaetal details, the species is therefore treated as valid here, pending further information.

***Psamathe fusca* Johnston, 1836**

**Nomenclature:** *Kefersteinia cirrhata* (Keferstein, 1862)

***Syllidia armata* Quatrefages, 1866**

**Nomenclature:** *Magalia perarmata* Marion & Bobretzky, 1874 | *Syllidia armata* Quatrefages, 1866

**Iospilidae Bergström, 1914*****Phalacrophorus pictus* Greeff, 1879**

**Notes:** Questionable status. Reported from Greece by Antoniadou and Chintiroglou (2005) from a benthic sample of hard substrates at 3–15 m depth collected via SCUBA diving. No description or further information was provided for the taxon. Iospilidae are holopelagic and easily confused with benthic juvenile forms of other species (Jiménez-Cueto et al. 2006). Thus the record is considered doubtful here.

**Lacydoniidae Bergström, 1914*****Lacydonia laureci* Laubier, 1975**

**Notes:** Originally described from the Hellenic Trench (Matapan deep; 35°49'48"N, 22°20'42"E; 4690 m depth), no other records from Greece.

***Lacydonia miranda* Marion & Bobretzky, 1875**

**Notes:** Type locality: Mediterranean (Gulf of Marseille).

## Longosomatidae Hartman, 1944

### *Heterospio mediterranea* Laubier, Picard & Ramos, 1973

**Notes:** Type locality: Mediterranean (Algiers Bay).

## Lopadorrhynchidae Claparède, 1870

### *Lopadorrhynchus appendiculatus* Southern, 1909

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). Widely distributed in the Mediterranean (Wesenberg-Lund 1939a).

### *Lopadorrhynchus brevis* Grube, 1855

**Notes:** Reported from Greece by Wesenberg-Lund (1939a); three additional specimens in the Senckenberg collection management system (Cat. No. 11278, 34°26'41.4"N, 26°06'34.2"E, 4255–4308 m; Cat. No. 11279, 35°48'57.6"N, 25°15'46.8"E, 1875–1877 m, Cat. No. 11280, 35°50'23.4"N, 25°16'08.4"E, 1876 m, all det. D. Fiege). Type locality: Mediterranean.

### *Lopadorrhynchus krohnii* (Claparède, 1870)

**Notes:** Reported from Greece by Wesenberg-Lund (1939a); one additional specimen from Greece in the collections of the Natural History Museum London (NHM 1968.52, 37°49'02"N, 19°45'06"E, 0-500 m depth, det. N. Tebble). Type locality: Mediterranean (Gulf of Naples).

### *Lopadorrhynchus nationalis* Reibisch, 1893

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). *Lopadorrhynchus nationalis* is often listed in the literature as a synonym of *Lopadorrhynchus brevis* Grube, 1855, but Dales and Peter (1972) point out differences between the two species and consider them distinct. Widely distributed in the Mediterranean (Wesenberg-Lund 1939a).

### *Lopadorrhynchus uncinatus* Fauvel, 1915

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). In the Mediterranean also known from Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015), otherwise cosmopolitan in pelagic waters (Dales and Peter 1972).

***Maupasia coeca* Viguier, 1886**

**Notes:** Reported from Greece by NCMR (1989). Type locality: Mediterranean (Bay of Algiers).

***Pelagobia longicirrata* Greeff, 1879**

**Notes:** One specimen from Greece in the collections of the Senckenberg Museum (SMF 11264, 35°50'40.8"N, 22°19'49.8"E, 4754–4766 m, coll. date 1993-05-25, det. D. Fiege). In the Mediterranean also known from Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015). Present in the adjacent Sea of Marmara (Çınar et al. 2014). Otherwise cosmopolitan in pelagic waters (Dales and Peter 1972).

**Lumbrineridae Schmarda, 1861*****Gallardonis iberica* Martins, Carrera-Parra, Quintino & Rodrigues, 2012**

**Notes:** Reported from Greece by Katsiaras et al. (2017). In the Mediterranean also known from the central and western basins (Bertasi et al. 2014, Gómez et al. 2015, Mikac 2015, D'Alessandro et al. 2016). Its presence in the region was probably overlooked until its description.

***Hilbigneris gracilis* (Ehlers, 1868)**

**Nomenclature:** *Hilbigneris gracilis* (Ehlers, 1868) | *Lumbriconereis gracilis* Ehlers, 1868 | *Lumbrineris gracilis* (Ehlers, 1868)

**Notes:** Type locality: Mediterranean (Adriatic).

***Lumbricalus adriatica* (Fauvel, 1940)**

**Nomenclature:** *Lumbricalus adriatica* (Fauvel, 1940) | *Lumbrineris adriatica* (Fauvel, 1940)

**Notes:** Type locality: Mediterranean (Adriatic). Carrera-Parra (2004) redefined the genus, including *Lumbricalus adriatica*, using the maxillary apparatus in combination with chaetal types for differentiating species.

***Lumbrinerides acuta* (Verrill, 1875) sensu Ramos, 1976**

**Nomenclature:** *Lumbrineris acuta* Verrill, 1875 | *Lumbrinerides acuta* (Verrill, 1875)

**Notes:** Questionable status. Reported from Greece by Dounas (1986). The record refers to *Lumbrinerides acuta* sensu Ramos 1976. The original description by Verrill

(1875) is not detailed, but the redescription by Perkins (1979) based on specimens collected in the type locality (Rhode Island, USA) includes an accessory tooth on maxillary pair M1. The absence of an accessory tooth, as it is described in Ramos (1976), may indicate an undescribed species (Gómez et al. 2015). In the Mediterranean also known from Spain (Sardá et al. 1999), Italy (Castelli et al. 2008), Turkey (Çinar et al. 2014) and the Adriatic (Mikac 2015), but it is unclear whether these records refer to specimens having an accessory tooth or not. While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.

### ***Lumbrinerides amoueuxi* Miura, 1980**

**Notes:** In the Mediterranean known from Greece, Turkey (Çinar et al. 2014) and Cyprus (Çinar 2005), otherwise distributed along the Atlantic coast of Europe.

### ***Lumbrinerides carpinei* (Ramos, 1976)**

**Notes:** New record for Greece. Two specimens, Pachia Ammos, Crete, 35°06'39.6"N, 25°48'32.4"E, 1–5 m depth, fine to coarse sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Ramos (1976). Type locality: Western Mediterranean.

### ***Lumbrinerides laubieri* Miura, 1980**

**Notes:** In the Mediterranean only reported from Greece (Simboura and Zenetos 2005). Known from deep waters off the Atlantic coast of Europe.

### ***Lumbrinerides neogesae* Miura, 1981**

**Native status:** Non-native (casual)

**Notes:** New record for Greece. One specimen, Elafonisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Miura (1980), Gravina and Cantone (1991). In the Mediterranean also known from Italy (Gravina and Cantone 1991). Originally described from South Africa.

### ***Lumbrineriopsis paradoxa* (Saint-Joseph, 1888)**

**Nomenclature:** *Lumbriconereis paradoxa* Saint-Joseph, 1888 | *Lumbrineriopsis paradoxa* (Saint-Joseph, 1888) | *Lumbrineris paradoxa* Saint-Joseph, 1888

***Lumbrineris coccinea* (Renier, 1804)**

**Nomenclature:** *Lumbriconereis coccinea* (Renier, 1804) | *Lumbrineris coccinea* (Renier, 1804)

**Notes:** Type locality: Mediterranean. Carrera-Parra (2006) considers records from outside the Mediterranean Sea doubtful.

***Lumbrineris inflata* Moore, 1911**

**Notes:** Questionable status. Reported from Greece by Arvanitidis (2000a) and Kitsos (2003); the first Mediterranean record was by Giangrande et al. (1981). Arvanitidis (2000a) argues that the species may have been underreported in the Mediterranean due to its similarity with *Lumbrineris coccinea* (Renier, 1804). Carrera-Parra (2006) and D'Alessandro et al. (2016) consider Mediterranean records of the species doubtful based on its otherwise East Pacific distribution. Çınar (2009) argues that the description and illustrations of *Lumbrineris inflata* by Giangrande et al. (1981) fit the characters of *Lumbrineris perkinsi* Carrera-Parra, 2001. *Lumbrineris inflata* differs from *Lumbrineris perkinsi* mainly in the number of teeth of maxilla III (M III). Giangrande et al. (1981), as well as Arvanitidis (2000a) for Greece, report a maxillary formula which could correspond to *Lumbrineris perkinsi* ("M III with 3–4 teeth"). Although published descriptions and records from the Mediterranean indicate the presence of *Lumbrineris perkinsi*, re-examination of all the material is needed to draw conclusions about the identity of the Greek records.

***Lumbrineris latreilli* Audouin & Milne Edwards, 1834**

**Nomenclature:** *Lumbriconereis latreilli* Audouin & Milne Edwards, 1834 | *Lumbrineris latreilli* Audouin & Milne Edwards, 1834

**Notes:** Frequently reported from Greece, however, mainly on the basis of external and unreliable diagnostic characters. Species descriptions focusing on the maxillary apparatus and other unambiguous and size-independent characters have recently become available (Carrera-Parra 2006, Martins et al. 2012, D'Alessandro et al. 2016). Therefore, re-examination of material could reduce the frequency of records and even reveal the presence of overlooked species. In the Mediterranean, the species is widely recorded (e.g. Campoy 1982, Çınar 2005, Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015). Otherwise distributed in the North-East Atlantic; its presumed cosmopolitan distribution is questionable (Carrera-Parra 2006).

***Lumbrineris luciliae* Martins, Carrera-Parra, Quintino & Rodrigues, 2012**

**Notes:** New record for Greece. Specimens previously reported as as *Lumbrineris cingulata* Ehlers, 1897 by Chatzigeorgiou et al. (2016) from rocky shores in northern Crete were re-examined and found to belong to *Lumbrineris luciliae* as described by

Martins et al. (2012), although the reported slight distal curvature of the aciculae could not be observed. In the Mediterranean also known from Italy (D'Alessandro et al. 2016). Otherwise known from the Atlantic coast of the Iberian Peninsula (Martins et al. 2012).

### ***Lumbrineris nonatoi* Ramos, 1976**

**Notes:** Type locality: Mediterranean (Baie de Rosas, Spain). *Lumbrineris nonatoi* has several diagnostic characters in common with the – later described – *Gallardoneris iberica* Martins, Carrera-Parra, Quintino & Rodrigues, 2012. Due to this confusion, Bertasi et al. (2014) and Katsiaras et al. (2017) suggest that several literature records of *Lumbrineris nonatoi* could belong to the overlooked *Gallardoneris iberica*, although the overall presence of the former in the region has not been doubted.

### ***Ninoe armoricana* Glémarec, 1968**

#### ***Ninoe nigripes* Verrill, 1873**

**Nomenclature:** *Ninoe kinbergi* Ehlers, 1887 | *Ninoe nigripes* Verrill, 1873

**Notes:** Questionable status. Reported from Greece as *Ninoe kinbergi* by Vamvakas (1970) and Vamvakas (1971). Arvanitidis (2000a) lists these records under the name *Ninoe nigripes*, following the synonymy established by Pettibone (1963). In the Mediterranean reported from the central basin as *Ninoe kinbergi* (Castelli et al. 2008) and as *Ninoe nigripes* by Mikac (2015) from the Adriatic Sea. Otherwise distributed in the West Atlantic.

### ***Scoletoma emandibulata* (Pillai, 1961)**

**Nomenclature:** *Lumbrineris emandibulata* Pillai, 1961

**Notes:** Questionable status. Reported from Greece by Dando et al. (1995) and Akoumianaki (2004). The record by Akoumianaki (2004) actually refers to *Scoletoma emandibulata mabiti*, as Ramos 1976 is given as the authority; it remains unclear whether the same holds true for the specimen of Dando et al. (1995). Originally described from the Indian Ocean.

### ***Scoletoma emandibulata mabiti* (Ramos, 1976)**

**Nomenclature:** *Lumbrineris emandibulata mabiti* Ramos, 1976

**Notes:** Type locality: Mediterranean (Catalan Coast).



***Scoletoma fragilis* (O.F. Müller, 1776)**

**Nomenclature:** *Lumbriconereis fragilis* (O.F. Müller, 1776) | *Lumbrineris fragilis* (O.F. Müller, 1776) | *Scoletoma fragilis* (O.F. Müller, 1776)

**Notes:** Species complex. Frequently reported from Greece and other parts of the Mediterranean (Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015). Otherwise distributed in the Atlantic Ocean, Arctic Ocean and North Sea. However, molecular analyses revealed the existence of at least two putative cryptic species, one in the Atlantic and one in the Arctic Ocean (Carr et al. 2011).

***Scoletoma funchalensis* (Kinberg, 1865)**

**Nomenclature:** *Lumbriconereis funchalensis* Kinberg, 1865 | *Lumbrineris funchalensis* (Kinberg, 1865) | *Scoletoma funchalensis* (Kinberg, 1865)

**Notes:** In the Mediterranean also known from the western and central basins (Campoy 1982, Castelli et al. 2008), Tunisia (Ayari et al. 2009), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015). Otherwise distributed in the North-East Atlantic. However, Oug (2011) noted that *Scoletoma funchalensis* is poorly known and the described characters by Fauvel (1923) could be confused with juveniles of several species.

***Scoletoma impatiens* (Claparède, 1868)**

**Nomenclature:** *Lumbriconereis impatiens* (Claparède, 1868) | *Lumbrineris impatiens* Claparède, 1868 | *Scoletoma impatiens* (Claparède, 1868)

**Notes:** Type locality: Mediterranean (Gulf of Naples). George and Hartmann-Schröder (1985) synonymised *Scoletoma impatiens* with the South African *Scoletoma tetraura* (Schmarda, 1861), but several authors do not accept this synonymy (see also notes under *Scoletoma tetraura*).

***Scoletoma rovignensis* (Fauvel, 1940)**

**Nomenclature:** *Lumbrineris rovignensis* Fauvel, 1940 | *Scoletoma rovignensis* (Fauvel, 1940)

**Notes:** Questionable status. Reported from Greece by Dounas (1986). Type locality: Mediterranean (Adriatic). While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.

### ***Scoletoma tetraura* (Schmarda, 1861)**

**Nomenclature:** *Lumbrineris tetraura* (Schmarda, 1861) | *Scoletoma tetraura* (Schmarda, 1861)

**Notes:** Questionable status. Reported from Greece by Koukouras and Rousso (1991). In the Mediterranean also known from Turkey (Doğan et al. 2005; but considered questionable by Çınar et al. 2014), Cyprus (Çınar 2005), Italy (Castelli et al. 2008), Albania (Marzano et al. 2010) and the Adriatic (listed under *Scoletoma impatiens* (Claparède, 1968) by Mikac (2015)). Originally described from South Africa. George and Hartmann-Schröder (1985) synonymised the European species *Scoletoma impatiens* with *Scoletoma tetraura*. However, several authors question the presence of the latter in Europe and suggest using the name *Scoletoma impatiens* for specimens from the area until a more detailed revision becomes available (Martins et al. 2012, Çınar et al. 2014, Gómez et al. 2015, Mikac 2015, D'Alessandro et al. 2016). The main diagnostic character of the genus can be problematic, since several juvenile specimens of *Lumbrineris* and *Hilbigneris* species may lack composite hooded hooks and can be key out as *Scoletoma* (Oug 2011).

## **Magelonidae Cunningham & Ramage, 1888**

### ***Magelona alleni* Wilson, 1958**

**Notes:** Reported from Greece by Dounas (1986) and Arvanitidis (1994). In the Mediterranean also known from France (Fiege et al. 2000b), Italy (Castelli et al. 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015). Probably underreported, since older records of *Magelona cincta* Ehlers, 1908 may belong to *Magelona alleni* (Wilson 1958; see also notes under *Magelona cincta*).

### ***Magelona cincta* Ehlers, 1908**

**Notes:** Questionable status. Reported from Greece by Zarkanellas (1980) and Makra and Nicolaidou (2000), but specimens probably belong to *Magelona alleni* Wilson, 1958. Wilson (1958) described *Magelona alleni* from material misidentified as *Magelona cincta* and concluded that the latter does not occur in European coasts.

### ***Magelona equilamellae* Harmelin, 1964**

**Notes:** Type locality: Mediterranean (Villefranche-sur-Mer, France).

### ***Magelona filiformis* Wilson, 1959**

***Magelona minuta* Eliason, 1962*****Magelona mirabilis* (Johnston, 1865)**

**Notes:** Probably an underreported species, since older records of *Magelona papillicornis* F. Müller, 1858 may belong to *Magelona mirabilis* (Fiege et al. 2000b; see also notes under *Magelona papillicornis*)

***Magelona papillicornis* F. Müller, 1858**

**Notes:** Questionable status. Until 1977, all European magelonids with mucronate chaetae on chaetiger 9 were assigned to *Magelona papillicornis*, originally described from Brazil. However, Jones (1977) re-described *Magelona papillicornis* and clarified that it actually lacks this character and therefore questioned its presence in European waters. Fiege et al. (2000b) demonstrated the presence of two species bearing mucronate chaetae on chaetiger 9 in Europe; *Magelona mirabilis* (Johnston, 1865) and *Magelona johnstoni* Fiege, Licher & Mackie, 2000. Greek records of *Magelona papillicornis* could belong to either of these, although only *Magelona mirabilis* has been recorded up to now.

***Magelona wilsoni* Glémarec, 1966**

**Notes:** One specimen from Greece in the collections of the Senckenberg Museum (SMF 11286, 39°15'0.6"N, 23°42'32.4"E, 1243 m, coll. date 1998-01-01, det. D. Fiege). In the Mediterranean also known from the coasts of France and Spain (Labruno et al. 2007, Serrano-Samaniego 2012) and the Adriatic (Mikac 2015).

**Maldanidae Malmgren, 1867*****Axiiothella constricta* (Claparède, 1869)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Axiiothella rubrocincta* (Johnson, 1901)**

**Notes:** Questionable status. Species complex. In the Mediterranean only reported from Greece (Dando et al. 1995). Distributed along the Pacific coast of North America where at least two morphologically indistinguishable forms with different life histories occur (Wilson 1983).

***Chirimia biceps* (M. Sars, 1861)**

**Nomenclature:** *Asychis biceps* (M. Sars, 1861) | *Chirimia biceps* (M. Sars, 1861) | *Chirimia biceps biceps* (Sars, 1861)

***Clymenella cf. koellikeri* (McIntosh, 1885)**

**Notes:** In the Mediterranean only reported from Greece (Simboura 1996). The specimens differ from *Clymenella torquata* (Leidy 1855) (originally from the East coast of North America but occurring non-natively on the Atlantic coast of Europe) in the shape and number of acicular chaetae and from *Clymenella cincta* (Saint-Joseph, 1894) (occurring in the Mediterranean) in the shape of the collar and of the nuchal organs. The Greek specimens are most similar to the Pacific species *Clymenella koellikeri*, from which they differ in the following characters: Nuchal slits of *Clymenella koellikeri* reach up to the middle of the cephalic plate (until the lateral notches in the Greek material) and the rim or the collar of the 4<sup>th</sup> chaetiger is smooth or slightly wavy in *Clymenella koellikeri*, whereas one Greek specimen shows a small lateral recess at the rim of the collar (Simboura 1996). The species is found regularly in Greece but is usually reported at genus level due to its uncertain identity. It may be an overlooked or cryptic species.

***Euclymene collaris* (Claparède, 1869)**

**Nomenclature:** *Clymene collaris* (Claparède, 1869) | *Euclymene collaris* (Claparède, 1869)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Euclymene lombricoides* (Quatrefages, 1866)**

**Nomenclature:** *Clymene lombricoides* Quatrefages, 1866 | *Euclymene lombricoides* (Quatrefages, 1866)

***Euclymene oerstedii* (Claparède, 1863)**

**Nomenclature:** *Clymene oerstedii* Claparède, 1863 | *Euclymene oerstedii* (Claparède, 1863)

***Euclymene palermitana* (Grube, 1840)**

**Nomenclature:** *Clymene palermitana* Grube, 1840 | *Euclymene palermitana* (Grube, 1840) | *Praxillella palermitana* (Grube, 1840)

**Notes:** Type locality: Mediterranean (Palermo, Sicily, Italy).

***Heteroclymene robusta* Arwidsson, 1906**

**Nomenclature:** *Clymene robusta* (Arwidsson, 1906) | *Euclymene robusta* (Arwidsson, 1906) | *Heteroclymene robusta* Arwidsson, 1906

***Johnstonia clymenoides* Quatrefages, 1866**

**Notes:** Reported from Greece by Bogdanos and Satsmadjis (1983) and Xenopoulou (1987). In the Mediterranean also known from France, Spain and Israel (Mackie and Gobin 1993), Egypt (Abd-Elnaby 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed along the European and North African Atlantic coasts (Mackie and Gobin 1993).

***Leiochone leiopygos* (Grube, 1860)**

**Nomenclature:** *Clymenura clypeata* (Saint-Joseph, 1894) | *Leiochone clypeata* Saint-Joseph, 1894 | *Leiochone leiopygos* (Grube, 1860)

***Lumbriclymene minor* Arwidsson, 1906*****Macroclymene santandarensis* (Rioja, 1917)**

**Nomenclature:** *Clymene santandarensis* Rioja, 1917 | *Euclymene santandarensis* (Rioja, 1917) | *Macroclymene santandarensis* (Rioja, 1917)

***Maldane capensis* (Day, 1961)**

**Nomenclature:** *Asychis capensis* Day, 1961 | *Maldane capensis* (Day, 1961)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Papadopoulos 1986). Known from South Africa.

***Maldane glebifex* Grube, 1860*****Maldane sarsi* Malmgren, 1865*****Maldanella harai* (Izuka, 1902)**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Nicolaidou 1985, Bogdanos and Satsmadjis 1987, Nicolaidou and Papadopoulou 1989, Koulouri et al. 2015). Outside the Mediterranean reported from worldwide locations, mostly from bathyal and abyssal depths.

### ***Metasychis gotoi* (Izuka, 1902)**

**Nomenclature:** *Asychis gotoi* (Izuka, 1902) | *Metasychis gotoi* (Izuka, 1902)

**Native status:** Non-native (established)

**Notes:** Originally distributed in the Red Sea, the Indian Ocean and the North Pacific; commonly reported from locations throughout the Mediterranean (Simboura and Zenetos 2005). The first Mediterranean record of *Metasychis gotoi* is from the Northern Adriatic (Fauvel 1940), constituting one of the earliest records of non-native species in the basin. As *Metasychis gotoi* is reported from worldwide distributions, it is likely that the name refers to a complex of cryptic species. The Mediterranean material could belong to an overlooked native species (J. Langeneck, pers. comm.).

### ***Micromaldane ornithochaeta* Mesnil, 1897**

**Notes:** Reported from Greece by Eleftheriou et al. (1990) and NCMR (2000a). In the Mediterranean also known from Israel (Ben-Eliahu 1976b), Spain (Alós 1990), Italy (Gherardi et al. 2002), Egypt (Abd-Elnaby 2008) and the Adriatic (Mikac 2015). Present in the adjacent Sea of Marmara (Çınar et al. 2014). Otherwise distributed along the European Atlantic coasts (Gherardi et al. 2002).

### ***Nicomache lumbricalis* (Fabricius, 1780)**

**Notes:** Questionable status. Frequently reported from Greece and other Eastern Mediterranean locations; however, de Assis et al. (2007) in their review of the genus restrict its distribution to boreal regions.

### ***Nicomache maculata* Arwidsson, 1911**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (e.g. Bogdanos and Satsmadjis 1983, Eleftheriou et al. 1990, Tselepides 1992, Simboura 1996; full reference list in Suppl. material 2), de Assis et al. (2007) restrict its distribution to the Shetland Islands.

### ***Nicomache trispinata* Arwidsson, 1906**

**Notes:** Reported from Greece by Tselepides (1992). In the Mediterranean also known from France (Clausade 1969), Israel (Ben-Eliahu 1995) and Italy (Castelli et al. 2008), otherwise distributed along the Atlantic coasts of Europe and in the North Sea.

***Petaloproctus terricolus* Quatrefages, 1866*****Praxillella affinis* (M. Sars in G.O. Sars, 1872)**

**Nomenclature:** *Clymene affinis* M. Sars in G.O. Sars, 1872 | *Praxillella affinis* (M. Sars in G.O. Sars, 1872)

***Praxillella gracilis* (M. Sars, 1861)**

**Nomenclature:** *Clymene gracilis* Sars, 1861 | *Praxillella gracilis* (M. Sars, 1861)

***Praxillella lophoseta* (Orlandi, 1898)**

**Nomenclature:** *Clymene lophosetosa* Orlandi, 1898 | *Praxillella lophoseta* (Orlandi, 1898)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Praxillella praetermissa* (Malmgren, 1865)**

**Notes:** Species complex. At least three putative cryptic species exist, one each in the Arctic and West Atlantic (Carr et al. 2011) and another in Portugal (Lobo et al. 2016).

***Praxillura longissima* Arwidsson, 1906**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Eleftheriou et al. 1990). Commonly distributed in the North-East Atlantic.

***Rhodine gracilior* Tauber, 1879*****Rhodine loveni* Malmgren, 1865****Myzostomatidae Benham, 1896*****Myzostoma cirriferum* Leuckart, 1836**

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016) based on a single specimen. In the Mediterranean also known from Tunisia (Wesenberg-Lund 1939b), France (Laubier and Paris 1962), Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015), otherwise distributed along the Atlantic coasts of Europe (Nygren and Pleijel 2015).

## Nephtyidae Grube, 1850

### *Aglaophamus agilis* (Langerhans, 1880)

**Nomenclature:** *Aglaophamus rubella* (Michaelsen, 1897) | *Nephtys agilis* Langerhans, 1880 | *Nephtys rubella* Michaelsen, 1897

### *Aglaophamus malmgreni* (Théel, 1879)

**Nomenclature:** *Aglaophamus malmgreni* (Théel, 1879) | *Nephtys malmgreni* Théel, 1879

**Notes:** Questionable status. Reported from Greece by von Marenzeller (1902). Marenzeller's description differs from the current concept of *Aglaophamus malmgreni* in the first branchial chaetiger and the pharyngeal papillation. Marenzeller describes his specimens with branchiae starting at chaetigers 6–7 (in one specimen at chaetiger 10) and states that his specimens correspond to the description of *Nephtys malmgreni* by McIntosh (1900) (listed by McIntosh under the name *Nephtys longisetosa*) in the number of papillae. McIntosh describes 15 rows of 11–15 papillae. Branchial and papillation characters correspond to *Aglaophamus pulcher* (Rainer, 1991), but Marenzeller's material should be examined for confirmation. In the Mediterranean, *Aglaophamus malmgreni* has also been reported from the Adriatic Sea (Mikac 2015), otherwise it is distributed in circumpolar regions and records from southern Europe require confirmation (Ravara et al. 2010).

### *Aglaophamus pulcher* (Rainer, 1991)

**Nomenclature:** *Nephtys pulchra* Rainer, 1991

**Notes:** Reported from Greece by Arvanitidis (2000a) but specimens of *Aglaophamus malmgreni* (Théel, 1897) by von Marenzeller (1902) could belong to *Aglaophamus pulcher* (see notes there). In the Mediterranean also known from abyssal canyons and plains in the western basin (Ravara et al. 2010), otherwise distributed in the North East Atlantic. May have been confused in the past with *Nephtys hystricis* McIntosh, 1900 and *Nephtys incisa* Malmgren 1865 (Rainer 1991) and therefore been overlooked in the area.

### *Inermonephtys inermis* (Ehlers, 1887)

**Nomenclature:** *Aglaophamus inermis* (Ehlers, 1887) | *Inermonephtys inermis* (Ehlers, 1887) | *Nephtys inermis* Ehlers, 1887

**Notes:** Questionable status. Ravara et al. (2010) found morphological differences between European specimens identified as *Inermonephtys inermis* and the holotype from Florida and established a new species, *Inermonephtys foretmontardoii* Ravara,



Cunha & Pleijel, 2010 for the European populations. *Inermonephtys inermis* is restricted to the West Atlantic. It is highly likely that Greek specimens of *Inermonephtys inermis* belong to *Inermonephtys foretmontardoi*, but no specimens have been re-examined yet for confirmation.

### ***Micronephtys sphaerocirrata* (Wesenberg-Lund, 1949)**

**Nomenclature:** *Micronephtys sphaerocirrata* (Wesenberg-Lund, 1949)| *Nephtys sphaerocirrata* Wesenberg-Lund, 1949

**Notes:** Ravara et al. (2010) regard the presence of the species in the Mediterranean as doubtful due to its otherwise Indo-Pacific distribution. Greek specimens may also have been confused with *Micronephtys stammeri* (Augener, 1932) in the past.

### ***Micronephtys stammeri* (Augener, 1932)**

**Nomenclature:** *Micronephtys maryae* San Martín, 1982| *Micronephtys stammeri* (Augener, 1932)

**Notes:** Type locality: Mediterranean (*Micronephtys maryae*: Ibiza, *Micronephtys stammeri*: Adriatic).

### ***Nephtys assimilis* Ørsted, 1843**

**Notes:** Rarely reported from the Mediterranean, but was considered a synonym of *Nephtys hombergii* Savigny in Lamarck, 1818 until its resurrection and redescription by Rainer (1989) and could therefore be underreported.

### ***Nephtys caeca* (Fabricius, 1780)**

**Notes:** Ravara et al. (2010) re-examined material from the Mediterranean and identified it as *Nephtys caeca*, concluding that, although the species is commonly distributed in Arctic waters, it occasionally occurs in warmer waters of the Mediterranean as far as the Black Sea.

### ***Nephtys ciliata* (Müller, 1788)**

**Notes:** Questionable status. Reported from Greece by Vamvakopoulou (1991). In the Mediterranean also reported from the western basin (Fredj 1974) and the Adriatic (Mikac 2015). Commonly distributed in boreal areas; Mediterranean records require confirmation (Ravara et al. 2010).

### ***Nephtys cirrosa* Ehlers, 1868**

### ***Nephtys hombergii* Savigny in Lamarck, 1818**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Nephtys hystricis* McIntosh, 1900**

**Notes:** The description of *Nephtys hystricis* by Fauvel (1923) actually refers to *Nephtys incisa* Malmgren, 1865 and vice versa (Rainer 1990). Thus, specimens identified as *Nephtys hystricis* using Fauvel's key and description probably belong to *Nephtys incisa*. Both *Nephtys incisa* and *Nephtys hystricis* have been reported from Greece, thus they can both be considered as present, despite this confusion.

### ***Nephtys incisa* Malmgren, 1865**

**Notes:** See notes under *Nephtys hystricis* McIntosh, 1900.

## **Nereididae Blainville, 1818**

### ***Alitta succinea* (Leuckart, 1847)**

**Nomenclature:** *Neanthes succinea* (Leuckart, 1847) | *Nereis succinea* Leuckart, 1847

**Notes:** Fauvel (1923) considered *Nereis lamellosa* Ehlers 1868 a synonym of *Nereis succinea* but revoked this synonymy in a subsequent publication, describing the differentiating characters (Fauvel 1936b). Later authors (e.g. Day 1967, Campoy 1982) continued to treat the species as synonymous. Thus, many reports of *Alitta succinea* may actually contain specimens of *Nereis lamellosa*, which in turn is probably underreported in the Mediterranean (Arvanitidis 2000a).

### ***Alitta virens* (M. Sars, 1835)**

**Nomenclature:** *Neanthes virens* (M. Sars, 1835)

**Notes:** Questionable status. Species complex. In the Mediterranean only reported from Greece (Christomanos and Giannitsis 1962). Distributed in the North Atlantic and North Pacific and comprising at least two cryptic species with genetic and morphological differences in these two areas (Carr et al. 2011).

### ***Ceratonereis (Compositia) costae* (Grube, 1840)**

**Nomenclature:** *Ceratonereis (Compositia) costae* (Grube, 1840) | *Ceratonereis costae* (Grube, 1840) | *Nereis costae* Grube, 1840 | *Nereis (Ceratonereis) costae* Grube, 1840

**Notes:** Type locality: Mediterranean.

***Ceratonereis (Composetia) hircinicola* (Eisig, 1870)**

**Nomenclature:** *Ceratonereis hircinicola* (Eisig, 1870) | *Nereis hircinicola* (Eisig, 1870)

**Notes:** Type locality: Mediterranean.

***Ceratonereis (Composetia) vittata* Langerhans, 1884**

**Notes:** In the Mediterranean only reported from Greece (Maidanou et al. 2017). Originally described from Madeira; possibly underreported due to confusion with *Nereis rava* (Ehlers, 1864) or *Ceratonereis hircinicola* (Eisig, 1870) (Núñez and Brito 2002).

***Eunereis longissima* (Johnston, 1840)**

**Notes:** Reported from Greece by Arvanitidis (2000a) and Chatzigeorgiou et al. (2016). In the Mediterranean also known from France (Picard 1965), Spain (Núñez 2004), Italy (Castelli et al. 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015). Otherwise distributed along the Atlantic coasts of Europe.

***Hediste diversicolor* (O.F. Müller, 1776)**

**Nomenclature:** *Hediste diversicolor* (O.F. Müller, 1776) | *Nereis (Hediste) diversicolor* Müller, 1776 | *Nereis diversicolor* Müller, 1776

**Notes:** Species complex. Multiple genetic lineages exist in Europe (Nygren 2014, Lobo et al. 2016) which could correspond to cryptic species. Vasileiadou et al. (2016) found genetically highly isolated populations with unique haplotypes in Greek lagoons.

***Leonnates persicus* Wesenberg-Lund, 1949**

**Native status:** Non-native (casual)

**Notes:** New record for Greece. Two specimens, Argolikos Gulf, 37°33'34.9"N, 22°47'06"E, 16 m depth, sandy silt with biogenic detritus, collected during monitoring for the Water Framework Directive. Literature used for identification: Çınar et al. (2002). An Indo-Pacific species, in the Mediterranean also known from Israel (Ben-Eliahu 1991b) and Turkey (Çınar et al. 2002).

***Micronereis variegata* Claparède, 1863*****Neanthes acuminata* (Ehlers, 1868)**

**Nomenclature:** *Neanthes caudata* (sensu Delle Chiaje, 1827) | *Nereis caudata* (sensu Delle Chiaje, 1827)

**Notes:** Species complex. Reish et al. (2014) recovered four different putative cryptic species of *Neanthes acuminata* from worldwide locations (based on specimens traditionally referred to the species *Neanthes acuminata*, *Neanthes caudata*, and *Neanthes arenaceodentata* Moore, 1903, all of which are morphologically indistinguishable but could not be recovered by the analyses). From Europe, only specimens from Portugal were included in the analyses and it is currently unknown whether more cryptic species exist. Concerning nomenclature, *Neanthes acuminata* sensu lato should be used to refer to the species complex (Reish et al. 2014). While the name *Neanthes caudata* is often found in literature for the European species, it is not an available name. Delle Chiaje (1825) wrongly applied the name *Spio caudatus* Lamarck, 1818 to a nereidid, thus the name is based on a misidentification. The next available name is *Neanthes acuminata* (see detailed notes by G. Read for [S. caudatus Lamarck, 1818](#), [Spio caudatus sensu Delle Chiaje, 1827](#) and [N. caudata \(sensu Delle Chiaje, 1827\)](#)), a species described from the Gulf of Naples.

### ***Neanthes agulhana* (Day, 1963)**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016). Originally from South Africa, in the Mediterranean also reported from Spain (Núñez 2004) and Italy (Occhipinti-Ambrogi et al. 2011).

### ***Neanthes fucata* (Savigny in Lamarck, 1818)**

**Notes:** Reported from Greece by Keklikoglou et al. (2013). Type locality: Eastern Mediterranean.

### ***Neanthes kerguelensis* (McIntosh, 1885)**

**Nomenclature:** *Neanthes kerguelensis* (McIntosh, 1885) | *Nereis kerguelensis* McIntosh, 1885

### ***Neanthes nubila* (Savigny, 1822)**

**Nomenclature:** *Neanthes irrorata* (Malmgren, 1867) | *Neanthes nubila* (Savigny, 1822) | *Nereis irrorata* (Malmgren, 1867)

### ***Nereis falsa* Quatrefages, 1866**

**Nomenclature:** *Nereis falsa* Quatrefages, 1866 | *Nereis splendida* Grube, 1840

**Notes:** *Nereis falsa* is considered a synonym of *Nereis splendida* Grube, 1840 by Núñez (2004). However, a *Nereis splendida* Blainville, 1825 also exists in literature and in online databases (e.g. WoRMS) which should take priority over Grube's species and

render the latter a junior homonym. Gravina et al. (2015) therefore use *Nereis falsa* as the valid name, as *Nereis splendida* Grube, although having priority over *Nereis falsa* Quatrefages, is pre-occupied. However, the identity of the species described by Blainville (1825) species is confused and early authors (e.g. Quatrefages 1866:434, Rathke 1843:172) consider it indeterminable and probably belonging to *Nephtys*. In addition, Blainville (1825) creates a confusing situation of synonyms by applying the name *Nereis splendida* to a specimen he had received from Dr. Leach under the (probably unpublished) name *Nereis clava* (p. 439) and in the same publication transfers *Hesione splendida* Savigny 1818 to *Nereis*, creating an unresolved synonymy with *Nereis splendida* (p. 443). Until this confusion is resolved, the name *Nereis falsa* is used here, following Gravina et al. (2015). See also Salazar-Vallejo et al. (2017) and [discussions on the Annelida mailing list](#) for extensive clarifications on the identity and nomenclatural problems associated with *Nereis falsa* and *Nereis splendida*.

### ***Nereis lamellosa* Ehlers, 1864**

**Notes:** Reported from Greece by Arvanitidis (2000a) and Keklikoglou et al. (2013). Type locality: Mediterranean (Adriatic). Probably underreported, see remarks under *Alitta succinea* (Leuckart, 1947).

### ***Nereis pelagica* Linnaeus, 1758**

**Notes:** Species complex. At least three putative cryptic species exist in the Arctic, the Atlantic and Portugal (Lobo et al. 2016).

### ***Nereis perivisceralis* Claparède, 1868**

**Notes:** Reported from Greece by Keklikoglou et al. (2013) and Chatzigeorgiou et al. (2016). Type locality: Mediterranean (Gulf of Naples). Considered a synonym of *Nereis falsa* Quatrefages 1866 by Fauvel (1923), thus it may be underreported and records of *Nereis falsa* from Greece may include specimens of *Nereis perivisceralis*.

### ***Nereis pulsatoria* (Savigny, 1822)**

**Nomenclature:** *Nereis pulsatoria* (Savigny, 1822) | *Nereis zonata* Malmgren, 1867

**Notes:** The taxonomy and nomenclature of *Nereis pulsatoria* and *Nereis zonata* are confused. In literature, both *Nereis pulsatoria* (Savigny, 1822) and *Nereis pulsatoria* Audouin & Milne Edwards 1834 are used (the latter not as a new name but as a reference to Savigny's species, thus not a homonym). It is currently unclear whether these two actually refer to the same species. Fauvel (1923) considers *Nereis pulsatoria* Audouin & Milne Edwards 1834 a synonym of *Nereis zonata* but does not mention Savigny's species. Núñez (2004) synonymises *Nereis zonata* Malmgren 1867 and *Nereis pulsatoria* Audouin & Milne Edwards, 1834 with *Nereis pulsatoria* (Savigny 1822), but without any mention of studied material or reference to the original

description. Gravina et al. (2015) point out that the description of *Nereis pulsatoria* by Núñez (2004) is inconsistent when compared to the corresponding illustrations and that the described species differs from *Nereis zonata* in chaetal and paragnath arrangement. Gravina et al. (2015) consider only Savigny's species valid and restrict it to the Atlantic, but the authors base this statement on the respective species entries in WoRMS, which have not been reviewed recently, contain partly incorrect information and may constitute database inconsistencies. If the species are indeed synonymous, *Nereis pulsatoria*, being the older name, should have priority over *Nereis zonata*. Until this confusion is resolved, the name *Nereis pulsatoria* is kept here and *Nereis zonata* considered its synonym. Specimens from Greece have both been reported under the name *Nereis zonata* and *Nereis pulsatoria* sensu Núñez (2004).

### ***Nereis rava* Ehlers, 1864**

**Notes:** Type locality: Mediterranean (Adriatic).

### ***Perinereis cultrifera* (Grube, 1840)**

**Notes:** Species complex. Multiple records from Greece exist, but *Perinereis cultrifera* constitutes a complex of species in the Mediterranean (Scaps et al. 2000, Maltagliati et al. 2001). Specimens from Greece could belong to any of these or another cryptic species. Type locality: Mediterranean (Gulf of Naples).

### ***Perinereis oliveirae* (Horst, 1889)**

**Notes:** Reported from Greece by Koukouras et al. (1985) and Simboura (1987). In the Mediterranean also known from France (Bellan 1964a), Spain (Núñez 2004) and Italy (Castelli et al. 2008), otherwise distributed along the Atlantic coasts of Europe and North Africa. Italian records are, however, considered questionable by Castelli et al. (2008) as the species is easily confused with *Perinereis rullieri* Pilato, 1974.

### ***Perinereis tenuisetis* (Fauvel, 1915)**

**Notes:** One specimen from Greece in the collections of the Senckenberg Museum (SMF 5609, 36°36'N, 21°34'E, 3848 m depth, coll. date 1993-05-22 / 1993-05-23 det. M.N. Ben-Eliahu). Mediterranean species, type locality Sicily.

### ***Platynereis coccinea* (Delle Chiaje, 1822)**

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016). Type locality: Mediterranean (Italy). Previous reports of the species by Arvanitidis (2000a) and Simboura and Nicolaidou (2001) are based on a record by Ergen (1987) from the Turkish Aegean coast.

***Platynereis dumerilii* (Audouin & Milne Edwards, 1834)**

**Notes:** Lobo et al. (2016) show that specimens from Portugal and Italy genetically diverge, indicating potentially more than one cryptic species in Europe (pending further corroboration through analysis of more specimens). One of the sibling species of *Platynereis dumerilii* occurring near CO<sub>2</sub> vents in Ischia, Italy, was recently shown to correspond to *Platynereis massiliensis* (Moquin-Tandon, 1869) (Valvassori et al. 2015).

***Pseudonereis anomala* Gravier, 1900**

**Nomenclature:** *Nereis anomala* (Gravier, 1900) | *Pseudonereis anomala* Gravier, 1900

**Native status:** Non-native (established)

**Notes:** Frequently reported from Greece. Originally from the Red Sea, in the Mediterranean also known from Egypt (Fauvel 1937), Israel (Ben-Eliahu 1989), Turkey (Çınar and Ergen 2005) and Italy (D'Alessandro et al. 2015).

***Websterinereis glauca* (Claparède, 1870)**

**Nomenclature:** *Leptonereis glauca* (Claparède, 1870) | *Websterinereis glauca* (Claparède, 1870)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

**Oeononidae Kinberg, 1865*****Arabella coeca* Fauvel, 1940**

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016) based on a single specimen. Type locality: Mediterranean (Adriatic).

***Arabella geniculata* (Claparède, 1868)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Arabella iricolor* (Montagu, 1804)**

**Notes:** Species complex. Frequently reported from Greece and other parts of the Mediterranean (Çınar 2005, Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015) and considered cosmopolitan. May however constitute a species complex comprising two cryptic species or populations along the East Coast of North America, differing in the maxillary formula (Colbath 1989, Zanol and Ruta 2015).

### ***Drilonereis filum* (Claparède, 1868)**

**Nomenclature:** *Drilonereis filum* (Claparède, 1868) | *Drilonereis macrocephala* Saint-Joseph, 1888

**Notes:** Type locality: Mediterranean (Gulf of Naples). Often considered cosmopolitan, but Helgason et al. (1990) note the possibility of more than one species being reported under *Drilonereis filum*, since a wide variation of characters is found in literature.

### ***Notocirrus scoticus* McIntosh, 1869**

### ***Oligognathus bonelliae* Spengel, 1882**

**Parasite of:** *Bonellia viridis* Rolando, 1822

**Notes:** *Oligognathus bonelliae* can be found in host bodies of the echiurid *Bonellia viridis*. Type locality: Mediterranean (Gulf of Naples).

## **Onuphidae Kinberg, 1865**

### ***Aponuphis bilineata* (Baird, 1870)**

**Nomenclature:** *Aponuphis bilineata* (Baird, 1870) | *Hyalinoecia bilineata* Baird, 1870

**Notes:** Species complex. Two colour morphs exist (Arias and Paxton 2015a) and molecular evidence of sibling species within *Aponuphis bilineata* are provided by Borisova et al. (submitted).

### ***Aponuphis brementi* (Fauvel, 1916)**

**Nomenclature:** *Aponuphis brementi* (Fauvel, 1916) | *Aponuphis fauveli* (Rioja, 1918) | *Hyalinoecia brementi* Fauvel, 1916

**Notes:** Species complex. Arias and Paxton (2015a) consider *Aponuphis fauveli* a junior synonym of *A. brementi*, thus literature records of *Aponuphis fauveli* are here assigned to *Aponuphis brementi*. However, molecular evidence of sibling species within *Aponuphis brementi* were found by Borisova et al. (submitted). Type locality: Mediterranean (Bay of Roquebrun, France).

### ***Diopatra neapolitana* Delle Chiaje, 1841**

**Nomenclature:** *Diopatra neapolitana* Delle Chiaje, 1841 | *Diopatra neapolitana neapolitana* Delle Chiaje, 1841



**Notes:** *Diopatra neapolitana*, originally described from the Gulf of Naples, used to be the only *Diopatra* species recorded along European coasts. However, several recent publications have demonstrated that more *Diopatra* species (*Diopatra marocensis* Paxton, Fadlaoui & Lechapt, 1995, *Diopatra micrura* Pires, Paxton, Quintino & Rodrigues, 2010, *Diopatra biscayensis* Fauchald, Berke & Woodin, 2012) are present along European coasts (e.g. Fauchald et al. 2012, Arias and Paxton 2013), including the Turkish Aegean (Çınar et al. 2014). Therefore, at least some of the Greek specimens of *Diopatra neapolitana* could belong to other *Diopatra* species.

### ***Hyalinoecia tubicola* (O.F. Müller, 1776)**

**Notes:** Species complex. Zaâbi et al. (2015) present evidence of possible cryptic species within *Hyalinoecia tubicola* in Tunisia.

### ***Nothria conchylega* (Sars, 1835)**

**Nomenclature:** *Nothria conchylega* (Sars, 1835) | *Onuphis conchylega* Sars, 1835

**Notes:** Species complex. Frequently reported from Greece and other parts of the Mediterranean (e.g. Çınar 2005, Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015). Otherwise distributed in the Arctic, Atlantic, Indian and Pacific Oceans. However, many of these records could belong to other species. Budaeva and Paxton (2013) note that morphological variations of *Nothria* Malmgren, 1866 have been underestimated in earlier species descriptions, including those of *Nothria conchylega*. In addition, two putative cryptic species of *Nothria conchylega* were discovered in the Arctic and Pacific Ocean (Carr et al. 2011).

### ***Onuphis eremita* Audouin & Milne Edwards, 1833**

**Notes:** Species complex. Frequently reported from Greece and other parts of the Mediterranean (e.g. Campoy 1982, Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015) and considered cosmopolitan. However, the taxonomic history of the species includes doubtful synonymies, vague descriptions and redescriptions based on material from worldwide locations (Arias and Paxton 2014). The latter authors redefine the diagnostic characters, designate a neotype and confine its distribution to the East Atlantic and the Western and Central Mediterranean. In addition, they re-instate and re-describe *Onuphis pancerii* Claparède, 1868 from the *Onuphis eremita* complex and designate a neotype from Italy. Re-examination of specimens is required to determine the status of records of *Onuphis eremita* and possible other species of the complex in Greece.

### ***Paradiopatra calliopae* Arvanitidis & Koukouras, 1997**

**Notes:** Originally described from Greece by Arvanitidis and Koukouras (1997), mainly based on having equal antennae instead of a longer median one in *Paradiopatra bihanica* Intes & Le Loeuff, 1975. Arvanitidis and Koukouras (1997) examined the holotype of *Paradiopatra bihanica* but found it to be dried out and the tips of the antennae broken off. Budaeva and Fauchald (2011) considered the holotype to be in good condition and with equal antennae, placing *Paradiopatra calliopae* into synonymy with *Paradiopatra bihanica*. However, Paxton and Arias (2016), after re-examining the holotype of *Paradiopatra bihanica* and specimens of *Paradiopatra calliopae*, re-instated the latter as a valid species and doubt the presence of *Paradiopatra bihanica* in the Mediterranean. See also notes under *Paradiopatra quadricuspis* (M. Sars in G.O. Sars, 1872).

### ***Paradiopatra fragosa* (Ehlers, 1887)**

**Nomenclature:** *Sarsonuphis fragosa* (Ehlers, 1887)

**Notes:** Questionable status. Reported twice from Greece as *Sarsonuphis* cf. *fragosa* (Karakassis 1991, Tselepides 1992) and once as *Sarsonuphis fragosa* (Eleftheriou and Smith 2000). No other Mediterranean records, commonly distributed in the Caribbean Sea. The species is not included in any studies of *Paradiopatra* from southern Europe (e.g. Arvanitidis and Koukouras 1997, Arias and Paxton 2015b) and it is probably absent from the area. Literature records could belong to *Paradiopatra calliopae* Arvanitidis & Koukouras, 1997.

### ***Paradiopatra lepta* (Chamberlin, 1919)**

**Nomenclature:** *Onuphis lepta* Chamberlin, 1919

**Notes:** Questionable status. Reported from Greece by Vamvakas (1971). Arias and Paxton (2015b) doubt the validity of records from European coasts and suggest that this species has been confused with *Paradiopatra bihanica*, Intes & Le Loeuff, 1975 (Mediterranean records now accepted as *Paradiopatra calliopae* Arvanitidis & Koukouras, 1997) or *Nothria maremontana* Andre & Pleijel, 1989. Arvanitidis and Koukouras (1997) note that, where descriptions are available for onuphids recorded in the Mediterranean under the name *Onuphis lepta*, these differ from the original *Onuphis lepta* and are much closer to *Paradiopatra calliopae*. Although re-examination of material is required, the distribution in the region is considered doubtful.

### ***Paradiopatra quadricuspis* (M. Sars in G.O. Sars, 1872)**

**Nomenclature:** *Paradiopatra quadricuspis* (M. Sars in G.O. Sars, 1872) | *Sarsonuphis quadricuspis* (M. Sars in G.O. Sars, 1872)

**Notes:** Questionable status. Budaeva and Fauchald (2011) doubt the presence of the species in the Mediterranean and suggest that Greek records by Arvanitidis and Koukouras (1997) may represent a new species. Their argument of a limited distribution from Norway to Iceland has been accepted also by other authors (Mikac 2015, Arias and Paxton 2015b, but see Çınar et al. 2014). Simboura (1996) found specimens of *Sarsonuphis quadricuspis* from Greece to differ from the type material but also from French material described by Bellan (1963). She considered it an undescribed species designated as *Sarsonuphis* sp.; but later found it to match the description of *Paradiopatra calliopae* Arvanitidis & Koukouras, 1997. Re-examination of material is required to draw conclusions about the identity of the Greek records but the species' distribution in the region is doubtful.

### ***Rhamphobrachium brevibrachiatum* (Ehlers, 1874)**

**Nomenclature:** *Onuphis brevibranchiata* (Ehlers, 1874) | *Rhamphobrachium brevibrachiatum* (Ehlers, 1874)

**Notes:** Reported from Greece by Simboura (1996). In the Mediterranean also known from the central and western basins (Bellan 1964a, Castelli et al. 2008) and the Adriatic (Mikac 2015); otherwise distributed in the North Atlantic and Japan.

## **Opheliidae Malmgren, 1867**

### ***Armandia cirrhosa* Filippi, 1861**

**Notes:** Type locality: Mediterranean (Sardinia).

### ***Armandia polyophtalma* Kükenthal, 1887**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Ophelia barquii* Fauvel, 1927**

**Notes:** Reported from Greece by Papageorgiou et al. (2006). Type locality: Mediterranean (Agay, France). *Ophelia barquii* was initially described as a subspecies of *Ophelia radiata* (Delle Chiaje, 1828) by Fauvel (1927), who distinguished the European species of *Ophelia* (*Ophelia neglecta* Schneider, 1892, Savigny, 1822, *Ophelia radiata* (Delle Chiaje, 1828) and *Ophelia barquii*) on the basis of the number of gill pairs. This was regarded as an unreliable character by subsequent authors (e.g. Bellan 1964a) who considered *Ophelia radiata barquii* and *Ophelia radiata* synonyms of *Ophelia bicornis*. However, molecular and morphological analyses by Maltagliati et al. (2004) and Maltagliati et al. (2005) clearly differentiate *Ophelia barquii* from *Ophelia bicornis*, the constant morphological difference between the two species being the

number of nephridiopore pairs (five in *Ophelia barquii*, six in *Ophelia bicornis*). See also comments under *Ophelia bicornis*.

### ***Ophelia bicornis* Savigny in Lamarck, 1818**

**Nomenclature:** *Ophelia bicornis* Savigny in Lamarck, 1818 | *Ophelia radiata* (Delle Chiaje, 1828)

**Notes:** Species complex with a confused taxonomic history. Originally described from the Gulf of Naples. Fauvel (1927) considered *Ophelia bicornis* and *Ophelia radiata* as distinct based on the number of gill pairs (*Ophelia radiata*: 14 pairs; *Ophelia bicornis*: 15 pairs) and geographic distribution. However, both species and intermediate forms (with 14 gill pairs on one side and 15 on the other) have been found to co-occur by Britton-Davidian and Amoureux (1982), who also found genetic differences between the two forms in an Atlantic population, with asymmetric specimens belonging to either of the two genetic variants. Based on these findings, the authors consider the species distinct. However, they were not able to assign Mediterranean specimens to either of the two Atlantic forms. Maltagliati et al. (2005) consider the results inconclusive and refer to the complex as *Ophelia bicornis sensu lato*. Parapar (2012) studied individuals of both *Ophelia radiata* and *Ophelia bicornis* from the collections of the Natural History Museums in Copenhagen and Madrid and could not find any differences apart from the number of gill pairs — a character which he considered too unreliable for discrimination.

### ***Ophelia limacina* (Rathke, 1843)**

**Nomenclature:** *Ophelia borealis* Quatrefages, 1866

**Notes:** Reported from Greece by Maidanou et al. (2017). In the Mediterranean also known from Israel (Ben-Eliahu 1995), Italy (Castelli et al. 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed in the boreal regions of Europe.

### ***Ophelia roscoffensis* Augener, 1910**

**Notes:** Reported from Greece by Simboura (1996), Zenetos et al. (1997) (based on the same specimens); Identification confirmed by G. Bellan (pers. comm. in Simboura 1996). In the Mediterranean also known from France (Bellan 1964a) and Turkey (Çınar et al. 2014), otherwise known from the Atlantic coasts of France, Spain and from the British Isles. The Greek specimens differ from Atlantic ones by a reduced number of branchial pairs (Simboura 1996).

***Ophelina acuminata* Ørsted, 1843**

**Nomenclature:** *Ammotrypane aulogaster* Rathke, 1843 | *Ophelina acuminata* Ørsted, 1843 | *Ophelina aulogaster* (Rathke, 1843)

**Notes:** Species complex. Comprises at least two putative cryptic species, one in the Atlantic, one in the Arctic Ocean (Carr et al. 2011).

***Ophelina cylindricaudata* (Hansen, 1879)**

**Nomenclature:** *Ammotrypane cylindricaudatus* Hansen, 1879 | *Ophelina cylindricaudata* (Hansen, 1879)

***Polyophthalmus pictus* (Dujardin, 1839)*****Tachytrypane jeffreysii* McIntosh, 1879****Orbiniidae Hartman, 1942*****Leitoscoloplos kerguelensis* (McIntosh, 1885)**

**Notes:** Questionable status. Reported from Greece by Akoumianaki (2004). In the Mediterranean also reported (questionably) from Spain / France (Ramos 1976b). *Leitoscoloplos kerguelensis* is a widely reported but much confused species with several incorrect re-descriptions and many wrongly identified specimens (Mackie 1987). The species is probably restricted to the Kerguelen Islands area and the Mediterranean material belongs to *Leitoscoloplos kerguelensis* sensu Ramos, 1976 which constitutes an undescribed species (Blake 2017).

***Naineris laevigata* (Grube, 1855)**

**Notes:** Type locality: Mediterranean.

***Naineris quadraticeps* Day, 1965**

**Native status:** Non-native (questionable)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Harmelin 1969). Distributed in the Indo-Pacific; its occurrence in the Mediterranean is considered questionable (Zenetos et al. 2010).

***Orbinia latreillii* (Audouin & Milne Edwards, 1833)**

**Nomenclature:** *Orbinia latreillii* (Audouin & Milne Edwards, 1833) | *Phylo latreillii* (Audouin & Milne Edwards, 1833)

***Orbinia sertulata* (Savigny, 1822)**

**Nomenclature:** *Orbinia cuvierii* (Audouin & Milne Edwards, 1833)

***Phylo foetida* (Claparède, 1869)**

**Nomenclature:** *Aricia foetida* Claparède, 1869 | *Phylo foetida* (Claparède, 1869) | *Phylo foetida foetida* (Claparède, 1870)

**Notes:** Species complex. Bleidorn et al. (2009) recover two putative cryptic species in the Atlantic and the Mediterranean (Sicily). Originally described from the Gulf of Naples, it is unknown whether more cryptic species exist in the Mediterranean and whether the Greek specimens belong to *Phylo foetida* sensu stricto or to a different species.

***Phylo grubei* (McIntosh, 1910)**

**Nomenclature:** *Aricia grubei* McIntosh, 1919 | *Phylo grubei* (McIntosh, 1910)

***Phylo kupfferi* (Ehlers, 1874)**

**Nomenclature:** *Aricia kupfferi* Ehlers, 1874 | *Phylo kupfferi* (Ehlers, 1874)

***Phylo norvegica* (M. Sars in G.O. Sars, 1872)**

**Notes:** Reported from Greece by Papadopoulos (1986); two additional specimens from Greece in the collections of the Senckenberg Museum (SMF 15218, 35°31'37.2"N, 27°13'28.2"E, 122 m depth, coll. date 2005-08-30, det: D. Fiege; SMF 15227, 35°31'33.6"N, 27°13'24"E, 126–142 m depth, coll. date 2005-08-30, det: D. Fiege). Records of *Phylo norvegica* from the Adriatic are doubtful and probably refer to *Phylo foetida* (Mikac 2015).

***Protoaricia oerstedii* (Claparède, 1864)**

**Nomenclature:** *Protoaricia oerstedii* (Claparède, 1864) | *Theostoma oerstedii* (Claparède, 1864)

**Notes:** Type locality: Mediterranean (Port-Vendres, France).

***Scoloplos armiger* (Müller, 1776)**

**Notes:** Species complex. In addition to forming a complex of at least three putative cryptic species in northern Europe alone (Bleidorn et al. 2006), *Scoloplos armiger* is also easily confused with congeners (e.g. the two Mediterranean species *Scoloplos haasi* (Monro, 1937) and *Scoloplos typicus* (Eisig, 1914)).

***Scoloplos chevalieri candiensis* Harmelin, 1969**

**Notes:** Originally described from the south coast of Crete (Kaloi Limenes, 7–11 m depth; meadows of *Halophila stipulacea* and *Cymodocea nodosa*).

***Scoloplos haasi* (Monro, 1937)**

**Nomenclature:** *Scolaricia haasi* Monro, 1937 | *Scoloplos haasi* (Monro, 1937)

**Notes:** Questionable status. Reported from Greece by Dounas (1986). Type locality: Mediterranean (Israel). As the species is easily confused with *Scoloplos armiger* (Müller, 1776) (López 2009), records of *Scoloplos armiger* from Greece may comprise specimens belonging to *Scoloplos haasi*. While it is quite possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.

***Scoloplos typicus* (Eisig, 1914)**

**Nomenclature:** *Scolaricia typica* Eisig, 1914

**Notes:** As the species is easily confused with *Scoloplos armiger* (Müller, 1776) (Mikac 2015), many records of *Scoloplos armiger* from Greece may refer to *Scoloplos typicus*. Type locality: Mediterranean (Gulf of Naples).

**Oweniidae Rioja, 1917*****Galathowenia fragilis* (Nilsen & Holthe, 1985)**

**Nomenclature:** *Myriochele fragilis* Nilsen & Holthe, 1985

**Notes:** Previously only known from the Arctic, the species was found in deep waters (>4000 m) off Crete by Fiege et al. (2000a) in large numbers, constituting to date the only records from Greece and the Mediterranean.

### ***Galathowenia oculata* (Zachs, 1923)**

**Nomenclature:** *Galathowenia oculata* (Zachs, 1923) | *Myriochele oculata* Zachs, 1923

**Notes:** Martín (1989) and Simboura (1996) note some morphological variation (constant presence of the third mid-dorsal pygidial lobe) in Mediterranean specimens compared to Atlantic and Pacific material.

### ***Myriochele heeri* Malmgren, 1867**

**Notes:** Questionable status. Martín (1989) re-examined material identified as *Myriochele heeri* in the Mediterranean and assigned the material and the description of *Myriochele heeri* by Fauvel (1927) to *Galathowenia oculata* (Zachs, 1923). Parapar (2006) revised and re-described *Myriochele heeri*, as the original type series of *Myriochele heeri* constituted a mixture of different and confused species. The species is probably restricted to the boreal Atlantic and Arctic regions and Mediterranean records are questionable (Parapar 2006). Specimens identified using Fauvel's key are likely to belong to *Galathowenia oculata*.

### ***Owenia fusiformis* Delle Chiaje, 1844**

**Notes:** Species complex. *Owenia fusiformis* was long believed to be a cosmopolitan species but in fact contains a number of pseudo-cryptic species worldwide (e.g. Martín et al. 2006). *Owenia fusiformis* sensu stricto was originally described from Sicily and currently no evidence exists for the presence of more than one species of the complex in the Mediterranean.

## **Paralacydoniidae Pettibone, 1963**

### ***Paralacydonia paradoxa* Fauvel, 1913**

**Notes:** Type locality: Mediterranean (Monaco).

## **Paraonidae Cerruti, 1909**

### ***Aricidea (Acmira) assimilis* Tebble, 1959**

**Nomenclature:** *Aricidea assimilis* Tebble, 1959 | *Aricidea mutabilis* Laubier & Ramos, 1974

**Notes:** *Aricidea assimilis* was included in the checklist by Simboura and Nicolaidou (2001) as separate from *Aricidea mutabilis*, but the species are considered synonymous by Katzmann and Laubier (1975) and *Aricidea mutabilis* has not been re-



instated since. The species has often been misidentified as *Aricidea (Acmira) lopezi* Berkeley & Berkeley, 1956 (see notes there) and may in fact be a complex of species (Langeneck 2017). Type locality: Mediterranean (Israel).

### ***Aricidea (Acmira) catherinae* Laubier, 1967**

**Nomenclature:** *Aricidea catherinae* Laubier, 1967 | *Aricidea (Acmira) catherinae* Laubier, 1967

**Notes:** Species complex. Molecular and morphological analyses by Langeneck (2017) revealed that specimens from the Mediterranean identified as *Aricidea (Acmira) catherinae* contain at least three pseudocryptic species which can be differentiated genetically, morphologically and on the basis of ecological characters. Type locality: Mediterranean (Banyuls-sur-Mer, France).

### ***Aricidea (Acmira) cerrutii* Laubier, 1966**

**Nomenclature:** *Aricidea (Acmira) cerrutii* Laubier, 1966 | *Aricidea cerrutii* Laubier, 1966 | *Aricidea jeffreysii* (McIntosh, 1879) [sensu Cerruti / sensu Fauvel] | *Paraonis paucibranchiata* Cerruti, 1909

**Notes:** *Aricidea jeffreysii* sensu Cerruti, 1909 (non McIntosh, 1879), on which the description and illustrations of Fauvel (1927) are based, was referred to *Aricidea (Acmira) cerrutii* by Laubier (1967). *Paraonis paucibranchiata* is considered a juvenile form by Strelzov (1973) who placed it into synonymy with *Aricidea (Acmira) cerrutii*. However, *Paraonis paucibranchiata* in this case would have nomenclatural priority over *Aricidea cerrutii* — an issue which is not resolved here. Type locality: Mediterranean (Albères, France).

### ***Aricidea (Acmira) lopezi* Berkeley & Berkeley, 1956**

**Nomenclature:** *Aricidea fauveli* Hartman, 1957 | *Aricidea lopezi lopezi* Berkeley & Berkeley, 1956

**Notes:** Questionable status. Records from Greece exist almost exclusively under the name *Aricidea fauveli*, despite the fact that they were recorded after Strelzov (1973) had synonymised the two species. It is therefore likely that they were identified using the key by Day (1967) which only includes *Aricidea fauveli*. Castelli (1987) attributes several Mediterranean records of *Aricidea lopezi* to *Aricidea (Acmira) assimilis* Tebble, 1959, which he considers synonymous with *Aricidea fauveli*. In the Mediterranean, *Aricidea lopezi* / *Aricidea fauveli* used to be distinguished from *Aricidea assimilis* on the basis of the length of the antenna (short in *Aricidea lopezi*, long in *Aricidea assimilis*) but molecular analyses by Langeneck (2017) revealed two clades related to depth which were incongruent with the two different morphotypes. It is therefore likely that Mediterranean specimens of *Aricidea (Acmira) lopezi* belong to different species.

### ***Aricidea (Acmira) neosuecica* Hartman, 1965 sensu Laubier & Ramos, 1974**

**Nomenclature:** *Aricidea neosuecica* Hartman, 1965 | *Aricidea neosuecica* Hartman, 1965 sensu Laubier & Ramos, 1974

**Notes:** Questionable status. Amongst the Greek reports of the species, only Arvanitidis (1994) explicitly refers his specimens to *Aricidea neosuecica* sensu Laubier and Ramos 1974 (non Hartman, 1965). Laubier and Ramos' description of a single specimen of *Aricidea* cf. *neosuecica* differs substantially from that of Hartman (1965b) from New England (Arvanitidis 1994) and could constitute a juvenile of *Aricidea (Acmira) simonae* Laubier & Ramos 1974 (Hartley 1981), whereas Hartman's species is referred to *Aricidea (Acmira) simplex* Day, 1963 by Aguirrezabalaga (2012). The actual identity of the Greek specimens is currently unknown.

### ***Aricidea (Acmira) simonae* Laubier & Ramos, 1974**

**Nomenclature:** *Aricidea simonae* Laubier & Ramos, 1974 | *Aricidea (Acmira) simonae* Laubier & Ramos, 1974

**Notes:** Type locality: Mediterranean (Adriatic).

### ***Aricidea (Acmira) simplex* Day, 1963**

**Nomenclature:** *Aricidea (Acesta) simplex* Day, 1963 | *Aricidea (Acmira) simplex* Day, 1963 | *Aricidea simplex* Day, 1963 | *Aricidea suecica simplex* Day, 1963

**Notes:** The species is poorly defined and, as a result, has been reported from a variety of habitats and locations worldwide; Mediterranean specimens probably belong to *Aricidea simonae* (Langeneck 2017).

### ***Aricidea (Aricidea) capensis* Day, 1961**

**Nomenclature:** *Aricidea (Aricidea) capensis* Day, 1961 | *Aricidea capensis* Day, 1961

**Notes:** Questionable status. *Aricidea (Aricidea) capensis* is usually reported from the Mediterranean as the subspecies *Aricidea (Aricidea) capensis bansei* Laubier & Ramos, 1974, whereas *Aricidea capensis* is considered to be restricted to South Africa. Arvanitidis (1994) refers the Greek records of *Aricidea capensis* by Bogdanos and Satsmadjis (1983) and Karakassis (1991) to *Aricidea (Aricidea) capensis bansei*. However, *Aricidea capensis* is frequently reported from Greece at species level (Simboura and Nicolaidou 2001) and Conides et al. (1999) report both *Aricidea capensis* and *Aricidea capensis bansei* in the same study.

***Aricidea (Aricidea) capensis bansei* Laubier & Ramos, 1974**

**Nomenclature:** *Aricidea (Aricidea) capensis bansei* Day, 1961 | *Aricidea capensis bansei* Laubier & Ramos, 1974

**Notes:** Type locality: Mediterranean (Baie de Rosas, Spain).

***Aricidea (Aricidea) longicirrata* Hartmann-Schröder, 1965**

**Nomenclature:** *Aricidea longicirrata* Hartmann-Schröder, 1965

**Notes:** Questionable status. Reported from Greece by Xenopoulou (1987), no other Mediterranean records. Distributed along the Pacific coast of tropical and South America (de León-González et al. 2006).

***Aricidea (Aricidea) longobranchiata* Day, 1961**

**Nomenclature:** *Aricidea (Aricidea) longobranchiata* Day, 1961 | *Aricidea longobranchiata* Day, 1961

**Notes:** Questionable status. In the Mediterranean reported from Greece (e.g. Bogdanos and Satsmadjis 1983, Bogdanos and Satsmadjis 1987, Simboura 1996; full reference list in Suppl. material 2) and France (Grémare et al. 1998) Records from Turkey belong to *Aricidea (Acmira) assimilis* Tebble, 1959 (Çınar et al. 2014). Otherwise distributed in tropical waters of the Atlantic and Indian Ocean and on the coasts of South Africa.

***Aricidea (Aricidea) minima* Strelzov, 1973**

**Nomenclature:** *Aricidea minima* Strelzov, 1973

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Spiliotis 1991, Zenetos et al. 1997). Originally described from the Patagonian Shelf.

***Aricidea (Aricidea) minuta* Southward, 1956**

**Nomenclature:** *Aricidea (Aricidea) minuta* Southward, 1956 | *Aricidea minuta* Southward, 1956

**Notes:** Reported from Greece by Simboura (1996), in the Mediterranean also reported from the coasts of Spain (Laubier and Ramos 1974). Otherwise distributed along the Atlantic coasts of Europe.

### ***Aricidea (Aricidea) pseudoarticulata* Hobson, 1972**

**Nomenclature:** *Aricidea (Aricidea) fragilis mediterranea* Laubier & Ramos, 1974 | *Aricidea fragilis mediterranea* Laubier & Ramos, 1974 | *Aricidea pseudoarticulata* Hobson, 1972

**Notes:** *Aricidea fragilis mediterranea* – a Mediterranean species originally described from southern France – was synonymised with *Aricidea pseudoarticulata* by Aguirrezabalaga and Gil (2008) but without examination of type material. This synonymy extends the distribution range of *Aricidea pseudoarticulata* to nearly cosmopolitan, including the Pacific (native range of *Aricidea pseudoarticulata*) and Atlantic Oceans and the Mediterranean Sea.

### ***Aricidea (Aricidea) rosea* Reish, 1968**

**Nomenclature:** *Aricidea (Acmira) lopezi rosea* Reish, 1968 | *Aricidea lopezi rosea* Reish, 1968

**Notes:** Questionable status. In the Mediterranean only reported from Greece (NCMR 1989, Simboura 1996; based on the same specimens). Distributed along the Pacific coasts of America.

### ***Aricidea (Aricidea) wassi* Pettibone, 1965**

**Nomenclature:** *Aricidea wassi* Pettibone, 1965 | *Aricidea (Aricidea) wassi* Pettibone, 1965

**Notes:** Reported from Greece by Simboura and Zenetos (2005). In the Mediterranean also reported from Italy (Castelli et al. 2008), Egypt (Abd-Elnaby 2009), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise reported from the North Atlantic and North Pacific (Hartley 1981).

### ***Aricidea (Strelzovia) belgicae* (Fauvel, 1936)**

**Nomenclature:** *Aedicira belgicae* (Fauvel, 1936) | *Paraonis belgicae* Fauvel, 1936

**Notes:** Questionable status. The species has a complex taxonomic history. Most reports and taxonomic accounts are not based on the holotype, but on a re-description by Monro (1939), which was later found to be based on non-type specimens belonging to two different species (Hartley 1984, López 2008). *Aricidea (Strelzovia) belgicae* had been considered eurybathic and cosmopolitan, but is in fact considered restricted to the deep waters of the Antarctic and most reports (including those from Greece, see Suppl. material 2 for a full list) must be considered misidentifications (López 2008).

***Aricidea (Strelzovia) claudiae* Laubier, 1967**

**Nomenclature:** *Aricidea (Allia) claudiae* Laubier, 1967 | *Aricidea claudiae* Laubier, 1967

**Notes:** Type locality: Mediterranean (Banyuls-sur-Mer, France).

***Aricidea (Strelzovia) curviseta* Day, 1963**

**Nomenclature:** *Aricidea (Allia) curviseta* Day, 1963 | *Aricidea curviseta* Day, 1963

**Notes:** Questionable status. In the Mediterranean only reported from Greece (e.g. Bogdanos and Satsmadjis 1983, Bogdanos and Satsmadjis 1987, Simboursa 1996; full reference list in Suppl. material 2). Distributed in the Indian Ocean and tropical Atlantic. As the species shows a wide polymorphism of the prostomial antenna (Strelzov 1973), Mediterranean specimens could have been misidentified and belong in fact to *Aricidea (Strelzovia) claudiae* Laubier, 1967 (J. Langeneck, pers. comm).

***Aricidea (Strelzovia) hartmani* Strelzov, 1968**

**Nomenclature:** *Aricidea (Allia) hartmani* Strelzov, 1968 | *Aricidea hartmani* Strelzov, 1968

**Notes:** Questionable status. In the Mediterranean only reported from Greece (NCMR 1992, Simboursa 1996, Zenetos et al. 1997, Pancucci-Papadopoulou et al. 1999); these could in fact belong to the Mediterranean species *Aricidea (Strelzovia) mariannae* Katzmann & Laubier, 1975, which shows similarities to *Aricidea (Strelzovia) hartmani*, especially in shape and length of the antenna (J. Langeneck, pers. comm). *Aricidea (Strelzovia) hartmani* is commonly distributed in Arctic regions.

***Aricidea (Strelzovia) mediterranea* (Laubier & Ramos, 1974)**

**Nomenclature:** *Aedicira mediterranea* Laubier & Ramos, 1974

**Notes:** The species was described as including two different morphological forms (based on antennal shape) by Laubier and Ramos (1974) from the Western Mediterranean. Aguirrezabalaga and Gil (2009) revised the species and split *Aricidea (Strelzovia) mediterranea* into two species based on these forms, *Aricidea (Strelzovia) mediterranea* and *Aricidea (Strelzovia) sardai* Aguirrezabalaga & Gil, 2009. The records from Greece could belong to either of the two species and require further examination.

***Aricidea (Strelzovia) monicae* Laubier, 1967**

**Nomenclature:** *Aricidea (Allia) monicae* Laubier, 1967 | *Aricidea monicae* Laubier, 1967

**Notes:** Reported from Greece by Koulouri et al. (2015). Type locality: Mediterranean (Banyuls-sur-Mer, France).

### ***Aricidea (Strelzovia) quadrilobata* Webster & Benedict, 1887**

**Nomenclature:** *Aricidea quadrilobata* Webster & Benedict, 1887

**Notes:** Questionable status. Reported from Greece by Dounas (1986). In the Mediterranean also reported from Italy (Castelli et al. 2008), the Adriatic (Mikac 2015) and the Malta Escarpment (Langeneck et al. 2017b), otherwise known from the North Atlantic. *Aricidea (Strelzovia) quadrilobata* sensu Strelzov (1973) was referred to *Aricidea (Strelzovia) antennata* Annenkova, 1934 by Aguirrezabalaga and Gil (2009). Thus, if Strelzov's work was used to identify the Greek specimens of *Aricidea (Strelzovia) quadrilobata*, they may actually correspond to *Aricidea antennata*. In addition, as the species has so far only been reported in grey literature and has not been found in over three decades, its presence in Greece is here considered questionable.

### ***Aricidea (Strelzovia) suecica* Eliason, 1920**

**Nomenclature:** *Aricidea suecica* Eliason, 1920

**Notes:** Questionable status. Usually reported in the Mediterranean as the subspecies *Aricidea suecica meridionalis* Laubier & Ramos, 1974, but Dando et al. (1995) and Koulouri et al. (2015) report *Aricidea suecica* at species level.

### ***Aricidea (Strelzovia) suecica meridionalis* Laubier & Ramos, 1974**

**Nomenclature:** *Aricidea suecica meridionalis* Laubier & Ramos, 1974

**Notes:** Type locality: Mediterranean (Baie de Rosas, Spain).

### ***Cirrophorus branchiatus* Ehlers, 1908**

**Nomenclature:** *Cirrophorus branchiatus* Ehlers, 1908 | *Cirrophorus lyriformis* (Annenkova, 1934)

### ***Cirrophorus furcatus* (Hartman, 1957)**

**Notes:** Questionable status. Frequently reported from the Mediterranean and from Greece, but Mediterranean specimens probably belong to different species (Castelli et al. 2008), such as the recently described *Cirrophorus nikebianchii* Langeneck, Barbieri, Maltagliati & Castelli, 2017 or *Cirrophorus turcicus* Erdoğan-Dereli, Çınar & Dağlı, 2017.

***Cirrophorus nikebianchii* Langeneck, Barbieri, Maltagliati & Castelli, 2017**

**Notes:** Reported from Greece by Langeneck et al. (2017a). Type locality: Mediterranean (Sardinia).

***Levinsenia demiri* Çınar, Dağlı & Açık, 2011**

**Notes:** Reported from Greece by Katsiaras and Simboursa (2015). Type locality: Sea of Marmara.

***Levinsenia gracilis* (Tauber, 1879)**

**Nomenclature:** *Levinsenia gracilis* (Tauber, 1879) | *Paraonis gracilis* (Tauber, 1879) | *Paraonis gracilis gracilis* (Tauber, 1879) | *Tauberia gracilis* (Tauber, 1879)

***Levinsenia multibranchiata* (Hartman, 1957)**

**Nomenclature:** *Levinsenia multibranchiata* (Hartman, 1957) | *Tauberia multibranchiata* (Hartman, 1957)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Nicolaidou et al. 1993, IOFR 1984, Papadopoulos 1986). Originally described from southern California.

***Levinsenia oculata* (Hartman, 1957)**

**Nomenclature:** *Levinsenia oculata* (Hartman, 1957) | *Paraonis gracilis oculata* Hartman, 1957

**Notes:** Questionable status. In the Mediterranean reported from Greece (e.g. Bogdanos and Satsmadjis 1983, Bogdanos and Satsmadjis 1987, Simboursa 1996; full reference list in Suppl. material 2) and Italy (Castelli et al. 2008). Italian specimens probably belong to an undescribed species (Castelli et al. 2008). Originally described from southern California.

***Levinsenia reducta* (Hartman, 1965)**

**Nomenclature:** *Levinsenia reducta* (Hartman, 1965) | *Tauberia reducta* (Hartman, 1965)

**Notes:** Questionable status. In the Mediterranean reported from Greece (e.g. Simboursa 1987, Simboursa 1996, Makra and Nicolaidou 2000, Bogdanos et al. 2002; full reference list in Suppl. material 2) and Egypt (Abd-Elnaby 2009). Commonly distributed in the Caribbean and the tropical West Atlantic.

### ***Levinsenia tribranchiata* Çınar, Dağlı & Açıık, 2011**

**Notes:** Reported from Greece by Katsiaras and Simboura (2015). Type locality: Sea of Marmara.

### ***Paradoneis armata* Glémarec, 1966**

**Nomenclature:** *Cirrophorus harpagoneus* (Storch, 1967) | *Paradoneis armata* Glémarec, 1966

**Notes:** López et al. (1987) compared type material of *Paradoneis armata* to the description of *Cirrophorus harpagoneus* (the type material of which is lost). The presumably differentiating character (acicular chaetae with subdistal teeth) is present in both species, thus López et al. (1987) consider the two species synonymous. However, as the species were described from different biogeographic regions (*Paradoneis armata* : Atlantic, *Paradoneis harpagoneus*: Red Sea) and the type material of *Paradoneis harpagoneus* is lost, not all authors share this view and examination of material from the type locality would be needed to clarify the status of the taxa (Langeneck 2017).

### ***Paradoneis drachi* Laubier & Ramos, 1974**

**Notes:** Reported from Greece by Simboura (1996) based on a single specimen, identification confirmed by A. Castelli (pers. comm. in Simboura 1996). Type locality: Mediterranean (Baie de Rosas, Spain).

### ***Paradoneis ilvana* Castelli, 1985**

**Notes:** Reported from Greece by Arvanitidis (2000a). Type locality: Mediterranean (coast of Tuscany).

### ***Paradoneis lyra* (Southern, 1914)**

**Nomenclature:** *Cirrophorus lyra* (Southern, 1914) | *Paradoneis lyra* (Southern, 1914) | *Paraonis lyra* Southern, 1914

### ***Paraonis fulgens* (Levinsen, 1884)**

**Notes:** Reported from Greece by Xenopoulou (1987), also found by Papageorgiou et al. (2006) (unpublished data). In the Mediterranean also reported from the Gulf of Lions (Laubier and Ramos 1974), Spain (Torres-Gavila 2008), Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015), otherwise distributed along the Atlantic coasts of Europe and North America.



## Pectinariidae Quatrefages, 1866

### *Amphictene auricoma* (O.F. Müller, 1776)

**Nomenclature:** *Amphictene auricoma* (O.F. Müller, 1776) | *Pectinaria* (*Amphictene*) *auricoma* (O.F. Müller, 1776) | *Pectinaria auricoma* (O.F. Müller, 1776)

### *Amphictene capensis* (Pallas, 1766)

**Nomenclature:** *Pectinaria capensis* (Pallas, 1766)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (e.g. Bogdanos and Satsmadjis 1983, Simboursa et al. 1995a, Simboursa 1996; full reference list in Suppl. material 2). Commonly distributed along the coasts of South Africa and in the Indo-Pacific.

### *Lagis koreni* Malmgren, 1866

**Nomenclature:** *Lagis koreni* Malmgren, 1866 | *Pectinaria* (*Lagis*) *koreni* (Malmgren, 1866) | *Pectinaria koreni* (Malmgren, 1866)

**Notes:** Species complex. At least two cryptic species with a strong genetic divergence exist along the north coast of France (Jolly et al. 2005, Jolly et al. 2006).

### *Lagis neapolitana* Claparède, 1869

**Nomenclature:** *Pectinaria* (*Lagis*) *neapolitana* (Claparède, 1869) | *Pectinaria neapolitana* Claparède, 1869

**Notes:** Reported from Greece by Arvanitidis (2000a). Type locality: Mediterranean (Gulf of Naples). Considered a synonym of *Lagis koreni* Malmgren, 1866 by Fauvel (1927), thus it may have been overlooked in the area. Surugiu (2006) provides an overview of the differences between *Lagis koreni* and *Lagis neapolitana*.

### *Pectinaria belgica* (Pallas, 1766)

**Nomenclature:** *Pectinaria* (*Pectinaria*) *belgica* (Pallas, 1766)

### *Petta pusilla* Malmgren, 1866

**Notes:** Reported from worldwide locations, but records from outside the North-East Atlantic probably belong to other species (Hutchings and Peart 2002).

## Phyllodocidae Ørsted, 1843

### *Eteone flava* (Fabricius, 1780)

**Notes:** Reported from Greece by Arvanitidis (2000a) based on a single specimen, in the Mediterranean also known from the western basin (Fredj 1974) and the Adriatic (Mikac 2015); otherwise distributed in the boreal Atlantic.

### *Eteone longa* (Fabricius, 1780)

**Notes:** Species complex. Consists of at least four putative cryptic species in the Arctic Ocean (Carr et al. 2011). It is unknown whether the Greek specimens belong to any of these or to a different species.

### *Eulalia bilineata* (Johnston, 1840)

**Notes:** Questionable status. Reported from Greece by Georgiou (1983) from Mazoma lagoon, Amvrakikos Gulf. In the Mediterranean also reported from the western basin (Fauvel 1923, Bellan 1964a) and the Adriatic (Mikac 2015); otherwise distributed in the North Atlantic. Taking into account its North Atlantic distribution, the rarity of Mediterranean records and the fact that, despite regular monitoring of the specific lagoonal system, the species has never been found again, the record must be considered questionable.

### *Eulalia clavigera* (Audouin & Milne Edwards, 1833)

**Notes:** Reported from Greece by Keklikoglou et al. (2013) and Maidanou et al. (2017). In the Mediterranean also known from Cyprus (Çınar 2005), Tunisia (Zaâbi 2013), Turkey (Çınar et al. 2014) and Egypt (Dorgham et al. 2014), otherwise distributed along the Atlantic coasts of Europe. Possibly underreported, as Bonse et al. (1996) refer the description and illustrations of *Eulalia viridis* (Linnaeus, 1767) by Fauvel (1923) to *Eulalia clavigera*, thus individuals of *Eulalia viridis* identified using Fauvel's publication probably belong to *Eulalia clavigera*.

### *Eulalia tripunctata* McIntosh, 1874

**Notes:** Reported from Greece by Arvanitidis (2000a). In the Mediterranean also reported from France (e.g. Bellan 1964a), Spain (Campoy 1982), Israel (Fishelson 2000), Cyprus (Çınar 2005), Italy (Castelli et al. 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed along the Atlantic coasts of Europe.

***Eulalia viridis* (Linnaeus, 1767)**

**Notes:** Species complex. *Eulalia viridis* contains at least two different species in northern Europe (Bonse et al. 1996). Greek specimens could belong to either of these or to a different species; however, identifications made using Fauvel (1923) probably belong to *Eulalia clavigera* (Audouin & Milne Edwards, 1833) (see remarks there).

***Eumida punctifera* (Grube, 1860)**

**Nomenclature:** *Eulalia punctifera* (Grube, 1860) | *Eumida punctifera* (Grube, 1860) | *Pirakia punctifera* (Grube, 1860)

***Eumida sanguinea* (Ørsted, 1843)**

**Nomenclature:** *Eulalia sanguinea* Ørsted, 1843 | *Eumida sanguinea* (Ørsted, 1843)

**Notes:** Species complex. *Nygren and Pleijel (2011)* recovered ten cryptic species in European specimens identified as *Eumida sanguinea*. They restrict the distribution of *Eumida sanguinea* to the Skagerrak, Kattegat and southern England. Five cryptic species were discovered from the Mediterranean: *Eumida asterope* Nygren & Pleijel 2011, *Eumida maia* Nygren & Pleijel 2011, *Eumida merobe* Nygren & Pleijel 2011, *Eumida* sp. (all from southern France and morphologically identical to *Eumida sanguinea* except for their pigmentation patterns) and *Eumida taygete* Nygren & Pleijel 2011 from Croatia, (morphologically identical to *Eumida sanguinea sensu stricto*). These new species are almost impossible to identify based on their morphology, thus the authors recommend the use of the name "*Eumida sanguinea sensu lato*" or "*Eumida sanguinea complex*" for all records which are not resolved through molecular methods.

***Hesionura coineai* (Laubier, 1962)**

**Notes:** New record for Greece. Three specimens, Elafonisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand; 3 specimens, Pachia Ammos, Crete, 35°06'39.6"N, 25°48'32.4"E, 1–5 m depth, fine to coarse sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Alós (2004). Type locality: Mediterranean (Argèles, France).

***Hesionura elongata* (Southern, 1914)**

**Nomenclature:** *Hesionura elongata* (Southern, 1914) | *Pseudomystides elongata* (Southern, 1914)

***Hypereteone foliosa* (Quatrefages, 1865)**

**Nomenclature:** *Eteone lactea* Claparède, 1868

**Notes:** Reported from Greece by Simboura (1996) and NCMR (1997) as *Eteone lactea*, a Mediterranean species (type locality Gulf of Naples), which was synonymised with *Hypereteone foliosa* by Pleijel (1991).

***Krohnia lepidota* (Krohn, 1845)**

**Nomenclature:** *Callizonella lepidota* (Krohn, 1845)

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). Type locality: Mediterranean (Sicily).

***Mysta picta* (Quatrefages, 1866)**

**Nomenclature:** *Eteone picta* Quatrefages, 1866 | *Mysta picta* (Quatrefages, 1866)

***Mysta siphodonta* (Delle Chiaje, 1822)**

**Nomenclature:** *Eteone siphonodonta* Claparède, 1868 | *Eteone siphodonta* (Delle Chiaje, 1822) | *Eteone (Mysta) siphonodonta* (Claparède, 1868) | *Mysta siphonodonta* (Claparède, 1868)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Mystides borealis* Théel, 1879**

**Notes:** Questionable status. Frequently reported from the Mediterranean (e.g. Castelli et al. 2008, Mikac 2015), but Pleijel and Dales (1991) and Pleijel (1993) consider the species to be restricted to the Arctic and the Mediterranean specimens to belong to *Mystides caeca*. Specimens from Cyprus were re-identified as *Mystides caeca* Langerhans, 1880 by Çinar (2005).

***Mystides caeca* Langerhans, 1880**

**Notes:** Arvanitidis (2000a) lists Aegean records of *Mystides borealis* Théel, 1879 under the name *Mystides caeca*, following Pleijel and Dales (1991) and Pleijel (1993) who consider Mediterranean specimens to belong to *Mystides caeca*. No published reports of *Mystides caeca* exist from Greece but the presence of the species has been confirmed recently (K. Keklikoglou, unpublished data).

***Naiades cantrainii* Delle Chiaje, 1828**

**Nomenclature:** *Alciopa cantraini* (Delle Chiaje, 1828)

**Notes:** Reported from Greece by Wesenberg-Lund (1939a) and Gogou (2009). Type locality: Mediterranean (Gulf of Naples).

***Nereiphylla paretti* Blainville, 1828**

**Nomenclature:** *Nereiphylla paretti* Blainville, 1828 | *Phyllodoce paretti* (Blainville, 1828)

**Notes:** Reported from Greece by Vamvakas (1971) and NCMR (2000b). In the Mediterranean also known from France (Bellan 1964a), Spain (Méndez and Cardell 1996), Cyprus (Çınar 2005), Italy (Castelli et al. 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015); otherwise distributed along the Atlantic coasts of Europe and in the Black Sea.

***Nereiphylla pusilla* (Claparède, 1870)**

**Nomenclature:** *Genetyllis nana* (de Saint Joseph, 1908) | *Nereiphylla pusilla* (Claparède, 1870) | *Paranaitis pusilla* (Claparède, 1870)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Nereiphylla rubiginosa* (Saint-Joseph, 1888)**

**Nomenclature:** *Eulalia rubiginosa* (de Saint Joseph, 1888) | *Genetyllis rubiginosa* (de Saint-Joseph, 1888) | *Nereiphylla rubiginosa* (de Saint-Joseph, 1888) | *Phyllodoce rubiginosa* de Saint-Joseph, 1888

***Notophyllum foliosum* (Sars, 1835)**

**Notes:** Species complex. At least two cryptic species exist in Scandinavia, differentiated by depth preference and colour pattern (Nygren et al. 2010a). The deeper form was described as *Notophyllum crypticum* Nygren, Eklöf & Pleijel, 2010. The authors do not exclude further cryptic species in the complex.

***Paranaitis kosteriensis* (Malmgren, 1867)**

**Notes:** Reported from Greece by Eleftheriou et al. (1990) and Karakassis (1991). In the Mediterranean also known from France (Guille and Laubier 1966), Italy (Castelli et al. 2008) and Spain (Torres-Gavila 2008), otherwise distributed along the Atlantic coasts of Europe. Confirmed from Sicily by Kato and Pleijel (2003).

***Phyllodoce groenlandica* Ørsted, 1842**

**Notes:** Questionable status. Species complex. *Phyllodoce groenlandica* has been shown to comprise at least two different cryptic species in Canadian waters (Carr et al. 2011). Pleijel (1988) considers the species to be restricted to circumboreal areas.

***Phyllodoce laminosa* Savigny in Lamarck, 1818**

**Notes:** Type locality: Mediterranean (Nice, France).

***Phyllodoce lineata* (Claparède, 1870)**

**Nomenclature:** *Anaitides lineata* (Claparède, 1870) | *Paranaitis lineata* (Claparède, 1870) | *Phyllodoce (Anaitides) lineata* (Claparède, 1870) | *Phyllodoce (Paranaitis) lineata* (Claparède, 1870) | *Phyllodoce lineata* (Claparède, 1870)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Phyllodoce longipes* Kinberg, 1866**

**Notes:** Reported from Greece by Maidanou et al. (2017). In the Mediterranean also known from the Adriatic (Mikac 2015, records confirmed by F. Pleijel and A. Nygren). Otherwise widely distributed (Chile, Gulf of Mexico, North Atlantic) and, although specimens from these location do not differ morphologically, they probably belong to different species (Nygren and Pleijel 2015).

***Phyllodoce maculata* (Linnaeus, 1767)**

**Nomenclature:** *Anaitides maculata* (Linnaeus, 1767) | *Phyllodoce maculata* (Linnaeus, 1767)

**Notes:** Easily confused with *Phyllodoce mucosa* Ørsted, 1843; probably restricted to the Atlantic coasts of Europe, the Arctic and the Sea of Japan (Pleijel 1988).

***Phyllodoce madeirensis* Langerhans, 1880**

**Nomenclature:** *Anaitides madeirensis* (Langerhans, 1880) | *Phyllodoce madeirensis* Langerhans, 1880

**Notes:** Species complex. Recent molecular analyses by Ravara et al. (2017) recover three distinct putative cryptic species from specimens identified as *Phyllodoce madeirensis* in the North-East Atlantic.

***Phyllodoce mucosa* Ørsted, 1843**

**Nomenclature:** *Anaitides mucosa* (Ørsted, 1843) | *Phyllodoce* (*Anaitides*) *mucosa* Ørsted, 1843 | *Phyllodoce mucosa* Ørsted, 1843

**Notes:** Easily confused with *Phyllodoce maculata* (Linnaeus, 1767); probably restricted to the Atlantic coasts of Europe to the White Sea (Pleijel 1988).

***Phyllodoce rosea* (McIntosh, 1877)**

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016) based on a single specimen. In the Mediterranean also reported from Spain (San Martín et al. 1981a), Italy (Castelli et al. 2008), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015). Otherwise distributed along the Atlantic coasts of Europe (Pleijel 1988).

***Pseudomystides limbata* (Saint-Joseph, 1888)**

**Nomenclature:** *Mystides limbata* Saint-Joseph, 1888 | *Pseudomystides limbata* (Saint-Joseph, 1888)

***Pseudomystides limbata nigrolineata* (Rioja, 1925)**

**Notes:** Reported from Greece by Papadopoulos (1986) and Nicolaidou et al. (1993). In the Mediterranean also known from Spain (López García 1995); otherwise known from the Atlantic coast of the Iberian Peninsula. Reports of *Pseudomystides limbata* (Saint-Joseph, 1888) could comprise specimens belonging to *Pseudomystides limbata nigrolineata*.

***Pterocirrus macroceros* (Grube, 1860)**

**Nomenclature:** *Eulalia macroceros* (Grube, 1860) | *Eulalia* (*Pterocirrus*) *macroceros* (Grube, 1860) | *Pterocirrus macroceros* (Grube, 1860) |

**Notes:** Type locality: Mediterranean (Adriatic).

***Rhynchonereella moebii* (Apstein, 1893)**

**Nomenclature:** *Callizona moebii* Apstein, 1893

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). Type locality: Mediterranean (Sicily).

***Torrea candida* (Delle Chiaje, 1841)**

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). Type locality: Mediterranean (Sicily).

***Vanadis crystallina* Greeff, 1876**

**Notes:** Questionable status. Reported from Greece by Wesenberg-Lund (1939a). Type locality: Mediterranean (Gulf of Naples). However, Wesenberg-Lund's depiction of *Vanadis crystallina* actually shows *Vanadis minuta* Treadwell, 1906 (Dales 1957). Without re-examination of Wesenberg-Lund's specimens, it remains unclear which of them belong to *Vanadis crystallina* and which to *Vanadis minuta*.

***Vanadis formosa* Claparède, 1870**

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). Type locality: Mediterranean (Gulf of Naples).

**Pilargidae de Saint-Joseph, 1899*****Ancistrosyllis falcata* (Day, 1957)**

**Nomenclature:** *Ancistrosyllis falcata* (Day, 1957) | *Pilargis falcata* Day, 1957

**Notes:** Questionable status. In the Mediterranean only reported from Greece (e.g. Bogdanos and Satsmadjis 1983, Bogdanos and Nicolaidou 1985, Bogdanos and Satsmadjis 1987, Nicolaidou and Papadopoulou 1989; full reference list in Suppl. material 2). Commonly distributed in South Africa and the Indian Ocean.

***Ancistrosyllis fiononii* Fiege & Böggemann, 1999**

**Notes:** One specimen from Greece in the collections of the Senckenberg Museum (SMF 11218, 39°16'05.4"N, 23°43'04.8"E, 1252–1255 m depth, coll. date 1998-01-01, det. D. Fiege). No other Mediterranean records. Originally described from the North Sea.

***Ancistrosyllis groenlandica* McIntosh, 1879*****Ancistrosyllis hamata* (Hartman, 1960)**

**Nomenclature:** *Ancistargis hamata* (Hartman, 1960) | *Ancistrosyllis hamata* (Hartman, 1960)



**Notes:** Reported from Greece by Arvanitidis (1994) and Simboura (1996). In the Mediterranean also known from Spain (Campoy 1982), Israel (Ben-Eliahu 1995), Italy (Castelli et al. 2008) and Turkey (Çınar et al. 2014), otherwise known from the temperate and tropical Atlantic and from the coasts of California.

### ***Otopsis chardyi* Katzmann, 1974**

**Notes:** Reported from Greece by Arvanitidis (2000a) based on a single specimen. Type locality: Mediterranean (Adriatic).

### ***Pilargis verrucosa* Saint-Joseph, 1899**

### ***Pseudexogone dineti* (Katzmann, Laubier & Ramos, 1974)**

**Nomenclature:** *Synelmis dineti* Katzmann, Laubier & Ramos, 1974

**Notes:** Reported from Greece by Simboura (1996). Type locality: Mediterranean (Adriatic).

### ***Sigambra parva* (Day, 1963)**

**Nomenclature:** *Ancistrosyllis parva* Day, 1963 | *Sigambra parva* (Day, 1963)

**Native status:** Non-native (established)

**Notes:** Considered a synonym of *Sigambra tentaculata* (Treadwell, 1941) until it was re-established by Moreira and Parapar (2002) based on differences in the arrangement and number of the pharyngeal papillae. Still, a level of confusion concerning the species in the *Sigambra tentaculata* complex and their distribution remains. Moreira and Parapar (2002) consider *Sigambra tentaculata* to possibly be restricted to the East coast of North America and identify specimens from the Mediterranean and Atlantic coasts of Spain as *Sigambra parva* – a non-native species originally described from South Africa. On the other hand, specimens of *Sigambra parva* from the Turkish Aegean were re-identified as *Sigambra tentaculata* (Çınar et al. 2014). The taxonomic and native status of the genus in the Mediterranean is therefore in need of revision.

### ***Sigambra robusta* (Ehlers, 1908)**

**Nomenclature:** *Ancistrosyllis robusta* Ehlers, 1908

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Satsmadjis 1983, Bogdanos and Nicolaidou 1985, Bogdanos and Satsmadjis 1987, Bogdanos et al. 2002). Commonly distributed in the tropical Atlantic.

***Sigambra tentaculata* (Treadwell, 1941)**

Notes: See notes under *Sigambra parva*.

**Polycirridae Malmgren, 1867*****Amaeana trilobata* (Sars, 1863)**

Nomenclature: *Amaea trilobata* (Sars, 1863) | *Amaeana trilobata* (Sars, 1863)

***Polycirrus aurantiacus* Grube, 1860**

Notes: Type locality: Mediterranean (Cres, Croatia).

***Polycirrus denticulatus* Saint-Joseph, 1894**

Notes: Reported from Greece by Dando et al. (1995). In the Mediterranean also known from Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015). Otherwise distributed along the Atlantic coasts of Europe.

***Polycirrus latidens* Eliason, 1962**

Notes: Questionable status. In the Mediterranean only reported from Greece (Maidanou et al. 2017; identified using key and descriptions by Holthe (1986)). Distributed in the boreal regions of Europe.

***Polycirrus medusa* Grube, 1850**

Notes: Type locality: Mediterranean (neotype from Anse du Rouet, France). Specimens identified using the key and descriptions by Holthe (1986) probably belong to other species (Glasby and Hutchings 2014).

***Polycirrus plumosus* Wollebæk, 1912**

Notes: Questionable status. In the Mediterranean reported from Greece (Simboura 1987, Simboura 1996, Pancucci-Papadopoulou et al. 1999, Chatzigeorgiou et al. 2016) and the Adriatic (Mikac 2015), otherwise distributed in the North Atlantic. Several Greek specimens were re-examined and found to belong to *Polycirrus twisti* Potts, 1928 (Simboura et al. 2010), a Lessepsian migrant. It is likely that *Polycirrus plumosus* does not occur in Greece.

***Polycirrus twisti* Potts, 1928**

**Native status:** Non-native (established)

**Notes:** Originally described from the Suez Canal. Reported from Greece by Simboura (2011), found several times afterwards (HCMR, unpublished data). May have been misidentified in the past as *Polycirrus plumosus* Wollebæk, 1912 (see notes there).

**Poecilochaetidae Hannerz, 1956*****Poecilochaetus fauchaldi* Pilato & Cantone, 1976**

**Notes:** Reported from Greece by Arvanitidis (2000a). Type locality: Mediterranean (Sicily).

***Poecilochaetus fulgoris* Claparède in Ehlers, 1875*****Poecilochaetus serpens* Allen, 1904****Polynoidae Kinberg, 1856*****Acholoe squamosa* (Delle Chiaje, 1825)**

**Nomenclature:** *Acholoe astericola* (Delle Chiaje, 1841) | *Acholoe squamosa* (Delle Chiaje, 1825)

**Notes:** Opinions on the validity of the name *Acholoe squamosa* are divided. Delle Chiaje (1825) described *Nereis squamosa*, but when he moved the species to *Polynoe* he used the replacement name *Polynoe astericola* (Delle Chiaje 1841: 106) possibly to avoid confusion with Savigny's 1818 *Polynoe squamata* (Pettibone 1996). Thus, Pettibone (1996) and Barnich and Fiege (2003) use *Acholoe astericola* as the valid name. However, as Savigny's species is *Polynoe squamata* (Savigny 1818: 309), not *squamosa* and Delle Chiaje was apparently aware of this, as he uses the name *squamata* alongside *squamosa* (Delle Chiaje 1841: 57, 106). Thus, there was no secondary homonymy and, as *Nereis squamosa* is an available name, there was and is no need for a replacement name. Therefore, we here consider *Acholoe squamosa* to be the valid name (see also [WoRMS](#) for additional information). Reported from Greece by Arvanitidis (1994). Type locality: Mediterranean (Gulf of Naples).

***Adyte hyalina* (G.O. Sars, 1873)**

**Nomenclature:** *Adyte assimilis* (McIntosh, 1874)

**Notes:** Reported from Greece by Koulouri et al. (2015). In the Mediterranean also known from Italy (Bock et al. 2010), Spain (Núñez et al. 2011) and Turkey (Çınar et al. 2014), otherwise distributed in the North East Atlantic and the North Sea (Bock et al. 2010).

***Antinoe aequiseta* Kinberg, 1856**

**Nomenclature:** *Harmothoe aequiseta* (Kinberg, 1856)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Papadopoulos 1986). Distributed along the coasts of South Africa and in the Indian Ocean.

***Bathyfauvelia affinis* (Fauvel, 1914)**

**Nomenclature:** *Bathyfauvelia affinis* (Fauvel, 1914) | *Macellicephala annae* Reys, 1968

**Notes:** Reported as *Macellicephala annae* by Chardy et al. (1973) from deep waters (2000–3000 m) in the Aegean and Ionian Sea, without giving a precise location. It is highly likely that the species was found in the Aegean in Greek waters, as such depths do not occur near the Turkish coast. Pettibone (1976) synonymised the two species after having examined the holotype of *Macellicephala annae* and Arvanitidis (2000a), accordingly, uses the name *Bathyfauvelia affinis* in his checklist. Type locality: Western Mediterranean.

***Eunoe assimilis* McIntosh, 1924**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (NCMR 1992). Originally described from South Africa.

***Eunoe nodosa* (M. Sars, 1861)**

**Notes:** Species complex. The exact distribution of the species in the Mediterranean is still unknown. Frequently reported from Greece in ecological lists but Barnich and Fiege (2003) consider their record of the species from the Western Mediterranean to be the first from the region. *Barnich and Fiege (2010)* state the distribution of the species to be "Arctic Ocean, North Atlantic and North Pacific". *Eunoe nodosa* has recently been shown to constitute a species complex comprising at least two cryptic species in the Arctic Ocean (Carr et al. 2011); whether the Greek specimens belong to either of these or to a different species is currently unknown.

***Gattyana cirrhosa* (Pallas, 1766)**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Satsmadjis 1983, Bogdanos and Diapoulis 1984, NCMR 1989), Spain (Desbruyères et al. 1973) and Italy (Castelli et al. 2008) otherwise known from boreal regions and cold water areas along the French and Spanish Atlantic coasts (Dauvin et al. 2006, Núñez et al. 2012); its presence in the Mediterranean could not yet be confirmed (Barnich and Fiege 2003). It is easily confused with *Harmothoe antilopes* McIntosh, 1876, a species with similar elytral structures but different notochaetae (Barnich and Fiege 2000a).

***Harmothoe antilopes* McIntosh, 1876*****Harmothoe areolata* (Grube, 1860)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Harmothoe extenuata* (Grube, 1840)**

**Nomenclature:** *Harmothoe extenuata* (Grube, 1840) | *Lagisca extenuata* (Grube, 1840)

**Notes:** Easily confused with other species of *Harmothoe*, literature records from the Mediterranean and North-East Atlantic should be treated with care (Barnich and Fiege 2000a).

***Harmothoe fraserthomsoni* McIntosh, 1897**

**Notes:** Two specimens of Arvanitidis (1994) from the Aegean station 45 were re-identified as *Harmothoe impar* (Johnston 1839) and *Harmothoe imbricata* (Linnaeus, 1767) (Barnich and Fiege 2000a).

***Harmothoe gilchristi* Day, 1960**

**Notes:** Reported from Greece by Barnich and Fiege (2000a) based on two unidentified specimens from the 1964 Calypso expedition. However, *Harmothoe* species are easily confused and *Harmothoe gilchristi* could be underreported (Barnich and Fiege 2000a).

***Harmothoe imbricata* (Linnaeus, 1767)**

**Nomenclature:** *Harmothoe imbricata* (Linnaeus, 1767)

**Notes:** Species complex. At least 6 different putative cryptic species – of which two occur in the Atlantic, four in the Arctic and three in the Pacific – were found by Carr et al. (2011). Nygren et al. (2010b) showed that different colour morphs in Scandinavia all

belong to the same species. Whether the Mediterranean specimens belong to the same species as those from northern Europe is currently unknown.

### ***Harmothoe impar* (Johnston, 1839)**

**Nomenclature:** *Harmothoe impar* (Johnston, 1839) | *Harmothoe reticulata* (Claparède, 1870)

**Notes:** The description of *Harmothoe impar* by Fauvel (1923) partly refers to *Harmothoe pagenstecheri* Michaelsen, 1896. Thus, specimens identified using Fauvel's key could belong to either species (Barnich and Fiege 2000a). *Harmothoe impar* was confirmed from Greece by Barnich and Fiege (2000a). Type locality: Mediterranean (Gulf of Naples).

### ***Harmothoe longisetis* (Grube, 1863)**

**Notes:** The description of *Harmothoe longisetis* of Fauvel (1923) partly refers to *Harmothoe glabra* (Malmgren, 1866) (Barnich and Fiege 2003, Barnich and Fiege 2009). Thus, records identified using Fauvel's key could belong to either species. Recent records (e.g. Chatzigeorgiou et al. 2016) were identified using the key and descriptions by Barnich and Fiege (2003). Type locality: Mediterranean (Adriatic).

### ***Harmothoe spinifera* (Ehlers, 1864)**

**Notes:** Type locality: Mediterranean (Rijeka Bay, Adriatic).

### ***Lepidasthenia elegans* (Grube, 1840)**

**Notes:** Type locality: Mediterranean.

### ***Lepidasthenia maculata* Potts, 1910**

**Notes:** Questionable status. Easily confused with *Lepidasthenia brunnea* Day, 1960, therefore Barnich and Fiege (2003) consider the records of the species in Europe doubtful. The authors also refer the description of *Lepidasthenia maculata* by Fauvel (1923) to *Lepidasthenia brunnea*, thus it is likely that specimens identified with Fauvel's key in fact belong to *Lepidasthenia brunnea*.

### ***Lepidonotus clava* (Montagu, 1808)**

### ***Lepidonotus squamatus* (Linnaeus, 1758)**

**Notes:** Questionable status, species complex. At least two putative cryptic species exist in waters around Canada (Carr et al. 2011). Reported from Greece by NCMR

(1995) and Conides et al. (1999) but the species is easily confused with *Lepidonotus clava* (Montagu, 1808) or *Lepidonotus tenuisetosus* (Gravier, 1902) (Barnich and Fiege 2003). Records from the Adriatic could not be confirmed (Mikac 2015).

### ***Lepidonotus tenuisetosus* (Gravier, 1902)**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016). Originally from the Red Sea, in the Mediterranean also known from Israel and Egypt (Barnich and Fiege 2003), Turkey (Çinar et al. 2014) and the Adriatic (Mikac 2015).

### ***Macellicephalo laubieri* Reys, 1971**

**Notes:** Two specimens from Greece in the collections of the Senckenberg Museum (SMF 10528, 39°14'59.4"N, 23°42'31.8"E, 1208–1253 m depth, coll. date 1998-01-01, det. R. Barnich; SMF 10529, 39°14'59.4"N, 23°43'25.8"E; 1208–1253 m depth, coll. date 1997-12-31, det R. Barnich). Type locality: Mediterranean (deep waters North-East off the Balearic Islands).

### ***Malmgrenia andreapolis* McIntosh, 1874**

**Nomenclature:** *Harmothoe lunulata* var. *andreapolis* McIntosh, 1874 | *Malmgrenia andreapolis* McIntosh, 1874 | *Malmgreniella andreapolis* (McIntosh, 1874)

**Notes:** Reported from Greece by Arvanitidis (2000a) and Koulouri et al. (2015). In the Mediterranean also known from France (Barnich and Fiege 2000a), Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015), otherwise distributed in the North East Atlantic (Barnich and Fiege 2001). Easily confused with *Malmgrenia castanea* McIntosh, 1876 and *Malmgrenia lunulata* (Delle Chiaje, 1830), thus literature records should be treated with care (Barnich and Fiege 2001).

### ***Malmgrenia castanea* McIntosh, 1876**

**Nomenclature:** *Malmgreniella castanea* (McIntosh, 1876)

**Notes:** Easily confused with *Malmgrenia lunulata* (Delle Chiaje, 1830), thus literature records should be treated with care (Barnich and Fiege 2001).

### ***Malmgrenia darbouxii* (Pettibone, 1993)**

**Nomenclature:** *Malmgreniella darbouxii* Pettibone, 1993

**Notes:** Reported from Greece by Arvanitidis (2000a) based on a single specimen. Type locality: Mediterranean (Adriatic).

***Malmgrenia ljunmani* (Malmgren, 1867)**

**Nomenclature:** *Harmothoe ljunmani* (Malmgren, 1867) | *Malmgreniella ljunmani* (Malmgren, 1867)

***Malmgrenia lunulata* (Delle Chiaje, 1830)**

**Nomenclature:** *Harmothoe lunulata* (Delle Chiaje, 1830) | *Malmgreniella lunulata* (Delle Chiaje, 1830)

**Notes:** Easily confused with congeners, thus literature records should be treated with care (Barnich and Fiege 2001). Type locality: Mediterranean (Gulf of Naples).

***Paradyte cf. crinoidicola* (Potts, 1910)**

**Native status:** Non-native (questionable)

**Notes:** Only recorded twice from the Mediterranean: Barnich and Fiege (2003) examined an old specimen in bad condition from the 1964 Calypso expedition from near Kassos Island (Greek Aegean) and identified it as *Paradyte cf. crinoidicola*. It differs from *Paradyte crinoidicola* as described by Pettibone (1969) in having blunt notochaetal tips and lower neurochaetae with straight distal regions. The authors do not rule out the possibility that these differences represent intraspecific variations and that *Paradyte crinoidicola* is indeed present in the Mediterranean (in which case it should be regarded as a Lessepsian migrant). Another record by Radić (2009) from the Adriatic as *Paradyte crinoidicola* is considered questionable by Mikac (2015) who, based on the uncertainty of the Greek record, considers the species' presence in the Mediterranean doubtful.

***Polaruschakov reyssi* Pettibone, 1976**

**Notes:** Reported from Greece by Barnich and Fiege (2003). Type locality: Mediterranean (submarine canyon off Banyuls-sur-Mer, France).

***Polynoe scolopendrina* Savigny, 1822**

**Notes:** Reported from Greece by Bogdanos et al. (2002). In the Mediterranean known from France (Bellan 1963), Spain (Campoy 1982), Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015).

***Subadyte pellucida* (Ehlers, 1864)**

**Nomenclature:** *Adyte pellucida* (Ehlers, 1864) | *Scalisetosus fragilis* (Claparède, 1868) | *Scalisetosus pellucidus* (Ehlers, 1864) | *Subadyte pellucida* (Ehlers, 1864)

**Notes:** Type locality: Mediterranean (Adriatic).



## Protodrilidae Hatschek, 1888

### *Lindrilus flavocapitatus* (Uljanin, 1877)

**Nomenclature:** *Protodrilus flavocapitatus* (Uljanin, 1877)

**Notes:** Reported from Greece by Koukouras (1979) and Dounas (1988). In the Mediterranean also known from Italy (Castelli et al. 2008) and France (Martínez et al. 2017), otherwise mainly distributed in the Black Sea. Many genus-level records of protodrilids exist in literature and in unpublished datasets from Greece; therefore it is likely that *Lindrilus flavocapitatus* is more widespread in Greece than currently known.

## Sabellariidae Johnston, 1865

### *Lygdamis muratus* (Allen, 1904)

**Nomenclature:** *Lygdamis muratus* (Allen, 1904) | *Pallasia murata* Allen, 1904

### *Phalacrostemma cidariophilum* Marenzeller, 1895

**Notes:** Originally described from various locations in Greece (detailed stations and coordinates are listed in von Marenzeller (1902)). Subsequently reported from Greece by Bellan (1964b).

### *Sabellaria alcocki* Gravier, 1906

**Nomenclature:** *Sabellaria alcocki* Gravier, 1906 | *Sabellaria spinulosa alcocki* Gravier, 1906

**Notes:** Questionable status. The specimens and description of *Sabellaria spinulosa* var. *alcocki* by Fauvel (1927) were examined by da Cunha Lana and Gruet (1989) and found to contain several species, some belonging to *Sabellaria wilsoni* Lana & Gruet, 1989. Thus, specimens identified using Fauvel's key probably belong to other species. Lezzi et al. (2015) suggest that specimens recorded as *Sabellaria alcocki* in the Mediterranean belong to juveniles of *Sabellaria spinulosa* (Leuckart, 1849) and that *Sabellaria alcocki* is absent from the region. Schimmenti et al. (2015) show through molecular analyses that specimens identified as *Sabellaria alcocki* from Italy fully matched the cytochrome c oxidase subunit I (COI) sequences of *Sabellaria spinulosa*.

### ***Sabellaria eupomatoides* Augener, 1918**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Nicolaidou and Papadopoulou 1989). Distributed in the South-East Atlantic along the coasts of Africa.

### ***Sabellaria spinulosa* (Leuckart, 1849)**

**Notes:** Could comprise records of *Sabellaria alcocki* Gravier, 1906 (see remarks there).

## **Sabellidae Latreille, 1825**

### ***Acromegalomma vesiculosum* (Montagu, 1813)**

**Nomenclature:** *Branchiomma vesiculosum* (Montagu, 1813) | *Megalomma vesiculosum* (Montagu, 1813)

**Notes:** Questionable status. Giangrande and Licciano (2008) re-examined several specimens previously identified as *Megalomma vesiculosum* from the Adriatic and Tyrrhenian Sea and Mediterranean Spain and found all of them to belong to *Megalomma lanigera* (Grube, 1846) (now *Acromegalomma lanigerum*, see Gil and Nishi 2017). They conclude that *Megalomma vesiculosum* is absent from the Mediterranean and that all Mediterranean material of *Megalomma vesiculosum* should be assigned to *Megalomma lanigera*.

### ***Amphicorina armandi* (Claparède, 1864)**

**Nomenclature:** *Amphicorina armandi* (Claparède, 1864) | *Oridia armandi* (Claparède, 1864) | *Oriopsis armandi* (Claparède, 1864)

**Notes:** Type locality: Mediterranean (Port-Vendres, France).

### ***Amphiglana mediterranea* (Leydig, 1851)**

**Notes:** Type locality: Mediterranean.

### ***Bispira fabricii* (Krøyer, 1856)**

**Nomenclature:** *Bispira fabricii* (Krøyer, 1856) | *Sabella fabricii* Krøyer, 1856

**Notes:** Frequently reported from the Mediterranean, but Knight-Jones and Perkins (1998) consider the species to be absent from the Mediterranean and restricted to Arctic areas.

***Bispira melanostigma* (Schmarda, 1861)**

**Nomenclature:** *Bispira bipunctata* (Baird, 1865) | *Sabella bipunctata* Baird, 1865

**Notes:** Questionable status. Mediterranean records are rare and considered questionable (Giangrande 1989). Knight-Jones and Perkins (1998) likewise consider Mediterranean specimens to belong to other species (e.g. the record by Iroso (1921) probably is a juvenile *Bispira mariae* (Lo Bianco, 1893)) and restrict the distribution of the species to the wider Caribbean area.

***Bispira volutacornis* (Montagu, 1804)**

**Notes:** Knight-Jones and Perkins (1998) consider the distribution of the species insufficiently known but possibly restricted to the Atlantic coasts of Europe; several Mediterranean records could belong to other species.

***Branchiomma bairdi* (McIntosh, 1885)**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Corsini-Foka et al. (2015). A West Atlantic / East Pacific species which has been reported in the Mediterranean from Italy (Giangrande et al. 2012), Malta (Arias et al. 2013b), Turkey (Çınar et al. 2014) and Tunisia (Khedhri et al. 2017).

***Branchiomma bombyx* (Dalyell, 1853)**

**Nomenclature:** *Branchiomma bombyx* (Dalyell, 1853) | *Dasychone bombyx* (Dalyell, 1853)

***Branchiomma luctuosum* (Grube, 1870)**

**Native status:** Non-native (established)

**Notes:** Reported from Greece by Arvanitidis (2000a) based on a single specimen, found again afterwards (HCMR, unpublished data). A Red Sea species has been reported in the Mediterranean from Italy (Licciano and Giangrande 2008), Spain (El Haddad et al. 2009) and Turkey (Çınar et al. 2014).

***Branchiomma lucullanum* (Delle Chiaje, 1828)**

**Nomenclature:** *Branchiomma lucullanum* (Delle Chiaje, 1828) | *Dasychone lucullana* (Delle Chiaje, 1828)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Chone duneri* Malmgren, 1867**

**Notes:** Questionable status. Tovar-Hernández et al. (2007) and Tovar-Hernández (2007) revise several species of *Chone* and restrict *Chone duneri* to the Arctic Ocean. Mediterranean specimens recorded as *Chone duneri* probably all belong to *Dialychone dunerificta* Tovar-Hernández, Licciano & Giangrande, 2007. According to Wasson et al. (2017), the genus *Chone* is absent from the Mediterranean.

### ***Chone filicaudata* Southern, 1914**

**Nomenclature:** *Chone filicaudata* Southern, 1914 | *Paradialychone filicaudata* (Southern, 1914)

**Notes:** Questionable status. Until recently, the only known species in the genus *Chone* with a pygidial cirrus used to be *Chone filicaudata*; consequently all specimens possessing it were assigned to this species (Tovar-Hernández et al. 2007). However, nowadays more species with this character are known and historic records could belong to any of these species possessing a pygidial cirrus. *Chone filicaudata* is frequently reported from the Mediterranean but without accompanying descriptions or illustrations. The only available illustration by Giangrande (1992) does not correspond to the characters of the type material of *Chone filicaudata* (Tovar-Hernández et al. 2007). The species is currently believed to be restricted to the North-East Atlantic (Tovar-Hernández et al. 2007, Wasson et al. 2017); Mediterranean specimens likely belong to *Dialychone usticensis* (Giangrande, Licciano & Castriota, 2006).

### ***Claviramus candelus* (Grube, 1863)**

**Nomenclature:** *Jasmineira candela* (Grube, 1863)

**Notes:** Reported from Greece by Antoniadou et al. (2004). Type locality: Mediterranean (Adriatic).

### ***Desdemona ornata* Banse, 1957**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Panagopoulos and Nikolaidou (1989). Originally from the Indo-Pacific region, in the Mediterranean also known from Italy (Castelli et al. 2008) and the Adriatic (Mikac 2015). Present in the adjacent Sea of Marmara (Çinar et al. 2014).

### ***Dialychone acustica* Claparède, 1870**

**Nomenclature:** *Chone acustica* (Claparède, 1870) | *Dialychone acustica* Claparède, 1870

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Dialychone arenicola* (Langerhans, 1881)**

**Nomenclature:** *Chone arenicola* Langerhans, 1881

**Notes:** Reported from Greece by NCMR (1997) in the Mediterranean also known from Italy (Castelli et al. 2008) and Turkey (Çınar et al. 2014), otherwise known from the Atlantic Ocean (Madeira).

### ***Dialychone collaris* (Langerhans, 1881)**

**Nomenclature:** *Chone collaris* Langerhans, 1881 | *Dialychone collaris* (Langerhans, 1881)

### ***Dialychone longiseta* (Giangrande, 1992)**

**Nomenclature:** *Chone longiseta* Giangrande, 1992

**Notes:** Reported from Greece by Simboura (1996) and Zenetos et al. (1997) (based on the same specimens) and Chatzigeorgiou et al. (2016). Identification of the former specimens confirmed by A. Giangrande (pers. comm. in Simboura 1996). Type locality: Mediterranean (Gulf of Taranto, Ionian Sea).

### ***Euchone capensis* Day, 1961**

**Notes:** Questionable status. In the Mediterranean reported from Greece (Arvanitidis 2000a), Turkey (Çınar et al. 2014) and Italy (Cognetti-Varriale, unpublished data in Giangrande and Licciano 2006) However, Giangrande and Licciano (2006) consider the presence of *Euchone capensis* in the Mediterranean questionable, as the species is considered to be restricted to South Africa and the few reports of the species from the Mediterranean have never been verified.

### ***Euchone pararosea* Giangrande & Licciano, 2006**

**Notes:** Reported from Greece by Katsiaras and Simboura (2015). Type locality: Mediterranean (South Tyrrhenian Sea).

### ***Euchone rosea* Langerhans, 1884**

**Notes:** According to Giangrande and Licciano (2004) the species usually occurs on coralligenous habitats and soft-bottom records in the Mediterranean could belong to other species. However, Mikac (2015) confirms the species from soft bottoms in the North Adriatic.

***Euchone rubrocincta* (Sars, 1862)*****Euchone southerni incisa* Banse, 1970**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Arvanitidis 2000a) based on a single specimen. Giangrande and Licciano (2004) do not list the species amongst the Mediterranean species of *Euchone*. Originally described from the Islas Malvinas, South Atlantic.

***Euratella salmacidis* (Claparède, 1869)**

**Nomenclature:** *Euratella salmacidis* (Claparède, 1869) | *Laonome salmacidis* Claparède, 1869

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Hypsicomus stichophthalmos* (Grube, 1863)**

**Nomenclature:** *Hypsicomus stichophthalmos* (Grube, 1863) | *Potamilla stichophthalmos* (Grube, 1863)

**Notes:** Type locality: Mediterranean.

***Jasmineira caudata* Langerhans, 1880*****Jasmineira elegans* Saint-Joseph, 1894*****Laonome kroyeri* Malmgren, 1866*****Myxicola infundibulum* (Montagu, 1808)**

**Notes:** Species complex. *Myxicola infundibulum* is a taxon with a long list of synonyms and has until recently been considered cosmopolitan (Giangrande et al. 2012). However, specimens from different parts of the world (West Atlantic, Mediterranean, Pacific Canada) show genetic and morphological differences (Carr et al. 2011). Giangrande et al. 2012 found specimens of *Myxicola* from the Gulf of Naples and Lake Faro to morphologically differ from specimens of *Myxicola infundibulum* from the type locality in the North Adriatic Sea. It is therefore possible that the species constitutes a complex of cryptic species even in the Mediterranean.

***Notaulax phaeotaenia* (Schmarda, 1861)**

**Nomenclature:** *Hypsicomus phaeotaenia* (Schmarda, 1861) | *Notaulax phaeotenia* (Schmarda, 1861)

**Notes:** *Notaulax phaeotaenia* has a long list of synonyms, as a result, the species has been reported from worldwide locations. However, many of these synonyms probably constitute valid and distinct species (Perkins 1984).

### ***Parasabella langerhansi* (Knight-Jones, 1983)**

**Nomenclature:** *Demonax langerhansi* Knight-Jones, 1983

**Notes:** Reported from Greece by Arvanitidis (2000a) and Kitsos et al. (2005). In the Mediterranean also known from Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed along the Atlantic coasts of Europe and in Madeira (Giangrande 1994).

### ***Parasabella saxicola* (Grube, 1861)**

**Nomenclature:** *Demonax brachychona* (Claparède, 1870)

**Notes:** Knight-Jones et al. (1991) synonymise *Demonax saxicola* and *Demonax brachychona* (Claparède, 1870) but erroneously use *Demonax brachychona* as the preferred name, despite Grube's name being older and thus having priority. Tovar-Hernández and Harris (2010) re-instate the use of the name *Demonax saxicola* over *Demonax brachychona* and introduce *Parasabella* Bush, 1905 as a replacement name for *Demonax* Kinberg, 1867 (which is pre-occupied by the beetle genus *Demonax* Thomson, 1860). Type locality: Mediterranean (Adriatic).

### ***Parasabella tenuicollaris* (Grube, 1870)**

**Nomenclature:** *Demonax tenuicollaris* (Grube, 1870)

**Notes:** Reported from Greece by Arvanitidis (1994). Type locality: Mediterranean (Adriatic).

### ***Potamilla torelli* (Malmgren, 1866)**

**Notes:** Questionable status. Knight-Jones (1983) re-examined the specimens of *Potamilla torrelli* of Fauvel (1927) and referred them to *Demonax brachychona* (Claparède, 1870) (now accepted as *Parasabella saxicola* (Grube, 1861)). Thus, it is likely that all specimens under the name *Potamilla torelli* which were identified using Fauvel's work belong to *Potamilla saxicola*. Species of *Demonax* (now *Parasabella*) were usually overlooked or misidentified during the last century due to a very narrow description of the genus, as specimens with broadly hooded inferior thoracic chaetae used to be referred to *Potamilla*; consequently, records of *Potamilla torelli* from the Mediterranean are probably misidentifications Giangrande (1994). The presence of *Potamilla torelli* in the Mediterranean is furthermore doubted by Knight-Jones et al. (1991) who consider it to be distributed in Arctic regions.

### ***Pseudopotamilla reniformis* (Brugière, 1789)**

**Nomenclature:** *Potamilla reniformis* (Brugière, 1789) | *Pseudopotamilla reniformis* (Brugière, 1789)

**Notes:** Questionable status. Knight-Jones et al. (2017) investigate specimens of *Pseudopotamilla reniformis* from various locations in Europe, restrict the distribution of *Pseudopotamilla reniformis* to cold northern waters and re-instate *Pseudopotamilla saxicava* (Quatrefages 1866) for specimens from temperate and tropical waters, including the Mediterranean, noting that the description of *Pseudopotamilla reniformis* by Fauvel (1927) contains figures of both *Pseudopotamilla reniformis* and *Pseudopotamilla saxicava*. Greek records of *Pseudopotamilla reniformis* probably belong to *Pseudopotamilla saxicava*.

### ***Sabella discifera* Grube, 1874**

**Nomenclature:** *Megalomma linaresi* (Rioja, 1917) | *Sabella discifera* Grube, 1874

**Notes:** Reported from Greece by Arvanitidis (1994) as *Sabella discifera*, by Fassari (1982) as *Megalomma linaresi*. Type locality of *Sabella discifera*: Mediterranean (Adriatic).

### ***Sabella pavonina* Savigny, 1822**

### ***Sabella spallanzanii* (Gmelin, 1791)**

**Nomenclature:** *Sabella spallanzanii* (Gmelin, 1791) | *Spirographis spallanzanii* (Viviani, 1805)

**Notes:** Type locality: Mediterranean (neotype from Malta).

## **Saccocirridae Czerniavsky, 1881**

### ***Saccocirrus papillocercus* Bobretzky, 1872**

**Notes:** The genus *Saccocirrus* was recently split into the two genera *Saccocirrus* Bobretzky, 1872 (without pharynx, mainly occurring in the Mediterranean and along European coasts) and *Pharyngocirrus* Di Domenico, Martínez, Lana and Worsaae, 2014 (with pharynx, in tropical latitudes) based on morphological and molecular analyses (Di Domenico et al. 2014).



## Scalibregmatidae Malmgren, 1867

### *Asclerocheilus capensis* Day, 1963

**Notes:** Questionable status. In the Mediterranean only reported from Greece (NCMR 1992, Simboura 1996). Distributed in the tropical and South Atlantic and the Indian Ocean.

### *Asclerocheilus intermedius* (Saint-Joseph, 1894)

**Notes:** Reported from Greece by Fassari (1982) and Arvanitidis (1994). In the Mediterranean also known from Cyprus (Çınar 2005), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed in the Atlantic.

### *Hyboscolex longiseta* Schmarda, 1861

**Native status:** Non-native (questionable)

**Notes:** Questionable status. Reported from Greece by Fassari (1982). In the Mediterranean also known from Tunisia (Cantone et al. 1978) and Italy (Castelli et al. 2008), otherwise distributed in the Southern Hemisphere (Kudenov and Blake 1978). Its presence in the Mediterranean is considered doubtful by Arvanitidis (1994) and Simboura et al. (2010).

### *Polyphysia crassa fauveli* (Laubier, 1959)

**Notes:** Reported from Greece by Simboura (1996) as *Polyphysia crassa* (Ørsted, 1843) but belonging in fact to the Mediterranean subspecies *Polyphysia crassa fauveli*; *Polyphysia crassa* is restricted to northern Europe (Kudenov and Blake 1978). In some unpublished datasets from Greece, the name *Lipobranchus jeffreysii* (McIntosh, 1869) is used. Støp-Bowitz (1945) considers *Lipobranchus jeffreysii*, the type of *Lipobranchius*, to be the juvenile of *Polyphysia crassa* (see also Kudenov and Blake 1978). Nevertheless, *Lipobranchus jeffreysii* continues to be listed in some works as a valid species (e.g. Hartmann-Schröder 1996).

### *Scalibregma celticum* Mackie, 1991

### *Scalibregma inflatum* Rathke, 1843

**Notes:** Species complex. *Scalibregma inflatum*, often considered cosmopolitan, actually comprises a number of very similar or sibling species (e.g. Kudenov and Blake 1978). In Europe, at least four species of *Scalibregma* are present (Mackie 1991). It is

likely that some older records of *Scalibregma inflatum* from Greece belong to *Scalibregma celticum* Mackie, 1991 (Simboura 1996).

### ***Sclerocheilus minutus* Grube, 1863**

**Notes:** Type locality: Mediterranean (Croatia).

## **Serpulidae Rafinesque, 1815**

### ***Apomatus ampulliferus* Philippi, 1844**

**Nomenclature:** *Apomatus ampulliferus* Philippi, 1844 | *Protula ampullifera* (Philippi, 1844)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Apomatus similis* Marion & Bobretzky, 1875**

**Notes:** Type locality: Mediterranean (Marseille).

### ***Bathyvermilia langerhansi* (Fauvel, 1909)**

**Notes:** Questionable status. Reported from Greece by Tselepides (1992), but its presence in the Mediterranean is questioned by Ben-Eliahu and Fiege (1996), as Zibrowius (1973a) had studied all Mediterranean specimens available to him and found that they belong in fact to other species.

### ***Bushiella (Jugaria) granulata* (Linnaeus, 1767)**

**Nomenclature:** *Jugaria granulata* (Linnaeus, 1767)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Christomanos 1963). Commonly distributed in boreal waters (Jirkov 2001).

### ***Ditrupa arietina* (O. F. Müller, 1776)**

**Nomenclature:** *Ditrupa arietina* (O.F. Müller, 1776) | *Ditrupa subulata* (Deshayes, 1825) in Berkeley, 1835

**Notes:** Hartley (2014) points out different ecological preferences between populations of *Ditrupa arietina* in the Mediterranean and in the Atlantic and a sudden increase in population densities in the Western Mediterranean in the late 1980s, indicating the possibility of an unknown cryptic invasive species.

***Ficopomatus enigmaticus* (Fauvel, 1923)**

**Nomenclature:** *Ficopomatus enigmaticus* (Fauvel, 1923) | *Mercierella enigmatica* Fauvel, 1923

**Native status:** Non-native (established)

**Notes:** Species complex. A species with an unknown native range but believed to be an introduced species in shallow waters worldwide (Styan et al. 2017) and frequently reported from Greece. Styan et al. (2017), using molecular methods, find at least three putative cryptic species of *Ficopomatus enigmaticus* in Australia.

***Filograna implexa* Berkeley, 1835**

**Notes:** Reported from Greece by Arvanitidis (1994) and Chatzigeorgiou et al. (2016). The relationships within the species of the *Filograna* / *Salmacina* complex and the characters used to distinguish species and genera are still a matter of debate (Nogueira and ten Hove 2000). *Filograna* is usually considered operculate and *Salmacina* non-operculate, but mixed colonies have been found, questioning the validity of the character. Fauvel (1927) considers the three taxa *Filograna implexa*, *Salmacina dysteri* (Huxle, 1855) and *Salmacina incrustans* Claparède, 1870 to be separate species, a view which is also tentatively followed by Nogueira and ten Hove (2000) until more information becomes available. They recommend using locally applied names for each of the species. We follow this view here and list all three species as separate, pending further taxonomic investigation. Greek specimens were identified using the key of Bianchi (1981).

***Filigranula annulata* (O. G. Costa, 1861)**

**Notes:** Reported from Greece by Zibrowius (1979) and Gerovasileiou et al. (2015).  
Type locality: Mediterranean.

***Filigranula calyculata* (O. G. Costa, 1861)*****Filigranula gracilis* Langerhans, 1884*****Filigranula stellata* (Southward, 1963)*****Hyalopomatus variorugosus* Ben-Eliahu & Fiege, 1996**

**Notes:** Reported from Greece by Ben-Eliahu and Fiege (1996) (several paratypes from deep waters off Crete) and by Sanfilippo et al. (2017) as *Hyalopomatus* cf. *variorugosus* from two caves of Lesvos Island. Type locality: Mediterranean (Levantine Basin).

### ***Hydroides brachyacantha* Rioja, 1941**

**Native status:** Non-native (casual)

**Notes:** Species complex. Reported from Greece by Ulman et al. (2017). In the Mediterranean also known from Israel (Ben-Eliahu 1991a), Turkey (Çınar 2006) and Spain (Ulman et al. 2017). Probably originating from the Mexican Pacific (Ulman et al. 2017), but the species has been shown to constitute a complex of cryptic and pseudo-cryptic species (Bastida-Zavala and ten Hove 2003, Sun et al. 2016). The identity of Mediterranean specimens, their introduction status in the Mediterranean and the potential native range cannot be determined without molecular methods.

### ***Hydroides dianthus* (Verrill, 1873)**

**Notes:** Species complex. *Hydroides dianthus* was originally described from Connecticut and long believed to be non-native in the Mediterranean (Zibrowius 1973b, Zenetos et al. 2017). A recent study by Sun et al. (2017) shows, however, that *Hydroides dianthus* could originate from the Mediterranean, as the species shows the highest genetic diversity in the Mediterranean (nine out of ten haplotypes present, as opposed to only two haplotypes in the USA). The authors recovered also a putative cryptic species, *Hydroides* cf. *dianthus*, probably originating from the Gulf of Mexico and apparently recently introduced to the Black Sea.

### ***Hydroides dirampha* Mörch, 1863**

**Nomenclature:** *Hydroides dirampha* Mörch, 1863 | *Hydroides lunulifera* (Claparède, 1870)

**Native status:** Non-native (established)

**Notes:** Originally described from the Caribbean, established throughout the Mediterranean (Zibrowius 1973b).

### ***Hydroides elegans* (Haswell, 1883)**

**Native status:** Non-native (established)

**Notes:** May be underreported in Greece, as specimens identified as *Hydroides norvegica* Gunnerus, 1768 from harbour and biofouling environments probably belong to *Hydroides elegans* (Zibrowius 1971, Arvanitidis 1994, ten Hove and Kupriyanova 2009). The species is considered introduced to the Mediterranean, although its centre of origin is still unknown (ten Hove 2003).

***Hydroides helmata* (Iroso, 1921)**

**Notes:** Reported from Greece by Arvanitidis (1994). Type locality: Mediterranean (Gulf of Naples).

***Hydroides nigra* Zibrowius, 1971**

**Notes:** Type locality: Mediterranean (Tarbaka, Tunisia).

***Hydroides norvegica* Gunnerus, 1768**

**Notes:** Zibrowius (1971) shows that the non-native species *Hydroides elegans* (Haswell, 1883) has replaced *Hydroides norvegica* in harbour fouling communities of the Mediterranean and that *Hydroides norvegica* is restricted to deeper waters. ten Hove and Kupriyanova (2009) suggest that all (sub)tropical records of *Hydroides norvegica* should be attributed to *Hydroides elegans*. In Greece, *Hydroides norvegica* has been reported both from harbour and shallow coastal communities as well as from deep waters below 500 m (e.g. Tselepides 1992). Its presence in the port of Heraklion and other Mediterranean ports has been recently confirmed (Ulman et al. 2017).

***Hydroides pseudouncinata* Zibrowius, 1968**

**Nomenclature:** *Hydroides pseudouncinata* Zibrowius, 1968 | *Hydroides pseudouncinata pseudouncinata* Zibrowius, 1968

**Notes:** Type locality: Mediterranean (Marseille).

***Hydroides stoichadon* Zibrowius, 1971**

**Notes:** Reported from Greece by Simboura (1996) and Kitsos et al. (2005). Type locality: Mediterranean (off the French coast).

***Janita fimbriata* (Delle Chiaje, 1822)**

**Nomenclature:** *Janita fimbriata* (Delle Chiaje, 1822) | *Janita (Omphalopoma) fimbriata* (Delle Chiaje, 1822) | *Omphalopomopsis fimbriata* (Delle Chiaje, 1822)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Janua heterostropha* (Montagu, 1803)**

**Nomenclature:** *Janua (Dexiospira) pagenstecheri* (Quatrefages, 1866) | *Janua pagenstecheri* (Quatrefages, 1866) | *Janua pagenstecheri gnomonica* (Bailey, 1969) | *Spirorbis (Janua) gnomonicus* Bailey, 1969 | *Spirorbis (Janua) pagenstecheri* Quatrefages, 1866

## ***Josephella marenzelleri* Caullery & Mesnil, 1896**

### ***Metavermilia acanthophora* (Augener, 1914)**

**Nomenclature:** *Metavermilia acanthophora* (Augener, 1914) | *Vermiliopsis acanthophora* Augener, 1914

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Nicolaidou 1985, Bogdanos and Satsmadjis 1987). Commonly distributed in the Indo-Pacific region (Kupriyanova et al. 2015).

### ***Metavermilia multicristata* (Philippi, 1844)**

**Nomenclature:** *Metavermilia multicristata* (Philippi, 1844) | *Vermilia multicristata* Philippi, 1844

**Notes:** Type locality: Mediterranean.

### ***Metavermilia taenia* Zibrowius, 1971**

**Notes:** Questionable status. One specimen from Greece in the collections of the Senckenberg Museum (SMF 22101, 34°57'20.34"N, 26°11'04.62"E, 107–117 m depth, coll. date 2005-08-31, det. H Zibrowius, 2005). Identification uncertain, entry indicated with a question mark. No other Mediterranean records; originally described from off the Portuguese coast.

### ***Neodexiospira pseudocorrugata* (Bush, 1905)**

**Nomenclature:** *Neodexiospira pseudocorrugata* (Bush, 1905) | *Spirorbis (Janua) parvulus* Bailey, 1969

**Notes:** Records likely also comprise those recorded under the name *Spirorbis corrugatus* which were identified using Fauvel (1927) (see also entry for *Spirorbis corrugatus* (Montagu, 1803) in Table 1 for detailed notes on taxonomy and nomenclature).

### ***Nidificaria clavus* (Harris, 1968)**

**Nomenclature:** *Nidificaria clavus* (Harris, 1968) | *Spirorbis (Pileolaria) clavus* Harris, 1968

***Pileolaria heteropoma* (Zibrowius, 1968)**

**Nomenclature:** *Pileolaria heteropoma* (Zibrowius, 1968) | *Spirorbis* (*Pileolaria*) *heteropoma* Zibrowius, 1968

**Notes:** Type locality: Mediterranean (Marseille).

***Pileolaria militaris* Claparède, 1870**

**Nomenclature:** *Pileolaria militaris* Claparède, 1870 | *Spirorbis mediterraneus* Caullery & Mesnil, 1896 | *Spirorbis* (*Pileolaria*) *militaris* (Claparède, 1870)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Placostegus crystallinus non* (Scacchi, 1836) sensu Zibrowius, 1968**

**Notes:** Greek records all refer to the species as described by Zibrowius (1968). Type locality: Mediterranean (France).

***Placostegus tridentatus* (Fabricius, 1779)**

**Notes:** The species exhibits two different morphotypes in the Atlantic and the Mediterranean. Mediterranean specimens are smaller than Atlantic specimens and show differences in morphological details of the tube (for a detailed description see Sanfilippo (2003)).

***Protis arctica* (Hansen, 1879)**

**Notes:** Reported from deep waters of Greece and the Eastern Mediterranean by Ben-Eliahu and Fiege (1996).

***Protolaeospira striata* (Quiévreux, 1963)**

**Nomenclature:** *Protolaeospira* (*Protolaeospira*) *striata* (Quiévreux, 1963) | *Protolaeospira striata* (Quiévreux, 1963) | *Spirorbis* (*Marsipospira*) *striatus* (Quiévreux, 1963)

***Protula intestinum* (Lamarck, 1818)**

**Nomenclature:** *Protula intestinum* (Lamarck, 1818) | *Serpula graeca* Brullé, 1832

**Notes:** Type locality: Mediterranean.

***Protula tubularia* (Montagu, 1803)**

***Salmacina dysteri* (Huxley, 1855)**

**Notes:** Reported from Greece by Morri et al. (1999) and Bianchi and Morri (2000b). Reported from worldwide locations but probably constitutes a complex of species (Nogueira and ten Hove 2000). See also remarks under *Filograna implexa* Berkeley, 1835.

***Salmacina incrustans* Claparède, 1870**

**Notes:** See remarks under *Filograna implexa* Berkeley, 1835. Type locality: Mediterranean (Gulf of Naples).

***Semivermilia agglutinata* (Marenzeller, 1893)**

**Nomenclature:** *Semivermilia agglutinata* (Marenzeller, 1893) | *Vermilia agglutinata* Marenzeller, 1893

**Notes:** The original description is based on specimens collected in Libya and Greece (near Kythira Island and north of Crete).

***Semivermilia crenata* (O. G. Costa, 1861)**

**Notes:** Type locality: Mediterranean.

***Semivermilia cribrata* (O. G. Costa, 1861)**

**Notes:** Reported from Greece by Ben-Eliahu and Fiege (1996); additionally two specimens from Greece in the collections of the Senckenberg Museum (SMF 21066, 37°39'48"N, 23°56'42"E, 23 m depth, coll. date July 1995; SMF 19682, near Corfu, ca. 20m depth, coll. date 1958-07-10, both det. H. Zibrowius). Type locality: Mediterranean.

***Semivermilia pomatostegoides* (Zibrowius, 1969)*****Semivermilia torulosa* (Delle Chiaje, 1822)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Serpula cavernicola* Fassari & Mòllica, 1991**

**Notes:** Reported from Greece by Sanfilippo et al. (2017). Only dead specimens of *Serpula cavernicola* were found – although in several stations – in two caves of Lesvos Island. Type locality: Mediterranean (Sicily).

***Serpula concharum* Langerhans, 1880**



***Serpula lobiancoi* Rioja, 1917*****Serpula vermicularis* Linnaeus, 1767**

**Nomenclature:** *Serpula vermicularis* Linnaeus, 1767 | *Serpula vermicularis echinata* Gmelin, 1791

***Simplaria pseudomilitaris* (Thiriot-Quévieux, 1965)**

**Nomenclature:** *Simplaria pseudomilitaris* (Thiriot-Quévieux, 1965) | *Spirorbis* (*Pileolaria*) *pseudomilitaris* Thiriot-Quévieux, 1965)

**Notes:** Type locality: Mediterranean (Villefranche-sur-Mer).

***Spiraserpula massiliensis* (Zibrowius, 1968)**

**Nomenclature:** *Serpula massiliensis* Zibrowius, 1968 | *Spiraserpula massiliensis* (Zibrowius, 1968)

**Notes:** Type locality: Mediterranean (Marseille).

***Spirobranchus lamarcki* (Quatrefages, 1866)**

**Nomenclature:** *Pomatoceros lamarckii* (Quatrefages, 1866)

***Spirobranchus polytrema* (Philippi, 1844)**

**Nomenclature:** *Pomatostegus polytrema* (Philippi, 1844) | *Spirobranchus polytrema* (Philippi, 1844)

**Notes:** Type locality: Mediterranean.

***Spirobranchus tetraceros* (Schmarda, 1861)**

**Native status:** Non-native (established)

**Notes:** Species complex with a confused taxonomic status, comprising one species in the Caribbean and several in the Indo-Pacific (Ben-Eliahu and ten Hove 2011). Reported from Greece by Ben-Eliahu (1991a) and Ulman et al. (2017). Originally from the Red Sea, in the Mediterranean also known from Egypt (Ben-Eliahu and Fiege 1996), Israel (Fishelson 2000), Cyprus (Çınar 2005), Turkey (Çınar et al. 2014) and Italy (Ulman et al. 2017).

***Spirobranchus triqueter* (Linnaeus, 1758)**

**Nomenclature:** *Pomatoceros triqueter* (Linnaeus, 1758) | *Spirobranchus triqueter* (Linnaeus, 1758)

***Spirorbis cuneatus* Gee, 1964**

**Nomenclature:** *Spirorbis (Spirorbis) cuneatus* Gee, 1964 | *Spirorbis cuneatus* Gee, 1964

***Spirorbis marioni* Caullery & Mesnil, 1897**

**Native status:** Non-native (established)

**Notes:** Reported from Greece by NCMR (1997) and Kitsos (2003). Originally described from Panama; known from harbours throughout the Mediterranean (Galil et al. 2008).

***Vermiliopsis infundibulum* (Philippi, 1844)**

**Nomenclature:** *Vermilia multivaricosa* Mörch, 1863 | *Vermiliopsis infundibulum* (Philippi, 1844)

**Notes:** Type locality: Mediterranean.

***Vermiliopsis labiata* (O. G. Costa, 1861)**

**Nomenclature:** *Vermiliopsis labiata* (O. G. Costa, 1861) | *Vermiliopsis richardi* Fauvel, 1909

**Notes:** Type locality: Mediterranean.

***Vermiliopsis monodiscus* Zibrowius, 1968**

**Notes:** Type locality: Mediterranean.

***Vermiliopsis striaticeps* (Grube, 1862)**

**Notes:** Type locality: Mediterranean.

***Vinearria endoumensis* (Zibrowius, 1968)**

**Nomenclature:** *Spirorbis (Pileolaria) endoumensis* Zibrowius, 1968 | *Vinearria endoumensis* (Zibrowius, 1968)

**Notes:** Reported from Greece by Bailey (1969). Type locality: Mediterranean (Marseille).

***Vinearia koehleri* (Caullery & Mesnil, 1897)**

**Nomenclature:** *Spirorbis* (*Pileolaria*) *koehleri* (Caullery & Mesnil, 1897) | *Vinearia koehleri* (Caullery & Mesnil, 1897)

**Notes:** Type locality: Mediterranean (La Ciotat, France).

**Siboglinidae Caullery, 1914*****Lamellibrachia anaximandri* Southward, Andersen & Hourdez, 2011**

**Notes:** Originally described from the Anaximander Seamount (Eastern Mediterranean), paratypes from the Olimpi mud volcano, south of Crete, ca. 2000 m depth. Previously recorded as *Lamellibrachia* sp. by Olu-Le Roy et al. (2004) from the Olimpi mud volcano and by Hughes and Crawford (2006) from the wreck of the SS Persia, south-east of Crete.

***Siboglinum* cf. *carpinei* Ivanov, 1970**

**Notes:** Reported from Greece by Southward et al. (2011) from the Olimpi mud volcano, south of Crete. Type locality: Mediterranean (west of Corsica).

**Sigalionidae Malmgren, 1867*****Claparedepelogenia inclusa* (Claparède, 1868)**

**Nomenclature:** *Claparedepelogenia inclusa* (Claparède, 1868) | *Psammolyce inclusa* (Claparède, 1868)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Euthalenessa oculata* (Peters, 1854)**

**Nomenclature:** *Euthalenessa oculata* (Peters, 1854) | *Thalenessa dendrolepis* (Claparède, 1868)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Fimbriosthenelais minor* (Pruvot & Racovitza, 1895)**

**Nomenclature:** *Fimbriosthenelais minor* (Pruvot & Racovitza, 1895) | *Sthenelais minor* Pruvot & Racovitza, 1895

**Notes:** Type locality: Mediterranean (Banyuls-sur-Mer).

***Fimbriosthenelais zetlandica* (McIntosh, 1876)**

**Nomenclature:** *Sthenelais papillosa* Day, 1960

***Heteropelogenia articulata* (Day, 1960)**

**Nomenclature:** *Psammolyce articulata* Day, 1960

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Makra and Nicolaidou 2000). Distributed along the coasts of South Africa and in the Indian Ocean.

***Labioleanira yhleni* (Malmgren, 1867)**

**Nomenclature:** *Labioleanira yhleni* (Malmgren, 1867) | *Leanira yhleni* Malmgren, 1867 | *Sthenolepis yhleni* (Malmgren, 1867)

***Neoleanira tetragona* (Ørsted, 1845)**

**Notes:** Questionable status. Usually occurring in muddy sediments of deep-water habitats (down to 2200 m) and rarely recorded from the Mediterranean (Barnich and Fiege 2003). The two Greek records are from shallower waters: from max. 35 m depth (Chintiroglou 1987) and from coelenteron content of the anemone *Calliactis parasitica* (Couch, 1842) collected from depths down to 80 m (Chintiroglou and Koukouras 1991) and must therefore be considered questionable.

***Pelogenia arenosa* (Delle Chiaje, 1830)**

**Nomenclature:** *Pelogenia arenosa* (Delle Chiaje, 1830) | *Psammolyce arenosa* (Delle Chiaje, 1830)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Pholoe inornata* Johnston, 1839**

**Nomenclature:** *Pholoe inornata* Johnston, 1839 | *Pholoe synophthalmica* Claparède, 1868

**Notes:** Recent phylogenetic analysis of the Aphroditiformia, taking into account morphological and molecular information, recovered species previously assigned to Pholoidae as nested within Sigalionidae (Norlinder et al. 2012, Gonzalez et al. 2017). Pholoidae were demoted to subfamilial rank (Pholoinae) within Sigalionidae by Gonzalez et al. (2017).

***Pholoe minuta* (Fabricius, 1780)**

**Notes:** Questionable status. Barnich and Fiege (2003) doubt the presence of *Pholoe minuta* in the Mediterranean and suggest that records should be attributed to *Pholoe inornata* Johnston, 1839. *Pholoe minuta* is very likely absent from Greece. See notes under *Pholoe inornata* for information on higher classification.

***Pholoides dorsipapillatus* (Marenzeller, 1893)**

**Nomenclature:** *Pholoe dorsipapillata* Marenzeller, 1893 | *Pholoides dorsipapillatus* (Marenzeller, 1893)

**Notes:** The original description is based on specimens collected in Egypt and Greece (harbour of Santorini). See notes under *Pholoe inornata* Johnston, 1839 for information on higher classification.

***Pisione remota* (Southern, 1914)**

**Nomenclature:** *Pisione remota* (Southern, 1914) | *Praegeria remota* Southern, 1914

**Notes:** Species of *Pisione* are mainly distinguished on the basis of number and shape of male copulatory organs (Yamanishi 1998, Gonzalez et al. 2017b). Juvenile specimens are often erroneously assigned to *Pisione remota* (A. Martínez, pers. comm.) and the diversity of the genus in Europe might be underestimated (Martins et al. 2012b). As at least three species of *Pisione* are known to occur in the Mediterranean (Martins et al. 2012b), more than one species of the genus could in fact be present in Greece (R. Martins, pers. comm.).

***Sigalion mathildae* Audouin & Milne Edwards in Cuvier, 1830*****Sigalion squamosus* Delle Chiaje, 1830**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Sthenelais boa* (Johnston, 1833)**

**Nomenclature:** *Sthenelais boa* (Johnston, 1833) | *Sthenelais ctenolepis* Claparède, 1868

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Sthenelais limicola* (Ehlers, 1864)**

**Notes:** Type locality: Mediterranean (Croatia).

## Sphaerodoridae Malmgren, 1867

### *Sphaerodoridium claparedii* (Greeff, 1866)

**Nomenclature:** *Sphaerodoridium claparedii* (Greeff, 1866) | *Sphaerodorum claparedii* Greeff, 1866

### *Sphaerodoridium minutum* (Webster & Benedict, 1887)

**Nomenclature:** *Sphaerodoropsis minuta* (Webster & Benedict, 1887) | *Sphaerodoridium minutum* (Webster & Benedict, 1887) | *Sphaerodorum minutum* (Webster & Benedict, 1887)

### *Sphaerodorum gracilis* (Rathke, 1843)

**Nomenclature:** *Sphaerodorum flavum* Ørsted, 1843

**Notes:** Questionable status. Reported from Greece by Goumenaki (2005). In the Mediterranean also known from Italy (Castelli et al. 2008), Spain (Serrano-Samaniego 2012) and the Adriatic (Mikac 2015). Present in the adjacent Sea of Marmara (Çınar et al. 2014). Otherwise distributed in the North Atlantic. While it is possible that the species occurs in Greece, it has so far only been reported in grey literature; its presence is therefore preliminarily considered questionable, pending confirmation.

## Spintheridae Augener, 1913

### *Spinther arcticus* (M. Sars, 1851)

**Nomenclature:** *Spinther miniaceus* Grube, 1860

**Notes:** Several specimens from Greece in the collections of the Natural History Museum Vienna (8 specimens, Inv. No. 109, Acq. No. 929, Zante [=Zakynthos Island], acquisition date 1874; leg. E. v. Marenzeller; many specimens, Inv. No. 107, Acq. No. 12583, Zante [=Zakynthos Island], acquisition date 1887, det. R. Drasche). Type locality of *S. miniaceus*: Mediterranean (North Adriatic).

## Spionidae Grube, 1850

### *Aonidella cirrobranchiata* (Day, 1961)

**Nomenclature:** *Prionospio cirrobranchiata* Day, 1961

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Satsmadjis 1983). Commonly distributed in West Atlantic and South-West Africa. Arvanitidis (1994) regards its presence in the Mediterranean as doubtful. Day (1961) and Day (1967) include it in the subgroup of *Prionospio* species with apinnate branchiae. Maciolek in her unpublished dissertation (Maciolek Blake 1983) and Lopez-Jamar (1989) moved it to a new genus (*Aonidella*) on the basis of other unique diagnostic characters (for more information see Maciolek 2000). Re-examination of the material would be required to investigate the possible assignment of the Greek record to a *Prionospio* (*Minuspio*) species with apinnate branchiae.

### ***Aonides oxycephala* (Sars, 1862)**

**Notes:** Frequently reported from Greece and other parts of the Mediterranean (Çınar 2005, Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015). Originally described from the North Sea, reported from worldwide locations. However, Radashevsky (2015) questions its cosmopolitanism, noting the possibility of a series of sibling species.

### ***Aonides paucibranchiata* Southern, 1914**

**Notes:** Identifications of Greek specimens are based on descriptions of Fauvel (1927) and may have been easily confused with *Aonides oxycephala* (Sars, 1862) or other similar species. According to a note in Fauvel (1927), this small sized species could be a juvenile form of *Aonides oxycephala*, differing from it mainly by the smaller size, the number of gills and the smaller sized pygidial cirri, while the tridentate ventral hooks could be in fact a post-larval character, frequently observed in the Spionidae. Besides, the number of branchiae and the range of the ventral hooks may follow an intraspecific variation (Ramos 1976a, Johnson 1984 and personal observations from Greek material). According to F. Gravina and A. Somaschini (pers. comm.), it is a rare species.

### ***Atherospio guillei* (Laubier & Ramos, 1974)**

**Nomenclature:** *Polydora guillei* Laubier & Ramos, 1974

**Notes:** Type locality: Mediterranean (Baie de Rosas, Spain).

### ***Aurospio banyulensis* (Laubier, 1966)**

**Nomenclature:** *Prionospio banyulensis* Laubier, 1966

**Notes:** Type locality: Mediterranean (Banyuls-sur-Mer, France).

***Boccardia polybranchia* (Haswell, 1885)**

**Nomenclature:** *Boccardia polybranchia* (Haswell, 1885) | *Polydora* (*Boccardia*) *polybranchia* Haswell, 1885

**Notes:** Originally described from Australia and reported from worldwide locations. However, Radashevsky (2015) notes that the characters of the material from type locality are poorly known and that the insufficient original description could have led to reports from worldwide locations. Therefore several records possibly have been confused or constitute undescribed species.

***Boccardiella cf. ligerica* (Ferrounière, 1898)**

**Notes:** Reported from Greece by Papazacharias (1991). In the Mediterranean also known from the Adriatic (Casellato and Stefanon 2008). Originally described from the East Atlantic.

***Dipolydora armata* (Langerhans, 1880)**

**Nomenclature:** *Dipolydora armata* (Langerhans, 1880) | *Polydora armata* Langerhans, 1881

**Notes:** Originally described from Madeira, with a circumtropical distribution. However, reports of this species show a high polymorphism of specific characters (Radashevsky and Nogueira 2003). It has been suggested that at least two allopatric species are present under the name *Dipolydora armata* (Blake 1983). Bick (2002) distinguishes three morphological forms amongst Western Mediterranean material on the basis of notopodial spines, but Radashevsky and Nogueira (2003) attribute this variability to stages of asexual reproduction of the species.

***Dipolydora blakei* (Maciolek, 1984)**

**Native status:** Cryptogenic

**Notes:** Reported from Greece by Radashevsky and Simboursa (2013) who re-examined specimens reported by Simboursa (1996) and Zenetos et al. (1997) as *Polydora caulleryi* Mesnil, 1897. Originally described from New England, otherwise distributed in the North-West Atlantic and South America.

***Dipolydora coeca* (Ørsted, 1843)**

**Nomenclature:** *Dipolydora coeca* (Ørsted, 1843) | *Polydora caeca* (Ørsted, 1843)



***Dipolydora flava* (Claparède, 1870)**

**Nomenclature:** *Dipolydora flava* (Claparède, 1870) | *Polydora flava* Claparède, 1870

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Dipolydora quadrilobata* (Jacobi, 1883)**

**Nomenclature:** *Polydora quadrilobata* Jacobi, 1883

**Notes:** Reported from Greece by Akoumianaki (2004). In the Mediterranean also known from the Adriatic (Mikac 2015). Reported also from the Black Sea, Romania (Kurt Şahin and Çınar 2012), otherwise distributed in the North Atlantic.

***Dispio magnus* (Day, 1955)**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Simboura (1996). In the Mediterranean also known from the central basin (Zenetos et al. 2012). Originally described from South Africa.

***Laonice cirrata* (M. Sars, 1851)**

**Notes:** Questionable status. Species complex. At least two putative cryptic species exist in the Canadian Pacific (Carr et al. 2011). Records of *Laonice cirrata* in the Mediterranean may actually belong to *Laonice bahusiensis* Söderström 1920 (Mikac 2015, see Sikorski 2003 for differences between species). The former is restricted to circumpolar areas, the very similar *Laonice bahusiensis*, however, has a more southern distribution and is also present in the Central and Eastern Mediterranean (Sikorski 2003, Çınar et al. 2014). Fauvel (1927) synonymised *Laonice bahusiensis* with *Laonice cirrata*, including only *Laonice cirrata* in his key. *Laonice bahusiensis* may therefore be underreported or misidentified and all previous records of *Laonice cirrata* need to be investigated to confirm their identity.

***Laubieriellus salzi* (Laubier, 1970)**

**Nomenclature:** *Prionospio salzi* Laubier, 1970

**Notes:** Reported from Greece by Simboura and Zenetos (2005) and HCMR (2011). Type locality: Mediterranean (Israel).

***Malacoceros fuliginosus* (Claparède, 1870)**

**Nomenclature:** *Malacoceros fuliginosus* (Claparède, 1870) | *Scolelepis fuliginosa* (Claparède, 1870)

**Notes:** Species complex. Guerin and Kerambrun (1984) report three forms, on the basis of morphological, ecological and genetic characters. Two of these forms co-exist in the Mediterranean Sea, while the third was found only in the North East Atlantic. Type locality: Mediterranean Sea (Italy).

### ***Malacoceros tetracerus* (Schmarda, 1861)**

**Nomenclature:** *Malacoceros ciliatus* (Keferstein, 1862) | *Scolelepis ciliata* (Keferstein, 1862)

### ***Malacoceros vulgaris* Johnston, 1827**

**Nomenclature:** *Malacoceros girardi* Quatrefages, 1834 | *Malacoceros vulgaris* Johnston, 1827 | *Scolelepis girardii* (Quatrefages, 1834)

**Notes:** Reported from Greece by Dando et al. (1995) as *Malacoceros vulgaris*, but Arvanitidis (2000a) and Simboura and Nicolaidou (2001) use the name *Malacoceros girardi*. Pettibone (1963) synonymised *Malacoceros girardi* with *Malacoceros vulgaris* and only the latter is accepted as valid by several later authors (e.g. Dauvin et al. 2003, Delgado-Blas and Díaz-Díaz 2010, Mikac 2015). However, the name *Malacoceros girardi* has been recently used for records from the Mediterranean (e.g. Castelli et al. 2008, Dağlı 2008, Çınar et al. 2014). Herein, we follow the suggested synonymy by Pettibone (1963), since no contradicting evidence has been presented in the literature.

### ***Microspio mecznikowianus* (Claparède, 1869)**

**Nomenclature:** *Microspio mecznikowianus* (Claparède, 1869) | *Paraspio mecznikowianus* (Claparède, 1869)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Paraprionospio coora* Wilson, 1990**

**Native status:** Cryptogenic

**Notes:** Reported from Greece by Simboura et al. (2010) who report some slight morphological deviations from the original description and Katsiaras and Simboura (2015). In the Mediterranean also known from Turkey (Yokoyama et al. 2010, Çınar et al. 2014), Spain (Zenetos et al. 2010), Tunisia (Zaâbi et al. 2012) and the Adriatic (Mikac 2015), otherwise known from the Pacific Ocean (Australia, Far East).

### ***Paraprionospio pinnata* (Ehlers, 1901)**

**Nomenclature:** *Paraprionospio pinnata* (Ehlers, 1901) | *Prionospio pinnata* Ehlers, 1901

**Notes:** Questionable status. Several authors suggest that Mediterranean records of this species should be assigned to *Paraprionospio coora* (Yokoyama et al. 2010, Simboursa et al. 2010, Mikac 2015).

### ***Polydora ciliata* (Johnston, 1838)**

**Notes:** Species complex. Mustaquim (1986) and Mustaquim (1988) provided behavioural (boring and tube-dwelling) and genetic evidence of two forms present under *Polydora ciliata*; morphological differences between the forms are also reported. Radashevsky and Pankova (2006), regarding European material, assign the tube-dwelling form to *Polydora ciliata* and the boring form to *Polydora calcarea* (Templeton, 1836). The presence of *Polydora ciliata* in the Mediterranean was questioned by Çınar et al. (2005) and Dağlı et al. (2011); but see also Mikac (2015). The former authors suggest that *Polydora ciliata* could have been confused with *Polydora cornuta* Bosc, 1802 or *Polydora agassizii* Claparède, 1869; a Mediterranean species synonymised with *Polydora ciliata* by Carazzi (1893), but resurrected by Radashevsky and Hsieh (2000). Until re-examination of the Greek material is possible, these records are retained under *Polydora ciliata* in this checklist.

### ***Polydora cornuta* Bosc, 1802**

**Native status:** Non-native (casual)

**Notes:** Species complex. Rice et al. (2008) provide molecular evidence of at least three putative cryptic species present in North America. Reported from Greece by Simboursa et al. (2008). Originally described from South Carolina; in the Mediterranean also known from Spain (Tena et al. 2000), Turkey (Çınar et al. 2014) and from Adriatic lagoons (Bertasi 2016). Also reported from the Black Sea (Kurt Şahin and Çınar 2012, Radashevsky and Selifonova 2013).

### ***Polydora hoplura* Claparède, 1868**

**Notes:** Often found boring in oysters. Type locality: Mediterranean (Gulf of Naples). Sato-Okoshi et al. (2016), in agreement with the taxonomic observations of Radashevsky and Migotto (2016), provided molecular evidence that *Polydora uncinata* Sato-Okoshi, 1998 is a synonym of *Polydora hoplura*.

### ***Polydora spongicola* Berkeley & Berkeley, 1950**

**Native status:** Non-native (questionable)

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Arvanitidis 2000a). Originally described from Western Canada, distributed in the East and West Pacific. The species is associated with sponges, a habitat for polychaetes which is little investigated in Greece.

### ***Prionospio caspersi* Laubier, 1962**

**Nomenclature:** *Prionospio caspersi* Laubier, 1962 | *Prionospio (Prionospio) caspersi* Laubier, 1962

**Notes:** Type locality: Mediterranean (Venice Lagoon). Otherwise distributed in the North Atlantic and Pacific Ocean. Simboura (1996) notes that the Greek material differs slightly from the type material (Adriatic) and from material from Japan, suggesting that the Greek specimens may belong to an undescribed species.

### ***Prionospio cirrifera* Wirén, 1883**

**Nomenclature:** *Minuspio cirrifera* (Wirén, 1883) | *Prionospio (Minuspio) cirrifera* Wirén, 1883 | *Prionospio cirrifera* Wirén, 1883

**Notes:** Questionable status. Reported frequently from Greece and other parts of the Mediterranean (Castelli et al. 2008, Zaâbi et al. 2012, Çınar et al. 2014, Mikac 2015) and considered cosmopolitan. However, Mackie (1984) re-examined European material identified as *Prionospio cirrifera* and found many specimens to actually belong to a species that could not be distinguished from *Prionospio multibranchiata* Berkeley, 1927, based on the available information at that time. Sigvaldadóttir (2002), in agreement with Mackie (1984), suggests that *Prionospio cirrifera* is a North Sea species and probably does not occur south of Portugal. Mackie suggested that Mediterranean specimens probably belong to other, endemic species (Mackie pers. comm. in Simboura 1996) and later, Dağlı and Çınar (2011) described *Prionospio maciolekae* from Mediterranean specimens. Other specimens of *Prionospio cirrifera* from Italy (Giangrande and Gambi 1982) were re-examined by Dağlı and Çınar (2011) and shown to belong to the alien species *Prionospio pulchra* Imajima, 1990. See also remarks under *Prionospio maciolekae*, *Prionospio multibranchiata* and *Prionospio pulchra*.

### ***Prionospio decipiens* Söderström, 1920**

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Nicolaidou 1985). Originally described from Portugal. The material was not available for re-examination, but the species' presence in Greece is considered questionable.

### ***Prionospio dubia* Maciolek, 1985**

**Notes:** Synonymised with *Prionospio steenstrupi* Malmgren, 1867 soon after the original description (Day 1963). Maciolek (1985) re-instated *Prionospio dubia* as a valid species and Sigvaldadóttir and Mackie (1993) assigned the description of *Prionospio steenstrupi* by Fauvel (1927) in part to *Prionospio dubia* (Sigvaldadóttir 2002). *Prionospio steenstrupi* seems to be restricted to boreal areas, whereas material from

Greece was identified to belong to *Prionospio dubia* (Sigvaldadóttir pers. comm. in Simboursa 1996).

### ***Prionospio ehlersi* Fauvel, 1928**

**Nomenclature:** *Prionospio* (*Prionospio*) *ehlersi* Fauvel, 1928 | *Prionospio ehlersi* Fauvel, 1928

### ***Prionospio ergeni* Dağlı & Çınar, 2009**

**Notes:** New record for Greece. One specimen, Antikyra, Gulf of Corinth, 38°23'03.8"N, 22°39'14.5"E, 6.5 m depth, silty sand with biogenic detritus. Literature used for identification: Dağlı and Çınar (2009). Type locality: Mediterranean (Turkey).

### ***Prionospio fallax* Söderström, 1920**

**Notes:** Frequently reported from Greece and other parts of the Mediterranean (Castelli et al. 2008, Ayari et al. 2009, Çınar et al. 2014, Mikac 2015). Simboursa (1996), after examination of material from Greece and in communication with other experts (Sigvaldadóttir, Mackie), concludes that several specimens from Europe erroneously identified as *Prionospio malmgreni* Claparède, 1870 actually belong to *Prionospio fallax*. In the original description, *Prionospio malmgreni* was reported to have up to 9 pairs of branchiae, a character which was never found again in any specimen. The description of *Prionospio malmgreni* by Fauvel (1927) belongs to *Prionospio fallax*, leading to several misidentifications (Sigvaldadóttir and Mackie 1993). *Prionospio malmgreni* has been considered an indeterminate species (Maciolek 1985, Sigvaldadóttir and Mackie 1993) and it is suggested to assign previous Mediterranean records to *Prionospio fallax* (Çınar et al. 2014, Mikac 2015).

### ***Prionospio maciolekae* Dağlı & Çınar, 2011**

**Notes:** New record for Greece. Dağlı and Çınar (2011) described *Prionospio maciolekae* from material previously identified as *Prionospio multibranchiata* Berkeley, 1927 and question the presence of *Prionospio multibranchiata* in the Mediterranean. Re-examination of some of the Greek specimens of *Prionospio multibranchiata* revealed that they actually belong to *Prionospio maciolekae*. Type locality: Mediterranean (Turkey).

### ***Prionospio multibranchiata* Berkeley, 1927**

**Nomenclature:** *Minuspio multibranchiata* (Berkeley, 1927) | *Prionospio multibranchiata* Berkeley, 1927 | *Prionospio* (*Minuspio*) *multibranchiata* Berkeley, 1927

**Notes:** Questionable status. Species complex. Reported frequently from Greece and other parts of the Mediterranean (Çınar 2005, Castelli et al. 2008, Mikac 2015).

However, Mediterranean records probably belong to *Prionospio maciolekae* Dağlı and Çınar 2011 (see remarks there and under *Prionospio cirrifera* Wirén 1883). Otherwise distributed in the West Atlantic and East Pacific. Simboursa (1996) describes two forms of *Prionospio* cf. *multibranchiata*, the one with longer gills which was later re-identified as *Prionospio pulchra* Imajima, 1990.

### ***Prionospio pulchra* Imajima, 1990**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Pancucci-Papadopoulou et al. (2005). Originally from the Pacific Ocean, in the Mediterranean also known from Turkey (Çınar et al. 2014) and introduced to the East Atlantic (Moreira et al. 2000). Dağlı and Çınar (2011) suggest that the specimens of *Prionospio cirrifera* Wirén, 1883 collected from Italy by Giangrande and Gambi (1982) are identical to *Prionospio pulchra*. Mediterranean records of *Prionospio multibranchiata* Berkeley, 1927 and *Prionospio cirrifera* could comprise specimens of *Prionospio pulchra* (see Dağlı and Çınar 2011). See also remarks under *Prionospio cirrifera*.

### ***Prionospio steenstrupi* Malmgren, 1867**

**Nomenclature:** *Prionospio* (*Prionospio*) *steenstrupi* Malmgren, 1867 | *Prionospio steenstrupi* Malmgren, 1867

**Notes:** Questionable status. Often confused with *Prionospio dubia* Maciolek, 1985 and several Greek specimens identified as *Prionospio steenstrupi* in fact belong to the former (see discussion there). In addition, Sigvaldadóttir and Mackie (1993) restrict its distribution to the coast of Iceland and consider all other reports doubtful. However, Dağlı and Çınar (2009) confirm the presence of *Prionospio steenstrupi* in Turkey. Reported from the Atlantic, Pacific, Arctic and Indian Ocean.

### ***Pseudopolydora antennata* (Claparède, 1869)**

**Nomenclature:** *Polydora antennata* Claparède, 1869 | *Polydora* (*Carazzia*) *antennata* Claparède, 1869 | *Pseudopolydora antennata* (Claparède, 1869)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Pseudopolydora paucibranchiata* (Okuda, 1937)**

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Simboursa et al. (2010), found again afterwards (HCMR, unpublished data). Originally described from Japan, in the Mediterranean also known from Turkey (Dağlı and Çınar 2008) and the Adriatic (Mikac 2015).

***Pseudopolydora pulchra* (Carazzi, 1893)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Pygospio elegans* Claparède, 1863**

**Notes:** Species complex. Radashevsky et al. (2016) provide molecular evidence that at least two sibling species are present in the Pacific Ocean. However, high genetic similarity was found between European and West Atlantic specimens. Currently, no evidence exists for the presence of more than one species in the Mediterranean.

***Scolelepis bonnieri* (Mesnil, 1896)**

**Nomenclature:** *Nerine bonnieri* Mesnil, 1896 | *Scolelepis bonnieri* (Mesnil, 1896)

**Notes:** Reported from Greece by Harmelin (1969) and Antoniadou and Chintiroglou (2005). In the Mediterranean also reported from the central basin (Castelli et al. 2008), Cyprus (Çınar 2005) and Turkey (Çınar et al. 2014), otherwise known from the Atlantic coast of Europe.

***Scolelepis cantabra* (Rioja, 1918)**

**Nomenclature:** *Nerinides cantabra* Rioja, 1918 | *Pseudomalacoceros cantabra* (Rioja, 1918) | *Scolelepis cantabra* (Rioja, 1918)

**Notes:** The new species *Scolelepis neglecta* Surugiu, 2016 was described after re-examination of Mediterranean material which was previously identified as *Scolelepis squamata* (O.F. Müller, 1806) and *Scolelepis cantabra* (Surugiu 2016). Reports of *Scolelepis cantabra* from Greece could therefore also comprise specimens of *Scolelepis neglecta*.

***Scolelepis foliosa* (Audouin & Milne Edwards, 1833)**

**Nomenclature:** *Nerine foliosa* (Audouin & Milne Edwards, 1833) | *Scolelepis foliosa* (Audouin & Milne Edwards, 1833)

***Scolelepis gilchristi* (Day, 1961)**

**Nomenclature:** *Nerinides gilchristi* Day, 1961

**Notes:** Questionable status. Reported from Greece by Nicolaidou and Pitta (1986). In the Mediterranean also known from Egypt (Ben-Eliahu 1972a) and Italy (Cantone and Fassari 1982). However, as the species is not included in subsequent reviews / checklists from Italy (Lardicci 1989, Castelli et al. 2008), it was probably considered a misidentification by the later authors. Originally described from South Africa.

***Scolelepis squamata* (O.F. Müller, 1806)**

**Nomenclature:** *Nerine cirratulus* (Delle Chiaje, 1831) | *Scolelepis mesnili* (Bellan & Lagardère, 1971) | *Scolelepis squamata* (Müller, 1806)

**Notes:** Could comprise specimens of *Spiophanes neglecta* Surugiu, 2016 (see notes under *Scolelepis cantabra* (Rioja, 1918).

***Scolelepis tridentata* (Southern, 1914)**

**Nomenclature:** *Nerinides tridentata* Southern, 1914 | *Pseudomalacoceros tridentata* (Southern, 1914) | *Scolelepis tridentata* (Southern, 1914)

***Spio decoratus* Bobretzky, 1870*****Spio filicornis* (Müller, 1776)**

**Notes:** Questionable status. Frequently reported, but its presence in the Mediterranean has been questioned by Simboura (1996). In addition, Meißner et al. (2011) redescribed the species from the type locality (Ilulîarssuk, Greenland) and concluded that traditionally used diagnostic characters are inappropriate. The same authors limit the species' distribution to Greenland and encourage the re-examination of material from other locations.

***Spio multioculata* (Rioja, 1918)*****Spiophanes bombyx* (Claparède, 1870)**

**Notes:** Species complex. At least four morphologically similar species exist (Meißner and Blank 2009). All forms currently known have been formally described as different species in recent years. Specimens from the Turkish Aegean were studied by Meißner and Blank (2009) and found to belong to *Spiophanes bombyx*, thus it can be assumed that at least some of the Greek material belongs to *Spiophanes bombyx* sensu stricto. Type locality: Mediterranean (Gulf of Naples).

***Spiophanes duplex* (Chamberlin, 1919)**

**Nomenclature:** *Spiophanes soderstromi* Hartman, 1953

**Notes:** Questionable status. Reported from Greece by NCMR (1989). Meißner (2005) restricts the distribution of the species to the Atlantic and Pacific coast of America and the Greek record is probably a misidentification.



***Spiophanes kroyeri* Grube, 1860**

**Nomenclature:** *Spiophanes kroyeri* Grube, 1860 | *Spiophanes kroyeri kroyeri* Grube, 1860

**Notes:** Questionable status. Frequently reported from Greece and other parts of the Mediterranean (Castelli et al. 2008, Çınar et al. 2014). However, Meißner (2005) concluded that so far only specimens from the North Atlantic can be assigned to *Spiophanes kroyeri*. Mikac (2015) could only confirm the presence of *Spiophanes afer* Meißner, 2005 from the Adriatic Sea, whereas *Spiophanes kroyeri* could not be found. *Spiophanes afer* is also known from the Turkish Aegean Sea and the Sea of Marmara (Dağlı 2008, Dağlı et al. 2011).

***Spiophanes reyssi* Laubier, 1964**

**Nomenclature:** *Spiophanes kroyeri reyssi* Laubier, 1964

**Notes:** Meißner (2005) raised *Spiophanes reyssi* to species level and described *Spiophanes mediterraneus* Meißner, 2005 from the Eastern Mediterranean, a species morphologically very close to *Spiophanes reyssi*. Thus, older records of *Spiophanes kroyeri reyssi* need to be treated with care as they may belong to either of the two species. *Spiophanes kroyeri reyssi* is reported from Greece and the Mediterranean from bathyal zones (Simboura and Nicolaidou 2001); *Spiophanes mediterraneus* is likewise described from depths of 300–700 m, a habitat which distinguishes these two species from congeneric species such as *Spiophanes kroyeri* or *Spiophanes afer*. Type locality: Mediterranean (off France).

***Streblospio shrubsolii* (Buchanan, 1890)****Sternaspidae Carus, 1863*****Sternaspis scutata* Ranzani, 1817**

**Notes:** Sendall and Salazar-Vallejo (2013) re-instated *Sternaspis thalassemoides* Otto, 1821 from the Mediterranean alongside *Sternaspis scutata*, mainly based on morphological features of the fan (truncate, entire and not expanding beyond the posterolateral margins in *Sternaspis thalassemoides*; notched and expanded beyond the posterolateral margins in *Sternaspis scutata*). Specimens recorded in the past as *Sternaspis scutata* could belong to either of the two species. Type locality: Mediterranean (originally Adriatic Sea, neotype from Izmir, Turkey).

## Syllidae Grube, 1850

### *Amblyosyllis formosa* (Claparède, 1863)

**Nomenclature:** *Amblyosyllis dorsigera* Claparède, 1864 | *Amblyosyllis formosa* (Claparède, 1863) | *Pterosyllis formosa* Claparède, 1863

**Notes:** Three *Amblyosyllis* species have been recorded so far from the Mediterranean (*Amblyosyllis formosa*, *Amblyosyllis dorsigera* (Claparède, 1864) and *Amblyosyllis madeirensis* Langerhans, 1879); *Amblyosyllis dorsigera* is currently considered a synonym of *Amblyosyllis formosa*. However, the characters traditionally used to distinguish these three species (number and type of pharyngeal teeth) seem not to be consistent (Çinar and Ergen 2002) and the genus is in need of revision. At least two species of *Amblyosyllis* occur in Greece, as Arvanitidis (1994) reports both *Amblyosyllis formosa* and *Amblyosyllis madeirensis* in the same study; therefore both records are here retained as valid pending further analyses resolving their identity. Type locality: Mediterranean (Port-Vendres, France).

### *Amblyosyllis madeirensis* Langerhans, 1879

**Notes:** Reported from Greece by Arvanitidis (2000a). In the Mediterranean also known from Italy (Mikac 2015, Castelli et al. 2008) and Spain (San Martín 2003). See also notes for *Amblyosyllis formosa* (Claparède, 1863).

### *Anoplosyllis edentula* Claparède, 1868

**Nomenclature:** *Syllides edentulus* (Claparède, 1868)

**Notes:** Questionable status. Reported from Greece by IOFR (1984). Type locality: Mediterranean (Gulf of Naples). While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.

### *Autolytus rubrovittatus* Claparède, 1864

**Notes:** Questionable status. Reported from Greece by Karakassis (1991). Type locality: Mediterranean (Gulf of Naples). As type material has been lost and the original description is insufficient, the taxon is considered of unclear generic affiliation (Nygren 2004) and all subsequent records must be considered questionable.

***Branchiosyllis cirropunctata* (Michel, 1909)**

**Nomenclature:** *Syllis cirropunctata* Michel, 1909 | *Syllis* (*Typosyllis*) *cirropunctata* Michel, 1909 | *Typosyllis cirropunctata* (Michel, 1909)

**Notes:** Treated as a synonym of *Branchiosyllis exilis* (Gravier, 1900) for many years (e.g. Licher 1999, San Martín 2003) and resurrected by San Martín et al. (2008), thus it may be underreported in Greece. Type locality: Mediterranean (Gulf of Naples).

***Branchiosyllis exilis* (Gravier, 1900)**

**Notes:** Species complex. The complex of pseudo-cryptic species around *Branchiosyllis exilis* is confused and in need of revision and it is likely that specimens around the world recorded under the name *Branchiosyllis exilis* belong to different species (San Martín et al. 2008). Specimens from the Mediterranean recorded as *Branchiosyllis exilis* could also belong to the native Mediterranean species *Branchiosyllis cirropunctata* (Michel, 1909), as the two species differ mainly in the shape of the shaft head of the falcigers and *Branchiosyllis cirropunctata* was considered a synonym of *Branchiosyllis exilis* until recently.

***Brania arminii* (Langerhans, 1881)**

**Nomenclature:** *Brania arminii* (Langerhans, 1881) | *Brania oculata* (Hartmann-Schröder, 1960)

***Brania pusilla* (Dujardin, 1851)**

**Nomenclature:** *Brania pusilla* (Dujardin, 1851) | *Grubea pusilla* (Dujardin, 1851)

***Brevicirrosyllis weismanni* (Langerhans, 1879)**

**Nomenclature:** *Pionosyllis weismanni* Langerhans, 1879

***Epigamia alexandri* (Malmgren, 1867)**

**Nomenclature:** *Autolytus longeferiens* Saint Joseph, 1887

**Notes:** Questionable status. Reported from Greece by Kisseleva (1983). In the Mediterranean also reported from Egypt (Fauvel 1937) and Italy (Castelli et al. 2008) but Nygren (2004) restricts it to the Arctic, the North-West Atlantic and the North Pacific.

***Erinaceusyllis cryptica* (Ben-Eliahu, 1977)**

**Nomenclature:** *Erinaceusyllis cryptica* (Ben-Eliahu, 1977) | *Sphaerosyllis cryptica* Ben-Eliahu, 1977

**Notes:** Reported from Greece by Arvanitidis (1994), also found by Papageorgiou et al. (2006) (unpublished data). Widely distributed in the Mediterranean (Musco and Giangrande 2005).

***Erinaceusyllis erinaceus* (Claparède, 1863)**

**Nomenclature:** *Sphaerosyllis erinaceus* Claparède, 1863

***Eurysyllis tuberculata* Ehlers, 1864**

**Notes:** Type locality: Mediterranean (Kvarner Bay, Adriatic).

***Eusyllis assimilis* Marenzeller, 1875**

**Notes:** Type locality: Mediterranean (Lošinj, Adriatic).

***Eusyllis blomstrandii* Malmgren, 1867*****Eusyllis lamelligera* Marion & Bobretzky, 1875**

**Notes:** Type locality: Mediterranean (Marseille).

***Exogone brevi antennata* Hartmann-Schröder, 1959**

**Nomenclature:** *Exogone ovalis* Hartmann-Schröder, 1960

**Native status:** Non-native (casual)

**Notes:** Reported from Greece by Dando et al. (1995) in an ecological species list as *Exogone ovalis*. In the Mediterranean also known from Lebanon (Aguado and San Martín 2007) and Turkey (Çınar et al. 2014), otherwise it has a circumtropical distribution (e.g. San Martín 1991, Nogueira et al. 2004, Böggemann and Westheide 2004). The Eastern Mediterranean records make an occurrence of *Exogone brevi antennata* in Greece likely, however, as it is very similar to the Mediterranean *Exogone verugera* (Claparède, 1868), (distinguished from the latter mainly by the presence of a dorsal cirrus on the second chaetiger (Aguado and San Martín 2007)), additional specimens are needed to confirm its presence.

***Exogone dispar* (Webster, 1879)**

**Nomenclature:** *Exogone* (*Exogone*) *dispar* (Webster, 1879) | *Exogone dispar* (Webster, 1879)

***Exogone lopezi* San Martín, Ceberio & Aguirrezabalaga, 1996**

**Nomenclature:** *Exogone* (*Exogone*) *lopezi* San Martín, Ceberio & Aguirrezabalaga, 1996

**Notes:** Deep sea species, in the Mediterranean only known from the bathyal zone of the North Aegean (Simboura and Zenetos 2005). In the Mediterranean also reported from deep waters off Sardinia (Busoni 2013), otherwise known from the Atlantic coasts of the Iberian Peninsula.

***Exogone naidina* Ørsted, 1845**

**Nomenclature:** *Exogone* (*Exogone*) *naidina* Ørsted, 1845 | *Exogone gemmifera* Pagenstecher, 1862 | *Exogone naidina* Ørsted, 1845

***Exogone rostrata* Naville, 1933**

**Nomenclature:** *Exogone* (*Exogone*) *rostrata* Naville, 1933 | *Exogone rostrata* Naville, 1933

**Notes:** Type locality: Mediterranean (Banyuls-sur-Mer, France).

***Exogone sorbei* San Martín, Ceberio & Aguirrezabalaga, 1996**

**Nomenclature:** *Exogone sorbei* San Martín, Ceberio & Aguirrezabalaga, 1996 | *Exogone* (*Exogone*) *sorbei* San Martín, Ceberio & Aguirrezabalaga, 1996

**Notes:** Deep sea species, in the Mediterranean only known from the bathyal zone of the North Aegean (Simboura and Zenetos 2005). Otherwise known from the Atlantic coasts of the Iberian Peninsula.

***Exogone torulosa* (Claparède, 1864)**

**Nomenclature:** *Spermosyllis torulosa* Claparède, 1864

**Notes:** Questionable status. A poorly known species which is frequently reported in ecological lists, but no recent descriptions exist (the newest being by Fauvel (1923), who reproduces description and illustrations by Claparède (1864) and Soulier (1904)). The species is not included in the treatise of the family by San Martín (2003). Type locality: Mediterranean (Port-Vendres, France).

***Exogone verugera* (Claparède, 1868)**

**Nomenclature:** *Exogone* (*Exogone*) *verugera* (Claparède, 1868) | *Exogone verugera* (Claparède, 1868)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Haplosyllis spongicola* (Grube, 1855)**

**Nomenclature:** *Haplosyllis hamata* (Claparède, 1868) | *Haplosyllis spongicola* (Grube, 1855) | *Syllis spongicola* Grube, 1855 | *Syllis* (*Haplosyllis*) *spongicola* Grube, 1855

**Notes:** Species complex. The taxonomy of the species is confused, with a long list of synonyms. Originally described from the Northern Adriatic but reported from worldwide locations (Martín et al. 2003), with three different morphotypes in the Western Mediterranean. Currently, three well-described species of *Haplosyllis* are known the Mediterranean: *Haplosyllis spongicola*, *Haplosyllis chamaeleon* Laubier, 1960 and *Haplosyllis granulosa* (Lattig, San Martín & Martín, 2007). The description of *Haplosyllis spongicola* by San Martín (2003) is referred in part to *Haplosyllis granulosa* (Lattig et al. 2007). In addition, a number of *Haplosyllis* species have been described from the Mediterranean and the Black Sea which are currently regarded as *incertae sedis* due to lost type material and insufficient descriptions (Lattig and Martín 2009). Eastern Mediterranean records should also be examined for possible occurrences of the Red Sea species *Haplosyllis djiboutiensis* Gravier, 1900.

***Myrianida brachycephala* (Marenzeller, 1874)**

**Nomenclature:** *Autolytus brachycephalus* (Marenzeller, 1874)

**Notes:** Type locality: Mediterranean (Adriatic).

***Myrianida convoluta* (Cognetti, 1953)**

**Nomenclature:** *Autolytus convolutus* Cognetti, 1953 | *Myrianida convoluta* (Cognetti, 1953)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Myrianida edwarsi* (Saint Joseph, 1887)**

**Nomenclature:** *Autolytus edwarsi* Saint Joseph, 1887 | *Myrianida edwarsi* (Saint Joseph, 1887)

***Myrianida inermis* (Saint Joseph, 1887)**

**Notes:** Questionable status. Reported from Greece by Faulwetter et al. (2011a) based on a single specimen. In the Mediterranean also known from Italy (Somaschini 1988). Nygren (2004) restricts its distribution to the North-East Atlantic and North-East Pacific. Mediterranean specimens could be misidentifications in which the trepan was overlooked. In addition, the Greek specimen was identified using the key and description by San Martín (2003) which, according to Nygren (2004), corresponds only in part to *Myrianida inermis*.

***Myrianida pinnigera* (Montagu, 1808)**

**Notes:** Reported from Greece by Arvanitidis (2000a) based on a single specimen. Widely distributed in the Mediterranean (Nygren 2004).

***Myrianida prolifera* (O.F. Müller, 1788)**

**Nomenclature:** *Autolytus prolifer* (O.F. Müller, 1788) | *Myrianida prolifera* (O.F. Müller, 1788)

**Notes:** The name has been applied to different taxa in the past, including stolons (Gidholm 1967). Nygren (2004) restricts its distribution to the North Atlantic and cautions that preserved specimens are easily confused with other taxa.

***Myrianida quindecimdentata* (Langerhans, 1884)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Keklikoglou et al. (2013). Widely distributed in the Mediterranean (Musco and Giangrande 2005).

***Myrianida rodosensis* Çınar, 2015**

**Notes:** Originally described from Rhodes (36°20.13'N, 28°12.52'E; 1 m depth), no other records from Greece.

***Myrianida rubropunctata* (Grube, 1860)**

**Nomenclature:** *Autolytus rubropunctatus* (Grube, 1860)

**Notes:** Type locality: Mediterranean (Kraljevica, Croatia).

***Neopetitia amphophthalma* (Siewing, 1956)**

**Nomenclature:** *Petitia amphophthalma* Siewing, 1956

**Notes:** Species complex. Reported from Greece by von Soosten et al. (1998) in the framework of a molecular analysis. Greek specimens differ genetically from those from the type locality (Arcachon Bay), but the authors do not consider the differences significant enough to separate different species (von Soosten et al. 1998, Westheide and Hass-Cordes 2001).

***Nudisyllis divaricata* (Keferstein, 1862)**

**Nomenclature:** *Nudisyllis divaricata* (Keferstein, 1862) | *Pionosyllis divaricata* (Keferstein, 1862)

***Nudisyllis pulligera* (Krohn, 1852)**

**Nomenclature:** *Pionosyllis pulligera* (Krohn, 1852) | *Pionosyllis serrata* Southern, 1914

**Notes:** Type locality: Mediterranean.

***Odontosyllis ctenostoma* Claparède, 1868**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Odontosyllis dugesiana* Claparède, 1864**

**Notes:** Reported from Greece by Fassari (1982). Type locality: Mediterranean (Port-Vendres, France).

***Odontosyllis fulgurans* (Audouin & Milne Edwards, 1833)**

**Notes:** Type locality: Mediterranean.

***Odontosyllis gibba* Claparède, 1863**

***Opisthodonta longocirrata* (Saint-Joseph, 1887)**

**Nomenclature:** *Pionosyllis longocirrata* Saint Joseph, 1887

**Notes:** Questionable status. Reported from Greece by Xenopoulou (1987). Widely distributed throughout the Mediterranean (Musco and Giangrande 2005). While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.



***Opisthodonta morena* Langerhans, 1879**

**Notes:** New record for Greece. Six specimens, Othoni, off Corfu, 39°47'50.3"N, 19°28'13.6"E, 29 m depth, coarse biogenic sand and *Posidonia oceanica* fragments, collected in the framework of the COCONET project, Literature used for identification: San Martín (2003). Widely distributed throughout the Mediterranean (Musco and Giangrande 2005).

***Opisthosyllis brunnea* Langerhans, 1879**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Keklikoglou et al. (2013). Widely distributed in the Mediterranean (Musco and Giangrande 2005). Considered cosmopolitan, but Paresque et al. (2016) found morphological differences between specimens from different localities and raise the possibility of *Opisthosyllis brunnea* constituting a species complex.

***Paraehlersia dionisi* (Núñez & San Martín, 1991)**

**Nomenclature:** *Pionosyllis dionisi* Núñez & San Martín, 1991

**Notes:** Reported from Greece by NCMR (1995) and Simboura (1996), identification confirmed by G. San Martín (pers. comm. in Simboura 1996). In the Mediterranean also known from the Western Mediterranean (Musco and Giangrande 2005), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise known from the Canary Islands and Antarctica (San Martín et al. 2009).

***Paraehlersia ferrugina* (Langerhans, 1881)**

**Nomenclature:** *Ehlersia ferrugina* Langerhans, 1881 | *Langerhansia ferrugina* (Langerhans, 1881) | *Paraehlersia ferrugina* (Langerhans, 1881) | *Syllis ferrugina* (Langerhans, 1881)

***Parapionosyllis brevicirra* Day, 1954*****Parapionosyllis cabezali* Parapar, San Martín & Moreira, 2000*****Parapionosyllis elegans* (Pierantoni, 1903)**

**Notes:** Questionable status. Reported from Greece by IOFR (1984). Type locality: Mediterranean (Gulf of Naples). While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in over three decades, therefore its presence in Greece is here considered questionable.

***Parapionosyllis minuta* (Pierantoni, 1903)**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Parexogone campoyi* (San Martín, Ceberio & Aguirrezabalaga, 1996)**

**Nomenclature:** *Exogone* ( *Parexogone* ) *campoyi* San Martín, Ceberio & Aguirrezabalaga, 1996

**Notes:** Deep sea species, reported from Greece by Simboura and Zenetos (2005) from the bathyal zone of the North Aegean. In the Mediterranean also reported from deep waters in the Ionian Sea (Langeneck et al. 2017b), otherwise known from the Atlantic coasts of the Iberian Peninsula.

***Parexogone gambiae* Lanera, Sordino & San Martín, 1994**

**Nomenclature:** *Exogone gambiae* Lanera, Sordino & San Martín, 1994 | *Exogone* ( *Parexogone* ) *gambiae* Lanera, Sordino & San Martín, 1994

**Notes:** Type locality: Mediterranean (Ponza, Tyrrhenian Sea).

***Parexogone hebes* (Webster & Benedict, 1884)**

**Nomenclature:** *Exogone hebes* (Webster & Benedict, 1884) | *Exogone* ( *Parexogone* ) *hebes* (Webster & Benedict, 1884) | *Parexogone hebes* (Webster & Benedict, 1884)

***Parexogone meridionalis* Cognetti, 1955**

**Nomenclature:** *Exogone parahomoseta mediterranea* San Martín, 1984

**Notes:** New record for Greece. Three specimens, Elaionisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Cognetti (1955), San Martín (2003). Type locality: Mediterranean (Gulf of Naples).

***Parexogone wolffi* San Martín, 1991**

**Nomenclature:** *Exogone* ( *Parexogone* ) *wolffi* San Martín, 1991

**Notes:** Reported from Greece from the bathyal zone of the North Aegean (Simboura and Zenetos 2005). In the Mediterranean also known from Cyprus (Çınar et al. 2003), otherwise distributed on both sides of the Atlantic (San Martín 2003).

***Perkinsyllis anophthalma* (Capaccioni & San Martín, 1990)**

**Nomenclature:** *Pionosyllis anophthalma* Capaccioni & San Martín, 1990

**Notes:** Reported from Greece by Simboura et al. (2007). Type locality: Mediterranean (Catalonia, Spain).

***Plakosyllis brevipes* Hartmann-Schröder, 1956**

**Nomenclature:** *Plakosyllis brevipes* Hartmann-Schröder, 1956 | *Plakosyllis quadrioculata* Perkins, 1981

**Notes:** Type locality: Mediterranean.

***Proceraea aurantiaca* Claparède, 1868**

**Nomenclature:** *Autolytus aurantiacus* (Claparède, 1868) | *Proceraea aurantiaca* Claparède, 1868

**Notes:** Has often been confused with similar species, especially with *Proceraea paraurantiaca* Nygren, 2004 and older records should be treated with caution (Nygren 2004). Type locality: Mediterranean (Gulf of Naples).

***Proceraea picta* Ehlers, 1864**

**Nomenclature:** *Autolytus pictus* (Ehlers, 1864) | *Proceraea picta* Ehlers, 1864

**Notes:** As preserved specimens of *Proceraea picta* are often impossible to distinguish from other species, literature records need to be treated with care (Nygren 2004). Type locality: Mediterranean (Croatia).

***Procerastea nematodes* Langerhans, 1884**

**Nomenclature:** *Procerastea nematodes* Langerhans, 1884 | *Procerastea perrieri* Gravier, 1900

**Notes:** Reported from Greece as *Procerastea perrieri* by Nicolaidou et al. (1986) and Simboura et al. (1995b), subsequently listed as *Procerastea nematodes* by Arvanitidis (2000a) and Simboura and Nicolaidou (2001). In the Mediterranean also reported from Italy (Castelli et al. 2008), otherwise distributed along the Atlantic coasts of Europe. Nygren (2004) restricts the species' distribution to the North-East Atlantic and the North-East Pacific.

### ***Prosphaerosyllis brevicirra* (Hartmann-Schröder, 1960)**

**Nomenclature:** *Sphaerosyllis brevicirra* Hartmann-Schröder, 1960

**Notes:** Questionable status. Records of *Sphaerosyllis brevicirra* from the Western Mediterranean by Alós (1989) and from the Aegean Sea (Simboura 1996, Çınar 1999) belong to an undescribed *Prosphaerosyllis* species (San Martín 2003). Faulwetter et al. (2011a) examined the holotype of *Prosphaerosyllis brevicirra* from the Red Sea and found that the species bears a conspicuous papilla on each dorsal cirrus. All descriptions and illustrations of *Prosphaerosyllis brevicirra* from the Mediterranean lack these papillae. In addition, these specimen descriptions differ from *Prosphaerosyllis brevicirra* by the absence of dorsal cirri on chaetiger 2 (reported as present by Alós (1989) but in fact absent (San Martín 2003) and by thicker aciculae.

### ***Prosphaerosyllis campoyi* (San Martín, Acero, Contonente & Gomez, 1982)**

**Nomenclature:** *Prosphaerosyllis campoyi* (San Martín, Acero, Contonente & Gomez, 1982) | *Sphaerosyllis campoyi* San Martín, Acero, Contonente & Gomez, 1982

**Notes:** Reported from Greece by Faulwetter et al. (2011a), also found by Papageorgiou et al. (2006) (unpublished data). Type locality: Mediterranean (Andalusia, Spain).

### ***Prosphaerosyllis tetralix* (Eliason, 1920)**

**Nomenclature:** *Sphaerosyllis tetralix* Eliason, 1920

**Notes:** Questionable status. Reported from Greece by Dando et al. (1995); the record by Zenetos et al. (1997) probably belongs to a different species. Records from throughout the Mediterranean exist but at least some of them belong to other species (Faulwetter et al. 2011a) and the species is probably restricted to the colder waters of the North Sea and North-East Atlantic (Olivier et al. 2012). During the last years, several new *Prosphaerosyllis* species have been described from Europe and the Mediterranean and Mediterranean species of the genus need to be revised. The Greek record is from near hydrothermal vents where temperatures are probably unfavourable for the species and must be considered questionable.

### ***Prosphaerosyllis xarifae* (Hartmann-Schröder, 1960)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Widely distributed in the Mediterranean (Musco and Giangrande 2005).

### ***Pseudosyllis brevipennis* Grube, 1863**

**Nomenclature:** *Pseudosyllis brevipennis* Grube, 1863 | *Syllis brevipennis* (Grube, 1863) | *Trypanosyllis coeliaca* Claparède, 1868 | *Typosyllis brevipennis* (Grube, 1863)

**Notes:** *Pseudosyllis brevipennis* was considered a synonym of *Trypanosyllis coeliaca* Claparède, 1868 by San Martín (2003), although *Pseudosyllis brevipennis*, being the older name, should have priority over *Trypanosyllis coeliaca*. San Martín (2003), however, considered the former a nomen dubium and chose the latter as the preferred name (San Martín 2003, Álvarez-Campos et al. 2017). A molecular analysis using multiple loci (Álvarez-Campos et al. 2017) reveals that *Trypanosyllis coeliaca* differs distinctly from other *Trypanosyllis* species and is much closer related to *Xenosyllis*, *Eurysyllis* and *Plakosyllis*. These authors re-instate *Pseudosyllis brevipennis*, designate a neotype from Rovigno (Croatia) and place *Trypanosyllis coeliaca* into synonymy with *Pseudosyllis brevipennis*.

### ***Salvatoria alvaradoi* (San Martín, 1984)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Type locality: Mediterranean (Balearic Islands).

### ***Salvatoria clavata* (Claparède, 1863)**

**Nomenclature:** *Brania clavata* (Claparède, 1863) | *Grubea clavata* (Claparède, 1863) | *Grubeosyllis clavata* (Claparède, 1863) | *Salvatoria clavata* (Claparède, 1863)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Salvatoria euritmica* (Sardá, 1984)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Widely distributed in the Mediterranean Sea (Musco and Giangrande 2005), originally described from Gibraltar.

### ***Salvatoria limbata* (Claparède, 1868)**

**Nomenclature:** *Grubeosyllis limbata* (Claparède, 1868) | *Pseudobrania limbata* (Claparède, 1868) | *Salvatoria limbata* (Claparède, 1868)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Salvatoria neapolitana* (Goodrich, 1930)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Keklikoglou et al. (2013). Type locality: Mediterranean (Gulf of Naples).

### ***Salvatoria tenuicirrata* (Claparède, 1864)**

**Nomenclature:** *Brania tenuicirrata* (Claparède, 1864)

**Notes:** Questionable status. A poorly known species which is frequently reported in ecological lists, but no recent descriptions exist (the newest being Fauvel (1923), who reproduces description and illustrations by Claparède (1864)). Fauvel considers the species very close to *Salvatoria clavata* (Claparède, 1863) from which it differs in " *minor* and variable characters". The species is not included in the treatise of the family by San Martín (2003). Type locality: Mediterranean (Port-Vendres, France).

***Salvatoria vieitezi* (San Martín, 1984)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Type locality: Mediterranean (Balearic Islands).

***Salvatoria yraidae* (San Martín, 1984)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Type locality: Mediterranean (Balearic Islands).

***Sphaerosyllis austriaca* Banse, 1959**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Type locality: Western Mediterranean.

***Sphaerosyllis boeroi* Musco, Çınar & Giangrande, 2005**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) based on specimens previously identified as *Sphaerosyllis hystrix*. Type locality: Mediterranean (Porto Cesareo, Ionian Sea).

***Sphaerosyllis bulbosa* Southern, 1914**

***Sphaerosyllis capensis* Day, 1953**

**Notes:** Questionable status. Reported from Greece by Arvanitidis (2000a), in the Mediterranean also reported from Spain (San Martín et al. 1981b), otherwise distributed along the coast of South Africa and in the Indian Ocean. The specimens may have been confused with other Mediterranean species. *Sphaerosyllis capensis* is distinguished from many congeners by having all antennae in line. However, the position of the antennae in preserved material may be misleading if fixation has caused muscle contraction (Riser 1991).

***Sphaerosyllis claparedei* Ehlers, 1864**

**Notes:** Questionable status. *Sphaerosyllis claparedei* is a poorly known species. Ehler's original description is detailed overall, but not clear on all characters that are

used nowadays to distinguish species (e.g. papillation pattern, shape of chaetae, parapodial glands etc.). The species is not included in the treatise of the family by San Martín (2003). A character which allegedly distinguishes *Sphaerosyllis claparedei* from other species of the genus is the presence of a dorsal cirrus on the second chaetiger. However, other species were also originally described with this feature (e.g. *Sphaerosyllis hystrix* Claparède, 1863) although it is in fact absent (see also remarks under *Sphaerosyllis ovigera* Langerhans, 1879). Ehlers (1864) also describes the dorsal cirri as having a “knob-like shape” at their tip, at least in the anterior chaetigers. This could be an indication for retractile cirri, as seen in *Prosphaerosyllis*. As no recent re-descriptions exist which take into account characters used nowadays to distinguish species, the status of the species remains unclear and specimens should be revised. Type locality: Mediterranean (Porto di Lazaretto, Adriatic).

### ***Sphaerosyllis glandulata* Perkins, 1981**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). In the Mediterranean also known from Spain (Somaschini and San Martín 1994), Italy (Castelli et al. 2008) and Turkey (Çınar et al. 2014), otherwise distributed on both sides of the Atlantic (San Martín 2003).

### ***Sphaerosyllis gravinae* Somaschini & San Martín, 1994**

**Notes:** New record for Greece. Two specimens, Elafonisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Somaschini and San Martín (1994). Type locality: Mediterranean (Tyrrhenian Sea).

### ***Sphaerosyllis hystrix* Claparède, 1863**

**Notes:** The name *Sphaerosyllis hystrix* has been applied to a number of specimens worldwide which likely belong to other species. Several specimens from Greece previously identified as *Sphaerosyllis hystrix* were assigned to *Sphaerosyllis boeroi* Musco, Çınar & Giangrande, 2005 after re-examination (Faulwetter et al. 2011b). It is likely that other specimens recorded under the name *Sphaerosyllis hystrix* from Greece actually belong to different species. Type locality: Mediterranean (Gulf of Naples).

### ***Sphaerosyllis magnidentata* Perkins, 1981**

**Notes:** Questionable status. Reported from Greece by Koulouri et al. (2015), in the Mediterranean also known from Spain (Tena et al. 2000), otherwise distributed in the Caribbean. The Mediterranean occurrence record leads to the inclusion of the species in the key of Mediterranean *Sphaerosyllis* by Faulwetter et al. (2011b), but the rarity of records indicates that it may in fact not be present in the region. A re-examination of the specimens is required to verify its presence in Greece and the Mediterranean.

***Sphaerosyllis ovigera* Langerhans, 1879**

**Notes:** Questionable status. A poorly known species which is frequently reported in ecological lists, but no recent descriptions exist (the newest being Fauvel (1923), who reproduces description and illustrations by Langerhans (1879)). The species is not included in the treatise of the family by San Martín (2003). A character which allegedly distinguishes *Sphaerosyllis ovigera* from other species of the genus is the presence of a dorsal cirrus on the second chaetiger. However, other species were originally also described with this feature (e.g. *Sphaerosyllis hystrix*) although it is in fact absent. Non-type material of Langerhans in the Natural History Museum of Vienna (Inv. No. 2549, Acq. No. 3296, locality Madeira, coll. year 1884) was examined by S. Faulwetter; the dorsal cirrus on chaetiger 2 is absent. The status of the species remains unclear and specimens should be revised.

***Sphaerosyllis pirifera* Claparède, 1868**

**Notes:** Type locality: Mediterranean (Gulf of Naples).

***Sphaerosyllis riseri* Perkins, 1981**

**Notes:** New record for Greece. Two specimens, Elafonisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: Perkins (1981). No other Mediterranean records, otherwise distributed along the Atlantic coast of North America.

***Sphaerosyllis taylori* Perkins, 1981*****Sphaerosyllis thomasi* San Martín, 1984**

**Notes:** Reported from Greece by Arvanitidis (2000a). Type locality: Mediterranean (Balearic Islands).

***Streptodonta pterochaeta* (Southern, 1914)**

**Nomenclature:** *Opisthodonta pterochaeta* Southern, 1914

**Notes:** Reported from Greece by Simboura (1996) and Zenetos et al. (1997) (based on the same specimens), identification confirmed by G. San Martín (pers. comm. in Simboura 1996). In the Mediterranean also reported from Spain (Parapar et al. 1993b) and Egypt (Abd-Elnaby 2010), otherwise distributed along the Atlantic coasts of Europe.



***Streptosyllis bidentata* Southern, 1914**

**Notes:** Questionable status. Reported from Greece by Xenopoulou (1987), in the Mediterranean also known from Spain (San Martín 2003) and Tunisia (Zaâbi et al. 2012). Present in the adjacent Sea of Marmara (Çınar et al. 2014). Otherwise distributed in the North East Atlantic.

***Streptosyllis nunezi* Faulwetter, Vasileiadou, Papageorgiou & Arvanitidis, 2008**

**Notes:** Originally described from Crete (Pachia Ammos; 35°06'43"N 25°48'34"E; 1–5 m depth), no other records from Greece.

***Streptosyllis websteri* Southern, 1914*****Syllides bansei* Perkins, 1981**

**Notes:** Reported from Greece by Simboura (1996), identification confirmed by G. San Martín (pers. comm. in Simboura 1996). Also found by Papageorgiou et al. (2006) (unpublished data). In the Mediterranean also reported from Spain (Alós 1989), Italy (Castelli et al. 2008) and Turkey (Çınar et al. 2014), otherwise distributed on both sides of the Atlantic (San Martín 2003).

***Syllides benedicti* Banse, 1971**

**Notes:** In the Mediterranean only reported from Greece (Dando et al. 1995; also found by M. Simboura in the North Aegean (HCMR, unpublished data)). Commonly distributed in the North Atlantic.

***Syllides convolutus* Webster & Benedict, 1884**

**Notes:** New record for Greece. One specimen, Elafonisi, Crete, 35°16'20.7"N, 23°32'15.9"E, 1 m depth, fine sand, collected in the framework of the MEDCORE project (unpublished data from Papageorgiou et al. 2006). Literature used for identification: San Martín (1984), San Martín (2003). In the Mediterranean also known from the western Basin (Musco and Giangrande 2005), otherwise distributed along the Atlantic coast of North America.

***Syllides edentatus* Westheide, 1974**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). In the Mediterranean also known from Spain (San Martín 2003), Italy (Cosentino 2011) and Turkey (Çınar et al. 2014). Cited from worldwide locations.

***Syllides fulvus* (Marion & Bobretzky, 1875)**

**Notes:** Type locality: Mediterranean (Adriatic).

***Syllides japonicus* Imajima, 1966**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). In the Mediterranean also reported from Spain (San Martín 2003), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015). Cited from disjunct worldwide locations (San Martín 2003).

***Syllides longocirratus* (Ørsted, 1845)**

**Notes:** Questionable status. The species described as *Syllides longocirrata* by Fauvel (1923) actually belongs to *Syllides fulvus* (Marion & Bobretzky, 1875) (Parapar et al. 1992, Çınar and Ergen 2003), thus any specimens identified using Fauvel's key probably belong to *Syllides fulvus*. Campoy (1982) considers Mediterranean specimens from soft substrates to belong to *Syllides fulvus* and specimens from hard substrates to *Syllides edentatus*. Records from Greece are all from soft substrates, thus probably belong to *Syllides fulvus*.

***Syllis alternata* Moore, 1908**

**Notes:** Species complex. Consists of at least five putative cryptic species in the North American Pacific coasts, near the type locality (Carr et al. 2011). Specimens from Europe could belong to either of these cryptic species or to a different one. Reported from Greece by Faulwetter et al. (2011a) and Maidanou et al. (2017), widely distributed in the Mediterranean (Musco and Giangrande 2005).

***Syllis amica* Quatrefages, 1866*****Syllis armillaris* (O.F. Müller, 1776)**

**Nomenclature:** *Syllis armillaris* (O.F. Müller, 1776) | *Syllis* (*Typosyllis*) *armillaris* (O.F. Müller, 1776) | *Typosyllis armillaris* (O.F. Müller, 1776)

**Notes:** Considered cosmopolitan, but populations from different geographic locations show morphological differences and the species could actually constitute a complex of pseudo-cryptic species (López et al. 2001).

***Syllis beneliahuae* (Campoy & Alquézar, 1982)**

**Notes:** Type locality: Mediterranean (Spain).

***Syllis columbretensis* (Campoy, 1982)**

**Notes:** Licher (1999) considers *Syllis columbretensis* a synonym of *Syllis variegata* Grube, 1860. San Martín (2003) lists it as a valid species. Specimens from Greece reported by Faulwetter et al. (2011a) and Keklikoglou et al. (2013) were identified using the description and key by San Martín (2003), but those identified by Arvanitidis (2000a) could belong to either *Syllis corallicola* Verrill, 1900 or *Syllis columbretensis*, as he used both the description by Campoy (1982) and the key by San Martín (1984) to identify the specimens. *Syllis columbretensis* sensu San Martín was, however re-identified as *Syllis corallicola* (Licher 1999, San Martín 2003). Type locality: Mediterranean (Columbretes Islands, Spain).

***Syllis compacta* Gravier, 1900**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Keklikoglou et al. (2013). Widely distributed in the Mediterranean (Musco and Giangrande 2005). Licher (1999) considers *Syllis compacta* a synonym of *Syllis variegata* Grube, 1860, San Martín (2003) lists it as a valid species. Specimens from Greece were identified using the description and key by San Martín (2003).

***Syllis corallicola* Verrill, 1900**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Keklikoglou et al. (2013). Widely distributed in the Mediterranean (Musco and Giangrande 2005).

***Syllis cornuta* Rathke, 1843**

**Nomenclature:** *Ehlersia cornuta* (Rathke, 1843) | *Langerhansia cornuta* (Rathke, 1843) | *Syllis cornuta* Rathke, 1843

**Notes:** Questionable status. San Martín and Lopez (2000) suggest that all European records of *Syllis cornuta* should be assigned to *Syllis parapari* San Martín & Lopez, 2000. Mikac and Musco (2010) consider it possible that they could belong to any of the Mediterranean *Syllis* with pseudospinigerous chaetae (*Syllis beneliahuae* (Campoy & Alquézar, 1982), *Syllis rosea* Langerhans, 1879, *Syllis garciai* (Campoy, 1982) or *Syllis parapari*). Licher (1999) examined specimens identified as *Syllis cornuta* from Greece and found that none of these specimens actually belonged to *Syllis cornuta*. However, all Greek specimens would need to be re-examined to determine their true identity.

***Syllis cruzi* Núñez & San Martín, 1991**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). In the Mediterranean also known from Spain (San Martín 2003), Cyprus (Çınar and Ergen 2003), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015); otherwise known from the Canary Islands.

### ***Syllis ferrani* Alós & San Martín, 1987**

**Notes:** Rarely reported from Greece, identifications of specimens by Simboura (1996) were confirmed by G. San Martín (pers. comm.). Type locality: Mediterranean (Cap Creus, Spain).

### ***Syllis garciai* (Campoy, 1982)**

**Notes:** Type locality: Mediterranean (Balearic Islands).

### ***Syllis gerlachi* (Hartmann-Schröder, 1960)**

**Nomenclature:** *Syllis gerlachi* (Hartmann-Schröder, 1960) | *Syllis truncata cryptica* Ben-Eliahu, 1977 | *Syllis* (*Typosyllis*) *truncata cryptica* Ben-Eliahu, 1977

**Notes:** Licher (1999) synonymised *Syllis truncata cryptica* with *Syllis gerlachi*. San Martín et al. (2017) examined the holotype of *Syllis gerlachi* and found it to differ from Mediterranean specimens of *Syllis truncata cryptica* in the shape of the posterior aciculae and other chaetal features (no exact details given). They consider the synonymy incorrect, but as they do not formally re-instate *Syllis truncata cryptica*, we here report records of both species under the name *Syllis gerlachi* until further evidence is provided.

### ***Syllis gerundensis* (Alós & Campoy, 1981)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Maidanou et al. (2017). Type locality: Mediterranean (Gerona, Spain).

### ***Syllis gracilis* Grube, 1840**

**Notes:** Species complex. Originally described from the Gulf of Naples. Maltagliati et al. (2000) found two cryptic species of *Syllis gracilis* in Sardinia and western Italy, Álvarez-Campos et al. (2016) recover three cryptic species in the Mediterranean: one from Sicily and two from the North-Western Mediterranean. No material from the Eastern Mediterranean was studied, and specimens from Greece could belong to any of these or a different species.

### ***Syllis hyalina* Grube, 1863**

**Nomenclature:** *Syllis hyalina* Grube, 1863 | *Typosyllis hyalina* (Grube, 1863) | *Typosyllis* (*Syllis*) *hyalina* (Grube, 1863)

**Notes:** Type locality: Mediterranean (Croatia).

***Syllis jorgei* San Martín & López, 2000**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) under this name, previously reported as *Syllis lutea* (Hartmann-Schröder, 1960) by Arvanitidis (1994) (see also Table 1). Type locality: Mediterranean (Columbretes Islands, Spain).

***Syllis katzmanni* Arvanitidis, 2017, nomen novum**

**Nomenclature:** *Langerhansia caeca* Katzmann, 1973 | *Syllis caeca* (Katzmann, 1973) | *Syllis katzmanni* (Katzmann, 1973) [as a temporary name]

**Notes:** Reported from Greece by Arvanitidis (1994). The species was originally described as *Langerhansia caeca* by Katzmann (1973) from the Adriatic. San Martín (1984) synonymised *Langerhansia* with *Syllis*, thus *Langerhansia caeca* Katzmann, 1973 became *Syllis caeca* (Katzmann, 1973), but the name was preoccupied by *Syllis caeca* Monro 1933. Arvanitidis (1994) proposed the replacement name *Syllis katzmanni* in his PhD thesis, which was however never validly published under the International Code of Zoological Nomenclature (ICZN). Licher (1999) considered *Syllis* and *Typosyllis* two different genera and placed *Syllis caeca* (Katzmann, 1973) back into *Typosyllis*, removing the need for a replacement name and thus synonymising the temporary name *Syllis katzmanni* (Katzmann, 1973) with *Typosyllis caeca* (Katzmann, 1973). He also considered *Syllis caeca* Monro, 1933 a nomen dubium (which does not affect the availability of the name). San Martín (2003) re-synonymised *Typosyllis* with *Syllis*, so again a replacement name is needed to avoid synonymy with *Syllis caeca* Monro, 1933. *Syllis katzmanni* Arvanitidis, 2017 nom. nov. is herewith proposed as a replacement name for *Syllis caeca* (Katzmann, 1973), a previously unreplaced secondary junior homonym of *Syllis caeca* Monro, 1933. As no synonyms are available to choose a replacement name, the new epithet *katzmanni* was chosen in memory of Werner Katzmann (1943–2004), the author of the original name. Records from the Mediterranean given in the literature without an authority likely all belong to Katzmann's species as the type locality of Monro's species is in the Galapagos islands.

***Syllis krohnii* Ehlers, 1864**

**Nomenclature:** *Syllis krohnii* Ehlers, 1864 | *Typosyllis krohnii* (Ehlers, 1864)

**Notes:** Type locality: Mediterranean (Kvarner Gulf, Croatia).

***Syllis* cf. *mayeri* Musco & Giangrande, 2005**

**Notes:** Reported from Greece by Keklikoglou et al. (2013). *Syllis mayeri* is a Caribbean species which was first reported from the Mediterranean (Lebanon) by Aguado and San Martín (2007) as *Syllis* cf. *mayeri*. The authors compared their material to the type material and found slight differences in the length of the anterior dorsal cirri and the pharynx. They note that *Syllis mayeri* and other "cosmopolitan" *Syllis* species possibly

constitute species complexes. The Greek material corresponds exactly to the description of Aguado and San Martín (2007) and may belong to an undescribed species.

### ***Syllis nigricirris* Grube, 1863**

**Nomenclature:** *Syllis nigricirris* Grube, 1863 | *Syllis* (*Typosyllis*) *nigricirris* Grube, 1863

**Notes:** Licher (1999) considers the species as valid, but notes that the characteristic black colour of the dorsal cirri may disappear in ethanol-preserved specimens which would make an unambiguous identification difficult. Greek records are all from before 1987 and were probably identified using Fauvel (1923); the increased use of the keys by San Martín (1984) and San Martín (2003) which do not contain the species could explain the absence of records since then. Type locality: Mediterranean (Adriatic).

### ***Syllis parapari* San Martín & López, 2000**

**Notes:** Probably recorded under the name *Syllis cornuta* before the year 2000 (see discussion under *Syllis cornuta*). In the Mediterranean also known from Cyprus, (Çınar and Ergen 2003), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise known from the Atlantic coast of the Iberian Peninsula and from the British Isles (San Martín 2003, San Martín and Worsfold 2015).

### ***Syllis prolifera* Krohn, 1852**

**Nomenclature:** *Syllis prolifera* Krohn, 1852 | *Typosyllis prolifera* (Krohn, 1852)

**Notes:** Type locality: Mediterranean.

### ***Syllis pulvinata* (Langerhans, 1881)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a). Widely distributed in the Mediterranean (Musco and Giangrande 2005).

### ***Syllis rosea* (Langerhans, 1879)**

#### ***Syllis rosea* cf. *magna* (Westheide 1974) sensu Ben-Eliahu 1977**

**Notes:** Reported from Greece by Simboura (1996) who considers the species closer to *Syllis rosea rosea* than to the Pacific form of *Syllis rosea magna* described by Westheide (1974) from the Galapagos Islands. It is likely that the specimens from the Mediterranean belong to neither of the two species but to an undescribed species (Simboura 1996).

***Syllis torquata* Marion & Bobretzky, 1875**

**Notes:** Reported from Greece by Simboura (1996) and Zenetos et al. (1997) (based on the same specimens). Identification confirmed by G. San Martín (pers. comm. in Simboura 1996). Type locality: Mediterranean (Gulf or Marseille, France).

***Syllis tyrrhena* (Licher & Kuper, 1998)**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) based on a single specimen. Type locality: Mediterranean (Tyrrhenian Sea).

***Syllis variegata* Grube, 1860**

**Nomenclature:** *Syllis variegata* Grube, 1860 | *Syllis* (*Typosyllis*) *variegata* Grube, 1860 | *Typosyllis variegata* (Grube, 1860)

**Notes:** Type locality: Mediterranean (Adriatic).

***Syllis vittata* Grube, 1840**

**Notes:** Type locality: Mediterranean (Palermo, Sicily, Italy).

***Syllis westheidei* San Martín, 1984**

**Notes:** Reported from Greece by Faulwetter et al. (2011a) and Keklikoglou et al. (2013). Type locality: Mediterranean (Balearic Islands).

***Synmerosyllis lamelligera* (Saint-Joseph, 1886)**

**Nomenclature:** *Pionosyllis lamelligera* Saint Joseph, 1887 | *Synmerosyllis lamelligera* (Saint-Joseph, 1886)

***Trypanosyllis aeolis* Langerhans, 1879**

**Notes:** Reported from Greece by Chatzigeorgiou et al. (2016) under this name, previously reported as *Trypanosyllis gemmipara* Johnson, 1901 by Arvanitidis (2000a), based on works by San Martín (1984) and Campoy (1982). San Martín (2003) synonymises *Trypanosyllis gemmipara* with *Trypanosyllis aeolis* and also refers the two aforementioned descriptions to *Trypanosyllis aeolis*. Álvarez-Campos et al. (2017) reinstate *Trypanosyllis gemmipara* based on molecular analyses and assign them to the genus *Trypanedenta* Imajima & Hartman 1964. *Trypanedenta gemmipara* is probably restricted to the Indo-Pacific region (Alaska to Mexico, Japan, New Zealand, India) and reports from the Mediterranean belong to *Trypanosyllis aeolis* (Álvarez-Campos et al. 2017).

### ***Trypanosyllis zebra* (Grube, 1860)**

**Notes:** Species complex. Due to its distinct colouration (dark dorsal stripes), *Trypanosyllis zebra* has long been considered a cosmopolitan species. Álvarez-Campos et al. (2017) recovered seven cryptic species worldwide, five of which are described as new species as they can also be morphologically distinguished. The analysis recovered at least three clades in the Mediterranean: *Trypanosyllis krohnii* Claparède, 1864 (re-instated, designated as type species of the genus and used as the preferred name for Mediterranean species), *Trypanosyllis* sp. 2 (differing from *Trypanosyllis krohnii* in bathymetrical distribution) and a clade from Crete (originally identified as *Trypanosyllis zebra* by Chatzigeorgiou et al. (2016), labelled *Trypanosyllis* cf. *krohnii* in the analyses). Greek specimens did not have enough support in the analyses to be placed unambiguously. Unfortunately, no material from the type locality of *Trypanosyllis zebra* in the Adriatic Sea was included in the analyses, so it remains unclear how many clades exist in the Mediterranean and whether *Trypanosyllis zebra* and *Trypanosyllis krohnii* are synonymous or distinct species. Until these ambiguities are resolved, all material from Greece is listed under the name *Trypanosyllis zebra*.

### ***Virchowia clavata* Langerhans, 1879**

**Nomenclature:** *Umbellisyllis clavata* (Langerhans, 1879) | *Virchowia clavata* Langerhans, 1879

**Notes:** Reported from Greece by Simboura (1996) and Faulwetter et al. (2011a). In the Mediterranean also known from Spain (San Martín 2003), Italy (Castelli et al. 2008) and France (Nygren and Pleijel 2010), otherwise known only from Madeira.

### ***Xenosyllis scabra* (Ehlers, 1864)**

**Notes:** Possibly a complex of pseudo-cryptic species. Campoy (1982) found differences in chaetal features in specimens of *Xenosyllis scabra* from different localities of the Iberian Peninsula. San Martín et al. (2008) describe two new species from Australia. One of these, *Xenosyllis scabroides* San Martín, Hutchings & Aguado, 2008, differs from *Xenosyllis scabra* only in the shape of the compound chaetae. The authors suggest that material from outside the type locality (Kvarner Gulf, Adriatic) should be carefully examined as it may belong to different, undescribed species.



## Terebellidae Johnston, 1846

### *Amphitrite cirrata* Müller, 1776

### *Amphitrite rubra* (Risso, 1826)

### *Amphitrite variabilis* (Risso, 1826)

Notes: Type locality: Mediterranean.

### *Amphitritides gracilis* (Grube, 1860)

### *Amphitritides kuehlmanni* Arvanitidis & Koukouras, 1995

Notes: Originally described from the Evoikos Gulf (38°28'55"N, 23°29'10"E, 5 m depth), no other records from Greece.

### *Axionice conchilega* (Pallas, 1766)

Nomenclature: *Lanice conchilega* (Pallas, 1766)

Notes: The genus *Lanice* Malmgren, 1866 is considered a synonym of *Axionice* Malmgren, 1866 by Jirkov and Leontovich (2017) based on a phylogenetic analysis of morphological characters. Hutchings et al. (2017) consider the genus *Lanice* valid but could not take the publication by Jirkov and Leontovich (2017) into account as it was published almost concurrently. There is general agreement that several Terebellidae genera are in need of revision (e.g. Matos Nogueira et al. 2013, Jirkov and Leontovich 2017, Hutchings et al. 2017). We here follow the nomenclature of the most recent publication by Jirkov and Leontovich (2017) until more information on the generic affinity of several Terebellidae species becomes available.

### *Axionice cretacea* (Grube, 1860)

Nomenclature: *Pista cretacea* (Grube, 1860)

Notes: *Pista cretacea* is considered to belong to *Axionice* by Jirkov and Leontovich (2017). Type locality: Mediterranean (Kvarner Gulf, Adriatic).

### *Axionice medusa* (Savigny in Lamarck, 1818)

Nomenclature: *Loimia medusa* (Savigny in Lamarck, 1818)

Native status: Non-native (established)

**Notes:** Considered a species complex by several authors (e.g. Hutchings and Glasby 1995). Originally described from the Red Sea, considered established in the Mediterranean Sea (Zenetos et al. 2017). The genus *Loimia* Malmgren, 1866 is considered a synonym of *Axionice* Malmgren, 1866 by Jirkov and Leontovich (2017), but see remarks under *Axionice conchilega*, which apply also to *Loimia*.

### ***Eupolymnia nebulosa* (Montagu, 1819)**

**Nomenclature:** *Eupolymnia nebulosa* (Montagu, 1819) | *Polymnia nebulosa* (Montagu, 1819)

**Notes:** Species complex. Different populations of *Eupolymnia nebulosa* show different life cycles (brooding vs. free spawning, Bhaud and Lenaers 1992). Sromek et al. (2016), using molecular methods, recovered four cryptic species in Europe: one along the south coast of Norway and three in the Western Mediterranean (of which two also occur in the Atlantic).

### ***Eupolymnia nesidensis* (Delle Chiaje, 1828)**

**Nomenclature:** *Eupolymnia nesidensis* (Delle Chiaje, 1828) | *Polymnia nesidensis* (Delle Chiaje, 1828)

**Notes:** Type locality: Mediterranean (Gulf of Naples).

### ***Neoamphitrite affinis* (Malmgren, 1866)**

**Notes:** Questionable status. Reported from Greece by Papazacharias (1991) from the North Aegean. In the Mediterranean also reported from the Adriatic (Mikac 2015) and Spain (Alós 1984), otherwise distributed in Arctic waters.

### ***Neoamphitrite edwardsi* (Quatrefages, 1865)**

**Nomenclature:** *Amphitrite edwardsi* (Quatrefages, 1865) | *Neoamphitrite edwardsi* (Quatrefages, 1865)

**Notes:** Questionable status. Reported from Greece by Karlou and Melea (1977). In the Mediterranean also known from Spain (Tena et al. 1991), Turkey (Çınar et al. 2014) and the Adriatic (Mikac 2015), otherwise distributed along the Atlantic coasts of Europe. While it is possible that the species occurs in Greece, it has so far only been reported in grey literature and has not been found in the last four decades, therefore its presence in Greece is here considered questionable.

***Neoamphitrite figulus* (Dalyell, 1853)**

**Nomenclature:** *Amphitrite johnstoni* Malmgren, 1865 | *Neoamphitrite figulus* (Dalyell, 1853)

***Neoamphitrite groenlandica* (Malmgren, 1866)**

**Nomenclature:** *Amphitrite groenlandica* Malmgren, 1866

**Notes:** Questionable status. In the Mediterranean only reported from Greece (Bogdanos and Satsmadjjs 1983, Nicolaidou and Papadopoulou 1989). Distributed in the North Atlantic.

***Nicolea venustula* (Montagu, 1819)*****Nicolea zostericola* Ørsted, 1844*****Pista cristata* (Müller, 1776)**

**Nomenclature:** *Pista cristata* (Müller, 1776) | *Pista lornensis* (Pearson, 1969) | *Scionella lornensis* Pearson, 1969

**Notes:** *Scionella lornensis* was synonymised with *Pista cristata* (Müller, 1776) by Jirkov and Leontovich (2017). However, both *Scionella lornensis* and *Pista cristata* are reported in the same publication by Arvanitidis (2000a) (who identified them using the key and descriptions by Holthe (1986)). Mikac and Hutchings (2017), in a paper published concurrently with the work by Jirkov and Leontovich (2017), consider several European records of *Scionella lornensis* (including those from Greece) as potential misidentifications and possibly belonging to either *Pista adriatica* Mikac & Hutchings, 2017 or *Pistella rovigensis* Mikac & Hutchings, 2017.

***Proclea graffii* (Langerhans, 1884)**

**Notes:** Questionable status. Reported from Greece by Bogdanos and Satsmadjjs (1983) and NCMR (1995). In the Mediterranean also reported from the Adriatic (Mikac 2015), otherwise distributed in the Arctic. Given the rarity of records in the Mediterranean and the native distribution range of the species, the Greek records are considered questionable.

***Terebella ehrenbergi* Gravier, 1906**

**Native status:** Non-native (questionable)

**Notes:** Questionable status. Reported from Greece by Arvanitidis (1994). Originally from the Red Sea, in the Mediterranean also reported from the western Mediterranean (Pérès 1954), Cyprus (Ben-Eliahu 1972b) and Israel (Ben-Eliahu 1976b). The taxonomy of the species is confused, as two different descriptions exist in literature: *Terebella ehrenbergi* sensu Day, 1967 and *Terebella ehrenbergi* sensu Rullier, 1972, differing in the shape of the notochaetae (denticulated / smooth) and the number of abdominal chaetigers with notochaetae (posterior 20–40 segments without notochaetae vs. posterior 10–15 segments) (Ben-Eliahu 1976b). Ben-Eliahu (1976b) compared material from the Israeli coasts of the Mediterranean with museum specimens from the Red Sea identified by Gravier and by Fauvel and found the former to lack calyx-type chaetae on abdominal chaetigers. However, as this may be a size-dependent character, it is currently unknown whether Mediterranean species belong to *Terebella ehrenbergi* Gravier, 1906 or to a different species.

***Terebella lapidaria* Linnaeus, 1767**

**Telothelepodidae Nogueira, Fitzhugh & Hutchings, 2013**

***Parathelepus collaris* (Southern, 1914)**

**Thelepodidae Hesse, 1917**

***Streblosoma bairdi* Malmgren, 1866)**

***Thelepus cincinnatus* (Fabricius, 1780)**

***Thelepus setosus* (Quatrefages, 186)**

**Notes:** Often regarded as cosmopolitan, but probably many of the worldwide records are misidentifications (Hutchings and Glasby 1987).

***Thelepus triserialis* (Grube, 1855)**

**Notes:** Type locality: Mediterranean.

## Trichobranchidae Malmgren, 1866

### *Terebellides stroemii* Sars, 1835

**Notes:** Questionable status. Species complex. *Terebellides stroemii* is a species complex of pseudo-cryptic species (Parapar and Hutchings 2014). Çınar (2005) re-examined some specimens from Cyprus previously identified as *Terebellides stroemii* and compared them to material collected near the type locality. He concluded that two or three different species of *Terebellides* occur along the coast of Cyprus. Parapar et al. (2013) re-investigated material from the Adriatic, found *Terebellides gracilis* (Malm, 1874) to be present in the area and additionally described a new species, *Terebellides mediterranea* Parapar, Mikac & Fiege, 2013. Thus, specimens reported in the Mediterranean under the name *Terebellides stroemii* certainly belong to different species.

### *Trichobranchus glacialis* Malmgren, 1866

## Typhloscolecidae Uljanin, 1878

### *Travisiopsis lanceolata* Southern, 1910

**Notes:** Reported from Greece by Wesenberg-Lund (1939a). In the Mediterranean also known from the Tyrrhenian Sea (Wesenberg-Lund 1939a) and the Adriatic (Mikac 2015), otherwise distributed in the North and South Atlantic (Dales and Peter 1972).

## Analysis

The present work comprises an updated and annotated inventory of polychaete species from Greek waters, compiled from literature reports, online databases, museum collections and unpublished datasets. In total, 999 species-level taxa have been reported from Greece since the first studies in 1832. Of these, 84 are currently considered absent from the country's waters (Table 1), 79 are heterotypic synonyms and the presence of a further 142 species is considered questionable (Fig. 2). These questionable species are included in the present inventory until additional evidence becomes available to justify their exclusion. Fourteen species are reported here for the first time from Greek waters. At least 52 species in the present list constitute in fact a complex of cryptic or pseudo-cryptic species. Forty-seven species are considered non-native to the area, ten of which are reported here for the first time for the country (Table 2). In addition to the species-level taxa reported in this checklist, eleven genera have been recorded from Greece with no representatives identified to species level (Table 3).

Table 1.

Species reported from Greece in previous publications but in fact absent from the area, with unclear identity (e.g. nomina *dubia*), or wrongly applied names.

| Family       | Scientific name                                       | Reported as                   | Remarks  |
|--------------|---|-------------------------------|--|
| Acoetidae    | <i>Euarche tubifex</i><br>Ehlers, 1887                | <i>Euarche tubifex</i>        | Arvanitidis (2000a) lists Aegean Sea records of <i>Eupanthalis kinbergi</i> McIntosh, 1876 under the name <i>Euarche tubifex</i> ; subsequently the name was included in the Greek checklist by Simboura and Nicolaidou (2001). Arvanitidis (2000a) follows Pettibone (1989) who assigns many Mediterranean records of <i>Eupanthalis kinbergii</i> , including those of Fauvel (1923), to <i>Euarche tubifex</i> . While it is highly likely that <i>Euarche tubifex</i> is present in Greece, it could not yet be confirmed. |
| Ampharetidae | <i>Amage gallasii</i><br>Marion, 1875                 | <i>Amage gallasii</i>         | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Ergen 1987); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Capitellidae | <i>Dasybranchus carneus</i> Grube,<br>1870            | <i>Dasybranchus carneus</i>   | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Ergen 1987); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Cirratulidae | <i>Timarete anchylochaeta</i><br>(Schmarda, 1861)     | <i>Timarete anchylochaeta</i> | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Sea of Marmara (Rullier 1963); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Cirratulidae | <i>Timarete dasylophius</i><br>(Marenzeller,<br>1879) | <i>Timarete dasylophius</i>   | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Sea of Marmara (Rullier 1963); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Cossuridae   | <i>Cossura delta</i><br>Reish, 1958                   | <i>Cossura delta</i>          | Reported by Faulwetter (2010). Records of <i>Cossura soyeri</i> Laubier, 1964 were reported as <i>Cossura delta</i> , as this was the accepted name in <a href="#">WoRMS</a> at the time of compilation.   |

|             |   |                                  |   |
|-------------|---|----------------------------------|---|
| Eunicidae   | <i>Eunice oerstedii</i><br>Stimpson, 1853       | <i>Eunice oerstedii</i>          | European specimens could belong to <i>Eunice dubitata</i> Fauchald, 1974 or a closely-related species (Wisnens 1989). The available descriptions of <i>Eunice oerstedii</i> are confusing and probably based on different species, thus its status was considered indeterminable by Fauchald (1992). Not listed in the key to global <i>Eunice</i> species by Carrera-Parra and Salazar-Vallejo (1998). |
| Eunicidae   | <i>Eunice stigmatura</i><br>(Verrill, 1990)     | <i>Eunice stigmatura</i>         | Recorded from a single specimen by Chatzigeorgiou et al. (2016) which was re-examined and found to be an incomplete and unidentifiable specimen.  |
| Eunicidae   | <i>Marphysa disjuncta</i><br>Hartman, 1961      | <i>Marphysa disjuncta</i>        | Reported from Greece by Simboursa et al. (2010). However, Kurt Şahin (2014) doubts its presence in the Mediterranean after re-examination of available material, including some of the Greek specimens of the former authors.   |
| Eunicidae   | <i>Lysidice hebes</i><br>(Verrill, 1900)        | <i>Nematonereis hebes</i>        | Reported by Faulwetter (2010). Records of <i>Lysidice unicornis</i> (Grube, 1840) were reported as <i>Nematonereis hebes</i> , as this was the accepted name in <a href="#">WoRMS</a> at the time of compilation.   |
| Fabriciidae | <i>Pseudofabriciola capensis</i> (Monro, 1937)  | <i>Pseudofabriciola capensis</i> | Records in OBIS (Arvanitidis 2000b) could not be traced back to any source and are believed to be a misidentification or database artefact.   |
| Fabriciidae | <i>Pseudofabriciola filamentosa</i> (Day, 1963) | <i>Fabricia filamentosa</i>      | Fitzhugh et al. (1994) re-examined all specimens identified previously as <i>Fabricia filamentosa</i> from Greece (Simboursa 1990) and described them as a new species, <i>Pseudofabriciola longipyga</i> Fitzhugh, Giangrande & Simboursa, 1994.   |
| Glyceridae  | <i>Glycera alba adspersa</i> Fauvel, 1939       | <i>Glycera alba adspersa</i>     | Reported by Faulwetter (2010), but in fact has only been recorded by Rullier (1963) from the Bosphorus and not from Greece. Considered indeterminable by Böggemann (2002).  |

|                |   |                                   |  |
|----------------|---|-----------------------------------|--|
| Glyceridae     | <i>Glycera papillosa</i><br>Grube, 1857                       | <i>Glycera papillosa</i>          | Considered indeterminable by Böggemann (2002). The holotype is lost and, based on the original description, it could be a synonym of either <i>Glycera capitata</i> Ørsted, 1843 or <i>Glycera lapidum</i> Quatrefages, 1866.  |
| Hesionidae     | <i>Leocrates chinensis</i><br>Kinberg, 1866                   | <i>Leocrates chinensis</i>        | The Mediterranean species <i>Leocrates claparedii</i> (Costa in Claparède, 1868) was considered synonymous with <i>Leocrates chinensis</i> by Hartman (1965a), the synonymy was revoked by Pleijel (1998). Greek specimens were initially identified as <i>Leocrates claparedii</i> using the key by Fauvel (1923) but reported under the name <i>Leocrates chinensis</i> . The latter name has survived in checklists and databases of introduced species (e.g. Zenetos et al. 2010, <a href="#">ELNAIS</a> ), but the specimens behind these records actually belong to the native species <i>Leocrates claparedii</i> . |
| Hesionidae     | <i>Neogyptis rosea</i><br>(Malm, 1874)                        | <i>Gyptis rosea</i>               | Recorded by Chatzigeorgiou et al. (2016), but the specimen was re-examined and actually belongs to <i>Neogyptis mediterranea</i> (Pleijel, 1983).  |
| Hesionidae     | <i>Orseis pulla</i><br>Ehlers, 1864                           | <i>Orseis pulla</i>               | Considered indeterminable by Pleijel (1998).   |
| Longosomatidae | <i>Heterospio reducta</i> Laubier,<br>Picard & Ramos,<br>1973 | <i>Heterospio reducta</i>         | Records in OBIS (Arvanitidis 2000b) could not be traced back to any source and is believed to be a misidentification or database artefact.   |
| Lumbrineridae  | <i>Lumbrineris labrofimbriata</i><br>Saint-Joseph,<br>1888    | <i>Lumbrineris labrofimbriata</i> | Species considered a juvenile form and thus invalid by Carrera-Parra (2006). Not listed as a Mediterranean species in recent regional reviews (Carrera-Parra et al. 2011, D'Alessandro et al. 2016).   |
| Lumbrineridae  | <i>Lumbrineris cingulata</i> Ehlers,<br>1897                  | <i>Lumbrineris cingulata</i>      | Reported by Chatzigeorgiou et al. (2016), but specimens were re-examined and actually belong to <i>Lumbrineris luciliae</i> Martins, Carrera-Parra, Quintino & Rodrigues, 2012.  |



|               |  |                               |  |
|---------------|--|-------------------------------|--|
| Lumbrineridae | <i>Lumbrineris perkinsi</i> Carrera-Parra 2001 | <i>Lumbrinerides perkinsi</i> | Name applied by Simboura et al. (2010) to literature records of <i>Lumbrineris inflata</i> Moore, 1911, based on conclusions by Çınar (2009). However, specimens subsequently identified as <i>Lumbrineris perkinsi</i> (unpublished data) were re-examined and found to belong to <i>Gallardoneris iberica</i> thus the presence of <i>Lumbrineris perkinsi</i> in Greece is not confirmed. |
| Lumbrineridae | <i>Scoletoma debilis</i> (Grube, 1878)         | <i>Scoletoma debilis</i>      | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean and the Sea of Marmara (Ergen 1987, Rullier 1963); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).   |
| Nephtyidae    | <i>Nephtys longosetosa</i> Ørsted, 1842        | <i>Nephtys longosetosa</i>    | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Ergen 1987); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Nephtyidae    | <i>Nephtys paradoxa</i> Malm, 1874             | <i>Nephtys paradoxa</i>       | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Ergen 1987); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Nereididae    | <i>Neanthes arenaceodentata</i> (Moore, 1903)  | <i>Nereis arenaceodonta</i>   | Multiple records in OBIS which are database artefacts. Originally recorded as <i>Nereis caudata</i> (sensu Delle Chiaje, 1827), which was listed as a synonym of <i>Nereis arenaceodentata</i> in WoRMS when datasets were submitted to OBIS. The latter name was erroneously used to standardise the taxonomy.  |
| Oeonidae      | <i>Arabella semimaculata</i> (Moore, 1911)     | <i>Arabella semimaculata</i>  | Reported by Chatzigeorgiou et al. (2016), but the specimen was re-examined and actually belongs to <i>Arabella iricolor</i> (Montagu, 1804).   |
| Opheliidae    | <i>Ophelia neglecta</i> Schneider, 1892        | <i>Ophelia neglecta</i>       | Specimens reported in OBIS belong in fact to <i>Ophelia roscoffensis</i> Augener, 1910.  |

|               |   |   |   |
|---------------|---|---|---|
| Paraonidae    | <i>Aricidea nolani</i><br>Webster &<br>Benedict, 1887       | <i>Aricidea nolani</i>  | Considered indeterminable (Hartley 1981), as the type material consists of fragments of two different species.  |
| Paraonidae    | <i>Paraonides neapolitana</i><br>(Cerruti, 1909)            | <i>Cirrophorus neapolitanus</i> ;<br><i>Paraonides neapolitana</i> ;<br><i>Paraonis neapolitana</i> | Considered of uncertain identity, as the original description may be based on artefacts resulting from the fixation method and the type material is lost (see Langeneck et al. (2017a) for detailed information).   |
| Paraonidae    | <i>Levinsenia flava</i><br>(Strelzov, 1973)                 | <i>Levinsenia flava</i>   | Greek specimens were re-examined by A. Castelli and identified tentatively as <i>Levinsenia</i> cf. <i>oculata</i> (Hartman, 1957), but possibly belong to an undescribed species.  |
| Phyllodocidae | <i>Lugia pterophora</i><br>(Ehlers, 1864)                   | <i>Lugia pterophora</i>   | Considered indeterminable by Pleijel (1991) as the original description is probably based on juveniles and the type material is lost.   |
| Phyllodocidae | <i>Paranaitis capensis</i> (Day, 1960)                      | <i>Paranaitis capensis</i>  | Considered an indeterminable species of <i>Paranaitis</i> by Kato and Pleijel (2003) who examined the holotype and found it in a poor condition preventing its identification.  |
| Phyllodocidae | <i>Phyllodoce lamelligera</i><br>(Gmelin in Linnaeus, 1788) | <i>Anaitides lamelligera</i> ;<br><i>Phyllodoce lamelligera</i>                                     | The species and its authorship are confused. The species was described by Gmelin in Linnaeus (1788), citing a " <i>Nereis lamellifera</i> Pallas 1788" but introducing a misspelling ( <i>lamellifera</i> became <i>lamelligera</i> ). Fauvel (1923) lists "Johnston, 1865" as the author, although Johnston (1865) merely reports the species from the UK. Both original descriptions (Gmelin and Pallas) could fit a variety of species. Therefore, Pleijel (1991) considers the species (listed by him as <i>Nereis lamellifera</i> ) indeterminable (see also Mikac 2015). Alós (2004) lists <i>Phyllodoce lamelligera</i> Johnston 1865 as a synonym of <i>Phyllodoce madeirensis</i> Langerhans, 1880, presumably on the basis of the description by Fauvel (1923), as the same incorrect authority is given. |

|               |   |   |  |
|---------------|---|---|--|
| Phyllodocidae | <i>Phyllodoce schmardaei</i> Day, 1963            | <i>Phyllodoce macrophthalma</i>                               | <i>Phyllodoce macrophthalma</i> Schmarda, 1861 is a junior homonym to <i>P. macrophthalma</i> Grube 1856. Day (1963) described <i>Phyllodoce macrophthalma</i> Schmarda into synonymy with it. Pleijel (1991), however, considers both <i>Phyllodoce macrophthalma</i> Schmarda and <i>Phyllodoce macrophthalma</i> Grube nomina <i>dubia</i> .  |
| Phyllodocidae | <i>Protomystides bidentata</i> (Langerhans, 1880) | <i>Mystides bidentata</i> ;<br><i>Protomystides bidentata</i> | Considered indeterminable by Pleijel (1991) due to the absence of type material. Alós (2004) provides a description of <i>Phyllodoce bidentata</i> , but gives no reference to the basis of the description (no type material or material from the type locality seems to have been studied). Thus, it is not certain whether the species of Alós (2004) is actually the same species as the one described by Langerhans. All Greek records are from before 2004 and therefore need to be re-examined. |
| Phyllodocidae | <i>Pterocirrus limbatus</i> (Claparède, 1868)     | <i>Pterocirrus limbata</i>                                    | Controversial species with a confused history of descriptions and many misidentifications (Alós 2004). Considered a nomen dubium by Pleijel (1991), re-described by Alós (2004) but not on the basis of type material or material from the type locality. The single record from Greece (Simboura 1996) would need to be re-examined to confirm its identity.  |
| Pilargidae    | <i>Ancistrosyllis cingulata</i> (Korschelt, 1893) | <i>Ancistrosyllis cingulata</i>                               | Larval form, probably indeterminable even to genus according to modern experts ( <a href="#">G .Read in WoRMS</a> ). Fiege and Böggemann (1999) do not include it in their revision of the genus.  |
| Pilargidae    | <i>Sigambra constricta</i> (Southern, 1921)       | <i>Sigambra constricta</i>                                    | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean and the Sea of Marmara (Ergen 1987, Rullier 1963); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).   |

|                  |   |   |  |
|------------------|---|---|--|
| Poecilochaetidae | <i>Elicodasia mirabilis</i> Laubier & Ramos, 1973 | <i>Elicodasia mirabilis</i>   | Nomen dubium, may be a posterior end of <i>Poecilochaetus fauchaldi</i> Pilato & Cantone, 1976 (Santos and Mackie 2008).   |
| Polycirridae     | <i>Polycirrus haematodes</i> (Claparède, 1864)    | <i>Polycirrus haematodes</i>  | Considered of uncertain identity by Glasby and Hutchings (2014) as the type material was not deposited, Claparède's description is insufficient to delimit the species and several species were possibly confused under this name in the past.                     |
| Polycirridae     | <i>Polycirrus pallidus</i> (Claparède, 1864)      | <i>Polycirrus pallidus</i>  | Considered of uncertain identity by Glasby and Hutchings (2014) as the type material is lost or was never deposited.   |
| Polycirridae     | <i>Polycirrus tenuisetis</i> Langerhans, 1881     | <i>Polycirrus tenuisetis</i>  | Considered of uncertain identity by Glasby and Hutchings (2014) as the type material is lost or was never deposited.   |
| Polynoidae       | <i>Eunoe mammiloba</i> Czerniavsky, 1882          | <i>Eunoe mammiloba</i>  | Records in OBIS (Dounas 1988) but not contained in original report, databasing error is assumed.   |
| Polynoidae       | <i>Harmothoe johnstoni</i> (McIntosh, 1876)       | <i>Harmothoe johnstoni</i>  | Indeterminable as the original description and the holotype are incomplete (Barnich and Fiege 2000a, Barnich and Fiege 2009).  |
| Polynoidae       | <i>Harmothoe minuta</i> (Potts, 1910)             | <i>Harmothoe minuta</i>   | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean and the Sea of Marmara (Ergen 1976, Rullier 1963). Considered indeterminable by Barnich and Fiege (2000a) as the original description and the holotype are incomplete. |
| Sabellariidae    | <i>Sabellaria alcocki fauveli</i>                 | <i>Sabellaria alcocki fauveli</i>                                   | Proposed by Arvanitidis (1994) as a new subspecies, but was never validly published and therefore does not constitute an available name. Even if it were available, it would constitute a junior homonym of <i>Sabellaria fauveli</i> Gravier, 1908.               |
| Sabellariidae    | <i>Tetereus porrectus</i> (Ehlers, 1908)          | <i>Lygdamis porrectus</i> ;<br><i>Lygdamis (Pallasia) porrectus</i> | Reported by Pérès (1959), but the specimen belongs in fact to <i>Lygdamis muratus</i> Allen, 1904 (Bhaud 1975, G. Bellan pers. comm. to C. Arvanitidis).   |

|                  |  |   |  |
|------------------|--|---|--|
| Sabellidae       | <i>Bispira crassicornis</i> (Sars, 1851)     | <i>Bispira crassicornis</i> ; <i>Sabella crassicornis</i> | Arvanitidis (2000a) listed Aegean Sea records of <i>Sabella fabricii</i> Krøyer, 1856 under the name <i>Bispira crassicornis</i> (following Fauvel (1927) in the synonymy but giving priority to the older name). The record in OBIS by Antoniadou (1998) from the North Aegean was originally recorded as <i>Sabella fabricii</i> (Antoniadou 2003); the name <i>Bispira crassicornis</i> was applied during data management. <i>Bispira crassicornis</i> is nowadays considered distinct from <i>Sabella fabricii</i> (Knight-Jones and Perkins 1998). |
| Sabellidae       | <i>Bispira mariae</i> Lo Bianco, 1893        | <i>Bispira mariae</i>                                     | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Knight-Jones et al. 1991); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Sabellidae       | <i>Bispira viola</i> (Grube, 1863)           | <i>Bispira viola</i>                                      | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Knight-Jones et al. 1991); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Sabellidae       | <i>Branchiomma moebii</i> Knight-Jones, 1994 | <i>Branchiomma moebii</i>                                 | Reported by Arvanitidis (2000a) for the Aegean Sea based on records from the Turkish Aegean (Knight-Jones et al. 1991); subsequently erroneously included in the Greek checklist by Simboura and Nicolaidou (2001).  |
| Sabellidae       | <i>Chone kroyeri</i> Sars, 1862              | <i>Chone kroyeri</i>                                      | Records in OBIS (Zenetos 1986) were subsequently found to be a misidentification.  |
| Sabellidae       | <i>Potamilla neglecta</i> (Sars, 1851)       | <i>Potamilla neglecta</i>                                 | Reported by Koukouras et al. (1992), record was referred to <i>Demonax brachychona</i> (Claparède, 1870) by Arvanitidis (1994).  |
| Scalibregmatidae | <i>Polyphysia crassa</i> (Ørsted, 1843)      | <i>Polyphysia crassa</i>                                  | Reported by Simboura (1996) but the specimen belongs in fact to the Mediterranean subspecies <i>Polyphysia crassa fauveli</i> (Laubier, 1959).   |
| Serpulidae       | <i>Hydroides uncinata</i> (Phillipi, 1844)   | <i>Hydroides uncinata</i>                                 | Considered indeterminable by Read et al. (2017), as the original taxon description is insufficient to identify the species and the name has been applied to a number of different species in the past (see also <a href="#">notes in WoRMS</a> )   |

|              |   |   |  |
|--------------|---|---|--|
| Serpulidae   | <i>Spirorbis corrugatus</i> (Montagu, 1803) | <i>Janua corrugatus</i> ; <i>Spirorbis (Janua) corrugatus</i> ; <i>Spirorbis corrugatus</i> | <i>Spirorbis corrugatus</i> (Montagu, 1803) is an insufficiently described species with a sinistrally coiled tube and type material is not retained (Knight-Jones et al. 1975, Knight-Jones and Knight-Jones 1977). Most authors actually refer to the dextral species described by Caullery and Mesnil (1897) which was re-described by Bush (1905) as <i>Spirorbis pseudocorrugatus</i> (now <i>Neodexiospira pseudocorrugata</i> ). The description of <i>Spirorbis corrugatus</i> by Fauvel (1927) refers in fact to <i>Neodexiospira pseudocorrugata</i> .  |
| Sigalionidae | <i>Pholoe fauveli</i> Kirkegaard, 1983      | <i>Pholoe fauveli</i>   | Kirkegaard (1983) examined specimens of <i>Pholoe synophthalmica</i> Claparède, 1868 by Fauvel (1923) and assigned them to <i>Pholoe fauveli</i> . Consequently, Arvanitidis (2000a) listed all Greek records of <i>Pholoe synophthalmica</i> under the name <i>Pholoe fauveli</i> , assuming that all of them were identified using the key by Fauvel (1923). <i>Pholoe synophthalmica</i> is nowadays considered a synonym of <i>Pholoe inornata</i> Johnston, 1839 and <i>Pholoe fauveli</i> is considered absent from the Mediterranean and possibly a synonym of <i>Pholoe inornata</i> (Barnich and Fiege 2003). |
| Spionidae    | <i>Dipolydora caulleryi</i> (Mesnil, 1897)  | <i>Polydora caulleryi</i>   | Radashevsky and Simboursa (2013) examined all Greek material and assigned it to <i>Dipolydora blakei</i> (Maciolek, 1984).   |
| Spionidae    | <i>Prionospio malmgreni</i> Claparède, 1870 | <i>Prionospio malmgreni</i>   | Considered indeterminable by Maciolek (1985) and Sigvaldadóttir and Mackie (1993). Reported to have up to 9 pairs of branchiae in the original description, a character that was never observed again (Sigvaldadóttir and Mackie 1993). The description of <i>Prionospio malmgreni</i> by Fauvel (1927) actually belongs to <i>Prionospio fallax</i> Söderström, 1920 (Sigvaldadóttir and Mackie 1993), thus Mediterranean records probably belong to <i>Prionospio fallax</i> (Mikac 2015, Çınar et al. 2014).  |

|           |   |                                  |   |
|-----------|---|----------------------------------|---|
| Spionidae | <i>Spio martinensis</i><br>Mesnil, 1896             | <i>Spio martinensis</i>          | Record in OBIS (Zenetos 1986) is possibly a misidentification and was later removed from a revised version of the dataset; these corrections were not incorporated in the OBIS data.  |
| Syllidae  | <i>Autolytus roseus</i><br>Claparède, 1864          | <i>Autolytus roseus</i>          | Indeterminable, as types do not exist; species probably is the female stolon of <i>Proceraea madeirensis</i> Nygren, 2004 (Nygren 2004).  |
| Syllidae  | <i>Dioplosyllis cirrosa</i><br>Gidholm, 1962        | <i>Dioplosyllis cirrosa</i>      | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus (Çınar and Ergen 2003).   |
| Syllidae  | <i>Epigamia macrophtalma</i><br>(Marenzeller, 1875) | <i>Autolytus sardai</i>          | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus (Çınar et al. 2003).  |
| Syllidae  | <i>Erinaceusyllis belizensis</i><br>(Russel, 1989)  | <i>Erinaceusyllis belizensis</i> | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çınar and Ergen 2002). |

|          |   |   |   |
|----------|---|---|---|
| Syllidae | <i>Eusyllis kupfferi</i><br>Langerhans,<br>1879                       | <i>Eusyllis kupfferi</i>                  | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus (Çınar and Ergen 2003) and Turkey (Çınar et al. 2014).  |
| Syllidae | <i>Inermosyllis balearica</i> (San Martín, 1982)                      | <i>Inermosyllis balearica</i>             | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Lebanon (Aguado and San Martín 2007).  |
| Syllidae | <i>Myrianida dentalia</i> (Imajima, 1966)                             | <i>Autolytus dentalius</i>                | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus (Çınar et al. 2003). Nygren (2004) restricts the distribution to the North-West Atlantic and North Pacific. |
| Syllidae | <i>Opisthodonta serratisetosa</i> (López, San Martín & Jiménez, 1997) | <i>Pionosyllis serratisetosa</i>          | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus (Çınar and Ergen 2003).   |
| Syllidae | <i>Parapionosyllis gestans</i><br>Pierantoni, 1903                    | <i>Parapionosyllis</i> cf. <i>gestans</i> | Reported by Simboursa (1996) as <i>Parapionosyllis</i> cf. <i>gestans</i> sensu Campoy 1892. Campoy's species was later described as a new species, <i>Parapionosyllis cabezali</i> Parapar, San Martín & Moreira, 2000 and Simboursa and Nicolaidou (2001) referred the Greek records to the latter.   |



|          |   |                                    |   |
|----------|---|------------------------------------|---|
| Syllidae | <i>Parapionosyllis labronica</i><br>Cognetti, 1965                | <i>Parapionosyllis labronica</i>   | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çinar and Ergen 2002). |
| Syllidae | <i>Paraprocerastea crocantinae</i> San Martín & Alós, 1989        | <i>Paraprocerastea crocantinae</i> | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çinar and Ergen 2002). |
| Syllidae | <i>Parexogone caribensis</i> (San Martín, 1991)                   | <i>Exogone caribensis</i>          | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çinar and Ergen 2002). |
| Syllidae | <i>Parexogone cognettii</i> Castelli, Badalamenti & Lardici, 1987 | <i>Exogone cognettii</i>           | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çinar et al. 2014).    |

|          |  |  |   |
|----------|--|--|---|
| Syllidae | <i>Prosphaerosyllis longipapillata</i> (Hartmann-Schröder, 1979) | <i>Prosphaerosyllis longipapillata</i>                           | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus (Çinar et al. 2003) and Turkey (Çinar 2009).  |
| Syllidae | <i>Streptospinigera templadoi</i> (San Martín, 1984)             | <i>Streptosyllis templadoi</i>                                   | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çinar and Ergen 2002). |
| Syllidae | <i>Syllis heterochaeta</i> Moore, 1909                           | <i>Langerhansia heterochaeta</i> ;<br><i>Syllis heterochaeta</i> | Records in OBIS are database artefacts: the species recorded in the original datasets is <i>Tharyx heterochaeta</i> / <i>Monticellina heterochaeta</i> Laubier, 1961; during taxon standardisation, taxa were incorrectly matched only on the specific epithet.   |
| Syllidae | <i>Syllis hyllebergi</i> (Licher, 1999)                          | <i>Syllis hyllebergi</i>   | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from Cyprus and the Israeli coast (Licher 1999)   |
| Syllidae | <i>Syllis lutea</i> (Hartmann-Schröder, 1960)                    | <i>Syllis lutea</i>  | Reported by Arvanitidis (1994) who used the key and description by San Martín (1984) for identification which actually belong to <i>Syllis jorgei</i> San Martín & López, 2000. <i>Syllis lutea</i> is absent from the Mediterranean (San Martín 2003). Arvanitidis' specimens are therefore assigned to <i>Syllis jorgei</i> .   |

|          |  |                                |  |
|----------|--|--------------------------------|--|
| Syllidae | <i>Syllis pontxioi</i><br>San Martín & López, 2000 | <i>Syllis pontxioi</i>         | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean, Cyprus and Turkey. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ and the Turkish record was wrongly placed into the Greek EEZ in the North Aegean. The only Eastern Mediterranean records are from the Turkish Aegean (Çınar and Ergen 2002) and from Cyprus (Çınar and Ergen 2003). |
| Syllidae | <i>Syllis schulzi</i><br>(Hartmann-Schröder, 1960) | <i>Syllis schulzi</i>          | Reported by Zenetos et al. (2011) as an alien species, with a reference to its first report by Arvanitidis (1994). However, Arvanitidis (1994) lists the species to occur in the Western Mediterranean, not in Greece.   |
| Syllidae | <i>Syllis vivipara</i><br>Krohn, 1869              | <i>Syllis vivipara</i>         | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Eastern Mediterranean. During digitisation, the centre of the Eastern Mediterranean was georeferenced, which falls into the Greek EEZ. The only Eastern Mediterranean records are from the Italian Salento (Corriero et al. 2004).   |
| Syllidae | <i>Trypanedenta gemmipara</i><br>(Johnson, 1901)   | <i>Trypanosyllis gemmipara</i> | Considered a synonym of <i>Trypanosyllis aeolis</i> Langerhans, 1897 by San Martín (2003), re-instated by Álvarez-Campos et al. (2017) and possibly restricted to the Indo-Pacific area. The only Greek specimens reported under this name by Arvanitidis (1994) actually belong to <i>Trypanosyllis aeolis</i> (see discussion there).  |
| Syllidae | <i>Trypanosyllis gigantea</i><br>(McIntosh, 1885)  | <i>Trypanosyllis gigantea</i>  | Records in OBIS were digitised from Musco and Giangrande (2005) who list it to occur in the Greek Aegean, based on Licher (1999) who in turn cites Arvanitidis (1994) for this record. However, Arvanitidis (1994) lists the species to occur in the Adriatic, not in the Aegean. The record is believed to be erroneous.  |

|              |  |                       |   |
|--------------|--|-----------------------|---|
| Terebellidae | <i>Axionice arborea</i><br>(Moore, 1903) | <i>Loimia arborea</i> | Reported by Faulwetter (2010). Records of <i>Loimia medusa</i> (Savigny in Lamarck, 1818) were reported as <i>Loimia arborea</i> , as this was the accepted name in <a href="#">WoRMS</a> at the time of compilation. The genus <i>Loimia</i> Malmgren, 1866 is considered a synonym of <i>Axionice</i> Malmgren, 1866 by Jirkov and Leontovich (2017). |
|--------------|--|-----------------------|---|

Table 2.

New reports of non-native species in Greece. Assessment of the establishment status in Greece follows the terminology and definitions by Zenetos et al. (2011).

| Species                        | Reference                                  | Date and location of first detection in Greece        | Origin                    | Other Mediterranean reports  | Establishment status in Greece |
|--------------------------------|--|---|---------------------------|--|--------------------------------|
| <i>Eurythoe complanata</i>     | Chatzigeorgiou et al. (2016)               | 2008: Alykes, Northern Crete                          | Caribbean                 | Cyprus (Ben-Eliahu 1972b), Israel (Ben-Eliahu 1976a, Arias et al. 2013a), Turkey (Ergen and Çınar 1997), Alboran Sea (Arias et al. 2013a)  | casual                         |
| <i>Linopherus canariensis</i>  | Chatzigeorgiou et al. (2016)               | 2007: Alykes, Northern Crete; Elounda, Northern Crete | Atlantic (Canary Islands) | Lebanon (Laubier 1966), Turkey (Ergen and Çınar 1997) (as <i>Pseudeurythoe acarunculata</i> ), Cyprus (Çınar 2009), Italy (Cosentino and Giacobbe 2010), Egypt (Dorgham et al. 2014) | cryptogenic                    |
| <i>Leiocapitellides analis</i> | Akoumianaki (2004); Neofitou et al. (2014) | 2000-2001: Maliakos Gulf                              | Red Sea                   | Egypt (Abd-Elnaby 2009)  | casual                         |
| <i>Mediomastus capensis</i>    | Maidanou et al. (2017)                     | Elounda Bay, Northern Crete                           | South Africa              | Italy (Castelli et al. 2008), Cyprus (Çınar 2005), Spain (Capaccioni-Azzati and El-Haddad 2015)  | casual                         |
| <i>Timarete punctata</i>       | Maidanou et al. (2017)                     | 2006: Elounda Bay, Northern Crete                     | Caribbean                 | Turkey (Çınar et al. 2014)   | casual                         |

|                                 |  |   |                |   |        |
|---------------------------------|--|---|----------------|---|--------|
| <i>Lumbrinerides neogesae</i>   | Papageorgiou et al. 2006 (unpublished); this study | 2002-2003: Elaфонisi Island, South-West Crete         | South Africa   | Italy (Gravina and Cantone 1991)  | casual |
| <i>Leonnates persicus</i>       | this study   | 2013: Argolikos Gulf                                  | Indo-Pacific   | Israel (Ben-Eliahu 1991b), Turkey (Çınar et al. 2002)                                       | casual |
| <i>Neanthes agulhana</i>        | Chatzigeorgiou et al. (2016)                       | 2007: Alykes, Northern Crete; Elounda, Northern Crete | South Africa   | Spain (Núñez 2004), Italy (Occhipinti-Ambrogi et al. 2011)                                  | casual |
| <i>Lepidonotus tenuisetosus</i> | Chatzigeorgiou et al. (2016)                       | 2008: Alykes, Northern Crete                          | Red Sea        | Israel and Egypt (Barnich and Fiege 2003), Turkey (Çınar et al. 2014), Adriatic Mikac 2015) | casual |
| <i>Exogone brevantennata</i>    | Dando et al. (1995)                                | 1992: Milos   | circumtropical | Lebanon (Aguado and San Martín 2007), Turkey (Çınar et al. 2014)                            | casual |

Table 3.

Taxa reported at genus level with no species-level representatives known from Greece

| Genus   | Notes   |
|---|---|
| <i>Abyssoninoe</i><br>Orensanz, 1990          | New record for Greece. Two specimens, off Limnos island NE Aegean, 77–220 m depth, collected in the framework of the Perseus project (Suppl. material 1). |
| <i>Acanthicolepis</i><br>McIntosh, 1900       | Reported by Koukouras et al. (1985).  |
| <i>Branchiocapitella</i><br>Fauvel, 1932      | Reported by Vamvakopoulou (1991). Currently considered a synonym of <i>Capitella</i> Blainville, 1828 (Magalhães and Bailey-Brock 2012).                  |
| <i>Apistobranchus</i><br>Levinsen, 1883       | Reported by Eleftheriou et al. (1990), Simboura (1996).   |
| <i>Fauveliopsis</i><br>McIntosh, 1922         | Reported by Eleftheriou et al. (1990), Karakassis (1991), Simboura et al. (1995a), Simboura (1996).   |
| <i>Hipponoe</i> Audouin & Milne Edwards, 1830 | Reported by Simboura et al. (1995b).  |
| <i>Meiodorvillea</i><br>Jumars, 1974          | Reported by Simboura (1996).  |

|  |   |
|--|---|
| <i>Perolepis</i> Ehlers, 1908            | Reported by Simboura and Zenetos (2005).  |
| <i>Polygordius</i> Schneider, 1868       | Reported by NCMR (1995), NCMR (1997), NCMR (1998).  |
| <i>Tomopteris</i> Eschscholtz, 1825      | Reported by Ramfos et al. (2005), Ramfos et al. (2006), Isari et al. (2010).  |
| <i>Paraleiocyathella</i> Thomassin, 1970 | <i>Paraleiocyathella</i> is considered a synonym of <i>Notomastus</i> (Ewing, 1982), although Green (2002) argues that <i>Paraleiocyathella</i> should be differentiated from <i>Notomastus</i> based on chaetal characters. <i>Paraleiocyathella</i> is monotypic, thus if the generic identification of the Greek specimens (Nicolaidou and Papadopoulou 1989, Eleftheriou et al. 1990, Simboura 1996) is correct, they probably belong to <i>Notomastus mossambicus</i> (Thomassin, 1970), an alien species first reported from the Mediterranean by Çınar (2005). |




Figure 2. doi

Numbers of polychaete species per family in Greece and their status (present / questionable).

Of the 836 species-level taxa reported from Greece and not explicitly excluded from this checklist, Syllidae are the most species-rich family (133 species / 15.9% of all species), followed by the Serpulidae (62 / 7.4%), Spionidae (54 / 6.5%), Paraonidae (39 / 4.7%) , Sabellidae (38 / 4.5%), Phyllodocidae (37 / 4.4%), Polynoidea (31 / 3.7%) and Maldanidae (30 / 3.6%). Together, these eight families constitute 50.7% of the Greek polychaete species richness (Fig. 3). Twenty-six families are represented by less than 5 species and 17 families have never been reported from Greece: Aberrantidae Wolf, 1987, Alvinellidae Desbruyères & Laubier, 1986, Antonbruunidae Fauchald, 1977, Dinophilidae Remane, 1925, Hartmaniellidae Imajima, 1977, Histriobdellidae Vaillant, 1890, Ichthyotomidae Eisig, 1906, Iphionidae Kinberg, 1856, Laetmonectidae Buzhinskaya, 1986, Nerillidae Levinsen, 1883, Parergodrilidae Reisinger, 1925, Pontodoridae Bergström, 1914, Protodriloididae Purschke & Jouin, 1988, Psammodrilidae Swedmark, 1952, Traviisiidae Hartmann-Schröder, 1971, Trochochaetidae Pettibone, 1963 and Uncispionidae Green, 1982.




Figure 3. [doi](#)

Percentages of species numbers of the eight most species-rich polychaete families in Greek waters compared to the remaining families.

The Greek polychaete fauna was poorly studied until the 1960-70s, with only occasional publications and a total number of 108 species reported from Greek waters until 1960. Subsequently, marine research was performed much more systematically and studies on polychaete communities were published frequently, adding a substantial number of species to the national faunal inventory (Fig. 4). Up to date, a total of 18 species-level taxa new to science were described from the country’s waters (Table 4). Despite the continuous research on benthic communities, 285 species (34.1%) have been only reported once or twice from Greek waters and 90 species have not been reported since 1993 (Table 5, Suppl. material 2). Since the publication of the last checklist of Greek polychaetes by Simboura and Nicolaidou (2001), 102 species were newly recorded from Greek waters. Another 53 species, reported from Greece before 2001 but not listed by Simboura and Nicolaidou (2001), were recovered from literature as a result of increased availability of electronic resources and the inclusion of habitats not covered previously (deep sea, pelagic waters).

Table 4.

Species originally described from Greek waters (i.e. the Exclusive Economic Zone as defined by [marineregions.org](http://marineregions.org)).

| Family       | Described as                                      | Currently accepted as                              | Type locality  |
|--------------|---|--|--|
| Acrocirridae | <i>Flabelligella mediterranea</i><br>Kolmer, 1985 | <i>Flabelligena mediterranea</i><br>(Kolmer, 1985) | Hellenic Trench, Matapan Deep<br>4690 m, mud, 35°49.8'N, 22°<br>20.7'E |
| Ampharetidae | <i>Uschakovius enigmaticus</i><br>Laubier, 1973   | <i>Uschakovius enigmaticus</i><br>Laubier, 1973    | Hellenic Trench, Matapan Deep<br>3174 m, 36°01.8'N, 22°24.6'E          |

|                     |  |  |  |
|---------------------|--|--|--|
| Amphinomidae        | <i>Amphinome savigny</i> Brullé, 1832                                  | <i>Hermodice carunculata</i> (Pallas, 1776)                            | Methana, Saronikos Gulf, Greece  |
| Capitellidae        | <i>Peresiella clymenoides</i> Harmelin, 1968                           | <i>Peresiella clymenoides</i> Harmelin, 1968                           | Mediterranean Sea (Marseilles, Crete and Santorini)  |
| Fabriciidae         | <i>Pseudofabriciola longipyga</i> Fitzhugh, Giangrande & Simbora, 1994 | <i>Pseudofabriciola longipyga</i> Fitzhugh, Giangrande & Simbora, 1994 | Geras Gulf, Lesvos Island, 21 m, muddy sand  |
| Lacydoniidae        | <i>Lacydonia laureci</i> Laubier, 1975                                 | <i>Lacydonia laureci</i> Laubier, 1975                                 | Hellenic Trench, Matapan Deep 4690 m, mud, 35°49.8'N, 22° 20.7'E                                 |
| Asteriomyzostomidae | <i>Myzostoma asteriae</i> Marenzeller, 1895                            | <i>Asteriomyzostomum asteriae</i> (Marenzeller, 1895)                  | Santorini, Kythira and Samos   |
| Onuphidae           | <i>Paradiopatra calliopae</i> Arvanitidis & Koukouras, 1997            | <i>Paradiopatra calliopae</i> Arvanitidis & Koukouras, 1997            | Aegean Sea, off Mount Athos coast, 80-300 m, silty sand to silt, 40°20'25"N, 24°18'05"E          |
| Orbiniidae          | <i>Scoloplos chevalieri candiensis</i> Harmelin, 1969                  | <i>Scoloplos chevalieri candiensis</i> Harmelin, 1969                  | Kaloi Limenes, Crete, 7-11 m, <i>Halophila stipulacea</i> and <i>Cymodocea nodosa</i> meadows    |
| Sabellariidae       | <i>Phalacrostemma cidariophilum</i> Marenzeller, 1895                  | <i>Phalacrostemma cidariophilum</i> Marenzeller, 1895                  | Several locations near Crete and Kythira   |
| Serpulidae          | <i>Serpula graeca</i> Brullé, 1832                                     | <i>Protula intestinum</i> (Lamarck, 1818)                              | Methana, Saronikos Gulf, Greece  |
| Serpulidae          | <i>Spirorbis (Janua) gnomonicus</i> Bailey, 1969                       | <i>Janua heterostropha</i> (Montagu, 1803)                             | Emborios Cave, Chios, Aegean Sea, 8-10 m   |
| Serpulidae          | <i>Spirorbis (Janua) parvulus</i> Bailey, 1969                         | <i>Neodexiospira pseudocorrugata</i> (Bush, 1905)                      | Emborios Cave, Chios, Aegean Sea   |
| Serpulidae          | <i>Vermilia agglutinata</i> Marenzeller, 1893                          | <i>Semivermilia agglutinata</i> (Marenzeller, 1893)                    | Mediterranean Sea (Libya and Greece)   |
| Sigalionidae        | <i>Pholoe dorsipapillata</i> Marenzeller, 1893                         | <i>Pholoides dorsipapillatus</i> (Marenzeller, 1893)                   | Mediterranean Sea (Greece and Egypt)   |
| Syllidae            | <i>Myrianida rodosensis</i> Çınar, 2015                                | <i>Myrianida rodosensis</i> Çınar, 2015                                | Rhodes, Faliraki port, 1 m, photophilic algae, 36°20.13'N, 28°12.52'E,                           |
| Syllidae            | <i>Streptosyllis nunezi</i> Faulwetter et al., 2008                    | <i>Streptosyllis nunezi</i> Faulwetter et al., 2008                    | Pachia Ammos, Crete, 1-5 m, coarse sand, 35°06'43"N, 25° 48'34"E                                 |
| Terebellidae        | <i>Amphitritides kuehlmanni</i> Arvanitidis & Koukouras, 1995          | <i>Amphitritides kuehlmanni</i> Arvanitidis & Koukouras, 1995          | Evoikos Gulf, coast of Loukissia, 5 m, on rocks with <i>Cystoseira</i> , 38°28'55"N, 23° 29'10"E |



Table 5.

Species not found in Greece since 1993 (chosen as a cut-off date, as thorough investigations of the Greek polychaete fauna (e.g. Arvanitidis 1994, Simboursa 1996) followed afterwards. Except for museum specimens, dates are publication dates, not collection dates.

| Year of last report | Species  |
|---------------------|--|
| 1887                | <i>Spinther arcticus</i>   |
| 1895                | <i>Asteriomyzostomum asteriae</i>  |
| 1902                | <i>Aglaophamus malmgreni</i> , <i>Leocrates atlanticus</i>   |
| 1939                | <i>Krohnia lepidota</i> , <i>Lopadorrhynchus appendiculatus</i> , <i>Lopadorrhynchus brevis</i> , <i>Lopadorrhynchus nationalis</i> , <i>Lopadorrhynchus uncinatus</i> , <i>Rhynchonereella moebii</i> , <i>Torrea candida</i> , <i>Travisioipsis lanceolata</i> , <i>Vanadis crystallina</i> , <i>Vanadis formosa</i> |
| 1959                | <i>Aphrodita perarmata</i> , <i>Phyllochaetopterus gracilis</i>  |
| 1961                | <i>Alkmaria romijni</i>  |
| 1962                | <i>Alitta virens</i> , <i>Bushiella (Jugaria) granulata</i>  |
| 1964                | <i>Phalacrostemma cidariophilum</i>  |
| 1969                | <i>Naineris quadratriceps</i> , <i>Vinearina endoumensis</i>   |
| 1971                | <i>Ninoe nigripes</i> , <i>Paradiopatra lepta</i>  |
| 1973                | <i>Bathyaufvelia affinis</i> , <i>Uschakovius enigmaticus</i>  |
| 1975                | <i>Lacydonia laureci</i>   |
| 1977                | <i>Neoamphitrite edwardsi</i>  |
| 1982                | <i>Hyboscolex longiseta</i> , <i>Odontosyllis dugesiana</i>  |
| 1983                | <i>Aonidella cirrobranchiata</i> , <i>Epigamia alexandri</i> , <i>Eulalia bilineata</i> , <i>Eunice floridana</i> , <i>Eunice roussaei</i> , <i>Euphrosine myrtosa</i> , <i>Manayunkia aestuarina</i> , <i>Pareurythoe borealis</i>  |
| 1984                | <i>Anoplosyllis edentula</i> , <i>Ophryotrocha labronica</i> , <i>Parapionosyllis elegans</i>  |
| 1985                | <i>Prionospio decipiens</i>  |
| 1986                | <i>Antinoe aquiseta</i> , <i>Aricidea (Strelzovia) quadrilobata</i> , <i>Lumbrinerides acuta</i> , <i>Maldane capensis</i> , <i>Maupasia coeca</i> , <i>Phylo norvegica</i> , <i>Scolecopsis gilchristi</i> , <i>Scoloplos haasi</i> , <i>Spiophanes duplex</i>  |
| 1987                | <i>Aricidea (Aricidea) longicirrata</i> , <i>Goniadella gracilis</i> , <i>Johnstonia clymenoides</i> , <i>Metavermilia acanthophora</i> , <i>Opisthodonta longocirrata</i> , <i>Streptosyllis bidentata</i> , <i>Syllis nigricirris</i>  |
| 1988                | <i>Lindrilus flavocapitatus</i>  |
| 1989                | <i>Desdemona ornata</i> , <i>Eunice norvegica</i> , <i>Gattyana cirrhosa</i> , <i>Proclea graffii</i> , <i>Protocirrinieris chrysoderma</i> , <i>Sabellaria eupomatoides</i>   |

|      |   |
|------|---|
| 1990 | <i>Neoamphitrite groenlandica</i> , <i>Oxydromus agilis</i> , <i>Praxillura longissima</i> , <i>Syllides longocirratu</i> s   |
| 1991 | <i>Arabella geniculata</i> , <i>Autolytus rubrovittatus</i> , <i>Boccardiella cf. ligerica</i> , <i>Claparedepelogenia inclusa</i> , <i>Malacoceros tetracerus</i> , <i>Neoamphitrite affinis</i> , <i>Neoleanira tetragona</i> , <i>Nephtys ciliata</i> , <i>Notomastus lineatus</i> , <i>Paranaitis kosteriensis</i> , <i>Phylo kupperi</i> , <i>Protodorvillea atlantica</i> , <i>Scoletoma tetraura</i> |
| 1992 | <i>Ampharete grubei</i> , <i>Bathyeremia langerhansi</i> , <i>Eunice schizobranchia</i> , <i>Eunoe assimilis</i> , <i>Nicomache trispinata</i> , <i>Polyodontes frons</i>   |
| 1993 | <i>Diplocirrus hirsutus</i> , <i>Levinsonia multibranchiata</i> , <i>Pelagobia longicirrata</i> , <i>Perinereis tenuisetis</i> , <i>Pseudomystides limbata nigrolineata</i>   |




Figure 4. [doi](#)

Cumulative numbers of known polychaete records from the Greek EEZ (blue line), superimposed on the number of publications containing polychaete records (red bars). Checklists and other secondary works are excluded.

The present list reports 836 species from Greece (including those considered questionable). This corresponds to 75.6% of the species reported by Martin and Gil (2010) from the Mediterranean (1,105 species, standardised to currently accepted names). Simboura and Nicolaidou (2001) report 753 species from Greece vs. 1,015 species from the Mediterranean, corresponding to 74.18%. Martín and Gil (2010) list 359 species in the Mediterranean checklist that are not known from Greece, whereas the Greek checklist contains 67 species not contained in the Mediterranean checklist (excluding questionable species and those described after 2010). Compared to the most recent (and thus comparable) inventories from neighbouring countries and regional seas, Greece shows similar species numbers to Italy (872 species, Castelli et al. 2008) and slightly higher numbers than the Adriatic Sea (756 species, Mikac 2015) and the Mediterranean coasts of Turkey (excluding the Sea of Marmara; 641 species, Çınar et al. (2014); all lists standardised to currently accepted names and including species indicated as questionable). Species unique to each of these checklists (not contained in any of the other

three checklists) number 147 for Greece, 48 for the Adriatic, 135 for Italy and 70 for the Turkish Mediterranean (excluding the Sea of Marmara).

## Discussion

Advanced imaging and molecular techniques have given a boost to polychaete systematics and taxonomy. This has not only resulted in up-to-date taxonomic descriptions, reviews and identification keys but also in a refined understanding of regional polychaete diversity and biogeography. Many polychaete species used to be considered cosmopolitan or widely distributed (e.g. Hutchings 1998). Lately, however, there is increasing evidence that these widely distributed species actually comprise endemic forms and cryptic species. Thus, many species seem to be restricted to much smaller geographical areas than was previously assumed (examples include *Chone duneri* (Tovar-Hernández et al. 2007), *Cirratulus cirratus* (Weidhase et al. 2014), *Terebellides stroemii* (Parapar et al. 2013), *Marphysa sanguinea* (Hutchings et al. 2012), *Inermonephtys inermis* (Ravara et al. 2010), *Myriochele heeri* (Parapar 2006), *Pholoe minuta* (Barnich and Fiege 2003), *Magelona cincta* (Wilson 1958)) and many of these species are probably absent from the Mediterranean.

One reason for worldwide reports of many species and thus their assumed cosmopolitanism lies in the use of inadequate literature for the identification of species (Hutchings 1998, Klautau et al. 1999). This is also the case in the Mediterranean, where many of the reported “cosmopolitan” or boreal species were identified using literature covering non-Mediterranean species. Thus, especially reports from older literature need to be treated with care, as many of these specimens may in fact belong to different, possibly endemic, species. Several Mediterranean taxa display morphological differences compared to material from the native distribution range of the species they were originally assigned to, indicating the existence of endemic forms (e.g. *Ophelia roscoffensis* Augener, 1910, *Galathowenia oculata* (Zachs, 1923), *Syllis rosea magna* (Westheide 1974) sensu Ben-Eliahu 1977, *Aphelochaeta monilaris* (Hartman, 1960) or *Therochaeta flabellata* (Sars in Sars, 1872)). Increasingly, many of these taxa are being revised and either described as species new to science or re-instated as previously synonymised Mediterranean species (e.g. *Acromegalomma lanigerum* (Grube, 1846), *Sternaspis thalassemoides* Otto, 1821, *Pherusa mikacae* Salazar-Vallejo, 2014, *Prionospio maciolekae* Dağlı and Çınar 2011, *Syllis parapari* San Martín & López, 2000 or *Terebellides mediterranea* Parapar, Mikac & Fiege, 2013). Over the last years, the increasing availability of updated keys which employ reliable, unique and age-independent diagnostic characters have allowed for a much finer distinction between species and have led to many reports of species new for Greece or the Mediterranean.

Another factor causing uncertainty, concerning the diversity and distribution of polychaete species worldwide, is the increasing discovery of cryptic species through molecular methods (Nygren 2014). Carr et al. (2011) show that polychaete diversity in Canadian waters is highly underestimated, as molecular analyses have revealed one quarter of the

333 morphologically recognised species to actually comprise two or more putative cryptic species. Similarly, many species occurring in the Mediterranean have been shown to actually form complexes of genetically distinct species within the basin (e.g. *Eulalia viridis* (Linnaeus, 1767), *Eurythoe complanata* (Pallas, 1776), *Lysidice ninetta* Audouin & Milne Edwards, 1833, *Myxicola infundibulum* Montagu, 1808, *Ophryotrocha puerilis* Claparède & Mecznirow, 1869, *Perinereis cultrifera* Grube, 1840, *Platynereis dumerilii* (Audouin & Milne Edwards, 1834), *Syllis gracilis* Grube, 1840 or *Trypanosyllis zebra* (Grube, 1860)). As only few of these studies include material from Greece, it remains unknown whether Greek populations of these species actually belong to the nominal taxon or to a different sibling species within the complex. Currently, 51 names (6.1%) of the Greek inventory are considered to refer to species complexes and resolving them will lead to major advances concerning our knowledge of the polychaete fauna in Greece and the Mediterranean.

This increased research effort contributes to an enhanced understanding of polychaete diversity, systematics and distribution but also introduces a high level of uncertainty for legacy records. The re-instatement or description of new species from a complex of cryptic or pseudo-cryptic species invariably renders most literature records under this name questionable, as they could belong to either of the species within this complex (e.g. *S calibregma inflatum* Rathke, 1843, *Spiophanes reyssi* Laubier, 1964, *Stylarioides monilifer* Delle Chiaje, 1841, *Sternaspis scutata* Ranzani, 1817, *Branchiosyllis exilis* (Gravier, 1900) or *Haplosyllis spongicola* (Grube, 1855)). However, in some cases the identity of these records can be resolved *a posteriori*, if information on the literature used for identification is available. A general recommendation of good practice is, therefore, to always provide a list of the literature used for identification alongside any published list of species. Similarly, revisionary publications and species lists should, where possible, report not only the currently accepted name but also the subjective synonyms under which the species were originally recorded. Failing to do so may lead to incorrect distribution records if the synonymised species is re-instated later (e.g. see reports of the non-native species *Cossura delta* Reish, 1958, *Leocrates chinensis* Kinberg, 1866, *Loimia arborea* Moore, 1903 and *Nematonereis hebes* Verrill, 1900 from Greece (Table 1) which had originally been recorded as *Cossura soyeri* Laubier, 1964, *Leocrates claparedii* (Costa in Claparède, 1868), *Loimia medusa* (Savigny in Lamarck, 1818) and *Nematonereis unicornis* (Grube, 1840) but were reported from Greece using the former names which were accepted at that time).

Despite a high number of changes compared to the last published checklist (Simboura and Nicolaidou 2001), the Greek polychaete fauna continues to represent ca. 75% of the Mediterranean inventory (74.2% in 2001, 75.6% this study). However, many families remain understudied in Greece and have only been reported at genus level or not at all. Most of these families include species associated with certain habitats which in turn are understudied, such as deep sea, pelagic and interstitial habitats and hard substrates (especially coralligenous formations and caves). More research effort needs to be targeted towards these habitats, as the knowledge on these communities is of the utmost importance for ecological studies assessing ecosystem functioning, food webs, habitat conservation and changes caused by invasions of non-native species (to which Greece is

particularly exposed due to its geographic location). In addition, as morphology alone underestimates species richness (Nygren 2014), more molecular studies are needed to resolve the identity of species complexes in Greece.

The number of species from Greece is similar to those of neighbouring areas from which recent checklists are available. However, individual checklists are influenced by research effort, different criteria for the inclusion or exclusion of species and the application of different species concepts. Thus, these comparisons can only serve as rough estimates of biogeographic relatedness and species richness. Nevertheless, each of these inventories comprises a relatively large number of names not reported from other areas. This can be partly explained by the aforementioned factors, partly by the actual biogeographic distribution and partly by incorrect species identifications. The Greek checklist contains 40 species that are of non-Mediterranean origin and have – to the best of our knowledge – not been reported from other areas in the Mediterranean. Where these species have only been found once or only by the same authors, misidentifications or an author bias must be assumed.

In fact, regional patterns of polychaete diversity in the Mediterranean seem to be biased by researcher identity (Chatzigeorgiou et al. 2017) and existing inventories are characterised by a large number of questionable species (e.g. 23.7% of the Mediterranean inventory (Arvanitidis 1994); 16% of the current Greek inventory). Geographically inappropriate or outdated literature contributes to this confusion (e.g. Hutchings 1998). Several updated keys and revisions for southern European polychaetes have been published recently (e.g. the *Fauna Ibérica* series) and the internet greatly facilitates access to literature. Nevertheless, comprehensive revisions and updated identification keys are lacking for many taxa in the Mediterranean and the use of century-old treatises such as those by (Fauvel 1923, Fauvel 1927) as the main source for species identification is still common practice in many laboratories.

The Mediterranean Sea is a hotspot of biodiversity and at the same time subject to a range of vectors of change such as habitat loss and degradation, climate change, pollution and invasion of alien species (Coll et al. 2010). To estimate the impact of these changes on biodiversity and ecosystem functioning, accurate knowledge of species' spatio-temporal distribution patterns is of crucial importance. The observed high levels of uncertainty regarding the regional fauna provide obstacles to such investigations and need to be resolved. The problem could be addressed through the organisation of regular pan-Mediterranean training and intercalibration workshops. These ring tests would train technicians and scientists in the identification of difficult or rarely encountered groups, calibrate and improve identification skills across areas and taxonomic groups and should result in a list of recommended and frequently updated identification literature and resources.

## Acknowledgements

The authors would like to thank Dr. Nicolas Bailly (HCMR) for providing an initial, automatically compiled list of species which served as a starting point for this work, Dr. Nafsika Papageorgiou (HCMR) for providing an unpublished dataset of meiofaunal polychaetes from sandy beaches, Kleoniki Keklikoglou (HCMR) for help with literature and Dr Helmut Sattmann (Natural History Museum in Vienna) for providing support and access to the collections. Many thanks go to Ruth Barnich, Rômulo Barroso, Markus Böggemann, Alberto Castelli, Melih Çınar, Pat Hutchings, Joachim Langeneck, Marco Lezzi, Alejandro Martínez García, Roberto Martins, Jorge Núñez, Julio Parapar, Ascensão Ravara, Rossana Sanfillipo and Argyro Zenetos. This work was supported by the [LifeWatchGreece](#) Research Infrastructure (MIS 384676), funded by the Greek Government under the [General Secretariat of Research and Technology](#) (GSRT), [ESFRI](#) Projects, National Strategic Reference Framework (NSRF).

## References

- Abd-Elnaby FA (2008) New recorded species of Magelonidae and Maldanidae (Annelida: Polychaeta) from El-Tina Bay, Mediterranean coast of Egypt. *Egyptian Journal of Aquatic Biology & Fisheries* 12 (4): 119-141. URL: <http://www.vliz.be/imisdocs/publications/247382.pdf>
- Abd-Elnaby FA (2009) Polychaete Study in Northeastern Mediterranean Coast of Egypt. *World Journal of Fish and Marine Sciences* 1 (2): 85-93. URL: <http://www.vliz.be/imisdocs/publications/247412.pdf>
- Abd-Elnaby FA (2010) New record of Lessepsian syllid polychaetes in the North Mediterranean Coast of Egypt. *Egyptian Journal of Experimental Biology (Zoology)* 6 (2): 297-306. URL: <http://www.ejmanager.com/mnstemp/3/3-1430780151.pdf>
- Abd-Elnaby FA, Gab-Alla AA (2007) New records of *Pettiboneia* and *Protodorvillea* (Polychaeta: Dorvilleidae) from the eastern harbour of Alexandria. *Journal of the Egyptian German Society of Zoology* 54 (D): 159-174.
- Aguado MT, San Martín G (2007) Syllidae (Polychaeta) from Lebanon with two new reports for the Mediterranean Sea. *Cahiers de Biologie Marine* 48 (2): 207-224. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Aguirrezabalaga F (2012) Familia Paraonidae Cerruti, 1909. In: Ramos, M.A. et al. *Annelida Polychaeta III*. In: Parapar J, Alós C, Núñez J, Moreira J, López E, Aguirrezabalaga F, Besteiro C, Martínez A (Eds.). *Fauna Ibérica*. 36. Museo Nacional de Ciencias Naturales, CSIC, Madrid, 160-272 pp. [In Spanish].
- Aguirrezabalaga F, Gil J (2008) Paraonidae (Annelida: Polychaeta) of the inlet of Zarautz (Basque Coast, Bay of Biscay), with new records from the Atlantic and the Iberian Coasts. *Cahiers de Biologie Marine* 49: 37-57.
- Aguirrezabalaga F, Gil J (2009) Paraonidae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic) with the description of eight new species. *Scientia Marina* 73 (4): 631-666. <https://doi.org/10.3989/scimar.2009.73n4631>
- Ahrens JB, Borda E, Barroso R, Paiva PC, Campbell AM, Wolf A, Nugues MM, Rouse G, Schulze A (2013) The curious case of *Hermodice carunculata* (Annelida:

- Amphinomidae): evidence for genetic homogeneity throughout the Atlantic Ocean and adjacent basins. *Molecular Ecology* 22: 2280-2291. <https://doi.org/10.1111/mec.12263>
- Åkesson B (1984) Speciation in the genus *Ophryotrocha* (Polychaeta, Dorvilleidae). *Fortschritte der Zoologie* 29: 299-314.
  - Åkesson B, Paxton H (2005) Biogeography and incipient speciation in *Ophryotrocha labronica* (Polychaeta, Dorvilleidae). *Marine Biology Research* 1 (2): 127-139. <https://doi.org/10.1080/17451000510019088>
  - Ακουμιανάκι Ι (2004) Μακροβενθικές βιοκοινωνίες υποθαλάσσιων δελταϊκών αποθέσεων: παράδειγμα από έναν ημίκλειστο κόλπο της Ανατολικής Μεσογείου (Μαλιακός κόλπος, Αιγαίο Πέλαγος). [Macrobenthic communities of subaqueous deltaic depositions: a case study from a semi-enclosed bay in the Eastern Mediterranean (Maliakos Bay, Aegean Sea)]. PhD thesis. University of Crete, Heraklion, 401 pp. [In Greek]. <https://doi.org/10.12681/eadd/16001>
  - Ακουμιανάκι Ι, Hughes AJ (1997) The macrofauna of the shallow hydrothermal vents at Paleohori Bay, Milos. 5th Panhellenic Symposium of Oceanography and Fisheries, Kavala, 5-18 April 1997. 241–244 pp. URL: <http://symposia.ath.hcmr.gr/oldver/symposia5/241.pdf>
  - Alós C (1984) Anélidos Poliquetos del Cabo de Creus. I. Rizomas de *Posidonia oceanica*. *Cuadernos Marisqueiros (Actas del III Simposio Ibérico de Estudios del Bentos Marino)* 7: 89-102.
  - Alós C (1989) Adiciones a la fauna de anélidos poliquetos de la península Ibérica: familia Syllidae. *Cahiers de Biologie Marine* 30 (3): 329-339. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
  - Alós C (1990) Anélidos poliquetos del Cabo de Creus (ne de España). Facies de *Corallina elongata* Ellis & Solander y de *Cystoseira mediterranea* (J. Feldmann). *Miscellània Zoològica* 14: 17-28. URL: <http://www.raco.cat/index.php/Mzoologica/article/viewFile/90563/169042>
  - Alós C (2004) Familia Phyllodocidae. In: Viéitez J, Alós C, Parapar J, Besteiro C, Moreira J, Núñez J, Laborda A, San Martín J (Eds) Annelida, Polychaeta I. In: Ramos, M.A., et al. (Eds.), Fauna Ibérica. 25. Museo Nacional de Ciencias Naturales, CA, SAIC, Madrid, 105–209 pp.
  - Álvarez-Campos P, Giribet G, Riesgo A (2016) The *Syllis gracilis* species complex: a molecular approach to a difficult taxonomic problem (Annelida, Syllidae). *Molecular Phylogenetics and Evolution* 109: 138-150. <https://doi.org/10.1016/j.ympev.2016.12.036>
  - Álvarez-Campos P, Giribet G, San Martín G, Rouse G, Riesgo A (2017) Straightening the striped chaos: systematics and evolution of *Trypanosyllis* and the case of its pseudocryptic type species *Trypanosyllis krohnii* (Annelida, Syllidae). *Zoological Journal of the Linnean Society* 179 (3): 492-540. <https://doi.org/10.1111/zoj.12443>
  - Annenkova N (1929) Beiträge zur Kenntnis der Polychaeten-Fauna der USSR. I. Fam. Pectinariidae Quatrefages (Amphictenidae Malmgren) und Ampharetidae Malmgren. *Annuaire du Musée Zoologique de l'Académie des Sciences de l'URSS* 30 (3): 477-502.
  - Antoniadou C (1998) Macro- and megafauna from the North Aegean Sea from 1997-1998. Aristotle University of Thessaloniki, Department of Biology, Laboratory of Zoology, Greece. Release date: 2008-5-14. URL: [http://ipt.vliz.be/eurobis/resource?r=largenet\\_q1](http://ipt.vliz.be/eurobis/resource?r=largenet_q1)
  - Antoniadou C (2003) Δομή των συνευρέσεων του σκληρού υποστρώματος της κατώτερης υποπαραλιακής ζώνης στο βόρειο Αιγαίο. [Structure of hard substrate

- benthic assemblages at the lower infralittoral zone in the north Aegean]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 457 pp. [In Greek]. <https://doi.org/10.12681/eadd/14763>
- Antoniadou C, Chintiroglou C (2005) Response of polychaete populations to disturbance: an evaluation of methods in hard substratum. *Fresenius Environmental Bulletin* 14: 1066-1073. URL: <http://www.vliz.be/imisdocs/publications/137228.pdf>
  - Antoniadou C, Nicolaidou A, Chintiroglou C (2004) Polychaetes associated with the sciaphilic alga community in the northern Aegean Sea: spatial and temporal variability. *Helgoland Marine Research* 58 (3): 168-182. <https://doi.org/10.1007/s10152-004-0182-6>
  - Antoniadou C, Voultziadou E, Chintiroglou C (2006) Sublittoral megabenthos along cliffs of different profile (Aegean Sea, Eastern Mediterranean). *Belgian Journal of Zoology* 136: 69-79. URL: <http://users.auth.gr/elvoults/pdf/Sublittoral%20megabenthos%2006.pdf>
  - Argyrou M, Demetropoulos A, Hadjichristophorou M (1999) Expansion of the macroalga *Caulerpa racemosa* and changes in softbottom macrofaunal assemblages in Moni Bay, Cyprus. *Oceanologica Acta* 22 (5): 517-528. [https://doi.org/10.1016/s0399-1784\(00\)87684-7](https://doi.org/10.1016/s0399-1784(00)87684-7)
  - Arias A, Paxton H (2013) First record of the polychaetous annelid *Diopatra micrura* Pires et al., 2010 in the Mediterranean Sea. *Mediterranean Marine Science* 15 (1): 5. <https://doi.org/10.12681/mms.377>
  - Arias A, Paxton H (2014) Hidden diversity within the polychaete *Onuphis eremita* sensu lato (Annelida: Onuphidae)—redescription of *O. eremita* Audouin & Milne-Edwards, 1833 and reinstatement of *Onuphis pancerii* Claparède, 1868. *Zootaxa* 3861 (2): 145-149. <https://doi.org/10.11646/zootaxa.3861.2.3>
  - Arias A, Paxton H (2015a) *Onuphis* and *Aponuphis* (Annelida: Onuphidae) from southwestern Europe, with the description of a new species. *Zootaxa* 3949 (3): 345-369. <https://doi.org/10.15468/C2FAWC>
  - Arias A, Paxton H (2015b) *Paradiopatra* Ehlers, 1887 (Annelida: Onuphidae) from southwestern Europe with the description of a new species and new ultramorphological data for the genus. *Zootaxa* 4040 (2): 149-168. <https://doi.org/10.15468/MCX6WK>
  - Arias A, Barroso R, Anadón N, Paiva PC (2013) On the occurrence of the fireworm *Eurythoe complanata* complex (Annelida, Amphinomidae) in the Mediterranean Sea with an updated revision of the alien Mediterranean amphinomids. *ZooKeys* 33 (337): 19-33. <https://doi.org/10.3897/zookeys.337.5811>
  - Arias A, Fernandez-Alvarez FA, Martins R, Anadon N (2015) Rediscovery and redescription of *Leodice laurillardii* (Quatrefages, 1866) comb. nov. (Annelida: Eunicidae)—a rare European polychaete or just an overlooked species? *Zootaxa* 3964 (4): 475. <https://doi.org/10.11646/zootaxa.3964.4.6>
  - Arias A, Giangrande A, Gambi MC, Anadón N (2013) Biology and new records of the invasive species *Branchiomma bairdi* (Annelida: Sabellidae) in the Mediterranean Sea. *Mediterranean Marine Science* 14 (1): 162-171. <https://doi.org/10.12681/mms.363>
  - Arvanitidis C (1994) Συστηματική και βιολογική μελέτη των μακροβενθικών πολυχαίτων (Annelida Polychaeta) του Βορείου Αιγαίου. [Systematic and bionomic study of the macrobenthic polychaetes (Annelida, Polychaeta) of the North Aegean]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 512 pp. [In Greek]. <https://doi.org/10.12681/eadd/5045>



- Arvanitidis C (2000a) Polychaete fauna of the Aegean Sea: inventory and new information. *Bulletin of Marine Science* 66 (1): 73-96. URL: <http://www.ingentaconnect.com/contentone/umrsmas/bullmar/2000/00000066/00000001/art00010>
- Arvanitidis C (2000b) Aegean polychaetes dataset. Hellenic Centre for Marine Research, Greece. Release date: 2005-9-27. URL: <http://ipt.vliz.be/eurobis/resource?r=aegeanpolychaetes>
- Arvanitidis C, Koukouras A (1997) The genus *Paradiopatra* (Polychaeta, Onuphidae) in the Mediterranean with the description of *Paradiopatra calliopae* sp. nov. *Ophelia* 46 (1): 51-63. <https://doi.org/10.1080/00785326.1997.10432477>
- Ayari R, Muir A, Patterson G, Afli A, Aissa P (2009) An updated list of polychaetous annelids from Tunisian coasts (Western Mediterranean Sea). *Cahiers de Biologie Marine* 50 (1): 33-45. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Bailey J (1969) Spirorbinae (Polychaeta: Serpulidae) from Chios (Aegean Sea). *Zoological Journal of the Linnean Society* 48 (3): 363-385. <https://doi.org/10.1111/j.1096-3642.1969.tb00718.x>
- Bailly N, Gerovasileiou V, Arvanitidis C, Legakis A (2016) Introduction to the Greek Taxon Information System (GTIS) in LifeWatchGreece: the construction of the Preliminary Checklists of Species of Greece. *Biodiversity Data Journal* 4: e7959. <https://doi.org/10.3897/bdj.4.e7959>
- Barnich R, Fiege D (2000a) Revision of the Mediterranean species of *Harmothoe* Kinberg, 1856 and *Lagisca* Malmgren, 1865 (Polychaeta: Polynoidae: Polynoinae) with descriptions of a new genus and a new species. *Journal of Natural History* 34: 1889-1938. <https://doi.org/10.1080/00222930050144783>
- Barnich R, Fiege D (2000b) Review of the North East Atlantic and Mediterranean species of *Aphrodita* Linnaeus, 1758 and *Aphroditella* Roule, 1898 (Polychaeta: Aphroditidae). *Ophelia* 53 (2): 131-140. <https://doi.org/10.1080/00785236.2000.10409443>
- Barnich R, Fiege D (2001) The Mediterranean species of *Malmgreniella* Hartman, 1967 (Polychaeta: Polynoidae: Polynoinae), including the description of a new species. *Journal of Natural History* 35 (8): 1119-1142. <https://doi.org/10.1080/00222930152434463>
- Barnich R, Fiege D (2003) The Aphroditoidea (Annelida: Polychaeta) of the Mediterranean Sea. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 599: 1-167.
- Barnich R, Fiege D (2009) Revision of the genus *Harmothoe* Kinberg, 1865 (Polychaeta: Polynoidae) in the Northeast Atlantic. *Zootaxa* 2104: 1-76.
- Barnich R, Fiege D (2010) On the distinction of *Harmothoe globifera* (G.O. Sars, 1873) and some other easily confused polynoids in the NE Atlantic, with the description of a new species of *Acanthicolepis* Norman in McIntosh, 1900 (Polychaeta, Polynoidae). *Zootaxa* 2525: 1-18.
- Barroso R, Paiva PC (2011) A new deep-sea species of *Chloeia* (Polychaeta: Amphinomidae) from southern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 91 (02): 419-423. <https://doi.org/10.1017/s0025315410001499>
- Barroso R, Klautau M, Solé-Cava A, Paiva PCd (2010) *Eurythoe complanata* (Polychaeta: Amphinomidae), the 'cosmopolitan' fireworm, consists of at least three

- cryptic species. *Marine Biology* 157 (1): 69-80. <https://doi.org/10.1007/s00227-009-1296-9>
- Bastida-Zavala JR, ten Hove HA (2003) Revision of *Hydroides* Gunnerus, 1768 (Polychaeta: Serpulidae) from the eastern Pacific region and Hawaii. *Beaufortia* 53: 67-110.
  - Bellan G (1959) Campagne de la Calypso: mer d'Alboran. 2. Annélides Polychètes. *Annales de l'Institut Océanographique Paris* 37: 315-342.
  - Bellan G (1963) Nouvelle contribution à l'étude de la microfaune annélide de la région de Marseille. *Recueil des Travaux de la Station Marine d'Endoume. Bulletin* 29 (44): 43-57.
  - Bellan G (1964a) Contribution à l'étude systématique, bionomique et écologique des Annélides Polychètes de la Méditerranée. [Contribution to the systematic, bionomic and ecological study of the Annelida Polychaeta of the Mediterranean]. PhD thesis. Université d'Aix-Marseille, Marseille, 371 pp. [In French].
  - Bellan G (1964b) Campagne de la Calypso: Méditerranée Nord-Orientale. 6. Annélides Polychètes. *Annales de l'Institut Océanographique Paris* 41: 271-288.
  - Ben-Eliahu MN (1972a) Polychaeta Errantia of the Suez Canal. *Israel Journal of Zoology* 21: 189-237. URL: <http://www.tandfonline.com/doi/abs/10.1080/00212210.1972.10688363>
  - Ben-Eliahu MN (1972b) Littoral Polychaeta from Cyprus. *Tethys* 4: 85-94.
  - Ben-Eliahu MN (1976a) Errant polychaete cryptofauna (excluding Syllidae and Nereidae) from rims of similar intertidal vermetid reefs on the Mediterranean coast of Israel and in the Gulf of Elat. *Israel Journal of Zoology* 25: 156-177. URL: <http://www.tandfonline.com/doi/abs/10.1080/00212210.1976.10688435>
  - Ben-Eliahu MN (1976b) Polychaete cryptofauna from rims of similar intertidal vermetid reefs on the Mediterranean coast of Israel and in the Gulf of Elat: Sedentaria. *Israel Journal of Zoology* 25: 121-155. <https://doi.org/10.1080/00212210.1975.10688411>
  - Ben-Eliahu MN (1989) Lessepsian migration in Nereididae (Annelida: Polychaeta): some case histories. In: Spanier E, Steinberger Y, Luria M (Eds) *Environmental Quality and Ecosystem Stability. Vol. IV/B-Ecosystem Stability. Proc. 4th Int. Conference Israel Society of Ecology and Environmental Quality Sciences. June 4-8 1989. Jerusalem, 125-134 pp.*
  - Ben-Eliahu MN (1991a) Red Sea serpulids (Polychaeta) in the Eastern Mediterranean. *Ophelia Supplement* 5: 515-528.
  - Ben-Eliahu MN (1991b) Nereididae of the Suez Canal – potential Lessepsian migrants? *Bulletin of Marine Science* 48: 318-329.
  - Ben-Eliahu MN (1995) A list of Polychaeta from along the Levant coast. *Haasiana* 1: 78-93. URL: <http://www.vliz.be/imisdocs/publications/272909.pdf>
  - Ben-Eliahu MN, Fiege D (1995) Polychaeta from the continental shelf and slope of Israel collected by the 'Meteor' 5 expedition (1987). *Senckenbergiana Maritima* 25 (4-6): 85-105.
  - Ben-Eliahu MN, Fiege D (1996) Serpulid tube-worms (Annelida: Polychaeta) of the Central and Eastern Mediterranean with particular attention to the Levant Basin. *Senckenbergiana Maritima* 28: 1-51. <https://doi.org/10.1007/BF03042821>
  - Ben-Eliahu MN, ten Hove H (2011) Serpulidae (Annelida: Polychaeta) from the Suez Canal- From a Lessepsian migration perspective. *Zootaxa* 2848: 1-147. URL: <http://dare.uva.nl/document/2/107139>

- Bertasi F (2016) The occurrence of the alien species *Polydora cornuta* Bosc, 1802 (Polychaeta: Spionidae) in North Adriatic lagoons: an overlooked presence. Italian Journal of Zoology 83 (1): 77-88. <https://doi.org/10.1080/11250003.2016.1140839>
- Bertasi F, Lomiri S, Vani D, Trabucco B, Lamberti CV (2014) First record of genus *Gallardoneris* (Polychaeta: Lumbrineridae) in Mediterranean marine waters. Marine Biodiversity Records 7: e63. <https://doi.org/10.1017/s1755267214000670>
- Bhaud M (1998) Species of *Spiochaetopterus* (Polychaeta, Chaetopteridae) in the Atlantic-Mediterranean biogeographic area. Sarsia 83: 243-263. [https://doi.org/10.1016/S0378-1127\(97\)00316-2](https://doi.org/10.1016/S0378-1127(97)00316-2)
- Bhaud M, Martín D, Gil J (2003) *Spiochaetopterus creocanae*, a new species of Chaetopteridae (Polychaeta) from the Persian Gulf belonging to the *costarum* complex. Scientia Marina 67 (1): 99-105. <https://doi.org/10.3989/scimar.2003.67n199>
- Bhaud MM (1975) Nouvelles données sur les larves de Sabellariidae récoltées en Méditerranée. Annales de l'Institut Océanographique Paris 51 (2): 155-172.
- Bhaud MM, Lenaers G (1992) Molecular Phylogeny of Some Polychaete Annelids: An Initial Approach to the Atlantic-Mediterranean Speciation Problem. Journal of Molecular Evolution 35: 429-435. <https://doi.org/10.1007/BF00171821>
- Bianchi CN (1981) Policheti Serpuloidei. Guide per il riconoscimento delle specie animali delle acque lagunari e costiere italiane 5: 1-187.
- Bianchi CN, Morri C (2000a) Marine Biodiversity of the Mediterranean Sea: Situation, Problems and Prospects for Future Research. Marine Pollution Bulletin 40 (5): 367-376. [https://doi.org/10.1016/s0025-326x\(00\)00027-8](https://doi.org/10.1016/s0025-326x(00)00027-8)
- Bianchi CN, Morri C (2000b) Serpuloidea (Annelida: Polychaeta) from Milos, an island in the Aegean Sea with submarine hydrothermalism. Journal of the Marine Biological Association of the United Kingdom 80: 259-269. <https://doi.org/10.1017/S0025315400001831>
- Bick A (2002) The morphology and ecology of *Dipolydora armata* (Polychaeta, Spionidae) from the western Mediterranean Sea. Acta Zoologica 82 (3): 177-187. <https://doi.org/10.1046/j.1463-6395.2001.00078.x>
- Blainville Hd (1825) Dictionnaire des Sciences naturelles, dans lequel on traite méthodiquement des différens êtres de la nature, considérés soit en eux-mêmes, d'après l'état actuel de nos connaissances, soit relativement à l'utilité qu'en peuvent retirer la médecine, l'agriculture, le commerce et les arts. 34. FG Levrault, Strasbourg and Paris, 560 pp. [In French]. URL: <http://www.biodiversitylibrary.org/page/25303394>
- Blake J (1991) Revision of some genera and species of Cirratulidae (Polychaeta) from the western North Atlantic. Ophelia Supplement, 5: 17-30.
- Blake J (1996) Family Cirratulidae Ryckholdt, 1851. In: Blake J, Hilbig B, Scott P (Eds) Taxonomic Atlas of the Benthic Fauna of the Santa Maria Basin and Western Santa Barbara Channel. The Annelida Part 3 - Polychaeta: Orbiniidae to Cossuridae . Vol. 6. Santa Barbara Museum of Natural History, Santa Barbara, 263–284 pp. URL: [https://invertebrates.si.edu/boem/reports/CAMP\\_Taxa6.pdf](https://invertebrates.si.edu/boem/reports/CAMP_Taxa6.pdf)
- Blake J (2000) Family Capitellidae Grube, 1862. In: Blake JA, Hilbig B, Scott PH (Eds) Taxonomic atlas of the benthic fauna of the Santa Maria basin and Western Santa Barbara Channel. The Annelida Part 4 – Polychaeta: Flabelligeridae to Ampharetidae . Vol. 7. Santa Barbara Museum of Natural History, Santa Barbara, 47–96 pp. URL: [https://invertebrates.si.edu/boem/reports/CAMP\\_Taxa7.pdf](https://invertebrates.si.edu/boem/reports/CAMP_Taxa7.pdf)

- Blake J (2009) Redescription of *Capitella capitata* (Fabricius) from West Greenland and designation of a neotype (Polychaeta, Capitellidae). *Zoosymposia* 2: 55-80. <https://doi.org/10.11646/zoosymposia.2.1.7>
- Blake J (2016) *Kirkegaardia* (Polychaeta, Cirratulidae), new name for *Monticellina* Laubier, preoccupied in the Rhabdocoela, together with new records and descriptions of eight previously known and sixteen new species from the Atlantic, Pacific, and Southern Oceans. *Zootaxa* 4166 (1): 1-93. <https://doi.org/10.11646/zootaxa.4166.1.1>
- Blake J (2017) Polychaeta Orbiniidae from Antarctica, the Southern Ocean, the Abyssal Pacific Ocean, and off South America. *Zootaxa* 4812: 1-145. <https://doi.org/10.11646/zootaxa.4218.1.1>
- Blake J, Grassle JP, Eckelbarger KJ (2009) *Capitella teleta*, a new species designation for the opportunistic and experimental *Capitella* sp. I, with a review of the literature for confirmed records. *Zoosymposia* 2: 25-53. <https://doi.org/10.11646/zoosymposia.2.1.6>
- Blake JA (1983) Polychaetes of the family Spionidae from South America, Antarctica, and adjacent seas and islands. *Antarctic Research Series* 39 (3): 205 – 288-205 – 288.
- Bleidorn C, Hill N, Erséus C, Tiedemann R (2009) On the role of character loss in orbiniid phylogeny (Annelida): Molecules vs. morphology. *Molecular Phylogenetics and Evolution* 52 (1): 57-69. <https://doi.org/10.1016/j.ympev.2009.03.022>
- Bleidorn C, Kruse I, Albrecht S, Bartolomaeus T (2006) Mitochondrial sequence data expose the putative cosmopolitan polychaete *Scoloplos armiger* (Annelida, Orbiniidae) as a species complex. *BMC Evolutionary Biology* 6 <https://doi.org/10.1186/1471-2148-6-47>
- Bock G, Fiege D, Barnich R (2010) Revision of *Hermadion* Kinberg, 1856, with a redescription of *Hermadion magalhaensi* Kinberg, 1856, *Adyte hyalina* (G.O.Sars, 1873) n. comb. and *Neopolynoe acanellae* (Verrill, 1881) n. comb. (Polychaeta:Polynoidae). *Zootaxa* 2554: 45-61. <https://doi.org/10.3852/10-176>
- Bogdanos C, Diapoulis A (1984) Contribution to the study of the zoobenthos and phytobenthos in Messolonghi lagoon. *Thalassographica* 7: 39-52.
- Bogdanos C, Fredj G (1983) Sur la présence de *Cossura coasta* Kitamori, 1960 (Polychaeta, Cossuridae) dans les eaux côtières Grèques (Mer Égée, Mer Ionienne). *Thalassographica* 6: 5-15. URL: [https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/239/vol6\(1\)5\\_15.pdf](https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/239/vol6(1)5_15.pdf)
- Bogdanos C, Nicolaidou A (1985) The offshore benthic fauna of the Patraikos Gulf (Greece). *Biologia Gallo-Hellenica* 10: 31-41.
- Bogdanos C, Satsmadjis J (1983) The macrozoobenthos of an Aegean Embayment. *Thalassographica* 6: 77-105. URL: [https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/249/vol6%20%2077\\_105.pdf](https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/249/vol6%20%2077_105.pdf)
- Bogdanos C, Satsmadjis J (1987) The Patraikos Gulf bottom fauna. *Thalassographica* 10: 37-71. URL: [https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/271/vol10\(1\)%2037\\_71.pdf](https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/271/vol10(1)%2037_71.pdf)
- Bogdanos C, Simboura N, Zenetos A (2002) The benthic fauna of Geras gulf (Lesvos Isl., Greece): inventory, distribution and some zoogeographical considerations. *Hellenic Zoological Archives* 6: 1-22.
- Böggemann M (2002) Revision of the Glyceridae Grube 1850 (Annelida: Polychaeta). *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 555: 1-249.
- Böggemann M (2005) Revision of the Goniadidae (Annelida, Polychaeta). *Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 39: 1-354.

- Böggemann M (2009) Polychaetes (Annelida) of the abyssal SE Atlantic. *Organisms Diversity & Evolution* 9: 251-428. <https://doi.org/10.1016/j.ode.2009.10.001>
- Böggemann M (2015) Glyceriformia (Annelida) of the abyssal SW Atlantic and additional material from the SE Atlantic. *Marine Biodiversity* 46 (1): 227-241. <https://doi.org/10.1007/s12526-015-0354-4>
- Böggemann M, Westheide W (2004) Interstitial Syllidae (Annelida: Polychaeta) from Mahé (Seychelles). *Journal of Natural History* 38 (4): 403-446. <https://doi.org/10.1080/0022293021000033229>
- Bonse S, Schmidt H, Eibye-Jacobsen D, Westheide W (1996) *Eulalia viridis* (Polychaeta: Phyllodocidae) is a complex of two species in northern Europe: Results from biochemical and morphological analyses. *Cahiers de Biologie Marine* 37 (1): 33-48. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Britton-Davidian J, Amoureux L (1982) Biochemical systematics of two sibling species of polychaete annelids: *Ophelia bicornis* and *O. radiata*. *Biochemical Systematics and Ecology* 10 (4): 351-354. [https://doi.org/10.1016/0305-1978\(82\)90009-6](https://doi.org/10.1016/0305-1978(82)90009-6)
- Brullé M (1832) Annelides. In: Brullé M (Ed.) *Expéditions scientifiques de Morée*. 3. Levrault, Paris, 396-400 pp. URL: [https://archive.org/stream/bub\\_gb\\_3TD5Tzo5T4gC#page/n401/](https://archive.org/stream/bub_gb_3TD5Tzo5T4gC#page/n401/)
- Budaeva N, Fauchald K (2011) Phylogeny of the *Diopatra* generic complex with a revision of *Paradiopatra* Ehlers, 1887 (Polychaeta: Onuphidae). *Zoological Journal of the Linnean Society* 163 (2): 319-436. <https://doi.org/10.1111/j.1096-3642.2011.00701.x>
- Budaeva N, Paxton H (2013) *Nothria* and *Anchinothria* (Annelida: Onuphidae) from eastern Australian waters, with a discussion of ontogenetic variation of diagnostic characters. *Journal of the Marine Biological Association of the United Kingdom* 93 (6): 1481-1502. <https://doi.org/10.1017/s0025315412001956>
- Bush KJ (1905) Tubicolous annelids of the tribes Sabellides and Serpulides from the Pacific Ocean. *Harriman Alaska Expedition* 12: 169-346. URL: <http://biodiversitylibrary.org/page/21676213>
- Busoni G (2013) Indagine sulla polichetofauna di ambienti marini profondi del Mediterraneo Occidentale: considerazioni tassonomiche, zoogeografiche ed ecologiche. [Survey of the polychaete fauna of the deep marine environment in the Western Mediterranean: taxonomic, zoogeographic and ecological considerations]. MSc thesis. Università degli Studi di Pisa, Pisa, 53 pp. [In Italian]. URL: [https://etd.adm.unipi.it/theses/available/etd-11202013-083528/unrestricted/TESE\\_Giulio\\_Busoni.pdf](https://etd.adm.unipi.it/theses/available/etd-11202013-083528/unrestricted/TESE_Giulio_Busoni.pdf)
- Campoy A (1982) Fauna de España. Fauna de Anélidos Poliquetos de la Península Ibérica. [Fauna of Spain: Fauna of the Annelida Polychaeta of the Iberian Peninsula]. 7 (1). Servicio de Publicaciones de la Universidad de Navarra, Pamplona, 781 pp. [In Spanish]. URL: <http://hdl.handle.net/10171/11777>
- Cantone G, Fassari G (1982) Policheti dei fondi mobili del Golfo di Catania. *Animalia* 9: 226-237.
- Cantone G, Fassari G, Brigandi S (1978) Ricerche sui policheti della Tunisia. *Animalia* 5 (1-3): 51-78.
- Capaccioni-Azzati R, El-Haddad M (2015) Familia Capitellidae. In: Ramos MA (Ed.) *Annelida. Polychaeta IV. Fauna Iberica*, 41. Museo Nacional de Ciencias Naturales. CSIC, Madrid, 416 pp. [In Spanish]. [ISBN 978-84-00-10006-3].

- Capaccioni-Azzati R, Torres-Gavila FJGAM (1989) On the presence of *Microphthalmus pseudoaberrans* Campoy and Vieitez 1982 (Polychaeta: Hesionidae) and *Magelona equilamellae* Harmelin 1964 (Polychaeta: Magelonidae) on the East Coast of Spain. *Boletín de la Real Sociedad Española de Historia Natural* 84 (3-4): 235-246.
- Capaccioni-Azzati R, Torres-Gavila FJ, Tena J (1992) Dos nuevos géneros de anélidos poliuetos para la península ibérica: *Neopseudocapitella* Rullier and Amoureux, 1979 y *Demonax* Kinberg, 1867. *Thalassas* 10: 115-12.
- Carr CM, Hardy SM, Brown TM, Macdonald TA, Hebert PDN (2011) A tri-oceanic perspective: DNA barcoding reveals geographic structure and cryptic diversity in Canadian polychaetes. *PLoS ONE* 6 (7): . <https://doi.org/10.1371/journal.pone.0022232>
- Carrera-Parra LF (2004) Revision of *Lumbricalus* (Polychaeta: Lumbrineridae). *Journal of the Marine Biological Association of the UK* 84 (1): 81-91. <https://doi.org/10.1017/s002531540400894xh>
- Carrera-Parra LF (2006) Revision of *Lumbrineris* de Blainville, 1828 (Polychaeta: Lumbrineridae). *Zootaxa* 1336: 1-64.
- Carrera-Parra LF, Salazar-Vallejo SI (1998) A new genus and 12 new species of Eunicidae (Polychaeta) from the Caribbean Sea. *Journal of the Marine Biological Association of the United Kingdom* 78: 145-182. <https://doi.org/10.1017/S0025315400040005>
- Carrera-Parra LF, Çınar ME, Dağlı E (2011) Description of a new species of *Lumbrineris* (Polychaeta: Lumbrineridae) from the coasts of Turkey (eastern Mediterranean). *Marine Biodiversity* 41: 343-347. <https://doi.org/10.1007/s12526-010-0074-8>
- Casellato S, Stefanon A (2008) Coralligenous habitat in the northern Adriatic Sea: an overview. *Marine Ecology* 29 (3): 321-341. <https://doi.org/10.1111/j.1439-0485.2008.00236.x>
- Castelli A (1987) Censimento dei policheti dei mari italiani: Paraonidae Cerruti, 1909. *Atti della Società Toscana di Scienze Naturali Serie B* 94: 319-340. URL: <http://www.stsn.it/AttiB1987/castelli.pdf>
- Castelli A (1990) Censimento dei policheti dei Mari Italiani: Flabelligeridae Saint-Joseph, 1894. *Atti della Società Toscana di Scienze Naturali, Memorie, serie B* 96: 9-22. URL: <http://www.stsn.it/AttiB1989/castelli.pdf>
- Castelli A, Bianchi N, Cantone G, Çınar M, Gambi M, Giagrande A, Sareri D, Lanera P, Licciano M, Musco L, Sanfilippo R, Simonini R (2008) Annelida Polychaeta . *Biologia Marina Mediterranea* 15: 323-373. URL: [http://www.sibm.it/checklist/bmm%2015\(s1\)%202008%20checklist%20i/40%20annelida%20polychaeta.pdf](http://www.sibm.it/checklist/bmm%2015(s1)%202008%20checklist%20i/40%20annelida%20polychaeta.pdf)
- Caullery M, Mesnil F (1897) Études sur la morphologie compare et la phylogenie des espèces chez les Spirorbes. *Bulletin scientifique de la France et de la Belgique* 30: 185-233. URL: <http://biodiversitylibrary.org/page/10721821>
- Chambers S, Lanera P, Mikac B (2011) *Chaetozone carpenteri* McIntosh, 1911 from the Mediterranean Sea and records of other bi-tentaculate Cirratulids. *Italian Journal of Zoology* 78: 41-48. <https://doi.org/10.1080/11250003.2011.580565>
- Chambers SJ, Woodham A (2003) A new species of *Chaetozone* (Polychaeta: Cirratulidae) from deep water in the northeast Atlantic, with comments on the diversity of the genus in cold northern waters. *Hydrobiologia* 496: 41-48. [https://doi.org/10.1007/978-94-017-0655-1\\_5](https://doi.org/10.1007/978-94-017-0655-1_5)
- Chardy P, Laubier L, Reyss D, Sibuet M (1973) Dragages profonds en mer Ionienne-données préliminaires (1). *Rapport et Procès-Verbaux des Réunions de la Commission*

- Internationale pour l'Exploration Scientifique de la Mer Méditerranée 22: 103-105. URL: <http://archimer.ifremer.fr/doc/1973/publication-5132.pdf>
- Chatzigeorgiou G, Faulwetter S, Arvanitidis C (2016) Polychaetes from two subtidal rocky shores of the North coast of Crete, collected for the NaGISA project 2007-2008. V1.2. Hellenic Centre for Marine Research. Release date: 2016-4-07. URL: [http://ipt.medobis.eu/resource?r=nagisa\\_species\\_2007\\_2008](http://ipt.medobis.eu/resource?r=nagisa_species_2007_2008)
  - Chatzigeorgiou G, Keklikoglou K, Faulwetter S, Badalamenti F (2017) Midlittoral polychaete communities in the Eastern Mediterranean Sea: new information from the implementation of the Natural Geography in Shore Areas (NaGISA) protocol and comparisons at local and regional scales. *Marine Ecology* 38: e12339. <https://doi.org/10.1111/maec.12339>
  - Chatzigeorgiou G, Sarropoulou E, Vasileiadou K, Brown C, Faulwetter S, Kotoulas G, Arvanitidis C (2014) Community structure and population genetics of Eastern Mediterranean polychaetes. *Frontiers in Marine Science* 1: 1-8. <https://doi.org/10.3389/fmars.2014.00047>
  - Chintiroglou C (1987) Βιονομική μελέτη των θαλάσσιων ανεμώνων (Actinaria, Anthozoa) του Θερμαϊκού Κόλπου. [Bionomical study of the sea-anemones (Actinaria, Anthozoa) of the Thermaikos Gulf.]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 241 pp. [In Greek]. <https://doi.org/10.12681/eadd/0830>
  - Chintiroglou C, Koukouras A (1991) Observations on the feeding habits of *Calliactis parasitica* (Couch, 1842), Anthozoa, Cnidaria . *Oceanologica Acta* 14: 389-396. URL: <http://archimer.ifremer.fr/doc/00102/21375/18975.pdf>
  - Chintiroglou C, Antoniadou C, Damianidis P, Diapoulis A (2004) Contribution to the knowledge of the macrobenthic biodiversity of Vouliagmeni Lagoon (Attica, Greece). *Rapport et Procès-Verbaux des Réunions de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée* 37: 506. URL: [http://ciesm.org/online/archives/abstracts/pdf/37/CIESM\\_Congress\\_2004\\_Barcelona\\_article\\_0506.htm](http://ciesm.org/online/archives/abstracts/pdf/37/CIESM_Congress_2004_Barcelona_article_0506.htm)
  - Christomanos A (1963) Dredging results from the oceanographic cruise "Tethys II" during October 1962. *Marine Science* 1963 (1): 27-38.
  - Christomanos A, Giannitsis D (1962) Dredging results from the oceanographic cruise "Tethys" during August-September 1961. *Marine Scientific Pages* 1-7-11.
  - Çınar ME, Dagli E, Çaglar S, Albayrak S (2015) Polychaetes from the northern part of the Sea of Marmara with the description of a new species of *Polydora* (Annelida: Polychaeta: Spionidae). *Mediterranean Marine Science* 16 (3): 524. <https://doi.org/10.12681/mms.1226>
  - Çınar ME, Ergen Z, Dagli E, Petersen M (2005) Alien species of spionid polychaetes (*Streblospio gynobranchiata* and *Polydora cornuta*) in Izmir Bay, eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* 85 (4): 821-827. <https://doi.org/10.1017/s0025315405011768>
  - Çınar ME (1999) Türkiye'nin Ege Denizi sahillerinde dağılım gösteren Syllidae (Polychaeta-Annelida) türlerinin taksonomisi ve ekolojisi. [Taxonomy and Ecology of Syllidae (Polychaeta, Annelida) distributed along the Turkish Aegean Coast]. PhD thesis. Ege University, Izmir, 443 pp. [In Turkish]. URL: <http://ulusaltezmerkezi.com/turkiyenin-ege-denizi-sahillerinde-dagilim-gosteren-syllidae-polychaeta-annelido-turlerinin-taksonomisi-ve-ekolojisi/>

- Çınar ME (2005) Polychaetes from the coast of northern Cyprus (eastern Mediterranean Sea), with two new records for the Mediterranean Sea. *Cahiers de Biologie Marine* 46 (2): 143-159. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Çınar ME (2006) Serpulid species (Polychaeta: Serpulidae) from the Levantine coast of Turkey (eastern Mediterranean), with special emphasis on alien species. *Aquatic Invasions* 1 (4): 223-240. <https://doi.org/10.3391/ai.2006.1.4.6>
- Çınar ME (2009) Alien polychaete species (Annelida: Polychaeta) on the southern coast of Turkey (Levantine Sea, eastern Mediterranean), with 13 new records for the Mediterranean Sea. *Journal of Natural History* 43: 2283-2328. <https://doi.org/10.1080/00222930903094654>
- Çınar ME, Ergen Z (2002) Faunistic analysis of Syllidae (Annelida: Polychaeta) from the Aegean Sea. *Cahiers de Biologie Marine* 43 (2): 171-178. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Çınar ME, Ergen Z (2003) Eusyllinae and Syllinae (Annelida: Polychaeta) from northern Cyprus (Eastern Mediterranean Sea) with a checklist of species reported from the Levant Sea. *Bulletin of Marine Science* 72 (3): 769-793. URL: <http://www.ingentaconnect.com/content/umrsmas/bullmar/2003/00000072/00000003/art00010>
- Çınar ME, Ergen Z (2005) Lessepsian migrants expanding their distributional ranges; *Pseudonereis anomala* (Polychaeta: Nereididae) in Izmir Bay (Aegean Sea). *Journal of the Marine Biological Association of the UK* 85 (2): 313-321. <https://doi.org/10.1017/S0025315405011203h>
- Çınar ME, Ergen Z (2007) The presence of *Chaetozone corona* (Polychaeta: Cirratulidae) in the Mediterranean Sea: an alien or a native species? *Cahiers de Biologie Marine* 48 (4): 339-346. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Çınar ME, Dağlı E, Kurt Şahin G (2014) Check-List of Annelida from the Coasts of Turkey. *Turkish Journal of Zoology* 38: 1-31. <https://doi.org/10.3906/zoo-1405-72>
- Çınar ME, Ergen Z, Benli HA (2003) Autolytinae and Exogoninae (Polychaeta: Syllidae) from northern Cyprus (Eastern Mediterranean Sea) with a checklist of species reported from the Levant Sea. *Bulletin of Marine Science* 72 (3): 741-767. URL: <http://www.ingentaconnect.com/content/umrsmas/bullmar/2003/00000072/00000003/art00009>
- Çınar ME, Ergen Z, Dağlı E (2002) Occurrence of the lessepsian species *Leonnates persicus* (Polychaeta: Nereididae) in Izmir Bay, Aegean Sea. *Journal of the Marine Biological Association of the United Kingdom* 82 (5): 811-815. <https://doi.org/10.1017/S0025315402006173>
- Çınar ME, Bakir K, Öztürk B, Katağan T, Doğan A, Açık S, Kurt-Sahin G, Özcan T, Dağlı E, Bitlis-Bakir B, Koçak F, Kirkim F (2017) Macrobenthic fauna associated with the invasive alien species *Brachidontes pharaonis* (Mollusca: Bivalvia) in the Levantine Sea (Turkey). *Journal of the Marine Biological Association of the United Kingdom* 97: 613-628. <https://doi.org/10.1017/S0025315417000133>
- Claparède É (1864) Glanures Zootomiques parmi les Annélides de Port-Vendres (Pyrenées Orientales). *Mémoires de la Société de physique et d'histoire naturelle de Genève* 17 (2): 463-600. URL: <http://www.biodiversitylibrary.org/ia/glanureszootomiq00clapare#page/7/mode/1up>



- Clausade M (1969) Peuplement animal sessile des petits substrats solides récoltés dans trois biocoenoses des fonds détritiques des parages de Marseille. *Tethys* 1: 719-750. URL: [http://paleopolis.rediris.es/benthos/REF/som/T-pdf/1970\\_1-3-719.pdf](http://paleopolis.rediris.es/benthos/REF/som/T-pdf/1970_1-3-719.pdf)
- Cognetti G (1955) Ricerche sui Sillidi del Golfo di Napoli. IV. Osservazioni su specie criptiche e su nuove sotto- specie geografiche ed ecologiche. *Pubblicazioni della Stazione Zoologica di Napoli* 26: 1-11.
- Colbath GH (1989) A revision of *Arabella mutans* (Chamberlin, 1919) and related species (Polychaeta: Arbellidae). *Proceedings of the Biological Society of Washington* 102: 283-299. URL: [https://archive.org/details/cbarchive\\_133975\\_arevisionofarabellamutanschamb1882](https://archive.org/details/cbarchive_133975_arevisionofarabellamutanschamb1882)
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Rais Lasram FB, Aguzzi J, Ballesteros E, Bianchi CN, Corbera J, Dailianis T, Danovaro R, Estrada M, Froglija C, Galil B, Gasol J, Gertwagen R, Gil J, Guilhaumon F, Kesner-Reyes K, Kitsos M, Koukouras A, Lampadariou N, Laxamana E, la Cuadra CLd, Lotze H, Martin D, Mouillot D, Oro D, Raicevich S, Rius-Barile J, Saiz-Salinas JI, Vicente CS, Somot S, Templado J, Turon X, Vafidis D, Villanueva R, Voultsiadou E (2010) The Biodiversity of the Mediterranean Sea: Estimates, Patterns, and Threats. *PLoS ONE* 5 (8): e11842. <https://doi.org/10.1371/journal.pone.0011842>
- Conides A, Bogdanos C, Diapoulis A (1999) Seasonal ecological variations of phyto- and zoobenthic communities in the south of Nisyros Island, Greece. *Environmentalist* 19 (2): 109-127. <https://doi.org/10.1023/A:1006641000873>
- Corriero G, Gherardi M, Giangrande A, Longo C, Mercurio M, Musco L, Marzano CN (2004) Inventory and distribution of hard bottom fauna from the marine protected area of Porto Cesareo (Ionian Sea): Porifera and Polychaeta. *Italian Journal of Zoology* 71 (3): 237-245. <https://doi.org/10.1080/11250000409356578>
- Corsini-Foka M, Zenetos A, Crocetta F, Çınar ME, Koçak F, Golani D, Katsanevakis S, Tsiamis K, Cook E, Froglija C, Triandaphyllou M, Lakkis S, Kondylatos G, Tricarico E, Zuljevic A, Almeida M, Cardigos F, Çağlar S, Durucan F, Fernandes AD, Ferrario J, Haberle I, Louizidou P, Makris J, Marić M, Micu D, Mifsud C, Nall C, Kytinou E, Poursanidis D, Spigoli D, Stasolla G, Yapici S, Roy H (2015) Inventory of alien and cryptogenic species of the Dodecanese (Aegean Sea, Greece): collaboration through COST action training school. *Management of Biological Invasions* 6 (4): 351-366. <https://doi.org/10.3391/mbi.2015.6.4.04>
- Cosentino A (2011) Microhabitat selection in a local syllid assemblage with the first record of *Syllis hyllebergi* (Syllinae) in the central Mediterranean. *Italian Journal of Zoology* 78: 267-279. <https://doi.org/10.1080/11250003.2011.589171>
- Cosentino A, Giacobbe S (2010) The new potential invader *Linopherus canariensis* (Polychaeta: Amphinomidae) in a Mediterranean coastal lake: Colonization dynamics and morphological remarks. *Marine Pollution Bulletin* 62: 236-245. <https://doi.org/10.1016/j.marpolbul.2010.11.006>
- da Cunha Lana P, Gruet Y (1989) *Sabellaria wilsoni* sp. n. (Polychaeta, Sabellariidae) from the southeast coast of Brazil. *Zoologica Scripta* 18: 239-244. <https://doi.org/10.1111/j.1463-6409.1989.tb00449.x>
- Dağlı E (2008) Türkiye'nin Ege Denizi sublittoral bölgesinde dağılım gösteren Spionidae (Polychaeta-Annelida) türlerinin taksonomi ve ekolojisi. *Ege Üniversitesi, İzmir*.
- Dağlı E, Çınar ME (2009) Species of the subgenera *Aquilaspio* and *Prionospio* (Polychaeta: Spionidae: *Prionospio*) from the southern coast of Turkey (Levantine Sea,

- eastern Mediterranean), with description of a new species and two new reports for the Mediterranean fauna. *Zootaxa* 2275: 1-20.
- Dağlı E, Çınar ME (2008) Invasion of polluted soft substratum of Izmir Bay (Aegean Sea, eastern Mediterranean) by the spionid polychaete worm, *Pseudopolydora paucibranchiata* (Polychaeta: Spionidae). *Cahiers de Biologie Marine* 49 (1): 87-96. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
  - Dağlı E, Çınar ME (2011) Species of the subgenus *Minuspio* (Polychaeta: Spionidae: *Prionospio*) from the southern coast of Turkey (Levantine Sea, eastern Mediterranean) with the description of two new species. *Zootaxa* 3043: 35-53.
  - Dağlı E, Çınar ME, Ergen Z (2011) Spionidae (Annelida: Polychaeta) from the Aegean Sea (eastern Mediterranean). *Italian Journal of Zoology* 78: 49-64. <https://doi.org/10.1080/11250003.2011.567828>
  - Dales RP (1957) Pelagic polychaetes of the Pacific Ocean. *Bulletin of the Scripps Institution of Oceanography of the University of California* 7 (2): 99-168. URL: <http://escholarship.org/uc/item/51c8j5nj>
  - Dales RP, Peter G (1972) A synopsis of the Pelagic Polychaeta. *Journal of Natural History* 6 (1): 55-92. <https://doi.org/10.1080/00222937200770071>
  - D'Alessandro M, Castriota L, Consoli P, Romeo T, Andaloro F (2015) *Pseudonereis anomala* (Polychaeta, Nereididae) expands its range westward: first Italian record in Augusta and Siracusa harbours. *Marine Biodiversity* 46 (1): 129-133. <https://doi.org/10.1007/s12526-015-0334-8>
  - D'Alessandro M, Romeo T, Castriota L, Cosentino A, Perzia P, Martins R (2016) New records of Lumbrineridae (Annelida: Polychaeta) in the Mediterranean biogeographic province, with an updated taxonomic key. *Italian Journal of Zoology* 83 (2): 233-243. <https://doi.org/10.1080/11250003.2016.1154615>
  - Dando PR, Hughes JA, Thiermann F (1995) Preliminary observations on biological communities at shallow hydrothermal vents in the Aegean Sea. *Geological Society, London, Special Publications* 87 (1): 303-317. <https://doi.org/10.1144/GSL.SP.1995.087.01.23>
  - Dando PR, Stüben D, Varnavas SP (1999) Hydrothermalism in the Mediterranean Sea. *Progress in Oceanography* 44: 333-367. [https://doi.org/10.1016/s0079-6611\(99\)00032-4](https://doi.org/10.1016/s0079-6611(99)00032-4)
  - Danovaro R, Company JB, Corinaldesi C, D'Onghia G, Galil B, Gambi C, Gooday A, Lampadariou N, Luna GM, Morigi C, Olu K, Polymenakou P, Ramirez-Llodra E, Sabbatini A, Sardà F, Sibuet M, Tselepidis A (2010) Deep-Sea Biodiversity in the Mediterranean Sea: The Known, the Unknown, and the Unknowable. *PLoS ONE* 5 (8): e11832. <https://doi.org/10.1371/journal.pone.0011832>
  - Dauvin J, Bachelet G, Bellan G (2006) Biodiversity and biogeographic relationships of the polychaete fauna in French Atlantic and Mediterranean waters. *Scientia Marina* 70: 259-267. <https://doi.org/10.3989/scimar.2006.70s3259>
  - Dauvin JC, Dewarumez JM, Gentil F (2003) Liste actualisée des espèces d'Annélides Polychètes présentes en Manche. *Cahiers de Biologie Marine* 44 (1): 67-95.
  - Day JH (1961) The Polychaete Fauna of South Africa. Part 6. Sedentary species dredged off Cape coasts with a few new records from the shore. *Journal of the Linnean Society of London* 44 (299): 463-560. <https://doi.org/10.1111/j.1096-3642.1961.tb01623.x>

- Day JH (1963) The Polychaete fauna of South Africa. Part 8: New species and records from grab samples and dredgings. *Bulletin of the British Museum (Natural History)* 10 (7): 384-445. <https://doi.org/10.5962/bhl.part.20530>
- Day JH (1967) A monograph on the Polychaeta of Southern Africa. Part 1. Errantia. Trustees of the British Museum (Natural History), London, 498 pp. URL: <http://www.biodiversitylibrary.org/item/35415#page/7/mode/1up>
- de Assis JE, Alonso C, Christoffersen ML (2007) A catalogue and taxonomic keys of the Subfamily Nicomachinae (Polychaeta: Maldanidae) of the world. *Zootaxa* 1657 (1657): 41-55.
- de León-González JA, Hernández Guevara NA, Rodríguez-Valencia JA (2006) Paraonidae (Polychaeta) from western Mexico, with description of two new species. *Journal of the Marine Biological Association of the UK* 86 (02): 253-262. <https://doi.org/10.1017/s0025315406013105>
- Delgado-Blas VH, Díaz-Díaz OF (2010) Description of two new species of *Malacoceros* and *Rhynchospio* spionids (Polychaeta: Spionidae) from the Grand Caribbean region. *Revista Chilena de Historia Natural* 83: 249-249.
- Delle Chiaje S (1825) Memorie sulla storia e notomia degli animali senza vertebre del Regno di Napoli. Stamperia delle Societa Tipografica, Napoli 2: 185-444. <https://doi.org/10.5962/bhl.title.10021>
- Delle Chiaje S (1841) Descrizione e notomia degli animali invertebrati della Sicilia citeriore osservati vivi negli anni 1822-1830. 5. Stamperia della Societa' Tipografica, Naples, 1–142 pp. URL: <https://biodiversitylibrary.org/page/10689723>
- de Oliveira Almeida W, Lindsey Christoffersen M, de Souza Amorim D, Senna Garraffoni AR, Sene Silva G (2003) Polychaeta, Annelida, and Articulata are not monophyletic: articulating the Metameria (Metazoa, Coelomata). *Revista Brasileira de Zoologia* 20 (1): 23-57. <https://doi.org/10.1590/s0101-81752003000100006>
- Desbryères D, Guille A, Ramos JM (1973) Bionomie benthique du plateau continental de la côte catalane Espagnole. *Vie et milieu* 23 (2): 335-363.
- Diapoulis A, Bogdanos C (1983) Preliminary study of soft substrate macrozoobenthos and marine flora in the Bay of Gera (Lesvos Island, Greece). *Thalassographica* 6: 127-139. URL: <http://hdl.handle.net/123456789/256>
- Di Domenico M, Martínez A, Lana P, Worsaae K (2014) Molecular and morphological phylogeny of Saccocirridae (Annelida) reveals two cosmopolitan clades with specific habitat preferences. *Molecular Phylogenetics and Evolution* 75: 202-218. <https://doi.org/10.1016/j.ympev.2014.02.003>
- Doğan A, Çınar ME, Önen M, Katağan T (2005) Seasonal dynamics of soft bottom zoobenthic communities in polluted and unpolluted areas of Izmir Bay (Aegean Sea). *Senckenbergiana Maritima* 35: 133-14. <https://doi.org/10.1007/BF03043182>
- D'Onghia G, Capezzuto F, Cardone F, Carlucci R, Carluccio A, Chimienti G, Corriero G, Longo C, Maiorano P, Mastrototaro F, Panetta P, Rosso A, Sanfilippo R, Sion L, Tursi A (2015) Macro- and megafauna recorded in the submarine Bari Canyon (southern Adriatic, Mediterranean Sea) using different tools. *Mediterranean Marine Science* 16 (1): 180-196. <https://doi.org/10.12681/mms.1082>
- Dorgham MM, Hamdy R, El-Rashidy HH, Atta MM (2013) First records of polychaetes new to Egyptian Mediterranean waters. *Oceanologia* 55 (1): 235-267. <https://doi.org/10.5697/oc.55-1.235>

- Dorgham MM, Hamdy R, El Rashidy HH, Atta MM, Musco L (2014) Distribution patterns of shallow water polychaetes (Annelida) along the Alexandria coast, Egypt (eastern Mediterranean). *Mediterranean Marine Science* 15 (3): 635-649. <https://doi.org/10.12681/mms.680>
- Dounas C (1988) Benthos Cretan Continental Shelf. Hellenic Centre for Marine Research, Greece.. Release date: 2005-9-27. URL: <http://ipt.vliz.be/eurobis/resource?r=benthoscretan>
- Dounas CG (1986) Study of the macrofaunal assemblages of Strymonikos Gulf. University of Thessaloniki, Thessaloniki, 185 pp. [In Greek].
- Ehlers E (1864) Die Borstenwürmer (Annelida Chaetopoda) nach systematischen und anatomischen Untersuchungen dargestellt. [Bristle worms (Annelida Chaetopoda) according to systematic and anatomical studies]. I. Wilhelm Engelmann Verlag, Leipzig, 748 pp. [In German]. <https://doi.org/10.5962/bhl.title.2081>
- Eleftheriou A, Smith C (2000) Preliminary Investigations of the Benthic Ecosystem from the Aegean shelf (Eastern Mediterranean). In: Della Croce N (Ed.) Symposium Mediterranean Seas 2000. ISAM, Santa Marherita Ligure. 105-120 pp.
- Eleftheriou A, Dounas K, Papadopoulou K, Karakassis I, Anastasiadis P, Koutsoubas D, Lazaridou T, Labropoulou M (1990) Study of the coasts around Crete presenting ecological disturbances. II. Marine Section Final Report. University of Crete, Heraklion, 92 pp. [In Greek].
- El Haddad M, Capaccioni-Azzati R, García-Carrascosa AM (2009) *Branchiomma luctuosum* (Polychaeta: Sabellidae): a non-indigenous species at Valencia Port (western Mediterranean Sea, Spain). *Marine Biodiversity Records* 1: 1-7. <https://doi.org/10.1017/S1755267207006604>
- Ergen Z (1976) Investigations on the taxonomy and ecology of Polychaeta from Izmir Bay and its adjacent area. *Scientific Reports of the Faculty of Science, Ege University* 209: 1-66.
- Ergen Z (1987) Biological diversity in Turkey. Ringworms (Annelida). Environmental Problem Foundation of Turkey. 150-152.
- Ergen Z, Çınar ME (1997) Polychaeta of Antalya Bay (Mediterranean coast of Turkey). *Israel Journal of Zoology* 43: 229-241. <https://doi.org/10.1080/00212210.1997.10688908>
- Fassari G (1982) Anellidi Policheti del mar Egeo. *Animalia* 9: 109-121.
- Fauchald K (1977) The Polychaete worms. Definitions and keys to the Orders, families and Genera. Science Series, Natural History Museum of Los Angeles County 28: 1 – 188-1 – 188.
- Fauchald K (1992) A Review of the Genus *Eunice* (Polychaeta: Eunicidae) Based upon Type Material. *Smithsonian Contributions to Zoology* 523: 1-422. <https://doi.org/10.5479/si.00810282.523>
- Fauchald K, Berke SK, Woodin SA (2012) *Diopatra* (Onuphidae: Polychaeta) from intertidal sediments in southwestern Europe. *Zootaxa* 3395: 47-58.
- Fauchald KA, Granados-Barba A, Solís-Weiss V (2009) Polychaeta (Annelida) of the Gulf of Mexico. In: Felder DL, Camp DK (Eds) *Gulf of Mexico—Origins, Waters, and Biota. Biodiversity*. Texas A&M University Press, College Station, Texas, 751– 788 pp.
- Faulwetter S (2010) Check-list of marine Polychaeta from Greece. Aristotle University of Thessaloniki. Assembled in the framework of the EU FP7 PESI project. <http://>

[www.marinespecies.org/aphia.php?p=sourcedetails&id=142069](http://www.marinespecies.org/aphia.php?p=sourcedetails&id=142069). Accessed on: 2016-11-21.

- Faulwetter S, Chatzigeorgiou G, Galil B, Arvanitidis C (2011a) An account of the taxonomy and distribution of Syllidae (Annelida, Polychaetes) in the eastern Mediterranean, with notes on the genus *Prosphaerosyllis* San Martín, 1984 in the Mediterranean. *ZooKeys* 150: 281-326. <https://doi.org/10.3897/zookeys.150.2146>
- Faulwetter S, Chatzigeorgiou G, Galil B, Nicolaidou A, Arvanitidis C (2011b) *Sphaerosyllis levantina* sp. n. (Annelida) from the eastern Mediterranean, with notes on character variation in *Sphaerosyllis hystrix* Claparède, 1863. *ZooKeys* 150: 327-345. <https://doi.org/10.3897/zookeys.150.1877>
- Faulwetter S, Vasileiadou A, Kouratoras M, Dailianis T, Arvanitidis C (2013) Micro-computed tomography: Introducing new dimensions to taxonomy. *ZooKeys* 263: 1-45. <https://doi.org/10.3897/zookeys.263.4261>
- Fauvel P (1917) Annélides Polychètes de l'Australie méridionale. *Archive de Zoologie Expérimentale et Générale* 56: 159-277. URL: <http://biodiversitylibrary.org/page/6322379>
- Fauvel P (1923) Faune de France: Polychètes errantes. Paul Lechevalier, Paris, 488 pp. URL: [https://faunedefrance.org/bibliotheque/docs/P.FAUVEL\(FdeFr05\)Polychetes-errantes.pdf](https://faunedefrance.org/bibliotheque/docs/P.FAUVEL(FdeFr05)Polychetes-errantes.pdf)
- Fauvel P (1927) Faune de France. Polychètes sédentaires. Paul Lechevalier, Paris, 497 pp. URL: [http://www.faunedefrance.org/bibliotheque/docs/P.FAUVEL\(FdeFr16\)Polychetes-sentaires.pdf](http://www.faunedefrance.org/bibliotheque/docs/P.FAUVEL(FdeFr16)Polychetes-sentaires.pdf)
- Fauvel P (1936a) Contribution à la faune des Annélides Polychètes du Maroc. *Mémoires de la Société des Sciences Naturelles du Maroc* 43: 1-143. URL: [http://www.annelida.net/docs/Fauvel1936\\_ContrFaunaAnnPolychetesMarocOCR.pdf](http://www.annelida.net/docs/Fauvel1936_ContrFaunaAnnPolychetesMarocOCR.pdf)
- Fauvel P (1936b) Remarques sur les Néréidiens *Nereis succinea* Leuckart et *Nereis lamellosa* Ehlers. *Bulletin de la Société Zoologique de France* 61: 307-314. URL: <http://gallica.bnf.fr/ark:/12148/bpt6k57583633/f334>
- Fauvel P (1937) Les Fonds de Pêches près d'Alexandrie XI. Annélides Polychètes. *Direction des Recherches des Pêcheries, Notes et Mémoires* 90: 1-60.
- Fauvel P, Rullier F (1959) Annélides Polychètes. *Resultats Scientifiques des Campagnes de Calypso* 4. *Annales de l'Institut Océanographique Paris* 37: 143-205.
- Fiege D, Böggemann M (1999) *Ancistrosyllis fioronii*, a new species of Pilargidae from the North Sea, with a key and synoptic table of characters for all species of the genus *Ancistrosyllis* McIntosh, 1879 (Annelida, Polychaeta, Pilargidae). *Senckenbergiana biologica* 78 (135-140): .
- Fiege D, Kröncke I, Barnich R (2000a) High abundance of *Myriochele fragilis* Nilsen & Holthe, 1985 (Polychaeta: Oweniidae) in the deep sea of the Eastern Mediterranean. *Hydrobiologia* 426: 97-103. <https://doi.org/10.1023/A:1003947318925>
- Fiege D, Licher F, Mackie AS (2000b) A partial review of the European Magelonidae (Annelida: Polychaeta): *Magelona mirabilis* redefined and *M. johnstoni* sp. nov. distinguished. *Journal of the Marine Biological Association of the United Kingdom* 80 (2): 215-234. <https://doi.org/10.1017/s0025315499001800>
- Fishelson L (2000) Marine animal assemblages along the littoral of the Israeli Mediterranean seashore: The Red-Mediterranean Seas communities of species. *Italian Journal of Zoology* 67 (4): 393-415. <https://doi.org/10.1080/11250000009356345>

- Fitzhugh K, Giangrande A, Simboura N (1994) New species of *Pseudofabriciola* Fitzhugh, 1990 (Polychaeta: Sabellidae: Fabriciinae), from the Mediterranean Sea. *Zoological Journal of the Linnean Society* 110 (3): 219-241. <https://doi.org/10.1111/j.1096-3642.1994.tb02016.x>
- Forbes E (1842) On two remarkable marine Invertebrata inhabiting the Aegean Sea. *Annals and Magazine of Natural History* 8: 243-245. URL: <http://biodiversitylibrary.org/page/2338184>
- Fredj G (1974) Stockage et exploitation des données en ecologie marine. – C. Considerations biogéographiques sur le peuplement benthique de la Méditerranée. *Memoires de l'Institut Oceanographique de Monaco* 7: 1-88.
- Galil BS, Occhipinti-Ambrogi A, Gollasch S (2008) Biodiversity impacts of species introductions via marine vessels. In: Ameer A, Linden O (Eds) *Maritime traffic effects on biodiversity in the Mediterranean Sea: review of impacts, priority areas and mitigation measures*. IUCN Centre for Mediterranean Cooperation, Malaga, 117-158 pp. URL: [https://cmsdata.iucn.org/downloads/maritime\\_v1\\_lr.pdf](https://cmsdata.iucn.org/downloads/maritime_v1_lr.pdf)
- Gambi MC, Bremec CS, Conti G (1998) Polychaete distribution, diversity and seasonality related to seagrass cover in shallow soft bottoms of the Tyrrhenian Sea (Italy). *Scientia Marina* 62 (1-2): 1-17. URL: <http://scimar.icm.csic.es/scimar/pdf/62/sm62n10>
- Gamenick I, Abbiati M, Giere O (1998) Field distribution and sulphide tolerance of *Capitella capitata* (Annelida: Polychaeta) around shallow water hydrothermal vents off Milos (Aegean Sea). A new sibling species? *Marine Biology* 130 (3): 447-453. <https://doi.org/10.1007/s002270050265>
- George J, Hartmann-Schröder G (1985) Polychaetes: British Amphinomida, Spintherida & Eunicida. *Synopses of the British Fauna (new series)*. 32. Brill/Backhuys, London, 221 pp.
- Georgiou E (1983) Study of the polychaete fauna in Mazoma lagoon of Amvrakikos Gulf, and especially of the populations of the species *Nainereis laevigata* and *Harmothoe spinifera*. MSc thesis. National and Kapodistrian University of Athens, Athens, 100 pp. [In Greek].
- Gerovasileiou V, Chintiroglou CC, Vafidis D, Koutsoubas D, Sini M, Dailianis T, Issaris Y, Akritopoulou E, Dimarchopoulou D, Voultsiadou E (2015) Census of biodiversity in marine caves of the eastern Mediterranean Sea. *Mediterranean Marine Science* 16 (1): 245-265. <https://doi.org/10.12681/mms.1069>
- Gherardi M, Gravina MF, Giangrande A (2002) Note tassonomiche ed ecologiche su *Micromaldane ornithochaeta* (Polychaeta, Maldanidae), rinvenuta lungo le coste Italiane Meridionali. *Thalassia Salentina* (26)133-143. <https://doi.org/10.1285/i15910725v26p133>
- Giangrande A (1989) Censimento dei policheti dei mari Italiani: Sabellidae Malmgren, 1867. *Atti della Società Toscana di Scienze Naturali Memorie* 96: 153-189. URL: <http://www.stsn.it/AttiB1989/giangrande.pdf>
- Giangrande A (1992) The genus *Chone* (Polychaeta, Sabellidae) in the Mediterranean Sea with description of *C. longiseta* n. sp. *Bollettino di Zoologia* 59: 517-529. <https://doi.org/10.1080/11250009209386712>
- Giangrande A (1994) The genus *Demonax* (Polychaeta, Sabellidae) in the Mediterranean Sea, with description of *D. tommasi* n. sp. *Bollettino di Zoologia* 61 (3): 229-233. <https://doi.org/10.1080/11250009409355890>

- Giangrande A, Castelli A (1986) Occurrence of *Fabricia filamentosa* Day, 1963 (Polychaeta, Sabellidae, Fabriciinae) in the Mediterranean Sea. *Oebalia* 13: 119-122.
- Giangrande A, Gambi MC (1982) Distribuzione dei Policheti nei fondi mobili della Rada di Augusta (Sicilia). *Bollettino dei Musei e Degli Istituti Biologici dell'Università di Genova* 50: 218-222.
- Giangrande A, Licciano M (2004) Factors influencing latitudinal pattern of biodiversity: an example using Sabellidae (Annelida, Polychaeta). *Biodiversity and Conservation* 13 (9): 1633-1646. <https://doi.org/10.1023/b:bioc.0000029327.63397.6b>
- Giangrande A, Licciano M (2006) The genus *Euchone* (Polychaeta, Sabellidae) in the Mediterranean Sea, addition of two new species and discussion on some closely related taxa. *Journal of Natural History* 40 (21-22): 1301-1330. <https://doi.org/10.1080/00222930600901458>
- Giangrande A, Licciano M (2008) Revision of the species of *Megalomma* (Polychaeta, Sabellidae) from the Mediterranean Sea, with the description of *M. messapicum* n. sp. *Italian Journal of Zoology* 75 (2): 207-217. <https://doi.org/10.1080/11250000801913124>
- Giangrande A, Gambi MC, Fresi E (1981) Two species of polychaetes new to the Mediterranean fauna. *Bollettino di Zoologia* 48: 311-317. <https://doi.org/10.1080/11250008109439349>
- Giangrande A, Cosentino A, Lo Presti C, Licciano M (2012) Sabellidae (Annelida) from the Faro coastal lake (Messina, Ionian Sea), with the first record of the invasive species *Branchiomma bairdi* along the Italian coast. *Mediterranean Marine Science* 13 (2): 283-293. <https://doi.org/10.12681/mms.310>
- Giangrande A, Delos AL, Frascchetti S, Musco L, Licciano M, Terlizzi A (2003) Polychaete assemblages along a rocky shore on the South Adriatic coast (Mediterranean Sea): patterns of spatial distribution. *Marine Biology* 143 (6): 1109-1116. <https://doi.org/10.1007/s00227-003-1162-0>
- Gibson PH (1978) Systematics of *Dodecaceria* (Annelida: Polychaeta) and its relation to the reproduction of its species. *Zoological Journal of the Linnean Society* 63 (3): 275-287. <https://doi.org/10.1111/j.1096-3642.1978.tb02565.x>
- Gibson PH (2015) Augmented descriptions and nomenclature of the cirratulid polychaetes *Dodecaceria fimbriata* and *D. concharum*. *Journal of the Marine Biological Association of the United Kingdom* 95 (04): 697-702. <https://doi.org/10.1017/s0025315414001854>
- Gibson PH, Heppell D (1995) *Dodecaceria concharum* Örsted, 1843 and *Heterocirrus fimbriatus* Verrill, 1879 (currently *D. fimbriata*) (Annelida, Polychaeta): proposed conservation of the specific names by the designation of a neotype for *D. concharum*. *Bulletin of Zoological Nomenclature* 52: 27-33. <https://doi.org/10.5962/bhl.part.6721>
- Gidholm L (1967) A revision of Autolytinae (Syllidae, Polychaeta) with special reference to Scandinavian species, and with notes on external and internal morphology, reproduction and ecology. *Arkiv för Zoologi* 19: 157-213.
- Gil J, Nishi E (2017) Nomenclatural checklist for *Acromegalomma* species (Annelida, Sabellidae), a nomen novum replacement for the junior homonym *Megalomma* Johansson, 1926. *ZooKeys* 677: 131-150. <https://doi.org/10.3897/zookeys.677.12030>
- Glasby CJ, Hutchings P (2014) Revision of the taxonomy of *Polycirrus* Grube, 1850 (Annelida: Terebellida: Polycirridae). *Zootaxa* 3877: 1-117. <https://doi.org/10.11646/zootaxa.3877.1.1>

- Gogou M (2009) Μελέτη της επίδρασης του κυματισμού των πλοίων στις βενθικές βιοκοινότητες του φανερόγαμου *Posidonia oceanica* στην θαλάσσια περιοχή στο στενό της Μυτιλήνης. [Study of the influence of ship-induced wave action on the benthic communities of the phanerogam *Posidonia oceanica* in the marine region of the Strait of Mytilene.]. BSc thesis. University of the Aegean, Mytilini, 110 pp. [In Greek].
- Gómez SCG, Carrera-Parra L, Mas FA, Freitas R, Martins R (2015) Novel insights on the diversity and ecology of the Family Lumbrineridae (Polychaeta) along the Iberian Peninsula coasts. *Journal of the Marine Biological Association of the United Kingdom* 96 (7): 1427-1435. <https://doi.org/10.1017/s0025315415001861>
- Gonzalez B, Martínez A, Borda E, Iliffe T, Eibye-Jacobsen D, Worsaae K (2017a) Phylogeny and systematics of Aphroditiformia. *Cladistics* <https://doi.org/10.1111/cla.12202>
- Gonzalez B, Petersen HC, Domenico MD, Martínez A, Armenteros M, García-Machado E, Møller PR, Worsaae K (2017b) Phylogeny and biogeography of the scaleless scale worm *Pisione* (Sigalionidae, Annelida). *Ecology and Evolution* 7 (9): 2894-2915. <https://doi.org/10.1002/ece3.2853>
- Goumenaki P (2005) Δομή και εποχική μεταβλητότητα μεγαβενθικών βιοκοινοτήτων σκληρού υποστρώματος της ανώτερης υποπαραλιακής ζώνης στο στενό της Μυτιλήνης. [Structure and seasonal variation of megabenthic biocommunities of the hard substrates in the upper midlittoral zone in the Strait of Mytilene]. BSc thesis. University of the Aegean, Mytilini, 73 pp. [In Greek].
- Grassle JP, Grassle JP (1976) Sibling species in the marine pollution indicator *Capitella* (Polychaeta). *Science* 192 (4239): 567-569. <https://doi.org/10.1126/science.1257794>
- Gravina M (1991) Observations on the genus *Branchiomaldane* (Polychaeta: Arenicolidae) with a new record for the Italian fauna: *Branchiomaldane vicenti* Langerhans, 1881. *Oebalia* 17: 159-165.
- Gravina MF, Cantone G (1991) *Lumbrinerides neogesae* Miura, 1980, un Lumbrineridae (Annelida, Polychaeta) nuovo per il Mediterraneo. *Animalia* 18: 149-155.
- Gravina MF, Somaschini A (1990) Censimento dei Policheti dei Mari Italiani: Capitellidae Grube, 1862. *Atti della Società Toscana di Scienze Naturali Residente in Pisa, Memorie, Processi Verbal, Serie B* 97: 259-285. URL: <http://www.stsn.it/AttiB1990/gravina-somaschini.pdf>
- Gravina MF, Lezzi M, Bonifazi A, Giangrande A (2015) The genus *Nereis* L., 1758 (Polychaeta, Nereididae): State of the art for identification of Mediterranean species. *Atti della Società Toscana di Scienze Naturali Memorie Serie. B* 122: 147-164. <https://doi.org/10.2424/ASTSN.M.2015.13>
- Green KD (2002) Capitellidae (Polychaeta) from the Andaman Sea. *Phuket Marine Biological Center Special Publication* 24: 249-343.
- Grémare A, Amouroux JM, Vétion G (1998) Long-term comparison of macrobenthos within the soft bottoms of the Bay of Banyuls-sur-Mer (northwestern Mediterranean Sea). *Journal of Sea Research* 40 (3-4): 281-302. [https://doi.org/10.1016/S1385-1101\(98\)00032-X](https://doi.org/10.1016/S1385-1101(98)00032-X)
- Guérin J, Massé H (1976) Étude expérimentale sur le recrutement des espèces de la macrofaune benthique des substrats meubles. 1. Méthodologie - données qualitatives et quantitatives. *Tethys* 8 (2): 151-168.



- Guerin JP, Kerambrun P (1984) Role of reproductive characters in the taxonomy of Spionids and elements of speciation in the *Malacoceros fuliginosus* complex. *Fortschritte der Zoologie* 29: 317-333.
- Guille A, Laubier L (1966) Additions à la faune des Annélides Polychètes de Banyuls-sur-Mer. *Vie et milieu* 17 (1B): 259-282.
- Harmelin J (1969) Contribution à l'étude de l'endofaune des prairies d'*Halophila stipulacea* de Méditerranée Orientale. *Travaux de la Station marine d'Endoume* 61 (45): 305-320.
- Harmelin JG (1968) Note sur trois Capitellidae (Annélides Polychètes) récoltés en Méditerranée, avec description d'un nouveau genre: *Peresiella*. *Recueil des Travaux de la Station Marine d'Endoume* 43 (59): 253-259. URL: [http://paleopolis.rediris.es/benthos/REF/som/R-pdf/1968-43-59\\_253.pdf](http://paleopolis.rediris.es/benthos/REF/som/R-pdf/1968-43-59_253.pdf)
- Harris T (1970) The occurrence of *Manayunkia aestuarina* (Bourne) and *Mercierella enigmatica* Fauvel (Polychaeta) in non-brackish water in Britain. *Journal of Experimental Marine Biology and Ecology* 5: 105-112. [https://doi.org/10.1016/0022-0981\(70\)90011-0](https://doi.org/10.1016/0022-0981(70)90011-0)
- Hartley J (1981) The family Paraonidae (Polychaeta) in British waters, a new species and new records with a key to species. *Journal of the Marine Biological Association of the United Kingdom* 61 (1): 279-280. <https://doi.org/10.1017/S0025315400045975>
- Hartley J (1984) Cosmopolitan polychaete species: the status of *Aricidea belgicae* (Fauvel, 136) and notes on the identity of *A. suecica* Eliason, 1920 (Polychaeta: Paraonidae). In: Hutchings P (Ed.) *Proceedings of the First International Polychaete Conference*. Sydney. Linnean Society of New South Wales, Sydney, 7-20 pp.
- Hartley J (2014) A review of the occurrence and ecology of dense populations of *Ditrupa arietina* (Polychaeta: Serpulidae). *Memoirs of Museum Victoria* 71: 85-95. <https://doi.org/10.24199/j.mmv.2014.71.09>
- Hartmann-Schröder G (1996) *Annelida, Borstenwürmer, Polychaeta*. 2. Neubearbeitete Auflage. *Tierwelt Deutschlands*, 647 pp.
- Hartman O (1944) Polychaetous Annelids. Part V. Eunicea. *Allan Hancock Pacific Expeditions* 10 (1): 1-237. URL: <http://digitallibrary.usc.edu/cdm/ref/collection/p15799coll82/id/21633>
- Hartman O (1947) Polychaetous annelids. Part VII. Capitellidae. *Allan Hancock Pacific Expeditions* 10 (4/5): 391-481.
- Hartman O (1948) The marine annelids erected by Kinberg. With some notes on some other types in the Swedish State Museum. *Arkiv för Zoologi* 42A (1): 1-137.
- Hartman O (1965a) Catalogue of the Polychaetous Annelids of the world. Supplement 1960-1965 and Index. *Occasional Papers of the Allan Hancock Foundation* 23: 1-204. URL: <http://digitallibrary.usc.edu/cdm/ref/collection/p15799coll82/id/19512>
- Hartman O (1965b) Deep-water benthic polychaetous annelids off New England to Bermuda and other North Atlantic areas. *Occasional Papers of the Allan Hancock Foundation* 28: 1-378. URL: <https://repository.si.edu/handle/10088/3458>
- HCMR (2011) Μελέτη των επιπτώσεων της ρύπανσης που προκλήθηκε από το ατύχημα του κρουαζιερόπλοιου «Sea Diamond» στον όρμο Αθηνιός της νήσου Θήρας. [Study of the short-term effects from the accident of the cruise ship 'SEA DIAMOND' in Athinios Bay, Thera island. Technical report]. Hatzianestis J (Ed.) Ministry of Merchantile Marine, Anavyssos.

- Helgason G, Gardarsson A, Svavarsson J, Adalsteinsdottir K, Gudmundsson H (1990) Polychaetes new to the Icelandic Fauna, with remarks on some previously recorded species. *Sarsia* 75 (3): 203-212. <https://doi.org/10.1080/00364827.1990.10413448>
- Helm C, Bernhart S, zu Siederdisen CH, Nickel B, Bleidorn C (2012) Deep sequencing of small RNAs confirms an annelid affinity of Myzostomida. *Molecular Phylogenetics and Evolution* 64 (1): 198-203. <https://doi.org/10.1016/j.ympev.2012.03.017>
- Hilbig B, Blake JA (2006) Deep-sea polychaete communities in the Northeast Pacific Ocean off the Gulf of the Farallones, California. *Bulletin of Marine Science* 78: 243-269. URL: <http://www.ingentaconnect.com/contentone/umrsmas/bullmar/2006/00000078/00000002/art00002>
- Holthe T (1986) Polychaeta Terebellomorpha. *Marine Invertebrates of Scandinavia* 7: 1-194.
- Hughes DJ, Crawford M (2006) A new record of the vestimentiferan *Lamellibrachia* sp. (Polychaeta: Siboglinidae) from a deep shipwreck in the eastern Mediterranean. *Marine Biodiversity Records* 2: e21. <https://doi.org/10.1017/S1755267206001989>
- Hutchings P (1998) Biodiversity and functioning of polychaetes in benthic sediments. *Biodiversity and Conservation* 7 (9): 1133-1145. <https://doi.org/10.1023/A:1008871430178>
- Hutchings P, Glasby CJ (1987) The Thelepininae (Terebellidae) from Australia, with a discussion of the generic and specific characters of the subfamily. *Bulletin of the Biological Society of Washington* 7: 217-250.
- Hutchings P, Glasby CJ (1995) Description of the widely reported terebellid polychaetes *Loimia medusa* (Savigny) and *Amphitrite rubra* (Risso). *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 92: 149-154.
- Hutchings P, Peart R (2002) A review of the genera of Pectinariidae (Polychaeta) together with a description of the Australian Fauna. *Records of the Australian Museum* 54: 99-127. <https://doi.org/10.3853/j.0067-1975.54.2002.1356>
- Hutchings P, Glasby C, Wijnhoven S (2012) Note on additional diagnostic characters for *Marphysa sanguinea* (Montagu, 1813) (Annelida: Eunicida: Eunicidae), a recently introduced species in the Netherlands. *Aquatic Invasions* 7 (2): 277-282. <https://doi.org/10.3391/ai.2012.7.2.014>
- Hutchings P, Nogueira JM, Carrerette O (2017) Terebellidae s.l.: Polycirridae Malmgren, 1866, Terebellidae Johnston, 1846, Thelepodidae Hessle, 1917, Trichobranchidae Malmgren, 1866, and Telothelepodidae Nogueira, Fitzhugh & Hutchings, 2013. In: Westheide W, Purschke, G (eds). *Handbook of Zoology Online. A Natural History of the Phyla of the Animal Kingdom. Annelida: Polychaetes*. [https://www.degruyter.com/view/Zoology/bp\\_029147-6\\_68](https://www.degruyter.com/view/Zoology/bp_029147-6_68). Accessed on: 2017-12-01.
- Hutchings PA, Karageorgopoulos P (2003) Designation of a neotype of *Marphysa sanguinea* (Montagu, 1813) and a description of a new species of *Marphysa* from eastern Australia. *Hydrobiologia* 496: 87-94. <https://doi.org/10.1023/a:1026124310552>
- Hutchings PA, Rainer S (1981) Designation of a neotype of *Capitella filiformis* Claparède, 1864, type species of the genus *Heteromastus* (Polychaeta: Capitellidae). *Records of the Australian Museum* 34: 373-380. <https://doi.org/10.3853/j.0067-1975.34.1982.295>
- Iannotta M, Gambi M, Patti F (2009) Molecular evidence of intraspecific variability in *Lysidice ninetta* (Polychaeta: Eunicidae) in the Mediterranean Sea. *Aquatic Biology* 6: 121-132. <https://doi.org/10.3354/ab00160>

- Iannotta MA, Patti FP, Ambrosino M, Procaccini G, Gambi MC (2006) Phylogeography of two species of *Lysidice* (Polychaeta, Eunicidae) associated to the seagrass *Posidonia oceanica* in the Mediterranean Sea. *Marine Biology* 150 (6): 1115-1126. <https://doi.org/10.1007/s00227-006-0405-2>
- Ifremer (2016) REBENT- Réseau benthique. Institut Français pour la Recherche et l'exploitation de la mer. Release date: 2013-11-29. URL: <http://www.eurobis.org/imis?module=dataset&dasiid=4412>
- IOFR (1984) Technical report on the faunal and sediment analysis near oil drilling sites in the Gulf of Kavala. C Bogdanos (Ed). Institute of Fisheries Research, 20 pp. [In Greek].
- Iroso I (1921) Revisione dei Serpulidi e Sabellidi del Golfo di Napoli. *Pubblicazioni della Stazione Zoologica di Napoli* 3: 47-91.
- Isari S, Somarakis S, Christou ED, Fragopoulou N (2010) Summer mesozooplankton assemblages in the north-eastern Aegean Sea: the influence of Black Sea water and an associated anticyclonic eddy. *Journal of the Marine Biological Association of the United Kingdom* 91 (1): 51-63. <https://doi.org/10.1017/S0025315410000123>
- Issaris Y, Salomidi M (2012) Benthos and Pelagos. *Life of the Hellenic Seas*. Hellenic Post Edition, Athens, 257 pp.
- Jacquotte R (1963) Signification biocénotique des fonds a *Peysonnellia polymorpha* (Zan.) Schmitz des côtes de Provence (de la Baie de Marseille aux Îles d'Hyères). *Travaux de la Station Marine d'Endoume* 44 (29): 27-41.
- Jiménez-Cueto S, Suárez-Morales E, Salazar-Vallejo SI (2006) Iospilids (Polychaeta: Iospilidae) from the northwest Caribbean Sea, with observations on reproductive structures. *Zootaxa* 1211: 53-68.
- Jimi N, Salazar-Vallejo S, Kajihara H (2017) Designation of a neotype and redescription of *Hesione reticulata* von Marenzeller, 1879 from Japan (Annelida, Hesionidae). *ZooKeys* 657: 29-41. <https://doi.org/10.3897/zookeys.657.11064>
- Jirkov I (2001) Polikhety severnogo Ledovitogo Okeana. [Polychaeta of the Arctic Ocean]. Yanus-K Press, Moscow, 632 pp. [In Russian].
- Jirkov I (2008) Revision of Ampharetidae (Polychaeta) with modified thoracic notopodia. *Invertebrate Zoology* 5 (2): 111-132.
- Jirkov IA, Leontovich MK (2017) Review of genera within the *Axionice/ Pista* complex (Polychaeta, Terebellidae), with discussion of the taxonomic definition of other Terebellidae with large lateral lobes. *Journal of the Marine Biological Association of the United Kingdom* 97 (0): 911-934. <https://doi.org/10.1017/s0025315417000923>
- Johnson PG (1984) Family Spionidae Grube, 1850. In: Uebelacker JM, Johnson PG (Eds) *Taxonomic Guide to the Polychaetes of the Northern Gulf of Mexico*. Final Report to the Minerals Management Service, contract 14-12-001-29091. 7. Barry A. Vittor & Associates, Inc, Mobile, Alabama, 249 pp.
- Johnston G (1865) A catalogue of the British non-parasitical worms in the collection of the British Museum. British Museum, London, 365 pp. <https://doi.org/10.5962/bhl.title.12291>
- Jolly MT, Jollivet D, Gentil F, Thiébaud E, Viard F (2005) Sharp genetic break between Atlantic and English Channel populations of the polychaete *Pectinaria koreni*, along the North coast of France. *Heredity* 94 (1): 23-32. <https://doi.org/10.1038/sj.hdy.6800543>
- Jolly MT, Viard F, Gentil F, Thiébaud E, Jollivet D (2006) Comparative phylogeography of two coastal polychaete tubeworms in the Northeast Atlantic supports shared history and

- vicariant events. *Molecular Ecology* 15 (7): 1841-55. <https://doi.org/10.1111/j.1365-294X.2006.02910.x>
- Jones ML (1977) A redescription of *Magelona papillicornis* F. Müller. In: Reish DJ, Fauchald K (Eds) *Essays on polychaetous annelids in memory of Dr. Olga Hartman*. Allan Hancock Foundation, University of Southern California, Los Angeles, 247-266 pp.
  - Jumars PA (1974) A generic revision of the Dorvilleidae (Polychaeta), with six new species from the deep North Pacific. *Zoological Journal of the Linnean Society* 54 (2): 101-135. <https://doi.org/10.1111/j.1096-3642.1974.tb00794.x>
  - Karakassis I (1991) Συμβολή στη μελέτη του βενθικού οικοσυστήματος της υφαλοκρηπίδας της Κρήτης. [Contribution to the study of the benthic ecosystem of the continental shelf of Crete]. PhD thesis. University of Crete, Heraklion, 195 pp. [In Greek].
  - Karlou K, Melea A (1977) Συστηματική κατάταξη και οικολογικές παρατηρήσεις στους ζωικούς πληθυσμούς των σπόγγων. [Systematic classification and ecological observations of the animal populations of sponges]. Laboratory of Zoology, National and Kapodistrian University of Athens, Athens, 25 pp. [In Greek].
  - Kato T, Pleijel F (2003) A revision of *Paranaitis* Southern, 1914 (Polychaeta: Phyllodocidae). *Zoological Journal of the Linnean Society* 138 (4): 379-429. <https://doi.org/10.1046/j.1096-3642.2003.00069.x>
  - Katsiaras N, Simboura N (2015) Beyond WFD: New information on polychaete diversity, geographical and ecological distributions from the Hellenic Monitoring Network. 11th Panhellenic Symposium of Oceanography and Fisheries, Mytilini, Greece, May 2015. 585-558 pp. URL: <https://oceanos-dspace.hcmr.gr/bitstream/handle/123456789/2585/PanhellSympOceanFish11%2585-588%292015..pdf>
  - Katsiaras N, Simboura N, Koutsoubas D (2014) The rare subgroup C1 of *Marphysa* (Polychaeta, Eunicidae): re-description of species and first records in the Mediterranean Sea. *Zootaxa* 3873 (3): 201-217. <https://doi.org/10.11646/zootaxa.3873.3.1>
  - Katsiaras N, Rousou M, Carrera-Parra L, Garcia-Gomez SC, Simboura N, Louizidou P, Chintiroglou CC, Martins R (2017) Taxonomy, ecology and geographic distribution of *Gallardonneris iberica* (Polychaeta, Lumbrineridae) in southern Europe. *Journal of the Marine Biological Association of the United Kingdom* 1-10. <https://doi.org/10.1017/s0025315417001254>
  - Katzmann W (1973) Polychaeten von Sedimentböden der mittleren Adria (50–525 m). *Zoologische Jahrbücher, Abteilung für Systematik, Ökologie und Geographie der Tiere* 100 (4): 436-450.
  - Katzmann W, Laubier L (1975) Paraonidae (Polychètes sédentaires) de l'Adriatique. *Annalen des Naturhistorischen Museums in Wien* 79: 567-588. URL: <http://archimer.ifremer.fr/doc/1975/publication-5086.pdf>
  - Keklikoglou K, Faulwetter S, Chatzigeorgiou G, Badalamenti F, Kitsos MS, Arvanitidis C (2013) MidMedPol: Polychaetes from midlittoral rocky shores in Greece and Italy (Mediterranean Sea). *Biodiversity Data Journal* 1 <https://doi.org/10.3897/BDJ.1.e961>
  - Khedhri I, Tovar-Hernández M, Bonifácio P, Ahmed A, Aleya L (2017) First report of the invasive species *Branchiomma bairdi* McIntosh, 1885 (Annelida: Sabellidae) along the Tunisian coast (Mediterranean Sea). *BioInvasions Records* 6 (2): 139-145. <https://doi.org/10.3391/bir.2017.6.2.09>

- Kinberg JG (1857) Nya sl gten och arter af Annelider.  fversigt af Kongliga Vetenskaps-Akademiens F rhandlingar, Stockholm 14: 11-14. URL: <http://www.biodiversitylibrary.org/item/150443#page/19/mode/1up>
- Kirkegaard J (2001) Polychaetes of the families Glyceridae, Goniadidae, and Nereididae from the North Atlantic around Iceland. Sarsia 86 (1): 13-20. <https://doi.org/10.1080/00364827.2001.10420457>
- Kirkegaard JB (1983) Bathyal benthic polychaetes from the N.E. Atlantic Ocean, S.W. of the British Isles. Journal of the Marine Biological Association of the United Kingdom 63: 593-608. <https://doi.org/10.1017/S0025315400070909>
- Kisseleva M (1961) Qualitative and quantitative distribution of the benthos in the area surrounding Dardanelles, Aegean Sea. Studies of Sevastopol Biological Station 14: 135-146. [In Russian]. URL: [http://vuzlib.com.ua/articles/book/21788-Kachestvennoe\\_i\\_kolichestvenno/1.html](http://vuzlib.com.ua/articles/book/21788-Kachestvennoe_i_kolichestvenno/1.html)
- Kisseleva MI (1983) Comparative characteristics of the Benthos at some banks in the Aegean Sea. Thalassographica 6: 107-118. URL: <http://hdl.handle.net/123456789/250>
- Kitamori R (1960) Two new species of cirratulid and Nephthydidae (Annelida: Polychaeta). Bulletin of the Japanese Society for Scientific Fisheries 26: 1082-1085. <https://doi.org/10.2331/suisan.26.1082>
- Kitsos M (2003) Συμβολή στη μελέτη της βιοποικιλότητας των συνευρέσεων του σκληρού μεσοπαραλιακού και υπερπαραλιακού υποστρώματος. [Contribution to the study of the biodiversity of the midlittoral and supralittoral hard substratum assemblages]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 313 pp. [In Greek]. <https://doi.org/10.12681/eadd/14684>
- Kitsos M, Christodoulou M, Arvanitidis C, Mavidis M, Kirmitzoglou I, Koukouras A (2005) Composition of the organismic assemblage associated with *Caretta caretta*. Journal of the Marine Biological Association of the United Kingdom 85 (2): 257-261. <https://doi.org/10.1017/s0025315405011136h>
- Klautau M, Russo CM, Lazoski C, Boury-Esnault N, Thorpe J, Sole-Cava A (1999) Does cosmopolitanism result from overconservative systematics? A case study using the marine sponge *Chondrilla nucula*. Evolution 53 (5): 1414. <https://doi.org/10.2307/2640888>
- Knight-Jones P (1983) Contributions to the taxonomy of Sabellidae (Polychaeta). Zoological Journal of the Linnean Society 79: 245-295. <https://doi.org/10.1111/j.1096-3642.1983.tb01167.x>
- Knight-Jones P, Knight-Jones EW (1977) Taxonomy and Ecology of British Spirorbidae (Polychaeta). Journal of the Marine Biological Association of the United Kingdom 57 (2): 453-500. <https://doi.org/10.1017/S002531540002186X>
- Knight-Jones P, Perkins T (1998) A revision of *Sabella*, *Bispira* and *Stylomma* (Polychaeta: Sabellidae). Zoological Journal of the Linnean Society 123: 385-467. <https://doi.org/10.1111/j.1096-3642.1998.tb01370.x>
- Knight-Jones P, Knight-Jones EW, Ergen Z (1991) Sabelliiform polychaetes, mostly from Turkey's Aegean coast. Journal of Natural History 25: 837-858. <https://doi.org/10.1080/00222939100770561>
- Knight-Jones P, Knight-Jones EW, Kawahara T (1975) A review of the genus *Janua*, including *Dexiospira* (Polychaeta: Spirorbinae). Zoological Journal of the Linnean Society, London 56 (2): 91-129. <https://doi.org/10.1111/j.1096-3642.1975.tb00812.x>

- Knight-Jones P, Darbyshire T, Petersen ME, Tovar-Hernández M (2017) What is *Pseudopotamilla reniformis* (Sabellidae)? Comparisons of populations from Britain, Iceland and Canada with comments on *Eudistylia* and *Schizobranchia*. *Zootaxa* 4254 (2): 201. <https://doi.org/10.11646/zootaxa.4254.2.3>
- Koukouras A (1979) Βιονομική μελέτη της μακροπανίδας του κινητού υποστρώματος της μεσοπαραλιακής ζώνης στο Στρυμονικό και Θερμαϊκό κόλπο. [Bionomic study of the macrofauna of the mediolittoral soft substratum in Strymonikos and Thermaikos Gulfs]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 283 pp. <https://doi.org/10.12681/eadd/0168>
- Koukouras A, Rouso A (1991) Midlittoral soft substratum macrofaunal assemblages in the North Aegean Sea. *P.S.Z.N.I.: Marine Ecology* 12: 293-316. <https://doi.org/10.1111/j.1439-0485.1991.tb00260.x>
- Koukouras A, Voultziadou-Koukoura E, Chintiroglou C, Dounas C (1985) Benthic bionomy of the North Aegean Sea III. A comparison of the macrobenthic animal assemblages associated with seven sponge species. *Cahiers de Biologie Marine* 26 (3): 301-319. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Koukouras A, Russo A, Voultziadou-Koukoura E, Dounas C, Chintiroglou C (1992) Relationship of sponge macrofauna with the morphology of their hosts in the North Aegean Sea. *Internationale Revue der gesamten Hydrobiologie und Hydrographie* 77 (4): 609-619. <https://doi.org/10.1002/iroh.19920770406>
- Koulouri P, Dounas C, Arvanitidis C, Koutsoubas D, Tselepides A, Eleftheriou A (2015) A field experiment on trophic relations within the benthic boundary layer (BBL) over an oligotrophic continental shelf. *Estuarine, Coastal and Shelf Science* 164: 392-407. <https://doi.org/10.1016/j.ecss.2015.07.029>
- Kudenov J, Blake J (1978) A review of the genera and species of the Scalibregmidae (Polychaeta) with descriptions of one new genus and three new species from Australia. *Journal of Natural History* 12 (4): 427-444. <https://doi.org/10.1080/00222937800770291>
- Kudenov J, Borda E (2013) Review of lesser-known Amphinomidae (Annelida, Amphinomida) with the description of a new Genus. In: Hutchings PA (Ed.) 2013 International Polychaete Conference: Program and Abstract Handbook. Sydney, 4–9 August 2013. 83 pp. URL: [https://australianmuseum.net.au/uploads/documents/30456/ipc2013\\_program\\_abstract\\_handbook.pdf](https://australianmuseum.net.au/uploads/documents/30456/ipc2013_program_abstract_handbook.pdf)
- Kupriyanova E, Sun Y, ten Hove H, Wong E, Rouse G (2015) Serpulidae (Annelida) of Lizard Island, Great Barrier Reef, Australia. *Zootaxa* 4019 (1): 275-353. <https://doi.org/10.11646/zootaxa.4019.1.13>
- Kurt Şahin G (2014) *Marphysa cinari*, a new species of Eunicidae (Polychaeta) from the coasts of Turkey (eastern Mediterranean) and re-descriptions of *Marphysa kinbergi* McIntosh, 1910 and *Marphysa disjuncta* Hartman, 1961. *Journal of Natural History* 48: 1989-2006. <https://doi.org/10.1080/00222933.2014.905125>
- Kurt Şahin G, Çınar ME (2009) Eunicidae (Polychaeta) species in and around İskenderun Bay (Levantine Sea, Eastern Mediterranean) with a new alien species for the Mediterranean Sea and a re-description of *Lysidice collaris*. *Turkish Zoological Journal* 33: 331-347. URL: <http://journals.tubitak.gov.tr/zoology/issues/zoo-09-33-3/zoo-33-3-11-0806-19.pdf>
- Kurt Şahin G, Çınar ME (2012) A check-list of polychaete species (Annelida: Polychaeta) from the Black Sea. *Journal of the Black Sea/ Mediterranean Environment* 18 (1): 10-48. URL: <http://dergipark.ulakbim.gov.tr/jbme/article/view/5000144348>

- Kurt-Şahin G, Dagli E, Sezgin M (2017) Spatial and temporal variations of soft bottom polychaetes of Sinop Peninsula (southern Black Sea) with new records. *Turkish Journal of Zoology* 41: 89-101. <https://doi.org/10.3906/zoo-1510-15>
- Labruno C, Grémare A, Amouroux J, Sardá R, Gil J, Taboada S (2007) Assessment of soft-bottom polychaete assemblages in the Gulf of Lions (NW Mediterranean) based on a mesoscale survey. *Estuarine, Coastal and Shelf Science* 71: 133-147. <https://doi.org/10.1016/j.ecss.2006.07.007>
- Langeneck J (2017) Diversity and evolution in the family Paraonidae (Annelida, Polychaeta): a morphological and molecular perspective. Università di Pisa, Pisa, 187 pp. URL: <https://etd.adm.unipi.it/t/etd-02132017-153609/>
- Langeneck J, Barbieri M, Maltagliati F, Castelli A (2017a) A new species of *Cirrophorus* (Annelida: Paraonidae) from Mediterranean organically enriched coastal environments, with taxonomic notes on the family. *Journal of the Marine Biological Association of the United Kingdom* 97 (05): 871-880. <https://doi.org/10.1017/s0025315417000935>
- Langeneck J, Busoni G, Aliani S, Castelli A (2017b) Deep-sea polychaetes (Annelida) from the Malta Escarpment (western Ionian Sea). *European Zoological Journal* 84 (1): 142-152. <https://doi.org/10.1080/24750263.2017.1287964>
- Langerhans P (1879) Die Wurmfauna von Madeira. *Zeitschrift für Wissenschaftliche Zoologie* 32 (4): 513-592. URL: [http://www.zobodat.at/pdf/Zeitschrift-fuer-wiss-Zoologie\\_32\\_0513-0592.pdf](http://www.zobodat.at/pdf/Zeitschrift-fuer-wiss-Zoologie_32_0513-0592.pdf)
- Lardicci C (1989) Censimento dei policheti dei mari Italiani: Spionidae Grube, 1850. *Atti della Società Toscana di Scienze Naturali - Memorie serie B* 96: 121-152. URL: <http://www.stsn.it/AttiB1989/lardicci.pdf>
- Lattig P, Martín D (2009) A taxonomic revision of the genus *Haplosyllis* Langerhans, 1887 (Polychaeta, Syllidae, Syllinae). *Zootaxa* 2220: 1-40.
- Lattig P, San Martín G, Martín D (2007) Taxonomic and morphometric analyses of the *Haplosyllis spongicola* complex (Polychaeta: Syllidae: Syllinae) from Spanish seas, with re-description of the type species and descriptions of two new species. *Scientia Marina* 71 (3): 551-570. <https://doi.org/10.3989/scimar.2007.71n3551>
- Laubier L (1966) Sur quelques Annelides Polychètes de la région de Beyrouth. *Miscellaneous Papers of Natural Sciences of American University of Beirut* 5: 1-15.
- Laubier L (1967) Sur quelques *Aricidea* (Polychètes, Paraonidae) de Banyuls-sur-Mer. *Vie et Milieu* 18 (1): 99-132.
- Laubier L, Paris J (1962) Faune marine des Pyrénées-Orientales. IV. Annélides Polychètes. *Vie et Milieu* 13 (1): 1-80.
- Laubier L, Ramos JM (1974) Paraonidae (Polychètes sédentaires) de Méditerranée. *Bulletin du Muséum d'Histoire Naturelle, Paris* 113 (168): 1097-1148. URL: <http://archimer.ifremer.fr/doc/1973/publication-5122.pdf>
- Le Garrec V, Grall J, Chevalier C, Guyonnet B, Jourde J, Lavesque N, Bonifácio P, Blake J (2016) *Chaetozone corona* (Polychaeta, Cirratulidae) in the Bay of Biscay: a new alien species for the North-east Atlantic waters? *Journal of the Marine Biological Association of the United Kingdom* 97: 433-44. <https://doi.org/10.1017/s0025315416000540>
- Lewis C, Karageorgopoulos P (2008) A new species of *Marphysa* (Eunicidae) from the western Cape of South Africa. *Journal of the Marine Biological Association of the UK* 88 (02): . <https://doi.org/10.1017/s002531540800009x>

- Lezzi M (2017) *Caulleriella mediterranea*, a new species of polychaete (Annelida: Cirratulidae) from the central Mediterranean Sea. The European Zoological Journal 1-10. <https://doi.org/10.1080/24750263.2017.1343397>
- Lezzi M, Cardone F, Mikac B, Giangrande A (2015) Variation and ontogenetic changes of opercular paleae in a population of *Sabellaria spinulosa* (Polychaeta: Sabellariidae) from the South Adriatic Sea, with remarks on larval development. Scientia Marina 79: 1-14. <https://doi.org/10.3989/scimar.04127.19A>
- Licciano M, Giangrande A (2006) The genus *Novafabricia* Fitzhugh, 1990 (Polychaeta: Sabellidae: Fabriciinae) along the Italian coast (Mediterranean Sea) with a description of *N. posidoniae* n. sp. Scientia Marina 70 (4): 673-678. <https://doi.org/10.3989/scimar.2006.70n4673>
- Licciano M, Giangrande A (2008) The genus *Branchiomma* (Polychaeta: Sabellidae) in the Mediterranean Sea, with the description of *B. maerli* n. sp. Scientia Marina 72 (2): 383-391. URL: <http://scimar.icm.csic.es/scimar/index.php/seclid/6/ldArt/3717/>
- Licher F (1999) Revision der Gattung *Typosyllis* Langerhans, 1879 (Polychaeta: Syllidae). Morphologie, Taxonomie und Phylogenie. Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 551: 1-336.
- Linnaeus C (1788) Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus differentiis, synonymis, locis. Editio decima tertia, aucta reformata. S. Cura J.F. Gmelin. G.E. Beer. Lipsiae 1: 2225-3910.
- Liu Y, Hutchings P, Sun S (2017) Three new species of *Marphysa* Quatrefages, 1865 (Polychaeta: Eunicida: Eunicidae) from the south coast of China and redescription of *Marphysa sinensis* Monro, 1934. Zootaxa 4263 (2): 228-250. <https://doi.org/10.11646/zootaxa.4263.2.2>
- Lobo J, Teixeira MAL, Borges LMS, Ferreira MSG, Hollatz C, Gomes P, Sousa R, Ravara A, Costa M, Costa F (2016) Starting a DNA barcode reference library for shallow water polychaetes from the southern European Atlantic coast. Molecular Ecology Resources 16 (1): 298-313. <https://doi.org/10.1111/1755-0998.12441>
- López E (2008) Redescription of *Aricidea belgicae* (Fauvel, 1936), an Antarctic Paraonidae (Annelida: Polychaeta), with reassessment of its taxonomic status based on new material. Proceedings of the Biological Society of Washington 121 (3): 297-307. <https://doi.org/10.2988/08-05.1>
- López E (2009) First record of the Levantine species *Scoloplos haasi* (Orbiniidae: Scolecida: Annelida) from the Mediterranean coast of Spain. Journal of the Marine Biological Association of the United Kingdom 90 (5): 1051-1054. <https://doi.org/10.1017/S0025315409991305>
- López E, O'Connor BDS, Gonzalez G (1987) Demography and gametogenic cycle of *Paradoneis armata* Glemarec (Polychaeta, Paraonidae) in Ria de La Coruna, North-west Spain. Ophelia 27 (2): 127-136. <https://doi.org/10.1080/00785236.1987.10422016>
- López E, Martín D, Britayev T, San Martín G (2001) New symbiotic associations involving Syllidae (Annelida: Polychaeta), with taxonomic and biological remarks on *Pionosyllis magnifica* and Syllis cf. armillaris. Journal of the Marine Biological Association of the United Kingdom 81: 399-409. <https://doi.org/10.1017/S0025315401004015>
- López García E (1995) Anelidos poliquetos de sustratos duros de las islas Chafarinas. [Annelida Polychaeta from hard substrates of the Chafarinas Islands]. PhD thesis.



Universidad Autónoma de Madrid, Madrid, 672 pp. [In Spanish]. URL: <http://hdl.handle.net/10486/4940>

- Lopez-Jamar E (1989) Primera cita para el litoral de la Peninsula Iberica del genero *Aonidella* (Polychaeta: Spionidae), con una redescription de la especie *Aonidella dayi* Maciolek 1983. Boletín Instituto Español de Oceanografía 5: 107-110.
- Lykousis V, Chronis G, Tselepides A, Price NB, Theocharis A, Siokou-Frangou I, Wambeke FV, Danovaro R, Stavrakakis S, Duineveld G, Georgopoulos D, Ignatiades L, Souvermezoglou A, Voutsinou-Taliadouri F (2002) Major outputs of the recent multidisciplinary biogeochemical researches undertaken in the Aegean Sea. Journal of Marine Systems 33–34: 313-334. [https://doi.org/10.1016/s0924-7963\(02\)00064-7](https://doi.org/10.1016/s0924-7963(02)00064-7)
- Maciolek Blake NJ (1983) Systematics of Atlantic Spionidae (Annelida: Polychaeta) with special reference to deep-water species. Boston University, Boston, 400 pp.
- Maciolek N (1985) A revision of the genus *Prionospio* Malmgren, with special emphasis on species from the Atlantic Ocean, and new records of species belonging to the genera *Apoprionospio* Foster and *Paraprionospio* Caullery (Polychaeta, Annelida, Spionidae). Zoological Journal of the Linnean Society 84 (4): 325-383. <https://doi.org/10.1111/j.1096-3642.1985.tb01804.x>
- Maciolek NJ (2000) New species and records of *Aonidella*, *Laonice*, and *Spiophanes* (Polychaeta: Spionidae) from shelf and slope depths of the Western North Atlantic. Bulletin Marine Science 67: 529-547.
- Mackie A (1984) On the identity and zoogeography of *Prionospio cirrifera* Wiren, 1883 and *Prionospio multibrachiata* Berkeley, 1927 (Polychaeta, Spionidae). In: Hutchings PA (Ed.) Proceedings of the First International Polychaete Conference, Sydney. Sydney. The Linnean Society of New South Wales
- Mackie ASY (1987) A review of species currently assigned to the genus *Leitoscoloplos* Day, 1977 (Polychaeta: Orbiniidae), with descriptions of species newly referred to *Scoloplos* Blainville, 1828. Sarsia 72: 37-41. <https://doi.org/10.1080/00364827.1987.10419701>
- Mackie ASY (1991) *Scalibregma celticum* new species (Polychaeta: Scalibregmatidae) from Europe, with a redescription of *Scalibregma inflatum* Rathke, 1843 and comments on the genus *Sclerobregma* Hartman, 1965. Bulletin of Marine Science 48 (2): 268-276. URL: <http://www.ingentaconnect.com/contentone/umrsmas/bullmar/1991/00000048/00000002/art00015>
- Mackie ASY, Gobin JF (1993) A review of the genus *Johnstonia* Quatrefages, 1866 (Polychaeta, Maldanidae), with a description of a new species from Trinidad, West Indies. Zoologica Scripta 22 (3): 229-241. <https://doi.org/10.1111/j.1463-6409.1993.tb00354.x>
- Magalhães W, Bailey-Brock JH (2012) Capitellidae Grube, 1862 (Annelida: Polychaeta) from the Hawaiian Islands with description of two new species. Zootaxa 3581: 1-52. URL: <http://www.mapress.com/zootaxa/list/2012/3581.html>
- Magalhães W, Seixas VC, Paiva PC, Elias R (2014) The Multitentaculate Cirratulidae of the Genera Cirriformia and Timarete (Annelida: Polychaeta) from Shallow Waters of Brazil. PLoS ONE 9 (11): e112727. <https://doi.org/10.1371/journal.pone.0112727>
- Maidanou M, Koulouri P, Arvanitidis C, Koutsoubas D, Dounas C (2017) Macrobenthic assemblage structure associated with a *Caulerpa prolifera* meadow in the eastern Mediterranean Sea (Elounda Bay, Crete Island). Regional Studies in Marine Science 14: 1-14. <https://doi.org/10.1016/j.rsma.2017.04.004>

- Makra A, Nicolaidou A (2000) Benthic communities of the inner Argolikos Bay. Belgian Journal of Zoology 130: 61-67. URL: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.630.6776&rep=rep1&type=pdf>
- Maltagliati F, Casu M, Castellini A (2004) Morphological and genetic evidence supports the existence of two species in the genus *Ophelia* (Annelida, Polychaeta) from the Western Mediterranean. Biological Journal of the Linnean Society 83 (1): 101-113. <https://doi.org/10.1111/j.1095-8312.2004.00374.x>
- Maltagliati F, Camilli L, Lardicci C, Castelli A (2001) Evidence for morphological and genetic divergence in *Perinereis cultrifera* (Polychaeta: Nereididae) from two habitat types at Elba Island. Journal of the Marine Biological Association of the United Kingdom 81 (3): 411-414. <https://doi.org/10.1017/S0025315401004027>
- Maltagliati F, Peru AP, Casu M, Rossi F, Lardicci C, Curini-Galletti M, Castelli A (2000) Is *Syllis gracilis* (Polychaeta: Syllidae) a species complex? An allozyme perspective. Marine Biology 136: 871-879. <https://doi.org/10.1007/s002270000288>
- Maltagliati F, Casu M, Lai T, Sareri DI, Casu D, Galletti MC, Cantone G, Castelli A (2005) Taxonomic distinction of *Ophelia barquii* and *O. bicornis* (Annelida: Polychaeta) in the Mediterranean as revealed by ISSR markers and the number of nephridiopores. Journal of the Marine Biological Association of the United Kingdom 85 (4): 835-841. <https://doi.org/10.1017/s0025315405011781>
- Marinov T (1959) Beitrag zur Erforschung der Polychäten vom Ägäischen Meer. Bulletin de l'Institut Zoologique de l'Académie des Sciences de Bulgarie 8: 292-294.
- Martín D (1987) Anélidos poliuetos asociados a las concreciones de algas calcáreas del litoral catalán. Miscel·lània Zoològica 11: 61-75.
- Martín D (1989) Revision de las especies de Owenidae (Annelida Polychaeta) de la Península Iberica. Scientia Marina 53: 47-52. URL: <http://hdl.handle.net/10261/28545>
- Martín D, Gil J (2010) Checklist of class Polychaeta (Phylum Annelida). In: Coll et al. (2010) The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. PLoS ONE 5: Supplementary Material 2. <https://doi.org/10.1371/journal.pone.0011842.s002>
- Martín D, Britayev T, San Martín G, Gil J (2003) Inter-population variability and character description in the sponge-associated *Haplosyllis spongicola* complex (Polychaeta: Syllidae). Hydrobiologia 496: 145-162. <https://doi.org/10.1023/a:1026184529208>
- Martín D, Koh B, Bhaud M, Dutrieux E, Gil J (2006) The genus *Owenia* (Annelida: Polychaeta) in the Persian Gulf, with description of *Owenia persica* sp. nov. Organisms Diversity & Evolution 6 (4): 325-326. <https://doi.org/10.1016/j.ode.2006.01.001>
- Martínez A, Purschke G, Worsaae K (2017) Protodrilidae Hatschek, 1888. In Zoology Online (n.d.). Berlin, Boston: De Gruyter. [https://www.degruyter.com/view/Zoology/bp\\_029147-6\\_8](https://www.degruyter.com/view/Zoology/bp_029147-6_8). Accessed on: 2017-11-20.
- Martins R, Carrera-Parra L, Quintino V, Rodrigues AM (2012a) Lumbrineridae (Polychaeta) from the Portuguese continental shelf (NE Atlantic) with the description of four new species. Zootaxa 3416: 1-21. URL: <https://pdfs.semanticscholar.org/07cb/4e360a86b22adccf55c2a0399f4a7bd23bc0.pdf>
- Martins R, Martín GS, Rodrigues AM, Quintino V (2012b) On the diversity of the genus *Pisione* (Polychaeta, Pisionidae) along the Portuguese continental shelf, with a key to European species. Zootaxa 3450: 12-22.

- Marzano CN, Gravina MF, Fianchini A, Portacci G, Gherardi M, Tursi A, Corriero G (2010) The macrozoobenthos of the Karavasta lagoon system (Albania): local assemblages and geographical comparisons. *Marine Ecology* 31 (4): 622-632. <https://doi.org/10.1111/j.1439-0485.2010.00381.x>
- Mavraki D, Fanini L, Tsompanou M, Gerovasileiou V, Nikolopoulou S, Chatzinikolaou E, Plaitis W, Faulwetter S (2016) Rescuing biogeographic legacy data: The "Thor" Expedition, a historical oceanographic expedition to the Mediterranean Sea. *Biodiversity Data Journal* 4: e11054. <https://doi.org/10.3897/bdj.4.e11054>
- McIntosh WC (1895) Note on the results of the voyage of the Challenger. *Annelida and Nemerteans*. Natural Science, London July: 50-52.
- McIntosh WC (1900) Notes from the Gatty Marine Laboratory, St. Andrews. No. 20. 1. *Annals and Magazine of Natural History* 5: 254-268. URL: <http://biodiversitylibrary.org/page/16333676>
- Meißner K (2005) Revision of the genus *Spiophanes* (Polychaeta, Spionidae); with new synonymies, new records and descriptions of new species. *Mitteilungen aus dem Museum für Naturkunde in Berlin – Zoologische Reihe* 81 (1): 3-65. <https://doi.org/10.1002/mmz.200310001>
- Meißner K, Blank M (2009) *Spiophanes norrisi* sp. nov. (Polychaeta: Spionidae)—a new species from the NE Pacific coast, separated from the *Spiophanes bombyx* complex based on both morphological and genetic studies. *Zootaxa* 2278: 1-2.
- Meißner K, Bick A, Bastrop R (2011) On the identity of *Spio filicornis* (O.F. Mueller, 1776) - with the designation of a neotype, and the description of two new species from the North East Atlantic Ocean based on morphological and genetic studies. *Zootaxa* 2815: 1-27.
- Méndez N, Cardell MJ (1996) Littoral annelid polychaetes inhabiting soft bottoms of the Barcelonès (Catalonia, NE Spain). *Miscellània Zoològica* 19 (1): 119-147. URL: <http://other.museocienciasjournals.cat/files/MZ-vol-19-1-1996-pp-119-147.pdf>
- Mikac B (2015) A sea of worms: polychaete checklist of the Adriatic Sea. *Zootaxa* 3943 (1): 1-17. <https://doi.org/10.11646/zootaxa.3943.1.1>
- Mikac B, Hutchings P (2017) One new species of *Pista* Malmgren, 1866 (Annelida: Terebellidae) and one new species of *Pistella* Hartmann-Schröder, 1996 (Annelida: Terebellidae) from the Adriatic Sea (Mediterranean). *Journal of the Marine Biological Association of the United Kingdom* 97: 943-953. <https://doi.org/10.1017/S0025315417000868>
- Mikac B, Musco L (2010) Faunal and biogeographic analysis of Syllidae (Polychaeta) from Rovinj (Croatia, northern Adriatic Sea). *Scientia Marina* 74 (2): 353-370. <https://doi.org/10.3989/scimar.2010.74n2353>
- Miura T (1980) Lumbrineridae (Annelides Polychetes) abyssaux recoltes au cours de campagnes du Centre Oceanologique de Bretagne dans l'Atlantique et la Mediterranee. *Bulletin du Museum National d'Histoire Naturelle, Paris* 2: 1019-1057.
- Molina-Acevedo IC, Carrera-Parra LF (2015) Reinstatement of three species of the *Marphysa sanguinea* complex (Polychaeta: Eunicidae) from the Grand Caribbean Region. *Zootaxa* 3925 (1): 37. <https://doi.org/10.11646/zootaxa.3925.1.3>
- Monro C (1939) Polychaeta. B.A.N.Z. Antarctic Research Expedition Reports 4: 87-156.
- Moreira J, Parapar J (2002) Redescription of *Sigambra tentaculata* and re-establishment of *S. parva* (Polychaeta, Pilargidae) based upon type material. *Cahiers de Biologie Marine* 43 (2): 99-109. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>

- Moreira J, Parapar J, Troncoso JS (2000) On the presence of *Prionospio pulchra* (Polychaeta: Spionidae) in the Atlantic Ocean. *Cahiers de Biologie Marine* 41 (3): 233-239. URL: <http://hdl.handle.net/2183/17865>
- Morri C, Bianchi CN, Cocito S, Peirano A, De Biase AM, Aliani S, Pansini M, Boyer M, Ferdeghini F, Pestarino M, Dando P (1999) Biodiversity of marine sessile epifauna at an Aegean island subject to hydrothermal activity: Milos, eastern Mediterranean Sea. *Marine Biology* 135 (4): 729-739. <https://doi.org/10.1007/s002270050674>
- Munari C, Bocchi N, Parrella P, Granata T, Moruzzi L, Massara F, Donati MD, Mistri M (2017) The occurrence of two morphologically similar *Chaetozone* (Annelida: Polychaeta: Cirratulidae) species from the Italian seas: *Chaetozone corona* Berkeley & Berkeley, 1941 and *C. carpenteri* McIntosh, 1911. *The European Zoological Journal* 84 (1): 541-553. <https://doi.org/10.1080/24750263.2017.1393843>
- Musco L, Giangrande A (2005) Mediterranean Syllidae (Annelida: Polychaeta) revisited: biogeography, diversity and species fidelity to environmental features. *Marine Ecology Progress Series* 304: 143-153. <https://doi.org/10.3354/meps304143>
- Mustaquim J (1986) Morphological variation in *Polydora ciliata* complex (Polychaeta: Annelida). *Zoological Journal of the Linnean Society* 86 (1): 75-88. <https://doi.org/10.1111/j.1096-3642.1986.tb01808.x>
- Mustaquim J (1988) Electrophoretic variation of isozymes in *Polydora ciliata* complex (Polychaeta: Spionidae). *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* 91 (1): 197-205. [https://doi.org/10.1016/0305-0491\(88\)90135-6](https://doi.org/10.1016/0305-0491(88)90135-6)
- NCMR (1989) Επισκόπηση της βενθικής πανίδας στο Κεντρικό Αιγαίο Πέλαγος (Κυκλάδες- Σαρωνικός). [A survey of the benthic fauna in the Central Aegean Sea (Cyclades-Saronikos)]. Technical Report. Zenetos A (Ed.). National Centre for Marine Research, Athens, 106 pp. [In Greek].
- NCMR (1992) Περιβαλλοντική μελέτη του Βόρειου Ευβοϊκού κόλπου. [Environmental study of the Northern Evoikos Gulf.]. Technical Report. Papathanasiou E (Ed.). National Centre for Marine Research, Athens, 243 pp. [In Greek].
- NCMR (1995) Μελέτη επιπτώσεων στις βενθικές βιοκοινωνίες του Κορινθιακού Κόλπου από τα κατάλοιπα επεξεργασίας βωξίτη. [Study of the impact of dumping bauxite mining residues on the benthic communities of Korinthiakos Gulf]. Technical Report. Zenetos A (Ed.). National Centre for Marine Research, Athens, 93 pp. [In Greek].
- NCMR (1997) Trawling impact on benthic ecosystems Final Report, DG-XIV Contract Number 95/14. Technical Report. Zenetos A (Ed.). National Centre for Marine Research, Athens, 112 pp.
- NCMR (1998) Μελέτη Περιβαλλοντικών Επιπτώσεων από Απόρριψη Σκωρίας στην περιοχή Λάρυμνας. [Study of the environmental impact of dumping coarse metalliferous waste in the area off Laryrnna]. Technical Report. Simboura N (Ed.). National Centre for Marine Research, Athens, 156 pp. [In Greek].
- NCMR (2000a) Το θαλάσσιο σύστημα του Αγίου Κοσμά. Ερευνητικό πρόγραμμα γεωφυσικών διασκοπήσεων, μετρήσεων περιβαλλοντικών παραμέτρων, ποιότητας υδάτων και παρακολούθησης της θαλάσσιας χλωρίδας και πανίδας για το Ολυμπιακό Κέντρο Ιστιοπλοΐας Αγίου Κοσμά. [The marine system of Agios Kosmas. Geophysical survey, measurements of environmental parameters water quality and monitoring of the marine flora and fauna for the Agios

- Kosmas Olympic Sailing Center.]. Technical report. Karageorgis A & Panayotidis P (Eds). National Centre for Marine Research, Athens.
- NCMR (2000b) Παρακολούθηση του οικοσυστήματος της λιμνοθάλασσας Πάπας Ακρωτηρίου Αράξου/Αχαΐας και προτάσεις διαχείρισης και προστασίας αυτής. [Monitoring of the ecosystem of Papas cape lagoon Araxos/Achaia and suggestions for the management and protection of the area]. Technical report. Pagou K (Ed.). National Centre for Marine Research, Athens, 190 pp. [In Greek].
  - Nebra A, Caiola N, Ibáñez C (2011) Community structure of benthic macroinvertebrates inhabiting a highly stratified Mediterranean estuary. *Scientia Marina* 75: 577-584. <https://doi.org/10.3989/scimar.2011.75n3577>
  - Neofitou N, Charizopoulos N, Vafidis D, Skordas K, Tziantziou L, Neofitou C (2014) Mussel farming impacts on trophic status and benthic community structure in Maliakos Gulf (Eastern Mediterranean). *Aquaculture International* 22 (2): 843-857. <https://doi.org/10.1007/s10499-013-9712-z>
  - Nicolaidou A, Papadopoulou K- (1989) Factors Affecting the Distribution and Diversity of Polychaetes in Amvrakikos Bay, Greece. *Marine Ecology* 10 (3): 193-204. <https://doi.org/10.1111/j.1439-0485.1989.tb00472.x>
  - Nicolaidou A, Pitta P (1986) Fauna associated with marine plants in a brackish lagoon of Amvrakikos bay. *Biologica Gallo-hellenica* 12: 141-147.
  - Nicolaidou A, Bourgoutzani F, Zenetos A (1990) Effects of pollution on the distribution of molluscs and polychaetes in Messologhi lagoon and adjacent areas. MAP Technical Reports Series FAO 40: 45-61.
  - Nicolaidou A, Papadopoulou KN, Simboura N (1986) Taxonomic and ecological study of the polychaete fauna of the N. Sporades. Aegean Sea. Rapport et Procès-Verbaux des Réunions de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée 30 (2): 20.
  - Nicolaidou A, Zenetos A, Pancucci M, Simboura N (1993) Comparing ecological effects of two different types of pollution using multivariate techniques. *Marine Ecology* 14: 113-128. <https://doi.org/10.1111/j.1439-0485.1993.tb00370.x>
  - Nogueira JMdM, ten Hove H (2000) On a new species of *Salmacina* Claparède, 1870 (Polychaeta: Serpulidae) from São Paulo state, Brazil. *Beaufortia* 50: 151-161.
  - Nogueira JMdM, Fitzhugh K, Hutchings P (2013) The continuing challenge of phylogenetic relationships in Terebelliformia (Annelida: Polychaeta). *Invertebrate Systematics* 27 (2): 186-23. <https://doi.org/10.1071/is12062>
  - Nogueira JMdM, San Martín G, Fukuda MV (2004) On some exogonines (Polychaeta, Syllidae, Exogoninae) from the northern coast of the State of Sao Paulo, southeastern Brazil. Results of BIOTA/FAPESP/Bentos Marinho Project. *Meiofauna Marina* 13: 45-77.
  - Norlinder E, Nygren A, Wiklund H, Pleijel F (2012) Phylogeny of scale-worms (Aphroditiformia, Annelida), assessed from 18SrRNA, 28SrRNA, 16SrRNA, mitochondrial cytochrome c oxidase subunit I (COI), and morphology. *Molecular Phylogenetics and Evolution* 65 (2): 490-500. <https://doi.org/10.1016/j.ympev.2012.07.002>
  - Núñez J (2004) Familia Nereididae Blainville, 1818. In: Viéitez JM, Alós C, Parapar J, Besteiro C, Moreira J, Núñez J, Laborda AJ, San Martín JG (Eds) Annelida, Polychaeta I. In: Ramos MA, Alba J, Bellés X, Gosálbez J, Guerra A, Macpherson E, Martín F, Serrano J, Templado J (Eds.) *Fauna Ibérica*. 25. Museo Nacional de Ciencias Naturales, CA, SAIC, Madrid, 293–389 pp.

- Núñez J, Brito MC (2002) Nereididae (Annelida, Polychaeta) colectados en las campañas "Fauna II, III, IV" (Proyecto "Fauna Ibérica") y catálogo de las especies conocidas para el ámbito ibérico. *Graellsia* 58 (1): 39-47. <https://doi.org/10.3989/graellsia.2002.v58.i1.265>
- Núñez J, Maggio Y, Riera R (2013) Dorvilleidos (Polychaeta, Dorvilleidae) recolectados durante el proyecto "Fauna Ibérica" y catálogo de las especies ibero-baleares. *Graellsia* 69 (2): 275-282. <https://doi.org/10.3989/graellsia.2013.v69.095>
- Núñez J, Barnich R, Brito MdC, Fiege D (2012) Familias Aphroditidae, Polynoidae, Acoetidae, Sigalionidae y Pholoidae . In: Ramos MA et al. *Annelida Polychaeta IV*. Parapar J, Moreira J, Núñez J, Barnich R, Brito M del C, Fiege D, Capaccioni-Azzati R, El-Haddad M. *Fauna Ibérica*. 41. Museo Nacional de Ciencias Naturales. CSIC, Madrid, 87-257 pp.
- Núñez J, Barnich R, Santos L, Maggio Y (2011) Poliquetos Escamosos (Annelida, Polychaeta) Colectados en Las Campañas "Fauna II, III, IV" (Proyecto "Fauna Ibérica") y Catálogo de Las Especies Conocidas Para el Ámbito Íbero-Balear. *Graellsia* 67 (2): 187-197. <https://doi.org/10.3989/graellsia.2011.v67.045>
- Nygren A (2004) Revision of Autolytinae (Syllidae: Polychaeta). *Zootaxa* 680: 1-314.
- Nygren A (2014) Cryptic polychaete diversity: a review. *Zoologica Scripta* 43 (2): 172-183. <https://doi.org/10.1111/zsc.12044>
- Nygren A, Pleijel F (2010) Redescription of *Imajimaea draculai* —a rare syllid polychaete associated with the sea pen *Funiculina quadrangularis* . *Journal of the Marine Biological Association of the United Kingdom* 90 (7): 1441-1448. <https://doi.org/10.1017/S0025315409991536>
- Nygren A, Pleijel F (2011) From one to ten in a single stroke – resolving the European *Eumida sanguinea* (Phyllodocidae, Annelida) species complex. *Molecular phylogenetics and evolution* 58 (1): 132-41. <https://doi.org/10.1016/j.ympev.2010.10.010>
- Nygren A, Pleijel F (2015) Ringmaskar: Havsborstmaskar, Annelida: Polychaeta . Art-Databanken, SLU, Uppsala, 346 pp. [In Swedish]. URL: <http://www.artdatabanken.se/publikationer/bokverket-nationalnyckeln/bestall-nationalnycklar/bestamningsnyckel-havsborstmaskar/>
- Nygren A, Eklöf J, Pleijel F (2010a) Cryptic species of *Notophyllum* (Polychaeta: Phyllodocidae) in Scandinavian waters. *Organisms Diversity & Evolution* 10 (3): 193-204. <https://doi.org/10.1007/s13127-010-0014-2>
- Nygren A, Norlinder E, Panova M, Pleijel F (2010b) Colour polymorphism in the polychaete *Harmothoe imbricata* (Linnaeus, 1767). *Marine Biology Research* 7 (1): 54-62. <https://doi.org/10.1080/17451001003713555>
- Occhipinti-Ambrogi A, Marchini A, Cantone G, Castelli A, Chimenz C, Cormaci M, Frogliia C, Furnari G, Gambi MC, Giaccone G, Giangrande A, Gravili C, Mastrototaro F, Mazziotti C, Orsi-Relini L, Piraino S (2011) Alien species along the Italian coasts: An overview. *Biological Invasions* 13 (1): 215-237. <https://doi.org/10.1007/s10530-010-9803-y>
- O'Connor BS (1987) The Glyceridae (Polychaeta) of the North Atlantic and Mediterranean, with descriptions of two new species. *Journal of Natural History* 21 (1): 167-189. <https://doi.org/10.1080/00222938700770051>
- Olivier F, Grant C, San Martín G, Archambault P, McKindsey C (2012) Syllidae (Annelida: Polychaeta: Phyllodocida) from the Chausey Archipelago (English Channel,

- France), with a description of two new species of the Exogoninae *Prosphaerosyllis*. Marine Biodiversity 42 (1): 55-63. <https://doi.org/10.1007/s12526-011-0092-1>
- Olu-Le Roy K, Sibuet M, Fiala-Médioni A, Gofas S, Salas C, Mariotti A, Foucher JP, Woodside J (2004) Cold seep communities in the deep eastern Mediterranean Sea: Composition, symbiosis and spatial distribution on mud volcanoes. Deep-Sea Research Part I: Oceanographic Research Papers 51 (12): 1915-1936. <https://doi.org/10.1016/j.dsr.2004.07.004>
  - Oug E (2011) Guide to identification of Lumbrineridae (Polychaeta) in Norwegian and adjacent waters. [www.polychaeta.no](http://www.polychaeta.no). Accessed on: 2016-10-20.
  - Panagopoulos D, Nikolaidou A (1989) A population of *Desdemonia ornata* Banse (Polychaeta, Sabelliidea) settled in a fully marine habitat of the Mediterranean. Oebalia 16: 35-39.
  - Pancucci-Papadopoulou A, Simboura N, Zenetos A, Thessalou-Legaki M, Nicolaidou A (1999) Benthic invertebrate communities of NW Rodos (Rhodes) island (SE Aegean Sea) as related to hydrological regime and geographical location. Israel Journal of Zoology 45: 371-393. URL: <http://www.tandfonline.com/doi/abs/10.1080/00212210.1999.10689006>
  - Pancucci-Papadopoulou MA, Zenetos A, Corsini-Foka M, Politou CY (2005) Update of marine alien species in Hellenic waters. Mediterranean Marine Science 6 (2): 147-158. <https://doi.org/10.12681/mms.188>
  - Papadopoulos D (1986) The impact of dumping coarse metalliferous waste on the benthos of the Northern Evvoikos Gulf. MSc thesis. National and Kapodistrian University of Athens, Athens, 99 pp. [In Greek].
  - Papageorgiou N, Arvanitidis C, Eleftheriou A (2006) Multicausal environmental severity: A flexible framework for microtidal sandy beaches and the role of polychaetes as an indicator taxon. Estuarine, Coastal and Shelf Science 70 (4): 643-653. <https://doi.org/10.1016/j.ecss.2005.11.033>
  - Papazacharias A (1991) Οικολογική μελέτη της μακροπανίδας του κινητού υποστρώματος της υποπαραλιακής και περιπαραλιακής ζώνης του κόλπου της Καβάλας. [Ecological study of the soft substratum macrofauna in the infralittoral and circalittoral zone of the Gulf of Kavala.]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 204 pp. [In Greek].
  - Parapar J (2006) The genera *Myriochele* and *Myrioglobula* (Polychaeta, Oweniidae) in Icelandic waters with the revision of type material of *Myriochele heeri* Malmgren, 1867, and the description of a new species. Journal of Natural History 40: 523-547. <https://doi.org/10.1080/00222930600711758>
  - Parapar J (2012) Familia Opheliidae Malmgren, 1867. In: Parapar J, Alós C, Núñez J, Moreira J, López E, Aguirrezabalaga F, Besteiro C, Martínez A (Eds) Annelida Polychaeta III, Ramos MA et al. (Eds) Fauna Ibérica. 3. Museo Nacional de Ciencias Naturales, CSIC, Madrid, 284-332 pp. [In Spanish].
  - Parapar J, Harto I (2001) Sobre la presencia en aguas ibéricas y variabilidad morfológica en *Eunice roussaei* Quatrefages, 1866 (Polychaeta, Eunicidae). Boletín de la Real Sociedad Española de Historia Natural. Sección Biológica 96 (34): 165-17.
  - Parapar J, Hutchings PA (2014) Redescription of *Terebellides stroemii* (Polychaeta, Trichobranchidae) and designation of a neotype. Journal of the Marine Biological Association of the United Kingdom 95 (2): 323-337. <https://doi.org/10.1017/S0025315414000903>

- Parapar J, Moreira J (2015) Familia Glyceridae . In: Parapar J, Moreira J, Barnich R, Brito MdC, Fiege D, Capaccioni-Azzati R, El-Haddad M (Eds) Annelida Polychaeta IV, Fauna Iberica. 41. Museo Nacional de Ciencias Naturales Consejo Superior de Investigaciones Científicas, Madrid, 416 pp.
- Parapar J, Besteiro C, Ugorri V (1992) Nuevas aportaciones al conocimiento de los anélidos poliquetos en el litoral gallego (N.O. Península Ibérica). Nova Acta Científica Compostelana (Biología) 3: 109-123. URL: <https://minerva.usc.es/xmlui/handle/10347/6209>
- Parapar J, Besteiro C, Ugorri V (1993) Primera cita de *Ampharete finmarchica* (Sars, 1865) (Polychaeta: Ampharetidae) en el litoral de la península Ibérica. Boletín Instituto Español de Oceanografía 9: 367-372.
- Parapar J, Mikac B, Fiege D (2013) Diversity of the genus *Terebellides* (Polychaeta: Trichobranchidae) in the Adriatic Sea with the description of a new species. Zootaxa 3691 (3): 333. <https://doi.org/10.11646/zootaxa.3691.3.3>
- Parapar J, San Martín G, Ugorri V, Besteiro C (1993) Anélidos Poliquetos Mesopsámnicos Sublitorales de la Costa de Ceuta (España). Cahiers de Biologie Marine 34 (4): 363-381. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- Paresque K, Fukuda MV, Nogueira JMdm (2016) *Branchiosyllis*, *Haplosyllis*, *Opisthosyllis* and *Trypanosyllis* (Annelida: Syllidae) from Brazil, with the description of two new species. PloSone 11 (5): e0153442. <https://doi.org/10.1371/journal.pone.0153442>
- Paterson GJ, Glover A, Barrio Froján CS, Whitaker A, Budaeva N, Chimonides J, Doner S (2009) A census of abyssal polychaetes. Deep Sea Research Part II: Topical Studies in Oceanography 56: 1739-1746. <https://doi.org/10.1016/j.dsr2.2009.05.018>
- Paxton H, Åkesson B (2010) The *Ophryotrocha labronica* group (Annelida: Dorvilleidae) — with the description of seven new species. Zootaxa 2713: 1-24. URL: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.359.8889&rep=rep1&type=pdf>
- Paxton H, Arias A (2016) The identity of *Paradiopatra bihanica* (Annelida: Onuphidae) and reinstatement of *P. calliopae*. Zootaxa 4132 (2): 283-286. <https://doi.org/10.15468/HU6OVZ>
- Pérès J (1954) Contribution à l'étude des Annélides polychètes de la Méditerranée occidentale. Travaux de la Station marine d'Endoume 13 (8): 83-154.
- Pérès J (1959) Contribution à la connaissance des polychètes benthiques des profondeurs moyennes de la Méditerranée. Travaux de la Station Marine d'Endoume 26 (16): 103-135.
- Perkins T (1981) Syllidae, principally from Florida, with descriptions of a new genus and twenty-one new species. Proceedings of the Biological Society of Washington 93 (4): 1080-1172. URL: <http://www.vliz.be/imisdocs/publications/255356.pdf>
- Perkins T (1984) Revision of *Demonax* Kinberg, *Hypsicomus* Grube, and *Notaulax* Tauber, with a review of *Megalomma* Johansson from Florida (Polychaeta: Sabellidae). Proceedings of the Biological Society of Washington 97: 285-368. URL: <https://archive.org/details/biostor-75392>
- Perkins TH (1979) Lumbrineridae, Arabellidae, and Dorvilleidae (Polychaeta), principally from Florida, with descriptions of six new species. Proceedings of The Biological Society of Washington 92: 415-465.
- Petersen ME (1991) A review of asexual reproduction in the Cirratulidae (Annelida: Polychaeta), with redescription of *Cirratulus gayheadius* (Hartman, 1965), new



combination and emendation or reinstatement of some cirratulid genera. *Bulletin of Marine Science* 48 (2): 592.

- Petersen ME (1999) Reproduction and development in Cirratulidae (Annelida: Polychaeta). *Hydrobiologia* 402: 107-128. <https://doi.org/10.1023/A:100373640>
- Pettibone M (1963) Marine polychaete worms of the New England region. I. Aphroditidae through Trochochaetidae. *Bulletin of the United States National Museum* 227: 1-356. <https://doi.org/10.5479/si.03629236.227.1>
- Pettibone M (1969) Review of some species referred to *Scalisetosus* McIntosh (Polychaeta, Polynoidae). *Proceedings of the Biological Society of Washington* 82: 1-30. URL: <https://repository.si.edu/handle/10088/3412>
- Pettibone M (1976) Revision of the genus *Macellicephalo* McIntosh and the Subfamily Macellicephalinae Hartmann-Schröder (Polychaeta: Polynoidae). *Smithsonian Contributions to Zoology* 229: 1-71. <https://doi.org/10.5479/si.00810282.229>
- Pettibone M (1986) Additions to the family Eulepethidae Chamberlin (Polychaeta: Aphroditacea). *Smithsonian Contributions to Zoology* 441: 1-51. <https://doi.org/10.5479/si.00810282.441>
- Pettibone M (1996) Revision of the scaleworm genera *Acholoe* Claparède, *Arctonoella* Buzhinskaja, and *Intoshella* Darboux (Polychaeta: Polynoidae) with the erection of the new subfamily Acholoinae. *Proceedings of the Biological Society of Washington* 109: 629-644. URL: <https://www.biodiversitylibrary.org/part/46306>
- Pettibone MH (1963) Revision of some genera of polychaete worms of the family Spionidae, including the description of a new species of *Scolecopsis*. *Proceedings of the Biological Society of Washington* 76: 89-10.
- Pettibone MH (1989) Revision of the aphroditoid polychaetes of the family Acoetidae Kinberg (=Polyodontidae Augener) and reestablishment of *Acoetes* Audouin and Milne-Edwards, 1832, and *Euarche* Ehlers, 1887. *Smithsonian Contributions to Zoology* 464: 1-138. <https://doi.org/10.5479/si.00810282.464>
- Picard J (1965) Recherches qualitatives sur les biocénoses marines des substrats meubles dragables de la région marseillaise. *Recueil des Travaux de la Station Marine d'Endoume* (52)1-60.
- Pleijel F (1988) *Phyllodoce* (Polychaeta, Phyllodocidae) from northern Europe. *Zoologica Scripta* 17 (2): 141-153. <https://doi.org/10.1111/j.1463-6409.1988.tb00091.x>
- Pleijel F (1991) Phylogeny and classification of the Phyllodocidae (Polychaeta). *Zoologica Scripta* 20 (3): 225-261. <https://doi.org/10.1111/j.1463-6409.1991.tb00289.x>
- Pleijel F (1993) Polychaeta, Phyllodocidae. *Marine Invertebrates of Scandinavia*. 8. Scandinavian University Press, Oslo, 159 pp.
- Pleijel F (1998) Phylogeny and classification of Hesionidae (Polychaeta). *Zoologica Scripta* 27 (2): 89-163. <https://doi.org/10.1111/j.1463-6409.1998.tb00433.x>
- Pleijel F (2004) A revision of *Hesiospina* (Psamathini, Hesionidae, Polychaeta). *Journal of Natural History* 38 (20): 2547-2566. <https://doi.org/10.1080/00222930310001647433>
- Pleijel F (2005) Book Review. *Journal of Natural History* 39 (40): 3543-3544. <https://doi.org/10.1080/00222930500465372>
- Pleijel F, Dales RP (1991) Polychaetes: British Phyllodocoideans, Typhloscolecoideans and Tomopteroideans. *Synopses of the British Fauna (New Series)* 45: 1-202.
- Pleijel F, Rouse G, Nygren A (2012) A revision of *Nereimyra* (Psamathini, Hesionidae, Aciculata, Annelida). *Zoological Journal of the Linnean Society* 164 (1): 36-51. <https://doi.org/10.1111/j.1096-3642.2011.00756.x>

- Probert PK, Glasby CJ, Grove SL, Paavo BL (2009) Bathyal polychaete assemblages in the region of the Subtropical Front, Chatham Rise, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 43 (5): 1121-1135. <https://doi.org/10.1080/00288330.2009.9626535>
- Pruvot G, Racovitza EG (1895) Matériaux pour la faune des annélides de Banyuls. *Archives de Zoologie Experimentale et Generale* 3: 378-385.
- Quatrefages Ad (1866) Histoire naturelle des Annelés marins et d'eau douce. Annelides et Géphyriens. 1. Librairie Encyclopédique de Roret, Paris, 588 pp. <https://doi.org/10.5962/bhl.title.122818>
- Radashevsky V, Hsieh HL (2000) *Polydora* (Polychaeta: Spionidae) species from Taiwan. *Zoological Studies* 39 (3): 203-21.
- Radashevsky V, Migotto A (2016) First report of the polychaete *Polydora hoplura* (Annelida: Spionidae) from North and South America and Asian Pacific. *Marine Biodiversity* 47 (3): 859-868. <https://doi.org/10.1007/s12526-016-0515-0>
- Radashevsky V, Nogueira J (2003) Life history, morphology and distribution of *Dipolydora armata* (Polychaeta: Spionidae). *Journal of the Marine Biological Association of the United Kingdom* 83 (2): 375-384. <https://doi.org/10.1017/s0025315403007227h>
- Radashevsky V, Pankova V (2006) The morphology of two sibling sympatric *Polydora* species (Polychaeta: Spionidae) from the Sea of Japan. *Journal of the Marine Biological Association of the United Kingdom* 86 (02): 245. <https://doi.org/10.1017/s0025315406013099>
- Radashevsky V, Pankova V, Neretna TV, Stupnikova AN, Tzetlin AB (2016) Molecular analysis of the *Pygospio elegans* group of species (Annelida: Spionidae). *Zootaxa* 4083 (2): 239. <https://doi.org/10.11646/zootaxa.4083.2.4>
- Radashevsky VI (2015) Spionidae (Annelida) from Lizard Island, Great Barrier Reef, Australia: the genera *Aonides*, *Dipolydora*, *Polydorella*, *Prionospio*, *Pseudopolydora*, *Rhynchospio* and *Tripolydora*. *Zootaxa* 4019 (1): 635-694. <https://doi.org/10.11646/zootaxa.4019.1.22>
- Radashevsky VI, Selifonova ZP (2013) Records of *Polydora cornuta* and *Streblospio gynobranchiata* (Annelida, Spionidae) from the Black Sea. *Mediterranean Marine Science* 14 (2): 261-26. <https://doi.org/10.12681/mms.415>
- Radashevsky VI, Simboura N (2013) First record of *Dipolydora blakei* (Annelida: Spionidae) from Europe: Greece, Mediterranean Sea. *Mediterranean Marine Science* 14 (1): 19-23. <https://doi.org/10.12681/mms.322>
- Radić I (2009) Utjecaj tunogjilišta na faunu mnogočetinaša u biocenozi morske cvjetnice *Posidonia oceanica* u Jadranu. [Impact of tuna farming facilities on polychaetes in the *Posidonia oceanica* biocoenosis in the Adriatic Sea.]. University of Zagreb, Zagreb, 170 pp.
- Rainer SF (1989) Redescription of *Nephtys assimilis* and *N. kersivalensis* (Polychaeta, Phyllodocida) and a Key to *Nephtys* from Northern Europe. *Journal of the Marine Biological Association of the United Kingdom* 69 (4): 875-890. <https://doi.org/10.1017/S0025315400032227>
- Rainer SF (1990) The genus *Nephtys* (Polychaeta: Phyllodocida) in northern Europe: redescription of *N. hystricis* and *N. incisa*. *Journal of Natural History* 24: 361-372. <https://doi.org/10.1080/00222939000770271>
- Rainer SF (1991) The genus *Nephtys* (Polychaeta: Phyllodocida) of northern Europe: a review of species, including the description of *N. pulchra* sp. n. and a key to the

Nephtyidae . Helgoländer Meeresuntersuchungen 45: 65-96. <https://doi.org/10.1007/BF02365637>

- Ramfos A, Somarakis S, Koutsikopoulos C, Fragopoulou N (2005) Summer mesozooplankton distribution in coastal waters of central Greece (eastern Mediterranean). I. Hydrology and group composition. Journal of the Marine Biological Association of the United Kingdom 85 (4): 755-764. <https://doi.org/10.1017/S0025315405011665>
- Ramfos A, Isari S, Somarakis S, Georgopoulos D, Koutsikopoulos C, Fragopoulou N (2006) Mesozooplankton community structure in offshore and coastal waters of the Ionian Sea (eastern Mediterranean) during mixed and stratified conditions. Marine Biology 150 (1): 29-44. <https://doi.org/10.1007/s00227-006-0326-0>
- Ramos J (1976) Lumbrineridae (Polychètes errantes) de Méditerranée. Annales de l'Institut Océanographique Paris 52: 103-137.
- Ramos JM (1976a) *Aonides oxycephala* (Sars, 1862) Remarques taxonomiques. Vie Milieu 26: 11-20.
- Ramos JM (1976b) *Haploscoloplos kerguelensis* McIntosh, 1885 - Nouvel Orbiniidae en Méditerranée Occidentale. Vie et milieu 26 (1): 1-9.
- Rathke H (1843) Beiträge zur Fauna Norwegens. Nova Acta der Kaiserlichen Leopold-Carolin Deutschen Akademie der Naturforscher, Halle 20: 1-264. URL: <http://www.biodiversitylibrary.org/item/43961>
- Ravara A, Cunha MR, Pleijel F (2010) Nephtyidae (Annelida, Polychaeta) from southern Europe. Zootaxa 2682: 1-68. URL: [http://ria.ua.pt/bitstream/10773/6571/1/41\\_Ravara\\_et\\_al\\_2010b.pdf](http://ria.ua.pt/bitstream/10773/6571/1/41_Ravara_et_al_2010b.pdf)
- Ravara A, Ramos D, Teixeira ML, Costa F, Cunha M (2017) Taxonomy, distribution and ecology of the order Phyllodocida (Annelida, Polychaeta) in deep-sea habitats around the Iberian margin. Deep Sea Research Part II: Topical Studies in Oceanography 137: 207-231. <https://doi.org/10.1016/j.dsr2.2016.08.008>
- Read GB, Hove HA, Sun Y, Kupriyanova EK (2017) *Hydroides* Gunnerus, 1768 (Annelida, Serpulidae) is feminine: a nomenclatural checklist of updated names. ZooKeys 52: 1-52. <https://doi.org/10.3897/zookeys.642.10443>
- Reish D, Anderson F, Horn K, Hardege J (2014) Molecular phylogenetics of the *Neanthes acuminata* (Annelida: Nereididae) species complex. Memoirs of Museum Victoria 71: 271-278. <https://doi.org/10.24199/j.mmv.2014.71.20>
- Rice S, Karl S, Rice K (2008) The *Polydora cornuta* complex (Annelida: Polychaeta) contains populations that are reproductively isolated and genetically distinct. Invertebrate Biology 127 (1): 45-64. <https://doi.org/10.1111/j.1744-7410.2007.00104.x>
- Riser NW (1991) An evaluation of taxonomic characters in the genus *Sphaerosyllis* (Polychaeta: Syllidae). Ophelia Supplement 5: 209-218.
- Riser NW (2000) *Podarke aberrans* Webster & Benedict, 1887 - resolution, with description of two new species in the genus *Microphthalmus* (Annelida: Polychaeta). Proceedings of the Biological Society of Washington 113 (2): 514-225. URL: <http://biodiversitylibrary.org/page/35460342>
- Rizzo AE, Salazar-Vallejo SI (2014) Hesionidae Grube, 1850 (Annelida: Polychaeta) from South-Southeastern Brazil, with descriptions of four new species. Zootaxa 3856 (2): 267-291. <https://doi.org/10.11646/zootaxa.3856.2.7>
- Rouse G, Pleijel F (2001) Polychaetes. Oxford University Press, Oxford, 354 pp.

- Rullier F (1963) Les Annélides polychètes du Bosphore, de la Mer de Marmara et de la Mer Noire, en relation avec celles de la Méditerranée. Rapport et Procès-Verbaux des Réunions de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée 17: 161-260.
- Salazar-Vallejo S (2012) Revision of *Flabelligera* Sars, 1829 (Polychaeta: Flabelligeridae). *Zootaxa* 3203: 1-64.
- Salazar-Vallejo S (2013) Revision of *Therochaeta* Chamberlin, 1919 (Polychaeta: Flabelligeridae). *Zoosystema* 35 (2): 227-263. <https://doi.org/10.5252/z2013n2a7>
- Salazar-Vallejo S (2017) Revision of *Brada* Stimpson, 1853, and *Bradabyssa* Hartman, 1967 (Annelida, Flabelligeridae). *Zootaxa* 4343 (1): 1-98. <https://doi.org/10.11646/zootaxa.4343.1.1>
- Salazar-Vallejo SI (2011) Revision of *Stylarioides* delle Chiaje, 1831 (Annelida: Flabelligeridae). *Italian Journal of Zoology* 78: 163-200. <https://doi.org/10.1080/11250003.2011.606985>
- Salazar-Vallejo SI (2014) Revision of *Pherusa* Oken, 1807 (Polychaeta: Flabelligeridae). *Zootaxa* 3886: 1-61. <https://doi.org/10.11646/zootaxa.3886.1.1>
- Salazar-Vallejo SI, Gillet P, Surugiu V (2017) How false is *Nereis falsa*? (Annelida, Phyllodocida, Nereididae). *Revista de Biología Tropical* 65: 847-857. URL: <https://revistas.ucr.ac.cr/index.php/rbt/article/view/26635/29967>
- Sanfilippo R (2003) Climatic response in the genus *Placostegus* Philippi, 1844 from Plio-Pleistocene to Recent Mediterranean, with description of a new species (Polychaeta, Serpulidae). *Bollettino della Società Paleontologica Italiana* 42: 171-178.
- Sanfilippo R, Rosso A, Guido A, Gerovasileiou V (2017) Serpulid communities from two marine caves in the Aegean Sea, Eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* 97: 1059-1068. <https://doi.org/10.1017/S0025315417000297>
- San Martín G (1984) Estudio biogeográfico, faunístico y sistemático de los Poliquetos de la familia Sílidos (Syllidae: Polychaeta) en Baleares. Ediciones de la Universidad Complutense de Madrid, Madrid 187: 1-529.
- San Martín G (1991) *Grubeosyllis* and *Exogone* (Exogoninae, Syllidae, Polychaeta) from Cuba, the Gulf of Mexico, Florida and Puerto Rico, with a revision of *Exogone*. *Bulletin of Marine Science* 49: 715-740. URL: <http://www.ingentaconnect.com/contentone/umrsmas/bullmar/1991/00000049/00000003/art00004>
- San Martín G (2003) Annelida Polychaeta II: Syllidae. In: Ramos MA, et al. (Eds). *Fauna Ibérica* 21, Museo Nacional de Ciencias Naturales, CA, SAIC, Madrid: 544 pp.
- San Martín G, Lopez E (2000) Three new species of *Syllis* (Syllidae: Polychaeta) from Iberian coasts. *Cahiers de Biologie Marine* 41 (4): 425-433. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
- San Martín G, Worsfold T (2015) Guide and keys for the identification of Syllidae (Annelida, Phyllodocida) from the British Isles (reported and expected species). *ZooKeys* 488: 1-29. <https://doi.org/10.3897/zookeys.488.9061>
- San Martín G, Álvarez-Campos P, Hutchings P (2017) The genus *Syllis* Savigny in Lamarck, 1818 (Annelida: Syllidae: Syllinae) from Australia (second part): four new species and re-description of twelve previously described species. *Zootaxa* 4237 (2): 201-243. <https://doi.org/10.11646/zootaxa.4052.3.2>
- San Martín G, Hutchings P, Aguado MT (2008) Syllinae (Polychaeta: Syllidae) from Australia. Part 1. Genera *Branchiosyllis*, *Eurysyllis*, *Karroonsyllis*, *Parasphaerosyllis*,

- Plakosyllis*, *Rhopalosyllis*, *Tetrapalpia* n.gen., and *Xenosyllis*. Records of the Australian Museum 60 (2): 119-160. <https://doi.org/10.3853/j.0067-1975.60.2008.1494>
- San Martín G, López E, Aguado MT (2009) Revision of the genus *Pionosyllis* (Polychaeta: Syllidae: Eusyllinae), with a cladistic analysis, and the description of five new genera and two new species. Journal of the Marine Biological Association of the United Kingdom 89 (7): 1455-1498. <https://doi.org/10.1017/s0025315409003099>
  - San Martín G, Viéitez JM, Campoy A (1981a) Contribucion al estudio de la fauna de anelidos poliquetos de las costas españolas: Poliquetos errantes recolectados en la Bahía de Palma de Mallorca. Boletín Instituto Espanol de Oceanografía 6: 62-87.
  - San Martín G, Acero M, Contonente M, Gomez J (1981b) Una coleccion de anélidos poliquetos de las costas mediterraneas andaluzas. Actas II Simposio Ibérico de Estudios del Bentos Marino 3: 171-182.
  - Santos CG, Mackie ASY (2008) New Species of *Poecilochaetus* Claparède, 1875 (Polychaeta, Spionidae, Poecilochaetidae) from Parangaguá Bay, Southeastern Brazil. Zootaxa 1790: 53-68. URL: <http://mapress.com/zootaxa/2008/f/zt01790p068.pdf>
  - Sardá R (1991) Polychaete Communities Related to Plant Covering in the Médiolittoral and Infralittoral Zones of the Balearic Islands (Western Mediterranean). Marine Ecology 12 (4): 341-360. <https://doi.org/10.1111/j.1439-0485.1991.tb00263.x>
  - Sardá R, Pinedo S, Martín D (1999) Seasonal dynamics of macroinfaunal key species inhabiting shallow soft-bottoms in the Bay of Blanes (NW Mediterranean). Acta Oecologica 20 (4): 315-326. [https://doi.org/10.1016/s1146-609x\(99\)00135-6](https://doi.org/10.1016/s1146-609x(99)00135-6)
  - Sato-Okoshi W, Abe H, Nishitani G, Simon C (2016) And then there was one: *Polydora uncinata* and *Polydora hoplura* (Annelida: Spionidae), the problematic polydorid pest species represent a single species. Journal of the Marine Biological Association of the United Kingdom 1-10. <https://doi.org/10.1017/s002531541600093x>
  - Savigny J (1818) Classe Neuvième. Les Annélides. In: Lamarck JBd (Ed.) Histoire Naturelle des animaux sans vertèbres, presentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent... Tome 5. 5. Déterville, Pari, 274–374 pp. [In French]. URL: <https://biodiversitylibrary.org/page/12887111>
  - Scaps P, Rouabah A, Leprêtre A (2000) Morphological and biochemical evidence that *Perinereis cultrifera* (Polychaeta: Nereididae) is a complex of species. Journal of the Marine Biological Association of the United Kingdom 80 (4): 735-736. <https://doi.org/10.1017/S0025315400002587>
  - Schimmenti E, Lo Brutto S, Badalamenti F, Giangrande A, Mikac B, Musco L (2015) DNA Barcoding to solve the tricky case of co-occurring *Sabellaria* (Annelida) species in the Mediterranean Sea. Biologia Marina Mediterranea 22: 109-110. URL: <https://iris.unipa.it/retrieve/handle/10447/237810/442301/2015>
  - Schmidt H, Westheide W (1999) Genetic relationships (RaPD-PCR) between geographically separated populations of the "cosmopolitan" interstitial Polychaete *Hesionides gohari* (Hesionidae) and the evolutionary origin of the freshwater species *Hesionides riegerorum*. Biological Bulletin 216: 216-226. <https://doi.org/10.2307/1542567>
  - Schmidt H, Westheide W (2000) Are the meiofaunal polychaetes *Hesionides arenaria* and *Stygocapitella subterranea* true cosmopolitan species? - Results of RAPD-PCR investigations. Zoologica Scripta 29 (1): 17-27. <https://doi.org/10.1046/j.1463-6409.2000.00026.x>

- Seixas VC, Zanol J, Magalhães W, Paiva PC (2017) Genetic diversity of *Timarete punctata* (Annelida: Cirratulidae): Detection of pseudo-cryptic species and a potential biological invader. *Estuarine, Coastal and Shelf Science* 197: 214-220. <https://doi.org/10.1016/j.ecss.2017.08.039>
- Sendall K, Salazar-Vallejo S (2013) Revision of *Sternaspis* Otto, 1821 (Polychaeta, Sternaspidae). *ZooKeys* 74: 1-74. <https://doi.org/10.3897/zookeys.287.4438>
- Serrano-Samaniego L (2012) Distribution of soft-bottom polychaete assemblages at different scales in shallow waters of the Northern Mediterranean Spanish coast. PhD thesis. Universitat Politècnica de Catalunya, 172 pp. URL: <https://upcommons.upc.edu/bitstream/handle/2117/94672/TLSS1de1.pdf>
- Sigvaldadóttir E (2002) Polychaetes of the genera *Prionospio* and *Aurospio* (Spionidae, Polychaeta) from Icelandic waters. *Sarsia* 87 (3): 207-215. <https://doi.org/10.1080/00364820260294842>
- Sigvaldadóttir E, Mackie AY (1993) *Prionospio steenstrupi*, *P. fallax* and *P. dubia* (Polychaeta, Spionidae): Re-evaluation of identity and status. *Sarsia* 78: 203-219. <https://doi.org/10.1080/00364827.1993.10413535>
- Sikorski A (2003) *Laonice* (Polychaeta, Spionidae) in the Arctic and the North Atlantic. *Sarsia* 88 (5): 316-345. <https://doi.org/10.1080/00364820310002551>
- Simbourn N (1987) Study of the distribution and variations of the benthic polychaete fauna in three regions of the Aegean Sea and especially in the Island of Rhodes. MSc thesis. National and Kapodistrian University of Athens, Athens, 111 pp. [In Greek].
- Simbourn N (1990) *Fabricia filamentosa* Day, 1963 (Polychaeta, Sabellidae, Fabriciinae) a lessepsian migrant in the Mediterranean Sea. *Oebalia* 16: 129-133.
- Simbourn N (1996) Θαλάσσιοι μακροβενθικοί Πολύχαιτοι (Annelida Polychaeta) της Ελλάδας: Συστηματική, Οικολογία, Ζωογεωγραφία. [Marine macrobenthic Polychaetes (Annelida, Polychaeta) of Greece: Taxonomy, Ecology, Zoogeography]. PhD thesis. National and Kapodistrian University of Athens, Athens, 241 pp. [In Greek]. <https://doi.org/10.12681/eadd/5952>
- Simbourn N (2008) First occurrence of the Indo-Pacific polychaete species *Glycinde bonhourei* Gravier, 1904 in the Hellenic seas (Northern Evvoikos Gulf). *Mediterranean Marine Science* 9: 173-176. <https://doi.org/10.12681/mms.148>
- Simbourn N (2011) An overlooked alien species present on the coasts of Greece (Eastern Mediterranean): The polychaete *Polycirrus twisti* Potts (Polychaeta: Terebellidae). *Mediterranean Marine Science* 12 (1): 239-246. <https://doi.org/10.12681/mms.62>
- Simbourn N, Nicolaidou A (2001) The Polychaetes (Annelida, Polychaeta) of Greece: Checklist, Distribution and Ecological Characteristics. *Monographs on Marine Sciences* 4: 1-115. URL: <http://epublishing.ekt.gr/en/8735>
- Simbourn N, Zenetos A (2005) Increasing Polychaete diversity as a consequence of increasing research effort in Greek waters: new records and exotic species. *Mediterranean Marine Science* 6: 75-88. <https://doi.org/10.12681/mms.194>
- Simbourn N, Kurt Şahin G, Panagoulia A, Katsiaras N (2010) Four new alien species on the coasts of Greece (Eastern Mediterranean). *Mediterranean Marine Science* 11: 341-352. <https://doi.org/10.12681/mms.81>
- Simbourn N, Reizopoulou S, Arvanitidis C, Basset A (2007) New information on the biogeographical and ecological distribution of *Pionosyllis anophthalma* Capaccioni and

- San Martín, 1989 (Polychaeta, Syllidae). *Transitional Waters Bulletin* 1 (4): 53-58. <https://doi.org/10.1285/i1825229Xv1n4p53>
- Simboura N, Sigala K, Voutsinas E, Kalkan E (2008) First occurrence of the invasive alien species *Polydora cornuta* Bosc, 1802 (Polychaeta: Spionidae) on the coast of Greece (Elefsis Bay; Aegean Sea). *Mediterranean Marine Science* 9 (2): 119-124. <https://doi.org/10.12681/mms.138>
  - Simboura N, Zenetos A, Panayotidis P, Makra A (1995a) Changes in benthic community structure along an environmental pollution gradient. *Marine Pollution Bulletin* 30 (7): 470-474. [https://doi.org/10.1016/0025-326X\(95\)00237-H](https://doi.org/10.1016/0025-326X(95)00237-H)
  - Simboura N, Zenetos A, Thessalou-Legaki M, Pancucci M-, Nicolaidou A (1995b) Benthic Communities of the Infralittoral in the N. Sporades (Aegean Sea): a Variety of Biotopes Encountered and Analysed. *Marine Ecology* 16 (4): 283-306. <https://doi.org/10.1111/j.1439-0485.1995.tb00413.x>
  - Sini M, Katsanevakis S, Koukouroufli N, Gerovasileiou V, Dailianis T, Buhl-Mortensen L, Damalas D, Dendrinou P, Dimas X, Frantzis A, Gerakaris V, Giakoumi S, Gonzalez-Mirelis G, Hasiotis T, Issaris Y, Kavadas S, Koutsogiannopoulos D, Koutsoubas D, Manoutsoglou E, Markantonatou V, Mazaris A, Poursanidis D, Papatheodorou G, Salomidi M, Topouzellis K, Trygonis V, Vassilopoulou V, Zotou M (2017) Assembling Ecological Pieces to Reconstruct the Conservation Puzzle of the Aegean Sea. *Frontiers in Marine Science* 4: 347. <https://doi.org/10.3389/fmars.2017.00347>
  - SoHelME (2005) State of the Marine Environment Report. Papatheodorou E & Zenetos A (Eds). Hellenic Centre for Marine Research Publishing, Athens, 360 pp. URL: <http://epublishing.ekt.gr/sites/ektpublishing/files/ebooks/Sohelme.pdf>
  - Somaschini A (1988) Policheti della biocenosi ad alghe fotofile (facies a *Corallina elongata*) nel lazio settentrionale. *Atti della Società Toscana di Scienze Naturali Memorie* 95: 83-94. URL: <http://www.stsn.it/AttiB1988/somaschini.pdf>
  - Somaschini A, San Martín G (1994) Description of two new species of *Sphaerosyllis* (Polychaeta: Syllidae: Exogoninae) and first report of *Sphaerosyllis glandulata* for the Mediterranean Sea. *Cahiers de Biologie Marine* 35 (3): 357-367. URL: <http://application.sb-roscoff.fr/cbm/archives.htm>
  - Sordino P (1990) Censimento dei Policheti (Annelida) dei Mari Italiani: Hesionidae Sars, 1862. *Atti della Società Toscana di Scienze Naturali B* 96: 31-52. URL: <http://www.stsn.it/AttiB1989/sordino.pdf>
  - Soulier A (1904) Revision des Annélides de la région de Cette. pt.3. *Memoirs de la Académie des Sciences et Lettres de Montpellier* 3: 319-374.
  - Southern R (1914) Clare Island Survey. Archiannelida and Polychaeta. *Proceedings of the Royal Irish Academy* 31 (47): 1-160. URL: <http://www.jstor.org/stable/20520589>
  - Southward E, Andersen A, Hourdez S (2011) *Lamellibrachia anaximandri* n. sp., a new vestimentiferan tubeworm (Annelida) from the Mediterranean, with notes on frenulate tubeworms from the same habitat. *Zoosystema* 33: 245-279. <https://doi.org/10.5252/z2011n3a1>
  - Spiliotis J (1991) The evolution of a polluted benthic community after pollution abatement. MSc Thesis. National and Kapodistrian University of Athens, Athens, 102 pp. [In Greek].
  - Sromek L, Nygren A, Plyuscheva M, Martín D (2016) *Eupolymnia nebulosa* (Montagu, 1818) revisited: An homage to Michel Bhaud. 11th International Polychaete Conference, Sydney, Australia, 4–9 August 2013. 118 pp. URL: <https://>

- [australianmuseum.net.au/uploads/documents/30456/ipc2013\\_program\\_abstract\\_handbook.pdf](http://australianmuseum.net.au/uploads/documents/30456/ipc2013_program_abstract_handbook.pdf)
- Støp-Bowitz C (1945) Les Scalibregmiens de Norvège. Meddelelser fra det Zoologiske Museum 51: 181-250.
  - Strelzov V (1973) Многощетинковые черви семейства Paraonidae Cerruti, 1909. [Polychaete Worms of the Family Paraonidae Cerruti, 1909 (Polychaeta, Sedentaria)]. Akademia Nauk, Moscow, 170 pp. [In Russian]. URL: [http://herba.msu.ru/shipunov/school/books/streltsov1973\\_paraonidae.pdf](http://herba.msu.ru/shipunov/school/books/streltsov1973_paraonidae.pdf)
  - Stummer-Traunfels RRV (1903) Beiträge zur Anatomie und Histologie der Myzostomen. I. *Myzostoma asteriae* Marenz. Zeitschrift für wissenschaftliche Zoologie 75: 495-595. URL: [http://www.zobodat.at/pdf/Zeitschrift-fuer-wiss-Zoologie\\_75\\_0495-0595.pdf](http://www.zobodat.at/pdf/Zeitschrift-fuer-wiss-Zoologie_75_0495-0595.pdf)
  - Styan C, McCluskey C, Sun Y, Kupriyanova E (2017) Cryptic sympatric species across the Australian range of the global estuarine invader *Ficopomatus enigmaticus* (Fauvel, 1923) (Serpulidae, Annelida). Aquatic Invasions 12 (1): 53-65. <https://doi.org/10.3391/ai.2017.12.1.06>
  - Sun Y, Wong E, Keppel E, Williamson J, Kupriyanova E (2017) A global invader or a complex of regionally distributed species? Clarifying the status of an invasive calcareous tubeworm *Hydroides dianthus* (Verrill, 1873) (Polychaeta: Serpulidae) using DNA barcoding. Marine Biology 164 (1): . <https://doi.org/10.1007/s00227-016-3058-9>
  - Sun Y, Wong E, Tovar-Hernández MA, Williamson J, Kupriyanova E (2016) Is *Hydroides brachyacantha* (Serpulidae: Annelida) a widespread species? Invertebrate Systematics 30 (1): 41-5. <https://doi.org/10.1071/is15015>
  - Surugiu V (2006) New data on the polychaete fauna from the Romanian coast of the Black Sea. Lucrările Conferinței Naționale Biodiversitate și impact antropoc în Marea Neagră și în ecosistemele litorale ale Mării Negre Universitatea „Al.I. Cuza” Iași: 47-56. URL: [https://www.researchgate.net/profile/Victor\\_Surugiu/publication/248399018\\_New\\_data\\_on\\_the\\_polychaete\\_fauna\\_from\\_the\\_Romanian\\_coast\\_of\\_the\\_Black\\_Sea/links/0deec51dfaadf55bc0000000.pdf](https://www.researchgate.net/profile/Victor_Surugiu/publication/248399018_New_data_on_the_polychaete_fauna_from_the_Romanian_coast_of_the_Black_Sea/links/0deec51dfaadf55bc0000000.pdf)
  - Surugiu V (2016) On the taxonomic status of the European *Scolecopsis* (*Scolecopsis*) *squamata* (Polychaeta: Spionidae), with description of a new species from southern Europe. Zootaxa 4161 (2): 151-176. <https://doi.org/10.11646/zootaxa.4161.2.1>
  - Taboada S, Leiva C, Bas M, Schult N, McHugh D (2017) Cryptic species and colonization processes in *Ophryotrocha* (Annelida, Dorvilleidae) inhabiting vertebrate remains in the shallow-water Mediterranean. Zoologica Scripta 46 (5): 611-624. <https://doi.org/10.1111/zsc.12239>
  - Taviani M, Angeletti L, Canese S, Cannas R, Cardone F, Cau A, Cau AB, Follesa MC, Marchese F, Montagna P, Tessarolo C (2015) The “Sardinian cold-water coral province” in the context of the Mediterranean coral ecosystems. Deep Sea Research Part II: Topical Studies in Oceanography <https://doi.org/10.1016/j.dsr2.2015.12.008>
  - Tena J, Capaccioni-Azzati R, Porras R, Torre-Gavilá FJ (1991) Cuatro especies de Poliquetos nuevas para las costas Mediterráneas españolas en los sedimentos del antepuerto de Valencia. Miscellània Zoològica 15: 29-41.
  - Tena J, Capaccioni-Azzati R, Torres-Gavila FJ, Torres-Gavila FJ, García-Carrascosa AM (2000) Polychaetes associated with different facies of the photophilic algal community in the Chafarinas archipelago (SW Mediterranean). Bulletin of Marine Science 67 (1): 55-72. URL: <http://www.ingentaconnect.com/content/umrsmas/bullmar/2000/00000067/00000001/art00007>



- ten Hove H (2003) Possible origin of *Hydroides elegans* . <http://www.bio.net/bionet/mm/annelida/2003-May/002108.html>. Accessed on: 2017-9-06.
- ten Hove H, Kupriyanova E (2009) Taxonomy of Serpulidae (Annelida, Polychaeta): The state of affairs. *Zootaxa* 2036: 1-126. URL: <http://www.mapress.com/zootaxa/2009/zt02036p126.pdf>
- Thingstad TF, Krom MD, Mantoura RFC, Flaten GAF, Groom S, Herut B, Kress N, Law CS, Pasternak A, Pitta P, Psarra S, Rassoulzadegan F, Tanaka T, Tselepides A, Wassmann P, Woodward EMS, Riser CW, Zodiatis G, Zohary T (2005) Nature of Phosphorus Limitation in the Ultraoligotrophic Eastern Mediterranean. *Science* 309 (5737): 1068-1071. <https://doi.org/10.1126/science.1112632>
- Torres-Gavila FJ (2008) Estudio faunístico, ecológico y ambiental de la fauna de anélidos poliquetos de sustratos sueltos de las Islas Chafarinas (Mar de Alborán, S.W. Mediterráneo). [Faunal, ecological and environmental study of the fauna of the polychaete annelids from soft substrates of the Chafarinas Islands (Alboran Sea, S.W. Mediterranean)]. Universitat de València, Valencia, 695 pp. [In Spanish]. URL: <http://roderic.uv.es/handle/10550/15862>
- Tovar-Hernández MA, Licciano M, Giangrande A (2007) Revision of *Chone* Krøyer, 1856 (Polychaeta: Sabellidae) from the eastern central Atlantic and Mediterranean Sea with descriptions of two new species. *Scientia Marina* 71 (2): 315-338. <https://doi.org/10.3989/scimar.2007.71n2315>
- Tovar-Hernández MA (2007) On some species of *Chone* Krøyer, 1856 (Polychaeta: Sabellidae) from world-wide localities. *Zootaxa* 1518: 31-68.
- Tovar-Hernández MA, Harris LH (2010) *Parasabella* Bush, 1905, replacement name for the polychaete genus *Demonax* Kinberg, 1867 (Annelida, Polychaeta, Sabellidae). *ZooKeys* 19: 13-19. <https://doi.org/10.3897/zookeys.60.547>
- Tselepides A (1992) Οικολογική μελέτη του βαθύαλου οικοσυστήματος του Αιγαίου πελάγους. [Ecological study of the bathyal ecosystem of the Aegean Sea.]. PhD thesis. University of Crete, Heraklion, 243 pp. [In Greek].
- Tselepides A, Lampadariou N, Hatziyanni E (2004) Distribution of meiobenthos at bathyal depths in the Mediterranean Sea. A comparison between sites of contrasting productivity. *Scientia Marina* 68: 39-51. <https://doi.org/10.3989/scimar.2004.68s339>
- Tursi A, Mastrotoaro F, Matarrese A, Maiorano P, D'onghia G (2004) Biodiversity of the white coral reefs in the Ionian Sea (Central Mediterranean). *Chemistry and Ecology* 20: 107-116. <https://doi.org/10.1080/02757540310001629170>
- Ulman A, Ferrario J, Occhpinti-Ambrogi A, Arvanitidis C, Bandi A, Bertolino M, Bogi C, Chatzigeorgiou G, Çiçek BA, Deidun A, Ramos-Esplá A, Koçak C, Lorenti M, Martínez-Laiz G, Merlo G, Princisgh E, Scribano G, Marchini A (2017) A massive update of non-indigenous species records in Mediterranean marinas. *PeerJ* 5: e3954. <https://doi.org/10.7717/peerj.3954>
- Valvassori G, Massa-Gallucci A, Gambi MC (2015) Reappraisal of *Platynereis massiliensis* (Moquin-Tandon) (Annelida, Nereididae), a neglected sibling species of *Platynereis dumerilii* (Audouin & Milne Edwards). *Biologia Marina Mediterranea* 22: 113-116.
- Vamvakas C (1970) Peuplements benthiques des substrats meubles du sud de la mer Égée. *Tethys* 2 (1): 89-130. URL: <https://decapoda.nhm.org/pdfs/30169/30169.pdf>
- Vamvakas C (1971) Συμβολή εις την μελέτην των βενθικών βιοκοινωνιών μαλακού υποστρώματος των ελληνικών θαλασσών (περιοχή Δυτικού

Σαρωνικού Κόλπου). [Contribution to the study of benthic biocoenoses in soft substrata around the greek seas (region of the Saronikos Gulf)]. Hellenic Oceanol. Limnol. - Institute of Oceanography and Fisheries Research, Athens, 129-272 pp. [In Greek].

- Vamvakopoulou S (1991) Comparative study of the benthic fauna of three lagoons: Vivari, Pogonitsa, Tsopeli. MSc thesis. National and Kapodistrian University of Athens, Athens, 98 pp. [In Greek].
- Vasileiadou K, Pavloudi C, Sarropoulou E, Fragopoulou N, Kotoulas G, Arvanitidas C (2016) Unique COI haplotypes in *Hediste diversicolor* populations in lagoons adjoining the Ionian Sea. Aquatic Biology 25: 7-15. <https://doi.org/10.3354/ab00654>
- Verrill AE (1875) Results of dredging expeditions off the New England coast in 1874. American Journal of Science 54: 411-415. <https://doi.org/10.2475/ajs.s3-9.54.411>
- Vertino A, Savini A, Rosso A, Di Geronimo I (2010) Benthic habitat characterization and distribution from two representative sites of the deep-water SML Coral Province (Mediterranean). Deep Sea Research Part II: Topical Studies in Oceanography 57: 380-380. <https://doi.org/10.1016/j.dsr2.2009.08.023>
- von Marenzeller E (1893) Berichte der Commission für Oceanographische Erforschung des Östlichen Mittelmeeres. Zweite Reihe. VI. Zoologische Ergebnisse II. Polychäten des Grundes, gesammelt 1890, 1891, 1892. Denkschriften der Akademie der Wissenschaften, Wien 60: 25-48. URL: <http://www.biodiversitylibrary.org/item/31609#page/735/mode/1up>
- von Marenzeller E (1902) Berichte der Commission für Oceanographische Forschungen. Achte Reihe. Forschungen im östlichen Mittelmeere. Zoologische Ergebnisse XIII. Polychäten des Grundes, gesammelt 1893, 1894. Denkschriften der Akademie der Wissenschaften, Wien 74: 295-323. URL: [http://www.zobodat.at/pdf/DAKW\\_74B\\_0295-0323.pdf](http://www.zobodat.at/pdf/DAKW_74B_0295-0323.pdf)
- von Soosten C, Schmidt H, Westheide W (1998) Genetic variability and relationships among geographically widely separated populations of *Petitita amphophthalma* (Polychaeta: Syllidae). Results from RAPD-PCR investigations. Marine Biology 131 (4): 659-669. <https://doi.org/10.1007/s002270050358>
- Voultziadou E, Gerovasileiou V, Vandepitte L, Ganias K, Arvanitidis C (2017) Aristotle's scientific contributions to the classification, nomenclature and distribution of marine organisms. Mediterranean Marine Science 18: 468-478. <https://doi.org/10.12681/mms.13874>
- Wasson B, Giangrande A, Tovar-Hernández MA (2017) Re-establishment of *Chone filicaudata* Southern, 1914 (Annelida: Sabellidae) and the first record of the Mediterranean species *Dialychone dunerificta* (Tovar-Hernández et al., 2007) (Annelida: Sabellidae) in British waters. Cahiers de Biologie Marine 58: 371-37. <https://doi.org/10.21411/CBM.A.77998E0C>
- Wehe T, Fiege D (2002) Annotated checklist of the polychaete species of the seas surrounding the Arabian Peninsula: Red Sea, Gulf of Aden, Arabian Sea, Gulf of Oman, Arabian Gulf. Fauna of Arabia 19: 7-238. URL: <http://www.senckenberg.de/files/content/forschung/abteilung/aquazool/mev2/wehefiege.pdf>
- Weidhase M, Bleidorn C, Helm C (2014) Structure and anterior regeneration of musculature and nervous system in *Cirratulus* cf. *cirratulus* (Cirratulidae, Annelida). Journal of Morphology 275 (12): 1418-1430. <https://doi.org/10.1002/jmor.20316>

- Weigert A, Bleidorn C (2016) Current status of annelid phylogeny. *Organisms Diversity & Evolution* 16 (2): 345-362. <https://doi.org/10.1007/s13127-016-0265-7>
- Wesenberg-Lund E (1939a) Pelagic polychaetes of the families Aphroditidae, Phyllodocidae, Typhloscolecidae and Alciopidae . Report of the Danish Oceanographic Expedition to the Mediterranean and adjacent seas *Biology* 2: 1-46.
- Wesenberg-Lund E (1939b) Polychètes et Géphyriens de Tunisie. *Bulletin de la Station Océanographique de Salammbô* 39: 1-20.
- Westheide W (1974) Interstitielle Fauna von Galapagos. XI. Pisionidae, Hesionidae, Pilargidae, Syllidae (Polychaeta). *Mikrofauna des Meeresbodens* 44: 1-146.
- Westheide W, Hass-Cordes E (2001) Molecular taxonomy: description of a cryptic *Petitia* species (Polychaeta: Syllidae) from the Island of Mahé (Seychelles, Indian Ocean) using RAPD markers and ITS2 sequences. *Journal of Zoological Systematics and Evolutionary Research* 39: 103-111. <https://doi.org/10.1046/j.1439-0469.2001.00166.x>
- Wijnhoven S, Dekker A (2010) Records of a new alien polychaete worm species, *Marphysa sanguinea* (Montagu, 1815) (Eunicidae) in the Eastern Scheldt, the Netherlands. *Aquatic Invasions* 5 (4): 431-436. <https://doi.org/10.3391/ai.2010.5.4.13>
- Wilson DP (1958) The polychaete *Magelona alleni* n. sp. and a re-assessment of *Magelona cincta* Ehlers. *Journal of the Marine Biological Association of the United Kingdom* 37 (3): 617-626. <https://doi.org/10.1017/S002531540000566X>
- Wilson WHJ (1983) Life-history evidence for sibling species in *Axiiothella rubrocincta* (Polychaeta: Maldanidae). *Marine Biology* 76: 297-300. <https://doi.org/10.1007/BF00393032>
- Wisnens IM (1989) Eunicid polychaetes (Annelida) from Scandinavian and adjacent waters. Family Eunicidae . *Zoologica Scripta* 18 (4): 483-500. <https://doi.org/10.1111/j.1463-6409.1989.tb00142.x>
- Wolff W, Stegenga H (1975) *Hesionura augeneri*, *Goniadella bobretzkii*, and *Parapodrilus psammophilus* (Annelida, Polychaeta): new to the Netherlands. *Zoologische Bijdrage* 17 (12): 82-8.
- Worsfold T (2007) Identification guides for the NMBAQC Scheme: 2. Goniadidae, with notes on Glyceridae (Polychaeta) from shallow seas around the British Isles. Porcupine Marine Natural History Society Newsletter 22: 19-23. URL: [http://www.nmbaqcs.org/media/1412/goniadidae\\_june07.pdf](http://www.nmbaqcs.org/media/1412/goniadidae_june07.pdf)
- Xenopoulou D (1987) Study of the distribution and temporal variations of the soft substrate benthic fauna of the island of Skiathos. MSc thesis. University of Athens, Athens, 81 pp. [In Greek].
- Yamanishi R (1998) Ten Species of *Pisione* (Annelida : Polychaeta : Pisionidae) from Japan and Evolutionary Trends in the Genus Based on Comparison of Male Copulatory Apparatus. *Publications of the Seto Marine Biological Laboratory* 38: 83-145. <https://doi.org/10.5134/176280>
- Yáñez-Rivera B, Carrera-Parra LF (2012) Reestablishment of *Notopygos megalops* McIntosh, description of *N. caribea* sp. n. from the Greater Caribbean and barcoding of “amphiamerican” *Notopygos* species (Annelida, Amphinomidae). *ZooKeys* 223: 69-84. <https://doi.org/10.3897/zookeys.223.3561>
- Yáñez-Rivera B, Salazar-Vallejo S (2011) Revision of *Hermodice* Kinberg, 1857 (Polychaeta: Amphinomidae). *Scientia Marina* 75 (2): 251-262. <https://doi.org/10.3989/scimar.2011.75n2251>

- Yokoyama H, Daglı E, Çınar ME (2010) First record of *Paraprionospio coora* Wilson, 1990 (Polychaeta: Spionidae) from the Mediterranean Sea. *Mediterranean Marine Science* 11 (1): 133-142. <https://doi.org/10.12681/mms.96>
- Zaâbi S (2013) Étude faunistique et écologique des Annélides Polychètes de la côte Nord-Est de la Tunisie (Peninsule du Cap Bon, Méditerranée Ouest). [Faunistic and ecological study of the Annelida Polychaeta of the North-East coast of Tunisia (Cap Bon Peninsula, Western Mediterranean)]. Université de Carthage, Tunis, 386 pp. [In French]. URL: <http://hdl.handle.net/1834/5480>
- Zaâbi S, Gillet P, Afli A, Boumaiza M (2009) Biodiversity of polychaetous annelids from the peninsula of Cap Bon, northeast coast of Tunisia. *Zoosymposia* 2: 587-600. URL: <http://citeseerx.ist.psu.edu/viewdoc/download;jsessionid=487A95425C9E7160197D00FF7482C14B?doi=10.1.1.643.4776&rep=rep1&type=pdf>
- Zaâbi S, Gillet P, Chambers S, Afli A, Boumaiza M (2012) Inventory and new records of polychaete species from the Cap Bon Peninsula, north-east coast of Tunisia, Western Mediterranean Sea. *Mediterranean Marine Science* 13 (1): 36-48. <https://doi.org/10.12681/mms.22>
- Zaâbi S, Metais I, Gillet P, Afli A, Boumaiza M (2015) Preliminary Study of Population Genetic Diversity of *Hyalinoecia tubicola* (Polychaeta: Onuphidae) from the North East Coast of Tunisia (Western Mediterranean) using Random Amplified Polymorphic DNA Markers. *Journal of Coastal Zone Management* 18 (1): 397. <https://doi.org/10.4172/2473-3350.1000397>
- Zanol J, Bettoso N (2006) Identity of *Eunice roussaei* (Eunicidae: Polychaeta: Annelida) from the Adriatic and Mediterranean Seas. *Journal of the Marine Biological Association of the United Kingdom* 86 (5): 1017. <https://doi.org/10.1017/s0025315406013993>
- Zanol J, Ruta C (2015) New and previously known species of Oeonidae (Polychaeta: Annelida) from Lizard Island, Great Barrier Reef, Australia. *Zootaxa* 4019 (1): 745. <https://doi.org/10.11646/zootaxa.4019.1.26>
- Zanol J, da Silva TC, Hutchings P (2016) *Marphysa* (Eunicidae, polychaete, Annelida) species of the Sanguinea group from Australia, with comments on pseudo-cryptic species. *Invertebrate Biology* 135 (4): 328-344. <https://doi.org/10.1111/ivb.12146>
- Zanol J, Halanych K, Fauchald K (2013) Reconciling taxonomy and phylogeny in the bristlemoth family Eunicidae (polychaete, Annelida). *Zoologica Scripta* 43 (1): 79-100. <https://doi.org/10.1111/zsc.12034>
- Zarkanellas AJ (1980) Οικολογική μελέτη του μακροβένθους του Θερμαϊκού κόλπου. [Ecological study of the macrobenthos in Thermaikos Gulf]. PhD thesis. Aristotle University of Thessaloniki, Thessaloniki, 147 pp. [In Greek].
- Zenetos A (1986) Zoobenthos Kyklades, Aegean Sea dataset. Hellenic Centre for Marine Research, Greece.. Release date: 2005-11-23. URL: <http://ipt.medobis.eu/resource?r=kyklades>
- Zenetos A, Christianidis S, Pancucci AM, Simbours N, Tziavos C (1997) Oceanographic study of an open coastal area in the Ionian sea with emphasis on its benthic fauna and some zoogeographical remarks. *Oceanologica Acta* 202 (2): 437-451. URL: <http://archimer.ifremer.fr/doc/00093/20408/18075.pdf>
- Zenetos A, Çınar ME, Crocetta F, Golani D, Rosso A, Servello G, Shenkar N, Turon X, Verlaque M (2017) Uncertainties and validation of alien species catalogues: The

Mediterranean as an example. *Estuarine, Coastal and Shelf Science* 191: 171-187.

<https://doi.org/10.1016/j.ecss.2017.03.031>

- Zenetos A, Cinar M, Pancucci-Papadopoulou M, Harmelin J, Furnari G, Andaloro F, Bellou N, Streftaris N, Zibowius H (2005) Annotated list of marine alien species in the Mediterranean with records of the worst invasive species. *Mediterranean Marine Science* 6: 63-118.
- Zenetos A, Katsanevakis S, Poursanidis D, Crocetta F, Damalas D, Apostolopoulos G, Gravili C, Vardala-Theodorou E, Malaquias M (2011) Marine alien species in Greek Seas: Additions and amendments by 2010. *Mediterranean Marine Science* 12 (1): 95-120. <https://doi.org/10.12681/mms.55>
- Zenetos A, Gofas S, Verlaque M, Çinar ME, Garcia Raso JE, Bianchi CN, Morri C, Azzurro E, Bilecenoglu M, Froggia C, Siokou I, Violanti D, Sfriso A, San Martín G, Giangrande A, Mastrototaro F, Katagan T, Ballesteros E, Ramos-Espla A, Mastrototaro F, Ocana O, Zingone A, Gambi MC, Streftaris N (2010) Alien species in the Mediterranean Sea by 2010 . A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I . Spatial distribution. *Mediterranean Marine Science* 12 (1): 381-493. <https://doi.org/10.12681/mms.87>
- Zenetos A, Gofas S, Morri C, Rosso A, Violanti D, Garcia Raso JE, Çinar ME, Almogilabini A, Ates AS, Azzurro E, Ballesteros E, Bianchi CN, Bilecenoglu M, Gambi MC, Giangrande A, Gravili C, Hyams-Kaphzan O, Karachle PK, Katsanevakis S, Lipej L, Mastrototaro F, Mineur F, Pancucci-Papadopoulou MA, Ramos Espla A, Salas C, San Martín G, Sfriso A, Streftaris N, Verlaque M (2012) Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science* 13 (2): 328-352. <https://doi.org/10.12681/mms.327>
- Zibowius H (1968) Étude morphologique, systématique et écologique des Serpulidae (Annelida Polychaeta) de la région de Marseille. *Recueil des Travaux de la Station Marine d'Endoume, Bulletin* 43: 81-252.
- Zibowius H (1971) Les espèces Méditerranéennes du genre *Hydroïdes* (Polychaeta Serpulidae). Remarques sur le prétendu polymorphisme de *Hydroïdes uncinata* . *Tethys* 2: 691-746. URL: [http://paleopolis.rediris.es/benthos/REF/som/T-pdf/1970\\_2-3-691.pdf](http://paleopolis.rediris.es/benthos/REF/som/T-pdf/1970_2-3-691.pdf)
- Zibowius H (1973a) Revision of some Serpulidae (Annelida Polychaeta) from abyssal depths in the Atlantic and Pacific, collected by the 'Challenger' and Prince of Monaco Expeditions. *Bulletin of the British Museum (Natural History)* 24 (9): 427-439. URL: <http://biodiversitylibrary.org/page/26497963>
- Zibowius H (1973b) Remarques sur trois espèces de Serpulidae acclimatées en Méditerranée: *Hydroïdes dianthus* (Verrill, 1873), *Hydroïdes dirampha* Mörch, 1863, et *Hydroïdes elegans* (Haswell, 1883). Rapport et Procès-Verbaux des Réunions de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée 21: 683-686. URL: [http://ciesm.org/online/archives/abstracts/pdf/22/CIESM\\_Congress\\_1970\\_Rome\\_article\\_0020.pdf](http://ciesm.org/online/archives/abstracts/pdf/22/CIESM_Congress_1970_Rome_article_0020.pdf)
- Zibowius H (1979) Serpulidae (Annelida Polychaeta) de l'Océan Indien arrivés sur des coques de bateaux à Toulon (France, Méditerranée). Rapport et Procès-Verbaux des Réunions de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée 25: 133-134.

- Zrzavý J, Hyspa V, Tietz D (2001) Myzostomida are not annelids: molecular and morphological support for a clade of animals with anterior sperm flagella. *Cladistics* 17: 170-198. <https://doi.org/10.1006/clad.2000.0148>
- Zunarelli R (1967) Una nuova specie di *Microphthalmus* delle coste livornesi (Polychaeta Hesionidae). *Archivio Zoologico Italiano* 52: 217-220.

## Supplementary materials

### Suppl. material 1: List of unpublished / online resources [doi](#)

**Authors:** Nomiki Simboura, Sarah Faulwetter

**Data type:** Literature list

**Brief description:** A list of references (unpublished datasets by the Hellenic Centre for Marine Research) and online resources which were used to assess the occurrence status of many species. This list is merely for completeness and for the convenience of the reader.

**Filename:** AdditionalSources.pdf - [Download file](#) (22.18 kb)

### Suppl. material 2: List of species per literature resource [doi](#)

**Authors:** Sarah Faulwetter, Nomiki Simboura, Nikolaos Katsiaras, Georgios Chatzigeorgiou, Christos Arvanitidis

**Data type:** References for species occurrence records in Greece, \*.csv format, tab-separated

**Brief description:** Contains a list of all consulted literature references with polychaete records in Greece. Note that a) the list of polychaetes for a few sources may be incomplete as only rare or new records were noted and b) that records mentioned in two or more sources may in some cases pertain to the same specimen, either because the same dataset was used in several publications or because records were repeated in secondary literature (checklists, databases).

Explanation of columns: *Family*: Family the taxon belongs to; *ScientificName\_accepted*: Currently (December 2017) accepted scientific name for the taxon; *Rank*: taxonomic rank; *ScientificAuthority\_accepted*: Currently (December 2017) accepted scientific authority for the taxon; *ScientificName\_asReported*: Taxon as reported in the given literature /data resource, standardised with WoRMS to remove spelling mistakes and typographic errors; *ScientificAuthority\_asReported*: Scientific authority as reported in the given literature /data resource, standardised with WoRMS to remove spelling mistakes and typos; *Source\_authors*: Authors of literature/data resource; *Source\_year*: Year of literature/data resource; *Source\_reference*: Other bibliographic info (title, journal, URL, etc) of literature/data resource; *Status\_in\_Greece*: current status of the taxon in Greece (present/absent/questionable); *Occurrence\_record\_notes*: information on the specific taxon-resource combination, e.g. misidentifications of the specific specimens;

**Filename:** GreekPolychaetes\_Species-by-reference.csv - [Download file](#) (4.92 MB)