



Radula and Shell Microstructure Variations are Congruent with a Molecular Estimate of Shallow-Water Japanese Chitons

Author: Owada, Masato

Source: Zoological Science, 40(5) : 390-403

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zs220060>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Radula and Shell Microstructure Variations are Congruent With a Molecular Estimate of Shallow-Water Japanese Chitons

Masato Owada*

Faculty of Science, Department of Biological Sciences, Kanagawa University,
3-27-1 Rokkakubashi, Kanagawa-ku, Yokohama,
Kanagawa 221-8686, Japan

Variations of the radula and shell microstructures in 33 species of Japanese chiton were investigated along with molecular phylogenetic trees. The molecular phylogenetic trees indicated that Chitonida was composed of four clades, of which two clades formed Acanthochitonina and corresponded to Mopalioida and Cryptoplacoidea, respectively, and the other clades formed Chitonina. In the radula, the shapes of the central and centro-lateral teeth and the petaloid process varied greatly among species or genera and were useful for the identification of particular species or genera. The presence of accessory and petaloid processes and the cusp shape were relatively conserved and useful for recognizing particular genera or even suborders. In the valves, four to six shell layers were found at the section, but the ventral mesostracum was not observed in Acanthochitonina. The shell microstructures in the ventral sublayer of the tegmentum varied at suborder, but those in the other layers were almost constant. The megal aesthete chamber type varied at superfamily and was helpful to identify particular families or superfamilies. The characteristics of the shell layers and shell microstructures appear to be a synapomorphy shared by the members of Acanthochitonina. The classification within Chitonina needs to be reexamined because the variations of the cusp shape and megal aesthete chamber type were relatively large and did not correspond to the current classification. *Callochiton* formed a sister group with Chitonida and would be equally closely related to Chitonina and Acanthochitonina because of possessing a mosaic of characteristics from both.

Key words: Mollusca, Polyplacophora, chiton, radula, shell microstructure, molecular phylogeny, cytochrome c oxidase subunit I, 16S ribosomal RNA, 18S ribosomal RNA, 28S ribosomal RNA

INTRODUCTION

Chitons, one of the minor groups in Mollusca, are elongate-oval and flattened in shape and have eight shell valves like armor. These valves are surrounded by a girdle, and the foot is on the ventral side. They have no head, and the oral region lacks eyes; instead, sensory organs called aesthetes are embedded in the valves. Chitons inhabit only seawater, clinging to the surface of something solid such as rocks and dead shells, and most eat algae or minute organisms with the radula. Their fossil records may date back to the Late Cambrian (Smith, 1960; Runnegar et al., 1979; Puchalski et al., 2008; Pojeta et al., 2010), and an obvious chiton fossil would be *Echinochiton* from Ordovician (Pojeta et al., 2003; Pojeta and Dufoe, 2008).

Approximately 1200 species of chitons have been discovered from shallow to deep seas in the world (Eernisse and Reynolds, 1994; Sliker, 2000; Schwabe, 2005; MolluscaBase editors, 2021). Chitons have been mainly classified based on the characteristics of shell, girdle, and radula (Bergenhayn, 1955; Kaas and Van Belle, 1980, 1990, 1994). Furthermore, Sirenko (2006) indicated a new classification

by adding characteristics such as gills, glands, egg hull projections, and spermatozooids. The phylogenetic relationships of chitons have been examined using these, and other characters, such as DNA sequences (Okusu et al., 2003; Sigwart et al., 2010, 2013; Irisarri et al., 2014, 2020), aesthete canals and complexes (Fernandez et al., 2007; Vendrasco et al., 2008), and shell microstructures (Peebles et al., 2017). These phylogenetic relationships predominantly corresponded to each other in the higher taxa, but sometimes not in the lower taxa.

The characteristics of radulae are useful tools for the classification of gastropods (e.g., Sasaki, 2010; Ponder and Lindberg, 2020), and those of shell microstructures reflect the phylogenetic relationships of higher taxa, especially in bivalves (e.g., Taylor et al., 1969, 1973; Carter, 1990). As for chitons, Dall (1879) described the radula dentition of North Pacific chitons with comparing species. Thiele (1909–1910) provided monographs of worldwide chitons and emphasized the characteristics of the radula more than other chiton researchers. Sirenko has been actively publishing more than 100 publications covering most chiton taxa, and the majority of them include scanning electron microscope (SEM) images of the radula (e.g., Sirenko and Zhang, 2019; Sirenko, 2020). For Japanese species, Taki Is and Taki Iw published precise illustrations

* Corresponding author. E-mail: owada-bio@kanagawa-u.ac.jp
doi:10.2108/zs220060

of the radula without SEM (e.g., Taki Is and Taki Iw, 1929a; Taki Is, 1938). Saito (2004) analyzed the radula variations of many species in Cryptoplacoidea and reported that their characteristics were useful in identifying the species and genus and could be used to construct the phylogenetic relationships. Bergenhayn (1930) was the first to examine the shell microstructures of chitons, but the descriptions were not very detailed because SEM was not used. Recently, Peebles et al. (2017) described the mineralogy and shell microstructures of eight chiton species collected from New Zealand using SEM and discussed the phylogenetic relationships at family level. However, in both Saito (2004) and Peebles et al. (2017), the phylogenetic resolutions were not so clear. The characteristics of the radula alone were not enough to analyze the higher taxa, and the shell microstructures alone were not enough to analyze the lower taxa. Probably, in the phylogenetic analysis of chitons, combining the characteristics of the radula and shell microstructures is necessary.

More than 90 species of chitons have been discovered on or near the coasts of Japan. The classification of

Japanese chitons was well arranged by Taki Is and Taki Iw (1929a, 1929b, 1929c, 1930, 1931a, 1931b), Taki Is (1938), and Saito (1994, 1995, 1998, 2000, 2001, 2006, 2011, 2017). Most Japanese chitons can be identified according to the taxonomic keys created by them.

In the present study, Japanese chitons living in the shallow sea were investigated. They inhabit various environments, and the species diversity is relatively high. First, phylogenetic relationships were constructed using the DNA sequences of mitochondrial cytochrome c oxidase subunit I (COI) and 16S ribosomal RNA (16S), and nuclear 18S ribosomal RNA (18S) and 28S ribosomal RNA (28S). Radula and shell microstructures were observed by SEM. Based on these results, the relationships among the variations of the radula and shell microstructures and the molecular phylogenetic tree of Japanese chitons living in the shallow sea are discussed.

MATERIALS AND METHODS

Sampling

All investigated specimens, identified as 33 chiton species,

Table 1. Species names, sampling localities, sampling dates, and accession numbers. Newly or renewedly determined DNA sequences are in bold.

order	suborder	superfamily	family	species name	sampling locality	sampling date	accession number							
							CO1	16S	18S	28S				
Lepidopleurida	Lepidopleurina	Lepidopleuroidea	Leptochitonidae	<i>Leptochiton aequispinus</i>	Misaki	November, 2015	LC718179	LC718131	LC718147	LC718163				
				<i>Leptochiton hakodatensis</i>	Asamushi	May, 2016	LC214408	LC214397	LC214369	LC214386				
Callochitonida	Callochitonina	Callochitonoidea	Callochitonidae	<i>Callochiton foveolatus</i>	Shimoda	May, 2016	LC214414	LC214403	LC214380	LC214392				
Chitonida	Chitonina	Chitonoidea	Ischnochitonidae	<i>Ischnochiton boninensis</i>	Manazuru	May, 2014	LC071647	LC071575	LC214370	LC071611				
				<i>Ischnochiton comptus</i>	Manazuru	May, 2014	LC071627	LC071570	LC214371	LC071606				
				<i>Ischnochiton hakodadensis</i>	Hakodate	August, 2015	LC214409	LC214398	LC214372	LC214387				
				<i>Ischnochiton hayamii</i>	Zushi	May, 2015	LC214410	LC214399	LC214373	LC214388				
				<i>Ischnochiton manazuruensis</i>	Manazuru	April, 2016	LC071619	LC071565	LC214374	LC071601				
				<i>Ischnochiton paululus</i>	Asamushi	June, 2015	LC214411	LC214400	LC214375	LC214389				
				<i>Ischnochiton poppei</i>	Isso	August, 2016	LC214412	LC214401	LC214376	LC214390				
				<i>Stenoplax alata</i>	Isso	August, 2016	LC214413	LC214402	LC214377	LC214391				
				<i>Tripoplax albrechti</i>	Hakodate	August, 2015	LC718180	LC718132	LC718148	LC718164				
				<i>Lepidozonia coreanica</i>	Manazuru	May, 2014	LC071669	LC071582	LC214378	LC071618				
				Callistoplacidae			<i>Callistochiton jacobaeus</i>	Manazuru	May, 2014	LC071667	LC071580	LC214379	LC071616	
				Chitonidae				<i>Rhyssoplax kurodai</i>	Manazuru	May, 2014	LC071668	LC071581	LC214381	LC071617
								<i>Rhyssoplax komaiana</i>	Isso	August, 2016	LC718185	LC718137	LC718153	LC718169
								<i>Acanthopleura gemmata</i>	Onna	June, 2017	LC718175	LC718127	LC718143	LC718159
								<i>Acanthopleura loochooana</i>	Onna	June, 2017	LC718176	LC718128	LC718144	LC718160
<i>Liolophura japonica</i>	Manazuru	March, 2017	LC718181					LC718133	LC718149	LC718165				
<i>Lucilina amanda</i>	Zushi	September, 2019	LC718187					LC718139	LC718155	LC718171				
<i>Lucilina lamellosa</i>	Onna	June, 2017	LC718188					LC718140	LC718156	LC718172				
<i>Onithochiton hirasei</i>	Zushi	April, 2017	LC718183					LC718135	LC718151	LC718167				
Acanthochitonina	Mopalioidae		Tonicellidae	<i>Tonicella lineata</i>	Asamushi	May, 2016	LC214415	LC214404	LC214382	LC214393				
				<i>Tonicella zotini</i>	Abashiri	August, 2018	LC718189	LC718141	LC718157	LC718173				
			Schizoplacidae	<i>Schizoplax brandtii</i>	Shari	August, 2018	LC718186	LC718138	LC718154	LC718170				
				Mopaliidae	<i>Mopalia retifera</i>	Shimoda	May, 2016	LC214416	LC214405	LC214383	LC214394			
					<i>Placiphorella stimpsoni</i>	Shimoda	May, 2017	LC718184	LC718136	LC718152	LC718168			
Cryptoplacoidea	Acanthochitonidae			<i>Acanthochitona defilippii</i>	Manazuru	March, 2016	LC214417	LC214406	LC214384	LC214395				
				<i>Acanthochitona achates</i>	Manazuru	March, 2016	LC718178	LC718130	LC718146	LC718162				
				<i>Acanthochitona dissimilis</i>	Manazuru	July, 2017	LC718174	LC718126	LC718142	LC718158				
				<i>Notoplax conica</i>	Shimoda	May, 2017	LC718182	LC718134	LC718150	LC718166				
				<i>Cryptochiton stelleri</i>	Akkeshi	May, 2021	LC718177	LC718129	LC718145	LC718161				
Cryptoplacidae			<i>Cryptoplax japonica</i>	Manazuru	May, 2015	LC214418	LC214407	LC214385	LC214396					

were collected from the adjacent water of Japan. They were collected from rocks or boulders on the shore during low tide, except *Leptochiton aequispinus* (Bergenhayn, 1993) collected at a water depth of 210 m by dredging, and *Lept. hakodatensis* (Thiele, 1909), *Ischnochiton paululus* Is. Taki, 1938, *Tonicella lineata* (Wood, 1815) and *Cryptochiton stelleri* (Middendorff, 1847) collected at a water depth of 3–8 m by SCUBA diving. The dates and localities of the samplings were shown in Table 1 and Fig. 1. All collected specimens were preserved in 100% EtOH directly or after freezing at -20°C . The species identification was performed as described by Saito (2017), and their classification followed the MolluscaBase editors (2021).

Phylogenetic analysis

Total DNA was extracted from the foot or rectal muscle of each specimen using a DNeasy Blood & Tissue Kit (QIAGEN). COI, 16S, 18S, and 28S gene regions were amplified by polymerase chain reaction (PCR) using Premix Taq (Takara) and a thermal cyclor (T100, Bio-Rad). As for the 18S and 28S gene regions, two fragments that were separated into the upper and lower regions were amplified. The primers used are listed in Table 2. The conditions for PCR amplification were as follows: denaturation at 94°C for 30 s; annealing at 48°C (COI), 55°C (16S and 18S), or 58°C (28S) for 30 s; and extension at 72°C for 60 s. These steps were repeated 25–35 times. The PCR products were purified using ExoSAP-IT (Affymetrix), and cycle-sequencing reactions were performed using a BigDye Terminator v3.1 Cycle Sequencing Kit (ABI). DNA sequences were analyzed using a genetic analyzer (3130 Genetic Analyzer, ABI) in both 5' and 3' directions. All sequences were registered in the DNA Data Bank of Japan (DDBJ) (Table 1).

The alignment of DNA sequences for each gene region was performed using MAFFT v7.475 (Kato et al., 2005). Sites those contained gaps or were of questionable homology were trimmed using trimAl v1.4rev15 (Capella-Gutiérrez et al., 2009). A molecular phylogenetic tree was constructed from the concatenated sequence of the four gene regions by Maximum Likelihood (ML) and Bayesian methods. The ML method was performed using RAxML-NG v1.0.2 (Kozlov et al., 2019). A bootstrap test was performed 10,000 times. The model for the ML method was selected




Fig. 1. Sampling localities. Abashiri: $44^{\circ}03'N$, $144^{\circ}16'E$; Shari: $44^{\circ}03'N$, $144^{\circ}57'E$; Akkeshi: $43^{\circ}01'N$, $144^{\circ}49'E$; Hakodate: $41^{\circ}45'N$, $140^{\circ}43'E$; Asamushi: $40^{\circ}54'N$, $140^{\circ}51'E$; Zushi: $35^{\circ}16'N$, $139^{\circ}34'E$; Manazuru: $35^{\circ}09'N$, $139^{\circ}09'E$; Misaki: $35^{\circ}10'N$, $139^{\circ}32'E$; Shimoda: $34^{\circ}40'N$, $138^{\circ}56'E$; Isso: $30^{\circ}27'N$, $130^{\circ}30'E$; Onna: $26^{\circ}29'N$, $127^{\circ}50'E$.

Table 2. Primers used in the present study. Superscript numbers show the combination in PCR amplification.

region	direction	name	5'—3' sequence	reference
COI	forward ¹	CO1F	TCWACAAATCAYAAAGATATTGG	Owada et al. (2013)
	reverse ¹	CO1R	ACYTCMGGRTGMCCAAAAAATCA	Owada et al. (2013)
16S	forward ¹	primer A	CGCCTGTTTATCAAAAACAT	Xiong and Kocher (1991)
	reverse ¹	primer B	CTCCGGTTTGAACCTCAGATC	Xiong and Kocher (1991)
18S	forward ¹	1F	TACCTGGTTGATCCTGCCAGTAG	Giribet et al. (1996)
	forward ²	3F	GTTTCGATTCCGGAGAGGGA	Giribet et al. (1996)
	reverse	4R	GAATTACCGCGGCTGCTGG	Giribet et al. (1996)
	reverse ¹	5R	CTTGGCAAATGCTTTCGC	Giribet et al. (1996)
	forward	18S a2.0	ATGTTTGCAAAGCTGAAAC	Whiting et al. (1997)
	reverse	18S bi	GAGTCTCGTTCGTTATCGGA	Whiting et al. (1997)
	reverse ²	9R	GATCCTTCCGCAGGTTACCTAC	Giribet et al. (1996)
28S	forward ¹	28S a	GACCCGCTTGAAACACGGA	Whiting et al. (1997)
	reverse	28S b	TCGGAAGGAACCAGCTACTA	Whiting et al. (1997)
	forward	28S-NLF105-22	CCGAAGTTTCCCTCAGGATAGC	www.psb.ugent.be/rRNA
	forward ²	28S Rd4.8a	ACCTATTCTCAAACCTTTAAATGG	Whiting (2002)
	reverse ¹	28S rD5b	CCACAGCGCCAGTTCTGCTTAC	Whiting (2002)
	forward	28S-1600F	CCTGAAAATGGATGGCGCT	This study
	reverse	28S-1600R	AGCGCCATCCATTTTCAGG	Distel et al. (2011)
	reverse ²	28S rD7b1	GACTTCCCTTACCTACAT	Whiting (2002)

for the sequence of each gene region using ModelTest-NG v0.2.0 (Darriba et al., 2020). The Bayesian method was performed using MrBayes v3.1.2 (Ronquist and Huelsenbeck, 2003). This program was run for 50,000,000 generations sampling every 1000th generation. The model for the Bayesian method was selected for the sequence of each gene region using Kakusan4 (Tanabe, 2011). Convergence in the Bayesian method was assessed using Tracer v1.7.1 (Rambaut et al., 2018).

Radula

The radula was carefully extracted from the soft parts of each specimen, which were fixed in 100% ETOH or frozen at -20°C . Each radula was cleaned with a dissecting needle under a binocular microscope (Olympus, SZ40), washed with an ultrasonic washer (Velvo-Clear, VS-100III), and preserved in 100% ETOH. Afterward, the radula was coated with 50-nm-thick platinum by ion sputter (JEOL, JEC-3000FC) and observed using SEM (JEOL, JCM-5000) at 10 kV accelerating voltage. The terminology for the radula followed Saito (2004) and Schwabe (2010).

Shell microstructures and aesthete canals and complexes

The investigated specimen was soaked in 10% chlorine bleach solution (Kao, Kitchen Haiter) for 1–4 days to remove the soft parts. The remaining eight valves were washed with pure water and desiccated at 60°C for 24 h. Valves III–VI were fractured by hand or with

a micro-chisel and hammer. The sections along the antero-posterior axis were coated with 50-nm-thick platinum by ion sputter (JEOL, JEC-3000FC) and observed using SEM (JEOL, JCM-5000) at 10 kV accelerating voltage. The shell layers were identified as described by Connors et al. (2012); however, the anterior and posterior myostracums were not distinguished. The shell microstructures were identified according to Carter (1990). The aesthete canals and complexes were identified on the basis of Baxter and Jones (1981, 1984) and Currie (1992).

RESULTS

Molecular phylogenetic tree

The sequence lengths for the COI, 16S, 18S, and 28S gene regions were 557, 505–516, 1692–1709, and 1279–1313 base pairs (bp), respectively. After alignment and trimming, the analyzed lengths were 557, 499, 1693, and 1283 bp, respectively. The model for the ML method was GTR + G4 for the 16S, 18S, and 28S, TIM3 + I + G4 for the first base of the COI, TVM + I + G4 for the second base, and TIM2 + I + G4 for the third base. The likelihood index was $-\ln 26478.320061$. The model for the Bayesian method was HKY85 + G for the 16S and the third base of the COI, SYM + G + I for the 18S, and GTR + G + I for the 28S and the first and second bases.




Fig. 2. Molecular phylogenetic trees. Clade A: some Chitonina, including the Chitoninae part of Chitonidae; Clade B: the Toniciinae plus Acanthopleurinae part of Chitonidae; Clade C: Mopalioidae; Clade D: Cryptoplacoidea. Filled circles on each node indicate bootstrap > 50% in the ML tree and posterior probability > 0.90 in the Bayesian tree, and open circles indicate not so.

The ML and Bayesian trees are shown in Fig. 2. The topologies of the two trees mostly corresponded to each other, and the bootstrap values and posterior probabilities on the nodes were significantly high. However, there were a few differences in the positions of *Ischnochiton* spp. and *Cryptoc. stelleri*. In both trees, Chitonida was monophyletic and contained four clades. Clade A was composed of some Chitonina, including the Chitoninae part of Chitonidae and actually including *Ischnochiton*, *Rhyssoplax*, *Stenoplax*, *Tripoplax*, *Lepidozona*, and *Callistochiton*. Clade B was composed of the Toniciinae plus Acanthopleurinae part of Chitonidae and including *Acanthopleura*, *Liolophura*, *Lucilina*, and *Onithochiton*. Clade C was composed of Mopalioida and including *Tonicella*, *Mopalia*, *Cryptochiton*, *Placiphorella*, and *Schizoplax*. Clade D was composed of Cryptoplacoidea and including *Acanthochitona*, *Notoplax*, and *Cryptoplax*. Ischnochitonidae, Chitonidae, Mopaliidae, and Acanthochitonidae were polyphyletic, and additionally, *Ischnochiton* and *Acanthopleura* were not monophyletic. *Callochiton* formed a sister group with Chitonida.

Radula

SEM images of the representative radulae of the investigated specimens are shown in Fig. 3. The characteristics of the radula were more or less different among the species; however, these differences were relatively moderate within the genus, except for *Ischnochiton hakodadensis* Carpenter, 1893. The central teeth had various shapes, but they were grouped into six types: bulbous, cup-shaped, deltoid, hooked-needle, rectangular, and spatulate. The centro-lateral (first) teeth also varied, but they were grouped into six types: columnar, envelope-like, fingerlike, fungi-form, sphenoid, and winglike. In addition, the accessory process, which is a minute process on the centro-lateral tooth (Saito, 2004), was observed in members of clades A and B, and *Lept. aequispinus*. The major (second) lateral teeth were grouped into four types: unicuspid, bicuspid, tricuspid, and patellar. Although the type of the major lateral tooth largely varied in clade A, it was conserved in the other clades. Furthermore, the petaloid process, which is a process below the head of the major lateral tooth (Thiele,




Fig. 3. Continued.




Fig. 3. SEM images of radula. A: *Leptochiton aequispinus*; B: *Lept. hakodatensis*; C: *Callochiton foveolatus*; D: *Ischnochiton comptus*; E: *Rhyssoplax komaiana*; F: *Stenoplax alata*; G: *I. hakodadensis*; H: *Tripopla albrechti*; I: *Lepidozona coreanica*; J: *Callistochiton jacobaeus*; K: *Acanthopleura gemmata*; L: *Acanthop. loochooana*; M: *Lucilina amanda*; N: *Lu. lamellosa*; O: *Onithochiton hirasei*; P: *Tonicella lineata*; Q: *To. zotini*; R: *Mopalia retifera*; S: *Cryptochiton stelleri*; T: *Placiphorella stimpsoni*; U: *Schizoplax brandtii*; V: *Acanthochitona defilippii*; W: *Notoplax conica*; X: *Cryptoplax japonica*. a: central tooth; b: centro-lateral (first) tooth; c: major (second) lateral tooth; d: petaloid process; e: accessory process. Scale bar unit indicates micrometer.

1893; Bullock, 1988; Saito, 2004), was found in clades A and B and was grouped into three types: cornered, nodule, and securiform. The characteristics of the radula of each species are shown in Table 3.

Shell microstructures

In the investigated specimens belonging to *Callochiton*, clades A and B, and the outgroups, six shell layers (i.e., tegmentum, dorsal mesostracum, articulamentum, ventral mesostracum, myostracum, and hypostracum) were observed in the valves. As for clades C and D, five shell layers, excluding the dorsal mesostracum from the six layers, were observed. Except in *Cryptoc. stelleri*, the number of shell layers was constant within the clade. In the valve of *Cryptoc. stelleri*, only four layers (i.e., articulamentum, ventral mesostracum, myostracum, and hypostracum) were

found (Fig. 4).

The tegmentum of *Callochiton*, clades A and B, and *Placiphorella stimpsoni* (Gould, 1859) was composed of two layers. The dorsal sublayer was homogeneous (Ho) structure, and the ventral sublayer was mainly composite prismatic (CP) structure but partly Ho structure (Fig. 5A–H). In clades C and D, and the outgroups, the tegmentum was only one layer, the Ho structure (Fig. 5I–L). The dorsal and ventral mesostracums were constructed with crossed-lamellar (CL) structure (Fig. 5M–P). The articulamentum was composed of two layers in all investigated species. The dorsal sublayer was Ho structure, and the ventral sublayer was mainly CP structure but partly Ho structure (Fig. 5Q, R). In *Cryptoc. stelleri*, the dorsal sublayer often contained irregular simple prismatic (ISP) structure (Fig. 5S). The myostracum in all investigated species was observed between the

Table 3. Characteristics of radula.

clade	species name	central tooth	centro-lateral tooth		major lateral tooth	
		shape	shape	accessory process	cuspid shape	petaloid process
outgroup	<i>Leptochiton aequispinus</i>	rectangular	envelope-like	present	bicuspidate	
	<i>Leptochiton hakodatensis</i>	cup-shaped	winglike		bicuspidate	
<i>Callochiton</i>	<i>Callochiton foveolatus</i>	bulbous	fungiform		tricuspidate	
clade A: some Chitonina, including the Chitoninae part of Chitonidae	<i>Ischnochiton paululus</i>	bulbous	fingerlike		bicuspidate	cornered
	<i>Ischnochiton poppei</i>	spatulate	fungiform		bicuspidate	cornered
	<i>Ischnochiton hayamii</i>	bulbous	fingerlike		bicuspidate	cornered
	<i>Ischnochiton boninensis</i>	bulbous	fingerlike		bicuspidate	cornered
	<i>Ischnochiton comptus</i>	bulbous	fingerlike		bicuspidate	cornered
	<i>Ischnochiton manazuruensis</i>	bulbous	fingerlike		bicuspidate	cornered
	<i>Rhyssoplax kurodai</i>	hooked-needle	sphenoid	present	patellar	nodulous
	<i>Rhyssoplax komaiana</i>	hooked-needle	sphenoid	present	patellar	nodulous
	<i>Stenoplax alata</i>	bulbous	fingerlike	present	tricuspidate	cornered
	<i>Ischnochiton hakodadensis</i>	bulbous	fingerlike	present	bicuspidate	cornered
	<i>Tripoplax albrechti</i>	cup-shaped	columnar	present	bicuspidate	securiform
	<i>Lepidozona coreanica</i>	cup-shaped	fingerlike	present	bicuspidate	securiform
	<i>Callistochiton jacobaeus</i>	cup-shaped	fingerlike	present	unicuspidate	cornered
clade B: the Tonicinae + Acanthopleurinae part of Chitonidae	<i>Acanthopleura gemmata</i>	bulbous	fingerlike	present	patellar	cornered
	<i>Liolophura japonica</i>	bulbous	fingerlike	present	patellar	cornered
	<i>Acanthopleura loochooana</i>	bulbous	fingerlike	present	patellar	cornered
	<i>Lucilina amanda</i>	bulbous	sphenoid	present	patellar	securiform
	<i>Lucilina lamellosa</i>	bulbous	sphenoid	present	patellar	securiform
	<i>Onithochiton hirasei</i>	bulbous	sphenoid	present	patellar	cornered
clade C: Mopalioidae	<i>Tonicella lineata</i>	cup-shaped	fingerlike		tricuspidate	
	<i>Tonicella zotini</i>	spatulate	fingerlike		tricuspidate	
	<i>Mopalia retifera</i>	rectangular	columnar		tricuspidate	
	<i>Cryptochiton stelleri</i>	deltoid	fungiform		tricuspidate	
	<i>Placiphorella stimpsoni</i>	rectangular	envelope-like		tricuspidate	
	<i>Schizoplax brandtii</i>	cup-shaped	winglike		tricuspidate	
clade D: Cryptoplacoidea	<i>Acanthochitona defilippii</i>	rectangular	columnar		tricuspidate	
	<i>Acanthochitona achates</i>	rectangular	columnar		tricuspidate	
	<i>Acanthochitona dissimilis</i>	rectangular	columnar		tricuspidate	
	<i>Notoplax conica</i>	rectangular	envelope-like		tricuspidate	
	<i>Cryptoplax japonica</i>	spatulate	columnar		tricuspidate	

ventral mesostracum and the hypostracum and/or beneath the hypostracum. This layer was mainly constructed with Ho structure (Fig. 5T, U) but sometimes contained ISP structure (Fig. 5V). The hypostracum was constructed with CL structure (Fig. 5W, X). The apophyses, the inner shell layers extending beyond the tegmentum, were composed of three layers: articulamentum, ventral mesostracum, and myostracum (Fig. 4).

Aesthete canals and complexes

Multiple-branch aesthete canals, which are a type of aesthete canals and complexes and widely distributed in all valves (Baxter and Jones, 1981, 1984; Currie, 1992), were observed in the tegmentum of all investigated species, except *Cryptoc. stelleri*. The shapes of the megal aesthete chambers in the aesthete complexes were observed and

grouped into four types (Fig. 6). Type A had a cylindrical shape shortened more or less and was observed in members of clades A, B, and D. Type B had a sprawling shape into which microaesthete canals often merged before entering and was observed only in clade C. Type C had a wider shape in the middle and was observed in *Callochiton* and members of clades A and B. Type D had a bud-like shape with a stalk-like aesthete canal and was observed only in the outgroups. Table 4 lists the characteristics of the shell layers, shell microstructures, and megal aesthete chamber type.

The above results were listed and shown in relation to the phylogenetic tree in Fig. 7.




Fig. 4. Schematic of valve section. A: section along antero-posterior axis for species of *Callochiton* and clades A and B; B: clades C and D; C: *Cryptochiton stelleri*. D, E, F: close view of the box in the figure. In *Placiphorella stimpsoni*, the tegmentum was composed of two layers, homogeneous and composite prismatic structures.

DISCUSSION

Molecular phylogeny

The ML and Bayesian trees in the present study were highly robust, though the phylogenetic positions of *Ischnochiton* spp. and *Cryptoc. stelleri* were unclear. The topologies of these trees were predominantly similar to those in Irisarri et al. (2014, 2020), which reported that Chitonoida (Chitonina) formed a sister group with a clade comprising Mopalioida plus Cryptoplacoidea (Acanthochitonina), but not to those in Okusu et al. (2003). Additionally, the phylogenetic position of *Callochiton* in the present study corre-

sponded to those in Sigwart et al. (2010), Irisarri et al. (2014, 2020), and Moles et al. (2021), but not to that in Sigwart et al. (2013), which described that *Callochiton* formed a sister group with a clade comprising Mopalioida plus Cryptoplacoidea. The phylogenetic position of *Cryptoc. stelleri* in the present study made Acanthochitonidae and Mopaliidae polyphyletic groups. This position corresponded to those reported by Kelly and Eernisse (2008) and Irisarri et al. (2014), and the latter study considered *Cryptoc. stelleri* to be a member of Mopaliidae. Therefore, the present study also grouped *Cryptoc. stelleri* into Mopaliidae (Figs. 2, 7). *Ischnochiton hakodadensis*, the only species possessing an




Fig. 5. SEM images of shell microstructures. Dorsal and ventral sublayers of tegumentum in (A, B) *Callochiton foveolatus*, (C, D) *Ischnochiton comptus*, (E, F) *Liolophura japonica*, (G, H) *Placiphorella stimpsoni*. Tegumentum in (I) *Leptochiton aequispinus*, (J) *Tonicella zotini*, (K) *Schizoplax brandtii*, (L) *Acanthochiton defilippii*. Dorsal mesostracum in (M) *Callo foveolatus*, (N) *Stenoplax alata*. Ventral mesostracum in (O) *Sc. Brandtii*, (P) *Cryptoplax japonica*. Dorsal and ventral sublayers of articulamentum in *Lucilina lamellose* (Q, R). Dorsal sublayer of articulamentum in *Cryptochiton stelleri* (S). Myostracum between the ventral mesostracum and hypostracum in *Lepidozona coreanica* (T). Myostracum beneath the hypostracum in (U) *Tonicella lineata*, (V) *Rhyssoplax komaiana*. Hypostracum in (W) *I. boninensis*, (X) *Acanthoc. aches*. Ho: homogeneous structure; CP: composite prismatic structure; CL: crossed-lamellar structure; ISP: irregular simple prismatic structure. Scale bar indicates 10 μm . The upper direction of SEM images is dorsal side, and the lower direction is ventral side.

accessory process in the genus, made *Ischnochitonidae* a polyphyletic group, as reported by Owada (2016, 2018). Furthermore, the present study clarified that *Rhyssoplax*, which was classified as *Chitonidae* by Sirenko (2006), formed a sister group with *Ischnochiton*. However, the phylogenetic

position of *Rhyssoplax* did not correspond to that reported by Irisarri et al. (2020). The DNA sequence of *Rhyssoplax* used by Irisarri et al. (2020) was determined by Riesgo et al. (2012). If their identification was correct, it would be suggested that *Rhyssoplax* is a polyphyletic group. The present




Fig. 6. Megal aesthete chamber type in multiple-branch aesthete canals. (A) type A in *Acanthopleura gemmata*, (B) type B in *Placiphorella stimpsoni*, (C) type C in *Ischnochiton comptus*, (D) type D in *Leptochiton hakodatensis*. ac: aesthete canal, mc: megal aesthete chamber. White arrowhead indicates the aperture of megal aesthete canal, and black arrowhead indicates that of microaesthete canal. Scale bar indicates 100 µm in (A), 50 µm in (B, C, D).

study also clarified that *Lepidozonia coreanica* (Reeve, 1847), a common species along the coast of Japan, formed a sister group with *Callistochiton jacobaeus* (Gould, 1859) (Figs. 2, 7). These results indicate the need to reexamine the classification within Chitonina.

Radula

Saito (2004) suggested that the characteristics of the radula in chitons were useful not only for species identification but also for the assessment of the phylogenetic relationships at higher levels, emphasizing examples of Cryptoplacoidea in his comparisons. The present study indicated that the characteristics of the radula could identify the species (Table 3, Fig. 7), and support Saito (2004). The shapes of the central and centro-lateral teeth varied among species and genera and could be used for identification at least Japanese chiton species at those levels. Likewise, the shape of the petaloid process is informative at genus level, and considering all these tooth shapes helps to confirm identification. In contrast, the presence of accessory and petaloid processes and the major lateral cusp shape

Table 4. Characteristics of shell layers, shell microstructures, and megal aesthete chamber type.

clade	species name	tegumentum		dorsal	articulamentum		ventral	myostracum	hypostracum	megal aesthete chamber type
		dorsal	ventral	mesostracum	dorsal	ventral	mesostracum			
outgroup	<i>Leptochiton aequispinus</i>	Ho	Ho	CL	Ho	CP	CL	Ho	CL	D
	<i>Leptochiton hakodatensis</i>	Ho	Ho	CL	Ho	CP	CL	Ho	CL	D
<i>Callochiton</i>	<i>Callochiton foveolatus</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
clade A: some Chitonina, including the Chitoninae part of Chitonidae	<i>Ischnochiton paululus</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Ischnochiton poppei</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Ischnochiton hayamii</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Ischnochiton boninensis</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Ischnochiton comptus</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Ischnochiton manazuruensis</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Rhyssoplax kurodai</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Rhyssoplax komaiana</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Stenoplax alata</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Ischnochiton hakodatensis</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Tripoplax albrechti</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Lepidozonia coreanica</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Callistochiton jacobaeus</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
clade B: the Tonicinae + Acanthopleurinae part of Chitonidae	<i>Acanthopleura gemmata</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Liolophura japonica</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Acanthopleura loochooana</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
	<i>Lucilina amanda</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Lucilina lamellosa</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	C
	<i>Onithochiton hirasei</i>	Ho	CP	CL	Ho	CP	CL	Ho	CL	A
clade C: Mopaliioidea	<i>Tonicella lineata</i>	Ho	Ho		Ho	CP	CL	Ho	CL	B
	<i>Tonicella zotini</i>	Ho	Ho		Ho	CP	CL	Ho	CL	B
	<i>Mopalia retifera</i>	Ho	Ho		Ho	CP	CL	Ho	CL	B
	<i>Cryptochiton stelleri</i>				Ho	CP	CL	Ho	CL	
	<i>Placiphorella stimpsoni</i>	Ho	CP		Ho	CP	CL	Ho	CL	B
	<i>Schizoplax brandtii</i>	Ho	Ho		Ho	CP	CL	Ho	CL	B
	<i>Schizoplax brandtii</i>	Ho	Ho		Ho	CP	CL	Ho	CL	B
clade D: Cryptoplacoidea	<i>Acanthochitona defilippii</i>	Ho	Ho		Ho	CP	CL	Ho	CL	A
	<i>Acanthochitona achates</i>	Ho	Ho		Ho	CP	CL	Ho	CL	A
	<i>Acanthochitona dissimilis</i>	Ho	Ho		Ho	CP	CL	Ho	CL	A
	<i>Notoplax conica</i>	Ho	Ho		Ho	CP	CL	Ho	CL	A
	<i>Cryptoplax japonica</i>	Ho	Ho		Ho	CP	CL	Ho	CL	A




Fig. 7. Relationships among the molecular phylogenetic trees, the genera or species, the character state of radula, shell microstructures and megalaesthete chamber type, and the classification in the higher taxon. p.: patellar, Ho: homogeneous structure, CP: composite prismatic structure.

were found to be relatively conservative, but the cusp shape can be somewhat useful for distinguishing genera.

Thiele (1893) described the petaloid process as a diagnostic feature for groups corresponding to Chitonoida in the present study, but until now it had not been tested with molecular phylogenetic analysis. Based on molecular phylogenetic analysis, the present study indicated that the petaloid process only occurs in Chitonina. The accessory process is likely convergent in Chitonina and *Lept. aequispinus* (Fig. 7) because it was not found in other Lepidopleurida, Callochitonida, or Acanthochitonina within Chitonida.

Shell microstructures

Bergenhayn (1930) reported that chiton valves consisted of four shell layers (i.e., tegmentum, articulamentum, hypostracum, and mesostracum), excluding the periostracum. However, Haas (1972, 1976, 1981) did not distinguish between articulamentum and hypostracum and described chiton valves as being constructed from three shell layers (i.e., tegmentum, hypostracum, and myostracum). Poulicek and Kreuzsch (1986) supported Bergenhayn (1930) but did not recognize mesostracum, and Carter (1990) supported Haas (1972, 1976, 1981). Connors et al. (2012) defined seven shell layers: tegmentum, dorsal mesostracum, articulamentum, ventral mesostracum, anterior and posterior myostracums, and hypostracum; *Tonicella marmorea* (O. Fabricius, 1780) was an exception in lacking the dorsal mesostracum. The present study adopted the definition of Connors et al. (2012) but did not distinguish between anterior and posterior myostracums because they did not appear to vary in the assembled information. Peebles et al. (2017) reported that the valves of chitons were composed of four to seven shell layers according to Connors et al. (2012) and suggested that the

properties of the shell layers did not clearly reflect the phylogenetic relationships because of the large species variations. The present study indicated that the valves were constructed of four to six layers and that the tegmentum and articulamentum had sublayers. Additionally, the present study has extended the observation of Connors et al. (2012) that the dorsal mesostracum layer is absent in *To. marmorea*; it is also absent in the seven other Acanthochitonina genera, and other species of *Tonicella* examined in the present study (Fig. 7). The congruent distribution relative to the molecular result implies that a shared absence of the dorsal mesostracum layer is a synapomorphy of Acanthochitonina (Table 4, Fig. 7).

Differences in the shell microstructures were observed only in the ventral sublayer of the tegmentum. In *Callochiton*, Chitonina, and *P. stimpsoni*, the sublayer was mainly constructed with CP structure, and with Ho structure in the others. It is likely that a clade comprising *Callochiton* and Chitonida first obtained the Ho structure in the sublayer because the two species of *Leptochiton*, which are outgroups, had Ho structure. Presumably, the CP structure was independently acquired in *Callochiton*, Chitonina, and *P. stimpsoni*. In *Cryptoc. stelleri*, the structure of the valve differed greatly from the others because the tegmentum was not observed. This may be because the valves are always embedded in the soft part. Peebles et al. (2017) distinguished the CL structure in the mesostracum or hypostracum into at least two types, based on the size of the fibers that made the CL structure. However, the present study did not distinguish such types of CL structures because the difference between CP and Ho structures is much larger than that between the types of CL structures. Furthermore, individual variation in the size of the fibers was often observed.

Aesthete canals and complexes

Baxter and Jones (1981, 1984) and Currie (1992) observed aesthete canals and complexes in detail, and Fernandez et al. (2007) and Vendrasco et al. (2008) additionally constructed the phylogenetic relationships using their characteristics. Their phylogenetic conclusions are consistent with the present study. In the present study, the megal aesthete chamber type, which was applied as one of the characters for the phylogenetic analysis in Fernandez et al. (2007) and Vendrasco et al. (2008), was available for identification of particular families or superfamilies. However, in Chitonina, the variation in the type was relatively high compared with the others. Although the megal aesthete chamber type may be related to the life type, both *Acanthopleura* living on the surface of a rock in the intertidal zone and having a lens on the surface of the valve, and *Lepidozona* living on the back of a rock in the subtidal zone and lacking a lens, possessed the type A.

Conclusions

In the radula, the shapes of the central and centro-lateral teeth and the petaloid process were useful for the identification of particular species or genera. The presence of accessory and petaloid processes and the cusp shape were available for recognizing particular genera or even suborders. In the valves, the number of shell layers and the shell microstructure type in the ventral sublayer of tegmentum appear to be a synapomorphy shared by the members of suborder Acanthochitonina, and the megal aesthete chamber type was helpful for identifying particular families or superfamilies. These characteristics of the valve could also be useful for verifying the fossil records of chitons. The molecular phylogenetic tree implied that the above characteristics were significant and simultaneously indicated that the classification within Chitonina would need to be reexamined because the variations of the cusp shape and megal aesthete chamber type did not correspond to the current classification. *Callochiton*, which is contained in Callochitonida in the current classification, would be equally closely related to any member of Acanthochitonina and any member of Chitonina because of possessing a mosaic of characteristics from both. Buckland-Nicks and Hodgson (2000) investigated the characteristics of the gametes in *Callochiton castaneus* (W. Wood, 1815) treated as *Callochiton dentatus* (Spengler, 1797) in MolluscaBase editors (2021), and reported that *Callo. castaneus* was basal to Chitonida and closely related to both Chitonina and Acanthochitonina. This result is consistent with the present study.

The radula and shell microstructures have been broadly conserved in chitons living in the shallow sea. Especially, shell microstructure characters are among the relatively few morphological features that can be observed in fossil chitons. They were found to be phylogenetically informative at both geologically ancient and recent levels.

ACKNOWLEDGMENTS

I deeply thank Professor Akiya Hino (Kanagawa University) and Professor Hiroshi Hosoya (Kanagawa University) for letting me use the laboratory for DNA sequencing. I am grateful to Associate Professor Michiko Nemoto (Okayama University), Dr. Masaya Saito (Kanagawa University), and Mr. Shoichi Hamano (Hokkaido University)

for helping to collect the samples. I am indebted to Professor Douglas J. Eernisse (California State University Fullerton) for fruitful comments significantly improving the manuscript.

COMPETING INTERESTS

The author has no competing interests to declare.

REFERENCES

- Baxter JM, Jones AM (1981) Valve structure and growth in the Chiton *Lepidochiton cinereus* (Polyplacophora: Ischnochitonida). *J Mar Biol Ass UK* 61: 65–78
- Baxter JM, Jones AM (1984) The valve morphology of *Callochiton achatinus* (Mollusca: Polyplacophora: Ischnochitonidae). *J Zool* 202: 549–560
- Bergenhayn JRM (1930) Die Loricaten von Dr. Sixten Bocks Pazifik-Expedition 1917–1918, mit spezieller Berücksichtigung der Perinotumbildungen und der Schalenstruktur. Göteborgs Kungl Vetensk Samhälles Handl, Ser. B, (12) 1: 1–52, pls 1–3
- Bergenhayn JRM (1955) Die fossilen schwedischen Loricaten nebst einer vorläufigen Revision des Systems der ganzen Klasse Loricata. *Lunds Univ Arsskr N F Adv* 2 51: 1–41
- Buckland-Nicks J, Hodgson AN (2000) Fertilization in *Callochiton castaneus* (Mollusca). *Biol Bull* 199: 59–67
- Bullock RC (1988) The genus *Chiton* in the New World (Polyplacophora: Chitonidae). *Veliger* 31: 141–191
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool for automated alignment trimming in large scale phylogenetic analyses. *Bioinformatics* 25: 1972–1973
- Carter JG, Ed (1990) *Skeletal Biomineralization: Patterns, Processes and Evolutionary Trends*, 2 vols. Van Nostrand Reinhold, New York
- Connors MJ, Ehrlich H, Hog M, Godefroy C, Araya S, Kallai I, et al. (2012) Three-dimensional structure of the shell plate assembly of the chiton *Tonicella marmorea* and its biomechanical consequences. *J Struct Biol* 177: 314–328
- Currie DR (1992) Aesthete channel morphology in three species of Australian chitons (Mollusca: Polyplacophora). *J Malacol Soc Australia* 13: 3–14
- Dall WH (1879) Report on the limpets and chitons of the Alaskan and Arctic regions, with descriptions of genera and species believed to be new. *Proc US Natl Mus* 1: 281–344, pls 1–5
- Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T (2020) ModelTest-NG: a new and scalable tool for the selection of DNA and protein evolutionary models. *Mol Biol Evol* 37: 291–294
- Distel DL, Amin M, Burgoyne A, Linton E, Mamangkey G, Morrill W, et al. (2011) Molecular phylogeny of Pholadoidea Lamarck, 1809 supports a single origin for xylophagy (wood feeding) and xylophagous bacterial endosymbiosis in Bivalvia. *Mol Phylogenet Evol* 61: 245–254
- Eernisse DJ, Reynolds PD (1994) Polyplacophora. In “Microscopic Anatomy of Invertebrates. Volume 5, Mollusca 1” Ed by FW Harrison, AJ Kohn, Wiley-Liss, New York, pp 56–110
- Fernandez CZ, Vendrasco MJ, Runnegar B (2007) Aesthete canal morphology in twelve species of chiton (Polyplacophora). *Am Malacol Bull* 25: 51–69
- Giribet G, Carranza S, Baguña J, Riutort M, Ribera C (1996) First molecular evidence for the existence of a Tardigrada + Arthropoda clade. *Mol Biol Evol* 13: 76–84
- Haas W (1972) Untersuchungen über die Mikro- und Ultrastruktur der Polyplacophorenschale. *Biomineralisation* 5: 3–52
- Haas W (1976) Observations on the shell and mantle of the placophora. In “The Mechanisms of Mineralization in the Invertebrates and Plants” Ed by N Watabe, KM Wilbur, University of South Carolina Press, Columbia, pp 389–402
- Haas W (1981) Evolution of calcareous hardparts in primitive mol-

- luscs. *Malacologia* 21: 403–418
- Irisarri I, Eernisse DJ, Zardoya R (2014) Molecular phylogeny of Acanthochitonina (Mollusca: Polyplacophora: Chitonida): three new mitochondrial genomes, rearranged gene orders and systematics. *J Nat Hist* 48: 2825–2853
- Irisarri I, Uribe JE, Eernisse DJ, Zardoya R (2020) A mitogenomic phylogeny of chitons (Mollusca: Polyplacophora). *BMC Evol Biol* 20: 22
- Kaas P, Van Belle RA (1980) Catalogue of Living Chitons (Mollusca, Polyplacophora). 2nd revised ed, Backhuys Publishers, Leiden
- Kaas P, Van Belle RA (1990) Monograph of Living Chitons (Mollusca: Polyplacophora), Vol. 4. Suborder Ischnochitonina: Ischnochitonidae: Ischnochitoninae (continued). Additions to Vols 1, 2 and 3. EJ Brill, Leiden
- Kaas P, Van Belle RA (1994) Monograph of Living Chitons (Mollusca: Polyplacophora), Vol. 5. Suborder Ischnochitonina: Ischnochitonidae: Ischnochitoninae (concluded), Callistoplacinae; Mopaliidae. Additions to Vols 1–4. EJ Brill, Leiden
- Katoh K, Kuma K, Toh H, Miyata T (2005) MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res* 33: 511–518
- Kelly RP, Eernisse DJ (2008) Reconstructing a radiation: the chiton genus *Mopalia* in the North Pacific. *Invert Syst* 22: 17–28
- Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A (2019) RAxML-NG: A fast, scalable, and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35: 4453–4455
- Moles J, Cunha TJ, Lemer S, Combosch DJ, Giribet G (2021) Tightening the girdle: phylotranscriptomics of Polyplacophora. *J Molluscan Stud* 87: eyab019
- MolluscaBase editors (2021) MolluscaBase. Polyplacophora. Accessed at: <http://molluscabase.org/aphia.php?p=taxdetails&id=55> on 24 Jan 2023
- Okusu A, Schwabe E, Eernisse DJ, Giribet G (2003) Towards a phylogeny of chitons (Mollusca, Polyplacophora) based on combined analysis of five molecular loci. *Org Divers Evol* 3: 281–302
- Owada M (2016) A new cryptic species distinguished from *Ischnochiton comptus* (Gould, 1859) (Polyplacophora: Ischnochitonidae) in central Honshu, Japan. *Molluscan Res* 36: 255–263
- Owada M (2018) Phylogenetic relationships among Japanese species of the genus *Ischnochiton* (Polyplacophora: Ischnochitonidae), including a new species. *Zool Sci* 35: 281–291
- Owada M, Kasai Y, Sato S, Hong JS (2013) Shell morphology and genetic characteristics of *Corbula* (*Potamocorbula*) spp. (Bivalvia: Corbulidae) from the waters adjacent to Japan, Korea and San Francisco Bay. *Aquat Biol* 19: 297–305
- Peebles BA, Smith AM, Spencer HG (2017) Valve microstructure and phylomineralogy of New Zealand chitons. *J Struct Biol* 197: 250–259
- Pojeta J, Dufoe J (2008) New information about *Echinochiton dufoei*, the Ordovician spiny chiton. *Amer Malac Bull* 25: 25–34
- Pojeta J, Eernisse DJ, Hoare RD, Henderson MD (2003) *Echinochiton dufoei*: A new spiny Ordovician chiton. *Jour Paleont* 77: 646–654
- Pojeta J, Vendrasco MJ, Darrrough G (2010) Upper Cambrian chitons (Mollusca, Polyplacophora) from Missouri, USA. *Bull Am Paleontol* 379: 1–82
- Ponder WF, Lindberg DR (2020) Biology and Evolution of the Mollusca. Vol 2. CRC Press, Boca Raton
- Poulicek E, Kreusch B (1986) Evolutionary trends in the skeletal structures of the Polyplacophora. Proceedings of the 8th International Malacological Congress, Budapest, 207–212
- Puchalski SS, Eernisse DJ, Johnson CC (2008) The effect of sampling bias on the fossil record of chitons (Mollusca, Polyplacophora). *Am Malacol Bull* 25: 87–95
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Syst Biol* 67: 901–904
- Riesgo A, Andrade SC, Sharma PP, Novo M, Perez-Porro AR, Vahtera V, et al. (2012) Comparative description of ten transcriptomes of newly sequenced invertebrates and efficiency estimation of genomic sampling in non-model taxa. *Front Zool* 9: 33
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574
- Runnegar B, Pojeta J, Taylor ME, Collins D (1979) New species of the Cambrian and Ordovician chitons *Mathevia* and *Chelodes* from Wisconsin and Queensland: Evidence for the early history of polyplacophoran mollusks. *J Paleontol* 53: 1374–1394
- Saito H (1994) The shallow-water chiton fauna of eastern Hokkaido, Japan. *Mem Natn Sci Mus Tokyo* 27: 93–104
- Saito H (1995) The chiton fauna of Onagawa Bay, northeastern Honshu, Japan. *Mem Natn Sci Mus Tokyo* 28: 99–112
- Saito H (1998) The chitons (Mollusca: Polyplacophora) of the Goto islands, west coast of Kyushu, Japan. *Mem Natn Sci Mus Tokyo* 31: 145–163
- Saito H (2000) The chitons (Mollusca: Polyplacophora) of the Seto Inland Sea, Japan. *Mem Natn Sci Mus Tokyo* 33: 97–104
- Saito H (2001) Shallow-water chitons (Mollusca: Polyplacophora) of Hachijo Island, Izu Islands, Japan. *Mem Natn Sci Mus Tokyo* 37: 193–202
- Saito H (2004) Phylogenetic significance of the radula in chitons, with special reference to the Cryptoplacoidea (Mollusca: Polyplacophora). *Boll Malacol Suppl* 5: 83–104
- Saito H (2006) A preliminary list of chitons (Mollusca: Polyplacophora) from Sagami Sea. *Mem Natn Sci Mus Tokyo* 40: 203–224
- Saito H (2011) Chitons (Mollusca: Polyplacophora) from submarine banks off Izu Islands and Bōsō Peninsula, Japan. *Mem Natn Sci Mus Tokyo* 47: 65–81
- Saito H (2017) Class Polyplacophora. In “Marine Mollusks in Japan. 2nd ed” Ed by T Okutani, Tokai University Press, Tokyo, pp 728–738, pls 2–11
- Sasaki T (2010) Malacology. University of Tokyo Press, Tokyo
- Schwabe E (2005) A catalogue of Recent and fossil chitons (Mollusca: Polyplacophora) Addenda. *Novapex* 6: 89–105
- Schwabe E (2010) Illustrated summary of chiton terminology. *Spixiana* 33: 171–194
- Sigwart JD, Schwabe E, Saito H, Samadi S, Giribet G (2010) Evolution in the deep sea: a combined analysis of the earliest diverging living chitons (Mollusca: Polyplacophora: Lepidopleurida). *Invertebr Syst* 24: 560–572
- Sigwart JD, Stoeger I, Knebelberger T, Schwabe E (2013) Chiton phylogeny (Mollusca: Polyplacophora) and the placement of the enigmatic species *Chorioplax grayi* (H. Adams & Angas). *Invertebr Syst* 27: 603–621
- Sirenko B (2006) New outlook on the system of chitons (Mollusca: Polyplacophora). *Venus* 65: 27–49
- Sirenko B (2020) New Zealand and Australian species of the genus *Leptochiton* (Mollusca: Polyplacophora). *Zootaxa* 4851: 401–449
- Sirenko B, Zhang J (2019) Chitons (Mollusca: Polyplacophora) of Hainan Island and vicinity, South China Sea. *Zootaxa* 4564: 1–40
- Slieker FJA (2000) Chitons of the World – An Illustrated Synopsis of Recent Polyplacophora. L’Informatore Piceno, Ancona
- Smith AG (1960) Amphineura. In “Treatise on Invertebrate Paleontology. Part I, Mollusca 1” Ed by RC Moore, University of Kansas and Geological Society of America, Boulder and Lawrence, pp 41–76
- Taki Is (1938) Report of the biological survey of Mutsu Bay 31. Studies on chitons of Mutsu Bay with general discussion on chitons of Japan. *Sci Rep Tohoku Imp Univ Ser 4* 12: 323–423

- Taki Is, Taki Iw (1929a) Studies on Japanese Chitons (1). *Venus* 1: 47–54
- Taki Is, Taki Iw (1929b) Studies on Japanese Chitons (2). *Venus* 1: 141–153
- Taki Is, Taki Iw (1929c) Studies on Japanese Chitons (3). *Venus* 1: 157–164
- Taki Is, Taki Iw (1930) Studies on Japanese Chitons (4). *Venus* 2: 99–107
- Taki Is, Taki Iw (1931a) Studies on Japanese Chitons (5). *Venus* 2: 178–185
- Taki Is, Taki Iw (1931b) Studies on Japanese Chitons (6). *Venus* 2: 215–220
- Tanabe AS (2011) Kakusan4 and Aminosan: two programs for comparing nonpartitioned, proportional, and separate models for combined molecular phylogenetic analyses of multilocus sequence data. *Mol Ecol Resour* 11: 914–921
- Taylor JD, Kennedy JM, Hall A (1969) The shell structure and mineralogy of the Bivalvia. Introduction. *Nuculacea – Trigonacea*. *Bull Br Mus Nat Hist, Zool, Suppl* 3: 1–125
- Taylor JD, Kennedy JM, Hall A (1973) The shell structure and mineralogy of the Bivalvia. *Lucinacea – Clavagellacea*. *Conclusions*. *Bull Br Mus Nat Hist, Zool* 22: 253–294
- Thiele J (1893) Polyplacophora, Lepidoglossa, Schuppenzüngler. In "Das Gebiss der Schnecken zur Begründung einer natürlichen Classification. Vol. 2" Ed by FH Troschel, Berlin, Nicolaische Verlagsbuchhandlung, pp 336–409, pls 29–32
- Thiele J (1909–1910) Revision des Systems der Chitonen. *Zool Stuttg* 22:1–70, pls 1–6 (1909); 71–132, pls 7–10 (1910)
- Vendrasco MJ, Fernandez CZ, Eernisse DJ, Runnegar B (2008) Aesthete canal morphology in the Mopaliidae (Polyplacophora). *Am Malacol Bull* 25: 51–69
- Whiting, MF (2002) Mecoptera is paraphyletic: multiple genes and phylogeny of Mecoptera and Siphonaptera. *Zool Scr* 31: 93–104
- Whiting MF, Carpenter JM, Wheeler QD, Wheeler WC (1997) The Strepsiptera problem: phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. *Syst Biol* 46: 1–68
- Xiong B, Kocher TD (1991) Comparison of mitochondrial DNA sequences of seven morphospecies of black flies (Diptera: Simuliidae). *Genome* 34: 306–311

(Received July 21, 2022 / Accepted June 29, 2023 /
Published online August 28, 2023)