



Mysida and Lophogastrida of Greece: a preliminary checklist

Panayota Koulouri[‡], Vasilis Gerovasileiou[‡], Nicolas Bailly[‡]

[‡] Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine Research, Heraklion, Greece

Corresponding author: Panayota Koulouri (yo172@hcmr.gr)

Academic editor: Christos Arvanitidis

Received: 20 May 2016 | Accepted: 17 Jul 2016 | Published: 01 Nov 2016

Citation: Koulouri P, Gerovasileiou V, Bailly N (2016) Mysida and Lophogastrida of Greece: a preliminary checklist. Biodiversity Data Journal 4: e9288. <https://doi.org/10.3897/BDJ.4.e9288>

Abstract

Background

The checklist of Mysida and Lophogastrida of Greece was created within the framework of the Greek Taxon Information System (GTIS), which is one of the applications of the LifeWatchGreece Research Infrastructure (ESFRI) resuming efforts to develop a complete checklist of species recorded and reported from Greek waters. The objectives of the present study were to update and cross-check taxonomically all records of Mysida and Lophogastrida species known to occur in Greek waters in order to search for inaccuracies and omissions.

New information

The up-to-date checklist of Mysida and Lophogastrida of Greece comprises 49 species, classified to 25 genera.

Keywords

Mysida, Lophogastrida, Greece, Aegean Sea, Sea of Crete, Ionian Sea, Eastern Mediterranean, checklist

Introduction

The peracarid crustaceans Lophogastrida, Stygiomysida and Mysida were formerly grouped under the order "Mysidacea". Although these three monophyletic groups are now ranked as distinct orders of Peracarida (Meland and Willassen 2007, Meland et al. 2015), we have chosen to present mysids and lophogastrids under the same checklist, while no stygiomysids have been recorded so far from Greek waters. The most important contribution to the Greek records of the taxonomic group of "Mysidacea" has been made by Hatzakis (Hatzakis 1974, Hatzakis 1977, Hatzakis 1978, Hatzakis 1982) followed later on by Madurell and Cartes (2003) as well as Koulouri et al. (2013). "Mysidacea" constitute the dominant faunal component for the so-called hyperbenthos, which includes all swimming bottom-dependent organisms (mainly crustaceans) performing, with varying amplitude, intensity and regularity, seasonal or daily vertical migrations into the water column (Brunel et al. 1978). A large proportion of these animals are living within the benthic boundary layer (BBL) which is the portion of the sediment and the water column that is influenced by the sediment-water interface and characterized by strong gradients of flow as well as high concentrations of dissolved and particulate organic matter (Boudreau and Jørgensen 2001). Sampling these often highly mobile animals is not easy as they are caught occasionally by conventional benthic (e.g., grabs, corers) or pelagic (e.g. plankton nets) sampling gears (Eleftheriou 2013). Specially designed samplers, generally known as hyperbenthic or suprabenthic sledges, have been constructed, modified and used over the last 40 years for sampling these macrofaunal organisms, but there are still practical technical difficulties in the accessibility of this particular habitat (Mees and Jones 1997, Eleftheriou 2013, Koulouri et al. 2013). However, and when depth allows, these organisms can also be sampled by scuba diver operated "hunting" tools such as hand nets (e.g. Ledoyer 1968) or suction bottles (e.g. Chevaldonné et al. 2008).

A first attempt for creating a checklist of Mysida and Lophogastrida under the order of "Mysidacea" was carried out in the context of the "Greek Biodiversity Database" project (2005-2008), coordinated by the Department of Zoology, at the Aristotle University of Thessaloniki. The documented occurrence of these marine species in Greek waters was recorded in a database that was set up online in 2010. Although not completed, the database had not been updated since the closure of the project (2010). During the European project PESI, [WoRMS/ERMS](#) specially created the reference Koukouras (2010) for the list of marine species provided by the Greek Biodiversity Database. The aim of the present work was to present an updated checklist of Lophogastrida and Mysida of the Greek waters. For this purpose, older lists were updated and annotated according to the recent literature and current taxonomic status of the species.

Materials and methods

The Checklist of Mysida and Lophogastrida of Greece (Suppl. material 1) was created within the framework of the Greek Taxon Information System (GTIS), one of the applications of the [LifeWatchGreece](#) Research Infrastructure (ESFRI), aiming at developing a complete checklist of all species reported from Greek waters (Bailly et al. 2016, this special collection). The general principles used for elaborating this Preliminary Checklist are given in Bailly et al. (2016). The checklist of Mysida and Lophogastrida of Greece was constructed based on the species records extracted from the dataset of WoRMS/ERMS for marine species. Then, all recent publications were reviewed and the species reported to date have been added to the list. The classification followed in the present checklist is the one proposed by the World List of Lophogastrida, Stygiomysida and Mysida (Mees and Meland 2012).

Checklist of Mysida and Lophogastrida known to occur in Greek waters

Order Lophogastrida

Family Eucopiidae

Eucopia unguiculata (Willemoes-Suhm, 1875)

Family Lophogastridae

Lophogaster typicus M. Sars, 1857

Order Mysida

Family Mysidae

Acanthomysis longicornis (Milne Edwards, 1837)

Anchialina agilis (G.O. Sars, 1877)

Anchialina oculata Hoenigman, 1960

Boreomysis arctica (Krøyer, 1861)

- Boreomysis megalops* G.O. Sars, 1872
- Calyptomma puritani* W. Tattersall, 1909
- Dactylamblyops corberai* San Vicente & Cartes, 2011
- Diamysis bacescui* Wittmann & Ariani, 1998
- Diamysis cymodoceae* Wittmann & Ariani, 2012
- Diamysis lagunaris* Ariani & Wittmann, 2000
- Diamysis mesohalobia* Ariani & Wittmann, 2000
- Erythroops elegans* (G.O. Sars, 1863)
- Erythroops erythrophthalmus* (Goës, 1864)
- Erythroops neapolitanus* Colosi, 1929
- Euchaetomeropsis merolepis* (Illig, 1908)
- Gastrosaccus mediterraneus* Bacescu, 1970
- Gastrosaccus sanctus* (Van Beneden, 1861)
- Haplostylus bacescui* Hatzakis, 1977
- Haplostylus lobatus* (Nouvel, 1951)
- Haplostylus normani* (G.O. Sars, 1877)
- Harmelinella mariannae* Ledoyer, 1989
- Hemimysis margalefi* Alcaraz, Riera & Gili, 1986
- Heteromysis eideri* Bacescu, 1941
- Leptomysis buergii* Bacescu, 1966

- Leptomysis gracilis* (G.O. Sars, 1864)
- Leptomysis lingvura* (G.O. Sars, 1866)
- Leptomysis mediterranea* G.O. Sars, 1877
- Leptomysis megalops* Zimmer, 1915
- Leptomysis truncata* (Heller, 1863)
- Limnomysis benedeni* Czerniavsky, 1882
- Mesopodopsis aegyptia* Wittmann, 1992
- Mesopodopsis slabberi* (Van Beneden, 1861)
- Mysideis parva* Zimmer, 1915
- Mysidopsis angusta* G.O. Sars, 1864
- Mysidopsis gibbosa* G.O. Sars, 1864
- Paraleptomysis apiops* (G.O. Sars, 1877)
- Paramysis agigensis* Bacescu, 1940
- Paramysis helleri* (G.O. Sars, 1877)
- Paramysis kosswigi* Bacescu, 1948
- Parerythrops paucispinosus* Nouvel & Lagardère, 1976
- Pseudomma nanum* Holt & Tattersall, 1906
- Siriella armata* (Milne Edwards, 1837)
- Siriella clausii* G.O. Sars, 1877
- Siriella gracilipes* Nouvel, 1942

Siriella jaltensis* Czerniavsky, 1868**Siriella norvegica* G.O. Sars, 1869*****Siriella thompsonii* (H. Milne Edwards, 1837)****Discussion**

A total of 49 species, classified in 25 genera, makes up the updated checklist of Lophogastrida and Mysida of Greece. More specifically, the order of Lophogastrida comprises 2 species classified to 2 genera, while the order of Mysida comprises 47 species, classified to 23 genera. It is worth mentioning that *Dactylamblyops corberai* San Vicente & Cartes, 2011, a new mysid species was described from the deep Ionian Sea by San Vicente and Cartes (2011), revealing the limited sampling effort that has taken place for these particular taxa. It should also be noted that Chevaldonné et al. (2015) provided new molecular and distribution data on the poorly known cave mysid *Harmelinella mariannae* Ledoyer, 1989. A number of issues aroused during the thorough examination of the checklist of Mysida of Greece, specifically regarding the presence of the species *Diamysis bahirensis* (Sars G.O., 1877), *Haplostylus magnilobatus* (Bacescu & Schiecke, 1974) and *Paramysis festae* Colosi, 1921 in Greek waters. The material previously indicated as *D. bahirensis* by Hatzakis (1982) has been revised and assigned to *D. lagunaris* and *D. bacescui* (Wittmann and Ariani 1998, Wittmann and Ariani 2012). Therefore, further research is needed to verify the presence of this species in the Greek waters. The species *H. magnilobatus* has been included in the mysid species of the Aegean Sea according to Coll et al. (2010). However, the reference used for this record does not actually include this species in the Greek fauna (Hatzakis 1977). The species *P. festae* has been recorded in the Ionian coast of Sicily and in a brackish lagoon of Libya (Audzijonyte et al. 2008, Wittmann and Ariani 2011), but not yet in Greek waters. Moreover, further research is needed for the verification of the presence of the mysid species recorded by Koulouri et al. (2013) as *Erythrops* cf. *peterdorhni* Basescu & Schiecke, 1974 in the Greek waters. We should also note that when establishing the new rank *Siriella gracilipes* for part of what was called *S. jaltensis*, Wittmann and Wirtz (1998) reported the finding of *S. gracilipes* in the Aegean Sea. Finally, two provisional cryptic species of the complex taxon *Hemimysis margalefi sensu lato* were recorded from marine cave habitats from Corfu and Crete islands by using molecular techniques (Rastorgueff et al. 2014) and therefore, their taxonomic status needs to be further studied. As far as eastern Mediterranean is concerned, Mysida and Lophogastrida have received little attention mostly due to difficulties in sampling these particular taxa especially in oligotrophic shelf areas (Koulouri et al. 2013). However, it is expected that several species distributed in other Mediterranean regions will be recorded as elements of the Greek fauna in the future.

Acknowledgements

This work was supported by the LifeWatchGreece infrastructure (MIS 384676), funded by the Greek Government under the General Secretariat of Research and Technology (GSRT), ESFRI Projects, National Strategic Reference Framework (NSRF).

References

- Audzijonyte A, Daneliya M, Mugue N, Väinölä R (2008) Phylogeny of *Paramysis* (Crustacea: Mysida) and the origin of Ponto-Caspian endemic diversity: Resolving power from nuclear protein-coding genes. *Molecular Phylogenetics and Evolution* 46 (2): 738-759. <https://doi.org/10.1016/j.ympev.2007.11.009>
- Bailly N, Gerovasileiou V, Arvanitidis C, Legakis A (2016) Introduction to the Greek Taxon Information System (GTIS) in LifeWatchGreece: the construction of the Preliminary Checklists of species of Greece. *Biodiversity Data Journal* LifeWatchGreece: Research infrastructure (ESFRI) for biodiversity data and data observatories: in press.
- Boudreau BP, Jørgensen BB (2001) *The benthic boundary layer: Transport processes and biogeochemistry*. Oxford University Press, 440 pp.
- Brunel P, Besner M, Messier D, Poirier L, Granger D, Weinstein M (1978) Le traîneau suprabenthique MACER-GIROQ : appareil amélioré pour l'échantillonnage quantitatif étagé de la petite faune nageuse au voisinage du fond. *The MACER-GIROQ Suprabenthic Sled: An Improved Device for Quantitative Two-level Sampling of the Small Swimming Fauna near the Bottom*. *Internationale Revue der gesamten Hydrobiologie und Hydrographie* 63 (6): 815-829. <https://doi.org/10.1002/iroh.19780630612>
- Chevaldonné P, Rastorgueff P, Arslan D, Lejeusne C (2015) Molecular and distribution data on the poorly known, elusive, cave mysid *Harmelinella mariannae* (Crustacea: Mysida). *Marine Ecology* 36 (3): 305-317. <https://doi.org/10.1111/maec.12139>
- Chevaldonné P, Sket B, Marschal C, Lejeusne C, Calado R (2008) Improvements to the "sket bottle": a simple manual device for sampling small crustaceans from marine caves and other cryptic habitats. *Journal of Crustacean Biology* 28 (1): 185-188. <https://doi.org/10.1651/07-2923R.1>
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Lasram FBR, Aguzzi J, Ballesteros E, Bianchi CN, Corbera J, Dailianis T, Danovaro R, Estrada M, Froglià C, Galil B, Gasol J, Gertwagen R, Gil J, Guilhaumon F, Kesner-Reyes K, Kitsos M, Koukouras A, Lampadariou N, Laxamana E, López-Fé de la Cuadra C, Lotze H, Martin D, Mouillot D, Oro D, Raicevich S, Rius-Barile J, Saiz-Salinas JI, Vicente CS, Somot S, Templado J, Turon X, Vafidis D, Villanueva R, Voultsiadou E (2010) The Biodiversity of the Mediterranean Sea: Estimates, Patterns, and Threats. *PLoS ONE* 5 (8): e11842. <https://doi.org/10.1371/journal.pone.0011842>
- Eleftheriou A (2013) *Methods for the study of marine benthos*. 4th Edition. John Wiley & Sons Ltd, Oxford, 496 pp. URL: <https://doi.org/10.1002/9781118542392> [ISBN 978-0-470-67086-6] <https://doi.org/10.1002/9781118542392>

- Hatzakis A (1974) Mysidacés trouvés en mer Egée (partie sud) de 1966 à 1971. Rapports et Procès-verbaux des Réunions de la Commission internationale pour l'Exploration scientifique de la Mer Méditerranée 22: 81.
- Hatzakis A (1977) Contribution à l'étude des Gastrosaccinae (Crustacea, Mysidacea) de la Méditerranée. Description de *Haplostylus bacescui* n.sp. et révision de la nomenclature des *Haplostylus* et *Gastrosaccus* méditerranéens. Biologia Gallo-Hellenica 6: 271-286.
- Hatzakis A (1978) Les Mysidacés, in: Ecologie marine de la région de l' Eubée du Nord. III Evolution du zooplankton dans sept stations. Biologia Gallo-Hellenica 7: 142-144.
- Hatzakis A (1982) Mysidacea of Greek Seas (morphotaxonomy, eco-geography, bionomics). University of Athens, Greece, 208 pp. [In Greek].
- Koukouras A (2010) Check-list of marine species from Greece. Aristotle University of Thessaloniki, Greece. Assembled in the framework of the EU FP7 PESI project. In WoRMS. <http://www.marinespecies.org/aphia.php?p=sourcedetails&id=142068>
- Koulouri P, Dounas C, Eleftheriou A (2013) Hyperbenthic community structure over oligotrophic continental shelves and upper slopes: Crete (South Aegean Sea, NE Mediterranean). Estuarine, Coastal and Shelf Science 117: 188-198. <https://doi.org/10.1016/j.ecss.2012.11.015>
- Ledoyer M (1968) Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonome. (Région de Marseille principalement) IV. Synthèse de l'étude écologique. Recueil des Travaux de la Station marine d' Endoume 44 (60): 125-295.
- Madurell T, Cartes J (2003) The suprabenthic peracarid fauna collected at bathyal depths in the Ionian Sea (eastern Mediterranean). Crustaceana 76 (5): 611-624. <https://doi.org/10.1163/156854003322316245>
- Mees J, Jones MB (1997) The Hyperbenthos. Oceanography and Marine Biology: An Annual Review 35: 221-255.
- Mees J, Meland K (2012) World List of Lophogastrida, Stygiomysida and Mysida. <http://www.marinespecies.org/mysidacea/index.php>
- Meland K, Willassen E (2007) The disunity of "Mysidacea" (Crustacea). Molecular Phylogenetics and Evolution 44 (3): 1083-1104. <https://doi.org/10.1016/j.ympev.2007.02.009>
- Meland K, Mees J, Porter M, Wittmann K (2015) Taxonomic Review of the Orders Mysida and Stygiomysida (Crustacea, Peracarida). PLOS ONE 10 (4): e0124656. <https://doi.org/10.1371/journal.pone.0124656>
- Rastorgueff P, Chevaldonné P, Arslan D, Verna C, Lejeusne C (2014) Cryptic habitats and cryptic diversity: unexpected patterns of connectivity and phylogeographical breaks in a Mediterranean endemic marine cave mysid. Molecular Ecology 23 (11): 2825-2843. <https://doi.org/10.1111/mec.12776>
- San Vicente C, Cartes J (2011) *Dactylamblyops corberai* n. sp., a new mysid (Crustacea: Mysida) from the deep Mediterranean Sea. Scientia Marina 75 (3): 455-464. <https://doi.org/10.3989/scimar.2011.75n3455>
- Wittmann K, Ariani A (2011) An Adjusted Concept for a Problematic Taxon, *Paramysis festae* Colosi, 1921, with Notes on Morphology, Biomineralogy, and Biogeography of the Genus *Paramysis* Czerniavsky, 1882 (Mysida, Mysidae). Crustaceana 84 (7): 849-868. <https://doi.org/10.1163/001121611x574272>
- Wittmann K, Ariani A (2012) *Diamysis cymodoceae* sp. nov. from the Mediterranean, Marmora, and Black Sea basins, with notes on geographical distribution and ecology of

the genus (Mysida, Mysidae). *Crustaceana* 85 (3): 301-332. <https://doi.org/10.1163/156854012x623719>

- Wittmann KJ, Ariani AP (1998) *Diamysis bacescui* n. sp., a new benthopelagic mysid (Crustacea: Peracarida) from Mediterranean seagrass meadows: description and comments on statolith composition. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" XL*: 35-49.
- Wittmann KJ, Wirtz P (1998) A first inventory of the mysid fauna (Crustacea: Mysidacea) in coastal waters of the Madeira and Canary Archipelagos. *Boletim do Museu de História Natural do Funchal* 5: 511-533.

Supplementary material

Suppl. material 1: Checklist of Mysida and Lophogastrida (Arthropoda: Malacostraca) of Greece

Authors: Panayota Koulouri, Vasilis Gerovasileiou, Nicolas Bailly

Data type: Taxonomic checklist

Brief description: Checklist of Mysida and Lophogastrida known to occur in Greek waters.

Filename: GTIS_Mysida_Lophogastrida_Greece_Checklist.xls - [Download file](#) (304.00 kb)