



# Management in northeastern Brazil : faunal biodiversity

M.N. Paranaguá<sup>1</sup>, S. Neumann-Leitão<sup>2</sup>, R.L.S. Melo<sup>1</sup>,  
P.A. Coelho<sup>2</sup>, A.L. Vasconcelos Filho<sup>2</sup>, A.M.E. Oliveira<sup>2</sup>

*<sup>1</sup>Universidade Federal Rural de Pernambuco - Depto. Zoologia. R. Dom Manuel de Medeiros s/n - Recife - PE- Brazil 52171-900*

*Email: prpqpg@gir.npde.ufpe.br*

*<sup>2</sup>Universidade Federal de Pernambuco - Depto. Oceanografia. Campus Universitário, CDU - Recife - PE - Brazil 50670-901*

*Email: oceano@npd.ufpe.br and sigrid@zaz.com.br*

## Abstract

A total of 618 taxa were identified to Itamaracá ecosystem. Of these, 113 were benthic mollusks, 177 benthic crustaceans, 140 fish, 71 birds, 116 zooplankton and 1 species of sea mammal. Taxonomic biodiversity was high considering a mangrove estuarine area. The coexistence of these taxa confirm the importance of the estuarine complex of Itamaracá as a feeding, breeding, maturation and protection area, which in turn confers special interest in preserving the existent resources. In spite of numerous indiscriminate activities carried out in this area, the pollution bioindicators are generally restricted to the estuaries of the rivers, meaning that the area has high resilience. There is a strong marine influence which allows for the preservation of the biodiversity.

## 1 Introduction

The estuarine complex of Itamaracá is located at the north coast of the state of Pernambuco, northeastern Brazil (7°32' - 8°56'S and 34°49' - 35°11'W), and it consists of the Santa Cruz Channel, a U-shaped channel of 20 km length with two connections to the Atlantic Ocean and 5 small rivers draining into the channel (Figure 1). The system sustains 36 km<sup>2</sup> of mangrove forests, dominated



by *Rhizophora mangle*, *Laguncularia racemosa* and *Avicennia* spp. Mangrove forests are usually separated from terrestrial vegetation by a disrupted belt of saltflats, typically composed by *Sesuvium* sp. and gramineans. Towards the shelf, the estuarine system is delimited by arenite reefs. Between the reefs and Itamaracá Island, a shallow (0.5 to 2 m depth) coastal basin is formed. In these areas, patches of seagrass *Halodule wrightii* create a distinct environment (Medeiros & Kjerfve<sup>5</sup>; Schwamborn<sup>8</sup>).

Due to its ecological characteristics, there is a strong trophic interaction between the communities caused by the shallow water, the intense dynamics of the tides and the mangrove contribution of a high load of particulate and dissolved organic matter. In consequence, the social-economic value of this estuarine mangrove complex is very high, as expressed by the intense fishing carried out in this area, up to 800 tons.yr<sup>-1</sup>.

In this paper, the diversity of the zooplankton, mollusks, crustaceans and fish is used as tools to monitor the “health” of Itamaracá, to indicate the effects of various anthropogenic and other intrusions to access management procedures.




Figure 1. Localization of the studied area (after Schwamborn, 1997)

## 2 Results and Discussion

### 2.1 Zooplankton

The Itamaracá ecosystem serves as a nursery for several larvae of mollusks, crustaceans and fish (Paranaguá & Eskinazi<sup>7</sup>; Schwambom<sup>8</sup>). Zooplankton biodiversity in this ecosystem is high with a total of 116 taxa, considering the smallest unit possible to be identified for each group (Table 1). Copepoda was dominant (28 species) followed by Decapoda (19 species) and Tintinnida (15 species). The holoplankton predominated (65 taxa) in relation to meroplankton (40 taxa) and tycooplankton (11 taxa).

The zooplankton composition presented a low variability throughout the entire area. Differences were caused by small scale patchiness due transversal convection and temporal release of meroplanktonic larvae. The general pattern was an increase in diversity towards the coastal area, where abiotic conditions are more stable.

Protozoa was represented by Testacea, Radiolaria and Foraminiferida, this last one mostly composed of benthic species. Tintinnina presented 15 species and they are frequently found in coastal ecosystems of northeastern Brazil. Cnidaria presented 3 species, of which *Lyrioie tetraphyla* is frequent in coastal regions; *Ostrumovia inkermanica* and *Blackfordia virginica* occur frequently in estuaries of Itamaracá. Ctenophora was abundant all over the area, mainly the *Beroe* genus.

Platyhelminthes was present with a larval stage of a benthic specie spread out along the littoral area of the channel. Aschelminthes was composed by Nematoda and Rotifera. Nematoda had a high number of benthic species, mostly in areas of high organic matter decomposition. Species identification of this group is urgently needed. Ten Rotifera species were identified most of them restricted to the Botafogo River estuary, and according to Neumann-Leitão *et al.*<sup>7</sup> forming a group of organic pollution indicators.

Mollusca was represented by the larval stages of Gastropoda and Bivalvia (3 species each). Polychaeta was the only class of Annelida, mostly represented by larval stages of Spionidae and Nereiididae families.

Crustacea was dominant outranking Copepoda (28 species) and Decapoda larvae (19 taxa). Among the Copepoda *Paracalanus crassirostris*, *Acartia lilljeborgi*, *Oithona hebes*, and *Euterpina acutifrons* were abundant, all forming a typical group of the Brazilian estuaries.

Decapoda had high larval abundance of the commercial species *Penaeus* spp, *Callinectes* spp, *Aratus pisoni*, and *Ucides cordatus*. Insecta larvae was significative around the mangrove areas. Bryozoa, Phoronidea and Echinodermata were represented by a few larvae mostly found in the coastal area associated to the seagrass meadows.

Chaetognatha was present with *Sagitta tennuis*, more abundant along the channel. Chordata had 2 Larvacea species, and 1 Ascidiacea and 1



Cephalocordata species in the larval stage, all dominating in the channel outlets. Fish larvae and eggs were abundant all over the area.

Table 1 – Zooplankton from the Itamaracá ecosystem, Pernambuco (Brazil)

PROTOZOA	<i>Keratella americana</i>	<i>Euterpina acutifrons</i>
Testacea	<i>Lecane curvicornis</i>	<i>Farranula gracilis</i>
<i>Arcella vulgaris</i>	<i>Lecane closteroerca</i>	<i>Triguiopulus</i> sp
<i>Arcella dentata</i>	<i>Cephalodella</i> sp	<i>Metis</i> sp
<i>Centropyxis acureata</i>	<i>Trichocerca elongata</i>	Cirripedia
Radiolaria (Spumellaria)	Nematoda	<i>Balanus</i> sp (nauplius and cypris)
Foraminiferida	MOLLUSCA	Cumacea
<i>Textularia</i> sp	Gastropoda	Isopoda (larvae)
<i>Quinqueloculina</i> sp	<i>Bitium varium</i> (veliger)	Amphipoda
<i>Triloculina</i> sp	<i>Smaragdia viridis</i> (veliger)	Decapoda
<i>Discorbis mira</i>	<i>Tricolia affinis</i> (veliger)	<i>Acetes americanus</i> (larvae)
<i>Planispirulina denticulata</i>	Bivalvia	<i>Atya</i> sp (larvae)
<i>Globigerinoides ruber</i>	<i>Anomalocardia brasiliiana</i> (veliger)	<i>Penaeus</i> sp (mysis)
<i>Globorotalia menardii</i>	<i>Mytella falcata</i> (veliger)	<i>Lucifer faxoni</i> (larvae and adult)
<i>Tretomphalus bulloides</i>	<i>Crassostrea rhyzophorae</i> (veliger)	<i>Periclimenes longicaudatus</i> (larvae)
<i>Amphistegina</i> sp	ANNELIDA – Polychaeta	<i>Lysmata</i> spp (larvae)
Tintinnina	Spionidae (larvae)	<i>Callianassa</i> spp (larvae)
<i>Leptotintinnus nordqvisti</i>	<i>Sabellaria</i> sp (larvae)	<i>Paguristes</i> spp (larvae)
<i>Tintinnopsis directa</i>	<i>Nereis</i> sp (larvae)	<i>Petrolisthes</i> spp (1 larvae)
<i>Tintinnopsis brasiliensis</i>	CRUSTACEA	<i>Polyonyx</i> spp (larvae)
<i>Tintinnopsis compressa</i>	Ostracoda	<i>Porcellana</i> spp (larvae)
<i>Tintinnopsis mortensenii</i>	Copepoda	<i>Callinectes</i> spp (larvae)
<i>Tintinnopsis tocartinensis</i>	<i>Clausocalanus furcatus</i>	<i>Pinothores</i> spp (larvae)
<i>Tintinnopsis</i> sp	<i>Eucalamus pileatus</i>	<i>Sesarma rectum</i> (zoea)
<i>Codonellopsis morchella</i>	<i>Paracalanus quasimodo</i>	<i>Uca maracoani</i> (zoea)
<i>Coxliella annulata</i>	<i>Paracalanus crassirostris</i>	<i>Aratus pisoni</i> (zoea)
<i>Favella ehrenbergii</i>	<i>Centropages velificatus</i>	<i>Uca leptodactyla</i> (zoea)
<i>Rhabdonella spiralis</i>	<i>Temora turbinata</i>	<i>Uca burgersi</i> (zoea)
<i>Amphorellopsis acuta</i>	<i>Temora stylifera</i>	<i>Ucides cordatus</i> (zoea)
<i>Eutintinnus tenuis</i>	<i>Pseudodiaptomus acutus</i>	Syncarida ( <i>Batynella</i> sp)
<i>Epiplocypris</i> sp	<i>Pseudodiaptomus richardi</i>	INSECTA (larvae)
<i>Undella</i> sp	<i>Pseudodiaptomus marshi</i>	BRYOZOA (cyphonauta)
CNIDARIA	<i>Calanopia americana</i>	PHORONIDEA (actinotroca)
<i>Lyriope tetraphyla</i>	<i>Labidocera fluviatilis</i>	ECHINODERMATA
<i>Blackfordia virginica</i>	<i>Lucicutia flavicornis</i>	Echinoidea (pluteus)
<i>Ostrumovia inkermanica</i>	<i>Acartia lilljeborgi</i>	CHAETOGNATHA
CTENOPHORA – <i>Beroe</i> sp	<i>Oithona nana</i>	<i>Sagitta tenuis</i>
PLATYHELMINTHES	<i>Oithona hebes</i>	CHORDATA
<i>Convoluta</i> sp	<i>Oithona oswaldocruzi</i>	Larvacea
ASCHELMINTHES	<i>Hemicyclops talassius</i>	<i>Oikopleura longicauda</i>
Rotifera	<i>Oncaea</i> sp	<i>Oikopleura dioica</i>
<i>Rotaria rotatoria</i>	<i>Corycaeus giesbrechti</i>	Ascidiacea
<i>Rotaria</i> sp	<i>Corycaeus speciosus</i>	<i>Ciona</i> sp (larvae)
<i>Brachionus bidentata</i> f.	<i>Saphirina</i> sp	Cephalochordata
<i>inermis</i>		<i>Branchiostomus</i> (larvae)
<i>Brachionus plicatilis</i>	<i>Clyemnestra scutelata</i>	Osteichthyes (egg and larvae)
<i>Keratella tropica tropica</i>	<i>Microsetella rosea</i>	



## 2.2 Mollusca

The mollusks were represented by the Gastropoda (78 species), Bivalvia (32 species), Polyplacophora (1 specie) and Scaphopoda (2 species) (Table 2). The class Gastropoda Mesogastropoda presented the highest taxonomic diversity at Itamaracá ecosystem. *Neritina virginica*, *Nassarius vibex*, *Cerithium atratum* and *Bittium varium* were found associated to seagrass meadows. The *Littorina angulifera*, *L. lineolata* and *L. flava* occurred under mangrove leaves and stems; the Collumbellidae (9 species) and Olividae (6 species) under small rocks; and the Pyramidellidae (10 species) and Caecidae (5 species), in the muddy sediment under mangrove roots. Bivalvia had lower taxonomic diversity. Among the economical importance bivalves it can be cited: *Anomalocardia brasiliiana*, *Crassostrea rhizophorae*, *Mytella falcata*, *Tagelus plebeius*, *Iphigenia brasiliiana*, *Protothaca pectorina*, *Lucina pectinata* e *Tivela mactroides*. Species of the bivalve family Mytilidae was strongly associated with eutrophic environments at Itamaracá ecosystem.

## 2.3 Crustacea

The diversity of crustaceans was high and apart planktonic species a total of 177 species were identified (at least to genus level). The microcrustaceans or parasitic crustaceans like Cladocera, Ostracoda, Rhizocephala and Cumacea were just referred as a group. Taxonomic identification is presented to Copepoda, Thoracica, Stomatopoda, Amphipoda, Isopoda and Decapoda.

Non-planktonic Copepoda studies are still recent. For this reason, only the parasitic species or the meiobenthos are cited: *Acanthocolax sp.*, *Canuella sp.*, *Caligus minimus*, *C. elongatus*, *Cyclopina sp.*, *Ectinosoma sp.*, *Ergasilus atagonensis*, *E. caraguatatubensis*, *E. lizae*, *Heterolaeophonte sp.*, *Leptocaris sp.*, *Lernanthropus gisteri*, *Longipedia sp.*, *Robertsonia sp.*, *Stenhelia cf. Normani* and *Stenhelia sp.*

The Thoracica were attached to hard substracts or living in association with other animals. To Itamaracá ecosystem 11 species are known. The Archaeobalanidae family constituted by *Chirone (Striatobalanus) amaryllis* is usually covered by sponges or other cirripeds. Balanidae was represented by *Balanus amphitrite*, *B. improvisus*, *B. reticulatus*, *B. trigonus*, *B. venustus* and *Fistulobalanus citerosum*. Chthamalidae by *Chthamalus bisinatus*, *C. proteus* and *Euraphia rhizophorae*. All these species were found under mangroves trees or rocks. The Coronulidae was represented by *Chelonibia patual*, found under the shell of *Callinectes exasperatus*.

Stomatopoda, essentially marine, presented 2 species, one belonging to Squillidae (*Chloridopsis dubia*) and other to Pseudosquillidae (*Pseudosquilla ciliata*). Amphipoda was present with 11 estuarine species: *Paragrubia sp.*, *Cymadusa sp.*, *Sunamphitoe sp.*, *Hyale sp.*, *Hyale nilssoni*, *Talitrus saltator*, *Orchestria gamarella*, *O. montagui*, *O. platensi*, *Orchestia sp.* and *Allorchestes sp.*

Isopoda was presented with 13 species: *Rocinela signata*, *Aegathoa linguifrons*, *Anilocra laticauda*, *Cymothoa excisa*, *Livoneca redmanni*, *Cassinidea fluminensis*, *Pseudosphaeroma jakobii*, *P. mourei*, *Sphaeroma terebrans*, *Bopyrella slphe*, *Leidyia distorta*, *Probopyrus bithynis* and *Ligia offersii*. *Rocinela*, *Aegathoa*, *Anilocra*, *Cymothoa* and *Livoneca* are parasites of different species of fish and *Bopyrella* and *Probopyrus* are found in the branchial chambers of *Alpheus* and *Macrobachium*. *Leidyia* is also found in the respiratory chambers of the crab *Ucides cordatus*. The other species, except for *Ligia*, were found preying on mangroves trees.

Table 2 – Mollusca from the Itamaracá ecosystem, Pernambuco (Brazil).

GASTROPODA	<i>Costoanachis catenata</i> *	<i>Bulla striata</i>
<b><i>Smaragdia viridis</i></b>	<i>Parvanachis isabellei</i> *	BIVALVIA
<i>Tricolia bella</i>	<i>Olivella petiolita</i> **	<i>Diplodonta punctata</i>
<i>Tricolia affinis</i>	<i>Olivella minuta</i> **	<i>Ostrea equestris</i>
<i>Neritina virginea</i>	<i>Olivella mutica</i> **	<i>Crassostrea rhizophorae</i>
<i>Parviturbo rehderi</i>	<i>Olivella floralia</i> **	<i>Anadara notabilis</i>
<i>Parviturbooides interruptus</i>	<i>Olivella watermani</i> **	<i>Anadara ovalis</i>
<i>Episcynia inornata</i>	<i>Olivella nivea</i> **	<i>Barbatia candida</i>
<i>Rissoina bryerea</i>	<i>Volvarina avena</i>	<i>Arca umbonata</i>
<i>Rissoina catesbyana</i>	<i>Mangelia biconica</i>	<i>Arcopsis adamsi</i>
<i>Zebina browniana</i>	<i>Mangelia stellata</i>	<i>Tivela mactroides</i>
<i>Cerithium atratum</i>	<i>Kurtziella dorvillae</i>	<i>Anomalocardia brasiliiana</i>
<i>Bittium varium</i>	<i>Tenaturris decora</i>	<i>Protothaca pectorina</i>
<i>Cerithiopsis greenii</i>	<i>Buchema interpleura</i>	<i>Tranzenella cubaniana</i>
<i>Seila adamsi</i>	<i>Brachycytara galae</i>	<i>Chione cancellata</i>
<i>Alaba incerta</i>	<i>Tenaturris fulgens</i>	<i>Corbula caribaea</i>
<i>Caecum floridanum</i> ****	<i>Epitonium denticulatum</i>	<i>Iphigenia brasiliiana</i>
<i>Caecum ryssotitum</i> ****	<i>Epitonium candeanum</i>	<i>Tagelus plebeius</i>
<i>Caecum pulchellum</i> ****	<i>Epitonium novangliae</i>	<i>Tagelus divisus</i>
<i>Caecum imbricatum</i> ****	<i>Epitonium nautlae</i>	<i>Sanguinolaria cruenta</i>
<i>Caecum striatum</i> ****	<i>Thais haemastoma</i>	<i>Tellina lineata</i>
<i>Littorina angulifera</i>	<i>Nassarius vibex</i>	<i>Tellina punicea</i>
<i>Littorina lineolata</i>	<i>Turbonilla haycocki</i> ***	<i>Macoma constricta</i>
<i>Littorina flava</i>	<i>Turbonilla fasciata</i> ***	<i>Strigilla mirabilis</i>
<i>Natica menkeana</i>	<i>Turbonilla interrupta</i> ***	<i>Mytella falcata</i>
<i>Natica canrena</i>	<i>Turbonilla (Pyrgiscus) elegans</i> ***	<i>Modiolus americanus</i>
<i>Balcis intermedia</i>	<i>Turbonilla multicostata</i> ***	<i>Brachidontes solicianus</i>
<i>Triphora nigrocincta</i>	<i>Turbonilla (Turbonilla) aff. Turris</i> ***	<i>Brachidontes exustus</i>
<i>Triphora ornata</i>	<i>Odostomia jadisi</i> ***	<i>Divalinga quadrisulcata</i>
<i>Modulus modulus</i>	<i>Odostomia canaliculata</i> ***	<i>Lucina pectinata</i>
<i>Trivia pediculus</i>	<i>Odostomia laevigata</i> ***	<i>Trachycardium muricatum</i>
<i>Truncatella pulchella</i>	<i>Peristichia agria</i> ***	<i>Laevicardium laevigatum</i>
<i>Parvanachis obesa</i> *	<i>Iselica anomala</i>	<i>Ervilia nitens</i>
<i>Mitrella lunata</i> *	<i>Atys sandersoni</i>	<i>Ervilia subcancellata</i>
<i>Aesopus metcalfei</i> *	<i>Atys caribaea</i>	POLYPLACOPHORA
<i>Collumbella mercatoria</i> *	<i>Haminoea antillarum</i>	<b><i>Ischnochiton striolatus</i></b>
<i>Mitrella ocellata</i> *	<i>Acteocina bullata</i>	SCAPHOPODA
<i>Anachis sertulariarum</i> *	<i>Bursatella leachii</i>	<b><i>Dentalium americanum</i></b>
<i>Anachis avara</i> *	<i>Aplysia dactylomela</i>	<b><i>Graptacme calamus</i></b>

Collumbellidae\* Olividae\*\* Pyramidellidae \*\*\* Caecidae\*\*\*\*

Decapoda is the best known group in the area and sum up a high percentage of the fisheries at Itamaracá. Some species are parasites or live on a variety of invertebrates as polychaetes, equinoderms, mollusks and other crustaceans. Although high in diversity (124 species), this corresponds to 37.6% of the marine decapod cited by Coelho & Ramos-Porto<sup>2</sup> for the estuarine and marine area (including the continental shelf) of Pernambuco and neighboring states. Stenopodidea and Astacidea were not registered to Itamaracá.

Penaeidea includes 8 species, 5 of the Penaeidae family: *Penaeus notialis*, *P. schmitti*, *P. suntuilis*, *P. brasiliensis* and *Xiphopenaeus kroyeri*, this last rarely found in the environment, indicate marine influence. Sicyoniidae presented: *Sicyonia typica*, *S. laevigata* and *S. dorsalis*.

Caridea is represented by 6 families (35 species) of freshwater and typically marine species. In the Atyidae a freshwater species (*Potimirim potimirim*) was found. The Pasiphaeidae were represented by *Leptochela serratorbita*. Two other families, with few species, are the Hippolytidae (*Latreutes parvulus* and *Merguia rhizophorae*) and Processidae (*Ambidexter symmetricus*). Palaemonidae presented 9 species (*Leander paulensis*, *L. tenuicornis*, *Macrobrachium acanthurus*, *M. heterochirus*, *Palaemon northropi*, *P. pandaliformis*, *Periclimenes americanus*, *P. longicaudatus* and *Typton distinctus*) and Alpheidae 13 species (*Alpheus armillatus*, *A. bouvieri*, *A. chacei*, *A. estuariensis*, *A. floridanus*, *A. normanni*, *Automate evermanni*, *Leptalpheus petronii*, *Salmoneus ortmanni*, *Synalpheus apiocerus*, *S. brevicarpus*, *S. fritzmulleri* and *S. minus*). Palinuridea presented only *Panulirus argus* (Palinuridae).

Thalassinidea needs to be better studied, and probably there are a larger number of species, of which 3 Callisanassidae are known at present (*Lepidophthalmus siriboiajamaicense*, *Neocallichirus rathbunae* and *Sergio guassutinga*), one Laomediidae (*Ctenaxianassa australis*) and 2 Upogebiidae (*Upogebia noronhensis* and *U. omissa*), totaling 7 species.

Anomura are predominantly marine and is represented by 5 families with 12 species. Paguridae is represented by *Pagurus criniticornis*, Diogenidae by *Clibanarius antillensis*, *C. scolopettarius*, *C. vittatus*, *Dardanus venosus* and *Petrochirus diogenes*, Porcellanidae by *Minyocerus angustus*, *Pachycheles greeleyi*, *Petrolisthes armatus* and *P. galathinus*, Hippidae by *Emerita portoricensis* and Albuneidae by *Lepidopa richmondi*.

Brachyura with 61 species accounts for nearly one third of the known crustaceans and half of the Decapoda species. All habitats are inhabited at least by one specie of this group. Most marine families were represented by 1 or 2 species, such as Calappidae (*Calappa ocellata*), Leucosiidae (*Lithadia brasiliensis*), Parthenopidae (*Hepatus pudibundus* and *Mesorhoea sexspinosa*) and Eriphiidae (*Eriphia gonagra*). Exception is the Maiidae represented by 7 species (*Acanthonyx dissimulatus*, *Epialtus bituberculatus* *Hemus cristulipes*, *Microphrys bicornutus*, *M. interruptus*, *Mithrax hispidus* and *Notolopas brasiliensis*). Pilumnidae, best known in seawater environment, has 3 species in



the studied area (*Pilumnus caribaeus*, *P. dasypodus* and *P. reticulatus*). Xanthidae with 17 species is the largest group in number of species (*Cyrtoplax spinidentata*, *Eurypanopeus dissimilis*, *Euryplax nitida*, *Eurytium limosum*, *Hexapanopeus angustifrons*, *H. caribbaeus*, *H. hemphilli*, *H. paulensis*, *H. quinquedentatus*, *H. schmitti*, *Menippe nodifrons*, *Panopeus americanus*, *P. bermudensis*, *P. harttii*, *P. lacustris*, *P. occidentalis* and *Panoplax* sp.). In holes dug in finer and more fluid sediment by *C. spinidentata* a diverse fauna was found, including 2 species of the shrimp *Alpheus* and *Salmonella*, one stomatopod (*Cloridopsis*) and 3 fishes (*Gobionellus smaragdus*, *G. boleosoma* and *Erotelis smaragdus smaragdus*).

Portunidae was represented by *Arenaeus cribrarius* (very rare), *Callinectes bocourti*, *C. danae*, *C. exasperatus*, *C. larvatus* and *C. ornatus*. The Pinnotheridae with 6 species were associated to polichaetes, equinoderms, mollusks and other crustaceans (*Dissodactylus crinitichelis*, *Pinnixa aida*, *P. chaetoptera*, *P. leptodactyla*, *P. sayana* and *Zoops ostreum*).

The Ocypodidae with 10 species (*Ocypode quadrata*, *Uca burgersi*, *U. cumulanta*, *U. leptodactyla*, *U. maracoani*, *U. mordax*, *U. rapax*, *U. thayeri*, *U. vocator* and *Ucides cordatus*) and the Grapsidae, with 8 species (*Aratus pisonii*, *Armases angustipes*, *Cyclograpsus integer*, *Goniopsis cruentata*, *Pachygrapsus gracilis*, *P. transversus*, *Sesarma crassipes* and *S. rectum*) dominated the environment. Gecarcinidae family presented one specie, *Cardisoma guanhumi*.

## 2.4 Fishes

The ictiofauna of the Itamaracá ecosystem is composed of 140 species. Two species belong to the Chondrichthyes (*Narcine brasiliensis* and *Dasayatis guttata*). The low number of Chondrichthyes seems to be a characteristic of estuarine environments in northeastern Brazil. It is possible that the Chondrichthyes do not find adequate conditions in the local estuarine waters for their feeding and reproduction. All the other species belong to the Osteichthyes class. The most representative families were: Carangidae (9 species), Gerreidae and Scianenidae (8 species, each), Gobiidae (7), Engraulidae, Haemulidae and Lutjanidae (6 species, each), Ariidae and Paralichthyidae (5 species, each) and Eleotridae, Soleidae and Syngnathidae (4 species, each). All other families were represented by 3 or less species.

The fishes species were grouped into 3 categories: resident, saltwater dependent and saltwater visitor. The residents include 24 species (Table 3) and spend all their life cycle in the estuary, although they are sometimes found in coastal areas or in freshwater. The saltwater dependents include 43 species (Table 4) and enter the estuary for feeding or reproduction. The saltwater visitors include 73 species (Table 5) spend their life in the sea, occasionally or regularly venturing into estuarine waters.

The 138 Osteichthyes species are commonly found in tropical estuaries. Among the species found at Itamaracá ecosystem, *Poecilia vivipara* classified as



resident is the only specie of freshwater origin, although it is not a true freshwater species, because it reproduces in all salinities range from freshwater to seawater, and is usually referred as a complete eurihaline specie. Other residents, have their origin in the marine habitat. Some prefer freshwater to saltwater, such as Eleotridae and some Gobiidae. These families frequently move from the sea to freshwater. The rest of the species are commonly found in the coastal shallow waters (Eskinazi<sup>3</sup>; Vasconcelos Filho *et al.*<sup>9</sup>).

The saltwater dependent species, although smaller in number than the visitors are very important, because they characterize the estuarine ictiofauna, such as *Mugil* spp., *Centropomus* spp., *Diapterus* spp., *Eucinostomus* spp., *Eugerres brasilianus* and *Gerres cinereus*.

The visitors are usually young, and some have commercial value, such as Lutjanidae, Carangidae, Haemirhamphidae, Serranidae, Lobotidae, Haemulidae, Sparidae, Scianidae and Scombridae. These families show the importance of the Itamaracá area as a nursery ground. Some species inhabit rocky areas (*Gymnotorax* spp., *Pomocanthus paru*, *Abudefduf saxtilis*, *Chaetodon* spp., *Pomacentrus variabilis*, *Acanthurus* spp. and *Amanses (Cantherines pullus)*, a substract not common in estuarine environments.

Table 3 - Residents fishes at Itamaraca ecosystem, Pernambuco, Brazil

<i>Achirus achirus</i>	<i>Evorthodus lyricus</i>	<i>Shpoeroides splengeri</i>
<i>Achirus declives</i>	<i>Gobionellus boleosoma</i>	<i>Shpoeroides testudineus</i>
<i>Achirus lineatus</i>	<i>Gobionellus oceanicus</i>	<i>Symphurus plagusia</i>
<i>Bathygobius soporator</i>	<i>Gobionellus smaragdus</i>	<i>Arius herzbergii</i>
<i>Cathorops spixii</i>	<i>Gobionellus stigmaticus</i>	<i>Arius parkeri</i>
<i>Dormitator maculatus</i>	<i>Gobionellus stomatu</i>	<i>Arius proops</i>
<i>Erotelis smaragdus smaragdus</i>	<i>Guavina guavina</i>	<i>Trinectes maculatus</i>
<i>Erotelis civitatum</i>	<i>Poecilia vivipara</i>	<i>Xenomelaniris brasiliensis</i>

Table 4 - Marine dependent fishes at Itamaraca ecosystem, Pernambuco, Brazil

<i>Anchoa filifera</i>	<i>Eucinostomus gula</i>	<i>Oligoplites palometa</i>
<i>Anchoa januaria</i>	<i>Eucinostomus havana</i>	<i>Oligoplites saliens</i>
<i>Anchoa tricolor</i>	<i>Eucinostomus melanopterus</i>	<i>Oligoplites saurus</i>
<i>Anchovia clupeoides</i>	<i>Eucinostomus lefroy</i>	<i>Ophioscion microps</i>
<i>Bairdiella ronchus</i>	<i>Eugerres brasilianus</i>	<i>Paralichthys brasiliensis</i>
<i>Bothus ocellatus</i>	<i>Gerres cinereus</i>	<i>Paralichthys orbignyana</i>
<i>Centropomus parallelus</i>	<i>Hyporhamphus unifasciatus</i>	<i>Polydactylus virginicus</i>
<i>Centropomus undecimalis</i>	<i>Lile piquitinga</i>	<i>Sphyræna barracuda</i>
<i>Chaetodipterus faber</i>	<i>Lycengraulis grossidens</i>	<i>Stellifer brasiliensis</i>
<i>Citharichthys crossotus</i>	<i>Menticirrhus martinicensis</i>	<i>Stellifer rastrifer</i>
<i>Citharichthys spilopterus</i>	<i>Micropogonias furnieri</i>	<i>Strongylura marina</i>
<i>Cynoscion acoupa</i>	<i>Mugil curema</i>	<i>Strongylura timucu</i>
<i>Cynoscion leiarchus</i>	<i>Mugil liza</i>	<i>Syngium micrurum</i>
<i>Diapterus olisthostomus</i>	<i>Mugil trichodon</i>	<i>Trichiurus lepturus</i>
<i>Diapterus rhombeus</i>		

Table 5 – Marine visitors fishes at Itamaraca ecosystem, Pernambuco, Brazil

<i>Abudefduf saxatilis</i>	<i>Epinephelus itajara</i>	<i>Ogcocephalus vespertilio</i>
<i>Acanthurus bahianus</i>	<i>Fistularia tabacaria</i>	<i>Opisthonema oglinum</i>
<i>Acanthurus chirurgus</i>	<i>Gentiatremus luteus</i>	<i>Orthopristes ruber</i>
<i>Albula vulpes</i>	<i>Gymnothorax funebris</i>	<i>Phrynelox scaber</i>
<i>Ananases (Cantherines) pullus</i>	<i>Gymnothorax moringa</i>	<i>Pomacanthus paru</i>
<i>Amphichthys criptocentrus</i>	<i>Gymnothorax nigromarginatus</i>	<i>Pomacentrus variabilis</i>
<i>Anisotremus virginicus</i>	<i>Haemulon aurolineatum</i>	<i>Pomadasys corvinaeformis</i>
<i>Archosargus rhomboidalis</i>	<i>Haemulon parrai</i>	<i>Prionotus alipionis</i>
<i>Archosargus unimaculatus</i>	<i>Harengula clupeola</i>	<i>Prionotus punctatus</i>
<i>Bagre marinus</i>	<i>Hemirhamphus brasiliensis</i>	<i>Pseudupeneus maculatus</i>
<i>Canthigaster rostratus</i>	<i>Hippocampus hudsonius</i>	<i>Rypticus randalli</i>
<i>Caranx hippos</i>	<i>Hippocampus reidi</i>	<i>Scomberomorus brasiliensis</i>
<i>Caranx latus</i>	<i>Lactophrys trigonus</i>	<i>Scorpaena plumieri</i>
<i>Cetengraulis edentulus</i>	<i>Lactophrys triquetar</i>	<i>Selene setapinnis</i>
<i>Chaetodon aculeatus</i>	<i>Lobotes surinamensis</i>	<i>Selene vomer</i>
<i>Chaetodon striatus</i>	<i>Lutjanus analis</i>	<i>Sparisoma radians</i>
<i>Chilomycterus spinosus</i>	<i>Lutjanus apodus</i>	<i>Syngnathus duncheri</i>
<i>Chloroscombrus chrysurus</i>	<i>Lutjanus griseus</i>	<i>Syngnathus elucens</i>
<i>Colomesus psittacus</i>	<i>Lutjanus jocu</i>	<i>Synodus foetens</i>
<i>Cynopontius savana</i>	<i>Lutjanus synagris</i>	<i>Synodus poey</i>
<i>Dactylopterus volitans</i>	<i>Megalops atlanticus</i>	<i>Thalassophryne montevidensis</i>
<i>Dasyatis guttata</i>	<i>Myrichthys ocellatus</i>	<i>Thalassophryne nattereri</i>
<i>Diodon hystrix</i>	<i>Narcine brasiliensis</i>	<i>Tomicodon fasciatus fasciatus</i>
<i>Echeneis naucrates</i>	<i>Ocyurus crysurus</i>	<i>Trachinotus falcatus</i>
<i>Elops saurus</i>		

## 2.5 Bird and mammal

It was registered a total of 71 birds residents and migrants. The migrant birds use the Itamaracá ecosystem as a forage, plumage change and weight gain area before returning to the Arctic for reproduction, showing that this ecosystem has more than a local importance (Azevedo Junior<sup>1</sup>).

There is just one specie of marine mammal *Trichechus manatus* strongly associated with the seagrass meadows and it is now under a protection program to avoid its extinction.

## 3 Conclusions

The Itamaracá ecosystem is an important socio-economical area and the high species diversity is thought to be maintained by habitat heterogeneity. In spite of the exploitation of this ecosystem, the pollution bioindicators seem to be more restricted to the rivers estuaries, very polluted areas according to Macedo *et al*<sup>4</sup>. Many organisms are living in their limits and may be excluded due to further stress as the loss of natural habitat converted to other uses.

The magnitude of the problem of species decline and loss is so great and growing so rapidly that calls to save endangered species are starting to be replaced by calls to save endangered ecosystems. This sensible, if politically



difficult, approach requires a great deal of biological knowledge in order to be effective. Protecting Itamaracá ecosystem requires the development of ways to monitor ecosystem health; the most effective ways seem to resolve around monitoring assemblages of organisms within the ecosystem.

## 4 References

1. Azevedo Júnior, S. M. As aves do Canal de Santa Cruz, Pernambuco, Brasil. *Caderno Omega, Série Biologia*, 5, pp.35-50, 1998.
2. Coelho, P. A. & Ramos-Porto, M. Distribuição ecológica dos crustáceos decápodos marinhos no Nordeste do Brasil. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 23, pp. 113-127, 1994/95.
3. Eskinazi, A. M. Peixes do Canal de Santa Cruz, Pernambuco-Brasil. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 13, pp. 283-302, 1972.
4. Macêdo, S. J., Melo, H. N. S. & Costa, K. M. P. Estudo ecológico da região de Itamaracá, pernambuco, Brasil. XXIII. Condições hidrológicas do estuário do rio Botafogo. *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 17, pp. 81-122, 1982.
5. Medeiros, C. Q. & Kjerfve, B. Hydrology of a tropical estuarine system: Itamaracá, Brazil. *Estuaries and Coastal Shelf Science.*, 36, pp. 495-515, 1993.
6. Neumann-Leitão, S., Paranaguá, M. N. & Valentin, J. L. Ecology of planktonic rotifers of the estuarine lagunar complex at Suape, Pernambuco (Brazil). *Hydrobiologia*, 232, pp. 133-143, 1992.
7. Paranaguá, M. N. & Eskinazi-Leça, E. Ecology of a northern tropical estuary in Brazil and technological perspectives in fishculture, Chapter 28, *Fish community ecology in estuaries and coastal lagoons towards na ecosystem integration*, ed. A. Yañes-Arancibia., Mexico, pp.595-614, 1985.
8. Schwamborn, R. Influence of mangroves on community structure and nutrition of macrozooplankton in Northeast Brazil. *ZMT Bremen Contribution* 4, pp. 1-77, 1997.
9. Vasconcelos Filho, A . L.; Cavalcanti, E. F.; Souza, S. T. Composição e distribuição da fauna ictiológica no Canal de Santa Cruz (Parte Sul de Itamaracá). *Revista Nordestina de Zoologia*. 1(1), pp. 247-262, 1994.