

ECOLOGY AND  
BEHAVIOUR OF THE  
LADYBIRD BEETLES  
(COCCINELLIDAE)

# ECOLOGY AND BEHAVIOUR OF THE LADYBIRD BEETLES (COCCINELLIDAE)

*Edited by I. Hodek, H.F. van Emden and A. Honěk*

 WILEY-BLACKWELL

A John Wiley & Sons, Ltd., Publication

This edition first published 2012 © 2012 by Blackwell Publishing Ltd

Blackwell Publishing was acquired by John Wiley & Sons in February 2007. Blackwell's publishing program has been merged with Wiley's global Scientific, Technical and Medical business to form Wiley-Blackwell.

*Registered office:* John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

*Editorial offices:* 9600 Garsington Road, Oxford, OX4 2DQ, UK  
The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK  
111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at [www.wiley.com/wiley-blackwell](http://www.wiley.com/wiley-blackwell).

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

*Library of Congress Cataloging-in-Publication Data*

Ecology and behaviour of the ladybird beetles (Coccinellidae) / edited by I. Hodek, H.F. van Emden, and A. Honěk.  
p. cm.

Includes index.

ISBN 978-1-4051-8422-9 (cloth)

1. Ladybugs. I. Hodek, Ivo. II. Van Emden, Helmut Fritz. III. Honěk, A. (Alois)

QL596.C65E26 2012

595.76'9—dc23

2011045545

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Set in 9/11 pt PhotinaMT by Toppan Best-set Premedia Limited

To all those scientists  
who have now passed on  
and who laid the foundation  
of our present knowledge of Coccinellidae,  
particularly to Michael Majerus  
who intended to be one of authors  
of this volume.

# CONTENTS

Detailed contents, ix

Contributors, xvii

Preface, xviii

Introduction, xix

Taxonomic glossary, xx

## **1. PHYLOGENY AND CLASSIFICATION, 1**

*Oldřich Nedvěd and Ivo Kovář*

## **2. GENETIC STUDIES, 13**

*John J. Sloggett and Alois Honěk*

## **3. LIFE HISTORY AND DEVELOPMENT, 54**

*Oldřich Nedvěd and Alois Honěk*

## **4. DISTRIBUTION AND HABITATS, 110**

*Alois Honěk*

## **5. FOOD RELATIONSHIPS, 141**

*Ivo Hodek and Edward W. Evans*

## **6. DIAPAUSE/DORMANCY, 275**

*Ivo Hodek*

## **7. INTRAGUILD INTERACTIONS, 343**

*Éric Lucas*

## **8. NATURAL ENEMIES OF LADYBIRD BEETLES, 375**

*Piotr Ceryngier, Helen E. Roy and Remy L. Poland*

## **9. COCCINELLIDS AND SEMIOCHEMICALS, 444**

*Jan Pettersson*

## **10. QUANTIFYING THE IMPACT OF COCCINELLIDS ON THEIR PREY, 465**

*J. P. Michaud and James D. Harwood*

## **11. COCCINELLIDS IN BIOLOGICAL CONTROL, 488**

*J. P. Michaud*

## **12. RECENT PROGRESS AND POSSIBLE FUTURE TRENDS IN THE STUDY OF COCCINELLIDAE, 520**

*Helmut E. van Emden and Ivo Hodek*

Appendix: List of Genera in Tribes and Subfamilies, 526

*Oldřich Nedvěd and Ivo Kovář*

Subject index, 532

Colour plate pages fall between pp. 250 and pp. 251

# DETAILED CONTENTS

Contributors, xvii	1.4.9 Coccidulinae, 8
Preface, xviii	1.4.10 Scymninae, 9
Introduction, xix	1.5 Future Perspectives, 10
Taxonomic glossary, xx	References, 10

## 1. PHYLOGENY AND CLASSIFICATION, 1

*Ondřich Nedvěd and Ivo Kovář*

1.1 Position of the Family, 2	1.1.1 The Cerylonid complex, 2
	1.1.2 Sister families, 2
	1.1.3 Feeding habits, 2
	1.1.4 Monophyly of Coccinellidae, 2
1.2 Characteristics of the Family, 3	
1.3 Changes in the Classification of Subfamilies, 3	1.3.1 Morphologically based classifications, 3
	1.3.2 Split of Sticholotidinae, 3
	1.3.3 Monophyly of other subfamilies, 4
	1.3.3.1 Contribution of immature stages, 4
	1.3.4 Molecular analyses, 4
	1.3.4.1 Alternative molecular methods, 5
	1.3.5 Rejection of the monophyly of subfamilies, 5
1.4 Characteristics of the Subfamilies and Tribes, 5	
	1.4.1 Proposed classification, 5
	1.4.2 Microweiseinae, 5
	1.4.3 Sticholotidinae, 5
	1.4.4 Coccinellinae, 6
	1.4.5 Epilachninae, 7
	1.4.6 Exoplectrinae, 7
	1.4.7 Chilocorinae, 7
	1.4.8 Ortaliinae, 8

## 2. GENETIC STUDIES, 13

*John J. Sloggett and Alois Honěk*

2.1 Introduction, 14	
2.2 Genome Size, 14	
2.3 Chromosomes and Cytology, 14	2.3.1 Chromosome numbers and banding, 14
	2.3.2 Sex determination, 14
	2.3.3 Supernumerary (B) chromosomes, 15
	2.3.4 Cytogenetic changes, intraspecific cytogenetic variation and speciation, 16
2.4 Colour Pattern Variation, 18	
	2.4.1 The nature of colour patterns, 19
	2.4.2 Genetic determination of colour patterns, 20
	2.4.3 Geographic variation, 23
	2.4.4 Temporal variation, 25
	2.4.5 Significance and evolution, 25
2.5 The Inheritance of Other Traits, 27	
	2.5.1 Morphological characters: wing polymorphism, 27
	2.5.2 Life history characters: heritability, selection experiments and genetic trade-offs, 28
2.6 Molecular Genetic Studies, 29	
	2.6.1 Sequence evolution, 29
	2.6.1.1 Mitochondrial DNA and the inference of evolutionary history, 29
	2.6.1.2 The ITS1 region, 31

2.6.2	Molecular studies of coccinellid biology, 33	3.5.6.3	Female choice and melanism, 83
2.6.2.1	Species, population and strain identification, 33	3.5.6.4	Hybridization, 85
2.6.2.2	Phylogenetics, 37	3.5.7	Oviposition, 85
2.6.2.3	Population genetic and phylogeographic studies, 38	3.5.7.1	Oviposition substrate, 85
2.6.2.4	Reproductive success, paternity and sperm competition, 42	3.5.7.2	Oviposition rhythmicity, 86
2.7	Conclusions, 43	3.5.7.3	Oviposition rate, 86
	Acknowledgements and Dedication, 44	3.5.7.4	Oviposition period, 86
	References, 44	3.5.8	Fecundity, 88
		3.5.9	Longevity, 90
		3.5.9.1	Voltinism, 90
		3.5.9.2	Effect of temperature, photoperiod and humidity, 90
		3.5.9.3	Effect of food, 91
		3.5.9.4	Effect of sexual activity, 91
2.7		3.6	Temperature and Development, 91
		3.6.1	Thermal constants, 92
		3.6.2	Relationship between LDT and SET, 92
		3.6.3	Thermal window and development rate isomorphy, 95
		3.6.4	Other events affected by temperature, 95
		3.6.4.1	Tolerance to extreme temperatures, 96
			Acknowledgements, 97
			References, 97
3.1	Introduction, 55	4.1	Introduction, 111
3.2	Egg, 55	4.2	Sampling, 111
3.2.1	Egg morphology, 55	4.3	Factors Determining the Composition of Communities, 114
3.2.2	Egg size, 56	4.3.1	Local faunas, 114
3.2.3	Cluster size, 56	4.3.1.1	Geographic differences, 114
3.2.4	Hatching rate, 68	4.3.1.2	Invasion and extinction, 115
3.2.5	Trophic eggs, 69	4.3.1.3	Climatic changes, 115
3.3	Larva, 71	4.3.2	Locality determinants, 116
3.3.1	Larval morphology, 71	4.3.2.1	Prey, 116
3.3.2	Instars, 71	4.3.2.2	Host plant, 120
3.3.3	Development, 72	4.3.2.3	Microclimate, 124
3.3.4	Body size, 73	4.3.2.4	Landscape, 125
3.4	Pupa, 73		
3.4.1	Prepupal stage, 73		
3.4.2	Pupal morphology, 74		
3.4.3	Timing of pupation, 75		
3.4.4	Places of pupation, 75		
3.4.5	Pupal defence, 76		
3.4.6	Colouration and thermal melanism, 76		
3.5	Adult, 76		
3.5.1	Teneral development, 76		
3.5.2	Wings and flight, 77		
3.5.3	Pre-oviposition period, 77		
3.5.4	Size, 78		
3.5.5	Ovarioles, 78		
3.5.6	Mating, 82		
	3.5.6.1 Frequency and duration of mating, 82		
	3.5.6.2 Sperm competition, 83		

4.3.3	Dominance, diversity and niche differentiation, 127	5.2.12	Tritrophic studies, 196
4.4	Coccinellid Communities of Particular Habitats, 128	5.2.13	Food of phytophagous Coccinellidae, 198
4.5	Conclusion, 132	5.2.14	Food of mycophagous Coccinellidae, 200
Acknowledgements, 133		5.3	Quantitative Aspects of Food Relations, 201
References, 133		5.3.1	Effect of physical factors on consumption, 202
		5.3.1.1	Daily consumption rate, 202
		5.3.1.2	Total food consumption, 202
		5.3.2	Effect of prey density on consumption: functional response, 204
		5.3.3	Effects of consumption on growth and reproduction, 207
		5.3.3.1	Larval development, 207
		5.3.3.2	Adult performance, 208
		5.3.4	Conversion and utilization of consumed food, 209
		5.3.5	Aggregative numerical response, 211
		5.3.5.1	Temporal and spatial patterns, 211
		5.3.5.2	Modeling of aggregative responses, 211
		5.3.5.3	Factors other than focal prey density, 211
5.1	Introduction, 142	5.4	Food-Related Behaviour, 213
5.2	Food Specificity, 142	5.4.1	Foraging behaviour, 213
5.2.1	Food range, 142	5.4.1.1	Indirect factors in foraging, 215
	5.2.1.1 Methods for detection of food range, 144	5.4.1.2	The role of senses in foraging, 218
5.2.2	Nutritional suitability of food, 145	5.4.1.3	Finding an oviposition site, 226
5.2.3	Prey size-density hypothesis, 147	5.4.1.4	Foraging of first instars, 233
5.2.4	Euryphagous and stenophagous species / Generalist and specialist species, 150	5.4.1.5	Movement among habitats/patches in the landscape, 234
	5.2.4.1 Generalists, 152	5.4.1.6	Ladybird foraging and ants, 236
	5.2.4.2 Specialists, 154	5.4.2	Prey capture, 238
5.2.5	Mixed and combined diet, 155	5.4.3	Food intake, 241
	5.2.5.1 Complementation across stages, 156	5.5	Conclusions, 242
	5.2.5.2 Prey switching, 157	Acknowledgements, 242	
	5.2.5.3 Prey specialization through selection, 157	References, 243	
5.2.6	Lower quality prey (mostly aphids), 157		
	5.2.6.1 Toxic prey, 157		
	5.2.6.2 Rejected prey, 165		
	5.2.6.3 'Problematic' prey, 167		
5.2.7	Prey other than aphids/coccids, 169		
	5.2.7.1 Developmental stages of Holometabola, 169		
	5.2.7.2 Non-aphid hemipterans, 174		
5.2.8	Cannibalism, 175		
	5.2.8.1 Sibling egg cannibalism, 175		
	5.2.8.2 Non-sibling egg cannibalism, 177		
5.2.9	Non-insect food (pollen, nectar, spores of fungi), 180		
5.2.10	Substitute diets and food supplements (sprays), 185		
5.2.11	Essential foods, 187		

**6. DIAPAUSE/DORMANCY, 275**

Ivo Hodek

<p>6.1 Introduction: Mechanisms and Definitions, 276</p> <ul style="list-style-type: none"> <li>6.1.1 Hibernation and aestivation, 276</li> <li>6.1.2 Termination/completion of diapause, 277</li> <li>6.1.3 Phases of dormancy, 277</li> <li>6.1.4 Endocrinological aspects of adult diapause, 277</li> </ul> <p>6.2 Ecophysiological Regulation of Diapause in Coccinellids, 278</p> <ul style="list-style-type: none"> <li>6.2.1 <i>Coccinella septempunctata</i>, 278           <ul style="list-style-type: none"> <li>6.2.1.1 Central Europe, 278</li> <li>6.2.1.2 Western Europe: France, 282</li> <li>6.2.1.3 Northern Europe, 283</li> <li>6.2.1.4 Mediterranean region, 284</li> <li>6.2.1.5 Nearctic region, 285</li> <li>6.2.1.6 Potential multivoltines in univoltine populations, 286</li> </ul> </li> <li>6.2.2 <i>Coccinella septempunctata brucki</i>, 287           <ul style="list-style-type: none"> <li>6.2.2.1 Central Japan (central Honshu), 287</li> <li>6.2.2.2 Sapporo, Hokkaido (Japan), 288</li> <li>6.2.2.3 Northern Honshu (Japan), 289</li> </ul> </li> <li>6.2.3 <i>Coccinella novemnotata</i>, 289</li> <li>6.2.4 <i>Adalia bipunctata</i>, 290</li> <li>6.2.5 <i>Propylea quatuordecimpunctata</i> and <i>P. dissecta</i>, 291</li> <li>6.2.6 <i>Hippodamia tredecimpunctata</i>, 292</li> <li>6.2.7 <i>Hippodamia convergens</i>, 292</li> <li>6.2.8 <i>Ceratomegilla</i> (=<i>Semiadalia</i>) <i>undecimnotata</i>, 292</li> <li>6.2.9 <i>Harmonia axyridis</i>, 295</li> <li>6.2.10 <i>Coccinella leonina</i> (=<i>repanda</i>), 297</li> <li>6.2.11 <i>Apolinus</i> (=<i>Scymnoides</i>) <i>lividigaster</i> and <i>Illeis</i> (=<i>Leptothea</i>) <i>galbula</i>, 297</li> <li>6.2.12 <i>Harmonia sedecimnotata</i>, 297</li> <li>6.2.13 <i>Chilocorus</i> spp., 298</li> <li>6.2.14 <i>Stethorus punctum picipes</i> and <i>S. japonicus</i>, 299</li> <li>6.2.15 <i>Scymnus</i> (<i>Neopullus</i>) <i>sinuanodus</i>, 300</li> </ul>	<p>6.3 Behaviour Patterns Related to Diapause, 301</p> <ul style="list-style-type: none"> <li>6.3.1 Phases of dormancy behaviour, 301           <ul style="list-style-type: none"> <li>6.3.1.1 Pre-diapause, 301</li> <li>6.3.1.2 Migration, 302</li> <li>6.3.1.3 Flight and methods for its study, 303</li> <li>6.3.1.4 Aggregations, 305</li> <li>6.3.1.5 Emergence from dormancy sites, 309</li> </ul> </li> <li>6.3.2 Behaviour of individual species, 309           <ul style="list-style-type: none"> <li>6.3.2.1 <i>Ceratomegilla</i> (=<i>Semiadalia</i>) <i>undecimnotata</i>, 309</li> <li>6.3.2.2 Hypsotactic species <i>Harmonia axyridis</i>, <i>Har. conformis</i> and <i>Aiolocaria hexaspilota</i> (=<i>mirabilis</i>), 310</li> <li>6.3.2.3 <i>Coccinella septempunctata</i> and other species dormant in the litter, 311</li> <li>6.3.2.4 <i>Coleomegilla maculata</i>, 313</li> <li>6.3.2.5 <i>Adalia bipunctata</i>, 313</li> <li>6.3.2.6 <i>Myrrha octodecimguttata</i> and other forest species which hibernate in bark crevices, 314</li> <li>6.3.2.7 <i>Hippodamia convergens</i>, 314</li> <li>6.3.2.8 <i>Hippodamia quinquesignata</i>, 316</li> <li>6.3.2.9 Hibernation of mycophagous and phytophagous species, 316</li> </ul> </li> </ul> <p>6.4 Anatomical and Physiological Changes Related to Dormancy, 316</p> <ul style="list-style-type: none"> <li>6.4.1 Anatomical state, 316           <ul style="list-style-type: none"> <li>6.4.1.1 Fat body and digestive tract, 316</li> <li>6.4.1.2 Ovary, spermatheca, 317</li> <li>6.4.1.3 Male gonads, 321</li> <li>6.4.1.4 Flight muscles, 323</li> </ul> </li> <li>6.4.2 Metabolic changes related to diapause, 323           <ul style="list-style-type: none"> <li>6.4.2.1 Lipids, 323</li> <li>6.4.2.2 Glycogen, 325</li> </ul> </li> </ul>
---	--

---

6.4.2.3	Water, 326	7.9.1	Defensive mechanisms of coccinellids against intraguild predation, 355
6.4.2.4	Metabolic rate, 326	7.9.1.1	Defence of all stages, 355
6.4.3	<i>Corpora allata</i> and regulation of vitellogenesis, 327	7.9.1.2	Defence of eggs, 356
6.4.4	Cold-hardiness, 328	7.9.1.3	Defence of larvae, 356
6.5	Conclusions and Lacunae in Knowledge, 332	7.9.1.4	Defence of moulting individuals and pupae, 357
	Acknowledgements, 333	7.9.1.5	Defence of adults, 357
	References, 333	7.9.2	Intraguild predation on coccinellids, 358
<b>7. INTRAGUILD INTERACTIONS, 343</b>		7.9.2.1	Intraguild predation by intraguild predators, 358
Éric Lucas		7.9.2.2	Intraguild predation by intraguild parasitoids, 358
7.1	Scope, 344	7.9.2.3	Intraguild predation by intraguild pathogens, 359
7.2	Coccinellids as Guild Members, 344	7.10	Coccinellids Interacting with Intraguild Ants, 359
7.3	Coccinellids as Neutralists, 344	7.10.1	Non-myrmecophilous coccinellid species, 359
7.3.1	Temporal guild partition, 344	7.10.2	Myrmecophilous coccinellid species, 360
7.3.2	Spatial guild partition, 344	7.11	Applied Aspects of Intraguild Interactions, 360
7.3.3	Thermal guild partition, 344	7.11.1	Conservation, 360
7.3.4	Body size guild partition, 345	7.11.2	Biological control, 361
7.4	Coccinellids as Interacting Organisms, 345	7.11.2.1	Intraguild predation and biocontrol, 361
7.5	Coccinellids as Intraguild Commensalists and Mutualists, 345	7.11.2.2	Facilitation and biocontrol, 361
7.6	Coccinellids as Competitors, 347	7.11.2.3	Ants and biocontrol, 362
7.6.1	Exploitative competition, 347	7.11.2.4	Intraguild interactions and biocontrol approaches, 362
7.6.2	Apparent competition, 349	7.12	Interguild Effects, 362
7.7	Coccinellids as Vectors of Male-Killing Bacteria, 349	7.13	Conclusion, 363
7.8	Coccinellids as Intraguild Predators, 349	References, 364	
7.8.1	General rules of intraguild predation involving coccinellid predators, 349	<b>8. NATURAL ENEMIES OF LADYBIRD BEETLES, 375</b>	
7.8.2	Occurrence of intraguild predation by coccinellids, 350	Piotr Ceryngier, Helen E. Roy and Remy L. Poland	
7.8.3	Intraguild predation on intraguild coccinellids (by coccinellids), 351	8.1	Introduction, 376
7.8.4	Intraguild predation on intraguild non-coccinellid predators, 352	8.2	Predation and Related Phenomena, 376
7.8.4.1	Intraguild neuropterans, 352	8.2.1	Anti-predator defences, 376
7.8.4.2	Intraguild dipterans, 352	8.2.1.1	Aposematic colouration and other visual signals, 376
7.8.4.3	Intraguild hemipterans, 353	8.2.1.2	Reflex bleeding, 376
7.8.4.4	Other intraguild predators, 353		
7.8.5	Intraguild predation on intraguild parasitoids, 353		
7.8.6	Intraguild predation on intraguild pathogens, 354		
7.8.7	Coccinellids as top predators, 354		
7.9	Coccinellids as Intraguild Prey, 355		

	8.2.1.3	Morphological anti-predator adaptations, 377	8.4	Parasites and Pathogens, 411
8.2.2	Vertebrate predators, 377		8.4.1	Acarina, 411
8.2.3	Invertebrate predators, 377		8.4.1.1	Phoretic mites, 411
8.2.4	Hemiptera-tending ants, 379		8.4.1.2	<i>Coccipolipus</i> Husband (Prostigmata: Podapolipidae), 411
8.2.5	Social aphids with a soldier caste, 382		8.4.2	Nematodes, 414
8.3	Parasitoids, 383		8.4.2.1	Allantonematidae (Tylenchida), 414
8.3.1	General characteristics of parasitoids of ladybirds, 390		8.4.2.2	Mermithidae (Mermithida), 415
	8.3.1.1	Host specificity of parasitoids recorded from Coccinellidae, 390	8.4.3	Fungal pathogens, 415
	8.3.1.2	Parasitoids of different subfamilies of Coccinellidae, 390	8.4.3.1	Hypocreales (Ascomycota), 415
	8.3.1.3	Parasitism in different developmental stages of Coccinellidae, 390	8.4.3.2	<i>Hesperomyces</i> spp. (Ascomycota: Laboulbeniales, Laboulbeniaceae), 416
8.3.2	Review of the more important taxa, 391		8.4.3.3	Nosematidae (Microsporidia), 418
	8.3.2.1	<i>Dinocampus</i> Foerster (Hymenoptera: Braconidae, Euphorinae), 391	8.4.4	Protozoan pathogens, 419
	8.3.2.2	<i>Uga</i> Girault (Hymenoptera: Chalcididae), 399	8.4.4.1	Septate eugregarines (Apicomplexa: Eugregarinida: Septatorina), 419
	8.3.2.3	<i>Cowperia</i> Girault (Hymenoptera: Encyrtidae), 399	8.4.5	Bacteria, 421
	8.3.2.4	<i>Homalotylus</i> Mayr (Hymenoptera: Encyrtidae), 400	8.4.5.1	General pathogenic bacteria, 421
	8.3.2.5	<i>Nothoserphus</i> Brues (Hymenoptera: Proctotrupidae), 404	8.4.5.2	Male-killing bacteria, 421
	8.3.2.6	<i>Metastenus</i> Walker (Hymenoptera: Pteromalidae), 405	8.5	Impact of Natural Enemies on Ladybird Populations, 425
	8.3.2.7	<i>Phalacrotophora</i> Enderlein (Diptera: Phoridae), 406	8.5.1	Impact on phytophagous Coccinellidae, 426
	8.3.2.8	<i>Oomyzus scaposus</i> (Thomson) (Hymenoptera: Eulophidae, Tetrastichinae), 408	8.5.2	Impact on predatory Coccinellidae, 426
	8.3.2.9	<i>Pediobius foveolatus</i> (Crawford) (Hymenoptera: Eulophidae, Encyrtidae), 409	8.5.3	Concluding note, 428
				References, 428
				<b>9. COCCINELLIDS AND SEMIOCHEMICALS, 444</b>
				<i>Jan Pettersson</i>
			9.1	Introduction, 445
			9.2	Aposematism and Reflex Bleeding Chemistry, 445
			9.2.1	Reflex bleeding, 445
			9.2.2	Reflex bleeding substances, 445
			9.2.2.1	Experiments, 445
			9.2.2.2	Identification, 445
			9.2.2.3	Sources, 445

9.2.2.4	Production cost, 446	9.7	Hibernation and Aggregation, 457	
9.2.2.5	Age and stage modifications, 447	9.8	Habitat Preferences: Responses to Plants and Plant Volatiles, 457	
9.2.2.6	Species specific chemistry, 448	9.8.1	Habitat selection, 457	
9.2.3	Relation to enemies and competitors, 449	9.8.2	Plant stand traits, 457	
9.2.3.1	Parasitoids, 449	9.8.3	Avoided plants, 458	
9.2.3.2	Ants, 449	9.8.4	Plant–plant interactions/attractive plants, 458	
9.2.3.3	Spiders, 449	9.9	Conclusions and Future Challenges, 459	
9.3	Semiochemicals Related to Food, 449		References, 460	
9.3.1	Plant volatiles, 449	<b>10. QUANTIFYING THE IMPACT OF COCCINELLIDS ON THEIR PREY, 465</b>		
9.3.1.1	Herbivore-induced plant volatiles, 449	<i>J. P. Michaud and James D. Harwood</i>		
9.3.1.2	Adult receptors for food semiochemicals, 450	10.1	Introduction, 466	
9.3.1.3	Responses of larvae, 450	10.2	Assays of Consumption, 467	
9.3.1.4	Prey sex pheromones: predator kairomones, 451	10.3	Indirect Impacts, 468	
9.3.2	Prey alarm pheromones, 451	10.4	Traditional Approaches to the Study of Predation, 468	
9.3.2.1	Aphid alarm pheromone, 451	10.4.1	Selective exclusion, 469	
9.3.2.2	Mechanism for modified responses, 451	10.4.2	Field cages, 469	
9.3.2.3	EBF release: aphid individual risks, 451	10.4.3	Cage inclusion, 470	
9.3.3	Toxic substances in prey, 451	10.4.4	Manual removal, 471	
9.3.4	Feeding stimulants for phytophagous coccinellids, 452	10.5	Statistical Approaches, 471	
9.3.5	Learning, 452	10.6	Resolving Coccinellid Impact within Complex Communities, 472	
9.4	Mating and Sex Pheromones, 453	10.6.1	Multi-species combinations, 472	
9.4.1	Chemoreceptors on the antennae, 453	10.6.2	Demographic-based estimates, 473	
9.4.2	Hydrocarbons on elytra, 454	10.7	Post-Mortem Analysis of Predation, 474	
9.4.3	Ultrastructure of the integumentary glands, 454	10.7.1	Antibody-based analysis of predation, 474	
9.5	Oviposition, 454	10.7.2	Protein marking for predation analysis, 477	
9.5.1	Oviposition deterrence: oviposition deterrence pheromones, 454	10.7.3	Detection of prey-specific DNA, 478	
9.5.2	Species-specificity of oviposition deterrence pheromone substances, 454	10.8	Conclusions, 481	
9.5.3	Active substances in oviposition deterrence pheromone tracks, 455		References, 482	
9.5.4	Relation to other aphid enemies, 455	<b>11. COCCINELLIDS IN BIOLOGICAL CONTROL, 488</b>		
9.5.5	Aphid abundance, 456	<i>J. P. Michaud</i>		
9.6	Egg and Pupa Protection, 456	11.1	Introduction, 489	
9.6.1	Protection of eggs, 456	11.2	The Roles of Coccinellidae in Biological Control, 489	
9.6.2	Protection of pupae, 456	11.2.1	Prey specificity, 489	
		11.2.2	Generalist coccinellids, 490	
		11.2.3	Intraguild predation, 490	

- 11.3 Scale Insects Versus Aphids as Targets of Exotic Introductions, 491  
    11.3.1 Coccidophagous coccinellids, 491  
    11.3.2 Aphidophagous coccinellids, 492  
    11.3.3 Invasive coccinellids, 493  
    11.3.4 Competitive displacement, 494
- 11.4 Augmentation of Coccinellids, 495  
    11.4.1 The mealybug destroyer, 496  
    11.4.2 Redistribution of coccinellids, 496  
    11.4.3 Selection of source material for augmentation, 497  
    11.4.4 Voltinism and diapause, 497  
    11.4.5 Dietary requirements, 497  
    11.4.6 Life stages for release, 498
- 11.5 Conservation, 498  
    11.5.1 Alternative or supplementary food, 499  
    11.5.2 Hibernation refuges, 500  
    11.5.3 Habitat management, 500  
        11.5.3.1 Strip-harvesting, 501  
        11.5.3.2 Floral diversity (non-crop plants), 501  
        11.5.3.3 Intercropping, 501  
        11.5.3.4 Reduced tillage, 502
- 11.6 Ancillary Factors Influencing Biological Control by Coccinellids, 502  
    11.6.1 Ant-attendance of aphid colonies, 502  
    11.6.2 Timing of arrival in annual crops, 503
- 11.6.3 Interaction of biological control by coccinellids with plant structure and chemistry, 504  
    11.6.4 Selective use of pesticides, 506
- 11.7 Conclusions, 509  
References, 509
- 12. RECENT PROGRESS AND POSSIBLE FUTURE TRENDS IN THE STUDY OF COCCINELLIDAE, 520**  
*Helmut E van Emden and Ivo Hodek*  
Acknowledgements, 525  
References, 525
- APPENDIX: LIST OF GENERA IN TRIBES AND SUBFAMILIES, 526**  
*Ondřich Nedvěd and Ivo Kovář*
- Coccinellidae Latreille, 1807, 527  
Chilocorinae Mulsant, 1846 [2:25], 527  
Coccidulinae Mulsant, 1846 [6:46], 527  
Coccinellinae Latreille, 1807 [5:94], 528  
Epilachninae Mulsant, 1846 [4:24], 529  
Exoplectrinae Crotch, 1874 [2:24], 529  
Microweiseinae Leng, 1920 [4:23], 529  
Ortaliinae Mulsant, 1850 [2:14], 530  
Scymninae Mulsant, 1846 [11:51], 530  
Sticholotidinae Weise, 1901 [6:58], 531
- Subject index, 532
- Colour plate pages fall between pp. 250 and pp. 251

# CONTRIBUTORS

**PIOTR CERYNGIER** *Centre for Ecological Research,  
Polish Academy of Sciences, Dziekanow Lesny, 05-092  
Lomianki, POLAND*

**HELMUT F. VAN EMDEN** *School of Biological Sciences,  
University of Reading, Whiteknights, Reading RG6  
6AS UK*

**EDWARD W. EVANS** *Department of Biology, Utah State University, Logan, UT 84322 USA*

**JAMES D. HARWOOD** *Department of Entomology,  
University of Kentucky, Lexington, KY 40546-0091  
USA*

**IVO HODEK** *Institute of Entomology, Academy of Sciences, CZ 37005 České Budějovice, CZECH REPUBLIC*

**ALOIS HONĚK** *Department of Entomology, Crop Research Institute, CZ 16106 Prague 6, CZECH REPUBLIC*

**IVO KOVÁŘ** *Emer. Scientist of the National Museum, Prague; Current address: Zichovec, CZECH REPUBLIC*

**ERIC LUCAS** *Département des Sciences Biologiques, Université du Québec à Montréal, C.P. 8888 Succ. Centre-ville, Montréal, Québec H3C 3P8 CANADA*

**J.P. MICHAUD** *Department of Entomology, Kansas State University, 1232 240th Ave., Hays, KS 67601 USA*

**OLDRICH NEDVĚD** *Faculty of Science, University of South Bohemia and Institute of Entomology, Academy of Sciences, CZ 37005 České Budějovice, CZECH REPUBLIC*

**JAN PETTERSSON** *Department of Ecology, Swedish University of Agricultural Sciences, Box 7044, SE-750 07 Uppsala, SWEDEN*

**REMY L. POLAND** *Department of Genetics, University of Cambridge, Cambridge CB2 3EH UK, recent address: Clifton College, 32 College Road, Clifton, Bristol, BS8 3JH, UK*

**HELEN E. ROY** *NERC Centre for Ecology and Hydrology, Crowmarsh Gifford, Oxfordshire OX10 8BB UK*

**JOHN J. SLOGGETT** *Maastricht Science Programme, Maastricht University, P.O. Box 616, 6200 MD Maastricht, THE NETHERLANDS*

# PREFACE

For more than a decade, no volume on the general aspects of ladybirds has been published, although the predaceous major part of this coleopteran family represents an important component of the natural enemies of Sternorrhyncha (aphids, coccids, aleyrodids and psyllids). These sucking insects are among the most dangerous pests of crops, as under suitable conditions their populations increase exponentially, especially when parthenogenesis and viviparity occur.

Although classical biological control with coccinellids has recently been mostly abandoned, the increasing concern about chemical pest control has increased the need for modern types of biological control, mostly involving conservation and augmentation, within the framework of integrated pest management. For the success of these sophisticated methods of control, precise knowledge of behaviour and ecological relations is indispensable.

Such knowledge, particularly in the areas of ethology and molecular genetics, has accumulated in the last decade and reviewing these areas is an important novel contribution of this book. We hope that, at this stage, it will help to improve conservation and augmentation control, but also that it will stimulate further research.

The book was very unfortunately interrupted by the premature death of Professor Michael Majerus, Department of Genetics, Cambridge University, who was our proposed author for Chapters 2 and 8. We were extremely lucky to find very able successors, John Sloggett (Chapter 2), and Helen Roy and Remy Poland

(for parasites and pathogens in Chapter 8), who fulfilled their task with great success. Many thanks!

We should like to thank the other authors of chapters for contributing their expertise so readily and for reacting so positively and constructively to our suggestions for revision. We are also grateful for the expert help we have received with taxonomic nomenclature: Dr O. Nedvěd of the University of South Bohemia, Czech Republic, and from the UK Drs V.F. Eastop, C.H. Lyal and D.J. Williams (Natural History Museum), Drs R.T.V. Fox and S.L. Jury (University of Reading) and Professor D.L.J. Quicke (Imperial College). Dr Lyal deserves additional thanks for his advice in relation to Chapter 1.

We should also like to thank Dr Ward Cooper of Wiley-Blackwell for enthusiastically agreeing to publish our book and Mr Kelvin Matthews for his help with the publication process. We should also like to thank the other Wiley-Blackwell staff who have worked so helpfully and efficiently on the production of this book.

**Ivo Hodek**  
**České Budějovice, Czech Republic**

**Helmut F. van Emden**  
**Reading, UK**

**Alois Honěk**  
**Prague, Czech Republic**

**March 2012**

# INTRODUCTION

The reader may find the following information helpful in order to use the book more easily.

To keep the flow of the text unbroken and to save repetition and space, taxonomic affiliations of organisms and species authorities are given only in the Taxonomic Glossary (following section, while the Subject Index is at the end of book, as usual). Readers should also note that the Latin names as given in this glossary are used in the text and tables, and therefore may not be the same as the older Latin names given in the original papers cited. However, because the species names

have usually not changed, and the older names are also listed in the glossary, any confusion will be avoided.

The very common generic names are always abbreviated as follows throughout the text of all chapters: *A.* for *Adalia*; *C.*, *Coccinella*; *Cer.*, *Ceratomegilla*; *Chil.*, *Chilocorus*; *Col.*, *Coleomegilla*; *Har.*, *Harmonia*; *Hip.*, *Hippodamia*; *P.*, *Propylea*.

Some phenomena are discussed in more than one chapter in the book, but from different angles; the reader's attention is directed to this by cross-references to the section number.

# TAXONOMIC GLOSSARY

With common names where appropriate

## **Coleoptera: Family Coccinellidae**

Synonyms [in square brackets] as well as currently valid names are listed in alphabetical order.

*Adalia bipunctata* (L.) – two spot ladybird [*Adalia fasciatopunctata* (Faldermann)]  
*Adalia conglomerata* (L.)  
*Adalia decempunctata* (L.) – ten spot ladybird  
*Adalia deficiens* Mulsant  
[*Adalia fasciatopunctata* (Faldermann)] = *Adalia bipunctata* (L.)  
[*Adalia flavomaculata* (De Geer)] = *Lioadalia flavomaculata* (De Geer)  
*Adalia tetraspilota* (Hope)  
[*Adonia*] = *Hippodamia*  
[*Adonia arctica* (Schneider)] = *Hippodamia arctica* (Schneider)  
[*Adonia variegata* (Goeze)] = *Hippodamia variegata* (Goeze)  
*Afidenta misera* (Weise) [*Afidenta mimetica* Dieke]  
*Aphidentula bisquadripunctata* (Gyllenhal) [*Epilachna bisquadripunctata* (Gyllenhal)]  
*Afissula rana* Kapur  
*Afissula sanscrita* (Crotch)  
*Aiolocaria hexaspilota* (Hope) [*Aiolocaria mirabilis* (Motschulsky)]  
[*Aiolocaria mirabilis* (Motschulsky)] = *Aiolocaria hexaspilota* (Hope)  
*Alloneda dodecaspilota* (Hope)  
*Anatis halonis* Lewis  
*Anatis labiculata* (Say) – fifteen spotted lady beetle  
*Anatis mali* (Say) – eyespotted lady beetle  
*Anatis ocellata* (L.) – eyed ladybird

*Anatis quindecimpunctata* (DeGeer)  
*Anegleis cardoni* (Weise)  
[*Anisolemnia dilatata* (F.)] = *Megalocaria dilatata* (F.)  
*Anisolemnia tetrasticta* Fairmaire  
*Anisosticta bitriangularis* (Say)  
*Anisosticta novemdecimpunctata* (L.) – water ladybird  
*Anisosticta sibirica* Bielawski  
*Aphidecta obliterate* (L.) – larch ladybird  
*Apolinus lividigaster* (Mulsant) [*Scymnodes lividigaster* (Mulsant)]  
*Axinoscymnus cardilobus* Ren & Pang  
[*Azya trinitatis* (Marshall)] = *Pseudoazyxa trinitatis* (Marshall)  
*Azya orbignera* Mulsant  
*Brachiacantha quadripunctata* Melsheimer  
*Brachiacantha ursina* (F.)  
*Brumoides septentrionis* (Weise)  
*Brumoides suturalis* (F.) – three-striped lady beetle  
[*Brumus suturalis* (Mani)]  
[*Brumus quadripustulatus* (L.)] = *Exochomus quadripustulatus* (L.)  
[*Brumus suturalis* (Mani)] = *Brumoides suturalis* (F.)  
*Bulaea lichatschovi* (Hummel)  
*Callicaria superba* (Mulsant)  
*Calvia albida* Bielawski  
*Calvia decemguttata* (L.)  
*Calvia duodecimmaculata* Gebler  
*Calvia muiri* Timberlake [*Eocaria muiri* Timberlake]  
*Calvia quatuordecimguttata* (L.) – cream-spot ladybird  
*Calvia quindecimguttata* (F.)  
*Calvia shiva* Kapur  
*Calvia shiva pasupati* Kapur  
*Calvia shiva pinaki* Kapur  
*Calvia shiva trilochana* Kapur  
*Ceratomegilla barovskii kiritschenkoi* (Semenov-Tian-Shanski) [*Spiladelpha barovskii kiritschenkoi* Semenov-Tian-Shanski]

- Ceratomegilla notata* (Laicharting)  
*Ceratomegilla undecimnotata* (Schneider) [*Semiadalia undecimnotata* (Schneider)]  
*Cheiromenes lunata* (F.)  
*Cheiromenes propinquua vicina* (Mulsant) [*Cheiromenes vicina* (Mulsant), *Cydonia vicina nilotica* Mulsant]  
[*Cheiromenes sexmaculata* (F.)] = *Menochilus sexmaculatus* (F.)  
*Cheiromenes sulphurea* (Olivier)  
[*Cheiromenes vicina*] = *Cheiromenes propinquua vicina* (Mulsant)  
[*Chilocorus baylei* (Blackburn)] = *Chilocorus malasiae* Crotch  
*Chilocorus bijugus* Mulsant  
*Chilocorus bipustulatus* (L.) – heather ladybird  
*Chilocorus braeti* Weise  
*Chilocorus cacti* (L.)  
*Chilocorus circumdatus* (Gyllenhal)  
*Chilocorus discoideus* Crotch  
*Chilocorus distigma* Klug  
*Chilocorus geminus* Zaslavsky  
*Chilocorus hauseri* Weise  
*Chilocorus hexacyclus* Smith  
*Chilocorus infernalis* Mulsant  
*Chilocorus inornatus* Weise  
*Chilocorus kuwanae* Silvestri  
*Chilocorus malasiae* Crotch [*Chilocorus baylei* (Blackburn)]  
*Chilocorus nigripes* Mader  
*Chilocorus nigritus* (F.)  
*Chilocorus orbus* Casey  
*Chilocorus quadrimaculatus* (Weise)  
*Chilocorus renipustulatus* (Scriba) – kidney-spot ladybird  
*Chilocorus rubidus* Hope  
*Chilocorus similis* (Rossi)  
*Chilocorus stigma* (Say) – twice-stabbed lady beetle  
*Chilocorus tricyclus* Smith  
*Chnootriba similis* (Thunberg)  
*Cleobora mellyi* (Mulsant) – Tasmanian ladybird  
*Clitostethus arcuatus* (Rossi)  
*Clitostethus oculatus* (Blatchley) [*Nephaspis oculatus* (Blatchley)]  
*Coccidophilus citricola* Brèthes  
*Coccidula rufa* (Herbst)  
*Coccidula scutellata* (Herbst)  
[*Coccinella algerica* Kovář] = *Coccinella septempunctata algerica* Kovář  
*Coccinella californica* Mannerheim  
*Coccinella explanata* Miyatake  
*Coccinella hieroglyphica* L. – hieroglyphic ladybird  
*Coccinella leonina transversalis* F. [*Coccinella repanda* Thunberg, *Coccinella transversalis* F.]  
*Coccinella luteopicta* (Mulsant)  
*Coccinella magnifica* Redtenbacher – scarce seven spot ladybird [*Coccinella divaricata* Olivier])  
*Coccinella monticola* Mulsant  
*Coccinella nigrovittata* Kapur [*Tytthaspis trilineata* (Weise)]  
*Coccinella novemnotata* Herbst – nine-spotted lady beetle  
*Coccinella quinquepunctata* L. – five spot ladybird  
*Coccinella reitteri* Weise  
[*Coccinella repanda* Thunberg] = *Coccinella leonina* F.  
*Coccinella septempunctata* L. – seven spot ladybird  
*Coccinella septempunctata algerica* Kovář [*Coccinella algerica* Kovář]  
*Coccinella septempunctata brucki* Mulsant  
[*Coccinella sinuatomarginata* Faldermann] = *Coccinula sinuatomarginata* (Faldermann)  
[*Coccinella transversalis* F.] = *Coccinella leonina transversalis* F.  
*Coccinella transversoguttata* Faldermann – transverse lady beetle  
*Coccinella transversoguttata richardsoni* Brown  
*Coccinella trifasciata* L. – three-banded lady beetle  
*Coccinella undecimpunctata* L. – eleven spot ladybird  
[*Coccinella undecimpunctata aegyptiaca* Reiche] = *Coccinella undecimpunctata menetriesi* Mulsant  
*Coccinella undecimpunctata menetriesi* Mulsant  
[*Coccinella undecimpunctata aegyptiaca* Reiche]  
*Coccinula crotchi* (Lewis)  
*Coccinula quatuordecimpustulata* (L.)  
*Coccinula redimita* (Weise)  
*Coccinula sinensis* (Weise)  
*Coccinula sinuatomarginata* (Faldermann) [*Coccinella sinuatomarginata* Faldermann]  
*Coelophora biplagiata* Swartz [*Lemnia biplagiata* (Swartz)]  
*Coelophora bissellata* Mulsant  
*Coelophora duvaucelii* (Mulsant)  
*Coelophora inaequalis* (F.) – common Australian lady beetle  
*Coelophora mulsanti* (Montrouzier)  
*Coelophora quadriplagiata* Fauvel  
*Coelophora saucia* Mulsant  
*Coleomegilla maculata* (DeGeer) – spotted lady beetle  
*Coleomegilla maculata fuscilabris* (Mulsant)

- Coleomegilla maculata lengi* Timberlake – twelve-spotted ladybeetle  
*Coleomegilla quadrifasciata* (Schoenherr)  
*Cryptognatha simillima* Sicard  
*Cryptognatha nodiceps* Marshall  
*Cryptognatha signata* Korschefsky  
*Cryptogonus ariasi* (Mulsant)  
*Cryptogonus kapuri* Ghorpade  
*Cryptogonus orbiculus* (Gyllenhal)  
*Cryptogonus postmedialis* Kapur  
*Cryptogonus quadriguttatus* (Weise)  
*Cryptolaemus montrouzieri* Mulsant – mealybug destroyer  
*Curinus coeruleus* Mulsant – metallic blue lady beetle  
*Cycloneida ancoralis* (Germar)  
*Cycloneida limbifer* Casey  
*Cycloneida munda* (Say) – polished lady beetle  
*Cycloneida polita* Casey  
*Cycloneida sanguinea* (L.) – blood-red lady beetle  
[*Cydonia vicina nilotica* Mulsant] = *Cheilomenes propinqua vicina* (Mulsant)  
*Declivitata* spp.  
*Delphastus catalinae* (Horn)  
*Delphastus pusillus* (LeConte)  
*Diomus austrinus* Gordon  
*Diomus hennesseyi* Fürsch  
*Diomus pumilio* Weise  
*Diomus seminulus* (Mulsant)  
*Diomus thoracicus* F.  
[*Eocaria muiri* Timberlake] = *Calvia muiri* (Timberlake)  
*Epilachna admirabilis* Crotch  
[*Epilachna bisquadripunctata* (Gyllenhal)] = *Aphidentula bisquadripunctata* (Gyllenhal)  
[*Epilachna boisduvali* (Mulsant)] = *Henosepilachna boisduvali* Mulsant  
*Epilachna borealis* (F.) – squash beetle  
*Epilachna canina* (F.)  
[*Epilachna 'chrysomelina'* (F.)]; used variably either for  
*Henosepilachna argus* (Geoffroy) or *Henosepilachna elaterii* (Rossi) or *Henosepilachna vigintioctopunctata* (F.)  
[*Epilachna cucurbitae*] = *Henosepilachna sumbana* Bielawski  
*Epilachna defecta* Mulsant  
*Epilachna dregei* Mulsant – potato ladybird  
*Epilachna dumerili* Mulsant  
*Epilachna eckloni* Mulsant  
[*Epilachna enneasticta* Mulsant] = *Henosepilachna enneasticta* (Mulsant)  
*Epilachna eusema* (Weise)
- Epilachna karisimbica* Weise  
*Epilachna marginella* (F.)  
*Epilachna marginicollis* (Hope)  
*Epilachna mexicana* (Guérin-Méneville)  
*Epilachna mystica* Mulsant  
*Epilachna nigrolimbata* Thomson  
[*Epilachna niponica* Lewis] = *Henosepilachna niponica* (Lewis)  
*Epilachna paenulata* (Germar)  
[*Epilachna philippinensis* Dieke] = *Henosepilachna vigintisexpunctata* (Boisduval)  
[*Epilachna pusillanima* Mulsant] = *Henosepilachna pusillanima* (Mulsant)  
[*Epilachna pustulosa* Kōno] = *Henosepilachna pustulosa* (Kōno)  
*Epilachna quadricollis* (Dieke)  
[*Epilachna septima* Dieke] = *Henosepilachna septima* (Dieke)  
[*Epilachna sparsa orientalis* Dieke] = *Henosepilachna vigintioctopunctata* (F.)  
*Epilachna undecimvariolata* (Boisduval)  
*Epilachna varivestis* Mulsant – mexican bean beetle  
[*Epilachna vigintioctomaculata* Motschulsky] = *Henosepilachna vigintioctomaculata* (Motschulsky)  
[*Epilachna vigintioctomaculata* Motschulsky] = *Henosepilachna vigintioctomaculata* (Motschulsky)  
*Epilachna vigintisexpunctata* (Boisduval)  
[*Epilachna yasutomii* (Katakura)] = *Henosepilachna yasutomii* Katakura  
*Epiverta chelonia* (Mader)  
*Eriopis connexa* (Germar)  
*Exochomus childreni* Mulsant  
[*Exochomus concavus* Fürsch] = *Parexochomus troberti concavus* (Fürsch)  
*Exochomus flavipes* (Thunberg)  
*Exochomus flavidiventris* Mader  
*Exochomus fulvimanus* Weise  
[*Exochomus lituratus* (Gorham)] = *Priscibrumus lituratus* (Gorham)  
[*Exochomus melanocephalus* (Zoubkoff)] = *Parexochomus melanocephalus* (Zoubkoff)  
[*Exochomus nigromaculatus* (Goeze)] = *Parexochomus nigromaculatus* (Goeze)  
*Exochomus quadripustulatus* (L.) – pine ladybird  
[*Brumus quadripustulatus* (L.)]  
[*Exochomus troberti* Mulsant] = *Parexochomus troberti* (Mulsant)  
*Halmus chalybeus* (Boisduval) – steelblue lady beetle  
[*Orcus chalybeus* (Boisduval)]  
[*Halyzia hauseri* (Mader)] = *Macroilleis hauseri* (Mader)

- Halyzia sanscrita* Mulsant  
*Halyzia sedecimguttata* (L.) – orange ladybird  
*Halyzia straminea* (Hope)  
*Halyzia tschitscherini* Semenov  
*Harmonia antipoda* (Mulsant in White) – antipodean ladybird  
*Harmonia axyridis* (Pallas) – harlequin ladybird or Asian multi-colored ladybeetle  
[*Harmonia breiti* Mader] = *Harmonia expalliatia* Sicard  
*Harmonia conformis* (Boisduval)  
*Harmonia dimidiata* (F.) [*Leis dimidiata* (F.)]  
*Harmonia eucharis* (Mulsant)  
*Harmonia expalliatia* Sicard [*Harmonia breiti* Mader]  
*Harmonia quadripunctata* (Pontoppidan) – cream-streaked ladybird  
*Harmonia octomaculata* (F.)  
*Harmonia sedecimnotata* (F.)  
*Harmonia yedoensis* (Takizawa)  
*Henosepilachna argus* (Geoffroy) – bryony ladybird  
*Henosepilachna bifasciata* (L.)  
*Henosepilachna boisduvali* (Mulsant) [*Epilachna boisduvali* Mulsant]  
*Henosepilachna dodecastigma* (Wiedemann)  
*Henosepilachna elaterii* (Rossi) [*Epilachna chrysomelina* (F.)]  
*Henosepilachna enneasticta* (Mulsant) [*Epilachna enneasticta* Mulsant]  
*Henosepilachna guttata* (F.)  
*Henosepilachna indica* (Mulsant)  
*Henosepilachna niponica* (Lewis) [*Epilachna niponica* Lewis]  
*Henosepilachna ocellata* (Redtenbacher)  
*Henosepilachna processa* Li  
*Henosepilachna pusillanima* (Mulsant) [*Epilachna pusillanima* Mulsant]  
*Henosepilachna pustulosa* (Kôno) [*Epilachna pustulosa* Kôno]  
*Henosepilachna septima* (Dieke) [*Epilachna septima* Dieke]  
*Henosepilachna sumbana* Bielawski [*Epilachna cucurbitae* Richards, *Henosepilachna cucurbitae* Richards]  
*Henosepilachna vigintioctomaculata* (Motschulsky)  
[*Epilachna vigintioctomaculata* Motschulsky]  
*Henosepilachna vigintioctopunctata* (F.) [*Epilachna vigintioctopunctata* (F.), *Epilachna sparsa orientalis* Dieke]  
*Henosepilachna vigintisexpunctata* (Boisduval)  
[*Epilachna philippinensis* Dieke]  
*Henosepilachna yasutomii* Katakura [*Epilachna yasutomii* (Katakura)]
- Hippodamia arctica* (Schneider) [*Adonia arctica* (Schneider)]  
*Hippodamia caseyi* Johnson  
*Hippodamia convergens* Guerin – convergent ladybeetle  
*Hippodamia glacialis* (F.) – glacial lady beetle  
*Hippodamia parenthesis* (Say) – parenthesis lady beetle  
*Hippodamia quinquesignata* (Kirby)  
*Hippodamia quinquesignata punctulata* Le Conte  
[*Hippodamia quinquesignata ambigua* LeConte]  
*Hippodamia septemmaculata* (DeGeer)  
*Hippodamia sinuata* Mulsant  
*Hippodamia tredecimpunctata* (L.) – thirteen spot ladybird  
*Hippodamia variegata* (Goeze) – Adonis ladybird, variegated lady beetle [*Adonia variegata* (Goeze)]  
*Hyperaspis aestimabilis* Mader  
*Hyperaspis bigeminata* (Randall)  
*Hyperaspis binotata* (Say)  
*Hyperaspis campestris* (Herbst)  
[*Hyperaspis congressis* Watson] = *Hyperaspis conviva* Casey  
*Hyperaspis conviva* Casey [*Hyperaspis congressis* Watson]  
*Hyperaspis desertorum* Weise  
*Hyperaspis lateralis* Mulsant  
*Hyperaspis notata* Mulsant  
*Hyperaspis pantherina* Fürsch  
*Hyperaspis raynevali* Mulsant  
*Hyperaspis reppensis* (Herbst)  
*Hyperaspis senegalensis* Mulsant  
*Hyperaspis senegalensis hottentotta* Mulsant  
*Hyperaspis sphaeridoides* Mulsant  
*Hyperaspis undulata* (Say)  
*Illeis bielawskii* Ghorpade  
*Illeis cincta* (F.)  
*Illeis galbula* (Mulsant) [*Leptothea galbula* (Mulsant)]  
*Illeis koebelei* Timberlake  
*Jauravia quadrinotata* Kapur  
[*Leis dimidiata* (F.)] = *Harmonia dimidiata* (F.)  
[*Leis*] = *Harmonia*  
[*Lemnia biplagiata* (Swartz)] = *Coelophora biplagiata* Swartz  
[*Leptothea galbula* (Mulsant)] = *Illeis galbula* (Mulsant)  
*Lindorus lophanthae* (Blaisdell) [*Rhyzobius lophantae* (Blaisdell), *Rhyzobius lorophantae* (Blaisdell)]  
*Liodalda flavomaculata* (De Geer) [*Adalia flavomaculata* (De Geer)]  
*Macrocheilus hauseri* (Mader) [*Halyzia hauseri* (Mader)]  
*Macronaemia hauseri* (Weise)

- Megalocaria dilatata* (F.) [*Anisolemnia dilatata* (F.)]  
 [*Menochilus quadriplagiatus* (Swartz)] = *Menochilus sexmaculatus* (F.)  
*Menochilus sexmaculatus* (F.) [*Cheilomenes sexmaculata* (F.), *Menochilus quadriplagiatus* (Swartz)]  
*Micraspis allardi* (Mulsant)  
*Micraspis discolor* (F.)  
*Microweisea* sp.  
*Mulsantina hudsonica* (Casey)  
*Mulsantina picta* (Randall)  
*Myrrha octodecimguttata* (L.) – eighteen spot ladybird  
*Myzia oblongoguttata* (L.) – striped ladybird  
*Myzia subvittata* (Mulsant)  
*Neda marginalis* Mulsant  
*Neocalvia anastomozans* Crotch  
 [*Nephaspis oculatus* (Blatchley)] = *Clitostethus oculatus* (Blatchley)  
*Nephus bilucernarius* (Mulsant)  
*Nephus bisignatus* (Bohemian)  
*Nephus flavifrons* (Melsheimer) [*Scymnus flavifrons* Melsheimer, North America, not *Scymnus flavifrons* Blackburn, Australia]  
*Nephus guttulatus* (LeConte) [*Scymnus guttulatus* LeConte]  
*Nephus includens* (Kirsch)  
*Nephus kiesenwetteri* (Mulsant) [*Scymnus kiesenwetteri* Mulsant]  
*Nephus ornatus* (LeConte) [*Scymnus ornatus* LeConte]  
*Nephus quadrimaculatus* (Herbst) [*Scymnus quadrimaculatus* (Herbst)]  
*Nephus redtenbacheri* (Mulsant)  
*Nephus soudanensis* (Sicard) [*Scymnus soudanensis* Sicard]  
*Oenopia billieti* (Mulsant)  
*Oenopia conglobata* (L.) [*Synharmonia conglobata* (L.)]  
*Oenopia kirbyi* Mulsant  
*Oenopia lyncea* (Olivier)  
*Oenopia sexareata* (Mulsant)  
 [*Olla abdominalis* (Say)] = *Olla v-nigrum* (Mulsant)  
*Olla v-nigrum* (Mulsant) – ash-gray lady beetle [*Olla abdominalis* (Say)]  
*Orcus australasiae* (Boisduval)  
 [*Orcus chalybeus* (Boisduval)] = *Halmus chalybeus* (Boisduval)  
*Palaeoneda auriculata* (Mulsant) [*Paleoneda miniata* (Hope)]  
 [*Paleoneda miniata* (Hope)] = *Palaeoneda auriculata* (Mulsant)  
*Pania luteopustulata* Mulsant  
*Paranaemia vittigera* (Mannerheim)
- Parastethorus nigripes* (Kapur) [*Stethorus loxtoni* Britton & Lee]  
*Parexochomus melanocephalus* (Zoubkoff) [*Exochomus melanocephalus* (Zoubkoff)]  
*Parexochomus nigromaculatus* (Goeze) [*Exochomus nigromaculatus* (Goeze)]  
*Parexochomus troberti* (Mulsant) [*Exochomus troberti* (Mulsant)]  
*Parexochomus troberti concavus* (Fuersch) [*Exochomus concavus* Fuersch]  
*Pentilia insidiosa* Mulsant  
*Pharoscymnus anchorago* (Fairmaire)  
*Pharoscymnus numidicus* (Pic)  
*Pharoscymnus ovoideus* Sicard  
*Phymatosternus lewisii* (Crotch)  
*Platynaspis luteorubra* (Goeze)  
*Priscibrumus lituratus* (Gorham) [*Exochomus lituratus* (Gorham)]  
*Priscibrumus uropygialis* (Mulsant)  
*Propylea dissecta* (Mulsant)  
*Propylea japonica* (Thunberg)  
*Propylea quatuordecimpunctata* (L.) – fourteen spot ladybird [*Propylaea quatuordecimpunctata* (L.)]  
*Pseudoazyia trinitatis* (Marshall) [*Azya trinitatis* (Marshall)]  
 [*Pseudoscydmus tsugae*] Sasaji & McClure] = *Sasajiscymnus tsugae* (Sasaji & McClure)  
*Pseudoscydmus kurohime* (Mityake) = *Sasajiscymnus kurohime* (Miyatake)  
*Psyllobora confluens* (F.)  
*Psyllobora vigintiduopunctata* (L.) – twenty two spot ladybird [*Thea vigintiduopunctata* (L.)]  
*Psyllobora vigintimaculata* (Say) – twenty-spotted lady beetle  
 [*Pullus auritus* (Thunberg)] = *Scymnus auritus* Thunberg  
*Pullus mediterraneus* (F.) = *Scymnus marinus* (Mulsant)  
*Pullus subvillosus* (Goeze) = *Scymnus subvillosus* (Goeze)  
*Rhyzobius litura* (F.)  
 [*Rhyzobius lophanthae* (Blaisdell)] = *Lindorus lophanthae* (Blaisdell)  
 [*Rhyzobius lorophanthae* (Blaisdell)] = *Lindorus lophanthae* (Blaisdell)  
*Rhyzobius ventralis* (Erichson) – black lady beetle  
*Rodatus major* (Blackburn)  
*Rodolia cardinalis* (Mulsant) – vedalia beetle  
*Rodolia fumida* Mulsant  
*Rodolia guerini* (Crotch)

- Rodolia iceryae* Janson  
*Rodolia occidentalis* Weise  
*Sasajiscymnus kurohime* (Miyatake) [*Pseudoscymnus kurohime* (Miyatake)]  
 [ *Sasajiscymnus ningshanensis* (Sasaji & McClure) ] =  
*Scymnus ningshanensis* Yu & Yao  
*Sasajiscymnus tsugae* (Sasaji & McClure)  
 [ *Pseudoscymnus tsugae* Sasaji & McClure ]  
[ *Scymnodes lividigaster* (Mulsant) ] = *Apolinus lividigaster* (Mulsant)  
*Scymnus abietis* (Paykull)  
[ *Scymnus aeneipennis* Sicard ] = *Zagloba aeneipennis* (Sicard)  
*Scymnus apetzi* Mulsant  
*Scymnus ater* Kugelann  
*Scymnus auritus* Thunberg [*Pullus auritus* (Thunberg)]  
*Scymnus coccivora* Ayyar  
*Scymnus creperus* Mulsant  
*Scymnus dorcatomoides* Weise  
*Scymnus flavifrons* Blackburn (Australia)  
[ *Scymnus flavifrons* Melsheimer ] (North America) = *Nephus flavifrons* (Melsheimer)  
*Scymnus frontalis* (F.)  
[ *Scymnus guttulatus* LeConte ] = *Nephus guttulatus* (LeConte)  
*Scymnus haemorrhoidalis* Herbst  
*Scymnus hilaris* Motschulsky  
*Scymnus hoffmanni* Weise  
*Scymnus impexus* Mulsant  
*Scymnus interruptus* (Goeze)  
[ *Scymnus kiesenwetteri* Mulsant ] = *Nephus kiesenwetteri* (Mulsant)  
*Scymnus lacustris* LeConte  
*Scymnus levallanti* Mulsant  
*Scymnus loewii* Mulsant – dusky lady beetle  
*Scymnus louisianae* Chapin  
*Scymnus marginicollis* Mannerheim  
*Scymnus marinus* (Mulsant) [*Scymnus mediterraneus* Iablokoff-Khnzorian, *Pullus mediterraneus* (Iablokoff-Khnzorian)]  
[ *Scymnus mediterraneus* ] = *Scymnus marinus* (Mulsant)  
*Scymnus moreletti* Mulsant  
*Scymnus nigrinus* Kugelann  
*Scymnus ningshanensis* Yu & Yao [*Sasajiscymnus ningshanensis* (Sasaji & McClure)]  
[ *Scymnus ornatus* LeConte ] = *Nephus ornatus* (LeConte)  
*Scymnus otohime* Kamiya  
*Scymnus posticalis* Sicard  
*Scymnus postpictus* Casey
- Scymnus pyrocheilus* Mulsant  
*Scymnus quadrillum* Motschulsky  
[ *Scymnus* (*Nephus*) *quadrimaculatus* (Herbst) ] = *Nephus quadrimaculatus* (Herbst)  
*Scymnus rubromaculatus* (Goeze)  
*Scymnus sinuanodus* Yu & Yao  
*Scymnus smithianus* Silvestri  
*Scymnus soudanensis* Sicard  
*Scymnus subvillosus* (Goeze) [*Pullus subvillosus* (Goeze)]  
*Scymnus suturalis* Thunberg  
*Scymnus syriacus* (Marsuel)  
*Scymnus tardus* Mulsant  
[ *Semiadalia undecimnotata* (Schneider) ] = *Ceratomegilla undecimnotata* (Schneider)  
*Serangium parcesetosum* Sicard  
[ *Sidis* ] = *Nephus*  
[ *Spiladelpha barovskii* kiritschenkoi Semenov-Tian-Shanski ] = *Ceratomegilla barovskii* kiritschenkoi (Semenov-Tian-Shanski)  
*Sospita vigintiguttata* (L.)  
*Stethorus bifidus* Kapur  
*Stethorus gilvifrons* (Mulsant)  
*Stethorus japonicus* Kamiya  
[ *Stethorus loxtoni* Britton & Lee ] = *Parastethorus nigripes* (Kapur)  
*Stethorus madecassus* Chazeau  
[ *Stethorus picipes* Casey ] = *Stethorus punctum picipes* Casey  
[ *Stethorus punctillum* Weise ] = *Stethorus pusillus* (Herbst)  
*Stethorus punctum* (LeConte) – spider-mite destroyer  
*Stethorus punctum picipes* Casey [*Stethorus picipes* Casey]  
*Stethorus pusillus* (Herbst) [*Stethorus punctillum* (Weise)]  
*Stethorus tridens* Gordon  
*Stethorus vegans* (Blackburn)  
*Subcoccinella vigintiquatuorpunctata* (L.) – twenty four spot ladybird  
[ *Synharmonia conglobata* (L.) ] = *Oenopia conglobata* (L.)  
*Synona obscura* Poorani, Ślipiński & Booth  
*Synonycha grandis* (Thunberg)  
*Thalassa saginata* Mulsant  
[ *Thea vigintiguopunctata* (L.) ] = *Psyllobora vigintiguopunctata* (L.)  
*Tytthaspis sedecimpunctata* (L.) – sixteen spot ladybird  
[ *Tytthaspis trilineata* (Weise) ] = *Coccinella nigrovittata* Kapur  
[ *Verania* ] = *Micraspis*

*Vibidia duodecimguttata* (Poda)  
*Zagloba aeneipennis* (Sicard) [*Scymnus aeneipennis*  
 Sicard]

## OTHER ORGANISMS

In alphabetical order within each taxon, and with Family given in italics within brackets after the authority

## OTHER INSECTS

### Dermaptera

*Anechura harmandi* (Burr) (*Forficulidae*) – hump earwig  
*Forficula auricularia* L. (*Forficulidae*) – European earwig

### Hemiptera: Heteroptera

*Calocoris norvegicus* (Gmelin) (*Miridae*) – strawberry bug  
*Campylomma verbasci* (Meyer) (*Miridae*) – mullein bug  
*Caternaultiella rugosa* (Schoutenden) (*Plataspidae*)  
*Eurygaster integriceps* Puton (*Pentatomidae*) – sunn pest or corn bug  
*Geocoris punctipes* (Say) (*Lygaeidae*) – big-eyed bug  
*Hyaliodes vitripennis* (Say) (*Miridae*)  
*Lygus hesperus* (Knight) (*Miridae*) – western plant bug  
*Lygus lineolaris* (Palidot de Beauvois) (*Miridae*) – tarnished plant bug  
*Lygus* (*Miridae*)  
*Nabis (Reduvius) americiferus* Carayon (*Nabidae*) – common damsel bug  
*Nysius huttoni* White (*Lygaeidae*) – wheat bug  
*Orius insidiosus* (Say) (*Anthocoridae*) – insidious flower bug  
*Podisus maculiventris* (Say) (*Pentatomidae*) – spined soldier bug  
*Pyrrhocoris apterus* (L.) (*Pyrrhocoridae*) – firebug  
*Sidnia kinbergi* (Stål) (*Miridae*)

### Hemiptera: Auchenorrhyncha

*Homalodisca vitripennis* (Germar) (*Cicadellidae*) – glassy-winged sharpshooter

*Nilaparvata lugens* (Stål) (*Delphacidae*) – brown planthopper  
*Philaenus spumarius* (L.) (*Cercopidae*) – common froghopper or meadow spittlebug

### Hemiptera: Sternorrhyncha: Aphidoidea

(Synonyms are in square brackets)

*Acyrthosiphon caraganae* (Cholodkovsky) (*Aphididae*)  
*Acyrthosiphon ignotum* Mordvilko (*Aphididae*)  
*Acyrthosiphon kondoi* Shinji (*Aphididae*) – blue alfalfa aphid  
 [Acyrthosiphon nipponicum (Essig & Kuwana)] = *Neoaulacorthum nipponicum* (Essig & Kuwana) (*Aphididae*)  
*Acyrthosiphon pisum* (Harris) (*Aphididae*) – pea aphid  
*Adelges cooleyi* (Gillette) (*Adelgidae*) – Cooley spruce gall adelgid  
*Adelges laricis* Vallot (*Adelgidae*) – larch adelgid  
*Adelges nordmannianae* (Eckstein) (*Adelgidae*)  
 [Adelges nusslini (Boerner)] = *Adelges nordmannianae* (Eckstein) (*Adelgidae*)  
*Adelges piceae* (Ratzeburg) (*Adelgidae*) – balsam woolly adelgid  
*Adelges tsugae* Annand (*Adelgidae*) – hemlock woolly adelgid  
*Aphis carduella* Walsh (*Aphididae*)  
 [Aphis cirsiiacanthoidis Boerner] = *Aphis fabae* cirsiiacanthoidis Scopoli (*Aphididae*)  
*Aphis craccivora* Koch (*Aphididae*) – cowpea aphid or groundnut aphid  
*Aphis cytisorum* Hartig (*Aphididae*)  
*Aphis fabae* Scopoli (*Aphididae*) – black bean aphid  
*Aphis fabae cirsiiacanthoidis* Scopoli (*Aphididae*)  
*Aphis farinosa* J.F.Gmelin (*Aphididae*)  
*Aphis glycines* Matsumura (*Aphididae*) – soybean aphid  
*Aphis gossypii* Glover (*Aphididae*) – cotton aphid or melon aphid  
*Aphis hederae* Kaltenbach (*Aphididae*)  
 [Aphis helianthi Monell] = *Aphis carduella* Walsh (*Aphididae*)  
*Aphis jacobaeae* Schrank (*Aphididae*)  
 [Aphis laburni Kaltenbach] = *Aphis cytisorum* cytisorum Hartig (*Aphididae*)  
*Aphis nerii* Boyer de Fonscolombe (*Aphididae*) – oleander aphid  
*Aphis pomi* De Geer (*Aphididae*) – green apple aphid  
*Aphis punicae* Passerini (*Aphididae*)

- Aphis sambuci* L. (*Aphididae*) – elder aphid  
*Aphis spiraecola* Patch (*Aphididae*) – spiraea aphid or green citrus aphid  
*Aphis spiraephaga* F.P. Müller (*Aphididae*)  
*Aphis spiraephila* Patch (*Aphididae*)  
*Aphis urticata* J.F. Gmelin (*Aphididae*)  
[*Aulacorthum magnoliae* (Essig & Kuwana)] = *Neoaulacorthum magnoliae* (Essig & Kuwana) (*Aphididae*)  
*Aulacorthum solani* (Kaltenbach) (*Aphididae*) – glasshouse potato aphid or foxglove aphid  
*Betulaphis brevipilosus* Boerner (*Aphididae*)  
*Betulaphis quadriflora* (Kaltenbach) (*Aphididae*)  
*Brachycaudus helichrysi* (Kaltenbach) (*Aphididae*) – leaf-curling plum aphid  
*Brachycaudus persicae* (Passerini) (*Aphididae*) – black peach aphid  
*Brachycaudus prunicola* (Kaltenbach) (*Aphididae*)  
*Brachycaudus tragopogonis* (Kaltenbach) (*Aphididae*)  
*Brevicoryne brassicae* (L.) (*Aphididae*) – cabbage aphid  
*Callipterella calliptera* (Hartig) (*Aphididae*)  
*Capitophorus elaeagni* (Del Guercio) (*Aphididae*)  
*Cavariella konoii* Takahashi (*Aphididae*)  
*Ceratovacuna lanigera* Zehntner (*Aphididae*) – sugar cane woolly aphid  
*Cervaphis quercus* Takahashi (*Aphididae*)  
*Chaetosiphon fragaefolii* (Cockerell) (*Aphididae*) – strawberry aphid  
*Chaitophorus capreae* (Mosley) (*Aphididae*)  
*Chaitophorus leucomelas* Koch (*Aphididae*)  
[*Chaitophorus versicolor* Koch] = *Chaitophorus leucomelas* Koch (*Aphididae*)  
*Chromaphis juglandicola* (Kaltenbach) (*Aphididae*) – walnut aphid  
*Cinara palaestinensis* Hille Ris Lambers (*Aphididae*)  
*Delphiniobium junackianum* (Karsch) (*Aphididae*)  
*Diuraphis noxia* (Kurdjumov) (*Aphididae*) – Russian wheat aphid  
*Drepanosiphum platanoidis* (Schrank) (*Aphididae*) – sycamore aphid  
*Dysaphis crataegi* (Kaltenbach) (*Aphididae*) – hawthorn–parsnip aphid  
*Dysaphis devecta* (Walker) (*Aphididae*) – rosy leaf-curling aphid  
*Dysaphis plantaginea* (Passerini) (*Aphididae*) – rosy apple aphid  
*Elatobium abietinum* (Walker) (*Aphididae*) – spruce aphid  
*Eriosoma lanigerum* (Hausmann) (*Aphididae*) – woolly apple aphid  
*Eucallipterus tiliae* (L.) (*Aphididae*) – lime aphid  
*Euceraphis betulae* (Koch) (*Aphididae*)  
*Euceraphis punctipennis* (Zetterstedt) (*Aphididae*)  
*Hyalopterus pruni* (Geoffroy) (*Aphididae*) – mealy plum aphid  
*Hyperomyzus carduellinus* (Theobald) (*Aphididae*)  
*Hyperomyzus lactucae* (L.) (*Aphididae*) – blackcurrant-sowthistle aphid  
*Laingia psammiae* Theobald (*Aphididae*)  
*Liosomaphis berberidis* (Kaltenbach) (*Aphididae*)  
*Lipaphis pseudobrassicae* (Davis) (*Aphididae*) – turnip aphid or mustard aphid  
[*Longiunguis donacis* (Passerini)] = *Melanaphis donacis* (Passerini) (*Aphididae*)  
*Macrosiphoniella artemisiae* (Boyer de Fonscolombe) (*Aphididae*)  
*Macrosiphoniella sanborni* (Gillette) (*Aphididae*) – chrysanthemum aphid  
*Macrosiphum albifrons* Essig (*Aphididae*) – lupin aphid  
*Macrosiphum euphorbiae* (Thomas) (*Aphididae*) – potato aphid  
[*Macrosiphum ibarae* (Matsumura)] = *Sitobion ibarae* (Matsumura) (*Aphididae*)  
*Macrosiphum rosae* (L.) (*Aphididae*) – rose aphid  
*Megoura viciae* Buckton (*Aphididae*) – vetch aphid  
*Melanaphis donacis* (Passerini) (*Aphididae*)  
*Metopolophium dirhodum* (Walker) (*Aphididae*) – rose-grain aphid.  
*Metopolophium festucae* (Theobald) (*Aphididae*) – fescue aphid  
*Microlophium carnosum* (Buckton) (*Aphididae*)  
*Microsiphoniella artemisiae* (Gillette) (*Aphididae*)  
*Mindarus abietinus* Koch (*Aphididae*)  
*Myzocallis boernerii* Stroyan (*Aphididae*)  
*Myzocallis carpini* (Koch) (*Aphididae*)  
*Myzocallis castanicola* Baker (*Aphididae*)  
*Myzocallis coryli* (Goetze) (*Aphididae*) – hazel aphid or filbert aphid  
*Myzus cerasi* (F.) (*Aphididae*) – cherry blackfly  
*Myzus persicae* (Sulzer) (*Aphididae*) – peach-potato aphid  
*Myzus persicae nicotianae* Blackman (*Aphididae*)  
*Neoaulacorthum magnoliae* (Essig & Kuwana) (*Aphididae*)  
*Neoaulacorthum nipponicum* (Essig & Kuwana) (*Aphididae*)  
*Neomyzus circumflexus* (Buckton) (*Aphididae*)  
*Neophyllaphis podocarpi* Takahashi (*Aphididae*)  
*Periphyllus californiensis* (Shinji) (*Aphididae*)  
*Periphyllus lyropictus* (Kessler) (*Aphididae*)

*Periphyllus testudinaceus* (Fernie) (*Aphididae*)  
*Phorodon humuli* (Schrank) (*Aphididae*) – damson–hop aphid  
*Phyllaphis fagi* (L.) (*Aphididae*)  
*Phylloxera glabra* (von Heyden) (*Phylloxeridae*)  
*Pineus pini* (Macquart) (*Adelgidae*)  
*Pseudoregma alexanderi* (Takahashi) (*Aphididae*)  
*Pseudoregma bambucicola* (Takahashi) (*Aphididae*)  
*Pterocallis alni* (deGeer) (*Aphididae*)  
*Rhopalosiphum maidis* (Fitch) (*Aphididae*) – corn leaf aphid  
*Rhopalosiphum padi* (L.) (*Aphididae*) – bird cherry–oat aphid  
*Schizaphis graminum* (Rondani) (*Aphididae*) – greenbug  
*Schizolachnus pineti* (F.) (*Aphididae*)  
*Schizolachnus piniradiatae* (Davidson) (*Aphididae*)  
*Sitobion akebiae* (Shinji) (*Aphididae*)  
*Sitobion avenae* (F.) (*Aphididae*) – grain aphid  
*Sitobion ibarae* (Matsumura) (*Aphididae*)  
*Symydobius oblongus* (von Heyden) (*Aphididae*)  
*Thelaxes dryophila* (Schrank) (*Aphididae*)  
[*Theroaphis maculata* (Buckton)] = *Theroaphis trifolii* (Monell) (*Aphididae*) – spotted alfalfa aphid  
*Theroaphis trifolii* (Monell) (*Aphididae*) – spotted alfalfa aphid  
*Toxoptera aurantii* (Boyer de Fonscolombe) (*Aphididae*) – black citrus aphid or tea aphid  
*Toxoptera citricidus* (Kirkaldy) (*Aphididae*) – brown citrus aphid  
[*Toxoptera graminum* (Rondani)] = *Schizaphis graminum* (Rondani) (*Aphididae*) – greenbug  
*Tuberculatus annulatus* (Hartig) (*Aphididae*)  
*Tuberolachnus salignus* (J.F. Gmelin) (*Aphididae*) – willow aphid  
*Uroleucon aeneum* (Hille Ris Lambers) (*Aphididae*)  
*Uroleucon ambrosiae* (Thomas) (*Aphididae*)  
*Uroleucon cichorii* (Koch) (*Aphididae*)  
*Uroleucon cirsii* (L.) (*Aphididae*)  
*Uroleucon compositae* (Theobald) (*Aphididae*)  
*Uroleucon formosanum* (Takahashi) (*Aphididae*)  
*Uroleucon jaceae* (L.) (*Aphididae*)  
*Uroleucon nigrotuberculatum* (Olive) (*Aphididae*)  
*Vesiculaphis caricis* (Fullaway) (*Aphididae*)

### Hemiptera: other Sternorrhyncha

*Abgrallaspis cyanophylli* (Signoret) (*Diaspididae*)  
*Acutaspis umbonifera* (Newstead) (*Diaspididae*)

*Agonoscena pistaciae* Burckhardt et Lauterer (*Psyllidae*)  
*Aleurodicus cocois* (Curtis) (*Aleyrodidae*)  
*Aleurodicus dispersus* Russell (*Aleyrodidae*) – spiralling whitefly  
*Aleurotuba jelinekii* (Frauenfeld) (*Aleyrodidae*)  
*Aleyrodes proletella* (L.) (*Aleyrodidae*)  
*Aonidiella aurantii* (Maskell) (*Diaspididae*) – California red scale  
*Aonidiella orientalis* (Newstead) (*Diaspididae*)  
*Aonidimytilus albus* (Cockerell) (*Diaspididae*) – cassava scale  
*Aspidiotus destructor* Signoret (*Diaspididae*) – coconut scale  
*Aspidiotus nerii* Bouché (*Diaspididae*)  
*Asterolecanium* sp. (*Asterolecaniidae*)  
*Aulacaspis tegalensis* (Zehntner) (*Diaspididae*) – sugar cane scale  
*Aulacaspis tubercularis* Newstead (*Diaspididae*)  
*Bemisia tabaci* (Gennadius) (*Aleyrodidae*)  
[*Bemisia argentifolii* Bellows] = *Bemisia tabaci* Gennadius (*Aleyrodidae*)  
*Chionaspis alnus* Kuwana (*Diaspididae*)  
*Chionaspis salicis* (L.) (*Diaspididae*)  
*Chrysomphalus aonidum* (L.) (*Diaspididae*) – Florida red scale  
*Chrysomphalus bifasciculatus* Ferris (*Diaspididae*)  
*Coccus hesperidum* L. (*Coccidae*) – soft brown scale  
*Coccus viridis* (Green) (*Coccidae*) – green coffee scale or soft green scale  
*Dactylopius opuntiae* (Cockerell) (*Dactylopiidae*)  
*Diaphorina citri* Kuwayama (*Psyllidae*) – Asian citrus psyllid  
*Diaspidiotus perniciosus* (Comstock) (*Diaspididae*)  
*Dysmicoccus* (*Pseudococcidae*)  
*Eriococcus coriaceus* Maskell (*Eriococcidae*)  
*Eulecanium caraganae* Borchsenius (*Coccidae*)  
*Ferrisia virgata* (Cockerell) (*Pseudococcidae*) – striped mealybug  
*Hemberlesia lataniae* (Signoret) (*Diaspididae*)  
*Heteropsylla cubana* Crawford (*Psyllidae*)  
*Icerya purchasi* Maskell (*Monophlebidae*) – cottony cushion scale  
*Lepidosaphes beckii* (Newman) (*Diaspididae*) – citrus mussel scale  
*Lepidosaphes cornutus* Ramakrishna Ayyar (*Diaspididae*)  
*Lepidosaphes ulmi* (L.) (*Diaspididae*) – mussel scale or oystershell scale  
*Maconellicoccus hirsutus* (Green) (*Pseudococcidae*) – pink hibiscus mealybug

*Matsucoccus feytaudi* Ducasse (*Matsucoccidae*)  
*Matsucoccus josephi* Bodenheimer & Harpaz  
   (*Matsucoccidae*)  
*Matsucoccus matsumurae* (Kuwana) (*Matsucoccidae*)  
*Melanaspis glomerata* (Green) (*Diaspididae*)  
*Monophlebulus pilosior* (Maskell) (*Monophlebidae*)  
*Orthezia urticae* (L.) (*Ortheziidae*)  
*Parlatoria blanchardi* (Tergioni Tozzetti) (*Diaspididae*)  
*Phenacoccus herreni* Cox & Williams (*Pseudococcidae*)  
*Phenacoccus madeirensis* Green (*Pseudococcidae*)  
*Phenacoccus manihoti* Matile-Ferrero (*Pseudococcidae*)  
   – cassava mealybug  
*Phoenicococcus marlatti* (Cockerell) (*Phoenicococcidae*)  
*Pinnaspis buxi* (Bouché) (*Diaspididae*)  
*Planococcus citri* (Risso) (*Pseudococcidae*) – citrus  
   mealybug  
*Planococcus minor* (Maskell) (*Pseudococcidae*)  
*Pseudochermes fraxini* (Kaltenbach) (*Eriococcidae*)  
*Pseudococcus cryptus* Hempel (*Pseudococcidae*)  
*Pseudococcus maritimus* (Ehrhorn) (*Pseudococcidae*)  
*Pseudococcus viburni* (Signoret) (*Pseudococcidae*)  
*Psylla alni* (L.) (*Psyllidae*)  
*Psylla jucunda* Tuthill (*Psyllidae*)  
*Psylla mali* (Schmidberger) (*Psyllidae*) – apple sucker  
*Psylla ulmi* Forster (*Psyllidae*)  
*Psylla uncatoides* (Ferris et Clyver) (*Psyllidae*)  
*Pulvinaria psidii* Maskell (*Coccidae*) – green shield scale  
*Pulvinaria urbicola* (Cockerell) (*Coccidae*)  
*Pulvinaria vitis* (L.) (*Coccidae*) – woolly vine scale  
*Rastrococcus invadens* Williams (*Pseudococcidae*)  
*Saissetia coffeae* (Walker) (*Coccidae*) – helmet scale  
*Saissetia oleae* (Olivier) (*Coccidae*) – olive scale  
*Siphoninus phillyreae* (Haliday) (*Aleyrodidae*)  
*Trialeurodes vaporariorum* (Westwood) (*Aleyrodidae*)  
   – glasshouse whitefly  
*Unaspis citri* (Comstock) (*Diaspididae*) – citrus snow scale  
*Unaspis euonymi* (Comstock) (*Diaspididae*)  
*Unaspis yanensis* Kuwana (*Diaspididae*)

## Thysanoptera

*Thrips tabaci* Lindeman (*Thripidae*) – onion thrips or  
   tobacco thrips

## Neuroptera

*Chrysopa* Leach (*Chrysopidae*)  
*Chrysopa oculata* Say (*Chrysopidae*) – golden-eyed  
   lacewing

*Chrysopa perla* (L.) (*Chrysopidae*)  
*Chrysopa sinica* (Tjeder) (*Chrysopidae*)  
*Chrysoperla carnea* (Stephens) (*Chrysopidae*) – green  
   lacewing  
*Chrysoperla plorabunda* Fitch (*Chrysopidae*)  
*Chrysoperla rufilabris* (Burmeister) (*Chrysopidae*)

## Lepidoptera

*Acraea encedon* (L.) (*Nymphalidae*)  
[*Anagasta kuehniella* (Zeller)] = *Ephestia kuehniella*  
   Zeller (*Pyralidae*)  
*Arctia* (*Arctiidae*) – tiger moths  
*Choristoneura pinus* Freeman (*Tortricidae*) – jack pine  
   budworm  
*Danaus plexippus* (L.) (*Nymphalidae*) – monarch  
   butterfly  
*Ephestia kuehniella* Zeller (*Pyralidae*) – Mediterranean  
   flour moth  
*Galleria mellonella* (L.) (*Pyralidae*) – wax moth  
*Helicoverpa armigera* (Hübner) (*Noctuidae*) – cotton  
   bollworm  
*Helicoverpa zea* (Boddie) (*Noctuidae*) – corn earworm  
*Heliothis virescens* (F.) (*Noctuidae*) – tobacco budworm  
*Hyphantria cunea* (Drury) (*Arctiidae*) – fall webworm  
*Ostrinia nubilalis* (Hübner) (*Crambidae*) – European  
   corn borer  
*Pectinophora gossypiella* (Saunders) (*Gelechiidae*) – pink  
   bollworm  
*Phthorimaea operculella* (Zeller) (*Gelechiidae*) – potato  
   tuber moth  
*Pieris rapae* (L.) (*Pieridae*) – small cabbage white  
   butterfly  
*Plutella xylostella* (L.) (*Plutellidae*) – diamondback  
   moth  
*Sitotroga cerealella* (Olivier) (*Gelechiidae*) – Angoumois  
   grain moth  
*Spodoptera litura* (F.) (*Noctuidae*) – common  
   cutworm  
*Trichoplusia ni* (Hübner) (*Noctuidae*) – cabbage looper  
   or ni moth  
*Tyria* (*Arctiidae*) – cinnabar moths

## Diptera

*Anopheles quadrimaculatus* Say (*Culicidae*)  
*Aphidoletes aphidimyza* (Rondani) (*Cecidomyiidae*)  
*Boettcheria latisterna* Parker (*Sarcophagidae*)

*Chetogena claripennis* (Macquart) (Tachinidae)  
*Chrysotachina slossonae* (Coquillett) (Tachinidae)  
*Contarinia nasturtii* (Kieffer) (Cecidomyiidae) – swede midge  
*Cryptochaetum iceryae* (Williston) (Cryptochaetidae)  
*Culex quinquefasciatus* Say (Culicidae) – southern house mosquito  
[*Degeeria luctuosa* (Meigen)] = *Medina luctuosa* (Meigen) (Tachinidae)  
[*Doryphorophaga doryphorae*] = *Myiopharus doryphorae* (Riley) (Tachinidae)  
*Drosophila melanogaster* Meigen (Drosophilidae)  
*Episyrphus balteatus* (DeGeer) (Syrphidae) – marmalade hover fly  
*Euthelyconychia epilachnae* (Aldrich) (Tachinidae)  
[*Exoristoides slossonae* Coquillett] = *Chrysotachina slossonae* (Coquillett) (Tachinidae)  
*Helicobia rapax* (Walker) (Sarcophagidae)  
*Lydinolydella metallica* Townsend (Tachinidae)  
[*Lypha slossonae* Coquillett] = *Chrysotachina slossonae* (Coquillett) (Tachinidae)  
*Medina collaris* (Fallén) (Tachinidae)  
*Medina funebris* (Meigen) (Tachinidae)  
*Medina luctuosa* (Meigen) (Tachinidae)  
*Medina melania* (Meigen) (Tachinidae)  
*Medina separata* (Meigen) (Tachinidae)  
*Megaselia* (Phoridae)  
*Myiopharus doryphorae* (Riley) (Tachinidae)  
[*Paradexodes epilachnae* Aldrich] = *Euthelyconychia epilachnae* (Aldrich) (Tachinidae)  
*Phalacrotophora Enderlein* (Phoridae)  
*Phalacrotophora berolinensis* Schmitz (Phoridae)  
*Phalacrotophora beuki* Disney (Phoridae)  
*Phalacrotophora decimaculata* Liu (Phoridae)  
*Phalacrotophora delageae* Disney (Phoridae)  
*Phalacrotophora fasciata* (Fallén) (Phoridae)  
*Phalacrotophora indiana* Colyer (Phoridae)  
*Phalacrotophora nedae* (Malloch) (Phoridae)  
*Phalacrotophora philaxyridis* Disney (Phoridae)  
*Phalacrotophora quadrimaculata* Schmitz (Phoridae)  
*Policheta unicolor* (Fallén) (Tachinidae)  
*Pseudebenia epilachnae* Shima & Han (Tachinidae)  
*Ravinia errabunda* (Wulp) (Sarcophagidae)  
[*Sarcophaga helicis* Townsend] = *Helicobia rapax* (Walker) (Sarcophagidae)  
[*Sarcophaga latisterna* (Parker)] = *Boettcheria latisterna* Parker (Sarcophagidae)  
[*Sarcophaga reinhardii* Hall] = *Ravinia errabunda* (Wulp) (Sarcophagidae)  
*Strongygaster triangulifera* (Loew) (Tachinidae)

## Hymenoptera

[*Aminellus Masi*] = *Cowperia* Girault (Encyrtidae)  
[*Aminellus sumatraensis* Kerrich] = *Cowperia sumatraensis* (Kerrich) (Encyrtidae)  
*Anagyrus* Howard (Encyrtidae)  
*Anagyrus australiensis* (Howard) (Encyrtidae)  
*Anagyrus kamali* Moursi (Encyrtidae)  
*Anagyrus lopezi* (De Santis) (Encyrtidae)  
*Anagyrus pseudococcii* (Girault) (Encyrtidae)  
*Anastatus* Motschulsky (Eupelmidae)  
[*Anisotylus Timberlake*] = *Homalotylus* Mayr (Encyrtidae)  
*Aphanogmus* Thomson (Ceraphronidae)  
*Aphidius colemani* Viereck (Braconidae)  
*Aphidius eadyi* Starý (Braconidae)  
*Aphidius ervi* Haliday (Braconidae)  
*Apis mellifera* L. (Apidae) – European honey bee  
*Aprostocetus* Westwood (Eulophidae)  
*Aprostocetus esurus* (Riley) (Eulophidae)  
*Aprostocetus neglectus* (Domenichini) (Eulophidae)  
[*Atritomellus Kieffer*] = *Dendrocerus* (Megaspilidae)  
*Astroterobia* Girault (Pteromalidae)  
*Aximopsis* Ashmead (Eurytomidae)  
*Azteca instabilis* (Smith) (Formicidae)  
*Baryscapus* Foerster (Eulophidae)  
*Baryscapus thanasimi* (Ashmead) (Eulophidae)  
*Brachymeria carinatifrons* Gahan (Chalcididae)  
*Catolaccus* Thomson (Pteromalidae)  
*Centistes scymni* Ferrière (Braconidae)  
*Centistes subsulcatus* (Thomson) (Braconidae)  
*Centistina nipponicus* (Belokobylskij) (Braconidae)  
*Cerchysiella* Girault [*Zeteticontus Silvestri*] (Encyrtidae)  
*Chartocerus subaeneus* (Foerster) (Signiphoridae)  
*Cheiloneurus carinatus* Compere (Encyrtidae)  
*Cheiloneurus cyanonotus* Waterston (Encyrtidae)  
*Cheiloneurus liorhipnusi* (Risbec) (Encyrtidae)  
*Cheiloneurus orbitalis* Compere (Encyrtidae)  
*Chrysocharis johnsoni* Subba Rao (Eulophidae)  
*Chrysonotomyia appannai* (Chandy Kurian) (Eulophidae)  
*Coccidoctonus trinidadensis* Crawford (Encyrtidae)  
*Conura* Spinola (Chalcididae)  
*Conura paranensis* (Schrottky) (Chalcididae)  
*Conura petioliventris* (Cameron) (Chalcididae)  
*Conura porteri* (Brèthes) (Chalcididae)  
*Cowperia* Girault (Encyrtidae)  
*Cowperia areolata* (Walker) (Encyrtidae)  
*Cowperia indica* (Kerrich) (Encyrtidae)  
*Cowperia punctata* Girault (Encyrtidae)

*Cowperia subnigra* Li (Encyrtidae)  
*Cowperia sumatraensis* (Kerrich) (Encyrtidae)  
*Crematogaster* Lund (Formicidae)  
*Crematogaster lineolata* (Say) (Formicidae)  
*Dendrocerus* (Megaspilidae)  
*Dendrocerus ergensis* (Guesquiere) (Megaspilidae)  
*Dibrachys cavus* (Walker) (Pteromalidae)  
*Dinocampus* Foerster (Braconidae)  
*Dinocampus coccinellae* (Schrank) (Braconidae)  
[*Dinocampus nipponicus* Belokobylskij] = *Centistina nipponicus* (Belokobylskij) (Braconidae)  
[*Dinocampus terminatus* (Nees)] = *Dinocampus coccinellae* (Schrank) (Braconidae)  
*Dolichoderus bident* L. (Formicidae)  
[*Echthroplectis* Foerster] = *Homalotylus* Mayr (Encyrtidae)  
*Elasmus* Westwood (Eulophidae)  
*Encarsia sophia* (Girault & Dodd) (Aphelinidae)  
[*Epidinocarsus lopezi* (De Santis)] = *Anagyrus lopezi* (De Santis) (Encyrtidae)  
*Eretmocerus mundus* Mercet (Aphelinidae)  
*Eupelmus* Dalman (Eupelmidae)  
*Eupelmus urozonus* Dalman (Eupelmidae)  
*Eupelmus vermai* (Bhatnagar) (Eupelmidae)  
[*Eupteronotus* Kurdjumov] = *Trichomalopsis* Crawford (Pteromalidae)  
*Formica* L. (Formicidae)  
*Formica obscuripes* Forel (Formicidae)  
*Formica polyctena* Foerster (Formicidae) – European red wood ant  
*Formica rufa* L. (Formicidae) – wood ant  
*Formica rufibarbis* F. (Formicidae) – red-barbed ant  
*Gelis* Thunberg (Ichneumonidae)  
*Gelis agilis* (F.) (Ichneumonidae)  
[*Gelis instabilis* (Förster)] = *Gelis agilis* (F.) (Ichneumonidae)  
*Gelis melanocephalus* (Schrank) (Ichneumonidae)  
[*Hemaenasioidea* Girault] = *Homalotylus* Mayr (Encyrtidae)  
*Homalotyloloidea dahlbomii* (Westwood) (Encyrtidae)  
*Homalotylus* Mayr (Encyrtidae)  
*Homalotylus affinis* Timberlake (Encyrtidae)  
*Homalotylus africanus* Timberlake (Encyrtidae)  
*Homalotylus agarwali* Anis & Hayat (Encyrtidae)  
*Homalotylus albiclavatus* (Agarwal) (Encyrtidae)  
*Homalotylus albifrons* (Ishii) (Encyrtidae)  
*Homalotylus albatarsus* Gahan (Encyrtidae)  
*Homalotylus aligarhensis* (Shafee & Rizvi) (Encyrtidae)  
*Homalotylus balchanensis* Myartseva (Encyrtidae)  
*Homalotylus brevicauda* Timberlake (Encyrtidae)

[*Homalotylus californicus* (Say)] = *Homalotylus terminalis* (Say) (Encyrtidae)  
*Homalotylus cockerelli* Timberlake (Encyrtidae)  
*Homalotylus ephippium* (Ruschka) (Encyrtidae)  
*Homalotylus eytellei* (Ratzeburg) (Encyrtidae)  
*Homalotylus ferrierei* Hayat, Alam & Agarwal (Encyrtidae)  
*Homalotylus flaminius* Dalman (Encyrtidae)  
*Homalotylus hemipterinus* (De Stefani) (Encyrtidae)  
*Homalotylus formosus* Anis & Hayat (Encyrtidae)  
*Homalotylus himalayensis* Liao (Encyrtidae)  
*Homalotylus hybridus* Hoffer (Encyrtidae)  
*Homalotylus hyperaspicola* Tachikawa (Encyrtidae)  
*Homalotylus hyperaspidis* Timberlake (Encyrtidae)  
*Homalotylus hypnos* Noyes  
*Homalotylus indicus* (Agarwal) (Encyrtidae)  
*Homalotylus latipes* Girault (Encyrtidae)  
*Homalotylus longicaudus* Xu & He (Encyrtidae)  
*Homalotylus longipedicellus* (Shafee & Fatma) (Encyrtidae)  
*Homalotylus mexicanus* Timberlake (Encyrtidae)  
*Homalotylus mirabilis* (Brèthes) (Encyrtidae)  
*Homalotylus mundus* Gahan (Encyrtidae)  
*Homalotylus nigricornis* Mercet (Encyrtidae)  
*Homalotylus oculatus* (Girault) (Encyrtidae)  
*Homalotylus pallentipes* (Timberlake) (Encyrtidae)  
*Homalotylus platynaspidis* Hoffer (Encyrtidae)  
*Homalotylus punctifrons* Timberlake (Encyrtidae)  
*Homalotylus quaylei* Timberlake (Encyrtidae)  
*Homalotylus rubricatus* Sharkov (Encyrtidae)  
*Homalotylus scutellaris* Tan & Zhao (Encyrtidae)  
*Homalotylus scymnivorus* Tachikawa (Encyrtidae)  
*Homalotylus shuvakhinae* Trjapitzin & Triapitsyn (Encyrtidae)  
*Homalotylus similis* Ashmead (Encyrtidae)  
*Homalotylus sinensis* Xu & He (Encyrtidae)  
*Homalotylus singularis* Hoffer (Encyrtidae)  
*Homalotylus terminalis* (Say) (Encyrtidae)  
*Homalotylus trisubalbus* Xu & He (Encyrtidae)  
*Homalotylus turkmenicus* Myartseva (Encyrtidae)  
*Homalotylus vicinus* Silvestri (Encyrtidae)  
*Homalotylus yunnanensis* Tan & Zhao (Encyrtidae)  
*Homalotylus zhaoi* Xu & He (Encyrtidae)  
*Inkaka quadridentata* Girault (Pteromalidae)  
*Iridomyrmex* Mayr (Formicidae)  
*Isodromus niger* Ashmead (Encyrtidae)  
*Lasius claviger* (Roger) (Formicidae)  
*Lasius japonicus* Santschi (Formicidae) – Japanese ant  
*Lasius niger* (L.) (Formicidae) – black ant or garden ant  
*Lasius umbratus* (Nylander) (Formicidae)

[*Lepidaphytes* Blanchard] = *Homalotylus* Mayr  
   (*Encyrtidae*)  
*Linepithema humile* (Mayr) (*Formicidae*) – Argentine ant  
[i]*Lygocerus* Foerster] = *Dendrocerus* (*Megaspilidae*)  
*Lysiphlebus fabarum* (Marshall) (*Braconidae*)  
*Lysiphlebus testaceipes* (Cresson) (*Braconidae*)  
[*Mendozaniella* Brethes] = *Homalotylus* Mayr  
   (*Encyrtidae*)  
*Merismoclea rojasii* De Santis (*Pteromalidae*)  
*Mesopolobus* Westwood (*Pteromalidae*)  
*Mesopolobus secundus* (Crawford) (*Pteromalidae*)  
[*Mestocharis lividus* Girault] = *Pediobius foveolatus*  
   (Crawford) (*Eulophidae*)  
*Metastenus* Walker (*Pteromalidae*)  
*Metastenus caliginosus* Szelényi (*Pteromalidae*)  
*Metastenus concinnus* Walker (*Pteromalidae*)  
*Metastenus indicus* Sureshan & Narendran  
   (*Pteromalidae*)  
*Metastenus sulcatus* (Dodd) (*Pteromalidae*)  
*Metastenus townsendi* (Ashmead) (*Pteromalidae*)  
*Microctonus Wesmael* (*Braconidae*)  
*Monomorium minimum* (Buckley) (*Formicidae*) – little black ant  
*Myrmica ruginodis* Nylander (*Formicidae*)  
*Myrmica rugulosa* Nylander (*Formicidae*)  
[*Neoaenasioidea* Agarwal] = *Homalotylus* Mayr  
   (*Encyrtidae*)  
[*Neotainania* Husain & Agarwal] = *Uga* Girault  
   (*Chalcididae*)  
[*Nobrimus* Thomson] = *Homalotylus* Mayr (*Encyrtidae*)  
*Nothoserphus* Brues (*Proctotrupidae*)  
*Nothoserphus admirabilis* Lin (*Proctotrupidae*)  
*Nothoserphus aequalis* Townes (*Proctotrupidae*)  
*Nothoserphus afissae* (Watanabe) (*Proctotrupidae*)  
*Nothoserphus boops* (Thomson) (*Proctotrupidae*)  
*Nothoserphus debilis* Townes (*Proctotrupidae*)  
*Nothoserphus pilachnae* (Pschorn-Walcher)  
   (*Proctotrupidae*)  
*Nothoserphus fuscipes* Lin (*Proctotrupidae*)  
*Nothoserphus mirabilis* Brues (*Proctotrupidae*)  
*Nothoserphus partitus* Lin (*Proctotrupidae*)  
*Nothoserphus scymni* (Ashmead) (*Proctotrupidae*)  
*Nothoserphus townesi* Lin (*Proctotrupidae*)  
*Omphale* Haliday (*Eulophidae*)  
[*Omphale* *epilachni* Singh & Khan] = *Chrysonotomyia appannai* (Chandy Kurian) (*Eulophidae*)  
*Ooencyrtus azul* Prinsloo (*Encyrtidae*)  
*Ooencyrtus bedfordi* Prinsloo (*Encyrtidae*)  
*Ooencyrtus camerounensis* (Risbec) (*Encyrtidae*)  
*Ooencyrtus distatus* Prinsloo (*Encyrtidae*)

*Ooencyrtus pilachnae* Annecke (*Encyrtidae*)  
[*Ooencyrtus epulus* Annecke] = *Ooencyrtus camerounensis* (Risbec) (*Encyrtidae*)  
*Ooencyrtus polyphagus* (Risbec) (*Encyrtidae*)  
*Ooencyrtus puparum* Prinsloo (*Encyrtidae*)  
*Ooencyrtus sinis* Prinsloo (*Encyrtidae*)  
*Oomyzus Rondani* (*Eulophidae*)  
*Oomyzus mashhoodi* (Khan & Shafee) (*Eulophidae*)  
*Oomyzus scaposus* (Thomson) (*Eulophidae*)  
*Oomyzus sempronius* (Erdoes) (*Eulophidae*)  
*Ophelosia bifasciata* Girault (*Pteromalidae*)  
*Oricoruna Boucek* (*Pteromalidae*)  
*Ophelosia crawfordi* Riley (*Pteromalidae*)  
*Oricoruna orientalis* (Crawford) (*Pteromalidae*)  
*Pachyneuron* Walker (*Pteromalidae*)  
*Pachyneuron albutilus* Walker (*Pteromalidae*)  
*Pachyneuron altiscuta* Howard (*Pteromalidae*)  
*Pachyneuron chilocori* Domenichini (*Pteromalidae*)  
[*Pachyneuron concolor* Förster] = *Pachyneuron muscarum* (L.) (*Pteromalidae*)  
*Pachyneuron muscarum* (L.) (*Pteromalidae*)  
[*Pachyneuron siculum* Delucchi] = *Pachyneuron muscarum* (L.) (*Pteromalidae*)  
*Pachyneuron solitarium* (Hartig) (*Pteromalidae*)  
[*Pachyneuron syrphi* (Ashmead)] = *Pachyneuron albutilus* Walker (*Pteromalidae*)  
*Parachrysocharis* Girault (*Eulophidae*)  
*Paratrechina* Motschulsky (*Formicidae*)  
*Pediobius* Walker (*Eulophidae*)  
*Pediobius amaurocoelus* (Waterston) (*Eulophidae*)  
[*Pediobius* *epilachnae* (Rohwer)] = *Pediobius foveolatus* (Crawford) (*Eulophidae*)  
*Pediobius* *foveolatus* (Crawford) (*Eulophidae*)  
[*Pediobius* *mediopunctata* (Waterston)] = *Pediobius foveolatus* (Crawford) (*Eulophidae*)  
*Pediobius nishidai* Hansson (*Eulophidae*)  
[*Pediobius simiolus* (Takahashi)] = *Pediobius foveolatus* (Crawford) (*Eulophidae*)  
[*Perilitus americanus* Riley] = *Dinocampus coccinellae* (Schrank) (*Braconidae*)  
[*Perilitus coccinellae* (Schrank)] = *Dinocampus coccinellae* (Schrank) (*Braconidae*)  
*Perilitus rutilus* (Nees) (*Braconidae*)  
*Perilitus stuardoi* Porter (*Braconidae*)  
[*Perilitus terminatus* (Nees)] = *Dinocampus coccinellae* (Schrank) (*Braconidae*)  
*Pheidole megacephala* (F.) (*Formicidae*) – big-headed ant  
*Phygadeuon subfuscus* Cresson (*Ichneumonidae*)  
[*Pleurotropis* Foerster] = *Pediobius* Walker (*Eulophidae*)  
*Pnigalio agraules* (Walker) (*Eulophidae*)

*Praon volucre* (Haliday) (Braconidae)  
*Pristomyrmex pungens* Mayr (Formicidae)  
*Prochiloneurus aegyptiacus* (Mercet) (Encyrtidae)  
*Prochiloneurus nigriflagellum* (Girault) (Encyrtidae)  
*Pseudocatolaccus* Masi (Pteromalidae)  
*Quadrastichus ovulorum* (Ferrière) (Eulophidae)  
[*Scymnophagus* Ashmead] = *Metastenus* Walker  
(Pteromalidae)  
[*Scymnophagus mesnili* Ferrière] = *Metastenus concinnus* Walker (Pteromalidae)  
*Sigmoepilachna indica* Khan, Agnihotri & Sushil  
(Eulophidae)  
*Solenopsis invicta* Buren (Formicidae) – red imported fire ant  
[*Syntomosphyrus taprobanes* Waterston] = *Oomyzus scapulos* (Thomson) (Eulophidae)  
*Syrphoctonus tarsatorius* (Panzer) (Ichneumonidae)  
*Tamarixia radiata* (Waterston) (Eulophidae)  
*Tapinoma nigerrimum* (Nylander) (Formicidae)  
*Tetramorium caespitum* (L.) (Formicidae)  
*Tetrastichus Haliday* (Eulophidae)  
[*Tetrastichus coccinellae* Kurdjumov] = *Oomyzus scapulos* (Thomson) (Eulophidae)  
*Tetrastichus cydoniae* Risbec (Eulophidae)  
*Tetrastichus decrescens* Graham (Eulophidae)  
*Tetrastichus epilachnae* (Giard) (Eulophidae)  
[*Tetrastichus melanis* Burks] = *Oomyzus scapulos* (Thomson) (Eulophidae)  
*Tetrastichus orissaensis* Husain & Khan (Eulophidae)  
[*Tetrastichus sexmaculatus* Chandy Kurian] = *Oomyzus scapulos* (Thomson) (Eulophidae)  
[*Thomsonina Hellen*] = *Nothoserphus* Brues  
(Proctotrupidae)  
*Trichogramma* Westwood (Trichogrammatidae)  
*Trichogramma evanescens* Westwood  
(Trichogrammatidae)  
*Trichomalopsis* Crawford (Pteromalidae)  
*Trichomalopsis acuminata* (Graham) (Pteromalidae)  
*Trichomalopsis dubia* (Ashmead) (Pteromalidae)  
[*Tripolycystus* Dodd] = *Metastenus* Walker  
(Pteromalidae)  
*Uga* Girault [*Neotainania* Husain & Agarwal]  
(Chalcididae)  
*Uga colliscutellum* (Girault) (Chalcididae)  
*Uga coriacea* Kerrich (Chalcididae)  
*Uga digitata* Qian & He (Chalcididae)  
*Uga hemicarinata* Qian & Li (Chalcididae)  
*Uga javanica* Kerrich (Chalcididae)  
*Uga menoni* Kerrich (Chalcididae)  
*Uga sinensis* Kerrich (Chalcididae)  
*Wasmannia auropunctata* (Roger) (Formicidae)

[*Watanabeia* Masner] = *Nothoserphus* Brues  
(Proctotrupidae)

### Coleoptera (other than Coccinellidae)

*Agelastica coerulea* (Baly) (Chrysomelidae) – alder leaf beetle  
*Agonum dorsale* (Pontoppidan) (Carabidae)  
*Chrysomela populi* L. (Chrysomelidae) – poplar leaf beetle  
*Chrysophtarta bimaculata* (Olivier) (Chrysomelidae) – eucalyptus leaf beetle  
*Diabrotica virgifera* LeConte (Chrysomelidae) – western corn rootworm  
*Dicranolaius bellulus* (Guérin-Méneville) (Melyridae)  
*Dytiscus* sp. (Dytiscidae)  
*Galeruca interrupta arminiana* Weise (Chrysomelidae)  
*Galerucella lineola* (F.) (Chrysomelidae)  
*Galerucella pusilla* Duftschmid (Chrysomelidae) – golden loosestrife beetle  
*Galerucella sagittariae* Gyllenhal (Chrysomelidae) – cloudberry beetle  
*Harpalus pensylvanicus* (De Geer) (Carabidae)  
*Hypera postica* (Gyllenhal) (Curculionidae) – alfalfa weevil  
*Laricobius nigrinus* Fender (Derodontidae)  
*Leptinotarsa decemlineata* (Say) (Chrysomelidae) – Colorado potato beetle  
[*Melasoma populi* (L.)] = *Chrysomela populi* L. (Chrysomelidae)  
*Melolontha* sp. (Scarabaeidae)  
*Plagiodesma versicolora* (Laicharting) (Chrysomelidae)  
*Platynus dorsalis* (Pontopiddan) (Carabidae)  
*Pterostichus melanarius* (Illiger) (Carabidae)  
[*Pyrhalta luteola* (Muller)] = *Xanthogaleruca luteola* (Muller) (Chrysomelidae)  
*Sitona discoideus* Gyllenhal (Curculionidae)  
*Stenotarsus rotundus* Arrow (Endomychidae)  
*Tachyporus* sp. (Staphylinidae)  
*Xanthogaleruca luteola* (Muller) (Chrysomelidae) – elm leaf beetle

### PARASITIC MITES

*Coccipolipus* Husband (Podapolipidae)  
*Coccipolipus africanae* Husband (Podapolipidae)  
*Coccipolipus arturi* Haitlinger (Podapolipidae)  
*Coccipolipus benoiti* Husband (Podapolipidae)  
*Coccipolipus bifasciatae* Husband (Podapolipidae)

*Coccipolipus cacti* Husband (*Podapolipidae*)  
*Coccipolipus camerouni* Husband (*Podapolipidae*)  
*Coccipolipus chilocori* Husband (*Podapolipidae*)  
*Coccipolipus cooremani* Husband (*Podapolipidae*)  
*Coccipolipus epilachnae* Smiley (*Podapolipidae*)  
*Coccipolipus hippodamiae* (McDaniel & Morrill)  
     (*Podapolipidae*)  
*Coccipolipus macfarlanei* Husband (*Podapolipidae*)  
*Coccipolipus micraspisi* Husband (*Podapolipidae*)  
*Coccipolipus oconnori* Husband (*Podapolipidae*)  
*Coccipolipus solanophilae* (Cooreman) (*Podapolipidae*)  
*Hemisarcoptes lignieres* (*Hemisarcoptidae*)  
*Hemisarcoptes cooremani* Thomas (*Hemisarcoptidae*)  
*Leptus ignotus* (Oudemans) (*Erythraeidae*)

### **ARACHNIDA (other than parasitic mites)**

*Agistemus longisetus* Gonzalez (*Stigmaeidae*)  
*Amblyseius andersoni* (Chant) (*Phytoseiidae*)  
*Amblyseius fallacis* (Garman) (*Phytoseiidae*)  
*Amphitetranychus viennensis* (Zacher) (*Tetranychidae*)  
     – hawthorn spider mite or fruit tree spider mite  
*Araneus diadematus* (Clerck) (*Araneidae*)  
*Bryobia praetiosa* Koch (*Tetranychidae*) – clover mite  
*Bryobia rubrioculus* (Scheutten) (*Tetranychidae*)  
     – brown mite or brown apple mite  
*Cheiracanthium* (*Miturgidae*)  
*Clubiona reclusa* O.P.-Cambridge (*Clubionidae*)  
[*Metatetranychus ulmi* (Koch)] = *Panonychus ulmi* (Koch) (*Tetranychidae*)  
*Misumenops tricuspidatus* (F.) (*Thomisidae*)  
*Panonychus mori* Yokovama (*Tetranychidae*)  
*Panonychus ulmi* (Koch) (*Tetranychidae*) – red spider mite or European red mite  
*Tetranychus evansi* Baker & Pritchard (*Tetranychidae*)  
*Tetranychus lintearius* Dufour (*Tetranychidae*) – gorse spider mite  
*Tetranychus mcdanieli* McGregor (*Tetranychidae*)  
[*Tetranychus telarius* (L.)] = *Tetranychus urticae* Koch (*Tetranychidae*)  
*Tetranychus urticae* Koch (*Tetranychidae*) – two-spotted spider mite  
*Thanatus* Koch (*Philodromidae*)

### **NEMATODES**

[*Coccinellimermis Rubtzov*] = *Hexameris Steiner* (*Mermithidae*)

*Hexameris* Steiner (*Mermithidae*)  
*Howardula Cobb* (*Allantonematidae*)  
*Mermis* Dujardin (*Mermithidae*)  
*Parasitlenchus coccinellinae* Iberti & van Waerebeke  
     (*Allantonematidae*)

### **MOLLUSCA**

*Arion ater* (L.) (*Arionidae*) – European black slug  
*Arion hortensis* (Férussac) (*Arionidae*) – garden slug  
*Deroceras reticulatum* (Müller) (*Limacidae*) – grey field slug or grey garden slug  
*Tandonia budapestensis* (Hazay) (*Milacidae*) – Budapest slug

### **AVES**

*Acrocephalus schoenobaenus* (L.) (*Sylviidae*) – sedge warbler  
*Alauda arvensis* L. (*Alaudidae*) – sky lark  
*Anthus campestris* (L.) (*Motacillidae*) – tawny pipit  
*Anthus pratensis* (L.) (*Motacillidae*) – meadow pipit  
*Anthus trivialis* (L.) (*Motacillidae*) – tree pipit  
*Apus apus* (L.) (*Apodidae*) – common swift  
*Calandrella cinerea* (Gmelin) (*Alaudidae*) – red-capped lark  
*Coturnix coturnix* L. (*Phasianidae*) – quail  
*Coturnix japonicus* Temminck & Schlegel (*Phasianidae*) – Japanese quail  
*Cuculus canorus* L. (*Cuculidae*) – common cuckoo  
*Cyanistes caeruleus* (L.) (*Paridae*) – blue tit  
*Delichon urbica* (L.) (*Hirundinidae*) – house martin  
*Dendrocopos medius* (L.) (*Picidae*) – middle spotted woodpecker  
*Ficedula parva* (Bechstein) (*Muscicapidae*) – red-headed flycatcher  
*Hippolais icterina* (Vieillot) (*Sylviidae*) – icterine warbler  
*Hirundo rustica* L. (*Hirundinidae*) – barn swallow  
*Luscinia luscinia* (L.) (*Muscicapidae*) – thrush nightingale  
*Luscinia svecica* (L.) (*Muscicapidae*) – bluethroat  
*Melanocorypha calandra* (L.) (*Alaudidae*) – Calandra lark  
*Motacilla flava* L. (*Motacillidae*) – blue-headed wagtail  
*Muscicapa striata* (Pallas) (*Muscicapidae*) – spotted flycatcher  
*Oenanthe oenanthe* (L.) (*Muscicapidae*) – northern wheatear

*Parus major* L. (Paridae)  
*Passer domesticus* (L.) (Passeridae) – house sparrow  
*Passer montanus* (L.) (Passeridae) – tree sparrow  
*Phoenicurus ochruros* (Gmelin) (Muscicapidae) – black redstart  
*Phoenicurus phoenicurus* (L.) (Muscicapidae) – common redstart  
*Phylloscopus collybita* (Vieillot) (Phylloscopidae) – common chiffchaff  
*Phylloscopus trochilus* (L.) (Phylloscopidae) – willow warbler  
*Saxicola rubicola* (L.) (Muscicapidae) – common stonechat  
*Sitta europaea* L. (Sittidae) – wood nuthatch  
*Sturnus vulgaris* L. (Sturnidae) – common starling  
*Sylvia atricapilla* (L.) (Sylviidae) – blackcap  
*Sylvia borin* (Boddaert) (Sylviidae) – garden warbler  
*Sylvia communis* Latham (Sylviidae) – common whitethroat  
*Sylvia curruca* (L.) (Sylviidae) – lesser whitethroat  
*Sylvia nisoria* (Bechstein) (Sylviidae) – barred warbler  
*Turdus philomelos* (Turton) (Turdidae) – song thrush  
*Turdus merula* L. (Turdidae) – Eurasian blackbird

## FUNGI

*Alternaria* Nees (Pleosporaceae)  
*Beauveria bassiana* (Balsamo) Vuillemin (Cordycipitaceae)  
*Cladosporium* Link (Davidiellaceae)  
*Erysiphe polygoni* DC. (Erysiphaceae) – powdery mildew  
*Erysiphe cichoracearum* DC (Erysiphaceae)  
*Hesperomyces Thaxter* (Laboulbeniaceae)  
*Hesperomyces chilomenis* (Thaxter) Thaxter (Laboulbeniaceae)  
*Hesperomyces coccinelloides* (Thaxter) Thaxter (Laboulbeniaceae)  
*Hesperomyces hyperaspidis* Thaxter (Laboulbeniaceae)  
*Hesperomyces virescens* Thaxter (Laboulbeniaceae)  
*Isaria farinosa* (Holmskiold ex S. F. Gray) Fries (Cordycipitaceae)  
*Isaria fumosorosea* Wize (Cordycipitaceae)  
*Lecanicillium lecanii* (Zimmerman) Zare & Gams (Cordycipitaceae)  
[*Lecanicillium longisporum* (Zimmerman) Zare & Gams – as ‘Vertalec’] = *Lecanicillium lecanii* (Zimmerman) Zare & Gams (Cordycipitaceae)  
*Metarhizium anisopliae* (Metschnikoff) Sorokin (Clavicipitaceae)

*Microsphaera alphitoides* Griffon & Maublanc (Erysiphaceae)  
*Microsphaera pulchra* Cooke & Peck (Erysiphaceae)  
*Neotyphodium lolii* Latch et al. (Clavicipitaceae)  
*Nosema coccinellae* Lipa (Nosematidae)  
*Nosema epilachnae* Brooks, Hazard & Becnel (Nosematidae)  
*Nosema henosepilachnae* Toguebaye & Marchand (Nosematidae)  
*Nosema hippodamiae* Lipa & Steinhaus (Nosematidae)  
*Nosema tracheophila* Cali & Briggs (Nosematidae)  
*Nosema varivestis* Brooks, Hazard & Becnel (Nosematidae)  
*Tubulinosema hippodamiae* Bjornson, Le, Saito & Wang (Nosematidae)  
*Oidium monilioides* (Nees) Link (Erysiphaceae)  
[*Paecilomyces farinosus* (Holmskiold ex S. F. Gray) Brown & Smith] = *Isaria farinosa* (Holmskiold ex S. F. Gray) Fries (Cordycipitaceae)  
[*Paecilomyces funosoroseus* (Wize) Brown & Smith] = *Isaria fumosorosea* Wize (Cordycipitaceae)  
*Pandora neoaphidis* (Remaudière & Hennebert) Humber (Entomophthoraceae)  
*Phyllactinia moricola* (Hennebert) Homma (Erysiphaceae)  
*Podosphaera leucotricha* (Ellis & Everhart) E.S. Salmon (Erysiphaceae) – apple powdery mildew  
*Podosphaera tridactyla* (Wallroth) de Bary (Erysiphaceae) – plum powdery mildew  
*Puccinia* Pers. (Pucciniaceae)  
*Saccharomyces fragilis* Kudrjawzew (Saccharomycetaceae)  
*Sphaerotheca castagñe* Léverrier (Erysiphaceae) – hop powdery mildew  
*Sphaerotheca cucurbitae* (Jaczewski) Z.Y. Zhao (Erysiphaceae)  
*Sphaerotheca pannosa* (Wallroth) de Bary (Erysiphaceae) – rose powdery mildew  
[*Verticillium lecanii* (Zimmerman) Viegas] = *Lecanicillium lecanii* (Zimmerman) Zare & Gams (Cordycipitaceae)

## PROTISTS

*Anisolobus indicus* Haldar, Ray & Bose (Gregarinidae)  
*Brustiospora indicola* Kundu & Haldar (Brustiophoridae)  
*Gregarina* Dufour (Gregarinidae)  
*Gregarina barbarara* Watson (Gregarinidae)

*Gregarina californica* Lipa (*Gregarinidae*)  
*Gregarina chilocori* Obata (*Gregarinidae*)  
*Gregarina coccinellae* Lipa (*Gregarinidae*)  
*Gregarina dasguptai* Mandal, Rai, Pranhan, Gurung,  
 Sharma, Rai & Mandal (*Gregarinidae*)  
*Gregarina fragilis* Watson (*Gregarinidae*)  
*Gregarina hyashii* Sengupta & Haldar (*Gregarinidae*)  
*Gregarina katherina* Watson (*Gregarinidae*)  
*Gregarina ruszkowskii* Lipa (*Gregarinidae*)  
*Gregarina straeleni* Theodorides & Jolivet  
 (*Gregarinidae*)

**BACTERIA**

*Alcaligenes paradoxus* Davis (*Alcaligenaceae*)  
*Bacillus thuringiensis* Berliner (*Bacillaceae*)  
*Flavobacterium* Bergey (*Flavobacteriaceae*)  
*Rickettsia* da Rocha-Lima (*Rickettsiaceae*)  
*Spiroplasma* Saglio, Lhospital, Lafleche, Dupont, Bove,  
 Tully & Freundt (*Spiroplasmataceae*)  
*Wolbachia* Hertig (*Rickettsiaceae*)

**PLANTS**

*Abies balsamea* (L.) Miller (*Pinaceae*) – balsam fir  
*Abutilon theophrasti* Medikus (*Malvaceae*) – Chinese  
 jute or Indian Mallow  
*Acacia* Miller (*Fabaceae*)  
*Acalypha ostryaefolia* Riddell (*Euphorbiaceae*)  
*Aconitum* L. (*Ranunculaceae*)  
*Aframomum melegueta* (Roscoe) K. Schum  
 (*Zingiberaceae*) – grains of paradise or melegueta  
 pepper  
*Agropyron desertorum* (Link) Schultes (*Poaceae*)  
 – crested wheatgrass  
*Alnus japonica* (Thunberg) Steudel (*Betulaceae*)  
*Amaranthus hybridus* L. (*Amaranthaceae*)  
*Ambrosia artemisiifolia* L. (*Asteraceae*) – hogbrake  
*Anethum graveolens* L. (*Apiaceae*) – dill  
*Artemisia tridentata* Nuttall (*Asteraceae*) – sagebrush  
*Artemisia vulgaris* L. (*Asteraceae*) – wormwood  
*Atriplex sagittata* Borkhausen (*Chenopodiaceae*)  
*Benthamidia florida* (L.) Spach (*Cornaceae*)  
*Berberis vulgaris* L. (*Berberidaceae*) – common  
 barberry  
*Beta vulgaris* L. (*Chenopodiaceae*) – sugar beet and  
 garden beet

*Betula populifolia* Marshall (*Betulaceae*)  
*Brassica campestris* L. (*Brassicaceae*)  
*Brassica napus* L. (*Brassicaceae*) – rape  
*Brassica napus* L. subsp. *oleifera* DC Metzger  
 (*Brassicaceae*) – oil seed rape  
*Brassica nigra* (L.) Koch (*Brassicaceae*) – black (or  
 brown) mustard  
*Brassica oleracea* L. *Italica* group (*Brassicaceae*)  
 – broccoli  
*Brassica oleracea* L. (*Brassicaceae*)  
*Calotropis procera* (Aiton) Aiton (*Apocynaceae*)  
*Caltha palustris* L. (*Ranunculaceae*)  
*Cannabis sativa* L. (*Cannabaceae*) – Indian hemp  
*Carduus crispus* L. (*Asteraceae*)  
*Carpinus caroliniana* Walter (*Betulaceae*) – American  
 hickory  
*Caulophyllum robustum* Maximowicz (*Berberidaceae*)  
*Centaurea jacea* L. (*Asteraceae*)  
*Centrosema pubescens* Benthham (*Fabaceae*)  
*Chamerion angustifolium* (L.) Holub (*Onagraceae*)  
 – fireweed  
*Chenopodium* L. (*Chenopodiaceae*)  
*Cirsium arvense* (L.) Scopoli (*Asteraceae*) – creeping  
 thistle  
*Cirsium kagamontanum* Nakai (*Asteraceae*)  
*Cirsium kamtschaticum* De Candolle (*Asteraceae*)  
*Citrus sinensis* (L.) Osbeck (*Rutaceae*) – sweet orange  
*Clematis* L. (*Ranunculaceae*)  
*Coriandrum sativum* L. (*Apiaceae*) – coriander  
*Cosmos* (*Asteraceae*)  
*Cotoneaster integerrima* Medikus (*Rosaceae*)  
*Cotoneaster tomentosus* Lindley (*Rosaceae*)  
*Crotalaria striata* De Candolle (*Fabaceae*)  
*Cucurbita maxima* Duchesne (*Cucurbitaceae*) – squash  
 or pumpkin  
*Cymbopogon citratus* (De Candolle) Stapf (*Poaceae*)  
 – lemon grass  
*Dendrocalamus giganteus* Munro (*Poaceae*) – giant  
 bamboo  
*Elytrigia repens* (L.) Nevski (*Poaceae*) – couch grass  
*Erythrina corallodendron* L. (*Fabaceae*)  
*Euonymus japonicus* Thunberg (*Celastraceae*)  
*Euphorbia* L. (*Euphorbiaceae*)  
*Eurotia ceratoides* Meyer (*Chenopodiaceae*)  
*Genista* L. (*Fabaceae*)  
*Glochidion ferdinandi* (Müller Argiovensis) Bailey  
 (*Euphorbiaceae*)  
*Helianthus annuus* L. (*Asteraceae*) – sunflower  
*Hibiscus* L. (*Malvaceae*)  
*Juglans cinerea* L. (*Juglandaceae*) – butternut

- Juniperus virginiana* L. (Cupressaceae) – red cedar or Virginian pencil cedar
- Lapsana communis* L. (Asteraceae) – nipplewort
- Leontopodium alpinum* Cassini (Asteraceae) – edelweiss
- Leucaena* Benthams (Fabaceae)
- Ligustrum* L. (Oleaceae)
- Lolium multiflorum* Lamarck (Poaceae) – Italian rye grass
- Lolium perenne* L. (Poaceae) – rye grass
- Lonicera periclymenum* L. (Caprifoliaceae) – honeysuckle
- Lupinus luteus* L. (Fabaceae) – yellow lupin
- Lupinus mutabilis* Sweet (Fabaceae)
- Malus pumila* Miller (Rosaceae) – apple
- Manihot esculenta* Crantz (Euphorbiaceae)
- Manilkara zapota* (L.) P. Royen (Sapotaceae) – sapodilla
- Medicago sativa* L. (Fabaceae) – alfalfa (or often lucerne in UK)
- Mercurialis annua* L. (Euphorbiaceae) – annual mercury
- Momordica charantia* L. (Cucurbitaceae) – bitter melon
- Morus australis* Poir. (Moraceae)
- Nerium oleander* L. (Apocynaceae) – oleander
- Nitraria* L. (Nitrariaceae)
- Onoclea sensibilis* L. (Onocleaceae)
- Oryzopsis hymenoides* (Roemer & Schultes) Piper (Poaceae) – Indian ricegrass
- Paederia foetida* L. (Rutaceae)
- Pastinaca sativa* L. (Apiaceae) – parsnip
- Phaseolus vulgaris* L. (Fabaceae) – French bean or haricot bean
- Physalis alkekengi* L. (Solanaceae) – Chinese lantern
- Picea schrenkiana* Fischer & Meyer (Pinaceae)
- Pinus armandii* Franchet (Pinaceae) – Chinese white pine
- Pinus sylvestris* L. (Pinaceae) – Scots pine
- Pisonia* L. (Nyctaginaceae)
- Pisum sativum* L. (Fabaceae) – garden pea
- Pittosporum tobira* Aiton (Pittosporaceae)
- Populus* L. (Salicaceae)
- Prunus avium* (L.) L. (Rosaceae) – gean or wild cherry
- Prunus cerasus* L. (Rosaceae) – morello cherry or amarelle cherry
- Prunus persica* (L.) (Rosaceae) – peach or nectarine Batsch
- Pterostyrax hispidus* Siebold & Zuccarini (Styracaceae)
- Pyracantha coccinea* M.H. Roemer (Rosaceae) – firethorn
- Quercus rubra* L. (Fagaceae) – red oak
- Raphanus sativus* L. (Brassicaceae) – radish
- Robinia pseudoacacia* L. (Fabaceae)
- Rosa multiflora* Thunberg (Rosaceae)
- Rubus occidentalis* L. (Rosaceae)
- Salix* L. (Salicaceae)
- Sambucus nigra* L. (Caprifoliaceae) – elder
- Sambucus racemosa* L. subsp. *sieboldiana* (Miquel) H. Hara (Caprifoliaceae)
- Schizopepon bryoniaefolius* Maximowicz (Cucurbitaceae)
- Senecio jacobaea* L. (Asteraceae) – ragwort
- Senecio vulgaris* L. (Asteraceae) – groundsel
- Sinapis alba* L. (Brassicaceae) – white (or yellow) mustard
- Solanum japonense* Nakai (Solanaceae)
- Solanum nigrum* L. (Solanaceae) – black nightshade
- Solanum tuberosum* L. (Solanaceae) – potato
- Solidago canadensis* L. (Asteraceae)
- Spartium junceum* L. (Fabaceae) – Spanish broom
- Spiraea* (Rosaceae)
- Styrax* L. (Styracaceae)
- Symporicarpos rivularis* Sucksdorf (Caprifoliaceae)
- Talinum triangulare* (Jacquin) Willdenow (Portulacaceae)
- Tamarix* L. (Tamaricaceae)
- Tanacetum vulgare* L. (Asteraceae) – tansy
- Taraxacum officinale* Weber (Asteraceae) – dandelion
- Trichosanthes kirilowii* Maximowicz (Cucurbitaceae)
- Tripleurospermum maritimum* (L.) Koch (Asteraceae)
- Triticum* L. (Poaceae) – wheat
- Tsuga Carrière* (Pinaceae)
- Typha latifolia* L. (Typhaceae) – bullrush
- Ulmus* (Ulmaceae)
- Urtica dioica* L. (Urticaceae) – stinging nettle
- Verbascum thapsus* L. (Scrophulariaceae) – Aaron's rod
- Vicia faba* L. (Fabaceae) – broad bean or faba bean
- Vicia sativa* L. (Fabaceae) – vetch or tare
- Vigna unguiculata* (L.) Walpers subsp. *cylindrica* (L.) Verdcourt (Fabaceae)
- Withania somnifera* (L.) Dunal (Solanaceae)
- Zea mays* L. (Poaceae) – maize (corn in USA)
- Zingiber officinale* Roscoe (Zingiberaceae) – ginger