

A Checklist of the Heteroptera of the Canary Islands (Insecta)¹

B. AUKEMA, J.P. DUFFELS & M. BÁEZ

Abstract: New records of Heteroptera from the Canary Islands are listed and an updated checklist is presented. A new species of the genus *Noualhieria* (Lygaeidae) from El Hierro is described, 12 species are recorded from the archipelago for the first time, and 121 new island records are listed.

Key words: Canary Islands, checklist, distribution, Heteroptera, *Noualhieria heissi* nov.sp.

Introduction

The first checklist of the Heteroptera of the Canary Islands was published by LINDBERG (1953) and supplemented by LINDBERG (1960) and LINDBERG & WAGNER (1965). The second list was compiled by HEISS & BÁEZ (1990), followed by five supplements: HEISS & RIBES (1992), HEISS & WOUDESTRA (1993), HEISS et al. (1996), RIBES & RIBES (1997) and HEISS (1997). After the last supplement 372 species were listed from the islands. The next list (BÁEZ & ZURITA 2001) was included in a book on the biodiversity of the Canary Islands (IZQUIERDO et al. 2001), followed by an update (BÁEZ et al. 2004) in a second edition of the book (IZQUIERDO et al. 2004). The present contribution is based on fourteen short collecting trips since 1997, listed in Table 1, and some additional material deposited in the collection of the Zoological Museum, Amsterdam, the collection of the Zoology Department of the University of La Laguna, Tenerife and in the collection of E. Heiss, Innsbruck.

Seven species listed by BÁEZ et al. (2004) are not included in the checklist presented here, because their occurrence on the Canary Islands needs to be specified or confirmed. It concerns the following species. *Phytocoris parvulus* REUTER 1880 was cited erroneously from the Canary Islands by KERZHNER & JOSIFOV (1999). The citation of *Pyrrhocoris apterus* (LINNAEUS 1758) from

Tenerife by MOULET (1995) is erroneous (KERZHNER 2001). *Dicranocephalus medius* (MULSANT & REY 1870) was cited from the Canary Islands by MOULET (1995) and already questioned by HEISS (1997). *Stictopleurus abutilon* (ROSSI 1790) was cited from the Canary Islands by GÖLLNER-SCHIEDING (1975), *Odontotarsus intermedius* HORVÁTH 1923 by GÖLLNER-SCHIEDING (1990). *Lygus pratensis* (LINNAEUS 1758) and *Holcostethus strictus vernalis* (WOLFF 1804) [listed as *Dryocoris strictus* (FABRICIUS 1803) by BÁEZ et al. (2004)] were removed from the list by HEISS & RIBES (1992). *Phytocoris* (*Compserocoris*) *sanctipetri* CARAPEZZA 1985 is removed from the list because all the specimens listed by HEISS et al. (1996) belong to *Ph.* (*C.*) *riegeri* J. RIBES & HEISS 2001. *Cercinthus elegans* (BRULLÉ 1839) was described from the Canary Islands without locality and is only known from the recently rediscovered holotype (MOULET & PLUOT-SIGWALT 1998). The record of this species from Gomera in BÁEZ et al. (2004) is erroneous.

Doubtful species records are listed separately by BÁEZ et al. (2004) and are not repeated here (see also HEISS & RIBES (1992), HEISS & WOUDESTRA (1993) and RIBES & RIBES (1997)). The presence of *Arenocoris fallenii* (SCHILLING 1829), however, is confirmed. *Byrsinus flavicornis* (FABRICIUS 1794), *Crocistethus waltianus* (FIEBER 1837) and *Ochetostethus opacus* (SCHOLTZ 1847) were

¹ This paper is dedicated to Prof. Dr. Ernst Heiss, Innsbruck, for his 70th birthday.

Table 1: Collecting trips 1997-2006.

El Hierro	10-19.iii.2002	A.C. Ellis-Adam
El Hierro	5-17.iii.2005	B. Aukema
El Hierro	14.xii.2005-16.i.2006	J. Bruers & G.M. Viskens
La Palma	21.ii-5.iii.1998	J.P. Duffels
La Gomera	11-24.iii.2000	B. Aukema
La Gomera	17-28.ii.2001	B. Aukema
Tenerife	20.iii-3.iv.2006	J.P. Duffels
Gran Canaria	15-25.ii.2000	A.C. & W.N. Ellis
Fuerteventura	8-20.iii.2002	B. Aukema
Fuerteventura	14.iii-1.iv.2004	J.P. Duffels
Fuerteventura	23.iii-5.iv.2006	B. Aukema
Fuerteventura/Lanzarote	14.iii-3.iv.2005	J.P. Duffels
Lanzarote	17-22.ii.1997	B. Aukema
Lanzarote	28.ii-14.iii.1999	B. Aukema

cited from the Canary Islands by LIS (1999). *Aelia acuminata* (LINNAEUS 1758), *Aelia gemari* KÜSTER 1852 and *Aelia klugii* HAHN 1833 were cited from the Canary Islands by DERJANSCHI & PÉRICART (2005). The presence of these species on the Canary Islands needs to be confirmed and specified before they can be included in the list.

The records of *Hydrometra stagnorum* (LINNAEUS 1758) from Fuerteventura and Lanzarote in BAENA & BÁEZ (1990), *Tingis ajugarum* (FREY-GESSNER 1872) from Fuerteventura, *Reuterista instabilis* (FIEBER 1861) from La Gomera and *Ancyrosoma leucogrammes* (GMELIN 1790) from Lanzarote in BÁEZ et al. (2004) are erroneous and excluded from the list. Records from the smaller islands are not listed separately, but are included under Fuerteventura (Isla de Lobos) and Lanzarote (Alegranza, Graciosa and Montaña Clara). BÁEZ & ZURITA (2001) and BÁEZ et al. (2004) also included a number of additional island records, partly from literature, but mainly based on unpublished data. As far as available, references or details of these records are included in the present contribution. Details of the remaining additional records were not available to the authors, but will be published elsewhere.

New records for the Archipelago, new island records and an updated checklist with island records are presented. Nomenclature and sequence of the taxa are based on the Catalogue of the Heteroptera of the Palaearctic Region (AUKEMA & RIEGER 1995-2006). Four types of records are included: specified records from the literature, i.e. records of which collecting information and

depositories are published (● in Table 2), specified additional records from BÁEZ & ZURITA (2001) and BÁEZ et al. (2004) (▲), unspecified additional records from BÁEZ & ZURITA (2001) and BÁEZ et al. (2004) (□), and new records, specified here (■). The following abbreviations are used. AE: A.C. Ellis-Adam; BA: B. Aukema; JB: J. Bruers; JD: J.P. Duffels; MD: M.J. Duffels-Van Egmond; WE: W.N. Ellis; EH: E. Heiss; GV: G.M. Viskens; KBIN: Koninklijk Belgisch Instituut voor Natuurwetenschappen, Brussel; ULCI: Departamento Biología Animal, Universidad de La Laguna, Tenerife, Canary Islands; ZMAN: Zoölogisch Museum, Amsterdam. Unless stated otherwise the material is deposited in the collection of the collectors.

New records for the archipelago

Miridae

Campylomma verbasci (MEYER-DÜR 1843)

La Gomera: San Sebastián, 15.iii.2000, 1 ♀, BA.
Lanzarote: El Golfo, 20.ii.1996, 1 ♀, JD (ZMAN); W of Playa Blanca, Punta Pechiguera, 20.ii.1996, 3 ♂♂, 1 ♀, JD (ZMAN); Papagayo, Playa Mujeres, 28.ii.1996, 1 ♂, JD (ZMAN).

A widespread Palaearctic species (KERZHNER & JOSIFOV 1999).

Stenoparia putoni FIEBER 1870

Fuerteventura: Tierra Dorada, Bco de Mal Nombre, 2.iv.2005, 22 ♂♂, 17 ♀♀, JD (ZMAN).

A mediterranean species known from Spain, Italy (Sardinia, Sicily, Aeolian Islands), Greece, Turkey, Iraq, Israel, Saudi Arabia, Egypt, Lybia, Algeria, Tunisia and Morocco (KERZHNER & JOSIFOV 1999).

Reduviidae

Stenolemus novaki HORVÁTH 1888

Lanzarote: Playa Blanca, 14.iii.1999, 2 ♀♀, BA.

A mediterranean species known from Spain, France, Italy, Croatia, Israel, Egypt, Saudi Arabia and Sudan (PUTSHKOV & PUTSHKOV 1996).

Oncocephalus vescerae DISPONS 1965

Tenerife: Baranco del Infierno, 28.ix.1971, 1 ♂, H.G.M. Teunissen, det. M. Baena (ZMAN); Los Cristianos, 10-20.iii.1980, 1 ♂, 1 ♀, J.B. Wol-schrijn, det. M. Baena (ZMAN).

A rare species, known from the Middle East (Iran, Iraq and Israel) and from North Africa (Egypt and Algeria) (Putshkov & Putshkov 1996).

Lygaeidae

Cosmopleurus fulvipes (DALLAS 1852)

Fuerteventura: E of Morro Jable, gardens, 26.iii.2004, 1♂, JD (ZMAN).

A mediterranean species known from Spain, Croatia, Malta, the Middle East, North and tropical Africa and Pakistan (PÉRICART 2001).

Marmottania simonis

PUTON 1887 (Fig. 1)

Lanzarote: Playa Blanca, 3.iii.1999, BA, 1♀.

A rare east and south mediterranean species known from Iran, Israel, Yemen, Egypt and Algeria (PÉRICART 2001).

Noualhieria heissi nov.sp. (Figs 2-3)

The genus *Noualhieria* PUTON 1899 is endemic to the Canary Islands and represented by three species: *N. herbanica* J. RIBES 1976, *N. pieltaini* GÓMEZ MENOR 1924 and *Noualhieria quadripunctata* (BRULLÉ 1838). Specimens collected on El Hierro represent a new species, described here.

Description: Submacropterous. Male (Fig. 2). Head shiny blackish brown, shagreened and 1.2x as long as wide. Ocular index 2.0. Antennae long and slender, dark brown, total length 6.2 mm, 0.9x as long as the body. Proportions between antennal joints 34:50:39:45, first segment slightly darker, 1.2x as long as diatone and about as long as head. Rostrum long and slender, reaching to hind coxae, first segment blackish brown, remainder light brown. Pronotum shiny blackish brown with some light brownish spots along posterior margin, callar region shagreened, anterior margin and posterior half densely and coarsely punctate, 0.7x as long as wide. Scutellum shiny blackish brown, densely and coarsely punctate. Legs long and slender. Coxae, trochanters and basal part of femora white, apical part of femora broadly dark brown. Inner margin of fore femora with a short row of small teeth. Tibiae brown with lighter coloured base and tarsi light brown. Inner margin of meso- and metatibia with short semi-erect yellowish

pubescence and two rows of short spines, not longer as the diameter of the tibia. Evaporative area dull blackish brown, shagreened, with an irregular row of punctures bordering its upper anterior part. Hemelytra dull dark brown with small yellowish white lateral margins. Greatest width behind middle. Clavus with 4-5 irregular rows of punctures along the scutellum and a single row of punctures along the claval suture. Corium with a double row of punctures along the claval suture, scattered punctures on the endocorium and coarsely punctated on the exocorium. Membrane concolorous with corium, with four swollen veins, the inner two forming a closed cell. Abdomen shiny blackish brown with short yellowish pubescence, more dense on the last segments. Male genitalia, pygofer and paramere: Fig. 3. Length 6.9 mm, greatest width 2.1 mm.

Female. As male, but larger and broader with the following measurements. Head 1.2x as long as wide. Ocular index 2.1. Antennae 6.7 mm long, 0.8x as long as body. Proportions between antennal segments 39:59:44:39, first segment 1.3x as long as diatone and 1.1x as long as the head. Prono-



Fig. 1: *Marmottania simonis*, female (Photo: Th. Heijerman).



Fig. 2: *Noualhieria heissi* nov.sp., male (Photo: Th. Heijerman).




Fig. 3: *Noualhieria heissi*, (a) pygofores, ventral view (b) pygofores, lateral view (c) paramere.

tum 0.6x as long as wide. Length 8.3 mm, greatest width 2.9 mm.

Diagnosis: *Noualhieria heissi* nov.sp. can easily be distinguished from the other species of the genus. It differs from *N. pieltaini* (Fig. 4) in the absence of the long erect pubescence on head, pronotum and scutellum

and in the colour of the posterior margin of the pronotum, which is small yellowish in *N. pieltaini*. It differs from both *N. herbanica* and *N. quadripunctata* (Fig. 5) by the bicoloured femora.

Material Examined: **Holotype:** ♂, Canary Islands, El Hierro, Tanajara, 31.xii.2005, leg. J.M. Bruers (ZMAN). **Paratypes:** 1 ♀, same data as holotype (ZMAN); 1 ♀, Canary Islands, El Hierro, La Frontera, Tigaday, 5.iii.2005, leg. B. Aukema; 1♂, Canary Islands, El Hierro, Tanajara, 25.xii.2005,



Fig. 4: *Noualhieria pieltaini*, male (Photo: Th. Heijerman).



Fig. 5: *Noualhieria quadripunctata*, male (Photo: Th.



Fig. 6: *Tenosius proletarius*, male (Photo: Th. Heijerman).

leg. J.M. Bruers (KBIN); 1 ♀, Canary Islands, El Hierro, Tanajara, 26.xii.2005, leg. J.M. Bruers (KBIN).

Etymology: The species is named after Ernst Heiss, who greatly stimulated our interest in the Heteroptera fauna of the Canary Islands.

Rhopalidae

Agraphopus suturalis REUTER 1900

La Palma: El Mudo, E of Garafía, 350m, 1.iii.1998, 1 ♀, JD (ZMAN). **Fuerteventura:** Jandía, Morro Jable, swimming pool, 26.iii.2006, 1♂, 9 ♀♀, 27.iii.2006, 2♂♂, 8 ♀♀ and 28.iii.2006, 1 ♀, BA.

A species known from North Africa (Egypt, Libya, Algeria and Tunisia), tropical Africa (Sudan) and Asia (DOLLING 2006).

Alydidae

Euthetus humilis HORVÁTH 1907

Fuerteventura: Jandía, Jandía Playa, coastal dunes, 25.iii.2006, 2♂♂, 5 ♀♀, 28.iii.2006, 7♂♂, 13 ♀♀ and 30.iii.2006, 2♂♂, 1 ♀, BA; Jandía, Playa Esquinzo, coastal dunes, 1.iv.2006, 2 ♀♀, BA.

Euthetus humilis is known from Israel, Iran, Libya, Tunisia and tropical Africa (DOLLING 2006).

Tenosius proletarius (SCHAUM 1853) (Fig. 6)

Fuerteventura: Casas Risco del Paso, 3.iv.2005, 1 larva V, JD (ZMAN); Jandía, Jandía Playa, coastal dunes, 28.iii.2006, 2♂♂ and 7.iv.2006, 1♂, 2 ♀♀ ex larva, BA; Jandía, Playa Esmeralda, 29.iii.2006, 1 larva V, BA; Jandía, Playa de Esquinzo, 1.iv.2006, 1 ex., BA. **Lanzarote:** Tías, 8-24.ix.1996, 2♂♂, 1 ♀, AE & WE (ZMAN).

The species is known from Israel, Iran, Oman, Saudi Arabia, Yemen, Libya and tropical Africa (DOLLING 2006).

Note: Figure 62b in Moulet (1995) is strongly misleading in presenting a specimen with extremely short fourth antennal segments (about as long as the first segment) and a total antennal length of only about 1.9x the width of the pronotal base. The actual length of the fourth segment in our specimens equals the length of the first three segments together and the total length is about 2.8x the width of the pronotal base. This is in accordance with the data presented by Linnavuori (1987).



Fig. 7: *Arenocoris intermedius*, male (Photo: Th. Heijerman).

Coreidae

Arenocoris intermedius (JAKOVLEV 1883) (Fig. 7)

Fuerteventura: Vega de Río Palmas, Bco de las Peñitas, 11.iii.2002, 1♂, 3 ♀♀, 18.iii.2002, 2♂♂, 1 ♀, 13.iv.2002, 1 ♀ ex larva and 4.iv.2006, 1 ♀, BA; Jandía, Morro Jable, swimming pool, 1 ♀, BA; Jandía, Puerto de Morro Jable, 3.iv.2006, 1♂, 1 ♀ and 7.iv.2006, 2♂♂ ex larva from *Erodium*, BA; Jandía, Jandía Playa, coastal dunes, 3.iv.2006, 1 ♀ from *Erodium*, BA.

A mediterranean species, known from Portugal, Spain, Malta, Greece, the Middle East, Egypt, Libya, Morocco and tropical Africa (DOLLING 2006).

Phyllomorpha laciniata (VILLERS 1789)

Fuerteventura: 1♂ trapped in pitfall, D.R. Collins (det. W.R. Dolling 1981).

A widespread mediterranean species (DOLLING 2006).

Pentatomidae

Acrosternum millierei (MULSANT & REY 1866)

La Gomera: San Sebastián, 11.iii.2000, 1 ♀, 13.iii.2000, 1 ♀, 14.iii.2000, 1 ♀ at light, 15.iii.2000, 2 ♀♀, 19.iii.2000, 2 ♀♀ and 28.ii.2001, 4♂♂, 2 ♀♀, BA. **Fuerteventura:** Corralejo, 16.iii.2002, 7♂♂, 2 ♀♀ and 17.iii.2002, 1♂, BA; Corralejo, El Jable, 8.iii.2002, 2♂♂, BA; Jandía, Puerto Morro Jable, 28.iii.2006, 1♂, BA; Jandía, Playa Esquinzo, coastal dunes, 1♂, 2 ♀♀, 1.iv.2006, BA; Jandía, Morro Jable, 31.iii.2006, 2♂♂, 1 ♀ from *Pistacia atlantica* Desf., BA.

A widespread mediterranean species, also known from tropical Africa (RIDER 2006).

New island records

Corixidae

Sigara (Halicorixa) selecta (FIEBER 1848)

Lanzarote: Salinas de Janubio, 3.iii.1999, 10♂♂, 10♀♀, BA.

Notonectidae

Anisops sardeus sardeus

HERRICH-SCHAEFFER 1849

La Gomera: Embalse La Palmita, 12.iii.2000, 2♀♀ and 22.iii.2000, 1♂, 4♀♀, BA; Agulo, water reservoir, 14.iii.2000, 3♂♂, BA.

Gerridae

Gerris thoracicus SCHUMMEL 1832

El Hierro: Between S de Los Reyes and Mirador de Bascos, 7.iii.2005, 2♂♂, 1♀, BA.

Saldidae

Saldula palustris (DOUGLAS 1874)

Fuerteventura: Jandía, Morro Jable, swimming pool, 26.iii.2006, 1♂, BA.

Tingidae

Tingis (Tingis) denudata HORVÁTH 1906

El Hierro: La Caleta, Pte de La Caleta, 15.iii.2005, 18♂♂, 18♀♀ on *Volutaria canariensis*, BA.

Miridae

Nesidiocoris tenuis (REUTER 1895)

El Hierro: La Frontera, El Matorral, 12.iii.2005, 1♂, BA; Pozo de las Calcosas, 22.xii.2005, 1♂, JB (KBIN).

Note: The record from El Hierro in BÁEZ et al. (2001) was erroneous.

Creontiades pallidus (RAMBUR 1839)

La Gomera: San Sebastián, 15.iii.2000, 2♂♂, 1♀, BA. **Fuerteventura:** E of Morro Jable, gardens, 26.iii.2004, 1♂, JD (ZMAN).

Eurystylus bellevoeyi (REUTER 1879)

El Hierro: La Maceta, 26.xii.2005, 1♀, GV (KBIN).

Reuterista instabilis (FIEBER 1861)

El Hierro: 4 km NE of La Frontera, Las Puntas, sea level, 16.iii.2002, 1♂, AE (ZMAN); La Frontera, El Matorral, 12.iii.2005, 2♂♂, BA.

Dolichomiris linearis REUTER 1882

Lanzarote: Arrieta, Playa de la Garita, 23.ii.1997, 6♂♂, 13♀♀, BA.

Trigonotylus tenuis REUTER 1893

Lanzarote: Tías, 8-24.ix.1996, 2♂♂, 1♀, AE & WN (ZMAN).

Canariocoris pinicola (LINDBERG 1953)

El Hierro: Between S de Los Reyes and Mirador de Bascos, 7.iii.2005, 1♂, BA; S. de los Reyes, 9.i.2006, 1♂, 3♀♀, GV (KBIN).

Orthotylus (Melanotrichus) flavosparsus (C.R. SAHLBERG 1841)

El Hierro: La Frontera, El Matorral, 12.iii.2005, 1♀, BA; La Restinga, El Calderetón, 14.iii.2005, 2♀♀, BA; La Caleta, Pte de La Caleta, 15.iii.2005, BA, 1♂, 4♀♀. **Lanzarote:** Tías, 8-24.ix.1996, 2♀♀, AE & WE (ZMAN); Guatiza, salinas, 23.iii.2005, 14♂♂, 12♀♀, JD (ZMAN); El Papagayo, dunes, 25.iii.2005, 2♂♂, JD (ZMAN); Punta de Mujeres, N of Arrieta, rocks at seafront, 26-29.iii.2005, 8♂♂, 10♀♀, JD (ZMAN); Orzola, ruderal land, 27.iii.2005, 4♀♀, JD (ZMAN).

Orthotylus (Melanotrichus) salicorniae LINDBERG 1953

Lanzarote: Guatiza, Salinas los Cocoteros, 20.ii.1997, 2♀♀, BA.

Platycranus (Platycranus) lindbergi WAGNER 1954

La Gomera: Between El Cercado and Retamal, 20.ii.2001, 7♂♂, 3♀♀, BA.

Amblytulus scutellaris HORVÁTH 1905

Tenerife: S of Candelaria, rocky coast, 30.iii.2006, 1♂ and 1.iv.2006, 5♂♂, 1♀, MD & JD (ZMAN); El Médano, Playa de la Tejita, 2.iv.2006, 1♂, JD (ZMAN).

Badezorus signaticornis (REUTER 1904)

La Gomera: Pajarito, 18.iii.2000, 3♂♂, 2♀♀, BA; San Sebastian, 15.iii.2000, 2♀♀, BA.

Lindbergopsallus hyperici (LINDBERG 1953)

La Gomera: Between Las Rosas en Agulo, 22.ii.2001, 3♂♂, BA; Las Rosas, CdV Juego de Bolas, 23.ii.2001, 1♂, 1♀, BA.

Lindbergopsallus laureti (LINDBERG 1936)

La Gomera: Chorros de Epina, iv.1981, 2♂♂, 2♀♀, G. Gyllerfors (EH).

Lindbergopsallus rumicis (LINDBERG 1953)

La Gomera: Barranco del Cedro, between El Cedro and San Pedro, 25.ii.2001, 11♂♂, 6♀♀ from *Rumex lunaria*, BA.

***Maurodactylus nigrigenis* (REUTER 1890)**

Fuerteventura: Vega de Río Palmas, vi.1989, 1♂, 1♀, G. Gyllerfors (EH).

***Tuponia (Chlorotuponia) oculata* WAGNER 1954**

Lanzarote: Arrieta, Playa de la Garita, 21.ii.1997, 4♂♂, 3♀♀ from *Tamarix*, BA.

***Tytthus parviceps* (REUTER 1890)**

La Gomera: San Sebastián, 15.iii.2000, 6♂♂, 6♀♀, BA; Embalse de Agulo, 26.ii.2001, 1♂, BA. **Lanzarote:** Tías, 8-24.ix.1996, 3♀♀, AE & WE (ZMAN).

Nabidae

***Prostemma (Prostemma) guttula guttula* (FABRICIUS 1787)**

La Gomera: Juel, 1.v.1995, 1♀, P. Oromí (EH). **Fuerteventura:** Las Mercedes, 800 m, 20.v.1972, 1♀, Richter (EH)

***Nabis (Halonabis) occidentalis* (KERZHNER 1963)**

Lanzarote: Playa Blanca, 3.iii.1999, 1♂, BA.

Anthocoridae

***Anthocoris nemoralis* (FABRICIUS 1794)**

El Hierro: La Caleta, Punta de la Caleta, 15.iii.2005, 1♂, 8♀♀, BA. **La Gomera:** San Sebastián, 24.iii.2000, 7♂♂, 8♀♀, BA. **Lanzarote:** Arrieta, Playa de Garita, 17.ii.1997, 6♂♂, 4♀♀ and 21.ii.1997, 2♂♂, 5♀♀, BA; Playa Blanca, El Berrugo, 1.iii.1999, 2♂♂, 6♀♀, BA; Los Valles, NE of Teguisse, 26.iii.2005, 4♀♀ and 28.iii.2005, 2♂♂, 1♀, JD (ZMAN).

***Elatophilus (Euhadrocerus) pilosicornis* LINDBERG 1953**

El Hierro: Mirador de las Playas, 15.i.2006, 2♀♀, JB (BA, KBIN).

***Montandoniola moraguesi* (PUTON 1896)**

Fuerteventura: Jandía, Morro Jable, 25.iii.2006, 1♀, 31.iii.2006, 1♂, 2♀♀, 2.iv.2006, 2♂♂, 3♀♀ ex larva and 5.iv.2006, 6♂♂, 7♀♀, all from *Ficus* infested with *Gynaikothrips ficorum* (MARCHAL 1908), BA.

***Orius (Dimorphella) albidipennis* (REUTER 1884)**

El Hierro: Sabinosa, Pl Arenas Blancas, 8.iii.2005, 2♂♂, BA; La Frontera, El Matorral, 12.iii.2005, 2♀♀, BA. **La Gomera:** San Sebastián, 28.ii.2001, 1♂, BA. **Lanzarote:** Tías, 8-24.ix.1996, 1♂, 2♀♀, AE & WE (ZMAN); Playa Blanca, Salinas de Berrugo, 16-18.i.1996, 1♀,

AE & WE (ZMAN); Guatiza, salinas, 23.iii.2005, 1♀, JD (ZMAN); El Papagayo, dunes, 25.iii.2005, 1♂, 4♀♀, JD (ZMAN); Punta de Mujeres, N of Arrieta, rocks at seafront, 26-29.iii.2005, 6♀♀, JD (ZMAN); Orzola, ruderal land, 27.iii.2005, 1♀, JD (ZMAN).

***Orius (Orius) laevigatus maderensis* (REUTER 1884)**

Fuerteventura: Jandía, Morro Jable, 23.iii.2006, 4♂♂, BA.

Note: The record in BÁEZ et al. (2004) was based on NOUALHIER (1893) and had to be confirmed.

***Cardiastethus nazarenus* REUTER 1884**

La Gomera: NW El Cedro, 16.iii.2000, 1♀, BA. **Fuerteventura:** Jandía, Morro Jable, 31.iii.2006, 2♂♂, 8♀♀, 2.iv.2006, 1♀ ex larva and 5.iv.2006, 1♂, 1♀, BA.

Reduviidae

***Ploiaria chilensis* (PHILIPPI 1862)**

Gran Canaria: El Palmital, ix.1988, 1♀, G. Gyllerfors (EH).

***Empicoris rubromaculatus* (BLACKBURN 1889)**

Fuerteventura: Corralejo, 8.iii.2002, 10♂♂, 3♀♀ and 10.iii.2002, 4♂♂, 5♀♀, BA; Jandía, Morro Jable, 1♀ and 1 larva from *Ficus* infested with *Gynaikothrips ficorum* (MARCHAL 1908) and 31.iii.2006, 7♂♂, 17♀♀, BA.

***Oncocephalus pilicornis* REUTER 1882**

La Gomera: Embalse de Agulo, 26.ii.2001, 1♂, 8.iv.2001, 1♂ ex larva, and 10.iv.2001, 1♀ ex larva, BA.

***Coranus (Coranus) griseus* (ROSSI 1790)**

El Hierro: S. de los Reyes, 9.i.2006, 1♂, JB (KBIN). **Fuerteventura:** Jandía, iii.1989, 1♂, 1♀, G. Gyllerfors (EH).

Lygaeidae

***Caenocoris nerii* (GERMAR 1847)**

El Hierro: La Caleta, 13.i.2006, 1♂, JB (KBIN). **Fuerteventura:** Jandía, Morro Jable, 27.iii.2006, 1♀ and 2.iv.2006, 1♀, BA. **Lanzarote:** Playa Blanca, El Berrugo, 1.iii.1999, 1♂, BA.

***Lygaeosoma sardeum sardeum* SPINOLA 1837**

El Hierro: La Frontera, Tigaday, 13.iii.2005, 2♀♀ and 14.iii.2005, 2♀♀, BA. **Fuerteventura:** Castillo de Fuste, 12 km S Puerto del Rosario, 15.iii.2004, 1♂, JD (ZMAN).



Fig. 8: *Geocoris megagephalus*, female (Photo: Th. Heijerman).

***Spilostethus saxatilis* (SCOPOLI 1763)**

Fuerteventura: Corralejo, iii. 1989, 2♂♂, 2 ♀♀, G. Gyllerfors (EH); Corralejo, dunes, 20.iii.2002, BA, 1 ♀ and 30.xii.2004, H.T. Edzes, 3♂♂, 4 ♀♀ (BA).

***Orsillus pinicanariensis* LINDBERG 1953**

El Hierro: Mirador de los Playas, 5.i.2006, 1♂, JB (BA), 1 ♀ (KBIN).

***Engistus exsanguis exsanguis* STÅL 1872**

La Palma: La Fajana Franceses, W of Barlovento, 400 m, 24.ii.1998, 1 ♀, JD (ZMAN); La Fajana Franceses, W of Barlovento, 0-100 m, 28.ii.1998, 2♂♂, 3 ♀♀, JD (ZMAN).

***Henestaris laticeps wagneri* LINDBERG 1960**

El Hierro: Echedo, 22.xii.2005-3.i.2006, 8♂♂, 2 ♀♀ from *Frankenia laevis* L., JB & GV (BA, KBIN). **Fuerteventura:** Castillo de Fuste, 12 km S Puerto del Rosario, 15.iii.2004, 1 ♀, JD (ZMAN); Jandía, Punta del Matorral, 24.vi.2006, 12♂♂, 7 ♀♀ from *Frankenia laevis* L., BA.

***Geocoris (Eilatus) chloroticus* PUTON 1888**

La Palma: La Fajana Franceses, W of Barlovento, 0-100 m, 28.ii.1998, 1♂, JD (ZMAN).

***Geocoris (Geocoris) megagephalus* (ROSSI 1790) (Fig. 8)**

Fuerteventura: Bco de los Molinos, water reservoir, 20.iii.2004, 3♂♂, 3 ♀♀, JD (ZMAN); Casas de las Salinas, 15 km S Puerto del Rosario, 15.iii.2004, 1 ♀, JD (ZMAN); Corralejo, dunes El Jable, 11.iii.1999, 1 ♀, BA; Corralejo, 6.iii.2002, 17♂♂, 11 ♀♀ and 17.iii.2002, 1 ♀, BA; Los Molinos, 13.iii.2002, 2♂♂, BA; Jandía, Puerto de Morro Jable, 3.iv.2006, 1 ♀, BA;

Jandía, Jandía Playa, coastal dunes, 3.iv.2006, 1 ♀, BA; Jandía, Playa de Esquinzo, 1.iv.2006, 2♂, 3 ♀♀, BA.

Note: Records of *G. pallidipennis pallidipennis* (A. COSTA 1843) may also refer to this species.

***Heterogaster urticae* (FABRICIUS 1775)**

Fuerteventura: Vega de Río Palmas, Bco de las Peñitas, 4.vi.2006, 2 ♀♀ from *Forsskaolea angustifolia*, BA.

***Leptodemus minutus* (JAKOVLEV 1874)**

Fuerteventura: Jandía, Jandía Playa, coastal dunes, 25.iii.2006, 1♂, BA; Jandía, Playa Esmeralda, coastal dunes, 29.iii.2006, 4♂♂, 5 ♀♀, BA.

***Macropternella inermis* (FIEBER 1851)**

Fuerteventura: Jandía, Costa Calma, Beach Hotel Sunrise, dunes, 1.iv.2004, 3♂♂, 3 ♀♀, JD (ZMAN).

***Microplax plagiata* (FIEBER 1837)**

Lanzarote: Arrieta, Playa de la Garita, 23.ii.1997, 1♂, 5 ♀♀, BA.

***Emblethis angustus* MONTANDON 1890**

El Hierro: Malpaso, Chorro del Jable, 9.iii.2005, 1♂, 2 ♀♀, BA; La Frontera, Tigaday, 10.iii.2005, 4♂♂, BA; Las Casillas, El Fayal, 11.iii.2005, 1♂, BA.

***Emblethis denticollis* HORVÁTH 1878**

El Hierro: La Frontera, El Matorral, 12.iii.2005, 1 ♀, BA; La Frontera, Tigaday, 5.iii.2005, 1♂ and 10.iii.2005, 3♂♂, 2 ♀♀, BA.

***Camptocera glaberrima* (WALKER 1872)**

Gran Canaria: Artenara, xi.1981, G. Gyllerfors (EH); San Mateo, xii.1984, 1♂, 1 ♀, G. Gyllerfors (EH); Las Lagunetas, xi.1988, 1 ♀, G. Gyllerfors (EH).

***Lamprodema maura* (FABRICIUS 1803)**

La Gomera: San Sebastián, 19.iii.2000, 1♂, 4 ♀♀, 21.iii.2000, 3♂♂ and 23.iii.2000, 7♂♂, 4 ♀♀, BA.

***Megalonotus praetextatus* (HERRICH-SCHAEFFER 1835)**

La Gomera: Alojera, xii.1981, 1 ♀, G. Gyllerfors (EH).

***Megalonotus sabulicola* (THOMSON 1870)**

La Palma: El Mudo, E of Garaffa, 350 m, 1.iii.1998, 1 ♀, JD (ZMAN); Puntallana, Playa de Nogales, N of Santa Cruz, 0-300 m, 21.ii.1998, 1 ♀, JD (ZMAN). **La Gomera:** Embalse La Palmita, 22.iii.2000, 2 ♀♀, BA.

Remaudiereana annulipes

(BÄRENSPRUNG 1859)

El Hierro: Sabinosa, 19.xii.2005, 1♂, GV. **Lanzarote:** Guatiza, Salinas los Cocoteros, 20.ii.1997, 3♂♂, 12 ♀♀, BA; Playa Blanca, 3.iii.1999, 1♂ and 14.iii.1999, 1 ♀, BA.

Dieuches schmitzi REUTER 1893

La Gomera: San Sebastián, 13-24.iii.2000, 9♂♂, 17 ♀♀ and 28.ii.2001, 3♂♂, 6 ♀♀, BA.

Stygnocoris fuliginus (GEOFFROY 1785)

Gran Canaria: Cruz de Tejeda, vi.1983, 1♂, G. Gyllerfors (EH).

Berytidae

Berytinus (Lizinus) geniculatus

HORVÁTH 1885

El Hierro: El Pinar, 8.i.2006, 1♂, JB. **Fuerteventura:** Casas Risco del Paso, 2.iv.2005, 5♂♂, 10 ♀♀, JD (ZMAN); Jandía, Playa de Butihondo, 25.iii.2006, 3♂♂, 2 ♀♀, BA; Jandía, Playa Esmeralda, coastal dunes, 29.iii.2006, 3♂♂, 9 ♀♀, BA; Jandía, Jandía Playa, coastal dunes, 30.iii.2006, 1♂ and 3.iv.2006, 1♂, 1 ♀, BA; Jandía, Playa de Esquinzo, 1.iv.2006, 11♂♂, 9 ♀♀ from *Ononis*, BA.

Berytinus (Lizinus) montivagus MEYER-DÜR 1841 (Fig. 9)

Fuerteventura: Embalse de Los Molinos, 13.iii.2002, 1 ♀, BA.

Stenocephalidae

Dicranocephalus agilis (SCOPOLI 1763)

Fuerteventura: El Jable NW Tegui, iii.1995, 1 ♀, R. Grimm (EH).

Dicranocephalus pallidus (SIGNORET 1879)

Fuerteventura: Jandía, Morro Jable, swimming pool, 26.iii.2006, 1 ♀ and 30.iii.2006, 1 ♀, BA.

Rhopalidae

Brachycarenum tigrinus (SCHILLING 1829)

Lanzarote: Tías, 8-24.ix.1996, 1 ♀, AE & WE (ZMAN).

Coreidae

Arenocoris fallenii (SCHILLING 1829)

Fuerteventura: Jandía, Morro Jable, swimming pool, 26.iii.2006, 3♂♂, 3 ♀♀, BA.

Note: The record from Gran Canaria in BÁEZ et al. (2004) is erroneous. The species was listed from the Canary Islands by MOULET (1995), but up till now no record was published.



Fig. 9: *Berytinus montivagus*, female (Photo: Th. Heijerman).

Cercinthus lehmanni (KOLENATTI 1857) (Fig. 10)

La Gomera: San Sebastián, 18.ii.2001, 1♂, 1 ♀ and 28.ii.2001, 4♂♂, BA.

Gonocerus juniperi (Herrich-Schaeffer 1839)

La Gomera: San Sebastián, 19.iii.2000, 1♂, 1 ♀ and 24.iii.2000, 1 ♀, BA.



Fig. 10: *Cercinthus lehmanni*, male (Photo: Th. Heijerman).

Fig. 11: *Chroantha ornatula*, male and egg batch (Photo: Th. Heijerman).



Cydnidae

Alonips intrusus (WAGNER 1953)

Fuerteventura: Corralejo, iii.1989, 1♂, G. Gyllerfors (EH).

Byrsinus pilosulus (KLUG 1845)

La Gomera: San Sebastián, beach, 23.iii.2000, 1♂, 2 ♀♀, BA.

Geotomus elongatus (HERRICH-SCHAEFFER 1840)

Gran Canaria: N San Agustín, i.1997, 2 ♀♀, R. Grimm (EH). **Lanzarote:** Playa Blanca, Montaña Roja, 6.iii.1999, 1♂, 10 ♀♀ and 9.iii.1999, 10♂♂, 19 ♀♀, BA.

Microporus nigrita (FABRICIUS 1794)

La Gomera: San Sebastián, beach, 19.iii.2000, 1 ♀, BA.

Linospa orbicularis (JAKOVLEV 1885)

Tenerife: El Médano, Playa de Tejita, 1♂, JD (ZMAN).

Canthophorus melanopterus melanopterus (HERRICH-SCHAEFFER 1835)

La Gomera: San Sebastián, 23.iii.2000, 1 ♀, BA.

Ochetostethus tarsalis (MULSANT & REY 1852)

La Gomera: Between Imada and Casas Agalan, 23.ii.2001, 1♂, BA.

Acanthosomatidae

Cyphostethus tristriatus (FABRICIUS 1787)

El Hierro: S de Los Reyes, 7.iii.2005, 1♂, 1 ♀, BA; **La Frontera:** 6.i.2006, 1♂, 1 ♀, JB (KBIN).

Scutelleridae

Odontoscelis dorsalis (FABRICIUS 1798)

Fuerteventura: Jandía, iii.1989, G. Gyllerfors (EH); Jandía, Morro Jable, 25.iii.2006, 1♂ from swimming pool, BA; Jandía Playa, coastal dunes, 25.iii.2006, 6♂♂ and 28.iii.2006, 1 ♀, BA.

Pentatomidae

Chroantha ornatula (HERRICH-SCHAEFFER 1842) (Fig. 11)

La Gomera: San Sebastián, 19.iii.2000, 1 ♀ and 24.iii.2000, 2♂♂, 3 ♀♀, BA.

Eysarcoris ventralis (WESTWOOD 1837)

Lanzarote: Playa Blanca, 28.ii.1999, 2♂♂, 11 ♀♀ and 9.iii.1999, 1♂, BA.

Mecidea lindbergi WAGNER 1954

El Hierro: La Restinga, Calderetón, 14.iii.2005, 2♂♂, 3 ♀♀, BA; **La Frontera:** Tigaday, 14.iii.2005, 2♂♂, BA; **Guarazoca:** 27-29.xii.2005, 2♂♂, 3 ♀♀, JB & GV (KBIN); **La Frontera:** 1-2.i.2006, 2 ♀♀, JB & GV (KBIN); **Sabinosa:** 4.i.2006, 1 ♀, JB (KBIN); **La Caleta:** 16.i.2006, 1♂, JB (KBIN).

Mecidea pallidissima JENSEN-HAARUP 1922

La Palma: Garafía, Puerto, 350m, 2.iii.1998, 2 ♀♀, JD (ZMAN). **Fuerteventura:** Bco de las Peñitas, 20.iii.2002, 2♂♂, BA. **Lanzarote:** 1 km S Mirador del Río, 19.ii.1996, 1 ♀, JD (ZMAN).

Acrosternum rubescens (NOUALHIER 1893)

El Hierro: El Pinar, Hoya del Gallego, 13.iii.2005, 1♂, BA; **La Frontera:** Guinea, 17.iii.2005, 2♂♂, 2 ♀♀, BA; **Los Llanillos:** Charco Azul, 17.iii.2005, 2♂♂, BA; **Sabinosa:** 19.xii.205, 1♂, JB (KBIN); **Frontera:** 30.xii.2005, 1♂, JB. **Fuerteventura:** Corralejo, xi.2004, EH.

Piezodorus teretipes (STÅL 1865)

La Gomera: San Sebastián, 13.iii.2000, 1 ♀, BA.

Menaccarus dohrnianus (MULSANT & REY 1866)

Fuerteventura: Jandía, iii.1989, 1♂, G. Gyllerfors (EH); **Playa Blanca:** 12.iii.2002, 1 ♀, BA; **Jandía:** Playa Esmeralda, coastal dunes, 29.iii.2006, 4♂♂, 10 ♀♀ and 2.iv.2006, 4♂♂, 8 ♀♀, BA.

Sciocoris (Neosciocoris) maculatus
FIEBER 1851

La Palma: DERJANSCHI & PÉRICART (2005).

Sciocoris (Parasciocoris) canariensis
LINDBERG 1953

El Hierro: La Restinga, Calderetón, 14.iii.2005, 1♂, BA.

Ventocoris modestus (JAKOVLEV 1880)

Fuerteventura: Bco de las Peñitas, 11.iii.2002, 1♀, 18.iii.2002, 1♂, 1♀ and 30.iii.2002, 1♂, 1♀ ex larva, BA; Jandía Playa, coastal dunes, 30.iii.2006, 1 larva, BA. Lanzarote: Arrieta, Playa de la Garita, 23.ii.1997, 2♂♂, 2♀♀, BA.

Data of additional island records included in BÁEZ & ZURITA (2001) and BÁEZ et al (2004)

Meoveliidae

Mesovelia vittigera HORVÁTH 1895

Cran Canaria: NILSSON et al. (1998).

Veliidae

Microvelia (Microvelia) gracillima
REUTER 1882

Cran Canaria: NILSSON et al. (1998).

Tingidae

Derephysia (Derephysia) foliacea foliacea (FALLÉN 1807)

Tenerife: BAENA & MORALES (1993); OROMÍ et al. (2002).

Lygaeidae

Horvathiolus canariensis (WAGNER 1954)

La Gomera: Juego de Bolas, 29.x.2001, 1 ex., P. Oromí (ULCI).

Horvathiolus superbus (POLLICH 1781)

Tenerife: PÉRICART (1999).

Kleidocerys truncatulus (WALKER 1872)

Fuerteventura: La Oliva, 6.xii.1979, 1 ex., R. García (ULCI).

Geocoris (Geocoris) megacephalus
(ROSSI 1790)

Lanzarote: Montaña Clara, Llano del Aljibe (PÉREZ et al. 2003).

Oxycarenus (Oxycarenus) lavaterae
(FABRICIUS 1787)

Fuerteventura: CARNERO HERNÁNDEZ et al. (1991).

Ischnocoris mundus (WALKER 1872)

Lanzarote: Montaña Clara, Cresta (PÉREZ et al. 2003).

Megalonotus opaconotum
(LINDBERG 1953)

La Palma: La Grama, 10.ix.1997, 1 ex., R. García (ULCI).

Xanthochilus saturnius (ROSSI 1790)

Fuerteventura: La Matilla, 20.ii.1995, 1 ex., P. Oromí (ULCI); Betancuria, 9.ii.1997, 2 exx., P. Oromí (ULCI).

Reduviidae

Empicoris rubromaculatus
(BLACKBURN 1889)

Lanzarote: Montaña Clara, Caldera (PÉREZ et al. 2003).

Ectomocoris (Ectomocoris) ululans
(ROSSI 1790)

La Palma: Juan Adalid, 15.i.1994, 2 exx., P. Oromí (ULCI).

Reduvius personatus (LINNAEUS 1758)

La Gomera: Hermigua, 8.viii.1991, 1 ex. P. Oromí (ULCI).

Rhopalidae

Liorhyssus hyalinus (FABRICIUS 1794)

El Hierro: MARTIN (1992)

Cydnidae

Byrsinus albipennis (A. COSTA 1853)

Fuerteventura: Jandía, El Paso, 11.ii.1997, 3 exx., P. Oromí (ULCI).

Concluding remarks

A total number of 385 species of Heteroptera are listed from the Canary Islands. One new species is described and twelve species are recorded for the first time from the archipelago. Altogether, 121 new islands records are presented as follows (between brackets the total number of known species): El Hierro 24 (127), La Palma 6 (186), La Gomera 27 (169), Tenerife 3 (289), Gran Canaria 4 (235), Fuerteventura 35 (165) and Lanzarote 22 (137). 94 species (24.4%) are endemic (e.g., Figs 12-17) and of those 25 are restricted to one of the Canarian Islands (Table 2). By far the most endemics (71) are found on Tenerife, followed by Gran Canaria (53), La Palma (46) and La Gomera (42). Fuerteventura and Lanzarote, the two islands closest to the African continent, show a sig-



nificant lower level of endemism (20 species each), about half the level of the other islands (Figs 18-23).

Acknowledgements

Thanks are due to R. Beenen, J. Bruers, H.T. Edzes, A.C. Ellis-Adam, W.N. Ellis, H.G.M. P. Oromí, Teunissen, G.M. Viskens and J.B. Wolschrijn for their collecting efforts and to Th. Heijerman for making photographs. The *Oncocephalus* material was identified by M. Baena. E. Heiss is thanked for his support and advice.

Zusammenfassung

Neue Heteropteren-Funde von den Kanarischen Inseln werden aufgelistet und eine aktuelle Checkliste präsentiert. Eine neue Art der Gattung *Noualhieria* (Lygaeidae) wird von El Hierro beschrieben, 12 Arten werden erstmals von den Inseln genannt und 121 Arten werden erstmals für eine der Inseln genannt.

Figs 12-17: Endemic Heteroptera of the Canary Islands (12) *Cyrtopeltis canariensis* (LINDBERG 1936) (13) *Dicyphus bolivari atlanticus* WAGNER 1951 (14) *Aetorhinella parviceps* NOUALHIER 1893 (15) *Oncocephalus milleri* LINDBERG 1953 (16) *Stygnocoris subglaber* (PUTON 1889) (17) *Aneurus tagasastei* ENDERLEIN 1931 (Photos: M. Báez).



Figs 18-23: The Canary Islands, a fascinating spot of habitat and species diversity and beauty (18) Tenerife, Las Cañadas (19) Gran Canaria, Guayadeque (20) Hierro, Valverde (21) Fuerteventura, Jandia (22) La Palma, La Caldera (23) La Gomera, El Cedro (Photos: M. Báez).

Table 2: The Heteroptera of the Canary Islands. Abbreviations used: H – El Hierro, P – La Palma, G – La Gomera, T – Tenerife, C – Gran Canaria, F – Fuerteventura, L – Lanzarote. *: additional species; ^ε: endemic species; ●: published records; ▲: additional specified island records from BAEZ & ZURITA (2001) and BAEZ et al. (2004); □: additional unspecified island records from BAEZ & ZURITA (2001) and BAEZ et al. (2004); ■: new records.

	H	P	G	T	C	F	L
Corixidae							
<i>Corixa affinis</i> LEACH 1817	●	●	●	●	●	●	●
<i>Helocoris vermiculata</i> (PUTON 1874)	-	-	-	-	●	●	●
<i>Sigara (Halicorixa) selecta</i> (FIEBER 1848)	-	-	-	□	●	-	■
<i>Sigara (Tropocorixa) hoggarica</i> POISSON 1929	-	-	-	-	●	-	-
<i>Sigara (Vermicorixa) lateralis</i> (LEACH 1817)	●	●	●	●	●	●	●
Notonectidae							
<i>Anisops debilis canariensis</i> NOUALHIER 1893	●	●	●	●	●	●	-
<i>Anisops sardus sardus</i> HERRICH-SCHAEFFER 1849	●	●	■	●	●	●	●
^ε <i>Notonecta (Notonecta) canariensis</i> KIRKALDY 1897	-	-	●	●	●	-	-
Mesoveliidae							
<i>Mesovelia vittigera</i> HORVÁTH 1895	-	□	-	●	▲	□	-
Hebridae							
^ε <i>Hebrus (Hebrus) pusillus canariensis</i> POISSON 1954	-	-	●	●	●	-	-
<i>Merragata hebroides</i> F.B. WHITE 1877	●	□	-	●	●	●	-
Hydrometridae							
<i>Hydrometra stagnorum</i> (LINNAEUS 1758)	●	●	●	●	●	-	-
Veliidae							
<i>Microvelia (Microvelia) gracillima</i> REUTER 1882	-	▲	●	●	●	-	-
^ε <i>Velia (Plesiovelia) lindbergi</i> TAMAMINI 1954	-	●	●	●	●	-	-
Gerridae							
<i>Gerris (Gerris) thoracicus</i> SCHUMMEL 1832	■	-	●	●	●	-	-
Saldidae							
<i>Saldula amplipollis</i> (REUTER 1891)	-	●	-	-	-	-	-
<i>Saldula arenicola arenicola</i> (SCHOLTZ 1847)	-	-	●	●	-	●	-
<i>Saldula pallipes</i> (FABRICIUS 1794)	-	●	●	●	●	●	-
<i>Saldula palustris</i> (DOUGLAS 1874)	-	□	-	-	●	■	●
<i>Saldula pilosella</i> (THOMSON 1871)	-	-	-	-	●	-	-
Leptopodiidae							
<i>Leptopus hispanus</i> RAMBUR 1840	-	-	-	●	●	-	-
<i>Patapius spinosus</i> (ROSSI 1790)	-	-	-	●	●	-	-
Tingidae							
<i>Cantacader quadricornis</i> (LEPELETIER & SERVILLE 1828)	-	-	-	●	-	-	-
<i>Acalypta hellenica</i> REUTER 1888	-	-	●	●	●	-	-
<i>Acalypta parvula</i> (FALLÉN 1807)	-	●	●	●	-	-	-
<i>Derephysia (Derephysia) foliacea foliacea</i> (FALLÉN 1807)	-	-	-	▲	●	-	-
<i>Dictyla indigena</i> (WOLLASTON 1858)	●	●	●	●	●	-	-
<i>Dictyla nassata</i> (PUTON 1874)	●	●	●	●	●	●	●
^ε <i>Dictyonota atlantica</i> PÉRICART 1981	-	-	-	-	-	●	●
Dictyonota teydensis LINDBERG 1936							
Galeatus scrophicus SAUNDERS 1876							
Kalama moralesi (J. RIBES 1975)							
Kalama oromii (J. RIBES 1978)							
Tingis (Tingis) canariensis PÉRICART 1981							
Tingis (Tingis) cardui (LINNAEUS 1758)							
Tingis (Tingis) denudata HORVÁTH 1906							
Tingis (Tingis) maderensis (REUTER 1890)							
Tingis (Tropidocheila) ajugarum (FREY-GESSNER 1872)							
Tingis (Tropidocheila) insularis (HORVÁTH 1902)							
Tingis (Tropidocheila) liturata (FIEBER 1844)							
Microphysidae							
Loricula meinanderi PÉRICART 1972							
Loricula stenocephala J. RIBES 1985							
Miridae							
Campyloneuropsis fulva J. RIBES & E. RIBES 2001							
Cyrtopeltis (Canariesia) salviae CARAPEZZA 1988							
Cyrtopeltis (Cyrtopeltis) canariensis (LINDBERG 1936)							
Dicyphus (Brachyceroea) heissi J. RIBES & BAENA 2006							
Dicyphus (Dicyphus) baezi J. RIBES 1983							
Dicyphus (Dicyphus) bolivari atlanticus WAGNER 1951							
Dicyphus (Dicyphus) rubicundus BLÖTE 1929							
Macrolophus melanotoma (A. COSTA 1853)							
Macrolophus pericarti J. RIBES & HEISS 1998							
Nesidiocoris tenuis (REUTER 1895)							
Singhalesia tenuissima (LINDBERG 1959)							
Deraeocoris (Camptobrochis) serenus (DGL. & SC. 1868)							
Deraeocoris (Plexaris) martini martini (PUTON 1887)							
Closterotomus norvegicus (GMELIN 1790)							
Creontiades pallidus (RAMBUR 1839)							
Cyphodema instabilis (LUCAS 1849)							
Eurystylus bellevoeyi (REUTER 1879)							
Lygus maritimus WAGNER 1949							
Megacoleum zollikoferiae (LINDBERG 1953)							
Miridius quadrivirgatus (A. COSTA 1853)							
Miris striatus (LINNAEUS 1758)							
Orthops (Orthops) basalis (A. COSTA 1853)							
Phytocoris (Compserocoris) retamae REUTER 1900							

	H	P	G	T	C	F	L
♯ <i>Phytocoris (Comperocoris) riegeri</i> J. RIBES & HEISS 2001	-	-	-	-	●	-	-
♯ <i>Phytocoris (Ecklerleinius) meinanderi</i> WAGNER 1968	-	-	-	-	-	●	●
<i>Phytocoris (Leptophytocoris) azrouensis</i> WAGNER 1959	-	-	-	-	-	●	-
<i>Phytocoris (Leptophytocoris) miridoioides</i> (LETHIERRY 1877)	-	-	-	●	●	●	-
<i>Pinalitus conspurcatus</i> (REUTER 1875)	-	□	●	●	●	●	●
<i>Pinalitus insularis</i> (REUTER 1895)	●	●	●	●	●	●	-
<i>Pinalitus parvulus</i> (REUTER 1879)	-	-	-	●	-	-	-
<i>Polymerus (Poeciloscytus) unifasciatus</i> (FABRICIUS 1794)	-	-	-	-	●	-	-
<i>Reuterista instabilis</i> (FIEBER 1861)	■	-	-	●	●	●	●
<i>Taylorilygus apicalis</i> (FIEBER 1861)	-	●	●	●	●	●	●
<i>Acetropis (Acetropis) gimmerthalii</i> <i>gimmerthalii</i> (FLOR 1860)	●	●	●	-	-	-	-
<i>Dolichomiris (Dolichomiris) linearis</i> REUTER 1882	□	●	●	●	●	●	■
<i>Pithanulus marshalli</i> DOUGLAS & SCOTT 1868	-	-	-	-	-	●	-
<i>Trigonotylus tenuis</i> REUTER 1893	-	-	-	●	●	●	■
<i>Pachytomella passerinii</i> (A. COSTA 1842)	-	●	-	●	●	-	-
♯ <i>Aetorhinella parviceps</i> NOUALHIER 1893	●	●	●	●	●	●	-
♯ <i>Canariocoris antennalis</i> (REUTER 1904)	-	-	-	●	●	●	-
♯ <i>Canariocoris ater</i> HEISS 1996	●	-	-	-	-	-	-
♯ <i>Canariocoris euphorbiae</i> LINDBERG 1951	-	●	-	●	-	-	-
♯ <i>Canariocoris hyperici</i> LINDBERG 1951	●	-	●	-	-	-	-
♯ <i>Canariocoris pinicola</i> (LINDBERG 1953)	■	●	-	●	●	-	-
♯ <i>Canariocoris punctatus</i> (NOUALHIER 1893)	-	-	●	●	●	●	●
♯ <i>Canariocoris viburni</i> (LINDBERG 1953)	-	●	●	●	●	●	●
<i>Heterocordylus (Heterocordylus) tibialis</i> (HAHN 1833)	-	-	●	●	●	-	-
<i>Orthotylus (Melanotrichus) divinus</i> LINNAVUORI 1961	-	-	-	-	-	-	●
<i>Orthotylus (Melanotrichus) fieberi</i> <i>punctipes</i> (REUTER 1895)	-	-	-	-	-	-	●
<i>Orthotylus (Melanotrichus) flavosparsus</i> (C.R. SAHLBERG 1841)	■	-	●	●	●	●	■
♯ <i>Orthotylus (Melanotrichus) halophilus</i> LINDBERG 1953	-	-	-	-	●	●	●
♯ <i>Orthotylus (Melanotrichus) salicorniae</i> LINDBERG 1953	-	-	-	-	-	●	■
♯ <i>Platycranus (Platycranus) lindbergi</i> WAGNER 1954	-	●	■	●	-	-	-
<i>Haliolapus concolor</i> (REUTER 1890)	-	-	●	●	-	-	-
<i>Mimocoris</i> sp.	-	-	●	-	-	-	-
♯ <i>Acrotelus canariensis</i> WAGNER 1943	-	-	-	-	-	●	●
<i>Amblytulus scutellaris</i> HORVATH 1905	-	-	-	■	●	-	-
<i>Atomoscelis onusta</i> (FIEBER 1861)	-	-	-	●	●	●	●
♯ <i>Auchenocrepis nigricornis</i> WAGNER 1954	-	-	-	-	●	-	-
<i>Auchenocrepis similis</i> WAGNER 1954	-	-	-	-	-	●	-
<i>Badezorus signaticornis</i> (REUTER 1904)	●	●	■	●	●	●	●
♯ <i>Campylomma verbasci</i> (MEYER-DÜR 1843)	-	-	■	-	-	-	■
<i>Compsidolon (Apsinthophylus) balachowskyi</i> (WAGNER 1958)	-	-	-	-	-	●	-
♯ <i>Compsidolon (Chamaeliops) collare</i> WAGNER 1976	-	-	-	-	-	●	-
♯ <i>Compsidolon (Coniortodes) anagae</i> GYLLENSVÄRD 1968	-	-	-	-	-	●	-
♯ <i>Compsidolon (Coniortodes) beckeri</i> (REUTER 1904)	-	●	-	-	-	-	-
♯ <i>Compsidolon (Coniortodes) cytisellum</i> (LINDBERG 1953)	-	-	-	-	-	●	-
♯ <i>Compsidolon (Coniortodes) cytisi</i> (LINDBERG 1953)	-	-	-	-	-	●	-
♯ <i>Compsidolon (Coniortodes) freyi</i> (WAGNER 1954)	-	●	●	●	●	●	-
♯ <i>Compsidolon (Coniortodes) galbanus</i> GYLLENSVÄRD 1968	-	-	-	-	-	●	-
♯ <i>Compsidolon (Coniortodes) hierroense</i> (WAGNER 1954)	●	●	●	●	●	●	-
♯ <i>Compsidolon (Coniortodes) longiceps</i> (REUTER 1904)	-	●	-	●	●	●	-
♯ <i>Compsidolon (Coniortodes) parviceps</i> (WAGNER 1954)	-	●	●	●	●	●	-
♯ <i>Compsidolon (Coniortodes) verbenae</i> (WAGNER 1954)	-	●	-	●	●	●	-
♯ <i>Compsidolon longiceps</i> (WAGNER 1965)	-	-	-	-	-	-	●
♯ <i>Conostethus venustus longicornis</i> WAGNER 1965	●	●	●	●	●	●	●
<i>Hadrophyes sulphurella</i> PUTON 1874	-	-	-	-	-	-	●
♯ <i>Lindbergopsallus hyperici</i> (LINDBERG 1953)	●	●	■	●	●	●	-
♯ <i>Lindbergopsallus impunctatus</i> WAGNER 1965	-	-	-	-	-	●	-
♯ <i>Lindbergopsallus instabilis</i> (REUTER 1904)	●	●	●	●	●	●	-
♯ <i>Lindbergopsallus laureti</i> (LINDBERG 1936)	●	-	■	●	●	●	-
♯ <i>Lindbergopsallus rumicis</i> (LINDBERG 1953)	●	-	■	●	●	●	-
<i>Macrotylus (Alloeonycha) nigricornis</i> FIEBER 1864	-	-	-	●	●	●	-
<i>Macrotylus (Alloeonycha) paykullii</i> (FALLEN 1807)	-	-	-	-	-	●	●
♯ <i>Macrotylus (Alloeonycha) spergulariae</i> LINDBERG 1953-	-	-	-	-	-	●	●
<i>Maurodactylus alutaceus</i> (FIEBER 1870)	-	-	-	-	-	●	-
<i>Maurodactylus nigrigenis</i> (REUTER 1890)	-	-	-	●	●	■	●
♯ <i>Moissonia halophila</i> (LINDBERG 1953)	-	-	-	-	-	●	●
<i>Pastocoris putonii</i> (REUTER 1875)	-	-	-	-	-	-	●
* <i>Stenoparia putoni</i> FIEBER 1870	-	-	-	-	-	-	■
<i>Tuponia (Chlorotuponia) longipennis</i> HORVATH 1909	-	-	-	-	-	●	●
♯ <i>Tuponia (Chlorotuponia) oculata</i> WAGNER 1954	-	●	●	●	●	●	■
<i>Tuponia (Tuponia) mixticolor</i> (A. COSTA 1862)	-	-	-	-	-	●	●
♯ <i>Tuponia (Tuponia) rubella</i> PUTON 1889	●	●	●	●	●	●	-
<i>Tyttthus parviceps</i> (REUTER 1890)	-	-	-	■	●	●	■
Nabidae							
<i>Prostemma (Prostemma) guttula</i> <i>guttula</i> (FABRICIUS 1787)	-	-	-	-	■	●	-
<i>Himacerus (Anaptus) mirmicoideus</i> (A. COSTA 1842)	-	●	-	-	-	●	-
<i>Himacerus (Aptus) mirmicoideus</i> (O. COSTA 1834)	-	-	-	-	-	●	-
<i>Nabis (Aspilaspis) viridulus</i> SPINOLA 1837	-	-	-	-	-	●	●
<i>Nabis (Halonabis) occidentalis</i> KERZHNER 1963	-	-	-	-	-	●	●
<i>Nabis (Nabis) pseudoferus ibericus</i> REMANE 1949	●	●	●	●	●	●	-

	H	P	G	T	C	F	L
<i>Nabis (Nabis) punctatus</i> A. COSTA 1847	-	□	-	●	-	-	-
<i>Nabis (Tropiconabis) capsiformis</i> GERMAR 1838	●	●	●	●	●	●	●
Anthocoridae							
<i>Anthocoris alienus</i> (F.B. WHITE 1880)	●	●	●	●	●	-	-
<i>Anthocoris nemoralis</i> (FABRICIUS 1794)	■	-	■	●	●	●	■
ε <i>Anthocoris salicis</i> LINDBERG 1953	-	●	●	●	-	-	-
<i>Anthocoris sarothamni</i> DOUGLAS & SCOTT 1865	-	-	-	-	●	-	-
ε <i>Elatophilus (Euhadrocerus) pilosicornis</i> LINDBERG 1953	■	□	-	●	●	-	-
<i>Montandoniella moraguesi</i> (PUTON 1896)	-	●	-	●	-	■	-
<i>Orius (Dimorphella) albidipennis</i> (REUTER 1884)	■	-	■	●	●	●	■
<i>Orius (Dimorphella) canariensis</i> WAGNER 1952	-	-	-	-	●	●	●
ε <i>Orius (Microtrachelia) retamae</i> (NOUALHIER 1893)	-	●	-	●	●	-	-
<i>Orius (Orius) laevigatus maderensis</i> (REUTER 1884)	-	●	●	●	●	■	●
<i>Orius (Orius) limbatus</i> WAGNER 1952	●	●	●	●	●	●	-
<i>Orius (Orius) lindbergi</i> WAGNER 1952	●	●	-	●	●	●	●
<i>Orius (Orius) niger</i> (WOLFF 1811)	-	-	-	●	●	-	-
<i>Orius (Orius) pallidicornis</i> (REUTER 1884)	-	●	-	-	-	-	-
<i>Orius (Orius) piceicollis</i> (LINDBERG 1936)	-	●	-	●	●	●	●
<i>Wollastoniella obesula</i> (WOLLASTON 1858)	-	-	-	●	-	●	-
<i>Brachysteles parvicornis</i> (A. COSTA 1847)	-	-	-	●	-	-	-
<i>Brachysteles wollastoni</i> F.B. WHITE 1880	-	-	-	●	●	●	-
<i>Buchananiella continua</i> (F.B. WHITE 1880)	-	●	-	-	-	-	-
<i>Cardiastethus nazarenus</i> REUTER 1884	-	-	-	■	●	-	■
<i>Dufouriellus ater</i> (DUFOUR 1833)	-	□	-	-	-	●	-
<i>Lycotocoris (Lycotocoris) campestris</i> (FABRICIUS 1794)	-	-	-	●	●	●	●
<i>Lycotocoris (Lycotocoris) dimidiatus</i> (SPINOLA 1837)	-	-	-	●	-	-	-
ε ⁷ <i>Lycotocoris (Lycotocoris) uytenboogaarti</i> BLÖTE 1929	□	●	□	●	●	●	●
ε <i>Lycotocoris (Paralycotocoris) menieri</i> CARAYON 1971	-	-	-	●	-	-	-
<i>Scoloposcelis obscurella</i> (ZETTERSTEDT 1838)	-	-	-	-	●	-	-
<i>Xylocoris (Arrostelus) flavipes</i> (REUTER 1875)	-	-	-	●	●	●	●
<i>Xylocoris (Proxylucoris) galactinus</i> (FIEBER 1836)	-	-	-	●	●	-	●
<i>Xylocoris (Xylocoris) canariensis</i> WAGNER 1954	-	□	-	●	-	-	-
ε <i>Xylocoris (Xylocoris) configuus</i> WAGNER 1954	●	●	●	●	●	●	-
<i>Xylocoris (Xylocoris) obliquus</i> A. COSTA 1853	-	-	-	●	-	-	-
Cimicidae							
<i>Cimex lectularius</i> LINNAEUS 1758	-	-	-	●	●	-	-
Reduviidae							
ε <i>Collartida anophthalma</i> ESPAÑOL & J. RIBES 1983	●	-	-	-	-	-	-
ε <i>Collartida tanausu</i> J. RIBES, OROMÍ & E. RIBES 1998	-	●	-	-	-	-	-
* <i>Stenolemus novaki</i> HORVÁTH 1888	-	-	-	-	-	-	■
<i>Ploiaria chilensis</i> (PHILIPPI 1862)	●	□	●	●	■	-	-
<i>Empicoris brevispinus</i> (PUTON 1889)	-	●	-	●	-	-	-
<i>Empicoris rubromaculatus</i> (BLACKBURN 1889)	-	●	-	●	-	■	▲
<i>Empicoris salinus</i> (LINDBERG 1932)	-	-	-	●	-	-	●
<i>Ectomocoris (Ectomocoris) chiragra</i> (FABRICIUS 1803)	-	□	-	●	●	-	-
<i>Ectomocoris (Ectomocoris) ululans</i> (ROSSI 1790)	-	▲	-	●	●	●	●
<i>Peirates strepitans</i> (RAMBUR 1839)	-	-	-	●	●	●	-
<i>Pasira lewisi</i> (MILLER 1951)	-	●	-	●	●	-	-
<i>Reduvius personatus</i> (LINNAEUS 1758)	●	□	▲	●	●	●	-
<i>Onccephalus pilicornis</i> REUTER 1882	-	-	■	●	●	-	-
<i>Onccephalus squalidus</i> (ROSSI 1790)	-	●	-	●	-	-	-
* <i>Onccephalus veserae</i> DISPONS 1965	-	-	-	-	■	-	-
ε <i>Onccephalus milleri</i> LINDBERG 1953	-	-	-	-	●	●	●
<i>Sastrapada baerensprungi</i> (STÄL 1859)	-	□	-	●	●	-	-
<i>Coranus (Coranus) aegyptius</i> (FABRICIUS 1775)	●	●	●	●	●	●	●
<i>Coranus (Coranus) griseus</i> (ROSSI 1790)	■	●	●	●	●	■	-
<i>Coranus (Coranus) kiritshenkoi</i> (BERGEVIN 1932)	-	-	-	-	●	●	●
Aradidae							
ε <i>Aneurus (Aneurodes) tagastei</i> ENDERLEIN 1931	●	●	●	●	-	-	-
ε <i>Aradus canariensis</i> KORMILIEV 1954	●	-	-	●	-	-	-
<i>Aradus flavicornis</i> DALMAN 1823	-	□	-	●	-	-	-
ε <i>Aradus lauri</i> NOUALHIER 1893	-	□	-	●	-	-	-
<i>Aradus reuterianus</i> PUTON 1875	-	●	-	●	●	-	-
Lygaeidae							
<i>Caenocoris nerii</i> (GERMAR 1847)	■	●	●	●	●	■	■
* <i>Cosmopleurus fulvipes</i> (DALLAS 1852)	-	-	-	-	-	-	-
ε <i>Horvathiolus albomaculata</i> (LINDBERG 1960)	-	-	-	-	●	-	-
<i>Horvathiolus canariensis</i> (WAGNER 1954)	-	-	-	▲	●	-	-
<i>Horvathiolus superbus</i> (POLLICH 1781)	-	-	-	-	▲	-	-
<i>Lygaeosoma sardeum sardeum</i> SPINOLA 1837	■	●	●	●	●	■	-
<i>Spilostethus pandurus</i> (SCOPOLI 1763)	●	●	●	●	●	●	●
<i>Spilostethus saxatilis</i> (SCOPOLI 1763)	-	-	-	-	●	■	●
<i>Nysius cymoides</i> (SPINOLA 1837)	●	●	●	●	●	●	●
<i>Nysius ericae ericae</i> (SCHILLING 1829)	-	●	-	●	-	-	-
ε <i>Nysius glorioae</i> BAENA & GARCÍA 1999	-	●	-	●	-	-	-
<i>Nysius immunis</i> (WALKER 1872)	●	●	●	●	●	●	●
<i>Nysius thymi latus</i> WAGNER 1958	□	●	●	●	●	●	●
ε <i>Orsillus pinicanariensis</i> LINDBERG 1953	■	●	●	●	●	●	●
<i>Kleidocerys truncatulus</i> (WALKER 1872)	●	●	●	●	●	●	▲
<i>Cymus clavicularis</i> (FALLÉN 1807)	-	-	-	●	●	-	-
<i>Cymus gracilicornis</i> VIDAL 1940	-	-	-	●	●	-	-
ε <i>Cymus remanei</i> HEISS & PERICART 1999	-	-	-	●	-	-	-
<i>Engistus boops boops</i> (DUFOUR 1857)	-	●	-	●	●	●	●
<i>Engistus exsanguis exsanguis</i> STÄL 1872	-	■	-	●	-	●	●

	H	P	G	T	C	F	L
€ Henestaris laticeps wagneri LINDBERG 1960	■	-	-	●	-	■	●
Geocoris (Eilatius) chloroticus PUTON 1888	-	■	-	●	●	●	●
Geocoris (Geocoris) collaris PUTON 1878	-	-	-	-	-	●	●
Geocoris (Geocoris) lineola lineola (RAMBUR 1839)	●	●	●	●	●	●	●
Geocoris (Geocoris) megacephalus (ROSSI 1790)	-	-	-	-	-	■	▲
Geocoris (Geocoris) pallidipennis pallidipennis (A. COSTA 1843)	-	-	-	-	-	●	●
Geocoris (Geocoris) phaeopterus (GERMAR 1838)	-	-	-	●	●	●	●
Geocoris (Geocoris) pubescens (JAKOVLEV 1871)	●	●	●	●	●	●	●
€ Heterogaster canariensis LINDBERG 1960	●	●	●	●	●	●	-
Heterogaster urticae (FABRICIUS 1775)	●	●	-	●	●	■	●
Platylax inermis (RAMBUR 1839)	●	-	●	●	●	-	●
€ Bethylimorphus leucophaeus LINDBERG 1953	-	●	●	●	-	-	-
Leptodemus minutus (JAKOVLEV 1874)	-	-	-	-	-	■	-
Macroplox fasciata fasciata (HERRICH-SCHAEFFER 1835)	-	-	-	-	-	●	●
Macroplox vicina PUTON 1889	●	●	●	●	●	●	-
Macropternella inermis (FIEBER 1851)	-	-	-	●	●	■	●
Microplax interrupta (FIEBER 1837)	●	●	●	●	●	-	-
Microplax plagiata (FIEBER 1837)	-	-	-	-	-	●	■
€ Neocampotelus aeonii (LINDBERG 1953)	●	-	●	●	●	-	-
Oxycaenus (Euoxycarenus) pallens (HERRICH-SCHAEFFER 1850)	-	-	-	●	●	-	-
Oxycaenus (Oxycaenus) hyalinipennis (A. COSTA 1843)	-	-	-	●	●	●	-
Oxycaenus (Oxycaenus) lavaterae (FABRICIUS 1787)	●	●	●	●	●	▲	●
Tropistethus seminitens PUTON 1889	●	●	●	●	●	-	●
Eremocoris maderensis (WOLLASTON 1858)	●	●	●	●	●	-	-
Ischnocoris mundus (WALKER 1872)	●	●	●	●	●	-	▲
Notochilus damryi PUTON 1871	-	-	-	-	-	-	-
Scolopostethus pilosus maderensis REUTER 1881	-	●	●	●	●	-	-
Aphanus rolandri (LINNAEUS 1758)	●	●	●	●	●	●	●
Emblethis angustus MONTANDON 1890	■	●	●	●	-	-	-
Emblethis denticollis HORVATH 1878	■	●	●	●	-	-	-
Emblethis duplicatus SEIDENSTÜCKER 1963	-	●	-	●	●	-	-
Emblethis griseus (WOLFF 1802)	●	●	●	●	●	-	-
Emblethis verbasci (FABRICIUS 1803)	-	□	-	●	-	-	-
Gonionotus galactodermus (FIEBER 1861)	●	●	●	●	●	●	●
Neurocladus brachiidens (DUFOUR 1851)	-	-	-	-	-	-	-
€ Pronosomus alticola LINDBERG 1953	-	-	-	●	●	-	-
Campocera glaberrima (WALKER 1872)	-	●	-	●	■	●	●
Lethaeus cf. fulvovarius PUTON 1884	-	●	-	-	-	-	-
Lethaeus lethierryi (PUTON 1869)	-	-	●	●	●	●	●
Lamprodema maura (FABRICIUS 1803)	●	□	■	●	●	●	●
Megalonotus emarginatus (REY 1888)	-	-	-	-	-	-	-
Megalonotus mixtus (HORVATH 1887)	-	-	-	-	-	-	-
€ Megalonotus opaconotum (LINDBERG 1953)	-	-	▲	-	●	●	-
Megalonotus praetextatus (HERRICH-SCHAEFFER 1835)	-	●	●	■	●	●	-
Megalonotus puncticollis (LUCAS 1849)	-	-	-	●	●	●	●
Megalonotus sabulicola (THOMSON 1870)	-	-	■	■	●	-	-
Proderes suberythropus (A. COSTA 1842)	-	●	●	-	●	●	-
Remaudiereana annulipes (BÄRENSPRUNG 1859)	■	●	●	●	●	●	■
* Marmottania simonis PUTON 1887	-	-	-	-	-	-	■
€ Noualhieria herbanica J. RIBES 1976	-	-	-	-	-	●	●
€ Noualhieria pietaini GÓMEZ-MENOR 1924	●	●	●	●	●	-	-
€ Noualhieria quadripunctata (BRULLÉ 1838)	●	-	-	●	●	●	●
* € Noualhieria heissi nov.sp.	■	-	-	-	-	-	-
Plinthius (Plinthius) canariensis WAGNER 1963	●	●	●	●	●	●	-
Beosus maritimus (SCOPOLI 1763)	●	●	●	●	●	●	-
Beosus quadripunctatus (MÜLLER 1766)	-	-	-	-	-	-	-
Dieuches schmitzi REUTER 1893	-	●	■	●	●	●	-
Liolobus pallidicornis (REUTER 1891)	-	-	-	-	-	-	-
Peritrechus gracilicornis PUTON 1877	-	-	●	-	●	-	-
Ragilus alboacuminatus (GOEZE 1778)	-	●	●	●	●	●	-
Rhyparochromus pini (LINNAEUS 1758)	-	-	-	-	-	-	-
Xanthochilus saturnius (ROSSI 1790)	●	●	●	●	●	▲	-
Hyalochilus ovatulus (A. COSTA 1853)	●	●	●	●	●	●	-
€ Stygnocoris barbieri PENICART 1993	□	●	●	●	●	●	-
Stygnocoris fuliginosus (GEOFFROY 1785)	-	-	-	-	-	■	-
€ Stygnocoris subglaber (PUTON 1889)	●	●	●	●	●	●	●
Stygnocoris truncatus (HORVATH 1893)	●	-	-	-	-	-	-
Piesmatidae							
Parapiesma rotundatum (HORVATH 1906)	-	-	-	-	-	-	●
Berytidae							
Neidus aduncus FIEBER 1859	●	●	●	●	●	●	-
€ Neides gomeranus HEISS 1978	-	-	-	●	-	-	-
Berytinus (Berytinus) hirticornis pilipes (PUTON 1875)	●	●	●	●	●	-	-
Berytinus (Lizinus) distinguendus (FERRARI 1874)	-	-	-	-	-	-	-
Berytinus (Lizinus) geniculatus (HORVATH 1885)	■	-	-	●	-	-	●
Berytinus (Lizinus) montivagus (MEYER-DÜR 1841)	●	●	●	●	●	■	●
€ Gampsocoris gomeranus WAGNER 1965	-	-	-	●	-	-	-
Gampsocoris punctipes punctipes (GERMAR 1822)	●	-	-	-	●	●	-
Pyrrhocoridae							
Scantius aegyptius aegyptius (LINNAEUS 1758)	●	●	●	●	●	●	●
Stenocephalidae							
Dicranoccephalus agilis (SCOPOLI 1763)	●	□	■	●	●	●	●

	H	P	G	T	C	F	L
<i>Dicranocephalus insularis</i> (DALLAS 1852) (= <i>bianchii</i> JAKOVLEV 1901)	-	-	-	-	●	●	-
<i>Dicranocephalus pallidus</i> (SIGNORET 1879) (= <i>berlandi</i> VILLIERS 1951)	-	□	-	●	-	■	-
Rhopalidae							
<i>Brachycarenum tigrinus</i> (SCHILLING 1829)	●	●	●	●	●	-	■
<i>Corizus hyoscyami nigridorsum</i> (PUTON 1874)	●	●	●	●	●	-	●
<i>Liothyssus hyalinus</i> (FABRICIUS 1794)	▲	●	●	●	●	●	●
<i>Rhopalus parumpunctatus</i> SCHILLING 1829	-	-	-	-	-	-	-
<i>Stictopleurus pictus</i> (FIEBER 1861)	-	●	●	●	●	●	-
<i>Stictopleurus ribesi</i> GÖLLNER-SCHIEDING 1975	●	□	-	●	●	-	-
<i>Stictopleurus subtomentosus</i> (REY 1888)	-	-	-	-	-	-	-
<i>Stictopleurus synavei</i> GÖLLNER-SCHIEDING 1975	-	-	-	●	-	-	-
* <i>Agraphopus suturalis</i> REUTER 1900	-	■	-	-	-	■	-
Alydidae							
<i>Camptopus lateralis</i> (GERMAR 1817)	●	●	●	●	●	●	-
* <i>Euthetus humilis</i> HORVÁTH 1907	-	-	-	-	-	-	■
<i>Nemausus sordidatus</i> (STÅL 1858) (= <i>N. simplex</i> HORVÁTH 1911)	-	-	-	-	-	●	●
* <i>Tenosius proletrarius</i> (SCHAUM 1853)	-	-	-	-	-	■	■
Coreidae							
<i>Arenocoris failenii</i> (SCHILLING 1829)	-	-	-	-	-	■	-
* <i>Arenocoris intermedius</i> (JAKOVLEV 1883)	-	-	-	-	-	■	-
<i>Arenocoris waltii</i> (HERRICH-SCHAEFFER 1835)	●	●	●	●	●	●	●
<i>Ceraleptus obtusus</i> (BRULLÉ 1839)	●	●	●	●	-	-	-
<i>Coriomeris affinis</i> (HERRICH-SCHAEFFER 1839)	●	●	●	●	●	●	●
<i>Leptoglossus gonagra</i> (FABRICIUS 1775)	-	-	-	●	●	-	-
<i>Centrocoris variegatus</i> KOLENATI 1845	□	-	●	●	●	-	-
‡ <i>Cercinthus elegans</i> (BRULLÉ 1839)	-	-	-	-	-	-	-
<i>Cercinthus lehmanni</i> (KOLENATI 1857)	-	-	■	●	●	●	●
‡ <i>Haploprocta incognita</i> HEISS & MOULET 1994	-	-	-	-	-	-	-
<i>Haploprocta sulcicornis</i> (FABRICIUS 1794)	●	●	●	●	●	●	●
<i>Syromastus rhombeus</i> (LINNAEUS 1767)	●	●	●	●	●	●	-
<i>Gonocerus insidiator</i> (FABRICIUS 1787)	-	-	-	●	-	-	-
<i>Gonocerus juniperi</i> HERRICH-SCHAEFFER 1839	-	□	-	●	●	-	-
<i>Plinactus imitator</i> (REUTER 1891)	●	-	-	-	-	-	-
* <i>Phylomorpha laciniata</i> (VILLERS 1789)	-	-	-	-	-	■	-
<i>Priotonylus brevicornis</i> (MULSANT & REY 1852)	-	-	-	-	-	●	-
Cydnidae							
<i>Cydnus aterrimus</i> (FORSTER 1771)	●	●	-	●	●	●	●
<i>Alonips intrusus</i> (WAGNER 1953)	-	-	-	●	-	■	-
<i>Byrsinus albipennis</i> (A. COSTA 1853)	-	-	-	-	●	▲	-
<i>Byrsinus laticollis</i> (WAGNER 1954)	-	-	-	-	●	●	●
<i>Byrsinus pallidus</i> (PUTON 1887)	-	-	-	-	●	-	-
<i>Byrsinus pilosulus</i> (KLUG 1845)	□	●	■	●	●	●	●
<i>Geotomus elongatus</i> (HERRICH-SCHAEFFER 1840)	-	-	-	●	■	●	■
<i>Geotomus longicornis</i> WAGNER 1953	-	-	-	-	-	-	-
<i>Geotomus punctulatus</i> (A. COSTA 1847)	-	-	-	-	-	-	-
<i>Macrocytus brunneus</i> (FABRICIUS 1803)	●	●	●	●	●	●	●
<i>Microporus nigrita</i> (FABRICIUS 1794)	-	□	■	●	●	-	-
<i>Amaurocoris curtus</i> (BRULLÉ 1839)	-	-	-	●	●	●	●
<i>Linospa orbicularis</i> (JAKOVLEV 1885)	-	-	-	-	■	●	-
<i>Canthophorus melanopterus melanopterus</i> (HERRICH-SCHAEFFER 1835)	-	-	-	■	-	-	●
<i>Crocistethus aeneus</i> (BRULLÉ 1839)	-	●	-	●	●	●	●
<i>Ochetostethus pygmaeus</i> (RAMBUR 1839)	-	●	●	●	●	-	-
<i>Ochetostethus tarsalis</i> (MULSANT & REY 1852)	-	-	-	■	-	-	●
<i>Sehirus morio</i> (LINNAEUS 1761)	-	-	-	-	●	-	-
Acanthosomatidae							
<i>Cyphostethus tristriatus</i> (FABRICIUS)	■	□	●	●	-	-	-
Scutelleridae							
<i>Odontotarsus caudatus</i> (BURMEISTER 1835)	●	●	●	●	●	●	●
<i>Odontoscelis dorsalis</i> (FABRICIUS 1798)	●	●	●	●	●	■	●
<i>Pascata granulata</i> (A. COSTA 1847) (= <i>marmottani</i> PUTON 1887)	-	-	-	-	●	●	●
Pentatomidae							
<i>Dryadocoris apicalis</i> (HERRICH-SCHAEFFER 1842) (= <i>analis</i> A. COSTA 1847)	-	-	-	●	-	-	-
<i>Holcogaster fibulata</i> (GERMAR 1831)	●	●	●	●	●	-	-
<i>Brachynema cinctum</i> (FABRICIUS 1775)	-	-	-	●	●	●	●
<i>Brachynema germarii</i> (KOLENATI 1846)	-	-	-	-	●	-	-
<i>Chroantha ornata</i> (HERRICH-SCHAEFFER 1842)	-	□	■	●	●	●	●
<i>Codophila varia varia</i> (FABRICIUS 1787)	●	●	●	●	●	●	●
<i>Dolycoris numidicus</i> HORVÁTH 1907	-	●	●	●	-	-	-
<i>Eudolycoris alluaudi</i> (NOUALHIER 1893)	●	●	●	●	●	-	-
<i>Eysarcoris ventralis</i> (WESTWOOD 1837)	-	●	●	●	●	●	■
<i>Mecidea lindbergi</i> WAGNER 1954	■	●	●	●	●	●	●
<i>Mecidea pallidissima</i> JENSEN-HAARUP 1922	-	■	-	●	-	■	■
<i>Acrosternum heegeri</i> FIEBER 1861	-	-	-	-	●	-	-
* <i>Acrosternum millierei</i> (MULSANT & REY 1866)	-	-	-	■	-	-	-
‡ <i>Acrosternum rubescens</i> (NOUALHIER 1893)	■	●	●	●	●	■	●
<i>Nezara viridula</i> (LINNAEUS 1758)	●	●	●	●	●	●	●
<i>Piezodorus punctipes</i> PUTON 1889	●	●	●	●	●	-	-
<i>Piezodorus teretipes</i> (STÅL 1865)	-	-	-	-	■	-	●

	H	P	G	T	C	F	L
<i>Menacarus dohrnianus</i> (MULSANT & REY 1866) (= <i>ovalis</i> PUTON 1873)	-	-	-	-	●	■	-
<i>Sciocoris</i> (<i>Neosciocoris</i>) <i>conspurcatus</i> <i>majusculus</i> LINNAVUORI 1964	-	-	-	-	-	●	-
<i>Sciocoris</i> (<i>Neosciocoris</i>) <i>maculatus</i> FIEBER 1851	-	■	-	-	-	●	-
<i>Sciocoris</i> (<i>Neosciocoris</i>) <i>pallens</i> KLUG 1845 (= <i>fissus</i> MULSANT & REY 1866)	-	-	-	●	●	-	●
<i>Sciocoris</i> (<i>Neosciocoris</i>) <i>sideritidis</i> WOLLASTON 1858	●	●	●	●	●	●	-
<i>Sciocoris</i> (<i>Parasciocoris</i>) <i>canariensis</i> LINDBERG 1953	■	●	●	●	●	●	●
<i>Sciocoris</i> (<i>Sciocoris</i>) <i>angularis</i> PUTON 1889	●	●	●	●	●	-	●
<i>Sciocoris</i> (<i>Sciocoris</i>) <i>helfferii</i> FIEBER 1851	-	●	-	●	-	●	-
<i>Eurydema</i> (<i>Eurydema</i>) <i>lundbladi</i> LINDBERG 1960	-	●	-	●	-	-	-
<i>Eurydema</i> (<i>Eurydema</i>) <i>ornata</i> (LINNAEUS 1758)	●	●	●	●	●	-	-
385 species/subspecies	127	186	169	289	235	165	137
94 Canarian endemics	34	46	42	71	53	20	20
% Canarian endemics	26.8	24.7	24.9	24.6	22.6	12.1	14.6
25 island endemics	3	2	2	12	4	2	1
% island endemics	2.4	1.1	1.2	4.2	1.7	1.2	0.7

References

- AUKEMA B. & C. RIEGER (Eds) (1995): Catalogue of the Heteroptera of the Palaearctic Region. Vol. **1**. — Netherlands Entomol. Soc., Amsterdam: i-xxvi, 1-222.
- AUKEMA B. & C. RIEGER (Eds) (1996): Catalogue of the Heteroptera of the Palaearctic Region. Vol. **2**. — Netherlands Entomol. Soc., Amsterdam: i-xiv, 1-361.
- AUKEMA B. & C. RIEGER (Eds) (1999): Catalogue of the Heteroptera of the Palaearctic Region. Vol. **3**. — Netherlands Entomol. Soc., Amsterdam: i-xiv, 1-577.
- AUKEMA B. & C. RIEGER (Eds) (2001): Catalogue of the Heteroptera of the Palaearctic Region. Vol. **4**. — Netherlands Entomol. Soc., Amsterdam: i-xiv, 1-346.
- AUKEMA B. & C. RIEGER (Eds) (2006): Catalogue of the Heteroptera of the Palaearctic Region. Vol. **5**. — Netherlands Entomol. Soc., Amsterdam: i-xiii, 1-550.
- BAENA M. & M. BÁEZ (1990): Los Heterópteros acuáticos de las Islas Canarias (Heteroptera, Nepomorpha, Gerromorpha). — *Vieraea* **19**: 233-244.
- BAENA M. & M. MORALES (1993): Contribución al estudio de los Tingidos canarios (Heteroptera: Tingidae). — *Vieraea* **22**: 127-132.
- BÁEZ M. & N. ZURITA (2001): Hemiptera Heteroptera. — In: IZQUIERDO I., MARTIN J.L., ZURITA N. & M. ARECHAVALA (Eds), Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres). Consejería de Política Territorial y Medio Ambiente Gobierno de Canarias: 182-197.
- BÁEZ M., MARTIN E. & N. ZURITA (2004): Hemiptera Heteroptera. — In: IZQUIERDO I., MARTIN J.L., ZURITA N. & M. ARECHAVALA (Eds), Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres). Consejería de Medio Ambiente y Ordenación Territorial Gobierno de Canarias: 192-207.
- CARNERO HERNÁNDEZ A., PÉREZ PADRÓN F. & G. PÉREZ GUERRA (1991): Una aproximación a las plagas de los cultivos en las Islas Canarias. — In: Homnaje al profesor Dr. Telesforo Bravo I. Secretariado de Publicaciones, Universidad de La Laguna: 125-160.
- DERJANSCHI V.V. & J. PÉRICART (2005): Hémiptères Pentatomoidea euro-méditerranéens 1. Généralités. Systématique: Première Partie. — *Faune de France* **90**: 1-494.
- DOLLING W.R. (2006): Superfamily Coreoidea LEACH, 1815. — In: AUKEMA B. & C. RIEGER (Eds), Catalogue of the Heteroptera of the Palaearctic Region. Vol. **5**. Netherlands Entomol. Soc., Amsterdam: 1-101.
- GÖLLNER-SCHIEDING U. (1975): Revision der Gattung *Stictopleurus* STÅL, 1872 (Heteroptera, Rhopalidae). — *Dtsch. Entomol. Z. (N.F.)* **22**: 1-60.
- GÖLLNER-SCHIEDING U. (1990): Revision der Gattung *Odontotarsus* LAPORTE DE CASTELNAU, 1832 (Heteroptera: Scutelleridae). — *Mitt. Zool. Mus. Berlin* **66**: 333-370.

- HEISS E. (1997): Nachtrag zur Heteropterenfauna der Kanarischen Inseln V (Insecta, Heteroptera). — Ber. nat.-med. Verein Innsbruck **84**: 359-369.
- HEISS E. & M. BÁEZ (1990): A preliminar catalog of the Heteroptera of the Canary Islands. — Vieraea **18**: 281-315.
- HEISS E. & J. RIBES (1992): Additions to the Heteroptera-fauna of the Canary Islands I. — Boletim do Museu Municipal do Funchal **44**: 77-102.
- HEISS E. & J.H. WOULDSTRA (1993): Additions to the Heteroptera-fauna of the Canary Islands II. — Boletim do Museu Municipal do Funchal **45**: 67-81.
- HEISS E., HEYDEN T.V.D., RIBES J. & C. RIEGER (1996): Nachtrag zur Heteropterenfauna der Kanarischen Inseln IV. (Insecta, Heteroptera). — Linzer biol. Beitr. **28** (2): 1117-1148.
- IZQUIERDO I., MARTIN J.L., ZURITA N. & M. ARECHAVALETA (Eds) (2001): Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres). — Consejería de Política Territorial y Medio Ambiente Gobierno de Canarias: 1-437.
- IZQUIERDO I., MARTIN J.L., ZURITA N. & M. ARECHAVALETA (Eds) (2004): Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres). — Consejería de Medio Ambiente y Ordenación Territorial Gobierno de Canarias: 1-500.
- KERZHNER I.M. (2001): Family Pyrrhocoridae AMYOT & SERVILLE, 1843. — In: AUKEMA B. & C. RIEGER (Eds), Catalogue of the Heteroptera of the Palaearctic Region. Vol. **4**. Netherlands Entomol. Soc., Amsterdam: 248-258.
- KERZHNER I. M. & M. JOSIFOV (1999): Family Miridae HAHN, 1833. — In: AUKEMA B. & C. RIEGER (Eds), Catalogue of the Heteroptera of the Palaearctic Region. Vol. **3**. Netherlands Entomol. Soc., Amsterdam: 1-576.
- LINDBERG H. (1953): Hemiptera Insularum Canariensium (Systematik, Ökologie und Verbreitung der Kanarischen Heteropteren und Cicadinen. — Comm. Biol. **14** (1): 1-304.
- LINDBERG H. (1960): Supplementum Hemiptera Insularum Canariensium. — Comm. Biol. **22** (6): 1-20.
- LINDBERG H. & E. WAGNER (1965): Supplementum secundum ad cognitionem Hemipterorum Insularum Canariensium. — Comm. Biol. **28** (10): 1-14.
- LINNAVUORI R.E. (1987): Alydidae, Stenocephalidae and Rhopalidae of West and Central Africa. — Acta Entomol. Fenn. **49**: 1-36.
- LIS J. (1999): Burrower bugs of the Old World – a catalogue (Hemiptera: Heteroptera: Cydnidae). — Genus **10**: 165-249.
- MARTIN J.L. (1992): Caracterización ecológica y evolución de las comunidades subterráneas en las islas de Tenerife, El Hierro, y La Palma. Thesis Universidad de Laguna: 1-342.
- MOULET P. (1995): Hémiptères Coreoidea (Coreidae, Rhopalidae, Alydidae), Pyrrhocoridae, Stenocephalidae euro-méditerranéens. — Faune de France **81**: 1-336.
- MOULET P. & D. PLUOT-SIGWALT (1998): Une espèce mythique, *Cercinthus elegans* (BRULLÉ, 1839): redescription du type (Heteroptera, Coreidae). — Revue fr. Ent. (N.S.) **20**: 109-114.
- NILSSON A.N., MALMQUIST B., BÁEZ M., BLACKBURN J.H. & P. ARMITAGE (1998): Stream insects and gastropods in the island of Gran Canaria. — Ann. Limnol. **34**: 413-435.
- NOUALHIER M. (1893): Voyage de M. Ch. Alluaud aux îles Canaries (Novembre 1889-Juin 1890). 2^e Mémoire. Hémiptères Gymnocérates & Hydrocorises. — Ann. Soc. Entomol. Fr. **62**: 5-18.
- OROMÍ P., ZURITA N., ARECHAVALETA M. & A. CAMACHO (2002): Fauna de invertebrados del Parque Nacional del Teide. — Organismo Autónomo Parques Nacionales: 1-421.
- PÉREZ A.J., MORALES E., OROMÍ P. & H. LÓPEZ (2003): Fauna de artrópodos de Montaña Clara (islas Canarias) II: Hexápodos (no coleópteros). — Vierea **31**: 237-251.
- PÉRICART J. (1999): Hémiptères Lygaeidae euro-méditerranéens. 1. — Faune de France **84A**: i-xx, 1-468.
- PÉRICART J. (2001): Family Lygaeidae SCHILLING, 1829 – seed-bugs. — In: AUKEMA B. & C. RIEGER (Eds), Catalogue of the Heteroptera of the Palaearctic Region. Vol. **4**. Netherlands Entomol. Soc., Amsterdam: 35-220.
- PUTSHKOV P.V. & V.G. PUTSHKOV (1996): Family Reduviidae LATREILLE, 1807 – assassin bugs. — In: AUKEMA B. & C. RIEGER (Eds), Catalogue of the Heteroptera of the Palaearctic Region. Vol. **2**. Netherlands Entomol. Soc., Amsterdam: 148-265.
- RIBES J. & E. HEISS (2001): On Ibero-Canarian *Phytocoris* subgen. *Compsocercocoris* with description of a new species from Canary Islands (Heteroptera, Miridae). — Linzer biol. Beitr. **33**: 625-636.
- RIBES J. & E. RIBES (1997): Adiciones a los heterópteros de las Islas Canarias III. — Ses. Entom. ICHN-SCL **9** (1995): 161-174.
- RIDER D.A. (2006): Family Pentatomidae LEACH, 1815. — In: AUKEMA B. & C. RIEGER (Eds), Catalogue of the Heteroptera of the Palaearctic Region. Vol. **5**. Netherlands Entomol. Soc., Amsterdam: 232-403.

Addresses of the Authors:

Dr. Berend AUKEMA & Dr. J.P. DUFFELS
 Zoölogisch Museum
 Afdeling Entomologie
 Plantage Middenlaan 64
 1018 DH Amsterdam, The Netherlands
 E-Mail: b.aukema@freeler.nl
 duffels@science.uva.nl

Dr. Manuel BÁEZ
 Departamento de Biología
 Universidad de La Laguna
 38200 Tenerife, Islas Canarias, Spain
 E-Mail: marbaez@ull.es

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Denisia](#)

Jahr/Year: 2006

Band/Volume: [0019](#)

Autor(en)/Author(s): Aukema Berend, Duffels Johannes P., Baez Manuel

Artikel/Article: [A Checklist of the Heteroptera of the Canary Islands \(Insecta\) 755-774](#)