

## Research article

urn:lsid:zoobank.org:pub:A519A06A-6C20-45A7-9120-B0C795E41E75

# Holocene ostracods (Crustacea) from a whale-fall excavation site from the Chao Phraya delta, Central Thailand

Anisong CHITNARIN<sup>1,\*</sup>, Marie-Béatrice FOREL<sup>2</sup> & Prachya TEPNARONG<sup>3</sup>

<sup>1,3</sup>School of Geotechnology, Institute of Engineering, Suranaree University of Technology, Mueang District, Nakhon Ratchasima, 30000, Thailand.

<sup>2</sup>Muséum national d'histoire naturelle, CR2P, MNHN-CNRS-SU, 75005 Paris, France.

\* Corresponding author: [anisong@sut.ac.th](mailto:anisong@sut.ac.th)

<sup>2</sup> Email: [marie-beatrice.forel@mnhn.fr](mailto:marie-beatrice.forel@mnhn.fr)

<sup>3</sup> Email: [prachya@sut-ac.th](mailto:prachya@sut-ac.th)

<sup>1</sup> urn:lsid:zoobank.org:author:804F4F3D-F368-4EA7-9D7C-7E0530A5FE42

<sup>2</sup> urn:lsid:zoobank.org:author:063C1F7E-6D26-48F9-A8B1-2AFA496B5FB8

<sup>3</sup> urn:lsid:zoobank.org:author:3ABB571B-791D-480C-B053-89B59FEB21CA

**Abstract.** Late Holocene ostracods were recovered from marine sediments of the Chao Phraya delta at a whale-fall excavation site located fifteen kilometers on land in the Am Pang Subdistrict, Ban Paew District, Samut Sakhon Province, north of the Gulf of Thailand. Thirteen species belonging to seven genera are identified. The deposition environment of the succession is for the first time characterized. The ostracod assemblages suggest that the entire succession associated with the whale-fall deposited in a shallow marine environment such as estuary, bay, inner shelf, subtidal, under less than 20 meters water-depth, in brackish to normal salinity with high mud content and turbidity, on a muddy substrate. This analysis is an important step toward the first in-depth study of ostracods associated with modern and fossil shallow-water whale-falls.

**Keywords.** Ostracods, shallow-water whale-fall, Holocene, Bangkok Clay, Chao Phraya delta.

Chitnarin A., Forel M.-B. & Tepnarong P. 2023. Holocene ostracods (Crustacea) from a whale-fall excavation site from the Chao Phraya delta, Central Thailand. *European Journal of Taxonomy* 856: 120–151.  
<https://doi.org/10.5852/ejt.2023.856.2033>

## Introduction

The discovery of a large whale skeleton in a privately excavated pit in Samut Sakhon Province in November 2020 raised questions about how this enormous animal deposited far away from the sea. The site was located about 15 kilometers north of the Gulf of Thailand shoreline and adjacent areas were covered by agricultural area and villages. The land owner reported the discovery to the Royal Thai Department of Mineral Resources and allowed geologists and their teams to perform detailed investigations and samplings. The first part of the skeleton that eroded out of the unconsolidated sediments at the bottom of the pit was caudal vertebrae, later 142 pieces in total were excavated and conserved (Fig. 1). The whole

12.5-meter-long skeleton has been identified as *Balaenoptera edeni* Anderson, 1879, and radiocarbon dating provided the age of  $3380 \pm 30$  years BP (Kawira & Saethien 2021; Saethien 2021).

The sediments making up the pit wall are a part of the Holocene marine sediments of the Chao Phraya delta, also known as Bangkok Clay (Sinsakul 2000). The Holocene sediments unconformably overlay the Late Pleistocene Bangkok Stiff Clay (Rau & Nutalaya 1983). The Holocene sediments, including transgressive peaty (mangrove swamp) sediments in the lower part and regressive deltaic sediments in the upper part, were deposited in the paleo-Gulf of Ayutthaya. Sea water invaded northward as far as 100 km around 8000–7000 years BP, then the sea water retreated and delta and shoreline prograded southward forming the present delta plain and shoreline (Rau & Nutalaya 1983; Sinsakul 2000; Woodroffe 2000). Lithostratigraphy, paleontology and evolution of the Holocene deposits have been discussed elsewhere and the reader is referred to the following references for details (Somboon 1988; Somboon & Thiramongkol 1992; Sinsakul 2000; Songtham *et al.* 2000; Woodroffe 2000; Tanabe *et al.* 2003). Fossils of vertebrates, invertebrates as well as palynological materials embedded in the Holocene sediments provide evidence of the great variety of depositional environments within the Chao Phraya delta: from infralittoral, tidal flat, intertidal, mangrove swamps to shallow sea (8 to 10 meters deep) according to mollusk communities and crabs (Robba *et al.* 1993; Songtham *et al.* 2000; Negri 2009), from flood plain, transitional swamp, back swamp, mangrove forest to paleo gulf environments according to palynology (Hutangkura 2012; Songtham *et al.* 2015).

In this study, we examine ostracods recovered from silty clays collected from the whale-fall excavation site in order to better constrain the environment of deposition. At present, studies of Recent marine



**Fig. 1.** Whale-fall excavation site, fifteen kilometers away from recent shoreline in Samut Sakhon Province, north of the Gulf of Thailand. **A.** First appearance of large skeletons eroded off the pit wall. **B.** The excavation during November–December 2020. **C.** Nearly complete, well preserved whale skeleton. All photos from ThaiWhales ([https://www.facebook.com/thaiwhales/photos/?ref=page\\_internal](https://www.facebook.com/thaiwhales/photos/?ref=page_internal)).

ostracods along the coasts of Thailand are rare (Montenegro *et al.* 2004; Yamada *et al.* 2014; Forel 2021). Here, we provide taxonomic information on the Holocene marine ostracods and further discuss the change in shallow-water ostracod assemblages of the northern Gulf of Thailand. Whale-falls are rarely encountered on the sea-floor and associated ostracods remain practically unknown as they are in most cases reported without taxonomic details (e.g., Furushima *et al.* 2007; Danise *et al.* 2014). This contribution is therefore an important step in the knowledge of ostracods associated with shallow-water whale-falls. It is important to note that only one of the 10 samples studied here is co-eval with the whale deposition. The entire succession provides the baseline of the background ostracod assemblages in this area during the Holocene, that will be of crucial significance to understand the ostracod communities directly associated with and dwelling around the whale-fall in an ongoing in-depth study.

## Material and methods

The sediment samples were collected from the pit which is situated approximately 10 kilometers northwest of Samut Sakhon city or 15 kilometers north of the shoreline (Fig. 2A–B). This area is located in the Recent delta plain of central Thailand, very close to the Tha Chin River. The lithostratigraphy of the area around the whale excavation consists of four sedimentary units, in ascending order (Fig. 3A): Pleistocene Stiff Clay found at 12.5 meters from ground surface; Shallow Marine Clay, SMC found at 3.5 meters depth (10–10.5 meters thick); Old Tidal Flat sediments, OTF found at 1 meter depth (2–2.5 meters thick); Topsoil about 1 meter thick (Nuammim *et al.* 2021). The whale skeleton was embedded in the SMC unit, also known as the Bangkok Clay. The excavation site was on the eastern wall of a large pit developed for surface water reservoirs (Fig. 2C). The land surface was at 5 meters below mean sea level, the whale skeleton was exposed at about 1.5 meters depth, thus the whale was deposited at 6.5 meters below mean sea level. The pit was dug deeper than the skeleton to allow paleontological preparation. The lowest level of the pit was at 7 meters below mean sea level (Figs 2C, 3B).

Ten unconsolidated samples, labelled 20SS01A to 20SS06, were collected from the pit wall just below and above the skeleton level at regular intervals of 25 centimeters (Figs 3B, 4). The sediments were dense, greenish gray silty mudstone with small shell fragments. The 200-gram samples of wet clay were washed through a 125-micron sieve and dried. Seven of the ten samples yielded ostracods (Fig. 4). Ostracod specimens were picked and initially identified under a stereoscopic microscope. The specimens then were viewed using a JEOL/JSM-6010LV scanning electron microscope at the Suranaree University of Technology (Nakhon Ratchasima, Thailand) for detailed identification.

The ostracod classification in this study follows the general classification given in the *Treatise on Invertebrate Paleontology, Part Q, Ostracoda* (Moore 1961), Hartmann & Puri (1974) and Martens & Horne (2009). Thirteen species belonging to seven genera are identified (Fig. 4). Selected specimens of genera *Propontocypris* Sylvester-Bradley, 1947 (Pontocyprididae Müller, 1894), *Aglaioocypris* Sylvester-Bradley, 1947 (Candonidae Kaufmann, 1900), *Sinocytheridea* Hou, 1982 (Cytherideidae Sars, 1925) are shown in Fig. 5. Members of Schizocytheridae Howe, 1961 (*Neomonoceratina* Kingma, 1948) are shown in Fig. 6. Selected specimens of genera *Stigmatocythere* Siddiqui, 1971, *Keijella* Ruggieri, 1967 and *Pistocythereis* Gou, 1983 (Trachyleberididae Sylvester-Bradley, 1948) are shown in Fig. 7. Series of juvenile and adult specimens of *Keijella multisulcus* Whatley & Zhao, 1988 and *K. gonia* Zhao & Whatley, 1989 are shown in Figs 8–10.


## Material repository

All specimens are stored in the Micropaleontology Collections of Suranaree University of Technology, Nakhon Ratchasima, Thailand, under collection numbers SUT-20SS-C001 to SUT-20SS-C350.




**Anatomical abbreviations**

- AB = anterior border
- ADB = anterodorsal border
- AVB = anteroventral border
- DB = dorsal border
- H = height
- Hmax = maximum of height
- L = length
- Lmax = maximum of length
- LV = left valve
- midH = mid-height
- midL = mid-length




**Fig. 2.** Map of the studied section. **A–B.** Position of the study area at the northern coast of the Gulf of Thailand. **C.** Location of the whale-fall excavation site in Samut Sakhon Province (Google Map, 2021).





**Fig. 3.** Lithostratigraphy of the studied section at whale-fall excavation site in Samut Sakhon Province. **A.** Lithologic log modified after Nuamnim *et al.* (2021). **B.** Lithologic log of the studied section (this study).

- PB = posterior border
- PDB = posterodorsal border
- PVB = posteroventral border
- RV = right valve
- VB = ventral border



**Fig. 4.** Distribution of Holocene ostracods below and above the whale skeleton level at the whale-fall excavation site in Samut Sakhon Province, north of Gulf of Thailand.



## Results

### *Taxonomic descriptions*

Class Ostracoda Latreille, 1806  
Subclass Podocopa Müller, 1894  
Order Podocopida Müller 1894  
Sub-order Podocopina Sars, 1866  
Superfamily Cypridoidea Baird, 1845  
Family Candonidae Kaufmann, 1900

Genus *Aglaiocypris* Sylvester-Bradley, 1947

### Type species

*Aglaiocypris pulchella* (Brady, 1868) subsequently designated by Sylvester-Bradley (1947).

*Aglaiocypris pellucida* Mostafawi, 2003  
Fig. 5A–C

*Aglaiocypris pellucida* Mostafawi, 2003: 71, figs 51a–c.

*Aglaiocypris?* sp. 5 – Maddocks 1969: 7, figs 31r–t.

*Aglaiocypris* sp. – Paik 1977: 28: 42, pl. 7 figs 3, 141–143, pl. 10 fig. 181.

*Bythocypris* sp. A. – Jain 1978: 94, fig. 2d.

*Aglaiocypris pellucida* – Wang *et al.* 2018: 324, figs 2l–o.

### Dimensions

L = 0.542–0.569 millimeters; H = 0.250–0.275 millimeters; H/L = 0.45–0.49.

### Distribution

Modern distribution: West coast of India, Recent (Jain 1978); Persian Gulf, Recent (Maddocks 1969; Paik 1977; Mostafawi 2003)

Fossil distribution: Hang Hau Formation, Lei Yue Mun, Hong Kong, Holocene (Wang *et al.* 2018); Bangkok Clay (samples 20SS01A, 02A, 02B, 03A), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

### Remarks

*Aglaiocypris pellucida* Mostafawi, 2003 is diagnosed by the following characters: thin, flat and obtuse carapace with triangular lateral outline, largely rounded AB and PB, Hmax located centrally, DB convex at LV, slightly angulated at RV, moderate overlapping of RV on LV all around, VB straight at RV, concave at LV, numerous normal pores. The present record is the first occurrence of *Aglaiocypris pellucida* in Thailand as, until now, this species was only known from the Persian Gulf (Maddocks 1969; Paik 1977; Mostafawi 2003), western coast of India (Jain 1978) and in Holocene deposits of Hong Kong (Wang *et al.* 2018). The size of the examined specimens is similar to the Holocene specimens recovered in Hong Kong (Wang *et al.* 2018) but they are smaller than the extant type material from the Persian Gulf (Mostafawi 2003).

Superfamily Pontocypridoidea Müller, 1894  
Family Pontocyprididae Müller, 1894

Genus *Propontocypris* Sylvester-Bradley, 1947

### Type species

*Pontocypris trigonella* Sars, 1866 by original designation.

*Propontocypris bengalensis* Maddocks, 1969  
Fig. 5D–F

*Propontocypris (Schedopontocypris) bengalensis* Maddocks, 1969: 34, fig. 31a, c, f.

*Pontocypris bengalensis* – Mostafawi 2003: 71, fig. 48a–b. — Wang *et al.* 2018: 325, fig. 2h–i.

*Pontocypris* sp. A. – Bate 1971: 246, pl. 1 figs 1g, 2g. — Bonaduce *et al.* 1983: pl. 5 figs 7–8.

*Propontocypris* sp. C. – Paik 1977: 42, pl. 7 fig. 140, pl. 10 fig. 179.

### Dimensions

L = 0.467–0.493 millimeters; H = 0.213–0.246 millimeters; H/L = 0.44–0.51.

### Distribution

Modern distribution: Persian Gulf (Maddocks 1969; Mostafawi 2003); Bay of Bengal and Sri Lanka (Maddocks 1969); Persian Gulf (Bate 1971; Paik 1977); Red Sea (Bonaduce *et al.* 1983).

Fossil distribution: Hang Hau Formation, Lei Yue Mun, Hong Kong, Holocene (Wang *et al.* 2018); Bangkok Clay (samples 20SS01A, 03A, 04A), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work).

### Remarks

*Propontocypris bengalensis* Maddocks, 1969 is diagnosed by its small compressed carapace, suboval outline, Hmax located anterior to midL, long ABD, round AB with small radius of curvature, straight VB, round PB with large radius of curvature, slight overlapping of LV on LV around the carapace. Carapaces are delicate, usually yellow, white or transparent. The examined specimens are smaller than those from the Holocene of Hong Kong (L = 0.52–0.53 millimeters; H = 0.27–0.28 millimeters) reported by Wang *et al.* (2018).

*Propontocypris clara* Zhao, 1988  
Fig. 5G–H

*Propontocypris clara* Zhao in Wang *et al.* 1988: 230, pl. 36 figs 3–5.

*Propontocypris clara* – Ruan 1989: 118, pl. 20 fig. 24. — Tanaka *et al.* 2009: pl. 1 fig. 1. — Wang *et al.* 2018: 326, fig. 3a–d.


### Dimensions

L = 0.38–0.493 millimeters; H = 0.213–0.246 millimeters; H/L = 0.44–0.51.

### Distribution

Modern distribution: East China Sea (Wang *et al.* 1988); Northern Xisha Trench, South China Sea, China (Ruan 1989); Northeastern coast of Vietnam, Vietnam (Tanaka *et al.* 2009).





**Fig. 5.** Holocene ostracods from Samut Sakhon Province, Central Thailand. **A–C.** *Aglaiocypris pellucida* Mostafawi, 2003. **A.** Carapace, right lateral view, SUT-20SS-C003. **B.** Carapace, left lateral view, SUT-20SS-C007. **C.** Carapace, left lateral view, SUT-20SS-C003. – **D–H.** *Propontocypris bengalensis* Maddocks, 1969. **D.** Carapace, left lateral view, SUT-20SS-C012. **E.** Carapace, left lateral view, SUT-20SS-C020. **F.** Carapace, left lateral view, SUT-20SS-C014. **G.** Carapace, right lateral view, SUT-20SS-C027. **H.** Carapace, left lateral view, SUT-20SS-C022. – **I–O.** *Sinocytheridea impressa* (Brady, 1869). **I.** Carapace, female, right lateral view, SUT-20SS-C063. **J.** Carapace, male, right lateral view, SUT-20SS-C040. **K.** Internal view of right valve, male, SUT-20SS-C049. **L.** Carapace, female, left lateral view, SUT-20SS-C071. **M.** Carapace, male, left lateral view, SUT-20SS-C071. **N.** Internal view of left valve, male, SUT-20SS-C048. **O.** Internal view of left valve, female, SUT-20SS-C064. Scale bars = 0.1. mm.

Fossil distribution: Hang Hau Formation, Lei Yue Mun, Hong Kong, Holocene (Wang *et al.* 2018); Bangkok Clay (samples 20SS01A, 02A, 02B, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

### Remarks

*Propontocypris clara* Zhao, 1988 is characterized by its small and delicate carapace, subtriangular in lateral view, with Hmax located in front of midL, rounded AB, slightly concave VB, narrowly rounded PB, round normal pores. *Propontocypris clara* is distinguished from *P. bengalensis* by the location of Hmax. The examined specimens are smaller than those reported from Holocene sediments of Hong Kong (L = 0.497–0.526 millimeters; H = 0.280–0.274 millimeters; Wang *et al.* 2018).

Suborder Cytherocopina Baird, 1850  
Superfamily Cytheroidea Baird, 1850  
Family Cytherideidae Sars, 1925

Genus *Sinocytheridea* Hou, 1982

### Type species

*Sinocytheridea latiovata* Hou in Hou *et al.*, 1982, junior synonym of *Sinocytheridea impressa* (Brady, 1869) following the revision of Whatley & Zhao (1988a).

*Sinocytheridea impressa* (Brady, 1869)

Fig. 5I–O

*Cytheridea impressa* Brady, 1869: 158, pl. 16 figs 13–14.

*Cyprideis yehi* Hu & Yeh, 1978: 157–159, pl. 3 figs 10–13.

*Sinocytheridea sinensis* Hou in Guan *et al.*, 1978: 240, pl. 65 figs 1–5.

*Sinocytheridea latiovata* Hou & Chen in Hou *et al.*, 1982: 164–165, pl. 72 figs 10–20.

*Sinocytheridea longa* Hou & Chen in Hou *et al.*, 1982: 165, pl. 72 figs 1–9.

*Eucytheridea sinobesani* Hu, 1984: 76, pl.10, figs. 27, 28.

*Sinocytheridea impressa* – Whatley & Zhao 1987: 24, pl. 1 figs 8–10. — Montenegro *et al.* 2004: pl. 2 figs 6–7. — Alberti *et al.* 2013: 341. — Yamada *et al.* 2014: 110. — Hong *et al.* 2017: 58; 2019: 596. — Cheung *et al.* 2019: 9. — Tanaka *et al.* 2019: fig. 7. — Tan *et al.* 2021: fig. 2(10).

### Dimensions

L = 0.515–0.567 millimeters, H = 0.250–0.317 millimeters, H/L = 0.45–0.49 (males; Fig. 5J–K, M–N).

L = 0.477–0.700 millimeters, H = 0.246–0.410 millimeters, H/L = 0.52–0.59 (females; Fig. 5I, L, O).

### Distribution

Tanaka *et al.* (2019) recently reviewed and presented the distribution of extant and fossil *Sinocytheridea impressa* around the eastern margin of Eurasia, from the Sea of Japan to South China Sea. The reader is referred to their work for details. Here, we list some occurrences of the species.

Modern distribution: Japan (Tanaka *et al.* 2019); China (Zhao & Wang 1988, 1990; Cheung *et al.* 2019); Hong Kong (Brady 1869; Hong *et al.* 2017, 2019); Central Vietnam (Tan *et al.* 2021). Mae Khlong estuary, NW Gulf of Thailand, Thailand (Montenegro *et al.* 2004); Klong Thom, Krabi, Andaman estuary, Thailand (Yamada *et al.* 2014).



Fossil distribution: Upper Pliocene, Japan (Yamada *et al.* 2002); Middle Pleistocene, Japan (Ishizaki 1990; Irizuki *et al.* 2005). Pliocene, China (Hou & Gou 2007); Pliocene to Pleistocene, China (Lee & Paik 1992; Hu & Tao 2008); Quaternary, China (Huang 1985; Hou *et al.* 1982; Alberti *et al.* 2013); Pleistocene to Holocene, Hong Kong (Cao, 1998); Bangkok Clay (samples 20SS01A, 01B, 02A, 02B, 03A, 04A, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

### Remarks

Carapaces of *Sinocytheridea impressa* (Brady, 1869) are characterized by flat, elongate to oval lateral outline, long hinge, slightly concave VB, with scattered sieve type pores on carapace surface. Sexual dimorphism clear: males are longer and slender with AB and PB nearly of equal size, Hmax located anteriorly but not distinct; females are shorter, with AB larger than PB, Hmax located anterior to midL and distinct. The juvenile carapaces are very delicate and transparent.

Family Schizocytheridae Howe in Moore, 1961  
Subfamily Schizocytherinae Mandelstam, 1960

Genus *Neomonoceratina* Kingma, 1948

### Type species

*Neomonoceratina columbiformis* Kingma, 1948 by original designation.

### Preliminary remarks

Members of the genus *Neomonoceratina* can be classified into five groups based on the ornamentation of their carapaces (Zhao & Whatley 1988): *N. columbiformis* group, *N. iniqua* group, *N. koeningswaldi* group, *N. macropora* group and *N. spinosa* group. The specimens investigated here belong to the *N. columbiformis* and *N. iniqua* groups. The *N. columbiformis* group is characterized by a smooth or finely punctate intercostal surface and includes *N. columbiformis*, *N. mediterranea* (Ruggieri, 1953), *N. mediterranea malayensis* Zhao & Whatley, 1988. Conversely, the *N. iniqua* group is characterized by a reticulate surface consisting of polygon fossae, thin muri and finely punctate sola, fringed AB with small denticles. In the present material, *N. iniqua* (Brady, 1868), *N. bataviana* (Brady, 1868), *N. delicata* Ishizaki & Kato, 1976, *N. chena* Zhao & Whatley, 1988 are representatives of this *N. iniqua* group.

*Neomonoceratina iniqua* (Brady, 1868)  
Fig. 6A–C

*Cytherura iniqua* Brady 1868: 64, pl. 8 figs. 3–6.

*Trachyleberis? vjetnamica* Schneider, 1971: 261, figs e–f.

*Neomonoceratina diptera* Hu & Yang, 1975: 108, pl. 1 figs 19–20.

*Cytherura iniqua* – Brady 1886: 130, pl. 39 figs 31–33.

*Neomonoceratina* sp. A – Paik 1977: 42, pl. 2 figs 24–28, pl. 8 fig. 148.

*Neomonoceratina iniqua* – Zhao & Whatley 1988: 566, pl. 1 figs 7–12; 1989: 171. — Dewi 1993: 60. — Hussain 1998: 4, pl. 1 fig. 11 — Al-Jumaily & Al-Sheikhly 1999: 217, fig. 15. — Mostafawi 2003: 56, fig. 6. — Montenegro *et al.* 2004: pl. 2 fig. 4. — Gopalakrishna *et al.* 2007: pl. 1 fig. 10. — Fauzielly *et al.* 2012: fig. 3(6); 2013: fig. 6(10). — Baskar *et al.* 2013: fig. 3(4). — Hussain & Kalaiyarasi 2013: fig. 11.2(a–b). — Forel 2021: fig. 4d–e. — Tan *et al.* 2021: fig. 2(7).

### Dimensions

L = 0.542–0.592 millimeters; H = 0.267–0.292 millimeters, H/L = 0.48–0.50.

**Distribution**

Modern distribution: Java Sea (Brady 1868; Dewi 1993, 2000; Fauzielly 2013; Fauzielly *et al.* 2012, 2013); Iraq (Al-Jumaily & Al-Sheikhly 1999); east India (e.g., Hussain & Mohan 2001; Hussain *et al.* 2007); southeast India (e.g., Hussain 1998; Hussain *et al.* 2004, 2007, 2013a; Baskar *et al.* 2013; Hussain & Kalaiyarasi 2013); southwest India (e.g., Hussain *et al.* 2013b; Gopalakrishna *et al.* 2007); west India (e.g., Bhatia & Kumar 1979); Sri Lanka (Iwatani *et al.* 2014); Persian Gulf (Paik 1977; Mostafawi 2003; Mostafawi *et al.* 2010); Malaysia (e.g., Zhao & Whatley 1989; Ramlan & Noraswana 2009, 2010); Vietnam (Tan *et al.* 2021); Mae Khlong river mouth, north west Gulf of Thailand (Montenegro *et al.* 2004); southwestern coast of Peninsular Thailand, Ao Nun, Satun Province, Andaman Sea (Forel 2021);

Fossil distribution: Bangkok Clay (samples 20SS01A, 02A, 02B, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

**Remarks**

*Neomonoceratina iniqua* (Brady, 1868) is characterized by its reticulate surface with thin muri and polygon fossae, indistinct and short posterodoral rib, long median rib from anterior to posteroventral region, and venterolateral rib terminating into a simple spine. Sexual dimorphism is well expressed with longer and slender male carapaces while female carapaces are shorter and higher. This is the first discovery of the species from Holocene sediments.

*Neomonoceratina rhomboidea* (Brady, 1968)

Fig. 6D–I

*Cytheropteron rhomboideum* Brady 1968: 65, pl. 8 figs 10–12.

*Neomonoceratina rhomboidea* – Hanai *et al.* 1980: 154. — Zhao & Whatley 1988: 569, pl. I figs 20–21, pl. II figs 1–3.

**Dimensions**

L = 0.400–0.778 millimeters, H = 0.206–0.467 millimeters, H/L = 0.40–0.67.

**Distribution**

Modern distribution: Batavia, Java, Indonesia (Brady 1968); Jason Bay, southeast Malaysia (Zhao & Whatley 1988).

Fossil distribution: Bangkok Clay (samples 20SS01A, 01B, 02A, 02B, 03A, 04A, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work).

**Remark**

*Neomonoceratina rhomboidea* (Brady, 1968) can be recognized by the inflated carapace, a weak and shallow reticulation, a thin median rib and alalike posteroventral inflation and relatively large, sieve type, normal pore canals.


*Neomonoceratina columbiformis* Kingma, 1948

Fig. 6J

*Neomonoceratina columbiformis* – Kingma 1948: 95, pl. 10 fig. 8a–f.

*Neomonoceratina columbiformis* – Keij 1979: 61, pl. 1 figs 1–4, pl. 2 figs 5–6. — Zhao & Whatley 1988: 565, pl. 1 fig. 1.

non *Neomonoceratina columbiformis* – Keij 1979: 166, pl. 1 fig. 11.



**Fig. 6.** Holocene ostracods from Samut Sakhon Province, Central Thailand. **A–C.** *Neomonoceratina iniqua* (Brady, 1868). **A.** Valve, male, right lateral view, SUT-20SS-C097. **B.** Valve, male, left lateral view, SUT-20SS-C095. **C.** Valve, male, right lateral view, SUT-20SS-C093. – **D–I.** *Neomonoceratina rhomboidei* (Brady, 1968). **D.** Carapace, male, right lateral view, SUT-20SS-C113. **E.** Carapace, male, right lateral view, SUT-20SS-C120. **F.** Carapace, female, right lateral view, SUT-20SS-C099. **G.** Carapace, female, left lateral view, SUT-20SS-C107. **H.** Carapace, female, dorsal view, SUT-20SS-C114. **I.** Sieve pores. – **J.** *Neomonoceratina columbiformis* Kingma, 1948. Carapace, left lateral view, SUT-20SS-C181. – **K–N.** *Neomonoceratina mediterranea mediterranea* (Ruggieri, 1953). **K.** Carapace, female, right lateral view, SUT-20SS-C167. **L.** Carapace, female, right lateral view, SUT-20SS-C173. **M.** Carapace, female, left lateral view, SUT-20SS-C164. **N.** Carapace, female, dorsal view, SUT-20SS-C174. – **O–T.** *Neomonoceratina mediterranea malayensis* Zhao & Whatley, 1988. **O.** Carapace, male, right lateral view, SUT-20SS-C159. **P.** Carapace, male, left lateral view, SUT-20SS-C157. **Q.** Carapace, male, dorsal view, SUT-20SS-C161. **R.** Carapace, male, left lateral view, SUT-20SS-C152. **S.** Carapace, male, left lateral view, SUT-20SS-C141. **T.** Valve, male, internal view of left valve, SUT-20SS-C160. Scale bars = 1 mm.



**Dimensions**

L = 0.440 millimeters, H = 0.240 millimeters, H/L = 0.55.

**Distribution**

Modern distribution: Malacca Strait (Zhao & Whatley 1988); Jakarta Bay, Indonesia (Fauzielly *et al.* 2013); Klong Thom, Krabi, Andaman estuary, Thailand (Yamada *et al.* 2014).

Fossil distribution: Pliocene, Sumatra (Kingma 1948; Keij 1979); Bangkok Clay (sample 20SS04A), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work).

**Remark**

*Neomonoceratina columbiformis* Kingma, 1948 is very rare in our material but it can be recognized by its non-ornamented carapace with curved posterodorsal rib and oblique median rib.

***Neomonoceratina mediterranea mediterranea* (Ruggieri, 1953)**

Fig. 6K–N

*Paijenborchella (Neomonoceratina) mediterranea* Ruggieri, 1953: 4–7, figs 1–5.

*Paijenborchella (Neomonoceratina) mediterranea* – Keij 1954: 288, pl. 5 fig. 15, pl. 16 fig. 12, 361, pl. 3 figs 12–13. — Morales 1966: 80, pl. 7 fig. 2a–c.

*Neomonoceratina* sp. Swain 1955: 643, pl. 64 fig. 14.

*Neomonoceratina mediterranea* – Morkhoven Van 1963: 369, fig. 604. — Teeter 1975: 473, fig. 17k. — Gou *et al.* 1981: 171, pl. 82 figs 11–12. — Hou *et al.* 1982: 219, pl. 80 figs 24–30. — McKenzie & Pickett 1984: fig. 4y–z.

*Neomonoceratina mediterranea mediterranea* – Zhao & Whatley 1988: 565, pl. 1 figs 2–3.

*Neomonoceratina* sp. (pars) – Forel 2021: 7, fig. 4g–h

**Dimensions**

L = 0.400–0.778 millimeters; H = 0.206–0.467 millimeters; H/L = 0.46–0.60.

**Distribution**

Modern distribution: Southwestern coast of Peninsular Thailand, Ao Nun, Satun Province, Andaman Sea (Forel 2021).

Fossil distribution: Pliocene of southeast China, Quaternary of east China, East of Australia, Recent of Eastern Mediterranean, the Philippines, Indonesia, Australia, the Caribbean and Gulf of Mexico (see details in Zhao & Whatley 1989); Bangkok Clay (samples 20SS01A, 01B, 02A, 02B, 03A, 04A, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

**Remarks**

*Neomonoceratina mediterranea mediterranea* (Ruggieri, 1953) is recognized by its small carapace with two short oblique posterodorsal ribs, a long median rib, one venterolateral rib and one ventral rib. The posterodorsal and venterodorsal ribs connect with median rib in the posterior area. The carapace surface is finely punctate. Two of the three specimens identified as *Neomonoceratina* sp. in Forel (2021) are here re-attributed to *N. mediterranea mediterranea* (Forel 2021: fig. 4g–h).

*Neomonoceratina mediterranea malayensis* Zhao & Whatley, 1988  
Fig. 6O–T

*Neomonoceratina mediterranea malayensis* Zhao & Whatley, 1988: 566, pl 1 figs 4–6.

*Neomonoceratina* sp. – Forel 2021: 7, fig. 4f.

**Dimensions**

L = 0.433–0.480 millimeters; H = 0.187–0.213 millimeters; H/L = 0.39–0.47.

**Distribution**

Recent distribution: Jason Bay, southeastern Malay Peninsula, Malaysia (Zhao & Whatley 1989); Southwestern coast of Peninsular Thailand, Ao Nun, Satun Province, Andaman Sea (Forel 2021).

Fossil distribution: Bangkok Clay (samples 20SS01A, 01B, 02A, 02B, 03A, 04A, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

**Remarks**

The size and H/L ratio of *N. mediterranea malayensis* are smaller than those of *N. mediterranea mediterranea*. Punctae on the surface are restricted to the base of the ribs while they are rare on the smooth intercostal surface. *Neomonoceratina mediterranea malayensis* was first recovered from shallow water sediments of the Jason Bay, in Malay Peninsula (Zhao & Whatley 1988). This species was also found along the Andaman coast of Thailand (Forel 2021: fig. 4f).

Family Trachyleberididae Sylvester-Bradley, 1948

Genus *Stigmatocythere* Siddiqui, 1971

**Type species**

*Stigmatocythere obliqua* Siddiqui, 1971 by original designation

*Stigmatocythere bona* Chen, 1982  
Fig. 7G–I

*Stigmatocythere bona* Chen in Hou *et al.*, 1982: figs 153–154.

*Stigmatocythere bona* – Whatley & Zhao 1988b: 9, pl. 6 fig. 19. — Dewi 1993: 69, figs 153–154. — Montenegro *et al.* 2004: pl. 2 fig. 8. — Forel 2021: 8, fig. 5d–e.


**Dimensions**

L = 0.407–0.678 millimeters, H = 0.221–0.360 millimeters, H/L = 0.41–0.58.

**Distribution**

Modern distribution: Java Sea (Dewi 1993, 2000); Malacca Straits (Whatley & Zhao 1988); Sedili River, Jason Bay (Zhao & Whatley 1989); Vietnam (Tan *et al.* 2021); Sri Lanka (Iwatani *et al.* 2014); Mae Khlong River mouth, north west Gulf of Thailand (Montenegro *et al.* 2004); Southwestern coast of Peninsular Thailand, Ao Nun, Satun Province, Andaman Sea (Forel 2021).

Fossil distribution: East China, Pliocene and Quaternary (Hou *et al.* 1982; Gou *et al.* 1983); Bangkok Clay (samples 20SS01A, 02B, 03A), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).



**Fig. 7.** Holocene ostracods from Samut Sakhon Province, Central Thailand. **A–E.** *Keijella gonia* Zhao & Whatley, 1989. **A.** Carapace, left lateral view, SUT-20SS-C310. **B.** Carapace, left lateral view, SUT-20SS-C311. **C.** Carapace, left lateral view, SUT-20SS-C312. **D.** Carapace, right lateral view, SUT-20SS-C288. **E.** Carapace, right lateral view, SUT-20SS-C286. – **F.** *Pistocythereis* sp. Carapace, right lateral view, SUT-20SS-C305. – **G–I.** *Stigmatocythere bona* Chen in Hou, Chen, Yang, Ho, Zhou & Tian, 1982. **G.** Carapace, right lateral view, SUT-20SS-C201. **H.** Carapace, left lateral view, SUT-20SS-C200. **I.** Valve, internal view of right valve, SUT-20SS-C190. – **J–O.** *Keijella multisulcus* Whatley & Zhao, 1988. **J.** Valve, juvenile, right lateral view, SUT-20SS-C231. **K.** Valve, juvenile, internal view of right valve, SUT-20SS-C213. **L.** Carapace, male, left lateral view, SUT-20SS-C225. **M.** Carapace, female, left lateral view, SUT-20SS-C242. **N.** Carapace, female, left lateral view, SUT-20SS-C206. **O.** Carapace, female, left lateral view, SUT-20SS-C243. Scale bars = 0.1. mm.



Genus *Keijella* Ruggieri, 1967

**Type species**

*Cythere hodgii* Brady, 1866 subsequently designated by Ruggieri (1967).

*Keijella multisulcus* Whatley & Zhao, 1988  
Figs 7J–O, 8A–I, 9

*Keijella multisulcus* Whatley & Zao, 1988: 15, fig. d.

**Dimensions**

L = 0.325–0.875 millimeters; H = 0.175–0.444 millimeters; H/L = 0.45–0.59 (Fig. 9).

**Distribution**

Modern distribution: Malacca Strait (Whatley & Zhao 1988b); Malaysia (Omar *et al.* 2017); Mae Khlong River mouth, north west Gulf of Thailand (Montenegro *et al.* 2004); southwestern coast of Peninsular Thailand, Ao Nun, Satun Province, Andaman Sea (Forel 2021).

Fossil distribution: Bangkok Clay (samples 20SS01B, 02A, 02B, 03A, 04A, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

**Remark**

Dimorphism of *Keijella multisulcus* can be recognized by the subrectangular shape with H/L ratio less than 0.50 in the male (Fig. 7L) and the subovate shape with a higher H/L ratio in the female (Fig. 7M–O). The juvenile carapaces are sub-triangular in lateral view with Hmax located anteriorly at one third of L, Lmax is located below midH in both valves. The ventral longitudinal carina is prominent in young juveniles (Fig. 8G–H) and extends into a small spine (posteroventral spine). The ventral longitudinal carina is faint and becomes obscured in the larger juvenile stage (Fig. 8A–F). Small conical posterior marginal denticles and a large terminal posteroventral spine are observed in juveniles. Figure 9 shows H and L plot of the species which clearly demonstrates that sexual dimorphism can be differentiated in adult specimens.

*Keijella gonia* Zhao & Whatley, 1989  
Figs. 7A–E, 10

*Keijella gonia* Zhao & Whatley, 1989: 181, pl. 3 figs 7–10, 16.

*Keijella gonia* – Montenegro *et al.* 2004: pl. 1 fig. 8. — Forel 2021: 8, fig. 5h–i.

**Dimensions**

L = 0.380–0.710 millimeters; H = 0.187–0.380 millimeters; H/L = 0.49–0.62.


**Distribution**

Modern distribution: Mae Khlong River mouth, north west Gulf of Thailand (Montenegro *et al.* 2004); Sedili River, Jason Bay (Zhao & Whatley 1989); Central Vietnam (Tan *et al.* 2021); Indonesia (Fauzielly 2013); Southwestern coast of Peninsular Thailand, Ao Nun, Satun Province, Andaman Sea (Forel 2021).

Fossil distribution: Bangkok Clay (samples 20SS01B, 02A, 02B, 03A, 04A, 04B), whale excavation site, Samut Sakhon Province, Thailand, Late Holocene (this work, Fig. 4).

**Remarks**

*Keijella gonia* is easily recognized by the distinct angle formed by the posteroventral intersection of vertical and horizontal muri extending into a spine, sometimes sharp and long. The carapace is sub-rectangular in lateral view, strongly reticulated. Well-preserved specimens show laterally compressed anterior and posterior marginal denticles, especially along AVB. Prominent postero-marginal spines are distinct and located below the marginal denticles especially in juvenile specimens (Fig. 7A–C). Figure 10 shows H and L plots of *K. gonia* compared to the types from Malaysia (Zhao & Whatley 1989), the specimens from Thailand consist of adults and juveniles.



**Fig. 8.** Holocene ostracods from Samut Sakhon Province, Central Thailand. **A–I.** *Keijella multisulcus* Whatley & Zhao, 1988. **A.** Valve, juvenile, right lateral view, SUT-20SS-C254. **B.** Valve, female, left lateral view, SUT-20SS-C269. **C.** Carapace, juvenile, left lateral view, SUT-20SS-C270. **D.** Valve, juvenile, right lateral view, SUT-20SS-C267. **E.** Valve, juvenile, left lateral view, SUT-20SS-C286. **F.** Valve, juvenile, right lateral view, SUT-20SS-C253. **G.** Valve, juvenile, right lateral view, SUT-20SS-C272. **H.** Valve, juvenile, left lateral view, SUT-20SS-C273. **I.** Valve, female, right lateral view, SUT-20SS-C257. – **J–L.** *Ammonia tepida* (Cushman, 1926), spiral side, SUT-20SS-F001. – **M.** *Spiroloculina* sp., side view, SUT-20SS-F002. – **N.** *Quinqueloculina* sp., side view, SUT-20SS-F003. – **O–Q.** *Asterorotalia pulchella* (d’Orbigny, 1839), side view, SUT-20SS-F004-6. Scale bars = 0.1 mm.

Genus *Pistocythereis* Gou, 1983

*Pistocythereis* sp.

Fig. 7F

**Dimensions**

L = 0.670–0.778 millimeters; H = 0.330–0.400 millimeters; H/L = 0.49–0.54.


**Remarks**

Five specimens of *Pistocythereis* sp. were found from samples 20SS02A, 02B, 03A. They can be differentiated from *Keijella gonia* by the larger carapace without posteroventral spine at intersection of vertical and horizontal muri.

**Discussion**

**Taxonomic composition and diversity**

Seven of the 10 silty clay samples, each weighing 200 grams, from the total thickness of 2 meters at the whale-excavation site yielded ostracods (Fig. 4). The specimens studied here are both disarticulated valves and complete carapaces. Thirteen species belonging to seven genera and five families are here identified. To discuss the composition and diversity of the ostracod assemblages, the specimens of each species were counted articulated plus a higher number of left or right valves (Nützel & Kaim 2014; Haussmann & Nützel 2015). 1869 specimens were sorted, the relative abundance of each taxon through the entire section was calculated. The family Trachyberididae is the most abundant (52.71% of the specimens) including *Keijella multisulcus* (31.38%), *Keijella gonia* (19.62%), *Stigmatocythere bona*




**Fig. 9.** Height and length scatter plot of *Keijella multisulcus* Whatley & Zhao, 1988 found at the whale-fall excavation site in Samut Sakhon Province, north of the Gulf of Thailand. Scale bars = 0.1 mm.



(1.58%) and *Pistocythereis* sp. (0.26%). The family Schizocytheridae is the second most abundant, accounting for 24.83% of the species; it is only represented by the genus *Neomonoceratina* including: *N. rhomboidae* (14.77%), *N. mediterranea* (4.69%), *N. mediterranea malayensis* (3.69%), *N. iniqua* (1.69%), and *N. columbiformis* (0.05%). *Sinocythere impressa* (16.89%) is the only representative of the family Cytherideidae. The family Pontocyprididae (*Propontocypris bengalensis* and *P. clara*) is 4.52% and the family Candonidae (*Agelaiocypris pellucida*) is 1.05% of the specimens.

### Stratigraphical and geographical distributions

The ostracods reported in this study are from sediments around the whale skeleton which is dated to  $3380 \pm 30$  years (Kawira & Saethien 2021; Saethien 2021). All seven genera including *Agelaiocypris*, *Propontocypris*, *Sinocytheridae*, *Neomonoceratina*, *Stigmatocythere*, *Keijella* and *Pistocythereis* are frequent components of Cenozoic to Recent ostracod faunas from the Indo-Pacific and South China region (e.g., Hong *et al.* 2019; Tanaka *et al.* 2019; Forel 2021; Tan *et al.* 2021), Indian Ocean (e.g., Hussain *et al.* 2004; Nishath *et al.* 2015) and Persian Gulf (Mostafawi 2003). In terms of species, Table 1 summarizes the spatial distributions of the species which are already known from previous works. Of these, six species have previously been reported from Recent sediments of Thailand: *K. multisulcus*, *K. gonia*, *S. bona*, *N. iniqua*, *S. impressa* from Mae Khlong River mouth, Phetchaburi Province (Montenegro *et al.* 2004); *K. multisulcus*, *S. impressa*, *N. iniqua*, *N. columbiformis*, *S. bona* from Klong Thom estuary, Krabi Province (Yamada *et al.* 2014); *K. gonia*, *S. bona*, *N. iniqua*, from Ao Nun, Satun Province (Forel 2021). It should be noted that the works of Yamada *et al.* (2014) and Forel (2021) were conducted on the Andaman Sea, not the Gulf of Thailand.



**Fig. 10.** Height and length scatter plots of *Keijella gonia* Zhao & Whatley, 1989 from Malaysia (grey circles) compared with specimens found from the whale-fall excavation site in Samut Sakhon Province, north of the Gulf of Thailand (blue circles). Scale bars = 0.1 mm.

Species of the family Trachyberididae are the most abundant and observed in this study, and they are also common species found in the Gulf of Thailand, Malacca Strait and Java Sea. *Keijella multisulcus* is restricted to Thailand and Malaysia (Zhao & Whatley 1988a, 1988b; Montenegro *et al.* 2004; Yamada *et al.* 2014; Forel 2021). *Keijella gonia* has been reported from Thailand and Malaysia and extended to Vietnam and Indonesia (Fauzielly 2013; Tan *et al.* 2021). *Stigmatocythere bona* is widely dispersed from Thailand, Malaysia, Indonesia (Dewi 1997) and Sri Lanka (Iwatani *et al.* 2014).

The family Schizocytheridae is the second abundant one in this study. They are very diversified in China and Malaysia (Zhao & Whatley 1988a, 1988b). Five species, namely *Neomonoceratina rhomboidae*, *N. mediterranea mediterranea*, *N. mediterranea malayensis*, *N. columbiformis* and *N. iniqua* are recovered in this study. *Neomonoceratina columbiformis* was reported from the Andaman Sea coast (Yamada *et al.* 2014). *Neomonoceratina iniqua* has been reported from the Andaman Sea coasts (Yamada *et al.* 2014; Forel 2021), Vietnam (Tan *et al.* 2021) and Japan (Ishizaki & Kato 1976).

From Table 1 *Sinocytheridae impressa* is the only species linked with the fauna from China Sea. Hong *et al.* (2019) classified the living *S. impressa* to the Subtropical Group which is dispersed from the East China Sea to the Indo-Pacific area. Tanaka *et al.* (2019) critically reviewed the occurrences of *S. impressa* from the Sea of Japan to the South China Sea and expressed that the species has records back to the Late Pliocene. *Sinocytheridae impressa* has been recovered from Recent marine sediments of Vietnam (Tanaka *et al.* 2009; Tan *et al.* 2021) and Thailand (Montenegro *et al.* 2004; Yamada *et al.* 2014). The discovery of *S. impressa* in this study reveals that *S. impressa* has inhabited in the Gulf of Thailand since at least 3300 years ago.

### **Paleoenvironmental interpretation**

Most of the ostracod species found in this study are typical South China-Indo-Pacific shallow water taxa. The composition of the successive assemblages is quite consistent from the lower to the upper part of the section (2 meters) as shown in Figs 3–4. Thus, there should not be any abrupt change of lithology and/or ostracods before and after the whale was deposited. However, there is still a question why the samples 20SS03B, 20SS05, 20SS06 were barren of ostracods. In this study, 1187 carapaces and 1455 valves in total were sorted. The relative proportion of carapaces and valves is relatively stable across the section, ranging from 47% to 57% of complete carapaces (Table 2). Carapace features such as surface ornamentation, spines, muscle scars can be observed in many specimens suggesting that they were buried quickly after death and had been shortly transported (Oertli 1971; Frenzel & Boomer 2005). All assemblages are composed of a mixture of adults of both sexes, juveniles (*K. multisulcus*, *K. gonia*, *S. impressa*) as well as complete carapaces of tiny species such as *N. mediterranea mediterranea* and *N. mediterranea malayensis*, and delicate and transparent carapaces of *A. pellucida*, *P. bengalensis* and *P. clara*. These characteristics all indicate a relatively low energy biocoenosis (Whatley 1983; Frenzel & Boomer 2005).

The environmental distribution of *S. impressa* within the South China Sea and the Indo-Pacific region is relatively well characterized (Zhao & Whatley 1988; Yasuhara & Seto 2006; Tanaka *et al.* 2009, 2012, 2019; Alberti *et al.* 2013; Wang *et al.* 2018; Hong *et al.* 2019). Accordingly, *S. impressa* prefers euryhaline, eurythermal, and usually correlates with a nutrient rich mud substrate. It can be tolerant to turbid and low dissolved oxygen conditions. It preferentially occurs in shallow water (< 20 meters) and thrives in a wide range of environments such as inner shelf, sublittoral to intertidal, estuary and brackish water. *Sinocytheridae impressa* is also a good bioindicator of Recent benthic marine ecosystems as it has strong correlations with mud and nutrient-rich muddy to fine sandy environments (Hong *et al.* 2021a, 2021b; Tan *et al.* 2021).

**Table 1.** Geographical distributions of Holocene ostracods from the marine Bangkok Clay in Samut Sakhon Province, north of the Gulf of Thailand.

Region	References/Species	<i>Aglaioocypris pellicida</i>	<i>Propontocypris clara</i>	<i>P. bengalensis</i>	<i>Neomonoceratina columbiformis</i>	<i>N. mediterranea mediterranea</i>	<i>N. mediterranea malayaensis</i>	<i>N. iniqua</i>	<i>N. rhomboidea</i>	<i>Sinocytheridae impressa</i>	<i>Keijella multisulcus</i>	<i>K. gonia</i>	<i>Stigmatocypris bona</i>
Japan	Ishizako & Kato 1976						*						
	Tanaka <i>et al.</i> 2019									*			
China	Cheung <i>et al.</i> 2019									*			
	Gu <i>et al.</i> 2019									*			
	Tanaka <i>et al.</i> 2019									*			
	Yan <i>et al.</i> 2020									*			
Hong Kong	Whatley & Zhao 1987									*			
	Yim <i>et al.</i> 1988									*			
	Wang <i>et al.</i> 2018	*	*	*						*			
	Hong <i>et al.</i> 2019									*			
Vietnam	Tanaka <i>et al.</i> 2009		*							*			
	Tan <i>et al.</i> 2021						*			*		*	
Thailand	Montenegro <i>et al.</i> 2004						*			*	*	*	*
	Yamada <i>et al.</i> 2014				*		*			*	*		
	Forel 2021						*			*	*	*	*
Malaysia	Zhao & Whatley 1988a, 1988b				*	*	*	*	*	*	*		
	Zhao & Whatley 1989				*	*	*			*	*	*	*
	Ramlan & Noraswana 2009; 2010												
	Noraswana <i>et al.</i> 2014												
	Omar <i>et al.</i> 2017										*		
Indonesia	Dewi 1997						*						*
	Fauzielly 2013				*		*					*	
Sri Lanka	Iwatani <i>et al.</i> 2014						*						*
India	e.g., Bhatia & Kumar 1979; Hussain <i>et al.</i> 2004; 2007; 2013a; 2013b						*						
Persia Gulf	Mostafawi 2003	*		*			*						
	Paik 1977						*						

Compared with previous research in Thailand, the studied ostracod assemblage is similar to those from the Mae Klong River Mouth (Montenegro *et al.* 2004), located about 30 km southwest of the studied section (Fig. 2, Table 1). In terms of bathymetry, Montenegro *et al.* (2004) and Pugliese *et al.* (2006) considered that *K. multisulcus*, *K. gonia*, *N. iniqua*, *S. impressa* and *S. bona* correspond to a group of species that have a wide depth range because of their extensive occurrence at depth less than 18 meters. *Neomonoceratina iniqua* and *S. impressa* tolerate diverse substrate types but are absent from sand



**Table 2.** Relative proportions of carapaces and valves per productive sample.

Sample	Carapaces (%)	Valves (%)
20SS04_B	43	57
20SS04_A	53	47
20SS03_A	48	52
20SS02_B	37	63
20SS02_A	45	55
20SS01_B	46	54
20SS01_A	53	47

whereas *K. gonia* and *S. bona* prefer silty sand to silt. They also suggested that autochthonous ostracods were not found at stations with a salinity of 20–22‰, the rich and abundant assemblages being present at high values of salinity, dissolved oxygen content and pH. But the ostracods were not recovered from sediments collected in the Mae Khlong River channel and on the tidal flat.

The assemblage studied here from about 3000 years ago may represent conditions prior to the onset of significant anthropogenic contamination. The abundance of *S. impressa* in this study should correspond to coastal and estuarine areas with a wide range of salinity and muddy and silty substrates (e.g., Zhao & Wang 1988; Tanaka *et al.* 2009, 2019; Hong *et al.* 2019, 2021b) and the infralittoral zone (Pugliese *et al.* 2006).

The ostracod assemblage from the whale-fall excavation site therefore points to a shallow-water marine environment such as estuary, bay, inner shelf, subtidal zone, (< 20 meters depth), with a brackish to normal salinity (> 22‰), a high mud content and turbidity, on muddy substrates. The depositional environment was stable and clam before and after the whale sank. The radiocarbon dating provided an age of  $3380 \pm 30$  years for the whale itself (Kawira & Saethien 2021; Saethien 2021), while the sediments below and above were a little older and younger, respectively. Good preservation of the skeleton owes to suitable bathymetric depth in calm condition and rapid depositional rate during sea water regression (Tanabe *et al.* 2003). In our samples, the ostracods, foraminifers and micro-mollusks are also well-preserved (Figs 4 and 8). Benthic foraminifers such as *Ammonia tepida* (Cushman, 1926) (Fig. 8J–L), *Spiroloculina* sp. (Fig. 8M), *Quinqueloculina* sp. (Fig. 8N), *Asterorotalia pulchella* (d’Orbigny, 1839) (Fig. 8O–P) are abundant in all samples. *Asterorotalia pulchella* is known to be associated with a water depth of 21–25 m in the Strait of Malacca (Minhat *et al.* 2021) and the inner shelf of the Sunda Shelf (Szarek *et al.* 2006). *Ammonia tepida* prefers areas with a high organic matter composition (Minhat *et al.* 2021).

In this study, we examined ostracods from Holocene sediments below and above the whale-fall skeleton found in Am Pang Subdistrict, Ban Paew District, Samut Sakhon Province. The locality is onshore, about fifteen kilometers away from the Recent shoreline of northern Gulf of Thailand. Numerous carapaces and valves were recovered and identified. Thirteen species were recognized. The present analysis of the ostracod assemblage provides the first characterization of the Holocene environment associated with the whale-fall. The ostracods typically correspond to a shallow marine assemblage indicating infralittoral zone at water depth less than 20 meters, brackish to normal salinity, high mud content and turbidity. The depositional environment might be estuary, bay, or inner shelf.

## Acknowledgements

This project was financially supported by Suranaree University of Technology (SUT7-719-60-12-64). We are grateful to Fossil Protection Division, Royal Thai Department of Mineral Resources and the Bright Blue Water Corporation Limited for allowance to visit the whale excavation site and collect samples. We are grateful to Dr Moriaki Yasuhara (University of Hong-Kong) and Dr Gengo Tanaka (Kanazawa University) for their very constructive advices that greatly improved an earlier version of this manuscript.

## References

- Alberti M., Hethke M., Fürsich F. & Liu C. 2013. Macro-versus microfauna: resolution potential of bivalves, gastropods, foraminifera and ostracods in reconstructing the Holocene palaeoenvironmental evolution of the Pearl River delta, China. *Palaeobiodiversity and Palaeoenvironments* 93: 327–353. <https://doi.org/10.1007/s12549-012-0101-9>
- Al-Jumaily W.A.K. & Al-Sheikhly S.S. 1999. Palaeozoogeography of shallow marine Ostracoda from Holocene sediments-Southern Iraq. *Qatar University Science Journal* 18: 215–230.
- Baird W. 1845. Arrangement of the British Entomostraca, with a list of species, particularly noticing those which have as yet been discovered within the bounds of the Club. *History of the Berwickshire Naturalists' Club* 2: 145–158. Available from <https://www.biodiversitylibrary.org/page/2408349> [accessed 2 Jan. 2023].
- Baird W. 1850. *The Natural History of the British Entomostraca*. Ray Society, London. <https://doi.org/10.5962/bhl.title.39641>
- Baskar K., Sridhar S.G.D., Hussain S.M., Solai A. & Kalaiivanan R. 2013. Distribution of Recent benthic Ostracoda off Rameswaram, Palk Strait, Tamil nadu, South East Coast of India. *Special Publication of the Geological Society of India* 1: 195–212.
- Bate R.H. 1971. The distribution of Recent Ostracoda in the Abu Dhabi Lagoon, Persian Gulf. In: Oertli H.J. (ed.) *Paléoécologie des Ostracodes*: 239–256. Société nationale des pétroles d'Aquitaine, Centre de recherche.
- Bhatia S.B. & Kumar S. 1979. Recent Ostracoda from off Karwar, West Coast of India. In: Krstic N. (ed.) *Proceedings of the VII<sup>th</sup> International Symposium on Ostracodes – Taxonomy, Biostratigraphy and Distribution of Ostracodes*: 173–178. The Serbin Geological Society, Beograd.
- Bonaduce G., Ciliberto B., Minichell G., Masoli M. & Pugliese N. 1983. The Red Sea benthic ostracodes and their geographical distribution. In: Maddocks R.E. (ed.) *Applications of Ostracoda. Proceedings of the 8<sup>th</sup> International Symposium on Ostracoda*: 472–491. Houston.
- Brady G.S. 1866. On new or imperfectly known species of marine Ostracoda. *Transactions of the Zoological Society of London* 5: 359–393. <https://doi.org/10.1111/j.1096-3642.1866.tb00649.x>
- Brady G.S. 1868. Description of Ostracoda. In: Folin L. de & Périer L. (eds) *Les Fonds de la Mer. Étude sur les Particularités nouvelles des Régions sous-marines. Part 1*: 49–112. Savy-Librairie Editeur, Paris. <https://doi.org/10.5962/bhl.title.13261>
- Brady G.S. 1869. Descriptions of Ostracoda. In: Folin L. de & Périer L. (eds) *Les Fonds de la Mer. Étude sur les Particularités nouvelles des Régions sous-marines Part 2*: 113–176. Savy-Librairie Editeur, Paris. <https://doi.org/10.5962/bhl.title.13261>
- Cao M. 1998. Ostracoda from Quaternary Hang Hau Formation, Lei Yue Mun, Hong Kong. In: Li Z. (ed.) *Paleontology and Stratigraphy in Hong Kong (Lower Volume)*: 171–183. Science Press, Beijing.

- Chen T.-C. 1978. Descriptions of ostracodes. In: *Zhongnan diqu gushengwu tuce (Fossils of Central Southern China)* 1: 111–325, 2: 682–710.
- Cheung R.C.W., Yasuhara M., Iwatani H., Wet C.-L. & Dong Y.-W. 2019. Benthic community history in the Changjiang (Yangtze River) mega-delta: damming, urbanization, and environmental control. *Paleobiology* 45 (3): 469–483. <https://doi.org/10.1017/pab.2019.21>
- Danise S., Dominici S., Glover A.G. & Dahlgren T.G. 2014. Molluscs from a shallow-water whale-fall and their affinities with adjacent benthic communities on the Swedish west coast. *Marine Biology Research* 10: 3–16. <https://doi.org/10.1080/17451000.2013.793811>
- Dewi K.T. 1993. *Ostracoda from the Java Sea, West of Bawean Island, Indonesia*. Master Thesis, University of Wollongong, Australia.
- Dewi K.T. 2000. Distribution of ostracoda from South of Tanjung Selatan, South Kalimantan. *Bulletin of Marine Geology* 15 (1): 1–14.
- Fauzielly L. 2013. Distribusi vertikal Ostracoda dan hubungannya dengan perubahan lingkungan di perairan teluk Jakarta. *Bulletin of Scientific Contribution* 11 (2): 108–117.
- Fauzielly L., Irizuki T. & Samp Y. 2012. Vertical changes of recent ostracode assemblages and environment in the inner part of Jakarta Bay, Indonesia. *Journal of Coastal Development* 16 (1): 11–24.
- Fauzielly L., Irizuki T. & Sampei Y. 2013. Spatial distribution of recent ostracode assemblages and depositional environments in Jakarta Bay, Indonesia, with relation to environmental factors. *Paleontological Research* 16 (4): 267–281. <https://doi.org/10.2517/1342-8144-16.4.267>
- Forel M. 2021. Recent ostracods (Crustacea) from the southwestern coast of Peninsular Thailand (Satun Province), Andaman Sea. *Revue de Micropaléontologie* 72: 100526. <https://doi.org/10.1016/j.revmic.2021.100526>
- Frenzel P. & Boomer I. 2005. The use of ostracods from marginal marine, brackish waters as bioindicators of modern and Quaternary environmental change. *Palaeogeography, Palaeoclimatology, Palaeoecology* 225: 68–92. <https://doi.org/10.1016/j.palaeo.2004.02.051>
- Furushima Y., Okoshi K., Miyake H., Miyazaki M., Nogi Y., Yatabe A. & Okutani T. 2007. Three-year investigations into sperm whale-fall ecosystems in Japan. *Marine Ecology* 28: 219–232. <https://doi.org/10.1111/j.1439-0485.2007.00150.x>
- Gopalakrishna K., Hussain S.M., Mahesh Bilwa L. & Ayisha V.A., 2007. Recent benthic Ostracoda from the inner-shelf off the Malabar coast, Malabar, Kerala, southwest coast of India. *Journal of the Palaeontological Society of India* 52 (1): 59–68.
- Gou Y., Chen T., Guan S. & Jian Y. 1981. Ostracoda. In: China, South sea branch of Petroleum Corporation of the Peoples Republic (ed.) *Tertiary Paleontology of North Continental Shelf of South China Sea*: 138–187. Guangdong Science & Technology Press, Guangzhou.
- Gou Y., Zheng S. & Huang B. 1983. Pliocene Ostracode fauna of Leizhou Peninsula and northern Hainan Island, Guangdong Province. *Palaeontologia Sinica Series B* 162 (18): 1–157.
- Guan S.Z., Sun Q.Y., Jiang Y.W., Li L.L., Zhao B.Q., Zhang X.Q., Yang R.L. & Feng B.Y. 1978. Ostracoda. In: *Paleontological atlas of Central and South China Vol. 4*. Geological Publishing House, Beijing.
- Guha D K. 1966. Young Cenozoic marine Ostracoda from subcrops of south India. *Journal of the Geological Society of India Special No.*: 30–31.
- Hanai T., Ikeya N. & Yajima M. 1980. Checklist of Ostracoda from southeast Asia. *Bulletin University Museum University of Tokyo* 17: 1–236.



- Hartmann G. & Puri H.S. 1974. Summary of neontological and paleontological classification of ostracoda. *Mitteilungen aus dem hamburgischen zoologischen Museum und Institut* 70: 7–73.
- Hausmann I.M. & Nützel A. 2015. Diversity and palaeoecology of a highly diverse Late Triassic marine biota from the Cassian Formation of north Italy. *Lethaia* 48: 235–255. <https://doi.org/10.1111/let.12102>
- Hong Y., Yasuhara M., Iwatani H., Seto K., Yokoyama Y., Yoshioka K. & Mamo R. 2017. Freshwater reservoir construction by damming a marine inlet in Hong Kong: paleoecological evidence of local community change. *Marine Micropaleontology* 132: 53–59. <https://doi.org/10.1016/j.marmicro.2017.04.003>
- Hong Y., Yasuhara M., Iwatani H. & Mamo B. 2019. Baseline for ostracod-based northwestern Pacific and Indo-Pacific shallow-marine paleoenvironmental reconstructions: ecological modeling of species distributions. *Biogeosciences* 16 (2): 585–604. <https://doi.org/10.5194/bg-16-585-2019>
- Hong Y., Yasuhara M., Iwatani H., Chao A., Harnik P.G. & Wei C.-L. 2021a. Ecosystem turnover in an urbanized subtropical seascape driven by climate and pollution. *Anthropocene* 36: 100304. <https://doi.org/10.1016/j.ancene.2021.100304>
- Hong Y., Yasuhara M., Iwatani H., Harnik P.G., Chao A., Cybulski J.D., Liu Y., Ruan Y., Li X. & Wei C.-L. 2021b. Benthic ostracod diversity and biogeography in an urbanized seascape. *Marine Micropaleontology* 174: 102067. <https://doi.org/10.1016/j.marmicro.2021.102067>
- Hou Y. & Gou Y. 2007. *Fossil Ostracoda of China, 2: Cytheracea and Cytherellidae*. Science Press, Beijing.
- Hou Y., Chen T., Yang H., Ho J., Zhou Q. & Tian M.-Q. 1982. *Cretaceous-Quaternary Ostracode Fauna from Jiangsu*. Geological Publishing House, Beijing.
- Hu C.-H. 1984. New fossil ostracod faunas from Hengchun Peninsula, southern Taiwan. *Journal of Taiwan Museum* 37 (1): 65–129.
- Hu C.-H. & Yang L.-C. 1975. Studies on Pliocene ostracodes from the Chinshui Shale, Miaoli district, Taiwan. *Proceedings of the Geological Society of China*. 18: 103–114.
- Hu C.-H. & Yeh K.-Y. 1978. Ostracod faunas from the Pleistocene Liushuang Formation in the Tainan area, Taiwan. *Proceedings of the Geological Society of China* 21: 151–162.
- Huang B. 1985. Ostracoda from the column samples under surface deposit on the bottom of north Bohai Sea. *Journal of Oceanography of Huanghai and Bohai Seas* 3: 42–53.
- Hussain S.M. 1998. Recent benthic Ostracoda from the Gulf of Mannar, off Tuticorin, southeast coast of India. *Journal of the Palaeontological Society of India* 43: 1–22.
- Hussain S.M. & Kalaiyarasi A. 2013. Distribution of Ostracoda in the Mullipallam Lagoon, near Muthupet, Tamil Nadu, Southeast Coast of India – Implications on Microenvironment. In: Sundaresan J., Sreekesh S., Ramanathan A.L., Leonard S. & Ram B. (eds) *Climate Change and Island and Coastal Vulnerability*: 166–176. Capital Publishing Company. [https://doi.org/10.1007/978-94-007-6016-5\\_11](https://doi.org/10.1007/978-94-007-6016-5_11)
- Hussain S.M. & Mohan S.P. 2001. Distribution of recent benthic Ostracoda in Adyar river estuary, east coast of India. *Indian Journal of Marine Sciences* 30: 53–56.
- Hussain S.M., Ravi G. Mohan S.P. & Rajeshwara Rao N. 2004. Recent benthic Ostracoda from the inner shelf off Chennai, southeast coast of India – implication on microenvironments. *Environmental Micropaleontology, Microbiology and Meiobenthology* 1: 105–121.
- Hussain S.M., Ganesan P., Ravi G., Mohan S.P. & Sridhar S.G.D. 2007. Distribution of Ostracoda in marine and marginal marine habitats off Tamil Nadu and adjoining areas, southern east coast of India and Andaman Islands: Environmental implications. *Indian Journal of Marine Sciences* 36 (4): 369–377.

- Hussain S.M., Elumalai K., Scott Immanuel Dhas C. & Mohan S.P., 2013a. Distribution of Ostracoda and the sediment characteristics of Ennore Creek, Tamil Nadu, southeast coast of India. *In: Rao D.V., Kosygin L. & Dash S. (eds) Estuaries of India: Biodiversity, Ecology, Conservation and Management*: 203–218. Nature Books India, Kolkata.
- Hussain S.M., Kuleen E.C., Jisha K., Elumalai K. & Ravi G. 2013b. Distribution of Benthic Ostracoda in surface and subsurface backwater sediments of Ernakulam, Kerala, Southwest coast of India: microenvironmental implications. *Special Publication of the Geological Society of India* 1: 213–224.
- Hutangkura T. 2012. *Pollen Analysis of the Holocene Sedimentary Sequences from the Lower Central Plain of Thailand and Its Implications for Understanding Palaeo-environmental and Phytogeographical Changes*. Ph.D. Thesis, Université de Nice Sophie Antipolis.
- Hu C.-H. & Tao H.-J. 2008. Studies on the ostracod fauna of Taiwan and its adjacent seas. *Journal National Taiwan Museum Special Publication Series* 13 (Parts I and II): 1–655 (Part I), 657–910 (Part II).
- Irizuki T., Matsubara T. & Matsumoto H. 2005. Middle Pleistocene Ostracoda from the Takatsukayama Member of the Meimi Formation, Hyogo Prefecture, western Japan: significance of the occurrence of *Sinocytheridea impressa*. *Paleontological Research* 9: 37–54. <https://doi.org/10.2517/prpsj.9.37>
- Iwatani H., Young S.M., Irizuki T., Sampei Y. & Ishiga H. 2014. Spatial variations in recent ostracode assemblages and bottom environments in Trincomalee Bay, northeast coast of Sri Lanka. *Micropaleontology* 60 (6): 509–518. <https://doi.org/10.47894/mpal.60.6.02>
- Ishizaki K. & Kato M. 1976. The basin development of the Diluvium Furuya Mud Basin, Shizuoka prefecture, Japan, based on faunal analysis of fossil ostracodes. *In: Takayanagi Y. & Saito T. (eds) Progress in Micropaleontology: Selected Papers in Honour of Professor Kiyoshi Asano*: 118–143. Micropaleontology Press, New York.
- Ishizaki K. 1990. Sea level change in mid-Pleistocene and effects on Japanese ostracode faunas. *Bulletin of Marine Science* 47: 213–220.
- Jain S.E. 1978. Recent Ostracoda from Mandvi Beach, west coast of India. *Bulletin of the Indian geologists Association* 11 (2): 89–139.
- Kaufmann A. 1900. Cypriden und Darwinuliden der Schweiz. *Revue suisse de Zoologie* 8: 209–423. <https://doi.org/10.5962/bhl.part.10584>
- Kawira A. & Saethien P. 2021. Excavation and taxonomic study on Holocene whale from Am Pang Subdistrict, Ban Peaw District, Samut Sakhon Province. *In: Suvapak I., Apsorn S., Thawatchai C., Prachya B., Siripond S., Denchak M., Angsumalin P., Sakda K., Veerachat V., Metha Y., Prodit N., Phornpen C. & Sirirat P. (eds) Abstract Book of Geothai Webinar 2021*: 47. Royal Department of Mineral Resources of Thailand, Bangkok.
- Keij A.J. 1953. Preliminary note on the Recent Ostracoda of the Snellius Expedition. *Proceedings of the Section of Sciences, Koninklijke Nederlandse Akademie van Wetenschappen te Amsterdam* 56: 155–168.
- Keij A.J. 1954. Some recent ostracods of Manila, Philippines. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B Physical Sciences* 57 (3): 351–363.
- Keij A.J. 1957. Eocene and Oligocene Ostracoda of Belgium. *Mémoires de l'Institut royal des Sciences naturelles de Belgique* 6 (23): 1–210.
- Keij A.J. 1979. Brief review of type species of genera from the Kingma collection. *In: Krstic N. (ed.) Proceedings of the VII<sup>th</sup> International Symposium on Ostracodes – Taxonomy, Biostratigraphy and Distribution of Ostracodes*: 59–62. The Serbin Geological Society, Beograd.

- Kingma J.T. 1948. *Contributions to the knowledge of the Young-Cenozoic Ostracoda from the Malayan region*. University of Utrecht. Available from <http://dspace.library.uu.nl/handle/1874/236218> [accessed 2 Jan. 2023].
- Latreille P.A. 1806. *Genera Crustaceorum et Insectorum: secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata. Tomus 1*. Koenig, Paris.  
<https://doi.org/10.5962/bhl.title.65741>
- Lee E.-H. & Paik K.-H. 1992. Late Cenozoic ostracod fauna and paleoenvironments of the marine sedimentary strata in the Cheju Island, Korea. *Paleontological Society of Korea Special Publication 1*: 121–160.
- Maddocks R.F. 1969. Recent ostracodes of the Family Pontocyprididae chiefly from the Indian Ocean. *Smithsonian Contributions to Zoology 7*: 1–56. <https://doi.org/10.5479/si.00810282.7>
- Mandelstam M.J. 1960. *Ostracoda*. In: Orlov J.A. (ed.) *Fundamentals of Palaeontology Vol. 8, The Arthropoda. Trilobitomorpha and Crustaceomorpha*: 264–421. Moscow.
- Martens K. & Horne D.J. 2009. Ostracoda. In: Likens G.E. (ed.) *Encyclopedia of Inland Waters*: 405–414. Elsevier, Oxford. <https://doi.org/10.1016/B978-012370626-3.00184-8>
- McKenzie K.G. & Pickett J.W. 1984. Environmental interpretations of Late Pleistocene ostracode assemblages from the Richmond River Valley, New South Wales. *Proceedings of the Royal Society of Victoria 96*: 227–242.
- Minhat F.I., Ghandhi S.M., Ahzan N.S.M., Haq N.A., Manaf O.A.R.A., Sabohi S.M., Lee L.H, Akhir M.F. & Abdullah M.M. 2021. The occurrence and distribution of benthic Foraminifera in tropical waters along the Strait of Malacca. *Frontiers in Marine Science 8*: 647531.  
<https://doi.org/10.3389/fmars.2021.647531>
- Moore R.C. 1961. *Treatise on Invertebrate Paleontology, Part Q, Arthropoda 3*. Geological Society of America and University of Kansas Press, Lawrence and Boulder.
- Montenegro M.E., Pugliese N. & Sciuto F. 2004. Shallow water ostracods near the Mae Khlong River mouth (NW Gulf of Thailand). *Bollettino della Società Paleontologica Italiana 43* (1–2): 225–234.
- Morkhoven Van F.P.C.M. 1963. *Post-paleozoic Ostracoda. Their Morphology. Taxonomy and Economic Use*. Elsevier Publishing Company, Amsterdam.
- Mostafawi N. 2003. Recent ostracods from the Persian Gulf. *Senckenbergiana Maritima 32* (1/2): 51–75. <https://doi.org/10.1007/BF03043085>
- Mostafawi N., Nabavi S.M.B. & Moghaddasi B. 2010. Ostracods from the Strait of Hormuz and Gulf of Oman, Northern Arabian Sea. *Revista Española de Micropaleontología 42* (2): 243–265.
- Müller G.W. 1894. Die Ostracoden des Golfes von Neapel und der angrenzenden Meeres Abschnilte. *Fauna und flora des golfes von Neapel 21*: 1–404. <https://doi.org/10.5962/bhl.title.7419>
- Negri M.P. 2009. Fossil mollusc-faunas: their bearing on the Holocene evolution of the lower central plain of Bangkok (Thailand). *Journal of Asian Earth Sciences 35*: 524–544.  
<https://doi.org/10.1016/j.jseaes.2009.04.003>
- Nishath N.M., Hussain S.M. & Rajkumar A. 2015. Distribution of Ostracoda in the sediments of the northwestern part of the Bay of Bengal, India-implications for microenvironment. *Journal of the Palaeontological Society of India 60* (2): 27–33.
- Noraswana N.F., Ramlan O. & Norashikin S. 2014. Distribution of recent Ostracoda in offshore sediment of selected stations in the Sulu Sea, Sabah. *Malays. Applied Biology 43* (2): 49–57.



- Nuamnim W., Sakha P. & Chaimani N. 2021. Lithostratigraphy of sediments at whale excavation site, in Ban Paew. Technical report. In: Suvapak I., Apsorn S., Thawatchai C., Prachya B., Siripond S., Denchak M., Angsumalin P., Sakda K., Veerachat V., Metha Y., Prodit N., Phornpen C. & Sirirat P. (eds) *Abstract Book of Geothai Webinar 2021*: 61–64. Royal Department of Mineral Resources of Thailand. Bangkok.
- Nützel A. & Kaim A. 2014. Diversity, palaeoecology and systematics of a marine fossil assemblage from the Late Triassic Cassian Formation at Settsass Scharte, N Italy. *Paläontologische Zeitschrift* 88 (4): 405–431. <https://doi.org/10.1007/s12542-013-0205-1>
- Oertli H.J. 1971. The aspect of ostracode faunas – A possible new tool in petroleum sedimentology. *Bulletin du Centre de Recherches de Pau-SNPA* 5: 137–151.
- Omar R., Faiz N.N. & Yusoff M.N.A. 2017. Recent benthic Ostracoda in offshore sediment of Pulau Perhentian, Terengganu. *Malaysian Applied Biology* 46 (1): 15–19.
- Paik K.H. 1977. Regionale Untersuchungen zur Verteilung der Ostracoden im Persischen Golf und im Golf von Oman. “Meteor” *Forschungsergebnisse, Reihe C: Geologie und Geophysik* 23: 37–76.
- Pugliese N., Montenegro M.E., Sciuto F. & Chaimanee N. 2006. Environmental monitoring through the shallow marine ostracods of Phetchaburi area (NW Gulf of Thailand). In: Coccioni R. & Marsili A. (eds) *Proceedings of the Second and Third Italian Meetings on Environmental Micropaleontology, Grzybowski Foundation Special Publication* 11: 85–90.
- Ramlan O. & Noraswana N.F. 2009. Distribution of ostracods in offshore sediment around Pulau Tioman, Pahang. *Malaysian Applied Biology* 38 (1): 11–19.
- Ramlan O. & Noraswana N.F. 2010. Distribution of recent Ostracoda in offshore sediment around Pulau Besar, Johor. *Sains Malaysiana* 39 (2): 199–207.
- Rau J.L. & Nutalaya, P. 1983. Geology of the Bangkok clay. *Geological Society of Malaysia Bulletin* 16: 99–116. <https://doi.org/10.7186/bgsm16198309>
- Robba E., Chaimanee N., Dheeradilok P., Jongkanjanasontorn Y., Piccoli G. & Boyd A.P. 1993. Late Quaternary Mollusca communities from the Bangkok Clay, Thailand. In: Thanasuthipitak T. (ed.) *Proceedings of the International Symposium on Biostratigraphy of Mainland Southeast Asia: Facies & Paleontology* 1: 427–437.
- Ruan P.H. 1989. Ostracoda. In: Hao Y.C. (ed.) *Quaternary Microbiotas and their Geological Significance from Northern Xisha Trench of South China Sea*. China University of Geosciences Press, Wuhan.
- Ruggieri G. 1953. Ostracodi del genere *Paijenborchella* vivienti nel Mediterraneo. *Atti della Società Italiana di Scienze Naturali (Milano)* 92: 10–14.
- Ruggieri G. 1967. Due Ostracofaune del Miocene Alloctono della Val Marecchia (Appenino Settentrionale). *Rivista Italiana di Paleontologia e Stratigrafia* 73 (1): 351–384.
- Saethien P. 2021. Protection and conservation management of Holocene whale skeleton from Am Pang Subdistrict, Ban Paew District, Samut Sakhon Province. In: Suvapak I., Apsorn S., Thawatchai C., Prachya B., Siripond S., Denchak M., Angsumalin P., Sakda K., Veerachat V., Metha Y., Prodit N., Phornpen C. & Sirirat P. (eds) *Abstract Book of Geothai Webinar 2021*: 207. Royal Department of Mineral Resources of Thailand. Bangkok.
- Sars G.O. 1866. Oversigt af Norges marine Ostracoder. *Forhandlinger i Videnskabs-Selskabet i Christiania* 1865 (1): 1–130.
- Sars G.O. 1922–1928. *An account of the Crustacea of Norway, with short descriptions and figures of all the species. Vol. 9. Ostracoda*. Bergen Museum, Oslo. <https://doi.org/10.5962/bhl.title.1164>

- Schneider G.F. 1971. Ostracodes from Quaternary deposits of North Vietnam. *Paleontologicheskij Zhurnal* 1971: 259–262.
- Siddiqui Q.A. 1971. Early Tertiary Ostracoda of the family Trachyleberididae from West Pakistan. *Bulletin of the British Museum (Natural History), Geology, Supplement* 9: 1–98. <https://doi.org/10.5962/p.310417>
- Sinsakul S. 2000. Late Quaternary geology of the lower central plain, Thailand. *Journal of Asian Earth Sciences* 18: 415–426. [https://doi.org/10.1016/S1367-9120\(99\)00075-9](https://doi.org/10.1016/S1367-9120(99)00075-9)
- Somboon J.R.P. 1988. Paleontological study of the recent marine sediments in the lower central plain, Thailand. *Journal of Southeast Asian Earth Sciences* 2: 201–210. [https://doi.org/10.1016/0743-9547\(88\)90031-1](https://doi.org/10.1016/0743-9547(88)90031-1)
- Somboon J.R.P. & Thiramongkol N. 1992. Holocene highstand shoreline of the Chao Phraya Delta, Thailand. *Journal of Southeast Asian Earth Sciences* 7: 53–60. [https://doi.org/10.1016/0743-9547\(92\)90014-3](https://doi.org/10.1016/0743-9547(92)90014-3)
- Songtham W., Watanasak M. & Insai P. 2000. Holocene marine crabs and further evidence of a sea-level peak at 6000 years BP in Thailand. In: Sinsakul S., Chaimanee N. & Tiyaipairach S. (eds) *Proceedings of the Thai-Japanese Geological Meeting, the Comprehensive Assessment on Impacts of Sea Level Rise*: 89–97. Department of Mineral and Resources, Bangkok.
- Songtham W., Musika S., Mildenhall D.C., Cochran U.A. & Kojevnikova D. 2015. Development of the lower central plain of Thailand with history of human settlements: evidence from pollen, spore and diatoms. *Journal of Geological Resources and Engineering* 2: 98–107.
- Swain F.M. 1955. Ostracoda of San Antonia Bay Texas. *Journal of Palaeontology* 29: 561–646.
- Sylvester-Bradley P.C. 1947. Some ostracod genotypes. *Annals and Magazine of Natural History, Series II* 13 (99): 192–199. <https://doi.org/10.1080/00222934608654541>
- Sylvester-Bradley P.C. 1948. The ostracod genus *Cythereis*. *Journal of Paleontology* 22 (6): 792–797.
- Szarek R., Kuhnt W., Kawamura H. & Kitazato H. 2006. Distribution of recent benthic foraminifera on the Sunda Shelf (South China Sea). *Marine Micropaleontology* 61: 171–195. <https://doi.org/10.1016/j.marmicro.2006.06.005>
- Tan C.W.J., Gouramanis C., Pham T.D., Hoang D.Q. & Switzer A.D. 2021. Ostracods as pollution indicators in Lap An Lagoon, central Vietnam. *Environmental Pollution* 278: 116762. <https://doi.org/10.1016/j.envpol.2021.116762>
- Tanabe S., Saito Y., Sato Y., Suzuki Y., Sinsakul S., Tiyaipairach S. & Chaimanee N. 2003. Stratigraphy and Holocene evolution of the mud-dominated Chao Phraya delta, Thailand. *Quaternary Science Reviews* 22: 789–807. [https://doi.org/10.1016/S0277-3791\(02\)00242-1](https://doi.org/10.1016/S0277-3791(02)00242-1)
- Tanaka G., Komatsu T. & Phong N. 2009. Recent ostracod assemblages from the northeastern coast of Vietnam and the biogeographical significance of the euryhaline species. *Micropaleontology* 55: 365–382. <https://doi.org/10.47894/mpal.55.4.03>
- Tanaka G., Matsushima Y. & Maeda H. 2012. Holocene ostracods from the borehole core at Oppama Park, Yokosuka City, Kanagawa Prefecture, central Japan: paleoenvironmental analysis and the discovery of a fossil ostracod with three-dimensionally preserved soft parts. *Paleontological Research* 16: 1–18. <https://doi.org/10.2517/1342-8144-16.1.001>
- Tanaka G., Henmi Y., Masuada T., Moriwaki H., Komatsu T., Zhou B., Maekawa T., Niiyama S., Nguyen P.D., Doan H.D. & Ikeya N. 2019. Recent ostracod distribution in western Kyushu, Japan, related to the migration of Chinese continental faunal elements. *Marine Micropaleontology* 146: 1–38. <https://doi.org/10.1016/j.marmicro.2018.12.002>

- Teeter J.W. 1975. Distribution of Holocene marine Ostracoda from Belize. *American Association of Petroleum Geologists Studies in Geology* 2: 400–499.
- Wang H., Zhang H., Cao M. & Horne D. 2018. Holocene ostracods from the Hang Hau Formation in Lei Yue Mun, Hong Kong, and their palaeoenvironmental implications. *Alcheringa: An Australasian Journal of Palaeontology* 43 (2): 320–333. <https://doi.org/10.1080/03115518.2018.1511830>
- Wang P.X., Zhang J.J., Zhao Q.H., Min Q.B., Bian Y.H., Zheng L.F., Cheng X.R. & Chen R.H. 1988. *Foraminifera and Ostracoda in Bottom Sediments of the East China Sea*. China Ocean Press, Beijing.
- Whatley R.C. 1983. The application of Ostracoda to palaeoenvironmental analysis. In: Maddocks R.F. (ed.) *Proceedings of the Eighth International Symposium on Ostracoda*: 51–77. University Houston Geoscience, Houston.
- Whatley R.C. & Zhao Q. 1987. A revision of Brady's 1869 study of the Ostracoda of Hong Kong. *Journal of Micropalaeontology* 6: 21–29. <https://doi.org/10.1144/jm.7.1.21>
- Whatley R.C. & Zhao Q. 1988a. Recent Ostracoda of the Malacca Straits, Part 1. *Revista Española de Micropaleontología* 19: 327–366.
- Whatley R.C. & Zhao Q. 1988b. Recent Ostracoda of the Malacca Straits. Part 2. *Revista Española de Micropaleontología* 20: 5–73.
- Woodroffe C.D. 2000. Deltaic and estuarine environments and their Late Quaternary dynamics on the Sunda and Sahul shelves. *Journal of Asian Earth Sciences* 18: 393–413. [https://doi.org/10.1016/S1367-9120\(99\)00074-7](https://doi.org/10.1016/S1367-9120(99)00074-7)
- Yamada K., Irizuki T. & Tanaka Y. 2002. Cyclic sea-level changes based on fossil ostracode faunas from the upper Pliocene Sasaoka Formation, Akita Prefecture, northeast Japan. *Palaeogeography, Palaeoclimatology, Palaeoecology* 185: 115–132. [https://doi.org/10.1016/S0031-0182\(02\)00281-X](https://doi.org/10.1016/S0031-0182(02)00281-X)
- Yamada K., Terakura M. & Tsukawaki S. 2014. The impact on bottom sediments and ostracods in the Khlong Thom River mouth following the 2004 Indian Ocean tsunami. *Paleontological Research* 18 (2): 104–117. <https://doi.org/10.2517/2014PR011>
- Yan D., Wünnemann S. & Gao S. 2020. Early Holocene tidal flat evolution in a western embayment of East China Sea in response to sea level rise episodes. *Quaternary Science Reviews* 250: 106642. <https://doi.org/10.1016/j.quascirev.2020.106642>
- Yasuhara M. & Seto K. 2006. Holocene relative sea-level change in Hiroshima Bay, Japan: a semi-quantitative reconstruction based on ostracodes. *Paleontological Research* 10: 99–116. <https://doi.org/10.2517/prpsj.10.99>
- Yim S.S.W., Zhang L.X. & Wang Q. 1988. Holocene ostracoda in Hong Kong and their palaeoenvironmental significance. In: *Proceedings 2<sup>nd</sup> Conference on Palaeoenvironment of East Asia from the Mid-Tertiary Vol. II Oceanography, Palaeozoology and Palaeoanthropology*: 810–827. Centre of Asian Studies, University of Hong Kong.
- Zhao Q. & Wang P. 1988. Distribution of modern Ostracoda in the shelf seas off China. In: Hanai T., Ikeya N. & Ishizaki K. (eds) *Evolutionary Biology of Ostracoda: its Fundamentals and Applications*: 805–821. Kodansha, Tokyo. [https://doi.org/10.1016/S0920-5446\(08\)70223-1](https://doi.org/10.1016/S0920-5446(08)70223-1)
- Zhao Q.H. & Wang P.X. 1990. Modern Ostracoda in shelf seas off China: zoogeographical zonation. *Oceanologia et Limnologia Sinica* 21: 458–464.
- Zhao Q.H. & Whatley R.C. 1988. The genus *Neomonoceratina* (Crustacea: Ostracoda) from the Cainozoic of the west Pacific margins. *Acta Oceanologica Sinica* 7: 562–577.



Zhao Q.H. & Whatley R.C. 1989. Recent podocopid Ostracoda of the Sedili River and Jason Bay, southeastern Malay Peninsula. *Micropaleontology* 35: 168–187. <https://doi.org/10.2307/1485467>

*Manuscript received: 4 August 2021*

*Manuscript accepted: 10 December 2021*

*Published on: 31 January 2023*

*Topic editor: Tony Robillard*

*Section editor: Christian de Muizon*

*Desk editor: Pepe Fernández*

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum, Prague, Czech Republic.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2022

Band/Volume: [0856](#)

Autor(en)/Author(s): Chitnarin Anisong, Forel Marie-Béatrice, Tepnarong Prachya

Artikel/Article: [Holocene ostracods \(Crustacea\) from a whale-fall excavation site from the Chao Phraya delta, Central Thailand 120-151](#)