

the International Union of Biological Sciences. London (ITZN), pp. i-xxix, 1-306. TRUTNAU, L. (1994): Terraristik. Stuttgart (Ulmer), 320 pp.

KEY WORDS: Reptilia: Testudines: Testudinidae: *Testudo weissingeri* BOUR, 1995: taxonomy, valid name

SUBMITTED: February 9, 2004

AUTHOR: Jarmo PERÄLÄ, Ph. D., Department of Biological and Environmental Sciences, University of Helsinki, PO Box 65 (Biocenter 3, Viikinkaari 1), FIN-00014 Finland. <jarmo.perala@helsinki.fi >

New data on fringe-toed lizards of the *Acanthodactylus scutellatus* group in Mauritania

According to CROCHET et al. (2003), five species of the *Acanthodactylus scutellatus* group occur in Mauritania: *Acanthodactylus senegalensis* CHABANAUD, 1918, *A. dumerili* (MILNE EDWARDS, 1829), *A. longipes* BOULENGER, 1918, *A. aureus* GÜNTHER, 1903 and *A. taghitensis* GENIEZ & FOUCART, 1995.

In November 1995 one of the authors (PL) found and photographed a large male of *Acanthodactylus scutellatus audouini* BOULENGER, 1918 (fig. 1), east of Ouadane (ca. 20°59'N / 11°26'W), in the north-eastern Adrar area of Mauritania (fig. 2). This specimen can be identified undoubtedly as *A. s. audouini* by its large size, the strongly



Fig. 1: Male *Acanthodactylus scutellatus audouini* BOULENGER, 1918 from west of Ouadane (Mauritania) (Photo: P. LLUCH).

swollen base of its tail, the large dark spots on its pileus, the strongly contrasted black and disjointed reticulation on its dorsum and its relatively large and carinate dorsal scales. *Acanthodactylus s. audouini* was previously known only from the eastern and central Sahara, (north-eastern Sudan and Egypt in the east to south-eastern Tunisia in the north and eastern Mali in the south-west (CROCHET et al. 2003). The present observation adds a new country (Mauritania) to the range of *A. scutellatus* and extends its distribution 1,000 km towards the West. The only species of the




Fig. 2: Western part of the distribution of *Acanthodactylus scutellatus audouini* BOULENGER, 1918 based on CROCHET et al. 2003. The asterisk represents the new Mauritanian locality.



Fig. 3: Adult male *Acanthodactylus taghitensis* GENIEZ & FOUCART, 1995 from Ouan n'Namour, Mauritania (BEV.7785) (Photo: PH. GENIEZ).




Fig. 4: Distribution of *Acanthodactylus taghitensis* GENIEZ & FOUCART, 1995. The list of localities (new record marked by an asterisk) is available in the text.

Acanthodactylus scutellatus group so far known from the Adrar area was *A. longipes*. With six species, Mauritania became the richest country in terms of representatives of the *Acanthodactylus scutellatus* group.

Acanthodactylus taghitensis is a little known species closely related to *A. aureus* but with a non-coastal distribution. On 29 November 2001 one of the authors (PL) caught an adult male *A. taghitensis* (fig. 3) at Ouan n'Namour, Mauritania, in a reg formation from the Northern Tagenzé of Guelb Richat (21°22'51"N / 11°24'36"W). This specimen is preserved at Montpellier (France) in the collection of the Laboratoire de Biogéographie et Ecologie des Vertébrés de l'École Pratique des Hautes Etudes (n° BEV.7785). The present observation extends the known distribution of *A. taghitensis* 200 km towards the south-east. This species is thus presently known from five localities (fig. 4):

- 5 km from Taghit towards Beni Abbès (Algeria) [30°53'53"N / 2°4'51"W] (GENIEZ & FOUCART 1995, locality 1 in fig. 4);
- 36 km from Taghit towards Beni Abbès (Algeria) [30°39'15"N / 2°11'4"W] (GENIEZ & FOUCART 1995, locality 2 in fig. 4);
- El Gor near Tindouf (Algeria) [27°28'N / 7°56'W], (DONAIRE et al. 2000, locality 3 in fig. 4);
- Fderik (= Fort-Gouraud) (Mauritania) [22°41'20"N / 12°42'34"W] (CROCHET et al. 2003, locality 4 in fig. 4);

- Northern Tagenzé, Ouan n'Namour (Mauritania) [21°22'51"N / 11°24'36"W], (present work, locality 5 in fig. 4).

Acanthodactylus taghitensis may be present in eastern parts of Western Sahara (under Moroccan administration) as well but no records are available for this country yet (GENIEZ et al. in press).

REFERENCES: CROCHET, P.-A. & GENIEZ, P. & INEICH, I. (2003): A multivariate analysis of the *Acanthodactylus scutellatus* group (Squamata: Lacertidae): systematics and biogeographical implications.- Zool. J. Linnean Soc., London: 137 (1): 117-155. DONAIRE, D. & MATEO, J.A. & HASI, M. & GENIEZ, PH. (2000): Nuevos datos sobre la fauna reptiliana de la Hammada de Tindouf (Argelia).- Bol. Asoc. Herpetol. Española, Barcelona; 11: 8-12. GENIEZ, PH. & FOUCART, A. (1995): Un nouvel *Acanthodactylus* en Algérie: *Acanthodactylus taghitensis* n.sp. (Reptilia, Sauria, Lacertidae).- Bull. Mus. natn. Hist. nat., Paris: (ser. 4) 17: 3-9. GENIEZ, PH. & MATEO, J.-A. & GENIEZ, M. & PETHERS, J. (in press): The amphibians and reptiles of the Western Sahara (former Spanish Sahara) and adjacent regions. Frankfurt am Main, Chimaira. SALVADOR, A. (1982): A revision of the lizards of the genus *Acanthodactylus* (Sauria: Lacertidae).- Bonner Zoologische Monographien, Bonn; 16: 1-167.

KEY WORDS: Reptilia: Squamata: Lacertidae: *Acanthodactylus scutellatus audouini*, *Acanthodactylus taghitensis*, Sahara, Mauritania, new country record, chorology.

SUBMITTED: October 20, 2003.

AUTHORS: Philippe GENIEZ, Laboratoire de Biogéographie et Ecologie des Vertébrés de l'École Pratique des Hautes Etudes (E.P.H.E.), Université Montpellier II, F-34095 Montpellier cedex 5, France < geniez@univ-montp2.fr >; Pascal LLUCH, Mas Salançon, F-30570 Notre-Dame de la Rouvière, France < pascal@hommes-et-montagnes.fr >

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2004

Band/Volume: [17_1_2](#)

Autor(en)/Author(s): Geniez Philippe, Lluch Pascal

Artikel/Article: [New data on fringe-toed lizards of the *Acanthodactylus scutellatus* group in Mauritania 88-89](#)