

Endococcus incrassatus, a new lichenicolous fungus (*Dothideales*)

J. ETAYO

Navarro Villoslada 16-3° dcha.

E-31003 Pamplona, Spain

Email jetayosa@pnte.cfnavarra.es

O. BREUSS

Naturhistorisches Museum Wien, Botanische Abteilung

Burgring 7, Postf. 417

A-1014 Wien, Austria

Email obreuss@g9-asn-wien.ac.at

Received 28. 7. 2001

Key words: Lichenicolous Ascomycetes, *Dothideales*, *Endococcus*, *Endococcus incrassatus* spec. nova. – Taxonomy, systematics. – Mycoflora of Mexico.

Abstract: *Endococcus incrassatus*, a parasite on the lichen *Placiopsis cinerascens*, is described as new.

Zusammenfassung: *Endococcus incrassatus* wird als Parasit auf *Placiopsis cinerascens* neu beschrieben.

Endococcus NYL. is a Dothidealean genus of lichenicolous fungi comprising about 15 species occurring mostly on saxicolous crustose lichens. It is characterized by perithecioid ascomata, a hemiamyloid hymenial gel, soon gelatinizing paraphysoids, brown, 1-septate ascospores, and the lichenicolous habit. After the study by HAWKSWORTH (1979), two important contributions dealed with the genus: TRIEBEL (1989) presented a rather broad species concept joining several entities growing on very different hosts, whereas SÉRUSIAUX & al. (1999) adopted a much narrower species concept for *Endococcus* resulting in a larger number of mostly host-specific taxa. As there are significant differences in spore size and several fine aspects of ascomatal anatomy between taxa on different hosts, this narrower species concept seems to be more appropriate.

During a collecting trip to Baja California in the course of the Sonoran Desert Lichen Flora project under the leadership of Prof. Dr THOMAS H. NASH (Tempe) in 1997, the second author collected a specimen of *Placiopsis cinerascens* (NYL.) BREUSS that is heavily infested by an *Endococcus* species which is described as new below.

***Endococcus incrassatus* ETAYO & BREUSS, spec. nova (Fig. 1)**

Diagnosis latina: Ascomata superficiales in thallo *Placiidiopsis cinerascentis*, subcylindrica, apicibus irregulariter incrassatis. Ascosporeae maturitate uniseptatae, fuscae, laeves, septo lamella media atro-brunnea, multiguttulatae, 10,5-14 x 6-8 µm.

Typus: Mexico, Baja California, lower slopes of Cerro los Enjambres ca. 2 km WNW of Laguna Chapala, Vizcaino Region of the Sonoran Desert, 29°23'39''N, 114°23'30''W, 700 m s. m., on *Placiidiopsis cinerascens*, 30. 12. 1997, O. BREUSS 14447 (LI, holotype).

Merkmale:

Ascomata perithecioid, lichenicolous, dispersed in the thallus of *Placiidiopsis cinerascens*, at first immersed, soon superficial, black, non-setose, irregularly subcylindrical, with thickenings in the upper part, 120-200 µm in diameter. Perithecial wall brown, lighter at the base and the inner parts, K+ blackish, apically with irregular thickenings, 20-25 µm thick at the base, 40-60 µm above, paraplectenchymatous, of several rows of tangentially compressed cells 7-11 x 2-3 µm, the inner ones with thin and hyaline walls, the outer ones with thick, dark walls. Periphysoids of two kinds: the lower ones long and septate, thin-walled, 16-22 x 1 µm, the upper ones short, simple, with thicker walls, 7-12 x 2(-4) µm. Paraphysoids simple, septate, soon disappearing. Hymenial gel I+ orange, KI+ blue. Asci clavate, 50-60 x 8-10 µm, 8-spored, wall distinctly thickened apically. Ascospores ellipsoidal to ovoid, with obtuse (rarely pointed) ends, soon brownish, dark brown when mature, at first non-septate and ornamented with very small granulation, then one-septate and smooth, without a distinct perispore, not or slightly constricted at the septum, cells unequal in size and shape, septum dark brown with a darker median lamella, each cell containing numerous small lipid drops, 10,5-14 x 6-8 µm. Anamorph unknown.

Notes:

Endococcus incrassatus is well distinguished from other species of the genus by its almost superficial ascomata with irregular thickenings in their ostiolar region, dark brown ascospores with a thick lamella in the septum and in growing on the terricolous squamulose lichen *Placiidiopsis cinerascens*.

References:

- HAWKSWORTH, D. L., 1979: Studies in the genus *Endococcus* (*Ascomycotina, Dothideales*). – Bot. Not. 132: 283-290.
SÉRUSIAUX, E., DIEDERICH, P., BRAND, A. M., VAN DEN BOOM, P., 1999: New or interesting lichens and lichenicolous fungi from Belgium and Luxembourg. VIII. – Lejeunia, n. s. 162: 1-95.
TRIEBEL, D., 1989: Lecideicole Ascomyceten. – Biblioth. Lichenol. 35: 1-278.




Fig. 1. *Endococcus incrassatus* (holotype). A Ascomata. B Section of apical part of ascromatal wall with periphysoids. C Section of ascoma showing apical thickenings. D Ascospores. – Bars: A = 100 µm, B = 5 µm, C = 150 µm, D = 10 µm.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Österreichische Zeitschrift für Pilzkunde](#)

Jahr/Year: 2001

Band/Volume: [10](#)

Autor(en)/Author(s): Etayo Javier, Breuss Othmar

Artikel/Article: [Endococcus incrassatus, a new lichenicolous fungus
\(Dothideales\). 315-317](#)