

Phyton (Horn, Austria)	Vol. 37	Fasc. 1	61-70	10. 9. 1997
------------------------	---------	---------	-------	-------------

**New Species or Interesting Records of Foliicolous  
Lichens. III.  
*Arthonia crystallifera* spec. nova (Lichenized  
Ascomycetes: *Arthoniaceae*), with a World-Wide Key  
to the Foliicolous *Arthoniaceae***

By

Lidia Itati FERRARO\*) and Robert LÜCKING\*\*)

With 2 Figures

Received December 23, 1996

Keywords: *Arthonia crystallifera* L. I. FERRARO & R. LÜCKING spec. nova, *Arthonia*, *Arthoniaceae*, *Eremothecella cingulata* (R. SANT.) L. I. FERRARO & R. LÜCKING comb. nova; *Ascomycetes*. – Taxonomy. – Foliicolous lichens.

Summary

FERRARO L. I. & LÜCKING R. 1997. New species or interesting records of foliicolous lichens. III. *Arthonia crystallifera* spec. nova (Lichenized Ascomycetes: *Arthoniaceae*), with a world-wide key to the foliicolous *Arthoniaceae*. *Phyton* (Horn, Austria) 37 (1): 61-70, 2 figures. – English with Spanish and German summary.

A new foliicolous species of *Arthonia* from northern Argentina and Brazil is described: *A. crystallifera* L. I. FERRARO & R. LÜCKING is distinguished from other foliicolous species of the genus by the presence of large, conspicuous, shiny crystals on the ascocarp surface. It is further characterized by the brownish colour of its ascocarps, the macrocephalic, 2-septate, colourless ascospores, and the radiate phyco-biont cells, indicating a close relationship to *A. accolens* STIRT. In order to facilitate the delimitation of foliicolous *Arthoniaceae*, a key to all hitherto known species (28) is presented. The new combination *Eremothecella cingulata* (R. SANT.) L. I. FERRARO & R. LÜCKING comb. nova is proposed (Bas.: *Arthothelium cingulatum* R. SANT.).

\*) .... Lidia Itati FERRARO, Instituto de Botanica del Nordeste (IBONE), C. C. 209, 3400 Corrientes, Argentina

\*\*\*) .... Dr. Robert LÜCKING, Abteilung Spezielle Botanik, Universität Ulm, D-89069 Ulm, Germany

## Resumen

FERRARO L. I. & LÜCKING R. 1997. Nuevas especies o comunicaciones interesantes de líquenes foliícolas. III. *Arthonia crystallifera* spec. nova (Ascomycetes Liquenizadas: *Arthoniaceae*), con una clave mundial para las especies foliícolas de *Arthoniaceae*. *Phyton* (Horn, Austria) 37 (1): 61–70, 2 figuras. – Inglés con resumen Español y Alemán.

Se describe una nueva especie foliícola del género *Arthonia* del norte de Argentina y de Brazil: *A. crystallifera* L. I. FERRARO & R. LÜCKING se distingue de otras especies foliícolas del género por la presencia de cristales brillantes grandes en la superficie de los ascocarpos. Además, está caracterizada por el color marrón de sus ascocarpos, por las esporas macrocefálicas, 2-septadas y incoloradas, y por las células radiantes del ficobionte, lo que indica una relación con *A. accolens* STIRT. Para facilitar la delimitación de las *Arthoniaceae* foliícolas, se presenta una clave para todas las especies conocidas (28) hasta la fecha. Además, se propone la nueva combinación *Eremothecella cingulata* (R. SANT.) L. I. FERRARO & R. LÜCKING comb. nova (Bas.: *Arthothelium cingulatum* R. SANT.).

## Zusammenfassung

FERRARO L. I. & LÜCKING R. 1997. Neue Arten oder interessante Nachweise foliicoler Flechten. III. *Arthonia crystallifera* spec. nova (lichenisierte Ascomyceten: *Arthoniaceae*), mit einem weltweiten Schlüssel für foliicole *Arthoniaceae*. – *Phyton* (Horn, Austria) 37(1): 61–70, 2 Abbildungen. – Englisch mit spanischer und deutscher Zusammenfassung.

Eine neue foliicole *Arthonia*-Art aus N Argentinien und Brasilien wird beschrieben: *A. crystallifera* L. I. FERRARO & R. LÜCKING ist von anderen foliicolen Arten der Gattung durch den Besitz großer, deutlicher, glänzender Kristalle auf der Ascocarp-Oberfläche verschieden. Weiters sind bräunliche Ascocarp-Farbe, macrocephalische, 2-septierte, farblose Ascosporen und strahlige Phycobionten-Zellen charakteristisch, was auf nahe Verwandtschaft mit *A. accolens* STIRT deutet. Um die Abgrenzung foliicoler *Arthoniaceae* zu erleichtern, wird ein Schlüssel aller bisher bekannten Arten (28) vorgelegt. Die Neukombination *Eremothecella cingulata* (R. SANT.) L. I. FERRARO & R. LÜCKING (Bas.: *Arthothelium cingulatum* R. SANT.) wird vorgeschlagen.

## 1. Introduction

Although the *Arthoniaceae* play a major role in cryptogam communities of tropical forest ecosystems, their diversity on leaves is moderate as compared to other groups, such as the *Trichotheliaceae*, *Gomphillaceae*, or *Pilocarpaceae* (SANTESSON 1952; FARKAS & SIPMAN 1993). Nevertheless, particularly in lowland rain forests, members of *Arthoniaceae* are an important and conspicuous element of the foliicolous lichen flora (LÜCKING 1992a, b, 1995, 1997a). The present paper deals with the new species *Arthonia crystallifera*, which was found on leaves in a rather restricted area in northern Argentina and southern Brazil. The species, which is most closely related to *A. accolens* STIRT, is particularly characterized by the

large, conspicuous crystals which are deposited on the ascocarp surface and easily visible under lens magnification.

In addition to the 19 foliicolous *Arthoniaceae* treated by SANTESSON 1952, together with *A. crystallifera* ten new species and one variety have been described in recent years (XAVIER FILHO 1964; UPADHYAY 1964; BATISTA & MAIA 1967; KANTVILAS & VĚZDA 1988; APTROOT & SIPMAN 1991; LÜCKING 1991, 1995; BECKER & LÜCKING 1995; LÜCKING & LÜCKING 1995; LÜCKING & al. 1997). Three species, viz. *A. lividofusca* MÜLL. ARG., *A. anisolocularis* L. XAVIER & TALTASSE, and *A. opegraphina* R. LÜCKING, have later been synonymized with *A. aciniformis* STIRT., *A. cyanea* MÜLL. ARG., and *A. orbygniae* (H. B. P. UPADHYAY) MATZER, respectively (LÜCKING 1995; MATZER 1996; LÜCKING & al. 1997). Furthermore, considerable generic rearrangements have been made with the reinstatement of *Eremothecella* and the introduction of *Amazonomyces* as new generic name (BATISTA & PERES 1964; SÉRUSIAUX 1992; LÜCKING & al. 1997; THOR & al. 1997). Thus, while SANTESSON 1952 listed two species of *Stirtonia*, one species of *Cryptothecia*, 15 species of *Arthonia*, and one species of *Arthothelium*, in the present paper, one species of *Cryptothecia*, two species of *Amazonomyces*, 20 species and one variety of *Arthonia*, and five species of *Eremothecella* are accepted, leaving the genera *Stirtonia* and *Arthothelium* without foliicolous representants. In order to facilitate access to all taxonomic and systematic alterations, an key is presented to all world-wide known foliicolous, lichenized species of *Arthoniaceae*.

## 2. *Arthonia crystallifera* L. I. FERRARO & R. LÜCKING spec. nova.

Thallus epiphyllus, tenuis, continuous vel dispersus, 5–15 mm diam., 10–15 µm crassus, flavovirens, opacus; prothallo nullo. Apothecia rotundata, 0.3–0.5 mm lata, 35–45 µm alta, leviter prominentia, fusca, marginibus crystallis nitidis instructa, K+ rubrofusca. Stratum hypotheciodeum rubrofuscum. Stratum ascigerum incoloratum vel leviter fuscum. Asci ovati, 20–30 × 15–20 µm. Ascopora 8-nae, 2-septatae, oblongae, macrocephalae, incoloratae, 9–14 × 2.5–3 µm. Pycnidia rotundata, 0.1 mm lata., appanata vel leviter conica, fusca; conidia simplices, ellipsoides, 2.5–3 × 0.7 µm (microconidia?) vel bacillares, 4–6 × 1–1.5 µm (macroconidia?). Alga ad genus *Phycopeltis* pertinens, cellulis rectangularibus, 12–15 × 3–5 µm.

Typus: Argentina. Prov. Misiones: Depto. Iguazú, Isla Grande frente al Puerto Canoas, sobre hojas de *Pilocarpus pennatifolius* (*Rutaceae*), III. 1995, L. FERRARÓ, V. MARUÑAK & R. VANNI 5137 (CTES, holotypus; ULM, isotypus).

Thallus (Fig. 1) rounded or irregular in outline, continuous or marginally dispersed, 5–15 mm diam., 10–15 µm thick, yellowish-green. No visible prothallus, but sometimes a thin brownish zone when bordered by

other lichens. Ascocarps (Fig. 1) rounded, 0.3–0.5 mm diam., 35–45  $\mu\text{m}$  high, chestnut-brown, K+ slightly reddish brown (better observed in microscopic sections), their surface especially in lateral parts encrusted with large, conspicuous, very nitidous,  $\pm$  tetraedric crystals. Hypothecoid layer 5  $\mu\text{m}$  high, reddish brown. Ascigerous layer 25–35  $\mu\text{m}$  high, colourless to pale brown. Asci (Fig. 2) ovate, 20–30  $\times$  15–20  $\mu\text{m}$ . Ascospores 8 per ascus, 2-septate, colourless, macrocephalic, 9–14  $\times$  2.5–3  $\mu\text{m}$ . Pycnidia rounded, 0.1 mm diam., applanate to slightly conical, brown. Conidia (Fig. 2) simple, either ellipsoid and 2.5–3  $\times$  0.7  $\mu\text{m}$  (microconidia?) or bacillar, 4–6  $\times$  1–1.5  $\mu\text{m}$ . Phycobiont a species of *Phycopeltis*, with rectangular, 12–15  $\times$  3–5  $\mu\text{m}$  large cells, forming continuous, radiate plates, encrusted with scattered crystallites.




Fig. 1. *Arthonia crystallifera* (isotypes). General habit; note the large crystals on the ascocarp surface. Scale = 1 mm.

*A. crystallifera* is easily recognized by its large, brilliant crystals distributed on the lateral surface of the ascocarps. Although the production of crystals, especially calcium-oxalate, is rather frequent in foliicolous lichens, no species is hitherto known with a type of crystals similar to that of *A. crystallifera*. Particular, needle-shaped crystals on the lateral apothecial surface are known from *Echinoplaca furcata* and *E. verrucifera* (SÉRUSIAUX 1989; LÜCKING 1997b), but their structure and chemistry is certainly very different from that of the crystals found in *A. crystallifera*.

The brownish ascocarps, the 2-septate, macrocephalic ascospores, the pycnidia, and the phycobiont suggest a close relationship of the new species to *A. accolens* STIRT. However, the very slight reddish K-reaction of the ascocarps, which is best observed under the microscope, points also towards *A. mira* R. SANT. The latter has slightly larger ascospores, and the ascocarps are laterally orange brown and react K+ deep purple.




Fig. 2. *Arthonia crystallifera* (holotype). (A) Hymenium with asci and large crystals deposited in the epithecioid layer; (B) Ascospores. (C) Conidia. (D) Cells of the phycobiont. (E) Crystals from the ascocarp surface. Scale = 10 µm.

The new species seems to be restricted to northern Argentina and adjacent areas and was found in humid spots with rather high foliicolous lichen diversity. Interestingly, in both Corrientes and Misiones, one of the identified phorophytes belonged to the same species, viz. the *Rutaceae* *Pilocarpus pennatifolius*.

Additional specimens examined: Argentina. Prov. Misiones: Depto. Iguazú, Parque Nacional Iguazú, Isla Grande frente al Puerto Canoas, foliícola sobre hojas de *Pilocarpus pennatifolius* (*Rutaceae*), III. 1995, L. FERRARO, V. MARUŃAK & R. VANNI 5140 (CTES, ULM); *ibid.*, L. FERRARO & al. 5143 (CTES); *ibid.*, Paseos Inferiores, sobre *Sorocea bonplandii* (*Moraceae*), II. 1996, R. VANNI, D. KURTZ, O. POPOFF & J. HERRERA 3640 (CTES). – Prov. Corrientes: Depto. Ituzaingó, camino a San Carlos, sobre *Pilocarpus pennatifolius* (*Rutaceae*), 1980, M. M. Arbo & al. 2394 (CTES, LG). – Brazil: Santa Catarina, Ytayubá, Restinga Municipio Barra Velha, sobre hojas de *Rudgea villiflora* (*Rubiaceae*), 1988, A. KRAPOVICKAS & al. 43558b (CTES).

### 3. Key to the foliicolous, lichenized *Arthoniaceae*

The following key includes all presently known species of foliicolous, lichenized *Arthoniaceae* (28 species). The various lichenicolous taxa are not included; the worker is here referred to the keys provided by MATZER 1996. For practical reasons, the key is artificially structured; a natural key



- upright campylidia; conidia septate, filiform,  $30-40 \times 1 \mu\text{m}$ ; Neotropics and tropical Australasia . . . . . *Arthonia aciniformis* STIRT.  
[Syn.: *Arthonia lividofusca* MÜLL. ARG. (LÜCKING 1995)]
- 8 Ascospores microcephalic or isolocular, often with one of the median cells slightly enlarged (*Opegrapha*-type), 3-septate; ascocarps (except for *Arthonia obesa*) black, very small (0.1–0.2 mm diam.) . . . . . 9
- 8\* Ascospores macrocephalic, with the distal cell distinctly enlarged (*Arthonia*-type), 2–3(–5)-septate; ascocarps variously coloured, medium-sized to large (0.3–1.0 mm diam.) . . . . . 12
- 9 Ascocarps dark brown with a purplish tinge, often with a thin, pruinose marginal cover, rather large (0.5–1.5 mm diam.); phycobiont cells in irregularly net-like plates; Neotropics and tropical Africa . . . . . *Arthonia obesa* (MÜLL. ARG.) R. SANT.
- 9\* Ascocarps black, non-pruinose, very small (0.1–0.2 mm diam.); phycobiont cells in radiate plates . . . . . 10
- 10 Ascospores very narrowly fusiform,  $18-24' \times 2-3.5 \mu\text{m}$ , 6–9 times as long as broad; Neotropics . . . . . *Arthonia epidendri* (Rehm) R. SANT.
- 10\* Ascospores fusiform-ellipsoid,  $8-16 \times 3-5 \mu\text{m}$ , 2.5–4 times as long as broad . . . . . 11
- 11 Ascospores  $8-12 \times 3-4 \mu\text{m}$ ; pantropical . . . . .  
. . . . . *Arthonia nigratula* (MÜLL. ARG.) R. SANT.
- 11\* Ascospores  $14-16 \times 4-5 \mu\text{m}$ ; Vietnam . . . . .  
. . . . . *Arthonia microcephala* VĚZDA
- 12 Ascospores predominantly 2-septate. . . . . 13
- 12\* Ascospores predominantly 3(–5)-septate. . . . . 22
- 13 Mature ascospores greyish brown . . . . . 14
- 13\* Mature ascospores colourless. . . . . 17
- 14 Ascocarps brownish but marginally with an orange tinge, K+ purple. . . . . *Arthonia mira* R. SANT.
- 14\* Ascocarps dark brown to black, K or K+ greenish. . . . . 5
- 15 Thallus with scattered, yellowish to pale reddish verrucae; tropical Africa . . . . . *Arthonia flavoverrucosa* BECKER & R. LÜCKING 1995
- 15\* Thallus smooth . . . . . 16
- 16 Ascocarps blackish brown to black, non-pruinose (very rarely with a slight marginal pruina); ascospores  $11-19(-24) \times (3-4)4.5-6.5 (-9) \mu\text{m}$ , 2.5–4 times as long as broad; pantropical . . . . .  
. . . . . *Arthonia trilocularis* MÜLL. ARG.
- 16\* Ascocarps dark brown with a purplish tinge, and with a thin marginal pruina; ascospores  $13-19 \times 2.5-4.5 \mu\text{m}$ , 3–5 times as long as broad; Paleotropics . . . . .  
. . . . . *Arthonia fuscocyanea* BECKER & R. LÜCKING 1995
- 17 Ascocarps brown with an orange margin or orange red, K+ purple 18

- 17\* Ascocarps light to dark brown or with a bluish tinge, K- . . . . . 19
- 18 Ascocarps orange-red to dark red; tropical Australasia . . . . .  
 . . . . . *Arthonia ramosii* (RÄS.) R. SANT.
- 18\* Ascocarps brown but with the marginal parts having an orange  
 tinge; Neotropics . . . . . *Arthonia mira* R. SANT.
- 19 Ascocarps with numerous large crystals on their lateral surface,  
 easily visible under lens magnification; northern Argentina and  
 Brazil . . . . .  
 . . . . . *Arthonia crystallifera* L. I. FERRARO & R. LÜCKING (this paper)
- 19\* Ascocarp surface without such crystals . . . . . 20
- 20 Ascocarps greenish brown to bluish grey, with a white pruina gi-  
 ving them a bluish appearance . . . . .  
 . . . . . *Arthonia cyanea* MÜLL. ARG. var. *cyanea* p.p.
- 20\* Ascocarps light to dark brown, non-pruinose . . . . . 21
- 21 Ascocarps light to dark brown, up to 1.3 mm diam., sharply deli-  
 mited; pantropical . . . . . *Arthonia accolens* STIRT.
- 21\* Ascocarps light brown, up to 0.5 mm diam., margins diffuse; tropi-  
 cal Australasia . . . . . *Arthonia lividula* VAIN.
- 22 Ascocarps greenish brown to bluish grey, with a white pruina  
 giving them a bluish appearance. . . . . 23
- 22\* Ascocarps blackish brown to black, non-pruinose. . . . . 24
- 23 Ascospores regularly (2-)3-septate, colourless (very rarely slightly  
 greyish brown); pantropical . . . . .  
 . . . . . *Arthonia cyanea* MÜLL. ARG. var. *cyanea* p.p.  
 [Syn.: *Arthonia anisolocularis* L. XAVIER & TALTASSE (LÜCKING & al. 1997)]
- 23\* Ascospores irregularly (2-)3-4-septate, colourless to greyish brown  
 in the same hymenium; Cocos Island (Costa Rica) . . . . .  
*Arthonia cyanea* var. *cocosensis* R. LÜCKING (LÜCKING & LÜCKING 1995)
- 24 Ascocarps 0.3-1.0 mm diam.; ascospores 3-5-septate, 16-25 × 4-  
 8 µm; pantropical . . . . . *Arthonia palmulacea* (MÜLL. ARG.) R. SANT.
- 24\* Ascocarps 0.2-0.5 mm diam.; ascospores (2-)3-septate, 10-16 × 3-  
 6 µm . . . . . 25
- 25 Thallus reduced, phycobiont *Trentepohlia*; epithecium K+ pur-  
 plish; Australia (Tasmania) . . . . .  
 . . . . . *Arthonia apteropteridis* KANTVILAS & VĚZDA
- 25\* Thallus normally developed, phycobiont *Phycopeltis*, in irregu-  
 larly net-like plates; epithecium K; Neotropics (Brazil). . . . .  
*Arthonia lecythidicola* (BAT. & H. MAIA) R. LÜCKING & SÉRUS.  
 (LÜCKING & al. 1997)  
 [Bas.: *Ameropeltomyces lecythidicola* BAT. & H. MAIA 1967]
- 26 Ascocarps whitish pruinose to pale yellowish or bluish white . . . . 27
- 26\* Ascocarps dark brown to almost black. . . . . 28



- 27 Ascocarps whitish pruinose, not covered by thallus tissue; phycobiont *Phycopeltis*; ascospores macrocephalic, with one large cell at the distal end, vermiform and tapering towards the proximal end; pycnidia black; tropical Australasia . . . . .  
*Eremothecella macrocephala* (R. SANT.) THOR, SÉRUS., R. LÜCKING  
 & MATSUMOTO (THOR & al. 1997)
- 27\* Ascocarps pale bluish white, laterally covered by algiferous thallus tissue; phycobiont *Trentepohlia*; ascospores microcephalic, with one large cell near the middle, fusiform; pycnidia pale bluish to greyish white, with a darker zone in the centre; Neotropics (Amazon basin) . . . . .  
*Amazonomyces sprucei* (R. SANT.) R. LÜCKING & SÉRUS. (LÜCKING & al. 1997)
- [Bas.: *Stirtonia sprucei* R. SANT.]
- 28 Ascospores 7–11-septate, 50–90 × 8–12 µm; tropical Asia . . . . .  
*Eremothecella macrosperma* (ZAHLEBR.) SÉRUS.
- 28\* Ascospores 5–7-septate, 28–45 × 7–10 µm . . . . . 29
- 29 Ascocarps K+ purple; tropical Asia (New Guinea) . . . . .  
*Eremothecella variatae* (SIPMAN & APTROOT) SÉRUS.
- 29\* Ascocarps K–; pantropical. . . . . *Eremothecella calamicola* SYD.
- 30 Ascocarps dark brown to almost black; pycnidia ± rounded, 0.15 mm diam.; Neotropics . . . . .  
*Eremothecella cingulata* (R. SANT.) L. I. FERRARO & R. LÜCKING comb. nova.
- [Bas.: *Arthothelium cingulatum* R. SANT., Symb. Bot. Ups. 12: 93 (1952).  
 – Type: Chile, Valdivia, 1905, THAXTER s.n. (MICH, holotype)]
- 30\* Ascocarps white to pale yellowish . . . . . 31
- 31 Ascocarps unorganized, cotton-like, pure white, without covering thallus layer; asco-spores regularly muriform; pycnidia unknown . . . . .  
*Cryptothecia candida* (Kremp.) R. SANT.
- 31\* Ascospores organized, smooth, pale yellowish to greenish white, laterally covered by an algiferous thallus layer; ascospores with a large, undivided cell near the middle; pycnidia oval, with a pale margin and dark centre, producing long, filiform conidia . . . . .  
*Amazonomyces farkasiae* (R. LÜCKING) R. LÜCKING & SÉRUS. (LÜCKING & al. 1997)
- [Bas.: *Cryptothecia farkasiae* R. LÜCKING]

## 4. References

- APTROOT A. & SIPMAN H. J. M. 1991. New lichens and lichen records from New Guinea. – Willdenowia 20: 221–256.
- BATISTA A. C. & PERES G. E. P. 1964. Líquens imperfeitos: novos gêneros e espécies de Manaus. – Anais Congr. Soc. Bot. Brasil 14: 89–102.

- & MAIA H. DA S. 1967. Novos líquens imperfeitos do Amazonas e de Pernambuco. – Atas Inst. Micol. Univ. Fed. Pern. 5: 55–71.
- BECKER U. & LÜCKING R. 1995. Foliikole Flechten aus dem Tai-Nationalpark, Elfenbeinküste (Tropisches Afrika). I. Neue Arten. In: DANIELS F. J. A., SCHULZ M. & PEINE J. (eds): Flechten FOLLMANN. Contributions to Lichenology in Honour of Gerhard FOLLMANN, p. 161–173. – Geobotanical and Phytotaxonomical Study Group, Botanical Institute, University of Cologne.
- FARKAS E. & SIPMAN H. J. M. 1993. Bibliography and checklist of foliicolous lichens up to 1992. – Tropical Bryology 7: 93–148.
- KANTVILAS G. & VĚZDA A. 1988. A new lichenised species of *Arthonia* from south-western Tasmania. – Aust. Syst. Bot. 1: 189–190.
- LÜCKING R. 1991. Neue Arten foliikoler Flechten aus Costa Rica, Zentralamerika. – Nova Hedwigia 52: 267–304.
- 1992a. Zur Verbreitungsökologie foliikoler Flechten in Costa Rica, Zentralamerika. Teil I. – Nova Hedwigia 54: 309–353.
- 1992b. Zur Verbreitungsökologie foliikoler Flechten in Costa Rica, Zentralamerika. Teil II. – Herzogia 9: 81–112.
- 1995. Additions and corrections to the foliicolous lichen flora of Costa Rica. The family *Arthoniaceae*, with notes on the genus *Stirtonia*. – Lichenologist 27: 127–153.
- 1997a. The use of foliicolous lichens as bioindicators in the tropics, with special reference to the microclimate. In: FARKAS E. É & Pócs T. (eds): Cryptogams in the Phyllosphere: Systematics, Distribution, Ecology, and Use. – Abstracta botanica 21 (in press).
- 1997b. Additions and corrections to the knowledge of the foliicolous lichen flora of Costa Rica, Central America. The family *Gomphillaceae*. – Bibl. Lichenol. 65: 1–109.
- & LÜCKING A. 1995. Foliicolous lichens and bryophytes from Cocos Island, Costa Rica. A taxonomical & ecogeographical study. I. Lichens. – Herzogia 11: 143–174.
- SÉRUSIAUX E., MAIA L. C. & PEREIRA E. C. G. 1997. A taxonomic revision of names of foliicolous, lichenized fungi published by BATISTA and co-workers between 1960 and 1975. – Lichenologist (subm.).
- MATZER M. 1996. Lichenicolous ascomycetes with fissitunicate asci on foliicolous lichens. – Mycological Papers 171: i–x, 1–202.
- SANTESSON R. 1952. Foliicolous lichens. I. A revision of the obligately foliicolous, lichenized fungi. – Symb. bot. ups. 12 (1): 1–590.
- SÉRUSIAUX E. 1989. *Echinoplaca furcata*, a new species of foliicolous lichen (*Gomphillaceae*) from Rwanda. – Mycotaxon 35: 237–242.
- 1992. Reinstatement of the lichenized genus *Eremothecella* SYDOW. Systema Ascomycetum 11: 39–47.
- THOR G., LÜCKING R. & MATSUMOTO T. 1997. The foliicolous lichen flora of Japan. – Opera botanica (subm.).
- UPADHYAY H. B. P. 1964. A new *Opegrapha* species from the Federal Territory of Rondônia. – Publicações, Inst. Micol. Univ. Recife 410: 1–6.
- XAVIER FILHO L. 1964. Um novo *Arthonia* e outros líquens estudados no IMUR. – Anais Congr. Soc. Bot. Brasil 13: 462–468.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Phyton, Annales Rei Botanicae, Horn](#)

Jahr/Year: 1997

Band/Volume: [37\\_1](#)

Autor(en)/Author(s): Ferraro Lidia Itati, Lücking Robert

Artikel/Article: [New Species or Interesting Records of Foliicolous Lichens III. \*Arthonia crystallifera\* spec. nova \(Lichenized Ascomycetes: Arthoniaceae\), with a World-Wide Key to the Foliicolous Arthoniaceae. 61-70](#)