

Phyton (Horn, Austria)	Vol. 41	Fasc. 2	277-293	28. 12. 2001
------------------------	---------	---------	---------	--------------

## Chances and Limits of Floristic Island Inventories – the Dionysades Group (South Aegean, Greece) Re-visited

By

Erwin BERGMEIER\*) and Panayotis DIMOPOULOS\*\*)

Received May 7, 2001

Keywords: Floristic inventory, species census, islands, monitoring. – Crete, Aegean, Greece.

### Summary

BERGMEIER E. & DIMOPOULOS P. 2001. Chances and limits of floristic island inventories – the Dionysades group (South-Aegean, Greece) re-visited. – *Phyton* (Horn, Austria) 41(2): 277-293, 1 figure. – English with German summary.

A new, annotated and critical flora inventory for the three South Aegean Dionysades islands is presented, differentiated for each island, and based on 240 own vascular plant records and all available data from literature. The total list for the island group comprises now 277 species of which 51 are reported for the first time. *Astragalus sesameus* was found new for the South Aegean (Cretan area), and *Medicago arborea* is new to the native flora of wider Crete. The distribution of *M. arborea* and other cliff species into phrygantic habitats on the two major islands was not noticed by previous researchers and is, since grazing by domestic herbivores was given up, probably a rather recent phenomenon. A comparison between the first (GANDOGGER, ca. 1917) and our recent records (in 2000) revealed few cases of taxa which are now likely to be extinct. They are weeds that bear witness of former local cultivation on the two bigger islands. The great majority of the new findings is not owing to recent establishment but to previous observation gaps. There is but one clear case of recent introduction: *Oxalis pes-caprae*. The question is raised whether conventional species census is an adequate tool for island monitoring. It is suggested to install and perform a more consistent and standardized way of island monitoring, including appropriate sampling on community and population level.

---

\*) Dr. Erwin BERGMEIER, Institut für Biologie II (Geobotanik), Albert-Ludwigs-Universität Freiburg, Schänzlestraße 1, D-79104 Freiburg, Germany; e-Mail: bergmeie@uni-freiburg.de

\*\*) Ass.-Prof. Panayotis DIMOPOULOS, Department of Environmental and Natural Resources Management, University of Ioannina, Seferi 2, GR-30100 Agrinio, Greece; e-mail: pdimopul@cc.uoi.gr

### Zusammenfassung

BERGMEIER E. & DIMOPOULOS P. 2001. Möglichkeiten und Grenzen bei der floristischen Erforschung von Inseln – die Dionysaden-Gruppe (Süd-Ägäis, Griechenland) neu untersucht. – *Phyton* (Horn, Austria) 41(2): 277–293, 1 Abbildung. – Englisch mit deutscher Zusammenfassung.

Auf der Grundlage von 240 eigenen Nachweisen und allen verfügbaren Literatordaten wird ein neues, kommentiertes und kritisches Verzeichnis der Gefäßpflanzen der südägäischen Dionysaden-Inselgruppe vorgelegt, mit Einzelangaben für jede der drei Inseln. Die Florenliste umfasst 277 Arten, von denen 51 neu für die Inselgruppe sind. *Astragalus sesameus* ist neu für die Südägäis, und *Medicago arborea* wird erstmals als indigene Art im Gebiet Kretas (einschließlich vorgelagerter Inseln) nachgewiesen. Die Ausbreitung von *M. arborea* und anderer Kliffspezialisten in Phrygana-Habitats wird von früheren Bearbeitern nicht erwähnt und ist wahrscheinlich ein junges Phänomen seit der Aufgabe der Weidenutzung. Ein Vergleich zwischen den ersten Pflanzennachweisen (GANDOGGER, um 1917) und unseren (2000) zeigt, dass seither offenbar nur wenige, nämlich ruderale Arten, Zeugen des früheren Feldbaus auf den beiden größeren Inseln, ausgestorben sind. Ein Vergleich der Liste von CHRISTODOULAKIS & al. 1990 mit unserer legt nahe, dass die große Mehrheit der Neunachweise auf Beobachtungslücken und nicht auf kürzlicher Etablierung beruht. Letztere ist nur für *Oxalis pes-caprae* anzunehmen. Es wird bezweifelt, dass konventionelle Artenzählungen für ein Insel-Monitoring methodisch angemessen sind, und daher vorgeschlagen, sie um standardisierte Probenahmen auf Gesellschafts- und Populationsebene zu ergänzen.

### 1. Introduction

The Dionysades Islands, 7–12 km off the north coast of eastern Crete, harbour a considerable number of Aegean endemic and other rare plant species, and they are also significant for a unique herpeto- and ornithofauna. The islands were consequently included in the European network of conservation areas, 'Natura 2000', currently under construction (DAFIS & al. 1996). It was in this framework that our fieldwork on the Dionysades was made possible, with the task of sampling and mapping the vegetation. A floristic inventory was also made, as a by-product of our investigations on all and all over the three islands, Janisada, Dragonada, and Paximada. Some geographic parameters of the Dionysades are summarized in Table 1; Fig. 1 shows the position of the archipelago. More details are available in the account of CHRISTODOULAKIS & al. 1990, together with an outline of the islands' phytogeography and vegetation.

Previous floristic data on the Dionysades had been sampled by GANDOGGER in 1917 and by RECHINGER in 1942, and were published by RECHINGER 1943a, 1943b, 1951 and in various later revisional contributions. RUNEMARK and co-workers, on their visits in 1960 and 1962, and GREUTER in 1966, collected a limited number of specimens from Janisada and Dragonada. Their materials were to a major extent published by GREUTER 1973a and in various monographs and revisional papers, including GREU-




Fig. 1. Geographical position of the Dionysades Islands.

Table 1

Some geographic parameters of the Dionysades islands.

Island	Area (ha)	Altitude (m)	Max. extension (km)	Geology	Nearest distance to Crete (km)
Janisada	208	150	3,2	Plattenkalk	7
Dragonada	284	125	3,5		8,5
Paximada	30	136	1,25	Phyllite	12

TER 1968, 1973b, STRID 1970, STORK 1972, BOTHMER 1974, RUNEMARK 1980, SNOGERUP & al. 1990, MEUSEL & KÄSTNER 1990. In 1987/88, a group of botanists from the University of Patras visited the three islands, attempting to provide a full floristic inventory subsequently published by CHRISTODOULAKIS & al. 1990. Later, KYPRIOTAKIS 1998 added a short list of records from Paximada. Our own fieldwork was in March/May 2000 (Table 2).

Table 2

Botanical research on the Dionysades islands in the 20th century. J = Janisada, D = Dragonada, P = Paximada.

Researcher	Year	Month/Day	Islands
GANDOGER	ca.1917	?	J, D
RECHINGER	1942	5/13-14	J, D, P
RUNEMARK & NORDENSTAM	1960	7/4	D, P
RUNEMARK & S. SNOGERUP	1962	5/19	J, (D, P)
GREUTER	1966	10/31	J, D
CHRISTODOULAKIS, GEORGIADIS, IATROU	1987	9/26-27	J, D, P
ECONOMIDOU, GEORGIADIS, IATROU	1988	4/2	
KYPRIOTAKIS	? (-1997)	?	P
BERGMEIER & DIMOPOULOS	2000	3/29	J
BERGMEIER	2000	3/31+5/11	D, P

A broad range of floristic studies presenting Aegean island inventories has accumulated which may be used in general-purpose island biogeography as well as in the field of regional phytogeography (GREUTER 1991, 1995; STRID 1996). To date, however, there are only few examples of repeated floristic investigations on Mediterranean islands (SNOGERUP & SNOGERUP 1987, RAUS 1988, HÖNER 1991), although species inventories that permit comparison along the time scale are urgently needed if questions concerning dynamics and turnover are to be addressed.

The Aegean archipelago has been termed a "natural laboratory" for studies on biogeography and speciation (GREUTER 1995, STRID 1996). Our new data on the uninhabited Dionysades, together with the materials taken from previous accounts, may serve as an example, along which the chances and limits of current floristic island census in the Aegean and elsewhere may be discussed.

In the present paper, the following questions are addressed:

1. Is there evidence for species turnover on the Dionysades in the course of the 20<sup>th</sup> century?
2. Are the available data sufficient for numerical comparison of island floras in space and time?
3. Is the species census an appropriate tool for island monitoring?

## 2. Methods

We visited Janisada on March 29, 2000, Dragonada on March 31 and May 5, and Paximada also on May 5, 2000. The time was limited and chiefly spent for vegetation sampling but floristic recording was performed simultaneously. The collections (73 vouchers) were stored in the herbarium of the first author, with the exception of that of *Astragalus sesameus* which was given to the Botanische Staatssammlung München (M). A full list of our vascular plant taxon records of the Dionysades was provided, specified for each island and author, and all available data from literature were included. Literature data were critically reviewed, i.e., all data were taken into consideration, and records considered unlikely were discussed but not used for the numerical evaluation. The latter is based on accepted taxa on the levels of species and subspecies. The nomenclature of the taxa follows JAHN & SCHÖNFELDER 1995, except for the genera *Prospero* and *Charybdis* (SPETA 2000, 2001). The island names in the list are abbreviated by the first letter J, D, P, and the authors/collectors as follows: G – GANDOGER; R – RECHINGER; RU – RUNEMARK & NORDENSTAM, RUNEMARK & SNOGERUP; GR – GREUTER; CH – CHRISTODOULAKIS, GEORGIADIS, ECONOMIDOU, IATROU & TZANOUDAKIS; K – KYPRIOTAKIS; B – BERGMEIER & DIMOPOULOS. Records based on our collections are indicated by !.

## 3. Floristic Inventory of the Dionysades Islands

The list is arranged with the major groups in systematic order, and the families and species in alphabetic order. The names of unlikely literature records not accepted by us are in square brackets. Synonyms from literature and misapplied names (the latter in inverted commas) are given if used by CHRISTODOULAKIS & al. 1990.

### *Pteridophyta*

#### *Sinopteridaceae*

*Cheilanthes acrostica* (“*C. pteridioides*”) – J [CH,B!]; D [CH,B]

*Cosentinia vellea* (*Cheilanthes vellea*) – J [CH]

### *Gymnospermae*

#### *Cupressaceae*

*Juniperus phoenicea* – J [B]; D [G,R,CH,B]

### *Angiospermae, Dicotyledones*

#### *Aizoaceae*

*Mesembryanthemum nodiflorum* – D [R,CH]; P [R,CH,B]

### *Anacardiaceae*

*Pistacia lentiscus* – J [B]; D [G,R,CH,B]

### *Boraginaceae*

*Anchusa aegyptiaca* – J [R,CH,B]; D [G,CH,B]; P [CH]

*Buglossoides arvensis* – J [B!]

*Echium angustifolium* – J [B]; D [G]

*Echium arenarium* – D [R,B]; P [CH,B!]

*Echium parviflorum* – J [R]; D [G,B]

*Neatostema apulum* – J [R,B]; D [B]

### *Campanulaceae*

*Campanula erinus* – D [G,CH]

[*Campanula spatulata* subsp. *spruneriana* was given for Janisada by

RECHINGER 1943a, based on a GANDOGGER record; considered erroneous by CHRISTODOULAKIS & al. 1990 and presumed to be *C. spatulata* subsp. *filicaulis*, but neither taxon was found since GANDOGGER. In fact, the occurrence of *C. spatulata* s.l. on the Dionysades islands is unlikely, and the species should be regarded as absent.]

### Capparaceae

*Capparis spinosa* subsp. *rupestris* –  
D [G,CH,B]; P [R,CH, K,B!]

### Caryophyllaceae

*Arenaria aegaea* – J [RU,CH]; D [CH];  
P [K]

*Dianthus fruticosus* subsp. *amarginus* –  
J [Ru]; D [Ru,GR]; P [K,B!]

*Herniaria hirsuta* – D [CH,B]; in the absence of a herbarium specimen, the identity of the *Herniaria* species on Dragonada remains to be confirmed.

*Paronychia macrosepala* – J [G,R,CH,B];  
D [G,R,CH,B]

*Polycarpon tetraphyllum* – J [B];  
D [G,CH,B]; P [CH,B]

*Sagina maritima* – P [R]

*Silene apetala* – D [B]

*Silene dichotoma* – J [G]; since GANDOGGER the species was never observed again. It is now absent on the islands and may have had the status of a casual weed on Janisada.

*Silene gallica* – J [CH]

*Silene nocturna* – J [B]; D [B]

*Silene sedoides* – J [R,B]; D [R,CH,B];  
P [R,CH,B!]

*Silene vulgaris* subsp. *macrocarpa* –  
J [R]; D [B]

*Stellaria pallida* – D [B]

### Chenopodiaceae

*Arthrocnemum macrostachyum* –  
J [CH,B]; D [R,CH,B]; P [R,CH, K,B!]

*Atriplex halimus* – J [R,CH,B];  
D [G,R,CH,B]; P [R,CH, K,B]

*Atriplex recurva* – P [R,CH]

*Chenopodium murale* – J [B];  
D [R,CH,B]; P [CH,B]

*Chenopodium opulifolium* – D [G]; the species was never observed by later botanists and seems to be absent from the Dionysades islands today.

*Salsola aegaea* – P [R,CH, K,B!]

*Salsola carpatha* – J [G,CH];  
D [G,CH,B!]; P [Ru,CH, K,B!]

[*Sarcocornia fruticosa* was recorded from Paximada by RECHINGER 1951, however, its presence in the Cretan area is not sufficiently documented and since misidentification with *Arthrocnemum macrostachyum* is quite possible, the record was not included in the map provided by TAN 1997.]

*Suaeda vera* – P [R,CH,B!]

### Cistaceae

*Cistus creticus* – D [CH]

*Fumana arabica* – J [CH,B]; D [CH,B]

*Fumana thymifolia* – J [CH,B]; D [G,B]

*Helianthemum salicifolium* –

J [R,CH,B]; D [R,CH,B]

*Tuberaria guttata* s.l. – J [CH]

### Compositae

*Aetheorhiza bulbosa* subsp. *microcephala* – J [B]; D [G,B]

*Anthemis ammanthus* subsp. *ammanthus* – J [G,R,RU,CH,B];

D [R,RU,CH,B!]; P [R,CH,K,B!]

*Atractylis cancellata* – J [R,CH,B];

D [R,CH,B]

*Calendula arvensis* – D [B]; P [CH]

*Carlina diae* – J [GR]; D [RU,GR,CH,B!]

*Carlina graeca* (*C. corymbosa* subsp. *graeca*) – J [G,CH,B]; D [G,CH,B]

*Chrysanthemum coronarium* – D [G,B]

*Cichorium pumilum* – D [R]

*Crepis commutata* – J [B]; D [G]

*Crepis cretica* – J [G,R,CH,B];

D [G,CH,B]

*Crepis multiflora* – J [G,R]; D [G,R,B];

P [B]



- Crupina crupinastrum* – D [B]  
*Cynara cornigera* – J [R, CH, B];  
 D [G, R, CH, B]  
*Filago aegaea* subsp. *aegaea* – J [G, B];  
 D [R, B!]  
*Filago pygmaea* – D [CH]  
*Hedypnois rhagadioloides* – J [CH, B];  
 D [G, CH, B]; the plants found may be  
 assigned to subsp. *tubaeformis* (gi-  
 ven by CHRISTODOULAKIS & al. 1990:  
 431) and to subsp. *monspeliensis*.  
*Helichrysum conglobatum* (*H. barre-*  
*lieri*) – J [CH, B]; D [G, CH, B]  
*Hyoseris scabra* – J [CH, B]; D [CH, B]  
*Hypochaeris achyrophorus* –  
 J [G, CH, B]; D [CH, B]; P [CH]  
*Lamyropsis cynaroides* – D [CH]  
*Leontodon tuberosus* – J [G, CH, B];  
 D [G, CH, B]  
*Phagnalon graecum* – J [CH, B];  
 D [R, CH, B!]; P [CH, K, B]  
*Ptilostemon chamaepeuce* – J [GR]  
*Reichardia intermedia* – J [B]  
*Reichardia orientalis* – J [R, RU, CH, B!];  
 D [R, CH, B]; P [R, CH, B!]  
*Reichardia picroides* – J [G]; D [B]  
*Rhagadiolus stellatus* – D [CH]  
*Scolymus hispanicus* – J [R, CH]  
*Senecio gnaphalodes* – D [RU, B!]  
*Senecio leucanthemifolius* (“*S. galli-*  
*cus*”) – J [B]; D [R, CH, B!]; P [CH, B]  
*Senecio pygmaeus* – J [B]; this taxon  
 which was known in regional litera-  
 ture for the western part of Crete  
 only, is treated as a variety of *S. leu-*  
*canthemifolius* by ALEXANDER 1979  
 but differs clearly in leaf shape, ab-  
 sence of ligulate florets, and quanti-  
 tative characters of the calyculus  
 and phyllaries. No intermediates  
 were seen in co-occurring populati-  
 ons elsewhere in Crète.  
*Senecio vulgaris* – J [B]; D [B]  
*Sonchus oleraceus* – J [B]; D [G, CH, B];  
 P [CH, K, B]  
 [*Sonchus tenerrimus*; due to frequent  
 confusion with forms of *S. oleraceus*  
 the record for Dragonada (RECHIN-  
 GER 1943b) needs confirmation.]  
*Tragopogon sinuatus* – J [B]; D [CH, B]  
*Urospermum picroides* – J [R, B];  
 D [G, CH, B]; P [B]
- Convolvulaceae*  
*Convolvulus althaeoides* – J [CH, B];  
 D [G, R, CH, B]  
*Convolvulus siculus* – J [B!]; D [B]  
*Cuscuta palaestina* – J [B]; D [CH, B];  
 P [B!]
- Crassulaceae*  
*Crassula alata* – J [B]  
*Sedum litoreum* – J [G, R, CH, B!];  
 D [G, R, CH, B]; P [CH, K, B]  
*Sedum rubens* – J [B]; D [B]  
*Umbilicus horizontalis* – J [R, CH, B];  
 P [CH, B]
- Cruciferae*  
*Alyssum strigosum* – J [CH, B!]  
*Biscutella didyma* – J [CH, B]; D [CH, B]  
*Brassica cretica* subsp. *aegaea* –  
 J [R, RU, CH, B!]; D [RU, B]; P [CH, K, B]  
 [*Cakile maritima*; there is no suitable  
 habitat (sandy beaches) for this spe-  
 cies on the Dionysades; hence, in  
 the absence of a herbarium speci-  
 men, the record for Dragonada  
 (CHRISTODOULAKIS & al. 1990) is im-  
 probable.]  
*Clypeola jonthlaspi* – J [CH, B]  
*Didesmus aegyptius* – J [R, B]; D [B];  
 P [R, B]  
*Fibigia lunarioides* – J [RU, CH];  
 D [GR, CH, B!]; P [RU, CH, K, B]  
*Malcolmia chia* – J [RU, CH, B!]  
*Malcolmia flexuosa* – J [G]; D [R, CH, B!];  
 P [R, CH, K, B!]  
*Matthiola sinuata* subsp. *glandulosa* –  
 J [GR, CH]; P [CH, K, B!]
- Cucurbitaceae*  
*Bryonia cretica* – J [B]

*Ericaceae**Erica manipuliflora* – D [B]*Euphorbiaceae**Euphorbia acanthothamnos* – J [B];

D [G,CH,B]

*Euphorbia dendroides* – J [G,R,CH,B];

D [G,R,CH,B]; P [CH,B]

*Euphorbia exigua* – D [B]*Euphorbia peplus* – J [CH,B]; D [B]*Mercurialis annua* – J [CH,B];

D [G,CH,B]; P [CH,B]

*Frankeniaceae**Frankenia hirsuta* – J [G]; D [R,CH,B!];

P [CH, K]

*Frankenia pulverulenta* – D [R,CH]*Gentianaceae**Blackstonia perfoliata* – J [CH]; D [CH]*Centaurium pulchellum* – J [G]*Centaurium tenuiflorum* – D [G,CH,B!]*Geraniaceae**Erodium cicutarium* – J [CH,B]; D [B]*Erodium gruinum* – D [B]

[*Erodium laciniatum*; the occurrence of this psammophilous species on Dragonada (RECHINGER 1943a, based on GANDOGER) is highly unlikely due to ecological reasons. GANDOGER might have seen *E. neuradifolium* instead.]

*Erodium malacoides* – J [B]; D [B!]*Erodium neuradifolium* – J [CH];

D [R,CH]; P [CH,B!]

*Geranium molle* subsp. *molle* – J [CH];

D [G,CH]; P [CH]

*Geranium rotundifolium* – J [CH,B];

D [CH,B]

*Labiatae**Coridothymus capitatus* – J [G,R,CH,B];

D [G,R,CH,B]

*Lamium amplexicaule* – J [CH,B]; D [B];

P [CH]

*Marrubium vulgare* – D [G,B]*Prasium majus* – J [B]; D [R,CH,B]; P [B]*Salvia verbenaca* – J [B]; D [G,CH,B]*Salvia viridis* – J [RU,CH,B]; D [G,CH,B]*Sideritis curvidens* – J [R,CH];

D [G,R,CH,B]

*Stachys spinosa* – J [G,R,CH,B];

D [CH,B]

*Teucrium brevifolium* – J [R,CH,B];

D [G,CH,B]

[*Teucrium cuneifolium* was said to occur on Dragonada by GANDOGER (RECHINGER 1943a) but has never been confirmed. It might have been mistaken for *T. divaricatum* or a form of *T. gracile*. The occurrence of *T. cuneifolium* was already considered doubtful by CHRISTODOULAKIS & al. 1990.]

*Teucrium divaricatum* – D [B]*Teucrium gracile* – J [R,B]; D [G,CH,B!]*Leguminosae**Anagyris foetida* – J [B]*Anthyllis hermanniae* – D [CH]*Anthyllis vulneraria* subsp. *rubriflora* – D [CH]*Astragalus hamosus* – J [CH,B];

D [CH,B!]

*Astragalus sesameus* – D [B]; first record for the Cretan area as defined by GREUTER & al. 1989, and second for the Aegean.

*Bituminaria bituminosa* (*Aspalathium bituminosum*) – J [B]; D [CH,B]*Calicotome villosa* – D [G,CH,B]*Coronilla scorpioides* – J [B]; D [CH,B!]*Genista acanthoclada* – D [G,CH,B]*Hedysarum spinosissimum* – D [CH,B]*Hippocrepis biflora* – J [CH]; D [R,CH,B]*Hippocrepis ciliata* – J [B!]; D [CH,B]*Hippocrepis cyclocarpa* – D [B!]*Hymenocarpus circinnatus* – J [CH,B]; D [B]*Lotus cytisoides* – J [G,R,CH,B!];

D [G,R,CH,B]

*Lotus edulis* – J [B]; D [G,CH,B]*Lotus ornithopodioides* – J [CH,B]*Lotus peregrinus* – J [CH,B]; D [CH,B]*Medicago arborea* – J [B]; D [B!]; this is the first record of a clearly native



population in the area of Crete and surrounding islands.

*Medicago coronata* – J [CH,B]; D [CH,B]  
[*Medicago littoralis*; another GANDOGGER record (from Dragonada, RECHINGER 1943a) that has never been confirmed. The occurrence of this chiefly psammophilous species on the Dionysades must be considered doubtful for ecological reasons. It may well have been confused with *M. truncatula*.]

*Medicago monspeliaca* (*Trigonella monspeliaca*) – J [B]; D [G,R,CH,B];  
P [CH,B]

*Medicago orbicularis* – J [CH,B]; D [CH]

*Medicago praecox* – J [B!]; P [CH,B!]

*Medicago truncatula* – J [B!]; D [B!];  
P [CH,B!]

*Ononis ornithopodioides* – D [R,CH,B]

*Ononis reclinata* – J [R,CH,B];  
D [G,CH,B]

*Scorpiurus muricatus* – J [CH,B];  
D [G,CH,B]; the plants we found belong to var. *subvillosus*.

*Trifolium angustifolium* s.l. – J [CH];  
D [B]

*Trifolium campestre* – J [CH,B];  
D [G,R,CH,B]

*Trifolium scabrum* – J [B]; D [R,CH,B]

*Trifolium stellatum* – J [B]; D [CH,B]

*Trifolium tomentosum* – D [B]

*Trigonella rechingeri* – J [B]; D [R,CH];  
P [R,CH,B!]

*Tripodion tetraphyllum* (*Anthyllis tetraphylla*) – J [CH]; D [B]

*Vicia cretica* – J [B]; D [B]

#### Linaceae

*Linum strictum* subsp. *spicatum* –  
J [G,CH,B]; D [G,R,CH,B]

#### Malvaceae

*Malva aegyptia* – J [R,B]; D [B]

*Malva cretica* – J [R,CH,B]; D [G,CH,B]

*Malva parviflora* – J [B]; D [R,B];  
P [CH,B!]

*Malva sylvestris* – D [G]; never recorded since GANDOGGER; the ruderal species may have been a casual introduction on Dragonada.

#### Orobanchaceae

*Cistanche phelypaea* – P [Ru]; the record was erroneously assigned to Janisada by CHRISTODOULAKIS & al. 1990.

*Orobanche ramosa* L. subsp. *mutellii* –  
J [CH,B]; D [CH,B]

*Orobanche schultzei* – J [Ru]; specimen (LD) seen by H. Uhlich.

#### Oxalidaceae

*Oxalis pes-caprae* – J [B]; D [B]

#### Papaveraceae

*Fumaria macrocarpa* – J [B!]; D [B];  
P [B]

[*Fumaria officinalis*; the GANDOGGER record from Dragonada reported by RECHINGER 1943a is doubtful. The species was never observed since and is likely to be confused with other *Fumaria* species.]

*Fumaria parviflora* – J [B!]

*Papaver hybridum* – D [G,B]

*Papaver purpureomarginatum* ("P. dubium") – J [CH,B!]; D [G]

*Papaver rhoeas* – J [B]; D [G]

#### Plantaginaceae

*Plantago afra* – J [CH,B]; D [B];  
P [R,CH,B]

*Plantago albicans* – J [B]

*Plantago amplexicaulis* – J [R,CH,B];  
D [G,CH,B]

*Plantago cretica* – J [B]; D [CH,B]

*Plantago lagopus* – J [CH,B];  
D [G,R,CH,B]

*Plantago weldenii* – D [R,CH,B]; P [CH]

#### Plumbaginaceae

*Limonium aucheri* – J [R]

*Limonium echioides* – D [CH,B!]

*Limonium frederici* – D [B]; P [CH,B]

*Limonium graecum* – J [G]; D [CH,B];  
this species (but not *L. frederici*)  
was given for Paximada by KYPRIO-  
TAKIS 1998; confirmation is required.

[*Limonium hyssopifolium*; there is a  
record from Dragonada based on  
GANDOGGER (RECHINGER 1943a) which  
was never confirmed since. The spe-  
cies is known to be difficult to sepa-  
rate from *L. graecum* and *L. siti-  
acum* both of which occur on the is-  
land.]

*Limonium sitiicum* – J [R,B!]; D [R,B!]

### *Polygonaceae*

*Emex spinosa* – D [B]

*Rumex pulcher* subsp. *woodsii* – J [B]

### *Primulaceae*

*Anagallis arvensis* – J [CH,B];  
D [G,CH,B]; P [CH,B]; both red- and  
blue-flowering (var. *caerulea*) plants  
were seen on Janisada. The record  
of *Anagallis foemina* for Dragonada  
by GANDOGGER (RECHINGER 1943a)  
most probably belongs to *A. arvensis*  
var. *caerulea* and is assigned here.

*Asterolinon linum-stellatum* – J [B];  
D [B]

### *Ranunculaceae*

*Adonis microcarpa* subsp. *cretica* –  
J [B!]; D [CH]; we found yellow- and  
red-flowering plants on Janisada.

*Anemone coronaria* – J [B]; D [B]

*Nigella doerfleri* – J [Ru,B]; D [B!]

*Nigella fumariifolia* – J [Ru,CH]

### *Resedaceae*

*Reseda alba* – J [B!]; D [B!]

*Reseda lutea* – J [R,CH,B]; D [G,R,CH]

### *Rhamnaceae*

*Rhamnus lycioides* subsp. *oleoides* –  
J [G,B]; D [CH,B]

### *Rosaceae*

*Sanguisorba minor* subsp. *verrucosa* –  
D [B]

*Sarcopoterium spinosum* –  
J [G,R,CH,B]; D [G,R,CH,B]

### *Rubiaceae*

*Asperula rigida* – J [R]; D [B]

*Asperula tournefortii* – J [R]; P [K]

*Crucianella latifolia* – J [R,CH,B];  
D [CH,B]

*Galium aparine* – J [B]; D [CH,B]

*Galium graecum* subsp. *pseudocanum* –  
J [R, Ru CH]

*Galium murale* – J [CH,B]; D [G,B]

*Galium setaceum* – J [R,CH,B]; D [G,B]

*Sherardia arvensis* – J [CH,B]; D [CH,B]

*Valantia hispida* – J [G,CH,B];  
D [G,CH,B]; P [B]

*Valantia muralis* – D [CH]; P [CH]

### *Rutaceae*

*Ruta chalepensis* subsp. *fumariifolia* –  
J [G,CH,B!]; D [R,CH,B]

### *Santalaceae*

*Thesium bergeri* – J [R]; D [B]

*Thesium humile* – J [B]; D [B]

### *Scrophulariaceae*

*Kickxia elatine* subsp. *sieberi* – D [G]

*Linaria simplex* – J [B]; D [B]

*Misopates orontium* – J [CH,B]; D [B]

*Verbascum sinuatum* – J [B]

*Veronica cymbalaria* – J [B!]

### *Solanaceae*

*Mandragora autumnalis* – J [CH,B];  
D [G,CH,B]; P [CH,B]

### *Theligonaceae*

*Theligonum cynocrambe* – J [CH,B];  
D [B]

### *Thymelaeaceae*

*Thymelaea hirsuta* – J [G,B!];  
D [R,CH,B]

*Umbelliferae*

- Bupleurum gracile* – J [RU,B]; D [B!]  
*Bupleurum semicompositum* –  
 J [R, RU]; D [R,CH,B!]; P [B!]  
*Crithmum maritimum* – J [R,CH,B]  
*Lagoecia cuminoides* – J [CH,B]; D [B]  
*Scaligeria halophila* – J [B!]; D [R,B]  
*Scandix australis* – J [CH,B]; D [B]  
*Scandix pecten-veneris* – J [B]; D [CH]  
*Tordylium apulum* – J [CH,B]; D [CH,B]  
*Torilis nodosa* – J [G,CH,B]; D [G,CH,B]

*Urticaceae*

- Parietaria cretica* – J [G,CH,B];  
 D [G,CH,B]; P [CH, K,B]  
*Parietaria judaica* – D [CH]; P [CH]  
*Urtica urens* – D [G]; the species has never been observed since GANDOGGER and is probably not present anymore on Dragonada.

*Valerianaceae*

- Centranthus calcitrapae* – J [B!];  
 D [CH,B]  
*Valerianella discoidea* – J [CH,B!];  
 D [CH,B]

*Angiospermae, Monocotyledones**Amaryllidaceae*

- Narcissus tazetta* – J [B]; D [CH,B]

*Araceae*

- Arisarum vulgare* – J [G,CH,B];  
 D [CH,B]; P [CH,B]

*Gramineae*

- Aegilops markgrafii* – D [B!]  
*Avena barbata* – J [B]; D [G,R,CH,B];  
 P [CH,B]  
*Brachypodium distachyon* (*Trachynia distachya*) – J [G,CH,B]; D [CH,B];  
 P [CH,B]  
*Brachypodium retusum* – J [CH]; D [G,CH,B]; P [CH]; GANDOGGER recorded *Brachypodium phoenicoides* on Dragonada (RECHINGER 1943a), almost with certainty in error for *B.*

*retusum* to which the record is assigned here.

- Briza maxima* – J [B]; P [CH]  
*Bromus fasciculatus* – J [R,CH,B];  
 D [G,CH,B]; P [CH]  
*Bromus intermedius* – J [G,CH,B];  
 D [CH,B]  
*Bromus madritensis* – D [B]  
*Bromus rubens* – J [CH]; P [CH, K,B]  
*Catapodium marinum* – D [R,CH,B]; in the absence of herbarium specimens the identity of the plants assigned to *C. marinum* remains to be verified.  
*Catapodium rigidum* – J [G,CH,B];  
 D [CH,B]  
*Dactylis glomerata* subsp. *hispanica* –  
 J [G,R,CH,B]; D [B]  
*Echinaria capitata* – D [B]  
*Gastridium phleoides* (“*G. ventricosum*”) – J [B]; D [G,CH,B]; the genus seems to be represented on the Dionysades by *G. phleoides* only. The latter was not consistently separated from *G. ventricosum* by previous workers.  
*Hordeum leporinum* s.l. – D [G,R,CH,B];  
 P [CH]; plants from Dragonada were assigned to subsp. *glaucum* (RECHINGER 1943b).  
*Hyparrhenia hirta* – J [R,CH,B];  
 D [G,CH,B]  
*Lagurus ovatus* – J [B]  
*Lolium temulentum* – D [B]  
*Lolium rigidum* – J [B]  
*Melica minuta* (*M. ramosa*) – J [G,CH,B];  
 D [G,R,CH,B]  
*Parapholis marginata* – D [R,CH]  
*Phalaris minor* – J [B!]; D [G,CH,B]  
*Piptatherum coerulescens* – J [B];  
 D [CH,B]  
*Piptatherum miliaceum* subsp. *miliaceum* – J [CH,B]; D [G,CH,B]  
*Psilurus incurvus* – J [B]; D [B]  
*Rostraria cristata* (*Lophochloa cristata*) – J [G,CH,B]; D [G,CH,B]; P [R,CH,B]  
*Stipa capensis* – J [CH,B]; D [G,R,CH,B];  
 P [CH,B]

*Iridaceae*

*Crocus tournefortii* – J [GR]; D [B]; sterile plants seen on Dragonada were tentatively placed here.

*Gynandriris monophylla* – D [B]

*Gynandriris sisyrinchium* – J [B];  
D [CH,B]

*Liliaceae* s.l.

*Allium commutatum* – J [B]; P [RU,CH]

*Allium rubrovittatum* – D [CH,B]

*Allium tardans* – J [Gr,B]; D [B]; sterile *Allium* plants seen on Dragonada were tentatively assigned to *A. tardans*; confirmation is required.

*Asparagus aphyllus* subsp. *orientalis* – J [B!]; D [B]

*Asparagus horridus* (*Asparagus stipularis*) – D [R,CH,B]; P [R,CH, K,B!]

*Asphodeline lutea* – D [CH]

*Asphodelus ramosus* (*A. aestivus*) – J [CH,B]; D [G,R,CH,B]

*Charybdis maritima* (*Drimia maritima*) – J [G,R,CH,B]; D [CH,B]

*Colchicum* spec. – D [B]; sterile plants seen have shining, up to 3 cm wide leaves.

*Muscari comosum* – J [B]; D [B]

*Muscari dionysicum* – P [R,CH,B]

*Muscari spreitzenhoferi* – J [R,B];  
D [R,CH,B]

*Ornithogalum creticum* – J [R];  
D [R,B!]; P [B!]

*Prospero autumnale* s.l. – J [B]; D [B]

*Orchidaceae*

*Anacamptis pyramidalis* – J [CH,B];  
D [CH,B]

*Orchis collina* – J [B]; D [B]

*Ophrys fusca* s.l. – D [CH]

*Ophrys lutea* s.l. – D [CH]

## 4. Discussion

The fundamentals of the phytogeographic pattern in the Aegean were described for the first time by RECHINGER 1951 and later corroborated and refined by the studies of GREUTER and RUNEMARK and his co-workers. Similarly, on the Dionysades, it was also RECHINGER who was the first to provide reliable floristic data, based on GANDOGGER's and his own collections and observations (RECHINGER 1943a, 1943b, 1951). RUNEMARK and co-workers, as well as GREUTER, added data chiefly on species of particular phytogeographical interest. Taxonomic and phytogeographical insight had been the scientific motive so far. In contrast, working out a full species list as was undertaken by CHRISTODOULAKIS & al. 1990 for the Dionysades, was motivated rather by the idea of providing a basis for floristic comparison, both between different islands and on the same island but between different years. The latter approach is tested here.

Table 3 shows the number of taxa recorded per island and author. The total number of vascular plant taxa currently known on the Dionysades was found to comprise 277 species. We added 51 to the 226 known from literature. The respective figures found for each island were: Janisada (213; 71 + 142); Dragonada (235; 63 + 172); Paximada (72; 9 + 63). As far as numbers deviate slightly from those given by CHRISTODOULAKIS & al. 1990, this is due to the following reasons: a small number of records from literature were overlooked by the latter authors; doubtful previous records were rejected by us; there are two more recent additions to the flora of

Table 3  
Number of taxa per island recorded per author and in total.

	Janisada	Dragonada	Paximada	Dionysades total
GANDOGGER	36	78	—	97
RECHINGER	49	57	18	94
RUNEMARK & al.; GREUTER	19	8	4	26
CHRISTODOULAKIS & al.	107	145	58	188
KYPRIOTAKIS	—	—	20	20
BERGMEIER & DIMOPOULOS	174	201	53	240
Taxa total	213	235	72	277

Paximada by KYPRIOTAKIS. Floristic knowledge increased considerably along with any new study. In the first inventory (CHRISTODOULAKIS & al. 1990), 66 (42 %) taxa were found as new to the Dionysades. The respective figures found for each island were: Janisada 54 (62 %); Dragonada 55 (47 %); Paximada 39 (205 %). Our investigations resulted in another substantial increase, with 51 (23 %) new taxa for the Dionysades, 71 (50 %) for Janisada, 63 (37 %) for Dragonada, and 9 (15 %) for Paximada. Compared to the present total, the list of CHRISTODOULAKIS & al. 1990, comprises 82 % of the Dionysades flora, and 67 %, 73 %, and 83 %, for Janisada, Dragonada and Paximada, respectively.

Not surprisingly, the majority of the new records consists of small annuals. Due to the marked seasonality, such species are easily overlooked. Among these, the findings of *Astragalus sesameus* on Dragonada (first record for the Cretan area), and of *Hippocrepis cyclocarpa* (a plant of chiefly North-African distribution known in the wider area from the Karpathos archipelago and south-eastern Crete) are of particular phytogeographical interest. More surprising are the new records of as conspicuous taxa as *Pistacia lentiscus*, *Juniperus phoenicea*, *Euphorbia acanthothamnus*, *Anagyris foetida* and *Medicago arborea* for Janisada; and *Erica manipuliflora* and again *Medicago arborea* for Dragonada. Most of these records are based on only one or few individuals which were most probably overlooked, rather than established recently.

*Medicago arborea* deserves special attention, as no native population has ever been found before in the area of Crete and surrounding islands. On the main island, *M. arborea* is frequently planted as an ornamental and is locally naturalized. Our finding confirms the assumption of this species being native on small Aegean islands until it was distributed by man. A few shrubs of *M. arborea* were observed in the coastal cliffs of both Janisada and Dragonada. A single shrub was found on Janisada in phryganic vegetation well in front of the cliff-top, and several shrubs in similarly accessible position on Dragonada. Since both islands used to be grazed by

domestic herbivores, and *M. arborea* being very palatable and subject to browsing whenever possible, it seems likely that the species is currently distributing from the cliff refugia into the phryganic habitats. The former effects of grazing would explain why previous researchers up to now had overlooked the attractive and conspicuous shrub.

*Brassica cretica*, on the Dionysades represented by subsp. *aegaea* (SNOGERUP & al. 1990), is another palatable species which is about to become a conspicuous phrygana plant while in Crete it is almost exclusively found at inaccessible sites in gorges. It is still rare in phryganic habitats in Janisada but already almost common in the western part of Dragonada. In the absence of grazing, also other species of the marine cliffs, such as *Fibigia lunarioides*, *Salsola carpatha*, *Carlina diae* and *Senecio gnaphalodes*, occur in accessible sites on top of the cliffs of western Dragonada, indicating the potential of distributing beyond their classical cliff refugia.

On the two major islands, Janisada and Dragonada, limited areas of former arable fields could be traced. The small fields are abandoned since decades but GANDOGGER might have seen them cultivated. This would explain a number of weedy species recorded by him, such as *Silene dichotoma*, *Chenopodium opulifolium*, *Malva sylvestris* and *Urtica urens*, which have never been found again. GANDOGGER's records are known to be not perfectly reliable but we succeeded to confirm other ruderals noted by GANDOGGER, such as *Chrysanthemum coronarium*, *Marrubium vulgare*, *Papaver rhoeas* and *P. hybridum*. There is no evidence of the continuous presence of these weeds since GANDOGGER's times but re-introduction seems to be a rare event which is documented for the Dionysades in only a single case here: *Oxalis pes-caprae*, with only few plants found to date.

## 5. Conclusions

Comparison of island floras over time, whether numerical or qualitative, requires a high degree of completeness in floristic recording, a condition which, as we found, was obviously not the case prior to the present study. The evaluation of the inventories provided by CHRISTODOULAKIS & al. 1990 and in the present paper reflects that a great many of taxa had been neglected by previous workers, rather than being turnover phenomena based on flora dynamics. On species level, numerical interpretation of the island inventories would be hazardous due to the high degree of species that may be expected to have been present but were overlooked. The reasons for this could be, seasonality of species, incompatibility of research and optimum growth period, insufficient area-time ratio and perhaps recording knowhow. These are standardization deficits that have to be dealt with in almost any study of this kind. Conditions will rarely be as favour-

able as with SNOGERUP & SNOGERUP 1987 and HÖNER 1991, who succeeded on the basis of two major advantages: the small size of their study areas (islets), and the field studies in different years performed by the same experienced scientists. The latter study is at the same time a good example of an extensive work based not only on floristic data but also on population dynamics.

The incompatibility of the island inventories from the Dionysades is the more irritating as we indeed found evidence for remarkable vegetation dynamics currently in progress on the Dionysades, chiefly on population and community level. For the Dionysades, as for other "natural laboratories", monitoring is required but should not be based on floristic census alone. Standardized sampling techniques are recommended (O'LEARY 1993) which may include a floristic inventory of the islands once within a 10-yr period, a carefully designed permanent-plot approach in selected plant communities to be sampled every few years, and studies on the population dynamics of selected species with different trends in population size.

#### 6. Acknowledgements

We thank R. JAHN, Radebeul, for putting the Dionysades part of his database of Cretan plant records at our disposal, H. UHLICH for information on *Orobanche* specimens, and D. PODLECH, München, for reviewing and confirming *A. sesameus*. The research was funded by the Greek Ministry of Environment, Regional Planning and Public Works (YPEXODE), in the framework of the Greek Natura 2000 Habitat Types Mapping and Documentation Project. Logistic support from the Athens University team (Department of Biology, Section of Botany) is gratefully acknowledged.

#### 7. References

- ALEXANDER J. C. M. 1979. The Mediterranean species of *Senecio* sections *Senecio* and *Delphinifolius*. – Notes roy. bot. Garden Edinburgh 37: 387–428.
- BOTHMER R. VON 1974. Studies in the Aegean flora XXI. Biosystematic studies in the *Allium ampeloprasum* complex. – Opera bot. 34.
- CHRISTODOULAKIS D., GEORGIADIS TH., ECONOMIDOU E., IATROU G. & TZANOUDAKIS D. 1990. Flora und Vegetation der Dionysaden-Inseln (Südägäis, Griechenland). – Willdenowia 19: 425–443.
- DAFIS S., PAPASTERGIADOU E., GEORGHIOU K., BABALONAS D., GEORGIADIS TH., PAPA-GEORGIIOU M., LAZARIDOU T. & TSIAOISSI V. (eds.) 1996. Directive 92/43/EEC. The Greek Habitat Project Natura 2000: An overview. ivxx + 893 pp. – Thessaloniki.
- GREUTER W. 1968. Contributio floristica austro-aegaea 13. – Candollea 23: 143–150.
- 1973a. Additions to the flora of Crete, 1938–1972. – Ann. Mus. Goulandris 1: 15–83.
- 1973b. Monographie der Gattung *Ptilostemon* (*Compositae*). – Boissiera 22: 1–215.



- 1991. Botanical diversity, endemism, rarity, and extinction in the Mediterranean area: an analysis based on the published volumes of Med-Checklist. – Bot. Chron. 10: 63–79.
- 1995. Origin and peculiarities of Mediterranean island floras. – Ecol. medit. 21: 1–10.
- , BURDET H. & LONG G. 1989. Med-Checklist. A critical inventory of vascular plants of the circummediterranean countries 3. – Genève.
- HÖNER D. 1991. Mehrjährige Beobachtungen kleiner Vegetationsflächen im Raume von Karpathos (Nomos Dhodhekanisou, Griechenland). Ein Beitrag zur Klärung des "Kleininselphänomens". – Diss. bot. 173.
- JAHN R. & SCHÖNFELDER P. 1995. Exkursionsflora für Kreta. – Stuttgart.
- TAN K. 1997. *Sarcocornia* A. J. Scott. – In: STRID A. & TAN K. (eds.), Flora Hellenica 1: 131–132. – Königstein.
- KYPRIOTAKIS Z. 1998. Contribution to the study of the chasmophytic flora of Crete and to its utilization as a natural resource, to the direction of the ecotourism, the floriculture, the ethnobotany and the protection of the threatened plant species and their biotopes. – Ph.D Thesis. Patras. (in Greek).
- MEUSEL H. & KÄSTNER A. 1990. Lebensgeschichte der Gold- und Silberdisteln. Monographie der mediterran-mitteleuropäischen Compositen-Gattung *Carlina*. Band 1. Merkmalspektren und Lebensräume der Gattung. – Österr. Akad. Wiss., math.-naturw. Kl., Denkschr. 127.
- O'LEARY J. F. 1993. Towards greater uniformity of species diversity studies in Mediterranean-type ecosystems. – Landscape and Urban Planning 24: 185–190.
- RAUS Th. 1988. Vascular plant colonization and vegetation development on sea-born volcanic islands in the Aegean (Greece). – Vegetatio 77: 139–147.
- RECHINGER K. H. 1943a. Flora Aegaea. – Akad. Wiss. Wien, math.-naturw. Kl., Denkschr. 105(1).
- 1943b. Neue Beiträge zur Flora von Kreta. – Akad. Wiss. Wien, math.-naturw. Kl., Denkschr. 105(2,1).
- & RECHINGER-MOSER F. 1951. Phytogeographia aegaea. – Akad. Wiss. Wien, math.-naturw. Kl., Denkschr. 105(2,2).
- RUNEMARK H. 1980. Studies in the Aegean flora XXIII. The *Dianthus fruticosus* complex (*Caryophyllaceae*). – Bot. Not. 133: 475–490.
- , SNOGERUP S. & NORDENSTAM B. 1960. Studies in the Aegean flora I. Floristic notes. – Bot. Not. 113: 421–450.
- SNOGERUP S., GUSTAFSSON M. & BOTHMER R. VON 1990. *Brassica* sect. *Brassica* (*Brassicaceae*). I. Taxonomy and variation. – Willdenowia 19: 271–365.
- & SNOGERUP B. 1987. Repeated floristical observations on islets in the Aegean. – Pl. Syst. Evol. 155: 143–164.
- SPETA F. 2000. Beitrag zur Kenntnis der Gattung *Prospero* SALISB. (*Hyacinthaceae*) auf der griechischen Insel Kreta. – Linzer biol. Beitr. 32: 1323–1326.
- 2001. Die Echte und die Falsche Meerzwiebel: *Charybdis* SPETA und *Stellaroides Medicus* (*Hyacinthaceae*), mit Neubeschreibungen und Neukombinationen im Anhang. – Stapfia 75: 139–176.
- STORK A. L. 1972. Studies in the Aegean flora XX. Biosystematics of the *Malcolmia maritima* complex. – Opera bot. 33.

- STRID A. 1970. Studies in the Aegean flora XVI. Biosystematics of the *Nigella arvensis* complex with special reference to the problem of non-adaptive radiation. – Opera bot. 28.
- 1996. Phytogeographia Aegaea and the Flora Hellenica Database. – Ann. naturhist. Mus. Wien 98 B Suppl.: 279–289.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Phyton, Annales Rei Botanicae, Horn](#)

Jahr/Year: 2001

Band/Volume: [41\\_2](#)

Autor(en)/Author(s): Bergmeier Erwin, Dimopoulos Panayotis Panayotis

Artikel/Article: [Chances and Limits of Floristic Island Inventories - the Dionysades Group \(South Aegean, Greece\) Re-visited. 277-293](#)