

Lichenicolous Biota (Nos 121–150)

Josef HAFELLNER*

HAFELLNER J. 2012: Lichenicolous Biota (Nos 121–150). – Fritschiana (Graz) 74: 19–41. - ISSN 1024-0306.

Abstract: The sixth fascicle (30 numbers) of the exsiccata 'Lichenicolous Biota' is published. The issue contains material of 29 non-lichenized fungal taxa (17 ascomycetes, 6 basidiomycetes, 6 anamorphic fungi), including isotype material of *Abrothallus parmotrematis* Diederich (no. 131), *Endococcus thelommatis* Kocourk. & K.Knudsen (no. 136), *Laetisaria lichenicola* Diederich et al. (no. 137), *Trimmatostroma dendrographae* Diederich et al. (no. 129), and of *Trimmatostroma quercicola* Diederich et al. (no. 149), as well as isoneotype material of *Briancoppinsia cytospora* (Vouaux) Diederich et al. (no. 133), representing the type species of *Briancoppinsia*, and paratype material of *Trimmatostroma dendrographae* (no 130). Collections of the type species of *Cercidospora* (*C. macrospora*, syn. *C. ulothii*), *Marchandiobasidium* (*M. aurantiacum*), *Phaeopyxis* (*P. punctum*), *Rhagadostoma* (*R. lichenicola*, syn. *R. corrugatum*), *Stigmatidium* (*S. schaeferi*), *Telogalla* (*T. olivieri*), and *Xanthoriicola* (*X. physciae*) are also included in this fascicle.

*Institut für Pflanzenwissenschaften, Karl-Franzens-Universität Graz,
Holteigasse 6, A-8010 Graz, AUSTRIA.

e-mail: josef.hafellner@uni-graz.at

Introduction

The exsiccata 'Lichenicolous Biota' is continued with fascicle 6 containing 30 numbers.

The exsiccata covers all lichenicolous biota. The exsiccata is open to non-lichenized and lichenized fungi, but also to myxomycetes, bacteria, and even animals, whenever they cause a characteristic symptom on their host (e.g., discoloration or galls). Consequently, the exsiccata contains both highly host-specific and plurivorous species, as long as the individuals clearly grow upon a lichen and the collection is homogeneous, so that identical duplicates can be prepared.

The five complete sets are sent to herbaria of the following regions: Central Europe (Graz [GZU]), Northern Europe (Uppsala [UPS]), Western Europe (Bruxelles [BR]), North America (New York [NY]), Australasia (Canberra [CANB]). Incomplete sets will preferably be distributed to Barcelona [BCN], Edinburgh [E], Leningrad [LE], Munich [M], and Prague [PRM] (herbarium acronyms sec. HOLM-

GREN et al. 1990, continued and updated as electronic database by THIERS 2010, onwards and hosted at New York Botanical Garden (<http://sciweb.nybg.org/science2/IndexHerbariorum.asp>). It is planned to publish at least one fascicle per year, consisting of a variable number of decades.

For the 6th issue, I gratefully acknowledge the contribution of 13 collections by Paul Diederich (partly together with Damien Ertz), 2 collections by Walter Obermayer, 2 collections by Toby Spribille (together with L. Yakovchenko), and 1 collection each by Brian Coppins, Damien Ertz, Siegfried Huneck(†), Kerry Knudsen (together with S. Chaney, sent by Jana Kocourková), James Lendemer, and Rolf Santesson. In fieldwork I received support by Roderick Rogers. Brian Coppins, Paul Diederich, Damien Ertz, Jana Kocourková, James Lendemer, and Toby Spribille contributed to the scientific content of the fascicle by the identification of either lichenicolous fungi or hosts. Christian Scheuer, Walter Obermayer, and Helmut Mayrhofer are thanked for critically reading the manuscript.

I would be much obliged to colleagues who send material of lichenicolous biota for distribution in future fascicles. The collections should be divided up into at least 5 (up to 10) duplicates, preferably already prepared. Unprepared collections should be rich enough to obtain at least 5 duplicates.

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

121. *Dactylospora lobariella* (Nyl.) Hafellner

in Nova Hedwigia Beih. 62: 118 (1979). – Bas.: *Lecidea lobariella* Nyl. in Flora (Regensburg) 52: 296 (1869). – Syn.: *Buellia lobariella* (Nyl.) Arnold in Flora (Regensburg) 53: 478 (1870). – *Karschia lobariella* (Nyl.) Arnold in Flora (Regensburg) 57: 103 (1874). – *Abrothallus lobariellus* (Nyl.) Zopf in Hedwigia 35: 363 (1896).

Host: *Lobaria* spec. (thallus)

Asia, Russia: Khabarovskiy Krai, Komsomolsk-De Kastru route, Khomi Mountains, c. 20 air km E of Chernyy Mys, hills on north bank of the Salasu River, 51°05'55"N / 138°46'17"E, 328 m alt.; mixed coniferous-broad-leaved mature forest, at the base of a trunk of *Picea jezoensis*.

Note: The type host of *Dactylospora lobariella* is *Lobaria pulmonaria*.

12. VII. 2009 leg. T. Spribille (30509)
(excursion together with L. Yakovchenko, C. Printzen, B. Kanz & E. Malashkina)
det. J. Hafellner

distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

122. *Marchandiobasidium aurantiacum* Diederich & Schultheis

in Diederich et al., Mycol. Res. 107: 524 (2003).

Hosts: *Physcia adscendens* (thallus) and *Physcia tenella* (thallus)

Europe, Luxembourg: Dudelange, in the centre of the city, along Schwaarze Wee, 49°28'40"N / 06°05'02"E, c. 290 m alt., IFBL: M8.54.42, UTM: KV.88; trees on a parking area, on bark of *Platanus* spec.

Note: *Physcia tenella* is the type host of *Marchandiobasidium aurantiacum*.

15. II. 2010 leg. P. Diederich (16901), det. P. Diederich

distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

123. *Nectriopsis parmeliae* (Berk. & M.A.Curtis) M.S.Cole & D.Hawksw.

in Mycotaxon 77: 321 (2001). – Bas.: *Diplodia parmeliae* Berk. & M.A.Curtis in Berkeley, Grevillea 3(25): 3 (1874). – Syn.: *Nectria parmeliae* (Berk. & M.A.Curtis) D.Hawksw. in Bull. British Mus. Nat. Hist., Bot. ser., 9(1): 76 (1981).

Host: *Physcia subtilis* (thallus)

North America, U.S.A.: Pennsylvania, Lancaster County, E slopes above Susquehanna River, 0.25–0.5 miles N of confluence of Tucquan Creek and Susquehanna River, 39°51'30"N / 76°20'55"W, c. 100 m alt.; schist outcrops with *Quercus-Nyssa-Pinus* forest, on rock.

Note 1: In the protologue the host is only given on genus level (*Parmelia*).

Note 2: Hawksworth (l.c.) has reinvestigated the holotype in K and identified the host lichen as *Punctelia* cf. *subrudecta* (sub *Parmelia*).

08. VIII. 2009 leg. J. C. Lendemer (19351), det. J. C. Lendemer
distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

124. *Sagediopsis fissurisedens* Hafellner

in Herzogia 9(3–4): 757 (1993).

Host: *Aspilidea myrinii* (thallus)

Europe, Sweden: Härjedalen, Tännäs par., Mt. Gruvvålen, 0.3 km SSW of Glimsjön, c. 1.5 km ENE of Ramundbergets Fjällgård, 62°42'40"N / 12°25'E, c. 920 m alt.; on open siliceous rocks in a subalpine birch forest.

Note: *Aspilidea myrinii* is the type host of *Sagediopsis fissurisedens*.

11. VIII. 1987 leg. R. Santesson (31955), det. R. Santesson
distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

125. *Sphinctrina leucopoda* Nyl.

in Synopsis Meth. Lich. 1: 144 (1860).

Host: *Pertusaria* spec. (thallus, apothecial verrucae)

Australia: Queensland, Tandora, about 25 km ENE of Maryborough, 25°27'S / 152°52'E, at sea level; mangroves.

Note: The type host of *Sphinctrina leucopoda* is a North American *Pertusaria*. Nylander (l.c.) argued it might be *P. leioplaca*.

23. VIII. 1986 leg. J. Hafellner (18211) & R. Rogers, det. Hafellner
distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

126. *Tremella dendrographae* Diederich & Tehler

in Diederich, Biblioth. Lichenol. 61: 74 (1996).

Host: *Dendrographa minor* (syn. *Dendrographa leucophaea* f. *minor*) (thallus)

North America, U.S.A.: California, Monterey Co., SW of Monterey, 17-Mile Drive, Crocker Grove, 36°34'40"N / 121°58'20"W, c. 15 m alt.; coniferous forest close to the sea, on *Cupressus macrocarpa*.

Note: *Dendrographa minor* is the type host of the species.

19. VII. 2008 leg. P. Diederich (16794) & D. Ertz, det. P. Diederich
distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

127. *Tremella ramalinae* Diederich

in Biblioth. Lichenol. 61: 152 (1996).

Host: *Ramalina fraxinea* (thallus)

Europe, Austria: Steiermark (Styria), Northern Alps, Dachsteingruppe, Ramsau, 4.6 km NNW of the centre of Schladming, 0.28 km NE of the church of Ramsau-Ort and 0.5 km SE of the farmstead 'Feldlhof', 47°25'26"N / 13°39'28"E, c. 1150 m alt., GF 8547/4; alley, on twigs of *Acer pseudoplatanus* and *Fraxinus excelsior*.

Note: The type host of *Tremella ramalinae* is *Ramalina lacera*.

19. II. 2012

leg. W. Obermayer (12390), det. J. Hafellner

distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

128. *Trichonectria pertusariae* Etayo & P.Boom

in Nova Hedwigia 81: 160 (2005).

Host: *Pertusaria* spec. (thallus, apothecial margins)

Africa, Canary Islands: La Gomera, Arure, Mirador de Alojera, 28°09'04"N / 17°18'30"W, c. 980 m alt.; brezal (*Erica arborea* stand), on bark of *Erica arborea*.

Note: The type host of *Trichonectria pertusariae* is *Pertusaria ophthalmiza*.

2. IV. 2011

leg. D. Ertz (16252), det. D. Ertz

distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

**129. *Trimmatostroma dendrographae* Diederich, Ertz, U.Braun
& Heuchert **Isotype****

in Diederich et al., Bull. Soc. Naturalistes Luxemburgois 111: 49 (2010).

Host: *Dendrographa* spec. (thallus)

North America, U.S.A.: California, Monterey Co., Carmel, Point Lobos State Reserve, 36°31'18"N / 121°57'09"W, c. 30 m alt.; coniferous forest close to the sea, on *Cupressus macrocarpa*.

Note 1: The host thallus consists of a crustose primary thallus and a secondary fruticose one; both are infected.

Note 2: Paratype material is distributed under Lichenicolous Biota no. 130.

18. VII. 2008 leg. P. Diederich (16789) & D. Ertz, det. P. Diederich
distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

**130. *Trimmatostroma dendrographae* Diederich, Ertz, U.Braun
& Heuchert **Paratype****

in Diederich et al., Bull. Soc. Naturalistes Luxemburgois 111: 49 (2010).

Host: *Dendrographa* spec. (thallus)

North America, U.S.A.: California, Monterey Co., SW of Monterey, 17-Mile Drive, Crocker Grove, 36°34'40"N / 121°58'20"W, c. 15 m alt.; coniferous forest close to the sea, on *Cupressus macrocarpa*.

Note 1: The type host of the species is a so far undetermined *Dendrographa* likely to be identical with that of the paratype.

Note 2: The host thallus consists of a crustose primary thallus and a secondary fruticose one; both are infected.

Note 3: Isotype material is distributed under Lichenicolous Biota no. 129.

19. VII. 2008 leg. P. Diederich (16767) & D. Ertz, det. P. Diederich
distributed to: BCN, BR, CANB, E, GZU, LE, M, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

131. ***Abrothallus parmotrema*** Diederich **Isotype**

in Bull. Soc. Naturalistes Luxembourgeois 112: 29 (2011).

Host: *Parmotrema crinitum* (thallus)

Europe, United Kingdom: Scotland, Inner Hebrides, Isle of Skye, SSE of Broadford, Ardnearcan [on the shore of Loch na Dal], 57°10'05"N / 5° 46'50"W, c. 30 m alt., NG 7 1; deciduous forest with dominant *Corylus avellana*, on bark of unidentified tree.

Note: The name was originally introduced in Clauzade et al., Bull. Soc. Linn. Provence, Num. Spéc. 1: 24 (1989) but the species was not formally described at that time.

27. V. 1987

leg. P. Diederich (8301), det. P. Diederich

distributed to: BR, CANB, E, GZU, M, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

132. ***Abrothallus peyritschii*** (Stein) I.Kotte

in Centralbl. Bakt. Paras., 2. Abt., 24: 76 (1909). – Bas.: *Abrothallus parmeliarum* var. *peyritschii* Stein, Flechten in Cohn, Kryptfl. Schles. 2(2): 211 (1879). – Syn.: *Abrothallus parmeliarum* f. *peyritschii* (Stein) Arnold in Flora (Regensburg) 68: 221 (1885).

Host: *Vulpicida pinastris* (thallus)

Asia, Russia: Khabarovskiy Krai, Bureinskiy Zapovednik, upper reach of the Pravaya Bureya River, Tsarskaya Doroga, ca. 650 m N of patrol cabin 'Staraya Medvezhka', 25 km SE of Sofiysk and 2.6 km N (upstream) of the patrol cabin 'Novaya Medvezhka'. 52°09'00"N / 134°19'00"E, c. 870 m alt.; mossy *Larix gmelinii*-*Vaccinium vitis-idaea* forest, on branches of *Pinus pumila*.

Note: *Vulpicida pinastris* is the type host of *Abrothallus peyritschii*.

2. VIII. 2009

leg. T. Spribille (31780) with L. Yakovchenko, det. T. Spribille

distributed to: BCN, BR, CANB, GZU, LE, M, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

133. *Briancoppinsia cytospora* (Vouaux) Diederich, Ertz, Lawrey & P.Boom *Isonotype*

in Fungal Diversity 52(1): 8 (2011). – Bas.: *Phyllosticta cytospora* Vouaux in Bull. Soc. Mycol. France 30: 193 (1914). – Syn.: *Phoma cytospora* (Vouaux) D.Hawksw. in Trans. Brit. Mycol. Soc. 67: 56 (1976).

Host: *Lecanora conizaeoides* (thallus)

Europe, Luxembourg: N of Reckange (Mersch), Enelter Kapelle, 49°45'35"N / 6°04'42"E, c. 265 m alt., IFBL: L8.24.44; UTM: KA.81; deciduous trees near the chapel, on old trunk of *Aesculus hippocastanum*.

Note 1: According to the protologue, the type host of *Phyllosticta cytospora* is *Flavoparmelia caperata*.

Note 2: The identification of the host in the exsiccata distributed here is uncertain.

Note 3: Diederich et al. (Fungal Diversity 52(1): 1–12, 2011) have shown that the coelomycete *Briancoppinsia cytospora* is likely to be a member of the family *Arthoniaceae* belonging to the Arthoniales-Arthoniomycetes.

Note 4: The host is partly killed by *Athelia arachnoidea* (e.g. specimen in GZU).

3. IX. 2009 leg. P. Diederich (16849), det. P. Diederich
distributed to: BR, CANB, E, GZU, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

134. *Cercidospora macrospora* (Uloth) Hafellner & Nav.-Ros.

in Navarro-Rosinés et al., Lich. Fl. Greater Sonoran Desert Region 2: 638 (2004). – Bas.: *Phacopsis macrospora* Uloth in Flora (Regensburg) 44: 653 (1861). – Syn.: *Cercidospora ulothii* Körb. in Parerga Lich.: 465 (1865).

Host: *Protoparmeliopsis muralis* sensu auct. (thallus)

Europe, Germany: Sachsen-Anhalt, Wimmelburg c. 2.5 km WSW of Eisleben, mining area N of the village, 51°31'25"N / 11°30'55"E, c. 190 m alt., MTB 4435/3; mining waste tip of copper-rich slate, on small stones.

Note 1: *Protoparmeliopsis muralis* sensu auct. (syn. *Lecanora muralis*) is the type host of *Cercidospora macrospora* and its syn. *C. ulothii*.

Note 2: The genus *Cercidospora* is based on *C. ulothii* Körb., a later heterotypic synonym of *Phacopsis macrospora* Uloth.

10. III. 1991 leg. S. Huneck, det. J. Hafellner
distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

135. *Cornutispora pyramidalis* Etayo

in Opuscula Philolichenum 8: 134 (2010).

Host: *Parmotrema robustum* (thallus)

Europe, Azores: São Miguel, Furnas, on the shore of Lagoa das Furnas, 37° 46'06"N / 25°19'54"W, c. 285 m alt.; planted trees on the lakeshore, on bark of *Cryptomeria japonica*.

Note: The type host of *Cornutispora pyramidalis* is *Hypotrachyna revoluta*.

30. VII. 2010

leg. P. Diederich (16968), det. P. Diederich

distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

**136. *Endococcus thelommatis* Kocourk. & K.Knudsen
Isotype**

in Biblioth. Lichenol. 106: 174 (2011).

Host: *Thelomma santessonii* (thallus, apothecial margins)

North America, U.S.A.: California, Santa Barbara Co., Santa Rosa Island, Channel Islands National Park, South Point, 33°54'27"N / 120°07'27"W, 299 m alt.; coastal sage shrub, on boulders of sandstone.

Note: *Thelomma santessonii* is the type host of the species.

14. VI. 2009

leg. K. Knudsen (11421) & S. Chaney, det. J. Kocourkova

distributed to: BCN, BR, CANB, GZU, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

**137. *Laetisaria lichenicola* Diederich, Lawrey & Van den Broeck
Isotype**

in Diederich et al., Mycologia 103(3): 530 (2011).

Hosts: *Physcia adscendens* (thallus) and *Physcia tenella* (thallus)

Europe, Luxembourg: Dudelage, in the centre of the city, along Schwaarze Wee, 49°28'40"N / 06°05'02"E, c. 290 m alt., IFBL: M8.54.42, UTM: KV.88; trees on a parking area, on bark of *Platanus* spec.

Note 1: All specimens are sterile.

Note 2: *Laetisaria lichenicola* also attacks *Phaeophyscia orbicularis* (specimen in BR) and *Xanthoria parietina* (specimen in GZU).

Note 3: The specimen in GZU is also infected with *Marchandiobasidium aurantiacum*. Material of that species is distributed under Lichenicolous Biota no. 122.

15. II. 2010 leg. P. Diederich (16900), det. P. Diederich
distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

138. *Minutoexcipula mariana* V. Atienza

in Biblioth. Lichenol. 82: 142 (2002).

Host: *Pertusaria* spec. (thallus, apothecial verrucae)

Europe, Azores: Pico, Lajes do Pico, along main road inside village, 38°23' 37"N / 28°15'03"W, c. 15 m alt.; row of trees beside the road, on bark of *Dracaena* spec..

Note: The type host of *Minutoexcipula mariana* is *Pertusaria heterochroa*.

25. VII. 2010 leg. P. Diederich (17023), det. P. Diederich
distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

139. ***Muellerella hospitans*** Stizenb.

in Nova Acta Acad. Leop.-Carol. 30(3): 51 (1863). – Syn.: *Muellerella polyspora* var. *hospitans* (Stizenb.) Keissl. in Ark. Bot. 18(16): 4 (1923).

Host: *Bacidia* spec. (apothecia)

Europe, Albania: Northern Albania, Lezhë distr., at the coast of the Adriatic Sea SW of the town Lezhë, Kuna Vain lagoon S of the inlet of the river Drin, 41°44'20"N / 19°34'50"E, c. 5 m alt.; remnants of coastal forest c. 200 m behind the shore, on bark of old *Fraxinus* spec.

Note: The type host of *Muellerella hospitans* is *Bacidia rubella*.

16. VIII. 2007 leg. J. Hafellner (80823), det. J. Hafellner
(excursion together with M. Tretiach, L. Muggia, M. Piccotto & J. Marka)

distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

140. ***Phaeopyxis punctum*** (A.Massal.) Rambold, Triebel & Coppins

in Rambold & Triebel, Notes Roy. Bot. Garden Edinburgh 46: 384 (1990). – Bas.: *Nesolechia punctum* A.Massal. in Sched. Crit. Lich. Exs. Italiae 5: 96 (1856). – Syn.: *Lecidea punctum* (A.Massal.) Jatta in Syll. Lich. Ital.: 353 (1900).

Host: *Cladonia digitata* (thallus) and *C. coniocraea* (thallus)

Europe, Austria: Vorarlberg, Eastern Alps, Verwall-Gruppe, Silbertal c. 8 km east of the town Schruns, above the inn Fellimännle, 47°04'45"N / 10°01'30"E, c. 1100 m alt., GF 8926/1; talus of forest road on slope exposed to the W, on mossy soil.

Note 1: In the protologue the host is only given to genus level (*Cladonia*).

Note 2: Type material has been distributed in Massalongo, Lich. Ital. Exs. 153. Rambold & Triebel (l.c.) revised two isotypes stored in M but did not mention the type host.

30. VIII. 2008 leg. J. Hafellner (80842), det. J. Hafellner

distributed to: BR, CANB, GZU, M, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

141. *Plectocarpon macaronesiae* Diederich, Etayo & Sérusiaux

in Diederich & Etayo, Nordic J. Bot. 14: 592 (1994).

Host: *Lobaria macaronesica* (thallus)

Europe, Azores: Pico, between Lajes do Pico and São Roque do Pico, Bosque da Junqueira, 1 km S of a junction with a road going to the E, 38°27'56"N / 28°17'57"W, c. 650 m alt.; laurel forest, on *Vaccinium cylindraceum*.

Note 1: The type host of *Plectocarpon macaronesiae* is *Lobaria* cf. *meridionalis*.

Note 2: The specimen in GZU is also infected by *Tremella lobariacearum*.

26. VII. 2010

leg. P. Diederich (17076), det. P. Diederich

distributed to: BCN, BR, CANB, GZU, LE, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

142. *Polycoccum minutulum* Kocourk. & F. Berger

in Czech Mycology 51: 171 (1999).

Host: *Trapelia placodioides* (thallus)

Europe, Austria: Steiermark (Styria), Eastern Alps, Steirisches Randgebirge, Grazer Bergland, in the gorge Raabklamm W of the town Weiz, orographically right slope E below the village Kapellenfeld, rocky ridge c. 0.6 km NW of the castle Gutenberg, 47°13'00"N / 15°33'50"E, c. 580 m alt., GF 8759/3; sparse mixed forest with small outcrops of siliceous schist, on loose plates lying on the ground.

Note: *Trapelia placodioides* is the type host of *Polycoccum minutulum*.

9. IX. 2009

leg. J. Hafellner (74056), det. J. Hafellner

distributed to: BCN, BR, CANB, GZU, NY, PRM, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

143. ***Rhagadostoma lichenicola*** (De Not.) Keissl.

in Rabenh. Krypt.-Fl., 2. Aufl. 8: 320 (1930). – Bas.: *Bertia lichenicola* De Not. in Erb. Crittog. Ital. no. 1190 (1864).

Host: *Solorina crocea* (thallus)

Europe, Austria: Kärnten (Carinthia), Eastern Alps, Saualpe W of the town Wolfsberg, short SSW of the saddle between the mountains Forstalpe and Geierkogel, 46°54'30"N / 14°40'05"E, c. 1850 m alt., GF 9054/3; outcrops of eclogite along the crest, exposed to the N, on soil.

Note 1: *Solorina crocea* is the type host of *Rhagadostoma lichenicola*.

Note 2: The genus *Rhagadostoma* is based on *R. corrugatum* Körb. a later heterotypic synonym of *Bertia lichenicola* De Not.

4. IX. 2011

leg. J. Hafellner (78506), det. J. Hafellner

distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

144. ***Stigmidium schaeferi*** (A.Massal.) Trevis.

in Conspectus Verruc.: 17 (1860). – Bas.: *Sphaeria schaeferi* A.Massal. in Sulla *Lecidea hookeri* nota: 8 (1853) '*schaererii*'. – Syn.: *Pharcidia schaeferi* (A.Massal.) Arnold in Verh. Zool.-Bot. Ges. Wien 19: 638 (1896). – *Verrucaria schaeferi* (A. Massal.) Nyl. in Flora (Regensburg) 53: 358 (1865). – *Sphaerella schaeferi* (A. Massal.) Anzi in Atti Soc Ital. Sci. Nat. 11: 180 (1868). – *Epicymatia schaeferi* (A.Massal.) Sacc. in Syll. Fung. 1: 571 (1882). – *Sphaerulina schaeferi* (A.Massal.) Sacc. & D.Sacc. in Syll. Fung. 17: 695 (1905).

Host: *Dacampia hookeri* (thallus)

Europe, Austria: Salzburg, Pinzgau, Eastern Alps, Hohe Tauern, Glockner-Gruppe, northern slopes of Kitzsteinhorn, c. 0.5 km W of Bundessportheim, 47°12'35"N / 12°41'10"E, c. 2450 m alt., GF 8742/3; open alpine vegetation over calcareous schist, on soil.

Note 1: *Dacampia hookeri* is the type host of *Stigmidium schaeferi*.

Note 2: *Dacampia hookeri* is lichenicolous itself, usually upon species of the *Solorina bisporea* group (see Henssen, Crypt. Bot. 5: 149–158, 1995) but often, as is the case in the material distributed here, the primary host is not recognizable anymore.

27. VIII. 1996

leg. J. Hafellner (37986), det. J. Hafellner

distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

145. ***Stigmatidium tabacinae*** (Arnold) Triebel

in Biblioth. Lichenol. 35: 236 (1989). – Bas.: *Pharcidia tabacinae* Arnold in Flora (Regensburg) 64: 176 (1881). – Syn.: *Sphaerulina tabacinae* (Arnold) Vouaux in Bull. Soc. Mycol. France 29: 34 (1913).

Host: *Toninia tristis* (thallus)

North America, U.S.A.: Alaska, Yukon-Koyukuk Census Area, Brooks Range, Endicott Mountains, Sukakpak Mountain, foothills on the W side, 67°35'35"N / 149°46'00"W, c. 580 m alt.; boulders of calcareous sandstone in open boreal forest, on soil.

Note: *Toninia tristis* (sub syn. *Psora tabacina*) is the type host of *Stigmatidium tabacinae*.

21. VIII. 2010 leg. J. Hafellner (79969), det. J. Hafellner
(excursion together with T. Spribille, L. Muggia and C. Hampton-Miller)

distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

146. ***Teloggalla olivieri*** (Vouaux) Nik.Hoffm. & Hafellner

in Biblioth. Lichenol. 77: 109 (2000). – Bas.: *Laestadia olivieri* Vouaux in Bull. Soc. Mycol. France 28: 216 (1912) nomen novum for *Verrucaria xanthoriae* Wedd. f. *megaspora* H.Olivier in Princ. Paras. Lich. Fr. Suppl. 1: 14 (1907).

Host: *Xanthoria parietina* (thallus)

Europe, Luxembourg: SSW of Belvaux, Kiemreech, 49°29'57"N / 5°53'38"E, c. 370–390 m alt., IFBL M8.51.21; ancient mining area, on twigs of *Salix caprea*.

Note: *Xanthoria parietina* is the type host of *Teloggalla olivieri*.

8. IX. 2012 leg. P. Diederich (17474), det. P. Diederich

distributed to: BR, CANB, GZU, LE, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

147. *Tremella caloplacae* (Zahlbr.) Diederich

in Sérusiaux et al., *Lejeunia* 173: 31 (2003). – Bas.: *Lindauopsis caloplacae* Zahlbr. in Ber. Deutsch. Bot. Ges. 24: 145 (1906).

Host: *Xanthoria parietina* (apothecia)

Europe, Austria: Steiermark (Styria), Dachsteingruppe (S-facing slopes), 4.6 km NNW of the centre of Schladming, Ramsau, 1.4 km W of the church of Ramsau-Ort, road to Vorberg near an underpass, 47°25'16"N / 13°38'10"E, c. 1170 m alt., GF 8747/4; roadside trees, on twigs of *Acer pseudoplatanus*.

Note: The type host of *Tremella caloplacae* is *Caloplaca aurantia*.

21. II. 2012

leg. W. Obermayer (12446), det. J. Hafellner

distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

148. *Tremella lobariacearum* Diederich & M.S.Christ.

in Diederich, *Biblioth. Lichenol.* 61: 103 (1996).

Host: *Lobaria pulmonaria* (thallus)

Europe, Azores: São Miguel, Furnas, Terra Nostra park, 37°46'15"N / 25°18'42"W, c. 220 m alt.; arboretum, on various exotic trees.

Note: *Lobaria pulmonaria* is the type host of the species.

30. VII. 2010

leg. P. Diederich (16979), det. P. Diederich

distributed to: BCN, BR, CANB, GZU, M, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

149. ***Trimmatostroma quercicola*** Diederich, U.Braun & Heuchert
Isotype

in Diederich et al., Bull. Soc. Naturalistes Luxembourgeois 111: 52 (2010)

Host: bark of phorophyte, algal colonies or sterile crustose lichen thallus

Europe, Luxembourg: Vogelsmühle, Halerbaach valley, on the orographically left river bank, 49°49'22"N / 06°18'51"E, c. 250 m alt. IFBL: L9.11.11, UTM: LA.02; deciduous forest, on bark of *Quercus* spec.

Note: The biology of the species remains unclear.

9. XII. 2007 leg. P. Diederich (16723), det. P. Diederich
distributed to: BR, CANB, GZU, NY, UPS

Hafellner J. 2012: Lichenicolous Biota (Nos 121–150). - Fritschiana 74: 19–41.

150. ***Xanthoriicola physciae*** (Kalchbr.) D.Hawksw.

in Hawksworth & Punithalingam, Trans. Brit. Mycol. Soc. 61: 67 (1973). – Bas.: *Gymnosporium physciae* Kalchbr. in Math. Term. Közlem. 3: 299 (1865). – Syn.: *Coniosporium physciae* (Kalchbr.) Sacc. in Syll. Fung. 4: 246 (1886).

Host: *Xanthoria parietina* (apothecia)

Europe, United Kingdom: Scotland, East Lothian (VC 82), Dunbar, Belhaven Bay, 56°00'6"N / 02°33'46"W, c. 5 m alt.; at edge of conifer plantation on coastal sand dunes, on twigs of *Sambucus nigra*.

Note1: *Xanthoria parietina* is the type host of the species.

Note 2: Ruibal et al. (IMA Fungus 2(1): 97–103, 2011) have shown that the dematiaceous hyphomycete *Xanthoriicola physciae* is likely to be a member of the family *Teratosphaeriaceae* belonging to the *Capnodiales*-Dothideomycetes.

18. III. 2012 leg. B. J. Coppins (23901), det. B. J. Coppins
distributed to: BR, CANB, GZU, NY, UPS

References

- BRUMMITT, R.K. 2001: World geographical scheme for recording plant distributions. Edition 2. Plant Taxonomic Database Standards No. 2. Edition 2, August 2001. Published for the International Working Group on Taxonomic Databases For Plant Sciences (TDWG) by the Hunt Institute for Botanical Documentation, Carnegie Mellon University, Pittsburgh. 137 pp.
- HOLMGREN, P.K., HOLMGREN, N.H. & BARNETT, L.C. (eds.) 1990: Index herbariorum. Part I. The herbaria of the world. 8th edition. - Bronx, New York: New York Botanical Garden for the International Association for Plant Taxonomy. Regnum Vegetabile 120: 693 pp.
- THIERS, B. 2010 [continuously updated]: Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. - URL: <http://sweetgum.nybg.org/ih/> [23. XI. 2012].

Bibliographic data of fascicles so far issued

- HAFELLNER, J. 2007: Lichenicolous Biota (Nos 1–20). - Fritschiana (Graz) 60: 35–49. URL: <http://www-classic.uni-graz.at/botwww/journals/fritschiana/fritschiana-60/lichenicolous-biota-nos-1-20.pdf>
- HAFELLNER, J. 2008: Lichenicolous Biota (Nos 21–60). - Fritschiana (Graz) 61: 1–28. URL: <http://www-classic.uni-graz.at/botwww/journals/fritschiana/fritschiana-61/hafellner-2008-lichenicolous-biota-21-to-60.pdf>
- HAFELLNER, J. 2009: Lichenicolous Biota (Nos 61–80). - Fritschiana (Graz) 65: 33–46. URL: <http://www-classic.uni-graz.at/botwww/journals/fritschiana/fritschiana-65/lichenicolous-biota-nos-61-80.pdf>
- HAFELLNER, J. 2010: Lichenicolous Biota (Nos 81–100). – Fritschiana (Graz) 67: 11–26. URL: <http://www-classic.uni-graz.at/botwww/journals/fritschiana/fritschiana-67/hafellner-2010-lichenicolous-biota-nos-81-100.pdf>
- HAFELLNER, J. 2012: Lichenicolous Biota (Nos 101–120). – Fritschiana (Graz) 74: 1–17. URL: <http://www-classic.uni-graz.at/botwww/journals/fritschiana/fritschiana-74/hafellner-2012-lichenicolous-biota-nos-101-120.pdf>

Taxon Synopsis:

Taxon	Exs. no.
Ascomycota	
Arthoniomycetes	
<i>Abrothallus parmotrematis</i>	131
<i>Abrothallus peyritschii</i>	132
<i>Briancoppinsia cytospora</i>	133
<i>Plectocarpon macaronesiae</i>	141
Lecanoromycetes	
<i>Dactylospora lobariella</i>	121
Leotiomycetes	
<i>Phaeopyxis punctum</i>	140
Sordariomycetes (incl. Hypocreales)	
<i>Nectriopsis parmeliae</i>	123
<i>Rhagadostoma lichenicola</i>	143
<i>Trichonectria pertusariae</i>	128
Eurotiomycetes (incl. Verrucariales and Mycocaliciales)	
<i>Endococcus thelommatis</i>	136
<i>Muellerella hospitans</i>	139
<i>Sphinctrina leucopoda</i>	125
<i>Telogalla olivieri</i>	146
Dothideomycetes	
<i>Cercidospora macrospora</i>	134
<i>Dacampia hookeri</i>	144
<i>Polycoccum minutulum</i>	142
<i>Sagediopsis fissurisedens</i>	124
<i>Stigmatidium schaereri</i>	144
<i>Stigmatidium tabacinae</i>	145
<i>Xanthoriicola physciae</i>	150
Anamorphic Fungi (unclassified)	
Hyphomycetes	
<i>Trimmatostroma dendrographae</i>	129, 130
<i>Trimmatostroma quercicola</i>	149
Coelomycetes	
<i>Cornutispora pyramidalis</i>	135
<i>Minutoexcipula mariana</i>	138
Basidiomycota	
Agaricomycetes	
<i>Athelia arachnoidea</i>	133
<i>Laetisaria lichenicola</i>	137
<i>Marchandiobasidium aurantiacum</i>	122

Pucciniomycetes

Tremellomycetes

<i>Tremella caloplacae</i>	147
<i>Tremella dendrographae</i>	126
<i>Tremella lobariacearum</i>	141, 148
<i>Tremella ramalinae</i>	127

Host Index:

Host taxon	Lichenicolous taxon	Exs. no.
<i>Aspilidea myrinii</i>	<i>Sagediopsis fissurisedens</i>	124
<i>Bacidia</i> spec.	<i>Muellerella hospitans</i>	139
<i>Cladonia coniocraea</i>	<i>Phaeopyxis punctum</i>	140
<i>Cladonia digitata</i>	<i>Phaeopyxis punctum</i>	140
<i>Dacampia hookeri</i>	<i>Stigmidium schaeeri</i>	144
<i>Dendrographa minor</i>	<i>Tremella dendrographae</i>	126
<i>Dendrographa</i> spec.	<i>Trimmatostroma dendrographae</i>	129, 130
<i>Lecanora conizaeoides</i>	<i>Briancoppinsia cytospora</i>	133
<i>Lecanora conizaeoides</i>	<i>Athelia arachnoidea</i>	133
<i>Lobaria macaronesica</i>	<i>Plectocarpon macaronesicae</i>	141
<i>Lobaria macaronesica</i>	<i>Tremella lobariacearum</i>	141
<i>Lobaria pulmonaria</i>	<i>Tremella lobariacearum</i>	148
<i>Lobaria</i> spec.	<i>Dactylospora lobariella</i>	121
<i>Parmotrema crinitum</i>	<i>Abrothallus parmotrematis</i>	131
<i>Parmotrema robustum</i>	<i>Cornutispora pyramidalis</i>	135
<i>Pertusaria</i> spec.	<i>Sphinctrina leucopoda</i>	125
<i>Pertusaria</i> spec.	<i>Trichonectria pertusariae</i>	128
<i>Pertusaria</i> spec.	<i>Minutoexcipula mariana</i>	138
<i>Physcia adscendens</i>	<i>Laetisaria lichenicola</i>	137
<i>Physcia adscendens</i>	<i>Marchandiobasidium aurantiacum</i>	122
<i>Physcia subtilis</i>	<i>Nectriopsis parmeliae</i>	123
<i>Physcia tenella</i>	<i>Laetisaria lichenicola</i>	137
<i>Physcia tenella</i>	<i>Marchandiobasidium aurantiacum</i>	122
<i>Protoparmeliopsis muralis</i>	<i>Cercidospora macrospora</i>	134
<i>Ramalina fraxinea</i>	<i>Tremella ramalinae</i>	127
<i>Solorina crocea</i>	<i>Rhagadostoma lichenicola</i>	143
<i>Thelomma santessonii</i>	<i>Endococcus thelommatis</i>	136
<i>Toninia tabacina</i>	<i>Stigmidium tabacinae</i>	145
<i>Trapelia placodioides</i>	<i>Polycoccum minutulum</i>	142
<i>Vulpicida pinastri</i>	<i>Abrothallus peyritschii</i>	132
<i>Xanthoria parietina</i>	<i>Teloggalla olivieri</i>	146
<i>Xanthoria parietina</i>	<i>Tremella caloplacae</i>	147
<i>Xanthoria parietina</i>	<i>Xanthoriicola physciae</i>	150
Lichen indet.	<i>Trimmatostroma quercicola</i>	149

Geographic Index:**BIOGEOGRAPHIC UNITS (see BRUMMITT 2001)**

Country (or Archipelago)	Lichenicolous taxon	Exs. no.
1. EUROPE		
Albania	<i>Muellerella hospitans</i>	139
Austria	<i>Dacampia hookeri</i>	144
	<i>Phaeopyxis punctum</i>	140
	<i>Polycoccum minutulum</i>	142
	<i>Rhagadostoma lichenicola</i>	143
	<i>Stigmidium schaeereri</i>	144
	<i>Tremella caloplacae</i>	147
	<i>Tremella ramalinae</i>	127
Azores	<i>Cornutispora pyramidalis</i>	135
	<i>Minutoexcipula mariana</i>	138
	<i>Plectocarpon macaronesiae</i>	141
	<i>Tremella lobariacearum</i>	141, 148
Germany	<i>Cercidospora macrospora</i>	134
Luxembourg	<i>Athelia arachnoidea</i>	133
	<i>Briancoppinsia cytospora</i>	133
	<i>Laetisaria lichenicola</i>	137
	<i>Marchandiobasidium aurantiacum</i>	122
	<i>Telogalla olivieri</i>	146
	<i>Trimmatostroma quercicola</i>	149
Portugal (see under Europe, Azores)		
Spain (see under Africa, Canary Islands)		
Sweden	<i>Sagediopsis fissurisedens</i>	124
United Kingdom	<i>Abrothallus parmotrematis</i>	131
	<i>Xanthoriicola physciae</i>	150
2. AFRICA		
Canary Islands	<i>Trichonectria pertusariae</i>	128
3. ASIA TEMPERATE		
Russia (Asian part)	<i>Abrothallus peyritschii</i>	132
	<i>Dactylospora lobariella</i>	121
4. ASIA TROPICAL		
5. AUSTRALASIA		
Australia	<i>Sphinctrina leucopoda</i>	125

6. PACIFIC

7. NORTHERN AMERICA

U.S.A.

<i>Endococcus thelommatis</i>	136
<i>Nectriopsis parmeliae</i>	123
<i>Stigmidium tabacinae</i>	145
<i>Tremella dendrographae</i>	126
<i>Trimmatostroma dendrographae</i>	129, 130

8. SOUTHERN AMERICA

9. ANTARCTIC

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Fritschiana](#)

Jahr/Year: 2012

Band/Volume: [74](#)

Autor(en)/Author(s): Hafellner Josef

Artikel/Article: [Lichenicolous Biota \(Nos 121-150\) 19-41](#)