

PLANTERS

10% GRANULAR NEMATOCIDE

A SOIL FUMIGANT FOR USE IN CONTROLLING
MANY PARASITIC NEMATODES

ACCEPTED
APR 19 1968

ACTIVE INGREDIENTS:		
1,2-dibromo-3-chloropropane and other halogenated CG compounds	10.0%	
INERT INGREDIENTS	90.0%	
Total	100.0%	
Contains 10 pounds of active ingredient per 100 pounds		

Not For Use Or Storage In Or Around The Home

WARNING KEEP OUT OF REACH OF CHILDREN

EPA REG. NO. 1868-204

See Side Panels For Precautionary and Warning Statements



BEANS (Snap). For overall treatment, preplanting, at time of planting or postplanting application to mineral soils, apply 240 lbs. per acre. For preplanting application, wait 7-14 days before planting. Do not feed vines to livestock.

BROCCOLI, CABBAGE, CAULIFLOWER, CELERY, LETTUCE, OKRA and RADISHES: For overall treatment, preplanting, at time of planting or postplanting application to mineral soils, apply 240 lbs. per acre. If preplanting application is made, wait 7-14 days before planting.

COTTON, EGGPLANT and SQUASH (Summer): For overall treatment, preplanting, at time of planting or postplanting application to mineral soils, apply 260 lbs. per acre. For preplanting use, wait 7-14 days before planting.

MELONS: For overall treatment, preplanting or at any time of planting treatment, apply 260 lbs. per acre. For preplanting use, wait 7-14 days before planting.

For row treatment only, preplanting or at time of planting treatment, apply 90 lbs per 8,712 linear feet of row. For preplanting use, wait 7-14 days before planting.

TOMATOES: For overall treatment, preplanting application only, apply 260 lbs. per acre. Wait 7-14 days before planting.

For row treatment only, preplanting, at planting or postplanting for direct seeded tomatoes, or preplanting or at planting for transplants, apply 180 lbs. per 13,068 linear feet of row. Set transplants 4 inches from injection furrows. Wait 7-14 days before planting or preplanting treatments.

CUCUMBERS: For overall treatment preplanting application only to mineral soils, apply 260 lbs. per acre. For preplanting use, wait 7-14 days before planting.

For row treatment only, preplanting, at planting or postplanting application, apply 150 lbs. per 8,712 linear feet of row.

CARROTS and PARSNIPS: For overall treatment preplanting, at time of planting or postplanting to mineral soils, apply 240 lbs. per acre. For preplanting use, wait 7-14 days before planting. Do not feed plant parts to livestock.

PEANUTS: For overall treatment, preplanting at planting treatment, apply 260 lbs. per acre. Do not feed soil or residues into commerce. Hay bails taken from treated soil. Forage crops grown on treated soil should not be used as feed for dairy cows or animals being finished for slaughter for 1-2 years after last treatment or 1 year after overall treatment.

STRAWBERRIES: For overall treatment, preplanting, at time of planting or postplanting application to mineral soil, apply 260 lbs. per acre. For preplanting use, wait 7-14 days before planting. Avoid at least 10 days between treatment and harvest.

APPLICATION DIRECTIONS

Apply PLANTERS 10% GRANULAR NEMATOCIDE when soil temperatures at a depth of 6 to 8 inches are in the 50° to 70° F. range. Soil should be moist and contain adequate moisture for good seed germination. Do not use on wet soils. Do not use on soils that are highly acidic. Apply to mineral soils only. Do not use on soils that are highly saline. Do not use on soils that are highly alkaline. Do not use on soils that are highly acidic. Do not use on soils that are highly alkaline. Do not use on soils that are highly acidic. Do not use on soils that are highly alkaline.

PRE-PLANT TREATMENTS: In the row. Apply in the row at a depth of 6 to 8 inches, taking care to seal the material in the soil with appropriate equipment. Use care to avoid contact with the suggested application equipment. See the next paragraph.

OVERALL: Apply evenly in the row 12 to 14 inches apart at a depth of 6 to 8 inches, taking care to seal the material in the soil with appropriate equipment. Use care to avoid contact with the suggested application equipment. See the next paragraph.

SOIL PREPARATION

VENTILATION AND PREPARATION OF SOIL BEFORE PLANTING: For preplanting application, the soil should be aerated by deep cultivation at the end of the exposure period. If heavy rains accompanied by low temperatures occur during the exposure period, additional aeration is required and the soil should be worked several times before planting. Aeration is usually complete when the odor of the fumigant is no longer evident. Shallow rooted crops can usually be planted about one week after the end of the exposure period.

PRECAUTIONS

To avoid reinfestation of treated soil, do not use transplants, tools or crop residues that may be contaminated with soil-borne pests.

Use PLANTERS 10% GRANULAR NEMATOCIDE only according to label directions.

This product is not recommended for use on extremely heavy soil.

WARNING

HARMFUL VAPOR
CAUSES SKIN IRRITATION AND BLISTERS ON PROLONGED CONTACT
MAY BE ABSORBED THROUGH SKIN
COMBUSTIBLE MIXTURE
HARMFUL IF SWALLOWED

Do not breathe vapor. Keep container closed. Do not get in eyes, on skin, or on clothing.

Do not take internally. Use only in a well ventilated area.

Keep away from heat and open flame. Keep out of reach of children and animals.

In case of contact, immediately remove contaminated clothing and wash skin with soap and water. Flush eyes with plenty of water for at least 15 minutes. Get medical attention. Do not induce vomiting. Do not use ABSOLUTELY NO OTHER REMEDIES.

If necessary, seek medical attention. Do not use on animals or children.

If swallowed, call a doctor and show container. Do not induce vomiting. Get medical attention. Do not use on animals or children.

Do not use on animals or children. Do not use on animals or children.