



400-577

03-18-2010

11

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	EPA Registration Number: 400-577	Date of Issuance: MAR 18 2010
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)	
	Term of Issuance: Conditional	
Name of Pesticide Product: Belmont 2.7 FS		
Name and Address of Registrant (include ZIP Code): Chemtura Corporation Attn: Crop Registration 199 Benson RD (2-5) Middlebury, CT 06749		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3 (c) (7) (A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit/cite all data required for the registration/ reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the following label changes: <p>Page 1 – EPA Registration Number: Change the EPA Registration Number to 400-577</p> <p>(Continued)</p>		
Signature of Approving Official:  Mary L. Waller, Product Manager 21 Fungicide Branch Registration Division (7504P)	Date: 3/18/2010	

Continued from first page

EPA Registration Number 400-577

Should you wish to retain a reference to the company's website on your label, please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

As an alternative, you may refer consumers to the company's phone number or e-mail address.

3. Provide the following as conditions of registration:

- a. OPPTS number 830.6317 - storage stability - due July 1, 2011
- b. OPPTS number 830.6320 - corrosion characteristics - July 1, 2011

4. Submit one (1) copy of the revised final printed label before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions, contact Lisa Jones of my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely,



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped copy of label
Product chemistry review dated 22 January 2010
Acute toxicity review dated 25 January 2010

Belmont™ 2.7 FS

Seed Treatment Fungicide
for control of seed rot and damping-off diseases of certain crops

Active Ingredient:	
metalaxyl (CAS No. 57837-19-1).....	28.98%*
Other Ingredients:.....	71.02%
Total:.....	100.00%

*Contains 2.65 lbs. metalaxyl per gallon, or 317 grams per liter.

KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. EMERGENCY PHONE 800-292-5898 SAFETY DATA AND INFORMATION 866-430-2775 TRANSPORTATION EMERGENCY (CHEMTREC) 800-424-9300	

Net Contents:

EPA FILE SYMBOL 400-LTT
EPA EST. NO.
001/

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 18 2010**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**



Manufactured for:
Chemtura Corporation
199 Benson Road
Middlebury, CT 06749

400-577

www.chemtura.com

5/11

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING. Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes. Do not get on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Coveralls over short-sleeved shirt and short pants, socks and chemical-resistant footwear and chemical-resistant gloves (such as natural rubber > 14 mils in thickness). When mixing and loading, and when cleaning equipment, wear a chemical-resistant apron. **NOTE:** Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 24 hours**. *Exception:* Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no worker contact with the soil subsurface or treated seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; socks and chemical-resistant footwear; chemical resistant gloves; and protective eyewear.

GENERAL

BELMONT™ 2.7 FS is a systemic fungicide seed treatment specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soilborne diseases, use in combination with other seed treatment fungicides. VITAVAX® and RANCONA™ products are compatible with BELMONT 2.7 FS. Do not use BELMONT 2.7 FS in combination with other seed treatment products unless compatibility and safety to the crop has been verified. Do not mix with any pesticide that has a prohibition against tank mixing. Observe the most restrictive of labeling limitations and precautions of all products used in mixtures.

BELMONT 2.7 FS may be applied on its own, as water-based slurry, or in combination with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: BELMONT 2.7 FS is a systemic fungicide with a specific mode of action. BELMONT 2.7 FS could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Chemtura cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible BELMONT 2.7 FS resistant strains of fungi which may occur.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

USE PRECAUTIONS AND RESTRICTIONS

- Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved colorant before being sold.
- IN CALIFORNIA: ONLY for use in commercial seed-treatment establishments. NOT allowed for "on-farm" use in California.
- IMPORTANT: MAXIMUM USAGE when applying multiple metalaxyl and mefenoxam containing products to the same crop within the same season: DO NOT apply more than the maximum seasonal total for the active ingredient as stated on the label of the lowest seasonal total on that crop.
- Do not use other metalaxyl or mefenoxam products for control of *Pythium* in water-seeded rice.
- Do not apply BELMONT 2.7 FS to spinach which is destined to be grown in the greenhouse.

TREATED SEED LABELING

Seeds treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This package or bag contains seed that has been treated with metalaxyl. Do not use treated seed for food, feed, oil production or any other purpose except planting. Store away from feed and foodstuffs. Do not allow children, pets or livestock to have access to treated seeds."

"Wear a long pants, long-sleeved shirt, shoes and socks, and protective gloves opening bag or handling treated seed."

"Do not use other metalaxyl or mefenoxam products for control of *Pythium* in water-seeded rice."

"Do not apply metalaxyl to spinach which is destined to be grown in the greenhouse."

"Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess seed by burying away from bodies of water." "Do not contaminate bodies of water when disposing of planting equipment wash water."

"Dispose of seed packaging or containers in accordance with local requirements."

SEED TREATMENT DIRECTIONS

Canola*, Mustard seed*, Rapeseed* *Not registered for use in California	For <i>Pythium</i> damping-off control: Apply BELMONT™ 2.7 FS as a seed treatment at 0.25 to 0.5 fl. oz. per 100 lbs. of seed.
Cotton	For <i>Pythium</i> seed rot and damping-off: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 to 1.5 fl. oz. per 100 lbs. of seed.
Soybeans	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 to 1.5 fl. oz. per 100 lbs. of seed.
Peanuts* *Not registered for use in California	For <i>Pythium</i> seed rot and damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Sorghum	For control of <i>Pythium</i> damping-off, apply 0.375 to 0.75 fl. oz. of BELMONT 2.7 FS per 100 lbs. of seed as a seed treatment. For control of systemic downy mildew (races susceptible to chemical control), apply 0.75 to 1.5 fl. oz. BELMONT 2.7 FS per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew, apply 1.5 to 3.0 fl. oz. of BELMONT 2.7 FS per 100 lbs. of seed as a seed treatment. Only high quality seed (at least 90% germination) should be treated with BELMONT 2.7 FS.
Sunflowers	For control of systemic downy mildew: Apply BELMONT 2.7 FS as a seed treatment at the rate of 1.5 to 3.0 fl. oz. per 100 lbs. of seed.
Forage Legumes	Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage). For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 to 1.5 fl. oz. per 100 lbs. of seed.

<p>Grain Crops</p>	<p>Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale: For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For systemic downy mildew control of sweet corn: Apply BELMONT 2.7 FS as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
<p>Rice* *Not registered for use in California</p>	<p>For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 to 1.5 fl. oz. per 100 lbs. of seed. If using BELMONT 2.7 FS treated rice, do not use other metalaxyl or mefenoxam products for control of <i>Pythium</i> in water-seeded rice. Containers or bags containing BELMONT 2.7 FS treated rice must be labeled with this restriction.</p>
<p>Seed and Pod Vegetables</p>	<p>Black-eyed peas, chickpeas (garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils. For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For early season <i>Phytophthora</i> control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply BELMONT 2.7 FS as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
<p>Brassica (Cole) Leafy Vegetables</p>	<p>Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens. For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p>
<p>Cucurbit Vegetables</p>	<p>Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuzza, Chinese okra, and hechima), <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), <i>Cucumis mixtar</i>, <i>Cucumis pepo</i> (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of <i>Citrullus lanatus</i>). For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p>
<p>Leafy Vegetables</p>	<p>Amaranth (leafy, Chinese spinach, tampala, cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finochio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including New Zealand and vine), and Swiss chard. *Not registered for use in California. For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.1 to 0.75 fl. oz. per 100 lbs. of seed. NOTE: Do not apply BELMONT 2.7 FS to spinach seed which is destined to be grown in the greenhouse. Containers or bags containing BELMONT 2.7 FS –treated spinach seed must be labeled with this restriction.</p>
<p>Fruiting Vegetables* *Not registered for use in California</p>	<p>Eggplant, groundcherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato: For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.1 to 0.75 fl. oz. per 100 lbs. of seed.</p>

Onions* *Not registered for use in California	Onion (dry bulb and green): For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables* (except potatoes) *Not registered for use in California	Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanager, turmeric, turnip, yam bean (jicama, manioc pea), yam: For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 fl. oz./100 lbs. of seed. NOTE: DO NOT apply BELMONT 2.7 FS to potatoes.
Turf Grasses	For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply BELMONT 2.7 FS as a seed treated at the rate of 1.5 fl. oz. per 100 lbs of seed.
Forage Grasses	Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage: For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.

SEED TREATMENT DIRECTIONS:

USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

BELMONT 2.7 FS systemic fungicide seed dressing may be applied at the reduced rates shown in the table below and still provide good *Pythium* control when used in combination with broad spectrum fungicides. Apply only in combination at EPA registered rates of broad-spectrum seed treatment fungicides for which compatibility and crop safety have been verified. Do not mix with any pesticide that has a prohibition against tank mixing. BELMONT 2.7 FS may be applied on its own or as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

Corn (field), Wheat, Oats, Barley, Rye	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply BELMONT 2.7 FS as a commercial seed treatment at the rate of 0.10 to 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Peanuts* *Not registered for use in California	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply BELMONT 2.7 FS as a commercial seed treatment at the rate of 0.20 to 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Rice* *Not registered for use in California	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply BELMONT 2.7 FS as a commercial seed treatment at the rate of 0.10 to 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Soybeans, Field beans, Kidney beans, Navy beans, Pinto beans, Lentils, Peas	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply BELMONT 2.7 FS as a commercial seed treatment at the rate of 0.20 to 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Sunflowers	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply BELMONT 2.7 FS as a commercial seed treatment at the rate of 0.10 to 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.

SEED TREATMENT DIRECTIONS - FOR EXPORT USE ONLY:

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation. The treater is responsible for confirming the acceptability of his treated seed in the country to which he is exporting.

Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.75 to 1.50 fl. oz. per 100 lbs. of seed.
Canola	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at the rate of 0.50 to 1.70 fl. oz. per 100 lbs. of seed. (32 to 110 ml per 100 kg of seed)
Corn	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply BELMONT 2.7 FS as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed.
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed.
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed.
Millet	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply BELMONT 2.7 FS as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed.
Onions	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed.
Peas	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply BELMONT 2.7 FS as a seed treatment at 3.5 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed. DO NOT apply BELMONT 2.7 FS to potatoes.
Sorghum	For <i>Pythium</i> damping-off control: Apply BELMONT 2.7 FS as a seed treatment at 0.75 to 1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply BELMONT 2.7 FS as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed.
Sunflowers	For systemic downy mildew control: Apply BELMONT 2.7 FS as a seed treatment at 9.0 fl. oz. per 100 lbs. seed.

TREATED SEED LABELING:

Seed treated in accordance with the Export Use pattern must be labeled as follows:
"FOR EXPORT ONLY - NOT FOR DOMESTIC SALES OR USE".

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal or by cleaning of equipment.

PESTICIDE STORAGE: Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container

Plastic Containers: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: *For containers with capacity equal to or less than 5 gallons or 50 pounds:* Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume - and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

For containers with capacities greater than 5 gallons or 50 pounds: Empty the remaining contents into application equipment or a mix tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

Recycling: Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

Refillable container

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

IMPORTANT: READ BEFORE USE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and, to the extent consistent with applicable law, the buyer assumes the risk of any such use.

11
11

Treatment of highly mechanically damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond control of Chemtura, Chemtura makes no claims of guarantees as to germination of carry-over seed.

BELMONT™, VITAVAX® and RANCONA™ are trademarks of Chemtura Corporation
©Copyright 2010, Chemtura Corporation

version 03-18-2010