

SEVIN 10 SULFUR 50 DUST

This product is formulated for application by ground or air.

RECOMMENDATIONS FOR USE:

ALFALFA and CLOVERS: Use 15 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, JAPANESE BEETLE, ALFALFA CATERPILLAR, ARMYWORMS, CORN EARWORMS and STINK BUGS.

BEANS: Use 20 lbs. per acre to control MEXICAN BEAN BEETLE, LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR, LYGUS and STINK BUGS. Do not feed treated vines to meat or dairy animals.

CORN: Use 20 lbs. per acre to control CORN EARWORM, EUROPEAN CORN BORER, FALL ARMYWORM, SAP BEETLES and JAPANESE BEETLE. Do not apply within 7 days of harvest for forage.

For control of CORN EAR WORM; apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re-apply immediately after rain.

For control of European CORN BORER, consult your state agricultural experiment station for proper timing of treatments.

For control of CORN EARWORMS on sweet corn: Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

COTTON: Use 25 lbs. per acre to control THRIPS, FLEA BEETLE, FLEAHOPPER, LEAFWORM, BOLLWORM, FALL ARMYWORM, LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLAR, STINK BUGS and GRASSHOPPERS. Do not graze treated fields or feed cotton trash to dairy animals or meat animals.

GRAPES: Use 10 to 20 lbs. per acre to control GRAPE LEAF FOLDER, GRAPE LEAFHOPPER, GRAPE BERRY MOTH and JAPANESE BEETLE. No time limitation.

EGGPLANTS, PEPPERS and TOMATOES: Use 20 lbs. per acre to control TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES, LEAFHOPPER, TOMATO HORNWORM, EUROPEAN CORN BORER, STINK BUGS, COLORADO POTATO BEETLE and LYGUS BUGS. No time limitation.

POTATOES: Use 15 to 20 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO POTATO BEETLE. Apply to foliage only.

STRAWBERRIES: Use 20 lbs. per acre to control MEADOW SPITTLEBUG and LEAFROLLER. Do not apply within 1 day of harvest.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) 10%
Sulfur 50%

INERT INGREDIENTS 40%

EPA Reg. No. 2935-308-AA 100%

KEEP OUT OF REACH OF CHILDREN!

CAUTION: Harmful if inhaled or swallowed. Do not breathe dust.

Do not take internally. Avoid breathing dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Carbaryl can be injurious to tender foliage. Do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

Do not reuse empty container. Containers should be disposed by burying with wastes or burning (keep out of smoke).

Note For Physicians: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

© Sevin is a trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

TO ATTACHED LABELS

SEVIN 10 SULFUR 50 DUST

This product is formulated for application by ground or air.

RECOMMENDATIONS FOR USE:

ALFALFA and CLOVER: Use 15 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, JAPANESE BEETLE, ALFALFA CATERPILLAR, ARMYWORMS, CORN EARWORMS and STINK BUGS.

BEANS: Use 20 lbs. per acre to control MEXICAN BEAN BEETLE, LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR, LYGUS and STINK BUGS. Do not feed treated vines to meat or dairy animals.

CORN: Use 20 lbs. per acre to control CORN EARWORM, EUROPEAN CORN BORER, FALL ARMYWORM, SAP BEETLES and JAPANESE BEETLE. Do not apply within 7 days of harvest for forage.

For control of CORN EAR WORM, apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re-apply immediately after rain.

For control of European CORN BORER, consult your state agricultural experiment station for proper timing of treatments.

For control of CORN EARWORMS on sweet corn: Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

COTTON: Use 25 lbs. per acre to control THRIPS, FLEA BEETLE, FLEAHOPPER, LEAFWORM, BOLLWORM, FALL ARMYWORM, LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLAR, STINK BUGS and GRASSHOPPERS. Do not graze treated fields or feed cotton trash to dairy animals or meat animals.

GRAPES: Use 10 to 20 lbs. per acre to control GRAPE LEAF FOLDER, GRAPE LEAFHOPPER, GRAPE BERRY MOTH and JAPANESE BEETLE. No time limitation.

SOGPLANTS, PEPPERS and TOMATOES: Use 20 lbs. per acre to control TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES, LEAFHOPPER, TOMATO HORNWORM, EUROPEAN CORN BORER, STINK BUGS, COLORADO POTATO BEETLE and LYGUS BUGS. No time limitation.

POTATOES: Use 15 to 20 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO POTATO BEETLE. Apply to foliage only.

STRAWBERRIES: Use 20 lbs. per acre to control MEADOW SPITTLEBUG and LEAFROLLER. Do not apply within 1 day of harvest.

F-874 EPA Est. 2935-CA-1

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate)	10%
Sulfur	50%
INERT INGREDIENTS	40%

EPA Reg. No. 2935-308-AA 100%

KEEP OUT OF REACH OF CHILDREN I

CAUTION: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid breathing dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Carbaryl can be injurious to tender foliage. Do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

Do not reuse empty container. Containers should be disposed by burying with wastes or burning (keep out of smoke).

Note For Physicians: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

©Sevin is a trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.