

# Emerging *Corynebacterium diphtheriae* Species Complex Infections, Réunion Island, France, 2015–2020

Thomas Garrigos,<sup>1</sup> Anais Grimal,<sup>1</sup> Edgar Badell, Nicolas Traversier, Sandrine Picot, Anne Lignereux, Mahery Ramiandrisoa, Céline Ben Cimon, Marie-Christine Jaffar-Bandjee, Houssein Gbaguidi-Haore, Julie Toubiana, Sylvain Brisse, Guillaume Miltgen,<sup>2</sup> Olivier Belmonte<sup>2</sup>

Clinical, epidemiologic, and microbiologic analyses revealed emergence of 26 cases of *Corynebacterium diphtheriae* species complex infections on Réunion Island, France, during 2015–2020. Isolates were genetically diverse, indicating circulation and local transmission of several diphtheria sublineages. Clinicians should remain aware of the risk for diphtheria and improve diagnostic methods and patient management.

Diphtheria is a contagious, potentially fatal infection caused by toxin-producing bacteria of the *Corynebacterium diphtheriae* species complex, which includes *C. diphtheriae*, *C. ulcerans*, *C. pseudotuberculosis*, *C. rouxii*, *C. belfantii*, and *C. silvaticum*. Infection is localized principally in the upper respiratory tract, and production of diphtheria toxin (encoded by the *tox* gene) can cause systemic complications. Cutaneous diphtheria

and diphtheria endocarditis can also act as sources of respiratory infections (1–4). Diphtheria surveillance has traditionally focused on respiratory illness caused by toxigenic *C. diphtheriae* but has been expanded in some countries to include all *C. diphtheriae* species complex infections irrespective of species, infection site, or toxigenicity, enabling broader disease monitoring. *C. diphtheriae* spreads via human-to-human contact; *C. ulcerans* and *C. pseudotuberculosis* are transmitted to humans primarily through animal contact.

Diphtheria was once a major cause of infant death, but global incidence has declined over the past century, largely because of mass vaccination. Consequently, diphtheria is now often considered a forgotten disease (5). Nevertheless, diphtheria reemergence has been reported in high-income countries and is closely related to patient travel history. Diphtheria is considered endemic in Madagascar, Comoros, and Mayotte in the southwest Indian Ocean, but few cases have been reported on other islands, including Réunion Island, an overseas department of France, where cases emerged in 2015 (6,7). Vaccination coverage is poorer in Mayotte (45% for 7- to 11-year-old children) than in Réunion Island (96% for children 11 months of age). Recent improvements in laboratory diagnostic capabilities, such as mass spectrometry use, have increased reports of *C. diphtheriae* species complex infections (8). However, knowledge of prevalence and origin of those infections is limited in this region. The aims of this study were to review the clinical, epidemiologic, and microbiologic characteristics of *C. diphtheriae* species complex infections on

Author affiliations: Félix Guyon University Hospital Center, Saint-Denis, Réunion Island, France (T. Garrigos, A. Grimal, N. Traversier, M.-C. Jaffar-Bandjee, G. Miltgen, O. Belmonte); Processus Infectieux en Milieu Insulaire Tropical, University of La Réunion, Inserm, CNRS, IRD, Saint-Denis, Réunion Island, France (T. Garrigos, G. Miltgen); Institut Pasteur University of Paris Cité, Paris, France (E. Badell, J. Toubiana, S. Brisse); Institut Pasteur National Reference Center for Corynebacteria of the Diphtheriae Complex, Paris (E. Badell, J. Toubiana, S. Brisse); University Hospital of La Réunion, Saint Pierre, Réunion Island, France (S. Picot); Hospital Center West Réunion, Saint Paul, Réunion Island, France (A. Lignereux); Cereb Alliance Laboratory, Le Port, Réunion Island, France (M. Ramiandrisoa); Inovie RéuniLAB Laboratory, Sainte-Clotilde (C. Ben Cimon); University Hospital Center and University of Bourgogne Franche-Comté, Besançon, France (H. Gbaguidi-Haore); Necker-Enfants malades Hospital, Paris (J. Toubiana).

DOI: <https://doi.org/10.3201/eid2908.230106>

<sup>1</sup>These first authors contributed equally to this article.

<sup>2</sup>These authors contributed equally to this article.

Réunion Island during 2015–2020 and identify possible links with cases on other islands in the region.

### The Study


We included all cases of *C. diphtheriae* species complex infections reported to the regional health agency and recorded at Réunion Island University Hospital during 2015–2020. We analyzed medical records and extracted age, sex, country of residence, recent travel, contact with animals, socioeconomic status, and diphtheria vaccination status for each case. We performed antimicrobial susceptibility testing; identified co-infecting strains; and determined *tox* gene presence, diphtheria toxin production, and biovar and sequence type (ST). We sent each isolate to the National Reference Center for Corynebacteria of the *diphtheriae* Complex (Institut Pasteur, Paris, France) to confirm species identity through multiplex PCR and biotyping as previously described (8–10). We detected the *tox* gene by using conventional PCR or, since 2019, by using multiplex real-time PCR (10). We assessed toxin production by using a modified Elek test (11). We determined antimicrobial drug susceptibility by using disk diffusion or by determining MICs (E-test; bioMérieux, <https://www.biomerieux.com>), in accordance with CASFM/EUCAST2021 (<https://www.sfm-microbiologie.org/2021/04/23/casfm-avril-2021-v1-0>) recommendations for benzylpenicillin, amoxicillin, cefotaxime, clindamycin, rifampin, and ciprofloxacin. We genotyped each isolate by using multilocus sequence typing (MLST) (12).

A total of 26 cases of *C. diphtheriae* species complex infections were recorded, from which 27 *C. diphtheriae* and 2 *C. ulcerans* isolates were cultured. Most (88.5%) infected patients were male; median age was 60 (interquartile range 32.5–67) years. Fourteen (50%) patients lived on Réunion Island, 3 (11.5%) in

Mayotte, 4 (19.2%) in mainland France, 3 (11.5%) in Comoros, and 2 (7.8%) in Madagascar. Most (84.6%) patients had skin manifestations, and 16 patients were vaccinated (Table 1, <https://wwwnc.cdc.gov/EID/article/29/8/23-0106-T1.htm>; Appendix Figure, <https://wwwnc.cdc.gov/EID/article/29/8/23-0106.pdf>). Of 24 *C. diphtheriae* infections, 8 occurred in patients who had recently traveled to or originated from Madagascar, 4 who traveled to or originated from Mayotte, and 3 who traveled to or originated from Comoros. Since 2018, a total of 9 cases on Réunion Island have been considered locally acquired; all of those patients lived in poor socioeconomic conditions. *C. ulcerans* infections occurred in 2 patients living on Réunion Island who had not traveled recently but had contact with animals (Table 1; Figure). We performed a Spearman rank correlation to compare locally acquired strains isolated during 2015–2018 and 2019–2020; a 75% increase in locally acquired *C. diphtheriae* infections occurred in 2019–2020 ( $\rho = 0.8452$ ;  $p = 0.0341$ ).

Isolates were obtained from cutaneous lesion ( $n = 24$ ), bone ( $n = 4$ ), and respiratory ( $n = 1$ ) samples. Eight of 27 *C. diphtheriae* isolates were toxigenic, yielding positive Elek test results. The 2 *C. ulcerans* isolates were nontoxigenic. *C. diphtheriae* isolates were characterized as biovars Mitis ( $n = 20$ ) and Gravis ( $n = 7$ ).

Patient isolates were co-infected most frequently with *Staphylococcus aureus* ( $n = 17$ ) and *Streptococcus pyogenes* ( $n = 18$ ). Benzylpenicillin resistance was observed in 80% of isolates according to CASFM/EUCAST2021 recommendations, but isolates were categorized as susceptible increased exposure according to EUCAST version 13.0 proposed breakpoints ([https://www.eucast.org/clinical\\_breakpoints](https://www.eucast.org/clinical_breakpoints)) (Appendix Table). One (3.5%) *C. diphtheriae* isolate was resistant to amoxicillin (CD8/FRC0402; MIC 1.5 mg/L),



**Figure.** Number of cases diagnosed per year in study of emerging *Corynebacterium diphtheriae* species complex infections, Réunion Island, France, 2015–2020. Number of cases were classified according to geographic origin (A) or travel history of patients (B). Dotted lines indicate linear trends.

and 1 was resistant to rifampin. Both *C. ulcerans* isolates were resistant to clindamycin (100%, natural low susceptibility), whereas clindamycin resistance was observed for only 1 *C. diphtheriae* isolate.

We identified 21 STs by MLST analysis, including ST88 for *C. diphtheriae* isolates from 4 patients and ST339 for both *C. ulcerans* isolates (Table 2). All *C. diphtheriae* STs had 2–5 mismatches, except ST87 and ST237, which had 1 mismatch between them. ST339 (*C. ulcerans*) had 7 mismatches with all *C. diphtheriae* STs.

## Conclusions

We report increased prevalence of cutaneous *C. diphtheriae* species complex infections on Réunion Island during 2015–2020. Introduction of mass spectrometry analysis in hospital laboratories and increased clinician awareness might have led to increased case reporting. Our study confirms that *C. diphtheriae* species complex members are circulating and are likely underestimated in the southwest Indian Ocean (7,13). Moreover, we observed emergence of locally acquired cutaneous *C. diphtheriae*

infections on Réunion Island since 2019. The number of imported cases in 2020 was probably limited because of the COVID-19 pandemic, which reduced travel. Indeed, all *C. diphtheriae* cases identified during 2015–2018 occurred in patients who had traveled from other islands in the Indian Ocean. In addition, cutaneous diphtheria appeared to be associated with poor socioeconomic living conditions, in which alcoholism, drug dependence, and homelessness are factors that increase risk for human-to-human transmission and virulence (14).

A total of 8 (30%) *C. diphtheriae* isolates were toxigenic and caused cutaneous infections. Nontoxigenic *C. diphtheriae* isolates (70%, n = 19) were obtained from cutaneous lesions, respiratory samples, and bone samples. Clinicians should be aware that nontoxigenic *C. diphtheriae* can potentially cause severe disease (1,14,15). Moreover, all isolates were co-infected with pyogenic bacteria, suggesting diphtheria infection should be considered under polymicrobial conditions.

MLST analysis identified 21 different STs; most were unrelated ( $\geq 2$  mismatches) reflecting marked

**Table 2.** Characteristics of isolates from 26 patients in study of emerging *Corynebacterium diphtheriae* species complex infections, Réunion Island, France, 2015–2020\*

Patient no.	Isolate	Year	Isolation site	Species	Biovar	tox gene	Elek test	ST†	Co-infections‡
1	CD1/FRC0304	2015	Cutaneous	<i>C. diphtheriae</i>	Gravis	Negative	NA	102	<i>S. pyogenes</i> , <i>S. aureus</i> , <i>A. haemolyticum</i>
	CD2/FRC0316	2015	Respiratory	<i>C. diphtheriae</i>	Mitis	Negative	NA	95	<i>S. aureus</i>
2	CD3/FRC0314	2015	Cutaneous	<i>C. diphtheriae</i>	Mitis	Positive	Positive	421	<i>S. aureus</i>
3	CD4/FRC0376	2015	Cutaneous	<i>C. diphtheriae</i>	Gravis	Positive	Positive	388	<i>S. pyogenes</i>
4	CD5/FRC0383	2016	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	423	<i>S. pyogenes</i> , <i>S. aureus</i>
	CD6/FRC0393	2016	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	423	<i>S. pyogenes</i> , <i>S. aureus</i>
5	CD7/FRC0385	2016	Cutaneous	<i>C. diphtheriae</i>	Mitis	Positive	Positive	91	<i>S. pyogenes</i>
6	CU1/FRC0391	2016	Cutaneous	<i>C. ulcerans</i>	NA	Negative	NA	<b>339</b>	<i>S. dysgalactiae</i>
7	CD8/FRC0402	2016	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	410	<i>S. dysgalactiae</i>
8	CD9/FRC0410	2016	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	415	<i>S. pyogenes</i>
9	CD10/FRC0423	2016	Cutaneous	<i>C. diphtheriae</i>	Gravis	Negative	NA	101	<i>S. aureus</i>
10	CD11/FRC0477	2017	Cutaneous	<i>C. diphtheriae</i>	Gravis	Negative	NA	481	<i>S. pyogenes</i> , <i>S. aureus</i> , <i>A. haemolyticum</i>
11	CD12/FRC0501	2017	Cutaneous	<i>C. diphtheriae</i>	Gravis	Positive	Positive	521	<i>S. pyogenes</i>
12	CD13/FRC0624	2018	Bone	<i>C. diphtheriae</i>	Mitis	Negative	NA	237	<i>S. aureus</i>
13	CD14/FRC0630	2018	Cutaneous	<i>C. diphtheriae</i>	Gravis	Negative	NA	606	<i>S. pyogenes</i> , <i>S. aureus</i>
14	CD15/FRC0733	2019	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	351	<i>S. pyogenes</i>
15	CD16/FRC0782	2019	Cutaneous	<i>C. diphtheriae</i>	Mitis	Positive	Positive	688	<i>S. pyogenes</i> , <i>S. aureus</i>
	CD17/FRC0809	2019	Cutaneous	<i>C. diphtheriae</i>	Mitis	Positive	Positive	688	<i>S. aureus</i>
16	CD18/FRC0819	2019	Cutaneous	<i>C. diphtheriae</i>	Gravis	Positive	Positive	87	<i>S. pyogenes</i> , <i>A. haemolyticum</i>
17	CU2/FRC0820	2019	Bone	<i>C. ulcerans</i>	NA	Negative	NA	<b>339</b>	<i>S. aureus</i>
18	CD19/FRC0849	2019	Cutaneous	<i>C. diphtheriae</i>	Mitis	Positive	Positive	426	<i>S. pyogenes</i> , <i>S. aureus</i>
19	CD20/FRC0865	2020	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	102	<i>S. pyogenes</i> , <i>S. aureus</i>
20	CD21/FRC0875	2020	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	707	<i>S. pyogenes</i>
21	CD22/FRC0893	2020	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	708	<i>S. pyogenes</i>
22	CD23/FRC0928	2020	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	<b>88</b>	<i>S. pyogenes</i>
23	CD24/FRC0970	2020	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	<b>88</b>	<i>S. pyogenes</i> , <i>S. aureus</i> , <i>A. haemolyticum</i>
24	CD25/FRC0975	2020	Cutaneous	<i>C. diphtheriae</i>	Mitis	Negative	NA	<b>88</b>	<i>S. aureus</i>
25	CD26/FRC1050	2020	Bone	<i>C. diphtheriae</i>	Mitis	Negative	NA	771	<i>A. haemolyticum</i>
26	CD27/FRC1065	2020	Bone	<i>C. diphtheriae</i>	Mitis	Negative	NA	<b>88</b>	<i>S. aureus</i>

\*CD, *Corynebacterium diphtheriae*; CU, *C. ulcerans*; NA, not applicable; ST, sequence type.

†Numbers in bold indicate a common ST shared among strains from different patients.

‡Co-infections with *Arcanobacterium haemolyticum*, *Staphylococcus aureus*, *Streptococcus dysgalactiae*, or *Streptococcus pyogenes*.

genetic diversity of isolates. ST88 was found in 4 patients living on Réunion Island who had not traveled recently, indicating probable local acquisition. ST88 had previously been reported only in patients from Mayotte. Therefore, our results show that multiple *C. diphtheriae* species complex clones are circulating in the southwest Indian Ocean (8). Both *C. ulcerans* strains belonged to ST339. The National Reference Center reported that ST339 is the predominant *C. ulcerans* ST found in animals in France. Although considerable ST diversity was revealed, whole-genome sequencing will be required to further evaluate circulating *C. diphtheriae* clones in this region.

In conclusion, we describe emergence of locally acquired *C. diphtheriae* species complex infections on Réunion Island during 2019–2020. Local clinicians and microbiologists should remain aware of this neglected infection; improvements should be made in diagnostic methods and management of infected patients, such as maintaining availability of diphtheria antitoxin.

The National Reference Center for Corynebacteria of the *diphtheriae* Complex is supported financially by Santé publique France (Saint-Maurice, France).

## About the Author

Dr. Garrigos is a research scientist in the microbiology department of Félix Guyon University Hospital of Réunion Island, France. His research interests focus on bacterial diseases, antimicrobial resistance, cystic fibrosis patients, and emerging infectious diseases.

## References

1. Patey O, Bimet F, Riegel P, Halioua B, Emond JP, Estrangin E, et al.; Coryne Study Group. Clinical and molecular study of *Corynebacterium diphtheriae* systemic infections in France. *J Clin Microbiol*. 1997;35:441–5. <https://doi.org/10.1128/jcm.35.2.441-445.1997>
2. Hadfield TL, McEvoy P, Polotsky Y, Tzinslering VA, Yakovlev AA. The pathology of diphtheria. *J Infect Dis*. 2000;181:S116–20. <https://doi.org/10.1086/315551>
3. Levi LL, Barbut F, Chopin D, Rondeau P, Lalande V, Jolivet S, et al. Cutaneous diphtheria: three case reports to discuss determinants of re-emergence in resource-rich settings. *Emerg Microbes Infect*. 2021;10:2300–2. <https://doi.org/10.1080/22221751.2021.2008774>
4. Sangal V, Hoskisson PA. Evolution, epidemiology and diversity of *Corynebacterium diphtheriae*: new perspectives on an old foe. *Infect Genet Evol*. 2016;43:364–70. <https://doi.org/10.1016/j.meegid.2016.06.024>
5. Sharma NC, Efstratiou A, Mokrousov I, Mutreja A, Das B, Ramamurthy T. Diphtheria. *Nat Rev Dis Primers*. 2019;5:81. <https://doi.org/10.1038/s41572-019-0131-y>
6. Scheifer C, Rolland-Debord C, Badell E, Reibel F, Aubry A, Perignon A, et al. Re-emergence of *Corynebacterium diphtheriae*. *Med Mal Infect*. 2019;49:463–6. <https://doi.org/10.1016/j.medmal.2018.12.001>
7. Belchior E, Henry S, Badell E, Collet L, Benoit-Cattin T, de Montera AM, et al. Diphtheria in Mayotte, 2007–2015. *Emerg Infect Dis*. 2017;23:1218–20. <https://doi.org/10.3201/eid2307.170262>
8. Hennart M, Panunzi LG, Rodrigues C, Gaday Q, Baines SL, Barros-Pinkelng M, et al. Population genomics and antimicrobial resistance in *Corynebacterium diphtheriae*. *Genome Med*. 2020;12:107. <https://doi.org/10.1186/s13073-020-00805-7>
9. Dazas M, Badell E, Carmi-Leroy A, Criscuolo A, Brisse S. Taxonomic status of *Corynebacterium diphtheriae* biovar Belfanti and proposal of *Corynebacterium belfantii* sp. nov. *Int J Syst Evol Microbiol*. 2018;68:3826–31. <https://doi.org/10.1099/ijsem.0.003069>
10. Badell E, Guillot S, Tulliez M, Pascal M, Panunzi LG, Rose S, et al. Improved quadruplex real-time PCR assay for the diagnosis of diphtheria. *J Med Microbiol*. 2019;68:1455–65. <https://doi.org/10.1099/jmm.0.001070>
11. Engler KH, Glushkevich T, Mazurova IK, George RC, Efstratiou A. A modified Elek test for detection of toxigenic corynebacteria in the diagnostic laboratory. *J Clin Microbiol*. 1997;35:495–8. <https://doi.org/10.1128/jcm.35.2.495-498.1997>
12. Bolt F, Cassiday P, Tondella ML, Dezoysa A, Efstratiou A, Sing A, et al. Multilocus sequence typing identifies evidence for recombination and two distinct lineages of *Corynebacterium diphtheriae*. *J Clin Microbiol*. 2010;48:4177–85. <https://doi.org/10.1128/JCM.00274-10>
13. Rakotomalala RS, Andrianirina ZZ, Ratsima E, Randrianandraina P, Randrianirina F, Edosoa GT, et al. *Corynebacterium diphtheriae* infection in Mahajanga, Madagascar: first case report. *J Trop Pediatr*. 2021; 67:fmaa064. <https://doi.org/10.1093/tropej/fmaa064>
14. Badenschier F, Berger A, Dangel A, Sprenger A, Hobmaier B, Sievers C, et al. Outbreak of imported diphtheria with *Corynebacterium diphtheriae* among migrants arriving in Germany, 2022. *Euro Surveill*. 2022;27:2200849. <https://doi.org/10.2807/1560-7917.ES.2022.27.46.2200849>
15. Farfour E, Badell E, Zasada A, Hotzel H, Tomaso H, Guillot S, et al. Characterization and comparison of invasive *Corynebacterium diphtheriae* isolates from France and Poland. *J Clin Microbiol*. 2012;50:173–5. <https://doi.org/10.1128/JCM.05811-11>

Address for correspondence: Thomas Garrigos, Laboratoire de Microbiologie, Hôpital Universitaire Félix Guyon, Allée des Topazes, 97400 Saint-Denis, La Réunion, France; email: thomas.garrigos@chu-reunion.fr

Article DOI: <https://doi.org/10.3201/eid2908.230106>

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

# Emerging *Corynebacterium diphtheriae* Species Complex Infections, Réunion Island, France, 2015–2020

## Appendix

**Appendix Table.** Sensitivity of different patient isolates to antimicrobial drugs\*

Isolates	Benzylpenicillin	Amoxicillin	Clindamycin	Rifampin	Ciprofloxacin
CD1/FRC0304	SIE	S	S	S	S
CD2/FRC0316	SIE	S	S	S	S
CD3/FRC0314	SIE	S	S	S	S
CD4/FRC0376	SIE	S	S	S	S
CD5/FRC0383	SIE	S	S	S	S
CD6/FRC0393	SIE	S	S	S	S
CD7/FRC0385	SIE	S	S	S	S
CU1/FRC0391	SIE	S	R	S	S
CD8/FRC0402	SIE	R	R	S	S
CD9/FRC0410	SIE	S	S	S	S
CD10/FRC0423	SIE	S	S	S	S
CD11/FRC0477	SIE	S	S	S	S
CD12/FRC0501	SIE	S	S	S	S
CD13/FRC0624	SIE	S	S	S	S
CD14/FRC0630	SIE	S	S	R	S
CD15/FRC0733	SIE	S	S	S	S
CD16/FRC0782	SIE	S	S	S	S
CD17/FRC0809	SIE	S	S	S	S
CD18/FRC0819	SIE	S	S	S	S
CU2/FRC0820	SIE	S	R	S	S
CD19/FRC0849	SIE	S	S	S	S
CD20/FRC0865	SIE	S	S	S	S
CD21/FRC0875	SIE	S	S	S	R
CD22/FRC0893	SIE	S	S	S	S
CD23/FRC0928	SIE	S	S	S	S
CD24/FRC0970	SIE	S	S	S	S
CD25/FRC0975	SIE	S	S	S	S
CD26/FRC1050	SIE	S	S	S	S
CD27/FRC1065	SIE	S	S	S	S

\*CD, *Corynebacterium diphtheriae*; CU, *C. ulcerans*; R, resistant; S, sensitive; SIE, susceptible increased exposure.



**Appendix Figure.** Image of right ankle cutaneous ulcer from patient with *Corynebacterium diphtheriae* species complex infection, Réunion Island, France, 2015–2020.