

The Fauna of Parasitic Barnacles (Cirripedia: Rhizocephala) in Korea

Jibom Jung^{1,2} and Joong-Ki Park^{1,*}

¹Division of EcoScience, Ewha Womans University, Seoul 03760, South Korea. *Correspondence: E-mail: jkpark@ewha.ac.kr (Park)

²School of Biological Sciences, Seoul National University, Seoul 08826, South Korea. E-mail: apociv@gmail.com (Jung)

Received 31 August 2023 / Accepted 21 December 2023 / Published 15 March 2024
Communicated by Benny K.K. Chan

Rhizocephala is a group of endoparasitic barnacles, the morphological characteristics of which are degenerated, and which has recently undergone active molecular identification. Despite several recent studies of Korean rhizocephalans, a comprehensive analysis of rhizocephalan fauna has not yet been conducted. In this study, we analyzed morphological and molecular characteristics of 64 rhizocephalan samples from 15 decapod hosts sampled across the Korean coast. We found 16 Rhizocephala species of six genera from four Rhizocephala families, *i.e.*, Peltogasterellidae, Peltogastridae, Polyascidae, and Sacculinidae. Combining morphological examination and molecular analysis of mitochondrial cytochrome *c* oxidase I revealed three new species candidates, *i.e.*, *Peltogasterella* sp., *Peltogaster* sp., and *Parasacculina* sp. 1, and three rhizocephalans that have expanded their distribution range to the Korean coast, *i.e.*, *Parasacculina oblonga*, *Sacculina angulata*, and *Sacculina gracilis*, whose expanded their distribution range to Korean coast. A synthetic update of Korean Rhizocephala fauna including a species checklist and remarks regarding distribution and taxonomy is also presented.

Key words: Rhizocephala, Cirripedia, Parasitic barnacles, Decapod hosts, Distribution pattern, Korean coast

BACKGROUND

Rhizocephala is a group of parasitic barnacles that infect both the internal (endoparasitic) and external body (ectoparasitic) of crustacean hosts, mainly decapods. The organism has a complicated life cycles, beginning with a planktonic larval stage, followed by an internal parasitic stage, and concluding with an external reproductive stage (Høeg 1992). In contrast to other crustacean species, they possess an invisible and branch-shaped internal parasitic organ (interna) and a visible and simplified external reproductive organ (externa) during the parasitic stage, lacking both segmentation and appendages (Høeg and Lützen 1995). Other organ systems such as the respiratory, digestive, sensory, and excretory systems are degenerate in adult rhizocephalans (Høeg 1992; Øksnebjerg 2000). As a result of this simplified external adult morphology, the previous

taxonomy system of rhizocephalans relied on the examination of morphological characteristics of larva and histological characteristics of adult externa (Yoshida et al. 2011; Kobayashi et al. 2018), and this system was further verified and partially modified through DNA barcode and molecular phylogenetic analysis (Yoshida et al. 2012 2014; Høeg et al. 2019; Chan et al. 2021).

Since the first report of parasitic barnacles *Parasacculina pinnotherae* in Korea (Shiino 1943), several systematic studies have been conducted on Korean parasitic barnacles (Lützen et al. 2016; Jung et al. 2019 2021). However, the rhizocephalan fauna in Korea has not yet been fully identified, as only a limited number of hosts have been investigated (Jung et al. 2019 2021). For example, *Parasacculina oblonga* (Lützen and Yamaguchi 1999) was first reported from the host crab *Cyclograpsus intermedius* in Japan, but parasitism of *P. oblonga* was not reported in Korean *C. intermedius*.

Therefore, we also collected Korean rhizocephalans and identified them based on morphological examination and molecular analyses of mitochondrial cytochrome *c* oxidase subunit I (*coxI*). In addition, we made a checklist included detailed biological characteristics about each Korean rhizocephalan to help future research about Rhizocephala.

MATERIALS AND METHODS

We examined 64 Korean rhizocephalans from 15 host decapods collected from 21 sampling sites in Korea (Table 1). Voucher specimens in this study were deposited in the National Institute of Biological Resources (NIBR) and the Honam National Institute of Biological Resources (HNIBR). All rhizocephalan specimens were fixed in 70–95% ethanol and subjected to morphological examination and molecular analysis. For morphological analysis, the externa and mantle aperture were examined using an MZ8 dissection microscope (Leica, Wetzlar, Germany). Photographs were taken with a D200 digital camera (Nikon, Tokyo, Japan). The shield length (hermit crab) and carapace length (crab) of the host decapod were measured as the length from the tip of the rostrum to the midpoint of the posterior margin of the carapace using a CD6CSX digital caliper (Mitutoyo, Kawasaki, Japan) to the nearest 1 mm.

For molecular analysis, the lateral end of the externa tissue of each rhizocephalan specimen was excised for total genomic DNA extraction using the QIAamp DNA Micro Kit (QIAGEN, Hilden, Germany). Universal primers LCO1490 (5'-GGTCAACAAAT CATAAAGATATTGG-3') and HCO2198 (5'-TAAA CTTCAGGGTGACCAAAAAATCA-3') were used to amplify a fragment of *coxI* (Folmer et al. 1994). Polymerase chain reaction (PCR) was performed in reaction volumes of 50 μ L that included 2 μ L DNA template, 5 μ L 10 x Ex Taq Buffer, 2 μ L of each primer (10 μ M), 0.25 μ L Go Taq DNA polymerase (Promega, Madison City, WI, USA), 5 μ L dNTP mix (10 mM), and 35.75 μ L distilled H₂O. PCR amplification was performed using the following steps: 5 min denaturation at 94°C followed by 35 cycles of 30 sec at 94°C, 1 min at 52°C, 1 min at 72°C, and a final extension of 7 min at 72°C. PCR products were visualized on 1% agarose gels and sequenced with an ABI PRISM 3730xl DNA analyzer (Applied Biosystems, Foster City, CA, USA). Nucleotide sequences of mtDNA *coxI* fragments were analyzed and edited using Geneious v. 9.1.8 (Kearse et al. 2012). *coxI* sequences in this study were deposited in GenBank (Table 1). Molecular identification of each sample was conducted through BLAST results of the

sequences.

To determine the fauna of Korean rhizocephalans, the results of morphological and molecular identification of the 64 individuals were combined with taxonomic information from previous studies of Korean Rhizocephala.

RESULTS

Based on the morphological examination of externae from 64 samples and information from 46 *coxI* sequences, we identified 16 rhizocephalan species belonging to six genera, and four families *i.e.*, Peltogasterellidae, Peltogastridae, Polyascidae, and Sacculinidae (Table 1). This study found that three species of rhizocephalans *i.e.*, *Parasacculina oblonga*, *Sacculina angulata*, and *Sacculina gracilis*, had an expanded distribution range and were newly found in Korea. In addition, we found three new species candidates, *i.e.*, *Peltogasterella* sp., *Peltogaster* sp., and *Parasacculina* sp. 1. As a result of including rhizocephalans reported in previous studies (Shiino 1943; Kim and Hong 2002; Lützen et al. 2016; Jung et al. 2019 2021), the presence of 22 rhizocephalans in Korea were identified. The distribution of Rhizocephala in South Korea is shown in figure 1, and a checklist was prepared for 16 species with confirmed species names and molecular data. Brief information on the externa morphology of 22 Korean rhizocephalans is presented in table 2. We provide brief remarks about noteworthy biological characteristics of Korean Rhizocephala, including comparisons to previous studies. Diagnoses and distribution information identical to previous studies (Jung et al. 2019 2021) are omitted.

Checklist of Korean Rhizocephala

- Phylum Arthropoda von Siebold, 1848
- Class Thecostraca Gruvel, 1905
- Subclass Cirripedia Burmeister, 1834
- Infraclass Rhizocephala Müller, 1862
- Family Peltogasterellidae Høeg & Glenner in Høeg, Noever, Rees, Crandall & Glenner, 2019
- Genus *Peltogasterella* Krüger, 1912
 1. *Peltogasterella gracilis* (Boschma, 1927)
- Family Peltogastridae Lilljeborg, 1861
- Genus *Peltogaster* Rathke, 1842
 2. *Peltogaster lineata* Shiino, 1943
 3. *Peltogaster postica* Yoshida & Osawa in Yoshida, Osawa, Hirose & Hirose, 2011
 4. *Peltogaster reticulata* Shiino, 1943
- Family Polyascidae Høeg & Glenner in Høeg, Noever,




Fig. 1. Distribution map of the eight most common Korean rhizocephalans. Blue dotted lines indicate divisions between ecoregions mentioned in Muhammad et al. (2022) and Kim et al. (2020 2022).

Table 1. Geographical locations and sample information of Korean rhizocephalans in this study in order of species name, host species, location, and specimen number

Species	Host Species	Location (Korea)	Specimen Number	coxI Accession Number	
<i>Heterosaccus papillosus</i>	<i>Charybdis (Charybdis) japonica</i>	Incheon	NIBRIV0000895329	OR481955	
			NIBRIV0000895330		
<i>Parasacculina imberbis</i>	<i>Pachygrapsus crassipes</i>	Jeju	Koreakr10	OR481956	
			Seogwipo	Koreax8	OR481963
			Tongyeong	Korearr54	OR481962
<i>Parasacculina oblonga</i>	<i>Cyclograpsus intermedius</i>	Ulleung	DSEVIV0000000050	OR481965	
			DSEVIV0000000051	OR481966	
			DSEVIV0000000054	OR481967	
			DSEVIV0000000055	OR481968	
			DSEVIV0000000056	OR481969	

Table 1. (Continued)

Species	Host Species	Location (Korea)	Specimen Number	<i>coxI</i> Accession Number
<i>Parasacculina pinnotherae</i>	<i>Arcotheres sinensis</i>	Jindo	KoreaJD1	
<i>Parasacculina pilosella</i>	<i>Pugettia intermedia</i>	Goseong	GS 2381	
		Sokcho	Korearr1	
<i>Parasacculina yatsui</i>	<i>Cyclograpsus intermedius</i>	Ulleung	Korearr40	
	<i>Hemigrapsus penicillatus</i>	Gangneung	Korearr10	
		Goseong	GS12	
	<i>Hemigrapsus sanguineus</i>	Goheung	KoreaN14	OR481973
		Goseong	Korea71	OR481975
		Tongyeong	Korea39	OR481974
	<i>Pachygrapsus crassipes</i>	Gangneung	Korearr9	OR481977
		Jeju	Koreakr9	OR481976
		Samcheok	Korearr25	OR481978
		Yeongdeok	Korea72	
<i>Parasacculina shiinoi</i>	<i>Upogebia major</i>	Namhae	Korea56	
<i>Parasacculina</i> sp. 1	<i>Hemigrapsus sanguineus</i>	Goheung	Korea6N	OR481970
			Korea13N	OR481971
			Korea16N	OR481972
<i>Peltogasterella gracilis</i>	<i>Pagurus maculosus</i>	Shinan	Korea61	OR481992
<i>Peltogasterella</i> sp.	<i>Porcellanopagurus nihonkaiensis</i>	Jeju	MADBK 160730_002	
		Seogwipo	KoreaPN1	OR481994
<i>Peltogaster reticulata</i>	<i>Pagurus minutus</i>	Namhae	MADBK 160706_065	MK604151*
			MADBK 160706_125	
	<i>Pagurus proximus</i>	Shinan	MADBK 160718_040	
<i>Peltogaster</i> sp. 1	<i>Pagurus lanuginosus</i>	Gangneung	DSEVIV0000000053	OR481985
	<i>Pagurus maculosus</i>	Seogwipo	DSEVIV0000000057	OR481986
		Shinan	DSEVIV0000000052	OR481987
			KoreaHS1	OR481988
			KoreaHS2	OR481989
			KoreaHS3	OR481990
			KoreaHS4	OR481991
<i>Polyascus</i> cf. <i>gregarius</i>	<i>Hemigrapsus sanguineus</i>	Busan	Korearr60	OR482023
		Goheung	Korea1N	OR481995
		Jindo	Koreakr1	OR482016
		Shinan	Korearr58	OR482022
		Tongyeong	Korearr56	OR482021
		Wando	Korea47	OR482011
		Yeosu	Korea45	OR482010
<i>Sacculina angulata</i>	<i>Thalamita sima</i>	Busan	NIBRIV0000895328	OR482032
		Sacheon	NIBRIV0000895331	
<i>Sacculina confragosa</i>	<i>Gaetice depressus</i>	Goheung	Korea52	OR482037
	<i>Hemigrapsus sanguineus</i>	Gangneung	Korearr22	OR482050
		Goheung	Korea7N	OR482033
		Jindo	Koreakr2	OR482038
		Pohang	Korearr59	OR482062
	<i>Pachygrapsus crassipes</i>	Gangneung	Korearr8	OR482049
		Jeju	Korea74	
		Samcheok	Korearr53	OR482061
		Seogwipo	Koreakr11	OR482041
		Sokcho	Korea73	
		Uljin	Korearr28	OR482053
<i>Sacculina gracilis</i>	<i>Thalamita sima</i>	Seogwipo	NIBRIV0000895332	OR482063
		Tongyeong	NIBRIV0000895334	OR482064
<i>Sacculina upogebiae</i>	<i>Upogebia major</i>	Boryeong	NIBRIV0000453552	
		Incheon	NIBRIV0000453567	

*: NCBI sequence.

Rees, Crandall & Glenner, 2019
 Genus *Parasacculina* Høeg & Glenner in Høeg, Noever, Rees, Crandall & Glenner, 2019
 5. *Parasacculina imberbis* (Shiino, 1943)
 6. *Parasacculina pilosella* (Van Kampen & Boschma, 1925)
 7. *Parasacculina pinnotherae* (Shiino, 1943)
 8. *Parasacculina shiinoi* (Lützen, Itani, Jespersen, Hong, Rees & Glenner, 2016)
 9. *Parasacculina yatsui* (Boschma, 1936)
 10. *Parasacculina oblonga* (Lützen & Yamaguchi, 1999)
 Genus *Polyascus* Glenner, Lützen & Takahashi, 2003

11. *Polyascus* cf. *gregarius* (Okada & Miyashita, 1935)

Family Sacculinidae Lilljeborg, 1861
 Genus *Heterosaccus* Smith, 1906
 12. *Heterosaccus papillosus* (Boschma, 1933)
 Genus *Sacculina* Thompson, 1836
 13. *Sacculina upogebiae* Shiino, 1943
 14. *Sacculina confragosa* Boschma, 1933
 15. *Sacculina angulata* Van Kampen & Boschma, 1925
 16. *Sacculina gracilis* Boschma, 1931

Table 2. Morphological characteristics of externa of 22 Korean Rhizocephala

Species	Externa number	Externa shape	Mantle aperture			Chitinous shield	Host family
			Projection	Opening	Position		
<i>Peltogasterella gracilis</i>	Gregarious	Elongate banana-shaped	Elevated	Circular	Terminal	No	Paguridae
<i>Peltogasterella</i> sp.	1–3	Elongate	Vestigial	Circular	Terminal	No	Paguridae
<i>Peltogaster lineata</i>	1	Elongate oval	Slightly elevated or elevated tube	Circular	Terminal or subterminal	Yes	Paguridae
<i>Peltogaster postica</i>	1–2	Irregularly elongate	Elevated tube	U-shaped	Terminal	Yes	Paguridae
<i>Peltogaster reticulata</i>	1	Banana-shaped	Elevated tube	X-shaped	Terminal	Yes	Paguridae
<i>Peltogaster</i> sp. 1	1	Irregularly elongate	Slightly elevated	Circular	Terminal or subterminal	Yes	Paguridae
<i>Peltogaster</i> sp. 2	1	Irregularly elongate			Terminal or subterminal	Yes	Diogenidae
<i>Parasacculina imberbis</i>	1	Smooth round-rectangular or bilobed	Slightly elevated	X-shaped	Terminal	No	Grapsidae
<i>Parasacculina pilosella</i>	1	Smooth and slightly flat oval	Vestigial	Circular	Terminal	No	Epialtidae
<i>Parasacculina pinnotherae</i>	1–2	Smooth or slightly wrinkled flat oval or cordiform	Elevated	Dot shaped	Terminal	No	Pinnotheridae
<i>Parasacculina shiinoi</i>	1	Smooth oval	Vestigial	Circular	Terminal	No	Upogebiidae
<i>Parasacculina yatsui</i>	1	Slightly wrinkled flat oval or flat cordiform	Elevated	Slit shaped	Terminal	No	Grapsidae, Varunidae
<i>Parasacculina oblonga</i>	1	Smooth or slightly wrinkled oval	Elevated	Y-shaped	Terminal	No	Varunidae
<i>Parasacculina</i> sp. 1	1	Smooth or slightly wrinkled flat cordiform	Slightly elevated	Circular	Terminal	No	Varunidae
<i>Parasacculina</i> sp. 2	1	Smooth or slightly wrinkled oval	Elevated	Circular	Terminal	No	Xanthidae
<i>Parasacculina</i> sp. 3	1	Smooth oval	Slightly elevated	Circular	Terminal	No	Xanthidae
<i>Polyascus</i> cf. <i>gregarius</i>	1–3	Smooth or slightly wrinkled flat cordiform	Elevated	Slit shaped	Terminal	No	Varunidae
<i>Heterosaccus papillosus</i>	1	Smooth or slightly wrinkled reversed cordiform	Slightly elevated or elevated	Large circular	Terminal	No	Portunidae
<i>Sacculina upogebiae</i>	1	Oval with a very shallow groove along the dorsal margin	Vestigial	Circular	Terminal	No	Upogebiidae
<i>Sacculina confragosa</i>	1–2	Wrinkled flat cordiform	Elevated tube	Circular	Terminal	No	Grapsidae, Varunidae
<i>Sacculina angulata</i>	1–3	Slightly wrinkled irregular pentagon	Elevated tube	Large circular	Terminal	No	Portunidae
<i>Sacculina gracilis</i>	1	Smooth round fusiform	Slightly elevated	Y-shaped	Terminal	No	Portunidae

**Peltogasterellidae Høeg & Glenner in Høeg,
Noever, Rees, Crandall & Glenner, 2019**

***Peltogasterella* Krüger, 1912**

***Peltogasterella gracilis* (Boschma, 1927)**

(Fig. 2A, see Jung et al. 2019)

Additional material examined: On *Pagurus maculosus*: 1 individual, Sinan, 34°40'55.1"N, 125°26'38.1"E, 26 Aug. 2021, collected by Jung J., Korea61, host: males, shield length 8 mm.

Host species: *Discorsopagurus schmitti*, *Labidochirus splendescens*, *Pagurus aleuticus*, *Pagurus dalli*, *Pagurus edwardsii*, *Pagurus filholi*, *Pagurus hemphilli*, *Pagurus hirsutiusculus*, *Pagurus lanuginosus*, *P. maculosus*, *Pagurus middendorffii*, *Pagurus nigrivittatus*, *Pagurus ochotensis*, *Pagurus pectinatus*, *Pagurus spina* (Anomura).

***Peltogasterella* sp.**

(Fig. 2B)

Material examined: on *Porcellanopagurus nihonkaiensis*: 2 individuals (single individual with 2 externae), Jeju, 33°25'17.44"N, 126°9'38.96"E, Scuba, 24 Sep. 2011, collected by Lee SK., MADBK 160730_002, host: males, shield length 2 mm; 1 individual (3 externae), Seogwipo, 33°13'30.2"N, 126°34'11.9"E, Scuba 60 m, 11 Nov. 2020, collected by Lee SH., KoreaPN1, host: shield length 1 mm.

Host species: *P. nihonkaiensis* (Anomura).

Distribution: Korea.

Diagnosis: Whole externa single to triple, smooth, and elongated oval in shape. Mantle aperture not well developed, and vertically small circular opening at the extremity.

Peltogastridae Lilljeborg, 1861

***Peltogaster* Rathke, 1842**

***Peltogaster lineata* Shiino, 1943**

(see Jung et al. 2019)

Host species: *Pagurus brachiomastus*, *Pagurus filholi*, *Pagurus nigrivittatus*, *Pagurus maculosus* (Anomura; Paguridae).

***Peltogaster postica* Yoshida & Osawa in
Yoshida, Osawa, Hirose & Hirose, 2011**

(see Jung et al. 2019)

Host species: *Pagurus filholi*, *P. minutus*, *P. nigrivittatus*, *Pagurus angustus* (Anomura; Paguridae).

***Peltogaster reticulata* Shiino, 1943**

(see *P. aff. reticulata* and *Peltogaster* sp. 4 of Jung et al. 2019)

Material reexamined: on *P. minutus*: 1 individual, Namhae, 34°52'40.83"N, 127°56'43.61"E, 14 Nov. 2012, collected by Jung J., MADBK 160706_065, host: female, shield length 4 mm; 1 individual (2 externae), Namhae, 34°52'40.83"N, 127°56'43.61"E, 12 May 2018, collected by Jung J., MADBK 160706_125, host: female, shield length 3 mm.

On *P. proximus*: 1 individual, Shinan, 34°35'49.24"N, 125°45'58.02"E, 16 Oct. 2008, collected by Hong J., MADBK 160718_040, host: female, shield length 3 mm.

Host species: *Pagurus constans*, *P. minutus*, *Pagurus proximus* (Anomura; Paguridae).

Distribution: Japan (type locality), Russia, Korea.

***Peltogaster* sp. 1**

(Fig. 2C)

Material examined: on *P. lanuginosus*: 1 individual, Gangneung, 37°44'04.3"N, 128°59'25.8"E, 23 Oct. 2021, collected by Jung J., DSEVIV0000000053. On *P. maculosus*: 1 individual, Sinan, 34°40'55.1"N, 125°26'38.1"E, 26 Aug. 2021, collected by Jung J., DSEVIV0000000052, KoreaHS1-KoreaHS4, host: females, shield length 8 mm; Seogwipo, 33°14'05.1"N, 126°21'30.1"E, 6 Sep. 2021, DSEVIV0000000057, host: shield length 9 mm.

Host species: *P. lanuginosus*, *P. maculosus* (Anomura; Paguridae).

Distribution: Korea.

Diagnosis: Whole externa single, smooth, and irregular oval in shape. Mantle aperture slightly elevated or elevated tube, and vertically circular opening at the extremity.

***Peltogaster* sp. 2**

(see *Peltogaster* sp. 3 at Jung et al. 2019)

Host species: *Paguristes ortmanni* (Anomura; Diogenidae).

**Polyascidae Høeg & Glenner in Høeg, Noever,
Rees, Crandall & Glenner, 2019**

***Parasacculina* Høeg & Glenner in Høeg, Noever,
Rees, Crandall & Glenner, 2019**

***Parasacculina imberbis* (Shiino, 1943)**

(Fig. 2D, see Jung et al. 2021)

Additional materials examined: on *Pachygrapsus crassipes*: 1 individual, Seogwipo, 33°14'09.6"N,

126°21'35.4"E, 6 Sep. 2021, collected by Jung J., Koreax8, host: ♀, carapace length 14 mm; 1 individual, Tongyeong, 34°45'31.3"N, 128°30'59.4"E, 29 Oct. 2021, collected by Jung J., Korearr54; 1 individual, Jeju, 33°30'57.3"N, 126°30'38.5"E, 23 Aug. 2022, collected by Jung J., Koreakr10.

Host species: *P. crassipes* (Brachyura; Grapsidae).

Diagnosis: Whole externa single, smooth or slightly wrinkled, and slightly bilobed with half-oval-shaped left and right lobes divided by a mid-notch. Mantle aperture slightly elevated as short tube shape and vertically circular opening at the extremity.

Color: yellow.

***Parasacculina pilosella* (Van Kampen & Boschma, 1925)**

(Fig. 2E, see Jung et al. 2021)

Additional materials examined: on *Pugettia intermedia*: 1 individual, Sokcho, 38°13'38.1"N, 128°35'16.1"E, 8 Sep. 2020, Korearr1; 1 individual, Goseong, 38°27'01.3"N, 128°28'03.9"E, 25 Sep. 2021, GS 2381.

Host species: *P. intermedia*, *Pugettia* aff. *ferox* (Brachyura; Epiplatidae).

***Parasacculina pinnotherae* (Shiino, 1943)**

(see Jung et al. 2021)

Additional materials examined: on *Arcotheres sinensis*: 3 individuals, Jindo, 34°23'16.9"N, 126°17'20.8"E, 23 July 1994, KoreaJD1.

Host species: *A. sinensis* (Brachyura; Pinnotheridae).

***Parasacculina shiinoi* (Lützen, Itani, Jespersen, Hong, Rees & Glenner, 2016)**

(see Jung et al. 2021)

Additional materials examined: on *Upogebia major*: 1 individual, Namhae, 34°52'40.83"N, 127°56'43.61"E, 26 Sep. 2021, collected by Jung J., Korea56.

Host species: *U. major* (Gebiidea; Upogebiidae).

Diagnosis: Whole externa single, smooth, and irregular oval shape with a shallow median groove on dorsal margin. Mantle aperture vestigial and vertically large circular opening with radiated ridges at the extremity.

***Parasacculina yatsui* (Boschma, 1936)**

(Fig. 2F, see Jung et al. 2021)

Additional materials examined: on *Cyclograpsus*

intermedius: 1 individual, Ulleung, 37°28'32.4"N, 130°53'51.5"E, 19 Sep. 2019, collected by Jung J., Korearr40.

On *Hemigrapsus penicillatus*: 1 individual, Goseong, 38°16'11.5"N, 128°33'27.2"E, 26 Sep. 2021, collected by Jung J., GS12, host: ♂, carapace length 20 mm; 1 individual, Gangneung, 37°44'04.3"N, 128°59'25.8"E, 23 Oct. 2021, collected by Jung J., Korearr10. On *Hemigrapsus sanguineus*: 1 individual, Goheung, 34°30'22.2"N, 127°28'56.0"E, 9 Oct. 2022, collected by Jung J., KoreaN14; 1 individual, Goseong, 38°27'01.3"N, 128°28'03.9"E, 25 Sep. 2021, collected by Jung J., Korea71, host: ♀, carapace length 19 mm; 1 ind, Tongyeong, 34°38'24.9"N, 128°34'06.1"E, 19 Aug. 2019, collected by Jung J., Korea39, host: ♂, carapace length 14 mm. On *P. crassipes*: 1 individual, Yeongdeok, 36°16'22.0"N, 129°22'44.3"E, 29 Aug. 2021, collected by Jung J., Korea72, host: ♀, carapace length 22 mm; 1 individual, Gangneung, 37°44'04.3"N, 128°59'25.8"E, 23 Oct. 2021, collected by Jung J., Korearr9; 1 individual, Samcheok, 37°12'50.5"N, 129°20'46.5"E, 19 May 2022, collected by Jung J., Korearr25; 1 individual, Jeju, 33°30'57.3"N, 126°30'38.5"E, 23 Aug. 2022, collected by Jung J., Koreakr9.

Host species: *P. crassipes* (Brachyura; Grapsidae), *H. sanguineus*, *H. penicillatus*, *C. intermedius* (Brachyura; Varunidae).

***Parasacculina oblonga* (Lützen & Yamaguchi, 1999)**

(Korean name: Bi-Dan-Ge-Yeop-Ju-Meo-Ni-Beol-Rae) (Fig. 2G)

Material examined: on *C. intermedius*: 1 individual, Ulleung, 37°28'32.4"N, 130°53'51.5"E, 3 June 2021, collected by Jung J., DSEVIV0000000050 host: ♀, carapace length 12 mm; 1 individual, Ulleung, 37°28'32.4"N, 130°53'51.5"E, 3 June 2021, collected by Jung J., DSEVIV0000000051, host: ♀, carapace length 12 mm; 1 individual, Ulleung, 37°28'32.4"N, 130°53'51.5"E, 19 Sep. 2019, collected by Jung J., DSEVIV0000000054; 1 individual, Ulleung, 37°30'49.9"N, 130°47'45.0"E, 18 Sep. 2019, collected by Jung J., DSEVIV0000000055; 1 individual, Ulleung, 37°30'49.9"N, 130°47'45.0"E, 18 Sep. 2019, collected by Jung J., DSEVIV0000000056, host: ♂, carapace length 14 mm.

Host species: *C. intermedius* (Brachyura; Varunidae).

Distribution: Japan (type locality), Korea.

Diagnosis: Whole externa single, smooth or slightly wrinkled, and oval in shape. Mantle aperture large, elevated, and vertically Y-shape opening at the

extremity.

***Parasacculina* sp. 1**
(Fig. 2H)

Material examined: on *H. sanguineus*: 1 individual, Goheung, 34°31'7.0"N, 127°26'12.2"E, 8 Oct. 2022, collected by Jung J and Park J-K., HVBNI0000000005, host: ♀, carapace length 24 mm; 1 individual, Goheung, 34°30'22.2"N, 127°28'56.0"E,

9 Oct. 2022, collected by Jung J., Korea13N, host: ♀, carapace length 33 mm; 1 individual, Goheung, 34°30'22.2"N, 127°28'56.0"E, 9 Oct. 2022, collected by Jung J. and Park J-K., HVBNI0000000006, host: ♀, carapace length 25 mm.

Host species: *H. sanguineus* (Brachyura; Varunidae).

Distribution: Korea.

Diagnosis: whole externa smooth, single, and cordiform with flat half-oval-shaped left and right




Fig. 2. Externae of Korean rhizocephalans in this study. Blue arrow: mantle aperture, scale bar: 2 mm for H and J, 1mm for A-G, I, K, L. A, *Peltogasterella gracilis*; B, *Peltogasterella* sp.; C, *Peltogaster* sp. 1; D, *Parasacculina imberbis*; E, *Parasacculina pilosella*; F, *Parasacculina yatsui*; G, *Parasacculina oblonga*; H, *Parasacculina* sp. 1; I, *Polyascus* cf. *gregarius*; J, *Heterosaccus papillosus*; K, *Sacculina angulata*; L, *Sacculina gracilis*. Externae of C and H are incomplete, because they were sampled for molecular analysis.

lobes divided by outer mid-groove and inner mid-ridge. Mantle aperture slightly elevated, and slightly wrinkled with circular opening at extremity.

Parasacculina sp. 2

(see *Parasacculina* sp. 1 of Jung et al. 2021)

Host species: Macromedaeus distinguendus (Brachyura; Xanthidae).

Parasacculina sp. 3

(see *Parasacculina* sp. 2 of Jung et al. 2021)

Host species: Macromedaeus distinguendus (Brachyura; Xanthidae).

Polyascus Glenner, Lützen & Takahashi, 2003
***Polyascus* cf. *gregarius* (Okada & Miyashita, 1935)**

(Fig. 2I, see *Polyascus* cf. *gregarius* of Jung et al. 2021)

Additional materials examined: on *H. sanguineus*: 1 individual, Yeosu, 34°29'22.4"N, 127°48'49.5"E, 11 May 2021, collected by Jung J., Korea45, 1 individual, Wando, 34°17'47.4"N, 126°42'05.1"E, 24 May 2021, collected by Jung J., Korea47, host: ♂, carapace length 34 mm; 1 individual, Tongyeong, 34°45'31.3"N, 128°30'59.4"E, 29 Oct. 2021, collected by Jung J., Korearr56; 1 individual, Shinan, 24 May 2016, collected by Jung J., Korearr58; 1 individual, Busan, 13 May 2020, collected by Jung J., Korearr60; 1 ind, Jindo, 34°31'24.4"N, 126°12'56.5"E, 6 July 2022, collected by Jung J., Koreakr1; 1 individual, Goheung, 34°31'7.0"N, 127°26'12.2"E, 8 Oct. 2022, collected by Jung J., Korea1N.

Host species: H. sanguineus, H. takanoi, Eriocheir japonica (Brachyura; Varunidae).

Sacculinidae Lilljeborg, 1861
***Heterosaccus* Smith, 1906**
***Heterosaccus papillosus* (Boschma, 1933)**

(Fig. 2J)

Material examined: on *Charybdis* (*Charybdis japonica*): 1 individual, Incheon, 37°15'54.2"N, 126°29'51.8"E, 23 Aug. 2012, NIBRIV0000895329; 1 individual, Incheon, 37°24'31.4"N, 126°38'41.0"E, 18 Aug. 1998, NIBRIV0000895330.

Host species: Charybdis (*Gonioneptunus bimaculata*, *C. (C.) japonica* Brachyura; (Portunidae).

Distribution: Philippines (type locality), Japan, Korea.

Diagnosis: Whole externa single, smooth or slightly wrinkled, and reversed cordiform with left and

right lobes divided by a mid-notch. Mantle aperture slightly elevated or elevated, radically wrinkled, and vertically large circular opening at the extremity.

***Sacculina* Thompson, 1836**
***Sacculina confragosa* Boschma, 1933**

(see Jung et al. 2021)

Additional materials examined: on *Gaeticte depressus*: 1 individual, Goheung, 34°24'56.9"N, 127°29'36.5"E, 12 May 2021, collected by Jung J., Korea52, host: ♂, carapace length 18 mm.

on *P. crassipes*: 1 individual, Sokcho, 38°12'50.9"N, 128°36'05.5"E, 6 Sep 2021, collected by Jung J., Korea73, host: ♀, carapace length 17 mm; 1 individual, Jeju, 33°56'40.8"N, 126°20'03.5"E, 7 Sep 2021, collected by Jung J., Korea74, host: ♀, carapace length 14 mm; 1 individual, Gangneung, 37°44'04.3"N, 128°59'25.8"E, 23 Oct. 2021, collected by Jung J., Korearr8; 1 individual, Uljin, 37°03'32.7"N, 129°25'46.0"E, 24 Oct. 2022, collected by Jung J., Korearr28; 1 individual, Samcheok, 37°08'51.2"N, 129°21'48.9"E, 19 May 2022, collected by Jung J., Korearr53; 1 ind, Seogwipo, 33°12'28.3"N, 126°17'30.7"E, 21 Aug. 2022, collected by Jung J., Koreakr11.

on *H. sanguineus*: 1 individual, Gangneung, 37°44'04.3"N, 128°59'25.8"E, 18 May 2022, collected by Jung J., Korearr22; 1 individual, Pohang, 36°00'14.4"N, 129°34'32.1"E, 11 May 2020, collected by Jung J., Korearr59; 1 ind, Jindo, 34°31'25.0"N, 126°12'06.4"E, 6 July 2022, collected by Jung J., Koreakr2; 1 ind, Goheung, 34°31'7.0"N, 127°26'12.2"E, 8 Oct. 2022, collected by Jung J., Korea7N.

Host species: G. depressus, P. crassipes (Brachyura; Grapsidae), *C. intermedius, H. sanguineus* (Brachyura; Varunidae).

***Sacculina angulate* Van Kampen & Boschma, 1925**

(Korean name: O-Gak-Ge-Ju-Meo-Ni-Beol-Rae)
(Fig. 2K)

Material examined: on *Thalamita sima*: 1 individual (3 externae, 1 externa removed for molecular analysis), Busan, 35°9'6.4"N, 129°10'43.1"E, 25 Apr. 2018, collected by SH Kim, NIBRIV0000895328 host: ♂, carapace length 23 mm; 1 individual, Sacheon, 34°53'2.0"N, 128°4'34.6"E, 16 July 2014, collected by SH Kim, NIBRIV0000895331, host: ♀, carapace length 25 mm.

Host: Xiphonectes longispinosus, C. (G.) bimaculata, T. sima (Brachyura; Portunidae).

Distribution: Indonesia (type locality), Thailand,

Taiwan, Korea.

Diagnosis: Whole externa single or triple, smooth or slightly wrinkled, and irregular pentagonal in shape. Mantle aperture large, elevated, radically wrinkled, and vertically large circle opening at the extremity.

***Sacculina gracilis* Boschma, 1931**

(Korean name: Mae-Kkeun-Ge-Ju-Meo-Ni-Beol-Rae)
(Fig. 2L)

Material examined: on *T. sima*: 1 individual, Seogwipo, 33°12'38.6"N, 126°15'35.5"E, 12 Aug. 2010, collected by HS Ko, NIBRIV0000895332, host: ♂, carapace length 21 mm; 1 individual, Tongyeong, 34°50'14.5"N, 128°9'29.9"E, 9 June 2013, collected by HY Suk, NIBRIV0000895334, host: ♂, carapace length 16 mm.

Host species: *Xiphonectes longispinosus*, *T. sima*, *Thalamita picta*, *Thranita danae*, (Brachyura; Portunidae) *Heteropilumnus setosus*, *Ser fukiensis*, (Brachyura; Pilumnidae) *Notonyx vitreus* (Brachyura; Goneplacidae).

Distribution: Thailand (type locality), Indonesia, Singapore, Philippines, China, Korea.

Diagnosis: Whole externa single, smooth, and round fusiform in shape. Mantle aperture small, slightly elevated, and vertically Y-shape opening at the extremity.

***Sacculina upogebiae* Shiino, 1943**

(see Lützen et al. 2016)

Materials examined: on *U. major*: 1 individual, Boryeong, 36°21'19.7"N, 126°32'40.2"E, 6 June 2012, NIBRIV0000453552; 1 individual, Incheon, 37°13'50.9"N, 126°31'56.9"E, 21 Mar. 2014, NIBRIV0000453567.

Host species: *U. major* (Gebiidea; Upogebiidae).

DISCUSSION

In this study, we compared morphological and molecular information of Korean Rhizocephala species with previously published studies and discussed details of their morphological features and sequence divergences, and taxonomic consideration.

***Peltogasterella* sp.:** The absence of a chitinous shield is a clear morphological indicator that this species is a member of the genus *Peltogasterella*. In addition, the elongated externa on the present specimens are similar to the characteristics of *P. gracilis*. However, the mantle aperture is not elevated, the number of externa is 1 to 3, and the *cox1* sequence (OR481994)

is different from that of *P. gracilis* and the similarity is below 74.4%. Moreover, this species is found only on the abdomen of *P. nihonkaiensis*. Therefore, we declare that these specimens are new species candidates of the genus *Peltogasterella*.

***Peltogaster reticulata*:** MADBK 160718_040 has morphological characteristics such as a single banana-shaped externa, an elevated mantle aperture, and an X-shaped mantle aperture opening, which correspond with the original description (Shiino 1943). In addition, the hosts of these specimens were identical to previous studies of *P. reticulata* in Russia (Kashenko and Korn 2003; Korn et al. 2020). Furthermore, the *cox1* sequence of MADBK 160706_065 has high DNA sequence similarity to the Russian *P. reticulata* (98.9–99.4% to MN193580–MN193582, Korn et al. 2020). Therefore, the Korean *P. aff. reticulata* and *Peltogaster* sp. 4 in Jung et al. (2019) were identified as *P. reticulata*.

***Peltogaster* sp. 1:** This species is regarded as a *Peltogaster* species because of the presence of a chitinous shield on the externa which is the main morphological characteristic of *Peltogaster* and *cox1* sequences of this species nested within *Peltogaster* species. In addition, this species has single oval externa identical to those of *P. lineata*, another parasite of *P. maculosus* in Japan. However, this species has irregular oval externa and a well-developed mantle aperture, as opposed to *P. lineata*, which has regular oval externa and a vestigial mantle aperture. In addition, *cox1* sequences of this species (OR481985–OR481991) significantly differ from those of *P. lineata* and other *Peltogaster* species and the similarity is below 78.2%. Therefore, we declare that these specimens are new species candidates of the genus *Peltogaster*.

***Parasacculina imberbis*:** The morphological characters of the Korean *P. imberbis*, which could not be observed due to immature specimens in Jung et al. (2021), were observed through additional specimens in this study. These morphological characters such as a yellow and bilobed single externa and a slightly elevated mantle aperture tube are identical to the original description (Shiino 1943).

***Parasacculina shiinoi*:** The morphological characters of Korean *P. shiinoi* which could not be observed due to immature specimens in Jung et al. (2021), were observed through additional specimens in this study. These morphological characteristics such as irregular oval externa and a vestigial mantle aperture tube are identical to the original description (Lützen et al. 2016).

***Parasacculina yatsui*:** Jung et al. (2021) noted that Korean *P. yatsui* infects only *H. sanguineus*. The present study revealed that Korean *P. yatsui* also infects *P. crassipes*, *H. penicillatus*, and *C. intermedius*. This

is the first report that *P. yatsui* is parasitized on *H. penicillatus*.

Parasacculina oblonga: These samples are identical in the host species (*C. intermedius*) and morphological characters to the original description (Lützen and Yamaguchi 1999) such as single round externa, and a large and elevated mantle aperture. Furthermore, the BLAST result showed that the sequences of this specimen (OR481965–OR481969) have very high sequence similarity to the sequences of the same GenBank species (DQ059780) at 99.4–99.8%. Therefore, we identified these specimens as *P. oblonga*. This study noted the expanded northward distribution of *P. oblonga*.

Parasacculina sp. 1: *Parasacculina yatsui* and *Polyascus* cf. *gregarius* have been reported to be parasites of *H. sanguineus*. However, these specimens differ in morphology and molecular sequences from *P. yatsui* and *P. cf. gregarius*. This species has a single smooth externa, whereas *P. cf. gregarius* has 1–2 smooth externae and *P. yatsui* has a single externa with a wrinkled outermost part (Jung et al. 2021). In addition, this species has a circular mantle aperture opening whereas *P. yatsui* and *P. cf. gregarius* have slit mantle aperture openings (Jung et al. 2021). BLAST results showed that *cox1* sequences of this species (OR481970–OR481972) are different from those of *P. yatsui* and *P. cf. gregarius* and all other polyascids and the similarity is below 74.9%. Therefore, we considered this species to be a new species candidate of the genus *Parasacculina*.

Heterosaccus papillosus: This species parasite on the *C. (C.) japonica* was first reported in Korea by Kim and Hong (2002). However, the morphology and the reference specimen of Korean *H. papillosus* were not presented in Kim and Hong (2002). In this study, the morphology and molecular characteristics of Korean *H. papillosus* were presented for the first time. The host (*C. (C.) japonica*) and morphological characteristics of these specimens such as a single reversed heart shape externa and a well-elevated, pillar-shaped mantle aperture were consistent with the original description (Boschma 1933). In addition, the *cox1* sequence of Korean *H. papillosus* (OR481955) was distinct from the *cox1* sequences of other *Heterosaccus* species, yet still more similar to other *Heterosaccus* than to other genera in the BLAST test (94.9% similar to *Heterosaccus lunatus* DQ059778).

Sacculina confragosa: Jung et al. (2021) mentioned that Korean *S. confragosa* infects only *G. depressus*. This study revealed that Korean *S. confragosa* also infects *P. crassipes* and *H. sanguineus*. This is the first report that *S. confragosa* is parasitized on *H. sanguineus*.

Sacculina angulata: Morphological characteristics of the examined specimen such as an irregular pentagonal externa and an elevated large mantle aperture correspond with the original description of *S. angulata* (Van Kampen and Boschma 1925). In addition, the host of these specimens (*T. sima*) is identical to the host in the previous study (Boschma 1931). Furthermore, the BLAST result showed that the sequence of this species (OR482032) is similar to that of *Sacculina* sp. (EF521402) at 83.0%. Therefore, we identified these specimens as *S. angulata*. This study noted that the distribution of *S. angulata* expanded northeastward.

Sacculina gracilis: The morphological characteristics of the examined specimen such as a single fusiform externa and a slightly elevated small mantle aperture correspond with the original description of *S. gracilis* (Boschma 1931). In addition, the host (*T. sima*) of these specimens was identical to the previous study of Boschma (1955). However, the BLAST result showed that the sequences of this species (OR482063–OR482064) are most similar to *Parasacculina beauforti* (KX426583) at 80.2%. A taxonomical revision of the genus of *S. gracilis* is required by further molecular analysis using additional DNA markers. This is the third report on *S. gracilis* after the original description (Boschma 1931) and Boschma (1955) and since previous studies, we noted that the species' distribution has expanded eastward.

CONCLUSIONS

Through this study, a checklist of 16 Korean parasitic barnacles was presented, three rhizocephalans were found to have an expanded distribution range, and three new species candidates in Korea were identified. In addition, we noted brief remarks on Korean parasitic barnacles that were not reported in previous studies. This study suggests that further biodiversity studies of Rhizocephala should be conducted by examining more diverse hosts beyond those previously reported. In addition, this study will be a reference for examining the fauna of East Asian Rhizocephala and conducting related further studies on parasitic barnacles.

List of abbreviations

mitochondrial cytochrome *c* oxidase subunit I, *coxI*.

Acknowledgments: This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR202203202) and the

Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2021R1A6A1A10039823, 2021R1A6A3A01086374). This research was supported by Learning & Academic research institution for Master's PhD students, and Postdocs (LAMP) Program of the National Research Foundation of Korea (NRF) grant funded by the Ministry of Education (No. RS-2023-00301976). We appreciate Jongwoo Jung (Ewha Womans University), who helped sample the materials analyzed in this study.

Authors' contributions: Jibom Jung contributed to the study design, fieldworks, acquisition, data analysis, interpretation of the sequence data, and manuscript draft. Joong-Ki Park contributed to the study design, acquisition, data analysis, interpretation of the sequence data, and drafting the manuscript.

Competing interests: JJ and J-KP declare that they have no conflict of interest.

Availability of data and materials: The *cox1* data in this study were deposited in GenBank (Table 1). Voucher specimens in this study were deposited in the National Institute of Biological Resources (NIBR) and the Honam National Institute of Biological Resources (HNIBR).

Consent for publication: Not applicable.

Ethics approval consent to participate: Not applicable.

REFERENCES

- Boschma H. 1927. On the larval forms of Rhizocephala. Proc Sect Sci K Ned Akad Wet Amst **30**:293–297.
- Boschma H. 1931. Papers from Dr. Th. Mortensen's Pacific Expedition 1914–16. LV. Rhizocephala. Vidensk Meddel Dansk Naturhist Foren Kjøbenhavn **89**:297–380.
- Boschma H. 1933. The Rhizocephala in the collection of the British Museum. J Linn Soc **38**:473–552, 1 pl.
- Boschma H. 1936. The specific characters of *Sacculina rotundata* Miers and *Sacculina yatsui* nov. spec. Zool Meded **19**:1–22.
- Boschma H. 1955. Rhizocephalan parasites of the crab *Pugettia brevisstris*, with notes on *Sacculina gracilis*. Zool Meded **33**(24):237–249.
- Burmeister H. 1834. Beiträge zur Naturgeschichte der Rankenfüßer (Cirripedia). G. Reimer, Berlin, Germany.
- Chan BKK, Dreyer N, Gale AS, Glenner H, Ewers-Saucedo C, Pérez-Losada M, Kolbasov GA, Crandall KA, Høeg JT. 2021. The evolutionary diversity of barnacles, with an updated classification of fossil and living forms. Zool J Linn Soc **193**(3):789–846. doi:10.1093/zoolinnean/zlaa160.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. Mol Mar Biol Biotechnol **3**(5):294–299.
- Glenner H, Lützen J, Takahashi T. 2003. Molecular and morphological evidence for a monophyletic clade of asexually reproducing Rhizocephala: *Polyascus*, new genus (Cirripedia). J Crustac Biol **23**:548–557. doi:10.1651/C-2361.
- Gravel A. 1905. Monographie des Cirripèdes ou Thécostracés. Masson et Cie, Éditeurs, Paris, France.
- Høeg JT. 1992. Rhizocephala. In: Harrison FW, Humes AG (ed) Microscopic anatomy of invertebrates, vol 9, Crustacea. Wiley, New York, pp. 313–345.
- Høeg JT, Lützen J. 1995. Life cycle and reproduction in the Cirripedia Rhizocephala. Oceanogr Mar Biol **33**:427–485.
- Høeg JT, Noever C, Rees DA, Crandall KA, Glenner H. 2019. A new molecular phylogeny-based taxonomy of parasitic barnacles (Crustacea: Cirripedia: Rhizocephala). Zool J Linn Soc **190**(2):632–653. doi:10.1093/zoolinnean/zlz140.
- Jung J, Yoshida R, Kim W. 2019. Diversity of parasitic Peltogastrid barnacles (Crustacea: Cirripedia: Rhizocephala) on hermit crabs in Korea. Zool Stud **58**:33. doi:10.6620/ZS.2019.58-33.
- Jung J, Yoshida R, Lee D, Park J-K. 2021. Morphological and molecular analyses of parasitic barnacles (Crustacea: Cirripedia: Rhizocephala) in Korea: preliminary data for the taxonomy and host ranges of Korean species. PeerJ **9**:e12281. doi:10.7717/peerj.12281.
- Kashenko SD, Korn OM. 2003. Combined effects of seawater temperature and salinity on development of the larvae of the rhizocephalan *Peltogaster reticulatus* (Crustacea: Cirripedia). Russ J Mar Biol **29**:150–155. doi:10.1023/A:1024664631438.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analyses of sequence data. Bioinformatics **28**:1647–1649. doi:10.1093/bioinformatics/bts199.
- Kim HK, Chan BKK, Lee SK, Kim W. 2020. Biogeography of intertidal and subtidal native and invasive barnacles in Korea in relation to oceanographic current ecoregions and global climatic changes. J Mar Biol Assoc U.K. **100**:1079–1091. doi:10.1017/S0025315420001009.
- Kim HK, Chan BKK, Yi C, Kim IH, Choi YN. 2022. Barnacle epibiosis on sea turtles in Korea: A west Pacific Region with low occurrence and intensity of *Chelonibia testudinaria* (Cirripedia: Chelonibiidae). Front Ecol Evol **10**:785692. doi:10.3389/fevo.2022.785692.
- Kim KB, Hong SY. 2002. Parasitism of the portunid crab, *Charybdis japonica* by the rhizocephalan barnacle, *Heterosaccus papillosus* in the western coast of Korea. Proc Korean Soc Fish Technol Conf **2002**:504–505.
- Kobayashi M, Wong YH, Oguro-Okano M, Dreyer N, Høeg JT, Yoshida R, Okano K. 2018. Identification, characterization, and larval biology of a rhizocephalan barnacle, *Sacculina yatsui* Boschma, 1936, from northwestern Japan (Cirripedia: Sacculinidae). J Crustac Biol **38**(3):329–340. doi:10.1093/jcbl/ry020.
- Korn OM, Golubinskaya DD, Sharina SN. 2020. The parasitic barnacle *Peltogaster reticulata* Shiino, 1943 (Rhizocephala, Peltogastridae) from Russian waters of the Sea of Japan: morphological description, molecular identification and complete larval development. Zootaxa **4768**(1):6–24. doi:10.11646/ZOOTAXA.4768.1.2.
- Krüger P. 1912. Über ostasiatische Rhizocephalen. Anhang: Über einige Vertreter der Cirripedia Thoracica. Abh Math-Phy. K., K Bayer Akad Wiss **2**:1–8.

- Muhammad BL, Seo Y, Chae J, Ki JS. 2022. Population genetic structure and demographic expansion of the exotic jellyfish *Carybdea brevipedalia* (Cubozoa, Cnidaria) into the Korean coast. *Zool Stud* **61**:48. doi:10.6620/ZS.2022.61-48.
- Müller F. 1862. Die rhizocephalen, eine neue gruppe schmarotzender kruster. *Arch Naturgesch* **28**:1–9, pl. 1.
- Lilljeborg W. 1861. Supplément au mémoire sur les genres Liriope et *Peltogaster*, H. Rathke. *Nova Acta Regiae Soc Sci Upsal* **3**:74–102, pl. 6.
- Lützen J, Itani G, Jespersen Å, Hong JS, Rees D, Glenner H. 2016. On a new species of parasitic barnacle (Crustacea: Rhizocephala), *Sacculina shiinoi* sp. nov., parasitizing Japanese mud shrimps *Upogebia* spp. (Decapoda: Thalassinidea: Upogebiidae), including a description of a novel morphological structure in the Rhizocephala. *Zool Sci* **33**(2):204–212. doi:10.2108/zs150112.
- Lützen J, Yamaguchi T. 1999. *Sacculina oblonga*, new species (Cirripedia: Rhizocephala), parasitic on the grapsid crab *Cyclograpsus intermedius* (Decapoda: Brachyura) from Japan. *J Crustac Biol* **19**:886–890. doi:10.1163/193724099X00565.
- Okada YK, Miyashita Y. 1935. Sacculinization in *Eriocheir japonicus* de Haan, with remarks on the occurrence of complete sex-reversal in parasitized male crabs. *Mem Coll Sci Kyoto Imp Univ, Ser* **10**(3):169–208.
- Øksnebjerg B. 2000. The Rhizocephala (Crustacea: Cirripedia) of the Mediterranean and Black Seas: taxonomy, biogeography, and ecology. *Isr J Zool* **46**:1–102. doi:10.1560/RCLC-NM2U-HV5L-6Q52.
- Rathke H. 1842. Beiträge zur vergleichenden Anatomie und Physiologie, Reisebemerkungen aus Skandinavien, nebst einem Anhang über die rückschreitende Metamorphose der Thiere. *Nueste Schrift Naturf Ges Danzig* **3**:i–v + 1–162, pls. 1–6.
- Shiino SM. 1943. Rhizocephala of Japan. *J Sigenkagaku Kenkyusyo* **1**:1–36.
- Smith G. 1906. Rhizocephala. *Fauna Flora Golfes Neapel* **29**:1–123, pls. 1–18.
- Thompson JV. 1836. Natural history and metamorphosis of an anomalous crustacean parasite of *Carcinus maenas*, the *Sacculina carcini*. *Entomol Mon Mag* **3**:452–456.
- Van Kampen PN, Boschma H. 1925. Die Rhizocephala der Siboga-Expedition. *Siboga-Exped* **31bis**:1–62, pls. 1–3.
- Von Siebold CT. 1848. Lehrbuch der vergleichenden Anatomie der Wirbellosen Thiere. Erster Theil. *In*: von Siebold CT, Stannius H (ed) *Lehrbuch der vergleichenden Anatomie*. Verlag von Veit & Comp, Berlin, pp. 1–679.
- Yoshida R, Hirose M, Hirose E. 2014. Hermit crab host prevalence by species of Peltogastridae (Cirripedia: Rhizocephala): hosts vary with locations on the Pacific coast in mainland Japan. *J Crust Biol* **34**(4):467–480. doi:10.1163/1937240X-00002246.
- Yoshida R, Hirose M, Mok HK, Hirose E. 2012. The first records of peltogastrid rhizocephalans (Crustacea: Cirripedia: Rhizocephala) on hermit crabs (Paguroidea) in Taiwan and differences in prevalences among collection sites. *Zool Stud* **51**(7):1027–1039.
- Yoshida R, Osawa M, Hirose M, Hirose E. 2011. A new genus and two new species of Peltogastridae (Crustacea: Cirripedia: Rhizocephala) parasitizing hermit crabs from Okinawa Island (Ryukyu Archipelago, Japan), and their DNA-barcodes. *Zool Sci* **28**:853–862. doi:10.2108/zsj.28.853.