## **ERRATA** for the Solutions Manual of T. Shifrin's *Multivariable Mathematics: Linear Algebra, Multivariable Calculus, and Manifolds*

## Note: All of these have been corrected in the second printing, June, 2017.

Solutions Manual, p. 23, **1.4.28d**. There is no Corollary 2.2. One must use the analogous reasoning with the rows of P to deduce that  $PP^{\mathsf{T}} = I$  as well.

Solutions Manual, p. 27, **1.5.4**. The vector  $A\mathbf{b}_2 = \begin{bmatrix} a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$ .

Solutions Manual, p. 40, 2.2.13. min should be max.

Solutions Manual, p. 40, **2.2.14a**.  $|x_{k_j} - x_0| \le |b - a|/2^j \to 0$ .

Solutions Manual, p. 71, **3.6.6**. In the fourth line from the end,  $\frac{\partial x}{\partial v}$  and  $\frac{\partial y}{\partial v}$  are missing; the third line from the end should be deleted.

Solutions Manual, p. 85, 4.1.21. Delete "A is singular, and so."

Solutions Manual, p. 126, **5.2.14**. The upper limit on the summation should be k, not l.

Solutions Manual, p. 138, **5.4.10**. The four critical points on the unit circle should have  $1/\sqrt{2}$  in front of them. Now the maximum occurs at  $\begin{bmatrix} 0\\0\\1 \end{bmatrix}$  and the minimum at  $\pm \frac{1}{\sqrt{2}} \begin{bmatrix} -1\\1\\0 \end{bmatrix}$ .

Solutions Manual, p. 168, 6.2.1d. The entries of the final matrix should be  $-e^{x_0}$  and  $-e^{y_0}$ .

Solutions Manual, p. 170, 6.2.3c. A minus sign got dropped at the very last entry.

Solutions Manual, p. 170, 6.2.3e. A factor of 1/2 was dropped in computing  $D\phi(\mathbf{x}_0)$ .

Solutions Manual, p. 176, **6.3.10**. We should have  $D\mathbf{F}(\mathbf{p}) = \begin{bmatrix} 2 & 2 & -2 & 2\\ 1 & 1 & 1 & -1 \end{bmatrix}$  and, resultingly, the basis for the tangent space should be given by  $\left\{ \begin{bmatrix} -1\\1\\0\\0 \end{bmatrix}, \begin{bmatrix} 0\\0\\1\\1 \end{bmatrix} \right\}$ .

Solutions Manual, p. 184, 7.1.1d. The correct answer is 15/2.

Solutions Manual, p. 184, 7.1.2a. The upper limit on the inner integral should be y.

Solutions Manual, p. 187, 7.2.9.  $0 \le x \le y/2$ , and the correct answer is 32/3.

Solutions Manual, p. 189, **7.2.12c**. We should have  $|x| \le z \le 1$ .

Solutions Manual, p. 194, **7.3.3**. The correct answer is  $3\pi/2$ .

Solutions Manual, p. 195, **7.3.10**. The correct answer is  $\pi/2$ .

Solutions Manual, p. 196, **7.3.14**. The final integrand should be  $1 - |\cos^3 \theta|$ . The answer is correct.

Solutions Manual, p. 196, **7.3.16**. The upper limit on the z integral should be  $\sqrt{a^2 - r^2}$ .

Solutions Manual, p. 197, **7.3.21b**. We should have  $\pi/\sqrt{a}$ , not  $\sqrt{\pi a}$ , and, similarly, the final answer is  $\pi^{3/2}/\sqrt{6}$ .

Solutions Manual, p. 203, 7.4.27b. A factor of G is missing in the final answer.

Solutions Manual, p. 222, 8.2.2f.  $(x^2 + y^2 + z^2)^{-1}$ 

Solutions Manual, p. 227, 8.3.4. We need  $z = |\sin t|$ , and the correct answer is  $-8/3 - \pi$ .

Solutions Manual, p. 230, 8.3.16b. The correct answer is  $21(15 + \frac{9}{4}\pi)$ .

Solutions Manual, p. 235, **8.4.4**. The final integral should be  $8 \int_0^{\pi/2} \int_0^{2\cos\theta} dz \, d\theta = 16$ .

Solutions Manual, p. 236, 8.4.6b. That det T = 1 is a red herring; what is relevant is that  $T^* \sigma = \sigma$ .

Solutions Manual, p. 238, 8.4.16b. A factor of  $a^4$  is missing.

Solutions Manual, p. 238, 8.4.16c.  $\mathbf{g}^* \omega = \cdots - 1 d\theta \wedge dr$ ; answer is  $-\pi/2$ .

Solutions Manual, p. 239, **8.4.18b**. Delete the  $\frac{1}{a}$  at the beginning of the second line.

Solutions Manual, p. 239, 8.4.19a. This is off by a factor of -1 because of orientation.

Solutions Manual, p. 245, 8.5.15b. The coefficient of  $dx \wedge dy$  should be  $(1 - z^2)$ .

Solutions Manual, p. 245, **8.5.16a**.  $\cos(t/2)$  should be  $\cos(\theta/2)$ .

Solutions Manual, p. 247, **8.5.21c**. The 1-form given does not give the area of a subset in the sphere. We need a 1-form  $\mathbf{g}^{-1*}\eta$  where  $d\eta = \mathbf{g}^*\sigma$ . Its existence is guaranteed by Exercise 8.7.12.

Solutions Manual, p. 288, **9.3.20c**.  $(x - a)^2$  should be  $(t - a)^2$ .

Solutions Manual, p. 296, **9.4.19d**.  $\frac{1}{\sqrt{3}}y_3$  should be  $\sqrt{3}y_3$ .

—July, 2017