

Taxonomy and systematics

## A new species of blindsnake from Jalisco, Mexico (Squamata: Leptotyphlopidae)

### *Una especie nueva de serpiente agujilla de Jalisco, México (Squamata: Leptotyphlopidae)*

Oscar A. Flores-Villela <sup>a, \*</sup>, Eric N. Smith <sup>b</sup>, Luis Canseco-Márquez <sup>a</sup>,  
Jonathan A. Campbell <sup>b</sup>

<sup>a</sup> Universidad Nacional Autónoma de México, Facultad de Ciencias, Museo de Zoología, Circuito Exterior s/n, Ciudad Universitaria, Coyoacán 04510 Ciudad de México, Mexico

<sup>b</sup> University of Texas, Department of Biology, 701 S. Nedderman Drive, Arlington, Texas 76019-0498, USA

\*Corresponding author: ofv@unam.mx (O.A. Flores-Villela)

Received: 23 February 2021; accepted: 19 April 2021

<http://zoobank.org/urn:lsid:zoobank.org:pub:8050D5B1-99C3-4C7C-AFF0-74511C90DC28>

#### Abstract

A new snake species of the genus *Rena* is described from northern Jalisco, Mexico. The new species represents an isolated member of the *R. dulcis* group in the extreme southwest Mesa Central of the country. We redefine the *R. dulcis* and *R. humilis* groups within the genus *Rena*. The status of the other species allocated to these groups is discussed.

*Keywords:* Blind snakes; Taxonomy; Nomenclature; Squamata; Serpentes

#### Resumen

Una especie nueva de serpiente perteneciente al género *Rena* se describe del norte de Jalisco, México. La nueva especie representa un miembro aislado del grupo de *R. dulcis*, en el extremo suroeste de la Mesa Central del país. Redefinimos los grupos *R. dulcis* y *R. humilis* dentro del género *Rena*. Se discute el status de otras especies dentro de estos grupos.

*Palabras clave:* Agujillas; Taxonomía; Nomenclatura; Squamata; Serpientes

## Introduction

While surveying amphibians and reptiles and their parasites in northern Jalisco during the summer of 2003, we obtained a specimen of blindsnake east of the town of Bolaños. This specimen belongs to the *dulcis* group of *Leptotyphlops* as defined by Klauber (1940), now allocated to the genus *Rena* (Adalsteinsson et al., 2009). This group is characterized by having a cream-colored ventral surface with scant dark pigmentation, lacking a sharply contrasting white spot on the snout or tail tip, and lacking a pattern of longitudinal lines on the dorsum. Following Klauber (1940), two subgroups usually have been recognized within the genus, the *R. dulcis* and the *R. humilis* subgroups, and considered as separated groups by us. For reasons discussed below we refer to the *R. dulcis* group those species having 10 scale rows around the tail. The only known specimen of the new species was collected 220 km (straight-line) from the closest known locality in San Luis Potosí for any other member of the *R. dulcis* group. In view of its geographical isolation and distinctive characters, we propose that it be recognized as a new species within the *R. dulcis* group. The validity of this taxon was tested by Adalsteinsson et al. (2009) using molecular techniques, referred as *Rena* sp. B.

## Materials and methods

Scale nomenclature and the method for determining numbers of middorsal and subcaudal scales follow Klauber (1940), except for nomenclature of the anterior midorsal head scales that we follow Wallach (2003). Counts were done using a dissecting microscope, and measurements were taken with a ruler (to nearest 0.1 mm) or an electronic digital caliper (to nearest 0.01 mm). Values for asymmetric head characters are given in left/right order. Museum specimens examined are listed in the appendix.

The map for the species of the *Rena dulcis* group was generated with information gathered from the following collections: AMNH, ANSP, AUM, BMNH, BYU, CAS, CM, CNAR, CU, UF (FLMNH), EHT-HMS, FMNH, INHS, KU, LACM, LSU, MCZ, MPM, MSB, MSUM, MVZ, MWSU (Midwestern State University), MZFC, NLU, OMNH, OS, PSM, SDMNH, SM (BCB), SNIB (Sistema Nacional de Información sobre Biodiversidad de México), SRSU, TCWC, TNHC, TTU, UAZ, UCM, UIMNH, UMMZ, UNL (Universidad de Nuevo León), UOMZ, USNM, UTA, UTEP, WTSU (West Texas A&M University), YPM. Localities were obtained directly from museum collections, the HerpNet database (October 2009), from Dixon and Vaughan (2003). Museum acronyms can be found in Sabaj (2016).

## Description

The discovery of a snake allied to *R. dulcis* in northern Jalisco, and the comparison of this snake with other taxa in the *Rena dulcis* group has led us to conclude that it represents a new species, here named as:

*Rena klauberi* new species. Fig. 1.

<http://zoobank.org/urn:lsid:zoobank.org:act:17770C36-1C8E-4171-A8D0-4E4682496071>

*Leptotyphlops* sp. B. Adalsteinsson et al. (2009): 7 (Fig. 3), 48 (App. 1).

*Rena* sp. B. Adalsteinsson et al. (2009): 20, 31 (Fig. 12), 48 (App. 1).

*Holotype*: MZFC-17047 from Mexico, Jalisco, Río Cartagena, adjacent to the road between Santa María de Los Ángeles and Bolaños, 1,602 m (21°59'0.24" N, 103°20'27.816" W). Original field number JAC 23308, collected by Eric Smith and Jesse Meik, on 10 June 2003.

*Diagnosis*. A species of the *R. dulcis* group that differs from *R. maxima*, in having 10 rows of scales around the tail (vs. 12); from *R. bressoni*, *R. myopica*, and *R. dissecta* in having an undivided anterior supralabial (vs. divided); from *R. dulcis* in having nine scales on the dorsal surface with dark pigment (vs. seven); from *R. humilis* in having supraoculars (vs. absent); and by having 10 rows of scales around the tail (12 in *R. humilis* group members, except for *R. h. segregata* and *R. h. tenuicula* that have 10). *Rena klauberi* differs from all other Mexican *Rena*, except the holotype of *R. d. iversoni*, in having a darkly pigmented cloacal plate, while the remainder of the venter is immaculate.

*Description* of the holotype. Middorsal scales 252 between the rostral scale and the tail tip; 14 rows of smooth, imbricate scales around body, constant between head and level in front of cloaca; 10 rows of scales around midpoint of tail; subcaudals 14, not including tail tip; cloacal plate undivided; rostral scale curving over snout, posterodorsal end rounded; nasal divided horizontally, nostril between upper and lower nasals, located medially between the suture; lower nasals extend to lip; four scales bordering mouth on each side behind rostral (lower nasal, anterior supralabial, ocular, and posterior supralabial); anterior supralabials 1/1, undivided; posterior supralabials 1/1, higher than wide, barely touching corner of mouth; parietals large, three times higher than wide, contacting posterior supralabials; supraoculars 1/1, equal in size to the frontal; occipitals 1/1, subequal to parietals, undivided; temporals 1/1, preventing contact between occipitals and posterior supralabials; postfrontal slightly wider than frontal, and separating supraoculars; interparietal slightly wider than postfrontal and interoccipital (Fig. 2); mental scale about

three and a half times wider than long; infralabials 4/4, first pair separated medially by postmental, second pair larger than first and third and separated from each other by three median chin shields, fourth pair slender; body cylindrical from head to tail, with ultimate scale terminating in a well-defined spine.

**Coloration.** After preservation, nasal openings surrounded by pale coloration; central area of rostral with a pale transverse marking; eyes visible through skin (Figs. 1, 2); seven dorsal scale rows uniformly dark brown; adjacent scale row slightly paler (total nine dark rows), pattern extending along entire length of body and tail (Fig. 1); venter pinkish on anterior half of body, grading to cream on posterior half; cloacal plate covered with dark pigmentation (Fig. 3); midventral row of subcaudal scales cream; paraventral subcaudal rows brown near cloaca and cream color distally. In life, dorsal coloration similar to that in preservative.

**Measurements.** Total length 265 mm; tail length 14.4 mm; relative tail length (or tail/LOA) 5.4%; horizontal diameter of body at head 3.8 mm, diameter at midbody 4.7 mm, diameter of body at base of tail 4.4 mm; diameter of head at interocular level 3.4 mm; relative rostral width 1.4; relative body proportion (or length/width ratio) 56.4; tail length/tail width ratio 3.27; rostral width/head width ratio 0.71.



Figure 1. Photograph in life of holotype of *R. klauberi* MZFC-17047.

#### *Taxonomic summary*

**Distribution.** *Rena klauberi* is known only from the type-locality (Fig. 4) in dry scrub forest. The holotype was found crawling just after sunset on a dirt path near the outer periphery of a riparian zone consisting of grasses, sedges, and scattered oak trees (Fig. 5). It is likely that this species may also occur in nearby southern Zacatecas, which possesses similar habitat. Several reptile species,




Figure 2. Drawing of dorsal (top) and lateral (bottom) views of head of holotype of *R. klauberi* MZFC-17047.



Figure 3. Photograph of ventral surface of tail of *R. klauberi* MZFC-17047, showing coloration of cloacal plate and tail.

apparently endemic to the area, have been reported for both southern Zacatecas and northern Jalisco (Ponce-Campos et al., 2001).

**Etymology.** This species is named in honor of the late Laurence Monroe Klauber, in recognition of his valuable contributions to North American herpetology, and especially to our knowledge of the genus *Leptotyphlops* (= *Rena*) from North America.

#### Remarks

Based on morphology, *R. klauberi* appears to be related to *R. dulcis*. Both *R. klauberi* and *R. dulcis* can be distinguished from all other species in the *R. dulcis* group by having undivided anterior supralabials (typical condition in *R. dulcis*), 10 scale rows around the tail, and supraocular scales. Adalsteinsson et al. (2009), used DNA to show a close phylogenetic relationship between *R. dulcis* and *R. dissecta* (here considered synonymous with *R. dulcis*), to the exclusion of *R. klauberi*. Using mitochondrial cytochrome b sequences used by Adalsteinsson et al. (2009), deposited in Genbank, we calculated raw genetic distances among the species they examined. The central Texas *R. dulcis* and southeastern Arizona *R. dissecta* show only a 5% sequence difference, a very small distance for samples coming from populations more than 630 km apart, regarded here as within species variation. Between *R. dissecta* or *R. dulcis* and *R. klauberi* there is an 11% sequence divergence, which we attribute to a species level difference. Species from the *R. dulcis* group differed from those within the *R. humilis* group by between 14-15%, genetic distance. Additionally, Adalsteinsson et al. (2009) tree (their figure 3) supports the monophyly of the *R. humilis* and *R. dulcis* groups.

The taxonomy of the blindsnakes allied to the genus *Rena*, was reviewed by Adalsteinsson et al. (2009), Dixon and Vaughan (2003), Hahn (1979a, 1979b, 1980a), Klauber (1940) and Smith and Chiszar (1993). These authors reached different conclusions on the status of the taxa associated with the genus and the species that comprise it. Currently, the genus *Rena* is composed of 10 to 11 species (Table 1), 2 of them with 2 (*R. dulcis*) and 4 (*R. humilis*) subspecies respectively (Uetz et al., 2022; Wallach et al., 2014).

For a long time, the species within this genus were placed in the widespread genus *Leptotyphlops*, until Adalsteinsson et al. (2009) found that this genus was composed of 2 clades, one of them containing New World species, restoring the name *Rena* (Baird & Girard, 1853) for this group of taxa. Nevertheless, the taxonomy of the genus *Rena* is still controversial and a thorough taxonomic revision is urgently needed. Currently, the genus *Rena* is distributed in North America from central and southern Mexico to southwestern United States (see below).

To clarify the status of the taxa allocated to the genus *Rena*, we reviewed characters that have been used, mainly by Klauber (1940). Klauber (1940) recognized the *dulcis* and *humilis* subgroups, which are currently placed in the genus *Rena*, recognizing four characters differentiating both subgroups (considered separate groups by us); number of middorsal scales (236-246 *dulcis*; 254-275 *humilis*), division of the anterior supralabials (undivided in *R. dulcis* group), total length (*R. dulcis* under 300 mm; *R. humilis* over 300 mm), and the presence of supraocular scales in the *R. dulcis* group, lacking in *R. humilis*. All of these characters, however, show considerable variation within each group and even within recognized species




Figure 4. Geographic distribution of members of the *R. dulcis* subgroup in Mexico and United States. Inset shows type locality of *R. klauberi*.

(Table 2). Therefore, we redefine these groups with a more consistent character, the number of scales around the tail, the *R. dulcis* group having 10 and the *R. humilis* group having 12 scales around the tail (Table 2). We adopt this new diagnostic character, the number of scales around the tail, as more important than others in defining

these groups in the past (e.g., number of supraoculars); this character defines and coincides with the relationships recovered and depicted by Adalsteinsson et al. (2009). This lepidotic count is consistent within each known species and apports the taxa to the east and west along the geographically important Cochise Filter Barrier. The Continental Divide has been important in isolating the biota of the Sonoran and Chihuahuan deserts (Castoe et al., 2007; Morafka, 1977; Myers et al., 2017; Provost et al., 2018, and references cited therein), and the formation of this barrier appears to be responsible for separating the snakes of the *R. dulcis-humilis* groups, partitioning them into 2 distinct taxonomic units. *Rena humilis* with 12 or more scales around tail, includes species occurring west of the Cochise Filter Barrier: and the *R. dulcis* group is characterized by 10 scales around tail and occurs east of the Cochise Filter Barrier.

Below we discuss the status and validity of some taxa in the *R. dulcis-humilis* groups, particularly as impacted by our newly rearranged groups, and the new species described here. The *R. humilis* group is presented first for ease of discussion.

*Rena dugesii* (Bocourt, 1881). Lemos-Espinal et al. (2004) elevated *R. h. dugesii* to full species status,



Figure 5. Photograph of the type-locality of *R. klauberi* taken June 10, 2003.

Table 1

Species currently recognized by the two main sources of information about snakes species in the world. Species are sorted by group based on the results of this publication, see text for more details.

Species	The Reptile Data Base	Wallach et al. (2014)
<i>R. dulcis</i> group		
<i>R. dulcis</i> (2 ssp.)	Valid	Valid
<i>R. dissecta</i>	Valid	Valid
<i>R. iversoni</i>	Valid	Valid
<i>R. myopica</i>	Valid	Valid
<i>R. bressoni</i>	Valid	Valid
<i>R. unguirostris</i>	Valid	As <i>Siagonodon unguirostris</i>
<i>R. humilis</i> group		
<i>R. boetgeri</i>	Valid	Synonym of <i>R. humilis</i>
<i>R. dugesii</i>	Valid	Valid
<i>R. humilis</i> (4 ssp.)	Valid	Valid
<i>R. segregata</i>	Valid	Valid
<i>R. maxima</i>	Valid	Valid

comparing this taxon to *R. h. segregata*, in Chihuahua. We agree with Lemos-Espinal and Smith (2007) in restricting the distribution of *R. dugesii* to the Pacific slopes from Sonora to Colima, considering a Guanajuato record erroneous. That record was based on a specimen sent to the U.S. National Museum by Dugès. There is another specimen at the Dugès Museum in Guanajuato (MADUGHE 90) from Colima; Dugès did not record this species from Guanajuato (Dugès, 1890, 1895), so it is unlikely that it occurs there. The closest record of this species to Guanajuato is that of Peterson et al. (1995) from Chapala, Jalisco.

*Rena humilis tenuicula* (Garman, 1883 [1884]). *Stenostoma tenuiculum* was described from a single specimen from San Luis Potosí. It was diagnosed by having 10 scales around the tail and less than 250 middorsal scales (Klauber, 1940). Brown and Brown (1967) reported a second specimen (BCB 11560) from Llera, Tamaulipas. This second specimen belongs to a population existing in geographical proximity to localities where *R. iversoni* (Smith, van Breukelen, Auth & Chiszar, 1998) has been collected. Smith et al. (1998) made this specimen (BCB 11560) part of the type series of *R. iversoni*, arguing

that BCB 11560 has divided anterior supralabials, a low number of middorsals, and that the only other known specimen, the holotype of *R. humilis tenuicula*, comes from the “semiarid plateau in San Luis Potosí”. Klauber (1940) commented that it was not clear whether the type-locality “San Luis Potosí” refers to the state or the city, and as Brown and Brown (1967) recognized, the type-locality may be situated 60-200 miles (96-320 km) from Llera, Tamaulipas, the general area occupied by *R. iversoni*. It is worth noting that the holotype of *R. tenuicula* lacks divided anterior supralabials, but has 10 scales around the tail. Based on our examination of the holotype of *R. h. tenuicula* (*Stenostoma tenuiculum*, MCZ 4519), and the considerable variation in middorsal scale counts, supraoculars and anterior supralabials exhibited by *R. dulcis* (Table 2), we consider *R. tenuicula* as belonging to the *R. dulcis* group. If proven to be a valid taxon, a matter that needs more investigation; its relationships to other members of the *R. dulcis* group, should be considered. For the present we place it in the synonymy of *R. dulcis*.

Klauber (1940), thought *R. tenuicula* might be related to *R. dugesii*, pending the revision of *R. dugesii* and confirmation of the number of scales around tail. Uetz et al. (2022) and Wallach et al. (2014) considered *R. tenuicula* as a synonym of *R. dugesii*. Our examination of the type of *Catodon dugesii* (Bocourt, 1881) revealed that it has 12 scale rows around the tail. Therefore, considering this taxon as a synonym of *R. dugesii* by Uetz et al. (2022) and Wallach et al. (2014) is unjustified.

*Rena humilis chihuahuensis* (Tanner, 1985). This taxon was synonymized by Lemos-Espinal and Smith (2007) based on variation of middorsal scale counts provided by Klauber (1940) for populations of *R. humilis segregata* from Arizona (254-280). We agree with these authors in relegating this taxon to *R. segregata* (see *R. h. segregata* account below). Therefore it should not be considered as a valid subspecies of *R. humilis*, as indicated by Uetz et al. (2022).

*Rena humilis segregata* (Klauber, 1939). This taxon is characterized by having 10 rows of scales around the tail, as in other members of the *R. dulcis* group, but it lacks supraocular scales, similar to members of the *R. humilis* group. Nevertheless, its distribution falls mostly to the east of the Cochise Filter Barrier and has undivided supralabials, similar to *R. dulcis* and *R. klauberi* of the *R. dulcis* group. *Rena segregata* is a distinct species from *R. humilis* and unique within the *R. dulcis* group, therefore we recognize it as a valid taxon and part of the *R. dulcis* group.

*Rena dulcis* (Baird & Girard, 1853). This species differs from other members in this group, except *R. klauberi*, by having supraoculars and undivided anterior supralabials (Dixon & Vaughan, 2003). Klauber (1940) recognized 3

Table 2

Selected characters used to diagnose species discussed in this paper. Data modified from Klauber (1940), Dixon and Vaughan (2003), and specimens examined (Appendix). Taxa we recognize appear in bold type. Typical condition in number of scales, excluding number of middorsals, appears first and is defined as the condition present in the largest number of individuals examined or reported in the references.

Species	Middorsals	Scales around body	Scales around tail	Supraoculars	Anterior supralabials	Parietal-posterior supralabial contact	Number of pigmented dorsals (adding half scales)	Dorsal color (as reported for live or freshly preserved)	Maximum size
<b><i>Rena dulcis</i> group</b>									
<b><i>R. dulcis</i></b>	<b>192-257</b>	<b>14</b>	<b>10</b>	<b>1/1 1/0 0/0</b>	<b>1/1 2/2 1/2 2/1 0/1 0/0</b>	<b>Yes</b>	<b>7 (5-8)</b>	<b>Pinkish, light to medium, and dark brown to black</b>	<b>272</b>
<i>R. d. rubella</i>	222-257	14	10	1/1	1/1	Yes	7 (6-7)	Pale brown	191
<b><i>R. dissecta</i></b>	<b>213-255</b>	<b>14</b>	<b>10</b>	<b>1/1</b>	<b>2/2 1/2</b>	<b>Yes</b>	<b>7</b>	<b>Pinkish</b>	<b>272</b>
<b><i>R. iversoni</i></b>	<b>202-226</b>	<b>14</b>	<b>10</b>	<b>0/0 1/0 1/1</b>	<b>2/2</b>	<b>Yes</b>	<b>7 (5-8)</b>	<b>Medium to dark brown</b>	<b>196</b>
<i>R. myopica</i>	192-236	14	10	1/1	2/2 1/2 1/1	Yes	7	Dark brown to black	227
<b><i>R. bressoni</i></b>	<b>227-246</b>	<b>14</b>	<b>10</b>	<b>1/1</b>	<b>2/2 1/2</b>	<b>No</b>	<b>7</b>	<b>Pale brown</b>	<b>265</b>
<b><i>R. klauberi</i></b>	<b>252</b>	<b>14</b>	<b>10</b>	<b>1/1</b>	<b>1/1</b>	<b>Yes</b>	<b>9</b>	<b>Dark brown</b>	<b>265</b>
<b><i>R. segregata</i></b>	<b>253-287</b>	<b>14</b>	<b>10</b>	<b>0/0 0/1</b>	<b>1/1</b>	<b>Yes</b>	<b>7</b>	<b>Pale to dark brown, grey</b>	<b>319</b>
<b><i>Rena humilis</i> group</b>									
<b><i>R. boettgeri</i></b>	<b>244-269</b>	<b>14</b>	<b>12</b>	<b>0/0</b>	<b>1/1</b>	<b>Yes</b>	<b>5</b>	<b>Pale to dark brown</b>	<b>253</b>
<b><i>R. dugesii</i></b>	<b>231-259</b>	<b>14</b>	<b>12</b>	<b>0/0</b>	<b>1/1</b>	<b>Yes</b>	<b>7 (7-9)</b>	<b>Medium brown</b>	<b>187</b>
<b><i>R. h. cahuilae</i></b>	<b>280-305</b>	<b>14</b>	<b>12</b>	<b>0/0</b>	<b>1/1</b>	<b>Yes</b>	<b>7 (5-7)</b>	<b>Almost white (pink), brown to orange</b>	<b>389</b>
<b><i>R. h. humilis</i></b>	<b>253-291</b>	<b>14</b>	<b>12</b>	<b>0/0</b>	<b>1/1</b>	<b>Yes</b>	<b>7 (7-9)</b>	<b>Medium to dark brown</b>	<b>315</b>
<b><i>R. h. utahensis</i></b>	<b>289-308</b>	<b>14</b>	<b>12</b>	<b>0/0</b>	<b>1/1</b>	<b>Yes</b>	<b>7</b>	<b>Pale gray-brown</b>	<b>322</b>
<i>R. h. chihuahuensis</i>	253-257	14	10	0/0	1/1	Yes	7	Pigmented	133
<i>R. humilis tenuicula</i>	244	14	10	0/0	1/1	Yes	7 (6-7)	Pale gray-brown	117
<b><i>R. maxima</i></b>	<b>216-235</b>	<b>14</b>	<b>12</b>	<b>1/1</b>	<b>1/1</b>	<b>Yes</b>	<b>7</b>	<b>Dark brown to orange</b>	<b>300</b>

subspecies: *R. d. dulcis*, *R. d. myopica* and *R. d. dissecta*. This arrangement was followed by Hahn (1979a, 1980b), McDiarmid et al. (1999) and Smith et al. (1998). Smith et al. (1998) described an additional subspecies, *R. d. iversoni* (see below). Dixon and Vaughan (2003) elevated to species rank the subspecies of *R. dulcis* recognized by

Klauber (1940), and recognized the taxon *R. d. iversoni* as a subspecies of *R. myopica*, on the basis of middorsal scale counts. Dixon and Vaughan (2003) also recognized *R. rubella* as a subspecies of *R. dulcis* (see below) and assigned the name to populations from southwestern Texas and adjacent Mexico. Dixon and Vaughan (2003) also

recognized an unnamed population of *Rena dulcis* ssp. from Seminole, Oklahoma, USA, which needs further confirmation.

*Rena dulcis rubella* (Garman, 1883 [1884]). Stejneger (1891) synonymized *R. rubella* with *R. dulcis*, demonstrating that the original description of Garman (1883 [1884]), based on a single specimen, had numerous errors. Garman reported erroneously 15 dorsal scale rows in *R. dulcis*, instead of 14 (as in other members of the genus *Rena* and most Leptotyphlopids); the rostral separating the nasals as a diagnostic character (as all Leptotyphlopids); 5 infralabials instead of 4 (as in all members of the genus *Rena*); and the anterior parietal (= parietal) being the only scale contacting the posterior labial, implying that in *R. dulcis* the posterior parietal (= occipital) also contacts the posterior labial (Stejneger, 1891). The latter condition, as mentioned by Stejneger, is only found on the left side of the holotype of *R. rubella*, and is variable in *R. dulcis*, contact only by the parietal being the prevalent condition. Klauber (1940) concurred with Stejneger (1891) and maintained *R. rubella* in the synonymy of *R. dulcis*, adding that Garman misinterpreted the original description of Baird and Girard (1853), and pointing out that the type-locality of *R. rubella* is within the geographical distribution of *R. dulcis*. This taxon was resurrected as a subspecies of *R. dulcis* by Dixon and Vaughan (2003). In spite of broad overlap in ranges of variation, the number of middorsal scale rows was one of the criteria used by Dixon and Vaughan (2003: 14, 22: key) to distinguish 3 subspecies of *R. dulcis*: *R. d. dulcis* (210-246), *R. d. rubella* (222-257), and an unnamed subspecies from Oklahoma (202-228). The other diagnostic character was dorsal coloration; “pinkish” for *R. d. dulcis* and “light to medium brown” for *R. d. rubella*. No dorsal color pattern was described for the unnamed subspecies. Because these 2 taxa show considerable overlap in middorsal counts, their distributional ranges are parapatric as delimited by Dixon and Vaughan (2003), and their middorsal scale counts decrease in a north to south clinal pattern, we find no valid argument to recognize any subspecies within *R. dulcis*. Current examination of the holotype of *R. rubella* (MZC 4584) does not provide evidence of its original color (see Table 2) and it seems necessary to have other evidence, besides scale variation and dorsal coloration, to arrive at a taxonomic decision involving this name. Our examination of *R. dulcis* from throughout its range in Texas reveals considerable variation in dorsal coloration, from pink to medium brown to black.

Additionally, Dixon and Vaughan (2003) reported a population of *Rena* from Querétaro and Hidalgo (their Population 5a, central Mexico) which they associated with *R. dulcis*. We have examined additional material

from central Mexico, and from a locality in northeastern Zacatecas. The Zacatecas locality is geographically intermediate between the central Mexico population and those in Tamaulipas and Nuevo León. The Zacatecas and central Mexico specimens coincide in middorsal scale counts with those to the north. We consider *R. rubella* as a junior synonym of *R. dulcis* and populations in central Mexico (Querétaro, Hidalgo, and Zacatecas) to represent *R. dulcis* populations.

*Rena myopica* (Garman, 1883 [1884]). This species was distinguished from other species in the *R. dulcis* group (including specimens previously assigned to *R. dissecta*) by possessing the following combination of characters (Dixon & Vaughan, 2003; Klauber, 1940): divided anterior supralabials; no postocular; parietals contacting posterior supralabials; and 192-255 middorsals. We have examined specimens in which the postoculars and the anterior supralabials are variable; the middorsal scale count falls within the range of *R. dulcis*, the lower count may be just part of a north-south cline found by Dixon and Vaughan (2003). Therefore, we do not consider this taxon valid, and it should be considered a junior synonym of *R. dulcis*. Dixon and Vaughan (2003) recognized 2 subspecies, *R. m. myopica* and *R. m. iversoni* (status of this taxon discussed below). Wallach et al. (2014), on the basis of Pinto (2010) unpublished dissertation, recognized this taxon as valid as well as Uetz et al. (2022).

*Rena dissecta* (Cope, 1896). This taxon was distinguished from other members of the *R. dulcis* subgroup by the following combination of characters: divided anterior supralabials, no postocular, parietals contacting posterior supralabials, middorsal scale counts, and dorsal coloration (pinkish, orange, or pale to dark brown Dixon and Vaughan [2003]). Klauber (1940) stated that *R. d. dissecta* can be consistently differentiated from *R. d. myopica* by number of middorsal scales, but Dixon and Vaughan (2003) showed that middorsal scale counts overlap in these taxa. Klauber (1940) considered 2 additional characters to further differentiate these taxa: in *R. dissecta* the fifth dorsal scale is wider than the fourth (equal width in *R. myopica*) and the occipital scales are divided on at least one side of the head in many individuals of *R. dissecta* (59% of the specimens he examined vs. always complete in *R. myopica*). Dixon and Vaughan (2003) stated that such characters are highly variable and should not be used in diagnoses, and we agree after examination of additional specimens. In 2 specimens of *R. myopica* that we examined, the fifth dorsal scale is wider than the fourth and the occipitals are single. As currently recognized, *R. dissecta* occurs in the southwestern United States (Arizona, New Mexico, Oklahoma, and Texas) and



northern Mexico (Chihuahua and Coahuila), it seems to be mostly allopatric from *R. dulcis* and *R. myopica*, with only limited distributional overlap (see map in Hahn [1979a] and Dixon and Vaughan [2003]). We do not agree with Dixon and Vaughan (2003) in recognizing *R. dissecta* as a separate species from *R. myopica*. These authors based the recognition of these as 2 separate taxa primarily on their supposed differences in dorsal color, pinkish in *R. dissecta* and brown to black in *R. myopica*. Klauber (1940) reports specimens of *R. dissecta* ranging from pale to medium brown. We have examined additional specimens of *R. dissecta* not seen by Dixon and Vaughan (2003), and we observed grey (e.g., UTA R-54613) and brown (UTA R-45091) dorsal colors. We have also observed pinkish (UTA R-3149) and medium brown (UTA R-54555) specimens of *R. myopica* from Nuevo León and Tamaulipas, respectively. The number of middorsal scales was also stated by Dixon and Vaughan (2003) as significant in differentiating these taxa. Examination of additional material considerably expands the lower limit in the range of these scales for *R. dissecta*, from 220 to 213, providing considerable overlap between *R. dissecta* and *R. myopica*. Based on the lack of differences between the 2 taxa, we consider *R. dissecta* a junior synonym of *R. dulcis* (Table 2).

*Rena iversoni* (Smith, van Breukelen, Auth & Chiszar, 1998). This taxon is of special interest because, unlike all other populations formerly in the *R. dulcis* group, it lacks supraoculars in most known specimens (10 out of 13). Smith et al. (1998) described this taxon as a subspecies of *R. dulcis*, based on it having 10 scales around the tail. We agree that the affinities of this taxon are with the *R. dulcis* group. The absence of supraoculars in *R. iversoni* may be the result of fusion of the ocular and supraocular scales. We consider the diagnostic features of this taxon are variable, although they may have systematic value. The matter needs to be resolved using molecular techniques. Wallach et al. (2014) on the basis of Pinto (2010) unpublished dissertation recognized this taxon as valid, as well as Uetz et al. (2022). We provisionally accept the status as valid species, pending more evidence.

In summary, we recognize 9 species in the *Rena dulcis-humilis* groups, including *R. bressoni* and *R. maxima* (Table 3). We do not recognize any subspecies for *R. dulcis*. Within *R. humilis* we recognize tentatively 2 subspecies, both with 12 scales around the tail and lacking supraoculars (Tables 2, 3). The *R. dulcis* and *R. humilis* groups are mainly separated from each other, in the northern part of their distribution, by the Cochise Filter Barrier between the Sonoran and Chihuahuan deserts.

Table 3

Taxonomic changes proposed in this paper.

Species previously recognized	Species recognized in this work
<i>R. dulcis</i> group	
<i>R. d. dulcis</i>	<i>R. dulcis</i>
<i>R. d. rubella</i>	<i>R. dulcis</i>
<i>R. dissecta</i>	<i>R. dulcis</i>
<i>R. h. tenuicula</i>	<i>R. dulcis</i>
<i>R. myopica</i>	<i>R. dulcis</i>
<i>R. iversoni</i>	<i>R. iversoni</i>
<i>R. klauberi</i>	<i>R. klauberi</i>
<i>R. bressoni</i>	<i>R. bressoni</i>
<i>R. h. chihuahuensis</i>	<i>R. segregata</i>
<i>R. segregata</i>	<i>R. segregata</i>
<i>R. humilis</i> group	
<i>R. boettgeri</i>	<i>R. boettgeri</i>
<i>R. dugesii</i>	<i>R. dugesii</i>
<i>R. h. humilis</i>	<i>R. humilis</i>
<i>R. h. utahensis</i>	<i>R. h. utahensis</i> , not treated here
<i>R. maxima</i>	<i>R. maxima</i>

### Acknowledgements

We thank J. Meik, J. Malone, T. Devitt, and P. Ponce-Campos for field assistance, J. Rosado and V. Wallach, Museum of Comparative Zoology, for providing us with information related to the type of *L. rubellum* and photographs of the specimen. We thank for the loan of comparative material or providing specimen information: T. Crumpton of Baylor University, Mayborn Museum Complex (Strecker Museum); G. Schneider of the University of Michigan Museum of Zoology; D. R. Frost of the American Museum of Natural History; S. Rogers of the Carnegie Museum; J. T. Giermakowski of the Museum of Southern Biology; V. H. Reynoso of the Instituto de Biología, UNAM; R. McDiarmid and S. Gotte of the Smithsonian Institution; R. G. Webb and C. Lieb from University of Texas, El Paso; N. Vidal from MNHN-Paris; G. Magaña-Cota from the Museo Alfredo Dugès. We thank F. Wallach and J. Boundy for sending relevant information. For preparing specimen loans we thank E. Pérez Ramos, MZFC, and C. Franklin, UTA. J.

Meik and R. Pinto made extensive valuable suggestions. J. A. Hernández Gómez helped with the figures. This paper is based in part upon work supported by the National Science Foundation under grant No. DEB-0613802 to J. A. Campbell; ENS participation was possible by NSF

grant DEB-0416160 and a grant from Bioclon, Mexico. Necessary permits were issued by Semarnap to OFV. The final revision of this paper was done while OFV was on sabbatical at UTA supported by the Conacyt and DGAPA, UNAM, Mexico.

#### Appendix. Specimens examined.

*Rena bressoni*: Mexico, Michoacán: MZFC-12356–12357: Coalcomán, 2 km SE of Coalcomán; MZFC-12358: Aguililla, 5 km W of Aguililla; MZFC-12390: Coalcomán, 4 km N Puerto de la Zarzamora, 1.5 km NW of C. El Laurel.

*Rena dissecta*: USA, Arizona: UTA R-26544: Cochise, Portal road, 0.5 km E Portal; UTA R-50598: Cochise, Forest Service barracks; Texas: UTA R-33758: Jeff Davis, 3.0 mi E of end of FM 1832 on FM 1832; UTA R-54613: Terrel.

*Rena dulcis*: USA, Texas: UTA R-32303: Bosque, 2.5 mi S of jct W FM 927 and Texas 144, W of 144 on AW Vickrey property; UTA R-1596: Cameron, South most Palm Grove, E Brownsville; UTA R-9166-9170: Cooke, 15.8 mi S Gainesville on FM 51; UTA R-36890: Crockett, 8 mi SE of Barnhart; UTA R-57102-103: Crockett, Dirth road to the south of Hwy 67 runs paralell to train tracks; UTA R-896-898: Dallas, Oak Cliff; UTA R-1444: Dallas, Hwy 67 at Hampton Road; UTA R-10273: Dallas, Dallas, Dallas Zoo; UTA R-9171: Denton, 19.4 mi S Gainesville on FM 51; UTA R-16799-800: Denton, N of Aubrey, Main Street ca. 2 km N jct Blackjack Road; UTA R-15680: Denton, unknown locality; UTA R-38466: Eastland, 1.5 mi N Cisco on 380; UTA R-15681–15682: Frio, 9.6 km NE Dilley near jct Leona and Frio Rivers; UTA R-16200: Frio, Panther Hollow Ranch, ca. 11 km NW Dilley; UTA R-10503: Garza, NE Post; UTA R-30067: Goliad, Goliad, grounds of courthouse; UTA R-26552: Henderson, no other data; UTA R-398-399: Hidalgo, E McAllen; UTA R-33378: Jeff Davis, 3.0 mi E of end of FM 1832 on FM 1832; UTA R-32815: Jim Hogg; S of Hebronville on FM 1017, 8.4 mi from jct FM 1017 and FM 285; UTA R-16415-420: Johnson, Johnson Ranch, 21.7 km W Rio Vista; UTA R-38402: Kendall, on 743, 5 mi N of Sisterdale; UTA R-11123-24, 12649-651: Llano, 12.8 km S Cherokee on St Hwy 16, Houston Ranch; UTA R-11075: McMullen, 11.1 km S Tilden on Texas Hwy 16 at creek; UTA R-15176: Montague, Sundagne Ranch, S side Sandy Creek, 7.7 km S, 4.2 km E jct US Hwy 287 and FM 1125; UTA R-9172: Palo Pinto, Birdwell Ranch, 1.5 mi S Palo Pinto on FM 4; UTA R-1234: Parker, 3 mi E Cresson; UTA R-1440: Parker, 4.8 km NE of Aledo Mary's Creek area; UTA R-5043: Parker, FM rd 2376 2 mi S Aledo, behind graveyard; UTA R-9173-74: Parker, 2.5 mi S Aledo, on road off of FM 2376; UTA R-26545-547: Parker, ca. 3 air km NW Wheatland, along Bear Creek; UTA R-7890-91: San Saba, 2.9 mi S, 1.0 mi E Bend; UTA R-55708-709: Shackelford, Hwy 351 S junction with Hwy 180; UTA R-599: Tarrant, Arlington (College Farm); UTA R-600: Tarrant, Arlington (4 blks N of Turnpike on Fielder Road); UTA R-863: Tarrant, Arlington; UTA R-18688: Tarrant, Arlington, UTA campus; UTA R-25669: Tarrant, Arlington, UTA Central Services Building, 1225 West Mitchell; UTA R-25712: Tarrant, Arlington, 1426 S West Street; UTA R-31418: Tarrant, University of Texas at Arlington campus, crossing path near Student Union Building; UTA R-32380: Tarrant, Arlington, in house near UTA campus; UTA R-38873: Tarrant, Arlington, UT Arlington campus, Lipscomb Hall; UTA R-55402: Tarrant, Arlington Veteran's Park; UTA R-19327: Tarrant, Benbrook-Aledo Road; UTA R-19328: Tarrant, ca. 1 km NW jct Loop 820 and Rufe Snow; UTA R-18272: Tarrant, Euless, 402 Huntington Drive; UTA R-28691: Tarrant, Euless, 607 Bent Tree Court; UTA R-5696: Tarrant, Ft. Worth, 800 block of Sylvania street near Trinity River; UTA R-14727-729: Tarrant; Fort Worth, 600 Congress; UTA R-40737: Tarrant, Fort Worth; UTA R-1233: Tarrant, Lake Worth; UTA R-1930: Tarrant, N. Richland Hill; UTA R-1585-86: Tarrant, Trinity River and Tex 360; UTA R-9175-77: Tarrant, 2.9 km S Jct FM 1886 on White Settlement Road; UTA R-394, 1232, 45050: Travis, Austin; UTA R-1235: Val Verde, Devil's River, 23.5 mi N Comstock; UTA R-28907: Val Verde, 8.7 km N Baker's Crossing on St Hwy 163; UTA R-1599: Williamson, Georgetown Country Club; UTA R-8633-37, 8650-52: Wise, 4.0 mi NNW Decatur; UTA R-10010-12: Wise, LBJ National Grassland; UTA R-14726, 15356-57: Wise, 8.0 Km N Decatur, L.B. Johnson National Grassland; UTA R-26549: Wise, 0.3 air km SW Flat Rock Cemetery; UTA R-31419: Wise, 1.6-2.0 air mi S Flat Rock Cemetery, 0.4 air mi W old Decatur Road; UTA R-31420-423: Wise, 3.0 mi N Runaway Bay, Sid Richardson Scout Ranch; UTA R-32459-462: Wise, Sid Richardson Scout Ranch; UTA R-10008: Zapata, 3.7 mi NE jct. US 83 on FM 3169; Mexico, Hidalgo: MZFC-8022: Metztlán, Metztlán; Querétaro: MZFC-8482: Arroyo Seco, N Concá-Santa María River Bridge; MZFC-6257–6258: Mesa de León; MZFC-6259: Rancho Nuevo; MZFC-6260: Nopalera Boye; CM-90296: San Juan Del Río, 2.2 mi S.; Zacatecas: CM-59982: 16 mi NE Concepción Del Oro turnoff on Mex Hwy 54.

*Rena dugesii*: Mexico, Colima: MNHN-RA-0.1651.

*Rena dulcis iversoni*: Mexico, Tamaulipas: BU-MMC 11522: 20.9 km NE Ignacio Zaragoza, Río Guayalejo (formerly SM 11522); BU-MMC 11560: 1.6 km N Llera; BU-MMC 15094-15096 1.6 km E Llera (all paratypes, all formerly BCB collection).

*Rena humilis tenuicula*: Mexico, San Luis Potosí: MCZ-R 4519: San Luis Potosí (Holotype).

*Rena maxima*: Mexico, Guerrero: MZFC-03826: Ixcateopan de Cuauhtémoc, Ixcateopan; Morelos: MZFC-01931: Mazatepec, Mazatepec; Puebla: MZFC-04710: Zapotitlán de las Salinas, Zapotitlán de las Salinas; UTA R-12227: 9 km SSW Zapotitlan Salinas; UTA R-12229: 5.6 km SSW Zapotitlán Salinas; UTA R-14533: vicinity of Zapotitlán Salinas.

*Rena myopica*: Mexico, Hidalgo: MZFC-7793: 2 km W Arroyo Blanco; Nuevo León: UTA R-3149: Monterrey; Tamaulipas: UTA R-54555: Carretera Estación Calles-Hacienda Acuña, 583 m, 23.18565 N 98.49051 W.

*Rena rubella*: Mexico, Coahuila: MCZ-R 4584: San Pedro (Holotype).

*Rena segregata*: USA, Texas: UTA R-17767: Brewster, W side Black Hill, ca. 15 km E, ca. 2 km N jct Dove Mountain Road, and US Hwy 385; UTA R-33761: Brewster, 100 yds. W of Big Bend National Park entrance on TX 118; UTA R-2881: Hudspeth, Eagle Mtns., 7 mi S Hot Wells, UTA R-33759: Presidio, 19.4 mi W of Lajitas on FM 170; UTA R-33760: Presidio, 26.7 mi W of Lajitas on FM 170; UTA R-54619: Terrel, TX 349, 4.2 miles N of Dryden; UTA R-45091: Ward, On Hwy 329, ca. 4.8 km E Grandfalls.

## References

- Adalsteinsson, S. A., Branch, W. R., Trape, S., Vitt, L. J., & Hedges, S. B. (2009). Molecular phylogeny, classification, and biogeography of snakes of the Family Leptotyphlopidae (Reptilia, Squamata). *Zootaxa*, 2244, 1–50. <https://doi.org/10.11646/zootaxa.2244.1.1>
- Baird, S. F., & Girard, C. (1853). *Catalogue of North American reptiles in the Museum of the Smithsonian Institution*. Part I - Serpents. Washington D.C., Government Printing Office. <https://doi.org/10.5962/bhl.title.5513>
- Brown, B. C., & Brown, L. M. (1967). Notable records of Tamaulipan snakes. *Texas Journal of Science*, 19, 323–326.
- Castoe, T. A., Spencer, C. L., & Parkinson, C. L. (2007). Phylogeographic structure and historical demography of the western diamondback rattlesnake (*Crotalus atrox*): A perspective on North American desert biogeography. *Molecular Phylogenetics and Evolution*, 42, 193–212. <https://doi.org/10.1016/j.ympev.2006.07.002>
- Dixon, J. R., & Vaughan, R. K. (2003). The status of Mexican and southern United States blind snakes allied with *Leptotyphlops dulcis* (Serpentes: Leptotyphlopidae). *Texas Journal of Science*, 55, 3–24.
- Dugès, A. (1890). Fauna del estado de Guajuato. In A. L. Velasco (Ed.), *Geografía y estadística del estado de Guajuato*. *Geografía y estadística de la República Mexicana*, Vol. 5 (pp. 287–295). Ciudad de México: Oficina Tipográfica de la Secretaría de Fomento.
- Dugès, A. (1895). Fauna del estado de Guajuato. In *Memoria sobre la administración pública del estado de Guajuato presentada al congreso del mismo por el C. Gobernador constitucional Lic. Joaquín Obregón González*, el 1 de abril de 1895. Morelia: Imprenta y litografía de la Escuela IM Porfirio Díaz.
- Garman, S. (1883). The reptiles and batrachians of North American reptiles, part 1, Ophidia. *Memoirs of the Museum of Comparative Zoology Harvard*, 8, 1–185. <https://doi.org/10.5962/bhl.title.50435>
- Hahn, D. E. (1979a). *Leptotyphlops dulcis* (Baird & Girard) Texas blind snake. *Catalogue of American Amphibians and Reptiles*, 231, 1–2.
- Hahn, D. E. (1979b). *Leptotyphlops humilis* (Baird & Girard) Western blind snake. *Catalogue of American Amphibians and Reptiles*, 232, 1–4.
- Hahn, D. E. (1980a). *Leptotyphlops maximus* Loveridge. *Catalogue of American Amphibians and Reptiles*, 244, 1.
- Hahn, D. E. (1980b). *Liste der rezenten Amphibien und Reptilien Anomalepididae, Leptotyphlopidae, Typhlopidae*. Das Tierreich, eine Zusammenstellung und Kennzeichnung der rezenten tierformen. Berlin: Walter de Gruyter.
- HerpNet database. Accessed 24 October 2009. <http://www.herpnet.org>
- Klauber, L. M. (1940). The worm snakes of the genus *Leptotyphlops* in the United States and Northern Mexico. *Transactions of the San Diego Society of Natural History*, 9, 87–162.
- Lemos-Espinal, J. A., & Smith, H. M. (2007). *Anfibios y reptiles del estado de Chihuahua, México*. México D.F.: UNAM/Conabio.
- Lemos-Espinal, J. A., Smith, H. M., Chiszar, D., & Woolrich-Piña, G. (2004). 2003. Snakes from Chihuahua and adjacent states of Mexico. *Bulletin of the Chicago Herpetological Society*, 39, 206–213.
- McDiarmid, R. W., Campbell, J. A., & Touré, T. A. (1999). *Snake species of the world, a taxonomic and geographic reference, Vol. 1*. Washington, DC: The Herpetologists League.
- Morafka, D. J. (1977). *A biogeographical analysis of the Chihuahuan Desert through its herpetofauna*. Dordrecht: Springer. <https://doi.org/10.1007/978-94-010-1318-5>
- Myers, E. A., Burgoon, J. L., Ray, J. M., Martínez-Gómez, J. E., Matías-Ferrer, N., Mulcahy, D. G. et al. (2017). Coalescent species tree inference of *Coluber* and *Masticophis*. *Copeia*, 105, 640–648. <https://doi.org/10.1643/CH-16-552>
- Peterson, H. W., Smith, H. M., & Chiszar, D. (1995). Some noteworthy amphibians and reptiles from the region of Chapala, Jalisco, Mexico. *Bulletin of the Chicago Herpetological Society*, 30, 90–91.
- Pinto, R. R. (2010). *Revisão Sistemática da Subtribo Reni (Serpentes: Leptotyphlopidae) (PhD. Thesis)*. Universidade Federal do Rio de Janeiro, Brazil.
- Ponce-Campos, P., Huerta-Ortega, S. M., Nogueira-Gonzalez, C., & Smith, H. M. (2001). Natural history notes on the Southern Plateau night lizard, *Xantusia sanchezi*. *Bulletin of the Maryland Herpetological Society*, 37, 18–21.
- Provost, K. L., Mauck III, W. M., & Smith, B. T. (2018). Genomic divergence in allopatric Northern Cardinals of the North American warm deserts is linked to behavioral differentiation. *Ecology and Evolution*, 8, 12456–12478. <https://doi.org/10.1002/ece3.4596>
- Sabaj, M. H. (2016). Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an online reference. Version 6.5 (16 August 2016). American

- Society of Ichthyologists and Herpetologists, Washington, DC. Electronically accessible at <http://www.asih.org/>
- Smith, H. M., & Chiszar, D. (1993). Apparent intergradation in Texas between the subspecies of the Texas blind snake (*Leptotyphlops dulcis*). *Bulletin of the Maryland Herpetological Society*, 29, 143–155.
- Smith, H. M., van Breukelen, F., Auth, D. L., & Chiszar, D. (1998). A subspecies of the Texas blind snake (*Leptotyphlops dulcis*) without supraoculars. *The Southwestern Naturalist*, 43, 437–440.
- Stejneger, L. (1891). Notes on some North American snakes. *Proceedings of the United States National Museum*, 14, 501–505. <https://doi.org/10.5479/si.00963801.876.501>
- Uetz, P., Freed, P., & Hošek, J. (Eds.). (2022). The Reptile Database. Accessed January, 10, 2022. <http://www.reptile-database.org>
- Wallach, V. (2003). *Scolecophidia miscellanea. Hamadryad*, 27, 227–245.
- Wallach, V., Williams, K. L., & Boundy, J. (2014). *Snakes of the world, a catalogue of living and extinct species*. Boca Raton, Florida: CRC Press. <https://doi.org/10.1086/679952>