



The effect of island type on lizard reproductive traits

Maria Novosolov* and Shai Meiri

Department of Zoology, Tel Aviv University,
6997801, Tel Aviv, Israel

ABSTRACT

Aim The origins of islands influence island colonization and radiation dynamics, thus exerting differential selection pressures on the species that inhabit them. The occurrence of lower numbers of predator and competitor species on islands than the mainland selects for ‘slow’ life-history attributes (the ‘island syndrome’). Animals colonizing, and radiating on, oceanic islands probably face more novel environments than do those inhabiting continental fragment and land-bridge islands. We hypothesized that oceanic island endemics will show the slowest life histories, whereas land-bridge island species will resemble mainland species the most. We predicted that species on old, small and isolated islands will also have slow life histories.

Location World-wide.

Methods We assembled life-history data for 540 mainland and 319 insular endemic lizard species. We tested whether clutch size, brood frequency, hatchling mass and productivity differed between islands of different origin and between islands and the mainland. We controlled for female size, for latitude and for phylogenetic relationship using the R package CAPER. In addition, we tested the influences of island age, area and isolation on species life histories.

Results Oceanic island endemics have the smallest clutches and the largest offspring, and, together with continental fragment island endemics, lay most frequently. Clutch size, brood frequency and productivity increase with increasing island age. Isolation and area have little effect on lizard life history.

Main conclusions Our findings support the proposition that selection pressure differs across island type. The predator-poor environments on oceanic islands select for few, large offspring, while the predator-rich environments of the mainland and land-bridge islands select for many, small offspring. Island geological origin creates the environment within which evolution takes place, and thus plays a major role in life-history evolution. As islands grow older, lizards adapt by increasing their yearly reproductive effort.

Keywords

Evolution, island biogeography, island endemism, island syndrome, island type, life history, lizards, reproduction.

*Correspondence: Maria Novosolov,
Department of Zoology, Tel Aviv University,
6997801, Tel Aviv, Israel.
E-mail: marianovosolov@gmail.com

INTRODUCTION

Darwin (1845) and Wallace (1902) described the unique faunas and floras that can be found on islands, and these island systems have been essential in understanding evolutionary and ecological processes ever since (MacArthur & Wilson, 1967; Whittaker & Fernández-Palacios, 2007; Lomolino *et al.*,

2010). Islands contain a high proportion of endemic species and are considered excellent ‘natural laboratories’ for evolutionary research (MacArthur & Wilson, 1967; Mayr, 1967; Whittaker & Fernández-Palacios, 2007). The isolation of an island and the differences between mainland and insular environments can lead to founder effects, drift and strong selection to adapt to a novel environment (Thomas *et al.*,

2009). Such processes will often lead to speciation and diversification (Blondel, 2000; Schluter, 2001). Islands contain fewer species than mainland regions of equivalent area (Whittaker & Fernández-Palacios, 2007; Foufopoulos *et al.*, 2011). They thus usually have fewer predator and competitor species, which may decrease interspecific competition and predation risk (but see Meiri *et al.*, 2007). This in turn can select for the expansion of species niches through adaptive radiation to include resources that are used by other species on the mainland (Wallace, 1902; Van Valen, 1965; MacArthur & Wilson, 1967; Whittaker & Fernández-Palacios, 2007; Losos & Ricklefs, 2009, 2010; cf. Meiri *et al.*, 2005a).

Adler & Levins (1994) described changes in reproduction and behaviour between mainland and insular populations of rodents and named this phenomenon the 'island syndrome'. According to this model, insular animals evolve increased body size and lower aggression, attain higher population densities, and shift their reproductive traits towards the slow end of the fast–slow continuum. Thus they evolve larger offspring, smaller clutches and litters, and low productivity rates (Adler & Levins, 1994; Blondel, 2000; Goltsman *et al.*, 2005; Novosolov *et al.*, 2013). Although originally based on data from rodent populations, similar trait shifts have also been described in other insular taxa. Insular passerine bird populations, for example, have been found to have evolved to lay smaller clutches (Blondel, 2000), and insular populations of the arctic fox (*Vulpes lagopus*) have low fertility and dispersal ability (Goltsman *et al.*, 2005). Recently, we showed that lizards mostly follow the island syndrome (Novosolov *et al.*, 2013): insular endemic species have smaller clutches, higher brood frequency, larger offspring and denser populations than mainland species.

Islands of different geological origins may have different selective environments that can, in turn, affect the evolution of colonizing and *in situ* evolving species (Whittaker & Fernández-Palacios, 2007). There are three main geological types of islands: (1) oceanic islands – islands that were never connected to the mainland or to large islands, and are usually formed by volcanic activity; (2) land-bridge islands – islands that were connected to the mainland and became isolated when sea levels rose (some have experienced repeated episodes of connection and isolation), which for most islands occurred for the last time after the Last Glacial Maximum, between 12,000 and 6000 years ago (depending on the depth of the intervening channel and other factors); and (3) continental fragment islands – islands that are fragments of continental tectonic plates (Whittaker & Fernández-Palacios, 2007; and see below).

Oceanic islands usually lack large predators and have fewer species than other types of islands (Williamson, 1981). Having no species upon formation, the first propagules that arrive on these islands must adapt to a highly novel environment, and thus probably undergo great trait shifts (Mayr, 1967; Carlquist, 1974; Thomas *et al.*, 2009; Raia & Meiri, 2011). Because all species on such islands arrive by dispersal or radiate *in situ*, oceanic islands have unique faunas and

floras, very different from those of the mainland (Carlquist, 1974). Land-bridge and continental fragment islands start their insular 'life' with an existing ecological community, similar to that of the mainland that they were once part of (Karr, 1982). Species on such islands may therefore be relatively well adapted to the environment at the time of isolation and need only 'keep up' with the new selection pressures brought about when the insular community adapts to its isolated state (Wilcox, 1978). The degree of such adaptation, however, is likely to differ between continental fragment and land-bridge islands, as the former are usually millions of years old, and the latter have been isolated for only a few thousand years. Land-bridge islands usually have more predator, competitor and prey species than oceanic or continental fragment islands of similar size because they were recently isolated and are close to the landmass that they were once part of (McNab, 2002). Selection forces may therefore be more similar between the mainland and land-bridge islands than between the mainland and continental fragment islands (Watson, 2009). Biotas of continental fragment islands probably occupy an intermediate position along the spectrum between oceanic and land-bridge island biotas (Whittaker & Fernández-Palacios, 2007). Although continental fragment islands were once part of a mainland system, their environment and biota have been long separated and have been greatly influenced by the species that colonized them, and radiated on them, through time. This may result in traits of continental fragment island species being intermediate between those of land-bridge and oceanic islands (Carlquist, 1974). One should also bear in mind that mainland regions are dynamic, and their faunas have undergone substantial evolutionary changes after continental fragment islands became isolated from them (Gordon, 1986). Some of the differences between insular and mainland species thus probably stem from the evolution of the latter.

Island age, area and isolation are thought to affect the number and composition of species, and the evolution of traits (Meiri *et al.*, 2005b; Losos & Ricklefs, 2010). As islands age, their faunas go through evolutionary changes, often resulting in a greater degree of endemism. Thus it is expected that life history 'slows down' through the lapse of evolutionary time on islands (MacArthur & Wilson, 1967; Chapter 7, pp. 156–172). Because species richness decreases with increasing isolation, isolated island environments are thought to offer more opportunities for diversification and adaptation, resulting in a higher degree of endemism (Carlquist, 1974; Case & Cody, 1987; Rosindell & Phillimore, 2011). Island area is likely to have the opposite effect, with larger islands being more mainland-like in terms of the number of competitor, predator and prey species, population densities, etc.

Lizards have diversified greatly on islands across the planet. Because lizards, in general, show the adaptations predicted by the island syndrome (Novosolov *et al.*, 2013) they are an excellent model system in which to test the effects of island type, isolation, age and area. We hypothesize that

island type plays a fundamental role in the evolution of reproductive traits. Specifically, we make the following predictions: (1) Species on oceanic islands will manifest the island syndrome most strongly, and thus have smaller clutch sizes, higher brood frequencies and larger offspring than species on other types of islands or the mainland. (2) Species on land-bridge islands will most resemble species on the mainland in their reproductive traits – and have large clutch sizes, low brood frequencies, and small offspring. (3) Species on continental fragment islands will show intermediate traits. (4) The productivity rate (clutch size \times brood frequency \times offspring mass; Meiri *et al.*, 2012) will be similar across different types of islands because of a trade-off between its components. (5) Increasing island ages and isolation will enhance the expression of traits related to the island syndrome, whereas increasing island area will diminish their expression.

MATERIALS AND METHODS

We assembled a database that included 859 lizard species – 540 mainland species, 177 oceanic island endemics, 45 land-bridge island endemics and 97 species endemic to continental fragment islands (Table 1). We excluded species endemic to islands larger than 50,000 km² (e.g. Madagascar, New Guinea, Sulawesi) because such large islands often have many lizard predators and competitors, and their ecological settings therefore resemble those of the mainland (Badano *et al.*, 2005). We examined the primary literature, field guides and the reptile database (<http://reptile-database.reptarium.cz/>) to determine if species were found only on islands (insular endemics), or not (henceforth mainland species, even if they are found both on islands and on the mainland: we derived data for these 'mainland' species only from mainland populations), and to collect life-history data for clutch size, brood frequency and adult and offspring size (Table 2, and see Appendix S1 in Supporting Information). For each insular endemic species we recorded the type, age and distance from the nearest landmass (larger than 50,000 km²) of the largest island it inhabits. Island areas were obtained from the National Imagery and Mapping Agency (NIMA, 1997), the UN island directory (<http://islands.unep.ch/lindex.htm>), and the primary literature.

We classified island type as follows:

1. Oceanic islands – islands that formed over oceanic plates, or as a result of plate collision, and have never been connected to continental landmasses. These include islands of volcanic or uplift origin that were connected to a larger oceanic island (larger than the island in question but smaller than 50,000 km²). For example, the volcanic Bequia Island in the Grenadines was connected to the larger, volcanic island of St Vincent during periods of low sea levels (Carstensen *et al.*, 2012), and we therefore consider both islands to be oceanic.
2. Land-bridge islands – islands that were connected to a landmass larger than 50,000 km² during periods of low sea level (mostly at the end of the Pleistocene, e.g. Bali to Java).

We also classified islands that were connected to continental fragment islands as land-bridge islands (e.g. Roti Island, formed by rock and limestone uplift, was connected to Timor, which fragmented from the Australian plate; Voris, 2000; Carstensen *et al.*, 2012).

3. Continental fragment islands – islands that broke off from a larger landmass by tectonic activity and which were never subsequently connected to a landmass larger than 50,000 km² (e.g. Timor, see above).

Data for island type, isolation and age were collected from the *Encyclopedia of Islands* (Gillespie & Clague, 2009), from the primary literature, and from sources on the web (Appendix S1). For islands that were once connected to the mainland we used the age of isolation (hereafter 'island age'). Island age is frequently (for 38 of 143 islands in our data set) reported as a geological era. In such cases we defined age as the end of the era (e.g. Gillespie & Clague, 2009, give the Miocene as the age of the main rock in Martinique, and we therefore used 5 Ma as its age), because this is the only age at which the island certainly existed. We define isolation as the shortest distance between an island and a landmass larger than 50,000 km². We calculated isolation using Google Earth and the distance calculator tool in ArcGIS 9.3.1 (ESRI, Redlands, CA, USA).

We used family-specific equations to estimate the adult and offspring masses from mean snout–vent lengths (SVLs, Appendix S1). For *Liolaemus* and *Phymaturus* we used equations from Pincheira-Donoso *et al.* (2011). For gecko clades and for *Anolis* we used the equations in Novosolov *et al.* (2013). For legged anguids we used data from Meiri *et al.* (2013). For all other species we used family-specific equations from Meiri (2010), differentiating between legless, leg-reduced and legged species. For 87 species we had no data for female SVL and we therefore used species-specific SVLs. For seven species we used offspring mass rather than SVL, as SVLs were unavailable. We used mean clutch/litter sizes and frequencies where possible. Where more than one mean was reported for a species we used the midpoint of the range of means. Where means were unavailable, we used the midpoint of the reported trait range.

Meiri *et al.* (2012) have recently shown that lizard productivity is best quantified as a rate – biomass produced per unit time. We thus define productivity as the product of brood frequency, clutch size and offspring mass, in units of grams per year.

To correct for possible phylogenetic effects in the data, we assembled a composite species-level phylogeny from the literature, following the broad-scale squamate phylogenetic relationships reported by Wiens *et al.* (2010) and the taxonomy of the reptile database (<http://reptile-database.reptarium.cz/>, downloaded 17 January 2013). In assembling the tree we gave priority to recently published phylogenies that are based on nuclear DNA, then on mitochondrial DNA sequences. For species for which no genetic phylogeny was available, we relied on phylogenies based on morphological data. Where phylogenies were unresolved at the intrageneric rank we sunk

Table 1 Sample sizes of lizard species across different types of islands and the mainland in a global study.

Family	Oceanic islands	Land-bridge islands	Continental fragment islands	Mainland	Total no. of species
Agamidae	2	6	2	44	54
Anguidae	1	0	1	16	18
Anniellidae	0	0	0	1	1
Bipedidae	0	0	0	3	3
Carphodactylidae	0	0	0	5	5
Chamaeleonidae	2	0	2	11	15
Cordylidae	0	0	0	3	3
Corytophanidae	0	0	0	2	2
Crotaphytidae	0	0	0	5	5
Dactyloidae	23	1	7	12	43
Diplodactylidae	0	2	23	12	47
Eublepharidae	2	1	0	8	11
Gekkonidae	29	10	10	31	80
Gerrhosauridae	0	0	0	1	1
Gymnophthalmidae	0	1	0	5	6
Helodermatidae	0	0	0	2	2
Iguanidae	12	2	0	6	20
Lacertidae	13	14	2	88	117
Leiocephalidae	3	0	0	0	3
Leiosauridae	0	0	0	1	1
Liolaemidae	0	0	0	20	20
Phrynosomatidae	3	0	0	51	54
Phyllodactylidae	5	0	3	13	21
Polychrotidae	0	0	0	1	1
Pygopodidae	0	0	0	6	6
Scincidae	56	6	37	109	198
Sphaerodactylidae	6	1	7	7	21
Teiidae	8	0	2	35	45
Trogonophiidae	0	0	0	1	1
Tropiduridae	3	0	0	12	15
Varanidae	8	1	1	23	33
Xantusiidae	1	0	0	5	6
Xenosauridae	0	0	0	1	1
Total	177	45	97	540	859

Table 2 The numbers of lizard species for which data were available for various reproductive traits on different types of islands and the mainland in a global study.

Reproductive trait	Oceanic islands	Land-bridge islands	Continental fragment islands	Mainland	Total no. of species
Clutch size	156	42	94	540	832
Brood frequency	45	21	11	540	617
Offspring mass	111	21	38	540	710
Productivity	37	11	9	540	597
All species	177	45	97	540	859

species into a polytomy within their genus. The phylogenetic relationships between the species and the sources of phylogenetic data for each are given in Appendix S1.

The phylogenetic tree is based on many phylogenetic hypotheses, and differences between source trees prevent us from recording actual branch lengths. We therefore scaled branches to make the tree ultrametric using the cladogram transform in FIGTREE 1.3.1 (Rambaut, 2010). To account

for possible effects of climate we mapped the geographical ranges of the various species in ArcGIS 9.3.1 using published data on lizard distributions (Appendix S2). We used the absolute value of the latitudinal centroid for each species as a proxy for climate. Latitude was used to correct for the tendency of tropical species to produce smaller, and more frequent, clutches than temperate species (Meiri *et al.*, 2012).

Statistical analyses

All the data, except latitude, were \log_{10} -transformed in all analyses in order to normalize the residual distribution and reduce heteroscedasticity. When analysing the relationship between island type and reproductive traits we used the clutch size, number of yearly broods, offspring mass and productivity rate of each species as response variables and regressed them against female mass (g) and latitude as covariates and island type as a main effect. To examine the effect of island attributes on lizard reproductive traits we used a backward stepwise model selection. Because we were interested mainly in differences between island types we retained island type in the model even when it was non-significantly correlated with the life-history trait examined. We first analysed only insular species, to test the effects of island area, isolation and age. We then compared islands with the mainland without these terms.

All four sets of analyses were duplicated to account for phylogenetic non-independence by using phylogenetic generalized least squares (PGLS) regression (Freckleton *et al.*, 2002). We adjusted the strength of phylogenetic non-independence using the maximum likelihood value of the scaling parameter λ implemented in the R package CAPER (Orme *et al.*, 2012). Pagel's λ is a multiplier of the off-diagonal elements of the variance-covariance matrix, which provides the best fit of the Brownian motion model to the tip data.

Sensitivity analyses

We conducted several types of analyses excluding different groups of species from our database to better determine the extent of the adaptive nature of our results. Geckos and anoles lay fixed, small clutches of one or two eggs and usually have a high brood frequency. Therefore we excluded them from the database and repeated our analyses. We then repeated the analyses for geckos alone (our anole sample was not large enough to permit meaningful analysis). Because our tree included some genus-level polytomies, we repeated the phylogenetic analyses on a pruned tree from which these polytomies had been removed (see Appendix S3). We only report the results of these analyses in the text when they differ from the results obtained with the complete database.

RESULTS

The best models for each reproductive characteristic are reported in Appendix S3.

Clutch size

Species on oceanic islands have the smallest clutches, those on continental fragment islands have larger clutches, and land-bridge island endemics have the largest clutches. All two-way contrasts except the one between oceanic and continental frag-

ment islands were significant in both the non-phylogenetic and the phylogenetic model (at $\alpha \leq 0.05$; Fig. 1a, Table 3). Species on all types of islands have significantly smaller clutches than mainland species (Table 4). Excluding geckos and anoles from the analyses, the difference between species on continental fragment and land-bridge islands becomes non-significant (Appendix S3). On islands, clutch size increases with increasing island age (phylogenetic model: slope = 0.03 ± 0.01 , $t = 2.24$, $P = 0.02$, but note that in the non-phylogenetic model: slope = 0.03 ± 0.02 , $t = 1.29$, $P = 0.19$). Clutch size is independent of island area (phylogenetic model: slope = 0.0005 ± 0.01 , $t = 0.05$, $P = 0.95$; non-phylogenetic model: slope = 0.001 ± 0.01 , $t = 0.09$, $P = 0.92$) and isolation (phylogenetic model: slope = -0.01 ± 0.02 , $t = -0.56$, $P = 0.57$, but note that in the non-phylogenetic model: slope = 0.08 ± 0.02 , $t = 3.15$, $P = 0.001$).

Brood frequency

Brood frequency was not significantly different between different types of islands or the mainland. However, the non-phylogenetic model showed that brood frequency is higher on continental fragment islands than on land-bridge islands (Fig. 1b, Table 5). Mainland species have a lower brood frequency than continental fragment and oceanic island species, but a higher brood frequency than land-bridge species (Table 4). Brood frequency increases with island age (slope = 0.06 ± 0.03 , $t = 2.35$, $P = 0.02$; note that in the non-phylogenetic model: slope = 0.06 ± 0.04 , $t = 1.51$, $P = 0.13$). Isolation and area are not significant predictors of brood frequency (island isolation phylogenetic model: slope = 0.06 ± 0.03 , $t = 2.35$, $P = 0.25$; non-phylogenetic model: slope = 0.04 ± 0.06 , $t = 0.59$, $P = 0.55$; island area phylogenetic model: slope = 0.02 ± 0.02 , $t = 0.92$, $P = 0.36$; non-phylogenetic model: slope = 0.04 ± 0.03 , $t = 1.24$, $P = 0.22$).

Offspring mass

Species on oceanic and land-bridge islands have larger offspring than species on the mainland (Table 4). There were no significant differences in offspring mass between the different types of islands in the phylogenetic model. However, in the non-phylogenetic model, species on land-bridge islands have larger offspring than those on continental fragment islands (Fig. 1c, Table 6). This difference disappears when we exclude geckos and anoles from the analyses (Appendix S3). Area, age and isolation are uncorrelated with offspring mass (phylogenetic model island area: slope = -0.01 ± 0.02 , $t = -0.88$, $P = 0.37$; non-phylogenetic model: slope = -0.02 ± 0.02 , $t = -1.03$, $P = 0.30$; island age phylogenetic model: slope = -0.01 ± 0.02 , $t = -0.46$, $P = 0.64$; non-phylogenetic model: slope = -0.01 ± 0.02 , $t = -0.52$, $P = 0.61$; island isolation phylogenetic model: slope = -0.02 ± 0.04 , $t = -0.60$, $P = 0.54$; non-phylogenetic model: slope = -0.01 ± 0.03 , $t = -0.38$, $P = 0.7$).




Figure 1 Relationship between log (reproductive trait) and log (adult body mass, g) of lizards on oceanic (light grey squares, dashed line), land-bridge (black diamonds, two-dash line) and continental fragment (dark grey circles, solid line) islands and the mainland (open circles, dotted line). In cases of insignificance we provide only a regression line of the model. (a) Clutch size, (b) brood frequency, (c) offspring mass (g), (d) productivity. See tables for statistics and sample sizes.

Table 3 Clutch sizes of lizards on different types of islands and the mainland in a global study. The intercepts of each type of island are reported on the diagonal; $n = 292$. Above the diagonal: the non-phylogenetic model t -values and P -values for the relationships between the different types of islands (e.g. clutch size on oceanic islands is significantly smaller than clutch size on land-bridge islands, with a t -value of -3.72 and P -value of 0.002); $R^2 = 0.46$. Below the diagonal: the phylogenetic model t -values and P -values for the relationships between the different types of islands; $\lambda = 0.90$; $R^2 = 0.2$.

	Oceanic islands ($n = 156$)	Land-bridge islands ($n = 42$)	Continental fragment islands ($n = 94$)
Oceanic islands	Non-phylogenetic intercept: -0.18 ± 0.07 Phylogenetic intercept: 0.19 ± 0.03	$t = -3.72$ $P < 0.002$	$t = -0.92$ $P = 0.36$
Land-bridge islands	$t = 2.70$ $P = 0.007$	Non-phylogenetic intercept: -0.02 ± 0.06 Phylogenetic intercept: 0.32 ± 0.06	$t = 2.78$ $P = 0.005$
Continental fragment islands	$t = -0.79$ $P = 0.42$	$t = -2.66$ $P = 0.008$	Non-phylogenetic intercept: -0.16 ± 0.07 Phylogenetic intercept: 0.18 ± 0.11

Productivity

Productivity rates were similar across all island types and were not significantly different from the mainland (Fig. 1d, Table 7). Productivity rates increase with increasing island age but do not vary with island isolation or island area (phylogenetic mod-

els followed by non-phylogenetic ones – island age phylogenetic model: slope = 0.11 ± 0.03 , $t = 3.30$, $P = 0.001$; non-phylogenetic model: slope = 0.09 ± 0.04 , $t = 2.41$, $P = 0.02$; island isolation phylogenetic model: slope = 0.009 ± 0.08 , $t = 0.11$, $P = 0.91$; non-phylogenetic model: slope = 0.02 ± 0.09 , $t = 0.21$, $P = 0.84$; island area phylogenetic model: slope =

Table 4 Results for various reproductive traits in lizards on different types of islands compared with the mainland in a global study. The intercepts for mainland in the phylogenetic and non-phylogenetic models are shown. Each cell shows the value for the intercept of the island and the *t*-value and *P*-value of the island when compared with the mainland.

	Non-phylogenetic model				Phylogenetic model			
	Mainland intercept	Oceanic islands	Land-bridge islands	Continental fragment islands	Mainland intercept	Oceanic islands	Land-bridge islands	Continental fragment islands
Clutch size	0.22 ± 0.03	0.02 ± 0.02 <i>t</i> = -8.59 <i>P</i> < 0.002 <i>n</i> = 156	0.13 ± 0.04 <i>t</i> = -2.32 <i>P</i> = 0.02 <i>n</i> = 42	0.06 ± 0.04 <i>t</i> = -4.99 <i>P</i> < 0.002 <i>n</i> = 94	0.26 ± 0.09	0.14 ± 0.02 <i>t</i> = -6.37 <i>P</i> < 0.002 <i>n</i> = 156	0.19 ± 0.03 <i>t</i> = -2.68 <i>P</i> = 0.007 <i>n</i> = 42	0.12 ± 0.03 <i>t</i> = -4.79 <i>P</i> < 0.002 <i>n</i> = 94
Brood frequency	0.60 ± 0.03	0.74 ± 0.04 <i>t</i> = 3.62 <i>P</i> < 0.002 <i>n</i> = 45	0.56 ± 0.06 <i>t</i> = -0.73 <i>P</i> = 0.46 <i>n</i> = 21	0.85 ± 0.08 <i>t</i> = 3.26 <i>P</i> = 0.001 <i>n</i> = 11	0.46 ± 0.1	0.46 ± 0.03 <i>t</i> = -0.08 <i>P</i> = 0.93 <i>n</i> = 45	0.32 ± 0.04 <i>t</i> = -1.94 <i>P</i> = 0.052 <i>n</i> = 21	0.55 ± 0.06 <i>t</i> = 1.33 <i>P</i> = 0.18 <i>n</i> = 11
Offspring mass	-0.92 ± 0.01	-0.87 ± 0.02 <i>t</i> = 2.12 <i>P</i> = 0.03 <i>n</i> = 111	-0.79 ± 0.05 <i>t</i> = 2.55 <i>P</i> = 0.01 <i>n</i> = 21	-0.92 ± 0.04 <i>t</i> = 0.02 <i>P</i> = 0.97 <i>n</i> = 38	-0.87 ± 0.08	-0.79 ± 0.02 <i>t</i> = 3.37 <i>P</i> < 0.002 <i>n</i> = 111	-0.75 ± 0.04 <i>t</i> = 3.06 <i>P</i> = 0.002 <i>n</i> = 21	-0.81 ± 0.04 <i>t</i> = 1.55 <i>P</i> = 0.12 <i>n</i> = 38
Productivity	-0.24 ± 0.02	-0.19 ± 0.05 <i>t</i> = 1.00 <i>P</i> = 0.32 <i>n</i> = 37	-0.25 ± 0.1 <i>t</i> = -0.18 <i>P</i> = 0.85 <i>n</i> = 11	-0.14 ± 0.02 <i>t</i> = 0.93 <i>P</i> = 0.35 <i>n</i> = 9	-0.02 ± 0.01	-0.022 ± 0.005 <i>t</i> = -0.49 <i>P</i> = 0.62 <i>n</i> = 37	-0.023 ± 0.008 <i>t</i> = -0.38 <i>P</i> = 0.70 <i>n</i> = 11	-0.022 ± 0.009 <i>t</i> = -0.0003 <i>P</i> = 0.99 <i>n</i> = 9

Table 5 Brood frequencies of lizards on different types of islands and the mainland in a global study. The intercept of each type of island is reported on the diagonal. Above the diagonal: the non-phylogenetic model *t*-values and *P*-values for the relationships between the different types of islands (e.g. the brood frequency on oceanic islands is significantly larger than the brood frequency on land-bridge islands, with a *t*-value of 2.75 and *P*-value of 0.006); *n* = 77; *R*² = 0.5. Below the diagonal: the phylogenetic model *t*-values and *P*-values for the relationships between the different types of islands; *n* = 77; *λ* = 0.89; *R*² = 0.3.

	Oceanic islands (<i>n</i> = 45)	Land-bridge islands (<i>n</i> = 21)	Continental fragment islands (<i>n</i> = 11)
Oceanic islands	Non-phylogenetic intercept: 1.11 ± 0.09 Phylogenetic intercept: 0.70 ± 0.07	<i>t</i> = 1.89 <i>P</i> = 0.06	<i>t</i> = -0.69 <i>P</i> = 0.49
Land-bridge islands	<i>t</i> = 0.87 <i>P</i> = 0.38	Non-phylogenetic intercept: 0.96 ± 0.11 Phylogenetic intercept: 0.77 ± 0.09	<i>t</i> = -2.01 <i>P</i> = 0.05
Continental fragment islands	<i>t</i> = 1.37 <i>P</i> = 0.17	<i>t</i> = 0.42 <i>P</i> = 0.67	Non-phylogenetic intercept: 1.18 ± 0.13 Phylogenetic intercept: 0.80 ± 0.17

-0.014 ± 0.04, *t* = -0.37, *P* = 0.71; non-phylogenetic model: slope = -0.02 ± 0.05, *t* = -0.59, *P* = 0.55).

DISCUSSION

Reproductive traits vary across island types, mostly according to our predictions. Oceanic and continental fragment island endemic lizards lay smaller clutches but reproduce more frequently than do lizards on land-bridge islands and the mainland. However, the difference in brood frequency between different island types disappears when accounting for phylogenetic relationships. Offspring mass was similar across

island types, with oceanic and land-bridge island species having larger offspring than mainland species. Island age was positively correlated with brood frequency but, contrary to our prediction, it was also positively correlated with clutch size and productivity. Interestingly, island area and isolation were mostly uncorrelated with reproductive traits (Appendix S3). This suggests that species richness per se has little to do with the traits we examined. Furthermore, because clutch size differs between mainland and land-bridge island species, despite the short time span since land bridges were severed from the mainland, we argue that some life-history traits, such as clutch size, evolve quickly in response to insularity.

Table 6 Offspring masses of lizards on different types of islands and the mainland in a global study. The intercept of each type of island is reported on the diagonal; $n = 170$. Above the diagonal: the non-phylogenetic model t -values and P -values for the relationships between the different types of islands (e.g. the offspring mass on oceanic islands is not significantly different from the offspring mass on land-bridge islands, with a t -value of 1.45 and P -value of 0.14); $R^2 = 0.9$. Below the diagonal: the phylogenetic model t -values and P -values for the relationships between the different types of islands; $\lambda = 0.73$; $R^2 = 0.8$.

	Oceanic islands ($n = 111$)	Land-bridge islands ($n = 21$)	Continental fragment islands ($n = 38$)
Oceanic islands	Non-phylogenetic intercept: -0.89 ± 0.04 Phylogenetic intercept: -0.76 ± 0.05	$t = -1.63$ $P = 0.10$	$t = 1.25$ $P = 0.21$
Land-bridge islands	$t = 0.64$ $P = 0.52$	Non-phylogenetic intercept: -0.81 ± 0.06 Phylogenetic intercept: -0.74 ± 0.06	$t = 2.28$ $P = 0.02$
Continental fragment islands	$t = -0.84$ $P = 0.39$	$t = -1.15$ $P = 0.25$	Non-phylogenetic intercept: -0.94 ± 0.04 Phylogenetic intercept: -0.81 ± 0.09

Table 7 Productivity of lizards on different types of islands and the mainland in a global study. The intercept of each type of island is reported on the diagonal; $n = 51$. Above the diagonal: the non-phylogenetic model t -values and P -values for the relationships between the different types of islands (e.g. productivity on oceanic islands is not significantly different from the productivity on land-bridge islands, with a t -value of -1.58 and P -value of 0.11); $R^2 = 0.93$. Below the diagonal: the phylogenetic model t -values and P -values for the relationships between the different types of islands; $\lambda = 0.48$; $R^2 = 0.88$.

	Oceanic islands ($n = 33$)	Land-bridge islands ($n = 11$)	Continental fragment islands ($n = 9$)
Oceanic islands	Non-phylogenetic intercept: 0.21 ± 0.1 Phylogenetic intercept: 0.06 ± 0.09	$t = -1.58$ $P = 0.11$	$t = 0.32$ $P = 0.74$
Land-bridge islands	$t = 1.65$ $P = 0.10$	Non-phylogenetic intercept: 0.39 ± 0.14 Phylogenetic intercept: 0.23 ± 0.11	$t = 1.50$ $P = 0.13$
Continental fragment islands	$t = -0.07$ $P = 0.94$	$t = -1.43$ $P = 0.16$	Non-phylogenetic intercept: 0.18 ± 0.15 Phylogenetic intercept: 0.06 ± 0.17

Species endemic to oceanic islands differ most in their reproductive traits from mainland species, with species on land-bridge and continental fragment islands having intermediate trait values. Species on oceanic islands show the most extreme manifestation of the island syndrome, laying frequent, small clutches of large offspring compared with mainland species of the same size and at the same latitude. Investing in smaller clutches may allow oceanic island endemics to produce larger offspring, which may be better adapted to an insular environment with weak predation and interspecific competition pressures, but dense populations and strong intraspecific competition (MacArthur *et al.*, 1972; Williamson, 1981; Blondel, 2000; Pafilis *et al.*, 2009).

The reproductive traits of land-bridge island species most closely resemble those of mainland species. Most land-bridge islands are relatively young and were formed when sea levels rose at the end of the Pleistocene (Whittaker & Fernández-

Palacios, 2007: there are only four exceptions in our data set, see Appendix S1). The lizard populations stranded on these islands face relatively similar selective pressures to those acting on the mainland. Nevertheless, land-bridge island endemics do lay smaller clutches than mainland species (but larger than on oceanic and continental fragment islands), while their brood frequencies and offspring masses are similar to those of mainland species. These results remain valid even when we exclude species with set clutch sizes (i.e. anoles and geckos; Appendix S3). This may indicate that clutch size is one of the first attributes to be affected by the selection regime in isolated environments.

Ashmole (1963) hypothesized that island birds have smaller clutch sizes than mainland ones owing to a climatically more stable environment, which he proposed offers fewer resources during the year. Dabool (2013) empirically established that islands inhabited by lizards are climatically less

seasonal than mainland areas of similar latitudes. We found that clutch size is smaller on all types of islands when compared with the mainland and is the smallest on oceanic and continental fragment islands. We hypothesize that reduced climatic seasonality selects for lower clutch sizes. The climate on land-bridge islands may be intermediate, selecting for intermediate clutch sizes. Intriguingly, after decreasing upon insularity, lizard clutch size increases with island age. A possible explanation for this apparent contradiction may be that endemic species are descendants of mainland species that were selected for decreased clutch size by insular environments with lower resources. However, with time, species endemic to oceanic and continental fragment islands better adapt to the insular environment and are able to better utilize resources and increase their clutch size and reproduction. Island age is not always an accurate proxy for the age of the lineages that inhabit it (Heads, 2011). Inferences regarding the effects of island age on the evolutionary dynamics of such lineages should be made with caution.

Anoles (*Anolis*) are proportionally over-represented on oceanic islands. From our data set, 13% of the oceanic island endemics in our data are anoles, compared with 2% of the mainland species ($\chi^2 = 82.12$, $P < 0.002$). Similarly, geckos dominate continental fragment island faunas (46% geckos compared with 17% on the mainland; $\chi^2 = 43.08$, $P < 0.002$). All anoles and geckos lay small (one egg is the mode in anoles, usually one or two in geckos) and usually frequent clutches both on islands and on the mainland. Without accounting for phylogeny, the high relative richness of insular anoles and geckos results in higher clutch frequencies on islands. Our phylogenetic analyses, however, reveal no differences between the brood frequencies of mainland and island taxa. This need not mean that high brood frequencies on islands are not adaptive. It may be that these attributes contribute to the successful colonization and high speciation rates of insular anoles and geckos (i.e. that these attributes differ between islands and the mainland owing to sorting rather than to evolution *in situ*). Brood frequency of geckos is higher on oceanic and continental fragment islands than on the mainland (intercepts: mainland 0.69 ± 0.07 ; continental fragment 1.03 ± 0.13 , $t = 2.70$, $P = 0.007$; oceanic 0.98 ± 0.11 , $t = 2.83$, $P = 0.005$; corrected for adult mass and latitude; we did not have enough data for brood frequency of geckos on land-bridge islands). No such difference was found in lizard taxa with variable clutch sizes (phylogenetic intercept: mainland 0.31 ± 0.09 ; land-bridge 0.26 ± 0.04 , $t = -1.32$, $P = 0.18$; continental fragment islands 0.26 ± 0.07 , $t = -0.70$, $P = 0.48$; oceanic islands 0.29 ± 0.03 , $t = -0.62$, $P = 0.54$; $\lambda = 0.72$; corrected for adult mass and latitude).

Wang *et al.* (2011) showed that brood frequency plays a role in the trade-off between offspring size and number in lizards. They showed that, after the clutch size is set, females of some species direct their energy towards increasing offspring mass, while others direct it towards another brood. Although islands are postulated to have relatively few resources (MacArthur & Wilson, 1967; cf. Meiri & Raia, 2010) they may also

be characterized by decreased climatic seasonality, which increases the length of the reproductive season (Ashmole, 1963; Hanski, 1977; Williamson, 1981; Cronk, 1997). In addition, continental fragment and oceanic islands are characterized by low interspecific competition and predation (Carlquist, 1974). Based on the results of Wang *et al.* (2011) we suggest that the ability to increase the brood frequency is a compensation mechanism developed only in species with a small clutch size. Thus, geckos may adapt to the insular environment by increasing their brood frequency, while other lizard taxa adapt by increasing their offspring mass (Appendix S3). The relative success (i.e. high relative species richness) of geckos on continental fragment islands and of anoles on oceanic islands may be a result of the adaptive advantage of high brood frequency and set clutch size on these types of islands. This advantage may be expressed by the ability of these species to use the longer reproductive season found on islands by increasing their brood frequency. Alternatively, the relative richness of members of such clades on islands may be attributed to factors that are unrelated to the pace of their lives (e.g. to egg-gluing in geckos).

Lack (1971) hypothesized that a uniform climate selects for small clutch sizes because food supplies are limited throughout the year. However, extremely seasonal climates are usually characterized by a season of high abundance (Geist, 1987), when clutches can safely be large. Latitude is a useful proxy of temperature seasonality, and our mainland sample generally stems from higher latitudes. We also found that, as expected, clutch size increases with latitude, and brood frequency decreases (Appendix S3). However, our findings of small clutch sizes and (especially in geckos) high brood frequencies, on oceanic and continental fragment islands compared with their relatives on the mainland, hold when latitudinal effects are accounted for (Appendix S3).

Insular lizard populations usually have high densities (Novosolov *et al.*, 2013), which can lead to strong intraspecific competition (MacArthur & Wilson, 1967; Pafilis *et al.*, 2009). This in turn could select for infrequent, small clutches of larger offspring, resulting in lower productivity rates (Andrews, 1979; Pafilis *et al.*, 2011). However, we did not find differences in productivity rates in lizards endemic to different types of islands and between insular and mainland species. These results, coupled to our finding that insular species have smaller clutches, are compatible with the hypotheses of Ashmole (1963) and Lack (1971) that less seasonal climates select for small clutch sizes. Food availability is often thought to be lower on islands than on the mainland (McNab, 1994a,b). Our results (similar yearly productivity), however, suggest that this is not the case. We suggest that overall similar amounts of resources are more evenly available through the year on islands. Thus species on islands probably adapt to this environment by dividing their energy more evenly than mainland species owing to a less seasonal climate or because of low predation and interspecific competition rates. Mainland species, living in a more seasonal environment, may have a temporally more peaked resource

abundance, and can enlarge their clutch size to match the food peak but can produce fewer clutches. This results in similar overall productivity rates.

We have shown that much of the variation in the evolution of clutch sizes, frequencies and their trade-off in insular endemic lizards can be attributed to island type. We therefore infer that species evolving on islands of different types are subjected to different selective pressures. Species on oceanic islands evolve 'slow' life-history attributes, while land-bridge island species are 'faster', but still slower than mainland species. Furthermore, our results indicate that the rate of reproduction of lizards on islands increases with island age (as defined above), suggesting a possible compensation of insular endemic species for the adaptations in reproductive traits made by their ancestors.

ACKNOWLEDGEMENTS

Erez Maza and members of the Global Assessment of Reptile Distribution (GARD) group were instrumental in obtaining data on lizard distributions. We thank three anonymous referees, the editor Kostas Triantis, and Jonathan Belmaker, Ofer Ovadia and, especially, Pasquale Raia for constructive comments on an earlier version of this work. We thank Salvador Carranza for valuable discussion. This study is funded by ISF grant number 1005/12 to S.M.

REFERENCES

- Adler, G.H. & Levins, R. (1994) The island syndrome in rodent populations. *Quarterly Review of Biology*, **69**, 473–490.
- Andrews, R.M. (1979) Evolution of life histories: a comparison of *Anolis* lizards from matched island and mainland habitats. *Breviora*, **454**, 1–51.
- Ashmole, N.P. (1963) The regulation of numbers of tropical oceanic birds. *Ibis*, **103b**, 458–473.
- Badano, E.I., Regidor, H.A., Núñez, H.A., Acosta, R. & Gianoli, E. (2005) Species richness and structure of ant communities in a dynamic archipelago: effects of island area and age. *Journal of Biogeography*, **32**, 221–227.
- Blondel, J. (2000) Evolution and ecology of birds on islands: trends and perspectives. *Vie et Milieu*, **50**, 205–220.
- Carlquist, S. (1974) *Island biology*. Columbia University Press, New York.
- Carstensen, D.W., Dalsgaard, B., Svenning, J.-C., Rahbek, C., Fjeldså, J., Sutherland, W.J. & Olesen, J.M. (2012) Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. *Journal of Biogeography*, **39**, 739–749.
- Case, T.J. & Cody, M.L. (1987) Testing theories of island biogeography. *American Scientist*, **75**, 402–411.
- Cronk, Q.C.B. (1997) Islands: stability, diversity, conservation. *Biodiversity and Conservation*, **6**, 477–493.
- Dabool, L. (2013) *Reproductive strategies of reptiles – a macroecological perspective*. MSc Thesis, Tel Aviv University, Tel Aviv.
- Darwin, C. (1845) *The voyage of the Beagle*. Smith, Elder and Co., London.
- Foufopoulos, J., Kilpatrick, A.M. & Ives, A.R. (2011) Climate change and elevated extinction rates of reptiles from Mediterranean islands. *The American Naturalist*, **177**, 119–129.
- Freckleton, R.P., Harvey, P.H. & Pagel, M. (2002) Phylogenetic analysis and comparative data: a test and review of evidence. *The American Naturalist*, **160**, 712–726.
- Geist, V. (1987) Bergmann's rule is invalid. *Canadian Journal of Zoology*, **65**, 1035–1038.
- Gillespie, R.G. & Clague, D.A. (eds) (2009) *Encyclopedia of islands*. University of California Press, Berkeley, CA.
- Goltsman, M., Kruchenkova, E.P., Sergeev, S., Volodin, I. & Macdonald, D.W. (2005) 'Island syndrome' in a population of Arctic foxes (*Alopex lagopus*) from Mednyi Island. *Journal of Zoology*, **267**, 405–418.
- Gordon, K.R. (1986) Insular evolutionary body size trends in *Ursus*. *Journal of Mammalogy*, **67**, 395–399.
- Hanski, I. (1977) Biogeography and ecology of carrion flies in the Canary Islands. *Annales Entomologici Fennici*, **43**, 101–107.
- Heads, M. (2011) Old taxa on young islands: a critique of the use of island age to date island-endemic clades and calibrate phylogenies. *Systematic Biology*, **60**, 204–218.
- Karr, J.R. (1982) Population variability and extinction in the avifauna of a tropical land-bridge island. *Ecology*, **63**, 1975–1978.
- Lack, D. (1971) Island birds. *Adaptive aspects of insular evolution* (ed. by W.L. Stern), pp. 29–31. Washington State University Press, Pullman, WA.
- Lomolino, M.V., Brown, J.H. & Sax, D.F. (2010) Island biogeography theory. *The theory of island biogeography revisited* (ed. by J.B. Losos and R.E. Ricklefs), pp. 13–51. Princeton University Press, Princeton, NJ.
- Losos, J.B. & Ricklefs, R.E. (2009) Adaptation and diversification on islands. *Nature*, **457**, 830–836.
- Losos, J.B. & Ricklefs, R.E. (eds) (2010) *The theory of island biogeography revisited*. Princeton University Press, Princeton, NJ.
- MacArthur, R.H. & Wilson, E.O. (1967) *The theory of island biogeography*. Princeton University Press, Princeton, NJ.
- MacArthur, R.H., Diamond, J.M. & Karr, J.R. (1972) Density compensation in island faunas. *Ecology*, **53**, 330–342.
- Mayr, E. (1967) The challenge of island faunas. *Australian Natural History*, **15**, 369–374.
- McNab, B.K. (1994a) Energy conservation and the evolution of flightless in birds. *The American Naturalist*, **144**, 628–642.
- McNab, B.K. (1994b) Resource use and the survival of land and freshwater vertebrates on oceanic islands. *The American Naturalist*, **144**, 643–660.
- McNab, B.K. (2002) *The physiological ecology of vertebrates: a view from energetics*. Cornell University Press, Ithaca, NY.
- Meiri, S. (2010) Length–weight allometries in lizards. *Journal of Zoology*, **281**, 218–226.

- Meiri, S. & Raia, P. (2010) Reptilian all the way? *Proceedings of the National Academy of Sciences USA*, **107**, E27.
- Meiri, S., Dayan, T. & Simberloff, D. (2005a) Variability and sexual size dimorphism in carnivores: testing the niche variation hypothesis. *Ecology*, **86**, 1432–1440.
- Meiri, S., Dayan, T. & Simberloff, D. (2005b) Insular carnivore biogeography: island area and mammalian optimal body size. *The American Naturalist*, **165**, 505–514.
- Meiri, S., Dayan, T. & Simberloff, D. (2007) Guild composition and mustelid morphology – character displacement but no character release. *Journal of Biogeography*, **34**, 2148–2158.
- Meiri, S., Brown, J.H. & Sibly, R.M. (2012) The ecology of lizard reproductive output. *Global Ecology and Biogeography*, **21**, 592–602.
- Meiri, S., Bauer, A., Chirio, L., Colli, G., Das, I., Doan, T., Feldman, A., Herrera, F.-C., Novosolov, M., Pafilis, P., Pincheira-Donoso, D., Powney, G., Torres-Carvajal, O., Uetz, P. & Van Damme, R. (2013) Are lizards feeling the heat? A tale of ecology and evolution under two temperatures. *Global Ecology and Biogeography*, **22**, 834–845.
- NIMA (1997) *National Imagery and Mapping Agency (NIMA), Vector Map Level 0 (VMAP0)*, 4th edn. NIMA, Washington, DC.
- Novosolov, M., Raia, P. & Meiri, S. (2013) The island syndrome in lizards. *Global Ecology and Biogeography*, **22**, 184–191.
- Orme, C.D.L., Freckleton, R.P., Thomas, G.H., Petzoldt, T., Fritz, S.A., Isaac, N.J.B. & Pearse, W.D. (2012) *caper: comparative analyses of phylogenetics and evolution in R*. R package version 0.5. Available at: <http://CRAN.R-project.org/package=caper/>.
- Pafilis, P., Meiri, S., Fougopoulos, J. & Valakos, E. (2009) Intraspecific competition and high food availability are associated with insular gigantism in a lizard. *Naturwissenschaften*, **96**, 1107–1113.
- Pafilis, P., Fougopoulos, J., Sagonas, K., Runemark, A., Svensson, E. & Valakos, E.D. (2011) Reproductive biology of insular reptiles: marine subsidies modulate expression of the ‘island syndrome’. *Copeia*, **2011**, 545–552.
- Pincheira-Donoso, D., Fox, S.F., Scolaro, J.A., Ibarquengoytia, N., Acosta, J.C., Corbalán, V., Medina, M., Boretto, J., Villavicencio, H.J. & Hodgson, D.J. (2011) Body size dimensions in lizard ecological and evolutionary research: exploring the predictive power of mass estimation equations in two Liolaemidae radiations. *Herpetological Journal*, **21**, 35–42.
- Raia, P. & Meiri, S. (2011) The tempo and mode of evolution: body sizes of island mammals. *Evolution*, **65**, 1927–1934.
- Rambaut, A. (2010) *FigTree, version 1.3.1*. Institute of Evolutionary Biology, University of Edinburgh. Available at: <http://tree.bio.ed.ac.uk/software/figtree/>.
- Rosindell, J. & Phillimore, A.B. (2011) A unified model of island biogeography sheds light on the zone of radiation. *Ecology Letters*, **14**, 552–560.
- Schluter, D. (2001) Ecology and the origin of species. *Trends in Ecology and Evolution*, **16**, 372–380.
- Thomas, G.H., Meiri, S. & Phillimore, A.B. (2009) Body size diversification in *Anolis*: novel environment and island effects. *Evolution*, **63**, 2017–2030.
- Van Valen, L. (1965) Morphological variation and width of ecological niche. *The American Naturalist*, **99**, 377–390.
- Voris, H.K. (2000) Maps of Pleistocene sea levels in South-east Asia: shorelines, river systems and time durations. *Journal of Biogeography*, **27**, 1153–1167.
- Wallace, A.R. (1902) *Island life*, 3rd edn. Macmillan, London.
- Wang, Z., Xia, Y. & Ji, X. (2011) Clutch frequency affects the offspring size–number trade-off in lizards. *PLoS ONE*, **6**, e16585.
- Watson, M.D. (2009) Continental islands. *Encyclopedia of islands* (ed. by R.G. Gillespie and D.A. Clague), pp. 180–187. University of California Press, Berkeley, CA.
- Whittaker, R.J. & Fernández-Palacios, J.M. (2007) *Island biogeography: ecology, evolution, and conservation*, 2nd edn. Oxford University Press, Oxford.
- Wiens, J.J., Kuczynski, C.A., Townsend, T., Reeder, T.W., Mulcahy, D.G. & Sites, J.W. (2010) Combining phylogenomics and fossils in higher-level squamate reptile phylogeny: molecular data change the placement of fossil taxa. *Systematic Biology*, **59**, 674–688.
- Wilcox, B.A. (1978) Supersaturated island faunas: a species–age relationship for lizards on post-Pleistocene land-bridge islands. *Science*, **199**, 996–998.
- Williamson, M.H. (1981) *Island populations*. Oxford University Press, Oxford.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Appendix S1 The data set and metadata (literature source), mass equations and phylogenetic relationships used for the analysis of lizard reproductive traits.

Appendix S2 Reference list for species mapping sources of species used in the analysis.

Appendix S3 Results of sensitivity analyses and the best model for each life-history trait.

BIOSKETCHES

Maria Novosolov is a PhD student studying the evolution of life-history traits in island endemic reptiles.

Shai Meiri studies the biogeography of animal traits in different vertebrate clades, evolutionary responses to insularity, and the patterns, drivers and consequences of the global distribution of reptiles.

Editor: Kostas Triantis

SUPPORTING INFORMATION

The effect of island type on lizard reproductive traits

Maria Novosolov and Shai Meiri

APPENDIX S1 The dataset, the mass equations, the phylogeny used in the analyses and the metadata (literature source).

- A. Species data
- B. Data of island attributes
- C. The distribution in island area, age and isolation across different island types
- D. Literature sources used to create the datasets and to determine the distribution of the species
- E. Equations used to estimate (\log_{10}) mass from (\log_{10}) snout–vent lengths in clades
- F. The phylogenetic tree
- G. The reference used for each species in the phylogenetic tree
- H. Literature source for the phylogenetic tree

A. Data that were used for the analyses.

Adult and hatchling mass are in grams.

Clutch size (eggs per clutch/neonates per litter) is the midpoint of the highest and lowest reported averages. When data on averages were unavailable clutch size is the mean of the largest and smallest reported clutches.

Brood frequency is calculated as the midpoint of the highest and lowest reported number of clutches or litters a female lays/give birth to in a year.

Productivity is hatchling mass*clutch size*brood frequency (units: $\text{gram} \cdot \text{year}^{-1}$).

Island area (in km^2) is the area of the largest island an insular endemic species is found on.

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Acanthocercus atricollis</i>	no	NA	NA	NA	mainland	-15	39.32	11.36	1	0.78	8.87	Spawls et al. 2002, Fitzsimons 1943, Reaney and Whiting 2002, Loveridge 1942, Branch 1998, Parker 1936, Jeffery 1993, Schmidt et al. 1919, Auerbach 1987, Manthey and Schuster 1996, Branch 2005, Lagen and Spawls 2006, Vonesh 1998, Barts 2003, Kohler 2005, Lagen and Spawls 2010, Pienaar 1966, Haagner et al. 2000, Curry-Lindahl 1979, Jacobsen 1982 Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1982, Spawls et al. 2002, Perry and Garland 2002, Loveridge 1936, Razzetti and Msuya 2002, Greenbaum and Carr 2005, Hughes 1988, Bohme et al. 1996, Schmidt et al. 1919, Rogner 1997a, Le Berre 1989, Werner 1908, Anderson 1898, Manthey and Schuster 1996, Lonnberg 1911, Branch 2005, Chirio and LeBreton 2007, Dunham et al. 1988, Leache et al. 2006, Clusella-Trullas et al. 2008, Zug 1987, Fitch 1982, Pauwels and Vande weghe 2008, Radder et al. 2008, Koul and Duda 1977, Sinervo et al. 2010, Kohler 2005, Lagen and Spawls 2010, Jackson and Blackburn 2010, Turner 1977, Bowker 1984, Damuth 1987, Western 1974,
Agamidae	<i>Agama agama</i>	no	NA	NA	NA	mainland	10	16.92	8.00	2	0.64	10.31	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Tinkle 1967, Cisse and Karns 1978, Heideman 1994, Van Wilgen and Richardson 2012, Mediannikov et al. 2012, Trape et al. 2012
Agamidae	<i>Agama impalearis</i>	no	NA	NA	NA	mainland	28	25.98	11.80	2	0.46	10.82	Fitch 1970, Schleich et al. 1996, Geniez et al. 2004, Papenfuss 1969, Rogner 1997a, Le Berre 1989, Bons and Geniez 1996, Brown et al. 1999, Nagy et al. 1999, Waltner 1991, Brown and Nagy 2007, El Mouden et al. 1999, Radder et al. 2008, Kohler 2005, Znari and El Mouden 1997, Znari and Nagy 1997, Trape et al. 2012 Cogger 2000, Wilson and Swan 2003, Greer 1989, James and Shine 1988, Uller et al. 2009, Fischer and Lindenmayer 2005, Bustard 1978, Sinervo et al. 2010, Kohler 2005, Stuart-Smith et al. 2008, Michael and Lindenmayer 2010, Warner et al. 2008, Heatwole and Taylor 1987
Agamidae	<i>Amphibolurus muricatus</i>	no	NA	NA	NA	mainland	-32	17.45	5.55	2.5	0.64	8.94	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Calotes mystaceus</i>	no	NA	NA	NA	mainland	12	25.64	7.00	1	0.33	2.33	Fitch 1970, Inger and Greenberg 1966, Smith 1935, Tikader and Sharma 1992, Taylor 1963, Stuart et al. 2006, Cox et al. 1998, Inger and Colwell 1977, Manthey and Schuster 1996, Pauwels et al. 2003, Manthey 2008, Stuart and Emmett 2006, Taylor and Elbel 1958, Das 2010, Cox et al. 2010, Lalremsanga et al. 2010, Ahmed et al. 2009, Teynie and David 2010, Van Wilgen and Richardson 2012
Agamidae	<i>Calotes nemoricola</i>	no	NA	NA	NA	mainland	12	18.27	13.71	1	0.34	4.69	Tikader and Sharma 1992, Inger et al. 1984, Dunham et al. 1988, Manthey 2008, Das 2002, Kohler 2005, Subba Rao and Rajabai 1972, Turner 1977, Balakrishna et al. 2012, Tinkle et al. 1970, Fitch 1970, Inger and Greenberg 1966, Smith 1935, Ji et al. 2002, Anderson 1999, Minton 1966, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Taylor 1963, Henkel and Schmidt 2000,
Agamidae	<i>Calotes versicolor</i>	no	NA	NA	NA	mainland	23	15.24	13.65	2.5	0.26	9.02	Ranawana and Bambaradenyia 1998, Schmidt 1927, Manthey and Grossmann 1997, Bahir and Maduwage 2005, Rogner 1997a, Cox et al. 1998, Khan 2006, Deraniyagala 1953, Karsen et al. 1986, Inger and Colwell 1977, Inger et al. 1984, Vinson and Vinson 1969, Manthey and Schuster 1996, Clark 1990, Murthy

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Chlamydosaurus kingii</i>	no	NA	NA	NA	mainland	-18	253.09	11.99	1.5	1.64	29.45	1995, Pauwels et al. 2003, Shrestha 2001, Dunham et al. 1988, Manthey 2008, Grismer et al. 2007, Radder et al. 2008, Koul and Duda 1977, Somaweera and Somaweera 2009, Das 2002, Sinervo et al. 2010, Kohler 2005, Das 2010, Cox et al. 2010, Pandav et al. 2010, Das and de Silva 2011, Shanbhag 2002, Ahmed et al. 2009, Grismer 2011, Aryal et al. 2010, Meek et al. 2005, Castanet 1994, Teynie and David 2010, Van Wilgen and Richardson 2012, Ahmed et al. 2009, Masroor 2012 Cogger 2000, Perry and Garland 2002, Wilson and Swan 2003, Allison 2006, de Rooij 1915, Greer 1989, Manthey and Schuster 1996, Nagy et al. 1999, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Allison 2007, Garrick 2008, Wilson and Swan 2008, Bedford et al. 1993, Sinervo et al. 2010, Kohler 2005, Heatwole and Taylor 1987, Swanson 2007, Amey and Whittier 2000 Cogger 2000, Cox et al. 2003, Wilson and Swan 2003, Johnston 2005, Wilson and Swan 2008, Johnston 1999, Swanson 2007
Agamidae	<i>Ctenophorus fionni</i>	no	NA	NA	NA	mainland	-33	7.97	3.58	1.25	0.67	2.99	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Ctenophorus fordi</i>	no	NA	NA	NA	mainland	-31	3.15	2.30	2.5	0.27	1.58	Cogger 2000, Melville et al. 2006, Huey and Pianka 1981, Pianka 1986, Wilson and Swan 2003, Henle 1991, Greer 1989, Dunham et al. 1988, Wilson and Swan 2008, Warne and Charnov 2008, Huey et al. 2001, Kohler 2005, Uller and Olsson 2010, Huey and Pianka 2007, Abensperg-Traun and Steven 1997, Witten 1993, Turner 1977, Heatwole and Taylor 1987, Swan and Watharow 2005
Agamidae	<i>Ctenophorus isolepis</i>	no	NA	NA	NA	mainland	-24	4.89	3.82	2.5	0.42	4.02	Cogger 2000, Melville et al. 2006, Dunham and Miles 1985, Huey and Pianka 1981, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Uetz 2006, Dunham et al. 1988, Wilson and Swan 2008, Warne and Charnov 2008, Huey et al. 2001, Todd 2008, Daly et al. 2008, Melville and Schlute 2001, Turner 1977, Sinervo et al. 2010, Kohler 2005, Znari and El Mouden 1997, Huey and Pianka 2007, Abensperg-Traun and Steven 1997, Heatwole and Pianka 1993, Heatwole and Taylor 1987, Swanson 2007, Morton and James 1988, Pianka 1971, Gordon et al. 2010
Agamidae	<i>Ctenophorus maculosus</i>	no	NA	NA	NA	mainland	-30	4.49	3.21	2	0.45	2.86	Cogger 2000, Wilson and Swan 2003, Henle 1991, Olsson 1995, Jessop et al. 2009, Devi Stuart-Fox, pers. Comm. 01 January 2010,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Mitchell 1973, Kohler 2005, Heatwole and Taylor 1987
Agamidae	<i>Ctenophorus nuchalis</i>	no	NA	NA	NA	mainland	-25	15.02	3.65	2.5	0.89	8.16	Cogger 2000, Melville et al. 2006, Dunham and Miles 1985, Huey and Pianka 1981, Cooper and Vitt 2002, Pianka 1986, Wilson and Swan 2003, Henle 1991, Pianka 1971, Rogner 1997a, Greer 1989, Manthey and Schuster 1996, Nagy et al. 1999, Uetz 2006, Dunham et al. 1988, Brown and Nagy 2007, Garland and Else 1987, Wilson and Swan 2008, Warne and Charnov 2008, Huey et al. 2001, Daly et al. 2008, Melville and Schlute 2001, Sinervo et al. 2010, Schlesinger et al. 2010, Witten 1993, Bradshaw and Main 1968, MacMillen et al. 1989, Heatwole and Taylor 1987, Swanson 2007, Licht et al. 1966, Light et al. 1966, Gordon et al. 2010 Clobert et al. 1998, Cogger 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Dunham et al. 1988, Warne and Charnov 2008, Barbour et al. 2002, Chapman and Dell 1985, Kohler 2005, Bush et al. 2010, Witten 1993, Bradshaw and Main 1968, Heatwole and Taylor 1987, Licht et al. 1966
Agamidae	<i>Ctenophorus ornatus</i>	no	NA	NA	NA	mainland	-31	11.79	3.13	2	0.94	5.85	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Diporiphora nobbi</i>	no	NA	NA	NA	mainland	-28	7.15	6.00	1.5	0.35	3.18	Cogger 2000, Wilson and Swan 2003, Greer 1989, Kohler 2005, Michael and Lindenmayer 2010, Witten 1993, Rauhala 1993, Swanson 2007, Swan and Watharow 2005
Agamidae	<i>Draco biaro</i>	yes	Sangihe Archipelago	Biaro	26.50	Oceanic	2	8.95	2.00	NA	NA	NA	McGuire et al. 2007, Manthey 2008, Lazell 1987
Agamidae	<i>Draco bourouniensis</i>	yes	Maluku Islands	Buru	8473.20	Oceanic	-3	11.82	2.50	NA	2.80	NA	Musters 1983
Agamidae	<i>Draco palawanensis</i>	yes	Philippine Islands	Palawan	12188.60	Continental	3	12.57	3.00	NA	NA	NA	Das 2004, McGuire and Alcala 2000, Manthey 2008, McGuire and Alcala 2000, McGuire et al. 2007, Musters 1983, Manthey 2008, Auffenberg 1980
Agamidae	<i>Draco reticulatus</i>	yes	Philippine Islands	Samar	12849.40	Land bridge	11	12.91	2.50	NA	NA	NA	Musters 1983, Manthey 2008, Kaiser et al. 2011, Sanchez et al. 2012, O'Shea et al. 2012, Clobert et al. 1998, Fitch 1970, Taylor 1963, Brown and Alcala 1961, Manthey and Grossmann 1997, de Rooij 1915, Rogner 1997a, Cox et al. 1998, Musters 1983, Manthey 2008, Vitt and Price 1982, Mori and Hikida 1993, Mori and Hikida 1994, Inger 1983, Sinervo et al. 2010, Kohler 2005, Das 2010, Cox et al. 2010, Alcala 1986, Smith 1993, Avery 1982, Alcala 1966
Agamidae	<i>Draco timorensis</i>	yes	Lesser Sunda Islands	Timor	28418.10	Continental	-9	12.28	2.50	NA	NA	NA	Manthey and Grossmann 1997, Rogner 1997a, Cox et al. 1998, Manthey and Schuster 1996, Manthey 2010, Onn et al. 2010, Das 2010, Cox et al. 2010, Grismer 2011
Agamidae	<i>Draco volans</i>	no	NA	NA	NA	mainland	-2	10.98	4.00	1	0.19	0.78	Manthey and Grossmann 1997, Rogner 1997a, Cox et al. 1998, Musters 1983, Manthey 2008, Vitt and Price 1982, Mori and Hikida 1993, Mori and Hikida 1994, Inger 1983, Sinervo et al. 2010, Kohler 2005, Das 2010, Cox et al. 2010, Alcala 1986, Smith 1993, Avery 1982, Alcala 1966
Agamidae	<i>Gonocephalus bellii</i>	no	NA	NA	NA	mainland	4	34.29	4.00	5	0.33	6.65	Manthey and Grossmann 1997, Rogner 1997a, Cox et al. 1998, Manthey and Schuster 1996, Manthey 2010, Onn et al. 2010, Das 2010, Cox et al. 2010, Grismer 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Gonocephalus bornensis</i>	no	NA	NA	NA	mainland	3	31.56	4.00	3.5	1.02	14.28	Taylor 1963, Das 2004, Manthey and Grossmann 1997, Manthey and Schuster 1996, Malkmus et al. 2002, Manthey 2010, Das 2010, Das 2011 Fitch 1970, Inger and Greenberg 1966, Hendrickson 1966, Taylor 1963, Manthey and Grossmann 1997, Rogner 1997a, Cox et al. 1998, Manthey and Schuster 1996, Malkmus et al. 2002, Fitch 1982, Kohler 2005, Manthey 2010, Teynie et al. 2010, Das 2010, Cox et al. 2010, Das 2011, Grismer 2011, Grismer 2011b, Teynie and David 2010 Cogger 2000, Wilson and Swan 2003, Rogner 1997a, Greer 1989, Manthey and Schuster 1996, Bauer and Jackman 2008, Andrews and Pough 1980, Wilson and Swan 2008, Kohler 2005, Michael and Lindenmayer 2010, Heatwole and Pianka 1993, Heatwole and Taylor 1987, Swanson 2007 Cox et al. 2003, Huang 1998b, Huang 2007, Kohler 2005, Manthey 2010, Huang 1997 Schleich and Kastle 2002, Tikader and Sharma 1992, Khan 2006, Manthey 2010 Ota et al. 1998, Huang 2007, Manthey 2010 Ota 1989, Huang 2007, Manthey 2010
Agamidae	<i>Gonocephalus grandis</i>	no	NA	NA	NA	mainland	1	48.38	3.50	7	0.82	20.03	
Agamidae	<i>Intelligama lesueurii</i>	no	NA	NA	NA	mainland	-27	205.89	12.00	1.5	2.63	47.34	
Agamidae	<i>Japalura brevipes</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	23	5.71	5.30	1.5	NA	NA	
Agamidae	<i>Japalura kumaonensis</i>	no	NA	NA	NA	mainland	31	5.03	11.50	1.5	0.13	2.21	
Agamidae	<i>Japalura luei</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	25	7.26	5.50	NA	NA	NA	
Agamidae	<i>Japalura makii</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	23	6.60	6.30	NA	NA	NA	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Japalura polygonata</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	24	4.24	3.40	NA	NA	NA	Goris and Maeda 2004, Manthey and Schuster 1996, Huang 2007, Manthey 2010, Tanaka 1986, Rummery et al. 1995
Agamidae	<i>Japalura swinhonis</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	24	7.10	3.95	2.5	0.19	1.92	Clobert et al. 1998, Perry and Garland 2002, Huang 2006, Ota 1991, Dunham et al. 1988, Warne and Charnov 2008, Huang 1998b, Huang 2007, Manthey 2010, Das 2010, Kuo et al. 2009, Huang 2010, Huang 1997, Norval et al. 2011, Charnov et al. 2007, Fitch 1970, Smith 1935, Szczerbak 2003, Anderson 1999, Perry and Garland 2002, Baran and Atatur 1998, Leviton and Anderson 1970, Greene 1982, Rogner 1997a, Anderson and Leviton 1969, Reed and Marx 1959, Khan 2006, Manthey and Schuster 1996, Clark 1990, Ananjeva and Tuniev 1994, Waltner 1991, Ahmadzadeh et al. 2008, Weber 1960, Das 2002, Kohler 2005, Arakelyan et al. 2011
Agamidae	<i>Laudakia caucasica</i>	no	NA	NA	NA	mainland	35	33.82	9.00	1.5	0.94	12.63	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Laudakia stellio</i>	no	NA	NA	NA	mainland	35	27.96	7.70	2.5	0.71	13.67	Frankenberg and Werner 1992, Amitai and Bouskila 2001, Arbel 1984, Arnold and Ovenden 2004, Perry and Garland 2002, Disi et al. 2001, Baran and Atatur 1998, Zinner 1967, Flower 1933, El Din 2006, Kumlutas et al. 2004, Reed and Marx 1959, Le Berre 1989, Manthey and Schuster 1996, Atatur and Gocmen 2001, Ananjeva and Tuniev 1994, Valakos et al. 2008, Valakos et al. 2004, Terbish et al. 2006, Garrick 2008, Kwet 2009, McElroy et al. 2008, Kohler 2005, Baier et al. 2009, Lachman et al. 2006, Almog et al. 2005, Panov and Zykova 1997, Hertz and Nevo 1981, Bar and Haimovitch 2012, Van Wilgen and Richardson 2012, Kopan and Yom-Tov 1982, Yanai and Dabool 2012
Agamidae	<i>Leiolepis reevesii</i>	no	NA	NA	NA	mainland	23	27.45	3.72	1.5	1.22	6.81	Zug et al. 2001, Taylor 1963, Cox et al. 1998, Manthey 2010, Das 2010, Cox et al. 2010, Du et al. 2011, Teynie and David 2010
Agamidae	<i>Lophognathus longirostris</i>	no	NA	NA	NA	mainland	-23	14.04	4.70	2	0.42	3.96	Huey and Pianka 1981, Cogger 2000, Melville et al. 2006, Pianka 1986, Wilson and Swan 2003, Greer 1989, Manthey and Schuster 1996, Clusella-Trullas et al. 2008, Wilson and Swan 2008, Huey et al. 2001, Melville and Schlute 2001, Sinervo et al. 2010, Kohler 2005, Huey and Pianka 2007, Heatwole and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Pianka 1993, Curry-Lindahl 1979, Bush et al. 2007, Light et al. 1966
Agamidae	<i>Moloch horridus</i>	no	NA	NA	NA	mainland	-25	19.35	7.25	1	0.86	6.20	Dunham and Miles 1985, Fitch 1970, Huey and Pianka 1981, Cogger 2000, Melville et al. 2006, Pianka and Parker 1975, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Manthey and Schuster 1996, Dunham et al. 1988, Brown and Nagy 2007, Vitt and Price 1982, Wilson and Swan 2008, Huey et al. 2001, Chapman and Dell 1985, Sinervo et al. 2010, Kohler 2005, Bush 1992, Huey and Pianka 2007, Abensperg-Traun and Steven 1997, Heatwole and Taylor 1987, Swanson 2007, Curry-Lindahl 1979, Bush et al. 2007, Light et al. 1966
Agamidae	<i>Phrynocephalus guttatus</i>	no	NA	NA	NA	mainland	47	2.36	2.50	2.5	0.22	1.40	Szczerbak 2003, Manthey and Schuster 1996, Shenbrot and Semenov 1986, Rogovin and Semenov 2004, Kohler 2005, Liu et al. 2012
Agamidae	<i>Phrynocephalus helioscopus</i>	no	NA	NA	NA	mainland	45	2.51	6.00	2.5	0.12	1.82	Tinkle et al. 1970, Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Rogner 1997a, Terbish et al. 2006, Shenbrot and Semenov 1986, Turner 1977, Sinervo et al. 2010, Kohler 2005, Clemann et al. 2008

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Phrynocephalus mystaceus</i>	no	NA	NA	NA	mainland	43	8.58	5.00	1.5	0.58	4.37	Szczerbak 2003, Nikolsky 1915, Manthey and Schuster 1996, Clark 1990, Shenbrot and Semenov 1986, Sinervo et al. 2010, Kohler 2005, Clemann et al. 2008
Agamidae	<i>Phrynocephalus theobaldi</i>	no	NA	NA	NA	mainland	34	2.73	2.00	1	0.34	0.69	Smith 1935, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Shrestha 2001, Das 2002
Agamidae	<i>Phrynocephalus versicolor</i>	no	NA	NA	NA	mainland	43	4.57	2.50	1.5	0.22	0.84	Szczerbak 2003, Perry and Garland 2002, Rogner 1997a, Rogovin et al. 2000, Terbish et al. 2006, Shenbrot and Semenov 1986, Kohler 2005, Tang et al. 2012
Agamidae	<i>Physignathus cocincinus</i>	no	NA	NA	NA	mainland	22	93.98	10.50	1.5	2.02	31.89	Smith 1935, Taylor 1963, Stuart et al. 2006, Rogner 1997a, Ziegler 2002, Cox et al. 1998, Inger and Colwell 1977, Manthey and Schuster 1996, Grismer et al. 2007, Stuart and Emmett 2006, Greer 1989, Kohler 2005, Taylor and Elbel 1958, Manthey 2010, Das 2010, Cox et al. 2010, Grismer 2011, Meek 1999, Teynie and David 2010
Agamidae	<i>Pogona barbata</i>	no	NA	NA	NA	mainland	-30	109.73	16.75	2	0.94	31.33	Fitch 1970, Cogger 2000, Cooper and Vitt 2002, Wilson and Swan 2003, Rogner 1997a, Greer 1989, Bustard 1966, Manthey and Schuster 1996, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Wilson and Swan 2008, Radder et al. 2008, Sinervo et al. 2010, Michael and Lindenmayer 2010, Wilson 2003, Heatwole and Pianka

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1993, Bartlett and Bartlett 2009, Heatwole and Taylor 1987, Swanson 2007, Amey and Whittier 2000, Michael et al. 2011, Light et al. 1966, Swan and Watharow 2005
Agamidae	<i>Pogona minor</i>	no	NA	NA	NA	mainland	-24	29.59	6.50	2	0.89	11.63	Cogger 2000, Melville et al. 2006, Cooper and Vitt 2002, Pianka 1986, Wilson and Swan 2003, Henle 1991, Greer 1989, Manthey and Schuster 1996, Huey et al. 2001, Chapman and Dell 1985, Todd 2008, Davidge 1979, Kohler 2005, Bush et al. 2010, Bush 1992, Huey and Pianka 2007, Bradshaw and Main 1968, Pianka 1986, Heatwole and Taylor 1987, Swanson 2007, Davidge 1979, Light et al. 1966, Moro and MacAulay 2010, Heatwole and Butler 1981 Cogger 2000, Melville et al. 2006, Wilson and Swan 2003, Rogner 1997a, Greer 1989, Manthey and Schuster 1996, Clusella-Trullas et al. 2008, Henle 1989c, Melville and Schlute 2001, Kohler 2005, Bartlett and Bartlett 2009, MacMillen et al. 1989, Heatwole and Taylor 1987, Swanson 2007, Gordon et al. 2010, Swan and Watharow 2005
Agamidae	<i>Pogona vitticeps</i>	no	NA	NA	NA	mainland	-27	132.64	20.75	3.5	1.36	98.50	Schleich and Kastle 2002, Manthey 2010
Agamidae	<i>Sitana fusca</i>	no	NA	NA	NA	mainland	27	2.42	7.20	2	0.07	1.00	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Sitana ponticeriana</i>	no	NA	NA	NA	mainland	24	3.37	14.30	2.5	0.07	2.47	Fitch 1970, Smith 1935, Daniel 1983, Tikader and Sharma 1992, Boulenger 1890, Erdelen 1998, Rao 1998, Deraniyagala 1953, Manthey and Schuster 1996, Shrestha 2001, Dunham et al. 1988, Radder et al. 2008, Somaweera and Somaweera 2009, Das 2002, Kohler 2005, Manthey 2010, Das and de Silva 2011, Radder and Shanbhag 2003, Subba Rao and Rajabai 1972, Turner 1977, Pal et al. 2010, Pal et al. 2011
Agamidae	<i>Sitana sivalensis</i>	no	NA	NA	NA	mainland	28	1.96	6.86	2.5	0.06	0.96	Schleich and Kastle 2002, Manthey 2010
Agamidae	<i>Trapelus mutabilis</i>	no	NA	NA	NA	mainland	26	9.07	8.50	2	0.58	9.90	Schleich et al. 1996, Geniez et al. 2004, Flower 1933, El Din 2006, Le Berre 1989, Bons and Geniez 1996, Kohler 2005, Wagner et al. 2011, Trape et al. 2012, Szczerbak 2003, Anderson 1999, Minton 1966, Disi et al. 2001, Baran and Atatur 1998, Leviton et al. 1992, Rogner 1997a, Reed and Marx 1959, Khan 2006, Anderson 1963, Clark 1990, Ahmadzadeh et al. 2008, Moravec and Modry 1994b, Anderson 1963 (blanfordi), Weber 1960, Kohler 2005, Fathinia and Rastegar-Pouyani 2011, Hertz and Nevo 1981 (as pallida), Fathinia et al. 2009 (as persicus)
Agamidae	<i>Trapelus ruderatus</i>	no	NA	NA	NA	mainland	34	10.15	8.00	2.5	0.42	8.43	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Agamidae	<i>Trapelus sanguinolentus</i>	no	NA	NA	NA	mainland	43	14.04	10.50	3	0.52	16.53	Fitch 1970, Szczerbak 2003, Manthey and Schuster 1996, Ananjeva and Tselarius 1986, Sinervo et al. 2010, Kohler 2005, Clemann et al. 2008
Agamidae	<i>Uromastyx acanthinura</i>	no	NA	NA	NA	mainland	27	247.50	13.00	1.375	2.80	50.02	Clobert et al. 1998, Fitch 1970, Schleich et al. 1996, Perry and Garland 2002, Geniez et al. 2004, Pianka and Vitt 2003, Rogner 1997a, Le Berre 1989, Manthey and Schuster 1996, Bons and Geniez 1996, Garrick 2008, Turner 1977, Kohler 2005, Damuth 1987, Trape et al. 2012, Wilms 2005
Agamidae	<i>Uromastyx aegyptia</i>	no	NA	NA	NA	mainland	27	826.45	22.00	0.875	4.65	89.59	Amitai and Bouskila 2001, Anderson 1999, Perry and Garland 2002, Disi et al. 2001, Flower 1933, Gallagher 1971, Arnold 1984, El Din 2006, Arnold 1980, Schatti and Desvoignes 1999, Le Berre 1989, Jongbloed 2000, Hornby 1996, Arbel 1984, Wilms and Bohme 2000, Kohler 2005, Nemtzov 2008, Zari 1991, van der Kooij 2001, Bar and Haimovitch 2012, Bringsoe 1998, Rapoport 1974, Wilms 2005, Wilms et al. 2011
Agamidae	<i>Uromastyx ornata</i>	no	NA	NA	NA	mainland	24	86.57	11.00	1	3.35	36.82	Amitai and Bouskila 2001, Flower 1933, El Din 2006, Zari 1996, Sinervo et al. 2010, Kohler 2005, Nemtzov 2008, Bar and Haimovitch 2012, Wilms 2005
Anguidae	<i>Anguis cephalonica</i>	no	NA	NA	NA	mainland	37	4.40	10.50	1	1.09	11.43	Arnold and Ovenden 2004, Grillitsch and Cabela 1990,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Valakos et al. 2008
Anguidae	<i>Anguis fragilis</i>	no	NA	NA	NA	mainland	52	9.13	9.36	0.75	0.39	2.76	Tinkle et al. 1970, Fitch 1970, Wiens and Slingluff 2001, Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Spellerberg 2002, Zug et al. 2001, Cooper and Habegger 2000, Street 1979, Rogner 1997b, Greene et al. 2006, Sindaco et al. 2006, Corti and Cascio 2002, Dunham et al. 1988, Andrews and Pough 1980, Valakos et al. 2008, Galan and Salvador 2006, Radder et al. 2008, Cooper and Bradley 2009, AL-Sadoon and Spellerberg 1985, Cree and Guillette 1995, Ibaruengoytia and Casalins 2007, Van Wyk 1991, Kwet 2009, Malkmus 2004, Sinervo et al. 2010, Necas et al. 1997, Brown and Roberts 2008, Meek 2005, Arakelyan et al. 2011, Hailey and Elliot 1995, Maso and Pijoan 2011
Anguidae	<i>Barisia herrerae</i>	no	NA	NA	NA	mainland	19	22.18	6.00	1	0.37	2.20	Zaldivar-Riveron and de Oca 2002
Anguidae	<i>Barisia imbricata</i>	no	NA	NA	NA	mainland	23	21.80	7.30	1	0.21	1.56	Fitch 1970, Wiens and Slingluff 2001, Zaldivar-Riveron and de Oca 2002, Guillette and Smith 1982, Greene et al. 2006, McCranie and Wilson 2001, Duellman 1961, Martinez-Torres et al. 2003, Radder et al. 2008, Guillette and Casas-Andreu 1987, Cree and Guillette 1995, Lemos-Espinal and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Anguidae	<i>Diploglossus millepunctatus</i>	yes	None	Malpelo	7.65	Oceanic	4	306.42	NA	NA	3.50	NA	Smith 2007, Davis and Smith 1953, Lemos-Espinal and Smith 2007b, Duellman 1965, Dixon and Lemos-Espinal 2010
Anguidae	<i>Diploglossus pleii</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	18.57	3.00	NA	NA	NA	Kiester 1975, Avery 1982, Lopez-Victoria 2006, Lopez-Victoria et al. 2011
Anguidae	<i>Dopasia gracilis</i>	no	NA	NA	NA	mainland	25	10.96	5.50	1	0.37	2.06	Wiens and Slingsluff 2001, Schwartz and Henderson 1991, Greene et al. 2006, Henderson and Powell 2009, Rivero 1998
Anguidae	<i>Elgaria coerulea</i>	no	NA	NA	NA	mainland	46	12.75	5.40	1	0.25	1.35	Fitch 1970, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Cox et al. 1998, Greene et al. 2006, Das 2010, Cox et al. 2010, Lalremsanga et al. 2010, Ahmed et al. 2009, Ahmed 2009
													Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1985, Wiens and Slingsluff 2001, Stebbins 2003, Smith 1946, Pianka and Vitt 2003, Rogner 1997b, Greene et al. 2006, Greene et al. 2006, Van Denburgh 1922, Dunham et al. 1988, Vitt and Price 1982, Warne and Charnov 2008, Radder et al. 2008, Cooper and Bradley 2009, Lais 1976, Sinervo et al. 2010, Jones and Lovich 2009, McBrayer and Anderson 2007, Rutherford 2004, Stewart 1985, Vitt 1974, Stewart 1984, Stebbins and McGinnis 2012, St. John 2002

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Anguidae	<i>Elgaria multicarinata</i>	no	NA	NA	NA	mainland	39	29.25	10.90	2	0.30	6.48	Tinkle et al. 1970, Wiens and Slingluff 2001, Fitch 1985, Stebbins 2003, Grismer 2002, Pianka and Vitt 2003, Greene 1982, Smith 1946, Linsdale 1932, Rogner 1997b, Greene et al. 2006, Van Denburgh 1922, Dunham et al. 1988, Andrews and Pough 1980, Brown and Nagy 2007, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Goldberg 1972, Brattstrom 196, Kingsbury 1995, Stebbins and McGinnis 2012, Mulroy and Wiseman 2012, Lemm 2006, St. John 2002
Anguidae	<i>Gerrhonotus infernalis</i>	no	NA	NA	NA	mainland	26	85.79	18.00	1	0.33	5.95	Smith 1946, Greene et al. 2006, Werler 1949, Werler 1951, Lemos-Espinal and Smith 2007, Flury 1949, Kohler 2005, Jones and Lovich 2009, Burkett 1962
Anguidae	<i>Mesaspis gadovii</i>	no	NA	NA	NA	mainland	17	8.49	7.25	1	0.20	1.47	Fitch 1970, Tihen 1954, Davis and Dixon 1961, Ramirez-Pinilla et al. 2009, Goldberg 2011
Anguidae	<i>Mesaspis juarezi</i>	no	NA	NA	NA	mainland	18	4.55	3.00	1	0.13	0.38	Cooper and Habegger 2000, Karges and Wright 1987, Ramirez-Pinilla et al. 2009
Anguidae	<i>Mesaspis monticola</i>	no	NA	NA	NA	mainland	9	5.62	4.39	0.5	0.13	0.28	Fitch 1970, 1973, Cox et al. 2003, Kohler 2003, Tihen 1954, Greene et al. 2006, Dunham et al. 1988, Fitch 1982, Cree and Guillette 1995, Van Wyk 1991, Vial and Stewart 1985, Pough et al. 2003

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Anguidae	<i>Ophiodes striatus</i>	no	NA	NA	NA	mainland	-20	22.31	5.55	1	0.35	1.95	Wiens and Slingluff 2001, Nogueira et al. 2005, Achaval and Olmos 2003, Colli et al. 2002, Pizzatto 2005, Wiens et al. 2006, Carreira et al. 2005, Shine and Wall 2008, Montechiaro et al. 2011 Fitch 1970, 1985, Cox et al. 2003, Conant and Collins 1998, McConkey 1954, Pianka and Vitt 2003, Greene et al. 2006, Johnson and Voigt 1978, Cooper and Bradley 2009, Sinervo et al. 2010, Kohler 2005, Jensen et al. 2008, Beane et al. 2010, Fitch 1956, Brattstrom 196 Conant and Collins 1998, McConkey 1954, Pianka and Vitt 2003, Bartlett 1985, Greene et al. 2006, Holman 1971, Jensen et al. 2008, Kohler 2005, Beane et al. 2010
Anguidae	<i>Ophisaurus attenuatus</i>	no	NA	NA	NA	mainland	34	18.35	9.74	1	0.65	6.34	Wiens and Slingluff 2001, Conant and Collins 1998, McConkey 1954, Smith 1946, Pianka and Vitt 2003, Schwartz and Henderson 1991, Rogner 1997b, Greene et al. 2006, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Cooper and Bradley 2009, Holman 1971, Kohler 2005, Jensen et al. 2008, Shine and Wall 2008, Beane et al. 2010, Kamel and Gatten 1983
Anguidae	<i>Ophisaurus compressus</i>	no	NA	NA	NA	mainland	29	7.96	11.00	1	0.30	3.30	
Anguidae	<i>Ophisaurus ventralis</i>	no	NA	NA	NA	mainland	31	15.23	11.00	1	0.41	4.56	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Anguidae	<i>Pseudopus apodus</i>	no	NA	NA	NA	mainland	40	157.55	8.00	1	4.21	33.65	Fitch 1970, Amitai and Bouskila 2001, Szczerbak 2003, Anderson 1999, Disi et al. 2001, Baran and Atatur 1998, Leviton et al. 1992, Pianka and Vitt 2003, Flower 1933, Rogner 1997b, Reed and Marx 1959, Greene et al. 2006, Corti and Cascio 2002, Ahmadzadeh et al. 2008, Valakos et al. 2008, Valakos et al. 2004, Kwet 2009, Kohler 2005, De Magalhaes and Costa 2009, Arakelyan et al. 2011, Meek 1986, Bar and Haimovitch 2012, Rifai et al. 2005
Anniellidae	<i>Anniella pulchra</i>	no	NA	NA	NA	mainland	35	3.98	1.65	0.75	0.39	0.49	Fitch 1970, Wiens and Slingluff 2001, Stebbins 2003, Grismer 2002, Pianka and Vitt 2003, Smith 1946, Linsdale 1932, Van Denburgh 1922, Andrews and Pough 1980, Hunt 2008, Hunt 2006, Sinervo et al. 2010, Goldberg and Miller 1985, Turner 1977, Kamel and Gatten 1983, Hailey and Elliot 1995, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006
Bipedidae	<i>Bipes biperus</i>	no	NA	NA	NA	mainland	26	5.63	2.08	0.5	0.78	0.81	Wiens et al. 2006, Andrade et al. 2006, Kearney 2003, Vega 2001, Papenfuss 1982, Bernardo-Silva et al. 2006
Bipedidae	<i>Bipes canaliculatus</i>	no	NA	NA	NA	mainland	18	8.51	2.80	0.5	0.86	1.21	Wiens et al. 2006, Andrade et al. 2006, Vega 2001, Papenfuss 1982, Davis and Dixon 1961, Hodges and Perez-Ramos 2001

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Bipedidae	<i>Bipes tridactylus</i>	no	NA	NA	NA	mainland	17	2.18	2.20	1	0.24	0.52	Wiens et al. 2006, Andrade et al. 2006, Vega 2001, Papenfuss 1982
Carphodactylidae	<i>Carphodactylus laevis</i>	no	NA	NA	NA	mainland	-17	41.06	2.00	5	1.76	17.60	Cogger 2000, Wilson and Swan 2003, Wilson and Swan 2008, Henkel 2010, Rosler 2005, Wilson 2005, Swanson 2007
Carphodactylidae	<i>Nephrurus levis</i>	no	NA	NA	NA	mainland	-25	8.07	2.00	2.5	0.84	4.21	Cogger 2000, Withers et al. 2000, Pianka 1986, Wilson and Swan 2003, Henle 1991, Greer 1989, Cree 1994, Vitt and Price 1982, Huey et al. 2001, Storr et al. 1990, How et al. 1990, Read 1999, Bauer 1990, Sinervo et al. 2010, Köhler 2005, Henkel 2010, Rosler 2005, Daza et al. 2009, Werner and Whitaker 1978, Heatwole and Taylor 1987, Wilson 2005, Swanson 2007, Morton and James 1988, Gordon et al. 2010, Henkel and Schmidt 1995
Carphodactylidae	<i>Phyllurus platurus</i>	no	NA	NA	NA	mainland	-34	12.00	2.00	1.5	0.85	2.56	Cogger 2000, Cox et al. 2003, Doughty and Shine 1995, Wilson and Swan 2003, Greer 1989, Couper et al. 1993, Henkel 2010, Daza et al. 2009, Swanson 2007, Lezzi 2008, Henkel and Schmidt 1995, Werner et al. 1993
Carphodactylidae	<i>Saltuarius cornutus</i>	no	NA	NA	NA	mainland	-17	65.58	2.00	1	2.24	4.48	Cogger 2000, Wilson and Swan 2003, Rogner 1997a, Greer 1989, Couper et al. 1993, Schaffer 2004, Couper et al. 1993, Köhler 2005, Henkel 2010, Rosler 2005, Daza et al. 2009, Carey and Judge 2000, Wilson 2005, Werner et al. 1993

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Carphodactylidae	<i>Underwoodisaurus milii</i>	no	NA	NA	NA	mainland	-31	9.31	2.00	2.5	0.93	4.65	Cogger 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Cree 1994, Wilson and Swan 2008, Storr et al. 1990, Chapman and Dell 1985, Bustard 1967, Angilletta and Werner 1998, How et al. 1990, Read 1999, Bauer 1990, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Bush et al. 2010, Rosler 2005, Wilson and Swan 2010, Daza et al. 2009, Michael and Lindenmayer 2010, Shah 2002, Werner and Whitaker 1978, Heatwole and Taylor 1987, Swanson 2007, Wilson 2005, Michael et al. 2011, Bush et al. 2007, Light et al. 1966, Swan and Watharow 2005, Werner et al. 1993
Chamaeleonidae	<i>Archaius tigris</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	16.38	7.50	NA	NA	NA	Henkel and Schmidt 2000, Necas 1999, Cheke 1984, Gerlach 2008
Chamaeleonidae	<i>Bradypodion pumilum</i>	no	NA	NA	NA	mainland	-34	11.41	12.50	3	0.35	13.20	Fitch 1970, Clobert et al. 1998, Fitzsimons 1943, Branch 1998, LeBerre et al. 2000, Necas 1999, Rogner 1997a, Andrews 2008, Dunham et al. 1988, Tolley and Burger 2007, Warne and Charnov 2008, Sinervo et al. 2010, Tilbury 2010, Turner 1977, Carey and Judge 2000, Avery 1982
Chamaeleonidae	<i>Bradypodion ventrale</i>	no	NA	NA	NA	mainland	-32	16.13	13.00	1.5	0.48	9.28	Fitch 1970, Fitzsimons 1943, Branch 1998, Necas 1999, Tilbury 2010

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Chamaeleonidae	<i>Chamaeleo calyptratus</i>	no	NA	NA	NA	mainland	16	106.55	52.00	2.5	0.56	73.02	LeBerre et al. 2000, Necas 1999, Rogner 1997a, Schatti and Desvoignes 1999, Andrews 2008, Sinervo et al. 2010, Kohler 2005, Tilbury 2010, Van Wilgen and Richardson 2012, Schleich et al. 1996, Amitai and Bouskila 2001, Arnold and Ovenden 2004, Disi et al. 2001, Baran and Atatur 1998, Geniez et al. 2004, Flower 1933, El Din 2006, LeBerre et al. 2000, Kumlutas et al. 2004, Necas 1999, Rogner 1997a, Le Berre 1989, Bons and Geniez 1996, Atatur and Cascio 2002, Andrews 2008, Valakos et al. 2008, Arbel 1984, Kwet 2009, Malkmus 2004, Sinervo et al. 2010, Kohler 2005, Baier et al. 2009, Tilbury 2010, Cuadrado 2010, Bar and Haimovitch 2012, Dimaki et al. 2000, Bogin and Werner 1995, Trape et al. 2012, Maso and Pijoan 2011, Fitch 1970, Spawls et al. 2002, Fitzsimons 1943, Loveridge 1936, 1942, Branch 1998, Razzetti and Msuya 2002, Pianka 1986, Pianka and Vitt 2003, Broadley 1971, Parker 1936, Jeffery 1993, Schmidt et al. 1919, Necas 1999, Rogner 1997a, Loveridge 1953, Auerbach 1987, Barbour and Loveridge 1928, Branch
Chamaeleonidae	<i>Chamaeleo chamaeleon</i>	no	NA	NA	NA	mainland	32	35.07	30.00	1	0.59	17.77	
Chamaeleonidae	<i>Chamaeleo dilepis</i>	no	NA	NA	NA	mainland	-14	29.84	37.50	1	1.04	39.13	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Chamaeleonidae	<i>Chamaeleo monachus</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	13	61.29	42.00	NA	0.92	NA	2005, Chirio and LeBreton 2007, Andrews 2008, Graham and Marais 2007, Clusella-Trullas et al. 2008, Pauwels and Vande weghe 2008, Sinervo et al. 2010, Kohler 2005, Largen and Spawls 2010, Tilbury 2010, Pienaar 1966, Jackson and Blackburn 2010, Haagner et al. 2000, Avery 1982, Patrick et al. 2011, Brattstrom 196 Schatti and Desvoignes 1999, Tilbury 2010, Razzetti et al. 2011, Rosler and Wrantik 2004
Chamaeleonidae	<i>Chamaeleo namaquensis</i>	no	NA	NA	NA	mainland	-25	37.60	13.20	2.5	0.74	24.52	Fitzsimons 1943, Perry and Garland 2002, Pianka and Vitt 2003, Necas 1999, Andrews 2008, Graham and Marais 2007, Dunham et al. 1988, Warne and Charnov 2008, Branch 1988, Sinervo et al. 2010, Kohler 2005, Tilbury 2010, Dimaki et al. 2001, Avery 1982, Brattstrom 196
Chamaeleonidae	<i>Furcifer cephalolepis</i>	yes	Comoro Islands	Grande Comore	1148.00	Oceanic	-12	5.48	5.50	NA	NA	NA	Henkel and Schmidt 2000, Necas 1999, Rogner 1997a, Kohler 2005, Meirte 2004
Chamaeleonidae	<i>Furcifer polleni</i>	yes	Comoro Islands	Anjouan	424.00	Oceanic	-13	6.27	9.00	NA	NA	NA	Henkel and Schmidt 2000, Necas 1999, Rogner 1997a, Kohler 2005, Meirte 2004
Chamaeleonidae	<i>Rhampholeon marshalli</i>	no	NA	NA	NA	mainland	-19	8.87	14.00	1	0.31	4.32	Fitch 1970, Fitzsimons 1943, Branch 1998, Necas 1999, Raxworthy 1991, Kohler 2005, Tilbury 2010
Chamaeleonidae	<i>Trioceros ellioti</i>	no	NA	NA	NA	mainland	-1	17.65	10.00	3	0.62	18.72	Spawls et al. 2002, Loveridge 1942, Necas 1999, Rogner 1997a, Andrews 2008, Clusella-Trullas et al. 2008,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Chamaeleonidae	<i>Trioceros hoehnelii</i>	no	NA	NA	NA	mainland	0	12.68	11.70	2	0.50	11.78	Vonesh 1998, Sinervo et al. 2010, Tilbury 2010
Chamaeleonidae	<i>Trioceros jacksonii</i>	no	NA	NA	NA	mainland	-1	26.29	16.95	1.5	0.56	14.28	Tinkle et al. 1970, Fitch 1970, Spawls et al. 2002, Loveridge 1936, Necas 1999, Rogner 1997a, Branch 2005, Andrews 2008, Lin & Nelson 1980, Sinervo et al. 2010, Tilbury 2010, Dimaki et al. 2000, Hebrard et al. 1982, Spawls et al. 2002, Razzetti and Msuya 2002, LeBerre et al. 2000, Necas 1999, Rogner 1997a, Branch 2005, Andrews 2008, Clusella-Trullas et al. 2008, Lin & Nelson 1980, Sinervo et al. 2010, Tilbury 2010, Boulenger 1896, Carey and Judge 2000, Goldberg and Kraus 2011, Van Wilgen and Richardson 2012
Chamaeleonidae	<i>Trioceros quadricornis</i>	no	NA	NA	NA	mainland	6	26.97	13.00	2.5	2.26	73.31	LeBerre et al. 2000, Necas 1999, Klaver and Bohme 1992, Bohme and Klaver 1981, Chirio and LeBreton 2007, Hofer et al. 2003, Gonwouo et al. 2006, Kohler 2005, Tilbury 2010, Trape et al. 2012
Cordylidae	<i>Hemicordylus capensis</i>	no	NA	NA	NA	mainland	-33	23.21	2.00	1	1.50	3.00	FitzSimons 1943, Branch 1998, Costandius and Mouton 2006, Rogner 1997b, Graham and Marais 2007, Curtin et al. 2005, Mouton et al. 2010

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Cordylidae	<i>Pseudocordylus melanotus</i>	no	NA	NA	NA	mainland	-28	28.30	3.45	1	1.10	3.79	FitzSimons 1943, Cox et al. 2003, Branch 1998, Costandius and Mouton 2006, Rogner 1997b, Loveridge 1944, Boretto and Ibarguengoytia 2006, Graham and Marais 2007, Flemming 1993, McConnachie et al. 2009, Sinervo et al. 2010, Kohler 2005, Flemming and Mouton 2002, Van Wilgen and Richardson 2012
Cordylidae	<i>Smaug giganteus</i>	no	NA	NA	NA	mainland	-28	260.98	2.70	0.625	9.50	16.03	Fitch 1970, Branch 1998, Costandius and Mouton 2006, Rogner 1997b, Loveridge 1944, Boretto and Ibarguengoytia 2006, Graham and Marais 2007, Flemming 1993, Cree and Guillette 1995, Ibarguengoytia and Casalins 2007, Van Wyk 1991, Flemming and Mouton 2002, Mouton et al. 2010, Van Wyk 1994, Goldberg 2006, Ibarguengoytia and Cussac 1996, Carey and Judge 2000, Van Wilgen and Richardson 2012, Honegger 1969
Corytophanidae	<i>Basiliscus basiliscus</i>	no	NA	NA	NA	mainland	8	130.31	9.20	6.5	1.45	86.93	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Savage 2002, Cooper and Vitt 2002, Kohler 2003, Fitch 1973, Rand and Myers 1990, Duellman 1990, Evans 1947, Rogner 1997a, Dunham et al. 1988, Lotzkat 2007, Warne and Charnov 2008, Radder et al. 2008, Kohler 2008, Van Devender 1983, Sinervo et al. 2010, Kohler

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Corytophanidae	<i>Basiliscus vittatus</i>	no	NA	NA	NA	mainland	16	48.21	5.40	3	1.18	19.10	2005, Van Wilgen and Richardson 2012, Shine and Charnov 1992, Brattstrom 1965, Pike et al. 2008, Tinkle et al. 1970, Fitch 1970, 1973, 1985, Perry and Garland 2002, Campbell 1999, Stafford and Meyer 2000, Conant and Collins 1998, Savage 2002, Lee 2000, Kohler 2003, Duellman 1990, Canseco-Marquez et al. 2000, Rogner 1997a, Kohler 1996, McCranie and Castaneda 2005, McCranie et al. 2005, Kohler et al. 2006, Guyer and Donnelly 2005, Duellman 1961, Vitt and Zani 1998, Rand 1957, Kohler 2008, Davis and Dixon 1961, Lopez and Gonzalez 1997, Sinervo et al. 2010, Kohler 2005, Duellman 1963, Duellman 1965, Damuth 1987, Tinkle et al. 1967, Hirth 1963, Van Wilgen and Richardson 2012, Hirth 1965, Leenders and Watkins-Colwell 2004, Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Vitt et al. 1978, Melville 2006, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Grismer 2002, Degenhardt et al. 1996, Smith 1946, McGuire 1996, Linsdale 1932, Rogner 1997a, Van Denburgh 1922, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas
Crotaphytidae	<i>Crotaphytus collaris</i>	no	NA	NA	NA	mainland	33	20.57	6.00	2	1.13	13.55	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Crotaphytid ae	<i>Crotaphytus grimeri</i>	no	NA	NA	NA	mainland	32	15.21	3.00	1.5	3.00	13.49	et al. 2008, Sexton et al. 1992, Garrick 2008, Vitt and Price 1982, Warne and Charnov 2008, Cooper et al. 2001, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Case 1975, Telemeco and Baird 2011, Muchlinski et al. 1995, Rorabaugh 2008, Werner and Whitaker 1978, Fitch 1967, Brennan and Holycross 2009, Fitch 1956, Shine and Charnov 1992, Brattstrom 1965, Goldberg 2011, Shine and Schwarzkopf 1992, Stebbins 2003, Grismer 2002, McGuire 1996, McGuire 1994
Crotaphytid ae	<i>Crotaphytus reticulatus</i>	no	NA	NA	NA	mainland	27	16.75	9.80	2	0.86	16.82	Conant and Collins 1998, Pianka and Vitt 2003, Smith 1946, Greene 1982, Montanucci 1971, Werler 1951, Lemos-Espinal and Smith 2007, Husak and Ackland 2003, Montanucci 1976, Kohler 2005
Crotaphytid ae	<i>Gambelia sila</i>	no	NA	NA	NA	mainland	36	26.14	3.10	2.5	2.54	19.67	Lappin and Swinney 1999, Fitch 1985, Perry and Garland 2002, Stebbins 2003, Smith 1946, McGuire 1996, Van Denburgh 1922, Warne and Charnov 2008, Todd 2008, Warrick et al. 1998, Jennings 1995, Turner et al. 1969, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Curry-Lindahl 1979, Brattstrom 1965, Stebbins

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													and McGinnis 2012
Crotaphytidae	<i>Gambelia wislizenii</i>	no	NA	NA	NA	mainland	36	38.09	5.15	1.25	1.70	10.96	Clobert et al. 1998, Huey and Pianka 1981, Melville et al. 2006, Tinkle et al. 1970, Fitch 1970, 1985, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Grismer 2002, Pianka 1986, Degenhardt et al. 1996, Pianka and Vitt 2003, Smith 1946, Greene 1982, McGuire 1996, Linsdale 1932, Rogner 1997a, Van Denburgh 1922, Dunham et al. 1988, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne and Charnov 2008, Huey et al. 2001, Todd 2008, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Turner 1977, Turner et al. 1969, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Case 1975, Huey and Pianka 2007, Turner 1977, Krekorian 1983, Morton and James 1988, Bury 1982, Brennan and Holycross 2009, Cunningham 1966, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006, St. John 2002

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Dactyloidae	<i>Anolis acutus</i>	yes	Virgin Islands	St. Croix	214.40	Oceanic	18	1.71	1.00	6	0.19	1.16	Clobert et al. 1998, Dunham and Miles 1985, Perry and Garland 2002, Andrews 1979, Schwartz and Henderson 1991, Dunham et al. 1988, Andrews and Pough 1980, Warne and Charnov 2008, Henderson and Powell 2009, Turner 1977, Ruibal and Philibosian 1974, Stamps et al. 1997, Charnov et al. 2007, Schoener and Gorman 1968, Perry and Garland 2002, Schwartz and Henderson 1991, Beebe 1944b, Herrel et al. 2004, Lazell 1972, Simmons et al. 2005, Murphy 1997, Henderson and Powell 2009, Losos 2009, McTaggart et al. 2011, see references in John et al. 2012
Dactyloidae	<i>Anolis aeneus</i>	yes	Windward Islands	Grenada	322.70	Oceanic	6	2.14	1.00	NA	0.25	NA	Rand et al. 1975, Losos 2009, Lopez-Victoria 2006, Lopez-Victoria et al. 2011
Dactyloidae	<i>Anolis agassizi</i>	yes	None	Malpelo	7.65	Oceanic	4	9.49	1.00	NA	0.35	NA	Cox et al. 2003, Savage 2002, Kohler 2003, Fitch et al. 1976, Campbell 1973, Losos 2009, Marquez and Marquez 2009
Dactyloidae	<i>Anolis aquaticus</i>	no	NA	NA	NA	mainland	9	4.35	1.13	9	0.33	3.31	Schwartz and Henderson 1991, Herrel et al. 2004, Lazell 1972, Powell et al. 2005, Malhotra and Thorpe 1999, Kohler 2005, Henderson and Powell 2009, Losos 2009
Dactyloidae	<i>Anolis bimaculatus</i>	yes	Lesser Antilles	Antigua	277.00	Oceanic	17	4.76	1.50	NA	0.37	NA	Bennett and Gorman 1979, van Buurt 2005, Andrews and Pough 1980, Ruthven 1923, Sinervo et al. 2010, Damuth 1987
Dactyloidae	<i>Anolis bonairensis</i>	yes	Lesser Antilles	Bonaire	282.50	Oceanic	14	3.43	1.00	NA	NA	NA	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Dactyloidae	<i>Anolis carolinensis</i>	no	NA	NA	NA	mainland	32	2.54	1.30	10.5	0.23	3.13	Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, Perry and Garland 2002, Stafford and Meyer 2000, Conant and Collins 1998, Smith 1946, Rogner 1997a, Hodge et al. 2003, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Radder et al. 2008, Andrews and Rand 1974, Sinervo et al. 2010, Kohler 2005, Henderson and Powell 2009, Losos 2009, Jensen et al. 2008, van Berkum 1988, Rodda and Dean-Bradley 2001, Johnson et al. 2010, Damuth 1987, Carey and Judge 2000, Beane et al. 2010, McCoid 1994, Van Wilgen and Richardson 2012
Dactyloidae	<i>Anolis conspersus</i>	yes	Cayman Islands	Grand Cayman	196.30	Oceanic	19	1.52	1.00	NA	0.24	NA	Cox et al. 2003, Todd 2008, Henderson and Powell 2009, Losos et al. 1993
Dactyloidae	<i>Anolis cooki</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	1.95	1.00	NA	NA	NA	Schwartz and Henderson 1991, Williams 1983, Jenssen 1990, Henderson and Powell 2009, Sinervo et al. 2010, Henderson and Powell 2009, Hertz et al. 1993, Huey and Webster 1976
Dactyloidae	<i>Anolis cristatellus</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	20	2.80	1.00	16	0.24	3.80	Perry and Garland 2002, Conant and Collins 1998, Lee 2000, Kohler 2003, Andrews 1979, Rodda et al. 2001, Schwartz and Henderson 1991, Williams 1983, Huang and Tu 2008, Clusella-Trullas et al. 2008, Kohler 2008, c, Schoener and Schoener

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1971b, Sinervo et al. 2010, Kohler 2005, Henderson and Powell 2009, Johnson et al. 2008, Hertz et al. 1993, Huey and Webster 1976, Hertz 1983, Rivero 1998, Van Wilgen and Richardson 2012
Dactyloidae	<i>Anolis cupreus</i>	no	NA	NA	NA	mainland	13	1.46	1.00	18	0.13	2.37	Clobert et al. 1998, Perry and Garland 2002, Cox et al. 2003, Savage 2002, Fitch 1973, Andrews 1979, Stuart 1955, Dunham et al. 1988, Losos 2009, van Berkum 1986, van Berkum 1988, Clark 1973
Dactyloidae	<i>Anolis cuvieri</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	34.40	1.00	NA	1.62	NA	Perry and Garland 2002, Greene 1982, Schwartz and Henderson 1991, Rogner 1997a, Herrel et al. 2004, Williams 1983, Henderson and Powell 2009, Tuli et al. 2009, Rivero 1998
Dactyloidae	<i>Anolis evermanni</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	2.28	1.00	NA	0.18	NA	Schwartz and Henderson 1991, Williams 1983, Schoener and Schoener 1971b, Henderson and Powell 2009, Hertz 1983, Rivero 1998
Dactyloidae	<i>Anolis garmani</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	20	11.01	1.00	NA	0.43	NA	Cox et al. 2003, Conant and Collins 1998, Schwartz and Henderson 1991, Rogner 1997a, Herrel et al. 2004, Williams 1983, Rand 1968, Russel and Bauer 1991, Kohler 2005, Henderson and Powell 2009, Marquez and Marquez 2009, Rand 1967

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Dactyloidae	<i>Anolis gingivinus</i>	yes	Leeward Islands	Saint Martin	91.90	Oceanic	18	2.35	1.00	NA	NA	NA	Schwartz and Henderson 1991, Lazell 1972, Powell et al. 2005, Hodge et al. 2003, Malhotra and Thorpe 1999, Breuil 2002, Sinervo et al. 2010, Henderson and Powell 2009, see Powell and Bauer 2012, Lazell and Williams 1962
Dactyloidae	<i>Anolis grahami</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	1.88	1.00	NA	0.14	NA	Cox et al. 2003, Schwartz and Henderson 1991, Rogner 1997a, Herrel et al. 2004, Cope 1895, Williams 1983, Rand 1968, Schoener and Schoener 1971, Russel and Bauer 1991, Henderson and Powell 2009, Sinervo et al. 2010, Johnson et al. 2010, Rand 1967
Dactyloidae	<i>Anolis gundlachi</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	1.85	1.00	NA	0.12	NA	Andrews 1979, Hertz 1981, Schwartz and Henderson 1991, Rogner 1997a, Williams 1983, Huang and Tu 2008, Clusella-Trullas et al. 2008, Huey et al. 2009, Flores et al. 1994, Schoener and Schoener 1971b, Henderson and Powell 2009, Sinervo et al. 2010, Tuli et al. 2009, Turner 1977, Hertz et al. 1993, Huey and Webster 1976, Rand 1964, Hertz 1983, Vega-Castillo and Puente-Rolon 2011, Rivero 1998
Dactyloidae	<i>Anolis intermedius</i>	no	NA	NA	NA	mainland	10	1.99	1.00	11	0.13	1.45	Clobert et al. 1998, Savage 2002, Fitch 1973, Pounds 1988, Losos 2009, van Berkum 1986, Clark 1973

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Dactyloidae	<i>Anolis krugi</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	1.10	1.00	NA	0.09	NA	Schwartz and Henderson 1991, Williams 1983, Schoener and Schoener 1971b, Henderson and Powell 2009, Rand 1964, Vega-Castillo and Puente-Rolon 2011, Rivero 1998, Hertz 1979
Dactyloidae	<i>Anolis leachii</i>	yes	Lesser Antilles	Antigua	277.00	Oceanic	17	9.10	NA	NA	0.27	NA	Lazell 1972, Kolbe et al. 2008, Irschick et al. 1996, Henderson and Powell 2009, Tinkle et al. 1970, Clobert et al. 1998, Fitch 1970, Fitch 1973, Perry and Garland 2002, Savage 2002, Kohler 2003, Rand and Myers 1990, Duellman 1990, Andrews 1979, Evans 1947, Guyer and Donnelly 2005, Dunham et al. 1988, Andrews and Pough 1980, Fitch et al. 1976, Vitt and Zani 1998, Huey et al. 2001, Todd 2008, Andrews and Rand 1974, Watling et al. 2005, Lopez and Gonzalez 1997, Sinervo et al. 2010, Kohler 2005, Tuli et al. 2009, Losos 2009, Marquez and Marquez 2009, Whitfield et al. 2007, Heatwole and Sexton 1966, Ruibal and Philibosian 1974, Tinkle et al. 1967, Clark 1973, Perry and Garland 2002, Cox et al. 2003, Andrews 1979, Schwartz and Henderson 1991, Rogner 1997a, Herrel et al. 2004, Williams 1983, Todd 2008, Rand 1968, Schoener and Schoener 1971, Schoener 1977, Sinervo et al.
Dactyloidae	<i>Anolis limifrons</i>	no	NA	NA	NA	mainland	11	1.56	1.50	21	0.08	2.54	
Dactyloidae	<i>Anolis lineatopus</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	2.10	1.00	18	0.21	3.78	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2010, Kohler 2005, Henderson and Powell 2009, Johnson et al. 2010, Rand 1967, Jenssen 1973, Rand 1967
Dactyloidae	<i>Anolis lionotus</i>	no	NA	NA	NA	mainland	11	5.12	1.00	2.5	0.27	0.68	Cox et al. 2003, Kohler 2003, Fitch 1973, Rand and Myers 1990, Duellman 1990, Rogner 1997a, Guyer and Donnelly 2005, Vitt and Zani 1998, Huey et al. 2001, Campbell 1973, Losos 2009, Leal et al. 2002, van Berkum 1986, Marquez and Marquez 2009
Dactyloidae	<i>Anolis lividus</i>	yes	Leeward Islands	Montserrat	124.10	Oceanic	17	2.28	1.00	NA	0.18	NA	Schwartz and Henderson 1991, Lazell 1972, Henderson and Powell 2009
Dactyloidae	<i>Anolis longiceps</i>	yes	Greater Antilles	Navassa	5.20	Oceanic	18	9.10	1.00	NA	NA	NA	Schwartz and Henderson 1991, Thomas 1966, Sinervo et al. 2010, Henderson and Powell 2009
Dactyloidae	<i>Anolis luciae</i>	yes	Lesser Antilles	St. Lucia	639.80	Oceanic	14	3.61	1.00	NA	0.27	NA	Schwartz and Henderson 1991, Rogner 1997a, Lazell 1972, Henderson and Powell 2009, Daltry 2009, Corke 1987
Dactyloidae	<i>Anolis monensis</i>	yes	Greater Antilles	Mona	57.00	Oceanic	18	1.65	NA	NA	0.18	NA	Schwartz and Henderson 1991, Herrel et al. 2004, Uetz 2006, Gorman and Stamm 1975, Sinervo et al. 2010, Henderson and Powell 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Dactyloidae	<i>Anolis nebulosus</i>	no	NA	NA	NA	mainland	22	1.27	2.00	25.5	0.13	6.70	Fitch 1970, Perry and Garland 2002, Ramirez-Bautista and Vitt 1997, Andrews 1979, McCranie and Wilson 2001, Duellman 1961, Hardy and McDiarmid 1969, Davis and Dixon 1961, Davis and Smith 1953, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Duellman 1965, Losos 2009, Rorabaugh 2008, Ruibal and Philibosian 1974, Clark 1973
Dactyloidae	<i>Anolis oculatus</i>	yes	Windward Islands	Dominica	787.30	Oceanic	15	2.98	1.05	NA	0.25	NA	Andrews 1979, Schwartz and Henderson 1991, Rogner 1997a, Lazell 1972, Dunham et al. 1988, Todd 2008, Malhotra and Thorpe 1999, Somma and Brooks 1976, Andrews and Rand 1974, Sinervo et al. 2010, Kohler 2005, Brooks 1968, Bullock and Evans 1990, Lazell and Williams 1962
Dactyloidae	<i>Anolis opalinus</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	1.62	1.00	28	0.16	4.52	Cox et al. 2003, Schwartz and Henderson 1991, Rogner 1997a, Cope 1895, Williams 1983, Todd 2008, Rand 1968, Schoener and Schoener 1971, Russel and Bauer 1991, Schoener 1977, Kohler 2005, Henderson and Powell 2009, Jenssen and Nunez 1994, Jenssen 1973, Rand 1967
Dactyloidae	<i>Anolis oxylophus</i>	no	NA	NA	NA	mainland	11	4.30	1.00	26	0.39	10.08	Vitt et al. 1995, Irschik et al. 1997, Cox et al. 2003, Kohler 2005, Leal et al. 2002, Losos 2009, Savage 2002
Dactyloidae	<i>Anolis pogus</i>	yes	Leeward Islands	Saint Martin	91.90	Oceanic	18	1.43	1.00	NA	NA	NA	Lazell 1972, Powell et al. 2005, Malhotra and Thorpe 1999, Breuil 2002, Henderson

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source and Powell 2009
Dactyloidae	<i>Anolis polylepis</i>	no	NA	NA	NA	mainland	9	2.00	1.00	27.5	0.15	4.24	Perry and Garland 2002, Cox et al. 2003, Savage 2002, Kohler 2003, Andrews 1979, Andrews 1983, Perry 1996, Andrews and Rand 1974, Sinervo et al. 2010, Losos 2009, Marquez and Marquez 2009, Clark 1973
Dactyloidae	<i>Anolis pulchellus</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	1.03	1.00	NA	NA	NA	Andrews 1979, Schwartz and Henderson 1991, Williams 1983, Radder et al. 2008, Henderson and Powell 2009, Losos 2009, Rand 1964, Rivero 1998, Hertz 1979
Dactyloidae	<i>Anolis reconditus</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	12.21	1.00	NA	0.48	NA	Schwartz and Henderson 1991, Williams 1983, Kohler 2005, Henderson and Powell 2009, Losos 2009
Dactyloidae	<i>Anolis richardii</i>	yes	Lesser Antilles	St. Vincent	381.00	Oceanic	12	7.54	1.00	NA	0.59	NA	Schoener and Gorman 1968, Cox et al. 2003, Greene 1982, Schwartz and Henderson 1991, Herrel et al. 2004, Lazell 1972, Daudin and de Silva 2007, Simmons et al. 2005, Murphy 1997, Todd 2008, Henderson and Powell 2009, Losos 2009
Dactyloidae	<i>Anolis roquet</i>	no	NA	NA	NA	mainland	15	3.03	1.00	25.5	0.24	6.05	Clobert et al. 1998, Hertz 1981, Schwartz and Henderson 1991, Rogner 1997a, Lazell 1972, Huang and Tu 2008, Kohler 2005, Henderson and Powell 2009, Hertz 1983
Dactyloidae	<i>Anolis sabanus</i>	yes	Leeward Islands	Saba	13.00	Oceanic	18	2.43	1.00	20	0.27	5.41	Schwartz and Henderson 1991, Lazell 1972, Powell et al. 2005, Henderson and Powell 2009, Schall and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Staats 2002
Dactyloidae	<i>Anolis sagrei</i>	no	NA	NA	NA	mainland	19	2.30	1.50	20	0.09	2.66	Perry and Garland 2002, Cox et al. 2003, Campbell 1999, Stafford and Meyer 2000, Conant and Collins 1998, Lee 2000, Kohler 2003, Schettino 1999, Smith 1946, Schwartz and Henderson 1991, Beebe 1944b, Rogner 1997a, Stuart 1955, Williams 1983, Mccranie et al. 2005, Radder et al. 2008, Schoener and Schoener 1971, Lopez and Gonzalez 1997, Sinervo et al. 2010, Duellman 1965, Henderson and Powell 2009, Jensen et al. 2008, Losos 2009, Schettino et al. 2010, Oliver 1948, Damuth 1987, Ruibal 1961, Losos et al. 1993
Dactyloidae	<i>Anolis trinitatis</i>	yes	Lesser Antilles	St. Lucia	639.80	Oceanic	13	2.09	1.00	NA	0.14	NA	Cox et al. 2003, Schwartz and Henderson 1991, Licht and Gorman 1970, Rogner 1997a, Lazell 1972, Murphy 1997, Todd 2008, Malhotra and Thorpe 1999, Kohler 2005, Henderson and Powell 2009, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Savage 2002, Kohler 2003, Fitch 1973, Dunham et al. 1988, Pounds 1988, Turner 1977, Losos 2009, van Berkum 1986, van Berkum 1988
Dactyloidae	<i>Anolis tropidolepis</i>	no	NA	NA	NA	mainland	10	2.78	1.00	9	0.19	1.73	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Dactyloidae	<i>Anolis valencienni</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	4.52	1.00	10	0.27	2.70	Perry and Garland 2002, Schwartz and Henderson 1991, Herrel et al. 2004, Williams 1983, Rand 1968, Schoener and Schoener 1971, Henderson and Powell 2009, Losos 2009, Johnson et al. 2010
Dactyloidae	<i>Anolis wattsi</i>	yes	Trinidad and Tobago	Trinidad	5008.70	Land bridge	17	1.70	1.00	NA	NA	NA	Schwartz and Henderson 1991, Lazell 1972, Powell et al. 2005, Kolbe et al. 2008, Henderson and Powell 2009, Henderson and Powell 2009, Losos 2009, Daltry 2009, Fitch 1970, Cogger 2000, Wilson and Swan 2003, Greer 1989, Werner and Seifan 2006, Hoskin and Higgie 2008, Bustard 1967, Doughty 1997, Kohler 2005, Henkel 2010, Rosler 2005, Swanson 2007, Wilson 2005, Schlesinger and Shine 1994, Henkel and Schmidt 1995, Werner et al. 1993
Diplodactylidae	<i>Amalosia lesueurii</i>	no	NA	NA	NA	mainland	-31	4.85	2.00	1.5	0.41	1.24	Bauer and Sadlier 2000, Bauer and Vindum 1990, Daza et al. 2009
Diplodactylidae	<i>Bavayia crassicollis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	8.90	2.00	NA	NA	NA	Bauer and DeVaney 1987, Rogner 1997a, Bauer and Sadlier 2000, Bauer and Vindum 1990, Kohler 2005, Daza et al. 2009, Werner et al. 1993
Diplodactylidae	<i>Bavayia cyclura</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	3.25	2.00	NA	NA	NA	Bauer and Sadlier 2000, Daza et al. 2009, Whittaker et al. 2004
Diplodactylidae	<i>Bavayia exsuccida</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	1.95	2.00	NA	NA	NA	Wright et al. 2000, Bauer and Sadlier 2000, Daza et al. 2009
Diplodactylidae	<i>Bavayia geitaina</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	5.54	2.00	NA	NA	NA	Wright et al. 2000, Bauer and Sadlier 2000, Daza et al. 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Diplodactylidae	<i>Bavayia montana</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	8.23	1.50	NA	NA	NA	Bauer and Sadlier 2000, Bauer and Vindum 1990, Kohler 2005, Rosler 2005, Henkel and Schmidt 1995, Whittaker et al. 2004
Diplodactylidae	<i>Bavayia ornata</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	6.16	2.00	NA	NA	NA	Bauer and Sadlier 2000
Diplodactylidae	<i>Bavayia pulchella</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	2.21	2.00	NA	NA	NA	Bauer and Sadlier 2000
Diplodactylidae	<i>Bavayia sauvagii</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	3.28	2.00	NA	NA	NA	Bauer and DeVaney 1987, Rogner 1997a, Bauer and Sadlier 2000, Bauer and Vindum 1990, Kohler 2005, Daza et al. 2009, Henkel and Schmidt 1995
Diplodactylidae	<i>Bavayia septuiclavus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	1.95	2.00	NA	NA	NA	Bauer and Sadlier 2000, Sadlier 1988
Diplodactylidae	<i>Correlophus ciliatus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	28.82	2.00	NA	1.20	NA	Badger 2003, Bauer and Sadlier 2000, Uetz 2006, Kohler 2005, Rosler 2005, Daza et al. 2009
Diplodactylidae	<i>Correlophus sarasinorum</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	41.06	2.00	NA	NA	NA	Rogner 1997a, Bauer and Sadlier 2000, Boulenger 1897, Kohler 2005, Rosler 2005, Daza et al. 2009
Diplodactylidae	<i>Crenadactylus ocellatus</i>	no	NA	NA	NA	mainland	-23	0.61	2.00	2	0.06	0.26	Cogger 2000, Withers et al. 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Wilson and Swan 2008, Storr et al. 1990, Chapman and Dell 1985, Dixon and Kluge 1964, Kohler 2005, Henkel 2010, Bush et al. 2010, Rosler 2005, Bush 1992, Wilson and Swan 2010, Wilson 2005, Bush et al. 2007, Gordon et al. 2010
Diplodactylidae	<i>Dierogecko insularis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	1.01	2.00	NA	NA	NA	Bauer et al. 2006
Diplodactylidae	<i>Dierogecko nehoueensis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	0.97	2.00	NA	NA	NA	Bauer et al. 2006

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Diplodactylidae	<i>Dierogekko poumensis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	0.82	2.00	NA	NA	NA	Bauer et al. 2006
Diplodactylidae	<i>Dierogekko validiclavus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	1.50	2.00	NA	NA	NA	Bauer and Sadlier 2000, Bauer et al. 2006, Sadlier 1988
Diplodactylidae	<i>Diplodactylus tessellatus</i>	no	NA	NA	NA	mainland	-27	2.97	2.00	2	0.23	0.92	Clobert et al. 1998, Cogger 2000, Wilson and Swan 2003, Henle 1991, Warne and Charnov 2008, Henle 1990, Read 1999, Henkel 2010, Rosler 2005, Daza et al. 2009, Michael and Lindenmayer 2010, Swanson 2007
Diplodactylidae	<i>Eurydactylodes agricolae</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	2.64	2.00	NA	0.29	NA	Henkel and Bohme 2001, Rosler 2005
Diplodactylidae	<i>Eurydactylodes symmetricus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	4.12	2.00	NA	NA	NA	Bauer and Sadlier 2000, Kohler 2005, Rosler 2005
Diplodactylidae	<i>Eurydactylodes vieillardii</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	2.49	2.00	NA	0.27	NA	Bauer and Vindum 1990, Bauer et al. 2009, Kohler 2005, Daza et al. 2009, Henkel and Schmidt 1995, Whittaker et al. 2004
Diplodactylidae	<i>Hesperoedura reticulata</i>	no	NA	NA	NA	mainland	-33	4.98	2.06	1	0.33	0.68	Clobert et al. 1998, Cogger 2000, Withers et al. 2000, Wilson and Swan 2003, Henle 1991, Cree 1994, Werner and Seifan 2006, Dunham et al. 1988, Warne and Charnov 2008, Storr et al. 1990, Chapman and Dell 1985, Angilletta and Werner 1998, How and Kitchener 1983, Henkel 2010, Bush et al. 2010, Daza et al. 2009, Heatwole and Taylor 1987, Henle 1990, Kitchener et al. 1988, Werner et al. 1993

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Diplodactylidae	<i>Lucasium damaeum</i>	no	NA	NA	NA	mainland	-29	2.84	2.00	2	0.20	0.80	Clobert et al. 1998, Cogger 2000, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Wilson and Swan 2008, Warne and Charnov 2008, Storr et al. 1990, Henle 1990, Read 1999, Henkel 2010, Wilson and Swan 2010, Daza et al. 2009, Michael and Lindenmayer 2010, Wilson 2005, Swanson 2007, Vucko 2008, Gordon et al. 2010, Swan and Watharow 2005, Werner et al. 1993
Diplodactylidae	<i>Mniarogekko chahoua</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	57.42	2.00	NA	1.20	NA	Bauer 1985, Rogner 1997a, Bauer and Sadlier 2000, Bauer and Vindum 1990, Kohler 2005, Rosler 2005, Henkel and Schmidt 1995, Whittaker et al. 2004, Werner et al. 1993
Diplodactylidae	<i>Oedodera marmorata</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	3.40	2.00	NA	NA	NA	Bauer et al. 2006
Diplodactylidae	<i>Oedura castelnaui</i>	no	NA	NA	NA	mainland	-17	17.09	2.00	1	0.88	1.76	Fitch 1970, Cogger 2000, Wilson and Swan 2003, Greer 1989, Wilson and Swan 2008, Schostakowski 2001, Bustard 1967, Kohler 2005, Henkel 2010, Rosler 2005, Swanson 2007, Wilson 2005, Henkel and Schmidt 1995
Diplodactylidae	<i>Oedura monilis</i>	no	NA	NA	NA	mainland	-24	9.53	2.00	1.5	0.74	2.22	Clobert et al. 1998, Dunham and Miles 1985, Cogger 2000, Wilson and Swan 2003, Henle 1991, Rogner 1997a, Greer 1989, Cree 1994, Warne and Charnov 2008, Bustard 1967, Bustard 1968, Kohler 2005, Henkel 2010, Rosler 2005, Heatwole and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Taylor 1987, Swanson 2007, Wilson 2005, Henle 1990, Bustard 1968b
Diplodactylidae	<i>Oedura tryoni</i>	no	NA	NA	NA	mainland	-28	9.60	2.00	2	0.81	3.23	Fitch 1970, Cogger 2000, Wilson and Swan 2003, Greer 1989, Dunham et al. 1988, Hoskin and Higgie 2008, Warne and Charnov 2008, Bustard 1967, Kohler 2005, Henkel 2010, Rosler 2005, Daza et al. 2009, Swanson 2007, Wilson 2005
Diplodactylidae	<i>Rhacodactylus auriculatus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	30.73	2.00	4.5	1.39	12.55	Cooper and Vitt 2002, Bauer and DeVaney 1987, Rogner 1997a, Bauer and Sadlier 2000, Bauer and Vindum 1990, Bauer 1990, Kohler 2005, Rosler 2005, Daza et al. 2009, Whittaker et al. 2004, Werner et al. 1993
Diplodactylidae	<i>Rhacodactylus leachianus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	150.27	1.50	6	4.16	37.42	Pianka and Vitt 2003, Bauer 1985, Bauer and DeVaney 1987, Bauer and Sadlier 2000, Werner and Seifan 2006, Bauer 1990, Kohler 2005, Rosler 2005, Daza et al. 2009, Honegger 1969, Henkel and Schmidt 1995, Bauer and Sadlier 1994, Werner et al. 1993
Diplodactylidae	<i>Rhacodactylus trachyrhynchus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	66.88	2.00	NA	3.48	NA	Pianka and Vitt 2003, Bauer 1985, Rogner 1997a, Bauer and Sadlier 2000, Daza et al. 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Diplodactylidae	<i>Strophurus ciliaris</i>	no	NA	NA	NA	mainland	-23	7.21	2.00	6	0.51	6.14	Cogger 2000, Cox et al. 2003, Withers et al. 2000, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, How et al. 1986, Henle 1991, Rogner 1997a, Cree 1994, Vitt and Price 1982, Huey et al. 2001, Storr et al. 1990, Read 1999, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Rosler 2005, Daza et al. 2009, Huey and Pianka 2007, Werner and Whitaker 1978, Heatwole and Taylor 1987, Swanson 2007, Wilson 2005, Morton and James 1988, Henle 1990, Sadlier 1990
Diplodactylidae	<i>Strophurus elderi</i>	no	NA	NA	NA	mainland	-26	1.55	2.00	2	0.13	0.52	Cogger 2000, Cox et al. 2003, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, How et al. 1986, Henle 1991, Greer 1989, Werner and Seifan 2006, Wilson and Swan 2008, Vitt and Price 1982, Huey et al. 2001, Storr et al. 1990, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Rosler 2005, Wilson and Swan 2010, Daza et al. 2009, Werner and Whitaker 1978, Heatwole and Taylor 1987, Swanson 2007, Werner 1976, Wilson 2005, Bush et al. 2007, Gordon et al. 2010, Werner et al. 1993
Diplodactylidae	<i>Strophurus intermedius</i>	no	NA	NA	NA	mainland	-32	5.43	2.00	2	0.36	1.45	Cogger 2000, Cox et al. 2003, Wilson and Swan 2003, How et al. 1986, Henle 1991, Greer 1989, Wilson and Swan 2008, Storr et al. 1990, Henkel 2010, Daza et al. 2009,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Michael and Lindenmayer 2010, Swanson 2007, Michael et al. 2011, Bush et al. 2007, Swan and Watharow 2005
Diplodactylidae	<i>Strophurus williamsi</i>	no	NA	NA	NA	mainland	-25	5.15	2.00	3	0.33	1.99	Cogger 2000, Wilson and Swan 2003, Henle 1991, Rogner 1997a, Greer 1989, Bustard 1968, Kohler 2005, Henkel 2010, Rosler 2005, Swanson 2007, Wilson 2005, Bustard 1968, Werner et al. 1993
Diplodactylidae	<i>Toropuku stephensi</i>	yes	New Zealand	Maud	3.09	Land bridge	-40	8.04	1.93	0.4	NA	NA	Robb 1980, Uetz 2006, Hare 2005, Jewell 2008, Iburguengoytia and Casalins 2007, Hare and Cree 2005, Hare et al. 2010
Diplodactylidae	<i>Tukutuku rakiurae</i>	yes	New Zealand	Stewart	1814.70	Land bridge	-47	4.92	2.00	0.415	NA	NA	Thomas 1981, Jewell 2008
Eublepharidae	<i>Aeluroscalabotes felinus</i>	no	NA	NA	NA	mainland	3	23.95	2.00	2	1.75	7.00	Fitch 1970, Inger and Greenberg 1966, Taylor 1963, Kratochvil and Frynta 2006, Das 2004, Seuffer et al. 2005, Cox et al. 1998, Cree 1994, Uetz 2006, Inger and Lian 1996, Das 2010, Cox et al. 2010, Das 2011, Rosler 2005, Grismer 2011, Grismer 2006
Eublepharidae	<i>Coleonyx brevis</i>	no	NA	NA	NA	mainland	29	2.62	2.00	2.5	0.17	0.85	Fitch 1970, Conant and Collins 1998, Stebbins 2003, Dial and Grismer 1992, Dial 1975, Kratochvil and Frynta 2006, Degenhardt et al. 1996, Smith 1946, Rogner 1997a, Seuffer et al. 2005, Werler 1951, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													and Lovich 2009, Rosler 2005, Daza et al. 2009, Huey et al. 1989, Avery 1982, Brattstrom 196
Eublepharid ae	<i>Coleonyx elegans</i>	no	NA	NA	NA	mainland	18	12.37	2.00	3	0.88	5.25	Fitch 1970, Campbell 1999, Stafford and Meyer 2000, Lee 2000, Dial and Grismer 1992, Kohler 2003, Kratochvil and Frynta 2006, Seufer et al. 2005, Kohler et al. 2006, Davis and Dixon 1961, Davis and Smith 1953, Sinervo et al. 2010, Kohler 2005, Duellman 1965, Rosler 2005
Eublepharid ae	<i>Coleonyx reticulatus</i>	no	NA	NA	NA	mainland	28	13.23	2.00	2	1.03	4.14	Conant and Collins 1998, Dial and Grismer 1992, Dial 1975, Kratochvil and Frynta 2006, Seufer et al. 2005, Lemos-Espinal and Smith 2007, Sinervo et al. 2010, Jones and Lovich 2009, Clobert et al. 1998, Fitch 1970, Vitt et al. 1978, Stebbins 2003, Grismer 2002, Pianka 1986, Dial and Grismer 1992, Kratochvil and Frynta 2006, Degenhardt et al. 1996, Smith 1946, Rogner 1997a, Seufer et al. 2005, Cree 1994, Van Denburgh 1922, Andrews and Pough 1980, Vitt and Price 1982, Warne and Charnov 2008, Huey et al. 2001, Parker 1972, Hardy and McDiarmid 1969, McElroy et al. 2008, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009,
Eublepharid ae	<i>Coleonyx variegatus</i>	no	NA	NA	NA	mainland	32	3.84	2.00	2	0.31	1.22	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Eublepharid ae	<i>Eublepharis macularius</i>	no	NA	NA	NA	mainland	27	35.92	2.50	2.5	3.01	18.78	Rosler 2005, Daza et al. 2009, Case 1975, Huey and Pianka 2007, Werner and Whitaker 1978, Carey and Judge 2000, Werner 1976, Huey et al. 1989, Brennan and Holycross 2009, Cunningham 1966, Henkel and Schmidt 1995, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006, Werner et al. 1993, Fitch 1970, Smith 1935, Anderson 1999, Minton 1966, Szczerbak and Golubev 1996, Daniel 1983, Tikader and Sharma 1992, Dial and Grismer 1992, Kratochvil and Frynta 2006, Rogner 1997a, Seufer et al. 2005, Anderson and Leviton 1969, Khan 2006, Anderson 1963, Garrick 2008, Bauer 1990, McElroy et al. 2008, Das 2002, Sinervo et al. 2010, Kohler 2005, De Magalhaes and Costa 2009, Rosler 2005, Daza et al. 2009, Henkel and Schmidt 1995, Masroor 2012, Werner et al. 1993, Szczerbak 2003, Anderson 1999, Szczerbak and Golubev 1996, Kratochvil and Frynta 2006, Seufer et al. 2005, Ananjeva et al. 2006, Bauer 1990, Kohler 2005, Rosler 2005, Daza et al. 2009, Seufer et al. 2005, Kratochvil, pers. Comm. To Shai Meiri (September 2009), Das 2010, Rosler 2005
Eublepharid ae	<i>Eublepharis turcomenicus</i>	no	NA	NA	NA	mainland	38	45.40	2.00	2	6.70	26.80	
Eublepharid ae	<i>Goniurosaurus araneus</i>	no	NA	NA	NA	mainland	23	27.76	2.00	3	1.81	10.86	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Eublepharidae	<i>Goniurosaurus hainanensis</i>	yes	None	Hainan	33209.80	Land bridge	20	17.11	2.00	NA	NA	NA	Seufer et al. 2005
Eublepharidae	<i>Goniurosaurus kuroiwae</i>	yes	Ryukyu Islands	Okinawajima	1199.50	Oceanic	27	13.06	2.00	3	0.55	3.32	Goris and Maeda 2004, Dial and Grismer 1992, Ota 1989, Tanaka and Nishihira 1989, Seufer et al. 2005, Sinervo et al. 2010, Kohler 2005, Henkel and Schmidt 1995, Werner et al. 2006
Eublepharidae	<i>Goniurosaurus splendens</i>	yes	Ryukyu Islands	Tokunoshima	248.87	Oceanic	28	8.73	2.00	NA	0.53	NA	Seufer et al. 2005, Rosler 2005
Gekkonidae	<i>Ailuronyx seychellensis</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	16.14	1.50	NA	0.69	NA	Henkel and Schmidt 2000, Rogner 1997a, Gerlach and Canning 1996, Cheke 1984, Bowler 2006, Gerlach 2004, Schonecker 2008, Kohler 2005, Rosler 2005, Gerlach 2008, Brooke and Houston 1983, Evans and Evans 1980, Henkel and Schmidt 1995, Werner et al. 1993
Gekkonidae	<i>Ailuronyx tachyscopaeus</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	6.82	1.50	NA	0.69	NA	Gerlach and Canning 1996, Bowler 2006, Gerlach 2004, Schonecker 2008, Rosler 2005, Gerlach 2008
Gekkonidae	<i>Ailuronyx trachygaster</i>	yes	Seychelles Islands	Praslin	26.00	Continental	-4	68.63	NA	NA	2.94	NA	Bowler 2006, Gerlach 2004, Gerlach 2002, Schonecker 2008, Gerlach 2008
Gekkonidae	<i>Alsophylax laevis</i>	no	NA	NA	NA	mainland	39	0.90	1.50	2	0.16	0.48	Szczerbak 2003, Rogner 1997a, Szczerbak and Golubev 1996, Kohler 2005, Rosler 2005
Gekkonidae	<i>Alsophylax loricatus</i>	no	NA	NA	NA	mainland	41	0.59	1.50	1.5	0.12	0.27	Szczerbak 2003, Szczerbak and Golubev 1996
Gekkonidae	<i>Alsophylax pipiens</i>	no	NA	NA	NA	mainland	47	0.95	1.50	1.5	0.14	0.32	Szczerbak 2003, Szczerbak and Golubev 1996, Rogner 1997a, Terbish et al. 2006, Kohler 2005, Rosler 2005
Gekkonidae	<i>Alsophylax szczerbaki</i>	no	NA	NA	NA	mainland	41	0.56	1.50	1.5	0.12	0.27	Szczerbak 2003, Szczerbak and Golubev 1996, Rosler 2005

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Chondrodactylus angulifer</i>	no	NA	NA	NA	mainland	-27	9.93	1.81	3	1.32	7.16	Fitzsimons 1943, Branch 1998, Pianka 1986, Loveridge 1947, Parker 1936, Rogner 1997a, Auerbach 1987, Cree 1994, Graham and Marais 2007, Vitt and Price 1982, Huey et al. 2001, Pianka and Huey 1978, Bauer 1990, Sinervo et al. 2010, Kohler 2005, Goldberg 2006, Pianka 1971, Rosler 2005, Daza et al. 2009, Huey and Pianka 2007, Werner and Whitaker 1978, Henkel and Schmidt 1995, Brattstrom 1965
Gekkonidae	<i>Christinus guentheri</i>	yes	None	Norfolk	36.80	Oceanic	-32	6.56	1.30	NA	0.60	NA	Cogger 2000, Pianka and Vitt 2003, Wilson and Swan 2003, Greer 1989, Cogger et al. 1983, Wilson and Swan 2008, Kohler 2005, Henkel 2010, Wells and Wellington 1983, Wilson and Swan 2010, Daza et al. 2009, King and Horner 1993
Gekkonidae	<i>Christinus marmoratus</i>	no	NA	NA	NA	mainland	-34	3.07	1.77	1	0.12	0.21	Cogger 2000, Withers et al. 2000, Wilson and Swan 2003, Greer 1989, Cree 1994, Wilson and Swan 2008, Storr et al. 1990, Fischer and Lindenmayer 2005, Angilletta and Werner 1998, Doughty and Thompson 1998, How et al. 1987, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Michael et al. 2010, Bush et al. 2010, Rosler 2005, Wilson and Swan 2010, Michael and Lindenmayer 2010, Werner and Whitaker 1978, Heatwole and Taylor 1987, Swanson 2007, Michael et al. 2011,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Light et al. 1966, Swan and Watharow 2005
Gekkonidae	<i>Cnemaspis affinis</i>	yes	None	Pinang	295.30	Land bridge	5	2.52	2.00	NA	NA	NA	Taylor 1963, de Rooij 1915, Das 2010, Grismer 2011
Gekkonidae	<i>Cnemaspis limi</i>	yes	Seribuat Archipelago	Tioman	132.00	Land bridge	3	4.75	1.50	NA	NA	NA	Das and Grismer 2003, Grismer and Pan 2008, Grismer et al. 2009, Das 2010, Grismer 2011, Grismer 2011b, Grismer 2006
Gekkonidae	<i>Cnemaspis modiglianii</i>	yes	Mentawai Archipelago	Enggano	397.00	Oceanic	-5	0.82	NA	NA	0.23	NA	Das 2005, Grismer and Das 2006, Grismer and Onn 2008, Grismer and Tri 2007, Grismer et al. 2008b, Grismer et al. 2010, Das 2010
Gekkonidae	<i>Cnemaspis monachorum</i>	yes	None	Langkawi	363.00	Land bridge	6	0.81	1.00	NA	NA	NA	Grismer et al. 2009, Grismer 2011
Gekkonidae	<i>Cnemaspis perhentianensis</i>	yes	Perhentian Islands	Perhentian Besar	11.51	Land bridge	6	2.17	2.00	NA	NA	NA	Grismer and Onn 2008, Das 2010, Grismer 2011
Gekkonidae	<i>Cnemaspis whittendorum</i>	yes	Mentawai Archipelago	Siberut	3828.50	Oceanic	-1	0.79	NA	NA	0.08	NA	Das 2010
Gekkonidae	<i>Crossobamon eversmanni</i>	no	NA	NA	NA	mainland	40	2.22	1.50	2.5	0.35	1.31	Szczerbak 2003, Anderson 1999, Minton 1966, Szczerbak and Golubev 1996, Rogner 1997a, Khan 2006, Sharma 2002, Clark 1990, Kohler 2005, Rosler 2005
Gekkonidae	<i>Cyrtodactylus jarakensis</i>	yes	None	Jarak	14.90	Land bridge	4	6.31	2.00	NA	NA	NA	Grismer et al. 2008c, Das 2010, Grismer 2011
Gekkonidae	<i>Cyrtodactylus laevigatus</i>	yes	Lesser Sunda Islands	Komodo	330.00	Oceanic	-9	1.96	2.00	NA	0.41	NA	Darevsky 1964, Auffenberg 1980
Gekkonidae	<i>Cyrtodactylus leegrimeri</i>	yes	None	Tenggol	4.03	Land bridge	5	14.26	2.00	NA	NA	NA	Onn and Ahmad 2010, Grismer 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Cyrtodactylus peguensis</i>	no	NA	NA	NA	mainland	0	10.93	1.50	3.5	0.57	2.98	Taylor 1963, Bauer et al. 2003, Rogner 1997a, Cox et al. 1998, Sharma 2002, Kohler 2005, Das 2010, Cox et al. 2010, Rosler 2005, Henkel and Schmidt 1995
Gekkonidae	<i>Cyrtodactylus redimiculus</i>	yes	Philippine Islands	Palawan	12188.60	Continental	9	11.16	2.00	NA	NA	NA	Brown and Alcalá 1978, Alcalá 1986
Gekkonidae	<i>Cyrtodactylus sadleiri</i>	yes	None	Christmas	135.00	Oceanic	-10	10.30	2.00	NA	NA	NA	Cogger 2000, Wells 2002, Henkel 2010
Gekkonidae	<i>Cyrtodactylus seribuatensis</i>	yes	None	Seribuat	5.42	Land bridge	2	6.11	2.00	NA	1.42	NA	Youmans and Grismer 2006, Grismer and Pan 2008, Grismer 2011, Grismer 2011b, Grismer 2006
Gekkonidae	<i>Cyrtodactylus tiomanensis</i>	yes	Seribuat Archipelago	Tioman	132.00	Land bridge	3	11.67	2.00	NA	NA	NA	Grismer and Pan 2008, Das 2010, Grismer 2011, Grismer 2011b, Grismer 2006
Gekkonidae	<i>Cyrtopodion caspium</i>	no	NA	NA	NA	mainland	40	3.72	1.50	2	0.27	0.82	Szczerbak 2003, Szczerbak and Golubev 1996, Rogner 1997a, Anderson and Leviton 1969, Clark 1990, Ahmadzadeh et al. 2008, Bauer 1990, Kohler 2005, Rosler 2005, Daza et al. 2009, Trubcheninova et al. 1977, Arakelyan et al. 2011, Henkel and Schmidt 1995
Gekkonidae	<i>Cyrtopodion fedtschenkoi</i>	no	NA	NA	NA	mainland	37	4.40	1.78	2	0.29	1.04	Szczerbak 2003, Minton 1966, Szczerbak and Golubev 1996, Tikader and Sharma 1992, Anderson and Leviton 1969, Sharma 2002, Kohler 2005, Rosler 2005

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Gehyra variegata</i>	no	NA	NA	NA	mainland	-29	2.70	1.50	1.5	0.33	0.74	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Shine and Greer 1991, Cogger 2000, Withers et al. 2000, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Cree 1994, Marquet et al. 1990, Dunham et al. 1988, Wilson and Swan 2008, Vitt and Price 1982, Warne and Charnov 2008, Huey et al. 2001, Chapman and Dell 1985, Angilletta and Werner 1998, Henle 1990, Bustard 1968, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Frankenberg 1978, Bush et al. 2010, Rosler 2005, Wilson and Swan 2010, Daza et al. 2009, Huey and Pianka 2007, Michael and Lindenmayer 2010, King and Horner 1993, Werner and Whitaker 1978, Heatwole and Taylor 1987, Swanson 2007, Henle 1990, Bustard 1968, Gordon et al. 2010, Moro and MacAulay 2010, Kitchener et al. 1988, Swan and Watharow 2005, Werner et al. 1993
Gekkonidae	<i>Gekko athymus</i>	yes	Philippine Islands	Palawan	12188.60	Continental	9	20.50	NA	NA	0.92	NA	Brown and Alcalá 1978, Alcalá 1986
Gekkonidae	<i>Gekko ernstkelleri</i>	yes	Philippine Islands	Panay	12011.10	Oceanic	12	10.01	2.00	NA	1.06	NA	Roselr et al. 2006, Gaulke 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Gekko gecko</i>	no	NA	NA	NA	mainland	23	39.14	2.00	2	2.10	8.41	Fitch 1970, Smith 1935, Conant and Collins 1998, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Taylor 1963, Das 2004, Teynie 2004, Manthey and Grossmann 1997, Rogner 1997a, Ziegler 2002, Cox et al. 1998, Brown and Alcala 1978, Brown et al. 1996, Taylor 1922, Karsen et al. 1986, Inger and Colwell 1977, Werner and Seifan 2006, Karsen et al. 1998, Pauwels et al. 2003, Andrews and Pough 1980, Garrick 2008, Radder et al. 2008, Bauer 1990, Das 2002, Kohler 2005, Das 2010, Cox et al. 2010, Alcala 1986, Das 2011, Rosler 2005, Ahmed et al. 2009, Daza et al. 2009, Grismer 2011, Auffenberg 1980, Kaiser et al. 2011, Werner 1976, Teynie and David 2010, Van Wilgen and Richardson 2012, Honegger 1969, Henkel and Schmidt 1995, Gaulke 2011, Ahmed 2009, Werner et al. 1993, Goris and Maeda 2004, Ota 1989, Huang 2006, Taylor 1962, Huang 2007, Zhang et al. 2009, Huang 2010
Gekkonidae	<i>Gekko hokouensis</i>	no	NA	NA	NA	mainland	28	7.52	2.00	1.5	0.75	2.25	Fitch 1970, Goris and Maeda 2004, Ota 1989, Stejneger 1907, Song 1987, Zhang et al. 2009, Das 2010, Ikeuchi 2004
Gekkonidae	<i>Gekko japonicus</i>	no	NA	NA	NA	mainland	30	5.99	2.00	1.5	0.53	1.58	
Gekkonidae	<i>Gekko shibatai</i>	yes	Ryukyu Islands	Takarajima	7.14	Oceanic	29	6.14	2.00	NA	0.58	NA	Toda et al. 2008

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Gekko smithii</i>	no	NA	NA	NA	mainland	4	56.64	2.00	12	3.20	76.80	Tikader and Sharma 1992, Taylor 1963, Das 2004, Rogner 1997a, Cox et al. 1998, Das 2002, Kohler 2005, Onn et al. 2010, Das 2010, Cox et al. 2010, Goldberg 2009, Das 2011, Rosler 2005, Grismer 2011, Grismer 2011b, Werner et al. 1993
Gekkonidae	<i>Gekko vertebralis</i>	yes	Ryukyu Islands	Amami Oshima	712.35	Oceanic	28	4.56	NA	NA	0.33	NA	Toda et al. 2008
Gekkonidae	<i>Gekko yakuensis</i>	yes	Japan Islands	Kyushu	37437.20	Land bridge	31	6.57	1.50	NA	0.67	NA	Goris and Maeda 2004, Ota 1989
Gekkonidae	<i>Hemidactylus bowieri</i>	yes	Cape Verde	Santo Antao	787.30	Oceanic	16	1.98	NA	NA	0.19	NA	Loveridge 1947, Arnold et al. 2008, Werner et al. 1993
Gekkonidae	<i>Hemidactylus bowringii</i>	no	NA	NA	NA	mainland	23	2.33	1.90	1.5	0.18	0.50	Goris and Maeda 2004, Schleich and Kastle 2002, Tikader and Sharma 1992, Ota 1989, Teynie 2004, Karsen et al. 1986, Zug et al. 1998, Zug et al. 2007, McMahan and Zug 2007, Xu and Ji 2007, Das 2002, Das 2010, Teynie and David 2010, Ahmed 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Hemidactylus brookii</i>	no	NA	NA	NA	mainland	19	2.74	2.00	2.5	0.34	1.71	Fitch 1970, Spawls et al. 2002, Minton 1966, Loveridge 1936, Schleich and Kastle 2002, Daniel 1983, Geniez et al. 2004, Tikader and Sharma 1992, Dunger 1968, Greenbaum and Carr 2005, Joger and Lambert 1996, Das 2004, Papenfuss 1969, Loveridge 1947, Schwartz and Henderson 1991, Lanza and Vanni 1976, Bauer 2006, Hughes 1988, Bohme et al. 1996, Schmidt et al. 1919, Parker 1942, Rogner 1997a, Brown and Alcalá 1978, Taylor 1922, Le Berre 1989, Khan 2006, Deraniyagala 1953, Karsen et al. 1986, Branch 2005, Chirio and LeBreton 2007, Rugiero et al. 2007, Zug et al. 2007, Pauwels and Vande weghe 2008, Reid 1986, Radder et al. 2008, Schonecker 2008, Kohler 2008, Bauer 1990, Somaweera and Somaweera 2009, Das 2002, Kohler 2005, Das 2010, Largen and Spawls 2010, Alcalá 1986, Das 2011, Das and de Silva 2011, Shanbhag 2002, Ahmed et al. 2009, Daza et al. 2009, Grismer 2011, Damuth 1987, Carey and Judge 2000, Avery 1981, Cisse and Karns 1978, Ahmed 2009, Masroor 2012, Werner et al. 1993
Gekkonidae	<i>Hemidactylus dracaenacolus</i>	yes	Socotra Archipelago	Socotra	3606.70	Continent al	13	6.85	2.00	NA	NA	NA	Rosler and Wranik 1999, Razzetti et al. 2011, Rosler and Wranik 2004, Rosler and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Wranik 2005
Gekkonidae	<i>Hemidactylus flaviviridis</i>	no	NA	NA	NA	mainland	25	5.30	2.00	2	0.44	1.75	Fitch 1970, Andeson 1999, Minton 1966, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Leviton et al. 1992, Parker 1932, Loveridge 1947, Gallagher 1971, Arnold 1984, El Din 2006, Parker 1942, Rogner 1997a, Anderson and Leviton 1969, Arnold 1980, Schatti and Desvoignes 1999, Le Berre 1989, Khan 2006, Anderson 1895, Murthy 1995, Jongbloed 2000, Shrestha 2001, Radder et al. 2008, Hornby 1996, Das 2002, Largen and Spawls 2010, Rosler 2005, Werner 1973, Daza et al. 2009, Gholamifard et al. 2010, Rosler and Wranik 2004, Zari 1997, Ahmed 2009, Masroor 2012, Werner et al. 1993, Fitch 1970, 1982, Hendrickson 1966, Cogger 2000, Spawls et al. 2002, Minton 1966, Goris and Maeda 2004, Savage 2002, Grismer 2002, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Taylor 1963, Ota 1989, Rand and Myers 1990, Glaw and Vences 1994, Henkel and Schmidt 2000, Das 2004, Wilson and Swan 2003, Loveridge 1947, Bauer and Gunther 1992, Allison 2006,
Gekkonidae	<i>Hemidactylus frenatus</i>	no	NA	NA	NA	mainland	4	3.25	1.95	14.5	0.19	5.50	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Goonewardene et al. 2003, Rogner 1997a, Bauer and Sadlier 2000, McCoy 1980, Ziegler 2002, Cox et al. 1998, Brown and Alcalá 1978, Brown et al. 1996, Bauer and Vindum 1990, Taylor 1922, Khan 2006, Cogger et al. 1983, Deraniyagala 1953, Cree 1994, Inger and Colwell 1977, Towns 1994, Inger et al. 1984, Vinson and Vinson 1969, Cogger et al. 1983, Werner and Seifan 2006, Malkmus et al. 2002, Mccranie et al. 2005, Kohler et al. 2006, Glaw and Vences 2007, McCoy 2006, Bowler 2006, Pauwels et al. 2003, Zug 1991, Dunham et al. 1988, Gans et al. 1965, Zug et al. 2007, Wilson and Swan 2008, Schwaner 1980, Radder et al. 2008, Schonecker 2008, Kohler 2008, Irschick et al. 1996, Sabath 1981, Krysko et al. 2003, Somaweera and Somaweera 2009, Das 2002, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Morrison 2003, Das 2010, Cox et al. 2010, Alcalá 1986, Das 2011, Rosler 2005, Ahmed et al. 2009, Wilson and Swan 2010, Daza et al. 2009, Grismer 2011, Feder and Feder 1981, Grismer 2011, Rodda and Dean-Bradley 2001, Auffenberg 1980, Chandramouli and Ganesh 2011, Kaiser et al. 2011,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Hemidactylus mabouia</i>	no	NA	NA	NA	mainland	-7	3.68	2.00	6.5	0.24	3.12	Swanson 2007, Huey et al. 1989, Ineich 2011, Meirte 2004, McCoid 1994, Dixon and Lemos-Espinal 2010, Teynie and David 2010, Van Wilgen and Richardson 2012, Werner 1980, Henkel and Schmidt 1995, Gaulke 2011, Ahmed 2009, Werner et al. 1993 Fitch 1970, Spawls et al. 2002, Fitzsimons 1943, Branch 1998, Duellman and Mendelson 1995, Loveridge 1936, Razzetti and Msuya 2002, Avila-Pires 1995, Vitt 1986, Beebe 1944b, Glaw and Vences 1994, Rodda et al. 2001, Joger and Lambert 1996, Henkel and Schmidt 2000, Broadley 1971, Loveridge 1947, Schwartz and Henderson 1991, Bauer 2006, Hughes 1988, Cei 1993, Parker 1936, Jeffery 1993, Dixon and Soini 1986, Schmidt et al. 1919, Hoogmoed 1973, Vitt 2000, Achaval and Olmos 2003, Laurent 1964, Loveridge 1953, Auerbach 1987, Cree 1994, Bartlett and Bartlett 2003, Werner and Seifan 2006, Barbour and Loveridge 1928, Powell et al. 2005, Daudin and de Silva 2007, Branch 2005, Van Buurt 2005, Chirio and LeBreton 2007, Colli et al. 2002, Hodge et al. 2003, Rugiero et al. 2007, Rodrigues 2003, Gans

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Hemidactylus maculatus</i>	no	NA	NA	NA	mainland	13	21.82	2.00	2	0.75	3.01	et al. 1965, Rodrigues 1996, Leache et al. 2006, de Witte 1953, Vonesh 1998, Pauwels and Vande weghe 2008, Murphy 1997, Reid 1986, Vitt et al. 1999, Vitt 1995, Huey et al. 2001, Bartlett and Bartlett 2003, Kohler 2008, Heatwole and Veron 1977, Fuenmayor et al. 2005, Krysko et al. 2003, Bauer 1990, Carreira et al. 2005, Vanzolini et al. 1980, Malonza et al. 2006, Sinervo et al. 2010, Kohler 2005, Henderson and Powell 2009, Ullenbruch et al. 2010, Ugueto and Rivas 2010, Rosler 2005, Daza et al. 2009, Daltry 2009, Haagner et al. 2000, Almeida-Gomes et al. 2008, Gasc 1990, Meirte 2004, Cisse and Karns 1978, Van Wilgen and Richardson 2012, Henkel and Schmidt 1995, Trape et al. 2012, Winck and Rocha 2012, Werner et al. 1993 Daniel 1983, Tikader and Sharma 1992, Rogner 1997a, Deraniyagala 1953, Bauer 1990, Das 2002, Kohler 2005, Das and de Silva 2011, Daza et al. 2009, Taylor 1953, Chandramouli and Ganesh 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Hemidactylus turcicus</i>	no	NA	NA	NA	mainland	31	2.28	2.00	3	0.41	2.49	Schleich et al. 1996, Amitai and Bouskila 2001, Arnold and Ovenden 2004, Anderson 1999, Minton 1966, Disi et al. 2001, Baran and Atatur 1998, Conant and Collins 1998, Flower 1933, Smith 1946, Loveridge 1947, Schwartz and Henderson 1991, Arnold 1984, El Din 2006, Rogner 1997a, Schatti and Desvoignes 1999, Le Berre 1989, Khan 2006, Cree 1994, Anderson 1898, Bons and Geniez 1996, Atatur and Gocmen 2001, Sindaco et al. 2006, Corti and Cascio 2002, Jongbloed 2000, Valakos et al. 2008, Valakos et al. 2004, Moravec & Bohme 1997, Cooper et al. 2001, Werner 1987, Kohler 2008, Hornby 1996, Weber 1960, Lemos-Espinal and Smith 2007, Bauer 1990, Kwet 2009, Lemos-Espinal and Smith 2007b, Malkmus 2004, Sinervo et al. 2010, Kohler 2005, Baier et al. 2009, Frankenberg 1978, Jensen et al. 2008, Rosler 2005, Daza et al. 2009, Werner 1989, Girling et al. 1998, Sindaco et al. 2010, van der Kooij 2001, Degenhardt et al. 1996, Beane et al. 2010, Brennan and Holycross 2009, Bar and Haimovitch 2012, Van Wilgen and Richardson 2012, Henkel and Schmidt 1995, Maso and Pijoan 2011,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Heteronotia binoei</i>	no	NA	NA	NA	mainland	-25	2.59	2.00	2	0.21	0.84	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Cogger 2000, Withers et al. 2000, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997a, Greer 1989, Cree 1994, Werner and Seifan 2006, Dunham et al. 1988, Wilson and Swan 2008, Vitt and Price 1982, Warne and Charnov 2008, Huey et al. 2001, Radder et al. 2008, Storr et al. 1990, Chapman and Dell 1985, Henle 1990, Read 1999, Bustard 1968, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Bush et al. 2010, Wilson and Swan 2010, Daza et al. 2009, King and Horner 1993, Werner and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Whitaker 1978, Heatwole and Taylor 1987, Swanson 2007, Bush et al. 2007, Kutt et al. 2011, Bustard 1968, Sadlier 1990, Light et al. 1966, Gordon et al. 2010, Moro and MacAulay 2010, Heatwole and Butler 1981, Swan and Watharow 2005, Werner et al. 1993
Gekkonidae	<i>Lepidodactylus euaensis</i>	yes	Tonga Islands	Eua	86.70	Oceanic	-21	2.32	2.00	NA	0.26	NA	Gibbons and Brown 1988
Gekkonidae	<i>Lepidodactylus herrei</i>	yes	Philippine Islands	Cebu	4467.50	Oceanic	10	1.96	NA	NA	0.15	NA	http://www.angelfire.com/empire/usc_brg/Cebu_reptiles.htm , Brown and Alcalá 1978, Alcalá 1986
Gekkonidae	<i>Lepidodactylus listeri</i>	yes	None	Christmas	135.00	Oceanic	-10	1.90	2.00	NA	NA	NA	Cogger et al. 1983, Cogger 2000, Henkel 2010 Fitch 1970, Cogger 2000, Perry and Garland 2002, Savage 2002, Zweifel 1980, Kohler 2003, Ota 1989, Rodda et al. 2001, Henkel and Schmidt 2000, Das 2004, Wilson and Swan 2003, Donoso-Barros 1966, Allison 2006, Crombie and Pregill 1999, Loveridge 1948, de Rooij 1915, Rogner 1997a, Bauer and Sadlier 2000, Greer 1989, McCoy 1980, Brown and Alcalá 1978, Bauer and Vindum 1990, Cogger et al. 1983, Deraniyagala 1953, Cree 1994, Towns 1994, Cogger et al. 1983, Buden 2000, McCoy
Gekkonidae	<i>Lepidodactylus lugubris</i>	no	NA	NA	NA	mainland	-8	1.58	1.50	5.5	0.14	1.18	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2006, Zug 1991, Dunham et al. 1988, Wilson and Swan 2008, Vitt and Price 1982, Schwaner 1980, Radder et al. 2008, Schonecker 2008, Kohler 2008, Irschick et al. 1996, Sabath 1981, Somaweera and Somaweera 2009, Das 2002, Sinervo et al. 2010, Kohler 2005, Henkel 2010, Morrison 2003, Das 2010, Alcalá 1986, Das 2011, Das and de Silva 2011, Rosler 2005, Wilson and Swan 2010, Daza et al. 2009, Grismer 2011, Feder and Feder 1981, Grismer 2011, Rodda and Dean-Bradley 2001, Auffenberg 1980, Swanson 2007, Huey et al. 1989, Ineich 2011, McCoid 1994, Van Wilgen and Richardson 2012, Werner 1980, Henkel and Schmidt 1995, Gaulke 2011
Gekkonidae	<i>Lepidodactylus manni</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-18	1.81	2.00	NA	0.13	NA	Zug 1991, Morrison 2003
Gekkonidae	<i>Lepidodactylus vanuatuensis</i>	yes	New Hebrides	Espiritu Santo	3955.50	Oceanic	-16	2.01	2.00	NA	0.15	NA	Ineich 2008, Ota et al. 1998, Ineich 2011
Gekkonidae	<i>Luperosaurus macgregori</i>	yes	Philippine Islands	Calayan	494.53	Oceanic	19	2.36	NA	NA	0.37	NA	Oliveros et al. 2011
Gekkonidae	<i>Lygodactylus klugei</i>	no	NA	NA	NA	mainland	-11	0.67	2.00	1.5	0.08	0.25	Cox et al. 2003, Vitt 1986, Pianka and Vitt 2003, Cree 1994, Dunham et al. 1988, Rodrigues 2003, Rodrigues 1996, Vitt 1995, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler 2005, Rocha et al. 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Mediodactylus kotschy</i>	no	NA	NA	NA	mainland	39	1.59	2.13	3	0.15	0.95	Fitch 1970, Amitai and Bouskila 2001, Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Szczerbak and Golubev 1996, Arnold 1987, Loveridge 1947, Rogner 1997a, Atatur and Gocmen 2001, Sindaco et al. 2006, Corti and Cascio 2002, Valakos et al. 2008, Valakos et al. 2004, Weber 1960, Kwet 2009, Werner 1930, Sinervo et al. 2010, Kohler 2005, Baier et al. 2009, Werner 1993, Frankenberg 1978, Rosler 2005, Werner 1989, Sindaco et al. 2010, Haxhiu 1998, Arnold 1987, Goldberg 2012, Bar and Haimovitch 2012
Gekkonidae	<i>Mediodactylus russowii</i>	no	NA	NA	NA	mainland	42	1.74	1.50	1.5	0.13	0.30	Szczerbak 2003, Anderson 1999, Kohler 2005, Szczerbak and Golubev 1996, Rosler 2005
Gekkonidae	<i>Pachydactylus mariquensis</i>	no	NA	NA	NA	mainland	-31	3.66	2.00	2	0.39	1.57	Fitzsimons 1943, Branch 1998, Loveridge 1947, Graham and Marais 2007, Rosler 2005, Bauer et al. 2011
Gekkonidae	<i>Perochirus ateles</i>	yes	None	Guam	541.00	Oceanic	11	3.08	2.10	NA	0.22	NA	Cox et al. 2003, Goris and Maeda 2004, Buden 2000, Buden 1998, Buden 2007, Sabath 1981, Kohler 2005, Ineich 2011, McCoid 1994, Buden 1999
Gekkonidae	<i>Phelsuma andamanense</i>	yes	Andaman and Nicobar Islands	North Andaman	2780.70	Oceanic	11	2.94	1.50	NA	NA	NA	Smith 1935, Daniel 1983, Tikader and Sharma 1992, Henkel and Schmidt 2000, Rogner 1997a, Lerner 2004, Stoliczka 1873, Schonecker 2008, Das 2002, Kohler 2005,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Rosler 2005
Gekkonidae	<i>Phelsuma astriata</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	2.74	2.00	NA	NA	NA	Henkel and Schmidt 2000, Rogner 1997a, Murphy and Myers 1996, Cheke 1984, Bowler 2006, Schonecker 2008, Kohler 2005, Noble et al. 2011, Brooke and Houston 1983, Evans and Evans 1980, Crawford and Thorpe 1979, Werner et al. 1993
Gekkonidae	<i>Phelsuma borbonica</i>	yes	Mascarene Archipelago	Reunion	2535.20	Oceanic	-16	4.34	2.00	NA	NA	NA	Glaw and Vences 1994, Henkel and Schmidt 2000, Vinson and Vinson 1969, Uetz 2006, Schonecker 2008, Kohler 2005, Rosler 2005
Gekkonidae	<i>Phelsuma cepediana</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-16	2.66	2.00	NA	0.05	NA	Glaw and Vences 1994, Henkel and Schmidt 2000, Rogner 1997a, Vinson and Vinson 1969, Glaw and Vences 2007, Schonecker 2008, Harmon et al. 2007, Kohler 2005, Rosler 2005, Henkel and Schmidt 1995, Werner et al. 1993
Gekkonidae	<i>Phelsuma comorensis</i>	yes	Comoro Islands	Grande Comore	1148.00	Oceanic	-12	3.85	2.00	NA	NA	NA	Henkel and Schmidt 2000, Uetz 2006, Schonecker 2008, Kohler 2005, Rosler 2005, Meirte 2004
Gekkonidae	<i>Phelsuma dubia</i>	no	NA	NA	NA	mainland	-7	3.30	2.00	9.5	0.26	4.87	Spawls et al. 2002, Glaw and Vences 1994, Henkel and Schmidt 2000, Loveridge 1947, Raxworthy and Nussbaum 1994, Rogner 1997a, Lerner 2004, Glaw and Vences 2007, Loveridge 1955, Schonecker 2008, Kohler 2005, Rosler 2005, Meirte 2004

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Phelsuma guentheri</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-20	28.06	1.50	NA	1.27	NA	Glaw and Vences 1994, Henkel and Schmidt 2000, Vinson and Vinson 1969, Uetz 2006, Schonecker 2008, Harmon et al. 2007, Rosler 2005, Bullock 1986, Werner et al. 1993
Gekkonidae	<i>Phelsuma guimbeaui</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-20	2.30	1.50	NA	0.22	NA	Glaw and Vences 1994, Henkel and Schmidt 2000, Rogner 1997a, Vinson and Vinson 1969, Harmon et al. 2007, Kohler 2005, Rosler 2005, Werner et al. 1993
Gekkonidae	<i>Phelsuma inexpectata</i>	yes	Mascarene Archipelago	Reunion	2535.20	Oceanic	-21	2.51	1.50	NA	NA	NA	Henkel and Schmidt 2000, Vinson and Vinson 1969, Schonecker 2008, Kohler 2005, Rosler 2005
Gekkonidae	<i>Phelsuma nigristriata</i>	yes	Comoro Islands	Mayotte	314.00	Oceanic	-13	2.10	1.50	NA	NA	NA	Henkel and Schmidt 2000, Schonecker 2008, Kohler 2005, Meirte 2004
Gekkonidae	<i>Phelsuma ornata</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-20	2.15	1.50	NA	0.17	NA	Glaw and Vences 1994, Henkel and Schmidt 2000, Rogner 1997a, Alvarez 2004, Mertens 1963b, Vinson and Vinson 1969, Bullock et al. 1985, Nyhagen et al. 2001, Whitaker 1987, Schonecker 2008, Harmon et al. 2007, Kohler 2005, Rosler 2005, Bullock 1986
Gekkonidae	<i>Phelsuma parkeri</i>	yes	None	Pemba	889.90	Continental	-5	5.80	1.50	NA	NA	NA	Henkel and Schmidt 2000, Loveridge 1947, Schonecker 2008, Rosler 2005
Gekkonidae	<i>Phelsuma robertmertensi</i>	yes	Comoro Islands	Mayotte	314.00	Oceanic	-13	1.52	1.50	NA	NA	NA	Henkel and Schmidt 2000, Rogner 1997a, Schonecker 2008, Kohler 2005, Rosler 2005, Meirte 2004
Gekkonidae	<i>Phelsuma rosagularis</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-20	3.94	2.00	NA	NA	NA	Henkel and Schmidt 2000, Vinson and Vinson 1969, Schonecker 2008, Harmon et al. 2007, Rosler 2005

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gekkonidae	<i>Phelsuma sundbergi</i>	yes	Seychelles Islands	Praslin	26.00	Continental	-5	8.00	1.50	NA	0.86	NA	Henkel and Schmidt 2000, Rogner 1997a, Cheke 1984, Bowler 2006, Schonecker 2008, Kohler 2005, Rosler 2005, Noble et al. 2011, Gerlach 2008, Werner et al. 1993
Gekkonidae	<i>Phelsuma v-nigra</i>	yes	Comoro Islands	Grande Comore	1148.00	Oceanic	-12	2.37	1.50	NA	0.14	NA	Henkel and Schmidt 2000, Rogner 1997a, Schonecker 2008, Kohler 2005, Rosler 2005, Meirte 2004
Gekkonidae	<i>Pseudogekko smaragdinus</i>	yes	Philippine Islands	Polillo	628.90	Land bridge	15	3.66	2.00	NA	0.37	NA	Brown and Alcalá 1978, Kohler 2005, Alcalá 1986, Rosler 2005
Gekkonidae	<i>Ptenopus garrulus</i>	no	NA	NA	NA	mainland	-27	2.23	1.00	1.5	0.33	0.49	Fitzsimons 1943, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Loveridge 1947, Hibbits et al. 2005, Auerbach 1987, Vitt and Price 1982, Huey et al. 2001, Pianka and Huey 1978, Goldberg 2006, Pianka 1971, Goldberg 2008, Rosler 2005, Daza et al. 2009, Huey and Pianka 2007, Werner and Whitaker 1978, Avery 1982, Hibbits 2005, Brain 1962, Werner et al. 1993, Hibbits et al. 2012
Gekkonidae	<i>Ptychozoon nicobarensis</i>	yes	Andaman and Nicobar Islands	Camorta	188.00	Oceanic	8	15.38	2.00	NA	0.50	NA	Das and Vijayakumar 2009
Gekkonidae	<i>Stenodactylus doriae</i>	no	NA	NA	NA	mainland	25	5.38	2.00	3.5	0.62	4.37	Amitai and Bouskila 2001, Arbel 1984, Anderson 1999, Disi et al. 2001, Arnold 1984, Arnold 1980, Schatti and Desvoignes 1999, Werner and Seifan 2006, Jongbloed 2000, Werner 1987, Hornby 1996, Kohler 2005, Arnold 1980, Frankenberg 1978, Rosler

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2005, Werner 1989, van der Kooij 2001, Bar and Haimovitch 2012, Werner et al. 1993
Gekkonidae	<i>Urocotyledon inexpectata</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-4	1.52	2.00	NA	NA	NA	Loveridge 1947, Cheke 1984, Bowler 2006, Schonecker 2008, Gerlach 2008
Gerrhosauridae	<i>Gerrhosaurus skoogi</i>	no	NA	NA	NA	mainland	-19	75.08	2.00	1	5.04	10.08	Branch 1998, Zug et al. 2001, Cooper and Vitt 2002, Greene 1982, Pianka and Vitt 2003, Rogner 1997b, Nagy et al. 1999, Graham and Marais 2007
Gymnophthalmidae	<i>Cercosaura schreibersii</i>	no	NA	NA	NA	mainland	-19	1.19	2.00	1.5	0.04	0.13	Fitzgerald et al. 1999, Wiens et al. 2006, Avila-Pires 1995, Cei 1986, Cei 1993, Nogueira et al. 2005, Achaval and Olmos 2003, Colli et al. 2002, Vitt 1991, Milstead 1961, Carreira et al. 2005, Kohler 2005, Balestrin 2008, Balestrin et al. 2010, Dos Santos et al. 2012, Doan and Lamar 2012
Gymnophthalmidae	<i>Euspondylus chasqui</i>	no	NA	NA	NA	mainland	-13	5.10	2.00	2	0.21	0.83	Chavez et al. 2011
Gymnophthalmidae	<i>Gymnophthalmus speciosus</i>	no	NA	NA	NA	mainland	9	1.46	1.87	2.5	0.08	0.39	Clobert et al. 1998, Dunham and Miles 1985, Stafford and Meyer 2000, Savage 2002, Lee 2000, Cole et al. 1990, Kohler 2003, Rand and Myers 1990, Duellman 1990, Molina et al. 2004, Mccranie et al. 2005, Kohler et al. 2006, van Buurt 2005, Dunham et al. 1988, Stuart 1939, Warne and Charnov 2008, Murphy 1997, Rand 1957, Kohler 2008,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Gymnophthalmidae	<i>Leposoma rugiceps</i>	no	NA	NA	NA	mainland	8	1.29	2.09	1.5	0.08	0.24	Kohler 2005, Leenders and Watkins-Colwell 2004, Kohler 2003, Dunham et al. 1988, Warne and Charnov 2008, Telford 1971, Kohler 2008, Kohler 2005, Duellman and Mendelson 1995, Avila-Pires 1995, Sherbrooke 1975, Fitch 1970, Shine and Greer 1991, Cox et al. 2003, Uzzell 1966, Duellman 1978, Duellman 1990, Pianka and Vitt 2003, Dixon and Soini 1986, Vitt 2000, Bartlett and Bartlett 2003, Doan 2008, Dunham et al. 1988, Warne and Charnov 2008, Vitt and Zani 1996b, Vitt et al. 1998, Vitt et al. 1999, Huey et al. 2001, Sinervo et al. 2010, Kohler 2005, Anaya-Rojas et al. 2010, Fitch 1968, Rocha et al. 2009, Vitt and Avila-Pires 1998
Gymnophthalmidae	<i>Potamites ecleopus</i>	no	NA	NA	NA	mainland	-6	4.46	2.00	2	0.16	0.64	Murphy 1997
Gymnophthalmidae	<i>Riama shrevei</i>	yes	Trinidad and Tobago	Trinidad	5008.70	Land bridge	11	1.77	2.00	NA	NA	NA	Fitch 1970, Perry and Garland 2002, Zug et al. 2001, Pianka and King 2004, Kohler 2003, Pianka and Vitt 2003, Rogner 1997b, Bogert & Oliver 1945, Duellman 1961, Kohler 2008, Hardy and McDiarmid 1969, Lemos-Espinal and Smith 2007b, Eidenmueller and Philippen 2008, Sinervo et al. 2010, Kohler 2005, Duellman 1965, Van Wilgen and Richardson 2012
Helodermatidae	<i>Heloderma horridum</i>	no	NA	NA	NA	mainland	21	991.06	8.00	1	26.48	211.83	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Helodermatidae	<i>Heloderma suspectum</i>	no	NA	NA	NA	mainland	32	538.49	5.35	1	45.39	242.85	Fitch 1970, Perry and Garland 2002, Stebbins 2003, Zug et al. 2001, Pianka and Vitt 2003, Smith 1946, Rogner 1997b, Van Denburgh 1922, Goldberg and Lowe 1997, Hardy and McDiarmid 1969, Beck 1990, Eidenmueller and Philippen 2008, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Brennan and Holycross 2009, Brattstrom 1965, Stebbins and McGinnis 2012
Iguanidae	<i>Amblyrhynchus cristatus</i>	yes	Galapagos Archipelago	Isabela	4588.00	Oceanic	-1	1403.80	2.25	0.75	59.55	100.49	Tinkle et al. 1970, Clobert et al. 1998, Fitch 1982, Wiewandt 1982, Cooper and Vitt 2002, Pianka and Vitt 2003, Wikelski and Carbone 2004, Ord and Blumstein 2002, Van Denburgh and Slevin 1913, Clusella-Trullas et al. 2008, Fitch 1982, Warne and Charnov 2008, Cree and Guillette 1995, Sinervo et al. 2010, Kohler 2005, Wikelski 2005, Turner 1977, Curry-Lindahl 1979, Brattstrom 1965
Iguanidae	<i>Brachylophus fasciatus</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-18	169.13	3.89	1.5	18.09	105.54	Gibbons and Watkins 1982, Ord and Blumstein 2002, Rogner 1997a, Zug 1991, Stacy et al. 2008, Kohler 2005, Morrison 2003, Carey and Judge 2000, Gibbons 1981
Iguanidae	<i>Brachylophus vitiensis</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-17	371.01	4.00	3	20.75	248.99	Gibbons and Watkins 1982, Zug 1991, Keogh et al. 2008, Kohler 2005, Boylan 1998, Gibbons 1984, Gibbons 1981, Reidpath

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Iguanidae	<i>Conolophus pallidus</i>	yes	Galapagos Archipelago	Barrington	24.00	Oceanic	-1	1590.77	10.00	1	44.26	442.59	Perry and Garland 2002, Fitch 1982, Cooper and Vitt 2002, Christian and Tracy 1985, Van Denburgh and Slevin 1913, Clusella-Trullas et al. 2008, Carpenter 1969, Sinervo et al. 2010, Snell and Christian 1985, Clobert et al. 1998, Fitch 1982, Werner 1982, Ord and Blumstein 2002, Van Denburgh and Slevin 1913, Heller 1903, Warne and Charnov 2008, Carpenter 1969, Costantini et al. 2005, Sinervo et al. 2010, Kohler 2005, Snell and Christian 1985, Werner 1983, Kohler 2003, Kohler 1996, Mccranie et al. 2005, Kohler 2005
Iguanidae	<i>Conolophus subcristatus</i>	yes	Galapagos Archipelago	Isabela	4588.00	Oceanic	-1	2385.51	13.50	1	43.60	588.66	Kohler 2003, Kohler 1996, Mccranie et al. 2005, Kohler 2005, de Queiroz 1990, Kohler 2005
Iguanidae	<i>Ctenosaura bakeri</i>	yes	Bay Islands	Utila	42.08	Oceanic	16	310.22	10.00	NA	NA	NA	Fitch 1970, Conant and Collins 1998, Cooper and Vitt 2002, Kohler 2003, Pianka and Vitt 2003, Alvarez 2004, Duellman 1961, Radder et al. 2008, Evans 1951, Durtsche 2000, Hardy and McDiarmid 1969, Davis and Dixon 1961, Davis and Smith 1953, Kohler 2005, Duellman 1965, Throckmorton 1973, Van Wilgen and Richardson 2012
Iguanidae	<i>Ctenosaura oedirhina</i>	yes	Bay Islands	Roatan	156.60	Oceanic	16	355.06	7.00	NA	NA	NA	
Iguanidae	<i>Ctenosaura pectinata</i>	no	NA	NA	NA	mainland	18	939.96	44.50	1	8.15	362.85	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Iguanidae	<i>Ctenosaura similis</i>	no	NA	NA	NA	mainland	14	1217.14	32.70	1	6.71	219.33	Clobert et al. 1998, Campbell 1999, Stafford and Meyer 2000, Savage 2002, Lee 2000, Fitch 1973, 1982, Wiewandt 1982, Van Devender 1982, Cooper and Vitt 2002, Kohler 2003, Pianka and Vitt 2003, Schwartz and Henderson 1991, Ord and Blumstein 2002, Rogner 1997a, Alvarez 2004, Mccranie et al. 2005, Kohler et al. 2006, Dunham et al. 1988, Garrick 2008, Fitch 1982, Warne and Charnov 2008, Rand 1957, Kohler 2008, Fitch and Hackforth-Jones 1983, Lopez and Gonzalez 1997, Kohler 2005, Fitch and Henderson 1978, Leenders and Watkins-Colwell 2004
Iguanidae	<i>Cyclura carinata</i>	yes	Bahamas	Grand Caicos	289.30	Oceanic	22	516.29	4.30	1	22.63	97.32	Clobert et al. 1998, Perry and Garland 2002, Wiewandt 1982, Cooper and Vitt 2002, Pianka and Vitt 2003, Schwartz and Henderson 1991, Bissell and Martins 2004, Ord and Blumstein 2002, Powell 1999, Alvarez 2004, Dunham et al. 1988, Warne and Charnov 2008, Kohler 2005, Henderson and Powell 2009, Shine and Charnov 1992
Iguanidae	<i>Cyclura collei</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	2303.06	17.00	NA	NA	NA	Schwartz and Henderson 1991, Kohler 2005, Henderson and Powell 2009, Carey 1975
Iguanidae	<i>Cyclura cyclura</i>	yes	Bahamas	North Andros	3439.40	Oceanic	24	3039.76	7.15	1	30.78	220.10	Schwartz and Henderson 1991, Iverson et al. 2006, Iverson et al. 2005,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Iguanidae	<i>Cyclura pinguis</i>	yes	Leeward Islands	Anegada	38.00	Land bridge	19	4101.17	12.30	1	41.05	504.92	Henderson and Powell 2009, Perez-Buitrago et al. 2010, Carey 1975
Iguanidae	<i>Cyclura rileyi</i>	yes	Bahamas	Crooked	282.10	Oceanic	23	492.82	3.10	NA	17.02	NA	Schwartz and Henderson 1991, Hayes et al. 2004, Alvarez 2004, Henderson and Powell 2009
Iguanidae	<i>Dipsosaurus dorsalis</i>	no	NA	NA	NA	mainland	32	65.33	4.50	1	4.69	21.12	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Melville et al. 2006, Perry and Garland 2002, Stebbins 2003, Grismer 2002, Ortega-Rubio et al. 1995, Zug et al. 2001, Wiewandt 1982, Cooper and Vitt 2002, Pianka 1986, Pianka and Vitt 2003, Smith 1946, Tracy 2004, Ord and Blumstein 2002, Pianka 1971, Linsdale 1932, Rogner 1997a, Van Denburgh 1922, Nagy et al. 1999, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Huey et al. 2001, Hardy and McDiarmid 1969, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Case 1975, Huey and Pianka 2007, Muchlinski et al. 1995, Krekorian 1983, Curry-Lindahl 1979, Bury 1982, Banks and Farmer 1963,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Asplund 1967, Brennan and Holycross 2009, Cunningham 1966, Brattstrom 1965, Charnov et al. 2007, Stebbins and McGinnis 2012, Lemm 2006
Iguanidae	<i>Iguana delicatissima</i>	yes	Windward Islands	Martinique	1166.60	Oceanic	17	1879.26	17.00	1.5	18.82	479.98	Fitch 1985, Lazell 1973, Schwartz and Henderson 1991, Hodge et al. 2003, Malhotra and Thorpe 1999, Breuil 2002, Kohler 2005, Henderson and Powell 2009, Lorgelec et al. 2007, Rivero 1998

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Iguanidae	<i>Iguana iguana</i>	no	NA	NA	NA	mainland	-6	1382.40	28.80	1	20.75	597.58	Tinkle et al. 1970, Fitch 1970, 1985, Shine and Greer 1991, Perry and Garland 2002, Campbell 1999, Stafford and Meyer 2000, Conant and Collins 1998, Savage 2002, Lee 2000, Fitch 1973, 1982, Avila-Pires 1995, Wiewandt 1982, Van Devender 1982, Cooper and Vitt 2002, Lazell 1973, Rand and Myers 1990, Zimmerman and Rodrigues 1990, Duellman 1990, Schwartz and Henderson 1991, Evans 1947, Mesquita et al. 2006b, Beebe 1944b, Dixon and Soini 1986, Hoogmoed 1973, Ord and Blumstein 2002, Molina et al. 2004, Rogner 1997a, Alvarez 2004, Bartlett and Bartlett 2003, Mccranie et al. 2005, Kohler et al. 2006, Powell et al. 2005, Daudin and de Silva 2007, Guyer and Donnelly 2005, Van Buurt 2005, Colli et al. 2002, Hodge et al. 2003, Andrews and Pough 1980, Rodrigues 2003, Rodrigues 1996, Duellman 1961, Garrick 2008, Lotzkat 2007, Fitch 1982, Murphy 1997, Kohler 2008, Hardy and McDiarmid 1969, Fuenmayor et al. 2005, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler 2005, De Magalhaes and Costa 2009, Avila-Pires et al. 2010, Ugueto and Rivas 2010, Gasc 1990, Castanet 1994, Rivero 1998, Van

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Wilgen and Richardson 2012, Honegger 1969, Brattstrom 1965, Leenders and Watkins-Colwell 2004
Iguanidae	<i>Sauromalus ater</i>	no	NA	NA	NA	mainland	33	178.91	7.75	0.835	6.52	42.20	Cooper and Vitt 2002, Shaw 1945, Linsdale 1932, Van Denburgh 1922, Warne and Charnov 2008, Jones and Lovich 2009, Brennan and Holycross 2009, Brattstrom 1965 (as <i>obesus</i>), Charnov et al. 2007, Stebbins and McGinnis 2012, Lemm 2006 Fitch 1985, Perry and Garland 2002, Grismer 2002, Wiewandt 1982, Case 1982, Cooper and Vitt 2002, Shaw 1945, Van Denburgh 1922, Andrews and Pough 1980, Kohler 2005, Smits 1985, Smits et al. 1986, Goldberg and Beaman 2012
Iguanidae	<i>Sauromalus hispidus</i>	yes	Baja California	Angel de la Guarda	945.20	Land bridge	29	837.96	22.00	1	17.37	382.14	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Iguanidae	<i>Sauromalus varius</i>	no	NA	NA	NA	mainland	29	1124.69	23.40	0.625	18.82	275.28	Fitch 1985, Grismer 2002, Wiewandt 1982, Shaw 1945, Van Denburgh 1922, Kohler 2005, Goldberg and Beaman 2012
Lacertidae	<i>Acanthodactylus beershebensis</i>	no	NA	NA	NA	mainland	31	6.95	4.50	1	0.38	1.72	Amitai and Bouskila 2001, Moravec et al. 1999, Werner 2004, Arbel 1984, Hawlena et al. 2010, Hawlena and Perez-Mellado 2009, Duvdevani and Borut 1974, Bar and Haimovitch 2012
Lacertidae	<i>Acanthodactylus boskianus</i>	no	NA	NA	NA	mainland	26	5.09	5.00	1	0.72	3.61	Frankenberg and Werner 1992, Schleich et al. 1996, Amitai and Bouskila 2001, Arbel 1984, Anderson 1999, Perry and Garland 2002, Disi et al. 2001, Baran and Atatur 1998, Perez-Mellado 1992, Leviton et al. 1992, Geniez et al. 2004, Flower 1933, Papenfuss 1969, Dunger 1967, Arnold 1984, El Din 2006, Vanhooydonck and Van Damme 1999, Rogner 1997b, Arnold 1980, Reed and Marx 1959, Schatti and Desvoignes 1999, Le Berre 1989, Anderson 1898, Bons and Geniez 1996, Jongbloed 2000, Andrews and Pough 1980, Hornby 1996, AL-Johany and Spellerberg 1989, Seifan et al. 2009, McElroy et al. 2008, Sinervo et al. 2010, Kohler 2005, Largen and Spawls 2010, van der Kooij 2001, Duvdevani and Borut 1974, Duvdevani 1971, Verwaijen and Van Damme 2007, Perry et al. 1990, Bar

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Acanthodactylus erythrurus</i>	no	NA	NA	NA	mainland	37	7.51	3.95	1.5	0.59	3.51	and Haimovitch 2012, Bar 2003, Trape et al. 2012, Fathinia et al. 2009 Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Cox et al. 2003, Cooper and Vitt 2002, Arnold 1987, Pianka and Vitt 2003, Papenfuss 1969, Rogner 1997b, Le Berre 1989, Boulenger 1921, Verwaijen and Van Damme 2008, Andrews and Pough 1980, Belliure 2006, Warne and Charnov 2008, Radder et al. 2008, Schleich et al. 1996, Kwet 2009, Malkmus 2004, Amat 2008, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Valakos 1986, Rouag et al. 2007, Busack 1976, Carretero and Llorente 1993, Cisse et al. 1977, Maso and Pijoan 2011 Clobert et al. 1998, Fitch 1970, Frankenberg and Werner 1992, Flower 1933, El Din 2006, Vanhooydonck and Van Damme 1999, Rogner 1997b, Moravec et al. 1999, Werner 2004, Le Berre 1989, Anderson 1898, Nagy et al. 1999, Andrews and Pough 1980, Brown and Nagy 2007, Schleich et al. 1996, Arbel 1984, Sinervo et al. 2010, Kohler 2005, Turner 1977, Castanet 1994, Cisse et al. 1977
Lacertidae	<i>Acanthodactylus pardalis</i>	no	NA	NA	NA	mainland	31	5.32	3.65	2	0.65	4.78	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Acanthodactylus schmidti</i>	no	NA	NA	NA	mainland	25	26.40	3.00	2	0.95	5.68	Anderson 1999, Perry and Garland 2002, Disi et al. 2001, Gallagher 1971, Arnold 1984, Arnold 1980, Schatti and Desvoignes 1999, Anderson 1963, Jongbloed 2000, Hornby 1996, AL-Sadoon and Spellerberg 1985, AL-Sadoon and Abdo 1991, AL-Johany and Spellerberg 1989, Arnold 1994, van der Kooij 2001, Al-Johany and Spellerberg 1988
Lacertidae	<i>Acanthodactylus schreiberi</i>	no	NA	NA	NA	mainland	34	7.79	2.60	1	0.79	2.06	Frankenberg and Werner 1992, Amitai and Bouskila 2001, Arbel 1984, Zinner 1967, Reed and Marx 1959, Atatur and Gocmen 2001, Andrews and Pough 1980, Martens 1997, Sinervo et al. 2010, Kohler 2005, Baier et al. 2009, Duvdevani and Borut 1974, Perry et al. 1990, Bar and Haimovitch 2012, Yalcinkaya and Gocmen 2012, Zotos et al. 2012
Lacertidae	<i>Acanthodactylus scutellatus</i>	no	NA	NA	NA	mainland	26	3.43	2.60	2.5	0.35	2.27	Frankenberg and Werner 1992, Schleich et al. 1996, Disi et al. 2001, Perez-Mellado 1992, Flower 1933, El Din 2006, Vanhooydonck and Van Damme 1999, Le Berre 1989, Anderson 1898, Arbel 1984, Amitai and Bouskila 2001, Ibrahim 2008, Subach et al. 2009, Sinervo et al. 2010, Rifai et al. 2003, Baha El Din 1996, Tomasevic-Kolarov et al. 2010, Al-Hashem 2009, Duvdevani and Borut 1974,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Perry 1990, Perry et al. 1990, Bar and Haimovitch 2012, Bar 2003, Trape et al. 2012
Lacertidae	<i>Acanthodactylus spinicauda</i>	no	NA	NA	NA	mainland	33	3.32	4.00	2	0.87	6.94	Schleich et al. 1996
Lacertidae	<i>Algyroides fitzingeri</i>	yes	None	Sardinia	23949.00	Land bridge	41	0.88	2.50	1	0.19	0.49	Arnold and Ovenden 2004, Vanhooydonck and Van Damme 1999, Rogner 1997b, Sindaco et al. 2006, In Den Bosch and Bout 1998, Kwet 2009, Kohler 2005, Sindaco et al. 2010, Capula and Luiselli 1994, Capula et al. 2002
Lacertidae	<i>Algyroides marchi</i>	no	NA	NA	NA	mainland	38	2.57	2.50	2	0.26	1.31	Arnold and Ovenden 2004, Arnold 1987, Rogner 1997b, Rubio and Palacios 1986, Rubio and Carrascal 1994, Valverde 1958, Amat 2008, Kohler 2005, Siliceo and Diaz 2010, Cardenete and Cardenete 2010, Maso and Pijoan 2011
Lacertidae	<i>Algyroides moreoticus</i>	no	NA	NA	NA	mainland	38	2.17	3.30	4.5	0.21	3.09	Arnold and Ovenden 2004, Arnold 1987, Rogner 1997b, In Den Bosch and Bout 1998, Valakos et al. 2008, Pafilis et al. 2009, Kwet 2009, Amat 2008, Kohler 2005
Lacertidae	<i>Algyroides nigropunctatus</i>	no	NA	NA	NA	mainland	42	3.61	3.50	1.5	0.26	1.38	Arnold and Ovenden 2004, Arnold 1987, Vanhooydonck and Van Damme 1999, Sindaco et al. 2006, Corti and Cascio 2002, Valakos et al. 2008, Pafilis et al. 2009,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Kwet 2009, Kohler 2005, Sindaco et al. 2010, Haxhiu 1998, Arnold 1987
Lacertidae	<i>Anatololacerta anatolica</i>	no	NA	NA	NA	mainland	39	3.51	5.50	1	0.28	1.54	Arnold and Ovenden 2004, Valakos et al. 2008, Kohler 2005
Lacertidae	<i>Anatololacerta oertzeni</i>	no	NA	NA	NA	mainland	37	5.73	5.00	1	0.38	1.89	Rogner 1997b, Valakos et al. 2008, Pafilis et al. 2009, Wilson and Grillitsch 2009, Arnold and Ovenden 2004, Cooper and Vitt 2002, Vanhooydonck and Van Damme 1999, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Kwet 2009, Schneider 1984, Sinervo et al. 2010, Kohler 2005, Sindaco et al. 2010, Bauwens et al. 1990, Bombi and Vignoli 2004
Lacertidae	<i>Archaeolacerta bedriagae</i>	yes	None	Sardinia	23949.00	Land bridge	41	8.24	4.50	NA	0.76	NA	Schleich et al. 1996, Arnold 1998, Bons and Geniez 1996, Amat 2008, Kohler 2005, Busack 1987, Galan and Vicente 2003, Carretero et al. 2006
Lacertidae	<i>Atlantolacerta andreanskyi</i>	no	NA	NA	NA	mainland	32	2.31	1.85	3.5	0.27	1.74	Arnold and Ovenden 2004, Arnold 1987, Arnold 1989, Vanhooydonck and Van Damme 1999, Rogner 1997b, Verwaijen and Van Damme 2008, Kwet 2009, Amat 2008, Bischoff 1984, Sinervo et al. 2010, Kohler 2005, Arnold 1987, Verwaijen and Van Damme 2007
Lacertidae	<i>Dalmatolacerta oxycephala</i>	no	NA	NA	NA	mainland	43	4.12	3.40	2	0.31	2.14	Szczerbak 2003, Anderson 1999, Vanhooydonck and Van Damme 1999, Arnold 1998, Kohler 2005

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Darevskia derjugini</i>	no	NA	NA	NA	mainland	42	4.26	4.50	2.5	0.23	2.55	Szczerbak 2003, Baran and Atatur 1998, Arnold 1998, Rogner 1997b, Orlova and Bischoff 1984, Roitberg 1999, Kohler 2005
Lacertidae	<i>Darevskia parvula</i>	no	NA	NA	NA	mainland	41	3.09	3.00	1	0.31	0.93	Szczerbak 2003, Baran and Atatur 1998, Rogner 1997b, Darevskii 1978, Ilgaz 2009
Lacertidae	<i>Darevskia portschinskii</i>	no	NA	NA	NA	mainland	41	3.32	3.50	1	0.38	1.34	Szczerbak 2003, Arakelyan et al. 2011
Lacertidae	<i>Darevskia praticola</i>	no	NA	NA	NA	mainland	43	3.15	5.71	1.5	0.26	2.26	Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Street 1979, Arnold 1998, Rogner 1997b, Valakos et al. 2008, Ljubisavljevic et al. 2008, Kwet 2009, Roitberg 1999, Kohler 2005, Tuniyev et al. 2011, Arakelyan et al. 2011, Bogin et al. 1999
Lacertidae	<i>Darevskia raddei</i>	no	NA	NA	NA	mainland	39	5.06	4.00	1	0.41	1.66	Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Rogner 1997b, Darevskii 1978, Ahmadzadeh et al. 2008, Rastegar-Pouyani et al. 2011
Lacertidae	<i>Darevskia rostombekovi</i>	no	NA	NA	NA	mainland	41	3.23	3.00	1	0.28	0.84	Szczerbak 2003, Murphy et al. 2000, Darevskii 1978, Arakelyan et al. 2011
Lacertidae	<i>Darevskia rudis</i>	no	NA	NA	NA	mainland	42	7.54	5.00	1	0.45	2.26	Szczerbak 2003, Baran and Atatur 1998, Rogner 1997b, Darevsky and Tuniyev 1997, Darevskii 1978, Bohme and Bischoff 1984, Kohler 2005
Lacertidae	<i>Darevskia saxicola</i>	no	NA	NA	NA	mainland	44	4.52	4.00	2	0.34	2.74	Arnold and Ovenden 2004, Szczerbak 2003, Arnold 1998, Rogner 1997b, Darevskii 1978, Kohler 2005, Bodenheimer 1944

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Darevskia unisexualis</i>	no	NA	NA	NA	mainland	40	4.78	5.00	1.5	0.43	3.22	Szczerbak 2003, Baran and Atatur 1998, Murphy et al. 2000, Schmidler et al. 1994, Rogner 1997b, Darevskii 1978, Kohler 2005, Arakelyan et al. 2011
Lacertidae	<i>Darevskia valentini</i>	no	NA	NA	NA	mainland	39	7.46	5.20	1	0.53	2.73	Szczerbak 2003, Baran and Atatur 1998, Schmidler et al. 1994, Darevskii 1978, Tayhan et al. 2011, Arakelyan et al. 2011
Lacertidae	<i>Dinarolacerta mosorensis</i>	no	NA	NA	NA	mainland	43	6.11	4.75	1	0.41	1.94	Arnold and Ovenden 2004, Arnold 1987, Arnold 1998, Rogner 1997b, Ljubisavljevi, et al. 2007, Kwet 2009, Bischoff 1984, Kohler 2005, Tomasevic-Kolarov et al. 2010, Arnold 1987
Lacertidae	<i>Eremias arguta</i>	no	NA	NA	NA	mainland	47	8.67	3.50	1.5	0.38	2.01	Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Perry and Garland 2002, Terbish et al. 2006, Kotenko 1986, Turner 1977, Kohler 2005, Turner 1977, Arakelyan et al. 2011, Tertyshnikov 1976
Lacertidae	<i>Eremias grammica</i>	no	NA	NA	NA	mainland	43	4.70	1.50	2.5	0.41	1.52	Szczerbak 2003, Anderson 1999, Perry and Garland 2002, Clark 1990
Lacertidae	<i>Eremias intermedia</i>	no	NA	NA	NA	mainland	43	3.32	2.00	2	0.48	1.92	Szczerbak 2003, Anderson 1999, Clark 1990
Lacertidae	<i>Eremias lineolata</i>	no	NA	NA	NA	mainland	41	1.77	2.00	1	0.38	0.76	Szczerbak 2003, Anderson 1999, Clark 1990
Lacertidae	<i>Eremias nigrocellata</i>	no	NA	NA	NA	mainland	36	6.89	5.50	2.5	0.48	6.60	Szczerbak 2003, Anderson and Leviton 1969, Clark 1990
Lacertidae	<i>Eremias nikolskii</i>	no	NA	NA	NA	mainland	41	6.41	3.50	1	0.38	1.34	Szczerbak 2003, Eremchenko 2007
Lacertidae	<i>Eremias persica</i>	no	NA	NA	NA	mainland	33	14.90	5.00	1	0.79	3.96	Szczerbak 2003, Anderson 1999, Minton 1966, Leviton et al. 1992, Vanhooydonck and Van Damme 1999,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Anderson and Leviton 1969, Khan 2006, Clark 1990
Lacertidae	<i>Eremias pleskei</i>	no	NA	NA	NA	mainland	39	3.81	4.00	2	0.30	2.39	Fitch 1970, Szczerbak 2003, Andeson 1999, Baran and Atatur 1998, Rogner 1997b, Tadevosyan 2007, Kohler 2005, Curry-Lindahl 1979, Arakelyan et al. 2011
Lacertidae	<i>Eremias regeli</i>	no	NA	NA	NA	mainland	38	5.32	2.50	2	0.43	2.15	Szczerbak 2003, Clark 1990
Lacertidae	<i>Eremias strauchi</i>	no	NA	NA	NA	mainland	38	6.41	4.50	2	0.53	4.81	Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Rogner 1997b, Franzen and Heckes 1999, Ahmadzadeh et al. 2008, Tadevosyan 2007, Kohler 2005, Ahmadzadeh et al. 2009, Arakelyan et al. 2011
Lacertidae	<i>Eremias velox</i>	no	NA	NA	NA	mainland	40	7.98	4.00	3	0.43	5.15	Fitch 1970, Smith 1935, Szczerbak 2003, Anderson 1999, Perry and Garland 2002, Vanhooydonck and Van Damme 1999, Clark 1990, Kohler 2005
Lacertidae	<i>Gallotia atlantica</i>	yes	Canary Islands	Fuerteventura	1633.30	Oceanic	29	5.55	2.20	2.5	0.66	3.64	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Rogner 1997b, Alvarez 2004, Valido and Nogales 2003, Nagy et al. 1999, Salvador 2008, Brown and Nagy 2007, Kohler 2005, Siliceo and Diaz 2010, Castanet 1994, Maso and Pijoan 2011
Lacertidae	<i>Gallotia bravoana</i>	yes	Canary Islands	Gomera	359.10	Oceanic	28	37.08	5.00	1	2.03	10.13	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Salvador 2007, Maso and Pijoan 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Gallotia caesaris</i>	yes	Canary Islands	Gomera	359.10	Oceanic	28	9.03	3.15	1.5	0.88	4.17	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Rogner 1997b, Salvador 2007, Roca 1999, Amat 2008, Molina-Borja et al. 2010, Siliceo and Diaz 2010, Maso and Pijoan 2011
Lacertidae	<i>Gallotia galloti</i>	yes	Canary Islands	Tenerife	2007.80	Oceanic	28	15.50	3.83	1.5	1.26	7.26	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Pianka and Vitt 2003, Vanhooydonck and Van Damme 1999, Rogner 1997b, Alvarez 2004, Valido and Nogales 2003, Nagy et al. 1999, Salvador 2009, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Tersa et al. 2010, Castanet 1994, de los Santos and de Nicolas 2008, Garcia et al. 2007, Maso and Pijoan 2011
Lacertidae	<i>Gallotia intermedia</i>	yes	Canary Islands	Tenerife	2007.80	Oceanic	28	57.29	7.40	NA	3.03	NA	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Hernandez et al. 2000, Salvador 2009
Lacertidae	<i>Gallotia simonyi</i>	yes	Canary Islands	El Hierro	290.50	Oceanic	28	91.01	10.15	1.5	3.13	47.71	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Salvador 2007, Rodriguez-Dominguez and Molina-Borja 1998, Amat 2008, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Maso and Pijoan 2011
Lacertidae	<i>Gallotia stehlini</i>	yes	Canary Islands	Gran Canaria	1529.90	Oceanic	28	142.33	9.80	1	1.96	19.20	Arnold and Ovenden 2004, Molina-Borja and Rodriguez-Dominguez 2004, Rogner 1997b, Nagy et al. 1999, Salvador 2007, Garrick 2008, Rodriguez-Dominguez and Molina-Borja 1998, Kohler

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2005, Siliceo and Diaz 2010, Tersa et al. 2010, Castanet 1994, Maso and Pijoan 2011
Lacertidae	<i>Heliobolus lugubris</i>	no	NA	NA	NA	mainland	-22	4.46	3.63	1.5	0.14	0.79	Huey and Pianka 1981, FitzSimons 1943, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Broadley 1971, Auerbach 1987, Nagy et al. 1999, McBrayer 2004, Brown and Nagy 2007, Vitt and Price 1982, Huey et al. 2001, Amat 2008, Sinervo et al. 2010, Pienaar 1966, Goldberg 2006, Huey and Pianka 2007, Huey and Pianka 1977, Verwajjen and Van Damme 2007
Lacertidae	<i>Hellenolacerta graeca</i>	no	NA	NA	NA	mainland	38	6.12	3.50	1	0.56	1.97	Arnold and Ovenden 2004, Arnold 1987, Rogner 1997b, Valakos et al. 2008, Pafilis et al. 2009, Kwet 2009, Bohme 1984, Arnold 1987
Lacertidae	<i>Iberolacerta aranica</i>	no	NA	NA	NA	mainland	43	4.37	2.99	1	0.40	1.20	Arnold and Ovenden 2004, Amat et al. 2008, Arribas 2008, Ibarguengoytia and Casalins 2007, Galan 2005, Amat 2008, Siliceo and Diaz 2010, Arribas 2009, Maso and Pijoan 2011
Lacertidae	<i>Iberolacerta aurelioi</i>	no	NA	NA	NA	mainland	43	3.93	2.52	1	0.54	1.36	Arnold and Ovenden 2004, Amat et al. 2008, Arribas 2008, Ibarguengoytia and Casalins 2007, Galan 2005, Kwet 2009, Amat 2008, Siliceo and Diaz 2010, Maso and Pijoan 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Iberolacerta bomnali</i>	no	NA	NA	NA	mainland	43	4.11	3.03	1	0.35	1.07	Arnold and Ovenden 2004, Amat et al. 2008, Arribas 2008, Iburguengoytia and Casalins 2007, Galan 2005, Amat 2008, Siliceo and Diaz 2010, Maso and Pijoan 2011
Lacertidae	<i>Iberolacerta cyreni</i>	no	NA	NA	NA	mainland	41	9.68	5.95	1	0.35	2.09	Salvador et al. 2008, Amo et al. 2007, Martin 2008, Kwet 2009, Amat 2008, Siliceo and Diaz 2010, Monasterio et al. 2009, Maso and Pijoan 2011
Lacertidae	<i>Iberolacerta horvathi</i>	no	NA	NA	NA	mainland	45	4.98	3.70	1	0.38	1.41	Arnold and Ovenden 2004, Arnold 1987, Arnold 1998, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Kwet 2009, Amat 2008, Bischoff 1984, Sindaco et al. 2010, Arnold 1987
Lacertidae	<i>Iberolacerta monticola</i>	no	NA	NA	NA	mainland	43	6.78	6.30	2	0.34	4.24	Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Perry and Garland 2002, Arnold 1987, Arnold 1998, Rogner 1997b, Arribas and Carranza 2004, Verwaijen and Van Damme 2008, Dunham et al. 1988, Arribas 2006, Martin 2008, Brana et al. 1992, Iburguengoytia and Casalins 2007, Kwet 2009, Malkmus 2004, Amat 2008, Salvador 1984, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Castilla and Bauwens 2000, Valakos 1986, Martín and Salvador 1997, Arnold 1987, Verwaijen and Van Damme 2007, Maso and Pijoan 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Ichnotropis capensis</i>	no	NA	NA	NA	mainland	-20	3.69	6.14	1.5	0.20	1.82	Clobert et al. 1998, Fitch 1970, Branch 1998, Broadley 1971, Vanhooydonck and Van Damme 1999, Rogner 1997b, Loveridge 1953, Auerbach 1987, Graham and Marais 2007, Jacobsen et al. 2010, Pienaar 1966, Haagner et al. 2000, Jacobsen 1982, Branch and McCartney 1992 Huey and Pianka 1981, Fitch 1970, Spawls et al. 2002, FitzSimons 1943, Loveridge 1942, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Broadley 1971, Jeffery 1993, Loveridge 1953, Auerbach 1987, Graham and Marais 2007, Huey et al. 2001, Sinervo et al. 2010, Pienaar 1966, Goldberg 2008, Huey and Pianka 2007, Huey and Pianka 1977, Verwaijen and Van Damme 2007
Lacertidae	<i>Ichnotropis squamulosa</i>	no	NA	NA	NA	mainland	-20	4.20	4.00	1.5	0.59	3.55	Tinkle et al. 1970, Clobert et al. 1998, Fitch 1970, Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Szczerbak 2003, Perry and Garland 2002, Baran and Atatur 1998, Spellerberg 2002, Arnold 1987, Street 1979, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Valakos et al. 2008, Terbish et al. 2006, Amat 2008, Warne and Charnov 2008, Strijbosch 1986, Korsos 1986, Roitberg 2007, AL-Sadoon and Spellerberg 1985, Kwet 2009, Amat 2008,
Lacertidae	<i>Lacerta agilis</i>	no	NA	NA	NA	mainland	52	10.80	7.65	1.5	0.31	3.56	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Bischoff 1984, Sinervo et al. 2010, Kohler 2005, Necas et al. 1997, Siliceo and Diaz 2010, Bauwens 1999, Sindaco et al. 2010, Turner 1977, Haxhiu 1998, Castanet 1994, Arakelyan et al. 2011, Verwaijen and Van Damme 2007, Tertyshnikov 1976, Maso and Pijoan 2011
Lacertidae	<i>Lacerta schreiberi</i>	no	NA	NA	NA	mainland	41	29.08	13.70	2	0.55	15.10	Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Perry and Garland 2002, Arnold 1987, Rogner 1997b, Verwaijen and Van Damme 2008, Marco 2008, Kwet 2009, Malkmus 2004, Amat 2008, Salvador 1984, Kohler 2005, Siliceo and Diaz 2010, Bauwens 1999, Norrie and Langerwerf 1987, Martin and Lopez 2010, Verwaijen and Van Damme 2007, Maso and Pijoan 2011
Lacertidae	<i>Lacerta strigata</i>	no	NA	NA	NA	mainland	41	18.74	10.50	1.5	0.69	10.88	Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Darewskij 1984, Kohler 2005, Arakelyan et al. 2011
Lacertidae	<i>Lacerta trilineata</i>	no	NA	NA	NA	mainland	40	33.30	16.50	1.5	0.95	23.44	Arnold and Ovenden 2004, Szczerbak 2003, Baran and Atatur 1998, Arnold 1987, Kumlutas et al. 2004, Rogner 1997b, Andrews and Pough 1980, Valakos et al. 2008, Valakos et al. 2004, Kwet 2009, Nettmann and Rykena 1984, Kohler 2005, Haxhiu 1998

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Lacerta viridis</i>	no	NA	NA	NA	mainland	45	24.54	8.05	1.5	0.64	7.75	Clobert et al. 1998, Fitch 1970, Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Szczerbak 2003, Perry and Garland 2002, Baran and Atatur 1998, Spellerberg 2002, Arnold 1987, Vanhooydonck and Van Damme 1999, Street 1979, Rogner 1997b, Herczeg et al. 2007, Nagy et al. 1999, Sindaco et al. 2006, Corti and Cascio 2002, Andrews and Pough 1980, Valakos et al. 2008, Brown and Nagy 2007, Korsos 1986, Radder et al. 2008, Kwet 2009, Werner 1930, Nettmann and Rykena 1984, Sinervo et al. 2010, Kohler 2005, Necas et al. 1997, Sindaco et al. 2010, Valakos 1986, Haxhiu 1998, Castanet 1994, Maura et al. 2011
Lacertidae	<i>Meroles anchietae</i>	no	NA	NA	NA	mainland	-23	2.03	1.30	2.5	0.45	1.48	Dunham and Miles 1985, Huey and Pianka 1981, FitzSimons 1943, Cox et al. 2003, Branch 1998, Cooper and Vitt 2002, Pianka and Vitt 2003, Greene 1982, Arnold 1998, Rogner 1997b, Nagy et al. 1999, Dunham et al. 1988, Vitt and Price 1982, Warne and Charnov 2008, Nkosi et al. 2004, Amat 2008, Goldberg and Robinson 1979, Arnold 1994, Sinervo et al. 2010, Brain 1962

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Meroles cuneirostris</i>	no	NA	NA	NA	mainland	-29	2.78	2.90	1.5	0.47	2.03	Dunham and Miles 1985, FitzSimons 1943, Cox et al. 2003, Branch 1998, Pianka and Vitt 2003, Arnold 1998, Rogner 1997b, Dunham et al. 1988, Vitt and Price 1982, Nkosi et al. 2004, Amat 2008, Goldberg and Robinson 1979, Goldberg 2006
Lacertidae	<i>Meroles suborbitalis</i>	no	NA	NA	NA	mainland	-29	4.04	4.11	1.5	0.45	2.80	Huey and Pianka 1981, FitzSimons 1943, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Rogner 1997b, Auerbach 1987, McBrayer 2004, Huey et al. 2001, Sinervo et al. 2010, Goldberg 2006, Huey and Pianka 2007, Curry-Lindahl 1979, Huey and Pianka 1977, Verwaijen and Van Damme 2007, Brattstrom 1965
Lacertidae	<i>Mesalina guttulata</i>	no	NA	NA	NA	mainland	30	2.24	4.40	2	0.25	2.23	Frankenberg and Werner 1992, Smith 1935, Schleich et al. 1996, Amitai and Bouskila 2001, Fitch 1970, Szczerbak 2003, Minton 1966, Perry and Garland 2002, Disi et al. 2001, Perez-Mellado 1992, Leviton et al. 1992, Geniez et al. 2004, Flower 1933, El Din 2006, Parker 1942, Arnold 1998, Vanhooydonck and Van Damme 1999, Schatti and Desvoignes 1999, Le Berre 1989, Bons and Geniez 1996, Moravec and Modry 1994b, Arbel 1984, Kohler 2005, Perry et al. 1990, Bar and Haimovitch 2012, Goldberg 2012, Trape et al. 2012

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Mesalina olivieri</i>	no	NA	NA	NA	mainland	31	1.63	4.50	2	0.28	2.52	Schleich et al. 1996, Clobert et al. 1998, Amitai and Bouskila 2001, Disi et al. 2001, Perez-Mellado 1992, Geniez et al. 2004, Pianka and Vitt 2003, El Din 2006, Le Berre 1989, Bons and Geniez 1996, Brown and Nagy 2007, Castanet 1994, Cisse and Karns 1978, Bar and Haimovitch 2012, Trape et al. 2012
Lacertidae	<i>Mesalina pasteuri</i>	no	NA	NA	NA	mainland	21	2.03	3.00	2	0.28	1.68	Schleich et al. 1996, Geniez et al. 2004, El Din 2006, Bons and Geniez 1996, Trape et al. 2012
Lacertidae	<i>Mesalina rubropunctata</i>	no	NA	NA	NA	mainland	26	2.69	4.50	3	0.43	5.79	Schleich et al. 1996, Geniez et al. 2004, Flower 1933, Papenfuss 1969, El Din 2006, Le Berre 1989, Werner and Ashkenazi 2010, Trape et al. 2012
Lacertidae	<i>Nucras lalandii</i>	no	NA	NA	NA	mainland	-30	16.26	7.00	1	0.20	1.38	Fitch 1970, FitzSimons 1943, Branch 1998, Kohler 2005, van der Meer et al. 2010, Bogin et al. 1999
Lacertidae	<i>Nucras taeniolata</i>	no	NA	NA	NA	mainland	-22	7.72	4.79	1	0.79	3.79	Spawls et al. 2002, Branch 1998, Broadley 1971, Auerbach 1987, van der Meer et al. 2010, Jacobsen 1982, Huey and Pianka 1981, FitzSimons 1943, Branch 1998, Pianka and Vitt 2003, Pianka 1986, Pianka 1986, Pianka and Vitt 2003, Auerbach 1987, Vitt and Price 1982, Huey et al. 2001, Sinervo et al. 2010, Pienaar 1966, van der Meer et al. 2010, Huey and Pianka 2007, Huey and Pianka 1977,
Lacertidae	<i>Nucras tessellata</i>	no	NA	NA	NA	mainland	-31	8.32	3.96	1	0.79	3.14	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Verwaijen and Van Damme 2007
Lacertidae	<i>Ophisops elegans</i>	no	NA	NA	NA	mainland	36	2.52	4.00	3	0.22	2.68	Frankenberg and Werner 1992, Schleich et al. 1996, Amitai and Bouskila 2001, Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Disi et al. 2001, Baran and Atatur 1998, El Din 2006, Rogner 1997b, Reed and Marx 1959, Le Berre 1989, Khan 2006, Atatur and Gocmen 2001, Ahmadzadeh et al. 2008, Valakos et al. 2008, Valakos et al. 2004, Pafilis et al. 2009, Moravec 1998, Arbel 1984, Weber 1960, Kwet 2009, Werner 1930, Kohler 2005, Baier et al. 2009, Arakelyan et al. 2011, Verwaijen and Van Damme 2007, Bar and Haimovitch 2012, Fathinia et al. 2009
Lacertidae	<i>Parvilacerta parva</i>	no	NA	NA	NA	mainland	39	3.61	3.00	2	0.35	2.08	Szczerbak 2003, Baran and Atatur 1998, Vanhooydonck and Van Damme 1999, Arnold 1998, Rogner 1997b, Kumluta et al. 2004, Kohler 2005, Arakelyan et al. 2011
Lacertidae	<i>Pedioplanis burchelli</i>	no	NA	NA	NA	mainland	-31	3.73	4.50	1.5	0.30	2.03	FitzSimons 1943, Branch 1998, Uetz 2006, Nkosi et al. 2004, Amat 2008, Kohler 2005, Goldberg 2006

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Pedioplanis namaquensis</i>	no	NA	NA	NA	mainland	-30	2.97	3.85	1.5	0.23	1.32	Huey and Pianka 1981, FitzSimons 1943, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Parker 1936, Rogner 1997b, Auerbach 1987, McBrayer 2004, Huey et al. 2001, Kohler 2005, Goldberg 2006, Huey and Pianka 2007, Curry-Lindahl 1979, Huey and Pianka 1977, Verwajen and Van Damme 2007
Lacertidae	<i>Phoenicolacerta laevis</i>	no	NA	NA	NA	mainland	35	6.25	5.30	3	0.39	6.22	Frankenberg and Werner 1992, Amitai and Bouskila 2001, Arbel 1984, Disi et al. 2001, Baran and Atatur 1998, Zinner 1967, Arnold 1998, Rogner 1997b, Atatur and Gocmen 2001, Amat 2008, Kohler 2005, Baier et al. 2009, In den Bosch and Zandee 2001, Perry et al. 1990, Bar and Haimovitch 2012, Bogin et al. 1999
Lacertidae	<i>Phoenicolacerta troodica</i>	yes	None	Cyprus	9234.40	Land bridge	35	5.14	5.00	2.5	0.41	5.07	Baier et al. 2009
Lacertidae	<i>Podarcis bocagei</i>	no	NA	NA	NA	mainland	43	3.44	3.40	2.5	0.28	2.38	Clobert et al. 1998, Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Perry and Garland 2002, Arnold 1987, Rogner 1997b, Kaliontzopoulou et al. 2008, Perez-Mellado 1981, Galan 2008, Brana et al. 1992, Galan 1997, Kwet 2009, Malkmus 2004, Amat 2008, Kohler 2005, Siliceo and Diaz 2010, Bauwens 1999, Castilla and Bauwens 2000, Galan and Vicente 2003, Arnold 1987, Maso and Pijoan 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Podarcis carbonelli</i>	no	NA	NA	NA	mainland	40	3.30	2.15	2	0.23	0.97	Arnold and Ovenden 2004, Sa-Sousa 2008, Galan 1997, Amat 2008, Kaliontzopoulou et al. 2010, Siliceo and Diaz 2010, Galan and Vicente 2003, Bowker et al. 2010, Maso and Pijoan 2011
Lacertidae	<i>Podarcis cretensis</i>	yes	Greece	Crete	8260.00	Land bridge	35	4.31	3.50	2	0.45	3.18	Lymberakis et al. 2008
Lacertidae	<i>Podarcis erhardii</i>	no	NA	NA	NA	mainland	39	5.42	2.53	2	0.52	2.65	Arnold and Ovenden 2004, Perry and Garland 2002, Arnold 1987, Vanhooydonck and Van Damme 1999, Rogner 1997b, Uetz 2006, Valakos et al. 2008, Pafilis et al. 2009, Maragou et al. 1999, Kwet 2009, Amat 2008, Kohler 2005, Valakos 1986, Tsasi et al. 2009, Adamopoulo et al. 1999
Lacertidae	<i>Podarcis filfolensis</i>	yes	Maltese Archipelago	Malta	245.70	Land bridge	36	5.64	2.50	1.5	0.37	1.40	Arnold and Ovenden 2004, Cooper and Vitt 2002, Vanhooydonck and Van Damme 1999, Sindaco et al. 2006, Corti and Cascio 2002, Lo Cascio et al. 2006, Carretero et al. 2010, Sindaco et al. 2010, Cascio 2010
Lacertidae	<i>Podarcis gaiageae</i>	yes	Greece	Skyros	209.00	Land bridge	39	5.50	2.47	2.5	0.52	3.23	Arnold and Ovenden 2004, Valakos et al. 2008, Pafilis et al. 2008, Pafilis et al. 2009, Pafilis, personal communication, February 2009, Meiri, unpublished, Adamopoulo et al. 1999, Pafilis et al. 2011, Sagonas et al. 2013
Lacertidae	<i>Podarcis hispanicus</i>	no	NA	NA	NA	mainland	41	3.50	2.85	3	0.32	2.72	Bauwens and Diaz-Uriarte 1997, Castilla and Bauwens 2000, Arnold and Ovenden 2004, Perry and Garland

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Podarcis lilfordi</i>	yes	Balearic Islands	Ibiza	576.90	Continental	40	4.90	2.50	2.5	0.75	4.66	2002, Cooper and Vitt 2002, Arnold 1987, Rogner 1997b, Alvarez 2004, Bons and Geniez 1996, Verwaijen and Van Damme 2008, Schleich et al. 1996, Kwet 2009, Malkmus 2004, Amat 2008, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Castilla and Bauwens 2000, Galan and Vicente 2003, Valakos 1986, Castanet 1994, Arnold 1987, Verwaijen and Van Damme 2007, Maso and Pijoan 2011 Arnold and Ovenden 2004, Olesen and Valido 2003, Cooper and Vitt 2002, Rogner 1997b, Alvarez 2004, Nagy et al. 1999, Nyhagen et al. 2001, Sazima et al. 2005, Salvador 2008, Brown and Nagy 2007, Roca 1999, AL-Sadoon and Spellerberg 1985, Amat 2008, Kohler 2005, Siliceo and Diaz 2010, Castilla and Bauwens 2000, Salvador 2008, Brooke and Houston 1983, Maso and Pijoan 2011 Bejakovic et al. 1995, Arnold and Ovenden 2004, Arnold 1987, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Verwaijen and Van Damme 2008, Kwet 2009, Amat 2008, Sinervo et al. 2010, Kohler 2005, Vervust 2011, Sindaco et al. 2010, Verwaijen and Van Damme 2007
Lacertidae	<i>Podarcis melisellensis</i>	no	NA	NA	NA	mainland	44	4.90	4.30	3	0.36	4.65	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Podarcis milensis</i>	yes	Greece	Milos	150.60	Oceanic	37	3.03	1.73	1.5	0.38	1.00	Arnold and Ovenden 2004, Arnold 1987, In Den Bosch and Bout 1998, Valakos et al. 2008, Pafilis et al. 2009, Adamopoulou and Valakos 2000, Sinervo et al. 2010, Kohler 2005, Adamopoulou and Legakis 2002, Adamopoulo et al. 1999, Adamopoulou and Valakos 2005, Arnold 1987, Clobert et al. 1998, Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Perry and Garland 2002, Baran and Atatur 1998, Cox et al. 2003, Spellerberg 2002, Cooper and Vitt 2002, Arnold 1987, Arnold 1987, Vanhooydonck and Van Damme 1999, Street 1979, Rogner 1997b, Herczeg et al. 2007, Sindaco et al. 2006, Corti and Cascio 2002, Verwajen and Van Damme 2008, Valakos et al. 2008, Clusella-Trullas et al. 2008, Rasilla 2008, Radder et al. 2008, Pafilis et al. 2009, Brana et al. 1992, Kwet 2009, Amat 2008, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Bauwens 1999, Castilla and Bauwens 2000, Sindaco et al. 2010, Galan and Vicente 2003, Haxhiu 1998, Castilla and Bauwens 1991, Monasterio et al. 2009, Castanet 1994, Verwajen and Van Damme 2007, Van Wilgen and
Lacertidae	<i>Podarcis muralis</i>	no	NA	NA	NA	mainland	45	5.14	5.48	2	0.36	3.92	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Richardson 2012, Maso and Pijoan 2011, Bogin et al. 1999
Lacertidae	<i>Podarcis peloponnesiacus</i>	no	NA	NA	NA	mainland	37	7.16	3.48	1.5	0.55	2.85	Arnold and Ovenden 2004, Arnold 1987, Rogner 1997b, Verwaijen and Van Damme 2008, Valakos et al. 2008, Pafilis et al. 2009, Maragou et al. 1999, Kwet 2009, Amat 2008, Kohler 2005, Maragou et al. 1999, Arnold 1987, Verwaijen and Van Damme 2007
Lacertidae	<i>Podarcis pityusensis</i>	yes	Balearic Islands	Ibiza	576.90	Continental	39	4.46	2.65	1	0.59	1.57	Arnold and Ovenden 2004, Olesen and Valido 2003, Cooper and Vitt 2002, Rogner 1997b, Alvarez 2004, Salvador 2009, Roca 1999, Kwet 2009, Amat 2008, Kohler 2005, Siliceo and Diaz 2010, Galan and Vicente 2003, Arnold 1987, Maso and Pijoan 2011
Lacertidae	<i>Podarcis raffoneae</i>	yes	Lipari (Eolie) Islands	Lipari	37.00	Oceanic	39	14.15	6.00	2	NA	NA	Arnold and Ovenden 2004, Sindaco et al. 2006, Corti and Cascio 2002, Sindaco et al. 2010, Cascio and Capula 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Podarcis siculus</i>	no	NA	NA	NA	mainland	42	5.54	4.60	3	0.59	8.17	Clobert et al. 1998, Arnold and Ovenden 2004, Baran and Atatur 1998, Arnold 1987, Vanhooydonck and Van Damme 1999, Street 1979, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Andrews and Pough 1980, Herrel et al. 2008, Salvador 2006, Kwet 2009, Amat 2008, Sinervo et al. 2010, Kohler 2005, Vervust 2011, Vervust et al. 2007, Vervust et al. 2008, Vervust et al. 2009, Vervust et al. 2010, Sindaco et al. 2010, Gibbs et al. 2007, Arnold 1987, Van Wilgen and Richardson 2012
Lacertidae	<i>Podarcis tauricus</i>	no	NA	NA	NA	mainland	43	5.51	4.79	2	0.46	4.44	Clobert et al. 1998, Dunham and Miles 1985, Arnold and Ovenden 2004, Szczerbak 2003, Perry and Garland 2002, Baran and Atatur 1998, Arnold 1987, Vanhooydonck and Van Damme 1999, Street 1979, Rogner 1997b, Dunham et al. 1988, Valakos et al. 2008, Pafilis et al. 2009, Maragou et al. 1999, Kwet 2009, Amat 2008, Kohler 2005, Turner 1977, Arnold 1987
Lacertidae	<i>Podarcis tiliguerta</i>	yes	None	Sardinia	23949.00	Land bridge	41	4.23	9.00	NA	0.45	NA	Arnold and Ovenden 2004, Vanhooydonck and Van Damme 1999, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Verwaijen and Van Damme 2008, Kwet 2009, Sinervo et al. 2010, Kohler 2005, Sindaco et al. 2010, Capula and Luiselli

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1994, Verwaijen and Van Damme 2007, Van Damme et al. 1989
Lacertidae	<i>Podarcis waglerianus</i>	yes	None	Sicily	25662.40	Land bridge	38	3.81	4.00	1.5	0.44	2.63	Arnold and Ovenden 2004, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Kwet 2009, Kohler 2005, Sindaco et al. 2010, Bauwens and Diaz-Uriarte 1997, Schleich et al. 1996, Arnold and Ovenden 2004, Perry and Garland 2002, Arnold 1987, Pianka and Vitt 2003, Rogner 1997b, Le Berre 1989, Bons and Geniez 1996, Sindaco et al. 2006, Corti and Cascio 2002, Verwaijen and Van Damme 2008, Diaz et al. 2007, AL-Sadoon and Spellerberg 1985, Kwet 2009, Malkmus 2004, Amat 2008, Sinervo et al. 2010, Kohler 2005, Siliceo and Diaz 2010, Salvador 2010, Sindaco et al. 2010, Valakos 1986, Rouag et al. 2007, Verwaijen and Van Damme 2007, Maso and Pijoan 2011
Lacertidae	<i>Psammodromus algirus</i>	no	NA	NA	NA	mainland	37	6.73	5.35	2	0.45	4.80	Schleich et al. 1996, Kohler 2005
Lacertidae	<i>Psammodromus blanci</i>	no	NA	NA	NA	mainland	36	1.83	3.00	4.5	0.23	3.08	Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Arnold 1987, Rogner 1997b, Verwaijen and Van Damme 2008, Kwet 2009, Malkmus 2004, Amat 2008, Kohler 2005, Siliceo and Diaz
Lacertidae	<i>Psammodromus hispanicus</i>	no	NA	NA	NA	mainland	40	2.69	3.25	1.5	0.23	1.11	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2010, Carretero and Llorente 1993, Verwaijen and Van Damme 2007
Lacertidae	<i>Scelarcis perspicillata</i>	no	NA	NA	NA	mainland	34	3.96	2.50	2.5	0.35	2.21	Schleich et al. 1996, Arnold and Ovenden 2004, Arnold 1998, Rogner 1997b, Bons and Geniez 1996, Perera 2007, Ana Perera, pers comm. 31.03.2009, Kohler 2005, Perera 2009, Galan and Vicente 2003, Carretero et al. 2006, Maso and Pijoan 2011
Lacertidae	<i>Takydromus amurensis</i>	no	NA	NA	NA	mainland	45	8.32	5.00	2	0.30	2.99	Szczerbak 2003, Goris and Maeda 2004, Takenaka 1989, Huang 2006, Shannon 1956, Arnold 1997, Szyndlar 1984, Huang 1998b, Dixon 1956
Lacertidae	<i>Takydromus dorsalis</i>	yes	Ryukyu Islands	Iriomotejima	289.00	Oceanic	24	6.53	1.40	NA	0.45	NA	Goris and Maeda 2004, Takenaka 1989, Huang 2006, Arnold 1997
Lacertidae	<i>Takydromus formosanus</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	24	2.72	2.00	NA	0.79	NA	Perry and Garland 2002, Takenaka 1989, Huang 2006, Arnold 1997, Uetz 2006, Huang and Tu 2008, Huang 1998b
Lacertidae	<i>Takydromus hsuehshanensis</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	24	5.32	3.23	1.5	NA	NA	Cox et al. 2003, Takenaka 1989, Huang 2006, Arnold 1997, Uetz 2006, Huang and Tu 2008, Todd 2008, Huang 1998, Huang 1998b, Kohler 2005
Lacertidae	<i>Takydromus luyeanus</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	23	2.70	2.50	2	NA	NA	Lue and Lin 2008
Lacertidae	<i>Takydromus sauteri</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	23	6.63	2.00	2	NA	NA	Huang 2006, Arnold 1997, Amat 2008

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Takydromus septentrionalis</i>	no	NA	NA	NA	mainland	31	5.24	3.30	3	0.31	3.11	Bauwens and Diaz-Uriarte 1997, Du et al. 2005b, Takenaka 1989, Huang 2006, Pope 1929, Arnold 1997, Song 1987, Du et al. 2006, Huang 1998b, Amat 2008, Sinervo et al. 2010, Kohler 2005, Du et al. 2005, Zhang and Ji 2004, Lai-Gao et al. 2010, Wang et al. 2011 Fitch 1970, Smith 1935, Tikader and Sharma 1992, Taylor 1963, Takenaka 1989, Stuart et al. 2006, Huang 2006, Das 2004, Schmidt 1927, Pope 1929, Manthey and Grossmann 1997, Vanhooydonck and Van Damme 1999, Arnold 1997, de Rooij 1915, Rogner 1997b, Cox et al. 1998, Karsen et al. 1986, Inger and Colwell 1977, Pauwels et al. 2003, Grismer et al. 2008, Stuart and Emmett 2006, Zhang and Ji 2004, Huang 1998b, Kohler 2005, Teynie et al. 2010, Das 2010, Cox et al. 2010, Das 2011, Grismer 2011, Verwajen and Van Damme 2007, Teynie and David 2010
Lacertidae	<i>Takydromus sexlineatus</i>	no	NA	NA	NA	mainland	14	4.09	2.20	4.5	0.17	1.68	Goris and Maeda 2004, Takenaka 1989, Huang 2006, Arnold 1997, Rogner 1997b, Huang 1998b, Kohler 2005
Lacertidae	<i>Takydromus smaragdinus</i>	yes	Ryukyu Islands	Okinawajima	1199.50	Oceanic	27	3.03	2.00	NA	0.34	NA	Takenaka 1989, Huang 2006, Arnold 1997, Rogner 1997b, Huang 1998b, Kohler 2005
Lacertidae	<i>Takydromus stejnegeri</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	24	3.79	2.20	2.5	NA	NA	Takenaka 1989, Huang 2006, Arnold 1997, Huang and Tu 2008, Huang 1998b
Lacertidae	<i>Takydromus toyamai</i>	yes	Miyako Islands	Miyakojima	158.70	Oceanic	25	3.89	2.00	2	NA	NA	Cox et al. 2003, Goris and Maeda 2004, Huang 2006, Arnold 1997

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Takydromus viridipunctatus</i>	yes	Taiwan	Taiwan	34506.60	Land bridge	25	3.14	2.50	2	NA	NA	Lue and Lin 2008
Lacertidae	<i>Takydromus wolteri</i>	no	NA	NA	NA	mainland	39	3.13	4.00	3	0.26	3.07	Szczerbak 2003, Takenaka 1989, Huang 2006, Arnold 1997, Szyndlar 1984, Chen et al. 2003, Huang 1998b, Dixon 1956, Das 2010
Lacertidae	<i>Teira dugesii</i>	yes	Maderia Archipelago	Madeira	755.00	Oceanic	36	5.28	2.50	2	0.71	3.54	Arnold and Ovenden 2004, Cooper and Vitt 2002, Molina-Borja and Rodriguez-Dominguez 2004, Arnold 1998, Rogner 1997b, Malkmus 2004, Amat 2008, Sinervo et al. 2010, Kohler 2005, Galan and Vicente 2003, Crisp et al. 1979, Maso and Pijoan 2011, Jesus 2012
Lacertidae	<i>Timon lepidus</i>	no	NA	NA	NA	mainland	41	75.55	13.70	1.5	1.49	30.64	Clobert et al. 1998, Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Perry and Garland 2002, Cooper and Vitt 2002, Arnold 1987, Greene 1982, Street 1979, Rogner 1997b, Alvarez 2004, Le Berre 1989, Sindaco et al. 2006, Corti and Cascio 2002, Mateo 2008, Garrick 2008, Warne and Charnov 2008, Rodriguez-Dominguez and Molina-Borja 1998, Kwet 2009, Malkmus 2004, Amat 2008, Bischoff et al. 1984, Kohler 2005, De Magalhaes and Costa 2009, Siliceo and Diaz 2010, Bauwens 1999, Sindaco et al. 2010, Valakos 1986, De Magalhaes & Costa 2009, Busack and Visnaw 1989, Castanet 1994, Maso and Pijoan 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Lacertidae	<i>Timon pater</i>	no	NA	NA	NA	mainland	34	63.07	11.00	2.5	1.77	48.59	Schleich et al. 1996, Vanhooydonck and Van Damme 1999, Street 1979, Rogner 1997b, Bons and Geniez 1996, Kohler 2005 Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Bauwens and Diaz-Uriarte 1997, Arnold and Ovenden 2004, Szczerbak 2003, Perry and Garland 2002, Cox et al. 2003, Goris and Maeda 2004, Spellerberg 2002, Arnold 1987, Vanhooydonck and Van Damme 1999, Street 1979, Arnold 1998, Rogner 1997b, Uetz 2006, Sindaco et al. 2006, Chamaille-Jammes et al. 2006, Corti and Cascio 2002, Verwaijen and Van Damme 2008, Dunham et al. 1988, Andrews and Pough 1980, Terbish et al. 2006, Warne and Charnov 2008, Strijbosch 1986, Radder et al. 2008, AL-Sadoon and Spellerberg 1985, Kwet 2009, Amat 2008, Dely and Bohme 1984, Sinervo et al. 2010, Necas et al. 1997, Siliceo and Diaz 2010, Bauwens 1999, Sindaco et al. 2010, Turner 1977, Castilla and Bauwens 1991, Castanet 1994, Verwaijen and Van Damme 2007, Shine and Charnov 1992, Maso and Pijoan 2011 IUCN, Schwartz and Henderson 1991, Powell 1999, Thomas 1966
Lacertidae	<i>Zootoca vivipara</i>	no	NA	NA	NA	mainland	56	4.53	6.07	1.5	0.14	1.32	
Leiocephali dae	<i>Leiocephalus eremitus</i>	yes	Greater Antilles	Navassa	5.20	Oceanic	18	8.22	3.00	NA	NA	NA	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Leiocephali dae	<i>Leiocephalus inaguae</i>	yes	Bahamas	Great Inagua	1615.30	Oceanic	21	10.28	4.00	NA	NA	NA	Schoener et al. 1982, Cooper and Vitt 2002, Schwartz and Henderson 1991, Noble and Klingel 1932, Henderson and Powell 2009
Leiocephali dae	<i>Leiocephalus psammodromus</i>	yes	Bahamas	Grand Caicos	289.30	Oceanic	22	17.31	1.71	1.5	1.29	3.30	Schwartz and Henderson 1991, Smith and Iverson 1993, Smith 1992, Henderson and Powell 2009, Reynolds 2011
Leiosaurida e	<i>Enyalius perditus</i>	no	NA	NA	NA	mainland	-21	11.34	10.20	1	0.58	5.89	Rodrigues et al. 2006, Sturaro and da Silva 2010, Lima and de Sousa 2006
Liolaemida e	<i>Liolaemus andinus</i>	no	NA	NA	NA	mainland	-23	20.23	5.80	1	0.64	3.69	Cruz et al. 2005, Schulte et al. 2000, Donoso-Barros 1966, Cei 1993, Ramirez Leyton and Pincheira Donoso 2005, Waltner 1991, Dunham et al. 1988, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Pincheira-Donoso et al. 2009
Liolaemida e	<i>Liolaemus bellii</i>	no	NA	NA	NA	mainland	-33	8.06	4.22	1	0.71	2.99	Web and Greer 1969, Schulte et al. 2004, Jaksic et al. 1980, Cei 1986, Schulte et al. 2000, Donoso-Barros 1966, Pincheira-Donoso and Scolaro 2007, Scolaro 2006, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Naya et al. 2008, Pincheira-Donoso et al. 2009, Sinervo et al. 2010, Pincheira-Donoso and Tregenza 2011, Rodrigues-Serrano et al. 2009, Ibarquengoytia 2008, Labra et al. 2008, Carothers et al. 1998

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Liolaemidae	<i>Liolaemus bibronii</i>	no	NA	NA	NA	mainland	-45	4.67	2.75	1	0.57	1.56	Schulte et al. 2004, Cei et al. 2003, Cruz et al. 2005, Cei 1986, Schulte et al. 2000, Scolaro 2005, Cei 1982, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Medina et al. 2009, Halloy et al. 2006, Pincheira-Donoso et al. 2009, Sinervo et al. 2010, Kohler 2005, Espinoza et al. 2004, Medina and Ibarquengoytia 2010, Tuli et al. 2009, Labra et al. 2008, Pincheira-Donoso, pers. Comm. 8.6.2011, Ibarquengoytia et al. 2010
Liolaemidae	<i>Liolaemus boulengeri</i>	no	NA	NA	NA	mainland	-42	6.87	4.08	1	0.44	1.81	Cei et al. 2003, Cruz et al. 2005, Cei 1986, Schulte et al. 2000, Abdala 2006, Scolaro 2005, Pincheira-Donoso et al. 2008, Halloy et al. 2006, Pincheira-Donoso et al. 2009, Sinervo et al. 2010, Medina and Ibarquengoytia 2010, Pincheira-Donoso and Tregenza 2011, Pincheira-Donoso, pers. Comm. 8.6.2011
Liolaemidae	<i>Liolaemus elongatus</i>	no	NA	NA	NA	mainland	-39	14.30	3.90	0.75	0.35	1.03	Cruz et al. 2005, Cei 1986, Schulte et al. 2000, Avilla et al. 2003, Espinoza et al. 2000, Espinoza and Lobo 2003, Boretto and Ibarquengoytia 2006, Scolaro 2005, Cei et al. 1983, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Ibarquengoytia and Casalins 2007, Ibarquengoytia and Cussac 1998, Halloy et al. 2007, Halloy et al. 2006, Pincheira-

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Liolaemidae	<i>Liolaemus huacahuasicus</i>	no	NA	NA	NA	mainland	-27	7.32	4.59	1	0.51	2.32	Donoso et al. 2009, Sinervo et al. 2010, Medina and Ibarguengoytia 2010, Tuli et al. 2009, Pincheira-Donoso and Tregenza 2011, Ibarguengoytia 2008, Labra et al. 2008, Pincheira-Donoso, pers. Comm. 8.6.2011, Ibarguengoytia et al. 2010, Ibarguengoytia and Cussac 1999, Ibarguengoytia et al. 2007, Ibarguengoytia 2005, Cruz et al. 2009
Liolaemidae	<i>Liolaemus koslowskyi</i>	no	NA	NA	NA	mainland	-28	5.92	4.89	1	0.30	1.47	Cei 1993, Pincheira-Donoso et al. 2008, Halloy and Halloy 1997, Espinoza et al. 2004, Sinervo et al. 2010, Tuli et al. 2009, Pincheira-Donoso and Tregenza 2011, Pincheira-Donoso, pers. Comm. 8.6.2011 Schulte et al. 2004, Cruz et al. 2005, Schulte et al. 2000, Pincheira-Donoso et al. 2008, Halloy et al. 2006, Pincheira-Donoso et al. 2009, Espinoza et al. 2004, Sinervo et al. 2010, Tuli et al. 2009, Martori and Aun 2010, Pincheira-Donoso and Tregenza 2011, Labra et al. 2008, Cruz et al. 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Liolaemidae	<i>Liolaemus lemniscatus</i>	no	NA	NA	NA	mainland	-36	3.50	4.50	1	0.30	1.36	Web and Greer 1969, Schulte et al. 2004, Jaksic et al. 1980, Cei 1986, Schulte et al. 2000, Donoso-Barros 1966, Scolaro 2006, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Jaksic et al. 1982, Halloy et al. 2006, Sinervo et al. 2010, Kohler 2005, Espinoza et al. 2004, Pincheira-Donoso and Tregenza 2011, Rodrigues-Serrano et al. 2009, Ibarquengoytia 2008, Vidal and Labra 2008, Labra et al. 2008, Pincheira-Donoso, pers. Comm. 8.6.2011, Fuentes and Jaksic 1979, Carothers et al. 1998
Liolaemidae	<i>Liolaemus lineomaculatus</i>	no	NA	NA	NA	mainland	-47	5.30	3.56	1	0.45	1.59	Fitch 1970, Cei et al. 2003, Cei 1986, Schulte et al. 2000, Donoso-Barros 1966, Scolaro 2005, Pincheira-Donoso et al. 2008, Pincheira-Donoso et al. 2009, Medina and Ibarquengoytia 2010, Tuli et al. 2009, Pincheira-Donoso and Tregenza 2011, Ibarquengoytia 2008, Bonino et al. 2011, Minoli et al. 2010 Perry and Garland 2002, Cooper and Vitt 2002, Schulte et al. 2000, Rocha 2000, Verrastro et al. 2003, Pincheira-Donoso et al. 2008, Halloy et al. 2006, Sinervo et al. 2010, Kohler 2005, Rand 1982, Martori and Aun 2010, Pincheira-Donoso and Tregenza 2011, Labra et al. 2008, Verrastro 2004, Rocha
Liolaemidae	<i>Liolaemus lutzae</i>	no	NA	NA	NA	mainland	-23	7.88	2.75	1	0.87	2.40	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													et al. 2009, Winck and Rocha 2012
Liolaemidae	<i>Liolaemus nigromaculatus</i>	no	NA	NA	NA	mainland	-27	5.91	3.09	2	2.40	14.85	Jaksic et al. 1980, Donoso-Barros 1966, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Espinoza et al. 2004, Sinervo et al. 2010, Rodrigues-Serrano et al. 2009, Ibarquengoytia 2008, Labra et al. 2008, Pincheira-Donoso, pers. Comm. 8.6.2011, Fuentes and Jaksic 1979 Web and Greer 1969, Vidal et al. 2006, Schulte et al. 2004, Cei 1986, Schulte et al. 2000, Greene 1982, Donoso-Barros 1966, Boretto and Ibarquengoytia 2006, Scolaro 2005, Scolaro 2006, Avila et al. 2006, Pincheira-Donoso et al. 2008, Pincheira Donoso and Nunez 2005, Boretto and Ibarquengoytia 2009, Ibarquengoytia and Casalins 2007, Ibarquengoytia and Cussa 1998, Halloy et al. 2006, Pincheira-Donoso et al. 2009, Espinoza et al. 2004, Sinervo et al. 2010, Tuli et al. 2009, Pincheira-Donoso and Tregenza 2011, Rodrigues-Serrano et al. 2009, Ibarquengoytia 2008, Vidal and Labra 2008, Labra et al. 2008, Pincheira-Donoso, pers.
Liolaemidae	<i>Liolaemus pictus</i>	no	NA	NA	NA	mainland	-39	7.15	3.95	0.415	0.60	0.99	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Comm. 8.6.2011, Ibaruengoytia and Cussac 1996, Ibaruengoytia and Cussac 1999
Liolaemidae	<i>Liolaemus scolaroi</i>	no	NA	NA	NA	mainland	-47	7.43	3.00	1	0.63	1.89	Scolaro 2006, Pincheira-Donoso and Nunez 2005, Minoli et al. 2010 Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, Donoso-Barros 1966, Boulenger 1901, Ramirez Leyton and Pincheira Donoso 2005, Pincheira-Donoso et al. 2008, Warne and Charnov 2008, Pincheira Donoso and Nunez 2005, Pinilla 1991, Espinoza et al. 2004, Sinervo et al. 2010, Pincheira-Donoso and Tregenza 2011, Ibaruengoytia 2008, Pincheira-Donoso, pers. Comm. 8.6.2011
Liolaemidae	<i>Liolaemus signifer</i>	no	NA	NA	NA	mainland	-18	23.11	6.10	1	0.64	3.88	Fitzgerald et al. 1999, Cei 1986, Schulte et al. 2000, Cei 1993, Achaval and Olmos 2003, Verrastro et al. 2003, Scolaro 2006, Pincheira-Donoso et al. 2008, Carreira et al. 2005, Halloy et al. 2006, Pincheira-Donoso et al. 2009, Kohler 2005, Rand 1982, Martori and Aun 2010,
Liolaemidae	<i>Liolaemus wiegmannii</i>	no	NA	NA	NA	mainland	-33	4.38	4.60	1.5	0.22	1.55	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Pincheira-Donoso and Tregenza 2011, Labra et al. 2008, Loveridge 1959
Liolaemidae	<i>Liolaemus zullyae</i>	no	NA	NA	NA	mainland	-47	8.77	2.00	1	0.64	1.29	Cei and Scolaro 1996, Pincheira-Donoso et al. 2008, Pincheira-Donoso and Nunez 2005, Bonino et al. 2011, Minoli et al. 2010
Liolaemidae	<i>Phymaturus palluma</i>	no	NA	NA	NA	mainland	-35	33.45	2.75	0.5	4.40	6.06	Fitch 1970, Cooper and Vitt 2002, Cei 1986, Schulte et al. 2000, Donoso-Barros 1966, Cei et al. 1983, Pincheira-Donoso et al. 2008, Sinervo et al. 2010, Cabezas et al. 2010, Cruz et al. 2009
Liolaemidae	<i>Phymaturus patagonicus</i>	no	NA	NA	NA	mainland	-44	31.83	2.00	0.5	1.01	1.01	Ibarguengoytia 2004, Cei 1986, Cei and Castro 1973, Piantoni et al. 2006, Uetz 2006, Boretto and Ibarguengoytia 2006, Scolaro 2005, Pincheira-Donoso et al. 2008, Ibarguengoytia and Casalins 2007, Espinoza et al. 2004, Labra et al. 2008, Ibarguengoytia 2005, Cruz et al. 2009
Liolaemidae	<i>Phymaturus punae</i>	no	NA	NA	NA	mainland	-30	31.83	1.50	0.5	3.82	2.87	Cei 1986, Boretto et al. 2007, Cei et al. 1983, Pincheira-Donoso et al. 2008, Ibarguengoytia et al. 2008, Boretto and Ibarguengoytia 2009, Espinoza et al. 2004, Cabezas et al. 2010, Ibarguengoytia 2008

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Liolaemidae	<i>Phymaturus zapalensis</i>	no	NA	NA	NA	mainland	-40	26.00	1.00	0.75	3.40	2.55	Cei 1986, Boretto and Ibarguengoytia 2006, Cei and Castro 1973, Scolaro 2006, Pincheira-Donoso et al. 2008, Cei 1986, Boretto et al. 2007, Cei et al. 1983, Pincheira-Donoso et al. 2008, Boretto and Ibarguengoytia 2009, Espinoza et al. 2004, Sinervo et al. 2010, Cabezas et al. 2010, Ibarguengoytia 2008, Ibarguengoytia et al. 2008, Cruz et al. 2009
Phrynosomatidae	<i>Callisaurus draconoides</i>	no	NA	NA	NA	mainland	33	13.50	4.30	3	1.15	14.81	Dunham and Miles 1985, Fitch 1970, 1985, Huey and Pianka 1981, Melville et al. 2006, Cox et al. 2003, Stebbins 2003, Grismer 2002, Ortega-Rubio et al. 1995, Pianka 1986, Degenhardt et al. 1996, Pianka and Vitt 2003, Smith 1946, Linsdale 1932, Rogner 1997a, Judd 1976, Van Denburgh 1922, Nagy et al. 1999, Dunham et al. 1988, Maisano 2001, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Warne and Charnov 2008, Huey et al. 2001, Bergmann et al. 2009, Hardy and McDiarmid 1969, Turner 1977, Sinervo et al. 2010, Kohler 2005, Grismer 1994, Jones and Lovich 2009, Case 1975, Huey and Pianka 2007, Turner 1977, Krekorian 1983, Morton and James 1988, Bury 1982, Asplund 1967, Brennan and Holycross 2009, Soule 1963, Cunningham 1966,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Brattstrom 1965, Shine and Schwarzkopf 1992, Stebbins and McGinnis 2012, Lemm 2006, St. John 2002
Phrynosomatidae	<i>Cophosaurus texanus</i>	no	NA	NA	NA	mainland	30	6.65	4.55	3.5	0.65	10.41	Clobert et al. 1998, Dunham and Miles 1985, Vitt et al. 1978, Melville et al. 2006, Tinkle et al. 1970, Fitch 1985, Perry and Garland 2002, Cox et al. 2003, Conant and Collins 1998, Stebbins 2003, Degenhardt et al. 1996, Judd 1976, Punzo 2007, Van Denburgh 1922, Dunham et al. 1988, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Cooper et al. 2001, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Turner 1977, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Curry-Lindahl 1979, Castanet 1994, Brennan and Holycross 2009, Barbault and Maury 1981
Phrynosomatidae	<i>Holbrookia lacerata</i>	no	NA	NA	NA	mainland	29	6.37	8.00	2	0.42	6.79	Fitch 1970, Conant and Collins 1998, Rogner 1997a, Axtell 1956, Lemos-Espinal and Smith 2007, Kohler 2005

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Holbrookia maculata</i>	no	NA	NA	NA	mainland	34	6.11	5.72	2	0.48	5.53	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Vitt et al. 1978, Melville et al. 2006, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Degenhardt et al. 1996, Smith 1946, Greene 1982, Schmidt 1921, Axtell 1956, Judd 1976, Van Denburgh 1922, McCranie and Wilson 2001, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne and Charnov 2008, Hardy and McDiarmid 1969, Lemos-Espinal and Smith 2007, Ibarquengoytia and Casalins 2007, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Curry-Lindahl 1979, Tanner 1987, Brennan and Holycross 2009, Barbault and Maury 1981
Phrynosomatidae	<i>Holbrookia propinqua</i>	no	NA	NA	NA	mainland	28	5.38	3.86	3	0.86	9.92	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1985, Cox et al. 2003, Conant and Collins 1998, Smith 1946, Cooper and Guillette 1991, Judd 1976, Judd and Ross 1978, Dunham et al. 1988, Cooper et al. 2001, Judd 1975, Axtell 1983, Sinervo et al. 2010, Kohler 2005
Phrynosomatidae	<i>Phrynosoma blainvillii</i>	no	NA	NA	NA	mainland	35	14.66	13.00	1	1.15	14.92	Van Denburgh 1922, Jones and Lovich 2009, Cowles and Bogert 1944, Gerson 2011, Goldberg 2011, Stebbins and McGinnis 2012, Woolrich-Pina et al. 2012

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Phrynosoma cornutum</i>	no	NA	NA	NA	mainland	31	15.19	25.25	1.5	0.48	18.32	Fitch 1970, 1985, Vitt et al. 1978, Perry and Garland 2002, Stebbins 2003, Grismer 2002, Pianka and Parker 1975, Sherbrooke 2003, Degenhardt et al. 1996, Pianka and Vitt 2003, Smith 1946, Rogner 1997a, Van Denburgh 1922, Lemos-Espinal et al. 2004, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne and Charnov 2008, Bergmann et al. 2009, Lemos-Espinal and Smith 2007, Price 1990, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Beane et al. 2010, Brennan and Holycross 2009, Barbault and Maury 1981, Van Wilgen and Richardson 2012, Brattstrom 1965, Woolrich-Pina et al. 2012
Phrynosomatidae	<i>Phrynosoma coronatum</i>	no	NA	NA	NA	mainland	33	18.19	11.80	1.5	1.37	24.29	Fitch 1970, Melville et al. 2006, Conant and Collins 1998, Stebbins 2003, Pianka and Parker 1975, Sherbrooke 2003, Smith 1946, Linsdale 1932, Rogner 1997a, Van Denburgh 1922, Vitt and Price 1982, Sinervo et al. 2010, Kohler 2005, Goldberg 1983, Case 1975, Cunningham 1966, Brattstrom 1965, Goldberg 2011, Woolrich-Pina et al. 2012, Lemm 2006

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Phrynosoma ditmarsii</i>	no	NA	NA	NA	mainland	30	12.73	7.50	1	0.69	5.19	Sherbrooke 1997, 1998, Pianka and Parker 1975, Sherbrooke 2003, Smith 1946, Rogner 1997a, Zamudio & Parra-Olea 2000, Bergmann et al. 2009, Hodges 1995, Montanucci 1989
Phrynosomatidae	<i>Phrynosoma douglassii</i>	no	NA	NA	NA	mainland	44	12.83	15.00	1	0.62	9.26	Fitch 1970, 1985, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Pianka and Parker 1975, Sherbrooke 2003, Degenhardt et al. 1996, Smith 1946, Rogner 1997a, Van Denburgh 1922, Zamudio & Parra-Olea 2000, Dunham et al. 1988, Andrews and Pough 1980, Vitt and Price 1982, Warne and Charnov 2008, Bergmann et al. 2009, Ibarguengoytia and Casalins 2007, Sinervo et al. 2010, Jones and Lovich 2009, Shine and Charnov 1992, Christian 1988, Stebbins and McGinnis 2012, Woolrich-Pina et al. 2012, St. John 2002
Phrynosomatidae	<i>Phrynosoma hernandesi</i>	no	NA	NA	NA	mainland	41	27.50	18.00	1	0.86	15.42	Conant and Collins 1998, Stebbins 2003, Sherbrooke 2003, Smith 1946, Van Denburgh 1922, Zamudio & Parra-Olea 2000, Lemos-Espinal and Smith 2007b, Jones and Lovich 2009, Russell and Bauer 2000, Brennan and Holycross 2009, St. John 2002

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Phrynosoma mcallii</i>	no	NA	NA	NA	mainland	33	11.50	4.95	1.5	1.26	9.33	Fitch 1970, Stebbins 2003, Grismer 2002, Pianka and Parker 1975, Sherbrooke 2003, Smith 1946, Rogner 1997a, Van Denburgh 1922, Andrews and Pough 1980, Vitt and Price 1982, Bergmann et al. 2009, Wone and Beauchamp 2003, Funk 1981, Mcgrann et al. 2006, Grant 2005, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Curry-Lindahl 1979, Goldberg 2011, Brennan and Holycross 2009, Stebbins and McGinnis 2012, Woolrich-Pina et al. 2012, Lemm 2006
Phrynosomatidae	<i>Phrynosoma modestum</i>	no	NA	NA	NA	mainland	30	7.41	11.00	1.5	0.37	6.11	Fitch 1970, Vitt et al. 1978, Melville et al. 2006, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Sherbrooke 2002, Pianka and Parker 1975, Sherbrooke 2003, Degenhardt et al. 1996, Smith 1946, Rogner 1997a, Van Denburgh 1922, Lemos-Espinal et al. 2004, McCranie and Wilson 2001, Vitt and Price 1982, Warne and Charnov 2008, Bergmann et al. 2009, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Kohler 2005, Jones and Lovich 2009, Brennan and Holycross 2009, Barbault and Maury 1981, Lemos-Espinal et al. 1997, Charnov et al. 2007, Woolrich-Pina et al. 2012

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Phrynosoma platyrhinos</i>	no	NA	NA	NA	mainland	38	17.31	7.32	1.5	0.77	8.46	Clobert et al. 1998, Dunham and Mills 1985, Fitch 1970, 1985, Huey and Pianka 1981, Melville et al. 2006, Perry and Garland 2002, Cox et al. 2003, Stebbins 2003, Grismer 2002, Pianka and Parker 1975, Pianka 1986, Sherbrooke 2003, Smith 1946, Greene 1982, Rogner 1997a, Van Denburgh 1922, Nagy et al. 1999, Dunham et al. 1988, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne and Charnov 2008, Huey et al. 2001, Bergmann et al. 2009, Turner 1977, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Huey and Pianka 2007, Turner 1977, Morton and James 1988, Bury 1982, Brennan and Holycross 2009, Stebbins and McGinnis 2012, Goldberg 2012, Woolrich-Pina et al. 2012, Lemm 2006, St. John 2002
Phrynosomatidae	<i>Phrynosoma solare</i>	no	NA	NA	NA	mainland	29	28.44	18.60	1.5	1.31	36.65	Fitch 1970, Melville et al. 2006, Perry and Garland 2002, Stebbins 2003, Grismer 2002, Pianka and Parker 1975, Sherbrooke 2003, Degenhardt et al. 1996, Degenhardt et al. 1996, Smith 1946, Rogner 1997a, Van Denburgh 1922, Bogert and Oliver 1945, Vitt and Price 1982, Warne and Charnov 2008, Bergmann et al. 2009, Hardy and McDiarmid 1969, Sinervo et al. 2010, Kohler

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2005, Jones and Lovich 2009, Brennan and Holycross 2009, Parker 1971
Phrynosomatidae	<i>Sceloporus aeneus</i>	no	NA	NA	NA	mainland	20	4.09	5.61	2	0.62	6.92	Fitch 1985, Cox et al. 2003, Fitch 1978, Benabib et al. 1997, Duellman 1961, Radder et al. 2008, Davis and Smith 1953, Sinervo et al. 2010, Kohler 2005, Duellman 1965, Andrews et al. 1999, Guillette 1982, Dixon and Lemos-Espinal 2010
Phrynosomatidae	<i>Sceloporus angustus</i>	yes	Galapagos Archipelago	Santa Cruz	985.60	Oceanic	25	9.87	6.00	NA	NA	NA	Herrel et al. 2002, Grismer 2002, Van Denburgh 1922 Conant and Collins 1998, Stebbins 2003, Degenhardt et al. 1996, Jones and Lovich 2009
Phrynosomatidae	<i>Sceloporus arenicolus</i>	no	NA	NA	NA	mainland	33	6.03	4.50	1.5	0.57	3.88	Clobert et al. 1998, Dunham and Miles 1985, Vitt et al. 1978, Melville et al. 2006, Fitch 1978, 1985, Cox et al. 2003, Stebbins 2003, Grismer 2002, Degenhardt et al. 1996, Smith 1939, Smith 1946, Martins 1993, Benabib 1994, Van Denburgh 1922, Valdez-Gonzalez and Ramirez-Bautista 2002, McCranie and Wilson 2001, Dunham et al. 1988, Bogert and Oliver 1945, Vitt and Price 1982, Warne and Charnov 2008, Cooper et al. 2001, Bergmann et al. 2009, Hardy and
Phrynosomatidae	<i>Sceloporus clarkii</i>	no	NA	NA	NA	mainland	29	25.46	13.80	1.5	1.31	27.19	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus consobrinus</i>	no	NA	NA	NA	mainland	30	9.84	7.70	2.5	0.55	10.55	McDiarmid 1969, Lemos-Espinal and Smith 2007b, Kohler 2005, Jones and Lovich 2009, Rorabaugh 2008, Brennan and Holycross 2009, Charnov et al. 2007, Smith 1946, Van Denburgh 1922, Warne and Charnov 2008, Vinegar 1975, Sinervo et al. 2010, Jones and Lovich 2009, Charnov et al. 2007, Bogert 1949
Phrynosomatidae	<i>Sceloporus cozumelae</i>	no	NA	NA	NA	mainland	21	3.83	1.80	1.5	0.62	1.67	Fitch 1970, 1978, 1985, Lee 2000, Kohler 2003, Smith 1939, Ramirez-Bautista & Olivera-Becerril 2004, Valdez-Gonzalez and Ramirez-Bautista 2002, Kohler 2008, Lopez and Gonzalez 1997, Sinervo et al. 2010, Duellman 1965, Bustos-Zagal et al. 2011, Hews 2012
Phrynosomatidae	<i>Sceloporus cyanogenys</i>	no	NA	NA	NA	mainland	26	37.14	13.00	1	1.13	14.65	Conant and Collins 1998, Bartlett and Bartlett 2006, Warne and Charnov 2008, Dunham et al. 1988, Hunsaker 1959, Lemos-Espinal and Smith 2007, Mendez-de la Cruz et al. 1998, Ramirez-Pinilla et al. 2009, Sinervo et al. 2010, Sinervo et al. 2010, Ramirez-Bautista and Davila-Ulloa 2009
Phrynosomatidae	<i>Sceloporus formosus</i>	no	NA	NA	NA	mainland	17	11.99	7.02	1	1.11	7.76	Fitch 1970, 1978, Cox et al. 2002, Savage 2002, Smith 1939, Martins 1993, Rogner 1997a, Boretto and Ibarquengoytia 2006, Davis and Dixon 1961, Perez & de

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus gadoviae</i>	no	NA	NA	NA	mainland	18	5.55	3.75	1.5	0.69	3.89	La Riva 2008, Mendez-de la Cruz et al. 1998, Ramirez-Pinilla et al. 2009, Acevedo 2009 Ramirez-Bautista and Gutierrez-Mayen 2003, Ramirez-Bautista et al. 2005, Smith 1939, Ramirez-Bautista & Olivera-Becerril 2004, Valdez-Gonzalez and Ramirez-Bautista 2002, Duellman 1961, Lemos-espinal et al. 1999, Davis and Dixon 1961, Davis and Smith 1953, Serrano-Cardozo et al. 2008, Sinervo et al. 2010, Duellman 1965, Bustos-Zagal et al. 2011, Lemos-Espinal et al. 1997, Lemos-Espinal et al. 1997, Woolrich-Pina et al. 2012 Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1978, 1985, Vitt et al. 1978, Melville et al. 2006, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Degenhardt et al. 1996, Smith 1946, Woodbury 1932, Benabib 1994, Van Denburgh 1922, Nagy et al. 1999, Valdez-Gonzalez and Ramirez-Bautista 2002, Dunham et al. 1988, Andrews and Pough 1980, Brown and Nagy 2007, Vitt and Price 1982, Warne and Charnov 2008, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Andrews 1998,
Phrynosomatidae	<i>Sceloporus graciosus</i>	no	NA	NA	NA	mainland	41	6.37	4.35	1.5	0.77	5.03	Benabib 1994, Van Denburgh 1922, Nagy et al. 1999, Valdez-Gonzalez and Ramirez-Bautista 2002, Dunham et al. 1988, Andrews and Pough 1980, Brown and Nagy 2007, Vitt and Price 1982, Warne and Charnov 2008, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Andrews 1998,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													van Berkum 1988, Turner 1977, Tinkle et al. 1967, Brennan and Holycross 2009, Cunningham 1966, Bustos-Zagal et al. 2011, Shine and Charnov 1992, Brattstrom 1965, St. John 2002
Phrynosomatidae	<i>Sceloporus grammicus</i>	no	NA	NA	NA	mainland	24	4.85	4.76	1	0.32	1.52	Clobert et al. 1998, Fitch 1970, 1978, 1985, Perry and Garland 2002, Herrel et al. 2002, Barbault et al. 1985, JimeNez-Cruz et al. 2005, Ramirez-Bautista et al. 2004, Smith 1939, Smith 1946, Lara-Gongora 1983, McCranie and Wilson 2001, Dunham et al. 1988, Duellman 1961, Radder et al. 2008, Werler 1951, Lemos-Espinal and Smith 2007, Davis and Dixon 1961, Davis and Smith 1953, Mendez-de la Cruz et al. 1998, Hernandez-Salinas et al. 2010, Sinervo et al. 2010, Duellman 1965, Andrews 1998, Dixon and Lemos-Espinal 2010, Bogert 1949

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus horridus</i>	no	NA	NA	NA	mainland	22	18.43	11.90	2	0.62	14.69	Fitch 1970, Baker et al. 1967, Ramirez-Bautista and Gutierrez-Mayen 2003, Valdez-Gonzalez and Ramirez-Bautista 2002, Smith 1939, Martins 1993, McCranie and Wilson 2001, Duellman 1961, Todd 2008, Hardy and McDiarmid 1969, Davis and Dixon 1961, Davis and Smith 1953, Lemos-Espinal and Smith 2007b, Serrano-Cardozo et al. 2008, Sinervo et al. 2010, Duellman 1965, Lemos-Espinal et al. 2001, Lemos-Espinal et al. 1997, Ramirez-Bautista et al. 2012
Phrynosomatidae	<i>Sceloporus jarrovii</i>	no	NA	NA	NA	mainland	27	13.77	6.70	1	1.10	7.34	Tinkle et al. 1970, Clobert et al. 1998, Fitch 1970, 1985, Melville et al. 2006, Perry and Garland 2002, Cox et al. 2003, Stebbins 2003, Herrel et al. 2002, Barbault et al. 1985, Degenhardt et al. 1996, Smith 1946, Axtell and Axtell 1971, Benabib 1994, Rogner 1997a, Van Denburgh 1922, Nagy et al. 1999, McCranie and Wilson 2001, Dunham et al. 1988, Brown and Nagy 2007, Vitt and Price 1982, Cooper et al. 2001, Radder et al. 2008, Bergmann et al. 2009, Hardy and McDiarmid 1969, Lemos-Espinal and Smith 2007, Ibarquengoytia and Casalins 2007, Davis and Smith 1953, Mendez-de la Cruz et al. 1998, Lemos-Espinal and Smith 2007b,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Ramirez-Pinilla et al. 2009, Jones and Lovich 2009, Sinervo et al. 2010, Andrews 1998, Brennan and Holycross 2009, Ramirez-Bautista and Davila-Ulloa 2009, Van Wilgen and Richardson 2012, Shine and Charnov 1992, Brattstrom 1965, Goldberg and Beaman 2012
Phrynosomatidae	<i>Sceloporus magister</i>	no	NA	NA	NA	mainland	33	25.39	9.30	2	1.90	35.32	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1978, 1985, Vitt et al. 1978, Melville et al. 2006, Conant and Collins 1998, Stebbins 2003, Herrel et al. 2002, Grismer 2002, Pianka 1986, Degenhardt et al. 1996, Smith 1939, Smith 1946, Greene 1982, Martins 1993, Linsdale 1932, Rogner 1997a, Van Denburgh 1922, Valdez-Gonzalez and Ramirez-Bautista 2002, Dunham et al. 1988, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne and Charnov 2008, Cooper et al. 2001, Huey et al. 2001, Bergmann et al. 2009, Hardy and McDiarmid 1969, Lemos-Espinal and Smith 2007, Vitt & Ohmart 1975, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Case 1975,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus malachiticus</i>	no	NA	NA	NA	mainland	14	12.25	5.25	1	1.15	6.03	Huey and Pianka 2007, Andrews 1998, Rorabaugh 2008, Morton and James 1988, Bury 1982, Brennan and Holycross 2009, Bustos-Zagal et al. 2011, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006, Bogert 1949, Goldberg 2012 (Sceloporus uniformis), St. John 2002 Dunham and Miles 1985, Fitch 1970, 1973, 1978, 1985, Savage 2002, Herrel et al. 2002, Kohler 2003, Smith 1939, Rogner 1997a, McCranie and Castaneda 2005, Dunham et al. 1988, Alvarez del Toro and Smith 1962, Clusella-Trullas et al. 2008, Fitch 1982, Rand 1957, Kohler 2008, Todd 2008, Robinson 1983, Cree and Guillette 1995, Andrews and Rand 1974, Perez & de La Riva 2008, Mendez-de la Cruz et al. 1998, McElroy et al. 2008, Sinervo et al. 2010, Bueter and Haas 2008, Andrews 1998, van Berkum 1988, Brattstrom 1965, Leenders and Watkins-Colwell 2004, Bogert 1949

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus merriami</i>	no	NA	NA	NA	mainland	29	4.36	4.11	1.5	0.55	3.37	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1978, 1985, Perry and Garland 2002, Conant and Collins 1998, Smith 1939, Smith 1946, Benabib 1994, Dunham et al. 1988, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Andrews 1998, Dunham et al. 1989, Bogert 1949
Phrynosomatidae	<i>Sceloporus mucronatus</i>	no	NA	NA	NA	mainland	19	18.46	4.69	1	0.77	3.62	Fitch 1978, Boretto and Ibarquengoytia 2006, Radder et al. 2008, Mendez-de la Cruz et al. 1992, Ortega-Leon et al. 2007, Todd 2008, Werler 1949, Webb et al. 2002, Mendez-de la Cruz et al. 1998, Ramirez-Pinilla et al. 2009, Sinervo et al. 2010, Ramirez-Bautista and Davila-Ulloa 2009
Phrynosomatidae	<i>Sceloporus occidentalis</i>	no	NA	NA	NA	mainland	40	13.55	10.10	2	0.95	19.15	Tinkle et al. 1970, Clobert et al. 1998, Fitch 1970, 1978, 1985, Huey and Pianka 1981, Melville et al. 2006, Perry and Garland 2002, Stebbins 2003, Herrel et al. 2002, Grismer 2002, Smith 1946, Martins 1993, Linsdale 1932, Rogner 1997a, Van Denburgh 1922, Nagy et al. 1999, Roe et al. 2005, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Radder et al. 2008, Sinervo et al. 2010,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus olivaceus</i>	no	NA	NA	NA	mainland	29	24.45	14.65	2.5	0.69	25.33	Kohler 2005, Jones and Lovich 2009, Andrews 1998, van Berkum 1988, Cunningham 1966, Vitt 1974, Brattstrom 1965, Lemm 2006, St. John 2002
Phrynosomatidae	<i>Sceloporus omiltemanus</i>	no	NA	NA	NA	mainland	18	18.69	6.23	1	1.11	6.89	Tinkle et al. 1970, Clobert et al. 1998, Fitch 1970, 1978, Perry and Garland 2002, Conant and Collins 1998, Smith 1939, Smith 1946, Martins 1993, Benabib 1994, Valdez-Gonzalez and Ramirez-Bautista 2002, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Bergmann et al. 2009, Lemos-Espinal and Smith 2007, Kenedy 1973, Dutton et al. 1975, Jones and Lovich 2009, Sinervo et al. 2010, Andrews 1998, Damuth 1987, Tinkle et al. 1967, Bustos-Zagal et al. 2011, Charnov et al. 2007, Burstein et al. 1974, Davis and Dixon 1961, Ramirez-Pinilla et al. 2009, Sinervo et al. 2010

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus orcutti</i>	no	NA	NA	NA	mainland	30	24.41	10.10	1	1.62	16.38	Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1978, Perry and Garland 2002, Stebbins 2003, Grismer 2002, Smith 1939, Smith 1946, Greene 1982, Linsdale 1932, Van Denburgh 1922, Valdez-Gonzalez and Ramirez-Bautista 2002, Dunham et al. 1988, Clusella-Trullas et al. 2008, Todd 2008, Hall and Smith 1979, Weintraub 1980, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Case 1975, Andrews 1998, Muchlinski et al. 1995, Turner 1977, Soule 1963, Cunningham 1966, Bustos-Zagal et al. 2011, Stebbins and McGinnis 2012, Lemm 2006
Phrynosomatidae	<i>Sceloporus poinsettii</i>	no	NA	NA	NA	mainland	29	29.41	10.40	1	1.49	15.54	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1978, 1985, Conant and Collins 1998, Stebbins 2003, Barbault et al. 1985, Cooper and Vitt 2002, Degenhardt et al. 1996, Smith 1946, Greene 1982, Benabib 1994, Rogner 1997a, Van Denburgh 1922, Dunham et al. 1988, Warne and Charnov 2008, Lemos-Espinal and Smith 2007, Webb 2006, Webb 2008, Mendez-de la Cruz et al. 1998, Lemos-Espinal and Smith 2007b, Ramirez-Pinilla et al. 2009, Jones and Lovich 2009, Sinervo et al. 2010, Andrews 1998, Turner 1977, Ramirez-Bautista and Davila-

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Ulloa 2009, Shine and Charnov 1992, Charnov et al. 2007, Bogert 1949
Phrynosomatidae	<i>Sceloporus scalaris</i>	no	NA	NA	NA	mainland	24	5.45	8.74	1.5	0.42	5.56	Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1978, Vitt et al. 1978, Melville et al. 2006, Perry and Garland 2002, Barbault et al. 1985, Degenhardt et al. 1996, Smith 1939, Benabib et al. 1997, Benabib 1994, Van Denburgh 1922, Ramirez-Bautista & Olivera-Becerril 2004, Valdez-Gonzalez and Ramirez-Bautista 2002, McCranie and Wilson 2001, Dunham et al. 1988, Duellman 1961, Vitt and Price 1982, Warne and Charnov 2008, Davis and Smith 1953, Sinervo et al. 2010, Kohler 2005, Duellman 1965, Andrews 1998, Dixon and Lemos-Espinal 2010, Shine and Charnov 1992, Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1978, 1985, Campbell 1999, Stafford and Meyer 2000, Conant and Collins 1998, Herrel et al. 2002, Lee 2000, Kohler 2003, Smith 1939, Smith 1942, Rogner 1997a,
Phrynosomatidae	<i>Sceloporus serrifer</i>	no	NA	NA	NA	mainland	23	30.38	15.25	1	1.21	18.49	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus slevini</i>	no	NA	NA	NA	mainland	28	4.55	8.50	1	0.42	3.61	Dunham et al. 1988, Alvarez del Toro and Smith 1962, Goldberg et al. 1994, Kohler 2008, Sinervo et al. 2010, Dixon and Lemos-Espinal 2010
Phrynosomatidae	<i>Sceloporus torquatus</i>	no	NA	NA	NA	mainland	21	24.76	8.05	1	1.15	9.24	Stebbins 2003, Smith 1946, Watkins-Colwell et al. 2003, Kohler 2005, Jones and Lovich 2009, Tanner 1987, Brennan and Holycross 2009, Fitch 1970, Herrel et al. 2002, Fitch 1978, Ramirez-Bautista and Gonzalez-Romero 2002, Boretto and Ibarguengoytia 2006, McCranie and Wilson 2001, Duellman 1961, Radder et al. 2008, Werler 1951, Davis and Smith 1953, Mendez-de la Cruz et al. 1998, Ramirez-Pinilla et al. 2009, Sinervo et al. 2010, Duellman 1965, Dixon and Lemos-Espinal 2010, Ramirez-Bautista and Davila-Ulloa 2009
Phrynosomatidae	<i>Sceloporus undulatus</i>	no	NA	NA	NA	mainland	35	8.97	9.13	3	0.79	21.70	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Vitt et al. 1978, Perry and Garland 2002, Conant and Collins 2002, Stebbins 2003, Fitch 1978, Degenhardt et al. 1996, Smith 1946, Greene 1982, Woodbury 1932, Martins 1993, Benabib 1994, Bowker et al. 1986, Van Denburgh 1922, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus utiformis</i>	no	NA	NA	NA	mainland	20	8.68	7.10	1	0.77	5.48	and Charnov 2008, Radder et al. 2008, Lemos-Espinal and Smith 2007, Ibarquengoytia and Casalins 2007, Vinegar 1975, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, De Magalhaes and Costa 2009, Jones and Lovich 2009, Jensen et al. 2008, Andrews 1998, Turner 1977, Heatwole and Taylor 1987, Beane et al. 2010, Gibbs et al. 2007, Brennan and Holycross 2009, Barbault and Maury 1981, Shine and Charnov 1992, St. John 2002 Ramirez-Bautista and Gutierrez-Mayen 2003, Fitch 1978, Smith 1939, Ramirez-Bautista & Olivera-Becerril 2004, Valdez-Gonzalez and Ramirez-Bautista 2002, Duellman 1961, Hardy and McDiarmid 1969, Duellman 1965 Clobert et al. 1998, Fitch 1970, 1973, 1985, Stafford and Meyer 2000, Conant and Collins 1998, Savage 2002, Lee 2000, Fitch 1978, Vitt et al. 1993, Kohler 2003, Smith 1946, Benabib 1994, Nagy et al. 1999, Kohler et al. 2006, Ramirez-Bautista & Olivera-Becerril 2004, Valdez-Gonzalez and Ramirez-Bautista 2002, Janzen 1973, Brown and Nagy 2007, Rand 1957, Werler 1951, Lemos-Espinal and Smith 2007,
Phrynosomatidae	<i>Sceloporus variabilis</i>	no	NA	NA	NA	mainland	20	6.12	3.85	5	0.80	15.33	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Sceloporus virgatus</i>	no	NA	NA	NA	mainland	29	6.35	9.83	1	0.52	5.06	Sinervo et al. 2010, Kohler 2005, Andrews 1998, van Berkum 1988, Lemos-Espinal et al. 2001, Dixon and Lemos-Espinal 2010, Leenders and Watkins-Colwell 2004, Townsend and Wilson 2008, Bogert 1949
Phrynosomatidae	<i>Sceloporus woodi</i>	no	NA	NA	NA	mainland	28	6.37	4.81	2.5	0.48	5.81	Clobert et al. 1998, Dunham and Miles 1985, Vitt et al. 1978, Melville et al. 2006, Fitch 1985, Perry and Garland 2002, Stebbins 2003, Fitch 1978, Degenhardt et al. 1996, Benabib 1994, Nagy et al. 1999, Dunham et al. 1988, Vitt and Price 1982, Warne and Charnov 2008, Cooper et al. 2001, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Andrews 1998, Turner 1977, Brennan and Holycross 2009, Shine and Charnov 1992, Fitch 1985, Conant and Collins 1998, Fitch 1978, Smith 1946, McCoy et al. 2004, Greene 1982, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Todd 2008, Lee & Funderburg 1977, Sinervo et al. 2010, Kohler 2005, Andrews 1998, Turner 1977, Curry-Lindahl 1979, Charnov et al. 2007, Bogert 1949

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Uma exsul</i>	no	NA	NA	NA	mainland	26	10.54	3.00	1.5	1.49	6.72	Schmidt and Bogert 1947, Gadsden et al. 2006, Commins and Savitzky 1973, Lemos-Espinal and Smith 2007, Sinervo et al. 2010, Barbault and Maury 1981, Pough et al. 1978 Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1985, Stebbins 2003, Grismer 2002, Ortega-Rubio et al. 1995, Smith 1946, Rogner 1997a, Van Denburgh 1922, Dunham et al. 1988, Bergmann et al. 2009, Todd 2008, Jones and Lovich 2009, Sinervo et al. 2010, Curry-Lindahl 1979, Brennan and Holycross 2009, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006, Goldberg 2012 Fitch 1970, Cox et al. 2003, Baker et al. 1967, Ramirez-Bautista and Vitt 1998, Kohler 2003, Franco and de la Torre 1990, McCranie and Wilson 2001, Duellman 1961, Kohler 2008, Hardy and McDiarmid 1969, Davis and Dixon 1961, Davis and Smith 1953, Lemos-Espinal and Smith 2007b, Duellman 1965 Brattstrom 1982, Brattstrom 1955, Sinervo et al. 2010
Phrynosomatidae	<i>Uma notata</i>	no	NA	NA	NA	mainland	32	17.07	2.30	2.5	2.28	13.13	
Phrynosomatidae	<i>Urosaurus bicarinatus</i>	no	NA	NA	NA	mainland	20	3.63	6.55	1.5	0.32	3.15	
Phrynosomatidae	<i>Urosaurus clarionensis</i>	yes	Revillagigedo Islands	Clarion	19.80	Oceanic	18	5.34	NA	NA	1.15	NA	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phrynosomatidae	<i>Urosaurus graciosus</i>	no	NA	NA	NA	mainland	34	5.09	4.55	1.5	0.69	4.72	Dunham and Miles 1985, Vitt et al. 1978, Melville et al. 2006, Fitch 1985, Stebbins 2003, Herrel et al. 2002, Grismer 2002, Pianka 1986, Pianka and Vitt 2003, Smith 1946, Greene 1982, Rogner 1997a, Van Denburgh 1922, Vitt and Price 1982, Huey et al. 2001, Vitt and Dickson 1988, Vitt & Ohmart 1975, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Huey and Pianka 2007, Rorabaugh 2008, Brennan and Holycross 2009, Stebbins and McGinnis 2012, Goldberg 2012, Lemm 2006 Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Vitt et al. 1978, Melville et al. 2006, Perry and Garland 2002, Cox et al. 2003, Conant and Collins 1998, Stebbins 2003, Herrel et al. 2002, Grismer 2002, Pianka and Vitt 2003, Degenhardt et al. 1996, Smith 1946, Rogner 1997a, Van Denburgh 1922, Dunham et al. 1988, Bogert and Oliver 1945, Clusella-Trullas et al. 2008, Vitt and Price 1982, Warne and Charnov 2008, Ballinger 1977, Cooper et al. 2001, Radder et al. 2008, Hardy and McDiarmid 1969, Lemos-Espinal and Smith 2007, Vitt & Ohmart 1975, Lemos-Espinal and Smith 2007b, Turner 1977, Sinervo
Phrynosomatidae	<i>Urosaurus ornatus</i>	no	NA	NA	NA	mainland	33	3.98	7.40	3.5	0.47	12.13	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													et al. 2010, Kohler 2005, Jones and Lovich 2009, Rorabaugh 2008, Brennan and Holycross 2009, Shine and Charnov 1992, Stebbins and McGinnis 2012, St. John 2002
Phrynosomatidae	<i>Uta palmeri</i>	yes	Baja California	San Pedro Martir	1.22	Oceanic	29	8.31	3.00	1.5	NA	NA	Grismer 2002, Case 2002, Sinervo et al. 2010, Grismer 1994 Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Soule 1966, Huey and Pianka 1981, Melville et al. 2006, Stebbins 2003, Grismer 2002, Ortega-Rubio et al. 1995, Pianka 1986, Sinervo and Licht 1991, Pianka and Vitt 2003, Degenhardt et al. 1996, Smith 1946, Woodbury 1932, Benabib 1994, Linsdale 1932, Rogner 1997a, Dunham et al. 1988, Andrews and Pough 1980, Maisano 2001, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Vitt and Price 1982, Warne and Charnov 2008, Cooper et al. 2001, Huey et al. 2001, Huey et al. 2001, Lemos-Espinal and Smith 2007, Andrews and Rand 1974, Lemos-Espinal and Smith 2007b, Turner
Phrynosomatidae	<i>Uta stansburiana</i>	no	NA	NA	NA	mainland	35	3.92	3.83	4	0.52	7.88	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1977, Sinervo et al. 2010, Kohler 2005, Grismer 1994, Jones and Lovich 2009, Milton et al. 2004, Case 1975, Huey and Pianka 2007, Turner 1977, Damuth 1987, Morton and James 1988, Curry-Lindahl 1979, Bury 1982, Brennan and Holycross 2009, Barbault and Maury 1981, Soule 1963, Scoular et al. 2011, Cunningham 1966, Shine and Charnov 1992, Stebbins and McGinnis 2012, Goldberg 2012, Lemm 2006, St. John 2002
Phyllodactylidae	<i>Gymnodactylus amarali</i>	no	NA	NA	NA	mainland	-13	1.56	1.65	1	0.21	0.34	Cassimiro & Rodrigues 2009, Colli et al. 2003, Vanzolini 2005, Vitt et al. 2007, Sinervo et al. 2010, Rocha et al. 2009 Fitch 1970, Colli et al. 2003, Cox et al. 2003, Vitt 1986, Pianka and Vitt 2003, Mesquita et al. 2006b, Rogner 1997a, Cree 1994, Colli et al. 2002, Vitt et al. 2007, Dunham et al. 1988, Rodrigues 2003, Rodrigues 1996, Warne and Charnov 2008, Murphy 1997, Vitt 1995, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler 2005, Rosler 2005, Daza et al. 2009, Pellegrino et al. 2005
Phyllodactylidae	<i>Gymnodactylus geckoides</i>	no	NA	NA	NA	mainland	-8	1.86	1.35	3	0.22	0.90	Loveridge 1947, Schatti and Desvoignes 1999, Rosler 2005, Razzetti et al. 2011,
Phyllodactylidae	<i>Haemodracon riebeckii</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	13	54.11	1.50	NA	0.90	NA	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Rosler and Wranik 2004, Rosler and Wranik 2005
Phyllodactylidae	<i>Haemodracon trachyrhinus</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	13	2.84	1.00	NA	0.21	NA	Loveridge 1947, Schatti and Desvoignes 1999, Boulenger 1889, Rosler 2005, Razzetti et al. 2011, Rosler and Wranik 2005
Phyllodactylidae	<i>Homonota darwinii</i>	no	NA	NA	NA	mainland	-44	1.35	1.00	0.75	0.25	0.19	Cox et al. 2003, Cei 1986, Kluge 1964, Scolaro 2005, Boretto and Ibaranguoytia 2009, Ibaranguoytia and Casalins 2007, Ibaranguoytia 2008, Aguilar and Cruz 2010, Ibaranguoytia et al. 2007, Kubisch et al. 2012
Phyllodactylidae	<i>Homonota fasciata</i>	no	NA	NA	NA	mainland	-23	2.68	1.25	2	0.47	1.18	Fitzgerald et al. 1999, Scolaro 2006, Ibaranguoytia and Casalins 2007, Largen and Spawls 2010, Ibaranguoytia 2008, Aguilar and Cruz 2010, Werner et al. 1993
Phyllodactylidae	<i>Phyllodactylus bugastrolepis</i>	yes	Baja California	Santa Catalina	194.20	Continental	26	2.68	1.50	NA	NA	NA	Grismer 2002
Phyllodactylidae	<i>Phyllodactylus lanei</i>	no	NA	NA	NA	mainland	19	5.70	2.00	3	0.27	1.63	Ramirez-Sandoval et al. 2006, Dixon 1964, Franco and de la Torre 1990, Duellman 1961, Herzog and Drummond 1984, Davis and Dixon 1961, Mautz 1982
Phyllodactylidae	<i>Phyllopezus pollicaris</i>	no	NA	NA	NA	mainland	-18	8.06	2.00	1.5	0.68	2.03	Cox et al. 2003, Vitt 1986, Pianka and Vitt 2003, Cei 1986, Cei 1993, Cree 1994, Colli et al. 2002, Dunham et al. 1988, Rodrigues 2003, Rodrigues 1996, Vitt 1995, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler 2005, Rocha and Rodrigues 2005, Recoder et al. 2012, Winck

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													and Rocha 2012, Righi et al. 2012
Phyllodactylidae	<i>Ptyodactylus guttatus</i>	no	NA	NA	NA	mainland	29	6.67	2.00	4	0.50	3.99	Amitai and Bouskila 2001, Arbel 1984, Disi et al. 2001, El Din 2006, Le Berre 1989, Johnston and Bouskila 2007, Werner 1986, Arad 1995, Werner and Sivan 1993, Werner and Sivan 1994, Frankenberg 1978, Werner 1989, Bar and Haimovitch 2012, Werner et al. 1993
Phyllodactylidae	<i>Ptyodactylus oudrii</i>	no	NA	NA	NA	mainland	32	2.79	1.50	6	0.54	4.87	Schleich et al. 1996, Geniez et al. 2004, Loveridge 1947, Le Berre 1989, Bons and Geniez 1996, Bons and Geniez 1996, Trape et al. 2012, Werner et al. 1993
Phyllodactylidae	<i>Ptyodactylus ragazzii</i>	no	NA	NA	NA	mainland	18	11.71	2.00	4.5	0.54	4.87	Schleich et al. 1996, Bauer 2006, Bohme et al. 1996, El Din 2006, Chirio and LeBreton 2007, Ineich 1999, Chirio 2009, Kohler 2005, Largen and Spawls 2010, Rosler 2005, Henkel and Schmidt 1995, Trape et al. 2012, Werner et al. 1993
Phyllodactylidae	<i>Tarentola angustimentalis</i>	yes	Canary Islands	Fuerteventura	1633.30	Oceanic	29	4.91	1.92	7.5	0.34	4.88	Arnold and Ovenden 2004, Rogner 1997a, Salvador 2007, Roca 1999, Kohler 2005, Rosler 2005, Maso and Pijoan 2011
Phyllodactylidae	<i>Tarentola annularis</i>	no	NA	NA	NA	mainland	22	20.08	1.50	5	1.25	9.37	Schleich et al. 1996, Geniez et al. 2004, Flower 1933, Papenfuss 1969, Loveridge 1947, El Din 2006, Le Berre 1989, Bons and Geniez 1996,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phyllodactylidae	<i>Tarentola boettgeri</i>	yes	Canary Islands	Gran Canaria	1529.90	Oceanic	28	4.57	1.50	4.5	0.57	3.84	Chirio and LeBreton 2007, Ineich 1999, Largen and Spawls 2006, Kohler 2005, Crochet and Renoult 2008, Largen and Spawls 2010, Rosler 2005, Van Wilgen and Richardson 2012, Trape et al. 2012, Werner et al. 1993 Arnold and Ovenden 2004, Rogner 1997a, Salvador and Brown 2007, Roca 1999, Sinervo et al. 2010, Kohler 2005, Rosler 2005, Brown 1996, Maso and Pijoan 2011 Schleich et al. 1996, Geniez et al. 2004, Rogner 1997a, Le Berre 1989, Bons and Geniez 1996, Kohler 2005, Rosler 2005, Henkel and Schmidt 1995, Trape et al. 2012
Phyllodactylidae	<i>Tarentola chazaliae</i>	no	NA	NA	NA	mainland	26	3.54	1.50	4.5	0.24	1.60	Kohler 2005, Rosler 2005
Phyllodactylidae	<i>Tarentola darwini</i>	yes	Cape Verde	Sao Tiago	991.00	Oceanic	15	4.08	1.00	NA	0.39	NA	Arnold and Ovenden 2004, Loveridge 1947, Hughes 1988, Rogner 1997a, Roca 1999, Kohler 2005, Rosler 2005, Salvador 2009, Maso and Pijoan 2011, Werner et al. 1993
Phyllodactylidae	<i>Tarentola delalandii</i>	yes	Canary Islands	Tenerife	2007.80	Oceanic	28	8.15	1.50	7	0.37	3.88	Loveridge 1947, Bons and Geniez 1996, Kohler 2005, Rosler 2005, Trape et al. 2012, Thili et al. 2012
Phyllodactylidae	<i>Tarentola deserti</i>	no	NA	NA	NA	mainland	33	11.31	2.00	5	1.11	11.11	Arnold and Ovenden 2004, Rogner 1997a, Salvador 2007, Roca 1999, Kohler 2005, Rosler 2005, Maso and Pijoan 2011
Phyllodactylidae	<i>Tarentola gomerensis</i>	yes	Canary Islands	Gomera	359.10	Oceanic	29	5.26	1.50	6	0.21	1.86	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Phyllodactylidae	<i>Tarentola mauritanica</i>	no	NA	NA	NA	mainland	37	5.72	2.00	8.5	0.21	3.51	Arnold and Ovenden 2004, Schleich et al. 1996, Loveridge 1947, Achaval and Olmos 2003, Rogner 1997a, Le Berre 1989, Bons and Geniez 1996, Sindaco et al. 2006, Corti and Cascio 2002, Valakos et al. 2008, Salvador 2008, Radder et al. 2008, Bauer 1990, Kwet 2009, Malkmus 2004, Ibrahim 2008, Carreira et al. 2005, Sinervo et al. 2010, Kohler 2005, Frankenberg 1978, Rosler 2005, Werner 1973, Daza et al. 2009, Werner 1989, Sindaco et al. 2010, Carretero 2008, Castanet 1994, Van Wilgen and Richardson 2012, Henkel and Schmidt 1995, Gil et al. 1994, Arad et al. 1997, Hoder et al. 2006, Trape et al. 2012, Maso and Pijoan 2011, Werner et al. 1993
Polychrotidae	<i>Polychrus acutirostris</i>	no	NA	NA	NA	mainland	-17	30.85	14.75	1	0.83	12.26	Clobert et al. 1998, Avila-Pires 1995, Cooper and Vitt 2002, Cei 1993, Mesquita et al. 2006b, Gainsbury and Colli 2003, Nogueira et al. 2005, Rogner 1997a, Herrel et al. 2004, Bartlett and Bartlett 2003, Colli et al. 2002, Dunham et al. 1988, Rodrigues 2003, Rodrigues 1996, Vitt and Price 1982, Fitch 1982, Vitt 1995, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler 2005, Rand 1982, Tuli et al. 2009, Vanzolini 1983, Garda et al.

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2012, Winck and Rocha 2012
Pygopodidae	<i>Aprasia pulchella</i>	no	NA	NA	NA	mainland	-34	5.80	2.00	1	1.74	3.48	Cox et al. 2003, Webb and Shine 1994, Wiens et al. 2006, Pianka and Vitt 2003, Wilson and Swan 2003, Greer 1989, Wiens et al. 2006, Henkel 2010, Bush et al. 2010
Pygopodidae	<i>Aprasia repens</i>	no	NA	NA	NA	mainland	-32	5.70	2.00	1	1.59	3.17	Cogger 2000, Cox et al. 2003, Webb and Shine 1994, Pianka and Vitt 2003, Wilson and Swan 2003, Greer 1989, Kohler 2005, Henkel 2010, Bush et al. 2010
Pygopodidae	<i>Delma fraseri</i>	no	NA	NA	NA	mainland	-31	5.80	2.00	1	1.64	3.27	Cogger 2000, Patchell and Shine 1986, Pianka 1986, Wilson and Swan 2003, Henle 1991, Rogner 1997a, Greer 1989, Chapman and Dell 1985, Todd 2008, Kohler 2005, Henkel 2010, Bush et al. 2010, Hailey and Elliot 1995
Pygopodidae	<i>Delma grayii</i>	no	NA	NA	NA	mainland	-26	5.14	2.00	1	1.66	3.33	Cogger 2000, Wiens et al. 2006, Wilson and Swan 2003, Henle 1991, Greer 1989, Wilson and Swan 2008, Chapman and Dell 1985, Kohler 2005, Henkel 2010, Bush et al. 2010, Wilson and Swan 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Pygopodidae	<i>Lialis burtonis</i>	no	NA	NA	NA	mainland	-25	12.99	1.85	1	2.91	5.39	Cogger 2000, Patchell and Shine 1986, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Allison 2006, Henle 1991, Greer 1989, Neill 1957, Allison 2007, Wilson and Swan 2008, Huey et al. 2001, Trudgen 1999, Chapman and Dell 1985, Henle 1989c, Bustard 1968, Kohler 2005, Henkel 2010, Bush et al. 2010, Daza et al. 2009, Wilson and Swan 2011, Michael and Lindenmayer 2010, Wilson 2003, Hutchinson 1993, Heatwole and Taylor 1987, Swanson 2007, Pianka 2011, Wilson 2005, Bush et al. 2007, Allison 1982, Sadlier 1990, Moro and MacAulay 2010, Heatwole and Butler 1981, Swan and Watharow 2005 Cogger 2000, Patchell and Shine 1986, Webb and Shine 1994, Wilson and Swan 2003, Rogner 1997a, Greer 1989, Wilson and Swan 2008, Todd 2008, Kohler 2005, Henkel 2010, Bush et al. 2010, Rosler 2005, Bush 1992, Daza et al. 2009, Wilson and Swan 2011, Shea 1993, Swanson 2007, Bush et al. 2007, Swan and Watharow 2005 Arnold and Ovenden 2004, Baran and Atatur 1998, Street 1979, Rogner 1997b, Atatur and Gocmen 2001, Herczeg et al. 2007, Valakos et al. 2008, Valakos et al. 2004, Arbel
Pygopodidae	<i>Pygopus lepidopodus</i>	no	NA	NA	NA	mainland	-32	11.23	2.00	1	2.45	4.90	
Scincidae	<i>Ablepharus kitaibelii</i>	no	NA	NA	NA	mainland	40	1.45	4.00	1	0.12	0.48	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1984, Kwet 2009, Kohler 2005, Schmidtler 1997
Scincidae	<i>Ablepharus rueppellii</i>	no	NA	NA	NA	mainland	32	0.81	4.00	1	0.12	0.48	Amitai and Bouskila 2001, Arbel 1984, Disi et al. 2001, El Din 2006, Bar and Haimovitch 2012, Shine and Greer 1991, FitzSimons 1943, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Greer 1977, Auerbach 1987, Cree 1994, Dunham et al. 1988, Huey et al. 1974, Goldberg 2006, Pianka 1971, Broadley 1968, Fitch 1970, Branch 1998, Pianka 1986, Pianka and Vitt 2003, Broadley 1971, Auerbach 1987, Cree 1994, Dunham et al. 1988, Huey et al. 1974, Goldberg 2006, Cree and Guillette 1995, Pianka 1971, Broadley 1968
Scincidae	<i>Acontias garipeensis</i>	no	NA	NA	NA	mainland	-27	3.97	1.00	1	0.71	0.71	Fuhn 1972
Scincidae	<i>Acontias kgalagadi</i>	no	NA	NA	NA	mainland	-23	4.80	1.48	0.75	0.68	0.75	Mausfeld and Lotters 2001, Harvey et al. 2008, Miralles et al. 2009
Scincidae	<i>Afroablepharus annobonensis</i>	yes	Sao Tome and Principe	Annobon	15.70	Oceanic	-1	0.62	NA	NA	0.07	NA	Tikader and Sharma 1992, Inger and Colwell 1977, Schleich and Kastle 2002, Daniel 1983, Shrestha 2001, Ouboter 1986, Das 2002, Aryal et al. 2010
Scincidae	<i>Aspronema cochabambae</i>	no	NA	NA	NA	mainland	-18	7.78	4.55	1	0.28	1.26	
Scincidae	<i>Asymblepharus sikimensis</i>	no	NA	NA	NA	mainland	27	2.08	5.00	1	0.10	0.50	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Bassiana duperreyi</i>	no	NA	NA	NA	mainland	-37	5.55	5.30	1	0.20	1.06	Cogger 2000, Shine and Greer 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Radder and Shine 2007, Shine and Thomas 2005, Wilson and Swan 2008, James and Shine 1988, Radder et al. 2008, Uller et al. 2009, Fischer and Lindenmayer 2005, Kohler 2005, Wilson and Swan 2011, Michael and Lindenmayer 2010, Greer 1982, Swanson 2007
Scincidae	<i>Bassiana trilineata</i>	no	NA	NA	NA	mainland	-33	5.17	5.15	1	0.31	1.62	Cogger 2000, Shine and Greer 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Shine 1983, Wilson and Swan 2008, James and Shine 1988, Taylor 1985, Kohler 2005, Bush et al. 2010, Wilson and Swan 2011, Heatwole and Taylor 1987, Greer 1982, Bush et al. 2007
Scincidae	<i>Brachymeles cebuensis</i>	yes	Philippine Islands	Cebu	4467.50	Oceanic	10	1.66	2.00	NA	NA	NA	Brown and Alcala 1980, Alcala 1986
Scincidae	<i>Brachymeles minimus</i>	yes	Philippine Islands	Catanduanes	1522.90	Land bridge	14	0.88	NA	NA	0.27	NA	Brown and Alcala 1995, Siler et al. 2010
Scincidae	<i>Brachymeles taylori</i>	yes	Philippine Islands	Negros	13074.50	Oceanic	10	12.21	2.50	2.5	NA	NA	Siler and Brown 2010, Gaulke 2011
Scincidae	<i>Brachymeles tridactylus</i>	yes	Philippine Islands	Negros	13074.50	Oceanic	12	1.09	2.00	NA	NA	NA	Brown and Alcala 1980, Brown 1956, Shine and Wall 2008, Gaulke 2011
Scincidae	<i>Brasiliscincus agilis</i>	no	NA	NA	NA	mainland	-20	6.33	4.45	1	0.54	2.42	Huang 2006b, Vrcibradic and Rocha 1996, Rocha et al. 2004, Rocha et al. 2002, Vrcibradic & Rocha 1996, Miralles et al. 2009, Sinervo et al. 2010, Ramírez-Pinilla et al. 2002, Rocha et al. 2009, Vrcibradic and Rocha 2011,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Brasiliscincus heathi</i>	no	NA	NA	NA	mainland	-8	7.84	5.02	1.5	0.53	3.97	Fitch 1985, Cox et al. 2003, Pianka and Vitt 2003, Mesquita et al. 2006b, Cree 1994, Dunham et al. 1988, Rodrigues 1996, Vitt and Price 1982, Warne and Charnov 2008, Blackburn and Vitt 1992, Vitt 1995, Schmidt and Inger 1951, Blackburn et al. 1984, Vitt 1986, Fitch 1982, Vanzolini et al. 1980, Sinervo et al. 2010, Ramirez-Pinilla et al. 2002, Rocha and Rodrigues 2005, Cooper and Whiting 2000, Vrcibradic and Rocha 2011
Scincidae	<i>Caledoniscincus aquilonius</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	2.15	3.00	NA	NA	NA	Bauer and Sadlier 2000
Scincidae	<i>Caledoniscincus atropunctatus</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	2.77	2.00	NA	NA	NA	Bauer and Sadlier 2000, Bauer and Vindum 1990, Ineich 2011, Bauer and Sadlier 1994
Scincidae	<i>Caledoniscincus auratus</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	2.45	3.00	NA	NA	NA	Bauer and Sadlier 2000
Scincidae	<i>Caledoniscincus austrocaledonicus</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	1.88	3.50	NA	0.19	NA	Bauer and DeVaney 1987, Bauer and Sadlier 2000
Scincidae	<i>Caledoniscincus chazeaui</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	1.41	3.00	NA	NA	NA	Bauer and Sadlier 2000
Scincidae	<i>Caledoniscincus festivus</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	5.49	5.00	NA	NA	NA	Bauer and Sadlier 2000, Shea et al. 2009
Scincidae	<i>Caledoniscincus haplorhinus</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	3.12	3.50	NA	NA	NA	Bauer and Sadlier 2000
Scincidae	<i>Caledoniscincus orestes</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	5.36	6.50	NA	NA	NA	Bauer and Sadlier 2000, Shea et al. 2009
Scincidae	<i>Caledoniscincus renevieri</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	2.45	3.00	NA	NA	NA	Bauer and Sadlier 2000
Scincidae	<i>Caledoniscincus terma</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-20	2.30	2.50	NA	NA	NA	Bauer and Sadlier 2000

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Carlia longipes</i>	no	NA	NA	NA	mainland	-14	3.81	2.00	2	0.14	0.56	Cogger 2000, Tinkle et al. 1970, Wilson and Swan 2003, Allison 2006, Greer 2005, Ingram and Covacevich 1989, Cree 1994, Wilson and Swan 2008, Goodman and Isaac 2008, Goodman 2007, Todd 2008, Goodman et al. 2009, Horner 1999, Wilhoft and Reiter 1965, Heatwole and Taylor 1987
Scincidae	<i>Carlia rhomboidalis</i>	no	NA	NA	NA	mainland	-19	2.19	2.00	2.5	0.10	0.50	Fitch 1970, Cogger 2000, Tinkle et al. 1970, Wilson and Swan 2003, Greer 1989, Ingram and Covacevich 1989, Cree 1994, Wilson and Swan 2008, James and Shine 1988, Radder et al. 2008, Goodman 2007, Taylor 1985, Goodman et al. 2009, Sinervo et al. 2010, Kohler 2005, Heatwole and Taylor 1987, Swanson 2007, Wilhoft 1961, Brattstrom 1965
Scincidae	<i>Carlia rostralis</i>	no	NA	NA	NA	mainland	-18	3.93	2.00	3.5	0.40	2.77	Cogger 2000, Wilson and Swan 2003, Ingram and Covacevich 1989, Goodman and Isaac 2008, Goodman 2007, Morelli 2000, Goodman et al. 2009, Manicom and Schwarzkopf 2010, Kutt et al. 2011
Scincidae	<i>Carlia rubrigularis</i>	no	NA	NA	NA	mainland	-18	2.76	2.00	1	0.16	0.32	Cogger 2000, Wilson and Swan 2003, Goodman 2006, Ingram and Covacevich 1989, Goodman 2006b, Goodman and Isaac 2008, Goodman 2007, Goodman et al. 2009, Manicom and Schwarzkopf 2010, Swanson 2007, Andersson et al. 2010, Kutt et

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source al. 2011
Scincidae	<i>Carlia tetradactyla</i>	no	NA	NA	NA	mainland	-32	2.64	2.00	2	0.24	0.98	Shine and Greer 1991, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Ingram and Covacevich 1989, Wilson and Swan 2008, James and Shine 1988, Fischer and Lindenmayer 2005, Kohler 2005, Michael et al. 2010, Michael and Lindenmayer 2010, Michael et al. 2011 Bauer and Sadlier 2000, Bauer and Vindum 1990, Sadlier et al. 2006, Bauer and Sadlier 1994, Whittaker et al. 2004
Scincidae	<i>Celatiscincus euryotis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-23	1.12	2.50	NA	NA	NA	Sadlier et al. 2006
Scincidae	<i>Celatiscincus similis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	1.69	2.00	NA	NA	NA	Arnold and Oviden 2004, Rogner 1997b, Pollo 2008, Greer et al. 1998, Galan 2003, Ibarquengoytia and Casalins 2007, Kwet 2009, Malkmus 2004, Hailey et al. 1987, Bogaerts 2006, Hailey and Elliot 1995, Maso and Pijoan 2011
Scincidae	<i>Chalcides bedriagai</i>	no	NA	NA	NA	mainland	39	8.63	2.44	1	0.26	0.64	Greer 2001, Scleich et al. 1996, Arnold and Oviden 2004, Street 1979, Rogner 1997b, Sindaco et al. 2006, Corti and Cascio 2002, Carranza et al. 2008, Greer et al. 1998, Kwet 2009, Shine and Wall 2008, Sindaco et al.
Scincidae	<i>Chalcides chalcides</i>	no	NA	NA	NA	mainland	39	12.92	7.00	1	0.76	5.31	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2010, Busack 1986, Guarino 2010
Scincidae	<i>Chalcides coeruleopunctatus</i>	yes	Canary Islands	Gomera	359.10	Oceanic	28	13.73	3.00	1	1.05	3.15	Carranza et al. 2008, Salvador 2008, Maso and Pijoan 2011 Fitch 1970, Scleich et al. 1996, Amitai and Bouskila 2001, Spawls et al. 2002, Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Minton 1966, Disi et al. 2001, Baran and Atatur 1998, Flower 1933, Parker 1932, Arnold 1984, El Din 2006, Rogner 1997b, Arnold 1980, Schatti and Desvoignes 1999, Le Berre 1989, Khan 2006, Bons and Geniez 1996, Atatur and Gocmen 2001, Sindaco et al. 2006, Corti and Cascio 2002, Jongbloed 2000,
Scincidae	<i>Chalcides ocellatus</i>	no	NA	NA	NA	mainland	29	50.68	7.00	1.5	1.12	11.73	Andrews and Pough 1980, Valakos et al. 2008, Gans 1965, Arnold 1986, Hornby 1996, Daut and Andrews 1993, Kwet 2009, Somaweera and Somaweera 2009, Sinervo et al. 2010, Baier et al. 2009, Largen and Spawls 2010, Carretero et al. 2010, Sindaco et al. 2010, Capula and Luiselli 1994, van der Kooij 2001, Cascio 2010, Bogaerts 2006, Hailey and Elliot 1995, Bar and Haimovitch 2012, Trape et al. 2012, Kalboussi and Nouira 2004

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Chalcides sepsoides</i>	no	NA	NA	NA	mainland	29	6.36	6.00	1	0.53	3.19	Schleich et al. 1996, Amitai and Bouskila 2001, Disi et al. 2001, Flower 1933, El Din 2006, Rogner 1997b, Le Berre 1989, Andrews and Pough 1980, Werner 1968, Carranza et al. 2008, Werner 1987, Bar and Haimovitch 2012, Trape et al. 2012
Scincidae	<i>Chalcides sexlineatus</i>	yes	Canary Islands	Gran Canaria	1529.90	Oceanic	28	7.09	4.50	NA	0.86	NA	Arnold and Ovenden 2004, Rogner 1997b, Nagy et al. 1999, Salvador and Brown 2007, Brown and Nagy 2007, Bogaerts 2006, Maso and Pijoan 2011
Scincidae	<i>Chalcides viridanus</i>	yes	Canary Islands	Tenerife	2007.80	Oceanic	28	26.45	4.00	1	0.79	3.17	Arnold and Ovenden 2004, Salvador 2007, Salvador 2008, Greer et al. 1998, Bogaerts 2006, Maso and Pijoan 2011
Scincidae	<i>Chioninia coctei</i>	yes	Cape Verde	Santa Luzia	36.70	Oceanic	17	443.03	2.00	NA	NA	NA	Greer 2001, Carranza et al. 2001, Case et al. 1998, Andreone and Guarino 2003
Scincidae	<i>Chioninia spinalis</i>	yes	Cape Verde	Sao Tiago	991.00	Oceanic	16	5.40	NA	NA	0.49	NA	Joger 1993
Scincidae	<i>Chioninia vaillantii</i>	yes	Cape Verde	Sao Tiago	991.00	Oceanic	15	29.63	NA	NA	0.54	NA	Carranza et al. 2001
Scincidae	<i>Coeranoscincus reticulatus</i>	no	NA	NA	NA	mainland	-28	15.87	3.85	1	1.43	5.51	Greer 2001, Wilson and Swan 2003, Greer 1989, Greer 2005, Wilson and Swan 2008, Kohler 2005, Wilson and Swan 2011, Hutchinson 1993, Swanson 2007

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Copeoglossum nigropunctatum</i>	no	NA	NA	NA	mainland	-7	13.30	4.70	1	0.95	4.45	Duellman and Mendelson 1995, Avila-Pires 1995, Pianka and Vitt 2003, Huang 2006b, Mesquita et al. 2006b, Gainsbury and Colli 2003, Nogueira et al. 2005, Vitt 2000, Molina et al. 2004, Vitt and Zani 1998, Duellman 2005, Bartlett and Bartlett 2003, Colli et al. 2002, Lotzkat 2007, Vitt and Zani 1996b, Vitt et al. 1997, Vitt et al. 1999, Huey et al. 2001, Fuenmayor et al. 2005, Sinervo et al. 2010, Avila-Pires et al. 2010, Ugueto and Rivas 2010, Cooper and Whiting 2000, Rocha et al. 2009, Vrcibradic and Rocha 2011, Winck and Rocha 2012
Scincidae	<i>Corucia zebrata</i>	yes	Solomon Islands	Bougainville	8590.10	Oceanic	-8	747.52	1.00	0.75	78.02	58.52	Greer 2001, Cooper and Vitt 2002, Pianka and Vitt 2003, Badger 2003, Rogner 1997b, McCoy 1980, Mann and Meek 2004, Sinervo et al. 2010, Hagen and Bull 2011, Hagen 2011, Parker 1983
Scincidae	<i>Cryptoblepharus egeriae</i>	yes	None	Christmas	135.00	Oceanic	-10	1.58	2.00	NA	NA	NA	Cogger et al. 1983, Cogger 2000, Smith et al. 2012
Scincidae	<i>Cryptoblepharus eximius</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-18	0.79	1.00	NA	0.12	NA	Zug 1991, Horner 2007, Morrison 2003
Scincidae	<i>Cryptoblepharus novocaledonicus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	1.09	1.50	NA	NA	NA	Bauer and Sadlier 2000, Bauer and Vindum 1990, Horner 2007, Whittaker et al. 2004
Scincidae	<i>Cryptoblepharus poecilopleurus</i>	yes	Bismarck Archipelago	New Britain	35144.60	Oceanic	-7	1.22	2.00	NA	0.12	NA	Rodda et al. 1991, McCoy 1980, Towns 1994, McCoy 2006, Horner 2007, Iburguengoytia 2008, McCoid 1994, Goldberg and Kraus 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Cryptoblepharus renschi</i>	yes	Lesser Sunda Islands	Flores	14154.40	Oceanic	-9	1.03	NA	NA	0.07	NA	Horner 2007, Das 2010
Scincidae	<i>Ctenotus brooksi</i>	no	NA	NA	NA	mainland	-27	1.74	2.35	2	0.21	1.01	Huey and Pianka 1981, Cogger 2000, Cox et al. 2003, Pianka 1986, Wilson and Swan 2003, Greer 2005, Wilson and Swan 2008, Huey et al. 2001, Todd 2008, Horner 1999, James 1991, Sinervo et al. 2010, Heatwole and Taylor 1987, Swan and Watharow 2005
Scincidae	<i>Ctenotus lanceolini</i>	yes	None	Lancelin	0.09	Land bridge	-31	10.06	NA	1	NA	NA	Cogger 2000, Wilson and Swan 2003, Greer 2005
Scincidae	<i>Ctenotus pantherinus</i>	no	NA	NA	NA	mainland	-24	10.06	5.10	2	0.66	6.74	Huey and Pianka 1981, Cogger 2000, Cox et al. 2003, Pianka 1986, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Vitt and Price 1982, Huey et al. 2001, Chapman and Dell 1985, Horner 1999, Sinervo et al. 2010, Kohler 2005, Bush et al. 2010, Huey and Pianka 2007, James 1991, (Oecologia), Heatwole and Taylor 1987, Swanson 2007, Goodyear and Pianka 2011, Gordon et al. 2010, Moro and MacAulay 2010, Heatwole and Butler 1981
Scincidae	<i>Ctenotus robustus</i>	no	NA	NA	NA	mainland	-25	20.57	5.75	1	0.73	4.17	Cogger 2000, Brown 1991, Wilson and Swan 2003, Allison 2006, Greer 2005, James and Shine 1988, Bjursell 2001, Goodman 2006, Taylor 1985, Fischer and Lindenmayer 2005, Sinervo et al. 2010, Michael et al. 2010, Heatwole and Taylor 1987, Swanson 2007,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Michael et al. 2011, Taylor 2004, Swan and Watharow 2005
Scincidae	<i>Ctenotus taeniolatus</i>	no	NA	NA	NA	mainland	-26	6.54	4.15	1	0.44	1.83	Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Cree 1994, Dunham et al. 1988, Huey and Bennett 1987, Warne and Charnov 2008, James and Shine 1988, Goodman 2006, Fischer and Lindenmayer 2005, Sinervo et al. 2010, Kohler 2005, Michael and Lindenmayer 2010, Heatwole and Pianka 1993, Hutchinson 1993, Heatwole and Taylor 1987, Swanson 2007, Kutt et al. 2011, Taylor 2004, Greer 2001, Cogger 2000, Fitch 1970, Shine and Greer 1991, Wilson and Swan 2003, Rogner 1997b, Greer 1989, Greer 2005, Shea 1995, Heatwole and Taylor 1987, Bennett and John-Alder, 1986
Scincidae	<i>Cyclodomorphus casuarinae</i>	no	NA	NA	NA	mainland	-42	42.73	5.80	1	1.44	8.34	Cogger 2000, Wilson and Swan 2003, Greer 2005, Shea 1995, Heatwole and Taylor 1987, Bennett and John-Alder, 1986
Scincidae	<i>Cyclodomorphus celatus</i>	no	NA	NA	NA	mainland	-27	21.74	3.80	1	1.12	4.24	Cogger 2000, Wilson and Swan 2003, Greer 2005, Shea and Miller 1995, Bush et al. 2010
Scincidae	<i>Cyclodomorphus melanops</i>	no	NA	NA	NA	mainland	-25	18.34	2.45	1	1.16	2.85	Cogger 2000, Wilson and Swan 2003, Greer 2005, Shea and Miller 1995, Pianka 2011
Scincidae	<i>Cyclodomorphus michaeli</i>	no	NA	NA	NA	mainland	-32	44.54	7.65	1	1.38	10.59	Greer 2001, Cogger 2000, Wilson and Swan 2003, Greer 2005, Shea 1995, Swanson 2007

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Dasia olivacea</i>	no	NA	NA	NA	mainland	3	27.40	6.00	1.5	0.73	6.53	Smith 1935, Tikader and Sharma 1992, Taylor 1963, Das 2004, Manthey and Grossmann 1997, de Rooij 1915, Rogner 1997b, Cox et al. 1998, Inger and Colwell 1977, Malkmus et al. 2002, Inger and Lian 1996, Stuart and Emmett 2006, Das 2002, Onn et al. 2010, Teynie et al. 2010, Das 2010, Cox et al. 2010, Das 2011, Neang et al. 2010, Grismer 2011, Grismer 2011b
Scincidae	<i>Egernia cunninghami</i>	no	NA	NA	NA	mainland	-34	171.04	4.40	1	4.97	21.86	Fitch 1970, Chapple 2003, Cogger 2000, Brown 1991, Cooper and Vitt 2002, Pianka and Vitt 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, Greer 1989, Greer 2005, Duffield and Bull 1998, Wilson and Swan 2008, Arena and Wooler 2003, Taylor 1985, Sinervo et al. 2010, Michael et al. 2010, Michael and Lindenmayer 2010, Heatwole and Pianka 1993, Hutchinson 1993, Heatwole and Taylor 1987, Swanson 2007, Michael et al. 2011, Light et al. 1966
Scincidae	<i>Egernia hosmeri</i>	no	NA	NA	NA	mainland	-19	143.64	2.00	1.5	6.65	19.95	Greer 2001, Chapple 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, Greer 2005, Wilson and Swan 2008, Horner 1999, Swanson 2007, Post 2000
Scincidae	<i>Egernia kingii</i>	no	NA	NA	NA	mainland	-32	254.94	5.00	1	6.80	34.02	Greer 2001, Chapple 2003, Arena and Wooler 2003, Wilson and Swan 2003, Greer 1989, Greer 2005, Wilson and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Swan 2008, Warne and Charnov 2008, Duffield and Bull 1996, Bush et al. 2010
Scincidae	<i>Egernia stokesii</i>	no	NA	NA	NA	mainland	-28	149.52	4.40	0.875	8.69	33.45	Greer 2001, Chapple 2003, Cogger 2000, Storr 1978, Cooper and Vitt 2002, Pianka and Vitt 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, Greer 2005, Duffield and Bull 1998, Wilson and Swan 2008, Warne and Charnov 2008, Chapple 2005, Duffield and Bull 1996, Sinervo et al. 2010, Hutchinson 1993, Heatwole and Taylor 1987, Swanson 2007, Light et al. 1966
Scincidae	<i>Egernia striolata</i>	no	NA	NA	NA	mainland	-28	26.78	3.80	1	2.37	9.01	Chapple 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997b, Greer 1989, Greer 2005, Chapple 2005, Fischer and Lindenmayer 2005, Henle 1989c, Bustard 1968, Bull and Bonnett 2004, Milton 1987, Sinervo et al. 2010, Michael et al. 2010, Michael and Lindenmayer 2010, Wilson 2003, Hutchinson 1993, Turner 1977, Heatwole and Taylor 1987, Swanson 2007, Michael et al. 2011, Bustard 1968
Scincidae	<i>Emoia adspersa</i>	yes	Samoa Islands	Savai'i	1717.60	Oceanic	-14	8.47	1.90	NA	NA	NA	Brown 1991, Dunham et al. 1988, Schwaner 1980
Scincidae	<i>Emoia aneityumensis</i>	yes	New Hebrides	Aneityum	159.20	Oceanic	-20	10.40	4.50	NA	NA	NA	Brown 1991, Hamilton et al. 2008

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Emoia atrocostata</i>	no	NA	NA	NA	mainland	-6	11.70	1.95	3	0.66	3.87	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Cogger 2000, Greer 1968, Goris and Maeda 2004, Zweifel 1980, Cox et al. 2003, Grismer et al. 2006, Pianka and Vitt 2003, Das 2004, Wilson and Swan 2003, Allison 2006, Crombie and Pregill 1999, Brown and Alcala 1980, Manthey and Grossmann 1997, Rodda et al. 1991, Greer 1989, McCoy 1980, Cox et al. 1998, Heatwole 1975, Brown 1991, Taylor 1922, Cogger et al. 1983, Cree 1994, McCoy 2006, Dunham et al. 1988, Youmans and Grismer 2006, Wilson and Swan 2008, Hamilton et al. 2008, Warne and Charnov 2008, Alcala and Brown 1967, Kohler 2005, Das 2010, Cox et al. 2010, Alcala 1986, Das 2011, Oliveros et al. 2011, Grismer 2011, Koch et al. 2010, Auffenberg and Auffenberg 1989, Grismer 2011b, Ineich 2011, McCoid 1994, Huang 2011, Gaulke 2011
Scincidae	<i>Emoia campbelli</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-18	9.06	2.00	NA	NA	NA	Brown 1991, Zug 1991, Morrison 2003, Brown and Gibbons 1986
Scincidae	<i>Emoia concolor</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-18	5.91	2.00	NA	0.32	NA	Brown 1991, Zug 1991, Hamilton et al. 2008, Morrison 2003, Brown and Gibbons 1986

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Emoia impar</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-5	1.59	NA	NA	0.17	NA	Crombie and Pregill 1999, Bruna et al. 1996, Buden 2000, Zug 1991, Buden 2007, Buden 1998, Zug 1991, Morrison 2003, McElroy 2007, Van Wilgen and Richardson 2012, Buden 1999
Scincidae	<i>Emoia isolata</i>	yes	Solomon Islands	Bellona	19.74	Oceanic	-11	2.30	2.00	NA	NA	NA	Brown 1991, McCoy 2006
Scincidae	<i>Emoia loyaltiensis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	5.77	3.50	NA	NA	NA	Bauer and Sadlier 2000, Brown 1991
Scincidae	<i>Emoia maculata</i>	yes	Solomon Islands	Makira	3089.80	Oceanic	-11	1.89	2.00	NA	NA	NA	McCoy 1980, Brown 1991, McCoy 2006
Scincidae	<i>Emoia nativittatis</i>	yes	None	Christmas	135.00	Oceanic	-10	6.33	2.00	NA	NA	NA	Fitch 1970, Greer 1968, Brown 1991, Cogger et al. 1983, Cogger 2000, Kohler 2005
Scincidae	<i>Emoia nigromarginata</i>	yes	New Hebrides	Espiritu Santo	3955.50	Oceanic	-16	4.84	2.50	NA	NA	NA	Brown 1991, Hamilton et al. 2008, Ineich 2011
Scincidae	<i>Emoia parkeri</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-18	1.97	2.00	NA	0.23	NA	Brown 1991, Hamilton et al. 2008, Morrison 2003
Scincidae	<i>Emoia ponapea</i>	yes	Caroline Islands	Pohnpei	334.00	Oceanic	7	1.85	2.00	NA	0.14	NA	Brown 1991, Buden 2000, Kiester 1982
Scincidae	<i>Emoia reimschiesseli</i>	yes	Maluku Islands	Halmahera	18039.60	Oceanic	-1	3.79	2.00	NA	NA	NA	Brown 1991
Scincidae	<i>Emoia sanfordi</i>	yes	New Hebrides	Espiritu Santo	3955.50	Oceanic	-16	21.42	5.00	NA	NA	NA	McCoy 1980, McCoy 2006, Hamilton et al. 2008, Hamilton et al. 2007
Scincidae	<i>Emoia schmidtii</i>	yes	Solomon Islands	New Georgia	2043.50	Oceanic	-8	3.09	2.00	NA	NA	NA	Brown 1991, McCoy 2006
Scincidae	<i>Emoia slevini</i>	yes	Mariana Islands	Guam	541.00	Oceanic	15	6.51	NA	NA	0.54	NA	Brown 1991, McCoid 1994
Scincidae	<i>Emoia sorex</i>	yes	Maluku Islands	Halmahera	18039.60	Oceanic	0	3.12	2.00	NA	NA	NA	Fitch 1970, Greer 1968, Brown 1991, Kohler 2005, Setiadi and Hamidy 2006
Scincidae	<i>Emoia taumakoensis</i>	yes	Solomon Islands	Taumako	10.00	Oceanic	-9	2.15	NA	NA	0.20	NA	Brown 1991, McCoy 2006, McCoy and Webber 1984
Scincidae	<i>Emoia tongana</i>	yes	Samoa Islands	Savai'i	1717.60	Oceanic	-15	4.05	2.00	NA	NA	NA	Burt 1930, Brown 1991, Hamilton et al. 2008, Gill 1993

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Emoia trossula</i>	yes	Fiji Islands	Viti Levu	10531.00	Oceanic	-17	15.13	3.50	NA	0.60	NA	Brown 1991, Bruna et al. 1996, Zug 1991, Hamilton et al. 2008, Morrison 2003, Brown and Gibbons 1986, Crombie and Steadman 1986, Clobert et al. 1998, Cogger 2000, Pianka 1986, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Wilson and Swan 2008, James and Losos 1991, Henle 1989, Horner 1999, Kohler 2005, Bush et al. 2010, Wilson 2003, Swanson 2007, Pianka 2011, Bush et al. 2007
Scincidae	<i>Eremiascincus richardsonii</i>	no	NA	NA	NA	mainland	-26	12.64	4.75	2.5	0.54	6.45	Shine and Greer 1991, Cogger 2001, Wilson and Swan 2003, Greer 2005, Goodman and Isaac 2008, Goodman 2007, Greer 1992, Swanson 2007, Schwarzkopf et al. 2010, Kutt et al. 2011
Scincidae	<i>Eulamprus brachyosoma</i>	no	NA	NA	NA	mainland	-21	6.93	3.90	1	0.58	2.25	Shine and Greer 1991, Cogger 2000, Fitch 1970, Pianka and Vitt 2003, Wilson and Swan 2003, Greer 1989, Greer 2005, Bauer and Jackman 2008, Huang and Tu 2008, Clusella-Trullas et al. 2008, Huey and Bennett 1987, Wilson and Swan 2008, Uller et al. 2009, Goodman and Isaac 2008, Goodman 2007, Watt 2002, Veron 1969, Heatwole and Veron 1977, Taylor 1985, Goodman et al. 2009, Salked 2004, McElroy et al. 2008, Sinervo et al. 2010, Heatwole and Pianka 1993, Heatwole and Taylor 1987, Swanson 2007,
Scincidae	<i>Eulamprus quoyii</i>	no	NA	NA	NA	mainland	-29	23.46	5.35	1	1.00	5.33	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Bennett and John-Alder 1986, Veron and Heatwole 1970
Scincidae	<i>Eulamprus tympanum</i>	no	NA	NA	NA	mainland	-37	14.30	3.28	0.75	0.76	1.87	Cogger 2000, Brown 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Cree 1994, Bauer and Jackman 2008, Huang and Tu 2008, Shine 1983, Wilson and Swan 2008, Warne and Charnov 2008, Uller et al. 2009, Rohr 1997, Taylor 1985, Fischer and Lindenmayer 2005, Sinervo et al. 2010, Cree and Hare 2010, Schwarzkopf 1992, Michael and Lindenmayer 2010, Hutchinson 1993, Schwarzkopf 1993, Heatwole and Taylor 1987, Greer 2001, Schleich et al. 1996, Amitai and Bouskila 2001, Szczerbak 2003, Anderson 1999, Disi et al. 2001, Baran and Atatur 1998, Flower 1933, El Din 2006, Rogner 1997b, Reed and Marx 1959, Le Berre 1989, Khan 2006, Atatur and Gocmen 2001, Ahmadzadeh et al. 2008, Cooper et al. 2000, McElroy et al. 2008, Kohler 2005, Baier et al. 2009, Arakelyan et al. 2011, Bar and Haimovitch 2012,
Scincidae	<i>Eumeces schneideri</i>	no	NA	NA	NA	mainland	33	69.70	11.00	1	3.71	40.79	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Fathinia et al. 2009
Scincidae	<i>Eutropis carinata</i>	no	NA	NA	NA	mainland	22	14.51	5.25	2	0.44	4.63	Inger and Greenberg 1966, Smith 1935, Schleich and Kastle 2002, Daniel 1983, Tikader and Sharma 1992, Rogner 1997b, Deraniyagala 1953, Inger et al. 1984, Murthy 1995, Shrestha 2001, Radder et al. 2008, Somaweera and Somaweera 2009, Das 2002, Das and de Silva 2011, Aryal et al. 2010, Chandramouli and Ganesh 2011, Ahmed 2009 Taylor 1963, Huang 2006a, 2006b, Manthey and Grossmann 1997, Ziegler 2002, Cox et al. 1998, Karsen et al. 1986, Inger and Colwell 1977, Grismer et al. 2008, Radder et al. 2008, Huang 2007, Huang 2007b, Taylor and Elbel 1958, Das 2010, Cox et al. 2010, Grismer 2011, Huang 2010, Grismer 2011b, Pauwels et al. 2009, Teynie and David 2010
Scincidae	<i>Eutropis longicaudata</i>	no	NA	NA	NA	mainland	24	27.18	6.85	1.5	1.63	16.79	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Eutropis multifasciata</i>	no	NA	NA	NA	mainland	16	19.81	5.05	2.5	0.95	12.00	Tinkle et al. 1970, Fitch 1970, Inger and Greenberg 1966, Smith 1935, Hendrickson 1966, Tikader and Sharma 1992, Taylor 1963, Grismer et al. 2006, Smedley 1931, Das 2004, Anderson 1889, Teynie 2004, Brown and Alcala 1961, Brown and Alcala 1980, Manthey and Grossmann 1997, Rogner 1997b, Ziegler 2002, Cox et al. 1998, Taylor 1922, Inger and Colwell 1977, Malkmus et al. 2002, Pauwels et al. 2003, Grismer 2007, Grismer et al. 2008, Grismer et al. 2007, Somma and Brooks 1976, Ji et al. 2006, Onn et al. 2009, Das 2002, Sinervo et al. 2010, Taylor and Elbel 1958, Onn et al. 2010, Das 2010, Cree and Hare 2010, Cox et al. 2010, Alcala 1986, Das 2011, Ahmed et al. 2009, Grismer 2011, Setiadi and Hamidy 2006, Auffenberg and Auffenberg 1989, Auffenberg and Auffenberg 1988, Grismer 2011b, Auffenberg 1980, Zhang and Ji 2004, Inger 1959, Alcala 1966, Teynie and David 2010, Van Wilgen and Richardson 2012, Brattstrom 1965, Gaulke 2011, Du et al. 2012
Scincidae	<i>Gongylomorphus bojerii</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-20	2.99	2.00	NA	0.44	NA	Vinson and Vinson 1969, Bullock et al. 1985, Case et al. 1998, Ross et al. 2008, Bullock 1986

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Hemiergis peronii</i>	no	NA	NA	NA	mainland	-33	1.71	2.68	1	0.20	0.52	Fitch 1970, Shine and Greer 1991, Cogger 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Cree 1994, Huey and Bennett 1987, Taylor 1985, Sinervo et al. 2010, Bush 1992, Heatwole and Taylor 1987, Bennett and John-Alder 1986 Fitch 1970, Cogger 2000, Greer 2001, Pianka and Vitt 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, Greer 2005, Wilson and Swan 2008, Koetz 2000, Heatwole and Pianka 1993, Hutchinson 1993, Swanson 2007, Hauschild 1988, Jones 1987 Brown and Alcala 1961, Brown and Alcala 1980, Sinervo et al. 2010, Alcala 1986, Turner 1977, Alcala and Brown 1966, Gaulke 2011
Scincidae	<i>Hemisphaeriodon gerrardii</i>	no	NA	NA	NA	mainland	-26	186.00	21.65	1	2.94	63.73	Linkem et al. 2010
Scincidae	<i>Insulasaurus arborens</i>	yes	Philippine Islands	Panay	12011.10	Oceanic	11	3.11	1.50	NA	0.20	NA	Cheke 1984, Bowler 2006, Rocha et al. 2010, Gerlach 2008, Evans and Evans 1980 Sadlier et al. 2004, Sadlier et al. 2009, Whittaker et al. 2004
Scincidae	<i>Insulasaurus traanorum</i>	yes	Philippine Islands	Palawan	12188.60	Continental	9	2.36	1.00	NA	NA	NA	Shine and Greer 1991, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Cree 1994, Towns 1994, Dunham et al. 1988, Shine 1983, James and Shine 1988, Greer 1997, Burgin 1992, Fischer and Lindenmayer 2005, Goodman
Scincidae	<i>Janetaescincus braueri</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	1.11	2.00	NA	NA	NA	
Scincidae	<i>Kanakysaurus viviparus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-20	7.29	3.00	NA	0.73	NA	
Scincidae	<i>Lampropholis delicata</i>	no	NA	NA	NA	mainland	-29	1.12	3.35	1	0.06	0.21	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													et al. 2009, Jewell 2008, Hoser 2009, Kohler 2005, Michael and Lindenmayer 2010, Heatwole and Taylor 1987, Swanson 2007, Swan and Watharow 2005
Scincidae	<i>Lampropholis guichenoti</i>	no	NA	NA	NA	mainland	-34	1.16	2.80	1.5	0.10	0.42	Fitch 1970, Shine and Greer 1991, Cogger 2000, Brown 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Shine 1983, Vitt and Price 1982, Qualls and Shine 1998, James and Shine 1988, Greer 1997, Burgin 1992, Todd 2008, Taylor 1985, Fischer and Lindenmayer 2005, Simbotwe 1985, Kohler 2005, Michael and Lindenmayer 2010, Hutchinson 1993, Swanson 2007, Michael et al. 2011
Scincidae	<i>Larutia seribuatensis</i>	yes	Seribuat Archipelago	Tioman	132.00	Land bridge	3	5.13	NA	NA	1.25	NA	Grismer et al. 2003, Grismer and Pan 2008, Das 2010, Grismer 2011, Grismer 2011b
Scincidae	<i>Leiopisma telfairii</i>	yes	Mascarene Archipelago	Mauritius	2040.00	Oceanic	-20	49.85	11.50	1	1.41	16.22	Greer 2001, Henkel and Schmidt 2000, Vinson and Vinson 1969, Kohler 2005, Bullock 1986, Pernetta et al. 2005
Scincidae	<i>Lerista borealis</i>	no	NA	NA	NA	mainland	-16	1.52	3.00	1	0.28	0.83	Cogger 2000, Wilson and Swan 2003, Greer 1989, Horner 1999, Kohler 2005, Wells 2012

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Lerista bougainvillii</i>	no	NA	NA	NA	mainland	-36	3.66	2.74	1	0.25	0.68	Shine and Greer 1991, Cogger 2000, Brown 1991, Qualls et al. 1994, Barden and Shine 1994, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Dunham et al. 1988, James and Shine 1988, Greer 1967, Sinervo et al. 2010, Kohler 2005, Michael et al. 2010, Michael and Lindenmayer 2010, Greenville and Dickman 2005, Heatwole and Taylor 1987, Swanson 2007, Michael et al. 2011, Hailey and Elliot 1995, Wells 2012, Swan and Watharow 2005
Scincidae	<i>Lerista labialis</i>	no	NA	NA	NA	mainland	-27	1.13	2.00	1.5	0.15	0.46	Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Henle 1989, Horner 1999, Kohler 2005, Schlesinger et al. 2010, Greenville and Dickman 2005, Gordon et al. 2010, Wells 2012
Scincidae	<i>Lerista punctatovittata</i>	no	NA	NA	NA	mainland	-30	4.84	2.15	2	0.53	2.29	Clobert et al. 1998, Cogger 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Henle 1989, Greer 1967, Bustard 1978, Kohler 2005, Michael and Lindenmayer 2010, Greenville and Dickman 2005, Swanson 2007, Bustard 1968, Wells 2012, Swan and Watharow 2005
Scincidae	<i>Lerista xanthura</i>	no	NA	NA	NA	mainland	-27	0.97	1.35	1.5	0.18	0.36	Clobert et al. 1998, Cogger 2001, Wilson and Swan 2003, Henle 1991, Greer 1989, Storr et al. 1999, Henle 1989, Henle 1989c, Horner 1999,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Greenville and Dickman 2005, Wells 2012
Scincidae	<i>Liopholis inornata</i>	no	NA	NA	NA	mainland	-28	7.34	2.50	2	0.63	3.15	Chapple 2003, Cogger 2000, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997b, Greer 1989, Greer 2005, Wilson and Swan 2008, Chapman and Dell 1985, Henle 1989c, Horner 1999, Abensperg-Traun and Steven 1997, Hutchinson 1993, Heatwole and Taylor 1987, Gordon et al. 2010, Swan and Watharow 2005 Greer 2001, Chapple 2003, Cogger 2000, Cooper and Vitt 2002, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Greer 1989, Greer 2005, Wilson and Swan 2008, Warne and Charnov 2008, Horner 1999, McAlpin et al. 2011, Heatwole and Taylor 1987
Scincidae	<i>Liopholis kintorei</i>	no	NA	NA	NA	mainland	-24	173.96	6.00	1	8.50	51.02	Chapple 2003, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Cree 1994, Warne and Charnov 2008, Todd 2008, Milton 1987, Swanson 2007 Pianka 1982, Chapple 2003, Cogger 2000, Pianka 1986, Pianka and Vitt 2003, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Wilson and Swan 2008, Huey et al. 2001, Horner 1999,
Scincidae	<i>Liopholis modesta</i>	no	NA	NA	NA	mainland	-29	22.41	2.70	1	1.33	3.59	Chapple 2003, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Cree 1994, Warne and Charnov 2008, Todd 2008, Milton 1987, Swanson 2007 Pianka 1982, Chapple 2003, Cogger 2000, Pianka 1986, Pianka and Vitt 2003, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Wilson and Swan 2008, Huey et al. 2001, Horner 1999,
Scincidae	<i>Liopholis striata</i>	no	NA	NA	NA	mainland	-24	13.48	2.65	1	1.21	3.21	Chapple 2003, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Wilson and Swan 2008, Huey et al. 2001, Horner 1999,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Pianka and Giles 1982, Huey and Pianka 2007, Hutchinson 1993, Swanson 2007
Scincidae	<i>Liopholis whitii</i>	no	NA	NA	NA	mainland	-34	14.17	2.90	1	1.12	3.24	Fitch 1970, Chapple 2003, Cogger 2000, Brown 1991, Wilson and Swan 2003, Rogner 1997b, Greer 1989, Greer 2005, Cree 1994, Dunham et al. 1988, Warne and Charnov 2008, Uller et al. 2009, Chapple 2005, Todd 2008, Fischer and Lindenmayer 2005, Milton 1987, Sinervo et al. 2010, Michael and Lindenmayer 2010, Heatwole and Taylor 1987, Swanson 2007, White et al. 2009
Scincidae	<i>Lioscincus greeri</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	4.36	2.00	NA	0.44	NA	Bauer and Sadlier 2000
Scincidae	<i>Lioscincus maruia</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	4.36	2.50	NA	NA	NA	Bauer and Sadlier 2000, Whittaker et al. 2004
Scincidae	<i>Lioscincus nigrofasciolum</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	31.04	8.50	NA	0.44	NA	Bauer and DeVaney 1987, Bauer and Sadlier 2000, Bauer and Vindum 1990, Shea et al. 2009, Meier 1979, Bauer and Sadlier 1994, Whittaker et al. 2004
Scincidae	<i>Lioscincus novaecaledoniae</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	6.20	4.00	NA	0.44	NA	Bauer and Sadlier 2000
Scincidae	<i>Lioscincus steindachneri</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	31.94	6.00	NA	NA	NA	Bauer and Sadlier 2000, Bauer and Jackman 2008, Shea et al. 2009
Scincidae	<i>Lipinia leptosoma</i>	yes	Palau Islands	Babeldaob	374.10	Oceanic	7	0.87	1.00	NA	NA	NA	Shine and Greer 1991, Greer 1977, Crombie and Pregill 1999, Greer 1974, Das and Austin 2007, Fitch 1982, Irschick et al. 1996

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Lipinia macrotympanum</i>	yes	Andaman and Nicobar Islands	Great Nicobar	955.70	Oceanic	12	1.63	2.00	NA	NA	NA	Smith 1935, Tikader and Sharma 1992, Das and Austin 2007, Stoliczka 1873, Das 2002, Das 1997
Scincidae	<i>Lipinia rabori</i>	yes	Philippine Islands	Negros	13074.50	Oceanic	9	2.22	1.50	NA	NA	NA	Brown and Alcala 1961, Brown and Alcala 1980, Alcala 1986, Gaulke 2011
Scincidae	<i>Lipinia rouxi</i>	yes	Bismarck Archipelago	New Ireland	7404.50	Oceanic	-4	0.83	1.00	NA	NA	NA	Shine and Greer 1991, Uetz 2006, Das and Austin 2007, Mys 1988, Shea 2007
Scincidae	<i>Lissolepis coventryi</i>	no	NA	NA	NA	mainland	-37	22.11	2.75	1	0.83	2.29	Chapple 2003, Storr 1978, Cogger 2000, Clemann et al. 2004, Wilson and Swan 2003, Greer 1989, Greer 2005, Shine 1983, Wilson and Swan 2008, Warne and Charnov 2008, Chapple 2005, Hutchinson 1993
Scincidae	<i>Mabuya mabouya</i>	no	NA	NA	NA	mainland	5	13.08	4.51	1	0.69	3.12	Fitch 1970, 1973, 1985, Duellman 1978, Beebe 1945, Huang 2006b, Rodda et al. 2001, Schwartz and Henderson 1991, Hoogmoed 1973, Dunn 1935, Dunn 1935, Dunham et al. 1988, Blackburn and Vitt 1992, Radder et al. 2008, Breuil 2002, Somma and Brooks 1976, Test et al. 1966, Henderson and Powell 2009, Sinervo et al. 2010, Daltry 2009, Duellman 1987, Gasc 1990, Fitch 1968, Hedges and Conn 2012, Bullock and Evans 1990

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Marisora unimarginata</i>	no	NA	NA	NA	mainland	15	6.93	4.00	1.5	0.54	3.26	Fitch 1970, 1973, 1985, Campbell 1999, Stafford and Meyer 2000, Savage 2002, Lee 2000, Kohler 2003, Rand and Myers 1990, Duellman 1990, Kohler 1996, Mccranie et al. 2005, Kohler et al. 2006, Guyer and Donnelly 2005, Cree 1994, Duellman 1961, Blackburn and Vitt 1992, Vitt and Zani 1998, Radder et al. 2008, Kohler 2008, Garcia-Vazquez and Feria-Ortiz 2006, Davis and Dixon 1961, Lopez and Gonzalez 1997, Duellman 1965, Duellman 1965, Goldberg 2009, Cooper and Whiting 2000, Leenders and Watkins-Colwell 2004
Scincidae	<i>Marmorosphax boulinda</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	2.77	2.00	NA	NA	NA	Sadlier et al. 2009
Scincidae	<i>Marmorosphax taom</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	2.94	2.00	NA	NA	NA	Sadlier et al. 2009
Scincidae	<i>Marmorosphax tricolor</i>	yes	New Caledonia	New Caledonia	18575 .00	Continental	-21	3.03	5.00	NA	0.16	NA	Bauer and DeVaney 1987, Bauer and Sadlier 2000, Bauer and Vindum 1990, Sadlier et al. 2009, Whittaker et al. 2004
Scincidae	<i>Menetia greyii</i>	no	NA	NA	NA	mainland	-25	0.54	1.55	1.5	0.04	0.09	Greer 2001, Cogger 2000, Cox et al. 2003, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Greer 1974, Cree 1994, Dunham et al. 1988, Fitch 1982, Warne and Charnov 2008, James and Shine 1988, Huey et al. 2001, Chapman and Dell 1985, Storr et al. 1999, Taylor 1985, Henle

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Morethia boulengeri</i>	no	NA	NA	NA	mainland	-25	1.92	3.00	2	0.09	0.56	1989c, Horner 1999, Kohler 2005, Bush et al. 2010, Bush 1992, Michael and Lindenmayer 2010, Heatwole and Taylor 1987, Michael et al. 2011, Davidge 1979, Sadlier 1990, Gordon et al. 2010, Swan and Watharow 2005 Clobert et al. 1998, Shine and Greer 1991, Cogger 2000, Brown 1991, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, Dunham et al. 1988, Warne and Charnov 2008, James and Shine 1988, Taylor 1985, Fischer and Lindenmayer 2005, Henle 1990, Henle 1989b, Henle 1989c, Henle 1991, Horner 1999, Read 1999, Sinervo et al. 2010, Kohler 2005, Michael et al. 2010, Michael and Lindenmayer 2010, Hutchinson 1993, Swanson 2007, Michael et al. 2011, Greer 1980, Swan and Watharow 2005 Shine and Greer 1991, Cogger 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, James and Shine 1988, Chapman and Dell 1985, Kohler 2005, Bush et al. 2010, Bush 1992, Swan and Watharow 2005
Scincidae	<i>Morethia obscura</i>	no	NA	NA	NA	mainland	-32	2.14	3.05	2	0.09	0.56	Shine and Greer 1991, Cogger 2000, Wilson and Swan 2003, Henle 1991, Greer 1989, Greer 2005, James and Shine 1988, Chapman and Dell 1985, Kohler 2005, Bush et al. 2010, Bush 1992, Swan and Watharow 2005
Scincidae	<i>Nannoscincus garrulus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	2.10	2.00	NA	NA	NA	Sadlier et al. 2006

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Nannoscincus gracilis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	0.58	2.50	NA	NA	NA	Greer 2001, Bauer and DeVaney 1987, Bauer and Vindum 1990, Sadlier et al. 2006
Scincidae	<i>Nannoscincus greeri</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	0.66	NA	NA	0.16	NA	Bauer and Sadlier 2000
Scincidae	<i>Nannoscincus maccoyi</i>	no	NA	NA	NA	mainland	-37	2.45	2.95	0.75	0.16	0.36	Shine and Greer 1991, Cogger 2000, Brown 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Cree 1994, Shine 1983, Vitt and Price 1982, Wilson and Swan 2008, James and Shine 1988, Taylor 1985, Fischer and Lindenmayer 2005, Kohler 2005, Shine and Wall 2008, Wilson and Swan 2011, Michael and Lindenmayer 2010, Hutchinson 1993, Heatwole and Taylor 1987, Swanson 2007, Downes and Shine 1999
Scincidae	<i>Nannoscincus mariei</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	1.03	1.50	NA	NA	NA	Bauer and DeVaney 1987, Bauer and Sadlier 2000, Bauer and Vindum 1990, Cogger 2000, Shine and Greer 1991, Wilson and Swan 2003, Greer 2005, Cree 1994, Melville and Swain 2000, Vitt and Price 1982, Wilson and Swan 2008, Fischer and Lindenmayer 2005, Michael and Lindenmayer 2010, Greer 1982, Murphy et al. 2006
Scincidae	<i>Niveoscincus coventryi</i>	no	NA	NA	NA	mainland	-36	1.40	2.55	1	0.08	0.20	Cogger 2000, Shine and Greer 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Towns 1994, Melville and Swain 2000, Radder et al. 2008, Sinervo et al. 2010, Cree and Hare 2010,
Scincidae	<i>Niveoscincus metallicus</i>	no	NA	NA	NA	mainland	-41	2.72	3.80	1	0.14	0.55	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Hutchinson 1993, Heatwole and Taylor 1987, Greer 1982, Swanson 2007
Scincidae	<i>Oligosoma fallai</i>	yes	New Zealand	Great Island	4.04	Continental	-34	38.79	4.50	1	NA	NA	Robb 1980, Alvarez 2004, Parrish and Gill 2003, Whitaker 1987, Towns et al. 2002, Jewell 2008
Scincidae	<i>Oligosoma homalonotum</i>	yes	New Zealand	Great Barrier	285.00	Land bridge	0	73.06	8.00	1	NA	NA	Robb 1980, Towns et al. 2002, Jewell 2008
Scincidae	<i>Oligosoma lichenigera</i>	yes	None	Lord Howe	15.04	Oceanic	-32	4.60	2.50	NA	0.44	NA	Cogger 2000, Wilson and Swan 2003, Greer 1989, Cogger et al. 1983, Wilson and Swan 2008, Kohler 2005
Scincidae	<i>Oligosoma nigriplantare</i>	yes	Chatham Islands	Pitt	744.60	Oceanic	-44	3.92	4.47	1	NA	NA	Robb 1980, Whitaker et al. 2002, Whitaker and Lyall 2004, Alvarez 2004, Patterson and Daugherty 1990, Cree 1994, Whitaker 1987, Towns et al. 2002, Hare 2005, Jewell 2008, Towns and Elliot 1996
Scincidae	<i>Oligosoma notosaurus</i>	yes	New Zealand	Stewart	1814.70	Land bridge	-47	6.20	2.50	1	0.24	0.61	Patterson and Daugherty 1990, Jewell 2008
Scincidae	<i>Oligosoma stenotis</i>	yes	New Zealand	Stewart	1814.70	Land bridge	-47	4.97	3.00	1	0.28	0.83	Patterson and Daugherty 1994, Jewell 2008
Scincidae	<i>Ophiomorus punctatissimus</i>	no	NA	NA	NA	mainland	38	1.70	3.50	1	0.27	0.93	Arnold and Ovenden 2004, Baran and Atatur 1998, Rogner 1997b, Valakos et al. 2008, Kwet 2009, Kohler 2005, In den Bosch 1988
Scincidae	<i>Pamelaescincus gardineri</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	6.80	2.00	NA	NA	NA	Cheke 1984, Bowler 2006, Rocha et al. 2010, Gerlach 2008, Evans and Evans 1980
Scincidae	<i>Panaspis kitsoni</i>	no	NA	NA	NA	mainland	0	1.51	1.92	6.5	0.21	2.68	Clobert et al. 1998, Fitch 1982, Dungen 1973, Greer et al. 1985, Fitch 1982, Turner 1977, Barbault 1974, Barbault 1974b

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Panaspis nimbaensis</i>	no	NA	NA	NA	mainland	8	1.63	2.51	5	0.08	1.06	Clobert et al. 1998, Fitch 1982, Hughes 1988, Fuhn 1972, Dunham et al. 1988, Fitch 1982, Barbault 1976, Barbault 1974, Leache 2005, Turner 1977, Bohme et al. 2011, Turner 1977, Damuth 1987, Barbault 1974 Fitch 1970, Fitzsimons 1943, Spawls et al. 2002, Loveridge 1936, 1942, Branch 1998, Razzetti and Msuya 2002, Broadley 1971, Parker 1942, Loveridge 1953, Auerbach 1987, Jacobson and Broadley 2000, Barbour and Loveridge 1928, Branch 2005, Graham and Marais 2007, Loveridge 1955, Fitch 1982, Sinervo et al. 2010, Kohler 2005, Lagen and Spawls 2010, Pienaar 1966, Haagner et al. 2000, Bowker 1984, Jacobsen 1982
Scincidae	<i>Panaspis wahlbergi</i>	no	NA	NA	NA	mainland	-9	1.43	4.45	1.5	0.05	0.32	Ferner et al. 1997, Gaulke 2011
Scincidae	<i>Parvosцинus sisoni</i>	yes	Philippine Islands	Panay	12011.10	Oceanic	11	0.64	1.00	NA	0.12	NA	Conant and Collins 1998, Smith 1946, Telford 1955, Clausen 1938, Lemay and Marsiglia 1952, Hotchkin et al. 2001, Walley 1998, Sinervo et al. 2010, Smith and Smith 1952, Youssef et al. 2008, Jensen et al. 2008, Beane et al. 2010, Gibbs et al. 2007, Brattstrom 1965
Scincidae	<i>Plestiodon anthracinus</i>	no	NA	NA	NA	mainland	35	4.34	8.50	1	0.12	1.01	Dunham and Miles 1985, Cree 1994, Dunham et al. 1988, Duellman 1961, Radder et al. 2008, Guillette 1983, Cree and Guillette 1995, Garcia-Vazquez and Feria-
Scincidae	<i>Plestiodon copei</i>	no	NA	NA	NA	mainland	19	6.02	3.53	1	0.24	0.83	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Ortiz 2006, Davis and Smith 1953, Ramirez-Pinilla et al. 2009, Duellman 1965, Lemos-Espinal et al. 1997
Scincidae	<i>Plestiodon egregius</i>	no	NA	NA	NA	mainland	31	0.76	4.80	1	0.13	0.64	Tinkle et al. 1970, Conant and Collins 1998, Smith 1946, Dunham et al. 1988, Warne and Charnov 2008, Griffith 1990, Mount 1968, Sinervo et al. 2010, Mount 1961, Mount 1963, Youssef et al. 2008, Jensen et al. 2008, Tinkle et al. 1967, Hailey and Elliot 1995
Scincidae	<i>Plestiodon fasciatus</i>	no	NA	NA	NA	mainland	37	6.17	7.70	1	0.17	1.34	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Radtkey et al. 1997, Anderson and Vitt 1990, Walker 1981, Perry and Garland 2002, Conant and Collins 1998, Cox et al. 2003, Smith 1946, Rogner 1997b, Cree 1994, Dunham et al. 1988, Andrews and Pough 1980, Warne and Charnov 2008, Cooper et al. 2000, Radder et al. 2008, Sinervo et al. 2010, Kohler 2005, Mount 1961, Youssef et al. 2008, Jensen et al. 2008, Werner and Whitaker 1978, Damuth 1987, Beane et al. 2010, Gibbs et al. 2007, Tinkle et al. 1967, Fitch 1955, Watson 2005, Light et al. 1966, Fitch 1956, Brattstrom 1965

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Plestiodon inexpectatus</i>	no	NA	NA	NA	mainland	33	7.04	7.50	1	0.16	1.22	Fitch 1970, Conant and Collins 1998, Mushinsky 1992, Rogner 1997b, Andrews and Pough 1980, Youssef et al. 2008, Sinervo et al. 2010, , Jensen et al. 2008, Beane et al. 2010, Brattstrom 1965
Scincidae	<i>Plestiodon kishinouyei</i>	yes	Ryukyu Islands	Iriomotejima	289.00	Oceanic	24	90.49	3.00	NA	NA	NA	Goris and Maeda 2004
Scincidae	<i>Plestiodon laticeps</i>	no	NA	NA	NA	mainland	35	20.77	13.70	1	0.40	5.54	Fitch 1970, Conant and Collins 1998, Cox et al. 2003, Smith 1946, Pianka and Vitt 2003, Vitt and Cooper 1986, Rogner 1997b, Dunham et al. 1988, Warne and Charnov 2008, Cooper et al. 2001, Vitt and Cooper 1985, Griffith 1990, Sinervo et al. 2010, Kohler 2005, Youssef et al. 2008, Jensen et al. 2008, Beane et al. 2010
Scincidae	<i>Plestiodon lynxe</i>	no	NA	NA	NA	mainland	22	4.34	4.35	1	0.21	0.93	Fitch 1970, Cox et al. 2003, Ramirez-Bautista et al. 1998, Boretto and Ibarquengoytia 2006, McCranie and Wilson 2001, Guillette 1983, Werler 1949, Werler 1951, Garcia-Vazquez and Feria-Ortiz 2006, Dixon and Lemos-Espinal 2010
Scincidae	<i>Plestiodon obsoletus</i>	no	NA	NA	NA	mainland	33	35.69	12.65	0.75	0.91	8.60	Tinkle et al. 1970, Clobert et al. 1998, Fitch 1985, Perry and Garland 2002, Conant and Collins 1998, Degenhardt et al. 1996, Smith 1946, Rogner 1997b, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Griffith 1990, Lemos-

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Plestiodon okadae</i>	yes	Miyako Islands	Miyako-jima	158.70	Oceanic	34	10.90	7.95	0.75	0.30	1.76	Espinal and Smith 2007, Garcia-Vazquez and Feria-Ortiz 2006, Lemos-Espinal and Smith 2007b, Turner 1977, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Mount 1961, Youssef et al. 2008, Tinkle et al. 1967, Fitch 1955, Brennan and Holycross 2009, Fitch 1956, Van Wilgen and Richardson 2012, Clobert et al. 1998, Perry and Garland 2002, Cox et al. 2003, Cree 1994, Dunham et al. 1988, Hasegawa 1994, Hasegawa 1984, Cree and Guillette 1995, Van Wyk 1991, Hasegawa 1997, Hasegawa 1990
Scincidae	<i>Plestiodon reynoldsi</i>	no	NA	NA	NA	mainland	28	3.22	2.00	0.75	0.21	0.32	Tinkle et al. 1970, Fitch 1970, Conant and Collins 1998, Pianka and Vitt 2003, Smith 1946, Telford 1959, Cooper 1953, Ashton and Telford 2006, Telford 1969, Ashton 2005, Youssef et al. 2008, Sinervo et al. 2010, Andrews 1994
Scincidae	<i>Plestiodon septentrionalis</i>	no	NA	NA	NA	mainland	39	7.54	10.65	1	0.21	2.21	Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1985, Conant and Collins 1998, Smith 1946, Cree 1994, Somma 1985, Somma 1987, Somma 1987b, Somma 1990, Somma 1991, Sinervo et al. 2010, Kohler 2005, Youssef et al. 2008, Fitch 1956, Brattstrom 1965

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Plestiodon skiltonianus</i>	no	NA	NA	NA	mainland	42	4.98	4.90	1	0.24	1.20	Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1985, Stebbins 2003, Grismer 2002, Smith 1946, Linsdale 1932, Cree 1994, Van Denburgh 1922, Dunham et al. 1988, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Youssef et al. 2008, Goldberg 2005, Brennan and Holycross 2009, Cunningham 1966, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006, St. John 2002
Scincidae	<i>Plestiodon stimpsonii</i>	yes	Ryukyu Islands	Iriomotejima	289.00	Oceanic	24	6.05	9.00	NA	NA	NA	Goris and Maeda 2004
Scincidae	<i>Plestiodon tetragrammus</i>	no	NA	NA	NA	mainland	29	4.72	8.50	0.835	0.17	1.24	Fitch 1970, Conant and Collins 1998, Degenhardt et al. 1996, Smith 1946, Werler 1951, Lemos-Espinal and Smith 2007, Garcia-Vazquez and Feria-Ortiz 2006, Lieb 1985, Lieb 1990, Lemos-Espinal and Smith 2007b, Jones and Lovich 2009, Dixon and Lemos-Espinal 2010, Brattstrom 1965, Cogger 2000, Brown 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Huey and Bennett 1987, Melville and Swain 2000, Shine 1983, Vitt and Price 1982, Taylor 1985, Fischer and Lindenmayer 2005, Sinervo et al. 2010, Michael and Lindenmayer 2010, Heatwole and Taylor 1987, Hutchinson and Donnellan 1992, Greer 1982, Thompson et al. 1999,
Scincidae	<i>Pseudemoia entrecasteauxii</i>	no	NA	NA	NA	mainland	-37	2.78	4.05	1	0.20	0.81	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Bennett and John-Alder 1986, Murphy et al. 2006
Scincidae	<i>Pseudemoia pagenstecheri</i>	no	NA	NA	NA	mainland	-35	2.89	5.85	1	0.16	0.95	Cogger 2000, Wilson and Swan 2003, Greer 2005, Thompson et al. 1999, Michael and Lindenmayer 2010, Hutchinson and Donnellan 1992, Swanson 2007, Thompson et al. 1999
Scincidae	<i>Pseudemoia rawlinsoni</i>	no	NA	NA	NA	mainland	-37	2.80	5.60	1	0.10	0.57	Cogger 2000, Wilson and Swan 2003, Greer 2005, Taylor 1985, Michael and Lindenmayer 2010 Shine and Greer 1991, Cogger 2001, Brown 1991, Wilson and Swan 2003, Greer 1989, Greer 2005, Shine 1983, Wilson and Swan 2008, Taylor 1985, Fischer and Lindenmayer 2005, Michael and Lindenmayer 2010, Hutchinson 1993, Heatwole and Taylor 1987, Greer 1982, Swanson 2007, Thompson et al. 1999
Scincidae	<i>Pseudemoia spenceri</i>	no	NA	NA	NA	mainland	-37	3.44	2.10	1	0.24	0.50	Fitch 1985, Huang 2006b, Vrcibradic and Rocha 2005, Vrcibradic and Rocha 1996, Cree 1994, Blackburn and Vitt 1992, Vrcibradic & Rocha 1996b, Sinervo et al. 2010, Ramirez-Pinilla et al. 2002, Rocha et al. 2009, Vrcibradic and Rocha 2011, Winck and Rocha 2012
Scincidae	<i>Psychosaura macrorhyncha</i>	no	NA	NA	NA	mainland	-17	6.32	2.80	1	0.47	1.30	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Scincella lateralis</i>	no	NA	NA	NA	mainland	34	1.66	3.08	3.5	0.10	1.09	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Perry and Garland 2002, Conant and Collins 1998, Pianka and Vitt 2003, Smith 1946, Cree 1994, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Radder et al. 2008, Milstead 1961, Werler 1951, Lemos-Espinal and Smith 2007, Sinervo et al. 2010, Kohler 2005, Fitch and Greene 1965, Jones and Lovich 2009, Jensen et al. 2008, Turner 1977, Beane et al. 2010, Fitch 1956, Hailey and Elliot 1995, Brattstrom 1965
Scincidae	<i>Scincus mitranus</i>	no	NA	NA	NA	mainland	22	21.32	5.00	1	0.79	3.97	Leviton et al. 1992, Pianka and Vitt 2003, Arnold 1984, Arnold 1980, Schatti and Desvoignes 1999, Jongbloed 2000, Cooper et al. 2000, Al-Johany et al. 1997, Hornby 1996, Al-Sadoon et al. 1999, Al-Johany et al. 1999, Sinervo et al. 2010, van der Kooij 2001
Scincidae	<i>Scincus scincus</i>	no	NA	NA	NA	mainland	27	20.33	4.00	1	2.45	9.79	Schleich et al. 1996, Amitai and Bouskila 2001, Anderson 1999, Disi et al. 2001, Dunger 1973, Papenfuss 1969, Gallagher 1971, Arnold 1984, El Din 2006, Rogner 1997b, Schatti and Desvoignes 1999, Le Berre 1989, Anderson 1963, Werner 1987, Hornby 1996, Kohler 2005, Bar and Haimovitch 2012, Trape et al.

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Sigaloseps deplanchei</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	1.69	2.50	NA	NA	NA	Bauer and DeVaney 1987, Bauer and Sadlier 2000, Bauer and Vindum 1990
Scincidae	<i>Sigaloseps ruficauda</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	2.01	3.00	NA	0.24	NA	Bauer and Sadlier 2000
Scincidae	<i>Sphenomorphus concinnatus</i>	yes	Solomon Islands	Bougainville	8590.10	Oceanic	-8	3.92	2.00	NA	0.28	NA	McCoy 1980, McCoy 2006, Greer and Parker 1967, Greer and Parker 1968
Scincidae	<i>Sphenomorphus cranei</i>	yes	Solomon Islands	Bougainville	8590.10	Oceanic	-8	5.15	2.80	NA	0.40	NA	Greer and Shea 2004, McCoy 1980, Greer and Parker 1967, Greer and Parker 1974, McCoy 2006, Greer and Parker 1968, Smith 1935, Fitch 1970, Schleich and Kastle 2002, Ji et al. 2006, Tikader and Sharma 1992, Taylor 1963, Dejun 1989, Teynie 2004, Pope 1929, Manthey and Grossmann 1997, Ziegler 2002, Cox et al. 1998, Song 1987, Karsen et al. 1986, Inger and Colwell 1977, Shrestha 2001, Grismer et al. 2007, Shine 1985, Das 2002, Sinervo et al. 2010, Taylor and Elbel 1958, Das 2010, Cree and Hare 2010, Cox et al. 2010, Schleich and Kastle 2002, Inger et al. 1990, Huang 2010, Grismer 2011, Teynie and David 2010, Ahmed 2009
Scincidae	<i>Sphenomorphus indicus</i>	no	NA	NA	NA	mainland	28	9.99	7.27	1	0.30	2.15	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Sphenomorphus maculatus</i>	no	NA	NA	NA	mainland	25	3.36	4.50	1	0.35	1.59	Smith 1935, Schleich and Kastle 2002, Taylor 1963, Stuart et al. 2006, Bauer and Gunther 1992, Cox et al. 1998, Inger and Colwell 1977, Pauwels et al. 2003, Shrestha 2001, Stuart and Emmett 2006, Mahony and Ali Reza 2008, Das 2002, Taylor and Elbel 1958, Das 2010, Cox et al. 2010, Ahmed et al. 2009, Huang 2010, Grismer 2011, Teynie and David 2010, Ahmed 2009
Scincidae	<i>Sphenomorphus scutatus</i>	yes	Palau Islands	Babeldaob	374.10	Oceanic	7	1.12	2.00	NA	NA	NA	Greer and Shea 2004, Crombie and Pregill 1999, Greer and Parker 1974
Scincidae	<i>Tiliqua adelaidensis</i>	no	NA	NA	NA	mainland	-35	18.74	3.23	1	1.58	5.09	Cogger 2000, Wilson and Swan 2003, Greer 2005, Fenner et al. 2007, Milne et al. 2002, Wilson and Swan 2008, Shea 2006, Swanson 2007, Clarke 2000
Scincidae	<i>Tiliqua nigrolutea</i>	no	NA	NA	NA	mainland	-37	651.54	7.00	0.375	32.87	86.27	Greer 2001, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Boretto and Ibarquengoytia 2006, Edwards et al. 2002, Edwards 1999, Shea 2006, Ibarquengoytia and Casalins 2007, Michael and Lindenmayer 2010, Heatwole and Taylor 1987, Swanson 2007
Scincidae	<i>Tiliqua occipitalis</i>	no	NA	NA	NA	mainland	-30	549.32	5.00	0.75	23.31	87.43	Greer 2001, Cogger 2000, Wilson and Swan 2003, Greer 1989, Greer 2005, Shea 2006, Horner 1999, Sinervo et al. 2010, Bush et al. 2010, Bush 1992, Heatwole and Taylor 1987, Swanson 2007, Light et

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													al. 1966, Swan and Watharow 2005
Scincidae	<i>Tiliqua rugosa</i>	no	NA	NA	NA	mainland	-31	470.92	2.05	1	43.12	88.39	Clobert et al. 1998, Greer 2001, Fitch 1970, Cogger 2000, Perry and Garland 2002, Cox et al. 2003, Cooper and Vitt 2002, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997b, Greer 1989, Greer 2005, Clusella-Trullas et al. 2008, Garrick 2008, Uller et al. 2009, Shea 2006, Henle 1989c, Sinervo et al. 2010, Bush et al. 2010, Michael and Lindenmayer 2010, Hutchinson 1993, MacMillen et al. 1989, Heatwole and Taylor 1987, Swanson 2007, Bennett and John-Alder 1986, Light et al. 1966, Swan and Watharow 2005
Scincidae	<i>Tiliqua scincoides</i>	no	NA	NA	NA	mainland	-24	852.85	11.00	0.75	8.25	68.06	Fitch 1970, Cogger 2000, Wilson and Swan 2003, Rogner 1997b, Greer 2005, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Todd 2008, Shea 2006, Horner 1999, Sinervo et al. 2010, De Magalhaes and Costa 2009, Michael and Lindenmayer 2010, Koenig et al. 2001, Heatwole and Taylor 1987, Swanson 2007, Michael et al. 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Trachylepis affinis</i>	no	NA	NA	NA	mainland	8	4.98	2.72	7.5	0.21	4.20	Clobert et al. 1998, Fitch 1982, Greer et al. 1985, Hughes 1988, Dunger 1972, Schmidt et al. 1919, Chirio and LeBreton 2007, Akani et al. 2002, Leache et al. 2006, Fitch 1982, Pauwels and Vande weghe 2008, Reid 1986, Pauwels et al. 2004, Turner 1977, Chirio 2009, Ullenbruch et al. 2010, Barbault 1974, Scott 1982, Barbault 1974b, Cisse and Karns 1978, Trape et al. 2012
Scincidae	<i>Trachylepis aurata</i>	no	NA	NA	NA	mainland	34	8.14	5.00	1	0.44	2.21	Smith 1935, Arnold and Ovenden 2004, Szczerbak 2003, Anderson 1999, Baran and Atatur 1998, Reed and Marx 1959, Ahmadzadeh et al. 2008, Valakos et al. 2008, Weber 1960, Gallagher 1971, Rogner 1997b, Largen and Spawls 2010, van der Kooij 2001, Arakelyan et al. 2011, Fathinia et al. 2009
Scincidae	<i>Trachylepis buettneri</i>	no	NA	NA	NA	mainland	7	6.82	8.70	1.5	0.16	2.11	Clobert et al. 1998, Fitch 1982, Dunger 1973, Hughes 1988, Schmidt et al. 1919, Chirio and LeBreton 2007, Dunham et al. 1988, Leache et al. 2006, Fitch 1982, Barbault 1976, Barbault 1974, Turner 1977, Damuth 1987, Barbault 1974, Jacobsen 1982, Trape et al. 2012
Scincidae	<i>Trachylepis lavarambo</i>	yes	Madagascar	Nosy Be	290.30	Oceanic	-13	3.57	3.00	NA	NA	NA	Raselimanana and Rakotomalala 2004, Nussbaum and Raxworthy 1998, Glaw and Vences 2007

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Trachylepis maculilabris</i>	no	NA	NA	NA	mainland	5	7.52	7.04	5	0.21	7.55	Clobert et al. 1998, Fitch 1970, 1982, Loveridge 1936, Spawls et al. 2002, Branch 1998, Loveridge 1942, Broadley 1971, Hughes 1988, Parker 1936, Dunger 1972, Schmidt et al. 1919, Loveridge 1953, Broadley 2000, Barbour and Loveridge 1928, Branch 2005, Chirio and LeBreton 2007, Akani et al. 2002, Dunham et al. 1988, Largen and Spawls 2006, Leache et al. 2006, Vonesh 1998, Fitch 1982, Pauwels and Vande weghe 2008, Reid 1986, Barbault 1976, Turner 1977, Goldberg and Bursey 2010, Branch et al. 2005, Largen and Spawls 2010, Barbault 1974, Haagner et al. 2000, Damuth 1987, Curry-Lindahl 1979, Barbault 1974, Trape et al. 2012
Scincidae	<i>Trachylepis quinquetaeniata</i>	no	NA	NA	NA	mainland	13	11.21	4.80	2	0.53	5.11	Fitch 1970, FitzSimons 1943, Spawls et al. 2002, Branch 1998, Loveridge 1936, Flower 1933, Joger and Lambert 1996, Broadley 1971, Hughes 1988, Bohme et al. 1996, Dunger 1972, El Din 2006, Schmidt et al. 1919, Rogner 1997b, Le Berre 1989, Branch 2005, Chirio and LeBreton 2007, Dunham et al. 1988, Ineich 1999, Leache et al. 2006, Fitch 1982, Chirio 2009, Sinervo et al. 2010, Kohler 2005, Largen and Spawls 2010, Broadley and Bauer

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1999, Simbotwe 1980, Cooper and Whiting 2000, Bowker 1984, Damuth 1987, Western 1974, Curry-Lindahl 1979, Trape et al. 2012
Scincidae	<i>Trachylepis sechellensis</i>	yes	Seychelles Islands	Mahe	148.00	Continental	-5	11.80	4.00	NA	0.69	NA	Perry and Garland 2002, Henkel and Schmidt 2000, Cheke 1984, Bowler 2006, Somma 1991, Rocha et al. 2010, Brooke and Houston 1983, Evans and Evans 1980, Crawford and Thorpe 1979, Gerlach 2005
Scincidae	<i>Trachylepis sparsa</i>	no	NA	NA	NA	mainland	-28	11.91	5.15	1	0.35	1.82	Broadley 2000, Rocha et al. 2009, Goldberg 2007, Huey and Pianka 2007, Cooper and Whiting 2000, Vitt and Pianka 1977
Scincidae	<i>Trachylepis spilogaster</i>	no	NA	NA	NA	mainland	-28	8.66	4.95	0.75	0.26	0.97	Branch 1998, Cox et al. 2003, Pianka and Vitt 2003, Pianka 1986, Auerbach 1987, Broadley 2000, Huey et al. 2001, Sinervo et al. 2010, Goldberg 2006, Cooper and Whiting 2000, Vitt and Pianka 1977, Huey and Pianka 1977

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Trachylepis striata</i>	no	NA	NA	NA	mainland	-7	9.77	6.70	2.5	0.28	4.66	Tinkle et al. 1970, Fitch 1970, Spawls et al. 2002, FitzSimons 1943, Loveridge 1936, 1942, Razzetti and Msuya 2002, Pianka 1986, Huang 2006b, Pianka and Vitt 2003, Broadley 1971, Parker 1932, Jeffery 1993, Loveridge 1953, Patterson 1990, Auerbach 1987, Broadley 2000, Barbour and Loveridge 1928, Nagy et al. 1999, Branch 2005, Dunham et al. 1988, Ineich 1999, Gans et al. 1965, Loveridge 1955, Brown and Nagy 2007, Broadley 1992, Fitch 1982, Huey et al. 2001, Todd 2008, Meik et al. 2002, Sinervo et al. 2010, Largen and Spawls 2010, Pienaar 1966, Simbotwe 1980, Haagner et al. 2000, Curry-Lindahl 1979, Huey and Pianka 1977, Patterson 1991, Brattstrom 1965, Clobert et al. 1998, Fitch 1970, Spawls et al. 2002, FitzSimons 1943, Branch 1998, Loveridge 1936, Razzetti and Msuya 2002, Broadley 1971, Parker 1932, Schmidt 1943, Parker 1942, Loveridge 1953, Auerbach 1987, Broadley 2000, Barbour and Loveridge 1928, Branch 2005, Ineich 1999, Largen and Spawls 2006, Loveridge 1955, Meik et al. 2002, Sinervo et al. 2010, Kohler 2005, Goldberg and Bursley 2010, Branch et al.
Scincidae	<i>Trachylepis varia</i>	no	NA	NA	NA	mainland	-12	4.72	7.64	1	0.15	1.15	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2005, Largen and Spawls 2010, Pienaar 1966, Haagner et al. 2000, Jacobsen 1982, Kirchof et al. 2010, Hebrard et al. 1982
Scincidae	<i>Trachylepis vittata</i>	no	NA	NA	NA	mainland	34	5.20	5.50	1	0.33	1.83	Schleich et al. 1996, Amitai and Bouskila 2001, Arbel 1984, Anderson 1999, Disi et al. 2001, Baran and Atatur 1998, El Din 2006, Rogner 1997b, Le Berre 1989, Atatur and Gocmen 2001, Baier et al. 2009, Bar and Haimovitch 2012, Fathinia et al. 2009
Scincidae	<i>Trachylepis wrightii</i>	yes	Seychelles Islands	Frigate	2.19	Continental	-4	83.21	4.50	NA	1.58	NA	Greer 2001, Perry and Garland 2002, Henkel and Schmidt 2000, Cheke 1984, Bowler 2006, Somma 1991, Rocha et al. 2010, Gerlach 2008, Brooke and Houston 1983
Scincidae	<i>Tribolonotus blanchardi</i>	yes	Solomon Islands	Bougainville	8590.10	Oceanic	-7	0.63	1.00	NA	NA	NA	Greer 1977, McCoy 1980, Greer and Parker 1967, McCoy 2006, Greer and Parker 1968
Scincidae	<i>Tribolonotus brongersmai</i>	yes	Bismarck Archipelago	Manus	1940.20	Oceanic	-2	3.97	NA	NA	0.30	NA	Cogger 1972
Scincidae	<i>Tribolonotus ponceleti</i>	yes	Solomon Islands	Bougainville	8590.10	Oceanic	-7	31.04	1.00	NA	1.88	NA	McCoy 1980, McCoy 2006, Greer and Parker 1968
Scincidae	<i>Tribolonotus pseudoponceleti</i>	yes	Solomon Islands	Bougainville	8590.10	Oceanic	-6	2.53	1.00	NA	0.28	NA	Greer 1977, McCoy 1980, McCoy 2006, Greer and Parker 1968
Scincidae	<i>Tribolonotus schmidti</i>	yes	Solomon Islands	Guadalcanal	5280.10	Oceanic	-10	0.76	1.00	NA	NA	NA	Greer 2001, Greer 1977, McCoy 1980, McCoy 2006, Greer and Parker 1968

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Scincidae	<i>Tropidoscincus boreus</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-21	18.24	2.50	NA	NA	NA	Bauer and Sadlier 2000, Shea et al. 2009, Whittaker et al. 2004
Scincidae	<i>Tropidoscincus variabilis</i>	yes	New Caledonia	New Caledonia	18575.00	Continental	-22	9.73	2.80	NA	0.60	NA	Bauer and DeVaney 1987, Bauer and Sadlier 2000, Bauer and Vindum 1990, Shea et al. 2009
Scincidae	<i>Varzea bistriata</i>	no	NA	NA	NA	mainland	-4	16.82	4.70	1	0.78	3.67	Vitt and Blackburn 1991, Vitt and Caldwell 1993, Avila-Pires 1995, Martins 1991, Rodrigues and Cadle 1990, Zimmerman and Rodrigues 1990, Duellman 1990, Cree 1994, Warne and Charnov 2008, Murphy 1997, Blackburn and Vitt 1992, Malhotra and Thorpe 1999, Miralles et al. 2009, Ramirez-Pinilla et al. 2002
Sphaerodactylidae	<i>Aristelliger barbouri</i>	yes	Bahamas	Great Inagua	1615.30	Oceanic	21	2.32	1.00	NA	0.15	NA	Shine and Greer 1991, Schwartz and Henderson 1991, Nobel and Kleengel 1932, Bauer and Russell 1993, Henderson and Powell 2009, Rosler 2005
Sphaerodactylidae	<i>Aristelliger georgeensis</i>	no	NA	NA	NA	mainland	13	17.84	1.00	1	0.59	0.59	Stafford and Meyer 2000, Kohler 2003, Schwartz and Henderson 1991, Kohler 2008, Bauer and Russell 1993, Lopez and Gonzalez 1997, Duellman 1965, Rosler 2005
Sphaerodactylidae	<i>Aristelliger praesignis</i>	yes	Greater Antilles	Jamaica	11189.60	Oceanic	18	4.07	1.00	2	0.49	0.98	Schwartz and Henderson 1991, Bauer and Russell 1993, Henderson and Powell 2009, Rosler 2005, Henkel and Schmidt 1995

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Sphaerodactylidae	<i>Gonatodes albogularis</i>	no	NA	NA	NA	mainland	11	1.66	1.50	12.5	0.15	2.88	Clobert et al. 1998, Fitch 1970, Fitch 1973, Stafford and Meyer 2000, Conant and Collins 1998, Savage 2002, Lee 2000, Kohler 2003, Rand and Myers 1990, Beebe 1944b, Smith 1946, Schwartz and Henderson 1991, Evans 1947, Molina et al. 2004, Rogner 1997a, Taylor 1956, Werner and Seifan 2006, Kohler et al. 2006, Guyer and Donnelly 2005, Van Buurt 2005, Murphy 1997, Rand 1957, Kohler 2008, Todd 2008, Kohler 2005, Henderson and Powell 2009, Rosler 2005, Heatwole and Sexton 1966, Castro-Herrera et al. 2007, Tinkle et al. 1967, Henle 1990, Van Wilgen and Richardson 2012, Henkel and Schmidt 1995, Leenders and Watkins-Colwell 2004
Sphaerodactylidae	<i>Gonatodes daudini</i>	yes	Windward Islands	Union	8.49	Oceanic	13	0.39	1.00	NA	NA	NA	Powell and Henderson 2005, Daudin and de Silva 2007, Henderson and Powell 2009, Rojas-Runjaic et al. 2010, Bentz et al. 2011, John et al. 2012
Sphaerodactylidae	<i>Gonatodes ocellatus</i>	yes	Trinidad and Tobago	Tobago	308.80	Land bridge	11	1.96	1.00	NA	0.17	NA	Beebe 1944b, Murphy 1997, Rojas-Runjaic et al. 2010, Cox et al. 2003, Beebe 1944b, Werner and Seifan 2006, Lotzkat 2007, Murphy 1997, Fuenmayor et al. 2005, Test et al. 1966, Turner 1977, Ugueto and Rivas 2010, Rosler 2005, Henle 1990, Henkel and Schmidt 1995
Sphaerodactylidae	<i>Gonatodes vittatus</i>	no	NA	NA	NA	mainland	10	0.66	1.50	12	0.09	1.70	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Sphaerodactylidae	<i>Pristurus abdelkuri</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	12	1.16	1.00	NA	NA	NA	Schatti and Desvoignes 1999, Arnold 1993, Arnold 2009, Razzetti et al. 2011, Rosler and Wranik 2004, Rosler and Wranik 2006
Sphaerodactylidae	<i>Pristurus guichardi</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	13	0.98	1.00	NA	NA	NA	Schatti and Desvoignes 1999, Arnold 1986, Joger 2000, Arnold 2009, Razzetti et al. 2011, Rosler and Wranik 2004, Rosler and Wranik 2006
Sphaerodactylidae	<i>Pristurus insignis</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	12	3.20	1.00	NA	NA	NA	Schatti and Desvoignes 1999, Joger 2000, Arnold 1993, Arnold 2009, Razzetti et al. 2011, Rosler and Wranik 2004, Rosler and Wranik 2006
Sphaerodactylidae	<i>Pristurus insignoides</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	13	2.05	1.00	NA	NA	NA	Schatti and Desvoignes 1999, Arnold 1986, Razzetti et al. 2011, Rosler and Wranik 2004, Rosler and Wranik 2006
Sphaerodactylidae	<i>Pristurus rupestris</i>	no	NA	NA	NA	mainland	22	0.56	1.00	1.5	0.10	0.15	Smith 1935, Shine and Greer 1991, Anderson 1999, Disi et al. 2001, Leviton et al. 1992, Anderson 1896, Loveridge 1947, Gallagher 1971, Arnold 1984, Parker 1942, Arnold 1980, Schatti and Desvoignes 1999, Jongbloed 2000, Arnold 1993, Arnold 2009, Lagen and Spawls 2010, Rosler 2005, van der Kooij 2001
Sphaerodactylidae	<i>Pristurus sokotranus</i>	yes	Socotra Archipelago	Socotra	3606.70	Continental	13	1.11	1.00	NA	0.12	NA	Loveridge 1947, Schatti and Desvoignes 1999, Joger 2000, Uetz 2006, Arnold 1993, Arnold 2009, Carretero and Lo Cascio 2010, Rosler 2005, Razzetti et al. 2011, Rosler and Wranik 2004, Rosler and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Wranik 2006
Sphaerodactylidae	<i>Quedenfeldtia trachyblepharus</i>	no	NA	NA	NA	mainland	31	1.92	1.00	2.5	0.22	0.54	Schleich et al. 1996, Loveridge 1947, Rogner 1997a, Arnold 1990, Arnold 1993, Bons and Geniez 1996, Kohler 2005, Rosler 2005, Daza et al. 2009, Carretero et al. 2006, Arnold 1993
Sphaerodactylidae	<i>Sphaerodactylus argivus</i>	yes	Cayman Islands	Grand Cayman	196.30	Oceanic	19	0.43	1.00	NA	NA	NA	Schwartz and Henderson 1991, Henderson and Powell 2009,
Sphaerodactylidae	<i>Sphaerodactylus kirbyi</i>	yes	Windward Islands	Bequia	18.00	Oceanic	13	0.36	1.00	NA	0.08	NA	Daudin and de Silva 2007, Henderson and Powell 2009, Bentz et al. 2011
Sphaerodactylidae	<i>Sphaerodactylus nicholsi</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	0.21	1.00	10.5	0.03	0.27	Schwartz and Henderson 1991, Heatwole and Veron 1977, Huey et al. 2009, Lopez-Ortiz and Lewis 2002, Henderson and Powell 2009, Rivero 1998, Regalado 2006
Sphaerodactylidae	<i>Sphaerodactylus roosevelti</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	1.06	1.00	5	0.19	0.94	Schwartz and Henderson 1991, Huey et al. 2009, Henderson and Powell 2009, Rivero 1998, Regalado 2006
Sphaerodactylidae	<i>Sphaerodactylus vincenti</i>	yes	Windward Islands	Martinique	1166.60	Oceanic	14	0.53	1.00	NA	0.07	NA	Schwartz and Henderson 1992, Steinberg et al. 2007, Malhotra and Thorpe 1999, Steinberg et al. 2008, Henderson and Powell 2009, Rosler 2005, Bentz et al. 2011, Daltry 2009, Alexander 2007, Leclair and Leclair 2011
Sphaerodactylidae	<i>Teratoscincus microlepis</i>	no	NA	NA	NA	mainland	29	10.23	2.00	3	1.47	8.80	Anderson 1999, Minton 1966, Szczerbak and Golubev 1996, Rogner 1997a, Khan 2006, Sharma 2002, Kohler 2005, Rosler 2005, Werner et al.

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Sphaerodactylidae	<i>Teratoscincus scincus</i>	no	NA	NA	NA	mainland	41	11.13	1.50	2	1.52	4.57	Smith 1935, Szczerbak 2003, Anderson 1999, Minton 1966, Szczerbak and Golubev 1996, Greene 1982, Arnold 1984, Rogner 1997a, Anderson and Leviton 1969, Khan 2006, Sharma 2002, Werner and Seifan 2006, Clark 1990, Jongbloed 2000, Garrick 2008, Bauer 1990, Kohler 2005, Rosler 2005, Henkel and Schmidt 1995, Werner et al. 1993 Anderson and Vitt 1990, Fitch 1985, Perry and Garland 2002, Conant and Collins 1998, Savage 2002, Cox et al. 2003, Avila-Pires 1995, Martins 1991, Vitt and de Carvalho 1992, Vitt and de Carvalho 1995, Duellman 1978, Beebe 1945, Kohler 2003, Rand and Myers 1990, Rodrigues and Cadle 1990, Zimmerman and Rodrigues 1990, Duellman 1990, Pianka and Vitt 2003, Schwartz and Henderson 1991, Cei 1993, Mesquita et al. 2006b, Dixon and Soini 1986, Hoogmoed 1973, Gainsbury and Colli 2003, Nogueira et al. 2005, Vitt 2000, Molina et al. 2004, Rogner 1997b, Vitt and Zani 1998, Duellman 2005, Bartlett and Bartlett 2003, Daudin and de Silva 2007, Colli et al. 2002, Dunham et
Teiidae	<i>Ameiva ameiva</i>	no	NA	NA	NA	mainland	-6	52.73	5.05	3	1.59	24.08	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Ameiva erythrocephala</i>	yes	Leeward Islands	Nevis	92.30	Oceanic	17	12.00	4.00	NA	NA	NA	al. 1988, Janzen 1973, Rodrigues 2003, Rodrigues 1996, Lotzkat 2007, Murphy 1997, Colli 1991, Vitt 1991, Vitt et al. 1999, Vitt 1995, Vitt et al. 2000, Huey et al. 2001, Radder et al. 2008, Kohler 2008, Fuenmayor et al. 2005, Test et al. 1966, McElroy et al. 2008, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler 2005, Henderson and Powell 2009, Avila-Pires et al. 2010, Ugueto and Rivas 2010, Duellman 1987, Gasc 1990, Fitch 1968, Rocha et al. 2009, Van Wilgen and Richardson 2012, Sales et al. 2011, Brattstrom 1965, Sales et al. 2011 Schwartz and Henderson 1991, Powell et al. 2005, Kerr et al. 2005, Malhotra and Thorpe 1999, Kerr et al. 2006, Henderson and Powell 2009 Perry and Garland 2002, Cox et al. 2003, Rodda et al. 2001, Schwartz and Henderson 1991, Rodriguez-Ramirez and Lewis 1991, Lewis et al. 2000, Lopez-Ortiz and Lewis 2002, Sinervo et al. 2010, Kohler 2005, Henderson and Powell 2009, Rivero 1998
Teiidae	<i>Ameiva exsul</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	24.32	2.68	2	1.24	6.64	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Ameiva fuscata</i>	yes	Windward Islands	Dominica	787.30	Oceanic	15	58.02	4.00	2.5	3.66	36.61	Schwartz and Henderson 1991, Dunham et al. 1988, Pregill 1984, Malhotra and Thorpe 1999, Somma and Brooks 1976, Malhotra and Thorpe 1995, Sinervo et al. 2010, Kohler 2005, Henderson and Powell 2009, Brooks 1968, Alexander 2007, Bullock and Evans 1990
Teiidae	<i>Ameiva plei</i>	yes	Leeward Islands	Saint Martin	91.90	Oceanic	18	18.79	2.60	2	NA	NA	Perry and Garland 2002, Powell et al. 2005, Hodge et al. 2003, Pregill 1984, Censky 1995, Breuil 2002, Kohler 2005, Henderson and Powell 2009
Teiidae	<i>Ameiva polops</i>	yes	Virgin Islands	St. Croix	214.40	Oceanic	18	9.37	NA	NA	0.70	NA	Schwartz and Henderson 1991, Meier et al. 1993, Cope 1895, Henderson and Powell 2009, McNair and Lombard 2004
Teiidae	<i>Ameiva wetmorei</i>	yes	Greater Antilles	Puerto Rico	9099.80	Continental	18	2.63	1.00	2	NA	NA	Cox et al. 2003, Vitt and de Carvalho 1992, Schwartz and Henderson 1991, Rodriguez-Ramirez and Lewis 1991, Kohler 2005, Henderson and Powell 2009, Rivero 1998, Anderson and Vitt 1990, Fitch 1985, Fitzgerald et al. 1999, Vitt and de Carvalho 1992, Vitt and Breitenbach 1993, Dias and Rocha 2004, Cei 1993, Mesquita and Colli 2003, Colli et al. 2002, Colli et al. 2003, Dunham et al. 1988, Rodrigues 2003, Rodrigues 1996, Vitt and Price 1982, Vitt 1991, Vitt 1995, Vanzolini et al. 1980, Sinervo et al. 2010, Kohler
Teiidae	<i>Ameivula ocellifera</i>	no	NA	NA	NA	mainland	-19	7.28	2.15	4	0.51	4.36	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2005, Menezes and Rocha 2011, Rocha et al. 2009, Sales et al. 2012, Winck and Rocha 2012
Teiidae	<i>Aspidoscelis burti</i>	no	NA	NA	NA	mainland	30	31.80	4.33	1.5	0.95	6.14	Stebbins 2003, Degenhardt et al. 1996, Goldberg 1987, Jones and Lovich 2009, Brennan and Holycross 2009, Grismer 2002, Case 1982, Van Denburgh 1922, Walker et al. 1966, Sinervo et al. 2010, Woolrich-Pina et al. 2011, Soule 1963, Brattstrom 1965
Teiidae	<i>Aspidoscelis ceralbensis</i>	yes	Baja California	Ceralbo	150.50	Oceanic	24	7.60	2.50	NA	0.95	NA	Campbell 1999, Stafford and Meyer 2000, Lee 2000, Kohler 2003, Manriquez-Moran et al. 2008, Lopez and Gonzalez 1997, Walker 2012, Clobert et al. 1998, Fitch 1970, 1973, Mata-Silva and Ramirez-Bautista 2005, Vitt et al. 1993, Savage 2002, Lee 2000, Vitt and de Carvalho 1992, Vitt and Breitenbach 1993, Pianka and Vitt 2003, Kohler et al. 2006, Duellman 1961, Cooper et al. 2001, Rand 1957, Kohler 2008, Echternacht 1983, Davis and Dixon 1961, Davis and Smith 1953, Kohler 2005, Duellman 1965, Kennedy 1968, Woolrich-Pina et al. 2011, Leenders and Watkins-Colwell 2004
Teiidae	<i>Aspidoscelis cozumelae</i>	no	NA	NA	NA	mainland	18	7.22	1.80	1.5	0.58	1.57	
Teiidae	<i>Aspidoscelis deppei</i>	no	NA	NA	NA	mainland	16	8.51	2.30	4	0.74	6.79	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Aspidoscelis dixonii</i>	no	NA	NA	NA	mainland	31	17.50	3.83	1	1.04	3.97	Stebbins 2003, Degenhardt et al. 1996, Warne and Charnov 2008, Jones and Lovich 2009 Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, Conant and Collins 1998, Stebbins 2003, Vitt and Breitenbach 1993, Schall 1993, Degenhardt et al. 1996, Winne and Beck 2004, Medica 1967, Rogner 1997b, Eiffler and Passek 2000, Dunham et al. 1988, Vitt and Price 1982, Schall 1978, Stuart 1991, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Woolrich-Pina et al. 2011, Brennan and Holycross 2009
Teiidae	<i>Aspidoscelis exsanguis</i>	no	NA	NA	NA	mainland	32	15.20	3.55	1.5	0.78	4.14	Fitch 1985, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Vitt and Breitenbach 1993, Schall 1993, Degenhardt et al. 1996, Smith 1946, Rogner 1997b, Eiffler and Passek 2000, McCranie and Wilson 2001, Vitt and Price 1982, Vitt and Price 1982, Ballinger and Schrank 1972, Lemos-Espinal and Smith 2007, Schall 1978, Lemos-Espinal and Smith 2007b, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Turner 1977, Curry-Lindahl 1979, Siever and Paulissen 1996, Woolrich-Pina et al. 2011, Dixon and Lemos-Espinal 2010, Ramirez-Bautista et al.
Teiidae	<i>Aspidoscelis gularis</i>	no	NA	NA	NA	mainland	29	13.34	3.74	2	0.90	6.74	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2009, Brattstrom 1965
Teiidae	<i>Aspidoscelis guttata</i>	no	NA	NA	NA	mainland	18	23.44	4.50	2	1.14	10.22	Fitch 1970, Mata-Silva and Ramirez-Bautista 2005, Kohler 2003, Greene 1982, Davis and Dixon 1961, Kennedy 1968, Woolrich-Pina et al. 2011 Tinkle et al. 1970, Dunham and Miles 1985, Radtkey et al. 1997, Walker 1981, Perry and Garland 2002, Stebbins 2003, Vitt and Breitenbach 1993, Case 2002, Smith 1946, Linsdale 1932, Van Denburgh 1922, Nagy et al. 1999, Dunham et al. 1988, Brown and Nagy 2007, Warne and Charnov 2008, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Case 1975, Woolrich-Pina et al. 2011, Soule 1963, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006
Teiidae	<i>Aspidoscelis hyperythra</i>	no	NA	NA	NA	mainland	28	5.45	2.30	1.5	0.60	2.07	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Aspidoscelis inornata</i>	no	NA	NA	NA	mainland	30	4.58	2.65	1.5	0.82	3.25	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Congdon et al. 1978, Vitt et al. 1978, Walker 1981, Fitch 1985, Conant and Collins 1998, Stebbins 2003, Vitt and Breitenbach 1993, Schall 1993, Degenhardt et al. 1996, Greene 1982, Smith 1946, Winne and Beck 2004, Medica 1967, Milstead 1951, Rogner 1997b, Eifler and Passek 2000, Van Denburgh 1922, Dunham et al. 1988, Vitt and Price 1982, Warne and Charnov 2008, Manriquez-Moran et al. 2008, Radder et al. 2008, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Turner 1977, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Woolrich-Pina et al. 2011, Barbault and Maury 1981
Teiidae	<i>Aspidoscelis laredoensis</i>	no	NA	NA	NA	mainland	27	9.10	2.60	1.5	0.86	3.35	Conant and Collins 1998, Paulissen 1994, Paulissen et al. 1989, Uetz 2006, McKinney et al. 1973, Walker 1987, Kohler 2005, Siever and Paulissen 1996, Woolrich-Pina et al. 2011
Teiidae	<i>Aspidoscelis lineattissima</i>	no	NA	NA	NA	mainland	20	14.36	4.45	2.5	0.78	8.66	Fitch 1985, Ramirez-Bautista et al. 2000, Casas-Andreu and Gurrola-Hidalgo 1993, Duellman 1961, Kohler 2005, Duellman 1965, Rodriguez and Casas-Andreu 2011, Navarro-Garcia et al. 2008, Mata-Silva et al. 2010

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Aspidoscelis marmorata</i>	no	NA	NA	NA	mainland	30	15.37	2.60	1	1.14	2.95	Conant and Collins 1998, Lemos-Espinal and Smith 2007, Lemos-Espinal and Smith 2007b, Jones and Lovich 2009, Mata-Silva et al. 2010, Punzo 2001, Tinkle et al. 1970, Dunham and Miles 1985, Conant and Collins 1998, Stebbins 2003, Vitt and Breitenbach 1993, Degenhardt et al. 1996, Medica 1967, Dunham et al. 1988, Warne and Charnov 2008, Vitt and Breitenbach 1993, Manriquez-Moran et al. 2008, Turner 1977, Jones and Lovich 2009, Woolrich-Pina et al. 2011, Brennan and Holycross 2009
Teiidae	<i>Aspidoscelis neomexicana</i>	no	NA	NA	NA	mainland	33	8.75	1.95	1.5	1.01	2.96	Clobert et al. 1998, Duellman 1965, Jones and Lovich 2009, Castanet 1994, Barbault and Maury 1981
Teiidae	<i>Aspidoscelis scalaris</i>	no	NA	NA	NA	mainland	27	14.00	3.32	2	0.90	5.99	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, Conant and Collins 1998, Stebbins 2003, Vitt and Breitenbach 1993, Pianka and Vitt 2003, Degenhardt et al. 1996, Smith 1946, Rogner 1997b, Dunham et al. 1988, Clusella-Trullas et al. 2008, Warne and Charnov 2008, Cooper et al. 2001, Sinervo et al. 2010, Kohler 2005, Jones and Lovich 2009, Jensen et al. 2008, Taylor et al. 1999, Turner 1977, Beane et al. 2010, Fitch 1956, Shine and Charnov 1992, Brattstrom 1965, Pike et al.
Teiidae	<i>Aspidoscelis sexlineata</i>	no	NA	NA	NA	mainland	35	8.55	3.70	2	0.82	6.05	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source 2008
Teiidae	<i>Aspidoscelis sonorae</i>	no	NA	NA	NA	mainland	31	15.64	3.90	2.5	0.27	2.61	Congdon et al. 1978, Vitt et al. 1978, Stebbins 2003, Vitt and Breitenbach 1993, Degenhardt et al. 1996, Routman and Hulse 1984, Case 1983, Eifler and Passek 2000, Dunham et al. 1988, Vitt and Price 1982, Cooper et al. 2001, Lemos-Espinal and Smith 2007b, Kohler 2005, Jones and Lovich 2009, Brennan and Holycross 2009, Dunham and Miles 1985, Stebbins 2003, Vitt and Breitenbach 1993, Schall 1993, Degenhardt et al. 1996, Smith 1946, Winne and Beck 2004, Milstead 1951, Linsdale 1932, Eifler and Passek 2000, Van Denburgh 1922, Dunham et al. 1988, Vitt and Price 1982, Warne and Charnov 2008, Radder et al. 2008, Lemos-Espinal and Smith 2007, Price 1986, Schall 1978, Lemos-Espinal and Smith 2007b, Kohler 2005, Jones and Lovich 2009, Taylor et al. 1999, Woolrich-Pina et al. 2011, Mata-Silva et al. 2010, Punzo 2001, Taylor
Teiidae	<i>Aspidoscelis tessellata</i>	no	NA	NA	NA	mainland	33	13.10	4.07	1.5	1.93	11.76	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													et al. 2003, Brattstrom 1965, Walker et al. 2012, Mendoza-Quijano 2012
Teiidae	<i>Aspidoscelis tigris</i>	no	NA	NA	NA	mainland	34	15.58	2.70	2	0.74	4.00	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Fitch 1970, 1985, Radtkey et al. 1997, Anderson and Vitt 1990, Walker 1981, Perry and Garland 2002, Stebbins 2003, Cox et al. 2003, Grismer 2002, Ortega-Rubio et al. 1995, Pianka 1986, Vitt and Breitenbach 1993, Vitt and Breitenbach 1993, Schall 1993, Pianka and Vitt 2003, Case 2002, Degenhardt et al. 1996, Greene 1982, Winne and Beck 2004, Medina 1967, Milstead 1951, Rogner 1997b, Case 1983, Eifler and Passek 2000, Nagy et al. 1999, Dunham et al. 1988, Andrews and Pough 1980, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Vitt and Price 1982, Warne and Charnov 2008, Manriquez-Moran et al. 2008, Huey et al. 2001, Huey et al. 2001, Radder et al. 2008, Hardy and McDiarmid 1969, Schall 1978, Sinervo et al. 2010,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Aspidoscelis uniparens</i>	no	NA	NA	NA	mainland	32	7.10	3.22	2	0.60	3.86	Kohler 2005, Jones and Lovich 2009, Case 1975, Huey and Pianka 2007, van Berkum 1988, Turner 1977, Krekorian 1983, Morton and James 1988, Bury 1982, Woolrich-Pina et al. 2011, Brennan and Holycross 2009, Barbault and Maury 1981, Cunningham 1966, Shine and Charnov 1992, Brattstrom 1965, Lemos-Espinal et al. 1997, Pike et al. 2008, Stebbins and McGinnis 2012, Lemm 2006, St. John 2002 Congdon et al. 1978, Vitt et al. 1978, Perry and Garland 2002, Conant and Collins 1998, Stebbins 2003, Vitt and Breitenbach 1993, Cuellar 1993, Degenhardt et al. 1996, Eifler and Passek 2000, Dunham et al. 1988, Vitt and Price 1982, Warne and Charnov 2008, Cooper et al. 2001, Radder et al. 2008, Cuellar 1984, Lemos-Espinal and Smith 2007b, Jones and Lovich 2009, Woolrich-Pina et al. 2011, Brennan and Holycross 2009, Shine and Charnov 1992
Teiidae	<i>Aspidoscelis velox</i>	no	NA	NA	NA	mainland	36	11.65	4.00	1	1.14	4.54	Fitch 1970, Stebbins 2003, Degenhardt et al. 1996, Bowker et al. 1986, Vitt and Price 1982, Radder et al. 2008, Stuart 1998, Jones and Lovich 2009, Woolrich-Pina et al. 2011, Brennan and Holycross 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Cnemidophorus arubensis</i>	yes	Lesser Antilles	Aruba	181.10	Oceanic	13	14.05	1.00	2	NA	NA	Fitch 1985, Zug et al. 2001, Cooper and Vitt 2002, Vitt and de Carvalho 1992, Vitt and Breitenbach 1993, Pianka and Vitt 2003, Paulissen and Walker 1994, van Buurt 2005, Schall 1983, Sinervo et al. 2010, Kohler 2005, Schall 1973
Teiidae	<i>Cnemidophorus cryptus</i>	no	NA	NA	NA	mainland	-2	7.93	1.50	2	0.78	2.33	Avila-Pires 1995, Vitt et al. 1997, Mesquita and Colli 2003, Huey et al. 2001, Sinervo et al. 2010, Rocha et al. 2009
Teiidae	<i>Cnemidophorus gramivagus</i>	no	NA	NA	NA	mainland	5	8.16	1.54	1	0.86	1.32	Avila-Pires 1995, Mesquita and Colli 2003, Sinervo et al. 2010, Rocha et al. 2009, Mojica et al. 2003, Fitch 1985, Perry and Garland 2002, Stafford and Meyer 2000, Conant and Collins 1998, Savage 2002, Lee 2000, Cox et al. 2003, Avila-Pires 1995, Cooper and Vitt 2002, Vitt and de Carvalho 1992, Vitt and de Carvalho 1995, Beebe 1945, Kohler 2003, Vitt and Breitenbach 1993, Pianka and Vitt 2003, Vitt et al. 1997, Schwartz and Henderson 1991, Mesquita and Colli 2003, Hoogmoed 1973, Vitt 2000, Molina et al. 2004, Montgomery et al. 2007, Kohler 1996, McCranie and Castaneda 2005, McCranie et al. 2005, van Buurt 2005, Janzen 1973, Lotzkat 2007, Murphy 1997, Vitt et al. 1999, Huey et al. 2001, Radder et al. 2008,
Teiidae	<i>Cnemidophorus lemniscatus</i>	no	NA	NA	NA	mainland	6	7.47	2.05	2	0.45	1.85	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Kohler 2008, Fuenmayor et al. 2005, Test et al. 1966, Sinervo et al. 2010, Kohler 2005, Ugueto and Rivas 2010, Montgomery et al. 2011, Gasc 1990, Rocha et al. 2009, Van Wilgen and Richardson 2012, Brattstrom 1965, Schall 1973
Teiidae	<i>Cnemidophorus murinus</i>	yes	Lesser Antilles	Curacao	443.10	Oceanic	12	27.55	1.18	NA	2.00	NA	Cox et al. 2003, Olesen and Valido 2003, Cooper and Vitt 2002, Dearing and Schall 1994, Pianka and Vitt 2003, Paulissen and Walker 1994, Alvarez 2004, Bennett and Gorman 1979, van Buurt 2005, Andrews and Pough 1980, Sinervo et al. 2010, Kohler 2005, Damuth 1987, Schall and Dearing 1994, Schwartz and Henderson 1991, Malhotra and Thorpe 1999, Funk & Fa 2006, Dickinson and Fa 2000, Henderson and Powell 2009, Daltry 2009, Corke 1987, Cei 1986, Cei 1993, Achaval and Olmos 2003, Milstead 1961, Carreira et al. 2005, Balestrin 2008, Balestrin et al. 2010, Menezes and Rocha 2011, Ariani et al. 2012, Feltrim and De Lema 2000, Rezende-Pinto et al. 2009, Caruccio et al.
Teiidae	<i>Cnemidophorus vanzoi</i>	yes	Lesser Antilles	Maria Major	0.25	Oceanic	14	19.83	NA	NA	1.86	NA	
Teiidae	<i>Contomastix lacertoides</i>	no	NA	NA	NA	mainland	-18	6.74	3.75	1	0.48	1.78	
Teiidae	<i>Contomastix vacariensis</i>	no	NA	NA	NA	mainland	-28	9.16	4.15	2	0.64	5.33	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Holcosus festivus</i>	no	NA	NA	NA	mainland	11	19.55	2.75	3	1.35	11.13	Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1973, 1985, Campbell 1999, Stafford and Meyer 2000, Savage 2002, Lee 2000, Cox et al. 2003, Vitt and de Carvalho 1992, Kohler 2003, Rand and Myers 1990, Duellman 1990, McCranie and Castaneda 2005, Guyer and Donnelly 2005, Toledo et al. 2007, Hillman 1969, Dunham et al. 1988, Vitt and Zani 1996, Vitt and Zani 1998, Huey et al. 2001, Kohler 2008, Echternacht 1983, Somma and Brooks 1976, Smith 1968, Sinervo et al. 2010, Kohler 2005, Duellman 1963, van Berkum 1988, Hirth 1965, Sebastian-Gonzalez and Gomez 2012
Teiidae	<i>Holcosus quadrilineatus</i>	no	NA	NA	NA	mainland	9	9.39	2.05	2.5	0.70	3.60	Tinkle et al. 1970, Dunham and Miles 1985, Fitch 1970, 1973, Perry and Garland 2002, Savage 2002, Vitt and de Carvalho 1992, Kohler 2003, Duellman 1990, Hillman 1969, Dunham et al. 1988, Somma and Brooks 1976, Smith 1968, Sinervo et al. 2010, Turner 1977, Damuth 1987, Curry-Lindahl 1979, Tinkle et al. 1967, Hirth 1963, Brattstrom 1965, Hirth 1965, Sebastian-Gonzalez and Gomez 2012

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Kentropyx calcarata</i>	no	NA	NA	NA	mainland	-5	22.45	4.50	1	0.85	3.83	Fitch 1970, Vitt 1991, Cox et al. 2003, Pianka and Vitt 2003, Avila-Pires 1995, Martins 1991, Vitt and de Carvalho 1992, Beebe 1945, Zimmerman and Rodrigues 1990, Duellman 1990, Pianka and Vitt 2003, Dixon and Soini 1986, Hoogmoed 1973, Vitt 2000, Molina et al. 2004, Gallagher and Dixon 1992, Gasnier et al. 1994, Vitt et al. 1997, Vitt et al. 1999, Huey et al. 2001, Beebe 1945, Werneck et al. 2009, Sinervo et al. 2010, Kohler 2005, Avila-Pires et al. 2010, Gasc 1990, Fitch 1968, Rocha et al. 2009
Teiidae	<i>Kentropyx pelviceps</i>	no	NA	NA	NA	mainland	-5	32.28	5.17	1.5	1.14	8.80	Cox et al. 2003, Duellman and Mendelson 1995, Pianka and Vitt 2003, Avila-Pires 1995, Vitt and de Carvalho 1992, Duellman 1978, Duellman 1990, Pianka and Vitt 2003, Dixon and Soini 1986, Gallagher and Dixon 1992, Duellman 2005, Bartlett and Bartlett 2003, Vitt et al. 1995, Vitt and Zani 1996b, Vitt et al. 2000, Huey et al. 2001, Werneck et al. 2009, Sinervo et al. 2010, Kohler 2005, Duellman 1987 Perry and Garland 2002, Cox et al. 2003, Pianka and Vitt 2003, Avila-Pires 1995, Vitt and de Carvalho 1992, Vitt and de Carvalho 1995, Dixon and Soini 1986, Hoogmoed 1973, Molina et al. 2004,
Teiidae	<i>Kentropyx striata</i>	no	NA	NA	NA	mainland	4	15.46	4.92	2	0.51	4.98	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Teiidae	<i>Teius oculatus</i>	no	NA	NA	NA	mainland	-34	23.40	5.00	1	1.14	5.68	Gallagher and Dixon 1992, Murphy 1997, Werneck et al. 2009, Sinervo et al. 2010, Kohler 2005, Avila-Pires et al. 2010, Rocha et al. 2009, Martins 2006
Teiidae	<i>Teius teyou</i>	no	NA	NA	NA	mainland	-26	29.99	4.80	1	1.58	7.56	Cei 1986, Cei 1993, Avila 2002, Scolaro 2006, Carreira et al. 2005, Kohler 2005, Cappellari et al. 2007, Cappellari et al. 2011, Winck and Rocha 2012, Acosta and Martori 1990
Teiidae	<i>Tupinambis teguixin</i>	no	NA	NA	NA	mainland	-6	1290.98	11.95	1	14.85	177.41	Fitzgerald et al. 1999, Cei 1986, Cei 1993, Alvarez 2004, Avila 2002, Warne and Charnov 2008, Milstead 1961, Cruz et al. 1999, Kohler 2005, Shine and Wall 2008, Fitch 1970, Anderson and Vitt 1990, Fitzgerald et al. 1991, Zug et al. 2001, Pianka and Vitt 2003, Avila-Pires 1995, Martins 1991, Vitt and de Carvalho 1995, Duellman 1978, Beebe 1945, Rodrigues and Cadle 1990, Zimmerman and Rodrigues 1990, Duellman 1990, Pianka and Vitt 2003, Cei 1986, Cei 1993, Dixon and Soini 1986, Hoogmoed 1973, Rogner 1997b, Vitt and Zani 1998, Duellman 2005, Bartlett and Bartlett 2003, Nagy et al. 1999, Toledo et al. 2007, Rodrigues 1996, Brown and Nagy 2007, Lotzkat 2007, Murphy 1997, Vitt and Zani

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													1996b, Huey et al. 2001, Milstead 1961, Fuenmayor et al. 2005, Fitch 1982, McElroy et al. 2008, Vanzolini et al. 1980, Kohler 2005, Ugueto and Rivas 2010, Duellman 1987, Gasc 1990, Van Wilgen and Richardson 2012
Trogonophidae	<i>Trogonophis wiegmanni</i>	no	NA	NA	NA	mainland	34	2.87	5.00	0.75	0.39	1.44	Wiens et al. 2006, Andrade et al. 2006, Civantos et al. 2003, Schleich et al. 1996, Loveridge 1941, Sinervo et al. 2010, Rosler 2005, Lopez et al. 2002, Avery 1982, Martin et al. 2011, Martin et al. 2011b, Kamel and Gatten 1983, Honegger 1969, Tinkle et al. 1970, Fitch 1970, 1982, 1985, Perry and Garland 2002, Cox et al. 2003, Van Denburgh and Slevin 1913, Nagy et al. 1999, Dunham et al. 1988, Brown and Nagy 2007, Fitch 1982, Sinervo et al. 2010, Kohler 2005, Vitt and Goldberg 1983, Goldberg and Rodriguez 1986, Turner 1977, Curry-Lindahl 1979, Huey 1974
Tropiduridae	<i>Microlophus albemarlensis</i>	yes	Galapagos Archipelago	Isabela	4588.00	Oceanic	-1	13.76	2.23	NA	NA	NA	Donoso-Barros 1966, Ortiz-Zapata 1980, Sinervo et al. 2010, Ibarquengoytia 2008, Vidal and Labra 2008, Labra et al. 2008, Vidal et al. 2002
Tropiduridae	<i>Microlophus atacamensis</i>	no	NA	NA	NA	mainland	-25	20.88	4.50	1	1.91	8.58	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Tropiduridae	<i>Microlophus delanonis</i>	yes	Galapagos Archipelago	Espanola	60.00	Oceanic	-1	26.40	3.47	NA	NA	NA	Fitch 1985, Perry and Garland 2002, Cox et al. 2003, Van Denburgh and Slevin 1913, Kohler 2005
Tropiduridae	<i>Microlophus habelii</i>	yes	Galapagos Archipelago	Marchena	130.00	Oceanic	0	22.59	2.50	NA	NA	NA	Cox et al. 2003, Van Denburgh and Slevin 1913
Tropiduridae	<i>Microlophus quadrivittatus</i>	no	NA	NA	NA	mainland	-21	18.04	3.16	1.5	2.37	11.21	Donoso-Barros 1966, Ortiz-Zapata 1980, Sinervo et al. 2010, Kohler 2005, Goldberg and Rodriguez 1986, Iburguengoytia 2008, Vidal and Labra 2008
Tropiduridae	<i>Plica plica</i>	no	NA	NA	NA	mainland	-3	57.37	2.51	2	2.89	14.49	Fitch 1970, Cox et al. 2003, Duellman and Mendelson 1995, Avila-Pires 1995, Rodrigues and Cadle 1990, Zimmerman and Rodrigues 1990, Duellman 1990, Pianka and Vitt 2003, Beebe 1944b, Dixon and Soini 1986, Hoogmoed 1973, Vitt 2000, Molina et al. 2004, Rogner 1997a, Duellman 2005, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Bartlett and Bartlett 2003, Lotzkat 2007, Murphy 1997, Vitt 1991, Vitt et al. 1997, Vitt et al. 1999, Huey et al. 2001, Radder et al. 2008, Sinervo et al. 2010, Kohler 2005, Avila-Pires et al. 2010, Grizante et al. 2011, Brandt and Navas 2011, Duellman 1987, Gasc 1990

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Tropiduridae	<i>Plica umbra</i>	no	NA	NA	NA	mainland	-4	18.84	2.05	1.5	1.63	5.02	Fitch 1970, Cox et al. 2003, Duellman and Mendelson 1995, Avila-Pires 1995, Duellman 1978, Rodrigues and Cadle 1990, Zimmerman and Rodrigues 1990, Duellman 1990, Pianka and Vitt 2003, Beebe 1944b, Dixon and Soini 1986, Hoogmoed 1973, Vitt 2000, Molina et al. 2004, Duellman 2005, Harvey and Gutberlet 1998, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Bartlett and Bartlett 2003, Gasnier et al. 1994, Vitt and Zani 1996b, Vitt et al. 1997, Vitt et al. 1999, Huey et al. 2001, Sinervo et al. 2010, Kohler 2005, Rand 1982, Avila-Pires et al. 2010, Grizante et al. 2011, Brandt and Navas 2011, Duellman 1987, Gasc 1990, Rocha et al. 2009
Tropiduridae	<i>Stenocercus chrysopygus</i>	no	NA	NA	NA	mainland	-10	8.50	5.50	1	0.58	3.18	Uetz 2006, Torres-Carvajal 2007, Torres-Carvajal 2007b, Kohler 2005
Tropiduridae	<i>Stenocercus dumerilii</i>	no	NA	NA	NA	mainland	-2	39.76	4.00	1	1.28	5.10	Avila-Pires 1995, Torres-Carvajal 2007, Torres-Carvajal 2007b, Kohler 2005
Tropiduridae	<i>Tropidurus etheridgei</i>	no	NA	NA	NA	mainland	-21	9.78	5.58	2	0.68	7.62	Fitzgerald et al. 1999, Cei 1986, Cei 1993, Vitt 1991, Vitt et al. 1997, Van Sluys et al. 2010, Kohler 2005, Dutra et al. 2011, Grizante et al. 2011, Brandt and Navas 2011, Cruz 1997, Winck and Rocha 2012

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Tropiduridae	<i>Tropidurus hispidus</i>	no	NA	NA	NA	mainland	4	15.74	6.50	2.5	1.07	17.35	Cox et al. 2003, Avila-Pires 1995, Cooper and Vitt 2002, Vitt and de Carvalho 1995, Pianka and Vitt 2003, Hoogmoed 1973, Vitt and Zani 1998, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Kohlsdorf and Navas 2007, Van Sluys et al. 2004, Rodrigues 2003, Rodrigues 1996, Vitt et al. 1996, Vitt et al. 1997, Vitt 1995, Huey et al. 2001, Fuenmayor et al. 2005, Van Sluys et al. 2010, Kolodiuk et al. 2010, Sinervo et al. 2010, Kohler 2005, Vitt and Goldberg 1983, Goldberg and Rodriguez 1986, Ugueto and Rivas 2010, Dutra et al. 2011, Grizante et al. 2011, Brandt and Navas 2011, Rocha et al. 2009, Ribeiro and Freire 2011
Tropiduridae	<i>Tropidurus semitaeniatus</i>	no	NA	NA	NA	mainland	-9	11.49	2.00	3.5	0.27	1.86	Cox et al. 2003, Cooper and Vitt 2002, Pianka and Vitt 2003, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Rodrigues 2003, Rodrigues 1996, Vitt and Price 1982, Fitch 1982, Dunham et al. 1988, Warne and Charnov 2008, Vitt et al. 1997, Vitt 1995, Vanzolini et al. 1980, Van Sluys et al. 2010, Kolodiuk et al. 2010, Sinervo et al. 2010, Vitt and Goldberg 1983, Dutra et al. 2011, Grizante et al. 2011, Brandt and Navas 2011, Vitt 1981, Rocha et al. 2009, Ribeiro and Freire 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Tropiduridae	<i>Tropidurus spinulosus</i>	no	NA	NA	NA	mainland	-18	22.28	4.90	1.5	1.51	11.08	Fitzgerald et al. 1999, Cei 1986, Cei 1993, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Vitt 1991, Vitt et al. 1997, Van Sluys et al. 2010, Kohler 2005, Dutra et al. 2011, Grizante et al. 2011, Brandt and Navas 2011, Cruz et al. 1997, Cruz 1998, Cox et al. 2003, Cooper and Vitt 2002, Cei 1993, Beebe 1944b, Nogueira et al. 2005, Achaval and Olmos 2003, Rogner 1997a, Alvarez 2004, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Kohlsdorf and Navas 2007, Colli et al. 2002, Dunham et al. 1988, Vitt and Price 1982, Fitch 1982, Warne and Charnov 2008, Radder et al. 2008, Carreira et al. 2005, McElroy et al. 2008, Vanzolini et al. 1980, Van Sluys et al. 2010, Sinervo et al. 2010, Rand 1982, Vitt and Goldberg 1983, Kiefer et al. 2008, Dutra et al. 2011, Grizante et al. 2011, Brandt and Navas 2011, Vitt 1981, Gasc 1990, Rocha et al. 2009, Winck and Rocha 2012, Cox et al. 2003, Avila-Pires 1995, Duellman 1978, Duellman 1990, Pianka and Vitt 2003, Dixon and Soini 1986, Kohlsdorf et al. 2001, Kohlsdorf and Navas 2006, Bartlett and Bartlett 2003, Vitt 1991, Vitt and Zani 1996b, Vitt and Zani 1996c,
Tropiduridae	<i>Tropidurus torquatus</i>	no	NA	NA	NA	mainland	-9	12.10	4.71	2	0.65	6.13	
Tropiduridae	<i>Uracentron flaviceps</i>	no	NA	NA	NA	mainland	-2	22.84	2.00	1.5	1.63	4.90	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													Vitt et al. 1997, Huey et al. 2001, Sinervo et al. 2010, Kohler 2005, Rand 1982, Grizante et al. 2011, Brandt and Navas 2011
Varanidae	<i>Varanus acanthurus</i>	no	NA	NA	NA	mainland	-21	51.89	9.30	1	3.29	30.63	Pianka 1995, Thompson and Pianka 2001, Cogger 2000, James et al. 1992, Pianka and King 2004, Wilson and Swan 2003, Rogner 1997b, Greer 1989, De Lisle 1996, Nagy et al. 1999, Sweet and Pianka 2007, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Garrick 2008, Losos and Greene 1988, Horn and Visser 1997, Eidenmueller and Philippen 2008, Thompson et al. 2008, Kohler 2005, Frydlova and Frynta 2010, Clemente et al. 2009, Swanson 2007
Varanidae	<i>Varanus albigularis</i>	no	NA	NA	NA	mainland	-10	3023.67	32.00	1	25.00	800.04	Pianka 1995, Thompson and Pianka 2001, Spawls et al. 2002, FitzSimons 1943, Perry and Garland 2002, Branch 1998, Jeffery 1993, De Lisle 1996, Auerbach 1987, Branch 2005, Graham and Marais 2007, Andrews and Pough 1980, Meik et al. 2002, Phillips and Millar 1998, Horn and Visser 1997, Eidenmueller and Philippen 2008, Kohler 2005, Frydlova and Frynta 2010, Largen and Spawls 2010, Pienaar 1966, Phillips 1995, King and Green 1993, Jacobsen 1982

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus beccarii</i>	yes	Aru Islands	Trangan	2148.50	Land bridge	-6	421.27	3.80	1.83	7.03	48.86	Pianka 1995, Thompson and Pianka 2001, Badger 2003, Sweet and Pianka 2007, Horn and Visser 1997, Eidenmueller and Philippen 2008, Kohler 2005, Frydlova and Frynta 2010, Mendyk 2011 Fitch 1970, Pianka 1995, Thompson and Pianka 2001, Anderson 1999, Perry and Garland 2002, Schleich and Kastle 2002, Zug et al. 2001, Pianka and King 2004, Daniel 1983, Tikader and Sharma 1992, Taylor 1963, Pianka and Vitt 2003, Rogner 1997b, Cox et al. 1998, Anderson and Leviton 1969, De Lisle 1996, Khan 2006, Deraniyagala 1953, Clark 1990, Pauwels et al. 2003, Shrestha 2001, Andrews and Pough 1980, Ahsan and Abu Saeed 2004, Sweet and Pianka 2007, Brown and Nagy 2007, Losos and Greene 1988, Auffenberg et al. 1989, Horn and Visser 1997, Somaweera and Somaweera 2009, Eidenmueller and Philippen 2008, de Buffrenil and Rimblot-Baly 1999, Das 2002, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Das 2010, Cox et al. 2010, Das and de Silva 2011, Ahmed et al. 2009, King and Green 1993, Chandramouli and Ganesh 2011, Rathnayake et al. 2003,
Varanidae	<i>Varanus bengalensis</i>	no	NA	NA	NA	mainland	21	1180.48	20.20	1	9.40	189.97	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus boehmei</i>	yes	New Guinea	Waigeo	3153.70	Oceanic	0	352.73	4.00	NA	NA	NA	Jacobs 2003, Philipp and Philipp 2007, Eidenmueller and Philippen 2008, Frydlova and Frynta 2010
Varanidae	<i>Varanus caerulivirens</i>	yes	Maluku Islands	Halmahera	18039.60	Oceanic	1	829.83	2.00	NA	NA	NA	Pianka and King 2004, Philipp et al. 2007, Eidenmueller and Philippen 2008, Philipp et al. 2007, Weijola 2010, Frydlova and Frynta 2010, Setiadi and Hamidy 2006
Varanidae	<i>Varanus caudolineatus</i>	no	NA	NA	NA	mainland	-26	13.57	3.70	1	1.37	5.08	Pianka 1994, 1995, Thompson and Pianka 2001, Cogger 2000, James et al. 1992, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997b, Greer 1989, De Lisle 1996, Nagy et al. 1999, Philipp et al. 2007, Sweet and Pianka 2007, Brown and Nagy 2007, Eidenmueller and Philippen 2008, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2010, Clemente et al. 2009, King and Green 1993, Rathnayake et al. 2003
Varanidae	<i>Varanus cerambonensis</i>	yes	Maluku Islands	Seram	17454.20	Oceanic	-3	1405.80	NA	NA	13.82	NA	Pianka and King 2004, Eidenmueller and Philippen 2008, Philipp et al. 2007, Pianka 1995, Thompson and Pianka 2001, Perry and Garland 2002, Schleich and Kastle 2002, Pianka and King 2004, Daniel 1983, Tikader and Sharma 1992, Taylor 1963, Rogner 1997b, De Lisle 1996, Khan 2006, Shrestha 2001, Sweet and Pianka 2007, Sweet and Pianka 2007, Eidenmueller and Philippen 2008, Das 2002, Kohler 2005, Frydlova and Frynta 2010, Ahmed et al. 2009, Ahmed 2009
Varanidae	<i>Varanus flavescens</i>	no	NA	NA	NA	mainland	27	717.34	12.25	1	3.57	43.76	Pianka 1994, 1995, Thompson and Pianka 2001, Cogger 2000, Perry and Garland 2002, James et al. 1992, Pianka and King 2004, King et al. 1989, Wilson and Swan 2003, Rogner 1997b, Greer 1989, De Lisle 1996, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Losos and Greene 1988, Horn and Visser 1997, Eidenmueller and Philippen 2008, Heger and Heger 2007, Thompson et
Varanidae	<i>Varanus giganteus</i>	no	NA	NA	NA	mainland	-25	4072.03	9.00	1	55.02	495.19	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus gilleni</i>	no	NA	NA	NA	mainland	-25	26.83	4.28	1	2.53	10.83	al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Clemente et al. 2009, Heatwole and Pianka 1993, King and Green 1993, Heatwole and Taylor 1987, Swanson 2007, Rathnayake et al. 2003, Moro and MacAulay 2010 Pianka 1982, 1995, Thompson and Pianka 2001, Cox et al. 2003, James et al. 1992, Pianka and King 2004, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, De Lisle 1996, Sweet and Pianka 2007, Clusella-Trullas et al. 2008, Losos and Greene 1988, Horn and Visser 1997, Eidenmueller and Philippen 2008, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Clemente et al. 2009, King and Green 1993, Rathnayake et al. 2003 Pianka 1995, Cogger 2000, James et al. 1992, Wilson and Swan 2003, Sweet and Pianka 2007, Wilson and Swan 2008, Losos and Greene 1988, Eidenmueller and Philippen 2008, Thompson et al. 2008, Frydlova and Frynta 2010, Clemente et al. 2009, de Zeeuw 2010
Varanidae	<i>Varanus glauerti</i>	no	NA	NA	NA	mainland	-16	97.31	4.50	4	3.31	59.59	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus gouldii</i>	no	NA	NA	NA	mainland	-25	501.49	6.55	1	4.04	26.46	Pianka 1982, 1994, 1995, Cogger 2000, Shine 1986, Perry and Garland 2002, Cox et al. 2003, James et al. 1992, Pianka and King 2004, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997b, Greer 1989, De Lisle 1996, Pengilley 1981, Sweet and Pianka 2007, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Garrick 2008, Losos and Greene 1988, Huey et al. 2001, Chapman and Dell 1985, Horn and Visser 1997, Henle 1989c, Bustard 1978, Eidenmueller and Philippen 2008, Thompson et al. 2008, Sinervo et al. 2010, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Bush et al. 2010, Sutherland 2011, Huey and Pianka 2007, Michael and Lindenmayer 2010, Clemente et al. 2009, Wilson 2003, Heatwole and Pianka 1993, King and Green 1993, Christian and Weavers 1996, Bartholomew and Tucker 1964, Heatwole and Taylor 1987, Swanson 2007, Bustard 1968, Light et al. 1966, Rathnayake et al. 2003, Gordon et al. 2010, Swan and Watharow 2005

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus griseus</i>	no	NA	NA	NA	mainland	28	887.15	15.75	0.665	12.08	126.52	Pianka 1995, Thompson and Pianka 2001, Smith 1935, Schleich et al. 1996, Amitai and Bouskila 2001, Szczerbak 2003, Anderson 1999, Minton 1999, Perry and Garland 2002, Disi et al. 2001, Baran and Atatur 1998, Pianka and King 2004, Tikader and Sharma 1992, Geniez et al. 2004, Flower 1933, Pianka and Vitt 2003, Arnold 1984, El Din 2006, Rogner 1997b, De Lisle 1996, Schatti and Desvoignes 1999, Le Berre 1989, Khan 2006, Anderson 1963, Murthy 1995, Jongbloed 2000, Sweet and Pianka 2007, Losos and Greene 1988, Werner 1987, Hornby 1996, Horn and Visser 1997, Arbel 1984, Eidenmueller and Philippen 2008, Das 2002, Kohler 2005, Frydlova and Frynta 2010, King and Green 1993, van der Kooij 2001, Castanet 1994, Rathnayake et al. 2003, Bar and Haimovitch 2012, Trape et al. 2012, Fathinia et al. 2009
Varanidae	<i>Varanus kingorum</i>	no	NA	NA	NA	mainland	-16	11.27	4.50	8	1.04	37.31	Pianka 1995, James et al. 1992, Pianka and King 2004, Wilson and Swan 2003, Greer 1989, Sweet and Pianka 2007, Wilson and Swan 2008, Eidenmueller and Philippen 2008, Thompson et al. 2008, Kohler 2005, Mendyk 2011, Clemente et al. 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus komodoensis</i>	yes	Lesser Sunda Islands	Flores	14154.40	Oceanic	-9	32879.93	18.30	1.75	82.13	2630.22	Pianka 1995, Thompson and Pianka 2001, Perry and Garland 2002, Pianka and King 2004, Pianka and Vitt 2003, de Rooij 1915, De Lisle 1996, Sweet and Pianka 2007, Brown and Nagy 2007, Horn and Visser 1997, Auffenberg 1981, Eidenmueller and Philippen 2008, de Buffrenil and Rimblot-Baly 1999, Sinervo et al. 2010, Kohler 2005, De Magalhaes and Costa 2009, Frydlova and Frynta 2010, King and Green 1993, Auffenberg 1980, Ciofi and De Boer 2004, Damuth 1987, Harlow et al. 2010, Rathnayake et al. 2003, McNab and Auffenberg 1976
Varanidae	<i>Varanus lirungensis</i>	yes	Talaud Islands	Salibabu	86.87	Oceanic	4	810.67	2.50	NA	NA	NA	Koch et al. 2009
Varanidae	<i>Varanus mabitang</i>	yes	Philippine Islands	Panay	12011.10	Oceanic	12	3191.96	9.00	1	NA	NA	Pianka and King 2004, Sweet and Pianka 2007, Reyes et al. 2008, Philipp and Philipp 2007, Gaulke and Curio 2001, Gaulke et al. 2002, Eidenmueller and Philippen 2008, Gaulke et al. 2007, Gaulke 2011
Varanidae	<i>Varanus macraei</i>	yes	New Guinea	Batanta	455.90	Land bridge	-1	446.94	4.03	NA	10.31	NA	Pianka and King 2004, Sweet and Pianka 2007, Philipp and Philippen 2007, Eidenmueller and Philippen 2008, Ziegler et al. 2009, Frydlova and Frynta 2010
Varanidae	<i>Varanus melinus</i>	yes	Maluku Islands	Taliabu	2913.20	Oceanic	-2	1108.15	4.90	2.5	21.60	264.62	Pianka and King 2004, Philipp et al. 2007, Sweet and Pianka 2007, Eidenmueller and Philippen 2008, Kohler 2005, Frydlova and Frynta

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
													2010, Mendyk 2011, Ziegler et al. 2010
Varanidae	<i>Varanus mertensi</i>	no	NA	NA	NA	mainland	-17	981.43	9.70	2.5	15.49	375.52	Pianka 1995, Thompson and Pianka 2001, Shine 1986, Zug et al. 2001, James et al. 1992, Pianka and King 2004, Pianka and Vitt 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, De Lisle 1996, Nagy et al. 1999, Sweet and Pianka 2007, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Wilson and Swan 2008, Horn and Visser 1997, Eidenmueller and Philippen 2008, Mayes et al. 2007, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Mendyk 2011, Clemente et al. 2009, King and Green 1993, Christian and Weavers 1996, Swanson 2007, Rathnayake et al. 2003

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus niloticus</i>	no	NA	NA	NA	mainland	-1	2686.48	23.38	1	17.28	404.01	Fitch 1970, Pianka 1995, Thompson and Pianka 2001, Spawls et al. 2002, FitzSimons 1943, Loveridge 1936, Pianka and King 2004, Greenbaum and Carr 2005, Flower 1933, Pianka and Vitt 2003, Broadley 1971, Hughes 1988, Dunger 1967c, Jeffery 1993, El Din 2006, Schmidt et al. 1919, Rogner 1997b, De Lisle 1996, Le Berre 1989, Loveridge 1953, Auerbach 1987, Barbour and Loveridge 1928, Branch 2005, Chirio and LeBreton 2007, Bauer and Jackman 2008, Graham and Marais 2007, Largen and Spawls 2006, Sweet and Pianka 2007, Leache et al. 2006, Losos and Greene 1988, Reid 1986, Horn and Visser 1997, Eidenmueller and Philippen 2008, de Buffrenil and Rimblot-Baly 1999, Chirio 2009, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Largen and Spawls 2010, Pienaar 1966, King and Green 1993, Bowker 1984, Western 1974, Castanet 1994, Van Wilgen and Richardson 2012, Trape et al. 2012
Varanidae	<i>Varanus nuchalis</i>	yes	Philippine Islands	Negros	13074.50	Oceanic	11	2166.42	10.50	NA	NA	NA	Pianka and King 2004, Koch et al. 2007, Eidenmueller and Philippen 2008, Koch et al. 2010, Gaulke 2011

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus ornatus</i>	no	NA	NA	NA	mainland	6	3056.8 3	18.00	1	24.86	447.55	Spawls et al. 2002, Pianka and King 2004, Luiselli et al. 1999, Chirio and LeBreton 2007, Bauer and Jackman 2008, Sweet and Pianka 2007, Pauwels and Vande weghe 2008, Pauwels et al. 2004, Eidenmueller and Philippen 2008, Hennessy 2010, Mendyk 2011, Trape et al. 2012
Varanidae	<i>Varanus panoptes</i>	no	NA	NA	NA	mainland	-21	1307.1 1	8.80	1	20.09	176.76	Pianka 1995, Thompson and Pianka 2001, Cogger 2000, Shine 1986, James et al. 1992, Pianka and King 2004, Pianka and Vitt 2003, Wilson and Swan 2003, Allison 2006, Greer 1989, Sweet and Pianka 2007, Paden 2008, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Philipp and Philipp 2007, Eidenmueller and Philippen 2008, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Mendyk 2011, Clemente et al. 2009, Christian and Weavers 1996, Swanson 2007, Rathnayake et al. 2003
Varanidae	<i>Varanus prasinus</i>	no	NA	NA	NA	mainland	-5	269.30	4.35	2.375	9.23	95.37	Pianka 1995, Thompson and Pianka 2001, Zweifel 1980, Pianka and King 2004, Pianka and Vitt 2003, Wilson and Swan 2003, Allison 2006, Rogner 1997b, Greer 1989, De Lisle 1996, Sprackland 1991, Sweet and Pianka 2007, Allison 2007, Philipp and Philipp 2007, Losos and

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus rosenbergi</i>	no	NA	NA	NA	mainland	-33	638.14	7.47	1	4.04	30.16	Greene 1988, Horn and Visser 1997, Eidenmueller and Philippen 2008, Kohler 2005, Mendyk 2008, Frydlova and Frynta 2010, Mendyk 2011 Pianka 1995, Thompson and Pianka 2001, Cogger 2000, Perry and Garland 2002, James et al. 1992, Pianka and King 2004, Wilson and Swan 2003, Greer 1989, De Lisle 1996, Nagy et al. 1999, Sweet and Pianka 2007, Brown and Nagy 2007, Eidenmueller and Philippen 2008, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Bush et al. 2010, Sutherland 2011, Michael and Lindenmayer 2010, Clemente et al. 2009, King and Green 1993, Christian and Weavers 1996, Swanson 2007, Rathnayake et al. 2003, Swan and Watharow 2005 Pianka 1995, Thompson and Pianka 2001, Pianka and King 2004, Taylor 1963, Das 2004, Manthey and Grossmann 1997, Rogner 1997b, Cox et al. 1998, De Lisle 1996, Sweet and Pianka 2007, Losos and Greene 1988, Horn and Visser 1997, Eidenmueller and Philippen 2008, Kohler 2005, Das 2010, Das 2011, Grismer 2011, Pauwels et al. 2009
Varanidae	<i>Varanus rudicollis</i>	no	NA	NA	NA	mainland	3	744.23	9.20	2	14.76	271.53	

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus salvator</i>	no	NA	NA	NA	mainland	11	2683.88	9.00	2	26.62	479.10	Pianka 1995, Thompson and Pianka 2001, Smith 1935, Daniel 1983, Pianka and King 2004, Tikader and Sharma 1992, Taylor 1963, Das 2004, Anderson 1889, Erdelen 1998, Schmidt 1927, Manthey and Grossmann 1997, de Rooij 1915, Rogner 1997b, Ziegler 2002, Cox et al. 1998, Brown et al. 1996, De Lisle 1996, Taylor 1922, Deraniyagala 1953, Karsen et al. 1986, Malkmus et al. 2002, Murthy 1995, Pauwels et al. 2003, Bauer and Jackman 2008, Sweet and Pianka 2007, Brown and Nagy 2007, Auffenberg et al. 1989, Horn and Visser 1997, Somaweera and Somaweera 2009, Eidenmueller and Philippen 2008, Kirshner 2007, de Buffrenil and Rimblot-Baly 1999, Das 2002, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Das 2010, Cox et al. 2010, Alcalá 1986, Das 2011, Das and de Silva 2011, Ahmed et al. 2009, Grismer 2011, Mendyk 2011, Grismer 2011, King and Green 1993, Auffenberg 1980, Teynie and David 2010, Rathnayake et al. 2003, Van Wilgen and Richardson 2012, Honegger 1969, Ahmed 2009

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus scalaris</i>	no	NA	NA	NA	mainland	-16	151.84	5.85	1	3.31	19.37	Pianka 1995, Thompson and Pianka 2001, James et al. 1992, Pianka and King 2004, Wilson and Swan 2003, Nagy et al. 1999, Sweet and Pianka 2007, Sweet 2007, Clusella-Trullas et al. 2008, Brown and Nagy 2007, Wilson and Swan 2008, Eidenmueller and Philippen 2008, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Clemente et al. 2009, Swanson 2007, Christian and Bedford 1996
Varanidae	<i>Varanus spenceri</i>	no	NA	NA	NA	mainland	-21	1469.03	19.75	1	29.60	584.50	Pianka 1995, Thompson and Pianka 2001, Cogger 2000, James et al. 1992, Pianka and King 2004, Wilson and Swan 2003, Rogner 1997b, Greer 1989, De Lisle 1996, King et al. 1989, Pengilley 1981, Sweet and Pianka 2007, Clusella-Trullas et al. 2008, Horn and Visser 1997, Eidenmueller and Philippen 2008, Thompson et al. 2008, Kohler 2005, Frydlova and Frynta 2010
Varanidae	<i>Varanus tristis</i>	no	NA	NA	NA	mainland	-23	221.97	9.10	1	4.42	40.24	Pianka 1982, 1995, James et al. 1992, Pianka and King 2004, Pianka 1986, Pianka and Vitt 2003, Wilson and Swan 2003, Henle 1991, Rogner 1997b, Greer 1989, De Lisle 1996, Sweet and Pianka 2007, Sweet 2007, Moldovan 2008, Losos and Greene 1988, Huey et al. 2001, Horn and Visser 1997, Eidenmueller and Philippen

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Varanidae	<i>Varanus varius</i>	no	NA	NA	NA	mainland	-30	2242.28	7.55	2	17.28	260.93	2008, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Bush et al. 2010, Huey and Pianka 2007, Clemente et al. 2009, Wilson 2003, Heatwole and Pianka 1993, King and Green 1993, Bartholomew and Tucker 1964, Swanson 2007, Kutt et al. 2011, Rathnayake et al. 2003 Pianka 1995, Thompson and Pianka 2001, Cogger 2000, Perry and Garland 2002, James et al. 1992, Pianka and King 2004, Pianka and Vitt 2003, Wilson and Swan 2003, Rogner 1997b, Greer 1989, De Lisle 1996, Sweet and Pianka 2007, Brown and Nagy 2007, Wilson and Swan 2008, Horn and Visser 1997, Henle 1989c, Bustard 1978, Eidenmueller and Philippen 2008, Kirshner 2007, Thompson et al. 2008, Sinervo et al. 2010, Kohler 2005, Frydlova and Frynta 2010, Michael and Lindenmayer 2010, Clemente et al. 2009, Wilson 2003, King and Green 1993, Bartholomew and Tucker 1964, Heatwole and Taylor 1987, Swanson 2007, Michael et al. 2011, Bustard 1968

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Xantusiidae	<i>Lepidophyma flavimaculatum</i>	no	NA	NA	NA	mainland	15	15.22	5.00	1	0.62	3.10	Fitch 1970, Campbell 1999, Stafford and Meyer 2000, Lee 2000, Kohler 2003, Rand and Myers 1990, Duellman 1990, Bezy 1989, Pianka and Vitt 2003, Walker 1955, Bezy and Camarillo 2002, McCranie and Castaneda 2005, Guyer and Donnelly 2005, Ramirez-Bautista et al. 2008, Kohler 2008, McElroy et al. 2008, Goldberg 2009, Duellman 1963, Mendez-De la Cruz et al. 1999
Xantusiidae	<i>Xantusia arizonae</i>	no	NA	NA	NA	mainland	34	3.60	1.57	1	0.29	0.46	Amrein and Amrein 1951, Klauber 1938, Bezy 1967, Brattstrom 1952, Klauber 1931, Sinervo et al. 2010, Jones and Lovich 2009, Brattstrom 1965
Xantusiidae	<i>Xantusia henshawi</i>	no	NA	NA	NA	mainland	33	4.65	1.79	1	0.35	0.62	Stebbins 2003, Cox et al. 2003, Grismer 2002, Pianka and Vitt 2003, Smith 1946, Pianka and Vitt 2003, Mautz and Nagy 2000, Rogner 1997b, Van Denburgh 1922, Andrews and Pough 1980, Brown and Nagy 2007, Amrein and Amrein 1951, Lee 1976, Sinervo et al. 2010, Jones and Lovich 2009, Brattstrom 1965, Stebbins and McGinnis 2012, Lemm 2006

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Xantusiidae	<i>Xantusia riversiana</i>	yes	California Channel islands	San Clemente	151.80	Oceanic	33	11.38	4.40	0.75	0.51	1.68	Fitch 1970, 1985, Dunham and Miles 1985, Perry and Garland 2002, Stebbins 2003, Cooper and Vitt 2002, Pianka and Vitt 2003, Smith 1946, Greene 1982, Mautz and Nagy 2000, Rogner 1997b, Van Denburgh 1922, Dunham et al. 1988, Ramirez-Bautista et al. 2008, Andrews and Pough 1980, Brown and Nagy 2007, Vitt and Price 1982, Amrein and Amrein 1951, Brattstrom 1952, Cree and Guillette 1995, Fellers and Drost 1991, Jones and Lovich 2009, Sinervo et al. 2010, Goldberg 2006, Brattstrom 1965, Stebbins and McGinnis 2012
Xantusiidae	<i>Xantusia vigilis</i>	no	NA	NA	NA	mainland	34	1.37	1.84	1	0.20	0.36	Tinkle et al. 1970, Clobert et al. 1998, Dunham and Miles 1985, Stebbins 2003, Grismer 2002, Pianka 1986, Pianka and Vitt 2003, Smith 1946, Mautz and Nagy 2000, Rogner 1997b, Van Denburgh 1922, Dunham et al. 1988, Ramirez-Bautista et al. 2008, Andrews and Pough 1980, Brown and Nagy 2007, Vitt and Price 1982, Amrein and Amrein 1951, Bezy 1967, Warne and Charnov 2008, Huey et al. 2001, Radder et al. 2008, Jones and Lovich 2009, Sinervo et al. 2010, Case 1975, Huey and Pianka 2007, van Berkum 1988, Zweifel and Lowe 1966,

Family	Species	Insular	Archipelago	Largest island	Island area (sq km)	Island type	Latitude	Adult mass (g)	Clutch size (number of eggs in a clutch)	Brood frequency (number of clutches per year)	Hatchling mass (g)	Productivity (clutch size * brood frequency * hatchling mass)	Source
Xantusiidae	<i>Xantusia wigginsi</i>	no	NA	NA	NA	mainland	29	1.63	1.50	1	0.20	0.30	Rorabaugh 2008, Avery 1982, Brennan and Holycross 2009, Shine and Charnov 1992, Stebbins and McGinnis 2012, Lemm 2006
Xenosauridae	<i>Xenosaurus grandis</i>	no	NA	NA	NA	mainland	18	23.59	4.20	0.4165	1.11	1.95	Savage 1952, Jones and Lovich 2009, Stebbins and McGinnis 2012 Fitch 1970, Kohler 2003, Pianka and Vitt 2003, Pianka and Vitt 2003, Lemos-Espinal et al. 2003, Lemos-Espinal et al. 1996, Kohler 2008, Ballinger et al. 2000, Sinervo et al. 2010, Goldberg 2009, Zuniga-Vega 2011, Lemos-Espinal et al. 2003, Zamora-Abrego et al. 2007, Lemos-Espinal et al. 2012

B. Data of island attributes

Island type is divided to three categories – (1) oceanic (2) land bridge (3) continental. Other species are continental

Island area is in km² and isolation is in km (distance to the nearest landmass > 50,000 km² in area)

Island age is reported in million years.

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Amami Oshima	Ryukyu Islands	Oceanic	Kizaki, K. 1986. Geology and tectonics of the Ryukyu Islands. <i>Tectonophysics</i> , 125, 193-207.	712.35	Wikipedia	2.00	Hashimoto, J., Ohta, S., Fujikura, K. & Miura, T. 1995. Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the Mid-Okinawa Trough, Western Pacific. <i>Deep-Sea Research</i> , 42, 577-598.	618.5
Anegada	Leeward Islands	Land bridge	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	38.00	UNEP ISLANDS Web Site	0.01	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	412.2
Aneityum	New Hebrides	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . London, England	159.20	UNEP ISLANDS Web Site	3.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	1100.0
Angel de la Guarda	Baja California	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	945.20	UNEP ISLANDS Web Site	5.00	Case, T. J. , Cody, M. L. & Ezcurra, E. 2002. <i>A new island biogeography of the Sea of Cortez</i> . Oxford University Press, New York.	25.0
Anjouan	Comoro Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	424.00	University of California Press, Berkeley, CA.	11.50	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	388.3
Annobon	Sao Tome and Principe	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	15.70	UNEP ISLANDS Web Site	1.10	Schluter 2006 <i>Geological atlas of Africa</i> in google books	340.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Antigua	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	277.00	UNEP ISLANDS Web Site	2.58	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	699.2
Aruba	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	181.10	UNEP ISLANDS Web Site	40.00	James, K. H. 2005. A simple synthesis of Caribbean geology. Transactions of the 16th Caribbean Geological Conference, Barbados. Caribbean Journal of Earth Science, 39, 69-82. Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.; Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	24.0
Babeldaob	Palau Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	374.10	UNEP ISLANDS Web Site	20.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	870.4
Barrington	Galapagos Archipelago	Oceanic	Chiarucci, A., Bacaro, G., Triantis, K.A., Fernández-Palacios, J.M. 2011. Biogeographical determinants of pteridophytes and spermatophytes on oceanic archipelagos. Systematics and Biodiversity, 9, 191-201	24.00	UNEP ISLANDS Web Site	4.60	Cameron, R. A. D., Triantis, K. A., Guilhaumon, F., Alonso, M. R., Ibanez, M., de Frias Martins, A. M., Ladle, R. J. & Wittaker, R. J. 2013. Snails on oceanic islands: testing the general dynamic model of oceanic island biogeography using linear mixed effect models. Journal of Biogeography, 40, 117-130.	850.0
Batanta	New Guinea	Land bridge	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. Journal of Biogeography, 27, 1153-1167.	455.90	UNEP ISLANDS Web Site	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. Journal of Biogeography, 27, 1153-1167.	37.4
Bellona	Solomon Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	19.74	Dimond, J. M. & Mayr, E. 1976. Species-area relation for birds of the Solomon Archipelago. PNAS, 73, 262-266.	8.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	995.1

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Bequia	Windward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	18.00	Wikipedia	2.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	254.8
Biaro	Sangihe Archipelago	Oceanic	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	26.50	NIMA	3.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	45.7
Bonaire	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	282.50	UNEP ISLANDS Web Site	40.00	James, K. H. 2005. A simple synthesis of Caribbean geology. Transactions of the 16th Caribbean Geological Conference, Barbados. Caribbean Journal of Earth Science, 39, 69-82.	61.0
Bougainville	Solomon Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	8590.10	Dimond, J. M. & Mayr, E. 1976. Species-area relation for birds of the Solomon Archipelago. PNAS, 73, 262-266.	10.00	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	646.5
Buru	Maluku Islands	Oceanic	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. Journal of Biogeography, 39, 739-749.	8473.20	UNEP ISLANDS Web Site	4.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	470.0
Calayan	Philippine Islands	Oceanic	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	494.53	Wikipedia	5.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	77.9
Camorta	Andaman and Nicobar Islands	Oceanic	Curry 2005 Tectonics and history of the Andaman Sea region. Journal of Asian Earth Sciences, 25, 187-232.	188.00	Wikipedia	34.00	Curry 2005 Tectonics and history of the Andaman Sea region. Journal of Asian Earth Sciences, 25, 187-232.	329.7

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Catanduanes	Philippine Islands	Land bridge	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	1522.90	UNEP ISLANDS Web Site	0.01	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	8.4
Cebu	Philippine Islands	Oceanic	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	4467.50	UNEP ISLANDS Web Site	5.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	76.1
Ceralbo	Baja California	Oceanic	R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	150.50	Case 1975 UNEP ISLANDS Web Site	2.50	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	9.0
Christmas	None	Oceanic	Grimes, K. G. 2001. Karst Features of Christmas Island (Indian Ocean). Helictite, 37, 41-58.	135.00	UNEP ISLANDS Web Site	10.00	http://www.abc.net.au/nature/island/e/p2/about2.htm	346.0
Clarion	Revillagigedo Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	19.80	UNEP ISLANDS Web Site	10.00	Bryan, W. B. 1967. Geology and petrology of Clarion Islands, Mexico. Geological Society of America Bulletin, 78, 1461-1476.	698.4
Crete	Greece	Land bridge	Higgins in Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	8260.00	Welter-Schultes, F. W. & Williams, M. R. 2003. History, island area and habitat availability determine land snail species richness of Aegean islands. Journal of Biogeography, 26, 239-349.	5.50	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	100.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Crooked	Bahamas	Oceanic	Morrison 1998 A review of Bahamian ant (Hymenoptera: Formicidae) biogeography	282.10	UNEP ISLANDS Web Site	0.13	Erdman, J. S. & Key, M. M. 1997. Hydrogeology of the Cockburn town aquifer, San Salvador island, Bahamas, and the change in water quality resulting from the development of a resort community. Proceedings of the eighth symposium on the geology of the Bahamas and other carbonate regions, pp. 47-58.	224.5
Curacao	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	443.10	UNEP ISLANDS Web Site	40.00	James, K. H. 2005. A simple syntesis of Caribbean geology. Transactions of the 16th Caribbean Geological Conference, Barbados. Caribbean Journal of Earth Science, 39, 69-82.	60.0
Cyprus	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	9234.40	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	80.0
Dominica	Windward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	787.30	UNEP ISLANDS Web Site	2.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	500.0
El Hierro	Canary Islands	Oceanic	Hoernle & Carracedo in Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	290.50	UNEP ISLANDS Web Site	0.70	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	350.0
Enggano	Mentawai Archipelago	Oceanic	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	397.00	UNEP ISLANDS Web Site	1.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	114.4
Espanola	Galapagos Archipelago	Oceanic	Chiarucci, A., Bacaro, G., Triantis, K.A., Fernández-Palacios, J.M. 2011. Biogeographical determinants of pteridophytes and spermatophytes on oceanic archipelagos. Systematics and Biodiversity, 9, 191-201	60.00	http://www.galapaguide.com/islands_espanola.htm	5.60	Cameron, R. A. D., Triantis, K. A., Guilhaumon, F., Alonso, M. R., Ibanez, M., de Frias Martins, A. M., Ladle, R. J. & Wittaker, R. J. 2013. Snails on oceanic islands: testing the general dynamic model of oceanic island biogeography using linear mixed effect models. Journal of Biogeography, 40, 117-130.	850.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Espiritu Santo	New Hebrides	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	3955.50	UNEP ISLANDS Web Site	3.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1200.0
Eua	Tonga Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA. Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. Journal of Biogeography, 39, 739-749.	86.70	UNEP ISLANDS Web Site	50.00	Sterns, H. T. 1971. Geologic Setting of an Eocene Fossil Deposit on Eua Island, Tonga. Geological Society of America Bulletin, 82, 2541-2552.	1857.1
Flores	Lesser Sunda Islands	Oceanic	Plummer, Ph. S. & Belle, E. R. 1995. Mesozoic tectono-stratigraphic evolution of the Seychelles microcontinent. Sedimentary Geology, 96, 73-91.	14154.40	UNEP ISLANDS Web Site	4.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	288.2
Frigate	Seychelles Islands	Continental	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	2.19	Wikipedia	65.00	Plummer, P. 1998. Seychelles geology and the Shiva impact crater theory. Phelsuma, 6, 9-19.	1098.7
Fuerteventura	Canary Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1633.30	UNEP ISLANDS Web Site	20.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	90.0
Gomera	Canary Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	359.10	UNEP ISLANDS Web Site	12.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	310.0
Gran Canaria	Canary Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1529.90	UNEP ISLANDS Web Site	14.50	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	190.0
Grand Caicos	Bahamas	Oceanic	Morrison 1998 A review of Bahamian ant (Hymenoptera: Formicidae) biogeography	289.30	UNEP ISLANDS Web Site	0.13	Erdman, J. S. & Key, M. M. 1997. Hydrogeology of the Cockburn town aquifer, San Salvador island, Bahamas, and the change in water quality resulting from the development of a resort community. Proceedings of the eighth symposium on the geology of the Bahamas and other carbonate regions, pp. 47-58.	222.5
Grand Cayman	Cayman Islands	Oceanic	Brunt, M.A. & Davies, J.E. 1994. The Cayman Islands: natural history and biogeography. Kluwer Academic Publishers, Netherlands.	196.30	UNEP ISLANDS Web Site	10.00	Brunt, M.A. & Davies, J.E. 1994. The Cayman Islands: natural history and biogeography. Kluwer Academic Publishers, Netherlands.	297.3

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Grande Comore	Comoro Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1148.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA. Nichol, S. L., Lain, O. B. & Carter, C. H. 2003. Sheet-gravel evidence for a late Holocene tsunami run-up on beach dunes, Great Barrier Island, New Zealand. Sedimentary Geology, 155, 129-145.	0.50	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	450.0
Great Barrier	New Zealand	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	285.00	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	0.02	Erdman, J. S. & Key, M. M. 1997. Hydrogeology of the Cockburn town aquifer, San Salvador island, Bahamas, and the change in water quality resulting from the development of a resort community. Proceedings of the eighth symposium on the geology of the Bahamas and other carbonate regions, pp. 47-58.	16.2
Great Inagua	Bahamas	Oceanic	Morrison 1998 A review of Bahamian ant (Hymenoptera: Formicidae) biogeography Chapple, D. G. & Patterson, G. B. 2007. A new skink species (Oligosoma taumakae sp. nov.; Reptilia: Scincidae) from the Open Bay Islands, New Zealand, New Zealand Journal of Zoology, 34, 347-357.	1615.30	UNEP ISLANDS Web Site	0.13		86.5
Great Island	New Zealand	Continental	Curray 2005 Tectonics and history of the Andaman Sea region. Journal of Asian Earth Sciences, 25, 187-232.	4.04	Wikipedia	1.80	http://www.conservation.org.nz	57.8
Great Nicobar	Andaman and Nicobar Islands	Oceanic	Curray 2005 Tectonics and history of the Andaman Sea region. Journal of Asian Earth Sciences, 25, 187-232.	955.70	UNEP ISLANDS Web Site	34.00	Curray 2005 Tectonics and history of the Andaman Sea region. Journal of Asian Earth Sciences, 25, 187-232.	206.3

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Grenada	Windward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	322.70	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	150.0
Guadalcanal	Solomon Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	5280.10	UNEP ISLANDS Web Site	8.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	970.3
Guam	None	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	541.00	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	2060.0
Guam	Mariana Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	541.00	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	2060.0
Hainan	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	33209.80	UNEP ISLANDS Web Site	0.02	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	20.0
Halmahera	Maluku Islands	Oceanic	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. Journal of Biogeography, 39, 739-749.	18039.60	UNEP ISLANDS Web Site	5.00	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. Journal of Biogeography, 39, 739-749.	286.0
Ibiza	Balearic Islands	Continental	Rosenbaum, G., Lister, G.S. and Duboz, C. 2002 Reconstruction of the tectonic evolution of the western Mediterranean since the Oligocene. Journal of the Virtual Explorer, 8, 107-130.	576.90	UNEP ISLANDS Web Site	5.30	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	85.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Iriomotejima	Ryukyu Islands	Oceanic	Kizaki, K. 1986. Geology and tectonics of the Ryukyu Islands. <i>Tectonophysics</i> , 125, 193-207.	289.00	Wikipedia	2.00	Hashimoto, J., Ohta, S., Fujikura, K. & Miura, T. 1995. Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the Mid-Okinawa Trough, Western Pacific. <i>Deep-Sea Research</i> , 42, 577-598.	441.0
Isabela	Galapagos Archipelago	Oceanic	Chiarucci, A., Bacaro, G., Triantis, K.A., Fernández-Palacios, J.M. 2011. Biogeographical determinants of pteridophytes and spermatophytes on oceanic archipelagos. <i>Systematics and Biodiversity</i> , 9, 191-201	4588.00	UNEP ISLANDS Web Site	0.70	Ladle, R. J. & Wittaker, R. J. 2013. Snails on oceanic islands: testing the general dynamic model of oceanic island biogeography using linear mixed effect models. <i>Journal of Biogeography</i> , 40, 117-130.	850.0
Jamaica	Greater Antilles	Oceanic	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	11189.60	UNEP ISLANDS Web Site	10.00	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	148.0
Jarak	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	14.90	UNEP ISLANDS Web Site	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	61.7
Komodo	Lesser Sunda Islands	Oceanic	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	330.00	UNEP ISLANDS Web Site	4.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	302.2
Kyushu	Japan Islands	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	37437.20	UNEP ISLANDS Web Site	0.02	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. <i>Philosophical Transactions of the Royal Society B</i> , 363, 3293-3308.	2.4
Lancelin	None	Land bridge	Jonker, L. & Bullen, L. 2004. Turquoise coast island Nature Reserves. Management Plan. Department of Conservation and Land Management. Australia	0.09	NIMA	0.01	Jonker, L. & Bullen, L. 2004. Turquoise coast island Nature Reserves. Management Plan. Department of Conservation and Land Management. Australia	0.6

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Langkawi	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	363.00	UNEP ISLANDS Web Site	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. Journal of Biogeography, 27, 1153-1167.	15.0
Lipari	Lipari (Eolie) Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	37.00	Wikipedia	5.32	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	29.0
Lord Howe	None	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	15.04	NIMA	7.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	600.0
Madeira	Maderia Archipelago	Oceanic	Hoernle & Carracedo in Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA. Plummer, Ph. S. & Belle, E. R. 1995. Mesozoic tectono-stratigraphic evolution of the Seychelles microcontinent. Sedimentary Geology, 96, 73-91.	755.00	Wikipedia	4.60	Kostas Triantis personal communication	560.0
Mahe	Seychelles Islands	Continental		148.00	UNEP ISLANDS Web Site	65.00	Plummer, P. 1998. Seychelles geology and the Shiva impact crater theory. Phelsuma, 6, 9-19.	1051.1
Makira	Solomon Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	3089.80	UNEP ISLANDS Web Site	24.00	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	1151.0
Malpelo	None	Oceanic	Graham, J. B. 1975. The biological investigation of Malpelo island, Colombia. Smithsonian Institution Press.	7.65	NIMA	15.00	Graham, J. B. 1975. The biological investigation of Malpelo island, Colombia. Smithsonian Institution Press.	375.9
Malta	Maltese Archipelago	Land bridge	Micallef, A., Foglini, F., Le Bas, T., Angeletti, L., Maselli, V., Pasuto, A. & Taviani, M. 2013. The submerged paleolandscape of the Maltese Islands: Morphology, evolution and relation to Quaternary environmental change. Marine Geology, 335, 129-147.	245.70	UNEP ISLANDS Web Site	0.01	Micallef, A., Foglini, F., Le Bas, T., Angeletti, L., Maselli, V., Pasuto, A. & Taviani, M. 2013. The submerged paleolandscape of the Maltese Islands: Morphology, evolution and relation to Quaternary environmental change. Marine Geology, 335, 129-147.	91.3

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Manus	Bismarck Archipelago	Oceanic	Franz, L. & Wirth, R. 2000. Spinel inclusions in olivine of peridotite xenoliths from TUBAF seamount (Bismarck Archipelago/Papua New Guinea): evidence for the thermal and tectonic evolution of the oceanic lithosphere. <i>Contributions to Mineralogy and Petrology</i> , 140, 283-295.	1940.20	UNEP ISLANDS Web Site	5.00	Exon, N. F. and Tiffin, D. L. 1982. Geology of offshore New Ireland basin in northern Papua New Guinea, and its petroleum prospects. – In: <i>Transactions of the Third Circum-Pacific Energy and Mineral Resources Conference</i> , 1982.	272.3
Marchena	Galapagos Archipelago	Oceanic	Chiarucci, A., Bacaro, G., Triantis, K.A., Fernández-Palacios, J.M. 2011. Biogeographical determinants of pteridophytes and spermatophytes on oceanic archipelagos. <i>Systematics and Biodiversity</i> , 9, 191-201	130.00	Wikipedia http://www.fondors.com/world/caribbean/st-lucia/review-94961.html	0.70	Cameron, R. A. D., Triantis, K. A., Guilhaumon, F., Alonso, M. R., Ibanez, M., de Frias Martins, A. M., Ladle, R. J. & Wittaker, R. J. 2013. Snails on oceanic islands: testing the general dynamic model of oceanic island biogeography using linear mixed effect models. <i>Journal of Biogeography</i> , 40, 117-130.	850.0
Maria Major	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. <i>Proceedings of the Ocean Drilling Program, Scientific results</i> , 110, 29-44.	0.25		4.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	367.0
Martinique	Windward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. <i>Proceedings of the Ocean Drilling Program, Scientific results</i> , 110, 29-44.	1166.60	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	410.0
Maud	New Zealand	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	3.09	Bell, B. D., Pledger, S. & Cewhurst, P. L. 2004. The fate of a population of the endemic frog <i>Leiopelma pakeka</i> (Anura: Leiopelmatidae) translocated to restored habitat on Maud Island, New Zealand.	0.02	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. <i>Philosophical Transactions of the Royal Society B</i> , 363, 3293-3308.	1.5

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
					New Zealand Journal of Zoology, 31, 123-131.			
Mauritius	Mascarene Archipelago	Oceanic	Micheneau, C., Carlswald, B. S., Fay, M. F., Bytebier, B., Pailler, T. & Chase, M. W. 2008. Phylogenetics and biogeography of Mascarene angraecoid orchids (Vandaeae, Orchidaceae). Molecular Phylogenetics and Evolutions, 46, 908-922.	2040.00	Polhemus, D. A. & Polhemus, J. T. 2008. A new Indian Ocean species of Ochterus from the island of Mauritius (Hemiptera: Heteroptera: Ochteridae). Acta Entomologica Musei Nationalis Pragae, 48, 281-288.	7.00	McDougall, I. & Chamalaun, F. H. 1969. Isotopic dating and geomagnetic polarity studies on volcanic rocks from Mauritius, Indian Ocean. Geological Society of America Bulletin, 80, 1419-1442.	866.5
Mayotte	Comoro Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	314.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	10.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	301.1

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Milos	Greece	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	150.60	Welter-Schultes, F. W. & Williams, M. R. 2003. History, island area and habitat availability determine land snail species richness of Aegean islands. Journal of Biogeography, 26, 239-349.	4.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	102.0
Miyako-jima	Miyako Islands	Oceanic	Kizaki, K. 1986. Geology and tectonics of the Ryukyu Islands. Tectonophysics, 125, 193-207.	158.70	Wikipedia	2.00	Hashimoto, J., Ohta, S., Fujikura, K. & Miura, T. 1995. Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the Mid-Okinawa Trough, Western Pacific. Deep-Sea Research, 42, 577-598.	530.9
Mona	Greater Antilles	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	57.00	Wikipedia	2.00	Heatwole, H. & MacKenzie, F. 1967. Herpetogeography of Puerto Rico. 4. Paleogeography, Faunal Similarity and Endemism. Evolution, 21, 429-438. (Jstor online)	61.3
Montserrat	Leeward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	124.10	UNEP ISLANDS Web Site	1.16	EGS Inc. 2010. Final Report Geothermal Exploration in Montserrat, Caribbean. For: Minister of Communications and Works Government of Montserrat, Caribbean. Santa Rosa, California.	650.0
Navassa	Greater Antilles	Oceanic	Miller, M. W., Halley, R. B. & Gleason, C.R. 2008. Reef geology and biology of Navassa Island. Coral Reefs of the USA, 1, 407-433.	5.20	Wikipedia	2.00	http://www.eoearth.org/article/Navassa_Island	55.5

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Negros	Philippine Islands	Oceanic	Linkem, C. W., Hased, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	13074.50	UNEP ISLANDS Web Site	5.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	46.7
Nevis	Leeward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44. Franz, L. & Wirth, R. 2000. Spinel inclusions in olivine of peridotite xenoliths from TUBAF seamount (Bismark Archipelago/Papua New Guinea): evidence for the thermal and tectonic evolution of the oceanic lithosphere. Contributions to Mineralogy and Petrology, 140, 283-295.	92.30	UNEP ISLANDS Web Site	2.00	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. Journal of Biogeography, 39, 739-749.	622.2
New Britain	Bismarck Archipelago	Oceanic	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	35144.60	UNEP ISLANDS Web Site	2.50	Woodhead, J., et al. 1998. Magma genesis in the New Britain island arc: further insights into melting and mass transfer processes. - Journal of Petrology 39: 1641-1668.	89.4
New Caledonia	New Caledonia	Continental	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	18575.00	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	45.00	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	1200.0
New Georgia	Solomon Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	2043.50	Dimond, J. M. & Mayr, E. 1976. Species-area relation for birds of the Solomon Archipelago. PNAS, 73, 262-266.	15.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	739.2

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
New Ireland	Bismarck Archipelago	Oceanic	Franz, L. & Wirth, R. 2000. Spinel inclusions in olivine of peridotite xenoliths from TUBAF seamount (Bismarck Archipelago/Papua New Guinea): evidence for the thermal and tectonic evolution of the oceanic lithosphere. <i>Contributions to Mineralogy and Petrology</i> , 140, 283-295.	7404.50	UNEP ISLANDS Web Site	5.00	Exon, N. F. and Tiffin, D. L. 1982. Geology of offshore New Ireland basin in northern Papua New Guinea, and its petroleum prospects. – In: <i>Transactions of the Third Circum-Pacific Energy and Mineral Resources Conference</i> , 1982. Grandcolas, P., Murienne, J., Robillard, T., Desutter-Grandcolas, L., Jourdan, H., Guilbert, E. & Deharveng, L. 2008. New Caledonia: a very old Darwinian island? <i>Philosophical Transactions of the Royal Society B</i> , 363, 3309-3317. Pal, T., et al. 2003. Geodynamic evolution of the outer-arc-forearc belt in the Andaman Islands, the central part of the Burma-Java subduction complex. - <i>Geological Magazine</i> 140: 289-307.	508.2
Norfolk	None	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	36.80	UNEP ISLANDS Web Site	3.70		763.2
North Andaman	Andaman and Nicobar Islands	Oceanic	Curry 2005 <i>Tectonics and history of the Andaman Sea region</i> . <i>Journal of Asian Earth Sciences</i> , 25, 187-232.	2780.70	UNEP ISLANDS Web Site	15.00		260.0
North Andros	Bahamas	Oceanic	Morrison 1998 <i>A review of Bahamian ant (Hymenoptera: Formicidae) biogeography</i>	3439.40	UNEP ISLANDS Web Site	0.13	Erdman, J. S. & Key, M. M. 1997. Hydrogeology of the Cockburn town aquifer, San Salvador island, Bahamas, and the change in water quality resulting from the development of a resort community. <i>Proceedings of the eighth symposium on the geology of the Bahamas and other carbonate regions</i> , pp. 47-58.	147.0
Nosy Be	Madagascar	Oceanic	Beccaluva, L., Bianchini, G. & Wilson, M. 2011. <i>Volcanism and evolution of the African lithosphere</i> . The Geological Society of America, Colorado.	290.30	UNEP ISLANDS Web Site	7.32	Melluso, L. and Morra, V. 2000. Petrogenesis of Late Cenozoic mafic alkaline rocks of the Nosy Be archipelago (northern Madagascar): relationships with the Comorean magmatism. - <i>Journal of Volcanology and Geothermal Research</i> 96: 129-142.	12.3

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Okinawajima	Ryukyu Islands	Oceanic	Kizaki, K. 1986. Geology and tectonics of the Ryukyu Islands. <i>Tectonophysics</i> , 125, 193-207.	1199.50	UNEP ISLANDS Web Site	2.00	Hashimoto, J., Ohta, S., Fujikura, K. & Miura, T. 1995. Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the Mid-Okinawa Trough, Western Pacific. <i>Deep-Sea Research</i> , 42, 577-598.	600.0
Palawan	Philippine Islands	Continental	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	12188.60	UNEP ISLANDS Web Site	20.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	288.1
Palawan	Philippine Islands	Continental	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	12188.60	UNEP ISLANDS Web Site	20.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	288.1
Panay	Philippine Islands	Oceanic	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). <i>Molecular Phylogenetics and Evolution</i> , 56, 572-585.	12011.10	UNEP ISLANDS Web Site	5.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	43.6
Pemba	None	Continental	http://www.zanzinet.org/zanzibar/nature/miamba.html	889.90	UNEP ISLANDS Web Site	10.00	http://www.bbc.co.uk/oceans/locations/spiceislands/	50.6
Perhentian Besar	Perhentian Islands	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	11.51	NIMA	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	16.4
Pinang	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	295.30	UNEP ISLANDS Web Site	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	5.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Pitt	Chatham Islands	Oceanic	Liggins, L., Chapple, D. G., Daugherty, C. H. & Ritchie, P. A. 2008. Origin and post-colonization evolution of the Chatham islands skink (<i>Oligosoma nigriplantare nigriplantare</i>). <i>Molecular Ecology</i> , 17, 3290-3305.	744.60	UNEP ISLANDS Web Site	1.00	Liggins, L., Chapple, D. G., Daugherty, C. H. & Ritchie, P. A. 2008. Origin and post-colonization evolution of the Chatham islands skink (<i>Oligosoma nigriplantare nigriplantare</i>). <i>Molecular Ecology</i> , 17, 3290-3305.	716.1
Pohnpei	Caroline Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	334.00	UNEP ISLANDS Web Site	1.00	Spengler, S. R., Peterson, F. L. & Mink, J. F. 1992. Geology and hydrology of the island of Pohnpei, Federated States of Micronesia. Technical Report No. 189.	1884.5
Polillo	Philippine Islands	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA. Plummer, Ph. S. & Belle, E. R. 1995. Mesozoic tectono-stratigraphic evolution of the Seychelles microcontinent. <i>Sedimentary Geology</i> , 96, 73-91.	628.90	UNEP ISLANDS Web Site	0.01	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). <i>Molecular Phylogenetics and Evolution</i> , 56, 572-585.	17.7
Praslin	Seychelles Islands	Continental	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	26.00	UNEP ISLANDS Web Site	65.00	Plummer, P. 1998. Seychelles geology and the Shiva impact crater theory. <i>Phelsuma</i> , 6, 9-19.	1101.1
Puerto Rico	Greater Antilles	Continental	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	9099.80	UNEP ISLANDS Web Site	12.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	113.8
Reunion	Mascarene Archipelago	Oceanic	Micheneau, C., Carlswald, B. S., Fay, M. F., Bytebier, B., Pailler, T. & Chase, M. W. 2008. Phylogenetics and biogeography of Mascarene angraecoid orchids (Vandaeae, Orchidaceae). <i>Molecular Phylogenetics and Evolutions</i> , 46, 908-922.	2535.20	UNEP ISLANDS Web Site	3.00	Micheneau, C., Carlswald, B. S., Fay, M. F., Bytebier, B., Pailler, T. & Chase, M. W. 2008. Phylogenetics and biogeography of Mascarene angraecoid orchids (Vandaeae, Orchidaceae). <i>Molecular Phylogenetics and Evolutions</i> , 46, 908-922.	680.2

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Roatan	Bay Islands	Oceanic	McBirney, A. R., and Bass, M. N., 1969, Geology of Bay Islands, Gulf of Honduras. In: McBirney, A. R., ed., Tectonic relations of northern Central America and the western Caribbean —the Bonacca Expedition: American Association Petroleum Geologists, Mem. 11, p. 229-243.	156.60	NIMA	1.00	McBirney, A. R., and Bass, M. N., 1969, Geology of Bay Islands, Gulf of Honduras. In: McBirney, A. R., ed., Tectonic relations of northern Central America and the western Caribbean —the Bonacca Expedition: American Association Petroleum Geologists, Mem. 11, p. 229-243.	48.2
Saba	Leeward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	13.00	UNEP ISLANDS Web Site	0.36	http://www.caribbeanvolcanoes.com/saba/geology.htm	547.1
Saint Martin	Leeward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	91.90	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	548.8
Salibabu	Talau Islands	Oceanic	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	86.87	NIMA	20.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	892.0
Samar	Philippine Islands	Land bridge	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	12849.40	UNEP ISLANDS Web Site	0.01	Linkem, C. W., Hesed, K. M., Diesmos, A. C. & Brown, R. M. 2010. Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). Molecular Phylogenetics and Evolution, 56, 572-585.	18.9
San Clemente	California Channel islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	151.80	UNEP ISLANDS Web Site	14.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	33.9
San Pedro Martir	Baja California	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1.22	Case 1975	5.00	Case, T. J. , Cody, M. L. & Ezcurra, E. 2002. A new island biogeography of the Sea of Cortez. Oxford University Press, New York.	48.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Santa Catalina	Baja California	Continental	Case, T. J. , Cody, M. L. & Ezcurra, E. 2002. A new island biogeography of the Sea of Cortez. Oxford University Press, New York.	194.20	Wikipedia	2.50	Case, T. J. , Cody, M. L. & Ezcurra, E. 2002. A new island biogeography of the Sea of Cortez. Oxford University Press, New York.	22.9
Santa Cruz	Galapagos Archipelago	Oceanic	Chiarucci, A., Bacaro, G., Triantis, K.A., Fernández-Palacios, J.M. 2011. Biogeographical determinants of pteridophytes and spermatophytes on oceanic archipelagos. Systematics and Biodiversity, 9, 191-201	985.60	UNEP ISLANDS Web Site	3.60	Cameron, R. A. D., Triantis, K. A., Guilhaumon, F., Alonso, M. R., Ibanez, M., de Frias Martins, A. M., Ladle, R. J. & Wittaker, R. J. 2013. Snails on oceanic islands: testing the general dynamic model of oceanic island biogeography using linear mixed effect models. Journal of Biogeography, 40, 117-130.	850.0
Santa Luzia	Cape Verde	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	36.70	UNEP ISLANDS Web Site	2.00	Ramalho, R., Helffrich, G., Schmidt, D. N. & Vance, D. 2010. Tracers of uplift and subsidence in the Cape Verde archipelago. Journal of the Geological Society, 167, 519-538.	203.0
Santo Antao	Cape Verde	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	787.30	UNEP ISLANDS Web Site	7.57	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	620.0
Sao Tiago	Cape Verde	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	991.00	http://www.britannica.com/EBchecked/topic/523574/Sao-Tiago-Island	10.00	Chiarucci, A., Bacaro, G., Triantis, K.A., Fernández-Palacios, J.M. 2011. Biogeographical determinants of pteridophytes and spermatophytes on oceanic archipelagos. Systematics and Biodiversity, 9, 191-201	620.0
Sardinia	None	Land bridge	Azzaroli 1981 Cainozoic mammals and the biogeography of the island of Sardinia, Western Mediterranean. Palaeogeography, Palaeoclimatology, Palaeoecology, 36, 107-111.	23949.00	UNEP ISLANDS Web Site	0.01	Azzaroli 1981 Cainozoic mammals and the biogeography of the island of Sardinia, Western Mediterranean. Palaeogeography, Palaeoclimatology, Palaeoecology, 36, 107-111.	200.0
Savai'i	Samoa Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1717.60	UNEP ISLANDS Web Site	2.50	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	2705.8


Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Seram	Maluku Islands	Oceanic	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	17454.20	UNEP ISLANDS Web Site	5.00	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	142.9
Seribuat	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	5.42	NIMA	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	14.1
Siberut	Mentawai Archipelago	Oceanic	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	3828.50	UNEP ISLANDS Web Site	1.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. <i>Journal of Asian Earth Sciences</i> , 20, 353-431.	120.9
Sicily	None	Land bridge	Rosenbaum, G., Lister, G.S. and Duboz, C. 2002 Reconstruction of the tectonic evolution of the western Mediterranean since the Oligocene. <i>Journal of the Virtual Explorer</i> , 8, 107-130.	25662.40	UNEP ISLANDS Web Site	0.02	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	3.0
Skyros	Greece	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	209.00	Wikipedia	4.00	Kostas Triantis personal communication	86.4
Socotra	Socotra Archipelago	Continental	Mies, B. A. & Brown, G. 2012. <i>Vegetation ecology of Socotra</i> . Springer Dordrecht Heidelberg, New York.	3606.70	UNEP ISLANDS Web Site	18.00	Mies, B. A. & Brown, G. 2012. <i>Vegetation ecology of Socotra</i> . Springer Dordrecht Heidelberg, New York.	233.6
St. Croix	Virgin Islands	Oceanic	Heatwole, H. & MacKenzie, F. 1967. Herpetogeography of Puerto Rico. 4. Paleogeography, Faunal Similarity and Endemism. <i>Evolution</i> , 21, 429-438. (Jstor online)	214.40	UNEP ISLANDS Web Site	2.00	Heatwole, H. & MacKenzie, F. 1967. Herpetogeography of Puerto Rico. 4. Paleogeography, Faunal Similarity and Endemism. <i>Evolution</i> , 21, 429-438. (Jstor online)	382.7
St. Lucia	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. <i>Proceedings of the Ocean Drilling Program, Scientific results</i> , 110, 29-44.	639.80	UNEP ISLANDS Web Site	4.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	340.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
St. Vincent	Lesser Antilles	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. Proceedings of the Ocean Drilling Program, Scientific results, 110, 29-44.	381.00	UNEP ISLANDS Web Site	2.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	275.0
Stewart	New Zealand	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1814.70	UNEP ISLANDS Web Site	0.02	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	31.9
Taiwan	Taiwan	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	34506.60	UNEP ISLANDS Web Site	0.02	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	130.0
Takarajima	Ryukyu Islands	Oceanic	Kizaki, K. 1986. Geology and tectonics of the Ryukyu Islands. Tectonophysics, 125, 193-207. Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. Journal of Biogeography, 39, 739-749.	7.14	Wikipedia	2.00	Hashimoto, J., Ohta, S., Fujikura, K. & Miura, T. 1995. Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the Mid-Okinawa Trough, Western Pacific. Deep-Sea Research, 42, 577-598.	555.6
Taliabu	Maluku Islands	Oceanic		2913.20	UNEP ISLANDS Web Site	3.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA. Dickinson, W. R. 2006. Temper Sands in Prehistoric Oceanian Pottery: Geotectonics, Sedimentology, Petrography, Provenance. - Geological Society of America.	135.4
Taumako	Solomon Islands	Oceanic	http://antbase.org/ants/africa/personal/solomons/sols12.html	10.00	UNEP ISLANDS Web Site	5.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	1807.0
Tenerife	Canary Islands	Oceanic	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	2007.80	UNEP ISLANDS Web Site	8.00	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	275.0
Tenggol	None	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. Encyclopedia of islands. University of California Press, Berkeley, CA.	4.03	NIMA	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. Journal of Biogeography, 27, 1153-1167.	26.0


Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Timor	Lesser Sunda Islands	Continental	Carstensen et al. 2012 Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. <i>Journal of Biogeography</i> , 39, 739-749.	28418.10	UNEP ISLANDS Web Site http://www.su.peryacht-charters.com/motor/my_moecca/images/moecca_tiomandf	4.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	450.0
Tioman	Seribuat Archipelago	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	132.00	UNEP ISLANDS Web Site	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	38.2
Tobago	Trinidad and Tobago	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	308.80	UNEP ISLANDS Web Site	0.02	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	120.0
Tokunoshima	Ryukyu Islands	Oceanic	Kizaki, K. 1986. Geology and tectonics of the Ryukyu Islands. <i>Tectonophysics</i> , 125, 193-207.	248.87	NIMA	2.00	Hashimoto, J., Ohta, S., Fujikura, K. & Miura, T. 1995. Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the Mid-Okinawa Trough, Western Pacific. <i>Deep-Sea Research</i> , 42, 577-598.	692.0
Trangan	Aru Islands	Continental	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	2148.50	UNEP ISLANDS Web Site	0.02	Voris, H. K. 2001 Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time duration. <i>Journal of Biogeography</i> , 27, 1153-1167.	203.4
Trinidad	Trinidad and Tobago	Land bridge	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	5008.70	UNEP ISLANDS Web Site http://www.caribbeantravelweb.com/stvincen/areaguide.htm	0.02	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	11.0
Union	Windward Islands	Oceanic	Bouysse, P., Westercamp, D. & Andreieff, P. 1990. 4. The Lesser Antilles island arc. <i>Proceedings of the Ocean Drilling Program, Scientific results</i> , 110, 29-44.	8.49		2.00	Gillespie, R. G. & Clague, D. A. 2009. <i>Encyclopedia of islands</i> . University of California Press, Berkeley, CA.	207.0

Largest island	Archipelago	Island type	Island type source	Island area (km ²)	Island area source	island age (my)	Island age source	Isolation (km)
Utila	Bay Islands	Oceanic	McBirney, A. R., and Bass, M. N., 1969, Geology of Bay Islands, Gulf of Honduras. In: McBirney, A. R., ed., Tectonic relations of northern Central America and the western Caribbean —the Bonacca Expedition: American Association Petroleum Geologists, Mem. 11, p. 229-243.	42.08	NIMA	1.00	McBirney, A. R., and Bass, M. N., 1969, Geology of Bay Islands, Gulf of Honduras. In: McBirney, A. R., ed., Tectonic relations of northern Central America and the western Caribbean —the Bonacca Expedition: American Association Petroleum Geologists, Mem. 11, p. 229-243.	32.8
Viti Levu	Fiji Islands	Oceanic	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	10531.00	UNEP ISLANDS Web Site	10.00	Neall, V. E. & Trewick, S. A. 2008. The age and origin of the Pacific islands: a geological overview. Philosophical Transactions of the Royal Society B, 363, 3293-3308.	1856.8
Waigeo	New Guinea	Oceanic	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	3153.70	UNEP ISLANDS Web Site	20.00	Hall 2002 Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-431.	45.9


- C. The distribution in island area, age and isolation across different island types
1. The distribution of \log_{10} -transformed island areas



2. Histogram for the distribution of \log_{10} -transformed island age



3. The distribution of \log_{10} -transformed island isolation



D. Literature sources that were used to create the datasets and to determine the distribution of the species

- Abdala, C.S. (2006) Dos nuevas especies del genero *Liolaemus* (Iguania: Liolaemidae) y redescription de *Liolaemus boulengeri* (Koslowky, 1898). *Cuadernos de Herpetologia*, **19**, 3-33.
- Abensperg-Traun, M. & Steven, D. (1997) Ant- and termite-eating in Australian mammals and lizards: a comparison. *Australian Journal of Ecology*, **22**, 9-17.
- Adamopoulo, C., Valakos, E.D. & Pafilis, P. (1999) Summer diet of *Podarcis milensis*, *P. gaigeae* and *P. erhardii* (Sauria: Lacertidae). *Bonner Zoologische Beitrage*, **48**, 275-282.
- Adamopoulou, C. & Valakos, E.D. (2000) Small clutch size in a Mediterranean endemic lacertid (*Podarcis milensis*). *Copeia*, **2000**, 610-614.
- Adamopoulou, C. & Legakis, A. (2002) Diet of a lacertid lizard (*Podarcis milensis*) in an insular dune ecosystem. *Israel Journal of Zoology*, **48**, 207-219.
- Adamopoulou, C. & Valakos, E.D. (2005) Thermal ecology and activity cycle of *Podarcis milensis* in a sandy coastal area. *Israel Journal of Zoology*, **51**, 39-52.
- Aguilar, R. & Cruz, F.B. (2010) Refuge use in a Patagonian nocturnal lizard, *Homonota darwini*: the role of temperature. *Journal of Herpetology*, **44**, 236-241.
- Ahmadzadeh, F., Kiabi, B.H., Kami, H.G. & Hojjati, V. (2008) A preliminary study of the lizard fauna and their habitats in Northwestern Iran. *Asiatic Herpetological Research*, **11**, 1-9.
- Ahmadzadeh, F., Kami, H.G., Hojjati, V. & Rezazadeh, E. (2009) Contribution to the knowledge of *Eremias strauchi strauchi* Kessler, 1878 (Sauria: Lacertidae) from northwestern Iran. *Iranian Journal of Animal Biosystematics*, **5**, 17-24.
- Ahmed, M.F., Das, A. & Dutta, S.K. (2009) *Amphibians and Reptiles of Northeast India: A Photographic Guide*. Aaranyak.
- Ahsan, M.F. & Abu Saeed, M. (2004) Some aspects of breeding biology of the Bengal lizard (*Varanus bengalensis*) in Bangladesh. *Asiatic Herpetological Research*, **10**, 236-240.
- Akani, G.C., Capizzi, D. & Luiselli, L. (2002) Community ecology of scincid lizards in a swamp rainforest of south-eastern Nigeria. *Russian Journal of Herpetology*, **9**, 125-134.
- Alcala, A.C. (1966) Populations of three tropical lizards of Negros Island, Philippines. Stanford University
- Alcala, A.C. & Brown, W.C. (1967) Population ecology of the tropical scincoid lizard, *Emoia atrocostata*, in the Philippines. *Copeia*, **1967**, 596-604.
- Alexander, K. (2007) *A photographic field guide to the reptiles and amphibians of Dominica, West Indies*. Texas A&M University.
- Al-Hashem, M. (2009) Monitoring population abundance of the sand lizard *Acanthodactylus scutellatus* and their ant prey in oil polluted soils at Kuwait's greater Al-Burgan oil field. *Pakistan Journal of Biological Sciences*, **12**, 1425-1429.
- Al-Johany, A.M. & Spellerberg, I.F. (1988) Reproductive biology of the lizard *Acanthodactylus schmidti*, Weigmann in central Arabia. *Journal of Arid Environments*, **15**, 197-207.
- Al-Johany, A.M., Al-Sadoon, M.K. & Al-Farraj, S.A. (1997) Reproductive biology of the skink *Scincus mitranus* (Anderson, 1871) in the central region of Saudi Arabia. *Journal of Arid Environments*, **36**, 319-326.
- Al-Johany, A.M., Al-Sadoon, M.K. & Al-Farraj, S.A. (1999) Thermal ecology and activity of

- the sand fish lizard, *Scincus mitranus* (Scincidae) in central Arabia. *Journal of King Saud University*, **2**, 1-16.
- Allison, A. (2006) Reptiles and amphibians of the Trans-Fly region, New Guinea. *Contribution to the Pacific Biological Survey*, **39**, 1-50.
- Almeida-Gomes, M., Vrcibradic, D., Siqueira, C.C., Kiefer, M.C., Klaion, T., Almeida-Santos, P., Nascimento, D., Ariani, C.V., Borges-Junior, V.N., Freitas-Filho, R.F., van Sluys, M. & Rocha, C.F. (2008) Herpetofauna of an Atlantic rainforest area (Morro Sao Joao) in Rio de Janeiro State, Brazil. *Anais da Academia Brasileira de Ciencias*, **80**, 291-300.
- Almog, A., Bonen, H., Herman, K. & Werner, Y.L. (2005) Subspeciation or none? The hardun in the Aegean (Reptilia: Sauria: Agamidae: *Laudakia stellio*). *Journal of Natural History*, **39**, 567-586.
- Al-Sadoon, M.K. & Spellerberg, I.F. (1985) Effect of temperature on the oxygen consumption of lizards from different climatic regions. *Amphibia-Reptilia*, **6**, 241-258.
- Al-Sadoon, M.K. & Abdo, N.M. (1991) Temperature and body mass effects on the metabolic rate of *Acanthodactylus schmidti* Weigmann (Reptilia : Lacertidae). *Journal of Arid Environments*, **21**, 351-361.
- Al-Sadoon, M.K., Al-Johany, A.M. & Al-Farraj, S.A. (1999) Food and feeding habits of the sand fish lizard *Scincus mitranus*. *Saudi Journal of Biological Sciences*, **6**, 91-101.
- Alvarez, H.G. (2004) Pollination and seed dispersal by lizards: a review. *Revista Chilena de Historia Natural*, **77**, 569-577.
- Alvarez del Toro, M. & Smith, H.M. (1962) Notulae Herpetologicae Chiapasiae III. *Herpetologica*, **18**, 101-107.
- Amat, F. (2008) Exploring female reproductive tactics: trade-offs between clutch size, egg mass and newborn size in lacertid lizards. *Herpetological Journal*, **18**, 147-153.
- Amat, F., Perez-Mellado, V., Hernandez-Estevez, J.A. & Diez, T.G. (2008) Dietary strategy of a Pyrenean lizard, *Iberolacerta aurelioi*, living in a poor resources alpine environment. *Amphibia-Reptilia*, **29**, 329-336.
- Amey, A.P. & Whittier, J.M. (2000) The annual reproductive cycle and sperm storage in the bearded dragon, *Pogona barbata*. *Australian Journal of Zoology*, **48**, 411-419.
- Amo, K., Lopez, P. & Martin, J. (2007) Habitat deterioration affects body condition of lizards: A behavioral approach with *Iberolacerta cyreni* lizards inhabiting ski resorts. *Biological Conservation*, **135**, 77-85.
- Amrein, Y.U. & Amrein, M.B. (1951) The number of young found in the island night lizard. *Copeia*, **1951**, 180.
- Ananjeva, N.B. & Tuniev, B.S. (1994) Some aspects of historical biogeography of Asian rock agamids. *Russian Journal of Herpetology*, **1**, 42-52.
- Anaya-Rojas, J.M., Serrano-Cardozo, V.H. & Ramírez-Pinilla, M.P. (2010) Diet, microhabitat use, and thermal preferences of *Ptychoglossus bicolor* (Squamata: Gymnophthalmidae) in an organic coffee shade plantation in Colombia. *Papeis Avulsos de Zoologia*, **50**, 159-166.
- Anderson (1889) Report on the mammals, reptiles, and batrachians, chiefly from the Mergui Archipelago, collected for the Trustees of the Indian Museum. *Journal of the Linnean Society of London*, **20**, 331-350.
- Anderson (1895) On a collection of reptiles and batrachians made by Colonel Yerbury at Aden and its neighbourhood. *Proceedings of the Zoological Society of London*, **1895**, 635—663.
- Anderson, S.C. (1963) Amphibians and reptiles from Iran. *Proceedings of the California Academy of Sciences*, **31**, 417-498.

- Anderson, S.C. & Leviton, A.E. (1969) Amphibians and reptiles collected by the Street Expedition to Afghanistan, 1965. *Proceedings of the California Academy of Sciences*, **37**, 25-56.
- Anderson, R.A. & Vitt, L.J. (1990) Sexual selection versus alternative causes of sexual dimorphism in teiid lizards. *Oecologia*, **84**, 145-157.
- Andersson, M., Krockenberger, A. & Schwarzkopf, L. (2010) Experimental manipulation reveals the importance of refuge habitat temperature selected by lizards. *Austral Ecology*, **35**, 294-299.
- Andrade, D.V., Nascimento, L.B. & Abe, A.S. (2006) Habits hidden underground: a review on the reproduction of the *Amphisbaenia* with notes on four neotropical species. *Amphibia-Reptilia*, **27**, 207-217.
- Andreone, F. & Guarino, F.M. (2003) Giant and long-lived? Age structure in *Macroscincus coctei*, an extinct skink from Cape Verde. *Amphibia-Reptilia*, **24**, 459-470.
- Andreone, F., Glaw, F., Nussbaum, R., Raxworthy, C., Vences, M. & Randrianirina, J. (2003) The amphibians and reptiles of Nosy Be (NW Madagascar) and nearby islands: a case study of diversity and conservation of an insular fauna. *Journal of Natural History*, **37**, 2119-2149.
- Andrews, R.M. (1979) Evolution of life histories: a comparison of *Anolis* lizards from matched island and mainland habitats. *Breviora*, **454**, 1-51.
- Andrews, R.M. (1998) Geographic variation in field body temperature of *Sceloporus* lizards. *Journal of Thermal Biology*, **23**, 329-334.
- Andrews, R.M. (2008) Lizards in the slow lane: thermal biology of chameleons. *Journal of Thermal Biology*, **33**, 57-61.
- Andrews, R.M. & Rand, A.S. (1974) Reproductive effort in anoline lizards. *Ecology*, **55**, 1317-1327.
- Andrews, R.M. & Pough, F.H. (1985) Metabolism of squamate reptiles: allometric and ecological relationships. *Physiological Zoology*, **58**, 214-231.
- Andrews, R.M., Mathies, T.Q., C. P. & Qualls, F.J. (1999) Rates of embryonic development of *Sceloporus* lizards: do cold climates favor rapid development? *Copeia*, **1999**, 691-699.
- Angellucci, A., Civitelli, G., Funciello, R., Mariotti, G., Matteucci, R., Passeri, L., Piali, G., Praturlon, A. & Sirna, G. (1975) Preliminary report on the carbonate sedimentation at the Dahlak Islands (Red Sea, Ethiopia). *Geologica Roma*, **14**, 41-61.
- Angilletta, M.J. & Werner, Y.L. (1998) Australian geckos do not display diel variation in thermoregulatory behavior. *Copeia*, **1998**, 736-742.
- Aparicio, A., Arafia, V., Garcia, R. & Grachev, A. (1994) The origin of the Columbretes Islands' basanitic and phonolitic magmas (Western Mediterranean). *Mineralogical Magazine*, **58**, 21-22.
- Arad, Z. (1995) Physiological responses to increasing ambient temperature in three ecologically different, congeneric lizards (Gekkoninae: *Ptyodactylus*). *Comparative Biochemistry and Physiology*, **112A**, 305-311.
- Arena, P.C. & Wooller, R.D. (2003) The reproduction and diet of *Egernia kingii* (Reptilia: Scincidae) on Penguin Island, Western Australia. *Australian Journal of Zoology*, **51**, 495-504.
- Arnold, D.L. (1980) Geographic variation in *Anolis brevirostris* (Sauria: Iguanidae) in Hispaniola. *Breviora*, **461**, 1-31.
- Arnold, E.N. (1984) Ecology of lowland lizards in the eastern United Arab Emirates. *Journal of*

- Zoology*, **204**, 329-354.
- Arnold, E.N. (1986) A key and annotated checklist to the lizards and amphisbaenians of Arabia. *Fauna of Saudi Arabia*, **8**, 385-403.
- Arnold, E.N. (1987) Resource partition among lacertid lizards in southern Europe. *Journal of Zoology B*, **1**, 739-782.
- Arnold, E.N. (1989) Systematics and adaptive radiation of Equatorial African lizards assigned to the genera *Adolfus*, *Bedriagaia*, *Gastropholis*, *Holaspis* and *Lacerta* (Reptilia, Lacertidae). *Journal of Natural History*, **23**, 525-555.
- Arnold, E.N. (1990) The two species of Moroccan day-geckoes, *Quedenfeldtia* (Reptilia: Gekkonidae). *Journal of Natural History*, **24**, 757-762.
- Arnold, E.N. (1993) Historical changes in the ecology and behaviour of semaphore geckos (*Pristurus*, Gekkonidae) and their relatives. *Journal of Zoology*, **229**, 353-384.
- Arnold, E.N. (1994) Do ecological analogues assemble their as examples. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **344**, 277-290.
- Arnold, E.N. (1997) Interrelationships and evolution of the east Asian grass lizards, *Takydromus* (Squamata: Lacertidae). *Zoological Journal of the Linnean Society*, **119**, 267-296.
- Arnold, E.N. (1998) Structural niche, limb morphology and locomotion in lacertid lizards (Squamata, Lacertidae): a preliminary survey. *Bulletin of the Natural History Museum, London (Zoology)*, **64**, 63-89.
- Arnold, E.N. (2009) Relationships, evolution and biogeography of Semaphore geckos, *Pristurus* (Squamata, Sphaerodactylidae) based on morphology. *Zootaxa*, **2060**, 1-21.
- Arnold, E.N., Vasconcelos, R., Harris, D.J., Mateo, J.A. & Carranza, S. (2008) Systematics, biogeography and evolution of the endemic *Hemidactylus* geckos (Reptilia, Squamata, Gekkonidae) of the Cape Verde Islands: based on morphology and mitochondrial and nuclear DNA sequences. *Zoologica Scripta*, **37**, 619-636.
- Arribas, O. & Carranza, S. (2004) Morphological and genetic evidence of the full species status of *Iberolacerta cyreni martinezricai* (Arribas, 1996). *Zootaxa*, **634**, 1-24.
- Aryal, P.C., Pokhrel, G.K., Shah, K.B., Rijal, B., Kharel, S.C., Paudel, E., Suwal, M.K., Dhamala, M.K. & Bhurtel, B.P. (2010) *Inventory of herpetofaunal diversity in Nagarjun Forest of Shivapuri Nagarjun National Park*. Nepal Academy of Science and Technology (NAST), Lalitpur, Nepal.
- Ashton, K.G. (2005) Life history of a fossorial lizard, *Neoseps reynoldsi*. *Journal of Herpetology*, **39**, 389-395.
- Ashton, K.G. & Telford, S.R. (2006) Monthly and daily activity of a fossorial lizard, *Neoseps reynoldsi*. *Southeastern Naturalist*, **5**, 175-183.
- Asplund, K.K. (1967) Ecology of lizards in the relictual cape flora, Baja California. *American Midland Naturalist*, **77**, 462-475.
- Auffenberg, W. (1980) The herpetofauna of Komodo, with notes on adjacent areas. *Bulletin of the Florida State Museum Biological Sciences*, **25**, 39-156.
- Auffenberg, W. & Auffenberg, T. (1988) Resource partitioning in a community of Philippine skinks (Sauria: Scincidae). *Bulletin of the Florida State Museum, Biological Sciences*, **32**, 151-219.
- Auffenberg, W., Rahman, H., Iffat, F. & Perveen, Z. (1989) A study of *Varanus flavescens* (Hardwicke & Gray) (Sauria: Varanidae). *Journal of the Bombay Natural History Society*, **86**, 286-307.

- Avery, R.A. (1981) Feeding ecology of the nocturnal gecko *Hemidactylus brooki* in Ghana. *Amphibia-Reptilia*, **1**, 269-276.
- Avery, R.A. (1982) Field studies of body temperatures and thermoregulation. *Biology of the Reptilia*, **12**, 93-166.
- Avila, L.J. (2002) Geographic distribution of lizards of the Genus *Teius* (Squamata: Teiidae: Teiinae) in southern South America. *Biogeographica*, **78**, 15-33.
- Avila, L.J., Frutos, N., Morando, M., Perez, C.H.F. & Kozykariski, M. (2006) Reptilia, Iguania, Liolaemini, *Liolaemus petrophilus* and *Liolaemus pictus*: distribution extension, filling gaps, new records. *Check List*, **2**, 65-69.
- Avila-Pires, T.C.S., Hoogmoed, M.S. & da Rocha, W.A. (2010) Notes on the vertebrates of northern Para, Brazil: a forgotten part of the Guyanian region I. Herpetofauna. *Boletim do Museu Paraense Emilio Goeldi Ciencias Naturais*, **5**, 13-112.
- Axtell, R.W. (1956) A solution to the long neglected *Holbrookia lacerata* problem, and the description of two new subspecies of *Holbrookia*. *Bulletin of the Chicago Academy of Sciences*, **10**, 163-179.
- Axtell, R.W. (1983) *Holbrookia propinqua*. *Catalogue of American Amphibians and Reptiles*, **341**, 1-2.
- Axtell, R.W. & Axtell, C.A. (1971) A new lizard (*Sceloporus jarrovii cyanostictus*) from Sierra Madre of Coahuila, Mexico. *Copeia*, **1971**, 89-98.
- Baha El Din, S.M. (1996) The occurrence of *Acanthodactylus longipes* Boulenger, 1918 in Egypt, with remarks on its identification and ecology. *Zoology in the Middle East*, **12**, 53-58.
- Bahir, M.M. & Maduwage, K.P. (2005) *Calotes desilvai*, a new species of agamid lizard From Morningside forest, Sri Lanka. *Raffles Bulletin of Zoology*, **12 (supplement)**, 381-392.
- Baker, R.H., Webb, R.G. & Dalby, P. (1967) Notes on reptiles and mammals from southern Zacatecas. *American Midland Naturalist*, **77**, 223-226.
- Balestrin, R.L. (2008) *Historia natural de uma taxocenose de Squamata e redescricao de uma especie de anuro no escudo sul-riograndense, Brasil*. Doctoral Thesis, Porto Alegre.
- Balestrin, R.L., Cappellari, L.H. & Outeiral, A.B. (2010) Biologia reprodutiva de *Cercosaura schreibersii* (Squamata, Gymnophthalmidae) e *Cnemidophorus lacertoides* (Squamata, Teiidae) no Escudo Sul-Riograndense, Brasil. *Biota Neotropica*, **10**, 131-139.
- Ballinger, R.E. (1977) Reproductive strategies: food availability as a source of proximal variation in a lizard. *Ecology*, **58**, 628-635.
- Ballinger, R.E. & Schrank, G.D. (1972) Reproductive potential of female whiptail lizards, *Cnemidophorus gularis gularis*. *Herpetologica*, **28**, 217-222.
- Ballinger, R.E., Lemos-Espinal, J.A. & Smith, G.R. (2000) Reproduction in females of three species of crevice-dwelling lizards (genus *Xenosaurus*) from Mexico. *Studies on Neotropical Fauna and Environment*, **35**, 179-183.
- Banks, R.C. & Farmer, W.M. (1963) Observations on reptiles of Cerralvo Island, Baja California, Mexico. *Herpetologica*, **18**, 246-250.
- Barbault, R. (1974) Ecologie comparee des lezards *Mabuya blandingi* (Hallowell) et *Panaspis kitsoni* (Boulenger) dans les forets de Lamto (Cote d'Ivoire). *La Terre et La Vie*, **28**, 272-295.
- Barbault, R. (1974) Structure et dynamique d'un peuplement de lezards: les scincides de la savane de Lamoto (Cote d'Ivoire). *Terre Vie*, **28**, 352-428.
- Barbault, R. (1976) Population dynamics and reproductive patterns of three African skinks.

- Copeia*, **1976**, 483-490.
- Barbault, R. & Maury, M.E. (1981) Ecological organization of a Chihuahuan desert lizard community. *Oecologia*, **51**, 335-342.
- Barbault, R., Ortega, A. & Maury, M.E. (1985) Food partitioning and community organization in a mountain lizard guild of northern Mexico. *Oecologia*, **65**, 550-554.
- Barbour, T. & Loveridge, A. (1928) A comparative study of the herpetological fauna of the Uluguru and Usambara mountains, Tanzania Territory with descriptions of new species. *Memoires of the Museum of Comparative Zoology*, **50**, 85-265.
- Barbour, H.R., Archer, M.A., Hart, N.S., Thomas, N., Dunlop, S.A., Beazley, L.D. & Shand, J. (2002) Retinal characteristics of the ornate dragon lizard, *Ctenophorus ornatus*. *Journal of Comparative Neurology*, **450**, 334-344.
- Barden, G. & Shine, R. (1994) Effects of sex and reproductive mode on dietary composition of the reproductively bimodal scincid lizard, *Lerista bougainvillii*. *Australian Zoologist*, **29**, 225-228.
- Bartholomew, G.A. & Tucker, V.A. (1964) Size, body temperature, thermal conductance, oxygen consumption, and heart rate in Australian varanid lizards. *Physiological Zoology*, **37**, 341-354.
- Bartlett, R.D. (1985) Notes on the natural history and reproductive strategy of the island glass lizard, *Ophisaurus compressus*. *British Herpetological Society Bulletin*, **11**, 19-21.
- Barts, M. (2003) Die agamen des sudlichen Afrikas. *Draco*, **4**, 70-79.
- Bauer, A.M. (1985) Notes on the taxonomy, morphology and behavior of *Rhacodactylus chahoua* (Bavay) (Reptilia: Gekkonidae). *Banner zoologische Beitrage*, **36**, 81-94.
- Bauer, A.M. (1990) Gekkonid lizards as prey of invertebrates and predators of vertebrates. *Herpetological Review*, **21**, 83-87.
- Bauer, A.M. (2006) A review of the gekkotan lizards of Benin, with the description of a new species of *Hemidactylus* (Squamata: Gekkonidae). *Zootaxa*, **1242**, 1-20.
- Bauer, A.M. & DeVaney, K.D. (1987) Aspects of diet and habitat use in some New Caledonian lizards. *Amphibia-Reptilia*, **8**, 349-364.
- Bauer, A.M. & Vindum, J.V. (1990) A checklist and key to the herpetofauna of New Caledonia, with remarks on biogeography. *Proceedings of the California Academy of Sciences*, **47**, 17-45.
- Bauer, A.M. & Gunther, R. (1992) A preliminary report of the reptile fauna of the kingdom of Bhutan with the description of a new species of scincid lizard (Reptilia: Scincidae). *Asiatic Herpetological Research*, **4**, 23-36.
- Bauer, A.M. & Russell, A.P. (1993) *Aristelliger barbouri*. *Catalogue of American Amphibians and Reptiles*, **566**, 1.
- Bauer, A.M. & Jackman, T.R. (2008) Global diversity of lizards in freshwater (Reptilia: Lacertilia). *Hydrobiologia*, **595**, 581-586.
- Bauer, A.M., Shea, G. & Gunther, R. (2003) An annotated catalogue of the types of scincid lizards (Reptilia, Squamata, Scincidae) in the collection of the Museum fur Naturkunde der Humboldt-Universitat zu Berlin (ZMB). *Mitteilungen aus dem Museum fur Naturkunde in Berlin*, **79**, 253-321.
- Bauer, A.M., Barts, M. & Hulbert, F. (2006) A new species of the *Pachydactylus weberi* group (Reptilia: Squamata: Gekkonidae) from the Orange River, with comments on its natural history. *Salamandra*, **42**, 83-92.
- Bauer, A.M., Jackman, T., Sadlier, R.A. & Whitaker, A.H. (2009) Review and phylogeny of the

- New Caledonian diplodactylid gekkotan genus *Eurydactylodes* Wermuth, 1965, with the description of a new species. *Memoires du Museum National d'Histoire Naturelle*, **198**, 13-36.
- Bauer, A.M., Glaw, F., Gehring, P.-S. & Vences, M. (2011) New species of *Blaesodactylus* (Squamata: Gekkonidae) from Ankarafantsika National Park in north-western Madagascar. *Zootaxa*, **2942**, 57-68.
- Bauwens, D. (1999) Life-history variations in lacertid lizards. *Natura Croatica*, **8**, 239-252.
- Bauwens, D. & Diaz-Uriarte, R. (1997) Covariation of life-history traits in lacertid lizards: a comparative study. *American Naturalist*, **149**, 91-111.
- Bauwens, D., Castilla, A.M., Van Damme, R. & Verheyen, R.F. (1990) Field body temperatures and thermoregulatory behaviour of the high altitude lizard, *Lacerta bedriagae*. *Journal of Herpetology*, **24**, 88-91.
- Beccaluva, L., Bianchini, G. & Wilson, M. (2011) *Volcanism and Evolution of the African Lithosphere*. Geological Society of America.
- Beck, D.D. (1990) Ecology and behavior of the gila monster in southwestern Utah. *Journal of Herpetology*, **24**, 54-68.
- Beebe, W. (1945) Field notes on the lizards of Kartabo, British Guiana, and Caripito, Venezuela. Part 3, Teiidae, Amphisbaenidae and Scincidae. *Zoologica*, **30**, 7-32.
- Bejakovic, D., Kalezic, M.L., Aleksic, I., Dzukic, G. & CrnobrnjaIsailovic, J. (1995) Female reproductive cycle and clutch traits in the Dalmatian wall lizard (*Podarcis melisellensis*). *Folia Zoologica*, **44**, 371-380.
- Benabib, M. (1994) Reproduction and lipid utilization of tropical populations of *Sceloporus variabilis*. *Herpetological Monographs*, **8**, 160-180.
- Benabib, M., Kjer, K.M. & Sites, J.W. (1997) Mitochondrial DNA sequence-based phylogeny and the evolution of viviparity in the *Sceloporus scalaris* group (Reptilia, Squamata). *Evolution*, **51**, 1262-1275.
- Bennett, A.F. & Gorman, G.C. (1979) Population density and energetics of lizards on a tropical island. *Oecologia*, **42**, 339-358.
- Bennett, A.F. & John-Alder, H. (1986) Thermal relations of some Australian skinks (Sauria: Scincidae). *Copeia*, **1986**, 57-64.
- Bentz, E.J., Rodriguez, M.J.R., John, R.R., Henderson, R.W. & Powell, R. (2011) Population densities, activity, microhabitats, and thermal biology of a unique crevice- and litter-dwelling assemblage of reptiles on Union Island, St. Vincent and the Grenadines. *Herpetological Conservation and Biology*, **6**, 40-50.
- Bergmann, P.J., Meyers, J.J. & Irschick, D.J. (2009) Directional evolution of stockiness coevolves with ecology and locomotion in lizards. *Evolution*, **63**, 215-227.
- Bernardo-Silva, J.S., Von-Muhlen, E.M., Di-Bernardo, M. & Ketterl, J. (2006) Feeding ecology in the small neotropical amphisbaenid *Amphisbaena munoai* (Amphisbaenidae) in southern Brazil. *Iheringia, Serie Zoologia*, **96**, 487-489.
- Bezy, R.L. (1967) Variation, distribution, and taxonomic status of the Arizona night lizard (*Xantusia arizonae*). *Copeia*, **1967**, 653-661.
- Bezy, R.L. (1989) Morphological differentiation in unisexual and bisexual xantusiid lizards of the genus *Lepidophyma* in Central America. *Herpetological Monographs*, **3**, 61-80.
- Bezy, R.L. (2002) Systematics of xantusiid lizards of the genus *Lepidophyma*. *Los Angeles County Museum Contributions in Science*, **493**, 1-41.
- Bjursell, A. (2001) *Ctenotus robustus*. Australian Herpetological Directory, Herpetofauna of

North Queensland.

- Blackburn, D.G., Vitt, L.J. & Beuchat, C.A. (1984) Eutherian-like reproductive specializations in a viviparous reptile. *Proceedings of the National Academy of Sciences USA*, **81**, 4860-4863.
- Bodenheimer, F.S. (1944) Introduction into the knowledge of the amphibia and reptilia of Turkey. *Revue de la Faculte des Sciences de l'Universite d'Istanbul*, **9B**, 1-93.
- Bogaerts, S. (2006) First data on the reproduction of Lanza's skink, *Chalcides lanzai* Pasteur, 1967. *Podarcis*, **7**, 9-16.
- Bogdanov, N.A. (1984) *Origin and History of Marginal and Inland Seas: Proceedings of the 27th International Geological Congress, Moscow 4-14 August 1984*. VNU Science Press.
- Bogert, C.M. & Oliver, J.A. (1945) A preliminary analysis of the herpetofauna of Sonora. *Bulletin of the American Museum of Natural History*, **83**, 297-426.
- Bohme, W. & Klaver, C. (1981) Zur innerartlichen gliederung und zur artgeschichte von *Chamaeleo quadricornis* Tornier, 1899 (Sauria: Chamaeleonidae). *Amphibia-Reptilia*, **1**, 313-328.
- Bohme, W., Meinig, H. & Rodel, M.-O. (1996) New records of amphibians and reptiles from Burkina Faso and Mali. *British Herpetological Society Bulletin*, **56**, 7-26.
- Bohme, W., Rodel, M.-O., Brede, C. & Wagner, P. (2011) The reptiles (Testudines, Squamata, Crocodylia) of the forested southeast of the Republic of Guinea (Guinee forestiere), with a country-wide checklist. *Bonn Zoological Bulletin*, **60**, 35-61.
- Bombi, P. & Vignoli, L. (2004) Distribution, ecology and conservation of *Archaeolacerta bedriagae* in Sardinia (Reptilia, Lacertidae). *Italian Journal of Zoology*, **71**, 135-144.
- Bonino, M.N., Azocar, D.L.M., Tulli, M.J., Abdala, C.S., Perotti, M.G. & Cruz, F.B. (2011) Running in cold weather: morphology, thermal biology, and performance in the southernmost lizard clade in the world (*Liolaemus lineomaculatus* section: Liolaemini: Iguania). *Journal of Experimental Zoology*, **315**, 495-503.
- Boretto, J.M. & Ibarquengoytia, N.R. (2006) Asynchronous spermatogenesis and biennial female cycle of the viviparous lizard *Phymaturus antofagastensis* (Liolaemidae): reproductive responses to high altitudes and temperate climate of Catamarca, Argentina. *Amphibia-Reptilia*, **27**, 25-36.
- Boretto, J.M., Ibarquengoytia, N.R., Acosta, J.C., Blanco, G.M., Villavicencio, J. & Marinero, J.A. (2007) Reproductive biology and sexual dimorphism of a high-altitude population of the viviparous lizard *Phymaturus punae* from the Andes in Argentina. *Amphibia-Reptilia*, **28**, 427-432.
- Boulenger, G.A. (1889) On the reptiles and batrachians obtained in Morocco by M. Henry Vaucher. *Annals and Magazine of Natural History*, **3**, 303-307.
- Boulenger, G.A. (1890) First report on additions to the lizard collection in the British Museum (Natural History). *Proceedings of the Zoological Society of London*, **1890**, 77-86.
- Boulenger, G.A. (1896) Descriptions of new reptiles and batrachians from Colombia. *Annals and Magazine of Natural History*, **17**, 16-21.
- Boulenger, G.A. (1897) A catalogue of the reptiles and batrachians of Celebes with special reference to the collections made by Drs P & F Sarasin in 1893-1896. *Proceedings of the Zoological Society*, **1897**, 193-237.
- Boulenger, G.A. (1901) Further descriptions of new reptiles collected by Mr. P. O. Simons in Peru and Bolivia. *Annals and Magazine of Natural History*, **7**, 546-549.
- Bouysse, P., Westercamp, D. & Andreieff, P. (1990) The Lesser Antilles island arc.

- Proceedings of the Ocean Drilling Program* (ed by J.C. Moore, A. Mascle and E. Al.). College Station.
- Bowker, R.G. (1984) Precision of thermoregulation of some African lizards. *Physiological Zoology*, **57**, 401-412.
- Bowker, R.G., Wright, C.L. & Bowker, G.E. (2010) Patterns of body temperatures: Is lizard thermoregulation chaotic? *Journal of Thermal Biology*, **35**, 1-5.
- Bowker, R.G., Damschroder, S., Sweet, A.M. & Anderson, D.K. (1986) Thermoregulatory behavior of the north american lizards *Cnemidophorus velox* and *Sceloporus undulatus*. *Amphibia-Reptilia*, **7**, 335-346.
- Boylan, T. (1989) Reproduction of the Fijian crested iguana, *Brachylophus vitiensis* at Taronga Zoo. *International Zoo Yearbook*, **28**, 126-130.
- Bradshaw, S.D. & Main, A.R. (1968) Behavioural attitudes and regulation of temperature in *Amphibolurus* lizards. *Journal of Zoology*, **154**, 193-221.
- Brain, C.K. (1962) Observations on the temperature tolerance of lizards in the central Namib Desert, South West Africa. *Cimbebasia*, **4**, 1-5.
- Brana, F., Gonzalez, F. & Barahona, A. (1992) Relationship between ovarian and fat body weights during vitellogenesis for three species of lacertid lizards. *Journal of Herpetology*, **26**, 515-518.
- Branch, W.R. & McCartney, C.J. (1992) A report on a small collection of reptiles from southern Angola. *Journal of the Herpetological Association of Africa*, **41**, 1-3.
- Branch, W.R., Rodel, M.O. & Marais, J. (2005) A new species of rupicolous *Cordylus Laurenti* 1768 (Sauria: Cordylidae) from Northern Mozambique. *African Journal of Herpetology*, **54**, 131-138.
- Brandt, R. & Navas, C.A. (2011) Life-history evolution on tropidurinae lizards: influence of lineage, body size and climate. *PLoS ONE*, **6**, e20040.
- Brattstrom, B.H. (1952) The food of the nightlizards, genus *Xantusia*. *Copeia*, **1952**, 168-172.
- Broadley, D.G. (1968) A review of the African genus *Typhlosaurus* Wiegmann (Sauria: Scincidae). *Arnoldia*, **3**, 1-20.
- Broadley, D.G. (1971) The reptiles and amphibians of Zambia. *Puku*, **6**, 1-143.
- Broadley, D.G. (2000) A review of the genus *Mabuya* in southeastern Africa (Sauria: Scincidae). *African Journal of Herpetology*, **49**, 87-110.
- Broadley, D.G. & Bauer, A.M. (1999) A review of the *Mabuya quinquetaeniata* complex in East Africa (Sauria: Scincidae). *African Journal of Herpetology*, **47**, 43-58.
- Brooke, M.D.L. & Houston, D.C. (1983) The biology and biomass of the skinks *Mabuya sechellensis* and *M. wrightii* on Cousin Island, Seychelles. *Journal of Zoology*, **200**, 179-195.
- Brooks, G.R. (1968) Body temperatures of three lizards from Dominica, West Indies. *Herpetologica*, **24**, 209-214.
- Brown, W.C. (1953) Results of the Archbold Expeditions. No. 69. A review of New Guinea lizards allied to *Emoia baudini* and *Emoia physicae* (Scincidae). *American Museum Novitates*, **1627**, 1-26.
- Brown, W.C. (1956) A revision of the genus *Brachymeles* (Scincidae), with descriptions of new species and subspecies. *Breviora*, **54**, 1-19.
- Brown, G.W. (1991) Ecological feeding analysis of south-eastern Australian scincids (Reptilia: Lacertilia). *Australian Journal of Zoology*, **39**, 9-29.
- Brown, W.C. & Alcalá, A.C. (1961) Populations of amphibians and reptiles in the submontane

- and montane forests of Cuernos de Negros, Philippine Islands. *Ecology*, **42**, 628-636.
- Brown, W.C. & Gibbons, J.R.H. (1986) Species of the *Emoia samoensis* group of lizards (Scincidae) in the Fiji Islands, with descriptions of two new species. *Proceedings of the California Academy of Sciences*, **44**, 41-53.
- Brown, R.P. & Roberts, N. (2008) Feeding state and selected body temperatures in the slow-worm (*Anguis fragilis*). *Herpetological Journal*, **18**, 59-62.
- Brown, R.P., Znari, M., El Mouden, E.H. & Harris, P. (1999) Estimating asymptotic body size and testing geographic variation in *Agama impalearis*. *Ecography*, **22**, 277-283.
- Bruna, E.M., Fisher, R.N. & Case, T.J. (1996) New evidence of habitat segregation between two cryptic species of Pacific skinks (*Emoia cyanura* and *E. impar*). *Copeia*, **1996**, 998-1005.
- Bryan, W.B. (1967) Geology and petrology of Clarion Island, Mexico. *Geological Society of America Bulletin*, **78**, 1461-1476.
- Buden, D.W. (1998) Morphological variation and distributional ecology of the giant Micronesian gecko (*Perochirus scutellatus*) of Kapingamarangi Atoll. *Pacific Science*, **52**, 250-258.
- Buden, D.W. (1999) Reptiles, birds, and mammals of Oroluk Atoll, Eastern Caroline Islands. *Micronesica*, **31**, 289-300.
- Buden, D.W. (2000) The reptiles of Pohnpei, Federated States of Micronesia. *Micronesica*, **32**, 155-180.
- Buden, D.W. (2007) A new species of the genus *Lepidodactylus* Fitzinger (Squamata: Gekkonidae) from the Mortlock Islands, Chuuk State, Federated States of Micronesia. *Pacific Science*, **61**, 407-414.
- Bueter, C. & Haas, A. (2008) Living the high life: *Sceloporus malachiticus* from high elevations perform better at extreme temperatures. *Eukarion*, **4**, 112-114.
- Bull, C.M. & Bonnett, M. (2004) *Egernia striolata* (tree skink) reproduction. *Herpetological Review*, **35**, 389.
- Bullock, D.J. (1986) The ecology and conservation of reptiles on Round Island and Gunner's Quoin, Mauritius. *Biological Conservation*, **37**, 135-156.
- Bullock, D.J., Arnold, E.N. & Bloxam, Q. (1985) A new endemic gecko (Reptilia: Gekkonidae) from Mauritius. *Journal of Zoology*, **206**, 591-599.
- Burkett, R.D. (1962) Two clutches of eggs in the lizard, *Gerrhonotus liocephalus infernalis*. *Herpetologica*, **18**, 211.
- Burstein, N., Larsen, K.R. & Smith, H.M. (1974) A preliminary survey of dermatoglyphic variation in the lizard genus *Sceloporus*. *Journal of Herpetology*, **8**, 359-369.
- Burt, C.E. (1930) Herpetological results of the Whitney South Sea Expedition. 4, Descriptions of new species of lizards from the Pacific Islands (Scincidae). *American Museum Novitates*, **427**, 1-3.
- Busack, S.D. (1976) Activity cycles and body temperatures of *Acanthodactylus erythrurus*. *Copeia*, **1976**, 826-830.
- Busack, S.D. (1986) Taxonomic implications of biochemical and morphological differentiation in Spanish and Moroccan populations of three-toed skinks, *Chalcides chalcides* (Lacertilia, Scincidae). *Herpetologica*, **42**, 230-236.
- Busack, S.D. (1987) Notes on the biology of *Lacerta andreanszkyi* (Reptilia: Lacertidae). *Amphibia-Reptilia*, **8**, 231-236.
- Busack, S.D. & Visnaw, J.A. (1989) Observations on the natural history of *Lacerta lepida* in

- Cadiz Province, Spain. *Amphibia-Reptilia*, **10**, 201-213.
- Bush, B. (1992) Some records of reproduction in captive lizards and snakes. *Herpetofauna*, **22**, 26-30.
- Bush, A.M., Bambach, R.K. & Daley, G.M. (2007) Changes in theoretical ecospace utilization in marine fossil assemblages between the mid-Paleozoic and late Cenozoic. *Paleobiology*, **33**, 76-97.
- Bustard, H.R. (1966) Notes on the eggs, incubation and young of the bearded dragon *Amphibolurus barbatus* (Cuvier). *British Journal of Herpetology*, **3**, 252-259.
- Bustard, H.R. (1967) Defensive display behavior of the Australian gecko *Nephruurus asper*. *Herpetologica*, **23**, 126-129.
- Bustard, H.R. (1968) The reptiles of Merriwindi State Forest, Pilliga West, northern New South Wales, Australia. *Herpetologica*, **24**, 131-140.
- Bustos-Zagal, M.G., de la Cruz, F.R.M., Castro-Franco, R. & Cruz, M.V.-S. (2011) Reproductive cycle of *Sceloporus ochoterena* from Morelos State, Mexico. *Revista Mexicana de Biodiversidad*, **82**, 589-597.
- Cabezas, C.F., Boretto, J.M., Acosta, J.C., Jahn, G., Blanco, G., Laspiur, A. & Ibarquengoytia, N.R. (2010) Reproductive biology of *Phymaturus* cf. *palluma*: a vulnerable lizard from the highlands of the Andes, San Juan, Argentina. *Herpetological Conservation and Biology*, **5**, 430-440.
- Campbell, H.W. (1973) Ecological observations on *Anolis lionotus* and *Anolis poecilopus* (Reptilia, Sauria) in Panama. *American Museum Novitates*, **2516**, 1-29.
- Canseco-Marquez, L., Gutierrez-Mayen, G. & Salazar-Arenas, J. (2000) New records and range extensions for amphibians and reptiles from Puebla, Mexico. *Herpetological Review*, **31**, 259-263.
- Cappellari, L.H., de Lema, T., Prates, P. & da Rocha, C.F.D. (2007) Diet of *Teius oculatus* (Sauria, Teiidae) in southern Brazil (Dom Feliciano, Rio Grande do Sul). *Iheringia, Serie Zoologia*, **97**, 31-35.
- Cappellari, L.H., Balestrin, R.L., de Lema, T. & Rocha, C.F.D. (2011) Reproductive biology of *Teius oculatus* (Sauria, Teiidae) in Southern Brazil (Dom Feliciano, Rio Grande do Sul). *North-Western Journal of Zoology*, **7**, 270-276.
- Capula, M. & Luiselli, L. (1994) Resource partitioning in a Mediterranean lizard community. *Bolletino di Zoologia*, **61**, 173-177.
- Capula, M., Luiselli, L., Filippi, E. & Ceccarelli, A. (2002) Habitat characteristics and demography of high altitude populations of the lacertid lizard, *Algyroides fitzingeri*. *Italian Journal of Zoology*, **69**, 33-36.
- Carey, W.M. (1975) Rock iguana, *Cyclura pinguis*, on Anegada, British Virgin Islands, with notes on *Cyclura ricordi* and *Cyclura cornuta* on Hispaniola. *Bulletin of the Florida State Museum Biological Sciences*, **19**, 189-233.
- Carpenter, C.C. (1969) Behavioral and ecological notes on the Galapagos land iguanas. *Herpetologica*, **25**, 155-164.
- Carranza, S., Arnold, E.N., Mateo, J.A. & Lopez-Jurado, L.F. (2001) Parallel gigantism and complex colonization patterns in the Cape Verde scincid lizards *Mabuya* and *Macroscincus* (Reptilia : Scincidae) revealed by mitochondrial DNA sequences. *Proceedings of the Royal Society of London B: Biological Sciences*, **268**, 1595-1603.
- Carranza, S., Arnold, E.N., Geniez, P., Roca, J. & Mateo, J.A. (2008) Radiation, multiple dispersal and parallelism in the skinks, *Chalcides* and *Sphenops* (Squamata: Scincidae),

- with comments on *Scincus* and *Scincopus* and the age of the Sahara Desert. *Molecular Phylogenetics and Evolution*, **46**, 1071-1094.
- Carretero, M.A. (2008) An integrated assessment of a group with complex systematics: the Iberomaghrebian lizard genus *Podarcis* (Squamata, Lacertidae). *Integrative Zoology*, **4**, 247-266.
- Carretero, M.A. & Llorente, G.A. (1993) Feeding of two sympatric lacertids in a sandy coastal area (Ebro Delta, Spain). *Conference Proceedings*, 155-172.
- Carretero, M.A. & Lo Cascio, P. (2010) What do myrmecophagous geckos eat when ants are not available? Comparative diets of three *Socotran* species. *African Zoology*, **45**, 115-120.
- Carretero, M.A., Cascio, P.L., Corti, C. & Pasta, S. (2010) Sharing resources in a tiny Mediterranean island? Comparative diets of *Chalcides ocellatus* and *Podarcis filfolensis* in Lampione. *Bonn Zoological Bulletin*, **57**, 111-118.
- Carretero, M.A., Perera, A., Harris, D.J., Batista, V. & Pinho, C. (2006) Spring diet and resource partitioning in an alpine lizard community from Morocco. *African Zoology*, **41**, 113-122.
- Cascio, P.L. (2010) Field body temperatures in a micro-insular lizard community (Squamata Sauria). *Il Naturalista Siciliano*, **34**, 21-27.
- Cascio, P.L. & Capula, M. (2011) Does diet in lacertid lizards reflect prey availability? Evidence for selective predation in the Aeolian wall lizard, *Podarcis raffonei* (Mertens, 1952) (Reptilia, Lacertidae). *Biodiversity Journal*, **2**, 89-96.
- Case, T.J. (1975) Species numbers, density compensation, and colonizing ability of lizards on islands in the the Gulf of California. *Ecology*, **56**, 3-18.
- Case, T.J., Cody, M.L. & Ezcurra, E. (2002) *A new island biogeography of the Sea of Cortés*. Oxford University Press, USA.
- Cassimiro, J. & Rodrigues, M.T. (2009) A new species of lizard genus *Gymnodactylus* Spix, 1825 (Squamata: Gekkota: Phyllodactylidae) from Serra do Sincoro, northeastern Brazil, and the status of *G. carvalhoi* Vanzolini, 2005. *Zootaxa*, **2008**, 38-52.
- Castanet, J. (1994) Age estimation and longevity in reptiles. *Gerontology*, **40**, 174-192.
- Castilla, A.M. & Bauwens, D. (1991) Observations on the Natural History, Present Status, and Conservation of the Insular Lizard *Podarcis hispanica atrata* on the Columbretes Archipelago, Spain. *Biological Conservation*, **58**, 69-84.
- Castilla, A.M. & Bauwens, D. (2000) Reproductive characteristics of the island lacertid Lizard *Podarcis lilfordi*. *Journal of Herpetology*, **34**, 390-396.
- Castro-Herrera, F., Bolivar-Garcia, W. & Herrera-Montes, M.I. (2007) *Guia de los anfibios y reptiles del Bosque de Yotoco, Valle del Cauca – Colombia*. Universidad del Valle, Cali, Colombia.
- Cei, J.M. (1982) A new endemic lizard from Sierra Pie de Palo in western Argentina. *Journal of Herpetology*, **16**, 179-182.
- Cei, J.M. (1993) Reptiles del noroeste, nordeste y este de la Argentina. *Museo Regionale di Scienze Naturali, Torino, Monografie*, **14**, 1-949.
- Cei, J.M. & Castro, L.P. (1973) Taxonomic and serological researches on the *Phymaturus patagonicus* complex. *Journal of Herpetology*, **7**, 237-247.
- Cei, J.M. & Scolaro, J.A. (1996) A new species of *Liolaemus* of the *Archeفورus* group from the precordilleran valley of the Zeballos river, Santa Cruz Province, Argentina (Reptilia, Tropiduridae). *Museo Regionale di Scienze Naturali Bollettino (Torino)*, **14**, 389-401.
- Cei, J.M.V., F. & Vicente, L. (2003) From oviparity to viviparity: a preliminary note on the

- morphometric differentiation between oviparous and viviparous species assigned to the genus *Liolaemus* (Reptilia, Squamata, Liolaemidae). *Journal of Zoological Systematics and Evolutionary Research*, **41**, 152-156.
- Cei, J.M., Etheridge, R. & Videla, F. (1983) Especies nuevas de iguanidos del noroeste de la provincia de San Juan (Reserva provincial San Guillermo), Argentina. *Deserta*, **7**, 316-323.
- Censky, E.J. (1995) Reproduction in two Lesser Antillean populations of *Ameiva plei* (Teiidae). *Journal of Herpetology*, **29**, 553-560.
- Chamaille-Jammes, S., Massot, M., Aragon, P. & Clobert, J. (2006) Global warming and positive fitness response in mountain populations of common lizards *Lacerta vivipara*. *Global Change Biology*, **12**, 392-402.
- Chandramouli, S.R. & Ganesh, S.R. (2011) Herpetofauna of southern Western Ghats, India, reinvestigated after decades. *Taprobanica*, **2**, 72-85.
- Chapman, A. & Dell, J. (1985) Biology and zoogeography of the amphibians and reptiles of the Western Australian wheatbelt. *Records of the Western Australian Museum*, **12**, 1-46.
- Chapple, D.G. (2003) Ecology, life-history, and behavior in the Australian Scincid genus *Egernia*, with comments on the evolution of complex sociality in lizards. *Herpetological Monographs*, **17**, 145-180.
- Chapple, D.G. (2005) Life history and reproductive ecology of White's skink, *Egernia whitii*. *Australian Journal of Zoology*, **53**, 353-360.
- Chapple, D.G. & Patterson, G.B. (2007) A new skink species (*Oligosoma taumakae* sp. nov.; Reptilia: Scincidae) from the Open Bay Islands, New Zealand. *New Zealand Journal of Zoology*, **34**, 347-357.
- Chavez, G., Siu-Ting, K., Duran, V. & Venegas, P.J. (2011) Two new species of Andean gymnophthalmid lizards of the genus *Euspondylus* (Reptilia, Squamata) from central and southern Peru. *ZooKeys*, **109**, 1-17.
- Chen, X.-J., Xu, X.F. & Ji, X. (2003) Influence of body temperature on food assimilation and locomotor performance in white-striped grass lizards, *Takydromus wolteri* (Lacertidae). *Journal of Thermal Biology*, **28**, 385-391.
- Chirio, L. (2009) Inventaire des reptiles de la region de la Reserve de Biosphere Transfrontaliere du W (Niger/Benin/Burkina Faso: Afrique de l'Ouest). *Bulletin de la Societe Herpetologique de France*, **132**, 13-41.
- Christian, K.A. & Tracy, C.R. (1985) Physical and biotic determinants of space utilization by the Galapagos land iguana (*Conolophus pallidus*). *Oecologia*, **66**, 132-140.
- Christian, K.A. & Weavers, B.W. (1996) Thermoregulation of monitor lizards in Australia: an evaluation of methods in thermal biology. *Ecological Monographs*, **66**, 139-157.
- Ciofi, C. & De Boer, M. (2004) Distribution and conservation of the Komodo monitor *Varanus komodoensis*. *Herpetological Journal*, **14**, 99-107.
- Cisse, M. & Karns, D.R. (1978) Saurians from Senegal. *Bulletin de l'Institut Fondamental d'Afrique Noire Serie A Sciences Naturelles*, **40**, 144-211.
- Cisse, M., Karns, D.R. & Karns, K.C. (1977) Aspects of the ecology of *Acanthodactylus dumerili* Sauria Lacertidae in Senegal. *Bulletin de l'Institut Fondamental d'Afrique Noire Serie A Sciences Naturelles*, **39**, 190-218.
- Civantos, E., Martin, J. & Lopez, P. (2003) Fossorial life constrains microhabitat selection of the amphibaenian *Trogonophis wiegmanni*. *Canadian Journal of Zoology*, **81**, 1839-1844.
- Clark, D.R. (1973) Temperature responses of three Costa Rican lizards (*Anolis*). *Caribbean Journal of Science*, **13**, 199-206.

- Clark, R. (1990) A report on herpetological observations in Afghanistan. *British Herpetological Society Bulletin*, **33**, 20-24.
- Clarke, S. (2000) *Management of the pygmy bluetongue lizard (Tiliqua adelaidensis) on private grazing properties, Mid-North SA*. Pygmy Bluetongue Recovery Program, South Australian Museum, North Terrace, Adelaide, Australia.
- Clausen, R.T. (1938) Notes on Eumeces anthracinus in Central New York. *Copeia*, **1938**, 3-7.
- Clemann, N., Chapple, D.G. & Wainer, J. (2004) Sexual dimorphism, diet, and reproduction in the Swamp Skink, *Egernia coventryi*. *Journal of Herpetology*, **38**, 461-467.
- Clemente, C.J., Thompson, G.G. & Withers, P.C. (2009) Evolutionary relationships of sprint speed in Australian varanid lizards. *Journal of Zoology*, **278**, 270-280.
- Clobert, J., Garland, T. & Barbault, R. (1998) The evolution of demographic tactics in lizards: a test of some hypotheses concerning life history evolution. *Journal of Evolutionary Biology*, **11**, 329-364.
- Clusella-Trullas, S.T., J. S., Blackburn, T.M. & Chown, S.L. (2008) Testing the thermal melanism hypothesis: a macrophysiological approach. *Functional Ecology*, **22**, 232-238.
- Colli, G.R. (1991) Reproductive ecology of *Ameiva ameiva* (Sauria, Teiidae) in the Cerrado of central Brazil. *Copeia*, **1991**, 1002-1012.
- Colli, G.R., Caldwell, J.P., Costa, G.C., Gainsbury, A.M., Garda, A.A., Mesquita, D.M., Filho, C.M.M.R., Soares, A.H.B., Silva, V.N., Valdujo, P.H., Vieira, G.H.C., Vitt, L.J., Werneck, F.P., Wiederhecker, H.C. & Zatz, M.G. (2003) A new species of *Cnemidophorus* (Squamata, Teiidae) from the cerrado biome in central Brazil. *Occasional Papers of the Sam Noble Oklahoma Museum of Natural History, University of Oklahoma*, **14**, 1-14.
- Commins, M.L. & Savitzky, A.H. (1973) Field observations on a population of the sand lizard *Uma exsul*. *Journal of Herpetology*, **7**, 51-53.
- Congdon, J.D., Vitt, L.J. & Hadley, N.F. (1978) Parental investment: comparative reproductive energetics in bisexual and unisexual lizards, genus *Cnemidophorus*. *American Naturalist*, **112**, 595-608.
- Cooper, B.W. (1953) Notes on the life history of the lizard, *Neoseps reynoldsi* Stejneger. *Journal of the Florida Academy of Sciences*, **16**, 235-238.
- Cooper, W.E. & Guillette, L.J. (1991) Observations on activity, display behavior, coloration and androgen levels in the keeled earless lizard, *Holbrookia propinqua*. *Amphibia-Reptilia*, **12**, 57-66.
- Cooper, W.E. & Habegger, J.J. (2000) Lingually mediated discrimination of prey, but not plant chemicals, by the Central American anguid lizard, *Mesaspis moreletii*. *Amphibia-Reptilia*, **22**, 81-90.
- Cooper, W.E. & Whiting, M.J. (2000) Ambush and active foraging modes both occur in the scincid genus *Mabuya*. *Copeia*, **2000**, 112-118.
- Cooper, W.E. & Vitt, L.J. (2002) Distribution, extent, and evolution of plant consumption by lizards. *Journal of Zoology*, **257**, 487-517.
- Cooper, W.E. & Bradley, K.A. (2009) Prey chemical discrimination by a diploglossine lizard, the giant Hispaniolan galliwasp (*Celestus warreni*). *Amphibia-Reptilia*, **30**, 135-140.
- Cooper, W.E., Al-Johany, A.M., Vitt, L.J. & Habegger, J.J. (2000) Responses to chemical cues from animal and plant foods by actively foraging insectivorous and omnivorous scincine lizards. *Journal of Experimental Zoology*, **287**, 327-339.
- Cooper, W.E., Vitt, L.J., Caldwell, J.P. & Fox, S.F. (2001) Foraging modes of some American lizards: relationships among measurement variables and discreteness of modes.

- Herpetologica*, **57**, 65-76.
- Cope, E.D. (1895) The Batrachia and Reptilia of the University of Pennsylvania West Indian expedition of 1890 and 1891. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **46**, 429-442.
- Corke, D. (1987) Reptile conservation on the Maria Islands (St Lucia, West Indies). *Biological Conservation*, **40**, 263-279.
- Costandius, E. & Mouton, P.L.F.N. (2006) Sexual size dimorphism in montane cordylid lizards: a case study of the dwarf crag lizard, *Pseudocordylus nebulosus*. *African Zoology*, **41**, 103-112.
- Costantini, D., Dell'Omo, G., Casagrande, S., Fabiani, A., Carosi, M., Bertacche, V., Marquez, C., Snell, H., Snell, H., Tapia, W. & Gentile, G. (2005) Inter-population variation of carotenoids in Galapagos land iguanas (*Conolophus subcristatus*). *Comparative Biochemistry and Physiology B*, **142**, 239-244.
- Couper, P.J., Covacevich, J.A. & Moritz, C. (1993) A review of the leaf-tailed geckos endemic to eastern Australia: a new genus, four new species, and other new data. *Memoirs of the Queensland Museum*, **34**, 95-124.
- Cowles, R.B. & Bogert, C.M. (1944) A preliminary study of the thermal requirements of desert reptiles. *Bulletin of the American Museum of Natural History*, **83**, 261-296.
- Cox, R.M., Skelly, S.L. & John-Alder, H.B. (2003) A comparative test of adaptive hypotheses for sexual size dimorphism in lizards. *Evolution*, **57**, 1653-1669.
- Crawford, C.M. & Thorpe, R.S. (1979) Body temperatures of two geckos (*Phelsuma*) and a skink (Mabuya) in Praslin, Seychelles. *British Journal of Herpetology*, **6**, 25-31.
- Cree, A. (1994) Low annual reproductive output in female reptiles from New Zealand. *New Zealand Journal of Zoology*, **21**, 351-372.
- Cree, A. & Guillette, L.J. (1995) Biennial reproduction with a fourteen-month pregnancy in the gecko *Hoplodactylus maculatus* from southern New Zealand. *Journal of Herpetology*, **29**, 163-173.
- Cree, A. & Hare, K.M. (2010) Equal thermal opportunity does not result in equal gestation length in a cool-climate skink and gecko. *Herpetological Conservation and Biology*, **5**, 271-282.
- Crochet, P.-A. & Renoult, J.P. (2008) *Tarentola annularis annularis* (Geoffroy de Saint-Hilaire, 1827) preying on a mammal. *Herpetology Notes*, **1**, 58-59.
- Crombie, R. & Steadman, D.W. (1986) The lizards of Rarotonga and Mangaia, Cook Island group, Oceania. *Pacific Science*, **40**, 44-53.
- Crombie, R.I. & Pregill, G.K. (1999) A checklist of the herpetofauna of the Palau Islands (Republic of Belau), Oceania. *Herpetological Monographs*, **13**, 29-80.
- Cruz, F.B. (1997) Reproductive activity in *Tropidurus etheridgei* in the semiarid Chaco of Salta, Argentina. *Journal of Herpetology*, **31**, 444-450.
- Cruz, F.B. (1998) Natural history of *Tropidurus spinulosus* (Squamata: Tropiduridae) from the dry chaco of Salta, Argentina. *Herpetological Journal*, **8**, 107-110.
- Cruz, F.B., Teisaire, E. & Nieto, L. (1997) Reproductive biology of the lizard *Tropidurus spinulosus* in the Chaco of Salta, Argentina. *Studies on Neotropical Fauna and Environment*, **32**, 28-32.
- Cruz, F.B., Schulte, J.A. & Bellagamba, P. (1999) New distributional records and natural history notes for reptiles from Southern Argentina. *Herpetological Review*, **30**, 182-183.
- Cruz, F.B., Fitzgerald, L.A., Espinoza, R.E. & Schulte, J.A. (2005) The importance of

- phylogenetic scale in tests of Bergmann's and Rapoport's rules: lessons from a clade of South American lizards. *Journal of Evolutionary Biology*, **18**, 1559-1574.
- Cruz, F.B., Antenucci, D., Luna, F., Abdala, C.S. & Vega, L.E. (2011) Energetics in Liolaemini lizards: implications of a small body size and ecological conservatism. *Journal of Comparative Physiology B.*, **181**, 373-382.
- Cruz, F.B., Belver, L., Acosta, J.C., Villavicencio, H.J., Blanco, G. & Canovas, M.G. (2009) Thermal biology of *Phymaturus* lizards: evolutionary constraints or lack of environmental variation? *Zoology*, **112**, 425-432.
- Cuellar, O. (1984) Reproduction in a parthenogenetic lizard: with a discussion of optimal clutch size and a critique of the clutch weight/body weight ratio. *American Midland Naturalist*, **111**, 242-258.
- Cumberlidge, N. (2008) Insular species of Afrotropical freshwater crabs (Crustacea: Decapoda: Brachyura: Potamonautidae and Potamidae) with special reference to Madagascar and the Seychelles. *Contributions to Zoology*, **77**, 71-81.
- Cunningham, J.D. (1966) Additional observations on the body temperatures of reptiles. *Herpetologica*, **22**, 184-189.
- Curry-Lindahl, K. (1979) Thermal ecology of the tree agama (*Agama atricollis*) in Zaire with a review of heat tolerance in reptiles. *Journal of Zoology*, **188**, 185-220.
- Curtin, A.J., Mouton, P.I.F.N. & Chinsamy, A. (2005) Bone growth patterns in two cordylid lizards, *Cordylus cataphractus* and *C. capensis*. *African Zoology*, **40**, 1-7.
- Daltry, J.C. (2009) *The status and management of Saint Lucia's forest reptiles and amphibians*. FCG International Ltd, Helsinki, Finland.
- Daly, B.G., Dickman, C.R. & Crowther, M.S. (2008) Causes of habitat divergence in two species of agamid lizards in arid Central Australia. *Ecology*, **89**, 65-76.
- Damuth, J. (1987) Interspecific allometry of population density in mammals and other animals: the independence of body mass and population energy-use. *Biological Journal of the Linnean Society*, **31**, 193-246.
- Darevsky, I.S. (1964) New species of scincid lizards from the islands of Lesser Sundas Archipelago, East Indonesia. *Zoologicheskii Zhurnal*, **43**, 80-88.
- Darevsky, I.S. (1984) *Lacerta strigata* Eichwald 1831 - Kaspische Smaragdeidechse. *Handbuch der Reptilien und Amphibien Europas*, Band 2/1, Echsen II (ed. By W. Bohme), pp. 82 – 99, AULA Verlag, Wiesbaden.
- Darevsky, I.S. & Tuniyev, B.S. (1997) A new species from *Lacerta saxicola* group - *Lacerta dryada* sp. nov. (Sauria: Lacertidae) and some comments relative to *Lacerta clarkorum* Darevsky & Vedmederja 1977. *Russian Journal of Herpetology*, **4**, 1-7.
- Das, I. (1997) A new species of *Cyrtodactylus* from the Nicobar Islands, India. *Journal of Herpetology*, **31**, 375-382.
- Das, I. & Grismer, L.L. (2003) Two new species of *Cnemaspis* Strauch, 1887 (Squamata: Gekkonidae) from the Seribuat Archipelago, Pahang and Johor States, West Malaysia. *Herpetologica*, **59**, 544-552.
- Das, I. & Austin, C.C. (2007) New species of *Lipinia* (Squamata: Scincidae) from Borneo, revealed by molecular and morphological data. *Journal of Herpetology*, **41**, 61-71.
- Das, I. & Vijayakumar, S.P. (2009) New species of *Ptychozoon* (Sauria: Gekkonidae) from the Nicobar Archipelago, Indian Ocean. *Zootaxa*, **2095**, 8-20.
- Daudin, J. & de Silva, M. (2007) An annotated checklist of the amphibians and terrestrial reptiles of the Grenadines with notes on their local natural history and conservation. *Applied*

- Herpetology*, **4**, 163-175.
- Daut, E.F. & Andrews, R.M. (1993) The effect of pregnancy on thermoregulatory behavior of the viviparous lizard *Chalcides ocellatus*. *Journal of Herpetology*, **27**, 6-13.
- Davidge, C. (1979) A census of a community of small terrestrial vertebrates. *Australian Journal of Ecology*, **4**, 165-170.
- Davis, W.B. & Smith, H.M. (1953) Lizards and turtles of the Mexican state of Morelos. *Herpetologica*, **9**, 100-108.
- Davis, W.B. & Dixon, J.R. (1961) Reptiles (exclusive of snakes) of the Chilpancingo region, Mexico. *Proceedings of the Biological Society of Washington*, **74**, 37-56.
- Daza, J.D., Herrera, A., Thomas, R. & Claudio, H.J. (2009) Are you what you eat? A geometric morphometric analysis of gekkotan skull shape. *Biological Journal of the Linnean Society*, **97**, 677-707.
- de Buffrenil, V. & Rimblot-Baly, F. (1999) Female reproductive output in exploited Nile monitor lizard (*Varanus niloticus* L.) populations in Sahelian Africa. *Canadian Journal of Zoology*, **77**, 1530-1539.
- De Magalhaes, J.P. & Costa, J. (2009) A database of vertebrate longevity records and their relation to other life-history traits. *Journal of Evolutionary Biology*, **22**, 1770-1774.
- de Zeeuw, M. (2010) Husbandry and reproduction of *Varanus glauerti* in captivity. *Biawak*, **4**, 103-107.
- Dearing, M.D. & Schall, J.J. (1994) Atypical reproduction and sexual dimorphism of the tropical Bonaire Island whiptail lizard, *Cnemidophorus murinus*. *Copeia*, **1994**, 760-766.
- Dial, B.E. & Grismer, L.L. (1992) A phylogenetic analysis of physiological-ecological character evolution in the lizard genus *Coleonyx* and its implications for historical biogeographic reconstruction. *Systematic Biology*, **41**, 178-195.
- Dias, E.J.R. & Rocha, C.F.D. (2004) Thermal ecology, activity patterns, and microhabitat use by two sympatric whiptail lizards (*Cnemidophorus abaetensis* and *Cnemidophorus ocellifer*) from northeastern Brazil. *Journal of Herpetology*, **38**, 586-588.
- Diaz, J.A., Perez-Tris, J., Bauwens, D., Perez-Aranda, D., Carbonell, R., Santos, T. & Telleria, J.L. (2007) Reproductive performance of a lacertid lizard at the core and the periphery of the species' range. *Biological Journal of the Linnean Society*, **92**, 87-96.
- Dickinson, W.R. (2006) *Temper Sands in Prehistoric Oceanian Pottery: Geotectonics, Sedimentology, Petrography, Provenance*. Geological Society of America.
- Dixon, J.R. (1956) A collection of amphibians and reptiles from west central Korea. *Herpetologica*, **12**, 50-56.
- Dixon, J.R. (1964) Further data on the geckos (*Phyllodactylus*) of islands of the extreme southern Caribbean. *Southwestern Naturalist*, **9**, 203-205.
- Dixon, J.R. & Kluge, A.G. (1964) A new gekkonid lizard genus from Australia. *Copeia*, **1964**, 174-180.
- Doan, T.M. (2008) Dietary variation within the Andean lizard clade *Proctoporus* (Squamata: Gymnophthalmidae). *Journal of Herpetology*, **42**, 16-21.
- Donoso-Barros, R. (1966) Dos nuevos Gonatodes de Venezuela. *Publicacion Ocasional Museo Nacional de Historia Natural, Santiago (Chile)*, **11**, 3-32.
- Doughty, P. (1997) The effects of fixed clutch sizes on lizard life-histories: reproduction in the Australian velvet gecko, *Oedura lesueurii*. *Journal of Herpetology*, **31**, 266-272.
- Doughty, P. & Shine, R. (1995) Life in two dimensions: natural history of the southern leaf-tailed gecko, *Phyllurus platurus*. *Herpetologica*, **51**, 193-201.

- Doughty, P. & Thompson, M.B. (1998) Unusual reproductive patterns in the Australian marbled gecko (*Phyllodactylus marmoratus*). *Copeia*, **1998**, 747-752.
- Du, W., Ji, X. & Shine, R. (2005) Does body volume constrain reproductive output in lizards? *Biology Letters*, **1**, 98-100.
- Du, W.-G., Ji, X. & Zhang, Y.-P. (2006) Inter-population variation in life-history traits of a Chinese lizard (*Takydromus septentrionalis*, Lacertidae). *Herpetological Journal*, **16**, 233-237.
- Du, W., Ji, X., Zhang, Y., Xu, X. & Shine, R. (2005) Identifying sources of variation in reproductive and life history traits among five populations of a Chinese lizard (*Takydromus septentrionalis*, Lacertidae). *Biological Journal of the Linnean Society*, **85**, 443-453.
- Du, Y., Lin, C.-X., Lin, L.-H., Qiu, Q.-B. & Ji, X. (2011) Ontogenetic shifts in sexual dimorphism and female reproduction in the Reeves's butterfly lizard *Leiolepis reevesii* from Hainan, China. *Journal of Herpetology*, **45**, 399-405.
- Duellman, W.E. (1961) The amphibians and reptiles of Michoacan, Mexico. *University of Kansas Publications, Museum of Natural History*, **15**, 1-148.
- Duellman, W.E. (1963) Amphibians and reptiles of the rainforests of southern El Peten, Guatemala. *University of Kansas Publications, Museum of Natural History*, **15**, 205-249.
- Duellman, W.E. (1965) A biogeographic account of the herpetofauna of Michoacan, Mexico. *University of Kansas Publications, Museum of Natural History*, **15**, 627-709.
- Duellman, W.E. (1978) The biology of an equatorial herpetofauna in Amazonian Ecuador. *University of Kansas Museum of Natural History Miscellaneous Publications*, **65**, 1-352.
- Duellman, W.E. & Mendelson, J.R. (1995) Amphibians and reptiles from Northern Departamento Loreto, Peru: taxonomy and biogeography. *University of Kansas Science Bulletin*, **55**, 329-376.
- Duffield, G.A. & Bull, C.M. (1996) Characteristics of the litter of the gidgee skink, *Egernia stokesii*. *Wildlife Research*, **23**, 337-341.
- Duffield, G.A. & Bull, C.M. (1998) Seasonal and ontogenetic changes in the diet of the Australian skink *Egernia stokesii*. *Herpetologica*, **54**, 414-419.
- Dunger, G.T. (1967) The lizards and snakes of Nigeria. Part 1. The Chamaeleons of Nigeria. *Nigerian Field*, **32**, 53-74.
- Dunger, G.T. (1967) The lizards and snakes of Nigeria. Part 3. The monitors and a plated lizard. *Nigerian Field*, **32**, 170-178.
- Dunger, G.T. (1968) The lizards and snakes of Nigeria. Part 4. The geckos of Nigeria. *Nigerian Field*, **33**, 18-47.
- Dunger, G.T. (1972) The lizards and snakes of Nigeria. Part 6. The skinks of Nigeria. *Nigerian Field*, **37**, 99-120.
- Dunger, G.T. (1973) The lizards and snakes of Nigeria. Part 7. The skinks of Nigeria (continued and completed). *Nigerian Field*, **38**, 54-80.
- Dunham, A.E. & Miles, D.B. (1985) Patterns of covariation in life history traits of squamate reptiles: the effects of size and phylogeny reconsidered. *American Naturalist*, **126**, 231-257.
- Dunham, A.E., Grant, B.W. & Overall, K.L. (1989) Interfaces between biophysical and physiological ecology and the population ecology of terrestrial vertebrate ectotherms. *Physiological Zoology*, **62**, 335-355.
- Dunn, E.R. (1935) Notes on American Mabuyas. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **87**, 533-560.
- Durtsche, R.D. (2000) Ontogenetic plasticity of food habits in the Mexican spiny-tailed iguana,

- Ctenosaura pectinata*. *Oecologia*, **124**, 185-195.
- Dutra, G.F., Siqueira, C.C., Vrcibradic, D., Kiefer, M.C. & Rocha, C.F.D. (2011) Plant consumption of insular and mainland populations of a tropical lizard. *Herpetologica*, **67**, 32-45.
- Dutton, R.H., Fitzpatrick, L.C. & Hughes, J.L. (1975) Energetics of the rusty lizard *Sceloporus olivaceus*. *Ecology*, **56**, 1378-1387.
- Duvdevani, I. (1971) The water and electrolyte metabolism in lizards of the genus *Acanthodactylus* Wiegmann 1834 (Lacertidae). *Dissertation*,
- Duvdevani, I. & Borut, A. (1974) Mean body temperature and heat absorption in four species of *Acanthodactylus* lizards. *Herpetologica*, **30**, 176-181.
- Edwards, A. (1999) Steroids and reproductive biology in the blotched blue-tongued lizard, *Tiliqua nigrolutea*. PhD thesis, University of Tasmania.
- Edwards, A., Jones, S.M. & Wapstra, E. (2002) Multiennial reproduction in females of a viviparous, temperate-zone skink, *Tiliqua nigrolutea*. *Herpetologica*, **58**, 407-414.
- Eifler, D.A. & Paskell, K.M. (2000) Body size effects on pursuit success and interspecific diet differences in *Cnemidophorus* lizards. *Amphibia-Reptilia*, **21**, 477-484.
- El Mouden, E.H., Znari, M. & Brown, R.P. (1999) Skeletochronology and mark-recapture assessments of growth in the North African agamid lizard (*Agama impalearis*). *Journal of Zoology*, **249**, 455-461.
- Eremchenko, V.K. (2007) Pamir-Alai transboundary conservancy area - biodiversity information management system and GIS. *Report*,
- Espinoza, R.E. & Lobo, F. (2003) Two new species of *Liolaemus* lizards from northwestern Argentina: speciation within the northern subclade of the *elongatus* group (Iguania: Liolaemidae). *Herpetologica*, **59**, 89-105.
- Espinoza, R.E., Lobo, F. & Cruz, F.B. (2000) *Liolaemus heliodermis*, a new lizard from northwestern Argentina with remarks on the content of the *elongatus* group (Iguania: Tropiduridae). *Herpetologica*, **56**, 235-244.
- Espinoza, R.E., Wiens, J.J. & Tracy, C.R. (2004) Recurrent evolution of herbivory in small, cold-climate lizards: breaking the ecophysiological rules of reptilian herbivory. *Proceedings of the National Academy of Sciences, USA*, **101**, 16819-16824.
- Evans, H.E. (1947) Notes on Panamanian reptiles and amphibians. *Copeia*, **1947**, 166-170.
- Evans, L.T. (1951) Field study of social behavior of the black lizard, *Ctenosaura pectinata*. *American Museum Novitates*, **1493**, 1-26.
- Evans, P.G.H. & Evans, J.B. (1980) The ecology of lizards on Praslin Island, Seychelles. *Journal of Zoology*, **191**, 171-192.
- Exon, N.F. & Tiffin, D.L. (1982) *Geology of offshore New Ireland basin in northern Papua New Guinea, and it's petroleum prospects*. In: Circum-Pacific Council for Energy and Mineral Resources 2009 – Transactions of the Third Circum-Pacific Energy and Mineral Resources Conference, 1982.
- Fathinia, B. & Rastegar-Pouyani, N. (2011) Sexual dimorphism in *Trapelus ruderatus ruderatus* (Sauria: Agamidae) with notes on the natural history. *Amphibian and Reptile Conservation*, **5**, 15-22.
- Feder, M.E. & Feder, J.H. (1981) Diel variation of oxygen consumption in three species of Philippine gekkonid lizards. *Copeia*, **1981**, 204-209.
- Fellers, G.M. & Drost, C.A. (1991) *Xantusia riversianna*. *Catalogue of American Amphibians and Reptiles*, **518**, 1-4.

- Feltrim, A.C. & De Lema, T. (2000) Uma nova especie de *Cnemidophorus* Wagler, 1830 do estado do Rio Grande do sul Brasil (Sauria, Teiidae). *Biociencias, Porto Alegre*, **8**, 103-114.
- Fenner, A.L., Bull, C.M. & Hutchinson, M.N. (2007) Omnivorous diet of the endangered pygmy bluetongue lizard, *Tiliqua adelaidensis*. *Amphibia-Reptilia*, **28**, 560-565.
- Ferner, J.W., Brown, R.M. & Greer, A.E. (1997) A new genus and species of moist closed canopy forest skinks from the Philippines. *Journal of Herpetology*, **31**, 187-192.
- Fischer, J. & Lindenmayer, D.B. (2005) The sensitivity of lizards to elevation: case study from south-eastern Australia. *Diversity and Distributions*, **11**, 225-233.
- Fitch, H.S. (1955) Habits and adaptations of the Great Plains skink (*Eumeces obsoletus*). *Ecological Monographs*, **25**, 59-83.
- Fitch, H.S. (1956) Temperature responses in free-living amphibians and reptiles of northeastern Kansas. *University of Kansas Museum of Natural History Miscellaneous Publications*, **8**, 417-476.
- Fitch, H.S. (1968) Temperature and behavior of some equatorial lizards. *Herpetologica*, **24**, 35-38.
- Fitch, H.S. (1970) Reproductive cycles of lizards and snakes. *University of Kansas Museum of Natural History Miscellaneous Publications*, **52**, 1-247.
- Fitch, H.S. (1973) A field study of Costa Rican lizards. *University of Kansas Science Bulletin*, **50**, 39-126.
- Fitch, H.S. (1978) Sexual size differences in the genus *Sceloporus*. *University of Kansas Science Bulletin*, **51**, 441-461.
- Fitch, H.S. (1982) Reproductive cycles in tropical reptiles. *Occasional Papers of the Museum of Natural History, University of Kansas*, **96**, 1-53.
- Fitch, H.S. (1985) Variation in clutch and litter size in New World reptiles. *University of Kansas Museum of Natural History Miscellaneous Publications*, **76**, 1-76.
- Fitch, H.S. & Greene, H.W. (1965) Breeding cycle in the ground skink, *Lygosoma laterale*. *University of Kansas Publications, Museum of Natural History*, **15**, 565-575.
- Fitch, H.S. & Henderson, R.W. (1978) Ecology and exploitation of *Ctenosaura similis*. *University of Kansas Science Bulletin*, **51**, 483-500.
- Fitch, H.S., Echelle, A.F. & Echelle, A.A. (1976) Field observations on rare or little known mainland anoles. *University of Kansas Science Bulletin*, **51**, 91-128.
- Fitzgerald, L.A., Cook, J.A. & Aquino, A.L. (1999) Phylogenetics and conservation of *Tupinambis* (Sauria: Teiidae). *Copeia*, **1999**, 894-905.
- Flemming, A.F. (1993) The male reproductive cycle of the lizard *Pseudocordylus m. melanotus* (Sauria: Cordylidae). *Journal of Herpetology*, **27**, 473-478.
- Flemming, A.F. & Mouton, P.L.F.N. (2002) Reproduction in a group-living lizard, *Cordylus cataphractus* (Cordylidae), from South Africa. *Journal of Herpetology*, **36**, 691-696.
- Flores, F., Lenzycki, J.H. & Palumbo, J. (1994) An ecological study of the endemic hispaniolan Anoline lizard *Chamaelinorops barbouri* (Lacertilia: Iguanidae). *Breviora*, **499**, 1-23.
- Flower, S.S. (1933) Notes on the recent reptiles and amphibians of Egypt, with a list of the species recorded from the kingdom. *Proceedings of the Zoological Society of London*, **19**, 735-851.
- Flury, A. (1949) *Gerrhonotus liocephalus infernalis* Baird in Texas. *Herpetologica*, **5**, 65-67.
- Franco, R.C. & de la Torre, G.G. (1990) Reptiles de la Isla La Pena, Nayarit, Mexico. *Anales del Instituto de Biologia, Universidad Nacional Autonoma de Mexico, serie Zoologia*, **61**, 175-187.

- Frank, E.F., Wicks, C., Mylroie, J., Troester, J., Alexander, E.C. & Carew, J. (1998) Geology of Isla de Mona, Puerto Rico. *Journal of Caves and Karst Studies*, **60**, 69-72.
- Frankenberg, E. (1978) Interspecific and seasonal variation of daily activity times in gekkonid lizards (Reptilia, Lacertilia). *Journal of Herpetology*, **12**, 505-519.
- Frankenberg, E. & Werner, Y.L. (1992) Egg, clutch and maternal sizes in lizards: intra- and interspecific relations in near-eastern Agamidae and Lacertidae. *Herpetological Journal*, **2**, 7-18.
- Franzen, M. & Heckes, U. (1999) *Eremias suphani* Basoglu & Hellmich, 1968 und *Eremias strauchi* Kessler, 1878 in der ostlichen Turkei: Diagnostische Merkmale, Verbreitung und Lebensraume (Sauria: Lacertidae). *Salamandra*, **35**, 255-266.
- Frydlova, P. & Frynta, D. (2010) A test of Rensch's rule in varanid lizards. *Biological Journal of the Linnean Society*, **100**, 293-306.
- Fuenmayor, G.R., Ugueto, G., Rivero, R. & Miralles, A. (2005) The herpetofauna of Isla de Margarita, Venezuela: new records and comments. *Caribbean Journal of Science*, **41**, 346-351.
- Fuhn, I.E. (1972) Revision du phylum forestier du genre *Panaspis* Cope (Reptilia, Scincidae, Lygosominae). *Review Roumain de Biologie Serie de Zoologie*, **17**, 257-271.
- Funk, R.S. (1981) *Phrynosoma mcallii*. *Catalogue of American Amphibians and Reptiles*, **281**, 1.
- Gadsden, H., Davila-Carrasco, M.L. & Gil-Martinez, R. (2006) Reproduction in the arenicolous Mexican lizard *Uma exsul*. *Journal of Herpetology*, **40**, 117-122.
- Gainsbury, A.M. & Colli, G.R. (2003) Lizard assemblages from natural cerrado enclaves in southwestern Amazonia: the role of stochastic extinctions and isolation. *Biotropica*, **35**, 503-519.
- Galan, P. (1997) Reproductive ecology of the lacertid lizard *Podarcis bocagei*. *Ecography*, **20**, 197-209.
- Galan, P. (2003) Female reproductive characteristics of the viviparous skink *Chalcides bedriagai pistaciae* (Reptilia, Squamata, Scincidae) from an Atlantic beach in north-west Spain. *Amphibia-Reptilia*, **24**, 79-85.
- Galan, P. & Vicente, L. (2003) Reproductive characteristics of the insular lacertid *Teira dugesii*. *Herpetological Journal*, **13**, 149-154.
- Gallagher, D.S. & Dixon, J.R. (1992) Taxonomic revision of the South American lizard genus *Kentropyx Spix* (Sauria, Teiidae). *Bolletino del Museo Regionale di Scienze Naturali, Torino*, **10**, 125-171.
- Gans, C. (1965) Notes on amphisbaenids (Amphisbaenia, Reptilia). 17. A redescription and discussion of *Amphisbaena angustifrons* Cope and *Amphisbaena camura* Cope of southern South America. *American Museum Novitates*, **2225**, 1-32.
- Gans, C., Laurent, R.F. & Pandit, H. (1965) Notes on a herpetological collection from the Somali Republic. *Annales, Musee Royal de L'Afrique Central*, **134**, 1-93.
- Garcia-Vazquez, U. & Feria-Ortiz, M. (2006) Skinks of Mexico. *Reptilia*, **49**, 74-79.
- Garda, A.A., Costa, G.C., Franca, F.G.R., Giugliano, L.G., Leite, G.S., Mesquita, D.O., Nogueira, C., Tavares-Bastos, L., Vasconcellos, M.M., Vieira, G.H.C., Vitt, L.J., Werneck, F.P., Wiederhecker, H.C. & Colli, G.R. (2012) Reproduction, body size, and diet of *Polychrus acutirostris* (Squamata: Polychrotidae) in two contrasting environments in Brazil. *Journal of Herpetology*, **46**, 2-8.
- Garland, T. & Else, P.L. (1987) Seasonal, sexual, and individual variation in endurance and

- activity metabolism in lizards. *American Journal of Physiology*, **252**, R439-R449.
- Garrick, D. (2008) Body surface temperature and length in relation to the thermal biology of lizards. *Bioscience Horizons*, **1**, 136-142.
- Gasnier, T.R., Magnusson, W.E. & Lima, A.P. (1994) Foraging activity and diet of four sympatric lizard species in a tropical rainforest. *Journal of Herpetology*, **28**, 187-192.
- Gaulke, M. & Curio, E. (2001) A new monitor lizard from Panay Island, Philippines (Reptilia, Sauria, Varanidae). *Spixiana*, **24**, 275-286.
- Gaulke, M., Curio, E., Demegillo, A. & Paulino, N. (2002) *Varanus mabitang*, a rare monitor lizard from Panay Island and a new conservation target. *Silliman Journal*, **43**, 24-41.
- Gaulke, M., Altenbach, A.V., Demegillo, A. & Struck, U. (2007) On the diet of *Varanus mabitang*. *Mertensiella*, **16**, 228-239.
- Geissler, P., Nguyen, T.Q., Phung, T.M., Van Devender, R.W., Hartmann, T., Farkas, B., Ziegler, T. & Bohme, W. (2011) A review of Indochinese skinks of the genus *Lygosoma* Hardwicke & Gray, 1827 (Squamata: Scincidae), with natural history notes and an identification key. *Biologia*, **66**, 1159-1176.
- Geldmacher, J., Hoernle, K., van den Bogaard, P., Zankl, G. & Garbe-Schönberg, D. (2001) Earlier history of the 70-Ma-old Canary hotspot based on the temporal and geochemical evolution of the Selvagen Archipelago and neighboring seamounts in the eastern North Atlantic. *Journal of Volcanology and Geothermal Research*, **111**, 55-87.
- Gerlach, J. (2005) *Ailuroonyx trachygaster* – the best view yet! *Gekko*, **4**, 31.
- Gerlach, J. (2008) Population and conservation status of the reptiles of the Seychelles islands. *Phelsuma*, **16**, 30-48.
- Gerlach, J. & Canning, K.L. (1996) A new species of the western Indian Ocean gecko *Ailuroonyx* (Reptilia; Gekkonidae). *Herpetological Journal*, **6**, 27-42.
- Gholamifard, A., Gholamhosseini, A., Rastegar-Pouyani, N., Esmaeili, H.R. & Kami, H.-G. (2010) First records of *Tropiocolotes steudneri* Peters, 1869 and *Hemidactylus flaviviridis* Ruppell, 1840 (Sauria: Gekkonidae) from Fars Province, Iran. *Asian Herpetological Research*, **1**, 61-63.
- Gibbons, J.R.H. (1981) The biogeography of *Brachylophus* (Iguanidae) including the description of a new species, *B. vitiensis*, from Fiji. *Journal of Herpetology*, **15**, 255-273.
- Gibbons, J.R.H. (1984) Iguanas of the South Pacific. *Oryx*, **18**, 82-92.
- Gibbons, J.R.H. & Brown, W.C. (1988) A new *Lepidodactylus* (Gekkonidae) from Eua Island, Tonga. *Journal of Herpetology*, **22**, 356-360.
- Gill, B.J. (1993) The land reptiles of Western Samoa. *Journal of the Royal Society of New Zealand*, **23**, 79-89.
- Girling, J.E., Cree, A. & Guillette, L.J. (1998) Oviducal structure in four species of gekkonid lizard differing in parity mode and eggshell structure. *Reproduction, Fertility and Development*, **10**, 139-154.
- Goldberg, S.R. (1983) Reproduction of the coast horned lizard, *Phrynosoma coronatum*, in southern California. *Southwestern Naturalist*, **28**, 478-479.
- Goldberg, S.R. (1987) Reproductive cycle of the giant spotted whiptail, *Cnemidophorus burti stictogrammus*, in Arizona. *Southwestern Naturalist*, **32**, 510-511.
- Goldberg, S.R. (2005) Reproductive cycle of the Western skink, *Eumeces skiltonianus*. *Texas Journal of Science*, **57**, 295-301.
- Goldberg, S.R. (2006) Reproductive cycle of bushveld lizard *Heliobolus lugubris* (Squamata: Lacertidae) from southern Africa. *Salamandra*, **42**, 151-154.

- Goldberg, S.R. (2007) Reproductive cycle of the Namibian striped skink, *Trachylepis sparsa* (Squamata: Scincidae) from Southern Africa. *African Zoology*, **42**, 289-293.
- Goldberg, S.R. (2008) Reproduction in the Siamese leaf-toed gecko, *Dixonius siamensis* (Squamata, Gekkonidae) from Thailand. *Texas Journal of Science*, **60**, 233-238.
- Goldberg, S.R. (2009) Notes on reproduction of the knob-scaled lizard, *Xenosaurus grandis* (Squamata: Xenosauridae), from Veracruz, Mexico. *Texas Journal of Science*, **61**, 317-322.
- Goldberg, S.R. (2011) *Mesaspis gadovii* (Gadow's alligator lizard). Reproduction. *Herpetological Review*, **42**, 606.
- Goldberg, S.R. (2012) *Mesalina guttulata* (desert lacerta). Endoparasites. *Herpetological Review*, **43**, 136.
- Goldberg, S.R. & Robinson, M.D. (1979) Reproduction in two Namib Desert lacertid lizards (*Aporosaura anchietae* and *Meroles cuneirostris*). *Herpetologica*, **35**, 169-175.
- Goldberg, S.R. & Miller, C.M. (1985) Reproduction of the silvery legless lizard, *Anniella pulchra pulchra* (Anniellidae), in southern California. *Southwestern Naturalist*, **30**, 617-619.
- Goldberg, S.R. & Rodriguez, E. (1986) Reproductive cycles of two iguanid lizards from northern Chile, *Tropidurus quadrivittatus* and *Tropidurus theresioides*. *Journal of Arid Environments*, **10**, 147-151.
- Goldberg, S.R. & Lowe, C.H. (1997) Reproductive cycle of the gila monster, *Heloderma suspectum*, in Southern Arizona. *Journal of Herpetology*, **31**, 161-166.
- Goldberg, S.R. & Bursey, C.R. (2010) Helminths from eight Species of African skinks (Trachylepis: Scincidae). *Comparative Parasitology*, **77**, 236-241.
- Goldberg, S.R. & Kraus, F. (2011) Notes on reproduction of *Cryptoblepharus poecilopleurus* (Squamata: Scincidae) from the northern Mariana Islands, Western Pacific. *Current Herpetology*, **30**, 159-161.
- Goldberg, S.R., Bursey, C.R. & Holshuh, H.J. (1994) *Physocephalus* sp. (Spirurida, Spirocercidae) Larvae in Stomach Granulomas of the Blue Spiny Lizard, *Sceloporus serrifer* (Phrynosomatidae) from Texas. *Journal of Wildlife Diseases*, **30**, 274-276.
- Gonwouo, N.L., LeBreton, M., C., W., Chirio, L., Ngassam, P. & Tchamba, N.M. (2006) Geographic and ecological distribution of the endemic montane chameleons along the Cameroon mountain range. *Salamandra*, **42**, 209-226.
- Goodman, B.A. (2006) Clutch traits in the skink *Carlia vivax*. *Herpetofauna*, **36**, 27-30.
- Goodman, B.A. (2006) Costs of reproduction in a tropical invariant-clutch producing lizard (*Carlia rubrigularis*). *Journal of Zoology*, **270**, 236-243.
- Goodman, B.A. (2007) Divergent morphologies, performance and escape behaviour in two tropical rock-using lizards. *Biological Journal of the Linnean Society*, **91**, 85-98.
- Goodman, B.A. & Isaac, J.L. (2008) Convergent body flattening in a clade of tropical rock-using lizards (Scincidae: Lygosominae). *Biological Journal of the Linnean Society*, **94**, 399-411.
- Goodman, B.A., Hudson, S.C., Isaac, J.L. & Schwarzkopf, L. (2009) The evolution of body shape in response to habitat: is reproductive output reduced in flat lizards? *Evolution*, **63**, 1279-1291.
- Goodyear, S.E. & Pianka, E.R. (2011) Spatial and temporal variation in diets of sympatric lizards (Genus *Ctenotus*) in the Great Victoria Desert, Western Australia. *Journal of Herpetology*, **45**, 265-271.
- Goonewardene, S., Hawke, Z., Vanneck, V., Drion, A., de Silva, A., Jayarathne, R. & Perera, J.

- (2003) Diversity of Nilgala fire savannah, Sri Lanka: with special reference to its herpetofauna. *Report of Project Hoona*.
- Gordon, C.E., Dickman, C.R. & Thompson, M.B. (2010) Partitioning of temporal activity among desert lizards in relation to prey availability and temperature. *Austral Ecology*, **35**, 41-52.
- Greenbaum, E. & Carr, J.L. (2005) The herpetofauna of Upper Niger National Park, Guinea, West Africa. *Scientific Papers of the Natural History Museum, University of Kansas*, **37**, 1-21.
- Greene, H.W., Rodriguez, J.J.S. & Powell, B.J. (2006) Parental behavior in anguid lizards. *South American Journal of Herpetology*, **1**, 9-19.
- Greenville, A.C. & Dickman, C.R. (2005) The ecology of *Lerista labialis* (Scincidae) in the Simpson Desert: reproduction and diet. *Journal of Arid Environments*, **60**, 611-625.
- Greer, A.E. (1980) A new species of *Morethia* (Lacertilia: Scincidae) from northern Australia, with comments on the biology and relationships of the genus. *Records of the Australian Museum*, **33**, 89-122.
- Greer, A.E. & Biswas, S. (1967) A new generic arrangement for some Australian scincid lizards. *Breviora*, **267**, 1-19.
- Greer, A.E. & Parker, F. (1967) A second skink with fragmented head scales from Bougainville, Solomon Islands. *Breviora*, **279**, 1-12.
- Greer, A.E. & Biswas, S. (1968) Clutch size in the scincid genus *Emoia*. *Copeia*, **1968**, 417-418.
- Greer, A.E. & Parker, F. (1968) A new species of *Tribolonotus* (Lacertilia: Scincidae) from Bougainville and Buka, Solomon Islands, with comments on the biology of the genus. *Breviora*, **291**, 1-23.
- Greer, A.E. & Biswas, S. (1974) The generic relationships of the scincid lizard genus *Leiopisma* and its relatives. *Australian Journal of Zoology*, **31** (supplement), 1-67.
- Greer, A.E. & Parker, F.A. (1974) The *fasciatus* species group of *Sphenomorphus* (Lacertilia: Scincidae): notes on eight previously described species and descriptions of three new species. *Papua New Guinea Scientific Society Proceedings*, **25**, 31-61.
- Greer, A.E. & Biswas, S. (1977) On the adaptive significance of the loss of an oviduct in reptiles. *Proceedings of the Linnean Society of New South Wales*, **101**, 242-249.
- Greer, A.E. & Biswas, S. (1982) A new species of *Geomyersia* (Scincidae) from the Admiralty Islands, with a summary of the genus. *Journal of Herpetology*, **16**, 61-66.
- Greer, A.E. & Biswas, S. (1992) Revision of the species previously associated with the Australian scincid lizard *Eulamprus tenuis*. *Records of the Australian Museum*, **44**, 7-19.
- Greer, A.E. & Biswas, S. (1997) A new species of *Lampropholis* (Squamata: Scincidae) with a restricted, high altitude distribution in eastern Australia. *Australian Zoologist*, **30**, 360-368.
- Greer, A.E. & Biswas, S. (2001) Distribution of maximum snout-vent length among species of scincid lizards. *Journal of Herpetology*, **35**, 383-395.
- Greer, A.E. & Shea, G. (2004) A new character within the taxonomically difficult *Sphenomorphus* group of lygosomine skinks, with a description of a new species from New Guinea. *Journal of Herpetology*, **38**, 79-87.
- Greer, A.E. & Biswas, S. (2005) Encyclopedia of Australian reptiles. *online Encyclopedia*.
- Greer, A.E., David, P. & Teynie, A. (1985) A new species of *Lygosoma* (Lacertilia: Scincidae) from West Africa, with comments on its biology. *Journal of Herpetology*, **19**, 365-373.
- Greer, A.E., Caputo, V., Lanza, B. & Palmieri, R. (1998) Observations on limb reduction in the

- scincid lizard Genus *Chalcides*. *Journal of Herpetology*, **32**, 244-252.
- Griffith, H. (1990) Miniaturization and elongation in *Eumeces* (Sauria: Scincidae). *Copeia*, **1990**, 751-758.
- Grillitsch, H. & Cabela, A. (1990) Zum Systematischen Status der *Blindschleichen* (Squamata: Anguillidae) der Peloponnes und der sudlichen Ionischen Inseln. *Herpetozoa*, **2**, 131-153.
- Grismer, L.L. (1994) Three new species of intertidal side-blotched lizards (genus *Uta*) from the Gulf of California, Mexico. *Herpetologica*, **50**, 451-474.
- Grismer, L.L. (2007) A new species of small montane forest floor skink (genus *Sphenomorphus* Fitzinger 1843) from southern peninsular Malaysia. *Herpetologica*, **63**, 544-551.
- Grismer, L.L. & Onn, C.K. (2008) A new species of *Cnemaspis* Strauch 1887 (Squamata: Gekkonidae) from Pulau Perhentian Besar, Terengganu, Peninsular Malaysia. *Zootaxa*, **1771**, 1-15.
- Grismer, L.L. & Pan, K.A. (2008) Diversity, endemism, and conservation of the amphibians and reptiles of southern Peninsular Malaysia and its offshore islands. *Herpetological Review*, **39**, 270-281.
- Grismer, L.L., Ahmad, N. & Onn, C.K. (2009) A new, diminutive, upland *Sphenomorphus* Fitzinger 1843 (Squamata: Scincidae) from the Belum-Temengor forest complex, Peninsular Malaysia. *Zootaxa*, **2312**, 27-38.
- Grismer, L.L., Youmans, T.M., Wood, P.L. & Grismer, J.L. (2006) Checklist of the herpetofauna of the Seribu Archipelago, West Malaysia with comments on biogeography, natural history, and adaptive types. *Raffles Bulletin of Zoology*, **54**, 157-180.
- Grismer, L.L., Grismer, J.L., Wood, P.L. & Onn, C.K. (2008) The distribution, taxonomy, and redescription of the geckos *Cnemaspis affinis* (Stoliczka 1887) and *C. flavolineata* (Nicholls 1949) with descriptions of a new montane species and two new lowland, karst-dwelling species from Peninsular Malaysia. *Zootaxa*, **1931**, 1-24.
- Grismer, L.L., Onn, C.K., Grismer, J.L., Wood, P.L. & Belabut, D. (2008) Three new species of *Cyrtodactylus* (Squamata: Gekkonidae) from Peninsular Malaysia. *Zootaxa*, **1921**, 1-23.
- Grismer, L.L., Chav, T., Neang, T., Wood, P.L., Grismer, J.L., Youmans, T.M., Ponce, A., Daltry, J.C. & Kaiser, H. (2007) The herpetofauna of the Phnom Aural Wildlife Sanctuary and checklist of the herpetofauna of the Cardamom Mountains, Cambodia. *Hamadryad*, **31**, 216-241.
- Grizante, M.B., Navas, C.A., Garland, T. & Kohlsdorf, T. (2010) Morphological evolution in Tropicodurinae squamates: an integrated view along a continuum of ecological settings. *Journal of Evolutionary Biology*, **23**, 98-111.
- Gross, J.M. (1975) The Archaeology of Anegada Island. *Journal of the Virgin Islands Archaeological Society*, **2**, 12-16.
- Grossmann, W. & Harbig, P. (2010) Pflege und vermehrung des korpulenten schlangenskinks *Lygosoma corpulentum* Smith, 1921 im terrarium – erste ergebnisse. *Sauria*, **32**, 35-46.
- Guarino, F.M. (2010) Structure of the femora and autotomous (postpygal) caudal vertebrae in the three-toed skink *Chalcides chalcides* (Reptilia: Squamata: Scincidae) and its applicability for age and growth rate determination. *Zoologischer Anzeiger*, **248**, 273-283.
- Guillette, L.J. (1982) The evolution of viviparity and placentation in the high elevation, Mexican lizard *Sceloporus aeneus*. *Herpetologica*, **38**, 94-103.
- Guillette, L.J. (1983) Notes concerning reproduction of the montane skink, *Eumeces copei*. *Journal of Herpetology*, **17**, 144-148.
- Guillette, L.J. & Smith, H.M. (1982) A review of the Mexican lizard *Barisia imbricata*, and the

- description of a new subspecies. *Transactions of the Kansas Academy of Sciences*, **85**, 13-33.
- Guillette, J.L. & Casas-Andreu, G. (1987) The reproductive biology of the high elevation Mexican lizard *Barisia imbricata*. *Herpetologica*, **43**, 29-38.
- Haagner, G.V., Branch, W.R. & Haagner, A.J.F. (2000) Notes on a collection of reptiles from Zambia and adjacent areas of the Democratic Republic of the Congo. *Annals of the Eastern Cape Museum*, **1**, 1-25.
- Hadden, R.L. & Minson, S.G. (2010) The Geology of Haiti: An Annotated Bibliography of Haiti's Geology, Geography and Earth Science. In. DTIC Document
- Hagen, I.J. & Bull, C.M. (2011) Home ranges in the trees: radiotelemetry of the prehensile tailed skink, *Corucia zebrata*. *Journal of Herpetology*, **45**, 36-39.
- Hailey, A. & Elliot, M. (1995) Thermoregulation of the amphisbaenian *Zygaspis quadrifrons*. *Herpetological Journal*, **5**, 281-284.
- Hailey, A., Rose, C.A. & Pulford, E. (1987) Food consumption, thermoregulation and ecology of the skink *Chalcides bedriagai*. *Herpetological Journal*, **1**, 144-153.
- Hall, W.P. & Smith, H.M. (1979) Lizards of the *Sceloporus orcutti* complex of the Cape Region of Baja California. *Breviora*, **452**, 1-26.
- Halloy, M. & Halloy, S. (1997) An indirect form of parental care in a high altitude viviparous lizard, *Liolaemus huacahuasicus* (Tropiduridae). *Bulletin of the Maryland Herpetological Society*, **33**, 139.
- Halloy, M., Robles, C. & Cuezco, F. (2006) Diet in two syntopic neotropical lizard species of *Liolaemus* (Liolaemidae): interspecific and intersexual differences. *Revista Espanola de Herpetologia*, **20**, 47-56.
- Halloy, M., Boretto, J.M. & Ibargüengoytía, N.R. (2007) Signs of parental behavior in *Liolaemus elongatus* (Sauria: Liolaemidae), of Neuquen, Argentina. *South American Journal of Herpetology*, **2**, 141-147.
- Hamilton, A.M., Klein, E.R., Eckstut, M.E. & Hartfield, E.E. (2007) A simple, inexpensive method to capture arboreal lizards. *Herpetological Conservation and Biology*, **2**, 164-167.
- Hamilton, A.M., Eckstut, M.E., Klein, E.R. & Austin, C.C. (2008) Clutch size in the tropical scincid lizard *Emoia sanfordi*, a species endemic to the Vanuatu Archipelago. *Zoological Science*, **25**, 843-848.
- Hardy, L.M. & McDiarmid, R.W. (1969) The amphibians and reptiles of Sinaloa, Mexico. *Publications of the University of Kansas Museum of Natural History*, **18**, 39-252.
- Hare, K.M. (2005) The paradox of nocturnality in lizards. *Dissertation*,
- Hare, K.M. & Cree, A. (2005) Natural history of *Hoplodactylus stephensi* (Reptilia: Gekkonidae) on Stephens Island, Cook Strait, New Zealand. *New Zealand Journal of Ecology*, **29**, 137-142.
- Hare, K.M., Pledger, S., Thompson, M.B., Miller, J.H. & Daugherty, C.H. (2010) Nocturnal lizards from a cool-temperate environment have high metabolic rates at low temperatures. *Journal of Comparative Physiology B.*, **180**, 1173-1181.
- Harlow, H.J., Purwandana, D., Jessop, T.S. & Phillips, J.A. (2010) Body temperature and thermoregulation of Komodo dragons in the field. *Journal of Thermal Biology*, **35**, 338-347.
- Harmon, L.J., Harmon, L.L. & Jones, C.G. (2007) Competition and community structure in diurnal arboreal geckos (genus *Phelsuma*) in the Indian Ocean. *Oikos*, **116**, 1863-1878.
- Harvey, M.B. & Gutberlet, R.L.J. (1998) Lizards of the genus *Tropidurus* (Iguania: Tropiduridae) from the Serrania de Huanchaca, Bolivia: new species, natural history, and a

- key to the genus. *Herpetologica*, **54**, 493-520.
- Hasegawa, M. (1984) Biennial reproduction in the lizard *Eumeces okadae* on Miyake-Jima, Japan. *Herpetologica*, **40**, 194-199.
- Hasegawa, M. (1990) Demography of an island population of the lizard, *Eumeces okadae*, on Miyake-jima, Izu Islands. *Researches on Population Ecology*, **32**, 119-133.
- Hasegawa, M. (1994) Insular radiation in life history of the lizard *Eumeces okadae* in the Izu Islands, Japan. *Copeia*, **1994**, 732-747.
- Hasegawa, M. (1997) Density effects on life history traits of an island lizard population. *Ecological Research*, **12**, 111-118.
- Hauschild, A. (1988) Bemerkungen zur haltung und zucht des schneckenskinks *Tiliqua gerrardii*. *Salamandra*, **24**, 248-257.
- Hawlena, D. & Perez-Mellado, V. (2009) Change your diet or die: predator-induced shifts in insectivorous lizard feeding ecology. *Oecologia*, **161**, 411-419.
- Hawlena, D., Saltz, D., Abramsky, Z. & Bouskila, A. (2010) Ecological trap for desert lizards caused by anthropogenic changes in habitat structure that favor predator activity. *Conservation Biology*, **23**, 804-809.
- Haxhiu, I. (1998) The Reptilia of Albania: species composition, distribution, habitats. *Bonner Zoologische Beiträge*, **48**, 35-57.
- Heatwole, H. (1975) Biogeography of reptiles on some of the islands and cays of eastern Papua-New Guinea. *Atoll Research Bulletin*, **180**, 1-41.
- Heatwole, H. & Sexton, O.J. (1966) Herpetofaunal comparisons between two climatic zones in Panama. *American Midland Naturalist*, **75**, 45-60.
- Heatwole, H. & Veron, J.E.N. (1977) Vital limit and evaporative water loss in lizards (Reptilia: Lacertilia): a critique and new data. *Journal of Herpetology*, **11**, 341-348.
- Heger, N.A. & Heger, T.G. (2007) Behavior, ecology and thermal physiology of *Varanus giganteus*: a field study of Australia's largest monitor lizard. *Mertensiella*, **16**, 255-290.
- Heideman, N.J.L. (1994) Reproduction in *Agama aculeata aculeata* and *Agama planiceps planiceps* females from Windhoek, Namibia. *Amphibia-Reptilia*, **15**, 351-361.
- Heller, E. (1903) Papers from the Hopkins Stanford Galapagos Expedition, 1898-1899. WIV. Reptiles. *Proceedings of the Biological Society of Washington*, **4**, 39-98.
- Hendrickson, J.R. (1966) Observations on the Fauna of Pulau Tioman and Pulau Tulai. 5. The reptiles. *Bulletin of the National Museum of Singapore*, **34**, 53-71.
- Henkel, F.W. & Bohme, W. (2001) A new carphodactyline gecko of the New Caledonian genus *Eurydactylodes* (Sauria: Gekkonidae). *Salamandra*, **37**, 149-156.
- Henle, K. (1989) Ecological segregation in a subterranean reptile assemblage in arid Australia. *Amphibia-Reptilia*, **10**, 277-295.
- Henle, K. (1989) Ecological segregation in an assemblage of diurnal lizards in arid Australia. *Acta Oecologia Generalis*, **10**, 19-35.
- Henle, K. (1989) Population ecology and life history of the diurnal skink *Morethia boulengeri* in arid Australia. *Oecologia*, **78**, 521-532.
- Henle, K. (1990) Population ecology and life history of the arboreal gecko *Gehyra variegata* in arid Australia. *Herpetological Monographs*, **4**, 30-60.
- Henle, K. (1991) Life history patterns in lizards of the arid and semiarid zone of Australia. *Oecologia*, **88**, 347-358.
- Hennessy, J. (2010) Parthenogenesis in an ornate Nile monitor, *Varanus ornatus*. *Biawak*, **4**, 26-30.

- Herczeg, G., Torok, J. & Koros, Z. (2007) Size-dependent heating rates determine the spatial and temporal distribution of small-bodied lizards. *Amphibia-Reptilia*, **28**, 347-356.
- Hernandez, E., Nogales, M. & Martin, A. (2000) Discovery of a new lizard in the Canary Islands, with a multivariate analysis of *Gallotia* (Reptilia: Lacertidae). *Herpetologica*, **56**, 63-76.
- Hernandez-Salinas, U., Ramirez-Bautista, A., Leyte-Manrique, A. & Smith, G.R. (2010) Reproduction and sexual dimorphism in two populations of *Sceloporus grammicus* (Sauria: Phrynosomatidae) from Hidalgo, Mexico. *Herpetologica*, **66**, 12-22.
- Herrel, A., Meyers, J.J. & Vanhooydonck, B. (2002) Relations between microhabitat use and limb shape in phrynosomatid lizards. *Biological Journal of the Linnean Society*, **77**, 149-163.
- Herrel, A., Vanhooydonck, B. & Van Damme, R. (2004) Omnivory in lacertid lizards: adaptive evolution or constraint? *Journal of Evolutionary Biology*, **17**, 974-984.
- Herrel, A., Huyghe, K., Vanhooydonck, B., Backeljau, T., Breugelmans, K., Grbac, I., Van Damme, R. & Irschick, D.J. (2008) Rapid large-scale evolutionary divergence in morphology and performance associated with exploitation of a different dietary resource. *Proceedings of the National Academy of Sciences, USA*, **105**, 4792-4795.
- Hertz, P.E. (1979) Sensitivity to high temperatures in three west Indian grass anoles. *Comparative Biochemistry and Physiology A.*, **63**, 217-222.
- Hertz, P.E. (1981) Adaptation to altitude in two West Indian anoles (Reptilia: Iguanidae): field thermal biology and physiological ecology. *Journal of Zoology*, **195**, 25-37.
- Hertz, P.E. & Nevo, E. (1981) Thermal biology of four Israeli agamid lizards in early summer. *Israel Journal of Zoology*, **30**, 190-210.
- Hertz, P.E., Huey, R.B. & Stevenson, R.D. (1993) Evaluating temperature regulation by field-active ectotherms: the fallacy of the inappropriate question. *American Naturalist*, **142**, 796-818.
- Herzog, H.A. & Drummond, H. (1984) Tail autotomy inhibits tonic immobility in geckos. *Copeia*, **1984**, 763-764.
- Hibbits, T.J. (2005) Ecology and sexual selection of the common barking gecko (*Ptenopus garrulus*). PhD thesis, University of the Witwatersrand.
- Hibbits, T.J., Pianka, E.R., Huey, R.B. & Whiting, M.J. (2005) Ecology of the common barking gecko (*Ptenopus garrulus*) in Southern Africa. *Journal of Herpetology*, **39**, 509-515.
- Hillman, P.E. (1969) Habitat specificity in three sympatric species of *Ameiva* (Reptilia: Teiidae). *Ecology*, **50**, 476-481.
- Hirth, H.F. (1963) The ecology of two lizards on a tropical beach. *Ecological Monographs*, **33**, 1-3.
- Hodges, W.L. (1995) *Phrynosoma ditmarsii*. *Catalogue of American Amphibians and Reptiles*, **614**, 1-3.
- Hodges, W.L. & Perez-Ramos, E. (2001) New localities and natural history notes on *Bipes canaliculatus* in Guerrero, Mexico. *Herpetological Review*, **32**, 153-156.
- Hoernle, K., van den Bogaard, P., Werner, R., Lissinna, B., Hauff, F., Alvarado, G. & Garbe-Schönberg, D. (2002) Missing history (16–71 Ma) of the Galápagos hotspot: Implications for the tectonic and biological evolution of the Americas. *Geology*, **30**, 795-798.
- Hofer, U., Baur, H. & Bersier, L.F. (2003) Ecology of three sympatric species of the genus *Chamaeleo* in a tropical upland forest in Cameroon. *Journal of Herpetology*, **37**, 203-207.
- Holman, J.A. (1971) *Ophisaurus attenuatus*. *Catalogue of American Amphibians and Reptiles*, **111**, 1-3.

- Honegger, R.E. (1969) Notes on some amphibians and reptiles at Zurich Zoo. *International Zoo Yearbook*, **9**, 24-28.
- Horn, H.-G. & Visser, G.J. (1997) Review of reproduction of Monitor lizards in captivity II. *International Zoo Yearbook*, **35**, 227-246.
- Hornby, R.J. (1996) A checklist of amphibians and reptiles of the United Arab Emirates. *Tribulus*, **6**, 9-13.
- Horner, P. (2007) *Ctenotus quirinus* sp. nov. (Reptilia: Sauria: Scincidae) – a new species of skink from the Northern Territory, with the recognition of *C. brevipes* Storr, 1981 and *C. essingtonii* (Gray, 1842) as distinct species. *The Beagle*, **23**, 119-130.
- Hoser, R. (2009) A new genus and species of skink from Victoria. *Australasian Journal of Herpetology*, **3**, 1-6.
- Hoskin, C.J. & Higgie, M. (2008) A new species of velvet gecko (Diplodactylidae: *Oedura*) from north-east Queensland, Australia. *Zootaxa*, **1788**, 21-36.
- Hotchkiss, P.E., Camp, C.D. & Marshall, J.L. (2001) Aspects of the life history and ecology of the coal skink, *Eumeces anthracinus*, in Georgia. *Journal of Herpetology*, **35**, 145-148.
- How, R.A. & Kitchener, D.J. (1983) The biology of the gecko, *Oedura reticulata* Bustard, in a small habitat isolate in the western Australian wheatbelt. *Australian Wildlife Research*, **10**, 543-556.
- How, R.A., Dell, J. & Wellington, B.D. (1986) Comparative biology of eight species of *Diplodactylus* gecko in western Australia. *Herpetologica*, **42**, 471-482.
- How, R.A., Dell, J. & Gordon, S.J. (1987) Reproductive patterns in chromosomally distinct races of *Phyllodactylus marmoratus* (Lacertilia: Gekkonidae) in southwestern Australia. *Records of the Western Australian Museum*, **13**, 413-418.
- How, R.A., Dell, J. & Wellington, B.D. (1990) Reproductive and dietary biology of *Nephrurus* and *Underwoodisaurus* (Gekkonidae) in Western Australia. *Records of the Western Australian Museum*, **14**, 449-459.
- Huang, W.-S. (1997) Reproductive cycle of the oviparious lizard *Japalura brevipes* (Agamidae: Reptilia) in Taiwan, Republic of China. *Journal of Herpetology*, **31**, 22-29.
- Huang, W.-S. (1998) Reproductive cycles of the grass lizard, *Takydromus hsuehshanensis*, with comments on reproductive patterns of lizards from the central high elevation area of Taiwan. *Copeia*, **1998**, 866-873.
- Huang, W.-S. (1998) Sexual size dimorphism and microhabitat use of two sympatric lizards, *Sphenomorphus taiwanensis* and *Takydromus hsuehshanensis*, from the central highlands of Taiwan. *Zoological Studies*, **37**, 302-308.
- Huang, W.-S. (2006) Ecological characteristics of the skink, *Mabuya longicaudata*, on a tropical East Asian island. *Copeia*, **2006**, 293-300.
- Huang, W.-S. (2006) Ecology and reproductive patterns of the grass lizard, *Takydromus sauteri*, in a tropical rain forest of an east Asian island. *Journal of Herpetology*, **40**, 267-273.
- Huang, W.-S. (2007) Costs of egg caring in the skink, *Mabuya longicaudata*. *Ecological Research*, **22**, 659-664.
- Huang, W.-S. (2007) Ecology and reproductive patterns of the agamid lizard *Japalura swinhonis* on an East Asian island, with comments on the small clutch sizes of island lizards. *Zoological Science*, **24**, 181-188.
- Huang, W.-S. (2010) Ecology and reproductive characteristics of the skink *Sphenomorphus incognitus* on an East Asian Island, with comments on variations in clutch size with

- reproductive modes in *Sphenomorphus*. *Zoological Studies*, **49**, 779-788.
- Huang, W.-S. (2011) Ecology and reproductive patterns of the littoral skink *Emoia atrocostata* on an East Asian tropical rainforest island. *Zoological Studies*, **50**, 506-512.
- Huang, S.-P. & Tu, M.-C. (2008) Heat tolerance and altitudinal distribution of a mountainous lizard, *Takydromus hsuehshanensis*, in Taiwan. *Journal of Thermal Biology*, **33**, 48-56.
- Huey, R.B. & Webster, T.P. (1976) Thermal biology of *Anolis* lizards in a complex fauna: the christatellus group on Puerto Rico. *Ecology*, **57**, 985-994.
- Huey, R.B. & Pianka, E.R. (1977) Seasonal variation in thermoregulatory behavior and body temperature of diurnal Kalahari lizards. *Ecology*, **58**, 1066-1075.
- Huey, R.B. & Pianka, E.R. (1981) Ecological consequences of foraging mode. *Ecology*, **62**, 991-999.
- Huey, R.B. & Bennett, A.F. (1987) Phylogenetic studies of coadaptation: preferred temperatures versus optimal performance temperatures of lizards. *Evolution*, **41**, 1098-1115.
- Huey, R.B. & Pianka, E.R. (2007) Lizard thermal biology: do genders differ? *The American Naturalist*, **170**, 473-478.
- Huey, R.B., Pianka, E.R. & Vitt, L.J. (2001) How often do lizards run on empty? *Ecology*, **82**, 1-7.
- Huey, R.B., Pianka, E.R., Egan, M.E. & Coons, L.W. (1974) Ecological shifts in sympatry: Kalahari fossorial lizards (*Typhlosaurus*). *Ecology*, **55**, 304-316.
- Huey, R.B., Niewiarowski, P.H., Kaufmann, J. & Herron, J.C. (1989) Thermal biology of nocturnal ectotherms: is sprint performance of geckos maximal at low body temperatures? *Physiological Zoology*, **62**, 488-504.
- Huey, R.B., Deutsch, C.A., Tewksbury, J.J., Vitt, L.J., Hertz, P.E., Perez, H.J.A. & Garland, T. (2009) Why tropical forest lizards are vulnerable to climate warming. *Proceedings of the Royal Society of London B: Biological Sciences*, **276**, 1939-1948.
- Hughes, B. (1988) Herpetology on Ghana (West Africa). *British Herpetological Society Bulletin*, **25**, 29-38.
- Hunsaker, D. (1959) Birth and litter sizes of the blue spiny lizard *Sceloporus cyanogenys*. *Copeia*, **1959**, 260-261.
- Hunt, L.E. (2007) *Anniella pulchra*. *Catalogue of American Amphibians and Reptiles*, **849**, 1-3.
- Husak, J.F. & Ackland, E.N. (2003) Foraging mode of the reticulate collared lizard, *Crotaphytus reticulatus*. *Southwestern Naturalist*, **48**, 282-286.
- Hutchinson, M.N. & Donnellan, S.C. (1992) Taxonomy and genetic variation in the Australian lizards of the genus *Pseudemoia* (Scincidae: Lygosominae). *Journal of Natural History*, **26**, 215-264.
- Ibarguengoytia, N.R. (2004) Prolonged cycles as a common reproductive pattern in viviparous lizards from Patagonia, Argentina: reproductive cycle of *Phymaturus patagonicus*. *Journal of Herpetology*, **38**, 73-79.
- Ibarguengoytia, N.R. (2005) Field, selected bodytemperature and thermal tolerance of the syntopic lizards *Phymaturus patagonicus* and *Liolaemus elongatus* (Iguania: Liolaemidae). *Journal of Arid Environments*, **62**, 435-448.
- Ibarguengoytia, N.R. & Cussac, V.E. (1996) Reproductive biology of the viviparous lizard *Liolaemus pictus* (Tropiduridae): biennial female reproductive cycle? *Herpetological Journal*, **6**, 137-143.
- Ibarguengoytia, N.R. & Cussac, V.E. (1998) Reproduction of the viviparous lizard *Liolaemus elongatus* in the highlands of southern South America: plastic cycles in response to climate?

- Herpetological Journal*, **8**, 99-105.
- Ibarguengoytia, N.R. & Cussac, V.E. (1999) Male response to low frequency female reproduction in the viviparous lizard *Liolaemus* (Tropiduridae). *Herpetological Journal*, **9**, 111-117.
- Ibarguengoytia, N.R. & Casalins, L. (2007) Reproductive biology of the southernmost gecko *Homonota darwini*: convergent life-history patterns among southern hemisphere reptiles living in harsh environments. *Journal of Herpetology*, **41**, 71-79.
- Ibarguengoytia, N.R., Renner, M.L., Boretto, J.M., Piantoni, C. & Cussac, V.E. (2007) Thermal effects on locomotion in the nocturnal gecko *Homonota darwini* (Gekkonidae). *Amphibia-Reptilia*, **28**, 235-246.
- Ibarguengoytia, N.R., Ascota, J.C., Boretto, J.M., Villavicencio, H.J., Marinero, J.A. & Krenz, J.D. (2008) Field thermal biology in *Phymaturus* lizards: comparisons from the Andes to the Patagonian steppe in Argentina. *Journal of Arid Environments*, **72**, 1620-1630.
- Ibarguengoytia, N.R., Medina, S.M., Fernandez, J.B., Gutierrez, J.A., Tappari, F. & Sclaro, A. (2010) Thermal biology of the southernmost lizards in the world: *Liolaemus sarmientoi* and *Liolaemus magellanicus* from Patagonia, Argentina. *Journal of Thermal Biology*, **35**, 21-27.
- Ibrahim, A.A. (2008) Contribution to the herpetology of southern Libya. *Acta Herpetologica*, **3**, 35-49.
- Ikeuchi, I. (2004) Male and female reproductive cycles of the Japanese gecko, *Gekko japonicus*, in Kyoto, Japan. *Journal of Herpetology*, **2004**, 269-274.
- Ilgaz, C. (2009) Comparative morphology of *Darevskia parvula* (Lantz-Cyren 1936) (Sauria: Lacertidae) subspecies in Northeastern Anatolia, Turkey. *North-Western Journal of Zoology*, **5**, 263-280.
- In den Bosch, H.A.J. (1988) Erste daten zum fortpflanzungsmodus von *Ophiomorus punctatissimus* (Bibron & Bory, 1833). *Salamandra*, **24**, 53-58.
- In Den Bosch, H.A.J. & Bout, R.G. (1998) Relationships between maternal size, egg size, clutch size and hatchling size in European lacertid lizards. *Journal of Herpetology*, **32**, 410-417.
- In den Bosch, H.A.J. & Zandee, M. (2001) Courtship behaviour in lacertid lizards: phylogenetic interpretations of the *Lacerta kulzeri* complex (Reptilia: Lacertidae). *Netherlands Journal of Zoology*, **51**, 263-284.
- Ineich, I. (1999) Reptiles & amphibiens de la Republic de Djibouti. *Rapport sur la mission herpétologique de Ivan Ineich*.
- Ineich, I. (2008) A new arboreal *Lepidodactylus* (Reptilia: Gekkonidae) from Espiritu Santo Island, Vanuatu: from egg to holotype. *Zootaxa*, **1918**, 26-38.
- Ineich, I. (2011) Amphibians and reptiles. In: *Te natural history of Santo, Publications Scientifiques du Museum*, (eds. by P. Bouchet, H. Le Guyader, and O. Pascal), pp. 187-236, Museum National d'Histoire Naturelle, Paris.
- Inger, R.F. (1959) Temperature responses and ecological relations of two Bornean lizards. *Ecology*, **40**, 127-136.
- Inger, R.F. (1983) Morphological and ecological variation in the flying lizards (genus *Draco*). *Fieldiana Zoology, New Series*, **18**, 1-35.
- Inger, R.F. & Greenberg, B. (1966) Annual reproductive patterns of lizards from a Bornean rain forest. *Ecology*, **47**, 1007-1021.
- Inger, R.F. & Colwell, R.K. (1977) Organization of contiguous communities of amphibians and reptiles in Thailand. *Ecological Monographs*, **47**, 229-253.

- Inger, R.F., Bradley, S.H., Mammen, K. & Ramesh, B. (1984) A report on a collection of amphibians and reptiles from the Ponmudi, Kerala, South India. *Journal of the Bombay Natural History Society*, **81**, 551-570.
- Inger, R.F., Zhao, E.-M., Shaffer, H.B. & Wu, G.-F. (1990) Report on a collection of amphibians and reptiles from Sichuan, China. *Fieldiana, Zoology*, **58**, 1-24.
- Ingram, G.J. & Covacevich, J. (1989) Revision of the genus *Carlia* (Reptilia, Scincidae) in Australia with comments on *Carlia bicarinata* of New Guinea. *Memoires of the Queensland Museum*, **27**, 443-490.
- Irschick, D.J., Vitt, L.J., Zani, P.A. & Losos, J.B. (1997) A comparison of evolutionary radiations in mainland and Caribbean Anolis lizards. *Ecology*, **78**, 2191-2203.
- Irschick, D.J., Austin, C.C., Petren, K., Fisher, R.N., Losos, J.B. & Ellers, O. (1996) A comparative analysis of clinging ability among pad-bearing lizards. *Biological Journal of the Linnean Society*, **59**, 21-35.
- Iverson, J.B., Pasachnik, S., Knapp, C.R. & Buckner, S.D. (2005) *Cyclura cyclura*. *Catalogue of American Amphibians and Reptiles*, **810**, 1-8.
- Iverson, J.I., Converse, S.J., Smith, G.R. & Valiulis, J.M. (2006) Long-term trends in the demography of the Allen Cays rock iguana (*Cyclura cyclura inornata*): human disturbance and density-dependent effects. *Biological Conservation*, **132**, 300-310.
- J., C.C., Dessauer, H.C., Townsend, C.R. & Arnold, M.G. (1990) Unisexual lizards of the Genus *Gymnophthalmus* (Reptilia: Teiidae) in the Neotropics: genetics, origin, and systematics. *American Museum Novitates*, **2994**, 1-29.
- Jackson, K. & Blackburn, D.C. (2010) A survey of amphibians and reptiles at degraded sites near Pointe-Noire, Kouilou Province, Republic of Congo. *Herpetological Conservation and Biology*, **5**, 414-429.
- Jacobs, H.J. (2003) A further new emerald tree monitor lizard of the *Varanus prasinus* species group from Waigeo, West Irian (Squamata: Sauria: Varanidae). *Salamandra*, **39**, 65-74.
- Jacobsen, N.H.G., Pietersen, E.W. & Pietersen, D.W. (2010) A preliminary herpetological survey of the Vilanculos Coastal Wildlife Sanctuary on the San Sebastian Peninsula, Vilankulo, Mozambique. *Herpetology Notes*, **3**, 181-193.
- Jacobson, N.H.G. & Broadley, D.G. (2000) A new species of *Panaspis* Cope (Reptilia: Scincidae) from southern Africa. *African Journal of Herpetology*, **49**, 61-71.
- Jaksic, F.M., Nunez, H. & Ojeda, F.P. (1980) Body proportions, microhabitat selection, and adaptive radiation of *Liolaemus* lizards in central Chile. *Oecologia*, **45**, 178-181.
- Jaksic, F.M., Greene, H.W. & Schwenk, K.a.S., R. L. (1982) Predation upon reptiles in Mediterranean habitats of Chile, Spain and California: a comparative analysis. *Oecologia*, **53**, 152-159.
- James, C.D. (1991) Annual variation in reproductive cycles of scincid lizards, *Ctenotus*, in central Australia. *Copeia*, **1991**, 744-760.
- James, C. & Shine, R. (1988) Life-history strategies of Australian lizards: a comparison between the tropics and the temperate zone. *Oecologia*, **75**, 307-316.
- James, C.D. & Losos, J.B. (1991) Diet and reproductive biology of the Australian sand-swimming lizards, *Eremiascincus* (Scincidae). *Wildlife Research*, **18**, 641-654.
- James, C.D., Losos, J.B. & King, D.R. (1992) Reproductive biology and diets of goannas (Reptilia: Varanidae) from Australia. *Journal of Herpetology*, **26**, 128-136.
- Janzen, D.H. (1973) Sweep samples of tropical foliage insects: description of study sites, With data on species abundances and size distributions. *Ecology*, **54**, 659-686.

- Jennings, M.R. (1995) *Gambelia sila*. *Catalogue of American Amphibians and Reptiles*, **612**, 1-4.
- Jenssen, T.A. (1990) *Anolis cooki*. *Catalogue of American Amphibians and Reptiles*, **488**, 1-2.
- Jenssen, T.A. (1973) Shift in the structural habitat of *Anolis opalinus* due to congeneric competition. *Ecology*, **54**, 863-869.
- Jenssen, T.A. & Nunez, S.C. (1994) Male and female reproductive cycles of the Jamaican lizard, *Anolis opalinus*. *Copeia*, **1994**, 767-780.
- Jessop, T.S., Chan, R. & Stuart-Fox, D. (2009) Sex steroid correlates of female-specific colouration, behaviour and reproductive state in Lake Eyre dragon lizards, *Ctenophorus maculosus*. *Journal of Comparative Physiology*, **195**, 619-630.
- Ji, Q., Luo, Z.-X., Yuan, C.-X., . & Tabrum, A.R. (2006) A swimming mammaliaform from the Middle Jurassic and ecomorphological diversification of early mammals. *Science*, **311**, 1123-1127.
- Ji, X., Luo, Z.X., Yuan, C.X., Wible, J.R., Zhang, J.P. & Georgi, J.A. (2002) The earliest known eutherian mammal. *Nature*, **416**, 816-822.
- Jimenez-Cruz, E., Ramirez-Bautista, A., Marshall, J.C., Lizana-Avia, M. & De Oca, A.N.-M. (2005) Reproductive cycle of *Sceloporus grammicus* (Squamata: Phrynosomatidae) from Teotihuacan, Mexico. *Southwestern Naturalist*, **50**, 178-187.
- Joger, U. & Lambert, M.R.K. (1996) Analysis of the herpetofauna of the Republic of Mali, I. Annotated inventory, with description of a new *Uromastix* (Sauria: Agamidae). *Journal of African Zoology*, **110**, 21-51.
- Johnson, C.R. & Voigt, W.G. (1978) Observations on thermoregulation in the western slender glass lizard, *Ophisaurus attenuatus attenuatus* (Sauria: Anguidae). *Zoological Journal of the Linnean Society*, **63**, 305-307.
- Johnson, M.A., Caton, J.L., Cohen, R.E., Vandecar, J.R. & Wade, J. (2010) The burden of motherhood: the effect of reproductive load on female lizard locomotor, foraging, and social behavior. *Ethology*, **116**, 1217-1225.
- Johnson, M.A., Leal, M., Schettino, L.R., Lara, A.C., Revell, L.J. & Losos, J.B. (2008) A phylogenetic perspective on foraging mode evolution and habitat use in West Indian *Anolis* lizards. *Animal Behaviour*, **75**, 555-563.
- Johnston, G. (1999) Reproductive biology of the peninsula dragon lizard, *Ctenophorus fionni*. *Journal of Herpetology*, **33**, 694-698.
- Johnston, G. (2005) *Ctenophorus fionni* (peninsula dragon lizard) cannibalism. *Herpetological Review*, **36**, 175.
- Johnston, G. & Bouskila, A. (2007) Sexual dimorphism and ecology of the gecko, *Ptyodactylus guttatus*. *Journal of Herpetology*, **41**, 506-513.
- Jones, S. (1987) A report on a reproducible and sustainable system for the captive propagation of the genus *Tiliqua* Gray 1825. *Conference Proceedings*, 17-25.
- Jonker, L. & Bullen, L. (2004) This Turquoise Coast Island Nature Reserves Management Plan. In. Department of Conservation and Land Management, Perth, Western Australia.
- Judd, F.W. (1975) Activity and thermal ecology of the keeled earless lizard, *Holbrookia propinqua*. *Herpetologica*, **31**, 137-150.
- Judd, F.W. (1976) Food and feeding behavior of the keeled earless lizard, *Holbrookia propinqua*. *Southwestern Naturalist*, **21**, 17-26.
- Judd, F.W. (1978) Year-to-year variation in clutch size of island and mainland populations of *Holbrookia propinqua* (Reptilia, Lacertilia, Iguanidae). *Journal of Herpetology*, **12**, 203-

- Kaiser, H., Carvalho, V.L., Ceballos, J., Freed, P., Heacox, S., Lester, B., Richards, S.J., Trainor, C.R., Sanchez, C. & O'Shea, M. (2011) The herpetofauna of Timor-Leste: a first report. *ZooKeys*, **109**, 19-86.
- Kaliontzopoulou, A., Carretero, M.A. & Llorente, G.A. (2008) Head shape allometry and proximate causes of head sexual dimorphism in *Podarcis* lizards: joining linear and geometric morphometrics. *Biological Journal of the Linnean Society*, **93**, 111-124.
- Kaliontzopoulou, A., Carretero, M.A. & Llorente, G.A. (2010) Intraspecific ecomorphological variation: linear and geometric morphometrics reveal habitat-related patterns within *Podarcis bocagei* wall lizards. *Journal of Evolutionary Biology*, **23**, 1234-1244.
- Kamel, S. & Gatten, R.E. (1983) Aerobic and anaerobic activity metabolism of limbless and fossorial reptiles. *Physiological Zoology*, **56**, 419-429.
- Karges, J.P. & Wright, J.W. (1987) A new species of *Barisia* (Sauria, Anguillidae) from Oaxaca, Mexico. *Contributions in Science of the Natural History Museum of Los Angeles County*, **381**, 1-11.
- Kearney, M. (2003) Diet in the Amphisbaenian *Bipes biporus*. *Journal of Herpetology*, **37**, 404-408.
- Kennedy, J.P. (1968) Observations on the ecology and behavior of *Cnemidophorus guttatus* and *Cnemidophorus deppei* (Sauria, Teiidae) in Southern Veracruz. *Journal of Herpetology*, **2**, 87-96.
- Kennedy, J.P. (1973) *Sceloporus olivaceus*. *Catalogue of American Amphibians and Reptiles*, **143**, 1-4.
- Keogh, J.S., Edwards, D.L., Fisher, R.N. & Harlow, P.S. (2008) Molecular and morphological analysis of the critically endangered Fijian iguanas reveals cryptic diversity and a complex biogeographic history. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **363**, 3413-3426.
- Kerr, A.M., Powell, R. & Parmerlee, J.S. (2005) *Ameiva erythrocephala* (Teiidae) on Sint Eustatius, Netherlands Antilles: baseline data on a small population in a severely altered habitat. *Caribbean Journal of Science*, **41**, 162-169.
- Kiefer, M.C., Van Sluys, M. & Rocha, C.F.D. (2008) Clutch and egg size of the tropical lizard *Tropidurus torquatus* (Tropiduridae) along its geographic range in coastal eastern Brazil. *Canadian Journal of Zoology*, **86**, 1376-1388.
- Kiester, A.R. (1982) A new forest skink from Ponape. *Breviora*, **468**, 1-10.
- King, D., Green, B. & Butler, H. (1989) The activity pattern, temperature regulation and diet of *Varanus giganteus* on Barrow Island, Western Australia. *Australian Wildlife Research*, **16**, 41-47.
- Kirchhof, S., Linden, J., Rodder, D. & Richter, K. (2010) Daily activity patterns of *Australolacerta rupicola* (FitzSimons, 1933) (Sauria: Lacertidae) with comments on niche segregation within a syntopic lizard community. *North-Western Journal of Zoology*, **6**, 172-181.
- Kirshner, D.S. (2007) Multiclutching in captive lace monitors (*Varanus varius*). *Mertensiella*, **16**, 403-421.
- Klauber, L.M. (1931) A new species of *Xantusia* from Arizona, with a synopsis of the genus. *Transactions of the San Diego Society of Natural History*, **7**, 1-16.
- Klauber, L.M. (1938) Notes from a herpetological diary I. *Copeia*, **1938**, 191-197.
- Klaver, C. & Bohme, W. (1992) The species of *Chamaeleo cristatus* group from Cameroon and

- adjacent countries, West Africa. *Bonner Zoologische Beiträge*, **43**, 433-476.
- Kluge, A.G. (1964) A revision of the South American gekkonid lizard genus *Homonota* Gray. *American Museum Novitates*, **2193**, 1-42.
- Koch, A.A., M. & Ziegler, T. (2010) Updated checklist of the living monitor lizards of the world (Squamata: Varanidae). *Bonn Zoological Bulletin*, **57**, 127-136.
- Koch, A., Auliya, M., Schmitz, A., Kuch, U. & Bohme, W. (2007) Morphological studies on the systematics of south east Asian water monitors (*Varanus salvator* Complex): nominotypic populations and taxonomic overview. *Mertensiella*, **16**, 109-180.
- Koenig, J., Shine, R. & Shea, G. (2001) The ecology of an Australian reptile icon: how do blue-tongued lizards (*Tiliqua scincoides*) survive in suburbia? *Wildlife Research*, **28**, 215-227.
- Koetz, A. (2004) *Hemisphaeriodon gerrardii*. *Herpetofauna of North Queensland*,
- Kohler, G. (1996) A new species of anole of the *Norops pentaprion* group from Isla de Utila, Honduras. (Reptilia: Sauria: Iguanidae). *Senckenbergiana Biologica*, **75**, 23-31.
- Kohler, G. (2003) A new species of *Morunasaurus* from Peru (Reptilia, Squamata, Hoplocercidae). *Senckenbergiana Biologica*, **82**, 235-241.
- Kohlsdorf, T. & Navas, C.A. (2006) Ecological constraints on the evolutionary association between field and preferred temperatures in Tropicurinae lizards. *Evolutionary Ecology*, **20**, 549-564.
- Kohlsdorf, T., Garland, T. & Navas, C.A. (2001) Limb and tail lengths in relation to substrate usage in *Tropidurus* lizards. *Journal of Morphology*, **248**, 151-164.
- Kolbe, J.J., Colbert, P.L. & Smith, B.E. (2008) Niche relationships and interspecific interactions in Antigua lizard communities. *Copeia*, **2008**, 261-272.
- Kolodiuk, M.F., Ribeiro, L.B. & Freire, E.M.X. (2010) Diet and foraging behavior of two species of *Tropidurus* (Squamata, Tropicuridae) in the Caatinga of northeastern Brazil. *South American Journal of Herpetology*, **5**, 35-44.
- Koul, O. & Duda, P.L. (1977) Ovarian cycle in high altitude lizards from Kashmir. *Copeia*, **1977**, 751-754.
- Kratochvil, L. & Frynta, D. (2006) Body-size effect on egg size in eublepharid geckos (Squamata: Eublepharidae), lizards with invariant clutch size: negative allometry for egg size in ectotherms is not universal. *Biological Journal of the Linnean Society*, **88**, 527-532.
- Kraus, F. (2005) New species of blindsnake from Rossel Island, Papua New Guinea. *Journal of Herpetology*, **39**, 591-595.
- Krekorian, C.O. (1983) Population density of the desert iguana, *Dipsosaurus dorsalis* (Reptilia: Iguanidae), in Southern California. *Copeia*, **1983**, 268-271.
- Krysko, K.L.S.C.M. & Hooper, A.N. (2003) Interspecific communal oviposition and reproduction of four species of lizards (Sauria: Gekkonidae) in the lower Florida Keys. *Amphibia-Reptilia*, **24**, 390-396.
- Kumluta, Y., Durmu, S.H., Kaska, Y., M., O. & Tunc, M.R. (2004) A morphological and taxonomic study on *Lacerta parva* Boulenger, 1887 (Sauria: Lacertidae) from West Taurus, Turkey. *Asiatic Herpetological Research*, **10**, 202-207.
- Kumlutas, Y., Oz, M., Durmus, H., Tunc, M.R., Ozdemur, A. & Dusen, S. (2004) On some lizard species of the Western Taurus range. *Turkish Journal of Zoology*, **28**, 225-236.
- Kuo, C.-Y., Lin, Y.-T. & Yao-Sung Lin, Y.-S. (2009) Sexual size and shape dimorphism in an agamid lizard, *Japalura swinhonis* (Squamata: Lacertilia: Agamidae). *Zoological Studies*, **48**, 351-361.
- Kutt, A.S., Bateman, B.L. & Vanderduys, E.P. (2011) Lizard diversity on a rainforest-savanna

- altitude gradient in north-eastern Australia. *Australian Journal of Zoology*, **59**, 86-94.
- Lachman, E., Carmely, H. & Werner, Y.L. (2006) Subspeciation befogged by the ‘‘Seligmann effect’’: the case of *Laudakia stellio* (Reptilia: Sauria: Agamidae) in southern Sinai, Egypt. *Journal of Natural History*, **40**, 1259-1284.
- Lai-Gao, L., Guo-Hua, D. & Xiang, J. (2010) Income breeding and temperature-induced plasticity in reproductive traits in lizards. *Journal of Experimental Biology*, **213**, 2073-2078.
- Lais, P.M. (1976) *Gerrhonotus multicarinatus*. *Catalogue of American Amphibians and Reptiles*, **187**, 1-4.
- Lalremsanga, H.T., Khawlhiring, L. & Lalrotluanga (2010) Three additional lizard (Squamata: Sauria) records for Mizoram, India. *Journal of Threatened Taxa*, **2**, 718-720.
- Lanza, B. & Vanni, S. (1976) On a small collection of reptiles from north Zaire. *Monitore Zoologico Italiano*, **8**, 129-160.
- Lappin, A.K. & Swinney, E.J. (1999) Sexual dimorphism as it relates to natural history of leopard lizards (Crotaphytidae: Gambelia). *Copeia*, **1999**, 649-660.
- Lara-Gongora, G. (1983) Two new species of the lizard genus *Sceloporus* (Reptilia, Sauria, Iguanidae) from the Ajusco and Ocuilan Sierras, Mexico. *Bulletin of the Maryland Herpetological Society*, **19**, 1-14.
- Largen, M.J. & Spawls, S. (2006) Lizards of Ethiopia (Reptilia Sauria): an annotated checklist, bibliography, gazetteer and identification key. *Tropical Zoology*, **19**, 21-109.
- Laurent, R.F. (1964) Reptiles et amphibiens de l'Angola (Troisieme contribution). *Companhia de Diamantes de Angola (Diamang), Servicos Culturais, Museu do Dundo (Angola)*, **67**, 1-165.
- Lazell, J.D. (1972) The anoles (Sauria: Iguanidae) of the Lesser Antilles. *Bulletin of the Museum of Comparative Zoology*, **143**, 1-115.
- Lazell, J.D. (1973) The lizard genus *Iguana* in the Lesser Antilles. *Bulletin of the Museum of Comparative Zoology*, **145**, 1-28.
- Leache, A.D. (2005) Results of a herpetological survey in Ghana and a new country record. *Herpetological Review*, **36**, 16-19.
- Leache, A.D., Rodel, M.O., Linkem, C.W., Diaz, R.E., Hillers, A. & Fujita, M.K. (2006) Biodiversity in a forest island: reptiles and amphibians of the West African Togo Hills. *Amphibian and Reptile Conservation*, **4**, 22-45.
- Leal, M., Knox, A.K. & Losos, J.B. (2002) Lack of convergence in aquatic *Anolis* lizards. *Evolution*, **56**, 785-791.
- Leclair, R. & Leclair, M.H. (2011) Life-history traits in a population of the dwarf gecko, *Sphaerodactylus vincenti ronaldi*, from a xerophytic habitat in Martinique, West Indies. *Copeia*, **2011**, 566-576.
- Lee, J.C. (1976) *Xantusia henshawi*. *Catalogue of American Amphibians and Reptiles*, **189**, 1-2.
- Lee, D.S. & Funderburg, J.B. (1977) *Sceloporus woodi*. *Catalogue of American Amphibians and Reptiles*, **196**, 1-2.
- Lee, G.H., Lee, K. & Watkins, J.S. (2001) Geologic evolution of the Cuu Long and Nam Con Son basins, offshore southern Vietnam, South China Sea. *AAPG bulletin*, **85**, 1055-1082.
- Lemay, L. & Marsiglia, A.G. (1952) The coal skink, *Eumeces anthracinus* (Baird), in Maryland. *Copeia*, **1952**, 193.
- Lemm, J.M., Steward, S.W. & Schmidt, T.F. (2005) Reproduction of the critically endangered Anegada Island iguana *Cyclura pinguis* at San Diego Zoo. *International Zoo Yearbook*, **39**, 141-152.

- Lemos-Espinal, J.A., Smith, G.R. & Ballinger, R.E. (1996) Natural history of the Mexican knob-scaled lizard, *Xenosaurus rectocollaris*. *Herpetological Natural History*, **4**, 151-154.
- Lemos-Espinal, J.A., Smith, G.R. & Ballinger, R.E. (1999) Reproduction in Gadow's spiny lizard, *Sceloporus gadovae* (Phrynosomatidae), from arid tropical Mexico. *Southwestern Naturalist*, **44**, 57-63.
- Lemos-Espinal, J.A., Smith, G.R. & Ballinger, R.E. (2001) Sexual dimorphism and body temperatures of *Sceloporus siniferus* from Guerrero Mexico. *Western North American Naturalist*, **61**, 498-500.
- Lemos-Espinal, J.A., Smith, G.R. & Ballinger, R.E. (2003) Diets of three species of knob-scaled lizards (genus *Xenosaurus*) from Mexico. *Southwestern Naturalist*, **48**, 119-122.
- Lemos-Espinal, J.A., Smith, G.R. & Ballinger, R.E. (2004) Aspects of the ecology of a distinct population of *Xenosaurus platyceps* from Queretaro, Mexico. *Amphibia-Reptilia*, **25**, 204-210.
- Lerner, A. (2004) A new taxonomically isolated species of the genus *Phelsuma* GRAY, 1825 from the Ampasindava peninsula Madagascar. *Phelsuma*, **12**, 91-98.
- Levinson, S.C. (2008) Landscape, seascape and the ontology of places on Rossel Island, Papua New Guinea. *Language Sciences*, **30**, 256-290.
- Leviton, A.E. & Anderson, S.C. (1970) The amphibians and reptiles of Afghanistan, a checklist and key to the herpetofauna. *Proceedings of the California Academy of Sciences*, **38**, 163-206.
- Lewis, A.R., Tirado, G. & Sepulveda, J. (2000) Body size and paternity in a teiid lizard (*Ameiva exsul*). *Journal of Herpetology*, **34**, 110-120.
- Lezzi, A. (2008) *Husbandry manual for southern leaf-tailed gecko* *Phyllurus platurus* Reptilia: Gekkonidae. Captive Animals, Western Sydney Institute of TAFE, Richmond.
- Licht, P. & Gorman, G.C. (1970) Reproductive and fat cycles in Caribbean *Anolis* lizards. *University of California Publications in Zoology*, **35**, 1-52.
- Licht, P., Dawson, W.R. & Shoemaker, V.R. (1966) Heat resistance of some Australian lizards. *Copeia*, **1966**, 162-169.
- Lieb, C.S. (1985) Systematics and distribution of the skinks allied to *Eumeces tetragrammus* (Sauria: Scincidae). *Los Angeles County Museum Contributions in Science*, **357**, 1-19.
- Lieb, C.S. (1990) *Eumeces tetragrammus*. *Catalogue of American Amphibians and Reptiles*, **492**, 1-4.
- Light, P., Dawson, W.R., Shoemaker, V.H. & Main, A.R. (1966) Observations on the thermal relations of Western Australian lizards. *Copeia*, **1966**, 97-110.
- Lin, J.-Y. & Nelson, C. (1980) Comparative reproductive biology of two sympatric tropical lizards *Chamaeleo jacksonii* Boulenger and *Chamaeleo hoehnelii* Steindachner (Sauria: Chamaeleonidae). *Amphibia-Reptilia*, **1**, 287-311.
- Linkem, C.W., Diesmos, A.C. & Brown, R.M. (2010) A new species of scincid lizard (Genus *Sphenomorphus*) from Palawan Island, Philippines. *Journal Information*, **66**, 67-79.
- Linsdale, J.M. (1932) Amphibians and reptiles from lower California. *University of California Publications in Zoology*, **38**, 345-386.
- Ljubisavljevic, K., Dzukic, G. & Kalezeic, M.L. (2008) Female reproductive life history traits of the meadow lizard, *Darevskia praticola* (Eversmann 1834) from the westernmost boundary of the species range. *Polish Journal of Ecology*, **56**, 289-297.
- Lonnberg, E. (1911) Reptiles, batrachians and fishes collected by the Swedish Zoological Expedition to British East Africa 1911. *Kungliga Svenska Vetenskaps-Akademiens*

- Handlingar*, **47**, 1-24.
- Lopez, P., Civantos, E. & Martín, J. (2002) Body temperature regulation in the amphisbaenian *Trogonophis wiegmanni*. *Canadian Journal of Zoology*, **80**, 42-47.
- Lopez-Ortiz, R. & Lewis, A.R. (2002) Seasonal abundance of hatchlings and gravid females of *Sphaerodactylus nicholsi* in Cabo Rojo, Puerto Rico. *Journal of Herpetology*, **36**, 276-280.
- Lopez-Victoria, M. (2006) The lizards of Malpelo (Colombia): some topics on their ecology and threats. *Caldasia*, **28**, 129-134.
- Lorvelec, O., Pascal, M., Pavis, C. & Feldmann, P. (2007) Amphibians and reptiles of the French West Indies: inventory, threats and conservation. *Applied Herpetology*, **4**, 131-161.
- Losos, J.B. & Greene, H.W. (1988) Ecological and evolutionary implications of diet in monitor lizards. *Biological Journal of the Linnean Society*, **35**, 379-407.
- Losos, J.B., Marks, J.C. & Schoener, T.W. (1993) Habitat use and ecological interactions of an introduced and a native species of *Anolis* lizard on Grand Cayman, with a review of the outcomes of anole introductions. *Oecologia*, **95**, 525-532.
- Loveridge, A. (1936) African reptiles and amphibians in Field Museum of Natural History. *Publications of the Field Museum of Natural History, Zoology*, **22**, 1-111.
- Loveridge, A. (1941) *Bogertia lutzae* – A new genus and species of gecko from Bahia, Brazil. *Proceedings of the Biological Society of Washington*, **54**, 195-196.
- Loveridge, A. (1942) Scientific results of a fourth expedition to forested areas in east and central Africa. IV. Reptiles. *Bulletin of the Museum of Comparative Zoology*, **91**, 237-373.
- Loveridge, A. (1944) Revision of the African lizards of the family Cordylidae. *Bulletin of the Museum of Comparative Zoology*, **95**, 1-118.
- Loveridge, A. (1947) Revision of the African lizards of the family Gekkonidae. *Bulletin of the Museum of Comparative Zoology*, **98**, 1-469.
- Loveridge, A. (1948) New Guinean reptiles and amphibians in the Museum of Comparative Zoology and the United States National Museum. *Bulletin of the Museum of Comparative Zoology*, **101**, 305-430.
- Loveridge, A. (1953) Zoological Results of a fifth expedition to East Africa. III. Reptiles from Nyasaland and Tete. *Bulletin of the Museum of Comparative Zoology*, **110**, 142-322.
- Loveridge, A. (1955) On a second collection of reptiles and amphibians taken in Tanganyika Territory by C. J. P. Ionides. *Journal of the East African Natural History Society*, **22**, 169-198.
- Loveridge, A. (1959) Notes on the present herpetofauna of Ascension Island. *Copeia*, **1959**, 69-70.
- Lue, K.-Y. & Lin, S.-M. (2008) Two new cryptic species of *Takydromus* (Squamata: Lacertidae) from Taiwan. *Herpetologica*, **64**, 379-395.
- Luiselli, L., Akani, G.C. & Capizzi, D. (1998) Food resource partitioning of a community of snakes in a swamp rainforest of south-eastern Nigeria. *Journal of Zoology*, **246**, 125-133.
- Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. & Mylonas, M. (2008) Two new species of *Podarcis* (Squamata; Lacertidae) from Greece. *Systematics and Biodiversity*, **6**, 307-318.
- MacMillen, R.E., Augee, M.L. & Ellis, B.A. (1989) Thermal ecology and diet of some xerophilous lizards from western New South Wales. *Journal of Arid Environments*, **16**, 193-201.
- Mahony, S. & Ali Reza, A.H.M. (2008) A herpetofaunal collection from the Chittagong hill tracts, Bangladesh, with two new species records for the country. *Hamadryad*, **32**, 45-56.

- Maisano, J.A. (2001) A survey of state of ossification in neonatal squamates. *Herpetological Monographs*, **15**, 135-157.
- Malhotra, A. & Thorpe, R.S. (1995) *Ameiva fuscata*. *Catalogue of American Amphibians and Reptiles*, **606**, 1-3.
- Malonza, P.K., Wasonga, V.D., Muchai, V., Rotich, D., Bwong, B.A. & Bauer, A.M. (2006) Diversity and biogeography of herpetofauna of the Tana River Primate National Reserve, Kenya. *Journal of East African Natural History*, **95**, 95-109.
- Manicom, C. & Schwarzkopf, L. (2011) Diet and prey selection of sympatric tropical skinks. *Austral Ecology*, **6**, 485-496.
- Mann, S.L. & Meek, R. (2004) Understanding the relationship between body temperature and activity patterns in the giant Solomon Island skink, *Corucia zebrata*, as a contribution to the effectiveness of captive breeding programmes. *Applied Herpetology*, **1**, 287-298.
- Manriquez-Moran, N.L., VillagraN-Santa Cruz, M. & Mendez de la Cruz, F.R. (2005) Reproductive biology of the parthenogenetic lizard, *Aspidoscelis cozumela*. *Herpetologica*, **61**, 435-439.
- Maragou, P., Chondropoulos, B. & Valakos, E.D. (1999) Comparative data on reproduction in *Podarcis erhardii*, *Podarcis peloponnesiaca* and *Podarcis taurica* (Reptilia, Sauria, Lacertidae). *Israel Journal of Zoology*, **45**, 487-496.
- Marquet, P.A., Bozinovic, F., Medel, R.G., Werner, Y.L. & Jaksic, F.M. (1990) Ecology of *Garthia gaudichaudi*, a gecko endemic to the semiarid region of Chile. *Journal of Herpetology*, **24**, 431-434.
- Marquez, C.M.B. & Marquez, L.D.R. (2009) Reproductive biology in the wild and in captivity of *Anolis aquaticus* (Sauria: Polychrotidae) in Costa Rica. *Laboratorios IASA Sangolqui, Ecuador, Boletín Técnico 8, Serie Zoológica*, **4-5**, 50-73.
- Marshall, M. & Fosberg, F.R. (1975) The Natural History of Namoluk Atoll, Eastern Caroline Islands with Identifications of Vascular Flora by F. R. Fosberg. *Atoll Research Bulletin*, **189**, 69.
- Marshall, A.J. & Beehler, B.M. (2007) *Ecology of Indonesian Papua Part One*. Periplus Editions (HK) Ltd.
- Martín, J. & Salvador, A. (1997) Microhabitat selection by the Iberian rock-lizard *Lacerta monticola*: effects on density and spatial distribution of individuals. *Biological Conservation*, **79**, 303-307.
- Martens, H. (1997) A review of zoogeography of amphibians and reptiles of Syria, with additional new records (Herpetozoa 9 (1/2), 1996). *Herpetozoa*, **10**, 99-106.
- Martin, J. & Lopez, P. (2010) Thermal constraints of refuge use by Schreiber's green lizards, *Lacerta schreiberi*. *Behaviour*, **147**, 275-284.
- Martin, J., Polo-Cavia, N., Gonzalo, A., Lopez, P. & Civantos, E. (2011) Social aggregation behaviour in the North African amphisbaenian *Trogonophis wiegmanni*. *African Journal of Herpetology*,
- Martin, J., Polo-Cavia, N., Gonzalo, A., Lopez, P. & Civantos, E. (2011) Structure of a population of the amphisbaenian *Trogonophis wiegmanni* In North Africa. *Herpetologica*, **67**, 250-257.
- Martinez-Torres, M., Hernandez-Caballero, M.E., Alvarez-Rodriguez, C., Luis-Diaz, J.A. & Ortizz-Lopez, G. (2003) Luteal development and progesterone levels during pregnancy of the viviparous temperate lizard *Barisia imbricata imbricata* (Reptilia: Anguillidae). *General and Comparative Endocrinology*, **132**, 55-65.

- Martins, M. (1991) The lizards of Balbina, Central Amazonia, Brazil: a qualitative analysis of resource utilization. *Studies on Neotropical Fauna and Environment*, **26**, 179-190.
- Martins, E.P. (1993) A comparative study of the evolution of *Sceloporus* push-up displays. *American Naturalist*, **142**, 994-1018.
- Martins, M. (2006) Life in the water: ecology of the jacarerana lizard, *Crocodylurus amazonicus*. *Herpetological Journal*, **16**, 171-177.
- Martori, R. & Aun, L. (2010) Reproduccion y variacion de grupos de tamano en una poblacion de *Liolaemus koslowskyi* (Squamata: Liolaemini). *Cuadernos de Herpetologia*, **24**, 39-55.
- Mata-Silva, V., Ramirez-Bautista, A. & Johnson, J.D. (2010) Reproductive characteristics of two syntopic whiptail lizards, *Aspidoscelis marmorata* and *Aspidoscelis tessellata*, from the northern Chihuahuan Desert. *Southwestern Naturalist*, **55**, 125-129.
- Mautz, W.J. & Nagy, K.A. (2000) Xantusiid lizards have low energy, water, and food requirements. *Physiological and Biochemical Zoology*, **73**, 480-487.
- Mayes, P.J., Bradshaw, S.D. & Bradshaw, F.J. (2007) Reproductive seasonality in the semi-aquatic monitor *Varanus mertensi* (Reptilia: Varanidae). *Mertensiella*, **16**, 322-335.
- McAlpin, S., Duckett, P. & Stow, A. (2011) Lizards cooperatively tunnel to construct a long-term home for family members. *PLoS ONE*, **6**, e19041.
- McBrayer, L.D. (2004) The relationship between skull morphology, biting performance and foraging mode in Kalahari lacertid lizards. *Zoological Journal of the Linnean Society*, **140**, 403-416.
- McBrayer, L.D. & Anderson, R.A. (2007) Sexual size dimorphisms and bite force in the northern alligator lizard, *Elgaria coerulea*. *Journal of Herpetology*, **41**, 554-559.
- McCoid, M.J. (1994) Egg, hatchling sizes, and ovipositor sites of lizards on Guam, Mariana Islands. *Herpetological Review*, **25**, 98-100.
- McConkey, E.H. (1954) A systematic study of the North American lizards of the genus *Ophisaurus*. *American Midland Naturalist*, **51**, 133-171.
- McConnachie, S., Alexander, G.J. & Whiting, M.J. (2009) Selected body temperature and thermoregulatory behavior in the sit-and-wait foraging lizard *Pseudocordylus melanotus melanotus*. *Herpetological Monographs*, **23**, 108-122.
- McCoy, M. & Webber, P. (1984) Two new species of scincid lizards of the genus *Emoia* from Santa Cruz and Duff Islands, Solomon Islands. *Copeia*, **1984**, 571-578.
- McCoy, E.D., Hartmann, P.P. & Mushinsky, H.R. (2004) Population biology of the rare Florida scrub lizard in fragmented habitat. *Herpetologica*, **60**, 54-61.
- McCranie, J.R. & Wilson, L.D. (2001) Herpetofauna of the Mexican state of Aguascalientes. *Courier Forschungsinstitut Senckenberg*, **230**, 1-57.
- McCranie, J.R. & Castaneda, F.E. (2005) The herpetofauna of Parque Nacional Pico Bonito, Honduras. *Phyllomedusa*, **4**, 3-16.
- McElroy, E.J., Hickey, K.L. & Reilly, S.M. (2008) The correlated evolution of biomechanics, gait and foraging mode in lizards. *Journal of Experimental Biology*, **211**, 1029-1040.
- Mcgrann, M.C., Wright, G.R., Dial, R.J. & Mcgrann, A.M. (2006) Off-highway vehicle impact on the flat-tailed horned lizard, *Phrynosoma mcallii*, in the Colorado Desert of southern California. *California Fish and Game*, **92**, 67-80.
- McGuire, J.A. (1994) A new species of collared lizard (Iguania: Crotaphytidae) from northeastern Baja California, Mexico. *Herpetologica*, **50**, 438-450.
- McGuire, J.A. (1996) Phylogenetic systematics of crotaphytid lizards. *Bulletin of the Carnegie Museum*, **32**, 1-143.

- McGuire, J.A. & Alcalá, A.C. (2000) A taxonomic revision of the flying lizards of the Philippine Islands (Iguania: Agamidae: *Draco*), with a description of a new species. *Herpetological Monographs*, **14**, 81-138.
- McGUIRE, J.A. & Heang, K.B. (2001) Phylogenetic systematics of Southeast Asian flying lizards (Iguania: Agamidae: *Draco*) as inferred from mitochondrial DNA sequence data. *Biological Journal of the Linnean Society*, **72**, 203-229.
- McGuire, J.A., Brown, R.M., Mumpuni, R., A. & Andayani, N. (2007) The flying lizards of the *Draco lineatus* group (Squamata: Iguania: Agamidae): a taxonomic revision with descriptions of two new species. *Herpetological Monographs*, **21**, 179-212.
- McKinney, C.O., Kay, F.R. & Anderson, R.A. (1973) A new all-female species of the genus *Cnemidophorus*. *Herpetologica*, **29**, 361-366.
- McMahan, C.D. & Zug, G.R. (2007) Burmese *Hemidactylus* (Reptilia, Squamata, Gekkonidae): geographic variation in the morphology of *Hemidactylus bowringii* in Myanmar and Yunnan, China. *Proceedings of the California Academy of Sciences*, **58**, 485-509.
- McNab, B.K. & Auffenberg, W. (1976) The effect of large body size on the temperature regulation of the Komodo dragon, *Varanus komodoensis*. *Comparative Biochemistry and Physiology*, **55A**, 345-350.
- McTaggart, A.L., Quinn, D.P., Parmerlee, J.S., Henderson, R.W. & Powell, R. (2011) A rapid assessment of reptilian diversity on Union Island, St. Vincent and the Grenadines. *South American Journal of Herpetology*, **6**, 59-65.
- Medica, P.A. (1967) Food habits, habitat preference, reproduction, and diurnal activity in four sympatric species of whiptail lizards (*Cnemidophorus*) in south central New Mexico. *Bulletin of the Southern California Academy of Sciences*, **66**, 251-276.
- Medina, M. & Ibarquengoytia, N.R. (2010) How do viviparous and oviparous lizards reproduce in Patagonia? A comparative study of three species of *Liolaemus*. *Journal of Arid Environments*, **74**, 1024-1032.
- Medina, M., Gutierrez, J., Scolaro, A. & Ibarquengoytia, N. (2009) Thermal responses to environmental constraints in two populations of the oviparous lizard *Liolaemus bibronii* in Patagonia, Argentina. *Journal of Thermal Biology*, **34**, 32-40.
- Meek, R. (1986) Field body temperature of the glass lizard *Ophisaurus apodus* in Yugoslavia. *Amphibia-Reptilia*, **1986**, 43-49.
- Meek, R. (1999) Thermoregulation and activity patterns in captive water dragons, *Physignathus cocincinus*, in a naturalistic environment. *Herpetological Journal*, **9**, 137-146.
- Meek, R. (2005) Null models and the thermal biology of the anguillid lizard *Anguis fragilis*; evidence for thermoregulation? *Amphibia-Reptilia*, **26**, 445-450.
- Meek, R., Jolley, E., de Silva, A., Goonewardene, S., Drake, J., Chalalochani, H.M.N., Liyanage, P.L.C.L., Abeysekera, T.S., Mayadunna, M.D.I.P.K., Somathilaka, S. A. U. S. & Chandrarathna, W.P.R. (2005) Altitudinal differences in thermoregulatory behaviour in *Calotes versicolor* in the Knuckles region, Sri Lanka. *Lyriocephalus*, **6**, 83-93.
- Meier, H. (1979) Herpologische beoachtungen auf Neokaledonien. *Salamandra*, **15**, 113-139.
- Meik, J.M., Jeo, R.M., Mendelson, J.R. & Jenks, K.E. (2002) Effects of bush encroachment on an assemblage of diurnal lizard species in central Namibia. *Biological Conservation*, **106**, 29-36.
- Melluso, L. & Morra, V. (2000) Petrogenesis of Late Cenozoic mafic alkaline rocks of the Nosy Be archipelago (northern Madagascar): relationships with the Comorean magmatism. *Journal of Volcanology and Geothermal Research*, **96**, 129-142.

- Melville, J. & Swain, R. (2000) Evolutionary relationships between morphology, performance and habitat openness in the lizard genus *Niveoscincus* (Scincidae: Lygosominae). *Biological Journal of the Linnean Society*, **70**, 667-683.
- Melville, J. & Schlute, J.A. (2001) Correlates of active body temperatures and microhabitat occupation in nine species of central Australian agamid lizards. *Austral Ecology*, **26**, 660-669.
- Melville, J., Harmon, L.J. & Losos, J.B. (2006) Intercontinental community convergence of ecology and morphology in desert lizards. *Proceedings of the Royal Society of London B: Biological Sciences*, **273**, 557-563.
- Mendez-de la Cruz, F.R., Casas-Andreu, G. & Villagran Santa Cruz, M. (1992) Variacion anual en la alimentacion y condicion fisica de *Sceloporus mucronatus* (Sauria: Iguanidae) en la Sierra del Ajusco, Distrito Federal, Mexico. *Southwestern Naturalist*, **37**, 349-355.
- Mendez-de la Cruz, F.R., Villagran-Santa Cruz, M. & Andrews, R.M. (1998) Evolution of viviparity in the lizard genus *Sceloporus*. *Herpetologica*, **54**, 521-532.
- Mendez-De la Cruz, F.R., Villagran-Santa Cruz, M., Hernandez-Gallegos, O., Manriquez-Moran, N.L. & Rodriguez-Romero, F. (1999) Reproductive cycle of the tropical night lizard *Lepidophyma pajapanensis* from Veracruz, Mexico. *Journal of Herpetology*, **33**, 336-339.
- Mendyk, R.W. (2008) Remarks on osteological deformities in a captive-bred emerald tree monitor, *Varanus prasinus*. *Biawak*, **2**, 72-79.
- Mendyk, R.W. (2011) Reproduction of varanid lizards (Reptilia: Squamata: Varanidae) at the Bronx Zoo. *Zoo Biology*, **30**, 1-16.
- Menezes, V.A. & Rocha, C.F.D. (2011) Thermal ecology of five *Cnemidophorus* species (Squamata: Teiidae) in east coast of Brazil. *Journal of Thermal Biology*, **36**, 232-238.
- Mertens, R. (1963) The geckos of the genus *Phelsuma* on Mauritius and adjacent islands. *Mauritius Institute Bulletin*, **5**, 299-305.
- Mesquita, D.O. & Colli, G.R. (2003) Geographical variation in the ecology of populations of some Brazilian species of *Cnemidophorus* (Squamata, Teiidae). *Copeia*, **2003**, 285-298.
- Mesquita, D.O., Colli, G.R., Franca, F.G.R. & Vitt, L.J. (2006) Ecology of a cerrado lizard assemblage in the Jalapao region of Brazil. *Copeia*, **2006**, 460-471.
- Michael, D.R.C., R. B. & Lindenmayer, D.B. (2011) Regrowth and revegetation in temperate Australia presents a conservation challenge for reptile fauna in agricultural landscapes. *Biological Conservation*, **144**, 407-415.
- Michael, D.R., Cunningham, R.B. & Lindenmayer, D.B. (2010) Microhabitat relationships among five lizard species associated with granite outcrops in fragmented agricultural landscapes of south-eastern Australia. *Austral Ecology*, **35**, 214-225.
- Milne, T., Bull, C.M. & Hutchinson, M.N. (2002) Characteristics of litters and juvenile dispersal in the endangered Australian skink *Tiliqua adelaidensis*. *Journal of Herpetology*, **36**, 110-112.
- Milstead, W.W. (1951) Observations on the natural history of four species of whiptail lizard, *Cnemidophorus* (Sauria, Teiidae) in Trans-Pecos Texas. *Southwestern Naturalist*, **2**, 105-121.
- Milstead, W.W. (1961) Notes on teiid lizards in southern Brazil. *Copeia*, **1961**, 493-495.
- Milton, D.A. (1987) Reproduction of two closely related skinks, *Egernia modesta* and *Egernia whitii* (Lacertilia, Scincidae) in southeast Queensland. *Australian Journal of Zoology*, **35**, 35-41.
- Milton, S.J., Richard, W. , Dean, J. & Leuteritz, T.E.J. (2004) Opportunistic and multiple

- breeding attempts in plants and vertebrates of semi-deserts with unpredictable rainfall events through the year. *Transactions of the Royal Society of South Africa*, **9**, 43-53.
- Minoli, I., Kozykariski, M.L. & Avila, L.J. (2010) Observations on parturition in two *Liolaemus* species of the archeforus group (Iguania: Squamata: Liolaemidae). *Herpetology Notes*, **3**, 333-336.
- Minton, S.A. (1966) A contribution to the herpetology of West Pakistan. *Bulletin of the American Museum of Natural History*, **134**, 31-184.
- Miralles, A., Chaparro, J.C. & Harvey, M.B. (2009) Three rare and enigmatic South American skinks. *Zootaxa*, **2012**, 47-68.
- Mitchell, F.J. (1973) Studies on the ecology of the agamid lizard *Amphibolorus maculosus*. *Transactions of the Royal Society of South Australia*, **97**, 47-76.
- Mojica, B.H., Rey, B.H., Serrano, V.H. & Ramirez-Pinilla, M.P. (2003) Annual reproductive activity of a population of *Cnemidophorus lemniscatus* (Squamata : Teiidae). *Journal of Herpetology*, **37**, 35-42.
- Moldovan, D. (2008) Husbandry and captive reproduction of *Varanus (Odatria) similis*, Mertens, 1958. *Biawak*, **2**, 89-94.
- Molina, C., Senaris, J.C. & Rivas, G. (2004) Los reptiles del Delta del Orinoco, Venezuela. *Memoria de la Fundacion La Salle de Ciencias Naturales*, **159-160**, 235-264.
- Molina-Borja, M. & Rodriguez-Dominguez, M.A. (2004) Evolution of biometric and life-history traits in lizards (*Gallotia*) from the Canary Islands. *Journal of Zoological Systematics and Evolutionary Research*, **42**, 44-53.
- Molina-Borja, M., Rodriguez-Dominguez, M.A., Gonzalez-Ortega, C. & Bohorquez-Alonso, M.L. (2010) Sexual size and shape dimorphism variation in Caesar's lizard (*Gallotia caesaris*, Lacertidae) from different habitats. *Journal of Herpetology*, **44**, 1-12.
- Monasterio, C., Salvador, A., Iraeta, P. & Diaz, J.A. (2009) The effects of thermal biology and refuge availability on the restricted distribution of an alpine lizard. *Journal of Biogeography*, **36**, 1673-1684.
- Monroe, W.H. (1980) Some tropical landforms of Puerto Rico. In: *Geological Survey professional paper*. U.S. G.P.O
- Montanucci, R.R. (1971) Ecological and distributional data on *Crotaphytus reticulatus* (Sauria: Iguanidae). *Herpetologica*, **27**, 183-197.
- Montanucci, R.R. (1976) *Crotaphytus reticulatus* Baird. *Catalogue of American Amphibians and Reptiles*, **185**, 1-2.
- Montanucci, R.R. (1989) The reproduction and growth of *Phrynosoma ditmarsii* (Sauria: Iguanidae) in captivity. *Zoo Biology*, **8**, 139-149.
- Montechiaro, L., Kaefer, I.L., Quadros, F.C. & Cechin, S. (2011) Feeding habits and reproductive biology of the glass lizard *Ophiodes* cf. *striatus* from subtropical Brazil. *North-Western Journal of Zoology*, **7**, 63-71.
- Montgomery, C.E., Reed, R.N., Shaw, H.J., Boback, S.M. & Walker, J.M. (2007) Distribution, habitat, size, and color pattern of *Cnemidophorus lemniscatus* (Sauria: Teiidae) on Cayo Cochino Pequeno, Honduras. *Southwestern Naturalist*, **52**, 38-45.
- Montgomery, C.E., Boback, S.M., Green, S.E.W., Paulissen, M.A. & Walker, J.M. (2011) *Cnemidophorus lemniscatus* (Squamata: Teiidae) on Cayo Cochino Pequeno, Honduras: extent of island occupancy, natural history, and conservation status. *Herpetological Conservation and Biology*, **6**, 10-24.
- Moores, E.M. & Fairbridge, R.W. (1997) *Encyclopedia of European and Asian Regional*

Geology. Springer.

- Moravec, J. (1998) Taxonomic and faunistic notes on the herpetofauna of Syria (Reptilia). *Faunistische Abhandlungen des Museums fur Tierkunde Dresden*, **21**, 99-106.
- Moravec, J. & Modry, D. (1994) On the occurrence of *Cyrtopodion heterocercus mardinensis* and *Pseudocerastes persicus fieldi* in Syria. *Zoology in the Middle East*, **10**, 53-56.
- Moravec, J. & Bohme, W. (1997) A new subspecies of the Mediterranean gecko, *Hemidactylus turcicus* from the Syrian lava desert (Squamata: Sauria: Gekkonidae). *Herpetozoa*, **10**, 121-128.
- Moravec, J., El, D., S. B., Seligmann, H., Sivan, N. & Werner, Y.L. (1999) Systematics and distribution of the *Acanthodactylus pardalis* group (Lacertidae) in Egypt and Israel. *Zoology in the Middle East*, **17**, 21-50.
- Morelli, M. (2000) *Carlia rostralis*. *Herpetofauna of North Queensland*,
- Mori, A. & Hikida, T. (1993) Natural history observations of the flying lizard, *Draco volans sumatranus* (Agamidae, Squamata) from Sarawak Malaysia. *Raffles Bulletin of Zoology*, **41**, 83-94.
- Mori, A. & Hikida, T. (1994) Field observations on the social behavior of the flying lizard, *Draco volans sumatranus*, in Borneo. *Copeia*, **1994**, 124-130.
- Moro, D. & MacAulay, I. (2010) *A guide to the reptiles and amphibians of Barrow Island*. Chevron Australia Pty Limited, Australia.
- Morton, S.R. & James, C.D. (1988) The diversity and abundance of lizards in arid Australia: A new hypothesis. *American Naturalist*, **132**, 237-256.
- Mount, R.H. (1961) The natural history of the red-tailed skink, *Eumeces egregius* Baird. *Dissertation*.
- Mount, R.H. (1968) *Eumeces egregius*. *Catalogue of American Amphibians and Reptiles*, **73**, 1-2.
- Mouton, P.L.F.N., Van Rensburg, D.A.J. & Van Wyk, J.H. (2010) Epidermal glands in cordylid lizards, with special reference to generation glands. *Zoological Journal of the Linnean Society*, **158**, 312-324.
- Muchlinski, A.E., Estany, A. & Don, M.T. (1995) The response of *Anolis equestris* and *Oplurus cyclurus* (Reptilia: Iguanidae) to bacterial endotoxin. *Journal of Thermal Biology*, **20**, 315-320.
- Murphy, T.J. & Myers, A.A. (1996) The behavioral ecology of *Phelsuma astriata semicarinata* on Aride Island Nature Reserve, Seychelles. *Journal of Herpetology*, **30**, 117-123.
- Murphy, K., Hudson, S. & Shea, G. (2006) Reproductive seasonality of three cold-temperate viviparous skinks from southeastern Australia. *Journal of Herpetology*, **40**, 454-464.
- Murphy, R.W., Fu, J., Macculloch, R.D.D., I. S. & Kupriyanova, L.A. (2000) A fine line between sex and unisexuality: the phylogenetic constraints on parthenogenesis in lacertid lizards. *Zoological Journal of the Linnean Society*, **130**, 527-549.
- Mushinsky, H.R. (1992) Natural history and abundance of southeastern five-lined skinks, *Eumeces inexpectatus*, on a periodically burned sandhill in Florida. *Herpetologica*, **48**, 307-312.
- Musters, C.J.M. (1983) Taxonomy of the genus *Draco* L. (Agamidae, Lacertilia, Reptilia). *Zoologische Verhandlungen*, **199**, 1-120.
- Mys, B. (1988) The zoogeography of the scincid lizards from North Papua New Guinea (Reptilia: Scincidae). I. The distribution of the species. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie*, **58**, 127-183.

- Nagy, K.A., Girard, I.A. & Brown, T.K. (1999) Energetics of free-ranging mammals, reptiles and birds. *Annual Review of Nutrition*, **19**, 247-277.
- Navarro-Garcia, J.C., Garcia, A. & Mendez de la Cruz, F.R. (2008) Seasonality, thermoregulation effectiveness of *Aspidoscelis lineatissima* (Sauria: Teiidae) and the thermal quality of a seasonally dry tropical forest in Chamela, Jalisco, Mexico. *Revista Mexicana de Biodiversidad*, **79**, 413-419.
- Naya, D.E., Veloso, C. & Bozinovic, F. (2008) Physiological flexibility in the Andean lizard *Liolaemus bellii*: seasonal changes in energy acquisition, storage and expenditure. *Journal of Comparative Physiology B*, **178**, 1007-1015.
- Neall, V.E. & Trewick, S.A. (2008) The age and origin of the Pacific islands: a geological overview. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363**, 3293-3308.
- Neang, T., Grismer, L.L., Onn, C.K., Grismer, J.L., Wood, P.L. & Youmans, T.M. (2011) First report on the herpetofauna of Dalai mountain in Phnom Samkos Wildlife Sanctuary, southwestern Cardamom Mountains, Cambodia. *Cambodian Journal of Natural History*, **2010**, 127-143.
- Neill, W.T. (1957) Notes on the pygopodid lizards, *Lialis burtoni* and *L. jicari*. *Copeia*, **1957**, 230-232.
- Nemtzov, S.C. (2008) *Uromastyx lizards in Israel*. Ndf Workshop Case Studies.
- Ngo, V.T. & Grismer, L.L. (2012) A new endemic species of *Cyrtodactylus* Gray (Squamata: Gekkonidae) from Tho Chu Island, southwestern Vietnam. *Zootaxa*, **3228**, 48-60.
- Nkosi, W.T., Heidman, J.L. & van Wyk, J.H. (2004) Reproduction and sexual size dimorphism in the lacertid lizard *Pedioplanis burchelli* (Sauria: Lacertidae) in South Africa. *Journal of Herpetology*, **38**, 473-480.
- Noble, G.K. & Klingel, G.C. (1932) The reptiles of Great Inagua Island, British West Indies. *American Museum Novitates*, **549**, 1-25.
- Noble, T., Bunbury, N., Kaiser-Bunbury, C.N. & Bell, D.J. (2011) Ecology and co-existence of two endemic day gecko (*Phelsuma*) species in Seychelles native palm forest. *Journal of Zoology*, **283**, 73-80.
- Nogueira, C., Valdujo, P.H. & Franca, F.G.R. (2005) Habitat variation and lizard diversity in a Cerrado area of Central Brazil. *Studies on Neotropical Fauna and Environment*, **40**, 105-112.
- Norrie, S. & Langerwerf, B. (1987) *Lacerta schreiberi* in Spain, Portugal, and in captivity. In: J. Coote, ed., Reptiles. Proceedings of the 1986, U.K. pp. 97, Herpetological Societies Symposium on Captive Breeding. British Herpetological Society, London.
- Norval, G., Mao, J.-J. & Slater, K. (2011) Notes on the reproduction of the Swinhoe's tree lizard, *Japalura swinhonis* Gunther, 1864, (Squamata: Agamidae) from southwestern Taiwan. *Herpetology Notes*, **4**, 19-324.
- Nussbaum, R.A. & Raxworthy, C.J. (1998) A new species of *Mabuya* Fitzinger (Reptilia: Squamata: Scincidae) from the high Plateau (Isalo National Park) of south-central Madagascar. *Herpetologica*, **54**, 336-343.
- Nyandwi, N. & Kangwe, J. (2009) Ecological and engineering importance of the Bet el Ras beach-sandstone. *Tanzania Journal of Science*, **32**, 117-123.
- Nyhagen, D.F., Kragelund, C., Olesen, J.M. & Jones, C.G. (2001) Insular interactions between lizards and flowers: flower visitation by an endemic Mauritian gecko. *Journal of Tropical Ecology*, **17**, 755-761.

- Olesen, J.M. & Valido, A. (2003) Lizards as pollinators and seed dispersers: an island phenomenon. *Trends in Ecology and Evolution*, **18**, 177-181.
- Oliver, J.A. (1948) The anoline lizards of Bimini, Bahamas. *American Museum Novitates*, **1383**, 1-36.
- Oliveros, C.H., Ota, H., Crombie, R.I. & Brown, R.M. (2011) The Herpetofauna of the Babuyan group of islands, northern Philippines. *Scientific Publications of the Natural History Museum of the University of Kansas*, **43**, 1-20.
- Olmsted, F.H. (1958) *Geologic reconnaissance of San Clemente Island, California*. US Government Printing Office.
- Olsson, M. (1995) Forced copulation and costly female resistance behavior in the Lake Eyre dragon, *Ctenophorus maculosus*. *Herpetologica*, **51**, 19-24.
- Onn, C.K. & Ahmad, N. (2010) A new insular species of *Cyrtodactylus* (Squamata: Gekkonidae) from northeastern Peninsular Malaysia, Malaysia. *Zootaxa*, **2389**, 47-56.
- Onn, C.K., Grismer, L.L., Sharma, D.S., Belabut, D. & Ahmed, N. (2009) New herpetofaunal records for Perlis State Park and adjacent areas. *Malayan Nature Journal*, **61**, 255-262.
- Ord, T.J. & Blumstein, D.T. (2002) Size constraints and the evolution of display complexity: why do large lizards have simple displays? *Biological Journal of the Linnean Society*, **76**, 145-161.
- Ortega-Leon, A.M., Smith, E.R., Zuniga-Vega, J. & Mendez-de la Cruz, F. (2007) Growth and population demography of one population of the lizard *Sceloporus mucronatus mucronatus*. *Western North American Naturalist*, **67**, 492-502.
- Ortega-Rubio, A., Gonzalez-Romero, A. & Barbault, R. (1995) Food analysis and resource partitioning. in a lizard guild of the Sonoran Desert, Mexico. *Journal of Arid Environments*, **29**, 367-382.
- Ortiz-Zapata, J.C. (1980) Revision taxonomica del genero *Tropidurus* en Chile. *Reunion Iberoamericana de Conservacion y Zoologia de Vertebrados*, **1**, 355-377.
- Ota, H. (1989) A new species of *Japalura* (Agamidae: Lacertilia: Reptilia) from Taiwan. *Copeia*, **1989**, 569-576.
- Ota, H. (1991) Taxonomic redefinition of *Japalura swinhonis* Gunther (Agamidae: Squamata), with a description of a new subspecies of *J. polygonata* from Taiwan. *Herpetologica*, **47**, 280-294.
- Ota, H., Chen, S.-L. & Shang, G. (1998) *Japalura luei*: a new agamid lizard from Taiwan (Reptilia: Squamata). *Copeia*, **1998**, 649-656.
- Ouboter, P.E. (1986) A revision of the genus *Scincella* (Reptilia: Sauria: Scincidae) of Asia, with some notes on its evolution. *Zoologische Verhandelingen*, **229**, 1-66.
- Paden, L. (2008) *Varanus panoptes horni* (Argus Monitor) reproduction. *Biawak*, **2**, 95-96.
- Pafilis, P., Perez-Mellado, V. & Valakos, E. (2008) Postautotomy tail activity in the Balearic lizard, *Podarcis lilfordi*. *Naturwissenschaften*, **95**, 217-221.
- Pafilis, P., Foufopoulos, J., Poulakakis, N., Lymberakis, P. & Valakos, E. (2009) Tail shedding in island lizards [Lacertidae, Reptilia]: decline of antipredator defences in relaxed predation environments. *Evolution*, **63**, 1262-1278.
- Pafilis, P., Foufopoulos, J., Sagonas, K., Runemark, A., Svensson, E. & Valakos, E.D. (2011) Reproductive biology of insular reptiles: marine subsidies modulate expression of the "Island Syndrome". *Copeia*, **2011**, 545-552.
- Pal, A., Swain, M.M. & Rath, S. (2010) Observations on microhabitat use and activity patterns in *Sitana ponticeriana* (Sauria: Agamidae). *Russian Journal of Herpetology*, **17**, 22-30.

- Pal, A., Swain, M.M. & Rath, S. (2011) Reproduction and sexual dichromatism in *Sitana ponticeriana* (Reptilia: Draconinae: Agamidae). *Taprobanica*, **3**, 31-37.
- Pal, T., Chakraborty, P.P., Gupta, T.D. & Singh, C.D. (2003) Geodynamic evolution of the outer-arc–forearc belt in the Andaman Islands, the central part of the Burma–Java subduction complex. *Geological Magazine*, **140**, 289-307.
- Pandav, B.N., Shanbhag, B.A. & Saidapur, S.K. (2010) Growth patterns and reproductive strategies in the lizard, *Calotes versicolor* raised in captivity. *Acta Herpetologica*, **5**, 131-142.
- Panov, E.N. & Zykova, L.Y. (1997) Differentiation and interrelations of two representatives of *Laudakia stellio* complex (Reptilia: Agamidae) in Israel. *Russian Journal of Herpetology*, **4**, 102-114.
- Papenfuss, T.J. (1969) Preliminary analysis of the reptiles of arid central West Africa. *The Wasmann Journal of Biology*, **27**, 249-325.
- Papenfuss, T.J. (1982) The ecology and systematics of the amphisbaenian genus *Bipes*. *Occasional Papers of the California Academy of Sciences*, **136**, 1-42.
- Parker, H.W. (1932) The status of two Peruvian lizards. *Copeia*, **1932**, 178.
- Parker, H.W. (1936) Dr. Karl Jordan's expedition to South West Africa and Angola: herpetological collections. *Novitates Zoologicae*, **40**, 115-146.
- Parker, H.W. (1942) The lizards of British Somaliland. *Bulletin of the Museum of Comparative Zoology*, **91**, 1-101.
- Parker, W.S. (1971) Ecological observations on the regal horned lizard (*Phrynosoma solare*) in Arizona. *Herpetologica*, **27**, 333-338.
- Parker, W.S. (1972) Aspects of the ecology of a Sonoran Desert population of the western banded gecko, *Coleonyx variegatus* (Sauria, Eublepharinae). *American Midland Naturalist*, **88**, 209-224.
- Parrish, G.R. & Gill, B.J. (2003) Natural history of the lizards of the Three Kings Islands, New Zealand. *New Zealand Journal of Zoology*, **30**, 205-220.
- Patchell, F.C. & Shine, R. (1986) Food habits and reproductive biology of the Australian legless lizards (Pygopodidae). *Copeia*, **1986**, 30-39.
- Patterson, J.W. (1990) Female reproductive cycles in two subspecies of the tropical lizard *Mabuya striata*. *Oecologia*, **84**, 232-237.
- Patterson, J.W. (1991) Emergence, basking behaviour, mean selected temperature and critical thermal minimum in high and low altitude subspecies of the tropical lizard *Mabuya striata*. *African Journal of Ecology*, **29**, 330-339.
- Patterson, G.B. & Daugherty, C.H. (1990) Four new species and one new subspecies of skinks, genus *Leiolopisma* (Reptilia: Lacertilia: Scincidae) from New Zealand. *Journal of the Royal Society of New Zealand*, **20**, 65-84.
- Patterson, G.B. & Daugherty, C.H. (1994) *Leiolopisma stenotis*, n. sp.; (Reptilia: Lacertilia: Scincidae) from Stewart Island. *Journal of the Royal Society of New Zealand*, **24**, 125-132.
- Paulissen, M.A. (1994) Microhabitat use and escape behavior of syntopic clonal complexes of the parthenogenetic whiptail lizard *Cnemidophorus laredoensis*. *American Midland Naturalist*, **132**, 10-18.
- Paulissen, M.A. & Walker, J.M. (1994) Diet of the insular whiptail lizard *Cnemidophorus nigricolor* (Teiidae) from Grand Rocques Island, Venezuela. *Journal of Herpetology*, **28**, 524-526.
- Paulissen, M.A., Cordes, J.E. & Walker, J.M. (1989) Notes on the thermal biology of the

- Laredo whiptail *Cnemidophorus laredoensis* (Teiidae). *Texas Journal of Science*, **41**, 224-228.
- Pauwels, O.S.G., David, P., Chimsunchart, C. & Thirakhupt, K. (2003) Reptiles of Phetchaburi Province, Western Thailand: a list of species, with natural history notes, and a discussion on the biogeography at the Isthmus of Kra. *The Natural History Journal of Chulalongkorn University*, **3**, 23-53.
- Pauwels, O.S.G., Bauer, A.M., Sumontha, M. & Chanhom, L. (2004) *Cyrtodactylus thirakhupti* (Squamata: Gekkonidae), a new cave-dwelling gecko from southern Thailand. *Zootaxa*, **72**, 1-11.
- Pauwels, O.S.G., Chan-Ard, T., Wanchai, P., Kheowyoo, B. & Bauer, A.M. (2009) Reptile diversity of Phetchaburi Province, western Thailand: an update. *Hamadryad*, **34**, 9-21.
- Pellegrino, K.C.M., Rodrigues, M.T., Waite, A.N., Morando, M., Yassuda, Y.Y. & Sites, J.W. (2005) Phylogeography and species limits in the *Gymnodactylus darwinii* complex (Gekkonidae, Squamata): genetic structure coincides with river systems in the Brazilian Atlantic Forest. *Biological Journal of the Linnean Society*, **85**, 13-26.
- Pengilley, R. (1981) Notes on the biology of *Varanus spenceri* and *V. gouldii*, Barkly Tablelands, Northern Territory. *Australian Journal of Herpetology*, **1**, 23-26.
- Perez, E. & de La Riva, R.L.S. (2008) Morphological revision of lizards of the formosus group, genus *Sceloporus* (Squamata: Sauria) of southern Mexico, with description of a new species. *Bulletin of the Maryland Herpetological Society*, **44**, 77-97.
- Perez-Buitrago, N., Sabat, A.M. & McMillan, W.O. (2010) Spatial ecology of the endangered Mona Island Iguana *Cyclura cornuta stejnegeri*: Does territorial behavior regulate density? *Herpetological Monographs*, **24**, 86-110.
- Perez-Mellado, V. (1981) La lagartija de bocage, *Podarcis bocagei* (Seoane, 1884). Primeros datos sobre su distribución, colorido y ecología. *Amphibia-Reptilia*, **1**, 253-268.
- Pernetta, A.P., Bell, D.J. & Jones, C.G. (2005) Macro- and microhabitat use of Telfair's skink (*Leiolopisma telfairii*) on Round Island, Mauritius: implications for their translocation. *Acta Oecologica*, **28**, 313-323.
- Perry, G. (1996) The evolution of sexual dimorphism in the lizard *Anolis polylepis* (Iguania): evidence from intraspecific variation in foraging behavior and diet. *Canadian Journal of Zoology*, **74**, 1238-1245.
- Perry, G. & Garland, T. (2002) Lizard home ranges revisited: effects of sex, body size, diet, habitat, and phylogeny. *Ecology*, **83**, 1870-1885.
- Perry, G., Lampl, I., Lerner, A., Rothenstein, D., Shani, E., Sivan, N. & Werner, Y.L. (1990) Foraging mode in lacertid lizards: variations and correlates. *Amphibia-Reptilia*, **11**, 373-384.
- Philipp, K.M., Ziegler, T. & Böhme, W. (2007) Preliminary investigations of the natural diet of six monitor lizard species of the *Varanus (Euprepiosaurus) indicus* group. *Mertensiella*, **16**, 1-447.
- Phillips, J.A. (1995) Movement patterns and density of *Varanus albigularis*. *Journal of Herpetology*, **29**, 407-416.
- Phillips, J.A. & Millar, R.P. (1998) Reproductive biology of the white-throated savanna monitor, *Varanus albigularis*. *Journal of Herpetology*, **32**, 366-377.
- Pianka, E.R. (1971) Comparative ecology of two lizards. *Copeia*, **1971**, 129-138.
- Pianka, E.R. (1982) Observations on the ecology of *Varanus* in the Great Victoria desert. *Western Australian Naturalist*, **15**, 37-44.

- Pianka, E.R. (1994) Comparative ecology of *Varanus* in the Great Victoria desert. *Australian Journal of Ecology*, **19**, 395-408.
- Pianka, E.R. (1995) Evolution of body size: varanid lizards as a model system. *American Naturalist*, **146**, 298-414.
- Pianka, E.R. (2011) Notes on the ecology of some uncommon skinks in the Great Victoria Desert. *Western Australian Naturalist*, **28**, 50-60.
- Pianka, E.R. & Parker, W.S. (1975) Ecology of horned lizards: a review with special reference to *Phrynosoma platyrhinos*. *Copeia*, **1975**, 141-162.
- Pianka, E.R. & Huey, R.B. (1978) Comparative ecology, resource utilization and niche segregation among gekkonid lizards the southern Kalahari. *Copeia*, **1978**, 691-701.
- Pianka, E.R. & Giles, W.F. (1982) Notes on the biology of two species of nocturnal skinks, *Egernia inornata* and *Egernia striata*, in the Great Victoria desert. *Western Australian Naturalist*, **15**, 44-49.
- Piantoni, C., Ibarguengoytia, N.R. & Cussac, V.E. (2006) Age and growth of the Patagonian lizard *Phymaturus patagonicus*. *Amphibia-Reptilia*, **27**, 385-392.
- Pincheira-Donoso, D. & Scolaro, A. (2007) Iguanian species-richness in the Andes of boreal Patagonia: evidence for an additional new *Liolaemus* lizard from Argentina lacking precloacal glands (Iguania, Liolaeminae). *Zootaxa*, **1452**, 55-68.
- Pincheira-Donoso, D. & Tregenza, T. (2011) Fecundity selection and the evolution of reproductive output and sex-specific body size in the *Liolaemus* lizard adaptive radiation. *Evolutionary Biology*, **38**, 197-207.
- Pincheira-Donoso, D., Hodgson, D.J. & Tregenza, T. (2008) The evolution of body size under environmental gradients in ectotherms: why should Bergmann's rule apply to lizards? *BMC Evolutionary Biology*, **8**, 68.
- Pincheira-Donoso, D., Hodgson, D.J., Stipala, J. & Tregenza, T. (2009) A phylogenetic analysis of sex-specific evolution of ecological morphology in *Liolaemus* lizards. *Ecological Research*, **24**, 1223-1231.
- Pinilla, M.P.R. (1991) Reproductive and fat body cycles of the viviparous lizard *Liolaemus huacahuasicus*. *Journal of Herpetology*, **25**, 205-208.
- Pizzatto, L. (2005) Reproductive biology of the glass snake *Ophiodes fragilis* (Squamata, Anguillidae) in south-east Brazil. *Herpetological Journal*, **15**, 9-13.
- Pope, C.H. (1929) Notes on reptiles from Fukien and other Chinese provinces. *Bulletin of the American Museum of Natural History*, **58**, 335-487.
- Post, M.J. (2000) The captive husbandry and reproduction of the Hosmer's Skink *Egernia hosmeri*. *Herpetofauna*, **30**, 2-6.
- Pounds, J.A. (1988) Ecomorphology, locomotion, and microhabitat structure: patterns in a tropical mainland *Anolis* community. *Ecological Monographs*, **58**, 299-320.
- Powell, R. (1999) *Anolis longiceps*. *Catalogue of American Amphibians and Reptiles*, **693**, 1-2.
- Powell, M.A., Powell, R. & Henderson, R.W. (2005) *Anolis sabanus*. *Catalogue of American Amphibians and Reptiles*, **815**, 1-5.
- Pregill, G. (1984) Durophagous feeding adaptations in an amphisbaenid. *Journal of Herpetology*, **18**, 186-191.
- Price, A.H. (1986) *Cnemidophorus tessellatus*. *Catalogue of American Amphibians and Reptiles*, **398**, 1-2.
- Price, A.H. (1990) *Phrynosoma cornutum*. *Catalogue of American Amphibians and Reptiles*, **469**, 1-7.

- Punzo, F. (2001) Studies on the natural history and ecology of sympatric whiptail lizards (*Cnemidophorus marmoratus* and *C. tessellatus*) from Madera Canyon (Brewster County, Texas). *Texas Journal of Science*, **53**, 43-54.
- Punzo, F. (2007) Life history, demography, diet and habitat associations in the southwestern earless lizard, *Cophosaurus texanus scitulus* from northern and southern limits of its geographical range. *Amphibia-Reptilia*, **28**, 65-76.
- Qualls, F.J. & Shine, R. (1998) Geographic variation in lizard phenotypes: importance of the incubation environment. *Biological Journal of the Linnean Society*, **64**, 477-491.
- Qualls, C.P., Shine, R.D., S. & Hutchinson, M. (1994) The evolution of viviparity within the Australian scincid lizard *Lerista bougainvillii*. *Journal of Zoology*, **237**, 13-26.
- Radder, R.S. & Shanbhag, B.A. (2003) Interrelationships among reproductive traits of female lizard, *Sitana ponticeriana* (Cuvier). *Current Science*, **85**, 89-91.
- Radder, R. & Shine, R. (2007) Germinal bed condition in a polyautochronic single-clutched lizard, *Bassiana duperreyi* (Scincidae). *Amphibia-Reptilia*, **28**, 159-162.
- Radder, R.S., Pizzatto, L. & Shine, R. (2008) Morphological correlates of life-history variation: is lizard clutch size related to the number of germinal beds in the ovary? *Biological Journal of the Linnean Society*, **94**, 81-88.
- Radtkey, R.R., Fallon, S.M. & Case, T.J. (1997) Character displacement in some *Cnemidophorus* lizards revisited: a phylogenetic analysis. *Proceedings of the National Academy of Sciences, USA*, **94**, 9740-9745.
- Ramírez-Pinilla, M.P., Serrano, V.H. & Galeano, J.C. (2002) Annual reproductive activity of *Mabuya mabouya* (Squamata, Scincidae). *Journal of Herpetology*, **36**, 667-677.
- Ramalho, R.A.S. & Ramalho, R.A.d.S. (2011) *Tracers of Uplift and Subsidence in the Cape Verde Archipelago*. Building the Cape Verde Islands. pp. 79-138. Springer Berlin Heidelberg.
- Ramirez-Bautista, A. & Vitt, L.J. (1997) Reproduction in the lizard *Anolis nebulosus* (Polychrotidae) from the Pacific coast of Mexico. *Herpetologica*, **53**, 423-431.
- Ramirez-Bautista, A. & Vitt, L.J. (1998) Reproductive biology of *Urosaurus bicarinatus* (Sauria : Phrynosomatidae) from a tropical dry forest of Mexico. *Southwestern Naturalist*, **43**, 381-390.
- Ramirez-Bautista, A. & Gonzalez-Romero, A. (2002) Some reproductive and feeding characteristics of the viviparous Mexican lizard *Sceloporus torquatus* (Phrynosomatidae). *Southwestern Naturalist*, **47**, 98-102.
- Ramirez-Bautista, A. & Gutierrez-Mayen, G. (2003) Reproductive ecology of *Sceloporus utiformis* (Sauria : Phrynosomatidae) from a tropical dry forest of Mexico. *Journal of Herpetology*, **37**, 1-10.
- Ramirez-Bautista, A. & Olivera-Becerril, V. (2004) Reproduction in the boulder spiny lizard, *Sceloporus pyrocephalus* (Sauria: Phrynosomatidae), from a tropical dry forest of Mexico. *Journal of Herpetology*, **38**, 225-231.
- Ramirez-Bautista, A. & Davila-Ulloa, E.G. (2009) Reproductive characteristics of a population of *Sceloporus dugesii* (Squamata: Phrynosomatidae) from Michoacan, Mexico. *Southwestern Naturalist*, **54**, 400-408.
- Ramirez-Bautista, A., Barba-Torres, J. & Vitt, L.J. (1998) Reproductive cycle and brood size of *Eumeces lynxe* from Pinal de Amoles, Queretero, Mexico. *Journal of Herpetology*, **32**, 18-24.
- Ramirez-Bautista, A., Balderas-Valdivia, C. & Vitt, L.J. (2000) Reproductive ecology of the

- whiptail lizard *Cnemidophorus lineatissimus* (Squamata : Teiidae) in a tropical dry forest. *Copeia*, **2000**, 712-722.
- Ramirez-Bautista, A., Jimenez-Cruz, E. & Marshall, J.C. (2004) Comparative life history for populations of the *Sceloporus grammicus* complex (Squamata : Phrynosomatidae). *Western North American Naturalist*, **64**, 175-183.
- Ramirez-Bautista, A., Smith, G.R. & Hernandez-Ibarra, X. (2009) Reproduction and sexual dimorphism in the whiptail lizard *Aspidoscelis gularis* (Squamata: Teiidae) in Guadalupe, San Luis Potosi, Mexico. *Southwestern Naturalist*, **54**, 453-460.
- Ramirez-Bautista, A., Ortiz-Cruz, A.L., Arizmendi, M.D. & Campos, J. (2005) Reproductive characteristics of two syntopic lizard species, *Sceloporus gadoviae* and *Sceloporus jalapae* (Squamata : Phrynosomatidae), from Tehuacan Valley, Puebla, Mexico. *Western North American Naturalist*, **65**, 202-209.
- Ramirez-Bautista, A., Vitt, L.J., Ramirez-Hernandez, A., Quijano, F.M. & Smith, G.R. (2008) Reproduction and sexual dimorphism of *Lepidophyma sylvaticum* (Squamata: Xantusiidae), a tropical night lizard from Tlanchinol, Hidalgo, Mexico. *Amphibia-Reptilia*, **29**, 207-216.
- Ramirez-Pinilla, M.P., Calderon-Espinosa, M.L., Flores-Villela, O., Munoz-Alonso, A. & Mendez de la Cruz, F.R. (2009) Reproductive activity of three sympatric viviparous lizards at Omiltemi, Guerrero, Sierra Madre del Sur, Mexico. *Journal of Herpetology*, **43**, 409-420.
- Ramirez-Sandoval, E., Ramirez-Bautista, A. & Vitt, L.J. (2006) Reproduction in the lizard *Phyllodactylus lanei* (Squamata: Gekkonidae) from the Pacific coast of Mexico. *Copeia*, **2006**, 1-9.
- Rand, A.S. (1957) Notes on amphibians and reptiles from El Salvador. *Fieldiana Zoology*, **34**, 505-534.
- Rand, A.S. (1967) Ecology and social organization in the iguanid lizard *Anolis lineatopus*. *Proceedings of the U.S. National Museum*, **122**, 1-79.
- Rand, A.S. (1968) The ecological distribution of anoline lizards around Kingston, Jamaica. *Breviora*, **272**, 1-18.
- Rand, A.S. (1982) Clutch and egg size in Brazilian iguanid lizards. *Herpetologica*, **38**, 171-178.
- Rand, A.S., Gorman, G.C. & Rand, W.M. (1975) Natural history, behavior, and ecology of *Anolis agassizi*. *Smithsonian Contributions to Zoology*, **176**, 27-38.
- Rastegar-Pouyani, N., Heydari, N. & Faizi, H. (2011) Analysis of geographic variation in four populations of *Acanthodactylus blanfordi* Boulenger, 1918 (Sauria: Lacertidae) from southern and southeastern Iran. *Russian Journal of Herpetology*, **18**, 228-234.
- Rathnayake, N.D., Herath, N.D., Hewamathes, K.K. & Jayalath, S. (2003) The thermal behavior, diurnal activity pattern and body temperature of *Varanus salvator* in central Sri Lanka. *Hamadryad*, **27**, 179-184.
- Rauhala, M. (1993) Distribution and habitat of the nobby dragon *Amphibolurus nobbi* in the Australian Capital Territory. *ACT Parks and Conservation Service, Technical report*, **5**, 1-46.
- Raxworthy, C.J. (1991) Field observations on some dwarf chameleons (*Brookesia* spp.) from rain-forest areas of Madagascar, with the description of a new species. *Journal of Zoology*, **224**, 11-25.
- Raxworthy, C.J. & Nussbaum, R.A. (1994) A partial systematic revision of the day geckos, *Phelsuma* Gray, of Madagascar (Reptilia, Squamata, Gekkonidae). *Zoological Journal of the Linnean Society*, **112**, 321-335.
- Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-

- Rosa, D., Butikofer, L., Suleiman, A.S., Al-Aseily, B.A., Carugati, C., Boncompagni, E. & Fasola, M. (2011) Annotated checklist and distribution of the Socotran Archipelago Herpetofauna (Reptilia). *Zootaxa*, **2826**, 1-44.
- Read, J.L. (1999) Longevity, reproductive effort and movements of three sympatric Australian arid-zone geckos. *Australian Journal of Zoology*, **47**, 307-316.
- Reaney, L.T. & Whiting, M.J. (2002) Life on a limb: ecology of the tree agama (*Acanthocercus a. atricollis*) in southern Africa. *Journal of Zoology*, **257**, 439-448.
- Reed, C.A. & Marx, H. (1959) A herpetological collection from northeastern Iraq. *Transactions of the Kansas Academy of Sciences*, **62**, 91-122.
- Regalado, R. (2006) Reproduction and growth of seven species of dwarf geckos, *Sphaerodactylus* (Gekkonidae), in captivity. *Herpetological Review*, **37**, 13-20.
- Reyes, M.A.P., Bennett, D. & Oliveros, C. (2008) The monitor lizards of Camiguin Island, Northern Philippines. *Biawak*, **2**, 28-36.
- Reynolds, R.G. (2011) Islands, metapopulations, and archipelagos: genetic equilibrium and non-equilibrium dynamics of structured populations in the context of conservation. *Dissertation*.
- Rezende-Pinto, F.M., Verrastro, L., Zanotelli, J.C. & Barata, P.C.R. (2009) Reproductive biology and sexual dimorphism in *Cnemidophorus vacariensis* (Sauria, Teiidae) in the grasslands of the Araucaria Plateau, southern Brazil. *Iheringia, Serie Zoologia*, **99**, 82-91.
- Rifai, L., Modry, D., Necas, P. & Amr, Z.S. (2003) The occurrence of *Acanthodactylus hardyi* Haas, 1957 in the Hashemite Kingdom of Jordan and notes on its ecology. *Zoology in the Middle East*, **28**, 33-38.
- Roca, V. (1999) Relacion entre las faunas endoparasitas de reptiles y su tipo de alimentacion. *Revista Espanola de Herpetologia*, **13**, 101-121.
- Rocha, C.F.D. (2000) Selectivity in plant food consumption in the lizard *Liolaemus lutzae* from southeastern Brazil. *Studies on Neotropical Fauna and Environment*, **35**, 14-18.
- Rocha, P.L.B. & Rodrigues, M.T. (2005) Electivities and resource use by an assemblage of lizards endemic to the dunes of the Sao Francisco River, northeastern Brazil. *Papeis Avulsos de Zoologia*, **45**, 261-284.
- Rocha, C.F.D., Vrcibradic, D. & Van Sluys, M. (2004) Diet of the lizard *Mabuya agilis* (Sauria; Scincidae) in an insular habitat (Ilha Grande, Rj, Brazil). *Brazilian Journal of Biology*, **64**, 135-139.
- Rocha, S., Carretero, M.A. & Harris, D.J. (2010) Genetic diversity and phylogenetic relationships of *Mabuya* spp. (Squamata: Scincidae) from western Indian Ocean islands. *Amphibia-Reptilia*, **31**, 375-385.
- Rocha, C.F.D., Vrcibradic, D., Teixeira, R.L. & Cuzzuol, M.G.T. (2002) Interpopulational variation in litter size of the skink *Mabuya agilis* in southeastern Brazil. *Copeia*, **2002**, 857-864.
- Rocha, C.F.D., Van Sluys, M., Vrcibradic, D., Kiefer, M.C., Menezes, V.A. & Siqueira, C.C. (2009) Comportamento de termoregulacao em lagartos brasileiros. *Oecologia Brasiliensis*, **13**, 115-131.
- Rodda, G.H. & Dean-Bradley, K. (2001) *Inventory of the reptiles of the War in the Pacific National Historical Park, Guam*. Pacific Cooperative Studies Unit Technical Report 169. University of Hawaii at Manoa, Department of Botany. Honolulu, HI.
- Rodda, G.H., Fritts, T.H. & Reichel, J.D. (1991) The distributional patterns of reptiles and amphibians in the Mariana Islands. *Micronesica*, **24**, 195-210.

- Rodda, G.H., Perry, G., Rondeau, R.J. & Lazell, J. (2001) The densest terrestrial vertebrate. *Journal of Tropical Ecology*, **17**, 331-338.
- Rodrigues, M.T. (1996) Lizards, snakes, and amphisbaenians from the Quaternary sand dunes of the middle Rio Sao Francisco, Bahia, Brazil. *Journal of Herpetology*, **30**, 513-523.
- Rodrigues, M.T., Freitas, M.A., Silva, T.F.S. & Bertolotto, C.E.V. (2006) A new species of lizard genus *Enyalius* (Squamata, Leiosauridae) from the highlands of Chapada Diamantina, state of Bahia, Brazil, with a key to species. *Phyllomedusa*, **5**, 11-24.
- Rodrigues-Serrano, E., Navas, C.A. & Bozinovic, F. (2009) The comparative field body temperature among *Liolaemus* lizards: testing the static and the labile hypothesis. *Journal of Thermal Biology*, **34**, 306-309.
- Rodriguez, M.A.G. & Casas-Andreu, G. (2011) Facultative specialization in the diet of the twelve-lined whiptail, *Aspidoscelis lineatissima*. *Journal of Herpetology*, **45**, 287-290.
- Rodriguez-Dominguez, M.A. & Molina-Borja, M. (1998) Reproduction of the endangered Hierro Giant Lizard *Gallotia simonyi machadoi*. *Journal of Herpetology*, **32**, 498-504.
- Rodriguez-Ramirez, J. & Lewis, A.R. (1991) Reproduction in the Puerto Rican teiids *Ameiva exsul* and *A. wetmorei*. *Herpetologica*, **47**, 395-403.
- Roe, J.H., Hopkins, W.A. & Talent, L.G. (2005) Effects of body mass, feeding, and circadian cycles on metabolism in the lizard *Sceloporus occidentalis*. *Journal of Herpetology*, **39**, 595-603.
- Rogovin, K.A. & Semenov, D., . V. (2004) The toad-headed agama *Phrynocephalus guttatus* (Reptilia: Agamidae): the correlates of survival in a translocated population in the Southwestern Kalmyk Republic. *Russian Journal of Ecology*, **35**, 389-394.
- Rogovin, K.A., Semenov, D.V. & Shenbrot, G.I. (2000) Lizards of the northern Mongolian deserts: densities and community structure. *Asiatic Herpetological Research*, **9**, 1-9.
- Rohr, D.H. (1997) Demographic and life-history variation in two proximate populations of a viviparous skink separated by a steep altitudinal gradient. *Journal of Animal Ecology*, **66**, 567-578.
- Roitberg, E.S. (1999) Phenetic relationships between *Lacerta caucasica*, *L. daghestanica* and *L. praticola* (Reptilia, Lacertidae): a multivariate trend in external morphology. *Natura Croatica*, **8**, 189-200.
- Rojas-Runjaic, F.J.M., Infante-Rivero, E.E., Cabello, P. & Velozo, P. (2010) A new non-sexually dichromatic species of the genus *Gonatodes* (Sauria: Sphaerodactylidae) from Sierra de Perija, Venezuela. *Zootaxa*, **2671**, 1-16.
- Rorabaugh, J.C. (2008) An introduction to the herpetofauna of mainland Sonora, Mexico, with comments on conservation and management. *Journal of the Arizona-Nevada Academy of Science*, **40**, 20-65.
- Rosler, H. & Wranik, W. (1999) Beitrage zur herpetologie der republik Jemen. 5. Drei neue gecko-arten vom Sokotra-Archipel (Reptilia: Sauria: Gekkonidae). *Zoologische Abhandlungen, Staatliches Museum for Tierkunde, Dresden*, **50**, 249-265.
- Rosler, H. & Wranik, W. (2004) A key and annotated checklist to the reptiles of the Socotra archipelago. *Fauna of Arabia*, **20**, 505-534.
- Rosler, H., Siler, C.D., Brown, R.M.D., A. D. & Gaulke, M. (2006) *Gekko ernstkelleri* sp. n. – a new gekkonid lizard from Panay Island, Philippines. *Salamandra*, **42**, 197-211.
- Ross, T.N., Parnetta, A.P., Jones, C.G. & Bell, D.J. (2008) Sexual size dimorphism and microhabitat use of the orange-tail skink (*Gongylomorphus* spp.) on Flat Island, Mauritius: conservation implications. *Amphibia-Reptilia*, **29**, 349-359.

- Rouag, R., Djilali, H., Gueraiche, H. & Luiselli, L. (2007) Resource partitioning patterns between two sympatric lizard species from Algeria. *Journal of Arid Environments*, **69**, 158-168.
- Routman, E.J. & Hulse, A.C. (1984) Ecology and reproduction of a parthenogenetic lizard, *Cnemidophorus sonorae*. *Journal of Herpetology*, **18**, 381-386.
- Rubio, J.L. & Carrascal, L.M. (1994) Habitat selection and conservation of an endemic spanish lizard *Algyroides marchi* (Reptilia, Lacertidae). *Biological Conservation*, **70**, 245-250.
- Rugiero, L., Luiselli, L., Eniang, E.A. & Akani, G.C. (2007) Diet of a guild of geckos in a fragmented, human-altered African rainforest. *African Journal of Herpetology*, **56**, 91-96.
- Ruibal, R. (1961) Thermal relations of five species of tropical lizards. *Evolution*, **15**, 98-111.
- Ruibal, R. & Philibosian, R. (1974) The population ecology of the lizard *Anolis acutus*. *Ecology*, **55**, 525-537.
- Rummery, C.S., R., Houston, D.L. & Thompson, M.B. (1995) Thermal biology of the Australian forest dragon, *Hypsilurus spinipes* (Agamidae). *Copeia*, **1995**, 818-827.
- Russell, A.P. & Bauer, A.M. (1991) *Anolis garmani*. *Catalogue of American Amphibians and Reptiles*, **513**, 1-3.
- Russell, A.P. & Bauer, A.M. (1991) *Anolis opalinus*. *Catalogue of American Amphibians and Reptiles*, **515**, 1-2.
- Rutherford, P.L. (2004) Proximate mechanisms that contribute to female biased sexual size dimorphism in an anguid lizard. *Canadian Journal of Zoology*, **82**, 817-822.
- Sabath, M.D. (1981) Gekkonid lizards of Guam, Mariana Islands: reproduction and habitat preference. *Journal of Herpetology*, **15**, 71-75.
- Sadlier, R.A. (1990) A new species of scincid lizard from western Arnhem Land, Northern Territory. *The Beagle*, **7**, 29-33.
- Sadlier, R.A., Bauer, A.M. & Smith, S.A. (2006) A new species of *Nannoscincus* Gunther (Squamata: Scincidae) from high elevation forest in southern New Caledonia. *Records of the Australian Museum*, **58**, 29-36.
- Sadlier, R.A., Bauer, A.M., Whitaker, A.H. & Smith, S.A. (2004) Two new species of scincid lizards (Squamata) from the Massif de Kopeto, New Caledonia. *Proceedings of the California Academy of Sciences*, **55**, 208-221.
- Sadlier, R.A., Smith, S.A., Bauer, A.M. & Whitaker, A.H. (2009) Three new species of skink in the genus *Marmorosphax* Sadlier (Squamata: Scincidae) from New Caledonia. *Memoires du Museum national d'Histoire naturelle*, **198**, 373-390.
- Salked, D.J. (2004) *Eulamprus quoyii* (eastern water skink) predation. *Herpetological Review*, **35**, 389.
- Salvador, A., Díaz, J.A., Veiga, J.P., Bloor, P. & Brown, R.P. (2008) Correlates of reproductive success in male lizards of the alpine species *Iberolacerta cyreni*. *Behavioural Ecology*, **19**, 169-176.
- Savage, J.M. (1952) Studies on the lizard family Xantusiidae I. The systematic status of the Baja California night lizards allied to *Xantusia vigilis*, with the description of a new subspecies. *American Midland Naturalist*, **48**, 467-479.
- Sazima, I., Sazima, C. & Sazima, M. (2005) Little dragons prefer flowers to maidens: a lizard that laps nectar and pollinates trees. *Biota Neotropica*, **5**, 1-8.
- Schaffer, J. (2004) *Saltuarius cornutus*. *Herpetofauna of North Queensland*,
- Schall, J.J. (1983) Reproductive strategies in sympatric whiptail lizards (*Cnemidophorus*): two parthenogenetic and three bisexual species. *Copeia*, **1983**, 108-116.

- Schall, J.J. & Dearing, M.D. (1994) Body temperature of the herbivorous Bonaire Island whiptail lizard (*Cnemidophorus murinus*). *Journal of Herpetology*, **28**, 526-528.
- Schall, J.J. & Staats, C.M. (2002) Virulence of lizard malaria: three species of *Plasmodium* infecting *Anolis sabanus*, the endemic anole of Saba, Netherlands Antilles. *Copeia*, **2002**, 39-43.
- Schettino, L.R., Losos, J.B., Hertz, P.E., de Queiroz, K., Chamizo, A.R., Leal, M. & Gonzalez, V.R. (2010) The anoles of Soroa: aspects of their ecological relationships. *Breviora*, **520**, 1-22.
- Schlesinger, C.A. & Richard Shine, R. (1994) Selection of diurnal retreat sites by the nocturnal gekkonid lizard *Oedura lesueurii*. *Herpetologica*, **50**, 156-163.
- Schlesinger, C.A., Christian, K.A., James, C.D. & Morton, S.R. (2010) Seven lizard species and a blind snake: activity, body condition and growth of desert herpetofauna in relation to rainfall. *Australian Journal of Zoology*, **58**, 273-283.
- Schlüter, T. & Trauth, M.H. (2008) *Geological Atlas of Africa: With Notes on Stratigraphy, Tectonics, Economic Geology, Geohazards, Geosites and Geoscientific Education of Each Country*. Springer.
- Schmidt, K.P. (1921) New species of North American lizards of the genera *Holbrookia* and *Uta*. *American Museum Novitates*, **22**, 1-6.
- Schmidt, K.P. (1927) Notes on Chinese reptiles. *Bulletin of the American Museum of Natural History*, **54**, 467-551.
- Schmidt, K.P. (1943) Amphibians and reptiles from the Sudan. *Field Museum of Natural History, Zoology Series*, **24**, 331-338.
- Schmidt, K.P. & Bogert, C.M. (1947) A new fringe-footed sand lizard from Coahuila, Mexico. *American Museum Novitates*, **1339**, 1-9.
- Schmidt, K.P. & Inger, R.F. (1951) Amphibians and reptiles of the Hopkins-Branner expedition to Brazil. *Fieldiana Zoology*, **31**, 439-465.
- Schmidt, K.P., Lang, H. & Chapin, J.P. (1919) Contributions to the herpetology of the Belgian Congo based on the collection of the American Museum Congo Expedition, 1909-1915. Part 1, Turtles, crocodiles, lizards and chameleons. *Bulletin of the American Museum of Natural History*, **39**, 385-624.
- Schmidtler, J.F. (1997) Die *Ablepharus kitaibelii* – Gruppe in Süd-Anatolien und benachbarten Gebieten (Squamata: Sauria: Scinidae). *Herpetozoa*, **10**, 35-63.
- Schmidtler, J.F., Eiselt, J. & Darevsky, I.S. (1994) Untersuchungen an Feldeidechsen (Lacertesaxicola-Gruppe) in der östlichen Türkei: 3. Zwei neue parthogenetische Arten. *Salamandra*, **30**, 55-70.
- Schoener, T.W. (1977) Competition and the niche. *Biology of the Reptilia*, **7**, 35-136.
- Schoener, T.W. & Gorman, G.C. (1968) Some niche differences in three Lesser Antillean lizards of the genus *Anolis*. *Ecology*, **49**, 819-830.
- Schoener, T.W. & Schoener, A. (1971) Structural habitats of West Indian *Anolis* lizards I. Lowland Jamaica. *Breviora*, **368**, 1-53.
- Schoener, T.W. & Schoener, A. (1971) Structural habitats of West Indian *Anolis* Lizards II. Puerto Rican uplands. *Breviora*, **375**, 1-39.
- Schoener, T.W., Slade, J.B. & Stinson, C.H. (1982) Diet and sexual dimorphism in the very catholic lizard genus, *Leiocephalus*, of the Bahamas. *Oecologia*, **53**, 160-169.
- Schostakowski, E. (2001) *Oedura castelnaui*. *Herpetofauna of North Queensland*.
- Schulte, J.A., Macey, J.R., Espinoza, R.E. & Larson, A. (2000) Phylogenetic relationships in

- the iguanid lizard genus *Liolaemus*: multiple origins of viviparous reproduction and evidence for recurring Andean vicariance and dispersal. *Biological Journal of the Linnean Society*, **69**, 75-102.
- Schulte, J.A., Losos, J.B., Cruz, F.B. & Nunez, H. (2004) The relationship between morphology, escape behaviour and microhabitat occupation in the lizard clade *Liolaemus* (Iguanidae: Tropidurinae: Liolaemini). *Journal of Evolutionary Biology*, **17**, 408-420.
- Schwaneer, T.D. (1980) Reproductive biology of lizards on the American Samoan Islands. *Occasional Papers of the University of Kansas Museum of Natural History*, **86**, 1-53.
- Schwarzkopf, L. (1992) Annual variation of litter size and offspring size in a viviparous skink. *Herpetologica*, **48**, 390-395.
- Schwarzkopf, L. (1993) Costs of reproduction in water skinks. *Ecology*, **74**, 1970-1981.
- Schwarzkopf, L., Barnes, M. & Goodman, B. (2010) Belly up: reduced crevice accessibility as a cost of reproduction caused by increased girth in a rock-using lizard. *Austral Ecology*, **35**, 82-86.
- Scoular, K.M., Caffry, W.C., Tillman, J.L., Finan, E.S., Schwartz, S.K., Sinervo, B. & Zani, P.A. (2011) Multiyear home-range ecology of common sideblotched lizards in eastern Oregon with additional analysis of geographic variation in home-range size. *Herpetological Monographs*, **25**, 52-75.
- Seifan, M., Gilad, A., Klass, K. & Werner, Y.L. (2009) Ontogenetically stable dimorphism in a lacertid lizard (*Acanthodactylus boskianus*) with tests of methodology and comments on life-history. *Biological Journal of the Linnean Society*, **97**, 275-288.
- Serrano-Cardozo, V.H., Lemos-Espinal, J.A. & Smith, G.R. (2008) Comparative diet of three sympatric *Sceloporus* in the semiarid Zapotitlan Valley, Mexico. *Revista Mexicana de Biodiversidad*, **79**, 427-434.
- Setiadi, M.I. & Hamidy, A. (2006) *Jenis herpetofauna di Pulau Halmahera*. Center for Biodiversity and Conservation Studies Universitas Indonesia and Bogoriense Zoologicum Museum, Research Center for Biology Indonesian Institute of Sciences.
- Sexton, O.J., Andrews, R.M. & Bramble, J.E. (1992) Size and growth rate characteristics of a peripheral population of *Crotaphytus collaris* (Sauria: Crotaphytidae). *Copeia*, **1992**, 968-980.
- Shanbhag, B.A. (2002) Reproductive biology of Indian reptiles. *Proceedings of the Indian National Science Academy*, **68**, 497-528.
- Shannon, F.A. (1956) The reptiles and amphibians of Korea. *Herpetologica*, **12**, 22-49.
- Shaw, C.E. (1945) The chuckwalla, genus *Sauromalus*. *Transactions of the San Diego Society of Natural History*, **10**, 269-306.
- Shea, G.M. (1995) A taxonomic revision of the *Cyclodomorphus casuarinae* complex (Squamata: Scincidae). *Records of the Australian Museum*, **47**, 83-115.
- Shea, G.M. (2006) Diet of two species of bluetongue skink, *Tiliqua multifasciata* and *Tiliqua occipitalis* (Squamata: Scincidae). *Australian Zoologist*, **33**, 359-368.
- Shea, G.M. (2007) Range extension for *Lipinia nototaenia* (Boulenger, 1914) (Squamata : Scincidae) and the reproductive mode of the species. *Hamadryad*, **32**, 80-81.
- Shea, G.M. & Miller, B. (1995) A taxonomic revision of the *Cyclodomorphus branchialis* species group (Squamata: Scincidae). *Records of the Australian Museum*, **47**, 265-325.
- Shea, G., Jourdan, H., Sadlier, R. & Bauer, A. (2009) Natural history of the New Caledonian whiptailed skink *Tropidoscincus variabilis* (Bavay, 1869) (Squamata: Scincidae). *Amphibia-Reptilia*, **30**, 207-220.

- Sherbrooke, W.C. (1975) Reproductive cycle of a tropical teiid lizard, *Neusticurus ecleopus* Cope, in Peru. *Biotropica*, **7**, 194-207.
- Sherbrooke, W.C. (1997) Ditmars' horned lizard, or rock horned lizard: an historical update since rediscovery (1970). *Sonoran Herpetologist*, **10**, 6-8.
- Sherbrooke, W.C. (2002) *Phrynosoma modestum* (Round-tailed horned lizard). Death due to prey (beetle) ingestion. *Herpetological Review*, **33**, 312.
- Shine, R. (1983) Reptilian viviparity in cold climates: testing the assumptions of an evolutionary hypothesis. *Oecologia*, **57**, 397-405.
- Shine, R. (1986) Food habits, habitats and reproductive biology of four sympatric species of varanid lizards in tropical Australia. *Herpetologica*, **42**, 346-360.
- Shine, R. & Greer, A.E. (1991) Why are clutch sizes more variable in some species than in others? *Evolution*, **45**, 1696-1706.
- Shine, R. & Thomas, J. (2005) Do lizards and snakes really differ in their ability to take large prey? A study of relative prey mass and feeding tactics in lizards. *Oecologia*, **144**, 492-498.
- Shine, R. & Wall, M. (2008) Interactions between locomotion, feeding, and bodily elongation during the evolution of snakes. *Biological Journal of the Linnean Society*, **95**, 293-304.
- Sievert, L.M. & Paulissen, M.A. (1996) Temperature selection and thermoregulatory precision of bisexual and parthenogenetic *Cnemidophorus* lizards from southern Texas, USA. *Journal of Thermal Biology*, **21**, 15-20.
- Siliceo, I. & Diaz, J.A. (2010) A comparative study of clutch size, range size, and the conservation status of island vs. mainland lacertid lizards. *Biological Conservation*, **143**, 2601-2608.
- Simbotwe, M.P. (1980) Reproductive biology of the skinks *Mabuya striata* and *Mabuya quinquetaeniata* in Zambia. *Herpetologica*, **36**, 99-104.
- Simmons, P.M., Greene, B.T., Williamson, K.E., Powell, R. & Parmerlee, J.S. (2005) Ecological interactions within a lizard community on Grenada. *Herpetologica*, **61**, 124-134.
- Sinervo, B. & Licht, P. (1991) Proximate constraints on the evolution of egg size, number, and total clutch mass in lizards. *Science*, **252**, 1300-1302.
- Sinervo, B., Mendez-de-la-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagran-Santa Cruz, M., Lara-Resendiz, R., Martinez-Mendez, N., Calderon-Espinosa, M.L., Meza-Lazaro, R.N., Gadsden, H., Avila, L.J., Morando, M., De la Riva, I.J., Sepulveda, P.V., Rocha, C.F.D., Ibarquengoytia, N., Puntriano, C.A., Massot, M., Lepetz, V., Oksanen, T.A., Chapple, D.G., Bauer, A.M., Branch, W.R., Clobert, J. & Sites, J.W. (2010) Erosion of lizard diversity by climate change and altered thermal niches. *Science*, **328**, 894-899.
- Smedley, N. (1931) Amphibians and reptiles from the Cameron Highlands, Malay Peninsula. *Bulletin of the Raffles Museum*, **6**, 105-123.
- Smith, H.M. (1939) Notes on Mexican reptiles and amphibians. *Zoological Series of Field Museum of Natural History*, **24**, 15-35.
- Smith, H.M. (1942) Mexican herpetological miscellany. *Proceedings of the United States National Museum*, **92**, 349-395.
- Smith, H.M. (1968) Another new lizard from Mexico of the schiedi group of *Anolis*. *Southwestern Naturalist*, **13**, 368-370.
- Smith, F.A. (1992) Evolution of body size among woodrats from Baja California, Mexico. *Functional Ecology*, **6**, 265-273.
- Smith, B.E. (1993) Notes on a collection of squamate reptiles from eastern Mindanao, Philippine Islands part 1: Lacertilia. *Asiatic Herpetological Research*, **5**, 85-95.

- Smith, P.W. & Smith, H.M. (1952) Geographic variation in the lizard *Eumeces anthracinus*. *University of Kansas Science Bulletin*, **34**, 679-694.
- Smith, G.R. & Iverson, J.B. (1993) Reproduction in the curly-tailed lizard *Leiocephalus psammodromus* from the Caicos Islands. *Canadian Journal of Zoology*, **71**, 2147-2151.
- Smits, A.W. (1985) Behavioral and dietary responses to aridity in the chuckwalla, *Sauromalus hispidus*. *Journal of Herpetology*, **19**, 441-449.
- Smits, A.W., Ward, J. & Lillywhite, H. (1986) Effects of hyperkalemia on thermoregulatory and feeding behaviors of the lizard, *Sauromalus hispidus*. *Copeia*, **1986**, 518-520.
- Snell, H.L. & Christian, K.A. (1985) Energetics of Galapagos land iguanas: a comparison of two island populations. *Herpetologica*, **41**, 437-442.
- Somma, L.A. (1985) Notes on maternal behavior and post-brooding aggression in the prairie skink *Eumeces septentrionalis*. *Nebraska Herpetological Newsletter*, **6**, 9-12.
- Somma, L.A. (1987) Alloparental care in the prairie skink *Eumeces septentrionalis*: a case of mistaken identity? *Nebraska Herpetological Newsletter*, **8**, 5-8.
- Somma, L.A. (1987) Maternal care of neonates in the prairie skink, *Eumeces septentrionalis*. *Great Basin Naturalist*, **47**, 536-537.
- Somma, L.A. (1990) A categorization and bibliographic survey of parental behavior in lepidosaurian reptiles. *Smithsonian Herpetological Information Service*, **81**, 1-53.
- Somma, L.A. (1991) *Eumeces septentrionalis* (prairie skink). Piscivory. *Herpetological Review*, **22**, 58-59.
- Somma, C.A. & Brooks, G.R. (1976) Reproduction in *Anolis oculatus*, *Ameiva fuscata* and *Mabuya mabouya* from Dominica. *Copeia*, **1976**, 249-256.
- Song, M. (1987) Survey of the reptiles of southern Shaanxi. *Acta Herpetologica Sinica*, **6**, 59-64.
- Soule, M. (1963) Aspects of thermoregulation in nine species of lizards from Baja California. *Copeia*, **1963**, 107-115.
- Soule, M.E. (1966) Trends in insular radiation of a lizard. *American Naturalist*, **100**, 47-64.
- Sprackland, R.G. (1991) Taxonomic review of the *Varanus prasinus* group with descriptions of two new species. *Memoirs of the Queensland Museum*, **3**, 561-576.
- Stacy, B.A., Howard, L., Kinkaid, J., Vidal, J.D. & Papendick, R. (2008) Yolk coelomitis in Fiji Island banded iguanas (*Brachylophus fasciatus*). *Journal of Zoo and Wildlife Medicine*, **39**, 161-169.
- Stamps, J.A., Losos, J.B. & Andrews, R.M. (1997) A comparative study of population density and sexual size dimorphism in lizards. *American Naturalist*, **149**, 64-90.
- Steinberg, D.S., Hite, J.L., Powell, R. & Henderson, R.W. (2007) *Sphaerodactylus vincenti*. *Catalogue of American Amphibians and Reptiles*, **853**, 1-6.
- Steppan, S.J., Zawadzki, C. & Heaney, L.R. (2003) Molecular phylogeny of the endemic Philippine rodent *Apomys* (Muridae) and the dynamics of diversification in an oceanic archipelago. *Biological Journal of the Linnean Society*, **80**, 699-715.
- Stewart, J.R. (1985) Growth and survivorship in a California population of *Gerrhonotus coeruleus*, with comments on intraspecific variation in adult female size. *American Midland Naturalist*, **113**, 30-44.
- Stoliczka, F. (1873) Notes on some Andamese and Nicobarese reptiles, with the descriptions of three new species of lizards. *Journal of the Asiatic Society of Bengal*, **42**, 162-169.
- Storr, G.M. (1978) The genus *Egernia* (Lacertilia, Scincidae) in Western Australia. *Records of the Western Australian Museum*, **6**, 147-187.

- Stuart, L.C. (1939) A description of a new *Gymnophthalmus* from Guatemala, with notes on other members of the genus. *Occasional Papers of the Museum of Zoology, University of Michigan*, **409**, 1-12.
- Stuart, L.C. (1955) A brief review of the Guatemalan lizards of the genus *Anolis*. *Miscellaneous Papers of the Museum of Zoology, University of Michigan*, **91**, 1-31.
- Stuart, J.N. (1991) *Cnemidophorus exsanguis*. *Catalogue of American Amphibians and Reptiles*, **516**, 1-4.
- Stuart, J.N. (1998) *Cnemidophorus velox*. *Catalogue of American Amphibians and Reptiles*, **656**, 1-6.
- Stuart, B.L. & Emmett, D.A. (2006) A collection of amphibians and reptiles from the Cardamom Mountains, southwestern Cambodia. *Fieldiana*, **109**, 1-27.
- Stuart, B.L., Dugan, K.A., Allard, M.W. & Kearney, M. (2006) Extraction of nuclear DNA from bone of skeletonized and fluid-preserved museum specimens. *Systematics and Biodiversity*, **4**, 133-136.
- Stuart-Fox, D. (2008) A test of Rensch's rule in dwarf chameleons (*Bradypodion* spp.), a group with female-biased sexual size dimorphism. *Evolutionary Ecology*, **23**, 425-433.
- Stuart-Smith, J., Swain, R.S.-S., R. D. & Wapstra, E. (2008) Size dimorphism in *Rankinia* [*Tympanocryptis*] *diemensis* (Family Agamidae): sex-specific patterns and geographic variation. *Biological Journal of the Linnean Society*, **94**, 699-709.
- Sturaro, M.J. & da Silva, V.X. (2010) Natural history of the lizard *Enyalius perditus* (Squamata: Leiosauridae) from an Atlantic forest remnant in southeastern Brazil. *Journal of Natural History*, **44**, 1225-1238.
- Subach, A., Scharf, I. & Ovadia, O. (2009) Foraging behavior and predation success of the sand viper (*Cerastes vipera*). *Canadian Journal of Zoology*, **87**, 520-528.
- Subba Rao, M.V. & Rajabai, B.S. (1972) Reproduction in the ground lizard, *Sitana ponticeriana* and garden lizard, *Calotes nemoricola*. *British Journal of Herpetology*, **4**, 245-251.
- Sutherland, D.R. (2011) Dietary niche overlap and size partitioning in sympatric varanid lizards. *Herpetologica*, **67**, 146-153.
- Sweet, S.S. & Pianka, E.R. (2007) Monitors, mammals, and Wallace's Line. *Mertensiella*, **16**, 79-99.
- Szyndlar, Z. (1984) A description of a small collection of amphibians and reptiles from the People's Democratic Republic of Korea with notes on the distribution of the herpetofauna in that country. *Acta Zoologica Cracoviensia*, **27**, 1-18.
- Tadevosyan, T.L. (2007) The role of vegetation in microhabitat selection of syntopic lizards, *Phrynocephalus persicus*, *Eremias pleskei*, and *Eremias strauchi* from Armenia. *Amphibia-Reptilia*, **28**, 444-448.
- Tanaka, D. (1986) Thermal ecology of the forest-dwelling agamid lizard, *Japalura polygonata ishigakiensis*. *Journal of Herpetology*, **20**, 333-340.
- Tanner, W.W. (1987) Lizards and turtles of Western Chihuahua. *Great Basin Naturalist*, **47**, 383-421.
- Taran, Y.A., Fischer, T., Cienfuegos, E. & Morales, P. (2002) Geochemistry of hydrothermal fluids from an intraplate ocean island: Everman volcano, Socorro Island, Mexico. *Chemical geology*, **188**, 51-63.
- Tayhan, Y., Cicek, K., Ayaz, D. & Tok, C.V. (2011) Data on hatchlings of Caucasian rock lizard, *Darevskia valentini* (Boettger, 1892) in northeastern Anatolia. *Ecologia Balkanica*,

- 3, 75-78.
- Taylor, E.H. (1953) A review of the lizards of Ceylon. *University of Kansas Science Bulletin*, **35**, 1525-1585.
- Taylor, E.H. (1956) A review of the lizards of Costa Rica. *University of Kansas Science Bulletin*, **38**, 3-322.
- Taylor, E.H. (1962) New Oriental reptiles. *University of Kansas Science Bulletin*, **43**, 209-263.
- Taylor, E.H. (1963) The lizards of Thailand. *University of Kansas Science Bulletin*, **44**, 687-1077.
- Taylor, J.A. (1985) Reproductive biology of the Australian lizard *Ctenotus taeniolatus*. *Herpetologica*, **41**, 408-418.
- Taylor, J.E. (2004) Reproduction in sympatric lizards: comparison of two species of *Ctenotus* (Scincidae) in south-eastern Australia. *Australian Journal of Zoology*, **52**, 649-666.
- Taylor, E.H. & Elbel, R.E. (1958) Contribution to the herpetology of Thailand. *University of Kansas Science Bulletin*, **38**, 1033-1189.
- Taylor, H.L., Walker, J.M. & Cordes, J.E. (1999) Possible phylogenetic constraint on clutch size in the parthenogenetic teiid lizard *Cnemidophorus neotesselatus*. *Journal of Herpetology*, **33**, 319-323.
- Taylor, H.L., Lemos-Espinal, J.A. & Smith, H.M. (2003) Morphological characteristics of a newly discovered population of *Aspidoscelis tessellata* (Squamata: Teiidae) from Chihuahua, Mexico, the identity of an associated hybrid, and a pattern of geographic variation. *Southwestern Naturalist*, **48**, 692-700.
- Telemeco, R.S. & Baird, T.A. (2011) Capital energy drives production of multiple clutches whereas income energy fuels growth in female collared lizards *Crotaphytus collaris*. *Oikos*, **120**, 915-921.
- Telford, R. (1955) The lizard *Eumeces anthracinus* in central Virginia. *Copeia*, **1955**, 143.
- Telford, S.R. (1959) A study of the sand skink, *Neoseps reynoldsi* Stejneger. *Copeia*, **1959**, 110-119.
- Telford, S.R. (1969) *Neoseps*, *N. reynoldsi*. *Catalogue of American Amphibians and Reptiles*, **80**, 1-2.
- Telford, S.R. (1971) Reproductive patterns and relative abundance of two microteiid lizard species in Panama. *Copeia*, **1971**, 670-675.
- Terbish, K., Munkhbayar, K., Clark, E.L., Munkhbat, J., Monks, E.M., Munkhbaatar, M., Baillie, J.E.M., Borkin, L., Batsaikhan, N., Samiya, R. & Semenov, D.V. (2006) *Mongolian red list of reptiles and amphibians*. Mongolian red list of reptiles and amphibians. Regional Red List Series Vol. 5. Zoological Society of London, London.
- Tersa, E., Pether, J. & Mateo, J.A. (2010) Evaluación de las poblaciones de reptiles canarios introducidos en Fuerteventura (Islas Canarias). *Boletín de la Asociación Herpetológica Española*, **21**, 104-109.
- Test, F.H., Sexton, O.J. & Heatwole, H. (1966) Reptiles of Rancho Grande and vicinity, Estado Aragua, Venezuela. *Miscellaneous Publications of the Museum of Zoology, University of Michigan*, **128**, 1-63.
- Teynie, A. (2004) Notes on reptiles of Nam Lan conservation area in Phongsaly province of Lao PDR. *Report for Phongsaly Forest Conservation and rural Development Project*.
- Teynie, A., David, P. & Ohler, A. (2010) Note on a collection of amphibians and reptiles from western Sumatra (Indonesia), with the description of a new species of the genus *Bufo*. *Zootaxa*, **2416**, 1-43.

- Thomas, R. (1966) A reassessment of the herpetofauna of Navassa Island. *Journal of the Ohio Herpetological Society*, **5**, 73-89.
- Thomas, B.W. (1981) *Hoplodactylus rakiurae* n. sp. (Reptilia: Gekkonidae) from Stewart Island, New Zealand and comments on the taxonomic status of *Heteropholis nebulosus* McCann. *New Zealand Journal of Zoology*, **8**, 33-47.
- Thompson, G.G. & Pianka, E.R. (2001) Allometry of clutch and neonate sizes in monitor lizards (Varanidae: *Varanus*). *Copeia*, **2001**, 443-458.
- Thompson, M.B., Speake, B.K., Stewart, J.R., Russell, K.J. & McCartney, R.J. (1999) Placental transfer of nutrients during gestation in the viviparous lizard, *Pseudemoia spenceri*. *Journal of Comparative Physiology and Biochemistry*, **169**, 319-328.
- Thompson, G.G., Clemente, C.J., Withers, P.C., Fry, B.G. & Norman, J.A. (2008) Is body shape of varanid lizards linked with retreat choice? *Australian Journal of Zoology*, **56**, 351-362.
- Throckmorton, G. (1973) Digestive efficiency in the herbivorous lizard *Ctenosaura pectinata*. *Copeia*, **1973**, 431-435.
- Tihen, J.A. (1954) Gerrhonotine lizards recently added to the American Museum collection : with further revisions of the genus *Abronia*. *American Museum Novitates*, **1687**, 1-26.
- Tinkle, D.W. (1967) The life and demography of the side-blotched lizard, *Uta stansburiana*. *Miscellaneous Publications of the Museum of Zoology University of Michigan*, **132**, 1-182.
- Tinkle, D.W., Wilbur, H.M. & Tilley, S.G. (1970) Evolutionary strategies in lizard reproduction. *Evolution*, **24**, 55-74.
- Toda, M., Sengoku, S., Hikida, T. & Ota, H. (2008) Description of two new species of the genus *Gekko* (Squamata: Gekkonidae) from the Tokara and Amami Island groups in the Ryukyu Archipelago, Japan. *Copeia*, **2008**, 452-466.
- Todd, A.C. (2008) Using testis size to predict the mating systems of New Zealand geckos. *New Zealand Journal of Zoology*, **35**, 103-114.
- Toledo, L.F., Ribeiro, R.S. & Haddad, C.F.B. (2007) Anurans as prey: an exploratory analysis and size relationships between predators and their prey. *Journal of Zoology*, **271**, 170-177.
- Tomasevic-Kolarov, N., Ljubisavljevic, K., Polovic, L., Dzukic, G. & Kalezic, M. (2010) The body size, age structure and growth pattern of the endemic balkan mosor rock lizards, *Dinarolacerta mosorensis* (Kolombatovic, 1886). *Acta Zoologica Academiae Scientiarum Hungaricae*, **56**, 55-71.
- Torres-Carvajal, O. (2007) A taxonomic revision of south American *Stenocercus* (Squamata: Iguania) lizards. *Herpetological Monographs*, **21**, 76-178.
- Torres-Carvajal, O. (2007) New Andean species of *Stenocercus* (Squamata: Iguania) from the eastern Cordillera in Colombia. *Copeia*, **2007**, 56-61.
- Towns, D.R. (1994) The role of ecological restoration in the conservation of Whitaker's skink (*Cyclodina whitakeri*), a rare New Zealand lizard (Lacertilia: Scincidae). *New Zealand Journal of Zoology*, **21**, 457-471.
- Towns, D.R. (2002) Korapuki Island as a case study for restoration of insular ecosystems in New Zealand. *Journal of Biogeography*, **29**, 593-607.
- Towns, D.R. & Elliot, G.P. (1996) Effects of habitat structure on distribution and abundance of lizards at Pukerua Bay, Wellington, New Zealand. *New Zealand Journal of Ecology*, **20**, 191-206.
- Towns, D.R.N., K. A. & Whitaker, A.H. (2002) North Island *Oligosoma* spp. skink recovery plan 2002–2012. Threatened Species Recovery Plan 48. Department of Conservation,

Wellington.

- Trubcheninova, L.P., Khutoryansky, A.A., Svet-Moldavsky, G.J., Kuznetsova, L.E., Sokolov, P.P. & Belianchykova, N.I. (1977) Body temperature and tumor virus infection. I. Tumorigenicity of Rous sarcoma virus for reptiles. *Neoplasma*, **24**, 3-19.
- Trudgen, T. (1999) *Lialis burtonis*. *Herpetofauna of North Queensland*,
- Tsasi, G., Pafilis, P., Simou, C. & Valakos, E.D. (2009) Predation pressure, density-induced stress and tail regeneration: a casual-nexus situation or a bunch of independent factors? *Amphibia-Reptilia*, **30**, 471-482.
- Tulli, M.J., Cruz, F.B., Herrel, A., Vanhooydonck, B. & Abdala, V. (2009) The interplay between claw morphology and microhabitat use in neotropical iguanian lizards. *Zoology*, **112**, 379-392.
- Tuniyev, S.B., Doronin, I.A., Kidov, A.A. & Tuniyev, B.S. (2011) Systematic and geographical variability of meadow lizard, *Darevskia praticola* (Reptilia: Sauria) in the Caucasus. *Russian Journal of Herpetology*, **18**, 295-316.
- Turner, F.B., Jennrich, R.I. & Weintraub, J.D. (1969) Home ranges and body size of lizards. *Ecology*, **50**, 1076-1081.
- Ugueto, G.N., Harvey, M.B. & Rivas, G.A. (2009) Two new species of *Cnemidophorus* (Squamata: Teiidae) from islands of the northeastern coast of Venezuela. *Herpetological Monographs*, **23**, 123-153.
- Ullénbruch, K., Grell, O. & Böhme, W. (2010) Reptiles from southern Benin, West Africa, with the description of a new *Hemidactylus* (Gekkonidae), and a country-wide checklist. *Bonn Zoological Bulletin*, **57**, 31-54.
- Uller, T. & Olsson, M. (2010) Offspring size and timing of hatching determine survival and reproductive output in a lizard. *Oecologia*, **162**, 663-671.
- Uller, T., While, G.M., Wapstra, E., Warner, D.A., Goodman, B.A., Schwarzkopf, L., Langkilde, T., Doughty, P., Radder, R.S., Rohr, D.H., Bull, C.M., Shine, R. & Olsson, M. (2009) Evaluation of offspring size–number invariants in 12 species of lizard. *Journal of Evolutionary Biology*, **22**, 143-151.
- Uzzell, T.M. (1966) Teiid lizards of the genus *Neusticurus* (Reptilia, Sauria). *Bulletin of the American Museum of Natural History*, **132**, 277-328.
- Vacher, H.L. & Quinn, T.M. (1997) *Geology and Hydrogeology of Carbonate Islands*. Elsevier.
- Vagelli, A.A. (2011) *The Banggai Cardinalfish: Natural History, Conservation, and Culture of Pterapogon kauderni*. John Wiley & Sons.
- Valakos, E. (1986) The feeding ecology of *Podarcis erhardii* (Reptilia – Lacertidae) in a main insular ecosystem. *Herpetological Journal*, **1**, 118-121.
- Valdez-Gonzalez, M.A. & Ramirez-Bautista, A. (2002) Reproductive characteristics of the spiny lizards, *Sceloporus horridus* and *Sceloporus spinosus* (Squamata : Phrynosomatidae) from Mexico. *Journal of Herpetology*, **36**, 36-43.
- Valido, A. & Nogales, M. (2003) Digestive ecology of two omnivorous Canarian lizard species (*Gallotia*, Lacertidae). *Amphibia-Reptilia*, **24**, 331-344.
- Valverde, J.A. (1958) Una nueva lagartija del genero *Algyroides* Bribon, procedente de la Sierra de Cazorla (Sur de Espana). *Archivos del Instituto de Aclimatacion, Almeria*, **7**, 127-134.
- van Berkum, F.H. (1986) Evolutionary patterns of the thermal sensitivity of sprint speed in *Anolis* lizards. *Evolution*, **40**, 594-604.

- van Berkum, F.H. (1988) Latitudinal patterns of the thermal sensitivity of sprint speed in lizards. *American Naturalist*, **132**, 327-343.
- Van Denburgh, J. (1922) The reptiles of western North America. Volume I. Lizards. *Occasional Papers of the California Academy of Sciences*, **10**, 1-612.
- Van Denburgh, J. & Slevin, J.R. (1913) Expedition of the California Academy of Sciences to the Galapagos Islands, 1905-1906. IX. The Galapagoan lizards of the genus *Tropidurus* with notes on iguanas of the genera *Conolophus* and *Amblyrhynchus*. *Proceedings of the California Academy of Sciences*, **2**, 132-202.
- van der Kooij, J. (2001) The herpetofauna of the Sultanate of Oman. Part 1: The amphibians, worm lizards, agamas and chameleons. *Podarcis*, **1**, 70-82.
- van der Meer, M.H., Whiting, M.J. & Branch, W.R. (2010) Ecology of southern African Sandveld lizards (Lacertidae, *Nucras*). *Copeia*, **2010**, 568-577.
- Van Sluys, M., Ferreira, V.M. & Rocha, C.F.D. (2004) Natural history of the lizard *Enyalius brasiliensis* (Lesson, 1828) (Leiosauridae) from an atlantic forest of southeastern Brazil. *Brazilian Journal of Biology*, **64**, 353-356.
- Van Sluys, M., Martelotte, S.B., Kiefer, M.C. & Rocha, C.F.D. (2010) Reproduction in neotropical *Tropidurus* lizards (Tropiduridae): evaluating the effect of environmental factors on *T. torquatus*. *Amphibia-Reptilia*, **31**, 117-126.
- Van Wilgen, N.J. & Richardson, D.M. (2012) The roles of climate, phylogenetic relatedness, introduction effort, and reproductive traits in the establishment of non-native reptiles and amphibians. *Conservation Biology*, **26**, 267-277.
- Van Wyk, J.H. (1991) Biennial reproduction in the female viviparous lizard *Cordylus giganteus*. *Amphibia-Reptilia*, **12**, 329-342.
- van Wyk, J.H. (1994) Physiological changes during the female reproductive cycle of the viviparous lizard *Cordylus giganteus* (Sauria: Cordylidae). *Herpetologica*, **50**, 480-493.
- Vanhooydonck, B. & Van Damme, R. (1999) Evolutionary relationships between body shape and habitat use in lacertid lizards. *Evolutionary Ecology Research*, **1**, 785-805.
- Vanzolini, P.E. (2005) On *Gymnodactylus amarali* Barbour, 1925, with the description of a new species (Sauria, Gekkonidae). *Anais da Academia Brasileira de Ciencias*, **77**, 595-611.
- Vega, L.E. (2001) Reproductive and feeding ecology of the amphisbaenian *Anops kingii* in east-central Argentina. *Amphibia-Reptilia*, **22**, 447-454.
- Vega-Castillo, S.I. & Puente-Rolon, A.R. (2011) *Anolis* species. Frugivory. *Herpetological Review*, **42**, 598-599.
- Veron, J.E.N. (1969) The reproductive cycle of the water skink, *Sphenomorphus quoyii*. *Journal of Herpetology*, **3**, 55-63.
- Verrastro, L., Veronese, L., Bujes, C. & Dias Filho, M.M. (2003) A new species of *Liolaemus* from southern Brazil (Iguania: Tropiduridae). *Herpetologica*, **59**, 105-118.
- Vervust, B. (2011) Lizards on islands: a model system for studying fast phenotypic change. *Dissertation*.
- Vervust, B., Grbac, I. & Van Damme, R. (2007) Differences in morphology, performance and behaviour between recently diverged populations of *Podarcis sicula* mirror differences in predation pressure. *Oikos*, **116**, 1343-1352.
- Vervust, B., Lailvaux, S., Grbac, I. & Van Damme, R. (2008) Do morphological condition indices predict locomotor performance in the lizard *Podarcis sicula*? *Acta Oecologica*, **34**, 244-251.
- Vervust, B., Grbac, I., Brecko, J. & Tvrtkovic, N and Van Damme, R. (2009) Distribution of

- reptiles and amphibians in the nature park Lastovo Archipelago: possible underlying biotic and abiotic causes. *Natura Croatica*, **18**, 113-127.
- Vervust, B., Pafilis, P., Valakos, E.D. & Van Damme, R. (2010) Anatomical and physiological changes associated with a recent dietary shift in the lizard *Podarcis sicula*. *Physiological and Biochemical Zoology*, **83**, 632-642.
- Verwajen, D. & Van Damme, R. (2007) Correlated evolution of thermal characteristics and foraging strategy in lacertid lizards. *Journal of Thermal Biology*, **32**, 388-395.
- Verwajen, D. & Van Damme, R. (2008) Foraging mode and its flexibility in lacertid lizards from Europe. *Journal of Herpetology*, **42**, 124-133.
- Vial, J.L. & Stewart, J.R. (1985) The reproductive cycle of *Barisia monticola*: a unique variation among viviparous lizards. *Herpetologica*, **41**, 51-57.
- Vidal, M., Ortiz, J.C. & Labra, A. (2002) Sexual and age differences in ecological variables of the lizard *Microlophus atacamensis* (Tropiduridae) from northern Chile. *Revista Chilena de Historia Natural*, **75**, 283-292.
- Vidal, M.A., Veloso, A. & Mendez, M.A. (2006) Insular morphological divergence in the lizard *Liolaemus pictus* (Liolaemidae). *Amphibia-Reptilia*, **27**, 103-111.
- Vinegar, M.B. (1975) Life history phenomena in two populations of the lizard *Sceloporus undulatus* in southwestern New Mexico. *American Midland Naturalist*, **93**, 388-402.
- Vinson, J. & Vinson, J.-M. (1969) The saurian fauna of the Mascarene Islands. *Mauritius Institute Bulletin*, **6**, 203-320.
- Vitt, L.J. (1974) Reproductive effort and energy comparisons of adults, eggs, and neonates of *Gerrhonotus coeruleus principis*. *Journal of Herpetology*, **8**, 165-168.
- Vitt, L.J. (1981) Lizard reproduction: habitat specificity and constraints on relative clutch mass. *American Naturalist*, **117**, 506-514.
- Vitt, L.J. (1986) Reproductive tactics of sympatric gekkonid lizards with a comment on the evolutionary and ecological consequences of invariant clutch size. *Copeia*, **1986**, 773-786.
- Vitt, L.J. (1991) An introduction to the ecology of cerrado lizards. *Journal of Herpetology*, **25**, 79-90.
- Vitt, L.J. (1995) The ecology of tropical lizards in the caatinga of northeast Brazil. *Occasional Papers of the Oklahoma Museum of Natural History*, **1**, 1-29.
- Vitt, L.J. (2000) Ecological consequences of body size in neonatal and small-bodied lizards in the neotropics. *Herpetological Monographs*, **14**, 388-400.
- Vitt, L.J. & Ohmart, R.D. (1975) Ecology, reproduction, and reproductive effort of the iguanid lizard *Urosaurus graciosus* on the lower Colorado River. *Herpetologica*, **31**, 56-65.
- Vitt, L.J. & Pianka, E.R. (1977) Patterns of niche overlap among broadly sympatric versus narrowly sympatric Kalahari lizards (Scincidae: *Mabuya*). *Ecology*, **58**, 119-128.
- Vitt, L.J. & Price, H.J. (1982) Ecological and evolutionary determinants of relative clutch mass in lizards. *Herpetologica*, **38**, 237-255.
- Vitt, L.J. & Goldberg, S.R. (1983) Reproductive ecology of two tropical iguanid lizards: *Tropidurus torquatus* and *Platynotus semitaeniatus*. *Copeia*, **1983**, 131-141.
- Vitt, L.J. & Cooper, W.E. (1985) The evolution of sexual dimorphism in the skink *Eumeces laticeps*: an example of sexual selection. *Canadian Journal of Zoology*, **63**, 995-1002.
- Vitt, L.J. & Cooper, W.E. (1986) Foraging and diet of a diurnal predator (*Eumeces laticeps*) feeding on hidden prey. *Journal of Herpetology*, **20**, 408-415.
- Vitt, L.J. & Dickson, N.A. (1988) *Urosaurus graciosus*. *Catalogue of American Amphibians and Reptiles*, **448**, 1-3.

- Vitt, L.J. & Blackburn, D.G. (1991) Ecology and Life History of the viviparous lizard *Mabuya bistrriata* (Scincidae) in the Brazilian Amazon. *Copeia*, **1991**, 916-927.
- Vitt, L.J. & de Carvalho, C.M. (1992) Life in the trees: the ecology and life history of *Kentropyx striatus* (Teiidae) in the lavrado area of Roraima, Brazil, with comments on the life histories of tropical teiid lizards. *Canadian Journal of Zoology*, **70**, 1995-2006.
- Vitt, L.J. & Caldwell, J.P. (1993) Ecological observations on cerrado lizards in Rondonia, Brazil. *Journal of Herpetology*, **27**, 46-52.
- Vitt, L.J. & de Carvalho, C.M. (1995) Niche partitioning in a tropical wet season: lizards in the lavrado area of northern Brazil. *Copeia*, **1995**, 305-329.
- Vitt, L.J. & Zani, P.A. (1996) Ecology of the elusive tropical lizard *Tropidurus* [= *Uracentron*] *flaviceps* (Tropiduridae) in lowland rain forest of Ecuador. *Herpetologica*, **52**, 121-132.
- Vitt, L.J. & Zani, P.A. (1996) Ecology of the lizard *Ameiva festiva* (Teiidae) in southeastern Nicaragua. *Journal of Herpetology*, **30**, 110-117.
- Vitt, L.J. & Zani, P.A. (1996) Ecology of the South American lizard *Norops chrysolepis* (Polychrotidae). *Copeia*, **1996**, 56-68.
- Vitt, L.J. & Avila-Pires, T.C.S. (1998) Ecology of two sympatric species of *Neusticurus* (Sauria: Gymnophthalmidae) in the western Amazon of Brazil. *Copeia*, **1998**, 570-582.
- Vitt, L.J. & Zani, P.A. (1998) Ecological relationships among sympatric lizards in a transitional forest in the northern Amazon of Brazil. *Journal of Tropical Ecology*, **14**, 63-86.
- Vitt, L.J., Van Loben Sels, R.C. & Ohmart, R.D. (1978) Lizard reproduction: annual variation and environmental correlates in the iguanid lizard *Urosaurus graciosus*. *Herpetologica*, **34**, 241-253.
- Vitt, L.J., Zani, P.A. & Durtsche, R.D. (1995) Ecology of the lizard *Norops oxylophus* (Polychrotidae) in lowland forest of south-eastern Nicaragua. *Canadian Journal of Zoology*, **73**, 1918-1927.
- Vitt, L.J., Avila-Pires, T.C.S. & Zani, P.A. (1996) Observations on the ecology of the rare Amazonian lizard *Enyalius leechii* (Polychrotidae). *Herpetological Natural History*, **4**, 77-82.
- Vitt, L.J., Zani, P.A. & Avila-Pires, T.C.S. (1997) Ecology of the arboreal tropidurid lizard *Tropidurus* (= *Plica*) *umbra* in the Amazon region. *Canadian Journal of Zoology*, **75**, 1876-1882.
- Vitt, L.J., Zani, P.A. & Esposito, M.C. (1999) Historical ecology of Amazonian lizards: implications for community ecology. *Oikos*, **87**, 286-294.
- Vitt, L.J., Zani, P.A., Caldwell, J.P. & Durtsche, R.D. (1993) Ecology of the whiptail lizard *Cnemidophorus deppii* on a tropical beach. *Canadian Journal of Zoology*, **71**, 2391-2400.
- Vitt, L.J., Sartorius, S.S., Avila-Pires, T.C.S. & Esposito, M.C. (1998) Use of time, space, and food by the gymnophthalmid lizard *Prionodactylus eigenmanni* from the western Amazon of Brazil. *Canadian Journal of Zoology*, **76**, 1981-1988.
- Vitt, L.J., Sartorius, S.S., Avila-Pires, T.C.S., Esposito, M.C. & Miles, D.B. (2000) Niche segregation among sympatric Amazonian teiid lizards. *Oecologia*, **122**, 410-420.
- Vitt, L.J., Shepard, D.B., Caldwell, J.P., Vieira, G.H.C., Franca, F.G.R. & Colli, G.R. (2007) Living with your food: geckos in termitaria of Cantao. *Journal of Zoology*, **272**, 321-328.
- Vrcibradic, D. & Rocha, C.F.D. (1996) Ecological differences in tropical sympatric skinks (*Mabuya agilis* and *Mabuya macrorhyncha*) in southeastern Brazil. *Journal of Herpetology*, **30**, 60-67.
- Vrcibradic, D. & Rocha, C.F.D. (1998) Ecology of the skink *Mabuya frenata* in an area of rock

- outcrops in southeastern Brazil. *Journal of Herpetology*, **32**, 229-237.
- Vrcibradic, D. & Rocha, C.F.D. (1998) Reproductive cycle and life-history traits of the viviparous skink *Mabuya frenata* in southeastern Brazil. *Copeia*, **1998**, 612-619.
- Vrcibradic, D. & Rocha, C.F.D. (2005) Observations on the natural history of the lizard *Mabuya macrorhyncha* Hoge (Scincidae) in Queimada Grande Island, Sao Paulo, Brazil. *Revista Brasileira de Zoologia*, **22**, 1185-1190.
- Vrcibradic, D., Almeida-Gomes, M., Borges-Junior, V.N.T., Kiefer, M.C., Van Sluys, M. & Reptilia, R., C. F. D. (2006) Scincidae, *Mabuya frenata*: distribution extension. *Check List*, **2**, 57-58.
- Wagner, P., Melville, J., Wilms, T.M. & Schmitz, A. (2011) Opening a box of cryptic taxa – the first review of the North African desert lizards in the *Trapelus mutabilis* Merrem, 1820 complex (Squamata: Agamidae) with descriptions of new taxa. *Zoological Journal of the Linnean Society*, **163**, 884-912.
- Walker, C.F. (1955) Two new lizards of the genus *Lepidophyma* from Tamaulipas. *Occasional Papers of the Museum of Zoology, University of Michigan*, **564**, 1-10.
- Walker, J.M. (1981) Population structure and reproductive characteristics in *Cnemidophorus parvisocius* (Lacertilia: Teiidae). *American Midland Naturalist*, **105**, 217-224.
- Walker, J.M. (1987) Habitat and population destruction and recovery in the parthenogenetic whiptail lizard, *Cnemidophorus laredoensis* (Sauria: Teiidae) in southern Texas. *Texas Journal of Science*, **39**, 81-88.
- Walker, J.M., Taylor, H.L. & Maslin, T.P. (1966) Morphology and relations of the teiid lizard, *Cnemidophorus ceralbensis*. *Copeia*, **1966**, 585-588.
- Walley, H.D. (1998) *Eumeces anthracinus*. *Catalogue of American Amphibians and Reptiles*, **658**, 1-6.
- Waltner, R.C. (1991) Altitudinal ecology of *Agama tuberculata* Gray in the western Himalayas. *University of Kansas Museum of Natural History Miscellaneous Publications*, **83**, 1-74.
- Warne, R.W. & Charnov, E.L. (2008) Reproductive allometry and the size-number trade-off for lizards. *American Naturalist*, **172**, E80-E98.
- Warner, D.A., Bonnet, X., Hobson, K.A. & Shine, R. (2008) Lizards combine stored energy and recently acquired nutrients flexibly to fuel reproduction. *Journal of Animal Ecology*, **77**, 1242-1249.
- Warrick, G.D., Kato, T.T. & Rose, B.R. (1998) Microhabitat use and home range characteristics of blunt-nosed leopard lizards. *Journal of Herpetology*, **32**, 183-191.
- Watkins-Colwell, G.J., Smith, H.M. & Chiszar, D. (2003) *Sceloporus slevini*. *Catalogue of American Amphibians and Reptiles*, **771**, 1-6.
- Watling, J.I., Waddle, J.H., Kizirian, D. & Donnelly, M.A. (2005) Reproductive phenology of three lizard species in Costa Rica, with comments on seasonal reproduction of neotropical lizards. *Journal of Herpetology*, **39**, 341-348.
- Watson, C.M. (2005) *Eumeces fasciatus* (Five-lined Skink). Hatchlings. *Herpetological Review*, **36**, 318.
- Watt, C. (2002) *Eulamprus quoyii*. *Herpetofauna of North Queensland*,
- Webb, R.G. (2006) Variation in the crevice spiny lizard, *Sceloporus poinsettii* Baird and Girard. *Bulletin of the Maryland Herpetological Society*, **42**, 65-114.
- Webb, R.G. (2008) *Sceloporus poinsettii*. *Catalogue of American Amphibians and Reptiles*, **856**, 1-18.
- Webb, R.G. & Greer, J.K. (1969) Amphibians and reptiles from Malleco Province. *Publications*

- of the Museum, Michigan State University*, **4**, 193-226.
- Webb, J. & Shine, R. (1994) Feeding habits and reproductive biology of Australian pygopodid lizards of the genus *Aprasia*. *Copeia*, **1994**, 390-398.
- Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002) Phylogenies and community ecology. *Annual Review of Ecology and Systematics*, **33**, 475-505.
- Weber, N.A. (1960) Some Iraq amphibians and reptiles with notes on their food habits. *Copeia*, **1960**, 153-154.
- Weijola, V.S.-A. (2010) Geographical distribution and habitat use of monitor lizards of the North Moluccas. *Biawak*, **4**, 7-23.
- Wells, R.W. (2002) Taxonomic notes on the genus *Cyrtodactylus* (Reptilia: Gekkonidae) in Australia. *Australian Biodiversity Record*, **3**, 1-8.
- Wells, R.W. & Wellington, C.R. (1983) A synopsis of the class Reptilia in Australia. *Australian Journal of Herpetology*, **1**, 1-59.
- Werler, J.E. (1949) Reproduction of captive Texas and Mexican lizards. *Herpetologica*, **5**, 67-70.
- Werler, J.E. (1951) Miscellaneous notes on the eggs and young of Texan and Mexican reptiles. *Zoologica*, **36**, 37-55.
- Werneck, F.D.P., Giugliano, L.G., Collevatti, R.G. & Colli, G. (2009) Phylogeny, biogeography and evolution of clutch size in South American lizards of the genus *Kentropyx* (Squamata: Teiidae). *Molecular Ecology*, **18**, 262-278.
- Werner, F. (1908) Ergebnisse der mit subvention aus der erbschaft treitl unternommenen zoologischen Forschungsreise Dr. Franz Werner's nach dem Egyptischen Sudan und Norduganda. *Sitzungsber Akademie der Wissenschaften Wien*, **116**, 1823-1926.
- Werner, F. (1930) Contribution to the knowledge of the reptiles and amphibians of Greece, especially the Aegean islands. *Occasional Papers of the Museum of Zoology, University of Michigan*, **211**, 1-48.
- Werner, Y.L. (1968) Distribution of the Saharan *Sphenops sepsoides* (Reptilia: Scincidae) in Israel and Jordan. *Herpetologica*, **24**, 238-242.
- Werner, Y.L. (1973) *The reptiles of the Sinai Peninsula*. Hebrew University of Jerusalem (in Hebrew).
- Werner, Y.L. (1976) Optimal temperatures for inner-ear performance in gekkonid lizards. *Journal of Experimental Zoology*, **195**, 319-351.
- Werner, Y.L. (1980) Habitat-dependent thermal regimes of two Hawaiian geckos (Reptilia: Gekkonidae). *Journal of Thermal Biology*, **15**, 281-290.
- Werner, D.I. (1983) Reproduction in the iguana *Conolophus subcristatus* on Fernandina Island, Galapagos: clutch size and migration costs. *American Naturalist*, **121**, 757-775.
- Werner, Y.L. (1987) Ecological zoogeography of Saharo-Arabian, Saharan and Arabian reptiles in the sand deserts of southern Israel. *Conference Proceedings*, 272-295.
- Werner, Y.L. (1989) Egg size and egg shape in near-eastern gekkonid lizards. *Israel Journal of Zoology*, **35**, 199-213.
- Werner, Y.L. (1993) The paradoxical tree gecko of Israel. *Dactylus, Journal of the International Gecko Society*, **32**, 29-45.
- Werner, Y.L. (2004) A new species of the *Acanthodactylus pardalis* group (Reptilia: Lacertidae) from Jordan. *Zoology in the Middle East*, **32**, 39-46.
- Werner, Y.L. & Whitaker, A.H. (1978) Observations and comments on the body temperatures of some New Zealand reptiles. *New Zealand Journal of Zoology*, **5**, 375-393.

- Werner, Y.L. & Sivan, N. (1993) Systematics and zoogeography of *Ptyodactylus* (Reptilia: Sauria: Gekkonidae) in the Levant: 1. Biometry of three species in Israel. *Revista Espanola de Herpetologia*, **7**, 47-64.
- Werner, Y.L. & Seifan, T. (2006) Eye size in geckos: asymmetry, allometry, sexual dimorphism, and behavioral correlates. *Journal of Morphology*, **267**, 1486-1500.
- Werner, Y.L. & Ashkenazi, S. (2010) Notes on some Egyptian Lacertidae, including a new subspecies of *Mesalina*, involving the Seligmann effect. *Turkish Journal of Zoology*, **34**, 1-11.
- Western, D. (1974) The distribution, density and biomass density of lizards in a semi-arid environment of northern Kenya. *East African Wildlife Journal*, **12**, 49-62.
- Whitaker, A.H. (1987) The roles of lizards in New Zealand plant reproductive strategies. *New Zealand Journal of Botany*, **25**, 315-328.
- Whitaker, T. & Lyall, J. (2004) *Conservation of lizards in west coast Tai Poutini Conservancy*. Special Report by DOC Science Publishing, Science & Research Unit.
- Whitaker, A.H., Tocher, M. & Blair, T. (2002) *Conservation of lizards in Otago Conservancy, 2002-2007*. Special Report by DOC Science Publishing, Science & Research Unit.
- Whitfield, S.M., Bell, K.E., Philippi, T.S., M. Bolanos, F. Chaves, G., Savage, J.M. & Donnelly, M.A. (2007) Amphibian and reptile declines over 35 years at La Selva, Costa Rica. *Proceedings of the National Academy of Sciences, USA*, **104**, 8352-8356.
- Wiens, J.J. & Slingluff, J.L. (2001) How lizards turn into snakes: a phylogenetic analysis of body-form evolution in anguid lizards. *Evolution*, **55**, 2303-2318.
- Wiens, J.J., Brandley, M.C. & Reeder, T.W. (2006) Why does a trait evolve multiple times within a clade? Repeated evolution of snake-like body form in squamate reptiles. *Evolution*, **60**, 123-141.
- Wilhoft, D.C. (1961) Temperature responses in two tropical Australian skinks. *Herpetologica*, **17**, 109-113.
- Wilhoft, D.C. & Reiter, E.O. (1965) Sexual cycle of the lizard, *Leiopisma fuscum*, a tropical Australian skink. *Journal of Morphology*, **116**, 379-388.
- Wilms, T. & Bohme, W. (2000) A new *Uromastyx* species from south-eastern Arabia, with comments on the taxonomy of *Uromastyx aegyptia* (Forskal, 1775) (Squamata: Sauria: Agamidae). *Herpetozoa*, **13**, 133-148.
- Wilson, S. (2003) *Reptiles of the Southern Brigalow Belt*. WWF Australia.
- Wilson, M.J. & Grillitsch, H. (2009) The herpetofauna of Simi (Dodecanese, Greece) (Amphibia, Reptilia). *Herpetozoa*, **22**, 99-113.
- Winne, C.T. & Keck, M.B. (2004) Daily activity patterns of whiptail lizards (Squamata: Teiidae: *Aspidoscelis*): a proximate response to environmental conditions or an endogenous rhythm? *Functional Ecology*, **18**, 314-321.
- Withers, P.C., Aplin, K.P. & Werner, Y.L. (2000) Metabolism and evaporative water loss of Western Australian geckos (Reptilia: Sauria: Gekkonomorpha). *Australian Journal of Zoology*, **48**, 111-126.
- Wone, B. & Beauchamp, B. (2003) Movement, home range, and activity patterns of the horned lizard, *Phrynosoma mcallii*. *Journal of Herpetology*, **37**, 679-686.
- Woodbury, L.A. (1932) Notes on food habits of three species of lizards from Utah. *Copeia*, **1932**, 13-16.
- Woodhead, J., Eggins, S. & Johnson, R. (1998) Magma genesis in the New Britain island arc: further insights into melting and mass transfer processes. *Journal of Petrology*, **39**, 1641-

1668.

- Woolrich-Pina, G.A., Smith, G.R. & Lemos-Espinal, J.A. (2011) Body temperatures of two species of *Aspidoscelis* from Zapotitlan Salinas, Puebla, Mexico. *Herpetology Notes*, **4**, 387-390.
- Wright, J.L., Bauer, A.M. & Sadler, R.A. (2000) Two new gecko species allied to *Bavayia sauvagii* and *Bavayia cyclura* (Reptilia: Diplodactylidae) from New Caledonia. *Pacific Science*, **54**, 39-55.
- Xua, D.-D. & Ji, X. (2007) Sexual dimorphism, female reproduction and egg incubation in the oriental leaf-toed gecko (*Hemidactylus bowringii*) from southern China. *Zoology*, **110**, 20-27.
- Youmans, T.M. & Grismer, L.L. (2006) A new species of *Cyrtodactylus* (Reptilia: Squamata: Gekkonidae) from the Seribuat Archipelago, West Malaysia. *Herpetological Natural History*, **10**, 61-70.
- Youssef, M.K., Adolph, S.C. & Richmond, J.Q. (2008) Evolutionarily conserved thermal biology across continents: The North American lizard *Plestiodon gilberti* (Scincidae) compared to Asian *Plestiodon*. *Journal of Thermal Biology*, **33**, 308-312.
- Zaldivar-Riveron, A. & de Oca, A.N.-M. (2002) Variation in the rare lizard *Barisia rudicollis* (Wiegmann) (Anguidae) with description of a new species from central Mexico. *Herpetologica*, **58**, 313-326.
- Zamora-Abrego, J.G., Zuniga-Vega, J.J. & de Oca, A.-N. (2007) Variation in reproductive traits within the lizard genus *Xenosaurus*. *Journal of Herpetology*, **41**, 630-637.
- Zamudio, K.R. & Parra-Olea, G. (2000) Reproductive mode and female reproductive cycles of two endemic Mexican horned lizards (*Phrynosoma taurus* and *Phrynosoma braconnieri*). *Copeia*, **2000**, 222-229.
- Zari, T.A. (1991) The influence of body mass and temperature on the standard metabolic rate of the herbivorous desert lizard, *Uromastix microlepis*. *Journal of Thermal Biology*, **16**, 129-134.
- Zari, T.A. (1996) Effects of body mass and temperature on standard metabolic rate of the herbivorous desert lizard *Uromastix philbyi*. *Journal of Arid Environments*, **33**, 457-461.
- Zari, T.A. (1997) Effects of body mass, temperature, and season on resting metabolism of the nocturnal gecko *Hemidactylus flaviviridis*. *Zoology in the Middle East*, **14**, 77-85.
- Zhang, Y.-P. & Ji, X. (2004) The thermal dependence of food assimilation and locomotor performance in southern grass lizards, *Takydromus sexlineatus* (Lacertidae). *Journal of Thermal Biology*, **29**, 45-53.
- Zhang, Y.-P., Du, W.-G. & Zhu, L.-J. (2009) Differences in body size and female reproductive traits between two sympatric geckos, *Gekko japonicus* and *Gekko hokouensis*. *Folia Zoologica*, **58**, 113-122.
- Ziegler, T., Strauch, M., Pes, T., Konas, J., Jirasek, T., Rutz, N., Oberreuter, J. & Holst, S. (2009) First captive breeding of the blue tree monitor *Varanus macraei* Bohme & Jacobs, 2001 at the Plzen and Cologne Zoos. *Biawak*, **3**, 122-133.
- Ziegler, T., Nazarov, R., Orlov, N., Nguyen, T.Q., Vu, T.N., Dang, K.N., Dinh, T.H. & Schmitz, A. (2010) A third new *Cyrtodactylus* (Squamata: Gekkonidae) from Phong Nha-Ke Bang National Park, Truong Son Range, Vietnam. *Zootaxa*, **2413**, 20-36.
- Zinner, H. (1967) Herpetological collection trips to the Lebanon 1965 and 1966. *Israel Journal of Zoology*, **16**, 49-58.
- Znari, M. & El Mouden, E.H. (1997) Sexual dimorphism, reproductive and fat body cycles in

- Bibron's agama (*Agama impalearis*, Boettger, 1874) (Sauria: Agamidae). *Herpetologica*, **53**, 411-422.
- Znari, M. & Nagy, K.A. (1997) Field metabolic rate and water flux in free-living Bibron's agama (*Agama impalearis*, Boettger, 1874) in Morocco. *Herpetologica*, **53**, 81-88.
- Zug, G.R. (1987) Amphibians and reptiles of the Outamba-Kilimi region, Sierra Leone. *Herpetological Association of Africa Journal*, **33**, 1-4.
- Zug, G.R. (1991) The lizards of Fiji: natural history and systematics. *Bishop Museum Bulletin of Zoology*, **2**, 1-136.
- Zug, G.R., Vindum, J.V. & Koo, M.S. (2007) Burmese *Hemidactylus* (Reptilia, Squamata, Gekkonidae): taxonomic notes on tropical Asian *Hemidactylus*. *Proceedings of the California Academy of Sciences*, **58**, 387-405.
- Zug, G.R., Springer, V.G., Williams, J.T. & Johnson, D.G. (1988) The vertebrates of Rotuma and surrounding waters. *Atoll Research Bulletin*, **316**, 1-25.
- Zuniga-Vega, J.J. (2011) Estimating potential reproductive costs in the survival of a xenosaurid lizard. *Herpetological Journal*, **21**, 117-129.
- Zweifel, R.G. (1980) Results of the Archbold Expeditions No. 103. Frogs and lizards from the Huon Peninsula, Papua New Guinea. *Bulletin of the American Museum of Natural History*, **165**, 387-434.
- Zweifel, R.G. & Lowe, C.H. (1966) The ecology of a population of *Xantusia vigilis*, the desert night lizard. *American Museum Novitates*, **2247**, 1-58.

E. Equations used to estimate (\log_{10}) mass from (\log_{10}) snout–vent lengths in clades

Clade	intercept	slope
Agamidae	-4.774	3.073
Amphisbaenia	-5.858	2.943
Anguimorpha	-5.058	3.145
<i>Anolis</i> (Meiri, unpublished)	-4.574	2.94155
Carphodactylidae + Diplodactylidae (Meiri, unpublished)	-4.7159	2.9941
Chamaeleonidae	-3.997	2.68
Eublepharidae (Meiri, unpublished)	-5.201	3.255
Gekkonidae (Meiri, unpublished)	-4.2424	2.761
Gekkonidae + Phyllodactylidae+ Sphaerodactylidae (Meiri, unpublished)	-4.407	2.861
Gerrhosauridae	-4.783	3.085
Iguania	-5.033	3.243
Iguanidae	-4.298	2.972
Lacertidae	-4.543	2.951
leg reduced lizards	-4.207	2.472
Legged Cordylidae	-5.747	3.589
Legged Gymnophthalmidae	-5.178	3.302
legged Anguidae (Meiri, unpublished)	-5.765	3.48
Legged Scincidae	-5.125	3.229
Limbless lizards	-4.207	2.3
<i>Liolaemus</i> , (Pincheira-Donoso et al. 2011)	-4.678	3.097
Phrynosomatidae	-3.855	2.677
<i>Phymaturus</i> , (Pincheira-Donoso et al. 2011)	-5.04	3.323
Polychrotidae	-4.583	2.94
Pygopodidae	-2.039	1.371
Sphaerodactylidae (Meiri, unpublished)	-4.559	2.970
Teiidae	-4.747	3.11
Tropiduridae	-4.216	2.846
Tropiduridae <i>sensu stricto</i> , (Meiri, unpublished)	-3.505	2.521
Varanidae	-5.301	3.235
Xantusiidae	-4.796	3.048

Source is Meiri (2010) unless noted otehrwise.

Meiri, S. (2010) Length–weight allometries in lizards. *Journal of Zoology*, **281**, 218–226.

Pincheira-Donoso, D., Fox, S.F., Scolaro, J.A., Iburgüengoytía, N., Acosta, J.C., Corbalan, V., Medina, M., Boretto, J., Villavicencio, H. J. & Hodgson, D.J. (2011) Body size dimensions

in lizard ecological and evolutionary research: exploring the predictive power of mass estimation equations in two Liolaemidae radiations. *The Herpetological Journal*, **21**, 35-42.

F. The phylogenetic tree

(((((((Leiolepis_reevesii:11.0,(((Moloch_horridus:8.0,(((Ctenophorus_nuchalis:4.0,(((Ctenophorus_isolepis:1.0,Ctenophorus_fordi:1.0):1.0,Ctenophorus_fionni:2.0):1.0,Ctenophorus_ornatus:3.0):1.0):1.0,Ctenophorus_maculosus:5.0):1.0,(((Diporiphora_nobbi:3.0,(Pogona_minor:1.0,Pogona_barbata:1.0):1.0,Pogona_vitticeps:2.0):1.0):1.0,(Amphibolurus_muricatus:1.0,Chlamydosaurus_kingii:1.0):3.0):1.0,Lophognathus_longirostris:5.0):1.0):1.0,Intellagama_lesueurii:7.0):1.0):1.0,Physignathus_cocincinus:9.0):1.0,(((Draco_biaro:1.0,Draco_bourouniensis:1.0):2.0,((Draco_timorensis:1.0,Draco_volans:1.0):1.0,(Draco_palawanensis:1.0,Draco_reticulatus:1.0):1.0):1.0):3.0,(((Sitana_fusca:1.0,Sitana_ponticeriana:1.0,Sitana_sivalensis:1.0):3.0,((Gonocephalus_bellii:1.0,Gonocephalus_bornensis:1.0,Gonocephalus_grandis:1.0):2.0,((Calotes_myataceus:1.0,Calotes_versicolor:1.0):1.0,Calotes_nemorica:2.0):1.0):1.0,(((Japalura_polygonata:1.0,Japalura_swinhoni:1.0):1.0,Japalura_brevipes:2.0):1.0,Japalura_kumaonensis:3.0,Japalura_luei:3.0,Japalura_makii:3.0):2.0):1.0,(((Trapelus_ruderatus:2.0,(Trapelus_mutabilis:1.0,Trapelus_sanguinolentus:1.0):1.0):1.0,Acanthocercus_atricollis:3.0):3.0,(Agama_impalearis:1.0,Agama_agama:1.0):5.0,((Phrynocephalus_mystaceus:4.0,((Phrynocephalus_theobaldi:2.0,(Phrynocephalus_versicolor:1.0,Phrynocephalus_guttatus:1.0):1.0):1.0,Phrynocephalus_helioscopus:3.0):1.0):1.0,(Laudakia_caucasia:1.0,Laudakia_stellio:1.0):4.0):1.0):1.0):3.0):1.0,((Uromastix_ornata:1.0,Uromastix_aegyptia:1.0):1.0,Uromastix_acinthina:2.0):10.0):1.0,(((((((Trioceros_elliotti:1.0,Trioceros_hoehnelii:1.0):1.0,Trioceros_jacksonii:2.0):1.0,Trioceros_quadricornis:3.0):1.0,(Furcifer_cephalolepis:1.0,Furcifer_polleni:1.0):3.0):1.0,Archaius_tigris:5.0):1.0,(Bradypodion_pumilum:1.0,Bradypodion_ventrale:1.0):5.0):1.0,(((Chamaeleo_dilepis:2.0,(Chamaeleo_chamaeleon:1.0,Chamaeleo_calyptratus:1.0):1.0):1.0,Chamaeleo_namachus:3.0):1.0,Chamaeleo_namaquensis:4.0):3.0):1.0,Rhampholeon_marshalli:8.0):5.0):8.0,(((((((Phymaturus_palluma:1.0,Phymaturus_punae:1.0):1.0,(Phymaturus_patagonicus:1.0,Phymaturus_zapalensis:1.0):1.0):4.0,(((Liolaemus_pictus:1.0,Liolaemus_bellii:1.0):1.0,Liolaemus_bibronii:2.0):1.0,Liolaemus_elongatus:3.0):1.0,(Liolaemus_lemniscatus:1.0,Liolaemus_nigromaculatus:1.0):3.0):1.0,(((Liolaemus_boulengeri:1.0,Liolaemus_koslowkyi:1.0):1.0,(Liolaemus_lutzae:1.0,Liolaemus_wiegmannii:1.0):1.0):1.0,(Liolaemus_signifer:1.0,Liolaemus_huacahuasicus:1.0,Liolaemus_andinus:1.0):2.0):1.0,(Liolaemus_lineomaculatus:2.0,(Liolaemus_scolaroii:1.0,Liolaemus_zullyae:1.0):1.0):2.0):1.0):1.0,(Enyalium_perditus:4.0,(Gambelia_wislizenii:1.0,Gambelia_sila:1.0):2.0,((Crotaphytus_collaris:1.0,Crotaphytus_grismeri:1.0):1.0,Crotaphytus_reticulatus:2.0):1.0):1.0):3.0,(((Brachylophus_fasciatus:1.0,Brachylophus_vitiensis:1.0):7.0,(((Ctenosaura_oderhina:1.0,Ctenosaura_bakeri:1.0):1.0,Ctenosaura_pectinata:2.0,Ctenosaura_similis:2.0):1.0,((Conolophus_pallidus:1.0,Conolophus_subcristatus:1.0):1.0,Amblyrhynchus_cristatus:2.0):1.0):4.0,(((Iguana_delicatissima:1.0,Iguana_iguana:1.0):4.0,(((Cyclura_cyclura:1.0,Cyclura_rileyi:1.0):1.0,Cyclura_collei:2.0):1.0,Cyclura_carinata:3.0):1.0,Cyclura_pinguis:4.0):1.0):1.0,((Sauromalus_hispidus:1.0,Sauromalus_varius:1.0):1.0,Sauromalus_ater:2.0):4.0):1.0):1.0,Dipsosaurus_dorsalis:9.0):1.0,((Stenocercus_dumerilii:1.0,Stenocercus_chrysopygus:1.0):6.0,(((Microlophus_albemarlensis:1.0,Microlophus_delanonis:1.0):1.0,Microlophus_habelii:2.0):1.0,(Microlophus_quadrivittatus:1.0,Microlophus_atacamensis:1.0):2.0):3.0,(((Tropidurus_hispidus:1.0,Tropidurus_torquatus:1.0):1.0,Tropidurus_etheridgei:2.0):1.0,Tropidurus_semitaeniatus:3.0):1.0,Tropidurus_spinulosus:4.0):1.0,(Plica_plica:1.0,Plica_umbra:1.0):1.0,Uracentron_flaviceps:2.0):3.0):1.0):4.0):8.0,(((Leiocephalus_eremitus:1.0,Leiocephalus_inaguae:1.0,Leiocephalus_psammodomus:1.0):13.0,(((((((Anolis_acutus:1.0,Anolis_evermanni:1.0):3.0,((Anolis_gundlachi:2.0,(Anolis_krugi:1.0,Anolis_pulchellus:1.0):1.0):1.0,((Anolis_cooki:1.0,Anolis_monensis:1.0):1.0,Anolis_cristatellus:2.0):1.0):1.0):2.0,((Anolis_wattsii:1.0,Anolis_pogus:1.0):4.0,((Anolis_bimaculatus:1.0,Anolis_gingivinus:1.0):3.0,((Anolis_sabanus:1.0,Anolis_lividus:1.0):1.0,Anolis_oculatus:2.0):1.0,Anolis_leachii:3.0):1.0):1.0):3.0,(Anolis_sagrei:8.0,((Anolis_nebulosus:5.0,(Anolis_aquaticus:4.0,(((Anolis_lionotus:1.0,Anolis_oxyllophus:1.0):1.0,Anolis_limifrons:2.0):1.0,((Anolis_cupreus:1.0,Anolis_polylepis:1.0):1.0,(Anolis_intermedius:1.0,Anolis_tropidolepis:1.0):1.0):1.0):1.0):2.0,(((((((Anolis_conspersus:1.0,Anolis_grahami:1.0):1.0,Anolis_garmani:2.0):1.0,Anolis_opalinus:3.0):1.0,Anolis_reconditus:4.0):1.0,Anolis_lineatopus:5.0):1.0,Anolis_valencienni:6.0):1.0):1.0):1.0,(Anolis_carolinensis:1.0,Anolis_longiceps:1.0):9.0):1.0,Anolis_cuvieri:11.0):1.0,(((Anolis_bonairensis:1.0,Anolis_lucia:1.0):3.0,(Anolis_trinitatis:3.0,((Anolis_roquet:1.0,Anolis_aeneus:1.0):1.0,Anolis_richardii:2.0):1.0):1.0,Anolis_agassizi:5.0):7.0):1.0,(Basiliscus_basiliscus:1.0,Basiliscus_vittatus:1.0):12.0):1.0):4.0,(((Uta_stansburiana:1.0,Uta_palmeri:1.0):15.0,(((((((Sceloporus_arenicolus:1.0,Sceloporus_graciosus:1.0):9.0,(((((((Sceloporus_virgatus:3.0,((Sceloporus_undulatus:1.0,Sceloporus_woodi:1.0):1.0,Sceloporus_consobrinus:2.0):1.0):1.0,Sceloporus_occidentalis:4.0):1.0,Sceloporus_olivaceus:5.0):1.0,((Sceloporus_slevini:1.0,Sceloporus_aeneus:1.0):1.0,Sceloporus_scalaris:2.0):4.0):1.0,(Sceloporus_clarkii:6.0,((Sceloporus_torquatus:1.0,Sceloporus_jarrovii:1.0):3.0,((Sceloporus_omiltemanus:1.0,Sceloporus_mucronatus:1.0):2.0,((Sceloporus_cyanogenys:1.0,Sceloporus_serri:1.0):1.0,Sceloporus_poinsettii:2.0):1.0):1.0):1.0,Sceloporus_grammicus:5.0):1.0):1.0,Sceloporus_orcutti:1.0,Sceloporus_magister:1.0):7.0):1.0,((Sceloporus_malachiticus:1.0,Sceloporus_formosus:1.0):1.0,Sceloporus_horridus:2.0):7.0):1.0):1.0,Sceloporus_gadoviae:11.0):1.0,Sceloporus_merriami:12.0):1.0,(Sceloporus_angustus:1.0,Sceloporus_utiformis:1.0):12.0):1.0,(Sceloporus_cozumelae:1.0,Sceloporus_variablis:1.0):13.0):1.0,(Urosaurus_bicarinatus:3.0,((Urosaurus_clarionensis:1.0,Urosaurus_ornatus:1.0):1.0,Urosaurus_graciosus:2.0):1.0):12.0):1.0,(((Uma_exsul:1.0,Uma_notata:1.0):4.0,(((Holbrookia_maculata:1.0,Holbrookia_propinqua:1.0):1.0,Holbrookia_lacerata:2.0):1.0,Cophosaurus_texasus:3.0):1.0,Callisaurus_draconoides:4.0):1.0):1.0,(((((((Phrynosoma_blainvillii:1.0,Phrynosoma_coronatum:1.0):1.0,(Phrynosoma_mcallii:1.0,Phrynosoma_platyrhinos:1.0):1.0):1.0,Phrynosoma_solare:3.0):1.0,((Phrynosoma_ditmarsi:1.0,Phrynosoma_hernandesi:1.0):1.0,Phrynosoma_douglassii:2.0):1.0,Phrynosoma_modestum:3.0):1.0):1.0,Phrynosoma_cornutum:5.0):1.0):1.0):1.0,Polychrus_acutirostris:20.0):1.0,(((((((Ophisaurus_attenuatus:1.0,Ophisaurus_compressus:1.0,Ophisaurus_ventralis:1.0):1.0,Dopasia_gracilis:2.0):1.0,(Anguis_cephallonica:1.0,Anguis_fragilis:1.0):1.0,Pseudopus_apodus:2.0):1.0):2.0,(((Mesaspis_monticola:1.0,Mesaspis_juarezi:1.0):1.0,Mesaspis_gadovii:2.0):1.0,(Gerrhonotus_infernalis:2.0,(Barisia_herrerae:1.0,Barisia_imbricata:1.0):1.0):1.0,(Elgaria_multicarinata:1.0,Elgaria_coerulea:1.0):3.0):1.0):1.0,(Diploglossus_millepunctatus:2.0,(Diploglossus_pleii:1.0,Ophiodes_striatus:1.0):1.0):4.0):1.0,Anniella_pulchra:7.0):1.0,(Heloderma_horridum:1.0,Heloderma_suspectum:1.0):7.0):1.0,Xenosaurus_grandis:9.0):2.0,(((Varanus_niloticus:1.0,Varanus_ornatus:1.0):1.0,Varanus_griseus:2.0):1.0,Varanus_albigularis:3.0):7.0,(((Varanus_bengalensis:1.0,Varanus_flavescens:1.0):2.0,((Varanus_nuchalis:1.0,Varanus_salvator:1.0):1.0,Varanus_rudicollis:2.0):1.0):5.0,(((Varanus_glauerti:1.0,Varanus_tristis:1.0):1.0,Varanus_scalaris:2.0):1.0,((

Varanus_kingorum:1.0,Varanus_acanthurus:1.0):1.0,(Varanus_gilleni:1.0,Varanus_caudolineatus:1.0):1.0):4.0,((((Varanus_gouldii:1.0,Varanus_panoptes:1.0):1.0,Varanus_rosebergi:2.0):1.0,Varanus_giganteus:3.0):1.0,Varanus_mertensi:4.0):1.0,Varanus_spenceri:5.0):1.0,(Varanus_komodoensis:1.0,Varanus_varius:1.0):5.0):1.0):1.0,(((Varanus_beccarii:3.0,((Varanus_macraei:1.0,Varanus_prasinus:1.0):1.0,Varanus_boehmei:2.0):1.0):1.0,(Varanus_caerulivirens:3.0,(Varanus_lirungensis:2.0,(Varanus_cerambonensis:1.0,Varanus_melinus:1.0):1.0):1.0):1.0,Varanus_mabitang:5.0):4.0):1.0):11.0):1.0,(((Trogonophis_wiegmanni:3.0,((Bipes_biporus:1.0,Bipes_canaliculatus:1.0):1.0):1.0,Bipes_tridactylus:2.0):1.0):10.0,(((((((Takydromus_amurensis:1.0,Takydromus_sauteri:1.0):4.0,(((Takydromus_septentrionalis:1.0,Takydromus_stejnegeri:1.0):1.0,Takydromus_toyamai:2.0):2.0,(((Takydromus_formosanus:1.0,Takydromus_wolteri:1.0):1.0,(Takydromus_luyeanus:1.0,Takydromus_viridipunctatus:1.0):1.0):1.0,Takydromus_hsuehshanensis:3.0):1.0):1.0):1.0,(Takydromus_dorsalis:1.0,Takydromus_smaragdinus:1.0):5.0):1.0,Takydromus_sexlineatus:7.0):1.0,Zootoca_vivipara:8.0):2.0,((((Algyroides_moreoticus:1.0,Algyroides_nigropunctatus:1.0):2.0,((Algyroides_fitzingeri:1.0,Algyroides_marchi:1.0):1.0,Dinarolacerta_mosorensis:2.0):1.0):3.0,((Phoenicolacerta_laevis:1.0,Phoenicolacerta_troodica:1.0):4.0,(((Podarcis_gaigeae:1.0,Podarcis_milensis:1.0):1.0,Podarcis_melisellensis:2.0):1.0,Podarcis_tauricus:3.0):1.0,((Podarcis_filfolensis:2.0,(Podarcis_waglerianus:1.0,Podarcis_raffoneae:1.0):1.0):1.0,((Podarcis_lilfordi:1.0,Podarcis_pityusensis:1.0):1.0,Podarcis_tiligerta:2.0):1.0):1.0,Podarcis_muralis:1.0,Podarcis_siculus:1.0):3.0,Podarcis_bocagei:1.0,Podarcis_carbonelli:1.0,Podarcis_hispanicus:1.0):3.0,(Podarcis_cretensis:1.0,Podarcis_erhardii:1.0,Podarcis_peloponnesiacus:1.0):3.0):1.0):2.0,(((Iberolacerta_monticola:2.0,Iberolacerta_aranica:1.0,Iberolacerta_aurelioi:1.0,Iberolacerta_bonnali:1.0):1.0):1.0,(Iberolacerta_cyreni:1.0,Iberolacerta_horvathi:1.0):2.0):1.0,(Archaeolacerta_bedriagae:2.0,(Dalmatolacerta_oxyccephala:1.0,Hellenolacerta_graeca:1.0):1.0):2.0):3.0,((Parvilacerta_parva:2.0,(Anatololacerta_anatolica:1.0,Anatololacerta_oertzeni:1.0):1.0):4.0,(Darevskia_chlorogaster:1.0,Darevskia_derjugini:1.0,Darevskia_parvula:1.0,Darevskia_portschinskii:1.0,Darevskia_praticola:1.0,Darevskia_raddei:1.0,Darevskia_rostombekovi:1.0,Darevskia_rudis:1.0,Darevskia_saxicola:1.0,Darevskia_unisexualis:1.0,Darevskia_valentini:1.0):5.0,((Timon_lemidus:1.0,Timon_pater:1.0):4.0,(((Lacerta_agilis:1.0,Lacerta_viridis:1.0):1.0,Lacerta_trilineata:2.0):1.0,Lacerta_strigata:3.0):1.0,Lacerta_schreiberi:4.0):1.0):1.0):1.0,(((Scelarcis_perspicillata:1.0,Teira_dugesii:1.0):1.0,Atlantolacerta_andreaskyi:2.0):7.0):1.0,(((Meroles_suborbitalis:3.0,((Meroles_cuneirostris:1.0,Meroles_anchietae:1.0):1.0,Ichnotropis_squamulosa:2.0):1.0):1.0,Ichnotropis_capensis:4.0):1.0,(Pedioplanis_burchelli:1.0,Pedioplanis_namaquensis:1.0):4.0):1.0,((Nucras_lalandii:1.0,Nucras_taeniolata:1.0,Nucras_tessellata:1.0):1.0,Heliobolus_lugubris:2.0):4.0):2.0,(((Mesalina_olivieri:1.0,Mesalina_pasteuri:1.0):1.0,(Mesalina_rubropunctata:1.0,Mesalina_guttulata:1.0):1.0):1.0,Ophisops_elegans:3.0):4.0,((((Acanthodactylus_spinicauda:1.0,Acanthodactylus_pardalis:1.0):1.0,Acanthodactylus_erythrurus:2.0):2.0,(Acanthodactylus_schmidti:3.0,((Acanthodactylus_beershebensis:1.0,Acanthodactylus_boskianus:1.0):1.0,Acanthodactylus_schreiberi:2.0):1.0):1.0,Acanthodactylus_scutellatus:5.0):1.0,((((Eremias_strauchii:1.0,Eremias_velox:1.0):1.0,Eremias_arguta:2.0,Eremias_persica:2.0):1.0,Eremias_pleskei:3.0):1.0,Eremias_grammica:4.0):1.0,Eremias_intermedia:5.0,Eremias_lineolata:5.0,Eremias_nigrocellata:5.0,Eremias_nikolskii:5.0,Eremias_regeli:5.0):1.0):1.0):3.0,((Psammmodromus_hispanicus:1.0,Psammmodromus_blanci:1.0):1.0,Psammmodromus_algirus:2.0):4.0,((((Gallotia_bravoana:1.0,Gallotia_caesaris:1.0):1.0,Gallotia_galloti:2.0):1.0,(Gallotia_intermedia:1.0,Gallotia_simonyi:1.0):2.0):1.0,Gallotia_stehlini:4.0):1.0,Gallotia_atlantica:5.0):1.0):6.0):1.0):2.0,((((((((Aspidoscelis_inornata:1.0,Aspidoscelis_exsanguiis:1.0,Aspidoscelis_sonorae:1.0,Aspidoscelis_uniparens:1.0,Aspidoscelis_velox:1.0):1.0,Aspidoscelis_sexlineata:2.0):1.0,((Aspidoscelis_gularis:1.0,Aspidoscelis_laredoensis:1.0,Aspidoscelis_scalaris:1.0):1.0,Aspidoscelis_burti:2.0):1.0):4.0,((((Aspidoscelis_deppei:1.0,Aspidoscelis_lineatissima:1.0):1.0,Aspidoscelis_guttata:1.0,Aspidoscelis_cozumelae:3.0):1.0,(Aspidoscelis_ceralbensis:1.0,Aspidoscelis_hyperythra:1.0):3.0):1.0,(Aspidoscelis_marmorata:1.0,Aspidoscelis_tigris:1.0):4.0):1.0,(Aspidoscelis_dixoni:1.0,Aspidoscelis_neomexicana:1.0,Aspidoscelis_tessellata:1.0):5.0):1.0):1.0,Ameivula_ocellifera:8.0):1.0,((Cnemidophorus_murinus:1.0,Cnemidophorus_vanzoi:1.0):2.0,((Cnemidophorus_gramivagus:1.0,Cnemidophorus_cryptus:1.0,Cnemidophorus_lemniscatus:1.0):1.0,Cnemidophorus_arubensis:2.0):1.0):1.0,((Kentropyx_calcarata:1.0,Kentropyx_pelviceps:1.0):1.0,Kentropyx_striata:2.0):2.0):5.0):1.0,(Ameiva_ameiva:5.0,(Holcosus_festivus:1.0,Holcosus_quadrilineatus:1.0):3.0,((Ameiva_erythrocephala:1.0,Ameiva_fusca:1.0):1.0,Ameiva_plei:2.0):1.0,(Ameiva_poliops:1.0,Ameiva_wetmorei:1.0):1.0,Ameiva_exsul:2.0):1.0):1.0):5.0):1.0,(Contomastix_lacertoides:1.0,Contomastix_vacariensis:1.0):1.0):1.0,(Teius_oculatus:1.0,Teius_teyou:1.0):11.0):1.0,Tupinambis_teguixin:13.0):1.0,(Leptosoma_rugiceps:5.0,((Cercosaura_schreibersii:2.0,Euspondylus_chasqui:1.0,Potamites_eupleopus:1.0):1.0):1.0,Riama_shrevei:3.0):1.0,Gymnophthalmus_speciosus:4.0):1.0):9.0):1.0):8.0):1.0,(((Acontias_gariepensis:1.0,Acontias_kgalagadi:1.0):14.0,(((Cryptoblepharus_egeriae:1.0,Cryptoblepharus_eximius:1.0,Cryptoblepharus_no_vocaledonicus:1.0,Cryptoblepharus_poecilopleurus:1.0,Cryptoblepharus_renschi:1.0):2.0,((Afroablepharus_annobonensis:1.0,Panaspis_wahlbergi:1.0):1.0,(Panaspis_kitsoni:1.0,Panaspis_nimbaensis:1.0):1.0):1.0):11.0,(((Sphenomorphus_indicus:1.0,Sphenomorphus_maculatus:1.0):4.0,((Scincella_lateralis:3.0,(Insulasaurus_traanorum:1.0,Insulasaurus_arborens:1.0):1.0,(Lipinia_rouxi:1.0,Lipinia_leptosoma:1.0,Lipinia_macrotympanum:1.0,Lipinia_rabori:1.0):1.0):1.0,Larutia_seribuatensis:4.0):1.0):4.0,((((Eremiascincus_richardsonii:1.0,Hemiergis_peronii:1.0):1.0,(Coeranoscincus_reticulatus:1.0,Eulamprus_brachyosoma:1.0,Eulamprus_quoyii:1.0,Eulamprus_tympanum:1.0):1.0):4.0,(((Ctenotus_brooksi:2.0,(Ctenotus_taeniolatus:1.0,Ctenotus_robustus:1.0):1.0):1.0,Ctenotus_pantherinus:3.0):1.0,Ctenotus_lancelini:4.0):1.0,(Lerista_bougainvillii:4.0,(Lerista_punctatovittata:3.0,(Lerista_labialis:2.0,(Lerista_borealis:1.0,Lerista_xanthura:1.0):1.0):1.0):1.0):1.0,Parvosцинus_sisoni:7.0):1.0,((Sphenomorphus_scutatus:1.0,Sphenomorphus_cranei:1.0):1.0,Sphenomorphus_cincinnati:2.0):6.0):1.0):5.0,((Gongylomorphus_bojerii:6.0,((((Chalcides_coeruleopunctatus:1.0,Chalcides_sexlineatus:1.0):1.0,Chalcides_viridanus:2.0):1.0,Chalcides_ocellatus:3.0):1.0,(Chalcides_sepsoides:1.0,Chalcides_bedriagai:1.0):3.0):1.0,Chalcides_chalcides:5.0):1.0):1.0,(Janetaescincus_braueri:1.0,Pamelaescincus_gardineri:1.0):6.0):5.0,(((Scincus_mitrans:1.0,Scincus_scincus:1.0):1.0,Eumeces_schneideri:2.0):9.0,(((Plestiodon_stimpsonii:1.0,Plestiodon_okadae:1.0):8.0,((((Plestiodon_fasciatus:1.0,Plestiodon_septentrionalis:1.0):1.0,Plestiodon_tetragrammus:2.0):1.0,(Plestiodon_inexpectatus:1.0,Plestiodon_laticeps:1.0):2.0):1.0,Plestiodon_obsoletus:4.0):1.0,Plestiodon_anthracinus:5.0):1.0,(Plestiodon_egregius:1.0,Plestiodon_reynoldsi:1.0):5.0):1.0,Plestiodon_skiltonianus:7.0):1.0,(Plestiodon_copei:1.0,Plestiodon_lynxe:1.0):7.0):1.0):1.0,Plestiodon_kishinouyei:10.0):1.0,(((Brachymeles_minimus:1.0,Brachymeles_cebuensis:1.0):1.0,Brachymeles_tridactylus:2.0):1.0,Brachymeles_taylori:3.0):9.0):2.0,((((Carlia_rostralis:1.0,Carlia_tetradactyla:1.0):1.0,(Carlia_rhomboidalis:1.0,Carlia_rubrigularis:1.0):1.0):1.0,Carlia_longipes:3.0):6.0,(Menetia_greyii:8.0,Leiopisma_telfairii:8.0,((((Emoia_tongana:1.0,Emoia_concolor:1.0):3.0,((Emoia_trossula:1.0,Emoia_aneityumensis:1.0):1.0,Emoia_nigromarginata:2.0):1.0,Emoia_parkeri:3.0):1.0):1.0,(Emoia_loyalti:1.0,Emoia_sanfordi:1.0):4.0):1.0,((Emoia_adspersa:1.0,Emoia_atrocostata:1.0):1.0,Emoia_maculata:2.0,Emoia_ponapea:2.0):1.0,Emoia_impar:3.0):3.0):1.0,Emoia_campbelli:7.0,Emoia_isolata:7.0,Emoia_nativitatis:7.0,Emoia_reimschi:7.0,Emoia_schmidti:7.0,Emoia_slevini:7.0,Emoia_sorex:7.0,Emoia_taumakoensis:7.0):1.0):1.0.

0):3.0,((Dasia_olivacea:10.0,(((Copeoglossum_nigropunctatum:6.0,((Brasiliscincus_agilis:1.0,Brasiliscincus_heathi:1.0):4.0,((Varzea_bistria:3.0,((Marisora_unimarginata:1.0,Mabuya_mabouya:1.0):1.0,Aspronema_cochabambae:2.0):1.0):1.0,Psychosaura_macrorhyncha:4.0):1.0):1.0):1.0,((Trachylepis_aurata:1.0,Trachylepis_vittata:1.0):6.0):1.0,((Chioninia_coctei:1.0,Chioninia_spinalis:1.0):1.0,Chioninia_vallantii:2.0):6.0):1.0,((Trachylepis_sparsa:5.0,Trachylepis_lavrambo:5.0,Trachylepis_buettneri:5.0,(((Trachylepis_spilogaster:1.0,Trachylepis_striata:1.0):1.0,Trachylepis_varia:2.0):1.0,((Trachylepis_affinis:1.0,Trachylepis_quinquetaeniata:1.0):2.0):1.0,((Trachylepis_sechellensis:1.0,Trachylepis_wrightii:1.0):1.0,Trachylepis_maculilabris:2.0):2.0):1.0):4.0):1.0):1.0,((Eutropis_multifasciata:2.0,((Eutropis_carinata:1.0,Eutropis_longicaudata:1.0):1.0):9.0):1.0):1.0,((((((((Marmorosphax_taoam:1.0,Marmorosphax_boulinda:1.0):1.0,Marmorosphax_tricolor:2.0):1.0,Kanakysaurus_viviparus:3.0):1.0,(Lioscincus_greeri:1.0,Lioscincus_nigrofasciolatum:1.0):3.0):3.0,((Tropidoscincus_boreus:1.0,Tropidoscincus_variabilis:1.0):5.0,(((Caledoniscincus_austrocaledonicus:1.0,Caledoniscincus_haplorhinus:1.0):1.0,Caledoniscincus_festivus:2.0):1.0,Caledoniscincus_orestes:3.0):1.0,(((Caledoniscincus_aquiloniensis:1.0,Caledoniscincus_terma:1.0):1.0,Caledoniscincus_chazeaui:2.0):1.0,Caledoniscincus_atropunctatus:3.0):1.0):1.0,((Caledoniscincus_renevieri:1.0,Caledoniscincus_auratus:1.0):4.0):1.0):1.0,(((Celaticiscincus_euryotis:1.0,Celaticiscincus_similis:1.0):1.0,Lioscincus_steindachneri:2.0):1.0,(Sigaloseps_deplanchei:1.0,Sigaloseps_ruficauda:1.0):2.0):4.0):1.0,(((Nannoscincus_greeri:1.0,Nannoscincus_mariei:1.0):1.0,((Nannoscincus_gracilis:1.0,Nannoscincus_garrulus:1.0):1.0):1.0,Nannoscincus_maccoyi:3.0):5.0):1.0,(Lioscincus_novaeaeoloniae:5.0,(((Oligosoma_notosaurus:1.0,Oligosoma_stenotis:1.0):1.0,Oligosoma_nigriplantare:2.0):1.0,(Oligosoma_fallai:1.0,Oligosoma_homalonotum:1.0):2.0):1.0,Oligosoma_lichenigera:4.0):1.0,Lioscincus_maruia:5.0):4.0):1.0,(((Pseudemoia_entrecasteauxii:1.0,Pseudemoia_pagenstecheri:1.0):1.0,Pseudemoia_rawlinsoni:2.0):1.0,Pseudemoia_spenceri:3.0):1.0,(Morethia_boulengeri:1.0,Morethia_obscura:1.0):3.0,(Bassiana_dupperreyi:1.0,Bassiana_trilineata:1.0):3.0):1.0,(Niveoscincus_metallicus:1.0,Niveoscincus_coventryi:1.0):4.0):5.0):1.0,(((Ablepharus_kitaibelii:1.0,Ablepharus_ruepellii:1.0):1.0,Asymblepharus_sikimmensis:2.0):1.0,(Lampropholis_delicata:1.0,Lampropholis_olichenoti:1.0):2.0):8.0):1.0,(((Tribolonotus_ponceleti:1.0,Tribolonotus_pseudoponceleti:1.0):1.0,Tribolonotus_brongersmai:2.0):1.0,Tribolonotus_schmidti:3.0):1.0,Tribolonotus_blanchardi:4.0):4.0,((((Egernia_hosmeri:1.0,Hemispheeriodon_gerrardii:1.0):1.0,Egernia_kingii:2.0):1.0,(Egernia_cunninghami:1.0,Egernia_striolata:1.0):2.0):1.0,(((Cyclodomorphus_michaeli:1.0,Hemispheeriodon_gerrardii:1.0):1.0,Cyclodomorphus_casuarinae:2.0,Cyclodomorphus_celatus:2.0,Cyclodomorphus_melanops:2.0):1.0,Tiliqua_adelaidensis:3.0,Tiliqua_nigrolutea:3.0,Tiliqua_occupitalis:3.0,Tiliqua_rugosa:3.0,Tiliqua_scincoides:3.0):1.0):1.0,(((Liopholis_kintorei:1.0,Liopholis_striata:1.0):1.0,Liopholis_inornata:2.0):1.0,(Liopholis_whitii:1.0,Liopholis_modesta:1.0):2.0):2.0):1.0,Lissolalepis_coventryi:6.0):1.0,Corucia_zebrata:7.0):1.0):4.0):1.0):1.0,Ophiomoropus_punctatissimus:14.0):1.0):1.0,(((Xantusia_henshawi:4.0,Xantusia_riversiana:3.0,((Xantusia_vigilis:1.0,Xantusia_wigginsi:1.0):1.0,Xantusia_arizonae:2.0):1.0):1.0):1.0,Lepidophyma_flavimaculatum:5.0):1.0,(((Hemicordylus_capensis:1.0,Pseudocordylus_melanotus:1.0):1.0,Smaug_giganteus:2.0):1.0,Gerrhosaurus_skooigi:3.0):3.0):10.0):8.0):1.0,(((Phyllurus_platurus:1.0,Saltuarius_cornutus:1.0):2.0,((Nephrurus_levis:1.0,Underwoodisaurus_millii:1.0):1.0,Carphodactylus_laevis:2.0):1.0):2.0,((Delma_fraseri:1.0,Delma_grayii:1.0):3.0,((Pygopus_lepidopus:2.0,(Aprasia_repens:1.0,Aprasia_pulchella:1.0):1.0):1.0,Lialis_burtonis:3.0):1.0):5.0,((((Strophurus_elderi:3.0,((Strophurus_intermedius:1.0,Strophurus_williamsi:1.0):1.0,Strophurus_ciliaris:2.0):1.0):1.0,(Diplodactylus_tessellatus:1.0,Lucasmus_damaeum:1.0):3.0):1.0,Hesperoedura_reticulata:5.0):1.0,(Amalosia_lesueurii:3.0,((Oedura_monilis:1.0,Oedura_tryoni:1.0):1.0,Oedura_castelnaui:2.0):1.0):3.0):1.0,Crenadactylus_ocellatus:7.0):2.0,((((((((Bavayia_cyclura:1.0,Bavayia_crassicolis:1.0):1.0,Bavayia_montana:2.0):1.0,Bavayia_ornata:1.0,Bavayia_septuiclavis:1.0):2.0):1.0,((Bavayia_pulchella:1.0,Bavayia_sauvagii:1.0):1.0,((Bavayia_exsuccida:1.0,Bavayia_geitaini:1.0):1.0):2.0):1.0,(((Rhacodactylus_trachyrhynchus:1.0,Rhacodactylus_auriculatus:1.0):1.0,Rhacodactylus_leachianus:2.0):2.0,((Eurydactylodes_agricolae:1.0,Eurydactylodes_vieillardii:1.0):1.0,Eurydactylodes_symmetricus:2.0):1.0,Mniarogekko_chahoua:3.0):1.0):1.0,(Correlophus_ciliatus:1.0,Correlophus_sarasinorum:1.0):5.0):1.0,(((Dierogekko_nehoueensis:1.0,Dierogekko_validiclavus:1.0):1.0,Dierogekko_insularis:2.0):1.0,Dierogekko_poumensis:3.0):1.0,Oedodera_marmorata:4.0):3.0):1.0,(Tukutuku_rakiurae:1.0,Toropuku_stephensi:1.0):7.0):1.0):9.0,(((Coleonyx_elegans:3.0,((Coleonyx_brevis:1.0,Coleonyx_variegatus:1.0):1.0,Coleonyx_reticulatus:2.0):1.0):1.0,Aeluroscalabotes_felinus:4.0):1.0,(((Goniurosaurus_hainanensis:1.0,Goniurosaurus_araneus:1.0):1.0,(Goniurosaurus_kuroiwae:1.0,Goniurosaurus_splendens:1.0):1.0):1.0,(Eublepharis_macularius:1.0,Eublepharis_turcmenicus:1.0):2.0):2.0):13.0,((((Stenodactylus_doriae:2.0,(Mediodactylus_russowii:1.0,Mediodactylus_kotschyi:1.0):1.0):1.0,((Cyrtopodion_caspium:1.0,Cyrtopodion_fedtschenkoi:1.0):1.0,Crossobamon_eversmanni:2.0):1.0):12.0,(((Altophylax_loricatus:1.0,Altophylax_szczerebaki:1.0):1.0,Altophylax_laevis:2.0,Altophylax_pipiens:2.0):12.0,(((Ptenopus_garrulus:11.0,(((Chondrodactylus_angulifer:1.0,Pachydactylus_mariquensis:1.0):6.0,(Lygodactylus_klugei:6.0,((Phelsuma_parkeri:2.0,(Phelsuma_astriata:1.0,Phelsuma_sundbergi:1.0):1.0):3.0,(((Phelsuma_robertmertensi:1.0,Phelsuma_nigra:1.0):1.0,Phelsuma_comorensis:2.0):1.0,(Phelsuma_nigristriata:1.0,Phelsuma_dubia:1.0):2.0):2.0,Phelsuma_andamanense:5.0,(((Phelsuma_borbonica:1.0,Phelsuma_cepiana:1.0):1.0,Phelsuma_rosagularis:2.0):1.0,(Phelsuma_inexpectata:1.0,Phelsuma_ornata:1.0):1.0,Phelsuma_guimbeaui:2.0):1.0):1.0,Phelsuma_guentheri:4.0):1.0):1.0):1.0,(Christinus_guentheri:1.0,Christinus_marmoratus:1.0):7.0):1.0,Urocotyledon_inexpectata:9.0):1.0,(Ailuroonyx_sechellensis:1.0,Ailuroonyx_tachyscopaeus:1.0,Ailuroonyx_trachygaster:1.0):9.0):1.0):1.0,(Perochirus_ateles:1.0,Gehyra_variegata:1.0):11.0):1.0,((((Luperosaurus_macgregori:1.0,Lepidodactylus_vanuatuensis:1.0):1.0,Pseudogekko_smaragdinus:2.0):1.0,(Lepidodactylus_euaensis:1.0,Lepidodactylus_manni:1.0):2.0):1.0,(Lepidodactylus_listeri:1.0,Lepidodactylus_lugubris:1.0):3.0):1.0,Lepidodactylus_herrei:5.0):2.0,((((Gekko_hokouensis:1.0,Gekko_shibatai:1.0):1.0,Gekko_vertebralis:2.0):1.0,(Gekko_japonicus:1.0,Gekko_yakuensis:1.0):2.0):1.0,(Gekko_gecko:1.0,Gekko_smithii:1.0):3.0):1.0,Ptychozoon_nicobarensis:5.0):1.0,(Gekko_ernstkeileri:1.0,Gekko_athymus:1.0):5.0):1.0):1.0,Heteronotia_binoei:8.0):5.0):1.0,((Cyrtodactylus_leeigrismeri:5.0,Cyrtodactylus_laevigatus:5.0,Cyrtodactylus_jarakensis:5.0,(((Cyrtodactylus_redimiculus:2.0,(Cyrtodactylus_seribuatensis:1.0,Cyrtodactylus_tioanensis:1.0):1.0):1.0,Cyrtodactylus_sadlei:3.0):1.0,Cyrtodactylus_peguensis:4.0):1.0):1.0,((((Hemidactylus_brookii:1.0,Hemidactylus_frenatus:1.0):1.0,Hemidactylus_flaviviridis:2.0):1.0,Hemidactylus_maculatus:3.0):1.0,Hemidactylus_bowringii:4.0):1.0,(Hemidactylus_mabouia:1.0,Hemidactylus_bouvieri:1.0):1.0,(Hemidactylus_turcicus:1.0,Hemidactylus_dracaenaculus:1.0):1.0):3.0):1.0):8.0):1.0,(Cnemaspis_affinis:1.0,Cnemaspis_limi:1.0,Cnemaspis_modigliani:1.0,Cnemaspis_monachorum:1.0,Cnemaspis_perhentianensis:1.0,Cnemaspis_whittonorum:1.0):14.0):1.0,((Haemodracon_riebeckii:1.0,Haemodracon_trachyrhinus:1.0):8.0,((Ptyodactylus_guttatus:1.0,Ptyodactylus_oudrii:1.0,Ptyodactylus_ragazzii:1.0):7.0,(((Tarentola_deserti:2.0,(Tarentola_angustimentalis:1.0,Tarentola_mauritanica:1.0):1.0):4.0,((Tarentola_annularis:4.0,(Tarentola_darwini:2.0,(Tarentola_delalandii:1.0,Tarentola_gomerensis:1.0):1.0):1.0,Tarentola_chazaliae:3.0):1.0):1.0,Tarentola_botteri:5.0):1.0):1.0,(((Phyllodactylus_bugastrolepis:1.0,Phyllodactylus_lanei:1.0):1.0,(Homonota_darwini:1.0,Homonota_fasciata:1.0):1.0):1.0,((Gymnodactylus_amarali:1.0,Gymnodactylus_geckoides:1.0):1.0,Phyllolopezus_pollicaris:2.0):1.0):4.0):1.0):1.0):7.0):1.0,((Sphaerodactylus

s_vincenti:2.0,Sphaerodactylus_kirbyi:2.0,Sphaerodactylus_argivus:2.0,(Sphaerodactylus_nicholsi:1.0,Sphaerodactylus_roosevelti:1.0):1.0):2.0,(((Gonatodes_vittatus:1.0,Gonatodes_albogularis:1.0):1.0,Gonatodes_daudini:2.0):1.0,Gonatodes_ocellatus:3.0):1.0):2.0,(((Teratoscincus_s_cincus:1.0,Teratoscincus_microlepis:1.0):3.0,(Quedenfeldtia_trachylepharus:3.0,((Aristelliger_georgeensis:1.0,Aristelliger_praesignis:1.0):1.0,Aristelliger_barbouri:2.0):1.0):1.0,((Pristurus_rupestris:1.0,Pristurus_abdelkuri:1.0):2.0,((Pristurus_insignis:1.0,Pristurus_insignoides:1.0):1.0,(Pristurus_guichardi:1.0,Pristurus_sokotranus:1.0):1.0):1.0):2.0):1.0):1.0):6.0);

G. The reference used for each species

Family	species	Source for phylogenetic relationship
Agamidae	<i>Acanthocercus atricollis</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Agama agama</i>	Goncalves et al. 2012
Agamidae	<i>Agama impalearis</i>	Geniez et al. 2011, Mediannikov et al. 2012
Agamidae	<i>Amphibolurus muricatus</i>	Hugall et al. 2008
Agamidae	<i>Calotes mystaceus</i>	Zug et al. 2006
Agamidae	<i>Calotes nemoricola</i>	Zug et al. 2006, Taxonomy
Agamidae	<i>Calotes versicolor</i>	Schulte and Moreno-Roark 2010
Agamidae	<i>Chlamydosaurus kingii</i>	Melville et al. 2011
Agamidae	<i>Ctenophorus fionni</i>	Schulte and Moreno-Roark 2010
Agamidae	<i>Ctenophorus fordi</i>	Doughty et al. 2007
Agamidae	<i>Ctenophorus isolepis</i>	Doughty et al. 2007
Agamidae	<i>Ctenophorus maculosus</i>	Doughty et al. 2007
Agamidae	<i>Ctenophorus nuchalis</i>	Doughty et al. 2007
Agamidae	<i>Ctenophorus ornatus</i>	Melville et al. 2001
Agamidae	<i>Diporiphora nobbi</i>	Hugall et al. 2008
Agamidae	<i>Draco biaro</i>	McGuire & Heang 2001
Agamidae	<i>Draco bourouniensis</i>	McGuire & Heang 2001
Agamidae	<i>Draco palawanensis</i>	McGuire & Heang 2001
Agamidae	<i>Draco reticulatus</i>	McGuire & Heang 2001
Agamidae	<i>Draco timorensis</i>	McGuire & Heang 2001
Agamidae	<i>Draco volans</i>	Stuart-Fox & Owens 2003
Agamidae	<i>Gonocephalus bellii</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Gonocephalus bornensis</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Gonocephalus grandis</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Intellagama lesueurii</i>	Hugall et al. 2008
Agamidae	<i>Japalura brevipes</i>	Chou 2007
Agamidae	<i>Japalura kumaonensis</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Japalura luei</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Japalura makii</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Japalura polygonata</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Japalura swinhonis</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Laudakia caucasia</i>	Schulte and Moreno-Roark 2010, Melville et al. 2009, see also Edwards & Melville (2011).
Agamidae	<i>Laudakia stellio</i>	Schulte and Moreno-Roark 2010, Melville et al. 2009
Agamidae	<i>Leiolepis reevesii</i>	Srikulnath et al. 2010, Townsend et al. 2004 (subfamily placement)
Agamidae	<i>Lophognathus longirostris</i>	Hugall et al. 2008
Agamidae	<i>Moloch horridus</i>	Stuart-Fox & Owens 2003, Hugall et al. 2008
Agamidae	<i>Phrynocephalus guttatus</i>	Melville et al. 2009
Agamidae	<i>Phrynocephalus helioscopus</i>	Guo and Wang 2007, Solovyeva et al. 2011

Family	species	Source for phylogenetic relationship
Agamidae	<i>Phrynocephalus mystaceus</i>	Guo and Wang 2007, Stuart-Fox and Owens 2003
Agamidae	<i>Phrynocephalus theobaldi</i>	Jin et al. 2008, Ji et al. 2009, Guo et al. 2012
Agamidae	<i>Phrynocephalus versicolor</i>	Guo and Wang 2007, Stuart-Fox and Owens 2003
Agamidae	<i>Physignathus cocincinus</i>	Hugall et al. 2008
Agamidae	<i>Pogona barbata</i>	Schulte and Moreno-Roark 2010
Agamidae	<i>Pogona minor</i>	Melville et al. 2008
Agamidae	<i>Pogona vitticeps</i>	Melville et al. 2011
Agamidae	<i>Sitana fusca</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Sitana ponticeriana</i>	Schulte et al. 2004
Agamidae	<i>Sitana sivalensis</i>	Stuart-Fox & Owens 2003, Taxonomy
Agamidae	<i>Trapelus mutabilis</i>	Wagner et al. 2011
Agamidae	<i>Trapelus ruderatus</i>	Schulte and Moreno-Roark 2010
Agamidae	<i>Trapelus sanguinolentus</i>	Wagner et al. 2011, Melville et al. 2009
Agamidae	<i>Uromastyx acanthinura</i>	Townsend et al. 2004, Schulte and Moreno-Roark 2010
Agamidae	<i>Uromastyx aegyptia</i>	Townsend et al. 2004, Schulte and Moreno-Roark 2010
Agamidae	<i>Uromastyx ornata</i>	Wilms et al. 2009
Anguidae	<i>Anguis cephallonica</i>	Gvozdík et al. 2010
Anguidae	<i>Anguis fragilis</i>	Macey et al. 1999
Anguidae	<i>Barisia herrerae</i>	Bryson and Riddle 2012
Anguidae	<i>Barisia imbricata</i>	Bryson and Riddle 2012
Anguidae	<i>Diploglossus millepunctatus</i>	Macey et al. 1999, taxonomy
Anguidae	<i>Diploglossus pleii</i>	Conroy et al. 2005
Anguidae	<i>Dopasia gracilis</i>	Macey et al. 1999, Nguyen et al. 2011, taxonomy
Anguidae	<i>Elgaria coerulea</i>	Conroy et al. 2005
Anguidae	<i>Elgaria multicaerinata</i>	Conroy et al. 2005
Anguidae	<i>Gerrhonotus infernalis</i>	Conroy et al. 2005
Anguidae	<i>Mesaspis gadovii</i>	Macey et al. 1999, Good 1994
Anguidae	<i>Mesaspis juarezi</i>	Macey et al. 1999, Good 1994
Anguidae	<i>Mesaspis monticola</i>	Macey et al. 1999, Good 1994
Anguidae	<i>Ophiodes striatus</i>	Conroy et al. 2005
Anguidae	<i>Ophisaurus attenuatus</i>	Macey et al. 1999
Anguidae	<i>Ophisaurus compressus</i>	Macey et al. 1999, biogeography (closer to nearctic than to asian ophisaurus)
Anguidae	<i>Ophisaurus ventralis</i>	Macey et al. 1999
Anguidae	<i>Pseudopus apodus</i>	Macey et al. 1999
Anniellidae	<i>Anniella pulchra</i>	Macey et al. 1999, Townsend et al. 2004
Bipedidae	<i>Bipes biporus</i>	Townsend et al. 2004, Macey et al. 2004
Bipedidae	<i>Bipes canaliculatus</i>	Townsend et al. 2004, Macey et al. 2004
Bipedidae	<i>Bipes tridactylus</i>	Townsend et al. 2004, Macey et al. 2004
Carphodactylidae	<i>Carphodactylus laevis</i>	Gamble et al. 2012
Carphodactylidae	<i>Nephrurus levis</i>	Oliver and Bauer 2011
Carphodactylidae	<i>Phyllurus platurus</i>	Oliver and Bauer 2011
Carphodactylidae	<i>Saltuarius cornutus</i>	Oliver and Bauer 2011
Carphodactylidae	<i>Underwoodisaurus milii</i>	Oliver and Bauer 2011
Chamaeleonidae	<i>Archaius tigris</i>	Raxworthy et al. 2002
Chamaeleonidae	<i>Bradypodion pumilum</i>	Stuart-Fox et al. 2007
Chamaeleonidae	<i>Bradypodion ventrale</i>	Branch et al. 2006
Chamaeleonidae	<i>Chamaeleo calyptratus</i>	Townsend & Larson 2002
Chamaeleonidae	<i>Chamaeleo chamaeleon</i>	Townsend & Larson 2002

Family	species	Source for phylogenetic relationship
Chamaeleonidae	<i>Chamaeleo dilepis</i>	Townsend & Larson 2002
Chamaeleonidae	<i>Chamaeleo monachus</i>	Schulte and Moreno-Roark 2010
Chamaeleonidae	<i>Chamaeleo namaquensis</i>	Townsend & Larson 2002
Chamaeleonidae	<i>Furcifer cephalolepis</i>	Raxworthy et al. 2002
Chamaeleonidae	<i>Furcifer polleni</i>	Raxworthy et al. 2002
Chamaeleonidae	<i>Rhampholeon marshalli</i>	Matthee et al. 2004
Chamaeleonidae	<i>Trioceros ellioti</i>	Tilbury and Tolley 2009
Chamaeleonidae	<i>Trioceros hoehnelii</i>	Townsend & Larson 2002
Chamaeleonidae	<i>Trioceros jacksonii</i>	Tilbury and Tolley 2009
Chamaeleonidae	<i>Trioceros quadricornis</i>	Townsend & Larson 2002
Cordylidae	<i>Hemicordylus capensis</i>	Stanley et al. 2011
Cordylidae	<i>Pseudocordylus melanotus</i>	Stanley et al. 2011
Cordylidae	<i>Smaug giganteus</i>	Stanley et al. 2011
Corytophanidae	<i>Basiliscus basiliscus</i>	Schlute et al. 2003, Taxonomy
Corytophanidae	<i>Basiliscus vittatus</i>	Schulte and Moreno-Roark 2010
Crotaphytidae	<i>Crotaphytus collaris</i>	Schlute et al. 2003, McGuire et al. 2007
Crotaphytidae	<i>Crotaphytus grimeri</i>	Schlute et al. 2003, McGuire et al. 2007
Crotaphytidae	<i>Crotaphytus reticulatus</i>	Schlute et al. 2003, McGuire et al. 2007
Crotaphytidae	<i>Gambelia sila</i>	Schlute et al. 2003
Crotaphytidae	<i>Gambelia wislizenii</i>	Schlute et al. 2003
Dactyloidae	<i>Anolis acutus</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis aeneus</i>	Nicholson et al. 2005, Castaneda and de Queiroz 2011
Dactyloidae	<i>Anolis agassizi</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis aquaticus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis bimaculatus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis bonairensis</i>	Poe 2004, Castaneda and de Queiroz 2011
Dactyloidae	<i>Anolis carolinensis</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis conspersus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis cooki</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis cristatellus</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis cupreus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis cuvieri</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis evermanni</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis garmani</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis gingivinus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis grahami</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis gundlachi</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis intermedius</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis krugi</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis leachii</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis limifrons</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis lineatopus</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis lionotus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis lividus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis longiceps</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis luciae</i>	Castaneda and de Queiroz 2011
Dactyloidae	<i>Anolis monensis</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis nebulosus</i>	Nicholson et al. 2005, Poe 2004
Dactyloidae	<i>Anolis oculatus</i>	Nicholson et al. 2005

Family	species	Source for phylogenetic relationship
Dactyloidae	<i>Anolis opalinus</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis oxylophus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis pogus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis polylepis</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis pulchellus</i>	Nicholson et al. 2005, Castaneda and de Queiroz 2011
Dactyloidae	<i>Anolis reconditus</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis richardii</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis roquet</i>	Nicholson et al. 2005, Castaneda and de Queiroz 2011
Dactyloidae	<i>Anolis sabanus</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis sagrei</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis trinitatis</i>	Castaneda and de Queiroz 2011
Dactyloidae	<i>Anolis tropidolepis</i>	Nicholson et al. 2005
Dactyloidae	<i>Anolis valencienni</i>	Alfoldi et al. 2011
Dactyloidae	<i>Anolis watti</i>	Nicholson et al. 2005
Diplodactylidae	<i>Amalosia lesueurii</i>	Oliver et al. 2012
Diplodactylidae	<i>Bavayia crassicollis</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia cyclura</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia exsuccida</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia geitaina</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia montana</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia ornata</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia pulchella</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia sauvagii</i>	Bauer et al. 2012
Diplodactylidae	<i>Bavayia septuiclavis</i>	Bauer et al. 2012
Diplodactylidae	<i>Correlophus ciliatus</i>	Bauer et al. 2012
Diplodactylidae	<i>Correlophus sarasinorum</i>	Bauer et al. 2012
Diplodactylidae	<i>Crenadactylus ocellatus</i>	Melville et al. 2004
Diplodactylidae	<i>Dierogekko insularis</i>	Bauer et al. 2012
Diplodactylidae	<i>Dierogekko nehoueensis</i>	Bauer et al. 2012
Diplodactylidae	<i>Dierogekko poumensis</i>	Bauer et al. 2012
Diplodactylidae	<i>Dierogekko validiclavis</i>	Bauer et al. 2012
Diplodactylidae	<i>Diplodactylus tessellatus</i>	Oliver et al. 2009
Diplodactylidae	<i>Eurydactylodes agricolae</i>	Bauer et al. 2012
Diplodactylidae	<i>Eurydactylodes symmetricus</i>	Bauer et al. 2012
Diplodactylidae	<i>Eurydactylodes vieillardii</i>	Bauer et al. 2012
Diplodactylidae	<i>Hesperoedura reticulata</i>	Oliver et al. 2012
Diplodactylidae	<i>Lucasium damaeum</i>	Pepper et al. 2006
Diplodactylidae	<i>Mniarogekko chahoua</i>	Bauer et al. 2012
Diplodactylidae	<i>Oedodera marmorata</i>	Bauer et al. 2012
Diplodactylidae	<i>Oedura castelnaui</i>	Melville et al. 2004, Taxonomy
Diplodactylidae	<i>Oedura monilis</i>	Melville et al. 2004, Taxonomy
Diplodactylidae	<i>Oedura tryoni</i>	Melville et al. 2004, Taxonomy
Diplodactylidae	<i>Rhacodactylus auriculatus</i>	Bauer et al. 2012
Diplodactylidae	<i>Rhacodactylus leachianus</i>	Bauer et al. 2012
Diplodactylidae	<i>Rhacodactylus trachyrhynchus</i>	Bauer et al. 2012
Diplodactylidae	<i>Strophurus ciliaris</i>	Melville et al. 2004
Diplodactylidae	<i>Strophurus elderi</i>	Brown et al. 2012
Diplodactylidae	<i>Strophurus intermedius</i>	Melville et al. 2004
Diplodactylidae	<i>Strophurus williamsi</i>	Melville et al. 2004

Family	species	Source for phylogenetic relationship
Diplodactylidae	<i>Toropuku stephensi</i>	Nielsen et al. 2011
Diplodactylidae	<i>Tuketuku rakiurae</i>	Nielsen et al. 2011
Eublepharidae	<i>Aeluroscalabotes felinus</i>	Gamble et al. 2012
Eublepharidae	<i>Coleonyx brevis</i>	Ota et al. 1999, Gamble et al. 2012 (Genus)
Eublepharidae	<i>Coleonyx elegans</i>	Kratochvil & Frynta 2002
Eublepharidae	<i>Coleonyx reticulatus</i>	Kratochvil & Frynta 2002
Eublepharidae	<i>Coleonyx variegatus</i>	Ota et al. 1999
Eublepharidae	<i>Eublepharis macularius</i>	Starostova et al. 2008
Eublepharidae	<i>Eublepharis turcmenicus</i>	Kratochvil & Frynta 2002
Eublepharidae	<i>Goniurosaurus araneus</i>	Kratochvil & Frynta 2002, Gamble et al. 2012 (Genus)
Eublepharidae	<i>Goniurosaurus hainanensis</i>	Kratochvil & Frynta 2002
Eublepharidae	<i>Goniurosaurus kuroiwae</i>	Pokorna et al. 2010
Eublepharidae	<i>Goniurosaurus splendens</i>	Pokorna et al. 2010
Gekkonidae	<i>Ailuronyx seychellensis</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Ailuronyx tachyscopaeus</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Ailuronyx trachygaster</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Alsophylax laevis</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Alsophylax loricatus</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Alsophylax pipiens</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Alsophylax szczerbaki</i>	was considered conspecific with loricatus, Gamble et al. 2012 (genus)
Gekkonidae	<i>Chondrodactylus angulifer</i>	Lamb and Bauer 2006, Gamble et al. 2012 (genus)
Gekkonidae	<i>Christinus guentheri</i>	Gamble et al. 2012
Gekkonidae	<i>Christinus marmoratus</i>	Taxonomy (Gekkoninae)
Gekkonidae	<i>Cnemaspis affinis</i>	SE Asian Cnemaspis, Gamble et al. 2012
Gekkonidae	<i>Cnemaspis limi</i>	SE Asian Cnemaspis, Gamble et al. 2012
Gekkonidae	<i>Cnemaspis modiglianii</i>	SE Asian Cnemaspis, Gamble et al. 2012
Gekkonidae	<i>Cnemaspis monachorum</i>	SE Asian Cnemaspis, Gamble et al. 2012
Gekkonidae	<i>Cnemaspis perhentianensis</i>	SE Asian Cnemaspis, Gamble et al. 2012
Gekkonidae	<i>Cnemaspis whittenorum</i>	SE Asian Cnemaspis, Gamble et al. 2012
Gekkonidae	<i>Crossobamon evermanni</i>	Gamble et al. 2012 (genus)
Gekkonidae	<i>Cyrtodactylus jarakensis</i>	Taxonomy (Gekkoninae)
Gekkonidae	<i>Cyrtodactylus laevigatus</i>	Taxonomy (Gekkoninae)
Gekkonidae	<i>Cyrtodactylus leegrimeri</i>	Taxonomy (Gekkoninae)
Gekkonidae	<i>Cyrtodactylus peguensis</i>	Wood et al. 2012
Gekkonidae	<i>Cyrtodactylus redimiculus</i>	Wood et al. 2012
Gekkonidae	<i>Cyrtodactylus sadleiri</i>	Oliver et al. 2012
Gekkonidae	<i>Cyrtodactylus seribuatensis</i>	Wood et al. 2012
Gekkonidae	<i>Cyrtodactylus tiomanensis</i>	Wood et al. 2012
Gekkonidae	<i>Cyrtopodion caspium</i>	Gamble et al. 2012
Gekkonidae	<i>Cyrtopodion fedtschenkoi</i>	Bauer et al. 2013
Gekkonidae	<i>Gehyra variegata</i>	Oliver et al. 2010
Gekkonidae	<i>Gekko athymus</i>	Rosler et al. 2011, Siler et al. 2012
Gekkonidae	<i>Gekko ernstkeileri</i>	Siler et al. 2012
Gekkonidae	<i>Gekko gekko</i>	Rosler et al. 2011
Gekkonidae	<i>Gekko hokouensis</i>	Toda et al. 2008, Rosler et al. 2011
Gekkonidae	<i>Gekko japonicus</i>	Rosler et al. 2011
Gekkonidae	<i>Gekko shibatai</i>	Toda et al. 2008
Gekkonidae	<i>Gekko smithii</i>	Rosler et al. 2011

Family	species	Source for phylogenetic relationship
Gekkonidae	<i>Gekko vertebralis</i>	Toda et al. 2008
Gekkonidae	<i>Gekko yakuensis</i>	Toda et al. 2001
Gekkonidae	<i>Hemidactylus bouvieri</i>	Carranza and Arnold 2006
Gekkonidae	<i>Hemidactylus bowringii</i>	Carranza and Arnold 2006
Gekkonidae	<i>Hemidactylus brookii</i>	Carranza and Arnold 2006
Gekkonidae	<i>Hemidactylus dracaenacolus</i>	Gomez-Diaz et al. 2012
Gekkonidae	<i>Hemidactylus flaviviridis</i>	Carranza and Arnold 2006
Gekkonidae	<i>Hemidactylus frenatus</i>	Carranza and Arnold 2006
Gekkonidae	<i>Hemidactylus mabouia</i>	Carranza and Arnold 2006
Gekkonidae	<i>Hemidactylus maculatus</i>	Bauer et al. 2010
Gekkonidae	<i>Hemidactylus turcicus</i>	Carranza and Arnold 2012
Gekkonidae	<i>Heteronotia binoei</i>	Pepper et al. 2011
Gekkonidae	<i>Lepidodactylus euaensis</i>	Heinicke et al. 2012
Gekkonidae	<i>Lepidodactylus herrei</i>	Brown et al. 2012
Gekkonidae	<i>Lepidodactylus listeri</i>	Heinicke et al. 2012
Gekkonidae	<i>Lepidodactylus lugubris</i>	Heinicke et al. 2012
Gekkonidae	<i>Lepidodactylus manni</i>	Heinicke et al. 2012
Gekkonidae	<i>Lepidodactylus vanuatuensis</i>	Heinicke et al. 2012
Gekkonidae	<i>Luperosaurus macgregori</i>	Brown et al. 2000
Gekkonidae	<i>Lygodactylus klugei</i>	Gamble et al. 2011
Gekkonidae	<i>Mediodactylus kotschyi</i>	Bauer et al. 2013
Gekkonidae	<i>Mediodactylus russowii</i>	Bauer et al. 2013
Gekkonidae	<i>Pachydactylus mariquensis</i>	Lamb and Bauer 2006
Gekkonidae	<i>Perochirus ateles</i>	Heinicke et al. 2011, Genus Gamble et al. 2012
Gekkonidae	<i>Phelsuma andamanense</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma astriata</i>	Volobouev and Ineich 1994, Taxonomy (Gekkoninae)
Gekkonidae	<i>Phelsuma borbonica</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma cepediana</i>	Volobouev and Ineich 1994, Taxonomy (Gekkoninae)
Gekkonidae	<i>Phelsuma comorensis</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma dubia</i>	Taxonomy (Gekkoninae)
Gekkonidae	<i>Phelsuma guentheri</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma guimbeaui</i>	Austin et al. 2004
Gekkonidae	<i>Phelsuma inexpectata</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma nigristriata</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma ornata</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma parkeri</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma robertmertensi</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma rosagularis</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma sundbergi</i>	Rocha et al. 2009
Gekkonidae	<i>Phelsuma v-nigra</i>	Rocha et al. 2010
Gekkonidae	<i>Pseudogekko smaragdinus</i>	Heinicke et al. 2012
Gekkonidae	<i>Ptenopus garrulus</i>	Gamble et al. 2012, taxonomy
Gekkonidae	<i>Ptychozoon nicobarensis</i>	Brown et al. 2012b
Gekkonidae	<i>Stenodactylus doriae</i>	Fujita and Papenfuss 2011
Gekkonidae	<i>Urocotyledon inexpectata</i>	Gamble et al. 2012 (Genus)
Gerrhosauridae	<i>Gerrhosaurus skoogi</i>	Hugall et al. 2007, Townsend et al. 2004, Lamb et al. 2003
Gymnophthalmidae	<i>Cercosaura schreibersii</i>	Castoe et al. 2004, Doan 2003
Gymnophthalmidae	<i>Euspondylus chasqui</i>	Goicoechea et al. 2012
Gymnophthalmidae	<i>Gymnophthalmus speciosus</i>	Warne Charnov

Family	species	Source for phylogenetic relationship
Gymnophthalmidae	<i>Leposoma rugiceps</i>	Warne Charnov
Gymnophthalmidae	<i>Potamites ecleopus</i>	Castoe et al. 2004
Gymnophthalmidae	<i>Riama shrevei</i>	Doan 2003
Helodermatidae	<i>Heloderma horridum</i>	Townsend et al. 2004
Helodermatidae	<i>Heloderma suspectum</i>	Townsend et al. 2004
Iguanidae	<i>Amblyrhynchus cristatus</i>	Wiens & Hollingsworth 2000
Iguanidae	<i>Brachylophus fasciatus</i>	Keogh et al. 2008
Iguanidae	<i>Brachylophus vitiensis</i>	Keogh et al. 2008
Iguanidae	<i>Conolophus pallidus</i>	Wiens & Hollingsworth 2000
Iguanidae	<i>Conolophus subcristatus</i>	Wiens & Hollingsworth 2000
Iguanidae	<i>Ctenosaura bakeri</i>	Pasachnik et al. 2010
Iguanidae	<i>Ctenosaura oedirhina</i>	Pasachnik et al. 2010
Iguanidae	<i>Ctenosaura pectinata</i>	Wiens & Hollingsworth 2000, taxonomy
Iguanidae	<i>Ctenosaura similis</i>	Wiens & Hollingsworth 2000, taxonomy
Iguanidae	<i>Cyclura carinata</i>	Malone et al. 2000
Iguanidae	<i>Cyclura collei</i>	Malone et al. 2000
Iguanidae	<i>Cyclura cychlura</i>	Malone et al. 2000
Iguanidae	<i>Cyclura pinguis</i>	Malone et al. 2000
Iguanidae	<i>Cyclura rileyi</i>	Malone et al. 2000
Iguanidae	<i>Dipsosaurus dorsalis</i>	Wiens & Hollingsworth 2000, taxonomy
Iguanidae	<i>Iguana delicatissima</i>	taxonomy
Iguanidae	<i>Iguana iguana</i>	Wiens & Hollingsworth 2000
Iguanidae	<i>Sauromalus ater</i>	Wiens & Hollingsworth 2000, Petren and Case 1997, Petren and Case 2002
Iguanidae	<i>Sauromalus hispidus</i>	Wiens & Hollingsworth 2000, Petren and Case 1997, Petren and Case 2002
Iguanidae	<i>Sauromalus varius</i>	Wiens & Hollingsworth 2000, Petren and Case 1997, Petren and Case 2002
Lacertidae	<i>Acanthodactylus beershebensis</i>	Harris and Arnold 2000, Harris 2008
Lacertidae	<i>Acanthodactylus boskianus</i>	Harris and Arnold 2000, Harris 2008
Lacertidae	<i>Acanthodactylus erythrurus</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Acanthodactylus pardalis</i>	Mayer & Pavlicev 2007, Harris and Arnold 2000
Lacertidae	<i>Acanthodactylus schmidtii</i>	Harris and Arnold 2000
Lacertidae	<i>Acanthodactylus schreiberi</i>	Harris and Arnold 2000
Lacertidae	<i>Acanthodactylus scutellatus</i>	Mayer & Pavlicev 2007, Harris and Arnold 2000
Lacertidae	<i>Acanthodactylus spinicauda</i>	Mayer & Pavlicev 2007, Harris and Arnold 2000
Lacertidae	<i>Algyroides fitzingeri</i>	Pavlicev & Mayer 2009
Lacertidae	<i>Algyroides marchi</i>	Arnold et al. 2007, Harris et al. 1999
Lacertidae	<i>Algyroides moreoticus</i>	Mayer & Pavlicev 2007, Harris et al. 1999
Lacertidae	<i>Algyroides nigropunctatus</i>	Mayer & Pavlicev 2007, Harris et al. 1999
Lacertidae	<i>Anatololacerta anatolica</i>	Arnold et al. 2007, taxonomy
Lacertidae	<i>Anatololacerta oertzeni</i>	Arnold et al. 2007, taxonomy
Lacertidae	<i>Archaeolacerta bedriagae</i>	Arnold et al. 2007
Lacertidae	<i>Atlantolacerta andreanskyi</i>	Arnold et al. 2007
Lacertidae	<i>Dalmatolacerta oxycephala</i>	Arnold et al. 2007
Lacertidae	<i>Darevskia chlorogaster</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia derjugini</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia parvula</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia portschinskii</i>	Mayer & Pavlicev 2007, Taxonomy

Family	species	Source for phylogenetic relationship
Lacertidae	<i>Darevskia praticola</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia raddei</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia rostombekovi</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia rudis</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia saxicola</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia unisexualis</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Darevskia valentini</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Dinarolacerta mosorensis</i>	Pavlicev & Mayer 2009
Lacertidae	<i>Eremias arguta</i>	Guo et al. 2011
Lacertidae	<i>Eremias grammica</i>	Guo et al. 2011
Lacertidae	<i>Eremias intermedia</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Eremias lineolata</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Eremias nigrocellata</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Eremias nikolskii</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Eremias persica</i>	Guo et al. 2011
Lacertidae	<i>Eremias pleskei</i>	Guo et al. 2011
Lacertidae	<i>Eremias regeli</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Eremias strauchi</i>	Rastegar-Pouyani et al. 2010
Lacertidae	<i>Eremias velox</i>	Guo et al. 2011
Lacertidae	<i>Gallotia atlantica</i>	Arnold et al. 2007
Lacertidae	<i>Gallotia bravoana</i>	Arnold et al. 2007
Lacertidae	<i>Gallotia caesaris</i>	Arnold et al. 2007
Lacertidae	<i>Gallotia galloti</i>	Arnold et al. 2007
Lacertidae	<i>Gallotia intermedia</i>	Arnold et al. 2007
Lacertidae	<i>Gallotia simonyi</i>	Arnold et al. 2007
Lacertidae	<i>Gallotia stehlini</i>	Arnold et al. 2007
Lacertidae	<i>Heliobolus lugubris</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Hellenolacerta graeca</i>	Arnold et al. 2007
Lacertidae	<i>Iberolacerta aranica</i>	Arnold et al. 2007
Lacertidae	<i>Iberolacerta aurelioi</i>	Arnold et al. 2007
Lacertidae	<i>Iberolacerta bonnali</i>	Arnold et al. 2007
Lacertidae	<i>Iberolacerta cyreni</i>	Arnold et al. 2007
Lacertidae	<i>Iberolacerta horvathi</i>	Arnold et al. 2007
Lacertidae	<i>Iberolacerta monticola</i>	Arnold et al. 2007
Lacertidae	<i>Ichnotropis capensis</i>	Edwards et al. 2012
Lacertidae	<i>Ichnotropis squamulosa</i>	Edwards et al. 2012
Lacertidae	<i>Lacerta agilis</i>	Arnold et al. 2007, Godhino et al. 2005
Lacertidae	<i>Lacerta schreiberi</i>	Arnold et al. 2007, Godhino et al. 2005
Lacertidae	<i>Lacerta strigata</i>	Arnold et al. 2007, Godhino et al. 2005
Lacertidae	<i>Lacerta trilineata</i>	Arnold et al. 2007, Godhino et al. 2005
Lacertidae	<i>Lacerta viridis</i>	Arnold et al. 2007, Godhino et al. 2005
Lacertidae	<i>Meroles anchietae</i>	Lamb & Bauer 2003
Lacertidae	<i>Meroles cuneirostris</i>	Lamb & Bauer 2003
Lacertidae	<i>Meroles suborbitalis</i>	Edwards et al. 2012
Lacertidae	<i>Mesalina guttulata</i>	Arnold et al. 2007, Mayer & Pavlicev 2007, Joger and Mayer 2002, Kapli et al. 2008
Lacertidae	<i>Mesalina olivieri</i>	Arnold et al. 2007, Mayer & Pavlicev 2007, Joger and Mayer 2002, Kapli et al. 2008
Lacertidae	<i>Mesalina pasteuiri</i>	Arnold et al. 2007, Mayer & Pavlicev 2007, Joger and

Family	species	Source for phylogenetic relationship
		Mayer 2002, Kapli et al. 2008
Lacertidae	<i>Mesalina rubropunctata</i>	Arnold et al. 2007, Mayer & Pavlicev 2007, Joger and Mayer 2002, Kapli et al. 2008
Lacertidae	<i>Nucras lalandii</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Nucras taeniolata</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Nucras tessellata</i>	Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Ophisops elegans</i>	Arnold et al. 2007, Mayer & Pavlicev 2007, Taxonomy
Lacertidae	<i>Parvilacerta parva</i>	Arnold et al. 2007
Lacertidae	<i>Pedioplanis burchelli</i>	Makokha et al. 2007, Conradie et al. 2012
Lacertidae	<i>Pedioplanis namaquensis</i>	Makokha et al. 2007, Conradie et al. 2012
Lacertidae	<i>Phoenicolacerta laevis</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Phoenicolacerta troodica</i>	was synonym of laevis
Lacertidae	<i>Podarcis bocagei</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis carbonelli</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis cretensis</i>	Lymberakis et al. 2008
Lacertidae	<i>Podarcis erhardii</i>	Arnold et al. 2007, Poulakakis et al. 2005
Lacertidae	<i>Podarcis filfolensis</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis gaigeae</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis hispanicus</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis lilfordi</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis melisellensis</i>	Arnold et al. 2007, Poulakakis et al. 2005
Lacertidae	<i>Podarcis milensis</i>	Arnold et al. 2007, Poulakakis et al. 2005, Amat 2008
Lacertidae	<i>Podarcis muralis</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis peloponnesiacus</i>	Arnold et al. 2007, Poulakakis et al. 2005
Lacertidae	<i>Podarcis pityusensis</i>	Arnold et al. 2007
Lacertidae	<i>Podarcis raffoneae</i>	Harris et al. 2005
Lacertidae	<i>Podarcis siculus</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis tauricus</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Podarcis tiliguerta</i>	Arnold et al. 2007
Lacertidae	<i>Podarcis waglerianus</i>	Arnold et al. 2007, Harris et al. 2005
Lacertidae	<i>Psammodromus algirus</i>	Carranza et al. 2006, Arnold et al. 2007
Lacertidae	<i>Psammodromus blanci</i>	Carranza et al. 2006, Arnold et al. 2007
Lacertidae	<i>Psammodromus hispanicus</i>	Carranza et al. 2006, Arnold et al. 2007
Lacertidae	<i>Scelarcis perspicillata</i>	Arnold et al. 2007
Lacertidae	<i>Takydromus amurensis</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus dorsalis</i>	Lin et al. 2002
Lacertidae	<i>Takydromus formosanus</i>	Lin et al. 2002
Lacertidae	<i>Takydromus hsuehshanensis</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus luyeanus</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus sauteri</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus septentrionalis</i>	Lin et al. 2002
Lacertidae	<i>Takydromus sexlineatus</i>	Lin et al. 2002
Lacertidae	<i>Takydromus smaragdinus</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus stejnegeri</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus toyamai</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus viridipunctatus</i>	Lue and Lin 2008
Lacertidae	<i>Takydromus wolteri</i>	Lue and Lin 2008
Lacertidae	<i>Teira dugesii</i>	Arnold et al. 2007, Taxonomy
Lacertidae	<i>Timon lepidus</i>	Ahmadzadeh et al. 2012

Family	species	Source for phylogenetic relationship
Lacertidae	<i>Timon pater</i>	Ahmadzadeh et al. 2012
Lacertidae	<i>Zootoca vivipara</i>	Arnold et al. 2007
Leiocephalidae	<i>Leiocephalus eremitus</i>	Torres-Carvajal and de Queiroz 2009, taxonomy
Leiocephalidae	<i>Leiocephalus inaguae</i>	Torres-Carvajal and de Queiroz 2009, taxonomy
Leiocephalidae	<i>Leiocephalus psammodromus</i>	Torres-Carvajal and de Queiroz 2009, taxonomy
Leiosauridae	<i>Enyalius perditus</i>	Frost et al. 2001
Liolaemidae	<i>Liolaemus andinus</i>	Pincheira-Donoso et al. 2008
Liolaemidae	<i>Liolaemus bellii</i>	Pincheira-Donoso et al. 2008b
Liolaemidae	<i>Liolaemus bibronii</i>	Morando et al. 2007, Olave et al. 2011, Pincheira-Donoso et al. 2008
Liolaemidae	<i>Liolaemus boulengeri</i>	Pincheira-Donoso, pers. Comm, November 2011
Liolaemidae	<i>Liolaemus elongatus</i>	Moreno Azocar et al. 2012
Liolaemidae	<i>Liolaemus huacahuasicus</i>	Montanus clade. Pincheira-Donoso et al. 2008a
Liolaemidae	<i>Liolaemus koslowskyi</i>	Camargo et al. 2012
Liolaemidae	<i>Liolaemus lemniscatus</i>	Pincheira-Donoso et al. 2008b
Liolaemidae	<i>Liolaemus lineomaculatus</i>	Pincheira-Donoso et al. 2009
Liolaemidae	<i>Liolaemus lutzae</i>	Pincheira-Donoso et al. 2008
Liolaemidae	<i>Liolaemus nigromaculatus</i>	Labra et al. 2009
Liolaemidae	<i>Liolaemus pictus</i>	Pincheira-Donoso et al. 2008
Liolaemidae	<i>Liolaemus scolaroi</i>	Breitman et al. 2011
Liolaemidae	<i>Liolaemus signifer</i>	Pincheira-Donoso et al. 2008
Liolaemidae	<i>Liolaemus wiegmanni</i>	Moreno Azocar et al. 2012
Liolaemidae	<i>Liolaemus zullyae</i>	Breitman et al. 2011
Liolaemidae	<i>Phymaturus palluma</i>	Lobo et al. 2012, Morando et al. 2013
Liolaemidae	<i>Phymaturus patagonicus</i>	Lobo et al. 2012, Morando et al. 2013
Liolaemidae	<i>Phymaturus punae</i>	Lobo et al. 2012, Morando et al. 2013
Liolaemidae	<i>Phymaturus zapalensis</i>	Morando et al. 2013
Phrynosomatidae	<i>Callisaurus draconoides</i>	Wiens et al. 2010
Phrynosomatidae	<i>Cophosaurus texanus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Holbrookia lacerata</i>	Wiens et al. 2010
Phrynosomatidae	<i>Holbrookia maculata</i>	Wiens et al. 2010
Phrynosomatidae	<i>Holbrookia propinqua</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma blainvillii</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma cornutum</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma coronatum</i>	Leache & McGuire 2006
Phrynosomatidae	<i>Phrynosoma ditmarsii</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma douglassii</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma hernandesi</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma mcallii</i>	Leache & McGuire 2006
Phrynosomatidae	<i>Phrynosoma modestum</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma platyrhinus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Phrynosoma solare</i>	Leache & McGuire 2006
Phrynosomatidae	<i>Sceloporus aeneus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus angustus</i>	Wiens & Reeder 1997
Phrynosomatidae	<i>Sceloporus arenicolus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus clarkii</i>	Leache 2010
Phrynosomatidae	<i>Sceloporus consobrinus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus cozumelae</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus cyanogenys</i>	Wiens et al. 2010

Family	species	Source for phylogenetic relationship
Phrynosomatidae	<i>Sceloporus formosus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus gadoviae</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus graciosus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus grammicus</i>	Leache 2010
Phrynosomatidae	<i>Sceloporus horridus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus jarrovii</i>	Wiens & Reeder 1997, Martinez-Mendez and Mendez-de la Cruz 2007, Leache 2010
Phrynosomatidae	<i>Sceloporus magister</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus malachiticus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus merriami</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus mucronatus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus occidentalis</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus olivaceus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus omiltemanus</i>	Martinez-Mendez & Mendez-de la Cruz 2007
Phrynosomatidae	<i>Sceloporus orcutti</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus poinsettii</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus scalaris</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus serrifer</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus slevini</i>	Wiens & Reeder 1997, Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus torquatus</i>	Wiens & Reeder 1997, Martinez-Mendez and Mendez-de la Cruz 2007
Phrynosomatidae	<i>Sceloporus undulatus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus utiformis</i>	Wiens & Reeder 1997
Phrynosomatidae	<i>Sceloporus variabilis</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus virgatus</i>	Wiens et al. 2010
Phrynosomatidae	<i>Sceloporus woodi</i>	Wiens et al. 2010
Phrynosomatidae	<i>Uma exsul</i>	Schulte and Moreno-Roark 2010
Phrynosomatidae	<i>Uma notata</i>	Wiens et al. 2010
Phrynosomatidae	<i>Urosaurus bicarinatus</i>	Feldman et al. 2011
Phrynosomatidae	<i>Urosaurus clarionensis</i>	Feldman et al. 2011
Phrynosomatidae	<i>Urosaurus graciosus</i>	Feldman et al. 2011
Phrynosomatidae	<i>Urosaurus ornatus</i>	Feldman et al. 2011
Phrynosomatidae	<i>Uta palmeri</i>	Upton and Murphy 1997
Phrynosomatidae	<i>Uta stansburiana</i>	Bonine et al. 2005
Phyllodactylidae	<i>Gymnodactylus amarali</i>	Pellegrino et al. 2005, taxonomy
Phyllodactylidae	<i>Gymnodactylus geckoides</i>	Pellegrino et al. 2005
Phyllodactylidae	<i>Haemodracon riebeckii</i>	Gamble et al. 2011
Phyllodactylidae	<i>Haemodracon trachyrhinus</i>	Gamble et al. 2011
Phyllodactylidae	<i>Homonota darwinii</i>	Gamble et al. 2008, Taxonomy
Phyllodactylidae	<i>Homonota fasciata</i>	Gamble et al. 2008, Taxonomy
Phyllodactylidae	<i>Phyllodactylus bugastrolepis</i>	Gamble et al. 2011
Phyllodactylidae	<i>Phyllodactylus lanei</i>	Blair et al. 2009
Phyllodactylidae	<i>Phyllopezus pollicaris</i>	Gamble et al. 2012
Phyllodactylidae	<i>Ptyodactylus guttatus</i>	Gamble et al. 2008b, taxonomy
Phyllodactylidae	<i>Ptyodactylus oudrii</i>	Gamble et al. 2008b, taxonomy
Phyllodactylidae	<i>Ptyodactylus ragazzii</i>	Gamble et al. 2008b, taxonomy
Phyllodactylidae	<i>Tarentola angustimentalis</i>	Gamble et al. 2008, Carranza et al. 2002
Phyllodactylidae	<i>Tarentola annularis</i>	Gamble et al. 2008, Carranza et al. 2002
Phyllodactylidae	<i>Tarentola boettgeri</i>	Gamble et al. 2008, Carranza et al. 2002

Family	species	Source for phylogenetic relationship
Phyllodactylidae	<i>Tarentola chazaliae</i>	Gamble et al. 2008, Carranza et al. 2002
Phyllodactylidae	<i>Tarentola darwini</i>	Vasconcelos et al. 2012
Phyllodactylidae	<i>Tarentola delalandii</i>	Vasconcelos et al. 2012
Phyllodactylidae	<i>Tarentola deserti</i>	Rato et al. 2012
Phyllodactylidae	<i>Tarentola gomerensis</i>	Vasconcelos et al. 2012
Phyllodactylidae	<i>Tarentola mauritanica</i>	Gamble et al. 2008, Carranza et al. 2002
Polychrotidae	<i>Polychrus acutirostris</i>	Schlute et al. 2003, taxonomy
Pygopodidae	<i>Aprasia pulchella</i>	Gamble et al. 2008, Taxonomy
Pygopodidae	<i>Aprasia repens</i>	Gamble et al. 2008, Lee et al. 2009
Pygopodidae	<i>Delma fraseri</i>	Jennings et al. 2003
Pygopodidae	<i>Delma grayii</i>	Jennings et al. 2003
Pygopodidae	<i>Lialis burtonis</i>	Gamble et al. 2008, Lee et al. 2009
Pygopodidae	<i>Pygopus lepidopodus</i>	Oliver et al. 2010
Scincidae	<i>Ablepharus kitaibelii</i>	Macey et al. 2006, taxonomy
Scincidae	<i>Ablepharus rueppellii</i>	Macey et al. 2006, taxonomy
Scincidae	<i>Acontias garipeensis</i>	Daniels et al. 2006, Taxonomy
Scincidae	<i>Acontias kgalagadi</i>	Daniels et al. 2006, Taxonomy
Scincidae	<i>Afroablepharus annobonensis</i>	Jesus et al. 2007, taxonomy
Scincidae	<i>Aspronema cochabambae</i>	Miralles et al. 2009, Hedges and Conn 2012
Scincidae	<i>Asymblepharus sikimmensis</i>	Macey et al. 2006, taxonomy
Scincidae	<i>Bassiana duperreyi</i>	Skinner et al. 2011, taxonomy
Scincidae	<i>Bassiana trilineata</i>	Skinner et al. 2011
Scincidae	<i>Brachymeles cebuensis</i>	Siler et al. 2011
Scincidae	<i>Brachymeles minimus</i>	Siler et al. 2011
Scincidae	<i>Brachymeles taylori</i>	Siler et al. 2011
Scincidae	<i>Brachymeles tridactylus</i>	Siler et al. 2011
Scincidae	<i>Brasiliscincus agilis</i>	Miralles et al. 2009
Scincidae	<i>Brasiliscincus heathi</i>	Whiting et al. 2006
Scincidae	<i>Caledoniscincus aquilonius</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Caledoniscincus atropunctatus</i>	Sadlier et al. 2012
Scincidae	<i>Caledoniscincus auratus</i>	Smith et al. 2007
Scincidae	<i>Caledoniscincus austrocaledonicus</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Caledoniscincus chazeaui</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Caledoniscincus festivus</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Caledoniscincus haplorhinus</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Caledoniscincus orestes</i>	Sadlier et al. 2012
Scincidae	<i>Caledoniscincus renevieri</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Caledoniscincus terma</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Carlia longipes</i>	Dolman and Hugall 2008
Scincidae	<i>Carlia rhomboidalis</i>	Reeder 2003, Dolman and Hugall 2008
Scincidae	<i>Carlia rostralis</i>	Dolman and Hugall 2008
Scincidae	<i>Carlia rubrigularis</i>	Dolman and Hugall 2008
Scincidae	<i>Carlia tetradactyla</i>	Dolman and Hugall 2008
Scincidae	<i>Celatiscincus euryotis</i>	Smith et al. 2007, Sadlier et al. 2012
Scincidae	<i>Celatiscincus similis</i>	Sadlier et al. 2009
Scincidae	<i>Chalcides bedriagai</i>	Carranza et al. 2008
Scincidae	<i>Chalcides chalcides</i>	Carranza et al. 2008
Scincidae	<i>Chalcides coeruleopunctatus</i>	Carranza et al. 2008

Family	species	Source for phylogenetic relationship
Scincidae	<i>Chalcides ocellatus</i>	Carranza et al. 2008, Austin and Arnold 2006
Scincidae	<i>Chalcides sepsoides</i>	Carranza et al. 2008
Scincidae	<i>Chalcides sexlineatus</i>	Carranza et al. 2008
Scincidae	<i>Chalcides viridanus</i>	Carranza et al. 2008
Scincidae	<i>Chioninia coctei</i>	Miralles et al. 2011
Scincidae	<i>Chioninia spinalis</i>	Miralles et al. 2011
Scincidae	<i>Chioninia vaillantii</i>	Miralles et al. 2011
Scincidae	<i>Coeranoscincus reticulatus</i>	Skinner 2007
Scincidae	<i>Copeoglossum nigropunctatum</i>	Miralles et al. 2009, Hedges and Conn 2012
Scincidae	<i>Corucia zebrata</i>	Skinner et al. 2011
Scincidae	<i>Cryptoblepharus egeriae</i>	Schmitz et al. 2005, Taxonomy
Scincidae	<i>Cryptoblepharus eximius</i>	Schmitz et al. 2005, Taxonomy
Scincidae	<i>Cryptoblepharus novocaledonicus</i>	Schmitz et al. 2005, Taxonomy
Scincidae	<i>Cryptoblepharus poecilopleurus</i>	Schmitz et al. 2005, Taxonomy
Scincidae	<i>Cryptoblepharus renschi</i>	Schmitz et al. 2005, Taxonomy
Scincidae	<i>Ctenotus brooksi</i>	Rabosky et al. 2007
Scincidae	<i>Ctenotus lanceolini</i>	Kay and Keogh 2012
Scincidae	<i>Ctenotus pantherinus</i>	Rabosky et al. 2007
Scincidae	<i>Ctenotus robustus</i>	Rabosky et al. 2007
Scincidae	<i>Ctenotus taeniolatus</i>	Rabosky et al. 2007
Scincidae	<i>Cyclodomorphus casuarinae</i>	Gardner et al. 2008, taxonomy
Scincidae	<i>Cyclodomorphus celatus</i>	Gardner et al. 2008
Scincidae	<i>Cyclodomorphus melanops</i>	Gardner et al. 2008, taxonomy
Scincidae	<i>Cyclodomorphus michaeli</i>	Gardner et al. 2008, taxonomy
Scincidae	<i>Dasia olivacea</i>	Datta-Roy et al. 2012
Scincidae	<i>Egernia cunninghami</i>	Gardner et al. 2008
Scincidae	<i>Egernia hosmeri</i>	Doughty et al. 2011
Scincidae	<i>Egernia kingii</i>	Doughty et al. 2011
Scincidae	<i>Egernia stokesii</i>	Doughty et al. 2011
Scincidae	<i>Egernia striolata</i>	Gardner et al. 2008
Scincidae	<i>Emoia adspersa</i>	Hamilton 2008
Scincidae	<i>Emoia aneityumensis</i>	Reeder 2003, Hamilton 2008
Scincidae	<i>Emoia atrocostata</i>	Hamilton 2008
Scincidae	<i>Emoia campbelli</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia concolor</i>	Reeder 2003, Hamilton 2008
Scincidae	<i>Emoia impar</i>	Hamilton 2008
Scincidae	<i>Emoia isolata</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia loyaltiensis</i>	Reeder 2003, Hamilton 2008
Scincidae	<i>Emoia maculata</i>	Linkem et al. 2011
Scincidae	<i>Emoia nativittatis</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia nigromarginata</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia parkeri</i>	Reeder 2003, Hamilton 2008
Scincidae	<i>Emoia ponapea</i>	Hamilton 2008
Scincidae	<i>Emoia reimschiüsseli</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia sanfordi</i>	Hamilton 2008
Scincidae	<i>Emoia schmidti</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia slevini</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia sorex</i>	Reeder 2003, taxonomy

Family	species	Source for phylogenetic relationship
Scincidae	<i>Emoia taumakoensis</i>	Reeder 2003, taxonomy
Scincidae	<i>Emoia tongana</i>	Reeder 2003, Hamilton 2008
Scincidae	<i>Emoia trossula</i>	Zug et al. 2012
Scincidae	<i>Eremiascincus richardsonii</i>	Mecke et al. 2009
Scincidae	<i>Eulamprus brachyosoma</i>	Rabosky et al. 2007, taxonomy
Scincidae	<i>Eulamprus quoyii</i>	Rabosky et al. 2007, taxonomy
Scincidae	<i>Eulamprus tympanum</i>	Rabosky et al. 2007, taxonomy
Scincidae	<i>Eumeces schneideri</i>	Perera et al. 2011
Scincidae	<i>Eutropis carinata</i>	Datta-Roy et al. 2012
Scincidae	<i>Eutropis longicaudata</i>	Datta-Roy et al. 2012
Scincidae	<i>Eutropis multifasciata</i>	Datta-Roy et al. 2012
Scincidae	<i>Gongylomorphus bojerii</i>	Austin and Arnold 2006
Scincidae	<i>Hemiergis peronii</i>	Reeder and Reichert 2011
Scincidae	<i>Hemisphaeriodon gerrardii</i>	Gardner et al. 2008
Scincidae	<i>Insulasaurus arborens</i>	Linkem et al. 2011
Scincidae	<i>Insulasaurus traanorum</i>	Linkem et al. 2011
Scincidae	<i>Janetaescincus braueri</i>	Austin and Arnold 2006
Scincidae	<i>Kanakysaurus viviparus</i>	Sadlier et al. 2009
Scincidae	<i>Lampropholis delicata</i>	Rabosky et al. 2007, Reeder 2003, taxonomy
Scincidae	<i>Lampropholis guichenoti</i>	Rabosky et al. 2007, Reeder 2003, taxonomy
Scincidae	<i>Larutia seribuatensis</i>	Grismer et al. 2011
Scincidae	<i>Leiopisma telfairii</i>	Austin and Arnold 2006
Scincidae	<i>Lerista borealis</i>	Skinner and Lee 2010
Scincidae	<i>Lerista bougainvillii</i>	Skinner and Lee 2010
Scincidae	<i>Lerista labialis</i>	Skinner and Lee 2010
Scincidae	<i>Lerista punctatovittata</i>	Skinner and Lee 2010
Scincidae	<i>Lerista xanthura</i>	Skinner and Lee 2010
Scincidae	<i>Liopholis inornata</i>	Gardner et al. 2008
Scincidae	<i>Liopholis kintorei</i>	Gardner et al. 2008
Scincidae	<i>Liopholis modesta</i>	Chapple et al. 2008
Scincidae	<i>Liopholis striata</i>	Gardner et al. 2008
Scincidae	<i>Liopholis whitii</i>	Chapple et al. 2008
Scincidae	<i>Lioscincus greeri</i>	Smith et al. 2007
Scincidae	<i>Lioscincus maruia</i>	Smith et al. 2007
Scincidae	<i>Lioscincus nigrofasciolatum</i>	Smith et al. 2007
Scincidae	<i>Lioscincus novaecaledoniae</i>	Smith et al. 2007
Scincidae	<i>Lioscincus steindachneri</i>	Smith et al. 2007
Scincidae	<i>Lipinia leptosoma</i>	Skinner et al. 2011, taxonomy
Scincidae	<i>Lipinia macrotympanum</i>	Skinner et al. 2011, taxonomy
Scincidae	<i>Lipinia rabori</i>	Skinner et al. 2011, taxonomy
Scincidae	<i>Lipinia rouxi</i>	Austin 1998
Scincidae	<i>Lissolepis coventryi</i>	Warne Charnov
Scincidae	<i>Mabuya mabouya</i>	Whiting et al. 2006
Scincidae	<i>Marisora unimarginata</i>	Whiting et al. 2006, Honda et al. 2006, Miralles et al. 2006
Scincidae	<i>Marmorosphax boulinda</i>	Sadlier et al. 2009
Scincidae	<i>Marmorosphax taom</i>	Sadlier et al. 2009
Scincidae	<i>Marmorosphax tricolor</i>	Sadlier et al. 2009
Scincidae	<i>Menetia greyii</i>	Smith et al. 2007
Scincidae	<i>Morethia boulengeri</i>	Reeder 2003, taxonomy

Family	species	Source for phylogenetic relationship
Scincidae	<i>Morethia obscura</i>	Reeder 2003, taxonomy
Scincidae	<i>Nannoscincus garrulus</i>	Smith et al. 2007
Scincidae	<i>Nannoscincus gracilis</i>	Smith et al. 2007
Scincidae	<i>Nannoscincus greeri</i>	Smith et al. 2007
Scincidae	<i>Nannoscincus maccoyi</i>	Smith et al. 2007, taxonomy, biogeography (sister to New Caledonia skinks)
Scincidae	<i>Nannoscincus mariei</i>	Smith et al. 2007
Scincidae	<i>Niveoscincus coventryi</i>	Melville & Swain 2000
Scincidae	<i>Niveoscincus metallicus</i>	Melville & Swain 2000
Scincidae	<i>Oligosoma fallai</i>	Chapple et al. 2009
Scincidae	<i>Oligosoma homalonotum</i>	Chapple et al. 2009
Scincidae	<i>Oligosoma lichenigera</i>	Smith et al. 2007, Chapple et al. 2009
Scincidae	<i>Oligosoma nigriplantare</i>	Chapple et al. 2009
Scincidae	<i>Oligosoma notosaurus</i>	Chapple et al. 2011
Scincidae	<i>Oligosoma stenotis</i>	Chapple et al. 2011
Scincidae	<i>Ophiomorus punctatissimus</i>	Greer and Wilson 2001
Scincidae	<i>Pamelaescincus gardineri</i>	Austin and Arnold 2006
Scincidae	<i>Panaspis kitsoni</i>	Schmitz et al. 2005, Jesus et al. 2007, taxonomy
Scincidae	<i>Panaspis nimbaensis</i>	Schmitz et al. 2005, Jesus et al. 2007, taxonomy
Scincidae	<i>Panaspis wahlbergi</i>	Jesus et al. 2007, taxonomy
Scincidae	<i>Parvosincus sisoni</i>	Linkem et al. 2011
Scincidae	<i>Plestiodon anthracinus</i>	Brandley et al. 2012, Siler et al. 2011 for placement of the genus with Eumeces as sister to Brachymeles
Scincidae	<i>Plestiodon copei</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon egregius</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon fasciatus</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon inexpectatus</i>	Brandley et al. 2011
Scincidae	<i>Plestiodon kishinouyei</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon laticeps</i>	Richmond 2006
Scincidae	<i>Plestiodon lynxe</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon obsoletus</i>	Richmond 2006, Brandley et al. 2011
Scincidae	<i>Plestiodon okadae</i>	Okamoto & Hikida 2009: allied to laticutatus and japonicus
Scincidae	<i>Plestiodon reynoldsi</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon septentrionalis</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon skiltonianus</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon stimpsonii</i>	Brandley et al. 2012
Scincidae	<i>Plestiodon tetragrammus</i>	Brandley et al. 2012
Scincidae	<i>Pseudemoia entrecasteauxii</i>	Smith et al. 2007, taxonomy
Scincidae	<i>Pseudemoia pagenstecheri</i>	Smith et al. 2007, taxonomy
Scincidae	<i>Pseudemoia rawlinsoni</i>	Hutchinson & Donnellan 1992
Scincidae	<i>Pseudemoia spenceri</i>	Stewart and Thompson 2003
Scincidae	<i>Psychosaura macrorhyncha</i>	Hedges and Conn 2012
Scincidae	<i>Scincella lateralis</i>	Reeder 2003, Grismer et al. 2011
Scincidae	<i>Scincus mitranus</i>	Smith et al. 2007, taxonomy
Scincidae	<i>Scincus scincus</i>	Smith et al. 2007, taxonomy
Scincidae	<i>Sigaloseps deplanchei</i>	Smith et al. 2007
Scincidae	<i>Sigaloseps ruficauda</i>	Smith et al. 2007
Scincidae	<i>Sphenomorphus concinnatus</i>	Linkem et al. 2011
Scincidae	<i>Sphenomorphus cranei</i>	Linkem et al. 2011

Family	species	Source for phylogenetic relationship
Scincidae	<i>Sphenomorphus indicus</i>	Linkem et al. 2011
Scincidae	<i>Sphenomorphus maculatus</i>	Linkem et al. 2011
Scincidae	<i>Sphenomorphus scutatus</i>	Linkem et al. 2011
Scincidae	<i>Tiliqua adelaidensis</i>	Gardner et al. 2008
Scincidae	<i>Tiliqua nigrolutea</i>	Gardner et al. 2008
Scincidae	<i>Tiliqua occipitalis</i>	Gardner et al. 2008
Scincidae	<i>Tiliqua rugosa</i>	Gardner et al. 2008
Scincidae	<i>Tiliqua scincoides</i>	Gardner et al. 2008
Scincidae	<i>Trachylepis affinis</i>	Rocha et al. 2010
Scincidae	<i>Trachylepis aurata</i>	Sindaco et al. 2012
Scincidae	<i>Trachylepis buettneri</i>	Rocha et al. 2010, taxonomy
Scincidae	<i>Trachylepis lavarambo</i>	Rocha et al. 2010, taxonomy
Scincidae	<i>Trachylepis maculilabris</i>	Sindaco et al. 2012
Scincidae	<i>Trachylepis quinquetaeniata</i>	Sindaco et al. 2012
Scincidae	<i>Trachylepis sechellensis</i>	Sindaco et al. 2012
Scincidae	<i>Trachylepis sparsa</i>	Rocha et al. 2010, taxonomy
Scincidae	<i>Trachylepis spilogaster</i>	Rocha et al. 2010, Datta-Roy et al. 2012, Sindaco et al. 2012
Scincidae	<i>Trachylepis striata</i>	Rocha et al. 2010, Sindaco et al. 2012
Scincidae	<i>Trachylepis varia</i>	Sindaco et al. 2012
Scincidae	<i>Trachylepis vittata</i>	Sindaco et al. 2012
Scincidae	<i>Trachylepis wrightii</i>	Sindaco et al. 2012
Scincidae	<i>Tribolonotus blanchardi</i>	Austin et al. 2010
Scincidae	<i>Tribolonotus brongersmai</i>	Austin et al. 2010
Scincidae	<i>Tribolonotus ponceleti</i>	Austin et al. 2010
Scincidae	<i>Tribolonotus pseudoponceleti</i>	Austin et al. 2010
Scincidae	<i>Tribolonotus schmidti</i>	Austin et al. 2010
Scincidae	<i>Tropidoscincus boreus</i>	Smith et al. 2007
Scincidae	<i>Tropidoscincus variabilis</i>	Smith et al. 2007
Scincidae	<i>Varzea bistriata</i>	Whiting et al. 2006, Hedges and Conn 2012
Sphaerodactylidae	<i>Aristelliger barbouri</i>	Gamble et al. 2012, taxonomy
Sphaerodactylidae	<i>Aristelliger georgeensis</i>	Gamble et al. 2012
Sphaerodactylidae	<i>Aristelliger praesignis</i>	Gamble et al. 2012
Sphaerodactylidae	<i>Gonatodes albogularis</i>	Schargel et al. 2010
Sphaerodactylidae	<i>Gonatodes daudini</i>	Gamble et al. 2012
Sphaerodactylidae	<i>Gonatodes ocellatus</i>	Schargel et al. 2010
Sphaerodactylidae	<i>Gonatodes vittatus</i>	Schargel et al. 2010
Sphaerodactylidae	<i>Pristurus abdelkuri</i>	Papenfuss et al. 2009
Sphaerodactylidae	<i>Pristurus guichardi</i>	Papenfuss et al. 2009
Sphaerodactylidae	<i>Pristurus insignis</i>	Arnold 2009, Papenfuss et al. 2009
Sphaerodactylidae	<i>Pristurus insignoides</i>	Arnold 2009
Sphaerodactylidae	<i>Pristurus rupestris</i>	Gamble et al. 2011, Papenfuss et al. 2009
Sphaerodactylidae	<i>Pristurus sokotranus</i>	Papenfuss et al. 2009
Sphaerodactylidae	<i>Quedenfeldtia trachyblepharus</i>	Gamble et al. 2012
Sphaerodactylidae	<i>Sphaerodactylus argivus</i>	Geurgas et al. 2008, Gamble et al. 2008b, Taxonomy
Sphaerodactylidae	<i>Sphaerodactylus kirbyi</i>	Geurgas et al. 2008, Gamble et al. 2008b, Taxonomy
Sphaerodactylidae	<i>Sphaerodactylus nicholsi</i>	Gamble et al. 2012 (partial)
Sphaerodactylidae	<i>Sphaerodactylus roosevelti</i>	Gamble et al. 2012 (partial)
Sphaerodactylidae	<i>Sphaerodactylus vincenti</i>	Geurgas et al. 2008, Gamble et al. 2008b, Taxonomy
Sphaerodactylidae	<i>Teratoscincus microlepis</i>	Gamble et al. 2012

Family	species	Source for phylogenetic relationship
Sphaerodactylidae	<i>Teratoscincus scincus</i>	Gamble et al. 2012
Teiidae	<i>Ameiva ameiva</i>	Hower and Hedges 2003, Giugliano et al. 2007, Reeder et al. 2002,
Teiidae	<i>Ameiva erythrocephala</i>	Hower and Hedges 2003
Teiidae	<i>Ameiva exsul</i>	Hower and Hedges 2003
Teiidae	<i>Ameiva fuscata</i>	Hower and Hedges 2003
Teiidae	<i>Ameiva plei</i>	Hower and Hedges 2003
Teiidae	<i>Ameiva polops</i>	Hower and Hedges 2003
Teiidae	<i>Ameiva wetmorei</i>	Hower and Hedges 2003
Teiidae	<i>Ameivula ocellifera</i>	Giugliano et al. 2007
Teiidae	<i>Aspidoscelis burti</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis ceralbensis</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis cozumelae</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis deppei</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis dixonii</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis exsanguis</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis gularis</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis guttata</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis hyperythra</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis inornata</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis laredoensis</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis lineattissima</i>	Castiglia et al. 2010
Teiidae	<i>Aspidoscelis marmorata</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis neomexicana</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis scalaris</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis sexlineata</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis sonora</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis tessellata</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis tigris</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis uniparens</i>	Reeder et al. 2002
Teiidae	<i>Aspidoscelis velox</i>	Reeder et al. 2002
Teiidae	<i>Cnemidophorus arubensis</i>	Harvey et al. 2012
Teiidae	<i>Cnemidophorus cryptus</i>	Reeder et al. 2002
Teiidae	<i>Cnemidophorus gramivagus</i>	Reeder et al. 2002
Teiidae	<i>Cnemidophorus lemniscatus</i>	Reeder et al. 2002
Teiidae	<i>Cnemidophorus murinus</i>	Reeder et al. 2002
Teiidae	<i>Cnemidophorus vanzoi</i>	Harvey et al. 2012 (See Crawford 2008)
Teiidae	<i>Contomastix lacertoides</i>	Reeder et al. 2002
Teiidae	<i>Contomastix vacariensis</i>	Feltrim and De Lema 2000
Teiidae	<i>Holcosus festivus</i>	Hower and Hedges 2003
Teiidae	<i>Holcosus quadrilineatus</i>	Harvey et al. 2012
Teiidae	<i>Kentropyx calcarata</i>	Werneck et al. 2009
Teiidae	<i>Kentropyx pelviceps</i>	Werneck et al. 2009
Teiidae	<i>Kentropyx striata</i>	Werneck et al. 2009
Teiidae	<i>Teius oculatus</i>	Reeder et al. 2002, taxonomy
Teiidae	<i>Teius teyou</i>	Reeder et al. 2002
Teiidae	<i>Tupinambis teguixin</i>	Giugliano et al. 2007, Fitzgerald et al. 1999
Trogonophiidae	<i>Trogonophis wiegmanni</i>	Macey et al. 2004, Taxonomy
Tropiduridae	<i>Microlophus albemarlensis</i>	Benavides et al. 2007

Family	species	Source for phylogenetic relationship
Tropiduridae	<i>Microlophus atacamensis</i>	Benavides et al. 2007
Tropiduridae	<i>Microlophus delanonis</i>	Benavides et al. 2007
Tropiduridae	<i>Microlophus habelii</i>	Benavides et al. 2007
Tropiduridae	<i>Microlophus quadrivittatus</i>	Benavides et al. 2007
Tropiduridae	<i>Plica plica</i>	Schlute et al. 2003, taxonomy
Tropiduridae	<i>Plica umbra</i>	Schlute et al. 2003, taxonomy
Tropiduridae	<i>Stenocercus chrysopygus</i>	Torres-Carvajal 2007
Tropiduridae	<i>Stenocercus dumerilii</i>	Torres-Carvajal 2007
Tropiduridae	<i>Tropidurus etheridgei</i>	Frost et al. 2001
Tropiduridae	<i>Tropidurus hispidus</i>	Frost et al. 2001
Tropiduridae	<i>Tropidurus semitaeniatus</i>	Frost et al. 2001
Tropiduridae	<i>Tropidurus spinulosus</i>	Frost et al. 2001
Tropiduridae	<i>Tropidurus torquatus</i>	Frost et al. 2001
Tropiduridae	<i>Uracentron flaviceps</i>	Schlute et al. 2003, Warne & Charnov 2008, taxonomy
Varanidae	<i>Varanus acanthurus</i>	Thompson et al. 2008, Vidal et al. 2012
Varanidae	<i>Varanus albigularis</i>	Collar et al. 2011
Varanidae	<i>Varanus beccarii</i>	Ast 2001, Collar et al. 2011
Varanidae	<i>Varanus bengalensis</i>	Ast 2001
Varanidae	<i>Varanus boehmei</i>	Ziegler et al. 2007
Varanidae	<i>Varanus caerulivirens</i>	Ziegler et al. 2007
Varanidae	<i>Varanus caudolineatus</i>	Vidal et al. 2012
Varanidae	<i>Varanus cerambonensis</i>	Ziegler et al. 2007
Varanidae	<i>Varanus flavescens</i>	Ast 2001
Varanidae	<i>Varanus giganteus</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus gilleni</i>	Vidal et al. 2012
Varanidae	<i>Varanus glauerti</i>	Thompson et al. 2008, Vidal et al. 2012
Varanidae	<i>Varanus gouldii</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus griseus</i>	Collar et al. 2011, Portik and Papenfuss 2012
Varanidae	<i>Varanus kingorum</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus komodoensis</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus liruensis</i>	Koch et al. 2009
Varanidae	<i>Varanus mabitang</i>	Ziegler et al. 2005
Varanidae	<i>Varanus macraei</i>	Ziegler et al. 2007
Varanidae	<i>Varanus melinus</i>	Ast 2001
Varanidae	<i>Varanus mertensi</i>	Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus niloticus</i>	Ast 2001, Collar et al. 2011
Varanidae	<i>Varanus nuchalis</i>	considered subspecies of salvator
Varanidae	<i>Varanus ornatus</i>	considered subspecies of niloticus
Varanidae	<i>Varanus panoptes</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus prasinus</i>	Ast 2001, Collar et al. 2011
Varanidae	<i>Varanus rosenbergi</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus rudicollis</i>	Ast 2001, Vidal et al. 2012
Varanidae	<i>Varanus salvator</i>	Ast 2001, Vidal et al. 2012
Varanidae	<i>Varanus scalaris</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus spenceri</i>	Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus tristis</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Varanidae	<i>Varanus varius</i>	Thompson et al. 2008, Collar et al. 2011, Vidal et al. 2012
Xantusiidae	<i>Lepidophyma flavimaculatum</i>	taxonomy
Xantusiidae	<i>Xantusia arizonae</i>	Leavitt et al. 2007

Family	species	Source for phylogenetic relationship
Xantusiidae	<i>Xantusia henshawi</i>	Leavitt et al. 2007
Xantusiidae	<i>Xantusia riversiana</i>	Leavitt et al. 2007
Xantusiidae	<i>Xantusia vigilis</i>	Leavitt et al. 2007
Xantusiidae	<i>Xantusia wigginsi</i>	Leavitt et al. 2007, Sinclair et al. 2004
Xenosauridae	<i>Xenosaurus grandis</i>	Bhullar 2011

H. Literature source for the phylogenetic tree

- Ahmadzadeh, F., Carretero, M.A., Harris, D.J., Perera, A. & Bohme, W. (2012) A molecular phylogeny of the eastern group of ocellated lizard genus *Timon* (Sauria: Lacertidae) based on mitochondrial and nuclear DNA sequences. *Amphibia-Reptilia*, **33**, 1-10.
- Alfoldi, J., Di Palma, F., Grabherr, M., Williams, C., Kong, L., Mauceli, E., Russell, P., Lowe, C. B., Glor, R. E., Jaffe, J. D., Ray, D. A., Boissinot, S., Shedlock, A. M., Botka, C., Castoe, T. A., Colbourne, J. K., Fujita, M. K., Moreno, R. G., ten Hallers, B. F., Haussler, D., Heger, A., Heiman, D., Janes, D. E., Johnson, J., de Jong, P. J., Koriabine, M. Y., Lara, M., Novick, P. A., Organ, C. L., Peach, S. E., Poe, S., Pollock, D. D., de Queiroz, K., Sanger, T., Searle, S., Smith, J. D., Smith, Z., Swofford, R., Turner-Maier, J., Wade, J., Young, S., Zadissa, A., Edwards, S. V., Glenn, T. C., Schneider, C. J., Losos, J. B., Lander, E. S., Breen, M., Ponting, C. P. & Lindblad-Toh, K. (2011) The genome of the green anole lizard and a comparative analysis with birds and mammals. *Nature*, **477**, 587-591.
- Amat, F. (2008) Exploring female reproductive tactics: trade-offs between clutch size, egg mass and newborn size in lacertid lizards. *Herpetological Journal*, **18**, 147-153.
- Arnold, E.N., Arribas, O. & Carranza, S. (2007) Systematics of the Palearctic and Oriental lizard tribe *Lacertini* (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa*, **1430**, 1-86.
- Arnold, E.N. (2009) Relationships, evolution and biogeography of Semaphore geckos, *Pristurus* (Squamata, Sphaerodactylidae) based on morphology. *Zootaxa*, **2060**, 1-21.
- Ast, J.C. (2001) Mitochondrial DNA evidence and evolution in Varanoidea (Squamata). *Cladistics*, **17**, 211-226.
- Austin, C.C. (1998) Phylogenetic relationships of *Lipinia* (Scincidae) from New Guinea based on DNA sequence variation from the mitochondrial 12S rRNA and Nuclear C-MOS genes. *Hamadryad*, **23**, 93-102.
- Austin, C.C., Das, I. & de Silva, A. (2004) Higher-level molecular phylogenetic relationships of the endemic genus *Lankascincus* from Sri Lanka based on nuclear DNA sequences. *Lyriocephalus*, **5**, 11-22.
- Austin, J.J.A., E. N. (2006) Using ancient and recent DNA to explore relationships of extinct and endangered *Leiopisma* skinks (Reptilia: Scincidae) in the Mascarene islands. *Molecular Phylogenetics and Evolution*, **39**, 503-511.
- Austin, C.C., Rittmeyer, E. N., Richards, S. J. & Zug, G. R. (2010) Phylogeny, historical biogeography and body size evolution in Pacific island crocodile skinks *Tribolonotus* (Squamata; Scincidae). *Molecular Phylogenetics and Evolution*, **57**, 227-236.
- Bauer, A.M., Jackman, T. R., Greenbaum, E., Giri, V. B. & de Silva, A. (2010) South Asia

- supports a major endemic radiation of *Hemidactylus* geckos. *Molecular Phylogenetics and Evolution*, **57**, 343-352.
- Bauer, A.M., Jackman, T. R., Sadlier, R. A. & Whitaker, A. H. (2012) Revision of the giant geckos of New Caledonia (Reptilia: Diplodactylidae: *Rhacodactylus*). *Zootaxa*, **3404**, 1-52.
- Benavides, E., Baum, R., McClellan, D. & Sites, J. W. (2007) Molecular phylogenetics of the lizard genus *Microlophus* (Squamata: Tropiduridae): aligning and retrieving indel signal from nuclear introns. *Systematic Biology*, **56**, 776-797.
- Bhullar, B.-A.S. (2011) The power and utility of morphological characters in systematics: A fully resolved phylogeny of *Xenosaurus* and its fossil relatives (Squamata: Anguimorpha). *Bulletin of the Museum of Comparative Zoology*, **160**, 65-181.
- Blair, C., Mendez de la Cruz, F. R., Ngo, A., Lindell, J., Lathrop, A. & Murphy, R. W. (2009) Molecular phylogenetics and taxonomy of leaf-toed geckos (Phyllodactylidae: *Phyllodactylus*) inhabiting the peninsula of Baja California. *Zootaxa*, **2027**, 28-42.
- Bonine, K.E., Gleeson, T. T. & Garland, T. (2005) Muscle fiber-type variation in lizards (Squamata) and phylogenetic reconstruction of hypothesized ancestral states. *Journal of Experimental Biology*, **208**, 4529-4547.
- Branch, W.R., Tolley, K. A. & Tilbury, C. R. (2006) A new dwarf chameleon (Sauria: Bradypodion Figzinger, 1843) from the Cape Fold Mountains, South Africa. *African Journal of Herpetology*, **55**, 123-141.
- Brandley, M.C., Wang, Y., Guo, X., de Oca, A. N. M., Feria-Ortiz, M., Hikida, T. & Ota, H. (2011) Accommodating heterogenous rates of evolution in molecular divergence dating methods: an example using Intercontinental dispersal of *Plestiodon* (*Eumeces*) lizards. *Systematic Biology*, **60**, 3-15.
- Brandley, M.C., Ota, H., Hikida, T., De Oca, A. N., Feria-Ortiz, M., Guo, X. & Wang, Y. (2012) The phylogenetic systematics of blue-tailed skinks (*Plestiodon*) and the family Scincidae. *Zoological Journal of the Linnean Society*, **165**, 163-189.
- Breitman, M.F., Avila, L. J., Sites, J. W. & Morando, M. (2011) Lizards from the end of the world: Phylogenetic relationships of the *Liolaemus lineomaculatus* section (Squamata: Iguania: Liolaemini). *Molecular Phylogenetics and Evolution*, **59**, 364-376.
- Brown, J.H., Fox, B. J. & Kelt, D. A. (2000) Assembly rules: desert rodent communities are structured at scales from local to continental. *American Naturalist*, **156**, 314-321.
- Brown, D., Wilmer, J. W. & Macdonald, S. (2012) A revision of *Strophurus taenicauda* (Squamata; Diplodactylidae) with the description of two new subspecies from central Queensland and a southerly range extension. *Zootaxa*, **3243**, 1-28.
- Brown, D., Wilmer, J. W. & Macdonald, S. (2012) A revision of *Strophurus taenicauda* (Squamata; Diplodactylidae) with the description of two new subspecies from central Queensland and a southerly range extension. *Zootaxa*, **3243**, 1-28.
- Bryson, R.W.R., B. R. (2012) Tracing the origins of widespread highland species: a case of Neogene diversification across the Mexican sierras in an endemic lizard. *Biological Journal of the Linnean Society*, **105**, 382-394.
- Camargo, A., Avila, L. J., Morando, M., & Sites, J. W. (2012) Accuracy and precision of species trees: effects of locus, individual, and base pair sampling on inference of species trees in lizards of the *Liolaemus darwini* group (Squamata, Liolaemidae). *Systematic Biology*, **61**, 272-288.
- Carranza, S.A., E. N., Mateo, J. A. & Geniez, P. (2002) Relationships and evolution of the North African geckos, *Geckonia* and *Tarentola* (Reptilia: Gekkonidae), based on

- mitochondrial and nuclear DNA sequences. *Molecular Phylogenetics and Evolution*, **23**, 244-256.
- Carranza, S., Harris, D. J., Arnold, E. N., Batista, V. & Gonzalez de la Vega, J. P. (2006) Phylogeography of the lacertid lizard, *Psammmodromus algirus*, in Iberia and across the Strait of Gibraltar. *Journal of Biogeography*, **33**, 1279-1288.
- Carranza, S.A., E. N. (2006) Systematics, biogeography, and evolution of *Hemidactylus* geckos (Reptilia: Gekkonidae) elucidated using mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution*, **38**, 531-545.
- Carranza, S., Arnold, E. N., Geniez, P., Roca, J. & Mateo, J. A. (2008) Radiation, multiple dispersal and parallelism in the skinks, *Chalcides* and *Sphenops* (Squamata: Scincidae), with comments on *Scincus* and *Scincopus* and the age of the Sahara Desert. *Molecular Phylogenetics and Evolution*, **46**, 1071-1094.
- Carranza, S.A., E. N. (2012) A review of the geckos of the genus *Hemidactylus* (Squamata: Gekkonidae) from Oman based on morphology, mitochondrial and nuclear data, with descriptions of eight new species. *Zootaxa*, **3378**, 1-95.
- Castaneda, M.d.R.d.Q., K. (2011) Phylogenetic relationships of the dactyloa clade of *Anolis* lizards based on nuclear and mitochondrial DNA sequence data. *Molecular Phylogenetics and Evolution*, **61**, 784-800.
- Castiglia, R., Annesi, F., Bezerra, A. M. R., Garcia, A. & Flores-Villela, O. (2010) Cytotaxonomy and DNA taxonomy of lizards (Squamata, Sauria) from a tropical dry forest in the Chamela-Cuixmala Biosphere Reserve on the coast of Jalisco, Mexico. *Zootaxa*, **2508**, 1-29.
- Castoe, T.A., Doan, T. M. & Parkinson, C. L. (2004) Data partitions and complex models in Bayesian analysis: the phylogeny of Gymnophthalmid lizards. *Systematic Biology*, **53**, 448-469.
- Chapple, D.G., Daugherty, C. H. & Ritchie, P. A. (2008) Comparative phylogeography reveals pre-decline population structure of New Zealand *Cyclodina* (Reptilia: Scincidae) species. *Biological Journal of the Linnean Society*, **95**, 388-408.
- Chapple, D.G., Ritchie, P. A. & Daugherty, C. H. (2009) Origin, diversification, and systematics of the New Zealand skink fauna (Reptilia: Scincidae). *Molecular Phylogenetics and Evolution*, **52**, 470-487.
- Chapple, D.G., Bell, T. P., Chapple, S. N. J., Miller, K. A., Daugherty, C. H. & Patterson, C. H. (2011) Phylogeography and taxonomic revision of the New Zealand cryptic skink (*Oligosoma inconspicuum*; Reptilia: Scincidae) species complex. *Zootaxa*, **2782**, 1-33.
- Chou, C.E. (2007) *Phylogeography of the endemic species Japalura brevipes of Taiwan based on cytochrome b sequences and morphology*. National Sun Yat-sen University, Kaohsiung City.
- Collar, D.C., Schulte, J. A. & Losos, J. B. (2011) Evolution of extreme body size disparity in monitor lizards (*Varanus*). *Evolution*, **65**, 2664-2680.
- Conradie, W., Measey, G. J., Branch, W. R. & Tolley, K. A. (2012) Revised phylogeny of African sand lizards (*Pedioplanis*), with the description of two new species from southwestern Angola. *African Journal of Herpetology*, **61**, 91-112.
- Conroy, C.J., Bryson, R. W., Lazcano, D. & Knight, A. (2005) Phylogenetic placement of the pygmy alligator lizard based on mitochondrial DNA. *Journal of Herpetology*, **39**, 142-147.
- Daniels, S.R., Heideman, N. J. L., Hendricks, M. G. J. & Crandall, K. A. (2006) Taxonomic subdivisions within the fossorial skink subfamily Acontinae (Squamata: Scincidae)

- reconsidered: a multilocus perspective. *Zoologica Scripta*, **35**, 353-362.
- Datta-Roy, A., Singh, M., Srinivasulu, C. & Karanth, K. P. (2012) Phylogeny of the Asian *Eutropis* (Squamata: Scincidae) reveals an into India endemic Indian radiation. *Molecular Phylogenetics and Evolution*, **63**, 817-824.
- Doan, T.M. (2003) A new phylogenetic classification for the gymnophthalmid genera *Cercosaura*, *Pantodactylus* and *Prionodactylus* (Reptilia: Squamata). *Zoological Journal of the Linnean Society*, **137**, 101-115.
- Dolman, G.H., A. F. (2008) Combined mitochondrial and nuclear data enhance resolution of a rapid radiation of Australian rainbow skinks (Scincidae: *Carlia*). *Molecular Phylogenetics and Evolution*, **49**, 782-794.
- Doughty, P., Maryan, B., Melville, J. & Austin, J. (2007) A new species of *Ctenophorus* (Lacertilia: Agamidae) from Lake Disappointment, Western Australia. *Herpetologica*, **63**, 72-86.
- Doughty, P., Kealley, L. & Donnellan, S. C. (2011) Revision of the pygmy spiny-tailed skinks (*Egernia depressa* species-group) from Western Australia, with descriptions of three new species. *Records Of The Western Australian Museum*, **26**, 115-137.
- Edwards, D.L.M., J. (2011) Extensive phylogeographic and morphological diversity in *Diporiphora nobbi* (Agamidae) leads to a taxonomic review and a new species description. *Journal of Herpetology*, **45**, 530-546.
- Feldman, C.R., Flores-Villela, O. & Papenfuss, T. J. (2011) Phylogeny, biogeography, and display evolution in the tree and brush lizard genus *Urosaurus* (Squamata: Phrynosomatidae). *Molecular Phylogenetics and Evolution*, **61**, 714-725.
- Feltrim, A.C.D.L., T. (2000) Uma nova especie de *Cnemidophorus* Wagler, 1830 do estado do Rio Grande do sul Brasil (Sauria, Teiidae). *Biociencias, Porto Alegre*, **8**, 103-114.
- Fitzgerald, L.A., Cook, J. A. & Aquino, A. L. (1999) Phylogenetics and conservation of *Tupinambis* (Sauria: Teiidae). *Copeia*, **1999**, 894-905.
- Frost, D.R., Etheridge, R., Janies, D. & Titus, T. A. (2001) Total evidence, sequence alignment, evolution of Polychrotid lizards, and a reclassification of the Iguania (Squamata: Iguania). *American Museum Novitates*, **3343**, 1-38.
- Fujita, M.K.P., T. J. (2011) Molecular systematics of *Stenodactylus* (Gekkonidae), an Afro-Arabian gecko species complex. *Molecular Phylogenetics and Evolution*, **58**, 71-75.
- Gamble, T., Bauer, A. M. Greenbaum, E. & Jackman, T. R. (2008) Evidence for Gondwanan vicariance in an ancient clade of gecko lizards. *Journal of Biogeography*, **35**, 88-104.
- Gamble, T., Bauer, A. M. Greenbaum, E. & Jackman, T. R. (2008) Out of the blue: a novel, trans-Atlantic clade of geckos (Gekkota, Squamata). *Zoologica Scripta*, **37**, 355-366.
- Gamble, T., Bauer, A. M., Colli, G. R. Greenbaum, E., Vitt, L. J. & Simons, A. M. (2011) Coming to America: multiple origins of New World geckos. *Journal of Evolutionary Biology*, **24**, 231-244.
- Gamble, T., Colli, G. R., Rodrigues, M. T., Werneck, F. P. & Simons, A. M. (2012) Phylogeny and cryptic diversity in geckos (Phyllopezus; Phyllodactylidae; Gekkota) from South America's open biomes. *Molecular Phylogenetics and Evolution*, **62**, 943-953.
- Gardner, M.G., Hugall, A. F., Donnellan, S. C., Hutchinson, M. N. & Foster, R. (2008) Molecular systematics of social skinks: phylogeny and taxonomy of the *Egernia* group (Reptilia: Scincidae). *Zoological Journal of the Linnean Society*, **154**, 781-794.
- Geniez, P., Padial, J. M. & Crochet, P-A. (2011) Systematics of north African *Agama* (Reptilia: Agamidae): a new species from the central Saharan mountains. *Zootaxa*, **3098**, 26-46.

- Geurgas, S.R., Rodrigues, M. T. & Moritz, C. (2008) The genus *Coleodactylus* (Sphaerodactylinae, Gekkota) revisited: a molecular phylogenetic perspective. *Molecular Phylogenetics and Evolution*, **49**, 92-101.
- Giugliano, L.G., Collevatti, R. G. & Colli, G. R. (2007) Molecular dating and phylogenetic relationships among Teiidae (Squamata) inferred by molecular and morphological data. *Molecular Phylogenetics and Evolution*, **45**, 168-179.
- Godinho, R., Crespo, E. G., Ferrand, N. & Harris, D. J. (2005) Phylogeny and evolution of the green lizards, *Lacerta* spp. (Squamata: Lacertidae) based on mitochondrial and nuclear DNA sequences. *Amphibia-Reptilia*, **26**, 271-285.
- Goicoechea, N., Padial, J. M., Chaparro, J. C., Castroviejo-Fisher, S. & De la Riva, I. (2012) Molecular phylogenetics, species diversity, and biogeography of the Andean lizards of the genus *Proctoporus* (Squamata: Gymnophthalmidae). *Molecular Phylogenetics and Evolution*, **65**, 953-964.
- Gomez-Diaz, E., Sindaco, R., Pupin, F., Fasola, M. & Carranza, S. (2012) Origin and in situ diversification in *Hemidactylus* geckos of the Socotra Archipelago. *Molecular Ecology*, **16**, 4074-4092.
- Goncalves, D.V., Brito, J. C., Crochet, P-A., Geniez, P., Padial, J. M. & Harris, D. J. (2012) Phylogeny of North African *Agama* lizards (Reptilia: Agamidae) and the role of the Sahara desert in vertebrate speciation. *Molecular Phylogenetics and Evolution*, **64**, 582-591.
- Good, D.A. (1994) Species limits in the genus *Gerrhonotus* (Squamata: Anguinae). *Herpetological Monographs*, **8**, 180-202.
- Greer, A.E.W., G. D. F. (2001) Comments on the scincid lizard genus *Ophiomorus* with a cladistic analysis of the species. *Hamadryad*, **26**, 261-271.
- Grismer, L.L., Grismer, J. L., Wood, P. L., Ngo, V. T., Neang, T. & Chan, K. O. (2011) Herpetology on the fringes of the Sunda Shelf: a discussion of discovery, taxonomy, and biogeography. *Bonner Zoologische Monographien*, **57**, 57-97.
- Guo, X.W., Y. (2007) Partitioned Bayesian analyses, dispersal vicariance analysis, and the biogeography of Chinese toad-headed lizards (Agamidae: *Phrynocephalus*): a re-evaluation. *Molecular Phylogenetics and Evolution*, **45**, 643-662.
- Guo, X., Dai, X., Chen, D., Papenfuss, T. J., Ananjeva, N. B., Melnikov, D. A. & Wang, Y. (2011) Phylogeny and divergence times of some racerunner lizards (Lacertidae: *Eremias*) inferred from mitochondrial 16S rRNA gene segments. *Molecular Phylogenetics and Evolution*, **61**, 400-412.
- Guo, X., Liu, L. & Wang, Y. (2012) Phylogeography of the *Phrynocephalus vlangalii* species complex in the upper reaches of the Yellow River inferred from mtDNA ND4-tRNA^{LEU} segments. *Asian Herpetological Research*, **3**, 52-68.
- Gvozdkik, V., Jandzik, D., Lymberakis, P., Jablonski, D. & Moravec, J. (2010) Slow worm, *Anguis fragilis* (Reptilia: Anguinae) as a species complex: genetic structure reveals deep divergences. *Molecular Phylogenetics and Evolution*, **55**, 460-472.
- Hamilton, A.M. (2008) *Species boundaries, biogeography, and intra-archipelago genetic variation within the Emoia samoensis species group in the Vanuatu Archipelago and Oceania*. Louisiana State University,
- Harris, D.J., Arnold, E. N. & Thomas, R. H. (1999) A phylogeny of the European lizard genus *Algyroides* (Reptilia: Lacertidae) based on DNA sequences, with comments on the evolution of the group. *Journal of Zoology*, **249**, 49-60.
- Harris, D.J.A., E. N. (2000) Elucidation of the relationships of spiny-footed lizards,

- Acanthodactylus* spp. (Reptilia: Lacertidae) using mitochondrial DNA sequence, with comments on their biogeography and evolution. *Journal of Zoology*, **252**, 351-362.
- Harris, D.J., Pinho, C., Carretero, M. A., Corti, C. & Bohme, W. (2005) Determination of genetic diversity within the insular lizard *Podarcis tiliguerta* using mtDNA sequence data, with a reassessment of the phylogeny of *Podarcis*. *Amphibia-Reptilia*, **26**, 401-407.
- Harris, D.J. (2008) Taxonomic inflation and red lists: how have Mediterranean reptile conservation issues been affected? *Conference Proceedings*, **6**, 28.
- Hedges, S.B.C., C. E. (2012) A new skink fauna from Caribbean islands (Squamata, Mabuyidae, Mabuyinae). *Zootaxa*, **3288**, 1-244.
- Heinicke, M.P., Adderly, L. M., Bauer, A. M. & Jackman, T. R. (2011) A long-known new species of gecko allied to *Pachydactylus bicolor* (Squamata: Gekkonidae) from the central Namibian coast. *African Journal of Herpetology*, **60**, 113-129.
- Honda, M., Ota, H., Murphy, R. W. & Hikida, T. (2006) Phylogeny and biogeography of water skinks of the genus *Tropidophorus* (Reptilia: Scincidae): a molecular approach. *Zoologica Scripta*, **35**, 85-95.
- Hower, L.M.H., S. B. (2003) Molecular phylogeny and biogeography of West Indian teiid lizards of the genus *Ameiva*. *Caribbean Journal of Science*, **39**, 298-306.
- Hugall, A.F., Foster, R. & Lee, M. S. Y. (2007) Calibration choice, rate smoothing, and the pattern of tetrapod diversification according to the long nuclear gene RAG-1. *Systematic Biology*, **56**, 543-563.
- Hugall, A.F., Foster, R., Hutchinson, M. & Lee, S. Y. (2008) Phylogeny of Australasian agamid lizards based on nuclear and mitochondrial genes: implications for morphological evolution and biogeography. *Biological Journal of the Linnean Society*, **93**, 343-358.
- Hutchinson, M.N.D., S. C. (1992) Taxonomy and genetic variation in the Australian lizards of the genus *Pseudemoia* (Scincidae: Lygosominae). *Journal of Natural History*, **26**, 215-264.
- Jennings, W.B., Pianka, E. R. & Donnellan, S. (2003) Systematics of the lizard family Pygopodidae with implications for the diversification of Australian temperate biotas. *Systematic Biology*, **52**, 757-780.
- Jesus, J., Harris, D. J. & Brehm, A. (2007) Relationships of *Afroablepharus* Greer, 1974 skinks from the Gulf of Guinea islands based on mitochondrial and nuclear DNA: Patterns of colonization and comments on taxonomy. *Molecular Phylogenetics and Evolution*, **45**, 904-914.
- Ji, X., Wang, Y. & Wang, Z. (2009) New species of *Phrynocephalus* (Squamata, Agamidae) from Qinghai, Northwest China. *Zootaxa*, **2988**, 61-68.
- Jin, Y.-T., Brown, R. P. & Liu, N-F. (2008) Cladogenesis and phylogeography of the lizard *Phrynocephalus vlangalii* (Agamidae) on the Tibetan plateau. *Molecular Ecology*, **17**, 1971-1982.
- Joger, U.M., W. (2002) A new species of *Mesalina* from Abd al-Kuri, Socotra archipelago, Yemen, and a preliminary molecular phylogeny for the genus *Mesalina*. *Fauna of Saudi Arabia*, **19**, 497-505.
- Kapli, P., Lymberakis, P., Poulakakis, N., Mantziou, G., Parmakelis, A. & Mylonas, M. (2008) Molecular phylogeny of three *Mesalina* (Reptilia: Lacertidae) species (*M. guttulata*, *M. brevirostris* and *M. bahaeldini*) from North Africa and the Middle East: another case of paraphyly? *Molecular Phylogenetics and Evolution*, **49**, 102-110.
- Kay, G.M.K., J. S. (2012) Molecular phylogeny and morphological revision of the *Ctenotus labillardieri* (Reptilia: Squamata: Scincidae) species group and a new species of immediate

- conservation concern in the southwestern Australian biodiversity hotspot. *Zootaxa*, **3390**, 1-18.
- Keogh, J.S., Edwards, D. L., Fisher, R. N. & Harlow, P. S. (2008) Molecular and morphological analysis of the critically endangered Fijian iguanas reveals cryptic diversity and a complex biogeographic history. *Philosophical Transactions of the Royal Society of London B.*, **363**, 3413-3426.
- Koch, A., Arida, E., Riyanto, A. & Bohme, W. (2009) Islands between the realms: A revised checklist of the herpetofauna of the Talaud Archipelago, Indonesia, with a discussion about its biogeographic affinities. *Bonner Zoologische Beiträge*, **56**, 107-129.
- Kratochvil, L.F., D. (2002) Body size, male combat and the evolution of sexual dimorphism in eublepharid geckos (Squamata: Eublepharidae). *Biological Journal of the Linnean Society*, **76**, 303-314.
- Labra, A., Pienaar, J. & Hansen, T. F. (2009) Evolution of thermal physiology in *Liolaemus* lizards: adaptation, phylogenetic inertia, and niche tracking. *American Naturalist*, **174**, 204-220.
- Lamb, T.B., A. M. (2003) Meroles revisited: complementary systematic inference from additional mitochondrial genes and complete taxon sampling of southern Africa's desert lizards. *Molecular Phylogenetics and Evolution*, **29**, 360-364.
- Lamb, T., Meeker, A. M., Bauer, A. M. & Branch, W. R. (2003) On the systematic status of the desert plated lizard (*Angolosaurus skoogi*): phylogenetic inference from DNA sequence analysis of the African Gerrhosauridae. *Biological Journal of the Linnean Society*, **78**, 253-261.
- Lamb, T.B., A. M. (2006) Footprints in the sand: independent reduction of subdigital lamellae in the Namiba-Kalahari burrowing geckos. *Proceedings of the Royal Society of London B: Biological Sciences*, **273**, 855-864.
- Leache, A.D.M., J. A. (2006) Phylogenetic relationships of horned lizards (*Phrynosoma*) based on nuclear and mitochondrial data: Evidence for a misleading mitochondrial gene tree. *Molecular Phylogenetics and Evolution*, **39**, 628-644.
- Leache, A.D. (2010) Species trees for spiny lizards (Genus *Sceloporus*): Identifying points of concordance and conflict between nuclear and mitochondrial data. *Molecular Phylogenetics and Evolution*, **54**, 162-171.
- Leavitt, D.H., Bezy, R. L., Crandall, K. A. & Sites, J. W. (2007) Multi-locus DNA sequence data reveal a history of deep cryptic vicariance and habitat-driven convergence in the desert night lizard *Xantusia vigilis* species complex (Squamata: Xantusiidae). *Molecular Ecology*, **16**, 4455-4481.
- Lee, J.E., Janion, C., Marais, E., van Vuuren, B. J. & Chown, S. L. (2009) Physiological tolerances account for range limits and abundance structure in an invasive slug. *Proceedings of the Royal Society of London B: Biological Sciences*, **276**, 1459-1468.
- Lin, S.M., Chen, C. A. & Lue, K-Y. (2002) Molecular phylogeny and biogeography of the grass lizards genus *Takydromus* (Reptilia: Lacertidae) of East Asia. *Molecular Phylogenetics and Evolution*, **22**, 276-288.
- Linkem, C.W., Diesmos, A. C. & Brown, R. M. (2011) Molecular systematics of the Philippine forest skinks (Squamata: Scincidae: *Sphenomorphus*): testing morphological hypotheses of interspecific relationships. *Zoological Journal of the Linnean Society*, **163**, 1217-1243.
- Lobo, F., Abdala, C. & Valdecantos, S. (2012) Morphological diversity and phylogenetic relationships within a South-American clade of iguanian lizards (Liolaemidae:

- Phymaturus*). *Zootaxa*, **3315**, 1-41.
- Lue, K.-Y.L., S-M. (2008) Two new cryptic species of *Takydromus* (Squamata: Lacertidae) from Taiwan. *Herpetologica*, **64**, 379-395.
- Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. & Mylonas, M. (2008) Two new species of *Podarcis* (Squamata; Lacertidae) from Greece. *Systematics and Biodiversity*, **6**, 307-318.
- Macey, J.R., Schulte, J. A., Larson, I. A., Tuniyev, B. S., Orlov, N. & Papenfuss, T. J. (1999) Molecular phylogenetics, tRNA evolution, and historical biogeography in anguid lizards and related taxonomic families. *Molecular Phylogenetics and Evolution*, **12**, 250-272.
- Macey, J.R., Schulte, J. A., Larson, I. A., Ananjeva, N. B., Wang, Y., Pethiyagoda, R., Rastegar-Pouyani, N. & Papenfuss, T. J. (2004) Phylogenetic relationships among amphisbaenian reptiles based on complete mitochondrial genomic sequences. *Molecular Phylogenetics and Evolution*, **33**, 22-31.
- Macey, J.R., Schulte, J. A., Strasburg, J. L., Brisson, J. A., Larson, A., Ananjeva, N. B., Wang, Y., Parham, J. F. & Papenfuss, T. J. (2006) Assembly of the eastern North American herpetofauna: New evidence from lizards and frogs. *Biology Letters*, **2**, 388-392.
- Makokha, J.S., Bauer, A. A., Mayer, W. & Matthee, C. A. (2007) Nuclear and mtDNA-based phylogeny of southern African sand lizards, *Pedioplanis* (Sauria: Lacertidae). *Molecular Phylogenetics and Evolution*, **44**, 622-633.
- Malone, C.L., Wheeler, T., Taylor, J. F. & Davis, S. K. (2000) Phylogeography of the Caribbean rock iguana (*Cyclura*): implications for conservation and insights on the biogeographic history of the West Indies. *Molecular Phylogenetics and Evolution*, **17**, 269-279.
- Martinez-Mendez, N. and Mendez-de la Cruz, F. R. (2007) Molecular phylogeny of the *Sceloporus torquatus* species-group (Squamata: Phrynosomatidae). *Zootaxa*, **1609**, 53-68.
- Matthee, C.A., Tilbury, C. R. & Townsend, T. (2004) A phylogenetic review of the African leaf chameleons: genus *Rhampholeon* (Chamaeleonidae): the role of vicariance and climate change in speciation. *Proceedings of the Royal Society of London B: Biological Sciences*, **271**, 1967-1975.
- Mayer, W.P., M. (2007) The phylogeny of the family Lacertidae (Reptilia) based on nuclear DNA sequences: Convergent adaptations to arid habitats within the subfamily Eremiinae. *Molecular Phylogenetics and Evolution*, **44**, 1155-1163.
- McGuire, J.A.H., K. B. (2001) Phylogenetic systematics of Southeast Asian flying lizards (Iguania: Agamidae: *Draco*) as inferred from mitochondrial DNA sequence data. *Biological Journal of the Linnean Society*, **72**, 203-229.
- McGuire, J.A., Brown, R. M., Mumpuni, Riyanto, A. & Andayani, N. (2007) The flying lizards of the *Draco lineatus* group (Squamata: Iguania: Agamidae): a taxonomic revision with descriptions of two new species. *Herpetological Monographs*, **21**, 179-212.
- Mecke, S., Doughty, P. & Donnellan, S. C. (2009) A new species of *Eremiascincus* (Reptilia: Squamata: Scincidae) from the Great Sandy Desert and Pilbara Coast, Western Australia and reassignment of eight species from *Glaphyromorphus* to *Eremiascincus*. *Zootaxa*, **2246**, 1-20.
- Mediannikov, O., Trape, S. & Trape, J-F. (2012) Molecular study of the genus *Agama* (Squamata: Agamidae) in West Africa, with description of two new species and a review of the taxonomy, geographic distribution, and ecology of currently recognized species. *Russian Journal of Herpetology*, **19**, 115-142.

- Melville, J.S., R. (2000) Evolutionary relationships between morphology, performance and habitat openness in the lizard genus *Niveoscincus* (Scincidae: Lygosominae). *Biological Journal of the Linnean Society*, **70**, 667-683.
- Melville, J., Schulte, J. A. & Larson, A. (2001) A molecular phylogenetic study of ecological diversification in the Australian lizard genus *Ctenophorus*. *Journal of Experimental Zoology*, **291**, 339-353.
- Melville, J., Schulte, J. A. & Larson, A. (2004) A molecular study of phylogenetic relationships and evolution of antipredator strategies in the Australian *Diplodactylus* geckos, subgenus *Strophurus*. *Biological Journal of the Linnean Society*, **82**, 123-138.
- Melville, J., Shoo, L. P. & Doughty, P. (2008) Phylogenetic relationships of the heath dragons (*Rankinia adelaidensis* and *R. parviceps*) from the south-western Australian biodiversity hotspot. *Australian Journal of Zoology*, **56**, 159-171.
- Melville, J., Hale, J., Mantziou, G., Ananjeva, N. B., Milto, K. & Clemann, N. (2009) Historical biogeography, phylogenetic relationships and intraspecific diversity of agamid lizards in the Central Asian deserts of Kazakhstan and Uzbekistan. *Molecular Phylogenetics and Evolution*, **53**, 99-112.
- Melville, J., Ritchie, E. G., Chapple, S. N. J., Glor, R. E. & Schulte, J. A. (2011) Evolutionary origins and diversification of dragon lizards in Australia's tropical savannas. *Molecular Phylogenetics and Evolution*, **58**, 257-270.
- Miralles, A., Barrio-Amoros, C. L., Rivas, G. & Chaparro-Auza, J. C. (2006) Speciation in the Varzea flooded forest: a new *Mabuya* (Squamata, Scincidae) from Western Amazonia. *Zootaxa*, **1188**, 1-22.
- Miralles, A., Chaparro, J. C. & Harvey, M. B. (2009) Three rare and enigmatic South American skinks. *Zootaxa*, **2012**, 47-68.
- Miralles, A., Kohler, J., Glaw, F. & Vences, M. (2011) A molecular phylogeny of the *Madascincus polleni* species complex, with description of a new species of scincid lizard from the coastal dune area of northern Madagascar. *Zootaxa*, **2876**, 1-16.
- Morando, M., Avila, L. J., Turner, C. R. & Sites, J. W. (2007) Molecular evidence for a species complex in the patagonian lizard *Liolaemus bibronii* and phylogeography of the closely related *Liolaemus gracilis* (Squamata: Liolaemini). *Molecular Phylogenetics and Evolution*, **43**, 952-973.
- Morando, M., Avila, L. J., Perez, C. H. F., Hawkins, M. & Sites, J. W. (2013) A molecular phylogeny of the lizard genus *Phymaturus* (Squamata, Liolaemini): implications for species diversity and historical biogeography of southern South America. *Molecular Phylogenetics and Evolution*, **66**, 694-714.
- Moreno Azocar, D.L., Vanhooydonck, B., Bonino, M. F., Perotti, M. G., Abdala, C. S., Schulte, J. A. & Cruz, F. B. (2013) Chasing the Patagonian sun: comparative thermal biology of *Liolaemus* lizards. *Oecologia*, **171**, 773-788.
- Nguyen, T.Q., Stenke, R., Nguyen, H. X. & Ziegler, T. (2011) The terrestrial reptile fauna of the Biosphere Reserve Cat Ba Archipelago, Hai Phong, Vietnam. *Bonner Zoologische Monographien*, **57**, 99-115.
- Nicholson, K.E., Glor, R. E., Kolbe, J. J., Larson, A., Hedges, S. B. & Losos, J. B. (2005) Mainland colonization by island lizards. *Journal of Biogeography*, **32**, 929-938.
- Nielsen, S.V., Bauer, A. M., Jackman, T. R., Hitchmough, R. A. & Daugherty, C. H. (2011) New Zealand geckos (Diplodactylidae): Cryptic diversity in a post-Gondwanan lineage with trans-Tasman affinities. *Molecular Phylogenetics and Evolution*, **59**, 1-22.

- Okamoto, T.H., T. (2009) Three genetic lineages of the Japanese skink *Plestiodon japonicus* (Scincidae, Squamata) and the genetic composition of their contact zones. *Journal of Zoological Systematics and Evolutionary Research*, **47**, 181-188.
- Olave, M., Martinez, L. E., Avila, L. J., Sites, J. W. & Morando, M. (2011) Evidence of hybridization in the Argentinean lizards *Liolaemus gracilis* and *Liolaemus bibronii* (Iguania: Liolaemini): an integrative approach based on genes and morphology. *Molecular Phylogenetics and Evolution*, **61**, 381-391.
- Oliver, P.M., Adams, M., Lee, M. S. Y., Hutchinson, M. N. & Doughty, P. (2009) Cryptic diversity in vertebrates: molecular data double estimates of species diversity in a radiation of Australian lizards (*Diplodactylus*, Gekkota). *Proceedings of the Royal Society of London B: Biological Sciences*, **276**, 2001-2007.
- Oliver, P.M., Couper, P. & Amey, A. (2010) A new species of *Pygopus* (Pygopodidae; Gekkota; Squamata) from north-eastern Queensland. *Zootaxa*, **2578**, 47-61.
- Oliver, P.M.B., A. M. (2011) Systematics and evolution of the Australian knob-tail geckos (*Nephrurus*, Carphodactylidae, Gekkota): plesiomorphic grades and biome shifts through the Miocene. *Molecular Phylogenetics and Evolution*, **59**, 664-674.
- Oliver, P.M., Bauer, A. M., Greenbaum, E., Jackman, T. & Hobbie, T. (2012) Molecular phylogenetics of the arboreal Australian gecko genus *Oedura* Gray 1842 (Gekkota: Diplodactylidae): another plesiomorphic grade? *Molecular Phylogenetics and Evolution*, **63**, 255-264.
- Ota, H., Honda, M., Kobayashi, M., Sengoku, S. & Hikida, T. (1999) Phylogenetic relationships of eublepharid Geckos (Reptilia: Squamata): a molecular approach. *Zoological Science*, **16**, 659-666.
- Papenfuss, T.J., Jackman, T., Bauer, A. M., Stuart, B. L., Robinson, M. D. & Parham, J. F. (2009) Phylogenetic relationships among species in the sphaerodactylid lizard genus *Pristurus*. *Proceedings of the California Academy of Sciences*, **60**, 675-681.
- Pasachnik, S.A., Echternacht, A. C. & Fitzpatrick, B. M. (2010) Gene trees, species and species trees in the *Ctenosaura palearis* clade. *Conservation Genetics*, **11**, 1767-1781.
- Pavlicev, M.M., W. (2009) Fast radiation of the subfamily Lacertinae (Reptilia: Lacertidae): History or methodical artefact? *Molecular Phylogenetics and Evolution*, **52**, 727-734.
- Pellegrino, K.C.M., Rodrigues, M. T., Waite, A. N., Morando, M., Yassuda, Y. Y. & Sites, J. W. (2005) Phylogeography and species limits in the *Gymnodactylus darwini* complex (Gekkonidae, Squamata): genetic structure coincides with river systems in the Brazilian Atlantic Forest. *Biological Journal of the Linnean Society*, **85**, 13-26.
- Pepper, M., Doughty, P. & Keogh, J. S. (2006) Molecular phylogeny and phylogeography of the Australian *Diplodactylus stenodactylus* (Gekkota; Reptilia) species-group based on mitochondrial and nuclear genes reveals an ancient split between Pilbara and non-Pilbara *D. stenodactylus*. *Molecular Phylogenetics and Evolution*, **41**, 539-555.
- Pepper, M., Doughty, P., Hutchinson, M. N. & Keogh, J. S. (2011) Ancient drainages divide cryptic species in Australia's arid zone: Morphological and multi-gene evidence for four new species of beaked geckos (*Rhynchoedura*). *Molecular Phylogenetics and Evolution*, **61**, 810-822.
- Perera, A., Sampaio, F., Costa, S., Salvi, D. & Harris, D. J. (2012) Genetic variability and relationships within the skinks *Eumeces algeriensis* and *Eumeces schneideri* using mitochondrial markers. *African Journal of Herpetology*, **61**, 69-80.
- Petren, K.C., T. J. (1997) A phylogenetic analysis of body size evolution in chuckwallas

- (*Sauromalus*) and other iguanines. *Evolution*, **51**, 206-219.
- Petren, K.C., T. J. (2002) An updated mtDNA phylogeny for *Sauromalus* and implications for the evolution of gigantism. *edited book chapter*, 574-579.
- Pincheira-Donoso, D., Hodgson, D. J. & Tregenza, T. (2008) The evolution of body size under environmental gradients in ectotherms: why should Bergmann's rule apply to lizards? *BMC Evolutionary Biology*, **8**, 68.
- Pincheira-Donoso, D., Scolaro, J. A. & Sura, P. (2008) A monographic catalogue on the systematics and phylogeny of the South American iguanian lizard family Liolaemidae (Squamata, Iguania). *Zootaxa*, **1800**, 1-85.
- Pincheira-Donoso, D., Hodgson, D. J., Stipala, J. & Tregenza, T. (2009) A phylogenetic analysis of sex-specific evolution of ecological morphology in *Liolaemus* lizards. *Ecological Research*, **24**, 1223-1231.
- Poe, S. (2004) Phylogeny of anoles. *Herpetological Monographs*, **18**, 37-89.
- Pokorna, M., Rabova, M., Rab, P., Ferguson-Smith, M. A., Rens, W. & Kratochvil, L. (2010) Differentiation of sex chromosomes and karyotypic evolution in the eye-lid geckos (Squamata: Gekkota: Eublepharidae), a group with different modes of sex determination. *Chromosome Research*, **18**, 809-820.
- Portik, D.M.P., T. J. (2012) Monitors cross the Red Sea: The biogeographic history of *Varanus yemenensis*. *Molecular Phylogenetics and Evolution*, **62**, 561-565.
- Poulakakis, N., Lymberakis, P., Zouros, E., Pafilis, P., Valakos, E. D. & Mylonas, M. (2005) Phylogeography of Balkan wall lizard (*Podarcis taurica*) and its relatives inferred from mitochondrial DNA sequences. *Molecular Ecology*, **14**, 2433-2443.
- Rabosky, D.L., Donnellan, S. C., Talaba, A. L. & Lovette, I. J. (2007) Exceptional among-lineage variation in diversification rates during the radiation of Australia's most diverse vertebrate clade. *Proceedings of the Royal Society of London B: Biological Sciences*, **274**, 2915-2923.
- Rato, C., Carranza, S. & Harris, D. J. (2012) Evolutionary history of the genus *Tarentola* (Gekkota: Phyllodactylidae) from the Mediterranean Basin, estimated using multilocus sequence data. *BMC Evolutionary Biology*, **12**, 14.
- Raxworthy, C.J., Forstner, M. R. J. & Nussbaum, R. A. (2002) Chameleon radiation by oceanic dispersal. *Nature*, **415**, 784-787.
- Reeder, T.W., Cole, C. J. & Dessauer, H. C. (2002) Phylogenetic relationships of whiptail lizards of the genus *Cnemidophorus* (Squamata: Teiidae): A test of monophyly, reevaluation of karyotypic evolution, and review of hybrid origins. *American Museum Novitates*, **3365**, 1-61.
- Reeder, T.W. (2003) A phylogeny of the Australian *Sphenomorphus* group (Scincidae: Squamata) and the phylogenetic placement of the crocodile skinks (*Tribolonotus*): Bayesian approaches to assessing congruence and obtaining confidence in maximum likelihood inferred relationships. *Molecular Phylogenetics and Evolution*, **27**, 384-397.
- Reeder, T.W.R., J. D. (2011) Phylogenetic relationships within the Australian limb-reduced lizard genus *Hemiergis* (Scincidae: Squamata) as inferred from the Bayesian analysis of mitochondrial rRNA gene sequences. *Copeia*, **2011**, 113-120.
- Richmond, J.Q. (2006) Evolutionary basis of parallelism in North American scincid lizards. *Evolution & Development*, **8**, 477-490.
- Rocha, C.F.D., Van Sluys, M., Vricibadic, D., Kiefer, M. C., Menezes, V. A. & Siqueira, C.C. (2009) Comportamento de termorregulacao em lagartos brasileiros. *Oecologia*

- Brasiliensis*, **13**, 115-131.
- Rocha, S., Carretero, M. A. & Harris, D. J. (2010) Genetic diversity and phylogenetic relationships of *Mabuya* spp. (Squamata: Scincidae) from western Indian Ocean islands. *Amphibia-Reptilia*, **31**, 375-385.
- Rosler, H., Bauer, A. M., Heinicke, M. P., Greenbaum, E., Jackman, T., Nguyen, T. Q. & Ziegler, T. (2011) Phylogeny, taxonomy, and zoogeography of the genus *Gekko Laurenti*, 1768 with the revalidation of *G. reevesii* Gray, 1831 (Sauria: Gekkonidae). *Zootaxa*, **2989**, 1-50.
- Sadlier, R.A., Smith, S. A., Bauer, A. M. & Whitaker, A. H. (2009) Three new species of skink in the genus *Marmorosphax* Sadlier (Squamata: Scincidae) from New Caledonia. *Memoires du Museum national d'Histoire naturelle*, **198**, 373-390.
- Sadlier, R.A., Whitaker, T., Wood, P. L. & Bauer, A. M. (2012) A new species of scincid lizard in the genus *Caledoniscincus* (Reptilia: Scincidae) from northwest New Caledonia. *Zootaxa*, **3229**, 47-57.
- Schargel, W.E., Rivas, G., Makowsky, R., Senaris, J. C., Natera, M. A., Barros, T. R., Molina, C. R. & Barrio-Amoros, C. L. (2010) Phylogenetic systematics of the genus *Gonatodes* (Squamata: Sphaerodactylidae) in the Guayana region, with description of a new species from Venezuela. *Systematics and Biodiversity*, **8**, 321-339.
- Schmitz, A.I., I. & Chirio, L. (2005) Molecular review of the genus *Panaspis sensu lato* (Reptilia: Scincidae) in Cameroon, with special reference to the status of the proposed subgenera. *Zootaxa*, **1863**, 1-28.
- Schulte, J.A., Melville, J. & Larson, A. (2003) Molecular phylogenetic evidence for ancient divergence of lizard taxa on either side of Wallace's Line. *Proceedings of the Royal Society of London B: Biological Sciences*, **270**, 597-603.
- Schulte, J.A., Losos, J. B., Cruz, F. B. & Nunez, H. (2004) The relationship between morphology, escape behaviour and microhabitat occupation in the lizard clade *Liolaemus* (Iguanidae: Tropidurinae: Liolaemini). *Journal of Evolutionary Biology*, **17**, 408-420.
- Schulte, J.A.M.-R., F. (2010) Live birth among iguanian lizards predates Pliocene-Pleistocene glaciations. *Biology Letters*, **6**, 216-218.
- Siler, C.D., Crombie, R. I., Diesmos, A. C. & Brown, R. M. (2011) Redescriptions of two poorly known slender skinks, *Brachymeles bicolor* and *Brachymeles pathfinderi* (Reptilia: Squamata: Scincidae), from the Philippines. *Journal of Herpetology*, **45**, 355-369.
- Siler, C.D., Alviola, P., Baniqued, R. D. & Duya, M. R. (2012) *Brachymeles boulengeri*: diet. *Herpetological Review*, **43**, 130.
- Sinclair, E.A., Bezy, R. L., Bolles, K., Camarillo, J. L., Crandall, K. A. & Sites, J. W. (2004) Testing species boundaries in an ancient species complex with deep phylogeographic history: genus *Xantusia* (Squamata: Xantusiidae). *American Naturalist*, **164**, 396-414.
- Sindaco, R., Metallinou, M., Pupin, F., Fasola, M. & Carranza, S. (2012) Forgotten in the ocean: systematics, biogeography and evolution of the *Trachylepis* skinks of the Socotra Archipelago. *Zoologica Scripta*, **41**, 346-362.
- Skinner, A. (2007) Phylogenetic relationships and rate of early diversification of Australian *Sphenomorphus* group scincids (Scincoidea, Squamata). *Biological Journal of the Linnean Society*, **92**, 347-366.
- Skinner, A.L., M. S. Y. (2010) Plausibility of inferred ancestral phenotypes and the evaluation of alternative models of limb evolution in scincid lizards. *Biology Letters*, **6**, 354-358.
- Skinner, A., Hugall, A. F. & Hutchinson, M. N. (2011) Lygosomine phylogeny and the origins

- of Australian scincid lizards. *Journal of Biogeography*, **38**, 1044-1058.
- Smith, S.A., Sadler, R. A., Bauer, A. M., Austin, C. C. & Jackman, T. (2007) Molecular phylogeny of the scincid lizards of New Caledonia and adjacent areas: Evidence for a single origin of the endemic skinks of Tasmantis. *Molecular Phylogenetics and Evolution*, **43**, 1151-1166.
- Solovyeva, A.N., Poyarkov, N. A., Dunaev, E. A., Duysebayeva, T. N. & Bannikova, A. A. (2011) Molecular differentiation and taxonomy of the sunwatcher toad headed agama species complex *Phrynocephalus* superspecies *helioscopus* (Pallas 1771) (Reptilia: Agamidae). *Russian Journal of Genetics*, **47**, 842-856.
- Srikulnath, K., Matsubara, K., Uno, Y., Thongpan, A., Suputtitada, S., Nishida, C., Matsuda, Y. & Apisitwanich, S. (2010) Genetic relationship of three butterfly lizard species (*Leiolepis reevesii rubritaeniata*, *Leiolepis belliana belliana*, *Leiolepis boehmei*, Agamidae, Squamata) inferred from nuclear gene sequence analyses. *Kasetsart Journal: Natural Science*, **44**, 424-435.
- Stanley, E.L., Baue, A. M., Jackman, T. R., Branch, W. R. & Mouton, P. Le F. N. (2011) Between a rock and a hard polytomy: Rapid radiation in the *rupicolous* girdled lizards (Squamata: Cordylidae). *Molecular Phylogenetics and Evolution*, **58**, 53-70.
- Starostova, Z., Kratochvil, L. S. & Flajshans, M. (2008) Cell size does not always correspond to genome size: phylogenetic analysis in geckos questions optimal DNA theories of genome size evolution. *Zoology*, **111**, 377-384.
- Stewart, J.R.T., M. B. (2003) Evolutionary transformations of the fetal membranes of viviparous reptiles: a case study of two lineages. *Journal of Experimental Zoology*, **299A**, 13-32.
- Stuart-Fox, D.M.O., I. P. F. (2003) Species-richness in agamid lizards: chance, body size, ecology or sexual selection? *Journal of Evolutionary Biology*, **16**, 659-669.
- Stuart-Fox, D., Moussalli, A. & Whiting, M. J. (2007) Natural selection on social signals: signal efficacy and the evolution of chameleon display coloration. *American Naturalist*, **170**, 916-930.
- Thompson, G.G., Clemente, C. J., Withers, P. C., Fry, B. G. & Norman, J. A. (2008) Is body shape of varanid lizards linked with retreat choice? *Australian Journal of Zoology*, **56**, 351-362.
- Tilbury, C.R.T., K. A. (2009) A new species of dwarf chameleon (Sauria; Chamaeleonidae, *Bradypodion Fitzinger*) from KwaZulu Natal South Africa with notes on recent climatic shifts and their influence on speciation in the genus. *Zootaxa*, **2226**, 43-57.
- Toda, M., Okada, S., Ota, H. & Hikida, T. (2001) Biochemical assessment of evolution and taxonomy of the morphologically poorly diverged geckos, *Gekko yakuensis* and *G. hokouensis* (Reptilia: Squamata) in Japan, with special reference to their occasional hybridization. *Biological Journal of the Linnean Society*, **73**, 153-165.
- Toda, M., Sengoku, S., Hikida, T. & Ota, H. (2008) Description of two new species of the genus *Gekko* (Squamata: Gekkonidae) from the Tokara and Amami Island groups in the Ryukyu Archipelago, Japan. *Copeia*, **2008**, 452-466.
- Torres-Carvajal, O. (2007) A taxonomic revision of south American *Stenocercus* (Squamata: Iguania) lizards. *Herpetological Monographs*, **21**, 76-178.
- Torres-Carvajal, O.d.Q., K. (2009) Phylogeny of hoplocercine lizards (Squamata: Iguania) with estimates of relative divergence times. *Molecular Phylogenetics and Evolution*, **50**, 31-43.
- Townsend, T.L., A. (2002) Molecular phylogenetics and mitochondrial genomic evolution in

- the Chamaeleonidae (Reptilia, Squamata). *Molecular Phylogenetics and Evolution*, **23**, 22-36.
- Townsend, C.R., McCranie, J. R. & Wilson, L. D. (2004) *Corytophanes cristatus*. *Catalogue of American Amphibians and Reptiles*, **789**, 1-6.
- Upton, D.E.M., R. W. (1997) Phylogeny of the side-blotched lizards (Phrynosomatidae: *Uta*) based on mtDNA sequences: support for midpeninsular seaway in Baja California. *Molecular Phylogenetics and Evolution*, **8**, 104-113.
- Vasconcelos, R., Brito, J., Carvalho, S. B., Carranza, S. & Harris, D. J. (2012) Identifying priority areas for island endemics using genetic versus specific diversity - the case of terrestrial reptiles of the Cape Verde Islands. *Biological Conservation*, **153**, 276-286.
- Vidal, M.A., Ortiz, J. C., Marin, J. C., Poulin, E. & Moreno, P. I. (2012) Comparative phylogeography of two co-distributed species of lizards of the genus *Liolaemus* (Squamata: Tropicuridae) from southern Chile. *Amphibia-Reptilia*, **33**, 55-67.
- Volobouev, V.I., I. (1994) A chromosome banding study of *Ailuronyx seychellensis* (Reptilia, Gekkonidae) and its phylogenetic affinities. *Journal of Herpetology*, **28**, 267-270.
- Wagner, P., Freund, W., Modry, D., Schmitz, A. & Bohme, W. (2011) Studies on African Agama IX. New insights into *Agama finchi* Bohme et al., 2005 (Sauria: Agamidae), with the description of a new subspecies. *Bonn zoological Bulletin*, **60**, 25-34.
- Warne, R.W.C., E. L. (2008) Reproductive allometry and the size-number trade-off for lizards. *American Naturalist*, **172**, E80-E98.
- Werneck, F.D.P., Giugliano, L. G., Collevatti, R. G. & Colli, G. (2009) Phylogeny, biogeography and evolution of clutch size in South American lizards of the genus *Kentropyx* (Squamata: Teiidae). *Molecular Ecology*, **18**, 262-278.
- Whiting, A.S., Sites, J. W., Pellegrino, K. C. M. & Rodrigues, M. T. (2006) Comparing alignment methods for inferring the history of the new world lizard genus *Mabuya* (Squamata: Scincidae). *Molecular Phylogenetics and Evolution*, **38**, 719-730.
- Wiens, J.J.R., T. W. (1997) Phylogeny of the spiny lizards (*Sceloporus*) based on molecular and morphological evidence. *Herpetological Monographs*, **11**, 1-101.
- Wiens, J.J.H., B. D. (2000) War of the iguanas: conflicting molecular and morphological phylogenies and long-branch attraction in iguanid lizards. *Systematic Biology*, **49**, 143-159.
- Wiens, J.J., Kuczynski, C. A., Arif, S. & Reeder, T. W. (2010) Phylogenetic relationships of phrynosomatid lizards based on nuclear and mitochondrial data, and a revised phylogeny for *Sceloporus*. *Molecular Phylogenetics and Evolution*, **54**, 150-161.
- Wilms, T.M., Bohme, W., Wagner, P., Lutzmann, N. & Schmitz, A. (2009) On the phylogeny and taxonomy of the genus *Uromastyx* Merrem, 1820 (Reptilia: Squamata: Agamidae: Uromastycinae) resurrection of the genus *Saara* Gray, 1845. *Bonner zoologische Beitrage*, **56**, 55-99.
- Wood, P.L., Heinicke, M. P., Jackman, T. R. & Bauer, A. M. (2012) Phylogeny of bent-toed geckos (*Cyrtodactylus*) reveals a west to east pattern of diversification. *Molecular Phylogenetics and Evolution*, **65**, 992-1003.
- Ziegler, T., Gaulke, M. & Bohme, W. (2005) Genital morphology and systematics of *Varanus mabitang* Gaulke & Curio, 2001 (Squamata: Varanidae). *Current herpetology*, **35**, 209-226.
- Ziegler, T., Bohme, W. & Schmitz, A. (2007) A new species of the *Varanus indicus* group (Squamata, Varanidae) from Halmahera Island, Moluccas: morphological and molecular evidence. *Mitteilungen aus dem Museum fur Naturkunde in Berlin - Zoologische Reihe*, **83**,

109-119.

- Zug, G.R., Brown, H. H. K., Schulte, J. A. & Vindum, J. V. (2006) Systematics of the Garden Lizards, *Calotes versicolor* Group (Reptilia, Squamata, Agamidae), in Myanmar: Central Dry Zone Populations. *Proceedings of the California Academy of Sciences*, **57**, 35-68.
- Zug, G.R., Ineich, I., Pregill, G. & Hamilton, A. M. (2012) Lizards of Tonga with description of a new Tongan treeskink (Squamata: Scincidae: *Emoia samoensis* group). *Pacific Science*, **66**, 225-237.

APPENDIX S2 Distribution Mapping References

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Ablepharus kitaibelii</i>	Disi, A. M. Modrý, D., Necas, P. and Rifai, L. 2001. Amphibians and reptiles of the Hashemite kingdom of Jordan: an atlas and field guide. Edition Chimaira, Frankfurt am Main.			
<i>Ablepharus rueppellii</i>	IUCN			
<i>Acanthocercus atricollis</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Acanthodactylus beershebensis</i>	Moravec, J., El, Din, S. B., Seligmann, H., Sivan, N. and Werner, Y. L. 1999. Systematics and distribution of the <i>Acanthodactylus pardalis</i> group (Lacertidae) in Egypt and Israel. <i>Zoology in the Middle East</i> , 17: 21-50.			
<i>Acanthodactylus boskianus</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Acanthodactylus erythrurus</i>	Bons, J. and Geniez, P. 1995. Contribution to the systematics of the lizard <i>Acanthodactylus erythrurus</i> (Sauria: Lacertidae) in Morocco. <i>Herpetological Journal</i> 5: 271-280.			
<i>Acanthodactylus pardalis</i>	Le Berre, M. 1989. Faune du Sahara 1. Poissons, amphibiens, reptiles.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Lechevalier, R. Chabaud, Paris.			
<i>Acanthodactylus schmidti</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Acanthodactylus schreiberi</i>	Salvador, A. 1982. A revision of the lizards of the genus <i>Acanthodactylus</i> (Sauria: Lacertidae). Bonner Zoologische Monographien 16: 1-167.			
<i>Acanthodactylus scutellatus</i>	Crochet, P. A., Geniez, P. and Ineich, I. 2003. A multivariate analysis of the fringe-toed lizards of the <i>Acanthodactylus scutellatus</i> group (Squamata: Lacertidae): systematic and biogeographical implications. Zoological Journal of the Linnean Society 137: 117–155.			
<i>Acanthodactylus spinicauda</i>	Salvador, A. 1982. A revision of the lizards of the genus <i>Acanthodactylus</i> (Sauria: Lacertidae). Bonner Zoologische Monographien 16: 1-167.			
<i>Acontias gariepensis</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Acontias kgalagadi</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Aeluroscalabotes felinus</i>	Grismer maps			
<i>Afroablepharus annobonensis</i>	Olivier Powells Pers. Comm.			
<i>Agama agama</i>	Bohme, W., Wagner, P., Malonza, P., Lotters, S and Kohler, J. 2005. A new species of the Agama agama group (Squamata: Agamidae) from Western Kenya, East Africa, with comments on Agama lionotus Boulenger 1896. Russian Journal of Herpetology 12: 143-150.			
<i>Agama impalearis</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biog'ographique. Asociacion Herpetologica Espanola, Barcelone.			
<i>Ailuroonyx seychellensis</i>	Schonecker, P. 2008. Geckos of Madagascar, the Seychelles, Comoros and Mascarene Islands. Aqualog Verlag Gmbh			
<i>Ailuroonyx tachyscopaeus</i>	Schonecker, P. 2008. Geckos of Madagascar, the Seychelles, Comoros and Mascarene Islands. Aqualog Verlag Gmbh			
<i>Ailuroonyx trachygaster</i>	IUCN			
<i>Algyroides fitzingeri</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Verlag, Radebeul, Germany.			
<i>Algyroides marchi</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Algyroides moreoticus</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Algyroides nigropunctatus</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Alsophylax laevis</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Alsophylax loricatus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Alsophylax pipiens</i>	Ananjeva, N. B., Munkhbayar, K., Orlov, N. L., Orlova, V. F., Semenov, D. V. and Terbish, K. 1997. Amphibians			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	and reptiles of Mongolia. Russian Academy of Sciences, Moscow.			
<i>Alsophylax szczerbaki</i>	Szczerbak, N. 2003. Guide to the reptiles of the Eastern Palearctic. Krieger Publishing Company, Malabar.			
<i>Amalosia lesueurii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Amblyrhynchus cristatus</i>	Swash, A. and Still, R. 2005. Birds, mammals, and reptiles of the Galapagos Islands. 2nd edition. Christopher Helm Publishers Ltd.			
<i>Ameiva ameiva</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Ameiva erythrocephala</i>	Kerr, A. M., Powell, R. and Parmerlee, J. S. 2005. <i>Ameiva erythrocephala</i> (Teiidae) on Sint Eustatius, Netherlands Antilles: baseline data on a small population in a severely altered habitat. Caribbean Journal of Science, 41: 162-169.			
<i>Ameiva exsul</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Ameiva fuscata</i>	Malhotra, A. and Thorpe, R. S. 1995. Ameiva fuscata. Catalogue of American Amphibians and Reptiles 606: 1-3.			
<i>Ameiva plei</i>	Censky, E. J. 1998. Ameiva plei. Catalogue of American Amphibians and Reptiles 671: 1-3.			
<i>Ameiva polops</i>	Dodd, C. K. 1980. Ameiva polops. Catalogue of American Amphibians and Reptiles 240: 1-2.			
<i>Ameiva wetmorei</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Ameivula ocellifera</i>	Cei, J. M. 1993. Reptiles del noroeste, nordeste y este de la Argentina. Museo Regionale di Scienze Naturali, Torino, Monografie 14: 1-949.			
<i>Amphibolurus muricatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Anatololacerta anatolica</i>	Eiselt, J. and Schmidtler, J. F. 1986. Der Lacerta danfordi-komplex (Reptilia: Lacertidae). Spixiana 9: 289-328.			
<i>Anatololacerta oertzeni</i>	Eiselt, J. and Schmidtler, J. F. 1986. Der Lacerta danfordi-komplex (Reptilia: Lacertidae). Spixiana 9: 289-328.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Anguis cephalonica</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Anguis fragilis</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Anniella pulchra</i>	Hunt, L. E. 2007. <i>Anniella pulchra</i> . Catalogue of American Amphibians and Reptiles, 850: 1-14.			
<i>Anolis acutus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis aeneus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis agassizi</i>	Reptile database			
<i>Anolis aquaticus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Anolis bimaculatus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis bonairensis</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis carolinensis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Anolis conspersus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis cooki</i>	Jennsen, T. A. 1990. <i>Anolis cooki</i> . Catalogue of American Amphibians and Reptiles 488: 1-2.			
<i>Anolis cristatellus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Anolis cupreus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). -			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Senckenbergiana Biologica 82: 235-241.			
<i>Anolis cuvieri</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis evermanni</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis garmani</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Anolis gingivinus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis grahami</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis gundlachi</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Anolis intermedius</i>	Savage, J. M. 2002. The amphibians and reptiles of Costa Rica. The University of Chicago Press, Chicago.			
<i>Anolis krugi</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis leachii</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis limifrons</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Anolis lineatopus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis lionotus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Anolis lividus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Anolis longiceps</i>	Powell, R. 1999. <i>Anolis longiceps</i> . Catalogue of American Amphibians and Reptiles 693: 1-2.			
<i>Anolis luciae</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis monensis</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis nebulosus</i>	IUCN			
<i>Anolis oculatus</i>	Malhotra, A. and Thorpe, R. S. 1992. <i>Anolis oculatus</i> . Catalogue of American Amphibians and Reptiles 540: 1-4.			
<i>Anolis opalinus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis oxylophus</i>	Kohler, G. 2008. Reptiles of Central America. 2nd edition. Herpeton Verlag, Offenbach.			
<i>Anolis pogus</i>	IUCN			
<i>Anolis polylepsis</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). -			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Senckenbergiana Biologica 82: 235-241.			
<i>Anolis pulchellus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis reconditus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis richardii</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis roquet</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Anolis sabanus</i>	Powell, M. A., Powell, R. and Henderson, R. W. 2005. <i>Anolis sabanus</i> . Catalogue of American Amphibians and Reptiles, 815: 1-5.			
<i>Anolis sagrei</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Anolis trinitatis</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	West Indies. University of Florida Press, Gainesville.			
<i>Anolis tropidolepis</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Anolis valencienni</i>	Schwartz, A. and Henderson, R. W. 1991. <i>Amphibians and Reptiles of the West Indies</i> . University of Florida Press, Gainesville.			
<i>Anolis wattsi</i>	Hensley, R. L., Savit, A. Z. and Powell, R. 2005. <i>Anolis schwartzi</i> . <i>Catalogue of American Amphibians and Reptiles</i> , 816: 1-5.			
<i>Aprasia pulchella</i>	Cogger, H. G. 2000. <i>Reptiles and amphibians of Australia</i> . 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Aprasia repens</i>	Cogger, H. G. 2000. <i>Reptiles and amphibians of Australia</i> . 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Archaeolacerta bedriagae</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. <i>Lurche und kriechtiere Europas</i> . Neumann Verlag, Radebeul, Germany.			
<i>Archaius tigris</i>	Gerlach, J. 2008. Population and conservation status of the reptiles of the			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Seychelles islands. Phelsuma 16: 30-48.			
<i>Aristelliger barbouri</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Aristelliger georgeensis</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Aristelliger praesignis</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Aspidoscelis burti</i>	Stebbins, R. C. 2003. A field guide to western reptiles and amphibians. Third Edition. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis ceralbensis</i>	Grismer, L. L. 1999. Phylogeny, taxonomy, and biogeography of <i>Cnemidophorus hyperythrus</i> and <i>C. ceralbensis</i> (Squamata: Teiidae) in Baja California, México. <i>Herpetologica</i> 55: 28-42.			
<i>Aspidoscelis cozumelae</i>	Fritts, T. H. 1969. The systematics of the parthenogenetic lizards of the <i>Cnemidophorus cozumela</i> complex. <i>Copeia</i> , 1969: 519-535.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Aspidoscelis deppei</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Aspidoscelis dixoni</i>	Stebbins, R. C. 2003. A field guide to western reptiles and amphibians. Third Edition. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis exsanguis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis gularis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis guttata</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Aspidoscelis hyperythra</i>	Thompson, J. S., Crother, B. I. and Price, A. H. 1998. <i>Cnemidophorus hyperythrus</i> . <i>Catalogue of American Amphibians and Reptiles</i> 655: 1-6.			
<i>Aspidoscelis inornata</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis laredoensis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis lineattissima</i>	IUCN			
<i>Aspidoscelis marmorata</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis neomexicana</i>	Wright, J. W. 1971. <i>Cnemidophorus neomexicanus</i> . Catalogue of American Amphibians and Reptiles 109: 1-3.			
<i>Aspidoscelis scalaris</i>	Jones, L. and Lovich, R. 2009. Lizards of the American southwest: a photographic field guide. Rio Nuevo Publishers, Tuscon.			
<i>Aspidoscelis sexlineata</i>	Trauth, S. E. and McAllister, C. T. 1996. <i>Cnemidophorus sexlineatus</i> . Catalogue of American Amphibians and Reptiles 628: 1-12.			
<i>Aspidoscelis sonorae</i>	Stebbins, R. C. 2003. A field guide to western reptiles and amphibians. Third			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Edition. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis tessellata</i>	Price, A. H. 1986. <i>Cnemidophorus tessellatus</i> . Catalogue of American Amphibians and Reptiles 398: 1-2.			
<i>Aspidoscelis tigris</i>	Smith, H. M. 1946. Handbook of lizards. Lizards of the United States and Canada. Cornell University Press, Ithaca.			
<i>Aspidoscelis uniparens</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Aspidoscelis velox</i>	Stebbins, R. C. 2003. A field guide to western reptiles and amphibians. Third Edition. Houghton Mifflin Company, Boston.			
<i>Aspronema cochabambae</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Asymblepharus sikimmensis</i>	Ouboter, P. E. 1986. A revision of the genus <i>Scincella</i> (Reptilia: Sauria: Scincidae) of Asia, with some notes on its evolution. Zoologische Verhandelingen 229: 1-66.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Atlantolacerta andreanskyi</i>	Schleich, H. H., Kastle, W. and Kabisch, K. 1996. Amphibians and reptiles of North Africa. Biology, systematics, field guide. Koeltz Scientific, Koenigstein, Germany.			
<i>Barisia herrerae</i>	Bryson, R. W. and Riddle, B. R. 2012. Tracing the origins of widespread highland species: a case of Neogene diversification across the Mexican sierras in an endemic lizard. Biological Journal of the Linnean Society, 105: 382-394.			
<i>Barisia imbricata</i>	Guillette, J. L. and Casas-Andreu, G. 1987. The reproductive biology of the high elevation Mexican lizard <i>Barisia imbricata</i> . Herpetologica 43: 29-38.			
<i>Basiliscus basiliscus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Basiliscus vittatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Bassiana duperreyi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Bassiana trilineata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Bavayia crassicollis</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia cyclura</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia exsuccida</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia geitaina</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia montana</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia ornata</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Bavayia pulchella</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia sauvagii</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bavayia septuiclavus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Bipes bipes</i>	Papenfuss, T. J. 1982. The ecology and systematics of the amphisbaenian genus <i>Bipes</i> . Occasional Papers of the California Academy of Sciences 136: 1-42.			
<i>Bipes canaliculatus</i>	Hodges, W. L. and Perez-Ramos, E. 2001. New localities and natural history notes on <i>Bipes canaliculatus</i> in Guerrero, Mexico. Herpetological Review 32: 153-156.			
<i>Bipes tridactylus</i>	Papenfuss, T. J. 1982. The ecology and systematics of the amphisbaenian genus <i>Bipes</i> . Occasional Papers of the California Academy of Sciences 136: 1-42.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Brachylophus fasciatus</i>	Keogh, J. S., Edwards, D. L., Fisher, R. N. and Harlow, P. S. 2008. Molecular and morphological analysis of the critically endangered Fijian iguanas reveals cryptic diversity and a complex biogeographic history. Philosophical Transactions of the Royal Society of London B. 363: 3413-3426.			
<i>Brachylophus vitiensis</i>	Keogh, J. S., Edwards, D. L., Fisher, R. N. and Harlow, P. S. 2008. Molecular and morphological analysis of the critically endangered Fijian iguanas reveals cryptic diversity and a complex biogeographic history. Philosophical Transactions of the Royal Society of London B. 363: 3413-3426.			
<i>Brachymeles cebuensis</i>	Brown, W. C. and Rabor, D. S. 1967. Review of the genus <i>Brachymeles</i> (Scincidae), with descriptions of new species and subspecies. Proceedings of the California Academy of Sciences 34: 525-548.			
<i>Brachymeles minimus</i>	IUCN			
<i>Brachymeles taylori</i>	Siler, C. D. and Brown, R. M. 2010. Phylogeny-based species delimitation in Philippine loam-swimming skinks (Reptilia: Squamata: Scincidae: <i>Brachymeles</i>): taxonomic revision of the pentadactyl species groups and			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	description of three new species. Herpetological Monographs 24: 1-54.			
<i>Brachymeles tridactylus</i>	Brown, W. C. and Rabor, D. S. 1967. Review of the genus <i>Brachymeles</i> (Scincidae), with descriptions of new species and subspecies. Proceedings of the California Academy of Sciences 34: 525-548.			
<i>Bradypodion pumilum</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Bradypodion ventrale</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Brasiliscincus agilis</i>	Bischoff, W. 1984. <i>Lacerta agilis</i> Linnaeus 1758 - Zauneidechse. Pages 23-68 in W. Bohme, editor. Handbuch der reptilien und amphibien Europas, Band 2/I: Echsen II (<i>Lacerta</i>). Aula- Verlag, Wiesbaden.	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. <i>Lurche und kriechtiere</i> Europas. Neumann Verlag, Radebeul, Germany.	Gasc, J. P., Cabela A., Crnobrnja-Isailovic, J., Dolmen, D., Grossenbacher, K., Haffner, P., Lescure, J., Martens, H., Martínez Rica, J. P., Maurin, H., Oliveira, M. E., Sofianidou, T. S., Veith, M. and Zuiderwijk, A. (eds), 1997. Atlas of amphibians and reptiles in Europe. Collection Patrimoines Naturels, 29, Societas Europaea	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson- Gale, Detroit.

Species	Map source 1	Map source 2	Map source 3	Map source 4
			Herpetologica, Muséum National d'Histoire Naturelle & Service du Patrimoine Naturel, Paris, 496 pp.	
<i>Brasiliscincus heathi</i>	Nogueira, C. D. 2006. Diversidade e padrões de distribuição da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências, Universidade de São Paulo.	Vrcibradic, D., Almeida-Gomes, M., Borges-Junior, V. N. T., Kiefer, M. C., Van Sluys, M. and Reptilia, Rocha, C. F. D. 2006. Scincidae, Mabuya frenata: distribution extension. Check List 2: 57-58.		
<i>Caledoniscincus aquilonius</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus atropunctatus</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus auratus</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus austrocaledonicus</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Reptiles, St. Louis.			
<i>Caledoniscincus chazeaui</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus festivus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus haplorhinus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus orestes</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus renevieri</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Caledoniscincus terma</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Callisaurus draconoides</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Calotes mystaceus</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Calotes nemoricola</i>	Tikader, B. K. and Sharma, R. C. 1992. Handbook of Indian lizards. Zoological Survey of India, Calcutta.			
<i>Calotes versicolor</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Carlia longipes</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Carlia rhomboidalis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Carlia rostralis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Carlia rubrigularis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Carlia tetradactyla</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Island, FL : Ralph Curtis Publishing.			
<i>Carphodactylus laevis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Celatiscincus euryotis</i>	Sadlier, R. A., Bauer, A. M. and Smith, S. A. 2006. A new species of <i>Nannoscincus</i> Günther (Squamata: Scincidae) from high elevation forest in southern New Caledonia. Records of the Australian Museum 58: 29-36.			
<i>Celatiscincus similis</i>	Sadlier, R. A., Bauer, A. M. and Smith, S. A. 2006. A new species of <i>Nannoscincus</i> Günther (Squamata: Scincidae) from high elevation forest in southern New Caledonia. Records of the Australian Museum 58: 29-36.			
<i>Cercosaura schreibersii</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Chalcides bedriagai</i>	Carranza, S., Arnold, E. N., Geniez, P., Roca, J. and Mateo, J. A. 2008. Radiation, multiple dispersal and parallelism in the skinks, <i>Chalcides</i> and <i>Sphenops</i> (Squamata: Scincidae), with comments on <i>Scincus</i> and <i>Scincopus</i> and the age of the Sahara Desert. Molecular Phylogenetics and Evolution 46: 1071-1094.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Chalcides chalcides</i>	Carranza, S., Arnold, E. N., Geniez, P., Roca, J. and Mateo, J. A. 2008. Radiation, multiple dispersal and parallelism in the skinks, Chalcides and Sphenops (Squamata: Scincidae), with comments on Scincus and Scincopus and the age of the Sahara Desert. <i>Molecular Phylogenetics and Evolution</i> 46: 1071-1094.			
<i>Chalcides coeruleopunctatus</i>	Carranza, S., Arnold, E. N., Geniez, P., Roca, J. and Mateo, J. A. 2008. Radiation, multiple dispersal and parallelism in the skinks, Chalcides and Sphenops (Squamata: Scincidae), with comments on Scincus and Scincopus and the age of the Sahara Desert. <i>Molecular Phylogenetics and Evolution</i> 46: 1071-1094.			
<i>Chalcides ocellatus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Chalcides sepsoides</i>	Disi, A. M. Modrý, D., Necas, P. and Rifai, L. 2001. Amphibians and reptiles of the Hashemite kingdom of Jordan: an atlas and field guide. Edition Chimaira, Frankfurt am Main.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Chalcides sexlineatus</i>	Carranza, S., Arnold, E. N., Geniez, P., Roca, J. and Mateo, J. A. 2008. Radiation, multiple dispersal and parallelism in the skinks, Chalcides and Sphenops (Squamata: Scincidae), with comments on Scincus and Scincopus and the age of the Sahara Desert. Molecular Phylogenetics and Evolution 46: 1071-1094.			
<i>Chalcides viridanus</i>	Carranza, S., Arnold, E. N., Geniez, P., Roca, J. and Mateo, J. A. 2008. Radiation, multiple dispersal and parallelism in the skinks, Chalcides and Sphenops (Squamata: Scincidae), with comments on Scincus and Scincopus and the age of the Sahara Desert. Molecular Phylogenetics and Evolution 46: 1071-1094.			
<i>Chamaeleo calyptratus</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Chamaeleo chamaeleon</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biog*ographique. Asociacion Herpetologica Espanola, Barcelone.			
<i>Chamaeleo dilepis</i>	Chirio, L. and LeBreton, M. 2007. Atlas			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	des reptiles du Cameroun. Publications Scientifiques du Museum national d'Histoire naturelle, Paris.			
<i>Chamaeleo monachus</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. Zootaxa 2826: 1-44 (Reptilia)			
<i>Chamaeleo namaquensis</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad, Project Report.			
<i>Chioninia coctei</i>	Miralles, A., Kohler, J., Glaw, F. and Vences, M. 2011. A molecular phylogeny of the “ <i>Madascincus polleni</i> species complex”, with description of a new species of scincid lizard from the coastal dune area of northern Madagascar. Zootaxa 2876: 1-16.			
<i>Chioninia spinalis</i>	Miralles, A., Vasconcelos, R., Perera, A., Harris, D. J. and Carranza, S. 2011. An integrative taxonomic revision of the Cape Verdean skinks (Squamata,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Scincidae). Zoologica Scripta 40: 16-44.			
<i>Chioninia vaillantii</i>	Miralles, A., Vasconcelos, R., Perera, A., Harris, D. J. and Carranza, S. 2011. An integrative taxonomic revision of the Cape Verdean skinks (Squamata, Scincidae). Zoologica Scripta 40: 16-44.			
<i>Chlamydosaurus kingii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Chondrodactylus angulifer</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad, Project Report.			
<i>Christinus guentheri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Christinus marmoratus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Cnemaspis affinis</i>	Grismer, L. L., Onn, C. K., Nasir, N. and Sumontha, M. 2008. A new species of karst dwelling gecko (genus <i>Cnemaspis</i> Strauch 1887) from the border region of Thailand and Peninsular Malaysia. Zootaxa, 1875: 51-68.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Cnemaspis limi</i>	Grismer, L. L., Ahmad, N. and Onn, C. K. 2009. A new, diminutive, upland <i>Sphenomorphus</i> Fitzinger 1843 (Squamata: Scincidae) from the Belum-Temengor forest complex, Peninsular Malaysia. <i>Zootaxa</i> , 2312: 27-38.			
<i>Cnemaspis modiglianii</i>	Reptile database			
<i>Cnemaspis monachorum</i>	Grismer maps			
<i>Cnemaspis perhentianensis</i>	Grismer, L. L. and Onn, C. K. 2008. A new species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from Pulau Perhentian Besar, Terengganu, Peninsular Malaysia. <i>Zootaxa</i> 1771: 1-15.			
<i>Cnemaspis whittenorum</i>	Reptile database			
<i>Cnemidophorus arubensis</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Cnemidophorus cryptus</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Cnemidophorus gramivagus</i>	IUCN			
<i>Cnemidophorus lemniscatus</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Cnemidophorus murinus</i>	Ugueto, G. N. and Harvey, M. B. 2010. Southern Caribbean Cnemidophorus (Squamata: Teiidae): description of new species and taxonomic status of <i>C. murinus ruthveni</i> Burt. Herpetological Monographs 24: 111-148.			
<i>Cnemidophorus vanzoi</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Coeranoscincus reticulatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Coleonyx brevis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Coleonyx elegans</i>	Dial, B. E. and Grismer, L. L. 1992. A phylogenetic analysis of physiological-ecological character evolution in the lizard genus <i>Coleonyx</i> and its implications for historical biogeographic reconstruction. Systematic Biology 41: 178-195.			
<i>Coleonyx reticulatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Boston.			
<i>Coleonyx variegatus</i>	Dial, B. E. and Grismer, L. L. 1992. A phylogenetic analysis of physiological-ecological character evolution in the lizard genus <i>Coleonyx</i> and its implications for historical biogeographic reconstruction. <i>Systematic Biology</i> 41: 178-195.			
<i>Conolophus pallidus</i>	Swash, A. and Still, R. 2005. Birds, mammals, and reptiles of the Galapagos Islands. 2nd edition. Christopher Helm Publishers Ltd.			
<i>Conolophus subcristatus</i>	Swash, A. and Still, R. 2005. Birds, mammals, and reptiles of the Galapagos Islands. 2nd edition. Christopher Helm Publishers Ltd.			
<i>Contomastix lacertoides</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Contomastix vacariensis</i>	Guarino Colli, pers. Comm.			
<i>Copeoglossum nigropunctatum</i>	Avila-Pires 1995, Miralles and Carranza 2010, Miralles et al. 2005, Miralles et al. 2009, Nogueira 2006			
<i>Cophosaurus texanus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Correlophus ciliatus</i>	Bauer, A. M., Jackman, T. R., Sadlier, R. A. and Whitaker, A. H. 2012. Revision of the giant geckos of New Caledonia (Reptilia: Diplodactylidae: Rhacodactylus). Zootaxa 3404: 1-52.			
<i>Correlophus sarasinorum</i>	Bauer, A. M., Jackman, T. R., Sadlier, R. A. and Whitaker, A. H. 2012. Revision of the giant geckos of New Caledonia (Reptilia: Diplodactylidae: Rhacodactylus). Zootaxa 3404: 1-52.			
<i>Corucia zebrata</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Crenadactylus ocellatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Crossobamon evermanni</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Crotaphytus collaris</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Crotaphytus grismeri</i>	McGuire, J. A. 1994. A new species of collared lizard (Iguania: Crotaphytidae) from northeastern Baja California, Mexico. <i>Herpetologica</i> , 50: 438-450.			
<i>Crotaphytus reticulatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Cryptoblepharus egeriae</i>	Horner, P. 2007. <i>Ctenotus quirinus</i> sp. nov. (Reptilia: Sauria: Scincidae) – a new species of skink from the Northern Territory, with the recognition of <i>C. brevipes</i> Storr, 1981 and <i>C. essingtonii</i> (Gray, 1842) as distinct species. <i>The Beagle</i> 23: 119-130.			
<i>Cryptoblepharus eximius</i>	Horner, P. 2007. <i>Ctenotus quirinus</i> sp. nov. (Reptilia: Sauria: Scincidae) – a new species of skink from the Northern Territory, with the recognition of <i>C. brevipes</i> Storr, 1981 and <i>C. essingtonii</i> (Gray, 1842) as distinct species. <i>The Beagle</i> 23: 119-130.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Cryptoblepharus novocaledonicus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Cryptoblepharus poecilopleurus</i>	Horner, P. 2007. <i>Ctenotus quirinus</i> sp. nov. (Reptilia: Sauria: Scincidae) – a new species of skink from the Northern Territory, with the recognition of <i>C. brevipes</i> Storr, 1981 and <i>C. essingtonii</i> (Gray, 1842) as distinct species. The Beagle 23: 119-130.			
<i>Cryptoblepharus renschi</i>	Horner, P. 2007. <i>Ctenotus quirinus</i> sp. nov. (Reptilia: Sauria: Scincidae) – a new species of skink from the Northern Territory, with the recognition of <i>C. brevipes</i> Storr, 1981 and <i>C. essingtonii</i> (Gray, 1842) as distinct species. The Beagle 23: 119-130.			
<i>Ctenophorus fionni</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenophorus fordi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenophorus isolepis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Ctenophorus maculosus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenophorus nuchalis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenophorus ornatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenosaura bakeri</i>	de Queiroz, K. 1990. <i>Ctenosaura bakeri</i> . Catalogue of American Amphibians and Reptiles 465: 1-2.			
<i>Ctenosaura oedirhina</i>	de Queiroz, K. 1990. <i>Ctenosaura oedirhina</i> . Catalogue of American Amphibians and Reptiles 466: 1-2.			
<i>Ctenosaura pectinata</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Ctenosaura similis</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Ctenotus brooksi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Island, FL : Ralph Curtis Publishing.			
<i>Ctenotus lanceolini</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenotus pantherinus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenotus robustus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Ctenotus taeniolatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Cyclodomorphus casuarinae</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Cyclodomorphus celatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Cyclodomorphus melanops</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Cyclodomorphus michaeli</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Cyclura carinata</i>	Powell, R. 2000. <i>Cyclura onchiopsis</i> . Catalogue of American Amphibians and Reptiles, 710: 1-3.			
<i>Cyclura collei</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Cyclura cychlura</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Cyclura pinguis</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Cyclura rileyi</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Cyrtodactylus jarakensis</i>	Grismer, L. L., Onn, C. K., Grismer, J. L., Wood, P. L. and Belabut, D. 2008. Three new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from Peninsular Malaysia. <i>Zootaxa</i> 1921: 1- 23.			
<i>Cyrtodactylus laevigatus</i>	Auffenberg, W. 1980. The herpetofauna of Komodo, with notes on adjacent areas. <i>Bulletin of the Florida State</i>			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Museum Biological Sciences, 25: 39-156.			
<i>Cyrtodactylus leegrimeri</i>	Onn, C. K. and Ahmad, N. 2010. A new insular species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from northeastern Peninsular Malaysia, Malaysia. <i>Zootaxa</i> 2389: 47-56.			
<i>Cyrtodactylus peguensis</i>	natural history museum records			
<i>Cyrtodactylus redimiculus</i>	IUCN			
<i>Cyrtodactylus sadleiri</i>	Cogger, H., Sadler, R. and Cameron, E. 1983. The terrestrial reptiles of Australia's island territories. Australian National Parks and Wildlife Service Special Publication 8 (11) 1-79.			
<i>Cyrtodactylus seribuatensis</i>	Youmans, T. M. and Grismer, L. L. 2006. A new species of <i>Cyrtodactylus</i> (Reptilia: Squamata: Gekkonidae) from the Seribuat Archipelago, West Malaysia. <i>Herpetological Natural History</i> , 10: 61-70.			
<i>Cyrtodactylus tiomanensis</i>	Grismer maps			
<i>Cyrtopodion caspium</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Cyrtopodion fedtschenkoi</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Dalmatolacerta oxycephala</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Darevskia chlorogaster</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia derjugini</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia parvula</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia portschinskii</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Darevskia praticola</i>	northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia raddei</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia rostombekovi</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia rudis</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Darevskia saxicola</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia unisexualis</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Darevskia valentini</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Dasia olivacea</i>	Harikrishnan, S., Vasudevan, K., Chandramouli, S. R., Choudhury, B. C., Dutta, S. K. and Das, I. 2012. A new species of <i>Coryphophylax</i> Fitzinger in: Steindachner, 1867 (Sauria: Iguania: Agamidae) from the Andaman Islands, India. <i>Zootaxa</i> 3451: 31-45.			
<i>Delma fraseri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Delma grayii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Dierogecko insularis</i>	Bauer, A. M., Barts, M. and Hulbert, F. 2006. A new species of the Pachydactylus weberi group (Reptilia: Squamata: Gekkonidae) from the Orange River, with comments on its natural history. Salamandra 42: 83-92.			
<i>Dierogecko nehoueensis</i>	Bauer, A. M., Barts, M. and Hulbert, F. 2006. A new species of the Pachydactylus weberi group (Reptilia: Squamata: Gekkonidae) from the Orange River, with comments on its natural history. Salamandra 42: 83-92.			
<i>Dierogecko poumensis</i>	Bauer, A. M., Barts, M. and Hulbert, F. 2006. A new species of the Pachydactylus weberi group (Reptilia: Squamata: Gekkonidae) from the Orange River, with comments on its natural history. Salamandra 42: 83-92.			
<i>Dierogecko validiclavis</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Dinarolacerta mosorensis</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Verlag, Radebeul, Germany.			
<i>Diplodactylus tessellatus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Diploglossus millepunctatus</i>	Reptile database			
<i>Diploglossus pleii</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Diporiphora nobbi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Dipsosaurus dorsalis</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Dopasia gracilis</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Draco biaro</i>	Manthey, U. 2008. Agamid Lizards of southern Asia. Volume 1. Draconinae. Edition Chimaira, Frankfurt.			
<i>Draco bourouniensis</i>	Musters, C. J. M. 1983. Taxonomy of the genus <i>Draco</i> L. (Agamidae, Lacertilia, Reptilia). Zoologische Verhandelingen,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Leiden 199: 1-120.			
<i>Draco palawanensis</i>	Grismer maps			
<i>Draco reticulatus</i>	McGuire, J. A. and Alcala, A. C. 2000. A taxonomic revision of the flying lizards of the Philippine Islands (Iguania: Agamidae: Draco), with a description of a new species. Herpetological Monographs 14: 81-138.			
<i>Draco timorensis</i>	Musters, C. J. M. 1983. Taxonomy of the genus Draco L. (Agamidae, Lacertilia, Reptilia). Zoologische Verhandelingen, Leiden 199: 1-120.			
<i>Draco volans</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Egernia cunninghami</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Egernia hosmeri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Egernia kingii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Egernia stokesii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Egernia striolata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Elgaria coerulea</i>	Good, D. A. 1988. Allozyme variation and phylogenetic relationships among the species of <i>Elgaria</i> (Squamata: Anguinae). <i>Herpetologica</i> , 44: 154-162.			
<i>Elgaria multicarinata</i>	Good, D. A. 1988. Allozyme variation and phylogenetic relationships among the species of <i>Elgaria</i> (Squamata: Anguinae). <i>Herpetologica</i> , 44: 154-162.			
<i>Emoia adspersa</i>	IUCN			
<i>Emoia aneityumensis</i>	Allison A. pers. Comm.			
<i>Emoia atrocostata</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnamera). Pensoft.			
<i>Emoia campbelli</i>	Morrison, C. 2003. A field guide to the herpetofauna of Fiji. Institute of Applied Sciences, University of the South Pacific. Suva, Fiji.			
<i>Emoia concolor</i>	Morrison, C. 2003. A field guide to the herpetofauna of Fiji. Institute of Applied Sciences, University of the South			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Pacific. Suva, Fiji.			
<i>Emoia impar</i>	Allison A. pers. Comm.			
<i>Emoia isolata</i>	Allison A. pers. Comm.			
<i>Emoia loyaltiensis</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Emoia maculata</i>	Allison A. pers. Comm.			
<i>Emoia nativittatis</i>	Cogger, H., Sadlier, R. and Cameron, E. 1983. The terrestrial reptiles of Australia's island territories. Australian National Parks and Wildlife Service Special Publication 8 (11) 1-79.			
<i>Emoia nigromarginata</i>	Allison A. pers. Comm.			
<i>Emoia parkeri</i>	Morrison, C. 2003. A field guide to the herpetofauna of Fiji. Institute of Applied Sciences, University of the South Pacific. Suva, Fiji.			
<i>Emoia ponapea</i>	Allison A. pers. Comm.			
<i>Emoia reimschiisseli</i>	Allison A. pers. Comm.			
<i>Emoia sanfordi</i>	Allison A. pers. Comm.			
<i>Emoia schmidtii</i>	Allison A. pers. Comm.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Emoia slevini</i>	IUCN			
<i>Emoia sorex</i>	Allison A. pers. Comm.			
<i>Emoia taumakoensis</i>	Allison A. pers. Comm.			
<i>Emoia tongana</i>	Allison A. pers. Comm.			
<i>Emoia trossula</i>	Allison A. pers. Comm.			
<i>Enyalius perditus</i>	Guarino Colli, pers. Comm.			
<i>Eremias arguta</i>	Ananjeva, N. B., Munkhbayar, K., Orlov, N. L., Orlova, V. F., Semenov, D. V. and Terbish, K. 1997. Amphibians and reptiles of Mongolia. Russian Academy of Sciences, Moscow.			
<i>Eremias grammica</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias intermedia</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias lineolata</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Eremias nigrocellata</i>	northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias nikolskii</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias persica</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias pleskei</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Eremias regeli</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias strauchi</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremias velox</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eremiascincus richardsonii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Eublepharis macularius</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Eublepharis turcmenicus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Sofia.			
<i>Eulamprus brachyosoma</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Eulamprus quoyii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Eulamprus tympanum</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Eumeces schneideri</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Eurydactylodes agricolae</i>	Bauer, A. M., Jackman, T., Sadler, R. A. and Whitaker, A. H. 2009. Review and phylogeny of the New Caledonian diplodactylid gekkotan genus <i>Eurydactylodes</i> Wermuth, 1965, with the description of a new species. in Grandcolas P. (ed.), Zoologia Neocaledonica 7. Biodiversity studies in New Caledonia. Memoires du Muséum national d'Histoire naturelle 198: 13-36.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Eurydactylodes symmetricus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Eurydactylodes vieillardii</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Euspondylus chasqui</i>	Chavez, G., Siu-Ting, K., Duran, V. and Venegas, P. J. 2011. Two new species of Andean gymnophthalmid lizards of the genus <i>Euspondylus</i> (Reptilia, Squamata) from central and southern Peru. <i>ZooKeys</i> 109: 1-17.			
<i>Eutropis carinata</i>	Schleich, H. H. and Kastle, W. 2002. Amphibians and reptiles of Nepal. Gantner Verlag, Koenigstein.			
<i>Eutropis longicaudata</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Eutropis multifasciata</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Furcifer cephalolepis</i>	Hawlitschek, O., Bruckmann, B., Berger, J., Green, K. and Glaw, F. 2011. Integrating field surveys and remote sensing data to study distribution, habitat use and conservation status of the			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	herpetofauna of the Comoro Islands. ZooKeys 144: 21-79.			
<i>Furcifer polleni</i>	Hawlitschek, O., Bruckmann, B., Berger, J., Green, K. and Glaw, F. 2011. Integrating field surveys and remote sensing data to study distribution, habitat use and conservation status of the herpetofauna of the Comoro Islands. ZooKeys 144: 21-79.			
<i>Gallotia atlantica</i>	Hernandez, E., Nogales, M. and Martín, A. 2000. Discovery of a new lizard in the Canary Islands, with a multivariate analysis of Gallotia (Reptilia: Lacertidae). Herpetologica, 56: 63-76.			
<i>Gallotia bravoana</i>	Arnold, E. N. and Ovenden, D. W. 2004. A field guide to the reptiles and amphibians of Britain and Europe. 2nd edition. Collins, London.			
<i>Gallotia caesaris</i>	Arnold, E. N. and Ovenden, D. W. 2004. A field guide to the reptiles and amphibians of Britain and Europe. 2nd edition. Collins, London.			
<i>Gallotia galloti</i>	Hernandez, E., Nogales, M. and Martín, A. 2000. Discovery of a new lizard in the Canary Islands, with a multivariate analysis of Gallotia (Reptilia: Lacertidae). Herpetologica, 56: 63-76.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Gallotia intermedia</i>	Hernandez, E., Nogales, M. and Martín, A. 2000. Discovery of a new lizard in the Canary Islands, with a multivariate analysis of Gallotia (Reptilia: Lacertidae). Herpetologica, 56: 63-76.			
<i>Gallotia simonyi</i>	Hernandez, E., Nogales, M. and Martín, A. 2000. Discovery of a new lizard in the Canary Islands, with a multivariate analysis of Gallotia (Reptilia: Lacertidae). Herpetologica, 56: 63-76.			
<i>Gallotia stehlini</i>	Hernandez, E., Nogales, M. and Martín, A. 2000. Discovery of a new lizard in the Canary Islands, with a multivariate analysis of Gallotia (Reptilia: Lacertidae). Herpetologica, 56: 63-76.			
<i>Gambelia sila</i>	Jennings, M. R. 1995. <i>Gambelia sila</i> . Catalogue of American Amphibians and Reptiles 612: 1-4.			
<i>Gambelia wislizenii</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Gehyra variegata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Gekko athymus</i>	IUCN			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Gekko ernstkelleri</i>	IUCN			
<i>Gekko gecko</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Gekko hokouensis</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Gekko japonicus</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Gekko shibatai</i>	Toda, M., Sengoku, S., Hikida, T. and Ota, H. 2008. Description of two new species of the genus Gekko (Squamata: Gekkonidae) from the Tokara and Amami Island groups in the Ryukyu Archipelago, Japan. Copeia 2008: 452-466.			
<i>Gekko smithii</i>	Grismer maps			
<i>Gekko vertebralis</i>	Toda, M., Sengoku, S., Hikida, T. and Ota, H. 2008. Description of two new species of the genus Gekko (Squamata: Gekkonidae) from the Tokara and Amami Island groups in the Ryukyu Archipelago, Japan. Copeia 2008: 452-466.			
<i>Gekko yakuensis</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Krieger Publications, Malabar, FL.			
<i>Gerrhonotus infernalis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Gerrhosaurus skoogi</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad, Project Report.			
<i>Gonatodes albogularis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Gonatodes daudini</i>	Powell, R. and Henderson, R. W. 2005. A new species of <i>Gonatodes</i> (Squamata: Gekkonidae) from the West Indies. Caribbean Journal of Science, 41: 709-715.			
<i>Gonatodes ocellatus</i>	Schargel, W. E. 2008. Species limits and phylogenetic systematics of the diurnal geckos of the genus <i>Gonatodes</i> (Squamata: Sphaerodatyliidae). PhD Dissertation, University of Texas at Arlington.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Gonatodes vittatus</i>	Marinus Hoogmoed, pers. Comm.			
<i>Gongylomorphus bojerii</i>	Austin, J. J., Arnold, E. N. and Jones, C. G. 2009. Interrelationships and history of the slit-eared skinks (Gongylomorphus, Scincidae) of the Mascarene islands, based on mitochondrial DNA and nuclear gene sequences. <i>Zootaxa</i> 2153: 55-68.			
<i>Goniurosaurus araneus</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Goniurosaurus hainanensis</i>	Grismer, L. L., Viets, B. E. and Boyle, L. J. 1999. Two new continental species of <i>Goniurosaurus</i> (Squamata: Eublepharidae) with a phylogeny and evolutionary classification of the genus. <i>Journal of Herpetology</i> , 33: 382-393.			
<i>Goniurosaurus kuroiwae</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Goniurosaurus splendens</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Gonocephalus bellii</i>	Grismer maps			
<i>Gonocephalus bornensis</i>	Indraneil Das, pers. Comm.			
<i>Gonocephalus grandis</i>	Grismer maps			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Gymnodactylus amarali</i>	Vanzolini, P. E. 2005. On <i>Gymnodactylus amarali</i> Barbour, 1925, with the description of a new species (Sauria, Gekkonidae). <i>Anais da Academia Brasileira de Ciencias</i> 77: 595-611.			
<i>Gymnodactylus geckoides</i>	Nogueira, C. D. 2006. Diversidade e padroes de distribuicao da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências, Universidade de São Paulo.			
<i>Gymnophthalmus speciosus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Haemodracon riebeckii</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. <i>Zootaxa</i> 2826: 1-44 (Reptilia)			
<i>Haemodracon trachyrhinus</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. Zootaxa 2826: 1-44 (Reptilia)			
<i>Heliobolus lugubris</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Hellenolacerta graeca</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Heloderma horridum</i>	Kohler, G. 2003. A new species of Morunasaurus from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Heloderma suspectum</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Hemicordylus capensis</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Hemidactylus bouvieri</i>	Loveridge, A. 1947. Revision of the African lizards of the family			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Gekkonidae. Bulletin of the Museum of Comparative Zoology 98: 1-469.			
<i>Hemidactylus bowringii</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Hemidactylus brookii</i>	Branch, W. R. 2005. A Photographic guide to snakes other reptiles and amphibians of East Africa. Struik Nature, Capetown.			
<i>Hemidactylus dracaenacolus</i>	Gomez-Diaz, E., Sindaco, R., Pupin, F., Fasola, M. and Carranza, S. 2012. Origin and in situ diversification in <i>Hemidactylus</i> geckos of the Socotra Archipelago. Molecular Ecology 16: 4074-4092.			
<i>Hemidactylus flaviviridis</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Hemidactylus frenatus</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Hemidactylus mabouia</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Hemidactylus maculatus</i>	Tikader, B. K. and Sharma, R. C. 1992. Handbook of Indian lizards. Zoological			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Survey of India, Calcutta.			
<i>Hemidactylus turcicus</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Hemiergis peronii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Hemisphaeriodon gerrardii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Hesperoedura reticulata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Heteronotia binoei</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Holbrookia lacerata</i>	Axtell, R. W. 1956. A solution to the long neglected <i>Holbrookia lacerata</i> problem, and the description of two new subspecies of <i>Holbrookia</i> . Bulletin of the Chicago Academy of Sciences 10: 163-179.			
<i>Holbrookia maculata</i>	Axtell, R. W. 1956. A solution to the long neglected <i>Holbrookia lacerata</i> problem, and the description of two new subspecies of <i>Holbrookia</i> . Bulletin of the Chicago Academy of Sciences 10: 163-			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	179.			
<i>Holbrookia propinqua</i>	Axtell, R. W. 1983. <i>Holbrookia propinqua</i> . Catalogue of American Amphibians and Reptiles 341: 1-2.			
<i>Holcosus festivus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Holcosus quadrilineatus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Homonota darwinii</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Homonota fasciata</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Iberolacerta aranica</i>	IUCN			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Iberolacerta aurelioi</i>	Gasc, J. P., Cabela A., Crnobrnja-Isailovic, J., Dolmen, D., Grossenbacher, K., Haffner, P., Lescure, J., Martens, H., Martínez Rica, J. P., Maurin, H., Oliveira, M. E., Sofianidou, T. S., Veith, M. and Zuiderwijk, A. (eds), 1997. Atlas of amphibians and reptiles in Europe. Collection Patrimoines Naturels, 29, Societas Europaea Herpetologica, Muséum National d'Histoire Naturelle & Service du Patrimoine Naturel, Paris, 496 pp.			
<i>Iberolacerta bonnali</i>	Gasc, J. P., Cabela A., Crnobrnja-Isailovic, J., Dolmen, D., Grossenbacher, K., Haffner, P., Lescure, J., Martens, H., Martínez Rica, J. P., Maurin, H., Oliveira, M. E., Sofianidou, T. S., Veith, M. and Zuiderwijk, A. (eds), 1997. Atlas of amphibians and reptiles in Europe. Collection Patrimoines Naturels, 29, Societas Europaea Herpetologica, Muséum National d'Histoire Naturelle & Service du Patrimoine Naturel, Paris, 496 pp.			
<i>Iberolacerta cyreni</i>	IUCN			
<i>Iberolacerta horvathi</i>	Bischoff, W. 1984. Lacerta agilis Linnaeus 1758 - Zauneidechse. Pages 23-68 in W. Bohme, editor. Handbuch der reptilien und amphibien Europas, Band 2/I: Echsen II (Lacerta). Aula-			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Verlag, Wiesbaden.			
<i>Iberolacerta monticola</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Ichnotropis capensis</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Ichnotropis squamulosa</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Iguana delicatissima</i>	Pasachnik, S., Breuil, M. and Powell, R. 2005. Iguana delicatissima. Catalogue of American Amphibians and Reptiles, 811: 1-14.			
<i>Iguana iguana</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Insulasaurus arborens</i>	http://philbreo.lifedesks.org/taxa?page=2			
<i>Insulasaurus traanorum</i>	Linkem, C. A., Diesmos, A. C. and Brown, R. M. 2010. A new scincid lizard (Genus Sphenomorphus) from Palawan Island, Philippines. Herpetologica 66: 67-79.			
<i>Intellagama lesueurii</i>	Cogger, H. G. 2000. Reptiles and			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Janetaescincus braueri</i>	Gerlach, J. 2008. Population and conservation status of the reptiles of the Seychelles islands. Phelsuma 16: 30-48.			
<i>Japalura brevipes</i>	Shang, G. S. and Lin, S. L. 2008. A field guide to lizards in Taiwan. Commonwealth Publishing.			
<i>Japalura kumaonensis</i>	Schleich, H. H. and Kastle, W. 2002. Amphibians and reptiles of Nepal. Gantner Verlag, Koenigstein.			
<i>Japalura luei</i>	Shang, G. S. and Lin, S. L. 2008. A field guide to lizards in Taiwan. Commonwealth Publishing.			
<i>Japalura makii</i>	Shang, G. S. and Lin, S. L. 2008. A field guide to lizards in Taiwan. Commonwealth Publishing.			
<i>Japalura polygonata</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Japalura swinhonis</i>	Lin, C-I. and Cheng, H-Y. 1990. T'ai-wan hsi i chih [A synopsis of the lizards of Taiwan]. [Taiwan Museum], T'ai-pei. 176 pp.			
<i>Kanakysaurus viviparus</i>	Sadlier, R. A., Bauer, A. M., Whitaker, A. H. and Smith, S. A. 2004. Two new			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	species of scincid lizards (Squamata) from the Massif de Kopeto, New Caledonia. Proceedings of the California Academy of Sciences 55: 208-221.			
<i>Kentropyx calcarata</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Kentropyx pelviceps</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Kentropyx striata</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Lacerta agilis</i>	Ananjeva, N. B., Munkhbayar, K., Orlov, N. L., Orlova, V. F., Semenov, D. V. and Terbish, K. 1997. Amphibians and reptiles of Mongolia. Russian Academy of Sciences, Moscow.			
<i>Lacerta schreiberi</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Lacerta strigata</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Sofia.			
<i>Lacerta trilineata</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Lacerta viridis</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Lampropholis delicata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lampropholis guichenoti</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Larutia seribuatensis</i>	Grismer Pers. Comm.			
<i>Laudakia caucasia</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Laudakia stellio</i>	Disi, A. M. Modrý, D., Necas, P. and Rifai, L. 2001. Amphibians and reptiles of the Hashemite kingdom of Jordan: an atlas and field guide. Edition Chimaira, Frankfurt am Main.			
<i>Leiocephalus eremitus</i>	Powell, R. 1999. Anolis longiceps. Catalogue of American Amphibians and Reptiles 693: 1-2.			
<i>Leiocephalus inaguae</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Leiocephalus psammodromus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Leiolepis reevesii</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Leiopisma telfairii</i>	Cheke, A. and Hume, J. P. 2008. Lost land of the dodo: The ecological history of Mauritius, Reunion, and Rodrigues. Yale University Press,			
<i>Lepidodactylus euaensis</i>	Allison A. pers. Comm.			
<i>Lepidodactylus herrei</i>	Brown, W. C. and Alcala, A. C. 1978. Philippine lizards of the family Gekkonidae. Silliman University Press,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Dumaguete City, Philippines.			
<i>Lepidodactylus listeri</i>	Cogger, H., Sadler, R. and Cameron, E. 1983. The terrestrial reptiles of Australia's island territories. Australian National Parks and Wildlife Service Special Publication 8 (11) 1-79.			
<i>Lepidodactylus lugubris</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Lepidodactylus manni</i>	Zug, G. R. 1991. The lizards of Fiji: natural history and systematics. Bishop Museum Bulletin of Zoology 2: 1-136.			
<i>Lepidodactylus vanuatuensis</i>	Allison A. pers. Comm.			
<i>Lepidophyma flavimaculatum</i>	Bezy, R. L. and Camarillo, J. L. 2002. Systematics of xantusiid lizards of the genus <i>Lepidophyma</i> . Los Angeles County Museum Contributions in Science 493: 1-41			
<i>Leposoma rugiceps</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Lerista borealis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Lerista bougainvillii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lerista labialis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lerista punctatovittata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lerista xanthura</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lialis burtonis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Liolaemus andinus</i>	Abdala, C. S. and Quinteros, S. 2008. A new species of <i>Liolaemus</i> (Iguania: Liolaemidae), endemic to Sierra de Fiambala, Catamarca, Argentina. Cuadernos de Herpetologia, 22: 35-47.			
<i>Liolaemus bellii</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Liolaemus bibronii</i>	Quinteros, A. S. 2012. Taxonomy of the <i>Liolaemus alticolor</i> – <i>bibronii</i> group			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Liolaemus boulengeri</i>	<p>(Iguania: Liolaemidae), with descriptions of two new species. <i>Herpetologica</i>, 68: 100-120.</p> <p>Abdala, C. S., Semhan, R. V., Moreno Azocar, D. L., Bonino, M., Paz, M. M. and Cruz, F. 2012. Taxonomic study and morphology based phylogeny of the patagonic clade <i>Liolaemus melanops</i> group (Iguania: Liolaemidae), with the description of three new taxa.. <i>Zootaxa</i> 3163: 1-32.</p>			
<i>Liolaemus elongatus</i>	<p>Abdala, C. S., Quinteros, A. S., Scrocchi, G. J. and Stazonelli, J. C. 2010. Three new species of the <i>Liolaemus elongatus</i> group (Iguania: Liolaemidae) from Argentina. <i>Cuadernos de Herpetologia</i>, 24: 93-109.</p>			
<i>Liolaemus huacahuasicus</i>	<p>Abdala, C. S. and Quinteros, S. 2008. A new species of <i>Liolaemus</i> (Iguania: Liolaemidae), endemic to Sierra de Fiambala, Catamarca, Argentina. <i>Cuadernos de Herpetologia</i>, 22: 35-47.</p>			
<i>Liolaemus koslowskyi</i>	<p>Abdala, C. S., Quinteros, A. S., Arias, F., Portelli, S. and Palavecino, A. 2011. A new species of the <i>Liolaemus darwini</i> group (Iguania: Liolaemidae) from Salta Province, Argentina. <i>Zootaxa</i> 2968: 26-38.</p>			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Liolaemus lemniscatus</i>	Quinteros, A. S. 2012. Taxonomy of the <i>Liolaemus alticolor</i> – <i>bibronii</i> group (Iguania: Liolaemidae), with descriptions of two new species. <i>Herpetologica</i> , 68: 100-120.			
<i>Liolaemus lineomaculatus</i>	Breitman, M. F., Avila, L. J., Sites, J. W. and Morando, M. 2011. Lizards from the end of the world: Phylogenetic relationships of the <i>Liolaemus lineomaculatus</i> section (Squamata: Iguania: Liolaemini). <i>Molecular Phylogenetics and Evolution</i> 59: 364-376.			
<i>Liolaemus lutzae</i>	Avila, L. J., Morando, M., Perez, D. R. and Sites, J. W. 2009. A new species of <i>Liolaemus</i> from Anelo sand dunes, northern Patagonia, Neuquen, Argentina, and molecular phylogenetic relationships of the <i>Liolaemus wiegmannii</i> species group (Squamata, Iguania, Liolaemini). <i>Zootaxa</i> 2234: 39-55.			
<i>Liolaemus nigromaculatus</i>	Donoso-Barros, R. 1966. Dos nuevos Gonatodes de Venezuela. <i>Publicacion Ocasional Museo Nacional de Historia Natural, Santiago (Chile)</i> 11: 3-32.			
<i>Liolaemus pictus</i>	Avila, L. J., Frutos, N., Morando, M., Perez, C. H. F. and Kozykariski, M. 2006. <i>Reptilia, Iguania, Liolaemini, Liolaemus petrophilus</i> and <i>Liolaemus</i>			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	pictus: distribution extension, filling gaps, new records. Check List 2: 65-69.			
<i>Liolaemus scolaroi</i>	Daniel Pincheira-Donoso, pers comm			
<i>Liolaemus signifer</i>	Donoso-Barros, R. 1966. Dos nuevos Gonatodes de Venezuela. Publicacion Ocasional Museo Nacional de Historia Natural, Santiago (Chile) 11: 3-32.			
<i>Liolaemus wiegmannii</i>	Avila, L. J., Morando, M., Perez, D. R. and Sites, J. W. 2009. A new species of <i>Liolaemus</i> from Anelo sand dunes, northern Patagonia, Neuquen, Argentina, and molecular phylogenetic relationships of the <i>Liolaemus wiegmannii</i> species group (Squamata, Iguania, Liolaemini). Zootaxa 2234: 39-55.			
<i>Liolaemus zullyae</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Liopholis inornata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Liopholis kintorei</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Liopholis modesta</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Liopholis striata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Liopholis whitii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lioscincus greeri</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Lioscincus maruia</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Lioscincus nigrofasciolatum</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Lioscincus novaecaledoniae</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Lioscincus steindachneri</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Lipinia leptosoma</i>	Allison A. pers. Comm.			
<i>Lipinia macrotympanum</i>	Sharma, R. C. 2002. The fauna of India and the adjacent countries: Reptilia: Volume 2: Sauria. Fauna of India and the adjacent countries. Zoological Survey of India.			
<i>Lipinia rabori</i>	IUCN			
<i>Lipinia rouxi</i>	Allison A. pers. Comm.			
<i>Lisssolepis coventryi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lophognathus longirostris</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Lucasium damaeum</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Luperosaurus macgregori</i>	Brown, J. H., Fox, B. J. Kelt, D. A. 2000. Assembly rules: desert rodent communities are structured at scales from local to continental. American Naturalist 156: 314-321.			
<i>Lygodactylus klugei</i>	Guarino Colli, pers. Comm.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Mabuya mabouya</i>	Hoogmoed, M. S. 1973. Notes on the herpetofauna of Surinam IV: the lizards and amphisbaenians of Surinam. W. Junk Publishers, Den Haag.			
<i>Marisora unimarginata</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241., Kohler et al. 2006			
<i>Marmorosphax boulinda</i>	Sadlier, R. A., Smith, S. A., Bauer, A. M. and Whitaker, A. H. 2009. Three new species of skink in the genus <i>Marmorosphax</i> Sadlier (Squamata: Scincidae) from New Caledonia. in Grandcolas P. (ed.), <i>Zoologia Neocaledonica</i> 7. Biodiversity studies in New Caledonia. Mémoires du Muséum national d'Histoire naturelle 198: 373-390.			
<i>Marmorosphax taom</i>	Sadlier, R. A., Smith, S. A., Bauer, A. M. and Whitaker, A. H. 2009. Three new species of skink in the genus <i>Marmorosphax</i> Sadlier (Squamata: Scincidae) from New Caledonia. in Grandcolas P. (ed.), <i>Zoologia Neocaledonica</i> 7. Biodiversity studies in New Caledonia. Mémoires du Muséum national d'Histoire naturelle 198: 373-390.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Marmorosphax tricolor</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Mediodactylus kotschyi</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Mediodactylus russowii</i>	Shi, L. and Zhao, E. 2011. A new gecko in the genus <i>Cyrtopodion</i> Fitzinger, 1843 (Reptilia: Squamata: Gekkonidae) from western China. <i>Herpetologica</i> 67: 186-193.			
<i>Menetia greyii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Merole anchietae</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Merole cuneirostris</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Merole suborbitalis</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Consultants, Gaborone.			
<i>Mesalina guttulata</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociación Herpetologica Española, Barcelona.			
<i>Mesalina olivieri</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociación Herpetologica Española, Barcelona.			
<i>Mesalina pasteuri</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociación Herpetologica Española, Barcelona.			
<i>Mesalina rubropunctata</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociación Herpetologica Española, Barcelona.			
<i>Mesaspis gadovii</i>	Karges, J. P. and Wright, J. W. 1987. A new species of Barisia (Sauria, Anguinae) from Oaxaca, Mexico. Contributions in Science of the Natural History Museum of Los Angeles County 381: 1-11.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Mesaspis juarezi</i>	Karges, J. P. and Wright, J. W. 1987. A new species of Barisia (Sauria, Anguidae) from Oaxaca, Mexico. Contributions in Science of the Natural History Museum of Los Angeles County 381: 1-11.			
<i>Mesaspis monticola</i>	Kohler, G. 2003. A new species of Morunasaurus from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Microlophus albemarlensis</i>	Swash, A. and Still, R. 2005. Birds, mammals, and reptiles of the Galapagos Islands. 2nd edition. Christopher Helm Publishers Ltd.			
<i>Microlophus atacamensis</i>	Donoso-Barros, R. 1966. Dos nuevos Gonatodes de Venezuela. Publicacion Ocasional Museo Nacional de Historia Natural, Santiago (Chile) 11: 3-32.			
<i>Microlophus delanonis</i>	Swash, A. and Still, R. 2005. Birds, mammals, and reptiles of the Galapagos Islands. 2nd edition. Christopher Helm Publishers Ltd.			
<i>Microlophus habelii</i>	Swash, A. and Still, R. 2005. Birds, mammals, and reptiles of the Galapagos Islands. 2nd edition. Christopher Helm Publishers Ltd.			
<i>Microlophus quadrivittatus</i>	Donoso-Barros, R. 1966. Dos nuevos Gonatodes de Venezuela. Publicacion			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Ocasional Museo Nacional de Historia Natural, Santiago (Chile) 11: 3-32.			
<i>Mniarogekko chahoua</i>	Bauer, A. M., Jackman, T. R., Sadlier, R. A. and Whitaker, A. H. 2012. Revision of the giant geckos of New Caledonia (Reptilia: Diplodactylidae: Rhacodactylus). Zootaxa 3404: 1-52.			
<i>Moloch horridus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Morethia boulengeri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Morethia obscura</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Nannoscincus garrulus</i>	Sadlier, R. A., Bauer, A. M. and Smith, S. A. 2006. A new species of <i>Nannoscincus</i> Günther (Squamata: Scincidae) from high elevation forest in southern New Caledonia. Records of the Australian Museum 58: 29-36.			
<i>Nannoscincus gracilis</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Nannoscincus greeri</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Nannoscincus maccoyi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Nannoscincus mariei</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Nephrurus levis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Niveoscincus coventryi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Niveoscincus metallicus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Nucras lalandii</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Nucras taeniolata</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Consultants, Gaborone.			
<i>Nucras tessellata</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Oedodera marmorata</i>	Bauer, A. M., Barts, M. and Hulbert, F. 2006. A new species of the <i>Pachydactylus weberi</i> group (Reptilia: Squamata: Gekkonidae) from the Orange River, with comments on its natural history. <i>Salamandra</i> 42: 83-92.			
<i>Oedura castelnaui</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Oedura monilis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Oedura tryoni</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Oligosoma fallai</i>	Pickard, C. R. and Towns, D. R. 1988. Atlas of the amphibians and reptiles of New Zealand. Department of Conservation Publications.			
<i>Oligosoma homalonotum</i>	Pickard, C. R. and Towns, D. R. 1988. Atlas of the amphibians and reptiles of New Zealand. Department of			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Conservation Publications.			
<i>Oligosoma lichenigera</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Oligosoma nigriplantare</i>	Patterson, G. B. and Daugherty, C. H. 1990. Four new species and one new subspecies of skinks, genus <i>Leiolopisma</i> (Reptilia: Lacertilia: Scincidae) from New Zealand. <i>Journal of the Royal Society of New Zealand</i> 20: 65-84.			
<i>Oligosoma notosaurus</i>	Patterson, G. B. and Daugherty, C. H. 1990. Four new species and one new subspecies of skinks, genus <i>Leiolopisma</i> (Reptilia: Lacertilia: Scincidae) from New Zealand. <i>Journal of the Royal Society of New Zealand</i> 20: 65-84.			
<i>Oligosoma stenotis</i>	http://www.doc.govt.nz			
<i>Ophiodes striatus</i>	Nogueira, C. D. 2006. Diversidade e padroes de distribuicao da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências, Universidade de São Paulo.			
<i>Ophiomorus punctatissimus</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. <i>Lurche und kriechtiere Europas</i> . Neumann Verlag, Radebeul, Germany.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Ophisaurus attenuatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Ophisaurus compressus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Ophisaurus ventralis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Ophisops elegans</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Pachydactylus mariquensis</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad, Project Report.			
<i>Pamelaescincus gardineri</i>	Gerlach, J. 2008. Population and conservation status of the reptiles of the			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Seychelles islands. Phelsuma 16: 30-48.			
<i>Panaspis kitsoni</i>	Fuhn, I. E. 1972. Revision du phylum forestier du genre <i>Panaspis</i> Cope (Reptilia, Scincidae, Lygosominae). Review Roumain de Biologie Série de Zoologie 17: 257-271.			
<i>Panaspis nimbaensis</i>	Fuhn, I. E. 1972. Revision du phylum forestier du genre <i>Panaspis</i> Cope (Reptilia, Scincidae, Lygosominae). Review Roumain de Biologie Série de Zoologie 17: 257-271.			
<i>Panaspis wahlbergi</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Parvilacerta parva</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Parvosцинus sisoni</i>	Brown, W. C. and Alcala, A. C. 1978. Philippine lizards of the family Gekkonidae. Silliman University Press, Dumaguete City, Philippines.			
<i>Pedioplanis burchelli</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Curtis Books. ISBN 0883590425pa.			
<i>Pedioplanis namaquensis</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Perochirus ateles</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Phelsuma andamanense</i>	IUCN			
<i>Phelsuma astriata</i>	Gerlach, J. 2008. Population and conservation status of the reptiles of the Seychelles islands. <i>Phelsuma</i> 16: 30-48.			
<i>Phelsuma borbonica</i>	Meier, H. 1995 Neue nachweise von <i>Phelsuma borbonica</i> auf Reunion, Maskarenen, mit dem versuch einer taxonomischen einordnung. <i>Salamandra</i> 31: 33-40.			
<i>Phelsuma cepediana</i>	Glaw, F. and Vences, M. 1994. A field guide to the amphibians and reptiles of Madagascar. Second edition, Serpents Tale.			
<i>Phelsuma comorensis</i>	Greckhamer, A. 1995. Bemerkungen zur haltung und zucht sowie zum verhalten von <i>Phelsuma comorensis</i> Boettger, 1913. im Terrarium. <i>Herpetofauna</i> 17: 6-16.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Phelsuma dubia</i>	Glaw, F. and Vences, M. 1994. A field guide to the amphibians and reptiles of Madagascar. Second edition, Serpents Tale.			
<i>Phelsuma guentheri</i>	Glaw, F. and Vences, M. 1994. A field guide to the amphibians and reptiles of Madagascar. Second edition, Serpents Tale.	Henkel, F. W. and Schmidt, W. 2000. Amphibians and reptiles of Madagascar and the Mascarene, Seychelles, and Comoro Islands. Kreiger, Malabar, Florida.		
<i>Phelsuma guimbeaui</i>	Henkel, F. W. and Schmidt, W. 2000. Amphibians and reptiles of Madagascar and the Mascarene, Seychelles, and Comoro Islands. Kreiger, Malabar, Florida.			
<i>Phelsuma inexpectata</i>	Vinson, J. and Vinson, J.-M. 1969. The saurian fauna of the Mascarene Islands. Mauritius Institute Bulletin, 6: 203-320.			
<i>Phelsuma nigra</i>	Hawlitschek, O., Bruckmann, B., Berger, J., Green, K. and Glaw, F. 2011. Integrating field surveys and remote sensing data to study distribution, habitat use and conservation status of the herpetofauna of the Comoro Islands. ZooKeys 144: 21-79.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Phelsuma nigristriata</i>	Hawlitschek, O., Bruckmann, B., Berger, J., Green, K. and Glaw, F. 2011. Integrating field surveys and remote sensing data to study distribution, habitat use and conservation status of the herpetofauna of the Comoro Islands. ZooKeys 144: 21-79.			
<i>Phelsuma ornata</i>	Schonecker, P. 2008. Geckos of Madagascar, the Seychelles, Comoros and Mascarene Islands. Aqualog Verlag Gmbh			
<i>Phelsuma parkeri</i>	Henkel, F. W. and Schmidt, W. 2000. Amphibians and reptiles of Madagascar and the Mascarene, Seychelles, and Comoro Islands. Kreiger, Malabar, Florida.			
<i>Phelsuma robertmertensi</i>	Hawlitschek, O., Bruckmann, B., Berger, J., Green, K. and Glaw, F. 2011. Integrating field surveys and remote sensing data to study distribution, habitat use and conservation status of the herpetofauna of the Comoro Islands. ZooKeys 144: 21-79.			
<i>Phelsuma rosagularis</i>	Henkel, F. W. and Schmidt, W. 2000. Amphibians and reptiles of Madagascar and the Mascarene, Seychelles, and Comoro Islands. Kreiger, Malabar, Florida.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Phelsuma sundbergi</i>	Schonecker, P. 2008. Geckos of Madagascar, the Seychelles, Comoros and Mascarene Islands. Aqualog Verlag Gmbh			
<i>Phoenicolacerta laevis</i>	Disi, A. M. Modrý, D., Necas, P. and Rifai, L. 2001. Amphibians and reptiles of the Hashemite kingdom of Jordan: an atlas and field guide. Edition Chimaira, Frankfurt am Main.			
<i>Phoenicolacerta troodica</i>	Baier, F., Sparrow, J. D. and Wiedl, H. - J. 2009. The amphibians and reptiles of Cyprus. Edition Chimaira, Frankfurt.			
<i>Phrynocephalus guttatus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Phrynocephalus helioscopus</i>	Ananjeva, N. B., Munkhbayar, K., Orlov, N. L., Orlova, V. F., Semenov, D. V. and Terbish, K. 1997. Amphibians and reptiles of Mongolia. Russian Academy of Sciences, Moscow.			
<i>Phrynocephalus mystaceus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Sofia.			
<i>Phrynocephalus theobaldi</i>	Schleich, H. H. and Kastle, W. 2002. Amphibians and reptiles of Nepal. Gantner Verlag, Koenigstein.			
<i>Phrynocephalus versicolor</i>	Ananjeva, N. B., Munkhbayar, K., Orlov, N. L., Orlova, V. F., Semenov, D. V. and Terbish, K. 1997. Amphibians and reptiles of Mongolia. Russian Academy of Sciences, Moscow.			
<i>Phrynosoma blainvillii</i>	Leache, A. D. and McGuire, J. A. 2006. Phylogenetic relationships of horned lizards (Phrynosoma) based on nuclear and mitochondrial data: Evidence for a misleading mitochondrial gene tree. Molecular Phylogenetics and Evolution 39: 628-644.			
<i>Phrynosoma cornutum</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Phrynosoma coronatum</i>	Leache, A. D. and McGuire, J. A. 2006. Phylogenetic relationships of horned lizards (Phrynosoma) based on nuclear and mitochondrial data: Evidence for a misleading mitochondrial gene tree. Molecular Phylogenetics and Evolution			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	39: 628-644.			
<i>Phrynosoma ditmarsii</i>	Hodges, W. L. 1995. <i>Phrynosoma ditmarsii</i> . Catalogue of American Amphibians and Reptiles 614: 1-3.			
<i>Phrynosoma douglassii</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Phrynosoma hernandesi</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Phrynosoma mcallii</i>	Funk, R. S. 1981. <i>Phrynosoma mcallii</i> . Catalogue of American Amphibians and Reptiles 281: 1			
<i>Phrynosoma modestum</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Phrynosoma platyrhinos</i>	Leache, A. D. and McGuire, J. A. 2006. Phylogenetic relationships of horned lizards (<i>Phrynosoma</i>) based on nuclear and mitochondrial data: Evidence for a misleading mitochondrial gene tree.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Molecular Phylogenetics and Evolution 39: 628-644.			
<i>Phrynosoma solare</i>	Leache, A. D. and McGuire, J. A. 2006. Phylogenetic relationships of horned lizards (Phrynosoma) based on nuclear and mitochondrial data: Evidence for a misleading mitochondrial gene tree. Molecular Phylogenetics and Evolution 39: 628-644.			
<i>Phyllodactylus bugastrolepis</i>	IUCN			
<i>Phyllodactylus lanei</i>	Dixon, J. R. 1964. Further data on the geckos (Phyllodactylus) of islands of the extreme southern Caribbean. Southwestern Naturalist, 9: 203-205.			
<i>Phyllopezus pollicaris</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Phyllurus platurus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Phymaturus palluma</i>	Corbalan, V., Scolaro, A. and Debandi, G. 2009. A new species of the genus Phymaturus of the flagellifer group from Central-Western Mendoza, Argentina (Reptilia: Iguania: Liolaemidae).			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Zootaxa 2021: 42-56.			
<i>Phymaturus patagonicus</i>	Lobo, F., Abdala, S. and Valdecantos, S. 2010. Taxonomic studies of the genus <i>Phymaturus</i> (Iguania: Liolaemidae): description of four new species. South American Journal of Herpetology 5: 102-126.			
<i>Phymaturus punae</i>	Cei, J. M. 1986. Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas aridas y semiaridas. Museo Regionale di Scienze Naturali, Torino. 527 pp.			
<i>Phymaturus zapalensis</i>	Avila, L. J., Perez, C. H. F., Perez, D. R. and Morando, M. 2011. Two new mountain lizard species of the <i>Phymaturus</i> genus (Squamata: Iguania) from northwestern Patagonia, Argentina. Zootaxa 2924: 1-21.			
<i>Physignathus cocincinus</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Plestiodon anthracinus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Plestiodon copei</i>	IUCN			
<i>Plestiodon egregius</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Plestiodon fasciatus</i>	Richmond, J. Q. 2006. Evolutionary basis of parallelism in North American scincid lizards. <i>Evolution & Development</i> 8: 477-490.			
<i>Plestiodon inexpectatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Plestiodon kishinouyei</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Plestiodon laticeps</i>	Richmond, J. Q. 2006. Evolutionary basis of parallelism in North American scincid lizards. <i>Evolution & Development</i> 8: 477-490.			
<i>Plestiodon lynxe</i>	IUCN			
<i>Plestiodon obsoletus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Boston.			
<i>Plestiodon okadae</i>	Hasegawa, M. 1994. Insular radiation in life history of the lizard <i>Eumeces okadae</i> in the Izu Islands, Japan. <i>Copeia</i> , 1994: 732-747.			
<i>Plestiodon reynoldsi</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Plestiodon septentrionalis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Plestiodon skiltonianus</i>	Smith, H. M. 1946. Handbook of lizards. Lizards of the United States and Canada. Cornell University Press, Ithaca.			
<i>Plestiodon stimpsonii</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Plestiodon tetragrammus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Plica plica</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Plica umbra</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Podarcis bocagei</i>	Sa-Sousa, P. and Harris, D. J. 2002. Podarcis carbonelli Perez-Mellado, 1981 is a distinct species. Amphibia-Reptilia 23: 459-468.			
<i>Podarcis carbonelli</i>	Kaliontzopoulou, A., Carretero, M. A. and Llorente, G. A. 2010. Intraspecific ecomorphological variation: linear and geometric morphometrics reveal habitat-related patterns within Podarcis bocagei wall lizards. Journal of Evolutionary Biology, 23: 1234-1244.			
<i>Podarcis cretensis</i>	Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. and Mylonas, M. 2008. Two new species of Podarcis (Squamata; Lacertidae) from Greece. Systematics and Biodiversity 6: 307-318.			
<i>Podarcis erhardii</i>	Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. and Mylonas, M. 2008. Two new species of Podarcis (Squamata; Lacertidae) from Greece. Systematics and Biodiversity 6:			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	307-318.			
<i>Podarcis filfolensis</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Podarcis gaigeae</i>	Valakos, E. D., Pafilis, P., Sotiropoulos, K., Lymberakis, P., Maragou, P. and Foufopoulos, J. 2008. The amphibians and reptiles of Greece. Edition Chimaira, Frankfurt Am Main.			
<i>Podarcis hispanicus</i>	IUCN			
<i>Podarcis lilfordi</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Podarcis melisellensis</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Podarcis milensis</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Podarcis muralis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	expanded. Houghton Mifflin Company, Boston.			
<i>Podarcis peloponnesiacus</i>	Lymberakis, P., Poulakakis, N., Kaliontzopoulou, A., Valakos, E. and Mylonas, M. 2008. Two new species of Podarcis (Squamata; Lacertidae) from Greece. Systematics and Biodiversity 6: 307-318.			
<i>Podarcis pityusensis</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Podarcis raffoneae</i>	Gasc, J. P., Cabela A., Crnobrnja-Isailovic, J., Dolmen, D., Grossenbacher, K., Haffner, P., Lescure, J., Martens, H., Martínez Rica, J. P., Maurin, H., Oliveira, M. E., Sofianidou, T. S., Veith, M. and Zuiderwijk, A. (eds), 1997. Atlas of amphibians and reptiles in Europe. Collection Patrimoines Naturels, 29, Societas Europaea Herpetologica, Muséum National d'Histoire Naturelle & Service du Patrimoine Naturel, Paris, 496 pp.			
<i>Podarcis siculus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Podarcis tauricus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Podarcis tiliguerta</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Podarcis waglerianus</i>	Engelmann, W. E., Fritzsche, J., Gunther, R. and Obst, F. J. 1993. Lurche und kriechtiere Europas. Neumann Verlag, Radebeul, Germany.			
<i>Pogona barbata</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Pogona minor</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Pogona vitticeps</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Polychrus acutirostris</i>	Garda, A. A., Costa, G. C., Franca, F. G. R., Giugliano, L. G., Leite, G. S., Mesquita, D. O., Nogueira, C., Tavares-Bastos, L., Vasconcellos, M. M., Vieira, G. H. C., Vitt, L. J., Werneck, F. P.,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Wiederhecker, H. C. and Colli, G. R. 2012. Reproduction, body size, and diet of <i>Polychrus acutirostris</i> (Squamata: Polychrotidae) in two contrasting environments in Brazil. <i>Journal of Herpetology</i> , 46: 2-8.			
<i>Potamites ecpleopus</i>	Avila-Pires, T. C. S. 1995. <i>Lizards of Brazilian Amazonia</i> . Backhuys Publishers, Leiden			
<i>Pristurus abdelkuri</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. <i>Zootaxa</i> 2826: 1-44 (Reptilia)			
<i>Pristurus guichardi</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. <i>Zootaxa</i> 2826: 1-44 (Reptilia)			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Pristurus insignis</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. Zootaxa 2826: 1-44 (Reptilia)			
<i>Pristurus insignoides</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of the Socotran Archipelago Herpetofauna. Zootaxa 2826: 1-44 (Reptilia)			
<i>Pristurus rupestris</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Pristurus sokotranus</i>	Razzetti, E., Sindaco, R., Grieco, C., Pella, F., Ziliani, U., Pupin, F., Riservato, E., Pellitteri-Rosa, D., Butikofer, L., Suleiman, A. S., Al-Aseily, B. A., Carugati, C., Boncompagni, E. and Fasola, M. 2011. Annotated checklist and distribution of			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	the Socotran Archipelago Herpetofauna. Zootaxa 2826: 1-44 (Reptilia)			
<i>Psammodromus algirus</i>	Busack, S. D., Salvador, A. and Lawson, R. 2006. Two new species in the genus <i>Psammodromus</i> (Reptilia: Lacertidae) from the Iberian Peninsula. <i>Annals of Carnegie Museum</i> 75: 1-10.			
<i>Psammodromus blanci</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociación Herpetologica Española, Barcelona.			
<i>Psammodromus hispanicus</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociación Herpetologica Española, Barcelona.			
<i>Pseudemoia entrecasteauxii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Pseudemoia pagenstecheri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Pseudemoia rawlinsoni</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Island, FL : Ralph Curtis Publishing.			
<i>Pseudemoia spenceri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Pseudocordylus melanotus</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Pseudogekko smaragdinus</i>	IUCN			
<i>Pseudopus apodus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Psychosaura macrorhyncha</i>	Rodrigues, M. T. 2000. A new species of Mabuya (Squamata: Scincidae) from the semiarid Caatingas of northeastern Brazil. Papeis Avulsos de Zoologia 41: 313-328.			
<i>Ptenopus garrulus</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Ptychozoon nicobarensis</i>	Das, I. and Vijayakumar, S. P. 2009. New species of Ptychozoon (Sauria: Gekkonidae) from the Nicobar Archipelago, Indian Ocean. Zootaxa			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	2095: 8-20.			
<i>Ptyodactylus guttatus</i>	Disi, A. M. Modrý, D., Necas, P. and Rifai, L. 2001. Amphibians and reptiles of the Hashemite kingdom of Jordan: an atlas and field guide. Edition Chimaira, Frankfurt am Main.			
<i>Ptyodactylus oudrii</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biog*ographique. Asociacion Herpetologica Espanola, Barcelone.			
<i>Ptyodactylus ragazzii</i>	Chirio, L. and LeBreton, M. 2007. Atlas des reptiles du Cameroun. Publications Scientifiques du Museum national d'Histoire naturelle, Paris.			
<i>Pygopus lepidopodus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Quedenfeldtia trachyblepharus</i>	Arnold, E. N. 1990. The two species of Moroccan day-geckoes, <i>Quedenfeldtia</i> (Reptilia: Gekkonidae). Journal of Natural History 24: 757-762.			
<i>Rhacodactylus auriculatus</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Rhacodactylus leachianus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Rhacodactylus trachyrhynchus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Rhampholeon marshalli</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Riama shrevei</i>	Rivas, G., Schargel W. E. and Meik, J. M. 2005. A new species of Riama (Squamata: Gymnophthalmidae), endemic to the Península de Paria, Venezuela. Herpetologica 61: 461-468.			
<i>Saltuarius cornutus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Sauromalus ater</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Sauromalus hispidus</i>	Grismer, L. L, Beaman, K. R. and Lawler, H. E. 1995. Sauromalus hispidus. Catalogue of American			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Amphibians and Reptiles 615: 1-4.			
<i>Sauromalus varius</i>	Lawler, H. E., Beaman, K. R. and Grismer, L. L. 1995. <i>Sauromalus varius</i> . Catalogue of American Amphibians and Reptiles 616: 1-4.			
<i>Scelarcis perspicillata</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biogéographique. Asociacion Herpetologica Espanola, Barcelone.			
<i>Sceloporus aeneus</i>	Benabib, M., Kjer, K. M. and Sites, J. W. 1997. Mitochondrial DNA sequence-based phylogeny and the evolution of viviparity in the <i>Sceloporus scalaris</i> group (Reptilia, Squamata). <i>Evolution</i> 51: 1262-1275.			
<i>Sceloporus angustus</i>	IUCN			
<i>Sceloporus arenicolus</i>	Conant, R. and Collins, J. T. 1998. <i>Reptiles and amphibians of eastern and central North America</i> . 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus clarkii</i>	Smith, H. M. 1939. Notes on Mexican reptiles and amphibians. <i>Zoological Series of Field Museum of Natural History</i> 24: 15-35.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Sceloporus consobrinus</i>	Smith, H. M. 1946. Handbook of lizards. Lizards of the United States and Canada. Cornell University Press, Ithaca.			
<i>Sceloporus cozumelae</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - <i>Senckenbergiana Biologica</i> 82: 235-241.			
<i>Sceloporus cyanogenys</i>	Martinez-Mendez, N. and Mendez-de la Cruz, F. R. 2007. Molecular phylogeny of the <i>Sceloporus torquatus</i> species-group (Squamata: Phrynosomatidae). <i>Zootaxa</i> 1609: 53-68.			
<i>Sceloporus formosus</i>	Smith, H. M. 1939. Notes on Mexican reptiles and amphibians. Zoological Series of Field Museum of Natural History 24: 15-35.			
<i>Sceloporus gadoviae</i>	Smith, H. M. 1939. Notes on Mexican reptiles and amphibians. Zoological Series of Field Museum of Natural History 24: 15-35.			
<i>Sceloporus graciosus</i>	Censky, E. J. 1986. <i>Sceloporus graciosus</i> . Catalogue of American Amphibians and Reptiles 386: 1-4.			
<i>Sceloporus grammicus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Boston.			
<i>Sceloporus horridus</i>	Smith, H. M. 1939. Notes on Mexican reptiles and amphibians. Zoological Series of Field Museum of Natural History 24: 15-35.			
<i>Sceloporus jarrovi</i>	Martinez-Mendez, N. and Mendez-de la Cruz, F. R. 2007. Molecular phylogeny of the <i>Sceloporus torquatus</i> species-group (Squamata: Phrynosomatidae). Zootaxa 1609: 53-68.			
<i>Sceloporus magister</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus malachiticus</i>	Kohler, G. 2003. A new species of <i>Morunasaurus</i> from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Sceloporus merriami</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus mucronatus</i>	Lynch, J. D. and Smith, H. M. 1965. New or unusual amphibians and reptiles from Oaxaca, Mexico. I. Herpetologica			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	24: 168-177.			
<i>Sceloporus occidentalis</i>	Bell, E. L. and Price, A. H. 1996. <i>Sceloporus occidentalis</i> . Catalogue of American Amphibians and Reptiles 631: 1-17.			
<i>Sceloporus olivaceus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus omiltemanus</i>	Martinez-Mendez, N. and Mendez-de la Cruz, F. R. 2007. Molecular phylogeny of the <i>Sceloporus torquatus</i> species-group (Squamata: Phrynosomatidae). <i>Zootaxa</i> 1609: 53-68.			
<i>Sceloporus orcutti</i>	Smith, H. M. 1939. Notes on Mexican reptiles and amphibians. Zoological Series of Field Museum of Natural History 24: 15-35.			
<i>Sceloporus poinsettii</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus scalaris</i>	Benabib, M., Kjer, K. M. and Sites, J. W. 1997. Mitochondrial DNA sequence-based phylogeny and the evolution of viviparity in the <i>Sceloporus scalaris</i>			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	group (Reptilia, Squamata). Evolution 51: 1262-1275.			
<i>Sceloporus serrifer</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus slevini</i>	Smith, H. M. 1946. Handbook of lizards. Lizards of the United States and Canada. Cornell University Press, Ithaca.			
<i>Sceloporus torquatus</i>	Martinez-Mendez, N. and Mendez-de la Cruz, F. R. 2007. Molecular phylogeny of the <i>Sceloporus torquatus</i> species-group (Squamata: Phrynosomatidae). Zootaxa 1609: 53-68.			
<i>Sceloporus undulatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Sceloporus utiformis</i>	Smith, H. M. 1939. Notes on Mexican reptiles and amphibians. Zoological Series of Field Museum of Natural History 24: 15-35.			
<i>Sceloporus variabilis</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Boston.			
<i>Sceloporus virgatus</i>	Cole, C. J. 1968. <i>Sceloporus virgatus</i> . Catalogue of American Amphibians and Reptiles 72: 1-2.			
<i>Sceloporus woodi</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Scincella lateralis</i>	Brooks, G. R. 1975. <i>Scincella lateralis</i> . Catalogue of American Amphibians and Reptiles 169: 1-4.			
<i>Scincus mitranus</i>	Sindaco, R. and Jeremcenko, V. K. 2008. The reptiles of the Western Palearctic. 1. Annotated checklist and distributional atlas of the turtles, crocodiles, amphisbaenians and lizards of Europe, North Africa, Middle East and Central Asia. Edizioni Belvedere, Latina (Italy).			
<i>Scincus scincus</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Sigaloseps deplanchei</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Sigaloseps ruficauda</i>	Bauer, A. M. and Sadler, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Sitana fusca</i>	Schleich, H. H. and Kastle, W. 2002. Amphibians and reptiles of Nepal. Gantner Verlag, Koenigstein.			
<i>Sitana ponticeriana</i>	Tikader, B. K. and Sharma, R. C. 1992. Handbook of Indian lizards. Zoological Survey of India, Calcutta.			
<i>Sitana sivalensis</i>	Schleich, H. H. and Kastle, W. 2002. Amphibians and reptiles of Nepal. Gantner Verlag, Koenigstein.			
<i>Smaug giganteus</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad, Project Report.			
<i>Sphaerodactylus argivus</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Sphaerodactylus kirbyi</i>	Hite, J. L., Steinberg, D. S. and Powell, R. 2007. <i>Sphaerodactylus kirbyi</i> . Catalogue of American Amphibians and Reptiles, 852: 1-2.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Sphaerodactylus nicholsi</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Sphaerodactylus roosevelti</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Sphaerodactylus vincenti</i>	Schwartz, A. and Henderson, R. W. 1991. Amphibians and Reptiles of the West Indies. University of Florida Press, Gainesville.			
<i>Sphenomorphus concinnatus</i>	Allison A. pers. Comm.			
<i>Sphenomorphus cranei</i>	Greer, A. E. and Parker, F. A. 1974. The fasciatus species group of <i>Sphenomorphus</i> (Lacertilia: Scincidae): notes on eight previously described species and descriptions of three new species. Papua New Guinea Scientific Society Proceedings 25: 31-61.			
<i>Sphenomorphus indicus</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Sphenomorphus maculatus</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Sphenomorphus scutatus</i>	Allison A. pers. Comm.			
<i>Stenocercus chrysopygus</i>	Cadle, J. E. 1991. Systematics of lizards of the genus <i>Stenocercus</i> (Iguania Tropiduridae) from northern Peru. New species and comments on relationships and distribution patterns. Proceedings of the Academy of Natural Sciences of Philadelphia 143: 1-96.			
<i>Stenocercus dumerilii</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Stenodactylus doriae</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Strophurus ciliaris</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Strophurus elderi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Strophurus intermedius</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Strophurus williamsi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Takydromus amurensis</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Takydromus dorsalis</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Takydromus formosanus</i>	Lue, K-Y. and Lin, S-M. 2008. Two new cryptic species of <i>Takydromus</i> (Squamata: Lacertidae) from Taiwan. <i>Herpetologica</i> , 64: 379-395.			
<i>Takydromus hsuehshanensis</i>	Shang, G. S. and Lin, S. L. 2008. A field guide to lizards in Taiwan. Commonwealth Publishing.			
<i>Takydromus luyeanus</i>	Lue, K-Y. and Lin, S-M. 2008. Two new cryptic species of <i>Takydromus</i> (Squamata: Lacertidae) from Taiwan. <i>Herpetologica</i> , 64: 379-395.			
<i>Takydromus sauteri</i>	Lin, C-I. and Cheng, H-Y. 1990. T'ai-wan hsi i chih [A synopsis of the lizards of Taiwan]. [Taiwan Museum], T'ai-pei. 176 pp.			
<i>Takydromus septentrionalis</i>	Yuezhao Wang, pers. Com.			
<i>Takydromus sexlineatus</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian:			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Yashcheritsy Vietnam). Pensoft.			
<i>Takydromus smaragdinus</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Takydromus stejnegeri</i>	Lin, C-I. and Cheng, H-Y. 1990. T'ai-wan hsi i chih [A synopsis of the lizards of Taiwan]. [Taiwan Museum], T'ai-pei. 176 pp.			
<i>Takydromus toyamai</i>	Goris, R. C. and Maeda, N. 2004. Guide to the amphibians and reptiles of Japan. Krieger Publications, Malabar, FL.			
<i>Takydromus viridipunctatus</i>	Lue, K-Y. and Lin, S-M. 2008. Two new cryptic species of Takydromus (Squamata: Lacertidae) from Taiwan. Herpetologica, 64: 379-395.			
<i>Takydromus wolteri</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Tarentola angustimentalis</i>	Joger, U. 1984. Die radiation der gattung Tarentola in Makaronesien. Courier Forschungsinstitut Senckenberg 71: 91-111.			
<i>Tarentola annularis</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Occidental compris) Atlas Biog'ographique. Asociacion Herpetologica Espanola, Barcelone.			
<i>Tarentola boettgeri</i>	Arnold, E. N. and Ovenden, D. W. 2004. A field guide to the reptiles and amphibians of Britain and Europe. 2nd edition. Collins, London.			
<i>Tarentola chazaliae</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biog'ographique. Asociacion Herpetologica Espanola, Barcelone.	Le Berre, M. 1989. Faune du Sahara 1. Poissons, amphibiens, reptiles. Lechevalier, R. Chabaud, Paris.	Schleich, H. H., Kastle, W. and Kabisch, K. 1996. Amphibians and reptiles of North Africa. Biology, systematics, field guide. Koeltz Scientific, Koenigstein, Germany.	
<i>Tarentola darwini</i>	Vasconcelos, R., Brito, J., Carvalho, S. B., Carranza, S. and Harris, D. J. 2012. Identifying priority areas for island endemics using genetic versus specific diversity - the case of terrestrial reptiles of the Cape Verde Islands. Biological Conservation, 153, 276-286.			
<i>Tarentola delalandii</i>	Arnold, E. N. and Ovenden, D. W. 2004. A field guide to the reptiles and amphibians of Britain and Europe. 2nd edition. Collins, London.			
<i>Tarentola deserti</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biog'ographique. Asociacion			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Herpetologica Espanola, Barcelone.			
<i>Tarentola gomerensis</i>	Arnold, E. N. and Ovenden, D. W. 2004. A field guide to the reptiles and amphibians of Britain and Europe. 2nd edition. Collins, London.			
<i>Tarentola mauritanica</i>	Bons, J. and Geniez, P. 1996. Amphibiens et reptiles du Maroc (Sahara Occidental compris) Atlas Biog'ographique. Asociacion Herpetologica Espanola, Barcelone.			
<i>Teira dugesii</i>	Arnold, E. N. and Ovenden, D. W. 2004. A field guide to the reptiles and amphibians of Britain and Europe. 2nd edition. Collins, London.			
<i>Teius oculatus</i>	Avila, L. J. 2002. Geographic distribution of lizards of the Genus teius (Squamata: Teiidae: Teiinae) in southern South America. Biogeographica 78: 15-33.			
<i>Teius teyou</i>	Avila, L. J. 2002. Geographic distribution of lizards of the Genus teius (Squamata: Teiidae: Teiinae) in southern South America. Biogeographica 78: 15-33.			
<i>Teratoscincus microlepis</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Teratoscincus scincus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Tiliqua adelaidensis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Tiliqua nigrolutea</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Tiliqua occipitalis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Tiliqua rugosa</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Tiliqua scincoides</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Timon lepidus</i>	Bischoff, W. 1984. <i>Lacerta agilis</i> Linnaeus 1758 - Zauneidechse. Pages 23-68 in W. Bohme, editor. Handbuch der reptilien und amphibien Europas, Band 2/I: Echsen II (Lacerta). Aula-Verlag, Wiesbaden.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Timon pater</i>	Schleich, H. H., Kastle, W. and Kabisch, K. 1996. Amphibians and reptiles of North Africa. Biology, systematics, field guide. Koeltz Scientific, Koenigstein, Germany.			
<i>Toropuku stephensi</i>	Pickard, C. R. and Towns, D. R. 1988. Atlas of the amphibians and reptiles of New Zealand. Department of Conservation Publications.			
<i>Trachylepis affinis</i>	Chirio, L. and LeBreton, M. 2007. Atlas des reptiles du Cameroun. Publications Scientifiques du Museum national d'Histoire naturelle, Paris.			
<i>Trachylepis aurata</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Trachylepis buettneri</i>	Chirio, L. and LeBreton, M. 2007. Atlas des reptiles du Cameroun. Publications Scientifiques du Museum national d'Histoire naturelle, Paris.			
<i>Trachylepis lavarambo</i>	Ramanamanjato, J. B., Nussbaum, R. A. and Raxworthy, C. J. 1999. A new species of Mabuya Fitzinger (Reptilia: Squamata: Scincidae) from the Onilahy River of south-west Madagascar. Herpetological Journal 9: 65-72.			
<i>Trachylepis maculilabris</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Trachylepis quinquetaeniata</i>	Branch, W. R. 1998. Field guide to snakes and other reptiles of Southern Africa. 3rd ed. Sanibel Island, FL: Ralph Curtis Books. ISBN 0883590425pa.			
<i>Trachylepis sechellensis</i>	Gerlach, J. 2008. Population and conservation status of the reptiles of the Seychelles islands. Phelsuma 16: 30-48.			
<i>Trachylepis sparsa</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Trachylepis spilogaster</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Trachylepis striata</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Trachylepis varia</i>	Auerbach, R. D. 1987. The amphibians and reptiles of Botswana. Nokwepa Consultants, Gaborone.			
<i>Trachylepis vittata</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Trachylepis wrightii</i>	Gerlach, J. 2008. Population and conservation status of the reptiles of the			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Seychelles islands. Phelsuma 16: 30-48.			
<i>Trapelus mutabilis</i>	Schleich, H. H., Kastle, W. and Kabisch, K. 1996. Amphibians and reptiles of North Africa. Biology, systematics, field guide. Koeltz Scientific, Koenigstein, Germany.			
<i>Trapelus ruderatus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Trapelus sanguinolentus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Tribolonotus blanchardi</i>	Allison A. pers. Comm.			
<i>Tribolonotus brongersmai</i>	IUCN			
<i>Tribolonotus ponceleti</i>	Greer, A. E. and Parker, F. 1968. A new species of <i>Tribolonotus</i> (Lacertilia: Scincidae) from Bougainville and Buka, Solomon Islands, with comments on the biology of the genus. <i>Breviora</i> 291: 1-23.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Tribolonotus pseudoponceleti</i>	Greer, A. E. and Parker, F. 1968. A new species of <i>Tribolonotus</i> (Lacertilia: Scincidae) from Bougainville and Buka, Solomon Islands, with comments on the biology of the genus. <i>Breviora</i> 291: 1-23.			
<i>Tribolonotus schmidti</i>	Allison A. pers. Comm.			
<i>Trioceros ellioti</i>	Stipala, J., Lutzmann, N., Malonza, P. K., Borghesio, L., Wilkinson, P., Godley, B. and Evans, M. R. 2011. A new species of chameleon (Sauria: Chamaeleonidae) from the highlands of northwest Kenya. <i>Zootaxa</i> 3002: 1-16.			
<i>Trioceros hoehnelii</i>	Stipala, J., Lutzmann, N., Malonza, P. K., Borghesio, L., Wilkinson, P., Godley, B. and Evans, M. R. 2011. A new species of chameleon (Sauria: Chamaeleonidae) from the highlands of northwest Kenya. <i>Zootaxa</i> 3002: 1-16.			
<i>Trioceros jacksonii</i>	Branch, W. R. 2005. A Photographic guide to snakes other reptiles and amphibians of East Africa. Struik Nature, Capetown.			
<i>Trioceros quadricornis</i>	Chirio, L. and LeBreton, M. 2007. Atlas des reptiles du Cameroun. Publications Scientifiques du Museum national d'Histoire naturelle, Paris.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Trogonophis wiegmanni</i>	Sindaco, R. and Jeremcenko, V. K. 2008. The reptiles of the Western Palearctic. 1. Annotated checklist and distributional atlas of the turtles, crocodiles, amphisbaenians and lizards of Europe, North Africa, Middle East and Central Asia. Edizioni Belvedere, Latina (Italy).			
<i>Tropidoscincus boreus</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Tropidoscincus variabilis</i>	Bauer, A. M. and Sadlier, R. A. 2000. The herpetofauna of New Caledonia. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Tropidurus etheridgei</i>	Nogueira, C. D. 2006. Diversidade e padroes de distribuicao da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências, Universidade de São Paulo.			
<i>Tropidurus hispidus</i>	Nogueira, C. D. 2006. Diversidade e padroes de distribuicao da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências, Universidade de São Paulo.			
<i>Tropidurus semitaeniatus</i>	Nogueira, C. D. 2006. Diversidade e padroes de distribuicao da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Universidade de São Paulo.			
<i>Tropidurus spinulosus</i>	Nogueira, C. D. 2006. Diversidade e padroes de distribuicao da fauna de lagartos do cerrado. PhD Dissertation, USP-Instituto de Biociências, Universidade de São Paulo.			
<i>Tropidurus torquatus</i>	Cei, J. M. 1993. Reptiles del noroeste, nordeste y este de la Argentina. Museo Regionale di Scienze Naturali, Torino, Monografie 14: 1-949.			
<i>Tuketuku rakiurae</i>	Pickard, C. R. and Towns, D. R. 1988. Atlas of the amphibians and reptiles of New Zealand. Department of Conservation Publications.			
<i>Tupinambis teguixin</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Uma exsul</i>	Schmidt, K. P. and Bogert, C. M. 1947. A new fringe-footed sand lizard from Coahuila, Mexico. American Museum novitates 1339: 1-9.			
<i>Uma notata</i>	Pough, F. H. 1977. Uma notata. Catalogue of American Amphibians and Reptiles 197: 1-2.			
<i>Underwoodisaurus milii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Island, FL : Ralph Curtis Publishing.			
<i>Uracentron flaviceps</i>	Avila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia. Backhuys Publishers, Leiden			
<i>Urocotyledon inexpectata</i>	IUCN			
<i>Uromastix acanthinura</i>	Le Berre, M. 1989. Faune du Sahara 1. Poissons, amphibiens, reptiles. Lechevalier, R. Chabaud, Paris.			
<i>Uromastix aegyptia</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Uromastix ornata</i>	Nemtzov, S. C. 2008. Uromastix lizards in Israel. NDF workshop case studies, WG 7 – reptiles and amphibians, case study 5			
<i>Urosaurus bicarinatus</i>	Kohler, G. 2003. A new species of Morunasaurus from Peru (Reptilia, Squamata, Hoplocercidae). - Senckenbergiana Biologica 82: 235-241.			
<i>Urosaurus clarionensis</i>	Feldman, C. R., Flores-Villela, O. and Papenfuss, T. J. 2011. Phylogeny, biogeography, and display evolution in the tree and brush lizard genus Urosaurus (Squamata: Phrynosomatidae). Molecular Phylogenetics and Evolution 61: 714-			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	725.			
<i>Urosaurus graciosus</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Urosaurus ornatus</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Uta palmeri</i>	Grismer, L. L. 1994. Three new species of intertidal side-blotched lizards (genus Uta) from the Gulf of California, Mexico. Herpetologica 50: 451-474.			
<i>Uta stansburiana</i>	Conant, R. and Collins, J. T. 1998. Reptiles and amphibians of eastern and central North America. 3rd Edition expanded. Houghton Mifflin Company, Boston.			
<i>Varanus acanthurus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus albigularis</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad,			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	Project Report.			
<i>Varanus beccarii</i>	Eidenmueller, B. and Wicker, R. 2005. Eine weitere neue waranart aus dem Varanus prasinus-komplex von der insel Misol, Indonesien. Sauria 27: 3-8.			
<i>Varanus bengalensis</i>	Anderson, S. C. 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, St. Louis.			
<i>Varanus boehmei</i>	Eidenmueller, B. and Wicker, R. 2005. Eine weitere neue waranart aus dem Varanus prasinus-komplex von der insel Misol, Indonesien. Sauria 27: 3-8.			
<i>Varanus caerulivirens</i>	Pianka, E. R. and King, D. R. editors. 2004. Varanoid lizards of the world. Indiana University Press.			
<i>Varanus caudolineatus</i>	Aplin, K. P., Fitch, A. J. and King, D. J. 2006. A new species of Varanus Merrem (Squamata: Varanidae) from the Pilbara region of Western Australia, with observations on sexual dimorphism in closely related species. Zootaxa 1313: 1- 38.			
<i>Varanus cerambonensis</i>	Ziegler, T., Schmitz, A., Koch, A. and Bohme, W. 2007. A review of the subgenus Euprepiosaurus of Varanus (Squamata: Varanidae): morphological and molecular phylogeny, distribution			

Species	Map source 1	Map source 2	Map source 3	Map source 4
	and zoogeography, with an identification key for the members of the <i>V. indicus</i> and the <i>V. prasinus</i> species groups. Zootaxa 1472: 1-28.			
<i>Varanus flavescens</i>	Auffenberg, W., Rahman, H., Iffat, F. and Perveen, Z. 1989. A study of <i>Varanus flavescens</i> (Hardwicke & Gray) (Sauria: Varanidae). Journal of the Bombay Natural History Society 86: 286-307.			
<i>Varanus giganteus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus gilleni</i>	Aplin, K. P., Fitch, A. J. and King, D. J. 2006. A new species of <i>Varanus Merrem</i> (Squamata: Varanidae) from the Pilbara region of Western Australia, with observations on sexual dimorphism in closely related species. Zootaxa 1313: 1-38.			
<i>Varanus glauerti</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus gouldii</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Varanus griseus</i>	Ananjeva, N. B., Orlov, N. L., Khalikov, R. G., Darevsky, I. S., Ryabov, S. A. and Barabanov, A. V. 2006. The reptiles of northern Eurasia. Taxonomic diversity, distribution, conservation status. Pensoft, Sofia.			
<i>Varanus kingorum</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus komodoensis</i>	Murphy, J. B., Schlager, N., Trumpey, J. E. and Hutchins, M. 2003. Grzimek's Animal Life Encyclopedia, 2nd edition, Volume 7: Reptiles. Thomson-Gale, Detroit.			
<i>Varanus lirungensis</i>	Koch, A., Auliya, M., Schmitz, A., Kuch, U. and Bohme, W. 2007. Morphological studies on the systematics of south east Asian water monitors (Varanus salvator Complex): nominotypic populations and taxonomic overview. Mertensiella 16: 109-180.			
<i>Varanus mabitang</i>	Pianka, E. R. and King, D. R. editors. 2004. Varanoid lizards of the world. Indiana University Press.			
<i>Varanus macraei</i>	Eidenmueller, B. and Wicker, R. 2005. Eine weitere neue waranart aus dem Varanus prasinus-komplex von der insel Misol, Indonesien. Sauria 27: 3-8.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Varanus melinus</i>	Pianka, E. R. and King, D. R. editors. 2004. Varanoid lizards of the world. Indiana University Press.			
<i>Varanus mertensi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus niloticus</i>	Alexander, K. 2007. A photographic field guide to the reptiles and amphibians of Dominica, West Indies. Texas A&M University & Dominica Study Abroad, Project Report.			
<i>Varanus nuchalis</i>	Koch, A. Auliya, M. and Ziegler, T. 2010. Updated checklist of the living monitor lizards of the world (Squamata: Varanidae). Bonn Zoological Bulletin 57: 127-136.			
<i>Varanus ornatus</i>	Chirio, L. and LeBreton, M. 2007. Atlas des reptiles du Cameroun. Publications Scientifiques du Museum national d'Histoire naturelle, Paris.			
<i>Varanus panoptes</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus prasinus</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Varanus rosenbergi</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus rudicollis</i>	Grismer maps			
<i>Varanus salvator</i>	Bobrov, V. V. and Semenov, D. V. 2008. The lizards of Vietnam. (in Russian: Yashcheritsy Vietnama). Pensoft.			
<i>Varanus scalaris</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus spenceri</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus tristis</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varanus varius</i>	Cogger, H. G. 2000. Reptiles and amphibians of Australia. 6th ed. Sanibel Island, FL : Ralph Curtis Publishing.			
<i>Varzea bistriata</i>	Avila-Pires 1995, Miralles and Carranza 2010, Miralles et al. 2006			
<i>Xantusia arizonae</i>	Bezy, R. L., Bezy, K. B. and Bolles, K. 2008. Two new species of night lizards (Xantusia) from Mexico. Journal of Herpetology, 42: 680-688.			

Species	Map source 1	Map source 2	Map source 3	Map source 4
<i>Xantusia henshawi</i>	Bezy, R. L. and Flores Villela, O. 1999. A new species of Xantusia (Squamata: Xantusiidae) from Zacatecas, Mexico. <i>Herpetologica</i> 55: 174-184.			
<i>Xantusia riversiana</i>	Bezy, R. L. and Flores Villela, O. 1999. A new species of Xantusia (Squamata: Xantusiidae) from Zacatecas, Mexico. <i>Herpetologica</i> 55: 174-184.			
<i>Xantusia vigilis</i>	Bezy, R. L. 1982. <i>Xantusia vigilis</i> . <i>Catalogue of American Amphibians and Reptiles</i> 302: 1-4.			
<i>Xantusia wigginsi</i>	Bezy, R. L., Bezy, K. B. and Bolles, K. 2008. Two new species of night lizards (Xantusia) from Mexico. <i>Journal of Herpetology</i> , 42: 680-688.			
<i>Xenosaurus grandis</i>	Ballinger, R. E., Lemos-Espinal, J. A. and Smith, G. R. 2000. Reproduction in females of three species of crevice-dwelling lizards (genus <i>Xenosaurus</i>) from Mexico. <i>Studies on Neotropical Fauna and Environment</i> , 35: 179-183.			
<i>Zootoca vivipara</i>	Ananjeva, N. B., Munkhbayar, K., Orlov, N. L., Orlova, V. F., Semenov, D. V. and Terbish, K. 1997. <i>Amphibians and reptiles of Mongolia</i> . Russian Academy of Sciences, Moscow.			

APPENDIX S3 Parameter estimates for the best models

A. Parameter estimates for the best models for the relationship between island type and different life history traits for all families excluding geckos and anoles.

Clutch size

Non-phylogenetic

Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	0.15829	0.045011	3.517	0.000566
Female body mass	0.204622	0.016855	12.14	< 2e-16
Land-bridge	0.013835	0.050463	0.274	0.784306
Oceanic	-0.07066	0.037388	-1.89	0.060551
Latitude	0.007662	0.001577	4.86	2.74E-06

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	0.217448	0.038814	5.602	3.23E-08
Female body mass	0.222321	0.012071	18.418	< 2e-16
Land-bridge	0.041047	0.05464	0.751	0.45282
Mainland	0.106132	0.036183	2.933	0.00348
Oceanic	-0.09283	0.040727	-2.279	0.023
Latitude	0.003949	0.000856	4.612	4.88E-06

Phylogenetic

Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	0.143641	0.088998	1.614	0.108467
Female body mass	0.198817	0.025659	7.7483	9.42E-13
Latitude	0.006815	0.002071	3.2912	0.001223
Land-bridge	0.079395	0.059776	1.3282	0.185964
Oceanic	-0.0374	0.04698	-0.796	0.427176

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	0.122599	0.101727	1.2052	0.22848
Latitude	0.001795	0.000843	2.1297	0.03349
Female body mass	0.203925	0.014053	14.511	< 2.2e-16
Land-bridge	0.067271	0.037484	1.7947	0.07307

Mainland	0.146855	0.03036	4.8371	1.57E-06
Oceanic	0.016101	0.030982	0.5197	0.60342

Brood frequency
Non-phylogenetic
Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	0.574551	0.099016	5.803	3.53E-07
Female body mass	-0.1014	0.024019	-4.222	9.35E-05
Latitude	-0.0093	0.00225	-4.131	0.000126
Land-bridge	-0.00316	0.069356	-0.046	0.963866
Oceanic	-0.03831	0.06628	-0.578	0.565668

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	0.42506	0.087106	4.88	1.44E-06
Female body mass	-0.07745	0.012826	-6.038	3.09E-09
Latitude	-0.00465	0.000855	-5.436	8.62E-08
Land-bridge	-0.02016	0.093599	-0.215	0.83
Mainland	-0.05742	0.082091	-0.699	0.485
Oceanic	-0.03647	0.088771	-0.411	0.681

Phylogenetic
Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	0.458467	0.106346	4.3111	6.93E-05
Female body mass	-0.08395	0.028517	-2.9437	0.004773
Latitude	-0.00769	0.002274	-3.3795	0.001355
Land-bridge	0.008547	0.063807	0.1339	0.893942
Oceanic	-0.00582	0.061086	-0.0952	0.924513

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	0.264512	0.115662	2.2869	0.022627
Female body mass	-0.05889	0.017775	-3.3131	0.000991
Land-bridge	-0.00492	0.079186	-0.0621	0.950531
Mainland	0.051052	0.072597	0.7032	0.48225
Oceanic	0.029355	0.076848	0.382	0.702633
Latitude	-0.00453	0.001023	-4.4312	1.16E-05

Offspring mass
 Non-phylogenetic
 Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	-0.90158	0.06501	-13.869	<2e-16
Female body mass	0.69984	0.0243	28.798	<2e-16
Land-bridge	0.08898	0.08364	1.064	0.29
Oceanic	0.07556	0.06882	1.098	0.275

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	-0.89244	0.06352	-14.05	<2e-16
Female body mass	0.69002	0.0138	50.013	<2e-16
Land-bridge	0.08995	0.08535	1.054	0.292
Mainland	-0.04295	0.06331	-0.678	0.498
Oceanic	0.0798	0.06969	1.145	0.253

Phylogenetic
 Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	-0.75238	0.111512	-6.7471	1.48E-09
Female body mass	0.628058	0.036227	17.3369	< 2.2e-16
Land-bridge	-0.01514	0.086557	-0.1749	0.8615
Oceanic	0.030128	0.07675	0.3925	0.6956

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	-0.69851	0.12699	-5.5005	5.93E-08
Female body mass	0.621648	0.01999	31.0974	< 2.2e-16
Land-bridge	-0.02095	0.077714	-0.2695	0.78764
Mainland	-0.17215	0.067562	-2.548	0.01112
Oceanic	-0.00228	0.072296	-0.0316	0.97484

Productivity
 Non-phylogenetic
 Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	-0.35233	0.14647	-2.405	0.02142
Female body mass	0.8786	0.04182	21.007	< 2e-16
Land-bridge	0.14353	0.16473	0.871	0.38935

Oceanic	-0.05445	0.14831	-0.367	0.71565
Island age	0.11843	0.04243	2.791	0.00835

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	-0.28939	0.158605	-1.825	0.0687
Female body mass	0.848139	0.019635	43.196	<2e-16
Land-bridge	0.005362	0.182708	0.029	0.9766
Mainland	0.010502	0.157199	0.067	0.9468
Oceanic	-0.00438	0.168614	-0.026	0.9793

Phylogenetic

Between islands

	Estimate	Std. Error	t value	p value
(Intercept)	-0.41383	0.162239	-2.5508	0.015145
Female body mass	0.826428	0.053065	15.574	< 2.2e-16
Land-bridge	0.186658	0.134234	1.3905	0.172904
Oceanic	0.04513	0.126554	0.3566	0.723467
Island age	0.118844	0.03629	3.2748	0.002342

With mainland

	Estimate	Std. Error	t value	p value
(Intercept)	-0.45751	0.176729	-2.5888	0.009929
Female body mass	0.839477	0.026609	31.5481	< 2.2e-16
Land-bridge	0.080353	0.142736	0.5629	0.57374
Mainland	0.118341	0.127119	0.9309	0.352358
Oceanic	0.068549	0.137612	0.4981	0.618625

B. Parameter estimates for the best models for the relationship between island type and different life history traits for analyses of only geckos

Clutch size

	Estimate	Std. Error	t value	p value
(Intercept)	-0.06348	0.086409	-0.735	0.4656
Female body mass	0.115925	0.023871	4.856	9.71E-06
Land-bridge	0.101408	0.055613	1.823	0.0735
Oceanic	-0.00375	0.029794	-0.126	0.9003
Isolation	0.069807	0.030898	2.259	0.0277

Brood frequency

	Estimate	Std. Error	t value	p value
(Intercept)	1.04631	0.14422	7.255	4.45E-11
Female body mass	-0.05211	0.04658	-1.119	0.26548
Latitude	-0.0112	0.0025	-4.48	1.73E-05
Mainland	-0.35679	0.13188	-2.705	0.00782
Oceanic	-0.04973	0.16725	-0.297	0.76671

Offspring mass

	Estimate	Std. Error	t value	p value
(Intercept)	-0.96834	0.07599	-12.743	<2e-16
Female body mass	0.75713	0.02597	29.156	<2e-16
Land-bridge	-0.02416	0.18226	-0.133	0.895
Mainland	0.09946	0.07354	1.352	0.177
Oceanic	-0.09675	0.09189	-1.053	0.293

Productivity

	Estimate	Std. Error	t value	p value
(Intercept)	-4.98975	0.99869	-4.996	0.004118
Female body mass	0.52361	0.05831	8.98	0.000286
Latitude	0.10926	0.01983	5.51	0.002693
Oceanic	-0.24471	0.10673	-2.293	0.070391
Island area	0.72637	0.16684	4.354	0.007334

C. Parameter estimates for the best models for the relationship between island type and different life history traits after excluding the polytomies in the phylogenetic tree

Clutch size

	Estimate	Std. Error	t value	p value
(Intercept)	0.157805	0.103192	1.5292	0.12665
Female body mass	0.20121	0.014662	13.7235	< 2.2e-16
Land-bridge	0.084375	0.041289	2.0435	0.04137
Mainland	0.151869	0.032763	4.6354	4.24E-06
Oceanic	0.0246	0.03418	0.7197	0.47194

Brood frequency

	Estimate	Std. Error	t value	p value
(Intercept)	0.518304	0.127679	4.0594	5.66E-05
Female body mass	-0.06878	0.019819	-3.4703	0.000562
Latitude	-0.00623	0.001111	-5.6101	3.25E-08
Land-bridge	-0.17515	0.087307	-2.0061	0.045348
Mainland	-0.06613	0.07784	-0.8496	0.395931
Oceanic	-0.05479	0.080651	-0.6793	0.497226

Offspring mass

	Estimate	Std. Error	t value	p value
(Intercept)	-0.81268	0.104485	-7.778	3.15E-14
Female body mass	0.642721	0.01795	35.8068	< 2.2e-16
Land-bridge	0.10413	0.064841	1.6059	0.1088
Mainland	-0.05668	0.04904	-1.1557	0.2483
Oceanic	0.030441	0.051127	0.5954	0.5518

Productivity

	Estimate	Std. Error	t value	p value
(Intercept)	0.118672	0.216658	0.5477	0.58677
Female body mass	0.694034	0.050889	13.6383	< 2e-16
Latitude	-0.01451	0.005455	-2.6598	0.01102
Land-bridge	0.352233	0.149257	2.3599	0.023
Oceanic	0.105519	0.106022	0.9953	0.32531
Island age	0.088492	0.038575	2.2941	0.02686

D. Parameter estimates for the best models for the relationship between island type and different life history traits

Intercept: for continental islands. Type is the difference in intercept of oceanic islands, land bridge islands and the mainland, respectively, from the intercept for continental islands

Clutch size

Non-phylogenetic model

With island type ($R^2 = 0.46$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.16121	0.07947	-2.028	0.04343
Female body mass	0.22978	0.01573	14.611	< 2e-16
Latitude	0.00644	0.00139	4.633	5.48E-06
Land-bridge	0.13928	0.05007	2.782	0.00576
Oceanic	-0.02704	0.02944	-0.918	0.35926
Isolation	0.08028	0.02547	3.153	0.00179

With mainland ($R^2 = 0.45$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.075697	0.031126	2.432	0.0152
Female body mass	0.2601	0.01193	21.801	< 2e-16
Latitude	0.004494	0.000823	5.463	6.19E-08
Land-bridge	0.052759	0.04622	1.141	0.254
Mainland	0.14544	0.029096	4.999	7.05E-07
Oceanic	-0.0604	0.032465	-1.86	0.0632

Without island type ($R^2 = 0.43$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.06183	0.06472	-0.955	0.34
Female body mass	0.225381	0.015844	14.225	< 2e-16
Latitude	0.007492	0.001389	5.392	1.44E-07
Isolation	0.036132	0.021242	1.701	0.09

Phylogenetic model

With island type ($R^2 = 0.20$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.1221082	0.1057593	1.1546	0.249217
Female body mass	0.1592485	0.0204361	7.7925	1.20E-13
Latitude	0.0031551	0.0018482	1.7071	0.08888
Land-bridge	0.1536783	0.0574011	2.6773	0.007848
Oceanic	0.0213218	0.032775	0.6506	0.515854
Island age	0.0367809	0.0163736	2.2464	0.025439

With mainland ($R^2 = 0.24$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.124312	0.101898	1.22	0.22282
Latitude	0.001761	0.000841	2.0942	0.03654
Female body mass	0.204514	0.014012	14.5958	< 2.2e-16
Land-bridge	0.065899	0.037155	1.7736	0.0765
Mainland	0.143703	0.029979	4.7934	1.94E-06
Oceanic	0.015496	0.030353	0.5105	0.60981

Without island type ($R^2 = 0.17$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.231963	0.101299	2.2899	0.02274
Female body mass	0.159543	0.020745	7.6906	2.27E-13
Island age	-0.00058	0.011082	-0.0525	9.58E-01

Brood frequency

Non-phylogenetic model

With island type ($R^2 = 0.50$)

	Estimate	Std. Error	t value	p value
(Intercept)	1.177011	0.134218	8.769	5.62E-13
Female body mass	-0.2402	0.035362	-6.793	2.66E-09
Latitude	-0.01864	0.003677	-5.068	3.00E-06
Land-bridge	-0.2175	0.108051	-2.013	0.0479
Oceanic	-0.06551	0.095145	-0.689	0.4933

With mainland ($R^2 = 0.22$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.865104	0.083967	10.303	< 2e-16
Female body mass	-0.14919	0.014857	-10.042	< 2e-16
Latitude	-0.00901	0.000977	-9.225	< 2e-16
Land-bridge	-0.30148	0.097368	-3.096	0.00205
Mainland	-0.25892	0.079459	-3.259	0.00118
Oceanic	-0.10947	0.088052	-1.243	0.21425

Without island type ($R^2 = 0.44$)

	Estimate	Std. Error	t value	p value
(Intercept)	1.158474	0.121313	9.549	1.54E-14
Female body mass	-0.24406	0.035354	-6.903	1.49E-09
Latitude	-0.0216	0.003497	-6.177	3.22E-08

Phylogenetic model

With island type ($R^2 = 0.28$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.805899	0.175555	4.5906	1.86E-05
Female body mass	-0.1281	0.044063	-2.9071	0.004864
Latitude	-0.01183	0.003668	-3.2254	0.001903
Land-bridge	-0.038	0.090578	-0.4195	0.676128
Oceanic	-0.10202	0.074242	-1.3742	0.173708
Island age	0.065421	0.027783	2.3547	0.021307

With mainland ($R^2 = 0.07$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.556434	0.121925	4.5638	6.08E-06
Female body mass	-0.07099	0.019041	-3.7282	0.000211
Land-bridge	-0.17929	0.076189	-2.3532	0.01893
Mainland	-0.09042	0.067783	-1.334	0.182712
Oceanic	-0.09338	0.071901	-1.2987	0.194533
Latitude	-0.00653	0.001047	-6.2327	8.55E-10

Without island type ($R^2 = 0.26$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.751542	0.166867	4.5038	2.48E-05
Female body mass	-0.13744	0.043706	-3.1447	0.002405
Latitude	-0.01208	0.003664	-3.2959	0.001517
Island age	0.056927	0.022216	2.5624	0.012456

Offspring body mass

Non-phylogenetic model

With island type ($R^2 = 0.90$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.945	0.03869	-24.427	<2e-16
Female body mass	0.71501	0.01838	38.904	<2e-16
Land-bridge	0.13268	0.05805	2.286	0.0235
Oceanic	0.05004	0.04013	1.247	0.2142

With mainland ($R^2 = 0.82$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.92316	0.039945	-23.111	<2e-16
Female body mass	0.691724	0.012051	57.4	<2e-16
Land-bridge	0.132979	0.064217	2.071	0.0387
mainland	-0.00109	0.039653	-0.028	0.978
Oceanic	0.051097	0.044391	1.151	0.2501

Without island type ($R^2 = 0.90$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.90732	0.041065	-22.095	<2e-16
Female body mass	0.715882	0.018704	38.275	<2e-16
Latitude	0.00057	0.00168	0.339	0.735

Phylogenetic model

With island type ($R^2 = 0.84$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.89758	0.05891	-15.2364	<2e-16
Female body mass	0.685867	0.023551	29.1222	<2e-16

Land-bridge	0.070122	0.060954	1.1504	0.2516
Oceanic	0.03676	0.043427	0.8465	0.3985

With mainland ($R^2 = 0.67$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.80887	0.097988	-8.2548	8.88E-16
Female body mass	0.648152	0.017195	37.6953	< 2.2e-16
Land-bridge	0.066473	0.058996	1.1267	0.2602
Mainland	-0.0692	0.044624	-1.5508	0.1214
Oceanic	0.018565	0.046933	0.3956	0.6926

Without island type ($R^2 = 0.82$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.85557	0.053781	-15.908	< 2.2e-16
Female body mass	0.677194	0.024545	27.59	< 2.2e-16

Productivity

Non-phylogenetic model

With island type ($R^2 = 0.93$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.186739	0.158827	1.176	0.24516
Female body mass	0.739313	0.0393	18.812	< 2e-16
Latitude	-0.01419	0.004435	-3.2	0.00236
Land-bridge	0.218284	0.14509	1.504	0.13863
Oceanic	0.032689	0.101376	0.322	0.74843
Island age	0.094376	0.03904	2.417	0.01924

With mainland ($R^2 = 0.78$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.14627	0.10922	-1.339	0.181
Female body mass	0.83198	0.01819	45.748	<2e-16
Land-bridge	-0.11981	0.14531	-0.824	0.41
Mainland	-0.10143	0.10865	-0.934	0.351
Oceanic	-0.04532	0.12071	-0.375	0.707

Without island type ($R^2 = 0.90$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.13875	0.06387	-2.172	0.0342
Female body mass	0.78978	0.03531	22.369	<2e-16

Phylogenetic model

With island type ($R^2 = 0.88$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.065312	0.175027	0.3732	0.71058
Female body mass	0.717245	0.044154	16.2442	< 2.2e-16
Latitude	-0.01056	0.004495	-2.35	0.022682
Land-bridge	0.170237	0.118945	1.4312	0.158465
Oceanic	0.00695	0.091994	0.0755	0.940077
Island age	0.11203	0.033921	3.3027	0.001755

With mainland ($R^2 = 0.66$)

	Estimate	Std. Error	t value	p value
(Intercept)	-0.021	0.015	-1.4356	0.15165
Female body mass	0.082	0.0025	32.6529	< 2e-16
Latitude	-0.0003	0.00014	-2.1406	0.03271
Land-bridge	-0.0031	0.012	-0.2555	0.79844
Mainland	0.0000025	0.0098	0.0003	0.99979
Oceanic	-0.0026	0.018	-0.2465	0.80541

Without island type ($R^2 = 0.86$)

	Estimate	Std. Error	t value	p value
(Intercept)	0.22453	0.162363	1.3829	0.17239
Female body mass	0.704028	0.045844	15.3572	< 2e-16
Latitude	-0.01316	0.004436	-2.9657	0.00449