

ITALIAN FRESHWATER GASTROTRICHA

by Lara Pierboni, Loretta Guidi, Maria Balsamo

(October, 2005)

Order	Chaetonotida	Remane, 1925 [Rao & Clausen, 1970]
Suborder	Paucitubulatina	d'Hondt, 1971
Family	Chaetonotidae	Zelinka, 1889
Subfamily	Chaetonotinae	Zelinka, 1889 sensu Kisielowski, 1991

Genus *Aspidiophorus* Voigt, 1904

<i>Aspidiophorus heterodermus</i>	Saito, 1937
<i>Aspidiophorus oculifer</i>	Kisielowski, 1981
<i>Aspidiophorus ophiodermus</i>	Balsamo, 1982
<i>Aspidiophorus paradoxus</i>	(Voigt, 1902)
<i>Aspidiophorus squamulosus</i>	Roszczak, 1936

Genus *Chaetonotus* Ehrenberg, 1830

<i>Chaetonotus acanthocephalus</i>	Valkanov, 1937
<i>Chaetonotus acanthodes</i>	Stokes, 1887
<i>Chaetonotus acanthophorus</i>	Stokes, 1887
<i>Chaetonotus aemilianus</i>	Balsamo, 1978
<i>Chaetonotus arethusae</i>	Balsamo & Todaro, 1995
<i>Chaetonotus balsamoe</i>	Kisielowski, 1997
<i>Chaetonotus bisacer</i>	Greuter, 1917
<i>Chaetonotus benacensis</i>	Balsamo & Fregni, 1995
<i>Chaetonotus brachyurus</i>	Balsamo, 1981
<i>Chaetonotus brevispinosus</i>	Zelinka, 1889
<i>Chaetonotus cestacanthus</i>	Balsamo, 1990
<i>Chaetonotus chuni</i>	Voigt, 1901
<i>Chaetonotus daphnes</i>	Balsamo & Todaro, 1995
<i>Chaetonotus decemsetosus</i>	Marcolongo, 1910
<i>Chaetonotus disjunctus</i>	Greuter, 1917
<i>Chaetonotus dracunculus</i>	Balsamo, 1990
<i>Chaetonotus enormis</i>	Stokes, 1887
<i>Chaetonotus fluviatilis</i>	Balsamo & Kisielowski, 1986
<i>Chaetonotus heideri</i>	Brehm, 1917
<i>Chaetonotus heterospinosus</i>	Balsamo, 1978
<i>Chaetonotus hirsutus</i>	Marcolongo, 1910
<i>Chaetonotus hystrix</i>	Metschnikoff, 1865
<i>Chaetonotus italicus</i>	Balsamo & Todaro, 1995
<i>Chaetonotus laroides</i>	Marcolongo, 1910
<i>Chaetonotus larus</i>	(Müller, 1773)
<i>Chaetonotus longispinosus</i>	Stokes, 1887
<i>Chaetonotus lunatospinosus</i>	Balsamo, 1981
<i>Chaetonotus macrochaetus</i>	Zelinka, 1889
<i>Chaetonotus maximus</i>	Ehrenberg, 1830
<i>Chaetonotus microchaetus</i>	Preobrajenskaia, 1926
<i>Chaetonotus minimus</i>	Marcolongo, 1910
<i>Chaetonotus multispinosus</i>	Grünspan, 1908
<i>Chaetonotus mutinensis</i>	Balsamo, 1978

<i>Chaetonotus naiadis</i>	Balsamo & Todaro, 1995
<i>Chaetonotus octonarius</i>	Stokes, 1887
<i>Chaetonotus oculifer</i>	Kisielewski, 1981
<i>Chaetonotus ophiogaster</i>	Remane, 1927
<i>Chaetonotus oplites</i>	Balsamo & Fregni, 1995
<i>Chaetonotus paucisetosus</i>	Marcolongo, 1910
<i>Chaetonotus pentacanthus</i>	Balsamo, 1981
<i>Chaetonotus persetosus</i>	Zelinka, 1889
<i>Chaetonotus polyspinosus</i>	Greuter, 1917
<i>Chaetonotus poznaniensis</i>	Kisielewski, 1981
<i>Chaetonotus pungens</i>	Balsamo, 1990
<i>Chaetonotus robustus</i>	Davison, 1938
<i>Chaetonotus schultzei</i>	Metschnikoff, 1865
<i>Chaetonotus similis</i>	Zelinka, 1889
<i>Chaetonotus simrothi</i>	Voigt, 1909
<i>Chaetonotus sphagnophilus</i>	Kisielewski, 1981
<i>Chaetonotus spinulosus</i>	Stokes, 1887
<i>Chaetonotus trispinosus</i>	Balsamo, 1990
<i>Chaetonotus uncinus</i>	Voigt, 1902
Genus <i>Fluxiderma</i>	d'Hondt, 1974
<i>Fluxiderma concinnum</i>	(Stokes, 1887)
Genus <i>Heterolepidoderma</i>	Remane, 1927
<i>Heterolepidoderma brevitubulatum</i>	Kisielewski, 1981
<i>Heterolepidoderma gracile</i>	Remane, 1927
<i>Heterolepidoderma lamellatum</i>	Balsamo & Fregni, 1995
<i>Heterolepidoderma macrops</i>	Kisielewski, 1981
<i>Heterolepidoderma majus</i>	Remane, 1927
<i>Heterolepidoderma multiseriatum</i>	Balsamo, 1978
<i>Heterolepidoderma ocellatum</i>	(Metschnikoff, 1865)
<i>Heterolepidoderma pineisquamatum</i>	Balsamo, 1981
Genus <i>Ichthydium</i>	Ehrenberg, 1830
<i>Ichthydium diacanthum</i>	Balsamo & Todaro, 1995
<i>Ichthydium forcipatum</i>	Voigt, 1901
<i>Ichthydium forcifcula</i>	Remane, 1927
<i>Ichthydium fossae</i>	d'Hondt, 1971
<i>Ichthydium plicatum</i>	Balsamo & Fregni, 1995
<i>Ichthydium podura</i>	(Müller, 1773)
<i>Ichthydium squamigerum</i>	Balsamo & Fregni, 1995
<i>Ichthydium tanytrichum</i>	Balsamo, 1983
Genus <i>Lepidochaetus</i>	Kisielewski, 1991
<i>Lepidochaetus zelinkai</i>	(Grünspan, 1908)

Genus <i>Lepidodermella</i>	Blake, 1933
<i>Lepidodermella punctata</i>	(Greuter, 1917)
<i>Lepidodermella squamata</i>	(Dujardin, 1841)
Genus <i>Polymerurus</i>	Remane, 1927
<i>Polymerurus nodicaudus</i>	(Voigt, 1901)
<i>Polymerurus nodifurca</i>	(Marcolongo, 1910)
<i>Polymerurus rhomboides</i>	(Stokes, 1887)
<i>Polymerurus serraticaudus</i>	(Voigt, 1901)
Genus <i>Anacanthoderma</i>	Marcolongo, 1910
<i>Anacanthoderma paucisetosum</i>	Marcolongo, 1910
<i>Anacanthoderma punctatum</i>	Marcolongo, 1910
Genus <i>Chitonodytes</i>	Remane, 1967
<i>Chitonodytes longisetosus</i>	(Metschnikoff, 1865)
Genus <i>Dasydytes</i>	Gosse, 1851
<i>Dasydytes ornatus</i>	Voigt, 1909
Genus <i>Haltidytes</i>	Remane, 1936
<i>Haltidytes crassus</i>	(Greuter, 1917)
<i>Haltidytes saltitans</i>	(Stokes, 1887)
Genus <i>Setopus</i>	Grünspan, 1908
<i>Setopus tongiorgii</i>	(Balsamo, 1983)
Genus <i>Stylochaeta</i>	Hlava, 1904
<i>Stylochaeta fusiformis</i>	(Spencer, 1890)
<i>Stylochaeta stylifera</i>	(Voigt, 1901)
Genus <i>Dichaetura</i>	Lauterborn, 1913
<i>Dichaetura capricornia</i>	(Metschnikoff, 1865)
Genus <i>Neogossea</i>	Remane, 1927

Neogosseia antennigera

(Gosse, 1851)