

Cotylorhiza tuberculata (Macri, 1778)

Family Mastigiidae

Phyllorhiza punctata von Lendenfeld, 1884

Family Catostylidae

Catostylus tagi (Haeckel, 1869)

Family Rhizostomatidae

Rhizostoma pulmo (Macri, 1778)
Rhopilema rhopalophorum Haeckel, 1880

Order Stauromedusae

Family Depastridae

Craterolophus convolvulus (Johnson, 1835)

Depastrum cyathiforme (M. Sars, 1846)

Manania auricula (Fabricius, 1780)
= *Halimocyathus lagena hexaradiata* (Broch, 1907)
uchidai (Naumov, 1961)

Family Kishinouyeidae

Lucernariopsis campanulata (Lamouroux, 1815)
cruxmelitensis Corbin, 1978

Family Lipkeidae

Lipkea ruspoliana Vogt, 1886
sturzi (Antipa, 1893)

Family Lucernariidae

Haliclystus auricula (Rathke, 1806)
octoradiatus (Lamarck, 1816)
salpinx Clark, 1863
Lucernaria bathyphila Haeckel, 1880
haeckeli (Antipa, 1892)
infundibulum Haeckel, 1880
quadricornis O.F. Müller, 1776
sainthilarei (Radicorzew,)
walteri (Antipa, 1892)

Stylocoronella riedli Salvini-Plawen, 1966
variabilis Salvini-Plawen, 1987

Class incertae sedis

Family Tetraplatidae

Tetraplatia volitans Busch, 1851

Hydrozoa

Hydroida

Compiled by Jacob van der Land, W. Vervoort, S.D. Cairns and P. Schuchert

The European list was derived from URMO which was edited by W. Vervoort. S.D. Cairns provided the list of hydrocorals and P. Schuchert provided additions the list of hydroids.

Class Hydrozoa

Order Actinulida

Family Halammohydridae

Halammohydra adherens Swedmark & Teissier, 1958 A
coronata Clausen, 1967 A
intermedia Clausen, 1967 A
neglecta Clausen A
octopodides Remane, 1927 A M
schulzei Remane, 1927 A M
vermiformis Swedmark & Teissier, 1857 A

Family Otohydridae

Otohydra tremulans Lacassagne, 1973 A M
vagans Swedmark & Teissier, 1958 A M

Order Capitata

Family Acaulidae

Acaulis primarius Stimpson, 1854 A
Acauloidea ammisatum Bouillon, 1965 A
ilonae (Brinckmann-Voss, 1966) A M

Family Boreohydridae

Boreohydra simplex Westblad, 1937
Meiorhopalon arenicolum Von Salvini-Plawen, 1984 A M
Psammohydra nana Schulz, 1950 A M

Family Candelabridae

Candelabrum cocksii (Vigurs, 1849) A
minutum (Bonnievie, 1898) A
mitra (Bonnievie, 1898) A
phrygium (Fabricius, 1780) A
serpentarii Segonzac & Vervoort, 1995
verrucosum (Bonnievie, 1898) A
Monocoryne gigantea (Bonnievie, 1898) A

Family Cladocorynidae

Cladocoryne floccosa Rotch, 1871 A

Family Cladonematidae

Cladonema radiatum Dujardin, 1843 A

Family Corymorphidae

Amalthaea sarsi (Steenstrup, 1854) A
Branchiocerianthus italicus (Stechow, 1921) M
norvegicus Brattström, 1957 A
Corymorpha abyssalis Broch, 1909 A
annulicornis Sars, 1859 A
appeloeffi Bonnievie, 1901 A
glacialis Sars, 1859 A
groenlandica (Allman, 1876) A
nana Alder, 1857 A
nutans Sars, 1835 A M

<i>Eucodonium</i>			
<i>brownei</i> Hartlaub, 1907	A M		
<i>Eugotoea</i>			
<i>petalina</i> Margulis, 1989	M		
<i>Euphysora</i>			
<i>annulata</i> Kramp, 1928	M		
= <i>Corymorpha annulata</i>			
<i>bigelowi</i> Maas, 1905	A		
= <i>Corymorpha bigelowi</i>			
<i>Paragotoea</i>			
<i>bathybia</i> Kramp, 1942	A M		
<i>Plotocnide</i>			
<i>borealis</i> Wagner, 1885	A M		
<i>Vannuccia</i>			
<i>forbesi</i> (Mayer, 1894)	A		
Family Corynidae			
<i>Actigia</i>			
<i>vanbenedeni</i> (Hincks, 1869)	A M		
<i>Coryne</i>			
<i>brevicornis</i> Bonnevie, 1898	A		
<i>caespes</i> Allman, 1871	M		
<i>clavata</i> Graeffe, 1883	M		
<i>epizoica</i> Stechow, 1921	M		
<i>filiformis</i> Rees, 1936	A		
<i>fucicola</i> (de Filippi, 1866)	M		
<i>graeffei</i> Jickeli, 1883	M		
<i>hincksi</i> Bonnevie, 1898	A		
<i>muscoides</i> (Linnaeus, 1761)	A M		
<i>pintneri</i> Schneider, 1897	A M		
<i>pusilla</i> (Gaertner, 1774)	A M		
<i>Dicodonium</i>			
<i>adriaticum</i> Graeffe, 1884	M		
<i>ocellatum</i> (Busch, 1851)	M		
<i>Dipurena</i>			
<i>dolichogaster</i> (Haeckel, 1864)	M		
<i>halterata</i> (Forbes, 1846)	A		
<i>ophiogaster</i> (Haeckel, 1877)	A		
<i>reesi</i> Vannucci, 1956	A M		
<i>simulans</i> Bouillon, 1965	A		
<i>spongicola</i> Anger, 1972	A		
<i>Sarsia</i>			
<i>barentsi</i> Linko, 1905	A		
<i>eximia</i> (Allman, 1859)	A		
<i>gemmifera</i> Forbes, 1848	A M		
<i>lovenii</i> (M. Sars, 1846)	A		
<i>occulta</i> Edwards, 1978	A		
<i>pattersoni</i> Haddon, 1886	A		
<i>piriformis</i> Edwards, 1983	A		
<i>princeps</i> (Haeckel, 1879)	A		
<i>producta</i> (Wright, 1858)	A M		
<i>prolifera</i> Forbes, 1848	A M		
<i>striata</i> Edwards, 1983	A		
<i>tubulosa</i> (M. Sars, 1835)	A M		
<i>Sarsiella</i>			
<i>dinema</i> Hartlaub, 1907	A M		
Family Eleutheriidae			
<i>Eleutheria</i>			
<i>claparedei</i> Hartlaub, 1889	A M		
<i>dichotoma</i> De Quatrefages, 1842	A M		
<i>Staurocladia</i>			
<i>portmanni</i> Brinckmann, 1964	M		
Family Euphysidae			
<i>Euphysa</i>			
<i>aurata</i> Forbes, 1848	A M		
<i>flammea</i> (Linko, 1905)	A M		
<i>obvoluta</i> (Kramp, 1933)	A		
<i>tentaculata</i> Linko, 1905	A		
<i>Gymnogonos</i>			
<i>crassicornis</i> Bonnevie, 1898	A		
<i>Siphonohydra</i>			
<i>adriatica</i> Salvini-Plawen, 1966	A M		
Family Halocorynidae			
<i>Halocoryne</i>			
<i>epizoica</i> Hadzi, 1917	M		
Family Margelopsidae			
<i>Margelopsis</i>			
<i>haeckeli</i> Hartlaub, 1897	A		
<i>hartlaubi</i> Browne, 1903	A		
Family Moerisiidae			
<i>Moerisia</i>			
<i>lyonsi</i> Boulenger, 1908	M		
<i>pallasi</i> (Derzhavin, 1912)			
<i>Odessia</i>			
<i>maeotica</i> (Ostroumoff, 1896)	M		
<i>Ostroumovia</i>			
<i>inkermanica</i> (Paltschikowa-Ostroumova, 1925)	M		
= <i>Moerisia inkermanica</i>			
Family Paracorynidae			
<i>Paracoryne</i>			
<i>huvei</i> Picard, 1957	M		
Family Pennariidae			
<i>Pennaria</i>			
<i>disticha</i> (Goldfuss, 1820)	A		
Family Porpitiidae			
<i>Porpita</i>			
<i>porpita</i> (Linnaeus, 1758)	A		
<i>Veleva</i>			
<i>veleva</i> (Linnaeus, 1758)	A		
Family Protohydryidae			
<i>Protohydra</i>			
<i>leuckarti</i> Greeff, 1870	A M		
Family Rosalindidae			
<i>Rosalinda</i>			
<i>incrustans</i> (Kramp, 1947)	A M		
Family Sphaerocorynidae			
<i>Sphaerocoryne</i>			
<i>bedoti</i> Pictet, 1893	A		
Family Tubulariidae			
<i>Ectopleura</i>			
<i>dumortieri</i> (Van Beneden, 1844)	A		
<i>minerva</i> Mayer, 1900	A		
<i>petalina</i> (Margulis, 1989)	M		
= <i>Eogothoea petalina</i>			
<i>sacculifera</i> Kramp, 1957	A		
<i>wrightii</i> Petersen, 1979	M		
<i>Hybocodon</i>			
<i>prolifer</i> Agassiz, 1862	A M		
<i>Rhabdoon</i>			
<i>singularis</i> Keferstein & Ehlers, 1861	M		
<i>Tubularia</i>			
<i>bellis</i> Allman, 1865	A		
<i>ceratogyne</i> Pérez, 1920	A		
<i>cornucopia</i> Bonnevie, 1898	A		
<i>crocea</i> (Agassiz, 1862)	A		
= <i>Ectopleura crocea</i>			
<i>indivisa</i> Linnaeus, 1758	A M		
<i>larynx</i> Ellis & Solander, 1786	A M		
<i>mesembryanthemum</i> Allman, 1871			
<i>regalis</i> Boeck, 1860	A		
Family Zancleidae			
<i>Zanclea</i>			
<i>costata</i> Gegenbaur, 1857	A M		
<i>sessilis</i> (Gosse, 1853)	A M		
Order Conica			
Family Aequoreidae			
<i>Aequorea</i>			
<i>aequorea</i> (Forskål, 1775)	M		
<i>conica</i> Browne, 1905	M		
<i>forskalea</i> (Forskål, 1775)	A		
<i>macroactyla</i> (Brandt, 1835)	A		
<i>pensilis</i> (Eschscholtz, 1829)	A M		
<i>vitrina</i> Gosse, 1853	A		
Family Aglaopheniidae			
<i>Aglaophenia</i>			
<i>acacia</i> Allman, 1883	A M		

<i>elongata</i> Meneghini, 1845	A M	<i>huxleyi</i> (Haeckel, 1879)	A
<i>harpago</i> Schenck, 1965	M	<i>Dipleurosoma</i>	
<i>kirchenpaueri</i> (Heller, 1868)	A M	<i>typicum</i> Boeck, 1866	A
<i>latecarinata</i> Allman, 1877	A	Family Eirenidae	
<i>lophocarpa</i> Allman, 1877	A M	<i>Campanopsis</i>	
<i>octodonta</i> Heller, 1868	A M	<i>adriaticus</i> (Hadzi, 1915)	M
<i>parvula</i> Bale, 1882	A	= <i>Halanthus adriaticus</i>	
<i>picardi</i> Svoboda, 1979	A M	<i>Eirene</i>	
<i>pluma</i> (Linnaeus, 1767)	A M	<i>viridula</i> (Péron & Lesueur, 1809)	A M
<i>tubiformis</i> Marktanner-Turneretscher, 1890	A M	<i>Eugymnanthea</i>	
<i>tubulifera</i> (Hincks, 1861)	A M	<i>inquilina</i> Palombi, 1935	M
Cladocarpus		<i>Eutima</i>	
<i>bicuspis</i> (Sars, 1873)	A	<i>gegenbauri</i> (Haeckel, 1864)	A
<i>bonneviaei</i> Jäderholm, 1909	A	<i>gracilis</i> (Forbes & Goodsir, 1853)	A M
<i>boucheti</i> Ramil & Vervoort, 1992	A	<i>mira</i> McCrady, 1859	A
<i>cartieri</i> Bedot, 1921	A	<i>Eutonina</i>	
<i>cornutus</i> (Verrill, 1879)	A	<i>indicans</i> (Romanes, 1876)	A
<i>diana</i> Broch, 1918	A	<i>scintillans</i> (Bigelow, 1909)	A M
<i>formosus</i> Allman, 1877	A	<i>Helgicirrha</i>	
<i>integer</i> (Sars, 1873)	A	<i>cari</i> (Haeckel, 1864)	A M
<i>multiseptatus</i> (Bale, 1915)	A	<i>schulzei</i> Hartlaub, 1909	A M
<i>paraventricosus</i> Ramil & Vervoort, 1992	A	<i>Irenium</i>	
<i>sigma</i> (Allman, 1877)	A	<i>quadrigatum</i> Haeckel, 1879	A
<i>sinuosus</i> Vervoort, 1966		<i>Neotima</i>	
Gymnangium		<i>lucullana</i> Petersen	M
<i>montagui</i> (Billard, 1912)	A M	<i>Phialopsis</i>	
<i>sinosum</i> (Fraser, 1925)	A	<i>diegensis</i> Torrey, 1909	A
Lytocarpia		<i>Tima</i>	
<i>distans</i> (Allman, 1877)	A M	<i>bairdii</i> (Johnston, 1833)	A
<i>myriophyllum</i> (Linnaeus, 1758)	A M	<i>flavilabris</i> Eschscholtz, 1829	A M
<i>phyteuma</i> (Kirchenpauer, 1876)	A M	<i>lucullana</i> (Delle Chiaje, 1822)	M
		= <i>Neotima lucullana</i>	
Macrorhynchia		Family Eucheilotidae	
<i>philippina</i> Kirchenpauer, 1872	A	<i>Eucheilota</i>	
Nematocarpus		<i>flevensis</i> Van Kampen, 1922	A
<i>ramuliferus</i> (Allman, 1874)	A	<i>maasi</i> Neppi & Stiasny, 1911	M
Streptocaulus		<i>maculata</i> Hartlaub, 1894	A
<i>corneliusi</i> (Ramil & Vervoort, 1992)	A M	<i>paradoxa</i> Mayer, 1900	M
<i>dollfusi</i> (Billard, 1924)	A M	<i>ventricularis</i> McCrady, 1859	A M
= <i>Cladocarpus dollfusi</i>		Family Haleciidae	
<i>pectiniferus</i> (Allman, 1883)	A	<i>Campalecium</i>	
<i>sinuosus</i> (Vervoort, 1966)	A	<i>cirratum</i> (Haeckel, 1879)	M
Family Barcinididae		<i>medusiferum</i> Torrey, 1902	M
Barcino		<i>microtheca</i> (Hadzi, 1914)	M
<i>foixensis</i> Gili, Bouillon, Pagès, Palanques & Puig, 1999	M	= <i>Haleciella microtheca</i>	
Family Blackfordiidae		<i>Halecium</i>	
Blackfordia		<i>articulosum</i> Clark, 1875	A
<i>virginica</i> Mayer, 1910	A M	<i>banyulense</i> Motz-Kossowska, 1912	M
Family Campanulinidae		<i>beanei</i> (Johnston, 1838)	A M
Calycella		<i>birulai</i> Spassky, 1929	A
<i>gracilis</i> Hartlaub, 1897	A	<i>conicum</i> Stechow, 1919	M
<i>syringa</i> (Linnaeus, 1758)	A M	<i>corrugatum</i> Nutting, 1899	A
Campomma		<i>curvicaule</i> Lorenz, 1886	A
<i>hincksii</i> (Hartlaub, 1897)	A M	<i>delicatum</i> Coughtrey, 1876	A
Cuspidella		<i>groenlandicum</i> Kramp, 1911	A
<i>costata</i> Hincks, 1868	A M	<i>halecium</i> (Linnaeus, 1758)	A M
<i>grandis</i> Hincks, 1868		<i>labrosum</i> Alder, 1859	A M
<i>humilis</i> (Alder, 1862)	A M	<i>lankesteri</i> (Bourne, 1890)	A M
<i>mertensi</i> (Brandt, 1835)	M	= <i>H. robustum</i> Pieper, 1884	
<i>procumbens</i> Kramp, 1911	A	<i>liouvillei</i> Billard, 1934	A
Egmundella		<i>mediterraneum</i> Weismann, 1883	A M
<i>amirantensis</i> Millard & Bouillon, 1973	A M	<i>minimum</i> Schneider, 1898	M
<i>grimaldii</i> Leloup, 1940	A	<i>minutum</i> Broch, 1903	A
Lafoeina		<i>muricatum</i> (Ellis & Solander, 1786)	A M
<i>maxima</i> (Levinsen, 1893)	A	<i>nanum</i> Alder, 1859	A M
<i>tenuis</i> Sars, 1874	A M	<i>ornatum</i> Nutting, 1901	A
Opercularella		<i>petrosom</i> Stechow, 1919	A M
<i>lacerata</i> (Johnston, 1847)	A M	<i>profundum</i> Calder & Vervoort, 1998	A
<i>panicula</i> (Sars, 1873)	A M	<i>pusillum</i> Sars, 1856	A M
<i>pumila</i> Clark, 1875	A	<i>reflexum</i> Stechow, 1919	M
Tetrapoma		<i>sessile</i> Norman, 1866	A M
<i>quadridentatum</i> (Hincks, 1874)	A	<i>tenellum</i> Hincks, 1861	A M
Family Dipleurosomatidae		<i>undulatum</i> Billard, 1921	A
Cuvieria		<i>Hydranthea</i>	

<i>aloyisii</i> (Zoja, 1893)	M	<i>fruticosa</i> Sars, 1851	A M
<i>diaphana</i> (Hadzi, 1912)	M	= <i>L. dumosa</i>	
<i>margarica</i> (Hincks, 1863)	A M	<i>grandis</i> Hincks, 1874	A M
<i>Hydrodendron</i>		<i>poecilum</i> Hincks, 1868	
<i>gorgonoide</i> (Sars, 1873)	A	<i>Scandia</i>	
<i>mirabile</i> (Hincks, 1866)	A	<i>gigas</i> (Pieper, 1884)	A M
= <i>H. caciniforme</i>		<i>michaelsarsi</i> (Leloup, 1935)	A M
= <i>Ophiodissa mirabile</i>		<i>Zygophylax</i>	
Family Halopterididae		<i>bathyphila</i> Leloup, 1940	A
<i>Antennella</i>		<i>biarmata</i> Billard, 1905	A M
<i>secundaria</i> (Gmelin, 1791)	A M	<i>brownei</i> Billard, 1924	A M
= <i>A. paucinoda</i>		<i>echinata</i> Calder & Vervoort, 1998	
<i>siliquosa</i> (Hincks, 1877)	A M	<i>elegantula</i> Leloup, 1940	A
<i>Halopteris</i>		<i>leloupi</i> Ramil & Vervoort, 1992	A
<i>alternata</i> (Nutting, 1900)	A	<i>levinensi</i> (Saemundsson, 1911)	A
<i>catharina</i> (Johnston, 1833)	A M	= <i>Z. elegantula</i>	
<i>diaphana</i> (Heller, 1868)	A	<i>pinnata</i> (Sars, 1873)	A
<i>glutinosa</i> (Lamouroux, 1816)	A	<i>sibogae</i> Billard, 1918	A
<i>liechtensternii</i> (Marktanner-Turneretscher, 1890)	A M	Family Laodiceidae	
<i>Monostaechas</i>		<i>Krampella</i>	
<i>quadridens</i> (McCrary, 1859)	A	<i>dubia</i> Russell, 1957	A M
<i>Schizotricha</i>		<i>tardenti</i> Gili, Bouillon & Pagès, 1998	M
<i>frutescens</i> (Ellis & Solander, 1786)	A M	<i>Laodicea</i>	
<i>polaris</i> Naumov, 1960	A	<i>bigelowi</i> Neppi & Stiasny, 1911	M
<i>variabilis</i> Bonnevie, 1899	A	<i>chapmani</i> Günther, 1903	A
Family Kirchenpaueriidae		<i>neptuna</i> Mayer, 1900	M
<i>Kirchenpaueria</i>		<i>ocellata</i> Babnik, 1948	M
<i>echinulata</i> (Lamarck, 1816)	A M	<i>undulata</i> (Forbes & Goodsir, 1853)	A M
<i>pinnata</i> (Linnaeus, 1758)	A M	<i>Melicertissa</i>	
= <i>K. similis</i>		<i>adriatica</i> Neppi, 1915	M
<i>triangulata</i> (Totton, 1930)		<i>Ptychogena</i>	
= <i>K. bonnevieae</i>		<i>crocea</i> Kramp & Damas, 1925	A
<i>Ophinella</i>		<i>lactea</i> Agassiz, 1865	A
<i>parasitica</i> (Sars, 1873)	A	<i>Stauroidiscus</i>	
<i>Ventromma</i>		<i>tetrastaurus</i> Haeckel, 1879	
<i>halecioides</i> (Alder, 1859)	A M	<i>Staurostoma</i>	
= <i>Kirchenpaueria halecioides</i>		<i>mertensi</i> (Brandt, 1838)	A M
Family Lafocidae		= <i>Staurophora mertensi</i>	
<i>Acryptolaria</i>		<i>Teclaia</i>	
<i>conferta</i> (Allman, 1877)	A M	<i>recincola</i> Gili, Bouillon, Pagès, Palanques & Puig, 1999	M
<i>crassicaulis</i> (Allman, 1888)	A	<i>Toxorchis</i>	
<i>longitheca</i> (Allman, 1877)	A M	<i>arcuatus</i> Haeckel, 1879	
= <i>A. crassicaulis</i>		Family Lovenellidae	
<i>Anthohebella</i>		<i>Lovenella</i>	
<i>parasitica</i> (Ciamician, 1880)	A M	<i>chiquitita</i> Millard, 1959	M
= <i>Hebella parasitica</i>		<i>cirrata</i> (Haeckel, 1879)	A M
<i>Bedotella</i>		= <i>Campalecium? cirrata</i>	
<i>armata</i> (Pictet & Bedot, 1900)	A	<i>clausa</i> (Lovén, 1836)	A M
<i>Cryptolarella</i>		<i>gracilis</i> Clarke, 1882	A M
<i>abyssicola</i> (Allman, 1888)	A M	<i>paniculata</i> (Sars, 1873)	A M
= <i>C. diffusa</i>		<i>producta</i> (Sars, 1873)	A
= <i>C. humulis</i>		Family Malagazziidae	
<i>Cryptolaria</i>		<i>Octophialucium</i>	
<i>exserta</i> Busk, 1858	A	<i>funerarium</i> (Quoy & Gaimard, 1827)	A M
<i>pectinata</i> (Allman, 1888)	A	Family Melicertidae	
<i>Filellum</i>		<i>Melicertum</i>	
<i>disaggregatum</i> Peña-Cantero, García-Carrascosa & Vervoort, 1998	M	<i>octocostatum</i> (Sars, 1835)	A
<i>serpens</i> (Hassall, 1848)	A M	<i>Orchistomella</i>	
<i>serratatum</i> (Clarke, 1879)	A M	<i>graeffei</i> (Neppi & Stiasny, 1911)	M
<i>Grammaria</i>		Family Mitrocomidae	
<i>abietina</i> (Sars, 1851)	A	<i>Cosmetira</i>	
<i>immersa</i> Nutting, 1901	A	<i>pilosella</i> Forbes, 1848	A
<i>Hebella</i>		<i>Cyclocanna</i>	
<i>brochi</i> (Hadzi, 1913)	A M	<i>welshi</i> Bigelow, 1918	A
<i>scandens</i> (Bale, 1888)	A M	<i>Foersteria</i>	
= <i>H. besnardi</i>		<i>antoniae</i> Gili, Bouillon, Pagès, Palanques, Puig & Heussner, 1998	M
= <i>H. eximia</i>		<i>araiiae</i> Gili, Bouillon, Pagès, Palanquea & Puig, 1999.	M
= <i>H. michaelseni</i>		<i>Halopsis</i>	
= <i>H. sinuosa</i>		<i>ocellata</i> Agassiz, 1863	A
= <i>H. urceolata</i>		<i>Mitrocoma</i>	
<i>Lafocia</i>		<i>annae</i> Haeckel, 1864	M
<i>dumosa</i> (Fleming, 1820)	A M	<i>Mitrocomella</i>	
= <i>L. tenellula</i>			

<i>brownei</i> (Kramp, 1930)	A M	= <i>D. cornicina</i> McCrady, 1859	
<i>polydiademata</i> (Romanes, 1876)	A	<i>pumila</i> (Linnaeus, 1758)	A M
Mitrocomium		<i>quadridentata</i> (Ellis & Solander, 1786)	A M
<i>cirratum</i> Haeckel, 1879	A M	Hydrallmania	
Octogonade		<i>falcata</i> (Linnaeus, 1758)	A
<i>mediterranea</i> Zoja, 1896	M	Salacia	
= <i>Melicerta adriatica</i> Neppi, 1915		<i>desmoides</i> (Torrey, 1902)	A
Tiaropsidium		<i>dubia</i> (Billard, 1922)	A M
<i>atlanticum</i> Russell, 1956	A	Selaginopsis	
<i>mediterraneum</i> (Metschnikoff, 1886)	M	<i>fusca</i> (Johnston, 1847)	A
<i>vilaevelebiti</i> (Hadzi, 1915)	M	Sertularella	
= <i>Halanthus vilaevelebiti</i>		<i>arbuscula</i> (Lamouroux, 1816)	
Family Orchistomidae		<i>crassicaulis</i> (Heller, 1868)	M
Orchistoma		<i>cubica</i> Garcia Aguirre & Gonzalez, 1980	M
<i>agariciforme</i> Keller, 1884	M	<i>cylindritheca</i> (Allman, 1888)	A M
Family Phialellidae		= <i>Cnidosecyphus macrotheca</i>	
Phialella		<i>ellisii</i> (Deshayes & Milne Edwards, 1836)	A M
<i>quadrata</i> (Forbes, 1848)	A M	= <i>S. fusiformis</i>	
Family Plumulariidae		<i>fusiformis</i> (Hincks, 1861)	A M
Monothecha		<i>gayi</i> (Lamouroux, 1821)	A M
<i>margaretta</i> Nutting, 1900	A M	<i>mediterranea</i> Hartlaub, 1901	A M
<i>obliqua</i> (Johnston, 1847)	A M	= <i>S. gaudichaudi</i>	
<i>pulchella</i> (Bale, 1882)	A M	<i>ornata</i> Broch, 1933	M
Nematophorus		<i>picta</i> (Meyen, 1834)	A M
<i>clarkei</i> Nutting, 1900	A	<i>polyzonias</i> (Linnaeus, 1758)	A M
= <i>Macrorhynchia clarkei</i>		<i>rugosa</i> (Linnaeus, 1758)	A
Nemertesia		<i>simplex</i> (Hutton, 1873)	A
<i>antennina</i> (Linnaeus, 1758)	A M	<i>tenella</i> (Alder, 1857)	A M
<i>belini</i> Bedot, 1916	A	<i>unituba</i> Calder, 1991	A
<i>falculica</i> Ramil & Vervoort, 1992	A	Sertularia	
= <i>Plumularia falculica</i>		<i>albimaris</i> Merezchkovskii, 1878	A
<i>irregularis</i> (Quelch, 1885)	A M	<i>argentea</i> Linnaeus, 1758	A
<i>norvegica</i> (Sars, 1873)	A	<i>cupressina</i> Linnaeus, 1758	A
<i>perrieri</i> (Billard, 1901)	A M	<i>distans</i> Lamouroux, 1816	A M
<i>ramosa</i> (Lamarck, 1816)	A M	<i>fabricii</i> Levinsen, 1893	A
<i>tetrasticha</i> Meneghini, 1845	M	<i>gracilis</i> Hassall, 1848	
<i>ventriculiformis</i> (Marktanner-Turneretscher, 1890)	A M	<i>inflata</i> (Versluys, 1899)	M
Plumularia		<i>marginata</i> (Kirchenpauer, 1864)	A M
<i>canariensis</i> Izquierdo, Garcia-Corrales & Bacallado, 1986		<i>mediterranea</i> (Marktanner-Turneretscher, 1890)	M
<i>filicula</i> Allman, 1877	A M	<i>mirabilis</i> (Verrill, 1872)	A
<i>floridana</i> Nutting, 1900	A	<i>perpusilla</i> Stechow, 1919	M
<i>linkoi</i> Naumov, 1960	M	<i>plumosa</i> (Clark, 1876)	A
<i>obliqua</i> (Johnston, 1847)	A	<i>robusta</i> (Clark, 1876)	A
<i>setacea</i> (Linnaeus, 1758)	A M	<i>schmidti</i> Kudelin, 1914	A
<i>syriaca</i> Billard, 1931	M	<i>tenera</i> Sars, 1874	A
Polyplumaria		<i>turbinata</i> (Lamouroux, 1816)	A
<i>flabellata</i> (Sars, 1873)	A M	Symplectoscyphus	
= <i>P. billardi</i>		<i>amphorifera</i> (Allman, 1877)	A
<i>gracillima</i> (Sars, 1873)	A	<i>bathyalis</i> Vervoort, 1972	A
Pseudoplumaria		<i>tricuspidatus</i> (Alder, 1856)	A
<i>marocana</i> (Billard, 1930)	A M	Tamarisca	
<i>sabinae</i> Ramil & Vervoort, 1992	A	<i>tamarisca</i> (Linnaeus, 1758)	A M
Family Sertulariidae		Thutaria	
Abietinaria		<i>alternitheca</i> Levinsen, 1893	A
<i>abietina</i> (Linnaeus, 1758)	A M	<i>articulata</i> (Pallas, 1766)	A
<i>filicula</i> (Ellis & Solander, 1786)	A	= <i>Salacia articulata</i>	
<i>intersersa</i> (Pictet & Bedot, 1900)	A	<i>breitfussi</i> (Kudelin, 1914)	A
<i>pulchra</i> (Nutting, 1904)	A	<i>carica</i> Levinsen, 1893	A
Amphisbetia		<i>cupressoides</i> (Lepechin, 1783)	A
<i>operculata</i> (Linnaeus, 1758)	A M	<i>decemserialis</i> (Merezchkovskii, 1878)	A
Diphasia		<i>hippuris</i> Allman, 1874	A
<i>alata</i> (Hincks, 1855)	A	<i>lichenastrum</i> (Pallas, 1766)	A
<i>attenuata</i> (Hincks, 1866)	A M	<i>obsoleta</i> (Lepechin, 1781)	A
<i>delagei</i> (Billard, 1912)	A	<i>thuja</i> (Linnaeus, 1758)	A
<i>fallax</i> (Johnston, 1847)	A	<i>variabilis</i> (Broch, 1918)	A
<i>margareta</i> (Hassall, 1841)	A M	Family Syntheciidae	
<i>nigra</i> (Pallas, 1766)	A	Synthecium	
<i>pinastrum</i> (Cuvier, 1830)	A M	<i>evansi</i> (Ellis & Solander, 1786)	A M
<i>rosacea</i> (Linnaeus, 1758)	A M	Family Tiarannidae	
Dynamena		Modeeria	
<i>crisiodes</i> Lamouroux, 1824	A	<i>rotunda</i> (Quoy & Gaimard, 1827)	A M
<i>dalmasi</i> (Versluys, 1899)	A	= <i>Stegopoma fastigiata</i> (Alder, 1860)	
<i>disticha</i> (Bosc, 1802)	A	Stegolaria	
		<i>geniculata</i> (Allman, 1888)	A M
		Stegopoma	

<i>bathyale</i> Vervoort, 1966	A	<i>multicornis</i> (Forskål, 1775)	A M
<i>giganteum</i> Ramil & Vervoort, 1992	A	= <i>C. squamata</i>	
<i>plicatile</i> (Sars, 1863)	A	<i>Cordylophora</i>	
<i>Tiaranna</i>		<i>caspia</i> (Pallas, 1771)	A
<i>ducalis</i> (Forbes & Goodsir, 1853)		<i>inkermanica</i> Marfenin, 1983	M
<i>globulosa</i> (Forbes, 1848)		<i>pusilla</i> Motz-Kossowska, 1905	M
Family Tiaropsidae		<i>Corydendrium</i>	
<i>Tiaropsis</i>		<i>dispar</i> Kramp, 1935	A
<i>multicirrata</i> (Sars, 1835)	A	<i>parasiticum</i> (Linnaeus, 1767)	A
Order Filifera		<i>Merona</i>	
Family Bougainvilliidae		<i>cornucopiae</i> Norman, 1864	A
<i>Aselomaris</i>		<i>ibera</i> Medel, García-Gómez & Bouillon, 1993	M
<i>arenosa</i> (Alder, 1863)	A M	<i>nematophorum</i> (Antsulevich & Polteva, 1986)	A
<i>michaeli</i> Berrill, 1948	A M	<i>Oceania</i>	
= <i>Clavopsella michaeli</i>		<i>armata</i> Källiker, 1853	A
<i>Bimeria</i>		<i>Tubiclava</i>	
<i>vestita</i> Wright, 1859	A M	<i>fruticosa</i> Allman, 1872	A
<i>Bougainvillia</i>		<i>lucerna</i> Allman, 1863	A M
<i>britannica</i> (Forbes, 1841)	A M	<i>Turritopsis</i>	
<i>charcoti</i> Le Danois, 1913		<i>nutricula</i> McCrady, 1857	A
<i>flavida</i> Hartlaub, 1887		= <i>Corydendrium nutricula</i>	
<i>macloviana</i> Lesson, 1836	A	Family Cytaeidae	
<i>maniculata</i> Haeckel, 1864	A M	<i>Cytaeis</i>	
<i>multicilia</i> (Haeckel, 1879)	M	<i>pusilla</i> Gegenbaur, 1856	M
<i>muscoides</i> (Sars, 1846)	A	<i>tetrastyla</i> Eschscholtz, 1829	A
<i>muscus</i> (Van Beneden, 1844)	A	<i>Perarella</i>	
= <i>B. ramosa</i>		<i>propagulata</i> Bavestrello, 1987	M
<i>principis</i> (Steenstrup & Lütken, 1850)	A	<i>schneideri</i> (Motz-Kossowska, 1905)	M
<i>pyramidata</i> (Forbes & Goodsir, 1853)	A	Family Eudendriidae	
<i>rugosa</i> Clarke, 1882	A	<i>Eudendrium</i>	
<i>superciliaris</i> (Agassiz, 1849)	A	<i>album</i> Nutting, 1898	A
<i>Clavopsella</i>		<i>annulatum</i> Norman, 1864	A
<i>navis</i> (Millard, 1959)	A	<i>arbusculum</i> Wright, 1859	A M
<i>Dicoryne</i>		<i>armatum</i> Tichomiroff, 1890	A M
<i>conferta</i> (Alder, 1856)	A M	<i>calceolatum</i> Motz-Kossowska, 1905	M
<i>conybearei</i> (Browne, 1907)	A M	<i>capillare</i> Alder, 1856	A M
<i>flexuosa</i> Sars, 1873	A	<i>carneum</i> Clarke, 1882	A
<i>Garveia</i>		<i>cunninghami</i> Kirkpatrick, 1910	A M
<i>biscayana</i> (Browne, 1907)	A	<i>fragile</i> Motz-Kossowska, 1905	M
<i>franciscana</i> (Torrey, 1902)	A	<i>glomeratum</i> Picard, 1951	A
<i>grisea</i> (Motz-Kossowska, 1905)	M	<i>insigne</i> Hincks, 1861	A M
<i>mutans</i> Wright, 1859	A M	<i>merulum</i> Watson, 1985	M
<i>Koellikerina</i>		<i>motzkossowskiae</i> Picard, 1951	M
<i>fasciculata</i> (Péron & Lesueur, 1809)	A M	<i>planum</i> Bonnevie, 1898	A
<i>Lizzella</i>		<i>racemosum</i> (Cavolini, 1785)	A M
<i>hyalina</i> (Van Beneden, 1866)	A	<i>rameum</i> (Pallas, 1766)	A M
<i>Lizzia</i>		<i>ramosum</i> (Linnaeus, 1758)	A M
<i>blondina</i> Forbes, 1848	A M	<i>teissieri</i> Cabioch, 1970	A
<i>fulgurans</i> (Agassiz, 1865)	A M	<i>tenellum</i> Allman, 1877	A
<i>octostyla</i> (Haeckel, 1879)	M	<i>vaginatum</i> Allman, 1863	A
<i>Nemopsis</i>		= <i>E. annulatum</i>	
<i>bachei</i> Agassiz, 1849	A	<i>vervoorti</i> Marques & Migotto, 1998	A
<i>crucifera</i> (Forbes & Goodsir, 1853)	A	<i>Myrionema</i>	
<i>Nubiella</i>		<i>amboinense</i> Pictet, 1893	A
<i>mitra</i> Bouillon, 1980	M	Family Hydractiniidae	
<i>Pachycordyle</i>		<i>Hydractinia</i>	
<i>napolitana</i> Weismann, 1883	A M	<i>aculeata</i> (Wagner, 1833)	M
<i>Rhizorhagium</i>		<i>allmani</i> Bonnevie, 1898	A
<i>album</i> Rees, 1938	A	<i>arctica</i> (Jäderholm, 1902)	A
<i>formosum</i> (Fewkes, 1889)	A M	= <i>H. ingolfti</i>	
<i>roseum</i> Sars, 1874	A	= <i>Stylactaria arctica</i>	
<i>sarsii</i> (Bonnevie, 1898)	A	<i>calderi</i> Bouillon, Medel & Peña-Cantero, 1997	M
<i>Thamnostoma</i>		<i>carica</i> Bergh, 1887	A
<i>dibalium</i> (Busch, 1851)	M	<i>echinata</i> Fleming, 1823	A M
<i>russelli</i> Rees, 1939	A	<i>fucicola</i> (Sars, 1857)	A M
Family Calycopsidae		<i>humilis</i> Bonnevie, 1898	A
<i>Bythotia</i>		<i>monocarpa</i> Allman, 1847	A
<i>murrayi</i> Günther, 1903	A	<i>ornata</i> Bonnevie, 1899	A M
<i>Calycopsis</i>		<i>serrata</i> Kramp, 1943	A
<i>gara</i> Petersen, 1957	A	<i>Podocoryna</i>	
<i>krampi</i> Petersen, 1957	A	<i>areolata</i> (Alder, 1862)	A M
<i>simplex</i> Kramp & Damas, 1925	A M	= <i>P. hartlaubi</i>	
Family Clavidae		<i>borealis</i> (Mayer, 1900)	A M
<i>Clava</i>		<i>carnea</i> Sars, 1846	A M

<i>exigua</i> (Haeckel, 1879)	M	<i>affinis</i> Moseley, 1879	A
<i>minima</i> (Trinci, 1903)	A M	<i>medioatlantica</i> Zibrowius & Cairns, 1992	A
<i>minuta</i> (Mayer, 1900)	M	<i>tenuiseptata</i> Cairns, 1986	A
<i>proboscidea</i> Hincks, 1869	A	<i>vascomarquesi</i> Zibrowius & Cairns, 1992	A
<i>sarsi</i> Steenstrup, 1850	A	<i>Errina</i>	
Stylactis		<i>aspera</i> (Linnaeus, 1767)	A M
<i>claviformis</i> Bouillon, 1971	A	<i>atlantica</i> Hickson, 1912	A
<i>fuciola</i> (Sars, 1857)	M	<i>dabneyi</i> (Pourtalès, 1871)	A
= <i>Hydractinia fuciola</i>		<i>Lepidopora</i>	
<i>inermis</i> (Allman, 1872)	M	<i>eburnea</i> (Calvet, 1903)	A
<i>pruvoti</i> (Motz-Kossowska, 1905)	A M	<i>Pliobothrus</i>	
= <i>Hydractomma pruvoti</i>		<i>gracilis</i> Zibrowius & Cairns, 1992	A
<i>Tregoubovia</i>		<i>symmetricus</i> Pourtalès, 1868	A
<i>atentaculata</i> Picard, 1958	M	<i>Stenohelia</i>	
Family Hydrichthyidae		<i>maderensis</i> (Johnston, 1862)	A
Hydrichthys		<i>Stylaster</i>	
<i>cyclothonis</i> Damas, 1934	A	<i>erubescens</i> subsp. <i>britannicus</i> Zibrowius & Cairns, 1992	A
<i>sarcotretis</i> (Jungersen, 1911)	A	<i>erubescens</i> subsp. <i>groenlandicus</i> Zibrowius & Cairns, 1992	A
= <i>Ichthyocodium sarcotretis</i>		<i>erubescens</i> subsp. <i>meteroensis</i> Zibrowius & Cairns, 1992	A
Family Niobiidae		<i>gemmascens</i> (Esper, 1794)	A
Niobia		<i>ibericus</i> Zibrowius & Cairns, 1992	A
<i>dendrotentaculata</i> Mayer, 1900	A	<i>maroccanus</i> Zibrowius & Cairns, 1992	A
Family Pandeidae		<i>norvegicus</i> (Gunnerus, 1768)	A
Amphinema		Family Trichydridae	
<i>dinema</i> (Péron & Lesueur, 1810)	A	<i>Trichydra</i>	
<i>krampi</i> Russell, 1956	A	<i>pubica</i> Wright, 1858	A
<i>rubrum</i> (Kramp, 1957)	M		
<i>rugosum</i> (Mayer, 1900)	A	Order Laingiomedusae	
<i>turridum</i> (Mayer, 1900)	A	Family Laingiidae	
Annatiara		<i>Kantiella</i>	
<i>affinis</i> (Hartlaub, 1914)	A	<i>enigmatica</i> Bouillon, 1978	M
Catablema		Order Limnopolypae	
<i>vesicarium</i> (A. Agassiz, 1862)	A	Family Armorhydridae	
Codonorchis		<i>Armorhydra</i>	
<i>octaedrus</i> Haeckel, 1879	A M	<i>janowiczi</i> Swedmark & Teissier, 1958	A M
Halitholus		Family Microhydrulidae	
<i>pauper</i> Hartlaub, 1913	A	<i>Microhydrula</i>	
<i>yoldiaarcticae</i> (Birula, 1897)	A	<i>pontica</i> Valkanov, 1965	A M
Leuckartiara		<i>Rhaptapagis</i>	
<i>abyssi</i> (G.O. Sars, 1874)	A	<i>cantacuzenei</i> Bouillon & Deroux, 1967	A
<i>breviconis</i> (Murbach & Shearer, 1902)	A	Family Olindiasidae	
<i>nobilis</i> Hartlaub, 1913	A M	<i>Craspedacusta</i>	
<i>octona</i> (Fleming, 1823)	A	<i>sowerbyi</i> Lankester, 1880	A M
Merga		<i>Gonionemus</i>	
<i>galleri</i> Brinckmann, 1962	M	<i>vertens</i> A. Agassiz, 1862	A
<i>reesi</i> Russell, 1956	A	<i>vindobonensis</i> Joseph, 1918	M
<i>tergestina</i> (Neppi & Stiasny, 1912)	M	<i>Gossea</i>	
<i>tregoubovii</i> Picard, 1960	M	<i>corynetes</i> (Gosse, 1853)	A M
Neoturris		<i>faureae</i> Picard, 1952	A
<i>pileata</i> (Forskål, 1775)	A M	<i>Maeotias</i>	
Octotiarra		<i>inexpectata</i> Ostroumov, 1896	A M
<i>violacea</i> Kramp, 1959	A	<i>Monobrachium</i>	
Pandea		<i>parasiticum</i> Mereschkowsky, 1877	A M
<i>conica</i> (Quoy & Gaimard, 1827)	A	<i>Olindias</i>	
<i>rubra</i> Bigelow, 1913	A	<i>phosphorica</i> (Delle Chiaje, 1841)	A M
Stomotoca		Family Proboscicactylidae	
<i>geogrinae</i> (Hadzi, 1913)	M	<i>Fabienna</i>	
Family Protiaridae		<i>oligonema</i> (Kramp, 1953)	A M
Halitiara		= <i>Trichydra oligonema</i>	
<i>formosa</i> Fewkes, 1882	A	<i>Proboscicactyla</i>	
Paratiara		<i>flavicirrata</i> Brandt, 1835	A
<i>digitalis</i> Kramp & Damus, 1925	A	<i>ornata</i> (McCrary, 1859)	A
Protiara		<i>stellata</i> (Forbes, 1846)	A
<i>tetranema</i> (Péron & Lesueur, 1809)	A M	Order Narcomedusae	
Family Ptilocodiidae		Family Aeginidae	
Thecocodium		<i>Aegina</i>	
<i>brieni</i> Bouillon, 1967	A M	<i>citrea</i> Eschscholtz, 1829	A M
<i>penicillatum</i> Jarms, 1987	A	= <i>A. rosea</i>	
Family Rathkeidae		<i>Aeginopsis</i>	
Rathkea		<i>laurentii</i> Brandt, 1838	A
<i>octopunctata</i> (Sars, 1835)	A M		
Family Stylasteridae			
Crypthelia			

<i>Aeginura</i>		<i>everta</i> (Clark, 1876)	M
<i>grimaldii</i> Maas, 1904	A	<i>integra</i> (MacGillivray, 1842)	A M
<i>Solmundella</i>		<i>Phialidium</i>	
<i>bitentaculata</i> (Quoy & Gaimard, 1833)	A	<i>discoideum</i> (Mayer, 1900)	M
Family Cuninidae		= <i>Clytia discoideum</i>	
<i>Cunina</i>		<i>folleatum</i> McCrady, 1859	A M
<i>fowleri</i> (Browne, 1906)		= <i>P. pentata</i>	
= <i>Cunocantha fowleri</i>		<i>islandicum</i> Kramp, 1919	A
<i>frugifera</i> Kramp, 1948	A M	<i>mccradyi</i> (Brooks, 1888)	M
<i>globosa</i> Eschscholtz, 1829	A	<i>Rhizocaulus</i>	
<i>lativentris</i> Gegenbaur, 1856	A	<i>verticillatus</i> (Linnaeus, 1758)	A
<i>octonaria</i> McCrady, 1857	A	<i>Tulpa</i>	
<i>polygonia</i> (Haeckel, 1879)	M	<i>speciosa</i> (Clark, 1876)	A
<i>proboscidea</i> E. & L. Metschnikoff, 1871	M	Order Trachymedusae	
<i>simplex</i> Gili, Bouillon, Pagès, Palanques, Puig & Heussner, 1998	M	Family Geryoniidae	
<i>vitrea</i> Gegenbaur, 1856	M	<i>Geryonia</i>	
<i>Solmissus</i>		<i>proboscidalis</i> (Huxley, 1859)	A
<i>albescens</i> (Gegenbaur, 1856)	A M	<i>Liriope</i>	
<i>incisus</i> (Fewkes, 1886)	A	<i>tetraphylla</i> (Otto, 1823)	A
Family Solmarisidae		Family Halicreatidae	
<i>Pegantha</i>		<i>Botrynema</i>	
<i>clara</i> Bigelow, 1909	A	<i>brucei</i> Browne, 1908	A
<i>mollicina</i> (Forskål, 1775)	M	<i>ellinorae</i> (Hartlaub, 1909)	A
<i>rubiginosa</i> (Kölliker, 1853)	A M	<i>Halicreas</i>	
<i>triloba</i> Haeckel, 1879	A	<i>minimum</i> Fewkes, 1882	A
<i>zonaria</i> (Haeckel, 1879)	M	<i>papillosum</i> Vanhöffen, 1902	A
<i>Solmaris</i>		<i>Haliscera</i>	
<i>corona</i> (Keferstein & Ehlers, 1861)	A	<i>bigelowi</i> Kramp, 1947	A M
<i>flavescens</i> (Kölliker, 1853)	M	<i>conica</i> Vanhöffen, 1902	A
<i>leucostyla</i> (Will, 1844)	M	<i>racovitzae</i> (Maas, 1906)	M
<i>solmaris</i> (Gegenbaur, 1856)	M	= <i>Homoeonema racovitzae</i>	
<i>vanhoeffeni</i> Neppi & Stiasny, 1911	M	<i>Halitrephes</i>	
Order Proboscoida		<i>maasi</i> Bigelow, 1909	A
Family Bonneviellidae		<i>Homoeonema</i>	
<i>Bonneviella</i>		<i>platygonum</i> Browne, 1903	A
<i>grandis</i> (Allman, 1876)	A	Family Petasidae	
Family Campanulariidae		<i>Petasus</i>	
<i>Campanularia</i>		<i>atavus</i> Haeckel, 1879	A M
<i>crenata</i> (Hartlaub, 1901)	M	Family Ptychogastridae	
<i>hincksi</i> Alder, 1856	A M	<i>Ptychogastria</i>	
<i>volubilis</i> (Linnaeus, 1758)	A M	<i>asteroides</i> (Haeckel, 1879)	M
<i>Clytia</i>		<i>polaris</i> Allman, 1878	A
<i>gracilis</i> (Sars, 1850)	A M	<i>Tesserogastria</i>	
<i>hemisphaerica</i> (Linnaeus, 1767)	A M	<i>musculosa</i> Beyer, 1959	A
<i>islandicum</i> Kramp, 1919	A	Family Rhopalonematidae	
<i>linearis</i> (Thomeley, 1900)	A	<i>Aglantha</i>	
<i>maccradyi</i> (Brooks, 1888)	A	<i>digitale</i> (Müller, 1776)	A M
<i>noliformis</i> (McCrady, 1859)	A M	<i>elata</i> (Haeckel, 1879)	M
<i>paulensis</i> (Vanhöffen, 1910)	A M	<i>Aglaura</i>	
<i>pentata</i> (Mayer, 1900)	A	<i>hemistoma</i> Péron & Le Sueur, 1809	A
<i>raridentata</i> (Alder, 1862)	M	<i>Arctapodema</i>	
<i>serrulata</i> (Bale, 1888)	M	<i>ampla</i> (Vanhöffen, 1902)	A
<i>Gonothyraea</i>		<i>australe</i> (Vanhöffen, 1912)	M
<i>loveni</i> (Allman, 1859)	A M	<i>Colobonema</i>	
<i>Hartlaubella</i>		<i>sericeum</i> Vanhöffen, 1902	A
<i>gelatinosa</i> (Pallas, 1766)	A M	<i>Crossota</i>	
<i>Laomedea</i>		<i>norvegica</i> Vanhöffen, 1902	A
<i>angulata</i> Hincks, 1861	A M	<i>ryfobrunnea</i> (Kramp, 1913)	A
<i>calceolifera</i> (Hincks, 1871)	A M	<i>Homoeonema</i>	
<i>exigua</i> Sars, 1857	A M	<i>platygonon</i> Browne, 1903	A M
<i>flexuosa</i> Alder, 1857	A M	<i>Pantachogon</i>	
<i>neglecta</i> Alder, 1856	A M	<i>haeckeli</i> Maas, 1893	A M
<i>pseudodichotoma</i> Vervoort, 1959	A	<i>militare</i> (Maas, 1893)	A M
<i>Obelia</i>		<i>Persa</i>	
<i>bidentata</i> Clark, 1875	A M	<i>incolorata</i> McCrady, 1857	A
<i>dichotoma</i> (Linnaeus, 1758)	A M	<i>Ransonia</i>	
<i>fimbriata</i> (Dalyell, 1848)	A M	<i>krampi</i> (Ranson, 1932)	A M
<i>geniculata</i> (Linnaeus, 1758)	A M	<i>Rhopalonema</i>	
<i>longissima</i> (Pallas, 1766)	A M	<i>funerarium</i> Vanhöffen, 1902	A
<i>Orthopyxis</i>		<i>velatum</i> Gegenbaur, 1856	A
<i>asymmetrica</i> Stechow, 1919	M	<i>Sminthea</i>	
<i>crenata</i> Hartlaub, 1901	A	<i>arctica</i> Hartlaub, 1909	A
= <i>Campanularia crenata</i>		<i>eurygaster</i> Gegenbaur, 1856	A

Tetrorchis erythrogaster Bigelow, 1909

A

Siphonophora

Compiled by G. M. Mapstone

This list was initially drawn up from 'Siphonophores and Verellids' by P. A. Kirkpatrick and P. R. Pugh (1984), and additional species added from other publications covering species recorded within the ERMS geographic area. The reference list includes all taxonomic papers covering any new and redescribed species found in the area, but only those distributional papers published from 1990 onwards. A more comprehensive literature list will be obtainable from a monograph entitled 'Siphonophora (Cnidaria, Hydrozoa) of Canadian Pacific waters.' by G. M. Mapstone and M. N. Arai which is due to be published shortly.

Since the species list is a reflection of sampling effort, it is likely that more species will be added in the future; many Siphonophora species have a cosmopolitan distribution.

P. R. Pugh kindly advised on the validity of species included in a world list from which this ERMS was derived.

References

- Andersen, V., Sardou, J. and Nival, P. 1992. The diel migrations and vertical distributions of zooplankton and micronekton in the Northwestern Mediterranean Sea. 2. Siphonophores, hydromedusae and pyrosomids. *J. Plankton Res.* 8, 1155-1169.
- Bamstedt, U., Fossa, J. H., Martinussen, M. B. and Fossagen, A. 1998. Mass occurrence of the physonect siphonophore *Apoemia uvaria* (Lesueur) in Norwegian waters. *Sarsia* 83, 79-85.
- Buecher, E. 1999. Appearance of *Chelophyes appendiculata* and *Abylopsis tetragona* (Cnidaria, Siphonophora) in the Bay of Villefranche, northwestern Mediterranean. *Journal of Sea Research* 41, 295-307.
- Biggs, D. C., Pugh P. R. and Carré, C. 1978. *Rosacea flaccida* n. sp., a new species of siphonophore (Calycophorae Prayinae) from the North Atlantic Ocean. *Beaufortia* 27: 207-218.
- Gamulin, T. and Krsinic, F. 1993. Distribution and abundance of Calycophores (Siphonophora, Calycophorae) in the Mediterranean and Adriatic Sea. *Mar. Ecol. (Pubbl. Stn. Zool. Napoli I)*. 14 (2): 97-111.
- Gili, J-M., Pagès, F. and Fusté, X. 1991. Mesoscale coupling between spatial distribution of planktonic cnidarians and hydrographic features along the Galician Coast (Northwestern Iberian Peninsula). *Sci. Mar.* 55(2): 419-426.
- Mapstone, G. M. 1998. *Bargmannia lata*, An undescribed species of physonect siphonophore (Cnidaria, Hydrozoa) from Canadian Pacific waters. 141-147, figs 1-3. In *Commemorative volume for the 80th birthday of Willem Vervoort in 1997*. Edited by: J. C. den Hartog, A. C. van Bruggen, P. F. S. Cornelius and L. P. van Ofwegen. *Zool. Verh. Leiden*: 323: i-xii, 1-448.
- Mills, C. E., Pugh, P. R., Harbison, G. R., and Haddock, S. H. D. 1996. Medusae, siphonophores and ctenophores of the Alborán Sea, south western Mediterranean. In *Advances in Hydrozoan Biology*. Edited by S. Piraino, F. Boero, J. Bouillon, P. F. S. Cornelius and J.-M. Gili. *Sci. Mar.* 60 (1): 145-163.
- Kirkpatrick, P. A. and Pugh, P. R. 1984. Siphonophores and verellids. *Synopses of the British Fauna* (New Series). Leiden, E. J. Brill Number 29, Pp. 154.
- Pagès, F., González, H. E., and González, S. R. 1996. Diet of the gelatinous zooplankton in Hardangerfjord (Norway) and potential predatory impact by *Aglantha digitale* (Trachymedusae). *Mar. Ecol. Prog. Ser.* 139: 69-77.
- Pinca, S., and Dallot, S. 1995. Meso- and macrozooplankton composition patterns related to hydrodynamic structures in