

Index for Volume 52 (2003)

Syst. Biol. 52(6):876–911, 2003
Copyright © Society of Systematic Biologists
ISSN: 1063-5157 print / 1076-836X online
DOI: 10.1080/10635150390276444

COMPILED BY JOE MADIA

- 12S rDNA, 407, 727, 798–799
12S rRNA, 39, 42, 56–58, 161, 163–165, 167, 181, 206–207, 209–212, 215, 218, 297, 440, 443, 445–447, 451, 456, 460, 462–467, 605
16S mtDNA, 738
16S rDNA, 72–76, 407, 727, 798–799
16S rRNA, 206–207, 209–212, 215, 219, 297–298, 307, 440, 443, 445–447, 451, 456, 460, 462–467, 557, 559, 618–640, 758–759, 762
18S rDNA, 57, 72–75, 820, 823
18S rRNA, 283–295, 296, 298, 307, 311–333, 557, 559
26S rDNA, 820, 823
28S rDNA, 57, 72–75
28S rRNA, 285, 287, 298, 307, 557, 559, 618, 620–623, 745–756
- A2AB, 605, 614–615
A6, 674
Abdo, Zaid, *see* Minin, Vladimir, 674–683
Acacia
 koa, 820
Acaenosuchus, 241–243, 250–251
Acanthalictus, 24, 26, 32
Acari, 298, 430
Acaridae, 430
accD, 336
Accelerated likelihood surface exploration: the likelihood ratchet,
 R. A. Vos, 368–373
Accuracy, 55–65, 428–435
Achalinus, 451
 rufescens, 440
Achromorpha, 314–315, 328, 333
Acleisanthes, 334–351
 acutifolia, 341, 343
 angustifolia, 341–343, 349
 chenopodioides, 341
 crassifolia, 347
 lanceolata, 341, 348
 longiflora, 347
 parvifolia, 341, 347–348
 purpusiana, 341
 undulata, 341, 343, 349
 wrightii, 343, 347
Aclys, 758, 775
 concinna, 758
Acomastylis, 374
Acrocera bulla, 756
Acroceridae, 756
Acrochordidae, 439–440, 442, 447, 455, 457
Acrochordus, 439, 442–443, 451–452, 457
 granulatus, 440
 javanicus, 440
Adams consensus tree, 850
Adams, Dean C., *see* Naylor, Gavin J. P., 864–865
Adams, Richard J., *see* Johnson, Kevin P., 37–47
Adams tree, 463
Adaptation, 501–514
Adaptive radiation, 70, 89–92, 207
Advanced snakes, 439–459
Aepyroceros, 209, 212, 215
 melampus, 210
Aepycerotini, 209–210
Aetosaur, 253–255
 phylogeny, 852
Aetosauria, 239
Aetosauroides, 241, 243, 248, 250–252
 scagliai, 250
Aetosaurus, 241–243, 248, 250–252
Affine, 66–69
African elephants, 591
Agamidae, 757
Agamid lizards, 758
Agaricales, 430
Agkistrodon, 447, 451
 contortrix, 440
Agrionemis exsudans, 94
Agriocnemus, 94
Ailurus fulgens, 846
Akaike Information Criterion (AIC), 79, 89, 95, 824
Alethinophidia, 440
Alexeter sp., Alcini, 209
Algae, 430, 518, 524
Algorithm, 696–704
Alinkidae, 751
Alivon, Eliane, *see* Manuel, Michael, 311–333
Alcelaphini, 209–210
Alleles, 374
 mitochondrial, 41
 nuclear, 41, 103
Alleno pithecus nigroviridis, 736
Alligator, American, 388
Alligator, 391, 409
 mississippiensis, 286–287, 386, 388, 391, 405
 sinensis, 405
Alligatoridae, 394, 396
Alligatoridea, 403, 405, 408–409, 412, 415
Allodape, 5, 7–8
 mucronata, 2
 skaifeorum, 2
Allodapine, 2, 7–9, 11
 bees, 1
Allodapini, 1–14
Allodapula, 2, 9–10, 12
Allodapulodes, 9
Allopolyploidy, 374–385
Allothyrsus sp., 300, 306
Allozyme, 23, 504, 506, 508–509
 trees, 509
 α -lactalbumin, 206, 209–212, 215, 218–220

- Alsinidendron*, 832
 Alsiinoideae, 833
Alsophis, 447, 451
 portoricensis, 440
 Amato, George, *see* Gatesy, John, 403–422
 Amber fossils, 746
Amblonyx, 590
 cinereus, 573, 577
Amblyomma, 299, 304, 309–310
 americanum, 299
 cajennense, 299
 finbriatum, 299
 glebopalma, 299
 hebraeum, 299, 306
 latum, 299
 maculatum, 299
 s.l., 310
 varanensis, 299
 variegatum, 299
 vikirri, 299
 Amblyomminae, 299
 Amblypygi, 62
Ambystoma mexicanum, 287
 American alligator, 388
 Amino acids, 3, 6, 10, 95, 165, 296, 585, 594, 596, 605, 662
Ammocrypta, 491, 493, 496–499
 beanii, 491, 497, 499
 vivax, 491, 496–497, 499
 Among-site rate variation, 604–617
 Amphibians, 286, 524
 Amphimerycidae, 222
 Amphimeryx, 222
 Amphoriscidae, 314
Amphoriscus, 314–315, 332
 Analysis,
 Bayesian, 26, 30, 89, 96, 102, 133, 140, 152, 165, 167, 181–182, 211, 390,
 439–459, 477–487, 571, 576, 583, 604–617, 618, 621, 623, 629, 665,
 681, 708, 713–714, 760, 797, 801–803, 807
 Bayesian information, 377
 biogeographic, 258, 761, 824
 biotic element, 717–723
 bootstrap, 7, 26, 447, 580, 609, 611, 621, 697, 775, 824, 841
 Bremer support, 211
 canonical correspondence, 687–689, 693
 canonical variate, 190, 196, 198, 491, 496
 cladistic, 9, 81, 253, 403, 541, 572, 738
 cluster, 194, 205, 497, 722
 continuous jackknife function (CJF), 55–58, 61–63
 copath, 41
 cophylogenetic, 737
 correspondence, 687
 cytogenetic, 374
 decay (*see also* Bremer support), 128, 243, 407
 distance, 334, 343
 DNA, 551
 generalized Procrustes, 66, 68
 genetic, 197
 jackknife, 407
 Jungles, 39, 43, 45
 likelihood, 18, 263, 374, 708, 715, 803–804, 807
 local clock, 89, 715
 logDet, 3, 6–7, 11
 Markov Chain Monte Carlo, 140, 213, 486, 750, 801
 maximum likelihood, 1, 3, 7, 11, 26–27, 30, 42, 63, 76, 79, 89, 95–97,
 99, 166–167, 171, 182, 211–212, 215, 231, 318, 322, 327, 329, 341,
 377–378, 386, 389–90, 394–396, 407, 419, 715, 737, 740, 760, 797,
 801, 824, 850
 maximum parsimony, 3, 5–7, 10–11, 95, 166–167, 182, 211, 213, 215,
 275, 317, 322, 327, 737, 797, 801
 molecular, 103–104, 221, 323, 386, 397, 542
 molecular dating, 375
 morphological, 187, 213, 323, 501, 504, 541, 560, 605, 858
 morphometric, 197
 multigene, 604
 multivariate, 652
 neighbor-joining, 463, 530, 532
 nested-clade, 159–185
 NHML, 390
 nonparametric bootstrap, 166
 paleontological, 605
 parsimony, 7, 26, 30, 42, 45, 70, 76, 79, 83, 249, 251, 255, 257, 260, 262,
 271–280, 336, 377, 389–390, 392, 406, 412, 423, 506, 508, 530, 554,
 571, 575, 583, 618, 620–621, 623, 629, 727, 764, 802, 804, 824
 partitioned, 4, 396, 604–617
 partitioned likelihood, 386–402
 phylogeographic, 461
 principal component, 687
 principal coordinate, 860
 reconciliation, 39, 42
 redundancy, 687
 regression, 68
 relative warp, 198
 reweighted, 318
 sensitivity, 554–564
 sequence, 737
 SSR, 612
 stability, 554–564
 star topology, 481
 statistical, 131, 847
 supermatrix, 724, 726, 728, 853
 supertree, 725–726, 728, 853
 total evidence, 839
 tree-reconstruction, 182
 UPGMA cluster, 488, 491, 493, 496–497
 variable rate, 751
 Anamixilla, 314–315, 320, 322–324
 torresi, 316
 Anas platyrhynchos, 286
 Anatomical characters, 742
 Ancestral character state reconstruction, 820–841
 Ancient Allopolyploid speciation in Geinae (Rosaceae): evidence
 from nuclear granule-bound starch synthase (GBSSI) gene
 sequences, Jenny E. E. Smedmark, Torsten Eriksson, Rodger
 C. Evans, and Christopher S. Campbell, 374–385
 Andayani, Noviar, *see* Evans, Ben J., 794–823
 Anemonia sulcata, 316
 Angiosperms, 346, 348, 518, 524, 753, 784
 Animals, 89, 591, 736, 742, 820
 Anisopodidae, 756
 Anocentor, 305
 Anolis, 423, 105
 Anomalodesmata, 557
 Anomaluridae, 605–606, 614
 Anomaluroidea, 614–615
 Anomaluromorpha, 605–606
 Anonyx, 590
 capensis, 573–574, 576–577
 Anopheles, 746, 754
 gambiae, 745
 Anseranas semipalmata, 388
 Antedon, 317
 serrata, 316
 Antelope, 206
 Anthophyte, 787–788

- Anthropoids, 48, 53, 709, 711–713
- Antilocapra*, 207, 209, 212–213, 215, 218, 220–221
americana, 210, 219, 224
- Antilocapridae, 206–207, 209–210, 212–213, 218–219, 224
- Antilocaprinae, 210, 218
- Antilopini, 209–210
- Antipathes lata*, 316
- Ants, 23, 33
- Aotus* 740–742
nancymai, 736, 738, 742
- Aparallactinae, 447
- Aparallactus, 446, 451, 455–456
guentheri, 440, 459
wernerii, 440
- Apes, 524, 707–708
- Aphids, 23, 137, 139–140, 143, 146
- Aphroceras*, 314–315, 332
- Aphylophorales, 430
- Apidae, 1
- Apiocera haruspex*, 756
- Apioceridae, 746, 750
- Apis mellifera*, 26, 28, 296
- Aplocheiloid fishes, 549–553
- Aplocheilidae, 548, 552
- Aplocheiloidei, 552
- Apodontia rufa*, 606
- Apodontidae, 605–606, 608, 614
- Aplopappus, 33
- Aplopeltura*, 451
boa, 440, 446
- Aplysilla sulfurea*, 316
- Approximately unbiased test, 781–793
- Aprasia*, 757–758, 762, 764, 767, 770–771, 773, 776–777
aurita, 759
fusca, 759, 776
inaurita, 759
parapulchella, 759, 776
picturata, 759
pseudopulchella, 759, 776
pulchella, 759, 776
repens, 759, 776
rostrata, 758, 776
smithi, 759
striolata, 759, 770
- Apterygota, 328
- Apteryx*
australis, 286, 471–472, 474
mantelli, 471, 474
- Apus affinis*, 286
- Arachnids, 56, 61, 299
- Araneae, 62, 70, 300, 557
- Araneid, 56, 61
- Araneidae, 57
- Archaea, 659
- Archaeognatha, 557
- Archaeomerycidae, 222
- Archaeomeryx*, 222
- Archeoglobus fulgidus*, 659
- Archosauria, 403, 833
- Archosaurs, 239–240, 253–255
- Arctocyon*, 858
- Arctonyx*, 581, 590
collaris, 573, 577, 582
- Area of endemism, 271–280, 717–723
- Arecaceae, 346
- Are the dental data really at odds with the molecular data?
Morphological evidence for the whale phylogeny
(re)reexamined, Maureen A. O’Leary, John Gatesy, and
Michael J. Novacek, 853–866
- Argasidae, 300, 309
- Argasinae, 300
- Argas lagenoplastis*, 300
- Argia*, 94
sedula, 94
- Aris-Brosou, Stéphane, Least and most powerful phylogenetic tests to
elucidate the origin of the seed plants in the presence of
conflicting signals under misspecified models,
781–793
- Arnedo, Miquel, *see* Hormiga, Gustavo, 70–88
- Arthropeas magnum*, 756
- Arthropods, 70, 107, 296, 299–300, 304–305, 307, 518, 557
- Artibeus jamaicensis*, 462–464
- Articulata, 518, 520
- Artiodactyla, 15–22, 724–729, 854–855, 857–858
- Artiodactylans, 853, 857, 860
hippopotamid, 854
suine, 854
- Ascaltis*, 314–315, 332–333
- Ascandra*, 314–315, 332–333
falcata, 332
minchini, 332
- Ascomycetes, 430
- Asilidae, 750, 756
- Asiloidea, 750, 753
- Asioryctes*, 858
- Aspidelaps*, 451
scutatus, 440
- Assessing progress in systematics with continuous jackknife function
analysis, Jeremy A. Miller, 55–65
- Assumptions, 352–367
- Asteraceae, 348, 820, 823
- Astigmata, 430
- Asymptomatic enumeration, 110–118
- atp6*, 657, 659
atp8, 656–659
- Atelestidae, 756
- Atelestus pulicarius*, 756
- Atherinomorpha, 549
- Atractaspidae, 439–440, 455–456
- Atractaspinae, 440, 447
- Atractaspis*, 451, 455
bibronii, 440
corpulenta, 440
micropholis, 440
- Augochlorini, 23
- Australia, 757–780
- Australictus*, 24
- Austrevylaeus*, 24, 26, 30
- Autopomorphic characters, 865
- Aves, 37, 394, 403–422
- Avian DNA, 389
- Avila, Luciano J., *see* Morando, Mariana, 159–185
- Axial symmetry, 311–333
- Axinella polypoides*, 316
- Azemiops*, 451, 453
feae, 440, 453
- Azemiophinae, 440
- Badgers, 571–573
American, 571–573
Chinese ferret, 573
Eurasian, 573
hog, 573
honey, 572

- Old World, 577, 590
stink, 573
- Baeriida, 314, 317–318, 320, 323–324, 329–330
- Baeriidae, 314, 320, 323
- Balaenoptera*, 209, 211–213, 215
physalus, 210
- Balaenopteridae, 209–210
- Balaenoptera*, 706, 712
- Balanidae, 557
- Barker, Nigel P., *see* Kelly, Christopher M., 439–459
- Barker, Stephen C., *see* Murrell, Anna, 296–310
- Barnacles, 557
- Barriel, Véronique, *see* Hugot, Jean-Pierre, 735–744
- Basidiomycetes, 525
- Basidiomycotina, 430
- Bassariscus astutus, 573
- Bats, 57, 462, 472
short-tailed, 460
- Bayesian, 48–54
analysis, 26, 30, 89, 96, 102, 133, 140, 152, 165, 167, 181–182, 211, 390, 439–459, 477–487, 571, 576, 583, 604–617, 618, 621, 623, 629, 665, 671, 708, 713–714, 745–756, 760, 797, 801–803, 807
approach, 171, 791
consensus tree, 622
estimation, 131–158, 394, 714, 745
factors, 48–50, 52–54, 649–664
framework, 48–49, 95, 372, 650, 666, 714, 760, 790–791
hypothesis, 100
inference, 48, 386, 442, 444–446, 642, 665–673
information analysis, 377
information criteria (BIC), 675–676, 719
methods, 23, 48, 159, 477, 696–697, 705–716, 726, 748, 801
model, 49, 650, 791
model selection, 674–683
phylogenetics, 48, 54, 89–109, 652
posterior probability, 30, 102, 215, 396, 554, 559, 612–613, 665–673
relaxed clock, 206–228
support, 623
tests, 607, 615
theorem, 48, 136, 267, 650, 655
trees, 30, 100, 167, 455, 457, 589, 750, 762, 764, 767, 771, 773, 802
- Bears, 711–713, 794
- Bedfordia*, 91, 94, 96
sp., 102
- Bees, 1–14, 23–26
allodapine, 1
corbiculate, 23, 33
halictid, 23–24, 33
honey, 26
- Beetles,
carabid, 70, 103
Ips, 677
longhorn, 89
- Bennettitales, 753
- Bennett, Jonathan R., *see* Scotland, Robert W., 539–547
- Bernissartia*, 412
- Beroe*, 317
cucumis, 316
- Best-fit trees, 4
- β 2-microglobulin, 738
- β -casein, 207
- β -tubulin, 596, 598–600
- Bicatillus*, 619, 623, 639
extinctorum, 623
- Bidirectional sequencing, 2
- Bifurcating tree (*see also* binary tree), 642
- Bilateria, 311, 316–318, 320
- Binary
characters, 244, 247, 256–257, 335, 535
coding, 841
tree (*see also* bifurcating tree), 114
- Bininda-Emonds, Olaf R. P., Novel versus unsupported clades: assessing the qualitative support for clades in MRP supertrees, 839–851
- Bininda-Emonds, Olaf R. P., Kate E. Jones, Samantha A. Price, Richard Grenyer, Marcel Cardillo, Michael Habib, Andy Purvis, and John L. Gittleman, Supertrees are a necessary not-so-evil: a comment on Gatesy et al., 724–729
- Biochemical tree, 403
- Biodiversity crisis, 428–435
- Biogeographic, 552
analysis, 258, 761, 824
barriers, 618–640
history, 552, 722
inference, 618
pattern, 820
studies, 777
- Biogeography, 7, 70–88, 240, 255–258, 430, 509, 618, 622, 717–723, 757–780, 794, 812, 820–841
historical, 271–280
- Bioinformatics, 119–124
- Biostratigraphy, 240
- Biosynthetic pathways, 334–335, 337, 339, 342, 346, 348
- Biotic elements, 271–280, 717–723
- Biotic element analysis in biogeography, Bernhard Hausdorf and Christian Hennig, 717–723
- Birds, 41, 45–46, 230, 283–287, 293, 334, 391, 404, 549–551, 777–778
ratite, 549–553
- Birth–death
model, 641, 644–645, 647, 761
process, 132, 641–648
- Bitis*, 451
arietans, 440, 459
- Blackburnia*, 103
- Black caiman, 388
- Black ravens, 353
- Black-tailed deer, 210
- Blesbok, 210
- Blouin, Christian, *see* Susko, Edward, 594–603
- Blow flies, 745
- Boa*, 447, 451
constrictor, 440, 443, 447
- Boaedontinae, 455
- Body shape variation, 488–500
- Boidae, 440
- Boiga*, 447, 451, 457
cynodon, 440
dedrophila, 440
- Boiginae, 457
- Boleichthys, 492–493
- Boleostoma, 492–493
- Bollback, Jonathan P., *see* Harshman, John, 386–402
- Bollback, Jonathan P., *see* Huelsenbeck, John P., 131–158
- Bolyphantes alticeps*, 73, 88
- Bombinatoridae, 794
- Bombyliidae, 749–750, 754
- Bombylius*, 754
major, 756
- Bonferroni
adjustment, 723
correction, 16, 279, 686, 783, 807
procedure, 802
- Boophilus*, 299, 305–306, 310
annulatus, 299, 306

- decoloratus*, 299
geigyi, 299
kohlsi, 299
microplus, 298–299, 306
- Bookstein, Fred, *see* Rohlf, F. James, 66–69
- Bootstrap, 3, 39, 41, 43, 95, 99, 128–129, 554, 556, 559, 561, 575–576, 589, 600, 665–673, 737, 791, 841
 analysis, 7, 26, 447, 580, 609, 611, 621, 697, 775, 824, 842
 consensus, 30, 32, 802
 consensus trees, 609
 estimates, 599
 frequencies, 839, 842, 844, 847
 maximum likelihood, 667–668
 nonparametric, 166, 506, 594, 665–667, 669, 671–672, 726, 760, 790, 801
 parametric, 18, 598, 667, 726, 757–780, 781
 percentages, 102, 211, 249, 737, 762, 781–793
 proportion, 129
 RELL, 390
 replicates, 42, 44, 477, 479–481, 621, 823
 support, 4–8, 26, 30, 42, 55, 62, 95, 167, 218, 444, 446, 467–468, 508, 540, 576–577, 581–583, 608–609, 611, 614, 621, 623, 738, 764, 829, 842, 844
 trees, 251, 380–381, 451, 455, 465
 value, 5, 7, 45, 95, 101–102, 211, 289, 320, 322, 336, 378, 477, 480, 484, 504, 506, 508, 510, 557, 576, 588, 609–610, 614–615, 665, 667, 727, 738, 748, 760, 802, 826–829
- Borchiellini, Carole, *see* Manuel, Michael, 311–333
- Boreallodapini, 2
- Borealosuchus* ssp., 409
- Boronia*, 347
- Bos*, 209, 706
taurus, 210–212, 405
- Boselaphini, 209–210
- Boselaphus*, 209–212
tragocamelus, 210
- Bostrycapulus*, 619, 623, 631, 636, 639
 cf. *aculeata* sp.1, 631
- Bothriocroton*, 300, 304–307, 309
concolor, 300
undatum, 300
- Bothriocrotoninae, 300, 307
- Boulengerina, 447, 451
annulata, 440
fasciatus, 440
 sp., 440
- Boury-Esnault, Nicole, *see* Manuel, Michael, 311–333
- Bovidae, 206–228
- Bovinae, 210, 218–219
- Bovini, 209
- Brachycera, 745–756
- Brachylinga* sp., 756
- Branch-and-bound search, 368, 377, 390
- Branchiopoda, 557
- Branch and bound search, 479
- Branch length, 22, 41, 45, 48–50, 52, 76, 81–82, 96, 127–130, 133–135, 137, 156, 211–212, 223, 358–359, 377, 412, 415, 424, 427, 478, 481, 484–485, 508, 529–531, 535, 575, 589, 596, 608, 621, 629, 641, 644, 651–652, 655, 666–667, 671–672, 675, 692, 696–697, 699–700, 703, 709, 761, 782, 784–785, 791, 820, 825, 827, 831–832, 834–835, 841, 849–851, 865–866
 error, 679–681
 estimates, 396, 676–677, 682, 747
- Branch lengths and support, Mark Wilkinson, François-Joseph Lapointe, and David J. Gower, 127–130
- Branch lengths and support: revisited, David A. Morrison, 849–851
- Branch support (*see also* bootstrap, clade support, decay, posterior probability), 15, 26, 30, 211, 323, 407, 773
 partitioned, 15–22, 76, 79, 407, 409, 412
- Branch-swapping, 30, 368–370, 842, 857
 algorithm, 287, 371
 NNI, 372
 TBR, 39, 75, 166, 275, 318, 336, 377–378, 390, 407, 444–445, 463, 504, 506, 575, 607, 621–622, 666, 681, 748, 760, 801, 823–824
- Brassica*, 374
- Braun, Michael, *see* Harshman, John, 386–402
- Braunsapis*, 2, 5, 7
paradoxa, 2
protuberans, 2
unicolor, 2
vitrea, 2
- BRCA1, 605, 613–615
- Bremer
 analysis, 211
 support (*see also* decay), 4, 10–11, 55, 62, 75–77, 127, 307, 318, 320, 322, 554–564, 571, 575, 581–584, 588–589, 665, 842
- Brevineura*, 1, 7–8, 10–11
ploratula, 2
xanthochypeata, 2
- Breviostres, 390
- Briggs, John C., Fishes and birds: Gondwana life rafts reconsidered, 548–553
- Britton, Tom, *see* Erixon, Per, 665–674
- Brownian motion parameter, 714
- Brown, Rafe M., *see* Evans, Ben J., 794–823
- Bryant, David, *see* François-Joseph Lapointe, 865–868
- Bufo valliceps*, 287
- Bungarus*, 451
fasciatus, 440
- Bull, Nicholas J., *see* Schwarz, Michael P., 1–14
- Burn-in trees, 607
- Butterflies, 794
 nymphalid, 28
 swallowtail, 589
- Caenogastropod, 620, 623
- Caenophidia, 439–459
- Caiman*, 391, 396, 415
crocodilus, 405, 412, 417
latirostris, 405
yocare, 388
- Caimanae, 409
- Cairina moschata*,
- Calamariinae, 440, 457
- Calcarea, 311–333
- Calcaronea, 311, 313–315, 318, 320, 322–325, 328–329
- Calcinea, 311, 313–315, 318, 320, 322–325, 328–329
- Calcispongia, 311–333
- Callilictus, 24
- Callimico*, 740, 742
goeldii, 736
- Calliopheinae, 443
- Callithrix*, 706, 740, 742
geoffroyi, 736
jacchus, 736, 738
- Callitrichinae, 740, 742
- Calloselasma*, 447, 451
rhodostoma, 441
- Calyptraea*, 618–640
aspersa, 638
 cf. *conica*, 638
 cf. *lichen*, 638

- chinensis*, 623, 638
fastigata, 638
manillaris, 638
 Calyptraeid, 620, 629, 631–633
 biogeography, 631
 evolution, 632
 gastropods, 618–619
 larvae, 631
 taxonomy, 619
 Calyptraeidae, 618–619, 623, 632
 Camels, 20, 221, 857
 Camin–Sokal parsimony, 444, 446
 Campbell, Christopher S., *see* Smedmark, Jenny E. E., 374–385
 Campbell, Nick, J. H., *see* Murrell, Anna, 296–310
 Canary, 389
 Canidae, 846
 Canids, 707–708
Canis, 706
 Cannatella, David C., *see* Evans, Ben J., 794–823
 Canonical
 correspondence analysis (CCA), 687–689, 693
 ordination (CO), 687, 693
 phylogenetic ordination (CPO), 684, 687–689, 691–694
 tree, 113, 115
 variate analysis (CVA), 190, 196, 198, 491, 496
 Canonical phylogenetic ordination, Norberto P. Giannini, 684–695
Canthumeryx, 224
 Capreolini, 209
Capreolus, 209, 212, 215, 218–219
 capreolus, 210
 Caprini s.l., 209–210, 215, 225
 Capulidae, 639
 Capulids, 620, 623
Capulus ungaricus, 639
 Carabid beetles, 70, 103
 Cardillo, Marcel, *see* Bininda-Emonds, Olaf R. P., 724–729
 Cardini, Andrea, Geometry of the marmot (Rodentia: Sciuridae)
 mandible: phylogeny and patterns of morphological evolution,
 186–205
Carios capensis, 300
 Carnivore supertree, 845
 Carnivora, 571, 839–851
Carollia perspicillata, 472
 Cartesian transformation grids (CTGs), 488, 491–492, 497
 Caryophyllaceae, 820, 832
 Cassowaries, 550
Castor canadensis, 606
 Castoridae, 604–606, 608–611, 613–615
Casuarius, 389
 casuarius, 388
 Catarrhini, 738
 Catonotus, 492–493
 Causinae, 441, 453
Causus, 451
 defilippi, 441
 rhombeatus, 441
 Cattle, 206
 CB1, 604, 606–610, 613
 Cave salamanders, 501
Cavia, 649–664
 porcellus, 649, 656
 Cenozoic, 206, 631, 746
 Centipedes, 559
 Cephalocarida, 557
 Cephalophini, 209–210
Cephalophus, 209, 212, 215, 221
 dorsalis, 210
 maxwelli, 210
 Cerataphidini, 137, 139
 Ceratina, 8
 japonica, 2, 5
 Ceratinini, 1, 2, 9
Cercomegistus sp., 300
Cercopithecus
 hamlyni, 736
 nictitans, 736
Cerberus, 451
 rhynchops, 440, 446
Cermatobius, 559
 Cervidae, 206–207, 209–210, 212–213, 215, 217–222, 224
 Cervinae, 209
Cervus, 207, 209, 212, 215, 218–219
 elaphus, 210
 unicolor, 210
 Cetacea, 15–22, 209, 221, 724–729, 853–855, 857–858, 860
 Cetaceans, 853, 860
 phylogeny, 854
 Cetartiodactyla, 206, 221
Chaetodipus, 609, 614
 californicus, 606
Chalinolobus tuberculatus, 462
 Characters, 239–252, 851–852, 864
 adult, 8–9
 anatomical, 742
 autopomorphic, 865
 binary, 244, 247, 256–257, 335, 535
 chromatic, 178
 cladistic, 854
 coding, 334–351, 401, 405, 540
 complex, 245
 constant, 378
 construction, 239–241, 243, 246, 253
 convergent, 502, 510
 correlation, 131–158
 cranial, 187
 cytological, 311, 313, 326–328
 dental, 738, 853, 857, 860
 diagnostic, 323, 328, 545
 discrete, 136, 825
 distribution, 264–265
 DNA, 56, 348, 405, 509
 embryological, 311, 313, 327–328
 equally weighted, 318, 327, 358
 evolution, 56, 152, 263, 318, 328, 364, 416, 543, 647, 833
 foot, 858
 fruit, 374–375
 gap, 317–318
 genitalic, 8, 91
 homoplastic, 589, 767
 indel, 389, 823, 826, 829
 informative, 16, 378, 575, 583
 larval, 8–9, 92
 life-history, 9
 mandibular, 192
 mapping, 27, 131–158
 meristic, 506, 513
 molecular, 55, 70, 76–77, 131, 199, 209, 296, 298, 409, 738, 742
 morphological, 55, 57, 70, 72, 74, 87–88, 89, 91, 131–132, 135, 142,
 178, 181, 199, 209, 213, 217, 219, 298, 313–314, 317, 323, 327–328,
 340, 389–390, 400, 406–407, 409, 412, 420, 439, 442, 457, 474,
 501–503, 506, 509–510, 537, 539–543, 545, 559, 572, 604, 632, 738,
 742, 748, 760, 855–857

- multistate, 26, 75, 239, 255–257, 335
 osteological, 187, 760
 parasitological, 738, 742
 parsimony-informative, 4, 298, 323, 345, 446, 451, 529, 531, 773
 phylogenetically informative, 55–56, 826
 polymorphic, 505, 833
 primary, 725–726
 pupal, 8
 reductive, 245
 reweighted, 318, 327, 371
 shell, 619, 623
 skeletal, 197, 199, 222, 313, 403
 snout, 416, 621–622
 state, 87, 132, 149, 239, 245–246, 250, 253, 296, 309–310, 352, 354, 357, 400, 405, 427, 502–503, 505, 542, 760, 831, 835
 summary, 337, 342
 systematic, 243
 two-state, 236
 unambiguous, 539, 557
 uninformative, 76, 217, 575, 581, 826
 vertebral, 439
Charadrius semipalmatus, 286
Cheilea, 632, 639
 equestris, 639
 Cheirogaleidae, 711–713
Cheirogaleus, 706
 Chelicerata, 56, 296, 557
 Chevrotains, 206
 lesser Malay, 210
 Chicken, 388–389, 391, 551
Chilalictus, 24
 Chilopoda, 557
Chimeromyia, 749
 Chimpanzee, 52–53, 711–713, 715
Chinchilla laniger, 606
 Chinchillidae, 606
 Chinese
 muntjak, 210
 water deer, 210, 220
 Chipmunks, 194, 197
 Chippindale, Paul T., *see* Wiens, John J., 501–514
 Chironomidae, 430
 Chi-square tests, 307, 395, 397–398, 607
 Chloroplast
 DNA, 374–375, 382
 genome, 336
 phylogeny, 375
 Choanoflagellata, 316–317
Chordeiles acutipennis, 286
 Chromatic characters, 178
 Chromatograms, 94
Chrysops carbonarius, 756
 Cichlidae, 549, 552
 Cichlid fish, 549–553
Ciconia nigra, 286
Cirripathes, 317
 lutkeni, 316
 Cirripectida, 557
 Clades, 11, 15–17, 19–20, 23, 34, 55, 62, 70, 76–77, 79, 81, 100–101, 103, 105–106, 119, 122, 125, 140, 159, 165–167, 172, 178–179, 186–187, 220, 241, 249, 275, 278, 304–305, 318, 320, 323–324, 327, 329, 334, 344, 374–375, 377, 379, 383–384, 390, 403, 409, 439, 445–447, 452, 455, 463, 468, 494, 503, 508, 511, 529–531, 535, 542, 557, 560–561, 581, 583, 585, 590, 604, 614, 618, 623, 629, 631–632, 665, 667–669, 671–672, 688, 697, 711–712, 724, 749, 761–762, 767, 771, 776, 797, 802, 804, 807, 810–811, 827, 829, 832, 839–840, 843–847
 phylogenetic, 714
 Cladistic
 analysis, 8, 81, 253, 403, 541, 572, 738
 characters, 854
 information content (CIC), 241
 hypothesis, 92
 relationship, 70–71
 parsimony, 356, 362, 365
 Cladistics, 63, 70–88, 520
 molecular, 187
 Cladogenesis, 641–642, 644–645, 647, 746, 757, 761, 834
 Cladogenetic model, 641
 Cladogram, 5, 70, 73, 76–77, 79, 81, 83–84, 195, 255–258, 272, 359, 362, 406–407, 520, 525–526, 548–549, 684, 722, 742, 802
 coding, 255–259
 molecular, 187, 742
Claravis pretiosa, 38
 Clarifying Aetosaur phylogeny requires more fossils, not more trees—reply to Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), Andrew B. Heckert and Spencer G. Lucas, 253–254
Clarkia, 347
Clathrina, 314–315, 320, 329, 332–333
 cerebrum, 316
 clathrus, 329
 Clathrinida, 314, 318
 Clathrinidae, 314
 Clayton, Dale H., *see* Johnson, Kevin P., 37–47
Climacoceras, 224
 Clocklike trees, 233
 Clock model (*see* molecular clock)
 Cluster analysis, 194, 205, 497, 722
C-mos, 758, 762, 764
C-myc, 386–402
 Cnidaria, 311, 316–317, 320, 430, 520
Coahomasuchus, 241, 243, 250–251, 254
 Coding, 239–252, 851–852
 cladograms, 255–259
 indels, 444, 575
 Codons/codon positions, 3, 5, 7, 10, 20, 28, 391
 Cody, Jeannine H., *see* Rodman, James E., 428–435
 Coelenterates, 518
 Coenagrionidae, 89, 91–92
Coenomyia ferruginea, 756
 Coevolution, 37–47
 Coleoptera, 430
 Collin, Rachel, Phylogenetic relationships among Calyptraeid gastropods and their implications for the biogeography of marine speciation, 618–640
 Colonization, 70–88
Coluber, 451
 constrictor, 440
 Colubridae, 440, 456
 Colubrid snakes, 758
 Colubrinae, 439–440, 447, 452, 455, 457
 Colubrini, 457
 Colubroidea, 439–459
Columba
 guinea, 38
 livia, 38, 286
Columbicola, 37–39, 41–42, 45
 adamsi, 38
 bacillus, 38
 baculoides, 38
 columbae 1, 38

- columbae* 2, 38, 42
exilicornis, 38
extinctus, 38, 42
gracilicapitis, 38, 42
gymnopelidae, 38
macrourae 1, 38, 46
macrourae 4, 38
macrourae 5, 38
n. sp., 38
passerinae 1, 38, 42
passerinae 2, 38
theresae, 38, 46
timmermanni, 38
veigasimoni, 38
xavieri, 38
Columbiformes, 37–47
Columbina, 46
inca, 38, 46
passerina, 38, 46
Coluria, 374, 381–383
geoides, 374, 376, 381–382
Colurieae, 374, 377–378, 382
Combinatorics of tandem duplication trees, Olivier Gascuel, Michael D. Hendy, Alain Jean-Marie, and Robert McLachlan, 110–118
Combined data, 55–65
analysis, 96
evidence, 403–422
tree, 612
Combined support for wholesale taxic atavism in gavialine
Crocodylians, John Gatesy, George Amato, Mark Norell, Rob DeSalle, and Cheryl Hayashi, 403–422
Combining data, 528–538
Comparative
biology, 55–65
method, 23–36
Comparing bootstrap and posterior probability values in the four–taxon case, Michael P. Cummings, Scott A. Handley, Daniel S. Myers, David L. Reed, Antonis Rokas, and Katarina Winka, 477–487
Comparison of likelihood and Bayesian methods for estimating divergence times using multiple gene loci and calibration points, with application to a radiation of cute-looking mouse lemur species, Ziheng Yang and Anne D. Yoder, 705–716
Complex characters, 245
Component coding, three-item coding, and consensus methods, David M. Williams and Christopher J. Humphries, 255–258
Compsomelissa, 1–2, 4–5, 7–12
borneri, 2
Computer simulation (*see also* simulation and Monte Carlo simulation), 528–538, 696–704
Computing the uniform component of shape variation, F. James Rohlf and Fred Bookstein, 66–69
Conflict, 839–851
Congruence, 37, 55–65, 193, 329, 539–548, 555, 607, 613, 669, 717, 865–868
Conifers, 784
Consensus, 255–259, 849–851
efficiency (CE), 241
trees, 55, 128, 140, 241, 251, 255, 258, 344, 450, 531, 850, 855, 865–868
Consistency index, 3, 5–6, 10, 28, 76, 240, 247–248, 252, 304, 339, 345–346, 348, 392, 396, 577, 581–583, 762, 764, 826
Constrained tree, 16–18, 20, 22, 407, 761, 807, 847
Continuous
jackknife function analysis (CJF), 55–58, 61–63
time Markov model, 131–134, 147, 149, 651
time Markov process, 179, 644
Control region (CR), 460–476
Conus, 619
Convergence, 501–514, 614
sciurid, 185–205
Convergent
characters, 502, 510
evolution, 296, 497
Conway, Lindsay, *see* Danforth, Bryan N., 23–36
Co-occurrence, 271–280
Cooper, Steven J. B., *see* Schwarz, Michael P., 1–14
Copath
analysis, 41
tree, 45
Cophylogenetic analysis, 737
Cophylogeny, 735–744
Coracias caudata, 286
Coragyps atratus, 388
Corbiculate bees, 23, 33
Correspondence analysis, 687
Corroboration, 352–367
Cosmiomma, 305–306
Cospeciation, 37–47
Coturnix pectoralis, 286
Cows, 210
Crangonyctidae, 430
Cranial characters, 187
Credibility regions (CR), 390, 394
Crepidula, 618–640
aculeata, 636
adunca, 637
aff. depressa, 636
aff. marginalis, 637
aff. onyx, 638
aff. williamsi, 637
arenata, 637
argentina, 636
atrasolea, 636
cerithicola, 637
cf. aculeata, 636
cf. aculeata Cape Verde, 623, 636
cf. aphysioides, 637
cf. incurva, 637
cf. nivea, 637
cf. onyx, 638
cf. perforans, 637
complenata, 629, 631, 637
convexa, 629, 637
coquimbensis, 636
costata, 636
dd. onyx, 638
depressa, 636
excavata Mexico, 620, 632, 637
excavata Peru, 620, 637
fimbriata, 636
fornicata, 619, 632, 637
gibbosa, 637
grandis, 632, 637
gravispinosa, 636
immersa, 636
incurva, 632, 637
lessoni, 636–637
maculosa, 638
marginalis, 638
monoxyla, 636
naticarum, 638
navicula, 638

- norrisarum*, 638
 n. sp., 638
 n. sp. dd., 638
 n. sp. pt., 638
nummaria, 637
onyx, 619, 631, 638
perforans, 637
philippiana, 623, 637
plana, 637
porcellana, 629, 638
protea, 637
 s.l., 637
 s.s., 623, 631–632, 636
striolata, 637
unguiformis, 637
ustulatulina, 638
williamsi, 623, 637
Crepipatella, 623, 631, 636
capensis, 631–632, 636
dilatata, 632, 636
fecunda, 632, 636
lingulata, 636
 n. sp., 636
 Cretaceous, 23, 33, 403, 419, 550–552, 631, 746, 753, 861
 early-, 548
 mid-, 549, 552
 late-, 419, 549–552, 753
 upper-, 753
Cretaformia, 749
 Crickets, 70, 833
 Crocodile, 285
 African slender-snouted, 388
 Crocodylians, 240, 283, 286, 293, 389, 403
 Crocodylia, 386–402, 403–422
 Crocodyliformes, 386, 415
 Crocodylidae, 386, 392, 394
 Crocodylinae, 403, 405, 408–409, 412, 415, 417, 419
 Crocodyloidea, 391
Crocodylus, 386, 391, 396
 cataphractus, 388, 391, 405, 408–409, 415, 417
 intermedius, 405, 408, 417
 New World, 409
 niloticus, 285
 rhombifer, 405, 408
 Crotalinae, 441, 447, 453
Crotalus, 447, 451
 horridus, 441
 viridis, 441
Crotaphopeltis, 447, 451, 457
 hotamboeia, 440
Crucibulum, 618–640
 auricula, 638
 cf. *personatum*, 638
 cf. *serrata*, 638
 concamaratum, 623, 638
 lignarum, 638
 radiata, 638
 scutellatum, 638
 spinosum, 638
 tenuis, 638
Crunomys, 812
 Crustacea, 430, 432, 557
Crystallaria, 496–498
 asprella, 491, 496–499
 Ctenodactylidae, 605–606
 Ctenohystric, 605, 615
Ctenonomia, 24
 Ctenophora, 311, 316–318, 320, 327–328
Cuculus pallidus, 286
Culex pipiens, 745
 Cumaceans, 430
 Cummings, Michael P., Scott A. Handley, Daniel S. Myers, David L. Reed, Antonis Rokas, and Katarina Winka, Comparing bootstrap and posterior probability values in the four-taxon case, 477–487
 Culicomorpha, 754
 CXCR4/CCR5 coreceptor, 649–664
 Cycads, 784
Cyanoramphus, 471
Cyclorrhapha, 746, 749–751, 753
Cymatium parthenopeum, 640
Cynomys, 189, 194
 leucurus, 204
 ludovicianus, 189, 204
Cypraea spurca, 640
 Cyprinodontiformes, 548–459
Cypripedium, 346
 Cytogenetic analysis, 374
Cytochrome b, 2–5, 7–8, 10–11, 16–18, 20, 22, 39, 41–42, 161, 163–166, 171, 179, 181, 186, 193, 195, 198–199, 206–207, 209–212, 215, 297, 443, 445–447, 451, 455, 471–472, 488, 504, 508, 571–593, 605, 657, 659, 677, 680–681, 705–706, 709, 727, 861
 Cytochrome oxidase I (COI), 1–5, 7–8, 10–11, 23, 26–30, 32, 37, 39, 41–42, 45, 72–73, 139, 297–298, 557, 559, 618–640, 657–659, 674
 Cytochrome P-450, 206, 209–212, 219
 Cytochrome oxidase II, 33, 94, 95, 139, 207, 705–706, 709
 Cytological characters, 311, 313, 326–328

Dactylanthus taylorii, 473
Dalloapula, 9
Damaliscus, 209, 212, 215, 218
 pygargus, 210
 Damselflies, 70, 833
 Megalagrion, 89–109, 833
 Danforth, Bryan N., Lindsay Conway, and Shuqing Ji, Phylogeny of eusocial *Lasioglossum* reveals multiple losses of eusociality within a primitively eusocial clade of bees (Hymenoptera: Halictidae), 23–36
 Darters, 488–489, 494, 498
Dasypeltis, 447, 451, 457
 medici, 440
 scabra, 440, 459
 Data
 combination, 439–459
 conflict, 386–402
 non-independence, 724–729
 partitions, 386–402
 visualization, 55–65
Daubentonia, 706, 712
 Dayrat, Benoît, Roots of phylogeny: how did Haeckel build his trees?, 515–527
 DeBry, Ronald W., Identifying conflicting signal in a multigene analysis reveals a highly resolved tree: the phylogeny of Rodentia (Mammalia), 604–617
 Decay
 analysis (see also Bremer support), 12, 128, 243, 407
 index, 128, 336, 377, 381, 842
 Decision theory, 674–683
 Deduction, 259–271
 Deep sea mollusks, 430
 Deer,
 black-tailed, 210

- Chinese water, 210
 European roe, 210
 mouse, 206, 854, 857
 musk, 206–207, 210, 219–220
 red, 210
 Dei-Cas, Eduardo, *see* Hugot, Jean-Pierre, 735–744
 Deletion (*see* indel)
 Delma, 757–758, 762, 764, 767, 770–771, 773, 775, 777–778
 (*Aclys*) *concinna*, 759, 762, 764, 770, 775, 777–778
 australis, 759, 775, 778
 borea, 759, 775, 778
 borea haroldi, 759
 butleri, 759, 770, 778
 elegans, 758, 775
 fraseri, 757–759, 762, 775–778
 fraseri fraseri, 762, 775–776
 fraseri petersoni, 762, 775–776
 grayii, 757, 759, 762, 775–778
 impar, 759, 775, 778
 inornata, 759
 labialis, 759, 764, 770, 775, 777–778
 mitella, 759, 762, 770
 molleri, 759, 775, 778
 nasuta, 759, 778
 pax, 759, 762, 775
 petersoni, 757, 759, 777
 plebeia, 758, 775
 tincta, 759, 775, 778
 torquata, 759, 775, 778
 Demanche, Christine, *see* Hugot, Jean-Pierre, 735–744
 Demospongiae, 311, 314, 316–317, 320, 328–330
 Dendrelaphis, 447, 451, 457
 calligastrea, 440
 pictus, 440
 Dendroaspis, 447, 451, 456
 polylepis, 441
 sp., 441
 Dendrogram, 192–193, 195, 199
 mandibular, 195, 197
 UPGMA, 193
 Dendroica, 776
 Dental characters, 738, 853, 857, 860
 Dentition, 16
 Dermacentor, 299, 305–306, 310
 andersoni, 299
 marginatus, 299
 reticulatus, 299
 s.l., 299, 305
 variabilis, 299
 DeSalle, Rob, *see* Gatesy, John, 403–422
 Desmotosuchus, 241–243, 248–252, 254
 haplocerus, 254
 de Queiroz, Kevin and Steven Poe, Failed refutations: further
 comments on parsimony and likelihood methods and their
 relationship to Popper's degree of corroboration, 352–367
 DHPS, 735–738, 740, 742
 Diacodexis, 858
 Diagnostic characters, 323, 328, 545
 Dialictus, 24, 28, 32–33
 Diaphanoeca grandis, 316
 Diapsida, 239, 253
 Dicotyledons, 524
 Dicrocerus, 224
 Didelphis, 706, 708
 virginiana, 285, 709
 Diesmos, Arvin, *see* Evans, Ben J., 794–823
 Dinosauria, 839
 Dinosaurs, 253
 Diospyros sandwichensis, 820
 Dipodomys, 614
 spectabilis, 606
 Dipsas, 447, 451
 Dinodon, 447, 451
 semicarinatum, 440, 445
 Diogmites sp., 756
 Diphyllidae, 430
 Diplodactylinae geckos, 758, 760
 Diplodactylus, 757
 damaeus, 758–759
 tessellatus, 758–759
 Dipodidae, 604–606, 608, 613–614
 Dipsas catesbyi, 441
 Diptera, 430, 557, 745–756
 Dipus, 614
 sagitta, 606
 Dirichlet
 density, 711
 distribution, 607, 652, 655, 749
 Discete characters, 136, 825
 Discrete-gamma
 distribution (*see also* gamma distribution), 165, 445, 666, 698, 700,
 703
 estimate (DGE), 595–596, 599–601
 probability estimate (DGPE), 596–602
 Dispersal, 549–553, 794–819
 Distance analysis, 334, 343
 Divergence time, 705–716, 745–756
 DNA, 2–3, 74, 94, 110, 166, 212, 221, 298, 306, 334, 340, 342, 345, 370,
 382, 389, 391, 408–409, 419, 506, 509, 551–552, 606, 620, 725, 727,
 760, 857
 analysis, 551
 avian, 389
 characters, 56, 348, 405, 509
 chloroplast, 374–375, 382
 evolution, 443
 extraction, 26, 38, 462, 736
 fingerprints, 404
 genomic, 161, 315–316, 443, 550, 566, 574, 606, 758, 823
 hybridization, 550–551, 573
 isolation, 375
 mitochondrial, 3–4, 7, 12, 37, 48, 52, 89, 94–101, 103, 159–185, 296,
 386–387, 395, 407–408, 415, 418, 423, 439–459, 460–461, 464, 471,
 474, 503–504, 506, 508–509, 548, 551, 571–572, 583, 586, 589, 656,
 674, 677, 680–681, 735, 757–758, 760–762, 764, 767, 770–771, 773,
 775–776, 794, 798, 801, 808, 810–812
 nuclear, 37, 89, 95, 99, 103, 221, 374–375, 382–383, 407–408, 415, 418,
 423, 509, 571–593, 735, 757–758, 760–762, 764, 767, 770–771, 773,
 775–776
 plasmid, 316
 Pneumocystis, 735
 polymerase, 606
 sequence, 38–39, 49, 79, 133, 135, 140, 159, 209, 296, 298, 304,
 334–336, 339–340, 344, 346, 348, 387, 396, 400, 404–405, 418–419,
 477, 479, 488, 571, 574, 604, 619, 621, 623, 632, 651, 666, 735, 823,
 857
 sequence data, 23, 509, 530–531, 539, 542, 545, 674, 708, 758, 773, 849
 substitution, 135, 140, 575, 644, 647, 824
 substitution model, 444
 tree, 770, 773, 775
 Dog, 591, 711–713
 Dolichopodidae, 756
 Dollosuchus, 416

- Dolphins, 501
 Donnellan, Stephen, *see* Jennings, W. Bryan, 757–780
 Douzery, Emmanuel J. P., *see* Hassanin, Alexandre, 206–228
 Doves, 37, 41, 45
 New World ground, 39
 Draco, 810
 Dremotherium, 224
 Dromaius novaehollandiae, 286
 Drosophila, 2, 89, 383, 557, 745–746, 751, 753–754, 833
 melanogaster, 94, 745, 756
 yakuba, 94, 298
 Drosophilidae, 756
 Drosophilid flies, 70
 Dryadeae, 374
 Dryomys nitedula, 606
 Dwarf caiman, 388
- Echinodermata, 296, 520
 Echinometra sea urchins, 631
 Echmoidea, 288
 Echthralictus, 24
 Ecology, 430, 515, 633, 684
 Ecomorphology, 498, 684–695
 EF-1 α , 2–4, 8, 10–11, 23, 26–30, 32, 39, 41–42, 94–99, 101–102, 557, 596, 598, 677, 680
 Effect of phylogeny on interspecific body shape variation in darters (Pisces: Percidae), J. Michael Guill, David C. Heins, and Craig S. Hood, 488–500
 ϵ -globulin, 738
 Eichhornia,
 azurea, 154, 156
 crassipes, 154, 156
 diversifolia, 154, 156
 heterosperma, 154, 156
 meyeri, 154, 156
 paniculata, 154, 156
 paradoxa, 154, 156
 sp., 154, 156
 Eigenvectors, 69, 694
 Eigenvalues, 67
 Eira, 577, 581, 590
 barbara, 573, 581
 Elaphe, 451
 flavolineata, 440
 obsoleta, 440
 Elapidae, 439, 441, 443, 447, 455–456
 Elapinae, 441, 447
 Elapsoidea, 451
 nigra, 441
 Elephants, 794
 African, 591
 Ellipsoptera, 776
 Embriological characters, 311, 313, 327–328
 Empididae, 754
 Empidoidea, 746, 751
 Empis, 754
 Emu, 550–551
 Enallagma, 94
 geminatum, 94
 Endemism, area of, 271–280, 717–723
 Enhydra, 590
 lutris, 573, 583, 590
 Enhydris, 446, 451
 enhydris, 440
 plumbea, 440
 Eocene, 222, 224, 403, 419, 857–858
 early-, 795
 late-, 222, 224
 mid-, 218, 222, 224, 550
 Eoconodon, 858
 Eogavialis, 401
 Eotragus, 225
 Ephemeroptera, 106
 Ephydatia muelleri, 316
 Equal-rates Markov (ERM), 229–238
 model, 57
 trees, 233
 Equally
 parsimonious tree, 528, 530
 weighted character, 318, 327, 358
 weighted parsimony, 620–621
 weighted tree, 211
 Equus, 706, 857
 Eremoneura, 746, 750–751, 754
 Erethizon dorsatum, 606
 Erethizontidae, 606
 Eriksson, Torsten, *see* Smedmark, Jenny E. E., 374–385
 Erixon, Per, Bodil Svennblad, Tom Britton, and Bengt Oxelman, Reliability of Bayesian posterior possibilities and bootstrap frequencies in phylogenetics, 665–673
 Ermine, 573
 Erythrocoma, 374
 triflora, 374, 376, 379, 383
 Esastigmatobius, 559
 Escherichia, 660
 coli, 209, 383, 659
 Estimation of rates-across-sites distributions in phylogenetic substitution models, Edward Susko, Chris Field, Christian Blouin, and Andrew J. Roger, 594–603
 Etheostoma
 blennioides, 497, 500
 blennius, 491, 500
 caeruleum, 498, 500
 coosae, 492, 500
 crossopterum, 491, 500
 exile, 498, 500
 flabellare, 500
 fusiforme, 498, 500
 inscriptum, 497, 500
 jordani, 492, 498, 500
 juliae, 491–492, 498
 lynceum, 494, 497, 500
 nigrum, 494, 497–498, 500
 olmstedii, 491–492, 494, 500
 proeliare, 491, 500
 punctulatum, 500
 radiosum, 493–494
 simoterum atripinne, 492, 494
 simoterum simoterum, 492, 494
 s.l., 491, 497, 500
 spectabile, 500
 s.s., 492–493, 500
 stigmaeum, 497–498, 500
 whipplei, 491, 493–494, 498, 500
 zonale, 491, 494, 500
 Etheostomatinae, 488
 Eubacteria, 659
 Eucalyptus trees, 777
 Euclidean
 distances, 67, 337, 676
 geometry, 67
 space, 66–67

- Euconolylops*, 8–9
Eugeum, 383
Eulemur, 706
 macaco, 736
Eulonchus sp., 756
Eumeryx, 224
 Eumetazoa, 311, 318, 327, 330
 Eumycota, 316–317
Eurycea, 501–514
 bislineata, 505
 chisholmensis, 505, 513
 latitans, 504–506, 509, 513
 longicauda, 505
 multiplicata, 504–505, 513
 nana, 505, 509, 513
 naufragia, 504–506, 513
 neotenes, 505, 509, 513
 pterophila, 504–505, 509, 513
 quadridigitata, 505
 rathbuni, 501, 504–506, 508–510, 513–514
 robusta, 504, 509
 sosorum, 505, 509, 513
 sp., 509, 513
 tonkwae, 504–505, 513
 tridentifera, 501, 504–506, 508–510, 513–514
 trogodytes, 504–505, 508–509, 513
 wallacei, 505, 509,
 waterlooensis, 504, 509
 wilderae, 505
 European roe deer, 210
 Eusocial evolution, 23
 Eusociality, 12–13, 23–36
Eutamias, 205
 sibiricus, 189, 205
 Eutherian mammals, 249, 585, 860
Euthocodon arambourgi, 417
 Euthyneura, 328
 Evans, Ben J., Rafe M. Brown, Jimmy A. McGuire, Jatna Supriatna,
 Noviar Andayani, Arvin Diesmos, Djoko Iskandar, Don J.
 Melnick, and David C. Cannatella, Phylogenetics of fanged
 frogs: testing biogeographical hypotheses at the interface of
 the Asian and Australian faunal zones, 794–823
 Evans, Rodger C., *see* Smedmark, Jenny E. E., 374–385
 Even-toed ungulates, 724
 Evolution, 1–2, 12, 20, 28, 62, 89, 92, 101–103, 126, 149, 165, 233,
 239–252, 254, 296–310, 311–333, 349, 368, 374, 377, 383, 451, 511,
 515–527, 549, 551, 573, 576, 645, 658, 660, 725, 735, 740, 851–852
 calyptraeid, 632
 character, 152, 263, 318, 328, 364, 416, 543, 647, 833
 convergent, 296, 497
 DNA, 443
 eusocial, 23
 HIV, 649–664
 island, 820–841
 mammalian, 206
 minimum, 121, 287, 478, 489, 579–580
 molecular, 38, 41, 591, 594–603
 morphological, 186–205, 404, 542
 parallel, 296, 307
 protein, 602
 punctuational, 831
 rate of, 602, 641, 646–647, 677, 746, 833
 reticulate, 374–385
 social, 1–14, 23–36
 Evolution of the strategy of long-branch subdivision to improve the
 accuracy of phylogenetic methods, Steven Poe, 423–427
 Evolutionary
 biology, 501–502, 515, 565
 branches, 849
 characters, 404
 character transformation, 354
 convergence, 853
 divergence, 119, 572
 ethnology, 684
 history, 197, 328, 417, 423, 474, 518, 526, 861
 hypotheses, 159
 model, 126, 352, 364, 377, 615, 665, 668, 672, 692, 696–697, 705, 727,
 781–782, 788, 791–792, 801, 829, 850
 pattern, 852
 plasticity, 329
 processes, 89, 211, 861
 radiations, 70, 89, 207
 rate, 218, 287, 396–397, 401, 700, 705–716, 748, 753
 studies, 727
 theory, 264–265, 515, 524–525
 trees, 526, 602
 Evylaeus, 24, 32–33
Exoneura, 1, 12
 angophorae, 2
 robusta, 2
Exoneurella, 1, 7–8, 10–11, 32
 eremophila, 32
 setosa, 2, 32
 tridentata, 2
 Exoneuridia, 2, 10
 Exoneurines, 1, 9–10
 Exons, 376, 383, 391–392, 394
 External transcriber spacer (ETS), 820–841
 Failed refutations: further comments on parsimony and likelihood
 methods and their relationship to Popper's degree of
 corroboration, Kevin de Queiroz and Steven Poe,
 352–367
Fallugia, 374
 False discovery rate (FDR), 783, 790
 Falsification, 259–271
 Fanged frogs, 794
Farancia, 447, 451, 455
 abacura, 441
Farrea occa, 316
Fejervarya
 limnocharis, 799, 815
 vittigera, 799
 Felidae, 845–846
 Felids, 707–708
Felis, 706
 Felsenstein
 model, 463, 703, 708–709, 711–715, 748, 850
 zone, 229, 426, 641–648, 675
 FES, 574–577, 581, 583
 FIB7, 39, 41–42
Ficus carica, 349
 Field, Chris, *see* Susko, Edward, 594–603
 Filicinae, 524
 Fin whale, 210
 Fishes, 286, 491, 524, 619
 aplocheiloid, 548–553
 cichlid, 549–553
 cyprinodontiform, 549
 euryhaline, 548
 freshwater, 548–549
 Fisher, 573

- Fishes and birds: Gondwana life rafts reconsidered, John C. Briggs, 548–553
- Fitch
 algorithm, 234
 parsimony, 444, 446
- Fitch–Margoliash methods, 290–291, 293
- Flatworms, 307
- Flies,
 blow, 745
 drosophilid, 70
 flower, 745
 fruit, 557, 745
 hippoboscid, 37
 horse, 745–746
 house, 745
 snipe, 745–746
 soldier, 746
 stalk-eyed, 32
 xylophagid, 745
- Flower flies, 745
- Flowers, 336, 341, 343, 346–347, 349, 753
- Flowering plants, 820
- Foot characters, 858
- Fossils, 23, 48, 61, 253–255, 264, 377, 382, 403–422, 521, 524, 528–538, 614, 631, 707, 711, 713, 746–747, 751, 761, 835, 853, 855
 amber, 746
- Fossombroniinae, 430
- Fossorella* sp., 639
- Four o'clocks, 334–351
- Fractional weighting, 258
- Fragrance, 334–351
- Frequent inconsistency of parsimony under a simple model of cladogenesis, John P. Huelsenbeck and Katherine M. Lander, 641–648
- Frogs, 287, 334, 794
 fanged, 794
- Fruit, 383
 characters, 374–375
 flies, 557, 745
- F*-test, 687
- Fungi, 317, 429–430
- F*-values, 692
- G6PD, 738, 742
- Galago*, 706
- Galbula pastazae*, 286
- Galictis*, 590
vittata, 573–574, 576, 581, 589–590
- Galliformes, 551
- Gallirex porphyreolophus*, 286
- Gallus gallus*, 286, 396, 405, 408
- Gamma
 correlation, 727
 distribution (*see also* discrete-gamma distribution), 4, 42, 50, 95, 135–136, 293, 318, 377, 390, 463, 506, 576, 594–595, 601, 606, 608, 702, 709, 711, 748, 750, 801, 823
 model, 594–603, 750
 shape parameter, 621, 711
- γ -fibrinogen, 207
- Gammarus minus*, 509
- Gamopetalae, 524
- Gap character, 317–318
- Gascuel, Olivier, Michael D. Hendy, Alain Jean-Marie, and Robert McLachlan, Combinatorics of tandem duplication trees, 110–118
- Gascuel, Olivier, *see* Guindon, Stéphane, 696–704
- Gastrea, 311
- Gastreaeades, 520
- Gastropoda, 328
- Gastropods, 619, 631
 Calyptraeid, 618–619
- Gastropyxis*, 447, 451
smaragdina, 440
- Gatesy, John, George Amato, Mark Norell, Rob DeSalle, and Cheryl Hayashi, Combined support for wholesale taxic atavism in gavialine Crocodylians, 403–422
- Gatesy, John, *see* O'Leary, Maureen A., 853–866
- Gavial, 403–422
- Gavialidae, 398–399
- Gavialinae, 403, 416–418, 420
- Gavialis*, 386, 388, 391–392, 394, 396–400, 403
gangeticus, 386, 388, 390, 398, 403, 405–406, 408–409, 412, 415–419
- Gavialoidea, 401
- Gavialosuchus*
americanus, 412
eggenbergensis, 417
- Gazella*, 209, 212, 215
granti, 210, 221
thomsoni, 210
- Gazelle, 210
- Geckos, 758
 diplodactylid, 758, 760
- Geinae, 374–379, 382–384
- Gekkonidae, 757
- Gelocidae, 206
- Genealogical trees, 515, 517, 521, 524, 526
- Generalized least squares, 491
- General-time-reversible (GTR) model, 3–4, 7–8, 18–19, 26, 28–30, 42, 95, 97, 165–166, 212, 286, 318, 390, 392, 396, 445, 463, 576, 579, 585, 608–611, 613, 666–669, 674, 677, 679, 681–682, 737, 762, 784, 801–802, 825, 831, 834
- Genes, 296, 605, 609, 612, 652, 658
 A2AB, 605, 614–615
 A6, 674
accD, 336
 α -lactalbumin, 206, 209–212, 215, 218–220
 APOB, 574–577, 581, 583
atp6, 657, 659
atp8, 656–659
 β 2-microglobulin, 738
 β -casein, 207
 β -tubulin, 596, 598–600
 BRCA1, 605, 613–615
 CB1, 604, 606–619, 613
 CHRNA1, 574–577, 581, 583
 cMDH, 596, 598
c-mos, 758, 762, 764
c-myc, 386–402
 COI, 2–5, 7–8, 10–11, 23, 26–30, 32, 37, 39, 41–42, 45, 72–73, 139, 297–298, 557, 559, 618–640, 674
 COI-1, 1, 5, 657–659
 COI-2, 5, 657–659
 COI-3, 5, 657–659
 COII, 33, 94, 95, 139, 207
Cytochrome b, 2–5, 7–8, 10–11, 16–18, 20, 22, 39, 41–42, 161, 163–166, 171, 179, 181, 186, 193, 195, 198–199, 206–207, 209–212, 215, 297, 443, 445–447, 451, 455, 471–472, 488, 504, 508, 571–593, 605, 657, 659, 677, 680–681, 705–706, 709, 727
 cytochrome P-450, 206, 209–212, 215, 219
 DHPS, 735–738, 740, 742
 EF1 α , 2–4, 8, 10–11, 23, 26–30, 32, 39, 41–42, 94–99, 101–102, 557, 596, 598, 677, 680

- ϵ -globulin, 738
 eRF3, 596, 598
 FES, 574–577, 581, 583
 FIB7, 39, 41–42
 γ -fibrogen, 207
 G6PD, 738, 742
 GBSSI, 374–385
 GHR, 605, 613–615
 H3, 73, 557
 HBS1, 598
 HMG-CoA, 596, 598
 homeobox, 311
 HSP70, 596, 598, 600
 HSP90, 598
 IRBP, 604–614
 ITS, 73, 76–77, 298, 336, 370–371, 382
 κ -casein, 206–207, 209–212, 215
 lactoferrin, 206, 209–212, 215, 218, 220
 mitochondrial, 5–6, 10, 16, 32, 41, 74, 439, 451, 460, 462, 510, 572, 587, 649, 651, 656, 674, 677, 680–681, 725
 mtLSU-rRNA, 735–738, 740–741
 mtSSU-rRNA, 735–738, 740
 NADH, 460
 ND1, 73–74, 76, 657, 659
 ND2, 460, 462–468, 471, 657, 659, 758–759, 762, 777
 ND3, 657, 659
 ND4, 161, 163–165, 167, 181, 440, 443, 445–447, 451, 453, 455, 657, 659
 ND4l, 657, 659
 ND5, 656–657, 659
 ND6, 406, 657–659
 ndhF, 140
 nrDNA, 336
 nuclear, 16, 28–29, 32, 41, 74, 375, 386, 572
 opsin, 28–30
 PcP, 735–744
Pneumocystis, 742
 protein-coding, 23, 39, 74, 94, 297–298, 307, 358, 396, 405, 445, 460, 471, 506, 604, 609, 612, 656, 682, 705–706
psaA, 785, 787, 791
psaB, 785, 787–788, 791
 RAG-1, 405, 408
 RAG-2, 604, 606–610, 613–614
rbcl, 140, 336, 379, 696–704
 rDNA, 406, 408, 808
 RHO1, 574–577, 587, 583
 RNA polymerase II, 557
 rRNA, 283–284, 297, 307
 single-copy, 375
 tree, 63, 179, 193, 258, 374, 446, 583–584, 588, 725, 727, 776
 tRNA, 2, 52, 94–95, 139, 296–298, 300–301, 303–308, 383, 396, 406, 460, 465–466, 725, 727, 758, 762, 798
 tRNA^{Asp}, 674
 tRNAL, 73, 76
trnL-trnF, 380–382, 384
 U2 snRNA, 557
v-myc, 388
 vWF, 605, 614–615, 738
 Genes, categories, and species: the evolutionary and cognitive causes of the species problem, Hanno Sandvik (rev.), 565–567
 Genetalic
 characters, 8, 91
 morphology, 71, 82
 Genetic analysis, 197
 Genomes, 59, 110, 126, 286, 296, 301, 550, 591, 754
 chloroplast, 336
 mitochondrial, 56, 58–59, 238, 296–297, 301, 303, 307, 309, 462–463, 478, 577, 706
 nuclear, 374–375, 571, 604
 sequencing, 119, 126
 Genomic DNA, 161, 315–316, 443, 550, 566, 574, 606, 758, 823
 Genotypes, 735
 Pneumocystis, 735
 Geometric morphometrics, 66–69, 186–205, 488–500
 Geometry of the marmot (Rodentia: Sciuridae) mandible: phylogeny and patterns of morphological evolution, Andrea Cardini, 186–205
 Geomyidae, 606, 608–609
 Geomyoidea, 604–605, 608–611, 613–615
Geonoma, 346
Geotrygon montana, 38
Geum, 374–385
 andicola, 374, 381–382
 heterocarpum, 374, 376, 379–384
 montanum, 374, 376, 379, 382–384
 reptans, 374, 376, 383–384
 rivale, 374, 376, 379, 382–384
 schofieldii, 381
 urbanum, 374, 376, 379, 382–384
 vernium, 374, 376, 379, 382–384
 Gharials, 386, 398–399
 false, 388
 true, 388, 396
 GHR, 605, 613–615
 Giannini, Norberto P., Canonical phylogenetic ordination, 684–695
 Gibbon, 52–53
 Gillespie, Rosemary G., *see* Hormiga, Gustavo, 70–88
Ginko, 784
Giraffa, 207, 209, 212, 218, 221–222, 224
 camelopardalis, 210
 Giraffes, 206, 210, 269
 Giraffidae, 206–207, 209–210, 212–213, 218–222
 Giribet, Gonzalo, Stability in phylogenetic formulations and its relationship to nodal support, 554–564
 Gittleman, John L., *see* Bininda-Emonds, Olaf R. P., 724–729
 Gliridae, 604–606, 608–609, 611, 614–615
 Global clock, 711–712
Gloma fuscipennis, 756
Glossalictus, 24
 Gnetaleans, 753
 Gnetales, 781, 784, 787–788
 Goodness-of-fit test, 464, 471, 674
 Goose magpie, 388
 Gorilla, 52–53, 706–708
 Gower, David J., *see* Harris, Simon R., 239–252, 851–852
 Gower, David J., *see* Wilkinson, Mark, 127–130
Gradicrepidula, 619, 623, 632
Grantia, 314–315, 320, 322, 332
 compressa, 316, 329
 Grantiidae, 314, 322–323
Grantiopsis, 314–315, 320, 322, 324, 333
 sp., 316
 Granule-bound starch synthase (GBSSI) (*see also* waxy), 374–385
Graphiurus murinus, 606
Grayia, 447, 451, 457
 ornata, 440
 Greater grison, 573
 Grenyer, Richard, *see* Bininda-Emonds, Olaf R. P., 724–729
 Grey-headed chachalaca, 388
 Ground spiders, 430
 Ground squirrels, 186, 197

- Grus canadensis*, 286
Gryposuchus colombianus, 412
 Guill, J. Michael, David C. Heins, and Craig S. Hood, Effect of phylogeny on interspecific body shape variation in darters (Pisces: Percidae), 488–500
 Guillot, Jacques, *see* Hugot, JeanPierre, 735–744
 Guindon, Stéphane and Olivier Gascuel, Simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood, 696–704
 Guinea pig, 656–657, 659
Gulo, 577, 581, 590
gulo, 573–574
 Gymnophthalmidae, 689, 691
 Gymnospermae, 518, 524, 781, 787–788
- H3, 73, 557
Haasiella, 557
 Habib, Michael, *see* BinindaEmonds, Olaf R. P., 724–729
Hadropterus, 491, 493
 Hadziidae, 430
 Haemaphysalinae, 299
Haemaphysalis, 299, 304–305, 309–310
humerosa, 299
inermis, 299
leporispalustris, 299
longicornis, 299
 ssp., 310
Haideotriton, 505, 509
wallacei, 508
 Handley, Scott A., *see* Cummings, Michael P., 477–487
Halictus, 23–24, 32
rubicundus, 33
 Halictidae, 23
 Halictid bees, 23–24, 33
 Halictini, 24
Haliotis, 631
Halterapis, 1–2, 4–5, 8–10, 12
nigrinervis, 2, 12
Hapalemur, 706
griseus, 736
 Haplorhini, 738
 Haplotype, 5, 71, 73, 159–160, 165–166, 171–172, 180–181, 465, 467, 776
 tree, 159, 165–166
 Hard ticks, 296–298, 300–301, 304, 309
 Harris, Simon R., David J. Gower, and Mark Wilkinson, Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), 239–252
 Harris, Simon R., David J. Gower, and Mark Wilkinson, Phylogenetic methods and Aetosaur interrelationships: a rejoinder, 851–852
 Harshman, John, Christopher J. Huddleston, Jonathan P. Bollback, Thomas J. Parsons, and Michael J. Braun, True and false gharials: a nuclear gene phylogeny of Crocodylia, 386–402
 Harvest mouse, 677
 Hasegawa-Kishino-Yano (HKY) model, 4, 7, 49, 52–53, 121, 135, 140, 143, 215, 377, 396, 423, 444, 463–464, 466–468, 471, 506, 508, 577, 614, 681–682, 703, 737, 750, 785, 787
 Hassanin, Alexandre and Emmanuel J. P. Douzery, Molecular and morphological phylogenies of Ruminantia and the alternative position of the Moschidae, 206–228
 Hausdorf, Bernhard and Christian Hennig, Biotic element analysis in biogeography, 717–723
 Hawaii, 70–88, 89–109, 820–841
 Hawkmoths, 349
 Hayashi, Cheryl, *see* Gatesy, John, 403–422
 Heckert, Andrew B. and Spencer G. Lucas, Clarifying Aetosaur phylogeny requires more fossils, not more trees—reply to Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), 253–254
 Heins, David C., *see* Guill, J. Michael, 4880500
Helicops, 447, 451
angulatus, 441
pictiventris, 441
 Helminthes, 430, 520
 Hemichordata, 296
 Hemidactyliine salamanders, 509
 Hemihalictus, 23–24, 26, 32–33
Hemirhagerrhis, 451
nototaenia, 440
viperinus, 440, 459
 Hempel–Oppenheim model, 260, 263
 Hendy, M. D., *see* Holland, B. R., 229–238
 Hendy, Michael D., *see* Gascuel, Olivier, 110–118
 Henicopidae, 559
 Henicopinae, 559
 Hennig, Christian, *see* Hausdorf, Bernhard, 717–723
 Hennig’s auxiliary principle, 239, 245–246
 Hepaticophyta, 430
Heptodon, 857
 Herbivores, 339, 349, 746
 Herpestidae, 846
Heteranthera, 154, 156
dubia, 154, 156
limosa, 154, 156
oblongifolia, 154, 156
rotundifolia, 154, 156
seubertiana, 154, 156
zosterifolia, 154, 156
 Heterocoela, 311, 313, 318, 320, 322, 327–328
 Heterodactyla, 746, 750–751, 754
Heterodon, 447, 451, 455
nasicus, 441
simus, 441
platyrhinos, 287
 Heterodonta, 561
 Heteromyidae, 606, 608, 614
Heteropia, 314–315, 332–333
 Heteroptera, 103
 Heteroptiidae, 314, 322, 324, 329
Heterotropus senex, 756
 Heuristic
 parsimony search, 75
 search, 4, 39, 76, 95–96, 166, 318, 327, 336, 368, 372, 377, 504, 575, 607, 609, 611, 621, 666, 737, 823–824, 842, 864
 tree, 607
 Heuristics, 368–373, 696, 782, 841
 Hexactinellida, 311, 314, 316–318, 320, 328, 330
 Hexapoda, 328, 557
 Hierarchical phylogenetic models for analyzing multipartite sequence data, Marc A. Suchard, Christina M. R. Kitchen, Janet S. Sinsheimer, and Robert E. Weiss, 649–664
 Hillis, David M., David D. Pollock, Jimmy A. McGuire, and Derrick J. Zwickl, Is sparse taxon sampling a problem for phylogenetic inference?, 124–126
 Hillis, David M., *see* Wiens, John J., 501–514
 Hippoboscids, 37
 Hipponicidae, 639
 Hipponics, 620, 622, 632
 artiodactylans, 854

- Hipponix*
 sp. Australia, 639
 sp. California, 639
 Hippopotamidae, 209–210, 221, 855
Hippopotamus, 209, 211–213, 215, 222, 706, 712
amphibius, 210
 Hippos, 207, 210, 707–708, 854–855, 858
Hippospongia communis, 316
 Hippotragini, 209–210, 215
Hippotragus, 209, 212, 215, 218
niger, 210
Hispanomeryx, 220
 Historical
 biogeography, 271–280
 effects, 684–695
 HIV, 660, 662, 735
 evolution, 649–664
 Hood, Craig S., *see* Guill, J. Michael, 488–500
 Holarctic rodents, 186
 Holland, B. R., D. Penny, and M. D. Hendy, Outgroup misplacement and phylogenetic inaccuracy under a molecular clock—a simulation study, 229–238
 Holothyrina, 300, 309
Homalictus, 24
 Homalopsinae, 439, 452–453, 455, 457
 Homeobox gene, 311
 Hominoid, 711–713
 Homocoela, 311, 313, 318, 320, 322, 327–328
 Homogeneous model, 792
 Homoplastic characters, 589, 767
 Homoplasy, 16, 30, 55, 57, 61, 152, 154, 156, 211, 230, 239, 243, 245–246, 258, 297, 304, 306–307, 311, 339, 342, 348, 354, 359, 361–362, 398, 403, 415, 420, 442, 444, 460–476, 501–514, 542, 571–572, 575, 577, 581–585, 589, 604, 842, 860–861
 index, 762, 764
Homo, 706
sapiens, 286, 606, 736
Homoroselaps, 447, 451, 455–456
lacteus, 441
 Homoscleromorpha, 328
 Honey bees, 26, 296
 Honeycreepers, 89
 Hood, Craig S., *see* Guill, J. Michael, 488–500
 Hoplitomerycidae, 220–221
Hoplitomeryx, 220–221
Hoplobatrachus
occipitalis, 799, 815
rugulosus, 799, 815
 Horizontal gene transfer (HGT), 649–664
 Hormiga, Gustavo, Miquel Arnedo, and Rosemary G. Gillespie, Speciation on a conveyor belt: sequential colonization of the Hawaiian islands by *Orsonwelles* spider (Araneae, Linyphiidae), 70–88
Hormiphora sp., 316–317
 Horned soldiers, 137, 139–140, 143
 Horse flies, 745–746
 Horses, 707–708, 857
 Horseshoe crab, 300
 Host
 phylogeny, 740, 742
 tree, 737, 740
 Host–parasite coevolution, 735–744
 House flies, 745
 HSP70, 596, 600
 HSP90, 598
 Huddleston, Christopher J., *see* Harshman, John, 386–402
 Huelsenbeck, John P. and Katherine M. Lander, Frequent inconsistency of parsimony under a simple model of cladogenesis, 641–648
 Huelsenbeck, John P., Rasmus Nielsen, and Jonathan P. Bollback, Stochastic mapping of morphological characters, 131–158
 Hugall, A. F., *see* Lee, M. S. Y., 15–22
 Hugot, Jean-Pierre, Christine Demanche, Véronique Barriol, Eduardo Dei-Cas, and Jacques Guillot, Phylogenetic systematics and evolution of primate-derived *Pneumocystis* based on mitochondrial or nuclear DNA sequence comparison, 735–744
 Human, 52–53, 526, 565, 571, 591, 656707–708, 711–713, 715, 738
 Humphries, Christopher J., *see* Williams, David M., 255–258
 Hyaenidae, 846
Hyalomma, 299, 304–305, 310
aegyptium, 299
dromedarii, 299
marginatum, 299
truncatum, 299
 Hybridization, 63, 89, 92, 103, 379, 383, 820, 832
Hydrodynastes, 447, 451
bicinctus, 441
 Hydrophiinae, 441, 447
Hydrophis, 447, 451, 456
semperi, 441
Hydropotes, 206, 209, 212, 215, 218–220, 222, 224–225
inermis, 210
 Hydropotinae, 209
 Hydropsini, 456
Hydrothrix gardneri, 154, 156
Hyemoschus, 206, 219
Hylaeochampsa, 412
 Hymenoptera, 23, 296, 307
 Hypertragulidae, 207, 222
Hypertragulus, 221
Hyposmocoma, 89
 Hypothesis test, 48–54, 781–793
Hyracotherium, 857
 Hystricognathi, 604–605, 608–611, 613, 615

Ictonyx, 590
striatus, 581, 589–590
 Identifying conflicting signal in a multigene analysis reveals a highly resolved tree: the phylogeny of Rodentia (Mammalia), Ronald W. DeBry, 604–617
 Idiosyncratic markers, 296–310
 Impala, 210
 Ichthyosaurs, 501
 Incomplete taxa, 528–538
 Incongruence, 11, 15, 20, 28, 37, 39–41, 43, 45, 404, 451, 559, 571–572, 581, 587, 608, 613
 length difference, 75–77, 406, 408, 463
 length difference (ILD) test (*see also* partition homogeneity test), 3, 11, 15, 26, 28, 75–76, 418, 467, 604, 607–609, 613, 622, 773, 853–855, 860, 862
 Inconsistency, 229–238, 641–648
 Incorrect model, 674–683
 Indel-change ratio, 556–557
 Indels, 283–295, 300, 377–378, 392, 394, 397, 574–575, 651, 823, 827
 characters, 389, 823, 826, 829
 coding, 444, 575
 parsimony informative, 823
 Independent and identically distributed (IID), 842, 847
 Ingroup, 27, 56, 89, 91–92, 96–97, 99, 197, 229–230, 386–387, 389, 392, 394, 396–397, 443, 576, 620–622, 748, 750, 753, 757, 764, 775, 824, 826

- Informative characters, 16, 575, 583
- Inquilina*, 1
- dawsoni*, 2
- schwarzi*, 2
- Insecta, 37, 296, 430
- Insects, 23, 32, 34, 89, 106, 430
- Insertion (*see indel*)
- Internal transcriber spacer (ITS), 73, 76–77, 298, 336, 370–371, 382, 820–841
- Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), Simon R. Harris, David J. Gower, and Mark Wilkinson, 239–252
- Intraspecific phylogeny of the New Zealand short-tailed bat *Mystacina tuberculata* inferred from multiple mitochondrial gene sequences, Brian D. Lloyd, 460–476
- Intron–exon, 20, 26
- Introns, 376, 383, 391, 394, 571–593
- Invertebrates, 429, 618–619
- Ips* beetles, 677
- IRBP, 604–614, 738
- Irregularia, 328
- Ischnura*, 91, 94, 102
- aurora*, 94
- heterosticta*, 94
- Island evolution, 820–841
- Iskandar, Djoko, *see* Evans, Ben J., 794–823
- Isopoda, 430
- Is sparse taxon sampling a problem for phylogenetic inference?, David M. Hillis, David D. Pollock, Jimmy A. McGuire, and Derrick J. Zwickl, 124–126
- Ixodes*
- affinis*, 300
- auritulus*, 300
- hexagonus*, 299–300
- holocyclus*, 300
- pilosus*, 300
- simplex*, 300
- ss.*, 298, 300, 304, 307, 309
- tasmani*, 300, 304, 309
- uriae*, 300
- Ixodidae, 296–310
- Jacarea, 409, 412
- Jaccard distance, 718
- Jackknife, 55–65, 128–129, 409, 554, 556, 560–561, 575, 581
- analysis, 407
- parsimony, 129, 557, 559
- scores, 412
- support, 419, 540, 559, 561, 582–583
- values, 589, 665
- Janacus*, 619, 623, 632, 636, 639
- Janneria pustulata*, 640
- Japygidae, 557
- Jean-Marie, Alain, *see* Gascuel, Olivier, 110–118
- Jenkina*, 314–315
- Jenkinidae, 314
- Jennings, W. Bryan, Eric R. Pianka, and Stephen Donnellan, Systematics of the lizard family Pygopodidae with implications for the diversification of Australian temperate biotas, 757–780
- Ji, Shuqing, *see* Danforth, Bryan N., 23–36
- Johnson, Kevin P., Richard J. Adams, Roderic D. M. Page, and Dale H. Clayton, When do parasites fail to speciate in response to host speciation?, 37–47
- Jones, Kate E., *see* Bininda-Emonds, Olaf R. P., 724–729
- Jordan, Steve, Chris Simon, and Dan Polhemus, Molecular systematics and adaptive radiation of Hawaii's endemic damselfly genus *Megalagrion* (Odonata: Coenagrionidae), 89–109
- Joyce, Paul, *see* Minin, Vladimir, 674–683
- Jukes–Cantor
- distances, 425
- model, 133, 371, 424, 485, 530, 644, 666–669, 703, 708, 711–715, 784–785, 787–788, 820, 825, 831–832, 834
- Juliformia, 557
- Jungles
- analysis, 39, 43, 45
- optimizations, 42
- Jurassic, 746, 753
- late-, 549, 746
- lower, 746, 749
- mid-, 745
- upper, 753
- κ -casein, 206–207, 209–212, 215
- Kebira*, 333
- Kelly, Christopher M., R. Nigel P. Barker, and Martin H. Villet, Phylogenetics of advanced snakes (Caenophidia) based on four mitochondrial genes, 439–459
- Kendall's
- shape space, 66
- tangent space, 66–67
- Kimura three-parameter model, 577, 581
- Kimura two-parameter model, 133, 229, 231, 236, 485, 577, 700, 703, 762, 801, 823, 826, 850
- Kishino–Hasegawa test, 4, 10, 15–22, 212, 216, 390, 392, 394–396, 398, 445, 504, 781
- Kishino, Hirohisa, *see* Wiegmann, Brian M., 745–756
- Kitchen, Christina M. R., *see* Suchard, Marc A., 649–664
- Kiwi, 474, 550
- Kjer, Karl M., *see* Xia, Xuhua, 283–295
- Koepfli Klaus-Peter and Robert K. Wayne, Type I STS markers are more informative than cytochrome *b* in phylogenetic reconstruction of the Mustelidae (Mammalia: Carnivora), 571–593
- Kolmogorov–Smirnov test, 190
- Krebsia* sp., 639
- Kulczynski distance, 718
- Kullback–Leibler distance, 782, 785
- Kumar, Sudhir, *see* Rosenberg, Michael S., 119–123
- Labulla*, 71, 77
- thoracica*, 71–73, 88
- Lacanicephalidea, 430
- Lactoferrin, 206, 209–212, 215, 218, 220
- Lagomeryx*, 224
- Lagomorpha, 606, 839–851
- Laminacauda gigas*, 71
- Lamontia*, 314–315, 317, 323–324
- zona*, 333
- Lamprophiinae, 439, 441, 447, 455–456
- Lamprophis*
- fuliginosus*, 441
- inornatus*, 441, 445, 459
- Lander, Katherine M., *see* Huelsenbeck, John P., 641–648
- Janice conchylega*, 316
- Lapointe, François-Joseph, Mark Wilkinson, and David Bryant, Matrix representations with parsimony or with distances: two sides of the same coin?, 865–868
- Lapointe, François-Joseph, *see* Wilkinson, Mark, 127–130
- Larus glaucooides*, 286
- Larvae, 8, 618, 631
- Larval characters, 8–9
- Lasioglossum*, 23–28, 30, 32–33

- (Dialictus) cressonii*, 27
(Dialictus) figueresi, 27, 33
(Dialictus) gundlachii, 27
(Dialictus) hyalinum, 27
(Dialictus) imitatum, 27
(Dialictus) parvum, 27
(Dialictus) pilosum, 27
(Dialictus) rohweri, 27
(Dialictus) tegulare, 27
(Dialictus) umbripenne, 27
(Dialictus) vierecki, 27
(Dialictus) zephyrum, 24, 27, 33
(Evyllaesus) albipes, 26–27, 33
(Evyllaesus) apristum, 27
(Evyllaesus) boreale, 27
(Evyllaesus) brevicorne, 27
(Evyllaesus) calceatum, 24, 27, 33
(Evyllaesus) cinctipes, 27
(Evyllaesus) comagenense, 27
(Evyllaesus) duplex, 27
(Evyllaesus) fulvicorne, 27
(Evyllaesus) gattaca, 27
(Evyllaesus) inconditum, 27
(Evyllaesus) interruptum, 27
(Evyllaesus) laticeps, 24, 27
(Evyllaesus) limbellum, 27
(Evyllaesus) lineare, 27
(Evyllaesus) lucidulum, 27
(Evyllaesus) malachurum, 27
(Evyllaesus) marginatum, 24, 27, 33
(Evyllaesus) mediterraneum, 27
(Evyllaesus) morio, 27
(Evyllaesus) nigripes, 27
(Evyllaesus) pauxillum, 27
(Evyllaesus) pectorale, 27
(Evyllaesus) politum, 24, 27, 33
(Evyllaesus) puncticolle, 27
(Evyllaesus) quebecense, 27
(Evyllaesus) subtropicum, 27
(Evyllaesus) truncatum, 27
(Evyllaesus) villosulum, 26–27
(Hemihalictus) lustrans, 24, 27, 33
(Lasioglossum) athabascense, 27
(Lasioglossum) coriaceum, 26
(Lasioglossum) fuscipenne, 27
(Lasioglossum) leucozonium, 24
(Lasioglossum) pavonotum, 27
(Lasioglossum) sisymbrii, 27
(Lasioglossum) titusi, 27
(Paralictus) asteris, 27
(Sphecodogastra) noctivaga, 27
(Sphecodogastra) oenotherae, 27
(Sudila) alphenum, 27
Laticauda, 451
colubrina, 441
Latimeria chalumnae, 286
Laupala, 833
 Least and most powerful phylogenetic tests to elucidate the origin of the seed plants in the presence of conflicting signals under misspecified models, Stéphane Aris–Brosou, 781–793
 Least-squares
 algorithm, 866
 criterion, 865
 Lee, M. S. Y. and A. F. Hugall, Partitioned likelihood support and the evaluation of data set conflict, 15–22
 Leeches, 430
Leioheterodon, 451, 456
 madagascariensis, 441
Lelapia, 314–315, 324
 Lelapiidae, 314, 329
 Lemuridae, 711–713
 Lemuriform, 711–713
 Lemurs, 52–53, 524, 706,
 mouse, 705–706, 708, 711–715
 Le Parco, Yannick, *see* Manuel, Michael, 311–333
Lepilemur, 706
Lepthyphantes minutus, 73, 88
Leptictis, 858
 Leptomerycidae, 222
Leptomeryx, 222
Leptonetis perplexus, 639
Leptonycteris curasoae yerbabuena, 473
Leptotila, 41
 jamaicensis, 38, 41–42
 plumbeiceps, 38
 rufaxilla, 38
 verreauxi, 41
 verreauxi angelica, 38
 verreauxi fulviventris, 38
Leptynoma hessei, 756
Lestis, 2
 Leucaltidae, 314
Leucaltis, 314–315, 320, 324, 332
 clathria, 316
Leucandra, 314–315, 320, 322–324, 332–333
 aspera, 316
Leucascandra, 314–315, 332
 Leucascidae, 314
Leucetta, 314–315, 320, 322, 324, 330, 332–333
 chagosensis, 316
 Leucettidae, 314
Leucilla, 314–315, 332
Leucocrossuromys leucurus, 189
Leuconia, 314–315, 320, 322–324, 328, 330, 332–333
 johnstoni, 333
 nivea, 316, 329, 333
 ochotensis, 333
Leucopsila, 314–315, 317, 323
Leucosolenia, 314–315, 320, 322–323, 329, 332–333
 laxa, 332
 sp., 316
 variabilis, 329
 Leucosoleniida, 314, 320, 323
 Leucosoleniidae, 314
 Levin, Rachel A., Lucinda A. McDade, and Robert A. Raguso,
 Systematic utility of floral and vegetative fragrance in two
 genera of Nyctaginaceae, 334–351
Lialis, 757–758, 762, 764, 767, 770–771, 775
 burtonis, 759, 762
 jicari, 759
Liancalus sp., 756
 Lice, 37–47, 430
 Lichens, 430, 518
Ligeromeryx, 224
 Likelihood (*see* maximum likelihood), 352–367
 analysis, 18, 263, 374, 708, 715, 803–804, 807
 framework, 369
 methods, 286
 model, 17, 212, 356–358, 362, 481, 506, 555, 674, 681, 705
 ratio tests (LRT), 41–42, 48, 54, 63, 95, 352, 363–364, 377, 379, 390,
 394, 445, 575, 580, 598, 600, 604–617, 621, 655, 674–683, 714, 825,
 849, 801

- trees, 39, 506, 508, 762, 764, 802
weight test, 18–19, 22
- Limnonectes*, 794–795, 797–799, 802, 805–809, 811–812, 815
- acanthi*, 797, 802, 807, 809–811, 815
- arathooni*, 795, 797, 812, 815
- asperatus*, 797, 799, 815
- blythii*, 797, 808
- cf. *acanthi*, 802, 810, 815
- cf. *blythii* 1, 804, 815
- cf. *blythii* 2, 804, 815
- cf. *blythii* 3, 802, 804, 815
- cf. *heinrichi*, 812, 815
- cf. *kuhlii* 1, 795, 797, 808, 815
- cf. *kuhlii* 2, 795, 797, 808, 815
- cf. *kuhlii* 3, 795, 797, 808, 815
- cf. *magnus*, 797, 802, 807, 811, 815–16
- cf. *microtypanum* 1, 795, 797, 802, 812, 816
- cf. *microtypanum* 2, 795, 797, 802, 812, 816
- cf. *modestus*, 795, 797, 802, 812, 816
- cf. *modestus* complex 2, 797, 802, 812, 816
- dammermani*, 809
- diuatus*, 811
- finchi*, 802, 804, 816
- grunniens*, 795, 802, 804, 809, 816
- gyldenstolpei*, 797, 816
- heinrichi*, 795, 797, 816
- ibanorum*, 802, 804, 816
- ingeri*, 797, 804, 816
- kardasani*, 809, 816
- laticeps*, 797, 816
- leporinus*, 797, 816
- leytensis*, 797, 802, 807, 809–812, 816–817
- limborgii*, 797
- macrocephalus*, 797, 802, 807, 811, 817
- macrodon*, 797, 804, 816, 8167
- magnus*, 797, 802, 807, 811, 817
- malesianus*, 797, 800, 804, 808, 816, 817
- micrixalus*, 811
- microdiscus*, 795, 809, 817
- palavanensis*, 802, 804, 811, 817
- paramacrodon*, 802, 804, 808, 817
- parvov*, 802, 804–807, 811, 817
- pileata*, 797
- shompenorum*, 797
- sp. 1, 797, 817
- sp. 2, 797, 817
- sp. D, 797, 812, 817
- sp. G, 817–818
- sp. G2, 797, 812
- sp. I, 802, 804, 807, 812, 818
- sp. nov., 797
- sp. T, 797, 812, 818
- sp. V complex 1, 797, 812, 818
- sp. V complex 2, 797, 802, 819
- visayanus*, 797, 802, 807, 811, 819
- woodworthi*, 797, 802, 807, 811, 819
- Limulus polyphemus*, 300
- Linear model, 684–695
- Linnean parsimony, 342
- Linyphia*, 77
- triangularis*, 73, 88
- Linyphiidae, 70–88
- Linyphiini, 72, 77
- Liocheles waigiensis*, 300
- Liolaemidae, 159
- Liolaemus*, 159–185, 688
- austromendocinus*, 161, 163, 167, 178–179
- bibronii*, 164, 167
- buergeri*, 161, 178, 182
- capillitas*, 161, 163, 167, 179
- ceii*, 161, 178, 182
- ceii-kriegi* clade, 172
- christiani*, 161
- curis*, 161
- elongatus*, 159–161, 163, 166–167, 171–172, 178–179, 182
- elongatus-kriegi* complex, 159–161, 167, 182
- gracilis*, 164, 167
- helioderms*, 161
- kingii*, 164, 167
- kriegi*, 159–161, 164, 166–167, 171–172, 178, 182
- leopardinus*, 161
- lineomaculatus*, 164, 167
- neuquensis*, 164, 167
- petrophilus*, 161, 163, 166–167, 172, 178–179, 182
- pictus*, 164, 167
- pseudocnomalus*, 165, 167
- sp. 1, 163, 182
- sp. 2, 164, 179
- sp. 3, 163–164, 167
- sp. 4, 164, 179, 182
- sp. 5, 163, 171, 178, 182
- sp. 6, 163, 171, 178, 182
- sp. 7, 163, 171, 178, 182
- sp. 8, 164, 179, 182
- sp. A, 164, 179, 182
- sp. B, 164, 178, 182
- sp. C, 164, 178
- thermarum*, 161, 178
- vallecurensis*, 164, 167
- Liophis*, 447, 451
- reginae*, 441
- Lithonida, 314, 317, 323, 329–330
- Littorina littorea*, 640
- Liverworts, 430
- Lizards, 105, 159–185, 286, 306, 423, 689, 691, 693, 757–780
- agamid, 758
- New World, 160
- pygopodid, 757–780
- Llamas, 221
- Lloyd, Brian D., Intraspecific phylogeny of the New Zealand short-tailed bat *Mystacina tuberculata* inferred from multiple mitochondrial gene sequences, 460–476
- Lobelioids, 833
- Local clock analysis, 89
- Local molecular clock, 89–109, 705–716
- LogDet
- analysis, 3, 6–7, 11
- method, 784
- tree, 4, 7
- LogDet/Paralinear
- analysis, 614, 621
- distance (*see also* paralinear distance), 284, 287, 289, 291, 293, 397, 580, 610, 612, 676
- transformation, 579
- Log-likelihood, 596, 598–599, 614, 655, 782
- ratio test, 3–4
- scores, 445, 464, 607, 761
- values, 445, 761
- Long branch
- attraction, 63, 79, 82, 96, 327, 354–370, 386, 419, 423–428, 443, 478, 483, 510, 531, 537, 558, 590, 594
- effects, 2

- subdivision, 423–428
 Longhorn beetle, 89
 Longirostres, 390–391, 398, 400
Longosuchus, 241–243, 247–252
 Long wavelength (LW)
 opsin, 26–28
 rhodopsin, 23
Lontra, 577, 590
 canadensis, 573
 felina, 573
 longicaudis, 573
Lorancameryx, 224
Lordotus sp., 756
Loris, 706
 Lorisiform, 711–713
 Loxocemidae, 440
Loxocemus bicolour, 440
Loxodonta, 591
 Lucas, Spencer G., *see* Heckert, Andrew B., 253–255
Lucasuchus, 241–243, 250–251
Lutra, 590
 lutra, 573
 maculicollis, 583, 590
 Lutrinae, 571–573, 581, 590, 846
Lycodon, 447, 451, 456–457
 capucinus, 440
 laensis, 440
Lycodonomorphus, 456
 Lycodontinae, 456
 Lycodontini, 457

Macaca, 706, 741
 fascicularis, 736, 741
 mulatta, 736
 nemestrina, 736, 741
 Macaque, 52–53
Macroderma gigas, 472
Macrogalea, 1, 4, 7–8, 10–13
 candida, 2, 10, 12
 scaevolae, 12
 zanzibarica, 2, 10, 12
Macropisthodon, 447, 451, 457
 rudis, 440, 447, 456
 Macrostromata, 440
Madagascarophis, 451
 colubrine, 441
 Magnoliaceae, 346
 Magnolias, 346
 Magpie goose, 388
 Mahalanobis distance, 192, 491
 Malacostraca, 557
Malluvium sp., 639
 Maloideae, 375
Malpolon, 451
 monspessulanus, 441, 459
 Mammalia, 566, 571–593, 604–605, 861
 Mammalian phylogeny, 857
 Mammals, 16, 23, 56, 66, 106, 206, 219, 224, 261–262, 283–287, 293, 415,
 518, 524, 550, 571, 605, 794, 853, 861
 eutherian, 249, 585, 860
 placental, 591
 Mandible, 186–205
 Mandibular
 characters, 192
 dendograms, 195
 Mandibulata, 557
Manuelia, 8
 postica, 2
 Manuelliini, 1, 9
 Mantel
 matrix correlations, 493–494, 497
 statistics, 488, 491, 493
 test, 491
 Manuel, Michael, Carole Borchiellini, Eliane Alivon, Yannick Le
 Parco, Jean Vacelet, and Nicole Boury-Esnault, Phylogeny and
 evolution of Calcareous sponges: monophyly of Calcinea and
 Calcaronea, high level of morphological homoplasy, and the
 primitive nature of axial symmetry, 311–333
Maoricicada, 674
 campbelli, 474
Maoricrypta, 619, 623, 631–632, 636
 Markov chain (*see also* MCMC), 50, 133–134, 140, 165, 212, 377, 445,
 576, 666, 749–750, 760, 785
 continuous time, 179, 644
 model, 782, 785, 791, 841
 processes, 594
 Markov chain Monte Carlo (MCMC), 48–54, 95, 131–158, 159, 212,
 377, 445, 478, 484, 576, 604, 607, 649–664, 667, 674, 709, 711, 749,
 751, 760
 algorithms, 135–136, 665–666, 697, 711, 785
 analysis, 140, 213, 486, 750, 801
 approach, 702, 750
 Metropolis-coupled, 604, 607, 621, 623
 procedure, 748, 750
 processes, 825
 sampling, 477–487, 612
 simulations, 165
Marmota, 186–205
 baibacina, 189, 192–193, 196, 199, 201
 baibacina centralis, 201
 bobak, 189, 192, 196–197, 199, 202
 broweri, 186, 189–190, 192–193, 196–198, 202
 caligata, 186, 190, 192, 194–196, 198, 202
 caligata caligata, 202
 caligata cascadenis, 202
 caligata nivaria, 202
 caligata okanagana, 202
 caligata oxytona, 202
 caligata sheldoni, 202
 caligata vigilis, 202
 camtschatica, 186, 189, 192, 196, 198
 camtschatica bungei, 202
 caudata, 188–190, 192, 195–196, 199
 caudata aurea, 202
 flaviventris, 192, 194–197, 205
 flaviventris avara, 203
 flaviventris dacota, 203
 flaviventris engelhardti, 203
 flaviventris flaviventris, 203
 flaviventris luteola, 203
 flaviventris nosphora, 203
 flaviventris obscura, 203
 flaviventris parvula, 203
 flaviventris sierrae, 203
 himalayana, 188–189, 195–197, 203
 himalayana himalayana, 203
 himalayana robusta, 203
 marmota, 186–187, 189, 192, 196, 203
 marmota marmota, 203–204
 menzbieri, 188–189, 192, 195–197, 199, 204
 monax, 186, 189, 192–193, 196–197, 199, 204
 monax canadensis, 204
 monax ignava, 204
 monax monax, 204

- monax ochracea*, 204
monax petrensis, 204
monax prebolorum, 204
monax rufescens, 204
olympus, 186, 190, 192–198, 204
sibirica, 189–190, 192–193, 196, 199, 204
vancouverensis, 186, 188, 190, 192, 194, 196, 198, 204
Marmotini, 197
Marmots (*see also Marmota*), 186, 188–189, 192–195, 197, 205
Alaskan, 186, 193, 195, 198
Alpine, 186–187, 198–199
black-capped, 186, 193, 198–199
Eurasian, 186, 193, 198–199
hoary, 194
long-tailed, 195, 199
Menzbier, 189, 199
Olympic, 194–195, 197–198
Palaeartic, 187, 198–199
Vancouver Island, 189, 194, 196–199
yellow-bellied, 196–198, 205
Marsupials, 524
Martens, 571–573
American, 573
Martes, 577, 581, 590
americana, 573, 581, 590
pennanti, 573, 581, 590
Marsupials, 709
Mast, Austin R. and Reto Nyffeler, Using a null model to recognize significant co-occurrence prior to identifying candidate areas of endemism, 271–280
Maticora, 447, 451
bivigata, 441
Matrix correlation analysis,
nesting, 241
parsimony, 241
with distances (MRD), 865–868
with parsimony (MRP), 255, 439, 455–456, 839–851, 865–868
Matrix representations with parsimony or with distances: two sides of the same coin? François-Joseph Lapointe, Mark Wilkinson, and David Bryant, 865–868
Matrix-weighted
analysis, 622
parsimony, 623
Maximum likelihood, 16–18, 23, 26, 28, 39, 41, 63, 76, 95, 98–100, 121, 128, 133, 159, 229–238, 283, 285, 287, 293, 320, 363, 365, 368–373, 386, 439–459, 463, 468, 477–487, 506, 571, 585, 594–603, 604–617, 622, 641–642, 656, 665–666, 671–683, 696–704, 705–716, 726, 801
analysis, 1, 3, 7, 11, 26–27, 30, 42, 63, 76, 79, 89, 95–97, 99, 166–167, 171, 182, 211–212, 215, 231, 318, 322, 327, 329, 341, 377–378, 386, 389–90, 394–396, 407, 419, 715, 737, 740, 797, 801, 824, 829, 850
bootstrap support, 667–668
estimates, 166, 394, 594, 596, 601, 604, 669, 709, 711–712, 714, 748, 850
model, 11, 45, 96, 159, 220, 358, 397
searches, 95
tests, 614
trees, 12, 18, 20, 22, 30, 42–45, 81, 96, 128, 323, 328, 368, 371, 383, 396, 455–456, 479, 574, 576–577, 580–581, 583, 589–590, 607, 611, 666, 677, 681, 762, 767, 771, 781, 825, 827–829, 849
value, 140
Maximum parsimony (*see also* parsimony), 1, 12, 95, 98, 120, 159, 229–238, 293, 298, 306–307, 318, 327, 339–340, 425, 439–459, 463, 468, 478, 489, 607, 609–610, 613–614, 726, 801, 820, 823, 826, 833
analysis, 3, 5–7, 10–11, 95, 166–167, 182, 211, 213, 215, 275, 317, 322, 327, 447, 467, 571, 577, 580–581, 615, 737, 740, 797, 801, 824, 829, 850
reconstruction, 450
trees (*see also* most-parsimonious trees), 4, 110, 213, 215, 217, 240, 242–243, 248, 252, 275, 317, 320, 323, 329, 336, 339–341, 346, 359, 378, 383–384, 444, 446–447, 463, 574, 576–577, 579–581, 589–590, 607–608, 824, 826–828
McDade, Lucinda A., *see* Levin, Rachel A., 334–351
McGuire, Jimmy A., *see* Evans, Ben J., 794–823
McGuire, Jimmy A., *see* Hillis, David M., 124–126
McLachlan, Robert, *see* Gascuel, Olivier, 110–118
Megalagrion, 89–109, 833
adytum, 93, 103, 105
blackburni, 93, 103
calliphya, 93, 103
damsel flies, 89–109, 833
eudytum, 89, 93, 103, 105
hawaiiense, 93, 101–104, 106
heterogamias, 93, 101, 103–104, 106
jugorum, 103, 106
kauaiense, 93, 101, 103–104, 106
koelense, 92–93, 101, 104
leptodemas, 93, 103
mauka, 89, 93, 101, 103, 105
molokaiense, 106
nesiotes, 89, 93, 101, 103, 105–106
nigrohamatum, 93, 103
nigrolineatum, 93
oahuense, 89, 93, 101, 103, 105
oceanicum, 93, 103
oresitrophum, 89, 93, 99, 101, 103–104, 106
orobates, 89, 93, 99, 101, 103–104, 106
pacificum, 89, 92–93, 99–100, 103, 105–106
paludicola, 89, 93, 101, 103, 105–106
sp. nov., 93
vagabundum, 89, 93, 103, 105
williamsoni, 92, 103
xanthomelas, 89, 91–93, 103, 105
Megaloptera, 106
Megapodon, 314–315, 328, 333
Megophryidae, 794
Mehelya, 447, 451, 455–456
capensis, 441
nyassae, 441, 459
Melanesobasis, 91
Melanosuchus, 391, 396, 412
niger, 388, 405
Meles, 581, 590
meles, 573–574, 577, 582
Melinae, 571–573, 581
Mellivorinae, 572–573
Melnick, Don J., *see* Evans, Ben J., 794–823
Melogale, 589–590
moschata, 581–583, 590
Mephitinae, 572–573, 590, 846
Merostostomata, 300
Merycodontinae, 224
Mesonychia, 853–855, 858, 860
Mesostigmata, 300, 309
Mesozoic, 549–553, 745–746
Meta-optimality criterion, 554–564
Metastrata, 296, 299, 304, 307
Metazoa, 311–333, 520
Meteoriaceae, 430
Methanococcus jannaschii, 659
Metriopelia ceciliae, 38
Metropolis-coupled MCMC, 604, 607, 621, 623
Metropolis–Hastings

- algorithm, 662
 criterion, 50
 proposal, 654–655
- Metropolis-within-Gibbs cycle, 654–655
- Metrosideros polymorpha*, 820
- Microbats, 460
- Microbes, 429
- Microcebus*
berthae, 706
griseorufus, 706
murinus, 706
myoxinus, 706
ravelobensis, 706
rufus, 706
sambiranensis, 706
tavaratra, 706
- Microchiroptera, 460–476
- Microcionia prolifera*, 316
- Microevolutionary model, 692
- Microhylidae, 794
- Microlymphina*, 77
dana, 73, 88
- Micruroides*, 447, 451
euryxanthus, 441
- Micrurus*, 447, 451
diastema, 441
fulvius, 441
- Miller, Jeremy A., Assessing progress in systematics with continuous jackknife function analysis, 55–65
- Millipedes, 430
- Minchinella*, 314–315, 317, 323, 328–329, 332
- Minchinellidae, 314, 320, 328
- Minimum evolution, 121, 287, 478, 489, 579
 tree, 506, 580, 610
- Minimum-length trees, 415, 750
- Minin, Vladimir, Zaid Abdo, Paul Joyce, and Jack Sullivan, Performance-based selection of likelihood models for phylogeny estimation, 674–683
- Mink, American, 573
- Miocene, 207, 221, 224–225, 451, 549, 629, 761, 776
 early-, 206–207, 209, 216, 218, 224–225
 mid-, 206, 217–218, 224–225
 late-, 218, 224
- Mirabilis*, 334–351
alipes, 340
bigelovii, 336
greenii, 343–344
jalapa, 336
longiflora, 336
macfarlanei, 341, 343–344
multiflora, 347
pudica, 340
- Mirza, 706
- Misleading zones, 229–238
- Missing data, 528–538
- Missing data, incomplete taxa, and phylogenetic accuracy, John J. Wiens, 528–538
- Mites, 299–300, 430
- Mitochondrial
 alleles, 41
 DNA, 3–4, 7, 12, 37, 48, 52, 89, 94–101, 103, 159–185, 296, 386–387, 395, 407–408, 415, 418, 423, 439–459, 460–461, 464, 471, 474, 503–504, 506, 508–509, 548, 551, 571–572, 583, 586, 589, 656, 674, 677, 680–681, 735, 757–758, 760–762, 764, 767, 770–771, 773, 775–776, 794, 798, 801, 808, 810–812
- DNA tree, 178
- genes, 5–6, 10, 16, 32, 41, 74, 439, 451, 460, 462, 510, 572, 587, 649, 651, 656
- genomes, 56, 58–59, 238, 296–297, 301, 303, 307, 309, 462–463, 478, 577, 706
- RNA, 445
- rRNA, 615
- tRNA, 615
- Mnemiopsis*, 317
leidy, 316
- Models
 analysis, 542
 Bayesian, 49, 650
 birth–death, 641, 644–645, 647, 761
 cladogenetic, 641
 continuous–time Markov, 131–134, 147, 149, 651
 discrete, 594, 602
 DNA substitution, 444
 equal-rates, 57
 evolutionary, 126, 352, 652, 665, 668, 672, 692, 696–697, 705, 727, 781–782, 788, 791–792, 801, 829, 850
 Felsenstein, 463, 703, 708–709, 711–715, 748, 850
 gamma, 594–603, 750
 general-time-reversible (GTR), 3–4, 7–8, 18–19, 26, 28–30, 42, 95, 97, 165–166, 212, 286, 318, 390, 392, 396, 445, 463, 576, 579, 585, 608–611, 613, 666–669, 674, 677, 679, 681–682, 737, 762, 784, 801–802, 825, 831, 834
 Hasegawa-Kishino-Yano (HKY), 4, 7, 49, 52–53, 121, 135, 140, 143, 215, 377, 396, 423, 444, 463–464, 466–468, 471, 506, 508, 577, 614, 681–682, 703, 737, 750, 785, 787
 Hempel–Oppenheim, 260, 263
 homogeneous, 792
 incorrect, 674–683
 Jukes–Cantor, 133, 371, 424, 485, 530, 644, 666–669, 703, 708, 711–715, 784–785, 787–788, 820, 825, 831–832, 834
 Kimura three-parameter, 577
 Kimura two-parameter, 133, 229, 231, 236, 485, 577, 700, 703, 762, 801, 823, 826, 850
 likelihood, 17, 212, 356–358, 362, 481, 506, 555, 674, 681, 705
 linear, 684–695
 Markov, 147, 594–603, 782, 785, 791, 841
 maximum likelihood, 11, 45, 96, 159, 220, 358, 397
 microevolutionary, 692
 misspecification, 665–673
 molecular clock, 48
 nested, 606
 nonreversible nucleotide substitution, 48
 nuclear substitution, 215, 287, 377, 604–617, 658, 674–683, 708, 785
 null, 271–280, 480, 482, 484, 717–723
 parsimony, 361, 363
 phylogenetic, 133, 594, 649, 651, 654, 656, 661
 Poisson process, 134
 rates-across-sites, 595–596
 rate homogeneity, 463
 regression, 684, 686
 site-specific-rates (SSR), 4, 8, 26–28, 30, 463, 466–468, 610, 612, 674
 stochastic, 152, 705, 761, 833
 substitution, 3, 140, 293, 371, 463, 577
 Tamura and Nei, 703
 three-parameter TN93, 293
 time-homogeneous, 647
 tree, 231, 423, 427, 540, 542, 761, 841–842
 Wright–Fisher, 179
- Model-based Gaussian clustering (MBGC), 717–718, 722
- Models of evolution, 126, 352, 364, 377, 615, 665, 668, 672, 692, 696–697, 705, 727, 781–782, 788, 791–792, 801, 829, 850
- Molecular

- analysis, 103–104, 221, 323, 386, 397, 796
 characters, 55, 70, 76–77, 131, 199, 209, 296, 298
 cladograms, 187, 742
 cladistics, 187
 clock (*see also* Bayesian clocklike model, non-parametric rate smoothing), 11, 48, 50–53, 70, 81, 110, 229–231, 233, 386, 390, 395, 550, 641, 644, 647, 700, 705–716, 745, 754, 757–780, 782, 801–802, 825, 841
 data, 57, 62, 75, 82, 99, 103–105, 187, 300, 317, 320, 328–329, 339, 369, 386–389, 397, 401, 403, 406, 423, 502–504, 510–511, 539, 543, 726–727, 760, 764, 767, 770, 796, 853, 857
 dating analysis, 375
 evolution, 20, 38, 41, 591, 594–603, 785, 791
 local, 89–109
 model, 48
 partitions, 17
 phylogenetics, 187, 283–295, 311, 715
 phylogeny, 37–47, 374–385, 545, 781, 832
 studies, 206, 386, 501, 504, 604, 614, 727
 substitution, 37
 systematics, 430, 501–514, 571–572, 591, 745–756, 820–841
 tree, 103, 105–106, 386, 389–392, 394–396, 399, 403, 853, 858, 860
- Molecular and morphological phylogenies of Ruminantia and the alternative position of the Moschidae, Alexandre Hassanin and Emmanuel J. P. Douzery, 206–228
- Molecular phylogenetics of Allodapine bees, with implications for the evolution of sociality and progressive rearing, Michael P. Schwarz, Nicholas J. Bull, and Steven J. B. Cooper, 1–14
- Molecular systematics and adaptive radiation of Hawaii's endemic damselfly genus *Megalagrion* (Odonata: Coenagrionidae), Steve Jordan, Chris Simon, and Dan Polhemus, 89–109
- Molecules, 206–228
- Mollusks, 430, 520, 619, 633
 deep sea, 430
- Monkeys, 707–708, 811
 owl, 738
- Monochoria
cyanea, 154, 156
hastata, 154, 156
korsakovii, 154, 156
vaginalis, 154, 156
- Monocotyledones, 524
Monoplectroninia, 329
Monosiga brevicolis, 316
- Monte Carlo
 error, 750
 estimate, 50
 permutation, 684–695
 randomization test, 684
 simulation (*see also* computer simulation and simulation), 53, 234, 717–718
- Morando, Mariana, Luciano J. Avila, and Jack W. Sites Jr., Sampling strategies for delimiting species: genes, individuals, and populations in the *Liolaemus elongatus-kriegi* complex (Squamata: Liolaemidae) in Andean–Patagonian South America, 159–185
- Morphological
 analysis, 187, 213, 323, 501, 504, 541, 560, 605, 858
 characters, 55, 57, 70, 72, 74, 87–88, 89, 91, 131–132, 135, 142, 178, 181, 199, 209, 213, 217, 219, 298, 313–314, 317, 323, 327–328, 340, 389–390, 400, 406–407, 409, 412, 420, 439, 442, 457, 474, 501–503, 506, 509–510, 537, 539–543, 545, 559, 572, 604, 632, 738, 742, 748, 760, 855–857
 data, 16, 20, 74, 103, 143, 182, 186–187, 243, 305, 315, 323, 329, 339, 369, 387–388, 397, 400–401, 403, 406, 423, 439, 501–503, 505, 726–727, 750, 764, 767, 771, 773, 796, 853–854, 857
 evolution, 186–205, 404, 542
 phylogeny, 206
 studies, 727
 support, 614–615
 trees, 387, 392, 394, 396, 503, 860
- Morphology, 10, 12, 16–17, 23, 57, 206–228, 239–252, 253–254, 296, 307, 318, 323, 328, 333, 477, 488, 501–514, 539–548, 559, 572, 614, 619, 632, 691, 748, 775, 851–852
 genitalic, 71, 82
- Morphometric
 analysis, 197
 data, 726
 distance, 497
- Morphometrics
 geometric, 66–69, 186–205, 488–500
- Morrison, David A., Branch lengths and support: revisited, 849–851
- Moschidae, 206–228
- Moschinae, 220
- Moschiola*, 206
- Moschus*, 206–207, 209, 212, 215, 219–221, 224
moschiferus, 210, 219
- Mosses, 430
- Most-parsimonious,
 combined tree, 16
 trees, 3, 5–7, 15, 55–60, 63, 575, 609, 621, 801, 804, 809, 826, 857–857, 864
- Most recent common ancestor (MRCA), 48–49, 51–52, 179–180
- Moths, tortricine, 430
- Mourasuchus atopus*, 417
- Mouse, 290, 571, 591, 615
 harvest, 677
- Mouse deer, 206, 854, 857
- Mouse lemur, 705–706, 708, 711–715
- mtLSU-rRNA, 735–738, 740–741
- mtSSU-rRNA, 735–738, 740
- Multidimensional scaling (MDS), 190, 194, 693, 718–719, 721
- Multigene analysis, 604
- Multiple trees, 849–851
- Multistate characters, 26, 75, 239, 255–257, 335
- Multivariate analysis, 652
- Muntiacinae, 209
- Muntiacus*, 209, 212, 215, 218–219, 222
reevesi, 210
- Muridae, 606, 608
- Muroidea, 604–605, 609, 613–614
- Muroid phylogeny, 613
- Murrayona*, 314–315, 317, 324, 332–333
axinellae, 316
- Murrayonidae, 314, 318, 329
- Murrell, Anna, Nick J. H. Campbell, and Stephen C. Barker, Value of idiosyncratic markers and changes to conserved tRNA sequences from the mitochondrial genome of hard ticks (Acari: Ixodida: Ixodidae) for phylogenetic inference, 296–310
- Mus*
domesticus, 606
musculus, 286, 316, 736
- Musca*, 751, 753–754
domestica, 756
- Muscidae, 756
- Muscinae, 520
- Muscomorpha, 746, 750–751, 753
- Musk deer, 206–207, 210, 219–220
- Musophaga violacea*, 388
- Mussels, 59
- Mustela*, 590
erminea, 573

- frenata*, 573, 577
vison, 573, 577
 Mustelidae, 571–593, 846
 Mustelinae, 571–573, 581
Mycale fibrexillis, 316
Mydas sp., 756
Mydaus, 573, 590
 Mydidae, 750, 756
 Myers, Daniel S., *see* Cummings, Michael P., 477–487
 Myodonta, 604–606, 608, 610–611, 613, 615
 Myomorpha, 614
 Myriapoda, 557
Mystacina, 462, 464, 467, 473
 aupourica, 460, 474
 gigas, 473
 rhyacobia, 460, 474
 tuberculata, 460–463, 466, 470–474

 NADH, 460
Naja, 447, 451
 kaouthia, 441
 naja, 441
 National Science Foundation, 428–435
 Natricinae, 439–440, 447, 452, 455–457
Natrix, 447, 451
 natrix, 440
 Naylor, Gavin J. P., and Dean C. Adams, Total evidence versus relative evidence: a response to O’Leary et al. (2003), 864–865
ND1, 73–74, 76, 657, 659
ND2, 460, 462–468, 471, 657, 659, 758–759, 762, 777
ND3, 657, 659
ND4, 161, 163–165, 167, 181, 440, 443, 445–447, 451, 453, 455, 657, 659
ND4l, 657, 659
ND5, 656–657, 659
ND6, 406, 657–659
 Nearest Neighbor Interchange (NNI), 120, 698, 700
 branch-swapping, 371
 Near-optimal trees, 696
 Neighbor joining (NJ), 121, 229–238, 368, 371–372, 506, 697, 699–702
 analysis, 463, 530, 532
 methods, 290, 293
 tree, 211, 370, 372, 455, 462–463, 508, 702, 801
 Nematocera, 754
 Nematodes, 430
 Nemertean, 430
 Nembrinidae, 746, 753–754
 Nembrinids, 753
Neoetosauroides, 241, 243, 248, 250–251
Neoceratina, 2
 Neogene, 206
 Neologisms, 515, 517
Neorhynchocephalus volaticus, 756
Neotragus pygmaeus, 221
Neotrigonia, 559
Nephila plumipes, 300
 Nepokroeff, Molly, Kenneth J. Sytsma, Warren L. Wagner, and Elizabeth A. Zimmer, Reconstructing ancestral patterns of colonization and dispersal in the Hawaiian understory tree genus *Psychotria* (Rubiaceae): a comparison of parsimony and likelihood approaches, 820–841
Neriene, 72, 77, 82
 radiata, 73–74, 77, 88
 variabilis, 73–74, 77, 88
Nerita, 317

albicilla, 316
Nerodia, 447, 451
 fasciata, 440
 rhombifera, 440
 taxispilota, 440
Nesobasis, 91, 94, 96
Nesomantis thomasseti, 287
 Nested-clade analysis (NCA), 159–185
 Nested model, 606
 Newton–Raphson
 algorithm, 699
 method, 698, 703
 New World, 207, 548–549, 552
 Crocodylus, 409
 ground doves, 39
 lizards, 160
 New Zealand pigeon, 45
Nicotiana, 347
 Nielsen, Rasmus, *see* Huelsenbeck, John P., 131–158
Noctilio, 471
 albiventris, 464
 leporinus, 464
 spp., 463
 Noctilionoidea, 460, 464
 Nonhomogeneous maximum likelihood analysis, 390
 Nonparametric
 bootstrap, 506, 594, 665–667, 669, 671–672, 726, 760, 790, 801
 bootstrap analysis, 165
 rate smoothing method (NPRS), 76
 sign-rank test, 16
 test, 781
 Nontrivial tree, 485
 Norell, Mark, *see* Gatesy, John, 403–422
Nosomma, 305
 monstrosum, 299, 305
Nothofagus, 160
 Novacek, Michael J., *see* O’Leary, Maureen A., 853–866
 Novel versus unsupported clades: assessing the qualitative support for clades in MRP supertrees, Olaf R. P. Bininda-Emonds, 839–851
Novosieversia glacialis, 382
 nrDNA, 336
Nucella, 631
 Nuclear
 alleles, 41, 103
 DNA, 37, 89, 95, 99, 103, 221, 374–375, 382–383, 407–408, 415, 418, 423, 509, 571–593, 735, 757–758, 760–762, 764, 767, 770–771, 773, 775–776
 genes, 16, 28–29, 32, 41, 74, 375, 386, 572
 genome, 374–375, 571, 604
 sequence, 308
 substitution model, 215, 287, 377, 604–617, 658, 674–683, 708
 Nucleotides, 122, 124, 126, 296, 300–301, 303, 692
 data, 305
 sequence, 156, 697, 747
 substitution, 121, 135, 375, 445, 465, 478, 480, 621, 651, 655, 750
 substitution model, 377, 604–617, 658
 Null
 distribution, 30, 761
 hypothesis, 16, 28, 241, 249, 278–279, 363–364, 425, 464, 478, 480, 598, 600, 607, 609, 611, 613, 718, 720, 751, 761, 770, 775, 781–784, 788, 790–792, 801
 model, 271–280, 480, 482, 484, 717–723
 Nyctaginaceae, 334, 344, 346–348
 Nyffeler, Reto, *see* Mast, Austin R., 271–280
 Nymphalid butterflies, 28

- Occam's window, 50
Occidozyga alevis, 719, 815
Ochotona, 609
 princeps, 606
 Odocoileini, 209
Odocoileus, 209, 212, 215, 218–219
 hemionus, 210
 virginianus, 210
 Odonata, 89
Oena capensis, 38, 46
 Okapi, 206, 210
Okapia, 207, 209, 212, 218–219, 221–222, 224
 johnstoni, 210
 Old World, 206–207, 548–550, 552
 Arenaviridae, 740
 badgers, 577, 590
 O'Leary, Maureen A., John Gatesy, and Michael J. Novacek, Are the dental data really at odds with the molecular data? Morphological evidence for the whale phylogeny (re)reexamined, 853–866
 Oligocene, 224–225, 795
 early-, 206–207, 216–218, 224
 late-, 206–207, 216–218, 225
 Oligocephalus, 492
 Olmstead, Richard G., *see* Scotland, Robert W., 539–547
 Onagraceae, 347
Oncostylus, 382
 cocklaynei, 381
 leiospermus, 381
 One-tailed test, 28, 339
 Ontogeny, 430, 515–527
 Onycophora, 557
Opsacas minuta, 316
 Operational taxonomic units (OTU), 291, 620
Ophidiocephalus, 757–758, 764, 767, 770–771
 taeniatus, 757, 759
Ophiophagus, 451
 hannah, 441
 Opilioacarid, 300, 309
 Opilioacarida, 300
 Opilioacariformes, 300
 Opiliones, 557
 Opossum, 285, 656
 Opsin, 28–30
 Optimal tree, 4, 8, 17, 20, 121, 128, 320, 341, 357, 696, 760, 850
 Orangutan, 52–53, 794
 Orchidaceae, 346
Oreamnos, 225
Oreocalamus, 447, 451, 457
 hanitschi, 440
Oreotragus, 221
 Ornithodorinae, 300
Ornithorhynchus anatinus, 285
Oropsylla silantiewi, 187
Orsonwelles, 70–88, 103, 833
 ambersonorum, 73, 84, 88
 arcanus, 73, 79, 84, 88
 bellum, 71, 73, 88
 calx, 73, 84, 88
 falstaffius, 73, 79, 84, 88
 graphicus, 71, 73, 79, 84, 88
 indicium, 71, 73, 79, 88
 macbeth, 73, 77, 79, 84, 88
 malus, 71–73, 77, 79, 81, 88
 othello, 73, 77, 79, 84, 88
 polites, 72–73, 79, 84, 88
 torosus, 71, 76–77, 79, 81–82, 88
 ventus, 73, 79, 88
Ortalis,
 cinereiceps, 388
 guttata, 286
 Orthoptera, 307
Orthotylus, 103
Oryctolagus cuniculus, 286, 289, 606
 Osteoderms, 239–252, 253–255, 851–852
Osteolaemus, 386, 391, 409
 tetraspis, 388, 391, 405, 408–409, 415
 Osteological characters, 760
 Ostrich, 388
 Otariidae, 846
Otobius megnini, 300
Otospermophilus, 197
 variegatus, 189
 Otters, 571–573, 590
 Asian small-clawed, 573
 Cape clawless, 573
 Eurasian, 573
 giant, 573
 marine, 573
 Neotropical, 573
 North American, 573
 river, 577, 590
 sea, 573
 spotted-necked, 573
Ourebia ourebi, 221
 Outgroup, 9, 27, 39, 48, 52, 56, 60–61, 81–82, 88, 89, 92, 94, 96, 99, 102, 189, 192, 196, 205, 229–230, 232, 235, 237, 298, 301, 309, 386–387, 391–392, 404, 415, 443, 446, 463–464, 557, 574, 576, 618, 620–622, 660–661, 674, 707, 756–757, 762, 764, 767, 770–771, 775, 798–799, 815, 821, 826–827
 Outgroup misplacement and phylogenetic inaccuracy under a molecular clock—a simulation study, B. R. Holland, D. Penny, and M. D. Hendy, 229–238
Ovibos, 225
Ovis, 209, 215, 218, 225
 aries, 210, 212
 Owl monkeys, 738
 Oxelman, Bengt, *see* Erixon, Per, 665–674
Oxylipeurus chiniri, 39

Pachygaster leachii, 756
 Paedomorphic salamanders, 503
 Page, Roderic D. M., *see* Johnson, Kevin P., 37–47
 Pairwise tests, 339, 344
Palaeobolbomyia, 746, 750
 Palaeoheterodonta, 559–560
Palaeoplatypygus, 749
 Paleocene, 551
 late-, 550
 Paleontological analysis, 605
 Paleontology, 206–207, 265
Paleosuchus, 391, 409
 palpebrosus, 388, 405
 trigonatus, 405
Pan, 706
 Pantophthalmidae, 756
Pantophthalmus sp., 756
Papilio, 589
Paradelma, 758, 775
 orientalis, 758, 761
Paradialictus, 24, 26
Paralamyctes, 559

- Paralictus, 24
- Paralinear distances (*see also* LogDet/paralinear distance), 287, 291
- Paraleucilla*, 314–315, 320, 322, 324–325, 332
sp., 316
- Parallel evolution, 296, 307
- Parametric
bootstrap, 18, 598, 667, 726, 757–780
bootstrap test, 607, 802, 807, 809, 812
test, 790
- Paraplatypeza atra*, 756
- Parasites, 26, 37–43, 45–46, 406, 735, 737–738, 740–742
phylogeny, 740–741
tree, 740, 742
- Parasitiformes, 299
- Parasitoids wasps, 430
- Parasphcodes, 24
- Paratomistoma*, 399
- Paratyphothorax*, 241, 243, 248–252
- Paraxonia, 853, 861–862
- Parazoanthus*, 317
axinellae, 316
- Pareas*, 451
nuchalis, 440, 446
- Pareatinae, 439–440, 447, 452, 455
- Parisitological characters, 738, 742
- Parrots, 549–553
- Parsimony (*see also* maximum parsimony), 19, 23, 39, 41, 76, 122, 124, 126, 131–132, 140, 147, 236, 258, 265, 335, 362–367, 386, 394, 396, 501, 556, 584, 604, 632, 641–648, 656, 696, 750, 761, 809, 820, 827, 832, 834–835, 841
algorithms, 867
analysis, 7, 26, 30, 42, 45, 70, 76, 79, 83, 249, 251, 255, 257, 260, 262, 271–280, 336, 377, 389–390, 392, 406, 412, 423, 506, 508, 530, 554, 571, 575, 583, 618, 620–621, 623, 629, 727, 764, 802, 804, 824
analysis of endemcity (PAE), 271–280
Camin–Sokal, 444, 446
cladistic, 356, 365
equally weighted, 620–621
Fitch, 444, 446
informative characters, 4, 323, 345, 446, 451, 529, 531, 773
informative indels, 823
informative sites, 28, 30, 613
jackknife, 128
jackknife support, 59, 557, 559
linear, 342
matrix, 241
matrix-weighted, 623
model, 361, 363
permutation tail probability (PTP), 250, 252
ratchet 368–373
reconstruction, 629
search, 368, 842
squared-change, 337, 342
support, 623
trees, 9–10, 39, 42–43, 79, 286, 645, 762, 764, 771, 773, 802–804
uniformative characters, 298
weighted, 284
- Parsons, Thomas J., *see* Harshman, John, 386–402
- Partition analysis, 4
- Partitioned
analysis, 396, 604–607
branch support (PBS), 15–22, 76, 79, 307, 407, 409, 412
likelihood analysis, 386–402
likelihood support (PLS), 15–22,
- Partitioned likelihood support and the evaluation of data set conflict,
M. S. Y. Lee and A. F. Hugall, 15–22
- Partition homogeneity test (PHT; *see also* incongruence length difference [ILD] test), 6, 15, 41–42, 45, 344, 347
- Passeriformes, 552
- Patagioenas*
fasciata, 38
plumbea, 38
speciosa, 38
subvinacea, 38
- PcP, 735–744
- p*-distances, 5
- Pecora, 206–207, 209, 213, 218, 221–222, 224–225
- Pedetes*, 613
capensis, 606
- Pedetidae, 604–606, 610–611, 613–614
- Pelagomapouria*, 826, 829, 832–834
- Pelecorhynchidae, 753, 756
- Pelecorhynchus personatus*, 756
- Penny, D., *see* Holland, B. R., 229–238
- Peracarida, 430
- Percidae, 488
- Percina*, 491
caprodes, 497, 500
nigrofasciata, 491, 497, 500
ouachitae, 497, 500
roanoka, 491, 497–498, 500
sciera, 491, 497, 500
s.l., 494, 497
- Performance-based selection of likelihood models for phylogeny
estimation, Vladimir Minin, Zaid Abdo, Paul Joyce, and Jack Sullivan, 674–683
- Pericharax*, 314–315, 332
- Periods,
cenozoic, 206, 631, 746
cretaceous, 23, 33, 403, 419, 550–552, 631, 746, 753, 861
early Cretaceous, 548–549
early Eocene, 795
early Miocene, 206–207, 209, 217–218, 224–225
early Oligocene, 206–207, 216–217, 224–225
eocene, 222, 403, 419, 857–858
jurassic, 746, 751, 753
late Cretaceous, 419, 549–552
late Eocene, 222, 224
late Jurassic, 549, 746
late Miocene, 218, 224
late Oligocene, 225
late Paleocene, 550
late Pleistocene, 473
late Triassic, 549, 745, 751, 754
lower Cretaceous, 753
lower Jurassic, 746, 749
lower/middle Triassic, 746
mesozoic, 549–553, 745–746
mid-Cretaceous, 549, 552
mid-Eocene, 218, 222, 224–225
mid-Jurassic, 745
mid-Miocene, 206–207, 209, 217–218, 224–225
mid-Pleistocene, 198, 460, 473, 810
mid-Tertiary, 549
mid-Triassic, 548
miocene, 207, 221, 451, 549, 551, 629, 761, 776, 808
neogene, 206
oligocene, 224–225, 795
paleocene, 551
pleistocene, 186, 198, 460–476, 777, 797, 807–808, 812
pliocene, 224–225, 629
plio-pleistocene, 225

- quaternary, 552
 tertiary, 549–553, 746, 753
 triassic, 239–252, 403, 749, 751
 upper Cretaceous, 753
 upper Jurassic, 753
 upper Triassic, 753
Perissodactyla, 850, 857–858
 Permutation tail probabilities (PTPs), 250, 252
Petrobiona, 314–315, 317, 320, 322–324, 328–329, 332–333
 massiliana, 316
Petrobionidae, 314
Petromarmota, 186, 192–195, 197–198
 caligata, 189
 flaviventris, 189
 olympus, 189, 193
 vancouverensis, 189
Petrostoma, 329
Phapitreron
 amethystina, 38
 leucotis, 38
 Phenotypes, 241, 349
 Philippines, 794–819
 Philosophy, 352–367
 of science, 259–271, 352
Philothamnini, 457
Phocidae, 846–847
Phthiraptera, 37–47
Phyllostomoidea, 460
 Phylogenesis, 520
 Phylogenetically informative characters, 55–56, 826
 Phylogenetic
 accuracy, 121, 528–538
 artifacts, 594
 clade, 714
 comparative method (PCM), 684–695
 congruence, 165
 conservatism, 1
 data, 89, 248–249, 314, 649, 724, 726, 745, 747
 distance, 491
 effect, 488–500, 684, 686, 692, 694
 eigenvector regression (PVR), 694
 error, 119, 121, 124–125
 estimates, 159, 284, 530, 802, 850
 evidence, 503
 framework, 311, 728
 generalized least squares (PGLS),
 history, 91, 131–132, 337, 348–349, 501, 589–590, 641, 745
 hypothesis, 1, 55, 62, 212, 268, 283, 352, 354, 362–363, 401–406, 445,
 477, 488, 554, 556–557, 571, 632, 672, 689, 693, 725–726, 728, 740,
 742, 745, 764, 854, 858, 860–861
 inference, 75, 180, 187, 296, 306, 335, 344, 352, 359, 368–373, 463, 511,
 554, 612, 641, 696, 791
 information, 96, 296, 311, 327, 339, 589, 686, 747, 761
 level, 542
 markers, 864
 methods, 131, 229, 283, 285, 291, 293, 352, 358, 469, 480, 641–642,
 665, 677
 model, 133, 594, 649, 651, 654, 656, 661
 patterns, 342–344, 348
 reconstruction, 19, 39, 83, 287, 318, 375, 539–548, 571, 658, 665, 669,
 671, 685, 724–729, 860–861, 864
 relationships, 2, 23, 35, 71, 92, 102, 159, 179, 181, 186, 199, 213, 215,
 220, 239, 245, 255, 265, 335–336, 347, 439, 444, 447, 451, 471, 477,
 548, 554, 591, 623, 728, 745, 775, 797, 828
 signal, 39, 55, 63, 70, 79, 83, 186, 197, 199, 328, 334, 371–372, 403, 412,
 415–416, 445, 497, 511, 577, 581, 584, 591, 773, 853–855, 857, 860
 species, 833
 structure, 178, 181, 473, 686–687, 691
 studies, 32, 110, 188, 374–375, 545, 757–758, 821–822
 supertrees, 839–851
 systematics, 260, 510, 735
 taxonomy, 852
 tests, 781
 tools, 311
 tree, 119, 121, 126, 71, 112, 131, 134–135, 244, 289–290, 359, 364–365,
 427, 463, 515, 554–555, 669, 671, 685, 692, 715, 749, 760–761, 771,
 791, 824, 847, 849
 uncertainty, 39, 132, 488
 utility, 27, 30, 32
 Phylogenetic methods and Aetosaur interrelationships: a rejoinder,
 Simon R. Harris, David J. Gower, and Mark Wilkinson, 851–852
 Phylogenetic relationships among Calyptraeid gastropods and their
 implications for the biogeography of marine speciation, Rachel
 Collin, 618–640
 Phylogenetic systematics and evolution of primate-derived
 Pneumocystis based on mitochondrial or nuclear DNA
 sequence comparison, Jean-Pierre Hugot, Christine
 Demanche, Véronique Barriol, Eduardo Dei-Cas, and Jacques
 Guillot, 735–744
 Phylogenetics, 1–14, 48, 70–88, 95, 127, 136, 243, 247, 270, 334–351,
 352–367, 386, 477–487, 515–527, 557, 565, 594–603, 655, 665,
 782–783, 794, 847, 862
 Bayesian, 48, 54, 89–109
 molecular, 187, 283–295, 311, 715
 vertebrate, 388
 Phylogenetics of advanced snakes (Caenophidia) based on four
 mitochondrial genes, Christopher M. R. Kelly, Nigel P. Barker,
 and Martin H. Villet, 439–459
 Phylogenetics of fanged frogs: testing biogeographical hypotheses at
 the interface of the Asian and Australian faunal zones, Ben J.
 Evans, Rafe M. Brown, Jimmy A. McGuire, Jatna Supriatna,
 Noviar Andayani, Arvin Diesmos, Djoko Iskandar, Don J.
 Melnick, and David C. Cannatella, 794–823
 Phylogenomics, 119–124
 Phylogeny, 8–9, 23–36, 37–47, 48–54, 57, 60–61, 63, 66, 89, 92, 111, 114,
 121, 133, 136, 139, 152, 159–185, 186–205, 206–228, 229, 239,
 253–255, 264, 272, 280, 296–310, 311–333, 334–335, 345, 348, 368,
 374–385, 386–402, 430, 460–461, 471, 488, 493, 497–498, 501–514,
 515, 520–521, 525–526, 539, 543, 548, 571–593, 604–617, 632, 645,
 647, 649–664, 665–673, 674–683, 687, 692–693, 696–704, 714, 724,
 726, 728, 735, 737, 740, 751, 757–780, 792, 798, 801–802, 820, 855,
 864
 aetosaur, 852
 cetacean, 854
 chloroplast, 375
 estimates, 423–428, 674, 677, 682
 fly, 746
 gene, 374
 host, 740, 742
 hypotheses, 738
 mammalian, 857
 molecular, 37–47, 206, 545, 781, 832
 morphological, 206
 muroid, 613
 parasite, 740–741
 Pneumocystis, 737, 740
 reconstruction, 264, 349, 501, 511, 542–543, 545, 586
 rodent, 613
 tetrapod, 283–295
 whale, 853
 Phylogeny and evolution of Calcareous sponges: monophyly of
 Calcinea and *Calcaronea*, high level of morphological

- homoplasy, and the primitive nature of axial symmetry, Michael Manuel, Carole Borchellini, Eliane Alivon, Yannick Le Parco, Jean Vacelet, and Nicole Boury-Esnault, 311–333
- Phylogeny of eusocial *Lasioglossum* reveals multiple losses of eusociality within a primitively eusocial clade of bees (Hymenoptera: Halictidae), Bryan N. Danforth, Lindsay Conway, and Shuqing Ji, 23–36
- Phylogeny reconstruction: the role of morphology, Robert W. Scotland, Richard G. Olmstead, and Jonathan R. Bennett, 539–547
- Phylogeographic
analysis, 460
structure, 460–476
- Phylogeography, 92, 801, 812
- Phylogram, 100, 140, 621
- Phymaturus indistinctus*, 160, 164
- Physeter*, 706, 712
- Phytosaurs, 253
- Pianka, Eric R., *see* Jennings, W. Bryan, 757–780
- Pichia anomala*, 316
- Picoides pubescens*, 286
- Pigeons, 37, 41, 45
New Zealand, 45
- Pigs, 854
- Pinworm, 742
- Pithecia*, 737, 740, 742
pithecia, 736
- Pityohyphantes*, 77
costatus, 73, 88
- Placental mammals, 591
- Placozoa, 31
- Plagithmysus*, 89
- Plankton, 618
- Plants, 89, 106, 335, 347, 517, 541, 696, 746, 794, 820
flowering, 375, 518, 753, 820
seed, 542, 781
- Plasmid DNA, 316
- Platodes, 520
- Platymantinae, 794
- Platypezidae, 756
- Platypodid beetles, 23
- Platyrrhini, 738, 742, 735, 737, 741
- Plecoptera, 106
- Plectroninia*, 314–315, 317, 323, 328–329, 332
halli, 329
- Pleistocene, 186, 198, 777, 797, 807–808, 812
divergence, 460–476
late-, 473
mid-, 198, 460, 473, 810
- Plesiometacarpalia, 207, 215
- Plethodontid salamanders, 510
- Pletholax*
gracilis, 757, 759
hortulanus, 761
- Pliocene, 629, 629
- Plio-pleistocene, 225
- Pneumocystis*, 735–744
carinii, 735–736
carinii aotus, 736, 738, 742
carinii callimico, 736, 738
carinii callithrix, 736, 738, 740
carinii fascicularis, 736
carinii fuscicollis, 736, 738
carinii geoffroyi, 736, 738, 740
carinii griseus, 736
carinii hamlyni, 736
carinii imperator, 736, 738
carinii macaco, 736
carinii midas, 736, 738
carinii mulatta 1, 736
carinii mulatta 2, 736
carinii muris, 736
carinii nemestrina, 736
carinii nictitans, 736
carinii nigroviridis, 736
carinii oedipus, 736
carinii pithecia, 736, 738, 742
carinii sciureus, 736, 738, 742
DNA, 736
genes, 742
genotypes, 735
jiroveci, 736
phylogeny, 737, 740
pneumonia (PcP), 735–744
spp., 735, 738, 742
- Poe, Steven, Evolution of the strategy of long-branch subdivision to improve the accuracy of phylogenetic methods, 423–427
- Poe, Steven, *see* de Queiroz, Kevin, 352–367
- Poisson
distribution, 133
process model, 134
- Polhemus, Dan, *see* Jordan, Steve, 89–109
- Polinices didyma*, 640
- Pollock, David D., *see* Hillis, David M., 124–126
- Polychaetes, 430
- Polymerase chain reaction (PCR), 26, 39, 74, 94, 301, 316, 376, 383, 389, 404, 571, 574, 591, 606, 620, 736, 738, 758, 823
amplification, 2–3, 161, 375, 389, 736, 747
recombination, 384
- Polymorphic characters, 505, 833
- Pongo*, 706
- Pontederia*
cordata var. *cordata*, 154, 156
cordata var. *lancifolia*, 154, 156
cordata var. *ovalis*, 154, 156
rotundifolia, 154, 156
sagittata, 154, 156
- Pontederiaceae, 139–140, 143, 147, 154, 156
- Popper and systematics, Olivier Riappel, 259–270
- Popper, Karl, 259–271, 352–367
- Popper's degree of corroboration, 352, 355, 360, 363, 366
- Porifera, 311–333
- Potent antiretroviral therapy (PART), 660–661
- Prairie dog, 186,
- Prestosuchidae, 252
- Priapulus caudatus*, 316
- Price, Samantha E., *see* Bininda-Emonds, Olaf R. P., 724–729
- Primary characters, 725–726
- Primates, 52–53, 57, 549, 606, 707–708, 735–744
- Principle component analysis (PCA), 687
- Principle coordinates analysis, 860
- Pristichampsus*, 409
- Probability, 352–367
- Procervulus*, 224
- Procrustes, 66–69
distance, 67, 190, 192, 493
generalized analysis, 66, 68
geometry, 67–68
- Procyonidae, 573–574, 846
- Procyoninae, 573
- Procyon lotor*, 573–574

- Progressive rearing, 1–14
 Prokaryotes, 658–660
Propithecus, 706
 Prosobranchia, 328
 Prostriata, 296, 300, 304, 307
 Protein-coding, 391
 genes, 23, 39, 74, 94, 297–298, 307, 358, 396, 405, 445, 460, 471, 506,
 604, 609, 612, 656, 682, 705–706
 regions, 52, 165
 Protein evolution, 602
 Protists, 429–430
Protragocerus labidotus, 221
 Prunoideae, 378–379, 382
psaA, 785, 787, 791
psaB, 785, 787–788, 791
 Psammophiinae, 439–440, 452, 455
Psammophis, 451
 condenarus, 441
 mossambicus, 441, 459
Psammophylax, 451
 variabilis, 441, 459
Pseudagrion, 91, 94, 102
 civicum, 94
 pacificum, 94
 palauense, 94
Pseudemys scripta, 286–287
Pseudochilalictus, 24
 Pseudoboini, 456
 Pseudoreplicate, 55–65
 Pseudoxyrophiinae, 439, 441, 443, 452, 455–456
 Psittacidae, 551
 Psittiformes, 551
 Psychodomorpha, 756
Psychomyia,
Psychotria, 820–841
 cadigiensis, 821–822
 fauriei, 822, 826, 829, 833–834
 grandiflora, 822, 826, 828–829, 832, 834
 greentwelliae, 820, 822–823, 825–827, 829, 832–834
 hathewayi, 822, 826, 829, 833–834
 hawaiiensis subsp. *hawaiiensis*, 829
 hawaiiensis var. *hawaiiensis*, 822, 829, 833
 hawaiiensis var. *hillebrandii*, 822–823, 829
 hexandra, 822, 828–829, 832, 834
 hexandra subsp. *O'ahuensis*, 822
 hobdyi, 822, 828–829, 832, 834
 hombroiana, 821–822
 kaduana, 822, 829, 831, 833
 luzionensis, 821–822
 mariniana, 822, 825, 827, 829, 832–834
 mauiensis, 822, 829, 833
 pickeringii, 821–822
 sp. Fiji, 821–822
 sp. Guam, 822
 sp. Tinian Is., 821
 tahitiensis, 821–822, 826
 wawrae, 822, 829, 834
 Pteriomorpha, 561
Pteronura, 590
 brasiliensis, 581, 583, 589–590
 Pterygota, 328
Ptilinopus occipitalis, 38
Ptiolina fasciata, 756
 Punctuational evolution, 831
 Pupal characters, 8
 Purvis, Andy, *see* Bininda-Emonds, Olaf R. P., 724–729
 P-values, 18, 22, 28, 99, 132, 137, 147, 156–157, 190, 192, 273, 279, 345,
 394–395, 406, 425, 598–600, 692, 722, 761, 770, 783–785, 787–788,
 790–791
 adjustment, 781–793
 Pycnogonida, 557
 Pygopodidae, 757–780
 Pygopodid
 lizards, 757–780
 phylogeny, 758, 764
 tree, 757, 760, 770
Pygopus, 757–758, 762, 764, 767, 770–771, 775
 nigriceps, 759–761, 775
 (*Paradelma*) *orientalis*, 759
 schraderi, 758
 stealescottii, 758
 Pythonidae, 440
Python reticulatus, 440
 Quail, 551
 Qualitative support (QS) index, 839, 841, 844, 847
Quamoclidion, 336
 Quartet puzzling (QP) values, 558
 Quaternary, 552
 Rabbits, 289
 Raccoon, 573
 RAG-1, 405, 408
 RAG-2, 604, 606–610, 613–614
 Raguso, Robert A., *see* Levin, Rachel A., 334–351
Rana
 nigromaculata, 798
 signata, 810
 vittigera, 815
 Random
 generation, 110–118
 trees, 30, 115, 445, 681
Rangifer, 209, 212, 215, 218–219
 tarandus, 210
Rangiferini, 209
Ranodon sibiricus, 286
 Ranunculaceae, 347
Ranunculus, 347
 Rare genomic changes, 296
 Rate
 distribution, 594–603
 homogeneity model, 463
 of evolution, 602, 641, 646–647, 677, 746, 753, 833
 Rates-across-sites model, 595–596
 Ratite, 290
 birds, 549–553
 Rats, 290
Rattus norvegicus, 286, 606, 736
Rauisuchia, 241–243, 251
rbcl, 140, 336, 379, 696–704
 rDNA, 405, 408, 808, 820
 Recapitulation, 515–527
 Recognition, 110–118
 Reconciliation analysis, 39, 42
 Recursion, 110–118
 Reconstructing ancestral patterns of colonization and dispersal in the
 Hawaiian understory tree genus *Psychotria* (Rubiaceae): a
 comparison of parsimony and likelihood approaches, Molly
 Nepokroeff, Kenneth J. Sytsma, Warren L. Wagner, and
 Elizabeth A. Zimmer, 820–841
 RDPII project, 696–704
 Red deer, 210

- Redondasuchus*, 241–243, 248–252
 Red panda, 846
 Reductive characters, 245
Redunca, 209, 212, 215
 fulvorufula, 210
 Reduncini, 209–210
 Redundancy analysis (RDA), 687–689, 693
 Reed, David L., *see* Cummings, Michael P., 477–487
 Regression, 66–69
 analysis, 68
 model, 684, 686
 Regularia, 328
 Reindeer, 210
Reithrodontomys sumichrasti, 677
 Relative warp analysis (RWA), 198
 Reliability of Bayesian posterior possibilities and bootstrap frequencies in phylogenetics, Per Erixon, Bodil Svennblad, Tom Britton, and Bengt Oxelman, 665–673
 RELL bootstrapping, 390
 Remipedia, 557
 Reptiles, 524
 Reptilia, 239, 253
 Rescaled consistency index, 762, 764
 Retention index, 3, 5–6, 10, 28, 56, 76, 217, 304, 339, 345–346, 348, 577–578, 581–583, 762, 764, 826
 Reticulate evolution, 374–385
 Reweighted characters, 318, 327, 371
 Rhagionidae, 746, 756
Rhagoletis, 754
 Rhamnaceae, 375
Rhammus, 378–379
 cathartica, 377
 Rhamphiophis, 451
 acutus, 441, 459
 oxyrhynchus, 441
Rhamphotyphlops braminus, 440
Rhea americana, 550
 Rheas, 550
Rhingia nasica, 756
 Rhinoceros, 706–708, 794
Rhipicentor, 305–306
 nuttalli, 299
 Rhipicephalinae, 297–298, 305–307
 s.l., 296, 299, 304–305, 307, 309
Rhipicephalus, 299, 306
 appendiculatus, 299, 306, 310
 compositus, 299
 evertsi, 299, 310
 maculatus, 299
 pravus, 299, 310
 pulchellus, 299, 306, 310
 pumilio, 306
 punctatus, 299
 sanguineus, 299, 306, 310
 simus, 299
 turanicus, 299, 306
 zambeziensis, 299
 zumpti, 299
Rhizomys, 613
 pruinus, 606
 Rhynchosaur, 253
 Rieppel, Olivier, Popper and systematics, 259–270
 Ringtail, 573
 Rivulidae, 548, 552
Rivulus marmoratus, 549
 RNA, 555, 610
 mitochondrial, 445
 polymerase II, 557
 secondary structure, 283–295
 sequencing, 49, 651
 Rock squirrel, 194, 197
 Rodent, 656, 679–680
 Holarctic, 186
 phylogeny, 613
 tree, 615
 Rodentia, 186, 604–617
 Rodman, James E. and Jeannine H. Cody, Taxonomic impediment overcome: NSF's Partnerships for Enhancing Expertise in Taxonomy (PEET) as a model, 428–435
 Roger, Andrew J., *see* Susko, Edward, 594–603
 Rohlf, F. James and Fred Bookstein, Computing the uniform component of shape variation, 66–69
 Rokas, Antonis, *see* Cummings, Michael P., 477–487
 Rooted
 duplication tree (RDT), 110–111, 113, 115
 tree, 865
 Rooting, 386–402
 Roots of phylogeny: how did Haeckel build his trees?, Benoit Dayrat, 515–527
Rosa, 375
 multiflora, 375–377
 Rosaceae, 374–377, 379, 382
 Rosenberg, Michael S. and Sudhir Kumar, Taxon sampling, bioinformatics, and phylogenomics, 119–123
 Rosoideae, 378–379
 rRNA, 131, 283–285, 291, 293, 297, 307, 615
 Rubiaceae, 820
Rubus, 375
 odoratus, 376–377
 Ruminantia, 206–228
Rupicapra rupicapra, 221
 Rutaceae, 347

 Sable antelope, 210
Saccharomyces cerevisiae, 316
Saguinus, 740, 742
 fuscicollis, 736–737
 imperator, 736
 midas, 736, 738
 oedipus, 736
Saimiri, 740, 742
 sciureus, 736
 Salamanders, 287, 501–514
 cave, 501
 hemidactyliine, 509
 paedomorphic, 503
 plethodontid, 510
 Sampling design, 159–185
 Sampling strategies for delimiting species: genes, individuals, and populations in the *Liolaemus elongatus-kriegi* complex (Squamata: Liolaemidae) in Andean–Patagonian South America, Mariana Morando, Luciano J. Avila, and Jack W. Sites Jr., 159–185
 Sandvik, Hanno, Genes, categories, and species: the evolutionary and cognitive causes of the species problem (rev.), 565–567
 Sarcoscyphinae, 430
Sarona, 833
 Savage–Dickey ration, 48, 50, 53
 Scale, 271–280
 of being, 515–527
Sceloporus jarrovi, 160
 Scenopinidae, 750, 756

- Scent, 334–351
- Schiedea*, 820, 832
- Schistosoma*, 307
- Schizophora, 746, 749–751, 753–754
- Schwarz, Michael P., Nicholas J. Bull, and Steven J. B. Cooper,
Molecular phylogenetics of Allodapine bees, with implications
for the evolution of sociality and progressive rearing, 1–14
- Scincidae, 757
- Sciuridae, 186, 605–606, 608, 614–615
- Sciurid convergence, 186–205
- Sciurids, 188, 197
- Sciurini, 189
- Sciuroidea, 604–606, 608–610, 613, 615
- Sciuromorpha, 605, 614
- Sciurus*
infuscatus, 205
vulgaris, 189, 205
- Scobius*, 271–280, 717–723
aciculatifrons, 273
cognatus, 273
griseus, 273, 275
horni, 275
obesus, 275
oneill, 275
pullus, 275
viridis, 275
- Scolecophidia, 440
- Scolopendra cingulata*, 316
- Scorpiones, 300, 556–557
- Scorpionoidea, 556
- Scotland, Robert W., Richard G. Olmstead, and Jonathan R. Bennett,
Phylogeny reconstruction: the role of morphology, 539–547
- Scypha*, 325
ciliata, 317
- Sea anemones, 430
- Seastar, 139–140, 143
- Sea urchins
Echinometra, 631
- Seed plants, 542, 781
- Selenodontia, 221
- Sellalictus*, 24, 26, 30
- Semicassis pyrum*, 640
- Sensitivity analysis, 554–564
- Sequence analysis, 737
- Sequence-tagged sites, 571–593
- Serinus canaria*, 389
- Sex ratio, 24, 33
- Shape statistics, 66–69
- Sheep, 206, 210, 854, 857
- Shell characters, 619, 623
- Shimodaira–Hasegawa test, 15–22, 96, 98, 212, 216, 221, 444–445, 464,
468, 504, 508, 781–793
- Short-tailed bat, 460
- Shrimp, 23
- Synalpheus*, 631
- Sicista*, 614
tianshanica, 606
- Sieversia*, 374
- Sigapatella*, 619, 632, 639
novaezealandiae, 639
terraenovae, 639
- Significance test, 781–793
- Silicispongia, 320
- Silverswords, 89
- Simimeryx*, 222
- Simon, Chris, *see* Jordan, Steve, 89–109
- Simple, fast, and accurate algorithm to estimate large phylogenies by
maximum likelihood, Stéphane Guindon and Olivier Gascuel,
696–704
- Simulation (*see also* computer simulation and Monte Carlo
simulation), 423–428, 665–673
- Single-copy genes, 375
- Single-gene tree, 20
- Single-valve decomposition (SVD), 68–69
- Sinsheimer, Janet S., *see* Suchard, Marc A., 48–54, 649–664
- Siphopatella*, 619, 623, 639
walshi, 623, 639
- Sites-specific rates (SSR)
analysis, 612
model, 4, 8, 26–28, 30, 463, 466–468, 610, 612, 674
tree, 610–611
- Sites, Jack W., Jr., *see* Morando, Mariana, 159–185
- Skeletal characters, 199, 222, 313, 403
- Skunks, 572–573
- Smedmark, Jenny E. E., Torsten Eriksson, Rodger C. Evans, and
Christopher S. Campbell, Ancient Allopolyploid speciation in
Geinae (Rosaceae): evidence from nuclear granule-bound
starch synthase (GBSSI) gene sequences, 374–385
- Snails, 70
- Snakes, 286–287, 456
African, 456
advanced, 439–459
colubrid, 758
Madagascan, 456
- Snipe flies, 745–746
- Snout characters, 416
- Social
behavior, 23–36
evolution, 1–14, 23–36
- Soft ticks, 300
- Solanaceae, 347
- Soldier flies, 746
- Soleneiscidae, 314
- Soleneiscus*, 314–315, 317, 322, 329, 332–333
radovani, 316
- Sordariales, 430
- Source trees, 727, 840–844, 846
- South Africa, 717–723
- Southern cassowary, 388
- Spalax*, 613
zemni, 606
- Spearman's rank correlation test, 339, 346
- Speciation, 70–88
- Speciation on a conveyor belt: sequential colonization of the Hawaiian
islands by *Orsonwelles* spider (Araneae, Linyphiidae), Gustavo
Hormiga, Miquel Arnedo, and Rosemary G. Gillespie, 70–88
- Species
boundaries, 159–185
radiations, 89–109
trees, 63, 180, 725, 727
- Spermophilus*, 189, 194, 197
beldingi, 189, 204
citellus, 189, 204
eversmanni, 205
grammurus, 204
gravojerici, 204
karamani, 204
richardsoni, 189, 204
tridecemlineatus, 606
undulatus, 189, 204–205
variegatus, 194, 197, 204
- Sphécodogastra*, 24, 32

- Sphenodon punctatus*, 286
 Spiders, 56, 58, 60, 62, 70–88, 300, 430, 833, 557
 ground, 430
 orsonwelles, 70–88, 833
 tetragnathid, 70, 105
 Sponges, 525
 asconoid, 313, 332–333
 calcareous, 311, 320, 328, 330, 332–333
 leuconoid, 313, 325, 329
 syconoid, 313, 329, 332
 sylllebid, 313
 Spongiae, 520
Spongilla, 317
 lacustris, 316
 Squamata, 159
 Squared-change parsimony, 337, 342
 Squirrel monkey, 52–53
 Squirrels,
 ground, 186, 197
 rock, 194, 197
 terrestrial, 187–188
 tree, 188, 197
 Stability, 55–65
 analysis, 554–564
 Stability in phylogenetic formulations and its relationship to nodal support, Gonzalo Giribet, 554–564
Stagonolepis, 247–248
 robertsoni, 241, 243, 250–251
 wellesi, 241, 243, 250–251, 254
 Stalk-eyed flies, 32
 Staphylinidae, 430
 Star
 topology analysis, 481
 tree, 236, 236
 State (see character state)
 Statistical
 analysis, 131, 847
 test, 197
 Staurorrhaphidae, 314, 328, 333
Stenomphrale teutankameni, 756
Stephanocemas, 224
 Strepsirrhine, 711–713
 Stochastic mapping of morphological characters, John P. Huelsenbeck, Rasmus Nielsen, and Jonathan P. Bollback, 131–158
 Stochastic model, 152, 705, 761, 833
 Stratiomyidae, 756
 Stratiomyomorpha, 746, 750–751, 756
Straussia, 820, 826, 828–829, 832–833
 Strepsirhini, 738
Streptopelia
 capicola, 38, 46
 decaocto, 38
 senegalensis, 38, 46
 vinacea, 38, 46
 Strict component consensus (SCC), 241, 243, 248, 250
 tree, 251, 532–533, 535
 Strict consensus, 406
 tree, 57–58, 61, 323, 339, 344, 737, 826–827
 Strict reduced consensus (SRC) method, 240, 243, 247–248, 250
 tree, 250
Strobilanthes, 543
Struthio camelus, 388, 405
 Struthioniformes, 551
Sturnira lilium, 472
Stylipus, 383
Suberites, 317
 ficus, 316
 Substitution
 model, 3, 140, 293, 371, 463, 577
 rates, 471
 Suchard, Marc A., Christina M. R. Kitchen, Janet S. Sinsheimer, and Robert E. Weiss, Hierarchical phylogenetic models for analyzing multipartite sequence data, 649–664
 Suchard, Marc A., Robert E. Weiss, and Janet S. Sinsheimer, Testing a molecular clock without an outgroup: derivations of induced priors on branch-length restrictions in a Bayesian framework, 48–54
Sudila, 24, 32
 Suine artiodactylans, 854
 Sulawesi, 794–819
 Sullivan, Jack, see Minin, Vladimir, 674–683
 Summary characters, 337, 342
 Sum of squared deviations (SSD),
 Supermatrix, 724–729
 analysis, 724, 726, 728, 853
 Supertree, 439–459, 528–538, 724–729, 865
 analysis, 725–726, 728, 853
 carnivore, 845
 clade, 840–841
 consensus, 440, 447
 construction, 728, 839
 MRP, 439, 444
 phylogenetic, 839–851
 support, 55–65, 665–673, 839–851
 Supertrees are a necessary not-so-evil: a comment on Gatesy et al., Olaf R. P. Bininda-Emonds, Kate E. Jones, Samantha A. Price, Richard Grenyer, Marcel Cardillo, Michael Habib, Andy Purvis, and John L. Gittleman, 724–729
 Supriatna, Jatna, see Evans, Ben J., 794–823
 Susko, Edward, Chris Field, Christian Blouin, and Andrew J. Roger, Estimation of rates-across-sites distributions in phylogenetic substitution models, 594–603
 Svennblad, Bodil, see Erixon, Per, 665–674
 Swallowtail butterflies, 589
 Swofford-Olsen-Waddell-Hillis test (SOWH), 607, 611, 615
 Sycetta, 314–315, 332
 Sycettidae, 314
Sycettusa, 314–315, 320, 322–324, 328, 332–333
 sp., 316
Sycon, 311, 313–315, 317, 320, 323, 325, 329, 332
 calcaravis, 316–317, 323, 329
 ciliatum, 316–317, 322–323, 328–329
 ensiferum, 333
 raphanus, 316–317, 322–325, 329–330
 spp., 320
Syconessa, 314–315, 332–333
Sylvicola alternatus, 756
 Symbionts, 735, 743
 Symmetric-difference tree, 463
 Sympatric
 lineages, 46–476
 speciation, 618–640
Symphoromyia hirta, 756
 Symphyla, 557
Synalpheus shrimp, 631
Synechocystis, 659
 Syrphidae, 756
 Systematic characters, 243
 Systematics, 23–36, 55, 259–271, 403, 430, 515–527, 539, 565, 632, 674, 853

- molecular, 430, 501–514, 571–572, 591, 745–756, 820–841
 phylogenetic, 510
- Systematics of the lizard family Pygopodidae with implications for the diversification of Australian temperate biotas, W. Bryan Jennings, Eric R. Pianka, and Stephen Donnellan, 757–780
- Systematic utility of floral and vegetative fragrance in two genera of Nyctaginaceae, Rachel A. Levin, Lucinda A. McDade, and Robert A. Raguso, 334–351
- Sytsma, Kenneth J., *see* Nepokroeff, Molly, 820–841
- Tabanidae, 746, 753–754, 756
- Tabanomorpha, 745–746, 750–751, 756
- Tabanus*, 754
rufograter, 756
- Taihangia rupestris*, 382
- Tamiini, 189, 194
- Tamias*, 189, 205
sibiricus, 205
striatus, 189, 205
- Tamiasciurus hudsonicus*, 606
- Tamura and Nei model, 703
- Tandem duplication trees, 110–118
- Tapirs, 857
- Tarsier, 52–53
- Taxic-homology, 539–548
- Taxidea*, 590
taxus, 571–573, 581
- Taxidiinae, 571–573
- Taxonomic
 congruence, 853, 862
 impediment, 428–435
 sampling, 832
 training, 428–435
- Taxonomic impediment overcome: NSF's Partnerships for Enhancing Expertise in Taxonomy (PEET) as a model, James E. Rodman and Jeannine H. Cody, 428–435
- Taxonomy, 330, 390, 474, 539–548
 Calyptraeid, 619
- Taxon sampling, 71, 119–124, 124–126, 207, 306–307, 423, 442, 446, 451, 539, 542–543, 614, 619, 632–633, 641–642, 644–645, 802, 853
- Taxon sampling, bioinformatics, and phylogenomics, Michael S. Rosenberg and Sudhir Kumar, 119–123
- Tayra, 573
- Tegula*, 619, 631, 633
- Teiidae, 689
- Teinobasis*, 91, 94, 96
palauensis, 102
rufithorax, 102
- Telemetacarpalia, 207, 215, 219
- Templeton tests, 15–22, 339, 344–345
- Termites, 23, 33
- Terrestrial squirrels, 187–188
- Tertiary, 549–553, 746, 753
 mid-, 549
- Testing a molecular clock without an outgroup: derivations of induced priors on branch-length restrictions in a Bayesian framework, Marc A. Suchard, Robert E. Weiss, and Janet S. Sinsheimer, 48–54
- Tests
 approximately unbiased, 781–793
 Bayesian, 607, 615
 chi-square, 307, 395, 397–398, 607
 goodness-of-fit, 464, 471, 674
 hypothesis, 48–54, 781–793
 incongruence length difference (ILD) (see also partition homogeneity), 3, 11, 15, 26, 28, 75–76, 418, 467, 604, 607–609, 613, 622, 773, 853–855, 860, 862
 Kishino–Hasegawa, 4, 10, 15–22, 212, 216, 390, 392, 394–396, 398, 445, 504, 781
 Kolmogorov–Smirnov, 190
 likelihood ratio (LRT), 41–42, 48, 54, 63, 95, 352, 363–364, 377, 379, 390, 394, 445, 575, 580, 598, 600, 604–617, 621, 655, 674–683, 714, 801, 825, 849
 likelihood weight, 18–19, 22
 log-likelihood ratio, 3–4, 377, 379
 Mantel, 491
 maximum likelihood, 614
 Monte Carlo randomization, 684
 nonparametric, 781
 nonparametric signed-rank, 16
 one-tailed, 28, 339
 pairwise, 339, 344
 parametric bootstrap, 607, 802, 807, 809, 812
 parametric, 790
 partition homogeneity (see also incongruence length difference), 6, 15, 41–42, 45, 344, 347
 phylogenetic, 781
 Shimodaira–Hasegawa, 15–22, 96, 98, 212, 216, 221, 444–445, 464, 468, 504, 508, 781–793
 significance, 781–793
 Spearman's rank correlation, 339, 346
 statistical, 197
 Swofford–Olsen–Waddell–Hillis, 607, 611, 615
 Templeton, 15–22, 339, 344–345
 topological ILD, 75–77
 two-tailed, 339
 Wilcoxon rank sum, 337, 345
 Wilcoxon signed rank, 406, 668, 857
- Tetilla japonica*, 316
- Tetracerus*, 221
quadricornis, 221
- Tetraconata, 557
- Tetragnatha*, 103
- Tetragnathidae, 56–57
- Tetragnathid spiders, 70, 105
- Tetramolopium*, 820
- Tetraphyllidea, 430
- Tetrapods, 283, 285
 evolution,
 phylogeny, 283–295
- Thamnophiini, 447
- Thamnophis*, 447, 451
butleri, 440
elegans, 440
- Therevidae, 430, 750, 756
- Thin-plate spline, 189, 192–193, 488–500
- Thomomys bottae*, 606
- Thoracosaurus*, 398, 401, 417–418
- Thorne, Jeffrey L., *see* Wiegmann, Brian M., 745–756
- Three-item statements, 255–259
- Three-parameter TN92 model, 293
- Thrincolhalictus, 24
- Thrips, 23
- Ticks, 296–310
 hard, 296–298, 300–301, 304, 309
 metastrate, 297, 306
 prostrate, 297
 rhipicephaline, 304, 306
 soft, 300
- Tigers, 794

- Time flies, a new molecular time-scale for Brachyceran fly evolution without a clock, Brian M. Wiegmann, David K. Yeates, Jeffrey L. Thorne, and Hirohisa Kishino, 745–756
- Time-homogeneous model, 647
- Tipula (Platytipula) paterifera*, 756
- Tipulidae, 756
- Tipulomorpha, 756
- Toxus nasutus*, 286
- Tomistoma*, 386, 391–392, 394, 396–399
cairensis, 405
schlegelii, 386, 388, 403, 405–406, 408–409, 412, 415–419
- Tomistominae, 403, 405, 416–418
- Topological ILD test, 75–77
- Tortricine moths, 430
- Total evidence, 55–65, 864
 analysis, 839
- Total evidence versus relative evidence: a response to O’Leary et al. (2003), Gavin J. P. Naylor and Dean C. Adams, 864–865
- Trachemys scripta*, 286
- Tragelaphini, 209–210
- Tragelaphus*, 209, 212, 219
imberbis, 210
- Tragopan satyra*,
 Tragulidae, 206–207, 209–210, 213, 222
- Tragulina, 206–207, 209, 218, 222
- Tragulus*, 206, 209, 212–213, 215, 220
javanicus, 210
napu, 210
- Transition/transversion
 probabilities, 363
 (TI:TV) ratio, 4, 140, 372, 390, 444–445, 467, 555–556, 559, 577, 650–651, 656, 700, 708–709, 727, 748, 760, 851
- Tree-bisect-reconnect (TBR), 120
 branch-swapping, 39, 75, 166, 275, 318, 336, 377–378, 390, 407, 444–445, 463, 504, 506, 575, 607, 621–622, 666, 681, 748, 760, 801, 823–824
- Tree-construction analysis, 182
- Tree landscapes, 368–373
- Tree of Life, 431, 434, 525
- Tree squirrel, 188, 197
- Trees
 Adams consensus, 463, 850
 allozyme, 509
 Bayesian, 30, 100, 167, 455, 457, 589, 750, 762, 764, 767, 771, 773, 802
 Bayesian consensus, 622
 best-fit, 4
 bifurcating, 642
 binary, 114
 biochemical, 403
 bootstrap, 251, 380–381, 451, 455, 465
 bootstrap consensus, 609
 burn-in, 607
 canonical, 113, 115
 clocklike, 233
 combined data, 612
 consensus, 55, 128, 140, 241, 251, 255, 258, 344, 447, 531, 850, 855
 constrained, 15–18, 20, 22, 761
 constraint, 407, 847, 807
 copaths, 45
 DNA, 770, 773, 775
 equally-parsimonious, 528, 530
 equally weighted, 211
 equal rate, 233
 evolutionary, 526, 602
 gene, 63, 179, 193, 258, 374, 446, 583–584, 588, 725, 727, 776
 genealogical, 515, 517, 521, 524, 526
 haplotype, 159, 165–166
 heuristic, 607
 host, 737, 740
 likelihood, 39, 506, 508, 762, 764, 802
 logDet, 4, 7, 614, 621
 matrix-weighted, 622
 maximum likelihood, 12, 18, 20, 22, 30, 42–45, 81, 96, 128, 323, 328, 368, 371, 383, 396, 455–456, 479, 574, 576–577, 580–581, 583, 589–590, 607, 611, 666, 677, 681, 762, 767, 771, 781, 825, 827–829, 849
 maximum parsimony, 4, 110, 213, 215, 217, 240, 242–243, 248, 252, 275, 317, 320, 323, 329, 336, 339–341, 346, 359, 378, 383–384, 444, 446–447, 463, 574, 576–577, 579–581, 589–590, 607–608, 824, 826–828
 maximum parsimony combined, 16
 minimum evolution, 506, 580, 610
 minimum length, 415, 750
 mitochondrial DNA, 178
 model, 231, 423, 427, 540, 542, 761, 841–842
 molecular, 103, 105–106, 386, 389–392, 394–396, 399, 403, 853, 858, 860
 morphological, 387, 392, 394–396, 409, 503, 860
 most-parsimonious, 3, 5–7, 15, 55–60, 63, 575, 609, 621, 645, 801, 804, 809, 826, 857–857, 864
 multiple, 849–851
 near optimal, 696
 neighbor-joining, 211, 370, 455, 462–463, 508, 702, 801
 nontrivial, 485
 optimal, 4, 8, 17, 20, 121, 128, 320, 341, 357, 696, 760, 850
 parasite, 740, 742
 parsimony, 9–10, 39, 42–43, 79, 286, 645, 762, 764, 771, 773, 802–804
 phylogenetic, 119, 121, 126, 71, 112, 131, 134–135, 244, 289–290, 359, 364–365, 427, 463, 515, 554–555, 669, 671, 685, 692, 715, 749, 760–761, 771, 791, 824, 847, 849
 polyphyletic, 525
 pygopodid, 757, 760, 770, 775
 random, 30, 115, 445, 681
 rodent, 615
 rooted duplication, 110–111, 113, 115
 rooting, 757–780, 865
 single-gene, 20
 source, 727, 840–844, 846
 species, 63, 180, 725, 727
 SSR, 610–611
 star, 236
 strict component consensus, 251
 strict consensus, 57–58, 61, 323, 339, 344, 737, 826–827
 strict reduced consensus, 250
 symmetric-difference, 463
 tandem duplication, 110–118
 unconstrained, 761
 unrooted, 50, 865
 weighted, 866
- Triassic, 239–252, 749
 late-, 549, 745, 753
 lower/middle, 746
 mid-, 548
 upper, 746
- Trichogypsia*, 314–315, 317, 323
- Trichogypsiidae, 314
- Trichomycetes, 430
- Trichoptera, 106
- Trichotropidae*, 639
cancellata, 639
conica, 639
insignis, 639

- Trichotropids, 620, 623
 Trigonioidea, 559
Trimeresurus, 447, 451
 albolabris, 441
 tokarensis, 441
Tripedalia cystophora, 316
 Triticeae, 383
 tRNA, 2, 52, 94–95, 139, 296–298, 300–301, 303–308, 383, 396, 406, 460,
 615, 725, 727, 758, 762, 798
 sequences, 296, 298
 tRNA^{Asp}, 674
 tRNA^L, 73, 76
trnL-trnF, 380–382, 384
Trochita, 619, 623, 631, 639
 calyptraeformis, 631, 639
Trogon collaris, 286
 Tropidophiidae, 440
Tropidophis
 haitianus, 440
 wrighti, 440
Tropidurus, 689
 True and false gharials: a nuclear gene phylogeny of Crocodylia, John
 Harshman, Christopher J. Huddleston, Jonathan P. Bollback,
 Thomas J. Parsons, and Michael J. Braun, 386–402
 True sampling, 665–673
 Tuatara, 286
Tulearimia, 329
 Tunicata, 520
Turdus migratorius, 287
Turitella sp., 640
 Turkey, 551
Turnix sylvatica, 286
Turtur brehmeri, 38
 Two-state characters, 236
 Two-tailed test, 339
 Tylopoda, 221
 Type I error, 18, 42, 273, 279, 505, 615, 666–669, 671–672, 684, 689
 Type I STS markers are more informative than cytochrome *b* in
 phylogenetic reconstruction of the Mustelidae (Mammalia:
 Carnivora), Klaus-Peter Koepfli and Robert K. Wayne, 571–593
 Type II error, 273, 279, 505, 666, 671–672, 770
Typhlomolge, 504
 Typhlopidae, 440
Typhlops lumbricalis, 440
Typhlotriton spelaeus, 505
Typhothorax, 241, 243, 248–252
 coccinarium, 254
Tyrannus tyrannus, 286
 U2 snRNA, 557
 Ulocentra, 492–493
 Unambiguous characters, 539, 557
 Unbiased (see approximately unbiased test)
 Unconstrained
 ordination, 687
 trees, 761
 Uninformative characters, 76, 217, 575, 581, 826
 Unionoidea, 559
 Ungulates, 860
 Unrooted tree, 50, 865
 Unweighted
 pair-group average (UPGMA), 205, 229–239, 722
 UPGMA
 cluster analysis, 488, 491, 493, 496–497
 phenogram, 497
Upupa epops, 286
Urocolius macrourus, 286
 Uropygi, 557
 Ursidae, 846
Ursus, 706
 Using a null model to recognize significant co-occurrence prior to
 identifying candidate areas of endemism, Austin R. Mast and
 Reto Nyffeler, 271–280
Ute, 314–315
Uteopsis, 314–315
 Vacelet, Jean, *see* Manuel, Michael, 311–333
Vallesia, 391,
 Value of idiosyncratic markers and changes to conserved tRNA
 sequences from the mitochondrial genome of hard ticks (Acari:
 Ixodida: Ixodidae) for phylogenetic inference, Anna Murrell,
 Nick J. H. Campbell, and Stephen C. Barker, 296–310
Vanikoro, 622, 639
 sp. 1, 639
 sp. 2, 639
 Varanidae, 757
Varecia, 706
 Variable rate analysis, 751
 Vermileonidae, 756
 Vertebral characters, 439
 Vertebrate, 56, 58–59, 261–262, 264, 296, 388, 403, 515, 518, 524, 526,
 571, 706
 phylogenetics, 388
 Vespidae, 23
 Vicariance, 549–553, 717–723, 794–819
 Villet, Martin H., *see* Kelly, Christopher M., 439–459
 Violet turaco, 388
Vipera, 451
 ammodytes, 441
 ursinii, 441
 Viperidae, 439, 441, 447, 452–453, 455
 Viperinae, 441, 447, 453
 Viverridae, 846
v-myc, 388
 Volatiles, 334–351
Vombatus ursinus, 285
 Vos, R. A., Accelerated likelihood surface exploration: the likelihood
 ratchet, 368–373
Vosmaeropsis, 314–315, 320, 322–325, 328, 333
 sp., 316
 Vulture, black, 388
 vWF, 605, 614–615, 738
 Wagner, Warren L., *see* Nepokroeff, Molly, 820–841
Waldsteinia, 374, 381–383
 geoides, 374, 376, 381–382
 Wallace's Line, 794–819
 Wasps
 parasitoid, 430
 vespid, 23
 Watson-Crick pairing, 131
Waxy (see also Granule-bound starch synthase), 374–385
 Wayne, Robert K., *see* Koepfli Klaus-Peter, 571–593
 Weasels, 571–573
 long-tailed, 573
 Weevils, 717
 Weighted
 parsimony, 283
 tree, 866
 Weiss, Robert E., *see* Suchard, Marc A., 48–54, 649–664
 Whales, 707–708, 724, 853, 855, 858
 baleen, 707–708

- fin, 210
 phylogeny, 853
 toothed, 707–708
 When are phylogenetic analyses misled by convergence? a case study in Texas cave salamanders, John J. Wiens, Paul T. Chippindale, and David M. Hillis, 501–514
 When do parasites fail to speciate in response to host speciation?, Kevin P. Johnson, Richard J. Adams, Roderic D. M. Page, and Dale H. Clayton, 37–47
 Wiegmann, Brian M., David K. Yeates, Jeffrey L. Thorne, and Hirohisa Kishino, Time flies, a new molecular time-scale for Brachyceran fly evolution without a clock, 745–756
 Wiens, John J., Missing data, incomplete taxa, and phylogenetic accuracy, 528–538
 Wiens, John J., Paul T. Chippindale, and David M. Hillis, When are phylogenetic analyses misled by convergence? a case study in Texas cave salamanders, 501–514
 Wilcoxon
 rank sum, 337, 345
 signed rank test, 406, 668, 857
 T-distribution, 345
 Wilkinson, Mark, François-Joseph Lapointe, and David J. Gower, Branch lengths and support, 127–130
 Wilkinson, Mark, *see* François-Joseph Lapointe, 865–868
 Wilkinson, Mark, *see* Harris, Simon R., 239–252, 851–852
 Williams, David M. and Christopher J. Humphries, Component coding, three-item coding, and consensus methods, 255–258
 Winka, Katarina, *see* Cummings, Michael P., 477–487
 Winteraceae, 346
 Wishart distribution, 652
 Wolverine, 573
 Woodchuck, 193, 195, 197–198
 Wombat, 285
 Wright–Fisher model, 179

 Xenoderminae, 439, 441, 447, 452–453, 457
 Xenodermus, 451
 javanicus, 441
 Xenodon, 447, 451
 neuwoiedi, 441
 servus, 441

 Xenodontinae, 439, 441, 447, 452, 455–456
 Xenodontini, 447, 456
Xenophora pallidula, 640
Xenopus laevis, 286
Xenopus laevis, 286
 Xia, Xuhua, Zheng Xie, and Karl M. Kjer, 18S Ribosomal RNA and tetrapod phylogeny, 283–295
 Xie, Zheng, *see* Xia, Xuhua, 283–295
 Xiphosura, 557
Xylocopa, 8
 bombylans, 2, 5
 Xylocopinae, 1–2, 9
 Xylocopini, 1, 3, 9
Xylomya parens, 756
 Xylomyidae, 746, 756
 Xylophagidae, 756
 Xylophagid flies, 745
 Xylophagomorpha, 745–746, 750–751, 756

 Yacare caiman, 388
 Yang, Ziheng and Anne D. Yoder, Comparison of likelihood and Bayesian methods for estimating divergence times using multiple gene loci and calibration points, with application to a radiation of cute-looking mouse lemur species, 705–716
 Yeates, David K., *see* Wiegmann, Brian M., 745–756
 Yoder, Anne D., *see* Yang, Ziheng, 705–716

 Zapodidae, 614
Zapus, 614
 princeps, 606
Zeacrypta, 636
Zegalerus, 632, 639
 tenuis, 639
Zenaida, 45
 galapagoensis, 38, 45
 Zimmer, Elizabeth A., *see* Nepokroeff, Molly, 820–841
 Zorilla, 573
 Zwickl, Derrick J., *see* Hillis, David M., 124–126
 Zygethobiini, 559
Zygethobius, 559